

COMPOUNDS AND PHARMACEUTICAL
COMPOSITIONS FOR THE TREATMENT OF VIRAL INFECTIONS

CROSS REFERENCE TO RELATED APPLICATIONS

This patent applications claims the benefit of priority to 1) U.S. Provisional Appl. No. 60/877,944, filed December 28, 2006; 2) U.S Provisional Appl. No. 60/936,290, filed June 18, 2007; and 3) U.S. Provisional Application No. 60/985,891, filed November 6, 2007. The disclosures of the above referenced applications are incorporated by reference in their entirety herein.

FIELD

[0001] Provided herein are compounds, methods and pharmaceutical compositions, for use in treatment of viral infections, including hepatitis C virus infection, and hepatitis B virus infection in a host in need thereof. In a particular embodiment, phosphoroamidate or phosphonoamidate nucleoside compounds are provided which allow concentration of the drug in the liver.

BACKGROUND

Flaviviridae Viruses

[0002] The Flaviviridae family of viruses comprises at least three distinct genera: *pestiviruses*, which cause disease in cattle and pigs; *flaviviruses*, which are the primary cause of diseases such as dengue fever and yellow fever; and *hepaciviruses*, whose sole member is HCV. The flavivirus genus includes more than 68 members separated into groups on the basis of serological relatedness (Calisher *et al.*, *J. Gen. Virol.*, 1993, 70, 37-43). Clinical symptoms vary and include fever, encephalitis and hemorrhagic fever (*Fields Virology*, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, 1996, Chapter 31, 931-959). Flaviviruses of global concern that are associated with human disease include the dengue hemorrhagic fever viruses (DHF), yellow fever virus, shock syndrome and Japanese encephalitis virus (Halstead, S. B., *Rev. Infect. Dis.*, 1984, 6, 251-264; Halstead, S. B., *Science*, 239:476-481, 1988; Monath, T. P., *New Eng. J. Med.*, 1988, 319, 641-643).

[0003] The pestivirus genus includes bovine viral diarrhea virus (BVDV), classical swine fever virus (CSFV, also called hog cholera virus) and border disease

virus (BDV) of sheep (Moennig, V. *et al. Adv. Vir. Res.* 1992, 41, 53-98). Pestivirus infections of domesticated livestock (cattle, pigs and sheep) cause significant economic losses worldwide. BVDV causes mucosal disease in cattle and is of significant economic importance to the livestock industry (Meyers, G. and Thiel, H.-J., *Advances in Virus Research*, 1996, 47, 53-118; Moennig V., *et al, Adv. Vir. Res.* 1992, 41, 53-98). Human pestiviruses have not been as extensively characterized as the animal pestiviruses. However, serological surveys indicate considerable pestivirus exposure in humans.

[0004] *Pestiviruses* and *hepaciviruses* are closely related virus groups within the Flaviviridae family. Other closely related viruses in this family include the GB virus A, GB virus A-like agents, GB virus-B and GB virus-C (also called hepatitis G virus, HGV). The hepacivirus group (hepatitis C virus; HCV) consists of a number of closely related but genotypically distinguishable viruses that infect humans. There are approximately 6 HCV genotypes and more than 50 subtypes. Due to the similarities between pestiviruses and hepaciviruses, combined with the poor ability of hepaciviruses to grow efficiently in cell culture, bovine viral diarrhea virus (BVDV) is often used as a surrogate to study the HCV virus.

[0005] The genetic organization of pestiviruses and hepaciviruses is very similar. These positive stranded RNA viruses possess a single large open reading frame (ORF) encoding all the viral proteins necessary for virus replication. These proteins are expressed as a polyprotein that is co- and post-translationally processed by both cellular and virus-encoded proteinases to yield the mature viral proteins. The viral proteins responsible for the replication of the viral genome RNA are located within approximately the carboxy-terminal. Two-thirds of the ORF are termed nonstructural (NS) proteins. The genetic organization and polyprotein processing of the nonstructural protein portion of the ORF for pestiviruses and hepaciviruses is very similar. For both the pestiviruses and hepaciviruses, the mature nonstructural (NS) proteins, in sequential order from the amino-terminus of the nonstructural protein coding region to the carboxy-terminus of the ORF, consist of p7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

[0006] The NS proteins of pestiviruses and hepaciviruses share sequence domains that are characteristic of specific protein functions. For example, the NS3 proteins of viruses in both groups possess amino acid sequence motifs characteristic of serine

proteinases and of helicases (Gorbalenya *et al.* (1988) *Nature* 333:22; Bazan and Fletterick (1989) *Virology* 171:637-639; Gorbalenya *et al.* (1989) *Nucleic Acid Res.* 17:3889-3897). Similarly, the NS5B proteins of pestiviruses and hepaciviruses have the motifs characteristic of RNA-directed RNA polymerases (Koonin, E.V. and Dolja, V.V. (1993) *Crit. Rev. Biochem. Molec. Biol.* 28:375-430).

[0007] The actual roles and functions of the NS proteins of pestiviruses and hepaciviruses in the lifecycle of the viruses are directly analogous. In both cases, the NS3 serine proteinase is responsible for all proteolytic processing of polyprotein precursors downstream of its position in the ORF (Wiskerchen and Collett (1991) *Virology* 184:341-350; Bartenschlager *et al.* (1993) *J. Virol.* 67:3835-3844; Eckart *et al.* (1993) *Biochem. Biophys. Res. Comm.* 192:399-406; Grakoui *et al.* (1993) *J. Virol.* 67:2832-2843; Grakoui *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:10583-10587; Hijikata *et al.* (1993) *J. Virol.* 67:4665-4675; Tome *et al.* (1993) *J. Virol.* 67:4017-4026). The NS4A protein, in both cases, acts as a cofactor with the NS3 serine protease (Bartenschlager *et al.* (1994) *J. Virol.* 68:5045-5055; Failla *et al.* (1994) *J. Virol.* 68: 3753-3760; Lin *et al.* (1994) 68:8147-8157; Xu *et al.* (1997) *J. Virol.* 71:5312-5322). The NS3 protein of both viruses also functions as a helicase (Kim *et al.* (1995) *Biochem. Biophys. Res. Comm.* 215: 160-166; Jin and Peterson (1995) *Arch. Biochem. Biophys.*, 323:47-53; Warrenner and Collett (1995) *J. Virol.* 69:1720-1726). Finally, the NS5B proteins of pestiviruses and hepaciviruses have the predicted RNA-directed RNA polymerases activity (Behrens *et al.* (1996) *EMBO J.* 15:12-22; Lehmann *et al.* (1997) *J. Virol.* 71:8416-8428; Yuan *et al.* (1997) *Biochem. Biophys. Res. Comm.* 232:231-235; Hagedorn, PCT WO 97/12033; US patent nos. 5,981,247; 6,248,589 and 6,461,845 Zhong *et al.* (1998) *J. Virol.* 72:9365-9369).

Hepatitis C Virus

[0008] The hepatitis C virus (HCV) is the leading cause of chronic liver disease worldwide. (Boyer, N. *et al. J. Hepatol.* 32:98-112, 2000). HCV causes a slow growing viral infection and is the major cause of cirrhosis and hepatocellular carcinoma (Di Besceglie, A. M. and Bacon, B. R., *Scientific American*, Oct.: 80-85, (1999); Boyer, N. *et al. J. Hepatol.* 32:98-112, 2000). An estimated 170 million persons are infected with HCV worldwide. (Boyer, N. *et al. J. Hepatol.* 32:98-112, 2000). Cirrhosis caused by chronic hepatitis C infection accounts for 8,000-12,000

deaths per year in the United States, and HCV infection is the leading indication for liver transplantation.

[0009] HCV is known to cause at least 80% of posttransfusion hepatitis and a substantial proportion of sporadic acute hepatitis. Preliminary evidence also implicates HCV in many cases of "idiopathic" chronic hepatitis, "cryptogenic" cirrhosis, and probably hepatocellular carcinoma unrelated to other hepatitis viruses, such as Hepatitis B Virus (HBV). A small proportion of healthy persons appear to be chronic HCV carriers, varying with geography and other epidemiological factors. The numbers may substantially exceed those for HBV, though information is still preliminary; how many of these persons have subclinical chronic liver disease is unclear. (The Merck Manual, ch. 69, p. 901, 16th ed., (1992)).

[0010] HCV is an enveloped virus containing a positive-sense single-stranded RNA genome of approximately 9.4kb. The viral genome consists of a 5' untranslated region (UTR), a long open reading frame encoding a polyprotein precursor of approximately 3011 amino acids, and a short 3' UTR. The 5' UTR is the most highly conserved part of the HCV genome and is important for the initiation and control of polyprotein translation. Translation of the HCV genome is initiated by a cap-independent mechanism known as internal ribosome entry. This mechanism involves the binding of ribosomes to an RNA sequence known as the internal ribosome entry site (IRES). An RNA pseudoknot structure has recently been determined to be an essential structural element of the HCV IRES. Viral structural proteins include a nucleocapsid core protein (C) and two envelope glycoproteins, E1 and E2. HCV also encodes two proteinases, a zinc-dependent metalloproteinase encoded by the NS2-NS3 region and a serine proteinase encoded in the NS3 region. These proteinases are required for cleavage of specific regions of the precursor polyprotein into mature peptides. The carboxyl half of nonstructural protein 5, NS5B, contains the RNA-dependent RNA polymerase. The function of the remaining nonstructural proteins, NS4A and NS4B, and that of NS5A (the amino-terminal half of nonstructural protein 5) remain unknown.

[0011] A significant focus of current antiviral research is directed to the development of improved methods of treatment of chronic HCV infections in humans (Di Besceglie, A. M. and Bacon, B. R., *Scientific American*, Oct.: 80-85, (1999)).

[0012] In light of the fact that HCV infection has reached epidemic levels worldwide, and has tragic effects on the infected patient, there remains a strong need to provide new effective pharmaceutical agents to treat hepatitis C that have low toxicity to the host.

[0013] Further, given the rising threat of other flaviviridae infections, there remains a strong need to provide new effective pharmaceutical agents that have low toxicity to the host.

Hepatitis B

[0014] Hepatitis B virus has reached epidemic levels worldwide. After a two to six month incubation period in which the host is unaware of the infection, HBV infection can lead to acute hepatitis and liver damage, that causes abdominal pain, jaundice, and elevated blood levels of certain enzymes. HBV can cause fulminant hepatitis, a rapidly progressive, often fatal form of the disease in which massive sections of the liver are destroyed. Patients typically recover from acute viral hepatitis. In some patients, however, high levels of viral antigen persist in the blood for an extended, or indefinite, period, causing a chronic infection. Chronic infections can lead to chronic persistent hepatitis. Patients infected with chronic persistent HBV are most common in developing countries. Chronic persistent hepatitis can cause fatigue, cirrhosis of the liver and hepatocellular carcinoma, a primary liver cancer. In western industrialized countries, high risk groups for HBV infection include those in contact with HBV carriers or their blood samples. The epidemiology of HBV is in fact very similar to that of acquired immunodeficiency syndrome, which accounts for why HBV infection is common among patients with AIDS or HIV-associated infections. However, HBV is more contagious than HIV.

[0015] Daily treatments with α -interferon, a genetically engineered protein, have shown promise. A human serum-derived vaccine has also been developed to immunize patients against HBV. Vaccines have been produced through genetic engineering. While the vaccine has been found effective, production of the vaccine is troublesome because the supply of human serum from chronic carriers is limited, and the purification procedure is long and expensive. Further, each batch of vaccine prepared from different serum must be tested in chimpanzees to ensure safety. In addition, the vaccine does not help the patients already infected with the virus.

[0016] An essential step in the mode of action of purine and pyrimidine nucleosides against viral diseases, and in particular, HBV and HCV is their metabolic activation by cellular kinases, to yield the mono-, di- and triphosphate derivatives. The biologically active species of many nucleosides is the triphosphate form, which inhibits viral DNA polymerase, RNA polymerase, or reverse transcriptase, or causes chain termination.

[0017] In light of the fact that hepatitis B and C viruses have reached epidemic levels worldwide, and has severe and often tragic effects on the infected patient, there remains a strong need to provide new effective pharmaceutical agents to treat humans infected with the virus that have low toxicity to the host.

[0018] Therefore, there is a continuing need for effective treatments of HCV and HBV infections.

SUMMARY

[0019] Phosphoramidate and phosphonoamidate compounds of a variety of therapeutic agents are provided, as well as methods for their manufacture and use in the treatment of a variety of disorders including liver disorders. Such compounds can be used in some embodiments to permit concentration of the therapeutic agent in the liver. In one embodiment, the compound is a *S*-pivaloyl-2-thioethyl phosphoramidate, *S*-pivaloyl-2-thioethyl phosphonoamidate, *S*-hydroxypivaloyl-2-thioethyl phosphoramidate or *S*-hydroxypivaloyl-2-thioethyl phosphonoamidate.

[0020] Phosphoramidate or phosphonoamidate compounds of a variety of therapeutic agents are provided. As used herein, a "phosphoramidate or phosphonoamidate compound of a therapeutic agent" includes a therapeutic agent derivatized to include a phosphoramidate or phosphonoamidate group. The therapeutic agent is, for example, an anti-viral agent that includes, or has been derivatized to include, a reactive group, such as a hydroxyl, for attachment of the phosphoramidate or phosphonoamidate moiety. Such therapeutic agents include, but are not limited to nucleosides and nucleoside analogs including acyclic nucleosides. In some embodiments, phosphoramidates of nucleotides and nucleotide analogs are also provided, such as phosphoramidates of 1', 2', 3'-branched and 4'-branched nucleosides. Such compounds can be administered in an effective amount for the

treatment of liver disorders, including infectious diseases, such as hepatitis B and hepatitis C infection, including resistant strains thereof.

[0021] In certain embodiments, while not being limited to any theory, it is possible that the parent drug is obtained from selective metabolism of the phosphoramidate or phosphonoamidate compound in the liver, and thus the parent drug is capable of accumulating in the liver of a host. By selectively targeting and activating compounds in the liver, potentially undesired distribution of active compound in the gastrointestinal tract can be reduced. Moreover, therapeutic amounts of active compound at the site of infection in the liver can be increased.

[0022] In certain embodiments, a 5'-monophosphate or phosphonate of a parent nucleoside (or nucleoside derivative) drug is formed from metabolism of the phosphoramidate or phosphonoamidate compound in the liver, allowing the monophosphate or phosphonate to form and accumulate in the liver of a host. Thus, in certain embodiments, the phosphoramidate in effect provides a stabilized phosphate on the nucleoside or nucleoside analogue. In certain embodiments, where the compound needs to be triphosphorylated to be active, this advantageously can eliminate the requirement for the initial phosphorylation step, and promote more ready formation of the active triphosphate, which inhibits the target enzyme, and can enhance the overall activity of the nucleoside or nucleoside analog.

[0023] Without being limited to any theory, in one embodiment, a phosphoramidate of a nucleoside, such as a 2'-C-methyl-ribonucleoside, is provided, that is selectively concentrated in the liver after oral administration, and metabolized in the liver cell to yield a 5'-monophosphate that can be enzymatically converted to the active form of the 5'-triphosphate, which inhibits the HCV polymerase. Thus potentially therapeutic doses can be reduced in comparison to administering the nucleoside parent molecule.

[0024] Thus, in some embodiments, after oral administration of the phosphoramidate and phosphonoamidate compounds described herein, the compounds can advantageously concentrate in the liver cells at the site of infection and convert to the phosphate or phosphonate in the liver cell, which then is optionally further phosphorylated to implement its therapeutic effect.

[0025] Since these methods allow accumulation of the phosphoramidate or phosphonoamidate compounds disclosed herein in the liver of a host, the methods described herein can be useful, for example, for the treatment and/or prophylaxis of diseases or disorders of the liver, such as hepatitis B or C.

[0026] In certain embodiments, the compounds provided herein are useful in the prevention and treatment of *Flaviviridae* infections and other related conditions such as anti-*Flaviviridae* antibody positive and *Flaviviridae*-positive conditions, chronic liver inflammation caused by HCV, cirrhosis, fibrosis, acute hepatitis, fulminant hepatitis, chronic persistent hepatitis, and fatigue. These compounds or formulations can also be used prophylactically to prevent or retard the progression of clinical illness in individuals who are anti-*Flaviviridae* antibody or *Flaviviridae*-antigen positive or who have been exposed to a *Flaviviridae*. In one specific embodiment, the *Flaviviridae* is hepatitis C. In certain embodiments, the compound is used to treat any virus that replicates through an RNA-dependent RNA polymerase.


[0027] A method for the treatment of a *Flaviviridae* infection in a host, including a human, is also provided that includes administering an effective amount of a compound provided herein, administered either alone or in combination or alternation with another anti-*Flaviviridae* agent, optionally in a pharmaceutically acceptable carrier.

[0028] In certain embodiments, a method for treatment and/or prophylaxis of hepatitis B infections and other related conditions such as anti-HBV antibody positive and HBV-positive conditions, chronic liver inflammation caused by HBV, fibrosis, cirrhosis, acute hepatitis, fulminant hepatitis, chronic persistent hepatitis, and fatigue are provided herein.

[0029] In certain embodiments, phosphoramidate or phosphonoamidate compounds of a variety of pharmaceutical agents can be made and used therapeutically as described herein, to enhance delivery of the drug to the liver. In one embodiment, the compound is an *S*-acyl-2-thioethyl phosphoramidate or an *S*-acyl-2-thioethyl phosphonoamidate derivative, *e.g.*, a *S*-pivaloyl-2-thioethyl phosphoramidate or a *S*-hydroxypivaloyl-2-thioethyl phosphonoamidate derivative.


[0030] The phosphoramidate or phosphonoamidate compounds, as well as salts thereof, and compositions comprising the compounds, provided herein are useful for

treatment of disorders of the liver, such as HBV and/or HCV infections. In one embodiment, the compound provided herein is a compound of Formula I:



or a pharmaceutically acceptable salt, solvate, a stereoisomeric, tautomeric or polymorphic form thereof, wherein

X^a is



Z is O or S;

each W is independently O or S;

R^y and R^u each independently represent alkyl, alkenyl, alkynyl, aryl, aryl alkyl, cycloalkyl, cycloalkenyl, amino, aminoalkyl, hydroxyalkyl, alkoxy, heterocyclyl, or heteroaryl, all optionally substituted;

R^a and R^b are selected as follows:

i) R^a and R^b are each independently hydrogen, alkyl, carboxyalkyl, hydroxyalkyl, hydroxyarylalkyl, acyloxyalkyl, aminocarbonylalkyl, alkoxy carbonylalkyl, aryl, aryl alkyl, cycloalkyl, aryl, heteroaryl or heterocyclyl, all optionally substituted; or


ii) R^a and R^b together with the nitrogen atom on which they are substituted form a 3-7 membered heterocyclic or heteroaryl ring;

n is 0-3; n_2 is 1-4; and

R^1 is a moiety derivable by removal of a hydrogen from a hydroxy group of an anti-viral drug.

[0031] In another embodiment,

X^a is



Z is O, S, NH or NR^w , where R^w is, e.g., alkyl, alkyl, alkenyl, alkynyl, aryl, aryl alkyl, cycloalkyl, cycloalkenyl, amino, aminoalkyl, alkoxy, heterocyclyl, or heteroaryl, all optionally substituted;

each W is O, S, NH or NR^w, where R^w is, e.g., alkyl, alkyl, alkenyl, alkynyl, aryl, aryl alkyl, cycloalkyl, cycloalkenyl, amino, aminoalkyl, alkoxy, heterocyclyl, or heteroaryl, all optionally substituted;

R^y and R^u each independently represent alkyl, alkenyl, alkynyl, aryl, aryl alkyl, cycloalkyl, cycloalkenyl, amino, aminoalkyl, alkoxy, heterocyclyl, or heteroaryl, all optionally substituted;

R^a and R^b are selected as follows:

i) R^a and R^b are each independently hydrogen, alkyl, carboxyalkyl, hydroxyalkyl, hydroxyarylalkyl, acyloxyalkyl, aminocarbonylalkyl, alkoxy carbonylalkyl, aryl, aryl alkyl, cycloalkyl, heteroaryl or heterocyclyl, all optionally substituted; or

ii) R^a and R^b together with the nitrogen atom on which they are substituted form a 3-7 membered heterocyclic or heteroaryl ring;

n is 0-3; n₂ is 1-4; and

R¹ is as described herein.

[0032] Those of skill in the art will recognize that compounds of Formula I can be designed or prepared by reaction, e.g., at a hydroxy group of said anti-viral drug, for example, via condensation or dehydration. For convenience, in the description herein when substituents, such as exemplary R¹ groups are identified as a drug, those of skill in the art will recognize that the compound e.g. of Formula I comprises a derivative, e.g. a radical of the anti-viral drug. Those derivatives can for example be prepared by elimination of a hydrogen radical from a hydroxy group of the drug, for instance in a dehydration reaction. Where appropriate, certain derivatives can be prepared by modification of a phosphate or phosphonate of an anti-viral drug to yield a compound of formula I.

[0033] In certain embodiments of Formula I, R¹ is a nucleoside comprising a cyclic or acyclic sugar or an analog thereof.

[0034] In certain embodiments, R¹ is an anti-viral nucleoside analog useful for treatment of HCV virus infection selected from ribavirin, viremagine, 2'-C-methylcytidine, 2'-C-methylguanosine, valopicitabine (NM 283), MK-0608 and PSI-6130.

[0035] In certain embodiments, R¹ is an anti-viral nucleoside analog useful for treatment of HBV virus infection selected from lamivudine (Epivir-HBV, Zeffix, or


Heptodin), adefovir, entecavir (Baraclude), telbivudine (Tyzeka, Sebivo), emtricitabine (FTC), clevudine (L-FMAU), viread (Tenofovir), torcitabine, valtorcitabine (monoval LdC), amdoxovir (DAPD) and RCV (Racivir).

[0036] In certain embodiments, R^1 is a non-nucleoside anti-viral useful for treatment of HBV virus infection selected from resiquimod or celgosivir.

[0037] In certain embodiments according to Formula I, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are each independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^y is $-C(CH_3)_2CH_2OH$.

[0038] In certain embodiments, the compounds provided herein are selected such that R^1 is not 3'-azido-2',3'-dideoxythymidine.

[0039] In another embodiment, the compound provided herein is a compound of Formula IIa or IIb:



or a pharmaceutically acceptable salt, solvate, a stereoisomeric, tautomeric or polymorphic form thereof, wherein

R^y is alkyl, alkenyl, alkynyl, aryl, aryl alkyl, cycloalkyl, cycloalkenyl, amino, aminoalkyl, hydroxyalkyl, heterocyclyl or heteroaryl, all optionally substituted;

R^a and R^b are selected as follows:

i) R^a and R^b are each independently hydrogen, alkyl, carboxyalkyl, hydroxyalkyl, hydroxyarylalkyl, acyloxyalkyl, aminocarbonylalkyl,


alkoxycarbonylalkyl, aryl, aryl alkyl, cycloalkyl, aryl, heteroaryl or heterocyclyl, all optionally substituted; or

ii) R^a and R^b together with the nitrogen atom on which they are substituted form a 3-7 membered heterocyclic or heteroaryl ring; and

R^1 is an antiviral drug (as used herein where R^1 is an antiviral drug, that embodiment includes a moiety derivable by removal of a hydrogen from a hydroxy group of an anti-viral drug), such as a nucleoside or nucleoside analog.

[0040] In certain embodiments according to Formula IIa or IIb, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^y is $-C(CH_3)_2CH_2OH$.

[0041] In another embodiment, the compound provided herein is a compound of Formula:




wherein R^a , R^b and R^y are as described in Formula I and

wherein R^2 and R^3 are each independently H, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, aryl alkylsulfonyl, a lipid, such as a phospholipid; an amino acid; and amino acid residue, a carbohydrate; a peptide; cholesterol; or other pharmaceutically acceptable leaving group which is capable of providing a compound wherein R^2 and/or R^3 is independently H, for example when administered *in vivo*; or R^2 and R^3 are linked to form a cyclic group by an alkyl, ester or carbamate linkage; and wherein each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, aryl alkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments according to this paragraph, R^2 and R^3 are each H; R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen,

alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^2 and R^3 are each H; R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^2 and R^3 are each H; R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^2 and R^3 are each H; R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^2 and R^3 are each H; R^y is $-C(CH_3)_2CH_2OH$. In certain embodiments according to this paragraph, R^2 and R^3 are each hydrogen, R^a is hydrogen, R^b is $-CH_2-C_6H_5$ and R^y is $-C(CH_3)_2CH_2OH$.


[0042] In another embodiment, the compound provided herein is a compound of formula:




wherein R^a , R^b and R^y are as described in Formula I. R^d is selected from the group consisting of hydrogen, alkyl and alkoxy. In certain embodiments, R^d is hydrogen, methyl or methoxy. In certain embodiments according to this paragraph, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^y is $-OR^c$, $-C(R^c)_3$ or


-NHR^c where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^y is $\text{-C(CH}_3)_2\text{CH}_2\text{OH}$. In certain embodiments according to this paragraph, R^2 and R^3 are each hydrogen, R^a is hydrogen, R^b is $\text{-CH}_2\text{-C}_6\text{H}_5$ and R^y is $\text{-C(CH}_3)_2\text{CH}_2\text{OH}$. In certain embodiments according to this paragraph, R^a is hydrogen, R^b is $\text{-CH}_2\text{-C}_6\text{H}_5$ and R^y is $\text{-C(CH}_3)_2\text{CH}_2\text{OH}$.

[0043] In one embodiment, the nucleosides that can be derivatized to include a phosphoramidate or phosphonoamidate, *e.g.* at the 5' position include:




[0044] Examples of phosphoramidate or phosphonoamidate nucleoside compounds include:








and




[0045] In one embodiment, the nucleosides that can be derivatized to include a phosphoramidate or phosphonoamidate, *e.g.* at the 5' position include:




and




[0046] In one embodiment, phosphoramidate or phosphonoamidate nucleoside compounds include:






[0047] In one embodiment, the nucleosides that can be derivatized to include a phosphoramidate or phosphonoamidate, *e.g.* at the 5' position include:



[0048] In one embodiment, phosphoramidate or phosphonoamidate nucleoside compounds include:



[0049] In one aspect, the compounds described herein are provided or administered in combination with a second therapeutic agent, such as one useful for the treatment or prevention of HBV and/or HCV infections. Exemplary therapeutic agents are described in detail in the sections below.

[0050] In another aspect, provided are pharmaceutical compositions, single unit dosage forms, and kits suitable for use in treating or preventing disorders such as HBV and/or HCV infections which comprise a therapeutically or prophylactically effective amount of a compound described herein, e.g. of Formula I, IIa or IIb, and a therapeutically or prophylactically effective amount of a second therapeutic such as one useful for the treatment or prevention of HBV and/or HCV infections.

[0051] In certain embodiments, a method of treatment of a liver disorder is provided comprising administering to an individual in need thereof a treatment

effective amount of a phosphoramidate or phosphonoamidate derivative of a nucleoside or nucleoside analogue, wherein optionally the derivative is a *S*-pivaloyl-2-thioethyl phosphoramidate or *S*-pivaloyl-2-thioethyl phosphonoamidate derivative. The derivative is optionally selected from the compounds disclosed herein.

[0052] In some embodiments, provided herein are:

- (a) compounds as described herein, *e.g.* of Formula I, IIa or IIb, and pharmaceutically acceptable salts and compositions thereof;
- (b) compounds as described herein, *e.g.* of Formula I, IIa or IIb, and pharmaceutically acceptable salts and compositions thereof for use in the treatment and/or prophylaxis of a liver disorder including *Flaviviridae* infection, especially in individuals diagnosed as having a *Flaviviridae* infection or being at risk of becoming infected by hepatitis C;
- (c) processes for the preparation of compounds as described herein, *e.g.* of Formula I, IIa or IIb, as described in more detail below;
- (d) pharmaceutical formulations comprising a compound as described herein, *e.g.* of Formula I, IIa or IIb, or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent;
- (e) pharmaceutical formulations comprising a compound as described herein, *e.g.* of Formula I, IIa or IIb, or a pharmaceutically acceptable salt thereof together with one or more other effective anti-HCV agents, optionally in a pharmaceutically acceptable carrier or diluent;
- (f) a method for the treatment and/or prophylaxis of a host infected with *Flaviviridae* that includes the administration of an effective amount of a compound as described herein, *e.g.* of Formula I, IIa or IIb, its pharmaceutically acceptable salt or composition;
- (g) a method for the treatment and/or prophylaxis of a host infected with *Flaviviridae* that includes the administration of an effective amount of a compounds as described herein, *e.g.* of Formula I, IIa or IIb, its pharmaceutically acceptable salt or composition in combination and/or alternation with one or more effective anti-HCV agent;

- (h) compounds as described herein, *e.g.* of Formula I, IIa or IIb, and pharmaceutically acceptable salts and compositions thereof for use in the treatment and/or prophylaxis of a HBV infection, especially in individuals diagnosed as having an HBV infection or being at risk of becoming infected by hepatitis B;
- (i) pharmaceutical formulations comprising a compound as described herein, *e.g.* of Formula I, IIa or IIb, or a pharmaceutically acceptable salt thereof together with one or more other effective anti-HBV agents, optionally in a pharmaceutically acceptable carrier or diluent;
- (j) a method for the treatment and/or prophylaxis of hepatitis B infections and other related conditions such as anti-HBV antibody positive and HBV-positive conditions, chronic liver inflammation caused by HBV, cirrhosis, acute hepatitis, fulminant hepatitis, chronic persistent hepatitis, and fatigue that includes administering an effective amount of a compound as described herein, *e.g.* of Formula I, IIa or IIb, or its pharmaceutically acceptable salt or composition.
- (k) a prophylactic method to prevent or retard the progression of clinical illness in individuals who are anti-HBV antibody or HBV-antigen positive or who have been exposed to HBV.

[0053] *Flaviviridae* which can be treated are, *e.g.*, discussed generally in *Fields Virology*, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, Chapter 31, 1996. In a particular embodiment of the invention, the *Flaviviridae* is HCV. In an alternate embodiment, the *Flaviviridae* is a flavivirus or pestivirus. Specific flaviviruses include, without limitation: Absettarov, Alfuy, Apoi, Aroa, Bagaza, Banzi, Bouboui, Bussuquara, Cacipacore, Carey Island, Dakar bat, Dengue 1, Dengue 2, Dengue 3, Dengue 4, Edge Hill, Entebbe bat, Gadgets Gully, Hanzalova, Hypr, Ilheus, Israel turkey meningoencephalitis, Japanese encephalitis, Jugra, Jutiapa, Kadam, Karshi, Kedougou, Kokobera, Koutango, Kumlinge, Kunjin, Kyasanur Forest disease, Langat, Louping ill, Meaban, Modoc, Montana myotis leukoencephalitis, Murray valley encephalitis, Naranjal, Negishi, Ntaya, Omsk hemorrhagic fever, Phnom-Penh bat, Powassan, Rio Bravo, Rocio, Royal Farm, Russian spring-summer encephalitis, Saboya, St. Louis encephalitis, Sal

Vieja, San Perlita, Saumarez Reef, Sepik, Sokuluk, Spondweni, Stratford, Tembusu, Tyuleniy, Uganda S, Usutu, Wesselsbron, West Nile, Yaounde, Yellow fever, and Zika.

[0054] Pestiviruses which can be treated are discussed generally in *Fields Virology*, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, Chapter 33, 1996. Specific pestiviruses include, without limitation: bovine viral diarrhea virus ("BVDV"), classical swine fever virus ("CSFV," also called hog cholera virus), and border disease virus ("BDV").

BRIEF DESCRIPTION OF DRAWINGS

[0055] Figure 1 depicts depletion of NM108 hydroxySATE phosphoramidate (B299) after incubation with and without NADPH in monkey liver S9.

[0056] Figure 2 depicts depletion of NM107 hydroxySATE phosphoramidate (B102) after incubation with and without NADPH in monkey liver S9.

DESCRIPTION OF EXEMPLARY EMBODIMENTS

[0057] Provided herein are compounds, compositions and methods useful for treating liver disorders such as HBV and/or HCV infection in a subject. Further provided are dosage forms useful for such methods.

Definitions

[0058] When referring to the compounds provided herein, the following terms have the following meanings unless indicated otherwise.

[0059] The term "alkyl", as used herein, unless otherwise specified, includes a saturated straight, branched, or cyclic, primary, secondary, or tertiary hydrocarbon of typically C₁ to C₁₀, and specifically includes methyl, CF₃, CCl₃, CFCl₂, CF₂Cl, ethyl, CH₂CF₃, CF₂CF₃, propyl, isopropyl, cyclopropyl, butyl, isobutyl, secbutyl, *t*-butyl, pentyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, cyclohexylmethyl, 3-methylpentyl, 2,2-dimethylbutyl, and 2,3-dimethylbutyl. The term includes both substituted and unsubstituted alkyl groups, and particularly includes halogenated alkyl groups, and even more particularly fluorinated alkyl groups. Non-limiting examples of moieties with which the alkyl group can be substituted are selected from the group consisting of halogen (fluoro, chloro, bromo or iodo), hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or

protected as necessary, as known to those skilled in the art, for example, as taught in Greene, *et al.*, Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991, hereby incorporated by reference.

[0060] The term "lower alkyl", as used herein, and unless otherwise specified, includes a C₁ to C₄ saturated straight, branched, or if appropriate, a cyclic (for example, cyclopropyl) alkyl group, including both substituted and unsubstituted moieties.

[0061] "Alkylene" includes divalent saturated aliphatic hydrocarbon groups particularly having up to about 11 carbon atoms and more particularly 1 to 6 carbon atoms which can be straight-chained or branched. This term is exemplified by groups such as methylene (-CH₂-), ethylene (-CH₂CH₂-), the propylene isomers (*e.g.*, -CH₂CH₂CH₂- and -CH(CH₃)CH₂-) and the like.

[0062] "Alkenyl" includes monovalent olefinically unsaturated hydrocarbon groups, in certain embodiment, having up to about 11 carbon atoms, from 2 to 8 carbon atoms, or from 2 to 6 carbon atoms, which can be straight-chained or branched and having at least 1 or from 1 to 2 sites of olefinic unsaturation. Exemplary alkenyl groups include ethenyl (-CH=CH₂), n-propenyl (-CH₂CH=CH₂), isopropenyl (-C(CH₃)=CH₂), vinyl and substituted vinyl, and the like.

[0063] "Alkenylene" includes divalent olefinically unsaturated hydrocarbon groups, in certain embodiments, having up to about 11 carbon atoms or from 2 to 6 carbon atoms which can be straight-chained or branched and having at least 1 or from 1 to 2 sites of olefinic unsaturation. This term is exemplified by groups such as ethenylene (-CH=CH-), the propenylene isomers (*e.g.*, -CH=CHCH₂- and -C(CH₃)=CH- and -CH=C(CH₃)-) and the like.

[0064] "Alkynyl" includes acetylenically unsaturated hydrocarbon groups, in certain embodiments, having up to about 11 carbon atoms or from 2 to 6 carbon atoms which can be straight-chained or branched and having at least 1 or from 1 to 2 sites of alkynyl unsaturation. Non-limiting examples of alkynyl groups include acetylenic, ethynyl (-C≡CH), propargyl (-CH₂C≡CH), and the like.

[0065] The term "aryl", as used herein, and unless otherwise specified, includes phenyl, biphenyl, or naphthyl, and preferably phenyl. The term includes both substituted and unsubstituted moieties. The aryl group can be substituted with any

described moiety, including, but not limited to, one or more moieties selected from the group consisting of halogen (fluoro, chloro, bromo or iodo), alkyl, haloalkyl, hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene, *et al.*, *Protective Groups in Organic Synthesis*, John Wiley and Sons, Second Edition, 1991.

[0066] "Alkoxy" includes the group -OR where R is alkyl. Particular alkoxy groups include, by way of example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, tert-butoxy, sec-butoxy, n-pentoxy, n-hexoxy, 1,2-dimethylbutoxy, and the like.

[0067] "Alkoxycarbonyl" includes a radical -C(O)-alkoxy where alkoxy is as defined herein.

[0068] "Amino" includes the radical -NH₂.

[0069] "Carboxyl" includes the radical -C(O)OH.

[0070] The term "alkylamino" or "arylamino" includes an amino group that has one or two alkyl or aryl substituents, respectively. Unless otherwise specifically stated in this application, when alkyl is a suitable moiety, lower alkyl is preferred. Similarly, when alkyl or lower alkyl is a suitable moiety, unsubstituted alkyl or lower alkyl is preferred.

[0071] "Halogen" or "halo" includes chloro, bromo, fluoro or iodo.

[0072] "Monoalkylamino" includes the group alkyl-NR', wherein R' is selected from hydrogen and alkyl.

[0073] "Thioalkoxy" includes the group -SR where R is alkyl.

[0074] The term "protected" as used herein and unless otherwise defined refers to a group that is added to an oxygen, nitrogen, or phosphorus atom to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis.

[0075] "Pharmaceutically acceptable salt" includes any salt of a compound provided herein which retains its biological properties and which is not toxic or otherwise undesirable for pharmaceutical use. Such salts may be derived from a

variety of organic and inorganic counter-ions well known in the art. Such salts include: (1) acid addition salts formed with organic or inorganic acids such as hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, sulfamic, acetic, trifluoroacetic, trichloroacetic, propionic, hexanoic, cyclopentylpropionic, glycolic, glutaric, pyruvic, lactic, malonic, succinic, sorbic, ascorbic, malic, maleic, fumaric, tartaric, citric, benzoic, 3-(4-hydroxybenzoyl)benzoic, picric, cinnamic, mandelic, phthalic, lauric, methanesulfonic, ethanesulfonic, 1,2-ethane-disulfonic, 2-hydroxyethanesulfonic, benzenesulfonic, 4-chlorobenzenesulfonic, 2-naphthalenesulfonic, 4-toluenesulfonic, camphoric, camphorsulfonic, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic, glucoheptonic, 3-phenylpropionic, trimethylacetic, *tert*-butylacetic, lauryl sulfuric, gluconic, benzoic, glutamic, hydroxynaphthoic, salicylic, stearic, cyclohexylsulfamic, quinic, muconic acid and the like acids; or (2) salts formed when an acidic proton present in the parent compound either (a) is replaced by a metal ion, *e.g.*, an alkali metal ion, an alkaline earth ion or an aluminum ion, or alkali metal or alkaline earth metal hydroxides, such as sodium, potassium, calcium, magnesium, aluminum, lithium, zinc, and barium hydroxide, ammonia or (b) coordinates with an organic base, such as aliphatic, alicyclic, or aromatic organic amines, such as ammonia, methylamine, dimethylamine, diethylamine, picoline, ethanolamine, diethanolamine, triethanolamine, ethylenediamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylene-diamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethylamine, *N*-methylglucamine piperazine, tris(hydroxymethyl)-aminomethane, tetramethylammonium hydroxide, and the like.

[0076] Salts further include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium and the like, and when the compound contains a basic functionality, salts of non-toxic organic or inorganic acids, such as hydrohalides, *e.g.* hydrochloride and hydrobromide, sulfate, phosphate, sulfamate, nitrate, acetate, trifluoroacetate, trichloroacetate, propionate, hexanoate, cyclopentylpropionate, glycolate, glutarate, pyruvate, lactate, malonate, succinate, sorbate, ascorbate, malate, maleate, fumarate, tartarate, citrate, benzoate, 3-(4-hydroxybenzoyl)benzoate, picrate, cinnamate, mandelate, phthalate, laurate, methanesulfonate (mesylate), ethanesulfonate, 1,2-ethane-disulfonate, 2-hydroxyethanesulfonate, benzenesulfonate (besylate), 4-chlorobenzenesulfonate, 2-

naphthalenesulfonate, 4-toluenesulfonate, camphorate, camphorsulfonate, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylate, glucoheptonate, 3-phenylpropionate, trimethylacetate, *tert*-butylacetate, lauryl sulfate, gluconate, benzoate, glutamate, hydroxynaphthoate, salicylate, stearate, cyclohexylsulfamate, quinate, muconate and the like.

[0077] The term "alkaryl" or "alkylaryl" includes an aryl group with an alkyl substituent. The term aralkyl or arylalkyl includes an alkyl group with an aryl substituent.

[0078] The term "purine" or "pyrimidine" base includes, but is not limited to, adenine, N⁶-alkylpurines, N⁶-acylpurines (wherein acyl is C(O)(alkyl, aryl, alkylaryl, or arylalkyl), N⁶-benzylpurine, N⁶-halopurine, N⁶-vinylpurine, N⁶-acetylenic purine, N⁶-acyl purine, N⁶-hydroxyalkyl purine, N⁶-alkylaminopurine, N⁶-thioalkyl purine, N²-alkylpurines, N²-alkyl-6-thiopurines, thymine, cytosine, 5-fluorocytosine, 5-methylcytosine, 6-azapyrimidine, including 6-azacytosine, 2- and/or 4-mercaptopyrimidine, uracil, 5-halouracil, including 5-fluorouracil, C⁵-alkylpyrimidines, C⁵-benzylpyrimidines, C⁵-halopyrimidines, C⁵-vinylpyrimidine, C⁵-acetylenic pyrimidine, C⁵-acyl pyrimidine, C⁵-hydroxyalkyl purine, C⁵-amidopyrimidine, C⁵-cyanopyrimidine, C⁵-iodopyrimidine, C⁶-iodo-pyrimidine, C⁵-Br-vinyl pyrimidine, C⁶-Br-vinyl pyrimidine, C⁵-nitropyrimidine, C⁵-amino-pyrimidine, N²-alkylpurines, N²-alkyl-6-thiopurines, 5-azacytidinyl, 5-azauracil, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, and pyrazolopyrimidinyl. Purine bases include, but are not limited to, guanine, adenine, hypoxanthine, 7-deazaguanine, 7-deazaadenine, 2,6-diaminopurine, and 6-chloropurine. Functional oxygen and nitrogen groups on the base can be protected as necessary or desired. Suitable protecting groups are well known to those skilled in the art, and include trimethylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, and *t*-butyldiphenylsilyl, trityl, alkyl groups, and acyl groups such as acetyl and propionyl, methanesulfonyl, and *p*-toluenesulfonyl.

[0079] The term "acyl" or "O-linked ester" includes a group of the formula C(O)R', wherein R' is an straight, branched, or cyclic alkyl (including lower alkyl), carboxylate residue of amino acid, aryl including phenyl, alkaryl, arylalkyl including benzyl, alkoxyalkyl including methoxymethyl, aryloxyalkyl such as phenoxymethyl; or substituted alkyl (including lower alkyl), aryl including phenyl optionally

substituted with chloro, bromo, fluoro, iodo, C₁ to C₄ alkyl or C₁ to C₄ alkoxy, sulfonate esters such as alkyl or arylalkyl sulphonyl including methanesulfonyl, the mono, di or triphosphate ester, trityl or monomethoxy-trityl, substituted benzyl, alkaryl, arylalkyl including benzyl, alkoxyalkyl including methoxymethyl, aryloxyalkyl such as phenoxymethyl. Aryl groups in the esters optimally comprise a phenyl group. In particular, acyl groups include acetyl, trifluoroacetyl, methylacetyl, cyclopropylacetyl, propionyl, butyryl, hexanoyl, heptanoyl, octanoyl, neo-heptanoyl, phenylacetyl, 2-acetoxy-2-phenylacetyl, diphenylacetyl, α -methoxy- α -trifluoromethyl-phenylacetyl, bromoacetyl, 2-nitro-benzeneacetyl, 4-chloro-benzeneacetyl, 2-chloro-2,2-diphenylacetyl, 2-chloro-2-phenylacetyl, trimethylacetyl, chlorodifluoroacetyl, perfluoroacetyl, fluoroacetyl, bromodifluoroacetyl, methoxyacetyl, 2-thiopheneacetyl, chlorosulfonylacetyl, 3-methoxyphenylacetyl, phenoxyacetyl, tert-butylacetyl, trichloroacetyl, monochloro-acetyl, dichloroacetyl, 7H-dodecafluoro-heptanoyl, perfluoro-heptanoyl, 7H-dodeca-fluoroheptanoyl, 7-chlorododecafluoro-heptanoyl, 7-chloro-dodecafluoro-heptanoyl, 7H-dodecafluoroheptanoyl, 7H-dodeca-fluoroheptanoyl, nona-fluoro-3,6-dioxahexanoyl, nonafluoro-3,6-dioxahexanoyl, perfluoroheptanoyl, methoxybenzoyl, methyl 3-amino-5-phenylthiophene-2-carboxyl, 3,6-dichloro-2-methoxy-benzoyl, 4-(1,1,2,2-tetrafluoro-ethoxy)-benzoyl, 2-bromo-propionyl, omega-aminocapryl, decanoyl, n-pentadecanoyl, stearyl, 3-cyclopentyl-propionyl, 1-benzene-carboxyl, O-acetylmandelyl, pivaloyl acetyl, 1-adamantane-carboxyl, cyclohexane-carboxyl, 2,6-pyridinedicarboxyl, cyclopropane-carboxyl, cyclobutane-carboxyl, perfluorocyclohexyl carboxyl, 4-methylbenzoyl, chloromethyl isoxazolyl carbonyl, perfluorocyclohexyl carboxyl, crotonyl, 1-methyl-1H-indazole-3-carbonyl, 2-propenyl, isovaleryl, 1-pyrrolidinecarbonyl, 4-phenylbenzoyl.

[0080] The term "amino acid" includes naturally occurring and synthetic α , β γ or δ amino acids, and includes but is not limited to, amino acids found in proteins, *i.e.* glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine. In a preferred embodiment, the amino acid is in the L-configuration. Alternatively, the amino acid can be a derivative of alanyl, valinyl, leucinyl, isoleuccinyl, prolinyl, phenylalaninyl, tryptophanyl, methioninyl, glyciny, serinyl, threoninyl, cysteinyl, tyrosinyl, asparaginyl, glutaminyl, aspartoyl,

glutaroyl, lysinyl, argininy, histidinyl, β -alanyl, β -valinyl, β -leucinyl, β -isoleuccinyl, β -prolinyl, β -phenylalaninyl, β -tryptophanyl, β -methioninyl, β -glycinyl, β -serinyl, β -threoninyl, β -cysteinyl, β -tyrosinyl, β -asparaginyl, β -glutaminyl, β -aspartoyl, β -glutaroyl, β -lysiny, β -argininy or β -histidinyl.

[0081] As used herein, the term “substantially free of” or “substantially in the absence of” with respect to a nucleoside composition includes a nucleoside composition that includes at least 85 or 90% by weight, preferably 95%, 98 % , 99% or 100% by weight, of the designated enantiomer of that nucleoside. In a preferred embodiment, in the methods and compounds of this invention, the compounds are substantially free of enantiomers.

[0082] Similarly, the term “isolated” with respect to a nucleoside composition includes a nucleoside composition that includes at least 85, 90%, 95%, 98%, 99% to 100% by weight, of the nucleoside, the remainder comprising other chemical species or enantiomers.

[0083] “Solvate” includes a compound provided herein or a salt thereof, that further includes a stoichiometric or non-stoichiometric amount of solvent bound by non-covalent intermolecular forces. Where the solvent is water, the solvate is a hydrate.

[0084] The term “host”, as used herein, includes any unicellular or multicellular organism in which the virus can replicate, including cell lines and animals, and preferably a human. Alternatively, the host can be carrying a part of the *Flaviviridae* viral genome, whose replication or function can be altered by the compounds of the present invention. The term host specifically includes infected cells, cells transfected with all or part of the *Flaviviridae* genome and animals, in particular, primates (including chimpanzees) and humans. In most animal applications of the present invention, the host is a human patient. Veterinary applications, in certain indications, however, are clearly anticipated by the present invention (such as chimpanzees).

[0085] As used herein, the terms “subject” and “patient” are used interchangeably herein. The terms “subject” and “subjects” refer to an animal, such as a mammal including a non-primate (e.g., a cow, pig, horse, cat, dog, rat, and mouse) and a primate (e.g., a monkey such as a cynomolgous monkey, a chimpanzee and a human), and for example, a human. In one embodiment, the subject is refractory or non-

responsive to current treatments for hepatitis C infection. In another embodiment, the subject is a farm animal (*e.g.*, a horse, a cow, a pig, etc.) or a pet (*e.g.*, a dog or a cat). In one embodiment, the subject is a human.

[0086] As used herein, the terms “therapeutic agent” and “therapeutic agents” refer to any agent(s) which can be used in the treatment or prevention of a disorder or one or more symptoms thereof. In certain embodiments, the term “therapeutic agent” includes a compound provided herein. In one embodiment, a therapeutic agent is an agent which is known to be useful for, or has been or is currently being used for the treatment or prevention of a disorder or one or more symptoms thereof.

[0087] “Therapeutically effective amount” includes an amount of a compound or composition that, when administered to a subject for treating a disease, is sufficient to effect such treatment for the disease. A “therapeutically effective amount” can vary depending on, *inter alia*, the compound, the disease and its severity, and the age, weight, *etc.*, of the subject to be treated.

[0088] “Treating” or “treatment” of any disease or disorder refers, in one embodiment, to ameliorating a disease or disorder that exists in a subject. In another embodiment, “treating” or “treatment” includes ameliorating at least one physical parameter, which may be indiscernible by the subject. In yet another embodiment, “treating” or “treatment” includes modulating the disease or disorder, either physically (*e.g.*, stabilization of a discernible symptom) or physiologically (*e.g.*, stabilization of a physical parameter) or both. In yet another embodiment, “treating” or “treatment” includes delaying the onset of the disease or disorder.

[0089] As used herein, the terms “prophylactic agent” and “prophylactic agents” as used refer to any agent(s) which can be used in the prevention of a disorder or one or more symptoms thereof. In certain embodiments, the term “prophylactic agent” includes a compound provided herein. In certain other embodiments, the term “prophylactic agent” does not refer a compound provided herein. For example, a prophylactic agent is an agent which is known to be useful for, or has been or is currently being used to the prevent or impede the onset, development, progression and/or severity of a disorder.


[0090] As used herein, the phrase “prophylactically effective amount” includes the amount of a therapy (*e.g.*, prophylactic agent) which is sufficient to result in the

prevention or reduction of the development, recurrence or onset of one or more symptoms associated with a disorder (, or to enhance or improve the prophylactic effect(s) of another therapy (e.g., another prophylactic agent).


Compounds

[0091] Phosphoramidate and phosphonoamidate compounds of a variety of therapeutic agents can be formed using methods available in the art and those disclosed herein. Such compounds can be used in some embodiments to enhance delivery of the drug to the liver. In one embodiment, the compound is an *S*-acyl-2-thioethyl phosphoramidate or an *S*-acyl-2-thioethyl phosphonoamidate derivative, e.g., an *S*-pivaloyl-2-thioethyl phosphoroamidate, an *S*-hydroxypivaloyl-2-thioethyl phosphoroamidate, an *S*-pivaloyl-2-thioethyl phosphonoamidate or an *S*-hydroxypivaloyl-2-thioethyl phosphonoamidate. Therapeutic agents that can be derivatized to compound form include an anti-viral agent that includes, or has been derivatized to include a reactive group for attachment of the phosphoramidate or phosphonoamidate moiety, including but not limited to nucleosides and nucleoside analogues including acyclic nucleosides. Therapeutic agents that can be derivatized to compound form also include an anti-viral agent that includes, or has been derivatized to include a phosphate or phosphonate group that can be derivatized to form a phosphoramidate or phosphonoamidate moiety, including but not limited to nucleosides and nucleoside analogues including acyclic nucleosides.

[0092] Nucleosides that can be derivatized include any R^1 as disclosed herein. Examples of nucleosides that can be derivatized to include a phosphoramidate or phosphonoamidate, e.g. at the 5', 3' or 2' position include:




Examples of phosphoramidate or phosphonoamidate nucleoside compounds include:



[0093] Phosphoramidate or phosphonoamidate compounds of other nucleosides and nucleoside analogues described herein and known in the art can be formed as described herein and used for the treatment of liver disorders. The phosphoramidate or phosphonoamidate moiety can be e.g., at the 5' position.

[0094] In one embodiment, provided herein are compounds, as well as salts thereof, and compositions comprising the compounds, that are useful for treatment of disorders of the liver, including HBV and/or HCV infections. In one embodiment, the phosphoramidate or phosphonoamidate compound provided herein is a compound of Formula IIa or IIb:



or a pharmaceutically acceptable salt, solvate, a stereoisomeric, tautomeric or polymorphic form thereof, wherein;

R^y is alkyl, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, cycloalkenyl, amino, heterocyclyl or heteroaryl, all optionally substituted;

R^a and R^b are selected as follows:

i) R^a and R^b are each independently hydrogen, alkyl, carboxyalkyl, hydroxyalkyl, hydroxyarylalkyl, acyloxyalkyl, aminocarbonylalkyl, alkoxycarbonylalkyl, aryl, arylalkyl, cycloalkyl, heteroaryl or heterocyclyl, all optionally substituted; or

ii) R^a and R^b together with the nitrogen atom on which they are substituted form a 3-7 membered heterocyclic or heteroaryl ring; and

R¹ is an anti-viral drug (which includes a moiety derivable by removal of a hydrogen from a hydroxy group of an anti-viral drug).

In certain embodiments, the compound of Formula IIa or IIb is selected with a proviso that when R^y is *tert*-butyl or hydroxy-*tert*-butyl, then R¹ is not 3'-azido-2',3'-dideoxythymidine.

[0095] In certain embodiments, R¹, R^a, R^b and R^y are optionally substituted with one or more substituents as defined in the definitions.

[0096] In certain embodiments, the compounds are of Formula IIa or IIb, wherein R^y is alkyl, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, cycloalkenyl, amino, heterocyclyl or heteroaryl;





R^a and R^b are each independently hydrogen, alkyl, carboxyalkyl, hydroxyalkyl, hydroxyarylalkyl, acyloxyalkyl, aminocarbonylalkyl, alkoxycarbonylalkyl, aryl, arylalkyl, cycloalkyl, heteroaryl or heterocyclyl, all optionally substituted; and






R¹ is an anti-viral drug (which is meant to include a moiety derivable by removal of a hydrogen from a hydroxy group of an anti-viral drug).

[0097] In one embodiment, R¹ is a nucleoside comprising a cyclic or acyclic sugar or analog thereof, including any any nucleoside or analogue thereof described herein or known in the art.




[0098] Exemplary nucleoside drugs useful in the treatment of hepatitis C infection that can be derivatized as described herein are:

Name	Structure
------	-----------




Name	Structure
Ribavirin	
Viramidine	
Valopicitabine (NM283)	
2'-C-methylcytidine (NM107)	







Name	Structure
PSI-6130	
MK-0608	
7-Fluoro-MK-0608	
NM108	
	




[0099] Exemplary non-nucleoside drugs that can be derivatized as described herein are:

Name	Structure
Resiquimod	
Celgosivir	
Gliotoxin	


[00100] Exemplary nucleoside drugs useful in the treatment of hepatitis B infection that can be derivatized as described herein are:

Name	Structure
Lamivudine or 3TC or Epivir®	
Entecavir	
Telbivudine or L-dT	


Name	Structure
Racivir	
Emtricitabine or (-)FTC	
Clevudine or L-FMAU	
Amdoxovir	
Valtorcitabine	
Torcitabine (L-dC)	

Name	Structure
Tenofovir or PMPA	
Adefovir or PMEAs	
L-cytidine	



[00101] A phosphoramidate or phosphonoamidate compound of acyclovir, L-ddA or D-ddA can be administered for treatment of Hepatitis B, an example of which is shown below:




[00102] Where the nucleoside analog already includes a phosphonate, that phosphonate group can be incorporated in the phosphonoamidate moiety shown in the formulas herein, as shown by way of example in the phosphonoamidate of adefovir:



[00103] Thus, in certain embodiments of the compounds of Formula IIa below:



is derived from a drug that is an acyclic nucleoside phosphonate, i.e.:



which is e.g. PME A (9-[2-(phosphonomethoxy)ethyl]adenine (adefovir).

[00104] In certain embodiments according to Formula IIa or IIb, R^y is substituted alkyl, e.g. hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl.

[00105] In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^y is $-C(CH_3)_2CH_2OH$. In certain


embodiments according to this paragraph, R^2 and R^3 are each hydrogen, R^a is hydrogen, R^b is $-\text{CH}_2-\text{C}_6\text{H}_5$ and R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$.

[00106] In one embodiment, R^y is alkyl or hydroxyalkyl. In one embodiment, R^y is methyl, *tert*-butyl, hydroxy-*tert*-butyl or hydroxyethyl. In certain embodiments, R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$.

[00107] In one embodiment, R^a and R^b are each independently hydrogen, alkyl, carboxyalkyl, hydroxyalkyl, hydroxyarylalkyl, acyloxyalkyl, aminocarbonylalkyl, alkoxycarbonylalkyl, aryl or arylalkyl, wherein the alkyl groups can be further substituted with one or more substituents. In one embodiment, at least one of R^a or R^b is other than hydrogen. In one embodiment, R^a and R^b are each independently hydrogen, methyl or benzyl.

[00108] In certain embodiments, R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$ and R^a and R^b are each independently hydrogen, methyl or benzyl. In certain embodiments, R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$ and R^a is hydrogen and R^b is benzyl.

[00109] In another embodiment, the compound provided herein is a compound of formula:



wherein R^1 and R^y are as defined in Formula IIa or IIb.

In certain embodiments of Formula IIIa, b, c or d:

R^y is substituted alkyl, *e.g.*, hydroxyalkyl or aminoalkyl; and


In another embodiment, R^y is $-\text{OR}^c$, $-\text{C}(\text{R}^c)_3$ or $-\text{NHR}^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy-

or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or hydroxy-, amino-, alkyl-, haloalkyl- or trifluoromethyl- substituted benzyl. In certain embodiments, R^a and R^b together with the nitrogen atom on which they are substituted form a 3-7 membered heterocyclic or heteroaryl ring.

In one embodiment, R^y is alkyl or hydroxyalkyl. In one embodiment, R^y is methyl, *tert*-butyl, hydroxy-*tert*-butyl or hydroxyethyl. In one embodiment, R^y is $-C(CH_3)_2CH_2OH$.

[00110] In certain embodiments according to Formula IIIa or IIIb, R^y is substituted alkyl, e.g. hydroxyalkyl or aminoalkyl. In another embodiment, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^y is $-C(CH_3)_2CH_2OH$.

[00111] In another embodiment, the compound provided herein is a compound of formula:



wherein R^1 , R^a and R^b are e.g. as defined in Formula IIa or IIb.

[00112] In certain embodiments of Formula IVa or IVb:

R^1 is an antiviral drug, such as a nucleoside or nucleoside derivative; and
 R^a and R^b are each independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or hydroxy-, amino-, alkyl-, haloalkyl- or trifluoromethyl- substituted benzyl. In a further embodiment, R^a and R^b are independently H, benzyl or substituted alkyl. In certain embodiments, R^a and R^b together with the nitrogen atom on which they are substituted form a 3-7 membered heterocyclic or heteroaryl ring.


[00113] In certain embodiments of Formula IVa or IVb:

R^1 is an antiviral drug, such as a nucleoside or nucleoside derivative; and

R^a and R^b are each independently hydrogen, alkyl, carboxyalkyl, hydroxyalkyl, hydroxyarylalkyl, acyloxyalkyl, aminocarbonylalkyl, alkoxy carbonylalkyl, aryl, arylalkyl, cycloalkyl, heteroaryl, or heterocyclyl, all optionally substituted; and

wherein, in one embodiment, one of R^a and R^b is H and the other is alkyl optionally substituted with aryl, benzyl, or heteroaryl, each optionally substituted.

[00114] In another embodiment, the compound provided herein is a compound of formula:



wherein R^1 is as defined in Formula IIa or IIb.

[00115] In certain embodiments, R^1 is an anti-viral nucleoside analog useful for treatment of HCV virus infection selected from ribavirin, viramidine, 2'-C-methylcytidine, 2'-C-methylguanosine, valopicitabine (NM 283), MK-0608 and PSI-6130. As used herein, where R^1 is an analogue of a nucleoside, such as acyclovir, that itself includes a phosphonate group, that phosphonate can be in the form of the phosphonoamidate in the formulas disclosed herein. Thus, e.g., in formula Va or Vb, the $R^1P(O)O^-$ fragment is derived from the nucleoside analog that includes a phosphonate.

[00116] In certain embodiments, R^1 is an anti-viral nucleoside analog useful for treatment of HBV virus infection selected from lamivudine (Epivir-HBV, Zeffix, or Heptodin), adefovir, entecavir (Baraclude), telbivudine (Tyzeka, Sebivo), emtricitabine (FTC), clevudine (L-FMAU), viread (Tenofovir), torcitabine, valtorcitabine (monoval LdC), amdoxovir (DAPD) and RCV (Racivir).


[00117] Further exemplary anti-viral nucleoside analogs that can be used as R^1 are disclosed in International Publication Nos. WO2005021568; WO2006094347 and WO2006093987 and US Patent Publication No. US20050215510.

[00118] In certain embodiments, R^1 is a non-nucleoside anti-viral useful for treatment of HBV virus infection selected from resiquimod or celgosivir.

[00119] In one embodiment, R^1 is an immunosuppressant, such as combretastatin A-4, mycophenolic acid, pentostatin, nelarabine or mitoxantrone.

[00120] In one embodiment, R^1 is an interfering RNA (iRNA) based antivirals, including short interfering RNA (siRNA) based antivirals. Such compounds are described in International Patent Publication Nos. WO/03/070750 and WO 2005/012525, US patent Nos. 7,176,304; 7,109,165; 7,041,817; 7,034,009; 7,022,828; 6,852,535 and 6,849,726 and US Patent Publication No. US 2004/0209831.


[00121] In another embodiment, the compound provided herein is a compound of formula:



wherein R^a , R^b and R^y are as defined in Formula IIa or IIb and R^2 and R^3 are each independently H; straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, a lipid, such as a phospholipid; an amino acid; and amino acid residue, a carbohydrate; a peptide; cholesterol; or other pharmaceutically acceptable leaving group which is capable of providing a compound wherein R^2 and/or R^3 is

independently H or phosphate (including mono-, di- or triphosphate), for example when administered *in vivo*; or R^2 and R^3 are linked to form a cyclic group by an alkyl, ester or carbamate linkage. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments, R^2 and R^3 are each hydrogen, R^a is hydrogen, R^b is $-\text{CH}_2-\text{C}_6\text{H}_5$ and R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$. In certain embodiments according to this paragraph, R^2 and R^3 are each H; R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^2 and R^3 are each H; R^y is $-\text{OR}^c$, $-\text{C}(\text{R}^c)_3$ or $-\text{NHR}^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^2 and R^3 are each H; R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^2 and R^3 are each H; R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^2 and R^3 are each H; R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$.


[00122] In another embodiment, the compound provided herein is a compound of formula:



wherein R^a, R^b and R^y are as defined in Formula IIa or IIb and R^c is hydrogen or alkyl. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments, R^c is methyl, ethyl or propyl, R^a is hydrogen, R^b is -CH₂-C₆H₅ and R^y is -C(CH₃)₂CH₂OH. In certain embodiments according to this paragraph, R² and R³ are each H; R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^c is methyl, ethyl or propyl; R² and R³ are each H; R^y is -OR^c, -C(R^c)₃ or -NHR^c where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^c is methyl, ethyl or

propyl; R^2 and R^3 are each H; R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^2 and R^3 are each H; R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^e is methyl, ethyl or propyl; R^2 and R^3 are each H; R^y is $-C(CH_3)_2CH_2OH$. In certain embodiments according to this paragraph, R^1 is chosen from nucleosides described in U.S. Patent Application Publication No. US 2006/0111324 A1, the content of which is hereby incorporated by reference in its entirety.


[00123] In another embodiment, the compound provided herein is a compound of formula:



wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments according to this paragraph, each R^L is hydrogen, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl,


for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, each R^L is hydrogen, R^Y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, each R^L is hydrogen, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^Y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, each R^L is hydrogen and R^Y is $-C(CH_3)_2CH_2OH$.

[00124] In another embodiment, the compound provided herein is a compound of formula:



[00125] wherein R^a , R^b and R^Y are as defined in Formula IIa or IIb. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl,


arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments according to this paragraph, each R^L is hydrogen, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, each R^L is hydrogen, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, each R^L is hydrogen, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, each R^L is hydrogen and R^y is $-C(CH_3)_2CH_2OH$. In another embodiment, the compound provided herein is a compound of formula:



wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl,

wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments according to this paragraph, each R^L is hydrogen, R^Y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, each R^L is hydrogen, R^Y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, each R^L is hydrogen, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^Y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, each R^L is hydrogen and R^Y is $-C(CH_3)_2CH_2OH$.


[00126] In another embodiment, the compound provided herein is a compound of formula:



[00127] wherein R^a , R^b and R^Y are as defined in Formula IIa or IIb. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including


lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments according to this paragraph, each R^L is hydrogen, R^Y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, each R^L is hydrogen, R^Y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, each R^L is hydrogen, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^Y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, each R^L is hydrogen and R^Y is $-C(CH_3)_2CH_2OH$.

[00128] In another embodiment, the compound provided herein is a compound of formula:




[00129] wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments according to this paragraph, each R^L is hydrogen, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, each R^L is hydrogen, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, each R^L is hydrogen, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, each R^L is hydrogen and R^y is $-C(CH_3)_2CH_2OH$.

[00130] In another embodiment, the compound provided herein is a compound of formula:



[00131] wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments according to this paragraph, each R^L is hydrogen, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, each R^L is hydrogen, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, each R^L is hydrogen, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, each R^L is hydrogen and R^y is $-C(CH_3)_2CH_2OH$.


[00132] In another embodiment, the compound provided herein is a compound of formula:



[00133] wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including

lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments according to this paragraph, each R^L is hydrogen, R^Y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, each R^L is hydrogen, R^Y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, each R^L is hydrogen, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^Y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, each R^L is hydrogen and R^Y is $-C(CH_3)_2CH_2OH$.

[00134] In certain embodiments, 2-deoxy-2-fluoro-2-C-ethynyl- β -D-nucleosides can be formed and derived into phosphoramidate compounds to potentiate delivery of an active monophosphate to the liver of an individual inflicted with HCV, such as the compounds described herein by way of example. In certain embodiments, a compound of the following formula is provided:



wherein:

$T = O, S, CH_2, CH(hal)$ or $CH(hal)_2, S(O)_n$;

$n = 1, 2$;

hal = halogen;

$R = H$, acyl (with lower linear and non linear alkyl -C1 to 6-, aminoacid), monophosphate, diphosphate, triphosphate, monophosphate prodrug such as (alkyl-

O)₂PO, (tBuSate-O)₂PO, cyclic monophosphate prodrug, phosphoramidate prodrug (aromatic amine, aminoacid);

X and Y are independently H, OH, O-alkyl (lower), O-acyl, F, NH₂, NH-alkyl, N-dialkyl, NH-acyl, N-diacyl, or azido;

Z is H, alkyl, alkenyl, alkynyl, hydroxymethyl, fluoromethyl, or azido;


W is H, alkyl, alkenyl, alkynyl, hydroxymethyl, fluoromethyl, azido, carboxylic acid, CO₂-alkyl, cyano, or carboxamide;

A is H, alkyl, alkenyl, alkynyl, hydroxymethyl, fluoromethyl, azido, carboxylic acid, CO₂-alkyl, cyano, or carboxamide; and

Base is a natural or modified base.

Optionally the compounds include a chlorine atom at the 2'-position.


[00135] In another embodiment, the compound provided herein is a compound of formula:



wherein R^a, R^b and R^y are as defined in Formula IIa or IIb; A is H, alkyl, alkenyl, alkynyl, hydroxymethyl, fluoromethyl, azido, carboxylic acid, CO₂-alkyl, cyano, or carboxamide; and R^{b1} is halo, alkoxy or haloalkyl. Each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; an amino acid residue; or a carbohydrate. In certain embodiments according to this paragraph, each

R^L is hydrogen, R^Y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, each R^L is hydrogen, R^Y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, each R^L is hydrogen, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^Y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, each R^L is hydrogen and R^Y is $-C(CH_3)_2CH_2OH$.


[00136] In another embodiment, the compound provided herein is a compound of formula:



wherein R^a , R^b and R^Y are as defined in Formula IIa or IIb. In certain embodiments, R^a is hydrogen, R^b is $-CH_2-C_6H_5$ and R^Y is $-C(CH_3)_2CH_2OH$. In certain embodiments according to this paragraph, R^Y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In

another embodiment, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^y is $-C(CH_3)_2CH_2OH$.


[00137] In another embodiment, the compound provided herein is a compound of formula:



wherein X is halogen, R^a , R^b and R^y are as defined in Formula IIa or IIb and R^2 and R^3 are each independently H, straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, a lipid, such as a phospholipid; an amino acid; and amino acid residue, a carbohydrate; a peptide; cholesterol; or other pharmaceutically acceptable leaving group which is capable of providing a compound wherein R^2 and/or R^3 is independently H or phosphate (including mono-, di- or triphosphate), for example when administered *in vivo*; or R^2 and R^3 are linked to form a cyclic group by an alkyl, ester or carbamate linkage. R^L is hydrogen or any lipophilic group known to those of skill in the art. In certain embodiments, R^2 and R^3 are each hydrogen, R^a is hydrogen, R^b is $-CH_2-C_6H_5$ and R^y is $-C(CH_3)_2CH_2OH$. In certain embodiments said lipophilic group is selected from alkyl, alkenyl, cycloalkyl, aryl, heteroaryl, arylalkyl and heteroaryl-alkyl. In certain embodiments according to this paragraph, X is fluoro, R^L is hydrogen, R^2 and R^3 are each H, R^y is substituted alkyl, e.g. hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-


substituted alkyl or benzyl. In another embodiment, X is fluoro, R^L is hydrogen, R^2 and R^3 are each H, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, X is fluoro, R^L is hydrogen, R^2 and R^3 are each H, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, X is fluoro, R^L is hydrogen, R^2 and R^3 are each H, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, X is fluoro, R^L is hydrogen, R^2 and R^3 are each H, R^y is $-C(CH_3)_2CH_2OH$.

[00138] In another embodiment, the compound provided herein is a compound of formula:




wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. In certain embodiments, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^a is hydrogen, R^b is $-CH_2-C_6H_5$ and R^y is $-C(CH_3)_2CH_2OH$.

[00139] In another embodiment, the compound provided herein is a compound of formula:



[00140] wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. In certain embodiments, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^a is hydrogen, R^b is $-\text{CH}_2-\text{C}_6\text{H}_5$ and R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$.


[00141] In another embodiment, the compound provided herein is a compound of formula:



[00142] wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. In certain embodiments, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^y is $-OR^c$, $-C(R^c)_3$ or $-NHR^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further

embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^a is hydrogen, R^b is $-\text{CH}_2-\text{C}_6\text{H}_5$ and R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$.


[00143] In another embodiment, the compound provided herein is a compound of formula:



[00144] wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. In certain embodiments, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^y is $-\text{OR}^c$, $-\text{C}(\text{R}^c)_3$ or $-\text{NHR}^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^a is hydrogen, R^b is $-\text{CH}_2-\text{C}_6\text{H}_5$ and R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$.


[00145]

[00146] In another embodiment, the compound provided herein is a compound of formula:




wherein wherein R^a , R^b and R^y are as defined in Formula IIa or IIb. R^L is hydrogen or any lipophilic group known to those of skill in the art. In certain embodiments, R^a is hydrogen, R^b is $-\text{CH}_2-\text{C}_6\text{H}_5$ and R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$. In certain embodiments said lipophilic group is selected from alkyl, alkenyl, cycloalkyl, aryl, heteroaryl, arylalkyl and heteroaryl-alkyl. In certain embodiments according to this paragraph, R^y is substituted alkyl, *e.g.* hydroxyalkyl or aminoalkyl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In another embodiment, R^y is $-\text{C}(\text{R}^c)_3$ or $-\text{NHR}^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl, for instance hydroxy- or amino-substituted alkyl or aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl, for instance hydroxy- or amino-substituted alkyl or benzyl. In a further embodiment, R^a and R^b are independently benzyl or substituted alkyl. In a further embodiment, R^y is selected from the group consisting of alkyl and hydroxyalkyl. In certain embodiments, R^y is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$.

[00147] In another embodiment, the compound provided herein is a compound of formula:



wherein the variables are as described above.

[00148] In another embodiment, the compound provided herein is a compound of formula:



wherein the variables are as described above.

[00149] In one embodiment, R^1 is a natural nucleoside. In one embodiment, R^1 is a 2'- or 3'- prodrug of biologically active 1', 2', 3' or 4'C-branched β -D or β -L


nucleoside. The term 1', 2', 3' or 4'-C-branched, as used in this specification, includes a nucleoside that has an additional non-natural substituent in the 1', 2', 3' or 4'-position (*i.e.*, carbon is bound to four nonhydrogen substituents). The term 2'-prodrug, as used herein, includes a 1', 2', 3' or 4' C-branched- β -D or β -L nucleoside that has a biologically cleavable moiety at the 2'-position, including, but not limited to acyl, and in one embodiment, a natural or synthetic D or L amino acid, in one embodiment, an L-amino acid. The term 3'-prodrug, as used herein, includes a 1', 2', 3' or 4' C-branched- β -D or β -L nucleoside that has a biologically cleavable moiety at the 3'-position, including, but not limited to acyl, and in one embodiment, a natural or synthetic D or L amino acid, in one embodiment, an L-amino acid. In one embodiment, the amino acid is valine.

[00150] Examples of prodrugs (that can be further derivatized as described herein to include a phosphoramidate or phosphonoamidate moiety, for example, at the 5' position) include 2'-L-valine ester of β -D-2'-C-methyl-cytidine; 2'-L-valine ester of β -D-2'-C-methyl-thymidine; 2'-L-valine ester of β -D-2'-C-methyl-adenosine; 2'-L-valine ester of β -D-2'-C-methyl-guanosine; 2'-L-valine ester of β -D-2'-C-methyl-5-fluorocytidine; 2'-L-valine ester of β -D-2'-C-methyl-uridine; 2'-acetyl ester of β -D-2'-C-methyl-cytidine; 2'-acetyl ester of β -D-2'-C-methyl-thymidine; 2'-acetyl ester of β -D-2'-C-methyl-adenosine; 2'-acetyl ester of β -D-2'-C-methyl-guanosine; 2'-acetyl ester of β -D-2'-C-methyl-5-fluoro-cytidine; and 2'-esters of β -D-2'-C-methyl-(cytidine, 5-fluorocytidine, guanosine, uridine, adenosine, or thymidine) wherein (i) the 2' ester is an amino acid ester; or (ii) the 2' ester is an alkyl or aryl ester.

[00151] Further examples of prodrugs are 3'-L-valine ester of β -D-2'-C-methyl-cytidine; 3'-L-valine ester of β -D-2'-C-methyl-thymidine; 3'-L-valine ester of β -D-2'-C-methyl-adenosine; 3'-L-valine ester of β -D-2'-C-methyl-guanosine; 3'-L-valine ester of β -D-2'-C-methyl-5-fluorocytidine; 3'-L-valine ester of β -D-2'-C-methyl-uridine; 3'-acetyl ester of β -D-2'-C-methyl-cytidine; 3'-acetyl ester of β -D-2'-C-methyl-thymidine; 3'-acetyl ester of β -D-2'-C-methyl-adenosine; 3'-acetyl ester of β -D-2'-C-methyl-guanosine; 3'-acetyl ester of β -D-2'-C-methyl-5-fluoro-cytidine; and 3'-esters of β -D-2'-C-methyl-(cytidine, 5-fluorocytidine, guanosine, uridine, adenosine, or thymidine) wherein (i) the 3' ester is an amino acid ester; or (ii) the 3' ester is an alkyl or aryl ester.

[00152] Additional examples of prodrugs include 2',3'-L-divaline ester of β -D-2'-C-methyl-cytidine (dival-2'-Me-L-dC); 2',3'-L-divaline ester of β -D-2'-C-methyl-thymidine; 2',3'-L-divaline ester of β -D-2'-C-methyl-adenosine; 2',3'-L-divaline ester of β -D-2'-C-methyl-guanosine; 2',3'-L-divaline ester of β -D-2'-C-methyl-5-fluoro-cytidine; 2',3'-L-divaline ester of β -D-2'-C-methyl-uridine; 2',3'-diacetyl ester of β -D-2'-C-methyl-cytidine; 2',3'-diacetyl ester of β -D-2'-C-methyl-thymidine; 2',3'-diacetyl ester of β -D-2'-C-methyl-adenosine; 2',3'-diacetyl ester of β -D-2'-C-methyl-guanosine; 2',3'-diacetyl ester of β -D-2'-C-methyl-5-fluoro-cytidine; and 2',3'-diesters of β -D-2'-C-methyl-(cytidine, 5-fluorocytidine, guanosine, uridine, adenosine, or thymidine) wherein (i) the 2' ester is an amino acid ester and the 3' ester is an alkyl or aryl ester; (ii) both esters are amino acid esters; (iii) both esters are independently alkyl or aryl esters; or (iv) the 2' ester is an alkyl or aryl ester and the 3'-ester is an amino acid ester.

[00153] In one embodiment, R^1 is:



wherein Base is a natural or non-natural purine or pyrimidine base as defined herein;

R^6 is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, azido, cyano, Br-vinyl, alkoxy, acyloxy, alkoxycarbonyl, alkenyloxy, halo, NO_2 or $\text{NR}^{6a}\text{R}^{6b}$;

R^{6a} and R^{6b} are each independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, acyl, aryl, heteroaryl or heterocyclyl;

R^7 , R^9 , R^8 and R^{10} are selected as follows:

i) R^7 and R^9 are each independently hydrogen, OR^{7a} , hydroxy, alkyl, alkenyl, alkynyl, azido, cyano, Br-vinyl, alkylloxycarbonyl, acyloxy, halo, NO_2 or $\text{NR}^{6a}\text{R}^{6b}$;

ii) R^8 and R^{10} are each independently H, alkyl or halo; or


iii) each R^7 and R^9 , R^7 and R^{10} , R^8 and R^9 or R^8 and R^{10} together form a double bond;

R^{7a} is H; straight chained, branched or cyclic alkyl (including lower alkyl); acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl,

CO-substituted aryl, sulfonate ester including alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted with one or more substituents as described in the definition of aryl given herein; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, including a phospholipid; an amino acid; and amino acid residue, a carbohydrate; a peptide; cholesterol; or other pharmaceutically acceptable leaving group which is capable of providing a compound wherein R^{7a} is H or phosphate (including mono-, di- or triphosphate), for example, when administered *in vivo*; wherein in one embodiment R^{7a} is not phosphate (including mono-, di- or triphosphate or a stabilized phosphate prodrug), or two R^{7a} groups are linked to form a cyclic group by an alkyl, ester or carbamate linkage; and

X is O, S, SO_2 or CH_2 .

[00154] In one embodiment, R^1 has formula:



[00155] wherein R^2 and R^3 are each independently H; straight chained, branched or cyclic alkyl; acyl (including lower acyl); CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester such as alkyl or arylalkyl sulfonyl including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted; alkylsulfonyl, arylsulfonyl, arylalkylsulfonyl, a lipid, such as a phospholipid; an amino acid; and amino acid residue, a carbohydrate; a peptide; cholesterol; or other pharmaceutically acceptable leaving group which is capable of providing a compound wherein R^2 and/or R^3 is independently H or phosphate (including mono-, di- or triphosphate), for example when administered *in vivo*; or R^2 and R^3 are linked to form a cyclic group by an alkyl, ester or carbamate linkage;

[00156] wherein Y^1 is hydrogen, bromo, chloro, fluoro, iodo, CN, OH, OR^4 , NH_2 , NHR^4 , NR^4R^5 , SH or SR^4 ;

[00157] X^1 is a straight chained, branched or cyclic optionally substituted alkyl, CH_3 , CF_3 , $C(Y^3)_3$, 2-Br-ethyl, CH_2F , CH_2Cl , CH_2CF_3 , CF_2CF_3 , $C(Y^3)_2C(Y^3)_3$, CH_2OH , optionally substituted alkenyl, optionally substituted alkynyl, COOH,

COOR⁴, COO-alkyl, COO-aryl, CO-Oalkoxyalkyl, CONH₂, CONHR⁴, CON(R⁴)₂, chloro, bromo, fluoro, iodo, CN, N₃, OH, OR⁴, NH₂, NHR⁴, NR⁴R⁵, SH or SR⁵; and

[00158] X² is H, straight chained, branched or cyclic optionally substituted alkyl, CH₃, CF₃, C(Y³)₃, 2-Br-ethyl, CH₂F, CH₂Cl, CH₂CF₃, CF₂CF₃, C(Y³)₂C(Y³)₃, CH₂OH, optionally substituted alkenyl, optionally substituted alkynyl, COOH, COOR⁴, COO-alkyl, COO-aryl, CO-Oalkoxyalkyl, CONH₂, CONHR⁴, CON(R⁴)₂, chloro, bromo, fluoro, iodo, CN, N₃, OH, OR⁴, NH₂, NHR⁴, NR⁴R⁵, SH or SR⁵; and


[00159] wherein each Y³ is independently H, F, Cl, Br or I;

[00160] each R⁴ and R⁵ is independently hydrogen, acyl (including lower acyl), alkyl (including but not limited to methyl, ethyl, propyl and cyclopropyl), lower alkyl, alkenyl, alkynyl or cycloalkyl.

[00161] In the embodiments described herein, R² and/or R³ may be a pharmaceutically acceptable leaving group which is capable of providing a compound wherein R² and/or R³ is independently H or phosphate (including mono-, di- or triphosphate), for example when administered *in vivo*.


[00162] In another embodiment, each R² and R³ is independently hydrogen or acyl. In another embodiment, R² and R³ are linked to form a cyclic group by an alkyl, ester or carbamate linkage.

[00163] In another embodiment, R¹ is:




[00164] wherein R², R³, Y¹, Y³, X¹ and X² are as defined in Formula XIII.


[00165] In one embodiment, R¹ is:




wherein Base is selected from the group consisting of




(A)




(B)




(C)




(D)




(E)







(F)



(G)



(H)




(I)


(J)

(K)


(L)




(M)




(N)



(O)




(P)




(Q)


(R)




(S)




(T)




(U)




(V)



(W)







(Y)



(Z)

(X)




(AA)


(AB)

(AC)


(AD)




(AE)




(AF)




(AG)




(AH)




(AI)









(AJ)













(BA)











(BB)






(BQ)

(BR)





[00166] wherein each W^1 , W^2 , W^3 and W^4 is independently N, CH, CF, Cl, CBr, CCl, CCN, CCH₃, CCF₃, CCH₂CH₃, CC(O)NH₂, CC(O)NHR⁴, CC(O)N(R⁴)₂, CC(O)OH, CC(O)OR⁴ or CX³;

[00167] each W^* is independently O, S, NH or NR⁴;

[00168] X is O, S, SO₂, CH₂, CH₂OH, CHF, CF₂, C(Y³)₂, CHCN, C(CN)₂, CHR⁴ or C(R⁴)₂;

[00169] X* is CH, CF, CY³ or CR⁴;

[00170] X² is H, straight chained, branched or cyclic optionally substituted alkyl, CH₃, CF₃, C(Y³)₃, 2-Br-ethyl, CH₂F, CH₂Cl, CH₂CF₃, CF₂CF₃, C(Y³)₂C(Y³)₃, CH₂OH, optionally substituted alkenyl, optionally substituted alkynyl, COOH, COOR⁴, COO-alkyl, COO-aryl, CO-Oalkoxyalkyl, CONH₂, CONHR⁴, CON(R⁴)₂, chloro, bromo, fluoro, iodo, CN, N₃, OH, OR⁴, NH₂, NHR⁴, NR⁴R⁵, SH or SR⁵;

[00171] each X³ is independently a straight chained, branched or cyclic optionally substituted alkyl (including lower alkyl), CH₃, CH₂CN, CH₂N₃, CH₂NH₂,

CH₂NHCH₃, CH₂N(CH₃)₂, CH₂OH, halogenated alkyl (including halogenated lower alkyl), CF₃, C(Y³)₃, 2-Br-ethyl, CH₂F, CH₂Cl, CH₂CF₃, CF₂CF₃, C(Y³)₂C(Y³)₃, optionally substituted alkenyl, haloalkenyl, Br-vinyl, optionally substituted alkynyl, haloalkynyl, N₃, CN, -C(O)OH, -C(O)OR⁴, -C(O)O(lower alkyl), -C(O)NH₂, -C(O)NHR⁴, -C(O)NH(lower alkyl), -C(O)N(R⁴)₂, -C(O)N(lower alkyl)₂, OH, OR⁴, -O(acyl), -O(lower acyl), -O(alkyl), -O(lower alkyl), -O(alkenyl), -O(alkynyl), -O(arylalkyl), -O(cycloalkyl), -S(acyl), -S(lower acyl), -S(R⁴), -S(lower alkyl), -S(alkenyl), -S(alkynyl), -S(arylalkyl), -S(cycloalkyl), chloro, bromo, fluoro, iodo, NH₂, -NH(lower alkyl), -NHR⁴, -NR⁴R⁵, -NH(acyl), -N(lower alkyl)₂, -NH(alkenyl), -NH(alkynyl), -NH(arylalkyl), -NH(cycloalkyl), -N(acyl)₂;

[00172] each Y is independently selected from the group consisting of H, optionally substituted lower alkyl, cycloalkyl, alkenyl, alkynyl, CH₂OH, CH₂NH₂, CH₂NHCH₃, CH₂N(CH₃)₂, CH₂F, CH₂Cl, CH₂N₃, CH₂CN, CH₂CF₃, CF₃, CF₂CF₃, CH₂CO₂R, (CH₂)_mCOOH, (CH₂)_mCOOR, (CH₂)_mCONH₂, (CH₂)_mCONR₂, and (CH₂)_mCONHR;

[00173] wherein R is H, alkyl or acyl;

[00174] Y¹ is hydrogen, bromo, chloro, fluoro, iodo, CN, OH, OR⁴, NH₂, NHR⁴, NR⁴R⁵, SH or SR⁴;

[00175] each Y² is independently O, S, NH or NR⁴;

[00176] each Y³ is independently H, F, Cl, Br or I;

[00177] each R⁴ and R⁵ is independently hydrogen, acyl (including lower acyl), alkyl (including but not limited to methyl, ethyl, propyl and cyclopropyl), lower alkyl, alkenyl, alkynyl or cycloalkyl;

[00178] each R⁶ is independently an optionally substituted alkyl (including lower alkyl), CH₃, CH₂CN, CH₂N₃, CH₂NH₂, CH₂NHCH₃, CH₂N(CH₃)₂, CH₂OH, halogenated alkyl (including halogenated lower alkyl), CF₃, C(Y³)₃, 2-Br-ethyl, CH₂F, CH₂Cl, CH₂CF₃, CF₂CF₃, C(Y³)₂C(Y³)₃, optionally substituted alkenyl, haloalkenyl, Br-vinyl, optionally substituted alkynyl, haloalkynyl, -CH₂C(O)OH, -CH₂C(O)OR⁴, -CH₂C(O)O(lower alkyl), -CH₂C(O)NH₂, -CH₂C(O)NHR⁴, -CH₂C(O)NH(lower alkyl), -CH₂C(O)N(R⁴)₂, -CH₂C(O)N(lower alkyl)₂, -(CH₂)_mC(O)OH, -(CH₂)_mC(O)OR⁴, -(CH₂)_mC(O)O(lower alkyl), -(CH₂)_mC(O)NH₂, -(CH₂)_mC(O)NHR⁴, -(CH₂)_mC(O)NH(lower alkyl), -(CH₂)_mC(O)N(R⁴)₂,


$-(CH_2)_mC(O)N(lower\ alkyl)_2$, $-C(O)OH$, $-C(O)OR^4$, $-C(O)O(lower\ alkyl)$, $-C(O)NH_2$, $-C(O)NHR^4$, $-C(O)NH(lower\ alkyl)$, $-C(O)N(R^4)_2$, $-C(O)N(lower\ alkyl)_2$ or cyano;

[00179] each R^7 is independently H, OH, OR^2 , optionally substituted alkyl (including lower alkyl), CH_3 , CH_2CN , CH_2N_3 , CH_2NH_2 , CH_2NHCH_3 , $CH_2N(CH_3)_2$, CH_2OH , halogenated alkyl (including halogenated lower alkyl), CF_3 , $C(Y^3)_3$, 2-Br-ethyl, CH_2F , CH_2Cl , CH_2CF_3 , CF_2CF_3 , $C(Y^3)_2C(Y^3)_3$, optionally substituted alkenyl, haloalkenyl, Br-vinyl, optionally substituted alkynyl, haloalkynyl, optionally substituted carbocycle (for example, a 3-7 membered carbocyclic ring), optionally substituted heterocycle (for example, a 3-7 membered heterocyclic ring having one or more O, S and/or N), optionally substituted heteroaryl (for example, a 3-7 membered heteroaromatic ring having one or more O, S and/or N), $-CH_2C(O)OH$, $-CH_2C(O)OR^4$, $-CH_2C(O)O(lower\ alkyl)$, $-CH_2C(O)SH$, $-CH_2C(O)SR^4$, $-CH_2C(O)S(lower\ alkyl)$, $-CH_2C(O)NH_2$, $-CH_2C(O)NHR^4$, $-CH_2C(O)NH(lower\ alkyl)$, $-CH_2C(O)N(R^4)_2$, $-CH_2C(O)N(lower\ alkyl)_2$, $-(CH_2)_mC(O)OH$, $-(CH_2)_mC(O)OR^4$, $-(CH_2)_mC(O)O(lower\ alkyl)$, $-(CH_2)_mC(O)SH$, $-(CH_2)_mC(O)SR^4$, $-(CH_2)_mC(O)S(lower\ alkyl)$, $-(CH_2)_mC(O)NH_2$, $-(CH_2)_mC(O)NHR^4$, $-(CH_2)_mC(O)NH(lower\ alkyl)$, $-(CH_2)_mC(O)N(R^4)_2$, $-(CH_2)_mC(O)N(lower\ alkyl)_2$, $-C(O)OH$, $-C(O)OR^4$, $-C(O)O(lower\ alkyl)$, $-C(O)SH$, $-C(O)SR^4$, $-C(O)S(lower\ alkyl)$, $-C(O)NH_2$, $-C(O)NHR^4$, $-C(O)NH(lower\ alkyl)$, $-C(O)N(R^4)_2$, $-C(O)N(lower\ alkyl)_2$, $-O(acyl)$, $-O(lower\ acyl)$, $-O(R^4)$, $-O(alkyl)$, $-O(lower\ alkyl)$, $-O(alkenyl)$, $-O(alkynyl)$, $-O(arylalkyl)$, $-O(cycloalkyl)$, $-S(acyl)$, $-S(lower\ acyl)$, $-S(R^4)$, $-S(lower\ alkyl)$, $-S(alkenyl)$, $-S(alkynyl)$, $-S(arylalkyl)$, $-S(cycloalkyl)$, NO_2 , NH_2 , $-NH(lower\ alkyl)$, $-NHR^4$, $-NR^4R^5$, $-NH(acyl)$, $-N(lower\ alkyl)_2$, $-NH(alkenyl)$, $-NH(alkynyl)$, $-NH(arylalkyl)$, $-NH(cycloalkyl)$, $-N(acyl)_2$, azido, cyano, SCN, OCN, NCO or halo (fluoro, chloro, bromo, iodo);


[00180] alternatively, R^6 and R^7 can come together to form a spiro compound selected from the group consisting of optionally substituted carbocycle (for example, a 3-7 membered carbocyclic ring) or optionally substituted heterocycle (for example, a 3-7 membered heterocyclic ring having one or more O, S and/or N);

[00181] each m is independently 0, 1 or 2.


[00182] In one embodiment, the base is



[00183] In one embodiment, the base is



[00184] In another embodiment, R^1 is



XXX

XXXI

XXXII

[00185] wherein each R^6 and R^7 is as defined in Formulae XX, XXI or XXII above;

[00186] wherein each R^8 and R^{11} is independently hydrogen, an optionally substituted alkyl (including lower alkyl), CH_3 , CH_2CN , CH_2N_3 , CH_2NH_2 , CH_2NHCH_3 , $CH_2N(CH_3)_2$, CH_2OH , halogenated alkyl (including halogenated lower alkyl), CF_3 , $C(Y^3)_3$, 2-Br-ethyl, CH_2F , CH_2Cl , CH_2CF_3 , CF_2CF_3 , $C(Y^3)_2C(Y^3)_3$, optionally substituted alkenyl, haloalkenyl, Br-vinyl, optionally substituted alkynyl, haloalkynyl, $-CH_2C(O)OH$, $-CH_2C(O)OR^4$, $-CH_2C(O)O(\text{lower alkyl})$, $-CH_2C(O)NH_2$, $-CH_2C(O)NHR^4$, $-CH_2C(O)NH(\text{lower alkyl})$, $-CH_2C(O)N(R^4)_2$, $-CH_2C(O)N(\text{lower alkyl})_2$, $-(CH_2)_mC(O)OH$, $-(CH_2)_mC(O)OR^4$, $-(CH_2)_mC(O)O(\text{lower alkyl})$, $-(CH_2)_mC(O)NH_2$, $-(CH_2)_mC(O)NHR^4$, $-(CH_2)_mC(O)NH(\text{lower alkyl})$, $-(CH_2)_mC(O)N(R^4)_2$, $-(CH_2)_mC(O)N(\text{lower alkyl})_2$, $-C(O)OH$, $-C(O)OR^4$, $-C(O)O(\text{lower alkyl})$, $-C(O)NH_2$, $-C(O)NHR^4$, $-C(O)NH(\text{lower alkyl})$, $-C(O)N(R^4)_2$, $-C(O)N(\text{lower alkyl})_2$, cyano, azido, NH-acyl or $N(\text{acyl})_2$;


[00187] each R^9 and R^{10} are independently hydrogen, OH, OR^2 , optionally substituted alkyl (including lower alkyl), CH_3 , CH_2CN , CH_2N_3 , CH_2NH_2 , CH_2NHCH_3 , $CH_2N(CH_3)_2$, CH_2OH , halogenated alkyl (including halogenated lower alkyl), CF_3 , $C(Y^3)_3$, 2-Br-ethyl, CH_2F , CH_2Cl , CH_2CF_3 , CF_2CF_3 , $C(Y^3)_2C(Y^3)_3$, optionally substituted alkenyl, haloalkenyl, Br-vinyl, optionally substituted alkynyl, haloalkynyl, optionally substituted carbocycle (for example, a 3-7 membered carbocyclic ring), optionally substituted heterocycle (for example, a 3-7 membered heterocyclic ring having one or more O, S and/or N), optionally substituted heteroaryl (for example, a 3-7 membered heteroaromatic ring having one or more O, S and/or N), $-CH_2C(O)OH$, $-CH_2C(O)OR^4$, $-CH_2C(O)O(\text{lower alkyl})$, $-CH_2C(O)SH$, $-CH_2C(O)SR^4$, $-CH_2C(O)S(\text{lower alkyl})$, $-CH_2C(O)NH_2$, $-CH_2C(O)NHR^4$, $-CH_2C(O)NH(\text{lower alkyl})$, $-CH_2C(O)N(R^4)_2$, $-CH_2C(O)N(\text{lower alkyl})_2$, $-(CH_2)_mC(O)OH$, $-(CH_2)_mC(O)OR^4$, $-(CH_2)_mC(O)O(\text{lower alkyl})$, $-(CH_2)_mC(O)SH$, $-(CH_2)_mC(O)SR^4$, $-(CH_2)_mC(O)S(\text{lower alkyl})$, $-(CH_2)_mC(O)NH_2$, $-(CH_2)_mC(O)NHR^4$, $-(CH_2)_mC(O)NH(\text{lower alkyl})$, $-(CH_2)_mC(O)N(R^4)_2$, $-(CH_2)_mC(O)N(\text{lower alkyl})_2$, $-C(O)OH$, $-C(O)OR^4$, $-C(O)O(\text{lower alkyl})$, $-C(O)SH$, $-C(O)SR^4$, $-C(O)S(\text{lower alkyl})$, $-C(O)NH_2$, $-C(O)NHR^4$, $-C(O)NH(\text{lower alkyl})$, $-C(O)N(R^4)_2$, $-C(O)N(\text{lower alkyl})_2$, $-O(\text{acyl})$, $-O(\text{lower acyl})$, $-O(R^4)$, $-O(\text{alkyl})$, $-O(\text{lower alkyl})$, $-O(\text{alkenyl})$, $-O(\text{alkynyl})$, $-O(\text{arylalkyl})$, $-O(\text{cycloalkyl})$, $-S(\text{acyl})$, $-S(\text{lower acyl})$, $-S(R^4)$, $-S(\text{lower alkyl})$, $-S(\text{alkenyl})$, $-S(\text{alkynyl})$, $-S(\text{arylalkyl})$, $-S(\text{cycloalkyl})$, NO_2 , NH_2 , $-NH(\text{lower alkyl})$, $-NHR^4$, $-NR^4R^5$, $-NH(\text{acyl})$, $-N(\text{lower alkyl})_2$, $-NH(\text{alkenyl})$, $-NH(\text{alkynyl})$, $-NH(\text{arylalkyl})$, $-NH(\text{cycloalkyl})$, $-N(\text{acyl})_2$, azido, cyano, SCN, OCN, NCO or halo (fluoro, chloro, bromo, iodo);

[00188] each m is independently 0, 1 or 2;

[00189] alternatively, R^6 and R^{10} , R^7 and R^9 , R^8 and R^7 or R^9 and R^{11} can come together to form a bridged compound selected from the group consisting of optionally substituted carbocycle (for example, a 3-7 membered carbocyclic ring) or optionally substituted heterocycle (for example, a 3-7 membered heterocyclic ring having one or more O, S and/or N); or

[00190] alternatively, R^6 and R^7 or R^9 and R^{10} can come together to form a spiro compound selected from the group consisting of optionally substituted carbocycle (for example, a 3-7 membered carbocyclic ring) or optionally substituted heterocycle (for example, a 3-7 membered heterocyclic ring having one or more O, S and/or N).

[00191] In another embodiment, R¹ is:



wherein Base* is a purine or pyrimidine base as defined herein;

[00192] each R¹² is independently a substituted alkyl (including lower alkyl), CH₂CN, CH₂N₃, CH₂NH₂, CH₂NHCH₃, CH₂N(CH₃)₂, CH₂OH, halogenated alkyl (including halogenated lower alkyl), CF₃, C(Y³)₃, 2-Br-ethyl, CH₂F, CH₂Cl, CH₂CF₃, CF₂CF₃, C(Y³)₂C(Y³)₃, substituted alkenyl, haloalkenyl (but not Br-vinyl), substituted alkynyl, haloalkynyl, -CH₂C(O)OH, -CH₂C(O)OR⁴, -CH₂C(O)O(lower alkyl), -CH₂C(O)NH₂, -CH₂C(O)NHR⁴, -CH₂C(O)NH(lower alkyl), -CH₂C(O)N(R⁴)₂, -CH₂C(O)N(lower alkyl)₂, -(CH₂)_mC(O)OH, -(CH₂)_mC(O)OR⁴, -(CH₂)_mC(O)O(lower alkyl), -(CH₂)_mC(O)NH₂, -(CH₂)_mC(O)NHR⁴, -(CH₂)_mC(O)NH(lower alkyl), -(CH₂)_mC(O)N(R⁴)₂, -(CH₂)_mC(O)N(lower alkyl)₂, -C(O)OH, -C(O)OR⁴, -C(O)NH₂, -C(O)NHR⁴, -C(O)NH(lower alkyl), -C(O)N(R⁴)₂, -C(O)N(lower alkyl)₂;

[00193] each R¹³ is independently substituted alkyl (including lower alkyl), CH₂CN, CH₂N₃, CH₂NH₂, CH₂NHCH₃, CH₂N(CH₃)₂, CH₂OH, halogenated alkyl (including halogenated lower alkyl), CF₃, C(Y³)₃, 2-Br-ethyl, CH₂F, CH₂Cl, CH₂CF₃, CF₂CF₃, C(Y³)₂C(Y³)₃, substituted alkenyl, haloalkenyl (but not Br-vinyl), substituted alkynyl, haloalkynyl, optionally substituted carbocycle (for example, a 3-7 membered carbocyclic ring), optionally substituted heterocycle (for example, a 3-7 membered heterocyclic ring having one or more O, S and/or N), optionally substituted heteroaryl (for example, a 3-7 membered heteroaromatic ring having one or more O, S and/or N), -CH₂C(O)OH, -CH₂C(O)OR⁴, -CH₂C(O)O(lower alkyl), -CH₂C(O)SH, -CH₂C(O)SR⁴, -CH₂C(O)S(lower alkyl), -CH₂C(O)NH₂, -CH₂C(O)NHR⁴, -CH₂C(O)NH(lower alkyl), -CH₂C(O)N(R⁴)₂, -CH₂C(O)N(lower alkyl)₂, -(CH₂)_mC(O)OH, -(CH₂)_mC(O)OR⁴, -(CH₂)_mC(O)O(lower alkyl), -(CH₂)_mC(O)SH, -(CH₂)_mC(O)SR⁴, -(CH₂)_mC(O)S(lower alkyl), -(CH₂)_mC(O)NH₂, -(CH₂)_mC(O)NHR⁴, -(CH₂)_mC(O)NH(lower alkyl), -(CH₂)_mC(O)N(R⁴)₂, -(CH₂)_mC(O)N(lower alkyl)₂;


-C(O)OH, -C(O)OR⁴, -C(O)SH, -C(O)SR⁴, -C(O)S(lower alkyl), -C(O)NH₂,
 -C(O)NHR⁴, -C(O)NH(lower alkyl), -C(O)N(R⁴)₂, -C(O)N(lower alkyl)₂, -O(R⁴),
 -O(alkynyl), -O(arylalkyl), -O(cycloalkyl), -S(acyl), -S(lower acyl), -S(R⁴), -S(lower
 alkyl), -S(alkenyl), -S(alkynyl), -S(arylalkyl), -S(cycloalkyl), -NHR⁴, -NR⁴R⁵, -
 NH(alkenyl), -NH(alkynyl), -NH(arylalkyl), -NH(cycloalkyl), SCN, OCN, NCO or
 fluoro;

[00194] alternatively, R¹² and R¹³ can come together to form a spiro compound
 selected from the group consisting of optionally substituted carbocycle (for example,
 a 3-7 membered carbocyclic ring) or optionally substituted heterocycle (for example,
 a 3-7 membered heterocyclic ring having one or more O, S and/or N);

[00195] R² and R³ are according to Formula XII; and

[00196] each m is independently 0, 1 or 2.

[00197] In another embodiment, R is:



wherein Base* is a purine or pyrimidine base as described herein; and


[00198] each R⁸ and R¹¹ is independently hydrogen, an optionally substituted alkyl
 (including lower alkyl), CH₃, CH₂CN, CH₂N₃, CH₂NH₂, CH₂NHCH₃, CH₂N(CH₃)₂,
 CH₂OH, halogenated alkyl (including halogenated lower alkyl), CF₃, C(Y³)₃, 2-Br-
 ethyl, CH₂F, CH₂Cl, CH₂CF₃, CF₂CF₃, C(Y³)₂C(Y³)₃, optionally substituted alkenyl,
 haloalkenyl, Br-vinyl, optionally substituted alkynyl, haloalkynyl, -CH₂C(O)OH,
 -CH₂C(O)OR⁴, -CH₂C(O)O(lower alkyl), -CH₂C(O)NH₂, -CH₂C(O)NHR⁴,
 -CH₂C(O)NH(lower alkyl), -CH₂C(O)N(R⁴)₂, -CH₂C(O)N(lower alkyl)₂,
 -(CH₂)_mC(O)OH, -(CH₂)_mC(O)OR⁴, -(CH₂)_mC(O)O(lower alkyl), -(CH₂)_mC(O)NH₂,
 -(CH₂)_mC(O)NHR⁴, -(CH₂)_mC(O)NH(lower alkyl), -(CH₂)_mC(O)N(R⁴)₂,
 -(CH₂)_mC(O)N(lower alkyl)₂, -C(O)OH, -C(O)OR⁴, -C(O)O(lower alkyl), -C(O)NH₂,
 -C(O)NHR⁴, -C(O)NH(lower alkyl), -C(O)N(R⁴)₂, -C(O)N(lower alkyl)₂, cyano, NH-
 acyl or N(acyl)₂;

[00199] each R^9 and R^{10} are independently hydrogen, OH, OR^2 , optionally substituted alkyl (including lower alkyl), CH_3 , CH_2CN , CH_2N_3 , CH_2NH_2 , CH_2NHCH_3 , $CH_2N(CH_3)_2$, CH_2OH , halogenated alkyl (including halogenated lower alkyl), CF_3 , $C(Y^3)_3$, 2-Br-ethyl, CH_2F , CH_2Cl , CH_2CF_3 , CF_2CF_3 , $C(Y^3)_2C(Y^3)_3$, optionally substituted alkenyl, haloalkenyl, Br-vinyl, optionally substituted alkynyl, haloalkynyl, optionally substituted carbocycle (for example, a 3-7 membered carbocyclic ring), optionally substituted heterocycle (for example, a 3-7 membered heterocyclic ring having one or more O, S and/or N), optionally substituted heteroaryl (for example, a 3-7 membered heteroaromatic ring having one or more O, S and/or N), $-CH_2C(O)OH$, $-CH_2C(O)OR^4$, $-CH_2C(O)O(\text{lower alkyl})$, $-CH_2C(O)SH$, $-CH_2C(O)SR^4$, $-CH_2C(O)S(\text{lower alkyl})$, $-CH_2C(O)NH_2$, $-CH_2C(O)NHR^4$, $-CH_2C(O)NH(\text{lower alkyl})$, $-CH_2C(O)N(R^4)_2$, $-CH_2C(O)N(\text{lower alkyl})_2$, $-(CH_2)_mC(O)OH$, $-(CH_2)_mC(O)OR^4$, $-(CH_2)_mC(O)O(\text{lower alkyl})$, $-(CH_2)_mC(O)SH$, $-(CH_2)_mC(O)SR^4$, $-(CH_2)_mC(O)S(\text{lower alkyl})$, $-(CH_2)_mC(O)NH_2$, $-(CH_2)_mC(O)NHR^4$, $-(CH_2)_mC(O)NH(\text{lower alkyl})$, $-(CH_2)_mC(O)N(R^4)_2$, $-(CH_2)_mC(O)N(\text{lower alkyl})_2$, $-C(O)OH$, $-C(O)OR^4$, $-C(O)O(\text{lower alkyl})$, $-C(O)SH$, $-C(O)SR^4$, $-C(O)S(\text{lower alkyl})$, $-C(O)NH_2$, $-C(O)NHR^4$, $-C(O)NH(\text{lower alkyl})$, $-C(O)N(R^4)_2$, $-C(O)N(\text{lower alkyl})_2$, $-O(\text{acyl})$, $-O(\text{lower acyl})$, $-O(R^4)$, $-O(\text{alkyl})$, $-O(\text{lower alkyl})$, $-O(\text{alkenyl})$, $-O(\text{alkynyl})$, $-O(\text{arylalkyl})$, $-O(\text{cycloalkyl})$, $-S(\text{acyl})$, $-S(\text{lower acyl})$, $-S(R^4)$, $-S(\text{lower alkyl})$, $-S(\text{alkenyl})$, $-S(\text{alkynyl})$, $-S(\text{arylalkyl})$, $-S(\text{cycloalkyl})$, NO_2 , NH_2 , $-NH(\text{lower alkyl})$, $-NHR^4$, $-NR^4R^5$, $-NH(\text{acyl})$, $-N(\text{lower alkyl})_2$, $-NH(\text{alkenyl})$, $-NH(\text{alkynyl})$, $-NH(\text{arylalkyl})$, $-NH(\text{cycloalkyl})$, $-N(\text{acyl})_2$, azido, cyano, SCN, OCN, NCO or halo (fluoro, chloro, bromo, iodo);


[00200] each R^{12} is independently a substituted alkyl (including lower alkyl), CH_2CN , CH_2N_3 , CH_2NH_2 , CH_2NHCH_3 , $CH_2N(CH_3)_2$, CH_2OH , halogenated alkyl (including halogenated lower alkyl), CF_3 , $C(Y^3)_3$, 2-Br-ethyl, CH_2F , CH_2Cl , CH_2CF_3 , CF_2CF_3 , $C(Y^3)_2C(Y^3)_3$, substituted alkenyl, haloalkenyl (but not Br-vinyl), substituted alkynyl, haloalkynyl, $-CH_2C(O)OH$, $-CH_2C(O)OR^4$, $-CH_2C(O)O(\text{lower alkyl})$, $-CH_2C(O)NH_2$, $-CH_2C(O)NHR^4$, $-CH_2C(O)NH(\text{lower alkyl})$, $-CH_2C(O)N(R^4)_2$, $-CH_2C(O)N(\text{lower alkyl})_2$, $-(CH_2)_mC(O)OH$, $-(CH_2)_mC(O)OR^4$, $-(CH_2)_mC(O)O(\text{lower alkyl})$, $-(CH_2)_mC(O)NH_2$, $-(CH_2)_mC(O)NHR^4$, $-(CH_2)_mC(O)NH(\text{lower alkyl})$, $-(CH_2)_mC(O)N(R^4)_2$, $-(CH_2)_mC(O)N(\text{lower alkyl})_2$, $-C(O)OH$, $-C(O)OR^4$, $-C(O)NH_2$, $-C(O)NHR^4$, $-C(O)NH(\text{lower alkyl})$, $-C(O)N(R^4)_2$, $-C(O)N(\text{lower alkyl})_2$;

[00202] alternatively, R^8 and R^{13} , R^9 and R^{13} , R^9 and R^{11} or R^{10} and R^{12} can come together to form a bridged compound selected from the group consisting of optionally substituted carbocycle (for example, a 3-7 membered carbocyclic ring) or optionally substituted heterocycle (for example, a 3-7 membered heterocyclic ring having one or more O, S and/or N); or

[00204] In one aspect, R^1 is:



79



[00216] G and E independently are selected from the group consisting of CH_3 , CH_2OH , CH_2F , CH_2N_3 , CH_2CN , $(\text{CH}_2)_m\text{COOH}$, $(\text{CH}_2)_m\text{COOR}$, $(\text{CH}_2)_m\text{CONH}_2$, $(\text{CH}_2)_m\text{CONR}_2$, $(\text{CH}_2)_m\text{CONHR}$, N_3 and N-acyl;


[00217] m is 0 or 1;

[00218] R is H, alkyl or acyl; and

[00219] Base is as defined for Formula (XIII).



[00220] In one embodiment, at most one of G and E can further be hydrogen.

[00221] In another embodiment, R^1 is:



[00222] wherein M is selected from the group consisting of O, S, SO, and SO_2 ; and Base is as defined for Formula (XIII).

[00223] In certain embodiments, R^1 is:



[00224] wherein A is selected from the group consisting of optionally substituted lower alkyl, cycloalkyl, alkenyl, alkynyl, CH_2OH , CH_2NH_2 , CH_2NHCH_3 ,


$\text{CH}_2\text{N}(\text{CH}_3)_2$, CH_2F , CH_2Cl , CH_2N_3 , CH_2CN , CH_2CF_3 , CF_3 , CF_2CF_3 , $\text{CH}_2\text{CO}_2\text{R}$, $(\text{CH}_2)_m\text{COOH}$, $(\text{CH}_2)_m\text{COOR}$, $(\text{CH}_2)_m\text{CONH}_2$, $(\text{CH}_2)_m\text{CONR}_2$, and $(\text{CH}_2)_m\text{CONHR}$;

[00225] Y is selected from the group consisting of H, optionally substituted lower alkyl, cycloalkyl, alkenyl, alkynyl, CH_2OH , CH_2NH_2 , CH_2NHCH_3 , $\text{CH}_2\text{N}(\text{CH}_3)_2$, CH_2F , CH_2Cl , CH_2N_3 , CH_2CN , CH_2CF_3 , CF_3 , CF_2CF_3 , $\text{CH}_2\text{CO}_2\text{R}$, $(\text{CH}_2)_m\text{COOH}$, $(\text{CH}_2)_m\text{COOR}$, $(\text{CH}_2)_m\text{CONH}_2$, $(\text{CH}_2)_m\text{CONR}_2$, and $(\text{CH}_2)_m\text{CONHR}$;

[00226] X is selected from the group consisting of H, -OH, optionally substituted alkyl, cycloalkyl, alkenyl, alkynyl, -O-alkyl, -O-alkenyl, -O-alkynyl, -O-aryl, -O-arylalkyl, -O-cycloalkyl-, O-acyl, F, Cl, Br, I, CN, NC, SCN, OCN, NCO, NO_2 , NH_2 , N_3 , NH-acyl, NH-alkyl, N-dialkyl, NH-alkenyl, NH-alkynyl, NH-aryl, NH-arylalkyl, NH-cycloalkyl, SH, S-alkyl, S-alkenyl, S-alkynyl, S-aryl, S-arylalkyl, S-acyl, S-cycloalkyl, CO_2 -alkyl, CONH-alkyl, CON-dialkyl, CONH-alkenyl, CONH-alkynyl, CONH-arylalkyl, CONH-cycloalkyl, CH_2OH , CH_2NH_2 , CH_2NHCH_3 , $\text{CH}_2\text{N}(\text{CH}_3)_2$, CH_2F , CH_2Cl , CH_2N_3 , CH_2CN , CH_2CF_3 , CF_3 , CF_2CF_3 , $\text{CH}_2\text{CO}_2\text{R}$, $(\text{CH}_2)_m\text{COOH}$, $(\text{CH}_2)_m\text{COOR}$, $(\text{CH}_2)_m\text{CONH}_2$, $(\text{CH}_2)_m\text{CONR}_2$, $(\text{CH}_2)_m\text{CONHR}$, an optionally substituted 3-7 membered carbocyclic, and an optionally substituted 3-7 membered heterocyclic ring having O, S and/or N independently as a heteroatom taken alone or in combination;

[00227] m is 0 or 1;

[00228] R is H, alkyl or acyl; and Base is a non-natural base selected from the group of:



[00229] wherein each R', R'', R''' and R'''' is independently selected from the group consisting of H, OH, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, cycloalkyl, Br-vinyl, -O-alkyl, O-alkenyl, O-alkynyl, O-aryl, O-arylalkyl, -O-acyl, O-cycloalkyl, NH₂, NH-alkyl, N-dialkyl, NH-acyl, N-aryl, N-arylalkyl, NH-cycloalkyl, SH, S-alkyl, S-acyl, S-aryl, S-cycloalkyl, S-arylalkyl, F, Cl, Br, I, CN, COOH, CONH₂, CO₂-alkyl, CONH-alkyl, CON-dialkyl, OH, CF₃, CH₂OH, (CH₂)_mOH, (CH₂)_mNH₂, (CH₂)_mCOOH, (CH₂)_mCN, (CH₂)_mNO₂ and (CH₂)_mCONH₂;

[00230] m is 0 or 1;

[00231] each W is independently C-R'' or N;

[00232] T and V independently are CH or N;

[00233] Q is CH, -CCl, -CBr, -CF, -Cl, -CCN, -C-COOH, -C-CONH₂, or N;


[00234] Q₁ and Q₂ independently are N or C-R''''; and

[00235] Q₃, Q₄, Q₅ and Q₆ independently are N or CH;

[00236] with the proviso that in bases (g) and (i), R', R'''' are not H, OH, or NH₂; and Q, T, V, Q₂, Q₅ and Q₆ are not N.

[00237] In one embodiment, R¹ is a 2'-(alkyl or aryl) ester or 3'-(alkyl or aryl) ester of 1', 2', 3' or 4' C-branched-β-D or β-L nucleoside with any natural or non-natural purine or pyrimidine base. In one embodiment, R¹ is a 2' or 3'-(D or L)-amino acid ester of 1', 2', 3' or 4' C-branched -β-D or β-L nucleoside, wherein the amino acid is a natural or synthetic amino acid. In another embodiment, R¹ is a 3'-D or L-amino acid ester of 1', 2', 3' or 4' C-branched-β-D or β-L nucleoside, wherein the amino acid is a natural or synthetic amino acid. In one embodiment, the amino acid is an L-amino acid.

[00238] In one embodiment, the amino acid residue is of the formula




[00239] wherein R¹¹ is the side chain of an amino acid and wherein, R¹¹ can optionally be attached to R¹³ to form a ring structure; or alternatively, R¹¹ is an alkyl, aryl, heteroaryl or heterocyclic moiety;

[00240] R¹² is hydrogen, alkyl (including lower alkyl) or aryl; and

[00241] R^{13} and R^{14} are independently hydrogen, acyl (including an acyl derivative attached to R^{11}) or alkyl (including but not limited to methyl, ethyl, propyl, and cyclopropyl).


[00242] In another embodiment, at least one of R^2 and R^3 is an amino acid residue. In one embodiment, at least one of R^2 and R^3 is L-valinyl.

[00243] In one embodiment, R^1 is:




wherein R^6 , R^7 , R^8 , R^9 , R^{10} and Base^* are as defined in Formula XXX, XXXI, XL, XLI or XLII.

[00244] In one embodiment, R^1 is:



wherein R^2 , R^3 , R^6 and Base^* are as defined in Formula XXX, XXXI, XL, XLI or XLII.


[00245] In one embodiment, R^1 is:



wherein X^1 and X^2 are each independently hydrogen, alkyl, halo or amino; Y is hydrogen, amino, aminoalkyl, aminocycloalkyl, alkyl, cycloalkyl, hydroxy, alkoxy,


cycloalkoxy, SH or thioalkyl; X is O or S; and wherein R^6 , R^7 , R^8 , R^9 are as defined in Formula XXX, XXXI, XL, XLI or XLII.

[00246] In one embodiment, R^1 is:




wherein X^1 is hydrogen, alkyl, halo or amino; Y is hydrogen, amino, aminoalkyl, aminocycloalkyl, alkyl, cycloalkyl, hydroxy, alkoxy, cycloalkoxy, SH or thioalkyl; X is O or S; and wherein R^6 , R^7 , R^8 , R^9 are as defined in Formula XXX, XXXI, XL, XLI or XLII.

[00247] In one embodiment, R^1 is:




wherein R^2 , R^3 , and $Base^*$ are as defined in Formula XIII, XXX, XXXI, XL, XLI or XLII.

[00248] In one embodiment, R^1 is:




wherein R², R³, Y¹, Y³, X¹, and X² are as defined in Formula XIII.

[00249] In one embodiment, R¹ is:




wherein R², R³, Y¹, Y³, X¹, and X² are as defined in Formula XIII.

[00250] In one embodiment, R¹ is:




wherein R², R³, R⁶, Y, and X¹ are as defined in Formula XIII, XX, XXI or XXII.

[00251] In one embodiment, R¹ is:



wherein R^2 , R^3 , R^6 , R^7 , X and $Base^*$ are as defined in Formula XIII, XX, XXI, XXII, XL, XLI or XLII.


[00252] In one embodiment, R^1 is



wherein R^8 is alkyl, alkenyl or alkynyl; R^7 is OR^{7a} ;

R^9 is OR^{7a} ;

R^{7a} is H or




R^m is a side chain of any natural or non-natural amino acid; and

R^p is hydrogen, hydroxy, alkyl or alkoxy; and


$Base^*$ is as defined in Formula XL, XLI or XLII

In one embodiment, R^8 is methyl, ethyl, vinyl or ethynyl; R^7 is hydroxy or fluoro; R^9 is hydroxy and other variables are as described herein.


[00253] In one embodiment, R^1 is



[00254] In one embodiment, R^8 is methyl or ethyl. In one embodiment, R^{7a} is H or




[00255] In one embodiment, the phosphoramidate compound provided herein is:




or a pharmaceutically acceptable salt, solvate or hydrate thereof.


[00256] In one embodiment, the phosphoramidate compound provided herein is:




[00257] In one embodiment, the phosphoramidate compound provided herein is:




[00258] In one embodiment, the phosphoramidate compound provided herein is:




[00259] In one embodiment, the phosphoramidate compound provided herein is:




[00260] In one embodiment, the phosphoramidate compound provided herein is:




[00261] In one embodiment, the phosphoramidate compound provided herein is:




[00262] In one embodiment, the phosphoramidate compound provided herein is:




[00263] In one embodiment, the phosphoramidate compound provided herein is:




[00264] In one embodiment, the phosphoramidate compound provided herein is:




[00265] In one embodiment, the phosphoramidate compound provided herein is:




[00266] In one embodiment, the phosphoramidate compound provided herein is:




[00267] In one embodiment, the phosphoramidate compound provided herein is:




[00268] In one embodiment, the phosphoramidate compound provided herein is:



[00269] In one embodiment, the phosphoramidate compound provided herein is:



[00270] In one embodiment, the phosphonoamidate compound provided herein is a phosphonoamidate form of PMPA or PMEAs such as:



Optically Active Compounds

[00271] It is appreciated that compounds provided herein have several chiral centers and may exist in and be isolated in optically active and racemic forms. Some compounds may exhibit polymorphism. It is to be understood that any racemic, optically-active, diastereomeric, polymorphic, or stereoisomeric form, or mixtures thereof, of a compound provided herein, which possess the useful properties described herein is within the scope of the invention. It being well known in the art how to prepare optically active forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase).

[00272] In particular, since the 1' and 4' carbons of a nucleoside are chiral, their nonhydrogen substituents (the base and the CHOR groups, respectively) can be either cis (on the same side) or trans (on opposite sides) with respect to the sugar ring system. The four optical isomers therefore are represented by the following configurations (when orienting the sugar moiety in a horizontal plane such that the oxygen atom is in the back): cis (with both groups "up", which corresponds to the configuration of naturally occurring β -D nucleosides), cis (with both groups "down", which is a nonnaturally occurring β -L configuration), trans (with the C2' substituent "up" and the C4' substituent "down"), and trans (with the C2' substituent "down" and the C4' substituent "up"). The "D-nucleosides" are cis nucleosides in a natural configuration and the "L-nucleosides" are cis nucleosides in the non-naturally occurring configuration.

[00273] Likewise, most amino acids are chiral (designated as L or D, wherein the L enantiomer is the naturally occurring configuration) and can exist as separate enantiomers.

[00274] Examples of methods to obtain optically active materials are known in the art, and include at least the following.

- i) physical separation of crystals - a technique whereby macroscopic crystals of the individual enantiomers are manually separated. This technique can be used if crystals of the separate enantiomers exist, *i.e.*, the material is a conglomerate, and the crystals are visually distinct;
- ii) simultaneous crystallization - a technique whereby the individual enantiomers are separately crystallized from a solution of the racemate, possible only if the latter is a conglomerate in the solid state;
- iii) enzymatic resolutions - a technique whereby partial or complete separation of a racemate by virtue of differing rates of reaction for the enantiomers with an enzyme;
- iv) enzymatic asymmetric synthesis - a synthetic technique whereby at least one step of the synthesis uses an enzymatic reaction to obtain an enantiomerically pure or enriched synthetic precursor of the desired enantiomer;
- v) chemical asymmetric synthesis - a synthetic technique whereby the desired enantiomer is synthesized from an achiral precursor under conditions that produce asymmetry (*i.e.*, chirality) in the product, which may be achieved using chiral catalysts or chiral auxiliaries;
- vi) diastereomer separations - a technique whereby a racemic compound is reacted with an enantiomerically pure reagent (the chiral auxiliary) that converts the individual enantiomers to diastereomers. The resulting diastereomers are then separated by chromatography or crystallization by virtue of their now more distinct structural differences and the chiral auxiliary later removed to obtain the desired enantiomer;
- vii) first- and second-order asymmetric transformations - a technique whereby diastereomers from the racemate equilibrate to yield a preponderance in solution of the diastereomer from

the desired enantiomer or where preferential crystallization of the diastereomer from the desired enantiomer perturbs the equilibrium such that eventually in principle all the material is converted to the crystalline diastereomer from the desired enantiomer. The desired enantiomer is then released from the diastereomer;

- viii) kinetic resolutions - this technique refers to the achievement of partial or complete resolution of a racemate (or of a further resolution of a partially resolved compound) by virtue of unequal reaction rates of the enantiomers with a chiral, non-racemic reagent or catalyst under kinetic conditions;
- ix) enantiospecific synthesis from non-racemic precursors - a synthetic technique whereby the desired enantiomer is obtained from non-chiral starting materials and where the stereochemical integrity is not or is only minimally compromised over the course of the synthesis;
- x) chiral liquid chromatography - a technique whereby the enantiomers of a racemate are separated in a liquid mobile phase by virtue of their differing interactions with a stationary phase. The stationary phase can be made of chiral material or the mobile phase can contain an additional chiral material to provoke the differing interactions;
- xi) chiral gas chromatography - a technique whereby the racemate is volatilized and enantiomers are separated by virtue of their differing interactions in the gaseous mobile phase with a column containing a fixed non-racemic chiral adsorbent phase;
- xii) extraction with chiral solvents - a technique whereby the enantiomers are separated by virtue of preferential dissolution of one enantiomer into a particular chiral solvent;
- xiii) transport across chiral membranes - a technique whereby a racemate is placed in contact with a thin membrane barrier. The barrier typically separates two miscible fluids, one containing the racemate, and a driving force such as concentration or pressure differential causes preferential

transport across the membrane barrier. Separation occurs as a result of the non-racemic chiral nature of the membrane which allows only one enantiomer of the racemate to pass through.


[00275] In some embodiments, compositions of phosphonoamidate or phosphoramidate compounds are provided that are substantially free of a designated enantiomer of that nucleoside. In a preferred embodiment, in the methods and compounds of this invention, the compounds are substantially free of enantiomers. In some embodiments, the composition includes that includes a compound that is at least 85, 90%, 95%, 98%, 99% to 100% by weight, of the compound, the remainder comprising other chemical species or enantiomers.

Preparation of Compounds

[00276] The compounds provided herein can be prepared, isolated or obtained by any method apparent to those of skill in the art. Exemplary methods of preparation are described in detail in the examples below.

[00277] In certain embodiments, compounds provided herein can be prepared by coupling alcohols and H-phosphonate monoesters as illustrated in the reaction scheme below:


Scheme A



any reactive function on R^y , R^7 , R^8 , R^9 , R^{10} or on the base may be protected during the coupling reaction. A variety of coupling agents known to one of skill in the art can be used. Exemplary coupling agents for use in the reaction include, but are not limited to HOBt (N-Hydroxybenzotriazole), HBTU (2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate), DCC (N,N'-dicyclohexylcarbodiimide), BOP (Benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphoniumhexafluorophosphate), PyBOP (1H-benzotriazol-1-

xyloxytripyrrolidinophosphonium hexafluorophosphate) and others known to one of skill in the art.


[00278] A general scheme for the synthesis of hydroxytBuSATE N-benzylphosphoramidate nucleoside derivatives represented by **B** is provided in Schemes B1-B3 below.




B

[00279] where R = H, Tr, MMTr or DMTr in case of reactive amine; R¹, R², R⁴, R⁶ = H, alkyl or halo and R³ / R⁵ are both H or isopropylidene.


[00280] **Scheme B1: Synthesis of the H-phosphonate monoester reagent**



[00281] **Scheme B2: Synthesis of the protected nucleosides (R= DMTr and/or R³/R⁵ = isopropylidene)**



[00282] **Scheme B3: Coupling of (non)protected nucleosides with reagent 5, oxidative amination and deprotection step**



[00283] In addition, certain nucleosides and analogs thereof and prodrugs thereof can be prepared according to methods described in U.S. Patent Nos. 6,812,219; 7,105,493; 7,101,861; 6,914,054; 6,555,676; 7,202,224; 7,105,499; 6,777,395; 6,914,054; 7,192,936; US publication Nos. 2005203243; 2007087960; 2007060541; 2007060505; 2007060504; 2007060503; 2007060498; 2007042991; 2007042990; 2007042940; 2007042939 and 2007037735; International Publication Nos. WO 04/003000; WO 04/022999; WO 04/002422; WO 01/90121 and WO 01/92282. Other patents/patent applications disclosing nucleoside analogs to treat hepatitis C virus that can be derivatized as described herein include: PCT/CA00/01316 (WO 01/32153; filed November 3, 2000) and PCT/CA01/00197 (WO 01/60315; filed February 19, 2001) filed by BioChem Pharma, Inc. (now Shire Biochem, Inc.); PCT/US02/01531 (WO 02/057425; filed January 18, 2002); PCT/US02/03086 (WO 02/057287; filed January 18, 2002); US 7,202,224; 7,125,855; 7,105,499 and 6,777,395 by Merck & Co., Inc.; PCT/EP01/09633 (WO 02/18404; published August 21, 2001); US 2006/0040890; 2005/0038240; 2004/0121980; 6,846,810; 6,784,166 and 6,660,721 by Roche; PCT Publication Nos. WO 01/79246 (filed April 13, 2001), WO 02/32920 (filed October 18, 2001) and WO 02/48165; US 2005/0009737 and US 2005/0009737; 7,094,770 and 6,927,291 by Pharmasset, Ltd. Contents of these references are hereby incorporated by reference in their entireties.

Assay Methods

[00284] Compounds can be assayed for HBV activity according to any assay known to those of skill in the art. Compounds can be assayed for HCV activity according to any assay known to those of skill in the art.

[00285] Further, compounds can be assayed for accumulation in liver cells of a subject according to any assay known to those of skill in the art. In certain embodiments, a compound can be administered to the subject, and a liver cell of the

subject can be assayed for the compound or a derivative thereof, *e.g.* a nucleoside, nucleoside phosphate or nucleoside triphosphate derivative thereof.

[00286] In one embodiment, a phosphoramidate or phosphonoamidate nucleoside compound is administered to cells, such as liver cells, *in vivo* or *in vitro*, and the nucleoside triphosphate levels delivered intracellularly are measured, to indicate delivery of the compound and triphosphorylation in the cell. The levels of intracellular nucleoside triphosphate can be measured using analytical techniques known in the art. Methods of detecting ddATP are described herein below by way of example, but other nucleoside triphosphates can be readily detected using the appropriate controls, calibration samples and assay techniques.

[00287] In one embodiment, ddATP concentrations are measured in a sample by comparison to calibration standards made from control samples. The ddATP concentrations in a sample can be measured using an analytical method such as HPLC LC MS. In one embodiment, a test sample is compared to a calibration curve created with known concentrations of ddATP to thereby obtain the concentration of that sample.

[00288] In one embodiment, the samples are manipulated to remove impurities such as salts (Na^+ , K^+ , etc.) before analysis. In one embodiment, the lower limit of quantitation is about ~ 0.2 pmol / mL for hepatocyte cellular extracts particularly where reduced salt is present.

[00289] In one embodiment, the method allows successfully measuring triphosphate nucleotides formed at levels of 1 – 10,000 pmol per million cells in *e.g.* cultured hepatocytes and HepG2 cells.

Methods of Use

[00290] The phosphoramidate and phosphonoamidate compounds of a variety of therapeutic agents can be formed using methods available in the art and those disclosed herein. Such compounds can be used in some embodiments to enhance delivery of the drug to the liver.

[00291] In one embodiment, the compound comprises a *S*-acyl-2-thioethyl phosphoramidate or *S*-acyl-2-thioethyl phosphonoamidate, *e.g.*, a *S*-pivaloyl-2-thioethyl phosphoramidate or *S*-hydroxypivaloyl-2-thioethyl phosphonoamidate derivative. Therapeutic agents that can be derivatized to phosphoramidate or

phosphonoamidate compound form include any anti-viral agent that includes, or has been derivatized to include a reactive group for attachment of the phosphoramidate or phosphonoamidate moiety, including but not limited to nucleosides and nucleoside analogues including acyclic nucleosides.

[00292] Advantageously, such phosphoramidate and phosphonamidate compounds advantageously can have enhanced delivery to the liver. In some embodiments, the compounds permit delivery of an active 5'-monophosphate of a nucleoside to the liver, which can enhance the formation of active triphosphorylated compound.

[00293] In one embodiment, provided herein are methods for the treatment and/or prophylaxis of a host infected with *Flaviviridae* that includes the administration of an effective amount of a compounds provided herein, or a pharmaceutically acceptable salt thereof. In one embodiment, provided herein are methods for treating an HCV infection in a subject. In certain embodiments, the methods encompass the step of administering to the subject in need thereof an amount of a compound effective for the treatment or prevention of an HCV infection in combination with a second agent effective for the treatment or prevention of the infection. The compound can be any compound as described herein, and the second agent can be any second agent described in the art or herein. In certain embodiments, the compound is in the form of a pharmaceutical composition or dosage form, as described in the sections above.

[00294] *Flaviviridae* that can be treated are discussed generally in *Fields Virology*, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, Chapter 31, 1996. In a particular embodiment of the invention, the *Flaviviridae* is HCV. In an alternate embodiment of the invention, the *Flaviviridae* is a flavivirus or pestivirus. Specific flaviviruses include, without limitation:

Absettarov, Alfuy, Apoi, Aroa, Bagaza, Banzi, Bouboui, Bussuquara, Cacipacore, Carey Island, Dakar bat, Dengue 1, Dengue 2, Dengue 3, Dengue 4, Edge Hill, Entebbe bat, Gadgets Gully, Hanzalova, Hypr, Ilheus, Israel turkey meningoencephalitis, Japanese encephalitis, Jugra, Jutiapa, Kadam, Karshi, Kedougou, Kokobera, Koutango, Kumlinge, Kunjin, Kyasanur Forest disease, Langat, Louping ill, Meaban, Modoc, Montana myotis leukoencephalitis, Murray valley encephalitis, Naranjal, Negishi, Ntaya, Omsk hemorrhagic fever, Phnom-Penh bat, Powassan, Rio Bravo, Rocio, Royal Farm, Russian spring-summer encephalitis, Saboya, St. Louis encephalitis, Sal Vieja, San Perlita, Saumarez Reef, Sepik, Sokuluk,

Spondweni, Stratford, Tembusu, Tyuleniy, Uganda S, Usutu, Wesselsbron, West Nile, Yaounde, Yellow fever, and Zika.

[00295] Pestiviruses that can be treated are discussed generally in *Fields Virology*, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, Chapter 33, 1996. Specific pestiviruses include, without limitation: bovine viral diarrhea virus ("BVDV"), classical swine fever virus ("CSFV," also called hog cholera virus), and border disease virus ("BDV").

[00296] In certain embodiments, provided herein are methods for the treatment and/or prophylaxis of hepatitis B infections that includes administering an effective amount of a compound as described herein, *e.g.* of Formula I, IIa or IIb, its pharmaceutically acceptable salt or composition. In another embodiment, provided herein are methods of treatment and/prophylaxis of conditions related to hepatitis B infections, such as anti-HBV antibody positive and HBV-positive conditions, chronic liver inflammation caused by HBV, cirrhosis, acute hepatitis, fulminant hepatitis, chronic persistent hepatitis, and fatigue. In certain embodiments, provided herein are prophylactic methods to prevent or retard the progression of clinical illness in individuals who are anti-HBV antibody or HBV-antigen positive or who have been exposed to HBV.

[00297] In certain embodiments, the subject can be any subject infected with, or at risk for infection with, HCV and/or HBV. Infection or risk for infection can be determined according to any technique deemed suitable by the practitioner of skill in the art. In one embodiment, subjects are humans infected with HCV and/or HBV.

[00298] In certain embodiments, the subject has never received therapy or prophylaxis for an HCV and/or HBV infection. In further embodiments, the subject has previously received therapy or prophylaxis for an HCV and/or HBV infection. For instance, in certain embodiments, the subject has not responded to an HCV and/or HBV therapy. For example, under current interferon therapy, up to 50% or more HCV subjects do not respond to therapy. In certain embodiments, the subject can be a subject that received therapy but continued to suffer from viral infection or one or more symptoms thereof. In certain embodiments, the subject can be a subject that received therapy but failed to achieve a sustained virologic response. In certain embodiments, the subject has received therapy for an HCV and/or HBV infection but

has failed to show, for example, a 2 log₁₀ decline in HCV RNA levels after 12 weeks of therapy. It is believed that subjects who have not shown more than 2 log₁₀ reduction in serum HCV RNA after 12 weeks of therapy have a 97-100% chance of not responding.

[00299] In certain embodiments, the subject is a subject that discontinued an HCV and/or HBV therapy because of one or more adverse events associated with the therapy. In certain embodiments, the subject is a subject where current therapy is not indicated. For instance, certain therapies for HCV are associated with neuropsychiatric events. Interferon (IFN)-alfa plus ribavirin is associated with a high rate of depression. Depressive symptoms have been linked to a worse outcome in a number of medical disorders. Life-threatening or fatal neuropsychiatric events, including suicide, suicidal and homicidal ideation, depression, relapse of drug addiction/overdose, and aggressive behavior have occurred in subjects with and without a previous psychiatric disorder during HCV therapy. Interferon-induced depression is a limitation for the treatment of chronic hepatitis C, especially for subjects with psychiatric disorders. Psychiatric side effects are common with interferon therapy and responsible for about 10% to 20% of discontinuations of current therapy for HCV infection.

[00300] Accordingly, provided are methods of treating or preventing an HCV infection in subjects where the risk of neuropsychiatric events, such as depression, contraindicates treatment with current HCV therapy. In one embodiment, provided are methods of treating or preventing HCV infection in subjects where a neuropsychiatric event, such as depression, or risk of such indicates discontinuation of treatment with current HCV therapy. Further provided are methods of treating or preventing HCV infection in subjects where a neuropsychiatric event, such as depression, or risk of such indicates dose reduction of current HCV therapy.

[00301] Current therapy is also contraindicated in subjects that are hypersensitive to interferon or ribavirin, or both, or any other component of a pharmaceutical product for administration of interferon or ribavirin. Current therapy is not indicated in subjects with hemoglobinopathies (e.g., thalassemia major, sickle-cell anemia) and other subjects at risk from the hematologic side effects of current therapy. Common hematologic side effects include bone marrow suppression, neutropenia and thrombocytopenia. Furthermore, ribavirin is toxic to red blood cells and is associated

with hemolysis. Accordingly, in one embodiment, provided are methods of treating or preventing HCV infection in subjects hypersensitive to interferon or ribavirin, or both, subjects with a hemoglobinopathy, for instance thalassemia major subjects and sickle-cell anemia subjects, and other subjects at risk from the hematologic side effects of current therapy.

[00302] In certain embodiments, the subject has received an HCV and/or HBV therapy and discontinued that therapy prior to administration of a method provided herein. In further embodiments, the subject has received therapy and continues to receive that therapy along with administration of a method provided herein. The methods can be co-administered with other therapy for HBC and/or HCV according to the judgment of one of skill in the art. In certain embodiments, the methods or compositions provided herein can be co-administered with a reduced dose of the other therapy for HBC and/or HCV.

[00303] In certain embodiments, provided are methods of treating a subject that is refractory to treatment with interferon. For instance, in some embodiments, the subject can be a subject that has failed to respond to treatment with one or more agents selected from the group consisting of interferon, interferon α , pegylated interferon α , interferon plus ribavirin, interferon α plus ribavirin and pegylated interferon α plus ribavirin. In some embodiments, the subject can be a subject that has responded poorly to treatment with one or more agents selected from the group consisting of interferon, interferon α , pegylated interferon α , interferon plus ribavirin, interferon α plus ribavirin and pegylated interferon α plus ribavirin. A pro-drug form of ribavirin, such as taribavirin, may also be used.

[00304] In certain embodiments, the subject has, or is at risk for, co-infection of HCV with HIV. For instance, in the United States, 30% of HIV subjects are co-infected with HCV and evidence indicates that people infected with HIV have a much more rapid course of their hepatitis C infection. Maier and Wu, 2002, *World J Gastroenterol* 8:577-57. The methods provided herein can be used to treat or prevent HCV infection in such subjects. It is believed that elimination of HCV in these subjects will lower mortality due to end-stage liver disease. Indeed, the risk of progressive liver disease is higher in subjects with severe AIDS-defining immunodeficiency than in those without. See, e.g., Lesens *et al.*, 1999, *J Infect Dis* 179:1254-1258. In one embodiment, compounds provided herein have been shown to

suppress HIV in HIV subjects. Thus, in certain embodiments, provided are methods of treating or preventing HIV infection and HCV infection in subjects in need thereof.

[00305] In certain embodiments, the compounds or compositions are administered to a subject following liver transplant. Hepatitis C is a leading cause of liver transplantation in the U.S, and many subjects that undergo liver transplantation remain HCV positive following transplantation. In one embodiment, provided are methods of treating such recurrent HCV subjects with a compound or composition provided herein. In certain embodiments, provided are methods of treating a subject before, during or following liver transplant to prevent recurrent HCV infection.

[00306] In certain embodiments, provided herein are methods for the treatment and/or prophylaxis of hepatitis B infections and other related conditions such as anti-HBV antibody positive and HBV-positive conditions, chronic liver inflammation caused by HBV, cirrhosis, acute hepatitis, fulminant hepatitis, chronic persistent hepatitis, and fatigue that includes administering an effective amount of a compound or composition provided herein.

[00307] In one embodiment, provided herein are methods for treatment and/or prophylaxis of hepatitis B infections and other related conditions such as anti-HBV antibody positive and HBV-positive conditions, chronic liver inflammation caused by HBV, cirrhosis, acute hepatitis, fulminant hepatitis, chronic persistent hepatitis, and fatigue that includes administering an effective amount of a compound or composition provided herein.

Second Therapeutic Agents

[00308] In certain embodiments, the compounds and compositions provided herein are useful in methods of treatment of a liver disorder, that comprises further administration of a second agent effective for the treatment of the disorder, such as HCV and/or HBV infection in a subject in need thereof. The second agent can be any agent known to those of skill in the art to be effective for the treatment of the disorder, including those currently approved by the FDA.

[00309] In certain embodiments, a compound provided herein is administered in combination with one second agent. In further embodiments, a second agent is administered in combination with two second agents. In still further embodiments, a second agent is administered in combination with two or more second agents.

[00310] As used herein, the term "in combination" includes the use of more than one therapy (e.g., one or more prophylactic and/or therapeutic agents). The use of the term "in combination" does not restrict the order in which therapies (e.g., prophylactic and/or therapeutic agents) are administered to a subject with a disorder. A first therapy (e.g., a prophylactic or therapeutic agent such as a compound provided herein) can be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapy (e.g., a prophylactic or therapeutic agent) to a subject with a disorder.

[00311] As used herein, the term "synergistic" includes a combination of a compound provided herein and another therapy (e.g., a prophylactic or therapeutic agent) which has been or is currently being used to prevent, manage or treat a disorder, which is more effective than the additive effects of the therapies. A synergistic effect of a combination of therapies (e.g., a combination of prophylactic or therapeutic agents) permits the use of lower dosages of one or more of the therapies and/or less frequent administration of said therapies to a subject with a disorder. The ability to utilize lower dosages of a therapy (e.g., a prophylactic or therapeutic agent) and/or to administer said therapy less frequently reduces the toxicity associated with the administration of said therapy to a subject without reducing the efficacy of said therapy in the prevention or treatment of a disorder). In addition, a synergistic effect can result in improved efficacy of agents in the prevention or treatment of a disorder. Finally, a synergistic effect of a combination of therapies (e.g., a combination of prophylactic or therapeutic agents) may avoid or reduce adverse or unwanted side effects associated with the use of either therapy alone.

[00312] The active compounds provided herein can be administered in combination or alternation with another therapeutic agent, in particular an anti-HCV or hepatitis B agent. In combination therapy, effective dosages of two or more agents are administered together, whereas in alternation or sequential-step therapy, an effective dosage of each agent is administered serially or sequentially. The dosages

given will depend on absorption, inactivation and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted that dosage values will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens and schedules should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions. In certain embodiments, an anti-HCV (or anti-pestivirus or anti-flavivirus) compound that exhibits an EC₅₀ of 10-15 μ M, or preferably less than 1-5 μ M, is desirable.

[00313] It has been recognized that drug-resistant variants of flaviviruses, pestiviruses or HCV can emerge after prolonged treatment with an antiviral agent. Drug resistance most typically occurs by mutation of a gene that encodes for an enzyme used in viral replication. The efficacy of a drug against the viral infection can be prolonged, augmented, or restored by administering the compound in combination or alternation with a second, and perhaps third, antiviral compound that induces a different mutation from that caused by the principle drug. Alternatively, the pharmacokinetics, biodistribution or other parameter of the drug can be altered by such combination or alternation therapy. In general, combination therapy is typically preferred over alternation therapy because it induces multiple simultaneous stresses on the virus.

[00314] Any of the viral treatments described in the Background of the Invention can be used in combination or alternation with the compounds described in this specification. Nonlimiting examples of second agents include:

[00315] HCV Protease inhibitors: Examples include Medivir HCV Protease Inhibitor (Medivir/Tobotec); ITMN-191 (InterMune), SCH 503034 (Schering) and VX950 (Vertex). Further examples of protease inhibitors include substrate-based NS3 protease inhibitors (Attwood *et al.*, Antiviral peptide derivatives, PCT WO 98/22496, 1998; Attwood *et al.*, *Antiviral Chemistry and Chemotherapy* 1999, 10, 259-273; Attwood *et al.*, Preparation and use of amino acid derivatives as anti-viral agents, German Patent Pub. DE 19914474; Tung *et al.* Inhibitors of serine proteases, particularly hepatitis C virus NS3 protease, PCT WO 98/17679), including alphaketoamides and hydrazinoureas, and inhibitors that terminate in an electrophile such as a boronic acid or phosphonate (Llinas-Brunet *et al.*, Hepatitis C inhibitor

peptide analogues, PCT WO 99/07734); Non-substrate-based NS3 protease inhibitors such as 2,4,6-trihydroxy-3-nitro-benzamide derivatives (Sudo K. *et al.*, *Biochemical and Biophysical Research Communications*, 1997, 238, 643-647; Sudo K. *et al.* *Antiviral Chemistry and Chemotherapy*, 1998, 9, 186), including RD3-4082 and RD3-4078, the former substituted on the amide with a 14 carbon chain and the latter processing a para-phenoxyphenyl group; and Sch 68631, a phenanthrenequinone, an HCV protease inhibitor (Chu M. *et al.*, *Tetrahedron Letters* 37:7229-7232, 1996).

[00316] SCH 351633, isolated from the fungus *Penicillium griseofulvum*, was identified as a protease inhibitor (Chu M. *et al.*, *Bioorganic and Medicinal Chemistry Letters* 9:1949-1952). Eglin c, isolated from leech, is a potent inhibitor of several serine proteases such as *S. griseus* proteases A and B, α -chymotrypsin, chymase and subtilisin. Qasim M.A. *et al.*, *Biochemistry* 36:1598-1607, 1997.

[00317] U.S. patents disclosing protease inhibitors for the treatment of HCV include, for example, U.S. Patent No. 6,004,933 to Spruce *et al.* which discloses a class of cysteine protease inhibitors for inhibiting HCV endopeptidase 2; U.S. Patent No. 5,990,276 to Zhang *et al.* which discloses synthetic inhibitors of hepatitis C virus NS3 protease; U.S. Patent No. 5,538,865 to Reyes *et al.*; WO 02/008251 to Corvas International, Inc. and US 7,169,760, US 2005/176648, WO 02/08187 and WO 02/008256 to Schering Corporation. HCV inhibitor tripeptides are disclosed in US Patent Nos. 6,534,523, 6,410,531, and 6,420,380 to Boehringer Ingelheim and WO 02/060926 to Bristol Myers Squibb. Diaryl peptides as NS3 serine protease inhibitors of HCV are disclosed in WO 02/48172 and US 6,911,428 to Schering Corporation. Imidazoleidinones as NS3 serine protease inhibitors of HCV are disclosed in WO 02/08198 and US 6,838,475 to Schering Corporation and WO 02/48157 and US 6,727,366 to Bristol Myers Squibb. WO 98/17679 and US 6,265,380 to Vertex Pharmaceuticals and WO 02/48116 and US 6,653,295 to Bristol Myers Squibb also disclose HCV protease inhibitors. Further examples of HCV serine protease inhibitors are provided in US 6,872,805 (Bristol-Myers Squibb); WO 2006/000085 (Boehringer Ingelheim); US 7,208,600 (Vertex); US 2006/0046956 (Schering-Plough); WO 2007/001406 (Chiron); US 2005/0153877; WO 2006/119061 (Merck); WO 00/09543 (Boehringer Ingelheim), US 6,323,180 (Boehringer Ingelheim) WO 03/064456 (Boehringer Ingelheim), US 6,642,204 (Boehringer Ingelheim), WO 03/064416 (Boehringer Ingelheim), US 7,091,184 (Boehringer Ingelheim), WO

03/053349 (Bristol-Myers Squibb), US 6,867,185, WO 03/099316 (Bristol-Myers Squibb), US 6,869,964, WO 03/099274 (Bristol-Myers Squibb), US 6,995,174, WO 2004/032827 (Bristol-Myers Squibb), US 7,041,698, WO 2004/043339 and US 6,878,722 (Bristol-Myers Squibb).

[00318] Thiazolidine derivatives which show relevant inhibition in a reverse-phase HPLC assay with an NS3/4A fusion protein and NS5A/5B substrate (Sudo K. *et al.*, *Antiviral Research*, 1996, 32, 9-18), especially compound RD-1-6250, possessing a fused cinnamoyl moiety substituted with a long alkyl chain, RD4 6205 and RD4 6193;

[00319] Thiazolidines and benzanilides identified in Kakiuchi N. *et al.* J. EBS Letters 421, 217-220; Takeshita N. *et al.* *Analytical Biochemistry*, 1997, 247, 242-246;

[00320] A phenanthrenequinone possessing activity against protease in a SDS-PAGE and autoradiography assay isolated from the fermentation culture broth of *Streptomyces* sp., SCH 68631 (Chu M. *et al.*, *Tetrahedron Letters*, 1996, 37, 7229-7232), and SCH 351633, isolated from the fungus *Penicillium griseofulvum*, which demonstrates activity in a scintillation proximity assay (Chu M. *et al.*, *Bioorganic and Medicinal Chemistry Letters* 9, 1949-1952);

[00321] Helicase inhibitors (Diana G.D. *et al.*, Compounds, compositions and methods for treatment of hepatitis C, U.S. Pat. No. 5,633,358; Diana G.D. *et al.*, Piperidine derivatives, pharmaceutical compositions thereof and their use in the treatment of hepatitis C, PCT WO 97/36554);

[00322] Nucleotide polymerase inhibitors and gliotoxin (Ferrari R. *et al.* *Journal of Virology*, 1999, 73, 1649-1654), and the natural product cerulenin (Lohmann V. *et al.*, *Virology*, 1998, 249, 108-118);

[00323] Interfering RNA (iRNA) based antivirals, including short interfering RNA (siRNA) based antivirals, such as Sirna-034 and others described in International Patent Publication Nos. WO/03/070750 and WO 2005/012525, and US Patent Publication No. US 2004/0209831.

[00324] Antisense phosphorothioate oligodeoxynucleotides (S-ODN) complementary to sequence stretches in the 5' non-coding region (NCR) of the virus (Alt M. *et al.*, *Hepatology*, 1995, 22, 707-717), or nucleotides 326-348 comprising the

3' end of the NCR and nucleotides 371-388 located in the core coding region of the HCV RNA (Alt M. *et al.*, *Archives of Virology*, 1997, 142, 589-599; Galderisi U. *et al.*, *Journal of Cellular Physiology*, 1999, 181, 251-257);

[00325] Inhibitors of IRES-dependent translation (Ikeda N *et al.*, Agent for the prevention and treatment of hepatitis C, Japanese Patent Pub. JP-08268890; Kai Y. *et al.* Prevention and treatment of viral diseases, Japanese Patent Pub. JP-10101591);

[00326] Ribozymes, such as nuclease-resistant ribozymes (Maccjak, D. J. *et al.*, *Hepatology* 1999, 30, abstract 995) and those disclosed in U.S. Patent No. 6,043,077 to Barber *et al.*, and U.S. Patent Nos. 5,869,253 and 5,610,054 to Draper *et al.*; and

[00327] Nucleoside analogs have also been developed for the treatment of Flaviviridae infections.

[00328] In certain embodiments, the compounds provided herein can be administered in combination with any of the compounds described by Idenix Pharmaceuticals in International Publication Nos. WO 01/90121, WO 01/92282, WO 2004/003000, 2004/002422 and WO 2004/002999.

[00329] Other patent applications disclosing the use of certain nucleoside analogs that can be used as second agents to treat hepatitis C virus include: PCT/CA00/01316 (WO 01/32153; filed November 3, 2000) and PCT/CA01/00197 (WO 01/60315; filed February 19, 2001) filed by BioChem Pharma, Inc. (now Shire Biochem, Inc.); PCT/US02/01531 (WO 02/057425; filed January 18, 2002); PCT/US02/03086 (WO 02/057287; filed January 18, 2002); US 7,202,224; 7,125,855; 7,105,499 and 6,777,395 by Merck & Co., Inc.; PCT/EP01/09633 (WO 02/18404; published August 21, 2001); US 2006/0040890; 2005/0038240; 2004/0121980; 6,846,810; 6,784,166 and 6,660,721 by Roche; PCT Publication Nos. WO 01/79246 (filed April 13, 2001), WO 02/32920 (filed October 18, 2001) and WO 02/48165; US 2005/0009737; US 2005/0009737; 7,094,770 and 6,927,291 by Pharmasset, Ltd.

[00330] Further compounds that can be used as second agents to treat hepatitis C virus are disclosed in PCT Publication No. WO 99/43691 to Emory University, entitled "2'-Fluoronucleosides". The use of certain 2'-fluoronucleosides to treat HCV is disclosed.

[00331] Other miscellaneous compounds that can be used as second agents include 1-amino-alkylcyclohexanes (U.S. Patent No. 6,034,134 to Gold *et al.*), alkyl lipids

(U.S. Pat. No. 5,922,757 to Chojkier *et al.*), vitamin E and other antioxidants (U.S. Pat. No. 5,922,757 to Chojkier *et al.*), squalene, amantadine, bile acids (U.S. Pat. No. 5,846,964 to Ozeki *et al.*), N-(phosphonoacetyl)-L-aspartic acid, (U.S. Pat. No. 5,830,905 to Diana *et al.*), benzenedicarboxamides (U.S. Pat. No. 5,633,388 to Diana *et al.*), polyadenylic acid derivatives (U.S. Pat. No. 5,496,546 to Wang *et al.*), 2',3'-dideoxyinosine (U.S. Pat. No. 5,026,687 to Yarchoan *et al.*), benzimidazoles (U.S. Pat. No. 5,891,874 to Colacino *et al.*), plant extracts (U.S. Patent No. 5,837,257 to Tsai *et al.*, U.S. Patent No. 5,725,859 to Omer *et al.*, and U.S. Patent No. 6,056,961), and piperidenes (U.S. Patent No. 5,830,905 to Diana *et al.*).

Exemplary Second Agents for Treatment of HCV

[00332] In one embodiment, one or more compounds provided herein can be administered in combination or alternation with an anti-hepatitis C virus interferon, such as Intron A[®] (interferon alfa-2b) and Pegasys[®] (Peginterferon alfa-2a); Roferon A[®] (Recombinant interferon alfa-2a), Infergen[®] (consensus interferon;interferon alfacon-1), PEG-Intron[®] (pegylated interferon alfa-2b) and Pegasys[®] (pegylated interferon alfa-2a).

[00333] In one embodiment, the anti-hepatitis C virus interferon is infergen, IL-29 (PEG-Interferon lambda), R7025 (Maxy-alpha), Belerofon, Oral Interferon alpha, BLX-883 (Locteron), omega interferon, multiferon, medusa interferon, Albuferon or REBIF[®].

[00334] In one embodiment, one or more compounds provided herein can be administered in combination or alternation with an anti-hepatitis C virus polymerase inhibitor, such as ribavirin, viramidine, NM 283 (valopicitabine), PSI-6130, R1626, HCV-796 or R7128.

[00335] In certain embodiments, the one or more compounds provided herein can be administered in combination with ribavarin and an anti-hepatitis C virus interferon, such as Intron A[®] (interferon alfa-2b) and Pegasys[®] (Peginterferon alfa-2a); Roferon A[®] (Recombinant interferon alfa-2a), Infergen[®] (consensus interferon;interferon alfacon-1), PEG-Intron[®] (pegylated interferon alfa-2b) and Pegasys[®] (pegylated interferon alfa-2a).

[00336] In one embodiment, one or more compounds provided herein can be administered in combination or alternation with an anti-hepatitis C virus protease

inhibitor such as ITMN-191, SCH 503034, VX950 (telaprevir) or Medivir HCV Protease Inhibitor.

[00337] In one embodiment, one or more compounds provided herein can be administered in combination or alternation with an anti-hepatitis C virus vaccine, such as TG4040, PeviPRO™, CGI-5005, HCV/MF59, GV1001, IC41 or INNO0101 (E1).

[00338] In one embodiment, one or more compounds provided herein can be administered in combination or alternation with an anti-hepatitis C virus monoclonal antibody, such as AB68 or XTL-6865 (formerly HepX-C); or an anti-hepatitis C virus polyclonal antibody, such as cicavir.

[00339] In one embodiment, one or more compounds provided herein can be administered in combination or alternation with an anti-hepatitis C virus immunomodulator, such as Zadaxin® (thymalfasin), NOV-205 or Oglufanide.

[00340] In one embodiment, one or more compounds provided herein can be administered in combination or alternation with Nexavar, doxorubicin, PI-88, amantadine, JBK-122, VGX-410C, MX-3253 (Ceglosivir), Suvus (BIVN-401 or virostat), PF-03491390 (formerly IDN-6556), G126270, UT-231B, DEBIO-025, EMZ702, ACH-0137171, MitoQ, ANA975, AVI-4065, Bavituxinab (Tarvacin), Alinia (nitrazoxanide) or PYN17.

Exemplary Second Agents for Treatment of HBV

[00341] It has been recognized that drug-resistant variants of HBV can emerge after prolonged treatment with an antiviral agent. Drug resistance most typically occurs by mutation of a gene that encodes for an enzyme used in the viral life cycle, and most typically in the case of HBV, DNA polymerase. The efficacy of a drug against HBV infection can be prolonged, augmented, or restored by administering the compound in combination or alternation with a second, and perhaps third, antiviral compound that induces a different mutation from that caused by the principle drug. Alternatively, the pharmacokinetics, biodistribution, or other parameter of the drug can be altered by such combination or alternation therapy. In general, combination therapy is typically preferred over alternation therapy because it induces multiple simultaneous stresses on the virus.

[00342] The anti-hepatitis B viral activity of compounds provided herein can be enhanced by administering one or more of these further agents in combination or

alternation. Alternatively, for example, one or more compounds provided herein can be administered in combination or alternation with any other known anti-hepatitis B virus agent. Such agents include anti-hepatitis B virus interferons, such as Intron A[®] (interferon alfa-2b) and Pegasys[®] (Peginterferon alfa-2a); polymerase inhibitors, such as Epivir-HBV (lamivudine), Hepsera (adefovir dipivoxil), baraclude (entecavir), Tyzeka (telbivudine), Emtricitabine (FTC), Clevudine (L-FMAU), Viread (tenofovir), Valtorcitabine, Amdoxovir, ANA 380, Pradefovir (remofovir) and RCV (racivir); vaccines, such as Hi-8 HBV, HepaVaxx B and HBV Core Antigen vaccine; and other agents, such as HepX, SpecifEx-HepB, Zadaxin, EHT899, Bay 41-4109, UT 231-B, HepeX-B and NOV-205 or any other compound that exhibits an EC₅₀ of less than 15 micromolar in 2.2.15 cells; or their prodrugs or pharmaceutically acceptable salts. Several other examples of anti-HBV agents are provided in U.S. Application Publication No. 20050080034 and international publication no. WO 2004/096286, which are incorporated by reference in their entireties.

[00343] In one embodiment, one or more compounds provided herein can be administered in combination or alternation with anti-hepatitis B virus agent such as interferon α -2b, peginterferon α -2a, lamivudine, hepsera, baraclude, telbivudine, emtricitabine, clevudine, tenofovir, valtorcitabine, amdoxovir, ANA 380, remofovir, racivir, alinia, Hi-8 HBV and HepaVaxx B.

[00344] In another embodiment, a compound provided herein is administered in combination or alternation with an immune modulator or other pharmaceutically active modifier of viral replication, including a biological material such as a protein, peptide, oligonucleotide, or gamma globulin, including but not limited to interfereon, interleukin, or an antisense oligonucleotides to genes which express or regulate hepatitis B replication.

[00345] Any method of alternation can be used that provides treatment to the patient. Nonlimiting examples of alternation patterns include 1-6 weeks of administration of an effective amount of one agent followed by 1-6 weeks of administration of an effective amount of a second anti-HBV agent. The alternation schedule can include periods of no treatment. Combination therapy generally includes the simultaneous administration of an effective ratio of dosages of two or more anti-HBV agents.

[00346] In light of the fact that HBV is often found in patients who are also anti-HIV antibody or HIV-antigen positive or who have been exposed to HIV, the active anti-HBV compounds disclosed herein or their derivatives or prodrugs can be administered in the appropriate circumstance in combination or alternation with anti-HIV medications.

[00347] The compounds provided herein can also be administered in combination with antibiotics, other antiviral compounds, antifungal agents or other pharmaceutical agents administered for the treatment of secondary infections.

Pharmaceutical Compositions and Methods of Administration

[00348] Phosphoramidate and phosphonoamidate compounds of a variety of therapeutic agents can be formulated into pharmaceutical compositions using methods available in the art and those disclosed herein. Such compounds can be used in some embodiments to enhance delivery of the drug to the liver. In one embodiment, the compound comprises a *S*-acyl-2-thioethyl phosphoramidate or *S*-acyl-2-thioethyl phosphonoamidate, e.g., a *S*-pivaloyl-2-thioethyl phosphoramidate or *S*-hydroxypivaloyl-2-thioethyl phosphonoamidate derivative. Therapeutic agents that can be derivatized to phosphoramidate or phosphonoamidate compound form include any anti-viral agent that includes, or has been derivatized to include a reactive group for attachment of the phosphoramidate or phosphonoamidate moiety, including but not limited to nucleosides and nucleoside analogues including acyclic nucleosides. Any of the phosphoramidate or phosphonoamidate compounds disclosed herein can be provided in the appropriate pharmaceutical composition and be administered by a suitable route of administration.

[00349] The methods provided herein encompass administering pharmaceutical compositions containing at least one compound as described herein, including a compound of general Formula I, IIa or IIb, if appropriate in the salt form, either used alone or in the form of a combination with one or more compatible and pharmaceutically acceptable carriers, such as diluents or adjuvants, or with another anti-HCV or anti-HBV agent.

[00350] In certain embodiments, the second agent can be formulated or packaged with the compound provided herein. Of course, the second agent will only be formulated with the compound provided herein when, according to the judgment of

those of skill in the art, such co-formulation should not interfere with the activity of either agent or the method of administration. In certain embodiments, the compound provided herein and the second agent are formulated separately. They can be packaged together, or packaged separately, for the convenience of the practitioner of skill in the art.

[00351] In clinical practice the active agents provided herein may be administered by any conventional route, in particular orally, parenterally, rectally or by inhalation (*e.g.* in the form of aerosols). In certain embodiments, the compound provided herein is administered orally.

[00352] Use may be made, as solid compositions for oral administration, of tablets, pills, hard gelatin capsules, powders or granules. In these compositions, the active product is mixed with one or more inert diluents or adjuvants, such as sucrose, lactose or starch.

[00353] These compositions can comprise substances other than diluents, for example a lubricant, such as magnesium stearate, or a coating intended for controlled release.

[00354] Use may be made, as liquid compositions for oral administration, of solutions which are pharmaceutically acceptable, suspensions, emulsions, syrups and elixirs containing inert diluents, such as water or liquid paraffin. These compositions can also comprise substances other than diluents, for example wetting, sweetening or flavoring products.

[00355] The compositions for parenteral administration can be emulsions or sterile solutions. Use may be made, as solvent or vehicle, of propylene glycol, a polyethylene glycol, vegetable oils, in particular olive oil, or injectable organic esters, for example ethyl oleate. These compositions can also contain adjuvants, in particular wetting, isotonicizing, emulsifying, dispersing and stabilizing agents. Sterilization can be carried out in several ways, for example using a bacteriological filter, by radiation or by heating. They can also be prepared in the form of sterile solid compositions which can be dissolved at the time of use in sterile water or any other injectable sterile medium.

[00356] The compositions for rectal administration are suppositories or rectal capsules which contain, in addition to the active principle, excipients such as cocoa butter, semi-synthetic glycerides or polyethylene glycols.

[00357] The compositions can also be aerosols. For use in the form of liquid aerosols, the compositions can be stable sterile solutions or solid compositions dissolved at the time of use in apyrogenic sterile water, in saline or any other pharmaceutically acceptable vehicle. For use in the form of dry aerosols intended to be directly inhaled, the active principle is finely divided and combined with a water-soluble solid diluent or vehicle, for example dextran, mannitol or lactose.

[00358] In one embodiment, a composition provided herein is a pharmaceutical composition or a single unit dosage form. Pharmaceutical compositions and single unit dosage forms provided herein comprise a prophylactically or therapeutically effective amount of one or more prophylactic or therapeutic agents (*e.g.*, a compound provided herein, or other prophylactic or therapeutic agent), and a typically one or more pharmaceutically acceptable carriers or excipients. In a specific embodiment and in this context, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" includes a diluent, adjuvant (*e.g.*, Freund's adjuvant (complete and incomplete)), excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water can be used as a carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin.

[00359] Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well-known to those skilled in the art of pharmacy, and non limiting examples of suitable excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol,

water, ethanol and the like. Whether a particular excipient is suitable for incorporation into a pharmaceutical composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a subject and the specific active ingredients in the dosage form. The composition or single unit dosage form, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents.

[00360] Lactose free compositions provided herein can comprise excipients that are well known in the art and are listed, for example, in the U.S. Pharmacopia (USP) SP (XXI)/NF (XVI). In general, lactose free compositions comprise an active ingredient, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Exemplary lactose free dosage forms comprise an active ingredient, microcrystalline cellulose, pre gelatinized starch, and magnesium stearate.

[00361] Further encompassed herein are anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (*e.g.*, 5%) is widely accepted in the pharmaceutical arts as a means of simulating long term storage in order to determine characteristics such as shelf life or the stability of formulations over time. See, *e.g.*, Jens T. Carstensen, *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, NY, 1995, pp. 379 80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

[00362] Anhydrous pharmaceutical compositions and dosage forms provided herein can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine can be anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

[00363] An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions

can be packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (*e.g.*, vials), blister packs, and strip packs.

[00364] Further provided are pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as “stabilizers,” include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

[00365] The pharmaceutical compositions and single unit dosage forms can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Such compositions and dosage forms will contain a prophylactically or therapeutically effective amount of a prophylactic or therapeutic agent, in certain embodiments, in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the subject. The formulation should suit the mode of administration. In a certain embodiment, the pharmaceutical compositions or single unit dosage forms are sterile and in suitable form for administration to a subject, for example, an animal subject, such as a mammalian subject, for example, a human subject.

[00366] A pharmaceutical composition is formulated to be compatible with its intended route of administration. Examples of routes of administration include, but are not limited to, parenteral, *e.g.*, intravenous, intradermal, subcutaneous, intramuscular, subcutaneous, oral, buccal, sublingual, inhalation, intranasal, transdermal, topical, transmucosal, intra-tumoral, intra-synovial and rectal administration. In a specific embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous, subcutaneous, intramuscular, oral, intranasal or topical administration to human beings. In an embodiment, a pharmaceutical composition is formulated in accordance with routine procedures for subcutaneous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing

agent and a local anesthetic such as lignocaine to ease pain at the site of the injection.

[00367] Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; ointments; cataplasms (poultices); pastes; powders; dressings; creams; plasters; solutions; patches; aerosols (*e.g.*, nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a subject, including suspensions (*e.g.*, aqueous or non aqueous liquid suspensions, oil in water emulsions, or a water in oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a subject; and sterile solids (*e.g.*, crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a subject.

[00368] The composition, shape, and type of dosage forms provided herein will typically vary depending on their use. For example, a dosage form used in the initial treatment of viral infection may contain larger amounts of one or more of the active ingredients it comprises than a dosage form used in the maintenance treatment of the same infection. Similarly, a parenteral dosage form may contain smaller amounts of one or more of the active ingredients it comprises than an oral dosage form used to treat the same disease or disorder. These and other ways in which specific dosage forms encompassed herein will vary from one another will be readily apparent to those skilled in the art. See, *e.g.*, Remington's Pharmaceutical Sciences, 20th ed., Mack Publishing, Easton PA (2000).

[00369] Generally, the ingredients of compositions are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[00370] Typical dosage forms comprise a compound provided herein, or a pharmaceutically acceptable salt, solvate or hydrate thereof lie within the range of

from about 0.1 mg to about 1000 mg per day, given as a single once-a-day dose in the morning or as divided doses throughout the day taken with food. Particular dosage forms can have about 0.1, 0.2, 0.3, 0.4, 0.5, 1.0, 2.0, 2.5, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 100, 200, 250, 500 or 1000 mg of the active compound.

Oral Dosage Forms

[00371] Pharmaceutical compositions that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (*e.g.*, chewable tablets), caplets, capsules, and liquids (*e.g.*, flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. See generally, Remington's Pharmaceutical Sciences, 20th ed., Mack Publishing, Easton PA (2000).

[00372] In certain embodiments, the oral dosage forms are solid and prepared under anhydrous conditions with anhydrous ingredients, as described in detail in the sections above. However, the scope of the compositions provided herein extends beyond anhydrous, solid oral dosage forms. As such, further forms are described herein.

[00373] Typical oral dosage forms are prepared by combining the active ingredient(s) in an intimate admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use in solid oral dosage forms (*e.g.*, powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

[00374] Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and intimately admixing the active ingredients with liquid carriers, finely divided

solid carriers, or both, and then shaping the product into the desired presentation if necessary.

[00375] For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free flowing form such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

[00376] Examples of excipients that can be used in oral dosage forms include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (*e.g.*, ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre gelatinized starch, hydroxypropyl methyl cellulose, (*e.g.*, Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

[00377] Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (*e.g.*, granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions is typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

[00378] Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL PH 101, AVICEL PH 103 AVICEL RC 581, AVICEL PH 105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA), and mixtures thereof. An specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC 581. Suitable anhydrous or low moisture excipients or additives include AVICEL PH 103™ and Starch 1500 LM.

[00379] Disintegrants are used in the compositions to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much

disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, specifically from about 1 to about 5 weight percent of disintegrant.

[00380] Disintegrants that can be used in pharmaceutical compositions and dosage forms include, but are not limited to, agar agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, pre gelatinized starch, other starches, clays, other algin, other celluloses, gums, and mixtures thereof.

[00381] Lubricants that can be used in pharmaceutical compositions and dosage forms include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (*e.g.*, peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL 200, manufactured by W.R. Grace Co. of Baltimore, MD), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, TX), CAB O SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, MA), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

Delayed Release Dosage Forms

[00382] Active ingredients such as the compounds provided herein can be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719; 5,674,533; 5,059,595; 5,591,767; 5,120,548; 5,073,543; 5,639,476; 5,354,556; 5,639,480; 5,733,566; 5,739,108; 5,891,474; 5,922,356; 5,972,891; 5,980,945; 5,993,855; 6,045,830; 6,087,324; 6,113,943; 6,197,350; 6,248,363;

6,264,970; 6,267,981; 6,376,461; 6,419,961; 6,589,548; 6,613,358; 6,699,500 each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients provided herein. Thus encompassed herein are single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled release.

[00383] All controlled release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non controlled counterparts. Ideally, the use of an optimally designed controlled release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled release formulations include extended activity of the drug, reduced dosage frequency, and increased subject compliance. In addition, controlled release formulations can be used to affect the time of onset of action or other characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (*e.g.*, adverse) effects.

[00384] Most controlled release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

[00385] In certain embodiments, the drug may be administered using intravenous infusion, an implantable osmotic pump, a transdermal patch, liposomes, or other modes of administration. In one embodiment, a pump may be used (*see*, Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald *et al.*, *Surgery* 88:507 (1980);

Saudek *et al.*, *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used. In yet another embodiment, a controlled release system can be placed in a subject at an appropriate site determined by a practitioner of skill, *i.e.*, thus requiring only a fraction of the systemic dose (*see, e.g.*, Goodson, *Medical Applications of Controlled Release*, vol. 2, pp. 115-138 (1984)). Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)). The active ingredient can be dispersed in a solid inner matrix, *e.g.*, polymethylmethacrylate, polybutylmethacrylate, plasticized or unplasticized polyvinylchloride, plasticized nylon, plasticized polyethyleneterephthalate, natural rubber, polyisoprene, polyisobutylene, polybutadiene, polyethylene, ethylene-vinylacetate copolymers, silicone rubbers, polydimethylsiloxanes, silicone carbonate copolymers, hydrophilic polymers such as hydrogels of esters of acrylic and methacrylic acid, collagen, cross-linked polyvinylalcohol and cross-linked partially hydrolyzed polyvinyl acetate, that is surrounded by an outer polymeric membrane, *e.g.*, polyethylene, polypropylene, ethylene/propylene copolymers, ethylene/ethyl acrylate copolymers, ethylene/vinylacetate copolymers, silicone rubbers, polydimethyl siloxanes, neoprene rubber, chlorinated polyethylene, polyvinylchloride, vinylchloride copolymers with vinyl acetate, vinylidene chloride, ethylene and propylene, ionomer polyethylene terephthalate, butyl rubber epichlorohydrin rubbers, ethylene/vinyl alcohol copolymer, ethylene/vinyl acetate/vinyl alcohol terpolymer, and ethylene/vinyloxyethanol copolymer, that is insoluble in body fluids. The active ingredient then diffuses through the outer polymeric membrane in a release rate controlling step. The percentage of active ingredient in such parenteral compositions is highly dependent on the specific nature thereof, as well as the needs of the subject.

Parenteral Dosage Forms

[00386] In one embodiment, provided are parenteral dosage forms. Parenteral dosage forms can be administered to subjects by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses subjects' natural defenses against contaminants, parenteral dosage forms are typically, sterile or capable of being sterilized prior to administration to a subject. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry

products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

[00387] Suitable vehicles that can be used to provide parenteral dosage forms are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

[00388] Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms.

Transdermal, Topical & Mucosal Dosage Forms

[00389] Also provided are transdermal, topical, and mucosal dosage forms. Transdermal, topical, and mucosal dosage forms include, but are not limited to, ophthalmic solutions, sprays, aerosols, creams, lotions, ointments, gels, solutions, emulsions, suspensions, or other forms known to one of skill in the art. See, *e.g.*, Remington's Pharmaceutical Sciences, 16th, 18th and 20th eds., Mack Publishing, Easton PA (1980, 1990 & 2000); and Introduction to Pharmaceutical Dosage Forms, 4th ed., Lea & Febiger, Philadelphia (1985). Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or as oral gels. Further, transdermal dosage forms include "reservoir type" or "matrix type" patches, which can be applied to the skin and worn for a specific period of time to permit the penetration of a desired amount of active ingredients.

[00390] Suitable excipients (*e.g.*, carriers and diluents) and other materials that can be used to provide transdermal, topical, and mucosal dosage forms encompassed herein are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane 1,3 diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to form lotions, tinctures, creams, emulsions, gels or ointments, which are non toxic and

pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. See, *e.g.*, Remington's Pharmaceutical Sciences, 16th, 18th and 20th eds., Mack Publishing, Easton PA (1980, 1990 & 2000).

[00391] Depending on the specific tissue to be treated, additional components may be used prior to, in conjunction with, or subsequent to treatment with active ingredients provided. For example, penetration enhancers can be used to assist in delivering the active ingredients to the tissue. Suitable penetration enhancers include, but are not limited to: acetone; various alcohols such as ethanol, oleyl, and tetrahydrofuryl; alkyl sulfoxides such as dimethyl sulfoxide; dimethyl acetamide; dimethyl formamide; polyethylene glycol; pyrrolidones such as polyvinylpyrrolidone; Kollidon grades (Povidone, Polyvidone); urea; and various water soluble or insoluble sugar esters such as Tween 80 (polysorbate 80) and Span 60 (sorbitan monostearate).

[00392] The pH of a pharmaceutical composition or dosage form, or of the tissue to which the pharmaceutical composition or dosage form is applied, may also be adjusted to improve delivery of one or more active ingredients. Similarly, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In this regard, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, and as a delivery enhancing or penetration enhancing agent. Different salts, hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

Dosage and Unit Dosage Forms

[00393] In human therapeutics, the doctor will determine the posology which he considers most appropriate according to a preventive or curative treatment and according to the age, weight, stage of the infection and other factors specific to the subject to be treated. In certain embodiments, doses are from about 1 to about 1000 mg per day for an adult, or from about 5 to about 250 mg per day or from about 10 to 50 mg per day for an adult. In certain embodiments, doses are from about 5 to about

400 mg per day or 25 to 200 mg per day per adult. In certain embodiments, dose rates of from about 50 to about 500 mg per day are also contemplated.

[00394] In further aspects, provided are methods of treating or preventing an HCV and/or HBV infection in a subject by administering, to a subject in need thereof, an effective amount of a compound provided herein, or a pharmaceutically acceptable salt thereof. The amount of the compound or composition which will be effective in the prevention or treatment of a disorder or one or more symptoms thereof will vary with the nature and severity of the disease or condition, and the route by which the active ingredient is administered. The frequency and dosage will also vary according to factors specific for each subject depending on the specific therapy (*e.g.*, therapeutic or prophylactic agents) administered, the severity of the disorder, disease, or condition, the route of administration, as well as age, body, weight, response, and the past medical history of the subject. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[00395] In certain embodiments, exemplary doses of a composition include milligram or microgram amounts of the active compound per kilogram of subject or sample weight (*e.g.*, about 10 micrograms per kilogram to about 50 milligrams per kilogram, about 100 micrograms per kilogram to about 25 milligrams per kilogram, or about 100 microgram per kilogram to about 10 milligrams per kilogram). For compositions provided herein, in certain embodiments, the dosage administered to a subject is 0.140 mg/kg to 3 mg/kg of the subject's body weight, based on weight of the active compound. In certain embodiments, the dosage administered to a subject is between 0.20 mg/kg and 2.00 mg/kg, or between 0.30 mg/kg and 1.50 mg/kg of the subject's body weight.

[00396] In certain embodiments, the recommended daily dose range of a composition provided herein for the conditions described herein lie within the range of from about 0.1 mg to about 1000 mg per day, given as a single once-a-day dose or as divided doses throughout a day. In one embodiment, the daily dose is administered twice daily in equally divided doses. In certain embodiments, a daily dose range should be from about 10 mg to about 200 mg per day, in other embodiments, between about 10 mg and about 150 mg per day, in further embodiments, between about 25 and about 100 mg per day. It may be necessary to use dosages of the active ingredient outside the ranges disclosed herein in some cases, as will be apparent to those of

ordinary skill in the art. Furthermore, it is noted that the clinician or treating physician will know how and when to interrupt, adjust, or terminate therapy in conjunction with subject response.

[00397] Different therapeutically effective amounts may be applicable for different diseases and conditions, as will be readily known by those of ordinary skill in the art. Similarly, amounts sufficient to prevent, manage, treat or ameliorate such disorders, but insufficient to cause, or sufficient to reduce, adverse effects associated with the composition provided herein are also encompassed by the above described dosage amounts and dose frequency schedules. Further, when a subject is administered multiple dosages of a composition provided herein, not all of the dosages need be the same. For example, the dosage administered to the subject may be increased to improve the prophylactic or therapeutic effect of the composition or it may be decreased to reduce one or more side effects that a particular subject is experiencing.

[00398] In certain embodiment, the dosage of the composition provided herein, based on weight of the active compound, administered to prevent, treat, manage, or ameliorate a disorder, or one or more symptoms thereof in a subject is 0.1 mg/kg, 1 mg/kg, 2 mg/kg, 3 mg/kg, 4 mg/kg, 5 mg/kg, 6 mg/kg, 10 mg/kg, or 15 mg/kg or more of a subject's body weight. In another embodiment, the dosage of the composition or a composition provided herein administered to prevent, treat, manage, or ameliorate a disorder, or one or more symptoms thereof in a subject is a unit dose of 0.1 mg to 200 mg, 0.1 mg to 100 mg, 0.1 mg to 50 mg, 0.1 mg to 25 mg, 0.1 mg to 20 mg, 0.1 mg to 15 mg, 0.1 mg to 10 mg, 0.1 mg to 7.5 mg, 0.1 mg to 5 mg, 0.1 to 2.5 mg, 0.25 mg to 20 mg, 0.25 to 15 mg, 0.25 to 12 mg, 0.25 to 10 mg, 0.25 mg to 7.5 mg, 0.25 mg to 5 mg, 0.5 mg to 2.5 mg, 1 mg to 20 mg, 1 mg to 15 mg, 1 mg to 12 mg, 1 mg to 10 mg, 1 mg to 7.5 mg, 1 mg to 5 mg, or 1 mg to 2.5 mg.

[00399] In certain embodiments, treatment or prevention can be initiated with one or more loading doses of a compound or composition provided herein followed by one or more maintenance doses. In such embodiments, the loading dose can be, for instance, about 60 to about 400 mg per day, or about 100 to about 200 mg per day for one day to five weeks. The loading dose can be followed by one or more maintenance doses. In certain embodiments, each maintenance does is, independently, about from about 10 mg to about 200 mg per day, between about 25 mg and about 150 mg per day, or between about 25 and about 80 mg per day.

Maintenance doses can be administered daily and can be administered as single doses, or as divided doses.

[00400] In certain embodiments, a dose of a compound or composition provided herein can be administered to achieve a steady-state concentration of the active ingredient in blood or serum of the subject. The steady-state concentration can be determined by measurement according to techniques available to those of skill or can be based on the physical characteristics of the subject such as height, weight and age. In certain embodiments, a sufficient amount of a compound or composition provided herein is administered to achieve a steady-state concentration in blood or serum of the subject of from about 300 to about 4000 ng/mL, from about 400 to about 1600 ng/mL, or from about 600 to about 1200 ng/mL. In some embodiments, loading doses can be administered to achieve steady-state blood or serum concentrations of about 1200 to about 8000 ng/mL, or about 2000 to about 4000 ng/mL for one to five days. In certain embodiments, maintenance doses can be administered to achieve a steady-state concentration in blood or serum of the subject of from about 300 to about 4000 ng/mL, from about 400 to about 1600 ng/mL, or from about 600 to about 1200 ng/mL.

[00401] In certain embodiments, administration of the same composition may be repeated and the administrations may be separated by at least 1 day, 2 days, 3 days, 5 days, 10 days, 15 days, 30 days, 45 days, 2 months, 75 days, 3 months, or 6 months. In other embodiments, administration of the same prophylactic or therapeutic agent may be repeated and the administration may be separated by at least at least 1 day, 2 days, 3 days, 5 days, 10 days, 15 days, 30 days, 45 days, 2 months, 75 days, 3 months, or 6 months.

[00402] In certain aspects, provided herein are unit dosages comprising a compound, or a pharmaceutically acceptable salt thereof, in a form suitable for administration. Such forms are described in detail above. In certain embodiments, the unit dosage comprises 1 to 1000 mg, 5 to 250 mg or 10 to 50 mg active ingredient. In particular embodiments, the unit dosages comprise about 1, 5, 10, 25, 50, 100, 125, 250, 500 or 1000 mg active ingredient. Such unit dosages can be prepared according to techniques familiar to those of skill in the art.

[00403] The dosages of the second agents are to be used in the combination therapies provided herein. In certain embodiments, dosages lower than those which have been or are currently being used to prevent or treat HCV and/or HBV infection are used in the combination therapies provided herein. The recommended dosages of second agents can be obtained from the knowledge of those of skill. For those second agents that are approved for clinical use, recommended dosages are described in, for example, Hardman *et al.*, eds., 1996, Goodman & Gilman's The Pharmacological Basis Of Therapeutics 9th Ed, Mc-Graw-Hill, New York; Physician's Desk Reference (PDR) 57th Ed., 2003, Medical Economics Co., Inc., Montvale, NJ, which are incorporated herein by reference in its entirety.

[00404] In various embodiments, the therapies (*e.g.*, a compound provided herein and the second agent) are administered less than 5 minutes apart, less than 30 minutes apart, 1 hour apart, at about 1 hour apart, at about 1 to about 2 hours apart, at about 2 hours to about 3 hours apart, at about 3 hours to about 4 hours apart, at about 4 hours to about 5 hours apart, at about 5 hours to about 6 hours apart, at about 6 hours to about 7 hours apart, at about 7 hours to about 8 hours apart, at about 8 hours to about 9 hours apart, at about 9 hours to about 10 hours apart, at about 10 hours to about 11 hours apart, at about 11 hours to about 12 hours apart, at about 12 hours to 18 hours apart, 18 hours to 24 hours apart, 24 hours to 36 hours apart, 36 hours to 48 hours apart, 48 hours to 52 hours apart, 52 hours to 60 hours apart, 60 hours to 72 hours apart, 72 hours to 84 hours apart, 84 hours to 96 hours apart, or 96 hours to 120 hours apart. In various embodiments, the therapies are administered no more than 24 hours apart or no more than 48 hours apart. In certain embodiments, two or more therapies are administered within the same patient visit. In other embodiments, the compound provided herein and the second agent are administered concurrently.

[00405] In other embodiments, the compound provided herein and the second agent are administered at about 2 to 4 days apart, at about 4 to 6 days apart, at about 1 week part, at about 1 to 2 weeks apart, or more than 2 weeks apart.

[00406] In certain embodiments, administration of the same agent may be repeated and the administrations may be separated by at least 1 day, 2 days, 3 days, 5 days, 10 days, 15 days, 30 days, 45 days, 2 months, 75 days, 3 months, or 6 months. In other embodiments, administration of the same agent may be repeated and the

administration may be separated by at least at least 1 day, 2 days, 3 days, 5 days, 10 days, 15 days, 30 days, 45 days, 2 months, 75 days, 3 months, or 6 months.

[00407] In certain embodiments, a compound provided herein and a second agent are administered to a patient, for example, a mammal, such as a human, in a sequence and within a time interval such that the compound provided herein can act together with the other agent to provide an increased benefit than if they were administered otherwise. For example, the second active agent can be administered at the same time or sequentially in any order at different points in time; however, if not administered at the same time, they should be administered sufficiently close in time so as to provide the desired therapeutic or prophylactic effect. In one embodiment, the compound provided herein and the second active agent exert their effect at times which overlap. Each second active agent can be administered separately, in any appropriate form and by any suitable route. In other embodiments, the compound provided herein is administered before, concurrently or after administration of the second active agent.

[00408] In certain embodiments, the compound provided herein and the second agent are cyclically administered to a patient. Cycling therapy involves the administration of a first agent (*e.g.*, a first prophylactic or therapeutic agents) for a period of time, followed by the administration of a second agent and/or third agent (*e.g.*, a second and/or third prophylactic or therapeutic agents) for a period of time and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improve the efficacy of the treatment.

[00409] In certain embodiments, the compound provided herein and the second active agent are administered in a cycle of less than about 3 weeks, about once every two weeks, about once every 10 days or about once every week. One cycle can comprise the administration of a compound provided herein and the second agent by infusion over about 90 minutes every cycle, about 1 hour every cycle, about 45 minutes every cycle. Each cycle can comprise at least 1 week of rest, at least 2 weeks of rest, at least 3 weeks of rest. The number of cycles administered is from about 1 to about 12 cycles, more typically from about 2 to about 10 cycles, and more typically from about 2 to about 8 cycles.

[00410] In other embodiments, courses of treatment are administered concurrently to a patient, *i.e.*, individual doses of the second agent are administered separately yet within a time interval such that the compound provided herein can work together with the second active agent. For example, one component can be administered once per week in combination with the other components that can be administered once every two weeks or once every three weeks. In other words, the dosing regimens are carried out concurrently even if the therapeutics are not administered simultaneously or during the same day.

[00411] The second agent can act additively or synergistically with the compound provided herein. In one embodiment, the compound provided herein is administered concurrently with one or more second agents in the same pharmaceutical composition. In another embodiment, a compound provided herein is administered concurrently with one or more second agents in separate pharmaceutical compositions. In still another embodiment, a compound provided herein is administered prior to or subsequent to administration of a second agent. Also contemplated are administration of a compound provided herein and a second agent by the same or different routes of administration, *e.g.*, oral and parenteral. In certain embodiments, when the compound provided herein is administered concurrently with a second agent that potentially produces adverse side effects including, but not limited to, toxicity, the second active agent can advantageously be administered at a dose that falls below the threshold that the adverse side effect is elicited.

Kits

[00412] Also provided are kits for use in methods of treatment of a liver disorder such as HCV and/or HBV infections. The kits can include a compound or composition provided herein, a second agent or composition, and instructions providing information to a health care provider regarding usage for treating the disorder. Instructions may be provided in printed form or in the form of an electronic medium such as a floppy disc, CD, or DVD, or in the form of a website address where such instructions may be obtained. A unit dose of a compound or composition provided herein, or a second agent or composition, can include a dosage such that when administered to a subject, a therapeutically or prophylactically effective plasma level of the compound or composition can be maintained in the subject for at least 1

days. In some embodiments, a compound or composition can be included as a sterile aqueous pharmaceutical composition or dry powder (*e.g.*, lyophilized) composition.


[00413] In some embodiments, suitable packaging is provided. As used herein, “packaging” includes a solid matrix or material customarily used in a system and capable of holding within fixed limits a compound provided herein and/or a second agent suitable for administration to a subject. Such materials include glass and plastic (*e.g.*, polyethylene, polypropylene, and polycarbonate) bottles, vials, paper, plastic, and plastic-foil laminated envelopes and the like. If e-beam sterilization techniques are employed, the packaging should have sufficiently low density to permit sterilization of the contents.

[00414] The following Examples illustrate the synthesis of representative compounds provided herein. These examples are not intended, nor are they to be construed, as limiting the scope of the claimed subject matter. It will be clear that the scope of claimed subject matter may be practiced otherwise than as particularly described herein. Numerous modifications and variations of the subject matter are possible in view of the teachings herein and, therefore, are within the scope the claimed subject matter.

EXAMPLES


Example 1

Preparation of A550 (NM204), the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of L-2',3'-dideoxyadenosine L-ddA




NM204, A550

SYNTHETIC SCHEME:




Synthesis of carboxylic acid 2:



[00415] 2,2-Dimethyl-3-hydroxypropanoic acid methyl ester (965 μ L, 7.57 mmol) was added dropwise to a stirring solution of 4,4'-dimethoxytrityl chloride (2.82 g, 8.33 mmol) in anhydrous pyridine (7.6 mL) at room temperature. The reaction

mixture turned to a red solution quickly, then to an orange suspension (ca. 30 min), and this was left stirring overnight. The mixture was poured carefully over saturated aqueous NaHCO_3 solution (30 mL) and the product was extracted with Et_2O (3×20 mL). The combined organic extracts were washed with brine (20 mL), dried (Na_2SO_4) and the volatiles were removed under reduced pressure. The resulting oil was co-evaporated with toluene and the residue was quickly purified by flash column chromatography (SiO_2 , $\varnothing = 4$ cm, $H = 20$ cm) eluting with $5 \rightarrow 10 \rightarrow 20 \rightarrow 30\%$ Et_2O in petroleum ether (40-60). Evaporation of the fractions ($R_f = 0.25$, 30% Et_2O in petroleum ether (40-60)) afforded ether **1** as a yellow oil (3.11 g, 95%). This compound (3.00 g, 6.91 mmol) was dissolved in THF (35 mL) and an aqueous solution of NaOH (10%, 3.5 g in 35 mL H_2O) was then added at room temperature. The solution turned instantly dark orange and this was stirred for 2 days. The medium was then carefully neutralized by dropwise addition of HCl (1M). The product was extracted with Et_2O (4×50 mL) and the combined organic extracts were washed with brine (50 mL), dried (Na_2SO_4) and the volatiles were removed under reduced pressure. The crude yellow oil was quickly purified by flash column chromatography (SiO_2 , $\varnothing = 2$ cm, $H = 10$ cm) eluting with 50% Et_2O in petroleum ether (40-60). Evaporation of the fractions afforded carboxylic acid **2** as a white foam (2.23 g, 77%). $R_f = 0.50$ (50% Et_2O in petroleum ether (40-60)); $^1\text{H-NMR}$ (300 MHz, CDCl_3) 1.10 (s, 6H, $2 \times \text{CH}_3$), 3.06 (s, 2H, CH_2O), 3.65 (s, 6H, $2 \times \text{OCH}_3$), 6.62-6.79 (m, 4H, PhCH), 7.02-7.46 (stack, 8H, PhCH); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) 22.6 ($2 \times \text{CH}_3$), 43.5 ($\text{C}(\text{CH}_3)_2$), 55.1 ($2 \times \text{OCH}_3$), 85.9 (CPh_3), [125.3, 126.7, 127.7, 128.2, 129.1, 130.0, 136.0, 144.9, 158.4 (Ph), some overlap], 182.2 (C=O).


Synthesis of thioester **3**:



[00416] 1,1'-carbonyldiimidazole (830 mg, 5.12 mmol) was added to a stirring solution of carboxylic acid **2** in anhydrous PhMe/DMF (2/1, v/v, 2.7 mL) at room

temperature and the reaction mixture turned turbid instantly. After 30 min, the medium was diluted by adding anhydrous PhMe/DMF (93/7, v/v, 17 mL) and cooled to -10 °C. 2-Mercaptoethanol (359 μ L, 5.12 mmol) was then added dropwise and the solution was stirred for 1 h at this temperature. The reaction mixture was diluted with H₂O (60 mL) and the product was extracted with Et₂O (3 $\times$ 15 mL). The combined organic extracts were washed with brine (15 mL), dried (Na₂SO₄) and the volatiles were removed under reduced pressure (bath temperature not exceeding 20 °C). The residue was purified by flash column chromatography (SiO₂, $\varnothing$ = 4 cm, H = 15 cm, 1% Et₃N) eluting with 60 $\rightarrow$ 70% Et₂O in petroleum ether (40-60). Evaporation of the fractions afforded thioester **3** as a white syrup (1.74 g, 92%) that solidified upon storage at 4 °C. R_f = 0.35 (70% Et₂O in petroleum ether (40-60)); ¹H-NMR (300 MHz, CDCl₃) 1.16 (s, 6H, 2 $\times$ CH₃), 3.02 (t, J 6.0, 2H, CH₂S), 3.09 (s, 2H, CH₂O), 3.66 (t, J 6.0, 2H, CH₂OH), 3.72 (s, 6H, 2 $\times$ OCH₃), 6.74-6.78 (m, 4H, PhCH), 7.09-7.36 (stack, 8H, PhCH); ¹³C-NMR (75 MHz, CDCl₃) 22.9 (CH₃, 2 $\times$ CH₃), 31.7 (CH₂, CH₂S), 51.0 (quat. C, C(CH₃)₂), 55.2 (CH₃, 2 $\times$ OCH₃), 61.9 (CH₂, CH₂OH), 70.0 (CH₂, CH₂O), 85.8 (quat. C., CPh₃), [113.0 (CH, Ph), 126.7 (CH, Ph), 127.7 (CH, Ph), 128.2 (CH, Ph), 130.1 (CH, Ph), some overlap], [135.9 (quat. C, Ph), 144.8 (quat. C, Ph), 158.4 (quat. C, Ph), some overlap], 205.0 (quat. C, C=O).


Synthesis of H-phosphonate monoester **4**:



[00417] β -L-ddA (1.00 g, 4.25 mmol) was co-evaporated with anhydrous pyridine (3 $\times$ 10 mL) and then dissolved in anhydrous pyridine/DMF (1/1, v/v, 21 mL). Diphenyl phosphite (5.76 mL, 29.8 mmol) was then added dropwise to this solution at room temperature. The reaction mixture was stirred for 20 min upon which a mixture of Et₃N/H₂O (1/1, v/v, 8.5 mL) was added dropwise, and stirring was pursued for an additional 20 min. The reaction mixture was concentrated under reduced pressure to approximately 15-20 mL and this residue was directly purified by flash column chromatography (SiO₂, $\varnothing$ = 4 cm, H = 15 cm, 1% Et₃N) eluting slowly with CH₂Cl₂

(150 mL) then 5% (200 mL) → 10% (200 mL) → 15% (300 mL) MeOH in CH₂Cl₂. Evaporation of the fractions afforded H-phosphonate monoester **4** as a white foam (1.36 g, 80%) that could be kept for several weeks at 4 °C. *R*_f = 0.10 (Et₃N/MeOH/CH₂Cl₂, 1/10/89); ¹H-NMR (300 MHz, CDCl₃) 1.21 (t, *J* 7.4, 9H, 3 × NCH₂CH₃), 1.92-2.50 (stack, 4H, 2 × 2'-H, 2 × 3'-H), 3.02 (q, *J* 7.4, 6H, 3 × NCH₂CH₃), [3.96-4.03 and 4.18-4.30 (stacks, 3H, 4'-H, 2 × 5'-H), 6.28 (m, 1'-H), 6.91 (d, *J* 623, 1H, *P-H*), 7.05 (br s, 2H, NH₂), 8.21 (s, 1H), 8.54 (br s, 1H, OH), 8.57 (s, 1H).


Synthesis of phosphoramidate diester **5**



[00418] H-Phosphonate monoester **4** (1.03 g, 2.57 mmol) and alcohol **3** (1.66 g, 3.45 mmol) were co-evaporated with anhydrous pyridine (3 × 5 mL) and then dissolved in anhydrous pyridine (5 mL). PyBOP (1*H*-benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate, 1.60 g, 3.08 mmol) was then added in one portion and the reaction mixture was stirred for 15 min at room temperature. The solution was poured over saturated aqueous NaHCO₃ solution (30 mL) and the product was extracted with CH₂Cl₂ (4 × 15 mL). The combined organic extracts were washed with brine (10 mL), dried (Na₂SO₄) and concentrated under reduced pressure to leave the corresponding H-phosphonate diester as a slightly yellow oil (1.84 g, assuming 2.41 mmol). This was co-evaporated with anhydrous pyridine (3 × 5 mL; note: do not evaporate to dryness in order to help further solubilization), and the residue was dissolved in anhydrous CCl₄ (24 mL). Benzylamine (791 μL, 7.23 mmol) was added dropwise and the reaction mixture turned cloudy instantly (slight heat development was observed). The milky solution was stirred for 1 h at room temperature and poured over saturated aqueous NaHCO₃ solution (30 mL) and the product was extracted with CH₂Cl₂ (4 × 15 mL). The combined organic extracts were washed with brine (15 mL), dried (Na₂SO₄) and concentrated under reduced pressure to afford phosphoramidate diester **5** as a yellow

oil (2.00 g, assuming 2.31 mmol). This was used in the next step without any further purification. $R_f = 0.29$ (4% MeOH in CH_2Cl_2); $^1\text{H-NMR}$ (300 MHz, CDCl_3) 1.11 (s, 6H, $2 \times \text{CH}_3$), 1.91-2.05 (m, 2H), 2.31-2.59 (m, 2H), 3.06 (m, 2H, CH_2S), 3.08 (s, 2H, CH_2ODMTTr), 3.69 (s, 6H, $2 \times \text{OCH}_3$), 3.83-4.28 (stacks, 7H, CH_2O , NCH_2Ph , $4'\text{-H}$, $2 \times 5'\text{-H}$), 5.71 (br s, 1H, NH), 6.18 (m, 1H, $1'\text{-H}$), 6.69-6.80 (m, 4H, PhCH), 7.02-7.31 (stack, 13H, PhCH), 7.90 (s, 1H), 8.01 (s, 1H), 8.23 (s, 2H, NH_2); $^{13}\text{P-NMR}$ (61 MHz, CDCl_3) 8.82, 8.99.

Synthesis of NM204 (A550), the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of L-ddA:




[00419] Crude phosphoramidate diester **5** (2.00 g, assuming 2.31 mmol) was dissolved in dioxane/AcOH/ H_2O (25/17/25, v/v/v, 462 mL) and the solution was stirred for 3 d at room temperature. Evaporation of the volatiles under reduced pressure left a residue that was purified by flash column chromatography (SiO_2 , $\varnothing = 3$ cm, $H = 15$ cm) eluting with CH_2Cl_2 (100 mL) then 2% (100 mL) $\rightarrow$ 4% (100 mL) $\rightarrow$ 6% (100 mL) $\rightarrow$ 8% (150 mL) MeOH in CH_2Cl_2 . Evaporation of the fractions left NM 204 as a white foam that was dissolved in MeCN (5 mL). Upon addition of H_2O (5 mL), the solution turned turbid and required sonication before lyophilization. The resulting white powder was dried at room temperature (using P_2O_5 as a desiccant) under vacuum for 1 d. The title compound was obtained as a highly hygroscopic white powder (1:1 mixture of diastereoisomers as judged by $^{31}\text{P-NMR}$; 499 mg, 35% over 3 steps). $[\alpha]_D^{20} = +4.2^\circ$ (c 1.0, CHCl_3); $R_f = 0.29$ (4% MeOH in CH_2Cl_2); $^1\text{H-NMR}$ (300 MHz, $\text{DMSO-}d_6$) 1.10 (s, 6H, $2 \times \text{CH}_3$), 2.02-2.14 (m, 2H, $2 \times 3'\text{-H}$), 2.41-2.55 (m, 2H, $2 \times 2'\text{-H}$), 3.01 (t, J 6.4, 2H, CH_2S), 3.43 (d, J 5.0, 2H, CH_2OH), 3.75-4.07 and 4.18-4.29 (stacks, 7H, CH_2O , NCH_2Ph , $4'\text{-H}$, $2 \times 5'\text{-H}$), 5.02 (t, J 5.0, 1H, OH), 5.62 (m, 1H, NH), 6.25 (t, J 5.1, 1H, $1'\text{-H}$), 7.16-7.36 (stack, 7H, PhH , NH_2), 8.14 (s, 1H), 8.26 (s, 1H); $^{13}\text{C-NMR}$ (75 MHz, $\text{DMSO-}d_6$) 21.8 ($2 \times \text{CH}_3$), 25.9 and 26.0 (CH_2 , $3'\text{-C}$), 28.2 and 28.3 (CH_2 , CH_2S), 30.9 and 31.0 (CH_2 , $2'\text{-C}$), 44.2

(CH₂, NCH₂Ph), 51.7 (quat. C, C(CH₃)₂), 63.7 and 63.8 (CH₂, CH₂O), 66.8 (CH₂, m, 5'-C), 68.3 (CH₂, CH₂OH), 78.9 (CH, m, 4'-C), 84.2 (CH, 1'-C), 118.9 (quat. C), [126.5 (CH, Ph), 127.2 (CH, Ph), 128.1 (CH, Ph), some overlap], 138.8 and 138.9 (CH), 140.5 and 140.6 (quat. C), 148.9 (quat. C), 152.3 (CH), 155.0 (quat. C), 204.0 (quat. C, C=O); ¹³P-NMR (61 MHz, DMSO-*d*₆) 9.86, 9.95; *m/z* (FAB⁺) 563 (2), 306 (76), 153 (100); HRMS 565.2034 ([M+H]⁺. C₂₄H₃₄O₆N₆PS requires 565.1998); HPLC *t_R* = 3.52 min (20% TEAC 20 mM in MeCN); UV (EtOH 95%) λ_{max} = 259 (ε_{max} 15900), λ_{min} = 224 (ε_{min} 7200).

Example 2


Preparation of B102, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-*C*-methylcytidine:



B102

PROCEDURE A:

Synthesis of H-phosphonate monoester **5**



Synthesis of carboxylic acid **3**:

[00420] To a stirred solution of 2,2-dimethyl-3-hydroxypropanoic acid methyl ester (**1**, 15 ml, 117.6 mmol) in a mixture of anhydrous methylene chloride (590 ml) and triethylamine (23 ml), were added triphenylmethylene chloride (1.2 eq, 39.3 g) and 4-dimethylaminopyridine (0.1 eq, 1.44 g). The reaction mixture was left refluxing

overnight. The mixture was poured carefully over a saturated aqueous NaHCO_3 solution and the product was extracted with methylene chloride and washed with water. The combined organic extracts were evaporated under reduced pressure to give crude compound **2** which will be used for the next step without further purification. The resulting oil was dissolved in a mixture of dioxan (350 ml) and an aqueous solution of NaOH (30%, 350 ml). The heterogeneous mixture was refluxed for 16 hours. The reaction mixture was allowed to cool down to room temperature, the two phases were separated, and the organic phase carefully neutralized by dropwise addition of HCl (1M). The product was extracted with methylene chloride and the organic phases were evaporated under reduced pressure. The crude orange oil was recrystallized from methylene chloride to afford carboxylic acid **3** as white crystals (92%). $R_f = 0.50$ (70% diethyl ether in petroleum ether); $^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.24 (s, 6H, $2 \times \text{CH}_3$), 3.19 (s, 2H, CH_2O), 7.2-7.5 (m, 15H, C_6H_5).

Synthesis of H-phosphonate monoester **5**:

[00421] 1,1'-carbonyldiimidazole (1.3 eq, 1.17 g) was added to a stirring solution of carboxylic acid **3** (2 g, 5.56 mmol) in an anhydrous mixture of toluene and dimethylformamide (2/1, v/v, 4.5 ml) at room temperature, and the reaction mixture turned turbid instantly. After 30 min, the reaction mixture was diluted with a mixture of toluene and dimethylformamide (93/7, v/v, 28 ml), cooled to -10°C , and 2-mercaptoethanol (1.3 eq, 500 μL) was added. The solution was stirred for 3 h at this temperature. The volatiles were removed under reduced pressure (bath temperature not exceeding 25°C). The residue was dissolved in methylene chloride and washed with water. The organic phases were combined, dried over sodium sulphate (Na_2SO_4), filtered and evaporated to dryness to give compound **4** as a yellow oil. This compound will be coevaporated with anhydrous pyridine and used for the next step without further purification. $R_f = 0.71$ (70% Et_2O in petroleum ether); $^1\text{H-NMR}$ (400 MHz, CDCl_3) 1.20 (s, 6H, $2 \times \text{CH}_3$), 3.05 (t, $J = 6.4$ Hz, 2H, CH_2S), 3.15 (s, 2H, CH_2OTr), 3.69 (t, $J = 6.4$ Hz, 2H, CH_2OH), 7.3-7.9 (m, 15H, C_6H_5).

[00422] Phosphorus acid (10 eq, 4.1 g) was coevaporated two times with anhydrous pyridine, dissolved in that solvent (25 ml) and added to crude **4**. The reaction mixture was stirred at room temperature and a white precipitate appeared after few minutes. The reaction mixture was cooled down to 0°C and pivaloyl chloride (5.5 eq, 3.4 ml) was added. The reaction mixture was allowed to warm to


room temperature and stirred for 3h. The reaction was stopped by addition of a solution of triethylammonium bicarbonate (TEAB 1M, 10 ml) and diluted with ethyl acetate (EtOAc). After extraction with EtOAc and TEAB 0.5M, the organic phases were combined, dried over sodium sulphate, filtered and evaporated to dryness (bath temperature not exceeding 30 °C). The residue was purified by flash column chromatography eluting with 10% of methanol in methylene chloride + 1% triethylamine. Evaporation of the fractions afforded the H-phosphonate monoester **5** as a white syrup (90%). $R_f = 0.25$ (70% Et₂O in petroleum ether); ¹H-NMR (400 MHz, CDCl₃) 1.17 (m, 2 × CH₃ + excess (CH₃CH₂)₃N), 2.9 (m, excess (CH₃CH₂)₃N), 3.12 (t, $J = 6.8$ Hz, 2H, CH₂S), 3.37 (s, 2H, CH₂OTr), 3.90 (m, 2H, CH₂OP), 7.2-7.6 (m, 15H, C₆H₅), 9.9 (m, excess (CH₃CH₂)₃NH); ³¹P-NMR (161 MHz, CDCl₃) 3.85 (s).

Synthesis of B102, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-*C*-methylcytidine:

The following two strategies were used:

Strategy a

Synthesis of the protected nucleoside **7**




[00423] A mixture of 2'-*C*-methylcytidine (NM107) (10 g, 39.0 mmol), triethyl orthoformate (8.3 eq, 54 ml) and *p*-toluenesulfonic acid monohydrate (1 eq, 7.4 g) in anhydrous acetone (650 ml), was refluxed overnight under nitrogen atmosphere. The reaction mixture was neutralized with an aqueous ammonia solution (26%) and the precipitate filtered. The filtrate was evaporated under reduced pressure and coevaporated with ethanol. Purification of the crude mixture on silica gel column chromatography (eluant: stepwise gradient [0-10%] of methanol in methylene chloride) led to compound **6** as a pale-yellow solid (86%). $R_f = 0.30$ (20% MeOH in methylene chloride), ¹H-NMR (400 MHz, DMSO-*d*₆) 1.06 (s, 3H, CH₃), 1.33 (s, 3H, CH₃), 1.47 (s, 3H, CH₃), 3.6 (m, 2H, *H*-5', *H*-5''), 4.1 (m, 1H, *H*-4'), 4.41 (d, 1H, *H*-3', $J = 3.2$ Hz), 5.16 (t, 1H, *OH*-5', $J = 4.0$ Hz, *D*₂O exchangeable), 5.69 (d, 1H, *H*-5,

$J = 8.0$ Hz), 6.04 (s, 1H, $H-1'$), 7.14-7.19 (bd, 2H, NH_2 , D_2O exchangeable), 7.74 (d, 1H, $H-6$, $J = 8.0$ Hz); LC/MS Scan ES- 296 (M-H)⁻, Scan ES+ 298 (M+H)⁺, $\lambda_{max} = 280.7$ nm.

[00424] Compound **6** (4.4 g, 14.8 mmol) was dissolved in anhydrous pyridine (74 ml) and chlorotrimethylsilane (3 eq, 5.4 ml) was added. The reaction mixture was stirred at room temperature under nitrogen atmosphere for 2h, then 4,4'-dimethoxytrityl chloride (1.5 eq, 7.5 g) and 4-dimethylaminopyridine (0.5 eq, 900 mg) were successively added. The reaction mixture was stirred overnight at room temperature, then quenched with a saturated aqueous $NaHCO_3$ solution. The crude product was extracted with methylene chloride, washed with saturated aq $NaHCO_3$ solution, and water. The combined organic phases were concentrated under reduced pressure, then dissolved in a mixture of dioxan (160 ml) and aqueous ammonia (28%, 29 ml). The solution was heated at 70°C for 3h and evaporated to dryness. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient of methanol [1-5%] in methylene chloride) to give protected nucleoside **7** as a yellow solid (81%). $R_f = 0.16$ (30% EtOAc in CH_2Cl_2) ^1H -NMR (400 MHz, $DMSO-d_6$) 1.03 (s, 3H, CH_3), 1.30 (s, 3H, CH_3), 1.42 (s, 3H, CH_3), 3.5 (m, 2H, $H-5'$, $H-5''$), 3.71 (s, 6H, 2 x OCH_3), 4.0 (d, 1H, $H-4'$, $J = 3.2$ Hz), 4.36 (d, 1H, $H-3'$, $J = 2.8$ Hz), 5.1 (m, 1H, $OH-5'$, D_2O exchangeable), 5.90 (s, 1H, $H-1'$), 6.2 (m, 1H, $H-5$), 6.8-7.2 (m, 13H, DMTr), 7.6 (m, 1H, $H-6$), 8.32 (s, 1H, NH , D_2O exchangeable); LC/MS Scan ES- 598 (M-H)⁻, $\lambda_{max1} = 231.7$ nm, $\lambda_{max2} = 283.7$ nm.

Synthesis B102 (compound 10)



[00425] Compounds **7** (2.0 g, 3.34 mmol) and **5** (2.2 eq, 4.3 g) were coevaporated together with anhydrous pyridine and dissolved in this solvent (50 ml). Pivaloyl chloride (2.5 eq, 1 ml) was added dropwise and the solution stirred at room temperature for 2h30. The reaction mixture was diluted with methylene chloride and neutralized with an aqueous solution of ammonium chloride (NH₄Cl 0.5M). After extraction with methylene chloride / aq NH₄Cl 0.5M, the organic phases were combined, evaporated under reduced pressure (bath temperature not exceeding 30 °C) and coevaporated with toluene. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride + 2% acetic acid) to afford the desired product **8** which was coevaporated with toluene to give a beige foam (94%). *R_f* = 0.63 (5% MeOH in CH₂Cl₂); ¹H-NMR (400 MHz, CDCl₃) 1.21 (m, 9H, 3 CH₃), 1.42 (s, 3H, CH₃), 1.60 (s, 3H, CH₃), 3.13 (m, 2H, CH₂S), 3.17 (m, 2H, CH₂OTr), 3.79 (s, 6H, 2 x OCH₃), 4.1 (m, 2H, CH₂OP), 4.2-4.3 (m, 3H, *H*-5', *H*-5'', *H*-4'), 5.09 (d, 1H, *H*-3', *J* = 7.6 Hz), 5.89 (d, 1H, *H*-5, *J* = 5.6 Hz), 6.0 (m, 1H, *H*-1'), 6.8-7.7 (m, 29 H, Tr, DMTr, *H*-6); ¹³P-NMR (161 MHz, CDCl₃) 7.92, 8.55; LC/MS Scan ES+ 1066 (M+H)⁺, Scan ES- 1064 (M-H)⁻.


[00426] a solution of compound **8** (3.4 g, 3.15 mmol) in anhydrous carbon tetrachloride (30 ml) was added dropwise benzylamine (10 eq, 3.4 ml). The reaction mixture was stirred at room temperature for 1h30. A white precipitate appeared. The solution was diluted with methylene chloride and neutralized with an aqueous solution of hydrogen chloride (HCl 1M). After successive extractions with CH₂Cl₂ / HCl 1M and CH₂Cl₂ / aq NaHCO₃, the organic phases were combined, dried over Na₂SO₄, filtered and evaporated to dryness. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride) to give **9** as a yellow foam (87%). *R_f* = 0.35 (5% MeOH in methylene chloride); ¹H-NMR (400 MHz, CDCl₃) 1.1-1.2 (m, 9H, 3 CH₃), 1.40 (s, 3H, CH₃), 1.59 (s, 3H, CH₃), 2.9-3.2 (m, 4H, CH₂OTr, CH₂OS), 3.76 (s, 6H, 2 x OCH₃), 3.9-4.4 (m, 8H, CH₂OP, CH₂N, *H*-3', *H*-4', *H*-5', *H*-5''), 5.0 (m, 1H, *H*-5), 6.0 (2s, 1H, *H*-1'), 6.7-7.7 (m, 34 H, Tr, DMTr, C₆H₅CH₂, *H*-6); ¹³P-NMR (161 MHz, CDCl₃) 8.40, 8.8.68; LC/MS Scan ES+ 1171 (M+H)⁺.

[00427] Finally, compound **9** (2.39g, 2.04 mmol) was dissolved in a mixture of methylene chloride (10 ml) and an aqueous solution of trifluoroacetic acid (90%, 10 ml). The reaction mixture was stirred at 35-40°C for 2h, then diluted with ethanol

(140 ml). The volatiles were evaporated under reduced pressure and coevaporated with ethanol. The crude mixture was purified by silica gel column chromatography (eluant: stepwise gradient of methanol [0-30%] in methylene chloride), followed by a purification on reverse phase chromatography (eluant: stepwise gradient of acetonitrile [0-50%] in water), to give the desired product **10** (B102) (1:1 mixture of diastereoisomers as judged by ^{31}P -NMR, 36%) which was lyophilized from a mixture of dioxan / water. $R_f = 0.34$ (15% MeOH in methylene chloride); ^1H -NMR (400 MHz, $\text{DMSO}-d_6$) 0.92 (s, 3H, CH_3), 1.10 (s, 6H, $2 \times \text{CH}_3$), 3.0 (m, 2H, CH_2S), 3.33 (m, 1H, $H-3'$), 3.56 (s, 2H, CH_2OH), 3.8-4.0 and 4.05-4.25 (stacks, 7H, CH_2OP , NCH_2Ph , $H-4'$, $H-5'$ and $H-5''$), 4.9 (m, 1H, $\text{OH}-3'$, $J = 5.4$ Hz, D_2O exchangeable), 5.07 (s, 1H, $\text{OH}-2'$, D_2O exchangeable), 5.3 (m, 1H, CH_2OH , D_2O exchangeable), 5.6-5.7 (m, 2H, $H-5$ and NH , D_2O exchangeable), 5.91 (s, 1H, $H-1'$), 7.3-7.4 (stack, 7H, PhH , NH_2 , D_2O exchangeable), 7.6 (m, 1H, $H-6$); ^{13}P -NMR (161 MHz, $\text{DMSO}-d_6$) 9.71, 9.91; HPLC $t_R = 4.67$ min (0-100% acetonitrile over a period of 8 min), $\lambda_{\text{max}} = 274.9$; LC/MS Scan ES+ 587 ($\text{M}+\text{H}$) $^+$.

Strategy b:


Synthesis of protected nucleoside **11**



[00428] NM107 (10 g, 38.87 mmol) was dissolved in anhydrous pyridine (194 ml) and chlorotrimethylsilane (4.5 eq, 21.6 ml) was added. The reaction mixture was stirred at room temperature under nitrogen atmosphere for 2h30, then 4,4'-dimethoxytrityl chloride (1.5 eq, 19.8 g) and 4-dimethylaminopyridine (0.5 eq, 2.37 g) were successively added. The reaction mixture was stirred overnight at room temperature, then quenched with a saturated aqueous NaHCO_3 solution. The crude product was extracted with methylene chloride, washed with saturated aq NaHCO_3 solution, and water. The combined organic phases were concentrated under reduced pressure, then dissolved in tetrahydrofuran (110 ml). To that solution was added tetrabutylammonium fluoride 1M in THF (1 eq, 38.87 ml) and the reaction mixture was stirred for 30 min at room temperature. After extraction with EtOAc and water,

the organic phases were collected and evaporated to dryness. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient of methanol [0-10%] in methylene chloride) to give protected nucleoside **11** as a yellow solid (93%). $R_f = 0.32$ (10% MeOH in CH_2Cl_2) $^1\text{H-NMR}$ (400 MHz, $\text{DMSO}-d_6$) 0.79 (s, 3H, CH_3), 3.56 (m, 2H, $H-5'$, $H-5''$), 3.71 (s, 7H, 2 x OCH_3 , $H-4'$), 5.0 (m, 4H, $H-3'$, $\text{OH}-2'$, $\text{OH}-3'$, $\text{OH}-5'$, D_2O exchangeable), 5.72 (s, 1H, $H-1'$), 6.16 (m, 1H, $H-5$), 6.8-7.2 (m, 13H, DMTr), 7.82 (m, 1H, $H-6$), 8.24 (m, 1H, NH D_2O exchangeable); LC/MS Scan ES- 560 ($\text{M}+\text{H}^+$), ES- 558 ($\text{M}-\text{H}^-$), $\lambda_{\text{max}} = 284.7 \text{ nm}$.

Synthesis of protected phosphoramidate pronucleotide **13**, precursor of **10**




[00429] Compound **11** (7 g, 12.5 mmol) and **5** (1.5 eq, 11.0 g) were coevaporated together with anhydrous pyridine and dissolved in this solvent (187 ml). Pivaloyl chloride (2.0 eq, 3.08 ml) was added dropwise at -15°C and the solution stirred at this temperature for 1h30. The reaction mixture was diluted with methylene chloride and neutralized with an aqueous solution of ammonium chloride (NH_4Cl 0.5M). After extraction with methylene chloride / aq NH_4Cl 0.5M, the organic phases were combined, evaporated under reduced pressure (bath temperature not exceeding 30°C) and coevaporated with toluene. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride + 0.2% acetic acid) to afford the desired product **12** which was coevaporated with toluene to give a white foam (3.5 g, 27%). $R_f = 0.44$ (5% MeOH in CH_2Cl_2); $^1\text{H-NMR}$ (400 MHz, DMSO) 0.8 (m, 3H, CH_3), 1.14 and 1.06 (2s, 6H, 2 CH_3), 3.06 (m, 2H, CH_2S), 3.16 (m, 2H, CH_2OTr), 3.5 (m, 1H, $H-3'$), 3.70 (m, 6H, 2 OCH_3), 3.90 (m, 1H, $H-4'$), 4.03 (m, 2H, CH_2OP), 4.24 (m, 2H, $H-5'$, $H-5''$), 5.30 and 5.04 (2ms, 2H, $\text{OH}-2'$ and $\text{OH}-3'$, D_2O exchangeable), 5.78 (m, 1H, $H-1'$), 5.98 (m, 1H, $\text{P}-\text{H}$), 6.22

(m, 1H, *H*-5), 7.0-7.5 (m, 16 H, Tr), 8.32 (m, 1H, *H*-6); ¹³P-NMR (161 MHz, DMSO) 9.17, 9.65; LC/MS Scan ES+ 1026 (M+H)⁺, λ_{max} = 282.7 nm.

[00430] To a solution of compound **12** (500 mg, 0.49 mmol) in anhydrous carbon tetrachloride (4.9 ml) was added dropwise benzylamine (5 eq, 0.266 ml). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride) to afford compound **13** as a foam (75%). R_f = 0.25 (3% MeOH in methylene chloride); ¹H-NMR (400 MHz, DMSO) 0.79 (s, 3H, CH₃), 1.13 and 1.06 (2s, 6H, 2 CH₃), 3.05 (m, 4H, CH₂OTr, CH₂OS), 3.51 (m, 1H, *H*-3'), 3.69 (s, 6H, 2 x OCH₃), 3.87 (m, 3H, CH₂OP, CH₂N, *H*-3'), 4.08 (m, 2H, *H*-5', *H*-5''), 5.19 and 5.0 (2m, 2H, OH-2' and OH-3', D₂O exchangeable), 5.67 (m, 1H, NH, D₂O exchangeable), 5.75 (2s, 1H, *H*-1'), 6.21 (m, 1H, *H*-5), 6.7-7.5 (m, 34 H, Tr, DMTr, C₆H₅CH₂, *H*-6); ¹³P-NMR (161 MHz, DMSO) 9.84, 9.69; LC/MS Scan ES+ 1132 (M+H)⁺.

[00431] Compound **13** can be converted into the phosphoramidate prodrug **10** (B102) following experimental conditions described for the last step in Examples 3 (Procedure A) and in Example 4.


PROCEDURE B:



SYNTHETIC SCHEME:

[00432] B102 is synthesized as a mixture of phosphorous diastereomers in 1:1 ratio. Isolated overall yield from NM107 to B102 was 31%, as not all the coupled material produced was used for deprotection.

Step 1.1:



Material	Grade	FW gmol ⁻¹	Quantity	Density gml ⁻¹	Amount mol	Eq
NM107	99%	257.2	150g	-	0.583	1
PhB(OH) ₂	98%	122.1	78g	-	0.639	1.10
Pyridine anhydrous	98%	79.1	2.5L	0.978	-	-

[00433] NM107 was dissolved in pyridine under argon and benzenboronic acid was added. The stirred mixture was heated at reflux for 3h under argon. Distillation of the azeotrope was then performed, removing 1.2L (pyridine/water).


T head : 103°C → 113°C T mixture : 112°C → 116°C

The mixture was cooled to room temperature and the pyridine was evaporated under vacuum to get golden oil. The product was stored under vacuum overnight to be used for the next step. A ratio of 97:3 product:starting material was observed by ¹H-NMR (d₆-DMSO). Prior to Step 1.2, the crude was dissolved in 250mL anhydrous pyridine.

Alternatively the following conditions may be used:

- eq benzenboronic acid
- 5 eq pyridine
- 1.5 eq Na₂SO₄
- 5mL CH₃CN for 1g of NM107
- Heat at reflux for 1h-1h30. Cool to RT. Used for next reaction.
- 98-99% of conversion by proton NMR

Step 1.2:




Material	Grade	FW gmol ⁻¹	Quantity	Density gml ⁻¹	Amount mol	Eq
2,3-PhB-NM107	-	343.1	Solution	-	~0.583	1
Phosphonate 3	-	585.7	615g	-	1.049	1.8
EDCI.HCl	98%	191.7	570g	-	2.973	5.1
Acetonitrile anhydrous	98%	-	3L	-	-	-
Benzylamine	98%	107.2	445mL	0.98	4.0	7*
Carbon tetrachloride	98%	153.8	260mL	1.59	2.6	4*

*Extra equivalents of these reagents may be added (e.g. 15eq) if required (if P-OH visible by HPLC)

[00434] Phosphonate 3 was dissolved in 3L of acetonitrile under argon. A solution of 2,3-PhB-NM107 from Step 1.1 was added followed by EDCI.HCl. The mixture was stirred under argon at 41-46 °C for 4h after which time HPLC analysis indicated ~7:1 ratio of P-H product to NM107. The mixture was cooled to 18 °C and benzylamine was added dropwise followed by carbon tetrachloride. The reaction was slightly exothermic. Analysis by HPLC indicated complete conversion of P-H to phosphoramidate product. Ethyl acetate (1L) was added to the mixture which was then acidified to pH 4 with 3L of 20% citric acid. The aqueous phase was extracted with 2.5L of ethyl acetate. The organic phases were combined and washed with 3L of 10% citric acid. The organic phase was basified to pH 8 with 5L of aqueous sodium bicarbonate (saturated) and washed a second time with 2L of aqueous sodium bicarbonate (saturated). The organic phase was dried over sodium sulfate, filtered under vacuum and evaporated to give a yellow foam, 712g.

[00435] The crude residue was dissolved in dichloromethane (1L) and purified on silica plug (2.3Kg of silica). Eluted with : 5L 4% Methanol / DCM, 2*1L 4%, 3*1L 5%, 8*250mL 6%, 4*250mL 7%, 9*1L 7%. Evaporation of the relevant fractions gave 254g (HPLC purity: 98.5%, yield: 52%) and 73 g (HPLC purity: 87.6%, yield: 13%) of phosphoramidate 4.

Step 2 :



Material	Grade	FW Gmol ⁻¹	Quantity	Density gmL ⁻¹	Amount mol	Eq
Phosphoramidate 4	-	828.9	246g	-	0.291	1
AcCl	99%	78.5	62.6mL	1.105	1.049	3.0*
EtOH anhydrous	98%	-	3.5L*	-	-	-

*Subsequently 2.0eq AcCl and 1:10 w/v 4:EtOH ratio were used.

[00436] Phosphoramidate 4 was dissolved in anhydrous ethanol and acetyl chloride was added (exothermic: 18°C to 27°C) to the reaction mixture, under argon. The mixture was stirred at 60°C under argon. After 30min, HPLC analysis indicated complete conversion of the phosphoramidate 4 to deprotected product 5. The mixture was cooled to 25°C and solid sodium bicarbonate (1.04Kg) was added in several portions (foaming, pH ~5.5-6). The mixture was filtered through Celite and washed with two volumes of ethanol. The filtrate was evaporated under vacuum at 35°C. The residue was triturated with TBME (3L) for 1h and then filtered to remove the trityl by-product. The solid obtained was dried under vacuum to give 185g with 93% purity at 254nm by HPLC.

If required, any residual benzenboronic acid may be removed from the product by dissolution in water and treatment with Amberlite IRA-743 resin.

The following alternative reaction conditions (to avoid the possibility of acylating 4) are possible:

- 2.0eq AcCl in EtOH 1:10 v:v to generate HCl and consume all AcCl (exothermic)
- Phosphoramidate 4 in EtOH (to make 1:10 w:v total volume EtOH)
- Add HCl/EtOH solution to reaction mixture at 20°C
- 60°C under argon, 30-45min

The crude was purified by reverse-phase chromatography (1.5Kg of prepared Bakerbond 40µm C-18 RP-silica – washed with 100% acetonitrile gradient to 100% H₂O). The crude was dissolved in acetonitrile (58mL), H₂O (164mL) and saturated aqueous sodium bicarbonate solution (170mL).

Elution under gentle vacuum with a stepwise gradient of 3% MeCN / H₂O, 10%, 15%, 25% (pure product eluted) and evaporation of the relevant fractions gave 106g B102 (62% yield) with 98.6% purity at 254nm by HPLC.

Typical analytical data is shown below:

B102: C₂₄H₃₅N₄O₉PS 586.59gmol⁻¹

HPLC AUC (Method Test 20): 98.9% @ 254nm, Rt 3.34min

m/z (ESI +): 587.12 [M+H]⁺ 100%; 1173.62 [2M+H]⁺ 80%

ν_{max} (KBr disc) (cm⁻¹): 3343.1 br (O-H, N-H); 1647.2 br (C=O base, thioester)

KF: 2.02% H₂O content

Specific Rotation: [α]_D²⁰ +55.011 (c. 10.492mg cm⁻³ in DMSO)

Elemental analysis: Calculated: C 49.14%; H 6.01%; N 9.55%; S 5.47%; P 5.28%;

Found: C 48.74%; H 5.83%; N 9.41%; S 5.81%; P 5.33%

NMR: Analyzed using ¹H, ¹³C, ³¹P, COSY, DEPT, HSQC and HMBC experiments.

¹H NMR δ_H (400 MHz, d6-DMSO): 0.94 (3H, d, *J* 1.8Hz, CH₃), 1.11 (6H, s, (CH₃)₂C), 3.04 (2H, m, *J* 6.4Hz, CH₂S), 3.44 (2H, d, *J* 5.0Hz, CH₂OH), 3.60 (1H, br-m, *H*-3'), 3.82-4.01 (5H, m, *H*-4', CH₂O, CH₂Ph), 4.07-4.12 (1H, m, *H*-5'), 4.13-4.24 (1H, m, *H*-5''), 4.94 (1H, t, *J* 5.0Hz, CH₂OH), 5.07 (1H, d, *J* 1.8Hz, OH-2'), 5.26 (1H, t, *J* 6.8Hz, OH-3'), 5.64-5.76 (1H, m, P-N-H), 5.69, 5.70 (1H, 2 x d, 2 x *J* 7.6Hz, *H*-5), 5.93 (1H, br-s, *H*-1'), 7.13-7.20 (2H, 2 x br-s, NH₂), 7.20-7.25 (1H, m, Ar-H), 7.28-7.35 (4H, m, 4 x Ar-H), 7.53, 7.57 (1H, 2 x d, *J* 7.6Hz, *H*-6)

¹³C NMR δ_C (100 MHz, d6-DMSO): 19.81 (CH₃), 21.79 (C(CH₃)₂), 28.17, 28.24 (CH₂S), 44.18 (PhCH₂), 51.62 (C(CH₃)₂), 63.74, 63.79 (CH₂O), 64.21, 64.51 (C-5'), 68.29 (CH₂OH), 72.41, 72.57 (C-3'), 77.80, 77.85 (C-2'), 79.47, (C-4'), 91.66, (C-1'), 93.82 (C-5), 126.68, 127.09, 128.08, 128.09 (5 x Ar-C),


140.34, 140.38, 140.40 (Ar-C_{ipso}, C-6), 155.12, 165.21 (C-2, C-4), 203.85 (C=OS)


³¹P NMR δ_P (162 MHz, d₆-DMSO): 9.71, 9.91 (1P, 2 x s, ratio 1.00:1.07)

[00437] Synthetic Procedure A can be used for synthesizing nucleoside prodrugs such as B 102. Protection of the 2' and 3' hydroxyl groups as well as the amino group that may be present on the nucleoside base is preferred. In Strategy A, the 2' and 3' hydroxyl groups are protected, e.g., as the acetonide derivative and the amino group is protected, e.g., as the di-methoxytrityl derivative. Hydrolysis of the acetonide after the coupling of the nucleoside with the SATE intermediate is carried out using an acid such as TFA. This hydrolysis procedure can potentially produce by-products and give low yields, and the di-methoxytrityl chloride is disadvantageously expensive. Synthetic Procedure B, below, can overcome the such difficulties. An acid, such as a boronic acid, such as phenyl boronic acid is used to protect the 2' and 3' hydroxyl groups on the sugar moiety. Coupling the nucleoside phenyl boronate derivative with the SATE intermediate can give good yield, and the phenyl boronate deprotection conveniently takes place during the work-up of the reaction mixture, on washing with an acid such as an aqueous citric acid solution. The final removal of a protecting group such as a trityl group (on the Sate moiety) is mildly carried out using an organic solvent system, such as an acetyl chloride/ethanol mixture. This deprotection reaction can be consistently reproducible, scalable and gave significantly high yield.

Example 3

Preparation of B299, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-*C*-methylguanosine





[00438] 2'-C-methylguanosine (NM108) (3 g, 10.10 mmol) and compound **5** [for the synthesis of **5**, See Example 2] (6.48 g, 11.10 mmol) were coevaporated together with anhydrous pyridine and dissolved in this solvent (152 mL). Pivaloyl chloride (2.48 mL, 20.18 mmol) was added dropwise at -15°C and the solution was stirred at the same temperature for 2h. The reaction mixture was diluted with methylene chloride and neutralized with an aqueous solution of ammonium chloride (NH₄Cl 0.5M). After extraction with methylene chloride / aq NH₄Cl 0.5M, the organic phases were combined, dried over Na₂SO₄ evaporated under reduce pressure (bath temperature not exceeding 30°C) and coevaporated twice with toluene. The crude mixture was purified on silica gel flash column chromatography (eluant : stepwise gradient [0-10%] of methanol in methylene chloride + 0.2% acetic acid) to afford the desired product **6** (2.5 g, 32%). *R_f* = 0.34 (15% MeOH in CH₂Cl₂); ¹H-NMR (400 MHz, DMSO-*d*₆) 0.80 (s, 3H, CH₃), 1.13 (s, 6H, 2 x CH₃), 3.04 (m, 2H, CH₂OTr), 3.14 (m, 2H, CH₂S), 3.97-4.08 (m, 4H, *H*-3', *H*-4', CH₂OP), 4.28-4.38 (m, 2H, *H*-5', *H*-5'), 5.10-5.35 (m, 2H, OH-2', OH-3', D₂O exchangeable), 5.77 (s, 1H, *H*-1'), 6.52 (bs, 2H, NH₂, D₂O exchangeable), 7.11-7.42 (m, 15H, Tr), 7.75 (s, 1H, *H*-8), 10.67 (bs, 1H, NH, D₂O exchangeable); ¹³P-NMR (161 MHz, DMSO-*d*₆) 9.47, 9.20; LC/MS Scan ES+ 764 (M+H)⁺, Scan ES- 762 (M-H)⁻.


[00439] To a solution of compound **6** (2.5 g, 3.27 mmol) in anhydrous carbon tetrachloride (33 mL) was added dropwise benzylamine (5 eq, 1.79 mL). The reaction mixture was stirred at room temperature for 1h and evaporated under reduced pressure (bath temperature not exceeding 30°C). The crude mixture was purified on silica gel flash column chromatography (eluant : stepwise gradient [0-10%] of methanol in methylene chloride) to give compound **7** as a white foam (2.9 g, quantitative yield).

R_f = 0.27 (10% MeOH in methylene chloride); ¹H-NMR (400 MHz, DMSO-*d*₆) 0.81 (s, 3H, CH₃), 1.10 (s, 6H, 2 x CH₃), 2.99-3.08 (m, 4H, CH₂OTr, CH₂S), 3.87-4.30 (m, 8H, *H*-3', *H*-4', *H*-5', *H*-5'' CH₂OP, NCH₂Ph), 5.66 (m, 1H, NH, D₂O exchangeable), 5.76 (s, 1H, *H*-1'), 6.60 (bs, 2H, NH₂, D₂O exchangeable), 7.17-7.39 (m, 20H, Tr, C₆H₅CH₂), 7.77 (s, 1H, *H*-8); ¹³P-NMR (161 MHz, DMSO-*d*₆) 9.93, 9.78; LC/MS Scan ES+ 869 (M+H)⁺, Scan ES- 867 (M-H)⁻.

[00440] Compound **7** (2.84 g, 3.27 mmol) was dissolved in a mixture of trifluoroacetic acid (1.1 mL) and methylene chloride (11.4 mL). The reaction mixture was stirred 0.5h at room temperature. The solution was diluted with ethanol, evaporated under reduce pressure (bath temperature not exceeding 30°C) and coevaporated twice with toluene. The crude mixture was purified on silica gel flash column chromatography (eluant : stepwise gradient [0-30%] of methanol in methylene chloride) and then, on reverse phase column chromatography (eluant : stepwise gradient [0-100%] of acetonitrile in water) to give the desired product **8** (B299) (1:1 mixture of diastereoisomers according to ³¹P-NMR, 800 mg, 39%) which was lyophilized from a mixture of dioxan / water. R_f = 0.57 (20% MeOH in methylene chloride); ¹H-NMR (400 MHz, DMSO-*d*₆) 0.82 (s, 3H, CH₃), 1.09 (s, 6H, 2 x CH₃), 3.01 (m, 2H, CH₂S), 3.42 (d, 2H, CH₂OH, *J* = 8.0 Hz), 3.81-4.00 (m, 6H, *H*-3', *H*-4' CH₂OP, NCH₂Ph), 4.11-4.27 (m, 2H, *H*-5', *H*-5''), 4.92 (t, 1H, CH₂OH, *J* = 8.0 Hz, D₂O exchangeable), 5.16 (s, 1H, OH-2', D₂O exchangeable), 5.40 (m, 1H, OH-3', D₂O exchangeable), 5.64 (m, 1H, NH, D₂O exchangeable), 5.75 (s, 1H, *H*-1'), 6.50 (bs, 2H, NH₂, D₂O exchangeable), 7.19-7.32 (m, 5H, PhH), 7.77 (s, 1H, *H*-8), 10.61 (bs, 1H, NH, D₂O exchangeable); ¹³P-NMR (161 MHz, DMSO-*d*₆) 9.91, 9.78; HPLC t_R = 3.67 min (0-100% acetonitrile over a period of 8 min), λ_{max} = 251.3; LC/MS Scan ES+ 627 (M+H)⁺, Scan ES- 625 (M-H)⁻.


Example 4

Preparation of B208, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-*C*-methylthymidine



B208

SYNTHETIC SCHEME:



[00441] 2'-*C*-Methylthymidine (NM105) (700 mg, 2.57 mmol) and **5** [for the synthesis of **5**, See Example 2] (1.1 eq, 1.6 g) were coevaporated together with anhydrous pyridine and dissolved in this solvent (40 ml). Pivaloyl chloride (2.0 eq, 0.633 ml) was added dropwise at -15°C and the solution stirred at this temperature for 1h30. The reaction mixture was diluted with methylene chloride and neutralized with an aqueous solution of ammonium chloride (NH₄Cl 0.5M). After extraction with methylene chloride / aq NH₄Cl 0.5M, the organic phases were combined, evaporated under reduced pressure (bath temperature not exceeding 30 °C) and coevaporated with toluene. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-10%] of methanol in methylene chloride + 0.2% acetic

acid) to afford the desired product **6** which was coevaporated with toluene to give a white foam (942 mg, 50%). $R_f = 0.56$ (15% MeOH in CH_2Cl_2); $^1\text{H-NMR}$ (400 MHz, DMSO) 1.00 (s, 3H, CH_3), 1.13 (s, 6H, 2 CH_3), 1.77 (s, 3H, CH_3), 3.16 (m, 2H, CH_2S), 3.32 (m, 2H, CH_2OTr), 3.6 (m, 1H, $H-3'$), 3.9 (m, 1H, $H-4'$), 4.0 (m, 2H, CH_2OP), 4.2-4.3 (m, 2H, $H-5'$, $H-5''$), 5.21 (s, 1H, OH-2' , D_2O exchangeable), 5.40 (t, 1H, OH-3' , D_2O exchangeable), 5.83 (s, 1H, $H-1'$), 6.0 (s, 1H, $P-H$), 7.0-7.5 (m, 16 H, Tr, $H-6$); $^{13}\text{P-NMR}$ (161 MHz, DMSO) 9.29, 9.68; LC/MS Scan ES+ 761 ($\text{M}+\text{Na}$) $^+$.


[00442] To a solution of compound **6** (920 mg, 1.25 mmol) in anhydrous carbon tetrachloride (13 ml) was added dropwise benzylamine (10 eq, 1.4 ml). The reaction mixture was stirred at room temperature for 2h. A white precipitate appeared. The solution was diluted with methylene chloride and neutralized with an aqueous solution of hydrogen chloride (HCl 1M). After successive extractions with CH_2Cl_2 / HCl 1M and CH_2Cl_2 / aq NaHCO_3 , the organic phases were combined, dried over Na_2SO_4 , filtered and evaporated to dryness. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-10%] of methanol in methylene chloride) to give **7** as a white foam (875 mg, 83%). $R_f = 0.56$ (15% MeOH in CH_2Cl_2); $^1\text{H-NMR}$ (400 MHz, DMSO) 0.99 (s, 3H, CH_3), 1.12 (s, 6H, 2 CH_3), 1.75 (s, 3H, CH_3), 3.04 (m, 4H, CH_2OTr , CH_2S), 3.69 (m, 1H, $H-3'$), 3.8-4.0 (m, 5H, CH_2OP , CH_2N , $H-4'$), 4.0-4.2 (m, 2H, $H-5'$, $H-5''$), 5.17 (s, 1H, OH-2' , D_2O exchangeable), 5.3 (m, 1H, OH-3' , D_2O exchangeable), 5.7 (m, 1H, NH , D_2O exchangeable), 5.82 (s, 1H, $H-1'$), 7.1-7.5 (m, 21 H, Tr, $\text{C}_6\text{H}_5\text{CH}_2$, $H-6$); $^{13}\text{P-NMR}$ (161 MHz, DMSO) 9.95, 9.86; HPLC $t_R = 7.91$ min (0-100% acetonitrile over a period of 8 min), $\lambda_{\text{max}} = 266.7$ nm; LC/MS Scan ES+ 866 ($\text{M}+\text{Na}$) $^+$.

[00443] Finally, compound **7** (860 mg, 1.02 mmol) was dissolved in a mixture of methylene chloride (15 ml) and trifluoroacetic acid (0.51 ml). The reaction mixture was stirred at room temperature for 2h, then diluted with toluene. The volatiles were evaporated under reduced pressure and coevaporated with ethanol. The crude mixture was purified by silica gel column chromatography (eluant: stepwise gradient of methanol [0-10%] in methylene chloride), followed by a purification on reverse phase chromatography (eluant: stepwise gradient of acetonitrile [0-50%] in water), to give the desired product **8** (B208) (257 mg, 42%). $R_f = 0.31$ (10% MeOH in methylene chloride); $^1\text{H-NMR}$ (400 MHz, DMSO- d_6) 0.99 (s, 3H, CH_3), 1.10 (s, 6H, 2 $\times$ CH_3),

1.75 (s, 3H, CH₃), 3.0 (m, 2H, CH₂S), 3.42 (d, 2H, CH₂OH), 3.7 (m, 1H, H-3'), 3.8-4.0 (stack, 5H, CH₂OP, NCH₂Ph, H-4'), 4.0-4.3 (m, 2H, H-5' and H-5''), 4.9 (m, 1H, CH₂OH, D₂O exchangeable), 5.17 (s, 1H, OH-2', D₂O exchangeable), 5.3 (m, 1H, OH-3', D₂O exchangeable), 5.7 (m, 1H, NH, D₂O exchangeable), 5.81 (s, 1H, H-1'), 7.2-7.4 (stack, 6H, PhH, H-6); ¹³P-NMR (161 MHz, DMSO-*d*₆) 9.84, 9.90; HPLC *t*_R = 4.98 min (0-100% acetonitrile over a period of 8 min), λ_{max} = 269.0 nm; LC/MS Scan ES+ 602 (M+H)⁺.

Example 5


Preparation of B261, the Hydroxy-*t*BuSATE *N*-benzylphosphonamidate derivative of PME A



B261

PROCEDURE A:


Synthesis of intermediate 4:



[00444] A 500 mL triple-neck flask fitted with a condenser was charged with PME A (2.00 g, 7.25 mmol), CH₂Cl₂ (121 mL) and DMF (617 μL, 7.98 mmol). The resulting slurry was vigorously stirred and oxalyl chloride (2.21 mL, 25.4 mmol) was added dropwise at 0°C over 10 min (gas evolution). The slurry turned to a yellow solution (10 min) before turning turbid (10 min). This was further stirred for 3 h under

reflux and turned to a white, thick slurry. The products were schlenk-dried 1 h in situ by evaporation of all volatiles under reduced pressure, at room temperature. The resulting yellow solid could then be partially dissolved in CH₂Cl₂ (121 mL), and pyridine (1.17 mL, 14.5 mmol) was added dropwise at 0°C over 10 min. The white suspension turned to a blue solution, which was cooled to -78°C. A solution of alcohol 3 [for the synthesis of 3, See Example 1] (3.480 g, 7.25 mmol) and triethylamine (6.37 mL, 45.7 mmol) in CH₂Cl₂ (72 mL) was then added slowly (ca. 45 min), dropwise along the internal wall, and the reaction was stirred 10 h at -78°C. Benzylamine (2.37 mL, 21.7 mmol) was then added dropwise at -78°C and the solution was left stirring warming to r.t. over 1 h. NaHCO₃ (aq. sat., 200 mL) was poured over the reaction and the layers separated. The aqueous phase was extracted with CH₂Cl₂ (2 x 100 mL) and the combined organic extracts were dried with brine (50 mL) and Na₂SO₄. The solution was filtered and concentrated to afford ca. 6.5 g of crude yellow syrup. Purification by flash column chromatography (SiO₂, Ø = 3.5 cm, H = 11 cm) eluting with 4 → 8% MeOH in CH₂Cl₂ (1% Et₃N) afforded 3.70 g of a yellow foam (0.15 < R_f < 0.30, 10% MeOH in CH₂Cl₂) that were submitted to a second purification by flash column chromatography (SiO₂, Ø = 3.5 cm, H = 12 cm) eluting with 4 → 6% MeOH in CH₂Cl₂ (1% Et₃N) to afford 2.67 g of a yellow foam (0.16 < R_f < 0.25, 10% MeOH in CH₂Cl₂). This was submitted to a third purification by flash column chromatography (SiO₂, Ø = 3.5 cm, H = 12 cm) eluting with 4 → 6% MeOH in CH₂Cl₂ (1% Et₃N) to produce 165 mg of phosphoramidate 4 (ca. 2.7 %) as a white foam and 1.75 g of mixed compounds. These were submitted to a last purification by flash column chromatography (SiO₂, Ø = 3.5 cm, H = 12 cm) eluting with 4 → 6% MeOH in CH₂Cl₂ (1% Et₃N) afforded 353 mg of phosphoramidate 4 (ca. 5.9 %) as a white foam. Total yield : 8.6%. R_f = 0.21 (6% MeOH in CH₂Cl₂); ¹H-NMR (300 MHz, CDCl₃) 1.13 (s, 6H, 2 CH₃), 3.02-3.10 (m, 2H, CH₂S), 3.59 (t, J 7.5, 2H, CH₂), 3.58 (s, 6H, 2 x OCH₃), 3.73 (t, J 7.1, 2H, CH₂), 3.88-4.09 (stacks, 4H, 2 x CH₂), 4.21 (t, J 7.0, 2H, CH₂O), 5.50 (br s, 2H, NH₂), 6.67-6.78 (m, 4H, PhH), 7.04-7.38 (stack, 9H, PhH), 7.72 (s, 1H), 8.22 (s, 1H); ³¹P-NMR (121 MHz, CDCl₃) 25.0; m/z (FAB+) 825 (1), 303 (100); HRMS 825.3171 ([M+H]⁺. C₄₃H₅₀O₇N₆PS requires 825.3199).

Synthesis of compound 5 (B261):




[00445] Dichloroacetic acid (20% solution in CH_2Cl_2 , ca. 140 drops) was added dropwise to a solution of ether 4 (353 mg, 0.43 mmol) in CH_2Cl_2 (4.3 mL) at 0°C and this was stirred for 55 min. Solid NaHCO_3 (ca. 1.5 g) was then added and the slurry was stirred for 10 min before filtration and evaporation. Purification by flash column chromatography (SiO_2 , $\varnothing = 1.5\text{ cm}$, $H = 10\text{ cm}$) eluting with 4% $\rightarrow$ 10% MeOH in CH_2Cl_2 afforded pure phosphonamidate 5 (130 mg after lyophilization in THF/ H_2O and 3 day-stay in P_2O_5 desiccator, 58%). This reaction was also performed on 165 mg of ether 4 to produce 51 mg of phosphonamidate 5 (B261, 49%). $R_f = 0.20$ (10% MeOH in CH_2Cl_2); $^1\text{H-NMR}$ (300 MHz, DMSO-d_6) 1.10 (s, 6H, 2 x CH_3), 2.80 (t, J 7.0, 2H, CH_2S), 3.43 (d, J 5.5, 2H, CH_2OH), 3.69 (A of AB, J 4.8, 1H, 1 x CH_2P), 3.71 (B of AB, J 4.8, 1H, 1 x CH_2P), 3.75-3.88 (stacks, 4H, CH_2O , NCH_2), 3.88-4.07 (m, 2H, NCH_2Ph), 4.30 (t, J 7.0, 2H, CH_2O), 4.97 (t, J 6.1, 1H, OH), 5.31-5.42 (m, 1H, NH), 7.16-7.32 (stack, 7H, PhH, NH_2), 8.09 (s, 1H), 8.13 (s, 1H); $^{13}\text{C-NMR}$ (75 MHz, DMSO-d_6) 21.8 (2 x CH_3), 28.4 and 28.5 (CH_2 , CH_2S), 42.4 (CH_2 , NCH_2), 43.3 (CH_2 , NCH_2), 51.7 (quat. C, $\text{C}(\text{CH}_3)_2$), 61.7 and 61.8 (CH_2 , CH_2O), 64.6 (CH_2 , CH_2O), 68.4 (CH_2 , CH_2O), 118.5 (quat. C), [126.5 (CH, Ph), 127.0 (CH, Ph), 128.0 (CH, Ph), some overlap], 140.5 and 140.6 (quat. C), 141.0 (CH), 149.4 (quat. C), 152.3 (CH), 155.9 (quat. C), 203.9 (quat. C, $\text{C}=\text{O}$); $^{31}\text{P-NMR}$ (121 MHz, DMSO-d_6) 25.9; m/z (FAB+) 161 (32), 256 (42), 523 (100); HRMS 523.1899 ($[\text{M}+\text{H}]^+$. $\text{C}_{22}\text{H}_{32}\text{O}_5\text{N}_6\text{PS}$ requires 523.1892); HPLC (C18, flow: 0.5 mL/min, solution A = TEAC 20 mM, solution B = 20% TEAC 20 mM): $t_R = 5.04\text{ min}$ (60% A in B), $t_R = 27.24\text{ min}$ ($t = 0 \rightarrow 10\text{ min}$: 100% A; $t = 10 \rightarrow 30\text{ min}$: 0 $\rightarrow$ 50% B in A; $t =$

30 → 35 min: 50 → 100% B in A); UV (EtOH 95%) $\lambda_{\text{max}} = 205$ ($\epsilon_{\text{max}} 23900$), $\lambda_{\text{min}} = 228$ ($\epsilon_{\text{min}} 5400$).

PROCEDURE B:

[Improved preparation of the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative (B261, Compound **5) of PME A]**

SYNTHETIC SCHEME:



Step 1: Synthesis of intermediate **A**

[00446] To the suspension of PME A (2 g, 7.3 mmol) in 120 mL of DCM (anhydrous) was added DMF (640 mg, 1.2 eq), followed with oxalyl chloride (2.3 mL, 3.5eq) at room temperature. The mixture was heated to reflux for 1.5 hrs to give a thick yellow suspension. The mixture was concentrated to dryness through rotarvap to give the crude intermediate **2** as pale yellow solid. LCMS analysis of the aliquot of intermediate **2** in methanolic solution confirmed the structure of the product in good purity.

Step 2: Synthesis of intermediate **B**:

[00447] The crude intermediate **A** (7.33 mmol) was suspended into 100 mL of anhydrous DCM. The suspension was cooled to 0 °C. To this was added pyridine (1.2 mL, 14.6 mmol, 2 eq) at 0 °C. After the addition, the pale yellow suspension turned to a golden colored clear solution. This solution was cooled to -32 °C with ACN/dry ice bath. To this was added a solution of **3** (3.52 g, 7.33 mmol, 1eq) in 70 mL of anhydrous DCM that contained triethylamine (6.3 mL, 44 mmol, 6eq) drop-wise. The internal reaction temperature was maintained between -35°C ~ -30 °C during the addition. The bright golden colored solution turned to a green colored solution with some precipitate crashing out from the solution during the addition. The precipitate was presumably the triethylamine HCl salt. It took 20 minutes to complete the addition. After the addition, the mixture was stirred at -30°C ~ -10°C for 1hr. The reaction mixture was cooled back to -20°C. To this was added benzylamine (2.4 mL, 22 mmol, 3eq). The mixture was stirred at -20°C for 10 minutes. To the reaction mixture was added saturated NaHCO₃/H₂O and the mixture was stirred for 2 minutes. The DCM layer was separated, dried with Na₂SO₄ and concentrated to dryness to give the crude intermediate **B** as yellow viscous oil. HPLC analysis of the crude intermediate gave 62% purity at 272 nm.

Step 3: Synthesis of intermediate **4**:


[00448] The crude intermediate **B** (7.33 mmol) as viscous pale yellow oil was dissolved into 200 mL of MeOH. The reaction mixture was refluxed overnight. HPLC analysis of the reaction mixture indicated the complete conversion of the amidine to amine. [The retention time of amidine (RT=5.92 min) is very close to the amine (RT=5.98 min) by the current in-house HPLC method!] The mixture was cooled to RT and filtered. The filtrate was concentrated to dryness by rotar-vap. The obtained crude product was purified through silica gel column chromatography (120g silica gel combiflash column was used, 3-8% of MeOH in DCM as the eluent) to give 3.1 g pure product **4** as white foam in 51% isolated yield from 2g of **PMEA**. ¹H-NMR of the obtained **4** was consistent with the desired structure. HPLC analysis of the obtained **4** gave 96% purity (AUC).

Step 4: Synthesis of B261 (Compound **5**)


[00449] Intermediate **4** (300 mg, 0.36 mmol) was dissolved into EtOH (anhydrous, 5 mL). To this was added acetyl chloride (43 mg, 1.5 eq) in one portion at room temperature. The reaction should be operated in a closed reaction flask to avoid the loss of HCl gas. The reaction mixture was stirred at RT for 30 min. To this was added solid NaHCO₃ and the mixture was stirred for 15 min. pH of the reaction mixture was found to be around 7~8. The mixture was filtered and the filtrate was concentrated to dryness. The crude product was purified through silica gel column chromatography (5-10% MeOH in DCM as the eluent) to give 163 mg of **5** as clear viscous oil in 86% yield. ¹H-NMR of the obtained product was consistent with the desired structure. HPLC analysis of the obtained product gave 97.4% purity (AUC).

Example 6

Preparation of B263, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-*C*-methyladenosine




SYNTHETIC SCHEME:



[00450] The pronucleotide **B263** (94 mg, 6% overall yield) has been synthesized from its parent nucleoside 2'-*C*-methyl-6-*NH*-dimethoxytrityl-adenosine (1.59 g, 2.73 mmol) following a similar procedure than the one described for the synthesis of the pronucleotide prepared in the Example 2 (Procedure A, Strategy b), and isolated as a white lyophilized powder. ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 0.80 (s, 3H), 0.97-0.98 (d, *J* = 4.26 Hz, 6H), 3.02 (m, 2H), 3.34-3.35 (m, 2H), 3.76-3.96 (m, 4H), 4.03-4.05 (m, 2H), 4.15-4.17 (m, 2H), 4.76-4.79 (m, 1H), 5.32 (s, 1H), 5.34-5.36 (m, 1H), 5.45-5.55 (m, 1H), 5.93 (s, 1H), 7.1-7.4 (m, 7H), 8.14 (s, 1H), 8.21 (1H); ³¹P NMR (DMSO-*d*₆, 162 MHz) δ (ppm) 9.75 and 9.86 (2s); Scan ES⁺ 611 (M+H)⁺, λ_{max} = 258 nm; HPLC (0-100% ACN over a period of 8 min) t_R = 4.79 min. λ_{max} = 260.8 nm.

Example 7

Preparation of **B229**, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-*C*-methyluridine



Reagents and Conditions: (i) pTsOH.H₂O, CH(OEt)₃, acetone, r.t.; (ii) PivCl, pyr, r.t.; (iii) Benzylamine, CCl₄, rt; (iv) aq. 90% TFA, DCM, r.t.

[00451] The pronucleotide **6** (446 mg, 0.76 mmol, overall yield 9% over 4 steps) has been synthesized from its nucleoside parent **1** following a similar procedure described for the synthesis of the pronucleotide prepared in Example 2, Strategy A.

B229 **6**.

[00452] ^1H NMR (400 MHz, DMSO- d_6): δ 0.98 (s, 3H, CH₃); 1.10 (s, 6H, 2 x CH₃); 3.03 (m, 2H, CH₂S); 3.41 (m, 2H, CH₂OH, J 5.6 Hz); 3.61 (m, 1H, H-3'); 3.8-4.0 and 4.05-4.25 (stacks, 5H, NCH₂Ph, H-4', H-5' and H-5''); 4.05-4.25 (2 x 1H, 2 x m, CH₂OP); 4.91 (t, 1H, 3'-OH, D₂O exchangeable, J = 5.62 Hz); 5.20 (br-s, 1H, 2'-OH, D₂O exchangeable); 5.39 (a-t, 1H, CH₂OH, D₂O exchangeable, J = 7.32 Hz); 5.52 (m, 1H, H-5); 5.65 (m, 1H, PhNH, D₂O exchangeable); 5.8 (br-s, 1H, H-1'); 7.2-7.32 (m, 5H, ArH); 7.55 (a-dd, 1H, H-6); 11.37 (br-s, 1H, NH, D₂O exchangeable).

^{31}P NMR (161.8 MHz, DMSO- d_6): δ 9.73 and 9.98 (ratio of signals by integration of 52:48)

m/z (ES+) 588.11 (M+H)⁺.

HPLC (Method 20): chemical purity 99.2%, 3.48 mins.


CHN analysis:- Found: C, 49.29, H, 5.95, N, 6.88, P, 5.16; C₂₄H₃₄N₃O₁₀PS requires C, 49.06, H, 5.83, N, 7.15, P, 5.46.

$[\alpha]_D^{23}$ +26.3 (c, 0.571 in H₂O).

ν_{max} (KBr): 3373 (br, NH and OH), 1682 (C=O).


Example 8

Preparation of B186, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-*C*-methylinosine



B186


SYNTHETIC SCHEME:




[00453] The pronucleotide **B186** (314 mg, 8% overall yield) has been synthesized from its parent nucleoside 2',3'-*O*-isopropylidene-2'-*C*-methyl-inosine (2.0 g, 6.26 mmol) following a similar procedure than the one described for the synthesis of the pronucleotide prepared in the Example 2 (Procedure A, Strategy a), and isolated as a white lyophilized powder. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 0.79 (s, 3H), 1.09 (s, 6H), 3.01-3.04 (t, $J=6.53$ Hz, 2H), 3.42 (s, 2H), 3.84-3.91 (m, 2H), 3.94-4.03 (m, 3H), 4.05-4.09 (m, 1H), 4.15-4.26 (m, 2H), 4.92 (s, 1H), 5.36 (s, 1H), 5.43 (t, $J=6.54$ Hz, 1H), 5.62-5.71 (m, 1H), 5.94 (s, 1H), 7.18-7.22 (m, 1H), 7.25-7.30 (m, 4H), 8.08 (s, 1H), 8.10 (s, 1H), 12.15 (brs, 1H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.76-9.90 (2s); Scan $\text{ES}^+ 612(\text{M}+\text{H})^+$, $\lambda_{\text{max}} = 240.7$ nm; HPLC (0-100% ACN over a period of 8 min) $t_{\text{R}} = 4.72$ min $\lambda_{\text{max}} = 243.1$ nm.

Example 9

Preparation of B396, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 9-[2-*C*-methyl- β -ribofuranosyl]-6-chloropurine




SYNTHETIC SCHEME:



[00454] The pronucleotide **B396** (75 mg, 10% overall yield) has been synthesized from its parent nucleoside 9-[2-*C*-methyl- β -ribofuranosyl]-6-chloropurine (571 mg, 1.90 mmol) following a similar procedure than the one described for the synthesis of the pronucleotide prepared in the Example 4 and isolated as a white lyophilized powder. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ (ppm) 0.82 (d, $J=2.63$ Hz, 3H), 1.07 (s, 6H), 3.02 (m, 2H), 3.40-3.41 (q, $J=3.36$ Hz and $J=1.89$ Hz, 2H), 3.85-3.98 (m, 4H), 4.12 (s, 2H), 4.25 (m, 2H), 4.89-4.90 (m, 1H), 5.47 (s, 1H), 5.50 (s, 1H), 5.62-5.70 (m, 1H), 6.10 (d, $J=1.23$ Hz, 1H), 7.17-7.29 (m, 5H), 8.76 (s, 1H), 8.82 (s, 1H); ^{31}P NMR ($\text{DMSO}-d_6$, 162 MHz) δ (ppm) 9.91 and 9.79 (2s); Scan ES^+ 630 ($\text{M}+\text{H}$) $^+$, $\lambda_{\text{max}} = 260$ nm; HPLC (0-100% ACN over a period of 8 min) $t_R=4.42$ min $\lambda_{\text{max}} = 265$ nm.


Example 10

Preparation of B307, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2',3'-*O*-carbonate-2'-*C*-methylguanosine



B307

SYNTHETIC SCHEME:



***N*-Benzylaminyl-2',3'-*O*-carbonate-2'-*C*-methylguanosin-5'-yl-*O*-
(triphenylmethoxy-*tert*-butyl-*S*-acyl-2-thioethyl) phosphate (**C1**):**


[00455] Compound **B2** [See, Compound 7, Example 3, Procedure A] (250 mg, 0.288 mmol) was dissolved in dimethylformamide (3.5 mL) and treated with 1,1-carbonyldiimidazole (186.60 mg, 1.15 mmol). The mixture was stirred at room temperature for 4 h 30 and concentrated under reduced pressure (bath temperature not exceeding 30 °C). The crude residue was subjected to silica gel chromatography, eluting with a gradient 0-10 % methanol in dichloromethane, to give **C1** as a colorless oil. (68 mg, 26%). Compound **C1**: ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 10.80 (1s, 1H, NH), 7.80 (s, 1H, *H*-8), 7.33-7.18 (m, 20H, 4 C_6H_5), 6.66 (sl, 2H, NH_2), 6.30 (s, 1H, *H*-1'), 5.78 (m, 1H, PNH), 5.22 (m, 1H, *H*-3'), 4.47-4.30 (m, 2H, *H*-4' and *H*-5'a), 4.20-4.05 (m, 1H, *H*-5'b), 3.99-3.87 (m, 4H, CH_2O and CH_2N), 3.10-3.03 (m, 4H, CH_2S and CH_2OTr), 1.27 (s, 3H, CH_3), 1.11 (s, 6H, 2 CH_3). ^{31}P NMR (162 MHz, $\text{DMSO-}d_6$) δ 10.42 (s), 10.18 (s). LR LC/MS ($\text{M}+\text{H}^+$) 895.4 (5.57 min). UV: λ_{max} = 253 nm.

***N*-Benzylaminyl-*O*-(hydroxy-*tert*-butyl-*S*-acyl-2-thioethyl)-2',3'-*O*-carbonate-2'-*C*-methylguanosin-5'-yl phosphate B307 (Compound C2):**

[00456] Compound **C1** (65 mg, 0.073 mmol) was dissolved in dichloromethane (260 μ L) and treated with TFA (26 μ L). The mixture was stirred at room temperature for 15 min, then diluted with ethanol, evaporated to dryness (bath temperature not exceeding 30 °C) and coevaporated with toluene. The resulting residue was purified by reverse phase (C 18) silica gel column chromatography eluting with a gradient 0-100% acetonitrile in water and lyophilised from a mixture of water/dioxane to give **B307 (Compound C2)** (34 mg, 72%, white lyophilised powder). **B307 (Compound C2)**: ^1H NMR (400 MHz, DMSO- d_6) δ 10.84 (ls, 1H, NH), 7.80 (s, 1H, *H*-8), 7.32-7.20 (m, 5H, C_6H_5), 6.69 (ls, 2H, NH_2), 6.30 (s, 1H, *H*-1'), 5.77 (m, 1H, PNH), 5.25 (d, 1H, *H*-3', $J_{3',4'} = 20.0$ Hz), 4.92 (ls, 1H, OH), 4.50-4.41 (s, 2H, CH_2OH), 3.03 (t, 2H, CH_2S , $J_{\text{CH}_2\text{S}-\text{CH}_2\text{O}} = 8.0$ Hz), 1.30 (s, 3H, CH_3), 1.10 (s, 3H, CH_3), 1.08 (s, 3H, CH_3). ^{13}C NMR (100 MHz, DMSO- d_6): δ 204.4 (C=O), 154.5 (C-4), 153.1 (C-2), 150.7 (C-6), 140.9 (C_6H_5), 135.6 (C-8), 128.7-127.3 (5C, C_6H_5), 117.0 (C-5), 89.7 (C-1'), 83.7 and 83.6 (2C, C-2' and C-3'), 81.8 (C-4'), 68.8 (CH_2OH), 65.1 (CH_2O), 64.5 (C-5'), 52.2 ($\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$), 44.7 (CH_2N), 28.7 (CH_2S), 22.3 (2C, 2 CH_3), 18.3 (CH_3). ^{31}P NMR (162 MHz, DMSO- d_6) δ 10.39 (s), 10.15 (s). LR LC/MS ($2\text{M}+\text{H}^+$) 1305.4 ($\text{M}+\text{H}^+$) 653.2 ($2\text{M}-\text{H}^-$) 1303.8 ($\text{M}-\text{H}^-$) 651.4 (5.57 min). HRFAB-MS $\text{C}_{26}\text{H}_{34}\text{O}_{10}\text{N}_6\text{PS}$ ($\text{M}+\text{H}^+$) calculated 653.1795, found 653.1819. UV: $\lambda_{\text{max}} = 251$ nm. R_f 0.67 (MeOH/ CH_2Cl_2 , 20/80, v/v).


Example 11

Preparation of B242, the Hydroxy-*t*BuSATE *N*-(4-trifluoromethyl)benzylphosphoramidate derivative of 2'-*C*-methylguanosine



B242

SYNTHETIC SCHEME:



2'-C-Methylguanosin-5'-yl-N-(4-trifluoromethyl)-benzylaminyloxy-O-(triphenylmethoxy-*tert*-butyl-S-acyl-2-thioethyl) phosphate (D1**):**

[00457] To a solution of compound **B1** [See, Compound 7, Example 3, Procedure A] (355 mg, 0.465 mmol) in anhydrous carbon tetrachloride (4.65 mL) 4-trifluoromethylbenzylamine (331 μL , 2.324 mmol) was added. The reaction mixture was stirred at room temperature for 1 h 30 and concentrated under reduced pressure (bath temperature not exceeding 30 $^{\circ}\text{C}$). The resulting residue was subjected to silica gel chromatography, eluting with a gradient 0-10 % methanol in dichloromethane, to give **D1** as a white solid. (420 mg, 96%). Compound **D1**: ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.77-7.20 (m, 20H, 3 C_6H_5 , $\text{C}_6\text{H}_4\text{CF}_3$ and $\text{H}-8$), 6.57 (ls, 2H, NH_2), 5.84-5.75 (m, 2H, $\text{H}-1'$ and PNH), 5.50 (m, 1H, $\text{OH}-3'$), 4.26-3.86 (m, 8H, $\text{H}-3'$, $\text{H}-4'$, $\text{H}-5'$, CH_2O and CH_2N), 3.10 (t, 2H, CH_2S , $J_{\text{CH}_2\text{S}-\text{CH}_2\text{O}} = 4.0$ Hz), 3.03 (m, 2H, CH_2OTr), 1.11 (s, 6H, 2 CH_3), 0.82 (s, 3H, CH_3). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 204.0 ($\text{C}=\text{O}$), 157.2 (C-4), 154.2 (C-2), 151.3 (C-6), 145.8-143.9 (4C, 3 C_6H_5 and $\text{C}_6\text{H}_4\text{CF}_3$), 135.6 (C-8), 129.0-120.0 (20C, 3 C_6H_5 and $\text{C}_6\text{H}_4\text{CF}_3$), 117.0 (C-5), 91.0 (C-1'), 86.1 (C(C_6H_5)), 80.7 (C-3'), 78.7 (C-2'), 73.3 (C-4'), 70.0 (CH_2OTr), 65.9 (CH_2O), 64.4 (C-5'), 50.8 (C(CH_3) $_2\text{CH}_2\text{OTr}$), 44.2 (CH_2N), 28.8 (CH_2S), 22.7 (2C, 2


CH₃), 20.4 (CH₃). ³¹P NMR (162 MHz, DMSO-*d*₆): δ 9.80 (s), 9.64 (s). ¹⁹F NMR (376 MHz, DMSO-*d*₆): δ -60.8 (s). LR LC/MS (M+H⁺) 937.3 (M-H⁻) 935.4 (5.47 min). UV: λ_{max} = 254 nm. R_f 0.61 (MeOH/CH₂Cl, 15/85, v/v).

***O*-(Hydroxy-*tert*-butyl-*S*-acyl-2-thioethyl)-2'-*C*-methylguanosin-5'-yl-*N*-(4-trifluoromethyl)-benzylaminyl phosphate B242 (Compound D2):**

[00458] Compound **D1** (400 mg, 0.427 mmol) was dissolved in dichloromethane (1.6 mL) and treated with TFA (160 μL). The mixture was stirred at room temperature for 15 min, then diluted with ethanol, evaporated to dryness (bath temperature not exceeding 30 °C) and coevaporated with toluene. The resulting residue was subjected to silica gel chromatography, eluting with a gradient 0-15 % methanol in dichloromethane and then purified by reverse phase (C 18) silica gel column chromatography eluting with a gradient 0-100% acetonitrile in water and lyophilised from a mixture of water/dioxan to give compound **B2742 (Compound D2)** (90 mg, 30%, white lyophilised powder). **B242 (Compound D2)**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.54 (ls, 1H, NH), 7.75 (s, 1H, *H*-8), 7.75-7.52 (m, 4H, C₆H₄CF₃), 6.50 (sl, 2H, NH₂), 5.82-5.74 (m, 2H, *H*-1' and PNH), 5.40 (m, 1H, OH-3'), 5.17 (s, 1H, OH-2'), 4.92 (t, 1H, OH, J_{OH-CH2} = 4.0 Hz), 4.26-3.84 (m, 8H, *H*-3', *H*-4', *H*-5', CH₂O and CH₂N), 3.41 (d, 2H, CH₂OH, J_{CH2-OH} = 4.0 Hz), 3.03 (t, 2H, CH₂S, J_{CH2S-CH2O} = 8.0 Hz), 1.07 (s, 6H, 2 CH₃), 0.82 (s, 3H, CH₃). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 204.4 (C=O), 157.2 (C-4), 154.1 (C-2), 151.2 (C-6), 145.9 (C₆H₄CF₃), 135.8 (C-8), 128.3-125.4 (6C, C₆H₄CF₃), 117.0 (C-5), 90.9 (C-1'), 80.5 (C-3'), 78.7 (C-2'), 73.2 (C-4'), 68.8 (CH₂OH), 66.0 (CH₂O), 64.4 (C-5'), 52.2 (C(CH₃)₂CH₂OH), 44.3 (CH₂N), 28.7 (CH₂S), 22.3 (2C, 2 CH₃), 20.4 (CH₃). ³¹P NMR (162 MHz, DMSO-*d*₆): δ 9.62 (s), 9.77 (s). ¹⁹F NMR (376 MHz, DMSO-*d*₆): δ -60.8 (s). LR LC/MS (M+H⁺) 695.2 (M-H⁻) 693.4 (4.25 min). HRFAB-MS C₂₆H₃₅O₉N₆F₃PS (M+H⁺) calculated 695.1876, found 695.1874. UV: λ_{max} = 253 nm. R_f 0.43 (MeOH/CH₂Cl, 20/80, v/v).


Example 12

Preparation of B503, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 9-[(2R)2-deoxy-2-fluoro-2-*C*-ethynyl-β-D-*erythro*-furanosyl]-guanine



B503

SYNTHETIC SCHEME:



{9-[(2R)2-Deoxy-2-fluoro-2-C-ethynyl-β-D-erythro-furanosyl]-guanine}-5'-yl-O-(triphenylmethyloxy-*tert*-butyl-S-acyl-2-thioethyl) H-phosphonate (**G1**):

[00459] Compound **B5** [Unpublished results] (100 mg, 0.32 mmol) and compound **F3** [See Compound **5** of Example 2] (246 mg, 0.42 mmol) were coevaporated together with anhydrous pyridine and dissolved in this solvent (4.8 mL). Pivaloyl chloride (80 μL, 0.64 mmol) was added dropwise at – 15 °C and the solution was stirred at the same temperature for 2 h. The reaction mixture was diluted with dichloromethane and neutralised with an aqueous solution of NH₄Cl 0.5M. The mixture was partitioned between dichloromethane and aqueous NH₄Cl 0.5M, the organic phases were combined, dried over Na₂SO₄ evaporated under reduced pressure (bath temperature not exceeding 30 °C) and coevaporated twice with toluene. The crude mixture was purified by flash column chromatography eluting with a gradient

0-10 % methanol in dichloromethane + 0.2 % acetic acid) to afford the desired product **G1** as a colorless oil (68 mg, 28%). Compound **G1**: ^1H NMR (400 MHz, DMSO- d_6): δ 10.72 (ls, 1H, NH), 7.83 (s, 1H, *H*-8), 7.35-7.11 (m 15H, 3 C_6H_5), 6.59 (m, 2H, NH_2), 6.36 (d, 1H, OH-3', $J_{\text{OH}-3'} = 7.6$ Hz), 6.14 (d, 1H, *H*-1', $J_{1'-\text{F}} = 18.0$ Hz), 4.65 (m, 1H, *H*-3'), 4.40-4.33 (m, 2H, *H*-5'), 4.10-4.01 (m, 3H, *H*-4' and CH_2O), 3.93 (d, 1H, CCH, $^4J_{\text{H-F}} = 5.6$ Hz), 3.15-3.12 (m, 2H, CH_2S), 3.04 (s, 2H, CH_2OTr). ^{31}P NMR (162 MHz, DMSO- d_6): δ 9.50 (s), 9.22 (s). ^{19}F NMR (376 MHz, DMSO- d_6): δ -156.5 (m). LR LC/MS (B) ($\text{M}+\text{Na}^+$) 798.2 ($\text{M}-\text{H}^-$) 774.2 (4.93 min). UV: $\lambda_{\text{max}} = 254$ nm. R_f 0.48 (MeOH/ CH_2Cl_2 , 15/85, v/v).

[00460] *N*-Benzylaminyl-{9-[(2R)2-deoxy-2-fluoro-2-*C*-ethynyl- β -D-erythro-furanosyl]-guanin}-5'-yl-*O*-(triphenylmethyloxy-*tert*-butyl-*S*-acyl-2-thioethyl) phosphate (**G2**)

[00461] To a solution of compound **G1** (68 mg, 0.088 mmol) in anhydrous carbon tetrachloride (880 μL), benzylamine (48 μL , 0.44 mmol) was added dropwise. The reaction mixture was stirred at room temperature for 2 h and evaporated to dryness (bath temperature not exceeding 30 $^\circ\text{C}$). The crude mixture was filtered on a silica gel plug, eluting with a gradient 0-10 % methanol in dichloromethane to give compound **G2** as a white solid (80 mg, quantitative yield). Compound **G2**: ^{31}P NMR (162 MHz, DMSO- d_6): δ 9.95 (s) 9.80 (s). ^{19}F NMR (376 MHz, DMSO- d_6): δ -157.5 (m). LR LC/MS (B) ($\text{M}+\text{H}^+$) 881.3 ($\text{M}-\text{H}^-$) 879.4 (5.18 min). UV: $\lambda_{\text{max}} = 254$ nm. R_f 0.31 (MeOH/ CH_2Cl_2 , 15/85, v/v).


N-Benzylaminyl-{9-[(2R)2-deoxy-2-fluoro-2-*C*-ethynyl- β -D-erythro-furanosyl]-guanin}-5'-yl-*O*-(hydroxy-*tert*-butyl-*S*-acyl-2-thioethyl) phosphate B503 (Compound **G3**):

[00462] Compound **G2** (80 mg, 0.09 mmol) was dissolved in dichloromethane (320 μL) and treated with TFA (32 μL). The mixture was stirred at room temperature for 10 min, filtered through a solid phase extraction column eluting with a gradient 0-30 % methanol in dichloromethane, then purified by reverse phase (C 18) silica gel column chromatography eluting with a gradient 0-100 % acetonitrile in water and lyophilised from a mixture of water/dioxan to give compound B503 (Compound **G3**)

(15 mg, 26%, white lyophilised powder). **B503 (Compound G3)**: ^1H NMR (400 MHz, DMSO- d_6): δ 10.61 (ls, 1H, NH), 7.83 (s, 1H, H-8), 7.30-7.18 (m, 5H, C₆H₅), 6.60 (ls, 2H, NH₂), 6.32 (m, 1H, OH-3'), 6.11 and 6.12 (2 d, 2 x 1H, 2 H-1', J_{1'-F} = 18.0 Hz), 5.68 (m, 1H, PNH), 4.93 (t, 1H, OH, J_{OH-CH2} = 5.5 Hz), 4.61 (m, 1H, H-3'), 4.26-4.18 (m, 2H, H-5'), 4.08 (m, 1H, H-4'), 3.98-3.82 (m, 5H, CH₂O, CH₂N and CCH), 3.42 (d, 2H, CH₂OH, J_{CH2-OH} = 5.0 Hz), 3.01 (m, 2H, CH₂S), 1.09 (s, 6H, 2 CH₃). ^{31}P NMR (162 MHz, DMSO- d_6): δ 9.92 (s), 9.79(s). ^{19}F NMR (376 MHz, DMSO- d_6): δ -156.8 (m). LR LC/MS (B) (M+H⁺) 639.2 (M-H⁻) 637.3 (3.85 min). HRFAB-MS C₂₆H₃₃O₈N₆FPS (M+H⁺) calculated 639.1802, found 639.1816. UV: λ_{max} = 253 nm. R_f 0.46 (MeOH/CH₂Cl, 20/80, v/v).


The starting nucleoside was synthesized as follows:

[00463] Synthesis of 9-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-furanosyl]-guanine (D961, starting nucleoside of EXAMPLE 12), and synthesis of its triphosphate derivative B427)



D961

SYNTHETIC SCHEME:



9-[3,5-O-(1,3-Diisopropyl-1,1,3,3-tetraisopropoxydisiloxane)-ribo-furanosyl]-N²-

isobutyryl-guanine (B1): Hirao, I.; Ishikawa, M.; Miura, K. *Chem. Lett.* **1986**, *11*, 1929-1932.

[00464] 9-[3,5-O-(1,3-diisopropyl-1,1,3,3-tetraisopropoxydisiloxane)-2-C-trimethylsilyl-β-D-arabino-furanosyl]-N²-isobutyryl-guanine (**B2**): To a suspension of CrO₃ (11.07 g, 110.76 mmol) in dichloromethane (220 mL) at 0 °C, acetic anhydride (10.4 mL, 110.76 mmol) and anhydrous pyridine (17.82 mL, 221.52 mmol) were added. Compound **B1** (22 g, 36.92 mmol) in solution in dichloromethane (110 mL) was added dropwise. The cooling bath was removed and the resulting solution stirred at room temperature for 5 h. The reaction mixture was poured into cold ethyl acetate, filtered through a silica and celite gel plug, concentrated to dryness and coevaporated twice with toluene. The residue obtained was dissolved in dichloromethane and stirred with an excess of MgSO₄ overnight, filtered and evaporated to get the ketone. The trimethylsilylacetylene (12.5 mL, 88.60 mmol) was dissolved in anhydrous THF (98 mL) under argon. Butyllithium (55.4 mL, 1.6 M in hexanes) was added dropwise at - 78 °C. The reaction mixture was stirred for 30 min at - 78 °C and then allowed to warm up to - 55 °C. The ketone in solution in THF (49 mL) was added dropwise at - 78 °C. The reaction mixture was stirred for 1 h at - 78 °C and then allowed to warm up to - 30 °C and stirred for 3 h. The reaction was quenched by careful addition of aqueous saturated NH₄Cl (72 mL) at - 78 °C. After warming to room temperature, the mixture was diluted with ethyl acetate, washed twice with saturated brine, dried (Na₂SO₄) and concentrated to dryness. The crude material was purified using column chromatography eluting with 1.5 % MeOH in

dichloromethane to give compound **B2** (8.59 g, 34 %, 2 steps) as a pale yellow foam. Compound **B2**: NMR ^1H (250 MHz, DMSO- d_6): δ 12.10 (ls, 1H, NH), 11.69 (ls, 1H, NH), 7.91 (s, 1H, *H*-8), 6.69 (s, 1H, OH), 5.94 (s, 1H, *H*-1'), 4.29 (d, 1H, *H*-3', $J_{3',4'} = 5.5$ Hz), 3.85-3.95 (m, 3H, *H*-4', *H*-5' and *H*-5''), 2.46 (m, 1H, $\text{CH}(\text{CH}_3)_2$), 0.90-1.08 (m, 30H, iPr and $\text{CH}(\text{CH}_3)_3$), 0.00 (s, 9H, $\text{Si}(\text{CH}_3)_2$). LC/MS (A): ($\text{M}+\text{H}^+$) 692.4 (24.96 min). UV: $\lambda_{\text{max}1} = 254$ nm, $\lambda_{\text{max}2} = 281$ nm. R_f 0.34 (MeOH/ CH_2Cl_2 , 15/85, v/v).

9-[(2R)-2-Deoxy-2-fluoro-3,5-*O*(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethyl-silylethynyl- β -D-erythro-furanosyl]-*N*²-isobutyryl-guanine (B3**):**

[00465] Compound **B2** (2.00 g, 2.89 mmol) was dissolved in dried DCM (60 mL) under argon and pyridine (1.45 mL, 18.06 mmol) was added. The reaction mixture was cooled to -20 °C and DAST (4.11 mL, 31.35 mmol) was added dropwise. The cooling bath was removed after completion of the addition. Stirring was continued for 1 h 15 and the mixture was dissolved with ethyl acetate and poured into saturated NaHCO_3 and stirred for 5 min. The organic layer was washed with saturated brine, dried (Na_2SO_4), concentrated, and purified by silica gel chromatography eluting with ethyl acetate in DCM (2 %) to give the desired compound **B3** (1.41 g, 70 %) as a yellow oil. Compound **B3**: NMR ^1H (250 MHz, DMSO- d_6): δ 12.22 (s, 1H, NH), 8.09 (s, 1H, *H*-8), 6.21 (d, 1H, *H*-1', $J_{1',F} = 15.6$ Hz), 4.54 (dd, 1H, *H*-3', $J_{3',F} = 23.6$ Hz, $J_{3',4'} = 9.8$ Hz), 4.33 (m, 1H, *H*-5', $^2J_{5',5''} = 13.1$ Hz), 4.16 (m, 1H, *H*-5''), 2.81 (m, 1H, $\text{CH}(\text{CH}_3)_2$), 1.13-1.03 (m, 34 H, iPr and $\text{CH}(\text{CH}_3)_2$), 0.08 (s, 9H, $\text{Si}(\text{CH}_3)_3$, $^3J_{\text{H-H}} = 6.9$ Hz). NMR ^{19}F (235 MHz, DMSO- d_6): δ -160.26 (dd, $J_{F-1'} = 16.1$ Hz, $J_{F-3'} = 23.3$ Hz). LC/MS (A): ($\text{M}+\text{H}^+$) 694.7 (24.02 min). LRFAB-MS (GT): 694 ($\text{M}+\text{H}$)⁺, 692 ($\text{M}-\text{H}$)⁻. UV: $\lambda_{\text{max}} = 256$ nm. R_f 0.46 (MeOH/ CH_2Cl_2 , 05/95, v/v).


9-[(2R)-2-Deoxy-2-*C*-ethynyl-2-fluoro- β -D-erythro-furanosyl]-*N*²-isobutyryl-guanine (B4**):**

[00466] Compound **B3** (1.31 g, 1.89 mmol) was dissolved in methanol (13.8 mL) and ammonium fluoride (908.9 mg, 24.54 mmol) was added. The resulting solution was stirred at reflux for 1 h and evaporated to dryness. The crude material was purified on silica gel chromatography eluting with a stepwise gradient 6-10 % of


methanol in dichloromethane to yield compound **B4** (344 mg, 48 %) as a pale yellow oil. Compound **B4**: NMR ^1H (400 MHz, $\text{DMSO}-d_6$): δ 12.18 (ls, 1H, NH), 11.77 (ls, 1H, NH), 8.34 (s, 1H, H-8), 6.29 (d, 1H, OH-3', $J_{\text{OH-3'}} = 7.5$ Hz), 6.20 (d, 1H, H-1', $J_{1'-\text{F}} = 16.2$ Hz), 5.39 (t, 1H, OH-5', $J_{\text{OH-5'}} = 5.1$ Hz), 4.52 (dt, 1H, H-3', $J_{3'-\text{F}} = 22.9$ Hz), 3.98 (m, 1H, H-4'), 3.90-3.85 (m, 2H, H-5' and ethynyl), 3.72 (m, 1H, H-5''), 2.52 (m, 1H, $\text{CH}(\text{CH}_3)_2$), 1.14 (d, 6 H, $\text{CH}(\text{CH}_3)_2$, $^3J_{\text{H-H}} = 6.9$ Hz). NMR ^{13}C (100 MHz, $\text{DMSO}-d_6$): δ 180.7 (C-6), 155.3 (C-2), 148.9 (C-4), 137.3 (C-8), 120.4 (C-5), 95.8 (d, C-2', $^1J_{2'-\text{F}} = 182.1$ Hz), 87.7 (d, C-1', $^2J_{1'-\text{F}} = 39.2$ Hz), 83.4 (d, CCH, $^3J_{\text{C-F}} = 9.1$ Hz), 82.6 (C-4'), 75.9 (d, CCH, $^2J_{\text{C-F}} = 31.2$ Hz), 72.9 (d, C-3', $^2J_{3'-\text{F}} = 19.1$ Hz), 59.3 (C-5'). NMR ^{19}F (235 MHz, $\text{DMSO}-d_6$): δ -158.9 (m). LC/MS (A): (M+H⁺) 380.3 (8.34 min). UV: $\lambda_{\text{max}1} = 260$ nm, $\lambda_{\text{max}2} = 277$ nm. R_f 0.40 (MeOH/ CH_2Cl_2 , 15/85, v/v).

9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-furanosyl]-guanine D961 (Compound B5):

[00467] Compound **B4** (0.78 g, 1.33 mmol) was dissolved in saturated methanolic ammonia (62 mL) and stirred at room temperature for 20 h. The reaction mixture was then evaporated to dryness under reduced pressure. The residue was dissolved in water and washed twice with ethyl acetate. The aqueous layer was evaporated and purified on reverse phase column chromatography (C 18) eluting with a gradient 2-15 % of acetonitrile in water. The residue obtained was then dissolved in hot ethyl acetate, filtered and dried to give **D961 (Compound B5)** (134 mg, 33 %) as a yellow solid. NMR ^1H (400 MHz, $\text{DMSO}-d_6$): δ 10.70 (ls, 1H, NH), 7.98 (s, 1H, H-8), 6.60 (ls, 2H, NH_2), 6.21 (d, 1H, OH-3', $J_{\text{OH-3'}} = 7.6$ Hz), 5.83 (d, 1H, H-1', $J_{1'-\text{F}} = 16.9$ Hz), 5.29 (t, 1H, OH-5', $J_{\text{OH-5'}} = 5.2$ Hz), 4.50 (td, 1H, H-3', $J_{3'-\text{F}} = 22.8$ Hz, $J_{3'-4'} = 9.2$ Hz), 3.93-3.81 (m, 3H, H-4', H-5' and ethynyl), 3.70 (m, 1H, H-5''). NMR ^{13}C (100 MHz, $\text{DMSO}-d_6$): δ 157.2 (C-6), 154.3 (C-2), 151.05 (C-4), 135.1 (C-8), 116.7 (C-5), 96.4 (d, C-2', $^1J_{\text{C-F}} = 182.1$ Hz), 87.4 (d, C-1', $^2J_{\text{C-F}} = 39.2$ Hz), 83.1 (d, CCH, $J_{\text{C-F}} = 9.1$ Hz), 82.4 (C-4'), 76.2 (d, CCH, $^2J_{\text{C-F}} = 31.2$ Hz), 73.2 (d, C-3', $^2J_{\text{C-F}} = 20.1$ Hz), 59.5 (C-5'). NMR ^{19}F (235 MHz, $\text{DMSO}-d_6$): δ -158.5 (m). LC/MS (A): (M+H⁺) 310.1 (5.55 min). LRFAB-MS (GT): 619 (2M+H)⁺, 310 (M+H)⁺, 152 (B+H)⁺, 617 (2M-H)⁻, 308 (M-H)⁻. UV: $\lambda_{\text{max}} = 254$ nm.



SYNTHETIC SCHEME:



Standard procedure for preparation of nucleoside 5'-triphosphate: (Ludwig, J. *Acta Biochim. Biophys. Acad. Sci. Hung.* **1981**, *16*, 131-133.)


[00468] To a solution of nucleoside (0.286 mmol) in triethylphosphate (750 μ L), phosphoryle chloride (75 μ L, 0.807 mmol) was added at 0 °C. This reaction mixture A was stirred overnight at 5 °C. Tributylammonium pyrophosphate (PPI/ Bu_3N 1/1,5, 1g, 2.19 mmol) was dissolved in anhydrous DMF (2 mL). Tributylamine (420 μ L, 1.76 mmol) was added to the PPI and the resulting mixture was stirred for 15 min at 0 °C. 2.4 mL of this solution were added to the reaction mixture A. The reaction mixture was stirred at 0 °C for 1 min. The reaction was carefully quenched with TEAB 1M (pH = 7.5, 10 mL), stirred 20 min at 0 °C, then diluted with water and ethyl acetate. The aqueous phase was concentrated under reduced pressure. The crude material was subjected to DEAE-Sephadex chromatography eluting with a gradient 10^{-3} -1 M of TEAB). The desired fractions were combined, concentrated under reduced pressure and coevaporated with a mixture of water/methanol, and finally coevaporated with water. The resulting residue was purified on semipreparative HPLC. Fractions containing the expected product were concentrated under reduced pressure, coevaporated with a mixture of water/methanol and lyophilised from water. The

triethylammonium salt triphosphate was eluted three times with water on a Dowex Na⁺ resin column to yield after lyophilisation from water to the sodium salt.

[00469] 9-[(2R)2-deoxy-2-C-ethynyl-2-fluoro-β-D-erythro-furanosyl]-guanine 5'-triphosphate sodium salt (B427): ¹H NMR (400 MHz, D₂O): δ 7.97 (s, 1H, H-8), 6.19 (d, 1H, H-1', ³J_{1'-F} = 16.0 Hz), 4.70 (m, 1H under H₂O, H-3'), 4.39 (m, 1H, H-5'), 4.29-4.22 (m, 2H, H-4' and H-5''), 2.98 (d, 1H, ethynyl, ⁴J_{H-F} = 5.0 Hz). ³¹P NMR (162 MHz, D₂O): -10.50 (d, 1P, P_γ, J_{P_γ-P_β} = 19.4 Hz), -11.03 (d, 1P, P_α, J_{P_α-P_β} = 19.4 Hz), -22.38 (t, 1P, P_β, J_{P_β-P_γ} = J_{P_β-P_α} = 19.4 Hz). NMR ¹⁹F (376 MHz, DMSO-*d*₆): δ -159.1 (m). LRFAB-MS (GT): 638 (M+Na)⁺, 616 (M+H)⁺, 594 (M-Na+2H)⁺, 572 (M-2Na+3H)⁺, 550 (M-3Na+4H)⁺, 592 (M-Na)⁻, 570 (M-2Na+H)⁻, 548 (M-3Na+2H)⁻.


Example 13

Preparation of B306, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-C-methyl-5-aza-7-deaza-guanosine



B306


SYNTHETIC SCHEME:




[00470] The pronucleotide **B306** (25 mg, 6% overall yield) has been synthesized from its parent nucleoside 2'-C-methyl-5-aza-7-deaza-guanosine (200 mg, 0.67 mmol) following a similar procedure than the one described for the synthesis of the pronucleotide prepared in the Example 4 and isolated as a white lyophilized powder. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 0.90-0.91 (d, J = 2.56 Hz, 3H), 1.09 (d, J = 4.26 Hz, 6H), 3.07-3.10 (t, J = 6.66 Hz, 2H), 3.42 (d, J = 5.64 Hz, 2H), 3.86-3.99 (m, 6H), 4.10-4.15 (m, 1H), 4.15-4.20 (m, 1H), 4.90-4.93 (t, J = 5.64 Hz, 1H), 5.28 (s, 1H), 5.46-5.50 (m, 1H) 5.62-5.69 (m, 1H), 5.80 (s, 1H), 7.00 (s, 2H), 7.18-7.21 (m, 2H), 7.26-7.33 (m, 5H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.80-9.95 (2s); Scan ES $^+$ 627 (M+H) $^+$, λ_{max} = 261.7 nm; HPLC (0-100% ACN over a period of 8 min) t_{R} =3.18 min λ_{max} = 258.4 nm.

Example 14

Preparation of B389, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-C-methyl-7-deaza-guanosine




SYNTHETIC SCHEME:




[00471] The pronucleotide **B389** (80 mg, 21% overall yield) has been synthesized from its parent nucleoside 2'-C-methyl-7-deaza-guanosine (200 mg, 0.67 mmol) following a similar procedure than the one described for the synthesis of the pronucleotide prepared in the Example 4 and isolated as a white lyophilized powder. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ (ppm) 0.74 (s, 3H), 1.09 (s, 6H), 3.0 (t, $J = 6.10$ Hz, 2H), 3.42 (d, $J = 5.49$ Hz, 2H), 3.8-4.0 (2m, 6H), 4.04-4.11 (m, 1H), 4.24-4.17 (m, 1H), 4.90-4.93 (t, $J = 5.36$ Hz 1H), 4.96-4.98 (d, $J = 4.76$ Hz, 1H), 5.31-5.36 (m, 1H), 5.57-5.67 (m, 1H), 5.93 (s, 1H), 6.21-6.26 (m, 3H), 6.76 (d, $J = 22$ Hz, 1H), 7.19-7.23 (m, 1H), 7.27-7.32 (m, 4H), 10.34 (brs, 1H); ^{31}P NMR ($\text{DMSO}-d_6$, 162 MHz) δ (ppm) 9.77 and 9.90 (2s); Scan $\text{ES}^+ 626 (\text{M}+\text{H})^+$, $\lambda_{\text{max}} = 258.7$ nm; HPLC (0-100% ACN over a period of 8 min) $t_R = 3.84$ min $\lambda_{\text{max}} = 259.6$ nm.

Example 15

Preparation of **B288**, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 3'-C-methyluridine




SYNTHETIC SCHEME:



[00472] The pronucleotide **B288** (34 mg, 3% overall yield) was synthesized from its parent nucleoside 3'-*C*-methyl-uridine (513 mg, 1.99 mmol) following a similar procedure than the one described for the synthesis of the pronucleotide prepared in the Example 4 and isolated as a white lyophilized powder. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 1.09 (s, 6H), 1.15 (s, 3H), 3.00-3.05 (m, 2H), 3.30 (s, 1H), 3.42 (d, $J=6.13\text{Hz}$, 2H), 3.76-3.79 (m, 1H), 3.86-3.99 (m, 6H), 4.92-4.94 (t, $J=5.40\text{ Hz}$, 1H), 4.97 (s, 1H), 5.47 (m, 1H), 5.59-5.62 (m, 1H), 5.67-5.78 (m, 1H), 5.83-5.87 (m, 1H), 7.20-7.24 (m, 1H), 7.30 (m, 4H), 7.66-7.71 (m, 1H), 11.32 (brs, 1H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.66 and 9.95 (2s); Scan ES $^+$ 588 (M+H) $^+$, λ_{max} = 261.7 nm.


Example 16

Preparation of B350, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 3'-*C*-methylguanosine



B350

SYNTHETIC SCHEME:



3'-C-Methylguanosin-5'-yl-O-(triphenylmethoxy-*tert*-butyl-*S*-acyl-2-thioethyl) *H*-phosphonate (E1**):**

[00473] 3'-C-Methylguanosine (NM76) (233.7 mg, 0.79 mmol) and compound **A3** [See Compound **5** of Example 2] (504.9 mg, 0.87 mmol) were coevaporated together with anhydrous pyridine and dissolved in this solvent (11.8 mL). Pivaloyl chloride (193.7 μ L, 1.57 mmol) was added dropwise at -15°C and the solution was stirred at the same temperature for 2 h. The reaction mixture was diluted with dichloromethane and neutralized with an aqueous solution of NH₄Cl 0.5M. The mixture was partitioned between dichloromethane and aqueous NH₄Cl 0.5M, the organic phases were combined, dried over Na₂SO₄ evaporated under reduced pressure (bath temperature not exceeding 30 °C) and coevaporated twice with toluene. The crude mixture was filtered on a silica gel plug eluting with a gradient 0-10% methanol in dichloromethane + 0.2% acetic acid) to afford the desired product **E1** (250 mg, 42%). Compound **E1**: ³¹P NMR (162 MHz, DMSO-*d*₆): δ 9.93 (s), 9.13(s). LR LC/MS ($M+H^+$) 521.1 (5.88 min). UV: $\lambda_{\max}$ = 262 nm. *R*_f 0.21 (MeOH/CH₂Cl, 15/85, v/v).

***N*-Benzylaminy-3'-C-methylguanosin-5'-yl-O-(triphenylmethoxy-*tert*-butyl-*S*-acyl-2-thioethyl) phosphate (**E2**):**

[00474] To a solution of compound **E1** (250 mg, 0.33 mmol) in anhydrous carbon tetrachloride (3.3 mL), benzylamine (178 μ L, 1.637 mmol) was added dropwise. The reaction mixture was stirred at room temperature for 1 h 30 and evaporated to dryness (bath temperature not exceeding 30 °C). The crude mixture was filtered on a silica gel plug eluting with a gradient 0-30 % methanol in dichloromethane) to give compound


E2 as a white solid (290 mg, quantitative yield). Compound **E2**: ^{31}P NMR (162 MHz, DMSO- d_6) δ 9.91 (s), 9.74 (s). LR LC/MS ($\text{M}+\text{H}^+$) 869.3 ($\text{M}-\text{H}^-$) 867.7 (5.20 min). UV: λ_{max} = 253 nm. R_f 0.13 (MeOH/ CH_2Cl_2 , 10/90, v/v).

***N*-Benzylaminyl-*O*-(hydroxyl-*tert*-butyl-*S*-acyl-2-thioethyl)-3'-*C*-methylguanosin-5'-yl phosphate **B350** (Compound **E3**):**

[00475] Compound **E2** (290 mg, 0.33 mmol) was dissolved in dichloromethane (1.16 mL) and treated with TFA (113 μL). The mixture was stirred at room temperature for 10 min, then diluted with ethanol, evaporated to dryness (bath temperature not exceeding 30 $^\circ\text{C}$) and coevaporated with toluene. The resulting residue was subjected to silica gel chromatography, eluting with a gradient 0-30 % methanol in dichloromethane, then purified by reverse phase (C 18) silica gel column chromatography eluting with a gradient 0-100% acetonitrile in water and lyophilised from a mixture of water/dioxan to give **B350** (Compound **E3**) (15 mg, 7%, white lyophilised powder). **B350** (Compound **E3**): ^1H NMR (400 MHz, DMSO- d_6) δ 10.60 (m, 1H, NH), 7.90 (s, 1H, *H*-8), 7.30-7.19 (m, 5H, C_6H_5), 6.47 (ls, 2H, NH_2), 5.72-5.59 (m, 2H, *H*-1' and PNH), 5.51 (d, 1H, *OH*-2', $J_{\text{OH}2'-1'} = 8.0$ Hz), 4.94-4.92 (2H, *OH*-3' and *OH*), 4.28 (m, 1H, *H*-2'), 4.01-3.83 (m, 7H, *H*-4', *H*-5', CH_2O and CH_2N), 3.41 (m, 2H, CH_2OH), 3.02 (t, 2H, CH_2S , $J_{\text{CH}2\text{S}-\text{CH}2\text{O}} = 6.0$ Hz), 1.20 (s, 3H, CH_3), 1.09 (s, 6H, 2 CH_3). ^{31}P NMR (162 MHz, DMSO- d_6) δ 9.86 (s), 9.72 (s). LR LC/MS ($\text{M}+\text{H}^+$) 627.2 ($\text{M}-\text{H}^-$) 625.5 (3.87 min). HRFAB-MS $\text{C}_{25}\text{H}_{36}\text{O}_9\text{N}_6\text{PS}$ ($\text{M}+\text{H}^+$) calculated 627.2002, found 627.2014. UV: λ_{max} = 251 nm.


Example 17

Preparation of B305, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 1-[2-*C*-methyl- β -ribofuranosyl]-3-carboxamido-4-fluoro-pyrazole



B305


SYNTHETIC SCHEME:




[00476] The pronucleotide **B305** (28.3 mg, 8% overall yield) has been synthesized from its parent nucleoside 1-[2-*C*-methyl-β-ribofuranosyl]-pyrazolo-3-carboxamide-4-fluoro (180 mg, 0.65 mmol) following a similar procedure than the one described for the synthesis of the pronucleotide prepared in the Example 4 and isolated as a white lyophilized powder. ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 0.75 (s, 3H), 1.08-1.09 (d, *J* = 3.35 Hz, 6H), 2.98-3.02 (m, 2H), 3.40-3.42 (m, 2H), 3.85-4.03 (m, 5H), 4.16-4.19 (m, 2H), 4.89-4.92 (m, 1H), 5.25-5.29 (m, 2H), 5.55 (s, 1H), 5.56-5.64 (m, 1H), 7.19-7.22 (m, 1H), 7.26-7.50 (m, 7H), 8.05 (d, *J* = 4.32 Hz, 1H); ³¹P NMR (DMSO-*d*₆, 162 MHz) δ (ppm) 9.75 and 9.90 (2s); ¹⁹F NMR (*d*₆-DMSO, 235 MHz) δ (ppm) -170.70 (d, *J* = 61.74 Hz, 1F); Scan ES⁺ 605 (M+H)⁺, λ_{max} = 233.7 nm; HPLC (0-100% ACN over a period of 10 min) t_R = 4.56 min λ_{max} = 235.2 nm.

Example 18

Preparation of **B436**, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-*C*-methyl-7-deaza-7-fluoro-adenosine




SYNTHETIC SCHEME:



[00477] The pronucleotide **B436** (30 mg, 9% overall yield) has been synthesized from its parent nucleoside 2'-C-methyl-7-deaza-6-NH-dimethoxytrityl-adenosine (320 mg, 0.53 mmol) following a similar procedure than the one described for the synthesis of the pronucleotide prepared in the Example 2 (Procedure A, Strategy b), and isolated as a white lyophilized powder. ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 0.66 (s, 3H), 1.02 (s, 6H), 2.95-2.98 (t, *J* = 6.10 Hz, 2H), 3.35 (d, *J* = 5.49 Hz, 2H), 3.77-3.85 (m, 3H), 3.88-3.95 (m, 3H), 4.03-4.18 (m, 2H), 4.83-4.86 (t, *J* = 5.44 Hz, 1H), 5.14 (s, 1H), 5.21-5.25 (t, *J* = 7.40 Hz, 1H), 5.55-5.66 (m, 1H), 6.14 (s, 1H), 6.9-7.3 (m, 8H), 8.01 (s, 1H); ³¹P NMR (DMSO-*d*₆, 162 MHz) δ (ppm) 9.77 and 9.89 (2s); ¹⁹F NMR (*d*₆-DMSO, 235 MHz) δ (ppm) -166.85 (d, *J* = 14.16 Hz, 1F); Scan ES⁺ 628(M+H)⁺, λ_{max} = 280.7 nm; HPLC (0-100% ACN over a period of 10 min) t_R = 4.78 min λ_{max} = 280.8 nm.


Example 19

Preparation of **B589**, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 4'-C-methyluridine



B589


SYNTHETIC SCHEME:



[00478] Following the procedures described for Example 4, and starting from 4'-C-methyluridine (A437) (200 mg, 0.77 mmol), intermediate **P3** was first produced (62 mg, 11%. ³¹P NMR (DMSO-*d*₆, 162 MHz) δ (ppm) 9.15, 9.56 (2s); Scan ES⁺ 747 (M+Na)⁺, λ_{max} = 259.7 nm), then compound **P4** during the second step (28 mg, 39%. Scan ES⁺ 852 (M+Na)⁺), and finally the desired prodrug **B589** was obtained as a white powder after lyophilization from dioxane (21 mg, 58%). ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 1.10 (s, 6H), 1.13 (s, 3H), 3.03 (t, *J* = 6.42 Hz, 2H), 3.15 (d, *J* = 5.29 Hz, 1H), 3.42 (d, *J* = 5.67 Hz, 2H), 3.72-4.16 (m, 7H), 4.06-4.15 (m, 1H), 4.92 (t, *J* = 5.29 Hz, 1H), 5.22 (d, *J* = 5.29 Hz, 1H), 5.36-5.38 (2d, 1H), 5.57-5.60 (2d, 1H), 5.64-5.70 (m, 1H), 5.78-5.80 (2d, 1H), 7.20-7.31 (m, 5H), 7.60-7.64 (2d, 1H); ³¹P NMR (DMSO-*d*₆, 162 MHz) δ (ppm) 9.58, 9.77 (2s); Scan ES⁺ 610 (M+Na)⁺, λ_{max} = 260.7 nm.


Example 20

Preparation of B678, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 4'-C-fluoromethylguanosine



B678

SYNTHETIC SCHEME:




[00479] Following the procedures of the Procedure A described in Example 3, and starting from 4'-C-fluoromethylguanosine (A402) (69.4mg, 0.22 mmol), compound **P1** (67.5 mg, 39%) was obtained as intermediate after the first step. Scan ES⁻ 780 (M-H)⁻. Second step led to the formation of intermediate **P2** (57.5 mg; 76%). Compound **P2** (26.3 mg, 0.03 mmol) was dissolved in dichloromethane (1 ml) and treated with montmorillonite K10 (150 mg) and stirred at room temperature for 1 h. The reaction mixture was directly deposited on silica SPE tube and extracted with a gradient 0-100% MeOH in dichloromethane to give after lyophilization from dioxane/water **B678** as a white powder (7.7 mg, 40%). ¹H NMR (DMSO- *d*₆, 400

MHz) δ (ppm) 1.09 (2 ls, 6H), 3.03-3.05 (m, 2H), 3.42 (m, 2H), 3.87-4.00 (m, 6H), 4.15-4.24 (m, 2H), 4.43-4.66 (m, 2H), 4.74 (m, 1H), 4.93 (m, 1H), 5.52 (d, J = 4.36 Hz, 1H), 5.58 (m, 1H), 5.70-5.73 (m, 1H), 5.75-5.77 (d, J = 8.05 Hz, 1H), 6.52 (ls, 2H), 7.24-7.35 (m, 5H), 7.92 (2s, 1H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.70, 9.83 (2s); ^{19}F NMR (d_6 -DMSO, 235 MHz) δ (ppm) -235.92, -236.25 (2s); Scan ES^+ 645 ($\text{M}+\text{H}$) $^+$, λ_{max} = 250.7 nm HPLC (0-100% ACN over a period of 8 min) t_{R} = 3.91 min λ_{max} = 251.1 nm.


Example 21

Preparation of B704, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of acyclovir



B704

SYNTHETIC SCHEME:



Scheme 1

9-(2-Hydroxy-ethoxymethyl)-guanine-5'-yl-*O*-(triphenylmethyloxy-*tert*-butyl-*S*-acyl-2-thioethyl) *H*-phosphonate (**F1**):

[00480] Acyclovir (200 mg, 0.89 mmol) and compound **A3** [See Compound **5** of Example 2] (674.2 mg, 1.15 mmol) were coevaporated together with anhydrous pyridine and dissolved in this solvent (13.3 mL). Pivaloyl chloride (162 μ L, 1.15 mmol) was added dropwise at -15°C and the solution was stirred at room temperature for 2 h. The reaction mixture was diluted with dichloromethane and neutralized with an aqueous solution of NH₄Cl 0.5M. The mixture was partitioned between dichloromethane and aqueous NH₄Cl 0.5M, the organic phases were combined, dried over Na₂SO₄ evaporated under reduced pressure (bath temperature not exceeding 30 °C) and coevaporated twice with toluene. The crude mixture was filtered on a silica gel plug eluting with a gradient 0-15% methanol in dichloromethane + 0.2% acetic acid) to afford the desired product **F1** (602 mg, 98%). Compound **F1**: LR LC/MS ($M+H^+$) 691.9 ($M-H^-$) 690.0 (4.82 min). UV: λ_{max} = 254 nm.

N-Benzylaminyl-9-(2-hydroxy-ethoxymethyl)-guanine-5'-yl-*O*-(triphenylmethyloxy-*tert*-butyl-*S*-acyl-2-thioethyl) phosphate (**F2**):

[00481] To a solution of compound **F1** (602 mg, 0.87 mmol) in anhydrous carbon tetrachloride (8.7 mL), benzylamine (475 μ L, 4.35 mmol) was added dropwise. The reaction mixture was stirred at room temperature for 1 h 30 and evaporated to dryness (bath temperature not exceeding 30 °C). The crude mixture was subjected to silica gel chromatography, eluting with a gradient 0-10% methanol in dichloromethane to give compound **F2** as a white solid (550 mg, 79%). Compound **F2**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.77 (s, 1H, *H*-8), 7.58-7.17 (m, 20H, 4 C₆H₅), 6.68 (ls, 2H, NH₂), 5.59 (m, 1H, PNH), 5.32 (s, 2H, OCH₂N), 3.92-3.78 (m, 6H, CH₂SCH₂O, CH₂N, POCH₂CH₂O), 3.51 (t, 2H, POCH₂CH₂O, $J_{CH_2-CH_2}$ = 5.2 Hz), 3.16 (s, 2H, CH₂OTr), 3.00 (t, 2H, CH₂S, $J_{CH_2S-CH_2O}$ = 5.6 Hz), 1.12 (s, 6H, 2 CH₃). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 204.0 (C=O), 157.2 (C-4), 154.6 (C-2), 151.8 (C-6), 143.9 (4 C, 4 C₆H₅), 136.9 (C-8), 128.9-127.2 (20C, 4 C₆H₅), 117.0 (C-5), 86.3 (1C, C(C₆H₅)₃), 72.3 (OCH₂N), 70.0 (CH₂OTr), 68.2 (POCH₂CH₂O), 64.8 (POCH₂CH₂O), 64.2 (CH₂SCH₂O), 50.8 (C(CH₃)₂), 44.7 (CH₂N), 28.8 (CH₂S), 22.8 (2C, C(CH₃)₂). ³¹P


NMR (162 MHz, DMSO- d_6) δ 9.79 (s). LR LC/MS ($M+H^+$) 797.2 (5.15 min). UV: $\lambda_{\max}$ = 254 nm. R_f 0.57 (MeOH/ CH_2Cl_2 , 15/85, v/v).

***N*-Benzylaminyl-9-(2-hydroxy-ethoxymethyl)-guanine-5'-yl-*O*-(hydroxy-*tert*-butyl-*S*-acyl-2-thioethyl) phosphate **B704** (Compound **F3**):**

[00482] Compound **F2** (550 mg, 0.69 mmol) was dissolved in dichloromethane (2.2 mL) and treated with TFA (220 μ L). The mixture was stirred at room temperature for 15 min, filtered through a solid phase extraction column eluting with a gradient 0-15 % methanol in dichloromethane, then purified by reverse phase (C 18) silica gel column chromatography eluting with a gradient 0-100% acetonitrile in water and lyophilised from a mixture of water/dioxan to give **B704** (Compound **F3**), (103 mg, 27%, white lyophilised powder). **B704** (Compound **F3**): ^1H NMR (400 MHz, DMSO- d_6): δ 10.57 (ls, 1H, *NH*), 7.79 (s, 1H, *H*-8), 7.29-7.18 (m, 5H, C_6H_5), 6.49 (ls, 2H, NH_2), 5.55 (m, 1H, *PNH*), 5.33 (s, 2H, OCH_2N), 4.92 (t, 1H, *OH*, J_{OH-CH_2} = 5.2 Hz), 3.94-3.73 (m, 6H, CH_2SCH_2O , CH_2N , $POCH_2CH_2O$), 3.60 (t, 2H, $POCH_2CH_2O$, $J_{CH_2-CH_2}$ = 4.2 Hz), 3.42 (d, 2H, CH_2OH , J_{CH_2-OH} = 4.4 Hz), 3.00 (t, 2H, CH_2S , $J_{CH_2S-CH_2O}$ = 6.4 Hz), 1.10 (s, 6H, 2 CH_3). ^{13}C NMR (100 MHz, DMSO- d_6): δ 204.4 ($C=O$), 157.2 (*C*-4), 154.4 (*C*-2), 151.9 (*C*-6), 141.0 (1C, C_6H_5), 138.1 (*C*-8), 128.6-127.2 (5C, C_6H_5), 117.0 (*C*-5), 72.3 (OCH_2N), 68.8 (CH_2OH), 68.2 ($POCH_2CH_2O$), 64.7 ($POCH_2CH_2O$), 64.2 (CH_2SCH_2O), 52.2 ($C(CH_3)_2$), 44.7 (CH_2N), 28.7 (CH_2S), 22.3 (2C, $C(CH_3)_2$). ^{31}P NMR (162 MHz, DMSO- d_6): δ 9.76 (s). LR LC/MS ($M+H^+$) 555.9 ($M-H^-$) 553.9 (3.77 min). HRFAB-MS $C_{22}H_{32}O_7N_6PS$ ($M+H^+$) calculated 555.1791, found 555.1795. UV: $\lambda_{\max}$ = 250 nm.

Example 22

Preparation of B390, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-*C*-methyl-2',3'-di-*O*-acetyl-cytidine




The reaction scheme shows the conversion of compound **13** to compound **14**. Compound **13** is a nucleoside derivative with a 4-NHDMTr group, a 2,3-dihydroxy-4-methylribofuranose sugar, and a 5'-O-phosphate group linked to a 2,2,4,4-tetramethyl-3-pentenoate moiety via a phosphoramidite linkage. The reaction conditions are Ac_2O , DMAP, TEA, and CH_3CN . The product, compound **14**, is the corresponding 5'-O-acetate derivative, where the 2,3-hydroxyl groups are acetylated (OAc) and the phosphate group is converted to an acetate ester. The yield is quantitative. Compound **14** is then treated with $\text{aq TFA} / \text{CH}_2\text{Cl}_2$ to yield compound **15**, where the 2,3-OAc groups are removed, replaced by 2,3-dihydroxy groups, and the phosphate group is converted to a phosphonate ester. The yield for this step is 21%.

188

273.7 nm; HPLC (0-100% ACN over a period of 10 min) $t_R = 5.04$ min $\lambda_{max} = 233.7$ nm and 271.4 nm.

Example 23


Preparation of B302



B302

Synthesis of Hydroxy-*t*BuSATE *N*-benzylphosphoramidate 2', 3'-cyclic carbonate derivative of 2'-C-methylcytidine B302

The following strategy was used for the synthesis:



[00484] Protected phosphoramidate **13** (1.72 g, 1.52 mmol) was dissolved in anhydrous dichloromethane (17ml) under argon. 1,1'-Carbonyldiimidazole (251 mg, 1.55 mmol) was added and the reaction mixture was stirred at room temperature under argon for 1 h.

[00485] Analysis by TLC (8% MeOH in DCM) indicated incomplete conversion of starting material (R_f 0.35) to product (R_f 0.56). HPLC analysis (method Test20, 272

nm) confirmed the profile: 9% starting material (Rt 7.30 min) and 91% product (Rt 7.97 min).

Further portions of CDI (final total 299 mg, 1.84 mmol) were added and the reaction mixture was left to stir for an additional 24 h at room temperature after which time analysis by HPLC indicated 1.5% SM and 97.5% P.

[00486] The reaction mixture was evaporated *in vacuo* to give an off-white foam (1.97 g). Purification by silica gel plug column, eluting with ethyl acetate, and evaporation of the appropriate fractions gave the protected cyclic carbonate **14** ($\text{C}_{65}\text{H}_{65}\text{N}_4\text{O}_{12}\text{PS}$ 1157.27 g mol^{-1}) as a white foam (1.62 g, 92% yield). TLC (8% MeOH in DCM): Rf 0.56; HPLC Test20 AUC: 99.5% @ 254 nm, Rt 7.97 min; m/z (ESI-): 1155.9 $[\text{M}-\text{H}]^-$ 100%; m/z (ESI+): 1157.5 $[\text{M}+\text{H}]^+$ 100%, 1179.5 $[\text{M}+\text{Na}]^+$ 20%.

[00487] Protected cyclic carbonate **14** (1.50 g, 1.30 mmol) was dissolved in anhydrous dichloromethane (15 ml) at room temperature under argon. Neat trifluoroacetic acid (1.77 g, 15.5 mmol) was added dropwise to the reaction mixture which was then stirred for 45 min at room temperature. Analysis by HPLC (method Test20, 272 nm) indicated disappearance of starting material (Rt 7.97 min) and formation of product (Rt 3.80 min).

[00488] Anhydrous methanol (5 ml) was added to the reaction mixture and solvents (10 ml) were partially removed *in vacuo* at 25°C. Further methanol (7 ml) was added to the mixture which was then evaporated to give an orange residue. Trituration with hexane/TBME 3:2 (12 ml) for 20 min yielded a sticky residue plus an opaque supernatant which was decanted. Retrituration with hexane/TBME 3:2 (5 ml) for 1 h and removal of the second supernatant gave, after coevaporation with methanol (3 ml), a pale foam (1.18 g).


[00489] The crude foam was purified by reverse phase chromatography (loaded in 1 ml acetonitrile and eluted with 0%, 10%, 15%, 20%, 25%, 30% acetonitrile in water). Combination of the relevant fractions, evaporation of the solvents at 25°C and chasing with ethanol (1 ml) gave cyclic carbonate **15**, **B302**, as a white foamy solid (560 mg, 71% yield).

B302: C₂₅H₃₃N₄O₁₀PS 612.59g mol⁻¹

HPLC AUC (Method Test20): 99% @ 254nm, Rt 3.83min

m/z (ESI +): 613.1 [M+H]⁺ 100%; 1225.5 [2M+H]⁺ 100%; 453.1 [N+H]⁺ 95%

m/z (ESI -): 611.4 [M-H]⁻ 80%; 1223.9 [2M-H]⁻ 50%; 451.3 [N-H]⁻ 100%



v_{max} (KBr disc) (cm⁻¹): 3346.4, 3206.5 O-H, intermolecular H-bond; 1815.3 C=O cyclic 5-ring carbonate; 1650.9 br C=O base, thioester

KF: 1.54% H₂O content

Specific Rotation: [α]_D²⁰ +9.289 (c 10.104mg cm⁻³ in DMSO)

m.p.: 100-102°C contracts and softens, 104-106°C phase transition I, 127-135°C phase transition II to a sticky glass, 140-150°C partial melting to sticky residue, 200-210°C decomposes to a brown sticky material

Elemental Analysis: Calculated: C 49.02%; H 5.43%; N 9.15%

Found: C 49.30%; H 5.26%; N 9.30% - passed with 0.26% F present (from TFA)


NMR: Assigned using ¹H, ¹³C, ³¹P, COSY, TOCSY, DEPT, HSQC and HMBC
[00490] **¹H NMR δ_H (400 MHz, d6-DMSO):** 1.11 (6H, s, (CH₃)₂C), 1.30 (3H, br-s, CH₃), 3.04 (2H, m, CH₂S), 3.44 (2H, d, *J* 4Hz, CH₂OH), 3.87-3.92 (2H, m, CH₂O), 3.94-4.01 (2H, m, CH₂Ph), 4.15-4.25 (2H, m, *H*-5', *H*-5''), 4.37 (1H, br-s, *H*-4'), 4.95 (2H, br-s, *H*-3', CH₂OH), 5.75-5.77 (2H, 2 x d, *J* 7Hz, *H*-5, P-N-*H*), 6.07 (1H, br-s, *H*-1'), 7.22-7.25 (1H, m, Ar-*H*), 7.29-7.33 (4H, m, 4 x Ar-*H*), 7.39, 7.44 (2H, 2 x br-s, NH₂), 7.62 (1H, br-d, *J* 7Hz, *H*-6)

[00491] **¹³C NMR δ_C (100 MHz, d6-DMSO):** 17.72 (CH₃), 21.78 (C(CH₃)₂), 28.13, 28.21 (CH₂S), 44.17 (PhCH₂), 51.62 (C(CH₃)₂), 63.84, 63.89 (CH₂O), 64.55 (C-5'), 68.29 (CH₂OH), 94.23 (C-5), 126.70 (Ar-C_{para}), 127.08, 128.11 (2 x Ar-C_{meta}, 2 x Ar-C_{ortho}), 140.35, 140.38 (Ar-C_{ipso}), 152.73, 154.45 (C-2, C-4), 165.69 (C-6), 203.87 (C=OS). C-1', C-2', C-3', C-4' and C=O broadened into baseline and were not observed.

^{31}P NMR δ_{P} (162 MHz, d_6 -DMSO): 9.80, 9.94 (1P, 2 x s, ratio 1.15:1.00)



Example 24

Preparation of B234, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 3'-*O*-L-valinyl-2'-*C*-methylcytidine



B234

The following strategy was used for the synthesis:



[00492] Boc protected valine (6.72 g, 30.94 mmol) was dissolved in anhydrous DCM (50 ml) and 1,1'-carbonyldiimidazole (4.87 g, 30.01 mmol) was added at room temperature under argon. Vigorous evolution of gas was observed initially during the activation step and the mixture was stirred at room temperature for 30 min.

[00493] Protected phosphoramidate **13** (10.0 g, 8.84 mmol) was dissolved in anhydrous dichloromethane (50 ml) in a separate vessel under argon. The activated Boc-Val solution was added dropwise to the phosphoramidate solution and the resulting mixture was heated to 40°C under argon for 24 h.

[00494] Analysis by TLC (8% MeOH in DCM) indicated complete conversion of starting material **13** (R_f 0.35) to product (R_f 0.50). HPLC analysis (method Test20, 272 nm) confirmed the profile: starting material (R_t 7.30 min) and product (R_t 9.46 min).

[00495] The reaction mixture was evaporated *in vacuo* to give an off-white foam. Purification by silica gel column chromatography, loaded from DCM, eluting with ethyl acetate/hexane 1:1 then 100% ethyl acetate, and evaporation of the appropriate fractions gave the protected valine ester **16** (C₇₄H₈₄N₅O₁₄PS 1330.52 g mol⁻¹) as a white foam (10.3 g, 88% yield). TLC (ethyl acetate): R_f 0.24; HPLC Test20 AUC: 97% @ 272 nm, R_t 9.46 min; *m/z* (ESI⁻): 1329.29 [M-H]⁻ 100%; *m/z* (ESI⁺): 1331.68 [M+H]⁺ 25%, 303.16 [DMTr]⁺ 100%.

[00496] Protected valine ester **16** (3.0 g, 2.25 mmol) was dissolved in anhydrous dichloromethane (22.5 ml) at room temperature under argon. Neat trifluoroacetic acid (4.5 ml, 58.4 mmol) was added dropwise to the reaction mixture over 3 min which was then stirred for 1 h at room temperature. Analysis by HPLC (method Test20, 272 nm) indicated disappearance of starting material (R_t 9.46 min) and formation of product (R_t 3.33 min) along with significant Boc intermediate (R_t 4.60 min).

[00497] Additional neat trifluoroacetic acid (1.0 ml, 13.0 mmol) was added dropwise to the reaction mixture which was then stirred for a further 1 h at room temperature. Analysis by HPLC (method Test20, 272 nm) indicated disappearance of Boc intermediate (R_t 4.60 min) and formation of product (R_t 3.33 min).

[00498] The reaction mixture was cooled to 5°C and anhydrous methanol (50 ml) was added, stirring for 30 min. Solvents were removed *in vacuo* at 25°C. The residue was treated with TBME (50 ml x 3) and triturated, decanting the three TBME liquors.

[00499] The residual material was dissolved in anhydrous methanol (5 ml) and anhydrous DCM (10 ml) and solid sodium bicarbonate (5 g) was added, stirring for 30 min, to give pH 6. The clear liquid was passed through a syringe filter. The residual solid bicarbonate was washed with 25% methanol in DCM (anhydrous, 10 ml) and the

solution was again filtered. The combined filtrates were concentrated *in vacuo* to give crude **17** (2.15 g).

[00500] The crude material was purified by reverse phase chromatography (loaded in 15 ml water and 3 ml acetonitrile and eluted with 0%, 5%, 20%, 30% acetonitrile in water). Combination of the relevant fractions and evaporation of the solvents at 25°C gave valine ester **17**, **B234**, as a white foamy solid (737 mg, 48% yield).


B234: $C_{29}H_{44}N_5O_{10}PS$ 685.73 gmol⁻¹

HPLC AUC (Method Test20): 99% @ 254nm, Rt 3.33min

m/z (ESI +): 686.3 [M+H]⁺ 100%; 1371.6 [2M+H]⁺ 20%; 526.1 [N+H]⁺ 20%

m/z (ESI -): 744.4 [M+OAc]⁻ 35%; 1369.8 [2M-H]⁻ 35%; 1430.2 [2M+OAc]⁻ 15%; 524.5 [N-H]⁻ 100%

Fragment N =
Exactly similar fragmentation
is observed for B102
and B302.



Chemical Formula: $C_{22}H_{32}N_5O_8P$
Exact Mass: 525.1988

ν_{max} (KBr disc) (cm⁻¹): 3350.7, 3211.9 O-H, N-H; 1757.8 C=O ester; 1673.9, 1652.0 C=O thioester, base

KF: 1.94% H₂O content

Specific Rotation: $[\alpha]_D^{20} +44.370$ (c 10.033mg cm⁻³ in DMSO)

NMR: Assigned using ¹H, ¹³C, ³¹P, COSY, TOCSY, DEPT, HSQC and HMBC


[00501] ¹H NMR δ_H (400 MHz, d6-DMSO): 0.96, 0.98 (2 x 3H, 2 x s, (CH₃)₂CH), 1.03 (3H, br-s, CH₃), 1.11 (6H, s, (CH₃)₂C), 2.15 (1H, m, (CH₃)₂CH), 3.03 (2H, m, CH₂S), 3.44 (2H, a-s, CH₂OH), 3.85 (1H, a-d, *J* 4.8Hz, CHNH₂), 3.85-3.92 (2H, m, CH₂O), 3.92-4.00 (2H, m, CH₂Ph), 4.06-4.11 (1H, br-m, *H*-5'), 4.17-4.20 (1H, br-m, *H*-5''), 4.27-4.29 (1H, br-m, *H*-4'), 5.08 (1H, br-s, *H*-3'), 5.73 (1H, a-t, *J* 7.3Hz, *H*-5), 5.74-5.82 (1H, m, P-N-H), 5.92 (1H, br-s, *H*-1'), 7.22-7.25 (1H, m, Ar-H), 7.28-7.32 (4H, m, 4 x Ar-H), 7.60, 7.63 (2 x 0.5H, 2 x d, *J* 7.3Hz, *H*-6). 2 x O-H and 2 x NH₂ not observed.

[00502] ^{13}C NMR δ_{C} (100 MHz, d6-DMSO): 17.84, 17.96 ($\text{CH}(\text{CH}_3)_2$), 20.42, 20.48 (CH_3), 21.78, ($\text{C}(\text{CH}_3)_2$), 28.09, 28.16 (CH_2S), 29.72 ($\text{CH}(\text{CH}_3)_2$), 44.16 (PhCH_2), 51.62 ($\text{C}(\text{CH}_3)_2$), 57.84 (CHNH_2), 63.77, 63.81 (CH_2O , C-5'), 68.26 (CH_2OH), 74.67 (C-3'), 77.28 (C-4'), 78.09 (C-2'), 91.32 (C-1'), 94.22 (C-5), 126.71 ($\text{Ar-C}_{\text{para}}$), 127.04, 127.08, 128.11 (2 x $\text{Ar-C}_{\text{meta}}$, 2 x $\text{Ar-C}_{\text{ortho}}$), 140.23, 140.27, 140.32 ($\text{Ar-C}_{\text{ipso}}$, C-6), 155.06 (C-2), 165.32 (C-4), 169.65, 169.72 (CO_2R), 203.84 ($\text{C}=\text{OS}$). C-1', C-3', C-4' broadened into baseline but observable.


[00503] ^{31}P NMR δ_{P} (162 MHz, d6-DMSO): 9.63, 9.96 (1P, 2 x s, ratio 1.02:1.00)

Example 25

Preparation of B183, the Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of 2'-C-methyl-NH-4-acetyl-cytidine



SYNTHETIC SCHEME:




[00504] To a solution of B102 (See Example 2) (compound **10**, 200 mg, 0.34 mmol) in anhydrous dimethylformamide (3.4 ml) was added dropwise acetic anhydride (1.1 eq, 34 μl). The reaction mixture was stirred at room temperature for 4h and 10 μl of acetic anhydride were added again. The reaction mixture was stirred overnight and the solvent evaporated under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient of methanol [0-10%] in methylene chloride) to give the desired acetylated pronucleotide **B183**

(169 mg, 79%) isolated as a white lyophilized powder. ^1H NMR ($\text{DMSO-}d_6$, 400 MHz) δ (ppm) 0.93 (s, 3H), 1.09 (s, 6H), 2.09 (s, 3H), 3.01-3.04 (t, $J=6.54$ Hz, 2H), 3.40-3.42 (d, $J=5.10$ Hz, 1H), 3.53-3.62 (m, 2H), 3.83-3.91 (m, 1H), 3.94-4.01 (m, 4H), 4.10-4.15 (m, 1H), 4.20-4.25 (m, 1H), 4.88-4.91 (t, $J=5.20$ Hz, 1H), 5.23 (s, 1H), 5.33-5.37 (t, $J=7.19$ Hz, 1H), 5.67-5.78 (m, 1H), 5.93 (s, 1H), 7.18-7.21 (m, 1H), 7.27-7.32 (m, 5H), 7.96 and 8.03 (2d, $J=7.59$ Hz, 1H), 10.87 (s, 1H); ^{31}P NMR ($\text{DMSO-}d_6$, 162 MHz) δ (ppm) 9.74 and 9.98 (2s); Scan ES^+ 629(M+H) $^+$, $\lambda_{\text{max}} = 300.7$ nm; HPLC (0-100% ACN over a period of 8 min) $t_R = 4.89$ min $\lambda_{\text{max}} = 302.1$ nm


Example 26

Preparation of B187, the Hydroxy-*t*BuSATE *N*-(2-(trifluoromethyl)benzyl)phosphoramidate derivative of 2'-*C*-methylcytidine



B187


SYNTHETIC SCHEME:



[00505] To a solution of compound **8** (See Example 2, Procedure A, Strategy a) (1.4 g, 1.3 mmol) in anhydrous carbon tetrachloride (13 ml) was added dropwise *N*-2-(trifluoromethyl)benzylamine (10 eq, 2.3 g). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-3%] of methanol in methylene chloride) to afford the desired protected nucleoside as a foam (60%). This compound was converted into the phosphoramidate prodrug **B187** (245 mg, 35%) following experimental conditions described in the Example 2 Strategy A and isolated as a white lyophilized powder. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 0.92 (s, 3H), 1.09 (s, 6H), 3.05 (t, J = 6.45 Hz, 2H), 3.29 (s, 1H), 3.41 (d, J = 5.60 Hz, 2H), 3.91-3.93 (m, 3H), 4.17-4.21 (m, 4H), 4.91 (t, J = 5.59 Hz, 1H), 5.06 (d, J = 4.25 Hz, 1H), 5.23 (t, J = 7.50 Hz, 1H), 5.65-5.67 (m, 1H), 5.76-5.83 (m, 1H), 5.91 (s, 1H), 7.08 and 7.16 (2s, 2H), 7.45-7.79 (m, 5H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.57-9.78 (2s, 1P); ^{19}F NMR (d_6 -DMSO, 235 MHz) δ (ppm) -60.79 (s, 3F); Scan ES $^+$ 655 (M+H) $^+$, λ_{max} = 280.73 nm; HPLC (0-100% ACN over a period of 10 min) t_{R} = 5.08 min λ_{max} = 271.4 nm.


Example 27

Preparation of B399, the Hydroxy-*t*BuSATE *N*-(4-(trifluoromethyl)benzyl)phosphoramidate derivative of 2'-*C*-methylcytidine



B399


SYNTHETIC SCHEME:



[00506] To a solution of compound **8** (See Example 2, Procedure A, Strategy a) (1.0 g, 0.94 mmol) in anhydrous carbon tetrachloride (10 ml) was added dropwise N -4-trifluoromethylbenzylamine (5 eq, 670 μl). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride) to afford the desired protected nucleoside as a foam (84%). This compound was converted into the phosphoramidate prodrug **B399** (204 mg, 40%) following experimental conditions described in the Example 2 Strategy A and isolated as a white lyophilized powder. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ (ppm) 0.91-0.92 (d, $J = 2.09$ Hz, 3H), 1.09 (s, 6H), 3.02-3.06 (m, 2H), 3.41 (d, $J = 6.17$ Hz, 2H), 3.53-3.57 (m, 1H), 3.84-3.94 (m, 3H), 4.03-4.13 (m, 3H), 4.18-4.23 (m, 1H), 4.91-4.94 (t, $J = 5.48$ Hz, 1H), 5.06 (s, 1H), 5.23-5.27 (t, $J = 6.82$ Hz, 1H), 5.65-5.67 (m, 1H), 5.79-5.87 (m, 1H), 5.90 (s, 1H), 7.09 and 7.16 (2s, 2H), 7.48-7.55 (m, 3H), 7.64-7.67 (m, 2H); ^{19}F NMR (d_6 -DMSO, 235 MHz) δ (ppm) -60.79 (s, 3F); ^{31}P NMR ($\text{DMSO}-d_6$, 162 MHz) δ (ppm) 9.55 and 9.76 (2s); Scan $\text{ES}^+ 655 (\text{M}+\text{H})^+$, $\lambda_{\text{max}} = 270$ nm; HPLC (0-100% ACN over a period of 10 min) $t_R = 5.03$ min $\lambda_{\text{max}} = 271$ nm.


Example 28

Preparation of B204, the Hydroxy-*t*BuSATE *N*-(*n*-methyl-*n*-octyl-amine)phosphoramidate derivative of 2'-*C*-methylcytidine



B204

SYNTHETIC SCHEME:




[00507] To a solution of compound **8** (See Example 2, Procedure A, Strategy a) (950 mg, 0.89 mmol) in anhydrous carbon tetrachloride (9 ml) was added dropwise *n*-methyl-*n*-octylamine (10 eq, 1.28 g). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-3%] of methanol in methylene chloride) to afford the desired protected nucleoside as a foam (88%). This compound was converted into the phosphoramidate prodrug **B204** (52 mg, 7%) following experimental conditions described in the Example 2 Strategy A and isolated as a white lyophilized powder. ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 0.83 (m, 3H), 0.93-0.94 (d, *J* = 3.75 Hz, 3H), 1.10 (s, 6H), 1.22 (s, 10H), 1.44 (m, 2H), 2.56 (d, *J* = 8.2 Hz, 3H), 2.88-2.93 (m, 2H), 3.31 (m, 2H), 3.43 (d, *J* = 5.60 Hz, 2H), 3.50-3.53 (m, 1H), 3.91-3.93 (m, 3H), 4.04-4.07 (m, 1H), 4.13-4.16

(m, 1H), 4.91 (t, $J = 5.59$ Hz, 1H), 5.06 (s, 1H), 5.23 (m, 1H), 5.65-5.67 (m, 1H), 5.91 (s, 1H), 7.08-7.16 (m, 2H), 7.50-7.57 (m, 1H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 10.52 and 10.66 (2s); Scan ES $^+$ 623 (M+H) $^+$, $\lambda_{\text{max}} = 280.73$ nm; HPLC (0-100% ACN over a period of 8 min) $t_R = 6.07$ min $\lambda_{\text{max}} = 274.9$ nm.


Example 29

Preparation of B244, the Hydroxy-*t*BuSATE *N,N*-(dibutylamine)phosphoramidate derivative of 2'-*C*-methylcytidine



B244

SYNTHETIC SCHEME:




[00508] To a solution of compound 12 (See Example 2, Procedure A, Strategy b) (1.5 g, 1.46 mmol) in anhydrous carbon tetrachloride (15 ml) was added dropwise dibutylamine (10 eq, 2.5 ml). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride) to afford the desired protected nucleoside as a foam (61%). This compound was converted into the phosphoramidate prodrug **B244** (21

mg, 4%) following experimental conditions described in the Example 2 Strategy B and isolated as a white lyophilized powder. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 0.76-0.81 (td, $J=2.40$ Hz and $J=7.43$ Hz, 6 H), 0.86-0.87 (d, $J=5.51$ Hz, 3H), 1.05 (s, 6H), 1.11-1.19 (m, 4H), 1.33-1.39 (m, 4H), 2.80-2.87 (q, $J=9.50$ Hz, $J=8.67$ Hz, 4H), 3.01-3.04 (t, $J=6.23$ Hz, 2H), 3.42-3.43 (m, 2H), 3.50-3.60 (m, 1H), 3.81-3.88 (m, 3H), 3.97-4.01 (m, 1H), 4.07-4.10 (m, 1H), 4.84-4.87 (m, 1H), 5.06 (s, 1H), 5.23 and 5.29 (2d, $J=8.0$ Hz, 1H), 5.70 (s, 1H), 5.91 (brs, 1H), 7.10 and 7.17 (2s, 2H), 7.49 and 7.55 (2d, $J=8.0$ Hz, 1H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 10.44 and 10.56 (2s); Scan ES $^+$ 609 (M+H) $^+$, $\lambda_{\text{max}} = 279.7$ nm; HPLC (0-100% ACN over a period of 8 min) $t_R = 5.59$ min $\lambda_{\text{max}} = 274.9$ nm.


Example 30

Preparation of B308, the Hydroxy-*t*BuSATE *N*-methylbenzylphosphoramidate derivative of 2'-*C*-methyl-cytidine



B308


SYNTHETIC SCHEME:



[00509] To a solution of compound **12** (See Example 2, Procedure A, Strategy b) (2.7 g, 2.6 mmol) in anhydrous carbon tetrachloride (26 ml) was added dropwise *N*-benzylmethylamine (5 eq, 1.67 ml). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride) to afford the desired protected nucleoside as a foam (44%). This compound was converted into the phosphoramidate prodrug **B308** (43 mg, 2%) following experimental conditions described in the Example 2 Strategy B and isolated as a white lyophilized powder. ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 0.93-0.94 (s, 3 H), 1.10 (s, 6H), 2.43-2.45(d, *J*= 4.26 Hz, 3H), 3.13 (t, *J*= 6.23 Hz, 2H), 3.36-3.37 (d, *J*= 5.24 Hz, 2H), 3.56-3.60 (m, 2H), 3.97-4.01 (m, 3H), 4.07-4.21 (m, 3H), 4.92-4.94 (m, 1H), 5.08 (s, 1H), 5.30-5.32 (m, 1H), 5.59-5.67 (2d *J*= 8.0 Hz, 1H), 5.91 (s, 1H), 7.13 (m, 2H), 7.42-7.50 (m, 5H), 7.45-7.54 (2d *J*= 8.0 Hz, 1H); ³¹P NMR (DMSO-*d*₆, 162 MHz) δ (ppm) 10.53 and 10.34 (2s); Scan ES ⁺ 601 (M+H)⁺, λ_{max} = 268.7; HPLC (0-100% ACN over a period of 8 min) t_R=3.37 min λ_{max} = 274.9 nm.


Example 31

Preparation of B353, the Hydroxy-*t*BuSATE *N*-piperidinephosphoramidate derivative of 2'-*C*-methyl-cytidine



B353


SYNTHETIC SCHEME:



[00510] To a solution of compound **12** (See Example 2, Procedure A, Strategy b) (300 mg, 0.29 mmol) in anhydrous carbon tetrachloride (3 ml) was added dropwise piperidine (5 eq, 145 μl). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride) to afford the desired protected nucleoside as a foam (55%). This compound was converted into the phosphoramidate prodrug **B353** (19 mg, 22%) following experimental conditions described in the Example 2 Strategy B and isolated as a white lyophilized powder. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ (ppm) 0.92 (d, $J=2.56$, 3H), 1.10 (s, 6H), 1.44-1.43 (m, 4H), 1.50-1.53 (m, 2H), 2.97-3.02 (m, 4H), 3.07-3.10 (t, $J=6.66$ Hz, 2H), 3.42 (d, $J=5.64$ Hz, 2H), 3.56-3.60 (m, 1H), 3.89-3.94 (m, 3H), 4.04-4.10 (m, 1H), 4.13-4.20 (m, 1H), 4.91-4.93 (t, $J=5.64$ Hz, 1H), 5.06 (s, 1H), 5.25-5.31 (2d, $J=9.31$ Hz, 1H), 5.68 (m, 1H), 5.90 (s, 1H), 7.17 and 7.10 (2s, 2H), 7.50-7.55 (2d, $J=9.01$ Hz, 1H); ^{31}P NMR ($\text{DMSO}-d_6$, 162 MHz) δ (ppm) 8.75 and 8.59 (2s); Scan ES^+ 565 ($\text{M}+\text{H}$) $^+$, $\lambda_{\text{max}} = 275.7$ nm; HPLC (0-100% ACN over a period of 6 min) $t_{\text{R}} = 3.08$ min $\lambda_{\text{max}} = 273.7$ nm.


Example 32

Preparation of **B354**, the Hydroxy-*t*BuSATE *N*-cyclohexylaminephosphoramidate derivative of 2'-*C*-methyl-cytidine



B354

SYNTHETIC SCHEME:




[00511] To a solution of compound **12** (See Example 2, Procedure A, Strategy b) (300 mg, 0.29 mmol) in anhydrous carbon tetrachloride (3 ml) was added dropwise cyclohexylamine (5 eq, 170 μ l). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride) to afford the desired protected nucleoside as a foam (55%). This compound was converted into the phosphoramidate prodrug **B354** (44 mg, 50%) following experimental conditions described in the Example 2 Strategy B and isolated as a white lyophilized powder. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 0.92 (d, $J=2.56$, 3H), 1.10 (s, 6H), 1.13 (m, 5H), 1.46-1.47 (m, 1H), 1.62 (m, 2H), 1.76-1.78 (m, 2H), 2.80 (m, 1H), 3.07-3.10 (t, $J=6.66$ Hz, 2H), 3.42 (d, $J=5.64$ Hz, 2H), 3.56-3.60 (m, 1H), 3.89-3.94 (m, 3H), 4.04-4.10 (m, 1H), 4.13-4.20 (m, 1H), 4.91-4.93 (t, $J=5.64$ Hz, 1H), 5.06 (m, 2H), 5.25 and 5.31 (2d, $J=7.2$ Hz, 1H), 5.68-5.71 (m, 1H), 5.90 (s, 1H), 7.19 and 7.09 (2s, 2H), 7.50 and 7.55 (2d, $J=7.2$ Hz, 1H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.05 and 8.91 (2s) Scan ES^+ 579 (M+H) $^+$, λ

$\lambda_{\max} = 280.7 \text{ nm}$; HPLC (0-100% ACN over a period of 6 min) $t_R = 3.23 \text{ min}$ $\lambda_{\max} = 274.9 \text{ nm}$.


Example 33

Preparation of B391, the Hydroxy-*t*BuSATE *N*-morpholinophosphoramidate derivative of 2'-*C*-methylcytidine



B391

SYNTHETIC SCHEME:




[00512] To a solution of compound **12** (See Example 2, Procedure A, Strategy b) (350 mg, 0.34 mmol) in anhydrous carbon tetrachloride (3.4 ml) was added dropwise morpholine (10 eq, 300 μ l). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride) to afford the desired protected nucleoside as a foam (70%). This compound was converted into the phosphoramidate prodrug **B391** (53 mg, 49%) following experimental conditions described in the Example 2 Strategy B and isolated

as a white lyophilized powder. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 0.92 (d, $J=2.56$, 3H), 1.10 (s, 6H), 3.0 (m, 4H), 3.07-3.10 (t, $J=6.66$ Hz, 2H), 3.31 (s, 2H), 3.42 (d, $J=5.64$ Hz, 2H), 3.56-3.60 (m, 3H), 3.89-3.94 (m, 3H), 4.04-4.10 (m, 1H), 4.13-4.20 (m, 1H), 4.91-4.93 (t, $J=5.64$ Hz, 1H), 5.08 (s, 1H), 5.25-5.31 (m, 1H), 5.68-5.71 (d, $J=7.2$ Hz, 1H), 5.90 (s, 1H), 7.18 and 7.12 (2s, 2H), 7.52 and 7.50 (2d, $J=7.6$ Hz, 1H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 7.76 and 7.61 (2s); Scan ES $^+ 567(\text{M}+\text{H})^+$, $\lambda_{\text{max}} = 279.7$ nm; HPLC (0-100% ACN over a period of 10 min) $t_{\text{R}} = 3.42$ min $\lambda_{\text{max}} = 273.7$ nm.


Example 34

Preparation of B395, the Hydroxy-*t*BuSATE *N*-pyrrolidinephosphoramidate derivative of 2'-*C*-methylcytidine



B395

SYNTHETIC SCHEME:



[00513] To a solution of compound **12** (See Example 2, Procedure A, Strategy b) (500 mg, 0.49 mmol) in anhydrous carbon tetrachloride (5 ml) was added dropwise

pyrrolidine (5 eq, 200 μ l). The reaction mixture was stirred at room temperature for 3h and the solvent removed under reduced pressure. The crude mixture was purified on silica gel column chromatography (eluant: stepwise gradient [0-5%] of methanol in methylene chloride) to afford the desired protected nucleoside as a foam (87%). This compound was converted into the phosphoramidate prodrug **B395** (48 mg, 21%) following experimental conditions described in the Example 2 Strategy B and isolated as a white lyophilized powder. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 0.93-0.94 (d, J = 3.75 Hz, 3H), 1.10 (s, 6H), 1.78-1.79 (q, J = 5.80 Hz, 4H), 3.09-3.09 (m, 6H), 3.42 (s, 2H), 3.57-3.59 (m, 1H), 3.92-3.93 (m, 3H), 4.09-4.11 (m, 1H), 4.16-4.18 (m, 1H), 4.93 (brs, 1H), 5.10 (s, 1H), 5.28-5.32 (t, J = 8.00 Hz, 1H), 5.70 (d, J =8.0 Hz, 1H), 5.89 (s, 1H), 7.27 and 7.40 (2s, 2H), 7.55 and 7.61 (2d, J = 8.0 Hz, 1H); ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 7.56 and 7.69 (2s); Scan ES $^+$ 551(M+H) $^+$, λ_{max} = 275.7 nm; HPLC (0-100% ACN over a period of 10 min) t_{R} =3.88 min λ_{max} = 273.7 nm.

Example 35

Anti-HBV Activity

[00514] The compound of Example 1 (NM 204) (Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of L-ddA) (A550) was contacted with HBV-infected HepG2 cells. EC₅₀ values were measured according to standard techniques. As shown in the table below, the compound of Example 1 showed significant activity compared to parent molecule LddA.

Drug	N	HBV wt (HepG2)
		EC ₅₀ (μM)
LddA	3	> 10
Ex 1 (A550)	3	0.062 $\pm$ 0.018
LdT	3	0.26 $\pm$ 0.048
Lamivudine	3	0.022 $\pm$ 0.007

Example 36

Preparation of Calibration Curve for measurement of ddATP

[00515] Measurements of the concentration of 2'-3'-dideoxyadenosine-5'-triphosphate (ddATP) (the triphosphate nucleotide of 2'-3'-dideoxyadenosine (ddA) are performed by liquid chromatography tandem mass spectrometry (LC/MS/MS), e.g., of methanolic extracts of hepatocytes.

[00516] The concentration of ddATP is measured by comparison to a standard curve.

[00517] Working stock solutions of TP-ddA are prepared from a 100 pmol / μ l stock solution in de-ionized water of ddATP (tetrasodium salt of > 91% purity) purchased from Sigma Chemical Co as follows:

ddATP working stock solutions and Preparation of Standard Curve for ddATP.

1. Working stock#1

Test compound	Stock conc <u>pmol/μl</u>	Vol taken μ L	DIH ₂ O vol μ L	Total vol μ L	Conc <u>pmol/μl</u>	<u>mol per 10 μl</u>
TP-ddA	100	2000	2000	4000	50.0	500

2. Working stock#2

Test article	Stock conc <u>pmol/μl</u>	Vol taken μ L	DIH ₂ O vol μ L	Total vol μ L	Conc <u>pmol/μl</u>	
TP-ddA	100	1000	3000	4000	25.0	250

3. Working stock#4(prepared from stock#1)

Test article	Stock conc <u>pmol/μl</u>	Vol taken μ L	DIH ₂ O vol μ L	Total vol μ L	Conc <u>pmol/μl</u>	
TP-ddA	100	500	3500	4000	12.5	125

4. Working stock#5 (prepared from stock#1)

Test article	Stock conc <u>pmol/μl</u>	Vol taken μ L	DIH ₂ O vol μ L	Total vol μ L	Conc <u>pmol/μl</u>	
TP-ddA	100	200	3800	4000	5.0	50

5. Working stock#6 (prepared from stock#1)

Test article	Stock conc <u>pmol/μl</u>	Vol taken μ L	DIH ₂ O vol μ L	Total vol μ L	Conc <u>pmol/μl</u>	
TP-ddA	100	100	3900	4000	2.5	25

6. Working stock#7 (prepared from stock#1)

Test article	Stock conc <u>pmol/μl</u>	Vol taken μ L	DIH ₂ O vol μ L	Total vol μ L	Conc <u>pmol/μl</u>	
TP-ddA	100	40	3960	4000	1.0	10

[00518] Internal standard (ISTD) working stock are prepared from a 0.50 mg /mL stock solution of 2-deoxyadenosine 5-triphosphate purchased from Sigma Chemical Co.

ISTD	Stock conc	Vol taken	MeOH vol	Total vol	Conc	Conc
	$\mu\text{g/mL}$	μL	μL	μL	$\mu\text{g/mL}$	pmol / mL
dATP	500	200	9800	10000	10	500

[00519] In some embodiments, calibration standards are prepared as follows using

Preparation of cal stds:

<u>cal std#</u>	<u>std conc</u>	<u>liver wt</u>	<u>working</u>	<u>working stock con</u>	<u>working stock vol</u>	<u>ISTD vol</u>	<u>MeOH vol</u>	<u>total vol</u>
	<u>pmol/mL</u>	<u>G</u>	<u>stock#</u>	<u>$\text{pmol}/\mu\text{L}$</u>	<u>μL</u>	<u>μL</u>	<u>μL</u>	<u>μL</u>
Blk	0	0.1			0	50	940	990
#1	50	0.1	#5	5.0	10	50	940	1000
#2	125	0.1	#4	12.5	10	50	940	1000
#3	250	0.1	#3	25.0	10	50	940	1000
#4	500	0.1	#2	50.0	10	50	940	1000
#5	1000	0.1	#1	100.0	10	50	940	1000

liver samples:

[00520] In some embodiments the following HPLC conditions are used for the HPLC MS, e.g. HPLC Tandem MS analysis instrument method:

[00521] HPLC is conducted on Phenomenex Luna Amino 3 μ m 100A, 30x2mm column, with a mobile phase: A: 70% 10mM NH₄OAc 30% ACN pH 6.0; and B: 70% 1mM NH₄OAc 30% ACN pH 10.5 as follows:

Gradient elution program:

Step	Time(min)	Flow (μ L/min)	A (%)	B (%)
0	0	400	60	40
1	1.1	400	60	40
2	1.11	400	40	60
3	2.11	400	30	70
4	2.6	400	20	80
5	3.1	400	0	100
6	5.5	400	0	100
7	5.51	400	60	40
8	10	400	60	40

Injection volume: 50 μ L

Flow rate to MS: 0.400 mL/min, no splitting of flow

Multiple Reaction Monitoring (MRM) conditions: (API3000)

Ionization Mode:	Positive Ion Electrospray (ESI+)
IonSpray Voltage (IS):	5000 V
Temperature (TEM):	550°C
Turbo IS gas	8 L/min
Nebulizer (NEB):	14
CAD Gas Setting (CAD):	6
Declustering potential (DP)	68 V
Collision energy (CE)	27 eV
Entrance/ Exit potentials (EP/CXP)	10V / 11V

Compound	Precursor ion =>	Product Ion
ddA triphosphate	476.2 =>	135.9
ddA diphosphate	396.2 =>	135.9
dA triphosphate (ISTD)	460.2 =>	135.9

*Luna Amino column is directly connected on the inlet end to a "Security Guard" cartridge holder suitable for 2.1 mm Phenomenex columns, containing a C18 cartridge.

Example 37

In vitro Phosphorylation in Hepatocytes

[00522] Primary hepatocytes (Rat, Cynomolgus Monkey or human) were seeded at 0.8×10^6 in a collagen-coated 12-well plate and allowed to attach 4-6 hours after which time the seeding medium was replaced with serum-free culture medium and cells allowed to acclimatize to the new medium overnight. On the next day, cells were exposed for 1, 4, 8 and 24 hours to test article (NM204) (A550) at 10 and 50 μM prepared in fresh culture medium from stock solution in DMSO (final DMSO concentration was 0.1%). At each time point, an aliquot (500 μl) was collected and immediately added to 500 μl of acetonitrile and stored at -20°C until analysis. The remaining exposure medium was removed and the cell monolayer (stuck to dish) washed 2 times with ice-cold PBS. Any remaining PBS was carefully removed by aspiration and cells were harvested by scraping in 1 mL 70% ice-cold methanol. Cell samples were placed overnight at -20°C and cellular debris removed by centrifugation on the next day. The supernatants were removed and filtered prior to analysis by LC/MS. A standard curve was prepared by using untreated cells processed similarly except that prior to harvesting in 70% methanol, 10 μl of LddATP standard solutions prepared in methanol were added to the washed monolayers. These control samples were then processed and analyzed as described for test samples.

[00523] The results are shown below:

LddA-TP formation in hepatocytes			
Ex 1 (A550) 10μM		LddA TP Levels (pmol/million cells)	
Time (hour)	Rat	Monkey	Human
1	159.5	287.5	161.5
4	388.0	978.0	312.5
8	468.5	1230.0	352.5
24	422.0	344.0	366.0
Ex 1 (A550) 50μM		LddATP Levels (pmol/million cells)	
Time (hour)	Rat	Monkey	Human
1	393.0	2085.0	682.5
4	1212.0	5690.0	1480.0
8	1590.0	6030.0	1930.0

[00524] As indicated from the data, significant levels of L-ddATP were detected in the hepatocytes. In monkey hepatocytes, the levels appear to reach a maximum level at 8 hours followed by a rapid decline. In contrast, levels in both rat and human hepatocyte appear to level off after 8 hours.

Example 38

In vivo Studies in Rat

[00525] Distribution of NM-204 (the compound of Example 1 (Hydroxy-*t*BuSATE *N*-benzylphosphoramidate derivative of L-ddA) (A550) in the rat liver was evaluated following a single intravenous (I.V.) or oral administration of A550 (NM-204) at a dose of 20 (oral) or 10 (I.V.) mg/Kg body weight. The dose solutions were prepared on the same day prior to dose administration.

[00526] At the specified time point (1 and 3 hours for IV animals or 1, 3 and 8 hours for oral animals), each animal was euthanized by CO₂ gas followed by exsanguination *via* the abdominal vein. Livers were collected immediately after sacrifice, flash frozen in liquid nitrogen, placed on dry ice, and later stored at -70°C, before being analyzed.

Preparation of calibration standards from control liver extracts:

[00527] Control rat liver samples were taken from whole frozen livers (Bioreclamation, Inc. Hicksville, NY) with the aid of a tissue coring utensil (Harris Unicore, 8.0mm, VWR,). Each ~ 0.1 g sample was placed in individual 2 mL poly vials with 0.940 mL of 80% MeOH / 20% DIH₂O and homogenates were prepared using a mechanical tissue disruptor (Tissue Master, Omni-International, Inc, Marietta GA). The vials received a 10 µl aliquot of a working stock solution and a 50 µl aliquot of the ISTD before vortexing for ~30 sec. The mixtures were stored overnight at -20 °C and the next day were removed for 10min of centrifugation in a benchtop centrifuge. Each supernatant was transferred to individual centrifugation filtration units (0.45 µm) and the resulting filtrates were transferred to HPLC vials for the LC/MS/MS analysis. The final concentrations of ddATP in the calibration standards was 1000, 500, 250, 125, 50, and 0 pmol / ml. Each calibration standard was directly

injected in a 50 μ L volume onto the ion-exchange column for analysis. Standard curve analysis of calibration standards from control liver extracts was conducted.

[00528] Analysis of ddATP was done by an ion-exchange chromatography method with on-line positive ionization ESI-MS/MS detection in multiple reaction monitoring (MRM) detection mode. The peak areas obtained for 4 of the 5 calibrants allowed for construction of a standard curve that demonstrated good linearity ($R^2 = 0.9996$) over a 50 – 1000 pmol / ml concentration range. This is equivalent to a range of 5 – 100 pmol per gram liver by the sample preparation employed. The HPLC MS MS conditions described in Example 5 were utilized. The lower limit of quantitation demonstrated by the LC/MS/MS method is e.g., ~ 0.2 pmol / mL for hepatocyte cellular extracts which contain much less salt.

[00529] The results showing intracellular levels of A550 (NM204) (showing the compound entered the liver cells) and LddATP (showing cleaving of the phosphoramidate moiety and triphosphorylation of the ddA to the active triphosphate in the liver) are shown below:

A550 (Ex 1) and LddATP measured in livers of male rats dosed IV or O with A550 (Ex 1)				
Animal Number	Concentration Compound (A550) (Ex 1)	Timepoint (hrs)	Concentration. ddA-TP	
	(pmol / g liver)		(pmol / g liver)	(pmol / 10 ⁶ cells)*
IV dose (10 mg/kg)				
1M1	65.8	1	2025	17.8
1M2	89.1	1	1930	16.9
1M3	85.1	1	1355	11.9
Mean	80.0		1770	15.5
IV dose (10 mg/kg)				
2M1	28.3	3	1345	11.8
2M2	26.0	3	1940	17.0
2M3	29.3	3	2990	26.2
Mean	27.9		2092	18.3
Oral dose (20 mg/kg)				
3M1	411	1	210	1.8
3M2	272	1	575	5.0
3M3	70.2	1	400	3.5
Mean	251		395	3.5
Oral dose (20 mg/kg)				
4M1	360	3	200	1.8
4M2	92.1	3	330	2.9
4M3	161	3	405	3.6
Mean	204		312	2.7

Oral dose (20 mg/kg)				
5M1	16.4	8	280	2.5
5M2	28	8	805	5.2
5M3	16.2	8	275	2.4
Mean	20.1		382	3.3

*Hepatocellularity number for rat was 114×10^6 cells per gram liver (*Toxicology in Vitro* 20 (2005) 1582-1586).

[00530] Thus, these results show that the compound can be used to enhance concentration of the drug in the liver. These results also show the enhanced concentration of the active triphosphate which is formed in the liver cells.

Example 39

HCV Replicon Assay

[00531] Huh-7 cells containing HCV Con1 subgenomic replicon (GS4.1 cells), (C. Seeger; Fox Chase University, Philadelphia, PA, USA), are grown in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 2 mM L-glutamine, 110 mg/L sodium pyruvate, 1X non-essential amino acids, 100 U/mL penicillin- streptomycin and 0.5 mg/mL G418 (Invitrogen). For dose-response testing, the cells are seeded in 96-well plates at 7.5×10^3 cells/well in a volume of 50 μ L and incubated at 37°C/5% CO₂. Three hours after plating, 50 μ L of ten 2-fold serial dilutions of compounds (highest concentration, 75 μ M) are added and cell cultures were incubated at 37°C/5% CO₂ in the presence of 0.5% DMSO.

Alternatively, compounds are tested at a single concentration of 15 μ M. In all cases, Huh-7 cells lacking the HCV replicon served as negative control. The cells are incubated in the presence of compounds for 72 hours after which they were monitored for expression of the NS4A protein by enzyme-linked immunosorbent assay (ELISA). For this, the plates were then fixed for 1 min with 1:1 acetone:methanol, washed twice with phosphate-buffered saline (PBS), 0.1% Tween 20, blocked for 1 hour at room temperature with TNE buffer containing 10% FBS and then incubated for 2 h at 37°C with the anti-NS4A mouse monoclonal antibody A-236 (ViroGen) diluted in the same buffer. After washing three times with PBS, 0.1% Tween 20, the cells are incubated 1 hour at 37°C with anti-mouse immunoglobulin G- peroxidase conjugate in TNE, 10% FBS. After washing as described above, the reaction is developed with O-phenylenediamine (Zymed). The reaction is stopped after 30 minutes with 2 N H₂SO₄ and the absorbance is read at 492 nm using a Sunrise Tecan spectrophotometer. EC₅₀ values are determined from the % inhibition *versus* concentration data using a










sigmoidal non-linear regression analysis based on four parameters with Tecan Magellan software. When screening at a single concentration, the results are expressed as % inhibition at 15 μ M. For cytotoxicity evaluation, GS4.1 cells are treated with compounds as described above and cellular viability was monitored using a Cell Titer 96 AQueous One Solution Cell Proliferation Assay (Promega). CC₅₀ values are determined from the % cytotoxicity versus concentration data with Tecan Magellan software as described above.






Results

[00532] Compounds presented in the table below were assayed according to the replicon assay described above.

Compound Reference	Structure	HCV ELISA 2	
		EC ₅₀ (μM)	CC ₅₀ (μM)
NUCLEOSIDE PARENT: A634 (NM107)		++	+
EXAMPLE 2: B102		++	+
EXAMPLE 26: B187		++	+
EXAMPLE 27: B399		++	+
EXAMPLE 28: B204		++	+
EXAMPLE 29: B244		+	+
EXAMPLE 30: B308		++	+
EXAMPLE 31: B353		+	+

Compound Reference	Structure	HCV ELISA 2	
		EC ₅₀ (μM)	CC ₅₀ (μM)
EXAMPLE 32: B354		++	+
EXAMPLE 33: B391		+	+
EXAMPLE 34: B395		++	+
EXAMPLE 24: B234		++	+
EXAMPLE 23: B302		++	+
EXAMPLE 22: B390		+	+
EXAMPLE 25: B183		+	+
NUCLEOSIDE PARENT: A844 (NM108)		++	+
EXAMPLE 3: B299		+++	+

Compound Reference	Structure	HCV ELISA 2	
		EC ₅₀ (μM)	CC ₅₀ (μM)
EXAMPLE 11: B242		+++	+
EXAMPLE 10: B307		+++	+
NUCLEOSIDE PARENT: A374 (NM80)		+++	++
EXAMPLE 6: B263		++	+
NUCLEOSIDE PARENT: C809 (NM106)		+	+
EXAMPLE 7: B229		++	+
NUCLEOSIDE PARENT: A608		+	+
EXAMPLE 8: B186		++	+
NUCLEOSIDE PARENT: A849		+++	+

Compound Reference	Structure	HCV ELISA 2	
		EC ₅₀ (μM)	CC ₅₀ (μM)
EXAMPLE 9: B396		+++	++
NUCLEOSIDE PARENT: D961		+	+
EXAMPLE 12: B503		++	+
NUCLEOSIDE PARENT: E810		+++	+
EXAMPLE 18: B436		++	+

EC₅₀ in ELISA 2 assay is provided as follows:

+++ ≤ 1 μm, ++ > 1-10 μm and + > 10 μm

CC₅₀ is provided as follows:

++ ≤ 75 μm, + > 75 μm

Example 40

HBV Drug susceptibility assay

- Collagen-I coated cell culture plates were seeded with cells at a density of 0.25-0.5x10⁶ cells per well in 2 ml of growth/selection media.
- Drug stock solutions were made up freshly in 100% DMSO as 200x stocks. Seven 4-fold dilutions of test compound were prepared ranging from 2.5 μM to 0.0006 μM (final). Master drug dilutions were divided into 4 aliquots, and then stored at -20°C until used.
- One day after cells were seeded, drug treatments were initiated by adding 10 μl of drug dilution along with 2 ml of fresh growth/selection media.

Thus, the final DMSO concentration did not exceed 0.5%. The no-drug control wells received 10 µl of DMSO in fresh media.

d) Cells were treated every-other-day with 2 ml of fresh drug combinations/medium for a total of 8 days. Cell lysates were then collected on day 10 as described below and endogenous polymerase assay were performed.

Preparation of nucleocapsid-containing lysates for EPA analysis

- a) Two days after the final drug treatment, cells were harvested.
- b) Media was carefully aspirated and the cell monolayers were rinsed once with 1 ml of PBS.
- c) 1 ml of lysis buffer (50 mM Tris-HCl pH 7.5/150 mM NaCl/5 mM MgCl₂/0.2% NP-40) was added to each well. The detergent is required to strip the outer envelope from virions and to allow capture of the inner nucleocapsids. Plates were kept on ice for >30 min.
- d) Lysed cells were transferred to 1.5 ml-microfuge tubes.
- e) Lysates were clarified by spinning at room temperature for 5 min at 14,000 rpm.
- f) Clarified lysates were transferred to fresh tubes and immediately frozen on dry-ice, then stored at -80°C until endogenous polymerase assays can be performed as described below.

Endogenous polymerase assay (EPA) of cell lysates

- a) EPAs were performed essentially as described in Seifer, et al (1998). *J. Virol.* 72: 2765-2776. Clarified lysates were thawed at room temperature.
- b) Intracellular HBV nucleocapsids were immunoprecipitated from the cytoplasmic lysates overnight at 4°C with a polyclonal rabbit anti-HBcAg antibody and immobilized on protein A sepharose CL-4B beads.
- c) After 2 washes of the immobilized capsids with 1 ml of EPA wash buffer (75 mM NH₄Cl, 50 mM Tris-HCl pH 7.4, 1 mM EDTA), endogenous polymerase reactions were initiated by adding 50 µl of detergent-containing EPA cocktail (50 mM Tris-HCl pH7.4, 75 mM NH₄Cl, 1 mM EDTA, 20 mM MgCl₂, 0.1 mM β-ME, 0.5% NP-40, 100 µM cold dGTP, TTP, dCTP, and 50 nM ³³P-dATP) and incubated overnight at 37°C. The detergent is required to enhance permeability of the nucleocapsids.

- d) Following digestion with 1 mg/ml of Proteinase K for 1 hour at 37°C, endogenously ³³P-labeled HBV DNA was liberated via phenol/chloroform extraction.
- e) The nucleic acids were then precipitated with one volume of 5 M NH₄-acetate and 2.5 volumes 100% EtOH, and separated on a 1% native agarose gel in Tris-borate buffer.
- f) Gels were blotted onto positively charged nylon membrane overnight at room temperature via capillary transfer in 0.4 N NaOH.
- g) Dried membranes were exposed to a phosphorimager screen (GE Healthcare) overnight at room temperature, then scanned (Storm 860, GE Healthcare) and quantitated with ImageQuant software (GE Healthcare).
- h) Dose-response curves were generated using XLfit 4.1 software. The mean effective drug concentrations that inhibit endogenous HBV polymerase activity by 50% were calculated from several independent experiments.

Cytotoxicity determination

A standard *in vitro* cytotoxicity assay was performed in HepG2 cells. Cells were exposed to drug for 9 days. Cell viability was determined via MTS staining using a CellTiter 96 Aqueous One Solution cell proliferation assay according to the manufacturer's instructions.

- a) HepG2 cells were seeded in 96-well tissue culture plates in 100 µl fresh growth media at 7x10³ cells per well.
- b) Drug stock solutions were made up in 100% DMSO as 400x stock solutions and stored at -20°C until used.
- c) Four hours after cells were plated the drug dilutions were prepared and then added to the cells. Cells received up to 100 µM of drug in a total of 200 µl of fresh growth media containing 0.25% DMSO. Control wells received growth media with 0.25% DMSO growth media. Plates were incubated at 37°C, 5% CO₂.
- d) Cells were treated every-other-day with fresh growth media and fresh drug dilutions for a total of 8 days as described above.
- e) On day 9, cell viability of HepG2 cells was determined by adding 20 µl of MTS CellTiter 96 Aqueous One Solution. Following 4 hours of incubation at 37°C, absorbance was measured at A₄₉₀ nm in a Victor V plate reader (Perkin Elmer).

f) The CC₅₀ concentrations were determined using XLfit 4.1 software.

[00533] The antiviral *in vitro* activity of PMEA, B261 (hydroxy-tBuSATE N-benzylphosphoramidate derivative of PMEA as shown in Example 10 Table) along with LdT as control was tested in a total of 4 HBV drug susceptibility assays. The table below provides the results:

HBV Cell Assay (EPA Read-out)

Cytotoxicity		Antiviral Activity *	
Drug	(CC ₅₀ in μ M)	(EC ₅₀ in μ M)	SI
PMEA	>100	0.328 $\pm$ 0.082	>310
B261	19.6	0.016 $\pm$ 0.004	1,225
LdT	>100	0.366 $\pm$ 0.056	>273

Cytotoxicity and efficacy was determined on collagen plates.

Example 41

Determination of total metabolism in liver subcellular fractions (depletion of parent)

[00534] NADPH Incubations. Microsomal or S9 incubations were conducted in a final volume of 0.5 mL. Pooled liver microsomal or S9 protein (1.0 mg/mL), suspended in incubation buffer (100mM potassium phosphate, pH 7.4, 5 mM MgCl₂, and 0.1 mM EDTA) was preincubated for 5 min at 37°C with 10-50 μ M OHSATE phosphoramidate compound from a stock solution in DMSO (final DMSO concentration was 0.1%); the reaction was initiated by the addition of NADPH (3 mM final concentration). Incubations with no NADPH served as controls. At specific times (0-120 min), 0.1 mL samples were taken and the reaction terminated by the addition of 1 volume of stop solution (acetonitrile). The samples were vortex for 30 sec and then centrifuged at 1500g for 10 min. The supernatant was transferred to HPLC glass vials and analyzed without further processing by HPLC. Figures 1 and 2 depict depletion of NM108 SATE phosphoroamidate (B299) and NM107 SATE phosphoroamidate (B102), respectively, after incubation with NADPH in monkey liver S9.

HPLC system for medium samples-unchanged prodrug

HPLC: Agilent 1100

Column:

Phenomenex Luna C18(2), 20x2mm,

Mobile phases (MP):

MP(A) 10mM K₂HPO₄ pH5, MP(B) ACN

Gradient elution:

20 to 63% MP(B) run from 0 to 30 min

Runtime:	20 min
Flow rate:	1 mL/min
Injection volume:	10-20 μ L
UV:	252 nm-NM108SATE deriv (B299)
	272 nm-NM107SATE deriv (B102)

[00535] Thus, without being limited to any theory, since the metabolism is NADPH dependent, it is possible that the phosphoroamidate compound is preferentially activated by Cytochrome P450 in the liver.

Example 42

Determination of Triphosphate Levels in Cells

Preparation of primary hepatocyte cultures

[00536] Freshly isolated cells from animal and human liver were obtained in suspension on ice. Following receipt, cells were pelleted by centrifugation at 500 rpm (rat) or 700 rpm (monkey and human) and resuspended at 0.8 million cells per mL of plating medium (HPM). Multi-well collagen-coated plates (12-well) were then seeded by addition of 1 mL of cell suspension (0.8 million cells/ mL). The plates were gently shaken to evenly distribute the cells and placed in an incubator at 37°C for approximately 4 to 6 hours to allow cells to attach. Once cells have attached, the plating medium was removed and replaced with hepatocyte culture medium (HCM). Cells were left overnight in an incubator at 37°C to acclimatize to culture and the medium.

Incubations with test article

[00537] Hepatocyte incubations were conducted in a final volume of 1.0 mL HCM/well (0.8 million cells/ mL). HCM from overnight incubation of cells was removed and replaced with fresh HCM, pre-warmed to 37°C, containing 10 μ M test article from a stock solution in DMSO (final DMSO concentration was 0.1%). At specific times (up to 24 hrs), incubation medium was discarded and the cell monolayers were carefully washed two times with ice-cold PBS. Following the last wash, all PBS was carefully removed and 1 mL of extraction buffer (ice-cold 70% methanol) was added. Each well was sealed with parafilm immediately following addition of methanol. Once the entire plate was processed, additional parafilm was placed on entire plate forming a double seal to prevent evaporation during the extraction process. The cover lid was then placed on the plate and sealed with tape.

The plates were then stored at -20°C for a minimum of 24 hrs to allow for extraction of intracellular contents.

Preparation of Huh7 or HepG2 cultures

[00538] HepG2s or Huh7 cells were plated at 0.4×10^6 cells/well in collagen-coated 12-well plates. Cells were allowed to attach overnight. Culture medium from overnight incubation of cells was removed and replaced with fresh culture medium, pre-warmed to 37°C, containing 10 µM test article from a stock solution in DMSO (final DMSO concentration was 0.1%). After 24-72 hours, incubation medium was discarded and the cell monolayers were carefully washed two times with ice-cold PBS. Following the last wash, all PBS was carefully removed and 1 mL of extraction buffer (ice-cold 70% methanol) was added. Each well was sealed with parafilm immediately following addition of methanol. Once the entire plate was processed, additional parafilm was placed on entire plate forming a double seal to prevent evaporation during the extraction process. The cover lid was then placed on the plate and sealed with tape. The plates were then stored at -20°C for a minimum of 24 hrs to allow for extraction of intracellular contents.

Sample preparation for analysis

[00539] Cellular extracts were prepared by transferring 0.9 mL of extract into 2 mL microfuge tubes followed by centrifugation for 5 min at 14,000 rpm. Approximately 100µL of the supernatant was transferred to HPLC vials and triphosphate levels determined by LCMS/MS as described below.

[00540] HPLC conditions: NM107-triphosphate

HPLC:

Column:

30x2mm,

Mobile phases (MP):

6.0

10.5

Phenomenex Luna Amino 3 µm 100A,

(A) 70% 10mM NH₄OAc 30% ACN pH

(B) 70% 1mM NH₄OAc 30% ACN pH

Gradient elution:

Step	Time	Flow	A	B
0	0.00	400	80	20
1	0.10	400	80	20
2	0.11	400	40	60
3	0.21	400	40	60
4	2.60	400	10	90
5	2.61	400	0	100
6	5.60	400	0	100
7	5.61	400	80	20
8	9.00	400	80	20

Flow rate to MS: 0.400 mL/min, no split

Injection volume: 10 µL

Compound	Precursor ion	Product
ion		
NM107 triphosphate	498.0	112.0

[00541] Exemplary HPLC conditions: NM108-triphosphate

HPLC:

Column:

Phenomenex Luna Amino 3 µm 100A,

30x2mm,

Mobile phases (MP):

(A) 70% 10mM NH₄OAc 30% ACN pH

6.0

(B) 70% 1mM NH₄OAc 30% ACN pH

10.5


Gradient elution:

Step	Time	Flow	A	B
0	0.00	400	60	40
1	0.10	400	60	40
2	0.11	400	40	60
3	0.21	400	40	60
4	2.60	400	10	90
5	2.61	400	0	100
6	5.61	400	0	100
7	5.61	400	60	40
8	9.00	400	60	40

Flow rate to MS: 0.400 mL/min, no split

Injection volume: 10 µL

Compound	Precursor ion	Product
ion		
NM108 triphosphate	538.0	152.0



[00542] NM107 triphosphate and B102 triphosphate levels in cell extracts were observed as follows:

	Intracellular Triphosphate (pmol/million cells)			
drug in culture	Human	Monkey	HepG2*	Huh7*
B102	991	1838	1.5	9.2
NM107	19	10	17	37
24 hr incubation in 10 μ M drug				
*72 hr incubation in 10 μ M drug				

[00543] As seen from the data levels of intracellular triphosphate for B102 were higher compared to those for NM107.

Example 43

Demonstration of Potent Antiviral Activity of Second Generation Nucleoside Inhibitors, B102, in HCV-Infected Chimpanzees

[00544] Nucleoside analogs such as NM107 (2'-methyl cytidine, valopicitabine nucleoside component) have shown efficacy against HCV in the clinical setting and their 5'-triphosphates (TP) can be potent inhibitors of HCV NS5B polymerase. However, their wide systemic distribution and inefficient hepatic conversion to TP may lead to reduced safety and antiviral activity. The *in vivo* preclinical safety and antiviral activity of the nucleotide prodrugs, B102 were assessed.

[00545] Methods: For pharmacokinetic (PK) and toxicology studies, B102 were orally administered to rats or monkeys at doses from 20 to 300 mg/kg/day up to 14 days. Hepatic nucleoside TP levels were determined by LC-MS/MS. Compounds (10 mg/kg/day) were administered once daily by oral gavage for 4 days in chimpanzees chronically infected with HCV genotype 1. HCV viral loads were monitored before, during and after treatment by quantitative RT-PCR.

[00546] Results: PK studies in rat and monkey revealed that B102 has a first-pass hepatic extraction of >95% with low systemic exposure (<1%). Hepatic TP levels were 10-50-fold higher with nucleotide prodrugs versus a nucleoside counterpart. No toxicity was observed after administration of 50 mg/kg/day of A2 to monkeys for 14 days. No initial emesis or GI toxicity was observed. In HCV-infected chimpanzees, B102 produced a rapid and potent antiviral effect followed by a rebound to baseline after drug discontinuation. Mean viral load reductions ranged from 1.5 log₁₀ with B102 to over 4 days of drug exposure. An equivalent dose of valopicitabine led to a 0.7 log₁₀ viral reduction. No lab abnormalities or evidence of toxicities were observed in chimpanzees.

[00547] Thus, when orally administered, B102 generates high hepatic levels of triphosphates coupled with low systemic exposure, leading to rapid and potent inhibition of HCV replication in chimpanzees, thus demonstrating a promising *in vivo* preclinical safety profile and antiviral activity.

Example 44

[00548] Compounds were tested in an anti-HBV assay. Determination of anti-HBV activity in HBV virions and nucleocapsids via endogenous polymerase assays (EPA)

Drug susceptibility assay using a wild-type HBV producer cell line

1. 12-well collagen-I coated plates were seeded with producer cells expressing wild-type HBV at a density of 0.5×10^6 cells per well in 2 ml growth/selection media.
2. Drug stock solutions were made up freshly in 100% DMSO as 200x stocks. Five additional 4-fold dilutions were prepared from these 200x stock in 100% DMSO. For each experiment, 4 aliquots of each drug dilution series were stored at -20°C until used.
3. Once cells reached confluency (1 day after cells were seeded), drug treatment was initiated by adding 10 µl of drug dilution into 2 ml of fresh growth/selection media. Thus, the final DMSO concentration did not exceed 0.5%. The no-drug control wells received only 10 µl of DMSO in fresh media.
4. Cells were treated every-other-day with 2 ml of fresh drug/medium for a total of 8 days. Cell lysates were then collected on day 10 and subjected to EPA analysis as described below.

Preparation of nucleocapsid-containing lysates for EPA analysis

1. Cells were grown for 3 to 4 days in 12-well collagen-I coated plates until confluent.
2. Media was carefully aspirated and the cell monolayers were rinsed once with 1 ml of PBS.
3. One ml of lysis buffer (50 mM Tris-HCl pH 7.5/150 mM NaCl/5 mM $MgCl_2$ /0.2% NP-40) was added to each well. Plates were stored on ice for 30 min to 4 h.
4. Lysed cells were transferred to 1.5 ml-microfuge tubes.
5. Lysates were clarified by spinning at 14,000 rpm for 5 min at room temperature.
6. Clarified lysates were transferred to fresh tubes and immediately frozen on dry-ice, then stored at $-80^{\circ}C$ until endogenous polymerase assays were performed as described below.

Preparation of secreted virions from supernatant for EPA analysis

1. Cells were grown for 3 to 4 days in 12-well collagen-I coated plates until confluent.
2. Media was carefully aspirated and transferred to 1.5 ml-microfuge tubes.
3. Supernatants were clarified by spinning at 14,000 rpm for 5 min at room temperature.
4. Clarified supernatants were transferred to fresh tubes and immediately frozen on dry-ice, then stored at $-80^{\circ}C$ until endogenous polymerase assays were performed essentially as described below.





Endogenous polymerase assay (EPA) of cell lysates and supernatants






1. EPAs were performed essentially as described by Seifer et al (1998). Intracellular HBV nucleocapsids were immunoprecipitated from the cytoplasmic lysates overnight at $4^{\circ}C$ with a polyclonal rabbit anti-HBcAg antibody and immobilized on protein A sepharose CL-4B beads. Secreted virions were immunoprecipitated from clarified cell supernatants overnight at $4^{\circ}C$ with a monoclonal mouse anti-LS antibody (MA18/7) in the absence of detergent.
2. Following 3 washes of the immobilized capsids or virions with 1 ml of EPA wash buffer (75 mM NH_4Cl , 50 mM Tris-HCl pH 7.4, 1 mM EDTA), endogenous polymerase reactions were initiated by adding 50 μ l of detergent-containing EPA cocktail (50 mM Tris-HCl pH 7.4, 75 mM NH_4Cl , 1 mM EDTA, 20 mM $MgCl_2$, 0.1 mM β -ME, 0.5% NP-40, 100 μ M cold dGTP, TTP, dCTP, and 50 nM ^{33}P -dATP) and

incubated overnight at 37°C. The detergent is required to strip the outer envelope from immunoprecipitated virions as well as to enhance permeability of the nucleocapsids.

3. Following digestion with 1 mg/ml of proteinase K for 1h at 37°C, endogenously ³³P-labeled HBV DNA was liberated via phenol/chloroform extraction.
4. The nucleic acids were precipitated with 1 volume of 5M NH₄-acetate and 2.5 volumes of 100% EtOH, and then separated on a 1% native agarose gel in Tris-borate-EDTA buffer.
5. Gels were blotted onto positively charged nylon membrane overnight at room temperature via capillary transfer in 0.4 N NaOH.
6. Dried membranes were exposed to a PhosphorImager screen (GE Healthcare) overnight at room temperature, then scanned (Storm 860, GE Healthcare) and quantitated with ImageQuant software (GE Healthcare).
7. The 50% effective concentration (EC₅₀) values were calculated from the resulting best-fit equations determined by Xlfit, version 4.1 (IDBS).

[00549] The following results were obtained.

Compound Ref. No.	Structure	Virion EC ₅₀ (μM)	RI EC ₅₀ (μM)
A348 (NM 48)		+++	
A362 (NM 77)		++	++
A616 (NM 128)			++
C819 (NM 177)		++	++


A361 (NM 55)		+	++
A550 (NM 204)		+++	+++
C791		++	
B261			+++
PMEA			+++
L-dT			+++

EC₅₀ in HBV virion and RI is provided as follows:

+++ ≤ 1 μm, ++ > 1-10 μm and + > 10 μm

Example 45: Ethynyl nucleosides for the Treatment of HCV

Exemplary compound syntheses are described below:



29: {9-[(2R)2-Deoxy-2-fluoro-2-C-ethynyl-β-D-erythro-furanosyl]-guanin}-5'-yl-O-(triphenylmethyloxy-*tert*-butyl-S-acyl-2-thioethyl) *H*-phosphonate

[00550] To a stirred solution of **7j** (0.32mmol) and *S*-(2-Phosphite-ethyl) 2,2-dimethyl-3-triphenylmethyloxy-thiopropionate (0.42 mmol) in pyridine (5ml) at -15 °C was added dropwise pivaloyl chloride (0.64 mmol) under nitrogen. The reaction mixture was stirred at -15°C for 2 hours. Dichloromethane and NH₄Cl solution were added. Organic phase was separated washed with NH₄Cl solution, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude material was purified by flash column chromatography (DCM/MeOH) to yield the title compound **Brown powder**. **Molecular Formula** C₃₈H₃₉FN₅O₈PS ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 1.12 (s, 6H), 1.84 (m, 4H) 3.04 (s, 2H), 3.92 (d, *J* = 5.60 Hz, 1H), 4.01-4.10 (m, 3H), 4.33-4.39 (m, 2H), 4.60-4.66 (m, 1H), 6.13 (d, *J* = 18.00 Hz, 1H), 6.67 (s, 2H), 7.21-7.35 (m, 15H), 7.81 (s, 1H), 10.86 (brs, 1H)

30a: *N*-(4-fluoro-benzylaminy)-{9-[(2R)2-deoxy-2-fluoro-2-C-ethynyl-β-D-erythro-furanosyl]-guanin}-5'-yl-O-(triphenylmethyloxy-*tert*-butyl-S-acyl-2-thioethyl) phosphate

[00551] To a stirred solution of **29** (0.088 mmol) in anhydrous carbon tetrachloride (880 μL), 4-fluoro-benzylamine (0.44 mmol) was added dropwise. The reaction mixture was stirred at room temperature for 2 h and evaporated to dryness (bath temperature not exceeding 30 °C). The crude mixture was filtered on a silica gel plug, eluting with a gradient 0-10 % methanol in dichloromethane to yield the title

compound. White solid. Molecular Formula $C_{45}H_{45}F_2N_6O_8PS$ ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 1.09 (s, 6H), 3.03 (s, 2H), 3.39-3.41 (m, 2H), 3.90-3.93 (m, 5H), 4.05-4.08 (m, 1H), 4.20-4.23 (m, 2H), 4.62-4.65 (m, 1H), 5.74 (m, 1H), 6.08-6.14 (dd, $J=17.94$ Hz and $J=4.22$ Hz, 1H), 6.32 (m, 1H), 6.67 (s, 2H), 7.21-7.35 (m, 19H), 7.81 (s, 1H), 10.86 (brs, 1H) ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.83 (s, 1P) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -116.24 (s, 1F), -158.2 (s, 1F) Scan ES^+ 899 (M-H) $^+$, UV λ_{max} 255 nm

31a: *N*-(4-fluoro-benzylaminy)-{9-[(2R)2-deoxy-2-fluoro-2-C-ethynyl- β -D-erythro-furanosyl]-guanine}-5'-yl-*O*-(hydroxy-*tert*-butyl-S-acyl-2-thioethyl) phosphate

[00552] **30a** (0.09 mmol) was dissolved in dichloromethane (320 μ L) and treated with formic acid (32 μ L). The mixture was stirred at room temperature for 10 min, filtered through a solid phase extraction column eluting with a gradient 0-30 % methanol in dichloromethane, then purified by reverse phase (C 18) silica gel column chromatography eluting with a gradient 0-100 % acetonitrile in water and lyophilised from a mixture of water/dioxan to yield the title compound. White solid. Molecular Formula $C_{26}H_{31}F_2N_6O_8PS$ ^1H NMR (d_6 -DMSO, 400 MHz) δ (ppm) 1.09 (s, 6H), 3.03 (s, 2H), 3.39-3.41 (m, 2H), 3.90-3.93 (m, 5H), 4.05-4.08 (m, 1H), 4.20-4.23 (m, 2H), 4.62-4.65 (m, 1H), 4.92 (m, 1H), 5.74 (m, 1H), 6.08-6.14 (dd, $J=17.94$ Hz and $J=4.22$ Hz, 1H), 6.32 (m, 1H), 6.67 (s, 2H), 7.21-7.35 (m, 4H), 7.81 (s, 1H), 10.86 (brs, 1H) ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.66 (s, 1P) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -116.24 (s, 1F), -158.44 (s, 1F) Scan ES^+ 657 (M-H) $^+$, UV λ_{max} 254nm

30b: *N*-(4-methoxy-benzylaminy)-{9-[(2R)2-deoxy-2-fluoro-2-C-ethynyl- β -D-erythro-furanosyl]-guanine}-5'-yl-*O*-(triphenylmethoxy-*tert*-butyl-S-acyl-2-thioethyl) phosphate

[00553] **30b** was synthesized from **29** and 4-methoxy-benzylamine as described for **30a**. White solid. Molecular Formula $C_{46}H_{48}FN_6O_9PS$ ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 1.09 (s, 6H), 3.03 (m, 2H), 3.42 (d, $J=5.02$ Hz, 2H), 3.71 (d, $J=3.60$ Hz, 3H), 3.85-3.90 (m, 5H), 4.06-4.10 (m, 1H), 4.23-4.29 (m, 2H), 4.60-4.66 (m, 1H), 5.54-5.57 (m, 1H), 6.08-6.14 (dd, $J=17.94$ Hz and $J=4.22$ Hz, 1H), 6.28-6.33 (m, 1H), 6.60 (s, 2H), 6.80-6.85 (m, 2H), 7.18-7.20 (m, 2H), 7.23-7.25 (m, 15H), 7.82 (s, 1H), 10.56 (brs, 1H) ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.83 (s, 1P) ^{19}F

NMR (DMSO- d_6 , 235 MHz) δ (ppm) -116.24 (s, 1F), -158.2 (s, 1F) Scan ES⁺ 911 (M-H)⁺, UV $\lambda_{\max}$ 255 nm

31b: *N*-(4-methoxy-benzylaminy)-{9-[(2R)2-deoxy-2-fluoro-2-*C*-ethynyl- β -D-*erythro*-furanosyl]-guanine}-5'-yl-*O*-(hydroxy-*tert*-butyl-*S*-acyl-2-thioethyl) phosphate

[00554] **31b** was synthesized from **30b** as described for **31a**. White solid.

Molecular Formula $C_{27}H_{34}FN_6O_9PS$ ^1H NMR (d_6 -DMSO, 400 MHz) δ (ppm) 1.09 (s, 6H), 3.03 (m, 2H), 3.42 (d, $J=5.02$ Hz, 2H), 3.71 (d, $J=3.60$ Hz, 3H), 3.85-3.90 (m, 5H), 4.06-4.10 (m, 1H), 4.23-4.29 (m, 2H), 4.60-4.66 (m, 1H), 4.92 (t, $J=5.50$ Hz, 1H), 5.54-5.57 (m, 1H), 6.08-6.14 (dd, $J=17.94$ Hz and $J=4.22$ Hz, 1H), 6.28-6.33 (m, 1H), 6.60 (s, 2H), 6.80-6.85 (m, 2H), 7.18-7.20 (m, 2H), 7.82 (s, 1H), 10.56 (brs, 1H), ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.86 (s, 1P) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -158.24 (s, 1F) Scan ES⁺ 669 (M-H)⁺, UV $\lambda_{\max}$ 254nm

30c: *N*-(4-trifluoro-benzylaminy)-{9-[(2R)2-deoxy-2-fluoro-2-*C*-ethynyl- β -D-*erythro*-furanosyl]-guanine}-5'-yl-*O*-(triphenylmethoxy-*tert*-butyl-*S*-acyl-2-thioethyl) phosphate

[00555] **30c** was synthesized from **29** and 4-trifluoromethyl-benzylamine as described for **30a**. White solid. Molecular Formula $C_{46}H_{45}F_4N_6O_8PS$ ^1H NMR (d_6 -DMSO, 400 MHz) δ (ppm) 1.09 (s, 6H), 3.03 (t, $J=6.44$ Hz, 2H), 3.42 (s, 2H), 3.87-3.92 (m, 5H), 4.03-4.08 (m, 1H), 4.24-4.29 (m, 2H), 4.60-4.64 (m, 1H), 5.79-5.82 (m, 1H), 6.08-6.14 (dd, $J=17.94$ Hz and $J=4.22$ Hz, 1H), 6.28-6.33 (m, 1H), 6.60 (s, 2H), 7.23-7.25 (m, 15H), 7.50-7.70 (m, 4H), 8.25 (brs, 1H), 10.76 (brs, 1H), ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) 9.86 (s, 1P) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -158.20 (s, 1F) Scan ES⁺ 949 (M-H)⁺, UV $\lambda_{\max}$ 254nm

31c: *N*-(4-trifluoromethyl-benzylaminy)-{9-[(2R)2-deoxy-2-fluoro-2-*C*-ethynyl- β -D-*erythro*-furanosyl]-guanine}-5'-yl-*O*-(hydroxy-*tert*-butyl-*S*-acyl-2-thioethyl) phosphate

[00556] **31c** was synthesized from **30c** as described for **31a**. White solid.

Molecular Formula $C_{27}H_{31}F_4N_6O_8PS$ ^1H NMR (d_6 -DMSO, 400 MHz) δ (ppm) 1.09 (s, 6H), 3.03 (t, $J=6.44$ Hz, 2H), 3.42 (s, 2H), 3.87-3.92 (m, 5H), 4.03-4.08 (m, 1H), 4.24-4.29 (m, 2H), 4.60-4.64 (m, 1H), 4.91 (brs, 1H), 5.79-5.82 (m, 1H), 6.08-6.14 (dd, $J=17.94$ Hz and $J=4.22$ Hz, 1H), 6.28-6.33 (m, 1H), 6.60 (s, 2H), 7.50-7.70 (m, 4H), 8.25 (brs, 1H), 10.64 (brs, 1H), ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm)

9.86 (s, 1P) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -158.24 (s, 1F) Scan ES $^{+}$ 669 (M-H) $^{+}$, UV λ_{max} 254nm

[00557] Further exemplary compounds synthesized using procedures similar to those described herein are listed below. Note the following names for the compounds synthesized in the examples.

6a: 6-Chloro-9-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]purine

6b: N2-Isobutyryl-9-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine

6c: 1-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]uracile

6d: 1-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]thymine

6e: N4-Benzoyl-1-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]cytosine

6f: 5-Fluoro- 1-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]uracile

6g: 4-Chloro-7-[(2R)-2-deoxy-2-C-ethynyl- 2-fluoro- β -D-erythro-pentofuranosyl]pyrrolo[2,3-d]pyrimidine

7i: 9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]adenine

7j: 9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine

7k: 1-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]cytosine

7l: 4-Amino-7-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]pyrrolo[2,3-d]pyrimidine

11c: 1-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]-uracil-5'-yl-bis(S-pivaloyl-2-thioethylphosphate)




11f: 1-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]-5-fluorouracil-5'-yl-bis(S-pivaloyl-2-thioethylphosphate)




11k: 1-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]-cytosin-5'-yl-bis(S-pivaloyl-2-thioethylphosphate)

11l: 1-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]-4-aminopyrrolo[2,3-d]pyrimidin-5'-yl-bis(S-pivaloyl-2-thioethylphosphate)

16: 9-[(2R)-2,3-Dideoxy-2-C-ethynyl-2-fluoro- β -D-glycero-pentofuranosyl]guanine

- 17: 9-[(2R)-2,3-Dideoxy-2-C-ethynyl-2-fluoro- β -D-glycero-pentofuranosyl]guanine-5'-yl-bis(S-pivaloyl-2-thioethylphosphate)
- 20: 9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine-5'-yl-bis(S-pivaloyl-2-thioethylphosphate)
- 23: 9-[(2R)-2-Deoxy-3,5-di-O-isobutyryl-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine
- 24: N2-Isobutyryl-9-[(2R)-2-Deoxy-3,5-di-O-isobutyryl-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine
- 27i: 9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]adenine 5'-triphosphate sodium salt
- 27j: 9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine 5'-triphosphate sodium salt
- 28: 9-[(2R)-2,3-dideoxy-2-C-ethynyl-2-fluoro- β -D-glycero-pentofuranosyl]guanine 5'-triphosphate sodium salt





3a: 6-Chloro-9-[2-oxo-3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-β-D-ribofuranosyl]purine

[00558] 6-Chloro-[3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-β-D-ribofuranosyl]purine (18.84 mmol) was coevaporated twice with THF then dissolved in anhydrous THF (50 mL). Anhydrous DMSO (119.82 mmol) was added and the resulting solution was cooled down to between -40 °C and -30 °C. Trifluoroacetic anhydride (36.17 mmol) was added dropwise and the solution was stirred between -40 °C and -30 °C for 2 h after which EtN₃ (97.52 mmol) was added. The resulting solution was allowed to warm up to room temperature over 30 min while stirring, then diluted with diethyl ether and washed with H₂O, dried (Na₂SO₄) and evaporated to

dryness. The crude material was purified by column chromatography eluting with 1 % ethyl acetate in dichloromethane. The yellow oil obtained was dissolved in DCM and stirred with an excess of MgSO_4 at room temperature for 36 h, filtered and concentrated under reduced pressure to give the title compound. **Pale yellow foam.** **Molecular Formula** $\text{C}_{22}\text{H}_{35}\text{ClN}_4\text{O}_5\text{Si}_2$. ^1H NMR ($\text{DMSO}-d_6$, 250 MHz) δ (ppm) 9.01 (s, 1H, $H-8$), 8.61 (s, 1H, $H-2$), 6.35 (s, 1H, $H-1'$), 5.35 (d, 1H, $H-3'$, $J_{3',4'} = 9.7$ Hz), 4.31 (m, 1H, $H-4'$), 4.12-4.09 (m, 2H, $H-5'$, $H-5''$), 1.22-0.94 (m, 28H, iPr). **LRFAB-MS (GT):** 527 ($\text{M}+\text{H}$) $^+$, 525 ($\text{M}-\text{H}$) $^-$. UV λ_{max} 263 nm. R_f 0.17 (ethyl acetate/ CH_2Cl_2 , 10/90, v/v).

4a and 4'a: 6-Chloro-9-[3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropyl-disiloxane)-2-*C*-trimethylsilylethynyl- β -D-arabino-furanosyl]purine (**4a**) and 6-chloro-9-[3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropyl-disiloxane)-2-*C*-trimethylsilylethynyl- β -D-ribo-furanosyl]purine (**4'a**)

[00559] Trimethylsilylacetylene (59.20 mmol) was dissolved in anhydrous THF (70 mL). *n*-Butyllithium (37 mL, 1.6 M in hexanes) was added dropwise at -78°C . The reaction mixture was stirred for 30 min at -78°C and then allowed to warm up to -55°C . **3a** (11.84 mmol) in solution in THF (34 mL) was added dropwise at -78°C . The reaction mixture was stirred for 1 h at -78°C and then allowed to warm up to -30°C . The reaction was quenched by careful addition of aqueous saturated NH_4Cl (45 mL) at -78°C . After warming to room temperature, the mixture was diluted with diethyl ether, washed with saturated brine, dried (Na_2SO_4) and concentrated to dryness. The crude material was purified by silica gel chromatography eluting with 20 % Et_2O in petroleum ether to yield two compounds: **4a** (4.62 g, 62 %). **Pale yellow foam.** **Molecular Formula** $\text{C}_{27}\text{H}_{45}\text{ClN}_4\text{O}_5\text{Si}_3$. ^1H NMR ($\text{DMSO}-d_6$, 200 MHz) δ (ppm) 8.81 (s, 1H, $H-8$), 8.64 (s, 1H, $H-2$), 6.64 (s, 1H, $\text{OH}-2'$), 6.33 (s, 1H, $H-1'$), 4.57 (d, 1H, $H-3'$, $J_{3',4'} = 6.6$ Hz), 4.20-3.97 (m, 3H, $H-4'$, $H-5'$ and $H-5''$), 1.20-1.00 (m, 28H, iPr), 0.14 (s, 9H, $\text{Si}(\text{CH}_3)_3$). **LRFAB-MS (GT):** 625 ($\text{M}+\text{H}$) $^+$. R_f 0.72 (ethyl acetate/ CH_2Cl_2 , 10/90, v/v);

and **4'a** (0.75 g, 10 %). **Yellow oil.** **Molecular Formula** $\text{C}_{27}\text{H}_{45}\text{ClN}_4\text{O}_5\text{Si}_3$. ^1H NMR ($\text{DMSO}-d_6$, 200 MHz) δ (ppm) 8.80 (s, 1H, $H-8$), 8.73 (s, 1H, $H-2$), 6.64 (s, 1H, $\text{OH}-2'$), 6.55 (s, 1H, $H-1'$), 4.62 (d, 1H, $H-3'$, $^3J_{3',4'} = 9.1$ Hz), 4.39 (m, 1H, $H-4'$), 4.13 (dd, 1H, $H-5'$, $J_{5',4'} = 3.4$ Hz, $^2J_{5',5''} = 13.2$ Hz), 3.90 (dd, 1H, $H-5''$, $J_{5'',4'} = 2.6$ Hz,

$^2J_{5'',5'} = 13.2$ Hz), 1.15-1.00 (m, 28H, iPr), 0.10 (s, 9H, Si(CH₃)₃). LRFAB-MS (GT): 625 (M+H)⁺. R_f 0.64 (ethyl acetate/CH₂Cl, 10/90, v/v).

5a: 6-Chloro-9-[(2R)-2-deoxy-2-fluoro-3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-C-trimethylsilylethynyl-β-D-erythro-pentofuranosyl]purine

[00560] **4a** (6.78 mmol) was dissolved in dried toluene (31.8 mL) under argon and cooled to -20°C. DAST (40.68 mmol) was added dropwise and the cooling bath was removed after the addition was complete. Stirring was continued for 1.5 hour and the mixture was dissolved with ethyl acetate and poured into saturated NaHCO₃ and stirred for 5 min. The organic layer was washed with saturated brine, dried (Na₂SO₄), concentrated, and purified by silica gel chromatography eluting with 20 % Et₂O in petroleum ether to give the title compound (1.11 g, 26 %). Yellow oil. Molecular Formula C₂₇H₄₄ClN₄O₄Si₃. ¹H NMR (CDCl₃-d₆, 200 MHz,) δ(ppm) 8.79 (s, 1H, H-8), 8.48 (s, 1H, H-2), 6.48 (d, 1H, H-1', J_{1'-F} = 16.0 Hz), 4.74 (dd, 1H, H-3', J_{3'-4'} = 9.4 Hz, J_{3'-F} = 22.4 Hz), 4.36 (d, 1H, H-5', ²J_{5',5''} = 13.4 Hz), 4.20 (m, 1H, H-4'), 4.10 (dd, 1H, H-5'', ²J_{5'',5'} = 13.4 Hz, J_{5'',4'} = 2.6 Hz), 1.30-1.10 (m, 28H, iPr), 0.00 (s, 9H, Si(CH₃)₃). LRFAB-MS (GT): 627 (M+H)⁺. UV λ_{max} 263 nm. R_f 0.24 (diethyl ether/petroleum ether, 30/70, v/v).

6a: 6-Chloro-9-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro-β-D-erythro-pentofuranosyl]purine

[00561] A mixture of **5a** (3.65 mmol) and ammonium fluoride (47.45 mmol) in methanol (12.5 mL) was heated at reflux for 2 h. After cooling down to room temperature, the mixture was concentrated to dryness and purified on silica gel chromatography eluting with a stepwise gradient 2-4 % of methanol in DCM to provide the title compound (0.89 g, 78 %). Yellow solid. Molecular Formula C₁₂H₁₀ClFN₄O₃. ¹H NMR (DMSO-d₆, 200 MHz) δ(ppm) 9.02 (s, 1H, H-8), 8.89 (s, 1H, H-2), 6.55 (d, 1H, H-1', J_{1'-F} = 16.1 Hz), 6.34 (td, 1H, OH-3'), 5.38 (td, 1H, OH-5'), 4.64 (dt, 1H, H-3', J_{3'-4'} = 9.3 Hz, J_{3'-F} = 22.5 Hz), 4.07 (m, 1H, H-4'), 3.83 (m, 2H, H-5', H-5''), 3.76 (d, 1H, ethynyl, ⁴J_{H-F} = 5.3 Hz). ¹³C NMR (DMSO-d₆, 75 MHz,) δ(ppm) 152.0 (C-2), 151.2 (C-4), 149.5 (C-6), 144.7 (C-8), 130.9 (C-5), 95.1 (d, C-2', ¹J_{2'-F} = 182.3 Hz), 88.0 (d, C-1', ²J_{1'-F} = 39.8 Hz), 82.9 (d, CCH, J_{C-F} = 8.2 Hz), 82.5 (C-4'), 75.3 (d, CCH, J_{C-F} = 31.5 Hz), 72.7 (d, C-3', ²J_{3'-F} = 19.5 Hz), 59.0

(C-5'). ^{19}F NMR (DMSO- d_6 , 235 MHz) $\delta(\text{ppm})$ -159.0 (m). LC/MS: (M+H $^+$) 313.1 (8.29 min). UV λ_{max} 262 nm. R_f 0.21 (MeOH/CH $_2$ Cl, 7/93, v/v).

7i: 9-[(2R)2-Deoxy-2-C-ethynyl-2-fluoro- β -D-*erythro*-pentofuranosyl]adenine

[00562] **6a** (2.24 mmol) was dissolved in saturated ammoniacal methanol (80 mL) and heated for 4 h in a steel bomb at 90 °C. After cooling down to room temperature the mixture was coevaporated to dryness and purified by silica gel chromatography eluting with a gradient 5-8 % of methanol in DCM to yield the title compound (305 mg, 46 %). **Yellow solid. Molecular Formula** C $_{12}$ H $_{12}$ FN $_5$ O $_3$. ^1H NMR (DMSO- d_6 , 200 MHz) $\delta(\text{ppm})$ 8.40 (s, 1H, *H*-8), 8.17 (s, 1H, *H*-2), 7.38 (ls, 2H, NH $_2$) 6.35 (d, 1H, *H*-1', $^3J_{1'-F}$ = 17.1 Hz), 6.25 (m, 1H, OH-3'), 5.33 (lt, 1H, OH-5'), 4.68 (m, 1H, *H*-3'), 4.00-3.69 (m, 3 H, *H*-4', *H*-5', *H*-5''), 3.77 (d, 1H, ethynyl, $^4J_{H-F}$ = 5.4 Hz). ^{13}C NMR (DMSO- d_6 , 75 MHz) $\delta(\text{ppm})$ 155.8 (C-4), 152. (C-2), 149.0 (C-6), 138.7 (C-8), 118.5 (C-5), 95.4 (d, C-2', $^1J_{2'-F}$ = 180.8 Hz), 87.6 (d, C-1', $^2J_{1'-F}$ = 40.5 Hz), 82.5 (d, CCH, $^3J_{C-F}$ = 8.0 Hz), 82.0 (C-4'), 74.5 (d, CCH, $^2J_{C-F}$ = 31.0 Hz), 72.8 (d, C-3', $^2J_{3'-F}$ = 19.5 Hz), 59.2 (C-5'). ^{19}F NMR (DMSO- d_6 , 235 MHz) $\delta(\text{ppm})$ -158.0 (t). LC/MS: (M+H $^+$) 294.1 (5.74 min). UV λ_{max} 258 nm. R_f 0.33 (MeOH/CH $_2$ Cl, 15/85, v/v).

4b: *N* ^{2} -Isobutyryl-9-[3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-C-trimethylsilylethynyl- β -D-*arabino*-furanosyl]guanine

[00563] To a suspension of CrO $_3$ (110.76 mmol) in DCM (220 mL) at 0 °C, acetic anhydride (110.76 mmol) and anhydrous pyridine (221.52 mmol) were added. 9-[3,5-O-(1,3-Diyl-1,1,3,3-tetraisopropylidisiloxane)-*ribo*-furanosyl]-*N* ^{2} -isobutyrylguanine (36.92 mmol) in solution in DCM (110 mL) was added dropwise. The cooling bath was removed and the resulting solution stirred at room temperature for 5 h. The reaction mixture was poured into cold ethyl acetate, filtered through a silica and celite gel plug, concentrated to dryness and coevaporated twice with toluene. The residue obtained was dissolved in DCM and stirred with an excess of MgSO $_4$ overnight, filtered and evaporated to get the ketone. The trimethylsilylacetylene (88.60 mmol) was dissolved in anhydrous THF (98 mL) under argon. *n*-Butyllithium (55.4 mL, 1.6 M in hexanes) was added dropwise at - 78 °C. The reaction mixture was stirred for 30 min at - 78 °C and then allowed to warm up to - 55 °C. The ketone in solution in THF (49 mL) was added dropwise at - 78 °C. The reaction mixture was stirred for 1 h at -

78 °C and then allowed to warm up to - 30 °C and stirred for 3 h. The reaction was quenched by careful addition of aqueous saturated NH₄Cl (72 mL) at - 78 °C. After warming to room temperature, the mixture was diluted with ethyl acetate, washed twice with saturated brine, dried (Na₂SO₄) and concentrated to dryness. The crude material was purified using column chromatography eluting with 1.5 % MeOH in dichloromethane to give the title compound (8.59 g, 34 %, 2 steps). Pale yellow foam. Molecular Formula C₃₁H₅₅N₅O₆ Si₃. ¹H NMR (DMSO-*d*₆, 250 MHz) δ(ppm) 12.10 (ls, 1H, NH), 11.69 (ls, 1H, NH), 7.91 (s, 1H, *H*-8), 6.69 (s, 1H, OH), 5.94 (s, 1H, *H*-1'), 4.29 (d, 1H, *H*-3', J_{3',4'} = 5.5 Hz), 3.85-3.95 (m, 3H, *H*-4', *H*-5' and *H*-5''), 2.46 (m, 1H, CH(CH₃)₂), 0.90-1.08 (m, 30H, iPr and CH(CH₃)₃), 0.00 (s, 9H, Si(CH₃)₂). LC/MS: (M+H⁺) 692.4 (24.96 min). UV λ_{max1} 254 nm, λ_{max2} 281 nm. R_f 0.34 (MeOH/CH₂Cl, 15/85, v/v).

5b: *N*²-Isobutyryl-9-[(2R)-2-deoxy-2-fluoro-3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethyl-silylethynyl-β-D-erythro-pentofuranosyl]guanine

[00564] **4b** (2.89 mmol) was dissolved in dried DCM (60 mL) under argon and pyridine (18.06 mmol) was added. The reaction mixture was cooled to -20 °C and DAST (31.35 mmol) was added dropwise. The cooling bath was removed after completion of the addition. Stirring was continued for 1 h 15 and the mixture was dissolved with ethyl acetate and poured into saturated NaHCO₃ and stirred for 5 min. The organic layer was washed with saturated brine, dried (Na₂SO₄), concentrated, and purified by silica gel chromatography eluting with ethyl acetate in DCM (2 %) to give the title compound (1.41 g, 70 %). **Yellow oil. Molecular Formula C₃₁H₅₄FN₅O₅ Si₃.** ¹H NMR (DMSO-*d*₆, 250 MHz) δ(ppm) 12.22 (s, 1H, NH), 8.09 (s, 1H, *H*-8), 6.21 (d, 1H, *H*-1', J_{1',F} = 15.6 Hz), 4.54 (dd, 1H, *H*-3', J_{3',F} = 23.6 Hz, J_{3',4'} = 9.8 Hz), 4.33 (m, 1H, *H*-5', ²J_{5',5''} = 13.1 Hz), 4.16 (m, 1H, *H*-5''), 2.81 (m, 1H, CH(CH₃)₂), 1.13-1.03 (m, 34 H, iPr and CH(CH₃)₃), 0.08 (s, 9H, Si(CH₃)₃, ³J_{H-H} = 6.9 Hz). ¹⁹F NMR (DMSO-*d*₆, 235 MHz) δ(ppm) -160.26 (dd, J_{F-1'} = 16.1 Hz, J_{F-3'} = 23.3 Hz). LC/MS: (M+H⁺) 694.7 (24.02 min). LRFAB-MS (GT): 694 (M+H)⁺, 692 (M-H)⁻. UV λ_{max} 256 nm. R_f 0.46 (MeOH/CH₂Cl, 05/95, v/v).

6b: *N*²-Isobutyryl-9-[(2R)-2-deoxy-2-*C*-ethynyl-2-fluoro-β-D-erythro-pentofuranosyl]guanine.

[00565] **5b** (1.89 mmol) was dissolved in methanol (13.8 mL) and ammonium fluoride (24.54 mmol) was added. The resulting solution was stirred at reflux for 1 h and evaporated to dryness. The crude material was purified on silica gel chromatography eluting with a stepwise gradient 6-10 % of methanol in DCM to yield the title compound (344 mg, 48 %). **Pale yellow oil. Molecular Formula $C_{16}H_{20}FN_5O_4Si_3$.** ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 12.18 (ls, 1H, NH), 11.77 (ls, 1H, NH), 8.34 (s, 1H, H-8), 6.29 (d, 1H, OH-3', $J_{OH-3'} = 7.5$ Hz), 6.20 (d, 1H, H-1', $J_{1'-F} = 16.2$ Hz), 5.39 (t, 1H, OH-5', $J_{OH-5'} = 5.1$ Hz), 4.52 (dt, 1H, H-3', $J_{3'-F} = 22.9$ Hz), 3.98 (m, 1H, H-4'), 3.90-3.85 (m, 2H, H-5' and ethynyl), 3.72 (m, 1H, H-5''), 2.52 (m, 1H, CH(CH₃)₂), 1.14 (d, 6 H, CH(CH₃)₂, $^3J_{H-H} = 6.9$ Hz). ^{13}C NMR (DMSO- d_6 , 100 MHz) δ (ppm) 180.7 (C-6), 155.3 (C-2), 148.9 (C-4), 137.3 (C-8), 120.4 (C-5), 95.8 (d, C-2', $^1J_{2'-F} = 182.1$ Hz), 87.7 (d, C-1', $^2J_{1'-F} = 39.2$ Hz), 83.4 (d, CCH, $^3J_{C-F} = 9.1$ Hz), 82.6 (C-4'), 75.9 (d, CCH, $^2J_{C-F} = 31.2$ Hz), 72.9 (d, C-3', $^2J_{3'-F} = 19.1$ Hz), 59.3 (C-5'). ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -158.9 (m). LC/MS: (M+H⁺) 380.3 (8.34 min). UV λ_{max} 256 nm, R_f 0.40 (MeOH/CH₂Cl, 15/85, v/v).

7j: 9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine

[00566] **6b** (1.33 mmol) was dissolved in saturated methanolic ammonia (62 mL) and stirred at room temperature for 20 h. The reaction mixture was then evaporated to dryness under reduced pressure. The residue was dissolved in water and washed twice with ethyl acetate. The aqueous layer was evaporated and purified on reverse phase column chromatography (C 18) eluting with a gradient 2-15 % of acetonitrile in water. The residue obtained was then dissolved in hot ethyl acetate, filtered and dried to give the title compound (134 mg, 33 %). **Yellow solid. Molecular Formula $C_{12}H_{12}FN_5O_4$.** ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 10.70 (ls, 1H, NH), 7.98 (s, 1H, H-8), 6.60 (ls, 2H, NH₂), 6.21 (d, 1H, OH-3', $J_{OH-3'} = 7.6$ Hz), 5.83 (d, 1H, H-1', $J_{1'-F} = 16.9$ Hz), 5.29 (t, 1H, OH-5', $J_{OH-5'} = 5.2$ Hz), 4.50 (td, 1H, H-3', $J_{3'-F} = 22.8$ Hz, $J_{3'-4'} = 9.2$ Hz), 3.93-3.81 (m, 3H, H-4', H-5' and ethynyl), 3.70 (m, 1H, H-5''). ^{13}C NMR (DMSO- d_6 , 100 MHz) δ (ppm) 157.2 (C-6), 154.3 (C-2), 151.05 (C-4), 135.1 (C-8), 116.7 (C-5), 96.4 (d, C-2', $^1J_{C-F} = 182.1$ Hz), 87.4 (d, C-1', $^2J_{C-F} = 39.2$ Hz), 83.1 (d, CCH, $J_{C-F} = 9.1$ Hz), 82.4 (C-4'), 76.2 (d, CCH, $^2J_{C-F} = 31.2$ Hz), 73.2 (d, C-3', $^2J_{C-F} = 20.1$ Hz), 59.5 (C-5'). ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -158.5 (m). LC/MS (A): (M+H⁺) 310.1 (5.55 min). LRFAB-MS (GT): 619 (2M+H)⁺, 310 (M+H)⁺, 152 (B+H)⁺, 617 (2M-H)⁻, 308 (M-H)⁻. UV λ_{max} 253 nm

3c: 1-[2-Oxo-3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)- β -D-ribo-furanosyl]uracile

[00567] **3c** was synthesized from 1-[3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-ribo-furanosyl]uracile as described for **3a**. Pale yellow foam. Molecular Formula $C_{21}H_{36}N_2O_7Si_2$ ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 11.58 (ls, 1H, NH), 7.74 (d, 1H, *H*-6, $J_{6-5} = 8.0$ Hz), 5.68 (d, 1H, *H*-5, $J_{5-6} = 8.0$ Hz), 5.45 (s, 1H, *H*-1'), 4.97 (d, 1H, *H*-3', $J_{3'-4'} = 9.2$ Hz), 4.06-3.90 (m, 3H, *H*-4', *H*-5'), 1.14-0.87 (m, 28H, iPr). LR LC/MS: (M+H⁺) 485.1 (M-H⁻) 483.1 (5.53 min). UV λ_{max} 262 nm. R_f 0.40 (MeOH/CH₂Cl, 05/95, v/v).

4c: 1-[3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethylsilylethynyl- β -D-arabino-furanosyl]uracile Yoshimura, Y.; Iino, T.; Matsuda, A. *Tetrahedron Lett.* 1991, 32, 6003-6006.

[00568] Molecular Formula $C_{26}H_{46}N_2O_7Si_3$ ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 11.35 (ls, 1H, NH), 7.44 (d, 1H, *H*-6, $J_{6-5} = 8.0$ Hz), 6.54 (s, 1H, OH), 6.02 (s, 1H, *H*-1'), 5.54 (d, 1H, *H*-5, $J_{6-5} = 8.0$ Hz), 4.13-3.93 (m, 3H, *H*-3', *H*-5'), 3.75 (m, 1H, *H*-4'), 1.03-0.96 (m, 28H, iPr), 0.00 (s, 9H, Si(CH₃)₃). ^{13}C NMR (DMSO- d_6 , 100 MHz) δ (ppm) 163.4 (*C*-4), 150.8 (*C*-2), 141.6 (*C*-6), 103.6 (CCSi), 101.2 (*C*-5), 92.5 (CCSi), 87.3 (*C*-1'), 80.9 (*C*-4'), 77.9 (*C*-2'), 75.9 (*C*-3'), 61.8 (*C*-5'), 17.7-17.1 (8C, 4 SiC(CH₃)₂), 13.3-12.6 (4C, 4 SiC(CH₃)₂), 0.2 (3C, Si(CH₃)₃). LR LC/MS: (M+H⁺) 583.2 (M-H⁻) 581.2 (6.72 min). UV λ_{max} 261 nm. R_f 0.27 (Ethyl acetate/CH₂Cl, 10/90, v/v).

5c: 1-[(2R)-2-Deoxy-2-fluoro-3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethylsilylethynyl- β -D-erythro-pentofuranosyl]uracile

[00569] **5c** was synthesized from **4c** as described for **5a**. Yellow oil. Molecular Formula $C_{27}H_{49}FN_2O_6Si_3$ ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 11.62 (sl, 1H, NH), 7.43 (dl, 1H, *H*-6, $J_{6-5} = 8.0$ Hz), 6.12 (d, 1H, *H*-1', $J_{1'-F} = 16.8$ Hz), 5.68 (d, 1H, *H*-5, $J_{5-6} = 8.0$ Hz), 4.22-3.85 (m, 4H, *H*-3', *H*-4', *H*-5'), 1.16-1.00 (m, 28H, iPr), 0.00 (s, 9H, Si(CH₃)₃). ^{19}F NMR (DMSO- d_6 , 376 MHz) δ (ppm) -159.7. LR LC/MS: (M+H⁺) 585.2 (M-H⁻) 583.3 (6.47 min). UV λ_{max} 261 nm. R_f 0.52 (Ethyl acetate/CH₂Cl, 15/85, v/v).

6c: 1-[(2R)-2-Deoxy-2-*C*-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]uracile

[00570] A mixture of **5c** (0.56 mmol) and ammonium fluoride (7.31 mmol) were dissolved in methanol (10mL) stirred at reflux for 1 h and evaporated to dryness. The resulting residue was purified on silica gel flash column chromatography eluting with a gradient 0-20 % methanol in DCM and then, on reverse phase column chromatography eluting with a gradient 0-100 % acetonitrile in water to give the desired product which was lyophilised from water (47 mg, 31 %). **White lyophilised powder. Molecular Formula** $C_{11}H_{11}FN_2O_5$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 11.49 (sl, 1H, NH), 7.87 (d, 1H, *H*-6, $J_{6-5} = 8.0$ Hz), 6.18 (d, 1H, OH-3', $J_{OH-3'} = 7.2$ Hz), 6.10 (d, 1H, *H*-1', $J_{1'-F} = 18.0$ Hz), 5.69 (d, 1H, *H*-5, $J_{5-6} = 8.0$ Hz), 5.32 (m, 1H, OH-5'), 4.19-4.10 (m, 2H, *H*-3' and ethynyl), 3.85-3.75 (m, 2H, *H*-4' *H*-5'), 3.60 (m, 1H, *H*-5''). ^{13}C NMR (DMSO- d_6 , 100 MHz) δ (ppm) 163.3 (C-4), 150.6 (C-2), 140.1 (C-6), 102.5 (C-5), 95.5 (d, C-2', $J_{2'-F} = 186.1$ Hz), 87.1 (d, C-1', $^2J_{1'-F} = 40.2$ Hz), 83.2 (d, CCH, $^2J_{C-F} = 8.0$ Hz), 82.1 (C-4'), 76.5 (d, CCH, $^4J_{C-F} = 30.1$ Hz), 73.3 (d, C-3', $^2J_{C-F} = 19.1$ Hz), 58.7 (C-5'). ^{19}F NMR (DMSO- d_6 , 376 MHz) δ (ppm) -158.2. **LR LC/MS:** (M+H⁺) 271.1 (M-H⁻) 269.2 (1.12 min). **HRFAB-MS** $C_{11}H_{12}O_5N_2F$. (M+H⁺) calculated 271.0730, found 271.0739. UV λ_{max} 261 nm. R_f 0.33 (MeOH/CH₂Cl, 20/80, v/v).

2d: **1-[3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)- β -D-ribo-furanosyl]thymine**

[00571] The 1-(β -D-ribo-furanosyl)thymine (40.9 mmol) was dissolved in pyridine (435ml) and the mixture was cooled down to 0°C with an ice-bath for 25 minutes. Then, TIPSCl₂ (16.2ml) was added and after complete addition, the mixture was allowed to warm up to room temperature. The reaction mixture was stirred at room temperature for 3hrs, diluted with dichloromethane and water, washed with a saturated aqueous solution of NaHCO₃. The organic phases were combined, dried over Na₂SO₄, filtered and evaporated. The residue was coevaporated with toluene to remove pyridine. The resulting residue was purified by flash column chromatography eluting with a gradient 0-2 % of methanol in dichloromethane to give the title compound. **Off-white powder. Molecular Formula** $C_{22}H_{40}N_2O_7Si_2$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 0.94-1.04 (m, 28H), 1.73 (s, 3H), 3.86-3.96 (m, 1H), 4.06-4.13 (m, 2H), 4.14-4.20 (m, 1H), 5.44-5.48 (m, 1H), 5.53 (brs, 1H), 5.77 (brs, 1H) 7.42 (s, 1H), 11.35 (brs, 1H). UV λ_{max} 212 nm, 266 nm.

3d: 1-[2-Oxo-3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)- β -D-ribo-furanosyl]thymine

[00572] To a suspension of CrO_3 (60 mmol) in dichloromethane (200 mL) at 0 °C, acetic anhydride (59 mmol) and anhydrous pyridine (120 mmol) were added. **2d** (20 mmol) in solution in DCM was added dropwise. The cooling bath was removed and the resulting solution stirred at room temperature for 3 h. The reaction mixture was poured into cold ethyl acetate, filtered through a silica and celite gel plug, concentrated to dryness and coevaporated twice with toluene to give the title compound. Colorless oil. Molecular Formula $\text{C}_{22}\text{H}_{38}\text{N}_2\text{O}_7\text{Si}_2$

4d: 1-[3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethylsilylethynyl- β -D-arabino-furanosyl]thymine

[00573] **4d** was synthesized from **3d** and trimethylacetylene as described for **4a**. Brown solid. Molecular Formula $\text{C}_{27}\text{H}_{48}\text{N}_2\text{O}_7\text{Si}_3$. Scan $\text{ES}^+ 597 (\text{M}+\text{H})^+$, UV λ_{max} 265 nm.

5d: 1-[(2R)-2-Deoxy-2-fluoro-3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethyl-silylethynyl- β -D-erythro-pentofuranosyl]thymine

[00574] **5d** was synthesized from **4a** as described for **5a**. Brown solid. Molecular Formula $\text{C}_{27}\text{H}_{47}\text{FN}_2\text{O}_6\text{Si}_3$. ^1H NMR (CDCl_3 - d_6 , 400 MHz) δ (ppm) 0.1 (s, 9H), 1.05-1.14 (m, 28H), 1.92 (s, 3H), 3.99-4.13 (m, 1H), 4.44-4.9 (m, 3H), 6.35 (d, 1H, $J = 16.44$ Hz), 7.2 (s, 1H), 8.86 (s, 1H). Scan $\text{ES}^+ 599 (\text{M}+\text{H})^+$,

6d: 1-[(2R)-2-Deoxy-2-*C*-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]thymine

[00575] **6d** was synthesized from **5d** as described for **7i**. Molecular Formula $\text{C}_{12}\text{H}_{13}\text{FN}_2\text{O}_5$. ^1H NMR (CDCl_3 - d_6 , 400 MHz) δ (ppm) 1.75 (s, 3H), 3.6-3.65 (m, 1H), 3.82-3.84 (m, 2H), 4.07 (d, 1H, $J = 5.27$ Hz), 4.19 (m, 1H), 5.4 (brs, 1H), 6.08 (d, 1H, $J = 17.8$ Hz), 6.17 (brs, 1H), 7.8 (s, 1H), 11.46 (brs, 1H). Scan $\text{ES}^+ 285 (\text{M}+\text{H})^+$, UV λ_{max} 266 nm.

4e: N^4 -Benzoyl-1-[3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethylsilylethynyl- β -D-arabino-furanosyl]cytosine

[00576] **4e** was synthesized from **3e** and trimethylacetylene as described for **4a**. Brown solid. Molecular Formula $\text{C}_{33}\text{H}_{51}\text{N}_3\text{O}_7\text{Si}_3$. Scan $\text{ES}^+ 686 (\text{M}+\text{H})^+$, UV λ_{max} 260 nm, 310 nm.

5e: *N*⁴-benzoyl-1-[(2*R*)-2-Deoxy-2-fluoro-3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethyl-silylethynyl-β-*D*-erythro-pentofuranosyl]cytosine

[00577] **5e** was synthesized from *N*⁴-benzoyl-1-[3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethyl-silylethynyl-*D*-arabino-furanosyl]cytosine as described for **5a**. Yellow solid. Molecular Formula C₃₀H₄₂FN₃O₆Si₂. Scan ES⁺ 688 (M+H)⁺, UV λ_{max} 260 nm, 310 nm.

6e: *N*⁴-benzoyl-1-[(2*R*)-2-Deoxy-2-*C*-ethynyl-2-fluoro-β-*D*-erythro-pentofuranosyl]cytosine

[00578] **6e** was synthesized from **5e** as described for **7i**. White powder. Molecular Formula C₁₈H₁₆FN₃O₅. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm) 3.63-3.69 (m, 1H), 3.82-.393 (m, 2H), 4 (d, 1H, *J* = 5.27 Hz), 4.13-4.24 (m, 1H), 5.38 (brs, 1H), 6.23-6.28 (m, 2H), 7.32-7.36 (m, 1H), 7.49-7.53 (m, 2H), 7.6-7.64 (m, 1H), 7.99-8.01 (m, 2H), 8.34 (d, 1H, *J* = 7.32 Hz), 11.30 (brs, 1H). Scan ES⁺ 374 (M+H)⁺, UV λ_{max} 262 nm, 303 nm.

7k: 1-[(2*R*)-2-Deoxy-2-*C*-ethynyl-2-fluoro-β-*D*-erythro-pentofuranosyl]cytosine

[00579] Molecular Formula C₁₁H₁₂FN₃O₄. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm) 3.57-3.62 (m, 1H), 3.77-3.80 (m, 2H), 3.95 (d, 1H, *J* = 5.53 Hz), 4.03-4.16 (m, 1H), 5.2 (brs, 1H), 5.73 (d, 1H, *J* = 7.19 Hz), 6.06 (d, 1H, *J* = 7.19 Hz), 6.14-6.25 (m, 1H), 7.17-7.3 (2brs, 2H), 7.74 (d, 1H, *J* = 7.74 Hz). Scan ES⁺ 270 (M+H)⁺, UV λ_{max} 271 nm.

8k: *N*⁴-dimethoxytrityl-1-[(2*R*)-2-Deoxy-2-*C*-ethynyl-2-fluoro-β-*D*-erythro-pentofuranosyl]cytosine

[00580] To a stirred solution of **7k** (2.34 mmol) in pyridine (7.2 ml) was added trimethylsilyl chloride (9.36 mmol) at room temperature. The reaction mixture was stirred at room temperature for 2 hours. 4-dimethylamino pyridine (1.17 mmol) and dimethoxytrityl chloride (3.51 mmol) were then added. The mixture was stirred at room temperature for 16 hours. The reaction mixture was diluted with DCM and sat NaHCO₃ solution. The organic phase was washed twice with sat NaHCO₃ solution, dried over Na₂SO₄, filtered and evaporated. The crude material was dissolved in a solution NH₄OH/dioxin (2:1) and stirred for 4hrs. Solvent was evaporated and the residue purified by silica gel chromatography (DCM/EtOH) to yield the title

compound. White foam. Molecular Formula $C_{32}H_{30}FN_3O_6$. Scan ES^- 570 (M+H) $^-$, UV λ_{max} 277 nm

9k: 1-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]-4-N-dimethoxytrityl-cytosin-5'-yl-bis(S-pivaloyl-2-thioethylphosphate

[00581] To a stirred solution of **8k** (0.35 mmol) in anhyd THF/tetrazole solution (1.05 mmol) was added bis(S-pivaloyl-2-thioethyl) N,N-diisopropylphosphoramidite (0.42 mmol) at 0°C. The reaction mixture was stirred at room temperature for 3 hrs. The reaction mixture was cooled down to 0°C and tert-butyl hydroperoxide (0.7 ml/mmol) was added. The reaction mixture was stirred at room temperature for 2 hours. The mixture was diluted with DCM, neutralized with sat $Na_2S_2O_3$ solution. The organic phase was washed twice with H_2O , extracted, dried over Na_2SO_4 , filtered and evaporated. The crude material was purified by silica gel chromatography (DCM/EtOH) to yield the title compound. Glassy compound. Molecular Formula $C_{47}H_{59}FN_3O_{11}PS_2$. Scan ES^+ 938 (M+H) $^+$, UV λ_{max} 277 nm

11k: 1-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]-cytosin-5'-yl-bis(S-pivaloyl-2-thioethylphosphate

[00582] **9k** (34 mmol) was stirred in AcOH/MeOH/ H_2O (3/6/1) solution for 2 hrs and at 50°C for 4 hours. The reaction mixture was then evaporated and purified by silica gel chromatography (DCM/EtOH) to yield the title compound. White lyophilized powder. Molecular Formula $C_{25}H_{37}FN_3O_9PS_2$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 1.17 (s, 18H), 3.08-3.11 (t, J = 6.07 Hz, 4H), 3.99-4.08 (m, 7H), 4.22-4.28 (m, 2H), 5.73-5.75 (d, J = 7.30 Hz, 1H), 6.30 (brs, 2H), 7.26-7.31 (d, J = 17.30 Hz, 2H), 7.47-7.48 (d, J = 7.30 Hz, 1H) ^{19}F NMR (DMSO- d_6 , 376 MHz) δ (ppm) -156.48 (s, 1F) ^{31}P NMR (DMSO- d_6 , 162 MHz) δ (ppm) -1.96 (s, 1P). Scan ES^+ 638 (M+H) $^+$, UV λ_{max} 271 nm

12: N²-Methoxytrityl-9-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine

[00583] To a stirred solution of **7i** (5.18 mmol) in pyridine (7 ml/mmol) was added trimethylsilyl chloride at room temperature. The mixture was stirred at room temperature for 6 hours. Methoxytrityl chloride (6.21 mmol) was then added and the reaction mixture was stirred at room temperature for 16 hours and 2 hours with NH_4OH (4 ml/mmol). The mixture was diluted with ethyl acetate, washed with H_2O , sat $NaHCO_3$ solution and sat $NaCl$ solution, dried over Na_2SO_4 , filtered and

evaporated. The crude material was purified by silica gel chromatography (DCM/MeOH) to yield the title compound. Yellowish oil. Molecular Formula $C_{32}H_{28}FN_5O_5$.

13: *N*²-Methoxytrityl-9-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro-5-O-*tert*-butyldimethylsilyl- β -D-*erythro*-pentofuranosyl]guanine

[00584] To a stirred solution of **12** (2.29 mmol) in pyridine (5 ml) at 0°C, was added *tert*-butyldimethylsilyl chloride (2.75 mmol). The reaction mixture was stirred at room temperature for 24 hours. It was then diluted in DCM and washed twice with H₂O. The organic phase was extracted, dried over Na₂SO₄, filtered and evaporated. The crude material was purified by silica gel chromatography (DCM/MeOH) to yield the title compound. Yellowish oil. Molecular Formula $C_{38}H_{44}FN_5O_5Si$. Scan ES⁺ 696 (M+H)⁺, λ_{max} 260 nm. Scan ES⁻ 694 (M+H)⁻, UV λ_{max} 260 nm

14: *N*²-Methoxytrityl-9-[(2R)-2,3-dideoxy-2-C-ethynyl-2-fluoro-5-O-*tert*-butyldimethylsilyl- β -D-*glycero*-pentofuranosyl]guanine

[00585] To a stirred solution of **13** (0.14 mmol) in acetonitrile (47 ml/mmol) was added 4-dimethylamino pyridine (0.56 mmol) and phenyl chlorothionoformate (0.43 mmol) at room temperature. The reaction mixture was stirred at room temperature for 16 hours and was concentrated under reduced pressure. The residue obtained was dissolved in DCM, the organic phase was washed with H₂O, HCl (1N), dried over Na₂SO₄, filtered evaporated and co-evaporated with toluene.

[00586] The crude material was dissolved in toluene (12 ml/mmol), azo-bis-isobutyronitrile (0.02 mmol) and tributylstannane (0.24 mmol) were added at room temperature. The reaction mixture was stirred at 125°C for 2 hours and concentrated under reduced pressure. The crude material was purified by silica gel chromatography (DMC/MeOH) to yield the title compound. Yellowish oil. Molecular Formula $C_{38}H_{44}FN_5O_4Si$. Scan ES⁺ 680 (M+H)⁺, UV λ_{max} 260 nm

15: *N*²-Methoxytrityl-9-[(2R)-2,3-dideoxy-2-C-ethynyl-2-fluoro- β -D-*glycero*-pentofuranosyl]guanine

[00587] **14** (0.35 mmol) was dissolved in MeOH (20 ml/mmol). Ammonium fluoride (3.55 mmol) was then added at room temperature and the reaction mixture was stirred at 70°C for 2 hours. After concentration under reduced pressure, the crude material was purified by silica gel chromatography (DCM/MeOH) to yield the title

compound. Beige foam. Molecular Formula $C_{32}H_{30}FN_5O_4$. ^1H NMR ($CDCl_3-d_6$, 400 MHz) δ (ppm) 2.38-2.45 (m, 2H), 2.75 (brs, 2H), 3.64-3.67 (d, $J=12.20$ Hz, 2H), 3.77 (s, 4H), 4.20-4.23 (d, $J=11.7$ Hz, 1H), 4.41-4.42 (d, $J=8.4$ Hz, 1H), 5.83-5.87 (d, $J=16.24$ Hz, 1H), 6.80-6.82 (d, $J=8.12$ Hz, 4H), 7.26-7.31 (m, 11H), 7.84 (brs, 1H), 9.26 (brs, 1H)

16: 9-[(2R)-2,3-Dideoxy-2-C-ethynyl-2-fluoro- β -D-glycero-pentofuranosyl]guanine

[00588] **15** (0.09 mmol) was stirred in AcOH/THF/H₂O (3/6/1) solution at 50°C for 1 day. The reaction mixture was then concentrated under reduced pressure and purified by silica gel chromatography, C18 (H₂O/ACN) Beige lyophilised powder. Molecular Formula $C_{12}H_{12}FN_5O_5$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 2.57-2.74 (m, 2H), 3.56 (s, 1H), 3.61-3.64 (d, $J=12.10$ Hz, 1H), 3.79-3.82 (d, $J=12.10$ Hz, 1H), 3.91-3.93 (d, $J=5.40$ Hz, 1H), 4.32-4.35 (m, 1H), 5.25 (s, 1H), 6.06-6.10 (d, $J=18.20$ Hz, 1H), 6.64 (s, 1H), 8.01 (s, 1H), 10.82 (s, 1H) ^{19}F NMR (DMSO- d_6 , 376 MHz) δ (ppm) -138.4 (s, 1F). Scan ES⁻ 292 (M+H)⁻, Scan ES⁺ 316 (M+Na)⁺, UV λ_{max} 251 nm

17: 9-[(2R)-2,3-Dideoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]-guanine-5'-yl-bis(S-pivaloyl-2-thioethylphosphate)

[00589] **17** was synthesized from **14** (0.35 mmol) as described for **9**. The crude material was then stirred in AcOH/THF/H₂O (4/2/1) at 50°C, for 3 hours. The reaction mixture was concentrated under reduced pressure and purified by silica gel chromatography (DCM/MeOH) to yield the title compound. Beige solid. Molecular Formula $C_{26}H_{37}FN_5O_8PSi_2$ ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 2.48 (s, 18H), 2.65-2.68 (m, 2H), 3.06-3.10 (q, $J=3.71$ Hz, and $J=6.02$ Hz, 4H), 3.97-4.04 (m, 5H), 4.31-4.35 (m, 2H), 4.50-4.52 (m, 1H), 6.13-6.18 (d, $J=17.60$ Hz, 1H), 6.63 (s, 2H), 7.82 (s, 1H), 10.85 (s, 1H). ^{19}F NMR (DMSO- d_6 , 376 MHz) δ (ppm) -139.2 (s, 1F). Scan ES⁺ 662 (M+H)⁺, UV λ_{max} 254 nm. HPLC (0-100% ACN over a period of 8 min) $t_R=5.65$ min.

18: N^2 -Methoxytrityl-9-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro-5-O-tert-butylidimethylsilyl-3-O-tetrahydropyranyl- β -D-erythro-pentofuranosyl]guanine

[00590] To a stirred solution of **13** (0.8 mmol), in anhyd THF (20 ml/mmol), at room temperature, was added p-toluen sulfonic acid (0.12 mmol) and dihydropyran (2 ml/mmol). The reaction mixture was stirred at room temperature for 3 days and

neutralized with TEA. The mixture was diluted with DCM, washed twice with H₂O. The organic phase was dried over Na₂SO₄, filtered and evaporated. The crude material was purified by silica gel chromatography (DCM/MeOH) to yield the title compound.
Molecular Formula C₄₃H₅₂FN₅O₆Si. Scan ES⁺ 780 (M+H)⁺

19: *N*²-Methoxytrityl-9-[(2*R*)-2-deoxy-2-*C*-ethynyl-2-fluoro-3-*O*-tetrahydropyranyl-β-*D*-erythro-pentofuranosyl]guanine

[00591] **19** was synthesized from **18**, as described for **15**. **Molecular Formula** C₃₇H₃₈FN₅O₆. Scan ES⁺ 666 (M+H)⁺.

21: *N*²-Tetrahydropyranyl-9-[(2*R*)-2-deoxy-2-*C*-ethynyl-2-fluoro-5-*O*-tert-butyltrimethylsilyl-3-*O*-tetrahydropyranyl-β-*D*-erythro-pentofuranosyl]guanine

[00592] **21** was obtained from the purification of **18**. **Molecular Formula** C₂₈H₄₂FN₅O₆Si. Scan ES⁺ 592 (M+H)⁺, UV λ_{max} 273 nm

22: *N*²-Tetrahydropyranyl-9-[(2*R*)-2-deoxy-2-*C*-ethynyl-2-fluoro-3-*O*-tetrahydropyranyl-β-*D*-erythro-pentofuranosyl]guanine

[00593] **22** was synthesized from **21** (0.46 mmol), as described for **15**. **Molecular Formula** C₂₂H₂₈FN₅O₆

20: 9-[(2*R*)-2-Deoxy-2-*C*-ethynyl-2-fluoro-β-*D*-erythro-pentofuranosyl]guanin-5'-yl-bis(*S*-pivaloyl-2-thioethylphosphate

[00594] **20** was synthesized from **19** (0.09 mmol), as described for **9k**. The crude material was then stirred at room temperature in AcOH/THF/H₂O (4/2/1) solution overnight. The reaction mixture was concentrated under reduced pressure and purified by silica gel chromatography (DCM/MeOH) to yield the title compound. **White lyophilized powder**. **Molecular Formula** C₂₆H₃₇FN₅O₉PS₂ ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 1.16 (s, 20H), 3.06-3.09 (m, 4H), 3.90-3.91 (d, *J* = 5.40 Hz, 1H), 3.99-4.10 (q, *J* = 6.70 Hz and *J* = 7.00 Hz, 4H), 4.32-4.38 (m, 2H), 4.63 (m, 1H), 6.10-6.14 (d, *J* = 16.93 Hz, 1H), 6.69 (s, 2H), 7.79 (s, 1H), 10.96 (s, 1H) ³¹P NMR (DMSO-*d*₆, 162 MHz) δ (ppm) -1.91 (s, 1P) ¹⁹F NMR (DMSO-*d*₆, 376 MHz) δ (ppm) -156.82 (s, 1F) Scan ES⁺ 678 (M+H)⁺.

25: 1-[(2*R*)-2-Deoxy-2-fluoro-3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethylsilylethynyl-β-*D*-erythro-pentofuranosyl]-4-thiouracile

[00595] **5c** (820 mg, 1.40 mmol) was dissolved in anhydrous 1,2-dichloroethane (35 mL) and treated with Lawesson's reagent (1.13 g, 2.80 mmol). The reaction

mixture was stirred at reflux overnight and evaporated to dryness. The resulting residue was filtered on a silica gel plug eluting with a gradient 0-5 % of ethyl acetate in dichloromethane to give the title compound. **Yellow oil. Molecular Formula** $C_{26}H_{45}FN_2O_5SSi_3$ LR LC/MS: (M+H⁺) 601.3 (M-H⁻) 599.3 (7.03 min). UV λ_{max} 332 nm. R_f 0.71 (Ethyl acetate/CH₂Cl, 7/93, v/v).

26: 1-[(2R)-2-Deoxy-2-fluoro-3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-C-ethynyl- β -D-erythro-pentofuranosyl]cytosine

[00596] Crude **25** was dissolved in saturated ammoniacal methanol (9 mL). The resulting solution was heated by micro-waves at 120 °C for 20 min and concentrated under reduced pressure to give the title compound. **Oily residue. Molecular Formula** $C_{26}H_{46}FN_3O_5Si_3$ LR LC/MS (B): (M+H⁺) 512.3 (M-H⁻) 510.3 (5.33 min). UV λ_{max1} 242 nm, λ_{max2} 273 nm.

27i: 9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]adenine 5'-triphosphate sodium salt

[00597] To a solution of **7i** (0.286 mmol) in triethylphosphate (750 μ L), phosphoryl chloride (75 μ L, 0.807 mmol) was added at 0 °C. This reaction mixture A was stirred overnight at 5 °C. Tributylammonium pyrophosphate (PPI/Bu₃N 1/1,5, 1g, 2.19 mmol) was dissolved in anhydrous DMF (2 mL). Tributylamine (420 μ L, 1.76 mmol) was added to the PPI and the resulting mixture was stirred for 15 min at 0 °C. 2.4 mL of this solution were added to the reaction mixture A. The reaction mixture was stirred at 0 °C for 1 min. The reaction was carefully quenched with TEAB 1M (pH = 7,5, 10 mL), stirred 20 min at 0 °C, then diluted with water and ethyl acetate. The aqueous phase was concentrated under reduced pressure. The crude material was subjected to DEAE-Sephadex chromatography eluting with a gradient 10⁻³-1 M of TEAB). The desired fractions were combined, concentrated under reduced pressure and coevaporated with a mixture of water/methanol, and finally coevaporated with water. The resulting residue was purified on semipreparative HPLC. Fractions containing the expected product were concentrated under reduced pressure, coevaporated with a mixture of water/methanol and lyophilised from water. The triethylammonium salt triphosphate was eluted three times with water on a Dowex Na⁺ resin column to yield after lyophilisation from water to the sodium salt.

[00598] Molecular Formula $C_{12}H_{11}FN_5O_{12}P_3 \cdot 3Na$. ^1H NMR (D_2O , 300 MHz) δ (ppm) 8.31 (s, 1H, *H*-8), 8.14 (s, 1H, *H*-2), 6.28 (d, 1H, *H*-1', $^3J_{1'-F} = 15.6$ Hz), 4.64 (m, 1H, *H*-3'), 4.42 (m, 1H, *H*-5'), 4.35-4.25 (m, 2H, *H*-4' and *H*-5''), 2.82 (d, 1H, ethynyl, $^4J_{H-F} = 5.5$ Hz). ^{31}P NMR (D_2O , 121 MHz) δ (ppm) -10.27 (d, 1P, P_γ , $J_{P_\gamma-P\beta} = 19.4$ Hz), -11.03 (d, 1P, P_α , $J_{P_\alpha-P\beta} = 19.4$ Hz), -22.38 (t, 1P, P_β , $J_{P\beta-P_\gamma} = J_{P\beta-P_\alpha} = 19.4$ Hz). ^{19}F NMR (D_2O , 282 MHz) δ (ppm) -160.0 (m). LRFAB-MS (GT): 600 ($M+H$)⁺, 578 ($M-Na+2H$)⁺, 556 ($M-2Na+3H$), 598 ($M-H$)⁻, 576 ($M-Na$)⁻, 554 ($M-2Na+H$)⁻, 532 ($M-3Na+2H$)⁻.

27j: 9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine 5'-triphosphate sodium salt

[00599] **27j** was synthesized from **7i** as described for **27**. Molecular Formula $C_{12}H_{11}FN_5O_{13}P_3 \cdot 3Na$. ^1H NMR (D_2O , 400 MHz) δ (ppm) 7.97 (s, 1H, *H*-8), 6.19 (d, 1H, *H*-1', $^3J_{1'-F} = 16.0$ Hz), 4.70 (m, 1H under H_2O , *H*-3'), 4.39 (m, 1H, *H*-5'), 4.29-4.22 (m, 2H, *H*-4' and *H*-5''), 2.98 (d, 1H, ethynyl, $^4J_{H-F} = 5.0$ Hz). ^{31}P NMR (D_2O , 162 MHz): -10.50 (d, 1P, P_γ , $J_{P_\gamma-P\beta} = 19.4$ Hz), -11.03 (d, 1P, P_α , $J_{P_\alpha-P\beta} = 19.4$ Hz), -22.38 (t, 1P, P_β , $J_{P\beta-P_\gamma} = J_{P\beta-P_\alpha} = 19.4$ Hz). ^{19}F NMR ($DMSO-d_6$, 376 MHz) δ (ppm) -159.1 (m). LRFAB-MS (GT): 638 ($M+Na$)⁺, 616 ($M+H$)⁺, 594 ($M-Na+2H$)⁺, 572 ($M-2Na+3H$)⁺, 550 ($M-3Na+4H$)⁺, 592 ($M-Na$)⁻, 570 ($M-2Na+H$)⁻, 548 ($M-3Na+2H$)⁻.

2g: 4-Chloro-7-[3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)- β -D-ribo-furanosyl]pyrrolo[2,3-d]pyrimidine

[00600] **2g** was synthesized from 9-[β -D-ribo-furanosyl]-7-deaza-6-chloropurine, as described for intermediate 12. Yellow oil. Molecular Formula $C_{23}H_{38}ClN_3O_5Si_2$. ^1H NMR ($DMSO-d_6$, 400 MHz) δ (ppm) 0.96-1.04 (m, 28H), 3.92-3.95 (m, 3H), 4.41-4.58 (m, 2H), 5.65 (s, 1H), 6.08 (s, 1H), 6.71 (s, 1H), 7.83 (s, 1H), 8.62 (s, 1H)

3g: 4-Chloro-7-[2-oxo-3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)- β -D-erythro-pentofuranosyl]pyrrolo[2,3-d]pyrimidine

[00601] **3g** was synthesized from **2g** as described for **3d**. Brown solid. Molecular Formula $C_{23}H_{36}ClN_3O_5Si_2$. Scan ES^+ ($M+H$)⁺ 528, UV λ_{max} 271 nm

4g: 4-Chloro-[3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-C-trimethylsilylethynyl- β -D-arabino-furanosyl]pyrrolo[2,3-d]pyrimidine

[00602] **4g** was synthesized from **3g** as described for **4a**. Beige solid. Molecular Formula : $C_{28}H_{46}ClN_3O_5Si_3$. ^1H NMR ($DMSO-d_6$, 400 MHz) δ (ppm), 0.12 (s, 9H),

0.95-1.09 (m, 28H), 3.90-3.94 (m, 1H), 4.02-4.03 (m, 2H), 4.37-4.39 (d, $J=6.74$ Hz, 1H), 6.43 (s, 1H), 6.44 (s, 1H), 6.68 (d, $J=3.71$ Hz, 1H), 7.71-7.72 (d, $J=3.84$ Hz, 1H), 8.66 (s, 1H)

5g: 4-Chloro-7-[(2R)2-deoxy-2-fluoro-3,5-*O*-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-*C*-trimethylsilylethynyl- β -D-erythro-pentofuranosyl]pyrrolo[2,3-*d*]pyrimidine

[00603] **5g** was synthesized from **4g** as described for **5a**. Yellow oil. Molecular formula $C_{29}H_{45}ClFN_3O_5Si_3$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 0.33 (s, 9H), 1.02-1.13 (m, 28H), 4.0-4.03 (d, $J=13.42$ Hz, 1H), 4.12-4.14 (d, $J=9.43$ Hz, 1H), 4.27-4.31 (d, $J=14.00$ Hz, 1H), 4.71 (brs, 1H), 6.58-6.62 (d, $J=17.07$ Hz, 1H), 6.82-6.83 (d, $J=3.80$ Hz, 1H), 7.72 (d, $J=3.80$ Hz, 1H), 8.69 (s, 1H) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -159.6 (s, 1F)

6g: 4-Chloro-7-[(2R)2-deoxy-2-*C*-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]pyrrolo[2,3-*d*]pyrimidine

[00604] **6g** was synthesized from **5g** as described for **6a**. Yellow oil. Molecular Formula $C_{13}H_{11}ClFN_3O_3$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 3.60-3.65 (d, $J=5.44$ Hz, 1H), 3.68-3.71 (d, $J=12.35$ Hz, 1H), 3.85-3.88 (d, $J=12.35$ Hz, 1H), 3.95-3.97 (d, $J=8.90$ Hz, 1H), 4.46-4.54 (dd, $J=23.23$ Hz and $J=9.39$ Hz, 1H), 5.38 (s, 1H), 6.28 (s, 1H), 6.57-6.61 (d, $J=16.47$ Hz, 1H), 6.79 (d, $J=3.82$ Hz, 1H), 8.04 (d, $J=3.78$ Hz, 1H), 8.70 (s, 1H) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -158.30 (s, 1F). Scan ES⁺ 312 (M+H)⁺. Scan ES⁻ 356 (M+HCO₂)⁻.

7l: 4-Amino-7-[(2R)2-deoxy-2-*C*-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]pyrrolo[2,3-*d*]pyrimidine

[00605] **7l** was synthesized from **6g** as described for **6a**. White lyophilised powder. Molecular Formula $C_{13}H_{13}FN_4O_3$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 3.61 (d, $J=5.52$ Hz, 1H), 3.63-3.67 (m, 1H), 3.80-3.83 (d, $J=12.14$ Hz, 1H), 3.86-3.88 (d, $J=9.38$ Hz, 1H), 4.46-4.54 (dd, $J=23.23$ Hz and $J=9.39$ Hz, 1H), 5.30 (brs, 1H), 6.1 (brs, 1H), 6.41-6.47 (d, $J=16.47$ Hz, 1H), 6.57-6.61 (d, $J=16.47$ Hz, 1H), 7.04 (s, 2H), 7.37-7.38 (d, $J=3.65$ Hz, 1H), 8.05 (s, 1H) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -157.15 (s, 1F) Scan ES⁺ 293 (M+H)⁺, UV λ_{max} 275 nm

23: 9-[(2R)2-Deoxy-3,5-di-*O*-isobutyryl-2-*C*-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]guanine

[00606] A solution of 9-[(2R)-2-Deoxy-2-C-ethynyl-2-fluoro-β-D-*erythro*-furanosyl]guanine (0.16 mmol), 4-dimethylaminopyridine (0.01 mmol), triethylamine (0.48 mmol) and isobutyric anhydride (0.48 mmol), in acetonitrile (1ml) was stirred at room temperature for 6 hours. The reaction mixture was hydrolysed with a NaHCO₃ saturated solution. Ethyl acetate was added. The organic phase was separated, washed with NaCl saturated solution, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude material was purified by flash column chromatography (DCM/EtOH) to yield the title compound **White powder**. **Molecular Formula** C₂₀H₂₄FN₅O₆ ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 1.02-1.22 (m, 12H), 2.53-2.59 (m, 1H), 2.65-2.70 (m, 1H), 4.04 (d, *J* = 4.77 Hz, 1H), 4.35-4.40 (m, 3H), 5.88-5.94 (dd, *J* = 9.39 Hz and *J* = 8.21 Hz, 1H), 6.21-6.25 (d, *J* = 17.28 Hz, 1H), 6.58 (s, 2H), 7.09 (s, 1H), 10.82 (s, 1H). **Scan ES⁺** 450.0 (M+H)⁺, UV λ_{max} 251 nm

24: N-2-Isobutyryl-9-[(2R)-2-deoxy-3,5-di-O-isobutyryl-2-C-ethynyl-2-fluoro-β-D-*erythro*-pentofuranosyl]guanine

[00607] **24** was obtained from the purification of **23**. **White powder**. **Molecular Formula** C₂₄H₃₀FN₅O₇ ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 1.02-1.22 (m, 18H), 2.53-2.59 (m, 1H), 2.65-2.70 (m, 1H), 2.74-2.80 (m, 1H), 4.04 (d, *J* = 4.90 Hz, 1H), 4.35-4.40 (m, 3H), 5.73-5.80 (dd, *J* = 10.14 Hz and *J* = 7.80 Hz, 1H), 6.29-6.34 (d, *J* = 17.36 Hz, 1H), 8.23 (s, 1H), 11.80 (brs, 1H), 12.3 (brs, 1H). **Scan ES⁺** 520 (M+H)⁺, UV λ_{max} 257 nm

4f: 5-Fluoro-1-[3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-C-trimethylsilylethynyl-β-D-*arabino*-furanosyl]uracile

[00608] **4f** was synthesized from **3f** as described for **4a**. **Orange solid**. **Molecular Formula** C₂₆H₄₅FN₂O₇Si₃ ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 0.13 (s, 9H), 0.94-1.06 (m, 28 H), 3.75 (m, 1H), 3.96-4.09 (m, 3H), 5.99 (d, *J* = 1.53 Hz, 1H), 6.53 (s, 1H), 7.58-7.60 (d, *J* = 6.76 Hz, 1H), 11.8 (brs, 1H) **Scan ES⁻** 599 (M-H)⁻, UV λ_{max} 271 nm

5f: 5-Fluoro-1-[(2R)-2-deoxy-2-fluoro-3,5-O-(1,3-diyl-1,1,3,3-tetraisopropylidisiloxane)-2-C-trimethyl-silylethynyl-β-D-*erythro*-pentofuranosyl]uracile

[00609] **5f** was synthesized from **4f** as described for **5a**. **White solid**. **Molecular Formula** C₂₆H₄₄F₂N₂O₆Si₃. ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm) 0.13 (s, 9H),

0.94-1.06 (m, 28 H), 3.92-3.95 (d, $J = 12.47$ Hz, 1H), 3.96-4.09 (m, 1H), 4.21 (d, $J = 12.22$ Hz, 1H), 5.20 (brs, 1H), 6.10-6.15 (d, $J = 16.53$ Hz, 1H), 7.56 (s, 1H), 12.23 (brs, 1H) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -160.06 (s, 1F), -165.94 (s, 1F) Scan ES⁺ 603 (M-H)⁺, UV λ_{max} 272 nm

6f: 5-Fluoro-1-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]uracile

[00610] **6f** was synthesized from **5f** as described for **6a**. **White solid. Molecular Formula** $\text{C}_{11}\text{H}_{10}\text{F}_2\text{N}_2\text{O}_5$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm) 3.61-3.64 (d, $J = 12.55$ Hz, 1H), 3.81-3.85 (m, 2H), 4.11 (d, $J = 4.91$ Hz, 1H), 4.14-4.20 (m, 1H), 5.50 (s, 1H), 6.04-6.09 (d, $J = 16.91$ Hz, 1H), 6.20 (d, $J = 7.64$ Hz, 1H), 8.29 (d, $J = 7.09$ Hz, 1H), 12.05 (s, 1H) ^{19}F NMR (DMSO- d_6 , 235 MHz) δ (ppm) -158.74 (s, 1F), -166.27 (s, 1F) Scan ES⁺ 289.0 (M-H)⁺, UV λ_{max} 270 nm.

11f: 1-[(2R)-2-deoxy-2-C-ethynyl-2-fluoro- β -D-erythro-pentofuranosyl]-5-fluorouracil-5'-yl-bis(S-pivaloyl-2-thioethylphosphate)

[00611] **11f** was synthesized from **6f** as described **9k**. **White solid. Molecular Formula** $\text{C}_{25}\text{H}_{35}\text{F}_2\text{N}_2\text{O}_{10}\text{P}_2\text{S}_2$. ^1H NMR (DMSO- $d_6 + D_2O$, 400 MHz) δ (ppm) 1.15-1.17 (m, 18H), 3.10 (t, $J = 6.40$ Hz, 4H), 4-4.08 (m, 5H), 4.19 (d, $J = 5.39$ Hz, 1H), 4.24-4.39 (m, 3H), 6.12 (d, $J = 16.82$ Hz, 1H), 6.39 (d, $J = 6.07$ Hz, 1H), 7.86 (brs, 1H), 12.12 (brs, 1H).

28: 9-[(2R)-2,3-dideoxy-2-C-ethynyl-2-fluoro- β -D-glycero-pentofuranosyl]guanine 5'-triphosphate sodium


[00612] **28** was synthesized from **16** as described for **27i**. **White powder. Molecular Formula** $\text{C}_{12}\text{H}_{12}\text{FN}_5\text{Na}_3\text{O}_{12}\text{P}_3$. ^1H NMR (D_2O , 400 MHz) δ (ppm) 2.61-2.72 (m, 2H), 2.95-2.96 (m, 1H), 4.16-4.22 (m, 1H), 4.35-4.40 (m, 1H), 4.6-4.7 (m, 1H), 6.17 (d, $J = 16$ Hz, 1H), 8.02 (s, 1H). ^{19}F NMR (D_2O , 235 MHz) δ (ppm) (-138.95)-(-138.74) (m, 1F), ^{31}P NMR (D_2O , 162 MHz) δ (ppm) -10.66 (d, $J = 19.44$ Hz, 1P), -11.14 (d, $J = 19.44$ Hz, 1P), -22.82 (t, $J = 19.44$ Hz, 1P). Scan ES⁺ 599.6 (M-3Na)³⁺, UV λ_{max} 253 nm.

[00613] All publications and patent, applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. While the claimed subject matter has been described in terms of various embodiments, the

skilled artisan will appreciate that various modifications, substitutions, omissions, and changes may be made without departing from the spirit thereof. Accordingly, it is intended that the scope of the subject matter limited solely by the scope of the following claims, including equivalents thereof.

What is claimed is:

1. A compound of formula



or a pharmaceutically acceptable salt, a stereoisomeric, tautomeric or polymorphic form thereof, wherein;

R^y is optionally substituted alkyl, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, cycloalkenyl, hydroxyalkyl, amino, heterocyclyl or heteroaryl;

R^a and R^b are selected as follows:


i) R^a and R^b are each independently hydrogen or optionally substituted alkyl, carboxyalkyl, hydroxyalkyl, hydroxyarylalkyl, acyloxyalkyl, aminocarbonylalkyl, alkoxycarbonylalkyl, aryl, arylalkyl, cycloalkyl, heteroaryl or heterocyclyl; or

ii) R^a and R^b together with the nitrogen atom on which they are substituted form a 3-7 membered heterocyclic or heteroaryl ring; and

R¹ is a moiety derivable by removal of a hydrogen from a hydroxy group of an anti-viral drug,


and wherein optionally, when R^y is *tert*-butyl or hydroxy-*tert*-butyl, then R¹ is not 3'-azido-2',3'-dideoxythymidine.

2. The compound of claim 1 having the formula:




or a pharmaceutically acceptable salt, a stereoisomeric, tautomeric or polymorphic form thereof.

3. The compound of claim 1 having the formula:




or a pharmaceutically acceptable salt, a stereoisomeric, tautomeric or polymorphic form thereof.

4. The compound of claim 1 having the formula:



or a pharmaceutically acceptable salt, , a stereoisomeric, tautomeric or polymorphic form thereof.


5. The compound of claim 1 having the formula:



wherein R^2 and R^3 are each independently H, or R^2 and R^3 are linked to form a cyclic group by an alkyl, ester or carbamate linkage;

or a pharmaceutically acceptable salt, a stereoisomeric, tautomeric or polymorphic form thereof.


6. The compound of claim 1 having the formula:



wherein R^d is hydrogen, methyl or methoxy;

each R^L is independently H, carbamyl, straight chained, branched or cyclic alkyl; acyl; CO-alkyl, CO-aryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonate ester, a lipid, an amino acid; an amino acid residue; or a carbohydrate; or a pharmaceutically acceptable salt, a stereoisomeric, tautomeric or polymorphic form thereof.

7. The compound of claim 1 having the formula:



or a pharmaceutically acceptable salt, a stereoisomeric, tautomeric or polymorphic form thereof.


8. The compound of any of the previous claims wherein R^y is substituted alkyl and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl.


9. The compound of claim 7 wherein R^y is hydroxyalkyl or aminoalkyl.

10. The compound of any of the previous claims wherein R^y is $-\text{OR}^c$, $-\text{C}(\text{R}^c)_3$ or $-\text{NHR}^c$ where each R^c is independently alkyl, substituted alkyl, aryl or substituted aryl; and R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl.

11. The compound of any of the previous claims wherein R^a and R^b are independently hydrogen, benzyl or substituted alkyl.
12. The compound of any of the previous claims wherein R^y is selected from the group consisting of alkyl and hydroxyalkyl.
13. The compound of any of the previous claims wherein R^y is –
 $C(CH_3)_2CH_2OH$.
14. The compound of any of the previous claims wherein R^2 and R^3 are each hydrogen, R^a is hydrogen, R^b is benzyl and R^y is $-C(CH_3)_2CH_2OH$.
15. The compound of claim 1, wherein R^1 is selected from ribavirin, viramidine, valopicitabine, PSI-6130 and MK-0608.
16. The compound of claim 1, wherein R^1 is selected from resiquimod and celgosivir.
17. The compound of claim 1, wherein R^1 is selected from lamivudine, entecavir, telbivudine, racivir, emtricitabine, clevudine, amdoxovir, valtorcitabine, tenofovir and Adefovir


18. The compound of claim 1 selected from:






and

19. The compound of claim 1 having the structure:




20. A method for the treatment of a host infected with a *Flaviviridae* virus or hepatitis B virus, comprising administering an effective treatment amount of a compound of any of claims 1, 2, 3, 4 or 5.

21. The method of claim 20, wherein the virus is hepatitis C.
22. The method of claim 21, wherein the host is a human.
23. The method of claim 21 wherein the compound is



or



or a pharmaceutically acceptable salt or stereoisomer thereof.

24. The method of claim 21, wherein said administration directs a substantial amount of said compound or pharmaceutically acceptable salt or stereoisomer thereof to the liver of said host.

25. The method of claim 21, wherein said compound or pharmaceutically acceptable salt or stereoisomer thereof is administered in combination or alternation with a second anti-viral agent optionally selected from the group consisting of an interferon, a ribavirin, an interleukin, a NS3 protease inhibitor, a cysteine protease inhibitor, a phenanthrenequinone, a thiazolidine derivative, a thiazolidine, a benzanilide, a helicase inhibitor, a polymerase inhibitor, a nucleotide analogue, a gliotoxin, a cerulenin, an antisense phosphorothioate oligodeoxynucleotide, an inhibitor of IRES-dependent translation, and a ribozyme.

26. The method of claim 24, wherein said compound or pharmaceutically acceptable salt or stereoisomer thereof is administered in combination or alternation with a second anti-viral agent, optionally selected from the group consisting of an interferon, a ribavirin, an interleukin, a NS3 protease inhibitor, a cysteine protease inhibitor, a phenanthrenequinone, a thiazolidine derivative, a thiazolidine, a benzanilide, a helicase inhibitor, a polymerase inhibitor, a nucleotide analogue, a gliotoxin, a cerulenin, an antisense phosphorothioate oligodeoxynucleotide, an inhibitor of IRES-dependent translation, and a ribozyme.

27. The method of claim 26, wherein the second agent is selected from the group consisting of pegylated interferon alpha 2a, interferon alfacon-1, natural


interferon, albuferon, interferon beta-1a, omega interferon, interferon alpha, interferon gamma, interferon tau, interferon delta and interferon γ -1b.

28. The method of claim 24, wherein the host is a human.

29. The method of claim 28, wherein said administration directs a substantial amount of said compound or pharmaceutically acceptable salt or stereoisomer thereof to the liver of said host.

30. The method of claim 20 comprising treating a human host infected with hepatitis B virus.


31. The method of claim 30, wherein the compound is of the formula



wherein R^y is hydroxyalkyl and optionally is $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{OH}$;

wherein R^a and R^b are independently hydrogen, alkyl, substituted alkyl, benzyl or substituted benzyl; and

wherein optionally the compound is




and wherein optionally said compound or pharmaceutically acceptable salt thereof is administered in combination or alternation with a second anti-viral agent optionally selected from interferon alfa-2b, Peginterferon alfa-2a, lamivudine, entecavir, telbivudine, racivir, emtricitabine, clevudine, amdoxovir, valtorcitabine, tenofovir and adefovir.

32. The method of claim 31, wherein said administration directs a substantial amount of said compound or pharmaceutically acceptable salt thereof to the liver of said host.


33. A method of treatment of a hepatitis C or B infection, comprising administering to an individual in need thereof a treatment effective amount of a phosphoramidate or phosphonoamidate derivative of a nucleoside or nucleoside analogue or a pharmaceutically acceptable salt or stereoisomer thereof.

34. A pharmaceutical composition comprising a compound of any of claims 1, 2, 3, 4 or 5 and a pharmaceutically acceptable excipient, carrier or diluent.

35. The composition of claim 34, wherein the compound is







or




or a pharmaceutically acceptable salt or stereoisomer thereof.

36. The composition of claim 34, wherein the compound is selected from





and



or a pharmaceutically acceptable salt or stereoisomer thereof.

37. The composition of claim 34, wherein the composition is an oral formulation.


38. The composition of claim 36, wherein the composition is an oral formulation.

39. A use of compound of any of claims 1, 2, 3, 4 or 5 for the treatment of a host infected with a *Flaviviridae* virus or hepatitis B virus.

40. The use of claim 39, wherein the virus is hepatitis C.

41. The use of claim 40, wherein the host is a human.

42. The use of claim 39, wherein the compound is



or a pharmaceutically acceptable salt or stereoisomer thereof.

43. The use of claim 39, wherein a substantial amount of said compound or pharmaceutically acceptable salt or stereoisomer thereof is directed to the liver of said host.

44. The use of claim 39, wherein said compound or pharmaceutically acceptable salt or stereoisomer thereof is administered in combination or alternation with a second anti-viral agent optionally selected from the group consisting of an interferon, a ribavirin, an interleukin, a NS3 protease inhibitor, a cysteine protease inhibitor, a phenanthrenequinone, a thiazolidine derivative, a thiazolidine, a benzanilide, a helicase inhibitor, a polymerase inhibitor, a nucleotide analogue, a gliotoxin, a cerulenin, an antisense phosphorothioate oligodeoxynucleotide, an inhibitor of IRES-dependent translation, and a ribozyme.

45. The use of claim 44, wherein said compound or pharmaceutically acceptable salt or stereoisomer thereof is administered in combination or alternation

with a second anti-viral agent, selected from the group consisting of an interferon, a ribavirin, an interleukin, a NS3 protease inhibitor, a cysteine protease inhibitor, a phenanthrenequinone, a thiazolidine derivative, a thiazolidine, a benzanilide, a helicase inhibitor, a polymerase inhibitor, a nucleotide analogue, a gliotoxin, a cerulenin, an antisense phosphorothioate oligodeoxynucleotide, an inhibitor of IRES-dependent translation, and a ribozyme.

46. The use of claim 45, wherein the second agent is selected from the group consisting of pegylated interferon alpha 2a, interferon alphacon-1, natural interferon, albuferon, interferon beta-1a, omega interferon, interferon alpha, interferon gamma, interferon tau, interferon delta and interferon γ -1b.

47. The use of claim 43, wherein the host is a human.

48. The use of claim 47, wherein said administration directs a substantial amount of said compound or pharmaceutically acceptable salt or stereoisomer thereof to the liver of said host.

49. The use of claim 39, wherein the treatment comprises administering the compound to a human host infected with hepatitis B virus.

50. The use of claim 49, wherein said compound or pharmaceutically acceptable salt thereof is administered in combination or alternation with a second anti-viral agent optionally selected from interferon alfa-2b, Peginterferon alfa-2a, lamivudine, entecavir, telbivudine, racivir, emtricitabine, clevudine, amdoxovir, valtorcitabine, tenofovir and adefovir.

51. The use of claim 50, wherein a substantial amount of said compound or pharmaceutically acceptable salt thereof is directed to the liver of said host.

52. A use of a phosphoramidate or phosphonoamidate derivative of a nucleoside or nucleoside analogue or a pharmaceutically acceptable salt or stereoismer thereof for treatment of a hepatitis C or B infection.

53. A use of the compound of any of claims 1-19 for manufacturing a medicament for treating of a host infected with a *Flaviviridae* virus or hepatitis B virus.

54. The use of claim 53, wherein the virus is hepatitis C.