

Description

Title of Invention: TETRAHYDROQUINOLINE DERIVATIVES AS P2X7 RECEPTOR ANTAGONISTS

Technical Field

[0001] This invention relates to tetrahydroquinoline derivatives that act as modulators of the P2X7 receptor. The present invention also relates to processes for the preparation of the compounds, pharmaceutical compositions containing the compounds, and to their use in the treatment of a wide range of diseases, syndromes, and disorders, which are associated with P2X7 receptor activity such as diseases of the autoimmune and inflammatory system, diseases of the nervous and neuro-immune system, diseases involved with neuroinflammation of the Central Nervous System (CNS) or diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems.

Background Art

[0002] The P2X7 receptors (P2RX7) belong to the family of P2X ionotropic receptors that are activated by extracellular nucleotides, in particular adenosine triphosphate (ATP). P2X7 receptor is distinguished from other P2X family members by the high concentrations (mM range) of ATP required to activate it and by its ability to form a large pore upon prolonged or repeated stimulation (NPL 1 to NPL 3: North, R. A., *Physiol. Rev.* 2002, 82(4), 1013-67; Surprenant, A., Rassendren, F. et al., *Science* 1996, 272(5262), 735-8; Virginio, C., MacKenzie, A. et al., *J. Physiol.*, 1999, 519, 335-46). The P2X7 receptor is a ligand-gated ion channel and is present on a variety of cell types, largely those known to be involved in the inflammatory and/ or immune process, specifically, macrophages and monocytes in the periphery and predominantly in glial cells (microglia and astrocytes) of the CNS. (NPL 4 to NPL 6: Duan and Neary, *Glia* 2006, 54, 738-746; Skaper et al, *FASEB J* 2009, 24, 337-345; Surprenant and North, *Annu. Rev. Physiol.* 2009, 71, 333-359).

[0003] P2X7 receptor is located on many cell types, especially ones known to be involved in inflammatory and immune processes. Activation of the P2X7 receptor by extracellular nucleotides, in particular adenosine triphosphate, leads to the release of proinflammatory cytokines IL-1 beta and IL-18 (NPL 7: Muller, et al., *Am. J. Respir. Cell Mol. Biol.* 2011, 44, 456-464), giant cell formation (macrophages/ microglial cells), degranulation (mast cells) and L-selectin shedding (lymphocytes) (NPL 8 to NPL 9: Ferrari et al., *J. Immunol.* 2006, 176, 3877-3883; Surprenant and North, *Annu. Rev. Physiol.* 2009, 71, 333-359). P2X7 receptors are also located on antigen-presenting cells (keratinocytes, salivary acinar cells (parotid cells)), hepatocytes, erythrocytes, erythroleukaemic cells, monocytes, fibroblasts, bone marrow cells, neurones, and renal

mesangial cells.

[0004] The importance of P2X7 in the nervous system arises primarily from experiments using P2X7 knockout mice. These mice demonstrate the role of P2X7 in the development and maintenance of pain, as these mice are protected from the development of both adjuvant-induced inflammatory pain and partial nerve ligation-induced neuropathic pain (NPL 10: Chessell et al., *Pain* 2005, 114, 386-396). In addition, P2X7 knockout mice also exhibit an anti-depressant phenotype based on reduced immobility in forced swim and tail suspension tests (NPL 11: Basso et al., *Behav. Brain Res.* 2009, 198, 83-90). Moreover, the P2X7 pathway is linked to the release of the proinflammatory cytokine, IL-1 beta, which has been linked to precipitation of mood disorders in humans (NPL 12 to NPL 13: Dantzer, *Immunol. Allergy Clin. North Am.* 2009, 29, 247-264; Capuron and Miller, *Pharmacol. Ther.* 2011, 130, 226-238). In addition, in murine models of Alzheimer's disease, P2X7 was upregulated around amyloid plaques indicating a role of this target in such pathology as well (NPL 14: Parvathenani et al., *J. Biol. Chem.* 2003, 278, 13309-13317).

[0005] There is therapeutic rationale for the use of P2X7 ion channel blockers in the treatment of a variety of disease states. These include but are not limited to diseases associated with the central nervous system such as stroke or injury and diseases associated with neuro-degeneration and neuroinflammation such as Alzheimer's disease, Huntington's disease, epilepsy, Amyotrophic lateral sclerosis, acute spinal cord injury additionally to meningitis, sleep disorders, mood and anxiety disorders as well as chronic and neuropathic and inflammatory pain. Furthermore, peripheral inflammatory disorders and autoimmune diseases including but not limited to rheumatoid arthritis, osteoarthritis, psoriasis, allergic dermatitis, asthma, chronic obstructive pulmonary disease, airways hyper-responsiveness, septic shock, bronchitis, glomerulonephritis, irritable bowel disease, skin injury, lung emphysema, Limb girdle dystrophy type 2B, fibrosis, Syndrome of synovitis Acne Pustulosis, atherosclerosis, burn injury, spinal cord injury, Hyperostosis Osteitis, Crohn's disease, ulcerative colitis, growth and metastases of malignant cells, myoblastic leukaemia, diabetes, trauma, meningitis, osteoporosis, burn injury, ischemic heart disease, and varicose veins and trauma, are all examples where the involvement of P2X7 channels has been implicated. In addition, a recent report suggests a link between P2X7 receptor and chronic, inflammatory and neuropathic pain (NPL 15: Chessell, I. P., Hatcher, J. P. et al., *Pain*, 2005, 114(3), 386-96). Overall, these findings indicate a role for the P2X7 receptor in the process of neuronal synaptic transmission and therefore a potential role for P2X7 antagonists as novel therapeutic tools to treat neuropathic pain.

[0006] In view of the above observations, there is significant requirement for P2X7 antagonists that can be efficiently used in the treatment of a wide range of diseases,

syndromes, and disorders, which are associated with P2X7 receptor activity such as diseases of the autoimmune and inflammatory system, diseases of the nervous and neuro-immune system, diseases involved with neuroinflammation of the CNS or diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems.

- [0007] Several reviews on small molecule inhibitors of P2X7 which have been published are: NPL 16: Guile, S.D., et al., J. Med. Chem, 2009, 52, 3123-3141; and NPL 17: Gunosewoyo, H. and Kassiou, M., Exp Opin, 2010, 20, 625-646.
- [0008] International patent application PTL 1: WO 2013/108227 purportedly describes aza-bicyclic pyridine derivatives as a P2X7 receptor antagonist. The chemical structures are dihydrofuopyridine derivatives and dihydropyranopyridine derivatives, which are quite different from tetrahydroquinoline derivatives of the present invention. They neither disclose nor suggest teterhydroquinoline derivatives.
- [0009] Recently, PTL 2: WO 2016/039983 and PTL3: WO 2016/019228 also disclose aza-bicyclic compounds with P2X7 receptor antagonistic activities. Each chemical structure is triazolopyrazine derivative and indolizine derivative, respectively, which is quite different from a tetrahydroquinoline derivative of the present invention.

Citation List

Patent Literature

- [0010] {PTL 1} WO 2013/108227
 {PTL 2} WO 2016/039983
 {PTL 3} WO 2016/019228

Non Patent Literature

- [0011] {NPL 1} North, R. A., Physiol. Rev. 2002, 82(4), 1013-67
 {NPL 2} Surprenant, A., Rassendren, F. et al., Science 1996, 272(5262), 735-8
 {NPL 3} Virginio, C., MacKenzie, A. et al., J. Physiol., 1999, 519, 335-46
 {NPL 4} Duan and Neary, Glia 2006, 54, 738-746
 {NPL 5} Skaper et al, FASEB J 2009, 24, 337-345
 {NPL 6} Surprenant and North, Annu. Rev. Physiol. 2009, 71, 333-359
 {NPL 7} Muller, et al. Am. J. Respir. Cell Mol. Biol. 2011,44, 456-464
 {NPL 8} Ferrari et al., J. Immunol. 2006, 176, 3877-3883
 {NPL 9} Surprenant and North, Annu. Rev. Physiol. 2009, 71, 333-359
 {NPL 10} Chessell et al., Pain 2005, 114, 386-396
 {NPL 11} Basso et al., Behav. Brain Res. 2009, 198, 83-90
 {NPL 12} Dantzer, Immunol. Allergy Clin. North Am. 2009, 29, 247-264
 {NPL 13} Capuron and Miller, Pharmacol. Ther. 2011, 130, 226-238
 {NPL 14} Parvathenani et al., J. Biol. Chem. 2003, 278, 13309-13317
 {NPL 15} Chessell, I. P., Hatcher, J. P. et al., Pain, 2005, 114(3), 386-96


{NPL 16} Guile, S.D., et al., J. Med. Chem, 2009, 52, 3123-3141

{NPL 17} Gunosewoyo, H. and Kassiou, M., Exp Opin, 2010, 20, 625-646

Summary

{0011a} In one aspect of the present disclosure, there is provided a compound represented by the following formula (I):

{Chem. 1}



or a pharmaceutically acceptable salt thereof,

wherein:

X is N or N-oxide;

n is 0 or 1;

R¹ is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) hydroxyl, (4) -NH₂, (5) -NH-C₁₋₆ alkyl and (6) -S(O)_m-C₁₋₆ alkyl;

wherein m is independently 0, 1 or 2;

R² is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl; wherein the C₁₋₆ alkyl or the -O-C₁₋₆ alkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl, -CN, -NR^{9a}R^{10a}, -(C=O)-R^{9a}, -(C=O)-NR^{9a}R^{10a} and -S(O)_m-R^{9a}; wherein m is independently 0, 1 or 2;

R¹ may form =CH₂ or =O with R²; or

R¹ may form a 3 to 7 membered ring with R² which may contain one or more independently selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and carbonyl group; where the 3 to 7 membered ring is unsubstituted or substituted one or more with C₁₋₆ alkyl;

R³ is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R³ is the same or different;

R⁴ is selected from the group consisting of:

(1) hydrogen, (2) halogen and (3) hydroxyl;

R⁵ is hydrogen or C₁₋₆ alkyl;

R⁶ is selected from the group consisting of:

(1) hydrogen, (2) C₁₋₆ alkyl, (3) hydroxyC₁₋₆ alkyl, (4) C₁₋₆ alkoxy C₁₋₆ alkyl and (5) heterocyclyl C₁₋₆ alkyl;

R⁵ may form a saturated 3 to 7 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom, a sulfur atom or a double bond; or a saturated or unsaturated bicyclic 9 to 10 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom or a sulfur atom; wherein the saturated 3 to 7 membered ring or the saturated or unsaturated bicyclic 9 to 10 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) -O-aryl and (4) -O-C₁₋₆ alkylaryl;

R^{7a} and R^{7b} are independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) hydroxyl, (4) C₁₋₆ alkyl and (5) -NR^{9b}R^{10b};

R^{7a} may form a 3 to 7 membered ring with R⁵ which may contain a nitrogen atom or an oxygen atom; or

R^{7a} may form a 3 to 7 membered ring with R^{7b} which may contain a nitrogen atom or an oxygen atom;

q is 0 or 1;

R⁸ is selected from the group consisting of:

(1) hydrogen, (2) C₁₋₆ alkyl, (3) -O-C₁₋₆ alkyl, (4) C₂₋₆ alkenyl, (5) C₃₋₁₀ cycloalkyl, (6) -NR^{9b}R^{10b}: wherein the C₁₋₆ alkyl, the -O-C₁₋₆ alkyl, the C₂₋₆ alkenyl, the C₃₋₁₀ cycloalkyl or the -NR^{9b}R^{10b} is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen and hydroxyl; (7) heterocyclyl, (8) aryl, (9) -O-C₁₋₆ alkylaryl, (10) -O-aryl, (11) heteroaryl and (12) aryl-substituted heteroaryl: wherein the heterocyclyl, the aryl, the -O-C₁₋₆ alkylaryl, the -O-aryl, the heteroaryl or the aryl-substituted heteroaryl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -C₃₋₇ cycloalkyl, -O-C₃₋₇ cycloalkyl, hydroxyl-C₁₋₆ alkoxy, -CN, -NR^{9b}R^{10b}, -(C=O)-R^{9b}, -(C=O)-NR^{9b}R^{10b}, -NR^{9b}-(C=O)-R^{10b}, -NR¹¹-(C=O)-NR^{9b}R^{10b}, -NR^{9b}-(C=O)-OR^{10b},

$-\text{NR}^{9b}-\text{S}(\text{O})_m-\text{R}^{10b}$, $-\text{NR}^{11}-\text{S}(\text{O})_m-\text{NR}^{9b}\text{R}^{10b}$, $-\text{S}(\text{O})_m-\text{R}^{9b}$ and C_{1-6} alkyl which may be substituted one or more with halogen, hydroxyl, $-\text{O}-\text{C}_{1-6}$ alkyl or $\text{NR}^{9b}\text{R}^{10b}$; wherein m is independently 0, 1 or 2;

R^{9a} , R^{9b} , R^{10a} , R^{10b} or R^{11} is independently selected from the group consisting of:


(1) hydrogen, (2) hydroxyl, (3) C_{1-6} alkyl and (4) hydroxy C_{1-6} alkyl;

R^{9a} may form a 4 to 7 membered ring with R^{10a} which may contain one or more independently selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and double bond, wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) C_{1-6} alkyl and (4) $-\text{O}-\text{C}_{1-6}$ alkyl;

R^{9b} may form a 4 to 7 membered ring with R^{10b} which may contain one or more independently selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and double bond, wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) C_{1-6} alkyl.

{0011b} In another aspect of the present disclosure, there is provided a compound represented by the following formula (M):

{Chem. 2}



or a pharmaceutically acceptable salt thereof,

wherein:

n is 0 or 1;

R¹ is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) hydroxyl, (4) -NH₂, (5) -NH-C₁₋₆ alkyl and (6) -S(O)_m-C₁₋₆ alkyl; wherein m is independently 0, 1 or 2;

R² is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl; wherein the C₁₋₆ alkyl or the -O-C₁₋₆ alkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl, -CN, -NR^{9a}R^{10a}, -(C=O)-R^{9a}, -(C=O)-NR^{9a}R^{10a} and -S(O)_m-R^{9a}; wherein m is independently 0, 1 or 2;

R¹ may form =CH₂ or =O with R²; or

R¹ may form a 3 to 7 membered ring with R² which may contain one or more independently selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and carbonyl group; where the 3 to 7 membered ring is unsubstituted or substituted one or more with C₁₋₆ alkyl;

R³ is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R³ is the same or different;

R⁴ is selected from the group consisting of:

(1) hydrogen, (2) halogen and (3) hydroxyl;

R⁵ is hydrogen or C₁₋₆ alkyl;

R⁶ is selected from the group consisting of:

(1) hydrogen, (2) C₁₋₆ alkyl, (3) hydroxyC₁₋₆ alkyl, (4) C₁₋₆ alkoxy C₁₋₆ alkyl and (5) heterocyclyl C₁₋₆ alkyl;

R⁵ may form a saturated 3 to 7 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom, a sulfur atom or a double bond; wherein the saturated 3 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) -O-aryl and (4) -O-C₁₋₆ alkylaryl;

R^{9a}, R^{9b}, R^{10a} or R^{10b} is independently selected from the group consisting of:

(1) hydrogen, (2) hydroxyl, (3) C₁₋₆ alkyl and (4) hydroxyC₁₋₆ alkyl;

R^{9a} may form a 4 to 7 membered ring with R^{10a} which may contain a nitrogen atom or an oxygen atom; wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

R^{9b} may form a 4 to 7 membered ring with R^{10b} which may contain a nitrogen atom or an oxygen atom; wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) C₁₋₆ alkyl;

R¹², R¹³, R¹⁴ and R¹⁵ are independently selected from the group consisting of:

(1) hydrogen, (2) hydroxyl, (3) halogen, (4) C₁₋₆ alkyl, (5) -O-C₁₋₆ alkyl and (6) CN; wherein the C₁₋₆ alkyl or the -O-C₁₋₆ alkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl and NR^{9b}R^{10b}; or

R¹² may form a 5 to 7 membered ring with R⁵ which may contain one or more independently selected from the group consisting of: nitrogen atom and oxygen atom.

{0011c} In a further aspect of the present disclosure, there is provided a compound which is selected from the group following or a pharmaceutically acceptable salt thereof,

N-(2,4-dichloro-6-methylbenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,3-dichlorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(cycloheptylmethyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide

;

N-(2,3-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(methoxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-hydroxy-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 ;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-(methoxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-5,8-dihydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-5,8-dihydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 N-(2,3-dichlorobenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 2-(5-((2,4-dichloro-6-methylbenzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;
 2-(5-((2-chloro-3-(trifluoromethyl)benzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;
 (2-amino-2-oxoethyl)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-7-(hydroxymethyl)-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8S*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8S*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

8-(aminomethyl)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

8-(aminomethyl)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

8-(aminomethyl)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxy-3-methylazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxy-3-methylazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-8-amino-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8S*)-8-amino-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (3R*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide;
 (3S*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide;
 (5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(2,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,7S)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8R)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8R)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,7R)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

(5S,7R)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-fluoro-3-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1-(2,3,4-trichlorophenyl)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(3-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(5-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-bromo-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-6-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((R)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorophenethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3-chloro-5-(trifluoromethyl)pyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(3,4,5-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-cyano-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-4-(trifluoromethoxy)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-5-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-(4-methoxy-2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(3,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-((4-(4-chlorophenyl)thiazol-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-(2-(morpholinomethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-((1S,2R)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(6-chloro-2-fluoro-3-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-(2-(3-(trifluoromethyl)phenoxy)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-N-((1-(4-fluorophenyl)cyclopropyl)methyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(3,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-((1R,2S)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-N-(2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-3-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-((1S,2S)-2-(benzyloxy)cyclopentyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-((S)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(3,3-dimethylbutyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-phenoxyethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4,6-dichloro-2,3-dihydrobenzofuran-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(5,7-dichlorochroman-4-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(1-(adamantan-1-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-(4-chlorophenyl)-2-(4,4-difluoropiperidin-1-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(chroman-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-(4-chlorophenyl)propyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-5-fluoro-8-hydroxy-N-(2-morpholino-2-phenylethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-(4,4-difluoropiperidin-1-yl)-2-(4-methylthiazol-5-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-3,5-difluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-3,5-difluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-((2,4-dichlorobenzyl)carbamoyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline 1-oxide;

(R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-5-fluoro-8-oxo-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-8-methyl-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3,8-dimethyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-(cyanomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-(cyanomethyl)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)thio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-4-methyl-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(5S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-quinoline]-5'-carboxamide;

(5R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-quinoline]-5'-carboxamide;

(2S,5'R)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(2S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfinyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-dichloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorophenethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-methoxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-methoxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-(2-hydroxyethoxy)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-8-(2,3-dihydroxypropoxy)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8rac)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(methylsulfonyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8rac)-N-(2,4-dichlorobenzyl)-5-fluoro-8-((2-hydroxyethyl)sulfonyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;


(5R,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide and

(5R,8S)-N-(2,3-dichlorobenzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{0011d} In another aspect of the present disclosure, there is provided a compound represented by the following formula (L-a) or a pharmaceutically acceptable salt thereof:

{Chem. 3}



wherein:

R³ is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R³ is the same or different;

R¹⁶ is selected from the group consisting of:

(1) CN and (2) -CO₂R¹⁸;

R¹⁷ is selected from the group consisting of:

(1) fluoro and (2) hydroxyl; or,


R¹⁶ may form =O with R¹⁷;

R¹⁸ is selected from the group consisting of:

(1) hydrogen and (2) C₁₋₆ alkyl.

{0011e} In a further aspect of the present disclosure, there is provided a compound represented by the following formula (L-b) or a pharmaceutically acceptable salt thereof:

{Chem. 4}



wherein:

R³ is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R³ is the same or different;

R⁴ is selected from the group consisting of:

(1) hydrogen, (2) halogen and (3) hydroxyl.

{0011f} In another aspect of the present disclosure, there is provided a use of a compound disclosed herein or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of a condition or disorder mediated by P2X7 receptor antagonistic activity,

wherein the condition or disorder is selected from the group consisting of: diseases of the autoimmune and inflammatory system; diseases of the nervous and neuro-immune system; diseases involved with, and without, neuroinflammation of the Central Nervous System (CNS); diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems; skeletal disorders, diseases involving the secretory function of exocrine glands and glaucoma, Glomerulonephritis, Chaga's Disease, chlamydia, neuroblastoma, Tuberculosis, polycystic Kidney Disease, cancer, and acne; and combinations thereof.

{0011g} In a further aspect of the present disclosure, there is provided a compound disclosed herein or a pharmaceutically acceptable salt thereof for use in the treatment of a condition or disorder mediated by P2X7 receptor antagonistic activity.

{0011h} In another aspect of the present disclosure, there is provided a pharmaceutical composition comprising a compound or a pharmaceutically acceptable salt thereof, as disclosed herein, and a pharmaceutically acceptable carrier.

{0011i} In a further aspect of the present disclosure, there is provided a process for preparing a pharmaceutical composition, wherein the process comprises mixing a compound disclosed herein or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier or excipient.

Technical Problem

{0012} There is a need in the art for P2X7 antagonists that can be used to treat a disease, syndrome, or condition in a mammal including human, wherein the disease, syndrome, or condition is affected by the modulation of P2X7 receptors, such as diseases of the autoimmune and inflammatory system; diseases of the nervous and neuro-immune system; diseases involved with, and without, neuroinflammation of the CNS; diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems; skeletal disorders, diseases involving the secretory function of exocrine glands and glaucoma, Glomerulonephritis, Chaga's Disease, chlamydia, neuroblastoma, Tuberculosis, Polycystic Kidney Disease, cancer, and acne.

{0013} It is an objective of the invention to provide new P2X7 receptor antagonists that are good drug candidates. P2X7 antagonists should be well absorbed from the GI tract and should be metabolically stable and possess favorable pharmacokinetic properties. They should be non-toxic. Furthermore, the ideal drug candidate would exist in a physical form that is stable, non-hygroscopic and easily formulated. In particular, it has been desired that compounds would have to bind potently to the P2X7 receptor and show functional activity as antagonists. Also it has been desired that compounds would have favorable pharmacokinetic properties.

Solution to Problem

{0014} With respect to other compounds disclosed in the art, the compounds of the present invention may show less toxicity, good absorption and distribution, good solubility, less plasma protein binding, less drug-drug interaction, good metabolic stability. The present invention provides novel compounds which have excellent P2X7 antagonistic activities as well as excellent pharmacokinetic properties.

{0015} In addition, the tetrahydroquinoline derivatives of the present invention show an excellent selectivity for the P2X7 channel as compared with other P2X families, especially P2X1 channel. Involvement of P2X1 against auto regulation in kidney has been reported in P2X1 antagonist (NF 279) (Purinergic Signalling (2012) 8 : 375-417). Thus, selective P2X7 compounds of the present invention lead to improvements in the side-effect profile.

{0016} The present invention provides the following items.

{0017} [1] A compound represented by the following formula (I):

(II)

R⁵ is hydrogen or C₁₋₆ alkyl;

R⁶ is selected from the group consisting of:

(1) hydrogen, (2) C₁₋₆ alkyl, (3) hydroxyC₁₋₆ alkyl, (4) C₁₋₆ alkoxy C₁₋₆ alkyl and (5) heterocyclyl C₁₋₆ alkyl;

preferably R⁶ is selected from the group consisting of (1) hydrogen and (2) C₁₋₆ alkyl;

R⁵ may form a saturated 3 to 7 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom, a sulfur atom or a double bond; or a saturated or unsaturated bicyclic 9 to 10 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom or a sulfur atom; wherein the saturated 3 to 7 membered ring or the saturated or unsaturated bicyclic 9 to 10 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) -O-aryl and (4) -O-C₁₋₆ alkylaryl; preferably R⁵ may form a saturated or unsaturated bicyclic 9 to 10 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom or a sulfur atom; wherein the saturated or unsaturated bicyclic 9 to 10 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) -O-aryl;

R^{7a} and R^{7b} are independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) hydroxyl, (4) C₁₋₆ alkyl and (5) -NR^{9b}R^{10b};

preferably R^{7a} and R^{7b} are independently selected from the group consisting of (1) hydrogen, (4) C₁₋₆ alkyl and (5) -NR^{9b}R^{10b};

R^{7a} may form a 3 to 7 membered ring with R⁵ which may contain a nitrogen atom or an oxygen atom; or

R^{7a} may form a 3 to 7 membered ring with R^{7b} which may contain a nitrogen atom or an oxygen atom;

q is 0 or 1; preferably q is 0;

R⁸ is selected from the group consisting of:

(1) hydrogen, (2) C₁₋₆ alkyl, (3) -O-C₁₋₆ alkyl, (4) C₂₋₆ alkenyl, (5) C₃₋₁₀ cycloalkyl, (6) -NR^{9b}R^{10b}: wherein the C₁₋₆ alkyl, the -O-C₁₋₆ alkyl, the C₂₋₆ alkenyl, the C₃₋₁₀ cycloalkyl or the -NR^{9b}R^{10b} is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen and hydroxyl; (7) heterocyclyl, (8) aryl, (9) -O-C₁₋₆ alkylaryl, (10) -O-aryl, (11) heteroaryl and (12) aryl-substituted heteroaryl:

wherein the heterocyclyl, the aryl, the -O-C₁₋₆ alkylaryl, the -O-aryl, the heteroaryl or the aryl-substituted heteroaryl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -C₃₋₇ cycloalkyl, -O-C₃₋₇ cycloalkyl, hydroxyl-C₁₋₆ alkoxy, -CN, -NR^{9b}R^{10b}, -(C=O)-R^{9b}, -(C=O)-NR^{9b}R^{10b}, -NR^{9b}-(C=O)-R^{10b}, -NR¹¹-(C=O)-NR^{9b}R^{10b}, -NR^{9b}-(C=O)-OR^{10b}, -NR^{9b}-S(O)_m-R^{10b}, -NR¹¹-S(O)_m-NR^{9b}R^{10b}, -S(O)_m-R^{9b} and C₁₋₆ alkyl which may be substituted one or more with halogen, hydroxyl, -O-C₁₋₆ alkyl or

$\text{NR}^{9b}\text{R}^{10b}$;

wherein m is independently 0, 1 or 2;

preferably R^8 is selected from the group consisting of: (1) hydrogen, (2) C_{1-6} alkyl, (5) C_{3-10} cycloalkyl, wherein the C_{1-6} alkyl or the C_{3-10} cycloalkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen and hydroxyl; (7) heterocyclyl, (8) aryl, (9) $-\text{O}-\text{C}_{1-6}$ alkylaryl, (10) $-\text{O}-\text{aryl}$, (11) heteroaryl and (12) aryl-substituted heteroaryl, wherein the heterocyclyl, the aryl, the $-\text{O}-\text{C}_{1-6}$ alkylaryl, the $-\text{O}-\text{aryl}$, the heteroaryl or the aryl-substituted heteroaryl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, $-\text{O}-\text{C}_{1-6}$ alkyl, $-\text{O}-\text{C}_{1-6}$ haloalkyl, $-\text{CN}$ and C_{1-6} alkyl which may be substituted one or more with halogen, hydroxyl, $-\text{O}-\text{C}_{1-6}$ alkyl or $\text{NR}^{9b}\text{R}^{10b}$;

more preferably, R^8 is selected from the group consisting of:

2,4-dichloro-3-fluorophenyl, 2-chloro-3,4-difluorophenyl, 2,3,4-trifluorophenyl, 2-chloro-4-fluorophenyl, 2,4-dichlorophenyl, 4-chloro-2,6-difluorophenyl, 2-chloro-4,6-difluorophenyl, 2,4-dichloro-6-fluorophenyl, 2,3-dichlorophenyl, 2-chloro-3-(trifluoromethyl)phenyl, 2,4-dichloro-6-(hydroxymethyl)phenyl and 2-chloro-4-fluoro-6-(hydroxymethyl)phenyl;

R^{9a} , R^{9b} , R^{10a} , R^{10b} or R^{11} is independently selected from the group consisting of:

(1) hydrogen, (2) hydroxyl, (3) C_{1-6} alkyl and (4) hydroxy C_{1-6} alkyl;

R^{9a} may form a 4 to 7 membered ring with R^{10a} which may contain one or more independently selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and double bond, wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) C_{1-6} alkyl and (4) $-\text{O}-\text{C}_{1-6}$ alkyl;

R^{9b} may form a 4 to 7 membered ring with R^{10b} which may contain one or more independently selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and double bond, wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) C_{1-6} alkyl.

[0018] [2] The compound according to [1]:

or a prodrug thereof or a pharmaceutically acceptable salt thereof,
wherein:

X is N;

R^5 is hydrogen or C_{1-6} alkyl;

R^6 is selected from the group consisting of:

(1) hydrogen and (2) C_{1-6} alkyl;

R^5 may form a saturated or unsaturated bicyclic 9 to 10 membered ring with R^6

which may contain a nitrogen atom, an oxygen atom or a sulfur atom; wherein the saturated or unsaturated bicyclic 9 to 10 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) -O-aryl;

R^{7a} and R^{7b} are independently selected from the group consisting of: (1) hydrogen, (4) C_{1-6} alkyl and (5) $-NR^{9b}R^{10b}$;

R^{7a} may form a 3 to 7 membered ring with R^5 which may contain a nitrogen atom or an oxygen atom; or


R^{7a} may form a 3 to 7 membered ring with R^{7b} which may contain a nitrogen atom or an oxygen atom;

R^8 is selected from the group consisting of:

(1) hydrogen, (2) C_{1-6} alkyl, (5) C_{3-10} cycloalkyl, wherein the C_{1-6} alkyl or the C_{3-10} cycloalkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen and hydroxyl; (7) heterocyclyl, (8) aryl, (9) -O- C_{1-6} alkylaryl, (10) -O-aryl, (11) heteroaryl and (12) aryl-substituted heteroaryl, wherein the heterocyclyl, the aryl, the -O- C_{1-6} alkylaryl, the -O-aryl, the heteroaryl or the aryl-substituted heteroaryl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O- C_{1-6} alkyl, -O- C_{1-6} haloalkyl, -CN and C_{1-6} alkyl which may be substituted one or more with halogen, hydroxyl, -O- C_{1-6} alkyl or $NR^{9b}R^{10b}$.

[0019] [3] A compound represented by the following formula (M):

[Chem.2]



or a prodrug thereof or a pharmaceutically acceptable salt thereof,

wherein:

n is 0 or 1; preferably n is 1;

R^1 is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) hydroxyl, (4) $-NH_2$, (5) $-NH-C_{1-6}$ alkyl and (6) $-S(O)_m-C_{1-6}$ alkyl;

wherein m is independently 0, 1 or 2;

R² is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl; wherein the C₁₋₆ alkyl or the -O-C₁₋₆ alkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl, -CN, -NR^{9a}R^{10a}, -(C=O)-R^{9a}, -(C=O)-NR^{9a}R^{10a} and -S(O)_m-R^{9a};

wherein m is independently 0, 1 or 2;

R¹ may form =CH₂ or =O with R²; or

R¹ may form a 3 to 7 membered ring with R² which may contain one or more independently selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and carbonyl group; where the 3 to 7 membered ring is unsubstituted or substituted one or more with C₁₋₆ alkyl;

R³ is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

preferably R³ is hydrogen at 4-position against X;

p is 0, 1, 2 or 3; preferably p is 0 or 1;

when p is 2 or 3, each R³ is the same or different;

R⁴ is selected from the group consisting of:

(1) hydrogen, (2) halogen and (3) hydroxyl;

R⁵ is hydrogen or C₁₋₆ alkyl;

R⁶ is selected from the group consisting of:

(1) hydrogen, (2) C₁₋₆ alkyl, (3) hydroxyC₁₋₆ alkyl, (4) C₁₋₆ alkoxy C₁₋₆ alkyl and (5) heterocyclyl C₁₋₆ alkyl;

preferably R⁶ is selected from the group consisting of (1) hydrogen and (2) C₁₋₆ alkyl;

R⁵ may form a saturated 3 to 7 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom, a sulfur atom or a double bond; wherein the saturated 3 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) -O-aryl and (4) -O-C₁₋₆ alkylaryl;

preferably R⁵ may form a saturated 3 to 7 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom, a sulfur atom or a double bond; wherein the saturated 3 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) -O-aryl;

R^{9a}, R^{9b}, R^{10a} or R^{10b} is independently selected from the group consisting of:

(1) hydrogen, (2) hydroxyl, (3) C₁₋₆ alkyl and (4) hydroxyC₁₋₆ alkyl;

R^{9a} may form a 4 to 7 membered ring with R^{10a} which may contain a nitrogen atom or an oxygen atom; wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2)

halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

R^{9b} may form a 4 to 7 membered ring with R^{10b} which may contain a nitrogen atom or an oxygen atom; wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) C₁₋₆ alkyl;

R¹², R¹³, R¹⁴ and R¹⁵ are independently selected from the group consisting of:

(1) hydrogen, (2) hydroxyl, (3) halogen, (4) C₁₋₆ alkyl, (5) -O-C₁₋₆ alkyl and (6) CN; wherein the C₁₋₆ alkyl or the -O-C₁₋₆ alkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl and NR^{9b}R^{10b}; or

R¹² may form a 5 to 7 membered ring with R⁵ which may contain one or more independently selected from the group consisting of: nitrogen atom and oxygen atom.

[0020] [4] The compound according to [3]:

or a prodrug thereof or a pharmaceutically acceptable salt thereof,

wherein:

n is 1;

R¹ is hydrogen or hydroxyl;

R² is methyl which is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl, -CN and -NR^{9a}R^{10a};

p is 0;

R⁴ is hydrogen or fluoro;

R⁵ and R⁶ are independently selected from the group consisting of: (1) hydrogen and (2) C₁₋₆ alkyl;

R¹², R¹³ and R¹⁴ are independently selected from the group consisting of: (1) hydrogen, (3) halogen, and (4) C₁₋₃ alkyl which may be substituted one or more with hydroxyl;

R¹⁵ is hydrogen.

[0021] [5] A compound which is selected from the group following or a prodrug thereof or a pharmaceutically acceptable salt thereof,

N-(2,4-dichloro-6-methylbenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,3-dichlorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(cycloheptylmethyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

ide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,3-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(methoxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-hydroxy-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-(methoxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5,8-dihydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5,8-dihydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2,3-dichlorobenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

2-(5-((2,4-dichloro-6-methylbenzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;

2-(5-((2-chloro-3-(trifluoromethyl)benzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;

(2-amino-2-oxoethyl)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-7-(hydroxymethyl)-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8S*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8S*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

8-(aminomethyl)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

8-(aminomethyl)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,4-dichloro-6-methylbenzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

8-(aminomethyl)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8R*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxy-3-methylazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxy-3-methylazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8R*)-8-amino-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S*,8S*)-8-amino-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(3R*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide;

(3S*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
;
(5R,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
;
(5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(2,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
;
(5S,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,7S)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,7R)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

(5S,7R)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-fluoro-3-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1-(2,3,4-trichlorophenyl)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(3-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(5-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-bromo-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-6-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((R)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorophenethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3-chloro-5-(trifluoromethyl)pyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(3,4,5-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-cyano-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-4-(trifluoromethoxy)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-5-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(4-methoxy-2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((4-(4-chlorophenyl)thiazol-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-(morpholinomethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1S,2R)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(6-chloro-2-fluoro-3-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-(3-(trifluoromethyl)phenoxy)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-((1-(4-fluorophenyl)cyclopropyl)methyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1R,2S)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((1S,2S)-2-(benzyloxy)cyclopentyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,3-dimethylbutyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-phenoxyethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4,6-dichloro-2,3-dihydrobenzofuran-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(5,7-dichlorochroman-4-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(1-(adamantan-1-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-(4-chlorophenyl)-2-(4,4-difluoropiperidin-1-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(chroman-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-(4-chlorophenyl)propyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-morpholino-2-phenylethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-(4,4-difluoropiperidin-1-yl)-2-(4-methylthiazol-5-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-3,5-difluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-3,5-difluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-((2,4-dichlorobenzyl)carbamoyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline 1-oxide;

(R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

;

(S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

;

(S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-5-fluoro-8-oxo-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-8-methyl-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3,8-dimethyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-(cyanomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-(cyanomethyl)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)thio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-4-methyl-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(5S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-quinoline]-5'-carboxamide;

(5R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-quinoline]-5'-carboxamide;

(2S,5'R)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(2S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfinyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-dichloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorophenethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-methoxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-methoxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-(2-hydroxyethoxy)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-8-(2,3-dihydroxypropoxy)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8rac)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(methylsulfonyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8rac)-N-(2,4-dichlorobenzyl)-5-fluoro-8-((2-hydroxyethyl)sulfonyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichlorobenzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide and
 (5R,8S)-N-(2,3-dichlorobenzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide.

- [0022] [6] The compound according to [5], which is selected from the group following or a prodrug thereof or a pharmaceutically acceptable salt thereof,
- N-(2,3-dichlorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 - N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 - N-(2-chloro-3-(trifluoromethyl)benzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 - N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 - N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 - N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 - N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 - N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 - N-(2-chloro-3-(trifluoromethyl)benzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 - N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 - 2-(5-((2,4-dichloro-6-methylbenzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;
 - 2-(5-((2-chloro-3-(trifluoromethyl)benzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;
 - N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7

,8-tetrahydroquinoline-5-carboxamide;
 N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-7-(hydroxymethyl)-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 (5S*,8S*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxy-3-methylazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxy-3-methylazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (3R*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide;
 (3S*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide;
 (5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 ;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carb

oxamide;

(5S,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carbox

amide;

(5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-

5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

droquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5R,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-3,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-4-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-N-(3-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydro

quinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-bromo-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(3-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-((R)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(3-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(3-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-((3-chloro-5-(trifluoromethyl)pyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-((R)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(3,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-(2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-N-(3,4,5-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-cyano-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(3-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-

-carboxamide;

(5S,8S)-N-(3-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-4-(trifluoromethoxy)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1S,2R)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-(3-(trifluoromethyl)phenoxy)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1R,2S)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,3-dimethylbutyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-phenoxyethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4,6-dichloro-2,3-dihydrobenzofuran-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(chroman-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

mide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-morpholino-2-phenylethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-(4,4-difluoropiperidin-1-yl)-2-(4-methylthiazol-5-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-((2,4-dichlorobenzyl)carbamoyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline 1-oxide;

(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amin


o)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amin
 o)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methy
 l)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-8-(cyanomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydr
 oquinoline-5-carboxamide;
 (5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetr
 ahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetr
 ahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-8-(cyanomethyl)-5-fluoro-8-
 hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetrahydr
 oquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetr
 ahydroquinoline-5-carboxamide;
 (2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-4-methyl-6',7'-dihydro-5'H-spiro[morpholin
 e-2,8'-quinoline]-5'-carboxamide;
 (5S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,
 8'-quinoline]-5'-carboxamide;
 (5R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,
 8'-quinoline]-5'-carboxamide;
 (2S,5'R)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,
 8'-quinoline]-5'-carboxamide;
 (2S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,
 8'-quinoline]-5'-carboxamide;
 (2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,
 8'-quinoline]-5'-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfinyl)methyl)-5
 ,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)met
 hyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl
)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetra

ydroquinoline-5-carboxamide;
(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8R)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8R)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2,4-dichloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,

8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-5-fluoro-8-methoxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide and
 (5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

[0023] [7] A compound represented by the following formula (L-a) or a pharmaceutically acceptable salt thereof:

[Chem.3]



wherein:

R^3 is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C_{1-6} alkyl and (4) $-O-C_{1-6}$ alkyl;

preferably R^3 is (1) hydrogen;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R^3 is the same or different;

R^{16} is selected from the group consisting of:

(1) CN and (2) $-\text{CO}_2R^{18}$;

R^{17} is selected from the group consisting of:

(1) fluoro and (2) hydroxyl; or,

R^{16} may form $=\text{O}$ with R^{17} ;

R^{18} is selected from the group consisting of:

(1) hydrogen and (2) C_{1-6} alkyl.

[7-1]The compound according to [7] or a salt thereof, wherein R^{16} forms $=\text{O}$ with R^{17} .

[7-2]The compound according to [7] or a salt thereof, wherein R^{17} is fluoro.


[7-3]The compound according to [7-2] or a salt thereof, wherein R^{16} is $-\text{CN}$.

[7-4]The compound according to [7-2] or a salt thereof, wherein R^{16} is $-\text{CO}_2R^{18}$.

[7-5]The compound according to [7] or a salt thereof, wherein R^{16} is $-\text{CO}_2R^{18}$; R^{17} is hydroxyl.

[0024] [8] A compound represented by the following formula (L-b) or a pharmaceutically acceptable salt thereof:

[Chem.4]



wherein:

R^3 is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C_{1-6} alkyl and (4) $-\text{O}-\text{C}_{1-6}$ alkyl;

preferably R^3 is (1) hydrogen;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R^3 is the same or different;

R^4 is selected from the group consisting of:

(1) hydrogen, (2) halogen and (3) hydroxyl.

[0025] [9] A use of a compound described in any one of [1] to [6] or a pharmaceutically acceptable salt thereof or a prodrug thereof for the manufacture of a medicament for the treatment of a condition or disorder mediated by P2X7 receptor antagonistic activity.

[0026] [10] The use as described in [9], wherein the condition or disorder is selected from the group consisting of: diseases of the autoimmune and inflammatory system; diseases of the nervous and neuro-immune system; diseases involved with, and without, neuroinflammation of the Central Nervous System (CNS); diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems; skeletal disorders,

diseases involving the secretory function of exocrine glands and glaucoma, Glomerulonephritis, Chaga's Disease, chlamydia, neuroblastoma, Tuberculosis, Polycystic Kidney Disease, cancer, and acne; and combinations thereof.

- [0027] [11] A method for the treatment of a condition or disorder mediated by P2X7 receptor antagonistic activity in a mammalian subject, including a human, which comprises administering to a mammal in need of such treatment a therapeutically effective amount of a compound described in any one of [1] to [6] or a pharmaceutically acceptable salt thereof or a prodrug thereof.
- [0028] [12] The method as described in [11], wherein the condition or disorder is selected from the group consisting of: diseases of the autoimmune and inflammatory system; diseases of the nervous and neuro-immune system; diseases involved with, and without, neuroinflammation of the Central Nervous System (CNS); diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems; skeletal disorders, diseases involving the secretory function of exocrine glands and glaucoma, Glomerulonephritis, Chaga's Disease, chlamydia, neuroblastoma, Tuberculosis, Polycystic Kidney Disease, cancer, and acne; and combinations thereof.
- [0029] [13] A pharmaceutical composition comprising a compound or a pharmaceutically acceptable salt thereof or a prodrug thereof, as described in any one of [1] to [6], and a pharmaceutically acceptable carrier.
- [0030] [14] The pharmaceutical composition as described in [13], further comprising another pharmacologically active agent.
- [0031] [15] A compound described in any one of [1] to [6] or a pharmaceutically acceptable salt thereof or a prodrug thereof for use in the treatment of a condition or disorder mediated by P2X7 receptor antagonistic activity.
- [0032] [16] A process for preparing a pharmaceutical composition, wherein the process comprises mixing a compound described in any one of [1] to [6] or a pharmaceutically acceptable salt thereof or a prodrug thereof and a pharmaceutically acceptable carrier or excipient.
- [0033] Examples of conditions or disorders mediated by P2X7 receptor activity include, but are not limited to, P2X7 related diseases.

Advantageous Effects of Invention

- [0034] The tetrahydroquinoline derivatives of the present invention act as modulators of the P2X7 receptor and have a number of therapeutic applications, particularly in the treatment of diseases of the autoimmune and inflammatory system, diseases of the nervous and neuro-immune system, diseases involved with neuroinflammation of the CNS or diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems. More particularly, the tetrahydroquinoline derivatives of the invention are

selective P2X7 receptor antagonists. In the discussion that follows, the invention is exemplified by reference to the inhibition of P2X7 channel as the purinergic receptor.

[0035] The compounds of the present invention show the P2X7 receptor antagonistic activity. The compounds of the present invention may show less toxicity, good absorption, distribution, good solubility, less protein binding affinity other than P2X7 receptor, less drug-drug interaction, and good metabolic stability.

[0036] Especially, the present invention show excellent P2X7 antagonistic activities as well as excellent pharmacokinetic properties. In addition, they show selectivity for the P2X7 channel as compared with other P2X families, especially P2X1 channel.

Description of Embodiments

[0037] As used herein, the term "alkyl" as a group or part of a group e.g. alkoxy or hydroxyalkyl refers to a straight or branched alkyl group in all isomeric forms. The term "C₁-C₆ alkyl" refers to an alkyl group, as defined above, containing at least 1 and at most 6 carbon atoms. Examples of such alkyl groups include methyl, ethyl, propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, pentan-2-yl, pentan-3-yl, 2-methylbutyl, 3-methylbutan-2-yl, isopentyl, tert-pentyl, neopentyl, n-hexyl, hexan-2-yl, hexan-3-yl, 2-methylpentyl, 4-methylpentyl, 2-methylpentan-3-yl, 4-methylpentan-2-yl, 2-ethylbutyl, 3-methylpentyl, 3-methylpentan-2-yl, 3-methylpentan-3-yl, 2,3-dimethylbutyl, 2,3-dimethylbutan-2-yl, 3,3-dimethylbutyl, 3,3-dimethylbutan-2-yl, and 2,2-dimethylbutyl.

Examples of such alkoxy groups include methoxy, ethoxy, propoxy, iso-propoxy, butoxy, iso-butoxy, sec-butoxy, tert-butoxy, n-pentoxy, pentan-2-oxy, pentan-3-oxy, 2-methylbutoxy, 3-methylbutan-2-oxy, isopentoxy, tert-pentoxy, neopentoxy, n-hexoxy, hexan-2-oxy, hexan-3-oxy, 2-methylpentoxy, 4-methylpentoxy, 2-methylpentan-3-oxy, 4-methylpentan-2-oxy, 2-ethylbutoxy, 3-methylpentoxy, 3-methylpentan-2-oxy, 3-methylpentan-3-oxy, 2,3-dimethylbutoxy, 2,3-dimethylbutan-2-oxy, 3,3-dimethylbutoxy, 3,3-dimethylbutan-2-oxy, and 2,2-dimethylbutoxy.

[0038] The term "cycloalkyl", as used herein, means a mono-, bicyclic- or tricyclic ring, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, norbornyl, adamantyl groups and the like.

[0039] The term "alkenyl", as used herein, means a hydrocarbon radical having at least one double bond, which may be in a E- or a Z- arrangement, including, but not limited to, ethenyl, propenyl, 1-butenyl, 2-butenyl and the like.

[0040] The term "halogen" refers to fluoro (-F), chloro (-Cl), bromo (-Br) and iodo (-I).

[0041] The term "haloalkyl", as used herein, means an alkyl radical which is substituted by halogen atom(s) as defined above including, but not limited to, fluoromethyl, difluo-

romethyl, trifluoromethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2,2,2-trichloroethyl, 3-fluoropropyl, 4-fluorobutyl, chloromethyl, trichloromethyl, iodomethyl, bromomethyl groups and the like.

[0042] The term "heterocyclyl", as used herein, means a saturated 3- to 16- membered ring which comprises one or more heteroatoms selected from nitrogen, oxygen and sulfur. For purposes of this invention, the heterocyclyl may be a monocyclic, bicyclic or tricyclic ring system, which may include fused, bridged or spiro ring systems. Examples of such heterocyclyl groups include azetidiny, 1,4-dioxany, pyrrolidiny, piperidiny, piperaziny, morpholiny, tetrahydrofurany, thiomorpholiny, tetrahydrothieny, 2-oxo-pyrrolidiny, 2-oxo-piperidiny, 2-oxo-imidazolidiny, 2-oxo-oxazolidiny, quinuclidiny, azabicyclo[3.2.1]octyl, 2-oxa-6-azaspiro[3.4]octyl and N-oxides thereof and S-oxides thereof.

[0043] The term "aryl", as used herein, means unsaturated or partially saturated mono- or bicyclic 6-15 membered ring which consists of carbon atoms.

Examples of such aryl include, but are not limited to, phenyl, naphthyl, indanyl, indenyl, 1,2,3,4-tetrahydronaphthyl, 1,2-dihydronaphthyl, 2,3-dihydro-1H-indenyl, cyclohexenyl, cyclopentenyl, (1S,4S)-bicyclo[2.2.2]oct-2-enyl, and (1R,4S)-bicyclo[2.2.1]hept-2-enyl. Examples of -O-C₁₋₆ alkylaryl include, benzyloxy.

[0044] The term "heteroaryl" as used herein, means unsaturated and partially saturated mono- or bicyclic 5-15 membered ring, preferably 6-10 membered ring, which may contain 1-4 heteroatoms selected from O, N and S:

Examples of such heteroaryl include, but are not limited to, thiophenyl, thiazolyl, isoxazolyl, pyrazolyl, tetrazolyl, furanyl, pyrrolyl, imidazolyl, oxazolyl, isothiazolyl, triazolyl, thiadiazolyl, pyridyl, pyrimidyl, pyridaziny, pyraziny, triaziny, benzofuranyl, benzothiophenyl, benzotriazolyl, indolyl, indazolyl, benzoimidazolyl, pyrrolopyridyl, pyrrolopyrimidiny, pyrazolopyridyl, pyrazolopyrimidiny, imidazopyridiny, furopyridyl, benzoisoxazolyl, imidazopyraziny, imidazopyridaziny, imidazopyrimidiny, quinolyl, isoquinolyl, quinazolinyl, phthalaziny, quinoxaliny, naphthyridiny, pyridopyrimidiny, and N-oxides thereof and S-oxides thereof.

[0045] R⁵ may form a saturated or unsaturated bicyclic 9 to 10 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom or a sulfur atom; Examples of such bicyclic 9 to 10 membered ring include 1,2,3,4-tetrahydronaphthalenyl, 2,3-dihydro-1H-indenyl, chromanyl, 2,3-dihydrobenzofuranyl, 1,2,3,4-tetrahydroquinoliny, indoliny, thiochromanyl and 2,3-dihydrobenzo[b]thiophenyl.

[0046] The substituents on the ring of the compound of the present invention may exist on the any atoms if it is chemically allowed.

[0047] The term "protecting group", as used herein, means a hydroxy or amino protecting

group which is selected from typical hydroxy or amino protecting groups described in Protective Groups in Organic Synthesis Forth Edition edited by T. W. Greene et al. (John Wiley & Sons, 2006);

[0048] The terms "treating" and "treatment", as used herein, refer to curative, palliative and prophylactic treatment, including reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition.

[0049] As used herein, the article "a" or "an" refers to both the singular and plural form of the object to which it refers unless indicated otherwise.

[0050] In some cases, the symbol letter is written the corresponding English word in the present specification. For example, the symbols α , β , and δ are written alpha, beta, and delta, respectively.

[0051] Included within the scope of the "compounds of the invention" are all salts, solvates, hydrates, complexes, polymorphs, prodrugs, radiolabeled derivatives, stereoisomers and optical isomers of the compounds of the present invention.

[0052] The compounds of the present invention can form acid addition salts thereof. It would be appreciated that for use in medicine, the salts of the compounds of the present invention should be pharmaceutically acceptable. Suitable pharmaceutically acceptable salts would be apparent to those skilled in the art and include those described in J. Pharm. Sci, 66, 1-19, 1977, such as acid addition salts formed with inorganic acids e.g. hydrochloric, hydrobromic, sulfuric, nitric or phosphoric acid; and organic acids e.g. succinic, maleic, formic, acetic, trifluoroacetic, propionic, fumaric, citric, tartaric, benzoic, p-toluenesulfonic, methanesulfonic or naphthalenesulfonic acid. Certain of the compounds of the present invention may form acid addition salts with one or more equivalents of the acid. The present invention includes within its scope all possible stoichiometric and non-stoichiometric forms. In addition, certain compounds containing an acidic function such as a carboxy can be isolated in the form of their inorganic salt in which the counter ion can be selected from sodium, potassium, lithium, calcium, magnesium and the like, as well as from organic bases.

[0053] Also within the scope of the invention are so-called "prodrugs" of the compounds of the present invention. Thus, certain derivatives of compounds of the present invention which may have little or no pharmacological activity themselves, when administered into or onto the body, can be converted into compounds of the present invention having the desired activity, for example, by hydrolytic cleavage. Such derivatives are referred to as "prodrugs". Further information on the use of prodrugs may be found in Pro-drugs as Novel Delivery Systems, Vol. 14, ACS Symposium Series (T Higuchi and V Stella) and Bioreversible Carriers in Drug Design, Pergamon Press, 1987 (ed. E

B Roche, American Pharmaceutical Association).

[0054] Prodrugs in accordance with the invention, for example, can be produced by replacing appropriate functionalities present in the compounds of the present invention with certain moieties known to those skilled in the art as 'pro-moieties' as described, for example, in Design of Prodrugs by H Bundgaard (Elsevier, 1985). Some examples of prodrugs in accordance with the invention include:

[0055] (i) where the compound of the present invention contains an alcohol functionality (-OH), compounds wherein the hydroxy group is replaced with a moiety convertible in vivo into the hydroxy group. Said moiety convertible in vivo into the hydroxy group means a moiety transformable in vivo into a hydroxyl group by e.g. hydrolysis and/or by an enzyme, e.g. an esterase. Examples of said moiety include, but are not limited to, ester and ether groups which may be hydrolyzed easily in vivo. Preferred are the moieties replaced the hydrogen of hydroxy group with acyloxyalkyl, 1-(alkoxycarbonyloxy)alkyl, phthalidyl and acyloxyalkyloxycarbonyl such as pivaloyloxymethyloxycarbonyl; and

(ii) where the compound of the present invention contains an amino group, an amide derivative prepared by reacting with a suitable acid halide or a suitable acid anhydride is exemplified as a prodrug. A particularly preferred amide derivative as a prodrug is -NHCO(CH₂)₂OCH₃, -NHCOCH(NH₂)CH₃ or the like.

[0056] Further examples of replacement groups in accordance with the foregoing examples and examples of other prodrug types may be found in the aforementioned references.

[0057] The compounds of the present invention, salts thereof and prodrugs thereof may be prepared in crystalline or non-crystalline form and, if crystalline, may optionally be hydrated or solvated. This invention includes within its scope stoichiometric hydrates or solvates as well as compounds containing variable amounts of water and/or solvent.

[0058] Salts and solvates having non-pharmaceutically acceptable counter-ions or associated solvents are within the scope of the present invention, for example, for use as intermediates in the preparation of other compounds of the present invention and their pharmaceutically acceptable salts.

[0059] Additionally, the compounds of the present invention may be administered as prodrugs. As used herein, a "prodrug" of a compound of the present invention is a functional derivative of the compound which, upon administration to a patient, eventually liberates the compound of the present invention in vivo. Administration of a compound of the present invention as a prodrug may enable the skilled artisan to do one or more of the followings: (a) modify the onset of action of the compound in vivo; (b) modify the duration of action of the compound in vivo; (c) modify the transportation or distribution of the compound in vivo; (d) modify the solubility of the compound in vivo; and (e) overcome a side effect or other difficulty encountered with

the compound. Typical functional derivatives used to prepare prodrugs include modifications of the compound that are chemically or enzymatically cleaved in vivo. Such modifications, which include the preparation of phosphates, amides, esters, thioesters, carbonates, and carbamates, are well known to those skilled in the art.

- [0060] In certain of the compounds of the present invention, there may be some chiral carbon atoms. In such cases, compounds of the present invention exist as stereoisomers. The invention extends to all optical isomers such as stereoisomeric forms of the compounds of the present invention including enantiomers, diastereoisomers and mixtures thereof, such as racemates. The different stereoisomeric forms may be separated or resolved one from the other by conventional methods or any given isomer may be obtained by conventional stereoselective or asymmetric syntheses.
- [0061] Certain of the compounds herein can exist in various tautomeric forms and it is to be understood that the invention encompasses all such tautomeric forms.
- [0062] The invention also includes isotopically-labeled compounds, which are identical to those described herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, iodine, and chlorine, such as ^3H , ^{11}C , ^{14}C , ^{18}F , ^{123}I and ^{125}I . Compounds of the invention that contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of the present invention. Isotopically-labeled compounds of the present invention, for example those into which radioactive isotopes such as ^3H , ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. ^{11}C and ^{18}F isotopes are particularly useful in PET (positron emission tomography), and ^{125}I isotopes are particularly useful in SPECT (single photon emission computerized tomography), all useful in brain imaging. Further, substitution with heavier isotopes such as deuterium, i.e., ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labeled compounds of the invention can be generally prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples/Intermediates below, then substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.
- [0063] The potencies and efficacies of the compounds of this invention for P2X7 can be determined by calcium influx assay performed on the human cloned receptor as described herein. Compounds of the present invention have demonstrated antagonistic

activity at the P2X7 receptor, using the functional assay described herein.

[0064] Compounds of the present invention and pharmaceutically acceptable salts thereof are therefore of use in the treatment of conditions or disorders which are mediated via the P2X7 receptor. In particular the compounds of the present invention and pharmaceutically acceptable salts thereof are of use in the treatment of a wide range of diseases, syndromes, and disorders, in particular for the treatment of diseases of the autoimmune and inflammatory system; diseases of the nervous and neuro-immune system; diseases involved with, and without, neuroinflammation of the CNS; diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems; skeletal disorders, diseases involving the secretory function of exocrine glands and diseases such as glaucoma, Glomerulonephritis, Chaga's Disease, chlamydia, neuroblastoma, Tuberculosis, Polycystic Kidney Disease, cancer, and acne.

[0065] Examples of diseases of the autoimmune and inflammatory system include rheumatoid arthritis, osteoarthritis, interstitial cystitis, psoriasis, septic shock, sepsis, allergic dermatitis, idiopathic pulmonary fibrosis, allergic rhinitis, chronic obstructive pulmonary disease, airway hyper-responsiveness and asthma. Examples of asthma include allergic asthma, mild to severe asthma, and steroid resistant asthma.

[0066] Examples of diseases of the nervous and neuro-immune system include acute and chronic pain. Examples of acute and chronic pain include neuropathic pain, inflammatory pain, migraine, spontaneous pain. Examples of spontaneous pain include opioid induced pain, diabetic neuropathy, postherpetic neuralgia, low back pain, chemotherapy-induced neuropathic pain, fibromyalgia.

[0067] Examples of diseases involved with, and without, neuroinflammation of the CNS include cognition, sleep disorders, multiple sclerosis, epileptic seizures, Parkinson's disease, schizophrenia, Alzheimer's disease, Huntington's disease, Amyotrophic Lateral Sclerosis, autism, spinal cord injury and cerebral ischemia/traumatic brain injury, stress-related disorders, and mood disorders. Examples of mood disorders include major depression, major depressive disorder, treatment resistant depression, bipolar disorder, seasonal affective disorder, post natal depression, manic depression, bipolar depression, anxious depression, and anxiety. Examples of anxiety include social anxiety, post traumatic stress disorder, phobias, social phobia, special phobias, panic disorder, obsessive compulsive disorder, acute stress disorder, separation anxiety disorder, and generalized anxiety disorder.

[0068] Examples of diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems include diabetes, diabetes mellitus, thrombosis, irritable bowel disease, irritable bowel syndrome, Crohn's disease, cardiovascular diseases (such as hypertension, myocardial infarction, ischemic heart disease, ischemia), ureteric obstruction, lower urinary tract syndrome, lower urinary tract dysfunction such as in-

continence, and disease after cardiac transplantation.

- [0069] Examples of a neuropathic pain include pain due to a disease, syndrome, condition, disorder, or pain state including cancer, neurological disorders, spine and peripheral nerve surgery, brain tumor, traumatic brain injury (TBI), spinal cord trauma, chronic pain syndrome, fibromyalgia, chronic fatigue syndrome, neuralgias (trigeminal neuralgia, glossopharyngeal neuralgia, postherpetic neuralgia and causalgia), lupus, sarcoidosis, peripheral neuropathy, bilateral peripheral neuropathy, diabetic neuropathy, central pain, neuropathies associated with spinal cord injury, stroke, amyotrophic lateral sclerosis (ALS), Parkinson's disease, multiple sclerosis, sciatic neuritis, mandibular joint neuralgia, peripheral neuritis, polyneuritis, stump pain, phantom limb pain, bony fractures, oral neuropathic pain, Charcot's pain, complex regional pain syndrome I and II (CRPS I/II), radiculopathy, Guillain-Barre syndrome, meralgia paraesthetica, burning-mouth syndrome, optic neuritis, postfebrile neuritis, migrating neuritis, segmental neuritis, Gombault's neuritis, neuronitis, cervicobrachial neuralgia, cranial neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, migrainous neuralgia, idiopathic neuralgia, intercostal neuralgia, mammary neuralgia, Morton's neuralgia, nasociliary neuralgia, occipital neuralgia, red neuralgia, Sluder's neuralgia, sphenopalatine neuralgia, supraorbital neuralgia, vulvodynia, or vidian neuralgia.
- [0070] Antagonistic activities of the compound of the present invention against P2X7 and other P2X families can be confirmed in the suitable methods known to skilled in the art. For example, antagonistic activities of compounds of the present invention have been confirmed in Ca^{2+} influx assay and electrophysiology assay.
- [0071] Activities of the compound of the present invention for each disease, syndrome, and disorder described above can be confirmed in the suitable model known to skilled in the art. For example, activities of compounds of the present invention for pain are confirmed in rodent models of pain.
- [0072] It is to be understood that "treatment" as used herein includes prophylaxis as well as alleviation of established symptoms as described above.
- [0073] A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is usually adapted for oral, parenteral or rectal administration and, as such, may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable or infusible solutions or suspensions or suppositories. Orally administered compositions are generally preferred. Tablets and capsules for oral administration may be in unit dose form, and may contain conventional excipients, such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); tableting lubricants (e.g. magnesium stearate, talc or

silica); disintegrants (e.g. potato starch or sodium starch glycolate); and acceptable wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated according to methods well known in normal pharmaceutical practice.

[0074] Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents (e.g. sorbitol syrup, cellulose derivatives or hydrogenated edible fats), emulsifying agents (e.g. lecithin or acacia), non-aqueous vehicles (which may include edible oils e.g. almond oil, oily esters, ethyl alcohol or fractionated vegetable oils), preservatives (e.g. methyl or propyl-p-hydroxybenzoates or sorbic acid), and, if desired, conventional flavourings or colorants, buffer salts and sweetening agents as appropriate. Preparations for oral administration may be suitably formulated to give controlled release of the active compound or pharmaceutically acceptable salt thereof.

[0075] For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the present invention or pharmaceutically acceptable salt thereof and a sterile vehicle. Formulations for injection may be presented in unit dosage form e.g. in ampoules or in multi-dose, utilising a compound of the present invention or pharmaceutically acceptable salt thereof and a sterile vehicle, optionally with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use. The compound, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead of being dissolved, and sterilisation cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

[0076] Lotions may be formulated with an aqueous or oily base and in general will also contain one or more emulsifying agents, stabilising agents, dispersing agents, suspending agents, thickening agents, or colouring agents. Drops may be formulated

with an aqueous or non-aqueous base also comprising one or more dispersing agents, stabilising agents, solubilising agents or suspending agents. They may also contain a preservative.

[0077] The compounds of the present invention or pharmaceutically acceptable salts thereof may also be formulated in rectal compositions such as suppositories or retention enemas, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

[0078] The compounds of the present invention or pharmaceutically acceptable salts may also be formulated as depot preparations. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds of the present invention or pharmaceutically acceptable salts may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0079] For intranasal administration, the compounds the present invention or pharmaceutically acceptable salts thereof may be formulated as solutions for administration via a suitable metered or unitary dose device or alternatively as a powder mix with a suitable carrier for administration using a suitable delivery device. Thus, the compounds of the present invention or pharmaceutically acceptable salts thereof may be formulated for oral, buccal, parenteral, topical (including ophthalmic and nasal), depot or rectal administration or in a form suitable for administration by inhalation or insufflation (through either mouth or nose). The compounds of the present invention and pharmaceutically acceptable salts thereof may be formulated for topical administration in the form of ointments, creams, gels, lotions, pessaries, aerosols or drops (e.g. eye, ear or nose drops). Ointments and creams may for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Ointments for administration to the eye may be manufactured in a sterile manner using sterilized components.

[0080] A P2X7 antagonist may be usefully combined with another pharmacologically active compound, or two or more other pharmacologically active compounds, particularly in the treatment of inflammatory, pain and urological diseases or disorders. For example, a P2X7 antagonist, particularly a compound of the present invention, or a pharmaceutically acceptable salt or solvate thereof, as defined above, may be administered simultaneously, sequentially or separately in combination with one or more agents.

[0081] Such combinations offer significant advantages, including synergistic activity, in therapy.

[0082] The composition may contain from 0.1% to 99% by weight, preferably from 10% to

60% by weight, of the active material, depending on the method of administration. The dose of the compound used in the treatment of the aforementioned disorders may vary in the usual way with the seriousness of the disorders, the weight of the sufferer, and other similar factors.

[0083] A therapeutically effective amount of a compound of the present invention or a pharmaceutical composition thereof includes a dose range from about 0.05 mg to about 3000 mg, in particular from about 1 mg to about 1000 mg or, more particularly, from about 10 mg to about 500 mg of active ingredient in a regimen of about once a day or more than once a day, for example two, three or four times a day for an average (70 kg) human; although, it is apparent to one skilled in the art that the therapeutically effective amount for active compounds of the invention may vary as well the diseases, syndromes, conditions, and disorders being treated.

[0084] For oral administration, a pharmaceutical composition is preferably provided in the form of tablets containing about 0.01, about 0.1, about 1, about 10, about 50, about 100, about 150, about 200, about 250, and about 500 milligrams of the inventive compound as the active ingredient.

[0085] Advantageously, a compound of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three and four times daily.

[0086] Optimal dosages of a compound of the present invention to be administered may be readily determined and may vary with the particular compound used, the mode of administration, the strength of the preparation, and the advancement of the disease, syndrome, condition, or disorder. In addition, factors associated with the particular subject being treated, including subject age, weight, diet and time of administration, will result in the need to adjust the dose to achieve an appropriate therapeutic level.

[0087] The above dosages are thus exemplary of the average case. There can be individual instances wherein higher or lower dosage ranges are merited of course, and such are within the scope of this invention.

[0088] Compounds of the present invention may be administered in any of the foregoing compositions and dosage regimens or by means of those compositions and dosage regimens established in the art whenever use of a compound of the present invention is required for a subject in need thereof.

[0089] General Synthesis

Throughout the instant application, the following abbreviations are used with the following meanings:

[0090] AcOH: acetic acid

AIBN: 2,2'-azobis(isobutyronitrile)

BINAP: 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl

BzATP: 2'(3')-O-(4-benzoylbenzoyl)adenosine-5'-triphosphate
DAST N,N-diethylaminosulfur trifluoride
d.e.: diastereomeric excess
Deoxo-Fluor(trademark): bis(2-methoxyethyl)aminosulfur trifluoride
Dess-Martin periodinane: 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one
DMF: N,N-dimethylformamide
DMSO: dimethylsulfoxide
DMT-MM: 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium Chloride
e.e.: enantiomeric excess
EtOAc: ethyl acetate
EtOH: ethanol
ESI: electrospray ionization
Ex: example(s)
h: hour(s)
HBTU: N,N,N',N'-tetramethyl-O-(1H-benzotriazol-1-yl)uronium hexafluorophosphate
HPLC: high-performance liquid chromatography
(M)Hz: (mega)hertz
IM: intermediate(s)
LHMDS: lithium hexamethyldisilazide
mCPBA: m-chloroperbenzoic acid
MeCN: acetonitrile
MeOH: methanol
min: minute(s)
MS: mass spectrometry
n-Bu₃P: tri-n-butylphosphine
NIS: N-iodosuccinimide
NMP: N-methyl-2-pyrrolidinone
NMR: nuclear magnetic resonance
NaBH₄: sodium tetrahydroborate
NaH: sodium hydride
NaHMDS: sodium hexamethyldisilazide
NBS: N-bromosuccinimide
NFSI: N-fluoro-N-(phenylsulfonyl)benzenesulfonamide
NMO: N-methylmorpholine oxide
NH gel: amino bound silica gel
Pd-C: palladium on carbon
RuCl(p-cymene)[(S,S)-Ts-DPEN]:
[(S,S)-N-(2-amino-1,2-diphenylethyl)-p-toluenesulfonamide]chloro(p-cymene)rutheni

um(II)

RuCl(p-cymene)[(R,R)-Ts-DPEN]:

[(R,R)-N-(2-amino-1,2-diphenylethyl)-p-toluenesulfonamide]chloro(p-cymene)ruthenium(II)

SCX: super cation exchange resin

TBAF tetrabutylammonium fluoride

T₃P(trademark): propylphosphonic anhydride

TBS: tert-butyldimethylsilyl

TBSCl: tert-butyldimethylsilyl chloride

TEMPO: 2,2,6,6-tetramethylpiperidine N-oxide

TFA: trifluoroacetic acid

THF: tetrahydrofuran

TLC: thin layer chromatography

TMAD: 1,1'-azobis(N,N-dimethylformamide)

TMSCl: trimethylsilyl chloride

Togni reagent: 1-trifluoromethyl-3,3-dimethyl-1,2-benziodoxole

tR: retention time

UV: ultraviolet

Xantphos 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene


[0091] The compounds described herein are prepared using techniques known to one skilled in the art through the reaction sequences depicted in Schemes 1-18 as well as by other methods. Furthermore, in the following schemes, where specific acids, bases, reagents, coupling agents, solvents, etc. are mentioned, it is understood that other suitable acids, bases, reagents, coupling agents, solvents, etc. may be used and are included within the scope of the present invention.

[0092] The use of "protecting groups" (PG) is well known in the art (see for example "Protective Groups in Organic Synthesis Forth Edition edited by T. W. Greene et al. (John Wiley & Sons, 2006)". For the purposes of this discussion, it will be assumed that such protecting groups are as necessary in place.

[0093] All starting materials in the following general syntheses may be commercially available or obtained by conventional methods known to those skilled in the art.

[0094] All of the 5,6,7,8-tetrahydroquinoline and 6,7-dihydro-5H-cyclopenta[b]pyridine derivatives of the formula I can be prepared by the procedures described in the general methods presented below or by the specific methods described in the Examples section and the Preparations section, or by routine modifications thereof. The present invention also encompasses any one or more of these processes for preparing the compounds of formula I, in addition to any novel intermediates used therein.

[Chem.5]




[0095] In the following general methods, X, R¹, R², R³, R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, n, p, and q are as previously defined for 5,6,7,8-tetrahydroquinoline and 6,7-dihydro-5H-cyclopenta[b]pyridine derivatives of the formula I unless otherwise stated.

Syntheses detailing the preparation of the compounds of formula I in a sequential manner are presented in the following reaction schemes. The phrase "amidation" is used here, and it means coupling a carboxylic acid with an amine using a coupling agent such as HBTU, DMT-MM, and T₃P(trademark) in an inert solvent such as DMF and dichloromethane in the presence of a base such as trialkylamine at temperature of 0 to 50 °C.

[0096] The following illustrates a preparation of the desired compounds of formula I (Scheme 1). X is N (II and I-a), X is N-oxide (I-b), PG is a carboxyl-protecting group, R¹, R², R³, R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, n, p, and q are as defined above.

[Chem.6]




Scheme 1

[0097] The compounds of formula I-a can be prepared by deprotection of carboxyl-protecting group of the compounds formula II and then amidation between the compounds of formula III. Typical PG includes, but not limited to: methyl, ethyl, t-butyl. The typical deprotection methods of PG are described in above reference [Protective Groups in Organic Synthesis Forth Edition]. The following amidation method is described in above. Compounds of formula II can be prepared as described in the experimental part below. The compounds of formula I-b can be prepared by

oxidation of the compounds of formula I-a with an oxidizing reagent such as mCPBA in a solvent such as halogenated hydrocarbons at temperature of -10 to 40 °C.

[0098] The compounds of formula II use for the synthesis of compounds of the present invention are prepared as described in Schemes 2, 3, and 4. R^3 , R^4 , p , n , and PG are as defined above. The compounds of formula II can be prepared from the compounds of formula IV by the following steps; alkenylalkylation (Step 1), intramolecular Heck reaction (Step 2), alcoholysis (Step 3), nucleophilic addition (Step 4), oxidative cleavage (Step 5), and reduction (Step 6).

[0099] [Chem.7]




[0100] The compounds of formula V can be prepared by alkylation of the compounds of formula IV with an alkylation reagent such as alkenylalkylhalides in the presence of the base such as NaHMDS in the suitable solvent such as THF at temperature of -100 to -60 °C (Step 1). The compounds of formula VI can be prepared from the compounds of formula V by intramolecular Heck reaction which is carried out with a palladium catalyst and a base such as $Pd(Ac)_2$ and triethylamine in the presence of a phosphine ligand such as BINAP, Xantphos in a suitable solvent such as MeCN at temperature of 50- to 150 °C (Step 2). The compounds of formula II-e-1 can be prepared by alcoholysis of the compounds of formula VI with an acids such as TMSCl, HCl in the suitable (aqueous) alcohols such as MeOH at temperature of 50- to 100 °C (Step 3). The compounds of formula II-e-2 can be prepared by nucleophilic addition the compounds of formula II-e-1 with an electrophile such as NFSI or Togni reagent in the presence of a base such as NaHMDS in a suitable solvent such as THF at temperature of -100 to -60 °C (Step 4). The compounds of formula II-c can be prepared by oxidative cleavage reaction of the compounds of formula II-e-2 with an oxidizing reagent such as ozone and following treatment with a reducing agent such as dimethylsulfide or triphenylphosphine in a suitable solvent such as alcohol and/or halogenated hydrocarbons at temperature of -80 to -40 °C (Step 5). The compounds formula II-d can be prepared by reduction of the compounds formula II-c with a reducing reagent such as $NaBH_4$ in a suitable solvent such as MeOH, THF at temperature of -10 to 70 °C. The alternative method for the preparation of the compounds

of formula II-d from II-c by transfer hydrogenation with a hydrogen donor such as formic acid in the presence of metal catalyst such as RuCl(p-cymene)[Ts-DPEN] and a base such as triethylamine in a suitable solvent such as DMF at temperature of -10 to 70 °C (Step 6). The diastereo isomers can be separated by silica gel column chromatography or preparative TLC. In the Scheme 2, the reaction of Step 4 can be skipped to afford the compounds of formulae II-c and II-d, which are hydrogen as R⁴.

[0101] The following illustrates a preparation of the desired compounds of formula II-v (Scheme 3). The compounds of formula II-v can be prepared from the compounds of formula II-c by the following steps; Horner-Wadsworth-Emmons (HWE) reaction (Step 1), and hydrogenation (Step 2).

[Chem.8]




Scheme 3

[0102] The HWE reaction is carried out for preparation of the compounds of formula VII, the compounds of formula II-c treated with a trialkyl phosphono acetate such as t-butyl dimethylphosphonoacetate in the presence of a base in a suitable solvent such as THF at temperature of -40 to 50 °C (Step 1). The compounds of formula II-v can be prepared by hydrogenation of the compounds of formula VII with a hydrogen in the presence of a metal catalyst such as platinum oxide, palladium in a suitable solvent such as alcohols at temperature of 0 to 40 °C.

[0103] The following illustrates a preparation of the desired compounds of formulae II-s and II-y (Scheme 4).

[Chem.9]




Scheme 4

[0104] The compounds of formula II-s can be prepared by alkylation of the compounds of formula II-d with an alkylation reagent such as alkylhalides, alkenylalkylhalides in the presence of a base, and/or additives such as silver(I)oxide in a suitable solvent such as THF, NMP at temperature of -10 to 100 °C (Step 1a). The compounds of formula II-y

can be prepared by deoxygenative fluorination of the compounds of formula II-d with a deoxygenative fluorination reagent such as DAST, Deoxo-Fluor(trademark) in a suitable solvent such as halogenated hydrocarbons, ethers at temperature of -50 to 20 °C (Step 1b).

[0105] Compounds of formula III, if not commercially available, can be prepared by known procedures or following procedures outlined in Scheme 5. R⁵ and R⁶ are hydrogen, q is 0, R⁸ is aryl. The compounds formula III-a can be prepared from the compounds of formula VIII by the following steps; bromination (Step 1), fluorination (Step 2), and reduction (Step 3). The compounds of formula III-b can be prepared from the compounds of formula VIII by the following steps; bromination (Step 1), nucleophilic substitution (Step 4), and reduction (Step 3).

[0106] [Chem.10]




Scheme 5

[0107] The bromination is carried out for the compounds of formula VIII with a bromine source such as NBS in the presence of a radical initiator such as AIBN in a suitable solvent such as carbon tetrachloride at temperature of 50- to 100 °C to afford the compounds of formulae IX and X (Step 1). The fluorination to the compounds of formula IX is carried out with a fluorine source such as silver tetrafluoroborate to afford the compounds of formula XI (Step 2). The compounds of formula XII can be prepared by nucleophilic substitution of the compounds of formula X with a suitable nucleophile such as sodium acetate in a suitable solvent such as AcOH at temperature of 80- to 120 °C. The compounds of formulae III-a and III-b can be prepared by reduction of the compounds of formulae XI and XII with a suitable reducing reagent such as borane-THF complex in a suitable solvent such as THF at temperature of -10 to 80 °C.

[0108] The following illustrates a preparation of the desired compounds of formulae I-c and I-d (Scheme 6). R¹ and R² form ketone (I-c) and R¹ is hydrogen, R² is hydroxyl (I-d), R³, R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, n, p, and q are as defined above.

[0109]

[Chem.11]




Scheme 6

[0110] The compounds of formula I-d can be prepared by the same procedure as described in the Step 6 in Scheme 2. The diastereoisomers of the compounds of formula I-d can be separated by silica gel column chromatography or preparative TLC. The compounds of formula I-c can be prepared by oxidation of the compounds of formula I-d with an oxidizing reagent such as Dess-Martin periodinane, TEMPO, and N-t-butylphenylsulfinimidoyl chloride in a suitable solvent such as dichloromethane, toluene at temperature of -10 to 70 °C.

[0111] The following illustrates a preparation of the desired compounds of formulae I-c, and I-f (Scheme 7). R¹ and R² form methylene (I-e), R¹ and R² form ketone (I-c) and R¹ is hydroxyl, R² is hydroxymethyl (I-f), R³, R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, n, p, and q are as defined above.

[0112] [Chem.12]




Scheme 7

[0113] The compounds of formula I-c can be prepared by the same procedure as described in the Step 5 in Scheme 2. The compounds of formula I-f can be prepared by dihydroxylation of the compounds of formula I-e with an oxidizing reagent such as osmium tetroxide in the presence or absence of co-oxidants such as NMO in a suitable solvent such as alcohols and water at temperature of -10 to 60 °C.

[0114] The following illustrates a preparation of the desired compounds of formulae I-g, and I-h (Scheme 8). R¹ and R² form ketone (I-c), R¹ and R² form methylene (I-e), R¹ and R² form epoxide (I-g), and R¹ is hydroxyl, R² is nucleophilic group (Nu) such as

acyloxy groups, hydroxides, alkoxides, fluoride, cyano, amines, azides, sulfides (I-h), R³, R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, n, p, and q are as defined above.


[0115] [Chem.13]



[0116] The compounds of formula I-g can be prepared by epoxidation of the compounds of formula I-e with NBS in a suitable solvent such as aqueous tert-butanol at temperature of 30- to 80 °C and then treatment with a base such as NaOH at temperature of -10 to 10 °C. The compounds of formula I-g can be prepared by epoxidation of the compounds of formula I-c with a sulfur ylide reagent such as dimethyloxosulfonium methylide in a suitable solvent such as DMSO at temperature of -10 to 50 °C. The diastereo isomers can be separated by silica gel column chromatography or preparative TLC. The nucleophilic addition of the compounds of formula I-g for the synthesis of the compounds of formula I-h is carried out with the nucleophiles such as alkali metal hydroxide, alkali metal alkoxide, acyloxy groups, alkali metal cyanide, alkali metal fluoride, substituted or unsubstituted alkylsulfide, and amine or the corresponding ammonium salt of nucleophiles in a suitable solvent such as halogenated hydrocarbons, ethers, aromatic hydrocarbons, amides, inert amines, sulfoxides at temperature of -80 to 100 °C in the presence or in the absence of a catalyst such as copper salts or zinc salts.


[0117] The following illustrates a preparation of the desired compounds of formulae I-i, I-j, and I-k (Scheme 9). LvG and LvG¹ are leaving groups such as halogen, hydroxyl, sulfonate, azide, imidazole. In the case where R^{10a} is LvG¹ substituted C₁-C₂alkyl and Y¹ and Y² are hydrogen (I-h-1 and I-j) or R^{10a} is hydrogen and Y¹ and Y² form carbonyl (I-h-1, I-i, and I-k), R³, R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, R^{9a}, n, p, and q are as defined above.

[0118] [Chem.14]



[0119] The compounds of formula I-i can be prepared by amidation of the compounds of formula I-h-1 (R^{10a} is hydrogen) with a carboxylic acid XIII (LvG is hydroxyl) under “amidation” condition as described above, or with a activated carboxylic acid XIII such as acid halide (LvG is halogen) in a suitable base such as trialkylamine in a suitable solvent such as dichloromethane at temperature of -30 to 20 °C. In the case where n is 0, the cyclized compounds of formula I-k can be prepared in this step without following “cyclization” step. In the case where LvG¹ is not hydroxyl, the compounds of formulae I-j and I-k can be prepared by cyclization of the compounds of formula I-i or I-h-1 (R^{10a} is LvG¹ substituted C₁-C₂alkyl) in the presence of a suitable base such as potassium tert-butoxide in a suitable solvent such as alcohol and /or halogenated hydrocarbons at temperature of -20 to 50 °C. In the case where LvG¹ is hydroxyl, the compounds of formulae I-j and I-k can be prepared by intramolecular Mitsunobu reaction of the compounds of formula I-h-1 or I-i with a Mitsunobu reagents such as N,N,N',N'-tetraalkylazodicarboxamide or dialkylazodicarboxylate in the presence of trialkylphosphine or triarylphosphine in a suitable solvent such as THF at temperature of -40 to 80 °C.

[0120] The following illustrates a preparation of the desired compounds of formula I-l (Scheme 10). R^1 is hydroxyl, R^2 is C₁alkylthio- R^{9a} , m is 0 (I-h-2), or m is 1 or 2 (I-l), R^3 , R^4 , R^5 , R^6 , R^{7a} , R^{7b} , R^8 , R^{9a} , n, p, and q are as defined above.




Scheme 10

[0121] The compounds of formula I-l can be prepared by oxidation of the compounds formula I-h-2 with an oxidizing reagent such as mCPBA in a suitable solvent such as halogenated hydrocarbons at temperature of -20 to 50 °C.

[0122] The following illustrates a preparation of the desired compounds of formulae I-m and I-n (Scheme 11). R^1 is hydroxyl, R^2 is C₁alkylazide (I-h-3), or R^1 and R^2 form aziridine (I-m), or R^1 is amino, R^2 is hydroxyC₁alkyl (I-n), R^3 , R^4 , R^5 , R^6 , R^{7a} , R^{7b} , R^8 , R^{9a} , n, p, q, and PG are as defined above.

[0123] [Chem.16]




Scheme 11

[0124] The compounds of formula I-m can be prepared by aziridine formation of the compounds formula I-h-3 with a triphenylphosphine in a suitable solvent such as aqueous MeCN at temperature of 20- to 100 °C (Step 1). The compounds of formula I-n can be prepared by following reactions, protection of aziridine (Step 2), nucleophilic ring opening and hydrolysis (Step 3), and deprotection (Step 4). The suitable PG such as 2-nitrophenylsulfonyl group, the protection and deprotection reactions can be performed in the usual manner. The compounds of formula XV can be prepared by nucleophilic addition and then hydrolysis of the compounds of formula XIV with a metal carboxide such as alkali metal acetoxide in a suitable solvent such as DMF at temperature of 60- to 120 °C and followed by hydrolysis with an alkali metal hydroxide such as NaOH in a suitable solvent such as DMF at room temperature. The compounds of formula I-n can be prepared by deprotection of the compounds of formula XV by the usual manner such as 4-mercaptobenzoic acid in a suitable solvent such as DMF at temperature of 70- to 120 °C.

[0125] The following illustrates a preparation of the desired compounds of formula I-o (Scheme 12). R¹ is amino, R² is hydroxyC₁alkyl (I-n), R³, R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, R^{9a}, LvG, LvG¹, n, p, and q are as defined above.

[0126]


[Chem.17]



[0127] The compounds of formula I-o can be prepared by the same procedure as described in Scheme 9 from the compounds of formula I-n via the compounds of formula XVI.

[0128] The following illustrates a preparation of the desired compounds of formulae I-p, I-q, and I-r (Scheme 13). R¹ is hydrogen and R² is hydroxyl (I-d), R is methyl or phenyl, R² is thioester (I-p), R² is sulfide (I-q), R² is sulfoxide or sulfone (I-r), R³, R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, R^{9a}, m, n, p, and q are as defined above.

[0129] [Chem.18]



Scheme 13


[0130] The compounds of formula I-p can be prepared by Mitsunobu reaction of the compounds of formula I-d with a thiocarboxylic acid such as thioacetic acid, thiobenzoic acid under Mitsunobu reaction condition as described above.

The compounds of formula I-q can be prepared by S-alkylation of the compounds of formula I-p with an alkylation reagent such as dimethyl sulfonate, alkylhalides in the presence of alkali metal hydroxide in a suitable solvent such as halogenated hydrocarbons, ethers, alcohols at temperature of -20 to 50 °C. The compounds of formula I-r can be prepared by the same procedures as described in Scheme 10.

[0131] The following illustrates a preparation of the desired compounds of formulae I-s, I-t and I-u (Scheme 14). R¹ is hydrogen and R² is alkenylalkoxide (I-s), R² is hydrox-

yalkoxide (I-t), R² is hydroxy substituted hydroxyalkoxide (I-u), R⁴, R⁵, R⁶, R^{7a}, R^{7b}, and R⁸ are as defined above.

[0132] [Chem.19]




Scheme 14

[0133] The compounds of formula I-t can be prepared by oxidation of the compounds of formula I-s with an oxidizing reagent such as ozone at temperature of -100 to -60 °C and then treated with sodium tetrahydroborate in a suitable solvent such as halogenated hydrocarbons and/or alcohols. The compounds of formula I-u can be prepared by the same procedure of the “dihydroxylation” in Scheme 7.

[0134] The following illustrates a preparation of the desired compounds of formulae I-w, and I-x (Scheme 15). R¹ is hydrogen and R² is “Protected carboxylic acid” alkyl (I-v), R² is carboxylic acid alkyl (I-w), R² is aminocarboxylalkyl (I-x), R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, R^{9a} and R^{10a} are as defined above.

[0135] [Chem.20]




Scheme 15

[0136] The compounds of formula I-w can be prepared by deprotection of the PG from the compounds of formula I-v. The compounds of formula I-x can be prepared by “amidation” of the compounds of formula I-w with a coupling agent and amines. The general condition of “amidation” is described in above.

[0137] The following illustrates a preparation of the desired compounds of formula II-e-2 (Scheme 16). R⁴ is hydroxyl or fluoro, R³, p, n, and PG are as defined above. The compounds of formula II-e-2 also can be prepared from the compounds of formula XVII by the following steps; alkenylacylation (Step 1), intramolecular Heck reaction (Step 2), cyanohydration and deoxygenative fluorination (Step 3), and alcoholysis (Step 4).

[0138]

[Chem.21]




[0139] The compounds of formula XVIII can be prepared by alkenylacylation of the compounds of formula XVII, which is carried out by the following procedure. The preparation of the Grignard reagents of the compounds of formula XVII is carried out by the treatment of the compounds of formula XVII with a Turbo Grignard reagent such as 2-propylmagnesium chloride lithium chloride complex in the suitable solvent such as THF at temperature of -20 to 10 °C. The compounds of formula XVIII can be prepared by the following addition of an alkenylalkylacylhalides in the suitable solvent such as THF at temperature of -50 to 0 °C (Step 1). The compounds of formula XIX can be prepared from the compounds of formula XVIII by the same procedure as described in Scheme 2 Step 2 (Step 2). The compounds of formula XX can be prepared by cyanohydration and following deoxygenative fluorination of the compounds of formula XIX. The cyanohydration of the compounds of formula XIX is carried out with the cyanide source such as trimethylsilyl cyanide in the presence of a catalyst such as NMO in a suitable solvent such as THF at temperature of -10 to 30 °C. And then deoxygenative fluorination is carried out by the same procedure as described in Scheme 4 Step 1b (Step 3). The compounds of formula II-e-2 can be prepared from the compounds of formula XX by the same procedure as described in Scheme 2 Step 3 (Step 4).

[0140] The following illustrates a preparation of the desired compounds of formula II-e-2 (Scheme 17). R^3 , R^4 , p , n , and PG are as defined above. The compounds of formula II-e-2 (R^4 is fluoro) also can be prepared from the compounds of formula XVII by the following steps; alkenylalkylation (Step 1), intramolecular Heck reaction (Step 2), and deoxygenative fluorination (Step 3).

[0141]

[Chem.22]




Scheme 17

[0142] The compounds of formula XXII, II-e-2 (R⁴ is hydroxyl), and II-e-2 (R⁴ is fluoro) can be prepared by the same procedure as described in Scheme 16 Step 1 with an alkylation reagent of XXI (Step 1), Scheme 2 Step 2 (Step 2), and Scheme 4 Step 1b (Step 3).

[0143] The following illustrates a preparation of the desired compounds of formula I-g (Scheme 18). R³, R⁴, R⁵, R⁶, R^{7a}, R^{7b}, R⁸, p, n, q, and PG are as defined above. The compounds of formula I-g also can be prepared from the compounds of formula II-e-2 by the following steps; hydrolysis (Step 1), iodolactonization (Step 2), and nucleophilic amidation (Step 3).

[0144] [Chem.23]




Scheme 18

[0145] The compounds of formula XXIII can be prepared from the compounds of formula II-e-2 by hydrolysis with a base such as NaOH in a suitable solvent such as MeOH, THF at temperature 0 to 40 °C. The compounds of formula XXIII also can be prepared by hydrolysis with an enzyme such as lipase in a suitable solvent such as phosphate

buffer at temperature 20 to 40 °C (Step 1). The compounds of formula XXIV can be prepared from the compounds of formula XXIII by iodolactonization which is carried out in the presence of iodine or NIS and a base such as K_2CO_3 in a suitable solvent such as MeCN, DMF, DMSO at temperature 0 to 40 °C (Step 2). The Step 1 and Step 2 can be carried out in one-pot. The compounds of formula I-g can be prepared by nucleophilic amidation of the compounds of formula XXIV with an amine III in the presence of a base such as K_2CO_3 in a suitable solvent such as DMF, DMSO, MeCN at temperature of 0 to 40 °C (Step 3).

[0146] The following illustrates a preparation of the desired compounds of formula II-e-2 (Scheme 19). R^4 is fluoro, R^3 , p, n, and PG are as defined above. The compounds of formula II-e-2 (R^4 is fluoro) also can be prepared from the compounds of formula V by the following steps; reduction (Step 1), organocatalytic fluorination (Step 2), oxidation (Step 3), protection (Step 4), and intramolecular Heck reaction (Step 5).

[0147] [Chem.24]



The compounds of formula XXV can be prepared by reduction of the compounds of formula V with a reducing reagent such as Diisobutylaluminium hydride in a suitable solvent such as toluene, THF at temperature of -78 to 0 °C (Step 1). The compounds of formula XXVI can be prepared by organocatalytic fluorination of the compounds of formula XXV, which is carried out with an organocatalyst and fluorination reagent such as proline, imidazolidinone organocatalysts and NFSI, Selectfluor (registered trademark) in a suitable solvent such as DMF, THF, toluene, and ether at temperature -20 to 50 °C (Step 2). The compounds of formula XXVII can be prepared by oxidation of the compounds of formula XXVI with an oxidizing reagent such as sodium chlorite in a suitable solvent such as tert-BuOH at temperature of -10 to 40 °C (Step 3). The compound of formula XXVIII can be prepared by protection of the compound of formula XXVII (Step 4). The compound of formula II-e-2 (R^4 is fluoro) can be prepared from the compounds of formula XXVIII by cyclization with a palladium catalyst and a base such as $Pd(OAc)_2$ and N,N-Diisopropylethylamine in the presence

of a phosphine ligand such as BINAP, Xantphos in a suitable solvent such as DMF at temperature of 50 to 150 °C as described in the Step 5. The compound of formula II-e-2 (R⁴ is fluoro) can be also prepared from the compounds of formula V by the same procedure as described in the Step 2, Step 3, and Step 4 in Scheme 2.

[0148] The invention is illustrated in the following non-limiting examples in which, unless stated otherwise: all reagents are commercially available, all operations were carried out at room or ambient temperature, that is, in the range of about 18-25 °C; evaporation of solvent was carried out using a rotary evaporator under reduced pressure with a bath temperature of up to about 60 °C; reactions were monitored by thin layer chromatography (TLC) and reaction times are given for illustration only; the structure and purity of all isolated compounds were assured by at least one of the following techniques: TLC (Merck silica gel 60 F₂₅₄ precoated TLC plates or Merck NH₂ F₂₅₄₈ precoated HPTLC plates), mass spectrometry or nuclear magnetic resonance (NMR). Yields are given for illustrative purposes only. Flash column chromatography was carried out using Merck silica gel 60 (230-400 mesh ASTM) or Fuji Silysia Chromatorex(trademark) DU3050 (Amino Type, 30-50 microm) or Biotage silica (32-63 mm, KP-Sil) or Biotage amino bounded silica (35-75 mm, KP-NH). SCX cartridge column chromatography was carried out using Biotage ISOLUTE(trademark) SCX-2 (1 g, 6 mL) SPE column. The purification of compounds using HPLC was performed by the following apparatus and conditions; Apparatus; Waters MS-trigger AutoPurification(trademark) system Column; Waters XTerra C18, 19 x 50 mm, 5 mm particle, solvent systems; Methanol or acetonitrile / 0.05%(v/v) formic acid aqueous solution, or; methanol or acetonitrile / 0.01% (v/v) ammonia aqueous solution. Low-resolution mass spectral data (ESI) were obtained by the following apparatus and conditions: Apparatus; Waters Alliance HPLC system on ZQ or ZMD mass spectrometer and UV detector. NMR data was determined at 270 MHz (JEOL JNM-LA 270 spectrometer), 300 MHz (JEOL JNM-LA300), 400 MHz (JEOL ECZ 400S) using deuterated chloroform (99.8% D) or dimethylsulfoxide (99.9% D) as solvent unless indicated otherwise, relative to tetramethylsilane (TMS) as internal standard in parts per million (ppm); conventional abbreviations used are: s = singlet, d = doublet, t = triplet, q = quartet, quin = quintet, m = multiplet, br = broad, etc. Chemical symbols have their usual meanings; microL (microliter(s)), microg (microgram(s)), M (mol(s) per liter), L (liter(s)), mL (milliliter(s)), g (gram(s)), mg (milligram(s)), mol (moles), mmol (millimoles).

[0149] Conditions for determining HPLC retention time:

Method A:

Apparatus: Waters ACQUITY UPLC / ACQUITY PDA Detector / ZQ 2000

Column: Waters ACQUITY BEH C18, 2.1 x 100 mm

Column temperature: 60 °C

PDA detection (scan range): 200 - 400 nm

MS detection: ESI positive/negative mode

Solvents:

A1: 10 mM ammonium acetate aqueous solution

B1: acetonitrile

[0150]

Time(min)	A1(%)	B1(%)
0	95	5
0.1	95	5
1.8	5	95
2.3	95	5
Run time		3 min
Flow rate		0.7 mL/min

[0151] Method B:

Apparatus: Waters ACQUITY UPLC / ACQUITY PDA Detector / ZQ 2000

Column: YMC Meteoric core C18, 2.1 x 100 mm

Column temperature: 60 °C

PDA detection (scan range): 200 - 400 nm

MS detection: ESI positive/negative mode

Solvents:

A1: 10 mM ammonium acetate aqueous solution

B1: acetonitrile

[0152]

Time(min)	A1(%)	B1(%)
0	95	5
0.1	95	5
1.8	5	95
2.3	5	95
2.31	95	5
Run time		3 min
Flow rate		0.7 mL/min

[0153] Method C:

Apparatus: Waters ACQUITY UPLC / ACQUITY PDA Detector / ZQ 2000

Column: Waters ACQUITY BEH C18, 2.1 x 100 mm

Column temperature: 60 °C

PDA detection (scan range): 200 - 400 nm

MS detection: ESI positive/negative mode

Solvents:

A1: 10 mM ammonium acetate aqueous solution

B1: acetonitrile

[0154]	Time(min)	A1(%)	B1(%)
	0	95	5
	0.1	95	5
	1.8	5	95
	2.3	5	95
	2.31	95	5
	Run time	3 min	
	Flow rate	0.7 mL/min	

[0155] Method D:

Apparatus: Waters ACQUITY UPLC / ACQUITY PDA Detector / ZQ 2000

Column: YMC Triart C18, 2.1 x 100 mm, 1.9 microm particle

Column temperature: 60 °C

PDA detection (scan range): 200 - 400 nm MS detection: ESI positive/negative mode

Solvents:

A1: 10 mM ammonium acetate aqueous solution

B1: acetonitrile

[0156]	Time(min)	A1(%)	B1(%)
	0	90	10
	0.05	90	10
	1.9	5	95
	2.5	5	95
	2.51	90	10
	Run time	3 min	
	Flow rate	0.75 mL/min	

[0157] Method E

Apparatus: Waters Alliance 2695 / 2996 PDA

Column: DAICEL CHIRALCEL OJ-H, 4.6 mm x 250 mm

Column temperature: 40 °C

UV detection: 270 nm

Solvents: n-hexane/2-propanol/diethylamine = 95/5/0.1

Flow rate: 1 mL/min

Run time: 40 min

[0158] Method F

Apparatus: Waters Alliance 2695 / 2996 PDA

Column: DAICEL CHIRALPAK IC-3, 4.6 mm x 250 mm

Column temperature: 40 °C

UV detection: 265 nm

Solvents:

A2: 0.1% diethylamine in n-hexane

B2: 0.1% diethylamine in 2-propanol

[0159]

Time(min)	A2(%)	B2(%)
0	85	15
15	85	15
30	50	50
40	50	50
40.01	85	15
Run time		60 min
Flow rate		1 mL/min

[0160] Method G

Apparatus: Waters Alliance 2695 / 2996 PDA

Column: DAICEL CHIRALPAK AD-H 4.6 mm x 250 mm

Column temperature: 40 °C

UV detection: 265 nm

Solvents: n-hexane/ethanol/diethylamine = 94/6/0.1

Flow rate: 1 mL/min

Run time: 30 min

[0161] Method H

Apparatus: Waters Alliance 2695 / 2996 PDA

Column: DAICEL CHIRALCEL OD-H, 4.6 mm x 250 mm

Column temperature: 40 °C

UV detection: 270 nm

Solvents: n-hexane/2-propanol/diethylamine = 95/5/0.1

Flow rate: 1 mL/min

Run time: 20 min

[0162] Method I

Apparatus: Waters Alliance 2695 / 2996 PDA

Column: DAICEL CHIRALPAK AD-H, 4.6 x 250 mm

Column temperature: 40 °C

UV Detection: 265 nm

Solvents: n-hexane/ethanol/diethylamine = 88/12/0.1

Flow rate: 1 mL/min

Run time: 30 min

[0163] Method J

Apparatus: Waters Alliance 2695 / 2996 PDA

Column: DAICEL CHIRALCEL OD-H, 4.6 mm x 250 mm

Column temperature: 40 °C

UV detection: 265 nm

Solvents: n-hexane/2-propanol/diethylamine = 85/15/0.1

Flow rate: 1 mL/min

Run time: 50 min

[0164] Method K

Apparatus: Waters Alliance 2695 / 2996 PDA

Column: DAICEL CHIRALCEL OJ-H, 4.6 mm x 250 mm

Column temperature: 40 °C

UV detection: 265 nm

Solvents: n-hexane/ethanol/diethylamine = 90/10/0.1

Flow rate: 1 mL/min

Run time: 45 min

[0165] Method L

Apparatus: Waters Alliance 2695 / 2996 PDA

Column: Daicel CHIRALPAK IC

Column Temperature: 40 °C

UV detection: 245 nm

Solvents: n-hexane/2-propanol = 90/10

Flow rate: 1 mL/min

Run time: 35 min

[0166] Preparation of Intermediates II

General Procedure: Scheme 2, Step 1

To a solution of substrate (1.0 eq.) in THF was added dropwise 1.1 M NaHMDS in THF solution (1.2 eq.) at -78 °C under N₂ atmosphere. After addition, the mixture was stirred at -78 °C for 2 h. A THF solution of alkenylalkylhalide was added dropwise to the mixture at -78 °C and the mixture was stirred at this temperature for 1 h. The mixture was warmed to room temperature. After being stirred at room temperature until complete reaction, the mixture was quenched with water. The mixture was extracted with EtOAc twice and washed successively with 10% aq. citric acid, aq. NaHCO₃, and brine. The extracts were combined, dried over Na₂SO₄, and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography to afford the following intermediates (Table 1).

[0167]

[Table 1]

Intermediates	Structure	Chemical Name	Substrate	alkenylalkylhalide
V-1		2-(2-chloropyridin-3-yl) hex-5-enenitrile		
V-2		2-(2-chloropyridin-3-yl) pent-4-enenitrile		
V-3		2-(2-chloro-5-fluoropyridin-3-yl)hex-5-enenitrile		
V-4		2-(2-chloro-5-methylpyridin-3-yl)hex-5-enenitrile		

[0168] IM V-1

¹H NMR (CDCl₃) delta 8.40 (1 H, dd, J = 4.9, 1.8 Hz), 7.92 (1 H, dd, J = 7.3, 1.8 Hz), 7.34 (1 H, dd, J = 7.3, 4.9 Hz), 5.80 (1 H, ddt, J = 17.1, 10.4, 7.3 Hz), 5.16 (1 H, br d, J = 17.1 Hz), 5.11 (1 H, br d, J = 10.4 Hz), 4.27 (1 H, dd, J = 9.2, 4.9 Hz), 2.42-2.27 (2 H, m), 2.04-1.90 (2 H, m).

MS (ESI) m/z: 207.1 (M+H)⁺.

[0169] IM V-2

¹H NMR (CDCl₃) delta 8.40 (1 H, dd, J = 4.9, 1.8 Hz), 7.89 (1 H, dd, J = 7.9, 1.8 Hz), 7.34 (1 H, dd, J = 7.9, 4.9 Hz), 5.83 (1 H, ddt, J = 17.1, 9.8, 7.3 Hz), 5.24 (1 H, br d, J = 9.8 Hz), 5.21 (1 H, br d, J = 17.1 Hz), 4.35 (1 H, dd, J = 7.9, 4.9 Hz), 2.73 (1 H, m), 2.61 (1 H, m).

MS (ESI) m/z: 193.2 (M+H)⁺.

[0170] IM V-3

¹H NMR (CDCl₃) delta 8.28 (1 H, dd, J = 15.3, 3.1 Hz), 7.69 (1 H, dd, J = 7.9, 3.1 Hz), 5.79 (1 H, ddt, J = 17.1, 10.4, 6.1 Hz), 5.18 (1 H, br dd, J = 17.1, 1.2 Hz), 5.12 (1 H, br dd, J = 10.4, 1.2 Hz), 4.24 (1 H, dd, J = 9.8, 5.5 Hz), 2.44-2.28 (2 H, m), 2.08-1.92 (2 H, m).

MS (ESI) m/z: 225.0 (M+H)⁺.

[0171] IM V-4

^1H NMR (CDCl_3) δ 8.19 (1 H, d, $J = 1.8$ Hz), 7.71 (1 H, d, $J = 1.8$ Hz), 5.80 (1 H, ddt, $J = 17.1, 10.4, 6.1$ Hz), 5.16 (1 H, br d, $J = 17.1$ Hz), 5.10 (1 H, br d, $J = 10.4$ Hz), 4.23 (1 H, m), 2.41-2.25 (2 H, m), 2.37 (3 H, s), 2.06-1.90 (2 H, m).

MS (ESI) m/z : 221.2 ($\text{M}+\text{H}$) $^+$.

[0172] General Procedure: Scheme 2, Step 2

A mixture of substrate (1.0 eq.), triethylamine (3.0 eq.), (+/-)-BINAP (0.135 eq.), and $\text{Pd}(\text{OAc})_2$ (0.135 eq.) in MeCN was heated at reflux until complete reaction. After cooling at room temperature, NH gel was added to the mixture and the mixture was filtered. The filter cake was washed with EtOAc. The filtrate was concentrated in vacuo. EtOAc was added to the resulting residue and the insoluble material was removed by filtration. Water was added to the filtrate, the mixture was extracted with EtOAc twice. The extracts were washed with brine, dried over Na_2SO_4 , and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography to afford the following intermediates (Table 2).

[0173] [Table 2]

Intermediates	Structure	Chemical Name	Substrate
VI-1		8-methylene-5,6,7,8-tetrahydroquinoline-5-carbonitrile	
VI-2		7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carbonitrile	
VI-3		3-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carbonitrile	
VI-4		3-methyl-8-methylene-5,6,7,8-tetrahydroquinoline-5-carbonitrile	

[0174] IM VI-1

^1H NMR (CDCl_3) delta 8.57 (1 H, dd, $J = 4.2, 1.8$ Hz), 7.73 (1 H, dd, $J = 7.9, 1.8$ Hz), 7.23 (1 H, dd, $J = 7.9, 4.2$ Hz), 6.38 (1 H, d, $J = 1.8$ Hz), 5.25 (1 H, d, $J = 1.8$ Hz), 4.07 (1 H, dd, $J = 7.9, 4.9$ Hz), 2.92 (1 H, m), 2.70 (1 H, m), 2.32-2.15 (2 H, m).

MS (ESI) m/z : 171.1 ($\text{M}+\text{H}$) $^+$.

[0175] IM VI-2

^1H NMR (CDCl_3) delta 8.60 (1 H, br d, $J = 5.3$ Hz), 7.81 (1 H, dd, $J = 7.9, 1.3$ Hz), 7.26 (1 H, dd, $J = 7.9, 5.3$ Hz), 6.11 (1 H, br t, $J = 2.0$ Hz), 5.51 (1 H, br t, $J = 2.0$ Hz), 4.22 (1 H, dd, $J = 8.6, 5.9$ Hz), 3.34 (1 H, m), 3.17 (1 H, m).

MS (ESI) m/z : 157.2 ($\text{M}+\text{H}$) $^+$.

[0176] IM VI-3

^1H NMR (CDCl_3) delta 8.44 (1 H, d, $J = 15.9, 3.1$ Hz), 7.47 (1 H, dd, $J = 8.6, 3.1$ Hz), 6.28 (1 H, d, $J = 1.2$ Hz), 5.24 (1 H, d, $J = 1.2$ Hz), 4.07 (1 H, dd, $J = 8.6, 4.9$ Hz), 2.94-2.87 (1 H, m), 2.73-2.64 (1 H, m), 2.34-2.27 (1 H, m), 2.23-2.15 (1 H, m).

MS (ESI) m/z : 189.1 ($\text{M}+\text{H}$) $^+$.

[0177] IM VI-4

^1H NMR (CDCl_3) delta 8.38 (1 H, d, $J = 1.2$ Hz), 7.52 (1 H, d, $J = 1.2$ Hz), 6.29 (1 H, d, $J = 1.2$ Hz), 5.18 (1 H, d, $J = 1.2$ Hz), 4.02 (1 H, m), 2.88 (1 H, m), 2.66 (1 H, m), 2.36 (3 H, s), 2.31 (1 H, m), 2.17 (1 H, m).

MS (ESI) m/z : 185.2 ($\text{M}+\text{H}$) $^+$.

[0178] General Procedure: Scheme 2, Step 3

To a solution of substrate (1.0 eq.) in MeOH (0.2 M) was added TMSCl (30 eq.) at ambient temperature. The mixture was heated at reflux until complete reaction, and then cooled to room temperature. The mixture was basified with aq. NaHCO_3 and the mixture was concentrated in vacuo to remove volatile. The resulting residue was extracted with EtOAc twice and washed with brine. The extracts were combined, dried over Na_2SO_4 , and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography to afford the following intermediates (Table 3).

[0179]

[Table 3]

Intermediates	Structure	Chemical Name	Substrate
II-e-1-1		methyl 8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate	
II-e-1-2		methyl 7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxylate	
II-e-1-3		methyl 3-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate	
II-e-1-4		methyl 3-methyl-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate	

[0180] IM II-e-1-1

¹H NMR (CDCl₃) delta 8.51 (1 H, dd, J = 4.3, 1.8 Hz), 7.52 (1 H, dd, J = 7.9, 1.8 Hz), 7.14 (1 H, dd, J = 7.9, 4.3 Hz), 6.29 (1 H, d, J = 1.8 Hz), 5.18 (1 H, d, J = 1.8 Hz), 3.88 (1 H, dd, J = 5.5, 5.5 Hz), 3.73 (3 H, s), 2.81 (1 H, m), 2.65 (1 H, m), 2.30 (1 H, m), 2.07 (1 H, m).

MS (ESI) m/z: 204.1 (M+H)⁺.

[0181] IM II-e-1-2

¹H NMR (CDCl₃) delta 8.52 (1 H, dd, J = 4.9, 1.8 Hz), 7.79 (1 H, dd, J = 7.3, 1.8 Hz), 7.16 (1 H, dd, J = 7.3, 4.9 Hz), 6.04 (1 H, br t, J = 2.4 Hz), 5.25 (1 H, br t, J = 2.4 Hz), 4.13 (1 H, dd, J = 9.2, 5.5 Hz), 3.76 (3 H, s), 3.27 (1 H, m), 3.09 (1 H, m).

MS (ESI) m/z: 190.2 (M+H)⁺.

[0182] IM II-e-1-3

¹H NMR (CDCl₃) delta 8.35 (1 H, dd, J = 15.3, 3.1 Hz), 7.26 (1 H, dd, J = 7.9, 3.1 Hz), 6.19 (1 H, s), 5.15 (1 H, d, J = 1.8 Hz), 3.87 (1 H, dd, J = 5.5, 5.5 Hz), 3.73 (3 H, s), 2.77 (1 H, m), 2.63 (1 H, m), 2.28 (1 H, m), 2.06 (1 H, m).

MS (ESI) m/z: 222.1 (M+H)⁺.

[0183] IM II-e-1-4

¹H NMR (CDCl₃) delta 8.32 (1 H, br s), 7.29 (1 H, br s), 6.21 (1 H, br s), 5.10 (1 H, br s), 3.83 (1 H, m), 3.71 (3 H, s), 2.78 (1 H, m), 2.60 (1 H, m), 2.30 (3 H, s), 2.24 (1 H, m), 2.03 (1 H, m).

MS (ESI) m/z: 218.1 (M+H)⁺.

[0184] General Procedure: Scheme 2, Step 4-A

To a stirred solution of substrate (1.0 eq.) in THF was added 1.1 M THF solution of NaHMDS (1.3 eq.) at -78 °C under N₂ atmosphere. After being stirred at -78 °C for 30 min, NFSI (1.3 eq.) was added to the mixture. The mixture was stirred at -78 °C until complete reaction and then warmed to room temperature. The mixture was poured into water and extracted with EtOAc. The extract was dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography to afford the following intermediates (Table 4).

[0185] General Procedure: Scheme 2, Step 4-B

To a stirred solution of substrate (1.0 eq.) in THF was added 1.1 M THF solution of NaHMDS (1.2 eq.) at 0 °C under N₂ atmosphere. After being stirred at 0 °C for 20 min, Togni reagent (1.2 eq.) in THF was added to the mixture. The mixture was warmed to room temperature and stirred until complete reaction. The mixture was poured into water and extracted with EtOAc. The extract was dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by silica gel column chromatography to afford the following intermediates (Table 4).

[0186]

[Table 4]

Intermediates	Structure	Chemical Name	Substrate	General Procedure: Step 4
II-e-2-1		methyl 5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate		A
II-e-2-2		methyl 5-fluoro-7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxylate		A
II-e-2-3		methyl 3,5-difluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate		A
II-e-2-4		methyl 5-fluoro-3-methyl-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate		A
II-e-2-5		methyl 5-hydroxy-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate		B

[0187] IM II-e-2-1

¹H NMR (CDCl₃) delta 8.63 (1 H, ddd, J = 4.3, 1.8, 1.2 Hz), 7.68 (1 H, dd, J = 7.9, 1.8 Hz), 7.24 (1 H, ddd, J = 7.9, 4.3, 1.8 Hz), 6.37 (1 H, br s), 5.31 (1 H, d, J = 1.2 Hz), 3.81 (3 H, s), 2.89-2.73 (2 H, m), 2.48 (1 H, m), 2.32 (1 H, m).

MS (ESI) m/z: 222.1 (M+H)⁺.

[0188] IM II-e-2-2

¹H NMR (CDCl₃) delta 8.70 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.83 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.27 (1 H, dd, J = 7.9, 4.9 Hz), 6.20 (1 H, br t, J = 2.4 Hz), 5.36 (1 H, dd, J = 1.8, 1.2 Hz), 3.83 (3 H, s), 3.58 (1 H, m), 3.20 (1 H, m).

MS (ESI) m/z: 208.2 (M+H)⁺.

[0189] IM II-e-2-3

¹H NMR (CDCl₃) delta 8.48 (1 H, m), 7.42 (1 H, dd, J = 8.6, 1.8 Hz), 6.27 (1 H, s),

5.28 (1 H, s), 3.83 (3 H, s), 2.82 (2 H, m), 2.45 (1 H, m), 2.29 (1 H, m).

MS (ESI) m/z: 240.0 (M+H)⁺.

[0190] IM II-e-2-4

¹H NMR (CDCl₃) delta 8.37 (1 H, br s), 7.36 (1 H, br s), 6.23 (1 H, br s), 5.19 (1 H, br s), 3.76 (3 H, s), 3.73 (1 H, m), 2.40-2.24 (2 H, m), 2.30 (3 H, s), 2.08 (1 H, m).

MS (ESI) m/z: 236.2 (M+H)⁺.

[0191] IM II-e-2-5

¹H NMR (CDCl₃) delta 8.55 (1 H, dd, J = 4.6, 2.0 Hz), 7.55 (1 H, dd, J = 7.9, 2.0 Hz), 7.19 (1 H, dd, J = 7.9, 4.6 Hz), 6.31 (1 H, br s), 5.26 (1 H, d, J = 2.0 Hz), 3.93 (1 H, br), 3.77 (3 H, s), 2.92-2.73 (2 H, m), 2.30 (1 H, ddd, J = 13.2, 8.6, 4.3 Hz), 2.10 (1 H, ddd, J = 13.2, 7.9, 4.6 Hz).

MS (ESI) m/z: 220.2 (M+H)⁺.

[0192] General Procedure: Scheme 2, Step 5

O₃ was bubbled into a solution of substrate (1.0 eq.) in 50% CH₂Cl₂-MeOH at -78 °C until starting material consumed. N₂ was bubbled into the mixture to remove the excess of O₃ at -78 °C. The mixture was quenched with Me₂S (2.0 eq.) and the mixture was warmed to room temperature. The mixture was concentrated in vacuo and the resulting residue was purified by silica gel column chromatography to afford the following Intermediates (Table 5).

[0193]

[Table 5]

Intermediates	Structure	Chemical Name	Substrate
II-c-1		methyl 8-oxo-5,6,7,8-tetrahydroquinoline- 5-carboxylate	 IM II-e-1-1
II-c-2		methyl 5-fluoro-8-oxo-5,6,7,8-tetrahydroq uinoline-5-carboxylate	 IM II-e-2-1
II-c-3		methyl 3,5-difluoro-8-oxo-5,6,7,8-tetrahyd roquinoline-5-carboxylate	 IM II-e-2-3
II-c-4		methyl 5-fluoro-3-methyl-8-oxo-5,6,7,8-tet rahydroquinoline-5-carboxylate	 IM II-e-2-4

[0194] IM II-c-1

¹H NMR (CDCl₃) delta 8.79 (1 H, dd, J = 4.6, 1.3 Hz), 7.78 (1 H, br d, J = 7.9 Hz), 7.46 (1 H, dd, J = 7.9, 4.6 Hz), 4.05 (1 H, dd, J = 5.2, 4.6 Hz), 3.76 (3 H, s), 3.04 (1 H, m), 2.82 (1 H, m), 2.59 (1 H, m), 2.43 (1 H, m).

MS (ESI) m/z: 206.1 (M+H)⁺.

[0195] IM II-c-2

¹H NMR (CDCl₃) delta 8.91 (1 H, dd, J = 4.3, 1.2 Hz), 7.96 (1 H, dd, J = 7.9, 1.2 Hz), 7.57 (1 H, dd, J = 7.9, 4.3 Hz), 3.85 (3 H, s), 3.06-3.03 (2 H, m), 2.85 (1 H, m), 2.63 (1 H, m).

MS (ESI) m/z: 224.1 (M+H)⁺.

[0196] IM II-c-3

¹H NMR (CDCl₃) delta 8.73 (1 H, dd, J = 14.6, 2.4 Hz), 7.66 (1 H, dd, J = 7.9, 2.4 Hz), 3.86 (3 H, s), 3.06-2.99 (2 H, m), 2.85 (1 H, m), 2.62 (1 H, m).

MS (ESI) m/z: 242.0 (M+H)⁺.

[0197] IM II-c-4

^1H NMR (CDCl_3) delta 8.72 (1 H, d, $J = 1.4$ Hz), 7.73 (1 H, d, $J = 1.4$ Hz), 3.86 (3 H, s), 3.03-2.99 (2 H, m), 2.82 (1 H, m), 2.61 (1H, m), 2.48 (3 H, s).

MS (ESI) m/z : 238.2 ($\text{M}+\text{H}$) $^+$.

[0198] General Procedure: Scheme 2, Step 6

NaBH_4 (1.5 eq.) was added to a solution of substrate (1.0 eq.) in MeOH and the mixture was stirred at room temperature until complete reaction. The mixture was poured into water and extracted with EtOAc. The extract was dried over Na_2SO_4 and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography to afford the following intermediates (Table 6).

[0199] [Table 6]

Intermediates	Structure	Chemical Name	Substrate
II-d-1		methyl 8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxylate	
II-d-2		methyl 5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxylate	

[0200] IM II-d-1

^1H NMR (CDCl_3) delta 8.49 (1 H, d, $J = 4.6$ Hz), 7.61 (0.5 H, d, $J = 7.9$ Hz), 7.57 (0.5 H, d, $J = 7.9$ Hz), 7.20 (1 H, dd, $J = 7.9, 5.3$ Hz), 4.75 (0.5 H, dd, $J = 8.6, 5.3$ Hz), 4.67 (0.5 H, dd, $J = 9.3, 5.3$ Hz), 4.10 (1 H, br), 3.91 (0.5 H, dd, $J = 7.9, 6.6$ Hz), 3.82 (0.5 H, br), 3.75 (1.5 H, s), 3.72 (1.5 H, s), 2.44-1.99 (3.5 H, m), 1.81 (0.5 H, m).

MS (ESI) m/z : 208.2 ($\text{M}+\text{H}$) $^+$.

[0201] IM II-d-2

^1H NMR (CDCl_3) delta 8.64 (1 H, dd, $J = 4.6, 1.3$ Hz), 7.80 (0.5 H, d, $J = 8.6$ Hz), 7.69 (0.5 H, dd, $J = 7.9, 1.3$ Hz), 7.31 (1 H, dd, $J = 7.9, 4.6$ Hz), 4.80-4.71 (1 H, m), 4.31 (0.5 H, br), 3.84 (2 H, s), 3.80 (1.5 H, s), 2.75-2.63 (0.5 H, m), 2.53-2.00 (3.5 H, m).

MS (ESI) m/z : 226.2 ($\text{M}+\text{H}$) $^+$.

[0202] General Procedure: Scheme 3, Step 1

To a stirred solution of tert-butyl 2-(diethoxyphosphoryl)acetate (1.0 eq.) in THF was added NaH (60% oil dispersant, 1.0 eq.) at 0 °C. After being stirred at this temperature for 1 h, a solution of substrate (1.0 eq.) in THF was added to the mixture. The mixture was warmed to room temperature and stirred until complete reaction. The reaction mixture was poured into water and extracted with EtOAc. The extract was dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography to afford the following intermediates (Table 7).

[0203] [Table 7]

Intermediates	Structure	Chemical Name	Substrate
VII-1		methyl 8-(2-(tert-butoxy)-2-oxoethylidene)- 5,6,7,8-tetrahydroquinoline-5-car boxylate	
VII-2		methyl 8-(2-(tert-butoxy)-2-oxoethylidene)- 5-fluoro-5,6,7,8-tetrahydroquinoli ne-5-carboxylate	

[0204] IM VII-1; isomer A

¹H NMR (CDCl₃) delta 8.54 (1 H, dd, J = 4.6, 1.3 Hz), 7.54 (1 H, br d, J = 7.9 Hz), 7.23 (1 H, dd, J = 7.9, 4.6 Hz), 7.15 (1 H, s), 3.87 (1 H, t, J = 5.3 Hz), 3.72 (3 H, s), 3.39 (1 H, m), 3.19 (1 H, m), 2.29 (1 H, m), 2.05 (1 H, m), 1.51 (9 H, s).

MS (ESI) m/z: 304.2 (M+H)⁺.

[0205] IM VII-1; isomer B

¹H NMR (CDCl₃) delta 8.42 (1 H, dd, J = 4.6, 1.3 Hz), 7.54 (1 H, br d, J = 7.9 Hz), 7.15 (1 H, dd, J = 7.9, 4.6 Hz), 5.86 (1 H, s), 3.86 (1 H, t, J = 5.3 Hz), 3.70 (3 H, s), 2.80 (1 H, m), 2.56 (1 H, m), 2.32 (1 H, m), 2.13 (1 H, m), 1.56 (9 H, s).

MS (ESI) m/z: 304.2 (M+H)⁺.

[0206] IM VII-2

¹H NMR (CDCl₃) delta 8.65 (0.25 H, br d, J = 3.3 Hz), 8.54 (0.75 H, dd, J = 4.6, 2.0 Hz), 7.77 (0.25 H, br d, J = 7.9 Hz), 7.69 (0.75 H, br d, J = 7.3 Hz), 7.33 (0.25 H, dd, J = 7.9, 4.6 Hz), 7.24 (0.75 H, br d, J = 4.6 Hz), 5.98 (0.75 H, s), 5.30 (0.25 H, s), 3.81 (0.75 H, s), 3.80 (2.25 H, s), 3.42 (0.5 H, m), 2.78 (1.5 H, m), 2.30-2.10 (2 H, m), 1.58





(6.75 H, s), 1.52 (2.25 H, s).

MS (ESI) m/z : 322.1 (M+H)⁺.

[0207] General Procedure: Scheme 3, Step 2

10% Pd-C (0.2 eq.) was added to a solution of substrate (1.0 eq.) in MeOH and the mixture was stirred until complete reaction at room temperature under H₂ atmosphere. The reaction mixture was filtrated through a pad of cerite and the filtrate was concentrated in vacuo. The resulting residue was purified by silica gel column chromatography to afford the following intermediates (Table 8).

[0208] [Table 8]

Intermediates	Structure	Chemical Name	Substrate
II-v-1		methyl 8-(2-(tert-butoxy)-2-oxoethyl)-5,6, 7,8-tetrahydroquinoline-5-carboxyl ate	 IM VII-1
II-v-2		methyl 8-(2-(tert-butoxy)-2-oxoethyl)-5-flu oro-5,6,7,8-tetrahydroquinoline-5- carboxylate	 IM VII-2

[0209] IM II-v-1

¹H NMR (CDCl₃) delta 8.45 (1 H, dd, J = 4.6, 1.3 Hz), 7.50 (1 H, br d, J = 7.3 Hz), 7.09 (1 H, dd, J = 7.3, 4.6 Hz), 3.81 (1 H, t, J = 5.3 Hz), 3.72 (3 H, s), 3.34 (1 H, m), 3.13 (1 H, dd, J = 10.5, 4.6 Hz), 2.41 (1 H, dd, J = 15.8, 9.9 Hz), 2.31-1.72 (4 H, m), 1.46 (9 H, s).

MS (ESI) m/z : 306.1 (M+H)⁺.

[0210] IM II-v-2

¹H NMR (CDCl₃) delta 8.58 (0.5 H, d, J = 4.6 Hz), 8.49 (0.5 H, d, J = 4.6 Hz), 7.70 (0.5 H, d, J = 7.3 Hz), 7.53 (0.5 H, d, J = 7.9 Hz), 7.20 (0.5 H, dd, J = 7.9, 4.6 Hz), 7.13 (0.5 H, dd, J = 7.3, 4.6 Hz), 3.83 (1 H, t, J = 5.3 Hz), 3.74 (1.5 H, s), 3.72 (1.5 H, s), 3.55-3.30 (1 H, m), 3.15-2.94 (1 H, m), 2.57-2.41 (1 H, m), 2.4-1.7 (3 H, m), 1.47 (4.5 H, s), 1.45 (4.5 H, s).

MS (ESI) m/z : 324.1 (M+H)⁺.

[0211] Preparation of Intermediates III

General Procedure: Scheme 5, Step 1

To a solution of the substrate (1.0 eq.) in CCl_4 were added NBS (1.2 eq.) and AIBN (0.1 eq.) at 50 °C under N_2 atmosphere. The reaction mixture was heated at reflux and stirred for 2 h, another portion of AIBN (0.1 eq.) was added to the mixture. After being stirred at reflux for 16 h, the mixture was cooled to room temperature. The mixture was concentrated in vacuo and diethylether was added to the resulting residue. The insoluble materials were removed by filtration. The filtrate was washed with 2 N hydrochloric acid and brine. The extract was dried over Na_2SO_4 and concentrated in vacuo. The resulting residue was purified and separated by column chromatography to afford the following intermediates, di-bromide IX and mono-bromide X (Table 9).

[0212] [Table 9]

Intermediates	Structure	Chemical Name	Substrate
IX-1		2,4-dichloro-6-(dibromomethyl)benzonitrile	
X-1		2-(bromomethyl)-4,6-dichlorobenzonitrile	
IX-2		2-chloro-6-(dibromomethyl)-4-fluorobenzonitrile	
X-2		2-(bromomethyl)-6-chloro-4-fluorobenzonitrile	

[0213] IM IX1

^1H NMR (CDCl_3) δ 7.93 (1 H, d, $J = 2.0$ Hz), 7.52 (1 H, d, $J = 2.0$ Hz), 6.90 (1 H, s).

[0214] IM X-1

¹H NMR (CDCl₃) delta 7.49 (1 H, d, J = 2.0 Hz), 7.47 (1 H, d, J = 2.0 Hz), 4.57 (2 H, s).

[0215] IM IX-2

¹H NMR (CDCl₃) delta 7.69 (1 H, dd, J = 9.2, 2.4 Hz), 7.26 (1 H, dd, J = 9.2, 2.4 Hz), 6.93 (1 H, d, J = 1.2 Hz).

[0216] IM X-2


¹H NMR (CDCl₃) delta 7.23 (1 H, dd, J = 7.9, 2.4 Hz), 7.22 (1 H, dd, J = 7.9, 2.4 Hz), 4.59 (2 H, s).

MS (ESI) m/z: 250.9 (M+H)⁺.

[0217] Procedure: Scheme 5, Step 2

Intermediate (IM) XI-1, 2,4-dichloro-6-(difluoromethyl)benzonitrile

[Chem.25]




To a solution of 2,4-dichloro-6-(difluoromethyl)benzonitrile (350 mg, 1.018 mmol, IM IX-1) in CH₂Cl₂ (5 mL) was added silver tetrafluoroborate (495 mg, 2.54 mmol) at ambient temperature under N₂ atmosphere. After being stirred at room temperature for 3 h, the insoluble material was removed by filtration. The filtrate was concentrated in vacuo to afford 204 mg (90%) of the title compound.

¹H NMR (CDCl₃) delta 7.68 (2 H, br s), 6.89 (1 H, dd, J = 54.7, 54.0 Hz).

[0218] Procedure: Scheme 5, Step 4

Intermediate (IM) XII-1, 3-chloro-2-cyano-5-fluorobenzyl acetate

[Chem.26]







To a solution of the 2-(bromomethyl)-4,6-dichlorobenzonitrile (609 mg, 2.451 mmol, IM X-2) in AcOH (6.0 mL) was added NaOAc (1.0 g, 12.25 mmol) at ambient temperature. The mixture was heated at 100 °C for 6 h. The mixture was concentrated in vacuo and aq. NaHCO₃ was added to the resulting residue. The mixture was extracted with EtOAc and washed with brine. The extracts were dried over Na₂SO₄

and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography (5% EtOAc/n-hexane) to afford 505 mg (91%) of the title compound. MS (ESI) m/z: 245.0 (M+H₃O)⁺.

[0219] General Procedure: Scheme 5, Step 3

To a solution of the substrate (1.0 eq.) in AcOH was added NaOAc (5.0 eq.) at ambient temperature. The resulting mixture was heated at 100 °C until complete reaction. The mixture was concentrated in vacuo and aq. NaHCO₃ was added to the resulting residue. The mixture was extracted with EtOAc and washed with brine. The extracts were dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography to afford the following intermediates (Table 10).

[0220] [Table 10]

Intermediates	Structure	Chemical Name	Substrate
III-a-1		(2,4-dichloro-6-(difluoromethyl)phenyl)methanamine	 IM XI-1
III-b-1		(2-(aminomethyl)-3-chloro-5-fluorophenyl)methanol	 IM XII-1

[0221] IM III-a-1

¹H NMR (CDCl₃) delta 7.53 (1 H, br s), 7.48 (1 H, br s), 6.91 (1 H, dd, J = 82.0, 81.0 Hz), 4.02 (2 H, s), 1.48 (2 H, s).

MS (ESI) m/z: 226.0 (M+H)⁺.

[0222] IM III-b-1

¹H NMR (CDCl₃) delta 7.09 (1 H, dd, J = 8.6, 2.4 Hz), 7.03 (1 H, dd, J = 8.6, 2.4 Hz), 4.64 (2 H, s), 4.16 (2 H, s), 2.35 (3 H, br).

MS (ESI) m/z: 190.1 (M+H)⁺.

[0223] The following Examples and Intermediates were prepared by General Procedure A (Tables 11 and 13).

[0224] General Procedure A
















A mixture of substrate (1.0 eq.) and 2 N aq. NaOH (2.0 eq.) in MeOH was stirred at room temperature for 1.5 h, 2 N hydrochloric acid (2.2 eq.) was added to the mixture. The mixture was concentrated in vacuo to afford a glass. Toluene and MeCN were added to the mixture and concentrated in vacuo. This procedure was repeated 3 times to remove remaining water. The residual powder was dissolved with DMF and amine (1.5 eq.), triethylamine (3.0 eq.), and HBTU (1.3 eq.) were added to the mixture at ambient temperature. After overnight stirring, the mixture was poured into water and the mixture was extracted with CH₂Cl₂. The extract was concentrated in vacuo and the resulting residue was purified by silica gel column chromatography, and/or SCX cartridge column, preparative HPLC to afford the following Examples and Intermediates.

[0225]

[Table 11-1]

Examples	Structure	Chemical Name	Substrate	Amine
1		N-(2,4-dichloro-6-methylbenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		
2		N-(2-chloro-3-(trifluoromethyl)benzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		
3		N-(2,3-dichlorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		
4		N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		
5		N-(cycloheptylmethyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		
6		N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 11-2]

7		N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-d-1	
8		N-(2,3-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-d-1	
9		N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-d-1	
10		N-(2,4-dichloro-6-(methoxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-d-1	
11		N-(2,4-dichloro-6-methylbenzyl)-7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide	 IM II-e-1-2	

[0226] [Table 12]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
1	D	1.53	363.0	7	D	1.46	385.0
2	D	1.47	383.0	8	D	1.40	351.0
3	D	1.40	349.0	9	A	1.26	381.3
4	A	1.25	379.3	10	A	1.44	395.1
5	D	1.45	301.2	11	D	1.79	347.0
6	D	1.53	365.0				

[0227] Ex 2

¹H NMR (DMSO d6) delta 9.00 (1 H, br), 8.67 (1 H, m), 7.79 (2 H, br d, J = 7.9 Hz), 7.68 (1 H, d, J = 7.3 Hz), 7.68-7.54 (2 H, m), 4.48 (2 H, m), 4.12 (1 H, dd, J = 5.9, 5.2 Hz), 2.84 (1 H, m), 2.66 (1 H, m), 2.38-2.32 (2 H, m).

[0228] Ex 6

¹H NMR (DMSO d6) delta 8.42 (1 H, d, J = 4.6 Hz), 8.37 (1 H, br), 7.50 (1 H, s), 7.40-7.36 (2 H, m), 7.23 (1 H, dd, J = 7.9, 4.6 Hz), 5.18 (0.7 H, d, J = 4.0 Hz), 5.13 (0.3 H, d, J = 4.0 Hz), 4.51 (1 H, br), 4.44 (0.7 H, s), 4.42 (0.7 H, s), 4.39 (0.3 H, s), 4.37 (0.3 H, s), 3.74 (0.3 H, br), 3.65 (0.7 H, br), 2.40 (2.1 H, s), 2.37 (0.9 H, s), 2.20-1.90 (2 H, m), 1.82-1.76 (2 H, m).

[0229] Ex 7



















¹H NMR (DMSO d6) delta 8.79 (1 H, br), 8.43 (1 H, d, J = 4.6 Hz), 7.80 (1 H, d, J = 7.9 Hz), 7.68 (1 H, dd, J = 8.6, 6.6 Hz), 7.58 (1 H, dd, J = 7.9, 7.3 Hz), 7.46 (1 H, d, J = 7.3 Hz), 7.23 (1H, dd, J = 8.6, 5.3 Hz), 5.18 (1 H, br), 4.54 (1 H, br), 4.47 (2 H, m), 3.75 (1 H, m), 2.22-1.80 (4 H, m).

[0230]



















[Table 13-1]

Intermediates	Structure	Chemical Name	Substrate	Amine
I-e-1		N-(2,4-dichloro-6-methylbenzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-2		N-(2-chloro-3-(trifluoromethyl)benzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-3		N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-4		N-(2,4-dichloro-6-methylbenzyl)-5-hydroxy-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-5		N-(2-chloro-3-(trifluoromethyl)benzyl)-5-hydroxy-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-6		N-(2,4-dichlorobenzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 13-2]

I-e-7		N-(2-chloro-4-fluorobenzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-e-1-1	
I-e-8		N-(2,4-dichloro-6-fluorobenzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-e-1-1	
I-e-9		8-methylene-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-e-1-1	
I-e-10		8-methylene-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-e-1-1	
I-e-11		N-(2,4-difluorobenzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-e-1-1	
I-e-12		N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-e-1-1	










[Table 13-3]

I-e-13		N-(4-chloro-2-fluorobenzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-14		N-(4-bromo-2-fluorobenzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-15		N-(2-chloro-3,4-difluorobenzyl)-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-16		N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-17		N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-18		N-(2,3-dichlorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 13-4]

I-e-19		N-(2,4-dichlorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-20		N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-21		N-(2,4-dichloro-6-(methoxymethyl)benzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-22		N-(2-chloro-3-(trifluoromethyl)benzyl)-7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		
I-e-23		N-(2,3-dichlorobenzyl)-7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		
I-e-24		N-(2,4-dichlorobenzyl)-5-fluoro-7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		

[Table 13-5]

I-v-1		tert-butyl 2-(5-((2,4-dichloro-6-methylbenzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetate		
I-v-2		tert-butyl 2-(5-((2-chloro-3-(trifluoromethyl)benzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetate		
I-v-3		tert-butyl 2-(5-((2,4-dichloro-6-methylbenzyl)carbamoyl)-5-fluoro-5,6,7,8-tetrahydroquinolin-8-yl)acetate		

[0231] IM I-e-1

¹H NMR (CDCl₃) delta 8.54 (1 H, dd, J = 4.6, 2.0 Hz), 7.40 (1 H, d, J = 9.2 Hz), 7.21 (1 H, d, J = 2.0 Hz), 7.15 (1 H, dd, J = 9.2, 4.6 Hz), 7.09 (1 H, d, J = 2.0 Hz), 6.31 (1 H, s), 5.66 (1 H, br), 5.19 (1 H, s), 4.56 (1 H, dd, J = 14.5, 5.9 Hz), 4.44 (1 H, dd, J = 14.5, 5.9 Hz), 3.71 (1 H, br t, J = 5.3 Hz), 2.63 (2 H, m), 2.45 (3 H, s), 2.35 (1 H, m), 2.04 (1 H, m).

MS (ESI) m/z: 361.7 (M+H)⁺.

[0232] IM I-e-2

¹H NMR (CDCl₃) delta 8.56 (1 H, d, J = 4.6 Hz), 7.64 (1 H, d, J = 7.3 Hz), 7.55 (1 H, d, J = 7.3 Hz), 7.42 (1 H, d, J = 8.6 Hz), 7.35 (1 H, br t, J = 7.3 Hz), 7.17 (1 H, dd, J = 7.3, 4.6 Hz), 6.34 (1 H, s), 5.86 (1 H, br), 5.22 (1 H, s), 4.59 (1 H, dd, J = 14.5, 5.9 Hz), 4.51 (1 H, dd, J = 14.5, 5.9 Hz), 3.77 (1 H, br t, J = 5.3 Hz), 2.66 (2 H, m), 2.18 (1 H, m), 2.06 (1 H, m).

MS (ESI) m/z: 381.7 (M+H)⁺.

[0233] IM I-e-3

¹H NMR (CDCl₃) delta 8.60 (1 H, br dd, J = 4.6, 1.3 Hz), 7.70 (1 H, d, J = 7.9 Hz), 7.65 (1 H, d, J = 7.3 Hz), 7.47 (1 H, br dd, J = 7.9, 1.3 Hz), 7.40 (1 H, dd, J = 7.9, 7.3 Hz), 7.30 (1 H, br), 7.18 (1 H, dd, J = 7.9, 4.6 Hz), 6.35 (1 H, br s), 5.31 (1 H, d, J = 2.6 Hz), 4.73 (2 H, d, J = 5.9 Hz), 2.88-2.83 (2 H, m), 2.60-2.20 (2 H, m).

MS (ESI) m/z: 399.0 (M+H)⁺.

[0234] IM I-e-4

¹H NMR (DMSO d6) delta 8.48 (1 H, dd, J = 4.6, 2.0 Hz), 8.12 (1 H, br t, J = 5.3 Hz), 7.53 (1 H, dd, J = 7.9, 2.0 Hz), 7.47 (1 H, d, J = 2.0 Hz), 7.32 (1 H, d, J = 2.0 Hz), 7.26 (1 H, dd, J = 7.9, 4.6 Hz), 6.26 (1 H, s), 6.18 (1 H, s), 5.15 (1 H, br), 4.44 (2 H, d, J = 5.3 Hz), 2.74 (2 H, m), 2.40 (3 H, s), 2.15 (1 H, m), 1.86 (1 H, m).

MS (ESI) m/z: 377.3 (M+H)⁺.

[0235] IM I-c-5

¹H NMR (DMSO d6) delta 8.90 (1 H, br t, J = 6.6 Hz), 8.49 (1 H, dd, J = 4.6, 2.0 Hz), 7.78 (1 H, br d, J = 7.9 Hz), 7.67-7.54 (3 H, m), 7.29 (1 H, dd, J = 7.9, 4.6 Hz), 6.43 (1 H, s), 6.18 (1 H, br s), 5.17 (1 H, br s), 4.47 (2 H, d, J = 6.6 Hz), 2.78 (2 H, m), 2.23 (1 H, m), 1.96 (1 H, m).

MS (ESI) m/z: 397.3 (M+H)⁺.

[0236] IM I-c-6

¹H NMR (CDCl₃) delta 8.55 (1 H, dd, J = 4.9, 1.8 Hz), 7.43 (1 H, dd, J = 7.9, 1.8 Hz), 7.36 (1 H, d, J = 2.4 Hz), 7.30-7.26 (1 H, m), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 7.17 (1 H, dd, J = 8.0, 4.9 Hz), 6.33 (1 H, d, J = 1.8 Hz), 5.77 (1 H, br), 5.21 (1 H, d, J = 1.8 Hz), 4.47 (1 H, dd, J = 15.3, 6.1 Hz), 4.41 (1 H, dd, J = 15.3, 6.1 Hz), 3.75 (1 H, t, J = 4.9 Hz), 3.75 (2 H, br t, J = 4.9 Hz), 2.34 (1 H, m), 2.05 (1 H, m).

MS (ESI) m/z: 346.7 (M+H)⁺.

[0237] IM I-c-7

¹H NMR (CDCl₃) delta 8.55 (1 H, dd, J = 4.9, 1.2 Hz), 7.43 (1 H, dd, J = 7.9, 1.2 Hz), 7.32 (1 H, dd, J = 8.6, 6.1 Hz), 7.16 (1 H, dd, J = 7.9, 4.9 Hz), 7.09 (1 H, dd, J = 8.6, 2.4 Hz), 6.94 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 6.33 (1 H, d, J = 1.8 Hz), 5.76 (1 H, br), 5.21 (1 H, d, J = 1.8 Hz), 4.47 (1 H, dd, J = 15.3, 6.1 Hz), 4.41 (1 H, dd, J = 15.3, 6.1 Hz), 3.75 (1 H, t, J = 4.9 Hz), 2.66-2.62 (2 H, m), 2.34 (1 H, m), 2.05 (1 H, m).

MS (ESI) m/z: 330.8 (M+H)⁺.

[0238] IM I-c-8

¹H NMR (CDCl₃) delta 8.54 (1 H, dd, J = 4.9, 1.8 Hz), 7.42 (1 H, dd, J = 7.9, 1.8 Hz), 7.20 (1 H, m), 7.16 (1 H, dd, J = 7.9, 4.9 Hz), 7.03 (1 H, dd, J = 9.2, 1.8 Hz), 6.32 (1 H, d, J = 1.8 Hz), 5.64 (1 H, br), 5.20 (1 H, d, J = 1.8 Hz), 4.58 (1 H, ddd, J = 14.7, 5.5, 1.2 Hz), 4.49 (1 H, dd, J = 14.7, 5.5, 1.8 Hz), 3.74 (1 H, m), 2.65-2.61 (2 H, m), 2.31 (1 H, m), 2.03 (1 H, m).

MS (ESI) m/z: 364.7 (M+H)⁺.

[0239] IM I-c-9

¹H NMR (CDCl₃) delta 8.46 (1 H, dd, J = 4.8, 1.2 Hz), 7.39 (1 H, dd, J = 7.9, 1.2 Hz), 7.13 (1 H, dd, J = 7.9, 4.8 Hz), 6.99 (1 H, m), 6.89 (1 H, m), 6.27 (1 H, d, J = 1.8 Hz), 6.13 (1 H, br t, J = 5.5 Hz), 5.18 (1 H, d, J = 1.8 Hz), 4.41 (2 H, br d, J = 6.1 Hz), 3.73 (1 H, br t, J = 5.5 Hz), 2.65-2.61 (2 H, m), 2.26 (1 H, m), 2.03 (1 H, m).

MS (ESI) m/z: 333.2 (M+H)⁺.

[0240] IM I-c-10

¹H NMR (CDCl₃) delta 8.53 (1 H, dd, J = 4.3, 1.2 Hz), 7.41 (1 H, dd, J = 7.9, 1.2 Hz), 7.16 (1 H, dd, J = 7.9, 4.3 Hz), 6.64 (2 H, m), 6.31 (1 H, d, J = 1.8 Hz), 5.56 (1 H, br), 5.19 (1 H, d, J = 1.8 Hz), 4.49 (1 H, dd, J = 14.7, 6.1 Hz), 4.43 (1 H, dd, J = 14.7, 5.5 Hz), 3.73 (1 H, br t, J = 5.5 Hz), 2.65-2.61 (2 H, m), 2.31 (1 H, m), 2.04 (1 H, m).

MS (ESI) m/z: 333.1 (M+H)⁺.

[0241] IM I-c-11

¹H NMR (CDCl₃) delta 8.54 (1 H, dd, J = 4.9, 1.2 Hz), 7.42 (1 H, dd, J = 7.9, 1.2 Hz), 7.29-7.22 (1 H, m), 7.16 (1 H, dd, J = 7.9, 4.9 Hz), 6.85-6.74 (2 H, m), 6.32 (1 H, d, J = 1.8 Hz), 5.66 (1 H, br), 5.20 (1 H, d, J = 1.8 Hz), 4.42 (2 H, br d, J = 5.5 Hz), 3.75 (1 H, br t, J = 5.5 Hz), 2.66-2.63 (2 H, m), 2.33 (1 H, m), 2.06 (1 H, m).

MS (ESI) m/z: 315.2 (M+H)⁺.

[0242] IM I-c-12

¹H NMR (CDCl₃) delta 8.48 (1 H, dd, J = 4.2, 1.2 Hz), 7.45 (1 H, dd, J = 8.6, 5.5 Hz), 7.38 (1 H, dd, J = 7.9, 1.2 Hz), 7.30 (1 H, dd, J = 8.6, 2.4 Hz), 7.17 (1 H, dt, J = 8.6, 2.4 Hz), 7.12 (1 H, dd, J = 7.9, 4.2 Hz), 6.27 (1 H, d, J = 1.2 Hz), 6.07 (1 H, br t, J = 6.1 Hz), 5.16 (1 H, d, J = 1.2 Hz), 4.51 (2 H, br d, J = 6.1 Hz), 3.72 (1 H, br t, J = 5.5 Hz), 2.72-2.60 (2 H, m), 2.27 (1 H, m), 2.02 (1 H, m).

MS (ESI) m/z: 365.1 (M+H)⁺.

[0243] IM I-c-13

¹H NMR (CDCl₃) delta 8.44 (1 H, dd, J = 4.9, 1.8 Hz), 7.39 (1 H, dd, J = 7.9, 1.8 Hz), 7.18 (1 H, t, J = 7.9 Hz), 7.11 (1 H, dd, J = 7.9, 4.9 Hz), 7.08-7.00 (2 H, m), 6.25 (1 H, d, J = 1.2 Hz), 6.17 (1 H, br t, J = 5.5 Hz), 5.16 (1 H, d, J = 1.2 Hz), 4.38 (2 H, br d, J = 6.1 Hz), 3.71 (1 H, br t, J = 5.5 Hz), 2.69-2.55 (2 H, m), 2.24 (1 H, m), 2.02 (1 H, m).

MS (ESI) m/z: 331.1 (M+H)⁺.

[0244] IM I-c-14

¹H NMR (CDCl₃) delta 8.46 (1 H, dd, J = 4.9, 1.8 Hz), 7.38 (1 H, dd, J = 7.9, 1.8 Hz), 7.22-7.10 (4 H, m), 6.26 (1 H, d, J = 1.2 Hz), 6.16 (1 H, br t, J = 5.5 Hz), 5.17 (1 H, d, J = 1.2 Hz), 4.37 (2 H, br d, J = 5.5 Hz), 3.71 (1 H, br t, J = 5.5 Hz), 2.69-2.55 (2 H, m), 2.24 (1 H, m), 2.02 (1 H, m).

MS (ESI) m/z: 375.0 (M+H)⁺.

[0245] IM I-c-15

¹H NMR (CDCl₃) delta 8.52 (1 H, dd, J = 4.9, 1.8 Hz), 7.42 (1 H, dd, J = 7.9, 1.8 Hz), 7.16 (1 H, dd, J = 7.9, 4.9 Hz), 7.09-7.01 (2 H, m), 6.31 (1 H, d, J = 1.8 Hz), 5.92 (1 H, br t, J = 5.5 Hz), 5.20 (1 H, d, J = 1.8 Hz), 4.47 (1 H, dd, J = 15.3, 6.1 Hz), 4.41 (1 H, dd, J = 15.5, 6.1 Hz), 3.75 (1 H, br t, J = 5.5 Hz), 2.64 (2 H, m), 2.31 (1 H, m), 2.05 (1 H, m).

MS (ESI) m/z: 349.0 (M+H)⁺.

[0246] IM I-e-16

¹H NMR (CDCl₃) delta 8.61 (1 H, dd, J = 4.6, 2.0 Hz), 7.47 (1 H, ddd, J = 7.9, 2.0, 1.3 Hz), 7.32 (1H, d, J = 1.3 Hz), 7.19 (1 H, dd, J = 7.9, 4.6 Hz), 7.16 (1H, d, J = 1.3 Hz), 7.04 (1 H, br d, J = 5.3 Hz), 6.34 (1 H, s), 5.31 (1 H, s), 4.72 (1 H, dd, J = 13.8, 5.3 Hz), 4.63 (1 H, dd, J = 13.8, 5.3 Hz), 2.84 (2 H, m), 2.45 (1 H, m), 2.48 (3 H, s), 2.25 (1 H, m).

MS (ESI) m/z: 379.0 (M+H)⁺.

[0247] IM I-e-17

¹H NMR (CDCl₃) delta 8.58 (1 H, dd, J = 4.6, 2.0 Hz), 7.62 (1 H, br d, J = 5.3 Hz), 7.44 (1H, d, J = 2.0 Hz), 7.40 (1 H, ddd, J = 7.9, 2.0, 1.3 Hz), 7.34 (1H, d, J = 2.0 Hz), 7.16 (1 H, dd, J = 7.9, 4.6 Hz), 6.33 (1 H, s), 5.30 (1 H, s), 4.81-4.65 (4 H, m), 3.96 (1 H, br), 2.80 (2 H, m), 2.86-2.14 (2 H, m).

MS (ESI) m/z: 394.9 (M+H)⁺.

[0248] IM I-e-18

¹H NMR (CDCl₃) delta 8.61 (1 H, br dd, J = 4.6, 1.3 Hz), 7.51-7.45 (2 H, m), 7.35 (1 H, dd, J = 7.9, 1.3 Hz), 7.26-7.20 (2 H, m), 7.19 (1 H, dd, J = 7.9, 4.6 Hz), 6.35 (1 H, s), 5.31 (1 H, s), 4.69 (1 H, d, J = 5.9 Hz), 2.88-2.83 (2 H, m), 2.61-2.20 (2 H, m).

MS (ESI) m/z: 364.9 (M+H)⁺.

[0249] IM I-e-19

¹H NMR (CDCl₃) delta 8.61 (1 H, br dd, J = 4.9, 1.8 Hz), 7.47 (1 H, br dd, J = 7.9, 1.8 Hz), 7.45 (1 H, d, J = 1.8 Hz), 7.37 (1 H, d, J = 7.9 Hz), 7.26 (1 H, dd, J = 7.9, 1.8 Hz), 7.19 (1 H, br), 7.18 (1 H, dd, J = 7.9, 4.9 Hz), 6.35 (1 H, s), 5.31 (1 H, s), 4.65 (1 H, dd, J = 14.7, 5.5 Hz), 4.60 (1 H, dd, J = 14.7, 5.5 Hz), 2.86-2.83 (2 H, m), 2.46 (1 H, m), 2.26 (1 H, m).

MS (ESI) m/z: 365.0 (M+H)⁺.

[0250] IM I-e-20

¹H NMR (CDCl₃) delta 8.61 (1 H, ddd, J = 4.9, 1.8, 1.8 Hz), 7.47 (1 H, ddd, J = 7.9, 1.8, 1.8 Hz), 7.42 (1 H, dd, J = 8.6, 6.1 Hz), 7.24-7.13 (3 H, m), 7.00 (1 H, ddd, J = 8.6, 3.1, 3.1 Hz), 6.35 (1 H, s), 5.31 (1 H, s), 4.65 (1 H, dd, J = 15.3, 6.1 Hz), 4.60 (1 H, dd, J = 15.3, 6.1 Hz), 2.90-2.79 (2 H, m), 2.48 (1 H, m), 2.25 (1 H, m).

MS (ESI) m/z: 348.9 (M+H)⁺.

[0251] IM I-e-21

¹H NMR (CDCl₃) delta 8.60 (1 H, dd, J = 4.6, 2.0 Hz), 7.48 (1 H, dd, J = 7.3, 2.0 Hz), 7.45 (1H, d, J = 2.0 Hz), 7.33 (1H, d, J = 2.0 Hz), 7.29 (1 H, m), 7.18 (1 H, dd, J = 7.3, 4.6 Hz), 6.34 (1 H, s), 5.30 (1 H, s), 4.79 (1 H, dd, J = 14.5, 6.6 Hz), 4.60 (1 H, m), 4.58 (1 H, d, J = 11.2 Hz), 4.55 (1 H, d, J = 11.2 Hz), 3.44 (3 H, s), 2.83 (2 H, m), 2.59-2.15 (2 H, m).

MS (ESI) m/z: 408.9 (M+H)⁺.

[0252] IM I-e-22

MS (ESI) m/z: 367.0 (M+H)⁺.

[0253] IM I-e-23

MS (ESI) m/z: 333.1 (M+H)⁺.

[0254] IM I-e-24

¹H NMR (CDCl₃) delta 8.68 (1 H, ddd, J = 4.9, 4.0, 1.8 Hz), 7.61 (1 H, br dt, J = 7.9, 1.8 Hz), 7.45 (1 H, d, J = 1.8 Hz), 7.40-7.35 (1 H, m), 7.29-7.15 (3 H, m), 6.19 (1 H, br d, J = 1.8 Hz), 5.35 (1 H, br d, J = 1.8 Hz), 4.67 (1 H, dd, J = 14.7, 6.1 Hz), 4.59 (1 H, dd, J = 14.7, 6.1 Hz), 3.58 (1 H, m), 3.12 (1 H, m).

MS (ESI) m/z: 351.1 (M+H)⁺.

[0255] IM I-v-1

¹H NMR (CDCl₃) delta 8.44 (1 H, m), 7.36 (1 H, m), 7.28-7.02 (3 H, m), 6.34 (0.5 H, br), 5.73 (0.5 H, br), 4.59-4.39 (2 H, m), 3.75 (0.5 H, br), 3.66 (0.5 H, br t, J = 6.6 Hz), 3.40 (0.5 H, m), 3.15 (0.5 H, m), 3.10-2.91 (1.5 H, m), 2.73-2.65 (0.5 H, m), 2.50-2.20 (2 H, m), 2.55 (1.5 H, s), 2.37 (1.5 H, s), 2.07-1.95 (2 H, m), 1.44 (4.5 H, s), 1.26 (4.5H, s).

MS (ESI) m/z: 463.1 (M+H)⁺.

[0256] IM I-v-2

¹H NMR (CDCl₃) delta 8.47 (1 H, m), 7.62 (0.5 H, d, J = 8.6 Hz), 7.58 (0.5 H, d, J = 7.9 Hz), 7.46-7.24 (3 H, m), 7.09 (1 H, m), 6.88 (0.5 H, br), 5.90 (0.5 H, br), 4.68 (0.5 H, dd, J = 15.2, 6.6 Hz), 4.59-4.51 (1 H, m), 4.39 (0.5 H, dd, J = 15.8, 5.2 Hz), 3.75 (0.5 H, br d, J = 2.6 Hz), 3.72 (0.5 H, br t, J = 7.3 Hz), 3.40 (0.5 H, br), 3.28 (0.5 H, dd, J = 16.4, 6.0 Hz), 3.19 (0.5 H, br), 2.94 (0.5 H, m), 2.68 (0.5 H, dd, J = 16.4, 2.6 Hz), 2.46 (0.5 H, dd, J = 15.8, 9.2 Hz), 2.37-2.30 (0.5 H, m), 2.16-1.97 (3 H, m), 1.80-1.67 (0.5 H, m), 1.44 (4.5 H, s), 1.28 (4.5 H, s).

MS (ESI) m/z: 483.0 (M+H)⁺.

[0257] IM I-v-3

¹H NMR (CDCl₃) delta 8.55 (1 H, m), 7.47 (1 H, d, J = 7.9 Hz), 7.30 (1 H, d, J = 2.0 Hz), 7.14 (2 H, m), 7.02 (1 H, br), 4.69 (1 H, dd, J = 13.8, 5.3 Hz), 4.61 (1 H, dd, J = 14.5, 5.3 Hz), 3.45 (1 H, br), 2.92 (1 H, dd, J = 16.5, 4.6 Hz), 2.61-2.48 (1.5 H, m), 2.45 (3 H, s), 2.26-2.05 (2.5 H, m), 1.45 (9 H, s).

MS (ESI) m/z: 481.0 (M+H)⁺.

[0258] The following Examples and Intermediates were prepared by General Procedure B (Tables 14 and 16).

[0259] General Procedure B

The General Procedure B was carried out by the same procedure described in General Procedure: Scheme 2, Step 5 with/without following purification; SCX

cartridge column, preparative HPLC to afford the following Examples and Intermediates.

[0260] [Table 14]

Examples	Structure	Chemical Name	Substrate
12		N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-16
13		N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-3
14		N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-17
15		N-(2,4-dichloro-6-methylbenzyl)-5-hydroxy-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-4
16		N-(2,4-dichloro-6-methylbenzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide	 Ex 11
17		N-(2-chloro-3-(trifluoromethyl)benzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide	 IM I-e-22

[0261]

[Table 15]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
12	D	1.68	381.0	15	A	1.47	379.3
13	D	1.59	401.0	16	D	1.55	349.0
14	A	1.35	397.2	17	D	1.48	369.0

[0262] Ex 12

¹H NMR (CDCl₃) delta 8.87 (1 H, d, J = 4.6 Hz), 7.73 (1 H, d, J = 7.9 Hz), 7.52 (1 H, dd, J = 7.9, 4.6 Hz), 7.33 (1 H, d, J = 1.3 Hz), 7.16 (1 H, d, J = 1.3 Hz), 7.05 (1 H, br d, J = 5.9 Hz), 4.71 (1 H, dd, J = 14.5, 5.9 Hz), 4.64 (1 H, dd, J = 14.5, 5.9 Hz), 3.17 (1 H, m), 3.00 (1 H, m), 2.79 (1 H, m), 2.52 (1 H, m), 2.46 (3 H, s).

[0263] Ex 13

¹H NMR (CDCl₃) delta 8.88 (1 H, dd, J = 4.6, 1.3 Hz), 7.74 (1 H, d, J = 7.9 Hz), 7.71 (1 H, d, J = 7.9 Hz), 7.63 (1 H, d, J = 7.9 Hz), 7.51 (1 H, dd, J = 7.9, 4.6 Hz), 7.40 (1 H, dd, J = 7.9, 7.9 Hz), 7.33 (1 H, br d, J = 5.9 Hz), 4.72 (2 H, d, J = 5.9 Hz), 3.17 (1 H, m), 3.01 (1 H, m), 2.59 (1 H, m).

[0264] Ex 14

¹H NMR (CDCl₃) delta 8.88 (1 H, br dd, J = 4.6, 1.3 Hz), 7.69 (1 H, d, J = 7.9 Hz), 7.59 (1 H, br), 7.50 (1 H, dd, J = 7.9, 4.6 Hz), 7.45 (1 H, d, J = 2.0 Hz), 7.34 (1 H, d, J = 2.0 Hz), 4.85-4.64 (4 H, m), 3.60 (1 H, br), 3.18-2.92 (2 H, m), 2.76 (1 H, m), 2.54 (1 H, m).

[0265] Ex 15











¹H NMR (DMSO d₆) delta 8.70 (1 H, br dd, J = 4.6, 2.0 Hz), 8.23 (1 H, br), 7.86 (1 H, dd, J = 7.9, 2.0 Hz), 7.60 (1 H, dd, J = 7.9, 4.6 Hz), 7.46 (1 H, d, J = 2.0 Hz), 7.30 (1 H, d, J = 2.0 Hz), 6.64 (1 H, s), 4.42 (2 H, m), 2.89-2.69 (2 H, m), 2.43 (1 H, m), 2.37 (3 H, s), 2.22 (1 H, m).

[0266] Ex 16















¹H NMR (DMSO d₆) delta 8.82 (1 H, br), 8.77 (1 H, d, J = 4.6 Hz), 8.00 (1 H, d, J = 7.9 Hz), 7.63 (1 H, dd, J = 7.9, 4.6 Hz), 7.50 (1 H, br s), 7.36 (1 H, br s), 4.39 (2 H, br d J = 4.6 Hz), 4.29 (1 H, m), 2.80 (2 H, m), 2.37 (3 H, s).

[0267]

[Table 16-1]

Intermediates	Structure	Chemical Name	Substrate
I-c-1		N-(2,4-dichlorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-6
I-c-2		N-(2-chloro-4-fluorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-7
I-c-3		N-(2,4-dichloro-6-fluorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-8
I-c-4		8-oxo-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-9
I-c-5		8-oxo-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-10

[Table 16-2]

I-c-6		N-(2,4-difluorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-11
I-c-7		N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-12
I-c-8		N-(4-chloro-2-fluorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-13
I-c-9		N-(4-bromo-2-fluorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-14
I-c-10		N-(2-chloro-3,4-difluorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-15
I-c-11		N-(2-chloro-3-(trifluoromethyl)benzyl)-5-hydroxy-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-5
I-c-12		N-(2,3-dichlorobenzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide	 IM I-e-23

[0268] IM I-c-1

¹H NMR (CDCl₃) delta 8.60 (1 H, dd, J = 4.3, 1.8 Hz), 7.67 (1 H, dd, J = 7.9, 1.8 Hz), 7.40 (1 H, dd, J = 7.9, 4.9 Hz), 7.34 (1 H, d, J = 1.8 Hz), 7.29 (1 H, d, J = 7.9 Hz), 7.19 (1 H, dd, J = 7.9, 1.8 Hz), 6.94 (1 H, t, J = 5.5 Hz), 4.53 (1 H, dd, J = 14.6, 5.5 Hz), 4.47 (1 H, dd, J = 14.6, 5.5 Hz), 3.96 (1 H, t, J = 5.5 Hz), 2.96 (1 H, ddd, J = 17.7, 10.4, 4.9 Hz), 2.70 (1 H, ddd, J = 17.7, 7.3, 4.9 Hz), 2.53 (1 H, m), 2.40 (1 H, m).

MS (ESI) m/z: 349.0 (M+H)⁺.

[0269] IM I-c-2

¹H NMR (CDCl₃) delta 8.53 (1 H, dd, J = 4.3, 1.2 Hz), 7.69 (1 H, dd, J = 7.9, 1.2 Hz), 7.55 (1 H, br), 7.37 (1 H, dd, J = 7.9, 4.3 Hz), 7.30 (1 H, dd, J = 8.5, 6.1 Hz), 7.06 (1 H, dd, J = 8.5, 2.4 Hz), 6.89 (1 H, dt, J = 8.5, 2.4 Hz), 4.48 (1 H, dd, J = 9.8, 5.5 Hz), 4.47 (1 H, dd, J = 9.8, 5.5 Hz), 4.03 (1 H, br t, J = 5.5 Hz), 2.99 (1 H, ddd, J = 17.7, 9.8, 4.9 Hz), 2.67 (1 H, m), 2.48 (1 H, m), 2.36 (1 H, m).

MS (ESI) m/z: 333.1 (M+H)⁺.

[0270] IM I-c-3

¹H NMR (CDCl₃) delta 8.52 (1 H, m), 7.69 (1 H, dd, J = 7.9, 1.2 Hz), 7.38 (1 H, dd, J = 7.9, 4.3 Hz), 7.26 (1 H, br), 7.16 (1 H, d, J = 1.8 Hz), 6.97 (1 H, dd, J = 8.6, 1.8 Hz), 4.59 (1 H, dd, J = 15.9, 4.9 Hz), 4.55 (1 H, dd, J = 15.9, 4.9 Hz), 3.98 (1 H, br t, J = 5.5 Hz), 3.19 (1 H, ddd, J = 17.7, 10.4, 4.9 Hz), 2.66 (1 H, m), 2.48 (1 H, m), 2.36 (1 H, m).

MS (ESI) m/z: 367.0 (M+H)⁺.

[0271] IM I-c-4

¹H NMR (CDCl₃) delta 8.53 (1 H, dd, J = 4.9, 1.2 Hz), 7.99 (1 H, br t, J = 5.5 Hz), 7.70 (1 H, d, J = 7.9 Hz), 7.39 (1 H, dd, J = 7.9, 4.9 Hz), 7.06 (1 H, m), 6.89 (1 H, m), 4.45 (2 H, m), 4.04 (1 H, br t, J = 5.5 Hz), 3.03 (1 H, ddd, J = 17.7, 10.4, 4.9 Hz), 2.65 (1 H, m), 2.50-2.34 (2 H, m).

MS (ESI) m/z: 335.1 (M+H)⁺.

[0272] IM I-c-5

¹H NMR (CDCl₃) delta 8.49 (1 H, dd, J = 4.3, 1.2 Hz), 8.07 (1 H, br t, J = 5.5 Hz), 7.69 (1 H, d, J = 7.9 Hz), 7.39 (1 H, dd, J = 7.9, 4.3 Hz), 6.62 (2 H, m), 4.46 (2 H, m), 4.01 (1 H, br t, J = 5.5 Hz), 2.99 (1 H, m), 2.64 (1 H, m), 2.42-2.28 (2 H, m).

MS (ESI) m/z: 335.1 (M+H)⁺.

[0273] IM I-c-6

¹H NMR (CDCl₃) delta 8.50 (1 H, dd, J = 4.9, 1.2 Hz), 7.68 (1 H, dd, J = 7.9, 1.2 Hz), 7.60 (1 H, br t, J = 5.5 Hz), 7.36 (1 H, dd, J = 7.9, 4.9 Hz), 7.26 (1 H, ddd, J = 15.3, 8.6, 6.7 Hz), 6.80-6.70 (2 H, m), 4.45 (1 H, dd, J = 15.9, 5.5 Hz), 4.41 (1 H, dd, J = 15.9, 5.5 Hz), 4.02 (1 H, br t, J = 5.5 Hz), 2.99 (1 H, ddd, J = 17.7, 10.4, 4.9 Hz), 2.66 (1 H, ddd, J = 17.7, 7.3, 4.9 Hz), 2.50-2.32 (2 H, m).

MS (ESI) m/z: 317.1 (M+H)⁺.

[0274] IM I-c-7

¹H NMR (CDCl₃) delta 8.55 (1 H, dd, J = 4.9, 1.2 Hz), 7.66 (1 H, dd, J = 7.9, 1.2 Hz), 7.51 (1 H, dd, J = 8.5, 5.5 Hz), 7.38 (1 H, dd, J = 7.9, 4.9 Hz), 7.31 (1 H, dd, J = 8.5, 2.4 Hz), 7.18 (1 H, ddd, J = 8.5, 7.9, 2.4 Hz), 7.12 (1 H, br t, J = 6.1 Hz), 4.60 (1 H, dd, J = 15.9, 6.1 Hz), 4.55 (1 H, dd, J = 15.9, 6.1 Hz), 3.99 (1 H, br t, J = 5.5 Hz), 2.97 (1 H, ddd, J = 17.7, 9.8, 4.9 Hz), 2.68 (1 H, ddd, J = 17.7, 6.7, 4.9 Hz), 2.53-2.34

(2 H, m).

MS (ESI) m/z: 367.1 (M+H)⁺.

[0275] IM I-c-8

¹H NMR (CDCl₃) delta 8.49 (1 H, dd, J = 4.9, 1.2 Hz), 7.67 (1 H, dd, J = 7.9, 1.2 Hz), 7.63 (1 H, br t, J = 6.1 Hz), 7.36 (1 H, dd, J = 7.9, 4.9 Hz), 7.21 (1 H, dd, J = 8.5, 7.3 Hz), 7.02 (1 H, dd, J = 8.9, 2.4 Hz), 6.99 (1 H, dd, J = 7.3, 2.4 Hz), 4.43 (1 H, dd, J = 15.3, 6.1 Hz), 4.38 (1 H, dd, J = 15.3, 6.1 Hz), 4.00 (1 H, br t, J = 5.5 Hz), 2.99 (1 H, ddd, J = 17.7, 9.8, 4.9 Hz), 2.65 (1 H, ddd, J = 17.7, 6.7, 5.5 Hz), 2.46-2.31 (2 H, m).

MS (ESI) m/z: 333.1 (M+H)⁺.

[0276] IM I-c-9

¹H NMR (CDCl₃) delta 8.50 (1 H, dd, J = 4.9, 1.2 Hz), 7.68 (1 H, dd, J = 7.9, 1.2 Hz), 7.60 (1 H, br t, J = 6.1 Hz), 7.37 (1 H, dd, J = 7.9, 4.9 Hz), 7.22-7.14 (3 H, m), 4.42 (2 H, m), 4.00 (1 H, br t, J = 5.5 Hz), 2.99 (1 H, ddd, J = 17.7, 10.4, 4.9 Hz), 2.66 (1 H, ddd, J = 17.7, 7.3, 4.9 Hz), 2.49-2.31 (2 H, m).

MS (ESI) m/z: 377.0 (M+H)⁺.

[0277] IM I-c-10

¹H NMR (CDCl₃) delta 8.69 (1 H, d, J = 4.3 Hz), 7.65 (1 H, d, J = 7.9 Hz), 7.44 (1 H, dd, J = 7.9, 4.3 Hz), 7.15 (1 H, m), 7.07 (1 H, dd, J = 8.6, 7.3 Hz), 6.35 (1 H, br), 4.57 (1 H, dd, J = 15.3, 6.1 Hz), 4.49 (1 H, dd, J = 15.3, 5.5 Hz), 3.91 (1 H, br t, J = 5.5 Hz), 2.94 (1 H, ddd, J = 17.7, 10.4, 4.9 Hz), 2.75 (1 H, ddd, J = 17.7, 6.1, 4.9 Hz), 2.56 (1 H, ddd, J = 17.7, 11.7, 6.1 Hz), 2.42 (1 H, ddd, J = 17.7, 10.4, 4.9 Hz).

MS (ESI) m/z: 351.0 (M+H)⁺.

[0278] IM I-c-11

¹H NMR (DMSO d₆) delta 8.97 (1 H, br t, J = 5.9 Hz), 8.71 (1 H, d, J = 4.6 Hz), 7.98 (1 H, d, J = 7.9 Hz), 7.77 (1 H, dd, J = 5.9, 2.6 Hz), 7.64 (1 H, dd, J = 7.9, 4.6 Hz), 7.66-7.50 (2 H, m), 6.81 (1 H, s), 4.45 (2 H, br d J = 5.9 Hz), 2.88-2.70 (2 H, m), 2.50 (1 H, m), 2.34 (1 H, m).

MS (ESI) m/z: 399.3 (M+H)⁺.

[0279] IM I-c-12

MS (ESI) m/z: 335.5 (M+H)⁺.

[0280] The following Examples and Intermediates were prepared by General Procedure C, D, or E (Tables 17 and 19).

[0281] General Procedure C

O₃ was bubbled into a solution of substrate (1.0 eq.) in 50% CH₂Cl₂-MeOH at -78 °C until starting material consumed. N₂ were bubbled into the mixture to remove the excess of O₃ at -78 °C. The mixture was quenched with NaBH₄ (10.0 eq.) and the mixture was warmed to room temperature gradually. After being stirred at room temperature for 1 h, 10% aq. citric acid was added to the mixture. The mixture extracted

with CH_2Cl_2 3 times and the extracts were washed with water. The extracts were combined, dried over Na_2SO_4 , and concentrated in vacuo. The resulting residue was purified by NH gel column chromatography and/or SCX cartridge column, preparative HPLC to afford the following Examples and Intermediates.

[0282] General Procedure D

Sodium borohydride (1.5 eq.) was added to a solution of substrate (1.0 eq.) in MeOH and the mixture was stirred at room temperature until complete reaction. Water was added to the mixture and the volatile was removed by evaporation. The residue was extracted with EtOAc and washed with brine. The extract was dried over Na_2SO_4 and concentrated in vacuo. The resulting residue was purified by NH gel column chromatography and/or SCX cartridge column, preparative HPLC to afford the following Examples and Intermediates.

[0283] General Procedure E

To a solution of substrate in CH_2Cl_2 (0.065 M) was added TFA (0.065 M) at room temperature. The mixture was stirred at room temperature until complete reaction, the mixture was poured into water. The mixture was extracted with CH_2Cl_2 . The extract was washed with water, dried over Na_2SO_4 and concentrated in vacuo to afford the following Examples and Intermediates.

[0284]







[Table 17-1]

Examples	Structure	Chemical Name	Substrate	General Procedure
18		N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		D
19		N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		D
20		N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		D
21		N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		C
22		N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		C

[Table 17-2]

23		N-(2,4-dichloro-6-(methoxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-e-21	C
24		N-(2,4-dichloro-6-methylbenzyl)-5,8-dihydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		Ex 15	D
25		N-(2-chloro-3-(trifluoromethyl)benzyl)-5,8-dihydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-c-11	D
26		N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		Ex 16	D
27		N-(2-chloro-3-(trifluoromethyl)benzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		Ex 17	D
28		N-(2,3-dichlorobenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		IM I-c-12	D

[Table 17-3]

29		N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		C
30		2-(5-((2,4-dichloro-6-methylbenzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid		E
31		2-(5-((2-chloro-3-(trifluoromethyl)benzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid		E

[0285] [Table 18]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
18	D	1.66, 1.69	383.0	25	A	1.40	401.3
19	D	1.56, 1.59	403.0	26	D	1.51	351.0
20	A	1.35	399.2	27	D	1.40, 1.45	371.0
21	A	1.45	369.1	28	D	1.33, 1.38	337.0
22	A	1.49	369.1	29	D	1.52	355.1
23	A	1.57	413.1	30	A	1.27	407.1
24	A	1.46	381.3	31	A	1.23	427.2

[0286] Ex 18

¹H NMR (DMSO d₆) delta 8.74 (1 H, m), 8.60 (1 H, br s), 7.58 (0.4 H, d, J = 7.9 Hz), 7.55 (0.6 H, d, J = 7.9 Hz), 7.46 (1 H, br s), 7.38 (1 H, dd, J = 7.9, 4.6 Hz), 7.35 (1 H, br s), 5.41 (1 H, br s), 4.61 (1 H, m), 4.48 (0.8 H, d, J = 4.6 Hz), 4.43 (1.2 H, d, J = 4.6 Hz), 2.80-1.75 (4 H, m), 2.42 (1.2 H, s), 2.38 (1.8 H, s).

[0287] Ex 19

¹H NMR (CDCl₃) delta 8.60 (1 H, br), 7.71-7.62 (2 H, m), 7.51 (1 H, m), 7.39 (1 H, m), 7.30 (1 H, br), 7.30-7.20 (1 H, m), 4.78-4.69 (3 H, m), 4.33 (0.5 H, br s), 3.75 (0.5 H, br s), 2.76-1.80 (4 H, m).

[0288] Ex 20

¹H NMR (DMSO d₆) delta 8.69 (1 H, br), 8.62 (1 H, br), 7.58 (1 H, m), 7.56 (1 H, d,

J = 2.6 Hz), 7.48 (1 H, d, J = 2.6 Hz), 7.38 (1 H, dd, J = 7.9, 4.6 Hz), 5.53 (1 H, br), 5.43 (1 H, m), 4.71-4.55 (3 H, m), 4.52-4.40 (2 H, m), 2.40-1.72 (4 H, m).

[0289] Ex 24

¹H NMR (DMSO d6) delta 8.48 (1 H, br dd, J = 4.6, 2.0 Hz), 8.07 (1 H, m), 7.52-7.45 (2 H, m), 7.32 (1 H, J = 2.6 Hz), 7.26 (1 H, dd, J = 7.9, 4.6 Hz), 6.23 (0.5 H, s), 6.11 (0.5 H, s), 5.16 (1 H, m), 4.54-4.42 (3 H, m), 2.41 (1.5 H, s), 2.39 (1.5 H, s), 2.30-1.64 (4 H, m).

[0290] Ex 25

¹H NMR (DMSO d6) delta 8.86 (1 H, t, J = 6.6 Hz), 8.49 (1 H, br d, J = 4.6 Hz), 7.78 (1 H, m), 7.66-7.57 (3 H, m), 7.30 (1 H, dd, J = 7.9, 4.6 Hz), 6.39 (0.5 H, br), 6.27 (0.5 H, br), 5.20 (0.5 H, d, J = 2.6 Hz), 5.16 (0.5 H, d, J = 3.3 Hz), 4.56-4.44 (3 H, m), 2.33-1.75 (4 H, m).



[0291] Ex 26

¹H NMR (DMSO d6) delta 8.53 (1 H, br), 8.44 (1 H, d, J = 4.6 Hz), 7.52 (1 H, d, J = 7.9 Hz), 7.50 (1 H, d, J = 2.0 Hz), 7.36 (1 H, d, J = 2.0 Hz), 7.23 (1 H, dd, J = 7.9, 4.6 Hz), 5.48 (1 H, d, J = 6.6 Hz), 4.90 (1 H, m), 4.44 (1 H, d, J = 4.6 Hz), 4.42 (1 H, d, J = 4.6 Hz), 3.82 (1 H, m), 2.54 (1 H, m), 2.39 (3 H, s), 2.02 (1 H, m).

[0292] Ex 29

¹H NMR (CDCl₃) delta 8.69 (1 H, br dd, J = 4.9, 1.2 Hz), 7.62 (1 H, br dd, J = 7.9, 1.2 Hz), 7.44 (1 H, d, J = 1.8 Hz), 7.35-7.16 (4 H, m), 5.33 (1 H, dd, J = 6.1, 2.4 Hz), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 3.94 (1 H, d, J = 5.5 Hz), 2.83-2.66 (2 H, m).

[0293] [Table 19]

Intermediate	Structure	Chemical Name	Substrate	General Procedure
I-w-1		2-(5-((2,4-dichloro-6-methoxybenzyl)carbamoyl)-5-fluoro-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid	 IM I-v-3	E



[0294] IM I-w-1

MS (ESI) m/z: 425.0 (M+H)⁺.

[0295] The following Example was prepared by General Procedure A (Table 20).

[0296]

[Table 20]

Example	Structure	Chemical Name	Substrate	Amine
32		(2-amino-2-oxoethyl)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 Ex 30	NH ₄ Cl

[0297] Ex 32

LCMS (ESI) m/z: 426.2 (M+H)⁺, tR 1.39 min (Method A).













[0298] The following Examples were prepared by General Procedure F (Table 21).

[0299] General Procedure F

A mixture of substrate (1.0 eq.), 4% aq. OsO₄ (1.0 eq.), and NMO (15.0 eq.) in 50% aq. THF was stirred at room temperature until complete reaction. Aq. Na₂S₂O₃ was added to the mixture and the mixture was stirred for 10 min. The mixture was extracted with EtOAc twice. The extracts were washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography and/or SCX column, preparative HPLC to afford the following Examples.

[0300]

[Table 21]

Examples	Structure	Chemical Name	Substrate
33		N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-1
34		N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-16
35		N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-3
36		N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-18
37		N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-e-19
38		N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-7-(hydroxymethyl)-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide	 IM I-e-24

[0301] [Table 22]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
33	A	1.41	395.3	36	A	1.39	399.1
34	A	1.52	413.1	37	A	1.42	399.1
35	A	1.44	433.1	38	D	1.41	385.1

[0302] The following Intermediates were prepared by General Procedure G or H (Table 23).

[0303] General Procedure G

A substrate (1.0 eq.) and trimethylsulfoxonium iodide (1.2 eq.) were dissolved in DMSO. A solution of KOBu^t (1.2 eq.) in DMSO was added to the mixture at ambient temperature. After being stirred at room temperature until complete reaction, the mixture was poured into ice-water. The mixture was extracted with EtOAc twice and washed with brine. The extracts were combined, dried over Na₂SO₄, and concrete in vacuo. The resulting residue was purified by silica gel column chromatography to afford the following Intermediates. The each diastero isomer can be separated by silica gel column chromatography.

[0304] General Procedure H

To a solution of substrate (1.0 eq.) in water-tert-BuOH-THF (2:1:1) was added NBS (2.0 eq.) in portions at ambient temperature and then the mixture was heated at 50 °C until starting material consumed. After cooling to 0 °C, the reaction mixture was basified with 5 M aqueous NaOH and stirred at 0 °C until complete reaction. The reaction mixture was poured into water and extract with EtOAc twice. The extracts were washed with brine, and dried over Na₂SO₄. The combined extracts were evaporated and the resulting residue was purified by silica gel column chromatography to afford the following Intermediates. The each diastereo isomer can be separated by silica gel column chromatography.

[0305]

[Table 23]

Intermediates	Structure	Chemical Name	Substrate	General Procedure
I-g-1		N-(2,4-dichloro-6-methylbenzyl)-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		G
I-g-2		N-(2,4-dichloro-6-methylbenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		G
I-g-3		N-(2-chloro-3-(trifluoromethyl)benzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		G
I-g-4		(2S*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		H
I-g-5		(2R*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		
I-g-6		(2S*,5'S*)-N-(2-chloro-4-fluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		H
I-g-7		(2R*,5'S*)-N-(2-chloro-4-fluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		

[0306] IM I-g-1

¹H NMR (CDCl₃) delta 8.53 (1 H, dd, J = 4.6, 1.3 Hz), 7.45 (0.5 H, d, J = 7.9 Hz), 7.42 (0.5 H, dd, J = 7.9, 1.3 Hz), 7.24 (0.5 H, d, J = 2.0 Hz), 7.22 (0.5 H, d, J = 2.0 Hz), 7.18 (1 H, dd, J = 7.9, 4.6 Hz), 7.11 (0.5 H, d, J = 2.0 Hz), 7.10 (0.5 H, d, J = 2.0 Hz),

Hz), 5.94 (0.5 H, br t, J = 5.9 Hz), 5.79 (0.5 H, br t, J = 5.9 Hz), 4.63-4.41 (2 H, m), 3.75 (1 H, m), 3.73 (0.5 H, d, J = 5.9 Hz), 3.62 (0.5 H, d, J = 5.9 Hz), 2.98 (0.5 H, d, J = 5.9 Hz), 2.97 (0.5 H, d, J = 5.9 Hz), 2.47 (1.5 H, s), 2.46 (1.5 H, s), 2.50-1.80 (4 H, m).

MS (ESI) m/z: 377.0 (M+H)⁺.

[0307] IM I-g-2

MS (ESI) m/z: 395.0 (M+H)⁺.

[0308] IM I-g-3

¹H NMR (CDCl₃) delta 8.63 (1 H, dd, J = 4.6, 2.0 Hz), 7.67 (2 H, m), 7.44 (2 H, m), 7.30-7.04 (2 H, m), 4.75 (2 H, d, J = 5.9 Hz), 3.86 (1 H, d, J = 5.9 Hz), 3.03 (1 H, d, J = 5.9 Hz), 2.99-2.00 (4 H, m).

MS (ESI) m/z: 414.9 (M+H)⁺.

[0309] IM I-g-4

¹H NMR (CDCl₃) delta 8.63 (1 H, br dd, J = 4.9, 1.8 Hz), 7.50 (1 H, br dd, J = 7.9, 1.8 Hz), 7.46 (1 H, d, J = 2.4 Hz), 7.39 (1 H, d, J = 8.6 Hz), 7.27 (1 H, br dd, J = 7.9, 1.8 Hz), 7.23 (1 H, dd, J = 7.9, 4.9 Hz), 7.19 (1 H, br), 4.67 (1 H, dd, J = 14.7, 6.1 Hz), 4.62 (1 H, dd, J = 14.7, 6.1 Hz), 3.86 (1 H, d, J = 6.1 Hz), 3.03 (1 H, d, J = 6.1 Hz), 2.76 (1 H, m), 2.57 (1 H, m), 2.32 (1 H, m), 2.14 (1 H, m).

MS (ESI) m/z: 380.9 (M+H)⁺.

[0310] IM I-g-5

¹H NMR (CDCl₃) delta 8.63 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.51 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.46 (1 H, d, J = 1.8 Hz), 7.37 (1 H, d, J = 7.9 Hz), 7.27 (1 H, dd, J = 7.9, 1.8 Hz), 7.24 (1 H, br), 7.22 (1 H, dd, J = 7.9, 4.9 Hz), 4.65 (1 H, dd, J = 14.7, 6.1 Hz), 4.60 (1 H, dd, J = 14.7, 6.1 Hz), 3.51 (1 H, d, J = 6.1 Hz), 3.06 (1 H, d, J = 6.1 Hz), 2.65 (1 H, m), 2.50-2.39 (2 H, m), 2.19 (1 H, m).

MS (ESI) m/z: 380.9 (M+H)⁺.

[0311] IM I-g-6

¹H NMR (CDCl₃) delta 8.63 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.49 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.44 (1 H, dd, J = 8.6, 5.5 Hz), 7.23 (1 H, dd, J = 7.9, 4.9 Hz), 7.18 (1 H, dd, J = 8.6, 2.4 Hz), 7.17 (1 H, br), 7.01 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 4.67 (1 H, dd, J = 14.7, 6.1 Hz), 4.62 (1 H, dd, J = 14.7, 6.1 Hz), 3.86 (1 H, d, J = 6.1 Hz), 3.03 (1 H, d, J = 6.1 Hz), 2.77 (1 H, m), 2.57 (1 H, m), 2.32 (1 H, m), 2.13 (1 H, m).

MS (ESI) m/z: 365.1 (M+H)⁺.

[0312] IM I-g-7

¹H NMR (CDCl₃) delta 8.62 (1 H, br dd, J = 4.9, 1.8 Hz), 7.51 (1 H, br dd, J = 7.9, 1.8 Hz), 7.42 (1 H, dd, J = 8.5, 6.1 Hz), 7.23-7.18 (3 H, m), 7.00 (1 H, m), 4.66 (1 H, dd, J = 14.7, 6.1 Hz), 4.60 (1 H, dd, J = 14.7, 6.1 Hz), 3.51 (1 H, d, J = 6.1 Hz), 3.06 (1 H, d, J = 6.1 Hz), 2.66 (1 H, m), 2.51-2.39 (2 H, m), 2.19 (1 H, m).

MS (ESI) m/z: 364.7 (M+H).

[0313] The following Examples were prepared by General Procedure I, J, or K (Table 24).

[0314] General Procedure I

A solution of 1.0 M NaOMe in MeOH (1.3 eq.) was added to substrate (1.0 eq.) in small portions at ambient temperature. After being stirred at room temperature until complete reaction, the mixture was concentrated in vacuo to afford a glass. The residual glass was distributed between EtOAc and water. The extract was dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by SCX cartridge column and preparative HPLC to afford the following Examples.

[0315] General Procedure J









A solution of substrate (1.0 eq.) in MeOH was treated with amine (45 eq.) at room temperature until complete reaction. The mixture was concentrated in vacuo and the resulting residue was distributed between EtOAc and water. The extract was dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography and/or SCX column and preparative HPLC to afford the following Examples.

[0316] General Procedure K







To a stirred solution of substrate (1.0 eq.) in EtOH was added NaBH₄ (6.0 eq.) at ambient temperature. The mixture was stirred at room temperature until complete reaction, water was added to the mixture. The mixture was extracted with EtOAc and washed with aq. NaHCO₃. The extract was dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by SCX cartridge column and preparative HPLC to afford the following Examples.

[0317]

[Table 24-1]

Examples	Structure	Chemical Name	Substrate	General Procedure (/Amine)
39		N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		I
40		N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		I
41		(5S*,8S*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		I
42		(5S*,8S*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		I






[Table 24-2]

43		8-(aminomethyl)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-g-1	J /NH ₄ OH
44		N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		J /MeNH ₂ in MeOH
45		8-(aminomethyl)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		J /NH ₄ OH
46		N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		J /MeNH ₂ in MeOH
47		N-(2,4-dichloro-6-methylbenzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		J /Me ₂ NH in THF





[Table 24-3]

48		8-(aminomethyl)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		J /NH ₄ OH
49		N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		J /MeNH ₂ in MeOH
50		N-(2-chloro-3-(trifluoromethyl)benzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		J /Me ₂ NH in THF
51		(5S*,8R*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		J/
52		(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		J/

[Table 24-4]

53		(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	
54		(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxy-3-methylazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	
55		(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxy-3-methylazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	
56		(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((2-hydroxyethyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	

[Table 24-5]

57		(5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		J/ 
58		N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-g-3	K

[0318] [Table 25]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
39	A	1.67	427.1	47	A	1.50	440.2
40	A	1.57	447.1	48	A	1.36	432.1
43	A	1.35	394.2	49	A	1.38	446.1
44	A	1.36	408.1	50	A	1.43	460.1
45	A	1.42	412.2	58	A	1.58	417.1
46	A	1.45	426.2				

[0319] Ex 41

¹H NMR (DMSO d6) delta 9.27 (1 H, br s), 8.74 (1 H, d, J = 4.5 Hz), 7.75-7.80 (2 H, m), 7.62 (1 H, d, J = 2.0 Hz), 7.53 (1 H, dd, J = 8.0, 4.7 Hz), 7.45 (1 H, dd, J = 8.4, 2.1 Hz), 7.36 (1 H, d, J = 8.3 Hz), 4.40 (2 H, t, J = 5.3 Hz), 3.88 (1 H, d, J = 8.6 Hz), 3.53 (1 H, d, J = 8.4 Hz) 2.29-2.48 (5 H, m).

MS (ESI) m/z: 413.2 (M+H)⁺.

[0320] Ex 42

¹H NMR (CDCl₃) delta 8.61 (1 H, dt, J = 4.7, 1.6 Hz), 7.51 (1 H, dt, J = 7.8, 1.5 Hz), 7.40 (1 H, dd, J = 8.5, 6.1 Hz), 7.14-7.25 (3 H, m), 6.99 (1 H, td, J = 8.3, 2.6 Hz), 4.51-4.68 (2 H, m), 3.76 (2 H, dd, J = 9.5, 8.4 Hz), 3.34 - 3.37 (4 H, m), 2.57 - 2.72 (1 H, m), 2.38-2.49 (1 H, m) 2.18-2.32 (2 H, m).

MS (ESI) m/z: 397.2 (M+H)⁺.

[0321] Ex 51

¹H NMR (CDCl₃) delta 8.61 (1 H, br d, J = 4.7 Hz), 7.54 (1 H, d, J = 8.0 Hz), 7.44 (1 H, d, J = 2.1 Hz), 7.34 (1 H, d, J = 7.7 Hz), 7.27-7.30 (1 H, m), 7.19-7.25 (1 H, m), 4.48-4.65 (3 H, m), 3.99-4.13 (4 H, m), 3.89 (2 H, br s), 3.56 (1 H, br d, J = 13.0 Hz),

3.18 (1 H, br d, J = 13.1 Hz), 2.67-2.82 (1 H, m), 2.16-2.28 (3 H, m).

MS (ESI) m/z: 454.2 (M+H)⁺.

[0322] Ex 52

¹H NMR (CDCl₃) delta 8.62 (1 H, dt, J = 4.7, 1.6 Hz), 7.49 (1 H, dt, J = 7.8, 1.5 Hz), 7.41 (1 H, dd, J = 8.5, 6.1 Hz), 7.09-7.25 (3 H, m), 6.99 (1 H, td, J = 8.3, 2.6 Hz), 4.53-4.67 (2 H, m), 4.41 (1 H, quin, J = 5.78 Hz), 3.74-3.82 (1 H, m), 3.57 (1 H, br s), 3.07-3.16 (2 H, m), 3.00 (1 H, br t, J = 6.4 Hz), 2.85 (1 H, d, J = 12.8 Hz), 2.56-2.77 (1 H, m), 2.13-2.37 (4 H, m).

MS (ESI) m/z: 438.3 (M+H)⁺.

[0323] Ex 53

¹H NMR (CDCl₃) delta 8.61-8.64 (1 H, m), 7.49 (1 H, d, J = 7.6 Hz), 7.41 (1 H, dd, J = 8.6, 6.0 Hz), 7.09-7.25 (3 H, m), 6.99 (1 H, td, J = 8.3, 2.6 Hz), 4.54-4.66 (2 H, m), 4.03 (1 H, br t, J = 5.8 Hz), 3.66-3.84 (1 H, m), 3.58 (1 H, br s), 3.15-3.24 (4 H, m), 2.95-3.10 (1 H, m), 2.85 (1 H, br d, J = 12.8 Hz), 2.58-2.72 (1 H, m), 2.08-2.35 (3 H, m).

MS (ESI) m/z: 452.0 (M+H)⁺.

[0324] Ex 54

¹H NMR (CDCl₃) delta 8.62 (1 H, d, J = 5.0 Hz), 7.50 (1 H, d, J = 7.6 Hz), 7.41 (1 H, dd, J = 8.6, 6.0 Hz), 7.10-7.25 (3 H, m), 6.99 (1 H, td, J = 8.3, 2.6 Hz), 4.54-4.67 (2 H, m), 3.46 (1 H, d, J = 8.0 Hz), 3.22-3.39 (4 H, m), 2.90 (1 H, d, J = 12.8 Hz), 2.59-2.74 (1 H, m), 2.14-2.27 (3 H, m), 1.45 (3 H, s).

MS (ESI) m/z: 452.3 (M+H)⁺.

[0325] Ex 55

¹H -NMR (CDCl₃) delta 8.62 (1 H, dt, J = 4.7, 1.6 Hz), 7.48 (1 H, d, J = 7.6 Hz), 7.41 (1 H, t, J = 7.0 Hz), 7.08-7.24 (3 H, m), 6.99 (1 H, td, J = 8.3, 2.6 Hz), 4.53-4.67 (2 H, m), 3.14-3.41 (6 H, m), 3.10 (2 H, br s), 2.83 (1 H, br d, J = 12.6 Hz), 2.49-2.76 (1 H, m), 2.13-2.37 (3 H, m), 1.43 (3 H, s).

MS (ESI) m/z: 466.3 (M+H)⁺.

[0326] Ex 56

¹H -NMR (CDCl₃) delta 8.59 (1 H, dt, J = 4.7, 1.7 Hz), 7.50 (1 H, dt, J = 7.9, 1.5 Hz), 7.41 (1 H, dd, J = 8.4, 6.0 Hz), 7.13-7.25 (3 H, m), 6.99 (1 H, td, J = 8.3, 2.6 Hz), 4.55-4.67 (2 H, m), 3.64 (2 H, t, J = 5.2 Hz), 3.12 (1 H, d, J = 12.0 Hz), 2.92 (1 H, d, J = 12.0 Hz), 2.68-2.83 (2 H, m), 2.28-2.36 (1 H, m), 2.13 - 2.25 (3 H, m).

MS (ESI) m/z: 426.3 (M+H)⁺.

[0327] Ex 57

¹H -NMR (CDCl₃) delta 8.62 (1 H, dt, J = 4.7, 1.7 Hz), 7.51 (1 H, d, J = 7.6 Hz), 7.41 (1 H, dd, J = 8.5, 6.1 Hz), 7.10-7.25 (3 H, m), 6.99 (1 H, td, J = 8.3, 2.6 Hz), 4.54-4.67 (2 H, m), 3.43-3.66 (3 H, m), 3.03 (1 H, d, J = 14.1 Hz), 2.87 (1 H, br d, J = 14.2 Hz),

2.57-2.73 (3 H, m), 2.14-2.41 (6 H, m).





MS (ESI) m/z : 440.0 (M+H)⁺.

The following Intermediates were prepared by General Procedure L (Table 26).

[0328] General Procedure L

NaN₃ (3.0 eq.) and NH₄Cl (3.0 eq.) were added to a solution of substrate (1.0 eq.) in MeOH, and the mixture was stirred at 60 °C until complete reaction. The reaction was quenched with aq. NaHCO₃, and the resulting mixture was extracted with CHCl₃. The extract was concentrated under reduced pressure to afford the following Intermediates.

[0329] [Table 26]

Intermediates	Structure	Chemical Name	Substrate
I-h-3-1		(5S*,8R*)-8-(azidomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-g-4
I-h-3-2		(5S*,8S*)-8-(azidomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-g-6

[0330] IM I-h-3-1

¹H -NMR (CDCl₃) delta 8.65 (1 H, dt, J = 4.7, 1.71 Hz), 7.53 (1 H, dt, J = 7.9, 1.5 Hz), 7.45 (1 H, d, J = 2.1 Hz), 7.36 (1 H, d, J = 8.2 Hz), 7.27-7.31 (2 H, m), 7.16 (1 H, br d, J = 5.3 Hz), 4.54-4.67 (2 H, m), 3.70 (2 H, d, J = 1.1 Hz), 3.20 (s, 1 H), 2.59-2.74 (1 H, m), 2.36-2.46 (1 H, m), 2.13-2.33 (2 H, m).

MS (ESI) m/z : 424.2 (M+H)⁺.

[0331] IM I-h-3-2

¹H -NMR (CDCl₃) delta 8.60-8.66 (1 H, m), 7.52 (1 H, d, J = 8.2 Hz), 7.46 (1 H, s), 7.17-7.39 (6 H, m), 4.54-4.68 (2 H, m), 3.53-3.60 (1 H, m), 3.41 (1 H, d, J = 13.0 Hz), 2.41-2.55 (2 H, m), 2.26-2.39 (1 H, m), 2.01-2.21 (1 H, m).

MS (ESI) m/z : 424.2 (M+H)⁺.

[0332] General Procedure: Scheme 11, Step 1

To a stirred solution of substrate (1.0 eq.) in MeCN and water (100 eq.) at 60 °C was added portion wise triphenyl phosphine (2.0 eq.). The mixture was stirred at that temperature until complete reaction. The mixture was purified by silica gel column

chromatography to afford the following Intermediates (Table 27).

[0333] [Table 27]

Intermediates	Structure	Chemical Name	Substrate
I-m-1		(2S*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[aziridine-2,8'-quinoline]-5'-carboxamide	
I-m-2		(2R*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[aziridine-2,8'-quinoline]-5'-carboxamide	

[0334] IM I-m-1

¹H -NMR (CDCl₃) delta 8.46 (1 H, dt, J = 4.7, 1.9 Hz), 7.21-7.48 (6 H, m), 7.13 (1 H, ddd, J = 7.8, 4.8, 0.9 Hz), 4.57-4.72 (2 H, m), 2.57-2.78 (1 H, m), 2.45-2.55 (1 H, m), 2.34 (1 H, tdd, J = 14.1, 14.1, 4.3, 2.9 Hz), 2.16 (1 H, s) 1.91 (1 H, s), 1.71 (1 H, dt, J = 13.6, 3.7 Hz).

MS (ESI) m/z: 380.1 (M+H)⁺.

[0335] IM I-m-2

¹H -NMR (CDCl₃) delta 8.47 (1 H, dt, J = 4.7, 1.7 Hz), 7.63-7.70 (1 H, m), 7.32-7.49 (4 H, m), 7.19-7.29 (2 H, m), 7.13 (1 H, ddd, J = 7.9, 4.7, 0.7 Hz), 4.54-4.68 (2 H, m), 2.49-2.70 (3 H, m), 2.23-2.36 (2 H, m), 2.09 - 2.20 (1 H, m).





MS (ESI) m/z: 380.1 (M+H)⁺.

[0336] General Procedure: Scheme 11, Step 2

Triethylamine (1.5 eq.) and 2-nitrophenylsulfonyl chloride (1.2 eq.) were successively added to a dichloromethane solution of substrate (1.0 eq.) at 0 °C. After stirring at ambient temperature until complete reaction, the reaction was quenched with aq. NH₄Cl. The mixture was extracted with CHCl₃, dried over Na₂SO₄, and concentrated under reduced pressure to afford the following Intermediates (Table 28).

[0337]

[Table 28]

Intermediates	Structure	Chemical Name	Substrate
XIV-1		(2S*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-1-((2-nitrophenyl)sulfonyl)-6',7'-dihydro-5'H-spiro[aziridine-2,8'-quinoline]-5'-carboxamide	
XIV-2		(2R*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-1-((2-nitrophenyl)sulfonyl)-6',7'-dihydro-5'H-spiro[aziridine-2,8'-quinoline]-5'-carboxamide	

[0338] IM XIV-1

MS (ESI) m/z: 565.0 (M+H)⁺

[0339] IM XIV-2





MS (ESI) m/z: 565.0 (M+H)⁺.

[0340] General Procedure: Scheme 11, Step 3

To a solution of substrate (1.0 eq.) in DMF was added NaOAc (10 eq.) and the mixture was stirred at 100 °C until starting material consumed. Aq. NaOH (5 N, 5 eq.) was added to the mixture and stirred at ambient temperature until complete reaction. The reaction was quenched with aq. NH₄Cl, extracted with CHCl₃. The extract was concentrated under reduced pressure. The residue was purified by silica gel column chromatography to afford the following Intermediates (Table 29).

[0341]

[Table 29]

Intermediates	Structure	Chemical Name	Substrate
XV-1		(5S*,8R*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-8-(2-nitrophenylsulfonamido)-5,6,7,8-tetrahydroquinoline-5-carboxamide	
XV-2		(5S*,8S*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-8-(2-nitrophenylsulfonamido)-5,6,7,8-tetrahydroquinoline-5-carboxamide	

[0342] IM XVI-1

MS (ESI) m/z: 583.1 (M+H)⁺.

[0343] IM XV-2

MS (ESI) m/z: 583.1 (M+H)⁺.

[0344] General Procedure: Scheme 11, Step 4

The mixture of substrate (1.0 eq.), 4-mercaptobenzoic acid (2.0 eq.), and K₂CO₃ (4.0 eq.) in DMF was stirred at 80 °C until complete reaction. The solvent of the mixture was removed under reduced pressure. The residue was purified by SCX cartridge column to afford the following Examples (Table 30).

[0345]

[Table 30]

Examples	Structure	Chemical Name	Substrate
59		(5S*,8R*)-8-amino-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	
60		(5S*,8S*)-8-amino-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	

[0346] Ex 59

¹H -NMR (CDCl₃) delta 8.56-8.65 (1 H, m), 7.50 (1 H, br d, J = 8.1 Hz), 7.45 (1 H, d, J = 2.2 Hz), 7.30-7.39 (1 H, m), 7.13-7.29 (3 H, m), 4.60 (2 H, dd, J=10.0, 5.9 Hz), 2.90-3.09 (2 H, m), 2.29-2.45 (2 H, m), 2.05-2.17 (2 H, m).

MS (ESI) m/z: 398.2 (M+H)⁺.

[0347] EX 60

¹H -NMR (CDCl₃) delta 8.54 (1 H, dd, J = 2.7, 1.3 Hz), 7.52 (1 H, d, J = 7.3 Hz) 7.40-7.44 (1 H, m), 7.27-7.36 (2 H, m), 7.12-7.25 (3 H, m), 4.51-4.65 (2 H, m), 3.14-3.31 (2 H, m), 2.67-3.00 (2 H, m), 2.17-2.31 (2 H, m).

MS (ESI) m/z: 398.2 (M+H)⁺.

[0348] General Procedure: Scheme 12, Step 1

Chloroacetyl chloride (1.1 eq.) was added dropwise to a biphasic solution of substrate (1.0 eq.) in dichloromethane (0.3 M) and 0.5 N aq. NaOH (2.0 eq) at 0 ° C. The reaction mixture was warmed up to ambient temperature and stirred at that temperature until complete reaction. The mixture was extracted with dichloromethane 3 times, dried over Na₂SO₄, and concentrated under reduced pressure to afford the following Intermediates (Table 31).

[0349]

[Table 31]

Intermediates	Structure	Chemical Name	Substrate
XVI-1		(5S*,8R*)-8-(2-chloroacetamido)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	
XVI-2		(5S*,8S*)-8-(2-chloroacetamido)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	

[0350] IM XVI-1

MS (ESI) m/z: 474.1 (M+H)⁺.

[0351] IM XVI-2

MS (ESI) m/z: 474.1 (M+H)⁺.

[0352] General Procedure: Scheme 12, Step 2

To a solution of substrate (1.0 eq.) in 50% dichloromethane/2-propanol was added portion wise tert-BuOK (4.0 eq.) at 0 °C. The solution was allowed to warm to ambient temperature and stirred until complete reaction. The solvent was removed under reduced pressure and the resulting crude residue was purified by preparative HPLC to afford the following Examples (Table 32).

[0353]

[Table 32]

Examples	Structure	Chemical Name	Substrate
61		(3R*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide	
62		(3S*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide	

[0354] Ex 61

¹H -NMR (CDCl₃) delta 8.61-8.69 (1 H, m), 7.54 (1 H, d, J = 7.7 Hz), 7.46 (1 H, d, J = 2.0 Hz), 7.16-7.40 (6 H, m), 6.18 (1 H, br s), 4.56-4.69 (2 H, m), 4.34 (1 H, dd, J = 12.2, 2.2 Hz), 4.17 (2 H, dd, J = 17.5, 16.6 Hz), 3.37 (1 H, dd, J = 12.2, 2.9 Hz), 2.77-2.95 (1 H, m), 2.40-2.50 (1 H, m), 2.12-2.32 (2 H, m).

MS (ESI) m/z: 438.1 (M+H)⁺.

[0355] Ex 62

¹H -NMR (CDCl₃) delta 8.70 (1 H, d, J = 4.7 Hz), 7.57 (1 H, d, J = 8.2 Hz), 7.46 (1 H, d, J = 2.1 Hz), 7.27-7.35 (3 H, m), 7.14-7.22 (1 H, m), 5.95 (1 H, br s), 4.53-4.65 (2 H, m), 4.25-4.42 (2 H, m), 4.20 (1 H, d, J = 11.4 Hz), 3.43 (1 H, dd, J = 12.4, 4.0 Hz), 2.54 - 2.64 (1 H, m) 2.29-2.53(3 H, m), 1.25 (1 H, s).

MS (ESI) m/z: 438.1 (M+H)⁺.

[0356] The following Examples and Intermediates were prepared by General Procedure M (Tables 33 and 35).

[0357] General Procedure M

To a mixture of substrate (1.0 eq.), Chiral Ru Catalyst (5 mol%) and triethylamine (2.0 eq.) in DMF was added formic acid (5.0 eq.) at 0 °C. The mixture was stirred at 0 °C for 10 min, and then warm to room temperature. After being stirred at room temperature until complete reaction, aq. NaHCO₃ was added to the mixture. The mixture was extracted with EtOAc twice and washed with water and brine. The extracts were combined, dried over Na₂SO₄, and concentrated in vacuo. The resulting residue was

purified by silica gel column chromatography and the each diastereo isomer was separated by silica gel column chromatography and/or preparative TLC to afford following Examples.









[0358] [Table 33-1]

Examples	Structure	Chemical Name	Substrate	Chiral Ru Catalyst
63		(5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
64		(5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	Ex 1	
65		(5S,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
66		(5R,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	IM I-c-1	









[Table 33-2]

67		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
68		(5R,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
69		(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
70		(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
71		(5S,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]

[Table 33-3]

72		(5R,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
73		(5R,8S)-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
74		(5R,8S)-N-(2,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
75		(5S,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
76		(5R,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 33-4]

77		(5S,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
78		(5R,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	IM I-c-8	
79		(5S,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
80		(5R,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	IM I-c-9	
81		(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]








[Table 33-5]

82		(5R,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
83		(5S,7S)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]
84		(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(R,R)-Ts-DPEN]
85		(5S,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
86		(5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(S,S)-Ts-DPEN]




[Table 33-6]

87		(5R,8R)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(R,R)-Ts-DPEN]
88		(5S,8R)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
89		(5R,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(R,R)-Ts-DPEN]
90		(5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
91		(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(R,R)-Ts-DPEN]

[Table 33-7]

92		(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
93		(5R,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(R,R)-Ts-DPEN]
94		(5S,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	IM I-c-4	
95		(5R,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		RuCl(p-cymene)[(R,R)-Ts-DPEN]
96		(5S,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	IM I-c-8	

[Table 33-8]

97		(5R,7R)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		RuCl(p-cymene)[(R,R)-Ts-DPEN]
98		(5S,7R)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide		

[0359] [Table 34]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
63	A	1.48	365.3	81	C	1.31	352.9
64	A	1.47	365.3	82	C	1.31	352.9
65	C	1.40	351.0	83	D	1.46	351.0
66	C	1.39	351.0	84	A	1.48	365.3
67	C	1.28	335.0	85	A	1.47	365.2
68	C	1.28	335.0	86	A	1.42	385.3
69	C	1.39	369.0	87	C	1.40	351.0
70	C	1.39	369.0	88	C	1.40	351.0
71	C	1.25	337.0	89	C	1.28	335.0
72	C	1.25	337.0	90	C	1.28	335.0
73	C	1.20	337.0	91	C	1.39	369.0
74	C	1.19	319.0	92	C	1.38	369.0
75	C	1.36	369.0	93	C	1.25	337.0
76	C	1.36	369.0	94	C	1.25	337.0
77	C	1.30	335.0	95	C	1.30	335.0
78	C	1.30	335.0	96	C	1.30	335.0
79	C	1.33	378.9	97	D	1.46	351.0
80	C	1.33	378.9				

[0360] Ex 63

¹H NMR (DMSO d₆) delta 8.42 (1 H, d, J = 4.6 Hz), 8.37 (1 H, br), 7.49 (1 H, br s), 7.37 (1 H, d, J = 7.9 Hz), 7.36 (1 H, br s), 7.23 (1 H, dd, J = 7.9, 4.6 Hz), 5.18 (1 H, d,

J = 4.0 Hz), 4.52 (1 H, d, J = 4.0 Hz), 4.43 (2 H, d, J = 4.0 Hz), 3.66 (1 H, dd, J = 8.6, 5.9 Hz), 2.40 (3 H, s), 2.18-1.99 (2 H, m), 1.84-1.79 (2 H, m).

Chiral HPLC tR: 28.1 min (Method E), >99% e.e., d.e.

[0361] Ex 64

¹H NMR (DMSO d6) delta 8.42 (1 H, d, J = 4.6 Hz), 8.34 (1 H, br), 7.48 (1 H, br s), 7.39 (1 H, d, J = 7.9 Hz), 7.34 (1 H, br s), 7.22 (1 H, dd, J = 7.9, 4.6 Hz), 5.11 (1 H, d, J = 3.3 Hz), 4.55 (1 H, br), 4.37 (2 H, d, J = 4.6 Hz), 3.73 (1 H, dd, J = 5.9, 5.9 Hz), 2.36 (3 H, s), 2.22 (1 H, m), 2.05 (1 H, m), 1.84 (1 H, m), 1.65 (1 H, m).

Chiral HPLC tR: 15.0 min (Method E), >99% e.e., d.e.

[0362] Ex 65

¹H NMR (CDCl₃) delta 8.49 (1 H, dd, J = 4.9, 1.2 Hz), 7.42 (1 H, dd, J = 7.9, 1.2 Hz), 7.39 (1 H, d, J = 1.8 Hz), 7.31 (1 H, d, J = 8.6 Hz), 7.22 (1 H, dd, J = 8.6, 1.8 Hz), 7.18 (1 H, dd, J = 7.9, 4.9 Hz), 6.03 (1 H, br), 4.74 (1 H, dd, J = 8.6, 5.5 Hz), 4.51 (1 H, dd, J = 15.3, 6.1 Hz), 4.46 (1 H, dd, J = 15.3, 6.1 Hz), 3.72 (1 H, dd, J = 9.2, 6.1 Hz), 2.38-2.26 (2 H, m), 2.06 (1 H, m), 1.78 (1 H, m). A signal due to OH was not observed.

[0363] Ex 66

¹H NMR (CDCl₃) delta 8.52 (1 H, dd, J = 4.9, 1.2 Hz), 7.47 (1 H, dd, J = 7.9, 1.2 Hz), 7.36 (1 H, d, J = 1.8 Hz), 7.26 (1 H, d, J = 8.6 Hz), 7.21 (1 H, m), 7.20 (1 H, dd, J = 7.9, 4.9 Hz), 5.84 (1 H, br), 4.70 (1 H, dd, J = 9.2, 5.5 Hz), 4.49 (1 H, dd, J = 14.7, 6.1 Hz), 4.39 (1 H, dd, J = 14.7, 6.1 Hz), 3.72 (1 H, dd, J = 6.1, 4.3 Hz), 2.37 (1 H, m), 2.21 (1 H, m), 2.10 (1 H, m), 1.90 (1 H, m). A signal due to OH was not observed.

[0364] Ex 67

¹H NMR (CDCl₃) delta 8.49 (1 H, dd, J = 4.9, 1.2 Hz), 7.42 (1 H, d, J = 7.9 Hz), 7.37 (1 H, dd, J = 8.6, 6.1 Hz), 7.18 (1 H, dd, J = 7.9, 4.9 Hz), 7.13 (1 H, dd, J = 7.9, 2.4 Hz), 6.96 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 6.00 (1 H, br), 4.74 (1 H, dd, J = 9.2, 5.5 Hz), 4.51 (1 H, dd, J = 15.3, 6.1 Hz), 4.47 (1 H, dd, J = 15.3, 5.5 Hz), 4.12 (1 H, br), 3.73 (1 H, dd, J = 9.2, 6.1 Hz), 2.38-2.23 (2 H, m), 2.07 (1 H, m), 1.80 (1 H, m).

[0365] Ex 68

¹H NMR (CDCl₃) delta 8.52 (1 H, dd, J = 4.9, 1.8 Hz), 7.47 (1 H, dd, J = 7.9, 1.8 Hz), 7.32 (1 H, dd, J = 8.6, 6.1 Hz), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 7.09 (1 H, dd, J = 8.6, 2.4 Hz), 6.94 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 5.87 (1 H, br), 4.70 (1 H, dd, J = 8.6, 5.5 Hz), 4.49 (1 H, dd, J = 14.7, 6.1 Hz), 4.39 (1 H, dd, J = 14.7, 6.1 Hz), 4.10 (1 H, br), 3.71 (1 H, dd, J = 6.1, 4.9 Hz), 2.36 (1 H, m), 2.20 (1 H, m), 2.10 (1 H, m), 1.90 (1 H, m).

[0366] Ex 69

¹H NMR (CDCl₃) delta 8.49 (1 H, dd, J = 4.9, 1.8 Hz), 7.43 (1 H, d, J = 7.3 Hz), 7.24 (1 H, d, J = 1.8 Hz), 7.18 (1 H, dd, J = 7.3, 4.9 Hz), 7.06 (1 H, dd, J = 9.2, 1.8 Hz),

5.90 (1 H, br), 4.73 (1 H, dd, J = 9.2, 5.5 Hz), 4.63-4.54 (2 H, m), 4.12 (1 H, br), 3.70 (1 H, dd, J = 7.9, 7.3 Hz), 2.37-2.21 (2 H, m), 2.06 (1 H, m), 1.77 (1 H, m).

[0367] Ex 70

¹H NMR (CDCl₃) delta 8.51 (1 H, dd, J = 4.9, 1.2 Hz), 7.47 (1 H, dd, J = 7.9, 1.2 Hz), 7.22-7.19 (2 H, m), 7.03 (1 H, dd, J = 9.2, 1.8 Hz), 5.74 (1 H, br), 4.68 (1 H, dd, J = 9.2, 5.5 Hz), 4.60 (1 H, ddd, J = 14.7, 6.1, 1.2 Hz), 4.46 (1 H, ddd, J = 14.7, 5.5, 1.2 Hz), 4.03 (1 H, br), 3.69 (1 H, dd, J = 6.1, 4.3 Hz), 2.35 (1 H, m), 2.19 (1 H, m), 2.08 (1 H, m), 1.89 (1 H, m).

[0368] Ex 71

¹H NMR (CDCl₃) delta 8.50 (1 H, d, J = 4.9 Hz), 7.41 (1 H, d, J = 7.9 Hz), 7.19 (1 H, dd, J = 7.9, 4.9 Hz), 7.07 (1 H, m), 6.94 (1 H, ddd, J = 9.2, 6.7, 1.8 Hz), 5.94 (1 H, br), 4.74 (1 H, dd, J = 9.2, 5.5 Hz), 4.50 (1 H, dd, J = 14.7, 6.1 Hz), 4.46 (1 H, dd, J = 14.7, 6.1 Hz), 4.09 (1 H, br), 3.73 (1 H, dd, J = 9.2, 6.1 Hz), 2.39-2.23 (2 H, m), 2.06 (1 H, m), 1.79 (1 H, m).

[0369] Ex 72

¹H NMR (CDCl₃) delta 8.51 (1 H, dd, J = 4.9, 1.8 Hz), 7.47 (1 H, d, J = 7.3 Hz), 7.22 (1 H, dd, J = 7.3, 4.9 Hz), 7.01 (1 H, m), 6.93 (1 H, m), 5.97 (1 H, br), 4.71 (1 H, dd, J = 8.6, 5.5 Hz), 4.46 (1 H, dd, J = 15.3, 6.1 Hz), 4.40 (1 H, dd, J = 15.3, 6.1 Hz), 4.12 (1 H, br), 3.73 (1 H, dd, J = 6.1, 4.9 Hz), 2.34 (1 H, m), 2.22-2.04 (2 H, m), 1.93 (1 H, m).

[0370] Ex 73

¹H NMR (CDCl₃) delta 8.52 (1 H, d, J = 4.9 Hz), 7.47 (1 H, d, J = 7.9 Hz), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 6.66 (2 H, dd, J = 8.6, 7.9 Hz), 5.75 (1 H, br), 4.69 (1 H, dd, J = 8.6, 5.5 Hz), 4.52 (1 H, ddd, J = 14.7, 5.5 Hz), 4.42 (1 H, dd, J = 14.7, 5.5 Hz), 4.10 (1 H, br s), 3.70 (1 H, dd, J = 5.5, 5.5 Hz), 2.34 (1 H, m), 2.22 (1 H, m), 2.09 (1 H, m), 1.91 (1 H, m).

[0371] Ex 74

¹H NMR (CDCl₃) delta 8.46 (1 H, dd, J = 4.9, 1.8 Hz), 7.46 (1 H, d, J = 7.9 Hz), 7.25 (1 H, m), 7.18 (1 H, dd, J = 7.9, 4.9 Hz), 6.84-6.74 (2 H, m), 6.02 (1 H, dd, J = 5.5, 5.5 Hz), 4.69 (1 H, dd, J = 7.9, 4.9 Hz), 4.43 (1 H, dd, J = 15.3, 6.1 Hz), 4.37 (1 H, dd, J = 15.3, 6.1 Hz), 4.16 (1 H, br), 3.70 (1 H, dd, J = 5.5, 5.5 Hz), 2.33 (1 H, m), 2.17-2.04 (2 H, m), 1.94 (1 H, m).

[0372] Ex 75

¹H NMR (CDCl₃) delta 8.50 (1 H, dd, J = 4.9, 1.2 Hz), 7.57 (1 H, dd, J = 8.6, 5.5 Hz), 7.40 (1 H, dd, J = 7.9, 1.2 Hz), 7.37 (1 H, dd, J = 8.6, 2.4 Hz), 7.23 (1 H, ddd, J = 8.6, 8.6, 2.4 Hz), 7.18 (1 H, dd, J = 7.9, 4.9 Hz), 6.03 (1 H, br), 4.74 (1 H, dd, J = 8.6, 5.5 Hz), 4.62 (1 H, dd, J = 15.3, 6.7 Hz), 4.58 (1 H, dd, J = 15.3, 6.1 Hz), 4.03 (1 H, br), 3.72 (1 H, dd, J = 8.6, 6.1 Hz), 2.35 (1 H, m), 2.26 (1 H, m), 2.07 (1 H, m), 1.79 (1 H, m).

[0373] Ex 76

¹H NMR (CDCl₃) delta 8.52 (1 H, dd, J = 4.9, 1.2 Hz), 7.51 (1 H, dd, J = 8.6, 5.5 Hz), 7.45 (1 H, d, J = 7.9 Hz), 7.33 (1 H, dd, J = 8.6, 2.4 Hz), 7.23-7.19 (2 H, m), 5.74 (1 H, br), 4.70 (1 H, dd, J = 8.6, 5.5 Hz), 4.57 (1 H, dd, J = 15.3, 6.1 Hz), 4.52 (1 H, dd, J = 15.3, 6.1 Hz), 3.71 (1 H, dd, J = 6.1, 4.3 Hz), 2.37 (1 H, m), 2.21 (1 H, m), 2.10 (1 H, m), 1.89 (1 H, m). A signal due to OH was not observed.

[0374] Ex 77

¹H NMR (CDCl₃) delta 8.47 (1 H, dd, J = 4.9, 1.2 Hz), 7.41 (1 H, d, J = 7.9 Hz), 7.25 (1 H, dd, J = 7.9, 6.7 Hz), 7.17 (1 H, dd, J = 7.9, 4.9 Hz), 7.10 (1 H, dd, J = 7.9, 1.8 Hz), 7.08 (1 H, dd, J = 7.9, 1.8 Hz), 6.14 (1 H, dd, J = 6.1, 5.3 Hz), 4.73 (1 H, dd, J = 9.2, 5.5 Hz), 4.46 (1 H, dd, J = 15.3, 6.1 Hz), 4.43 (1 H, dd, J = 15.3, 6.1 Hz), 4.26 (1 H, br), 3.71 (1 H, dd, J = 8.6, 6.1 Hz), 2.37-2.21 (2 H, m), 2.05 (1 H, m), 1.76 (1 H, m).

[0375] Ex 78

¹H NMR (CDCl₃) delta 8.51 (1 H, dd, J = 4.9, 1.8 Hz), 7.47 (1 H, d, J = 7.9 Hz), 7.22 (1 H, dd, J = 7.9, 5.5 Hz), 7.20 (1 H, dd, J = 7.9, 1.8 Hz), 7.10-7.04 (2 H, m), 5.80 (1 H, br), 4.70 (1 H, dd, J = 8.6, 5.5 Hz), 4.45 (1 H, dd, J = 14.7, 6.1 Hz), 4.38 (1 H, dd, J = 14.7, 6.1 Hz), 4.11 (1 H, br), 3.71 (1 H, dd, J = 5.5, 4.9 Hz), 2.35 (1 H, m), 2.19 (1 H, m), 2.09 (1 H, m), 1.91 (1 H, m).

[0376] Ex 79

¹H NMR (CDCl₃) delta 8.49 (1 H, dd, J = 4.9, 1.2 Hz), 7.42 (1 H, d, J = 7.3 Hz), 7.27-7.18 (3 H, m), 7.18 (1 H, dd, J = 7.3, 4.9 Hz), 6.01 (1 H, br), 4.74 (1 H, dd, J = 8.6, 5.5 Hz), 4.47 (1 H, dd, J = 14.7, 6.1 Hz), 4.43 (1 H, dd, J = 14.7, 6.1 Hz), 4.15 (1 H, s), 3.72 (1 H, dd, J = 8.6, 6.1 Hz), 2.38-2.22 (2 H, m), 2.06 (1 H, m), 1.77 (1 H, m).

[0377] Ex 80

¹H NMR (CDCl₃) delta 8.50 (1 H, dd, J = 4.9, 1.2 Hz), 7.46 (1 H, d, J = 7.3 Hz), 7.26-7.13 (4 H, m), 5.83 (1 H, br), 4.69 (1 H, dd, J = 8.6, 7.9 Hz), 4.43 (1 H, dd, J = 15.3, 6.1 Hz), 4.36 (1 H, dd, J = 15.3, 6.1 Hz), 4.14 (1 H, s), 3.71 (1 H, dd, J = 5.5, 4.9 Hz), 2.35 (1 H, m), 2.24-2.04 (2 H, m), 1.90 (1 H, m).

[0378] Ex 81

¹H NMR (CDCl₃) delta 8.51 (1 H, d, J = 4.9 Hz), 7.44 (1 H, d, J = 7.9 Hz), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 7.16 (1 H, m), 7.09 (1 H, m), 5.95 (1 H, br), 4.76 (1 H, dd, J = 8.6, 4.9 Hz), 4.54 (1 H, dd, J = 14.7, 6.1 Hz), 4.50 (1 H, dd, J = 14.7, 6.1 Hz), 3.75 (1 H, dd, J = 8.6, 6.1 Hz), 3.49 (1 H, s), 2.39-2.25 (2 H, m), 2.07 (1 H, m), 1.81 (1 H, m).

[0379] Ex 82

¹H NMR (CDCl₃) delta 8.53 (1 H, dd, J = 4.9, 1.8 Hz), 7.48 (1 H, d, J = 7.9 Hz), 7.23 (1 H, dd, J = 7.9, 4.9 Hz), 7.12-7.02 (2 H, m), 5.86 (1 H, br), 4.72 (1 H, dd, J = 9.2, 5.5 Hz), 4.45 (1 H, dd, J = 15.3, 6.7 Hz), 4.40 (1 H, dd, J = 15.3, 6.1 Hz), 3.73 (1 H, dd, J = 6.1, 4.2 Hz), 2.90 (1 H, br), 2.36 (1 H, m), 2.28-2.09 (2 H, m), 1.92 (1 H, m).






- [0380] Ex 83
 ¹H NMR (DMSO d6) delta 8.44 (1 H, br), 8.43 (1 H, d, J = 4.6 Hz), 7.56 (1 H, d, J = 7.3 Hz), 7.49 (1 H, d, J = 2.0 Hz), 7.35 (1 H, d, J = 2.0 Hz), 7.22 (1 H, dd, J = 7.3, 4.6 Hz), 5.41 (1 H, d, J = 5.3 Hz), 5.09 (1 H, dd, J = 11.9, 5.3 Hz), 4.38 (2 H, d, J = 4.6 Hz), 4.08 (1 H, dd, J = 7.9, 5.3 Hz), 2.47 (1 H, m), 2.36 (3 H, s), 1.97 (1 H, m).
- [0381] Ex 84
 ¹H NMR was identified with the Example 63.
 Chiral HPLC tR: 25.1 min (Method E) , >99% e.e., d.e.
- [0382] Ex 85
 ¹H NMR was identified with the Example 64.
 Chiral HPLC tR: 17.9 min (Method E) , >99% e.e., d.e.
- [0383] Ex 86
 ¹H NMR (DMSO d6) delta 8.73 (1 H, br), 8.45 (1 H, d, J = 4.6 Hz), 7.79 (1 H, dd, J = 7.9, 1.3 Hz), 7.66 (1 H, d, J = 7.9 Hz), 7.57 (1 H, dd, J = 7.9, 7.3 Hz), 7.49 (1 H, d, J = 7.3 Hz), 7.24 (1 H, dd, J = 7.9, 4.6 Hz), 5.14 (1 H, d, J = 4.0 Hz), 4.56 (1 H, br), 4.52-4.43 (2 H, m), 3.85 (1 H, dd, J = 5.9, 5.3 Hz), 2.25-2.18 (2 H, m), 1.92 (1 H, m), 1.69 (1 H, m).
- [0384] Ex 87
 ¹H NMR was identified with the Example 65.
- [0385] Ex 88
 ¹H NMR was identified with the Example 66.
- [0386] Ex 89
 ¹H NMR was identified with the Example 67.
- [0387] Ex 90
 ¹H NMR was identified with the Example 68.
- [0388] Ex 91
 ¹H NMR was identified with the Example 69.
- [0389] Ex 92
 ¹H NMR was identified with the Example 70.
- [0390] Ex 93
 ¹H NMR was identified with the Example 71.
- [0391] Ex 94
 ¹H NMR was identified with the Example 72.
- [0392] Ex 95
 ¹H NMR was identified with the Example 77.
- [0393] Ex 96
 ¹H NMR was identified with the Example 78.
- [0394] Ex 97

¹H NMR was identified with the Example 83.

[0395] Ex 98

¹H NMR (DMSO d6) delta 8.53 (1 H, br), 8.44 (1 H, d, J = 4.6 Hz), 7.52 (1 H, d, J = 7.9 Hz), 7.50 (1 H, d, J = 2.0 Hz), 7.36 (1 H, d, J = 2.0 Hz), 7.24 (1 H, dd, J = 7.9, 4.6 Hz), 5.48 (1 H, d, J = 7.33 Hz), 4.90 (1 H, dd, J = 13.8, 6.6 Hz), 4.47 (1 H, dd, J = 14.5, 4.6 Hz), 4.40 (1 H, dd, J = 14.5, 4.6 Hz), 3.82 (1 H, dd, J = 7.3, 7.3 Hz), 2.57 (1 H, m), 2.39 (3 H, s), 2.02 (1 H, m).

[0396] [Table 35-1]

Intermediates	Structure	Chemical Name	Substrate	Chiral Ru Catalyst
II-d-3		(5R,8S)-methyl 5-fluoro-8-hydroxy-5,6, 7,8-tetrahydroquinoline -5-carboxylate		RuCl(p-cymen e)[(S,S)-Ts-DP EN]
II-d-4		(5S,8S)-methyl 5-fluoro-8-hydroxy-5,6, 7,8-tetrahydroquinoline -5-carboxylate		
II-d-5		(5S,8R)-methyl 5-fluoro-8-hydroxy-5,6, 7,8-tetrahydroquinoline -5-carboxylate		RuCl(p-cymen e)[(R,R)-Ts-D PEN]
II-d-6		(5R,8R)-methyl 5-fluoro-8-hydroxy-5,6, 7,8-tetrahydroquinoline -5-carboxylate		

[Table 35-2]

II-d-7		(5R,8S)-methyl 3,5-difluoro-8-hydroxy- 5,6,7,8-tetrahydroquino line-5-carboxylate		RuCl(p-cymen e)[(S,S)-Ts-DP EN]
II-d-8		(5S,8S)-methyl 3,5-difluoro-8-hydroxy- 5,6,7,8-tetrahydroquino line-5-carboxylate	IM II-c-3	
II-d-9		(5R,8S)-methyl 5-fluoro-8-hydroxy-3-m ethyl-5,6,7,8-tetrahydro quinoline-5-carboxylate		RuCl(p-cymen e)[(S,S)-Ts-DP EN]
II-d-10		(5S,8S)-methyl 5-fluoro-8-hydroxy-3-m ethyl-5,6,7,8-tetrahydro quinoline-5-carboxylate	IM II-c-4	

[0397] IM II-d-3

¹H NMR (CDCl₃) delta 8.64 (1 H, br dd, J = 4.9, 1.2 Hz), 7.69 (1 H, br dd, J = 7.9, 1.2 Hz), 7.31 (1 H, dd, J = 7.9, 4.9 Hz), 4.74 (1 H, m), 4.32 (1 H, br), 3.83, (3 H, s), 2.53-2.34 (3 H, m), 2.08-1.98 (1 H, m).

MS (ESI) m/z: 226.1 (M+H)⁺.

Chiral HPLC tR: 16.7 min (Method F), >99% e.e., d.e.

[0398] IM II-d-4

¹H NMR (CDCl₃) delta 8.62 (1 H, br d, J = 4.3 Hz), 7.79 (1 H, d, J = 7.9 Hz), 7.31 (1 H, dd, J = 7.9, 4.3 Hz), 4.80 (1 H, m), 4.06 (1 H, br), 3.81, (3 H, s), 2.74-2.65 (1 H, m), 2.45-2.39 (1 H, m), 2.27-2.13 (2 H, m).

MS (ESI) m/z: 226.1 (M+H)⁺.

Chiral HPLC tR: 34.6 min (Method F), >99% e.e., d.e.

[0399] IM II-d-5

¹H NMR and LCMS were identified with the IM II-d-3.

Chiral HPLC tR: 12.2 min (Method F), >99% e.e., d.e.

[0400] IM II-d-6

¹H NMR and LCMS were identified with the IM II-d-4.

Chiral HPLC tR: 13.9 min (Method F) , >99% e.e., d.e.

[0401] IM II-d-7

¹H NMR (CDCl₃) delta 8.50 (1 H, m), 7.43 (1 H, dd, J = 7.3, 2.4 Hz), 4.72 (1 H, m), 3.97 (1 H, br), 3.85 (3 H, s), 2.52-2.33 (3 H, m), 2.03 (1 H, m).

MS (ESI) m/z: 244.0 (M+H)⁺.

Chiral HPLC tR: 7.3 min (Method H) , 98.8% e.e., >99% d.e

[0402] IM II-d-8

¹H NMR (CDCl₃) delta 8.49 (1 H, m), 7.52 (1 H, dd, J = 8.5, 2.4 Hz), 4.78 (1 H, m), 3.83 (3 H, s), 3.69 (1 H, s), 2.54 (1 H, m), 2.41 (1 H, m), 2.26-2.11 (2 H, m).

MS (ESI) m/z: 244.0 (M+H)⁺.

Chiral HPLC tR: 9.6 min (Method H), 97.5% e.e., 97.1% d.e.

[0403] IM II-d-9

¹H NMR (CDCl₃) delta 8.44 (1 H, s), 7.49 (1 H, s), 4.73 (1 H, m), 4.34 (1 H, br), 3.83 (3 H, s), 2.48-2.32 (3 H, m), 2.35 (3 H, s), 2.02 (1 H, m).

MS (ESI) m/z: 238.2 (M+H)⁺.

Chiral HPLC tR: 15.4 min (Method G), 98.6% e.e., 97.0% d.e.

[0404] IM II-d-10

¹H NMR (CDCl₃) delta 8.43 (1 H, s), 7.56 (1 H, s), 4.79 (1 H, m), 3.82 (3 H, s), 2.68 (1 H, m), 2.37 (1 H, m), 2.35 (3 H, s), 2.24-2.11 (2 H, m) , 1.86 (1 H, br).

MS (ESI) m/z: 238.2 (M+H)⁺.

Chiral HPLC tR: 10.4 min (Method G), >99% e.e., 96.4% d.e.

[0405] The following Examples were prepared by General Procedure A (Table 36).

[0406]

[Table 36-1]

Examples	Structure	Chemical Name	Substrate	Amine
99		(5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
100		(5S,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
101		(5R,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
102		(5S,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
103		(5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 36-2]

104		(5S,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
105		(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
106		(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
107		(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
108		(5S,8R)-N-(2,3,4-trifluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
109		(5S,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		















[Table 36-3]

110		(5S,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
111		(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
112		(5R,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
113		(5S,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
114		(5R,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 36-4]

115		(5S,8R)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
116		(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
117		(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
118		(5S,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
119		(5S,8R)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
120		(5R,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 36-5]

121		(5S,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
122		(5S,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
123		(5S,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
124		(5S,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
125		(5S,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[0407]

[Table 37]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
99	A	1.61	383.0	114	A	1.61	419.1
101	A	1.52	403.1	115	A	1.61	419.1
102	A	1.52	403.1	116	C	1.50	386.9
105	A	1.51	369.1	117	C	1.49	386.9
106	A	1.51	369.1	118	A	1.54	413.0
107	A	1.39	353.1	119	A	1.58	383.2
108	C	1.35	354.9	120	A	1.47	369.1
109	C	1.30	337.0	121	A	1.47	369.1
110	C	1.47	386.9	122	A	1.45	349.2
111	C	1.41	352.9	123	A	1.42	371.2
112	A	1.62	437.0	124	A	1.54	403.1
113	A	1.62	437.1	125	A	1.47	349.2

[0408] Ex 99

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.6, 2.0, 1.3 Hz), 7.49 (1 H, ddd, J = 7.9, 2.0, 1.3 Hz), 7.33 (1 H, d, J = 2.0 Hz), 7.25 (1 H, dd, J = 7.9, 4.6 Hz), 7.16 (1 H, d, J = 2.0 Hz), 7.08 (1 H, br), 4.76 (1 H, dd, J = 5.3, 5.3 Hz), 4.73 (1 H, dd, J = 13.8, 5.3 Hz), 4.63 (1 H, dd, J = 13.8, 5.3 Hz), 4.35 (1 H, s), 2.66-2.23 (3 H, m), 2.48 (3 H, s), 2.00 (1 H, m).

Chiral HPLC tR: 11.5 min (Method I), >99% e.e., d.e.

[0409] Ex 100

¹H NMR and LCMS were identified with the Example 99.

Chiral HPLC tR: 15.2 min (Method I), >99% e.e., d.e.

[0410] Ex 101

¹H NMR (CDCl₃) delta 8.61 (1 H, br d, J = 4.6 Hz), 7.71 (1 H, d, J = 7.3 Hz), 7.65 (1 H, d, J = 7.9 Hz), 7.49 (1 H, ddd, J = 7.9, 2.0, 1.3 Hz), 7.41 (1 H, dd, J = 7.9, 7.3 Hz), 7.32 (1 H, br), 7.24 (1 H, dd, J = 7.9, 4.6 Hz), 4.79-4.71 (3 H, m), 4.35 (1 H, s), 2.57 (1 H, m), 2.44-2.27 (2 H, m), 2.03 (1 H, m).

Chiral HPLC tR: 20.7 min (Method K), >99% e.e., d.e.

[0411] Ex 102

¹H NMR was identified with the Example 101.

Chiral HPLC tR: 29.8 min (Method K), >99% e.e., d.e.

[0412] Ex 103

¹H NMR (CDCl₃) delta 8.60 (1 H, br d, J = 4.6 Hz), 7.65 (1 H, br), 7.45 (1 H, d, J = 2.0 Hz), 7.42 (1 H, d, J = 7.9 Hz), 7.34 (1 H, d, J = 2.0 Hz), 7.23 (1 H, dd, J = 7.9, 4.6 Hz), 4.83-4.65 (5 H, m), 4.32 (1 H, s), 3.82 (1 H, br), 2.52 (1 H, m), 2.42-2.21 (2 H,

m), 2.00 (1 H, m).

MS (ESI) m/z : 399.0 (M+H)⁺.

Chiral HPLC tR: 12.2 min (Method J), >99% e.e., d.e.

[0413] Ex 104

¹H NMR and LCMS were identified with the Example 103.

Chiral HPLC tR: 16.4 min (Method J), >99% e.e., d.e.

[0414] Ex 107

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.49 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.41 (1 H, dd, J = 8.6, 6.1 Hz), 7.25-7.22 (1 H, br), 7.24 (1 H, dd, J = 7.9, 4.9 Hz), 7.19 (1 H, dd, J = 8.6, 2.4 Hz), 7.00 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 4.74 (1 H, m), 4.65 (1 H, dd, J = 14.7, 6.1 Hz), 4.59 (1 H, dd, J = 14.7, 6.1 Hz), 4.46 (1 H, br s), 2.61-2.28 (3 H, m), 2.02 (1 H, m).

[0415] Ex 108

¹H NMR (CDCl₃) delta 8.61 (1 H, br d, J = 4.9 Hz), 7.48 (1 H, br d, J = 7.9 Hz), 7.26 (1 H, dd, J = 7.9, 4.9 Hz), 7.23 (1 H, br), 7.12 (1 H, m), 6.98 (1 H, m), 4.75 (1 H, m), 4.65 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 4.40 (1 H, br s), 2.61-2.28 (3 H, m), 2.02 (1 H, m).

[0416] Ex 109

¹H NMR (CDCl₃) delta 8.60 (1 H, br d, J = 4.9 Hz), 7.48 (1 H, br d, J = 7.9 Hz), 7.36 (1 H, m), 7.26-7.23 (1 H, br), 7.24 (1 H, dd, J = 7.9, 4.9 Hz), 6.91-6.84 (2 H, m), 4.75 (1 H, m), 4.62 (1 H, dd, J = 14.7, 6.1 Hz), 4.53 (1 H, dd, J = 14.7, 5.5 Hz), 4.49 (1 H, br s), 2.61-2.27 (3 H, m), 2.02 (1 H, m).

[0417] Ex 111

¹H NMR (CDCl₃) delta 8.60 (1 H, br d, J = 4.9 Hz), 7.48 (1 H, br d, J = 7.9 Hz), 7.32 (1 H, dd, J = 7.9, 7.9 Hz), 7.25 (1 H, dd, J = 7.9, 4.9 Hz), 7.21 (1 H, br), 7.16-7.14 (2 H, m), 4.74 (1 H, m), 4.62 (1 H, dd, J = 15.3, 6.1 Hz), 4.54 (1 H, dd, J = 15.3, 5.5 Hz), 4.43 (1 H, br s), 2.61-2.27 (3 H, m), 2.01 (1 H, m).

[0418] Ex 116

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.52 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.29 (1 H, dd, J = 1.8, 1.8 Hz), 7.25 (1 H, dd, J = 7.9, 4.9 Hz), 7.14-7.08 (1 H, br), 7.11 (1 H, ddd, J = 9.2, 1.8, 1.8 Hz), 4.79-4.63 (3 H, m), 4.46 (1 H, br), 2.61-2.26 (3 H, m), 2.02 (1 H, m).

[0419] Ex 117

¹H NMR was identified with the Example 116.

[0420] EX 119

¹H NMR (CDCl₃) delta 8.57 (1 H, br d, J = 4.9 Hz), 7.44 (1 H, d, J = 1.8 Hz), 7.43 (1 H, br d, J = 7.9 Hz), 7.37 (1 H, d, J = 8.6 Hz), 7.30 (1 H, dd, J = 8.6, 1.8 Hz), 7.20 (1 H, dd, J = 7.9, 4.9 Hz), 7.17 (1 H, br), 5.45 (1 H, m), 4.73 (1 H, m), 4.38 (1 H, br),

2.62-2.30 (3 H, m), 2.02 (1 H, m), 1.59 (3 H, d, J = 7.3 Hz).

[0421] The following Examples and Intermediates were prepared by General Procedure N (Tables 38 and 40).

[0422] General Procedure N

A mixture of substrate (1.0 eq.) and 2 N aq. NaOH (2.0 eq.) in MeOH was stirred at room temperature for 1.5 h, 2 N hydrochloric acid (2.2 eq.) was added to the mixture. The mixture was concentrated in vacuo. Toluene and MeCN was added to the resulting mixture and concentrated in vacuo. This procedure was repeated 3 times to remove remaining water. The residual powder was dissolved with THF and amine (1.1 eq.), triethylamine (1.3 eq.), and DMT-MM (1.8 eq.) were added to the mixture at ambient temperature. After being stirred until complete reaction, the insoluble material was removed by filtration, and the filtrate was concentrated in vacuo. The resulting residue was purified by silica gel column chromatography, and/or SCX cartridge column, and then preparative HPLC to afford the following Examples and Intermediates.

[0423]

[Table 38-1]

Examples	Structure	Chemical Name	Substrate	Amine
126		(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
127		(5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
128		(5R,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
129		(5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
130		(5R,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-2]

131		(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
132		(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
133		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
134		(5R,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
135		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
136		(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-3]

137		(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
138		(5R,8R)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
139		(5S,8S)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
140		(5R,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
141		(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-4]

142		(5S,8S)-N-(2-chloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
143		(5R,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
144		(5S,8S)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
145		(5R,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
146		(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-5]

147		(5R,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
148		(5S,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
149		(5S,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
150		(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
151		(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-6]

152		(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
153		(5R,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
154		(5S,8S)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
155		(5R,8R)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
156		(5R,8R)-N-(4-chloro-2-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-7]

157		(5S,8S)-N-(4-chloro-2-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
158		(5R,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
159		(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
160		(5R,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
161		(5S,8S)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-8]

162		(5R,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
163		(5S,8S)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
164		(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
165		(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
166		(5R,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-9]

167		(5S,8S)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-4
168		(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1-(2,3,4-trichlorophenyl)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-4
169		(5S,8S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-4
170		(5S,8S)-N-(2-chloro-3,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-4
171		(5S,8S)-N-(2-chloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-4

[Table 38-10]

172		(5S,8S)-N-(2-chloro-4-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
173		(5S,8S)-N-(2,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
174		(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
175		(5S,8S)-5-fluoro-N-(2-fluoro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
176		(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
177		(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-11]

178		(5S,8S)-5-fluoro-N-(3-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
179		(5S,8S)-N-(2-chloro-6-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
180		(5S,8S)-5-fluoro-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
181		(5S,8S)-N-(5-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
182		(5S,8S)-N-(4-bromo-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-12]

183		(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
184		(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
185		(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
186		(5S,8S)-5-fluoro-N-(2-fluoro-6-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
187		(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-13]

188		(5S,8S)-N-(3-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
189		(5S,8S)-5-fluoro-8-hydroxy-N-((R)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
190		(5S,8S)-N-(3-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
191		(5S,8S)-N-(2,4-dichlorophenethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
192		(5S,8S)-5-fluoro-8-hydroxy-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
















[Table 38-14]

193		(5S,8S)-N-(3-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
194		(5S,8S)-N-((3-chloro-5-(trifluoromethyl)pyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
195		(5S,8S)-N-(2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
196		(5S,8S)-N-((R)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
197		(5S,8S)-N-(3,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-15]

198		(5S,8S)-N-(2-chloro-6-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
199		(5S,8S)-5-fluoro-8-hydroxy-N-(2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
200		(5S,8S)-5-fluoro-8-hydroxy-N-(3,4,5-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
201		(5S,8S)-N-(4-cyano-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
202		(5S,8S)-N-(3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
















[Table 38-16]

203		(5S,8S)-N-(3-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
204		(5S,8S)-N-(2-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
205		(5S,8S)-N-(3-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
206		(5S,8S)-N-(4-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
207		(5S,8S)-5-fluoro-N-(2-fluoro-4-(trifluoromethoxy)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-17]

208		(5S,8S)-N-(2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
209		(5S,8S)-N-(2-chloro-5-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
210		(5S,8S)-N-(4-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
211		(5S,8S)-5-fluoro-8-hydroxy-N-(4-methoxy-2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
212		(5S,8S)-N-(3,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-18]

213		(5S,8S)-N-((4-(4-chlorophenyl)thiazol-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-d-4	
214		(5S,8S)-5-fluoro-8-hydroxy-N-(2-(morpholinomethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-d-4	
215		(5S,8S)-5-fluoro-8-hydroxy-N-((1S,2R)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-d-4	
216		(5S,8S)-N-(6-chloro-2-fluoro-3-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-d-4	
217		(5S,8S)-N-(2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-d-4	

[Table 38-19]

218		(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
219		(5S,8S)-5-fluoro-8-hydroxy-N-(2-(3-(trifluoromethyl)phenoxy)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
220		(5S,8S)-5-fluoro-N-((1-(4-fluorophenyl)cyclopropyl)methyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
221		(5S,8S)-N-(3,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
222		(5S,8S)-5-fluoro-8-hydroxy-N-((1R,2S)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
223		(5S,8S)-5-fluoro-N-(2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-20]

224		(5S,8S)-N-(2-chloro-3-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
225		(5S,8S)-N-((1S,2S)-2-(benzyloxy)cyclopentyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
226		(5S,8S)-N-((S)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
227		(5S,8S)-N-(3,3-dimethylbutyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
228		(5S,8S)-5-fluoro-8-hydroxy-N-(2-phenoxyethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
229		(5S,8S)-N-(4,6-dichloro-2,3-dihydrobenzofuran-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-21]

230		(5S,8S)-N-(5,7-dichlorochroman-4-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
231		(5S,8S)-N-(1-(adamantan-1-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
232		(5S,8S)-N-(2-(4-chlorophenyl)-2-(4,4-difluoropiperidin-1-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
233		(5S,8S)-N-(chroman-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
234		(5S,8S)-N-(2-(4-chlorophenyl)propyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-22]

235		(5S,8S)-5-fluoro-8-hydroxy-N-((rac)-2-morpholino-2-phenylethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
236		(5S,8S)-N-(2-(4,4-difluoropiperidin-1-yl)-2-(4-methylthiazol-5-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
237		(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
238		(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
239		(5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 38-23]

240		(5S,8S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-4	
241		(5S,8S)-N-(2,4-dichlorobenzyl)-3,5-difluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-8	
242		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-3,5-difluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-8	
243		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-10	
244		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM II-d-10	

[0424]

[Table 39-1]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
127	A	1.59	383.0	187	A	1.45	387.0
128	A	1.50	403.1	188	B	1.31	353.1
129	A	1.50	403.1	189	C	1.46	341.1
132	A	1.49	369.1	190	A	1.37	353.1
133	A	1.49	369.1	191	B	1.46	383.0
134	A	1.38	353.1	192	C	1.44	392.2
135	A	1.38	353.1	193	B	1.30	371.0
136	C	1.34	355.0	194	C	1.43	404.0
137	B	1.27	355.1	195	A	1.34	335.0
138	A	1.41	369.1	196	C	1.38	327.1
139	A	1.41	369.1	197	A	1.47	369.0
140	C	1.29	337.1	198	A	1.40	365.2
141	B	1.22	337.1	199	B	1.35	369.1
142	A	1.45	367.2	200	C	1.35	355.1
143	C	1.45	387.0	201	B	1.12	344.1
144	A	1.45	387.0	202	A	1.30	337.1
145	C	1.40	353.0	203	B	1.33	353.1
146	A	1.40	353.0	204	A	1.36	353.2
147	A	1.60	437.1	205	A	1.35	335.1
148	A	1.60	437.0	206	B	1.32	353.1
149	A	1.58	419.1	207	B	1.43	403.0
150	D	1.26	383.1	208	A	1.28	337.1
151	C	1.48	386.9	209	A	1.50	403.1
152	B	1.41	387.0	210	B	1.29	335.1
153	A	1.52	413.0	211	B	1.39	399.1
154	A	1.52	413.0	212	B	1.43	369.0
155	A	1.56	383.2	213	C	1.52	418.0
156	C	1.55	403.0	214	C	1.36	400.1
157	B	1.49	403.0	215	C	1.35	327.1
158	A	1.45	369.1	216	A	1.45	367.2
159	A	1.46	369.1	217	B	1.18	337.1

[Table 39-2]

160	A	1.44	349.2	218	C	1.46	341.1
161	A	1.44	349.2	219	C	1.48	399.1
162	A	1.41	371.2	220	C	1.44	359.1
163	A	1.41	371.2	221	B	1.23	337.1
164	A	1.51	403.1	222	C	1.35	327.1
165	A	1.52	403.0	223	B	1.18	319.1
166	A	1.46	349.2	224	B	1.24	365.0
167	A	1.46	349.2	225	C	1.48	385.2
168	A	1.64	417.1	226	C	1.38	327.1
169	A	1.36	353.2	227	C	1.40	295.2
170	A	1.35	371.2	228	C	1.29	331.1
171	A	1.34	353.1	229	C	1.49	397.1
172	A	1.38	365.2	230	C	1.55	411.0
173	A	1.45	369.1	231	C	1.78	373.2
174	B	1.34	370.9	232	D	1.69	468.1
175	A	1.44	387.0	233	C	1.34	343.1
176	C	1.37	370.9	234	C	1.48	363.1
177	B	1.29	370.0	235	C	1.23	400.2
178	C	1.42	387.1	236	D	1.30	455.2
179	A	1.46	403.0	237	D	1.50	367.1
180	B	1.22	355.1	238	D	1.68	395.1
181	A	1.48	413.0	239	D	1.51	367.1
182	B	1.35	397.0	240	D	1.57	453.1
183	C	1.42	370.9	241	C	1.56	387.0
184	C	1.40	370.9	242	C	1.45	371.1
185	C	1.41	370.9	243	D	1.67	383.1
186	B	1.33	387.0	244	D	1.54	367.1

[0425] Ex 126

¹H NMR (CDCl₃) delta 8.59 (1 H, d, J = 4.6 Hz), 7.53 (1 H, d, J = 7.9 Hz), 7.31 (1 H, br s), 7.25 (1 H, m), 7.14 (1 H, br s), 6.98 (1 H, br d, J = 5.3 Hz), 4.80 (1 H, dd, J = 5.3, 5.3 Hz), 4.70 (1 H, dd, J = 13.8, 5.9 Hz), 4.60 (1 H, dd, J = 13.8, 5.9 Hz), 3.79 (1 H, br s), 2.65 (1 H, m), 2.46 (3 H, s), 2.40-2.28 (2 H, m), 2.16 (1 H, m).

MS (ESI) m/z: 382.9 (M+H)⁺.

Chiral HPLC tR: 13.6 min (Method I), 99.0% e.e., >99% d.e.

[0426] Ex 127

¹H NMR was identified with the Example 126.

Chiral HPLC tR: 17.0 min (Method I), 98.2% e.e., >99% d.e.

[0427] Ex 128

¹H NMR (CDCl₃) delta 8.58 (1 H, ddd, J = 4.6, 2.0, 1.3 Hz), 7.70 (1 H, d, J = 7.9 Hz), 7.64 (1 H, d, J = 7.2 Hz), 7.54 (1 H, d, J = 7.9 Hz), 7.39 (1 H, dd, J = 7.9, 7.2 Hz), 7.24 (1 H, br), 7.23 (1 H, dd, J = 7.9, 4.6 Hz), 4.81 (1 H, dd, J = 5.3, 4.6 Hz), 4.71 (2 H, d, J = 5.9 Hz), 3.87 (1 H, br), 2.68 (1 H, m), 2.42-2.08 (3 H, m).

Chiral HPLC tR: 35.1 min (Method K), >99% e.e., d.e.

[0428] Ex 129

¹H NMR was identified with the Example 128.

Chiral HPLC tR: 23.9 min (Method K), >99% e.e., d.e.

[0429] Ex 130

¹H NMR (DMSO d₆) delta 8.69 (1 H, br), 8.61 (1 H, d, J = 4.6 Hz), 7.56 (1 H, d, J = 2.0 Hz), 7.55 (1 H, d, J = 7.9 Hz), 7.48 (1 H, d, J = 2.0 Hz), 7.38 (1 H, dd, J = 7.9, 4.6 Hz), 5.54 (1 H, t, J = 5.3 Hz), 5.43 (1 H, d, J = 4.0 Hz), 4.70 (2 H, d, J = 5.3 Hz), 4.63 (1 H, d, J = 4.0 Hz), 4.50 (2 H, m), 2.76-2.55 (1 H, m), 2.14-1.90 (3 H, m).

MS (ESI) m/z: 398.9 (M+H)⁺.

Chiral HPLC tR: 14.1 min (Method J), 98.2% e.e., >99% d.e.

[0430] Ex 131

¹H NMR and MS were identified with the Example 130.

Chiral HPLC tR: 39.1 min (Method J), >99% e.e., d.e.

[0431] Ex 132

¹H NMR (CDCl₃) delta 8.59 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.54 (1 H, dd, J = 7.9, 1.2 Hz), 7.45 (1 H, d, J = 1.8 Hz), 7.36 (1 H, d, J = 7.9 Hz), 7.31-7.23 (2 H, m), 7.18 (1 H, br), 4.80 (1 H, ddd, J = 6.7, 6.0, 1.2 Hz), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 3.79 (1 H, br s), 2.67 (1 H, m), 2.41-2.25 (2 H, m), 2.15 (1 H, m).

[0432] Ex 133

¹H NMR was identified with the Example 132.

[0433] Ex 134

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.54 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.41 (1 H, dd, J = 8.6, 6.1 Hz), 7.25 (1 H, dd, J = 7.9, 4.9 Hz), 7.19 (1 H, dd, J = 7.9, 2.4 Hz), 7.13 (1 H, br d, J = 5.5 Hz), 7.00 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 4.81 (1 H, dd, J = 5.5, 4.9 Hz), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 5.5 Hz), 3.74 (1 H, br s), 2.67 (1 H, m), 2.41-2.25 (2 H, m), 2.15 (1 H, m).

[0434] Ex 135

¹H NMR was identified with the Example 134.

[0435] Ex 150

¹H NMR (CDCl₃) delta 8.58 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.52 (1 H, br d, J = 4.9 Hz), 7.48 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.23 (1 H, dd, J = 7.9, 4.9 Hz), 7.17 (1 H, dd,

J = 7.9, 2.4 Hz), 7.09 (1 H, dd, J = 8.6, 2.4 Hz), 4.79-4.73 (4 H, m), 4.65 (1 H, dd, J = 14.7, 6.1 Hz), 3.82 (1 H, br), 3.67 (1 H, br), 2.61 (1 H, m), 2.37-2.21 (2 H, m), 2.11 (1 H, m).

Ex 158

¹H NMR (CDCl₃) delta 8.61 (1 H, br dd, J = 4.9, 1.6 Hz), 7.56 (1 H, d, J = 7.9 Hz), 7.41 (1 H, dd, J = 7.9, 1.6 Hz), 7.34 (1 H, dd, J = 7.6, 1.3 Hz), 7.28-7.15 (3 H, m), 4.81 (1 H, dd, J = 6.3, 4.9 Hz), 4.70 (1 H, dd, J = 14.5, 5.9 Hz), 4.63 (1 H, dd, J = 14.5, 6.3 Hz), 3.62 (1 H, s), 2.66 (1 H, m), 2.44-2.08 (3 H, m).

[0436] Ex 159

¹H NMR was identified with the Example 158.

[0437] Ex 191

¹H NMR (CDCl₃) delta 8.58 (1 H, m), 7.42 (1 H, d, J = 1.8 Hz), 7.39 (1 H, d, J = 7.9 Hz), 7.25-7.20 (2 H, m), 7.19 (1 H, d, J = 7.9 Hz), 6.79 (1 H, br), 4.80 (1 H, ddd, J = 6.4, 5.2, 1.2 Hz), 3.80 (1 H, br), 3.75-3.59 (2 H, m), 3.03 (2 H, dt, J = 6.7, 1.8 Hz), 2.62 (1 H, m), 2.38-2.22 (2 H, m), 2.10 (1 H, m).

[0438] Ex 192

¹H NMR (CDCl₃) delta 8.61 (1 H, br d, J = 4.9 Hz), 7.60 (1 H, d, J = 7.9 Hz), 7.28 (2 H, dd, J = 7.9, 4.9 Hz), 4.83 (1 H, dd, J = 6.1, 5.5 Hz), 3.76 (1 H, br), 3.69 (4 H, dd, J = 4.9, 4.3 Hz), 3.48 (1 H, dd, J = 14.1, 6.1 Hz), 3.41 (1 H, dd, J = 14.1, 5.5 Hz), 2.70 (1 H, m), 2.63 (4 H, dd, J = 4.9, 4.3 Hz), 2.38 (1 H, m), 2.28 (1 H, m), 2.16 (1 H, m), 1.70-1.52 (5 H, m), 1.47-1.37 (4 H, m), 1.24 (1 H, m).

[0439] Ex 237

¹H NMR (CDCl₃) delta 8.56 (1 H, br dd, J = 4.3, 1.8 Hz), 7.50 (1 H, br dd, J = 7.9, 1.8 Hz), 7.39 (1 H, dd, J = 8.6, 6.1 Hz), 7.21 (1 H, dd, J = 7.9, 4.3 Hz), 7.16 (1 H, dd, J = 7.9, 3.1 Hz), 7.08 (1 H, m), 7.02 (1 H, ddd, J = 8.6, 7.9, 3.1 Hz), 5.43 (1 H, m), 4.81 (1 H, br t, J = 5.8 Hz), 2.90 (1 H, br), 2.69 (1 H, m), 2.41-2.26 (2 H, m), 2.17 (1 H, m), 1.60 (3 H, d, J = 7.3 Hz).

[0440] Ex 238

¹H NMR (CDCl₃) delta 8.63 (1 H, m), 7.62 (1 H, br d, J = 7.3 Hz), 7.40 (0.5 H, d, J = 1.8 Hz), 7.39 (0.5 H, d, J = 1.8 Hz), 7.30 (0.5 H, dd, J = 7.9, 4.3 Hz), 7.29 (0.5 H, dd, J = 7.9, 4.3 Hz), 7.180 (0.5 H, dd, J = 8.6, 1.8 Hz), 7.179 (0.5 H, dd, J = 8.6, 1.8 Hz), 7.12 (0.5 H, d, J = 8.6 Hz), 7.11 (0.5 H, d, J = 8.6 Hz), 6.98 (1 H, br), 4.82 (1 H, br), 3.68 (1 H, br), 3.05 (1 H, m), 2.69 (1 H, m), 2.42-2.25 (3 H, m), 2.17 (1 H, m), 1.41-1.24 (2 H, m).

[0441] Ex 239

¹H NMR (CDCl₃) delta 8.61 (1 H, br dd, J = 4.9, 1.8 Hz), 7.65 (1 H, br dd, J = 7.9, 1.8 Hz), 7.35 (1 H, dd, J = 8.6, 6.1 Hz), 7.30 (1 H, dd, J = 7.9, 4.9 Hz), 7.16 (1 H, dd, J = 8.6, 2.4 Hz), 7.08 (1 H, br), 7.02 (1 H, ddd, J = 8.6, 6.1, 2.4 Hz), 5.41 (1 H, m), 4.80

(1 H, br), 3.75 (1 H, br), 2.60 (1 H, m), 2.37-2.07 (3 H, m), 1.60 (3 H, d, $J = 7.3$ Hz).

[0442] Ex 240

^1H NMR (CDCl_3) delta 8.58 (1 H, br dd, $J = 4.9, 1.8$ Hz), 7.47 (1 H, d, $J = 1.8$ Hz), 7.33 (1 H, dd, $J = 8.6, 1.8$ Hz), 7.29 (1 H, d, $J = 8.6$ Hz), 7.21 (1 H, dd, $J = 7.9, 4.9$ Hz), 7.15 (1H, br d, $J = 7.9$ Hz), 6.32 (1 H, br), 4.75 (1 H, br dd, $J = 5.5, 4.9$ Hz), 4.18 (1 H, dd, $J = 14.1, 6.1$ Hz), 3.95 (1 H, dd, $J = 14.1, 6.1$ Hz), 3.92-3.83 (2H, m), 3.74-3.55 (3 H, m), 2.49 (1 H, m), 2.40-2.55 (3 H, m), 2.24-2.08 (3 H, m), 2.01 (1 H, m).

[0443] [Table 40]

Intermediates	Structure	Chemical Name	Substrate	Amine
I-d-1		(5S,8S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-d-2		(5S,8S)-N-(2,3-dichloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-d-3		(5S,8S)-N-(2,4-dichloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-d-4		(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-d-5		(5S,8S)-N-((S)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[0444] IM I-d-1

MS (ESI) m/z : 383.2 ($\text{M}+\text{H}$) $^+$.

[0445] IM I-d-2

MS (ESI) m/z : 386.9 ($\text{M}+\text{H}$) $^+$.

[0446] IM I-d-3

MS (ESI) m/z: 386.9 (M+H)⁺.

[0447] IM I-d-4

MS (ESI) m/z: 403.0 (M+H)⁺.


[0448] IM I-d-5

MS (ESI) m/z: 383.2 (M+H)⁺.

[0449] Intermediate (IM) I-d-6:

(5S,8S)-N-(2-(((tert-butyldimethylsilyl)oxy)methyl)-6-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide

[Chem.27]



[0450] To a solution of

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide (174 mg, 0.455 mmol, Ex 150) in CH₂Cl₂ (1.0 mL) was added imidazole (93 mg, 1.364 mmol) at 0 °C. After 10 min stirring, TBSCl (82 mg, 0.545 mmol) was added to the mixture at 0 °C. Then the mixture was allowed to warm to room temperature and stirred for 3 h. The mixture was diluted with cold water and extracted with CH₂Cl₂. The extracts were washed with brine, dried over Na₂SO₄, and then concentrated in vacuo. The resulting residue was purified by silica gel column chromatography (30- to 100% EtOAc/n-hexane, gradient) to afford 165 mg (73%) of the title compound.

[0451] IM I-d-6


¹H NMR (CDCl₃) delta 8.58 (1 H, br d, J = 4.9 Hz), 7.55 (1 H, br d, J = 7.9 Hz), 7.24 (1 H, dd, J = 7.9, 4.9 Hz), 7.15 (1 H, dd, J = 9.8, 3.1 Hz), 7.12 (1 H, dd, J = 8.6, 3.1 Hz), 7.05 (1 H, br d, J = 5.5 Hz), 4.89 (1 H, d, J = 13.4 Hz), 4.82 (1 H, d, J = 13.4 Hz), 4.80 (1 H, m), 4.70 (1 H, dd, J = 14.1, 6.1 Hz), 4.57 (1 H, dd, J = 14.1, 5.5 Hz), 3.80 (1 H, br), 2.66 (1 H, m), 2.39-2.25 (2 H, m), 2.12 (1 H, m), 0.94 (9 H, s), 0.12 (3 H, s), 0.11 (3 H, s).

MS (ESI) m/z: 496.8 (M+H)⁺.

[0452] Example 245:

(5S,8S)-5-((2,4-dichlorobenzyl)carbamoyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline 1-oxide

[0453] [Chem.28]



[0454] To a stirred solution of (5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide (30 mg, 0.081 mmol, Ex 133) in CH_2Cl_2 (2 mL) was added mCPBA (17 mg, 0.099 mmol) at 0 °C. The mixture was stirred at 0 °C for 2 h, and then warmed to room temperature. After being stirred at room temperature for 16 h, aq. $\text{Na}_2\text{S}_2\text{O}_3$ was added to the mixture. The resulting mixture was stirred for 30 min vigorously, and then extracted with EtOAc. The extract was washed with aq. NaHCO_3 , water, and brine. The extract was dried over Na_2SO_4 and concentrated in vacuo. The resulting solid was washed with EtOAc to afford 22 mg (70%) of the title compound.

[0455] Ex 245

^1H NMR (CDCl_3) δ 8.27 (1 H, ddd, $J = 4.9, 1.8, 1.2$ Hz), 7.46 (1 H, d, $J = 1.8$ Hz), 7.38 (1 H, d, $J = 7.9$ Hz), 7.28-7.25 (2 H, m), 7.16-7.14 (2 H, m), 5.23 (1 H, br), 5.15 (1 H, br), 4.65 (1 H, dd, $J = 14.7, 6.1$ Hz), 4.59 (1 H, dd, $J = 14.7, 6.1$ Hz), 2.61 (1 H, m), 2.33-2.28 (2 H, m), 2.11 (1 H, m).

LCMS (ESI) m/z : 385.0 ($\text{M}+\text{H}$) $^+$, t_R 1.40 min (Method D).


[0456] Intermediate (IM) II-e-2-6:

(5R)-Methyl 5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate

Intermediate (IM) II-e-2-7:

(5S)-Methyl 5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate

[0457] [Chem.29]



[0458] The title compounds were prepared by chiral HPLC separation of methyl

5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate (IM II-e-2-1).

[0459] IM II-e-2-6

Chiral HPLC tR: 15.1 min (Method L), >98% e.e.

[0460] IM II-e-2-7

Chiral HPLC tR: 17.8 min (Method L), >98% e.e.

[0461] The following Intermediates were prepared by General Procedure A (Table 41),
except Intermediate I-e-35.

[0462] The Intermediate I-e-35 was prepared by General Procedure N (Table 41).

[0463] [Table 41-1]

Intermediates	Structure	Chemical Name	Substrate	Amine
I-e-25		(R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-26		(R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-27		(S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-28		(S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-29		(S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 41-2]

I-e-30		(S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-31		(S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-32		(S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-33		(S)-5-fluoro-8-methylene-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-34		(S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		
I-e-35		(S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[0464] IM I-e-25

¹H NMR and MS were identified with the IM I-e-3.

[0465] IM I-e-26













¹H NMR and MS were identified with the IM I-e-19.

[0466] IM I-e-27

¹H NMR and MS were identified with the IM I-e-3.

- [0467] IM I-e-28
¹H NMR and MS were identified with the IM I-e-19.
- [0468] IM I-e-29
MS (ESI) m/z: 384.6 (M+H)⁺.
- [0469] IM I-e-30
¹H NMR (CDCl₃) delta 8.61 (1 H, br dd, J = 4.3, 1.8 Hz), 7.48 (1 H, br dd, J = 7.9, 1.8 Hz), 7.24 (1 H, br), 7.24-7.17 (2 H, m), 7.10 (1 H, m), 6.35 (1 H, s), 5.31 (1 H, s), 4.66 (1 H, dd, J = 15.3, 6.1 Hz), 4.62 (1 H, dd, J = 15.3, 6.1 Hz), 2.88-2.82 (2 H, m), 2.47 (1 H, m), 2.27 (1 H, m).
MS (ESI) m/z: 367.1 (M+H)⁺.
- [0470] IM I-e-31
¹H NMR and MS were identified with the IM I-e-20.
- [0471] IM I-e-32
MS (ESI) m/z: 348.8 (M+H)⁺.
- [0472] IM I-e-33
¹H NMR (CDCl₃) delta 8.61 (1 H, br dd, J = 4.9, 1.8 Hz), 7.46 (1 H, br dd, J = 7.9, 1.8 Hz), 7.21-7.10 (3 H, m), 6.98 (1 H, m), 6.36 (1 H, s), 5.31 (1 H, s), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.58 (1 H, dd, J = 14.7, 6.1 Hz), 2.90-2.79 (2 H, m), 2.47 (1 H, m), 2.26 (1 H, m).
MS (ESI) m/z: 350.8 (M+H)⁺.
- [0473] IM I-e-34
¹H NMR and MS were identified with the IM I-e-18.
- [0474] IM I-e-35
¹H NMR and MS were identified with the IM I-e-17.
- [0475] The following Examples and Intermediates were prepared by General Procedure B or O (Tables 42 and 44).
- [0476] General Procedure O
To a solution of substrate (1.0 eq.) in CH₂Cl₂ was added Dess-Martin Periodinane (1.5 eq.) at ambient temperature. After being stirred until complete reaction, aq. Na₂S₂O₃ and aq. NaHCO₃ were added to the mixture. The mixture was extracted with CH₂Cl₂ and washed with water. The extract was dried over Na₂SO₄ and concentrated in vacuo to afford a solid. The residual solid was purified by silica gel column chromatography and preparative HPLC to afford following Examples and Intermediates.
- [0477]

[Table 42-1]

Examples	Structure	Chemical Name	Substrate	General Procedure
246		(R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM 1-e-25	B
247		(S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM 1-e-27	B
248		(S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM 1-e-28	B
249		(S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 Ex 107	O
250		(S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 Ex 169	O
251		(S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide	 Ex 117	O

[Table 42-2]

252		(S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		B
253		(S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
254		(S)-5-fluoro-8-oxo-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
255		(S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O

[0478] [Table 43]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
246	A	1.51	401.1	251	B	1.44	384.9
247	A	1.51	401.1	252	B	1.40	366.9
248	B	1.44	367.0	253	B	1.49	400.9
249	B	1.33	351.0	254	C	1.35	353.0
250	B	1.31	351.0	255	D	1.43	368.0

[0479] Ex 246

¹H NMR was identified with the Example 13.

[0480] Ex 247

¹H NMR was identified with the Example 13.

[0481] Ex 248

¹H NMR (DMSO d₆) delta 9.35 (1 H, br), 8.85 (1 H, d, J = 4.9 Hz), 7.95 (1 H, dd, J = 7.9, 1.8 Hz), 7.73 (1 H, dd, J = 7.9, 4.9 Hz), 7.62 (1 H, d, J = 1.8 Hz), 7.44 (1 H, dd, J = 7.9, 1.8 Hz), 7.34 (1 H, d, J = 7.9 Hz), 4.44 (1 H, dd, J = 15.9, 6.1 Hz), 4.39 (1 H, dd,

J = 15.9, 5.5 Hz), 2.89-2.85 (2 H, m), 2.77-2.58 (2 H, m).

[0482] Ex 249

¹H NMR (DMSO d6) delta 9.31 (1 H, br), 8.85 (1 H, dd, J = 4.9, 1.8 Hz), 7.95 (1 H, dd, J = 7.9, 1.8 Hz), 7.73 (1 H, dd, J = 7.9, 4.9 Hz), 7.45 (1 H, dd, J = 8.6, 3.1 Hz), 7.38 (1 H, dd, J = 8.6, 6.1 Hz), 7.23 (1 H, ddd, J = 8.6, 8.6, 3.1 Hz), 4.44 (1 H, dd, J = 15.9, 6.1 Hz), 4.39 (1 H, dd, J = 15.9, 5.5 Hz), 2.94-2.81 (2 H, m), 2.77-2.54 (2 H, m).

[0483] Ex 252

¹H NMR (CDCl₃) delta 8.90 (1 H, dd, J = 4.6, 1.3 Hz), 7.75 (1 H, dd, J = 7.9, 1.3 Hz), 7.52 (1 H, dd, J = 7.9, 4.6 Hz), 7.48 (1 H, dd, J = 7.9, 2.0 Hz), 7.33 (1 H, dd, J = 7.9, 2.0 Hz), 7.23-7.20 (1 H, br), 7.22 (1 H, dd, J = 7.9, 7.9 Hz), 4.68 (2 H, d, J = 5.9 Hz), 3.17 (1 H, m), 3.02 (1 H, m), 2.81 (1 H, m), 2.58 (1 H, m).

[0484] Ex 253

¹H NMR (DMSO d6) delta 9.35 (1 H, br), 8.84 (1 H, d, J = 4.3 Hz), 7.90 (1 H, dd, J = 7.9, 1.2 Hz), 7.71 (1 H, dd, J = 7.9, 4.3 Hz), 7.29 (1 H, m), 7.19 (1 H, m), 4.43 (1 H, dd, J = 15.3, 5.5 Hz), 4.39 (1 H, dd, J = 15.3, 5.5 Hz), 2.88-2.84 (2 H, m), 2.73-2.56 (2 H, m).

[0485]

[Table 44-1]

Intermediates	Structure	Chemical Name	Substrate	General Procedure
I-c-13		(S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-14		(S)-N-(2,3-dichloro-4-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-15		(S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-16		(S)-N-(2,4-dichloro-3-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-17		(S)-N-(2,4-difluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O

[Table 44-2]

I-c-18		(S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-19		(S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-20		(S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-21		(S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		B
I-c-22		(S)-N-(2-(((tert-butyl(dimethylsilyl)oxy)methyl)-6-chloro-4-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O

[Table 44-3]

I-c-23		(S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-d-1	O
I-c-24		(S)-N-((S)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-d-5	O
I-c-25		(S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		Ex 237	O
I-c-26		(S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		Ex 239	O
I-c-27		(S)-N-(2,4-dichlorophenylethyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		Ex 191	O

[Table 44-4]

I-c-28		(S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-29		(S)-5-fluoro-N-((1-morpholinocyclohexyl)methyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-30		(S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-31		(S)-N-(2,4-dichlorobenzyl)-5-fluoro-3-methyl-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O
I-c-32		(S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-3-methyl-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide		O

[0486] IM I-c-13
MS (ESI) m/z: 350.8 (M+H)⁺.

[0487] IM I-c-14
MS (ESI) m/z: 384.8 (M+H)⁺.

[0488] IM I-c-15
MS (ESI) m/z: 368.8 (M+H)⁺.

[0489] IM I-c-16
MS (ESI) m/z: 384.8 (M+H)⁺.

[0490] IM I-c-17
MS (ESI) m/z: 334.8 (M+H)⁺.

[0491] IM I-c-18
MS (ESI) m/z: 368.8 (M+H)⁺.

- [0492] IM I-c-19
MS (ESI) m/z: 368.8 (M+H)⁺.
- [0493] IM I-c-20
MS (ESI) m/z: 368.8 (M+H)⁺.
- [0494] IM I-c-21
¹H NMR and MS were identified with the Ex 14.
- [0495] IM I-c-22
¹H NMR (CDCl₃) delta 8.88 (1 H, ddd, J = 4.3, 1.8, 1.2 Hz), 7.74 (1 H, br d, J = 7.9 Hz), 7.50 (1 H, dd, J = 7.9, 4.3 Hz), 7.14 (3 H, br d, J = 7.9 Hz), 4.88 (1 H, d, J = 13.4 Hz), 4.82 (1 H, d, J = 13.4 Hz), 4.71 (1 H, dd, J = 14.1, 5.5 Hz), 4.62 (1 H, dd, J = 14.1, 5.5 Hz), 3.16 (1 H, m), 2.99 (1 H, m), 2.79 (1 H, m), 2.54 (1 H, m), 0.94 (9 H, s), 0.12 (3 H, s), 0.11 (3 H, s).
MS (ESI) m/z: 494.8 (M+H)⁺.
- [0496] IM I-c-23
¹H NMR (CDCl₃) delta 8.87 (1 H, ddd, J = 4.3, 1.8, 1.2 Hz), 7.67 (1 H, dd, J = 7.9, 2.4 Hz), 7.48 (1 H, dd, J = 7.9, 4.3 Hz), 7.43 (1 H, d, J = 1.8 Hz), 7.33 (1 H, d, J = 8.6 Hz), 7.29 (1 H, dd, J = 8.6, 2.4 Hz), 7.12 (1 H, br), 5.42 (1 H, m), 3.18 (1 H, m), 3.02 (1 H, m), 2.83 (1 H, m), 2.59 (1 H, m), 1.62 (3 H, d, J = 6.7 Hz).
MS (ESI) m/z: 381.4 (M+H)⁺.
- [0497] IM I-c-24
¹H NMR (DMSO d6) delta 9.34 (1 H, br d, J = 6.1 Hz), 8.85 (1 H, br d, J = 4.3 Hz), 7.97 (1 H, dd, J = 7.9, 1.8 Hz), 7.74 (1 H, dd, J = 7.9, 4.9 Hz), 7.59 (1 H, d, J = 1.8 Hz), 7.57 (1 H, d, J = 7.9 Hz), 7.48 (1 H, dd, J = 7.9, 1.8 Hz), 5.23 (1 H, m), 2.83 (1 H, m), 2.74-2.50 (3 H, m), 1.44 (3 H, d, J = 7.3 Hz).
MS (ESI) m/z: 381.0 (M+H)⁺.
- [0498] IM I-c-25
¹H NMR (DMSO d6) delta 9.28 (1 H, br d, J = 7.3 Hz), 8.81 (1 H, br d, J = 4.3 Hz), 7.84 (1 H, dd, J = 7.9, 1.8 Hz), 7.69 (1 H, dd, J = 7.9, 4.3 Hz), 7.59 (1 H, dd, J = 8.6, 6.1 Hz), 7.37 (1 H, dd, J = 8.6, 2.4 Hz), 7.24 (1 H, dt, J = 8.6, 2.4 Hz), 5.26 (1 H, m), 2.98-2.80 (2 H, m), 2.77-2.54 (2 H, m), 1.43 (3 H, d, J = 6.7 Hz).
MS (ESI) m/z: 365.0 (M+H)⁺.
- [0499] IM I-c-26
¹H NMR (DMSO d6) delta 9.32 (1 H, br d, J = 7.3 Hz), 8.85 (1 H, d, J = 4.3 Hz), 7.98 (1 H, br d, J = 7.9 Hz), 7.74 (1 H, dd, J = 7.9, 4.3 Hz), 7.60 (1 H, dd, J = 8.6, 6.1 Hz), 7.40 (1 H, br d, J = 9.2 Hz), 7.28 (1 H, dt, J = 8.6, 2.4 Hz), 5.26 (1 H, m), 2.83 (1 H, m), 2.76-2.51 (3 H, m), 1.44 (3 H, d, J = 7.3 Hz).
MS (ESI) m/z: 365.0 (M+H)⁺.
- [0500] IM I-c-27

¹H NMR (CDCl₃) delta 8.88 (1 H, br d, J = 4.3 Hz), 7.57 (1 H, d, J = 7.9 Hz), 7.49 (1 H, dd, J = 7.9, 4.3 Hz), 7.43 (1 H, d, J = 1.8 Hz), 7.23 (1 H, dd, J = 7.9, 1.8 Hz), 7.18 (1 H, d, J = 7.9 Hz), 6.85 (1 H, br), 3.77-3.62 (2 H, m), 3.17-2.94 (4 H, m), 2.74 (1 H, m), 2.52 (1 H, m).

MS (ESI) m/z: 381.0 (M+H)⁺.

[0501] IM I-c-28

¹H NMR (CDCl₃) delta 8.91 (1 H, m), 7.81 (1 H, dd, J = 7.9, 1.8 Hz), 7.57-7.53 (1 H, m), 7.403 (0.5 H, d, J = 2.4 Hz), 7.397 (0.5 H, d, J = 2.4 Hz), 7.189 (0.5 H, dd, J = 8.6, 1.8 Hz), 7.185 (0.5 H, dd, J = 8.6, 1.8 Hz), 7.09 (0.5 H, d, J = 8.6 Hz), 7.08 (0.5 H, d, J = 8.6 Hz), 7.06 (1 H, br), 3.23-3.14 (1 H, m), 3.10-2.96 (2 H, m), 2.89-2.76 (1 H, m), 2.64-2.54 (1 H, m), 2.36-2.29 (1 H, m), 1.44-1.30 (2 H, m).

MS (ESI) m/z: 393.3 (M+H)⁺.

[0502] IM I-c-29

¹H NMR (CDCl₃) delta 8.90 (1 H, ddd, J = 4.3, 1.8, 1.2 Hz), 7.77 (1 H, dd, J = 7.9, 1.2 Hz), 7.53 (1 H, dd, J = 7.9, 4.3 Hz), 7.46 (1 H, br), 3.73 (4 H, dd, J = 4.9, 4.3 Hz), 3.48 (1 H, dd, J = 13.8, 4.9 Hz), 3.45 (1 H, dd, J = 13.8, 4.9 Hz), 3.17 (1 H, m), 3.02 (1 H, m), 2.83 (1 H, m), 2.66 (4 H, dd, J = 4.9, 4.3 Hz), 2.57 (1 H, m), 1.76-1.22 (10 H, m).

MS (ESI) m/z: 390.4 (M+H)⁺.

[0503] IM I-c-30

¹H NMR (CDCl₃) delta 8.87 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.48 (1 H, d, J = 1.8 Hz), 7.46 (1 H, dd, J = 7.9, 4.9 Hz), 7.35-7.32 (2 H, m), 7.29 (1 H, d, J = 8.6 Hz), 6.40 (1 H, br d, J = 6.1 Hz), 4.16 (1 H, dd, J = 14.1, 6.1 Hz), 3.98 (1 H, dd, J = 14.1, 6.1 Hz), 3.92-3.83 (2 H, m), 3.73-3.61 (2 H, m), 3.06 (1 H, m), 2.93 (1 H, m), 2.61 (1 H, m), 2.47-2.36 (3 H, m), 2.13-2.08 (2 H, m).

MS (ESI) m/z: 450.9 (M+H)⁺.

[0504] IM I-c-31

¹H NMR (CDCl₃) delta 8.68 (1 H, br s), 7.462 (1 H, d, J = 1.8 Hz), 7.456 (1 H, d, J = 2.4 Hz), 7.37 (1 H, d, J = 7.9 Hz), 7.27 (1 H, dd, J = 7.9, 1.8 Hz), 7.23 (1 H, br), 4.64 (1 H, dd, J = 15.3, 6.1 Hz), 4.61 (1 H, dd, J = 15.3, 6.1 Hz), 3.11 (1 H, m), 2.97 (1 H, m), 2.79 (1 H, m), 2.54 (1 H, m), 2.40 (3 H, s).

MS (ESI) m/z: 381.0 (M+H)⁺.

[0505] IM I-c-32

¹H NMR (CDCl₃) delta 8.69 (1 H, br s), 7.46 (1 H, d, J = 1.2 Hz), 7.42 (1 H, dd, J = 8.6, 6.1 Hz), 7.20 (1 H, dd, J = 7.9, 2.4 Hz), 7.20 (1 H, br), 7.01 (1 H, dt, J = 8.6, 2.4 Hz), 4.64 (1 H, dd, J = 14.7, 6.1 Hz), 4.61 (1 H, dd, J = 14.7, 6.1 Hz), 3.10 (1 H, m), 2.97 (1 H, m), 2.79 (1 H, m), 2.54 (1 H, m), 2.40 (3 H, s).

MS (ESI) m/z: 365.2 (M+H)⁺.

[0506] The following Intermediates were prepared by General Procedure G or H (Table 45).

In the preparation of the Intermediate (IM) I-g-16, the TBS group was removed under this reaction condition.










[0507] [Table 45-1]

Intermediates	Structure	Chemical Name	Substrate	General Procedure
I-g-8		(2S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		G
I-g-9		(2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	Ex 284	
I-g-10		(2R,5'R)-N-(2,4-dichlorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		H
I-g-11		(2S,5'R)-N-(2,4-dichlorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	IM I-e-26	
I-g-12		(2S,5'S)-N-(2-chloro-4-fluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		G
I-g-13		(2R,5'S)-N-(2-chloro-4-fluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	Ex 249	











[Table 45-2]

I-g-14		(2S,5'S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		G
I-g-15		(2R,5'S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	IM I-c-21	
I-g-16		(2S,5'S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		G
I-g-17		(2S,5'S)-5'-fluoro-N-(2,3,4-trifluorobenzyl)-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		G
I-g-18		(2R,5'S)-5'-fluoro-N-(2,3,4-trifluorobenzyl)-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	Ex 254	

[Table 45-3]

I-g-19		(2S,5'S)-N-(2,4-difluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-17	G
I-g-20		(2S,5'S)-N-(4-chloro-2-fluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-13	G
I-g-21		(2R,5'S)-N-(4-chloro-2-fluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		
I-g-22		(2S,5'S)-N-(2,4-dichloro-6-fluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 EX 251	G
I-g-23		(2S,5'S)-N-(2,4-dichloro-3-fluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-16	G













[Table 45-4]

I-g-24		(2S,5'S)-N-(2,3-dichloro-4-fluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-14	G
I-g-25		(2S,5'S)-N-(2-chloro-4,6-difluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-20	G
I-g-26		(2S,5'S)-N-(4-chloro-2,3-difluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-15	G
I-g-27		(2S,5'S)-N-(3-chloro-2,4-difluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-18	G
I-g-28		(2S,5'S)-N-(4-chloro-2,6-difluorobenzyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-19	G

[Table 45-5]

l-g-29		(2S,5'S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		IM l-c-23	G
l-g-30		(2S,5'S)-N-((S)-1-(2,4-dichlorophenyl)ethyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		IM l-c-24	G
l-g-31		(2S,5'S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		IM l-c-25	G
l-g-32		(2R,5'S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		IM l-c-26	
l-g-33		(2S,5'S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		IM l-c-27	G
l-g-34		(2S,5'S)-N-((3,5-dichloropyridin-2-yl)methyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide		Ex 255	G

[Table 45-6]

I-g-35		(2S,5'S)-N-(2,4-dichlorophenyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-27	G
I-g-36		(2S,5'S)-N-(2-(2,4-dichlorophenyl)cyclopropyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-28	G
I-g-37		(5'S)-5'-fluoro-N-((1-morpholinocyclohexyl)methyl)-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-29	G
I-g-38		(2S,5'S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5'-fluoro-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-30	G
I-g-39		(2S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-3'-methyl-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-31	G
I-g-40		(2S,5'S)-N-(2-chloro-4-fluorobenzyl)-5'-fluoro-3'-methyl-6',7'-dihydro-5'H-spiro[oxirane-2,8'-quinoline]-5'-carboxamide	 IM I-c-32	G

[0508] IM I-g-8
¹H NMR and LCMS were identified with the IM I-g-4.

[0509] IM I-g-9
¹H NMR and LCMS were identified with the IM I-g-5.

[0510] IM I-g-10
¹H NMR and LCMS were identified with the IM I-g-4.

[0511] IM I-g-11
¹H NMR and LCMS were identified with the IM I-g-5.

- [0512] IM I-g-12
¹H NMR and LCMS were identified with the IM I-g-6.
- [0513] IM I-g-13
¹H NMR and LCMS were identified with the IM I-g-7.
- [0514] IM I-g-14
¹H NMR (CDCl₃) delta 8.61 (1 H, br dd, J = 4.9, 1.8 Hz), 7.59 (1 H, br), 7.45 (1 H, d, J = 1.8 Hz), 7.43 (1 H, br dd, J = 7.9, 1.8 Hz), 7.36 (1 H, d, J = 1.8 Hz), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 4.80 (1 H, dd, J = 14.7, 6.1 Hz), 4.78 (2 H, d, J = 6.1 Hz), 4.71 (1 H, dd, J = 14.7, 6.1 Hz), 3.85 (1 H, d, J = 6.1 Hz), 3.73 (1 H, t, J = 6.1 Hz), 3.02 (1 H, d, J = 6.1 Hz), 2.71 (1 H, m), 2.54 (1 H, m), 2.29 (1 H, m), 2.10 (1 H, m).
MS (ESI) m/z: 411.1 (M+H)⁺.
- [0515] IM I-g-15
¹H NMR (DMSO-d₆) delta 8.77 (1 H, br), 8.60 (1 H, d, J = 4.9 Hz), 7.64 (1 H, d, J = 7.9 Hz), 7.55 (1 H, d, J = 1.8 Hz), 7.48 (1 H, d, J = 1.8 Hz), 7.41 (1 H, dd, J = 7.9, 4.9 Hz), 5.52 (1 H, t, J = 5.5 Hz), 4.68 (2 H, d, J = 5.5 Hz), 4.51 (1 H, d, J = 14.7 Hz), 4.46 (1 H, d, J = 14.7 Hz), 3.43 (1 H, d, J = 6.1 Hz), 3.05 (1 H, d, J = 6.1 Hz), 2.50 (1 H, m), 2.33-2.23 (2 H, m), 2.13 (1 H, m).
MS (ESI) m/z: 410.7 (M+H)⁺.
- [0516] IM I-g-16
¹H NMR (CDCl₃) delta 8.61 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.57 (1 H, br d, J = 6.1 Hz), 7.43 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.20 (1 H, dd, J = 7.9, 4.9 Hz), 7.18 (1 H, dd, J = 7.9, 2.4 Hz), 7.10 (1 H, dd, J = 8.6, 2.4 Hz), 4.82-4.77 (3 H, m), 4.71 (1 H, dd, J = 14.7, 6.1 Hz), 3.85 (1 H, d, J = 6.1 Hz), 3.74 (1 H, t, J = 6.1 Hz), 3.02 (1 H, d, J = 6.1 Hz), 2.72 (1 H, m), 2.54 (1 H, m), 2.29 (1 H, m), 2.10 (1 H, m).
MS (ESI) m/z: 394.8 (M+H)⁺.
- [0517] IM I-g-17
¹H NMR (DMSO d₆) delta 9.32 (1 H, br), 8.61 (1 H, br dd, J = 4.9, 1.8 Hz), 7.80 (1 H, br dd, J = 7.9, 1.8 Hz), 7.61 (1 H, dd, J = 7.9, 4.9 Hz), 7.35-7.16 (2 H, m), 4.45 (2 H, m), 3.72 (1 H, d, J = 6.1 Hz), 3.03 (1 H, d, J = 6.1 Hz), 2.60-2.41 (2 H, m), 2.27 (1 H, m), 1.90 (1 H, m).
MS (ESI) m/z: 367.0 (M+H)⁺.
- [0518] IM I-g-18
¹H NMR (DMSO-d₆) delta 9.31 (1 H, br), 8.61 (1 H, br d, J = 4.9 Hz), 7.64 (1 H, d, J = 7.9 Hz), 7.41 (1 H, dd, J = 7.9, 4.9 Hz), 7.32 (1 H, m), 7.22 (1 H, m), 4.45 (1 H, dd, J = 15.3, 5.5 Hz), 4.41 (1 H, dd, J = 15.3, 5.5 Hz), 3.43 (1 H, d, J = 6.1 Hz), 3.05 (1 H, d, J = 6.1 Hz), 2.50 (1 H, m), 2.38-2.23 (2 H, m), 2.12 (1 H, m).
MS (ESI) m/z: 367.0 (M+H)⁺.
- [0519] IM I-g-19

¹H NMR (CDCl₃) delta 8.65 (1 H, br dd, J = 4.9, 1.8 Hz), 7.52 (1 H, br dd, J = 7.9, 1.8 Hz), 7.41 (1 H, m), 7.25 (1 H, br dd, J = 7.9, 4.9 Hz), 7.17 (1 H, br d, J = 6.1 Hz), 6.96-6.87 (2 H, m), 4.66 (1 H, dd, J = 14.7, 6.1 Hz), 4.58 (1 H, dd, J = 14.7, 6.1 Hz), 3.89 (1 H, d, J = 6.1 Hz), 3.05 (1 H, d, J = 6.1 Hz), 2.79 (1 H, m), 2.60 (1 H, m), 2.34 (1 H, m), 2.15 (1 H, m).

MS (ESI) m/z: 349.9 (M+H)⁺.

IM I-g-20

¹H NMR (CDCl₃) delta 8.63 (1 H, br dd, J = 4.9, 1.8 Hz), 7.49 (1 H, br dd, J = 7.9, 1.8 Hz), 7.35 (1 H, m), 7.23 (1 H, br dd, J = 7.9, 4.9 Hz), 7.19-7.11 (3 H, m), 4.64 (1 H, dd, J = 14.7, 6.1 Hz), 4.56 (1 H, dd, J = 14.7, 6.1 Hz), 3.87 (1 H, d, J = 6.1 Hz), 3.03 (1 H, d, J = 6.1 Hz), 2.77 (1 H, m), 2.58 (1 H, m), 2.32 (1 H, m), 2.13 (1 H, m).

MS (ESI) m/z: 364.8 (M+H)⁺.

[0520] IM I-g-21

¹H NMR (CDCl₃) delta 8.65 (1 H, br dd, J = 4.9, 1.8 Hz), 7.53 (1 H, br dd, J = 7.9, 1.8 Hz), 7.34 (1 H, dd, J = 7.9, 6.7 Hz), 7.25 (1 H, dd, J = 7.9, 4.9 Hz), 7.26-7.13 (3 H, m), 4.65 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 3.52 (1 H, d, J = 6.1 Hz), 3.08 (1 H, d, J = 6.1 Hz), 2.67 (1 H, m), 2.53-2.40 (2 H, m), 2.19 (1 H, m).

MS (ESI) m/z: 364.8 (M+H)⁺.

[0521] IM I-g-22

¹H NMR (CDCl₃) delta 8.62 (1 H, br dd, J = 4.9, 1.8 Hz), 7.53 (1 H, br dd, J = 7.9, 1.8 Hz), 7.31 (1 H, d, J = 1.8 Hz), 7.24 (1 H, br dd, J = 7.9, 4.9 Hz), 7.12 (1 H, dd, J = 8.6, 1.8 Hz), 7.02 (1 H, br), 4.78 (1 H, dd, J = 14.7, 6.1 Hz), 4.69 (1 H, dd, J = 14.7, 6.1 Hz), 3.85 (1 H, d, J = 6.1 Hz), 3.02 (1 H, d, J = 6.1 Hz), 2.76 (1 H, m), 2.56 (1 H, m), 2.31 (1 H, m), 2.14 (1 H, m).

MS (ESI) m/z: 398.7 (M+H)⁺.

[0522] IM I-g-23

¹H NMR (CDCl₃) delta 8.63 (1 H, br dd, J = 4.9, 1.8 Hz), 7.50 (1 H, br dd, J = 7.9, 1.8 Hz), 7.34 (1 H, br dd, J = 7.9, 4.9 Hz), 7.26-7.18 (3 H, m), 4.69 (1 H, dd, J = 14.7, 6.1 Hz), 4.65 (1 H, dd, J = 14.7, 6.1 Hz), 3.86 (1 H, d, J = 6.1 Hz), 3.03 (1 H, d, J = 6.1 Hz), 2.76 (1 H, m), 2.57 (1 H, m), 2.33 (1 H, m), 2.14 (1 H, m).

MS (ESI) m/z: 398.8 (M+H)⁺.

[0523] IM I-g-24

¹H NMR (CDCl₃) delta 8.63 (1 H, br dd, J = 4.9, 1.8 Hz), 7.49 (1 H, br dd, J = 7.9, 1.8 Hz), 7.38 (1 H, dd, J = 8.6, 5.5 Hz), 7.24 (1 H, br dd, J = 7.9, 4.9 Hz), 7.25-7.22 (1 H, br), 7.12 (1 H, dd, J = 8.6, 7.9 Hz), 4.69 (1 H, dd, J = 15.3, 6.1 Hz), 4.65 (1 H, dd, J = 15.3, 6.1 Hz), 3.86 (1 H, d, J = 6.1 Hz), 3.03 (1 H, d, J = 6.1 Hz), 2.76 (1 H, m), 2.57 (1 H, m), 2.32 (1 H, m), 2.14 (1 H, m).

MS (ESI) m/z: 398.8 (M+H)⁺.

[0524] IM I-g-25

¹H NMR (CDCl₃) delta 8.62 (1 H, br dd, J = 4.9, 1.8 Hz), 7.54 (1 H, br dd, J = 7.9, 1.8 Hz), 7.24 (1 H, m), 7.06 (1 H, br dd, J = 7.9, 4.9 Hz), 7.01 (1 H, br), 6.86 (1 H, m), 4.78 (1 H, dd, J = 14.7, 5.5 Hz), 4.68 (1 H, dd, J = 14.7, 5.5 Hz), 3.86 (1 H, d, J = 6.1 Hz), 3.02 (1 H, d, J = 6.1 Hz), 2.76 (1 H, m), 2.56 (1 H, m), 2.32 (1 H, m), 2.14 (1 H, m).

MS (ESI) m/z: 382.8 (M+H)⁺.

[0525] IM I-g-26

¹H NMR (CDCl₃) delta 8.63 (1 H, br dd, J = 4.9, 1.8 Hz), 7.50 (1 H, br dd, J = 7.9, 1.8 Hz), 7.28-7.08 (4 H, m), 4.66 (1 H, dd, J = 14.7, 6.7 Hz), 4.61 (1 H, dd, J = 14.7, 6.7 Hz), 3.87 (1 H, br d, J = 6.1 Hz), 3.03 (1 H, br d, J = 6.1 Hz), 2.76 (1 H, m), 2.57 (1 H, m), 2.33 (1 H, m), 2.15 (1 H, m).

MS (ESI) m/z: 382.8 (M+H)⁺.

[0526] IM I-g-27

¹H NMR (CDCl₃) delta 8.63 (1 H, br dd, J = 4.9, 1.8 Hz), 7.49 (1 H, br dd, J = 7.9, 1.8 Hz), 7.31 (1 H, m), 7.24 (1 H, br dd, J = 7.9, 4.9 Hz), 7.19 (1 H, br), 7.00 (1 H, ddd, J = 8.6, 4.3, 1.8 Hz), 4.64 (1 H, dd, J = 14.7, 6.1 Hz), 4.59 (1 H, dd, J = 14.7, 6.1 Hz), 3.87 (1 H, d, J = 6.1 Hz), 3.03 (1 H, d, J = 6.1 Hz), 2.76 (1 H, m), 2.57 (1 H, m), 2.32 (1 H, m), 2.13 (1 H, m).

MS (ESI) m/z: 382.8 (M+H)⁺.

[0527] IM I-g-28

¹H NMR (CDCl₃) delta 8.62 (1 H, br dd, J = 4.3, 1.8 Hz), 7.51 (1 H, br dd, J = 7.9, 1.8 Hz), 7.24 (1 H, br dd, J = 7.9, 4.3 Hz), 7.06-6.99 (3 H, m), 4.71 (1 H, dd, J = 14.7, 5.5 Hz), 4.61 (1 H, dd, J = 14.7, 4.9 Hz), 3.86 (1 H, d, J = 6.1 Hz), 3.02 (1 H, d, J = 6.1 Hz), 2.74 (1 H, m), 2.56 (1 H, m), 2.30 (1 H, m), 2.13 (1 H, m).

MS (ESI) m/z: 382.8 (M+H)⁺.

[0528] IM I-g-29

¹H NMR (CDCl₃) delta 8.60 (1 H, br dd, J = 4.9, 1.8 Hz), 7.45 (1 H, d, J = 1.8 Hz), 7.44 (1 H, br dd, J = 7.9, 1.8 Hz), 7.37 (1 H, d, J = 8.6 Hz), 7.30 (1 H, dd, J = 8.6, 1.8 Hz), 7.19 (1 H, dd, J = 7.9, 4.9 Hz), 7.13 (1 H, br), 5.47 (1 H, m), 3.85 (1 H, d, J = 6.1 Hz), 3.02 (1 H, d, J = 6.1 Hz), 2.78 (1 H, m), 2.58 (1 H, m), 2.35 (1 H, m), 2.16 (1 H, m), 1.60 (3 H, d, J = 7.3 Hz).

MS (ESI) m/z: 395.3 (M+H)⁺.

[0529] IM I-g-30

¹H NMR (CDCl₃) delta 8.64 (1 H, br dd, J = 4.9, 1.8 Hz), 7.61 (1 H, br dd, J = 7.9, 1.8 Hz), 7.42 (1 H, d, J = 1.8 Hz), 7.31-7.26 (3 H, m), 7.17 (1 H, dd, J = 6.7, 6.1 Hz), 5.45 (1 H, m), 3.86 (1 H, d, J = 6.1 Hz), 3.01 (1 H, d, J = 6.1 Hz), 2.77-2.51 (2 H, m), 2.27 (1 H, m), 2.10 (1 H, m), 1.62 (3 H, d, J = 7.3 Hz).

MS (ESI) m/z: 395.2 (M+H)⁺.

[0530] IM I-g-31

¹H NMR (CDCl₃) delta 8.59 (1 H, br dd, J = 4.3, 1.8 Hz), 7.45 (1 H, br dd, J = 7.9, 1.8 Hz), 7.41 (1 H, dd, J = 8.6, 6.1 Hz), 7.18 (1 H, dd, J = 7.9, 4.3 Hz), 7.16 (1 H, dd, J = 6.1, 3.1 Hz), 7.14 (1 H, br), 7.03 (1 H, ddd, J = 8.5, 7.9, 3.1 Hz), 5.46 (1 H, m), 3.85 (1 H, d, J = 6.1 Hz), 3.02 (1 H, d, J = 6.1 Hz), 2.77 (1 H, m), 2.57 (1 H, m), 2.34 (1 H, m), 2.15 (1 H, m), 1.60 (3 H, d, J = 6.7 Hz).

MS (ESI) m/z: 379.2 (M+H)⁺.

[0531] IM I-g-32

¹H NMR (CDCl₃) delta 8.59 (1 H, br dd, J = 4.9, 1.8 Hz), 7.45 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.40 (1 H, dd, J = 8.6, 6.1 Hz), 7.18 (1 H, dd, J = 8.6, 2.4 Hz), 7.16 (1 H, dd, J = 7.9, 4.9 Hz), 7.14 (1 H, br), 7.04 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 5.45 (1 H, m), 3.50 (1 H, d, J = 6.1 Hz), 3.05 (1 H, d, J = 6.1 Hz), 2.67 (1 H, m), 2.50-2.42 (2 H, m), 2.22 (1 H, m), 1.60 (3 H, d, J = 6.7 Hz).

MS (ESI) m/z: 379.2 (M+H)⁺.

[0532] IM I-g-33

¹H NMR (CDCl₃) delta 8.64 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.61 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.36 (1 H, dd, J = 8.6, 6.1 Hz), 7.28 (1 H, dd, J = 7.9, 4.9 Hz), 7.16 (1 H, dd, J = 8.6, 3.1 Hz), 7.13 (1 H, br), 7.02 (1 H, dt, J = 8.6, 3.1 Hz), 5.46 (1 H, m), 3.86 (1 H, d, J = 6.1 Hz), 3.02 (1 H, d, J = 6.1 Hz), 2.78-2.52 (2 H, m), 2.28 (1 H, m), 2.10 (1 H, m), 1.63 (3 H, d, J = 6.7 Hz).

MS (ESI) m/z: 378.9 (M+H)⁺.

[0533] IM I-g-34

¹H NMR (CDCl₃) delta 8.64 (1 H, br dd, J = 4.3, 1.8 Hz), 8.47 (1 H, d, J = 2.4 Hz), 8.10 (1 H, br), 7.78 (1 H, d, J = 2.4 Hz), 7.74 (1 H, dd, J = 7.9, 1.8 Hz), 7.26 (1 H, dd, J = 7.9, 4.3 Hz), 4.83 (1 H, dd, J = 18.3, 4.9 Hz), 4.71 (1 H, dd, J = 18.3, 4.9 Hz), 3.89 (1 H, d, J = 6.1 Hz), 3.05 (1 H, d, J = 6.1 Hz), 2.83 (1 H, m), 2.62 (1 H, m), 2.40 (1 H, m), 2.16 (1 H, m).

MS (ESI) m/z: 382.5 (M+H)⁺.

[0534] IM I-g-35

¹H NMR (CDCl₃) delta 8.61 (1 H, br d, J = 4.3 Hz), 7.43 (1 H, s), 7.33 (1 H, d, J = 7.9 Hz), 7.25-7.19 (2 H, m), 7.21 (1 H, dd, J = 7.9, 4.3 Hz), 6.86 (1 H, br d, J = 4.9 Hz), 3.86 (1 H, d, J = 6.1 Hz), 3.79-3.63 (2 H, m), 3.08-3.04 (2 H, m), 3.02 (1 H, d, J = 6.1 Hz), 2.72 (1 H, m), 2.55 (1 H, m), 2.26 (1 H, m), 2.12 (1 H, m).

MS (ESI) m/z: 395.0 (M+H)⁺.

[0535] IM I-g-36

¹H NMR (CDCl₃) delta 8.65 (1 H, m), 7.59 (1 H, m), 7.40 (1 H, d, J = 1.8 Hz), 7.27 (1 H, m), 7.19 (1 H, m), 7.12 (1 H, d, J = 8.6 Hz), 7.03 (1 H, br), 3.88 (0.5 H, d, J = 6.1

Hz), 3.87 (0.5 H, d, J = 6.1 Hz), 3.10 (1 H, m), 3.03 (1 H, d, J = 6.1 Hz), 2.79 (1 H, m), 2.60 (1 H, m), 2.39-2.29 (2 H, m), 2.15 (1 H, m), 1.43-1.31 (2 H, m).

MS (ESI) m/z: 407.3 (M+H)⁺.

[0536] IM I-g-37

¹H NMR (CDCl₃) delta 8.63 (1 H, ddd, J = 4.3, 1.8, 1.8 Hz), 7.55 (1 H, dd, J = 7.9, 1.8 Hz), 7.39 (1 H, br), 7.25 (1 H, m), 3.87 (0.7 H, d, J = 6.1 Hz), 3.7-3.69 (4 H, m), 3.54-3.42 (2.3 H, m), 3.06 (0.3 H, d, J = 6.1 Hz), 3.03 (0.7 H, d, J = 6.1 Hz), 2.85-2.38 (6 H, m), 2.31 (1 H, m), 2.15 (1 H, m), 1.76-1.46 (10 H, m).

MS (ESI) m/z: 404.2 (M+H)⁺.

[0537] IM I-g-38

¹H NMR (CDCl₃) delta 8.60 (1 H, br d, J = 4.9 Hz), 7.47 (1 H, d, J = 1.8 Hz), 7.34-7.29 (2 H, m), 7.18 (1 H, dd, J = 7.9, 4.9 Hz), 7.10 (1 H, br d, J = 7.9 Hz), 6.40 (1 H, br d, J = 6.7 Hz), 4.21 (1 H, dd, J = 14.1, 6.7 Hz), 3.99 (1 H, dd, J = 14.1, 6.1 Hz), 3.93-3.84 (2 H, m), 3.81 (1 H, d, J = 6.1 Hz), 3.74-3.62 (2 H, m), 2.99 (1 H, d, J = 6.1 Hz), 2.60 (1 H, m), 2.50-2.34 (3 H, m), 2.19-2.04 (4 H, m).

MS (ESI) m/z: 465.0 (M+H)⁺.

[0538] IM I-g-39

¹H NMR (CDCl₃) delta 8.44 (1 H, br), 7.46 (1 H, d, J = 1.8 Hz), 7.39 (1 H, m), 7.29-7.22 (3 H, m), 4.71-4.59 (2 H, m), 3.85 (1 H, d, J = 6.1 Hz), 3.00 (1 H, d, J = 6.1 Hz), 2.76 (1 H, m), 2.57 (1 H, m), 2.29 (1 H, m), 2.27 (3 H, s), 2.09 (1 H, m).

MS (ESI) m/z: 395.0 (M+H)⁺.

[0539] IM I-g-40

¹H NMR (CDCl₃) delta 8.44 (1 H, br), 7.46 (1 H, dd, J = 8.6, 6.1 Hz), 7.20-7.14 (3 H, m), 7.02 (1 H, ddd, J = 8.6, 7.9, 3.1 Hz), 4.69 (1 H, dd, J = 14.7, 6.1 Hz), 4.62 (1 H, dd, J = 14.7, 6.1 Hz), 3.86 (1 H, d, J = 6.1 Hz), 3.01 (1 H, d, J = 6.1 Hz), 2.76 (1 H, m), 2.58 (1 H, m), 2.29 (1 H, m), 2.27 (3 H, s), 2.07 (1 H, m).

MS (ESI) m/z: 379.4 (M+H)⁺.

[0540] The following Examples and Intermediates were prepared by General Procedure J, K, P, Q, R, or S (Tables 46 and 48).

[0541] General Procedure P

A mixture of substrate (1.0 eq.), LiClO₄ (1.5 eq.), and KCN (1.5 eq.) in MeCN was heated at reflux. After being stirred at reflux until complete reaction, the mixture was cooled to room temperature. Water was added to the mixture, and the mixture was extracted with EtOAc twice. The extracts were washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by silica gel column chromatography and preparative HPLC to afford following Examples.

[0542] General Procedure Q

The substrate (1.0 eq.) was dissolved with 1.0 M TBAF in THF (12.0 eq.) and the

mixture was heated at 70 °C until complete reaction. The mixture was cooled to room temperature and water was added to the mixture. The mixture was extracted with EtOAc and washed with brine. The extract was dried over Na₂SO₄ and concentrated in vacuo. The resulting mixture was purified by silica gel column chromatography and preparative HPLC to afford the following Examples.

[0543] General Procedure R

Fifteen percent of sodium thiomethoxide in water (3.0 eq.) was added to the solution of substrate (1.0 eq.) in THF. The mixture was stirred at 60 °C until complete reaction and then cooled to room temperature. Water was added to the mixture and extracted with EtOAc twice. The extracts were dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography and preparative HPLC to afford the following Examples and Intermediates.

[0544] General Procedure S











To a mixture of substrate (1.0 eq.) and K₂CO₃ (2.0 eq.) in acetone was added mercaptoethanol (4.4 eq.) at ambient temperature. The mixture was stirred at 70 °C until complete reaction and cooled to room temperature. The volatile was removed under reduced pressure and water was added to the mixture. The mixture was extracted with EtOAc, dried over Na₂SO₄, and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography and preparative HPLC to afford the following Examples and Intermediates.

[0545]

[Table 46-1]

Examples	Structure	Chemical Name	Substrate	General Procedure (/Amine)
256		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		K
257		(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		K
258		(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		K
259		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		K
260		(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		K

[Table 46-2]

261		(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-14	K
262		(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-16	K
263		(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-19	K
264		(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-20	K
265		(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-22	K

[Table 46-3]

266		(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM-I-g-25	K
267		(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-26	K
268		(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-27	K
269		(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-28	K
270		(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM-I-g-31	K











[Table 46-4]

271		(5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-33	K
272		(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-34	K
273		(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-36	K
274		(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-37	K
275		(5S,8R)-5-fluoro-8-hydroxy-8-methyl-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide			
276		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3,8-dimethyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-40	K

[Table 46-5]

277		(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-8	J/
278		(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-9	J/
279		(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-8	J/
280		(5S,8R)-8-(cyanomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-8	P
281		(5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-12	P

[Table 46-6]

282		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-g-13	P
283		(5S,8R)-8-(cyanomethyl)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-g-14	P
284		(5S,8R)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-g-16	P
285		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-g-8	Q
286		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-g-12	Q

[Table 46-7]

287		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-8	R
288		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-12	R
289		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)thio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-12	S

[0546] [Table 47]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
256	C	1.57	382.9	272	D	1.54	384.0
257	C	1.57	383.0	273	D	1.79	409.1
258	C	1.43	369.0	274	D	1.61	406.2
259	C	1.47	367.0	275	D	1.58	406.2
260	C	1.45	366.9	276	D	1.63	381.1
261	C	1.43	412.9	280	D	1.66	408.0
262	D	1.37	397.1	281	D	1.55	392.1
263	C	1.37	351.1	282	D	1.54	392.1
264	C	1.48	367.1	283	D	1.49	438.1
265	C	1.57	401.0	284	D	1.37	422.0
266	D	1.54	385.0	285	D	1.70	401.0
267	D	1.59	385.0	286	D	1.58	385.1
268	D	1.58	385.0	287	D	1.78	429.0
269	D	1.57	385.0	288	D	1.67	413.1
270	D	1.61	381.1	289	D	1.44	443.1
271	D	1.62	381.1				

[0547] Ex 256

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.49 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.44 (1 H, d, J = 2.4 Hz), 7.36 (1 H, d, J = 7.9 Hz), 7.23 (1 H, dd, J = 7.9, 2.4 Hz), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 7.13 (1 H, br d, J = 5.5 Hz), 4.62 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 3.26 (1 H, br s), 2.65 (1 H, m), 2.38 (1 H, m), 2.22-2.11 (2 H, m), 1.63 (3 H, s).

[0548] Ex 257

¹H NMR (CDCl₃) delta 8.60 (1 H, br dd, J = 4.9, 1.8 Hz), 7.46 (1 H, d, J = 1.8 Hz), 7.45 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.39-7.20 (4 H, m), 4.66 (1 H, dd, J = 14.7, 6.1 Hz), 4.59 (1 H, dd, J = 14.7, 6.1 Hz), 4.00 (1 H, br s), 2.60 (1 H, m), 2.36-2.23 (2 H, m), 2.16 (1 H, m), 1.56 (3 H, s).

[0549] Ex 258

¹H NMR (CDCl₃) delta 8.61 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.49 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.22 (1 H, dd, J = 7.9, 4.9 Hz), 7.14-7.08 (2 H, m), 6.97 (1 H, m), 4.61 (1 H, dd, J = 14.7, 6.1 Hz), 4.55 (1 H, dd, J = 14.7, 6.1 Hz), 3.30 (1 H, br s), 2.65 (1 H, m), 2.37 (1 H, m), 2.22-2.11 (2 H, m), 1.63 (3 H, s).

[0550] Ex 259

¹H NMR (CDCl₃) delta 8.61 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.50 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.41 (1 H, dd, J = 8.6, 5.5 Hz), 7.22 (1 H, dd, J = 7.9, 4.9 Hz), 7.18 (1 H, dd, J = 8.6, 2.4 Hz), 7.11 (1 H, br d, J = 5.5 Hz), 6.99 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 3.18 (1 H, br s), 2.65 (1 H, m), 2.39 (1 H, m), 2.23-2.10 (2 H, m), 1.63 (3 H, s).

[0551] Ex 260

¹H NMR (CDCl₃) delta 8.60 (1 H, br dd, J = 4.9, 1.8 Hz), 7.45-7.41 (2 H, m), 7.26-7.18 (3 H, m), 7.00 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 4.66 (1 H, dd, J = 14.7, 6.1 Hz), 4.59 (1 H, dd, J = 14.7, 5.5 Hz), 4.03 (1 H, s), 2.59 (1 H, m), 2.36-2.23 (2 H, m), 2.14 (1 H, m), 1.56 (3 H, s).

[0552] Ex 261

¹H NMR (CDCl₃) delta 8.57 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.52 (1 H, br d, J = 5.5 Hz), 7.43-7.41 (2 H, m), 7.33 (1 H, d, J = 1.8 Hz), 7.18 (1 H, dd, J = 7.9, 4.9 Hz), 4.76-4.69 (3 H, m), 4.64 (1 H, dd, J = 14.7, 6.1 Hz), 3.99 (1 H, br), 3.43 (1 H, br), 2.60 (1 H, m), 2.32 (1 H, m), 2.17-2.04 (2 H, m), 1.61 (3 H, s).

[0553] Ex 262

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.51 (1 H, br d, J = 5.5 Hz), 7.43 (1 H, br d, J = 7.9 Hz), 7.19 (1 H, dd, J = 7.9, 4.9 Hz), 7.17 (1 H, dd, J = 7.9, 2.4 Hz), 7.09 (1 H, dd, J = 8.6, 2.4 Hz), 4.79-4.62 (3 H, m), 4.66 (1 H, dd, J = 14.7, 6.1 Hz), 3.79 (1 H, br), 3.21 (1 H, br), 2.60 (1 H, m), 2.33 (1 H, m), 2.19-2.08 (2 H, m), 1.63 (3 H, s).

[0554] Ex 263

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.49 (1 H, d, J = 7.9 Hz), 7.36 (1 H, dd, J = 8.6, 6.1 Hz), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 7.07 (1 H, br d, J = 4.9 Hz), 6.90-6.84 (2 H, m), 4.59 (1 H, dd, J = 14.7, 6.1 Hz), 4.52 (1 H, dd, J = 14.7, 6.1 Hz), 3.28 (1 H, br), 2.66 (1 H, m), 2.38 (1 H, m), 2.22-2.11 (2 H, m), 1.63 (3 H, s).

[0555] Ex 264

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.49 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.31 (1 H, t, J = 7.9 Hz), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 7.16-7.08 (3 H, m), 4.59 (1 H, dd, J = 14.7, 6.1 Hz), 4.52 (1 H, dd, J = 14.7, 6.1 Hz), 3.31 (1 H, br s), 2.66 (1 H, m), 2.38 (1 H, m), 2.22-2.11 (2 H, m), 1.63 (3 H, s).

[0556] Ex 265

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.51 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.29 (1 H, dd, J = 1.8, 1.2 Hz), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 7.10 (1 H, dd, J = 9.2, 1.8 Hz), 6.97 (1 H, br d, J = 5.5 Hz), 4.74 (1 H, dd, J = 14.7, 6.1 Hz), 4.64 (1 H, dd, J = 14.7, 5.5 Hz), 3.29 (1 H, br s), 2.66 (1 H, m), 2.38 (1 H, m), 2.21-2.10 (2 H, m), 1.63 (3 H, s).

[0557] Ex 266

¹H NMR (CDCl₃) delta 8.59 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.52 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.21 (1 H, dd, J = 7.9, 4.9 Hz), 7.04 (1 H, br dd, J = 7.9, 2.4 Hz), 6.96 (1 H, br d, J = 5.5 Hz), 6.84 (1 H, br dt, J = 8.6, 2.4 Hz), 4.74 (1 H, dd, J = 14.7, 6.1 Hz), 4.64 (1 H, dd, J = 14.7, 5.5 Hz), 3.32 (1 H, br s), 2.67 (1 H, m), 2.32 (1 H, m), 2.21-2.10 (2 H, m), 1.63 (3 H, s).

[0558] Ex 267

¹H NMR (CDCl₃) delta 8.61 (1 H, m), 7.49 (1 H, dd, J = 7.9, 1.2 Hz), 7.25-7.08 (4 H, m), 4.62 (1 H, dd, J = 15.3, 6.1 Hz), 4.56 (1 H, dd, J = 15.3, 6.7 Hz), 3.30 (1 H, br), 2.64 (1 H, m), 2.38 (1 H, m), 2.22-2.11 (2 H, m), 1.63 (3 H, s).

[0559] Ex 268

¹H NMR (CDCl₃) delta 8.61 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.49 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.28 (1 H, m), 7.22 (1 H, dd, J = 7.9, 4.9 Hz), 7.11 (1 H, br), 6.98 (1 H, dt, J = 8.6, 1.8 Hz), 4.60 (1 H, dd, J = 14.7, 6.1 Hz), 4.55 (1 H, dd, J = 14.7, 6.1 Hz), 3.29 (1 H, br s), 2.65 (1 H, m), 2.38 (1 H, m), 2.25-2.11 (2 H, m), 1.63 (3 H, s).

[0560] Ex 269

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.50 (1 H, d, J = 7.9 Hz), 7.22 (1 H, dd, J = 7.9, 4.9 Hz), 7.04-6.95 (3 H, m), 4.66 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 5.5 Hz), 3.30 (1 H, br s), 2.65 (1 H, m), 2.37 (1 H, m), 2.20-2.09 (2 H, m), 1.63 (3 H, s).

[0561] Ex 270

¹H NMR (CDCl₃) delta 8.57 (1 H, br d, J = 4.9 Hz), 7.45 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.39 (1 H, dd, J = 8.6, 6.1 Hz), 7.17-7.14 (2 H, m), 7.06 (1 H, dd, J = 6.7, 5.5 Hz),

7.02 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 5.42 (1 H, dq, J = 6.7, 6.7 Hz), 3.33 (1 H, br s), 2.68 (1 H, m), 2.39 (1 H, m), 2.24-2.13 (2 H, m), 1.63 (3 H, s), 1.59 (3 H, d, J = 6.7 Hz).

[0562] Ex 271

¹H NMR (CDCl₃) delta 8.62 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.60 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.34 (1 H, dd, J = 8.6, 5.5 Hz), 7.26 (1 H, dd, J = 7.9, 4.9 Hz), 7.15 (1 H, dd, J = 8.6, 2.4 Hz), 7.07 (1 H, dd, J = 6.7, 5.5 Hz), 7.01 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 5.41 (1 H, dq, J = 7.3, 6.7 Hz), 3.28 (1 H, br s), 2.59 (1 H, m), 2.33 (1 H, m), 2.19-2.08 (2 H, m), 1.63 (3 H, s), 1.59 (3 H, d, J = 7.3 Hz).

[0563] Ex 272

¹H NMR (CDCl₃) delta 8.63 (1 H, br d, J = 4.9 Hz), 8.48 (1 H, d, J = 2.4 Hz), 8.03 (1 H, br d, J = 4.3 Hz), 7.77 (1 H, d, J = 2.4 Hz), 7.72 (1 H, d, J = 7.9 Hz), 7.24 (1 H, dd, J = 7.9, 4.9 Hz), 4.78 (1 H, dd, J = 18.3, 4.9 Hz), 4.67 (1 H, dd, J = 18.3, 4.3 Hz), 3.27 (1 H, br s), 2.72 (1 H, m), 2.40 (1 H, m), 2.29-2.18 (2 H, m), 1.66 (3 H, s).

[0564] Ex 277

¹H NMR (CDCl₃) delta 8.61 (1 H, dt, J = 4.7, 1.6 Hz), 7.51 (1 H, dt, J = 8.0, 1.5 Hz), 7.44 (1 H, s), 7.35 (1 H, d, J = 8.19 Hz), 7.16-7.29 (3 H, m), 4.53-4.66 (2 H, m), 3.46-3.64 (2 H, m), 3.03 (1 H, d, J = 14.1 Hz), 2.85 (1 H, d, J = 14.1 Hz), 2.49-2.75 (3 H, m), 2.13-2.40 (6 H, m) 1.99-2.10 (1 H, m).

MS (ESI) m/z: 456.2 (M+H)⁺.

[0565] Ex 278

¹H NMR and LCMS were identified with the Ex 277.

[0566] Ex 279

¹H NMR (CDCl₃) delta 8.61 (1 H, br d, J = 4.7 Hz), 7.54 (1 H, d, J = 8.0 Hz), 7.44 (1 H, d, J = 2.1 Hz), 7.34 (1 H, d, J = 7.7 Hz), 7.27-7.30 (1 H, m), 7.19-7.25 (1 H, m), 4.48-4.65 (3 H, m), 3.96-4.16 (2 H, m), 3.89 (1 H, br s), 3.56 (1 H, br d, J = 13.0 Hz), 3.18 (1 H, br d, J = 13.1 Hz), 2.67-2.82 (1 H, m), 2.16-2.28 (3 H, m).

MS (ESI) m/z: 454.2 (M+H)⁺.

[0567] Ex 280

¹H NMR (DMSO d₆) delta 9.25 (1 H, br), 8.70 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.67 (1 H, dd, J = 7.9, 1.8 Hz), 7.64 (1 H, d, J = 1.8 Hz), 7.50-7.42 (2 H, m), 7.39 (1 H, d, J = 6.7 Hz), 5.98 (1 H, s), 4.53 (1 H, dd, J = 15.3, 6.1 Hz), 4.44 (1 H, dd, J = 15.3, 5.5 Hz), 3.35 (1 H, d, J = 16.5 Hz), 3.24 (1 H, d, J = 16.5 Hz), 2.82-2.64 (2 H, m), 2.30-2.17 (2 H, m).

[0568] Ex 282

¹H NMR (CDCl₃) delta 8.63 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.56 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.40 (1 H, dd, J = 8.6, 5.5 Hz), 7.32 (1 H, dd, J = 7.9, 4.9 Hz), 7.23 (1 H, br), 7.20 (1 H, dd, J = 8.6, 2.4 Hz), 7.01 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 4.63 (1 H, dd,

J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 4.29 (1 H, s), 2.98 (1 H, d, J = 16.5 Hz), 2.89 (1 H, d, J = 16.5 Hz), 2.82-2.27 (4 H, m).

[0569] Ex 283

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.60 (1 H, br), 7.45 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.44 (1 H, d, J = 1.8 Hz), 7.33 (1 H, d, J = 1.8 Hz), 7.26 (1 H, m), 4.80-4.62 (4 H, m), 3.78 (1 H, dd, J = 6.1, 5.5 Hz), 3.71 (1 H, br s), 3.20 (1 H, d, J = 16.5 Hz), 3.06 (1 H, d, J = 16.5 Hz), 2.71 (1 H, m), 2.40-2.37 (2 H, m), 2.19 (1 H, m).

[0570] Ex 284

¹H NMR (CDCl₃) delta 8.60 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.59 (1 H, br d, J = 6.1 Hz), 7.45 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.26 (1 H, m), 7.18 (1 H, dd, J = 7.9, 2.4 Hz), 7.08 (1 H, dd, J = 8.6, 2.4 Hz), 4.87-4.71 (3 H, m), 4.65 (1 H, dd, J = 14.7, 6.1 Hz), 3.77 (1 H, br), 3.62 (1 H, br), 3.20 (1 H, d, J = 17.1 Hz), 3.06 (1 H, d, J = 17.1 Hz), 2.73 (1 H, m), 2.43-2.37 (2 H, m), 2.18 (1 H, m).

[0571] Ex 287

¹H NMR (CDCl₃) delta 8.62 (1 H, br d, J = 4.3 Hz), 7.49 (1 H, dd, J = 7.9, 1.2 Hz), 7.45 (1 H, d, J = 2.4 Hz), 7.37 (1 H, d, J = 7.9 Hz), 7.27 (1 H, dd, J = 7.9, 2.4 Hz), 7.26 (1 H, dd, J = 7.9, 4.3 Hz), 7.18 (1 H, br d, J = 5.5 Hz), 4.65 (1 H, dd, J = 14.7, 6.1 Hz), 4.58 (1 H, dd, J = 14.7, 6.1 Hz), 3.33 (1 H, br), 3.16 (1 H, d, J = 13.5 Hz), 3.12 (1 H, d, J = 13.5 Hz), 2.68 (1 H, m), 2.52 (1 H, m), 2.30 (1 H, m), 2.18 (1 H, m), 2.10 (3 H, s).

[0572] Ex 289

¹H NMR (CDCl₃) delta 8.62 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.51 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.41 (1 H, dd, J = 8.6, 6.1 Hz), 7.24 (1 H, dd, J = 7.9, 4.9 Hz), 7.19 (1 H, dd, J = 7.9, 3.1 Hz), 7.19 (1 H, br), 7.00 (1 H, ddd, J = 8.6, 7.9, 3.1 Hz), 4.64 (1 H, dd, J = 14.7, 6.1 Hz), 4.58 (1 H, dd, J = 14.7, 6.1 Hz), 3.73 (2 H, t, J = 6.1 Hz), 3.71 (1 H, br), 3.23 (1 H, d, J = 13.4 Hz), 3.18 (1 H, d, J = 13.4 Hz), 2.82-2.63 (3 H, m), 2.45 (1 H, m), 2.31 (1 H, m), 2.17 (1 H, m), 1.69 (1 H, br).

[0573]

[Table 48]

Intermediates	Structure	Chemical Name	Substrate	General Procedure (/Amine)
I-h-2-1		(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		R
I-h-2-2		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)thio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamid		S
I-h-1-1		(5R,8S)-8-(aminomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		J/ NH ₄ Cl
I-h-1-2		(5S,8S)-8-(aminomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		J/ NH ₄ Cl
I-h-1-3		(5S,8R)-8-(aminomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		J/ NH ₄ Cl

[0574] IM I-h-2-1

¹H NMR (CDCl₃) delta 8.62 (1 H, ddd, J = 4.3, 1.8, 1.2 Hz), 7.49 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.45 (1 H, d, J = 1.8 Hz), 7.36 (1 H, d, J = 8.6 Hz), 7.28-7.21 (3 H, m), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.58 (1 H, dd, J = 14.7, 6.1 Hz), 4.12 (1 H, s), 2.97 (2 H, s), 2.65-2.43 (2 H, m), 2.30 (1 H, m), 2.18 (3 H, s), 2.14 (1 H, m).

MS (ESI) m/z: 429.1 (M+H)⁺.

[0575] IM I-h-2-2

^1H NMR (CDCl_3) δ 8.61 (1 H, br dd, $J = 4.9, 1.8$ Hz), 7.50 (1 H, br dd, $J = 7.9, 1.8$ Hz), 7.45 (1 H, d, $J = 1.8$ Hz), 7.36 (1 H, d, $J = 7.9$ Hz), 7.27-7.22 (2 H, m), 7.20 (1 H, br), 4.64 (1 H, dd, $J = 14.7, 6.1$ Hz), 4.58 (1 H, dd, $J = 14.7, 6.1$ Hz), 3.82 (1 H, br), 3.72 (2 H, t, $J = 5.5$ Hz), 3.47 (1 H, s), 3.24 (1 H, d, $J = 13.5$ Hz), 3.17 (1 H, d, $J = 13.5$ Hz), 2.81-2.62 (3 H, m), 2.44 (1 H, m), 2.30 (1 H, m), 2.17 (1 H, m).

MS (ESI) m/z : 459.2 ($\text{M}+\text{H}$) $^+$.

[0576] IM I-h-1-1

MS (ESI) m/z : 398.1 ($\text{M}+\text{H}$) $^+$.

[0577] IM I-h-1-2

MS (ESI) m/z : 398.1 ($\text{M}+\text{H}$) $^+$.


[0578] IM I-h-1-3

MS (ESI) m/z : 398.1 ($\text{M}+\text{H}$) $^+$.

[0579] Example 290:

(2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-4-methyl-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide

[0580] [Chem.30]



[0581] To a stirred solution of substrate (1.0 eq.) and TMAD (1.5 eq.) in THF was added $n\text{-Bu}_3\text{P}$ (1.5 eq.) at ambient temperature. After being stirred at ambient temperature until complete reaction, the mixture was concentrated under reduced pressure. The crude residue was purified by silica gel column chromatography (EtOAc) to afford the title compound.

[0582] Ex 290

^1H NMR ($\text{DMSO}-d_6$) δ 9.22 (1 H, br s), 8.65 (1 H, dt, $J = 4.6, 1.6$ Hz), 7.60-7.70 (2 H, m), 7.36-7.49 (3 H, m), 4.35-4.53 (2 H, m), 3.79 (1 H, td, $J = 11.7, 2.5$ Hz), 3.61 (1 H, dd, $J = 11.6, 3.1$ Hz), 3.08 (1 H, d, $J = 11.3$ Hz), 2.84 (1 H, dt, $J = 11.8, 2.5$ Hz), 2.66 (1 H, br d, $J = 11.1$ Hz), 2.52-2.58 (1 H, m), 2.32-2.47 (1 H, m), 2.21 (3 H, s), 1.90-2.16 (3 H, m).

MS (ESI) m/z : 438.1 ($\text{M}+\text{H}$) $^+$.

[0583] The following Examples were prepared by General Procedure T (Table 49).

[0584] General Procedure T

The mixture of substrate (1.0 eq.) and 1,1'-carbonylbis-1H-imidazole (1.1 eq.) in THF was stirred at ambient temperature until complete reaction. The reaction mixture was purified by silica gel column chromatography (10% MeOH/EtOAc) to afford the following Examples.

[0585] [Table 49]

Examples	Structure	Chemical Name	Substrate
291		(5S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-quinoline]-5'-carboxamide	
292		(5R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-quinoline]-5'-carboxamide	

[0586] Ex 291

¹H NMR (DMSO d6) delta 9.27 (1 H, br s), 8.74 (1 H, d, J = 4.5 Hz), 7.75-7.80 (2 H, m), 7.62 (1 H, d, J = 2.1 Hz), 7.53 (1 H, dd, J = 8.0, 4.7 Hz), 7.45 (1 H, dd, J = 8.4, 2.1 Hz), 7.36 (1 H, d, J = 8.3 Hz), 4.40 (2 H, t, J = 5.3 Hz), 3.88 (1 H, d, J = 8.6 Hz), 3.53 (1 H, d, J = 8.4 Hz), 2.29-2.48 (5 H, m).

MS (ESI) m/z: 424.1 (M+H)⁺.

[0587] Ex 292

¹H NMR (DMSO d6) delta 9.30 (1 H, br s), 8.76 (1 H, d, J = 4.8 Hz), 7.79 (1 H, s), 7.71 (1 H, d, J = 8.2 Hz), 7.64 (1 H, d, J = 2.1 Hz), 7.54 (1 H, t, J = 6.2 Hz), 7.47 (1 H, d, J = 8.1 Hz), 7.40 (1 H, d, J = 8.1 Hz), 4.40-4.53 (2 H, m), 4.16 (1 H, d, J = 8.2 Hz), 3.45 (1 H, d, J = 8.7 Hz), 2.57-2.68 (1 H, m), 2.39-2.45 (1 H, m), 2.24-2.38 (2 H, m).

MS (ESI) m/z: 424.1 (M+H)⁺.

[0588] The following Intermediates were prepared by General Procedure U (Table 50).

[0589] General Procedure U

Chloroacetyl chloride (1.1 eq.) was added dropwise to a biphasic solution of substrate (1.0 eq.) in dichloromethane and 0.5 N aq. NaOH (2.0 eq.) at 0 °C. The reaction mixture was warmed up to ambient temperature and stirred until complete reaction. The mixture was extracted with dichloromethane 3 times, and the combined

extracts were dried over Na₂SO₄ and concentrated under reduced pressure to afford the following Intermediates.

[0590] [Table 50]

Intermediates	Structure	Chemical Name	Substrate
I-i-1		(5R,8S)-8-((2-chloroacetamido)methyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	
I-i-2		(5S,8S)-8-((2-chloroacetamido)methyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	
I-i-3		(5S,8R)-8-((2-chloroacetamido)methyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide	

[0591] IM I-i-1

MS (ESI) m/z: 474.1 (M+H)⁺.

[0592] IM I-i-2

MS (ESI) m/z: 474.1 (M+H)⁺.

[0593] IM I-i-3

MS (ESI) m/z: 474.1 (M+H)⁺.

[0594] The following Examples were prepared by General Procedure V (Table 51).

[0595] General Procedure V

To a solution of substrate (1.0 eq.) in 50% dichloromethane/2-propanol was added portion wise tert-BuOK (4.0 eq.) at 0 °C. The solution was allowed to warm to ambient temperature and stirred until complete reaction. The solvent of the reaction mixture was removed by evaporation. The crude residue was purified by silica gel column chromatography to afford the following Examples.

[0596]

[Table 51]

Examples	Structure	Chemical Name	Substrate
293		(2S,5'R)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide	 IM I-i-1
294		(2S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide	 IM I-i-2
295		(2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide	 IM I-i-3

[0597] Ex 293

¹H NMR (CDCl₃) delta 8.64-8.72 (1 H, m), 7.54 (1 H, d, J = 7.5 Hz), 7.27-7.47 (4 H, m), 7.09-7.25 (1 H, m), 4.56-4.70 (2 H, m), 4.34 (1 H, dd, J = 12.2, 2.3 Hz), 4.11-4.23 (2 H, m), 3.37 (1 H, dd, J = 12.2, 3.1 Hz), 2.77-2.99 (1 H, m), 2.40-2.51 (1 H, m), 2.09-2.28 (2 H, m).

MS (ESI) m/z: 438.1 (M+H)⁺.

[0598] Ex 294

¹H NMR (CDCl₃) delta 8.70 (1 H, dt, J = 4.7, 1.6 Hz), 7.57 (1 H, d, J = 7.6 Hz), 7.46 (1 H, d, J = 2.0 Hz), 7.19-7.35 (6 H, m), 6.04 (1 H, br s), 4.53-4.64 (2 H, m), 4.39 (1 H, d, J = 16.8 Hz), 4.28 (1 H, d, J = 15.2 Hz), 4.19 (1 H, dd, J = 12.5, 1.5 Hz), 3.43 (1 H, dd, J = 12.5, 3.9 Hz), 2.56-2.64 (1 H, m), 2.29-2.53 (3 H, m).

MS (ESI) m/z: 438.1 (M+H)⁺.

[0599] Ex 295

¹H NMR (CDCl₃) delta 8.66 (1 H, dt, J = 4.7, 1.8 Hz), 7.54 (1 H, d, J = 7.6 Hz), 7.46 (1 H, d, J = 2.1 Hz), 7.39 (1 H, d, J = 8.2 Hz), 7.27-7.33 (2 H, m), 7.13-7.25 (1 H, m),

6.10 (1 H, br s), 4.58-4.69 (2 H, m), 4.34 (1 H, dd, $J = 12.2, 2.3$ Hz), 4.17 (2 H, dd, $J = 16.8, 15.5$ Hz), 3.37 (1 H, dd, $J = 12.2, 3.0$ Hz) 2.77-2.95 (1 H, m), 2.41-2.49 (1 H, m), 2.08-2.30 (2 H, m).

MS (ESI) m/z : 438.1 (M+H)⁺.

[0600] The following Examples were prepared by General Procedure W or X (Table 52).

[0601] General Procedure W









To a solution of substrate (1.0 eq.) in dichloromethane was added mCPBA (1.05 eq.) at 0 °C. The mixture was stirred at 0 °C until complete reaction, 1:1 mixture of aq. Na₂S₂O₃ and aq. NaHCO₃ was added to the mixture. The resulting mixture was extracted with CH₂Cl₂ and the extract was dried over Na₂SO₄. The extract was concentrated in vacuo, and the resulting residue was purified by silica gel column chromatography and preparative HPLC to afford the following Examples.

[0602] General Procedure X





To a solution of substrate (1.0 eq.) in dichloromethane was added mCPBA (2.5 eq.) at 0 °C. The mixture was stirred at 0 °C until complete reaction, 1:1 mixture of aq. Na₂S₂O₃ and aq. NaHCO₃ was added to the mixture. The resulting mixture was extracted with CH₂Cl₂ and the extract was dried over Na₂SO₄. The extract was concentrated in vacuo, and the resulting residue was purified by silica gel column chromatography and preparative HPLC to afford the following Examples.

[0603]

[Table 52-1]

Examples	Structure	Chemical Name	Substrate	General Procedure
296		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 Ex 288	W
297		(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM 1-h-2-1	X
298		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 Ex 287	X
299		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 Ex 288	X

[Table 52-2]

300		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		X
301		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		X

[0604] [Table 53]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
296	D	1.29	429.0	299	D	1.44	445.0
297	D	1.58	461.0	300	D	1.47	491.0
298	D	1.56	461.0	301	D	1.36	475.0

[0605] Ex 296

¹H NMR (CDCl₃) delta 8.65-8.60 (1 H, m), 7.56-7.51 (1 H, m), 7.43-7.37 (1 H, m), 7.31-7.26 (1 H, m), 7.24-7.17 (2 H, m), 7.02-6.97 (1 H, m), 4.68-4.54 (2 H, m), 4.08 (1 H, br), 3.77 (0.5 H, d, J = 12.8 Hz), 3.58 (0.5 H, d, J = 13.5 Hz), 3.35 (0.5 H, d, J = 12.8 Hz), 3.11 (0.5 H, d, J = 13.5 Hz), 2.86 (0.5 H, m), 2.74 (1.5 H, s), 2.69 (1.5 H, s), 2.69 (0.5 H, m), 2.61-2.46 (2 H, m), 2.32-2.19 (1 H, m).

[0606] Ex 297

¹H NMR (CDCl₃) delta 8.62 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.57 (1 H, dd, J = 7.9, 1.2 Hz), 7.45 (1 H, d, J = 1.8 Hz), 7.35-7.31 (2 H, m), 7.27 (1 H, dd, J = 7.9, 4.9 Hz), 7.19 (1 H, br d, J = 4.9 Hz), 4.62 (1 H, dd, J = 14.7, 6.1 Hz), 4.573 (1 H, s), 4.566 (1 H, dd, J = 14.7, 6.1 Hz), 3.55 (1 H, d, J = 16.5 Hz), 3.51 (1 H, d, J = 16.5 Hz), 3.14 (3 H, s), 3.07 (1 H, m), 2.63 (1 H, m), 2.40 (1 H, m), 2.18 (1 H, m).

[0607] Ex 298

¹H NMR (CDCl₃) delta 8.62 (1 H, br d, J = 4.9 Hz), 7.55 (1 H, dd, J = 7.9, 1.2 Hz), 7.45 (1 H, d, J = 1.8 Hz), 7.35 (1 H, d, J = 7.9 Hz), 7.30 (1 H, dd, J = 7.9, 4.9 Hz), 7.26 (1 H, dd, J = 7.9, 1.8 Hz), 7.21 (1 H, br), 4.64 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 4.12 (1 H, d, J = 15.3 Hz), 4.03 (1 H, br), 3.37 (1 H, d, J = 15.3 Hz), 3.11 (3 H, s), 2.79-2.64 (3 H, m), 2.24 (1 H, m).

[0608] Ex 299

¹H NMR (CDCl₃) delta 8.62 (1 H, br d, J = 4.9 Hz), 7.55 (1 H, d, J = 7.9 Hz), 7.40 (1 H, dd, J = 8.6, 6.1 Hz), 7.30 (1 H, dd, J = 7.9, 4.9 Hz), 7.21 (1 H, br), 7.19 (1 H, dd, J = 8.6, 2.4 Hz), 7.00 (1 H, ddd, J = 8.6, 7.9, 1.8 Hz), 4.64 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 4.12 (1 H, d, J = 15.3 Hz), 4.02 (1 H, br), 3.37 (1 H, d, J = 15.3 Hz), 3.11 (3 H, s), 2.79-2.64 (3 H, m), 2.24 (1 H, m).

[0609] Ex 300

¹H NMR (CDCl₃) delta 8.58 (1 H, br s), 7.53 (1 H, d, J = 7.9 Hz), 7.44 (1 H, br s), 7.34-7.22 (4 H, m), 4.72 (1 H, br), 4.62 (1 H, dd, J = 15.3, 6.1 Hz), 4.34 (1 H, dd, J = 15.3, 6.1 Hz), 4.34 (1 H, d, J = 15.3 Hz), 4.09 (2 H, br), 3.62 (1 H, m), 3.43 (1 H, d, J = 15.3 Hz), 3.42 (1 H, br s), 3.25 (1 H, m), 2.77-2.66 (3 H, m), 2.21 (1 H, m).

[0610] Ex 301

¹H NMR (CDCl₃) delta 8.60 (1 H, br s), 7.54 (1 H, d, J = 7.9 Hz), 7.39 (1 H, dd, J = 7.9, 6.1 Hz), 7.29 (1 H, m), 7.26 (1 H, br), 7.19 (1 H, dd, J = 8.6, 2.4 Hz), 7.00 (1 H, dd, J = 8.6, 7.9 Hz), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.55 (1 H, dd, J = 14.7, 6.1 Hz), 4.54 (1 H, br), 4.34 (1 H, d, J = 15.3 Hz), 4.12 (2 H, br), 3.62 (1 H, m), 3.46 (1 H, d, J = 15.3 Hz), 3.44 (1 H, br s), 3.29 (1 H, m), 3.03-2.71 (3 H, m), 2.22 (1 H, m).

[0611] The following Examples were prepared by General Procedure Y or Z (Table 54).

[0612] General Procedure Y









To stirred solution of the substrate (1.0 eq.) in MeOH-tert-BuOH-water (1:1:1) was added AD-Mix alpha and/or beta (4 times weight of substrate) at ambient temperature. The mixture was stirred at room temperature until complete reaction, aq. Na₂S₂O₃ was added to the mixture and then the mixture was stirred for 2 h. The mixture was extracted with CH₂Cl₂ twice and the extracts were combined. The extracts were dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified and each diastereo isomer was separated by silica gel column chromatography to afford the following Examples.

[0613] General Procedure Z

A mixture of substrate (1.0 eq.) and 2 N aq. NaOH (15.0 eq.) in 50% 1,4-dioxane-water (0.02 M) was heated at 65 °C until complete reaction. The mixture was extracted with CH₂Cl₂ twice and the extracts were combined. The extracts were dried over Na₂SO₄ and concentrated in vacuo to afford a glass. The residual glass was purified by silica gel column chromatography and preparative HPLC to afford following Examples.

[0614]

[Table 54-1]

Examples	Structure	Chemical Name	Substrate	General Procedure
302		(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Y
303		(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	IM I-e-26	
304		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Y
305		(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	IM I-e-28	
306		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Y

[Table 54-2]

307		(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
308		(5S,8S)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Y
309		(5S,8R)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	IM I-e-33	
310		(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Y
311		(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	IM I-e-29	













[Table 54-3]

312		(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-e-30	Y
313		(5S,8R)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide			
314		(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-e-32	Y
315		(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide			
316		(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-14	Z

[Table 54-4]

317		(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-16	Z
318		(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-19	Z
319		(5S,8S)-N-(2,4-dichloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-23	Z
320		(5S,8S)-N-(2,3-dichloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-24	Z
321		(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-25	Z

[Table 54-5]

322		(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Z
323		(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Z
324		(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Z
325		(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide		Z
326		(5S,8S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Z
327		(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		Z

[Table 54-6]

328		(5S,8R)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-32	Z
329		(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-34	Z
330		(5S,8S)-N-(2,4-dichlorophenethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-35	Z
331		(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-36	Z
332		(5S,8S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		IM I-g-38	Z

[0615]

[Table 55]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
302	C	1.42	398.8	318	C	1.22	366.9
303	C	1.41	398.8	319	D	1.47	417.0
304	C	1.41	398.9	320	D	1.46	417.0
305	C	1.42	398.9	321	D	1.33	401.0
306	C	1.31	382.9	322	D	1.40	401.0
307	C	1.30	382.9	323	D	1.38	401.0
308	C	1.27	384.9	324	D	1.36	401.0
309	C	1.27	385.0	325	D	1.54	413.1
310	C	1.40	416.8	326	D	1.54	413.1
311	C	1.40	416.8	327	D	1.41	397.1
312	C	1.34	401.0	328	D	1.42	397.1
313	C	1.34	401.0	329	D	1.31	400.0
314	C	1.32	382.8	330	D	1.51	413.0
315	C	1.32	382.8	331	D	1.59	425.0
316	C	1.28	428.9	332	D	1.47	483.0
317	D	1.18	413.0				

[0616] Ex 302

¹H NMR (CDCl₃) delta 8.54 (1 H, br dd, J = 4.9, 1.8 Hz), 7.53 (1 H, br dd, J = 7.9, 1.8 Hz), 7.45 (1 H, d, J = 1.8 Hz), 7.38 (1 H, d, J = 8.6 Hz), 7.29 (1 H, dd, J = 7.9, 4.9 Hz), 7.20-7.14 (2 H, m), 5.00 (1 H, br), 4.66 (1 H, dd, J = 14.7, 6.1 Hz), 4.60 (1 H, dd, J = 14.7, 6.1 Hz), 3.99 (1 H, d, J = 11.6 Hz), 3.76-3.64 (2 H, m), 2.88 (1 H, m), 2.24-1.99 (3 H, m).

[0617] Ex 303

¹H NMR (DMSO d₆) delta 9.16 (1 H, br), 8.64 (1 H, br d, J = 4.9 Hz), 7.66 (1 H, d, J = 7.9 Hz), 7.62 (1 H, d, J = 2.4 Hz), 7.46 (1 H, dd, J = 7.9, 2.4 Hz), 7.39 (1 H, dd, J = 7.9, 4.9 Hz), 7.36 (1 H, J = 7.9 Hz), 4.93 (1 H, s), 4.75 (1 H, br), 4.44 (1 H, dd, J = 15.9, 6.1 Hz), 4.38 (1 H, dd, J = 15.9, 6.1 Hz), 3.68 (1 H, dd, J = 11.0, 6.1 Hz), 3.56 (1 H, dd, J = 11.0, 4.3 Hz), 2.54-2.22 (3 H, m), 1.87 (1 H, m).

[0618] Ex 304

¹H NMR was identified with the Example 302.

[0619] Ex 305

¹H NMR was identified with the Example 303.

[0620] Ex 306

¹H NMR (CDCl₃) delta 8.54 (1 H, br dd, J = 4.5, 1.8 Hz), 7.53 (1 H, br dd, J = 7.9, 1.8 Hz), 7.43 (1 H, dd, J = 8.6, 6.1 Hz), 7.28 (1 H, br dd, J = 4.9, 1.2 Hz), 7.19 (1 H,

dd, J = 8.6, 2.4 Hz), 7.17 (1 H, br), 7.00 (1 H, dt, J = 7.9, 2.4 Hz), 5.01 (1 H, br d, J = 9.8 Hz), 4.66 (1 H, dd, J = 14.7, 6.1 Hz), 4.60 (1 H, dd, J = 14.7, 5.5 Hz), 3.99 (1 H, d, J = 11.6 Hz), 3.79-3.63 (2 H, m), 2.88 (1 H, m), 2.36-1.99 (3 H, m).

[0621] Ex 307

¹H NMR (CDCl₃) delta 8.57 (1 H, br dd, J = 4.3, 1.8 Hz), 7.56 (1 H, d, J = 7.9 Hz), 7.39 (1 H, dd, J = 8.5, 6.1 Hz), 7.28 (1 H, dd, J = 7.9, 4.3 Hz), 7.19 (1 H, dd, J = 8.5, 2.4 Hz), 7.16 (1 H, br), 6.99 (1 H, ddd, J = 8.5, 7.9, 2.4 Hz), 4.62 (1 H, dd, J = 14.7, 6.1 Hz), 4.55 (1 H, dd, J = 14.7, 6.1 Hz), 4.01 (1 H, br), 3.85 (1 H, d, J = 11.6 Hz), 3.79 (1 H, br), 3.71 (1 H, d, J = 11.6 Hz), 2.54-2.36 (3 H, m), 2.02 (1 H, m).

[0622] Ex 308

¹H NMR (CDCl₃) delta 8.54 (1 H, br dd, J = 4.5, 1.8 Hz), 7.53 (1 H, br dd, J = 7.9, 1.8 Hz), 7.30 (1 H, dd, J = 7.9, 4.9 Hz), 7.20 (1 H, br d, J = 4.9 Hz), 7.13 (1 H, m), 6.98 (1 H, m), 5.02 (1 H, br), 4.64 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 3.97 (1 H, d, J = 11.6 Hz), 3.85 (1 H, br), 3.68 (1 H, d, J = 11.6 Hz), 2.87 (1 H, m), 2.23-1.99 (3 H, m).

[0623] Ex 309

¹H NMR (CDCl₃) delta 8.57 (1 H, br dd, J = 4.9, 1.2 Hz), 7.56 (1 H, br dd, J = 7.9, 1.2 Hz), 7.29 (1 H, dd, J = 7.9, 4.9 Hz), 7.16 (1 H, br), 7.09 (1 H, m), 6.97 (1 H, m), 4.60 (1 H, dd, J = 15.3, 6.1 Hz), 4.53 (1 H, dd, J = 15.3, 6.1 Hz), 4.01 (1 H, br), 3.84 (1 H, d, J = 11.0 Hz), 3.82 (1 H, br), 3.71 (1 H, d, J = 11.6 Hz), 2.44-2.04 (3 H, m), 2.04 (1 H, m).

[0624] Ex 310

¹H NMR (CDCl₃) delta 8.53 (1 H, m), 7.56 (1 H, d, J = 7.9 Hz), 7.33-7.28 (2 H, m), 7.11 (1 H, dd, J = 9.2, 1.8 Hz), 7.02 (1 H, br), 5.01 (1 H, br), 4.78 (1 H, dd, J = 14.7, 5.5 Hz), 4.66 (1 H, dd, J = 14.7, 5.5 Hz), 3.97 (1 H, d, J = 11.6 Hz), 3.76 (1 H, br), 3.68 (1 H, d, J = 11.6 Hz), 2.88 (1 H, m), 2.23-1.98 (3 H, m).

[0625] Ex 311

¹H NMR (CDCl₃) delta 8.56 (1 H, dd, J = 4.9, 1.2 Hz), 7.57 (1 H, dd, J = 7.9, 1.2 Hz), 7.29-7.27 (2 H, m), 7.11 (1 H, dd, J = 9.2, 2.4 Hz), 7.04 (1 H, br), 4.73 (1 H, dd, J = 14.7, 6.1 Hz), 4.63 (1 H, dd, J = 14.7, 5.5 Hz), 4.00 (1 H, br s), 3.84 (1 H, d, J = 11.6 Hz), 3.79 (1 H, br), 3.70 (1 H, d, J = 11.6 Hz), 2.52-2.37 (3 H, m), 2.02 (1 H, m).

[0626] Ex 312

¹H NMR (CDCl₃) delta 8.53 (1 H, dd, J = 4.9, 1.8 Hz), 7.52 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.29 (1 H, dd, J = 7.9, 4.9 Hz), 7.25-7.08 (3 H, m), 4.99 (1 H, br d, J = 10.4 Hz), 4.67 (1 H, dd, J = 14.7, 6.1 Hz), 4.62 (1 H, dd, J = 14.7, 5.5 Hz), 3.99 (1 H, d, J = 10.4 Hz), 3.71 (1 H, br s), 3.67 (1 H, d, J = 10.4 Hz), 2.88 (1 H, m), 2.24-1.99 (3 H, m).

[0627] Ex 313

¹H NMR (CDCl₃) delta 8.57 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.56 (1 H, dd, J = 7.9, 1.8 Hz), 7.29 (1 H, dd, J = 7.9, 4.9 Hz), 7.19-7.08 (3 H, m), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.57 (1 H, dd, J = 14.7, 5.5 Hz), 3.99 (1 H, s), 3.85 (1 H, d, J = 11.0 Hz), 3.76 (1 H, br s), 3.72 (1 H, d, J = 11.0 Hz), 2.50-2.40 (3 H, m), 2.03 (1 H, m).

[0628] Ex 314

¹H NMR (CDCl₃) delta 8.53 (1 H, br dd, J = 4.9, 1.8 Hz), 7.53 (1 H, br dd, J = 7.9, 1.8 Hz), 7.33 (1 H, t, J = 7.9 Hz), 7.29 (1 H, dd, J = 7.9, 4.9 Hz), 7.27-7.10 (2 H, m), 7.15 (1 H, d, J = 7.9 Hz), 5.02 (1 H, br), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.54 (1 H, dd, J = 14.7, 6.1 Hz), 3.98 (1 H, d, J = 11.6 Hz), 3.82 (1 H, br), 3.68 (1 H, d, J = 11.6 Hz), 2.87 (1 H, m), 2.18 (1 H, m), 2.09-1.99 (2 H, m).

[0629] Ex 315

¹H NMR (CDCl₃) delta 8.56 (1 H, br dd, J = 4.9, 1.8 Hz), 7.55 (1 H, d, J = 7.9 Hz), 7.35-7.27 (2 H, m), 7.18-7.09 (3 H, m), 4.57 (1 H, dd, J = 14.7, 6.1 Hz), 4.50 (1 H, dd, J = 14.7, 6.1 Hz), 4.06 (1 H, br), 3.83 (2 H, d, J = 11.0 Hz), 3.70 (1 H, d, J = 11.0 Hz), 2.52-2.34 (3 H, m), 2.04 (1 H, m).

[0630] Ex 316

¹H NMR (CDCl₃) delta 8.51 (1 H, br dd, J = 4.9, 1.8 Hz), 7.58 (1 H, br d, J = 5.5 Hz), 7.46 (1 H, br dd, J = 7.9, 1.8 Hz), 7.43 (1 H, d, J = 2.4 Hz), 7.34 (1 H, d, J = 2.4 Hz), 7.26 (1 H, dd, J = 7.9, 4.9 Hz), 5.03 (1 H, br), 4.82-4.64 (4 H, m), 3.97 (1 H, d, J = 11.0 Hz), 3.87 (2 H, br), 3.66 (1 H, d, J = 11.0 Hz), 2.81 (1 H, m), 2.19-1.95 (3 H, m).

[0631] Ex 317

¹H NMR (CDCl₃) delta 8.52 (1 H, br d, J = 4.9 Hz), 7.56 (1 H, br), 7.46 (1 H, dd, J = 7.9, 1.8 Hz), 7.25 (1 H, dd, J = 7.9, 4.9 Hz), 7.17 (1 H, dd, J = 8.6, 2.4 Hz), 7.10 (1 H, dd, J = 8.6, 2.4 Hz), 5.06 (1 H, br), 4.82-4.62 (4 H, m), 3.98 (1 H, d, J = 11.0 Hz), 3.81 (1 H, br), 3.66 (1 H, d, J = 11.0 Hz), 2.81 (1 H, m), 2.11 (1 H, m), 2.07-1.96 (2 H, m), 1.65 (1 H, br).

[0632] Ex 318

¹H NMR (CDCl₃) delta 8.53 (1 H, br dd, J = 4.9, 1.8 Hz), 7.53 (1 H, d, J = 7.9 Hz), 7.34 (1 H, m), 7.28 (1 H, dd, J = 7.9, 4.9 Hz), 7.16 (1 H, br), 6.90-6.85 (2 H, m), 5.30 (1 H, br), 4.62 (1 H, dd, J = 14.7, 6.1 Hz), 4.53 (1 H, dd, J = 14.7, 6.1 Hz), 3.97 (1 H, d, J = 11.6 Hz), 3.87 (1 H, br), 3.67 (1 H, d, J = 11.6 Hz), 2.88 (1 H, m), 2.23-1.99 (3 H, m).

[0633] Ex 319

¹H NMR (CDCl₃) delta 8.54 (1 H, br dd, J = 4.9, 1.8 Hz), 7.54 (1 H, br dd, J = 7.9, 1.8 Hz), 7.33 (1 H, dd, J = 8.6, 6.7 Hz), 7.30 (1 H, dd, J = 7.9, 4.9 Hz), 7.23 (1 H, br), 7.20 (1 H, dd, J = 8.6, 1.8 Hz), 4.99 (1 H, br d, J = 9.2 Hz), 4.68 (1 H, dd, J = 14.7, 6.1 Hz), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 4.00 (1 H, d, J = 11.0 Hz), 3.77 (1 H, br s), 3.68 (1 H, br), 2.87 (1 H, m), 2.24-1.99 (3 H, m).

[0634] Ex 320

¹H NMR (CDCl₃) delta 8.54 (1 H, br dd, J = 4.9, 1.8 Hz), 7.53 (1 H, br dd, J = 7.9, 1.8 Hz), 7.37 (1 H, dd, J = 8.6, 5.5 Hz), 7.29 (1 H, dd, J = 7.9, 4.9 Hz), 7.21 (1 H, br d, J = 6.1 Hz), 7.11 (1 H, dd, J = 8.6, 7.9 Hz), 5.00 (1 H, br), 4.68 (1 H, dd, J = 15.3, 6.1 Hz), 4.63 (1 H, dd, J = 14.7, 6.1 Hz), 3.99 (1 H, d, J = 11.0 Hz), 3.74 (1 H, br s), 3.68 (1 H, d, J = 11.0 Hz), 2.87 (1 H, m), 2.24-1.99 (3 H, m).

[0635] Ex 321

¹H NMR (CDCl₃) delta 8.54 (1 H, br dd, J = 4.9, 1.8 Hz), 7.57 (1 H, br dd, J = 7.9, 1.8 Hz), 7.29 (1 H, dd, J = 7.9, 4.9 Hz), 7.05 (1 H, ddd, J = 7.9, 2.4, 1.8 Hz), 6.99 (1 H, br), 6.85 (1 H, ddd, J = 9.8, 8.6, 2.4 Hz), 4.80 (1 H, br), 4.77 (1 H, dd, J = 14.7, 6.1 Hz), 4.66 (1 H, dd, J = 14.7, 6.1 Hz), 3.98 (1 H, d, J = 11.6 Hz), 3.74 (1 H, br), 3.68 (1 H, d, J = 11.6 Hz), 2.88 (1 H, m), 2.23-1.99 (3 H, m).

[0636] Ex 322

¹H NMR (CDCl₃) delta 8.55 (1 H, br dd, J = 4.9, 1.8 Hz), 7.54 (1 H, br dd, J = 7.9, 1.8 Hz), 7.30 (1 H, dd, J = 8.0, 5.5 Hz), 7.21-7.10 (3 H, m), 5.00 (1 H, br), 4.66 (1 H, dd, J = 14.7, 6.9 Hz), 4.59 (1 H, dd, J = 14.7, 6.9 Hz), 3.99 (1 H, d, J = 11.6 Hz), 3.78 (1 H, br), 3.68 (1 H, d, J = 11.6 Hz), 2.87 (1 H, m), 2.24-1.99 (3 H, m).

[0637] Ex 323

¹H NMR (CDCl₃) delta 8.55 (1 H, br dd, J = 4.3, 1.8 Hz), 7.53 (1 H, br dd, J = 7.9, 1.8 Hz), 7.34-7.28 (2 H, m), 7.16 (1 H, m), 6.99 (1 H, dt, J = 8.6, 1.8 Hz), 5.01 (1 H, br), 4.64 (1 H, dd, J = 14.7, 6.1 Hz), 4.50 (1 H, dd, J = 14.7, 6.1 Hz), 3.99 (1 H, d, J = 11.6 Hz), 3.77 (1 H, br), 3.68 (1 H, d, J = 11.6 Hz), 2.87 (1 H, m), 2.24-1.99 (3 H, m).

[0638] Ex 324

¹H NMR (CDCl₃) delta 8.54 (1 H, ddd, J = 4.3, 1.8, 1.2 Hz), 7.55 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.30 (1 H, dd, J = 7.9, 4.9 Hz), 7.02 (1 H, br), 7.01 (2 H, d, J = 7.3 Hz), 5.00 (1 H, br), 4.70 (1 H, dd, J = 14.7, 6.1 Hz), 4.60 (1 H, dd, J = 14.7, 5.5 Hz), 3.98 (1 H, d, J = 11.6 Hz), 3.71 (1 H, br), 3.68 (1 H, d, J = 11.6 Hz), 2.87 (1 H, m), 2.22-1.98 (3 H, m).

[0639] Ex 325

¹H NMR (CDCl₃) delta 8.35 (1 H, s), 7.46 (1 H, d, J = 2.4 Hz), 7.40 (1 H, d, J = 7.9 Hz), 7.28-7.25 (2 H, m), 7.19 (1 H, br d, J = 5.5 Hz), 5.10 (1 H, br), 4.67 (1 H, dd, J = 14.7, 6.1 Hz), 4.60 (1 H, dd, J = 14.7, 6.1 Hz), 3.97 (1 H, d, J = 11.6 Hz), 3.67 (1 H, d, J = 11.6 Hz), 3.65 (1 H, br s), 2.87 (1 H, m), 2.27 (3 H, s), 2.22-1.96 (3 H, m).

[0640] Ex 326

¹H NMR (CDCl₃) delta 8.54 (1 H, br dd, J = 4.3, 1.8 Hz), 7.47 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.44 (1 H, d, J = 1.8 Hz), 7.37 (1 H, d, J = 8.6 Hz), 7.30 (1 H, dd, J = 8.6, 1.8 Hz), 7.25 (1 H, dd, J = 7.9, 4.3 Hz), 7.12 (1 H, dd, J = 6.7, 6.1 Hz), 5.46 (1 H, dq, J = 7.3, 6.7 Hz), 3.98 (1 H, d, J = 11.6 Hz), 3.75 (1 H, br), 3.69 (1 H, d, J = 11.6 Hz), 3.48

(1 H, d, J = 1.8 Hz), 2.88 (1 H, m), 2.26-2.01 (3 H, m), 1.59 (3 H, d, J = 7.3 Hz).

[0641] Ex 327

¹H NMR (CDCl₃) delta 8.51 (1 H, ddd, J = 4.3, 1.8, 1.2 Hz), 7.48 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.41 (1 H, dd, J = 8.6, 6.1 Hz), 7.24 (1 H, dd, J = 7.9, 4.3 Hz), 7.18 (1 H, dd, J = 8.6, 2.4 Hz), 7.14 (1 H, br), 7.04 (1 H, ddd, J = 8.6, 7.9, 2.4 Hz), 5.48 (1 H, dq, J = 7.3, 6.7 Hz), 3.98 (1 H, d, J = 11.6 Hz), 3.69 (1 H, d, J = 11.6 Hz), 2.90 (1 H, m), 2.48 (2 H, br), 2.26-1.99 (3 H, m), 1.60 (3 H, d, J = 7.3 Hz).

[0642] Ex 328

¹H NMR (CDCl₃) delta 8.53 (1 H, ddd, J = 4.9, 1.8, 1.2 Hz), 7.53 (1 H, ddd, J = 7.9, 1.8, 1.2 Hz), 7.38 (1 H, dd, J = 8.6, 6.1 Hz), 7.24 (1 H, dd, J = 7.9, 4.9 Hz), 7.16 (1 H, dd, J = 8.6, 3.1 Hz), 7.10 (1 H, br), 7.02 (1 H, dt, J = 8.6, 3.1 Hz), 5.39 (1 H, m), 4.00 (1 H, br), 3.83 (1 H, d, J = 11.6 Hz), 3.69 (1 H, d, J = 11.6 Hz), 2.56-2.38 (3 H, m), 2.03 (1 H, m), 1.66 (1 H, br), 1.59 (3 H, d, J = 7.3 Hz).

[0643] Ex 329

¹H NMR (CDCl₃) delta 8.55 (1 H, br d, J = 4.3 Hz), 8.48 (1 H, d, J = 2.4 Hz), 8.08 (1 H, br d, J = 4.3 Hz), 7.79 (1 H, dd, J = 7.9, 1.8 Hz), 7.78 (1 H, d, J = 2.4 Hz), 7.31 (1 H, dd, J = 7.9, 4.3 Hz), 4.83 (1 H, dd, J = 18.3, 4.9 Hz), 4.70 (1 H, dd, J = 18.3, 4.3 Hz), 4.02 (1 H, d, J = 11.6 Hz), 3.72 (1 H, d, J = 11.6 Hz), 2.94 (1 H, m), 2.32-2.02 (3 H, m). The signals due to OH were not observed.

[0644] Ex 330

¹H NMR (CDCl₃) delta 8.54 (1 H, br dd, J = 4.9, 1.8 Hz), 7.42 (1 H, d, J = 1.8 Hz), 7.36 (1 H, br dd, J = 7.9, 1.8 Hz), 7.28-7.20 (3 H, m), 6.86 (1 H, br d, J = 5.5 Hz), 5.00 (1 H, br), 3.98 (1 H, d, J = 11.0 Hz), 3.77-3.62 (2 H, m), 3.68 (1 H, d, J = 11.0 Hz), 3.48 (1 H, s), 3.07-3.03 (2 H, m), 2.83 (1 H, m), 2.18-1.98 (3 H, m).

[0645] Ex 331







¹H NMR (CDCl₃) delta 8.57-8.54 (1 H, m), 7.63 (1 H, br d, J = 7.9, 1.8 Hz), 7.39 (1 H, d, J = 1.8 Hz), 7.33 (1 H, dd, J = 7.9, 4.9 Hz), 7.20-7.11 (2 H, m), 7.03 (1 H, br), 4.90 (1 H, br), 3.99 (1 H, d, J = 11.6 Hz), 3.74 (1 H, br), 3.70 (1 H, d, J = 11.6 Hz), 3.08 (1 H, m), 2.90 (1 H, m), 2.3-2.01 (4 H, m), 1.42-1.29 (2 H, m).

[0646] The following Intermediates were prepared by General Procedure AA (Table 56).

[0647] General Procedure AA

To a stirred solution of substrate (1.0 eq.), iodomethane (15 eq.), and silver(I)oxide (10 eq.) in THF was added one drop of dimethyl sulfide at ambient temperature in the dark. The mixture was stirred at room temperature for 1 h, and then heated to 50 °C. After being stirred at 50 °C until complete reaction, the mixture was cooled to room temperature. The insoluble material was removed by filtration, the resulting filtrate was concentrated in vacuo. The residue was purified by silica gel column chromatography to afford following Intermediates.

[0648] [Table 56]

Intermediates	Structure	Chemical Name	Substrate
II-s-1		(5R,8S)-methyl 5-fluoro-8-methoxy-5,6,7,8-tetrahydr oquinoline-5-carboxylate	 IM II-d-3
II-s-2		(5S,8S)-methyl 5-fluoro-8-methoxy-5,6,7,8-tetrahydr oquinoline-5-carboxylate	 IM II-d-4
II-s-3		(5S,8R)-methyl 5-fluoro-8-methoxy-5,6,7,8-tetrahydr oquinoline-5-carboxylate	 IM II-d-5

[0649] IM II-s-1

^1H NMR (CDCl_3) delta 8.69 (1 H, d, $J = 4.6$ Hz), 7.79 (1 H, d, $J = 7.9$ Hz), 7.31 (1 H, dd, $J = 7.9, 4.6$ Hz), 4.42 (1 H, m), 3.79 (3 H, s), 3.57, (3 H, s), 2.55 (1 H, m), 2.43-2.24 (3 H, m).

MS (ESI) m/z : 240.2 ($\text{M}+\text{H}$) $^+$.

[0650] IM II-s-2

^1H NMR (CDCl_3) delta 8.69 (1 H, br dd, $J = 4.6, 1.3$ Hz), 7.68 (1 H, dd, $J = 7.9, 1.3$ Hz), 7.32 (1 H, dd, $J = 7.9, 4.6$ Hz), 4.41 (1 H, m), 3.82 (3 H, s), 3.54 (3 H, s), 2.78 (1 H, m), 2.36-2.04 (3 H, m).

MS (ESI) m/z : 240.1 ($\text{M}+\text{H}$) $^+$.

IM II-s-3

^1H NMR (CDCl_3) delta 8.69 (1 H, br dd, $J = 4.6, 1.3$ Hz), 7.79 (1 H, dd, $J = 7.9, 1.3$ Hz), 7.31 (1 H, dd, $J = 7.9, 4.6$ Hz), 4.42 (1 H, m), 3.79 (3 H, s), 3.57, (3 H, s), 2.55 (1 H, m), 2.43-2.24 (3 H, m).

MS (ESI) m/z: 240.1 (M+H)⁺.

[0651] The following Examples were prepared by General Procedure A (Table 57).

[0652] [Table 57-1]

Examples	Structure	Chemical Name	Substrate	Amine
333		(5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
334		(5S,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
335		(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
336		(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
337		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[Table 57-2]

338		(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
339		(5S,8S)-5-fluoro-8-methoxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
340		(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		
341		(5S,8S)-5-fluoro-8-methoxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide		
342		(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[0653] [Table 58]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
333	A	1.45	413.1	338	B	1.63	417.0
334	A	1.44	413.1	339	C	1.46	368.9
335	B	1.58	400.9	340	C	1.52	366.9
337	B	1.54	383.0	341	C	1.42	368.9
337	B	1.47	367.1	342	C	1.41	351.0

[0654] Ex 333

¹H NMR (CDCl₃) delta 8.66 (1 H, br d, J = 4.6 Hz), 7.57 (1 H, br d, J = 5.9 Hz), 7.49

(1 H, d, J = 7.9 Hz), 7.44 (1 H, d, J = 2.0 Hz), 7.33 (1 H, d, J = 2.0 Hz), 7.23 (1 H, dd, J = 7.9, 4.6 Hz), 4.73-4.67 (4 H, m), 4.43 (1 H, m), 3.90 (1 H, br), 3.57 (3 H, s), 2.80-2.11 (4 H, m).


[0655] Ex 334

¹H NMR was identified with the Example 333.

[0656] Intermediate (IM) II-s-4:

(5S,8S)-allyl 8-(allyloxy)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxylate

[0657] [Chem.31]



[0658] A mixture of (5S,8S)-methyl

5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxylate (100 mg, 0.444 mmol, IM II-d-4) and 2 N aq. NaOH in MeOH (2.0 mL) was stirred at room temperature for 1 h, the resulting solution was concentrated in vacuo. The residue was added NMP (1.0 mL) and the mixture was added sodium hydride (60% oil dispersant, 5 mg, 0.222 mmol) at 0 °C. The mixture was warmed to room temperature and stirred for 20 min. Allyl bromide (0.192 mL, 2.22 mmol) was added to the reaction mixture and the mixture was heated at 60 °C for 6 h. The mixture was cooled to room temperature and stirred further 9 h. Water was added to the mixture and the mixture was extracted with EtOAc. The extract was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by silica gel column chromatography (0- to 50% EtOAc/n-hexane, gradient) to afford 25 mg (19%) of the title compound.

[0659] ¹H NMR (CDCl₃) delta 8.67 (1 H, br dd, J = 4.6, 1.8 Hz), 7.68 (1 H, br dd, J = 7.9, 1.8 Hz), 7.31 (1 H, dd, J = 7.9, 4.9 Hz), 5.95 (1 H, ddt, J = 17.1, 10.4, 5.5 Hz), 5.88 (1 H, ddt, J = 17.1, 10.4, 5.5 Hz), 5.32 (1 H, ddt, J = 17.1, 4.9, 1.8 Hz), 5.30 (1 H, ddt, J = 17.1, 3.1, 1.2 Hz), 5.25 (1 H, br d J = 10.4 Hz), 5.17 (1 H, ddt, J = 10.4, 3.1, 1.2 Hz), 4.76-4.66 (2 H, m), 4.56 (1 H, m), 4.30-4.21 (2 H, m), 2.89 (1 H, m), 2.33-2.15 (3 H, m).


MS (ESI) m/z: 291.9 (M+H)⁺.

[0660] Intermediate (IM) I-s-1:

(5R,8R)-8-(allyloxy)-N-(2,4-dichlorobenzyl)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide

[0661]

[Chem.32]



[0662] 60% oil dispersant NaH (6 mg, 0.244 mmol) was added to a solution of (5R,8R)-methyl 5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxylate (25 mg, 0.111 mmol, IM II-d-6) in THF (1.0 mL) under N₂ atmosphere at 0 °C. After being stirred at 0 °C, allyl iodide (0.025 mL, 0.278 mmol) was added to the mixture and the mixture was stirred for 1 h. 60% oil dispersant NaH (6 mg, 0.244 mmol) and allyl iodide (0.025 mL, 0.278 mmol) was added to the mixture and the mixture was stirred further 1 h. Water was added to the mixture and acidified with 2 N hydrochloric acid. The mixture was concentrated in vacuo. The resulting residue was diluted with MeCN and toluene and the mixture was concentrated in vacuo. This procedure was repeated 3 times to remove remaining water. The residue was dissolved in 25% MeOH-THF (2.0 mL) and 2,4-dichlorobenzylamine (20 mg, 0.111 mmol), triethylamine (0.025 mL, 0.179 mmol), and DMT-MM (40 mg, 0.167 mmol) were added to the mixture. After being stirred at room temperature for 16 h, water was added to the mixture. The mixture was extracted with EtOAc and washed with brine. The extract was dried over Na₂SO₄ and concentrated in vacuo to afford glass. The residual glass was purified by preparative TLC (70% EtOAc/n-hexane) to afford 15 mg (33%) of the title compound as a white solid.

¹H NMR (CDCl₃) delta 8.64 (1 H, br dd, J = 4.9, 1.8 Hz), 7.51 (1 H, br dd, J = 7.9, 1.8 Hz), 7.44 (1 H, d, J = 1.8 Hz), 7.37 (1 H, br d, J = 7.9 Hz), 7.27-7.22 (2 H, m), 7.16 (1 H, br), 5.92 (1 H, ddt, J = 17.1, 10.4, 5.5 Hz), 5.29 (1 H, ddt, J = 17.1, 3.7, 1.8 Hz), 5.14 (1 H, ddt, J = 17.1, 3.7, 1.8 Hz), 4.71-4.51 (3 H, m), 4.27-4.17 (2 H, m), 2.92 (1 H, m), 2.33-2.06 (3 H, m).

MS (ESI) m/z: 408.7 (M+H)⁺.

[0663] The following Intermediates were prepared from IM II-s-4 by General Procedure A (Table 59).

[0664]

[Table 59]

Intermediates	Structure	Chemical Name	Amine
I-s-2		(5S,8S)-8-(allyloxy)-N-(2-(((tert-butyl dimethylsilyl)oxy)methyl)-4,6-dichlorobenzyl)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide	
I-s-3		(5S,8S)-8-(allyloxy)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide	
I-s-4		(5S,8S)-8-(allyloxy)-5-fluoro-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	
I-s-5		(5S,8S)-8-(allyloxy)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide	
I-s-6		(5S,8S)-8-(allyloxy)-N-(2,4-difluorobenzyl)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide	
I-s-7		(5S,8S)-8-(allyloxy)-N-(2,4-dichlorobenzyl)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide	

[0665] IM I-s-2
MS (ESI) m/z: 552.4 (M+H)⁺.

[0666] IM I-s-3

MS (ESI) m/z: 392.7 (M+H)⁺.

[0667] IM I-s-4

MS (ESI) m/z: 394.8 (M+H)⁺.

[0668] IM I-s-5

MS (ESI) m/z: 392.8 (M+H)⁺.

[0669] IM I-s-6

MS (ESI) m/z: 376.8 (M+H)⁺.

[0670] IM I-s-7

MS (ESI) m/z: 408.7 (M+H)⁺.

[0671] The following Examples were prepared by General Procedure C (Table 60).

In preparation of Example 343, the TBS group was deprotected on the SCX cartridge column.

[0672]

[Table 60]

Examples	Structure	Chemical Name	Substrate
343		(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-s-2
344		(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-s-3
345		(5S,8S)-5-fluoro-8-(2-hydroxyethoxy)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-s-4
346		(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-s-5
347		(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-s-6
348		(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM I-s-1

[0673]


[Table 61]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
343	C	1.37	442.9	346	C	1.41	397.1
344	C	1.39	397.0	347	C	1.31	381.0
345	C	1.36	399.0	348	C	1.49	413.0

[0674] Example 349:

(5S,8S)-N-(2,4-dichlorobenzyl)-8-(2,3-dihydroxypropoxy)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide

[0675] [Chem.33]



[0676] A mixture of AD-Mix beta (250 mg) in 50% tert-BuOH-water (3.0 mL) was stirred at room temperature until the two clear phases formed. After cooling to 0 °C, a solution of (5S,8S)-8-(allyloxy)-N-(2,4-dichlorobenzyl)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide (24 mg, 0.059 mmol, IM I-s-7) in THF (1.0 mL) was added to the mixture. The resulting mixture was stirred at 0 °C overnight. The mixture was filtered through a pad of celite, the filter cake was washed with THF. The filtrate was evaporated in vacuo to remove the volatile. The residue was diluted with brine and extracted with EtOAc twice. The extracts were combined and dried over Na₂SO₄. After removal of the solvent, the residue was purified by silica gel column chromatography (50%EtOAc/n-hexane then 5% MeOH/EtOAc) to afford 24 mg (92%) of the title compound.

[0677] ¹H NMR (CDCl₃) delta 8.61 (1 H, m), 7.55 (1 H, m), 7.50 (1 H, d, J = 1.8 Hz), 7.36 (1 H, br d, J = 7.9 Hz), 7.31-7.23 (2 H, m), 7.20 (1 H, br d, J = 5.5 Hz), 4.67-4.56 (3 H, m), 3.92-3.52 (5 H, m), 2.82 (1 H, m), 2.7 (1 H, br), 2.36-2.07 (3 H, m), 1.8 (1 H, br).
LCMS (ESI) m/z: 443.2 (M+H)⁺, tR 1.45 min (Method D).


[0678] Intermediate (IM) I-p-1:

Procedure: Scheme 13, Step 1

S-((5S)-5-((2,4-dichlorobenzyl)carbamoyl)-5-fluoro-5,6,7,8-tetrahydroquinolin-8-yl)

benzothioate

[Chem.34]



[0679] Bis(2-methoxyethyl) azodicarboxylate (57 mg, 0.244 mmol) was added to a solution of triphenylphosphine (64 mg, 0.244 mmol) in THF (2.0 mL) at 0 °C under N₂ atmosphere. The reaction mixture was stirred at 0 °C for 30 min, and then a solution of (5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide (60 mg, 0.163 mmol, Ex 133) and thiobenzoic acid (34 mg, 0.244 mmol) in THF (1.0 mL) was added dropwise to the mixture. After being stirred at 0 °C for 20 h, water was added to the mixture and the mixture was extracted with EtOAc. The extract was washed with brine, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography (20- to 35% EtOAc/n-hexane, gradient) to afford 25 mg (31%) of the title compound.

¹H NMR (CDCl₃) delta 8.66-8.63 (1 H, m), 8.01-7.96 (2 H, m), 7.60-7.17 (9 H, m), 5.30 (0.4 H, br), 5.24 (0.6 H, m), 4.75-4.56 (2 H, m), 2.87-2.14 (4 H, m).


MS (ESI) m/z: 488.6 (M+H)⁺.

[0680] Intermediate (IM) I-q-1:

Procedure: Scheme 13, Step 2

(5S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(methylthio)-5,6,7,8-tetrahydroquinoline-5-carboxamide

[0681] [Chem.35]



[0682] A solution of S-((5S)-5-((2,4-dichlorobenzyl)carbamoyl)-5-fluoro-5,6,7,8-tetrahydroquinolin-8-yl)benzothioate (25 mg, 0.051 mmol, IM I-p-1) in methanol (1.0 mL) was added 1 N aq.

NaOH (0.06 mL, 0.06 mmol) and dimethyl sulfate (8 mg, 0.06 mmol) at room temperature. The mixture was stirred at room temperature for 2 h and concentrated in vacuo. Water was added to the resulting residue and extracted with EtOAc. The extract was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by SCX column to afford 18 mg (88%) of the title compound as a glass.

[0683] ¹H NMR (CDCl₃) delta 8.67-8.58 (1 H, m), 7.57-7.14 (6 H, m), 4.68-4.53 (2 H, m), 4.18-4.14 (1 H, m), 3.49 (3 H, s), 3.03-2.86 (0.4 H, m), 2.72-2.64 (0.6 H, m), 2.59-2.10 (3 H, m).


MS (ESI) m/z: 400.5 (M+H)⁺.

[0684] Intermediate (IM) I-q-2:

Procedure: Scheme 13, Step 2

(5S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-((2-hydroxyethyl)thio)-5,6,7,8-tetrahydroquinoline-5-carboxamide

[0685] [Chem.36]



[0686] A solution of S-((5S)-5-((2,4-dichlorobenzyl)carbamoyl)-5-fluoro-5,6,7,8-tetrahydroquinolin-8-yl) benzothioate (24 mg, 0.049 mmol, IM I-p-1) in methanol (1.0 mL) was added 1 N aq. NaOH (0.06 mL, 0.06 mmol) and (2-bromoethoxy)(tert-butyl)dimethylsilane (18 mg, 0.074 mmol) at room temperature. The mixture was stirred at room temperature for 2 h and concentrated in vacuo. Water was added to the residue and the mixture was extracted with EtOAc. The extract was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The resulting residue was purified by SCX column to afford 16 mg (76%) of the title compound as a glass.

[0687] ¹H NMR (CDCl₃) delta 8.57 (0.5 H, dd, J = 4.9, 1.8 Hz), 8.56 (0.5 H, dd, J = 4.9, 1.8 Hz), 7.53 (0.5 H, d, J = 7.9 Hz), 7.49 (0.5 H, d, J = 7.9 Hz), 7.45 (1 H, d, J = 1.8 Hz), 7.39 (0.5 H, d, J = 7.9 Hz), 7.34 (0.5 H, d, J = 7.9 Hz), 7.31-7.14 (3 H, m), 4.68-4.53 (2 H, m), 4.47 (0.5 H, br), 4.29 (0.5 H, br), 3.95-3.86 (2 H, m), 3.48 (1 H, br), 3.05-2.68 (3 H, m), 2.53-2.12 (3 H, m).

MS (ESI) m/z: 428.6 (M+H)⁺.

[0688] General Procedure: Scheme 13, Step 3

The following Examples were prepared by General Procedure X (Table 62).

[0689] [Table 62]

Examples	Structure	Chemical Name	Substrate
350		(5S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(methylsulfonyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	
			IM 1-q-1
351		(5S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-((2-hydroxyethyl)sulfonyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide	
			IM 1-q-2

[0690] Ex 350

¹H NMR (CDCl₃) delta 8.66-8.60 (1 H, m), 7.68 (1 H, dd, J = 7.9, 1.8 Hz), 7.45 (0.5 H, d, J = 1.8 Hz), 7.39 (0.5 H, d, J = 1.8 Hz), 7.37-7.21 (3 H, m), 7.15 (0.5 H, br), 7.00 (0.5 H, br), 4.62-4.50 (2 H, m), 4.44 (1 H, m), 3.30 (1.5 H, s), 3.16 (1.5 H, s), 3.03-2.62 (3 H, m), 2.42-2.27 (1 H, m).

LCMS (ESI) m/z: 430.5 (M+H)⁺, tR 1.58 min (Method C).

[0691] Ex 351

¹H NMR (CDCl₃) delta 8.59 (1 H, dd, J = 4.9, 1.2 Hz), 7.72 (1 H, dd, J = 7.9, 1.2 Hz), 7.45 (1 H, d, J = 2.4 Hz), 7.41-7.22 (3 H, m), 7.14 (1 H, br), 4.80 (1 H, m), 4.70 (1 H, br), 4.62-4.49 (2 H, m), 4.24-4.17 (2 H, m), 3.95-3.89 (1 H, m), 3.42-3.36 (1 H, m), 3.07-2.66 (3 H, m), 2.43-2.30 (1 H, m).

LCMS (ESI) m/z: 460.8 (M+H)⁺, tR 1.50 min (Method C).



[0692] The following Intermediate was prepared by General Procedure AB (Table 63).

[0693] General Procedure AB

Deoxo-Fluor(Trademark) (2.0 eq.) was added to a solution of substrate (1.0 eq.) in CH₂Cl₂ (0.15 M) and stirred for 1 h. The mixture was poured into water and extracted with EtOAc. The extract was dried over Na₂SO₄ and concentrated in vacuo. The resulting residue was purified by silica gel column chromatography.

[0694]

[Table 63]

Intermediate	Structure	Chemical Name	Substrate
II-y-1		(5R,8S)-methyl 5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxylate	 II-d-6








[0695] IM II-y-1

¹H NMR (CDCl₃) delta 8.74 (1 H, d, J = 4.6 Hz), 7.87 (1 H, d, J = 7.9 Hz), 7.39 (1 H, m), 5.61 (1 H, br d, J = 49.4 Hz), 3.80 (3 H, s), 2.57-2.40 (4 H, m).

MS (ESI) m/z: 228.1 (M+H).

[0696] The following Examples were prepared by General Procedure A (Table 64).

[0697] [Table 64]

Examples	Structure	Chemical Name	Substrate	Amine
352		(5R,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide	 IM II-y-1	
353		(5R,8S)-N-(2,4-dichlorobenzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide		
354		(5R,8S)-N-(2,3-dichlorobenzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide		

[0698]


[Table 65]

Examples	LC MS			Examples	LC MS		
	Method	tR (min)	[M+H] ⁺		Method	tR (min)	[M+H] ⁺
352	A	1.64	405.1	354	A	1.61	371.1
353	A	1.64	371.1				

[0699] Procedure: Scheme 16, Step 1

Intermediate (IM) XVIII-1, 1-(2-chloropyridin-3-yl)pent-4-en-1-one

[0700] [Chem.37]



[0701] To a solution of 3-bromo-2-chloro pyridine (4.0 g, 20.8 mmol) in THF (100 mL) was added dropwise 1.3 M 2-propyl magnesium chloride lithium chloride complex in THF solution (18.5 ml, 25.0 mmol) at -10 °C under Ar atmosphere. The reaction mixture was stirred at the same temperature for 15 min. Then a THF solution of pent-4-enoyl chloride (3.2 g, 27.0 mmol) was added to the mixture at -40 °C. After addition, the mixture was stirred at the same temperature for 1.5 h. The mixture was poured into aq. NH₄Cl at 0 °C. The mixture was extracted with EtOAc twice and washed with aq. NaHCO₃ and brine. The combined extracts were dried over Na₂SO₄ and concentrated in vacuo. The insoluble material of the resulting residue was removed by filtration and washed with EtOAc. An activated charcoal was added to the filtrate. The mixture was stirred at 45 °C for 1.0 h and filtered through a pad of celite. The filtrate was distilled at reduced pressure to afford 2.9 g (72%) of the title compound.

¹H NMR (CDCl₃) delta 8.49 (1 H, dd, J = 4.9, 2.0 Hz), 7.81 (1 H, dd, J = 7.6, 2.0 Hz), 7.34 (1 H, dd, J = 7.6, 4.8 Hz), 5.77-5.91 (1 H, m), 4.99-5.13 (3 H, m), 3.11 (2 H, t, J = 7.3 Hz, 2 H), 2.43-2.54 (2 H, m).


MS (ESI) m/z: 196.3 (M+H)⁺.

[0702] Procedure: Scheme 16, Step 2

Intermediate (IM) XIX-1, 8-methylene-7,8-dihydroquinolin-5(6H)-one

[0703]

[Chem.38]



[0704] A mixture of 1-(2-chloropyridin-3-yl)pent-4-en-1-one (300 mg, 1.53 mmol, IM XVIII-1), triethylamine (0.64 mL, 4.61 mmol), Xantphos (17.7 mg, 0.031 mmol), and Pd(OAc)₂ (6.9 mg, 0.031 mmol) in MeCN (1.53 mL) was heated at reflux. After being refluxed for 15 h, the mixture was cooled to room temperature. The mixture was filtered with celite and the filtrate was concentrated in vacuo. To the residue was added EtOAc/n-hexane and the mixture was stirred for 15 min at room temperature. The mixture was filtered with celite and the filtrate was washed with water, dried over Na₂SO₄. The residue was distilled at reduced pressure to afford 238 mg (97%) of the title compound.

[0705] ¹H NMR (CDCl₃) delta 8.76 (1 H, dd, J = 4.6, 1.8 Hz), 8.28 (1 H, dd, J = 8.0, 1.8 Hz), 7.34 (2 H, dd, J = 8.0, 4.7 Hz), 6.38 (1 H, s), 5.47 (1 H, d, J = 1.3 Hz), 2.89 - 2.98 (2 H, m), 2.75 - 2.85 (2 H, m).


MS (ESI) m/z: 160.3 (M+H).

[0706] Procedure: Scheme 16, Step 3

Intermediate (IM) XX-1,

5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carbonitrile

[0707] [Chem.39]



[0708] To a solution of 8-methylene-7,8-dihydroquinolin-5(6H)-one (300 mg, 1.9 mmol, IM XIX-1) in dichloromethane (15 mL) was added trimethylsilyl cyanide (243 microL, 2.45 mmol) and NMO (132 mg, 1.13 mmol) at 25 °C under Ar atmosphere. The reaction mixture was stirred at the same temperature for 4 h. Then Deoxo-Fluor(trademark) (382 microL 2.07 mmol) was added to the mixture at 0 °C. The mixture was stirred at the same temperature for 2 h and the mixture was poured into aq. NaHCO₃ at 0 °C. The mixture was extracted with dichloromethane. The organic layer was dried over Na₂SO₄ and removed solvent in vacuo. The resulting residue was

purified by silica gel column chromatography (30% EtOAc/n-hexane) to afford 256 mg (72%) of the title compound.

[0709] ^1H NMR (CDCl_3) δ 8.70 (1 H, dt, $J = 4.6, 1.6$ Hz), 8.02 (1 H, dt, $J = 8.0, 1.4$ Hz), 7.35 (1 H, dd, $J = 8.0, 4.7$ Hz), 6.45 (1 H, s), 5.37 (1 H, s), 2.77-2.97 (3 H, m), 2.42-2.62 (2 H, m).


MS (ESI) m/z : 189.4 ($\text{M}+\text{H}$).

[0710] Procedure: Scheme 16, Step 4

Intermediate (IM) II-e-2-1, methyl

5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate

[0711] [Chem.40]



[0712] The title compound was prepared according to the procedure as described in


General Procedure: Scheme 2, Step 3 using

5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carbonitrile (IM XX-1).

[0713] Procedure: Scheme 17, Step 1

Intermediate (IM) XXII-1, methyl 2-(2-chloropyridin-3-yl)-2-hydroxyhex-5-enoate

[0714] [Chem.41]



[0715] To a solution of 3-bromo-2-chloropyridine (1.79 g, 9.30 mmol) in THF (10 mL) was added dropwise 1.3 M 2-propylmagnesium chloride lithium chloride complex in THF (7.2 mL, 9.4 mmol) at -15 °C and the mixture was stirred at the same temperature for 1 h. A solution of methyl 2-oxohex-5-enoate (1.33 g, 9.36 mmol) in THF (3 mL) was added to the mixture at -40 °C. The mixture was stirred at the same temperature for 2 h. Then the reaction was quenched with sat. aq. NH_4Cl . The mixture was extracted with EtOAc twice. The combined organic layer was washed with brine, dried over Na_2SO_4 , and removed solvent in vacuo. The residue was purified by silica gel column chromatography (25% EtOAc/n-hexane) to afford 1.78 g (74%) of the title compound.


[0716] ^1H NMR (CDCl_3) delta 8.35 (1 H, dd, $J = 4.7, 1.8$ Hz), 7.97 (1 H, dd, $J = 7.8, 1.8$ Hz), 7.26-7.33 (1 H, m), 5.83 (1 H, ddt, $J = 17.0, 10.4, 6.4, 6.4$ Hz), 4.98-5.10 (2 H, m), 3.84 (1 H, s), 3.78 (3 H, s), 2.14-2.40 (3 H, m), 1.96-2.07 (1 H, m).

[0717] Procedure: Scheme 17, Step 2

Intermediate (IM) II-e-2-5, methyl

5-hydroxy-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate

[0718] [Chem.42]



[0719] The title compound was prepared according to the procedure as described in

General Procedure: Scheme 16, Step 2 using methyl


2-(2-chloropyridin-3-yl)-2-hydroxyhex-5-enoate (IM XXII-1).

[0720] Procedure: Scheme 18, Step 1

Intermediate (IM) XXIII-1,

5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylic acid

[0721] [Chem.43]



[0722] A mixture of methyl

5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylate (3.0 g, 13.6 mmol, IM II-e-2-1) and 2 N aq. NaOH (14 mL, 28.0 mmol) in MeOH (67 mL) was stirred at room temperature for 1.5 h, and then the mixture was concentrated in vacuo. 10% aq. citric acid (90 mL) was added to the mixture. The mixture was extracted with EtOAc. The organic layer was dried over Na_2SO_4 and concentrated in vacuo to afford 1.57 g (56%) of the title compound.

^1H NMR (CDCl_3) delta 8.73 (1 H, d, $J = 4.9$ Hz), 8.28 (2 H, br s), 7.95 (1 H, d, $J = 7.6$ Hz), 7.35 (1 H, dd, $J = 7.9, 5.0$ Hz), 6.24 (1 H, s), 5.39 (1 H, s), 2.75-2.91 (2 H, m), 2.44-2.65 (1 H, m), 2.26-2.40 (1 H, m).


MS (ESI) m/z : 208.1 ($\text{M}+\text{H}$) $^+$.

[0723] Procedure: Scheme 18, Step 2

Intermediate (IM) XXIV,

5-fluoro-8-(iodomethyl)-5,6,7,8-tetrahydro-8,5-(epoxymethano)quinolin-10-one

[0724] [Chem.44]



[0725] To a solution of 5-fluoro-8-methylene-5,6,7,8-tetrahydroquinoline-5-carboxylic acid (2.0 g, 9.65 mmol, IM XXIII-1) in MeCN (100 mL) was added NaHCO₃ (1.7 g, 19.6 mmol) and iodine (5.0 g, 19.6 mmol) at room temperature. The reaction mixture was stirred at the same temperature for 3 h. Then sat. aq. Na₂S₂O₃ was added to the mixture. The mixture was extracted with EtOAc and washed with brine. The extract was dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by silica gel column chromatography (30% EtOAc/n-hexane) to afford 2.6 g (81%) of the title compound.

[0726] ¹H NMR (CDCl₃) delta 8.59 (1 H, dd, J = 5.1, 1.5 Hz), 7.88 (1 H, dd, J = 7.6, 1.5 Hz), 7.43 (1 H, dd, J = 7.7, 5.0 Hz), 4.03-4.12 (2 H, m), 2.71-2.81 (1 H, m), 2.51 (1 H, tdd, J = 11.3, 11.3, 5.5, 3.4 Hz), 2.09 (1 H, tdd, J = 11.7, 11.7, 4.6, 3.2 Hz), 1.91-2.00 (1 H, m).

MS (ESI) m/z: 333.9 (M+H)⁺.

[0727] Pharmacological assays

The ability of the 5,6,7,8-tetrahydroquinoline and 6,7-dihydro-5H-cyclopenta[b]pyridine derivatives of the formula (I) to inhibit the P2X7 channel was measured by Ca²⁺ influx assay and electrophysiology assay described below.

[0728] human P2X7 Functional Assay

The functional activity of compounds was determined by measuring changes in intracellular calcium concentration using a Ca²⁺-sensitive fluorescent dye, Fluo-4 (Molecular Probes). The changes in fluorescent signal were monitored by the cell imaging technology by Hamamatsu Photonics's Functional Drug Screening System (FDSS). Increases in intracellular Ca²⁺ concentration were readily detected upon activation with BzATP.

Cell Maintenance:

HEK293 cells stably expressing human P2X7 (GenBank accession number BC011913) carrying a C-terminal FLAG tag were grown in T225 flasks, in a 5% CO₂

humidified incubator to about 80% confluence. Media composition consisted of Dulbecco's Modified Eagle Medium (high glucose), 10% fetal bovine serum (BSA), 100 units/microM Penicillin, 100 microg/mL Streptomycin and 250 microg/mL Geneticine.

[0729] Protocol:

Day One:

1. Plate-out HEK293-human P2X7 cells (40 microL medium containing 10,000 cells per well) into poly-D-lysine coated 384-well plates (Corning) at 24 h prior to assay.
2. Incubate at 37 °C in 5% CO₂.

[0730] Day Two:

1. Wash each well with 80 microL of assay buffer (20 mM HEPES, 1 x HBSS, pH 7.4 adjusted with NaOH) three times and leave 20 microL using plate washer, ELx-405 Select CW (BIO-TEK).
2. Add 20 microL of assay buffer containing 2.5 mM probenecid, 0.5 microM Fluo-4-AM (Molecular Probes) and 0.1% Pluronic F-127 to each well.
3. Incubate the plate at 37 °C in 5% CO₂ for 1 h.
4. Wash each well with 80 microL of assay buffer (see below) three times and leave 20 microL using plate washer, ELx-405 Select CW (BIO-TEK).
5. Test compounds were prepared at 100X the test concentration in DMSO by serial dilution with Biomek-FX liquid handling instrument. 33X diluted compound solutions in assay buffer were prepared in intermediate compound plate with Biomek-NX liquid handling instrument. A further 3X dilution occurred in below steps 6 and 7.
6. Add 20 microL of 33X diluted compound solutions into each well and leave the plate for 10 min under the dark at room temperature.
7. Measure activity by FDSS as follows:
 - Set the assay plate on the stacker of FDSS.
 - Start the detection of fluorescence intensity at 540 nm by 480 nm excitation.
 - After 30 seconds, add 20 microL of assay buffer containing 240 microM BzATP (final concentration 80 microM).

IC₅₀ values for compounds of the present invention were determined from 7-point dose-response studies. Curves were generated using the average of duplicate wells for each data point. Finally, the IC₅₀ values are calculated with the best-fit dose curve determined by XLfit (ID Business Solutions Ltd.).

[0731] Antagonistic activities with respect to the human P2X7 receptor (IC₅₀ values) of exemplified compounds are displayed in Table 66.

[0732]

[Table 66-1]

A: <50 nM, B: 50- to 100 nM, C: 101- to 300 nM, D: 301- to 1,000 nM, E: 1,001- to 3,000 nM

Examples	IC ₅₀	Examples	IC ₅₀	Examples	IC ₅₀	Examples	IC ₅₀
1	A	33	A	65	D	97	A
2	A	34	A	66	A	98	C
3	D	35	A	67	C	99	A
4	A	36	A	68	A	100	A
5	E	37	A	69	D	101	C
6	A	38	C	70	A	102	A
7	A	39	A	71	D	103	C
8	A	40	A	72	A	104	A
9	A	41	A	73	C	105	D
10	A	42	A	74	C	106	A
11	A	43	A	75	E	107	C
12	A	44	A	76	B	108	D
13	A	45	A	77	D	109	D
14	A	46	A	78	A	110	D
15	D	47	A	79	D	111	D
16	A	48	A	80	B	112	C
17	B	49	A	81	A	113	A
18	A	50	A	82	A	114	C
19	A	51	A	83	C	115	A
20	A	52	A	84	A	116	D
21	A	53	A	85	A	117	A
22	A	54	A	86	C	118	A
23	A	55	A	87	B	119	A
24	A	56	A	88	A	120	E
25	C	57	A	89	C	121	B
26	A	58	A	90	C	122	B
27	A	59	A	91	C	123	C
28	B	60	C	92	C	124	C
29	A	61	A	93	E	125	C
30	A	62	C	94	E	126	A
31	B	63	C	95	E	127	A
32	A	64	A	96	E	128	A

[Table 66-2]

129	A	164	C	199	B	234	D
130	A	165	A	200	B	235	E
131	A	166	C	201	B	236	E
132	A	167	A	202	B	237	A
133	A	168	C	203	B	238	A
134	C	169	A	204	B	239	C
135	A	170	A	205	B	240	D
136	C	171	A	206	C	241	A
137	A	172	C	207	C	242	A
138	C	173	B	208	C	243	D
139	A	174	A	209	C	244	E
140	D	175	A	210	C	245	A
141	A	176	A	211	C	246	A
142	A	177	A	212	C	247	A
143	D	178	A	213	C	248	A
144	A	179	A	214	C	249	B
145	C	180	A	215	C	250	C
146	A	181	A	216	C	251	A
147	A	182	A	217	C	252	A
148	A	183	A	218	D	253	A
149	A	184	A	219	D	254	D
150	A	185	A	220	D	255	C
151	A	186	A	221	D	256	A
152	A	187	A	222	D	257	C
153	A	188	A	223	D	258	A
154	A	189	A	224	D	259	A
155	B	190	A	225	D	260	D
156	C	191	A	226	E	261	A
157	A	192	A	227	E	262	A
158	A	193	A	228	E	263	A
159	A	194	A	229	A	264	A
160	C	195	A	230	A	265	A
161	A	196	A	231	A	266	A
162	C	197	A	232	A	267	A
163	A	198	A	233	C	268	A

[Table 66-3]

269	A	291	C	313	B	335	A
270	A	292	B	314	C	336	A
271	D	293	E	315	E	337	A
272	A	294	B	316	A	338	A
273	A	295	A	317	A	339	A
274	A	296	B	318	D	340	A
275	E	297	D	319	A	341	B
276	D	298	A	320	A	342	B
277	A	299	C	321	A	343	A
278	E	300	A	322	C	344	A
279	A	301	B	323	C	345	A
280	A	302	C	324	B	346	A
281	D	303	E	325	E	347	B
282	A	304	A	326	A	348	C
283	A	305	A	327	A	349	A
284	A	306	A	328	C	350	A
285	A	307	D	329	B	351	A
286	A	308	C	330	D	352	A
287	A	309	E	331	B	353	B
288	A	310	A	332	E	354	B
289	A	311	B	333	C		
290	A	312	A	334	A		

[0733] rat P2X7 Functional Assay

The functional activity of compounds was determined by measuring changes in intracellular calcium concentration using a Ca²⁺-sensitive fluorescent dye, Fluo-4 (Molecular Probes). The changes in fluorescent signal were monitored by the cell imaging technology by Hamamatsu Photonics's Functional Drug Screening System (FDSS). Increases in intracellular Ca²⁺ concentration were readily detected upon activation with BzATP.

Cell Maintenance:

HEK293 cells stably expressing rat P2X7 (GenBank accession number NM_019256) were grown in Corning CellBIND cell culture flasks, in a 5% CO₂ humidified incubator to about 80% confluence. Media composition consisted of Dulbecco's Modified Eagle Medium (high glucose), 10% fetal bovine serum (BSA), 100 units/microL Penicillin, 100 microg/mL Streptomycin and 250 microg/mL Geneticine.

[0734] Protocol:

Day One:

1. Plate-out HEK293-human P2X7 cells (40 microL medium containing 10,000 cells per well) into poly-D-lysine coated 384-well plates (Corning) at 24 h prior to assay. Plate-out HEK293-rat P2X7 cells (40 microL medium containing 5,000 cells per well) into poly-D-lysine coated 384-well plates (BD Falcon) at 24 h prior to assay.
2. Incubate at 37 °C in 5% CO₂.

[0735] Day Two:

1. Wash each well with 80 microL of assay buffer (20 mM HEPES, 1 x HBSS, pH 7.4 adjusted with NaOH) three times and leave 20 microL using plate washer, ELx-405 Select CW (BIO-TEK).
2. Add 20 microL of assay buffer containing 2.5 mM probenecid, 0.5 microM Fluo-4-AM (Molecular Probes) and 0.1% Pluronic F-127 to each well.
3. Incubate the plate at 37 °C in 5% CO₂ for 1 h.
4. Wash each well with 80 microL of assay buffer (see below) three times and leave 20 microL using plate washer, ELx-405 Select CW (BIO-TEK).
5. Test compounds were prepared at 100X the test concentration in DMSO by serial dilution with Biomek-FX liquid handling instrument. 33X diluted compound solutions in assay buffer were prepared in intermediate compound plate with Biomek-NX liquid handling instrument. A further 3X dilution occurred in below steps 6 and 7.
6. Add 20 microL of 33X diluted compound solutions into each well and leave the plate for 10 min under the dark at room temperature.
7. Measure activity by FDSS as follows:
 - Set the assay plate on the stacker of FDSS.
 - Start the detection of fluorescence intensity at 540 nm by 480 nm excitation.
 - After 30 seconds, add 20 microL of assay buffer containing 30 microM BzATP (final 10 microM).

IC₅₀ values for compounds of the present invention were determined from 7-point dose-response studies. Curves were generated using the average of duplicate wells for each data point. Finally, the IC₅₀ values are calculated with the best-fit dose curve determined by XLfit (ID Business Solutions Ltd.).

[0736] Antagonistic activities with respect to the rat P2X7 receptor (IC₅₀ <1 uM) of exemplified compounds are displayed in Table 67.

[0737]

[Table 67-1]

Examples	Examples	Examples	Examples	Examples	Examples	Examples
1	40	81	137	184	250	288
2	41	82	139	185	251	289
3	42	83	141	187	252	290
6	43	84	142	189	253	291
7	44	85	144	190	254	292
8	45	87	146	192	256	294
9	46	88	148	193	257	295
10	47	89	149	195	258	296
11	48	97	150	196	259	297
12	49	100	152	199	260	298
13	50	102	153	201	261	299
14	51	104	154	203	262	300
16	52	106	157	205	263	301
18	53	107	158	206	264	302
19	54	108	159	208	265	303
20	55	110	161	221	266	304
21	56	111	163	229	267	305
22	57	115	165	230	268	306
23	58	117	167	231	269	307
24	59	118	168	232	270	308
25	60	121	169	233	272	309
26	61	122	170	234	273	310
27	62	126	171	237	274	311
28	64	127	172	238	277	312
29	66	128	174	239	279	313
32	67	129	175	240	280	314
33	68	130	176	241	281	315
34	70	131	177	242	282	316
35	72	132	178	245	283	317
36	74	133	179	246	284	318
37	76	134	180	247	285	319
38	78	135	182	248	286	320
39	80	136	183	249	287	321

[Table 67-2]

322	327	334	338	342	346	351
323	328	335	339	343	347	352
324	329	336	340	344	349	
326	333	337	341	345	350	

[0738] All tested compounds of the invention show higher IC₅₀ values in human dofetilide binding than IC₅₀ values in P2X7 functional assay described above.

[0739] Metabolic stability assay:

[0740] Human Liver Microsomal Clearance of Test Compounds.

The in vitro elimination half-life estimates ($t_{1/2}$) and in vitro intrinsic clearance values ($hCL_{int,u}$) were observed from metabolic stability in human liver microsomes.

[0741] Incubation with liver microsomes

Stock solutions of test compound were prepared at 10 mM (as active compound) in DMSO. The stock solution was diluted immediately before use to 50 microM using 50% acetonitrile-water mixture solution (v/v) to produce working solution. The NADPH-regenerating solution was prepared on the day of analysis by diluting 1 volume of 80 mM NADP⁺ (ORIENTAL YEAST) with 1 volume of 240 mM MgCl₂ (WAKO) and 1 volume of 320 mM of glucose-6-phosphate (Sigma-Aldrich) and 1 volume of 32 U/mL of glucose-6-phosphate dehydrogenase (Sigma-Aldrich) and 2 volume of 200 mM UDP-GA (Nacalai) and 2 volume of 6.6 mM beta-NAD (ORIENTAL YEAST), respectively. Immediately prior to use, the reaction mixture was produced mixing 1 volume of the NADPH-regenerating solution with 6.8 volume of 125 mM potassium phosphate assay buffer. Human liver microsomes (XenoTech, pooled, mixed-gender human microsomes) were diluted to 2.5 mg protein/mL using 125 mM potassium phosphate assay buffer. Two microliters of working solution of each test compound and 78 microL of reaction mixture were added to 96 well cluster tubes (Micronic) in duplicate.

[0742] The tubes were placed in an incubator at 37°C for 5 minutes before adding the human liver microsomes. A 20 microL of aliquot of the human liver microsome solution (2.5 mg protein/mL) was added to each original well to initiate metabolism. Incubation was performed at 37°C. At 15 minutes, the plate was removed from the incubator and a solution containing internal standard (200 microL, 1 microM reserpine, 50 nM buspirone and 1 microM tolbutamide in 100% acetonitrile) was added to each well. The plate was then spun in a centrifuge at 3500 rpm for 15 minutes at 4°C. A supernatant was transferred from each well to a 96-well shallow plate and then diluted with 4 volume of the mobile phase (A).

[0743] LC-MS/MS Analysis

Quantitative analysis of test compound in quenched reaction mixture was performed using the LC-MS/MS system, which consisted of an Agilent 1100 series gradient HPLC pump (Agilent Technologies), a CTC HTS PAL Autosampler (AMR), and a Sciex API 3200 triple quadrupole mass spectrometer (Sciex) equipped with a turbo ionspray interface. The Chromatographic separation was achieved using reverse phase HPLC with an Inert Sustain RP C18 50 × 2.1 mm column (GL Science) or Capcell Pak RP C18 50 × 2.1 mm column (Shiseido). The column temperature was 40°C, and the flow rate was 0.4 mL/min. The mobile phase consisted of 2 solvents: (A) 0.1% formic acid in water and (B) acetonitrile or 0.1% formic acid in acetonitrile. The compounds were eluted with a step gradient achieving 5% to 90% of B in 0.7 min, 90% of B in 1.3 min and then returned to initial conditions for equilibration (1.5 or 1.6 min). The mass spectrometer was operated in multiple-reaction-monitoring mode. Integration of test compound and internal standard peak was performed using Analyst Software (version 1.6). The area ratio of each test compound was calculated by comparing the peak area of the compound to the peak area of an internal standard.

[0744] Calculation of Human Liver Microsomal Intrinsic Clearance ($hCL_{int,u}$)

The mean peak area ratios were calculated by averaging the peak area ratios ($n=2$) of a compound and internal standard for each sample. Metabolic stability was determined by plotting the natural logarithm of the mean peak area ratio of unchanged test compound as a function of time. Percent remaining was calculated by determining the ratio of the mean peak area ratio at incubation time to the mean peak area ratio of the time-zero samples. The rate of loss of test compound was calculated using the equation $k = [\ln(C_0) - \ln(C)] / \text{incubation time}$, where C_0 was the initial mean peak area ratio of the test compound, C was the mean peak area ratio of test compound remaining after incubation ($C = C_0 \times \text{remaining ratio}$), and the incubation time was 15 min. The $t_{1/2}$ was estimated using the equation $t_{1/2} = 0.693 / k$. The $hCL_{int,u}$ was estimated using the equation $hCL_{int,u} = k / (\text{microsomal protein concentration}) \times (\text{microsomal protein per gram of liver}) \times (\text{liver mass per kilogram of body mass}) / (\text{human microsomal fu})$, where the microsomal protein concentration was 0.5 mg/mL, and the physical and physiological scaling factors were used such as the microsomal protein per gram of liver (48.8 mg) and, the liver mass per kilogram of body mass (25.7 g), and the human microsomal fu was determined experimentally from the human liver microsomal binding assay.

[0745] The compounds of this invention show preferable stability, which show the above-mentioned practical use.

[0746] Drug-drug interaction assay

[0747] Cytochrome P450 Inhibition of Test Compounds.

Incubation with recombinant CYP and chemiluminescent probes

CYP inhibition assays (CYP1A2, 2B6, 2C8, 2C9, 2C19, 2D6 and 3A4) were performed with recombinant CYP enzyme (BD Gentest) and Promega assay kits (P450-Glo Assays) in 384 well plate (Corning). Each product's catalog number is shown in Table 68.

Stock solutions of test compound were prepared at 10 mM (as active compound) in DMSO. NADPH-regenerating solution for each well was prepared on the day of analysis by diluting 0.4 microL of NADPH-A reagent (BD Gentest), 0.08 microL of NADPH-B reagent (BD Gentest) and 3.52 microL of water, for CYP 1A2, 2B6, 2C8, 2C9, 2C19 and 2D6. For CYP3A4, 1.6 microL of 1 M KPO₄, 0.4 microL of NADPH-A reagent and 0.08 microL of NADPH-B reagent, and 1.92 microL of water was mixed for each well. CYP enzyme mixture was prepared by following mixing ratio: 0.96 microL of water, 0.8 microL of 1 M KPO₄, 0.16 microL of Luciferin-ME, CYP1A2 Enzyme 0.08 microL/well for CYP1A2, 1.176 microL of water, 0.8 microL of 1 M KPO₄, 0.008 microL of Luciferin-2B6, CYP2B6 Enzyme 0.016 microL/well for CYP2B6, 1.04 microL of water, 0.4 microL of 1 M KPO₄, 0.24 microL of Luciferin-ME, CYP2C8 Enzyme 0.32 microL/well for CYP2C8, 1.56 microL of water, 0.2 microL of 1 M KPO₄, 0.16 microL of Luciferin-H, CYP2C9 Enzyme 0.08 microL/well for CYP2C9, 1.552 microL of water, 0.4 microL of 1 M KPO₄, 0.008 microL of Luciferin-H EGE, CYP2C19 Enzyme 0.04 microL/well for CYP2C19, 1.136 microL of water, 0.8 microL of 1 M KPO₄, 0.024 microL of Luciferin-ME EGE, CYP2D6 Enzyme 0.04 microL/well for CYP2D6, 1.916 microL of 100 mM Tris-HCl, 0.004 microL of Luciferin-PPXE, CYP3A4 Enzyme 0.08 microL/well for CYP3A4. Four microL of NADPH-regenerating solution was placed in 384 well plate, and then 2 microL of stock solution of test compounds and 2 microL of CYP enzyme mixture were added into each well. The plate was spun down and incubated at each condition as shown in Table 69. After incubation, 8 microL of Luciferin Detection Reagent for each CYP enzyme was added into each well and stirred by plate shaker (BioShake XP, WAKEN B TECH) at 1000 rpm for 1 min. The plate was incubated for 30 min at room temperature, protected from light. Luminescence was measured by luminometer (Ultra, Tecan and EnVision, PerkinElmer). Luminescence signals were used to determine percent inhibition at 10 microM of test compound. A separate control incubation for chemiluminescence contained test compound (10 microM) and control CYPs.

[0748]

[Table 68]

CYP isoforms	P450-GloAssay Kit Cat No.	Human CYP Enzymes (BD Gentest) Cat No.
CYP1A2	V8772	456203
CYP2B6	V8322	456255
CYP2C8	V8782	456252
CYP2C9	V8792	456258
CYP2C19	V8882	456259
CYP2D6	V8892	456217
CYP3A4	V8912	456202

[0749] [Table 69]

CYP isoforms	Enzyme (pmol)	Substrate concentration (μ M)	Incubation time(min) / Temperature
CYP1A2	0.08	100	45 / room temperature
CYP2B6	0.016	3	30 / room temperature
CYP2C8	0.08	150	90 / 37°C
CYP2C9	0.08	100	30 / 37°C
CYP2C19	0.04	10	30 / room temperature
CYP2D6	0.04	30	30 / room temperature
CYP3A4	0.08	25	30 / room temperature

[0750] The compounds of this invention show preferable results, which show the above-mentioned practical use.

[0751] To summarize the above metabolism assays, all the compounds of the present invention show unexpectedly preferable results in the HLM assay and/or in the drug-drug interaction assay comparing with the closest compounds. Therefore, all the compounds of the present invention have excellent pharmacokinetic properties.

[0752] hERG assay

The hERG (Human ether-a-go-go-related gene) channel inhibitory activity and the QT prolonging action of the compound of the present invention can be confirmed in the suitable methods known to skilled in the art. For example, hERG channel inhibitory activity of compounds of the present invention have been confirmed in electrophysiology assay (Chanchin, M. et al., Folia Pharmacol. Jpn., 2002, 119, 345-351) .

[0753] All tested compounds of the invention show higher IC₅₀ values in hERG assay than IC₅₀ values in P2X7 functional assay described above.

[0754] All publications, including but not limited to, issued patents, patent applications,


and journal articles, cited in this application are each herein incorporated by reference in their entirety. Although the invention has been described above with reference to the disclosed embodiments, those skilled in the art would readily appreciate that the specific experiments detailed are only illustrative of the invention. It should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the claims.

{CLAIMS}

{Claim 1}

A compound represented by the following formula (I):

{Chem. 1}



or a pharmaceutically acceptable salt thereof,

wherein:

X is N or N-oxide;

n is 0 or 1;

R¹ is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) hydroxyl, (4) -NH₂, (5) -NH-C₁-₆ alkyl and (6) -S(O)ₘ-C₁-₆ alkyl;

wherein m is independently 0, 1 or 2;

R² is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁-₆ alkyl and (4) -O-C₁-₆ alkyl; wherein the C₁-₆ alkyl or the -

O-C₁-₆ alkyl is unsubstituted or substituted with one or more substituents independently

selected from the group consisting of: halogen, hydroxyl, -O-C₁-₆ alkyl, -CN, -NR⁹ᵃR¹⁰ᵃ, -

(C=O)-R⁹ᵃ, -(C=O)-NR⁹ᵃR¹⁰ᵃ and -S(O)ₘ-R⁹ᵃ; wherein m is independently 0, 1 or 2;

R¹ may form =CH₂ or =O with R²; or

R¹ may form a 3 to 7 membered ring with R² which may contain one or more independently

selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and carbonyl

group; where the 3 to 7 membered ring is unsubstituted or substituted one or more with C₁-₆

alkyl;

R³ is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁-₆ alkyl and (4) -O-C₁-₆ alkyl;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R³ is the same or different;

R⁴ is selected from the group consisting of:

(1) hydrogen, (2) halogen and (3) hydroxyl;

R⁵ is hydrogen or C₁₋₆ alkyl;

R⁶ is selected from the group consisting of:

(1) hydrogen, (2) C₁₋₆ alkyl, (3) hydroxyC₁₋₆ alkyl, (4) C₁₋₆ alkoxy C₁₋₆ alkyl and (5) heterocyclyl C₁₋₆ alkyl;

R⁵ may form a saturated 3 to 7 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom, a sulfur atom or a double bond; or a saturated or unsaturated bicyclic 9 to 10 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom or a sulfur atom; wherein the saturated 3 to 7 membered ring or the saturated or unsaturated bicyclic 9 to 10 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) -O-aryl and (4) -O-C₁₋₆ alkylaryl;

R^{7a} and R^{7b} are independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) hydroxyl, (4) C₁₋₆ alkyl and (5) -NR^{9b}R^{10b};

R^{7a} may form a 3 to 7 membered ring with R⁵ which may contain a nitrogen atom or an oxygen atom; or

R^{7a} may form a 3 to 7 membered ring with R^{7b} which may contain a nitrogen atom or an oxygen atom;

q is 0 or 1;

R⁸ is selected from the group consisting of:

(1) hydrogen, (2) C₁₋₆ alkyl, (3) -O-C₁₋₆ alkyl, (4) C₂₋₆ alkenyl, (5) C₃₋₁₀ cycloalkyl, (6) -NR^{9b}R^{10b}; wherein the C₁₋₆ alkyl, the -O-C₁₋₆ alkyl, the C₂₋₆ alkenyl, the C₃₋₁₀ cycloalkyl or the -NR^{9b}R^{10b} is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen and hydroxyl; (7) heterocyclyl, (8) aryl, (9) -O-C₁₋₆ alkylaryl, (10) -O-aryl, (11) heteroaryl and (12) aryl-substituted heteroaryl; wherein the heterocyclyl, the aryl, the -O-C₁₋₆ alkylaryl, the -O-aryl, the heteroaryl or the aryl-substituted heteroaryl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -C₃₋₇ cycloalkyl, -O-C₃₋₇ cycloalkyl, hydroxyl-C₁₋₆ alkoxy, -CN, -NR^{9b}R^{10b}, -(C=O)-R^{9b}, -(C=O)-NR^{9b}R^{10b}, -NR^{9b}-(C=O)-R^{10b}, -NR¹¹-(C=O)-NR^{9b}R^{10b}, -NR^{9b}-(C=O)-OR^{10b}, -NR^{9b}-S(O)_m-R^{10b}, -NR¹¹-S(O)_m-NR^{9b}R^{10b}, -S(O)_m-R^{9b} and C₁₋₆ alkyl which may be substituted one or more with halogen, hydroxyl, -O-C₁₋₆ alkyl or NR^{9b}R^{10b}; wherein m is independently 0, 1 or 2;

R^{9a}, R^{9b}, R^{10a}, R^{10b} or R¹¹ is independently selected from the group consisting of:

(1) hydrogen, (2) hydroxyl, (3) C₁₋₆ alkyl and (4) hydroxyC₁₋₆ alkyl;

R^{9a} may form a 4 to 7 membered ring with R^{10a} which may contain one or more independently selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and double bond, wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

R^{9b} may form a 4 to 7 membered ring with R^{10b} which may contain one or more independently

selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and double bond, wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) C₁₋₆ alkyl.

{Claim 2}

The compound according to claim 1:

or a pharmaceutically acceptable salt thereof,

wherein:

X is N;

R⁵ is hydrogen or C₁₋₆ alkyl;

R⁶ is selected from the group consisting of:

(1) hydrogen and (2) C₁₋₆ alkyl;

R⁵ may form a saturated or unsaturated bicyclic 9 to 10 membered ring with R⁶ which may contain a nitrogen atom, an oxygen atom or a sulfur atom; wherein the saturated or unsaturated bicyclic 9 to 10 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) -O-aryl;

R^{7a} and R^{7b} are independently selected from the group consisting of:

(1) hydrogen, (4) C₁₋₆ alkyl and (5) -NR^{9b}R^{10b};

R^{7a} may form a 3 to 7 membered ring with R⁵ which may contain a nitrogen atom or an oxygen atom; or

R^{7a} may form a 3 to 7 membered ring with R^{7b} which may contain a nitrogen atom or an oxygen atom;


R⁸ is selected from the group consisting of:

(1) hydrogen, (2) C₁₋₆ alkyl, (5) C₃₋₁₀ cycloalkyl, wherein the C₁₋₆ alkyl or the C₃₋₁₀ cycloalkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen and hydroxyl; (7) heterocyclyl, (8) aryl, (9) -O-C₁₋₆ alkylaryl, (10) -O-aryl, (11) heteroaryl and (12) aryl-substituted heteroaryl, wherein the heterocyclyl, the aryl, the -O-C₁₋₆ alkylaryl, the -O-aryl, the heteroaryl or the aryl-substituted heteroaryl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -CN and C₁₋₆ alkyl which may be substituted one or more with halogen, hydroxyl, -O-C₁₋₆ alkyl or NR^{9b}R^{10b}.

{Claim 3}

A compound represented by the following formula (M):

{Chem. 2}



or a pharmaceutically acceptable salt thereof,

wherein:

n is 0 or 1;

R^1 is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) hydroxyl, (4) $-NH_2$, (5) $-NH-C_{1-6}$ alkyl and (6) $-S(O)_m-C_{1-6}$ alkyl; wherein m is independently 0, 1 or 2;

R^2 is selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C_{1-6} alkyl and (4) $-O-C_{1-6}$ alkyl; wherein the C_{1-6} alkyl or the $-O-C_{1-6}$ alkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, $-O-C_{1-6}$ alkyl, $-CN$, $-NR^{9a}R^{10a}$, $-(C=O)-R^{9a}$, $-(C=O)-NR^{9a}R^{10a}$ and $-S(O)_m-R^{9a}$; wherein m is independently 0, 1 or 2;

R^1 may form $=CH_2$ or $=O$ with R^2 ; or

R^1 may form a 3 to 7 membered ring with R^2 which may contain one or more independently selected from the group consisting of: nitrogen atom, oxygen atom, sulfur atom and carbonyl group; where the 3 to 7 membered ring is unsubstituted or substituted one or more with C_{1-6} alkyl;

R^3 is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C_{1-6} alkyl and (4) $-O-C_{1-6}$ alkyl;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R^3 is the same or different;

R^4 is selected from the group consisting of:

(1) hydrogen, (2) halogen and (3) hydroxyl;

R^5 is hydrogen or C_{1-6} alkyl;

R^6 is selected from the group consisting of:

(1) hydrogen, (2) C_{1-6} alkyl, (3) hydroxy C_{1-6} alkyl, (4) C_{1-6} alkoxy C_{1-6} alkyl and (5) heterocyclyl C_{1-6} alkyl;

R^5 may form a saturated 3 to 7 membered ring with R^6 which may contain a nitrogen atom, an oxygen atom, a sulfur atom or a double bond; wherein the saturated 3 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group

consisting of: (1) hydroxyl, (2) halogen, (3) -O-aryl and (4) -O-C₁₋₆ alkylaryl;

R^{9a}, R^{9b}, R^{10a} or R^{10b} is independently selected from the group consisting of:

(1) hydrogen, (2) hydroxyl, (3) C₁₋₆ alkyl and (4) hydroxyC₁₋₆ alkyl;

R^{9a} may form a 4 to 7 membered ring with R^{10a} which may contain a nitrogen atom or an oxygen atom; wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

R^{9b} may form a 4 to 7 membered ring with R^{10b} which may contain a nitrogen atom or an oxygen atom; wherein the 4 to 7 membered ring is optionally substituted with 1 to 6 substituents independently selected from the group consisting of: (1) hydroxyl, (2) halogen and (3) C₁₋₆ alkyl;

R¹², R¹³, R¹⁴ and R¹⁵ are independently selected from the group consisting of:

(1) hydrogen, (2) hydroxyl, (3) halogen, (4) C₁₋₆ alkyl, (5) -O-C₁₋₆ alkyl and (6) CN; wherein the C₁₋₆ alkyl or the -O-C₁₋₆ alkyl is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl and NR^{9b}R^{10b}; or

R¹² may form a 5 to 7 membered ring with R⁵ which may contain one or more independently selected from the group consisting of: nitrogen atom and oxygen atom.

{Claim 4}

The compound according to claim 3:

or a pharmaceutically acceptable salt thereof,

wherein:

n is 1;

R¹ is hydrogen or hydroxyl;

R² is methyl which is unsubstituted or substituted with one or more substituents independently selected from the group consisting of: halogen, hydroxyl, -O-C₁₋₆ alkyl, -CN and -NR^{9a}R^{10a};

p is 0;

R⁴ is hydrogen or fluoro;

R⁵ and R⁶ are independently selected from the group consisting of: (1) hydrogen and (2) C₁₋₆ alkyl;

R¹², R¹³ and R¹⁴ are independently selected from the group consisting of: (1) hydrogen, (3) halogen, and (4) C₁₋₃ alkyl which may be substituted one or more with hydroxyl;

R¹⁵ is hydrogen.

{Claim 5}

A compound which is selected from the group following or a pharmaceutically acceptable salt thereof,

N-(2,4-dichloro-6-methylbenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

N-(2,3-dichlorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(cycloheptylmethyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,3-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-(methoxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-methylbenzyl)-7-methylene-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-methylbenzyl)-5-hydroxy-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-methylbenzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
N-(2-chloro-3-(trifluoromethyl)benzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-(methoxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-methylbenzyl)-5,8-dihydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2-chloro-3-(trifluoromethyl)benzyl)-5,8-dihydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

N-(2-chloro-3-(trifluoromethyl)benzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 N-(2,3-dichlorobenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 2-(5-((2,4-dichloro-6-methylbenzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;
 2-(5-((2-chloro-3-(trifluoromethyl)benzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;
 (2-amino-2-oxoethyl)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-7-(hydroxymethyl)-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8S*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8S*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 8-(aminomethyl)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 8-(aminomethyl)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-5,6,7,8-

tetrahydroquinoline-5-carboxamide;
 8-(aminomethyl)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-((methylamino)methyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-
 yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxyazetidin-1-
 yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxy-3-methylazetidin-1-
 yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxy-3-methylazetidin-1-
 yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-
 hydroxyethyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-
 hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-8-amino-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S*,8S*)-8-amino-N-(2,4-dichlorobenzyl)-5-fluoro-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (3R*,5S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-
 quinoline]-5'-carboxamide;
 (3S*,5S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-
 quinoline]-5'-carboxamide;
 (5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-
 carboxamide;
 (5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-
 carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,7S)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,7R)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 (5S,7R)-N-(2,4-dichloro-6-methylbenzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 (5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-

carboxamide;

(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-fluoro-3-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1-(2,3,4-trichlorophenyl)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-

carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(3-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(5-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-bromo-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-6-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((R)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorophenethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3-chloro-5-(trifluoromethyl)pyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(3,4,5-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-cyano-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide

(5S,8S)-N-(2-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-4-(trifluoromethoxy)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-5-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(4-methoxy-2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((4-(4-chlorophenyl)thiazol-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-(morpholinomethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1S,2R)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(6-chloro-2-fluoro-3-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-(3-(trifluoromethyl)phenoxy)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-((1-(4-fluorophenyl)cyclopropyl)methyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1R,2S)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((1S,2S)-2-(benzyloxy)cyclopentyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,3-dimethylbutyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-phenoxyethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4,6-dichloro-2,3-dihydrobenzofuran-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(5,7-dichlorochroman-4-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(1-(adamantan-1-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-(4-chlorophenyl)-2-(4,4-difluoropiperidin-1-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(chroman-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-(4-chlorophenyl)propyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-morpholino-2-phenylethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-(4,4-difluoropiperidin-1-yl)-2-(4-methylthiazol-5-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-3,5-difluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-3,5-difluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-((2,4-dichlorobenzyl)carbamoyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline 1-oxide;

(R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-5-fluoro-8-oxo-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-8-methyl-N-((1-morpholinocyclohexyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3,8-dimethyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-

hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-
 hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-8-(cyanomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8R)-8-(cyanomethyl)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)thio)methyl)-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-4-methyl-6',7'-dihydro-5'H-spiro[morpholine-2,8'-
 quinoline]-5'-carboxamide;
 (5S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-
 quinoline]-5'-carboxamide;
 (5R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-
 quinoline]-5'-carboxamide;
 (2S,5'R)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-
 quinoline]-5'-carboxamide;
 (2S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-
 quinoline]-5'-carboxamide;
 (2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-
 quinoline]-5'-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfinyl)methyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylthio)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-

5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,3-dichloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-3-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorophenethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((4-(2,4-dichlorophenyl)tetrahydro-2H-pyran-4-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-methoxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-methoxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-(2-hydroxyethoxy)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-8-(2,3-dihydroxypropoxy)-5-fluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8rac)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(methylsulfonyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8rac)-N-(2,4-dichlorobenzyl)-5-fluoro-8-((2-hydroxyethyl)sulfonyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide and

(5R,8S)-N-(2,3-dichlorobenzyl)-5,8-difluoro-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 6}

The compound according to claim 5, which is selected from the group following or a pharmaceutically acceptable salt thereof,

N-(2,3-dichlorobenzyl)-8-oxo-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-7-oxo-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 2-(5-((2,4-dichloro-6-methylbenzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;
 2-(5-((2-chloro-3-(trifluoromethyl)benzyl)carbamoyl)-5,6,7,8-tetrahydroquinolin-8-yl)acetic acid;
 N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichlorobenzyl)-5-fluoro-7-hydroxy-7-(hydroxymethyl)-6,7-dihydro-5H-cyclopenta[b]pyridine-5-carboxamide;
 (5S*,8S*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(methoxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 N-(2,4-dichloro-6-methylbenzyl)-8-((dimethylamino)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxyazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxy-3-methylazetidin-1-

yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((3-methoxy-3-methylazetidin-1-yl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S*,8R*)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (3R*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide;
 (3S*,5'S*)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-3,8'-quinoline]-5'-carboxamide;
 (5S,8S)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichlorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(2,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8S)-N-(4-bromo-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-3-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-

carboxamide;

(5R,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-

carboxamide;

(5S,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-3-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(4-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,6-dichloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-(difluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-bromo-2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-6-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-5-fluoro-8-hydroxy-N-(2,3,6-trichlorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-methylbenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(3-fluoro-2-(trifluoromethyl)benzyl)-8-hydroxy-5,6,7,8-

tetrahydroquinoline-5-carboxamide;
 (5S,8S)-5-fluoro-8-hydroxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-bromo-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-(trifluoromethyl)benzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-5-fluoro-8-hydroxy-N-((R)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((3-chloro-5-(trifluoromethyl)pyridin-2-yl)methyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((R)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-5-fluoro-8-hydroxy-N-(2-(trifluoromethyl)benzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-5-fluoro-8-hydroxy-N-(3,4,5-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-cyano-2-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3,4-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-5-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluoro-4-(trifluoromethoxy)benzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,3-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,5-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1S,2R)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,6-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((S)-1,2,3,4-tetrahydronaphthalen-1-yl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-(3-(trifluoromethyl)phenoxy)ethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,5-difluorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-((1R,2S)-2-phenylcyclopropyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-N-(2-fluorobenzyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-3-methoxybenzyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-2,3-dihydro-1H-inden-1-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(3,3-dimethylbutyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-phenoxyethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4,6-dichloro-2,3-dihydrobenzofuran-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(chroman-3-yl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-fluoro-8-hydroxy-N-(2-morpholino-2-phenylethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-(4,4-difluoropiperidin-1-yl)-2-(4-methylthiazol-5-yl)ethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-tetrahydroquinoline-5-

carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-3-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-5-((2,4-dichlorobenzyl)carbonyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline 1-oxide;

(5S,8S)-5-fluoro-8-hydroxy-8-methyl-N-(2,3,4-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-

carboxamide;

(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((S)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-

hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-

hydroxyethyl)(methyl)amino)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-((3-hydroxyazetidin-1-yl)methyl)-

5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-8-(cyanomethyl)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-

tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-8-(cyanomethyl)-5-fluoro-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-(fluoromethyl)-8-hydroxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-4-methyl-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(5S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-quinoline]-5'-carboxamide;

(5R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-2-oxo-6',7'-dihydro-5'H-spiro[oxazolidine-5,8'-quinoline]-5'-carboxamide;

(2S,5'R)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(2S,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfinyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-((methylsulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(((2-hydroxyethyl)sulfonyl)methyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5R,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8R)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-

tetrahydroquinoline-5-carboxamide;
 (5S,8S)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8R)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-N-(2,3,4-trifluorobenzyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4-fluoro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-
 (hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2,4-dichloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2,3-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(3-chloro-2,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-(4-chloro-2,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-
 tetrahydroquinoline-5-carboxamide;
 (5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;
 (5S,8R)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-
 5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-((trans)-2-(2,4-dichlorophenyl)cyclopropyl)-5-fluoro-8-hydroxy-8-

(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide; (5S,8S)-5-fluoro-8-methoxy-N-(2,4,6-trifluorobenzyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide;

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-methoxy-5,6,7,8-tetrahydroquinoline-5-carboxamide;


(5S,8S)-N-(4-chloro-2-fluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide and

(5S,8S)-N-(2,4-difluorobenzyl)-5-fluoro-8-(2-hydroxyethoxy)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 7}

A compound represented by the following formula (L-a) or a pharmaceutically acceptable salt thereof:

{Chem. 3}



wherein:

R³ is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R³ is the same or different;

R¹⁶ is selected from the group consisting of:

(1) CN and (2) -CO₂R¹⁸;

R¹⁷ is selected from the group consisting of:

(1) fluoro and (2) hydroxyl; or,

R¹⁶ may form =O with R¹⁷;


R¹⁸ is selected from the group consisting of:

(1) hydrogen and (2) C₁₋₆ alkyl.

{Claim 8}

A compound represented by the following formula (L-b) or a pharmaceutically acceptable salt thereof:

{Chem. 4}



wherein:

R³ is independently selected from the group consisting of:

(1) hydrogen, (2) halogen, (3) C₁₋₆ alkyl and (4) -O-C₁₋₆ alkyl;

p is 0, 1, 2 or 3;

when p is 2 or 3, each R³ is the same or different;

R⁴ is selected from the group consisting of:

(1) hydrogen, (2) halogen and (3) hydroxyl.

{Claim 9}

A use of a compound described in any one of claims 1 to 6 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of a condition or disorder mediated by P2X7 receptor antagonistic activity, wherein the condition or disorder is selected from the group consisting of: diseases of the autoimmune and inflammatory system; diseases of the nervous and neuro-immune system; diseases involved with, and without, neuroinflammation of the Central Nervous System (CNS); diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems; skeletal disorders, diseases involving the secretory function of exocrine glands and glaucoma, Glomerulonephritis, Chaga's Disease, chlamydia, neuroblastoma, Tuberculosis, Polycystic Kidney Disease, cancer, and acne; and combinations thereof.

{Claim 10}

A compound for use described in any one of claims 1 to 6 or a pharmaceutically acceptable salt thereof for use in the treatment of a condition or disorder mediated by P2X7 receptor antagonistic activity.

{Claim 11}

The compound as described in claim 10, wherein the condition or disorder is selected from the group consisting of: diseases of the autoimmune and inflammatory system; diseases of the nervous and neuro-immune system; diseases involved with, and without, neuroinflammation of the Central Nervous System (CNS); diseases of the cardiovascular, metabolic, gastrointestinal and urogenital systems; skeletal disorders, diseases involving the secretory function of exocrine glands and glaucoma, Glomerulonephritis, Chaga's Disease, chlamydia, neuroblastoma, Tuberculosis, Polycystic Kidney Disease, cancer, and acne; and combinations thereof.

{Claim 12}

A pharmaceutical composition comprising a compound or a pharmaceutically acceptable

salt thereof, as described in any one of claims 1 to 6, and a pharmaceutically acceptable carrier.

{Claim 13}

The pharmaceutical composition as described in claim 12, further comprising another pharmacologically active agent.

{Claim 14}

A process for preparing a pharmaceutical composition, wherein the process comprises mixing a compound described in any one of claims 1 to 6 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier or excipient.

{Claim 15}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-(2,4-dichloro-6-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide or a pharmaceutically acceptable salt thereof.

{Claim 16}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-(2-chloro-4,6-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 17}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-(2,4-dichloro-6-(hydroxymethyl)benzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 18}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-((R)-1-(2-chloro-4-fluorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 19}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 20}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-((R)-1-(2,4-dichlorophenyl)ethyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 21}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-(2-chloro-4-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 22}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-(2-chloro-3,4-difluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 23}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-(2,4-dichloro-3-fluorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 24}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-(2,4-dichlorobenzyl)-5-fluoro-8-hydroxy-8-(hydroxymethyl)-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 25}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (5S,8S)-N-((3,5-dichloropyridin-2-yl)methyl)-5-fluoro-8-hydroxy-8-methyl-5,6,7,8-tetrahydroquinoline-5-carboxamide.

{Claim 26}

The compound according to claim 6, or a pharmaceutically acceptable salt thereof, which is (2R,5'S)-N-(2,4-dichlorobenzyl)-5'-fluoro-5-oxo-6',7'-dihydro-5'H-spiro[morpholine-2,8'-quinoline]-5'-carboxamide.

{Claim 27}

A compound described in any one of claims 15 to 26 or a pharmaceutically acceptable salt thereof for use in the treatment of a condition or disorder mediated by P2X7 receptor antagonistic activity.

{Claim 28}

The compound for use as described in claim 27, wherein the condition or disorder is neuropathic pain.