

## BINDING MOLECULES AGAINST CHIKUNGUNYA VIRUS AND USES THEREOF

### FIELD OF THE INVENTION

5           The present invention relates to binding molecules against Chikungunya virus. Particularly, said binding molecules are capable of neutralizing Chikungunya virus infectivity. Said binding molecules can be used with therapeutic, diagnosis or research purposes, among others. The invention also relates to a pharmaceutical composition comprising said binding molecules and more particularly, to the use  
10           thereof in the treatment or prevention of Chikungunya fever.

### BACKGROUND OF THE INVENTION

          Chikungunya fever is an emerging, epidemic disease, caused by an arbovirus and transmitted by the *Aedes* mosquitoes, of much significance for  
15           WHO's South-East Asia Region. The disease has been reported from countries of South and East Africa, South Asia and South-East Asia. In WHO's South-East Asia Region, outbreaks have been reported from India, Indonesia, Myanmar, Sri Lanka, Thailand and Maldives. Massive outbreaks of Chikungunya fever have occurred in recent years in India and in the island countries of the Indian Ocean. Similarly,  
20           Maldives reported outbreaks of Chikungunya fever for the first time in December 2006. Although not a killer disease, high morbidity rates and prolonged polyarthrititis leading to considerable disability in a proportion of the affected population can cause substantial socio-economic impact in affected countries (WHO, guidelines for prevention&control of Chikungunya fever, 2009).

25           The Chikungunya virus (CHIKV) is a member of the genus *alphavirus* and family *Togaviridae* (reviewed by Strauss and Strauss, 1994, *Microbiol Rev* 58, 491-562). The alphaviruses are small enveloped single-stranded positive RNA viruses exhibiting a large cell tropism. The viral surfaces are covered in membrane-anchored spikes composed of triplets of heterodimers of the envelope E1 and E2  
30           glycoproteins. The viral spike proteins facilitate attachment to cell surfaces and viral entry. The E1 envelope glycoprotein is a class II fusion protein that mediates

low pH-triggered membrane fusion during virus infection. E2 is a 50 kDa type I transmembrane glycoprotein: the first 260 amino acids constitute the ectodomain, followed by about 100 amino acids that form the stem region, a spanning region of 30 amino acids, and a short cytoplasmic endodomain of 30 amino acids (Plenetv, et al., 2001, Cell 105, 127-136; Mukhopadhyay, et al., 2006, Structure 14, 63-73). pE2 (the 62-kDa precursor to the E3 and E2 proteins) and E1 are assembled as heterodimers in the endoplasmic reticulum (Strauss and Strauss, 1994, Microbiol Rev 58, 491-562). After the cleavage of pE2 in the Golgi apparatus to form E3 and E2, the E1-E2 complexes are transported to the plasma membrane (PM). The interaction of the cytoplasmic E2 endodomain with the preassembled nucleocapsid is one of the initial steps in the process of virus envelopment at the PM. Integrity of virion is maintained by direct interactions between E1 and E2 (Strauss and Strauss, 1994, Microbiol Rev 58, 491-562). During the course of *alphavirus* life cycle, the E2 glycoprotein is responsible for receptor binding. In general, neutralizing antibodies against alphaviruses recognize epitopes in E2 rather than E1 (Roehrig, J. T. 1986. The use of monoclonal antibodies in studies of the structural proteins of togaviruses and flaviviruses, p.251-278. In S. Schlesinger and M. J. Schlesinger (ed.), The Togaviridae and Flaviviridae. Plenum Publishing Corp., New York).

Biological diagnosis of CHIK virus infection is essentially based on quantitative real-time RT-PCR-based method during the initial viraemic phase (Edwards et al., 2007, J.Clin.Virol. 39, 271-275; Laurent et al., 2007, Clin. Chem. 53, 1408-1414; Parida et al., 2007, J. Clin. Microbiol. 45, 351-357). Serological methods detect anti-CHIK IgM early times after the first clinical manifestations and specific IgG after two weeks (Pialoux et al., 2007, Lancet Infect. Dis. 7, 319-327). However, ELISA and immunodetection assays are poorly specific and sensitive due the cross reactivity of Chikungunya virus with related members of the Semliki Forest (SF) antigenic complex (Greiser-Wilke et al., 1991, J. Clin. Microbiol. 29, 131-137).

More recently, Brehin and colleagues (Brehin et al., 2008, Virology 371, 185-195) have developed some monoclonal antibodies (mAbs) reactive to CHIKV E2 glycoprotein for diagnosis and research purposes, which were also described in

WO 2009/031045. The three anti-CHIKV-E2 mAbs showed cross-reactivity with the O'nyong-nyong viral strains Igbo-Ora and ONN-59. CHIKV, Igbo-Ora and ONN-59 are serologically classified in the SF antigenic complex (Strauss and Strauss, 1994, Microbiol Rev 58, 491-562). These monoclonal antibodies are of murine origin, and  
 5 although they have demonstrated significant reactivity with CHIKV-associated E2 glycoprotein they failed to neutralize CHIKV infection of primate cells *in vitro*. Besides, a first commercial Chikungunya virus indirect immunofluorescence test (IIFT) is commercialized by Euroimmun AG, Germany for analyzing the CHIKV specific immune response. This test was evaluated by Litzba et al. (Litzba et al.,  
 10 2008, Journal of virological methods, 149(1), 175-179).

In the recent years, there has been an explosive re-emergence of Chikungunya fever. Thus, although some advances have been undertaken to provide a reliable test allowing the detection and monitoring of CHIKV specific antibodies, there is still a need to develop new anti-CHIKV monoclonal antibodies  
 15 for diagnosis and research purposes on Chikungunya virus infection.

On the other hand, a specific treatment is not available and there is no approved vaccine for the prevention of Chikungunya fever. Currently, vector control is the only way to prevent and control the outbreaks. Vector control is not an easy task and insecticide spraying is not always effective and desirable (WHO,  
 20 guidelines for prevention&control of Chikungunya fever, 2009).

Symptomatic treatment is recommended after excluding more serious conditions. Symptomatic or supportive treatment basically comprises rest and use of acetaminophen or paracetamol to relieve fever and ibuprofen, naproxen or other non-steroidal anti-inflammatory agent (NSAID) to relieve the arthritic  
 25 component. Patients with persistent or chronic phase of arthritis who fail to respond to NSAID may show some response to chloroquine phosphate. The latter may act as a weak broad spectrum antiviral agent apart from being an anti-inflammatory agent. Use of corticosteroids in managing Chikungunya related arthropathy has in general been a contentious issue and has to be the last resort in  
 30 a clinical decision (WHO, guidelines for prevention&control of Chikungunya fever, 2009).

While there has been extensive work in vaccinology for several other alphaviruses (Rayner et al., 2002, Rev Med Virol 12, 279–296; Nalca et al., 2003, Antiviral Res 60, 153–174); Johnston & Davis, 2004, Arch Virol Suppl 18, 207–220), the history of vaccine development for CHIKV is short and none of these efforts have  
 5 yet resulted in a licensed vaccine. Recently, a Phase II study was performed with a serially passaged live chikungunya virus (Edelman et al., 2000, Am J Trop Med Hyg. 62(6), 681-5) with good immunogenicity and tolerance results. However, this vaccine seems not to have reached market authorization and further efforts are being made to find a Chikungunya vaccine, see for example the *in vivo* study on  
 10 the immunogenicity of consensus-based DNA vaccines against CHIKV performed by Muthumani et al. (Muthumani K. et al., 2008, Vaccine 26(40), 5128-34).

With regards to the immunotherapy strategies against Chikungunya virus infection, the use of a concentrate of human immunoglobulins (IgA, IgM and IgG) has been previously described, as well as F(ab)'2 and/or Fab fragments specific to  
 15 an arbovirus (i.e. Chikungunya virus) for use as a medicament in the treatment of arbovirosis, see WO 2007/118986.

Furthermore, studies with neutralizing mAbs have been reported for several alphaviruses including Sindbis virus (SIN), Venezuelan equine encephalitis virus (VEE), Ross River virus (RR), Semliki Forest virus (SF), Eastern equine encephalitis  
 20 virus (EEE) and Western equine encephalitis virus (WEE) (see, e.g. Roehrig, J. T. 1986. The use of monoclonal antibodies in studies of the structural proteins of togaviruses and flaviviruses, p.251-278. In S. Schlesinger and M. J. Schlesinger (ed.), The Togaviridae and Flaviviridae. Plenum Publishing Corp., New York). Recently, human polyvalent immunoglobulins were purified from plasma samples obtained  
 25 from donors in the convalescent phase of CHIKV infection, and the preventive and curative effects of these immunoglobulins were investigated (Couderc et al., 2009, J Infect Dis. 2009, 200(4), 489-91). However, to our knowledge, at present no neutralizing monoclonal antibodies against Chikungunya virus have been described, let alone fully human neutralizing antibodies.

30 Since at present there is no vaccine or specific treatment available on the market to combat the Chikungunya fever, several efforts are being undertaken to

obtain a safe and active therapy to be administered to patients suffering from Chikungunya fever and also to obtain a protective therapy against the virus infection. Accordingly, there is a need for providing therapies that are useful in the prevention and/or treatment of the Chikungunya fever.

5

#### **SUMMARY OF THE INVENTION**

The invention provides new binding molecules against Chikungunya virus (CHIKV). In particular, two monoclonal antibodies specifically binding to an epitope located in an antigenic site of one of the Chikungunya virus envelope proteins are provided. Furthermore, these antibodies are fully human antibodies with CHIKV neutralizing properties which make them particularly useful for the treatment or prevention of Chikungunya fever, thus avoiding all the secondary effects or unwanted reactions associated to murine, chimeric or humanized antibodies in the treatment of human patients.

15 In accordance with a first aspect of this invention, we provide an isolated binding protein that specifically binds to Chikungunya virus, comprising a heavy chain amino acid sequence comprising at least one of the CDR's selected from the group consisting of: (a) CDRH1's of SEQ ID NOs: 10 or 30, (b) CDRH2's of SEQ ID NOs: 12 or 32, and (c) CDRH3's of SEQ ID NOs: 14 or 34, and/or a light chain amino acid sequence comprising at least one of the CDR's selected from the group consisting of: (d) CDRL1's of SEQ ID NOs: 16 or 36, (e) CDRL2's of SEQ ID NOs: 18 or 38, and (f) CDRL3's of SEQ ID NOs: 20 or 40.

20 A second aspect of the present invention relates to a functional variant of a binding protein of the invention characterized in that said functional variant binds to Chikungunya virus.

25 A third aspect of the present invention refers to an immunoconjugate of a binding protein or a functional variant of the invention wherein said binding protein or functional variant is coupled to at least one labeling and/or effector group.

A further aspect of the present invention relates to an isolated nucleic acid molecule comprising a nucleic acid sequence encoding a binding protein, a functional variant or an immunoconjugate of the invention.

Another aspect of the present invention pertains to a vector comprising a nucleic acid molecule of the invention. A further aspect of the invention relates to a host cell comprising the nucleic acid molecule or the vector of the invention. In accordance to an additional aspect of this invention, we provide a method for producing a binding protein, a functional variant or an immunoconjugate of the invention, comprising the step of producing said binding protein, functional variant or immunoconjugate in a host cell of the invention and optionally, isolating said binding protein, functional variant or immunoconjugate.

In yet another aspect of the present invention, we provide a pharmaceutical composition comprising as an active agent at least one isolated binding protein, at least one functional variant or at least one immunoconjugate of the invention; and a pharmaceutically acceptable carrier, diluent or adjuvant.

A related aspect of the present invention refers to an isolated binding protein, a functional variant or an immunoconjugate of the invention for use as a medicament. Moreover, the present invention relates in a further aspect to a isolated binding protein, a functional variant, an immunoconjugate and/or the pharmaceutical composition of the invention for use in the prevention or treatment of the infection of an arbovirus from the Togaviridae family, preferably from the genus *alphavirus* more preferably the Chikungunya virus.

In even another aspect, the present invention relates to the use of a binding protein, a functional variant or an immunoconjugate of the invention for diagnostic or screening purposes.

A further aspect of the present invention relates to a kit comprising at least one isolated binding protein, at least one functional variant, at least one immunoconjugate and/or at least one pharmaceutical composition of the invention.

## BRIEF DESCRIPTION OF THE FIGURES

**Figure 1A** shows the nucleotide (SEQ ID N° 1) and corresponding amino acid (SEQ ID N° 50) sequences of the heavy chain variable domain (VH) and a partial region of the Constant Heavy 1 (CH1) domain of the **5F10F175E2** antibody and **Figure 1B** shows the nucleotide (SEQ ID N° 3) and corresponding amino acid (SEQ ID N° 51) sequences of whole light chain of the **5F10F175E2** antibody. Leader sequences are indicated in light grey. Variable domains (VH and VL, respectively) are highlighted in dark grey. The Complementary Determining Regions (CDR's) are also indicated for each of the variable domains.

**Figure 2A** shows the nucleotide (SEQ ID N° 21) and corresponding amino acid (SEQ ID N° 52) sequences of the heavy chain variable domain (VH) and a partial region of the Constant Heavy 1 (CH1) domain of the **8B10F8** antibody and **Figure 2B** shows the nucleotide (SEQ ID N° 23) and corresponding amino acid (SEQ ID N° 53) sequences of whole light chain of the **8B10F8** antibody. The leader sequence (5' Untranslated region; 5'UTR) is indicated in light grey. Variable domains (VH and VL, respectively) are highlighted in dark grey. The Complementary Determining Regions (CDR's) are also indicated for each of the variable domains.

**Figure 3** shows the results of the indirect immunofluorescence assay performed to assess the infection of HEK 293T cells by Chikungunya virus, confirming the *in vitro* neutralizing properties of the recombinant antibodies (**rec8B10F8** and **rec5F10F175E2**) and of the isolated antibodies (**8B10F8** and **5F10F175E2**) against the Chikungunya virus A226 and A226V strains. Images were obtained using a fluorescent microscope (x10 magnification, NIKON ECLIPSE TS 100).

**Figure 4** shows the dose-response neutralization curves of the recombinant mAbs (**rec8B10F8** and **rec5F10F175E2**) against Chikungunya virus A226 and A226V strains, as neutralization percentage (%) relative to an irrelevant IgG1. **rec5F10F175E2** antibody is represented as a square (-■-), **rec8B10F8** antibody is represented as a triangle (-▲-) and both antibodies used in combination as a dot (-●-).

**Figure 5** shows the results obtained by a western blot immunoassay on the recombinant antibodies (**rec8B10F8** and **rec5F10F175E2**) against infected cell lysates and purified viral particles of both A226 and A226V Chikungunya virus

strains. (E1/E2) represents the envelope glycoproteins E1 and E2 and (C) represents the viral capsid protein.

**Figure 6** shows the mouse signal peptide of Ig kappa chain (SEQ ID N° 48), followed by the first nucleotides/amino acids of the human IgG1 constant region sequence (SEQ ID N° 49). The Sal I and Nhe I sites are shown in bold case.

## DETAILED DESCRIPTION

### Binding proteins

In a first aspect, the present invention encompasses binding proteins capable of specifically binding to Chikungunya virus (CHIKV).

The term "specifically binding", as used herein, in reference to the interaction of a binding protein, e. g. an antibody, and its binding partner, e. g. an antigen, means that the interaction is dependent upon the presence of a particular structure, e. g. an antigenic determinant or epitope, on the binding partner. In other words, the antibody preferentially binds or recognizes the binding partner even when the binding partner is present in a mixture of other molecules or organisms. The binding may be mediated by covalent or noncovalent interactions or a combination of both. In yet other words, the term "specifically binding" means immunospecifically binding to an antigen or a fragment thereof and not immunospecifically binding to other antigens. A binding protein that immunospecifically binds to an antigen may bind to other peptides or polypeptides with lower affinity as determined by, e. g., radioimmunoassays (RIA), enzyme-linked immunosorbent assays (ELISA), BIACORE, or other assays known in the art. Binding proteins or fragments thereof that immunospecifically bind to an antigen may be cross-reactive with related antigens. Preferably, binding proteins or fragments thereof that immunospecifically bind to an antigen do not cross-react with other antigens.

Chikungunya virus is part of the genus *alphavirus*. According to Strauss and Strauss (Strauss and Strauss, 1994, Microbiol Rev 58, 491-562), in total, the genus *alphavirus* have 26 recognized members (i.e. genotypes). Alphaviruses have been classified in 4 main groups according to its serological cross-reaction, i.e., the

Venezuelan equine encephalitis (VEE/EEE) group, the Semliki Forest (SF) group, the Sindbis virus (SIN) group and the Western Equine Encephalitis (WEE) group. Table I below lists some of the currently recognized alphaviruses, together with their geographical distribution and their serological group. Besides binding to

5 Chikungunya virus, the binding proteins of the invention may also be capable of binding to other genotypes of the genus *alphavirus*, including but not limited to those listed in Table I.

Table I

| Group   | Virus                                | Geographic distribution          |
|---------|--------------------------------------|----------------------------------|
| VEE/EEE | Eastern equine encephalitis (EEE)    | America                          |
|         | Venezuelan equine encephalitis (VEE) | America                          |
|         | Everglades                           | America                          |
|         | Mucambo                              | America                          |
|         | Pixuna                               | America                          |
| SF      | Semliki Forest (SF)                  | Africa, Eurasia                  |
|         | Middelburg                           | Africa                           |
|         | <b>Chikungunya</b>                   | Africa, Asia                     |
|         | O'Nyong Nyong                        | Africa                           |
|         | Ross River                           | Australia                        |
|         | Barma Forest                         | Australia                        |
|         | Getah                                | Australia, Asia                  |
|         | Sagiyama                             | Japan                            |
|         | Bebaru                               | Malaysia                         |
|         | Mayaro                               | South America                    |
|         | Una                                  | South America                    |
|         | Ndumu                                | Africa                           |
| SIN     | Sindbis (SIN)                        | Africa, Asia, Europe, Australia, |

|     |                             |             |
|-----|-----------------------------|-------------|
|     |                             | Scandinavia |
|     | Aura                        | America     |
|     | Whataroa                    | New Zealand |
|     | Babanki                     | Africa      |
|     | Kyzylagash                  | Russia      |
| WEE | Western equine encephalitis | America     |

Furthermore, the binding proteins of the invention may even be capable of binding to viruses other than alphaviruses of the *Togaviridae* family, such as those belonging to the genus *rubivirus* or others. Further information on the *Togaviridae* family and its taxonomic structure and members can be found on the Index of Viruses - *Togaviridae* (2006). In: ICTVdB - The Universal Virus Database, version 4. Büchen-Osmond, C (Ed), Columbia University, New York, USA.

The binding proteins of the invention may be capable of specifically binding to Chikungunya virus in its natural form or in its inactivated/attenuated form. General viral inactivation methods well known to the skilled artisan such as inter alia pasteurization (wet heat), dry heat treatment, vapor heat treatment, treatment with low pH, treatment with organic solvent/detergent, nanofiltration and/or UV light irradiation may be used. Preferably, the inactivation is performed by heat-treatment for 1 hour at 56 °C.

The binding proteins of the invention may also be capable of specifically binding to one or more fragments of the Chikungunya virus such as inter alia a preparation of one or more proteins and/or (poly) peptides derived from Chikungunya virus or a cell transfected with a Chikungunya virus protein and/or (poly) peptide. For methods of treatment and/or prevention of Chikungunya virus the binding proteins of the invention are preferably capable of specifically binding to surface accessible proteins of Chikungunya virus such as the E1 or E2 glycoproteins (Strauss and Strauss, 1994, *Microbiol Rev* 58, 491-562). For diagnostic purposes the binding proteins of the invention may also be capable of specifically binding to proteins not present on the surface of Chikungunya virus. The amino acid sequence of surface accessible and internal proteins of various known strains

of Chikungunya virus can be found in the EMBL-database and/or other databases.

Preferably, the fragment at least comprises an antigenic determinant recognized by the binding proteins of the invention. An "antigenic determinant" as used herein is a moiety, such as a Chikungunya virus (poly) peptide, (glyco)protein, or analog or fragment thereof, that is capable of binding to a binding proteins of the invention with sufficiently high affinity to form a detectable antigen-binding protein complex.

Typically, binding proteins according to the invention can bind to their binding partners, i. e. Chikungunya virus or fragments thereof such as Chikungunya virus proteins, with an affinity constant ( $K_d$ -value) that is lower than  $0.2 \times 10^{-4}$  M,  $1.0 \times 10^{-5}$  M,  $1.0 \times 10^{-6}$  M,  $1.0 \times 10^{-7}$  M, preferably lower than  $1.0 \times 10^{-8}$  M, more preferably lower than  $1.0 \times 10^{-9}$  M, more preferably lower than  $1.0 \times 10^{-10}$  M, even more preferably lower than  $1.0 \times 10^{-11}$  M, and in particular lower than  $1.0 \times 10^{-12}$  M. The affinity constants can vary for different antibody isotypes. Affinity constants can for instance be measured using surface plasmon resonance, i.e. an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, in particular using the BIACORE system (GE Healthcare).

The binding proteins according to the invention may bind to Chikungunya virus in purified/isolated or non-purified/non-isolated form. The binding proteins may bind to Chikungunya virus in soluble form such as for instance in a sample or may bind to Chikungunya virus bound or attached to a carrier or substrate, e. g., microtiter plates, membranes and beads, etc. Carriers or substrates may be made of glass, plastic (e. g., polystyrene), polysaccharides, nylon, nitrocellulose, or teflon, etc. The surface of such supports may be solid or porous and of any convenient shape. Alternatively, the binding proteins may also bind to fragments of Chikungunya virus such as proteins or (poly) peptides of the Chikungunya virus. In an embodiment the binding proteins are capable of specifically binding to the Chikungunya virus E2 or E1 protein or to a fragment thereof. The Chikungunya virus proteins or (poly) peptides may either be in soluble form or may bind to

Chikungunya virus bound or attached to a carrier or substrate as described above. In another embodiment cells transfected with Chikungunya virus proteins or (poly) peptides may be used as binding partner for the binding proteins.

In one embodiment of the present invention, the isolated binding protein of the invention comprises a heavy chain amino acid sequence comprising at least one of the CDR's selected from the group consisting of: (a) CDRH1's of SEQ ID NOs: 10 or 30, (b) CDRH2's of SEQ ID NOs: 12 or 32, and (c) CDRH3's of SEQ ID NOs: 14 or 34, and/or a light chain amino acid sequence comprising at least one of the CDR's selected from the group consisting of: (d) CDRL1's of SEQ ID NOs: 16 or 36, (e) CDRL2's of SEQ ID NOs: 18 or 38, and (f) CDRL3's of SEQ ID NOs: 20 or 40. Preferably, the isolated binding protein comprises both the heavy chain amino acid sequence and the light chain amino acid sequence.

The term "complementarity determining regions" (CDR) as used herein means sequences within the variable regions of binding proteins, such as immunoglobulins, that usually contribute to a large extent to the antigen binding site which is complementary in shape and charge distribution to the epitope recognized on the antigen. The CDR regions can be specific for linear epitopes, discontinuous epitopes, or conformational epitopes of proteins or protein fragments, either as present on the protein in its native conformation or, in some cases, as present on the proteins as denatured, e. g., by solubilization in SDS. Epitopes may also consist of posttranslational modifications of proteins.

The CDR3 region of the variable domain of the heavy chain (CDRH3) provides typically with the greatest source of molecular diversity within the antibody binding-site (Xu et al., 2000, Immunity, 13(1):37-45). Accordingly, preferably, the isolated binding protein of the invention comprises a heavy chain amino acid sequence comprising at least one of the CDR's selected from SEQ ID NO: 14 and SEQ ID NO: 34.

In another embodiment of the present invention, the isolated binding protein of the invention comprises a heavy chain amino acid sequence that comprises a CDRH1 selected from SEQ ID NOs: 10 or 30, a CDRH2 selected from SEQ ID NOs: 12 or 32, and a CDRH3 selected from SEQ ID NOs: 14 or 34, and/or a

light chain amino acid sequence that comprises a CDRL1 selected from SEQ ID NOs: 16 or 36, a CDRL2 selected from SEQ ID NOs: 18 or 38, and a CDRL3 selected from SEQ ID NOs: 20 or 40. Preferably, said isolated binding protein comprises both the heavy chain amino acid sequence and the light chain amino acid sequence.

5 In a further embodiment, the isolated binding protein of the invention comprises a heavy chain amino acid sequence that comprises the CDRH1 of SEQ ID NO: 10, the CDRH2 of SEQ ID NO: 12, and the CDRH3 of SEQ ID NO: 14, and/or a light chain amino acid sequence that comprises the CDRL1 of SEQ ID NO: 16, the CDRL2 of SEQ ID NO: 18, and the CDRL3 of SEQ ID NO: 20.

10 In yet another embodiment, the isolated binding protein of the invention comprises a heavy chain amino acid sequence that comprises the CDRH1 of SEQ ID NO: 30, the CDRH2 of SEQ ID NO: 32, and the CDRH3 of SEQ ID NO: 34, and/or a light chain amino acid sequence that comprises the CDRL1 of SEQ ID NO: 36, the CDRL2 of SEQ ID NO: 38, and the CDRL3 of SEQ ID NO: 40.

15 In another embodiment, the isolated binding protein of the invention comprises a heavy chain amino acid sequence which comprises a VH domain amino acid sequence selected from the group consisting of SEQ ID Nos: 6 or 26, and/or a light chain amino acid sequence that comprises a VL domain amino acid sequence selected from the group consisting of SEQ ID NOs: 8 or 28. In a preferred embodiment, the isolated binding protein of the invention comprise the VH domain amino acid sequence of SEQ ID NO: 6 and the VL domain amino acid sequence of SEQ ID NO: 8. Alternatively, the isolated binding protein of the invention comprises the VH domain amino acid sequence of SEQ ID NO: 26 and the VL domain amino acid sequence of SEQ ID NO: 28.

25 In further embodiment, the isolated binding protein of the invention comprises a heavy chain amino acid sequence which comprises the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 22 and/or a light chain amino acid sequence that comprises the amino acid sequence of SEQ ID NO: 4 or SEQ ID NO: 24. In a preferred embodiment, the isolated binding protein of the invention comprises a heavy chain amino acid sequence which comprises the amino acid sequence of SEQ ID NO: 2 and a light chain amino acid sequence which

30

comprises the amino acid sequence of SEQ ID NO: 4. Alternatively, the isolated binding protein of the invention comprises a heavy chain amino acid sequence which comprises the amino acid sequence of SEQ ID NO: 22 and a light chain amino acid sequence which comprises the amino acid sequence of SEQ ID NO: 24.

In a particular embodiment, the binding protein of the invention is a scaffold protein having an antibody like binding activity or an antibody, i.e. an anti-CHIKV antibody.

Within the context of the present invention, the term "scaffold protein", as used herein, means a polypeptide or protein with exposed surface areas in which amino acid insertions, substitutions or deletions are highly tolerable. Examples of scaffold proteins that can be used in accordance with the present invention are protein A from *Staphylococcus aureus*, the bilin binding protein from *Pieris brassicae* or other lipocalins, ankyrin repeat proteins, and human fibronectin (reviewed in Binz and Plückthun, 2005, *Curr Opin Biotechnol* 16, 459-69 and in Plückthun, A., 2009, *Recombinant Antibodies for Immunotherapy: Alternative Scaffolds: Expanding the Options of Antibodies* (Little, M., ed), pp. 243-271, Cambridge University Press, New York). Engineering of a scaffold protein can be regarded as grafting or integrating an affinity function onto or into the structural framework of a stably folded protein. Affinity function means a protein binding affinity according to the present invention. A scaffold can be structurally separable from the amino acid sequences conferring binding specificity. A scaffold protein having an antibody like binding activity can for instance be derived from an acceptor polypeptide containing the scaffold domain, which can be grafted with binding domains of a donor polypeptide to confer the binding specificity of the donor polypeptide onto the scaffold domain containing the acceptor polypeptide. Said inserted binding domains may be, for example, the complementarity determining region (CDR) of an antibody, in particular an anti-CHIKV antibody. In one embodiment of the present invention, at least one of said inserted binding domains is one of the CDRs of the antibodies identified in the Examples (i.e. **8B10F8**, **5F10F175E2**, **rec8B10F8** and **rec5F10F175E2** antibodies), as

above described. Insertion can be accomplished by various methods known to those skilled in the art including, for example, polypeptide synthesis, nucleic acid synthesis of an encoding amino acid as well by various forms of recombinant methods well known to those skilled in the art.

5           In a preferred embodiment, the binding protein of the invention is an antibody. The term "antibody" or "anti-CHIKV antibody", as used herein, means a monoclonal antibody, a polyclonal antibody, a recombinant antibody, a humanized antibody (Jones et al., 1986, Nature 321, 522-525; Riechmann et al., 1988, Nature 332, 323-329; and Presta, 1992, Curr. Op. Struct. Biol. 2, 593-596), a  
10       chimeric antibody (Morrison et al., 1984, Proc. Natl. Acad. Sci. U.S.A. 81, 6851-6855), a multispecific antibody (e.g. a bispecific antibody) formed from at least two antibodies, or an antibody fragment thereof. The term "antibody fragment" comprises any portion of the afore-mentioned antibodies, preferably their antigen binding or variable regions. Examples of antibody fragments include Fab  
15       fragments, Fab' fragments, F(ab')<sub>2</sub> fragments, Fv fragments, diabodies (Hollinger et al., 1993, Proc. Natl. Acad. Sci. U.S.A. 90, 6444-6448), single chain antibody molecules (Plückthun, A., 1994, in The pharmacology of monoclonal antibodies: Antibodies from Escherichia coli (Rosenberg, M., and Moore, G. P., eds), Vol. 113, pp. 269-315, Springer Verlag, Berlin) and other fragments as long as they exhibit  
20       the desired capability of binding to Chikungunya virus.

          In addition, the term "antibody" or "anti-CHIKV antibody", as used herein, may include antibody-like molecules that contain engineered sub-domains of antibodies or naturally occurring antibody variants. These antibody-like molecules may be single-domain antibodies such as VH-only or VL-only domains derived  
25       either from natural sources such as camelids (Muyldermans et al., 2001, J Biotechnol. 74(4), 277-302) or through *in vitro* display of libraries from humans, camelids or other species (Holt et al., 2003, Trends Biotechnol. 21, 484-90 ).

          In another preferred embodiment the binding protein of the invention is an antibody fragment.

30           In accordance with the present invention, the "Fv fragment" is the minimum antibody fragment that contains a complete antigen-recognition and -binding

site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDR's of each variable domain interact to define an antigen- binding site on the surface of the VH-VL dimer. Collectively, the six CDR's confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDR's specific for an antigen) has the ability to recognize and bind the antigen, although usually at a lower affinity than the entire binding site. Derivatives of Fv fragments, such as scFv (single-chain Fv) and dsFv (disulfide-stabilized Fv), which have been modified to increase stability of the recombinant Fv fragment are also included. The "Fab fragment" also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. The "Fab fragment" differs from the "Fab' fragment" by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. The "F(ab')<sub>2</sub> fragment" originally is produced as a pair of "Fab' fragments" which have hinge cysteines between them. Methods of preparing such antibody fragments, such as papain or pepsin digestion, are known to those skilled in the art.

In a further preferred embodiment of the present invention, the anti-CHIKV antibody of the invention is of the IgA-, IgD-, IgE-, IgG- or IgM-type, preferably of the IgG- or IgM-type including, but not limited to, the IgG1-, IgG2-, IgG3-, IgG4-, IgM1- and IgM2-type. In most preferred embodiments, the antibody is of the IgG1-, IgG2- or IgG4- type. More preferably, of the IgG1- type.

In certain respects, e.g. in connection with the generation of antibodies as therapeutic candidates against Chikungunya virus, it may be desirable that the anti-CHIKV antibody of the invention is capable of fixing complement and participating in complement-dependent cytotoxicity (CDC). There are a number of isotypes of antibodies that are capable of the same including without limitations the following: murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, human IgG3, and human IgA. It will be appreciated that antibodies that are generated need not initially possess such an isotype but, rather the antibody as generated can possess any isotype and the antibody can be isotype

switched by appending the molecularly cloned V region genes or cDNA to molecularly cloned constant region genes or cDNAs in appropriate expression vectors using conventional molecular biology techniques that are well known in the art and then expressing the antibodies in host cells using techniques known in the art. The isotype-switched antibody may also possess an Fc region that has been molecularly engineered to possess superior CDC over naturally occurring variants (Idusogie et al., 2001, J Immunol. 166, 2571-2575) and expressed recombinantly in host cells using techniques known in the art. Such techniques include the use of direct recombinant techniques (see e.g. US 4,816,397), cell-cell fusion techniques (see e.g. US 5,916,771 and US 6,207,418), among others. In the cell-cell fusion technique, a myeloma or other cell line such as CHO is prepared that possesses a heavy chain with any desired isotype and another myeloma or other cell line such as CHO is prepared that possesses the light chain. Such cells can, thereafter, be fused and a cell line expressing an intact antibody can be isolated. By way of example, a human anti-CHIKV IgG4 antibody, that possesses the desired binding to the Chikungunya virus antigen, could be readily isotype switched to generate a human IgM, human IgG1 or human IgG3 isotype, while still possessing the same variable region. Such molecule might then be capable of fixing complement and participating in CDC.

Moreover, it may also be desirable for the anti-CHIKV antibody of the invention to be capable of binding to Fc receptors on effector cells, such as monocytes and natural killer (NK) cells, and participate in antibody- dependent cellular cytotoxicity (ADCC). There are a number of isotypes of antibodies that are capable of the same, including without limitations the following: murine IgG2a, murine IgG2b, murine IgG3, human IgG1 and human IgG3. It will be appreciated that antibodies that are generated need not initially possess such an isotype but, rather the antibody as generated can possess any isotype and the antibody can be isotype switched by appending the molecularly cloned V region genes or cDNA to molecularly cloned constant region genes or cDNAs in appropriate expression vectors using conventional molecular biological techniques that are well known in the art and then expressing the antibodies in host cells using

techniques known in the art. The isotype-switched antibody may also comprise an Fc region that has been molecularly engineered to possess superior ADCC over naturally occurring variants (Shields et al., 2001, J Biol Chem. 276, 6591- 6604) and expressed recombinantly in host cells using techniques known in the art. Such techniques include the use of direct recombinant techniques (see e.g. US 4,816,397), cell-cell fusion techniques (see e.g. US 5,916,771 and US 6,207,418), among others. Such molecule might then be capable of binding to FcγR on effectors cells and participating in ADCC.

Antibodies with superior ADCC activity may be obtained by modifying the oligosaccharides profile of IgG's, by using different strategies such as for example, glycosylation inhibition, genetic modifications, amino acid changes in FcR, transgenesis or production in cell lines producing naturally unfucosylated antibodies. Examples of particular genetic engineering strategies are those used by companies such as GLYCART, KYOWA or LFB. GLYCART BIOTECHNOLOGY AG (Zurich, CH) has expressed N-acetyl-glucosaminyltransferase III (GnTIII) which catalyzes the addition of the bisecting GlcNac residue to the N-linked oligosaccharide, in a CHO cell line, and showed a greater ADCC of the IgG1 antibody produced (WO 99/54342; WO 03/011878; WO 2005/044859). On the other hand, by removing or supplanting fucose from the Fc portion of the antibody, KYOWA HAKKO KOGYO (Tokyo, Japan) has enhanced Fc binding and improved ADCC, and thus the efficacy of the Mab (US 6,946,292). More recently, Laboratoire Francais du Fractionnement et des Biotechnologies (LFB) (France) showed that the ratio Fuc/Gal in Mab oligosaccharide should be equal or lower than 0.6 to get antibodies with a high ADCC (FR 2 861 080). Preferably, an antibody with a glycosylation profile providing enhanced cell-mediated effector functions is obtained by expressing said antibody in a cell line producing naturally unfucosylated antibodies, such as for example, an avian cell, preferably a duck cell. Particularly interesting is the production of an antibody of the invention in an avian embryonic derived stem cell line EBx® marketed by Vivalis (Nantes, France) as described in WO 2008/142124, which will provide the antibody with an optimized ADCC activity. Furthermore, an antibody produced in the avian

embryonic derived stem cell line EBx® will display a human-like glycosylation pattern.

Thus, according to a preferred embodiment, an antibody of the invention has an optimized ADCC activity. Preferably, said antibody has been produced in an avian embryonic derived stem cell line EBx® and thus, has a glycosylation profile providing it with enhanced cell-mediated effector functions. More preferably, the avian embryonic derived stem cell line EBx® is from chicken or duck, and more preferably are chicken EB 14 cells, duck EB 66 or duck EB 24 cells, which may be genetically engineered to express recombinant proteins.

The binding proteins of the invention can be obtained or derived by a variety of ways, specifically by using well known methods for obtaining antibodies or antibody fragments, such as generation of antibodies by the use of mouse hybridomas (see, for example, Köhler et al., 1975, *Nature* 256, 495-97), production of chimaeric (Hardman et al., 1989, *Int J Cancer.*, 44(3), 424-33) or humanized (Winter and Harris, 1993, *Immunology Today* 14 (6), 243-246) antibodies or antibody fragments by using recombinant DNA techniques. Preferably, fully human antibodies are obtained using technologies for the production of mAbs derived from human immunoglobulin gene sequences, such as, genetically engineered animals (e.g. Xenomouse® strains (Abgenix, Inc., Fremont, Canada), by using recombinant library methods, such as phage display, yeast display, ribosome display, *E.coli* display, etc. (See, e.g., Clackson et al., 1991, *Nature* 352, 624-628 ; Marks et al., 1991, *J. Mol. Biol.* 222, 581-597; Feldhaus and Siegel, 2004, *J Immunol Methods.* 290, 69-80; Groves and Osbourn, 2005, *Expert Opin Biol Ther.* 5(1), 125-135; and Jostock and Dubel, 2005, *Comb. Chem. High Throughput Screen.* 8, 127-133). Preferably, native human antibodies are obtained from recently developed technologies, including the use of human B cells directly by Human-Human hybridoma (Karpas et al., 2001, *Proc Natl Acad Sci U S A.* 98(4), 1799-804), Hybrid hybridoma (Schmidt E, 2001, *J Immunol Methods.* 255(1-2), 93-102), B cell immortalization and cloning (Lanzavecchia et al., *Curr Opin Biotech*, 2007, 18, 523-8), genetic programming of immortalized B cells (Kwakkenbos et al., 2009, *Nature*

Med 16(1):123-129) or Single-cell RT-PCR (Tiller et al., 2008, J Immunol Methods 329(1-2), 112-124; Wrammert et al., 2008, Nature 453(7195):667-71).

Particularly preferred is the generation of native human antibodies from human B cells, such as described in the Examples of the invention

5 Accordingly, the antibody of the invention could be an antibody of animal origin (e.g. murine), a chimaeric, humanized or fully human antibody.

A problem with murine antibodies derives from the fact that there are many sequence differences between rodent immunoglobulins and human immunoglobulins (Kabat EA, Wu TT, Perry HM, et al. Sequences of proteins of immunological interest. US Department of Health and Human Services, U.S. Government Printing Office, 1991.). Consequently, use of rodent monoclonal antibodies into a human recipient usually results in an antiglobulin response, detectable at about 8-12 days with a peak at about 20-30 days (Isaacs JD. The antiglobulin response to therapeutic antibodies. Seminars in Immunology 1990; 2: 10 449-456). The presence of this immunological response will usually render the treatment inoperative after 10 days. Furthermore, later retreatment is not possible, due to the rapid onset of a secondary response.

In order to avoid immune response from the patient, is thus important to use monoclonal antibodies that are as near as possible to human antibodies. Thus, by re-engineering and de-immunization techniques chimaeric antibodies and humanized antibodies may be obtained based on murine antibodies. In a chimaeric antibody, the whole of the variable regions of a mouse or rat (or of a non-human) antibody are expressed along with human constant regions. This will lead to an antibody with proper human effector functions, while decreasing the immunogenicity caused by the xenogeneic Fc region. In a humanized antibody, only the CDR's from the rodent antibody V-regions are combined with framework regions from human V-regions. It is expected that these antibodies should thus be less immunogenic than chimaeric antibodies.

In a preferred embodiment, the antibody of the invention is fully human. Preferably, the antibody of the invention is a native human antibody or antibody fragment.

A "fully human antibody" is an antibody containing exclusively human sequences. Thus, a fully human antibody shall not induce an immune response when administered to a human recipient. Preferably, a human antibody of the invention is a "native human antibody", in which the antibody is naturally occurring in a human, as opposed to a human antibody in which the individual heavy and light chains are isolated from humans but are assembled randomly (i.e. by using library methods such as phage display) creating all forms of natural and unnatural antibodies.

Specifically, "native human antibodies" are those that arise naturally as the result of the functioning of an intact human immune system. The utility of native antibodies for the treatment of human viral diseases has been established through experience with hyperimmune human globulins. Native antibodies, as a class, differ in some respects from those obtained by library methods (phage or transgenic mouse) and possess distinct properties that may make them ideal therapeutics for human diseases. (See Dessain et al, Exploring the Native Human Antibody Repertoire to Create Antiviral Therapeutics in Current Topics in Microbiology and Immunology 317: 155-183 (2008), (c) Springer-Verlag New York). Specifically, there is a specific advantage of native antibodies expressed from human B cells over phage-derived antibodies, due to the limitations in a phage approach to recreate all of the original or native heavy chain: light chain pairings, thus preventing important antibody structures from being incorporated into a phage-generated library. The term "native human antibodies" includes "native human antibody fragments" as described herein and in particular, Fv fragments (including derivatives thereof such as scFv and dsFv), Fab fragments, Fab' fragments or F(ab')<sub>2</sub> fragments. Specifically, the binding site of these "native human antibody fragments" will correspond to that of a native human antibody (i.e., particular combination of heavy and light chain sequences naturally occurring in a human).

Human antibodies of the invention, however, may contain residues or modifications (such as post-translational modifications) not found in a naturally occurring human antibody, including those modifications and variant sequences

described herein. These modifications are typically made to further refine or enhance antibody desired properties, such as those providing a better performance, increased antibody life-time, increased establiity, increased ADCC activity, etc.

5           In addition, the binding proteins of the invention preferably neutralize Chikungunya virus infectivity. Accordingly, in a further preferred embodiment, the invention relates to a binding protein of the invention having Chikungunya virus neutralizing activity. This may be achieved by preventing the attachment of Chikungunya virus to its receptors on host cells or inhibition of the release of RNA  
10   into the cytoplasm of the cell or prevention of RNA transcription or translation.

          Preferably, the binding proteins of the invention may also be capable of neutralizing other genotypes of the genus *alphavirus*. Furthermore, the binding proteins of the invention may even be capable of neutralizing infectivity of viruses other than alphaviruses of the *Togaviridae* family, such as those belonging to the  
15   genus *rubivirus* or others. Further information on the *Togaviridae* family and its taxonomic structure and members can be found on the Index of Viruses - *Togaviridae* (2006). In: ICTVdB - The Universal Virus Database, version 4. Büchen-Osmond, C (Ed), Columbia University, New York, USA.

          The binding proteins of the invention may prevent Chikungunya virus from  
20   infecting host cells by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to infection of host cells by Chikungunya virus in the absence of said binding proteins. The neutralizing activity of the binding protein may be measured,  
25   for instance, by a standard plaque reduction neutralization test (PRNT) as shown in Example 4 and Figure 3 or by other *in vitro* neutralization assays, such as rapid fluorescent focus inhibition test (RFFIT), (Vene et al., 1998, Journal of Virological Methods 73(1), Pages 71-75). The neutralizing activity could also be investigated by analyzing the cell supernatant for the presence of Chikungunya viral genome  
30   by quantitative real time PCR (qRT-PCR).

Alternatively, the binding protein may be characterized in that it has a Chikungunya virus neutralizing activity of at least 2500 IU/mg protein. More preferably, said binding protein has a Chikungunya virus neutralizing activity of at least 2800 IU/mg protein, at least 3000 IU/mg protein, at least 3200 IU/mg protein, at least 3400 IU/mg protein, at least 3600 IU/mg protein, at least 3800 IU/mg protein, at least 4000 IU/mg protein, at least 4200 IU/mg protein, at least 4400 IU/mg protein, at least 4600 IU/mg protein, at least 4800 IU/mg protein, at least 5000 IU/mg protein, at least 5200 IU/mg protein, at least 5400 IU/mg protein. The neutralizing activity of the binding protein may be measured, for instance, by as standard plaque reduction neutralization test (PRNT) as shown in Example 4 and Figure 3 or by other *in vitro* neutralization assays, such as rapid fluorescent focus inhibition test (RFFIT), (Vene et al., 1998, Journal of Virological Methods 73(1), Pages 71-75). The neutralizing activity could also be investigated by analyzing the cell supernatant for the presence of Chikungunya viral genome by quantitative real time PCR (qRT-PCR).

In a preferred embodiment, the neutralizing protein of the invention is selected from the anti-CHIKV neutralizing fully human antibodies shown in the Examples (**8B10F8**, **5F10F175E2**, **rec8B10F8** and **rec5F10F175E2**). More preferably, from the IgG1 type recombinant antibodies **rec8B10F8** and **rec5F10F175E2** which are characterized in Example 4.

#### Functional variants

A second aspect of the invention includes functional variants of binding proteins as defined herein.

The term "functional variant", as used herein, refers to a binding molecule that comprises a nucleotide and/or amino acid sequence that is altered by one or more nucleotides and/or amino acids compared to the nucleotide and/or amino acid sequences of the parent binding protein and that is still capable of competing for binding to the binding partner, e.g. Chikungunya virus or a fragment thereof, with the parent binding molecule. In other words, the modifications in the amino acid and/or nucleotide sequence of the parent

binding molecule do not significantly affect or alter the binding characteristics of the binding molecule encoded by the nucleotide sequence or containing the amino acid sequence, i. e. the binding molecule is still able to recognize and bind its target. Whether a modification in the amino acid sequence results in a functional binding protein (i.e. in a binding protein that binds to Chikungunya virus or a fragment thereof), can readily be determined by assaying the specific activity of the resulting binding protein in ELISA or FACS for binding to Chikungunya virus or a fragment thereof or other *in vitro* or *in vivo* functional assay. Preferably, the functional variants should also have Chikungunya virus neutralizing activity.

The functional variant may preferably have conservative sequence modifications including nucleotide and amino acid substitutions, additions and deletions. Furthermore, functional variants can comprise truncations of the amino acid sequence at either or both the amino or carboxy termini. These modifications can be introduced by standard techniques known in the art, such as site-directed mutagenesis and random PCR-mediated mutagenesis, and may comprise natural as well as non-natural nucleotides and amino acids.

Non-natural amino acids may include, for example, stereoisomers (e.g. D-amino acids) of the 20 conventional amino acids, unnatural amino acids such as  $\alpha$ -,  $\alpha$ -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids. Examples of unconventional amino acids, which may also be suitable components for the binding protein of the invention, include: 4-hydroxyproline,  $\gamma$ -carboxyglutamate,  $\epsilon$ -N,N,N-trimethyllysine,  $\epsilon$ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine,  $\sigma$ -N-methylarginine, and other similar amino acids and imino acids, e.g. 4-hydroxyproline.

Especially preferred variations in the nucleotide or amino acid sequences shown in SEQ ID NOs: 1-46 are those that lead to a reduced susceptibility to proteolysis or oxidation, alter glycosylation patterns or alter binding affinities or confer or modify other physicochemical or functional properties of the binding protein. In particular, conservative amino acid replacements are contemplated. Conservative amino acid substitutions include the ones in which the amino acid

residue is replaced with an amino acid residue having similar structural or chemical properties. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e. g., lysine, arginine, histidine), acidic side chains (e. g., aspartic acid, glutamic acid), uncharged polar side chains (e. g., asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (e. g., glycine, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e. g., threonine, valine, isoleucine) and aromatic side chains (e. g., tyrosine, phenylalanine, tryptophan). It will be clear to the skilled artisan that other classification of amino acid residue families than the one used above can also be employed. Furthermore, a variant may have non-conservative amino acid substitutions, e.g., replacement of an amino acid with an amino acid residue having different structural or chemical properties. Similar minor variations, may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing immunological activity may be found using computer programs well known in the art.

Functional variants according to the invention may have the same or different, either higher or lower, binding affinities compared to the parent binding molecule but are still capable of binding to Chikungunya virus or a fragment thereof and preferably, still capable of neutralizing Chikungunya virus. For instance, functional variants according to the invention may have increased or decreased binding affinities for Chikungunya virus or a fragment thereof compared to the parent binding proteins or may have a higher or lower Chikungunya virus neutralizing activity. Preferably, the amino acid sequences of the variable regions, including, but not limited to, framework regions, hypervariable regions, in particular the CDR3 regions, are modified. Functional variants intended to fall within the scope of the present invention have at least about 50% to about 99%, preferably at least about 60% to about 99%, more preferably at least about 70% to about 99%, even more preferably at least about 80% to about 99%, most preferably at least about 90% to about 99%, in particular

at least about 95% to about 99%, and in particular at least about 97% to about 99% amino acid sequence identity with the parent binding proteins as defined herein. Computer algorithms such as inter alia Gap or Bestfit known to a person skilled in the art can be used to optimally align amino acid sequences to be compared and to define similar or identical amino acid residues.

Moreover, functional variants also include derivatives that are substantially similar in primary structural sequence, but which contain e. g. *in vitro* or *in vivo* modifications, chemical and/or biochemical, that are not found in the parent binding protein. Such modifications include inter alia acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma- carboxylation, glycosylation, GPI-anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, ubiquitination, and the like.

#### Immunoconjugates

The binding protein or a functional variant of the invention may be coupled to at least one labeling and/or effector group. Thus, in yet a further aspect, the invention includes immunoconjugates, i.e. a binding protein or a functional variant of the invention which is coupled to at least one labeling and/or effector group.

In one embodiment of the invention, a binding protein or a functional variant of the invention is coupled to at least one effector group. Such an immunoconjugate is especially suitable for therapeutic applications. As used herein, the term "effector group" refers to a therapeutic group, a toxin, a cytotoxic group, an antigen or other effector group known in the art. In a particular embodiment, a binding protein or a functional variant of the invention may be

conjugated/attached to one or more antigens. Preferably, these antigens are antigens which are recognized by the immune system of a subject to which the binding molecule-antigen conjugate is administered to. The antigens may be identical but may also differ from each other. In another particular embodiment, said effector group is a therapeutic agent. Preferably, said therapeutic agent is suitable for the treatment of a viral infection or the symptoms associated to a viral disease, such as chloroquine phosphate, paracetamol, NSAID's (e.g., ibuprofen, naproxen, etc.) or corticosteroids. More preferably, said therapeutic agent is an anti-viral such as ribavirin or interferon-alpha or acyclovir. Alternatively, said therapeutic agent is useful for treating secondary effects related to the binding protein or functional variant of the invention. In certain respects, it may be desirable that the effector groups are attached by spacer arms of various lengths to reduce potential steric hindrance.

In another embodiment of the present invention, a binding protein or a functional variant of the invention is coupled to at least one labelling group. Such an immunoconjugate is particularly suitable for diagnostic applications, for example, assess if a subject has been infected with Chikungunya virus or monitor the development or progression of Chikungunya virus infection as part of a clinical testing procedure to, e. g., determine the efficacy of a given treatment regimen. However, they may also be used for other detection and/or analytical and/or diagnostic purposes. As used herein, the term "labelling group" refers to a detectable marker, e.g. a radiolabeled amino acid or biotinyl moiety that can be detected by marked avidin (e.g. streptavidin bound to a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). Various methods for labelling polypeptides and glycoproteins, such as antibodies, are known in the art and may be used in performing the present invention. Examples of suitable labelling groups include, but are not limited to, the following: radioisotopes or radionuclides (e.g.  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{35}\text{S}$ ,  $^{90}\text{Y}$ ,  $^{99}\text{Tc}$ ,  $^{111}\text{In}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ), fluorescent groups (e.g. FITC, rhodamine, lanthanide phosphors), enzymatic groups (e.g. horseradish peroxidase,  $\beta$ -galactosidase, luciferase, alkaline phosphatase), chemiluminescent groups, biotinyl groups, or predetermined

polypeptide epitopes recognized by a secondary reporter (e.g. leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In certain respects, it may be desirable that the labelling groups are attached by spacer arms of various lengths to reduce potential steric hindrance.

5           The labeling groups used to label the binding protein or the functional variant of the invention for detection and/or analytical and/or diagnostic purposes depend on the specific detection/analysis/diagnosis techniques and/or methods used such as inter alia immunohistochemical staining of (tissue) samples, flow cytometric detection, scanning laser cytometric detection, fluorescent  
10 immunoassays, enzyme-linked immunosorbent assays (ELISA's), radioimmunoassays (RIA's), bioassays (e. g., neutralization assays), Western blotting applications, etc.

          When the binding proteins or functional variants of the invention are used for *in vivo* diagnostic use, said binding molecules can also be made detectable by conjugation to e. g. magnetic resonance imaging (MRI) contrast agents, such  
15 as gadolinium diethylenetriaminepentaacetic acid, to ultrasound contrast agents or to X-ray contrast agents, or by radioisotopic labeling.

          Also contemplated in the present invention are mixtures of immunoconjugates according to the invention or mixtures of at least one of said immunoconjugates according to the invention and another molecule, such as a  
20 therapeutic agent or another binding molecule. In a further embodiment, the immunoconjugates of the invention may comprise one or more label and/or effector group. These can be the same or distinct from each other and can be joined/conjugated non-covalently to the binding molecules. The labelling and/or effector group (s) can also be joined/conjugated directly to the binding  
25 molecules through covalent bonding, including, but not limited to, disulfide bonding, hydrogen bonding, electrostatic bonding, recombinant fusion and conformational bonding. Alternatively, these can be joined/conjugated to the binding molecules by means of one or more linking compounds. Techniques for conjugating said groups to binding molecules are well known to the skilled artisan.

30           Furthermore, the binding proteins, functional variants thereof or immunoconjugates of the invention (i.e., binding molecules of the invention) can

also be attached to solid supports, which are particularly useful for *in vitro* immunoassays or purification of Chikungunya virus or a fragment thereof. Such solid supports might be porous or nonporous, planar or nonplanar and include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene supports. The binding molecules of the invention can also for example usefully be conjugated to filtration media, such as NHS-activated Sepharose or CNBr-activated Sepharose for purposes of immunoaffinity chromatography. They can also usefully be attached to paramagnetic microspheres, typically by biotin-streptavidin interaction. The microspheres can be used for isolation of Chikungunya virus or a fragment thereof from a sample containing Chikungunya virus or a fragment thereof. As another example, the binding molecules of the present invention can usefully be attached to the surface of a microtiter plate for ELISA.

#### 15 Nucleic acid molecules

Another aspect of the present invention relates to an isolated nucleic acid molecule encoding a binding protein, a functional variant, an immunoconjugate of the invention or a fragment thereof. Preferably, the nucleic acid molecule comprises one of the nucleotide sequences of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 42, 43, 44, 45 or 46.

In one embodiment, the isolated nucleic acid molecule of the invention comprises a nucleotide sequence encoding a heavy chain amino acid sequence of a binding protein of the invention comprising at least one of the sequences selected from the group consisting of SEQ ID NOs: 9, 11, 13, 29, 31, 33 or 46. Preferably, the nucleic acid molecule of the invention comprises a nucleotide sequence of SEQ ID NO: 13 or 33.

In another embodiment, the isolated nucleotide molecule of the invention comprises a nucleotide sequence encoding a light chain amino acid sequence of a binding protein of the invention comprising at least one of the sequences selected from the group consisting of SEQ ID NOs: 15, 17, 19, 35, 37 or 39.

In another embodiment, the isolated nucleic acid molecule of the invention comprises a nucleotide sequence encoding a heavy chain amino acid sequence of a binding protein of the invention comprising the sequences of SEQ ID NOs: 9 or 29, and SEQ ID NOs: 11 or 31 and SEQ ID NOs: 13 or 33. In a further embodiment, the nucleic acid molecule of the invention comprises a nucleotide sequence encoding a light chain amino acid sequence of a binding protein of the invention comprising the sequences of SEQ ID NOs: 15 or 35, and SEQ ID NOs: 17 or 37 and SEQ ID NOs: 19 or 39.

In a further embodiment, the isolated nucleic acid molecule of the invention comprises a nucleotide sequence encoding a heavy chain amino acid sequence of a binding protein of the invention comprising a sequence selected from the group consisting of SEQ ID NO: 5, 25 or 45. In another embodiment, the nucleic acid molecule of the invention comprises a nucleotide sequence encoding a heavy chain amino acid sequence of a binding protein of the invention comprising a sequence of SEQ ID NO: 1, 21, 41 or 43.

In an even further embodiment, the isolated nucleic acid molecule of the invention comprises a nucleotide sequence encoding a light chain amino acid sequence of a binding protein of the invention comprising a sequence selected from the group consisting of SEQ ID NO: 7 and 27. In another embodiment, the nucleic acid molecule of the invention comprises a nucleotide sequence encoding a light chain amino acid sequence of a binding protein of the invention comprising a sequence of SEQ ID NO: 3, 23, 42 or 44.

Within the context of the present invention, the term "isolated nucleic acid molecule", as used herein, means a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin, the "isolated nucleic acid molecule" (1) is not associated with all or a portion of a polynucleotide in which the "isolated polynucleotide" is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence. Further, the term "nucleic acid molecule", as referred to herein, means a polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides, or a modified form

of either type of nucleotide, such as nucleotides with modified or substituted sugar groups and the like. The term also includes single and double stranded forms of DNA.

In a one embodiment of the present invention, a nucleic acid molecule of the invention is operably linked to a control sequence. The term "control sequence", as used herein, refers to polynucleotide sequences that are necessary to effect the expression and processing of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism. In prokaryotes, such control sequences generally include promoters, ribosomal binding sites, and transcription termination sequences. In eukaryotes, generally, such control sequences include promoters and transcription termination sequences. In accordance with the present invention, the term "control sequence" is intended to include, at a minimum, all components whose presence is essential for expression and processing, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences. Furthermore, the term "operably linked", as used herein, refers to positions of components so described which are in a relationship permitting them to function in their intended manner. Moreover, according to the present invention, an expression control sequence operably linked to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the expression control sequence.

#### Vector

A further aspect of the present invention is a vector comprising one or more nucleic acid molecule that encodes a binding protein, functional variant or immunoconjugate of the invention. Vectors may be autonomously replicating or may replicate together with the chromosome into which they have been integrated. The nucleic acid molecule can be operably linked to a control sequence. Furthermore, the vector may additionally contain a replication origin or a selection marker gene. Examples of vectors that may be used in accordance with the present invention are e.g. plasmids, cosmids, phages, viruses, etc., Vectors

can be used for cloning and/or for expression of the binding molecules of the invention.

Preferably, said vectors are used in eukaryotic cells, preferably, in mammalian cells. Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) typically also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding anti-PSCA antibody. One useful transcription termination component is the bovine growth hormone polyadenylation region. See for example, WO 94/11026 and the expression vector disclosed therein.

#### 15 Host

Another aspect of the present invention relates to hosts containing one or more copies of the nucleic acid molecules or vectors mentioned above. Preferably, the hosts are host cells. Host cells include, but are not limited to, cells of mammalian, plant, insect, fungal or bacterial origin. Transformation could be done by any known method for introducing polynucleotides into a host cell, including for example packaging the polynucleotide in a virus (or into a viral vector) and transducing a host cell with the virus (or vector) or by transfection procedures known in the art, as exemplified by US 4,399,216, US 4,912,040, US 4,740,461, and US 4,959,455, which patents are hereby incorporated herein by reference. Particularly, methods for introducing heterologous polynucleotides into host cells are well known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei. Examples of host cells that may be used according to the present invention include but are not limited to eukaryotic cells such as mammalian cells, e.g. hamster, rabbit, rat, pig, mouse, etc.; avian cells,

e.g. duck, chicken, quail, etc.; insect cells or other animal cells; plant cells and fungal cells, e.g. corn, tobacco, *Saccharomyces cerevisiae*, *Pichia pastoris*; prokaryotic cells such as *E. coli*; and other cells used in the art for the production of antibodies and other binding proteins. Especially mammalian cell lines available as hosts for expression are well known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g. Hep G2), and a number of other cell lines. Preferably, the host cells are human cells. Examples of human cells are inter alia HeLa, 911, AT1080, A549, 293 and HEK293T cells. In another preferred embodiment, the host cells are non-immortalized cell lines, more preferably avian embryonic stem cells. Particularly preferred host cells are avian embryonic derived stem cell lines marketed under the trademark EBx® by Vivalis (Nantes, France). The avian embryonic derived stem cell lines EBx® are described, for example, in WO 03/076601 and in WO 2008/142124 which are hereby incorporated by reference. The avian embryonic derived stem cell lines EBx® are able to proliferate, in adherence or non-adherence conditions, in a basal medium in absence of growth factors, serum and/or feeder cells. Accordingly, EBx® cell lines are especially suitable for its industrial use in the obtaining of molecules of interest, in particular for the production of viral vaccines or proteins (e.g. the binding protein or functional variant of the present invention). Preferably, the EBx® cells are from chicken or duck, and more preferably are chicken EB 14 cells, duck EB 66 or duck EB 24 cells. Furthermore, as described in WO 2008/142124, antibodies expressed in avian embryonic derived stem cell line EBx®, e.g., EB 66 cell line, have a glycosylation profile providing the antibody with enhanced cell-mediated effector functions. Furthermore, said antibody will display a human-like glycosylation pattern.

A process for producing the binding protein, functional variant or immunoconjugate of the invention

A further aspect of the present invention relates to a method for producing a binding protein, functional variant or immunoconjugate of the invention (i.e., binding molecules of the invention) by preparing said molecule(s) of the invention from a host cell that secretes said molecule(s). Examples of host cells that may be used according to the present invention include but are not limited to eukaryotic cells such as mammalian cells, e.g. hamster, rabbit, rat, pig, mouse, etc.; avian cells, e.g. duck, chicken, quail, etc.; insect cells or other animal cells; plant cells and fungal cells, e.g. corn, tobacco, *Saccharomyces cerevisiae*, *Pichia pastoris*; prokaryotic cells such as *E. coli*; and other cells used in the art for the production of antibodies and other binding proteins as above described. Various methods for preparing and isolating binding proteins, such as scaffold proteins or antibodies, from host cells are known in the art and may be used in performing the present invention. Moreover, methods for preparing binding protein fragments, e.g. scaffold protein fragments or antibody fragments, such as papain or pepsin digestion, modern cloning techniques, techniques for preparing single chain antibody molecules (Pluckthun in: *The Pharmacology of Monoclonal Antibodies* 113, Rosenberg and Moore, EDS, Springer Verlag, N.Y. (1994), 269-315) and diabodies (Hollinger et al., 1993, *Proc. Natl. Acad. Sci. U.S.A.* 90, 6444-6448), are also known to those skilled in the art and may be used in performing the present invention.

In one embodiment of the present invention, a binding protein, functional variant or immunoconjugate of the invention is prepared from a hybridoma that secretes the binding protein. See e.g. Köhler et al., 1975, *Nature* 256, 495-97.

In a preferred embodiment of the present invention, the binding protein, functional variant or immunoconjugate of the invention (i.e., binding molecules of the invention) is prepared recombinantly by optimizing and/or amplifying expression of the binding molecule of the invention in a host cell and isolating the binding molecule from said host cell. To this end, the host cells are transformed or transfected with DNA encoding a binding protein or a vector containing DNA encoding the binding molecule and cultured under appropriate conditions to produce the binding molecule of the invention, see for example, US 4,816,567.

Another aspect of the present invention relates to a method for producing a binding protein, functional variant or immunoconjugate of the invention (binding molecules of the invention) by preparing said binding molecule of the invention from a tissue, product or secretion of an animal, plant or fungus transgenic for a nucleic acid molecule or nucleic acid molecules encoding the molecule(s) of the invention. Preferably, a binding protein, functional variant or immunoconjugate of the invention is prepared from the tissue, product or secretion of a transgenic animal such as cow, sheep, rabbit, chicken or other mammalian or avian species, a transgenic plant such as corn, tobacco or other plant, or a transgenic fungus such as *Aspergillus*, *Pichia* or other fungal species.

#### Nucleic acid molecule compositions

In yet a further aspect, the invention provides compositions comprising at least one nucleic acid molecule as defined in the present invention. The compositions may comprise aqueous solutions such as aqueous solutions containing salts (e.g., NaCl or salts as described above), detergents (e. g., SDS) and/or other suitable components.

#### Pharmaceutical compositions

Furthermore, in an additional aspect the present invention relates to pharmaceutical compositions comprising at least one binding protein, at least one functional variant, at least one immunoconjugate or at least one composition according to the invention, or combinations thereof, and a pharmaceutically acceptable carrier, excipient, or stabilizer (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980)). In certain embodiments, pharmaceutical formulations are prepared to enhance the stability of the polypeptide or antibody during storage, e.g., in the form of lyophilized formulations or aqueous solutions.

In accordance with the present invention, acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include, e.g., buffers such as acetate, Tris, phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine;

preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; surfactant such as polysorbate; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

RNA viruses such as Chikungunya virus make use of their own RNA polymerase during virus replication. These RNA polymerases tend to be error-prone. This leads to the formation of so-called quasi-species during a viral infection. Each quasi-species has a unique RNA genome, which could result in differences in amino acid composition of viral proteins. If such mutations occur in structural viral proteins, the virus could potentially escape from the host's immune system due to a change in T or B cell epitopes.

Accordingly, in a preferred embodiment the pharmaceutical composition according to the invention comprises at least one additional binding protein, functional variant or immunoconjugate of the invention (binding molecules of the invention), i. e. the pharmaceutical composition can be a cocktail/mixture of binding molecules of the invention. The pharmaceutical composition may comprise at least two binding molecules according to the invention or at least one binding molecule according to the invention and at least one further anti-CHIKV virus binding molecule. The binding molecules in the pharmaceutical composition should preferably be capable of reacting with different, non-competing epitopes of the Chikungunya virus, thus, minimizing the chance of the occurrence of Chikungunya escape viruses. Epitopes may be present for example on the E1 or E2

glycoproteins of Chikungunya virus and may be different, non-overlapping epitopes. The binding molecules should preferably be of high affinity and should preferably have a broad specificity. More preferably, they neutralize as many strains of Chikungunya virus as possible. Even more preferably, they also exhibit neutralizing activity towards other genotypes of the genus *alphavirus* or even with other viruses of the *Togaviridae* family, while exhibiting no cross-reactivity with other viruses or normal cellular proteins. Preferably, one of the binding molecules is capable of neutralizing escape variants of the other binding molecule in the cocktail.

In a preferred embodiment, the present invention relates to a pharmaceutical composition comprising at least two Chikungunya virus neutralizing binding molecules, preferably binding molecules according to the invention, characterized in that the binding molecules are capable of reacting with different, non-competing epitopes of the Chikungunya virus. In an embodiment the pharmaceutical composition comprises a first Chikungunya virus neutralizing binding molecule which is capable of reacting with an epitope located in an antigenic site of one of the Chikungunya virus envelope proteins (E1 or E2) and a second Chikungunya virus neutralizing binding molecule which is capable of reacting with an epitope located in a different antigenic site of the same Chikungunya virus envelope protein (e.g. E2) or alternatively in an antigenic site of the other envelope glycoprotein (e.g. E1). Epitope mapping and identification of the antigenic site can be performed by methods well known in the art, such as using truncated and/or mutated forms of recombinant virus protein(s), identification of the nucleotide substitution(s) in viral escape mutants, if any (Gal-Tanamy et al., 2008, PNAS 105(49):19450-19455), or crystal structure analysis of the antibody-antigen complex (Lescar et al., 1995, J Biol Chem 270(30):18067-76). Epitope mapping could also be performed by competition ELISA as described in The Protein Protocols Handbook (1996), p595-600).

Furthermore, the pharmaceutical composition according to the invention may comprise at least one other therapeutic and/or prophylactic agent for the particular indication, e.g. infection being treated, or to prevent undesired effects.

Preferably, the additional therapeutic and/or prophylactic agent has an activity complementary to that of the binding molecule, functional variant, immunoconjugate or composition according to the invention and do not adversely affect each other. In particular, said further therapeutic and/or prophylactic agent can be an agent suitable for the treatment of a viral infection or suitable for the treatment of the symptoms associated to a viral infection, such as chloroquine phosphate, paracetamol, NSAID's (e.g., ibuprofen, naproxen, etc.) or corticosteroids. More preferably, said therapeutic agent is an anti-viral such as ribavirin, acyclovir or interferon-alpha, or a vaccine against an *alphavirus*, in particular against the Chikungunya virus.. Alternatively, said therapeutic agent is useful for treating secondary effects related to the binding protein, functional variant or immunoconjugate of the invention.

The active ingredients, e.g., binding molecules, functional variants or immunoconjugates of the present invention and other therapeutic agents, may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and polymethylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano- particles and nanocapsules) or in macroemulsions. Such techniques are disclosed for example in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

The pharmaceutical composition of the invention will be formulated according to the chosen route of administration. The pharmaceutical composition of the invention can be administered by any suitable route, including but not limited to oral, rectal, transdermal, ophthalmic, nasal, topical, vaginal or parenteral. In a particular embodiment, the pharmaceutical composition is formulated in order to be suitable for parenteral administration to a patient, e.g., a human being, preferably by intravenous, intramuscular, intraperitoneal or subcutaneous administration. Illustrative, non limiting examples of suitable formulations for parenteral administration are solutions, suspensions, emulsions, lyophilized compositions and the like.

Furthermore, sustained-release compositions are also contemplated. Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of  
5 sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (US 3,773,919), copolymers of L-glutamic acid and [ $\gamma$ ] ethyl- L-glutamate, non-degradable ethylene -vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT(TM) (injectable microspheres composed of lactic acid-glycolic  
10 acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid.

The administration of the pharmaceutical composition of the invention to the patient in need thereof can be carried out by conventional means and delivery will take place using the appropriate equipments, apparatus, and devices which are known by the skilled person in art. The term "patient", as used herein,  
15 refers to a mammalian patient, i.e. any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the patient in need of treatment is a human patient.

The dosage and schedule of administration of the pharmaceutical  
20 composition of the invention will vary according to the particular formulation, the mode of administration, and the particular situs and infection being treated. Other factors like age, body weight, sex, diet, rate of excretion, condition of the subject, drug combinations, reaction sensitivities and severity of the infection shall be taken into account. Generally, a therapeutically effective amount of a binding  
25 molecule, functional variant or immunoconjugate of the invention is administered to a patient. The term "therapeutically effective amount" refers to an amount of drug effective to "prevent or treat" a disease or disorder in a patient, as defined below.

In particular embodiments, the amount of molecule administered will  
30 typically be in the range of about 0.01 mg/kg to about 50 mg/kg of patient body weight. Depending on the type and severity of the infection, about 0.01 mg/kg to

about 50 mg/kg body weight of a molecule of the invention can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. The progress of this therapy can be readily monitored by conventional methods and assays and  
5 based on criteria known to the physician or other persons of skill in the art.

#### Therapeutic uses

In a further aspect, the binding proteins, functional variants, immunoconjugates, compositions, or pharmaceutical compositions of the  
10 invention can be used as a medicament. In particular, the binding proteins, functional variants, immunoconjugates, compositions, or pharmaceutical compositions of the invention may be used in the prevention or treatment of an arbovirus infection. Preferably this arbovirus is a virus of the Togaviridae family, more preferably this virus is an *alphavirus*. The *alphavirus* can be a virus from any of  
15 the known genotypes, but is preferably a Chikungunya virus.

In accordance with the present invention, the term "prevention or treatment", when used herein, refers to both therapeutic treatment and prophylactic or preventative measures, wherein the patient in need is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need  
20 of prevention or treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented. The patient in need of prevention or treatment is a mammalian patient, i.e. any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits,  
25 etc. Preferably, the patient in need of treatment is a human patient.

In a preferred embodiment, a binding protein, functional variant or immunoconjugate of the invention (binding molecules of the invention) with an optimized ADCC activity is used in the prevention or treatment of an arbovirus infection. Preferably this arbovirus is a virus of the Togaviridae family, more  
30 preferably an *alphavirus*. The *alphavirus* can be a virus from any of the known genotypes, but is preferably a Chikungunya virus. In a further preferred

embodiment, said binding molecule of the invention is an antibody with a glycosylation profile providing it with enhanced cell-mediated effector functions. Preferably, said antibody of the invention has been obtained by expression in a cell line producing naturally unfucosylated antibodies, such as for example, an  
5 avian cell, preferably a duck cell. Preferably this binding molecule has been produced in avian embryonic derived stem cell line EBx® marketed by Vivalis (Nantes, France), as described for example in WO 2008/142124.

In a further aspect of the invention, binding proteins, functional variants, immunoconjugates, compositions, or pharmaceutical compositions of the  
10 invention can be used with other drugs to provide a combination therapy for the treatment of an arbovirus infection, as above described. The other drugs may form part of the same composition, or be provided as a separate pharmaceutical composition for administration at the same time or at a different time.

The identity of the other drug is not particularly limited. In a particular  
15 embodiment, said additional drug is administered simultaneously or sequentially to the binding proteins, functional variants, immunoconjugates, compositions, or pharmaceutical compositions of the invention, spaced out in time, in any order, i.e. first the molecules or compositions of the invention, then the additional drug can be administered, or first the additional drug and then the molecules or  
20 compositions of the invention. In another alternative embodiment the molecules or compositions of the invention and an additional drug are simultaneously administered.

Preferably, said combination therapy comprises at least two Chikungunya virus neutralizing binding molecules. More preferably said binding molecules are  
25 binding proteins, functional variants or immunoconjugates of the invention (binding molecules of the invention), characterized in that the binding molecules are capable of reacting with different, non-competing epitopes of the Chikungunya virus. The chance of the occurrence of Chikungunya escape viruses is thereby minimized. As a consequence thereof, the binding molecules of the  
30 invention preferably are capable of reacting with different, non-overlapping, non-

competing epitopes of the Chikungunya virus, such as epitopes on the virus glycoproteins E1 and E2.

Alternatively, the additional drug is a drug useful in the prevention and/or treatment of a viral infection or a drug useful for reducing undesired secondary effects of the molecules or compositions of the invention. For instance, the binding molecules, functional variants, immunoconjugates or pharmaceutical compositions of the invention can be co-administered with antiviral molecules such as ribavirin or interferon- $\alpha$  or with a vaccine against an *alphavirus*, in particular against the Chikungunya virus. The additional drug may also be administered before or after administration of the molecules or compositions of the invention, in any desired order.

#### Diagnostic use

The invention further pertains to a method of *in vitro* detecting a Chikungunya virus in an isolated sample, wherein the method comprises the steps of a) contacting a sample with a diagnostically effective amount of a binding protein, a functional variant or an immunoconjugate according to the invention (i.e., binding molecules of the invention), and b) determining whether the binding molecule of the invention specifically binds to a molecule of the sample. The sample may be a biological sample including, but not limited to blood, serum, tissue or other biological material from (potentially) infected subjects. The (potentially) infected patients may be human patients, but also animals that are suspected as carriers of Chikungunya virus might be tested for the presence of Chikungunya virus using the binding molecules of the invention. The sample may first be manipulated to make it more suitable for the method of detection. Manipulation means inter alia treating the sample suspected to contain and/or containing Chikungunya virus in such a way that the Chikungunya virus will disintegrate into antigenic components such as proteins, (poly) peptides or other antigenic fragments. Preferably, the binding molecules of the invention are contacted with the sample under conditions which allow the formation of an immunological complex between the binding molecules and Chikungunya virus or

antigenic components thereof that may be present in the sample. The formation of an immunological complex, if any, indicating the presence of Chikungunya virus in the sample, is then detected and measured by suitable means. Such methods include, inter alia, homogeneous and heterogeneous binding immunoassays, such as radioimmunoassays (RIA), enzyme-linked immunoassay (ELISA), immunofluorescence, immunohistochemistry, flow cytometry (e.g. FACS), surface plasmon resonance (e.g. BIACORE) and Western blot analyses. Preferably, the molecule of the invention is an immunoconjugate.

In addition, immunoconjugates of the invention may be used for example in the *in vivo* detection of the Chikungunya virus infection by the use of imaging and nuclear medicine techniques. Various nuclear medicine techniques are well known in the art and have been widely used for example to image foci of infection and inflammation (Corstens et al., 1993, Seminars in Nuclear Medicine 23(2), 148-164).

#### Screening uses

Furthermore, the binding protein, functional variant or immunoconjugate of the invention (binding molecules of the invention) can be used to identify epitopes of Chikungunya virus proteins such as the E1 or E2 glycoproteins. The epitopes can be linear, but also structural and/or conformational. In one embodiment, binding of the binding molecules of the invention to a series of overlapping peptides, such as 15-mer peptides, of a protein from Chikungunya virus such as the Chikungunya virus E1 or E2 glycoproteins can be analyzed by means of PEPSCAN analysis (see inter alia WO 84/03564, WO 93/09872, Slootstra et al., 1996, Mol Divers. 1(2), 87-96).

The binding of said binding molecules to each peptide can be tested in a PEPSCAN-based enzyme-linked immuno assay (ELISA). In another embodiment, a random peptide library comprising peptides from Chikungunya virus proteins can be screened for peptides capable of binding to the binding molecules of the invention. In the above assays the use of Chikungunya virus neutralizing binding molecules may identify one or more neutralizing epitopes. The peptides/epitopes found can be used as vaccines and for the diagnosis of Chikungunya infection.

Accordingly, in a further aspect, the invention provides a method of screening a binding molecule for specific binding to a different, preferably non-overlapping epitope of Chikungunya virus as the epitope bound by a binding molecule of the invention, wherein the method comprises the steps of a) contacting a binding molecule to be screened, a binding molecule of the invention and Chikungunya virus or a fragment thereof (such as for instance the Chikungunya virus E1 or E2 glycoproteins), b) measure if the binding molecule to be screened is capable of competing for specifically binding to the Chikungunya virus or fragment thereof with the binding molecule of the invention. If no competition is measured the binding molecules to be screened bind to a different epitope. In a specific embodiment of the above screening method, human binding molecules may be screened to identify human binding molecules capable of binding a different epitope than the epitope recognized by the binding molecule comprising the CDRH3 comprising the amino acid sequence of SEQ ID NO: 14 or 34. Preferably, the epitopes are non-overlapping or non-competing. It is clear to the skilled person that the above screening method can also be used to identify binding molecules capable of binding to the same epitope. In a further step it may be determined if the screened binding molecules that are not capable of competing for specifically binding to the Chikungunya virus or fragment thereof have neutralizing activity. It may also be determined if the screened binding molecules that are capable of competing for specifically binding to the Chikungunya virus or fragment thereof have neutralizing activity. Neutralizing anti-CHIKV virus binding molecules found in the screening method are another part of the present invention.

In the screening method "specifically binding to the same epitope" also contemplates specific binding to substantially or essentially the same epitope as the epitope bound by the binding molecules of the invention. The capacity to block, or compete with, the binding of the binding molecules of the invention to Chikungunya virus typically indicates that a binding molecule to be screened binds to an epitope or binding site on the Chikungunya virus that structurally overlaps with the binding site on the Chikungunya virus that is immunospecifically

recognized by the binding molecules of the invention. Alternatively, this can indicate that a binding molecule to be screened binds to an epitope or binding site which is sufficiently proximal to the binding site immunospecifically recognized by the binding molecules of the invention to sterically or otherwise inhibit binding of the binding molecules of the invention to Chikungunya virus or a fragment thereof.

In general, competitive inhibition is measured by means of an assay, wherein an antigen composition, i. e. a composition comprising Chikungunya virus or fragments thereof (such as E1 or E2 glycoproteins), is admixed with reference binding molecules and binding molecules to be screened. In an embodiment the reference binding molecule may be one of the human binding molecules of the invention and the binding molecule to be screened may be another human binding molecule of the invention. In another embodiment the reference binding molecule may be the binding molecule comprising the CDRH3 comprising the amino acid sequence of SEQ ID NO: 14 or 34 and the binding molecule to be screened may be one of the human binding molecules of the invention. In yet another embodiment the reference binding molecule may be one of the human binding molecule of the invention and the binding molecule to be screened may be the binding molecule comprising the CDRH3 comprising the amino acid sequence of SEQ ID NO: 14 or 34.

Usually, the binding molecules to be screened are present in excess. Protocols based upon ELISAs are suitable for use in such simple competition studies. In certain embodiments, one may pre-mix the reference binding molecules with varying amounts of the binding molecules to be screened (e. g., 1:10, 1: 20, 1: 30, 1: 40, 1: 50, 1: 60, 1: 70, 1: 80, 1: 90 or 1: 100) for a period of time prior to applying to the antigen composition. In other embodiments, the reference binding molecules and varying amounts of binding molecules to be screened can simply be admixed during exposure to the antigen composition. In any event, by using species or isotype secondary antibodies one will be able to detect only the bound reference binding molecules, the binding of which will be reduced by the

presence of a binding molecule to be screened that recognizes substantially the same epitope.

Binding molecules identified by these competition assays ("competitive binding molecules") include, but are not limited to, antibodies, antibody fragments and other binding agents that bind to an epitope or binding site bound by the reference binding molecule as well as antibodies, antibody fragments and other binding agents that bind to an epitope or binding site sufficiently proximal to an epitope bound by the reference binding molecule for competitive binding between the binding molecules to be screened and the reference binding molecule to occur. Preferably, competitive binding molecules of the invention will, when present in excess, inhibit specific binding of a reference binding molecule to a selected target species by at least 10%, preferably by at least 25%, more preferably by at least 50%, and most preferably by at least 75%-90% or even greater. The identification of one or more competitive binding molecules that bind to about, substantially, essentially or at the same epitope as the binding molecules of the invention is a straightforward technical matter. As the identification of competitive binding molecules is determined in comparison to a reference binding molecule, it will be understood that actually determining the epitope to which the reference binding molecule and the competitive binding molecule bind is not in any way required in order to identify a competitive binding molecule that binds to the same or substantially the same epitope as the reference binding molecule. Alternatively, binding molecules binding to different non-competing epitopes identified by these competition assays may also include, but are not limited to, antibodies, antibody fragments and other binding agents.

25

#### Kits

In a further aspect, the invention relates to kits comprising at least one binding protein according to the invention, at least one functional variant thereof according to the invention, at least one immunoconjugate according to the invention, at least one nucleic acid molecule according to the invention, at least one composition according to the invention, at least one pharmaceutical

30

composition according to the invention, at least one vector according to the invention, at least one host according to the invention or a combination thereof are also a part of the present invention. Optionally, the above described components of the kits of the invention are packed in suitable containers and  
5 labeled for diagnosis, prevention and/or treatment of the indicated conditions. The above-mentioned components may be stored in unit or multi-dose containers, for example, sealed ampoules, vials, bottles, syringes, and test tubes, as an aqueous, preferably sterile, solution or as a lyophilized, preferably sterile, formulation for reconstitution. The containers may be formed from a variety of  
10 materials such as glass or plastic and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The kit may further comprise more containers comprising a pharmaceutically acceptable buffer, such as phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other  
15 materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes and/or culture medium for one or more of the suitable hosts. Associated with the kits can be instructions customarily included in commercial packages of therapeutic, prophylactic or diagnostic products, that contain information about for example the indications, usage, dosage,  
20 manufacture, administration, contraindications and/or warnings concerning the use of such therapeutic, prophylactic or diagnostic products.

## EXAMPLES

The following examples, including the experiments conducted and the results  
25 achieved, further illustrate the invention. They should not, however, be interpreted as a limitation of the scope of the invention.

### EXAMPLE 1: Isolation of human mAbs neutralizing Chikungunya virus infection

The purpose of this assay was to isolate human monoclonal antibodies (mAbs)  
30 capable of neutralizing Chikungunya virus from an infected individual who has naturally developed antibody protection against the virus infection. In this study,

two different strains of Chikungunya Viruses (CHIKV), A226 and A226V strains, were used (both kindly provided by Dr Raymond Lin, National University Hospital, Singapore). The viral strains sequences have not yet been fully characterized except for the viral envelope protein 1 (E1) encoding sequences. One amino acid  
 5 difference has been identified between the E1 protein sequences of these viral isolates at the 226 amino acid position, either an alanine (A) in the A226 strain or a valine (V) in the A226V strain.

#### Isolation of B cells from an infected donor

10 50 ml of blood were obtained in ACD-A tubes (BD Biosciences, SG) from an individual (42 years-old woman) who had been infected by Chikungunya virus (CHIKV) six weeks before and had neutralizing titers of anti-Chikungunya antibodies in the plasma.

Peripheral blood mononuclear cells (PBMC), including monocytes, B and T  
 15 lymphocytes, were isolated by classical Ficoll™ (GE healthcare, SG) extraction. PBMC were counted and the percentage of B lymphocyte cells was determined by flow cytometry analysis (FacsCalibur, Beckton-Dickinson) after co-staining with anti-human CD45 (Mouse anti-Human CD45 Alexa Fluor 488-labeled, Invitrogen) and anti-human CD19 (Mouse anti-Human CD19 R-Phycoerythrin-labeled,  
 20 Invitrogen).

#### Immortalization of the B cells

The B cells were transformed with a suitable Epstein-Barr virus (EBV) strain and were immediately activated via the CD40 pathway.

25

The total PBMC was seeded into the wells of 96 well-plates (BD Falcon) in order to have three different densities of B cell populations: 10 B cells per well (6x 96-well plates), 30 B cells per well (7x 96-well plates) and 500 B cells per well (7x 96-well plates). The B cells were immediately activated via the CD40 pathway, either by  
 30 adding to the cell medium an antibody anti-Human CD40 (Mouse anti-CD40; Dendritics, France) at 0.5 µg/ml, or by adding to the wells 10000 lyophilized L cells

(fibroblastic cells; Dendritics, France) constitutively expressing at the cell surface the CD40-ligand. The B cells were also immediately immortalized by EBV infection by adding 100 µl/well of infectious cell supernatant obtained from B95.8 cells. B95.8 is a human lymphoblastoïde cell line chronically infected by the EBV and actively producing some infectious viral particles (ATCC Number VR-1492).

The immortalized B cells were grown in Dulbecco's modified Eagle's growth medium-F12 (DMEM-F12, Invitrogen) supplemented with 10% Fetal Calf Serum (FCS, Invitrogen), 4 mM L-Glutamine (Gibco-Invitrogen), 100 U/ml of penicillin (Gibco-Invitrogen), 100 µg/ml of streptomycin (Gibco-Invitrogen) and 2% of ADCM medium (Dentritics, France).

#### Characterization of the immortalized B cell populations by binding and neutralization tests

A couple of weeks after the B cells immortalization, the polyclonal B cell populations obtained were analyzed for their capacity to secrete antibodies specific for CHIKV by performing a binding test on fixed CHIKV-infected HEK 293T cells (ATCC Number CRL-N268).

Firstly, 48 h prior to undertaking the binding test on the immortalized B cell supernatants, 20000 HEK 293T cells were seeded per well of 96-well plates and 24 hours before the test, the cells were infected with A226V CHIKV strain at a Multiplicity Of Infection (MOI, or number of infectious viral particles per cell) of 0.1. The day of the test, the cells were extensively washed with phosphate buffered saline (PBS), permeabilized and fixed with an ethanol/acetone (70:30 v/v) fixation solution. The immortalized B cell supernatants were then analyzed for the presence of CHIKV-specific antibodies. For each B cells supernatant's containing well, 50 µl of the supernatant were applied to one well with CHIKV infected HEK 293T cells, and 50 µl were applied to a well with uninfected HEK 293T cells, as a negative control. The wells were incubated for 1 h at 37 °C and the binding of anti-CHIKV antibodies was then detected by immunofluorescent assay (IFA), using a cocktail of secondary antibodies consisting of anti-Human IgG (Alexa Fluor 488 goat anti-

human IgG (H+L), Invitrogen) at 2µg/ml, anti-human IgM (Alexa Fluor 488 goat anti-human IgM (µ), Invitrogen) at 2µg/ml, and anti-human IgA (FITC-goat anti-human IgA, Invitrogen) 1/1000. Fluorescence was analyzed under a fluorescent microscope (x10 magnification, NIKON ECLIPSE TS 100). 215 polyclonal B cell populations were identified as being specific for A226V CHIKV. Results of the binding tests are not shown.

The polyclonal B cell populations identified as positive for anti-CHIKV antibodies secretion were subjected to a neutralization test with the purpose to identify those secreting antibodies neutralizing CHIKV infection *in vitro*. 4000 plaque forming units (PFU) of A226V CHIKV were incubated for 1h at 37°C with 50 µl of CHIKV-specific polyclonal B cells supernatants. As positive control, 1/40 of anti-CHIKV human plasma (isolated from the donor patient) was incubated with the same amount of PFU. 1/40 of human serum AB (Gemini Bio-products) from normal healthy male donors was used as negative control. The mixtures were then added for 1.5 hours to 40,000 HEK 293T cells, which were seeded within 96-wells plates 24h before. 24 h after the infection, cells were extensively washed, fixed as described above, incubated for 1 h with anti-CHIKV plasma 1/200 as a primary antibody, and then incubated for 1 h with the fluorescent labeled-anti-human Ig as described above. Fluorescence was analyzed under a fluorescent microscope (x10 magnification, NIKON ECLIPSE TS 100). 91 of the 215 polyclonal B cell populations selected for being specific for A226V CHIKV displayed neutralizing activity against A226V.

#### Cloning of the B cells secreting CHIKV neutralizing antibodies

The B cells secreting antibodies neutralizing CHIKV infectivity were cloned by limiting dilutions within 96-well plates (1/5 serial dilutions) and, after expansion (for 15 days), the wells containing the highest cell dilution and remaining positive for CHIKV neutralization were further amplified to a culture volume of 0.5 L. During amplification, the cells were regularly analyzed for their specificity for CHIKV, as well as for their capacity to secrete CHIKV neutralizing antibodies, by binding and neutralization tests, as described above.

### Monoclonal antibodies purification and Ig subclass determination

After amplification, the monoclonal antibodies were purified from the cell culture supernatant.

The monoclonal B cells were cultivated for 7 days in DMEM F12 medium supplemented as described above but without FCS, in order to avoid any presence of undesired bovine immunoglobulins. After 7 days, the cell culture supernatant was harvested, filtered on 0.45 µm membranes (Corning) and 500 ml of culture supernatant were incubated overnight with 2 ml of protein G-conjugated agarose beads (Protein G Agarose, Millipore). The protein G-beads were then extensively washed with PBS buffer and subsequently the bound antibodies were eluted with a solution of Glycine 1M (Sigma-Aldrich), pH 2.8. After elution, the acidity of the elution buffer was immediately neutralized by the volume to volume addition of Tris-base buffer pH 8 (1st Base company). The purified antibodies were then dialyzed overnight in PBS buffer, using a Slide-A-Lyzer Dialysis Cassette, 10.000 MWCO (Thermoscientific).

Two different monoclonal antibodies capable of efficiently preventing Chikungunya virus infection *in vitro* have been isolated. They have been named **8B10F8** and **5F10F175E2**. The isotype class was determined by Flow Cytometry analysis (FACSCalibur, Beckton-Dickinson) using the Multiplex Bead Assay for Human Ig Isotyping (SouthernBiotech). Both monoclonal antibodies belong to the IgG1 subclass of immunoglobulins (results not shown).

### IC50 determination by Plaque Reduction Assay

The inhibition concentration 50 (IC50) is the antibody concentration leading to the neutralization of half of the infective viral particles. The IC50 of the two isolated human monoclonal antibodies was determined.

Serial dilutions of purified monoclonal antibodies (1/10 dilutions from 100 µg/ml to 0.1 ng/ml) were put in contact with 200 PFU (Plaque Forming Unit, in other words infectious viral particles) of CHIKV for 1 h at 37 °C and subsequently, said antibody/virus mixtures were applied onto Vero cells (ATCC CCL-81), which were seeded 24 h before within 6-well plates (BD Falcon), for 1 h at 37 °C to allow the

virus to bind to the cells. After the infection step, the culture medium (DMEM (Gibco-Invitrogen) supplemented with 10% FCS) was removed and replaced by 0.25% agarose-medium (Invitrogen), this was done in order to prevent secondary infections by spreading of neo-synthesized viral particles so that the PFUs eventually quantified directly correlate with the amount of viral particles initially used for the cell infection. 48 h after infection, the agarose-medium was removed and the viable cells were stained with a cristal violet solution (Sigma-Aldrich) in order to visualize the plaques resulting from each PFU. The Plaque Reduction Assay was performed with both CHIKV strains used in the study, namely the A226 and the A226V CHIKV strains. Within one same experiment, each condition was performed in duplicate. The neutralization potency of the **5F10F175E2** and **8B10F8** monoclonal antibodies, as single agents and in combination (at the same final antibody concentration as that used in the single antibodies' determination) was determined to investigate any synergic effect of the antibody association. The experiment was performed in parallel with the irrelevant human IgG1 antibody HA4 as a negative control for neutralization. The percentage of neutralization associated to each condition was determined by the number of PFU obtained, compared to that obtain with the irrelevant human IgG1. Both antibodies (**8B10F8** and **5F10F175E2**) were found to have an IC<sub>50</sub> of about 100 ng/ml as single agents. No significant synergistic effect on neutralization was observed when both antibodies were used in combination. Results are not shown.

#### **EXAMPLE 2: CHIKV neutralizing antibodies sequencing**

The variable domains of the heavy and light chains of the **5F10F175E2** and **8B10F8** monoclonal antibodies were sequenced by first extracting the total RNA of the corresponding monoclonal B cell populations. Total RNA was extracted using TRizol® reagent (Invitrogen) according to the provider's instructions. For each antibody, from the total RNAs, two independent reverse-transcriptions were performed using the reverse-transcriptase kit (Clontech), giving rise to 2 independent complementary DNAs (cDNAs). From each cDNA, a whole set of PCRs aiming to amplify all the possible heavy and light chains were performed

using the AdvantageR 2PCR kit (Clontech). The light chains were amplified using, as a forward primer, the 5'-PCR Primer II A provided with the PCR kit (Clontech), and as reverse primers, the whole set of reverse primers specific for the different constant regions of the different possible light chains. The Heavy chains were amplified using, subtype-specific (IgG, IgM, IgA) reverse primers in conjunction with forward primers corresponding to the signal peptide regions.

Depending on the obtained results, the same amplified fragments were mixed together and sequenced. The nucleotide sequences of the heavy and light chains of the **5F10F175E2** antibody are shown in Figures 2A and 2B, respectively; and the nucleotide sequences of the heavy and light chains of the **8B10F8** antibody are shown in Figures 3A and 3B, respectively.

### **EXAMPLE 3: Expression of recombinant mAbs**

The nucleotide sequences encoding the variable domain of the heavy chain (VH) and the whole light chain (L), were molecularly cloned into pPMhlgG1 expression plasmid (a gift kindly provided by Dr. John Wu from ProMab), using the T4 DNA ligase (New England Biolabs) according to the provider's instructions and the ligation product was transformed into competent bacteria (Library Efficiency DHF- $\alpha$ , Invitrogen).. The two plasmids encoding the light and the heavy chain of each antibody were co-transfected (0.5 $\mu$ g/1x10<sup>6</sup> cells for each plasmid) using lipofectamin 293™ (Invitrogen) into human HEK293TPM1 cells.

Silent point mutations were introduced in the recombinant sequences to add or remove restriction enzyme sites to facilitate the cloning into the expression vector. The heavy chain and light chain cDNA sequence of **rec5F10F175E2** is shown in SEQ ID NO: 41 and 42, respectively. The heavy chain and light chain cDNA sequence of **rec8B10F8** is shown in SEQ ID NO: 43 and 44, respectively. The protein sequences are identical between the "isolated" (**8B10F8** and **5F10F175E2**) and the "recombinant" (**rec8B10F8** and **rec5F10F175E2**) antibodies.

pPMhlgG1 is a mammalian cell expression vector. It contains a sequence encoding the signal peptide of mouse Ig kappa chain, followed by the sequence of the human IgG1 constant region (CH1-3) (SEQ ID NO: 47), cloned between the Nhe I and Not I sites. pPMhlgG1 allows the construction of the full length human IgG1 heavy chain by cloning the variable region (VH) between the Sal I and Nhe I sites. To construct the expression vector for the light chain, the entire light chain sequence is cloned between Sal I and Not I sites. Using this cloning strategy, the secretion of both heavy and light chains is directed by the mouse Ig kappa chain signal peptide (see Figure 6).

The recombinant antibodies (**rec8B10F8** and **rec5F10F175E2**) were purified from the transfected cells following the above described procedure.

#### **EXAMPLE 4: *in vitro* characterization of recombinant mAbs**

##### **Determination of the recombinant mAbs *in vitro* neutralizing properties**

The capacity of the purified recombinant antibodies (**rec8B10F8** and **rec5F10F175E2**) to neutralize Chikungunya virus infection *in vitro* was analyzed by a neutralization test in 96-wells plates. For comparison purposes, the test was also performed on the originally isolated antibodies (**8B10F8** and **5F10F175E2**).

4000 plaque forming units (PFU) of Chikungunya virus (A226 and A226V strains, respectively) were incubated for 1h at 37°C with 1 µg of antibody (isolated or recombinant). In addition, 1/40 of anti-Chikungunya virus human plasma (isolated from the donor patient) in 100 µl of PBS was used as a positive control, and 1 µg of purified irrelevant human IgG1 HA4 (human IgG1 Ab specific for H5N1 Influenza virus, kindly provided by DSO National Laboratories, Singapore) as a negative control. The mixtures were then added for 1.5 hours to 40,000 HEK 293T cells, which were seeded within 96-wells plates 24h before. 24 hours after infection, cells were extensively washed and fixed as described above. The infection of Chikungunya virus was assessed by incubating the cells for 1 hour with 100 µl of PBS-1/200 anti-Chikungunya virus human plasma as a primary antibody, followed by an anti-human IgG mouse antibody conjugated to Alexa-488 at 2µg/ml (Alexa Fluor 488

goat anti-human IgG(H+L), Invitrogen) as a secondary antibody. Fluorescence was analyzed under a fluorescent microscope (x10 magnification, NIKON ECLIPSE TS 100).

In Figure 3 is shown the neutralizing activity of the two antibodies isolated from immortalized B cells (**8B10F8** and **5F10F175E2**) and also the neutralizing activity of the two corresponding recombinant antibodies (**rec8B10F8** and **rec5F10F175E2**). It can be observed that the neutralization potential of the recombinant antibodies is similar to the corresponding B cell-purified antibodies.

#### 10 Dose-response curves of the recombinant mAbs against A226 and A226V strains

Once having demonstrated that the recombinant antibodies have neutralization characteristics similar to those previously shown for the corresponding B cell-purified antibodies (Figure 3), the recombinant antibodies were titrated against the Chikungunya virus strains (A226 and A226V) as single agents and in combination to obtain neutralization dose-response curves and determine the IC<sub>50</sub>. The dose-response curves were obtained by Plaque Reduction Assay performed as described above, and the experiments were performed by triplicate. Results are shown in Figure 4.

The IC<sub>50</sub> value of the recombinant antibodies against the A226V strain was estimated between 20 and 80 ng/ml and the IC<sub>50</sub> of the recombinant antibodies against the A226 strain was estimated between 200 and 800 ng/ml.

#### Determination of the recombinant mAbs specificity

To determine recombinant mAb specificity, Western Blot analysis was performed on both CHIKV-infected cells and CHIKV particles. For uninfected cell lysate preparation, 12x10<sup>6</sup> Vero cells (ATTC CCL-81) were lysed with 1 ml of lysis buffer (PBS-1% Triton X-100, VWR) with Complete Protease Inhibitor Cocktail (Roche). For infected cell lysate preparation, 6x10<sup>6</sup> Vero cells were infected with CHIKV (A226 or A226V, respectively) at a MOI of 1 and 24 h after infection the cells were lysed with 1 ml of lysis buffer. For purified viral particles preparation, 40x10<sup>6</sup> Vero cells were infected with CHIKV (A226 or A226V, respectively) at a MOI of 0.1 and 2 days

after infection the supernatant was harvested, concentrated on Vivaspin-100 kDa columns (Vivaspin20, Sartorius Stedim) and, after having inactivated the virus for 1 h at 56 °C, the viral particles were purified by ultracentrifugation (24,000 rpm for 3h) performed on a OptiPrep™ (Sigma) cushion. 0.5 ml of purified A226 CHIKV viral particles and 1.25 ml of purified A226V CHIKV viral particles were finally obtained.

15 15 µl of cell lysates and 1.5 µl of purified viral particles were mixed with NuPAGE® loading buffer (Invitrogen), heated for 5 min at 94 °C and loaded on a NuPAGE® 4-12 % Bis-Tris Gel (Invitrogen). The electrophoresis was performed in NuPAGE® MES running buffer (Invitrogen). As molecular weight marker, 5 µl of SeeBlue® marker  
 10 (Invitrogen) were loaded in parallel. After migration, the samples were electrotransferred onto a hydrophobic polyvinylidene difluoride (PVDF) membrane (Hybond™-P, Amersham, GE Healthcare). The membrane was blocked for 1h with PBS-milk 5%(cow powdered milk)-Tween 0.05% (Sigma) and probed for 1h with primary antibody diluted in PBS-milk 0.5%-Tween 0.05% (Sigma) at the following  
 15 concentrations: 1/250 for the α-CHIKV plasma, 1µg/ml for CHIKV mAb 5F10F175E2, 20 µg/ml for CHIKV mAb 8B10F8 and 20 µg/ml for irrelevant human IgG1 HA4 (human IgG1 Ab specific for H5N1 Influenza virus, kindly provided by DSO National Laboratories, Singapore). After extensive washing in PBS-Tween (Sigma) 0.1%, the membrane was probed for 1 h with, as a secondary antibody, the Peroxidase-  
 20 conjugated AffiniPure goat Anti-Human IgG (Fc) (Jackson ImmunoResearch Laboratories) at 40 ng/ml in PBS-milk 0.5%-Tween 0.05%. After extensive washing, the peroxidase activity was detected using ECL substrate solutions (Amersham ECL™ Western Blotting Detection Reagents, GE Healthcare).

Western blot immunoassay results (Figure 5) show that both recombinant  
 25 antibodies bind to a Chikungunya virus envelope protein (E1 and/or E2). Chikungunya virus envelope glycoproteins E1 and E2 have a similar molecular weight, around 50kDa (Strauss and Strauss, 1994, Microbiol Rev 58, 491-562). Accordingly, these two proteins cannot be distinguished on a western blot analysis.

## CLAIMS

1. An isolated antibody or antibody fragment that specifically binds to Chikungunya virus, comprising:
  - 5       a. a heavy chain amino acid sequence that comprises the CDRH1 of SEQ ID NO: 10, the CDRH2 of SEQ ID NO: 12, and the CDRH3 of SEQ ID NO: 14, and a light chain amino acid sequence that comprises the CDRL1 of SEQ ID NO: 16, the CDRL2 of SEQ ID NO: 18, and the CDRL3 of SEQ ID NO: 20; or
  - 10       b. a heavy chain amino acid sequence that comprises the CDRH1 of SEQ ID NO: 30, the CDRH2 of SEQ ID NO: 32, and the CDRH3 of SEQ ID NO: 34, and a light chain amino acid sequence that comprises the CDRL1 of SEQ ID NO: 36, the CDRL2 of SEQ ID NO: 38, and the CDRL3 of SEQ ID NO: 40.
2. The isolated antibody or antibody fragment of claim 1 comprising the VH domain amino acid sequence of SEQ ID NO: 6 and the VL domain amino acid sequence of SEQ ID NO: 8.
- 15   3. The isolated antibody or antibody fragment of claim 1 comprising the VH domain amino acid sequence of SEQ ID NO: 26 and the VL domain amino acid sequence of SEQ ID NO: 28.
4. The isolated antibody or antibody fragment of any of claims 1 to 3 which is a humanized antibody or antibody fragment.
5. The isolated antibody or antibody fragment of any one of claims 1 to 3 wherein said antibody is fully human.
- 20   6. An immunoconjugate of the antibody or antibody fragment of any of claims 1 to 5 wherein said antibody or antibody fragment is coupled to at least one labeling and/or effector group.
7. An isolated nucleic acid molecule comprising a nucleic acid sequence encoding the antibody or antibody fragment of any of claims 1 to 5 or the immunoconjugate of claim 6.
- 25   8. A vector comprising the nucleic acid molecule of claim 7.
9. A host cell comprising the nucleic acid molecule of claim 7 or the vector of claim 8.
10. A method for producing the isolated antibody or antibody fragment of any of claims 1 to 5 or the immunoconjugate of claim 6, comprising the step of producing said antibody or antibody fragment or immunoconjugate in a host cell of claim 9 and optionally, isolating said antibody or antibody fragment or immunoconjugate.
- 30

11. A pharmaceutical composition comprising as an active agent at least one isolated antibody or antibody fragment of any of claims 1 to 5 and/or at least one immunoconjugate of claim 6, and a pharmaceutically acceptable carrier, diluent or adjuvant.
12. The pharmaceutical composition of claim 11, which comprises the isolated antibody or antibody fragment according to claim 1.a and the isolated antibody or antibody fragment according to claim 1.b.
13. The isolated antibody or antibody fragment of any of claims 1 to 5 and/or the immunoconjugate of claim 6 as a medicament.
14. The isolated antibody or antibody fragment of any of claims 1 to 5, the immunoconjugate of claim 6 and/or the pharmaceutical composition of claim 11 or 12 for use in the prevention or treatment of the infection of the Chikungunya virus.
15. Use of the isolated antibody or antibody fragment of any of claims 1 to 5 or the immunoconjugate of claim 6 for *in vitro* diagnostic or screening purposes.
16. A kit comprising at least one isolated antibody or antibody fragment of any of claims 1 to 5, at least one immunoconjugate of claim 6 and/or at least one pharmaceutical composition of claim 11 or 12.

## SEQUENCE LISTING

<110> HUMALYS  
SINGAPORE IMMUNOLOGY NETWORK (SiGN)

<120> BINDING MOLECULES AGAINST CHIKUNGUNYA VIRUS AND USES THEREOF

<130> BRV 28 - PRIO

<160> 53

<170> PatentIn version 3.3

<210> 1

<211> 839

<212> DNA

<213> Artificial

<220>

<223> Heavy chain (DNA) of Purified 5F10F175E2

<400> 1

```

atggactgga cctggagcgt cctcttcttg gtggcagcag caacagggtgc cactcccag      60
gtgcaactgg tgcaatctgg gtcggagttg aagaagcctg ggcctcagt gaagggttcc      120
tgcaaggcct ctggatacac ctcactcgc tatgtatga cttgggtgcg acaggcccct      180
ggacaagggc ttgagtggat gggatggatc aacacctaca ctgggaaccc aacgtatgtc      240
cagggcttca caggacggtt tgtcttctcg ttggacacct ctgtcagcac ggcgtttctg      300
cacatcacca gcctaaaggc tgaggacact gccgtgtatt tctgtgcgag agaagggggc      360
gctcgggggt ttgactactg gggccaggga accctggtca ccgtctctc agcctccacc      420
aagggcccat cggctcttcc cctggcacc tctccaaga gcacctctgg gggcacagcg      480
gccctgggct gcctggtcaa ggactacttc cccgaaccgg tgacggtgtc gtggaactca      540
ggcgccctga ccagcggcgt gcacaccttc ccggctgtcc tacagtcctc aggactctac      600
tccctcagca gcgtggtgac cgtgccctcc agcagcttgg gcaccagac ctacatctgc      660
aacgtgaatc acaagcccag caacaccaag gtggacaaga gagttgagcc caaatcttgt      720
gacaaaactc acacatgcc accgtgccca gaacctgaac tctgggggg accgtcagtc      780
ttcctcttcc ccccaaaacc caaggacacc ctcatgatct cccggacccc tgaggtcac      839

```

<210> 2

<211> 260

<212> PRT

<213> Artificial

<220>

<223> Heavy chain (Protein )common to both Purified 5F10F175E2 and  
Recombinant 5F10F175E2

<400> 2

Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala  
 1 5 10 15  
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Leu Thr Arg Tyr  
 20 25 30  
 Ala Met Thr Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met  
 35 40 45  
 Gly Trp Ile Asn Thr Tyr Thr Gly Asn Pro Thr Tyr Val Gln Gly Phe  
 50 55 60  
 Thr Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Val Ser Thr Ala Phe  
 65 70 75 80  
 Leu His Ile Thr Ser Leu Lys Ala Glu Asp Thr Ala Val Tyr Phe Cys  
 85 90 95  
 Ala Arg Glu Gly Gly Ala Arg Gly Phe Asp Tyr Trp Gly Gln Gly Thr  
 100 105 110  
 Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
 115 120 125  
 Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
 130 135 140  
 Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
 145 150 155 160  
 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
 165 170 175  
 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
 180 185 190  
 Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
 195 200 205  
 Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr  
 210 215 220  
 His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
 225 230 235 240



His Asp Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Thr Leu  
 35 40 45

Leu Ile Tyr Val Asn Ser Asn Arg Pro Ser Gly Val Pro Asp Arg Phe  
 50 55 60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu  
 65 70 75 80

Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Asn  
 85 90 95

Leu Ser Gly Ser Ala Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu  
 100 105 110

Gly Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser  
 115 120 125

Glu Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp  
 130 135 140

Phe Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro  
 145 150 155 160

Val Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn  
 165 170 175

Lys Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys  
 180 185 190

Ser His Lys Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val  
 195 200 205

Glu Lys Thr Val Ala Pro Thr Glu Cys Ser  
 210 215

<210> 5  
 <211> 354  
 <212> DNA  
 <213> Artificial

<220>  
 <223> VH domain

<400> 5  
 caggtgcaac tgggtgcaatc tgggtcggag ttgaagaagc ctggggcctc agtgaaggtt 60  
 tcctgcaagg cctctggata caccctcact cgctatgcta tgacttgggt gcgacaggcc 120

```

cctggacaag ggcttgagtg gatgggatgg atcaacacct acactgggaa cccaacgtat      180
gtccaggggt tcacaggacg gtttgtcttc tcgttgagaca cctctgtcag cacggcggtt      240
ctgcacatca ccagcctaaa ggctgaggac actgccgtgt atttctgtgc gagagaaggg      300
ggcgctcggg gttttgacta ctggggccag ggaaccctgg tcaccgtctc ctca          354

```

```

<210> 6
<211> 118
<212> PRT
<213> Artificial

```

```

<220>
<223> VH domain

```

```

<400> 6

```

```

Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
1              5              10              15

```

```

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Leu Thr Arg Tyr
          20              25              30

```

```

Ala Met Thr Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
          35              40              45

```

```

Gly Trp Ile Asn Thr Tyr Thr Gly Asn Pro Thr Tyr Val Gln Gly Phe
          50              55              60

```

```

Thr Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Val Ser Thr Ala Phe
65              70              75              80

```

```

Leu His Ile Thr Ser Leu Lys Ala Glu Asp Thr Ala Val Tyr Phe Cys
          85              90              95

```

```

Ala Arg Glu Gly Gly Ala Arg Gly Phe Asp Tyr Trp Gly Gln Gly Thr
          100              105              110

```

```

Leu Val Thr Val Ser Ser
          115

```

```

<210> 7
<211> 336
<212> DNA
<213> Artificial

```

```

<220>
<223> VL domain

```

```

<400> 7
cagtctgtgc tgacgcagcc gccctcagtg totggggccc cagggcagag ggtcaccatc      60

```

tcttgcaactg ggagcagctc caacatcggg gcaagtcattg atgtacactg gtaccagcag 120  
 cttcctggaa cagccccac actcctcacc tatgttaaca gcaatcggcc ctcaggggtc 180  
 cctgaccgat tctctgggtc caagtctggc acctcagcct ccctggccat cactggggtc 240  
 caggctgagg atgaggctga ttattactgc cagtcctatg acagcaacct gagggttcg 300  
 gcggtgttcg gcggaggagc caagttgacc gtccta 336

<210> 8  
 <211> 112  
 <212> PRT  
 <213> Artificial

<220>  
 <223> VL domain

<400> 8

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln  
 1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Ser  
 20 25 30

His Asp Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Thr Leu  
 35 40 45

Leu Ile Tyr Val Asn Ser Asn Arg Pro Ser Gly Val Pro Asp Arg Phe  
 50 55 60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu  
 65 70 75 80

Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Asn  
 85 90 95

Leu Ser Gly Ser Ala Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu  
 100 105 110

<210> 9  
 <211> 15  
 <212> DNA  
 <213> Artificial

<220>  
 <223> CDRH1

<400> 9

cgctatgcta tgact

<210> 10  
 <211> 5  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRH1

<400> 10

Arg Tyr Ala Met Thr  
 1 5

<210> 11  
 <211> 51  
 <212> DNA  
 <213> Artificial

<220>  
 <223> CDRH2

<400> 11  
 tggatcaaca cctacactgg gaacccaacg tatgtccagg gcttcacagg a

51

<210> 12  
 <211> 17  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRH2

<400> 12

Trp Ile Asn Thr Tyr Thr Gly Asn Pro Thr Tyr Val Gln Gly Phe Thr  
 1 5 10 15

Gly

<210> 13  
 <211> 27  
 <212> DNA  
 <213> Artificial

<220>  
 <223> CDRH3

<400> 13  
 gaagggggcg ctcgggggtt tgactac

27

<210> 14  
 <211> 9  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRH3

<400> 14

Glu Gly Gly Ala Arg Gly Phe Asp Tyr  
 1 5

<210> 15  
 <211> 42  
 <212> DNA  
 <213> Artificial

<220>  
 <223> CDRL1

<400> 15  
 actgggagca gctccaacat cggggcaagt catgatgtac ac

42

<210> 16  
 <211> 14  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRL1

<400> 16

Thr Gly Ser Ser Ser Asn Ile Gly Ala Ser His Asp Val His  
 1 5 10

<210> 17  
 <211> 21  
 <212> DNA  
 <213> Artificial

<220>  
 <223> CDRL2

<400> 17  
 gttaacagca atcggccctc a

21

<210> 18  
 <211> 7  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRL2

<400> 18

Val Asn Ser Asn Arg Pro Ser  
 1 5

<210> 19  
 <211> 36  
 <212> DNA  
 <213> Artificial

<220>  
 <223> CDRL3

<400> 19  
 cagtcctatg acagcaacct gagtgggttcg gcggtg 36

<210> 20  
 <211> 12  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRL3

<400> 20

Gln Ser Tyr Asp Ser Asn Leu Ser Gly Ser Ala Val  
 1 5 10

<210> 21  
 <211> 854  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Heavy chain (DNA) of Purified 8B10F8

<400> 21  
 atgaagcayc tgtggttttt ccttctcctg gtggcagctc ccagatgggt cctgtcccag 60  
 gtgcagctgc aggagtcggg cccaggactg gtgaagcctt cggagaccct gtccctcacc 120  
 tgcactgtct ctggtgactc catcagtagt tactcctgga gctggatccg gcagccccca 180  
 ggggaaggac tggagtggat tggttatata cattacactg ggagcaccaa ctacaacccc 240  
 tccctcaaga gtcgactcac catatcagta gacgcgtcca agaaccagtt ctccctgaag 300  
 ctgagctctg tgaccgctgc ggacacggcc gtgtattact gtgcgagaga ttgggggggg 360  
 tatagcagca gctggacctc cggatatggac gtctggggcc aagggaccac ggtcacccgc 420  
 tcctcagcct ccaccaaggg cccatcggtc ttccccctgg caccctcctc caagagcacc 480  
 tctgggggca cagcggccct gggctgctg gtcaaggact acttccccga accggtgacg 540  
 gtgtcgtgga actcaggcgc cctgaccagc ggcgtgcaca ccttcccggc tgtcctacag 600  
 tcctcaggac tctactccct cagcagcgtg gtgaccgtgc cctccagcag cttggggcacc 660  
 cagacctaca tctgcaacgt gaatcacaag cccagcaaca ccaagggtgga caagagagtt 720  
 gagcccaaat cttgtgacaa aactcacaca tgcccaccgt gcccagcacc tgaactcctg 780

gggggaccgt cagtcttcct cttcccccca aaaccaagg acaccctcat gatytcccg 840

accctgagg tcac 854

<210> 22

<211> 266

<212> PRT

<213> Artificial

<220>

<223> Heavy chain (Protein) of common to both Purified 8B10F8 and Recombinant 8B10F8

<400> 22

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu  
1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Asp Ser Ile Ser Ser Tyr  
20 25 30

Ser Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile  
35 40 45

Gly Tyr Ile His Tyr Thr Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys  
50 55 60

Ser Arg Leu Thr Ile Ser Val Asp Ala Ser Lys Asn Gln Phe Ser Leu  
65 70 75 80

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala  
85 90 95

Arg Asp Trp Gly Gly Tyr Ser Ser Ser Trp Thr Tyr Gly Met Asp Val  
100 105 110

Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly  
115 120 125

Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly  
130 135 140

Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val  
145 150 155 160

Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe  
165 170 175

Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val  
 180 185 190

Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val  
 195 200 205

Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys  
 210 215 220

Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu  
 225 230 235 240

Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr  
 245 250 255

Leu Met Ile Ser Arg Thr Pro Glu Val Thr  
 260 265

<210> 23  
 <211> 708  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Light chain (DNA) of Purified 8B10F8

<400> 23  
 atggacatga ggggtccccgc tcagctcctg gggctcctgc tgctctggct cccaggtacc 60  
 agatgtgaca tccagatgac ccagtcctca tctcctctgt ctgcatctgt aggagacaga 120  
 gtcaccatca cttgccgggc gagtcagggc attagcaatt ctttagcctg gtatcagcag 180  
 aaaccagga aagcccctaa gctcctgctc tatgtgtcat ccagattgga aagtggggtc 240  
 ccatccaggt tcagtggcag tggatctggg acggattaca ctctcaccat cagcagcctg 300  
 cagcctgaag attttgcaac ttattactgt caacagtatt atagtacccc gtacactttt 360  
 ggccagggga ccaagctgga gatcaaacga actgtggctg caccatctgt cttcatcttc 420  
 ccgccatctg atgagcagtt gaaatctgga actgcctctg ttgtgtgcct gctgaataac 480  
 ttctatccca gagaggccaa agtacagtgg aaggtggata acgccctcca atcgggtaac 540  
 tcccaggaga gtgtcacaga gcaggacagc aaggacagca cctacagcct cagcagcacc 600  
 ctgacgctga gcaaagcaga ctacgagaaa cacaaagtct acgcctgcga agtcacccat 660  
 cagggcctga gctcgcccggt cacaaagagc ttcaacaggg gagagtgt 708

<210> 24  
 <211> 214  
 <212> PRT

<213> Artificial

<220>

<223> Light chain (Protein) common to both Purified 8B10F8 and Recombinant 8B10F8

<400> 24

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Asn Ser  
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Leu  
35 40 45

Tyr Ala Ala Ser Arg Leu Glu Ser Gly Val Pro Ser Arg Phe Ser Gly  
50 55 60

Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro  
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Tyr Ser Thr Pro Tyr  
85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala  
100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly  
115 120 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala  
130 135 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln  
145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser  
165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr  
180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser  
195 200 205

Phe Asn Arg Gly Glu Cys

210

<210> 25  
 <211> 369  
 <212> DNA  
 <213> Artificial

<220>  
 <223> VH domain of Purified 8B10F8

<400> 25  
 caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcggagac cctgtccctc 60  
 acctgcactg tctctggtga ctccatcagt agttactcct ggagctggat ccggcagccc 120  
 ccagggaagg gactggagtg gattggttat atccattaca ctgggagcac caactacaac 180  
 ccctccctca agagtcgact caccatatca gtagacgcgt ccaagaacca gttctccctg 240  
 aagctgagct ctgtgaccgc tgcggacacg gccgtgtatt actgtgcgag agattggggg 300  
 gggatatagca gcagctggac ctacggtatg gacgtctggg gccaaaggac cacggtcacc 360  
 gtctcctca 369

<210> 26  
 <211> 123  
 <212> PRT  
 <213> Artificial

<220>  
 <223> VH domain common to both Purified 8B10F8 and Recombinant 8B10F8

<400> 26

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu  
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Asp Ser Ile Ser Ser Tyr  
 20 25 30

Ser Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile  
 35 40 45

Gly Tyr Ile His Tyr Thr Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys  
 50 55 60

Ser Arg Leu Thr Ile Ser Val Asp Ala Ser Lys Asn Gln Phe Ser Leu  
 65 70 75 80

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala  
 85 90 95

Arg Asp Trp Gly Gly Tyr Ser Ser Ser Trp Thr Tyr Gly Met Asp Val  
 100 105 110

Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 115 120

<210> 27  
 <211> 321  
 <212> DNA  
 <213> Artificial

<220>  
 <223> VL domain

<400> 27  
 gacatccaga tgacccagtc tccatcctcc ctgtctgcat ctgtaggaga cagagtcacc 60  
 atcacttgcc gggcgagtca gggcattagc aattctttag cctgggtatca gcagaaacca 120  
 gggaaagccc ctaagctcct gctctatgct gcatccagat tggaaagtgg ggtcccatcc 180  
 aggttcagtg gcagtggatc tgggacggat tacactotca ccatcagcag cctgcagcct 240  
 gaagattttg caacttatta ctgtcaacag tattatagta ccccgtaac ttttggccag 300  
 gggaccaagc tggagatcaa a 321

<210> 28  
 <211> 107  
 <212> PRT  
 <213> Artificial

<220>  
 <223> VL domain

<400> 28

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Asn Ser  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Leu  
 35 40 45

Tyr Ala Ala Ser Arg Leu Glu Ser Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60

Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro  
 65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Tyr Ser Thr Pro Tyr

85

90

95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys  
 100 105

<210> 29  
 <211> 15  
 <212> DNA  
 <213> Artificial

<220>  
 <223> CDRH1

<400> 29  
 agttactcct ggagc

15

<210> 30  
 <211> 5  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRH1

<400> 30

Ser Tyr Ser Trp Ser  
 1 5

<210> 31  
 <211> 48  
 <212> DNA  
 <213> Artificial

<220>  
 <223> CDRH2 of Purified 8B10F8

<400> 31  
 tatatccatt aactgggag caccaactac aaccctccc tcaagagt

48

<210> 32  
 <211> 16  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRH2 common to both Purified 8B10F8 and Recombinant 8B10F8

<400> 32

Tyr Ile His Tyr Thr Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser  
 1 5 10 15

<210> 33  
 <211> 45

<212> DNA  
<213> Artificial

<220>  
<223> CDRH3

<400> 33  
gattgggggg ggtatagcag cagctggacc tacggtatgg acgtc

45

<210> 34  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> CDRH3

<400> 34

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Asp | Trp | Gly | Gly | Tyr | Ser | Ser | Ser | Trp | Thr | Tyr | Gly | Met | Asp | Val |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     | 15  |     |

<210> 35  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> CDRL1

<400> 35  
cgggcgagtc agggcattag caattcttta gcc

33

<210> 36  
<211> 11  
<212> PRT  
<213> Artificial

<220>  
<223> CDRL1

<400> 36

|     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Arg | Ala | Ser | Gln | Gly | Ile | Ser | Asn | Ser | Leu | Ala |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |

<210> 37  
<211> 21  
<212> DNA  
<213> Artificial

<220>  
<223> CDRL2

<400> 37  
gctgcatcca gattggaaag t

21

<210> 38  
 <211> 7  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRL2

<400> 38

Ala Ala Ser Arg Leu Glu Ser  
 1 5

<210> 39  
 <211> 27  
 <212> DNA  
 <213> Artificial

<220>  
 <223> CDRL3

<400> 39  
 caacagtatt atagtacccc gtacact

27

<210> 40  
 <211> 9  
 <212> PRT  
 <213> Artificial

<220>  
 <223> CDRL3

<400> 40

Gln Gln Tyr Tyr Ser Thr Pro Tyr Thr  
 1 5

<210> 41  
 <211> 842  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Heavy chain (DNA) Recombinant 5F10F175E2

<400> 41  
 atggagacag acacactcct gctatgggta ctgctgctct gggttccagg gtcgactggc 60  
 caggtgcaac tgggtgcaatc tgggtcggag ttgaagaagc ctggggcctc agtgaaggtt 120  
 tcctgcaagg cctctggata caccctcact cgctatgcta tgacttgggt gcgacaggcc 180  
 cctggacaag ggcttgagtg gatgggatgg atcaacacct aactgggaa cccaacgtat 240  
 gtccagggct tcacaggacg gtttgtcttc tcgttgga cctctgtcag cacggcgttt 300  
 ctgcacatca ccagcctaaa ggctgaggac actgccgtgt atttctgtgc gagagaaggg 360

```

ggcgctcggg gttttgacta ctggggccag ggaaccctgg tcaccgtctc ctcagcctcc 420
accaagggcc catcggtctt cccgctagca ccctcctcca agagcacctc tgggggcaca 480
gcgggccttg gctgcctggg caaggactac ttccccgaac cggtgacggg gtcgtggaac 540
tcaggcgccc tgaccagcgg cgtgcacacc ttcccggctg tcctacagtc ctcaggactc 600
tactcctca gcagcgtggg gaccgtgccc tccagcagct tgggcaccca gacctacatc 660
tgcaacgtga atcacaagcc cagcaacacc aaggtggaca agagagttga gcccaaattc 720
tgtgacaaaa ctacacatg cccaccgtgc ccagcacctg aactcctggg gggaccgtca 780
gtcttctct tcccccaaa acccaaggac accctcatga tctccggac ccctgaggtc 840
ac 842

```

```

<210> 42
<211> 714
<212> DNA
<213> Artificial

```

```
<220>
```

```
<223> Light chain (DNA) of Recombinant 5F10F175E2
```

```

<400> 42
atggagacag acacactcct gctatgggta ctgctgctct gggttccagg gtcgactggc 60
cagtctgtgc tgacgcagcc gccctcagtg tctggggccc cagggcagag ggtcaccatc 120
tcctgcactg ggagcagctc caacatcggg gcaagtcatg atgtacactg gtaccagcag 180
cttcctggaa cagccccac actcctcatc tatgttaaca gcaatcggcc ctcaggggtc 240
cctgaccgat tctctggctc caagtctggc acctcagcct ccctggccat cactgggctc 300
caggctgagg atgaggctga ttattactgc cagtcctatg acagcaacct gagtggttcg 360
gcggtgttcg gcggagggac caagttgacc gtcctaggtc agccaaggc tgccccctcg 420
gtcactctgt tcccgccctc ctctgaggag cttcaagcca acaaggccac actggtgtgt 480
ctcataagtg acttctaccc gggagccgtg acagtggcct ggaaggcaga tagcagcccc 540
gtcaaggcgg gagtggagac caccacaccc tccaaacaaa gcaacaacaa gtacgcggcc 600
agcagctacc tgagcctgac gcctgagcag tggaagtccc aaaaagcta cagctgccag 660
gtcacgcatg aaggagcac cgtggagaag acagtggccc ctacagaatg ttca 714

```

```

<210> 43
<211> 857
<212> DNA
<213> Artificial

```

```
<220>
```

```
<223> Heavy chain (DNA) of Recombinant 8B10F8
```

<400> 43  
 atggagacag acacactcct gctatgggta ctgctgctct gggttccagg gtcgactggc 60  
 cagggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcgagac cctgtccctc 120  
 acctgcactg tctctgggtga ctccatcagt agttactcct ggagctggat ccggcagccc 180  
 ccagggaagg gactggagtg gattggttat atccattaca ctgggagcac caactacaac 240  
 ccctccctca agagccgact caccatatca gtagacgcgt ccaagaacca gttctccctg 300  
 aagctgagct ctgtgaccgc tgcggacacg gccgtgtatt actgtgagag agattggggg 360  
 gggatatagca gcagctggac ctacggtatg gacgtctggg gccaaaggac cacggtcacc 420  
 gtctctcag cctccaccaa gggcccatcg gtcttcccg tagcaccctc ctccaagagc 480  
 acctctgggg gcacagcggc cctgggctgc ctggtcaagg actacttccc cgaaccggtg 540  
 acggtgtcgt ggaactcagg cgccctgacc agcggcgtgc acaccttccc ggctgtccta 600  
 cagtcctcag gactctactc cctcagcagc gtggtgaccg tgccctccag cagcttgggc 660  
 acccagacct acatctgcaa cgtgaatcac aagcccagca acaccaaggt ggacaagaga 720  
 gttgagccca aatcttgtga caaaactcac acatgccac cgtgccagc acctgaactc 780  
 ctggggggac cgtcagtctt cctcttcccc caaaaccca aggacaccct catgatytcc 840  
 cggaccctg aggtcac 857

<210> 44  
 <211> 702  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Light chain (DNA) of Recombinant 8B10F8

<400> 44  
 atggagacag acacactcct gctatgggta ctgctgctct gggttccagg gtcgactggc 60  
 gacatccaga tgaccagtc tccatcctcc ctgtctgcat ctgtaggaga cagagtcacc 120  
 atcacttgcc gggcgagtca gggcattagc aattcttttag cctggtatca gcagaaacca 180  
 gggaaagccc ctaagctcct gctctatgct gcatccagat tggaaagtgg ggtcccatcc 240  
 aggttcagtg gcagtggatc tgggacggat tacactctca ccacagcag cctgcagcct 300  
 gaagattttg caacttatta ctgtcaacag tattatagta cccgtacac ttttggccag 360  
 gggaccaagc tggagatcaa acgaactgtg gctgcacat ctgtcttcat cttcccgcca 420  
 tctgatgagc agttgaaatc tggaactgcc tctgttgtgt gcctgctgaa taacttctat 480  
 ccagagagg ccaaagtaca gtggaagggtg gataacgcc tccaatcggg taactcccag 540  
 gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg 600

ctgagcaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc 660  
 ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gt 702

<210> 45  
 <211> 369  
 <212> DNA  
 <213> Artificial

<220>  
 <223> VH domain(DNA) of Recombinant 8B10F8

<400> 45  
 caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcggagac cctgtccctc 60  
 acctgcactg tctctggtga ctccatcagt agttactcct ggagctggat ccggcagccc 120  
 ccaggaaggg gactggagtg gattggttat atccattaca ctgggagcac caactacaac 180  
 ccctccctca agagccgact caccatatca gtagacgcgt ccaagaacca gttctccctg 240  
 aagctgagct ctgtgaccgc tgcggacacg gccgtgtatt actgtgcgag agattggggg 300  
 gggatatagca gcagctggac ctacggtatg gacgtctggg gccaaaggac cacggtcacc 360  
 gtctcctca 369

<210> 46  
 <211> 48  
 <212> DNA  
 <213> Artificial

<220>  
 <223> VH domain(DNA) of Recombinant 8B10F8

<400> 46  
 tatatccatt acactgggag caccaactac aaccctccc tcaagagc 48

<210> 47  
 <211> 81  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Sequence of Figure 6

<400> 47  
 atggagacag acacactcct gctatgggta ctgctgctgt gggttccagg gtcgactggc 60  
 ttgctagcac cctcctccaa g 81

<210> 48  
 <211> 20  
 <212> PRT  
 <213> Artificial

&lt;220&gt;

&lt;223&gt; Mouse signal peptide (Figure 6)

&lt;400&gt; 48

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Glu | Thr | Asp | Thr | Leu | Leu | Leu | Trp | Val | Leu | Leu | Leu | Trp | Val | Pro |
| 1   |     |     |     | 5   |     |     |     | 10  |     |     |     |     | 15  |     |     |

|     |     |     |     |
|-----|-----|-----|-----|
| Gly | Ser | Thr | Gly |
|     |     |     | 20  |

&lt;210&gt; 49

&lt;211&gt; 6

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Human IgG1 constant region (Figure 6)

&lt;400&gt; 49

|     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|
| Leu | Ala | Pro | Ser | Ser | Lys |
| 1   |     |     | 5   |     |     |

&lt;210&gt; 50

&lt;211&gt; 280

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Heavy chain variable domain (VH) and a partial region of the Constant Heavy 1 (CH1) domain of 5F10F175E2 (figure 2A)

&lt;400&gt; 50

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Asp | Trp | Thr | Trp | Ser | Val | Leu | Phe | Leu | Val | Ala | Ala | Ala | Thr | Gly |
| 1   |     |     |     | 5   |     |     |     | 10  |     |     |     |     |     | 15  |     |

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ala | His | Ser | Gln | Val | Gln | Leu | Val | Gln | Ser | Gly | Ser | Glu | Leu | Lys | Lys |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Pro | Gly | Ala | Ser | Val | Lys | Val | Ser | Cys | Lys | Ala | Ser | Gly | Tyr | Thr | Leu |
|     |     | 35  |     |     |     |     | 40  |     |     |     |     | 45  |     |     |     |

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Thr | Arg | Tyr | Ala | Met | Thr | Trp | Val | Arg | Gln | Ala | Pro | Gly | Gln | Gly | Leu |
|     | 50  |     |     |     |     | 55  |     |     |     |     | 60  |     |     |     |     |

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Trp | Met | Gly | Trp | Ile | Asn | Thr | Tyr | Thr | Gly | Asn | Pro | Thr | Tyr | Val |
| 65  |     |     |     |     | 70  |     |     |     |     | 75  |     |     |     | 80  |     |

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Gln | Gly | Phe | Thr | Gly | Arg | Phe | Val | Phe | Ser | Leu | Asp | Thr | Ser | Val | Ser |
|     |     |     |     | 85  |     |     |     |     | 90  |     |     |     |     | 95  |     |

Thr Ala Phe Leu His Ile Thr Ser Leu Lys Ala Glu Asp Thr Ala Val  
 100 105 110

Tyr Phe Cys Ala Arg Glu Gly Gly Ala Arg Gly Phe Asp Tyr Trp Gly  
 115 120 125

Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser  
 130 135 140

Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala  
 145 150 155 160

Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val  
 165 170 175

Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala  
 180 185 190

Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val  
 195 200 205

Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His  
 210 215 220

Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys  
 225 230 235 240

Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly  
 245 250 255

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met  
 260 265 270

Ile Ser Arg Thr Pro Glu Val Thr  
 275 280

<210> 51  
 <211> 237  
 <212> PRT  
 <213> Artificial

<220>  
 <223> sequence of whole light chain of 5F10F175E2

<400> 51

Met Ala Trp Ser Pro Leu Leu Leu Thr Leu Leu Ala His Cys Thr Gly  
 1 5 10 15

Ser Trp Ala Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala  
 20 25 30

Pro Gly Gln Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile  
 35 40 45

Gly Ala Ser His Asp Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala  
 50 55 60

Pro Thr Leu Leu Ile Tyr Val Asn Ser Asn Arg Pro Ser Gly Val Pro  
 65 70 75 80

Asp Arg Phe Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile  
 85 90 95

Thr Gly Leu Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr  
 100 105 110

Asp Ser Asn Leu Ser Gly Ser Ala Val Phe Gly Gly Gly Thr Lys Leu  
 115 120 125

Thr Val Leu Gly Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro  
 130 135 140

Pro Ser Ser Glu Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu  
 145 150 155 160

Ile Ser Asp Phe Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp  
 165 170 175

Ser Ser Pro Val Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln  
 180 185 190

Ser Asn Asn Lys Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu  
 195 200 205

Gln Trp Lys Ser His Lys Ser Tyr Ser Cys Gln Val Thr His Glu Gly  
 210 215 220

Ser Thr Val Glu Lys Thr Val Ala Pro Thr Glu Cys Ser  
 225 230 235

<210> 52  
 <211> 285  
 <212> PRT

<213> Artificial

<220>

<223> Heavy chain variable domain (VH) and a partial region of the Constant Heavy 1 (CH1) domain of 8B10F8 antibody

<400> 52

Met Lys His Leu Trp Phe Phe Leu Leu Leu Val Ala Ala Pro Arg Trp  
1 5 10 15

Val Leu Ser Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys  
20 25 30

Pro Ser Glu Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Asp Ser Ile  
35 40 45

Ser Ser Tyr Ser Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu  
50 55 60

Glu Trp Ile Gly Tyr Ile His Tyr Thr Gly Ser Thr Asn Tyr Asn Pro  
65 70 75 80

Ser Leu Lys Ser Arg Leu Thr Ile Ser Val Asp Ala Ser Lys Asn Gln  
85 90 95

Phe Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr  
100 105 110

Tyr Cys Ala Arg Asp Trp Gly Gly Tyr Ser Ser Ser Trp Thr Tyr Gly  
115 120 125

Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser  
130 135 140

Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr  
145 150 155 160

Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro  
165 170 175

Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val  
180 185 190

His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser  
195 200 205

Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile

210                                      215                                      220  
 Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val  
 225                                      230                                      235                                      240  
 Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala  
                                     245                                      250                                      255  
 Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro  
                                     260                                      265                                      270  
 Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr  
                                     275                                      280                                      285  
  
 <210> 53  
 <211> 236  
 <212> PRT  
 <213> Artificial  
  
 <220>  
 <223> Whole light chain of 8B10F8 antibody  
  
 <400> 53  
  
 Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp  
 1                                      5                                      10                                      15  
  
 Leu Pro Gly Thr Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser  
                                     20                                      25                                      30  
  
 Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser  
                                     35                                      40                                      45  
  
 Gln Gly Ile Ser Asn Ser Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys  
                                     50                                      55                                      60  
  
 Ala Pro Lys Leu Leu Leu Tyr Ala Ala Ser Arg Leu Glu Ser Gly Val  
 65                                      70                                      75                                      80  
  
 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr  
                                     85                                      90                                      95  
  
 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln  
                                     100                                      105                                      110  
  
 Tyr Tyr Ser Thr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile  
                                     115                                      120                                      125

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp  
 130 135 140

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn  
 145 150 155 160

Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu  
 165 170 175

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp  
 180 185 190

Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr  
 195 200 205

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser  
 210 215 220

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
 225 230 235