

PHOSPHORUS-CONTAINING GROUP-SUBSTITUTED QUINOLINE, ITS PREPARATION PROCESS, MEDICAL COMPOSITION CONTAINING THE COMPOUND AND APPLICATION

Technical Field

The present invention relates to the fields of organic chemistry and medicinal chemistry, particularly relates to a phosphorus-containing group-substituted quinoline, its preparation process, pharmaceutical composition containing the compound and the application.

Background Art

Protein kinase is a kind of phosphotransferase, and it has the function of transferring γ -phosphate group of ATP to a specific amino acid residue of a substrate, making a protein phosphorylated and exhibit its physiological and biochemical functions. Protein kinase is a kind of important kinase, and it mainly has the following two functions in signal transduction: on one hand, it regulates the activity of proteins by phosphorylation; on the other hand, it amplifies the signals step by step through protein phosphorylation step by step, thereby causing cell response.

Protein kinase abnormal activity not only closely relates to abnormal state of a certain stage in series of intra- or extracellular signal transduction pathways such as tumor proliferation, apoptosis, metastasis, but also a main cause of incurring series of other human diseases associated with inflammation, or proliferation response, for example, rheumatoid arthritis, cardiovascular and neural system diseases, asthma, psoriasis. It is known by now that there are over 400 kinds of human diseases directly or indirectly related to protein kinase, which makes the protein kinase become another kind of important drug target following the G-protein coupled receptor.

Protein kinase as a big family consists of over 500 members, and is generally classified into two types, i.e., protein tyrosine kinases (PTKs) and serine-threonine kinases. According to the position of kinases in a cell, they are classified into receptor kinases and non-receptor kinases, also called intracellular kinases. The receptor

kinases generally belong to tyrosine kinases, also called receptor tyrosine kinases (RTKs), such receptor kinases consist of the portion outside cell membrane, transmembrane domain and the portion in the cytoplasm, the catalytic portion of the kinase is located in the cytoplasm. Most of the serine-threonine kinases are located inside the cell, and belong to non-receptor kinases, or called cytosolic kinases.

The typical representatives of the RTKs family are growth factor receptors, which have at least 19 sub-families, the following are the main sub-families:

(a) HER family receptor tyrosine kinases, which include EGFR (epithelial growth factor receptor), HER2, HER3 and HER4. EGFR is a target of synthetic small-molecule drugs such as Tarceva[®], Tykerb[®] and monoclonal antibody Erbitux[®] for treating non-small cell lung cancer.

(b) Insulin family consists of insulin receptor (IR), insulin-like growth factor I receptor (IGF-1R) and insulin receptor-related receptor (IRR); wherein IGF-1R is a well-known anti-cancer target, since it is too similar to IR, particularly the kinase portion inside the cell, their amino acid sequences are 100% identical, inhibition of the activity of IGF-1R generally also inhibits the activity of IR. It is proved that IR is also an effective anticancer target, since the inhibition of IR may have the risk of increasing blood sugar, and it needs to find a balance between efficacy and safe risk for using IR inhibitor as anticancer agent.

(c) Platelet-derived growth factor receptors (PDGFRs) family, which includes PDGFR- α , PDGFR- β , CSF1R, c-KIT and c-fms; wherein c-KIT is a molecule target of leukemia-treating drugs such as Gleevec[®] and also for treating gastrointestinal stromal tumors.

(d) Vascular endothelial growth factor receptors (VEGFRs) family, which includes FLT1 (Fms-like tyrosine kinase 1 or VEGFR1), KDR (or VEGFR-2) and FLT4 (or VEGFR3). The members among them are molecular target of Sutent[®] and Naxavar[®].

(e) Fibroblast growth factor receptors (FGFRs) family, which includes FGFR1, FGFR2, FGFR3 and FGFR4 and 7 ligands, FGF1, FGF2, FGF3, FGF4, FGF5, FGF6 and FGF7. The members among them are molecular targets of drugs currently

undergoing clinical tests.

(f) MET family, which includes c-Met or called human hepatocyte growth factor receptor (hHGFR), and RON; wherein c-Met plays an important role in the growth and metastasis of the initial tumor, drugs targarting the member of MET family are currently in clinical trials.

(g) RET family, RETs are receptors of a member from GDNF family and have isoforms of RET51, RET43 and RET9, drugs targeting the member of RET family are currently in clinical trials.

(h) Eph family, which is the biggest family of receptor tyrosine kinases, consists of 16 receptors (EPHA1, EPHA2, EPHA3, EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EPHA9, EPHA10, EPHB1, EPHB2, EPHB3, EPHB4, EPHB5, EPHB6) and 9 ligands (EFNA1, EFNA2, EFNA3, EFNA4, EFNA5, EFNB1, EFNB2, EFNB3). These members play important roles in development of animals, and some of them play roles in tumors.

Non-receptor kinases do not have the portion outside cell membrane and the transmembrane domain, and the whole kinase is in the cytoplasm. By now it is known that there are at least 24 non-receptor kinases, which are divided into 11 subfamilies, i.e. Src, Frk, Btk, CsK, Abl, Zap70, Fes, Fps, Fak, Jak and AcK subfamilies; wherein Src subfamily is the biggest and includes Src, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr, AUR1, AUR2 and Yrk kinases. For more detailed information, see: Neet, K.; Hunter, T. *Genes to Cells* 1996, 1, 147-169 and the documents cited therein. Although several non-receptor kinases belong to tyrosine kinases, most of the non-receptor kinases belong to serine-threonine kinases; wherein several members of them are molecular targets of leukemia-treating drugs such as Gleevec® and Sprycel®.

As stated above, receptor kinases and non-receptor kinases as antitumor targets have been fully proved in clinical and practical applications, and several antitumor drugs are approved to sale on the markets to be used for patients. in addition to treatment of tumors, inhibiting abnormal activity of receptor kinases and non-receptor kinases may also be used for treating the following diseases, which include, but not limited to: psoriasis, hepatic cirrhosis, diabetes, angiogenesis-related diseases,

restenosis-related diseases, eye diseases, age-related macular degeneration, rheumatoid arthritis and other inflammatory diseases, immune system diseases such as autoimmune disease, cardiovascular diseases such as atherosclerosis, or kidney disease etc. Thus it is needed to go on developing desired inhibitors of these kinases.

Contents of the Invention


One purpose of the present invention is to provide a phosphorus-containing group-substituted quinoline having the activity of inhibiting protein kinase, and the preparation process thereof.

Another purpose of the present invention is to provide use of the above phosphorus-containing group-substituted quinoline in manufacturing of medicaments for treating protein kinase abnormal activity-associated diseases.

Still another purpose of the present invention is to provide a pharmaceutical composition comprising the above phosphorus-containing group-substituted quinoline capable of treating protein kinase abnormal activity-associated diseases.

Technical Solution of the Invention

A phosphorus-containing group-substituted quinoline, which has a molecular structure represented by the following formula (I):




in which,

Z represents , V¹ and V² are each independently selected from hydrogen, halogen, -OCF₃, -CF₃, -NO₂, -CN, -OH, -NH₂, -NMe₂, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, or C₃₋₆ heterocycloalkoxy;

Either of R and R' represents phosphorus-containing substituent Q, the other is

selected from hydrogen, methoxyl, methoxyethoxyl, or phosphorus-containing substituent Q;

Wherein, the phosphorus-containing substituent Q represents



A is absent or represents O, NH, S(=O)_m, or C₁₋₆ alkyl, and A is optionally substituted with G¹;

L is absent or represents C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl, and L is optionally substituted with G²;

J is absent or represents O, NH, S(=O)_m, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl and J is optionally substituted with G³;

X is absent or represents -C(=O)-, -S(O)_m-, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl, and X is optionally substituted with G⁴;

Y is absent or represents C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl, and Y is optionally substituted with G⁵;

R¹ and R² are each independently selected from -OH, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, C₆ aryloxy, C₅₋₆ heteroaryloxy, or C₃₋₆ heterocycloalkoxy, and R¹ and R² are each optionally substituted with G⁶; R¹ and R² together with the phosphorus atom to which they are attached may form a C₃₋₆ heterocycloalkyl ring, this C₃₋₆ heterocycloalkyl ring may further comprise heteroatom(s) selected from O, N, or S(=O)_m;

wherein,


G¹, G², G³, G⁴, G⁵ and G⁶ are each independently selected from H, -CN, -CF₃, -CO₂H, halogen, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, R³O-, R³R⁴N-, R³S(=O)_m-, R³R⁴NS(=O)_m-, R⁵C(=O)-, R³R⁴NC(=O)-, R³OC(=O)-, R⁵C(=O)O-, R³R⁴NC(=O)O-, R⁵C(=O)NR³-, R³R⁴NC(=O)NR⁶-, R³OC(=O)NR⁶-, R³S(=O)_mNR⁶-, R³R⁴NS(=O)_mNR⁶-, R³R⁴NC(=NR⁷)NR⁶-, R³R⁴NC(=CHNO₂)NR⁶-, R³R⁴NC(=N-CN)NR⁶-, R³R⁴NC(=NR⁷)-, R³S(=O)(=NR⁷)NR⁶-, or R³R⁴NS(=O)(=NR⁷)-

R³, R⁴, R⁵, R⁶ and R⁷ are each independently selected from H, C₁₋₆ alkyl, C₃₋₆

cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl; when R³ and R⁴ are attached to the same nitrogen atom, R³ and R⁴ together with the nitrogen to which they are attached may form a C₃₋₆ heterocycloalkyl ring, this C₃₋₆ heterocycloalkyl ring may further comprise heteroatom(s) selected from O, N, or S(=O)_m; and R³, R⁴, R⁵, R⁶ and R⁷ may be optionally substituted with halogen, CN, C₁₋₆ alkyl, or C₃₋₆ cycloalkyl;

$$m = 0-2.$$

Provided herein is a phosphorus-containing group-substituted quinoline represented by formula (Ia) below:



in the above formula,

A represents O, NH, or S(=O)_m, and A is optionally substituted with G¹;

L represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and L is optionally substituted with G²;


J represents O, NH, or S(=O)_m, and J is optionally substituted with G³;

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and Y is optionally substituted with G⁵;

R¹ and R² are each independently selected from -OH, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, C₆ aryloxy, C₅₋₆ heteroaryloxy, or C₃₋₆ heterocycloalkoxy, and R¹ and R² are each optionally substituted with G⁶;

$$m = 0-2.$$

A phosphorus-containing group-substituted quinoline is represented by formula (Ib) below:



in the above formula,

A represents O, NH, or S(=O)_m, and A is optionally substituted with G¹;

L represents C₁₋₆ alkyl, and L is optionally substituted with G²;

J represents C₃₋₆ cycloalkyl, or C₃₋₆ heterocycloalkyl, and J is optionally substituted with G³;


X is absent or represents -C(=O)-, or -S(O)_m-;

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl; and Y is optionally substituted with G⁵;

R¹ and R² are each independently selected from -OH, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, C₆ aryloxy, C₅₋₆ heteroaryloxy, or C₃₋₆ heterocycloalkoxy, and R¹ and R² are each optionally substituted with G⁶;

m = 0-2.

A phosphorus-containing group-substituted quinoline is represented by formula (Ic) below:



in the above formula,

A represents O, NH, or S(=O)_m, and A is optionally substituted with G¹;

L represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and L is optionally substituted with G²;


J represents O, NH, or S(=O)_m, and J is optionally substituted with G³;

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and Y is optionally substituted with G⁵;

R¹ and R² are each independently selected from -OH, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, C₆ aryloxy, C₅₋₆ heteroaryloxy, or C₃₋₆ heterocycloalkoxy, and R¹ and R² are each optionally substituted with G⁶;

m = 0-2.

Provided herein is a phosphorus-containing group-substituted quinoline represented by formula (Id) below:



in the above formula,

A represents O, NH, or S(=O)_m, and A is optionally substituted with G¹;

L represents C₁₋₆ alkyl, and L is optionally substituted with G²;

J represents C₃₋₆ cycloalkyl, or C₃₋₆ heterocycloalkyl, and J is optionally substituted with G³;


X is absent or represents -C(=O)-, or -S(O)_m-;

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and Y is optionally substituted with G⁵;

R¹ and R² are each independently selected from -OH, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, C₆ aryloxy, C₅₋₆ heteroaryloxy, or C₃₋₆ heterocycloalkoxy, and R¹ and R² are each optionally substituted with G⁶;

m = 0-2.

Provided herein is a phosphorus-containing group-substituted quinoline represented by formula (Ie) below:




in the above formula,

L represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and L is optionally substituted with G²;

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and Y is optionally substituted with G⁵.

A phosphorus-containing group-substituted quinoline is represented by formula (If) below:



in the above formula,

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and Y is optionally substituted with G⁵.

The phosphorus-containing group-substituted quinoline is selected from the group consisting of one or more of the following compounds:

N1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(diethoxyphosphorylmethylamino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(diethoxyphosphorylmethyl)(methyl)amino]propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(*n*-propyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(methyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(*n*-propyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(acetyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(formyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(propionyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(isopropionyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(cyclopropylformyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(acetyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;
*N*1-[4-[[7-[3-(formyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(propionyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(isopropionyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

;

*N*1-[4-[[7-[3-(cyclopropylformyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[7-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethylaminocarbonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(methylaminocarbonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(*N,N*'-dimethylaminocarbonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[7-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethylaminocarbonyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(methylaminocarbonyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(*N,N*'-dimethylaminocarbonyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

xyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(methylsulfonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethylsulfonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethylamino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethylamino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethyl)(methyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethyl)(n-propyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethyl)(methyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethyl)(n-propyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(acetyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(formyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(propionyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(isopropionyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(cyclopropylformyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(acetyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(formyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(propionyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(isopropionyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

;

*N*1-[4-[[6-[3-(cyclopropylformyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethylaminocarbonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(methylaminocarbonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dic

arboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(*N,N'*-dimethylaminocarbonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethylaminocarbonyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(methylaminocarbonyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(*N,N'*-dimethylaminocarbonyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(methylsulfonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethylsulfonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[[1-(2-diethoxyphosphorylacetyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[[1-(2-diethoxyphosphorylacetyl)-4-piperidinyl]methoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide.

A phosphorus-containing group-substituted quinoline, which is selected from the group consisting of one or more of the following compounds:

*N*1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphoryl(*N*-methyl)methylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(acetyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;


*N*1-[4-[[7-[[1-(2-diethoxyphosphorylacetyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide.

A racemate or enantiomer of a phosphorus-containing group-substituted quinoline.

Provided herein is a preparation process of the phosphorus-containing group-substituted quinoline, consisting of the steps in the following Scheme 1:

Scheme 1



wherein

$t = 0-4$;


LG or GL represents common leaving groups in the organic chemistry field;

PG represents common protecting groups in the organic chemistry field;

RR₁, RR₁₁, RR₁₂, RR₂, RR₃, RR₄, or RR₅ is each selected from H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl, and is optionally substituted with H, -CN, -CF₃, -CO₂H, halogen, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, R³O-, R³R⁴N-, R³S(=O)_m-, R³R⁴NS(=O)_m-, R⁵C(=O)-, R³R⁴NC(=O)-, R³OC(=O)-, R⁵C(=O)O-, R³R⁴NC(=O)O-, R⁵C(=O)NR³-, R³R⁴NC(=O)NR⁶-, R³OC(=O)NR⁶-, R³S(=O)_mNR⁶-, R³R⁴NS(=O)_mNR⁶-, R³R⁴NC(=NR⁷)NR⁶-, R³R⁴NC(=CHNO₂)NR⁶-, R³R⁴NC(=N-CN)NR⁶-, R³R⁴NC(=NR⁷)-, R³S(=O)(=NR⁷)NR⁶-, or R³R⁴NS(=O)(=NR⁷), wherein R³, R⁴, R⁵, R⁶ and R⁷ are each independently selected from hydrogen, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl; when R³ and R⁴ are attached to the same nitrogen atom, R³ and R⁴


together with the nitrogen to which they are attached may form a C₃₋₆ heterocycloalkyl ring, this C₃₋₆ heterocycloalkyl ring may further comprise heteroatom(s) selected from O, N, S(=O)_m; and R³, R⁴, R⁵, R⁶ and R⁷ are each optionally substituted with halogen, CN, C₁₋₆ alkyl, or C₃₋₆ cycloalkyl; m = 0-2.

Provided herein is a preparation process of the phosphorus-containing group-substituted quinoline, consisting of the steps in the following Scheme 2:




wherein

PP' can be any one of PG, H and PP, and PG can be converted into H by deprotection. Provided herein is preparation process of the phosphorus-containing group-substituted quinoline, consisting of the steps in the following Scheme 3:




Provided herein is a preparation process of there phosphorus-containing

group-substituted quinoline, consisting of the steps in the following Scheme 4:



Provided herein is a preparation process of the phosphorus-containing group-substituted quinoline, consisting of the steps in the following Scheme 5:



Provided herein is the use of a pharmaceutical composition comprising a phosphorus-containing group-substituted quinoline, wherein the composition is used for treating protein kinase abnormal activity-associated diseases.

Provided herein is the use of a pharmaceutical composition comprising a phosphorus-containing group-substituted quinoline, wherein the protein kinase is c-Met, KDR, or VEGFR2, RET, PDGFR- β , c-KIT, Flt3, MEK5, DDR1, LOK, CSF1R, EPHA7, EPHA8, EPHB6, MKNK2, BLK, HIPK4, HCK, or Flt4.

Provided herein is the use of a pharmaceutical composition comprising a phosphorus-containing group-substituted quinoline, wherein the protein kinase is RON, ALK (or Anaplastic Lymphoma Kinase), EGF1R, HER2, HER3, HER4,

PDGFR- α , c-fms, FLT1, Src, Frk, Btk, CsK, Abl, Fes, Fps, Fak, AcK, Yes, Fyn, Lyn, Lck, Hck, Fgr, Yrk, PDK1, TAK1, Tie-2, Tie-1, YSK4, TRK A, TRK B, TRK C, SLK, PKN2, PLK4, MST1R, MAP4K, or DDR2.

Provided herein is the use of a pharmaceutical composition comprising a phosphorus-containing group-substituted quinoline, wherein the disease treated with the composition is psoriasis, hepatic cirrhosis, diabetes, angiogenesis-mediated diseases, eye diseases, immune system diseases, or cardiovascular diseases.

Provided herein is the use of a pharmaceutical composition comprising a phosphorus-containing group-substituted quinoline, wherein the disease treated with the composition is tumor, including solid and liquid tumors.

Provided herein is the use of a pharmaceutical composition comprising a phosphorus-containing group-substituted quinoline, wherein the disease treated with the composition is tumor, including lung cancer, bone cancer, pancreatic cancer, skin cancer, head and neck cancer, skin or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, anal region cancer, stomach cancer, colon cancer, breast cancer, fallopian tube cancer, endometrial cancer, cervical cancer, vaginal cancer, vulva cancer, Hodgkin's disease, esophageal cancer, intestinal cancer, endocrine system cancer, thyroid cancer, parathyroid cancer, soft tissue sarcoma, urethral cancer, penile cancer, prostate cancer, chronic or acute leukemia, bladder cancer, kidney or ureter cancer, renal cancer, central nervous system (CNS) tumors, spinal axis tumors, pituitary adenomas, gastrointestinal stromal tumors, colorectal cancer, non-small cell lung cancer, small cell lung cancer, mastocytosis, glioma, sarcoma, lymphoma

Provided herein is a medicament for treating protein kinase abnormal activity-associated diseases, which comprises any one of the above compounds or pharmaceutically acceptable salt, solvate, prodrug thereof, or a racemate or enantiomer of any one of the above compounds or pharmaceutically acceptable salt, solvate, prodrug thereof.

The above medicament further comprises at least one pharmaceutically acceptable carrier.

The above medicament is in a form of the following Formulations: (1) oral formulation; (2) injectable formulation; (3) anal suppository; (4) nasal inhalation; (5) eye drop; (6) skin patch.

A series of experiments confirm that, the phosphorus-containing

group-substituted quinolines according to the present invention have the following advantages: (1) the screening tests for inhibiting the kinase reveal that, the compound of present invention exhibits strong inhibiting effect on a series of protein kinases and mutants thereof; (2) the tests for inhibiting tumor reveal that, such phosphorus-containing group-substituted quinoline significantly inhibits tumors without apparent toxicity; (3) the compound of the present invention can be used in combination with other anti-tumor drugs and thus obtain synergistic or additive effect; (4) the compound of the present invention can be used together with other tumor therapies, for example, radiotherapy, interventional therapy etc. This shows that a phosphorus-containing group-substituted quinoline according to the invention may be used as a medicament for effective treatment of protein kinase abnormal activity-associated diseases.

In the protein kinase abnormal activity-associated diseases treated by the compound according to the invention, the kidney cancers are adrenal cancer, renal cell carcinoma or carcinoma of renal pelvis; the gliomas are brain stem neuroglioma, neuroendocrine glioma and neuroglioma.

Besides tumor, the protein kinase abnormal activity-associated diseases which the compound of the invention is used to treat further include psoriasis, hepatic cirrhosis, diabetes, angiogenesis-related diseases, restenosis-related diseases, eye diseases such as AMD, rheumatoid arthritis and other inflammatory diseases, immune system diseases such as autoimmune disease (eg. AIDS), cardiovascular diseases such as atherosclerosis, or kidney disease etc.

A pharmaceutical composition comprising the compound of the present invention may be used for treating protein kinase abnormal activity-associated diseases in mammals, such as human patients.

Through a preparation (formulation) process, the compound according to the invention (including racemate, enantiomer and other stereoisomer) or the pharmaceutically acceptable salt, hydrate, solvate or prodrug thereof together with proper pharmaceutically acceptable carriers and commonly used medical auxiliaries may be prepared into a pharmaceutical composition which is favorable for

administration.

The routes of administration of the medicament comprising the compound according to the invention may be: (1) oral administration, for example, tablet, capsule etc.; (2) injection, for example, intravenous, subcutaneous, intramuscular, eye injection, intraperitoneal injection etc.; (3), anal plug, for example, suppository, gels etc.; (4), nasal inhalation, for example spray, aerosol etc.; (5) eye drop; (6) skin patch. The drug release system, for example, liposome, slow release techniques etc. also may be used, wherein oral administration and injection are preferably used, particularly oral administration is more preferably used.

The various formulations of the pharmaceutical composition comprising the compound according to the present invention may be prepared by conventional methods in medical industry, for example, mixing, dissolving, granulating, grinding, emulsifying, encapsulating, sugar-coating, freeze-drying, freeze spraying etc.

The content of the compound according to the present invention present in the pharmaceutical composition is in a range of 0.001-100%. The effective amount of the pharmaceutical composition for administering in mammals including human beings is 0.1-500 mg / per kilogram body weight in each day, preferably in an amount of 1-100 mg / per kilogram body weight in each day. In such an effective amount range, the compound of the present invention exhibits the pharmaceutical actions of inhibiting protein kinase activity and treating protein kinase abnormal activity-associated diseases (for example, cancer).

The administration frequency of the medicament according to the present invention may vary, depending on the used compound or the pharmaceutical composition thereof and the diseases to be treated; the pharmaceutical composition according to the present invention is typically administered 1-6 times each day, optimized administration frequency is 1-3 times each day.

The packaging and storing of the medicament according to the present invention are similar to those of the general western medicines, for example, the medicament of solid formulation can be directly filled in a bottle made of glass, plastic, paper or metal, desiccant is preferably contained in the bottle to maintain the quality of the medicament; generally the medicament of liquid formulation is filled in a bottle made of glass, plastic or metal, or flexible pipe; and the medicament of fogging formulation is generally filled in a overpressure resistant container made of plastic or metal and

equipped with pressure-reducing valve etc.

Definition of Terms

The following are definitions of the terms mentioned in the present invention. The variable groups used in the present invention, for example R^a, R^b, m etc. are only applied to this section (i.e., the section of "Definition of Terms").

According to the common knowledge of a person skilled in this field, most of the chemical reactions need to be carried out in a solvent; the commonly used solvents for preparing the compound according to the invention include, but not limited to, water, methanol, ethanol, isopropanol, n-propanol, n-butanol, isobutanol, t-butanol, 2-methoxyethanol, 2,2,2-trifluoroethanol, dichloromethane, 1,2-dichloroethane, chloroform, THF, dioxane, DME, ethyl acetate, diethyl ether, methyl t-butyl ether, hexane, cyclohexane, toluene, acetonitrile, DMF, DMSO, or a mixture of any two or more of the above solvents. In some cases the chemical reactions need to be carried out in the presence of acid or base; the commonly used bases for preparing the compound according to the invention include, but not limited to, Et₃N, Me₃N, *i*-Pr₂NEt, pyridine, DBU, DABCO, tetramethyl guanidine, NaOH, KOH, Cs₂CO₃, Na₂CO₃, K₂CO₃, NaHCO₃, KF, CsF, K₃PO₃, K₂HPO₄, KH₂PO₄, NaH, *n*-BuLi, *s*-BuLi, *t*-BuLi, NaN(SiMe₃)₂, LiN(SiMe₃)₂, KN(SiMe₃)₂, or a mixture of any two or more of the above bases; and the commonly used acids include, but not limited to, HCO₂H, AcOH, TFA (trifluoroacetic acid), HCl (hydrochloric acid), H₂SO₄, HNO₃, H₃PO₄, *p*-TsOH, PhSO₃H, CSA, MsOH or Lewis acid such as ZnCl₂, AlCl₃, BF₃·OEt₂. In some cases the chemical reactions need to be carried out in the presence of coupling reagent; the commonly used coupling reagents for preparing the compound according to the invention include, but not limited to, DCC, EDC, HATU, TBTU, PyBOP, HCTU, BOP, DIC, HOBt, HOAt, CDI, DEPBT etc. In some steps of preparing the compound according to the invention, reduction reaction and reducing reagents are needed; the reducing reagents include, but not limited to, H₂+Pd/C, H₂+Pd(OH)₂, H₂+PtO₂, H₂+Raney Ni, H₂NNH₂+Raney Ni, Mg+MeOH, Fe+AcOH, Fe+HCl, Zn+AcOH, Zn+HCl, Zn+NH₄OAc, SnCl₂, LiAlH₄, NaBH₄, NaBH₃(CN), NaB(OAc)₃H, BH₃, etc. In some steps of preparing the compound according to the invention, deprotections are needed; when the protecting group is Boc, the commonly used deprotection reagents include, but not limited to, HCl, TFA, H₂SO₄ etc.; when the protecting group is CBZ, the commonly used deprotection reagents include, but not limited to, strong HCl, H₂+Pd/C etc.; when the protecting group is Bn, the

commonly used deprotection reagents include, but not limited to, $\text{H}_2+\text{Pd/C}$, $\text{H}_2+\text{Pd}(\text{OH})_2$, $\text{H}_2+\text{Pd/C}+\text{HCl}$ etc. The reaction for preparing the compound according to the invention is typically carried out at room temperature, however sometimes the temperature needs to be decreased to $-78\text{ }^\circ\text{C}$ or increased to $200\text{ }^\circ\text{C}$; typically the reaction is carried out at the conditions of the above solvents, and temperatures with conventionally stirring, however sometimes it needs to be carried out in a microwave oven; when the base, reagent, catalyst are sensitive to water or oxygen, the reaction needs to be carried out under the anhydrous and oxygen-free conditions; in such case protonic solvents can not be used.

The term “pharmaceutically acceptable salt” means a salt formed by the reaction of the compound according to the invention with inorganic acid, organic acid, inorganic base, or organic base; the salt has the same or similar biological activity and effectiveness of the compound according to the invention. The inorganic acid or organic acid may be hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, nitric acid, carbonic acid, phosphoric acid, perchloric acid, acetic acid, citric acid, oxalic acid, lactic acid, malic acid, salicylic acid, tartaric acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, acid substituted benzenesulfonic acid (for example, p-toluene sulfonic acid), isonicotinic acid, oleic acid, tannic acid, pantothenic acid, ascorbic acid, succinic acid, maleic acid, gentisic acid, fumaric acid, gluconic acid, uronic acid, glucaric acid or sugar acid, formic acid, benzoic acid, glutamic acid, pantoic acid, sorbic acid etc.; the inorganic base or organic base may be sodium hydroxide, potassium hydroxide, lithium hydroxide, iron hydroxide, calcium hydroxide, barium hydroxide, aluminum hydroxide, magnesium hydroxide, zinc hydroxide, ammonia, organic quaternary ammonium hydroxide, sodium carbonate, potassium carbonate, lithium carbonate, calcium carbonate, barium carbonate, magnesium carbonate, carbonated organic quaternary ammonium salts, sodium hydrogencarbonate, potassium hydrogencarbonate, lithium hydrogencarbonate, calcium hydrogencarbonate, barium hydrogencarbonate, magnesium hydrogencarbonate, hydrogencarbonated organic quaternary ammonium salt etc.

The term “solvate” means a stable substance formed by the compound according to the invention and a solvent commonly used in chemical reactions through covalent bond, hydrogen bond, ionic bond, van der Waals forces, complexation, inclusion etc.; the solvent may be methanol, ethanol, propanol, butanol, ethylene glycol, propanediol, polyethylene glycol, acetone, acetonitrile, diethyl ether, methyl tert-butyl ether etc.

The term “hydrate” means a solvate, in which the solvent is water.

The term “prodrug” means a compound which is obtained by conversion of the compound of the present invention by chemical synthesis or physical approach and is converted back into the compound according to the present invention in the body of the mammal after it is administrated to a mammal. Usually the “prodrug” approach is used to overcome the poor or bad physicochemical property or druggability of the medical compound itself.

The term “racemate, enantiomer and other stereoisomer” means compounds having the same molecular formula and molecular weight, but they are different due to the different bonding manners between atoms and different space arrangements, such compounds are called isomer or stereoisomer. When these stereoisomers mirror image for each other, i.e. look similar but can not superimpose completely, just like the left hand and right hand; these compounds are called enantiomers. The absolute configurations of enantiomers are typically represented by (*R*)- and (*S*)- or *R*- and *S*-. Particularly the rules of determining absolute configurations, see Chapter 4 of “Advanced Organic Chemistry,” 4th edition (by J. March, John Wiley and Sons, New York, 1992). (*R*)- and (*S*)-enantiomers rotate polarized light through opposite angles, i.e. rotating left and rotating right. When (*R*)- and (*S*)-enantiomers are mixed or present in a ratio of 1:1, the mixture has no rotating effect on polarized light, such a mixture is called racemate.

The compound according to the invention may have tautomers, rotamers, cis-trans isomers etc., these concepts can be found and understood in “Advanced Organic Chemistry,” 4th edition, by J. March. As long as these isomers have the same function of inhibiting the activity of protein kinase as the compound according to the invention, thus these isomers are covered by the present invention.

According to the common knowledge in the art, after the compound of the present invention is administrated to mammals (for example, human beings), it is likely to be metabolized into various metabolites in the body of the mammal by different enzymes, as long as these metabolites have similar function of inhibiting protein kinase activity to that of the compound according to the invention, they are also covered by the present invention.

The term “pharmaceutical composition” means a mixture obtained by mixing one or more of the compounds according to the invention, pharmaceutically acceptable salt or solvate, or hydrate or prodrug thereof with other chemical

ingredient (for example pharmaceutically acceptable carrier). The purpose of preparation of a pharmaceutical composition is to facilitate the administration to animals. The above pharmaceutical compositions, besides a pharmaceutically acceptable carrier, may further comprise pharmaceutically commonly used auxiliaries, for example, antibacterial agent, antifungal agent, antimicrobial agent, preservative agent, toner, solubilizer, thickening agent, surfactant, complexing agent, protein, amino acid, fat, carbohydrate, vitamins, minerals, trace elements, sweetening agent, pigment, essence or the combinations thereof.


The term “pharmaceutically acceptable carrier” refers to a non-active ingredients in the pharmaceutical composition, it may be calcium carbonate, calcium phosphate, a variety of sugars (for example lactose, mannitol etc.), starch, cyclodextrin, magnesium stearate, cellulose, magnesium carbonate, acrylic polymer, methyl, acrylic polymer, gel, water, polyethylene, glycol, propanediol, ethylene glycol, castor oil, hydrogenated castor oil, polyethoxyl hydrogenated castor oil, sesame oil, corn oil, peanut oil etc.

The term “alkyl” means a linear or branched saturated hydrocarbon group having specific number of carbon atoms, for example C₁₋₁₂ alkyl stands for linear or branched groups having 1-12 carbon atoms. C₀ alkyl stands for one covalent single bond. As used in the present invention, the alkyl includes, but is not limited to, methyl, ethyl, propyl, butyl, isopropyl, neopentyl, 2-methyl-1-hexyl etc. One or all hydrogen atoms of the alkyl may be substituted with the following groups: cycloalkyl, aryl, heteroaryl, heterocycloalkyl ring, halogen, amino, hydroxyl, cyano, nitro, carboxyl, thio, oxo, alkoxy, aryloxy, alkylthio, arylthio, carbonyl, thiocarbonyl, C-amide, N-amide, O-aminocarbonyloxy, N-aminocarbonyloxy, O-thioaminocarbonyloxy, N-thioaminocarbonyloxy, C-ester, O-ester and -NR^aR^b, wherein R^a and R^b are respectively selected from hydrogen, alkyl, cycloalkyl, aryl, acetyl, carbonyl, sulfonyl, and trifluoromethylsulfonyl, and R^a and R^b together with the nitrogen atom to which they are attached may form a 5- or 6-member heterocycloalkyl ring.

The term “cycloalkyl” or “cyclic alkyl” stands for a single-, double- or multiple-ring hydrocarbon group having specific number of carbon atoms, in case of double- or multiple-ring, they can be joined together in fused (two adjacent carbon atoms are shared by two or more rings) or spiro (one carbon atoms is shared by two or more rings) manner, for example C₁₋₁₂ cycloalkyl stands for single-, double- or multiple-ring hydrocarbon compound groups having 1-12 carbon atoms. C₀ cycloalkyl

stands for one covalent single bond. A cycloalkyl may comprise unsaturated double- or triple-bond, but does not have completely conjugated π electronic system. As used in the present invention, the cycloalkyl includes, but not limited to, cyclopropyl, cyclobutyl, cyclohexyl, cyclopentenyl, cycloheptatrienyl, adamantane (examples are shown in Table 1):

Table 1



One or all hydrogen atoms of the cycloalkyl or cycloalkane may be substituted with the following groups: alkyl, aryl, heteroaryl, heterocycloalkyl ring, halogen, amino, hydroxyl, cyano, nitro, carboxyl, thio, oxo, alkoxy, aryloxy, alkylthio, arylthio, carbonyl, thiocarbonyl, C-amide, N-amide, O-aminocarbonyloxy, N-aminocarbonyloxy, O-thioaminocarbonyloxy, N-thioaminocarbonyloxy, C-ester, O-ester and $-NR^aR^b$, wherein R^a and R^b are respectively selected from hydrogen, alkyl, cycloalkyl, aryl, acetyl, carbonyl, sulfonyl, trifluoromethylsulfonyl, and R^a and R^b together with the nitrogen atom to which they are attached may form a 5- or 6-member heterocycloalkyl ring.

The term “halogen” stands for fluorine, chlorine, bromine or iodine.

The term “alkoxy” or “alkoxyl” means that an alkyl having specified number of carbon atoms is attached to other group through oxygen atom. As used in the present invention, alkoxy includes, but is not limited to, methoxyl, ethoxyl, propoxy, butoxyl, cyclopentyloxy, cyclohexyloxy, isopropoxy, neopentyloxy, 2-methyl-1-hexyloxy etc.


The term “cycloalkoxy” or “cycloalkoxyl” means that a cycloalkyl having specified number of carbon atoms is attached to other group through oxygen atom. As used in the present invention, cycloalkoxy includes, but not limited to, cyclopropoxy, cyclobutoxy, cyclohexoxy, and the like.

The term “aryl” stands for a single-, double- or multiple-ring group consisting of 6-12 carbon atoms, wherein at least one ring has completely conjugated π electronic system and meets for the Rule $N+2$, i.e., having aromaticity; but the whole group is not necessary to conjugate completely. An aryl may also be present in form of arylene, i.e., there are two or more atoms attached to other groups in the structure of aryl. As

used in the present invention, aryl includes, but is not limited to, phenyl, naphthyl, indenyl, indanyl, tetrahydronaphthalene etc. One or all hydrogen atoms of the aryl may be substituted with the following groups: alkyl, cycloalkyl, heteroaryl, heterocycloalkyl ring, halogen, amino, hydroxyl, cyano, nitro, carboxyl, thio, oxo, alkoxy, aryloxy, alkylthio, arylthio, carbonyl, thiocarbonyl, C-amide, N-amide, O-aminocarbonyloxy, N-aminocarbonyloxy, O-thioaminocarbonyloxy, N-thioaminocarbonyloxy, C-ester, O-ester and $-NR^aR^b$, wherein R^a and R^b are respectively selected from the group consisting of hydrogen, alkyl, cycloalkyl, aryl, acetyl, carbonyl, sulfonyl and trifluoromethylsulfonyl, and R^a and R^b together with the nitrogen atom to which they are attached may form a 5- or 6-member heterocycloalkyl ring.

The term “heteroaryl” stands for a single-, double- or multiple-ring group consisting of 5-12 ring atoms other than hydrogens, wherein at least one atom is O, N or $S(=O)_m$ (wherein $m = 0-2$), and, wherein at least one ring has completely conjugated π electronic system and meets for the Rule $N+2$, i.e., having aromaticity; but the whole group is not necessary to conjugate completely; for example, C_5 heteroaryl stands for an aromatic ring group consisting of 5 ring atoms, wherein at least one ring atom is selected from O, N or $S(=O)_m$ (wherein $m = 0-2$). A heteroaryl may also be present in form of heteroarylene, i.e., there are two or more atoms attached to other groups in the structure of heteroaryl. As used in the present invention, heteroaryl includes, but not limited to, pyridinyl, pyridinonyl, tetrahydropyridinonyl, pyrimidinyl, pyrazinyl, pyridazinyl, imidazolyl, thiazolyl, thiophenyl, furanyl, indolyl, azaindolyl, benzimidazolyl, indolinyl, indolonyl, quinolinyl etc. (examples are shown in Table 2):


Table 2



One or all hydrogen atoms of the heteroaryl may be substituted with the following groups: alkyl, cycloalkyl, aryl, heterocycloalkyl ring, halogen, amino, hydroxyl, cyano, nitro, carboxyl, thio, oxo, alkoxy, aryloxy, alkylthio, arylthio, carbonyl, thiocarbonyl, C-amide, N-amide, O-aminocarbonyloxy, N-aminocarbonyloxy, O-thioaminocarbonyloxy, N-thioaminocarbonyloxy, C-ester, O-ester and $-NR^aR^b$, wherein R^a and R^b are respectively selected from the group consisting of hydrogen, alkyl, cycloalkyl, aryl, acetyl, carbonyl, sulfonyl, and trifluoromethylsulfonyl, and R^a and R^b together with the nitrogen atom to which they are attached may form a 5- or 6-member heterocycloalkyl ring.

The term “heterocycloalkyl or heterocycloalkyl ring” stands for a single-, double- or multiple-ring alkyl or alkane consisting of 3-12 ring atoms other than hydrogen atoms, wherein at least one atom is O, N or $S(=O)_m$ (wherein $m = 0-2$), for example, C_6 heterocycloalkyl stands for a single ring group consisting of 6 ring atoms, wherein at least one ring atom is selected from O, N or $S(=O)_m$ (wherein $m = 0-2$). Such ring may further comprise double or triple bonds other than single bonds, but these double or triple bonds do not form a aromatic structure. Such single-, double- or multiple-ring alkyls or alkanes may be present in form of fused ring, bridged ring or spiro ring. As used in the present invention, heterocycloalkyl or heterocycloalkyl ring includes, but not limited to, piperidinyl, morpholinyl, piperazinyl, pyrrolidinyl, indolinyl, tetrahydropyridinyl, tetrahydrofuranyl, tropinyl etc. (examples are shown in Table 3):

Table 3



One or all hydrogen atoms of the heterocycloalkyl or heterocycloalkyl ring may be substituted with the following groups: alkyl, cycloalkyl, aryl, heteroaryl, heterocycloalkyl ring, halogen, amino, hydroxyl, cyano, nitro, carboxyl, thio, oxo, alkoxy, aryloxy, alkylthio, arylthio, carbonyl, thiocarbonyl, C-amide, N-amide, O-aminocarbonyloxy, N-aminocarbonyloxy, O-thioaminocarbonyloxy, N-thioaminocarbonyloxy, C-ester, O-ester and $\text{-NR}^a\text{R}^b$, wherein R^a and R^b are respectively selected from the group consisting of: hydrogen, alkyl, cycloalkyl, aryl, acetyl, carbonyl, sulfonyl, trifluoromethylsulfonyl, and R^a and R^b together with the nitrogen atom to which they are attached may form a 5- or 6-member heterocycloalkyl ring.


The term “aryloxy” means that an aryl is attached to other group through oxygen atom. As used in the present invention, aryloxy includes, but not limited to, phenoxy, naphthyloxy etc.

The term “heteroaryloxy” means that a heteroaryl is attached to other group through oxygen atom. As used in the present invention, heteroaryloxy includes, but not limited to, 4-pyridinyloxy, 2-thienyloxy etc.

The term “amino” stands for $\text{H}_2\text{N-}$ or $\text{H}_2\text{N-}$ in which hydrogen atoms are substituted, i.e., $\text{R}^a\text{HN-}$ and $\text{R}^a\text{R}^b\text{N-}$.

The term “oxo” or “oxyl” stands for $=\text{O}$, i.e., the oxygen is attached to carbon or heteroatoms including N, S, P, through double bond. Examples of substance substituted by oxyl include, but not limited to, those shown in Table 4:

Table 4



“Hydroxyl” stands for -OH.

“Nitro” stands for -NO₂.

“Carboxyl” stands for -CO₂H.

“Thio” stands for -SH.

“Alkylthio” stands for alkyl-S-.

“Arylthio” stands for aryl-S-.

“Carbonyl” stands for -C(=O)-.

“Thiocarbonyl” stands for -C(=S)-.

“C-amide” stands for -C(=O)NR^aR^b.

“N-amide” stands for C(=O)NR^a-.

“O- aminocarbonyloxy” stands for -O-C(=O)NR^aR^b.

“N- aminocarbonyloxy” stands for O-C(=O)NR^a-.

“O- thioaminocarbonyloxy” stands for -O-C(=S)NR^aR^b.

“N- thioaminocarbonyloxy” stands for O-C(=S)NR^a-.

“C- ester” stands for -C(=O)OR^a.

“N- ester” stands for C(=O)O-.

“Acetyl” stands for CH₃C(=O)-.

“Sulfonyl” stands for -SO₂R^a.

“Trifluoromethylsulfonyl” stands for CF₃SO₂-.

Brief Description of the Drawings

Fig. 1 is a picture of tumors from the GTL-16 xenograft mouse models.

Fig. 2 is the curve of tumor sizes from the GTL-16 xenograft mouse models.

Fig. 3 is the curve showing the body weight changes in the GTL-16 xenograft mouse models.

Fig. 4 is the molecular structure of the compound according to the invention.

Specific Embodiments

The present invention is further described in the following examples, such that

the public may further understand the method of preparing the compound according to the invention and the advantageous effect thereof. But the examples should not be construed as limiting the scope of the present application.

The followings are the abbreviations used in the Examples and their corresponding meanings. If any abbreviation not included in the following list appears in the Examples, it stands for a generally accepted meaning.

DMSO: dimethyl sulfoxide

TMS: tetramethylsilane

DCM: dichloromethane

CDCl₃: deuterio-trichloromethane

CD₃OD: deuterio-methanol

DME: 1, 2-dimethoxyethane

THF: tetrahydrofuran

aq.: aqueous solution

TLC: thin-layer chromatography

LC-MS: liquid chromatography-mass spectrometry

g: gram

mg: milligram

mmol: milimole

μM: micromole

μL: microliter

nM: nanomole

M: molarity

N: normality (equivalent concentration)

m/z: mass-to-charge ratio

δ: chemical shift

DMAP: 4-dimethylaminopyridine

DIPEA: diisopropylethylamine

HATU: 2-(7-azabenzotriazol)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate

EDC.HCl: 1-ethyl-3- (3-dimethylaminopropyl)carbodiimide hydrochloride.

General experimental conditions:

The hydrogen and carbon spectra of Nuclear Magnetic Resonance are obtained


by using equipment Varian INOVA 500NB or Varian 400 MHz or Bruker 400 MHz (the solvent is deuterio-DMSO, deuterio-trichloromethane, or deuterio-methanol etc, and the internal standard is TMS). Mass spectrum is obtained using Liquid Chromatography-Mass Spectrometer (ESI or APCI ion source ZQ4000, Waters Corporation, USA). Ultraviolet spectrum is measured by UV-3010 ultraviolet spectrophotometer of Hitachi Corporation, Japan. Infrared spectrum is measured by NICOLET6700 infrared spectrometer (KBr discs). High Performance Liquid Chromatography is obtained by Waters 2695 ZORBAX High Performance Liquid Chromatograph (Bx-C₈ 5 μ 150 $\times$ 4.6 mm Column). Melting point is measured by Electrothermal Digital Melting Point Apparatus IA9100, without correction.

Starting materials, reagents and solvents are commercially available from: Beta-Pharma, Shanghai; Shanghai PI Chemicals; AndaChem, Taiyuan; Shanghai FWD Chemicals; Sigma-Aldrich, Milwaukee, WI, USA; Acros, Morris Plains, NJ, USA; Frontier Scientific, Logan, Utah, USA; Alfa Aesar, Ward Hill, MA, USA etc., or synthesized by methods reported in the literature. Unless otherwise indicated, generally a solvent from supplier is directly used without being dried or it is dried by molecular sieve.

Preparation methods of various intermediates (including intermediate A, intermediate B, intermediate C, intermediate D, intermediate F, intermediate G, intermediate H, intermediate I, intermediate J, intermediate K) needed for preparing the compound according to the invention are as follows:

Preparation method of intermediate A:


4-(2-fluoro-4-nitrophenoxy)-6-methoxyquinolyl-7-ol:



The Intermediate A is prepared according to the method of WO2008/035209, and the analytical data of the resulting product are as follows: ¹HNMR (400 MHz, DMSO-*d*₆): δ = 11.74 (s, br, 1 H), 8.76 (d, *J* = 6.8 Hz, 1 H), 8.47 (dd, *J* = 2.8 Hz, *J* = 10.4 Hz, 1 H), 8.23 (dd, *J* = 1.2 Hz, *J* = 8.8 Hz, 1 H), 7.81 (m, 1 H), 7.65 (s, 1 H), 7.49 (s, 1 H), 7.04 (d, *J* = 6.4 Hz, 1 H), 3.94 (s, 3 H).

Preparation method of intermediate B:


4-(2-fluoro-4-nitrophenoxy)-7-methoxyquinolyl-6-ol:



The Intermediate B is prepared according to the methods of documents WO2003/033472 and WO2004/039782, and the analytical data of the resulting product are as follows: mass spectrum m/z : 331.12 [M+H].

Preparation method of intermediate C: *tert*-butyl


4-[[4-(4-amino-2-fluorophenoxy)-6-methoxyl-7-quinolyl]oxymethyl]piperidine-1-carboxylate:



The Intermediate C is prepared according to the method of document WO2008/076415, and the analytical data of the resulting product are as follows: mass spectrum m/z : 498.21 [M+H].

Preparation method of intermediate D:


1-[(4-fluorophenyl)carbamoyl]cyclopropanecarboxylic acid:



The Intermediate C is prepared according to the method of document WO2005/030140, and the analytical data of the resulting product are as follows: ^1H NMR (400 MHz, DMSO- d_6): δ = 13.0 (s, 1 H), 10.6 (s, 1 H), 7.62-7.57 (m, 2 H), 7.15-7.09 (m, 2 H), 1.39 (s, 4 H).

Preparation method of intermediate E:


N-benzyl-3-chloro-*N*-(diethoxyphosphorylmethyl)propan-1-amine:



1.6 mL of 37% aqueous formaldehyde solution is added to a suspension of *N*-benzyl-3-chloropropan-1-amine hydrochloride (7.9 g or 35.9 mmol) in dioxane (60 mL), the resulting mixture is slightly heated until it becomes a solution. The solution is cooled to room temperature, 3.2 mL of diethyl phosphate is added thereto, stirring at room temperature for 30 min, and then stirring at 90-100 °C for 3 h, the reaction is completed. The solvent is removed under reduced pressure, and the residue is purified by column chromatography (eluent: 1-4% MeOH in DCM) to obtain 7.6 g target compound (yield: 63%). The analytical data of the obtained product are as follows: ¹HNMR (400 MHz, DMSO-*d*₆): δ= 7.30-7.21 (m, 5 H), 4.01-3.93 (m, 4 H), 3.69 (s, 2 H), 3.59 (t, *J* = 6.8 Hz, 2 H), 2.83(d, *J* = 10.0 Hz, 2 H), 2.61 (t, *J* = 6.8 Hz, 2 H), 1.85-1.82 (m, 2 H), 1.2 (t, *J* = 7.2 Hz, 6 H).

Preparation method of intermediate F:

4-[[7-[3-(benzyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenylamine:



Step 1: preparation of

N-benzyl-*N*-(diethoxyphosphorylmethyl)-3-[[4-(2-fluoro-4-nitrophenoxy)-6-methoxyl-7-quinolyl]oxyl]propan-1-amine: potassium carbonate (7.3 g or 53.2 mmol) is added to DMF solution (100 mL) of intermediate E (7.6 g or 22.8 mmol) and intermediate A (5.0 g or 15.2 mmol). The resulting mixture is stirred at 110 °C for 2 h. The reaction


mixture is diluted with 500 ml of ethyl acetate, inorganic salt is removed by filtering, the filtrate is concentrated, and the residue is purified by column chromatography (eluent: 1-4% MeOH in DCM) to obtain 3.6 g target compound (yield: 38%). The analytical data of the obtained product are as follows: ^1H NMR (400 MHz, DMSO- d_6): δ = 8.57 (d, J = 5.2 Hz, 1 H), 8.47 (dd, J = 2.4, 10.4 Hz, 1 H), 8.21-8.18 (m, 1 H), 7.63 (t, J = 8.8 Hz, 1 H), 7.41-7.18 (m, 8 H), 6.78 (d, J = 5.2 Hz, 1 H), 4.17 (t, J = 6.4 Hz, 2 H), 4.04-3.96 (m, 4 H), 3.86 (s, 3 H), 3.78 (s, 2 H), 2.91 (d, J = 10.0 Hz, 2 H), 2.73 (t, J = 6.4 Hz, 2 H), 2.00 (t, J = 6.4 Hz, 2 H), 1.24-1.19 (m, 6 H).

Step 2: preparation of 4-[[7-[3-(benzyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenylamine:

N-benzyl-*N*-(diethoxyphosphorylmethyl)-3-[[4-(2-fluoro-4-nitrophenoxy)-6-methoxyl-7-quinolyl]oxyl]propan-1-amine (3.6 g or 28.7 mmol) obtained from Step 1 is dissolved in 300 ml of THF, 5 g of Raney nickel is added thereto, the resulting mixture is stirred at 30 °C under 30 psi of hydrogen pressure for 2 h. Finally the reaction mixture is filtered, and the filtrate is concentrated under reduced pressure to obtain 2.9 g of target compound (yield: 83%). The analytical data of the obtained product are as follows: mass spectrum m/z : 598.01 [M+H].

Preparation method of intermediate G:


4-[[6-[3-(benzyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenylamine:



Starting from intermediate E and intermediate B, intermediate G is prepared in the same manner as the method of preparing intermediate F. The analytical data of the obtained product are as follows: mass spectrum m/z : 598.08 [M+H].

Preparation method of intermediate H:


3-chloro-*N*-(diethoxyphosphorylmethyl)-*N*-methylpropan-1-amine:



0.8 mL of 37% aqueous formaldehyde solution is added to suspension of 3-chloro-*N*-methylpropan-1-amine (1.4 g or 1.3 mmol) in dioxane (30 mL), the resulting mixture is slightly heated until it becomes a solution. The solution is cooled to room temperature, 1.6 mL of diethyl phosphate is added thereto, stirring at room temperature for 30 min, and then stirring at 90-100 °C for 3 h, the reaction is completed. The solvent is removed under reduced pressure, and the residue is purified by column chromatography (eluent: 1-4% MeOH in DCM) to obtain 2.3 g target compound (yield: 81%). The analytical data of the obtained product are as follows: ¹HNMR (400 MHz, CDCl₃): δ = 4.17-4.10 (m, 4 H), 3.74-3.53 (m, 4 H), 3.28-3.08 (m, 2 H), 2.80 (s, 3 H), 2.25-2.09 (m, 2 H), 1.29 (t, *J* = 6.8 Hz, 6 H).

Preparation method of intermediate I:

4-[[7-[3-(diethoxyphosphorylmethyl(methyl))amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenylamine:



Step 1: preparation of

N-(diethoxyphosphorylmethyl)-3-[[4-(2-fluoro-4-nitrophenoxy)-6-methoxyl-7-quinolyl]oxyl]-*N*-methyl-propan-1-amine: potassium carbonate (2.9 g or 21.0 mmol) is added to DMF solution (45 mL) of intermediate H (2.0 g or 6.0 mmol) and intermediate A (2.3 g or 9.0 mmol). The resulting mixture is stirred at 110 °C for 2 h. The reaction mixture is diluted with 100 ml of ethyl acetate, inorganic salt is removed by filtering, the filtrate is concentrated, and the residue is purified by column chromatography (eluent: 1-5% MeOH in DCM) to obtain 1.8 g target compound (yield: 55%). The analytical data of the obtained product are as follows: ¹HNMR (400

MHz, DMSO- d_6): δ = 8.55 (d, J = 5.2 Hz, 1 H), 8.46 (dd, J = 2.4, 10.8 Hz, 1 H), 8.22-8.15 (m, 1 H), 7.62 (t, J = 8.4 Hz, 1 H), 7.43 (d, J = 12.8 Hz, 2 H), 6.76 (d, J = 5.2 Hz, 1 H), 4.20 (t, J = 8.4 Hz, 2 H), 4.01-3.95 (m, 4 H), 3.91 (s, 3 H), 2.82 (d, J = 10.8 Hz, 2 H), 2.64 (t, J = 6.8 Hz, 2 H), 2.34 (s, 3 H), 1.96-1.93 (m, 3 H), 1.21 (t, J = 4.4 Hz, 6 H).


Step 2: preparation of

4-[[7-[3-(diethoxyphosphorylmethyl(methyl))amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenylamine:

N-(diethoxyphosphorylmethyl)-3-[[4-(2-fluoro-4-nitrophenoxy)-6-methoxyl-7-quinolyl]oxyl]-*N*-methyl-propan-1-amine (1.8 g or 3.3 mmol) obtained from Step 1 is dissolved in 170 ml of THF, 5 g of Raney nickel is added thereto, the resulting mixture is stirred at 30 °C under 30 psi of hydrogen pressure for 2 h. Finally the reaction mixture is filtered, and the filtrate is concentrated under reduced pressure to obtain 1.6 g of target compound (yield: 97%). The analytical data of the obtained product are as follows: mass spectrum m/z : 522.10 [M+H].

Preparation method of intermediate J:


4-[[6-[3-(diethoxyphosphorylmethyl)(methyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenylamine:



Starting from intermediate H and intermediate B, intermediate J is prepared in the same manner as the method of preparing intermediate I. The analytical data of the obtained product are as follows: mass spectrum m/z : 522.12 [M+H].

Preparation method of intermediate K:

*N*1-[3-fluoro-4-[[6-methoxyl-7-(4-piperidinylmethoxyl)-4-quinolyl]oxyl]phenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide dihydrochloride:




Step 1: preparation of *tert*-butyl 4-[[4-[2-fluoro-4-[[1-[(4-fluorophenyl)aminocarbonyl]cyclopropanecarbonyl]amino]phenoxy]-6-methoxy-7-quinolyl]oxymethyl]piperidine-1-carboxylate: a mixture of intermediate C (1.5 g or 3.0 mmol), intermediate D (1.7 g or 7.6 mmol), DIPEA (1.55 g or 12.0 mmol), HATU (2.3 g or 6.0 mmol), DMAP (0.183 g or 1.5 mmol) in DMF (60 mL) is stirred at 30-40 °C overnight. The mixture is concentrated under reduced pressure, and the residue is purified by column chromatography (eluent: 1-5% MeOH in DCM) to obtain 1.9 g target compound (yield: 90%). The analytical data of the obtained product are as follows: mass spectrum *m/z*: 703.30 [M+H].

Step 2: preparation of *tert*-butyl *N*1-[3-fluoro-4-[[6-methoxy-7-(4-piperidinylmethoxy)-4-quinolyl]oxyl]phenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide dihydrochloride: at 0 °C, saturated EtOAc solution (40 mL) of HCl is added to a solution of *tert*-butyl 4-[[4-[2-fluoro-4-[[1-[(4-fluorophenyl)aminocarbonyl]cyclopropanecarbonyl]amino]phenoxy]-6-methoxy-7-quinolyl]oxymethyl]piperidine-1-carboxylate (1.6 g or 2.3 mmol) obtained from Step 1 in EtOAc (40 mL), the resulting mixture is stirred at room temperature for 4 h. The mixture is filtered to collect the precipitate followed by washing with EtOAc to give 1.0 g of target compound (yield: 69%). The analytical data of the obtained product are as follows: mass spectrum *m/z*: 603.10 [M+H].

Example 1

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention, *N*1-[4-[[7-[3-(benzyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxy-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




A mixture of intermediate F (3.3 g or 5.5 mmol), intermediate D (3.1 g or 13.8 mmol), DIPEA (3.8 g or 22.1 mmol), HATU (4.2 g or 11.0 mmol), DMAP (0.337 g or 2.8 mmol) dissolved in DMF (150 mL) is stirred at 30-40 °C overnight. The mixture is concentrated under reduced pressure, and the residue is purified by column chromatography (eluent: 1-4% MeOH in DCM) to obtain 3.0g of the compound according to the invention (yield: 68%). The analytical data of the obtained product are as follows: mass spectrum m/z : 803.10 [M+H].

Example 2

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1-[4-[[6-[3-(benzyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




Using intermediate G and intermediate D, the above compound is prepared in the same manner as Example 1. The analytical data of the obtained product are as follows: mass spectrum m/z : 803.13 [M+H].

Example 3

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




*N*1-[4-[[7-[3-(benzyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxy]-4-quinolyl]oxy]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide obtained from Example 1 (1.0 g or 1.2 mmol) is dissolved in EtOH/THF (1:1, 250 mL), Pd(OH)₂ (1 g) is added thereto, and the resulting mixture is stirred at 45 °C under 50 psi of hydrogen pressure for 18 h. Finally, the reaction mixture is filtered, the filtrate is concentrated under reduced pressure, and the residue is purified by column chromatography (eluent: 1-4% MeOH in DCM) to give 0.368 g of the compound according to the invention (yield: 43%). The analytical data of the obtained product are as follows: mass spectrum *m/z*: 713.20 [M+H].

Example 4

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1-[4-[[6-[3-(diethoxyphosphorylmethylamino)propoxyl]-7-methoxy]-4-quinolyl]oxy]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




Using

*N*1-[4-[[6-[3-(benzyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxy]-4-quinolyl]oxy]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide obtained from Example 2, the above compound is prepared in the same manner as Example 3. The analytical data of the obtained product are as follows: mass spectrum *m/z*: 713.21 [M+H].

Example 5

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1-[4-[[7-[3-(acetyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




*N*1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide obtained from Example 3 (1.0 g or 1.4 mmol) and DIPEA (0.544 g or 4.2 mmol) are dissolved in DCM (50 mL), acetic anhydride (0.428 g or 4.2 mmol) is added thereto, and the resulting mixture is stirred at room temperature for 2 h. The reaction mixture is diluted with DCM, followed by washing with brine, and then concentrating, and the residue is purified by column chromatography (eluent: 1-3% MeOH in DCM) to give 0.856 g of the compound according to the invention (yield: 81%). The analytical data of the obtained product are as follows: mass spectrum *m/z*: 755.18 [*M*+*H*].

Example 6

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1-[4-[[6-[3-(acetyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




Using

*N*1-[4-[[6-[3-(diethoxyphosphorylmethylamino)propoxyl]-7-methoxyl-4-quinolyl]oxy]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide obtained from Example 4, the above compound is synthesized in the same manner as Example 5. The analytical data of the obtained product are as follows: mass spectrum *m/z*: 755.18 [M+H].

Example 7

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(methyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxy]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




A mixture of intermediate I (0.8 g or 1.5 mmol), intermediate D (0.856 g or 7.6 mmol), DIPEA (1.1 mL or 6.1 mmol), HATU (1.2 g or 3.1 mmol), DMAP (0.093 g or 0.8 mmol) in DMF (60 mL) is stirred at 30-40 °C overnight. The mixture is concentrated under reduced pressure, and the residue is purified by column chromatography (eluent: 1-5% MeOH in DCM) to give 0.352g of the compound according to the invention (yield: 31%). The analytical data of the obtained product are as follows: ¹HNMR (400 MHz, DMSO-*d*₆): δ = 10.37 (s, 1 H), 9.99 (s, 1 H), 8.46 (d, *J* = 5.2 Hz, 1 H), 7.90 (dd, *J* = 2.4, 13.2 Hz, 1 H), 7.64-7.60 (m, 2 H), 7.64-7.60 (m, 2 H), 7.51-7.49 (m, 2 H), 7.16-7.11 (m, 2 H), 6.41 (t, *J* = 4.4 Hz, 1 H), 4.19 (t, *J* = 6.0 Hz, 1 H), 4.04-3.96 (m, 4 H), 3.93 (s, 3 H), 2.90-2.81 (m, 2 H), 2.71-2.61 (m, 2 H), 2.06 (s, 3 H), 1.98-1.90 (m, 2 H), 1.46-1.42 (m, 4 H), 1.21 (t, *J* = 7.2 Hz, 6 H).

Example 8

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1-[4-[[6-[3-(diethoxyphosphorylmethyl)(methyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




Using intermediate J and intermediate D, the above compound is prepared in the same manner as Example 7. The analytical data of the obtained product are as follows: mass spectrum *m/z*: 727.20 [M+H].

Example 9

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1-[4-[[7-[[1-(2-diethoxyphosphorylacetyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




A mixture of intermediate K (0.5 g or 0.8 mmol), 2-diethoxyphosphoryl acetic acid (0.307 g or 1.6 mmol), DIPEA (0.404 g or 3.1 mmol), HATU (0.597 g or 1.6 mmol) in DMF (15 mL) is stirred at room temperature for 2 h. The mixture is concentrated under reduced pressure, and the residue is purified by column chromatography (eluent: 1-5% MeOH in DCM) to give 0.237 g the compound according to the invention (yield: 39%). The analytical data of the obtained product are as follows: ¹HNMR (400 MHz, DMSO-*d*₆): δ = 10.37 (s, 1 H), 9.99 (s, 1 H), 8.45 (d, *J* = 5.2 Hz, 1 H), 7.90 (dd, *J* = 2.0, 13.2 Hz, 1 H), 7.64-7.60 (m, 2 H), 7.50-7.48 (m, 2 H), 7.42-7.37 (m, 2 H), 7.16-7.11 (m, 2 H), 6.40 (dd, *J* = 1.2, 5.2 Hz, 1 H),

4.45-4.35 (m, 1 H), 4.05-3.98 (m, 7 H), 3.93 (s, 3 H), 3.20-3.00 (m, 3 H), 2.65-2.55 (m, 1 H), 2.20-2.00 (m, 1 H), 1.88-1.78 (m, 2 H), 1.45-1.44 (m, 4 H), 1.42-1.28 (m, 1 H), 1.23 (t, $J = 6.8$ Hz, 6 H), 1.18-1.08 (m, 1 H).

Example 10

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1-[4-[[7-[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




0.11 mL of 37% aqueous formaldehyde solution is added to the suspension of intermediate K (0.8 g or 1.3 mmol) in dioxane (10 mL), the resulting mixture is slightly heated until it becomes a solution. The solution is cooled to room temperature, 0.21 mL of diethyl phosphate is added thereto with stirring at room temperature for 30 min, and then stirring at 90-100 °C for 3 h, and the reaction is completed. The solvent is removed under reduced pressure, and the residue is purified by column chromatography (eluent: 1-4% MeOH in DCM) to give 0.354 g of the compound according to the invention (yield: 35%). The analytical data of the obtained product are as follows: ^1H NMR (400 MHz, DMSO- d_6): $\delta = 10.37$ (s, 1 H), 10.00 (s, 1 H), 8.45 (d, $J = 5.2$ Hz, 1 H), 7.90 (dd, $J = 2.0, 13.2$ Hz, 1 H), 7.64-7.60 (m, 2 H), 7.51-7.49 (m, 2 H), 7.42-7.36 (m, 2 H), 7.16-7.11 (m, 2 H), 6.39 (d, $J = 5.2$ Hz, 1 H), 4.05-3.97 (m, 6 H), 3.93 (s, 3 H), 3.06-3.03 (m, 2 H), 2.76-2.74 (m, 2 H), 2.15-2.09 (m, 2 H), 1.88-1.74 (m, 3 H), 1.45-1.33 (m, 6 H), 1.24 (t, $J = 6.8$ Hz, 6 H).

Example 11

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1'-[4-[[7-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




*N*1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide obtained from Example 3 (71.2 mg or 0.1 mmol) is dissolved in DCM (5 mL), then Me₃Si-NCO (23.0 mg or 0.20 mmol) and diisopropylethylamine (38.5 mg or 0.3 mmol) are added thereto, the resulting mixture is stirred at room temperature for 16 h, the solvent is removed under reduced pressure, and the crude product is purified by silica gel column chromatography (eluent: 1-4% MeOH in DCM) to give 57.5 mg of the compound according to the invention (yield: 76%). The analytical data of the obtained product are as follows: mass spectrum *m/z*: 756.12 [M+H].

Example 12

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

*N*1'-[4-[[6-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




Using

*N*1-[4-[[6-[3-(diethoxyphosphorylmethylamino)propoxyl]-7-methoxyl-4-quinolyl]oxy

l]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide obtained from Example 4, the above compound is prepared in the same manner as Example 11. The analytical data of the obtained product are as follows: mass spectrum m/z : 756.18 [M+H].

Example 13


Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,
N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethylaminocarbonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:



N1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide obtained from Example 3 (71.2 mg or 0.1 mmol) is dissolved in DCM (5 mL), then EtNCO (14.2 mg or 0.20 mmol) and diisopropylethylamine (38.5 mg or 0.3 mmol) are added thereto, the resulting mixture is stirred at room temperature for 16 h, the solvent is removed under reduced pressure, and the crude product is purified by silica gel column chromatography (eluent: 1-4% MeOH in DCM) to give 64.3 mg the compound according to the invention (yield: 82%). The analytical data of the obtained product are as follows: mass spectrum m/z : 784.22 [M+H].

Example 14

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,
N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(propionyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:




N1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N1'*-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (71.2 mg or 0.1 mmol) obtained from Example 3 is dissolved in DCM (5 mL), then propionic acid (14.8 mg or 0.20 mmol), EDC.HCl (38.4 mg or 0.2 mmol) and diisopropylethylamine (64.2 mg or 0.5 mmol) are added thereto, the resulting mixture is stirred at room temperature for 16 h, the solvent is removed under reduced pressure, and the crude product is purified by silica gel column chromatography (eluent: 1-4% MeOH in DCM) to give 50.7 mg the compound according to the invention (yield: 66%). The analytical data of the obtained product are as follows: mass spectrum *m/z*: 769.15 [M+H].

Example 15

Preparation of one of the phosphorus-containing group-substituted quinolines according to the present invention,

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(methylsulfonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N1*-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide:



N1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N1'*-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide

obtained from Example 3 (71.2 mg or 0.1 mmol) is dissolved in DCM (5 mL), and then methanesulfonyl chloride (22.9 mg or 0.20 mmol), EDC.HCl (38.4 mg or 0.2 mmol) and diisopropylethylamine (64.2 mg or 0.5 mmol) are added thereto, the resulting mixture is stirred at room temperature for 16 h, the solvent is removed under reduced pressure, and the crude product is purified by silica gel column chromatography (eluent: 1-4% MeOH in DCM) to give 44.3 mg the compound according to the invention (yield: 56%). The analytical data of the obtained product are as follows: mass spectrum *m/z*: 791.28 [M+H].

Example 16

The Screening Test for Inhibiting the Kinase:

In this example, the compound prepared in Example 10, *N*1-[4-[[7-[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide is tested for the inhibitory effect in the 442-kinase panel of KinomeScan™ (www.kinomescan.com) from DiscoverX Corporation (4215 Sorrento Valley Blvd, San Diego, CA 92121) at 1000 nM of the compound concentration, the results are shown in the following Table 5:

For the detailed information of the screening technique, see Fabian, M. A. et al, *Nat., Biotechnol.* **2005**, 23, 329 and Karaman, M. W. et al, *Nat., Biotechnol.* **2008**, 26, 127. It can be seen from the following representative results that, the compound according to the present invention has significant inhibitory effect on kinases including c-Met, KDR or VEGFR2, RET, PDGFR-β, c-KIT, Flt3, MEK5, DDR1, LOK, CSF1R, EPHA7, EPHA8, EPHB6, MKNK2, BLK, HIPK4, HCK, Flt4 and the mutants thereof.

Table 5

Kinase Target	Example 10	Kinase Target	Example 10
Ambit Gene Symbol	%Ctrl@1000nM	Ambit Gene Symbol	%Ctrl@1000nM
AAK1	91	MAPKAPK2	100
ABL1(E255K)-phosphorylated	63	MAPKAPK5	100
ABL1(F317I)-	42	MARK1	100

nonphosphorylated			
ABL1(F317I)-phosphorylated	70	MARK2	100
ABL1(F317L)- nonphosphorylated	10	MARK3	100
ABL1(F317L)- phosphorylated	69	MARK4	100
ABL1(H396P)- nonphosphorylated	1.9	MAST1	100
ABL1(H396P)- phosphorylated	63	MEK1	24
ABL1(M351T)- phosphorylated	42	MEK2	33
ABL1(Q252H)- nonphosphorylated	2.4	MEK3	100
ABL1(Q252H)- phosphorylated	84	MEK4	100
ABL1(T315I)- nonphosphorylated	3.8	MEK5	0.25
ABL1(T315I)-phosphorylated	75	MEK6	87
ABL1(Y253F)- phosphorylated	32	MELK	29
ABL1-nonphosphorylated	1.8	MERTK	2.1
ABL1-phosphorylated	33	MET	7.7
ABL2	28	MET(M1250T)	3
ACVR1	100	MET(Y1235D)	4.4
ACVR1B	100	MINK	100
ACVR2A	100	MKK7	100
ACVR2B	100	MKNK1	45
ACVRL1	100	MKNK2	0.05
ADCK3	67	MLCK	100
ADCK4	100	MLK1	83
AKT1	100	MLK2	78
AKT2	100	MLK3	63
AKT3	100	MRCKA	100

ALK	15	MRCKB	100
AMPK-alpha1	100	MST1	100
AMPK-alpha2	100	MST1R	4
ANKK1	100	MST2	100
ARK5	100	MST3	100
ASK1	100	MST4	80
ASK2	100	MTOR	100
AURKA	100	MUSK	13
AURKB	74	MYLK	100
AURKC	66	MYLK2	100
AXL	8	MYLK4	100
BIKE	100	MYO3A	93
BLK	0.5	MYO3B	100
BMPR1A	100	NDR1	100
BMPR1B	100	NDR2	82
BMPR2	100	NEK1	100
BMX	92	NEK11	100
BRAF	100	NEK2	100
BRAF(V600E)	100	NEK3	58
BRK	3.0	NEK4	100
BRSK1	100	NEK5	97
BRSK2	100	NEK6	100
BTK	51	NEK7	100
CAMK1	78	NEK9	66
CAMK1D	96	NIM1	100
CAMK1G	88	NLK	100
CAMK2A	100	CSR1	96
CAMK2B	100	P38-alpha	53
CAMK2D	94	P38-beta	87
CAMK2G	100	P38-delta	53
CAMK4	100	P38-gamma	100
CAMKK1	32	PAK1	37
CAMKK2	81	PAK2	54
CASK	100	PAK3	32

CDC2L1	100	PAK4	100
CDC2L2	100	PAK6	87
CDC2L5	100	PKA7	100
CDK11	67	PCTK1	100
CDK2	100	PCTK2	100
CDK3	100	PCTK3	100
CDK4-cyclinD1	100	PDGFRA	1.8
CDK4-cyclinD3	100	PDGFRB	0
CDK5	100	PDPK1	100
CDK7	15	PFCDPK1 (P.falciparum)	84
CDK8	76	PFPK5 (P.falciparum)	100
CDK9	100	PFTAIRE2	87
CDKL1	100	PFTK	100
CDKL2	100	PHKG1	100
CDKL3	100	PHKG2	96
CDKL5	100	PIK3C2B	100
CHEK1	100	PIK3C2G	98
CHEK2	35	PIK3CA	100
CIT	100	PIK3CA(C420R)	100
CLK1	89	PIK3CA(E542K)	100
CLK2	78	PIK3CA(E545A)	74
CLK3	100	PIK3CA(E545K)	100
CLK4	46	PIK3CA (H1047L)	100
CSF1R	0.25	PIK3CA (H1047Y)	63
CSK	14	PIK3CA(18DDL)	100
CSNK1A1	100	PIK3CA(M1043I)	100
CSNK1A1L	100	PIK3CA(Q546K)	100
CSNK1D	100	PIK3CB	100
CSNK1E	100	PIK3CD	100
CSNK1G1	100	PIK3CG	100
CSNK1G2	100	PIK4CB	100
CSNK1G3	100	PIM1	96

CSNK2A1	100	PIM2	67
CSNK2A2	100	PIM3	67
CTK	100	PIP5K1A	100
DAPK1	100	PIP5K1C	100
DAPK2	100	PIP5K2B	100
DAPK3	100	PIP5K2C	38
DCAMKL1	91	PAKC-alpha	100
DCAMKL2	100	PAKC-beta	100
DCAMKL3	100	PKMYT1	100
DDR1	0	PKN1	100
DDR2	7.8	PKN2	2.6
DLK	52	PKNB (M.tuberculosis)	96
DMPK	85	PLK1	100
DMPK2	76	PLK2	83
DRAK1	100	PLK3	100
DRAK2	100	PLK4	7.4
DYRK1A	100	PRKCD	100
DYRK1B	86	PRKCE	100
DYRK2	100	PRKCH	100
EGFR	100	PRKCI	100
EGFR(E746-A750del)	92	PRKCQ	100
EGFR(G719C)	75	PRKD1	100
EGFR(G719S)	81	PRKD2	100
EGFR(L747-E749del,A750P)	67	PRKD3	88
EGFR(L747-S752del,P753S)	67	PRKG1	100
EGFR(L747-T751del,Sins)	72	PRKG2	100
EGFR(L858R)	59	PRKR	100
EGFR(L858R,T790M)	100	PRKX	81
EGFR(L861Q)	84	PRP4	76
EGFR(S752-1759del)	82	PYK2	19
EGFR(T790M)	100	QSK	100
EIF2AK1	100	RAF1	100
EPHA1	4.6	RET	0.1

EPHA2	4.8	RET (M918T)	0.05
EPHA3	6	RET (V804L)	0.35
EPHA4	4	RET (V804M)	0.6
EPHA5	3.5	RIOK1	100
EPHA6	4.8	RIOK2	100
EPHA7	0.8	RIOK3	100
EPHA8	0.15	RIPK1	99
EPHB1	4.6	RIPK2	30
EPHB2	2.8	RIPK4	89
EPHB3	36	RIPK5	82
EPHB4	12	ROCK1	100
EPHB6	0.9	ROCK2	100
ERBB2	96	ROS1	29
ERBB3	98	RPS6KA4((kin.Dom.1-N-terminal))	76
ERBB4	100	RPS6KA4((kin.Dom.2-C-terminal))	100
ERK1	99	RPS6KA5((kin.Dom.1-N-terminal))	100
ERK2	100	RPS6KA5((kin.Dom.2-C-terminal))	83
ERK3	100	RSK1(Kin.Dom.1-N-terminal)	59
ERK4	100	RSK1(Kin.Dom.2-C-terminal)	100
ERK5	100	RSK2(Kin.Dom.1-N-terminal)	100
ERK8	95	RSK3(Kin.Dom.1-N-terminal)	100
ERN1	81	RSK3(Kin.Dom.2-C-terminal)	100
FAK	51	RSK4(Kin.Dom.1-N-terminal)	77
FER	50	RSK4(Kin.Dom.2-C-terminal)	91

		-terminal)	
FES	44	S6K1	89
FGFR1	100	SBK1	100
FGFR	76	SgK110	100
FGFR3	69	SGK3	75
FGFR3(G697C)	100	SIK	60
FGFR4	100	SIK2	100
FGR	35	SLK	7.4
FLT1	7.3	SNARK	100
FLT3	2.3	SNRK	100
FLT3(D835H)	0.6	SRC	3.2
FLT3(D835Y)	2.4	SRMS	8.9
FLT3(ITD)	2.8	SRPK1	70
FLT3(K663Q)	0.55	SRPK2	100
FLT3(N841L)	0	SRPK3	100
FLT3(R834Q)	16	STK16	77
FLT4	0.2	STK33	46
FRK	8.8	STK35	82
FYN	48	STK36	100
GAK	100	STK39	100
GCN2(kin.dom.2.S808G)	100	SYK	58
GRK1	100	TAK1	75
GRK4	78	TAOK1	100
GRK7	100	TAOK2	70
GSK3A	100	TAOK3	100
GSK3B	100	TBK1	79
HCK	0.35	TEC	100
HIPK1	92	TESK1	39
HIPK2	100	TGFBR1	100
HIPK3	66	TGFBR2	100
HIPK4	0.15	TIE1	4
HPK1	40	TIE2	2.4
HUNK	100	TLK1	88
ICK	95	TLK2	100

IGF1R	94	TNIK	70
IKK-alpha	100	TNK1	30
IKK-beta	100	TNK2	94
IKK-epsilon	94	TNNI3K	94
INSR	54	TRAK	1.8
INSRR	65	TRKB	4
IRAK1	100	TRKC	2.2
IRAK3	33	TRPM6	86
IRAK4	100	TSSK1B	100
ITK	34	TTK	84
JAK1(JH1domain-catalytic)	93	TXK	62
JAK1(JH2domain-pseudokinase)	100	TYK2(JH1domain-catalytic)	92
JAK2(JH1domain-catalytic)	100	TYK2(JH1domain-pseudokinase)	100
JAK3(JH1domain-catalytic)	100	TYRO3	41
JNK1	80	ULK1	100
JNK2	93	ULK2	100
JNK3	95	ULK3	82
KIT	0.2	VEGFR2	3.9
KIT(A829P)	64	VRK2	100
KIT(D816H)	63	WEE1	100
KIT(D816V)	18	WEE2	100
KIT(L576P)	0	YANK1	100
KIT(V559D)	0.15	YANK2	100
KIT(V559D,T670I)	1.4	YANK3	100
KIT(V559D,V654A)	4.1	YES	20
LATS1	100	YSK1	57
LATS2	96	YSK4	1.7
LCK	2.2	ZAK	36
LIMK1	92	ZAP7C	81
LIMK2	100		
LKB1	100		
LOK	0		

LRRK2	100		
LRRK2(G2019S)	100		
LTK	26		
LYN	18		
LZK	47		
MAK	100		
MAP3K1	80		
MAP3K15	100		
MAP3K2	100		
MAP3K3	100		
MAP3K4	100		
MAP4K2	100		
MAP4K3	20		
MAP4K4	100		
MAP4K5	2.8		

Example 17

Inhibitory activity on the kinases in tumor cells: In this example, the compound prepared in Example 10, *N*1-[4-[[7-[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide is tested in tumor cells by ProQinase GmbH corporation (Breisacher Str. 117, D-79106 Freiburg, Germany. www.proqinase.com) for the inhibitory activity on the following kinases: c-KIT, c-MET, VEGFR2 and PDGFR- β . The test includes the following steps:

Firstly, the compound according to present invention is dissolved in dimethyl sulfoxide (DMSO) to prepare a 10 mM stock solution, which is diluted serially according to the concentration range shown in Table 6:

Table 6

compound	concentration of the stock solution [M]	testing range [M]	testing range [M]	testing range [M]	testing range [M]
----------	---	----------------------	----------------------	----------------------	-------------------------

		c-KIT	c-MET	VEGFR2	PDGFR- β
the compound prepared in Example 10	1.0×10^{-2}	1.0×10^{-5} - 1.0×10^{-9}	1.0×10^{-5} - 1.0×10^{-9}	1.0×10^{-5} - 1.0×10^{-9}	1.0×10^{-5} - 1.0×10^{-9}

Cell Culture

c-KIT kinase: in the test experiment of KIT kinase phosphorylation in the cells, human acute megakaryocytic leukemia cell line M07e having endogenous high expression wild type c-KIT is used. When these cells are stimulated with human stem cell factor (SCF), the tyrosine of KIT receptor kinases is self-phosphorylated. M07e cells are inoculated in RPMI containing 20% fetal calf serum (FCS) and 10 ng/ml GM-CSF in a multi-well cell culture dish. The compound of the invention is added to the cells after overnight starvation of serum and growth factor, then the cells are cultured in a medium without serum.

c-MET kinase: in the test experiment of c-MET kinase phosphorylation in the cells, human gastric cancer cell line MKN45 having endogenous high expression wild type c-MET (wild-type c-MET) is used. Such high expression of c-MET causes the tyrosine of c-MET receptor kinases constantly self-phosphorylated without ligands. MKN45 cells are inoculated in DMEM culture solution containing 10% FCS in a multi-well cell culture dish. The compound of the invention is added to the cells after serum overnight starvation, then the cells are cultured in a medium without serum.

VEGFR2 kinase: in the test experiment of VEGFR2 kinase phosphorylation in the cells, a spontaneous immortalized human umbilical vein endothelial cell line HUE having endogenous high expression vascular endothelial growth factor-R2 (VEGFR2) is used. When these cells are simulated with human vascular endothelial growth factor-A (VEGF-A), it causes the tyrosine of VEGFR2 receptor kinases self-phosphorylated. HUE cells are inoculated in endothelial cell growth medium (ECGM) containing 10% FCS in a multi-well cell culture dish. The compound of the invention is added to the cells after serum overnight starvation in endothelial cell

growth medium (ECGM) containing 10% FCS, then the cells cultured in a ECGM culture solution without serum.

PDGFR- β kinase: in the test experiment of PDGFR-beta kinase phosphorylation in the cells, mouse embryo fibroblast cell line NIH3T3 having endogenous high expression wild type PDGFR- β (wild-type PDGFR- β) is used. When these cells are simulated with mouse platelet-derived growth factor BB (PDGF-BB), it causes the tyrosine of PDGFR-beta receptor kinases self-phosphorylated. NIH3T3 cells are inoculated in DMEM culture solution containing 10% FCS in a multi-well cell culture dish. The compound of the invention is added to the cells after serum overnight starvation, then the cells are cultured in a medium without serum.

Test Method

The diluted stock solution sample of the compound according to the invention is added to cell culture solution in a ratio of 1:100 until a final DMSO concentration of 1% is reached. After culturing at 37°C for 90 min, the cells are stimulated according to Table 7 (quantitative method: sandwich ELISA: Substrate phosphorylation is quantitated through 96-well sandwich ELISA method by using capture antibody of specific substrate and detection antibody of dephosphorylated tyrosine. IC₅₀ values are determined by using GraphPad Prism 5.01 software):

Table 7

target	ligand	concentration	time
c-KIT	SCF	100 ng/mL	3 min
c-MET	without stimulation		
PDGFR- β	PDGF-BB	100 ng/mL	3 min
VEGFR2	VEGF-A	100 ng/mL	3 min

Test Results

By drawing titration curve of N1-[4-[[[7-[[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide inhibiting c-KIT, c-MET, VEGFR2 and PDGFR- β in the cells, IC₅₀ values of the

compound inhibiting the above four kinases are obtained, shown in Table 8:

Table 8

compound	IC ₅₀ values in the cell*			
the compound	c-KIT	c-MET	VEGFR2	PDGFR-β
prepared in	27 nM	17 nM	0.43 nM	13 nM
Example 10				

* mean value of two testing results

It can be seen from the above data that, the compound according to the present invention has significant inhibitory effect on several kinases, i.e., c-Met, c-KIT, VEGFR2 and PDGFR-β in tumor cells, their IC₅₀ values are in the range of 0.43-27 nM.

Example 18

Anti-tumor experiments: in this example, *N*1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxy]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide obtained from Example 3 is used in the anti-tumor experiments in nude mice animal model of xenograft GTL-16 human gastric cancer cells, the methods and results are as follows:

Materials

7-8 weeks old SPF level female BALB/c-nu/nu mice with 16-20 g of body weights are purchased from Experimental Animal Center of Guangdong Province, inspection certificate number: NO:0072659. GTL-16 gastric cancer cells are purchased from the Cell Source Center of Shanghai Institutes for Biological Sciences. RPM I-1640 cell culture medium, fetal calf serum (FBS) and digestive juice of trypsin are from Gibco Corporation. All the antibiotics are purchased from Sigma Company.

Methods

Firstly culturing the GTL-16 gastric cancer cells: GTL-16 cell line is inoculated in RPM I-1640 culture solution containing 10% FBS, 100 U/ml penicillin, 100 U/ml streptomycin, and cultured at 37 °C, 5% CO₂, 100% humidity in a carbon dioxide cell

incubator, the cells enter into logarithmic growth phase after 24 hours, after 48 hours of inoculation, the cells grow to cover the bottom of the culture bottle. Then the GTL-16 cells covering 80% bottom of the culture bottle are digested, 5 minutes after centrifuging at 1000 r/min, the cells are diluted to 2×10^7 /ml and implanted subcutaneously in the right anterior axillary fossa with 0.1 ml per mouse. After 12 days after tumor cells were implanted, the tumor-bearing nude mice are weighted, the sizes of the tumors are measured and the mice with tumors in a size range of 40-160 mm³ are randomly divided into vehicle group (vehicle, 0.2% aqueous methylsulfonic acid solution) and treatment group using the compound prepared in Example 3, N1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (which is dissolved in 0.2% aqueous methylsulfonic acid solution with a concentration of 5 mg/mL), 8 mice per group; thereafter the nude mice are weighted each day and administrated orally in a mount of 0.1 mL/10 g body weight by gavage method (50 mg/kg qd *via* p.o), the sizes of tumors on each animal are measured every other days, after 14 days, the tumor-bearing mice are sacrificed with cervical spine articulation, and the tumors are taken out to weight. The calculation method of Tumor Growth Inhibition (TGI): the experimental data are expressed as mean $\pm$ SD; $TGI = [(V_{\text{vehicle group}} - V_{\text{treatment group}}) \div V_{\text{vehicle group}}] \times 100\%$. $V_{\text{vehicle group}}$ means tumor volume of vehicle group mice; $V_{\text{treatment group}}$ means tumor volume of treatment group mice.

Results

The tumor inhibitory effect of the compound according to present invention prepared in Example 3 on mice tumor in the treatment group: vehicle group and treatment group are respectively fed with vehicle and drug solution, after 14 days the average tumor volumes of the vehicle group and treatment group animals are 377.1 ± 244.5 mm³ (n=7) and 57.1 ± 61.2 (n=8) in respective; thus TGI of the treatment group is 85%. As compared with the initial tumor volumes, the tumor sizes of the treatment group are significantly reduced. The comparisons of tumor tissues are shown in Fig. 1, and the comparisons of the experimental curves are shown in Fig. 2.

Body weight changes for mice in vehicle group and treatment group are shown

in Fig. 3: after treating for 14 days, the body weight of mice in the treatment group using the compound of Example 3 had little change (<5%).

It can be seen from the results of the above anti-tumor experiments that, in nude mice xenograft animal models, the representative compound of the present invention exhibits significant anti-tumor effect, administrating orally in a dose of 50 mg/kg, once daily, after 14 days TGI reaches 85%. Body weight of the animals in treatment group substantively do not reduce, this demonstrates that the drug has no significant toxicity.

Example 19:

Drug Ingredients and Formulation

Tablet (mg/tablet)

The compound prepared in Example 1: 100;

Lactose, Ph EUR: 182.75;

Sodium carboxymethylcellulose: 12.0;

Corn starch slurry (5 w/v%): 2.25;

Magnesium stearate: 3.0;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 20:

Drug Ingredients and Formulation

Tablet (mg/tablet)

The compound prepared in Example 5: 100; the contents of other materials are the same as those in Example 19.

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 21:

Drug Ingredients and Formulation

Tablet (mg/tablet)

The compound prepared in Example 1: 50;

Lactose, Ph EUR: 223.75;

Sodium carboxymethylcellulose: 6.0;

Corn starch: 15.0;

Polyvinylpyrrolidone (5 w/v%): 2.25;

Magnesium stearate: 3.0;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 22:

Drug Ingredients and Formulation

Tablet (mg/tablet)

The compound prepared in Example 3: 50; the contents of other materials are the same as those in Example 21;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 23:

Drug Ingredients and Formulation

Tablet (mg/tablet)

The compound prepared in Example 7: 1.0;

Lactose, Ph EUR: 93.25;

Sodium carboxymethylcellulose: 4.0;

Corn starch slurry (5 w/v%): 0.75;

Magnesium stearate: 76;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 24:

Drug Ingredients and Formulation

Tablet (mg/tablet)

The compound prepared in Example 5: 1.0; the contents of other materials are the same as those in Example 23;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 25:

Drug Ingredients and Formulation

Capsule (mg/capsule)

The compound prepared in Example 7: 10.0;

Lactose, Ph EUR: 488.5;

Magnesium: 1.5;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 26:

Drug Ingredients and Formulation

Capsule (mg/capsule)

The compound prepared in Example 2: 10.0; the contents of other materials are the same as those in Example 25;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 27:

Drug Ingredients and Formulation

Injection (50 mg/ml)

The compound prepared in Example 6: 5%;

1 M Sodium hydroxide solution: 15%;

0.1 M hydrochloric acid solution (pH is adjusted to 7.6);

Polyenthylene glycol 400: 5%;

Adjusting to 100% with water for injection;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 28:

Drug Ingredients and Formulation

Injection (50 mg/ml)

The compound prepared in Example 12: 5%; the contents of other materials are the same as those in Example 27; finally adjusting to 100% with water for injection;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated

diseases.

Example 29:

Drug Ingredients and Formulation

Injection (10 mg/ml)

The compound prepared in Example 11: 1%;

Disodium hydrogen phosphate BP: 3.6%;

0.1M Sodium hydroxide solution: 15%;

Adjusting to 100% with water for injection;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 30:

Drug Ingredients and Formulation

Injection (10 mg/ml)

The compound prepared in Example 9: 1%; the contents of other materials are the same as those in Example 29; adjusting to 100% with water for injection;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 31:

Drug Ingredients and Formulation

Injection (1 mg/ml) (pH is adjusted to 6)

The compound prepared in Example 6: 0.1%;

Disodium hydrogen phosphate BP: 2.26%;

Citric acid: 0.38%;

Polyethylene glycol 400: 3.5%;

Water for injection (for adjusting to 100%);

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 32:

Drug Ingredients and Formulation

Injection (1 mg/ml) (pH is adjusted to 6)

The compound prepared in Example 10: 0.1%; the contents of other materials are

the same as those in Example 31; finally water for injection is used to adjust the content to 100%;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 33:

Drug Ingredients and Formulation

Aerosol (mg/ml)

The compound prepared in Example 1: 10;

Sorbitan oleate: 13.5;

Trichlorofluoromethane: 910.0;

Dichlorodifluoromethane: 490.0;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 34:

Drug Ingredients and Formulation

Aerosol (mg/ml)

The compound prepared in Example 3: 10; the contents of other materials are the same as those in Example 33;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 35:

Drug Ingredients and Formulation

Aerosol (mg/ml)

The compound prepared in Example 4: 0.2;

Sorbitan oleate: 0.27;

Trichlorofluoromethane: 70.0;

Dichlorodifluoromethane: 280.0;

Dichlorotetrafluoroethane: 1094.0;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 36:

Drug Ingredients and Formulation

Aerosol (mg/ml)

The compound prepared in Example 7: 0.2; the contents of other materials are the same as those in Example 35;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 37:

Drug Ingredients and Formulation

Aerosol (mg/ml)

The compound prepared in Example 8: 2.5;

Sorbitan oleate: 3.38;

Trichlorofluoromethane: 67.5;

Dichlorodifluoromethane: 1086.0;

Dichlorotetrafluoroethane: 191.60;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 38:

Drug Ingredients and Formulation

Aerosol (mg/ml)

The compound prepared in Example 11: 2.5; the contents of other materials are the same as those in Example 37;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 39:

Drug Ingredients and Formulation

Aerosol (mg/ml)

The compound prepared in Example 4: 2.5;

Soybean lecithin: 2.7;

Trichlorofluoromethane: 67.5;

Dichlorodifluoromethane: 1086.0;

Dichlorotetrafluoroethane: 191.60;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 40:

Drug Ingredients and Formulation

Aerosol (mg/ml)

The compound prepared in Example 13: 2.5; the contents of other materials are the same as those in Example 39;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 41:

Drug Ingredients and Formulation

Ointment (/ml)

The compound prepared in Example 1: 40 mg;

Ethanol: 300 μ L;

Water: 300 μ L;

1-dodecyl azacycloheptanone: 50 μ L;

Propanediol: to 1ml;

Applicable People

Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

Example 42:

Drug Ingredients and Formulation

Ointment (/ml)


The compound prepared in Example 7: 40 mg; the contents of other materials are the same as those in Example 41;

Applicable People


Applicable to patients suffering from protein kinase abnormal activity-associated diseases.

What is claimed is:

1. A phosphorus-containing group-substituted quinoline, which is represented by formula (I) below:




in formula (I),

Z represents , V¹ and V² are each independently selected from hydrogen, halogen, -OCF₃, -CF₃, -NO₂, -CN, -OH, -NH₂, -NMe₂, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, or C₃₋₆ heterocycloalkoxy;

Either of R and R' represents phosphorus-containing substituent Q, the other is selected from hydrogen, methoxyl, methoxyethoxyl, or phosphorus-containing substituent Q;

Wherein, the phosphorus-containing substituent Q represents



A is absent or represents O, NH, S(=O)_m, or C₁₋₆ alkyl, and A is optionally substituted with G¹;

L is absent or represents C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl, and L is optionally substituted with G²;

J is absent or represents O, NH, S(=O)_m, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl and J is optionally substituted with G³;

X is absent or represents -C(=O)-, -S(O)_m-, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl, and X is optionally substituted with G⁴;

Y is absent or represents C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, or C₃₋₆ heterocycloalkyl, and Y is optionally substituted with G⁵;

R^1 and R^2 are each independently selected from -OH, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_6 aryl, C_{5-6} heteroaryl, C_{3-6} heterocycloalkyl, C_{1-6} alkoxy, C_{3-6} cycloalkoxy, C_6 aryloxy, C_{5-6} heteroaryloxy, or C_{3-6} heterocycloalkoxy, and R^1 and R^2 are each optionally substituted with G^6 ; R^1 and R^2 together with the phosphorus atom to which they are attached may form a C_{3-6} heterocycloalkyl ring, this C_{3-6} heterocycloalkyl ring may further comprise heteroatom(s) selected from O, N, or S(=O)_m;


wherein,

G^1 , G^2 , G^3 , G^4 , G^5 and G^6 are each independently selected from H, -CN, -CF₃, -CO₂H, halogen, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_6 aryl, C_{5-6} heteroaryl, C_{3-6} heterocycloalkyl, R^3O- , R^3R^4N- , $R^3S(=O)_m-$, $R^3R^4NS(=O)_m-$, $R^5C(=O)-$, $R^3R^4NC(=O)-$, $R^3OC(=O)-$, $R^5C(=O)O-$, $R^3R^4NC(=O)O-$, $R^5C(=O)NR^3-$, $R^3R^4NC(=O)NR^6-$, $R^3OC(=O)NR^6-$, $R^3S(=O)_mNR^6-$, $R^3R^4NS(=O)_mNR^6-$, $R^3R^4NC(=NR^7)NR^6-$, $R^3R^4NC(=CHNO_2)NR^6-$, $R^3R^4NC(=N-CN)NR^6-$, $R^3R^4NC(=NR^7)-$, $R^3S(=O)(=NR^7)NR^6-$, or $R^3R^4NS(=O)(=NR^7)-$;

R^3 , R^4 , R^5 , R^6 and R^7 are each independently selected from H, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_6 aryl, C_{5-6} heteroaryl, or C_{3-6} heterocycloalkyl; when R^3 and R^4 are attached to the same nitrogen atom, R^3 and R^4 together with the nitrogen to which they are attached may form a C_{3-6} heterocycloalkyl ring, this C_{3-6} heterocycloalkyl ring may further comprise heteroatom(s) selected from O, N, or S(=O)_m; and R^3 , R^4 , R^5 , R^6 and R^7 may be optionally substituted with halogen, CN, C_{1-6} alkyl, or C_{3-6} cycloalkyl;

$m = 0-2$.

2. The phosphorus-containing group-substituted quinoline according to claim 1, wherein the compound is represented by formula (Ia) below:



in the above formula,

A represents O, NH, or S(=O)_m, and A is optionally substituted with G^1 ;

L represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and L is optionally substituted with G²;


J represents O, NH, or S(=O)_m, and J is optionally substituted with G³;

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and Y is optionally substituted with G⁵;

R¹ and R² are each independently selected from -OH, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, C₆ aryloxy, C₅₋₆ heteroaryloxy, or C₃₋₆ heterocycloalkoxy, and R¹ and R² are each optionally substituted with G⁶;

m = 0-2.

3. The phosphorus-containing group-substituted quinoline according to claim 1, wherein the compound is represented by formula (Ib) below:



in the above formula,

A represents O, NH, or S(=O)_m, and A is optionally substituted with G¹;

L represents C₁₋₆ alkyl, and L is optionally substituted with G²;

J represents C₃₋₆ cycloalkyl, or C₃₋₆ heterocycloalkyl, and J is optionally substituted with G³;


X is absent or represents -C(=O)-, or -S(O)_m-;

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl; and Y is optionally substituted with G⁵;

R¹ and R² are each independently selected from -OH, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, C₆ aryloxy, C₅₋₆ heteroaryloxy, or C₃₋₆ heterocycloalkoxy, and R¹ and R² are each optionally substituted with G⁶;

m = 0-2.

4. The phosphorus-containing group-substituted quinoline according to claim 1, wherein the compound is represented by formula (Ic) below:



in the above formula,

A represents O, NH, or S(=O)_m, and A is optionally substituted with G¹;

L represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and L is optionally substituted with G²;


J represents O, NH, or S(=O)_m, and J is optionally substituted with G³;

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and Y is optionally substituted with G⁵;

R¹ and R² are each independently selected from -OH, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₆ aryl, C₅₋₆ heteroaryl, C₃₋₆ heterocycloalkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkoxy, C₆ aryloxy, C₅₋₆ heteroaryloxy, or C₃₋₆ heterocycloalkoxy, and R¹ and R² are each optionally substituted with G⁶;

m = 0-2.

5. The phosphorus-containing group-substituted quinoline according to claim 1, wherein the compound is represented by formula (Id) below:



in the above formula,

A represents O, NH, or S(=O)_m, and A is optionally substituted with G¹;

L represents C₁₋₆ alkyl, and L is optionally substituted with G²;

J represents C₃₋₆ cycloalkyl, or C₃₋₆ heterocycloalkyl, and J is optionally

substituted with G^3 ;


X is absent or represents $-C(=O)-$, or $-S(O)_m-$;

Y represents C_{1-6} alkyl, or C_{3-6} cycloalkyl, and Y is optionally substituted with G^5 ;

R^1 and R^2 are each independently selected from $-OH$, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_6 aryl, C_{5-6} heteroaryl, C_{3-6} heterocycloalkyl, C_{1-6} alkoxy, C_{3-6} cycloalkoxy, C_6 aryloxy, C_{5-6} heteroaryloxy, or C_{3-6} heterocycloalkoxy, and R^1 and R^2 are each optionally substituted with G^6 ;

$m = 0-2$.

6. The phosphorus-containing group-substituted quinoline according to claim 1, wherein the compound is represented by formula (Ie) below:




in the above formula,

L represents C_{1-6} alkyl, or C_{3-6} cycloalkyl, and L is optionally substituted with G^2 ;

Y represents C_{1-6} alkyl, or C_{3-6} cycloalkyl, and Y is optionally substituted with G^5 .

7. The phosphorus-containing group-substituted quinoline according to claim 1, wherein the compound is represented by formula (If) below:



in the above formula,

Y represents C₁₋₆ alkyl, or C₃₋₆ cycloalkyl, and Y is optionally substituted with G⁵.

8. The phosphorus-containing group-substituted quinoline according to claim 1, wherein the compound is selected from the group consisting of one or more of the following compounds:

*N*1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethylamino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(methyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(n-propyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(methyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(n-propyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(acetyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

uinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(formyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(propionyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(isopropionyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(cyclopropylformyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(acetyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(formyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(propionyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(isopropionyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[7-[3-(cyclopropylformyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethylaminocarbonyl)amino)propoxyl]

-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(methylaminocarbonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(N,N'-dimethylaminocarbonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethylaminocarbonyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(methylaminocarbonyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(N,N'-dimethylaminocarbonyl)amino)ethoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(methylsulfonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1'-[4-[[7-[3-(diethoxyphosphorylmethyl)(ethylsulfonyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethylamino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

N1-[4-[[6-[3-(diethoxyphosphorylmethylamino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-N1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(diethoxyphosphorylmethyl)(methyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(diethoxyphosphorylmethyl)(*n*-propyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(diethoxyphosphorylmethyl)(methyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(diethoxyphosphorylmethyl)(*n*-propyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(acetyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(formyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(propionyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(isopropionyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(cyclopropylformyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(acetyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(formyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(propionyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(isopropionyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[3-(cyclopropylformyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethylaminocarbonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(methylaminocarbonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(*N,N*'-dimethylaminocarbonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(aminocarbonyl(diethoxyphosphorylmethyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

oxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethylaminocarbonyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(methylaminocarbonyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(*N,N'*-dimethylaminocarbonyl)amino)ethoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(methylsulfonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1'-[4-[[6-[3-(diethoxyphosphorylmethyl)(ethylsulfonyl)amino)propoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[[1-(2-diethoxyphosphorylacetyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[6-[[1-(2-diethoxyphosphorylacetyl)-4-piperidinyl]methoxyl]-7-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide.

9. The phosphorus-containing group-substituted quinoline according to claim 1, wherein the compound is selected from the group consisting of one or more of the

following compounds:

*N*1-[4-[[7-[3-(diethoxyphosphorylmethylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[3-(diethoxyphosphoryl(*N*-methyl)methylamino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;


*N*1-[4-[[7-[3-(acetyl(diethoxyphosphorylmethyl)amino)propoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[[1-(2-diethoxyphosphorylacetyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide;

*N*1-[4-[[7-[[1-(diethoxyphosphorylmethyl)-4-piperidinyl]methoxyl]-6-methoxyl-4-quinolyl]oxyl]-3-fluorophenyl]-*N*1'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide.

10. A racemate or enantiomer of the phosphorus-containing group-substituted quinolines according to any of claims 1-9.

11. A process of preparing the phosphorus-containing group-substituted quinolines according to any of claims 1-9, consisting of the steps in the following Scheme 1:

Fc1ccc(NC(=O)C2(C)C2C(=O)Nc3ccc(F)c(Oc4ccccc4)c3)c1

t = 0-4;


PG represents common protecting groups in the organic chemistry field;

80

together with the nitrogen to which they are attached may form a C₃₋₆ heterocycloalkyl ring, this C₃₋₆ heterocycloalkyl ring may further comprise heteroatom(s) selected from O, N, S(=O)_m; and R³, R⁴, R⁵, R⁶ and R⁷ are each optionally substituted with halogen, CN, C₁₋₆ alkyl, or C₃₋₆ cycloalkyl; m = 0-2.

12. A process of preparing the phosphorus-containing group-substituted quinolines according to any of claims 1-9, consisting of the steps in the following


Scheme 2:



wherein PP' can be any one of PG, H and PP, and PG can be converted into H by deprotection.


13. A process of preparing the phosphorus-containing group-substituted quinolines according to any of claims 1-9, consisting of the steps in the following Scheme 3:

Scheme 3




14. A process of preparing the phosphorus-containing group-substituted quinolines according to any of claims 1-9, consisting of the steps in the following Scheme 4:

Scheme 4



15. A process of preparing the phosphorus-containing group-substituted quinolines according to any of claims 1-9, consisting of the steps in the following Scheme 5:



16. Use of a pharmaceutical composition comprising the phosphorus-containing group-substituted quinolines according to any of claims 1-9, wherein the composition is used for treating diseases associated with the abnormal activities of protein kinases.

17. The use according to claim 16, wherein the protein kinase is c-Met, KDR or VEGFR2, RET, PDGFR- β , c-KIT, Flt3, MEK5, DDR1, LOK, CSF1R, EPHA7, EPHA8, EPHB6, MKNK2, BLK, HIPK4, HCK, or Flt4.

18. The use according to claim 16, wherein the protein kinase is RON, ALK (or Anaplastic Lymphoma Kinase), EGF1R, HER2, HER3, HER4, PDGFR- α , c-fms, FLT1, Src, Frk, Btk, CsK, Abl, Fes, Fps, Fak, AcK, Yes, Fyn, Lyn, Lck, Hck, Fgr, Yrk, PDK1, TAK1, Tie-2, Tie-1, YSK4, TRK A, TRK B, TRK C, SLK, PKN2, PLK4, MST1R, MAP4K, or DDR2.

19. The use according to claim 16, wherein the disease is psoriasis, hepatic cirrhosis, diabetes, angiogenesis-mediated diseases, eye diseases, immune system diseases, or cardiovascular diseases.

20. The use according to claim 16, wherein the disease is tumor comprising solid tumor and liquid tumors.

21. The use according to claim 16, wherein the tumor comprises: lung cancer, bone cancer, pancreatic cancer, skin cancer, head and neck cancer, skin or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, anal region cancer, stomach cancer, colon cancer, breast cancer, fallopian tube cancer, endometrial cancer, cervical cancer, vaginal cancer, vulva cancer, Hodgkin's disease, esophageal cancer, intestinal cancer, endocrine system cancer, thyroid cancer, parathyroid cancer, soft tissue sarcoma, urethral cancer, penile cancer, prostate cancer, chronic or acute leukemia, bladder cancer, kidney or ureter cancer, renal cancer, central nervous

system (CNS) tumors, spinal axis tumors, pituitary adenomas, gastrointestinal stromal tumors, colorectal cancer, non-small cell lung cancer, small cell lung cancer, mastocytosis, glioma, sarcoma, lymphoma, or a combination of any thereof.

22. A medicament for treating protein kinase abnormal activity-associated diseases, comprising the compound according to any of claims 1-9 or pharmaceutically acceptable salt, solvate, prodrug thereof, or a racemate or enantiomer of the compound according to any of claims 1-9 or pharmaceutically acceptable salt, solvate, prodrug thereof.

23. The medicament according to claim 22, wherein it further comprises at least one pharmaceutically acceptable carrier.

24. The medicament according to claim 22, wherein it is in a form of the following Formulations:

(1) oral formulation; (2) injectable formulation; (3) anal suppository; (4) nasal inhalation; (5) eye drop; (6) skin patch.