


CYCLOALKYLAMINE DERIVATIVES


The present invention is concerned with novel cycloalkylamine derivatives, their manufacture and their use as medicaments.

In particular, the invention relates to compounds of the formula (I)



5 wherein

R¹ is selected from



R², R³, R⁴, R⁵ and R⁶ are each independently selected from hydrogen, lower alkyl, halogenated lower alkyl or halogen; provided that R², R³, R⁴, R⁵ and R⁶ are not all hydrogen;

R⁷ is lower alkyl;

R⁸ is lower alkyl;

X is >C=O or >SO₂;

R⁹ and R¹¹ are hydrogen or together form a double bond;

R¹⁰ and R¹² are independently selected from hydrogen or lower alkyl;

m is 1 or 2;

n is 0, 1, or 2;

and pharmaceutically acceptable salts thereof for use in therapy.


The enzyme dipeptidyl peptidase IV (EC.3.4.14.5, abbreviated in the following as DPP-IV) is involved in the regulation of the activities of several hormones. In particular DPP-IV is degrading efficiently and rapidly glucagon like peptide 1 (GLP-1), which is one of the most potent stimulator of insulin production and secretion. Inhibiting DPP-IV would potentiate the effect of endogenous GLP-1, and lead to higher plasma insulin concentrations. In patients suffering from impaired glucose tolerance and type 2 diabetes mellitus, higher plasma insulin concentration would moderate the dangerous hyperglycaemia and accordingly reduce the risk of tissue damage. Consequently, DPP-IV inhibitors have been suggested as drug candidates for the treatment of impaired glucose tolerance and type 2 diabetes mellitus (e.g. Villhauer, WO98/19998). Other related state of the art can be found in WO 99/38501, DE 19616486, DE 19834591, WO 01/40180, WO 01/55105, US 6110949, WO 00/34241 and US6011155.

Furthermore, DPP IV contributes to the generation and modulation of a T cell immune response. DPP IV (also known as CD26) has an essential role in immune regulation as a T cell activation molecule and a regulator of chemokine function thus suggesting a role for DPP-IV in the pathophysiology of immune-mediated disorders as well as autoimmune diseases (Hosano O. et al., *Modern Rheumatology* 2003, 13(3), 199-204). Abnormal expression of DPP-IV is found in the case of autoimmune diseases, HIV-related diseases and cancer. Natural substrates for DPP-IV are involved in immunomodulation, psycho/neuronal modulation and physiol. processes in general (Boonacker E.; Van Noorden C. J. F, *European Journal of Cell Biology* 2003, 82(2), 53-73). Furthermore, it has been shown that there is a correlation between DPP-IV and the key nuclear protein topoisomerase alpha (Aytac U., Dang, N. H., *Current Drug Targets: Immune, Endocrine and Metabolic Disorders* 2004, 4(1), 11-18). Thus, DPP-IV inhibitors may be useful as medicaments for the treatment of various diseases in which DPP-IV is involved.

We have found novel DPP-IV inhibitors that very efficiently lower plasma glucose levels. Consequently, the compounds of the present invention are useful for the treatment and/or prophylaxis of diabetes, particularly non-insulin dependent diabetes mellitus, and/or impaired glucose tolerance, as well as other conditions wherein the amplification of action of a peptide normally inactivated by DPP-IV gives a therapeutic benefit. In addition, the compounds of the present invention can also be used in the treatment and/or prophylaxis of obesity, metabolic syndrome, β -cell protection, autoimmune diseases such as inflammatory bowel disease, encephalitis periaxialis scleroticans and rheumatoid arthritis, Colitis Ulcerosa, Morbus Crohn, psoriasis, lichen


planus and/or benign prostate hypertrophy. The compounds may also be useful for the prevention of AIDS (acquired immunodeficiency syndrome) or for preventing metastasis, particularly preventing metastasis of breast and prostate cancer to lung. Furthermore, the compounds of the present invention can be used as diuretic agents and for the treatment and/or prophylaxis of hypertension.

According to a first aspect, there is provided the use of compounds of the formula (I)



wherein

R¹ is selected from



R², R³, R⁴, R⁵ and R⁶ are each independently selected from hydrogen, C₁-C₆- alkyl, halogenated C₁-C₆- alkyl, C₁-C₆- alkoxy or halogen; provided that R², R³, R⁴, R⁵ and R⁶ are not all hydrogen;

R⁷ is C₁-C₆- alkyl;

R⁸ is C₁-C₆- alkyl;

X is >C=O or >SO₂;

R⁹ and R¹¹ are hydrogen or together form a double bond;

R¹⁰ and R¹² are independently selected from hydrogen or C₁-C₆- alkyl;


m is 1 or 2;

n is 0, 1, or 2;

and pharmaceutically acceptable salts thereof for the preparation of medicaments for the treatment and/or prophylaxis of diseases which are associated with DPP-IV.


According to a second aspect, there is provided the use according to the first aspect for the preparation of medicaments for the treatment and/or prophylaxis of diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, obesity, and/or metabolic syndrome or β -cell protection.

According to a third aspect, there is provided compounds of the formula (I)



wherein

R^1 is selected from



R^7 is C_1 - C_6 - alkyl;

R^8 is C_1 - C_6 - alkyl;

X is $>C=O$ or $>SO_2$;

R^9 and R^{11} are hydrogen or together form a double bond;

R^{10} and R^{12} are independently selected from hydrogen or C_1 - C_6 - alkyl;


m is 1 or 2;

n is 0, 1, or 2;

and pharmaceutically acceptable salts thereof.

According to a fourth aspect, there is provided process for the manufacturing of the compounds of formula I as defined above, comprising


a) a reductive amination of a ketone of formula II



II

wherein R¹ and n are as defined in above, or

b) a deprotection of a carbamic acid ester of formula III



III

wherein R¹ and n are as defined above and R^P is an amino protecting group.

According to a fifth aspect, there is provided pharmaceutical compositions comprising a compound as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

According to a sixth aspect, there is provided compounds as defined above for use as therapeutic active substances for the treatment and/or prophylaxis of diseases which are associated with DPP-IV.

According to a seventh aspect, there is provided the use of compounds as defined above for the preparation of medicaments for the treatment and/or prophylaxis of diseases which are associated with DPP-IV.

According to an eighth aspect, there is provided the use of compounds according to the third aspect for the preparation of medicaments for the treatment and/or prophylaxis of diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, obesity, and/or metabolic syndrome or β -cell protection.

Unexpectedly, the compounds of the present disclosure exhibit improved therapeutic and pharmacological properties compared to other DPP-IV inhibitors known in the art, such as e.g. in context of pharmacokinetics and bioavailability.

Objects of the embodiments of the present disclosure are compounds of formula I and their pharmaceutically acceptable salts per se and as pharmaceutically active substances, their manufacture, medicaments based on a compound of formula I and their production, as well as the use of the compounds of formula I in accordance with the invention in the control or prevention of illnesses of the aforementioned kind, and, respectively, for the production of corresponding medicaments.

Unless otherwise indicated, the following definitions are set forth to illustrate and define the meaning and scope of the various terms used to describe the invention herein.

In this specification the term "lower" is used to mean a group consisting of one to six, preferably of one to four carbon atom(s).

The term "halogen" refers to fluorine, chlorine, bromine and iodine, with fluorine, bromine and chlorine being preferred. Most preferred halogen is chlorine.


The term "alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent saturated aliphatic hydrocarbon radical of one to twenty carbon atoms, preferably one to sixteen carbon atoms, more preferably one to ten carbon atoms. The term "lower alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent alkyl radical of one to six carbon atoms, preferably one to four carbon atoms. This term is further exemplified by radicals such as methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, isobutyl, t-butyl, n-pentyl, 3-methylbutyl, n-hexyl, 2-ethylbutyl and the like. Preferable lower alkyl residues are methyl and ethyl, with methyl being especially preferred.

The term "halogenated lower alky" refers to a lower alkyl group wherein at least one of the hydrogen atoms of the lower alkyl group is replaced by a halogen atom, preferably fluoro or chloro, most preferably fluoro. Among the preferred halogenated lower alkyl groups are trifluoromethyl, difluoromethyl, fluoromethyl and chloromethyl, with fluoromethyl being especially preferred.

The term "alkoxy" refers to the group R'-O-, wherein R' is alkyl. The term "lower-alkoxy" refers to the group R'-O-, wherein R' is lower alkyl. Examples of lower alkoxy groups are e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy and hexyloxy, with methoxy being especially preferred.


- 5 The term "pharmaceutically acceptable salts" embraces salts of the compounds of formula (I) with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, nitric acid, sulphuric acid, phosphoric acid, citric acid, formic acid, maleic acid, acetic acid, fumaric acid, succinic acid, tartaric acid, methanesulphonic acid, salicylic acid, p-toluenesulphonic acid and the like, which are non toxic to living organisms. Preferred
10 salts with acids are formates, maleates, citrates, hydrochlorides, hydrobromides and methanesulfonic acid salts, with hydrochlorides being especially preferred.

In one embodiment, the present invention relates to compounds for use in therapy having the formula (I)



- 15 wherein

R¹ is selected from



- 20 R², R³, R⁴, R⁵ and R⁶ are each independently selected from hydrogen, lower alkyl, halogenated lower alkyl, lower alkoxy or halogen; provided that R², R³, R⁴, R⁵ and R⁶ are not all hydrogen;

R⁷ is lower alkyl;

R⁸ is lower alkyl;

X is >C=O or >SO₂;

R⁹ and R¹¹ are hydrogen or together form a double bond;


R^{10} and R^{12} are independently selected from hydrogen or lower alkyl;

m is 1 or 2;

n is 0, 1, or 2;

and pharmaceutically acceptable salts thereof.

- 5 In one further embodiment, the invention relates to compounds of formula (I) for use in therapy, wherein R^1 is



- wherein R^2 , R^3 , R^4 , R^5 and R^6 are each independently selected from hydrogen, lower alkyl, halogenated lower alkyl, lower alkoxy or halogen; provided that R^2 , R^3 , R^4 , R^5 and R^6 are not all hydrogen.

R^2 preferably has the meaning of hydrogen, lower alkyl or halogen, more preferably of hydrogen, methyl or chlorine.

- R^3 , R^4 , R^5 and R^6 are preferably selected from hydrogen, lower alkyl, lower alkoxy or halogen. Most preferred lower alkyl is methyl, most preferred lower alkoxy is methoxy and most preferred halogens are selected from fluorine, chlorine and bromine.

In one preferable embodiment, R^2 , R^3 , R^4 and R^5 are hydrogen and R^6 is lower alkyl, lower alkoxy or halogen, more preferably methyl, methoxy or chlorine.


In another preferable embodiment, R^2 , R^3 , R^5 and R^6 are hydrogen and R^4 is lower alkyl or halogen, more preferably methyl, fluorine, chlorine or bromine.

- In another preferable embodiment, R^2 , R^4 and R^5 are hydrogen and R^3 and R^6 are each independently lower alkyl or halogen, more preferably methyl, fluorine or chlorine.

Still in another preferable embodiment, R^3 , R^4 and R^5 are hydrogen and R^2 and R^6 are each independently lower alkyl or halogen, more preferably methyl or chlorine.

- In another embodiment the present invention relates to compounds of formula (I) for use in therapy, wherein R^1 is


- 6 -



wherein R^7 is lower alkyl.

Preferable lower alkyl residues R^7 are methyl and ethyl, with methyl being especially preferred.


- 5 In another embodiment the present invention relates to compounds of formula (I) for use in therapy, wherein R^1 is



wherein R^8 is lower alkyl.

Preferable lower alkyl residue R^8 is methyl.

- 10 In another embodiment the present invention relates to compounds of formula (I) for use in therapy, wherein R^1 is



wherein X is $>C=O$ or $>SO_2$;

R^9 and R^{11} are hydrogen or together form a double bond;

- 15 R^{10} and R^{12} are independently selected from hydrogen or lower alkyl and

m is 1 or 2.

In one preferable embodiment X is $>SO_2$, R^9 , R^{10} , R^{11} and R^{12} are hydrogen and m is 2.

- In another preferable embodiment X is $>C=O$, R^9 and R^{11} together form a double bond, R^{10} and R^{12} are hydrogen and m is 2.
- 20

In another preferable embodiment X is $>C=O$, R^9 and R^{11} are hydrogen or together form a double bond, R^{10} is lower alkyl, preferably methyl and R^{12} is hydrogen and m is 1 or 2, more preferably 1.

Preferred compound of formula I for use in therapy are those, wherein n is 1.

5 Compounds of formula I, wherein n is 2, are also preferred for use in therapy.


Preferred compounds of the general formula I for use in therapy are those selected from the group consisting of:

- (trans)-2-m-tolyl-cyclohexylamine,
- (cis)-2-m-tolyl-cyclohexylamine,
- 10 (trans)-2-o-tolyl-cyclohexylamine,
- (cis)-2-o-tolyl-cyclohexylamine,
- (trans)-2-(2-methoxy-phenyl)-cyclohexylamine,
- (trans)-2-(2,5-dichloro-phenyl)-cyclohexylamine,
- (cis)-2-(2,5-dichloro-phenyl)-cyclohexylamine,
- 15 (trans)-2-(2,4-dimethyl-phenyl)-cyclohexylamine,
- (cis)-2-(3-bromo-phenyl)-cyclohexylamine,
- (trans)-2-(3-bromo-phenyl)-cyclohexylamine,
- (trans)-2-(2-fluoro-5-methyl-phenyl)-cyclohexylamine,
- (cis)-2-(5-methyl-thiophen-2-yl)-cyclohexylamine,
- 20 (trans)-2-(5-methyl-thiophen-2-yl)-cyclohexylamine,
- (cis)-2-(2,4-dichloro-phenyl)-cyclohexylamine,
- (trans)-2-(2,4-dichloro-phenyl)-cyclohexylamine,
- (cis)-2-(3-fluoro-phenyl)-cyclohexylamine,
- (trans)-2-(2-chloro-phenyl)-cyclohexylamine,
- 25 (trans)-2-(2,5-dimethyl-phenyl)-cyclohexylamine,
- (cis/trans)-2-(2-fluoro-phenyl)-cyclohexylamine,
- (trans)-2-(2-fluoro-phenyl)-cyclohexylamine,
- (cis)-2-(3-chloro-phenyl)-cyclohexylamine,


- (trans)-2-(3-chloro-phenyl)-cyclohexylamine,
 (cis)-2-(2,5-dichloro-phenyl)-cycloheptylamine,
 (trans)-2-(2,5-dichloro-phenyl)-cycloheptylamine,
 (cis)-2-(2,5-dichloro-phenyl)-cyclopentylamine,
 5 (trans)-2-(3-methyl-pyrrol-1-yl)-cyclohexylamine,
 (trans)-2-(3-ethyl-pyrrol-1-yl)-cyclohexylamine,
 (trans)-2-(1,1-dioxo-[1,2]thiazinan-2-yl)-cyclohexylamine,
 (trans)-1-(2-amino-cyclohexyl)-5,6-dihydro-1H-pyridin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-4-methyl-1,5-dihydro-pyrrol-2-one,
 10 (trans)-1-(2-amino-cyclohexyl)-4-methyl-5,6-dihydro-1H-pyridin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-piperidin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-4-methyl-pyrrolidin-2-one, and
 pharmaceutically acceptable salts thereof.

The compounds of formula I have two or more asymmetric carbon atoms and can
 15 exist in the form of optically pure enantiomers, mixtures of diastereomers, racemates, or
 mixtures of diastereoisomeric racemates. The invention embraces all of these forms.

In a preferable embodiment, R^1 and the amino group in 1-position of the
 cycloalkylamine structure is in trans- configuration, i.e.




20 In a preferable embodiment, R^1 and the amino group in 1-position of the
 cycloalkylamine structure is in cis- configuration, i.e.



It will be appreciated, that the compounds of general formula (I) in this invention may be derivatised at functional groups to provide derivatives which are capable of converting back to the parent compound in vivo.


The present invention also relates to compounds of the formula (I)



5

wherein

R¹ is selected from



10 R², R³, R⁴, R⁵ and R⁶ are each independently selected from hydrogen, lower alkyl, halogenated lower alkyl or halogen; provided that R², R³, R⁴, R⁵ and R⁶ are not all hydrogen;

R⁷ is lower alkyl;

R⁸ is lower alkyl;

X is >C=O or >SO₂;

15 R⁹ and R¹¹ are hydrogen or together form a double bond;

R¹⁰ and R¹² are independently selected from hydrogen or lower alkyl;

m is 1 or 2;

n is 0, 1, or 2;

and pharmaceutically acceptable salts thereof,

20 with the further proviso that the following compounds are excluded:


2-(m-tolyl)-cyclohexylamine, 2-(p-tolyl)-cyclohexylamine, 2-(o-tolyl)-cyclohexylamine, 2-(2-chlorophenyl)-cyclohexylamine, 2-(3-chlorophenyl)-

cyclohexylamine, 2-(4-chlorophenyl)-cyclohexylamine, 2-(2-bromophenyl)-cyclohexylamine, 2-(o-tolyl)-cyclopentylamine, 2-(p-tolyl)-cyclopentylamine, 2-(4-chlorophenyl)-cyclopentylamine, 2-(3,5-difluorophenyl)-cyclopentylamine, 2-(3-fluorophenyl)-cyclopentylamine, 2-(4-fluorophenyl)-cyclopentylamine, 2-(4-bromophenyl)-cyclopentylamine, and 2-(4-tert-butylphenyl)-cyclopentylamine.

2-(m-Tolyl)-cyclohexylamine and 2-(p-tolyl)-cyclohexylamine are described as intermediates for the synthesis of phenanthridine derivatives in J. Chem. Soc. 1956, 4280-4283. The synthesis of all isomeric forms of 2-(o-tolyl)-cyclohexylamine, 2-(p-tolyl)-cyclohexylamine, 2-(2-chlorophenyl)-cyclohexylamine, 2-(3-chlorophenyl)-cyclohexylamine and 2-(4-chlorophenyl)-cyclohexylamine for study of their proton magnetic resonance spectra is disclosed in J. Org. Chem. 1962, 27, 3006-3010. In J. Org. Chem. 1971, 36, 3046-3048 the synthesis of all isomers of 2-(2-bromophenyl)-cyclohexylamine and their NMR spectra are described.

2-(o-Tolyl)-cyclopentylamine is known from WO 2004/016601 as reactant for the preparation of aminohydroxyalkylbenzothiazolones useful as β 3 adrenoreceptor agonists. 2-(p-Tolyl)-cyclopentylamine, 2-(4-chlorophenyl)-cyclopentylamine, 2-(3,5-difluorophenyl)-cyclopentylamine, 2-(3-fluorophenyl)-cyclopentylamine, 2-(4-fluorophenyl)-cyclopentylamine, 2-(4-bromophenyl)-cyclopentylamine and 2-(4-tert-butylphenyl)-cyclopentylamine are disclosed in WO 2001/042203 as intermediates for the synthesis of N-(phenylcyclopentyl)sulfonamides with glutamate receptor function potentiating activity.

Preferred compounds of formula I are those, wherein R^1 is



wherein R^2 , R^3 , R^4 , R^5 and R^6 are each independently selected from hydrogen, lower alkyl, halogenated lower alkyl, lower alkoxy or halogen; provided that R^2 , R^3 , R^4 , R^5 and R^6 are not all hydrogen.

R^2 preferably has the meaning of hydrogen, lower alkyl or halogen, more preferably of hydrogen, methyl or chlorine.

R^3 , R^4 , R^5 and R^6 are preferably selected from hydrogen, lower alkyl, lower alkoxy or halogen. Most preferred lower alkyl is methyl, most preferred lower alkoxy is methoxy and most preferred halogens are selected from fluorine, chlorine and bromine.


In one preferable embodiment, R^2 , R^3 , R^4 and R^5 are hydrogen and R^6 is lower alkyl, lower alkoxy or halogen, more preferably methyl, methoxy or chlorine.

In another preferable embodiment, R^2 , R^3 , R^5 and R^6 are hydrogen and R^4 is lower alkyl or halogen, more preferably methyl, fluorine, chlorine or bromine.

In another preferable embodiment, R^2 , R^4 and R^5 are hydrogen and R^3 and R^6 are each independently lower alkyl or halogen, more preferably methyl, fluorine or chlorine.

Still in another preferable embodiment, R^3 , R^4 and R^5 are hydrogen and R^2 and R^6 are each independently lower alkyl or halogen, more preferably methyl or chlorine.


Further preferred compounds of formula I of the present invention are those, wherein R^1 is



and wherein R^7 is lower alkyl.

Preferable lower alkyl residues R^7 are methyl and ethyl, with methyl being especially preferred.

In another embodiment of the present invention, compounds of formula I are those, wherein R^1 is




and wherein R^8 is lower alkyl.

Preferable lower alkyl residue R^8 is methyl.

Also preferred are compounds of formula I of the present invention, wherein R^1 is

- 12 -



wherein X is $>\text{C}=\text{O}$ or $>\text{SO}_2$;

R^9 and R^{11} are hydrogen or together form a double bond;

R^{10} and R^{12} are independently selected from hydrogen or lower alkyl and

5 m is 1 or 2.

In one preferable embodiment X is $>\text{SO}_2$, R^9 , R^{10} , R^{11} and R^{12} are hydrogen and m is 2.

In another preferable embodiment X is $>\text{C}=\text{O}$, R^9 and R^{11} together form a double bond, R^{10} and R^{12} are hydrogen and m is 2.

10 In another preferable embodiment X is $>\text{C}=\text{O}$, R^9 and R^{11} are hydrogen or together form a double bond, R^{10} is lower alkyl, preferably methyl and R^{12} is hydrogen and m is 1 or 2, more preferably 1.

Preferred compounds of formula I are those, wherein n is 1.

Compounds of formula I, wherein n is 2,

15 Especially preferred compounds of the general formula I are those selected from the group consisting of:

(trans)-2-(2-methoxy-phenyl)-cyclohexylamine,

(trans)-2-(2,5-dichloro-phenyl)-cyclohexylamine,

(cis)-2-(2,5-dichloro-phenyl)-cyclohexylamine,

20 (trans)-2-(2,4-dimethyl-phenyl)-cyclohexylamine,

(cis)-2-(3-bromo-phenyl)-cyclohexylamine,

(trans)-2-(3-bromo-phenyl)-cyclohexylamine,

(trans)-2-(2-fluoro-5-methyl-phenyl)-cyclohexylamine,

(cis)-2-(5-methyl-thiophen-2-yl)-cyclohexylamine,


25 (trans)-2-(5-methyl-thiophen-2-yl)-cyclohexylamine,

- (cis)-2-(2,4-dichloro-phenyl)-cyclohexylamine,
 (trans)-2-(2,4-dichloro-phenyl)-cyclohexylamine,
 (cis)-2-(3-fluoro-phenyl)-cyclohexylamine,
 (trans)-2-(2-chloro-phenyl)-cyclohexylamine,
 5 (trans)-2-(2,5-dimethyl-phenyl)-cyclohexylamine,
 (cis/trans)-2-(2-fluoro-phenyl)-cyclohexylamine,
 (trans)-2-(2-fluoro-phenyl)-cyclohexylamine,
 (cis)-2-(2,5-dichloro-phenyl)-cycloheptylamine,
 (trans)-2-(2,5-dichloro-phenyl)-cycloheptylamine,
 10 (cis)-2-(2,5-dichloro-phenyl)-cyclopentylamine,
 (trans)-2-(3-methyl-pyrrol-1-yl)-cyclohexylamine,
 (trans)-2-(3-ethyl-pyrrol-1-yl)-cyclohexylamine,
 (trans)-2-(1,1-dioxo-[1,2]thiazinan-2-yl)-cyclohexylamine,
 (trans)-1-(2-amino-cyclohexyl)-5,6-dihydro-1H-pyridin-2-one,
 15 (trans)-1-(2-amino-cyclohexyl)-4-methyl-1,5-dihydro-pyrrol-2-one,
 (trans)-1-(2-amino-cyclohexyl)-4-methyl-5,6-dihydro-1H-pyridin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-piperidin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-4-methyl-pyrrolidin-2-one, and
 pharmaceutically acceptable salts thereof.

- 20 The present invention also relates to a process for the manufacture of compounds of formula I.

In general the compounds of the formula I can be obtained either


a) by a reductive amination of a ketone of formula II



II

wherein R^1 and n are as defined above, or

b) by deprotection of a carbamic acid ester of formula III



III

wherein R^1 and n are as defined above and R^P is an amino protecting group.

- 5 R^P is a suitable amino protecting group such as benzyloxycarbonyl (Z or Cbz), allyloxycarbonyl (Aloc), 9-fluorenylmethoxycarbonyl (Fmoc), and preferably, tert-butoxycarbonyl (Boc).


The compounds of formula (I) can be manufactured by the methods given below, by the methods given in the Examples or by analogous methods. Appropriate reaction...
 10 conditions for the individual reaction steps are known to the person skilled in the art. Starting materials are either commercially available or can be prepared by methods analogous to the methods given below or in the Examples or by methods known in the art.

The compounds of the present invention can be prepared as illustrated in the
 15 schemes below:

Compounds of general formula Ia, in which R^1 is linked to the cycloalkane core through a carbon atom, are synthesized from a ketone II by methods known in the art such as by reductive amination using preferably ammonium acetate and sodium cyanoborohydride (Scheme 1)


20 Ketones of general formula II can be obtained from the respective alcohol e.g. by an oxidation using methods known in the art, preferably by using a Dess Martin Reagent. The alcohol itself can be obtained from the respective epoxide by methods known in the art, preferably using an organometallic reagent such as the suitable organo lithium reagent or the suitable Grignard reagent.

Scheme 1



Compounds of general formula Ib, in which R¹ is linked to the cycloalkane core through a nitrogen atom can be synthesized from a carbamic acid ester III by methods known in the art. When R^P is *tert*-butoxycarbonyl the reaction preferably is performed in the presence of hydrogen chloride in dioxane or with trifluoroacetic acid in dichloromethane. The carbamic acid ester III can be obtained from a carboxylic acid ester IV by hydrolysis and subsequent Curtius rearrangement, using methods known in the art (Scheme 2).


10 Scheme 2



R^a suitably is methyl or ethyl; R^P is a suitable amino protecting group such as benzyloxycarbonyl, allyloxycarbonyl, and preferably, *tert*-butoxycarbonyl.

The synthesis of lactam or sultam derivatives Ic starts from cycloalkylamine V as shown below in scheme 3. V is reacted with an acid chloride or a sulfonyl chloride VII in the presence of a base (e. g., triethylamine) to afford amide or sulfonamide V. Then, cyclisation of V using a base, e.g., sodium hydride, in a solvent such as N,N-dimethylformamide, optionally in the presence of sodium iodide, leads to Ic. The carbamic acid ester can be transformed to the free amine as shown in scheme 2.

Scheme 3




R^p is a suitable amino protecting group such as benzyloxycarbonyl, allyloxycarbonyl, and, preferably, tert-butoxycarbonyl; Hal is a halogen, preferably chlorine

5

The unsaturated lactams of the formula Ic wherein R⁹ and R¹⁰ form a double bond and X is >C=O can be synthesized from cycloalkylamine VI according to scheme 4. Thus, alkylation of VI with alkenyl halide IX (in the presence of a base, e. g., triethylamine), followed by acylation (in the presence of a base, e. g., triethylamine) with acyl halide IX, affords amide VIII. Compound VIII can then be subjected to ring-closing metathesis (Acc. Chem. Res. 2001, 34, 18), using a ruthenium catalyst, e. g., bis(tricyclohexylphosphine)-benzylidene ruthenium(IV)dichloride, and optionally a Lewis acid, e. g., tetraisopropyl-orthotitanate, to afford Id. The carbamic acid ester can be transformed to the free amine as shown in scheme 2.

Scheme 4




R^p is a suitable amino protecting group such as benzyloxycarbonyl, allyloxycarbonyl, and, preferably, tert-butoxycarbonyl; Hal is a halogen, preferably chlorine

The invention further relates to compounds of formula (I) as defined above, when
20 manufactured according to a process as defined above.

As described above, the compounds of formula I of the present invention can be used as medicaments for the treatment and/or prophylaxis of diseases which are


associated with DPP-IV such as diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, inflammatory bowel disease, Colitis Ulcerosa, Morbus Crohn, obesity, and/or metabolic syndrome or β -cell protection, preferably non-insulin dependent diabetes mellitus and/or impaired glucose tolerance. Furthermore, the compounds of the present invention can be used as diuretic agents or for the treatment and/or prophylaxis of hypertension.

The invention therefore also relates to pharmaceutical compositions comprising a compound of formula I



wherein

R^1 is selected from



R^2 , R^3 , R^4 , R^5 and R^6 are each independently selected from hydrogen, lower alkyl, halogenated lower alkyl, lower alkoxy or halogen; provided that R^2 , R^3 , R^4 , R^5 and R^6 are not all hydrogen;

R^7 is lower alkyl;

R^8 is lower alkyl;

X is $>\text{C}=\text{O}$ or $>\text{SO}_2$;

R^9 and R^{11} are hydrogen or together form a double bond;

R^{10} and R^{12} are independently selected from hydrogen or lower alkyl;

m is 1 or 2; and

n is 0, 1, or 2;

or a pharmaceutically acceptable salt thereof;
and a pharmaceutically acceptable carrier and/or adjuvant.

Further, the invention relates to compounds as defined above for use as therapeutic active substances, particularly as therapeutic active substances for the treatment and/or prophylaxis of diseases which are associated with DPP-IV such as diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, inflammatory bowel disease, Colitis Ulcerosa, Morbus Crohn, obesity, and/or metabolic syndrome or β -cell protection, preferably for use as therapeutic active substances for the treatment and/or prophylaxis of non-insulin dependent diabetes mellitus and/or impaired glucose tolerance. Furthermore, the invention relates to compounds as defined above for use as diuretic agents or for use as therapeutic active substances for the treatment and/or prophylaxis of hypertension.

In another embodiment, the invention relates to a method for the treatment and/or prophylaxis of diseases which are associated with DPP-IV such as diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, inflammatory bowel disease, Colitis Ulcerosa, Morbus Crohn, obesity, and/or metabolic syndrome or β -cell protection, preferably for the treatment and/or prophylaxis of non-insulin dependent diabetes mellitus and/or impaired glucose tolerance, which method comprises administering a compound as defined above to a human being or animal. Furthermore, the invention relates to a method for the treatment and/or prophylaxis as defined above, wherein the disease is hypertension or wherein a diuretic agent has a beneficial effect.

The invention further relates to the use of compounds as defined above for the treatment and/or prophylaxis of diseases which are associated with DPP-IV such as diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, inflammatory bowel disease, Colitis Ulcerosa, Morbus Crohn, obesity, and/or metabolic syndrome or β -cell protection, preferably for the treatment and/or prophylaxis of non-insulin dependent diabetes mellitus and/or impaired glucose tolerance. Furthermore, the invention relates to the use as defined above, wherein the disease is hypertension or to the use as diuretic agent.

In addition, the invention relates to the use of compounds as defined above for the preparation of medicaments for the treatment and/or prophylaxis of diseases which are associated with DPP-IV such as diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, inflammatory bowel disease, Colitis Ulcerosa, Morbus Crohn, obesity, and/or metabolic syndrome or β -cell protection, preferably for the treatment and/or prophylaxis of non-insulin dependent diabetes mellitus and/or impaired glucose tolerance. Such medicaments comprise a compound as defined above.

Furthermore, the invention relates to the use as defined above, wherein the disease is hypertension or the use for the preparation of diuretic agents.

In context with the methods and uses defined above, the following diseases relate to a preferred embodiment: diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, obesity, and/or metabolic syndrome or β -cell protection,
5 preferably non-insulin dependent diabetes mellitus and/or impaired glucose tolerance.

The following tests were carried out in order to determine the activity of the compounds of formula I.

Activity of DPP-IV inhibitors are tested with natural human DPP-IV derived from
10 a human plasma pool or with recombinant human DPP-IV. Human citrate plasma from different donors is pooled, filtered through a 0.2 micron membrane under sterile conditions and aliquots of 1 ml are shock frozen and stored at -120°C until used. In the colorimetric DPP-IV assay 5 to 10 μl human plasma and in the fluorometric assay 1.0 μl of human plasma in a total assay volume of 100 μl is used as an enzyme source. The
15 cDNA of the human DPP-IV sequence of amino acid 31 - to 766, restricted for the N-terminus and the transmembrane domain, is cloned into Pichia pastoris. Human DPP-IV is expressed and purified from the culture medium using conventional column chromatography including size exclusion and anion and cation chromatography. The purity of the final enzyme preparation of Coomassie blue SDS-PAGE is $> 95\%$. In the
20 colorimetric DPP-IV assay 20 ng rec.-h DPP-IV and in the fluorometric assay 2 ng rec.-h DPP-IV in a total assay volume of 100 μl is used as an enzyme source.

In the fluorogenic assay Ala-Pro-7-amido-4-trifluoromethylcoumarin (Calbiochem No 125510) is used as a substrate. A 20 mM stock solution in 10 % DMF/ H_2O is stored at -20°C until use. In IC_{50} determinations a final substrate concentration of 50 μM is used.
25 In assays to determine kinetic parameters as K_m , $V_{\max}$, K_i , the substrate concentration is varied between 10 μM and 500 μM .

In the colorimetric assay H-Ala-Pro-pNA.HCl (Bachem L-1115) is used as a substrate. A 10 mM stock solution in 10% MeOH/ H_2O is stored at -20°C until use. In IC_{50} determinations a final substrate concentration of 200 μM is used. In assays to
30 determine kinetic parameters as K_m , $V_{\max}$, K_i , the substrate concentration is varied between 100 μM and 2000 μM .

Fluorescence is detected in a Perkin Elmer Luminescence Spectrometer LS 50B at an excitation wavelength of 400 nm and an emission wavelength of 505 nm continuously

every 15 seconds for 10 to 30 minutes. Initial rate constants are calculated by best fit linear regression.

The absorption of pNA liberated from the colorimetric substrate is detected in a Packard SpectraCount at 405 nm continuously every 2 minutes for 30 to 120 minutes.

- 5 Initial rate constants are calculated by best fit linear regression.

- DPP-IV activity assays are performed in 96 well plates at 37°C in a total assay volume of 100 µl. The assay buffer consists of 50 mM Tris/HCl, pH 7.8 containing 0.1 mg/ml BSA and 100 mM NaCl. Test compounds are solved in 100 % DMSO, diluted to the desired concentration in 10% DMSO/H₂O. The final DMSO concentration in the
10 assay is 1 % (v/v). At this concentration enzyme inactivation by DMSO is < 5%. Compounds are with (10 minutes at 37°C) and without preincubation with the enzyme. Enzyme reactions are started with substrate application followed by immediate mixing.

IC₅₀ determinations of test compounds are calculated by non-linear best fit regression of the DPP-IV inhibition of at least 5 different compound concentrations.

- 15 Kinetic parameters of the enzyme reaction are calculated at at least 5 different substrate concentrations and at least 5 different test compound concentrations.

The compounds of the present invention exhibit IC₅₀ values of 0.1 µM to 50µM, more preferably of 0.1µM to 1 µM as shown in the following table:

Example	IC ₅₀ [µM]
7	0.13
18	0.16
24	0.72
30	0.73

- 20 The compounds of formula I and/or their pharmaceutically acceptable salts can be used as medicaments, e.g. in the form of pharmaceutical preparations for enteral, parenteral or topical administration. They can be administered, for example, perorally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions, rectally, e.g. in the form of suppositories,
25 parenterally, e.g. in the form of injection solutions or infusion solutions, or topically, e.g. in the form of ointments, creams or oils. Oral administration is preferred.

The production of the pharmaceutical preparations can be effected in a manner which will be familiar to any person skilled in the art by bringing the described compounds of formula I and/or their pharmaceutically acceptable salts, optionally in combination with other therapeutically valuable substances, into a galenical

- 5 administration form together with suitable, non-toxic, inert, therapeutically compatible solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

Suitable carrier materials are not only inorganic carrier materials, but also organic carrier materials. Thus, for example, lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used as carrier materials for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carrier materials for soft gelatine capsules are, for
10 example, vegetable oils, waxes, fats and semi-solid and liquid polyols (depending on the nature of the active ingredient no carriers might, however, be required in the case of soft gelatine capsules). Suitable carrier materials for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar and the like. Suitable carrier
15 materials for injection solutions are, for example, water, alcohols, polyols, glycerol and vegetable oils. Suitable carrier materials for suppositories are, for example, natural or hardened oils, waxes, fats and semi-liquid or liquid polyols. Suitable carrier materials for topical preparations are glycerides, semi-synthetic and synthetic glycerides, hydrogenated oils, liquid waxes, liquid paraffins, liquid fatty alcohols, sterols, polyethylene glycols and
20 cellulose derivatives.

Usual stabilizers, preservatives, wetting and emulsifying agents, consistency-improving agents, flavour-improving agents, salts for varying the osmotic pressure, buffer substances, solubilizers, colorants and masking agents and antioxidants come into consideration as pharmaceutical adjuvants.

- 25 The dosage of the compounds of formula I can vary within wide limits depending on the disease to be controlled, the age and the individual condition of the patient and the mode of administration, and will, of course, be fitted to the individual requirements in each particular case. For adult patients a daily dosage of about 1 to 1000 mg, especially about 1 to 100 mg, comes into consideration. Depending on severity of the disease and
30 the precise pharmacokinetic profile the compound could be administered with one or several daily dosage units, e.g. in 1 to 3 dosage units.

The pharmaceutical preparations conveniently contain about 1-500 mg, preferably 1-100 mg, of a compound of formula I.

The following Examples serve to illustrate the present invention in more detail.

- 35 They are, however, not intended to limit its scope in any manner.

Examples

Example 1 and 2

(trans)-2-m-tolyl-cyclohexylamine and (cis)-2-m-tolyl-cyclohexylamine

To a solution of 1-bromo-3-methyl-benzene in THF at -78°C, was added dropwise
5 a solution of nBuLi (1.6M in THF, 12.4 ml) and the reaction mixture was stirred at -78°C
for 30 minutes. After such time, 7-oxa-bicyclo[4.1.0]heptane (3.4g) was added slowly to
the reaction mixture followed by the addition of boron trifluoride etherate (2.5 ml). The
reaction mixture was stirred at -78 °C for another 2 hours before allowing it to warm up
to room temperature. The solution was then treated with solution of ammonium
10 chloride (25 ml), the phases were separated and then extracted twice with ethyl acetate.
The combined organic extracts were then washed with water, dried over sodium sulfate
and concentrated *in vacuo*. The residue was then purified by column chromatography to
give 2.9 g of (cis/trans) 2-o-tolyl-cyclohexanol MS(EI) 190.1 (M^+).

To (cis/trans) 2-o-tolyl-cyclohexanol (1g) in dichloromethane (30ml) was added
15 Dess-Martin periodinane (Aldrich 27,462-3) at room temperature. The reaction mixture
was allowed to stir at room temperature for 16 hours before diethyl ether was added. The
volume of solvent was reduced *in vacuo* to about one quarter of the initial amount. More
diethylether (87ml) was added and the solution was washed with a 10% solution of
sodium thiosulfate (87ml), a solution of saturated sodium bicarbonate (87ml), brine
20 (100mL) and water (100 ml). The organic phase was then dried over sodium sulfate and
concentrated *in vacuo*. The residue was then purified by column chromatography to give
0.54 g of 2-o-tolyl-cyclohexanone MS (EI) 188.2 (M^+).

To 2-o-tolyl-cyclohexanone (0.22g) in methanol (30 ml) was added ammonium
acetate (0.90g) and the reaction was allowed to stir at room temperature for 16 hours.
25 After such time sodiumcyanoborohydride (91mg) was added to the reaction mixture and
stirred for 10 minutes. The reaction mixture was then treated with a saturated aqueous
solution of NaHCO₃, and extracted with ethyl acetate (2 x 50 ml). The combined organic
phases were dried over sodium sulfate, concentrated *in vacuo* and purified by column
chromatography to yield (cis) 2-m-tolyl-cyclohexylamine (35mg) MS(ISP) 190.3
30 ($M+H$)⁺ and (trans) 2-m-tolyl-cyclohexylamine MS(ISP) 190.3 (M^+H)⁺.

The following examples were prepared in analogy to Examples 1 and 2:

Ex.	Systematic name	Starting material	MW	MW (found) (MH ⁺)
3	(trans)-2-o-Tolyl-cyclohexylamine	1-Bromo-2-methyl-benzene	189.3	190.2
4	(cis)-2-o-Tolyl-cyclohexylamine	1-Bromo-2-methyl-benzene	189.3	190.2
5	(trans)-2-(2-Methoxy-phenyl)-cyclohexylamine	1-Bromo-2-methoxy-benzene	205.3	206.1
6	(trans)-2-(2,5-Dichloro-phenyl)-cyclohexylamine	2-Bromo-1,4-dichloro-benzene	244.2	244.2
7	(cis)-2-(2,5-Dichloro-phenyl)-cyclohexylamine	2-Bromo-1,4-dichloro-benzene	244.2	244.2
8	(trans)-2-(2,4-Dimethyl-phenyl)-cyclohexylamine	1-Bromo-2,4-dimethyl-benzene	203.3	204.1
9	(cis)-2-(3-Bromo-phenyl)-cyclohexylamine	1,3-Dibromo-benzene	254.2	254.0
10	(trans)-2-(3-Bromo-phenyl)-cyclohexylamine	1,3-Dibromo-benzene	254.2	254.0
11	(trans)-2-(2-Fluoro-5-methyl-phenyl)-cyclohexylamine	2-Bromo-1-fluoro-4-methyl-benzene	207.3	244.2
12	(cis)-2-(5-Methyl-thiophen-2-yl)-cyclohexylamine	2-Bromo-5-methyl-thiophene	195.3	196.2
13	(trans)-2-(5-Methyl-thiophen-2-yl)-cyclohexylamine	2-Bromo-5-methyl-thiophene	195.3	196.1

Ex.	Systematic name	Starting material	MW	MW (found) (MH ⁺)
14	(cis)-2-(2,4-Dichloro-phenyl)-cyclohexylamine	1-Bromo-2,4-dichloro-benzene	244.2	244.2
15	(trans)-2-(2,4-Dichloro-phenyl)-cyclohexylamine	1-Bromo-2,4-dichloro-benzene	244.2	244.2
16	(cis)-2-(3-Fluoro-phenyl)-cyclohexylamine	1-Bromo-3-fluoro-benzene	193.3	194.2
17	(trans)-2-(2-Chloro-phenyl)-cyclohexylamine	1-Chloro-2-iodo-benzene	209.7	210.2
18	(trans)-2-(2,5-Dimethyl-phenyl)-cyclohexylamine	2-Bromo-1,4-dimethyl-benzene	203.3	204.3
19	(cis/trans)-2-(2-Fluoro-phenyl)-cyclohexylamine	1-Bromo-2-fluoro-benzene	193.3	194.3
20	(trans)-2-(2-Fluoro-phenyl)-cyclohexylamine	1-Bromo-2-fluoro-benzene	193.3	194.2
21	(cis)-2-(3-Chloro-phenyl)-cyclohexylamine	1-Bromo-3-chloro-benzene	209.7	210.2
22	(trans)-2-(3-Chloro-phenyl)-cyclohexylamine	1-Bromo-3-chloro-benzene	209.7	210.2
23	(cis)-2-(2,5-Dichloro-phenyl)-cycloheptylamine	2-Bromo-1,4-dichloro-benzene and 8-Oxa-bicyclo[5.1.0]octane	258.2	258.1
24	(trans)-2-(2,5-Dichloro-phenyl)-cycloheptylamine	2-Bromo-1,4-dichloro-benzene and 8-Oxa-bicyclo[5.1.0]octane	258.2	258.1

Ex.	Systematic name	Starting material	MW	MW (found) (MH ⁺)
25	(cis)-2-(2,5-Dichloro-phenyl)-cyclopentylamine	2-Bromo-1,4-dichloro-benzene and 6-Oxa-bicyclo[3.1.0]hexane	230.1	230.1

Example 26

(trans)-2-(3-Methyl-pyrrol-1-yl)-cyclohexylamine

a) trans-[2-(3-Formyl-pyrrol-1-yl)-cyclohexyl]-carbamic acid tert-butyl ester

5 A solution of trans-(2-amino-cyclohexyl)-carbamic acid tert-butyl ester (*Tetrahedron Lett.* 2000, 41, 9607; 400 mg, 1.87 mmol) and 2,5-dimethoxy-3-tetrahydrofurancarboxaldehyde (365 mg, 2.05 mmol) in pyridine (0.5 mL) and acetic acid (0.82 mL) was heated at 100 °C for 4.5 h. After cooling, the reaction mixture was partitioned between ethyl acetate and 10% aq. citric acid solution. The organic layer was washed with brine, dried (MgSO₄), and evaporated. Chromatography (SiO₂, heptane-ethyl acetate gradient) yielded the title compound (334 mg, 61%). Off-white solid, MS (ISP) 293.3 (M⁺H)⁺.

b) trans-2-(3-Methyl-pyrrol-1-yl)-cyclohexylamine

15 Triethylsilane (368 mg, 3.16 mmol) was added at 0 °C to a solution of trans-[2-(3-formyl-pyrrol-1-yl)-cyclohexyl]-carbamic acid tert-butyl ester (330 mg, 1.13 mmol) in trifluoroacetic acid (5.1 ml), then after 90 min the reaction mixture was evaporated and partitioned between ethyl acetate and 2 M aq. sodium hydroxide solution. The organic layer was washed with brine, dried (MgSO₄), and evaporated. Chromatography (SiO₂, CH₂Cl₂/MeOH/NH₄OH 90:10:0.25) yielded the title compound (163 mg, 81%). Yellow oil, MS (ISP) 179.1 (M⁺H)⁺.

Example 27

trans-2-(3-Ethyl-pyrrol-1-yl)-cyclohexylamine

a) trans-[2-(3-Ethyl-pyrrol-1-yl)-cyclohexyl]-carbamic acid tert-butyl ester

Methylmagnesium chloride solution (3 M in tetrahydrofuran, 0.23 ml, 0.68 mmol) was added at -78 °C to a solution of trans-[2-(3-formyl-pyrrol-1-yl)-cyclohexyl]-carbamic acid tert-butyl ester (example 26a, 100 mg, 0.34 mmol) in tetrahydrofuran (2 ml), then after 3.5 h the reaction was quenched by addition of saturated aqueous ammonium chloride solution and extracted with dichloromethane. The organic layer was washed with brine, dried (MgSO₄), and evaporated. This crude material (103 mg) was dissolved in dichloromethane (2 ml) and treated with triethylsilane (58 mg, 0.50 mmol) and trifluoroacetic acid (190 mg, 1.67 mmol). The reaction mixture was allowed to reach 0°C over 3 h, then evaporated and the residue chromatographed (SiO₂, heptane-ethyl acetate gradient) to afford the title compound (27 mg, 28%). Yellow solid, MS (ISP) 293.2 (M⁺H)⁺.

b) trans-2-(3-Ethyl-pyrrol-1-yl)-cyclohexylamine

A solution of trans-[2-(3-ethyl-pyrrol-1-yl)-cyclohexyl]-carbamic acid tert-butyl ester (24 mg, 82 µmol) in hydrochloric acid solution (4 M in 1,4-dioxane) was stirred for 90 min at room temperature, then evaporated. The residue was taken up in CH₂Cl₂/MeOH/NH₄OH 90:10:0.25 and the solution concentrated in vacuo. Chromatography (SiO₂, CH₂Cl₂/MeOH/NH₄OH 90:10:0.25) afforded the title compound (5 mg, 32%). Light yellow solid, MS (ISP) 193.4 (M⁺H)⁺.

Example 28

trans-2-(1,1-Dioxo-[1,2]thiazinan-2-yl)-cyclohexylamine

a) trans-[2-(4-Chloro-butane-1-sulfonylamino)-cyclohexyl]-carbamic acid tert-butyl ester

4-Chlorobutanesulfonyl chloride (178 mg, 0.93 mmol). was added at 0 °C to a solution of trans-(2-amino-cyclohexyl)-carbamic acid tert-butyl ester (200 mg, 0.93 mmol) and triethylamine (94 mg, 0.93 mmol) in dichloromethane (2 ml), and the reaction mixture was allowed to reach room temperature over 3 h, then partitioned between dichloromethane and 10% aq. citric acid solution. The organic layer was washed with 1 M aq. sodium carbonate solution and brine, dried (MgSO₄), and evaporated. Chromatography (SiO₂, heptane-ethyl acetate gradient) afforded the title compound (130 mg, 38%). Off-white solid, MS (ISP) 367.2 (M-H)⁻.

b) trans-2-(1,1-Dioxo-[1,2]thiazinan-2-yl)-cyclohexylamine

Sodium hydride (60% dispersion in mineral oil, 15 mg, 0.38 mmol) was added at 0°C to a solution of trans-[2-(4-chloro-butane-1-sulfonylamino)-cyclohexyl]-carbamic acid tert-butyl ester (125 mg, 0.34 mmol) and sodium iodide (51 mg, 0.34 mmol), and the reaction mixture was stirred at room temperature for 24 h, then another portion of sodium hydride (15 mg, 0.38 mmol) was added, and the reaction mixture was heated at 60 °C for 3 h. After cooling, the solution was partitioned between heptane/ethyl acetate (1:1) and water. The organic layer was washed with brine, evaporated, and chromatographed (SiO₂, heptane-ethyl acetate gradient). The title compound was obtained from this material in accordance with the general method of example 27b. Off-white solid, MS (ISP) 233.1 (M⁺H)⁺.

Example 29

trans-1-(2-Amino-cyclohexyl)-5,6-dihydro-1H-pyridin-2-one

a) trans-(2-But-3-enylamino-cyclohexyl)-carbamic acid tert-butyl ester

The title compound was produced in accordance with the general method of example 30a from trans-(2-amino-cyclohexyl)-carbamic acid tert-butyl ester and 4-bromo-1-butene. Brown solid, MS (ISP) 269.3 (M+H)⁺.

b) trans-[2-(Acryloyl-but-3-enyl-amino)-cyclohexyl]-carbamic acid tert-butyl ester

Acryloyl chloride (47 mg, 0.50 mmol) was added at 0 °C to a solution of trans-(2-but-3-enylamino-cyclohexyl)-carbamic acid tert-butyl ester (123 mg, 0.46 mmol) and triethylamine (51 mg, 0.51 mmol) in dichloromethane (3 ml), and the reaction mixture was allowed to reach room temperature over 3 h. After partitioning between dichloromethane and 10% aq. citric acid solution, the organic layer was washed with 1 M aq. sodium carbonate solution and brine, dried (MgSO₄), and evaporated. Chromatography (SiO₂, heptane-ethyl acetate gradient) afforded the title compound (103 mg, 70%). Light yellow solid, MS (ISP) 323.3 (M⁺H)⁺.

c) trans-[2-(6-Oxo-3,6-dihydro-2H-pyridin-1-yl)-cyclohexyl]-carbamic acid tert-butyl ester

Bis(tricyclohexylphosphine)benzylidene ruthenium(IV)dichloride (13 mg, 16 μmol) was added to a solution of trans-[2-(acryloyl-but-3-enyl-amino)-cyclohexyl]-carbamic acid tert-butyl ester (50 mg, 0.16 mmol) and tetraisopropyl orthotitanate (8.8 mg, 31 μmol) in dichloromethane (2.5 ml), and the reaction mixture was stirred for 1 h at room temperature. The solvent was then evaporated and the residue chromatographed

(SiO₂, heptane-ethyl acetate gradient) to produce the title compound (44 mg, 96%). Off-white solid, MS (ISP) 295.2 (M⁺H)⁺.

d) trans-1-(2-Amino-cyclohexyl)-5,6-dihydro-1H-pyridin-2-one

The title compound was produced in accordance with the general method of
5 example 27b from trans-[2-(6-oxo-3,6-dihydro-2H-pyridin-1-yl)-cyclohexyl]-carbamic acid tert-butyl ester. Light yellow liquid, MS (ISP) 195.3 (M⁺H)⁺.

Example 30

trans-1-(2-Amino-cyclohexyl)-4-methyl-1,5-dihydro-pyrrol-2-one

a) trans-[2-(2-Methyl-allylamino)-cyclohexyl]-carbamic acid tert-butyl ester

10 Methallyl bromide (139 mg, 1.03 mmol) was added at 0 °C to a solution of trans-(2-amino-cyclohexyl)-carbamic acid tert-butyl ester (200 mg, 0.93 mmol) and triethylamine (113 mg, 1.12 mmol) in tetrahydrofuran (4 ml). The reaction mixture was stirred for 16 h at room temperature, then partitioned between ethyl acetate and 1 M aq. sodium hydroxide solution. The organic layer was washed with brine, dried (MgSO₄),
15 and evaporated. Chromatography (SiO₂, CH₂Cl₂/MeOH/NH₄OH 90:10:0.25) afforded the title compound (177 mg, 71%). Orange solid, MS (ISP) 269.3 (M⁺H)⁺.

b) trans-[2-[Acryloyl-(2-methyl-allyl)-amino]-cyclohexyl]-carbamic acid tert-butyl ester

The title compound was produced in accordance with the general method of
20 example 29b from trans-[2-(2-methyl-allylamino)-cyclohexyl]-carbamic acid tert-butyl ester and acryloyl chloride. Off-white solid, MS (ISP) 323.3 (M⁺H)⁺.

c) trans-[2-(4-Methyl-2-oxo-2,5-dihydro-pyrrol-1-yl)-cyclohexyl]-carbamic acid tert-butyl ester

The title compound was produced in accordance with the general method of
25 example 31c from trans-[2-[acryloyl-(2-methyl-allyl)-amino]-cyclohexyl]-carbamic acid tert-butyl ester. Black solid, MS (ISP) 295.2 (M⁺H)⁺.

d) trans-1-(2-Amino-cyclohexyl)-4-methyl-1,5-dihydro-pyrrol-2-one

The title compound was produced in accordance with the general method of
example 27b from trans-[2-(4-methyl-2-oxo-2,5-dihydro-pyrrol-1-yl)-cyclohexyl]-
30 carbamic acid tert-butyl ester. Yellow solid, MS (ISP) 195.2 (M⁺H)⁺.

Example 31

trans-1-(2-Amino-cyclohexyl)-4-methyl-5,6-dihydro-1H-pyridin-2-one

a) trans-[2-(3-Methyl-but-3-enylamino)-cyclohexyl]-carbamic acid tert-butyl ester

The title compound was produced in accordance with the general method of
5 example 30a from trans-(2-amino-cyclohexyl)-carbamic acid tert-butyl ester and 4-bromo-2-methyl-1-butene (*J. Org. Chem.* 1997, 62, 1536). Brown solid, MS (ISP) 283.3 (M^+H)⁺.

b) trans-{2-[Acryloyl-(3-methyl-but-3-enyl)-amino]-cyclohexyl}-carbamic acid tert-butyl ester

10 The title compound was produced in accordance with the general method of example 29b from trans-[2-(3-methyl-but-3-enylamino)-cyclohexyl]-carbamic acid tert-butyl ester and acryloyl chloride. Off-white solid, MS (ISP) 337.4 (M^+H)⁺.

c) trans-[2-(4-Methyl-6-oxo-3,6-dihydro-2H-pyridin-1-yl)-cyclohexyl]-carbamic acid tert-butyl ester

15 Dichloro(1,3-dimesityl-4,5-dihydroimidazol-2-ylidene)(phenylmethylene)-(tricyclohexylphosphine)ruthenium (54 mg, 63 μ mol) was added to a solution of trans-{2-[acryloyl-(3-methyl-but-3-enyl)-amino]-cyclohexyl}-carbamic acid tert-butyl ester (212 mg, 0.63 mmol) and tetraisopropyl orthotitanate (36 mg, 0.13 mmol) in chloroform (11 ml), and the reaction mixture was heated at reflux for 72 h. The solvent was then
20 evaporated and the residue chromatographed (SiO₂, heptane-ethyl acetate gradient) to produce the title compound (120 mg, 62%). Off-white solid, MS (ISP) 309.1 (M^+H)⁺.

d) trans-1-(2-Amino-cyclohexyl)-4-methyl-5,6-dihydro-1H-pyridin-2-one

The title compound was produced in accordance with the general method of
example 27b from trans-[2-(4-methyl-6-oxo-3,6-dihydro-2H-pyridin-1-yl)-cyclohexyl]-
25 carbamic acid tert-butyl ester. Brown liquid, MS (ISP) 209.2 (M^+H)⁺.

Example 32

trans-1-(2-Amino-cyclohexyl)-piperidin-2-one

The title compound was produced in accordance with the general method of
example 28 from trans-(2-amino-cyclohexyl)-carbamic acid tert-butyl ester and 5-
30 chlorovaleroyl chloride. Colourless liquid, MS (ISP) 197.2 (M^+H)⁺.

Example 33

1-(2-Amino-cyclohexyl)-4-methyl-pyrrolidin-2-one

a) cis-2-(4-Chloro-3-methyl-butyrylamino)-cyclohexanecarboxylic acid ethyl ester

Preparation of ethyl cis-2-amino-1-cyclohexanecarboxylate:

5 Ethyl-cis-2-amino-1-cyclohexanecarboxylate hydrochloride (750 mg) was suspended in 1 N NaOH (pH = 12). The aqueous layer was extracted with CH₂Cl₂, washed with brine, dried over Na₂SO₄ and evaporated to give the crude ethyl-cis-2-amino-1-cyclohexanecarboxylate (570 mg).

The crude ethyl-cis-2-amino-1-cyclohexanecarboxylate (570 mg) was dissolved in
10 CH₂Cl₂ (15 ml) under argon and cooled to 0 °C by means of an ice bath. Triethylamine (0.51 ml) was then added dropwise over a period of 10 min. The mixture was then stirred for 30 min and then treated with 4-chloro-3-methyl-butyryl chloride (568 mg), synthesized according to Chem. Ber., 97, 1964, 2544-2550 dropwise over a period of 10 min; a white suspension was obtained. The resulting mixture was allowed to RT and
15 stirred for 30 min. The mixture was poured into ice/brine and the aqueous layer was extracted with ethyl acetate. The organic layer was separated, washed with brine, dried over Na₂SO₄ and evaporated. The residue was purified by flash chromatography (heptane/ethyl acetate 7:3) to give the product as a 1:1 mixture of epimers as a light yellow oil (913 mg). MS (ESI): 290.1 (M⁺H⁺).

20 b) trans-2-(4-Methyl-2-oxo-pyrrolidin-1-yl)-cyclohexanecarboxylic acid ethyl ester

cis-2-(4-Chloro-3-methyl-butyrylamino)-cyclohexanecarboxylic acid ethyl ester (895 mg) was dissolved in absolute DMF (20 ml) under argon at RT. Sodium iodide (463 mg) and sodium hydride (55 %) (270 mg) were added; a white suspension was obtained. The mixture was then stirred for 2 hours at room temperature. The reaction mixture was
25 poured into ice/water containing saturated NH₄Cl solution and the aqueous layer was extracted with ethyl acetate. The organic layer was separated, washed with brine, dried over Na₂SO₄ and evaporated. The residue was purified by flash chromatography (CH₂Cl₂/MeOH/NH₃ 95/5/0.5) to give the product as a 1:1 mixture of epimers as a light yellow liquid (402 mg). MS (ESI): 254.1 (M⁺H⁺).

30 c) trans- 2-(4-Methyl-2-oxo-pyrrolidin-1-yl)-cyclohexanecarboxylic acid

trans-2-(4-Methyl-2-oxo-pyrrolidin-1-yl)-cyclohexanecarboxylic acid ethyl ester (395 mg) was dissolved in absolute THF (15 ml) and 1 N lithium hydroxide solution (5.12 ml) was added. The resulting mixture was refluxed over night. The reaction

mixture was cooled to room temperature and then HCl conc. (1.80 ml) was added (pH=2). The mixture was evaporated and then diluted with toluene and evaporated to remove the water. The residue was purified by flash chromatography (AcOEt/MeOH 85/15) to give the product as a 1:1 mixture of epimers as light yellow foam (435 mg). MS (ESI): 224.3 (M^+H^+).

d) trans-2-(4-Methyl-2-oxo-pyrrolidin-1-yl)-cyclohexyl]-carbamic acid benzyl ester

trans-2-(4-Methyl-2-oxo-pyrrolidin-1-yl)-cyclohexanecarboxylic acid (100 mg), diphenylphosphorylazide (DPPA) (183 mg), benzylalcohol (0.686 ml) and triethylamine (0.062 ml) were dissolved in absolute toluene (1.0 ml) and the mixture was then heated to 80°C over night. The reaction mixture was then directly evaporated. The residue was purified by flash-chromatography (AcOEt/heptane 80/20) to give the compound as a 1:1 mixture of epimers as white foam (42 mg). MS (ESI): 331.2 (M^+H^+).

e) trans-1-(2-Amino-cyclohexyl)-4-methyl-pyrrolidin-2-one

To a solution of trans-[2-(4-methyl-2-oxo-pyrrolidin-1-yl)-cyclohexyl]-carbamic acid benzyl ester (34 mg) in absolute ethanol (4.0 ml) was added 10% Pd on charcoal (5 mg). A hydrogen atmosphere was introduced by repeated evacuation/gas introduction. The suspension was vigorously stirred over night. The catalyst was removed by filtration through dicalite and the filtrate was concentrated in vacuo. The residue was purified by flash chromatography (CH_2Cl_2 /MeOH/ NH_3 93/7/0.5) to give the product as a 1:1 mixture of epimers as a colorless liquid (15 mg). MS (ESI): 197.3 (M^+H^+).

Galenical Examples

Example A

Film coated tablets containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per tablet</u>	
Kernel:		
Compound of formula (I)	10.0 mg	200.0 mg
Microcrystalline cellulose	23.5 mg	43.5 mg
Lactose hydrous	60.0 mg	70.0 mg
Povidone K30	12.5 mg	15.0 mg
Sodium starch glycolate	12.5 mg	17.0 mg
Magnesium stearate	1.5 mg	4.5 mg
(Kernel Weight)	120.0 mg	350.0 mg
Film Coat:		
Hydroxypropyl methyl cellulose	3.5 mg	7.0 mg
Polyethylene glycol 6000	0.8 mg	1.6 mg
Talc	1.3 mg	2.6 mg
Iron oxide (yellow)	0.8 mg	1.6 mg
Titanium dioxide	0.8 mg	1.6 mg

5

The active ingredient is sieved and mixed with microcrystalline cellulose and the mixture is granulated with a solution of polyvinylpyrrolidone in water. The granulate is mixed with sodium starch glycolate and magnesium stearate and compressed to yield kernels of 120 or 350 mg respectively. The kernels are lacquered with an aq. solution /
 10 suspension of the above mentioned film coat.

Example B

Capsules containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per capsule</u>
Compound of formula (I)	25.0 mg
Lactose	150.0 mg
Maize starch	20.0 mg
Talc	5.0 mg

- 5 The components are sieved and mixed and filled into capsules of size 2.

Example C

Injection solutions can have the following composition:

<u>Ingredients</u>	
Compound of formula (I)	3.0 mg
Polyethylene Glycol 400	150.0 mg
Acetic Acid	q.s. ad pH 5.0
Water for injection solutions	ad 1.0 ml

- 10 The active ingredient is dissolved in a mixture of polyethylene glycol 400 and water for injection (part). The pH is adjusted to 5.0 by acetic acid. The volume is adjusted to 1.0 ml by addition of the residual amount of water. The solution is filtered, filled into vials using an appropriate overage and sterilized.

Example D

Soft gelatin capsules containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	
Capsule contents	
Compound of formula (I)	5.0 mg
Yellow wax	8.0 mg
Hydrogenated Soya bean oil	8.0 mg
Partially hydrogenated plant oils	34.0 mg
Soya bean oil	110.0 mg
Weight of capsule contents	165.0 mg

Gelatin capsule	
Gelatin	75.0 mg
Glycerol 85 %	32.0 mg
Karion 83	8.0 mg (dry matter)
Titanium dioxide	0.4 mg
Iron oxide yellow	1.1 mg

The active ingredient is dissolved in a warm melting of the other ingredients and the mixture is filled into soft gelatin capsules of appropriate size. The filled soft gelatin capsules are treated according to the usual procedures.

Example E


Sachets containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	
Compound of formula (I)	50.0 mg
Lactose, fine powder	1015.0 mg
Microcrystalline cellulose (AVICEL PH 102)	1400.0 mg
Sodium carboxymethyl cellulose	14.0 mg
Polyvinylpyrrolidone K 30	10.0 mg
Magnesium stearate	10.0 mg
Flavoring additives	1.0 mg

- 5 The active ingredient is mixed with lactose, microcrystalline cellulose and sodium carboxymethyl cellulose and granulated with a mixture of polyvinylpyrrolidone in water. The granulate is mixed with magnesium stearate and the flavouring additives and filled into sachets


Claims

1. The use of compounds of the formula (I)



wherein

R^1 is selected from



R^2 , R^3 , R^4 , R^5 and R^6 are each independently selected from hydrogen, C_1 - C_6 - alkyl, halogenated C_1 - C_6 - alkyl, C_1 - C_6 - alkoxy or halogen; provided that R^2 , R^3 , R^4 , R^5 and R^6 are not all hydrogen;

R^7 is C_1 - C_6 - alkyl;

R^8 is C_1 - C_6 - alkyl;

X is $>C=O$ or $>SO_2$;

R^9 and R^{11} are hydrogen or together form a double bond;


R^{10} and R^{12} are independently selected from hydrogen or C_1 - C_6 - alkyl;

m is 1 or 2;

n is 0, 1, or 2;


and pharmaceutically acceptable salts thereof for the preparation of medicaments for the treatment and/or prophylaxis of diseases which are associated with DPP-IV.

2. The use according to claim 1 of compounds of formula I, wherein R is




and wherein R^2 , R^3 , R^4 , R and R are each independently selected from hydrogen, C_1 - C_6 - alkyl, halogenated C_1 - C_6 - alkyl, C_1 - C_6 - alkoxy or halogen; provided that R^2 , R^3 , R^4 , R^5 and R^6 are not all hydrogen.

3. The use according to claim 1 of compounds of formula I, wherein R^1 is




and wherein R^7 is C_1 - C_6 - alkyl.

4. The use according to claim 1 of compounds of formula I, wherein R is



and wherein R is C_1 - C_6 - alkyl.

5. The use according to claim 1 of compounds of formula I, wherein R is



and wherein X is $>C=O$ or $>SO_2$;

R^9 and R^{11} are hydrogen or together form a double bond;

R^{10} and R^{12} are independently selected from hydrogen or C_1 - C_6 - alkyl and

m is 1 or 2.

6. The use according to claim 1 of compounds of formula I selected from

(cis)-2-(5-methyl-thiophen-2-yl)-cyclohexylamine,

(trans)-2-(5-methyl-thiophen-2-yl)-cyclohexylamine,


(trans)-2-(3-methyl-pyrrol-1-yl)-cyclohexylamine,

(trans)-2-(3-ethyl-pyrrol-1-yl)-cyclohexylamine,

(trans)-2-(1,1-dioxo-[1,2] thiazinan-2-yl)-cyclohexylamine,
 (trans)-1-(2-amino-cyclohexyl)-5,6-dihydro-1H-pyridin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-4-methyl-1,5-dihydro-pyrrol-2-one,
 (trans)-1-(2-amino-cyclohexyl)-4-methyl-5,6-dihydro-1H-pyridin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-piperidin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-4-methyl-pyrrolidin-2-one,


and pharmaceutically acceptable salts thereof.

7. Compounds of the formula (I)



wherein

R^1 is selected from



R^7 is C_1 - C_6 - alkyl;

R^8 is C_1 - C_6 - alkyl;

X is $>C=O$ or $>SO_2$;

R^9 and R^{11} are hydrogen or together form a double bond;


R^{10} and R^{12} are independently selected from hydrogen or C_1 - C_6 - alkyl;

m is 1 or 2;

n is 0, 1, or 2;


and pharmaceutically acceptable salts thereof.

8. Compounds of formula I according to claim 7, wherein R^1 is




and wherein R^7 is C_1 - C_6 - alkyl.

9. Compounds of formula I according to claim 7, wherein R^1 is



and wherein R^8 is C_1 - C_6 - alkyl.

10. Compounds of formula I according to claim 7, wherein R^1 is



X is $>C=O$ or $>SO_2$;

R^9 and R^{11} are hydrogen or together form a double bond;

R^{10} and R^{12} are independently selected from hydrogen or C_1 - C_6 - alkyl and m is 1 or 2.

11. Compounds of formula I according to claim 7 selected from

(cis)-2-(5-methyl-thiophen-2-yl)-cyclohexylamine,

(trans)-2-(5-methyl-thiophen-2-yl)-cyclohexylamine,

(trans)-2-(3-methyl-pyrrol-1-yl)-cyclohexylamine,

(trans)-2-(3-ethyl-pyrrol-1-yl)-cyclohexylamine,

(trans)-2-(1,1-dioxo-[1,2]thiazinan-2-yl)-cyclohexylamine,


(trans)-1-(2-amino-cyclohexyl)-5,6-dihydro-1H-pyridin-2-one,

(trans)-1-(2-amino-cyclohexyl)-4-methyl-1,5-dihydro-pyrrol-2-one,

(trans)-1-(2-amino-cyclohexyl)-4-methyl-5,6-dihydro-1H-pyridin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-piperidin-2-one,
 (trans)-1-(2-amino-cyclohexyl)-4-methyl-pyrrolidin-2-one, and
 pharmaceutically acceptable salts thereof.

12. Process for the manufacturing of the compounds of formula I as claimed in claims 7 to 11, comprising


a) a reductive amination of a ketone of formula II



II

wherein R¹ and n are as defined in claim 7, or

b) a deprotection of a carbamic acid ester of formula III



III

wherein R¹ and n are as defined in claim 7 and Rᵖ is an amino protecting group.

13. Pharmaceutical compositions comprising a compound according to any of claims 7 to 11 and a pharmaceutically acceptable carrier and/or adjuvant.

14. Compounds according to any of claims 7 to 11 for use as therapeutic active substances for the treatment and/or prophylaxis of diseases which are associated with DPP-IV.

15. The use of compounds according to any of claims 7 to 11 for the preparation of medicaments for the treatment and/or prophylaxis of diseases which are associated with DPP-IV.

16. The use of compounds according to any of claims 7 to 11 for the preparation of medicaments for the treatment and/or prophylaxis of diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, obesity, and/or metabolic syndrome or β -cell protection.

17. The use according to any of claims 1 to 6 for the preparation of medicaments for the treatment and/or prophylaxis of diabetes, particularly non-insulin dependent diabetes mellitus, impaired glucose tolerance, obesity, and/or metabolic syndrome or β -cell protection.