

Claims

1. An *in vitro* method for detecting HPV-induced high-grade precancerous lesions and HPV-induced invasive cancers and non HPV-induced gynaecological and anogenital cancers, said method comprising detection of hypermethylation in the GHSR and/or ZIC1 gene in a cell
5 whereby such hypermethylation indicates the presence of HPV-induced precursor lesions with invasive potential, HPV-induced invasive cancers and nonHPV-induced gynaecological and anogenital cancers.
2. Method according to claim 1, wherein said HPV-induced high-grade precancerous lesion or HPV-induced invasive carcinoma is a high-
10 grade premalignant cervical lesion or invasive cervical cancer.
3. Method according to claim 1 or 2, wherein said HPV-induced invasive cancer is a high-risk HPV-induced invasive cancer.
4. Method according to claim 1, wherein said nonHPV-induced gynaecological cancers is an endometrial cancer.
- 15 5. Method according to any one of the preceding claims, wherein said hypermethylation is detected in the CpG rich sequences as indicated in Figures 1 and 2.
6. An *in vitro* method of detecting HPV-induced high-grade precancerous lesion and HPV-induced invasive cancers and nonHPV-
20 induced gynaecological or anogenital cancer according to any of claims 1-5, wherein said hypermethylation is an increased methylation of GHSR and/or ZIC1 CpG rich promoter and/or gene sequences in the test cell as compared to the comparable normal cell
7. Method according to any of claims 1-6, wherein the reagent is a
25 methylation sensitive restriction endonuclease, chosen from the group consisting of BssHII, MspI, NotI and HpaII.