

- 1 -

3,7-DIAMINO-10H-PHENOTHIAZINE SALTS AND THEIR USERELATED APPLICATION

- 5 This application is related to United States patent application number 60/786,690 filed 29 March 2006; the contents of which are incorporated herein by reference in their entirety.

TECHNICAL FIELD

10

This invention pertains generally to the field of phenothiazine compounds, and more particularly to certain stably reduced phenothiazine compounds, specifically, certain 3,7-diamino-10H-phenothiazine (DAPTZ) compounds, for example, N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride) and N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen iodide). These compounds are
15 useful as drugs, for example, in the treatment of tauopathies, such as Alzheimer's disease, and also as prodrugs for the corresponding oxidized thioninium drugs (for example, methythioninium chloride, MTC).

20

BACKGROUND

A number of patents and publications are cited herein in order to more fully describe and disclose the invention and the state of the art to which the invention pertains. Each of these references is incorporated herein by reference in its entirety into the present
25 disclosure, to the same extent as if each individual reference was specifically and individually indicated to be incorporated by reference.

Throughout this specification, including the claims which follow, unless the context requires otherwise, the word "comprise," and variations such as "comprises" and
30 "comprising," will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

It must be noted that, as used in the specification and the appended claims, the singular
35 forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a pharmaceutical carrier" includes mixtures of two or more such carriers, and the like.

Ranges are often expressed herein as from "about" one particular value, and/or to "about"
40 another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values

- 2 -

are expressed as approximations, by the use of the antecedent "about," it will be understood that the particular value forms another embodiment.

5 Conditions of dementia are frequently characterised by a progressive accumulation of intracellular and/or extracellular deposits of proteinaceous structures such as β -amyloid plaques and neurofibrillary tangles (NFTs) in the brains of affected patients. The appearance of these lesions largely correlates with pathological neurofibrillary degeneration and brain atrophy, as well as with cognitive impairment (see, e.g., Mukaetova-Ladinska, E.B. et al., 2000, Am. J. Pathol., Vol. 157, No. 2, pp. 623-636).
10 Methythioninium chloride (MTC) and other diaminophenothiazines have been described as inhibitors of protein aggregation in such diseases, that is, diseases in which proteins aggregate pathologically (see, for example, WO 96/30766 and WO 02/055720).

Methythioninium chloride (MTC) is currently used to treat methemoglobinemia
15 (a condition that occurs when the blood cannot deliver oxygen where it is needed in the body). MTC is also used as a medical dye (for example, to stain certain parts of the body before or during surgery); a diagnostic (for example, as an indicator dye to detect certain compounds present in urine); a mild urinary antiseptic; a stimulant to mucous surfaces; a treatment and preventative for kidney stones; and in the diagnosis and treatment of
20 melanoma.

MTC has been used to treat malaria either singly (see, e.g., Guttman, P. and Ehrlich, P., 1891, "Über die Wirkung des methylenblau bei malaria," Berl. Klin. Wochenschr., Vol. 28, pp. 953-956) or in combination with chloroquine (see, e.g., Schirmer, H., et al., 2003,
25 "Methylene blue as an antimalarial agent," Redox Report, Vol. 8, pp. 272-275; Rengelshausen, J., et al., 2004, "Pharmacokinetic interaction of chloroquine and methylene blue combination against malaria," European Journal of Clinical Pharmacology, Vol. 60, pp. 709-715). Malaria in humans is caused by one of four protozoan species of the genus Plasmodium: *P. falciparum*, *P. vivax*, *P. ovale*, or *P.*
30 *malariae*. All species are transmitted by the bite of an infected female Anopheles mosquito. Occasionally, transmission occurs by blood transfusion, organ transplantation, needle-sharing, or congenitally from mother to fetus. Malaria causes 300-500 million infections worldwide and approximately 1 million deaths annually. Drug resistance, however, is a major concern and is greatest for *P. falciparum*, the species that accounts
35 for almost all malaria-related deaths. Drugs or drug combinations that are currently recommended for prophylaxis of malaria include chloroquine/proguanil hydrochloride, mefloquine, doxycycline, and primaquine.

MTC (under the name Virostat®, from Bioenvision Inc., New York) has shown potent
40 viricidal activity in vitro. Specifically Virostat® is effective against viruses such as HIV and West Nile Virus in laboratory tests. West Nile virus (WNV) is a potentially serious illness

- 3 -

affecting the central nervous system. The large majority of infected people will show no visible symptoms or mild flu-like symptoms such as fever and headache. About one in 150 will develop severe symptoms including tremors, convulsions, muscle weakness, vision loss, numbness, paralysis, or coma. Generally, WNV is spread by the bite of an infected mosquito, but can also spread through blood transfusions, organ transplants, breastfeeding or during pregnancy from mother to child.

Virostat® is also currently in clinical trials for the treatment of chronic Hepatitis C. Hepatitis C is a viral infection of the liver. The virus, HCV, is a major cause of acute hepatitis and chronic liver disease, including cirrhosis and liver cancer. HCV is spread primarily by direct contact with human blood. The major causes of HCV infection worldwide are use of unscreened blood transfusions, and re-use of needles and syringes that have not been adequately sterilized. The World Health Organization has declared hepatitis C a global health problem, with approximately 3% of the world's population infected with HCV and it varies considerably by region. The prevalence in the US is estimated at 1.3% or approximately 3.5 million people. Egypt has a population of approximately 62 million and contains the highest prevalence of hepatitis C in the world, estimated at over 20% of the nation's approximately 62 million people.

MTC, when combined with light, can prevent the replication of nucleic acid (DNA or RNA). Plasma, platelets and red blood cells do not contain nuclear DNA or RNA. When MTC is introduced into the blood components, it crosses bacterial cell walls or viral membrane then moves into the interior of the nucleic acid structure. When activated with light, the compound then binds to the nucleic acid of the viral or bacterial pathogen, preventing replication of the DNA or RNA. Because MTC can inactivate pathogens, it has the potential to reduce the risk of transmission of pathogens that would remain undetected by testing.


Oral and parenteral formulations of MTC are commercially available in the United States, usually under the name Urolene Blue®.

SUMMARY OF THE INVENTION

Disclosed are compounds, specifically, certain 3,7-diamino-10H-phenothiazine (DAPTZ) compounds, as described herein.

5

In one aspect, there is provided a compound selected from compounds of the following formula and pharmaceutically acceptable salts, solvates, and hydrates thereof:



wherein:

10

each of R^1 and R^9 is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

each of R^{3NA} and R^{3NB} is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

each of R^{7NA} and R^{7NB} is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

15

each of HX^1 and HX^2 is independently a protic acid.

Also disclosed is a composition comprising a DAPTZ compound as described herein and a pharmaceutically acceptable carrier or diluent.

20

Also disclosed is a pharmaceutical composition comprising a DAPTZ compound as described herein and a pharmaceutically acceptable carrier or diluent.

In another aspect, there is provided a method of preparing a pharmaceutical composition comprising admixing a DAPTZ compound as described herein and a pharmaceutically acceptable carrier or diluent.

25

In yet another aspect, there is provided a method of reversing and/or inhibiting the aggregation of a protein (e.g., a tau protein, a synuclein, etc.), for example, aggregation of a protein associated with a neurodegenerative disease and/or clinical dementia, comprising contacting the protein with an effective amount of a DAPTZ compound, as described herein. Such a method may be performed in vitro, or in vivo.

30

Also disclosed is a method of treatment or prophylaxis of a disease condition in a subject comprising administering to said subject a prophylactically or therapeutically effective amount of a DAPTZ compound, as described herein, preferably in the form of a pharmaceutical composition.

In yet another aspect, there is provided a DAPTZ compound as described herein for use in a method of treatment or prophylaxis (e.g., of a disease condition) of the human or animal body by therapy.

In yet another aspect, there is provided the use of a DAPTZ compound, as described herein, in the manufacture of a medicament for use in the treatment or prophylaxis of a disease condition.

In one embodiment, the disease condition is a disease of protein aggregation.

In one embodiment, the disease condition is a tauopathy, e.g., a neurodegenerative tauopathy, e.g., Alzheimer's disease.

In one embodiment, the disease condition is skin cancer, e.g., melanoma.

In one embodiment, the disease condition is a viral, bacterial or protozoal disease condition, e.g., Hepatitis C, HIV, West Nile Virus (WNV), or malaria.

- 5 In yet another aspect, there is provided an *in vitro* method of reversing and/or inhibiting the aggregation of a protein comprising contacting the protein with an effective amount of a compound as defined above.

- 10 In yet another aspect, there is provided a method of inactivating a pathogen in a sample (for example a blood or plasma sample), comprising the steps of introducing a DAPTZ compound, as described herein, into the sample, and then exposing the sample to light.

- Also disclosed is a kit comprising (a) a DAPTZ compound as described herein, preferably provided as a pharmaceutical composition and in a suitable container and/or with suitable
15 packaging; and (b) instructions for use, for example, written instructions on how to administer the compound.

- As will be appreciated by one of skill in the art, features and preferred embodiments of one aspect of the invention will also pertain to other aspect of the invention.
20

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph of the percent reduced form (%) versus time (minutes) for each of three compounds, B1 (MTC), B3 (N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride)), and B6 (N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen iodide)), as determined using absorbance at 665 nm.

Figure 2 is a graph of the percent reduced form (%) versus time (minutes) for each of three compounds, B1, B3, and B6, as determined using absorbance at 610 nm.

Figure 3A shows the UV/visible absorption spectra for aqueous samples of each of three compounds, B1 (open circles, maximum at 605 nm), B3 (open squares, maximum at 660 nm), and B6 (open triangles, maximum at 660 nm), after 20 minutes.

Figure 3B shows the UV/visible absorption spectra for aqueous samples of each of three compounds, B1 (open circles, maximum at 605 nm), B3 (open squares, maximum at 605 nm), and B6 (open triangles, maximum at 605 nm), after 3 hours.

Figure 3C shows the UV/visible absorption spectra for aqueous samples of each of three compounds, B1 (open circles, maximum at 605 nm), B3 (open squares, maximum at 605 nm), and B6 (open triangles, maximum at 605 nm), after 28 hours.

Figure 4 shows the crystal structure of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide).


Figure 5 shows the the side-on view of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide).

Figure 6 shows part of one helical column of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide) molecules in the crystal.

- 7 -


DETAILED DESCRIPTION OF THE INVENTION

Methythioninium Chloride (MTC) (also known as Methylene blue (MB); methythionine chloride; tetramethylthionine chloride; 3,7-bis(dimethylamino) phenothiazin-5-ium chloride; C.I. Basic Blue 9; tetramethylthionine chloride; 3,7-bis(dimethylamino) phenazathionium chloride; Swiss blue; C.I. 52015; C.I. Solvent Blue 8; aniline violet; and Urolene Blue®) is a low molecular weight (319.86), water soluble, tricyclic organic compound of the following formula:



Methythioninium Chloride (MTC) (also known as Methylene Blue), perhaps the most well known phenothiazine dye and redox indicator, has also been used as an optical probe of biophysical systems, as an intercalator in nanoporous materials, as a redox mediator, and in photoelectrochromic imaging.

MTC, a phenothiazin-5-ium salt, may conveniently be considered to be an "oxidized form" when considered in respect of the corresponding 10H-phenothiazine compound, N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine, which may conveniently be considered to be a "reduced form":



The "reduced form" (the "leuko form") is known to be unstable, and is readily and rapidly oxidized to give the corresponding "oxidized" form.

- 8 -

May et al. (Am J Physiol Cell Physiol, 2004, Vol. 286, pp. C1390-C1398) have shown that human erythrocytes sequentially reduce and take up MTC; that MTC itself is not taken up by the cells; that it is the reduced form of MTC that crosses the cell membrane; that the rate of uptake is enzyme dependent; and that both MTC and reduced MTC are
5 concentrated in cells (reduced MTC re-equilibrates once inside the cell to form MTC).

MTC and similar drugs are taken up in the gut and enter the bloodstream. Unabsorbed drug percolates down the alimentary canal, to the distal gut. One important undesired side-effect is the effect of the unabsorbed drug in the distal gut, for example, sensitisation
0 of the distal gut and/or antimicrobial effects of the unabsorbed drug on flora in the distal gut, both leading to diarrhoea. Therefore, it is desirable to minimize the amount of drug that percolates to the distal gut. By increasing the drug's uptake in the gut (i.e., by increasing the drug's bioavailability), dosage may be reduced, and the undesired side-effects, such as diarrhoea, may be ameliorated.


5 Since it is the reduced form of MTC that is taken up by cells, it would be desirable to administer the reduced form. This would also reduced reliance on the rate limiting step of enzymatic reduction.

20 The inventors have identified a class of compounds that may also be considered to be in the "reduced form" when considered in respect of MTC, and which are surprisingly and unexpectedly stable. The compounds may therefore be described as "stabilized reduced forms," for example, of MTC.


25 These compounds are themselves active as drugs, and may also serve as prodrugs, yielding, upon oxidation, the corresponding oxidized compounds (e.g., MTC), which are also active as drugs.

- 9 -

One representative member of this class of compounds is shown below.




Another representative member of this class of compounds is shown below.



- 10 -

The Compounds

In general, the present invention pertains certain 3,7-diamino-10H-phenothiazine compounds of the following formula (collectively referred to herein as "diamino-phenothiazine compounds" and/or "DAPTZ compounds"):



wherein:

each of R^1 and R^9 is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;


each of $\text{R}^{3\text{NA}}$ and $\text{R}^{3\text{NB}}$ is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

each of $\text{R}^{7\text{NA}}$ and $\text{R}^{7\text{NB}}$ is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

each of HX^1 and HX^2 is independently a protic acid;

and pharmaceutically acceptable salts, solvates, and hydrates thereof.

Without wishing to be bound to any particular theory, the inventors believe that it is possible, if not likely, that the compounds exist in the following form:



Although the DAPTZ compounds are themselves salts, they may also be provided in the form of a mixed salt (i.e., the DAPTZ in combination with another salt). Such mixed salts are intended to be encompassed by the term "and pharmaceutically acceptable salts thereof". Unless otherwise specified, a reference to a particular compound also includes salts thereof.

The DAPTZ compounds may also be provided in the form of a solvate or hydrate. The term "solvate" is used herein in the conventional sense to refer to a complex of solute (e.g., compound, salt of compound) and solvent. If the solvent is water, the solvate may be conveniently referred to as a hydrate, for example, a mono-hydrate, a di-hydrate, a

tri-hydrate, etc. Unless otherwise specified, a reference to a particular compound also includes solvate forms thereof.

5 In one embodiment, the C₁₋₄alkyl groups are selected from: linear C₁₋₄alkyl groups, such as -Me, -Et, -nPr, -iPr, and -nBu; branched C₃₋₄alkyl groups, such as -iPr, -iBu, -sBu, and -tBu; and cyclic C₃₋₄alkyl groups, such as -cPr and -cBu.

0 In one embodiment, the C₂₋₄alkenyl groups are selected from linear C₁₋₄alkenyl groups, such as -CH=CH₂ (vinyl) and -CH₂-CH=CH₂ (allyl).

0 In one embodiment, the halogenated C₁₋₄alkyl groups are selected from: -CF₃, -CH₂CF₃, and -CF₂CF₃.

15 *The Groups R¹ and R⁹*

In one embodiment, each of R¹ and R⁹ is independently -H, -Me, -Et, or -CF₃.

In one embodiment, each of R¹ and R⁹ is independently -H, -Me, or -Et.

In one embodiment, R¹ and R⁹ are the same.

20 In one embodiment, R¹ and R⁹ are different.

In one embodiment, each of R¹ and R⁹ is independently -H.

In one embodiment, each of R¹ and R⁹ is independently -Me.

In one embodiment, each of R¹ and R⁹ is independently -Et.

25

The Groups R^{3NA} and R^{3NB}

Each of R^{3NA} and R^{3NB} is independently selected from: -H, C₁₋₄alkyl, C₂₋₄alkenyl, and halogenated C₁₋₄alkyl.

30

In one embodiment, each of R^{3NA} and R^{3NB} is independently selected from: C₁₋₄alkyl, C₂₋₄alkenyl, and halogenated C₁₋₄alkyl.

In one embodiment, each of R^{3NA} and R^{3NB} is independently -Me, -Et, -nPr, -nBu,

35 -CH₂-CH=CH₂, or -CF₃.

In one embodiment, each of R^{3NA} and R^{3NB} is independently -Me, -nPr, -nBu,

-CH₂-CH=CH₂, or -CF₃.

40 In one embodiment, each of R^{3NA} and R^{3NB} is independently -Me or -Et.

- 12 -

In one embodiment, R^{3NA} and R^{3NB} are the same.

In one embodiment, R^{3NA} and R^{3NB} are different.

In one embodiment, each of R^{3NA} and R^{3NB} is independently -Me.

5 In one embodiment, each of R^{3NA} and R^{3NB} is independently -Et.

The Groups R^{7NA} and R^{7NB}

10 Each of R^{7NA} and R^{7NB} is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl.

In one embodiment, each of R^{7NA} and R^{7NB} is independently selected from: C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl.

15 In one embodiment, each of R^{7NA} and R^{7NB} is independently -Me, -Et, -nPr, -nBu, $-CH_2-CH=CH_2$, or $-CF_3$.

In one embodiment, each of R^{7NA} and R^{7NB} is independently -Me, -nPr, -nBu, $-CH_2-CH=CH_2$, or $-CF_3$.

20

In one embodiment, each of R^{7NA} and R^{7NB} is independently -Me or -Et.

In one embodiment, R^{7NA} and R^{7NB} are the same.

In one embodiment, R^{7NA} and R^{7NB} are different.

25

In one embodiment, each of R^{7NA} and R^{7NB} is independently -Me.

In one embodiment, each of R^{7NA} and R^{7NB} is independently -Et.

In one embodiment, R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} are the same.

30

In one embodiment, R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} are as defined herein, with the proviso that at least one of R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} is *other than* -Et.

Optional Provisos

35

In one embodiment, the compound is as defined herein, but with the proviso that: R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} are not each -Et.

In one embodiment, the compound is as defined herein, but with the proviso that:

40 if: each of R^1 and R^9 is -H;

then: R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} are not each -Et.

The Groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$

In one embodiment:

- 5 each of R^{3NA} and R^{3NB} is independently C_{1-4} alkyl, C_{2-4} alkenyl, or halogenated C_{1-4} alkyl; each of R^{7NA} and R^{7NB} is independently C_{1-4} alkyl, C_{2-4} alkenyl, or halogenated C_{1-4} alkyl; optionally with the proviso that at least one of R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} is other than -Et.

10 In one embodiment:

each of R^{3NA} and R^{3NB} is independently -Me, -Et, -nPr, -nBu, $-CH_2-CH=CH_2$, or $-CF_3$; each of R^{7NA} and R^{7NB} is independently -Me, -Et, -nPr, -nBu, $-CH_2-CH=CH_2$, or $-CF_3$; optionally with the proviso that at least one of R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} is other than -Et.

15

In one embodiment:

each of R^{3NA} and R^{3NB} is independently -Me or -Et; each of R^{7NA} and R^{7NB} is independently -Me or -Et; optionally with the proviso that at least one of R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} is other than -Et.

20

In one embodiment, the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ are the same.

In one embodiment, the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ are different.

25

In one embodiment, each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently selected from: $-NMe_2$, $-NEt_2$, $-N(nPr)_2$, $-N(Bu)_2$, $-NMeEt$, $-NMe(nPr)$, and $-N(CH_2CH=CH_2)_2$.

- 30 In one embodiment, the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ are the same, and are independently selected from: $-NMe_2$, $-NEt_2$, $-N(nPr)_2$, $-N(Bu)_2$, $-NMeEt$, $-NMe(nPr)$, and $-N(CH_2CH=CH_2)_2$.

- 35 In one embodiment, the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ are the same, and are independently selected from: $-NMe_2$ and $-NEt_2$.

In one embodiment, each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is: $-NMe_2^+$.

- 40 In one embodiment, at least one of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is other than $-NEt_2$.

- 14 -

In one embodiment, each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is *other than* $-NEt_2$.

- 5 For example, in one embodiment, the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ are the same, and are selected from: $-NMe_2$, $-N(nPr)_2$, $-N(Bu)_2$, $-NMeEt$, $-NMe(nPr)$, and $-N(CH_2CH=CH_2)_2$.

The Groups HX^1 and HX^2

- 10 Each of HX^1 and HX^2 is independently a protic acid.

Examples of protic acids include, for example, inorganic acids, such as hydrohalide acids (e.g., HCl, HBr, HI), nitric acid (HNO_3), sulphuric acid (H_2SO_4), and organic acids, such as carbonic acid (H_2CO_3) and acetic acid (CH_3COOH).

15

In one embodiment, each of HX^1 and HX^2 is independently a monoprotic acid.

In one embodiment, each of HX^1 and HX^2 is independently a hydrohalide acid (i.e., a hydrohalic acid)

20

In one embodiment, each of HX^1 and HX^2 is independently selected from HCl, HBr, and HI.

In one embodiment, HX^1 and HX^2 are the same.

- 25 In one embodiment, HX^1 and HX^2 are different.

In one embodiment, HX^1 and HX^2 are the same, and are independently selected from HCl, HBr, and HI. In this case, the compound (a diamino-phenothiazine compound) may conveniently be referred to as a "diamino-phenothiazine bis(hydrogen halide) salt".

30

In one embodiment, HX^1 and HX^2 are each HCl. In this case, the compound may conveniently be referred to as a "diamino-phenothiazine bis(hydrogen chloride) salt".

- 35 In one embodiment, HX^1 and HX^2 are each HBr. In this case, the compound may conveniently be referred to as a "diamino-phenothiazine bis(hydrogen bromide) salt".

In one embodiment, HX^1 and HX^2 are each HI. In this case, the compound may conveniently be referred to as a "diamino-phenothiazine bis(hydrogen iodide) salt".

- 15 -

Some Preferred Combinations

In one embodiment:

each of R^1 and R^9 is independently -H, -Me, or -Et; and

- 5 each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently $-NMe_2$ or $-NEt_2$.

In one embodiment:

each of R^1 and R^9 is independently -H, -Me, or -Et; and

- 10 each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently $-NMe_2$.

In one embodiment:

each of R^1 and R^9 is independently -H; and

- each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently $-NMe_2$ or $-NEt_2$.
- 15 In one embodiment:
- each of R^1 and R^9 is independently -H; and
- each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently $-NMe_2$.

In one embodiment:

- 20 each of R^1 and R^9 is independently -H, -Me, or -Et; and
- each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently $-NMe_2$ or $-NEt_2$;
- and
- each of HX^1 and HX^2 is independently selected from HCl, HBr, and HI.

- 25 In one embodiment:

each of R^1 and R^9 is independently -H, -Me, or -Et; and

each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently $-NMe_2$; and

each of HX^1 and HX^2 is independently selected from HCl, HBr, and HI.

- 30 In one embodiment:

each of R^1 and R^9 is independently -H; and

each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently $-NMe_2$ or $-NEt_2$;

and

each of HX^1 and HX^2 is independently selected from HCl, HBr, and HI.

35

In one embodiment:

each of R^1 and R^9 is independently -H; and

each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently $-NMe_2$; and

each of HX^1 and HX^2 is independently selected from HCl, HBr, and HI.

40


- 16 -

In one embodiment:

each of R^1 and R^9 is independently -H; and

each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently -NMe₂; and

each of HX¹ and HX² is HCl.




5

In one embodiment:

each of R^1 and R^9 is independently -H; and

each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently -NMe₂; and

each of HX¹ and HX² is HBr.




10

In one embodiment:

each of R^1 and R^9 is independently -H; and

each of the groups $-N(R^{3NA})(R^{3NB})$ and $-N(R^{7NA})(R^{7NB})$ is independently -NMe₂; and

each of HX¹ and HX² is HI.



15

Isotopic Variation

20

In one embodiment, one or more of the carbon atoms of the compound is ¹¹C, ¹³C, or ¹⁴C.

In one embodiment, one or more of the carbon atoms of the compound is ¹¹C.

In one embodiment, one or more of the carbon atoms of the compound is ¹³C.

In one embodiment, one or more of the carbon atoms of the compound is ¹⁴C.

- 17 -

In one embodiment, one or more of the nitrogen atoms of the compound is ^{15}N .

In one embodiment, one or more or all of the carbon atoms of one or more or all of the groups $\text{R}^{3\text{NA}}$, $\text{R}^{3\text{NB}}$, $\text{R}^{7\text{NA}}$, $\text{R}^{7\text{NB}}$, R^1 , R^9 , and R^{10} is ^{11}C . (Or ^{13}C .) (Or ^{14}C .)

5

In one embodiment, one or more or all of the carbon atoms of one or more or all of the groups $\text{R}^{3\text{NA}}$, $\text{R}^{3\text{NB}}$, $\text{R}^{7\text{NA}}$, and $\text{R}^{7\text{NB}}$ is ^{11}C . (Or ^{13}C .) (Or ^{14}C .)




10 In one embodiment, the groups $-\text{N}(\text{R}^{3\text{NA}})(\text{R}^{3\text{NB}})$ and $-\text{N}(\text{R}^{7\text{NA}})(\text{R}^{7\text{NB}})$ are the same, and are: $-\text{N}(\text{}^{11}\text{CH}_3)_2$. (Or $-\text{N}(\text{}^{13}\text{CH}_3)_2$.) (Or $-\text{N}(\text{}^{14}\text{CH}_3)_2$.)

Compatible Combinations







15 All compatible combinations of the embodiments described above are explicitly disclosed herein as if each combination was specifically and individually recited.






Some Preferred Embodiments

20 In one embodiment, the compound is selected from the following compounds, and pharmaceutically acceptable salts, solvates, and hydrates thereof.






1	
2	
3	

- 18 -







4	 HCl HCl
5	 HCl HCl
6	 HCl HCl
7	 HCl HCl
8	 HCl HCl
9	 HCl HCl

10	 HCl HCl
11	 HCl HCl
12	 HCl HCl
13	 HCl HCl
14	 HCl HCl




In one embodiment, the compound is selected from the following compounds, and pharmaceutically acceptable salts, solvates, and hydrates thereof.

15	 <chem>CN(C)c1ccc2c(c1)sc3ccncc32</chem>	HBr HBr
16	 <chem>CCN(CC)c1ccc2c(c1)sc3ccncc32</chem>	HBr HBr
17	 <chem>CCCN(CCC)c1ccc2c(c1)sc3ccncc32</chem>	HBr HBr
18	 <chem>CCCCN(CCC)c1ccc2c(c1)sc3ccncc32</chem>	HBr HBr
19	 <chem>C=CCN(CC=C)c1ccc2c(c1)sc3ccncc32</chem>	HBr HBr



- 21 -







20	 HBr HBr
21	 HBr HBr
22	 HBr HBr
23	 HBr HBr
24	 HBr HBr
25	 HBr HBr

- 22 -







26	 HBr HBr
27	 HBr HBr
28	 HBr HBr

In one embodiment, the compound is selected from the following compounds, and pharmaceutically acceptable salts, solvates, and hydrates thereof.

29	 HI HI
30	 HI HI

31	 HI HI
32	 HI HI
33	 HI HI
34	 HI HI
35	 HI HI
36	 HI HI

- 24 -

37	 HI HI
38	 HI HI
39	 HI HI
40	 HI HI
41	 HI HI
42	 HI HI

- 25 -

Purity

The DAPTZ compounds of the present invention may conveniently be described as being in a "stabilized reduced form". The compounds oxidize (e.g., autoxidize) to give the
5 corresponding oxidized forms. Thus, it is likely, if not inevitable, that compositions comprising the DAPTZ compounds of the present invention will contain, as an impurity, as least some of the corresponding oxidized compound.

Thus, another aspect of the present invention pertains to DAPTZ compounds, as
0 described herein, in substantially purified form and/or in a form substantially free from contaminants (e.g., the corresponding oxidized compound, other contaminants).

In one embodiment, the substantially purified form is at least 50% by weight pure, e.g., at least 60% by weight pure, e.g., at least 70% by weight pure, e.g., at least 80% by weight
5 pure, e.g., at least 90% by weight pure, e.g., at least 95% by weight pure, e.g., at least 97% by weight pure, e.g., at least 98% by weight pure, e.g., at least 99% by weight pure.

In one embodiment, the contaminants represent no more than 50% by weight, e.g., no more than 40% by weight, e.g., no more than 30% by weight, e.g., no more than 20% by
20 weight, e.g., no more than 10% by weight, e.g., no more than 5% by weight, e.g., no more than 3% by weight, e.g., no more than 2% by weight, e.g., no more than 1% by weight.

Product-by-Process


25 In one embodiment, the DAPTZ compound is one which is *obtained by, or is obtainable by*, a method as described herein.

Chemical Synthesis

Methods for the chemical synthesis of DAPTZ compounds of the present invention are described herein. These and/or other well known methods may be modified and/or adapted in known ways in order to facilitate the synthesis of additional DAPTZ compounds within the scope of the present invention.


For example, a suitable phenothiazine may be converted to the corresponding 3,7-dinitro-phenothiazine, for example, using sodium nitrite with acetic acid and chloroform. The ring amino group may then be protected, for example, as the acetate, for example, using acetic anhydride and pyridine. The nitro groups may then be reduced to amino groups, for example, using tin (II) chloride with ethanol. The amino groups may then be substituted, for example, disubstituted, for example, methyl disubstituted, for example, using methyl iodide, sodium hydroxide, DMSO, and tetra-n-butyl ammonium bromide. The amino group may then be deprotected, for example, the N-acetyl group may be removed, for example, using concentrated aqueous hydrochloride acid. The corresponding salt is then prepared, for example, using concentrated aqueous hydrochloric acid, for example, at the same time as deprotection. An example of such a method is illustrated in the following scheme.

20

Scheme 1

- 27 -

Thus, another aspect of the invention pertains to a method of preparing a 3,7-diamino-10H-phenothiazine (DAPTZ) compound of the following formula:



wherein R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} , R^{7NB} , HX^1 and HX^2 are as defined herein (for example, where HX^1 and HX^2 are each HCl), comprising the step of:

(vi) salt formation (SF).

In one embodiment, the method comprises the steps of:

(v) ring amino deprotection (DP); and

(vi) salt formation (SF).

In one embodiment, the method comprises the steps of:

(iv) amine substitution (AS),

optional (v) ring amino deprotection (DP), and

(vi) salt formation (SF).

In one embodiment, the method comprises the steps of

(iii) nitro reduction (NR),

(iv) amine substitution (AS),

(v) ring amino deprotection (DP), and

(vi) salt formation (SF).

In one embodiment, the method comprises the steps of

optional (ii) ring amino protection (AP),

(iii) nitro reduction (NR),

(iv) amine substitution (AS),

(v) ring amino deprotection (DP), and

(vi) salt formation (SF).

In one embodiment, the method comprises the steps of

(i) nitration (NO),

(ii) ring amino protection (AP),

(iii) nitro reduction (NR),

(iv) amine substitution (AS),

(v) ring amino deprotection (DP), and

- 28 -

(vi) salt formation (SF).

In one embodiment, the steps are performed in the order listed (i.e., any step in the list is performed at the same time as, or subsequent to, the preceding step in the list).


5

In one embodiment, the step of (v) ring amino deprotection (DP) and the step of (vi) salt formation (SF) are performed simultaneously (i.e., as one step).

In one embodiment, the nitration (NO) step is:

10

(i) nitration (NO), wherein a 10H-phenothiazine is converted to a 3,7-dinitro-10H-phenothiazine, for example:




15

In one embodiment, nitration is performed using a nitrite, for example, sodium nitrite, for example, sodium nitrite with acetic acid and chloroform. In one embodiment, R^{10} is -H.

In one embodiment, the ring amino protection (AP) step is:

20

(ii) ring amino protection (AP), wherein the ring amino group (-NH-) of a 3,7-dinitro-10H-phenothiazine is converted to a protected ring amino group (-NR^{prot}), for example:




25

In one embodiment, ring amino protection is achieved as an acetate, for example, using acetic anhydride, for example, using acetic anhydride and pyridine.

In one embodiment, the nitro reduction (NR) step is:

(iii) nitro reduction (NR), wherein each of the nitro (-NO₂) groups of a protected 3,7-dinitro-10H-phenothiazine is converted to an amino (-NH₂) group, for example:

- 29 -




In one embodiment, nitro reduction may be performed using, for example, tin (II) chloride, for example, tin (II) chloride with ethanol.

5

In one embodiment, the amine substitution (AS) step is:

(iv) amine substitution (AS), wherein each of the amino ($-NH_2$) groups of a protected 3,7-diamino-10H-phenothiazine is converted to disubstituted amino group, for example:




10

In one embodiment, amine substitution is performed using an alkyl halide, for example, an alkyl iodide, for example, methyl iodide, for example, methyl iodide with sodium hydroxide, DMSO, and tetra-n-butyl ammonium bromide.

15

In one embodiment, the ring amino deprotection (DP) step is:

(v) ring amino deprotection (DP), wherein the protecting group, R^{Prot} , is removed, for example:



20


In one embodiment, ring amino deprotection may be performed using acid, for example, hydrochloric acid, for example, concentrated aqueous hydrochloric acid.

In one embodiment, the step is:

25

(vi) salt formation (SF), wherein the corresponding salt is formed, for example:

- 30 -




In one embodiment, salt formation may be performed using acid, for example, hydrochloric acid, for example, concentrated aqueous hydrochloric acid.

5

In one embodiment, the steps of ring amine deprotection and salt formation are performed simultaneously (i.e., as one step), for example, compound (1) is formed from compound (2) in one step.


- 10 In another approach, a suitable thioninium chloride (for example, methylthioninium chloride, MTC, also known as Methylene Blue) is converted to the corresponding halide, for example, by reaction with potassium iodide, for example, aqueous potassium iodide. The resulting thioninium iodide is then reduced, for example, with ethyl iodide and ethanol, and the corresponding salt formed. A similar method is described in Drew, H.D.K. and Head, F.S.H., "Derivatives of Methylene-blue," Journal of the Chemical Society, 1933, pp. 248-253. An example of such a method is illustrated in the following scheme.
- 15



20

- 31 -

Thus, another aspect of the invention pertains to a method of preparing a 3,7-diamino-10H-phenothiazine (DAPTZ) compound of the following formula:



- wherein R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} , R^{7NB} , HX^1 and HX^2 are as defined herein (for example, where HX^1 and HX^2 are each HI), comprising the step of:
- (ii) reduction and iodide salt formation (RISF).


In one embodiment, the method comprises the steps of:

- (i) iodide exchange (IE); and
- (ii) reduction and iodide salt formation (RISF).

In one embodiment, the steps are performed in the order listed (i.e., any step in the list is performed at the same time as, or subsequent to, the preceding step in the list).

- 15 In one embodiment, the iodide exchange (IE) step is:

(i) iodide exchange (IE), wherein a 3,7-di(disubstituted amino)-thioninium salt is converted to the corresponding 3,7-di(disubstituted amino)-thioninium iodide, for example (where Y^- is an anionic counter ion, for example, halide, for example, chloride or bromide):




In one embodiment, iodide exchange (IE) is achieved by reaction with potassium iodide, for example, aqueous potassium iodide.

- 32 -

In one embodiment, the reduction and iodide salt formation (RISF) step is:

(ii) reduction and iodide salt formation (RISF), wherein a 3,7-di(disubstituted amino)-thioninium iodide is reduced and converted to the corresponding 3,7-diamino-10H-phenothiazine iodide compound, for example:




5

In one embodiment, reduction and iodide salt formation (RISF) is achieved by reaction with ethyl iodide, for example, ethyl iodide and ethanol.


- 10 In another approach, an appropriate thioninium salt, for example, ethyl thioninium semi zinc chloride, is simultaneously reduced and the ring amino group protected, for example, by reaction with phenylhydrazine, ethanol, acetic anhydride, and pyridine. The corresponding salt may then be prepared, for example, using concentrated aqueous hydrochloric acid, for example, at the same time as deprotection. An example of such a
- 15 method is illustrated in the following scheme.

Scheme 3



- 33 -

Thus, another aspect of the invention pertains to a method of preparing a 3,7-diamino-10H-phenothiazine (DAPTZ) compound of the following formula:



wherein R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} , R^{7NB} , HX^1 and HX^2 are as defined herein (for example, where HX^1 and HX^2 are each HI), comprising the step of:

comprising the step of:

(iv) salt formation (SF).

In one embodiment, the method comprises the steps of

(iii) ring amino deprotection (DP), and

(iv) salt formation (SF).

In one embodiment, the method comprises the steps of

(ii) ring amino protection (AP),

(iii) ring amino deprotection (DP), and

(iv) salt formation (SF).

In one embodiment, the method comprises the steps of

(i) reduction (RED)

(ii) ring amino protection (AP),

(iii) ring amino deprotection (DP), and

(iv) salt formation (SF).

In one embodiment, the steps are performed in the order listed (i.e., any step in the list is performed at the same time as, or subsequent to, the preceding step in the list).


In one embodiment, the step of (i) reduction (RED) and the step of (ii) ring amino protection (AP) are performed simultaneously (i.e., as one step).

For example, in one embodiment, the combined reduction (RED) step and ring amino protection (AP) step is:

(i) reduction (RED) and ring amino protection (AP), wherein a 3,7-di(disubstituted amino)-thionium salt is reduced to give the corresponding 3,7-di(disubstituted amino)-10H-phenothiazine, and the ring amino group (-NH-) of the 3,7-di(disubstituted amino)-

- 34 -

10H-phenothiazine is converted to a protected ring amino group ($-R^{Prot}$) to give the corresponding protected 3,7-di(disubstituted amino)-10H-phenothiazine, for example:



5 In one embodiment, Y represents Cl^- .

In one embodiment, the combined reduction (RED) step and ring amino protection (AP) step is achieved using phenylhydrazine and acetic anhydride, for example, phenylhydrazine, ethanol, acetic anhydride, and pyridine.


10

In one embodiment, the step of (iii) ring amino deprotection (DP) and the step of (iv) salt formation (SF) are performed simultaneously (i.e., as one step).

For example, in one embodiment, the combined ring amino deprotection (DP) step and salt formation (SF) step is:

15

(ii) ring amino deprotection (DP) and salt formation (SF), wherein the protecting group of a protected 3,7-di(disubstituted amino)-10H-phenothiazine is removed to give a 3,7-di(disubstituted amino)-10H-phenothiazine, and the corresponding salt is formed, for example:



20

In one embodiment, the combined ring amino deprotection (DP) step and salt formation (SF) step may be performed using acid, for example, hydrochloric acid, for example, concentrated aqueous hydrochloric acid.

25

In a similar approach, an appropriate thioninium chloride (e.g., methyl thioninium chloride, ethyl thioninium chloride) is first reduced and acetylated to give the corresponding 1-(3,7-bis-dimethylamino-phenothiazin-10-yl)-ethanone, for example, by reaction with hydrazine (NH_2NH_2), methyl hydrazine ($MeNHNH_2$), or sodium borohydride ($NaBH_4$); and acetic

- 35 -

anhydride ((H₃CCO)₂O); for example, in the presence of a suitable base, for example, pyridine (C₅H₅N) or Hunig's base (diisopropylethylamine, C₈H₁₉N), for example, in a suitable solvent, for example, ethanol or acetonitrile. The reduced and acetylated compound is then deprotected (by removing the acetyl group), for example, by reaction
5 with a suitable halic acid, for example, hydrochloric acid or hydrobromic acid, in a suitable solvent, for example, ethanol, and optionally with the addition of a suitable ether, for example, diethyl ether.

Compositions

10

Another aspect of the invention pertains to a composition comprising a DAPTZ compound, as described herein, and a pharmaceutically acceptable carrier or diluent.

Uses

15

Reversing and/or Inhibiting the Aggregation of a Protein

One aspect of the invention is the use of a DAPTZ compound, as described herein, to regulate (e.g., to reverse and/or inhibit) the aggregation of a protein, for example,
20 aggregation of a protein associated with a neurodegenerative disease and/or clinical dementia. The aggregation may be in vitro, or in vivo, and may be associated with a disease state as discussed below.

Thus, one aspect of the invention pertains to a method of regulating (e.g., reversing and/or inhibiting) the aggregation of a protein, for example, aggregation of a protein
25 associated with a neurodegenerative disease and/or clinical dementia, comprising contacting the protein with an effective amount of a DAPTZ compound, as described herein. The method may be performed in vitro, or in vivo.

Similarly, one aspect of the invention pertains to a method of regulating (e.g., reversing and/or inhibiting) the aggregation of a protein in the brain of a mammal, which
30 aggregation is associated with a disease state as described herein, the treatment comprising the step of administering to said mammal in need of said treatment, a prophylactically or therapeutically effective amount of a DAPTZ compound, as described
35 herein, that is an inhibitor of said aggregation.

- 36 -

Methods of Treatment

- Another aspect of the present invention pertains to a method of treatment comprising administering to a patient in need of treatment a prophylactically or therapeutically effective amount of a DAPTZ compound, as described herein, preferably in the form of a pharmaceutical composition.

Use in Methods of Therapy

- Another aspect of the present invention pertains to a DAPTZ compound, as described herein, for use in a method of treatment (e.g., of a disease condition) of the human or animal body by therapy.

Use in the Manufacture of Medicaments

- Another aspect of the present invention pertains to use of a DAPTZ compound, as described herein, in the manufacture of a medicament for use in treatment (e.g., of a disease condition).
- In one embodiment, the medicament comprises the DAPTZ compound.

Disease Conditions Treated - Diseases of Protein Aggregation

- The DAPTZ compounds of the present invention are useful in the treatment or prophylaxis of diseases of protein aggregation.

- Thus, in one embodiment, the disease condition is a disease of protein aggregation, and, for example, the treatment is with an amount of a DAPTZ compound, as described herein, sufficient to inhibit the aggregation of the protein associated with said disease condition.

- In general, the protein aggregation is that which arises from an induced conformational polymerisation interaction, i.e., one in which a conformational change of the protein, or in a fragment thereof, gives rise to templated binding and aggregation of further (precursor) protein molecules in a self-propagating manner. Once nucleation is initiated, an aggregation cascade may ensue which involves the induced conformational polymerisation of further protein molecules, leading to the formation of toxic product fragments in aggregates which are substantially resistant to further proteolysis. The protein aggregates thus formed are thought to be a proximal cause of disease states manifested as neurodegeneration, clinical dementia, and other pathological symptoms.

The following Table provides a listing of various disease-associated aggregating proteins and the corresponding diseases of protein aggregation.

Diseases of protein aggregation				
Protein	Disease	Aggregating domain and/or mutations	Fibril subunit size (kDa)	Reference
<i>Neuro-degenerative disorders</i>				
Prion protein	Prion diseases	<i>Inherited and sporadic forms</i>	27	Prusiner (1998)
	(CJD, nvCJD, Fatal familial insomnia, Gerstmann-Straussler-Scheinker syndrome, Kuru)	PrP-27-30; many mutations.		
		Fibrillogenic domains: 113-120, 178-191, 202-218.		Gasset et al. (1992)
Tau protein	Alzheimer's disease, Down's syndrome, FTDP-17, CBD, post-encephalitic parkinsonism, Pick's disease, parkinsonism with dementia complex of Guam	<i>Inherited and sporadic forms</i>	10-12	Wischnik et al. (1988)
		Truncated tau (tubulin-binding domain) 297-391.		
		Mutations in tau in FTDP-17.		Hutton et al. (1998)
		Many mutations in presenilin proteins.		Czech et al. (2000)
Amyloid β -protein	Alzheimer's disease, Down's syndrome	<i>Inherited and sporadic forms</i>	4	Glenner & Wong, (1984)
		Amyloid β -protein; 1-42(3).		
		11 mutations in APP in rare families.		Goate et al. (1991)

Diseases of protein aggregation				
Protein	Disease	Aggregating domain and/or mutations	Fibril subunit size (kDa)	Reference
Huntingtin	Huntington's disease	N-termini of protein with expanded glutamine repeats.	40	DiFiglia et al. (1997)
Ataxins (1, 2, 3, 7)	Spinocerebellar ataxias (SCA1, 2, 3, 7)	Proteins with expanded glutamine repeats.		Paulson et al. (1999)
Atrophin	Dentatorubropallidoluysian atrophy (DRPLA)	Proteins with expanded glutamine repeats.		Paulson et al. (1999)
Androgen receptor	Spinal and bulbar muscular atrophy	Proteins with expanded glutamine repeats.		Paulson et al. (1999)
Neuroserpin	Familial encephalopathy with neuronal inclusion bodies (FENIB)	Neuroserpin; S49P, S52R.	57	Davis et al. (1999)
α -Synuclein	Parkinson's disease, dementia with Lewy bodies, multiple system atrophy	<i>Inherited and sporadic forms</i>	19	Spillantini et al. (1998)
		A53T, A30P in rare autosomal-dominant PD families.		Polymeropoulos et al. (1997)
Cystatin C	Hereditary cerebral angiopathy (Icelandic)	Cystatin C less 10 residues; L68Q.	12-13	Abrahamson et al. (1992)
Superoxide dismutase 1	Amyotrophic lateral sclerosis	SOD1 mutations.		Shibata et al. (1996)
Non-neuro-degenerative disorders				
Haemoglobin	Sickle cell anaemia	Haemoglobin beta chain (S).		Carrell & Gooptu (1998)

Diseases of protein aggregation				
Protein	Disease	Aggregating domain and/or mutations	Fibril subunit size (kDa)	Reference
	Inclusion body haemolysis	Many mutations.		
Serpins	α 1-Antitrypsin deficiency (emphysema, cirrhosis)	Mutations		Lomas et al. (1992)
	Antithrombin deficiency (thromboembolic disease)	Mutations		Carrell & Gooptu (1998)
	C1-inhibitor deficiency (angioedema)	Mutations		Carrell & Gooptu (1998)
Immunoglobulin light chain	Plasma cell dyscrasias (primary systemic AL amyloidosis)	light chain or fragments.	0.5-25	Westermarck et al. (1985)
Serum amyloid A	Reactive, secondary systemic AA amyloidosis	76-residue fragment (critical residues 2-12).	4.5-7.5	Westermarck et al. (1985)
	Chronic inflammatory disease			
Transthyretin	Familial amyloid polyneuropathy (systemic; FAP I)	Tetramer dissociated to conformational monomer variant.	10-14	Gustavsson et al. (1991)
		Many mutations (some not associated with amyloid; several different types of disease).		
	Senile cardiac amyloidosis	Normal transthyretin	10-14	Gustavsson et al. (1991)
Gelsolin	Familial amyloidosis - Finnish type (FAP IV)	D187Q leads to truncated 173-225/243 (critical residues 182-192).	9.5	Maury & Baumann (1990)
β 2-Microglobulin	Haemodialysis amyloidosis	β 2-Microglobulin	12-25	Gorevic et al. (1985)
	Prostatic amyloid			

Diseases of protein aggregation				
Protein	Disease	Aggregating domain and/or mutations	Fibril subunit size (kDa)	Reference
Apolipoprotein AI	Familial amyloid polyneuropathy (systemic; FAP III)	N-terminal 83-93 residues; G26R, W50R, L60R	9	Booth et al. (1997)
Lysozyme	Familial visceral amyloidosis	Lysozyme or fragments (with or without I56T, D67H)	14	Pepys et al. (1993)
Amylin (Islet amyloid polypeptide)	Type II diabetes (NIDDM)	Fragments (critical core of 20-29); no mutations	3.9	Westermarck (1990)
Fibrinogen α -chain	Hereditary renal amyloidosis	Fibrinogen fragments	7-10	Uemichi et al. (1992)
Procalcitonin	Medullary carcinoma of thyroid	Calcitonin fragments	3.4	Sletten et al. (1976)
Atrial natriuretic factor	Cardiac amyloidosis	ANF, no mutants	3.5	Johansson et al. (1987)
Insulin	Injection localised amyloidosis	Insulin		Dische et al. (1988)
Other proteins forming amyloid	(<i>in vitro</i>)	Other proteins		Chiti et al. (1999)

References for the above Table:

- 5 Abrahamson, M., Jonsdottir, S., Olafsson, I. & Grubb, A. (1992) Hereditary cystatin C amyloid angiopathy identification of the disease-causing mutation and specific diagnosis by polymerase chain reaction based analysis. *Human Genetics* 89, 377-380.
- 10 Booth, D.R., Sunde, M., Bellotti, V., Robinson, C.V., Hutchinson, W.L., Fraser, P.E., Hawkins, P.N., Dobson, C.M., Radford, S.E., Blake, C.C.F. & Pepys, M.B. (1997) Instability, unfolding and aggregation of human lysozyme variants underlying amyloid fibrillogenesis. *Nature* 385, 787-793.
- Carrell, R.W. & Gooptu, B. (1998) Conformational changes and disease - serpins, prions and Alzheimer's. *Current Opinion in Structural Biology* 8, 799-809.

- Chiti, F., Webster, P., Taddei, N., Clark, A., Stefani, M., Ramponi, G. & Dobson, C. (1999) Designing conditions for in vitro formation of amyloid protofilaments and fibrils. *Proceedings of the National Academy of Sciences, USA* 96, 3590-3594.
- 5 Czech, C., Tremp, G. & Pradier, L. (2000) Presenilins and Alzheimer's disease: biological functions and pathogenic mechanisms. *Progress in Neurobiology* 60, 363-384.
- Davis, R.L., Shrimpton, A.E., Holohan, P.D., Bradshaw, C., Feiglin, D., Collins, G.H., Sonderegger, P., Kinter, J., Becker, L.M., Lacbawan, F., Krasnewich, D., Muenke, M., Lawrence, D.A., Yerby, M.S., Shaw, C.-M., Gooptu, B., Elliott, P.R., Finch, J.T., Carrell, R.W. & Lomas, D.A. (1999) Familial dementia caused by
10 polymerization of mutant neuroserpin. *Nature* 401, 376-379.
- DiFiglia, M., Sapp, E., Chase, K.O., Davies, S.W., Bates, G.P., Vonsattel, J.P. & Aronin, N. (1997) Aggregation of huntingtin in neuronal intranuclear inclusions and dystrophic neurites in brain. *Science* 277, 1990-1993.
- Dische, F.E., Wernstedt, C., Westermarck, G.T., Westermarck, P., Pepys, M.B., Rennie, J.A., Gilbey, S.G. & Watkins, P.J. (1988) Insulin as an amyloid-fibril protein at sites
15 of repeated insulin injections in a diabetic patient. *Diabetologia* 31, 158-161.
- Gasset, M., Bladwin, M.A., Lloyd, D.H., abriel, J.-M., Holtzman, D.M., Cohen, F.E., Fletterick, R. & Prusiner, S.B. (1992) Predicted α -helical region of the prion protein when synthesized as peptides form amyloid. *Proceedings of the National
20 Academy of Sciences, USA* 89, 10940-10944.
- Glenner, G.G. & Wong, C.W. (1984) Alzheimer's disease: initial report of the purification and characterisation of a novel cerebrovascular amyloid protein. *Biochemical and Biophysical Research Communications* 120, 885-890.
- Goate, A., Chartier-Harlin, M.-C., Mullan, M., Brown, J., Crawford, F., Fidani, L., Giuffra, L., Haynes, A., Irving, N., James, L., Mant, R., Newton, P., Rooke, K., Roques, P., Talbot, C., Pericak-Vance, M., Roses, A., Williamson, R., Rossor, M., Owen, M. & Hardy, J. (1991) Segregation of a missense mutation in the amyloid precursor
25 protein gene with familial Alzheimer's disease. *Nature* 349, 704-706.
- Gorevic, P.D., Casey, T.T., Stone, W.J., DiRaimondo, C.R., Prelli, F.C. & Frangione, B. (1985) b-2 Microglobulin is an amyloidogenic protein in man. *Journal of Clinical
30 Investigation* 76, 2425-2429.
- Gustavsson, A., Engström, U. & Westermarck, P. (1991) Normal transthyretin and synthetic transthyretin fragments form amyloid-like fibrils in vitro. *Biochemical and Biophysical Research Communications* 175, 1159-1164.
- 35 Hutton, M., Lendon, C., Rizzu, P., Baker, M., Froelich, S., Houlden, H., Pickering-Brown, S., Chakraborty, S., Isaacs, A., Grover, A., Hackett, J., Adamson, J., Lincoln, S., Dickson, D., Davies, P., Petersen, R.C., Stevens, M., de Graaf, E., Wauters, E., van Baren, J., Hillebrand, M., Joosse, M., Kwon, J.M., Nowotny, P., Che, L.K., Norton, J., Morris, J.C., Reed, L.A., Trojanowski, J.Q., Basun, H., Lannfelt, L., Neystat, M., Fahn, S., Dark, F., Tannenberg, T., Dodd, P.R., Hayward, N., Kwok,
40 J.B.J., Schofield, P.R., Andreadis, A., Snowden, J., Craufurd, D., Neary, D.,

- Owen, F., Oostra, B.A., Hardy, J., Goate, A., van Swieten, J., Mann, D., Lynch, T. & Heutink, P. (1998) Association of missense and 5'-splice-site mutations in *tau* with the inherited dementia FTDP-17. *Nature* 393, 702-705.
- 5 Johansson, B., Wernstedt, C. & Westermark, P. (1987) Atrial natriuretic peptide deposited as atrial amyloid fibrils. *Biochemical and Biophysical Research Communications* 148, 1087-1092.
- Lomas, D.A., Evans, D.L., Finch, J.T. & Carrell, R.W. (1992) The mechanism of Z α 1-antitrypsin accumulation in the liver. *Nature* 357, 605-607.
- 10 Maury, C.P. & Baumann, M. (1990) Isolation and characterization of cardiac amyloid in familial amyloid polyneuropathy type IV (Finnish): relation of the amyloid protein to variant gelsolin. *Biochimica et Biophysica Acta* 1096, 84-86.
- Paulson, H.L. (1999) Human genetics '99: trinucleotide repeats. *American Journal of Human Genetics* 64, 339-345.
- 15 Pepys, M.B., Hawkins, P.N., Booth, D.R., Vigushin, D.M., Tennent, G.A., Soutar, A.K., Totty, N., Nguyen, O., Blake, C.C.F., Terry, C.J., Feast, T.G., Zalin, A.M. & Hsuan, J.J. (1993) Human lysozyme gene mutations cause hereditary systemic amyloidosis. *Nature* 362, 553-557.
- Polymeropoulos, M.H., Lavedan, C., Leroy, E., Ide, S.E., Dehejia, A., Dutra, A., Pike, B., Root, H., Rubenstein, J., Boyer, R., Stenroos, E.S., Chandrasekharappa, S., 20 Athanassiadou, A., Papaetropoulos, T., Johnson, W.G., Lazzarini, A.M., Duvoisin, R.C., Di Iorio, G., Golbe, L.I. & Nussbaum, R.L. (1997) Mutation in the α -synuclein gene identified in families with Parkinson's disease. *Science* 276, 2045-2047.
- Prusiner, S.B., Scott, M.R., DeArmond, S.J. & Cohen, F.E. (1998) Prion protein biology. *Cell* 93, 337-348.
- 25 Shibata, N., Hirano, A., Kobayashi, M., Siddique, T., Deng, H.X., Hung, W.Y., Kato, T. & Asayama, K. (1996) Intense superoxide dismutase-1 immunoreactivity in intracytoplasmic hyaline inclusions of familial amyotrophic lateral sclerosis with posterior column involvement. *Journal of Neuropathology and Experimental Neurology* 55, 481-490.
- 30 Sletten, K., Westermark, P. & Natvig, J.B. (1976) Characterization of amyloid fibril proteins from medullary carcinoma of the thyroid. *Journal of Experimental Medicine* 143, 993-998.
- Spillantini, M.G., Crowther, R.A., Jakes, R., Hasegawa, M. & Goedert, M. (1998) α -Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and 35 dementia with Lewy bodies. *Proceedings of the National Academy of Sciences, USA* 95, 6469-6473.
- Uemichi, T., Liuepnicks, J.J. & Benson, M.D. (1994) Hereditary renal amyloidosis with a novel variant fibrinogen. *Journal of Clinical Investigation* 93, 731-736.
- 40 Westermark, P., Engstrom, U., Johnson, K.H., Westermark, G.T. & Betsholtz, C. (1990) Islet amyloid polypeptide: pinpointing amino acid residues linked to amyloid fibril

formation. *Proceedings of the National Academy of Sciences, USA* 87, 5036-5040.

Westermarck, P., Johnson, K.H., O'Brien, T.D. & Betsholtz, C. (1992) Islet amyloid polypeptide - a novel controversy in diabetes research. *Diabetologia* 35, 297-303.

5 Westermarck, P., Johnson, K.H. & Pitkanen, P. (1985) Systemic amyloidosis: A review with emphasis on pathogenesis. *Applied Physiology* 3, 55-68.

Wischik, C.M., Novak, M., Thøgersen, H.C., Edwards, P.C., Runswick, M.J., Jakes, R., Walker, J.E., Milstein, C., M., R. & Klug, A. (1988) Isolation of a fragment of tau derived from the core of the paired helical filament of Alzheimer's disease.

10 *Proceedings of the National Academy of Sciences, USA* 85, 4506-4510.

As described in WO 02/055720 and US patent application number 60/786,700 filed on 29 March 2006 (title: Inhibitors of Protein Aggregation), diaminophenothiazines have utility in the inhibition of such protein aggregating diseases.

15

Thus it will be appreciated that, except where context requires otherwise, description of embodiments with respect to tau protein or tau-like proteins (e.g., MAP2), should be taken as applying equally to the other proteins discussed herein (e.g., β -amyloid, synuclein, prion, etc.) or other proteins which may initiate or undergo a similar pathological aggregation by virtue of conformational change in a domain critical for propagation of the aggregation, or which imparts proteolytic stability to the aggregate this formed (see, e.g., the article by Wischik et al. in "Neurobiology of Alzheimer's Disease", 2nd Edition, 2000, Eds. Dawbarn, D. and Allen, S.J., The Molecular and Cellular Neurobiology Series, Bios Scientific Publishers, Oxford). All such proteins may be referred to herein as "aggregating disease proteins."

20
25

Likewise, where mention is made herein of "tau-tau aggregation", or the like, this may also be taken to be applicable to other "aggregating-protein aggregation", such as β -amyloid aggregation, prion aggregation, synuclein aggregation, etc. The same applies for "tau proteolytic degradation" etc.

30

Preferred Aggregating Disease Proteins

Preferred embodiments of the invention are based on tau protein. The term "tau protein," as used herein, refers generally to any protein of the tau protein family. Tau proteins are characterised as being one among a larger number of protein families which co-purify with microtubules during repeated cycles of assembly and disassembly (see, e.g., Shelanski et al., 1973, Proc. Natl. Acad. Sci. USA, Vol. 70, pp. 765-768), and are known as microtubule-associated-proteins (MAPs). Members of the tau family share the common features of having a characteristic N-terminal segment, sequences of approximately 50 amino acids inserted in the N-terminal segment, which are

35
40

- 44 -

developmentally regulated in the brain, a characteristic tandem repeat region consisting of 3 or 4 tandem repeats of 31-32 amino acids, and a C-terminal tail.

5 MAP2 is the predominant microtubule-associated protein in the somatodendritic compartment (see, e.g., Matus, A., in "*Microtubules*" [Hyams and Lloyd, Eds.] pp. 155-166, John Wiley and Sons, New York, USA). MAP2 isoforms are almost identical to tau protein in the tandem repeat region, but differ substantially both in the sequence and extent of the N-terminal domain (see, e.g., Kindler and Garner, 1994, *Mol. Brain Res.*, Vol. 26, pp. 218-224). Nevertheless, aggregation in the tandem-repeat region is not
10 selective for the tau repeat domain. Thus it will be appreciated that any discussion herein in relation to tau protein or tau-tau aggregation should be taken as relating also to tau-MAP2 aggregation, MAP2-MAP2 aggregation, and so on.

15 In one embodiment, the protein is tau protein.

In one embodiment, the protein is a synuclein, e.g., α - or β -synuclein.

Where the protein is tau protein, in one embodiment of the present invention, there is provided a method of inhibiting production of protein aggregates (e.g. in the form of
20 paired helical filaments (PHFs), optionally in neurofibrillary tangles (NFTs) in the brain of a mammal, the treatment being as described above.

Preferred Diseases of Protein Aggregation

25 Notably it is not only Alzheimer's disease (AD) in which tau protein (and aberrant function or processing thereof) may play a role. The pathogenesis of neurodegenerative disorders such as Pick's disease and Progressive Supranuclear Palsy (PSP) appears to correlate with an accumulation of pathological truncated tau aggregates in the dentate gyrus and stellate pyramidal cells of the neocortex, respectively. Other dementias include fronto-
30 temporal dementia (FTD); parkinsonism linked to chromosome 17 (FTDP-17); disinhibition-dementia-parkinsonism-amyotrophy complex (DDPAC); pallido-ponto-nigral degeneration (PPND); Guam-ALS syndrome; pallido-nigro-luysian degeneration (PNLD); cortico-basal degeneration (CBD) and others (see, e.g., the article by Wischik et al. in
35 "*Neurobiology of Alzheimer's Disease*", 2nd Edition, 2000, Eds. Dawbarn, D. and Allen, S.J., *The Molecular and Cellular Neurobiology Series*, Bios Scientific Publishers, Oxford; especially Table 5.1). All of these diseases, which are characterized primarily or partially by abnormal tau aggregation, are referred to herein as "tauopathies".

40 Thus, in one embodiment, the disease condition is a tauopathy.

In one embodiment, the disease condition is a neurodegenerative tauopathy.

- 45 -

In one embodiment, the disease condition is Alzheimer's disease.

5 In one embodiment, treatment (e.g., treatment of a neurodegenerative tauopathy, e.g., Alzheimer's disease) may optionally be in combination with one or more other agents, for example, one or more cholinesterase inhibitors (such as Donepezil (also known as Aricept™), Rivastigmine (also known as Exelon™), Galantamine (also known as Reminyl™), NMDA receptor antagonists (such as Memantine (also known as Ebixa™, Namenda™), muscarinic receptor agonists, and/or inhibitors of amyloid precursor protein
10 processing that leads to enhanced generation of beta-amyloid.

Disease Conditions Treated - Other Disease Conditions

15 In one embodiment, the disease condition is skin cancer.

In one embodiment, the disease condition is melanoma.

20 In one embodiment, the disease condition is a viral, bacterial or protozoal disease condition.

In one embodiment, the (protozoal) disease condition is malaria.

25 In this embodiment, treatment may be in combination with one or more antimicrobial agents, for example, chloroquine and/or atovaquone.

In one embodiment, the (viral) disease condition is caused by Hepatitis C, HIV, or West Nile Virus (WNV).

Other Uses

30 Another aspect of the present invention pertains to use of a DAPTZ compound, as described herein, in a method of inactivating a pathogen in a sample (for example a blood or plasma sample), comprising the steps of introducing the DAPTZ compound into the sample, and exposing the sample to light.

35 For example, in one embodiment, the method comprises the steps of introducing the DAPTZ compound into the sample, and then exposing the sample to light.

- 46 -

Use as Ligands

The DAPTZ compounds that are capable of inhibiting the aggregation of tau protein will also be capable of acting as ligands or labels of tau protein (or aggregated tau protein).

- 5 Thus, in one embodiment, the DAPTZ compound is a ligand of tau protein (or aggregated tau protein).

- 10 Such DAPTZ compounds (ligands) may incorporate, be conjugated to, be chelated with, or otherwise be associated with, other chemical groups, such as stable and unstable detectable isotopes, radioisotopes, positron-emitting atoms, magnetic resonance labels, dyes, fluorescent markers, antigenic groups, therapeutic moieties, or any other moiety that may aid in a prognostic, diagnostic, or therapeutic application.

- 15 For example, in one embodiment, the DAPTZ compound is as defined herein, but with the additional limitation that the compound incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels, for example, isotopes, radioisotopes, positron-emitting atoms, magnetic resonance labels, dyes, fluorescent markers, antigenic groups, or therapeutic moieties.

- 20 In one embodiment, the DAPTZ compound is a ligand as well as a label, e.g., a label for tau protein (or aggregated tau protein), and incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels.

- 25 For example, in one embodiment, the DAPTZ compound is as defined above, but with the additional limitation that the compound incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels.

- 30 Labelled DAPTZ compounds (e.g., when ligated to tau protein or aggregated tau protein) may be visualised or detected by any suitable means, and the skilled person will appreciate that any suitable detection means as is known in the art may be used.

- 35 For example, the DAPTZ compound (ligand-label) may be suitably detected by incorporating a positron-emitting atom (e.g., ^{11}C) (e.g., as a carbon atom of one or more alkyl group substituents, e.g., methyl group substituents) and detecting the compound using positron emission tomography (PET) as is known in the art.

Such ^{11}C labelled DAPTZ compounds may be prepared by adapting the methods described herein in known ways, for example, in analogy to the methods described in WO 02/075318 (see Figures 11a, 11b, 12) and WO 2005/030676.

- 47 -

Thus, another aspect of the present invention pertains to a method of labelling tau protein (or aggregated tau protein) comprising the step of: (i) contacting the tau protein (or aggregated tau protein) with a DAPTZ compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels.

Another aspect of the present invention pertains to a method of detecting tau protein (or aggregated tau protein) comprising the steps of: (i) contacting the tau protein (or aggregated tau protein) with a DAPTZ compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels, and (ii) detecting the presence and/or amount of said compound bound to tau protein (or aggregated tau protein).

Another aspect of the present invention pertains to a method of diagnosis or prognosis of a tau proteinopathy in a subject believed to suffer from the disease, comprising the steps of: (i) introducing into the subject a DAPTZ compound capable of labelling tau protein or aggregated tau protein, particularly tau protein (e.g., a DAPTZ compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels); (ii) determining the presence and/or amount of said compound bound to tau protein or aggregated tau protein in the brain of the subject; and (iii) correlating the result of the determination made in (ii) with the disease state of the subject.

Another aspect of the present invention pertains to a DAPTZ compound capable of labelling tau protein or aggregated tau protein (e.g., a DAPTZ compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels), for use in a method of diagnosis or prognosis of a tau proteinopathy.

Another aspect of the present invention pertains to use of a DAPTZ compound capable of labelling tau protein or aggregated tau protein, particularly tau protein (e.g., a DAPTZ compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels), in a method of manufacture of a diagnostic or prognostic reagent for use in the diagnosis or prognosis of a tau proteinopathy.

Those skilled in the art will appreciate that instead of administering DAPTZ ligands/labels directly, they could be administered in a precursor form, for conversion to the active form (e.g., ligating form, labelling form) by an activating agent present in, or administered to, the same subject.

- 48 -

The ligands disclosed herein may be used as part of a method of diagnosis or prognosis. It may be used to select a patient for treatment, or to assess the effectiveness of a treatment or a therapeutic (e.g., an inhibitor of tau protein aggregation) administered to the subject.

5

Treatment

The term "treatment," as used herein in the context of treating a condition, pertains generally to treatment and therapy, whether of a human or an animal (e.g., in veterinary applications), in which some desired therapeutic effect is achieved, for example, the inhibition of the progress of the condition, and includes a reduction in the rate of progress, a halt in the rate of progress, regression of the condition, amelioration of the condition, and cure of the condition. Treatment as a prophylactic measure (i.e., prophylaxis, prevention) is also included.

15

The term "therapeutically-effective amount," as used herein, pertains to that amount of a DAPTZ compound, or a material, composition or dosage from comprising a DAPTZ compound, which is effective for producing some desired therapeutic effect, commensurate with a reasonable benefit/risk ratio, when administered in accordance with a desired treatment regimen.

20

Similarly, the term "prophylactically effective amount," as used herein, pertains to that amount of a DAPTZ compound, or a material, composition or dosage from comprising a DAPTZ compound, which is effective for producing some desired prophylactic effect, commensurate with a reasonable benefit/risk ratio, when administered in accordance with a desired treatment regimen.

25

The term "treatment" includes combination treatments and therapies, in which two or more treatments or therapies are combined, for example, sequentially or simultaneously. Examples of treatments and therapies include, but are not limited to, chemotherapy (the administration of active agents, including, e.g., drugs, antibodies (e.g., as in immunotherapy), prodrugs (e.g., as in photodynamic therapy, GDEPT, ADEPT, etc.); surgery; radiation therapy; and gene therapy.

30

For example, it may be beneficial to combine treatment with a DAPTZ compound as described herein with one or more other (e.g., 1, 2, 3, 4) agents or therapies.

35

The particular combination would be at the discretion of the physician who would select dosages using his/her common general knowledge and dosing regimens known to a skilled practitioner.

40

- 49 -

The agents (i.e., a DAPTZ compound as described here, plus one or more other agents) may be administered simultaneously or sequentially, and may be administered in individually varying dose schedules and via different routes. For example, when administered sequentially, the agents can be administered at closely spaced intervals (e.g., over a period of 5-10 minutes) or at longer intervals (e.g., 1, 2, 3, 4 or more hours apart, or even longer periods apart where required), the precise dosage regimen being commensurate with the properties of the therapeutic agent(s).

The agents (i.e., a DAPTZ compound as described here, plus one or more other agents) may be formulated together in a single dosage form, or alternatively, the individual agents may be formulated separately and presented together in the form of a kit, optionally with instructions for their use.

Routes of Administration

15

The DAPTZ compound, or pharmaceutical composition comprising it, may be administered to a subject/patient by any convenient route of administration, whether systemically/peripherally or topically (i.e., at the site of desired action).

Routes of administration include, but are not limited to, oral (e.g., by ingestion); buccal; sublingual; transdermal (including, e.g., by a patch, plaster, etc.); transmucosal (including, e.g., by a patch, plaster, etc.); intranasal (e.g., by nasal spray); ocular (e.g., by eyedrops); pulmonary (e.g., by inhalation or insufflation therapy using, e.g., via an aerosol, e.g., through the mouth or nose); rectal (e.g., by suppository or enema); vaginal (e.g., by pessary); parenteral, for example, by injection, including subcutaneous, intradermal, intramuscular, intravenous, intraarterial, intracardiac, intrathecal, intraspinal, intracapsular, subcapsular, intraorbital, intraperitoneal, intratracheal, subcuticular, intraarticular, subarachnoid, and intrasternal (including, e.g., intracatheter injection into the brain); by implant of a depot or reservoir, for example, subcutaneously or intramuscularly.

30

The Subject/Patient

The subject/patient may be an animal, a mammal, a placental mammal, a rodent (e.g., a guinea pig, a hamster, a rat, a mouse), murine (e.g., a mouse), a lagomorph (e.g., a rabbit), avian (e.g., a bird), canine (e.g., a dog), feline (e.g., a cat), equine (e.g., a horse), porcine (e.g., a pig), ovine (e.g., a sheep), bovine (e.g., a cow), a primate, simian (e.g., a monkey or ape), a monkey (e.g., marmoset, baboon), an ape (e.g., gorilla, chimpanzee, orangutang, gibbon), or a human.

40

- 50 -

Furthermore, the subject/patient may be any of its forms of development, for example, a foetus.

In one preferred embodiment, the subject/patient is a human.

5

In one embodiment, the subject/patient is not a human.

Formulations

10 While it is possible for the DAPTZ compound to be used (e.g., administered) alone, it is often preferable to present it as a composition or formulation.

In one embodiment, the composition is a pharmaceutical composition (e.g., formulation, preparation, medicament) comprising a DAPTZ compound, as described herein, and a
15 pharmaceutically acceptable carrier, diluent, or excipient.

In one embodiment, the composition is a pharmaceutical composition comprising at least one DAPTZ compound, as described herein, together with one or more other
20 pharmaceutically acceptable ingredients well known to those skilled in the art, including, but not limited to, pharmaceutically acceptable carriers, diluents, excipients, adjuvants, fillers, buffers, preservatives, anti-oxidants, lubricants, stabilisers, solubilisers, surfactants (e.g., wetting agents), masking agents, colouring agents, flavouring agents, and sweetening agents.

25 In one embodiment, the composition further comprises other active agents, for example, other therapeutic or prophylactic agents.

Suitable carriers, diluents, excipients, etc. can be found in standard pharmaceutical texts. See, for example, Handbook of Pharmaceutical Additives, 2nd Edition (eds. M. Ash and I.
30 Ash), 2001 (Synapse Information Resources, Inc., Endicott, New York, USA), Remington's Pharmaceutical Sciences, 20th edition, pub. Lippincott, Williams & Wilkins, 2000; and Handbook of Pharmaceutical Excipients, 2nd edition, 1994.

Another aspect of the present invention pertains to methods of making a pharmaceutical
35 composition comprising admixing at least one [¹¹C]-radiolabelled DAPTZ compound, as defined herein, together with one or more other pharmaceutically acceptable ingredients well known to those skilled in the art, e.g., carriers, diluents, excipients, etc. If formulated as discrete units (e.g., tablets, etc.), each unit contains a predetermined amount (dosage) of the DAPTZ compound.

40

- 51 -

The term "pharmaceutically acceptable," as used herein, pertains to compounds, ingredients, materials, compositions, dosage forms, etc., which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of the subject in question (e.g., human) without excessive toxicity, irritation, allergic response, or other
5 problem or complication, commensurate with a reasonable benefit/risk ratio. Each carrier, diluent, excipient, etc. must also be "acceptable" in the sense of being compatible with the other ingredients of the formulation.

The formulations may be prepared by any methods well known in the art of pharmacy.
10 Such methods include the step of bringing into association the DAPTZ compound with a carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the DAPTZ compound with carriers (e.g., liquid carriers, finely divided solid carrier, etc.), and then shaping the product, if necessary.

15 The formulation may be prepared to provide for rapid or slow release; immediate, delayed, timed, or sustained release; or a combination thereof.

Formulations suitable for parenteral administration (e.g., by injection), include aqueous or
20 non-aqueous, isotonic, pyrogen-free, sterile liquids (e.g., solutions, suspensions), in which the DAPTZ compound is dissolved, suspended, or otherwise provided (e.g., in a liposome or other microparticulate). Such liquids may additionally contain other pharmaceutically acceptable ingredients, such as anti-oxidants, buffers, preservatives, stabilisers, bacteriostats, suspending agents, thickening agents, and solutes which render
25 the formulation isotonic with the blood (or other relevant bodily fluid) of the intended recipient. Examples of excipients include, for example, water, alcohols, polyols, glycerol, vegetable oils, and the like. Examples of suitable isotonic carriers for use in such formulations include Sodium Chloride Injection, Ringer's Solution, or Lactated Ringer's Injection. Typically, the concentration of the DAPTZ compound in the liquid is from about
30 1 ng/ml to about 10 µg/ml, for example from about 10 ng/ml to about 1 µg/ml. The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampoules and vials, and may be stored in a freeze-dried (lyophilised) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from
35 sterile powders, granules, and tablets.

Examples of Some Preferred Formulations

One aspect of the present invention pertains to a dosage unit (e.g., a pharmaceutical
40 tablet or capsule) comprising 20 to 300 mg of a DAPTZ compound as described herein

- 52 -

(e.g., obtained by, or obtainable by, a method as described herein; having a purity as described herein; etc.), and a pharmaceutically acceptable carrier, diluent, or excipient.

In one embodiment, the dosage unit is a tablet.

5 In one embodiment, the dosage unit is a capsule.

In one embodiment, the amount is 30 to 200 mg.

In one embodiment, the amount is about 30 mg.

In one embodiment, the amount is about 60 mg.

10 In one embodiment, the amount is about 100 mg.

In one embodiment, the amount is about 150 mg.

In one embodiment, the amount is about 200 mg.

15 In one embodiment, the pharmaceutically acceptable carrier, diluent, or excipient is or comprises one or both of a glyceride (e.g., Gelucire 44/14 ®; lauroyl macrogol-32 glycerides PhEur, USP) and colloidal silicon dioxide (e.g., 2% Aerosil 200 ®; Colliodal Silicon Dioxide PhEur, USP).

Dosage

20

It will be appreciated by one of skill in the art that appropriate dosages of the DAPTZ compound, and compositions comprising the DAPTZ compound, can vary from patient to patient. Determining the optimal dosage will generally involve the balancing of the level of therapeutic benefit against any risk or deleterious side effects. The selected dosage
25 level will depend on a variety of factors including, but not limited to, the activity of the particular compound, the route of administration, the time of administration, the rate of excretion of the compound, the duration of the treatment, other drugs, compounds, and/or materials used in combination, the severity of the condition, and the species, sex, age, weight, condition, general health, and prior medical history of the patient. The amount of
30 compound and route of administration will ultimately be at the discretion of the physician, veterinarian, or clinician, although generally the dosage will be selected to achieve local concentrations at the site of action which achieve the desired effect without causing substantial harmful or deleterious side-effects.

35 Administration can be effected in one dose, continuously or intermittently (e.g., in divided doses at appropriate intervals) throughout the course of treatment. Methods of determining the most effective means and dosage of administration are well known to those of skill in the art and will vary with the formulation used for therapy, the purpose of the therapy, the target cell(s) being treated, and the subject being treated. Single or
40 multiple administrations can be carried out with the dose level and pattern being selected by the treating physician, veterinarian, or clinician.

- 53 -

In general, a suitable dose of the DAPTZ compound is in the range of about 100 ng to about 25 mg (more typically about 1 μ g to about 10 mg) per kilogram body weight of the subject per day.

5

In one embodiment, the DAPTZ compound is administered to a human patient according to the following dosage regime: about 100 mg, 3 times daily.

10

In one embodiment, the DAPTZ compound is administered to a human patient according to the following dosage regime: about 150 mg, 2 times daily.

In one embodiment, the DAPTZ compound is administered to a human patient according to the following dosage regime: about 200 mg, 2 times daily.


EXAMPLES

The following examples are provided solely to illustrate the present invention and are not intended to limit the scope of the invention, as described herein.

5

Chemical SynthesisSynthesis 1

3-Nitro-10H-phenothiazine



10


Sodium nitrite (20.00 g, 210 mmol) was added to a mixture of 10H-phenothiazine (20.00 g, 50 mmol), chloroform (100 cm³), and acetic acid (20 cm³), and the mixture was stirred for 1 hour at room temperature. Acetic acid (20 cm³) was then added and the mixture was stirred for a further 18 hours. The suspension was filtered and washed with acetic acid, ethanol, water, and finally ethanol to give a purple/brown solid. The residue was dissolved in hot DMF and allowed to cool before filtering the di-nitro compound as a purple solid. Concentration of the DMF solution and washing the precipitate with water and methanol gave the title mono-nitro compound (15g, ~50%) as a brown solid; $\nu_{\max}$ (KBr)/cm⁻¹: 3328 (NH), 3278 (NH), 3229 (NH), 3119 (CH), 3049 (CH), 1557 (NO₂), 1531 (NO₂); δ_{H} (250 MHz; DMSO): 6.64 (5H, m, ArH), 7.68 (1H, d, *J* 2.5, ArH), 7.79-7.84 (1H, dd, *J* 2.75, 6.5, ArH); δ_{C} (62.9 MHz; DMSO): 113.3 (ArC), 115.3 (ArC), 116.9 (ArC), 121.8 (ArC), 123.6 (ArC), 123.7 (ArC), 124.6 (ArC), 126.4 (ArC), 128.1 (ArC), 138.8 (ArC), 141.0 (ArC), 147.8 (ArC).

20

25

Synthesis 2

3,11-Dinitro-10H-phenothiazine




30

The procedure for the synthesis of 3-nitro-10H-phenothiazine was followed using 3-nitro-10H-phenothiazine (10.00 g, 41 mmol), chloroform (40 cm³), acetic acid (2 x 10 cm³), and sodium nitrite (11.86 g, 173 mmol). The residue obtained was recrystallised from DMF to yield the title di-nitro compound (6.60 g 56%) as purple needles; $\nu_{\max}$ (KBr)/cm⁻¹: 3331 (NH), 3294 (NH), 3229 (NH), 3101 (CH), 3067 (CH), 1602 (NO₂), 1558 (NO₂); δ_{H} (250 MHz; DMSO): 6.73-6.76 (2H, d, *J* 9, ArH), 7.78 (2H, s, ArH), 7.89-7.85 (2H, d, *J* 9, ArH).

- 55 -

Synthesis 3


1-(3,7-Dinitro-phenothiazin-10-yl)-ethanone



- 5 A solution of 3,11-dinitro-10H-phenothiazine (3.00 g, 10.37 mmol), acetic anhydride (15.88 g, 155.50 mmol), and pyridine (30 cm³) was stirred at reflux for 18 hours. The warm solution was then carefully poured over ice water. A precipitate formed and was filtered, dissolved in dichloromethane, dried over magnesium sulphate, filtered, and concentrated to give a brown/orange solid, which was purified by column chromatography
- 10 (SiO₂, ethyl acetate : petroleum ether, 2:3, loaded as a dichloromethane solution) to give the title compound (2.46 g, 71%) as a light yellow solid which can be recrystallised from acetone to give light yellow needles; $\nu_{\max}$ (KBr)/cm⁻¹: 3091 (CH), 3063 (CH), 1680 (C=O), 1575 (NO₂), 1510 (NO₂); δ_{H} (250 MHz; CDCl₃): 2.28 (3H, s, CH₃), 7.65-7.69 (2H, d, *J* 9, *ArH*), 8.22-8.26 (2H; dd, *J* 2.75, 8.75, *ArH*), 8.33-8.32 (2H, d, *J* 2.5, *ArH*); δ_{C} (62.9 MHz; CDCl₃): 168.2 (C=O), 146.3 (ArC), 143.3 (ArC), 133.6 (ArC), 127.8 (ArC), 123.4 (ArC), 122.9 (ArC), 23.1 (CH₃); *m/z* (ES) 331.0 (80%, [M]⁺).
- 15

Synthesis 4

1-(3,7-Diamino-phenothiazin-10-yl)-ethanone




- 20 A mixture of 1-(3,7-dinitro-phenothiazin-10-yl)-ethanone (2 g, 6.04 mmol), tin (II) chloride dihydrate (14.17 g, 62.8 mmol), and ethanol (50 cm³) was heated to reflux and stirred at this temperature for 5 hours. The mixture was then cooled to room temperature and poured over ice water. The pH was adjusted to 7 with 5% sodium hydrogen carbonate
- 25 before the product was extracted with ethyl acetate (3 x 50 cm³). The extracts were washed with brine and dried over magnesium sulphate, filtered, and concentrated to give the title compound (1.64 g, 100%) as a purple blue solid; $\nu_{\max}$ (KBr)/cm⁻¹: 3445 (NH), 3424 (NH), 3368 (NH), 3322 (NH), 3203 (NH), 3054 (CH), 2995 (CH), 1706 (C=O), 1650 (NO₂), 1590 (NO₂); δ_{H} (250 MHz; CDCl₃): 2.01 (3H, s, CH₃), 5.09-5.43 (4H, brd s, NH), 6.47-6.51
- 30 (2H, dd, *J* 1.5, 8.25, *ArH*), 6.61 (2H, s, *ArH*), 7.11-7.15 (2H, d, *J* 8, *ArH*); δ_{C} (62.9 MHz; CDCl₃): 169.1 (C=O), 147.2 (ArC), 128.1 (ArC), 127.6 (ArC), 127.3 (ArC), 112.3 (ArC), 111.5 (ArC), 22.6 (CH₃); *m/z* (ES) 293.9 (95%, [M + H, Na]⁺), 272.0 (20%, [M + H]⁺), 227.9 (100%, [M + H, - Ac]⁺).

- 56 -

Synthesis 5


3,7-Diamino-phenothiazine bis(hydrogen chloride) (B4)



- 5 1-(3,7-Diamino-phenothiazin-10-yl)-ethanone (0.25 g, 0.921 mmol) was dissolved in aqueous hydrochloric acid (5 N, 10 cm³) and the solution was heated to reflux and stirred for 30 minutes. Concentration of the reaction mixture gave the title compound as a light blue solid. δ_H (250 MHz; D₂O): 6.60 (2H, brd d, ArH), 7.07 (4H, brd s, ArH).

Synthesis 6

1-(3,7-Bis-dimethylamino-phenothiazin-10-yl)-ethanone




- 10 1-(3,7-Diamino-phenothiazin-10-yl)-ethanone (0.25 g 0.92 mmol) was dissolved in DMSO (3 cm³). Toluene (10 cm³), iodomethane (1.96 g, 13.8 mmol), tetrabutylammonium bromide (50 mg), and finally aqueous sodium hydroxide solution (50%, 1.25 cm³) were added. The mixture was stirred at room temperature for 2 hours. Additional aqueous sodium hydroxide (50%, 1.25 cm³) and iodomethane (1.96 g, 13.8 mmol) were then added. The mixture was allowed to stir for a further 3 hours at room temperature before a third aliquot of aqueous sodium hydroxide (50%, 1.25 cm³) and iodomethane (1.96 g, 13.8 mmol) were added and the mixture stirred for a further 18 hours. The thick suspension was washed with water (3 x 75 cm³) and the toluene extract collected. The water was extracted with dichloromethane (3 x 50 cm³) and the extracts combined with the toluene, and dried over magnesium sulphate, filtered, and concentrated to give a deep purple solid. The residue was purified by column chromatography (SiO₂, ethyl acetate : petroleum ether, 2:3, loaded as a dichloromethane solution) to give the title compound product (0.12 g, 40%) as a light purple solid; ν_{max} (KBr)/cm⁻¹: 2910 (CH), 2876 (CH), 2856 (CH), 2799 (CH), 1659 (C=O), 1596 (NO₂), 1502 (NO₂); δ_H (250 MHz; CDCl₃): 2.16 (3H, s, CH₃), 2.93 (12H, s, NCH₃), 6.59-6.62 (2H, d, J 8.5, ArH), 6.69-6.71 (2H, d, J 2.75, ArH), 7.08-7.47 (2H, brd s, ArH); δ_C (62.9MHz; CDCl₃): 170.3 (C=O), 148.9 (ArC), 127.2 (ArC), 127.1 (ArC), 127.0 (ArC), 110.9 (ArC), 110.7 (ArC), 40.7 (NCH₃), 22.9 (CH₃).

- 57 -

Synthesis 7


N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride) (B3)



1-(3,7-Bis-dimethylamino-phenothiazin-10-yl)-ethanone (0.5 g, 1.84 mmol) was dissolved in aqueous hydrochloric acid (5 N, 15 cm³), and the solution was heated to reflux temperature and stirred for 30 minutes. Concentration of the reaction mixture gave the title compound as a green/blue solid; δ_H (250 MHz; D₂O): 3.18 (12H, s, NCH₃), 6.67 (2H, d, *J* 8.5, ArH), 7.16 (4H, brd s, ArH); δ_C (62.9 MHz; D₂O): 144.3 (ArC), 138.9 (ArC), 122.4 (ArC), 120.8 (ArC), 120.7 (ArC), 117.6 (ArC), 48.9 (NCH₃).

Synthesis 8


Methylthioninium iodide



To a round bottom flask was added methylthioninium chloride (MTC, Methylene Blue) (2 g, 6.25 mmol) and water (50 cm³) and the mixture stirred for 10 minutes or until the solid dissolved. Potassium iodide (1.56 g, 9.4 mmol) was then added to the mixture and a green black suspension formed. The reaction was heated to boiling and allowed to cool naturally giving the title compound (2.03 g, 79%) as bright green needles. Anal. Calcd for C₁₆H₁₈N₃SI: C, 46.72; H, 4.41; N, 10.22; S, 7.80; I, 30.85. Found: C, 46.30; H, 4.21; N, 10.14; S, 7.86; I, 29.34.

Synthesis 9

N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen iodide) (B6)



To a round bottom flask was added methylthioninium iodide (2 g, 4.86 mmol), ethanol (100 cm³) and ethyl iodide (75.8 g, 486 mmol) and the mixture was heated at reflux for 18 hours where the colour changed from green/blue to brown with a yellow precipitate. Once cooled to room temperature, the mixture was filtered and washed with diethylether (20 cm³) to give the title compound (1.99 g, 76%) as a light green solid. δ_H (250 MHz;


- 58 -

D₂O): 3.20 (12H, s, NCH₃), 6.76 (2H, d, *J* 8.5, ArH), 7.22 (2H, brd s, ArH); δ_c (62.9 MHz; D₂O): 145.0 (ArC), 139.3 (ArC), 122.6 (ArC), 121.1 (ArC), 120.9 (ArC), 117.9 (ArC), 48.9 (NCH₃).

5

Synthesis 10

1-(3,7-Bis-diethylamino-phenothiazin-10-yl)-ethanone




To a dry 25cm³ round bottom flask was added ethylthioninium zinc chloride (0.5 g, 1.13 mmol) and ethanol (10 cm³). Phenylhydrazine (0.134 g, 1.24 mmol) was then added dropwise under an atmosphere of nitrogen. The mixture was stirred 25°C for 1 hour and concentrated under high vacuum. Pyridine (50 cm³) and acetic anhydride was added and the mixture stirred for 18 hours at 60°C. The solution was opened to ice/water (250 cm³) and the organics were extracted into ethyl acetate (3 x 50 cm³). The extracts were washed with saturated copper sulphate solution and dried over magnesium sulphate, filtered, and concentrated to give the crude product as a brown oil, which was purified using flash column chromatography with an eluent of 40% ethylacetate : 60% petroleum spirit 40-60°C and silica 40-63 μ 60Å to give the title compound (0.18 g, 41%) as a green glassy solid. δ_H (250 MHz; CDCl₃): 7.0-7.5 (2H, brds, ArH), 6.64 (2H, s, ArH), 6.52 (2H, d, ArH), 3.35 (8H, q, 7, NCH₂), 2.18 (3H, s, CH₃), 1.16 (12H, t, 7, CH₃); δ_c (62.9 MHz; CDCl₃): 12.5 (CH₃), 22.9 (CH₃), 44.6 (NCH₂), 110.1 (ArC), 127.4 (ArC), 146.5 (ArC), 170.2 (C=O).

20

Synthesis 11

N,N,N',N'-Tetraethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride)



2 HCl

25

To a 25 cm³ round bottom flask was added 3,7-diethylamino-10-acetyl-phenothiazine (0.125 g, 0.33 mmol) and aqueous hydrochloric acid (5 M, 5 cm³). The mixture was heated at 100°C for 2 hours before cooling to room temperature and was concentrated to give the title compound (0.11 g, 81%) as a yellow green glassy solid. δ_H (250 MHz; CD₃OD): 7.07 (4H, brd, ArH), 6.65 (2H, brd, ArH), 3.35 (8H, brd, NCH₂), 0.97 (12H, brd, CH₃); δ_c (62.9 MHz; CD₃OD): 10.8 (CH₃), 55.1 (NCH₂), 116.6 (ArC), 120.4 (ArC), 121.5 (ArC), 123.6 (ArC), 132.6 (ArC), 144.5 (ArC).


30

- 60 -

precipitate. The precipitate was stirred for 45 minutes before it was filtered, washed with water (3 x 250 cm³), and air dried for 30 minutes. The crude material was crystallised from hot ethanol (2750 cm³) to yield the title compound (112.1 g, 64%) as a light grey solid. Mp 137°C; $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ 2910 (CH), 2876 (CH), 2856 (CH), 2799 (CH), 1659 (C=O), 1596 (NO₂), 1502 (NO₂); δ_{H} (250MHz; CDCl₃) 2.16 (3H, s, CH₃), 2.93 (12H, s, NCH₃), 6.59-6.62 (2H, d, *J* 8.5, ArH), 6.69-6.71 (2H, d, *J* 2.75, ArH), 7.08-7.47 (2H, brd s, ArH); δ_{C} (62.9MHz; CDCl₃) 170.3 (C=O), 148.9 (ArC), 127.2 (ArC), 127.1 (ArC), 127.0 (ArC), 110.9 (ArC), 110.7 (ArC), 40.7 (NCH₃), 22.9 (CH₃); *m/z* (ES) 284.2 (100%, [M - OAc]⁺), 328.1 (15%, [M + H]⁺), 350.1 (41%, [M + Na]⁺).

Synthesis 14


1-(3,7-Bis-dimethylamino-phenothiazin-10-yl)-ethanone



Synthesis using methylhydrazine/ pyridine in one pot. To a 250 cm³ round bottom flask under an atmosphere of nitrogen was added methylthionium chloride trihydrate (26.74 mmol, 10 g) and acetonitrile (50 cm³). Methylhydrazine (53.5 mmol, 2.46 g) was added in four equal portions over a 30 minutes time period. The temperature of the mixture was maintained at 35°C with a cold water bath and was stirred for 30 minutes. The yellow/green suspension had acetic anhydride (267 mmol, 27.3 g) and pyridine (80.2 mmol, 6.35 g) was added. The mixture was heated at 90°C for 2 hours. The cooled mixture was then poured carefully into ice water (200 cm³) in ten equal portions while stirring to give a precipitate. The precipitate was stirred for 30 minutes before it was filtered, washed with water (3 x 50 cm³) and air dried for 30 minutes. The crude material was crystallised from hot ethanol (120 cm³) to yield the title compound (5.97 g, 68%) as a light grey solid. Mp 137°C; $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ 2910 (CH), 2876 (CH), 2856 (CH), 2799 (CH), 1659 (C=O), 1596 (NO₂), 1502 (NO₂); δ_{H} (250MHz; CDCl₃) 2.16 (3H, s, CH₃), 2.93 (12H, s, NCH₃), 6.59-6.62 (2H, d, *J* 8.5, ArH), 6.69-6.71 (2H, d, *J* 2.75, ArH), 7.08-7.47 (2H, brd s, ArH); δ_{C} (62.9MHz; CDCl₃) 170.3 (C=O), 148.9 (ArC), 127.2 (ArC), 127.1 (ArC), 127.0 (ArC), 110.9 (ArC), 110.7 (ArC), 40.7 (NCH₃), 22.9 (CH₃); *m/z* (ES) 284.2 (100%, [M - OAc]⁺), 328.1 (15%, [M + H]⁺), 350.1 (41%, [M + Na]⁺).

Synthesis 15

1-(3,7-Bis-dimethylamino-phenothiazin-10-yl)-ethanone




Synthesis using sodium borohydride/ pyridine in one pot. To a 500 cm³ round bottom

5 flask under an atmosphere of nitrogen was added methylthionium chloride trihydrate (0.134 mol, 50 g) and acetonitrile (250 cm³). Sodium borohydride (0.174 mol, 6.6 g) was added in four equal portions over a 30 minute time period. The temperature of the mixture was maintained at 35°C with a cold water bath and was stirred for 30 minutes. The yellow/green suspension had acetic anhydride (0.535 mol, 55 g) and pyridine (0.174 mol, 13.76 g) added. The mixture was heated at 90°C for 2 hours. The cooled mixture was then poured carefully into ice water (250 cm³) in ten equal portions while stirring to give a precipitate. The precipitate was stirred for 30 minutes before it was filtered, washed with water (3 x 50 cm³), and air dried for 30 minutes. The crude material was crystallised from hot ethanol (500 cm³) to yield the title compound (26.7 g, 61%) as a light grey solid. Mp 137°C; $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ 2910 (CH), 2876 (CH), 2856 (CH), 2799 (CH), 1659 (C=O), 1596 (NO₂), 1502 (NO₂); δ_{H} (250MHz; CDCl₃) 2.16 (3H, s, CH₃), 2.93 (12H, s, NCH₃), 6.59-6.62 (2H, d, J 8.5, ArH), 6.69-6.71 (2H, d, J 2.75, ArH), 7.08-7.47 (2H, brd s, ArH); δ_{C} (62.9MHz; CDCl₃) 170.3 (C=O), 148.9 (ArC), 127.2 (ArC), 127.1 (ArC), 127.0 (ArC), 110.9 (ArC), 110.7 (ArC), 40.7 (NCH₃), 22.9 (CH₃); m/z (ES) 284.2 (100%, [M - OAc]⁺), 328.1 (15%, [M + H]⁺), 350.1 (41%, [M + Na]⁺).

Synthesis 16

1-(3,7-Bis-dimethylamino-phenothiazin-10-yl)-ethanone



25 Synthesis using sodium borohydride/Hunig's base in one pot. To a 500 cm³ round bottom flask under an atmosphere of nitrogen was added methylthionium chloride trihydrate (80.2 mmol, 30 g) and acetonitrile (150 cm³). Sodium borohydride (104 mmol, 3.94 g) was added in four equal portions over a 30 minute time period. The temperature of the mixture was maintained at 35°C with a cold water bath and was stirred for 30 minutes.


30 The yellow/green suspension had acetic anhydride (321 mmol, 32.75 g) and Hunig's base (diisopropylethylamine) (120 mmol, 15.55 g) added. The mixture was heated at 90°C for 2 hours. The cooled mixture was then poured carefully into ice water (200 cm³) in ten

- 62 -

equal portions while stirring to give a precipitate. The precipitate was stirred for 30 minutes before it was filtered, washed with water (3 x 50 cm³), and air dried for 30 minutes. The crude material was crystallised from hot ethanol (300 cm³) to yield the title compound (13.55g, 52%) as a light grey solid. Mp 137°C; $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ 2910 (CH), 2876 (CH), 2856 (CH), 2799 (CH), 1659 (C=O), 1596 (NO₂), 1502 (NO₂); δ_{H} (250MHz; CDCl₃) 2.16 (3H, s, CH₃), 2.93 (12H, s, NCH₃), 6.59-6.62 (2H, d, *J* 8.5, ArH), 6.69-6.71 (2H, d, *J* 2.75, ArH), 7.08-7.47 (2H, brd s, ArH); δ_{C} (62.9MHz; CDCl₃) 170.3 (C=O), 148.9 (ArC), 127.2 (ArC), 127.1 (ArC), 127.0 (ArC), 110.9 (ArC), 110.7 (ArC), 40.7 (NCH₃), 22.9 (CH₃); *m/z* (ES) 284.2 (100%, [M - OAc]⁺), 328.1 (15%, [M + H]⁺), 350.1 (41%, [M + Na]⁺).

Synthesis 17


1-(3,7-Bis-dimethylamino-phenothiazin-10-yl)-ethanone



- 15 Synthesis using hydrazine monohydrate/pyridine in one pot. To a 250 cm³ round bottom flask under an atmosphere of nitrogen was added methylthioninium chloride trihydrate (26.74 mmol, 10 g) and acetonitrile (50 cm³). Hydrazine monohydrate (58.8 mmol, 2.95 g) was added and the mixture was heated to reflux and stirred for 10 minutes before cooling to 25°C. The yellow/green suspension had acetic anhydride (424 mmol, 43.3 g)
- 20 and pyridine (124 mmol, 9.78 g) added. The mixture was heated at 90°C for 2 hours. The cooled mixture was then poured carefully into ice water (100 cm³) in ten equal portions while stirring to give a precipitate. The precipitate was stirred for 30 minutes before it was filtered, washed with water (3 x 50 cm³), and air dried for 30 minutes. The crude material was crystallised from hot ethanol (100 cm³) to yield the title compound
- 25 (4.87 g, 56%) as a light grey solid. Mp 137°C; $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ 2910 (CH), 2876 (CH), 2856 (CH), 2799 (CH), 1659 (C=O), 1596 (NO₂), 1502 (NO₂); δ_{H} (250MHz; CDCl₃) 2.16 (3H, s, CH₃), 2.93 (12H, s, NCH₃), 6.59-6.62 (2H, d, *J* 8.5, ArH), 6.69-6.71 (2H, d, *J* 2.75, ArH), 7.08-7.47 (2H, brd s, ArH); δ_{C} (62.9MHz; CDCl₃) 170.3 (C=O), 148.9 (ArC), 127.2 (ArC), 127.1 (ArC), 127.0 (ArC), 110.9 (ArC), 110.7 (ArC), 40.7 (NCH₃), 22.9 (CH₃); *m/z* (ES) 284.2 (100%, [M - OAc]⁺), 328.1 (15%, [M + H]⁺), 350.1 (41%, [M + Na]⁺).
- 30

Synthesis 18


1-(3,7-Bis-dimethylamino-phenothiazin-10-yl)-ethanone



- Synthesis using hydrazine monohydrate/Hunig's base in one pot. To a 250 cm³ round bottom flask under an atmosphere of nitrogen was added methylthioninium chloride trihydrate (80.2 mmol, 30 g) and acetonitrile (150 cm³). Hydrazine monohydrate (176.5 mmol, 8.84 g) was added and the mixture was heated to reflux and stirred for 10 minutes before cooling to 25°C. The yellow/green suspension had acetic anhydride (794 mmol, 81.2 g) and Hunig's base (diisopropylethylamine) (232 mmol, 29.97 g) added.
- The mixture was heated at 90°C for 2 hours. The cooled mixture was then poured carefully into ice water (400 cm³) in ten equal portions while stirring to give a precipitate. The precipitate was stirred for 30 minutes before it was filtered, washed with water (3 x 100 cm³), and air dried for 30 minutes. The crude material was crystallised from hot ethanol (400 cm³) to yield the title compound (17.15g, 65%) as a light grey solid. Mp 137°C; $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ 2910 (CH), 2876 (CH), 2856 (CH), 2799 (CH), 1659 (C=O), 1596 (NO₂), 1502 (NO₂); δ_{H} (250MHz; CDCl₃) 2.16 (3H, s, CH₃), 2.93 (12H, s, NCH₃), 6.59-6.62 (2H, d, *J* 8.5, ArH), 6.69-6.71 (2H, d, *J* 2.75, ArH), 7.08-7.47 (2H, brd s, ArH); δ_{C} (62.9MHz; CDCl₃) 170.3 (C=O), 148.9 (ArC), 127.2 (ArC), 127.1 (ArC), 127.0 (ArC), 110.9 (ArC), 110.7 (ArC), 40.7 (NCH₃), 22.9 (CH₃); *m/z* (ES) 284.2 (100%, [M - OAc]⁺), 328.1 (15%, [M + H]⁺), 350.1 (41%, [M + Na]⁺).

Synthesis 19

3,11-Dinitro-10H-phenothiazine



- 10H-Phenothiazine (20.00 g, 100 mmol), dichloromethane (100 cm³) and acetic acid (40 cm³) had sodium nitrite (20.07 g, 300 mmol) added and the mixture was stirred for 10 minutes at room temperature. Additional acetic acid (40 cm³), dichloromethane (100 cm³) and sodium nitrite (20.07 g, 300 mmol) were then added. A further 120 cm³ of acetic acid was added to try and break up the thick reaction mixture. The mixture was stirred for 3 hours. The suspension was filtered and washed with 100 cm³ each of ethanol, water, and finally ethanol to give a purple/brown solid. The residue was stirred in hot DMF and allowed to cool before filtering the dinitro product, which was washed with ethanol (150 cm³) and dried to give the title compound (24.88 g, 86%) as a brown solid;


- 64 -

$\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ 3331 (NH), 3294 (NH), 3229 (NH), 3101 (CH), 3067 (CH), 1602 (NO₂), 1558 (NO₂); δ_{H} (250MHz; DMSO) 6.73-6.76 (2H, d, *J* 9, ArH), 7.78 (2H, s, ArH), 7.89-7.85 (2H, d, *J* 9, ArH).

5

Synthesis 20


1-(3,7-Bis-diethylamino-phenothiazin-10-yl)-ethanone



- To a 250 cm³ round bottom flask under an atmosphere of nitrogen was added ethylthioninium nitrate monohydrate (7.13 mmol, 3 g) and acetonitrile (20 cm³). Hydrazine monohydrate (16.4 mmol, 0.82 g) was added and the mixture was heated to reflux and stirred for 10 minutes before cooling to 25°C. The brown solution had acetic anhydride (114 mmol, 11.65 g) and Hunig's base (diisopropylethylamine) (21.4 mmol, 2.77 g) was added. The mixture was heated at 90°C for 2 hours. The cooled mixture was then poured carefully into ice water (40 cm³) in ten equal portions while stirring to give a precipitate. The precipitate was stirred for 30 minutes before it was filtered, washed with water (3 x 25 cm³) and air dried for 30 minutes. The crude material was crystallised from hot ethanol (50 cm³) to yield the title compound (1.73 g, 63%) as a light grey solid. δ_{H} (250 MHz; CDCl₃) 7.0-7.5 (2H, brds, ArH), 6.64 (2H, s, ArH), 6.52 (2H, d, ArH), 3.35 (8H, q, 7, NCH₂), 2.18 (3H, s, CH₃), 1.16 (12H, t, 7, CH₃); δ_{C} (62.9 MHz; CDCl₃) 12.5 (CH₃), 22.9 (CH₃), 44.6 (NCH₂), 110.1 (ArC), 127.4 (ArC), 146.5 (ArC), 170.2 (C=O).

Synthesis 21

N,N,N',N'-Tetraethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride)




25

- To a round bottom flask was added 1-(3,7-bis-diethylamino-phenothiazin-10-yl)-ethanone (0.5 g, 1.30 mmol), ethanol (5 cm³), and hydrochloric acid (37%, 1.3 cm³) and the solution was heated at 80°C for 1 hour. Once cooled to room temperature, the mixture was concentrated giving the title compound (0.54 g, 100%) as a light green glass. δ_{H} (250 MHz; CD₃OD) 7.07 (4H, brd, ArH), 6.65 (2H, brd, ArH), 3.35 (8H, brd, NCH₂), 0.97 (12H, brd, CH₃); δ_{C} (62.9 MHz; CD₃OD) 10.8 (CH₃), 55.1 (NCH₂), 116.6 (ArC), 120.4 (ArC), 121.5 (ArC), 123.6 (ArC), 132.6 (ArC), 144.5 (ArC).

30

Synthesis 22


N,N,N',N'-Tetraethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide)



- 5 To a round bottom flask was added 1-(3,7-bis-diethylamino-phenothiazin-10-yl)-ethanone (0.5 g, 1.30 mmol), ethanol (5 cm³), and hydrobromic acid (48%, 0.75 cm³) and the solution was heated at 80°C for 1 hour. Once cooled to room temperature, the mixture was concentrated giving the title compound (0.65g, 100%) as a light yellow glass. δ_H (250 MHz; D₂O) 7.05 (4H, brd, ArH), 6.79 (2H, brd d, ArH), 3.43 (8H, brd, NCH₂), 1.05 (12H, brd t, CH₃); δ_C (62.9 MHz; D₂O) 12.3 (CH₃), 56.2 (NCH₂), 117.9 (ArC), 121.4 (ArC), 122.4 (ArC), 124.5 (ArC), 133.5 (ArC), 145.1 (ArC).

Synthesis 23

N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride)




- 15 To a round bottom flask was added 1-(3,7-bis-dimethylamino-phenothiazin-10-yl)-ethanone (1 g, 3.05 mmol), ethanol (10 cm³), and hydrochloric acid (37%, 3 cm³) and the solution was heated at 80°C for 1 hour. Once cooled to room temperature, diethyl ether was added while stirring until a constant turbid solution was obtained. After some time, a precipitate formed, which was filtered and washed with diethyl ether (10 cm³) giving the title compound (0.98 g, 90%) as a light green solid. Mp (dec) 230°C; ν_{max} (KBr)/cm⁻¹ 3500-3229 (NH), 3061 (CH), 3021 (CH), 2948 (CH), 2879 (CH), 2679 (CH), 2601 (CH), 1604 (CH), 1483 (CH), 1318 (CH); δ_H (250MHz; D₂O) 3.18 (12H, s, NCH₃), 6.67 (2H, d, J 8.5, ArH), 7.16 (4H, brd s, ArH); δ_C (62.9MHz; D₂O) 144.3 (ArC), 138.9 (ArC), 122.4 (ArC), 120.8 (ArC), 120.7 (ArC), 117.6 (ArC), 48.9 (NCH₃); m/z (ES) 286.1 (100%, [M - H, 2Cl]⁺), 285.1 (40%), 284.1 (41%, [M - 3H, 2Cl]⁺).

- 66 -

Synthesis 24


N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide)



To a round bottom flask was added 1-(3,7-bis-dimethylamino-phenothiazin-10-yl)-ethanone (1 g, 3.05 mmol), ethanol (10 cm³), and hydrobromic acid (48%, 4 cm³) and the solution was heated at 80°C for 1 hour. Once cooled to room temperature, a precipitate formed, which was filtered and washed with diethyl ether (10 cm³) giving the product (1.22 g, 89%) as a light mustard solid. Mp (dec) 230°C; $\nu_{\max}(\text{KBr})/\text{cm}^{-1}$ 3500-3229 (NH), 3061 (CH), 3021 (CH), 2948 (CH), 2879 (CH), 2679 (CH), 2601 (CH), 1604 (CH), 1483 (CH), 1318 (CH); δ_{H} (250MHz; D₂O) 3.18 (12H, s, NCH₃), 6.66 (2H, d, J 8.75, ArH), 7.15 (4H, s, ArH); δ_{C} (62.9MHz; D₂O) 144.3 (ArC), 138.9 (ArC), 122.4 (ArC), 120.8 (ArC), 120.7 (ArC), 117.6 (ArC), 48.9 (NCH₃).

Synthesis 25

N,N,N',N'-Tetraethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide)



To a round bottom flask was added 1-(3,7-bis-diethylamino-phenothiazin-10-yl)-ethanone (1.0 g, 2.60 mmol), methanol (10 cm³), and hydrobromic acid (48%, 2.94 cm³) and the solution was heated at 80°C for 1 hour. Once cooled to 5°C, the mixture had diethyl ether added, giving a cloudy solution. The solution was stirred for 30 minutes and gave the title compound (0.83 g, 63%) as a light yellow solid. δ_{H} (250 MHz; D₂O) 7.05 (4H, brd, ArH), 6.79 (2H, brd d, ArH), 3.43 (8H, brd, NCH₂), 1.05 (12H, brd t, CH₃); δ_{C} (62.9 MHz; D₂O) 12.3 (CH₃), 56.2 (NCH₂), 117.9 (ArC), 121.4 (ArC), 122.4 (ArC), 124.5 (ArC), 133.5 (ArC), 145.1 (ArC).

Stability Studies

The DAPTZ compounds of the present invention are stably reduced (i.e., are in a stably reduced form). For example, they are stable in the solid form, for example, for at least 1 week, e.g., at least 2 weeks, e.g., at least 1 month, e.g., at least 2 months, e.g., at least 1 year (e.g., at room temperature, e.g., 18-25°C, e.g., in a sealed container).

One sample of compound B3 (N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride)), in solid form, was found to be substantially reduced even after 2 years in storage.

Once the DAPTZ compounds are dissolved in water (i.e., in the form of an aqueous solution), they slowly oxidize (giving the solution a blue colour), typically over a period of 1 to 3 hours.

The stability of two DAPTZ compounds of the present invention was studied, specifically, B3 (N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride)), and B6 (N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen iodide)). MTC was used as a standard.

The compounds were weighed into universal containers. Enough water was added to yield a 1 mM solution, and the mixture stirred to dissolve the solid. The absorbance was determined at 610 nm and 665 nm for 50 µL samples (in triplicate) of each of the solutions at various time points. The initial time point taken was at 10 minutes as the compounds took time to dissolve completely. A UV/visible spectrum was also recorded at time points 20 minutes, 3 hours, and 18 hours.

The percent reduced form (%) was calculated assuming that the 10 minute reading for MTC represented 0% reduced, and that a blank represented 100% reduced (colourless).

Figure 1 is a graph of the percent reduced form (%) versus time (minutes) for each of three compounds, B1, B3, and B6, as determined using absorbance at 665 nm.

Figure 2 is a graph of the percent reduced form (%) versus time (minutes) for each of three compounds, B1, B3, and B6, as determined using absorbance at 610 nm.

Figures 3A, 3B, and 3C show the UV/visible absorption spectra for aqueous samples of each of three compounds, B1 (open circles), B3 (open squares), and B6 (open triangles), after 20 minutes (Figure 3A), 3 hours (Figure 3B), and 18 hours (Figure 3C).

- 68 -

These data demonstrate that the DAPTZ compounds (stabilized reduced forms) remain substantially stable (>50%) for at least 1 hour, and that compound B6 remains substantially stable (>50%) for almost 3 hours. However, after about 18 hours, the compounds are not significantly different from MTC. See, for example, Figure 3C, where
5 the spectra are almost indistinguishable.

Additionally, the rate of autoxidation was found to be slower for the "iodide" compound (Compound B6) as compared to the "chloride" compound (Compound B3), suggesting that the rate of autoxidation depends upon the counterion. Although the difference in rate
10 was small, it may be significant in drug formulation. Other salts may be more stable against oxidation.

A batch of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide) was prepared in April 2006 and analysed by NMR. After 10 months of storage in the dark
15 at room temperature, the solid material was analysed once more and the NMR data was found to be identical. The colour of the solid remained consistent over time. It appears that the molecule, in this form, is stable under these conditions over this time period.

Biological Studies

20

Methods: In vitro Assay for Establishing B50

These methods are described in detail in WO 96/30766. Briefly, a fragment of tau corresponding to the core repeat domain, which has been adsorbed to a solid phase
25 substrate, is able to capture soluble full-length tau and bind tau with high affinity. This association confers stability against proteolytic digestion of the aggregated tau molecules. The process is self-propagating, and can be blocked selectively by prototype pharmaceutical agents.

30 More specifically, truncated tau (residues 297-390; dGA) diluted in carbonate buffer (pH 9.6) was bound to the assay plate, and full-length tau (T40) was added in the aqueous phase. The aqueous phase binding buffer contained 0.05% Tween-20 and 1% gelatine in phosphate-buffered saline (pH 7.4). Bound tau was detected using mAb 499 that
35 recognises an N-terminal epitope within the aqueous phase full-length tau but that fails to recognise the solid phase-bound truncated tau fragment.

The concentration of compound required to inhibit the tau-tau binding by 50% is referred to as the B50 value.

Methods: Cell-based Assay for Establishing EC50

These methods are described in more detail in WO 02/055720. Briefly, fibroblast cells (3T6) express full-length tau ("T40") under control of an inducible promoter, and low constitutive levels of the PHF-core tau fragment (12 kD fragment). When T40 expression is induced, it undergoes aggregation-dependent truncation within the cell, N-terminally at ~ αα 295 and C-terminally at ~ αα 390, thereby producing higher levels of the 12 kD PHF-core domain fragment. Production of the 12 kD fragment can be blocked in a dose-dependent manner by tau-aggregation inhibitors. Indeed, the quantitation of inhibitory activity of compounds with respect to proteolytic generation of the 12 kD fragment within cells can be described entirely in terms of the same parameters that describe inhibition of tau-tau binding in vitro. That is, the extent of proteolytic generation of the 12 kD fragment within cells is determined entirely by the extent of tau-tau binding through the repeat domain. The availability of the relevant proteases within the cell is non-limiting.

Results are expressed as the concentration at which there is a 50% inhibition of generation of the 12 kD fragment. This is referred to as the EC50 value.

Methods: Toxicity in Cells (LD50) and Therapeutic Index (Rxl)

Toxicity of the compounds described herein was assessed in the cell based assay used to assess EC50. Toxicity was measured by cell numbers determined after 24 hours exposure to the compound using a lactate dehydrogenase assay kit TOX-7 (Sigma Biosciences) according to the manufacturer's instructions after lysis of remaining cells. Alternatively, a kit from Promega UK (CytoTox 96) was used, again according to the manufacturer's instructions.

The therapeutic index (Rxl) was calculated as: $Rxl = LD50 / EC50$.

The data are summarised in the following Table.

Biological Data				
Compound	B50	EC50	LD50	Rxl
B3	57.3	0.50	44.0	88
B4	23.5	2.23	-	-
B6	494	0.38	115	303
MTC	218	0.59	65.0	110

B3: N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride).

B4: 10H-phenothiazine-3,7-diamine bis(hydrogen chloride).

- 70 -

B6: N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen iodide).

MTC: Methylthioninium chloride.

Partition Coefficient Studies

5

It is well known that the partition coefficient for a drug in an organic phase/water system (typically an *n*-octanol/water system), usually reported as the logarithm (i.e., $\log_{10}P$), is a good indicator for the biological activity of that drug. See, e.g., Hansch, C., et al., 1964, J. Am. Chem. Soc., Vol. 86, pp. 1616-1626; Kubinyi, H., 1977, J. Med. Chem., Vol. 20, pp. 625-629. It is believed that this is because the absorption of a compound depends on its partition between the biological membrane and the aqueous phase. Partition coefficients are also useful in separation techniques and in the prediction of the solubility of drugs.

In the context of drug-like substances, hydrophobicity is related to absorption, bioavailability, hydrophobic drug-receptor interactions, metabolism, and toxicity. Low hydrophilicities, and therefore high $\log_{10}P$ values, may cause poor absorption or permeation. It has been shown for compounds to have a reasonable probability of being well absorbed, their $\log_{10}P$ value must not be greater than 5.0. The distribution of calculated $\log_{10}P$ values of more than 3000 drugs on the market underlines this fact.

20

Many methods for determining partition coefficients are known. In this study, aqueous solutions of selected compounds were shaken with *n*-octanol and aliquots concentrations in each phase determined by visible spectrophotometry. The $\log_{10}P$ was then calculated for each compound, using the following formula:

25

$$\log_{10}P = \log_{10} [\text{Drug}]_{\text{octanol}} - \log_{10} [\text{Drug}]_{\text{water}} = \log_{10} ([\text{Drug}]_{\text{octanol}} / [\text{Drug}]_{\text{water}})$$

The data are summarised in the following table. The DAPTZ compounds of the present invention were found to have $\log_{10}P$ values expected for drug-like molecules.

30

Partition Coefficient Data				
Compound	λ_{max} (octanol/ water)	Absorbance for <i>n</i> -octanol phase	Absorbance for water phase	$\log_{10}P$
MTC	665 / 660	0.217	4.83	-1.35
B3	664 / 660	0.083	0.111	-0.13
B6	658.4 / 662.4	0.179	0.563	-0.498

B3: N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen chloride).

B6: N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen iodide).

MTC: Methylthioninium chloride.

- 71 -


Crystal Structure

- Figure 4 shows the crystal structure of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide). The crystal structure shows three crystallographically distinct bromide ions. Br1 and Br2 occupy special positions with 2-fold symmetry, whereas the organic main molecule and Br3 occupy general positions. Hence, the overall stoichiometry is $C_{16}H_{21}N_3SBr_2$.
- Figure 5 shows the side-on view of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide) and reveals the non-planarity; the dihedral angle between the outer benzene rings is 11.0 (3) degrees.
- Figure 6 shows part of one helical column of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis(hydrogen bromide) molecules in the crystal.

- 72 -

CLAIMS

1. A compound selected from compounds of the following formula and pharmaceutically acceptable salts, solvates, and hydrates thereof:



wherein:

each of R^1 and R^9 is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

each of R^{3NA} and R^{3NB} is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

each of R^{7NA} and R^{7NB} is independently selected from: -H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

each of HX^1 and HX^2 is independently a protic acid.

2. A compound according to claim 1, wherein each of R^1 and R^9 is independently -H, -Me, -Et, or -CF₃.
3. A compound according to claim 1, wherein each of R^1 and R^9 is independently -H, -Me, or -Et.
4. A compound according to any one of claims 1 to 3, wherein R^1 and R^9 are the same.
5. A compound according to claim 1, wherein each of R^1 and R^9 is independently -H.
6. A compound according to any one of claims 1 to 5, wherein each of R^{3NA} and R^{3NB} is independently -Me, -Et, -nPr, -nBu, -CH₂-CH=CH₂, or -CF₃.
7. A compound according to any one of claims 1 to 6, wherein R^{3NA} and R^{3NB} are the same.
8. A compound according to any one of claims 1 to 7, wherein each of R^{7NA} and R^{7NB} is independently -Me, -Et, -nPr, -nBu, -CH₂-CH=CH₂, or -CF₃.

- 73 -

9. A compound according to any one of claims 1 to 8, wherein R^{7NA} and R^{7NB} are the same.
- 5 10. A compound according to any one of claims 1 to 9, with the proviso that:
at least one of R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} is *other than* -Et.
11. A compound according to any one of claims 1 to 9, with the proviso that:
if: each of R^1 and R^9 is -H;
then: R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} are not each -Et.
- 10 12. A compound according to any one of claims 1 to 5, wherein each of the groups
-N(R^{3NA})(R^{3NB}) and -N(R^{7NA})(R^{7NB}) is independently selected from: -NMe₂, -NEt₂,
-N(nPr)₂, -N(Bu)₂, -NMeEt, -NMe(nPr), and -N(CH₂CH=CH₂)₂.
- 15 13. A compound according to any one of claims 1 to 5, wherein the groups
-N(R^{3NA})(R^{3NB}) and -N(R^{7NA})(R^{7NB}) are the same, and are selected from: -NMe₂
and -NEt₂.
- 20 14. A compound according to any one of claims 1 to 5, 11, and 12, wherein the
groups -N(R^{3NA})(R^{3NB}) and -N(R^{7NA})(R^{7NB}) are the same.
15. A compound according to any one of claims 1 to 5 and 11 to 13, with the proviso
that: each of the groups -N(R^{3NA})(R^{3NB}) and -N(R^{7NA})(R^{7NB}) is *other than* -NEt₂.
- 25 16. A compound according to any one of claims 1 to 5 and 11 to 13, wherein the
groups -N(R^{3NA})(R^{3NB}) and -N(R^{7NA})(R^{7NB}) are the same, and are selected from:
-NMe₂, -N(nPr)₂, -N(Bu)₂, -NMeEt, -NMe(nPr), and -N(CH₂CH=CH₂)₂.
- 30 17. A compound according to any one of claims 1 to 5, wherein each of the groups
-N(R^{3NA})(R^{3NB}) and -N(R^{7NA})(R^{7NB}) is: -NMe₂.
18. A compound according to any one of claims 1 to 17, wherein each of HX¹ and HX²
is independently a monoprotic acid.
- 35 19. A compound according to any one of claims 1 to 17, wherein each of HX¹ and HX²
is independently a hydrohalide acid.
20. A compound according to any one of claims 1 to 17, wherein each of HX¹ and HX²
is independently selected from HCl, HBr, and HI.


- 74 -

21. A compound according to any one of claims 1 to 17, wherein HX^1 and HX^2 are each HCl.


22. A compound according to any one of claims 1 to 17, wherein HX^1 and HX^2 are each HBr.

23. A compound according to any one of claims 1 to 17, wherein HX^1 and HX^2 are each HI.


24. A compound according to claim 1, selected from compounds of the following formula, and pharmaceutically acceptable salts, solvates, and hydrates thereof:



25. A compound according to claim 1, selected from compounds of the following formula, and pharmaceutically acceptable salts, solvates, and hydrates thereof:



26. A compound according to claim 1, selected from compounds of the following formula, and pharmaceutically acceptable salts, solvates, and hydrates thereof:



27. A compound according to any one of claims 1 to 26, in substantially purified form.

- 75 -

28. A compound according to any one of claims 1 to 27, wherein one or more of the carbon atoms of the compound is ^{11}C .
- 5 29. A composition comprising a compound according to any one of claims 1 to 28, and a pharmaceutically acceptable carrier or diluent.
30. A pharmaceutical composition comprising a compound according to any one of claims 1 to 28, and a pharmaceutically acceptable carrier or diluent.
- 10 31. A method of preparing a pharmaceutical composition comprising admixing a compound according to any one of claims 1 to 28, and a pharmaceutically acceptable carrier or diluent.
- 15 32. A compound according to any one of claims 1 to 27 for use in a method of treatment or prophylaxis of the human or animal body by therapy.
33. A compound according to any one of claims 1 to 27 for use in a method of treatment or prophylaxis of a disease of protein aggregation.
- 20 34. A compound according to any one of claims 1 to 27 for use in a method of treatment or prophylaxis of a tauopathy.
35. A compound according to any one of claims 1 to 27 for use in a method of treatment or prophylaxis of a neurodegenerative tauopathy.
- 25 36. A compound according to any one of claims 1 to 27 for use in a method of treatment or prophylaxis of Alzheimer's disease.
- 30 37. A compound according to any one of claims 1 to 27 for use in a method of treatment or prophylaxis of skin cancer or melanoma.
38. A compound according to any one of claims 1 to 27 for use in a method of treatment or prophylaxis of a viral, bacterial or protozoal disease condition.
- 35 39. A compound according to any one of claims 1 to 27 for use in a method of treatment or prophylaxis of Hepatitis C, HIV, or West Nile Virus (WNV).
40. A compound according to any one of claims 1 to 27 for use in a method of treatment or prophylaxis of malaria.
- 40

41. Use of a compound according to any one of claims 1 to 27 in the manufacture of a medicament for use in the treatment or prophylaxis of a disease of protein aggregation.
42. Use of a compound according to any one of claims 1 to 27 in the manufacture of a medicament for use in the treatment or prophylaxis of a tauopathy.
43. Use of a compound according to any one of claims 1 to 27 in the manufacture of a medicament for use in the treatment or prophylaxis of a neurodegenerative tauopathy.
44. Use of a compound according to any one of claims 1 to 27 in the manufacture of a medicament for use in the treatment or prophylaxis of Alzheimer's disease.
45. Use of a compound according to any one of claims 1 to 27 in the manufacture of a medicament for use in the treatment or prophylaxis of skin cancer or melanoma.
46. Use of a compound according to any one of claims 1 to 27 in the manufacture of a medicament for use in the treatment or prophylaxis of a viral, bacterial or protozoal disease condition.
47. Use of a compound according to any one of claims 1 to 27 in the manufacture of a medicament for use in the treatment or prophylaxis of Hepatitis C, HIV, or West Nile Virus (WNV).
48. Use of a compound according to any one of claims 1 to 27 in the manufacture of a medicament for use in the treatment or prophylaxis of malaria.

49. An *in vitro* method of reversing and/ or inhibiting the aggregation of a protein comprising contacting the protein with an effective amount of a compound according to any one of claims 1 to 27.
50. A method of inactivating a pathogen in a sample, comprising the steps of introducing a compound according to any one of claims 1 to 27 into the sample, and then exposing the sample to light.