

COMPOUNDS AND USES THEREOF FOR THE MODULATION OF HEMOGLOBIN

FIELD OF THE INVENTION

[0001] This invention provides compounds and pharmaceutical compositions suitable as allosteric modulators of hemoglobin, methods and intermediates for their preparation, and methods for their use in treating disorders mediated by hemoglobin and disorders that would benefit from tissue and/or cellular oxygenation.

STATE OF THE ART


[0002] Sickle cell disease is a disorder of the red blood cells, found particularly among those of African and Mediterranean descent. The basis for sickle cell disease is found in sickle hemoglobin (HbS), which contains a point mutation relative to the prevalent peptide sequence of hemoglobin (Hb).

[0003] Hemoglobin (Hb) transports oxygen molecules from the lungs to various tissues and organs throughout the body. Hemoglobin binds and releases oxygen through conformational changes. Sickle hemoglobin (HbS) contains a point mutation where glutamic acid is replaced with valine, allowing HbS to become susceptible to polymerization to give the HbS containing red blood cells their characteristic sickle shape. The sickled cells are also more rigid than normal red blood cells, and their lack of flexibility can lead to blockage of blood vessels. U.S. 7,160,910 discloses compounds that are allosteric modulators of hemoglobin. However, a need exists for additional therapeutics that can treat disorders that are mediated by Hb or by abnormal Hb such as HbS.

SUMMARY OF THE INVENTION

[0004] This invention relates generally to compounds and pharmaceutical compositions suitable as allosteric modulators of hemoglobin. In some aspects, this invention relates to methods for treating disorders mediated by hemoglobin and disorders that would benefit from tissue and/or cellular oxygenation.

[0005] In certain aspects of the invention, a compound of formula (I) is provided:




or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein L^{10} is optionally substituted methylene or, preferably, a bond;

 is a single or a double bond;

each X and Y is independently $(CR^{20}R^{21})_e$, O, S, SO, SO₂, or NR²⁰; e is 1 to 4, preferably 1; each R²⁰ and R²¹ independently is hydrogen or C₁-C₃ alkyl optionally substituted with 1-3 halo, OH, or C₁-C₆ alkoxy, or $CR^{20}R^{21}$ is C=O, provided that if one of X and Y is O, S, SO, SO₂, then the other is not CO, and X and Y are both not heteroatoms or oxidized forms thereof;

V¹ and V² independently are C₁-C₆ alkoxy; or V¹ and V² together with the carbon atom they are attached to form a ring of formula:



wherein each V³ and V⁴ are independently O, S, or NH, provided that when one of V³ and V⁴ is S, the other is NH, and provided that V³ and V⁴ are both not NH; q is 1 or 2; each V⁵ is independently C₁-C₆ alkyl optionally substituted with 1-3 OH groups, or V⁵ is CO₂R⁶⁰, where each R⁶⁰ independently is C₁-C₆ alkyl or hydrogen; t is 0, 1, 2, or 4; or CV¹V² is C=V, wherein V is O, NOR⁸⁰, or NNR⁸¹R⁸²;

R^{80} is optionally substituted C_1 - C_6 alkyl;

R^{81} and R^{82} independently are selected from the group consisting of hydrogen, optionally substituted C_1 - C_6 alkyl, COR^{83} , or CO_2R^{84} ;

R^{83} is hydrogen or optionally substituted C_1 - C_6 alkyl; and

R^{84} is optionally substituted C_1 - C_6 alkyl;


provided that the compounds provided herein exclude those disclosed in U.S. patent application nos. 13/730,730 and 13/730,674; and

provided that the compounds provided herein exclude those in Table 1 hereinbelow; and A, B, and C are defined as follows.

[0006] In one instance,

ring A is C_6 - C_{10} aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, or heteroaryl is optionally substituted, preferably with 1-4 C_1 - C_6 alkyl and/or C_1 - C_6 alkoxy groups;

ring B is:



wherein ring B' including the -N-CO- moiety is a 5-6 membered heterocycle containing up to 3 heteroatoms selected from nitrogen, oxygen, and sulfur and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted, preferably with 1-4 C_1 - C_6 alkyl groups;

ring C is C_6 - C_{10} aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted, preferably with 1-4: halo, oxo, $-OR^1$, C_1 - C_6 alkyl, $-COOR^1$, and NR^5R^6 ,

R^1 is a hydrogen, C_1 - C_6 alkyl or a prodrug moiety; wherein the alkyl is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring

heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, which is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein the heteroaryl is optionally substituted with C₁-C₆ alkyl;

R⁵ and R⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl or -COOR³;

R³ is hydrogen or optionally substituted C₁-C₆ alkyl;

with the proviso that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;

then ring A excludes optionally substituted 5-10 membered heteroaryl;

and provided that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 5-10 membered heteroaryl;

then ring A is not optionally substituted 4-10 membered heterocycle.

[0007] In another instance:

ring A is C₆-C₁₀ aryl, a C₃-C₈ cycloalkyl, a 5-10 membered heteroaryl or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, heteroaryl, cycloalkyl, or heterocycle is optionally substituted, preferably with 1-4: halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, and/or C₃-C₁₀ cycloalkyl, wherein the C₁-C₆ alkyl is optionally substituted with 1-5 halo, C₁-C₆ alkoxy, and/or C₃-C₁₀ cycloalkyl;

ring B is a 5-10 membered heteroaryl or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted, preferably with 1-4: halo, C₁-C₆ alkyl and/or -CO-C₁-C₆ alkyl,



is a single or a double bond;

ring C is C₆-C₁₀ aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted, preferably with 1-4: halo, oxo, -OR¹, C₁-C₆ alkyl, -COOR¹, and/or C₁-C₆ alkoxy, wherein the C₁-C₆ alkyl is optionally substituted with 1-5 halo, C₁-C₆ alkoxy and/or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S; and

R¹ is hydrogen or a prodrug moiety;

provided that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;


then ring A excludes optionally substituted 5-10 membered heteroaryl;

and provided that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 5-10 membered heteroaryl;


then ring A is not optionally substituted 4-10 membered heterocycle.

[0008] In certain aspects of the invention, a compound of formula (X-I) is provided:




an N-oxide thereof, or a tautomer or each thereof, or a pharmaceutically acceptable salt of each of the preceding, wherein

ring A is phenyl optionally substituted with 1-3 halo and/or C₁-C₆ alkoxy, or is a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, optionally substituted, or is



wherein R^7 is C_1 - C_6 alkyl, optionally substituted with 3-5 fluoro groups, or is C_3 - C_6 cycloalkyl;

ring B is selected from the group consisting of



wherein R^8 is C_1 - C_6 alkyl, $-\text{CO}-C_1$ - C_6 alkyl or a prodrug moiety and wherein the pyridyl ring is optionally substituted with a halo or an $\text{NR}^{25}(\text{CH}_2)_2\text{N}(\text{R}^{25})_2$ group where each R^{25} is independently hydrogen or C_1 - C_6 alkyl;


X is O, S, SO, or SO_2 ;

 is a single or a double bond;

ring C is phenyl or a 6 membered nitrogen-containing heteroaryl, each of which is optionally substituted with 1-4: halo, oxo, $-\text{OR}^1$, C_1 - C_6 alkyl, $-\text{COOR}^1$ and/or C_1 - C_6 alkoxy, wherein the C_1 - C_6 alkyl is optionally substituted with 1-5 halo, C_1 - C_6 alkoxy and/or 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S; and

each R^1 is hydrogen or a prodrug moiety R;

V^1 and V^2 independently are C_1 - C_6 alkoxy; or V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:



wherein each V^3 and V^4 are independently O, S, or NH, provided that when one of V^3 and V^4 is S, the other is NH, and provided that V^3 and V^4 are both not NH; q is

1 or 2; each V^5 is independently C_1 - C_6 alkyl or CO_2R^{60} , where each R^{60} independently is C_1 - C_6 alkyl or hydrogen; t is 1, 2, or 4; or CV^1V^2 is $C=V$, wherein V is O, NOR^{80} , or $NNR^{81}R^{82}$;

wherein R^{80} is optionally substituted C_1 - C_6 alkyl;

R^{81} and R^{82} independently are selected from the group consisting of hydrogen, optionally substituted C_1 - C_6 alkyl, COR^{83} , or CO_2R^{84} ;

R^{83} is hydrogen or optionally substituted C_1 - C_6 alkyl;

R^{84} is optionally substituted C_1 - C_6 alkyl;

provided that when ring C is C_6 - C_{10} aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;

then ring A excludes optionally substituted 5-10 membered heteroaryl;

provided that when ring C is C_6 - C_{10} aryl;


and ring B is optionally substituted 5-10 membered heteroaryl;

then ring A is not optionally substituted 4-10 membered heterocycle;


provided that the compounds provided herein exclude those disclosed in U.S. patent application nos. 13/730,730 and 13/730,674; and

provided that the compounds provided herein exclude those in Table 1 hereinbelow.

[0009] Preferably, V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:




[0010] In some embodiments, V^1 and V^2 independently are C_1 - C_6 alkoxy; or V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:




wherein each V^3 and V^4 are independently O, S, or NH, provided that when one or V^3 and V^4 is S the other is NH, and provided that V^3 and V^4 are both not NH; q is 1 or 2; each V^5 is independently C_1 - C_6 alkyl or CO_2R^{60} , where each R^{60} independently is C_1 - C_6 alkyl or hydrogen; t is 0, 1, 2, or 4; or CV^1V^2 is $C=V$, wherein V is O, and wherein the remaining variables are defined herein.

[0011] In certain embodiments, a compound of formula (II) is provided:



wherein the remaining variables are defined herein.

[0012] In certain embodiments, a compound selected from formulas (IIA), (IIB) and (IIC) is provided:



wherein

R^9 is hydrogen, $-OR^1$, C_1 - C_6 alkoxy optionally substituted with 1-3 C_1 - C_6 alkoxy or 4-10 membered heterocycle containing up to 5 ring heteroatoms selected from N, O, S or oxidized forms thereof;

R^{10} is hydrogen, hydroxy, halo or C_1 - C_6 alkoxy;

R^{11} is hydrogen or C_1 - C_6 alkyl; and

R^{12} is $-OR^1$;

wherein R^1 is hydrogen or the prodrug moiety R.


[0013] In certain aspects of the invention, a compound of formula (I) is provided:

or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein

L^{10} is optionally substituted methylene or, preferably, a bond;

ring A is C_6 - C_{10} aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, or heteroaryl is optionally substituted with 1-4 C_1 - C_6 alkyl and/or C_1 - C_6 alkoxy groups;

ring B is:



wherein ring B' including the $-N-CO-$ moiety is a 5-6 membered heterocycle containing up to 3 heteroatoms selected from nitrogen, oxygen, and sulfur and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted with 1-4 C_1 - C_6 alkyl groups;

each X and Y is independently $(CR^{20}R^{21})_e$, O, S, SO, SO_2 , or NR^{20} ; e is 1 to 4, preferably 1; each R^{20} and R^{21} independently is hydrogen or C_1 - C_3 alkyl optionally substituted with 1-3 halo, OH, or C_1 - C_6 alkoxy, or $CR^{20}R^{21}$ is $C=O$, provided that if one of X and Y is O, S, SO, SO_2 , then the other is not CO, and X and Y are both not heteroatoms or oxidized forms thereof;


ring C is C_6 - C_{10} aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, $-OR^1$, C_1 - C_6 alkyl, $-COOR^5$, NR^5R^6 ,

R^1 is a hydrogen, C_1 - C_6 alkyl or a prodrug moiety; wherein the alkyl is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, which is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein the heteroaryl is optionally substituted with C_1 - C_6 alkyl;

R^5 and R^6 are each independently hydrogen, optionally substituted C_1 - C_6 alkyl or $-COOR^3$;

R^3 is hydrogen or optionally substituted C_1 - C_6 alkyl;

V^1 and V^2 independently are C_1 - C_6 alkoxy; or V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:



wherein each V^3 and V^4 are independently O, S, or NH, provided that when one of V^3 and V^4 is S, the other is NH, and provided that V^3 and V^4 are both not NH; q is 1 or 2; each V^5 is independently C_1 - C_6 alkyl optionally substituted with 1-3 OH groups, or V^5 is CO_2R^{60} , where each R^{60} independently is C_1 - C_6 alkyl or hydrogen; t is 0, 1, 2, or 4; or CV^1V^2 is $C=V$, wherein V is O, NOR^{80} , or $NNR^{81}R^{82}$;

R^{80} is optionally substituted C_1 - C_6 alkyl;

R^{81} and R^{82} independently are selected from the group consisting of hydrogen, optionally substituted C_1 - C_6 alkyl, COR^{83} , or CO_2R^{84} ;

R^{83} is hydrogen or optionally substituted C_1 - C_6 alkyl; and

R^{84} is optionally substituted C_1 - C_6 alkyl;

with the proviso that when ring C is C_6 - C_{10} aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;

then ring A excludes optionally substituted 5-10 membered heteroaryl;

and provided that when ring C is C₆-C₁₀ aryl;
and ring B is optionally substituted 5-10 membered heteroaryl;
then ring A is not optionally substituted 4-10 membered heterocycle.

[0014] In further aspects of the invention, a composition is provided comprising any of the compounds described herein, and at least a pharmaceutically acceptable excipient.

[0015] In still further aspects of the invention, a method is provided for increasing oxygen affinity of hemoglobin S in a subject, the method comprising administering to a subject in need thereof a therapeutically effective amount of any of the compounds or compositions described herein.

[0016] In further aspects of the invention, a method is provided for treating oxygen deficiency associated with sickle cell anemia, the method comprising administering to a subject in need thereof a therapeutically effective amount of any of the compounds or compositions described herein.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0017] It must be noted that as used herein and in the appended claims, the singular forms “a”, “an”, and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a solvent” includes a plurality of such solvents.

[0018] As used herein, the term “comprising” or “comprises” is intended to mean that the compositions and methods include the recited elements, but not excluding others. “Consisting essentially of” when used to define compositions and methods, shall mean excluding other elements of any essential significance to the combination for the stated purpose. Thus, a composition or process consisting essentially of the elements as defined herein would not exclude other materials or steps that do not materially affect the basic and novel characteristic(s) of the claimed invention. “Consisting of” shall mean excluding more than trace elements of other ingredients and substantial method steps. Embodiments defined by each of these transition terms are within the scope of this invention.

[0019] Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about.” Accordingly, unless indicated to the contrary, the numerical parameters set forth in the following specification and attached


claims are approximations. Each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. The term “about” when used before a numerical designation, *e.g.*, temperature, time, amount, and concentration, including range, indicates approximations which may vary by (+) or (-) 10 %, 5 % or 1 %.

[0020] As used herein, C_m-C_n, such as C₁-C₁₂, C₁-C₈, or C₁-C₆ when used before a group refers to that group containing m to n carbon atoms.

[0021] The term “alkoxy” refers to -O-alkyl.

[0022] The term “alkyl” refers to monovalent saturated aliphatic hydrocarbyl groups having from 1 to 12 carbon atoms (*i.e.*, C₁-C₁₂ alkyl) or 1 to 8 carbon atoms (*i.e.*, C₁-C₈ alkyl), or 1 to 4 carbon atoms. This term includes, by way of example, linear and branched hydrocarbyl groups such as methyl (CH₃-), ethyl (CH₃CH₂-), *n*-propyl (CH₃CH₂CH₂-), isopropyl ((CH₃)₂CH-), *n*-butyl (CH₃CH₂CH₂CH₂-), isobutyl ((CH₃)₂CHCH₂-), *sec*-butyl ((CH₃)(CH₃CH₂)CH-), *t*-butyl ((CH₃)₃C-), *n*-pentyl (CH₃CH₂CH₂CH₂CH₂-), and neopentyl ((CH₃)₃CCH₂-).

[0023] The term “aryl” refers to a monovalent, aromatic mono- or bicyclic ring having 6-10 ring carbon atoms. Examples of aryl include phenyl and naphthyl. The condensed ring may or may not be aromatic provided that the point of attachment is at an aromatic carbon atom. For example, and without limitation, the following is an aryl group:




[0024] The term “-CO₂H ester” refers to an ester formed between the -CO₂H group and an alcohol, preferably an aliphatic alcohol. A preferred example included -CO₂R^E, wherein R^E is alkyl or aryl group optionally substituted with an amino group.

[0025] The term “chiral moiety” refers to a moiety that is chiral. Such a moiety can possess one or more asymmetric centers. Preferably, the chiral moiety is enantiomerically enriched, and more preferably a single enantiomer. Non limiting examples of chiral moieties include chiral carboxylic acids, chiral amines, chiral amino acids, such as the naturally occurring amino acids, chiral alcohols including chiral steroids, and the likes.


[0026] The term “cycloalkyl” refers to a monovalent, preferably saturated, hydrocarbyl mono-, bi-, or tricyclic ring having 3-12 ring carbon atoms. While cycloalkyl, refers

preferably to saturated hydrocarbyl rings, as used herein, it also includes rings containing 1-2 carbon-carbon double bonds. Nonlimiting examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, and the like. The condensed rings may or may not be non-aromatic hydrocarbyl rings provided that the point of attachment is at a cycloalkyl carbon atom. For example, and without limitation, the following is a cycloalkyl group:




[0027] The term “halo” refers to F, Cl, Br, and/or I.

[0028] The term “heteroaryl” refers to a monovalent, aromatic mono-, bi-, or tricyclic ring having 2-16 ring carbon atoms and 1-8 ring heteroatoms selected preferably from N, O, S, and P and oxidized forms of N, S, and P, provided that the ring contains at least 5 ring atoms. Nonlimiting examples of heteroaryl include furan, imidazole, oxadiazole, oxazole, pyridine, quinoline, and the like. The condensed rings may or may not be a heteroatom containing aromatic ring provided that the point of attachment is a heteroaryl atom. For example, and without limitation, the following is a heteroaryl group:



[0029] The term “heterocyclyl” or heterocycle refers to a non-aromatic, mono-, bi-, or tricyclic ring containing 2-12 ring carbon atoms and 1-8 ring heteroatoms selected preferably from N, O, S, and P and oxidized forms of N, S, and P, provided that the ring contains at least 3 ring atoms. While heterocyclyl preferably refers to saturated ring systems, it also includes ring systems containing 1-3 double bonds, provided that the ring is non-aromatic. Nonlimiting examples of heterocyclyl include, azalactones, oxazoline, piperidinyl, piperazinyl, pyrrolidinyl, tetrahydrofuranyl, and tetrahydropyranyl. The condensed rings may or may not contain a non-aromatic heteroatom containing ring provided that the point of attachment is a heterocyclyl group. For example, and without limitation, the following is a heterocyclyl group:



[0030] The term “hydrolyzing” refers to breaking an $R^H-O-CO-$, $R^H-O-CS-$, or an R^H-O-SO_2- moiety to an R^H-OH , preferably by adding water across the broken bond. A hydrolyzing is performed using various methods well known to the skilled artisan, non limiting examples of which include acidic and basic hydrolysis.

[0031] The term “oxo” refers to a $C=O$ group, and to a substitution of 2 geminal hydrogen atoms with a $C=O$ group.

[0032] The term “optionally substituted,” unless defined otherwise, refers to a substituted or unsubstituted group. The group may be substituted with one or more substituents, such as e.g., 1, 2, 3, 4 or 5 substituents. Preferably, the substituents are selected from the group consisting of oxo, halo, $-CN$, NO_2 , $-N_2^+$, $-CO_2R^{100}$, $-OR^{100}$, $-SR^{100}$, $-SOR^{100}$, $-SO_2R^{100}$, $-NR^{101}R^{102}$, $-CONR^{101}R^{102}$, $-SO_2NR^{101}R^{102}$, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-CR^{100}=C(R^{100})_2$, $-CCR^{100}$, C_3-C_{10} cycloalkyl, C_3-C_{10} heterocyclyl, C_6-C_{12} aryl and C_2-C_{12} heteroaryl, wherein each R^{100} independently is hydrogen or C_1-C_8 alkyl; C_3-C_{12} cycloalkyl; C_3-C_{10} heterocyclyl; C_6-C_{12} aryl; or C_2-C_{12} heteroaryl; wherein each alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with 1-3 halo, 1-3 C_1-C_6 alkyl, 1-3 C_1-C_6 haloalkyl or 1-3 C_1-C_6 alkoxy groups. Preferably, the substituents are selected from the group consisting of chloro, fluoro, $-OCH_3$, methyl, ethyl, *iso*-propyl, cyclopropyl, vinyl, ethynyl, $-CO_2H$, $-CO_2CH_3$, $-OCF_3$, $-CF_3$ and $-OCHF_2$.

[0033] R^{101} and R^{102} independently is hydrogen; C_1-C_8 alkyl, optionally substituted with $-CO_2H$ or an ester thereof, C_1-C_6 alkoxy, oxo, $-CR^{103}=C(R^{103})_2$, $-CCR$, C_3-C_{10} cycloalkyl, C_3-C_{10} heterocyclyl, C_6-C_{12} aryl, or C_2-C_{12} heteroaryl, wherein each R^{103} independently is hydrogen or C_1-C_8 alkyl; C_3-C_{12} cycloalkyl; C_3-C_{10} heterocyclyl; C_6-C_{12} aryl; or C_2-C_{12} heteroaryl; wherein each cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with 1-3 alkyl groups or 1-3 halo groups, or R^{101} and R^{102} together with the nitrogen atom they are attached to form a 5-7 membered heterocycle.

[0034] The term “pharmaceutically acceptable” refers to safe and non-toxic for *in vivo*, preferably, human administration.

[0035] The term “pharmaceutically acceptable salt” refers to a salt that is pharmaceutically acceptable.

[0036] The term “salt” refers to an ionic compound formed between an acid and a base. When the compound provided herein contains an acidic functionality, such salts include, without limitation, alkali metal, alkaline earth metal, and ammonium salts. As used herein, ammonium salts include, salts containing protonated nitrogen bases and alkylated nitrogen bases. Exemplary, and non-limiting cations useful in pharmaceutically acceptable salts include Na, K, Rb, Cs, NH₄, Ca, Ba, imidazolium, and ammonium cations based on naturally occurring amino acids. When the compounds utilized herein contain basic functionality, such salts include, without limitation, salts of organic acids, such as carboxylic acids and sulfonic acids, and mineral acids, such as hydrogen halides, sulfuric acid, phosphoric acid, and the likes. Exemplary and non-limiting anions useful in pharmaceutically acceptable salts include oxalate, maleate, acetate, propionate, succinate, tartrate, chloride, sulfate, bisulfate, mono-, di-, and tribasic phosphate, mesylate, tosylate, and the likes.

[0037] The terms “treat”, “treating” or “treatment”, as used herein, include alleviating, abating or ameliorating a disease or condition or one or more symptoms thereof, preventing additional symptoms, ameliorating or preventing the underlying metabolic causes of symptoms, inhibiting the disease or condition, *e.g.*, arresting or suppressing the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or suppressing the symptoms of the disease or condition, and are intended to include prophylaxis. The terms also include relieving the disease or conditions, *e.g.*, causing the regression of clinical symptoms. The terms further include achieving a therapeutic benefit and/or a prophylactic benefit. By therapeutic benefit is meant eradication or amelioration of the underlying disorder being treated. Also, a therapeutic benefit is achieved with the eradication or amelioration of one or more of the physiological symptoms associated with the underlying disorder such that an improvement is observed in the individual, notwithstanding that the individual is still be afflicted with the underlying disorder. For prophylactic benefit, the compositions are administered to an individual at risk of developing a particular disease, or to an individual reporting one or more of the physiological symptoms of a disease, even though a diagnosis of this disease has not been made.

[0038] The terms “preventing” or “prevention” refer to a reduction in risk of acquiring a disease or disorder (*i.e.*, causing at least one of the clinical symptoms of the disease not to develop in a subject that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease). The terms further include causing the clinical

symptoms not to develop, for example in a subject at risk of suffering from such a disease or disorder, thereby substantially averting onset of the disease or disorder.


[0039] The term “effective amount” refers to an amount that is effective for the treatment of a condition or disorder by an intranasal administration of a compound or composition described herein. In some embodiments, an effective amount of any of the compositions or dosage forms described herein is the amount used to treat a disorder mediated by hemoglobin or a disorder that would benefit from tissue and/or cellular oxygenation of any of the compositions or dosage forms described herein to a subject in need thereof.

[0040] The term “carrier” as used herein, refers to relatively nontoxic chemical compounds or agents that facilitate the incorporation of a compound into cells, *e.g.*, red blood cells, or tissues.

[0041] As used herein, a “prodrug” is a compound that, after administration, is metabolized or otherwise converted to an active or more active form with respect to at least one property. To produce a prodrug, a pharmaceutically active compound can be modified chemically to render it less active or inactive, but the chemical modification is such that an active form of the compound is generated by metabolic or other biological processes. A prodrug may have, relative to the drug, altered metabolic stability or transport characteristics, fewer side effects or lower toxicity. For example, see the reference Nogrady, 1985, Medicinal Chemistry A Biochemical Approach, Oxford University Press, New York, pages 388-392. Prodrugs can also be prepared using compounds that are not drugs.

Compounds

[0042] In certain aspects of the invention, a compound of formula (I) is provided:



(I)


or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein L^{10} is optionally substituted methylene or, preferably, a bond;

ring A is C_6 - C_{10} aryl, a C_3 - C_8 cycloalkyl, a 5-10 membered heteroaryl or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, heteroaryl, cycloalkyl, or heterocycle is optionally substituted with 1-4: halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, and/or C_3 - C_{10} cycloalkyl, wherein the C_1 - C_6 alkyl is optionally substituted with 1-5 halo, C_1 - C_6 alkoxy, and/or C_3 - C_{10} cycloalkyl; or

ring A is C_6 - C_{10} aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, or heteroaryl is optionally substituted with 1-4 C_1 - C_6 alkyl and/or C_1 - C_6 alkoxy groups;

ring B is a 5-10 membered heteroaryl or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted with 1-4: halo, C_1 - C_6 alkyl and/or $-CO-C_1-C_6$ alkyl, or

ring B is:



wherein ring B' including the -N-CO- moiety is a 5-6 membered heterocycle containing up to 3 heteroatoms selected from nitrogen, oxygen, and sulfur and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted with 1-4 C₁-C₆ alkyl groups;

 is a single or a double bond;

each X and Y is independently (CR²⁰R²¹)_e, O, S, SO, SO₂, or NR²⁰; e is 1 to 4, preferably 1; each R²⁰ and R²¹ independently is hydrogen or C₁-C₃ alkyl optionally substituted with 1-3 halo, OH, or C₁-C₆ alkoxy, or CR²⁰R²¹ is C=O, provided that if one of X and Y is O, S, SO, SO₂, then the other is not CO, and X and Y are both not heteroatoms or oxidized forms thereof;

ring C is C₆-C₁₀ aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, -OR¹, C₁-C₆ alkyl, -COOR¹, and/or C₁-C₆ alkoxy, wherein the C₁-C₆ alkyl is optionally substituted with 1-5 halo, C₁-C₆ alkoxy, and/or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S; or

ring C is C₆-C₁₀ aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, -OR¹, C₁-C₆ alkyl, -COOR¹, NR⁵R⁶,


R¹ is a hydrogen, C₁-C₆ alkyl or a prodrug moiety; wherein the alkyl is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, which is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein

the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein the heteroaryl is optionally substituted with C₁-C₆ alkyl;

R⁵ and R⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl or -COOR³;

R³ is hydrogen or optionally substituted C₁-C₆ alkyl;

V¹ and V² independently are C₁-C₆ alkoxy; or V¹ and V² together with the carbon atom they are attached to form a ring of formula:



wherein each V³ and V⁴ are independently O, S, or NH, provided that when one of V³ and V⁴ is S, the other is NH, and provided that V³ and V⁴ are both not NH; q is 1 or 2; each V⁵ is independently C₁-C₆ alkyl optionally substituted with 1-3 OH groups, or V⁵ is CO₂R⁶⁰, where each R⁶⁰ independently is C₁-C₆ alkyl or hydrogen; t is 0, 1, 2, or 4; or CV¹V² is C=V, wherein V is O, NOR⁸⁰, or NNR⁸¹R⁸²;

R⁸⁰ is optionally substituted C₁-C₆ alkyl;

R⁸¹ and R⁸² independently are selected from the group consisting of hydrogen, optionally substituted C₁-C₆ alkyl, COR⁸³, or CO₂R⁸⁴;

R⁸³ is hydrogen or optionally substituted C₁-C₆ alkyl; and

R⁸⁴ is optionally substituted C₁-C₆ alkyl;

with the proviso that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;


then ring A excludes optionally substituted 5-10 membered heteroaryl;

and provided that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 5-10 membered heteroaryl;

then ring A is not optionally substituted 4-10 membered heterocycle.

[0043] In certain aspects of the invention, a compound of formula (X-I) is provided:



or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein ring A is C₆-C₁₀ aryl, a C₃-C₈ cycloalkyl, a 5-10 membered heteroaryl or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, heteroaryl, cycloalkyl, or heterocycle is optionally substituted with 1-4: halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, and/or C₃-C₁₀ cycloalkyl, wherein the C₁-C₆ alkyl is optionally substituted with 1-5 halo, C₁-C₆ alkoxy, and/or C₃-C₁₀ cycloalkyl;

ring B is a 5-10 membered heteroaryl or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted with 1-4: halo, C₁-C₆ alkyl and/or -CO-C₁-C₆ alkyl,

 is a single or a double bond;


X is O, S, SO, or SO₂;

ring C is C₆-C₁₀ aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, -ORⁱ, C₁-C₆ alkyl, -COORⁱ, and/or C₁-C₆ alkoxy, wherein the C₁-C₆ alkyl is optionally substituted with 1-5 halo, C₁-C₆ alkoxy and/or a 4-10 membered heterocycle containing up to 5 ring

heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S; and

R^1 is hydrogen or a prodrug moiety;

V^1 and V^2 independently are C_1 - C_6 alkoxy; or V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:



wherein each V^3 and V^4 are independently O, S, or NH, provided that when one of V^3 and V^4 is S, the other is NH, and provided that V^3 and V^4 are both not NH; q is 1 or 2; each V^5 is independently C_1 - C_6 alkyl or CO_2R^{60} , where each R^{60} independently is C_1 - C_6 alkyl or hydrogen; t is 0, 1, 2, or 4; or CV^1V^2 is $C=V$, wherein V is O, NOR^{80} , or $NNR^{81}R^{82}$;

R^{80} is optionally substituted C_1 - C_6 alkyl;

R^{81} and R^{82} independently are selected from the group consisting of hydrogen, optionally substituted C_1 - C_6 alkyl, COR^{83} , or CO_2R^{84} ;

R^{83} is hydrogen or optionally substituted C_1 - C_6 alkyl;

R^{84} is optionally substituted C_1 - C_6 alkyl;

provided that when ring C is C_6 - C_{10} aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;

then ring A excludes optionally substituted 5-10 membered heteroaryl;

and provided that when ring C is C_6 - C_{10} aryl;


and ring B is optionally substituted 5-10 membered heteroaryl;

then ring A is not optionally substituted 4-10 membered heterocycle.

[0044] In certain aspects of the invention, a compound of formula (X-I) is provided:


or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein

ring A is phenyl optionally substituted with 1-3 halo and/or C_1 - C_6 alkoxy, or is a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, optionally substituted, or is



wherein R^7 is C_1 - C_6 alkyl, optionally substituted with 3-5 fluoro groups, or is C_3 - C_6 cycloalkyl;

ring B is selected from the group consisting of



wherein R^8 is C_1 - C_6 alkyl, $-\text{CO}-C_1$ - C_6 alkyl or a prodrug moiety;


X is O, S, SO, or SO_2 ;

 is a single or a double bond;

ring C is phenyl or a 6 membered nitrogen-containing heteroaryl, each of which is optionally substituted with 1-4: halo, oxo, $-\text{OR}^1$, C_1 - C_6 alkyl, $-\text{COOR}^1$, and/or C_1 - C_6 alkoxy, wherein the C_1 - C_6 alkyl is optionally substituted with 1-5 halo, C_1 - C_6 alkoxy and/or 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S; and

each R^1 is hydrogen or a prodrug moiety R;


V^1 and V^2 independently are C_1 - C_6 alkoxy; or V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:




wherein each V^3 and V^4 are independently O, S, or NH, provided that when one of V^3 and V^4 is S, the other is NH, and provided that V^3 and V^4 are both not NH; q is 1 or 2; each V^5 is independently C_1 - C_6 alkyl or CO_2R^{60} , where each R^{60} independently is C_1 - C_6 alkyl or hydrogen; t is 0, 1, 2, or 4; or CV^1V^2 is $\text{C}=\text{V}$, wherein V is O, NOR^{80} , or $\text{NNR}^{81}\text{R}^{82}$;

R^{80} is optionally substituted C_1 - C_6 alkyl;
 R^{81} and R^{82} independently are selected from the group consisting of hydrogen,
 optionally substituted C_1 - C_6 alkyl, COR^{83} , or CO_2R^{84} ;
 R^{83} is hydrogen or optionally substituted C_1 - C_6 alkyl;
 R^{84} is optionally substituted C_1 - C_6 alkyl;
 provided that when ring C is C_6 - C_{10} aryl;
 and ring B is optionally substituted 4-10 membered heterocyclyl;
 then ring A excludes optionally substituted 5-10 membered heteroaryl;
 and provided that when ring C is C_6 - C_{10} aryl;
 and ring B is optionally substituted 5-10 membered heteroaryl;
 then ring A is not optionally substituted 4-10 membered heterocycle.

[0045] Preferably, V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:




[0046] In some embodiments, V^1 and V^2 independently are C_1 - C_6 alkoxy; or V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:




wherein each V^3 and V^4 are independently O, S, or NH, provided that when one or V^3 and V^4 is S the other is NH, and provided that V^3 and V^4 are both not NH; q is 1 or 2; each V^5 is independently C_1 - C_6 alkyl or CO_2R^{60} , where each R^{60} independently is C_1 - C_6 alkyl or hydrogen; t is 0, 1, 2, or 4; or CV^1V^2 is $C=V$, wherein V is O, and wherein the remaining variables are defined herein.

[0047] In certain embodiments, a compound of formula (III) is provided:




(III)

or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein ring A is phenyl optionally substituted with 1-3 halo and/or C₁-C₆ alkoxy, or is a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, optionally substituted, or is



wherein R⁷ is C₁-C₆ alkyl, optionally substituted with 3-5 fluoro groups, or is C₃-C₆ cycloalkyl;

ring B is selected from the group consisting of




wherein R⁸ is C₁-C₆ alkyl, -CO-C₁-C₆ alkyl or a prodrug moiety;

X is O, S, SO, or SO₂;

is a single or a double bond:

ring C is phenyl or a 6 membered nitrogen-containing heteroaryl, each of which is optionally substituted with 1-4: halo, oxo, -OR¹, C₁-C₆ alkyl -COOR¹, and/or C₁-C₆ alkoxy, wherein the C₁-C₆ alkyl is optionally substituted with 1-5 halo,

C₁-C₆ alkoxy and/or 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S; and
 each R¹ is hydrogen or a prodrug moiety R;
 V¹ and V² independently are C₁-C₆ alkoxy; or V¹ and V² together with the carbon atom they are attached to form a ring of formula:



wherein each V³ and V⁴ are independently O, S, or NH, provided that when one of V³ and V⁴ is S, the other is NH, and provided that V³ and V⁴ are both not NH; q is 1 or 2; each V⁵ is independently C₁-C₆ alkyl or CO₂R⁶⁰, where each R⁶⁰ independently is C₁-C₆ alkyl or hydrogen; t is 0, 1, 2, or 4; or CV¹V² is C=V, wherein V is O, NOR⁸⁰, or NNR⁸¹R⁸²;

R⁸⁰ is optionally substituted C₁-C₆ alkyl;

R⁸¹ and R⁸² independently are selected from the group consisting of hydrogen, optionally substituted C₁-C₆ alkyl, COR⁸³, or CO₂R⁸⁴;

R⁸³ is hydrogen or optionally substituted C₁-C₆ alkyl;

R⁸⁴ is optionally substituted C₁-C₆ alkyl;

provided that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;

then ring A excludes optionally substituted 5-10 membered heteroaryl;


and provided that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 5-10 membered heteroaryl;

then ring A is not optionally substituted 4-10 membered heterocycle.

[0048] In certain embodiments, t is 0. In certain embodiments, t is 1. In certain embodiments, t is 2. In certain embodiments, t is 3. In one embodiment, --X-Y- is --CH₂-O-. In another embodiment, -X-Y is -O-CH₂-.

[0049] In certain embodiments, a compound selected from formulas (IIIA), (IIIB) and (IIIC) is provided:



wherein

R^9 is hydrogen, $-OR^1$, C_1 - C_6 alkoxy optionally substituted with 1-3 C_1 - C_6 alkoxy or 4-10 membered heterocycle containing up to 5 ring heteroatoms selected from N, O, S or oxidized forms thereof;

R^{10} is hydrogen, halo, hydroxy or C_1 - C_6 alkoxy;

R^{11} is hydrogen or C_1 - C_6 alkyl; and

R^{12} is $-OR^1$;

wherein R^1 is hydrogen or the prodrug moiety R.


[0050] In certain embodiments, ring A is

phenyl substituted with 1-3 halo or C_1 - C_6 alkoxy, or









C_3 - C_8 heterocyclyl containing 1-3 heteroatoms, wherein the heterocycle is optionally substituted with 1-3 halo.

[0051] In certain embodiments, ring A is pyridyl, optionally substituted as defined herein.

[0052] In certain embodiments, compounds of formulas (II), (III), (IIIA), (IIIB) and (IIIC) are provided, wherein




is selected from the group consisting of:




and










[0053] In certain embodiments, compounds of formulas (II), (III), (IIIA), (IIIB) and (IIIC) are provided, wherein




















is



[0054] In certain embodiments, a compound is provided, wherein the compound selected from the group consisting of:









or an N oxides thereof, or a pharmaceutically acceptable salt of each thereof.


[0055] In certain embodiments, a compound is provided, wherein the compound selected from the group consisting of:







or an N oxides thereof, or a pharmaceutically acceptable salt of each thereof.

[0056] In certain aspects of the invention, a compound is provided, wherein the compound is selected from the group consisting of:








or a prodrug thereof, or a pharmaceutically acceptable salt of each thereof. Other compounds of this invention are illustrated in the Examples section.


[0057] In certain aspects of the invention, a compound of formula (I) is provided:



or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein L^{10} is optionally substituted methylene or, preferably, a bond;

ring A is C_6 - C_{10} aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, or heteroaryl is optionally substituted with 1-4 C_1 - C_6 alkyl and/or C_1 - C_6 alkoxy groups;

ring B is:



wherein ring B' including the -N-CO- moiety is a 5-6 membered heterocycle containing up to 3 heteroatoms selected from nitrogen, oxygen, and sulfur and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted with 1-4 C₁-C₆ alkyl groups;

each X and Y is independently (CR²⁰R²¹)_e, O, S, SO, SO₂, or NR²⁰; e is 1 to 4, preferably 1; each R²⁰ and R²¹ independently is hydrogen or C₁-C₃ alkyl optionally substituted with 1-3 halo, OH, or C₁-C₆ alkoxy, or CR²⁰R²¹ is C=O, provided that if one of X and Y is O, S, SO, SO₂, then the other is not CO, and X and Y are both not heteroatoms or oxidized forms thereof;


ring C is C₆-C₁₀ aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, -OR¹, C₁-C₆ alkyl, -COOR⁵, NR⁵R⁶,

R¹ is a hydrogen, C₁-C₆ alkyl or a prodrug moiety; wherein the alkyl is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, which is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein the heteroaryl is optionally substituted with C₁-C₆ alkyl;

R⁵ and R⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl or -COOR³;

R³ is hydrogen or optionally substituted C₁-C₆ alkyl;

V^1 and V^2 independently are C_1 - C_6 alkoxy; or V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:



wherein each V^3 and V^4 are independently O, S, or NH, provided that when one of V^3 and V^4 is S, the other is NH, and provided that V^3 and V^4 are both not NH; q is 1 or 2; each V^5 is independently C_1 - C_6 alkyl optionally substituted with 1-3 OH groups, or V^5 is CO_2R^{60} , where each R^{60} independently is C_1 - C_6 alkyl or hydrogen; t is 0, 1, 2, or 4; or CV^1V^2 is $C=V$, wherein V is O, NOR^{80} , or $NNR^{81}R^{82}$;

R^{80} is optionally substituted C_1 - C_6 alkyl;

R^{81} and R^{82} independently are selected from the group consisting of hydrogen, optionally substituted C_1 - C_6 alkyl, COR^{83} , or CO_2R^{84} ;

R^{83} is hydrogen or optionally substituted C_1 - C_6 alkyl; and

R^{84} is optionally substituted C_1 - C_6 alkyl;

with the proviso that when ring C is C_6 - C_{10} aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;


then ring A excludes optionally substituted 5-10 membered heteroaryl;

and provided that when ring C is C_6 - C_{10} aryl;

and ring B is optionally substituted 5-10 membered heteroaryl;


then ring A is not optionally substituted 4-10 membered heterocycle.

[0058] In certain aspects of the invention, a compound of formula (IV) is provided:



or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein ring A is C₆-C₁₀ aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl or heteroaryl is optionally substituted with 1-4 C₁-C₆ alkyl;

ring B is:



wherein ring B' including the -N-CO- moiety is a 5-6 membered heterocycle containing up to 3 heteroatoms selected from nitrogen, oxygen, and sulfur and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted with 1-4 C₁-C₆ alkyl groups;

each X and Y is independently CR²⁰R²¹, O, S, SO, SO₂, or NR¹⁰; each R²⁰ and R²¹ independently is hydrogen or C₁-C₃ alkyl optionally substituted with 1-3 halo, OH, or C₁-C₆ alkoxy, or CR²⁰R²¹ is C=O, provided that if one of X and Y is O, S, SO, SO₂, then the other is not CO, and X and Y are both not heteroatoms or oxidized forms thereof;


ring C is C₆-C₁₀ aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, -OR¹, C₁-C₆ alkyl, -COOR⁵, NR⁵R⁶,

R¹ is a hydrogen, C₁-C₆ alkyl or a prodrug moiety; wherein the alkyl is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, which is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein the heteroaryl is optionally substituted with C₁-C₆ alkyl;

R⁵ and R⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl or -COOR³;

R³ is hydrogen or optionally substituted C₁-C₆ alkyl;

V¹ and V² independently are C₁-C₆ alkoxy; or V¹ and V² together with the carbon atom they are attached to form a ring of formula:



wherein each V³ and V⁴ are independently O, S, or NH, provided that when one of V³ and V⁴ is S, the other is NH, and provided that V³ and V⁴ are both not NH; q is 1 or 2; each V⁵ is independently C₁-C₆ alkyl optionally substituted with 1-3 OH groups, or V⁵ is CO₂R⁶⁰, where each R⁶⁰ independently is C₁-C₆ alkyl or hydrogen; t is 0, 1, 2, or 4; or CV¹V² is C=V, wherein V is O, NOR⁸⁰, or NNR⁸¹R⁸²;

R⁸⁰ is optionally substituted C₁-C₆ alkyl;

R⁸¹ and R⁸² independently are selected from the group consisting of hydrogen, optionally substituted C₁-C₆ alkyl, COR⁸³, or CO₂R⁸⁴;

R⁸³ is hydrogen or optionally substituted C₁-C₆ alkyl; and

R⁸⁴ is optionally substituted C₁-C₆ alkyl.

[0059] In certain embodiments, ring C is substituted with at least one substituent selected from with 1-4: halo, $-OR^1$, C_1-C_6 alkyl, $-COOR^5$, NR^5R^6 .


[0060] In certain embodiments, X is CH_2 , O, S, SO, SO_2 or NH. In certain embodiments, X is O, S, SO or SO_2 . Preferably, X is O, and wherein the remaining variables are defined herein.

[0061] In certain embodiments, Y is $CR^{20}R^{21}$, O, S, SO, SO_2 , or NR^{10} ; wherein each R^{20} and R^{21} independently is hydrogen or C_1-C_3 alkyl. In certain embodiments, Y is $CR^{20}R^{21}$ wherein each R^{20} and R^{21} independently is hydrogen or C_1-C_3 alkyl. Preferably, Y is CH_2 , and wherein the remaining variables are defined herein.

[0062] In certain embodiments, t is 0. In certain embodiments, t is 1. In certain embodiments, t is 2. In certain embodiments, t is 3.


[0063] Preferably, CV^1V^2 is $C=V$, wherein V is O, and wherein the remaining variables are defined herein.

[0064] In certain embodiments, a compound of formula (V) is provided:



or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein ring A is C_6-C_{10} aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, or heteroaryl is optionally substituted with 1-4 C_1-C_6 alkyl and/or C_1-C_6 alkoxy groups;

ring B is:



wherein ring B' including the -N-CO- moiety is a 5-6 membered heterocycle containing up to 3 heteroatoms selected from nitrogen, oxygen, and sulfur and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted with 1-4 C₁-C₆ alkyl groups;

X is O, S, SO or SO₂;


ring C is C₆-C₁₀ aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, -OR¹, C₁-C₆ alkyl, -COOR⁵, NR⁵R⁶,

R¹ is a hydrogen, C₁-C₆ alkyl or a prodrug moiety R; wherein the alkyl is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, which is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein the heteroaryl is optionally substituted with C₁-C₆ alkyl;

R⁵ and R⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl or -COOR³; and

R³ is hydrogen or optionally substituted C₁-C₆ alkyl.

[0065] In certain embodiments, a compound of formula (VI) or (VII) is provided:



or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein ring A is C₆-C₁₀ aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, or heteroaryl is optionally substituted with 1-4 C₁-C₆ alkyl;

ring B is C₆-C₁₀ aryl, C₃-C₈ cycloalkyl, a 5-10 membered heteroaryl containing up to 5 ring heteroatoms or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, heteroaryl, cycloalkyl or heterocycle is optionally substituted with 1-4: halo, C₁-C₆ alkyl, or C₁-C₆ alkoxy, wherein the C₁-C₆ alkyl is optionally substituted with 1-5 halo, C₁-C₆ alkoxy, and/or C₃-C₁₀ cycloalkyl;

R⁴ is halo, oxo, -OR¹⁸, C₁-C₆ alkyl, C₁-C₆ alkoxy, -COOR⁵, and/or NR⁵R⁶;

R¹⁸ is hydrogen, substituted C₁-C₆ alkyl, or a prodrug moiety R;


R⁵ and R⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl or -COOR³; and

R³ is hydrogen, provided that the COOR³ is not joined to a nitrogen atom, or is optionally substituted C₁-C₆ alkyl.


[0066] In one embodiment, R⁴ is -OH. In another embodiment, R⁴ is NH₂. In one embodiment, R⁴ is NH(CH₃). In one embodiment, R⁴ is N(CH₃)₂. In one embodiment, R⁴ is NHC(O)OC(CH₃)₃. In one embodiment, R⁴ is COOH. In one embodiment, R⁴ is optionally

substituted dioxolan. In one embodiment, R^4 is a substituted pyridine. As used herein, R^3 is hydrogen, provided that the $COOR^3$ is not joined to a nitrogen atom.


[0067] In certain embodiments, ring B is selected from the group consisting of:



[0068] In certain embodiments, compounds of formula (V) are provided, wherein



is




[0069] In certain embodiments, a compound of formula (IV) as disclosed above is provided, with the proviso that:


when ring C is C_6 - C_{10} aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;

then ring A excludes optionally substituted 5-10 membered heteroaryl.

[0070] In certain embodiments, a compound is provided, wherein the compound is selected from the group consisting of:





or N oxides thereof, or a pharmaceutically acceptable salt of each thereof.

[0071] Compounds provided herein include those in the Examples section.

Prodrug Moiety

[0072] In one aspect, **R** is hydrogen, a phosphate or a diphosphate containing moiety, or another promoiety or prodrug moiety. Preferably the prodrug moiety imparts at least a 2 fold, more preferably a 4 fold, enhanced solubility and/or bioavailability to the active moiety (where **R** is hydrogen), and more preferably is hydrolyzed in vivo. The promoieties are structurally and functionally defined herein.

[0073] In one embodiment, **R** is $-\text{COR}^{90}$, CO_2R^{91} , or $\text{CONR}^{92}\text{R}^{93}$ wherein R^{90} and R^{91} independently are $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, 4-9 membered heterocycle, or a 5-10 membered heteroaryl, each containing at least 1 basic nitrogen moiety; and R^{92} and R^{93} independently are $\text{C}_1\text{-C}_6$ alkyl; $\text{C}_3\text{-C}_8$ cycloalkyl, 4-9 membered heterocycle, or a 5-10 membered heteroaryl, each containing at least 1 basic nitrogen moiety; or R^{92} and R^{93} together with the nitrogen atom they are bonded to for a 4-9 member heterocycle substituted with at least 1 amino, $\text{C}_1\text{-C}_6$ alkyl amino, or di $\text{C}_1\text{-C}_6$ alkylamino group.

[0074] In certain embodiments, **R** is $-\text{C}(\text{O})\text{R}^{31}$, $\text{C}(\text{O})\text{OR}^{31}$, or $\text{CON}(\text{R}^{13})_2$,
 each R^{31} is independently a $\text{C}_1\text{-C}_6$ alkyl; $\text{C}_3\text{-C}_8$ cycloalkyl, 4-9 membered heterocycle, or a 5-10 membered heteroaryl, containing at least 1 basic nitrogen moiety; and
 each R^{13} independently is $\text{C}_1\text{-C}_6$ alkyl; $\text{C}_3\text{-C}_8$ cycloalkyl, 4-9 membered heterocycle, or a 5-10 membered heteroaryl, containing at least 1 basic nitrogen moiety; or both R^{13}

together with the nitrogen atom they are bonded to for a 4-9 member heterocycle substituted with at least 1 amino, C₁-C₆ alkyl amino, or di C₁-C₆ alkylamino group.

[0075] In one aspect, **R** is C(O)OR³¹, C(S)OR³¹, C(O)SR³¹ or COR³¹, wherein R³¹ is as defined herein.

[0076] In one embodiment, R³¹ is a group of the formula (CR³²R³³)_eNR³⁴R³⁵, wherein each R³² and R³³ is independently H, a C₁-C₈ alkyl, C₃-C₉ heterocyclyl, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, C₃-C₉ heteroaryl or R³² and R³³ together with the carbon atom they are bond to form a C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, C₃-C₉ heterocyclyl or C₃-C₉ heteroaryl ring system, or 2 adjacent R³² moieties or 2 adjacent R³³ moieties together with the carbon atom they are bond to form a C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, C₃-C₉ heterocyclyl or C₃-C₉ heteroaryl ring system;

each R³⁴ and R³⁵ is a C₁-C₈ alkyl, C₃-C₉ heterocyclyl, C₃-C₈ cycloalkyl, or R³⁴ and R³⁵ together with the nitrogen atom they are bond to form a C₃-C₈ cycloalkyl or C₃-C₉ heterocyclyl ring system;

each heterocyclic and heteroaryl ring system is optionally substituted with C₁-C₃ alkyl, -OH, amino and carboxyl groups; and


e is an integer of from 1 to 4.

[0077] In some less preferred embodiments R³⁴ and R³⁵ can be hydrogen.

[0078] In one embodiment, the subscript e is preferably 2 and each R³² and R³³ is preferably independently selected from the group, H, CH₃, and a member in which R³² and R³³ are joined together to form a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, or 1,1-dioxo-hexahydro-1Δ⁶-thiopyran-4-yl or tetrahydropyran-4-yl group.

[0079] With regard to the prodrug group, preferred embodiments are compounds wherein NR³⁴R³⁵ is morpholino.

[0080] In one embodiment, **R** is:



wherein


each R^{32} and R^{33} is independently H, C_1 - C_8 alkyl, or optionally, if both present on the same substituent, may be joined together to form a C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, C_3 - C_9 heterocyclyl or C_3 - C_9 heteroaryl ring system.


[0081] Within this embodiment, each R^{32} and R^{33} is independently, H, CH_3 , or are joined together to form a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 1,1-dioxo-hexahydro-1 λ^6 -thiopyran-4-yl or tetrahydropyran-4-yl group.

[0082] In a preferred embodiment, linkage of the prodrug moiety to the rest of the active molecule is stable enough so that the serum half life of the prodrug is from about 8 to about 24 hours.








[0083] In an embodiment of the invention, the prodrug moiety comprises a tertiary amine having a pK_a near the physiological pH of 7.5. Any amines having a pK_a within 1 unit of 7.5 are suitable alternatives amines for this purpose. The amine may be provided by the amine of a morpholino group. This pK_a range of 6.5 to 8.5 allows for significant concentrations of the basic neutral amine to be present in the mildly alkaline small intestine. The basic, neutral form of the amine prodrug is lipophilic and is absorbed through the wall of the small intestine into the blood. Following absorption into the bloodstream, the prodrug moiety is cleaved by esterases which are naturally present in the serum to release an active compound.

[0084] Examples of R include, without limitation:






[0001] In another embodiment, R is as tabulated below:

R	m	R ³⁴	R ³⁵	NR ³⁴ R ³⁵
C(O)(CH ₂) _m NR ³⁴ R ³⁵	1	Me	Me	
C(O)(CH ₂) _m NR ³⁴ R ³⁵	2	Me	Me	
C(O)(CH ₂) _m NR ³⁴ R ³⁵	3	Me	Me	
C(O)(CH ₂) _m NR ³⁴ R ³⁵	4	Me	Me	
C(O)(CH ₂) _m NR ³⁴ R ³⁵	1			
C(O)(CH ₂) _m NR ³⁴ R ³⁵	2			
C(O)(CH ₂) _m NR ³⁴ R ³⁵	3			
C(O)(CH ₂) _m NR ³⁴ R ³⁵	4			
C(O)O(CH ₂) _m NR ³⁴ R ³⁵	2	Me	Me	
C(O)O(CH ₂) _m NR ³⁴ R ³⁵	3	Me	Me	
C(O)O(CH ₂) _m NR ³⁴ R ³⁵	4	Me	Me	
C(O)O(CH ₂) _m NR ³⁴ R ³⁵	2			
C(O)O(CH ₂) _m NR ³⁴ R ³⁵	3			
C(O)O(CH ₂) _m NR ³⁴ R ³⁵	4			
P(O)(OH) ₂				

an N oxide thereof, or a pharmaceutically acceptable salt of each thereof.


[0085] In another aspect, R is,



wherein

R³⁶ is lower alkyl (e.g. C₁-C₆ alkyl).

[0086] In yet another aspect, **R** is:



wherein X¹, Y¹ and X² are as defined herein.

[0087] In one embodiment, X¹ is selected from the group consisting of O, S and NR³⁷

wherein R³⁷ is hydrogen or C₁-C₆ alkyl;

Y^1 is $-C(R^{38})_2$ or a sugar moiety, wherein each R^{38} is independently hydrogen or C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl;

X^2 is selected from the group consisting of halogen, C_1 - C_6 alkoxy, diacylglycerol, amino, C_1 - C_6 alkylamino, C_1 - C_6 dialkylamino, C_1 - C_6 alkylthio, a PEG moiety, a bile acid moiety, a sugar moiety, an amino acid moiety, a di- or tri-peptide, a PEG carboxylic acid, and $-U-V$ wherein

U is O or S; and

V is selected from the group consisting of C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, C_3 - C_9 heteroaryl, $C(W^2)X^3$, $PO(X^3)_2$, and SO_2X^3 ;

wherein W^2 is O or NR^{39}

wherein R^{39} is hydrogen or C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl; and

each X^3 is independently amino, hydroxyl, mercapto, C_1 - C_6 alkyl, heteroalkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl, C_1 - C_6 alkoxy, C_1 - C_6 alkylamino, C_1 - C_6 dialkylamino, C_1 - C_6 alkylthio, a bile acid based alkoxy group, a sugar moiety, a PEG moiety, and $-O-CH_2-CH(OR^{40})CH_2X^4R^{40}$,

wherein:


X^4 is selected from the group consisting of O, S, $S=O$, and SO_2 ; and

each R^{40} is independently C_{10} - C_{22} alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl, C_1 - C_8 alkylene, or C_1 - C_8 heteroalkylene.

[0088] Each heterocyclic and heteroaryl ring system is optionally substituted with C_1 - C_3 alkyl, $-OH$, amino and carboxyl groups.


[0089] In one embodiment, the present invention utilizes the following Y^1 groups: CH_2 , $CHMe$, $CH(isopropyl)$, $CH(tertiarybutyl)$, $C(Me)_2$, $C(Et)_2$, $C(isopropyl)_2$, and $C(propyl)_2$.

[0090] In another embodiment, the present invention utilizes the following X^2 groups:



-OMe, -OEt, -O-isopropyl, O-isobutyl, O-tertiarybutyl, -O-COMe, -O-C(=O)(isopropyl), -O-C(=O)(isobutyl), -O-C(=O)(tertiarybutyl), -O-C(=O)-NMe₂, -O-C(=O)-NHMe, -O-C(=O)-NH₂, -O-C(=O)-N(H)-CH(R⁴¹)-CO₂Et wherein R⁴¹ is a side chain C₁-C₆ alkyl, or C₃-C₉ heterocyclyl group selected from the side chain groups present in essential amino acids; -O-P(=O)(OMe)₂, -O-P(=O)(O-isopropyl)₂, and -O-P(=O)(O-isobutyl)₂. Each heterocyclic is optionally substituted with one or more, preferably, 1-3, C₁-C₃ alkyl, -OH, amino and/or carboxyl groups.

[0091] In another embodiment, In one embodiment, R is:





wherein

X³ is independently C₁-C₆ alkyl, C₃-C₈ cycloalkyl, C₃-C₉ heterocyclyl, C₆-C₁₀ aryl, or C₃-C₉ heteroaryl; and

R⁴² is independently hydrogen or C₁-C₆ alkyl, C₃-C₈ cycloalkyl, C₃-C₉ heterocyclyl, C₆-C₁₀ aryl, or C₃-C₉ heteroaryl.

[0092] Each heterocyclic is optionally substituted with one or more, preferably, 1-3, C₁-C₃ alkyl, -OH, amino and/or carboxyl groups.

[0093] In one embodiment, R is:



wherein

each X^3 is independently amino, hydroxyl, mercapto, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl, C_1 - C_6 alkoxy, C_1 - C_6 alkylamino, C_1 - C_6 dialkylamino, C_1 - C_6 alkylthio, a bile acid based alkoxy group, a sugar moiety, a PEG moiety, and $-O-CH_2-CH(OR^{40})CH_2X^4R^{40}$,

wherein:


X^4 is selected from the group consisting of O, S, S=O, and SO_2 ; and


each R^{40} is independently C_{10} - C_{22} alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, C_3 - C_9 heteroaryl, C_1 - C_8 alkylene, or C_1 - C_8 heteroalkylene; and

R^{42} is independently hydrogen or C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl.


[0094] In some embodiments, R^{42} is independently hydrogen or C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl; and each X^3 independently is C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl, C_1 - C_6 alkoxy, C_1 - C_6 alkylamino, C_1 - C_6 dialkylamino, or C_1 - C_6 alkylthio.

[0095] In some embodiments, **R** is represented by the following structures:






wherein, in the above examples, R^{43} is $\text{C}_{10}\text{-C}_{22}$ alkyl or alkylene, R^{44} is H or $\text{C}_1\text{-C}_6$ alkyl and R^{45} represents side chain alkyl groups present in naturally occurring alpha amino acids;




wherein R^{46} is $(CH_2)_n$, $n=2-4$, and $CO-R^{47}-NH_2$ represents an aminoacyl group; or



wherein R^{46} is $(CH_2)_n$, $n=2-4$, R^{47} is $(CH_2)_n$, $n=1-3$ and R^{49} is O or NMe.


[0096] In one embodiment, **R** is:




[0097] In one aspect, **R** is $-C(R^{200}R^{201})O(R^{202}R^{203})P(O)OR^{204}NR^{205}R^{206}$, wherein each R^{200} , R^{201} , R^{202} , R^{203} , R^{204} , R^{205} and R^{206} is independently H, a C_1 - C_8 alkyl, C_3 - C_9 heterocyclyl, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, C_3 - C_9 heteroaryl, wherein each alkyl, heterocyclyl, cycloalkyl, aryl, and heteroaryl is optionally substituted.

[0098] In some embodiments, **R** is $-CH(R^{201})OCH_2P(O)OR^{204}NHR^{206}$, wherein R^{201} is C_1 - C_8 alkyl, R^{204} is phenyl, optionally substituted. In one embodiment, R^{206} is $-CHR^{207}C(O)OR^{208}$ wherein R^{207} is selected from the group consisting of the naturally occurring amino acid side chains and $-CO_2H$ esters thereof and R^{208} is C_1 - C_8 alkyl. In one embodiment, R^{206} is C_1 - C_6 alkyl, optionally substituted with 1-3, CO_2H , SH , NH_2 , C_6 - C_{10} aryl, and C_2 - C_{10} heteroaryl.

[0099] In one embodiment, **R** is:




[0100] In one embodiment, **R** is:



wherein Y^1 is $-C(R^{38})_2$, wherein each R^{38} is independently hydrogen or C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl.

[0101] Various polyethylene glycol (PEG) moieties and synthetic methods related to them that can be used or adapted to make compounds of the invention are described in U.S. Patent Nos. 6,608,076; 6,395,266; 6,194,580; 6,153,655; 6,127,355; 6,111,107; 5,965,566; 5,880,131; 5,840,900; 6,011,042 and 5,681,567.

[0102] In one embodiment, **R** is



wherein


R^{50} is $-OH$ or hydrogen;

R^{51} is $-OH$, or hydrogen;

W is- $\text{CH}(\text{CH}_3)\text{W}^1$;
 wherein W^1 is a substituted C_1 - C_8 alkyl group containing a moiety which is optionally negatively charged at physiological pH,
 said moiety is selected from the group consisting of CO_2H , SO_3H , SO_2H , $-\text{P}(\text{O})(\text{OR}^{52})(\text{OH})$, $-\text{OP}(\text{O})(\text{OR}^{52})(\text{OH})$, and OSO_3H ,
 wherein R^{52} is C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl.

[0103] Each heterocyclic and heteroaryl ring system is optionally substituted with one or more, preferably 1-3, C_1 - C_3 alkyl, $-\text{OH}$, amino and/or carboxyl groups.

[0104] In one embodiment, **R** is:



wherein R^{53} is H or C_1 - C_6 alkyl.

[0105] In another aspect, **R** is SO_3H .

[0106] In another aspect, **R** comprises a cleavable linker, wherein the term “cleavable linker” refers to a linker which has a short half life in vivo. The breakdown of the linker **Z** in a compound releases or generates the active compound. In one embodiment, the cleavable linker has a half life of less than ten hours. In one embodiment, the cleavable linker has a half life of less than an hour. In one embodiment, the half life of the cleavable linker is between one and fifteen minutes. In one embodiment, the cleavable linker has at least one connection with the structure: $C^*-C(=X^*)X^*-C^*$ wherein C^* is a substituted or unsubstituted methylene group, and X^* is S or O. In one embodiment, the cleavable linker has at least one $C^*-C(=O)O-C^*$ connection. In one embodiment, the cleavable linker has at least one $C^*-C(=O)S-C^*$ connection. In one embodiment, the cleavable linker has at least one $-C(=O)N^*-C^*-SO_2-N^*$ -connection, wherein N^* is $-NH-$ or C_1-C_6 alkylamino. In one embodiment, the cleavable linker is hydrolyzed by an esterase enzyme.

[0107] In one embodiment, the linker is a self-immolating linker, such as that disclosed in U.S. patent publication 2002/0147138, to Firestone; PCT Appl. No. US05/08161 and PCT Pub. No. 2004/087075. In another embodiment, the linker is a substrate for enzymes. See generally Rooseboom et al., 2004, Pharmacol. Rev. 56:53-102.

Pharmaceutical Compositions

[0108] In further aspects of the invention, a composition is provided comprising any of the compounds described herein, and at least a pharmaceutically acceptable excipient.

[0109] In another aspect, this invention provides a composition comprising any of the compounds described herein, and a pharmaceutically acceptable excipient.

[0110] Such compositions can be formulated for different routes of administration. Although compositions suitable for oral delivery will probably be used most frequently, other routes that may be used include transdermal, intravenous, intraarterial, pulmonary, rectal, nasal, vaginal, lingual, intramuscular, intraperitoneal, intracutaneous, intracranial, and subcutaneous routes. Suitable dosage forms for administering any of the compounds described herein include tablets, capsules, pills, powders, aerosols, suppositories, parenterals, and oral liquids, including suspensions, solutions and emulsions. Sustained release dosage forms may also be used, for example, in a transdermal patch form. All dosage forms may be prepared using methods that are standard in the art (see e.g., Remington's Pharmaceutical Sciences, 16th ed., A. Oslo editor, Easton Pa. 1980).

[0111] Pharmaceutically acceptable excipients are non-toxic, aid administration, and do not adversely affect the therapeutic benefit of the compound of this invention. Such excipients may be any solid, liquid, semi-solid or, in the case of an aerosol composition, gaseous excipient that is generally available to one of skill in the art. Pharmaceutical compositions in accordance with the invention are prepared by conventional means using methods known in the art.

[0112] The compositions disclosed herein may be used in conjunction with any of the vehicles and excipients commonly employed in pharmaceutical preparations, *e.g.*, talc, gum arabic, lactose, starch, magnesium stearate, cocoa butter, aqueous or non-aqueous solvents, oils, paraffin derivatives, glycols, etc. Coloring and flavoring agents may also be added to preparations, particularly to those for oral administration. Solutions can be prepared using water or physiologically compatible organic solvents such as ethanol, 1,2-propylene glycol, polyglycols, dimethylsulfoxide, fatty alcohols, triglycerides, partial esters of glycerin and the like.

[0113] Solid pharmaceutical excipients include starch, cellulose, hydroxypropyl cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk and the like. Liquid and semisolid excipients may be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, *e.g.*, peanut oil, soybean oil, mineral oil, sesame oil, etc. In certain embodiments, the compositions provided herein comprises one or more of α -tocopherol, gum arabic, and/or hydroxypropyl cellulose.

[0114] In one embodiment, this invention provides sustained release formulations such as drug depots or patches comprising an effective amount of a compound provided herein. In another embodiment, the patch further comprises gum Arabic or hydroxypropyl cellulose separately or in combination, in the presence of alpha-tocopherol. Preferably, the hydroxypropyl cellulose has an average MW of from 10,000 to 100,000. In a more preferred embodiment, the hydroxypropyl cellulose has an average MW of from 5,000 to 50,000.

[0115] Compounds and pharmaceutical compositions of this invention may be used alone or in combination with other compounds. When administered with another agent, the co-administration can be in any manner in which the pharmacological effects of both are

manifest in the patient at the same time. Thus, co-administration does not require that a single pharmaceutical composition, the same dosage form, or even the same route of administration be used for administration of both the compound of this invention and the other agent or that the two agents be administered at precisely the same time. However, co-administration will be accomplished most conveniently by the same dosage form and the same route of administration, at substantially the same time. Obviously, such administration most advantageously proceeds by delivering both active ingredients simultaneously in a novel pharmaceutical composition in accordance with the present invention.

Methods of Treatment

[0116] In aspects of the invention, a method is provided for increasing tissue and/or cellular oxygenation, the method comprising administering to a subject in need thereof a therapeutically effective amount of any of the compounds or compositions described herein.

[0117] In aspects of the invention, a method is provided for increasing oxygen affinity of hemoglobin S in a subject, the method comprising administering to a subject in need thereof a therapeutically effective amount of any of the compounds or compositions described herein.

[0118] In aspects of the invention, a method is provided for treating a condition associated with oxygen deficiency, the method comprising administering to a subject in need thereof a therapeutically effective amount of any of the compounds or compositions described herein.

[0119] In further aspects of the invention, a method is provided for treating oxygen deficiency associated with sickle cell anemia, the method comprising administering to a subject in need thereof a therapeutically effective amount of any of the compounds or compositions described herein.

[0120] In further aspects of the invention, a method is provided for treating sickle cell disease, the method comprising administering to a subject in need thereof a therapeutically effective amount of a compound of any of the compounds or compositions described herein. In still further aspects of the invention, a method is provided for treating cancer, a pulmonary disorder, stroke, high altitude sickness, an ulcer, a pressure sore, Alzheimer's disease, acute respiratory disease syndrome, and a wound, the method comprising administering to a subject

in need thereof a therapeutically effective amount of a compound of any of the compounds or compositions described herein.

Synthetic Methods

[0121] Certain methods for making the compounds described herein are also provided. The reactions are preferably carried out in a suitable inert solvent that will be apparent to the skilled artisan upon reading this disclosure, for a sufficient period of time to ensure substantial completion of the reaction as observed by thin layer chromatography, $^1\text{H-NMR}$, etc. If needed to speed up the reaction, the reaction mixture can be heated, as is well known to the skilled artisan. The final and the intermediate compounds are purified, if necessary, by various art known methods such as crystallization, precipitation, column chromatography, and the likes, as will be apparent to the skilled artisan upon reading this disclosure.

[0122] An illustrative and non-limiting method for synthesizing a compound of formula (I), is schematically shown below.

In the following Schemes, ,  and  refer to rings A, B and C as described herein;

A^5 and B^5 are independently NR^{14} , O, S, S(O)_x , NBOC , CH_2 , CHR^{14} , $\text{C(R}^{14})_2$ provided that when both A^5 and B^5 are present in a ring, both are not CH_2 , CHR^{14} , $\text{C(R}^{14})_2$, and that when only 1 A^5 or B^5 is present in a ring the A^5 or B^5 is not CH_2 , CHR^{14} , $\text{C(R}^{14})_2$;

R^{14} is $\text{C}_1\text{-C}_6$ alkyl, COR^{15} or COOR^{15} ; wherein R^{15} is optionally substituted $\text{C}_1\text{-C}_6$ alkyl, optionally substituted $\text{C}_6\text{-C}_{10}$ aryl, optionally substituted 5-10 membered heteroaryl containing up to 5 ring heteroatoms, or optionally substituted 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S;

X, and X^5 each represents a leaving group and are independently selected from Cl, Br, and I.

X^6 represents CR, N, O, S(O)_x ; wherein x is 0, 1, or 2;

R^{71} is $\text{C}_1\text{-C}_6$ alkyl;

Y^5 represents a leaving group selected from Cl, F, Br, I, $\text{OSO}_2\text{R}^{17}$ and OSO_2Ar ;


Ar is phenyl optionally substituted with 1-3 halo and/or C₁-C₄ alkyl;

n is 0, 1, or 2; and

Where variables already used in the structures hereinabove are used in the schemes, the context makes it unambiguous as to what the variable refers to.

General Synthetic Schemes

Scheme 1




[0123] General method A (Scheme 1) for preparing aryloxy/heteroarylether analogs (4a/4b) from substituted methylene alcohol (1) and hydroxyl (hetero)aryl aldehyde derivatives (3a/3b). A hydroxyl (hetero)arylaldehyde derivatives (3a/3b) (0.1-2 mmol) mixture with substituted methylene alcohol (1) (0.8 to 1.2eq) and PPh₃ (1-1.5eq) in anhydrous THF (1-10mL) was stirred under nitrogen until complete dissolution. The solution was cooled to 0 °C on ice bath and DIAD or DEAD (1.1 eq) in THF or toluene was added dropwise over a 1-20 min period. The ice cooling bath was allowed to expire over 90 min and the mixture was stirred at RT for 2-48 hours. The mixture was stirred for 10 min, then filtered through a pad of silica. The silica was washed with ethyl acetate 2-20mL. The combined filtrates were evaporated and the residue was dried on highvac. The residue was purified by preparative HPLC or flash silica gel chromatography.

[0124] General method A (Scheme 1) for preparing aryloxy/heteroarylether analogs (4a/4b) from substituted methylene halide (2) and hydroxyl (hetero)aryl aldehyde derivatives (3a/3b). A mixture of hydroxyl (hetero)arylaldehyde derivatives (**3a/3b**) (0.1-2 mmol, 1-4 eq.), substituted methylene chloride or bromide (**2**) (1eq), and K_2CO_3 (2-5 eq.) (catalytic amount of NaI or Bu_4NI may also be added) in DMF or acetonitrile (1 to 10 mL) was stirred at RT or heating up to 120 °C for 0.5-8 h under nitrogen atmosphere. In workup A, water was added to the reaction mixture, the precipitated product was collected, washed with water, and then subjected to preparative HPLC or flash silica gel chromatography purification. In workup B (for products that did not precipitate), diluted HCl or aqueous NH_4Cl was added at 0 °C to adjusted the pH to ~7, the reaction mixture was partitioned between ethyl acetate or dichloromethane and aqueous sodium chloride and the organic layer separated, dried, and solvent removed under vacuum to afford crude product which was purified by automated silica gel column chromatography using appropriate solvents mixture (e.g., ethyl acetate/hexanes).

[0125] General method C for preparing substituted methylene chloride (2a). To a solution of substituted methylene alcohol (**1**) (0.1 to 2 mmol) in DCM (1-10 mL) was added $SOCl_2$ dropwise (2eq to 5eq) at 0 °C or RT. The reaction mixture was stirred at RT for 10min to 6 h, or until reaction is judged complete (LC/MS). The reaction mixture is concentrated to dryness over a rotavap. The crude chloride residue was suspended in toluene, sonicated and concentrated to dryness. The process was repeated three times and dried under vacuum to give the substituted methylene chloride (**2**), usually as an off-white solid, which was used for next step without further purification. Alternatively, a solution of aqueous 1N Na_2CO_3 is then added to produce a solution of pH~ 8. the mixture was extracted with DCM (3 x10-50mL), dried over sodium sulfate, and concentrated to the crude substituted methylene chloride (**2a**), which is then purified by column chromatography on silica gel (0-100% ethyl acetate-hexanes).

[0126] General method D for preparing substituted methylene bromide (2b). To a solution of substituted methylene alcohol (**1**) (0.1 to 2 mmol) in DCM (1-10 mL) was added $Ph_3P Br_2$ dropwise (2eq to 5eq) at 0 °C or RT. The reaction mixture was stirred at RT for 10 min to 2 h, or until reaction is judged complete (LC/MS). The reaction mixture is concentrated to dryness over a rotavap. The residue purified by column chromatography on silica gel (0-100% ethyl acetate-hexanes) to afford the pure bromide **2b**.




[0127] **General method E (Scheme 2) for preparing heterocyclic methylene derivatives 9, 10, 12 and 13.** Condensation of heterocyclic ketone analog **5** with chloroformate or dialkyl carbonate gives (hetero)cyclic beta-ketone ester **6** (Step 1). The ketone ester **6** is converted to the triflate intermediate **7** by treating with a triflating agent (e.g., triflic anhydride) in the presence of an organic base such as Hunig's base (Step 2). Suzuki coupling of the triflate **7** with a boronic acid or ester affords heterocyclohexene carboxylate **8** (Step 3). Subsequent reduction of the ester group by LAH or DIBAL gives the corresponding alcohol **9-OH** (Step 4). Further reaction of the alcohol **9-OH** with thionyl chloride, Ph_3PBr_2 (or $\text{CBr}_4\text{-Ph}_3\text{P}$ or PBr_3), or alkyl/aryl sulfonyl chloride produces the corresponding **10-X** chloride, bromide or sulfonate (Step 5).

[0128] Alternatively, the double bond of heterocyclohexene carboxylate **8** is reduced to give the *cis*-heterocyclohexane **11-cis** carboxylate under palladium catalyzed hydrogenation conditions (Step 6). Reduction of the ester group of **11-cis** by LAH or DIBAL yields *cis*-alcohol **12-OH-cis** (Step 8). Conversion of the alcohol **12-OH-cis** to its chloride, bromide or


sulfonate (such as mesylate, tosylate) **13-X-cis** can be achieved by reacting with thionyl chloride, or Ph_3PBr_2 , or sulfonyl chloride (such as mesyl chloride or tosyl chloride) (Step 9). The *cis*-cyclohexane carboxylate **11-cis** can also be isomerized to the thermodynamically more stable *trans*-isomer **11-trans** by the treatment with an alcoholic alkoxide (e.g., ethoxide) solution. Analogously, transformation of **11-trans** ester to **12-trans** alcohol and **13-X-trans** halide is accomplished by applying conditions of Step 8 and Step 9 (Scheme 2) similar to these for the corresponding *cis*-isomers.

Scheme 3



[0129] Coupling of the (hetero)cyclic methylene derivatives **9**, **10**, **12** and **13** with hydroxyl (hetero)arylaldehyde derivatives (**3a/3b**) (Scheme 3) by general method A or B affords the corresponding aryloxy/heteroarylether analogs (**4c** and **4d**).

[0130] Similarly, N-linked heterocyclic analogs (compound **5**, Scheme 4) can also be synthesized from amination procedures developed by Buchwald and Hartwig.



[0131] Syntheses of the ester prodrugs start with the free carboxylic acid bearing the tertiary amine. The free acid is activated for ester formation in an aprotic solvent and then reacted with a free alcohol group in the presence of an inert base, such as triethyl amine, to provide the ester prodrug. Activating conditions for the carboxylic acid include forming the acid chloride using oxalyl chloride or thionyl chloride in an aprotic solvent, optionally with a catalytic amount of dimethyl formamide, followed by evaporation. Examples of aprotic solvents, include, but are not limited to methylene chloride, tetrahydrofuran, and the like. Alternatively, activations can be performed in situ by using reagents such as BOP (benzotriazol-1-yloxytris(dimethylamino) phosphonium hexafluorophosphate, and the like (see Nagy et al., 1993, Proc. Natl. Acad. Sci. USA 90:6373-6376) followed by reaction with the free alcohol. Isolation of the ester products can be affected by extraction with an organic solvent, such as ethyl acetate or methylene chloride, against a mildly acidic aqueous solution; followed by base treatment of the acidic aqueous phase so as to render it basic; followed by extraction with an organic solvent, for example ethyl acetate or methylene chloride; evaporation of the organic solvent layer; and recrystallization from a solvent, such as ethanol. Optionally, the solvent can be acidified with an acid, such as HCl or acetic acid to provide a pharmaceutically acceptable salt thereof. Alternatively the crude reaction can be passed over an ion exchange column bearing sulfonic acid groups in the protonated form, washed with deionized water, and eluted with aqueous ammonia; followed by evaporation.


[0132] Suitable free acids bearing the tertiary amine are commercially available, such as 2-(N-morpholino)-propionic acid, N,N- dimethyl-beta-alanine, and the like. Non-commercial acids can be synthesized in straightforward manner via standard literature procedures.

[0133] Carbonate and carbamate prodrugs can be prepared in an analogous way. For example, amino alcohols and diamines can be activated using activating agents such as phosgene or carbonyl diimidazole, to provide an activated carbonates, which in turn can react with the alcohol and/or the phenolic hydroxy group on the compounds utilized herein to provide carbonate and carbamate prodrugs.

[0134] Various protecting groups and synthetic methods related to them that can be used or adapted to make compounds of the invention can be adapted from the references Testa et al., Hydrolysis in Drug and Prodrug Metabolism, June 2003, Wiley- VCH, Zurich, 419-534 and Beaumont et al., Curr. Drug Metab. 2003, 4:461-85.

[0135] Scheme 5 below provides a method of synthesizing an acyloxymethyl version of a prodrug by adapting a method from the reference Sobolev et al., 2002, J. Org. Chem. 67:401-410.


Scheme 5



wherein R^{51} is C_1 - C_6 alkyl.


[0136] Scheme 6 below provides a method for synthesizing a phosphonooxymethyl version of a prodrug by adapting a method from Mantyla et al., 2004, J. Med. Chem. 47:188-195.

Scheme 6




[0137] Scheme 7 below provides a method of synthesizing an alkyloxymethyl version of a prodrug

Scheme 7



wherein R^{52} is C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_3 - C_9 heterocyclyl, C_6 - C_{10} aryl, or C_3 - C_9 heteroaryl.

Scheme 8



[0138] Compounds of structure **17** can be synthesized via general synthetic scheme 8. Reduction of carboxylic acid derivative **14** gives hydroxymethyl analog, which can be N-derivatized via copper-mediated N-arylation reaction (CuI, Ar-I, base such as N,N-dimethylethylenediamine and potassium phosphate, heat) to give key hydroxymethyl intermediate **15**. Coupling of **15** with phenol aldehyde **16** produces the desired aldehyde analog **17** via typical Mitsunobu conditions using either triphenylphosphine or polymer supported triphenylphosphine.

[0139] **General method step 1 – reduction of carboxylic acid derivative 14 to methyl alcohol 15:** To a suspension of carboxylic acid **14** (1-10 mmol) in MeOH or EtOH (2-10 mL) at 0 °C was added SOCl₂ (1.5 eq). After stirred at room temperature for 1-12 h, it was concentrated to remove all solvents, dried under high vacuum to give corresponding methyl or ethyl ester. The ester was dissolved in MeOH or EtOH (5-30 mL), to this solution, was added NaBH₄ (1-4 eq) at 0 °C, the mixture was warmed up to room temperature and stirred for additional 1-24 h. The mixture was quenched with Sat. NH₄Cl, filtered off the insolubles and the filtrate was concentrated to give crude product, which was purified by flash silica gel chromatography to give the corresponding hydroxymethylene compound **15**.

[0140] **General method step 2 – N-alkylation (14a to 14b):** The carboxylate **14a** (R₁=H) can be first alkylated and then reduced to give N-alkyl hydroxymethylene analog **14b** (R₁=alkyl). In a typical procedure, the carboxylate **14a** (1-10 mmol) is first dissolved in DMF (2-20 mL); to this was then added a base such as NaH or Cs₂CO₃ (1-1.2 eq), followed by the addition of alkyl halide (eg, BnBr) (0.9-1.5 eq). The reaction allowed to proceed at room temperature or heat at 40 to 115 °C for 0.5 to 24 h. In workup A, water was added to the reaction mixture, the precipitated product was collected, washed with water, and then subjected to preparative HPLC or flash silica gel chromatography purification. In workup B (for products that did not precipitate), diluted HCl or aqueous NH₄Cl was added at 0 °C to


adjusted the pH to ~7, the reaction mixture was partitioned between ethyl acetate or dichloromethane and aqueous sodium chloride and the organic layer separated, dried, and solvent removed under vacuum to afford crude product which was purified by automated silica gel column chromatography, reaction appropriate solvents mixture (e.g., ethyl acetate/hexanes).

[0141] General method step 3 – Copper-mediated N-arylation from 15a to 15c: For cyclic amines (X=H, H), to a solution of hydroxymethylene compound **15a** (1-10 mmol) and aryl/hetero iodide (1-1.5eq) in iPrOH (0.5-10 mL) was added ethylene diol (1.3eq) and CuI (6.7mol%), followed by K₃PO₄ (1.3eq), then it was degassed and heated at 88 °C for 6-24 h. Alternatively, for lactams (X=O), to a solution of hydroxymethylene compound **15a** (1-10mmol) and aryl/hetero iodide (1-1.5eq) in Dioxane (2-20 mL) was added CuI (0.17eq), N,N-dimethylethylenediamine (0.17eq), K₃PO₄ (1.7eq), then it was degassed and heated at 100 °C for 6-48 h.

[0142] Workup for both procedures: the reaction mixture was cooled to room temperature the mixture was diluted with EtOAc and water, organic layer was separated and the aqueous layer was extracted with EtOAc, organic layer was combined, washed with brine, dried and concentrated to give crude product, which was purified by flash silica gel chromatography to give N-aryl/heteroaryl compound **15c**.

[0143] General method C –Mitsunobu conditions A hydroxyl (hetero)arylaldehyde derivatives (**17**) (0.1-2 mmol) mixture with substituted methylene alcohol (**16**) (0.8 to 1.2eq) and (polymer-supported) PPh₃ (1-1.5eq) in anhydrous THF (1-10mL) was stirred under nitrogen until complete dissolution. The solution was cooled to 0 °C on ice bath and DIAD or DEAD (1.1 eq) in THF or toluene was added dropwise over a 1-20 min period. The ice cooling bath was allowed to expire over 90 min and the mixture was stirred at RT for 2-48 hours. The mixture was filtered through a pad of silica. The silica was washed with ethyl acetate 2-20mL. The combined filtrates were evaporated and the residue was dried on highvac. The residue was purified by preparative HPLC or flash silica gel chromatography.

Scheme 9



[0144] **General method step 1 (Scheme 9) for preparing substituted methylene alcohol (2) by Suzuki coupling reaction.** To a solution of (2-chloropyridin-3-yl)methanol or (2-bromopyridin-3-yl)methanol (1-100mmol) and appropriate carboxylic acid or ester (0.8 to 1.5 eq) in dioxane (2-200 mL) was added a solution of sodium bicarbonate (3 eq) in water (1-100 mL), followed by the addition of Pd(dppf)Cl₂ (5 to 10mol%). After heating at 100 °C for 4-24 h, the reaction mixture was cooled and diluted with EtOAc, organic layer was washed with water, brine, dried and concentrated to give crude product, which was purified by column chromatography.

[0145] **General method step 2 (Scheme 9) for preparing substituted methylene chloride (3a).** To a solution of substituted methylene alcohol (2) (0.1 to 2 mmol) in DCM (1-10 mL) was added SOCl₂ dropwise (2eq to 5eq) at 0 °C or rt. The reaction mixture was stirred at rt for 10min to 6 h, or until reaction is judged complete (LC/MS). The reaction mixture is concentrated to dryness over a rotavap. The crude chloride residue was suspended in toluene, sonicated and concentrated to dryness. The process was repeated three times and dried under vacuum to give the substituted methylene chloride (3a), usually as an off-white solid, which was used for next step without further purification. Alternatively, a solution of aqueous 1N Na₂CO₃ is then added to produce a solution of pH~ 8. the mixture was extracted with DCM (3 x10-50mL), dried over sodium sulfate, and concentrated to the crude substituted methylene chloride (3a), which is then purified by column chromatography on silica gel (0-100% ethyl acetate-hexanes).

[0146] **General method 2 (Scheme 9) for preparing substituted methylene bromide (3b).** To a solution of substituted methylene alcohol (2) (0.1 to 2 mmol) in DCM (1-10 mL) was added Ph₃P Br₂ dropwise (2eq to 5eq) at 0 °C or rt. The reaction mixture was stirred at

rt for 10 min to 2 h, or until reaction is judged complete (LC/MS). The reaction mixture is concentrated to dryness over a rotavap. The residue purified by column chromatography on silica gel (0-100% ethyl acetate-hexanes) to afford the pure bromide **3b**.

[0147] General method step 3 (Scheme 9) for preparing aryloxy/heteroarylether analogs (5) from substituted methylene alcohol (2) and hydroxyl (hetero)aryl aldehyde derivatives (4). A hydroxyl (hetero)arylaldehyde derivatives (4) (0.1-2 mmol) mixture with substituted methylene alcohol (2) (0.8 to 1.2eq) and (polymer-supported)/PPh₃ (1-1.5eq) in anhydrous THF (1-10mL) was stirred under nitrogen until complete dissolution. The solution was cooled to 0 °C on ice bath and DIAD or DEAD (1.1 eq) in THF or toluene was added drop wise over a 1-20 min period. The ice cooling bath was allowed to expire over 90 min and the mixture was stirred at RT for 2-48 hours. The mixture was stirred for 10 min, then filtered through a pad of silica. The silica was washed with ethyl acetate 2-20mL. The combined filtrates were evaporated and the residue was dried on highvac. The residue was purified by preparative HPLC or flash silica gel chromatography.

[0148] General method step 4 (Scheme 9) for preparing aryloxy/heteroarylether analogs (5) from substituted methylene halide (3) and hydroxyl (hetero)aryl aldehyde derivatives (4). A mixture of hydroxyl (hetero)arylaldehyde derivatives (4) (0.1-2 mmol, 1-4 eq.), substituted methylene chloride or bromide (3) (1eq), and K₂CO₃ (2-5 eq.) (catalytic amount of NaI or Bu₄NI may also be added) in DMF, acetonitrile, NMP or DMSO (1 to 10 mL) was stirred at RT or heating up to 120 °C for 1-24 h under nitrogen atmosphere. In workup A, water was added to the reaction mixture, the precipitated product was collected, washed with water, and then subjected to preparative HPLC or flash silica gel chromatography purification. In workup B (for products that did not precipitate), diluted HCl or aqueous NH₄Cl was added at 0 °C to adjusted the pH to ~7, the reaction mixture was partitioned between ethyl acetate or dichloromethane and aqueous sodium chloride and the organic layer separated, dried, and solvent removed under vacuum to afford crude product which was purified by automated silica gel column chromatography using appropriate solvents mixture (e.g., ethyl acetate/hexanes).

Examples

[0149] In the examples below as well as throughout the application, the following abbreviations have the following meanings. If not defined, the terms have their generally accepted meanings.


°C	=	degrees Celsius
RT	=	Room temperature
min	=	minute(s)
h	=	hour(s)
μL	=	Microliter
mL	=	Milliliter
mmol	=	Millimole
eq	=	Equivalent
mg	=	Milligram
ppm	=	Parts per million
atm	=	Atmospheric pressure
MS	=	Mass spectrometry
LC-MS	=	Liquid chromatography–mass spectrometry
HPLC	=	High performance liquid chromatography
NMR	=	Nuclear magnetic resonance
Sat./sat.	=	Saturated
MeOH	=	Methanol
EtOH	=	Ethanol
EtOAc	=	Ethyl acetate
Et ₃ N	=	Triethylamine
ACN	=	Acetonitrile
Ac ₂ O	=	Acetic anhydride
Na(OAc) ₃ BH	=	Sodium triacetoxo borohydride
PBr ₃	=	phosphorus tribromide
Ph ₃ P	=	Triphenylphosphine
Ph ₃ PBr ₂	=	Triphenylphosphine dibromide
CBr ₄	=	Tetrabromomethane
DMF	=	N, N-Dimethylformamide
DCM	=	Dichloromethane

LAH/ LiAlH ₄	=	Lithium aluminum hydride
THF	=	Tetrahydrofuran
DIBAL	=	Diisobutylaluminium hydride
DIAD	=	Diisopropyl azodicarboxylate
DEAD	=	Diethyl azodicarboxylate
DIPEA	=	N,N-Diisopropylethylamine
Tf ₂ O	=	Trifluoromethanesulfonic (triflic) anhydride
Pd(dppf)Cl ₂	=	[1,1'-Bis(diphenylphosphino)ferrocene] dichloropalladium(II), complex


[0150] The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion. The present examples, along with the methods described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses which are encompassed within the spirit of the invention as defined by the scope of the claims will occur to those skilled in the art.

Experimental Procedures for Intermediates

[0151] (E)-1-(3-(((tert-butyldimethylsilyl)oxy)methyl)pyridin-2-yl)-3-(dimethylamino)prop-2-en-1-one (**INT-1**)




Step 1



[0152] To a mixture of (2-bromopyridin-3-yl)methanol (20.0 g, 106.4 mmol, 1 eq.; refer to example 14) and imidazole (14.5 g, 212.8 mmol, 2 eq.) in DMF (50.0 mL) was added TBSCl


(19.2 g, 150.7 mmol, 1.2 eq.) at RT. The mixture was stirred at RT for 1 h and diluted with a mixture of water (100 mL) and EtOAc (300 mL). The organic layer was washed with $\text{NH}_4\text{Cl}_{(\text{sat.})}$ solution and brine, dried over Na_2SO_4 , concentrated, and purified on silica gel using 10% EtOAc/hexanes as eluent to give 2-bromo-3-((tert-butyl dimethylsilyloxy)methyl)pyridine (30.1 g, 94%) as a colorless oil. MS (ESI) m/z 302.0 $[\text{M}+\text{H}]^+$.

Step 2



[0153] A mixture of 2-bromo-3-((tert-butyl dimethylsilyloxy)methyl)pyridine (30.1 g, 100.0 mmol, 1 eq.) and $\text{Zn}(\text{CN})_2$ (23.5 g, 200.0 mmol, 2.0 eq.) in DMF (100.0 mL) was purged with N_2 for 5 min and added $\text{Pd}(\text{PPh}_3)_4$ (5.78 g, 5.0 mmol, 0.05 eq.). The mixture was heated at 120 °C for 2 h under N_2 , cooled, filtered, concentrated, and purified on silica gel using a mixture of EtOAc and hexanes as eluent to give 3-((tert-butyl dimethylsilyloxy)methyl)picolinonitrile (20.4 g, 82%) as a colorless oil. MS (ESI) m/z 249.1 $[\text{M}+\text{H}]^+$.


Step 3:



[0154] Methylmagnesium bromide (3M/ether, 41.0 mL, 123.4 mmol) was added to a stirred solution of 3-((tert-butyl dimethylsilyloxy)methyl)picolinonitrile (20.4 g, 82.25 mmol) in THF (100.0 mL) at -78 °C. The reaction mixture was warm to RT, quenched with aqueous citric acid solution, and extracted with EtOAc (50 mL) twice. The combined organic layers were washed with $\text{NaHCO}_3_{(\text{sat.})}$ solution and brine, dried over Na_2SO_4 , concentrated, and purified on silica gel using a mixture of EtOAc/hexanes as eluent to give 1-(3-((tert-


butyldimethylsilyloxy)methyl)pyridin-2-yl)ethanone (12.9 g, 59%) as a colorless oil. MS (ESI) m/z 266.2 $[M+H]^+$.

Step 4:




[0155] 1-(3-((tert-butyldimethylsilyloxy)methyl)pyridin-2-yl)ethanone (10.8 g, 40.75 mmol) in dimethoxy-N,N-dimethylmethanamine (15.0 mL) was heated to reflux for 3 days. The mixture was concentrated and used for next step without further purification. MS (ESI) m/z 321.1 $[M+H]^+$.

[0156] *Preparation of 3-(chloromethyl)-2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridine (INT-2).*




Step 1:




[0157] To (3,3,3-trifluoroethyl)hydrazine (25 g, 50% wt in water, 153.5 mmol, 1 eq.) in a RB flask (250 mL) was added HCl (12 N, 25.6 mL, 307.0 mmol, 2 eq.). The mixture was concentrated to give (3,3,3-trifluoroethyl)hydrazine dihydrochloride (1.07 g) as a yellow solid. MS (ESI) m/z 115.1 $[M+H]^+$.

Step 2:



[0158] To (*E*)-1-(3-((tert-butyldimethylsilyloxy)methyl)pyridin-2-yl)-3-(dimethylamino)prop-2-en-1-one (crude above, 5.91 g, 18.44 mmol, 1 eq.) in EtOH (20 mL) was added (3,3,3-trifluoroethyl)hydrazine dihydrochloride (4.13 g, crude above, 22.13 mmol, 1.2 eq.) at RT. The mixture was heated at 80 °C for 1 h, concentrated, and diluted with EtOAc (50 mL) and NaHCO_{3(sat)} solution (10 mL). The layers were separated and aqueous layer was extracted with EtOAc three times. The combined organic layers were dried over Na₂SO₄, concentrated, and purified on silica gel using a mixture of EtOAc and hexanes as eluent to give 3-((tert-butyldimethylsilyloxy)methyl)-2-(1-(3,3,3-trifluoroethyl)-1H-pyrazol-5-yl)pyridine (5.90 g; 86% for 2 steps). MS (ESI) *m/z* 372.2 [M+H]⁺.


Step 3:



[0159] To 3-((tert-butyldimethylsilyloxy)methyl)-2-(1-(3,3,3-trifluoroethyl)-1H-pyrazol-5-yl)pyridine (5.91 g, 15.93 mmol) in MeOH (20 mL) was added HCl (4 N, 8.0 mL). The mixture was stirred at RT for 1 h, concentrated, and diluted with EtOAc (50 mL) and NaHCO_{3(sat)} solution (10 mL). The layers were separated and aqueous layer was extracted with EtOAc three times. The combined organic layers were dried over Na₂SO₄, and concentrated to give (2-(1-(3,3,3-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methanol (4.1 g, quantitative yield) as colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 8.54 (dd, *J* = 4.7, 1.5 Hz, 1H), 7.92 (dd, *J* = 7.9, 1.2 Hz, 1H), 7.57 (d, *J* = 1.9 Hz, 1H), 7.30 (dd, *J* = 7.8, 4.8 Hz, 1H),


6.50 (d, $J = 1.9$ Hz, 1H), 5.09 (q, $J = 8.6$ Hz, 2H), 4.63 (s, 2H), 1.76 (s, 1H). MS (ESI) m/z 258.1 $[M+H]^+$.

Step 4:




[0160] To 2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-ylmethanol (408 mg, 1.59 mmol) in DCM (5 mL) was added SOCl_2 (1.5 mL) at RT. The reaction mixture was stirred at RT for 4 h and concentrated to dryness. The crude solid was suspended in toluene and concentrated to dryness. The process was repeated three times and dried under vacuum to give 3-(chloromethyl)-2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridine hydrochloride (498 mg) as an off-white solid, which was used for next step without further purification.

[0161] Preparation of 3-(chloromethyl)-2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridine (**INT-3**).




Step 1:



[0162] To a mixture of benzyl hydrazinecarboxylate (5.0 g, 30.3 mmol, 1 eq.) and DIEA (15.0 mL, 90.9 mmol, 3 eq.) in DMF (20 mL) was added 3,3,3-trifluoropropyl bromide (10.7 g 60.6 mmol, 2 eq.) at RT. The mixture was heated at 80 °C for 20 h, concentrated, and purified on silica gel using a mixture of EtOAc and hexanes as eluent to benzyl 2-(3,3,3-


trifluoropropyl) hydrazinecarboxylate (4.2 g; 53%) as a white solid. ^1H NMR (400 MHz, CDCl_3) δ 7.33 – 7.17 (m, 5H), 6.11 (s, 1H), 5.01 (s, 2H), 4.00 (s, 1H), 3.00 (dd, $J = 12.2, 7.1$ Hz, 2H), 2.17 (qt, $J = 10.8, 7.3$ Hz, 2H). MS (ESI) m/z 263.1 $[\text{M}+\text{H}]^+$.

Step 2:



[0163] To benzyl 2-(3,3,3-trifluoropropyl)hydrazinecarboxylate (1.7 g, 6.49 mmol, 1 eq.) in a mixture of EtOH (30 mL) were added Pd/C (1.0 g) and HCl (12 N, 2.0 mL). The mixture was charged with H_2 (60 psi), stirred at RT for 1 h, filtered, and concentrated to give (3,3,3-trifluoropropyl)hydrazine dihydrochloride (1.07 g) as a yellow solid. MS (ESI) m/z 129.1 $[\text{M}+\text{H}]^+$.


Step 3:



[0164] To (*E*)-1-(3-((tert-butyldimethylsilyloxy)methyl)pyridin-2-yl)-3-(dimethylamino)prop-2-en-1-one (crude above, 1.73 g, 5.41 mmol, 1 eq.) in EtOH (10 mL) was added (3,3,3-trifluoropropyl)hydrazine dihydrochloride (1.30 g, crude above, 6.49 mmol, 1.2 eq.) at RT. The mixture was heated at 80 °C for 1 h, concentrated, and diluted with EtOAc (50 mL) and $\text{NaHCO}_3(\text{sat})$ solution (10 mL). The layers were separated and aqueous layer was extracted with EtOAc three times. The combined organic layers were dried over Na_2SO_4 , concentrated, and purified on silica gel using a mixture of EtOAc and hexanes as eluent to give 3-((tert-butyldimethylsilyloxy)methyl)-2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridine (1.58 g; 76% for 2 steps). ^1H NMR (400 MHz, CDCl_3) δ 8.53 (dd, $J =$


4.7, 1.6 Hz, 1H), 7.96 – 7.88 (m, 1H), 7.51 (d, $J = 1.9$ Hz, 1H), 7.29 (dd, $J = 7.9, 4.7$ Hz, 1H), 6.34 (d, $J = 1.9$ Hz, 1H), 4.62 (s, 2H), 4.45 – 4.33 (m, 2H), 2.82 – 2.61 (m, 2H), 0.85 (s, 8H), -0.00 (s, 5H). MS (ESI) m/z 386.2 $[M+H]^+$.

Step 4:



[0165] To 3-((tert-butyldimethylsilyloxy)methyl)-2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridine (1.58 g, 4.1 mmol) in MeOH (20 mL) was added HCl (4 N, 4.0 mL). The mixture was stirred at RT for 1 h, concentrated, and diluted with EtOAc (50 mL) and $\text{NaHCO}_3(\text{sat})$ solution (10 mL). The layers were separated and aqueous layer was extracted with EtOAc three times. The combined organic layers were dried over Na_2SO_4 , and concentrated to give 2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-ylmethanol (1.1 g, 99%) as colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 8.64 (dd, $J = 4.7, 1.7$ Hz, 1H), 8.00 (dd, $J = 7.9, 1.7$ Hz, 1H), 7.57 (d, $J = 1.9$ Hz, 1H), 7.38 (dd, $J = 7.9, 4.8$ Hz, 1H), 6.48 (d, $J = 1.9$ Hz, 1H), 4.69 (s, 2H), 4.51 – 4.43 (m, 2H), 2.85 – 2.72 (m, 2H), 2.70 (s, 1H). MS (ESI) m/z 272.1 $[M+H]^+$.


Step 5:



[0166] To 2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-ylmethanol (140 mg, 0.52 mmol) in DCM (5 mL) was added SOCl_2 (2.0 mL) at RT. The reaction mixture was stirred at RT for 4 h and concentrated to dryness. The crude solid was suspended in toluene and concentrated to dryness. The process was repeated three times and dried under vacuum.

to give 3-(chloromethyl)-2-(1-(2,2,2-trifluoropropyl)-1H-pyrazol-5-yl)pyridine hydrochloride (498 mg) as an off-white solid, which was used for next step without further purification.


[0167] *Preparation of 3-(chloromethyl)-2-(1-isopropyl-1H-pyrazol-5-yl)pyridine (INT-4).*



[0168] **Step 1:** To a 500-mL flask containing the pyrazole boronate (9.0g, 38.1mmol), 2-chloropyridine (5.47g, 38.1mmol), Pd(dppf)Cl₂ ([1,1-bis(diphenylphosphino)ferrocene] dichloropalladium) (1.39g, 1.91mmol, 5%mol), and sodium bicarbonate (9.61g, 114.4mmol, 3 equiv) was added 100 mL of dioxane and 30 mL of water. The mixture was heated under nitrogen at 100 °C for 12 hrs. Then solvents were removed on a rotavap at 40 °C undervacuum. The resulting brown residue was suspended in 20%EtOAc/DCM (60mL), filtered through a pad of silica gel (15g); washed with 20%EtOAc/DCM (4x20mL). The combined filtrate were concentrated to afford a brown oil (13 g). The residue was dissolved 10% EtOAc/hexanes (20mL) and loaded on a Biotage 100g snap SiO₂ column and eluted with 0-50% EtOAc. (2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol was obtained as a light brown oil (3.32 g, 40%). MS (ESI) m/z 218 [M+H]⁺.

[0169] **Step 2:** To a solution of (2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol (440mg, 2.02mmol) in DCM (4 mL) was added SOCl₂ (2eq) at 0 °C. The reaction mixture was stirred at RT for 15mins and concentrated to dryness. The crude solid was suspended in toluene and concentrated to dryness. The process was repeated three times and dried under vacuum to give 3-(chloromethyl)-2-(1-isopropyl-1H-pyrazol-5-yl)pyridine hydrochloride (432 mg) as an off-white solid, which was used for next step without further purification. MS (ESI) m/z 236.5 [M+H]⁺.


[0170] Preparation of 3-(chloromethyl)-2-(1-cyclopentyl-1H-pyrazol-5-yl)pyridine (INT-5).




[0171] **Step 1:** To (E)-1-(3-((tert-butyldimethylsilyloxy)methyl)pyridin-2-yl)-3-(dimethylamino)prop-2-en-1-one (crude, 3.205g, 10.0 mmol, 1 eq.) in EtOH (30 mL) was added cyclopentylhydrazine HCl salt (1.639g, 12.0 mmol, 1.2eq) at RT. The mixture was heated at 80 °C for 2 h, concentrated, and purified on silica gel using a mixture of EtOAc and hexanes as eluent to give a mixture of regio-isomers, the less polar (2-(1-cyclopentyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol was obtained as a light brown oil (440mg). MS (ESI) m/z 244.2 $[M+H]^+$.

[0172] **Step 2:** To a solution of (2-(1-cyclopentyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol (301mg, 1.24mmol) in DCM (3 mL) was added SOCl_2 (3eq) at 0 °C. The reaction mixture was stirred at RT for 15mins (the reaction was done in 10mins by LCMS) and concentrated to dryness. The crude solid was suspended in toluene and concentrated to dryness. The process was repeated three times and dried under vacuum to give 3-(chloromethyl)-2-(1-cyclopentyl-1H-pyrazol-5-yl)pyridine hydrochloride (305 mg) as an off-white solid, which was used for next step without further purification. MS (ESI) m/z 262.2 $[M+H]^+$.

[0173] Preparation of 5-hydroxy-2-(2-methoxyethoxy)isonicotinaldehyde (INT-6).




Step 1




[0174] To a solution of 6-(benzyloxy)pyridin-3-ol (2.0 g, 10 mmol, 1 eq.) in DMF (20 mL) was added NaH (60% in mineral oil; 0.6 g, 15 mmol, 1.5eq.) at 0-5 °C portion-wise. Upon the completion of addition, the mixture was continued to stir at 0-5 °C for 15 min, added chloromethyl methyl ether (0.88 g, 11 mmol, 1.1 eq.), stirred at 0-5 °C for another 20 min, and quenched with $\text{NH}_4\text{Cl}_{(\text{sat.})}$ solution. The aqueous layer was extracted with EtOAc (3 x 20 mL) and the combined organic layers were washed with water and brine, dried over Na_2SO_4 , concentrated, and purified on silica gel using 25% EtOAc/hexanes as eluent to give 2-(benzyloxy)-5-(methoxymethoxy)pyridine (2.1 g, 87%) as a colorless oil. MS (ESI) m/z 246.1 $[\text{M}+\text{H}]^+$.

Step 2



[0175] To 2-(benzyloxy)-5-(methoxymethoxy)pyridine (1.8 g, 8.71 mol) in EtOH was added Pd/C (1.0 g). The mixture was charged with H_2 (15 psi), stirred at RT for 45 min, filtered, and concentrated to give 5-(methoxymethoxy)pyridin-2-ol (1.35 g, quantitative yield) as a pale yellow solid. MS (ESI) m/z 156.1 $[\text{M}+\text{H}]^+$.


Step 3



[0176] To a mixture of 5-(methoxymethoxy)pyridin-2-ol (1.35 g, 8.71 mmol, 1 eq.) and K_2CO_3 (6.01 g, 43.6 mmol, 5.0 eq.) in DMF (30.0 mL) was added 1-bromo-2-methoxyethane


(3.61 g, 26.1 mmol, 3eq.). The mixture was heated at 60 °C for 2 h, cooled, filtered, concentrated, and purified on silica gel using a mixture of EtOAc and hexanes as eluent to give 2-(2-methoxyethoxy)-5-(methoxymethoxy)pyridine (500 mg, 27%) as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.94 (d, *J* = 3.0 Hz, 1H), 7.35 (ddd, *J* = 8.9, 3.0, 1.0 Hz, 1H), 6.76 (dd, *J* = 8.9, 1.0 Hz, 1H), 5.11 (s, 2H), 4.48 – 4.40 (m, 2H), 3.79–3.71 (m, 2H), 3.50 (s, 3H), 3.45 (s, 3H). MS (ESI) *m/z* 214.1 [M+H]⁺.

Step 4




[0177] To a mixture of 2-(2-methoxyethoxy)-5-(methoxymethoxy)pyridine (1.34 g, 6.3 mol, 1 eq.) and diisopropylamine (17.5 uL, 0.13 mmol, 0.02 eq.) in THF (50 mL) was added methyl lithium (1.6 M/THF, 7 mL, 11.3 mol, 1.8 eq.) at -40 °C. Upon the completion of addition, the mixture was warmed to 0 °C, continued to stir at 0 °C for 3 h, cooled back down to -40 °C, and added DMF (0.83 mL, 11.3 mol, 1.8 eq.) slowly. The mixture was then stirred at -40 °C for 1 h, quenched with a mixture of HCl (12 N, 12 mL) and THF (28 mL), warmed to RT, and added water (20 mL). The pH of the mixture was adjusted to pH 8-9 with solid K₂CO₃. The aqueous layer was extracted with EtOAc (30 mL) twice. The combined organic layers were dried over Na₂SO₄, concentrated, and purified on silica gel using a mixture of EtOAc and hexanes as eluent to give a mixture of 2-(2-methoxyethoxy)-5-(methoxymethoxy)isonicotinaldehyde and 2-(2-methoxyethoxy)-5-(methoxymethoxy)nicotinaldehyde (5/1, 1.27 g, 83.6%) as a pale yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 10.45 (s, 1H), 8.23 (s, 1H), 7.16 (s, 1H), 5.27 (s, 2H), 4.46 (dd, *J* = 5.4, 3.9 Hz, 2H), 4.14 (q, *J* = 7.1 Hz, 1H), 3.77 – 3.71 (m, 2H), 3.56 (s, 3H), 3.46 (s, 3H) and ¹H NMR (400 MHz, CDCl₃) δ 10.41 (s, 1H), 8.18 (d, *J* = 3.2 Hz, 1H), 7.85 (d, *J* = 3.1 Hz, 1H), 5.16 (s, 2H), 4.64 – 4.57 (m, 2H), 3.85 – 3.79 (m, *J* = 5.4, 4.0 Hz, 2H), 3.50 (s, 3H), 3.46 (s, 3H); MS (ESI) *m/z* 242.1 [M+H]⁺.

Step 5



[0178] To a solution of 2-methoxy-5-(methoxymethoxy)isonicotinaldehyde (1.27 g, 5.29 mol) in THF (5 mL) was added HCl (3 N, 4 mL). The reaction was stirred at 50 °C for 1 h, cooled to RT, and diluted with water (5 mL). The mixture was neutralized to pH 7-8 with solid K₂CO₃ and the aqueous layer was extracted with EtOAc (100 mL) twice. The combined organic layers were dried over Na₂SO₄, concentrated, and purified on silica gel using a mixture of EtOAc and hexanes to give 5-hydroxy-2-(2-methoxyethoxy)isonicotinaldehyde (630 mg, 60%) and 5-hydroxy-2-(2-methoxyethoxy)nicotinaldehyde (120 mg, 11%). Data for 5-hydroxy-2-(2-methoxyethoxy)isonicotinaldehyde: ¹H NMR (400 MHz, CDCl₃) δ 9.98 (s, 1H), 9.50 (s, 1H), 8.07 (s, 1H), 7.02 (s, 1H), 4.51 – 4.39 (m, 2H), 3.81 – 3.72 (m, 2H), 3.47 (s, 3H). LRMS (M+H⁺) m/z 198.1. Data for 5-hydroxy-2-(2-methoxyethoxy)nicotinaldehyde: ¹H NMR (400 MHz, CDCl₃) δ 10.3 (s, 1H), 7.99 (d, *J* = 3.2 Hz, 1H), 7.58 (d, *J* = 3.2 Hz, 1H), 7.18 – 7.07 (br, 1H), 4.54 (dd, *J* = 5.4, 3.7 Hz, 2H), 3.84 (dd, *J* = 5.4, 3.7 Hz, 2H), 3.49 (s, 3H); MS (ESI) m/z 198.1 [M+H]⁺.

[0179] Preparation of 2,6-dihydroxybenzaldehyde (INT-7).




[0180] Into a 3000-mL three neck round-bottom flask, was placed a solution of AlCl₃ (240 g, 1.80 mol, 3.00 equiv) in dichloromethane (1200mL). A solution of 2,6-dimethoxybenzaldehyde (100 g, 601.78 mmol, 1.00 equiv) in dichloromethane (800ml) was added to the reaction mixture dropwise at 0°C. The resulting solution was stirred overnight at room temperature, and then it was quenched with 200 mL of diluted HCl (2M). The resulting solution was extracted with 2x200 mL of dichloromethane. The combined organic layers were concentrated under vacuum. The residue was applied onto a silica gel column with ethyl

acetate/petroleum ether (1:200-1:50) as eluent to furnish 40 g (48%) of 2,6-dihydroxybenzaldehyde as a yellow solid.

$^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.25(s, 2H), 10.25(s, 1H), 7.36(m, 1H), 6.36 (d, $J=8.4\text{ Hz}$ 2H); MS (ESI) m/z 139 $[\text{M}+\text{H}]^+$.

[0181] Preparation of 5-hydroxy-2-methoxyisonicotinaldehyde (**INT-8**).



[0182] Step 1: To a solution of 6-methoxypyridin-3-ol (20 g, 0.16 mol) in DMF (200 mL) was added NaH (60% in mineral oil; 9.6 g, 0.24 mol) at 0-5 °C portion-wise. Upon the completion of addition, the mixture was continued to stir at 0-5 °C for 15 min followed by additional of chloromethyl methyl ether. The mixture was stirred at 0-5 °C for another 20 min and quenched with aqueous $\text{NH}_4\text{Cl}_{(\text{sat})}$. The aqueous layer was extracted with EtOAc (3 x 100 mL) and the combined organic layer was washed with water and brine, dried over Na_2SO_4 , and concentrated under reduced pressure. The residue was purified on silica gel with 25% EtOAc/hexanes as eluent to give 2-methoxy-5-(methoxymethoxy)pyridine (24.1 g, 89.3%) as a colorless oil. $^1\text{H NMR}$ (400 MHz; CDCl_3) 7.97 (d, 1 H), 7.35 (dd, 1 H), 6.70 (d, 1 H), 5.12 (s, 2 H), 3.91 (s, 3 H), 3.51 (s, 3 H); MS (ESI) m/z 170.1 $[\text{M}+\text{H}]^+$.


[0183] Step 2: To a mixture of 2-methoxy-5-(methoxymethoxy)pyridine (30 g, 0.178 mol) and diisopropylamine (507 μL , 3.6 mmol) in THF (500 mL) was added methyl lithium (1.6 M/THF, 200 mL, 0.32 mol) at -40 °C. Upon the completion of addition, the mixture was warmed to 0 °C and continued to stir at 0 °C for 3 h. The reaction mixture was then cooled back down to -40 °C followed by addition of DMF (24.7 mL, 0.32 mol) slowly. The mixture was then stirred at -40 °C for 1 h and quenched with a mixture of HCl (12 N, 120 mL) and THF (280 mL). Water (200 mL) was added and the pH of the mixture was adjusted to pH 8-9 with solid K_2CO_3 . The mixture was extracted with EtOAc (300 mL) twice. The organic layer was combined, dried over Na_2SO_4 , and concentrated to give 2-methoxy-5-(methoxymethoxy)isonicotinaldehyde (33.5 g, 95.7%) as a brown solid, which was used for next step without further purification. $^1\text{H NMR}$ (400 MHz; CD_3OD) 7.90 (s, 1 H), 6.92 (s, 1 H), 5.64 (s, 1 H), 5.20 (s, 2 H), 3.84 (s, 3 H), 3.48 (s, 3 H); MS (ESI) m/z 198.1 $[\text{M}+\text{H}]^+$.

[0184] **Step 3:** To a solution of 2-methoxy-5-(methoxymethoxy)isonicotinaldehyde (33.5 g, 0.17 mol) in THF (150 mL) was added HCl (3 N, 250 mL). The reaction was stirred at 50 °C for 1 h, cooled to RT and diluted with water (500 mL). The mixture was neutralized to pH 7-8 with solid K₂CO₃. The pale yellow solid was collected, washed with water, and dried in vacuum oven (40 °C) overnight to give 5-hydroxy-2-methoxyisonicotinaldehyde (17.9 g, 74.6%). ¹H NMR (400 MHz; DMSO) δ= 10.31 (s, 1 H), 8.03 (s, 1 H), 6.89 (s, 1 H), 3.80 (s, 3 H); MS (ESI) m/z 154.0 [M+H]⁺.


Experimental procedures for Examples:


[0185] **GBT527** Preparation of 2-methoxy-5-[[2-[1-(2,2,2-trifluoroethyl)pyrazol-3-yl]pyridin-3-yl]methoxy]pyridine-4-carbaldehyde.

[0186] GBT527 was prepared using general method B from 5-hydroxy-2-methoxyisonicotinaldehyde and **INT-2**.




[0187] **GBT576** Preparation of 2-oxo-5-[[2-(2-propan-2-ylpyrazol-3-yl)pyridin-3-yl]methoxy]-1H-pyridine-4-carbaldehyde





Step 1:




[0188] To (*E*)-1-(3-((*tert*-butyldimethylsilyloxy)methyl)pyridin-2-yl)-3-(dimethylamino)prop-2-en-1-one (crude, 1.03 g, 3.22 mmol, 1 eq.; **INT-1**) in EtOH (10 mL) was added isopropylhydrazine hydrochloride (430 mg, 3.86 mmol, 1.2 eq.). The mixture was heated at 80 °C for 2 h, cooled, added HCl (6 N, 0.5 mL), and stirred O/N. The mixture was concentrated and diluted with EtOAc (80 mL) and NaHCO_{3(sat)} (10 mL) solution. The layers were separated and the aqueous layer was extracted with EtOAc three times. The combined organic layers were dried over Na₂SO₄, concentrated, and purified on silica gel using EtOAc as eluent to give (2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol (500 mg, 71%) and (2-(1-isopropyl-1H-pyrazol-3-yl)pyridin-5-yl)methanol (55 mg, 25%) as pale yellow oils. Data for 2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol: ¹H NMR (400 MHz, CDCl₃) δ 8.67 (dd, *J* = 4.7, 1.5 Hz, 1H), 8.0 (d, *J* = 7.8 Hz, 1H), 7.61 (d, *J* = 1.8 Hz, 1H), 7.39 (dd, *J* = 7.8, 4.8 Hz, 1H), 6.37 (d, *J* = 1.8 Hz, 1H), 4.67 (s, 2H), 4.55 (sep, *J* = 6.6 Hz 1H), 1.98-2.05 (br, 1H), 1.47 (d, *J* = 6.6 Hz, 6H). LRMS (M+H⁺) *m/z* 218.1 Data for (2-(1-isopropyl-1H-pyrazol-3-yl)pyridin-5-yl)methanol: ¹H NMR (400 MHz, CDCl₃) δ 8.62 (dd, *J* = 4.8, 1.6 Hz, 1H), 7.72 (d, *J* = 7.6 Hz, 1H), 7.55 (d, *J* = 2.4 Hz, 1H), 7.23 (dd, *J* = 7.6, 4.8 Hz, 1H), 6.99 (dd, *J* = 8.0, 6.5 Hz, 1H), 6.07 (t, *J* = 7.6 Hz, 1H), 4.67 (d, *J* = 7.6 Hz, 2H), 4.58 (sep, *J* = 6.7 Hz, 1H), 1.60 (d, *J* = 6.7 Hz, 1H). MS (ESI) *m/z* 218.1 [M+H]⁺.

Step 2:



[0189] To 2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-ylmethanol (560 mg, 2.58 mmol) in DCM (10 mL) was added SOCl_2 (3.0 mL) at RT. The reaction mixture was stirred at RT for 4 h and concentrated to dryness. The crude solid was suspended in toluene and concentrated to dryness. The process was repeated three times and dried under vacuum to give 3-(chloromethyl)-2-(1-isopropyl-1H-pyrazol-5-yl)pyridine hydrochloride (700 mg) as an off-white solid, which was used for next step without further purification.

Step 3:




[0190] A mixture of 5-hydroxy-2-methoxyisonicotinaldehyde (395 mg, 2.58 mmol, 1 eq.), 3-(chloromethyl)-2-(1-isopropyl-1H-pyrazol-5-yl)pyridine hydrochloride (700 mg, 2.58 mmol, 1 eq.), and K_2CO_3 (1.4 g, 10.32 mmol, 4 eq.) in DMF (10.0 mL) was heated at 70 °C for 2 h. The mixture was cooled, filtered, concentrated, and purified on silica gel using a mixture of EtOAc and hexanes as eluent to give 5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde (590 mg, 65%) as an off-white solid. ^1H NMR (400 MHz, CDCl_3) δ 10.41 (s, 1H), 8.76 (dd, J = 4.7, 1.6 Hz, 1H), 8.04 (dd, J = 7.9, 1.6 Hz, 1H), 7.90 (s, 1H), 7.61 (d, J = 1.8 Hz, 1H), 7.44 (dd, J = 7.9, 4.8 Hz, 1H), 7.10 (s, 1H), 6.37 (d, J = 1.8 Hz, 1H), 5.14 (s, 2H), 4.65 (sep, J = 6.6 Hz, 1H), 3.91 (s, 3H), 1.49 (d, J = 6.6 Hz, 6H); MS (ESI) m/z 353.1 $[\text{M}+\text{H}]^+$.

[0191] To 5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde (590 mg) suspended in water (5.0 mL) was added HCl (6 N, 4 mL). Once the mixture turned into a homogeneous solution, it was freeze at -78 °C to a solid and pump under high vacuum O/N. The yellow solid was continued to pump at 45 °C

for 20 h, dissolved in water (2.0 mL), and basified to pH 11 with NaOH (2 N). The aqueous layer was washed with DCM three times and the pH of the mixture was adjusted to pH 6-7. The solid was collected and dried to give 2-oxo-5-[[2-(2-propan-2-ylpyrazol-3-yl)pyridin-3-yl]methoxy]-1H-pyridine-4-carbaldehyde as a yellow solid. ^1H NMR (400 MHz, CDCl_3) δ 10.3 (s, 1H), 8.8 (dd, $J = 4.7, 1.6$ Hz, 1H), 8.1 (dd, $J = 7.9, 1.5$ Hz, 1H), 7.6 (s, 1H), 7.5 (d, $J = 1.8$ Hz, 1H), 7.1 (s, 1H), 7.0 (s, 1H), 6.6 (d, $J = 1.8$ Hz, 1H), 4.9 (s, 2H), 4.7 (sep, $J = 6.6$ Hz, 1H), 1.5 (d, $J = 6.6$ Hz, 6H); MS (ESI) m/z 339.4 $[\text{M}+\text{H}]^+$.

[0192] GBT779 Preparation of 2-(2-morpholin-4-ylethoxy)-5-[[2-(2-propan-2-ylpyrazol-3-yl)pyridin-3-yl]methoxy]pyridine-4-carbaldehyde


[0193] GTB779 was prepared according to general method B from 5-hydroxy-2-(2-morpholinoethoxy)isonicotinaldehyde and INT-4.



[0194] ^1H NMR (400 MHz, Chloroform- d) δ 10.33 (s, 1H), 8.68 (dd, $J = 4.8, 1.7$ Hz, 1H), 7.95 (dd, $J = 7.9, 1.7$ Hz, 1H), 7.79 (s, 1H), 7.53 (d, $J = 1.8$ Hz, 1H), 7.36 (dd, $J = 7.9, 4.7$ Hz, 1H), 7.04 (s, 1H), 6.28 (d, $J = 1.8$ Hz, 1H), 5.06 (s, 2H), 4.57 (s, 0H), 4.32 (t, $J = 5.7$ Hz, 2H), 3.69 – 3.62 (m, 4H), 2.70 (t, $J = 5.7$ Hz, 2H), 2.53 – 2.45 (m, 4H), 1.41 (d, $J = 6.6$ Hz, 6H); MS (ESI) m/z 452 $[\text{M}+\text{H}]^+$.

[0195] GBT832 Preparation of 2-(2-methoxyethoxy)-5-[[2-[2-(2,2,2-trifluoroethyl)pyrazol-3-yl]pyridin-3-yl]methoxy]pyridine-4-carbaldehyde.


[0196] GTB832 was prepared according to general method B from 5-hydroxy-2-(2-methoxyethoxy)isonicotinaldehyde (INT-5) and INT-2.



[0197] ^1H NMR (400 MHz, CDCl_3) δ 10.32 (s, 1H), 8.67 (dd, $J = 4.8, 1.6$ Hz, 1H), 7.97 (dd, $J = 7.9, 1.5$ Hz, 1H), 7.87 (s, 1H), 7.59 (d, $J = 1.9$ Hz, 1H), 7.38 (dd, $J = 7.9, 4.8$ Hz, 1H), 7.11 (s, 1H), 6.47 (d, $J = 1.9$ Hz, 1H), 5.17 (q, $J = 8.6$ Hz, 2H), 5.10 (s, 2H), 4.39 – 4.32 (m, 2H), 3.70 – 3.63 (m, 2H); MS (ESI) m/z 437 $[\text{M}+\text{H}]^+$.

[0198] **GBT835** Preparation of 6-methyl-3-[[2-[2-(2,2,2-trifluoroethyl)pyrazol-3-yl]pyridin-3-yl]methoxy]pyridine-2-carbaldehyde


[0199] **GBT835** was prepared according to general method B from 3-hydroxy-6-methylpicolinaldehyde and **INT-2**.



[0200] ^1H NMR (400 MHz, CDCl_3) δ 10.23 (s, 1H), 8.64 (dd, $J = 4.7, 1.6$ Hz, 1H), 8.16 (dd, $J = 7.9, 1.5$ Hz, 1H), 7.61 (d, $J = 1.9$ Hz, 1H), 7.38 (dd, $J = 7.9, 4.8$ Hz, 1H), 7.21 (d, $J = 8.6$ Hz, 1H), 7.10 (d, $J = 8.6$ Hz, 1H), 6.47 (d, $J = 1.9$ Hz, 1H), 5.19 (q, $J = 8.6$ Hz, 2H), 5.12 (d, $J = 6.1$ Hz, 2H), 2.51 (s, 3H); MS (ESI) m/z 377 $[\text{M}+\text{H}]^+$.

[0201] **GBT836** Preparation of 6-methyl-3-[[2-[2-(3,3,3-trifluoropropyl)pyrazol-3-yl]pyridin-3-yl]methoxy]pyridine-2-carbaldehyde


[0202] **GBT836** was prepared according to general method B from 3-hydroxy-6-methylpicolinaldehyde and **INT-3**.



[0203] ^1H NMR (400 MHz, CDCl_3) δ 10.31 (s, 1H), 8.75 (dd, $J = 4.7, 1.7$ Hz, 1H), 8.27 (dd, $J = 7.9, 1.6$ Hz, 1H), 7.62 (d, $J = 1.9$ Hz, 1H), 7.49 (dd, $J = 7.9, 4.8$ Hz, 1H), 7.33 (d, $J = 8.6$ Hz, 1H), 7.24 (d, $J = 8.6$ Hz, 1H), 6.46 (d, $J = 1.9$ Hz, 1H), 5.18 (s, 2H), 4.61 – 4.44 (m, 2H), 2.96 – 2.75 (m, 2H), 2.62 (s, 3H); MS (ESI) m/z 391 $[\text{M}+\text{H}]^+$.

[0204] **GBT839** Preparation of 3-[[2-[2-(2,2,2-trifluoroethyl)pyrazol-3-yl]pyridin-3-yl]methoxy]pyridine-2-carbaldehyde


[0205] GBT839 was prepared according to general method B from 3-hydroxypicolinaldehyde and **INT-2**.



[0206] ^1H NMR (400 MHz, CDCl_3) δ 10.26 (s, 1H), 8.65 (dd, $J = 4.7, 1.5$ Hz, 1H), 8.38 (dd, $J = 4.4, 1.0$ Hz, 1H), 8.19 (dd, $J = 7.9, 1.0$ Hz, 1H), 7.61 (d, $J = 1.9$ Hz, 1H), 7.43 – 7.33 (m, 2H), 7.21 (d, $J = 8.6$ Hz, 1H), 6.48 (d, $J = 1.9$ Hz, 1H), 5.19 (q, $J = 8.6$ Hz, 2H), 5.15 (s, 2H); MS (ESI) m/z 363.1 $[\text{M}+\text{H}]^+$.

[0207] **GBT840** Preparation of 3-[[2-[2-(3,3,3-trifluoropropyl)pyrazol-3-yl]pyridin-3-yl]methoxy]pyridine-2-carbaldehyde


[0208] GBT839 was prepared according to general method B from 3-hydroxypicolinaldehyde and **INT-3**.



[0209] ^1H NMR (400 MHz, CDCl_3) δ 10.24 (s, 1H), 8.66 (dd, $J = 4.7, 1.6$ Hz, 1H), 8.39 (dd, $J = 4.5, 1.1$ Hz, 1H), 8.21 (dd, $J = 7.9, 1.6$ Hz, 1H), 7.53 (d, $J = 1.9$ Hz, 1H), 7.44 – 7.37 (m, 2H), 7.26 (d, $J = 8.5$ Hz, 1H), 6.37 (d, $J = 1.9$ Hz, 1H), 5.13 (s, 2H), 4.49 – 4.40 (m, 2H), 2.87 – 2.64 (m, 2H); MS (ESI) m/z 377.1 $[\text{M}+\text{H}]^+$.

[0210] **GBT841** Preparation of 3-chloro-5-[[2-(2-propan-2-ylpyrazol-3-yl)pyridin-3-yl]methoxy]pyridine-4-carbaldehyde.


[0211] GBT841 was prepared according to general method B from 3-chloro-5-hydroxyisonicotinaldehyde and INT-4.



[0212] ^1H NMR (400 MHz, CDCl_3) δ 10.51 (s, 1H), 8.77 (dd, $J = 4.7, 1.6$ Hz, 1H), 8.41 (s, 1H), 8.28 (s, 1H), 8.13 (dd, $J = 7.9, 1.5$ Hz, 1H), 7.63 (d, $J = 1.8$ Hz, 1H), 7.47 (dd, $J = 7.9, 4.8$ Hz, 1H), 6.37 (d, $J = 1.8$ Hz, 1H), 5.23 (s, 2H), 4.66 (sep, $J = 6.6$ Hz, 1H), 1.49 (d, $J = 6.6$ Hz, 6H); MS (ESI) m/z 357 $[\text{M}+\text{H}]^+$.


[0213] **GBT844** Preparation of 3-chloro-5-[[2-[2-(2,2,2-trifluoroethyl)pyrazol-3-yl]pyridin-3-yl]methoxy]pyridine-4-carbaldehyde.

[0214] GBT844 was prepared according to general method B from 3-chloro-5-hydroxyisonicotinaldehyde and INT-2.




[0215] ^1H NMR (400 MHz, CDCl_3) δ 10.43 (s, 1H), 8.67 (dd, $J = 4.7, 1.5$ Hz, 1H), 8.35 (s, 1H), 8.26 (s, 1H), 8.06 (dd, $J = 7.9, 1.3$ Hz, 1H), 7.61 (d, $J = 1.9$ Hz, 1H), 7.40 (dd, $J = 7.9, 4.8$ Hz, 1H), 6.47 (d, $J = 1.9$ Hz, 1H), 5.21 – 5.10 (m, 4H); MS (ESI) m/z 397 $[\text{M}+\text{H}]^+$.


[0216] **GBT860** Preparation of *tert*-butyl 4-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)-5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydropyridine-1(2H)-carboxylate




[0217] **Step 1:** To a solution of 1-*tert*-butyl 4-ethyl 3-oxopiperidine-1,4-dicarboxylate (2.0g, 7.37 mmol) in DCM (45 mL) was added DIPEA (1.54 mL, 8.84 mmol) and TiF_2O (1.36 mL, 8.11 mmol) at -78°C , then the temperature was warmed up to room temperature and the solution was stirred at RT for 1.5 h, the mixture was diluted with DCM (100 mL), organic layer was washed with Sat. NaHCO_3 , brine, dried and concentrated to give 1-(*tert*-butyl) 4-ethyl 5-(((trifluoromethyl)sulfonyl)oxy)-3,6-dihydropyridine-1,4(2H)-dicarboxylate, which was used for next step without purification.




[0218] Step 2: To a solution of 1-tert-butyl 4-ethyl 3-(((trifluoromethyl)sulfonyl)oxy)-5,6-dihydropyridine-1,4(2H)-dicarboxylate (1.49 g, 3.7 mmol) and (1-isopropyl-1H-pyrazol-5-yl)boronic acid (0.57 g, 3.7 mmol) in dioxane (10 mL) was added Pd(dppf)Cl₂ (0.27 g, 0.37 mmol) and a solution of sodium carbonate (1.18 g, 11.10) in water (3 mL), the mixture was degassed with N₂ for 5 min, and was heated at 100 °C for 15 h, after cooling to room temperature the mixture was diluted with EtOAc and washed with Sat. NaHCO₃ and brine, organic layer was combined, dried and concentrated to give crude product, which was purified by column chromatography (Hexanes/EtOAc=3:1) to give desired product 830 mg (62%).



[0219] Step 3: To a solution of 1-(tert-butyl) 4-ethyl 5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydropyridine-1,4(2H)-dicarboxylate (450 mg, 1.24 mmol) in THF (6 mL) was added LiAlH₄ (1M in THF, 1.49 mL, 1.49 mmol) at -20 °C, the reaction was stirred at -20 °C for 30 min, and was quenched with Sat. NH₄Cl, the aqueous layer was extracted with EtOAc, the combined organics were washed with brine, dried and concentrated to give crude oil, which was purified by column (Hexanes/EtOAc= 100:0 to 40:60) to give tert-butyl 4-(hydroxymethyl)-5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydropyridine-1(2H)-carboxylate (370 mg, 91%).




[0220] Step 4: To a solution of give tert-butyl 4-(hydroxymethyl)-5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydropyridine-1(2H)-carboxylate (25 mg, 0.08 mmol) in DCM (1 mL) was added triphenylphosphine bromine adduct (40 mg, 0.09 mmol) at room temperature, after stirring for 30 min, it was diluted with DCM, washed with Sat. NaHCO₃, brine, dried and concentrated to give crude product, which was purified by column to give tert-butyl 4-(bromomethyl)-5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydropyridine-1(2H)-carboxylate (18 mg).




[0221] Step 5: To a solution of tert-butyl 4-(bromomethyl)-5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydropyridine-1(2H)-carboxylate (18 mg, 0.05 mmol) and 5-hydroxy-2-methoxyisonicotinaldehyde (10 mg, 0.06 mmol) in DMF (1 mL) was added K_2CO_3 (14 mg, 0.1 mmol). After stirred at room temperature for 1 h, it was diluted with water and EtOAc, organic layer was separated, and the aqueous layer was extracted with EtOAc, organic layer was combined, washed with brine, dried and concentrated to give crude product, which was purified by column (Hexanes/EtOAc=2:1) to give tert-butyl 4-((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)-3-(1-isopropyl-1H-pyrazol-5-yl)-5,6-dihydropyridine-1(2H)-carboxylate (7.2 mg). ^1H NMR (400 MHz, $CDCl_3$) (ppm) 10.39 (s, 1H), 7.79 (s, 1H), 7.56 (d, $J=1.6$ Hz, 1H), 7.05 (s, 1H), 6.11 (d, $J=1.6$ Hz, 1H), 4.40 (s, 2H), 4.38 (m, 1H), 4.01 (s, 2H), 3.88 (s, 3H), 3.66 (bs, 2H), 2.46 (bs, 2H), 1.48 (s, 9H), 1.43 (d, 6.4 Hz, 6H). MS (ESI) m/z 457.3 $[M+H]^+$.

[0222] GBT861 Preparation of 2-hydroxy-6-((5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydro-2H-pyran-4-yl)methoxy)benzaldehyde




[0223] Step 1: To a solution of ethyl 3-oxotetrahydro-2H-pyran-4-carboxylate (1.0 g, 5.81 mmol) in DCM (30 mL) was added DIPEA (1.22 mL, 6.97 mmol) and Tf_2O (1.08 mL, 6.39


mmol) at -78 °C, then it was warmed up to room temperature and stirred at room temperature for 2 h, the solution was diluted with DCM, washed with Sat. NaHCO₃, brine, dried and concentrated to give ethyl 5-(((trifluoromethyl)sulfonyl)oxy)-3,6-dihydro-2H-pyran-4-carboxylate as crude product (2 g).




[0224] **Step 2:** To a solution of ethyl 5-(((trifluoromethyl)sulfonyl)oxy)-3,6-dihydro-2H-pyran-4-carboxylate (crude from step 1) and 1-isopropyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (1.37 g, 5.82 mmol) in dioxane (20 ml) was added Pd(dppf)Cl₂ (430 mg, 0.58 mmol) and Na₂CO₃ (1.85 g, 17.46 mmol) in water (6 mL), the mixture was degassed with N₂ for 5 min, and was heated at 100 °C for 15 h, after cooling to room temperature the mixture was diluted with EtOAc and washed with Sat. NaHCO₃ and brine, organic layer was combined, dried and concentrated to give crude product, which was purified by column chromatography (Hexanes/EtOAc=3:1) to give ethyl 5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydro-2H-pyran-4-carboxylate (850 mg).



[0225] **Step 3:** To a solution of ethyl 5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydro-2H-pyran-4-carboxylate (600 mg, 2.27 mmol) in THF (10 mL) was added LiAlH₄ (1M in THF, 2.72 mL, 2.72 mmol) at -20 °C, the reaction was stirred at -20 °C for 30 min, and was quenched with Sat. NH₄Cl, the aqueous layer was extracted with EtOAc, the combined organics were washed with brine, dried and concentrated to give crude oil, which was purified by column (Hexanes/EtOAc= 100:0 to 20:80) to give (5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydro-2H-pyran-4-yl)methanol (500 mg).




[0226] **Step 4:** To a solution of 5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydro-2H-pyran-4-yl)methanol (300 mg, 1.35 mmol) in DCM (5 mL) was added dibromotriphenylphosphorane (630 mg, 1.35 mmol) at room temperature, after stirring for 30 min, it was diluted with DCM, organic layer was washed with Sat. NaHCO_3 , brine, dried and concentrated to give crude product, which was purified by column (Hexanes/ EtOAc = 4:1) to give 5-(4-(bromomethyl)-5,6-dihydro-2H-pyran-3-yl)-1-isopropyl-1H-pyrazole (360 mg).




[0227] **Step 5:** To a solution of 5-(4-(bromomethyl)-5,6-dihydro-2H-pyran-3-yl)-1-isopropyl-1H-pyrazole (110 mg, 0.38 mmol) and 2,6-dihydroxybenzaldehyde (100 mg, 0.76 mmol) in DMF (6 mL) was added K_2CO_3 (110 mg, 0.76 mmol). After stirred at room temperature for 1 h, it was diluted with water and EtOAc , organic layer was separated, and the aqueous layer was extracted with EtOAc . Organic layer was combined, washed with brine, dried and concentrated to give crude product, which was purified by column (Hexanes/ EtOAc = 1:1) to give 2-hydroxy-6-((5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydro-2H-pyran-4-yl)methoxy)benzaldehyde (90 mg). ^1H NMR (400 MHz, CDCl_3) δ (ppm) 11.89 (s, 1H), 10.33 (s, 1H), 7.53 (d, J = 1.6 Hz, 1H), 7.33 (t, J = 8.8 Hz, 1H), 6.51 (d, J = 8.8 Hz, 1H), 6.16 (d, J = 8.0 Hz, 1H), 6.08 (d, J = 2.0 Hz, 1H), 4.40 (dd, J = 12.8, 6.4 Hz, 1H), 4.35 (s, 2H), 4.18 (s, 2H), 3.97 (t, J = 5.2 Hz, 2H), 2.44 (s, 2H), 1.40 (d, J = 6.4 Hz, 6H); MS (ESI) m/z 343.3 $[\text{M}+\text{H}]^+$.


[0228] **GBT863** Preparation of 2-methoxy-5-[[5-(2-propan-2-ylpyrazol-3-yl)-3,6-dihydro-2H-pyran-4-yl]methoxy]pyridine-4-carbaldehyde



[0229] To a solution of 5-(4-(bromomethyl)-5,6-dihydro-2H-pyran-3-yl)-1-isopropyl-1H-pyrazole (50 mg, 0.19 mmol) (see the synthesis of **GBT861**) and 5-hydroxy-2-methoxyisonicotinaldehyde (30 mg, 0.23 mmol) in DMF (1 mL) was added K_2CO_3 (50 mg, 0.38 mmol). After stirred at room temperature for 3 h, it was diluted with water and EtOAc, organic layer was separated, and the aqueous layer was extracted with EtOAc, organic layer was combined, washed with brine, dried and concentrated to give crude product, which was purified by column (Hexanes/EtOAc=1:1) to give 5-((5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydro-2H-pyran-4-yl)methoxy)-2-methoxyisonicotinaldehyde (26 mg). ^1H NMR (400 MHz, $CDCl_3$) δ (ppm) 10.40 (s, 1H), 7.81 (s, 1H), 7.54 (d, $J=1.6$ Hz, 1H), 7.05 (s, 1H), 6.08 (d, $J=1.6$ Hz, 1H), 4.42 (s, 2H), 4.40 (m, 1H), 4.19 (s, 2H), 3.98 (t, $J=5.6$ Hz, 2H), 3.88 (s, 3H), 2.47 (s, 2H), 1.41 (d, $J=6.8$ Hz, 6H); MS (ESI) m/z 358.4 $[M+H]^+$.


[0230] **GBT864** Preparation of 6-methyl-3-[[5-(2-propan-2-ylpyrazol-3-yl)-3,6-dihydro-2H-pyran-4-yl]methoxy]pyridine-2-carbaldehyde






[0231] To a solution of 5-(4-(bromomethyl)-5,6-dihydro-2H-pyran-3-yl)-1-isopropyl-1H-pyrazole (50 mg, 0.19 mmol) (see the synthesis of **GBT861**) and 3-hydroxy-6-methylpicolinaldehyde (30 mg, 0.24 mmol) in DMF (1 mL) was added K_2CO_3 (50 mg, 0.38 mmol). After stirred at room temperature for 3 h, it was diluted with water and EtOAc, organic layer was separated, and the aqueous layer was extracted with EtOAc, organic layer was combined, washed with brine, dried and concentrated to give crude product, which was purified by column (Hexanes/EtOAc=40:60) to give 5-((5-(1-isopropyl-1H-pyrazol-5-yl)-3,6-dihydro-2H-pyran-4-yl)methoxy)-2-methoxyisonicotinaldehyde (37 mg). ^1H NMR (400 MHz, $CDCl_3$) δ (ppm) 10.30 (s, 1H), 7.54 (d, $J=1.6$ Hz, 1H), 7.24 (d, $J=8.4$ Hz, 1H), 7.09 (d, $J=9.2$ Hz, 1H), 6.08 (d, $J=2.0$ Hz, 1H), 4.42 (m, 1H), 4.38 (s, 2H), 4.18 (s, 2H), 3.98 (t, $J=5.6$ Hz, 2H), 2.56 (s, 3H), 2.51 (s, 2H), 1.39 (d, $J=6.4$ Hz, 6H); MS (ESI) m/z 342.4 $[M+H]^+$.

[0232] **GBT867** Preparation of 2-hydroxy-6-[(5-phenyl-3,6-dihydro-2H-pyran-4-yl)methoxy]benzaldehyde




[0233] **Step 1:** To a solution of ethyl 5-(((trifluoromethyl)sulfonyl)oxy)-3,6-dihydro-2H-pyran-4-carboxylate (1.77 g, 5.81 mmol) and phenylboronic acid (1.42 g, 11.62 mmol) in


dioxane (15 ml) was added Pd(dppf)Cl₂ (430 mg, 0.58 mmol) and Na₂CO₃ (1.85 g, 17.46 mmol) in water (4.5 mL), the mixture was degased with N₂ for 5 min, and was heated at 100 °C for 15 h, after cooling to room temperature the mixture was diluted with EtOAc and washed with Sat. NaHCO₃ and brine, organic layer was combined, dried and concentrated to give crude product, which was purified by column chromatography (Hexanes/EtOAc=4:1) to give ethyl 5-phenyl-3,6-dihydro-2H-pyran-4-carboxylate (1.05 g, 78%).



[0234] **Step 2:** To a solution of ethyl 5-phenyl-3,6-dihydro-2H-pyran-4-carboxylate (1.05 g, 4.52 mmol) in THF (20 mL) was added LiAlH₄ (1M in THF, 5.42 mL, 5.42 mmol) at -20 °C, the reaction was stirred at -20 °C for 30 min, and was quenched with Sat. NH₄Cl, the aqueous layer was extracted with EtOAc, the combined organics were washed with brine, dried and concentrated to give crude oil, which was purified by column (Hexanes/EtOAc= 100:0 to 35:65) to give (5-phenyl-3,6-dihydro-2H-pyran-4-yl)methanol (720 mg).




[0235] **Step 3:** To a solution of (5-phenyl-3,6-dihydro-2H-pyran-4-yl)methanol (360 mg, 1.89 mmol) in DCM (6 mL) was added dibromotriphenylphosphorane (880 mg, 2.08 mmol) at room temperature, after stirring for 30 min, it was diluted with DCM, organic layer was washed with Sat. NaHCO₃, brine, dried and concentrated to give crude product, which was purified by column (Hexanes/EtOAc= 9:1) to give 4-(bromomethyl)-5-phenyl-3,6-dihydro-2H-pyran (380 mg).



[0236] Step 4: To a solution of 4-(bromomethyl)-5-phenyl-3,6-dihydro-2H-pyran (110 mg, 0.45 mmol) and 2,6-dihydroxybenzaldehyde (120 mg, 0.90 mmol) in DMF (3 mL) was added K_2CO_3 (120 mg, 0.90 mmol). After stirred at room temperature for 1 h, it was diluted with water and EtOAc, organic layer was separated, and the aqueous layer was extracted with EtOAc. Organic layer was combined, washed with brine, dried and concentrated to give crude product, which was purified by column (Hexanes/EtOAc=3:1) to give 2-hydroxy-6-((5-phenyl-3,6-dihydro-2H-pyran-4-yl)methoxy)benzaldehyde (120 mg). ^1H NMR (400 MHz, $CDCl_3$) δ (ppm) 11.92 (s, 1H), 10.36 (s, 1H), 7.35 (m, 4H), 7.18 (m, 2H), 6.49 (d, $J=8.0$ Hz, 1H), 6.13 (d, $J=8.0$ Hz, 1H), 4.48 (s, 2H), 4.32 (s, 2H), 3.95 (t, $J=5.6$ Hz, 2H), 2.41 (m, 2H); MS (ESI) m/z 309.


[0237] GBT868 Preparation of 3-methoxy-5-[[2-[2-(2,2,2-trifluoroethyl)pyrazol-3-yl]pyridin-3-yl]methoxy]pyridine-4-carbaldehyde



[0238] To a solution of 3-hydroxy-5-methoxyisonicotinaldehyde (0.13 g, 0.88 mmol) in DMF was added 3-(chloromethyl)-2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridine (0.24 g, 0.88 mmol) (**INT-2**) and potassium carbonate (0.49 g, 3.52 mmol) and the reaction mixture was heated (60 °C). After 3 hours, the reaction mixture was filtered through a plug of silica (MeOH/ CH_2Cl_2 , 0-20%). Purification of the resulting residue by Prep-HPLC, provided 2-methoxy-6-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde (12 mg, 5% yield). ^1H NMR (400 MHz, Chloroform- d) δ 10.54 (s, 1H), 8.71 (dd, $J = 5.0, 1.8$ Hz, 1H), 8.23 (s, 1H), 8.21 (ddd, $J = 7.9, 1.7, 0.7$ Hz, 1H), 8.10 (s, 1H), 7.67 (dd, $J = 1.9, 0.5$

Hz, 1H), 7.46 (dd, $J = 8.0, 4.5$ Hz, 1H), 7.26 (d, $J = 0.5$ Hz, 3H), 6.56 (dd, $J = 1.9, 0.5$ Hz, 1H), 5.23 (s, 2H), 5.28 – 5.15 (m, 2H), 4.04 (s, 3H); MS (ESI) m/z 393 $[M+H]^+$.

[0239] GBT870 Preparation of 2-methoxy-5-[[2-(2-methoxyphenyl)pyridin-3-yl]methoxy]pyridine-4-carbaldehyde




[0240] Step 1: Into a 50-mL round-bottom flask, was placed a solution of (2-chloropyridin-3-yl)methanol (500 mg, 3.48 mmol, 1.00 equiv) in a solvent mixture of dioxane and H_2O (10/10 mL). (2-Methoxyphenyl)boronic acid (532 mg, 3.50 mmol, 1.20 equiv), sodium bicarbonate (882 mg, 10.50 mmol, 3.00 equiv), and $Pd(dppf)Cl_2$ (286 mg, 0.39 mmol, 0.10 equiv) were added to the reaction mixture. The resulting solution was stirred for 2 h at $100^\circ C$, and then it was diluted with 100 mL of H_2O . The resulting solution was extracted with 2x100 mL of ethyl acetate, and the combined organic layers were concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:5) as eluent to furnish 650 mg (87%) of [2-(2-methoxyphenyl)pyridin-3-yl]methanol as a yellow solid.

[0241] Step 2: Into a 50-mL round-bottom flask, was placed a solution of [2-(2-methoxyphenyl)pyridin-3-yl]methanol (600 mg, 2.79 mmol, 1.00 equiv) in thionyl chloride (10 mL). The resulting solution was heated to reflux for 2 hr, and then it was concentrated under vacuum. This resulted in 600 mg (92%) of 3-(chloromethyl)-2-(2-methoxyphenyl)pyridine as a yellow solid.

[0242] Step 3: Into a 100-mL round-bottom flask, was placed a solution of 3-(chloromethyl)-2-(2-methoxyphenyl)pyridine (306 mg, 1.31 mmol, 1.00 equiv) in CH_3CN (20 mL). 5-Hydroxy-2-methoxypyridine-4-carbaldehyde (200 mg, 1.31 mmol, 1.00 equiv), potassium carbonate (364 mg, 2.63 mmol, 2.00 equiv), and KI (44 mg, 0.27 mmol, 0.20 equiv) were added to the reaction mixture. The resulting solution was stirred for 5 h at $60^\circ C$, and then it was concentrated under vacuum. The crude product (200 mg) was purified by

Prep-HPLC with the following conditions (Prep-HPLC-010): Column, SunFire Prep C18 OBD Column, 5µm, 19*150mm.; mobile phase, water with 0.1% HCOOH and MeCN (10.0% MeCN up to 40.0% in 10 min, up to 95.0% in 2 min, down to 10.0% in 2 min); Detector, Waters2545 UV Detector 254&220nm. This resulted in 65 mg (9%) of 2-methoxy-5-[[2-(2-methoxyphenyl)pyridin-3-yl]methoxy]pyridine-4-carbaldehyde bis(trifluoroacetic acid) as a yellow solid. The compound exhibited a melting point of 105-107°C. ¹H NMR(300MHz, CDCl₃) δ 10.32 (s, 1H), 8.69 (s, 1H), 7.93 (m, 2H), 7.36(m, 3H), 6.99 (m, 3H), 5.35 (s, 2H), 3.86(m, 6H); MS (ESI) m/z 351 [M+H]⁺.

[0243] GBT871 Preparation of 2-methoxy-5-[[2-(3-methoxyphenyl)pyridin-3-yl]methoxy]pyridine-4-carbaldehyde



[0244] Step 1: Into a 50-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of (3-methoxyphenyl)boronic acid (1.6 g, 10.53 mmol, 1.20 equiv), (2-chloropyridin-3-yl)methanol (1 g, 6.97 mmol, 1.00 equiv), sodium bicarbonate (1.7 g, 20.24 mmol, 3.00 equiv), Pd(dppf)Cl₂ (0.57 g, 0.10 equiv) in a solvent mixture of dioxane (10 mL) and water (10 mL). The resulting solution was stirred for 1.5 h at 100°C, and then it was diluted with 20 mL of H₂O. The resulting solution was extracted with 2x50 mL of ethyl acetate, and the combined organic layers were concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:50-1:1) as eluent to yield 1.3 g (87%) of [2-(3-methoxyphenyl)pyridin-3-yl]methanol as a colorless oil.


[0245] Step 2: Into a 50-mL round-bottom flask, was placed a solution of [2-(3-methoxyphenyl)pyridin-3-yl]methanol (1 g, 4.65 mmol, 1.00 equiv) in thionyl chloride (20 mL). The resulting solution was stirred for 2 h at reflux. The resulting mixture was

concentrated under vacuum to furnish 600 mg (55%) of 3-(chloromethyl)-2-(3-methoxyphenyl)pyridine as a white solid.


[0246] **Step 3:** Into a 100-mL round-bottom flask, was placed a solution of 3-(chloromethyl)-2-(3-methoxyphenyl)pyridine (234 mg, 1.00 mmol, 1.00 equiv), 5-hydroxy-2-methoxypyridine-4-carbaldehyde (153 mg, 1.00 mmol, 1.00 equiv), and potassium carbonate (278 mg, 2.01 mmol, 2.00 equiv) in CH₃CN (30 mL). The resulting solution was stirred for 4 h at 70°C, and then it was concentrated under vacuum. The crude product (200 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-010): Column, SunFire Prep C18 OBD Column, 5µm, 19*150mm; mobile phase, water with 0.1% TFA and MeCN (20% MeCN up to 40% in 10 min, up to 95% in 2 min, down to 20% in 1 min); Detector, Waters 2545 UV Detector 254&220nm. This resulted in 100.8 mg (17%) of 2-methoxy-5-[[2-(3-methoxyphenyl)pyridin-3-yl]methoxy]pyridine-4-carbaldehyde bis(trifluoroacetic acid) as a yellow solid.

¹H NMR (300 MHz, DMSO-*d*₆) δ 10.01 (s, 1H), 8.65 (m, 1H), 8.39 (s, 1H), 8.10 (m, 2H), 7.57 (d, J = 9 Hz, 2H), 7.42 (m, 1H), 6.97 (m, 3H), 5.33 (s, 2H), 3.80 (m, 6H); MS (ESI) m/z 351 [M+H]⁺.

[0247] **GBT874** Preparation of 2-hydroxy-6-[(1-methyl-5-phenyl-3,6-dihydro-2H-pyridin-4-yl)methoxy]benzaldehyde




[0248] **Step 1:** To a solid of tert-butyl 4-(hydroxymethyl)-3-phenyl-5,6-dihydropyridine-1(2H)-carboxylate (300 mg, 1.04 mmol) in round bottom flask was added 4N HCl in dioxane (6 mL) at room temperature, after stirring for 1 h, the mixture was concentrated and dried under high vacuum to give (5-phenyl-1,2,3,6-tetrahydropyridin-4-yl)methanol as HCl salt.



[0249] Step 2: To a solution of (5-phenyl-1,2,3,6-tetrahydropyridin-4-yl)methanol hydrochloride (230 mg, 1.04 mmol) in ACN (10 mL) was added Et₃N (0.15 mL, 1.04 mmol) followed by formalin (340 mg, 4.16 mmol). After stirred at room temperature for 10 min, it was added Na(OAc)₃BH (440 mg, 2.08 mmol) and was stirred for 30 min, the mixture was concentrated to remove most of the ACN, and the residue was diluted with CHCl₃, organic layer was washed with Sat. NaHCO₃, brine, dried and concentrated to give crude product, which was purified by column (DCM/MeOH=9:1) to give (1-methyl-5-phenyl-1,2,3,6-tetrahydropyridin-4-yl)methanol (140 mg).




[0250] Step 3: To a solution of (1-methyl-5-phenyl-1,2,3,6-tetrahydropyridin-4-yl)methanol (130 mg, 0.64 mmol) in DCM (4 mL) was added SOCl₂ (1.16 mL, 16 mmol) at room temperature, after stirred at room temperature for 30 min, the mixture was concentrated, dried under high vacuum to give 4-(chloromethyl)-1-methyl-5-phenyl-1,2,3,6-tetrahydropyridine as crude HCl salt.



[0251] Step 4: To a suspension of K₂CO₃ (350 mg, 2.56 mmol) and 2,6-dihydroxybenzaldehyde (180 mg, 1.28 mmol) in DMF (3 mL) was added a solution of 4-(chloromethyl)-1-methyl-5-phenyl-1,2,3,6-tetrahydropyridine (140 mg, 0.64 mmol) in DMF (4 mL), the mixture was heated at 50 °C for 3 h, cooled to room temperature, and was diluted

with EtOAc, organic layer was separated and aqueous layer was extracted with EtOAc. EtOAc layers were combined, washed with Sat. NaHCO_3 , brine, dried over Na_2SO_4 , and was concentrated to give crude oil, which was purified by column (Hexane/EtOAc= 1:1 followed by DCM/MeOH= 90:10) to give 2-hydroxy-6-((1-methyl-5-phenyl-1,2,3,6-tetrahydropyridin-4-yl)methoxy)benzaldehyde (55 mg). ^1H NMR (400 MHz, CDCl_3) δ (ppm) 11.92 (s, 1H), 10.35 (s, 1H), 7.34 (m, 5H), 7.19 (dd, $J = 8.4, 8.0$ Hz, 1H), 6.46 (d, $J = 8.4$ Hz, 1H), 6.16 (d, $J = 8.0$ Hz, 1H), 4.45 (s, 2H), 3.20 (s, 2H), 2.68 (t, $J = 5.6$ Hz, 2H), 2.47 (m, 2H), 2.42 (s, 3H); MS (ESI) m/z 324.3 $[\text{M}+\text{H}]^+$.

[0252] GBT875 Preparation of 2-methoxy-5-[[2-(4-methoxyphenyl)pyridin-3-yl]methoxy]pyridine-4-carbaldehyde



[0253] Step 1: Into a 50-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed a solution of (4-methoxyphenyl)boronic acid (1.6 g, 10.53 mmol, 1.20 equiv), (2-chloropyridin-3-yl)methanol (1 g, 6.97 mmol, 1.00 equiv), sodium bicarbonate (1.7 g, 20.24 mmol, 3.00 equiv), $\text{Pd}(\text{dppf})\text{Cl}_2$ (0.57 g, 0.10 equiv) in a solvent mixture of dioxane (10 mL) and water (10 mL). The resulting solution was stirred for 1.5 h at 100°C , and then it was diluted with 20 mL of H_2O . The resulting solution was extracted with 2x50 mL of ethyl acetate, and the combined organic layers were concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:50-1:1) as eluent to furnish 1 g (67%) of [2-(4-methoxyphenyl)pyridin-3-yl]methanol as a colorless oil.


[0254] Step 2: Into a 50-mL round-bottom flask, was placed a solution of [2-(4-methoxyphenyl)pyridin-3-yl]methanol (1 g, 4.65 mmol, 1.00 equiv) in thionyl chloride (20 mL). The resulting solution was stirred for 2 h at reflux. The resulting mixture was

concentrated under vacuum to yield 600 mg (55%) of 3-(chloromethyl)-2-(4-methoxyphenyl)pyridine as a white solid.

[0255] Step 3: Into a 50-mL round-bottom flask, was placed a solution of 3-(chloromethyl)-2-(2-methoxyphenyl)pyridine (234 mg, 1.00 mmol, 1.00 equiv), 5-hydroxy-2-methoxypyridine-4-carbaldehyde (153 mg, 1.00 mmol, 1.00 equiv), and potassium carbonate (278 mg, 2.01 mmol, 2.00 equiv) in CH₃CN (20 mL). The resulting solution was stirred for 4 h at 70°C, and then it was concentrated under vacuum. The crude product (300 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-010): Column, SunFire Prep C18 OBD Column, 5μm, 19*150mm; mobile phase, water with 0.1%TFA and MeCN (20.0% MeCN up to 50.0% in 10 min, up to 95.0% in 2 min, down to 20.0% in 1 min); Detector, Waters2545 UvDetector 254&220nm. This resulted in 265.1 mg (46%) of 2-methoxy-5-[[2-(4-methoxyphenyl)pyridin-3-yl]methoxy]pyridine-4-carbaldehyde; bis(trifluoroacetic acid) as a brown oil.

¹HNMR(300MHz, DMSO-*d*₆) δ 10.08(s, 1H), 8.69 (m, 1H), 8.15(m, 2H), 7.50(m, 1H), 7.37(m, 1H), 7.18(m, 2H), 7.16(m, 1H), 6.99(m, 1H), 5.34(s, 2H), 3.86(s, 3H), 3.77(s, 3H); MS (ESI) m/z 351 [M+H]⁺.

[0256] GBT877 Preparation of 5-[[2-(2-chlorophenyl)pyridin-3-yl]methoxy]-2-methoxypyridine-4-carbaldehyde



[0257] Step 1: Into a 50-mL round-bottom flask, was placed a solution of (2-chlorophenyl)boronic acid (1.6 g, 10.23 mmol, 1.20 equiv), (2-chloropyridin-3-yl)methanol (1 g, 6.97 mmol, 1.00 equiv), Pd(dppf)Cl₂ (570 mg, 0.78 mmol, 0.10 equiv), and sodium bicarbonate (1.7 g, 20.24 mmol, 3.00 equiv) in a solvent mixture of dioxane (10 mL) and water(10 mL). The resulting solution was stirred for 3 h at 70°C, and then it was diluted with


20 mL of H₂O. The resulting solution was extracted with 2x20 mL of dichloromethane, and the combined organic layers were concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:100-1:5) as eluent to furnish 1 g (65%) of [2-(2-chlorophenyl)pyridin-3-yl]methanol as a white solid.

[0258] Step 2: Into a 25-mL round-bottom flask, was placed a solution of [2-(2-chlorophenyl)pyridin-3-yl]methanol (1 g, 4.55 mmol, 1.00 equiv) in thionyl chloride (5 mL). The resulting solution was stirred for 1 h at reflux. The resulting mixture was concentrated under vacuum to yield 1 g (92%) of 3-(chloromethyl)-2-(2-chlorophenyl)pyridine as a white solid.


[0259] Step 3: Into a 50-mL round-bottom flask, was placed a solution of 3-(chloromethyl)-2-(2-chlorophenyl)pyridine (309 mg, 1.30 mmol, 1.00 equiv), 5-hydroxy-2-methoxypyridine-4-carbaldehyde (200 mg, 1.31 mmol, 1.00 equiv), and potassium carbonate (361 mg, 2.61 mmol, 1.50 equiv) in CH₃CN (20 mL). The resulting solution was stirred for 4 h at 70°C, and then it was concentrated under vacuum. The residue was purified by prep-HPLC. This resulted in 86.2 mg (11%) of 5-[[2-(2-chlorophenyl)pyridin-3-yl]methoxy]-2-methoxypyridine-4-carbaldehyde; bis(trifluoroacetic acid) as a brown oil.

¹HNMR(300MHz, DMSO-*d*₆) δ 10.06(s, 1H), 8.69 (m, 1H), 8.19(m, 1H), 8.05(s, 1H), 7.56(m, 2H), 7.41(m, 3H), 6.92(s, 1H), 5.14(m, 2H), 3.81(s, 3H); MS (ESI) *m/z* 355 [M+H]⁺.


[0260] GBT878 Preparation of 2-[(1-acetyl-5-phenyl-3,6-dihydro-2H-pyridin-4-yl)methoxy]-6-hydroxybenzaldehyde



[0261] **Step 1:** To a solution of (5-phenyl-1,2,3,6-tetrahydropyridin-4-yl)methanol hydrochloride (90 mg, 0.38 mmol) in DCM (2 mL) at 0 °C was added Et₃N (0.11 mL, 0.76 mmol) and a solution of Ac₂O (0.04 mL, 0.38 mmol) in DCM (0.4 mL), after stirred for 15 min, it was diluted with Sat. NH₄Cl and EtOAc, organic layer was separated and the aqueous layer was further extracted with EtOAc, organic layers were combined, washed with Sat. NaHCO₃, brine, dried over Na₂SO₄, and was concentrated to give 1-(4-(hydroxymethyl)-5-phenyl-3,6-dihydropyridin-1(2H)-yl)ethan-1-one as crude product (95 mg).




[0262] **Step 2:** To a solution of 1-(4-(hydroxymethyl)-3-phenyl-5,6-dihydropyridin-1(2H)-yl)ethanone (86 mg, 0.37 mmol) in DCM (2 mL) was added SOCl₂ (0.67 mL, 9.25 mmol). After stirred at RT for 15 min, the mixture was concentrated and was diluted with Sat. NaHCO₃ and EtOAc, organic layer was separated and the aqueous layer was extracted with EtOAc, organic layer were combined, washed with brine, dried and concentrated to give crude oil, which was purified by column (Hexanes/EtOAc=100:0 to 25:75) to give 1-(4-(chloromethyl)-5-phenyl-3,6-dihydropyridin-1(2H)-yl)ethan-1-one (35 mg).




[0263] **Step 3:** To a suspension of K₂CO₃ (40 mg, 0.28 mmol) and 2,6-dihydroxybenzaldehyde (40 mg, 0.28 mmol) in DMF (1 mL) was added a solution of 1-(4-(chloromethyl)-5-phenyl-3,6-dihydropyridin-1(2H)-yl)ethan-1-one (35 mg, 0.14 mmol) in DMF (1 mL), the mixture was heated at 50 °C for 3 h, cooled to room temperature, and was diluted with EtOAc, organic layer was separated and aqueous layer was extracted with EtOAc. EtOAc layers were combined, washed with Sat. NaHCO₃, brine, dried over Na₂SO₄, and was concentrated to give crude oil, which was purified by column (DCM/MeOH= 90:10) to give 2-((1-acetyl-5-phenyl-1,2,3,6-tetrahydropyridin-4-yl)methoxy)-6-

hydroxybenzaldehyde (17 mg). ^1H NMR (400 MHz, CDCl_3 , NMR shows rotamer exist, only one set of signal was reported) δ (ppm) 11.93 (s, 1H), 10.36 (s, 1H), 7.34 (m, 5H), 7.22 (m, 1H), 6.49 (d, $J = 8.8$ Hz, 1H), 6.10 (d, $J = 8.8$ Hz, 1H), 4.47 (s, 2H), 4.32 (s, 2H), 3.68 (t, $J = 6.0$ Hz, 2H), 2.47 (m, 2H), 2.18 (s, 3H); MS (ESI) m/z 352.5 $[\text{M}+\text{H}]^+$.

[0264] GBT881 Preparation of 2-[(1-acetyl-4-phenyl-3,6-dihydro-2H-pyridin-5-yl)methoxy]-6-hydroxybenzaldehyde




[0265] Step 1: To a solution of 1-tert-butyl 3-methyl 4-oxopiperidine-1,3-dicarboxylate (2.50 g, 9.72 mmol) in DCM (50 mL) was added DIPEA (2.03 mL, 11.66 mmol) and TiF_2O (1.80 mL, 10.69 mmol) at -78°C , and then it was warmed up to room temperature and stirred further for 2 h, the solution was diluted with DCM and the organic layer was washed with Sat. NaHCO_3 , dried and concentrated to give 1-tert-butyl 3-methyl 4-(((trifluoromethyl)sulfonyl)oxy)-5,6-dihydropyridine-1,3-dicarboxylate as crude product (4.4 g).




[0266] Step 2: To a solution of 1-tert-butyl 3-methyl 4-(((trifluoromethyl)sulfonyl)oxy)-5,6-dihydropyridine-1,3-dicarboxylate (1.95 g, 5 mmol) and phenylboronic acid (1.22 g, 10 mmol) in Dioxane (20 mL) was added $\text{Pd}(\text{dppf})\text{Cl}_2$ and a solution of Na_2CO_3 (3.18 g, 30 mmol) in water (6 mL), after degassed with N_2 for 5 min, the reaction was heated at 100°C for 15 h, the mixture was cooled to room temperature, diluted with EtOAc, organic layer was washed with water, brine, dried and concentrated to give crude product, which was purified


by column (Hexanes/EtOAc= 3:1) to give 1-tert-butyl 3-methyl 4-phenyl-5,6-dihydropyridine-1,3(2H)-dicarboxylate (740 mg).




[0267] Step 3: To a solution of 1-tert-butyl 3-methyl 4-phenyl-5,6-dihydropyridine-1,3(2H)-dicarboxylate (740 mg, 2.33 mmol) in THF (7.2 mL) was added 1M LiAlH₄ in THF (2.80 mL, 2.80 mmol) at -20 °C dropwise, after stirring at -20 °C for 30 min, it was quenched with Sat. NH₄Cl, the mixture was extracted with EtOAc. Organic layers were combined, washed with brine, dried and concentrated to give crude product, which was purified by column (Hexanes/EtOAc= 60:40) to give tert-butyl 5-(hydroxymethyl)-4-phenyl-3,6-dihydropyridine-1(2H)-carboxylate (512 mg).




[0268] Step 4: To tert-butyl 3-(hydroxymethyl)-4-phenyl-5,6-dihydropyridine-1(2H)-carboxylate (510 mg, 1.76 mmol) was added 4N HCl in Dioxane (3 ml), after stirring at room temperature for 1h, it was concentrated to give (4-phenyl-1,2,5,6-tetrahydropyridin-3-yl)methanol as HCl salt.



[0269] Step 5: To a solution of (4-phenyl-1,2,5,6-tetrahydropyridin-3-yl)methanol hydrochloride (110 mg, 0.49 mmol) in DCM (2 mL) was added DIPEA (0.17 mL, 0.98 mmol) and Ac₂O (0.05 g, 0.49 mmol), 15 min later, it was diluted with water and extracted with DCM. Organic layers were combined, dried and concentrated, the resulting crude oil was purified by column (EtOAc followed by DCM/MeOH =9:1) to give 1-(5-(hydroxymethyl)-4-phenyl-3,6-dihydropyridin-1(2H)-yl)ethan-1-one (88 mg).





[0270] Step 6: To a solution of 1-(3-(hydroxymethyl)-4-phenyl-5,6-dihydropyridin-1(2H)-yl)ethanone (88 mg, 0.38 mmol) in DCM (2 mL) was added SOCl_2 (0.67 mL, 9.50 mmol) at 0 °C. After stirring at 0 °C for 15 min, the solution was concentrated to remove SOCl_2 , dried under high vacuum to give 1-(5-(chloromethyl)-4-phenyl-3,6-dihydropyridin-1(2H)-yl)ethanone as crude product.



[0271] Step 7: To a solution of 1-(3-(chloromethyl)-4-phenyl-5,6-dihydropyridin-1(2H)-yl)ethanone (100 mg, 0.40 mmol) and 2,6-dihydroxybenzaldehyde (110 mg, 0.80 mmol) in DMF (2.5 mL) was added K_2CO_3 (170 mg, 1.20 mmol), after heated at 50 degree for 2 h, the reaction was diluted with EtOAc, organic layer was separated and aqueous layer was extracted with EtOAc. EtOAc layers were combined, washed with Sat. NaHCO_3 , brine, dried over Na_2SO_4 , and was concentrated to give crude oil, which was purified by preparative HPLC (eluted with ACN/ H_2O) to give 2-((1-acetyl-4-phenyl-1,2,5,6-tetrahydropyridin-3-yl)methoxy)-6-hydroxybenzaldehyde (26 mg). ^1H NMR (400 MHz, CDCl_3 , NMR shows rotamers exist, only one set of signal was reported) δ (ppm) 11.97 (s, 1H), 10.34 (s, 1H), 7.34 (m, 4H), 7.17 (m, 2H), 6.49 (d, $J = 8.0$ Hz, 1H), 6.11 (d, $J = 8.8$ Hz, 1H), 4.48 (s, 2H), 4.33 (s, 2H), 3.69 (t, $J = 6.0$ Hz, 2H), 2.55 (m, 2H), 2.18 (s, 3H); MS (ESI) m/z 352.3 $[\text{M}+\text{H}]^+$.


[0272] GBT887 Preparation of 2-((2-(1-cyclopentyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-6-hydroxybenzaldehyde






[0273] To a mixture of 3-(chloromethyl)-2-(1-cyclopentyl-1H-pyrazol-5-yl)pyridine hydrochloride (44.7mg, 0.15mmol) and 2,6-dihydroxybenzaldehyde (83mg, 0.6mmol, 4eq) and potassium carbonate (41.5mg, 0.3mmol, 2eq) were added 1mL anhydrous DMF. The mixture was heated to 80 °C for 40mins. The reaction was almost done by LCMS. Solvent was removed at 50 °C on a rotavap. Water 3mL and 0.3mL of formic acid were added to the resulting brown residue, the mixture was sonicated to make sure all carbonate was neutralized. Solvents were then removed at 45 °C on a rotavap. DCM (4x1ml) was added to the yellow residue, the mixture was sonicated and filtered. The filtrate was concentrated to give the crude product as a yellow-light brown film. It contains the product, 2,6-dihydroxybenzaldehyde, and some starting chloride, no bis-alkylation product was observed. The residue was taken up in 2ml DCM, filtered and loaded on a 4g ZAP SiO₂ column. It was purified on Biotage Isolera One system eluted with 5%-100% EtOAc (the product came out around 25% EtOAc, 2nd peak; the 1st peak is dihydroxybenzaldehyde). The product as a yellow film was obtained after removing solvents, the residue was re-dissolved in 0.3mL CH₃CN and to this was added 0.5mL of water. This suspension was freeze-dried and put on a lyophilizer over the weekend. The product was obtained as a light brown film (18.6mg, 34% yield). ¹H NMR (400 MHz, CDCl₃-d) δ 11.94 (s, 1H), 10.37 (s, 1H), 8.75 (dd, *J* = 4.8, 1.7 Hz, 1H), 7.97 (dd, *J* = 8.0, 1.4 Hz, 1H), 7.59 (d, *J* = 2.0 Hz, 1H), 7.42 (dd, *J* = 7.7, 4.8 Hz, 1H), 7.37 (t, *J* = 8.3 Hz, 1H), 6.56 (d, *J* = 8.6 Hz, 1H), 6.35 (d, *J* = 1.9 Hz, 1H), 6.25 (d, *J* = 8.3 Hz, 1H), 5.07 (s, 2H), 4.79 – 4.67 (m, 1H), 2.18 – 1.95 (m, 4H), 1.95 – 1.84 (m, 2H), 1.66 – 1.50 (m, 2H); MS (ESI) *m/z* 364.3 [M+H]⁺.

[0274] **GBT888** Preparation of 2-hydroxy-6-[[[(2S)-1-phenylpyrrolidin-2-yl]methoxy]benzaldehyde






[0275] **Step 1:** To a solution of (S)-pyrrolidin-2-ylmethanol (1.52 g, 15 mmol) and CuI (190 mg, 1mmol) in iPrOH (10 mL) was added (CH₂OH)₂ (1.11 mL, 20 mmol), iodobenzene (2.04 g, 20 mmol) and K₃PO₄ (4.25 g, 20 mmol), after degassed with N₂, the mixture was heated at 88 °C for 15 h. Water and ether was added, organic layer was separated and aqueous layer was further extracted with ether. Organic layers were combined, concentrated and the resulting crude oil was purified by column (hexanes/EtOAc=2:1) to give (S)-(1-phenylpyrrolidin-2-yl)methanol (1.6 g).

[0276] **Step 2:** To a solution of (S)-(1-phenylpyrrolidin-2-yl)methanol (45 mg, 0.23 mmol) and 2,6-dihydroxybenzaldehyde (60 mg, 0.46 mmol) in THF (1 ml) was added PPh₃ (0.12 g, 0.46 mmol), followed by DIAD (90 mg, 0.46 mmol) at room temperature. After stirrer for 10 min, the mixture was concentrated and the residue was purified by column (Hexanes/EtOAc=9:1) to give (S)-2-hydroxy-6-((1-phenylpyrrolidin-2-yl)methoxy)benzaldehyde (14 mg). ¹H NMR (400 MHz, CDCl₃) (ppm) 11.96 (s, 1H), 10.37 (s, 1H), 7.35 (t, J = 8.0 Hz, 1H), 7.25 (m, 2H), 6.73 (m, 3H), 6.53 (d, J = 8.4 Hz, 1H), 6.33 (d, J = 9.2 Hz, 1H), 4.21 (m, 1H), 4.15 (d, J = 3.6 Hz, 1H), 3.83 (t, J = 8.0 Hz, 1H), 3.53 (m, 1H), 3.22 (m, 1H), 2.11 (m, 4H); MS (ESI) m/z 298.4

[0277] **GBT892** Preparation of 5-[[2-(3-chlorophenyl)pyridin-3-yl]methoxy]-2-methoxypyridine-4-carbaldehyde




[0278] **Step 1:** Into a 50-mL round-bottom flask, was placed a solution of (3-chlorophenyl)boronic acid (1.6 g, 10.23 mmol, 1.20 equiv), (2-chloropyridin-3-yl)methanol (1 g, 6.97 mmol, 1.00 equiv), Pd(dppf)Cl₂ (570 mg, 0.78 mmol, 0.10 equiv), and sodium bicarbonate (1.7 g, 20.24 mmol, 3.00 equiv) in a solvent mixture of dioxane (10 mL) and water (10 mL). The resulting solution was stirred for 3 h at 70°C, and then it was diluted with 20 mL of H₂O. The resulting solution was extracted with 2x20 mL of dichloromethane, and the combined organic layers were concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:100-1:5) as eluent to yield 1.2 g (78%) of [2-(3-chlorophenyl)pyridin-3-yl]methanol as a white solid.

[0279] **Step 2:** Into a 50-mL round-bottom flask, was placed a solution of [2-(3-chlorophenyl)pyridin-3-yl]methanol (600 mg, 2.73 mmol, 1.00 equiv) in thionyl chloride (10 mL). The resulting solution was stirred for 1 h at reflux. The resulting mixture was concentrated under vacuum. This resulted in 500 mg (77%) of 3-(chloromethyl)-2-(3-chlorophenyl)pyridine as a white solid.

[0280] **Step 3:** Into a 50-mL round-bottom flask, was placed a solution of 3-(chloromethyl)-2-(3-chlorophenyl)pyridine (309 mg, 1.30 mmol, 1.00 equiv), 5-hydroxy-2-methoxypyridine-4-carbaldehyde (200 mg, 1.31 mmol, 1.00 equiv), and potassium carbonate (361 mg, 2.61 mmol, 2.00 equiv) in CH₃CN (20 mL). The resulting solution was stirred for 4 h at 70°C, and then it was concentrated under vacuum. The crude product (300 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-010): Column, SunFire Prep C18 OBD Column, 5µm, 19*150mm; mobile phase, water with 0.05% TFA and MeCN (20.0% MeCN up to 60.0% in 10 min, up to 95.0% in 2 min, down to 20.0% in 1 min); Detector, Waters 2545 UV Detector 254&220nm. This resulted in 71 mg (9%) of 5-[[2-(3-chlorophenyl)pyridin-3-yl]methoxy]-2-methoxypyridine-4-carbaldehyde; bis(trifluoroacetic acid) as a yellow solid.

¹H NMR (400 MHz, DMSO-*d*₆) δ 10.07(s, 1H), 8.72 (m, 1H), 8.20(m, 2H), 7.79(s, 1H), 7.60(m, 4H), 6.95(m, 1H), 5.21(m, 2H), 3.85(s, 3H); MS (ESI) m/z 355 [M+H]⁺.

[0281] **GBT893** Preparation of 5-[[2-(4-chlorophenyl)pyridin-3-yl]methoxy]-2-methoxypyridine-4-carbaldehyde



[0282] Step 1: Into a 100-mL round-bottom flask, was placed a solution of (4-chlorophenyl)boronic acid (1.6 g, 10.23 mmol, 1.20 equiv), (2-chloropyridin-3-yl)methanol (1 g, 6.97 mmol, 1.00 equiv), Pd(dppf)Cl₂ (570 mg, 0.78 mmol, 0.10 equiv), and sodium bicarbonate (1.7 g, 20.24 mmol, 3.00 equiv) in a solvent mixture of dioxane (10 mL) and water (10 mL). The resulting solution was stirred for 4 h at 70°C, and then it was diluted with 100 mL of H₂O. The resulting solution was extracted with 2x200 mL of dichloromethane, and the combined organic layers were concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:100-1:5) as eluent to yield 1 g (65%) of [2-(4-chlorophenyl)pyridin-3-yl]methanol as a light yellow oil.


[0283] Step 2: Into a 25-mL round-bottom flask, was placed a solution of [2-(4-chlorophenyl)pyridin-3-yl]methanol (1 g, 4.55 mmol, 1.00 equiv) in thionyl chloride (5 mL). The resulting solution was stirred for 1 h at reflux. The resulting mixture was concentrated under vacuum. This resulted in 1 g (92%) of 3-(chloromethyl)-2-(4-chlorophenyl)pyridine as a white solid.

[0284] Step 3: Into a 50-mL round-bottom flask, was placed a solution of 3-(chloromethyl)-2-(4-chlorophenyl)pyridine (309 mg, 1.30 mmol, 1.00 equiv), 5-hydroxy-2-methoxypyridine-4-carbaldehyde (200 mg, 1.31 mmol, 1.00 equiv), and potassium carbonate (361 mg, 2.61 mmol, 2.00 equiv) in CH₃CN (20 mL). The resulting solution was stirred for 4 h at 70°C, and then it was concentrated under vacuum. The crude product (300 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-010): Column, SunFire Prep C18 OBD Column, 5μm, 19*150mm; mobile phase, water with 0.05% TFA and MeCN (20.0% MeCN up to 60.0% in 10 min, up to 95.0% in 2 min, down to 20.0% in 1 min); Detector, Waters 2545 UvDector 254&220nm. This resulted in 148.2 mg (20%) of 5-[[2-(4-chlorophenyl)pyridin-3-yl]methoxy]-2-methoxypyridine-4-carbaldehyde; bis(trifluoroacetic

acid) as a yellow solid.

^1H NMR (300MHz, DMSO- d_6) δ 10.05(s, 1H), 8.69 (m, 1H), 8.16(m, 2H), 7.64(m, 2H), 7.50(m, 3H), 5.32(s, 2H), 3.81(s, 3H); MS (ESI) m/z 355 $[\text{M}+\text{H}]^+$.


GBT903



(*R*)-2-hydroxy-6-((1-phenylpyrrolidin-2-yl)methoxy)benzaldehyde

[0285] **GBT903 - Preparation of (*R*)-2-hydroxy-6-((1-phenylpyrrolidin-2-yl)methoxy)benzaldehyde.** The compound was prepared from (*R*)-pyrrolidin-2-ylmethanol and iodobenzene according to scheme 8, reaction steps 3 and 4. ^1H NMR (400 MHz, Chloroform- d) δ 11.96 (d, J = 0.4 Hz, 1H), 10.37 (s, 1H), 7.37 (td, J = 8.4, 0.4 Hz, 1H), 7.31 – 7.18 (m, 2H), 6.77 – 6.64 (m, 3H), 6.53 (dt, J = 8.5, 0.7 Hz, 1H), 6.33 (dd, J = 8.3, 0.8 Hz, 1H), 4.25 – 4.12 (m, 2H), 3.88 – 3.78 (m, 1H), 3.53 (dt, J = 8.8, 4.4 Hz, 1H), 3.27 – 3.16 (m, 1H), 2.11 (dqt, J = 13.0, 6.4, 2.8 Hz, 4H). MS (M+H) $^+$ found for $\text{C}_{18}\text{H}_{19}\text{NO}_3$: 298.2.

GBT904




2-hydroxy-6-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde


[0286] **GBT904 - Preparation of 2-hydroxy-6-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde.** The compound was prepared from (2-chloropyridin-3-yl)methanol and 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole according to scheme 9, reaction steps 1, 2 and 4. In step 4, 2-hydroxy-6-(methoxymethoxy)benzaldehyde was used; the MOM ether protecting group fell off after the reaction to give the final product. ^1H NMR (400 MHz, Chloroform- d) δ 11.94 (s, 1H), 10.36

(s, 1H), 8.75 (dd, $J = 4.8, 1.7$ Hz, 1H), 7.98 (d, $J = 8.0$ Hz, 1H), 7.54 (d, $J = 1.9$ Hz, 1H), 7.41 (dd, $J = 8.0, 4.0$ Hz, 1H), 7.38 (t, $J = 8.0$ Hz, 1H), 6.58 (dt, $J = 8.5, 0.7$ Hz, 1H), 6.39 (d, $J = 1.9$ Hz, 1H), 6.28 (dd, $J = 8.3, 0.8$ Hz, 1H), 5.12 (s, 2H), 3.95 (s, 3H). MS (M+H) found for $C_{17}H_{15}N_3O_3$: 310.3.

GBT907 and GBT908



(S)-2-hydroxy-6-((1-(2-methoxyphenyl)pyrrolidin-2-yl)methoxy)benzaldehyde




2-hydroxy-6-((1-(2-methoxyphenyl)piperidin-3-yl)oxy)benzaldehyde

[0287] GBT907 and GBT908 - Preparation of (S)-2-hydroxy-6-((1-(2-methoxyphenyl)pyrrolidin-2-yl)methoxy)benzaldehyde (GBT907) and 2-hydroxy-6-((1-(2-methoxyphenyl)piperidin-3-yl)oxy)benzaldehyde (GBT908). The reaction of (S)-pyrrolidin-2-ylmethanol with 2-iodoanisole, and subsequent Mitsunobu reaction, according to scheme 8, gave a mixture of GBT907 and GBT908 in 3:2 ratio, which were separated by reverse-phase prep HPLC.

[0288] GBT907- (S)-2-hydroxy-6-((1-(2-methoxyphenyl)pyrrolidin-2-yl)methoxy)benzaldehyde ^1H NMR (400 MHz, Chloroform- d) δ 11.96 (d, $J = 0.4$ Hz, 1H), 10.37 (d, $J = 0.6$ Hz, 1H), 7.39 (td, $J = 8.4, 0.4$ Hz, 1H), 7.04 – 6.87 (m, 3H), 6.84 (dd, $J = 8.0, 1.4$ Hz, 1H), 6.52 (ddt, $J = 13.7, 8.4, 0.7$ Hz, 2H), 4.66 (tt, $J = 8.2, 3.9$ Hz, 1H), 3.80 (s, 3H), 3.67 – 3.58 (m, 1H), 3.29 (dt, $J = 9.9, 4.3$ Hz, 1H), 2.85 (dd, $J = 11.3, 8.3$ Hz, 1H), 2.82 – 2.74 (m, 1H), 2.20 (dd, $J = 12.4, 4.9$ Hz, 1H), 2.00 (dp, $J = 13.0, 4.6$ Hz, 1H), 1.94 – 1.79 (m, 1H), 1.72 (dddd, $J = 13.0, 10.7, 9.0, 4.3$ Hz, 1H). MS (M+H) $^+$ found for $C_{19}H_{21}NO_4$: 328.3).

[0289] GBT908- 2-hydroxy-6-((1-(2-methoxyphenyl)piperidin-3-yl)oxy)benzaldehyde (40 mg, ^1H NMR (400 MHz, Chloroform- d) δ 11.91 (s, 1H), 10.09 (d, $J = 0.6$ Hz, 1H), 7.32 (d, $J = 8.4$ Hz, 1H), 6.93 – 6.80 (m, 4H), 6.46 (dt, $J = 8.5, 0.7$ Hz, 1H), 6.23 (dd, $J = 8.4, 0.8$ Hz, 1H), 4.49 (tt, $J = 7.2, 4.7$ Hz, 1H), 4.06 (dd, $J = 9.3, 4.4$ Hz, 1H), 3.88 – 3.78 (m, 1H), 3.80 (s, 3H), 3.63 (ddd, $J = 9.1, 7.3, 6.3$ Hz, 1H), 3.21 – 3.11 (m, 1H), 2.35 – 2.22 (m, 1H), 2.09 – 1.86 (m, 3H), MS (M+H) $^+$ found for $C_{19}H_{21}NO_4$: 328.3).


GBT912



2-(((3*S*,4*R*)-1-acetyl-4-(2,5-difluorophenyl)pyrrolidin-3-yl)methoxy)-6-hydroxybenzaldehyde


[0290] GBT912- 2-(((3*S*,4*R*)-1-acetyl-4-(2,5-difluorophenyl)pyrrolidin-3-yl)methoxy)-6-hydroxybenzaldehyde. The compound was prepared from -(2,5-difluorophenyl)-4-(hydroxymethyl)pyrrolidin-1-yl)ethan-1-one and 2,6-dihydroxybenzaldehyde (INT-7) using general method A (Mitsunobu conditions). ¹H NMR (400 MHz, Chloroform-*d*) δ 11.90 (s, 1H), 9.88 (s, 1H), 7.37 (dd, *J* = 8.4, 6.4 Hz, 1H), 7.01 (m, 3H), 6.53 (t, *J* = 8.6 Hz, 1H), 6.27 (dd, *J* = 8.3, 0.8 Hz, 1H), 4.02 (m, 4H), 3.55 (m, 3H), 2.96 (m, 1H), 2.11 (s, 3H). MS (M+H)⁺ found for C₂₀H₁₉F₂NO₄: 376.3.

[0291] Preparation of -(2,5-difluorophenyl)-4-(hydroxymethyl)pyrrolidin-1-yl)ethan-1-one




[0292] To a solution of ((3*S*,4*R*)-4-(2,5-difluorophenyl)pyrrolidin-3-yl)methanol hydrochloride (200 mg, 0.8 mmol) in DCM (2 mL) at 0 °C was added DIPEA (0.3 mL, 1.68 mmol) and Ac₂O (90 mg, 0.84 mmol), after stirred for 30 min, the solution was diluted with DCM, organic layer was washed with Sat. NaHCO₃, brine, dried over MgSO₄ and was concentrated to give 1-((3*R*,4*S*)-3-(2,5-difluorophenyl)-4-(hydroxymethyl)pyrrolidin-1-yl)ethan-1-one as crude product (175 mg).

GBT913 and GBT914



(S)-2-hydroxy-6-((1-(3-methoxyphenyl)pyrrolidin-2-yl)methoxy)benzaldehyde




2-hydroxy-6-((1-(3-methoxyphenyl)piperidin-3-yl)oxy)benzaldehyde

[0293] GBT913 and GBT914 - Preparation of (S)-2-hydroxy-6-((1-(3-methoxyphenyl)pyrrolidin-2-yl)methoxy)benzaldehyde (GBT913) and 2-hydroxy-6-((1-(3-methoxyphenyl)piperidin-3-yl)oxy)benzaldehyde (GBT914). The reaction of (S)-pyrrolidin-2-ylmethanol with 3-iodoanisole, and subsequent Mitsunobu reaction, according to scheme 8, gave a mixture of GBT913 and GBT914 in 5:4 ratio, which were separated by reverse-phase prep HPLC.

[0294] GBT913- (S)-2-hydroxy-6-((1-(3-methoxyphenyl)pyrrolidin-2-yl)methoxy)benzaldehyde ^1H NMR (400 MHz, Chloroform-*d*) δ 11.95 (s, 1H), 10.37 (t, J = 0.5 Hz, 1H), 7.42 – 7.31 (m, 1H), 7.16 (t, J = 8.2 Hz, 1H), 6.53 (dq, J = 8.4, 0.6 Hz, 1H), 6.36 – 6.24 (m, 3H), 6.23 (t, J = 2.4 Hz, 1H), 4.23 – 4.12 (m, 2H), 3.79 (s, 4H), 3.50 (ddd, J = 9.0, 5.6, 3.5 Hz, 1H), 3.27 – 3.16 (m, 1H), 2.17 – 1.98 (m, 4H). MS (M+H)⁺ found for $\text{C}_{19}\text{H}_{21}\text{NO}_4$: 328.3


GBT914 - 2-hydroxy-6-((1-(3-methoxyphenyl)piperidin-3-yl)oxy)benzaldehyde ^1H NMR (400 MHz, Chloroform-*d*) δ 11.94 (d, J = 0.4 Hz, 1H), 10.25 (d, J = 0.6 Hz, 1H), 7.39 (td, J = 8.4, 0.4 Hz, 1H), 7.19 – 7.08 (m, 1H), 6.51 (dt, J = 8.5, 0.7 Hz, 2H), 6.48 – 6.37 (m, 3H), 4.58 (m, 1H), 3.78 (m, 1H), 3.77 (s, 3H), 3.74 – 3.64 (m, 1H), 3.39 (d, J = 5.6 Hz, 1H), 3.17 (dd, J = 12.4, 7.6 Hz, 1H), 3.11 – 3.01 (m, 1H), 2.14 (s, 1H), 2.02 – 1.92 (m, 1H), 1.86 – 1.74 (m, 1H). MS (M+H)⁺ found for $\text{C}_{19}\text{H}_{21}\text{NO}_4$: 328.4).

GBT916




[0295] GBT916 - 2-hydroxy-6-((2-(2-methoxyphenyl)pyridin-3-yl)methoxy)benzaldehyde. The compound was prepared from (2-chloropyridin-3-

yl)methanol and (2-methoxyphenyl)boronic acid according to scheme 9, reaction steps 1, 2 and 4. In step 4, alkylation with 2,6-dihydroxybenzaldehyde, the product of TFA salt was obtained after HPLC purification. ¹HNMR (300MHz, DMSO, ppm): 11.71 (s, 1H), 9.99 (s, 1H), 8.65 (m, 1H), 8.13 (d, J=7.5Hz, 1H), 7.50 (m, 3H), 7.31 (m, 1H), 7.04 (m, 2H), 6.47 (m, 1H) 6.38 (m, d, J=8.4Hz, 1H), 5.00 (s, 2H), 3.73 (s, 3H); MS (ES, *m/z*): 336 [M+1]⁺.

GBT917


[0296] **GBT917 - 2-hydroxy-6-((2'-methoxy-[2,3'-bipyridin]-3-yl)methoxy)benzaldehyde.** The compound was prepared from (2-chloropyridin-3-yl)methanol and (2-methoxypyridin-3-yl)boronic acid according to scheme 9, reaction steps 1, 2 and 4. In step 4, alkylation with 2,6-dihydroxybenzaldehyde, the product TFA salt was obtained after HPLC purification.

¹HNMR (300MHz, CDCl₃, ppm): 11.91 (s, 1H), 10.24 (s, 1H), 8.71 (t, 1H), 8.69 (m, 1H), 7.93 (d, 1H), 7.75 (d, 1H), 7.40 (m, 1H), 7.39 (m, 1H), 7.08 (m, 1H), 6.53 (d, 1H), 6.50 (d, 1H), 5.07 (s, 2H), 3.94 (s, 3H); MS (ES, *m/z*): 337 [M+1]⁺


GBT930

[0297] **GBT930- 2-((2-(1-cyclobutyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-6-hydroxybenzaldehyde.** The compound was prepared by Mitsunobu reaction (scheme 9, step 3) of (2-(1-cyclobutyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol (A) with 2,6-dihydroxybenzaldehyde (INT3). The product was obtained as pale yellow solid. ¹HNMR (300MHz, CDCl₃, ppm): 11.95 (s, 1H), 10.35 (s, 1H), 8.75 (m, 1H), 7.97 (d, J=7.2Hz, 1H), 7.62 (m, 1H), 7.45 (m, 1H), 7.36 (m, 1H), 6.56 (d, J=8.4Hz, 1H), 6.38 (m, 1H), 6.23 (d,

$J=8.4\text{Hz}$, 1H), 5.07 (s, 2H), 4.85 (m, 1H), 2.75 (m, 2H), 2.27 (m, 2H), 1.25 (m, 2H); (ES, m/z): 350 $[M+1]^+$



[0298] Preparation of intermediate A. Intermediate A was prepared from pyrazole in three steps according scheme below.




[0299] Step 1: Into a 500-mL round-bottom flask, was placed a solution of 1H-pyrazole (10 g, 146.89 mmol, 1.00 equiv), Cs_2CO_3 (95.9 g, 294.33 mmol, 2.00 equiv), and bromocyclobutane (29.7 g, 220.00 mmol, 1.50 equiv) in CH_3CN (150 mL). The resulting solution was stirred overnight at 80°C , and then it was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:400-1:200) as eluent to yield 8 g (45%) of 1-cyclobutyl-1H-pyrazole as a colorless liquid.

[0300] Step 2: Into a 250-mL round-bottom flask, was placed a solution of 1-cyclobutyl-1H-pyrazole (6.5 g, 53.21 mmol, 1.00 equiv) in tetrahydrofuran (100 mL). This was followed by the addition of BuLi (2.5 M, 22.7 mL, 1.10 equiv) dropwise with stirring at -30°C . The mixture was stirred for 30 min at the same temperature. To this was added 4,4,5,5-tetramethyl-2-(propan-2-yloxy)-1,3,2-dioxaborolane (14.9 g, 80.08 mmol, 1.50 equiv) dropwise with stirring at -78°C . The resulting solution was stirred for 3 h at -78°C , and then it was quenched by the addition of 30 mL of water. The resulting solution was extracted with 2x200 mL of dichloromethane. The combined organic layers were concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:200-1:30) as eluent to furnish 6.2 g (47%) of 1-cyclobutyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole as a colorless oil.

[0301] Step 3: Into a 100-mL round-bottom flask, was placed a solution of 1-cyclobutyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (791 mg, 3.19 mmol, 2.00 equiv), (2-

bromopyridin-3-yl)methanol (500 mg, 2.66 mmol, 1.00 equiv), Pd(dppf)Cl₂ (217 mg, 0.30 mmol, 0.10 equiv), and sodium carbonate (670 mg, 6.32 mmol, 3.00 equiv) in a solvent mixture of dioxane (10 mL) and water (10 mL). The resulting solution was stirred for 2 h at 80°C. The resulting solution was extracted with 3x20 mL of dichloromethane. The combined organic layers were concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:3-1:1) as eluent to provide 200 mg (33%) of [2-(1-cyclobutyl-1H-pyrazol-5-yl)pyridin-3-yl]methanol as a yellow oil.


GBT934



[0302] GBT934- 2-((2'-ethoxy-[2,3'-bipyridin]-3-yl)methoxy)-6-hydroxybenzaldehyde


The compound was prepared from (2-chloropyridin-3-yl)methanol and ((2-ethoxypyridin-3-yl)boronic acid according to scheme 9, reaction steps 1, 2 and 4. In step 4, alkylation with 2,6-dihydroxybenzaldehyde (INT3), the product TFA salt as white solid was obtained after HPLC purification. ¹HNMR (400MHz, CDCl₃, ppm): 11.91 (*br s*, 1H), 10.29 (*s*, 1H), 8.97 (*s*, 2H), 8.97 (*br s*, 1H), 8.46 (*d*, *J* = 8.0 Hz, 1H), 8.40 (*d*, *J* = 4.8 Hz, 1H), 7.87 (*s*, 1H), 7.81 (*d*, *J* = 6.8 Hz, 1H), 7.53 (*t*, *J* = 4.4 Hz, 1H), 7.12 (*t*, *J* = 6.0 Hz, 1H), 6.59 (*d*, *J* = 8.4 Hz, 1H), 6.14 (*d*, *J* = 8.4 Hz, 1H), 5.17 (*s*, 2H), 4.46 (*q*, *J* = 6.8 Hz, 2H), 1.32 (*t*, *J* = 6.8 Hz, 3H); MS (ES, *m/z*:) 351.1 [M+1]⁺

GBT948




[0303] GBT948- 3-chloro-2-hydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde. The compound was prepared using general reaction step 4 of scheme 9, by O-alkylation of 3-chloro-6-hydroxy-2-(methoxymethoxy)benzaldehyde (B) with 3-(chloromethyl)-2-(1-isopropyl-1H-pyrazol-5-yl)pyridine (A) and subsequent

deprotection of the MOM ether by treating with aqueous 6N HCl in THF. ^1H NMR (400MHz, CDCl_3 , ppm): 12.49(s, 1H), 10.34(s, 1H), 8.80(dd, $J=3.6\text{Hz}$, 1H), 8.00(dd, $J=5.7\text{Hz}$, 1H), 7.63(d, $J=1.2\text{Hz}$, 1H), 7.47(m, 2H), 6.36 (m, 2H), 5.11 (d, $J=10.8\text{Hz}$, 2H), 4.70(m, 1H), 4.61(m, 1H), 1.53(d, $J=4.5\text{Hz}$, 6H); MS (ES, m/z): 372[M+1] $^+$



[0304] The MOM-protected phenol aldehyde intermediate B was prepared according to the following synthetic scheme:




[0305] Step 1: Into a 2000-mL round-bottom flask, was placed a solution of 4-chlorobenzene-1,3-diol (50.0 g, 345.89 mmol, 1.00 equiv) and potassium carbonate (95 g, 687.36 mmol, 2.00 equiv) in acetone (1000 mL). This was followed by the addition of MOMBr (40 g, 320.10 mmol, 0.90 equiv) dropwise with stirring. The resulting solution was stirred for 60 h at room temperature. The solids were filtered out. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:10-1:2) as eluent to furnish 49 g (75%) of 4-chloro-3-(methoxymethoxy)phenol as a colorless oil.

[0306] Step 2: Into a 1000-mL round-bottom flask, was placed a solution of 4-chloro-3-(methoxymethoxy)phenol (49.0 g, 259.80 mmol, 1.00 equiv) and potassium carbonate (57.4 g, 415.31 mmol, 1.60 equiv) in acetone (500 mL). This was followed by the addition of BnBr (55 g, 321.58 mmol, 1.20 equiv) dropwise with stirring in 40 min. The resulting solution was stirred overnight at room temperature. The solids were filtered out. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:99-1:10-1:2) as eluent to furnish 40.0 g (55%) of 4-(benzyloxy)-1-chloro-2-(methoxymethoxy)benzene as a colorless oil.

[0307] Step 3: Into a 500-mL three neck round-bottom flask, was placed a solution of bis(propan-2-yl)amine (29.7 g, 293.51 mmol, 5.00 equiv) in tetrahydrofuran (70 mL). This was followed by the addition of BuLi (100 mL, 3.00 equiv) dropwise with stirring at -78 °C. The mixture was stirred for 10 min at -78 °C, then stirred for 10 min at 0°C. To this was added a solution of 4-(benzyloxy)-1-chloro-2-(methoxymethoxy)benzene (23.3 g, 83.59 mmol, 1.00 equiv) in tetrahydrofuran (70 mL) dropwise with stirring at -78 °C. The mixture was stirred for 1 h at -40 °C. To the mixture was added N,N-dimethylformamide (18.3 g, 250.38 mmol, 3.00 equiv) dropwise with stirring at -78°C. The resulting solution was stirred for 30 min at -78 °C. The reaction was then quenched by the addition of 50 mL of NH₄Cl (aq). The resulting solution was diluted with 50 mL of water. The resulting solution was extracted with 3x100 mL of ethyl acetate and the organic layers combined and dried over sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:5) as eluent. This resulted in 8.8 g (34%) of 6-(benzyloxy)-3-chloro-2-(methoxymethoxy)benzaldehyde as a yellow solid.


[0308] Step 4: Into a 500-mL round-bottom flask, was placed a solution of 6-(benzyloxy)-3-chloro-2-(methoxymethoxy)benzaldehyde (8.8 g, 28.69 mmol, 1.00 equiv) in ethyl acetate (100 mL). Rh/C (1.0 g) was added to the reaction. The resulting solution was stirred for 20 h at room temperature under 1 atm of hydrogen gas. The solids were filtered out. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:2) as eluent. This resulted in 5.2 g (84%) of 3-chloro-6-hydroxy-2-(methoxymethoxy)benzaldehyde as a light yellow solid.

GBT954




[0309] **GBT954- 4-hydroxy-2-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde.** The compound was prepared by Mitsunobu reaction (scheme 9, step 3) of 2-hydroxy-4-(methoxymethoxy)benzaldehyde (**A**) with (2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol (**B**) and subsequent deprotection of the MOM ether by treating with aqueous 6N HCl in THF according the scheme below. ¹HNMR (400MHz,

DMSO, ppm): 10.70 (s, 1H), 10.09 (s, 1H), 8.75 (m, 1H), 8.22 (d, $J=8\text{Hz}$, 1H), 7.59 (m, 3H), 6.52 (m, 3H), 5.16 (s, 2H), 4.65 (m, 1H), 1.37 (m, 6H); (ES, m/z): 338 $[M+1]^+$




[0310] Preparation of intermediate 2-hydroxy-4-(methoxymethoxy)benzaldehyde




[0311] Into a 100-mL round-bottom flask, was placed a solution of 2,4-dihydroxybenzaldehyde (3 g, 21.72 mmol, 1.00 equiv), MOMBr (3.2 g, 25.60 mmol, 1.20 equiv), and potassium carbonate (3.9 g, 28.22 mmol, 1.30 equiv) in acetone (20 mL). The resulting solution was stirred overnight at room temperature. The resulting solution was diluted with 30 mL of H_2O . The resulting solution was extracted with 3×20 mL of dichloromethane and the organic layers combined and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:50-1:30) as eluent to furnish 2.6 g (66%) of 2-hydroxy-4-(methoxymethoxy)benzaldehyde as a white solid.

GBT967




[0312] GBT967- 2-((2',6'-dimethoxy-[2,3'-bipyridin]-3-yl)methoxy)-6-hydroxybenzaldehyde. The compound was prepared from (2-chloropyridin-3-yl)methanol and 2,6-dimethoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine according to scheme 9, reaction steps 1, 2 and 4. In step 4, alkylation with 2,6-dihydroxybenzaldehyde, the product of TFA salt as white solid was obtained after HPLC purification. ^1H NMR (400MHz, DMSO, ppm): 10.04 (s, 1H), 8.65 (m, 1H), 8.14 (d, $J=7.6\text{Hz}$, 1H), 7.73 (d, $J=8\text{Hz}$, 1H), 7.51 (m, 2H), 6.50 (m, 3H), 5.16 (m, 2H), 3.91 (s, 1H), 3.86 (s, 1H); (ES, m/z): 367 $[M+1]^+$

GBT000985




[0313] GBT985- 3-(difluoromethyl)-1-methyl-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-1H-pyrazole-4-carbaldehyde.




[0314] 3-(Difluoromethyl)-5-hydroxy-1-methyl-1H-pyrazole-4-carbaldehyde (100 mg, 0.568 mmol) was dissolved in DMF (2.8 ml). 3-(chloromethyl)-2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridine hydrochloride (0.195 g, 0.625 mmol) and potassium carbonate (0.235 g, 1.7 mmol) were then added and the mixture was stirred in a 60 °C heat block for 16 h. The reaction mixture was cooled and water (50 ml) and ethyl acetate (100 ml) were added. The phases were separated and the aqueous phase was extracted with ethyl acetate (2 x 50 ml). The combined organic phases were washed with water (20 ml) and a saturated aqueous sodium chloride solution (20 ml). After drying over sodium sulfate and evaporation, the residue was purified by silica gel chromatography (5 – 50 % ethyl acetate/hexanes) to give 86 mg (36%) of 3-(difluoromethyl)-1-methyl-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-1H-pyrazole-4-carbaldehyde as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.97 (s, 1H), 8.75 (d, J = 3.64 Hz, 1H), 8.08 (d, J = 7.31 Hz, 1H), 7.68 (d, J = 1.69 Hz, 1H), 7.44 (dd, J = 4.66, 7.83 Hz, 1H), 6.65 (t, J = 53.67 Hz, 1H), 6.61 (d, J = 1.78 Hz, 1H), 5.64 (s, 2H), 5.24 (q, J = 8.60 Hz, 2H), 3.54 (s, 3H). MS (ESI) m/z 416 [M+H]⁺.

GBT986




[0315] GBT986- 1-methyl-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-3-(trifluoromethyl)-1H-pyrazole-4-carbaldehyde




[0316] 5-Hydroxy-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carbaldehyde (100 mg, 0.515 mmol) was dissolved in DMF (2.5 ml). 3-(chloromethyl)-2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridine hydrochloride (0.177 g, 0.567 mmol) and potassium carbonate (0.213 g, 1.545 mmol) were then added and the mixture was stirred in a 60 °C heat block for 16 h. The reaction mixture was cooled and water (50 ml) and ethyl acetate (100 ml) were added. The phases were separated and the aqueous phase was extracted with ethyl acetate (2 x 50 ml). The combined organic phases were washed with water (20 ml) and a saturated aqueous sodium chloride solution (20 ml). After drying over sodium sulfate and evaporation, the residue was purified by silica gel chromatography (5 – 50 % ethyl acetate/hexanes) to give 86 mg (36%) of 1-methyl-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-3-(trifluoromethyl)-1H-pyrazole-4-carbaldehyde as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.87 (s, 1H), 8.75 (d, *J* = 4.33 Hz, 1H), 8.08 (d, *J* = 7.73 Hz, 1H), 7.68 (d, *J* = 1.74 Hz, 1H), 7.45 (dd, *J* = 4.64, 7.84 Hz, 1H), 6.59 (d, *J* = 1.70 Hz, 1H), 5.63 (s, 2H), 5.24 (q, *J* = 8.60 Hz, 2H), 3.57 (s, 3H). MS (ESI) *m/z* 434 [M+H]⁺.


GBT1065




[0317] **GBT1065 - 4-fluoro-7-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2,3-dihydro-1H-inden-1-one bis(2,2,2-trifluoroacetate).** The compound was prepared by Mitsunobu reaction of 4-fluoro-7-hydroxy-2,3-dihydro-1H-inden-1-one with [2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methanol according to scheme 9, reaction step 3. The product TFA salt as white solid was obtained after HPLC purification. ¹HNMR (300MHz, DMSO, *ppm*): 8.72 (m, 1H), 8.23 (m, 1H), 7.56(m, 1H), 7.51(d, J=1.8Hz, 1H), 7.43(m, 1H), 6.93 (m, 1H), 6.58(d, J=1.8Hz, 1H), 5.08(s, 2H), 4.63(m, 1H), 3.03 (m, 2H), 2.61 (m, 2H), 1.33 (d, J=6.6Hz, 6H); MS (ES, *m/z*): 366[M+1]⁺



GBT1133



[0318] **GBT1133- 2-(((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)oxy)methyl)benzaldehyde.** The compound was prepared from ethyl 2-(bromomethyl)benzonitrile in 2 steps according to reaction scheme below.




[0319] **Step 1:** Into a 100-mL round-bottom flask, was placed a solution of 2-(bromomethyl)benzonitrile (1.0 g, 5.10 mmol, 1.00 equiv) in dichloromethane (40 mL). This

was followed by the addition of DIBAL-H (5.5 mL, 1.10 equiv) at 0 °C. The resulting solution was stirred for 3.5 h at 0 °C. The reaction was then quenched by the addition of 10 mL of 5% HBr at 0 °C. The resulting solution was extracted with 3x30 mL of dichloromethane and the combined organic layers were dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:10) as eluent. This resulted in 500 mg (49%) of 2-(bromomethyl)benzaldehyde as a green oil.


[0320] Step 2: Into a 50-mL round-bottom flask, was placed a solution of 2-(bromomethyl)benzaldehyde (150 mg, 0.75 mmol, 1.00 equiv) in CH₃CN (25 mL). 2-[1-(Propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-ol (150 mg, 0.74 mmol, 1.00 equiv), potassium carbonate (210 mg, 1.52 mmol, 2.00 equiv), and KI (40 mg, 0.30 equiv) were added to the reaction. The resulting solution was heated to reflux for 6 h, and then it was cooled to rt. The resulting solution was diluted with 20 mL of H₂O, and then it was extracted with 3x20 mL of ethyl acetate. The combined organic layers were washed with 1x30 mL of brine and concentrated under vacuum. The crude product (200 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-010): Column, SunFire Prep C18 OBD Column, 5µm, 19*150mm,; mobile phase, water with 0.05%TFA and MeCN (38.0% MeCN up to 55.0% in 8 min); Detector, nm. This provided 98.6 mg (41%) of 2-[[2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]oxy)methyl]benzaldehyde as a light yellow solid; ¹HNMR (300MHz, CDCl₃, ppm): 10.01 (s, 1H), 8.43 (m, 1H), 7.88(m, 1H), 7.86 (m, 1H), 7.61-7.79(m, 6H), 6.61(d, *J*=2.1Hz, 1H), 5.60 (s, 2H), 4.69-4.78 (m, 1H), 1.46(d, *J*=6.6Hz, 6H); (ES, *m/z*): 322 [M+1]⁺

GBT1197




[0321] GBT1197- 2-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde. The compound was prepared by O-alkylation of 2-hydroxybenzaldehyde with 3-(chloromethyl)-2-(1-isopropyl-1H-pyrazol-5-yl)pyridine (INT-4). The product of TFA salt as white solid was obtained after HPLC purification. ¹HNMR (300MHz, CDCl₃, ppm): 10.49(s, 1H), 8.78(m, 1H), 8.16(m, 1H), 7.88(m, 1H), 7.69(d,


$J=6.0\text{Hz}$, 1H), 7.54(m, 2H), 7.13(m, 1H), 6.90(d, $J=8.4\text{Hz}$, 1H), 6.41(d, $J=1.8\text{Hz}$, 1H), 5.11 (s, 2H), 4.62 (m, 1H); (ES, m/z): 322[M+1]⁺

GBT1252

[0322] **GBT1252- 2-((6-bromo-2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-6-hydroxybenzaldehyde.** The compound was prepared by Mitsunobu reaction of (6-bromo-2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol (intermediate A) and 2,6-dihydroxybenzaldehyde (INT-3) according to scheme 9, reaction step 3. The product as white solid was obtained after flash column purification. ¹HNMR (300MHz, DMSO, ppm): 11.70 (s, 1H), 11.20 (s, 1H), 8.17 (d, $J = 8.1\text{ Hz}$, 1H), 7.81 (d, $J = 8.1\text{ Hz}$, 1H), 7.57 (s, 1H), 7.50 (t, $J = 8.4\text{ Hz}$, 1H), 6.60 (s, 1H), 6.55 (dd, $J = 8.4\text{ Hz}$, 3.6 Hz, 1H), 5.19 (s, 2H), 4.65-4.55 (m, 1H), 1.38 (d, $J = 6.6\text{ Hz}$, 6H); (ES, m/z): 418.1 [M+1]⁺



[0323] **Preparation of intermediate A.** Intermediate A was prepared from 2,6-dichloropyridine-3-carboxylic acid in five steps according to the synthetic scheme below.



[0324] Step 1: Into a 500-mL round-bottom flask, was placed a solution of 2,6-dichloropyridine-3-carboxylic acid (25 g, 130.21 mmol, 1.00 equiv) in methanol (350 mL). This was followed by the addition of *t*-BuOK (43.8 g, 390.34 mmol, 3.00 equiv), in portions at 0 °C. The resulting solution was heated to reflux for 3 days. The resulting mixture was concentrated under vacuum, and then it was diluted with 400 mL of water. The pH value of the solution was adjusted to 1 with hydrogen chloride aq. The solids were collected by filtration. The solid was dried in an oven under reduced pressure. This provided 20.5 g (84%) of 2-chloro-6-methoxypyridine-3-carboxylic acid as a white solid.

[0325] Step 2: Into a 100-mL round-bottom flask, was placed a solution of 2-chloro-6-methoxypyridine-3-carboxylic acid (11.3 g, 60.24 mmol, 1.00 equiv) in methanol (50 mL). This was followed by the addition of SOCl₂ (26 mL, 5.00 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 1 h at room temperature and for an additional 2 h at reflux. The resulting solution was diluted with 100 mL of water. The pH value of the solution was adjusted to 10 with 2M sodium carbonate aq. The resulting solution was extracted with 3x80 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 1x150 mL of water and 1x100 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:10) as eluent to furnish 7.8 g (64%) of methyl 2-chloro-6-methoxypyridine-3-carboxylate as a white solid.

[0326] Step 3: Into a 100-mL round-bottom flask, which was purged and maintained with an inert atmosphere of nitrogen, was placed methyl 2-chloro-6-methoxypyridine-3-

carboxylate (3.29 g, 16.32 mmol, 1.00 equiv), 1-(propan-2-yl)-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (4.63 g, 19.61 mmol, 1.20 equiv), toluene (45 mL), ethanol (15 mL), sodium carbonate (2M in H₂O) (15 mL). This was followed by the addition of Pd(dppf)(DCM)Cl₂ (665 mg, 0.05 equiv). The resulting solution was stirred for 20 h at 90 °C. The reaction was then quenched by the addition of 20 mL of water and 100 mL of ethyl acetate. The resulting mixture was washed with 2x50 mL of water and 1x50 mL of brine. The mixture was dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:25-1:15) as eluent to furnish 3.83 g (85%) of methyl 6-methoxy-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridine-3-carboxylate as a light yellow oil.

[0327] Step 4: Into a 250-mL round-bottom flask, was placed methyl 6-methoxy-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridine-3-carboxylate (3.6 g, 13.08 mmol, 1.00 equiv), C₆H₆ (60 mL), H₃PO₄(85%) (150 mg, 1.53 mmol, 0.10 equiv), PyHBr (208 mg, 1.30 mmol, 0.10 equiv), POBr₃ (11.5 g, 40.11 mmol, 3.00 equiv). The resulting solution was heated to reflux for 40 h. The reaction mixture was cooled to 0 °C with an ice bath. The pH value of the solution was adjusted to 10 with potassium carbonate sat. The resulting solution was extracted with 3x80 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 2x50 mL of water and 1x50 mL of brine. The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:10-1:8) as eluent. This provided 2.62 g (62%) of methyl 6-bromo-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridine-3-carboxylate as a yellow oil.

[0328] Step 5: Into a 50-mL round-bottom flask, was placed methyl 6-bromo-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridine-3-carboxylate (2.62 g, 8.08 mmol, 1.00 equiv), tetrahydrofuran (30 mL). This was followed by the addition of LiBH₄ (350 mg, 16.07 mmol, 2.00 equiv) at 0 °C. The resulting solution was stirred for 30 min at 0 °C and for an additional 1 h at room temperature. The reaction mixture was cooled to 0 °C with an ice bath. The reaction was then quenched by the addition of 20 mL of water. The resulting solution was extracted with 3x60 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 2x50 mL of water and 1x50 mL of brine. The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:8-1:5) as eluent to furnish 1.97 g


quenched by the addition of 100 mL of sodium hydroxide (aq). The pH value of the solution was adjusted to 12 with sodium hydroxide (aq). The resulting solution was extracted with 3x60 mL of ethyl acetate. The combined organic layers were washed with 1x60 mL of water and 1x60 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1-1:0) as eluent to yield 2.08 g (75%) of 4-(dimethylamino)-2,6-dimethoxybenzaldehyde (**3**) as a white solid.

[0332] Step 3: Into a 50-mL round-bottom flask, was placed a solution of 4-(dimethylamino)-2,6-dimethoxybenzaldehyde (**3**) (630 mg, 3.01 mmol, 1.00 equiv) in dichloromethane (25 mL). AlCl₃ (6 g, 45.11 mmol, 12.50 equiv) was added to the reaction. The resulting solution was heated to reflux for 24 h. The pH value of the solution was adjusted to 8-9 with sodium bicarbonate (aq). The resulting solution was extracted with 3x80 mL of ethyl acetate. The combined organic layers were washed with 1x60 mL of water and 1x60 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:4) as eluent to furnish 383 mg (70%) of 4-(dimethylamino)-2,6-dihydroxybenzaldehyde (**4**) as a light yellow solid.

[0333] Step 4: Into a 25-mL round-bottom flask, was placed a solution of [2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methanol (**5**) (132 mg, 0.61 mmol, 1.00 equiv), 4-(dimethylamino)-2,6-dihydroxybenzaldehyde (**4**) (110 mg, 0.61 mmol, 1.00 equiv), PPh₃ (207.3 mg, 0.79 mmol, 1.30 equiv) in tetrahydrofuran (10 mL). DIAD (160 mg, 0.79 mmol, 1.30 equiv) was added to the reaction dropwise dropwise at 0 °C. The resulting solution was stirred for 10 min at 0 °C and for an additional 2 h at room temperature. The reaction was then quenched with 10 mL of water. The resulting solution was extracted with 3x40 mL of ethyl acetate. The combined organic layers were washed with 3x20 mL of water and 1x20 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The crude product was purified by Prep-HPLC with the following conditions (Prep-HPLC-010): Column, SunFire Prep C18 OBD Column, Sun, 19*150mm,; mobile phase, water with 0.05%TFA and MeCN (30.0% MeCN up to 60.0% in 10 min, up to 95.0% in 4 min, down to 30.0% in 2 min); Detector, Waters2545 UvDector 254&220nm. This resulted in 60 mg (26%) of 4-(dimethylamino)-2-hydroxy-6-([2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methoxy)benzaldehyde (**GBT933**) as a light yellow solid.

(82%) of [6-bromo-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methanol as a light yellow solid.

GBT933




[0329] GBT933 - 4-(dimethylamino)-2-hydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde

[0330] Step 1: Into a 250-mL round-bottom flask, was placed a solution of 3,5-dimethoxyaniline (**1**) (4.6 g, 30.03 mmol, 1.00 equiv) and potassium carbonate (14.6 g, 105.64 mmol, 4.00 equiv) in N,N-dimethylformamide (80 mL). This was followed by the addition of $\text{SO}_2(\text{OMe})_2$ (8.4 g, 66.67 mmol, 2.00 equiv) dropwise with stirring at 0 °C. The mixture was stirred for 2h at 0 °C. The resulting solution was stirred overnight at room temperature, and then it was quenched by the addition of 100 mL of water. The resulting solution was extracted with 2x80 mL of ethyl acetate. The combined organic layers were washed with 1x50 mL of water and 1x50 mL of brine, dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with PE:EA (30:1-10:1) as eluent to furnish 2.8 g (51%) of 3,5-dimethoxy-N,N-dimethylaniline (**2**) as a white solid.

[0331] Step 2: Into a 50-mL round-bottom flask, was placed a solution of 3,5-dimethoxy-N,N-dimethylaniline (**2**) (2.4 g, 13.24 mmol, 1.00 equiv) in N,N-dimethylformamide (25 mL). This was followed by the addition of POCl_3 (2.57 g, 16.76 mmol, 1.30 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 15 min at 0 °C, and then it was

[0334] ^1H NMR (400MHz, CDCl_3 , ppm): 12.45 (*br s*, 1H), 9.87 (*s*, 1H), 8.70 (*d*, $J = 4.0$ Hz, 1H), 7.96 (*d*, $J = 7.2$ Hz, 1H), 7.58 (*s*, 1H), 7.42-7.39 (*m*, 1H), 6.32 (*s*, 1H), 5.68 (*s*, 1H), 5.43 (*s*, 1H), 4.94 (*s*, 2H), 4.58-4.51 (*m*, 1H), 2.96 (*s*, 6H), 1.40 (*d*, $J = 6.8$ Hz, 1H); MS (ES, m/z): 381.2 $[\text{M}-2\text{CF}_3\text{COOH}+1]^+$

GBT953

[0335] **GBT953 - (S)-2-hydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde**


[0336] Step 1: Into a 250-mL round-bottom flask, was placed a solution of methyl 3-hydroxybenzoate (3 g, 19.72 mmol, 1.00 equiv) in trifluoroacetic acid (100 mL). Urotropin (5.5 g, 39.29 mmol, 2.00 equiv) was added to the reaction. The resulting solution was heated to reflux for 2 hr. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:5) as eluent to furnish 0.5 g (14%) of methyl 2-formyl-3-hydroxybenzoate as a yellow solid.

[0337] Step 2: Into a 100-mL round-bottom flask, was placed a solution of methyl 2-formyl-3-hydroxybenzoate (400 mg, 2.22 mmol, 1.00 equiv) in CH_3CN (30 mL). 3-(Chloromethyl)-2-[(1-(propan-2-yl)-1H-pyrazol-5-yl)pyridin-3-yl]methoxybenzoate (523 mg, 2.22 mmol, 1.00 equiv), potassium carbonate (927 mg, 6.71 mmol, 3.00 equiv), and KI (40 mg, 0.24 mmol, 0.10 equiv) were added to the reaction. The resulting solution was stirred for 2 h at 60°C , and then it was diluted with 200 mL of EA. The resulting mixture was washed with 2x100 mL of brine, and then it was concentrated under vacuum. This provided 500 mg (59%) of methyl 2-formyl-3-((2-[(1-(propan-2-yl)-1H-pyrazol-5-yl)pyridin-3-yl]methoxy)benzoate as a yellow oil.

[0338] Step 3: Into a 100-mL round-bottom flask, was placed a solution of methyl 2-formyl-3-((2-[(1-(propan-2-yl)-1H-pyrazol-5-yl)pyridin-3-yl]methoxy)benzoate (100 mg, 0.26 mmol, 1.00 equiv) in tetrahydrofuran (10 mL). This was followed by the addition of a solution of sodium hydroxide (22 mg, 0.55 mmol, 2.00 equiv) in water (3 mL) dropwise with stirring at 0°C . The resulting solution was stirred for 3 h at room temperature. The pH value of the solution was adjusted to 6 with hydrogen chloride (3 mol/L). The resulting solution

was extracted with 2x100 mL of ethyl acetate and the organic layers combined and concentrated under vacuum. The crude product (100 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-010): Column, SunFire Prep C18 OBD Column, 5 μ m, 19*150mm; mobile phase, water with 0.05%TFA and MeCN (30% MeCN up to 60% in 10 min, up to 95% in 3 min, down to 30% in 2 min); Detector, Waters2545 UvDetector 254&220nm. This resulted in 30 mg (19%) of 2-formyl-3-([2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methoxy)benzoic acid; bis(trifluoroacetic acid) as a white solid. ¹H NMR (300MHz, DMSO, ppm): 8.75(dd, J=4.8Hz, 1H), 8.15 (m, 2H), 7.59(m,3H), 7.40(d, J=7.8Hz, 1H), 7.33(d, J=8.1Hz, 1H), 6.66((d, J=7.8Hz, 1H) , 6.60 (s, 1H), 5.18 (m, 2H), 4.70 (m, 1H), 1.35(m, 6H); MS (ES, *m/z*):366 [M+1]⁺

GBT963




[0339] GBT963 - (S)-2-hydroxy-6-((5-oxo-1-phenylpyrrolidin-2-yl)methoxy)benzaldehyde

[0340] Step 1: To a suspension of (S)-5-(hydroxymethyl)pyrrolidin-2-one (230 mg, 2 mmol) and iodobenzene (0.49 g, 2.4 mmol) in Dioxane (4 mL) was added CuI (0.08 g, 0.4 mmol), N,N-dimethylethylenediamine (0.05 mL, 0.4 mmol), K₂CO₃ (0.55 g, 4 mmol). After heating at 100 °C for 24 h, the mixture was cooled and was diluted with EtOAc, insoluble material was filtered off, the filtrate was concentrated and purified by column (Hexanes/EtOAc=100:0 to 0:100) to give (S)-5-(hydroxymethyl)-1-phenylpyrrolidin-2-one (280 mg).

[0341] Step 2: To a solution of (S)-5-(hydroxymethyl)-1-phenylpyrrolidin-2-one (100 mg, 0.52 mmol) and 2,6-dihydroxybenzaldehyde (0.09 g, 0.65 mmol) in THF (5 mL) at 0 °C was added PPh₃ (polymer supported, 650 mg, 0.78 mmol) and DIAD (0.16 mL, 0.78 mmol). After stirred for 1 h, it was diluted with AcCN, the insoluble material was filtered off and the filtrate was concentrated and purified by preparative HPLC to give (S)-2-hydroxy-6-((5-oxo-1-phenylpyrrolidin-2-yl)methoxy)benzaldehyde (86 mg). ¹H NMR (400 MHz, Chloroform-*d*)

δ 11.91 (d, $J = 0.4$ Hz, 1H), 9.93 (d, $J = 0.6$ Hz, 1H), 7.50 – 7.18 (m, 6H), 6.53 (dt, $J = 8.5$, 0.7 Hz, 1H), 6.21 (dd, $J = 8.3$, 0.8 Hz, 1H), 4.70 – 4.59 (m, 1H), 4.15 – 4.01 (m, 2H), 2.82 – 2.58 (m, 2H), 2.50 (ddt, $J = 13.1$, 9.9, 8.3 Hz, 1H), 2.16 (dddd, $J = 13.6$, 9.5, 5.1, 3.9 Hz, 1H). MS found for $C_{18}H_{17}NO_4$: 312.3.


GBT998

[0342] GBT998 - (S)-2-hydroxy-6-((5-oxo-1-phenylpyrrolidin-2-yl)methoxy)benzaldehyde

[0343] Step 1: To a suspension of (S)-5-(hydroxymethyl)pyrrolidin-2-one (230 mg, 2 mmol) and 1-iodo-2-methoxybenzene (0.56 g, 2.4 mmol) in Dioxane (4 mL) was added CuI (0.08 g, 0.4 mmol), N,N-dimethylethylenediamine (0.05 mL, 0.4 mmol), K_3PO_4 (0.85 g, 4 mmol). After heating at 100 °C for 24 h, the mixture was cooled and was diluted with EtOAc, insoluble material was filtered off, the filtrate was concentrated and purified by column (Hexanes/EtOAc=100:0 to 0:100) to give (S)-5-(hydroxymethyl)-1-(2-methoxyphenyl)pyrrolidin-2-one (110 mg).

[0344] Step 2: To a solution of (S)-5-(hydroxymethyl)-1-phenylpyrrolidin-2-one (115 mg, 0.54 mmol) and 2,6-dihydroxybenzaldehyde (0.10 g, 0.70 mmol) in THF (4 mL) at 0 °C was added PPh_3 (polymer supported, 675 mg, 0.81 mmol) and DIAD (0.16 mL, 0.81 mmol). After stirred for 1 h, it was diluted with AcCN, the insoluble material was filtered off and the filtrate was concentrated and purified by column (100% EtOAc) to give (S)-2-hydroxy-6-((1-(2-methoxyphenyl)-5-oxopyrrolidin-2-yl)methoxy)benzaldehyde (53 mg). ^1H NMR (400 MHz, Chloroform- d) δ 11.92 (s, 1H), 9.68 (t, $J = 0.5$ Hz, 1H), 7.38 – 7.30 (m, 1H), 7.30 – 7.22 (m, 1H), 7.14 (dd, $J = 7.7$, 1.7 Hz, 1H), 6.99 – 6.89 (m, 2H), 6.50 (dq, $J = 8.5$, 0.6 Hz, 1H), 6.25 – 6.18 (m, 1H), 4.55 (dtd, $J = 9.0$, 5.1, 4.0 Hz, 1H), 4.10 – 3.94 (m, 2H), 3.73 (s, 3H), 2.75 – 2.55 (m, 2H), 2.48 (dddd, $J = 13.0$, 9.6, 8.4, 7.3 Hz, 1H), 2.16 – 2.02 (m, 1H). MS found for $C_{19}H_{19}NO_5$: 342.3.

GBT1004



[0345] **GBT1004** - 2-amino-4-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy) nicotinaldehyde


[0346] Step 1: Into a 250-mL round-bottom flask, was placed a solution of 4-chloropyridin-2-amine (10 g, 77.78 mmol, 1.00 equiv) in tetrahydrofuran (150 mL). This was followed by the addition of LiHMDS(1M) (156 mL) dropwise with stirring at 0°C. The mixture was stirred for 20 min at 0 °C. To this was added Boc₂O (17.02 g, 77.98 mmol, 1.00 equiv). The resulting solution was stirred for 1 h at 0 °C. The reaction was then quenched by the addition of 100 mL of NH₄Cl. The resulting solution was extracted with 2x200 mL of ethyl acetate and the organic layers combined. The resulting mixture was washed with 1x150 mL of brine. The resulting mixture was concentrated under vacuum, washed with 2x20 mL of EA/hexane (3:7). This resulted in 12.5 g (70%) of tert-butyl N-(4-chloropyridin-2-yl)carbamate as a white solid.

[0347] Step 2: Into a 100-mL three neck round-bottom flask, was placed a solution of tert-butyl N-(4-chloropyridin-2-yl)carbamate (2 g, 8.75 mmol, 1.00 equiv) in tetrahydrofuran (50 mL). This was followed by the addition of BuLi (2.5M) (7.0 mL, 2.00 equiv) dropwise with stirring at -78 °C in 20 min. The mixture was stirred for 0.5h at -78 °C. To this was added N,N-dimethylformamide (2.0 mL, 3.00 equiv) dropwise with stirring at -78 °C in 5 min. The resulting solution was stirred for 1 h at -78 °C. The reaction was then quenched by the addition of 3 mL of hydrogen chloride (12M). The resulting mixture was concentrated under vacuum. The resulting solution was diluted with 50 mL of ethyl acetate. The resulting mixture was washed with 1x40 mL of 5% sodium bicarbonate and 1x30 mL of brine. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with EA:PE (1:4). This resulted in 1.46 g (65%) of tert-butyl N-(4-chloro-3-formylpyridin-2-yl)carbamate as a yellow solid.

[0348] Step 3: Into a 100-mL round-bottom flask, was placed a solution of [2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methanol (500 mg, 2.30 mmol, 1.20 equiv) in N,N-dimethylformamide (50 mL). This was followed by the addition of sodium hydride (190 mg, 7.92 mmol, 2.50 equiv) at 0 °C. The mixture was stirred for 20 min at 0 °C. To this was added tert-butyl N-(4-chloro-3-formylpyridin-2-yl)carbamate (500 mg, 1.95 mmol, 1.00 equiv). The resulting solution was stirred overnight at room temperature, and then it was quenched by the addition of 50 mL of water. The resulting solution was extracted with 8x20 mL of ethyl acetate and the organic layers combined and concentrated under vacuum. This resulted in 506 mg (59%) of tert-butyl N-[3-formyl-4-([2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methoxy)pyridin-2-yl]carbamate as a yellow oil.

[0349] Step 4: Into a 100-mL round-bottom flask, was placed a solution of trifluoroacetic acid (10 mL) in dichloromethane (60 mL), tert-butyl N-[3-formyl-4-([2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methoxy)pyridin-2-yl]carbamate (500 mg, 1.14 mmol, 1.00 equiv). The resulting solution was stirred overnight at room temperature. The resulting mixture was concentrated under vacuum. The crude product (500 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-010): Column, SunFire Prep C18 OBD Column, 5µm, 19*150mm; mobile phase, water with 0.05%TFA and MeCN (5.0% MeCN up to 26.0% in 10 min, up to 95.0% in 2 min, down to 5.0% in 2 min); Detector, Waters2545 UvDetector 254&220nm. This resulted in 122.9 mg (32%) of 2-amino-4-([2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methoxy)pyridine-3-carbaldehyde as a yellow solid. ¹HNMR (300MHz, DMSO, ppm): 10.12(s, 1H), 8.79(m, 1H), 8.24(m, 2H), 7.59(m, 2H), 6.76(d, J=5.1Hz, 1H), 6.55(d, J=1.2Hz, 1H), 5.43(m, 2H), 4.67(m, 1H), 1.36(m, 6H); MS (ES, m/z): 338 [M+1-3CF₃COOH]⁺

GBT1006




[0350] GBT1006 - 4-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-(methylamino)nicotinaldehyde

[0351] Step 1: Into a 100-mL round-bottom flask, was placed a solution of tert-butyl N-(4-chloropyridin-2-yl)carbamate (3.0 g, 13.12 mmol, 1.00 equiv) in tetrahydrofuran (50 mL). This was followed by the addition of sodium hydride (631 mg, 26.29 mmol, 1.20 equiv) at 0 °C. The mixture was stirred for 20 min at 0 °C. To this was added iodomethane (2.24 g, 15.78 mmol, 1.20 equiv) dropwise with stirring. The resulting solution was stirred for 6 h at room temperature. The reaction was then quenched by the addition of 30 mL of water. The resulting solution was extracted with 3x30 mL of ethyl acetate and the organic layers combined and concentrated under vacuum. This resulted in 3.01 g (95%) of tert-butyl N-(4-chloropyridin-2-yl)-N-methylcarbamate as a yellow oil.


[0352] Step 2: Into a 100-mL three neck round-bottom flask, was placed a solution of tert-butyl N-(4-chloropyridin-2-yl)-N-methylcarbamate (1.5 g, 6.18 mmol, 1.00 equiv) in tetrahydrofuran (50 mL). This was followed by the addition of BuLi (2.5M) (3.0 mL, 1.20 equiv) dropwise with stirring at -78 °C. The mixture was stirred for 30 mins at -78 °C. To this was added N,N-dimethylformamide (1.5 mL, 3.00 equiv) dropwise with stirring at -78 °C. The resulting solution was stirred for 1 h at -78 °C. The reaction was then quenched by the addition of 2.5 mL of hydrogen chloride (12M). The resulting mixture was concentrated under vacuum. The residue was dissolved in 40 mL of EA. The resulting mixture was washed with 1x30 mL of 5% sodium bicarbonate and 1x20 mL of brine. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with EA:PE (1:4). This resulted in 0.97 g (92%) of 4-chloro-2-(methylamino)pyridine-3-carbaldehyde as a yellow solid.

[0353] Steps 3 & 4: Into a 100-mL round-bottom flask, was placed a solution of [2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methanol (1.15 g, 5.29 mmol, 1.00 equiv) in N,N-dimethylformamide (40 mL). This was followed by the addition of sodium hydride (530 mg, 13.25 mmol, 2.50 equiv, 60%) at 0 °C. The mixture was stirred for 15 min at 0 °C. To this was added 4-chloro-2-(methylamino)pyridine-3-carbaldehyde (900 mg, 5.28 mmol, 1.00 equiv). The resulting solution was stirred overnight at room temperature. The reaction was then quenched by the addition of 30 mL of water. The resulting solution was extracted with 5x30 mL of ethyl acetate and the organic layers combined and concentrated under vacuum. The crude product (300 mg) was purified by Prep-HPLC with the following conditions (Prep-HPLC-020): Column, SunFire Prep C18 OBD Column, 5µm, 19*100mm.; mobile phase, water with 0.1% TFA and MeCN (3.0% MeCN up to 20.0% in 5 min, up to 95.0% in 2 min, down to 3.0% in 1 min); Detector, waters2489 254&220nm. This resulted in 107.1 mg (6%) of 2-

(methylamino)-4-([2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridin-3-yl]methoxy)pyridine-3-carbaldehyde as a yellow solid. ^1H NMR (400MHz, DMSO, *ppm*): 8.72 (m, 1H), 8.17 (s, 1H), 7.91 (m, 1H), 7.52 (m, 3H), 6.56 (s, 1H), 6.26(d, $J=4.2\text{Hz}$, 1H), 6.15(d, $J=3.3\text{Hz}$, 1H), 5.43 (m, 1H), 5.12 (m, 1H), 4.60 (m, 1H), 2.87(d, $J=3.3\text{Hz}$, 1H), 1.46(d, $J=5.1\text{Hz}$, 1H), 1.35(d, $J=5.1\text{Hz}$, 1H); (ES, m/z):352.1 $[\text{M}+1]^+$


GBT1007

[0354] GBT1007- 2-(4-(hydroxymethyl)-1,3-dioxolan-2-yl)-3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)phenol



[0355] 2-Hydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde (0.4 g, 1.19 mmol) was dissolved in a solution of propane-1,2,3-triol (5.8 ml, 79 mmol) and DMF (5 ml). Amberlyst 15 resin (80 mg) and 3A molecular sieves (1 g) were added and the mixture was stirred in a heat block at 70 °C for 18 h. After cooling the mixture was filtered and taken up in ethyl acetate (200 ml) and water (100 ml). The phases were separated and the aqueous phase was extracted with ethyl acetate (2 x 50 ml). The combined organic phases were washed with water (50 ml) and a saturated aqueous sodium chloride solution (50 ml), and dried over sodium sulfate. After evaporation, the residue was purified by silica gel chromatography (0 – 90% ethyl acetate/dichloromethane) to give 0.118 g (24%) of 2-(4-(hydroxymethyl)-1,3-dioxolan-2-yl)-3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)phenol as an off-white solid after lyophilization from acetonitrile/water. ^1H NMR (400 MHz, CDCl_3) δ 8.74 – 8.67 (m, 1H), 8.53 (s, 0.6H), 8.18 – 8.12 (m, 0.4H), 7.97 (d, $J=7.90\text{ Hz}$, 1H), 7.62 – 7.55 (m, 1H), 7.44 – 7.34 (m, 1H), 7.17 – 7.07 (m, 1H), 6.55 (d, $J=8.47\text{ Hz}$, 1H), 6.42 – 6.36 (m, 1H), 6.31 – 6.23 (m, 1.6H), 5.95 – 5.89 (m, 0.4H), 4.98 (s, 2H), 4.71 – 4.58 (m, 1H), 4.40 – 4.28 (m, 1.5H), 4.24 – 4.17 (m, 0.6H), 4.10 – 4.01 (m, 1H), 3.99 – 3.92 (m, 0.6H), 3.73 – 3.65 (m, 0.6H), 3.56 (dd, $J=10.09, 20.18\text{ Hz}$, 1H), 1.51 – 1.43 (m, 6H). MS (ESI) m/z 412 $[\text{M}+\text{H}]^+$.

GBT1090




[0356] GBT1090 - (S)-2-((1-benzyl-5-oxopyrrolidin-2-yl)methoxy)-6-hydroxybenzaldehyde

[0357] Step 1: To a solution of (S)-methyl 5-oxopyrrolidine-2-carboxylate (700 mg, 4.89 mmol) in DMF (5 mL) was added Cs_2CO_3 (1.97 g, 5.99 mmol) and BnBr (0.59 mL, 4.99 mmol). After heated at 60 °C for 15 h, it was diluted with EtOAc, organic layer was washed with water, brine, dried and concentrated to give crude product which was purified by column (Hexanes/EtOAc=40:60) to give methyl (S)-1-benzyl-5-oxopyrrolidine-2-carboxylate (240 mg). To a solution of methyl (S)-1-benzyl-5-oxopyrrolidine-2-carboxylate (240 mg, 1.0 mmol) in THF (2 mL) was added LiBH_4 (1M in THF, 1 mL, 1 mmol) at room temperature. After stirred at room temperature for 1 h, the solution was quenched with Sat. NH_4Cl , aqueous layer was extracted with EtOAc, EtOAc layer was combined, dried and concentrated to give crude product, which was purified by column (100% EtOAc) to give (S)-1-benzyl-5-(hydroxymethyl)pyrrolidin-2-one (170 mg).

[0358] Step 2: To a solution of (S)-1-benzyl-5-(hydroxymethyl)pyrrolidin-2-one (170 mg, 0.83 mmol) and 2,6-dihydroxybenzaldehyde (0.15 g, 1.08 mmol) in THF (6 mL) was added PPh_3 (polymer supported, 1.0 g, 1.25 mmol) and DIAD (0.24 mL, 1.25 mmol) at 0 °C. Then it was warmed up to room temperature and stirred for 1 h, AcCN was added to dilute the mixture, the insoluble material was filtered off, the filtrate was concentrated and was purified by preparative HPLC to give (S)-2-((1-benzyl-5-oxopyrrolidin-2-yl)methoxy)-6-hydroxybenzaldehyde (95 mg). ^1H NMR (400 MHz, Chloroform-*d*) δ 11.93 (t, J = 0.4 Hz, 1H), 10.17 (t, J = 0.5 Hz, 1H), 7.39 – 7.29 (m, 1H), 7.29 – 7.18 (m, 5H), 6.54 (dt, J = 8.5, 0.7 Hz, 1H), 6.17 (dd, J = 8.3, 0.9 Hz, 1H), 4.85 (d, J = 15.2 Hz, 1H), 4.28 (d, J = 15.2 Hz, 1H), 3.99 (d, J = 4.0 Hz, 2H), 3.92 (td, J = 7.8, 7.2, 3.6 Hz, 1H), 2.68 – 2.45 (m, 2H), 2.33 – 2.19 (m, 1H), 2.10 – 1.96 (m, 1H). MS ($\text{M}+\text{H}$) found for $\text{C}_{19}\text{H}_{19}\text{NO}_4$: 326.4.

GBT1093



[0359] GBT1093 - (S)-2-hydroxy-6-((5-oxo-4-phenylmorpholin-3-yl)methoxy)benzaldehyde

[0360] Step 1: To a solution of (S)-benzyl 2-amino-3-hydroxypropanoate hydrochloride (5 g, 21.58 mmol) in THF-water (1/1, 80 mL) was added K₂CO₃ (8.95 g, 64.74 mmol) and chloroacetyl chloride (2.92 mL, 36.69 mmol). After stirred for 1 h, it was diluted with EtOAc, organic layer was washed with water, brine, dried and concentrated to give benzyl (2-chloroacetyl)-L-serinate (5 g).

[0361] Step 2: A solution of (S)-benzyl 2-(2-chloroacetamido)-3-hydroxypropanoate (2.55 g, 9.39 mmol) in iPrOH (20 mL) was added to KOtBu (3.16 g, 28.17 mmol) in iPrOH (15 mL) at room temperature. After stirred at room temperature for 1 h, the mixture was quenched with 6N HCl (10 mL) at 0 °C, extracted with EtOAc, organic layers were combined, washed with brine, dried and concentrated to give isopropyl (2-chloroacetyl)-L-serinate as crude product.


[0362] Step 3: To a solution of (S)-benzyl 5-oxomorpholine-3-carboxylate in EtOH (7 mL) was added NaBH₄ (150 mg) at 0 °C, after stirred at room temperature for 3 h, it was quenched with NH₄Cl (220 mg in 0.6 mL water), and the insoluble material was filtered off, the filtrate was concentrated and was purified by column (DCM/MeOH=100:0 to 80:20) to give (R)-5-(hydroxymethyl)morpholin-3-one (100 mg).

[0363] Step 4: To a suspension of (R)-5-(hydroxymethyl)morpholin-3-one (100 mg, 2 mmol) and 3-iodopyridine (0.57 g, 2.8 mmol) in Dioxane (4 mL) was added CuI (0.08 g, 0.4 mmol), N,N-dimethylethylenediamine (0.05 mL, 0.4 mmol), K₃PO₄ (0.85 g, 4 mmol). After heating at 100 °C for 24 h, the mixture was cooled and was diluted with EtOAc, insoluble material was filtered off, the filtrate was concentrated and purified by column

(Hexanes/EtOAc=100:0 to 0:100) to give (S)-5-(hydroxymethyl)-1-(pyridin-3-yl)pyrrolidin-2-one (55 mg).

[0364] Step 5: To a solution of (S)-5-(hydroxymethyl)-1-(pyridin-3-yl)pyrrolidin-2-one (55 mg, 0.29 mmol) and 2,6-dihydroxybenzaldehyde (0.05 g, 0.38 mmol) in THF (2 mL) was added PPh₃ (polymer supported, 367 mg, 0.44 mmol) and DIAD (0.09 mL, 0.44 mmol) at 0 °C. Then it was warmed up to room temperature and stirred for 1 h, AcCN was added to dilute the mixture, the insoluble material was filtered off, the filtrate was concentrated and was purified by preparative HPLC to give (S)-2-hydroxy-6-((5-oxo-4-phenylmorpholin-3-yl)methoxy)benzaldehyde (29 mg). ¹H NMR (400 MHz, Chloroform-*d*) δ 11.88 (d, *J* = 0.4 Hz, 1H), 9.94 (d, *J* = 0.6 Hz, 1H), 7.53 – 7.40 (m, 2H), 7.40 – 7.30 (m, 2H), 7.26 (s, 2H), 6.53 (dt, *J* = 8.5, 0.7 Hz, 1H), 6.20 (dd, *J* = 8.3, 0.8 Hz, 1H), 4.47 (dd, *J* = 16.9, 0.9 Hz, 1H), 4.40 – 4.25 (m, 3H), 4.25 – 4.16 (m, 1H), 4.15 – 4.07 (m, 2H). MS (M+H) found for C₁₈H₁₇NO₅: 328.3.

GBT1123



[0365] GBT1123- *tert*-butyl (2-formyl-3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)phenyl)carbamate

[0366] Step 1: *tert*-Butyl (3-methoxyphenyl)carbamate (0.517 g, 2.3 mmol) was dissolved in dry diethyl ether (12 mL) and cooled to -40 °C in a solvent bath. *t*-Butyl lithium (4.1 mL of a 1.7 M pentane solution, 6.95 mmol) was added dropwise, the reaction was allowed to warm to -20 °C and stirred for 2 h more. The reaction was cooled to ~-78 °C, DMF (0.54 mL, 6.95

mmol) was added, and the reaction was allowed to gradually warm to 25 °C over 16 h. The reaction mixture was cooled in an ice bath and ammonium chloride solution (10 ml) was added. The reaction was extracted with ethyl acetate (3 x 80 ml), the combined organic phases were washed with a saturated aqueous sodium chloride solution (50 ml) and dried over sodium sulfate. After evaporation, the residue was purified by silica gel chromatography (0 – 30% ethyl acetate/hexanes) to give 0.46 g (79%) of *tert*-butyl (2-formyl-3-methoxyphenyl)carbamate as a lightly-colored solid. MS (ESI) m/z 252 $[M+H]^+$.


[0367] Step 2: *tert*-Butyl (2-formyl-3-methoxyphenyl)carbamate (0.38 g, 1.5 mmol) was dissolved in dichloromethane (10 ml) and added dropwise to a solution of aluminum chloride (1 g, 7.56 mmol) in dichloromethane (15 ml) while stirring in an ice bath. The yellowish solution was then stirred in a heat block at 50 °C. After 4 h, ice and water were added and the mixture stirred for 15 m. The reaction mixture was extracted with dichloromethane (3 x 50 ml), the combined organic phases were washed with water (30 ml) and a saturated aqueous sodium chloride solution (30 ml), and dried over sodium sulfate. After evaporation the crude residue was carried directly into the next step. MS (ESI) m/z 138 $[M+H]^+$.

[0368] Step 3: 2-Amino-6-hydroxybenzaldehyde (0.207 g, 1.5 mmol) was dissolved in THF (4 ml). Di-*tert*-butyldicarbonate (0.726 g, 3.3 mmol) and 4-dimethylaminopyridine (37 mg, 0.3 mmol) were added and the reaction was stirred for 18 h. The solution was evaporated and the residue was purified by silica gel chromatography (0 – 40% ethyl acetate/hexanes) to give 50 mg (14%) of *tert*-butyl (2-formyl-3-hydroxyphenyl)carbamate. MS (ESI) m/z 238 $[M+H]^+$.

[0369] Step 4: *tert*-Butyl (2-formyl-3-hydroxyphenyl)carbamate (50 mg, 0.21 mmol) and 3-(chloromethyl)-2-((1-isopropyl-1H-pyrazol-5-yl)pyridin-1-ium chloride (57 mg, 0.21 mmol) were dissolved in DMF (3 ml). The solution was purged with Ar gas. Potassium carbonate (116 mg 0.84 mmol) was added and the reaction mixture was stirred in a 60 °C heat block for 18 h. The reaction was cooled and water (50 ml) and ethyl acetate (50 ml) were added. The phases were separated and the aqueous phase was extracted with ethyl acetate (2 x 50 ml). The combined organic phases were washed with a saturated aqueous sodium chloride solution and dried over sodium sulfate. After evaporation the residue was purified by silica gel chromatography (5 – 50% ethyl acetate/hexanes) to give 5 mg (5%) of *tert*-butyl (2-formyl-3-((2-((1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)phenyl)carbamate as a white solid after lyophilization from acetonitrile/water. ^1H NMR (400 MHz, CDCl_3) δ 10.51 (s, 1H), 8.74 (d, J = 4.81 Hz, 1H), 8.06 (d, J = 7.84 Hz, 1H), 7.60 (s, 1H), 7.48 (t, J = 8.22

Hz, 1H), 7.44 – 7.39 (m, 1H), 6.91 – 6.81 (m, 2H), 6.36 (s, 1H), 5.10 (s, 2H), 4.64 (d, $J = 6.51$ Hz, 1H), 1.47 (d, $J = 4.57$ Hz, 6H), 1.41 (s, 9H). MS (ESI) m/z 437 $[M+H]^+$.

GBT1131



[0370] GBT1131- (S)-2-hydroxy-6-((6-oxo-1-phenylpiperidin-2-yl)methoxy)benzaldehyde


[0371] Step 1: To a suspension of (S)-6-oxopiperidine-2-carboxylic acid (1.0 g, 6.99 mmol) in EtOH (4 mL) at 0 °C was added SOCl_2 (0.61 mL, 8.39 mmol). After stirred at room temperature for 3 h, it was concentrated to remove all solvents, dried under high vacuum to give corresponding ethyl ester. The ester was dissolved in EtOH (15 mL) and was added NaBH_4 (300 mg) at 0 °C, the mixture was warmed up to room temperature and stirred for additional 15 h. The mixture was quenched with Sat. NH_4Cl , filtered off the insolubles and the filtrate was concentrated to give crude product, which was purified by column (DCM/MeOH=90:10) to give (S)-6-(hydroxymethyl)piperidin-2-one (450 mg) as white solid.

[0372] Step 2: To a suspension of (S)-6-(hydroxymethyl)piperidin-2-one (150 mg, 1.16 mmol) and iodobenzene (0.35 g, 1.74 mmol) in Dioxane (2 mL) was added CuI (0.09 g, 0.46 mmol), N,N-dimethylethylenediamine (0.04 mL, 0.35 mmol), K_3PO_4 (0.49 g, 2.32 mmol). After heating at 100 °C for 24 h, the mixture was cooled and was diluted with EtOAc, organic layer was washed with brine, dried (Na_2SO_4) and concentrated to give crude product, which was purified by column (Hexanes/EtOAc=100:0 to 0:100) to give (S)-6-(hydroxymethyl)-1-phenylpiperidin-2-one (85 mg).

[0373] Step 3: To a solution of (S)-6-(hydroxymethyl)-1-phenylpiperidin-2-one (85 mg, 0.44 mmol) and 2,6-dihydroxybenzaldehyde (0.08 g, 0.57 mmol) in THF (5 mL) was added PPh_3 (polymer supported, 550 mg, 0.66 mmol) and DIAD (0.13 mL, 0.66 mmol) at 0 °C. After warmed to room temperature and further stirred for 2 h, it was diluted with AcCN, the insoluble material was filtered off and the filtrate was concentrated and subjected to preparative HPLC to give (S)-2-hydroxy-6-((6-oxo-1-phenylpiperidin-2-yl)methoxy)benzaldehyde (31 mg). ^1H NMR (400 MHz, Chloroform- d) δ 11.90 (d, $J = 0.4$ Hz, 1H), 10.08 (d, $J = 0.6$ Hz, 1H), 7.44 – 7.36 (m, 2H), 7.36 – 7.27 (m, 2H), 7.22 – 7.14 (m,


2H), 6.60 – 6.43 (m, 1H), 6.11 (dd, $J = 8.3, 0.8$ Hz, 1H), 4.25 (qd, $J = 5.7, 4.2$ Hz, 1H), 4.04 – 3.81 (m, 2H), 2.77 – 2.53 (m, 2H), 2.29 – 1.87 (m, 4H). MS (M+H) found for $C_{19}H_{19}NO_4$: 326.5.

GBT001717




6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2,3-dimethoxybenzaldehyde

[0374] GBT1717- 6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2,3-dimethoxybenzaldehyde. The compound was prepared by O-alkylation of 6-hydroxy-2,3-dimethoxybenzaldehyde (A) and 3-(chloromethyl)-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridine hydrochloride salt (INT-2) according to scheme 9, reaction step 4. The product as white solid was obtained after flash column purification. ^1H NMR (400 MHz, Chloroform- d) δ 10.49 (t, $J = 0.4$ Hz, 1H), 8.69 (dd, $J = 4.8, 1.7$ Hz, 1H), 8.29 (ddd, $J = 8.0, 1.7, 0.9$ Hz, 1H), 7.61 (dd, $J = 1.9, 0.5$ Hz, 1H), 7.44 (dd, $J = 7.9, 4.8$ Hz, 1H), 7.03 (d, $J = 9.0$ Hz, 1H), 6.52 (d, $J = 7.9$ Hz, 1H), 6.36 (dd, $J = 1.9, 0.4$ Hz, 1H), 4.98 (s, 2H), 4.59 (hept, $J = 6.7$ Hz, 1H), 3.95 (d, $J = 0.4$ Hz, 3H), 3.84 (s, 3H), 1.46 (d, $J = 6.6$ Hz, 6H). MS (M+H) found for $C_{21}H_{23}N_3O_4$: 382.5.




Preparation of intermediate A



[0375] Step 1: To a suspension of 3,4-dimethoxyphenol (2.0 g, 12.97 mmol) in dihydropyran (2 mL, 21.87 mmol) at ambient temperature was added 1 drop of Conc. HCl. After stirred for 1 h, the solution was diluted with EtOAc, organic layer was washed with Sat. NaHCO₃, brine, dried and concentrated to give crude product, which was purified by column (Hexanes/EtOAc = 65:35) to give 2-(3,4-dimethoxyphenoxy)tetrahydro-2H-pyran (2.9 g).


[0376] Step 2: To a solution of 2-(3,4-dimethoxyphenoxy)tetrahydro-2H-pyran (1.0 g, 4.2 mmol) in THF (6 mL) at 0 °C was added TMEDA (0.72 mL, 4.83 mmol) and BuLi (2.5 M, 2.02 mL, 5.05 mmol). After stirred for 1.5 h at 0 °C, it was added DMF (1.3 mL). After stirred for 1 h, the mixture was added 6N HCl (4 mL), and was stirred for 1 h at ambient temperature, additional 12N HCl (2 mL) was added to drive the reaction to completion. The solution was diluted with EtOAc and water, organic layer was separated and washed with brine, dried and concentrated to give crude product, which was purified by column chromatography to give 6-hydroxy-2,3-dimethoxybenzaldehyde (490 mg).

GBT001659




2,3-dihydroxy-6-((2-((1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde

[0377] GBT1659- 2,3-dihydroxy-6-((2-((1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde.




[0378] To a solution of 6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2,3-dimethoxybenzaldehyde (24 mg, 0.05 mmol) was added BBr₃ (1M, 0.5 mL), after stirred at ambient temperature for 30 min, the mixture was concentrated and the crude product was purified by preparative HPLC to give 2,3-dihydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde (10 mg). ¹H NMR (400 MHz, Chloroform-*d*) δ 12.01 (s, 1H), 10.32 (d, *J* = 0.5 Hz, 1H), 8.74 (dd, *J* = 4.8, 1.7 Hz, 1H), 8.00 – 7.92 (m, 1H), 7.60 (dd, *J* = 1.9, 0.6 Hz, 1H), 7.41 (dd, *J* = 7.9, 4.8 Hz, 1H), 7.03 (d, *J* = 8.8 Hz, 1H), 6.34 (d, *J* = 1.9 Hz, 1H), 6.19 (d, *J* = 8.8 Hz, 1H), 5.29 (s, 1H), 5.02 (s, 2H), 4.64 (hept, *J* = 6.6 Hz, 1H), 1.47 (d, *J* = 6.6 Hz, 6H). MS (M+H) found for C₁₉H₁₉N₃O₄, 354.4.

GBT001718




2-hydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-3-methoxybenzaldehyde

[0379] GBT1718- 2-hydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-3-methoxybenzaldehyde




[0380] To a solution of 6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2,3-dimethoxybenzaldehyde (50.00 mg; 0.13 mmol) in DCM (0.8 mL) at ambient temperature was added boron tribromide (0.13 mL; 1.00 mol/L). After stirred for 5 min, the red mixture was concentrated and the crude product was purified by preparative HPLC to give 2-hydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-3-methoxybenzaldehyde (15 mg). ¹H NMR (400 MHz, Chloroform-*d*) δ 12.15 (d, *J* = 0.6 Hz, 1H), 10.35 (s, 1H), 8.74 (dd, *J* = 4.8, 1.7 Hz, 1H), 7.97 (ddd, *J* = 7.9, 1.6, 0.7 Hz, 1H), 7.60 (dd, *J* = 1.8, 0.5 Hz, 1H), 7.41 (dd, *J* = 7.9, 4.8 Hz, 1H), 6.98 – 6.94 (m, 1H), 6.34 (d, *J* = 1.9 Hz, 1H), 6.18 (d, *J* = 8.9 Hz, 1H), 5.02 (s, 2H), 4.72 – 4.57 (m, 1H), 3.84 (s, 3H), 1.47 (d, *J* = 6.6 Hz, 6H). MS (M+H) found for C₂₀H₂₁N₃O₄: 368.4.


GBT001723



[0381] GBT001723 - 2-((6-((2-(dimethylamino)ethyl)(methylamino)-2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-6-hydroxybenzaldehyde. The compound was prepared by Mitsunobu coupling of (6-((2-(dimethylamino)ethyl)(methylamino)-2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methanol (A) with 2,6-dihydroxybenzaldehyde (INT-3) according to scheme 9, reaction step 3. The product as green oil was obtained after flash column purification. ¹H NMR (400 MHz, CDCl₃, ppm): 11.87 (*br s*, 1H), 10.21 (*s*, 1H), 7.54 (d, *J* = 8.8 Hz, 1H), 7.44 (*s*, 1H), 7.26 (t, *J* = 8.4 Hz, 1H), 6.50 (d, *J* = 8.8 Hz, 1H), 6.41 (d, *J* = 8.4 Hz, 1H), 6.20 (d, *J* = 8.4 Hz, 1H), 6.18 (*s*, 1H), 4.82 (*s*, 2H), 4.90-4.60 (m, 1H), 3.61 (t, *J* = 11.2 Hz, 2H), 3.03 (*s*, 3H), 2.43 (t, *J* = 11.2 Hz, 2H), 2.20 (*s*, 6H), 1.39 (d, *J* = 6.8 Hz, 6H); MS(ES, *m/z*): 438.4 [M + 1]⁺



[0382] Intermediate A was prepared followed the scheme below:



[0383] Step 1: Into a 25-mL round-bottom flask, was placed a solution of methyl 6-bromo-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridine-3-carboxylate (1.3 g, 4.01 mmol, 1.00 equiv) in tetrahydrofuran (15 mL). DIPEA (1.55 g, 3.00 equiv) and [2-(dimethylamino)ethyl](methyl)amine (2.4 g, 23.49 mmol, 5.00 equiv) were added to the reaction mixture. The resulting solution was heated to reflux for 24 hr, and then it was quenched with 30 mL of water. The resulting solution was extracted with 3x50 mL of ethyl acetate. The combined organic layers were washed with 1x80 mL of water and 1x80 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/methanol (30:1-15:1) as eluent to furnish 1.22 g (88%) of methyl 6-[[2-(dimethylamino)ethyl](methyl)amino]-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridine-3-carboxylate as a light yellow oil.

Step2. Into a 100-mL round-bottom flask, was placed a solution of methyl 6-[[2-(dimethylamino)ethyl](methyl)amino]-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]pyridine-3-carboxylate (1.2 g, 3.47 mmol, 1.00 equiv) in tetrahydrofuran (20 mL). This was followed by the addition of LAH (330 mg, 8.70 mmol, 2.50 equiv), in portions at 0 °C. The resulting solution was stirred for 4 h at room temperature. The reaction was quenched by the addition of 0.5 mL water, 0.5 mL 2.5M sodium hydroxide (aq.). The resulting solution was diluted with 50 mL of ethyl acetate. The solids were filtered out and washed with THF for 3 times.

The resulting mixture was concentrated under vacuum. This resulted in 1.1 g (100%) of (6-[[2-(dimethylamino)ethyl][(methyl)amino]-2-[1-(propan-2-yl)-1H-pyrazol-5-yl]]pyridin-3-yl)methanol as a colorless oil.

[0384] From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention.





[0385] Throughout the description of this invention, reference is made to various patent applications and publications, each of which are herein incorporated by reference in their entirety.






[0256] As used herein, Table 1 includes compounds described below or tautomers or pharmaceutically acceptable salts thereof:






2-(imidazo[1,2-a]pyridin-8-ylmethoxy)-5-methoxybenzaldehyde,






2-(imidazo[1,2-a]pyridin-2-ylmethoxy)-5-methoxybenzaldehyde,






2-(imidazo[1,5-a]pyridin-8-ylmethoxy)-5-methoxybenzaldehyde,
5-methoxy-2-(quinolin-5-ylmethoxy)benzaldehyde,
5-methoxy-2-((1-methyl-1H-indazol-4-yl)methoxy)benzaldehyde,
5-methoxy-2-((8-methylimidazo[1,2-a]pyridin-2-yl)methoxy)benzaldehyde,
2-((1H-indazol-4-yl)methoxy)-5-methoxybenzaldehyde,
5-methoxy-2-(pyridin-3-ylmethoxy)benzaldehyde,
2-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-5-methoxybenzaldehyde,
2-hydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde,
2-((3-(2H-tetrazol-5-yl)benzyl)oxy)-6-hydroxybenzaldehyde,
2-((4-(2H-tetrazol-5-yl)benzyl)oxy)-6-hydroxybenzaldehyde,
methyl 4-((2-formylphenoxy)methyl)benzoate,
4-((2-formylphenoxy)methyl)benzoic acid,
methyl 3-((2-formylphenoxy)methyl)benzoate,
2-bromo-3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde,
2-hydroxy-6-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde,
2-hydroxy-6-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde,
2-fluoro-6-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde,
2-fluoro-6-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde,
2-fluoro-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde, and
1-(2-formyl-3-hydroxyphenethyl)piperidine-4-carboxylic acid, or a tautomer or
pharmaceutically acceptable salt thereof.






Compound	Structure	Name
1		2-(imidazo[1,2-a]pyridin-8-ylmethoxy)-5-methoxybenzaldehyde
2		4-formyl-3-(imidazo[1,2-a]pyridin-8-ylmethoxy)benzonitrile
3		2-(imidazo[1,2-a]pyridin-8-ylmethoxy)-4-methoxybenzaldehyde
4		2-(imidazo[1,2-a]pyridin-6-ylmethoxy)-5-methoxybenzaldehyde






5		2-(imidazo[1,2-a]pyridin-2-ylmethoxy)-5-methoxybenzaldehyde
6		2-(imidazo[1,5-a]pyridin-8-ylmethoxy)-4-methoxybenzaldehyde
7		2-(imidazo[1,5-a]pyridin-8-ylmethoxy)-5-methoxybenzaldehyde
8		2-(imidazo[1,2-a]pyridin-7-ylmethoxy)-5-methoxybenzaldehyde
9		2-(imidazo[1,2-a]pyridin-3-ylmethoxy)-5-methoxybenzaldehyde







10		5-methoxy-2-(quinolin-5-ylmethoxy)benzaldehyde
11		5-bromo-2-(imidazo[1,2-a]pyridin-8-ylmethoxy)benzaldehyde
12		4-chloro-2-(imidazo[1,2-a]pyridin-8-ylmethoxy)benzaldehyde
13		2-(imidazo[1,2-a]pyridin-8-ylmethoxy)benzaldehyde
14		4-fluoro-2-(imidazo[1,2-a]pyridin-8-ylmethoxy)benzaldehyde






15		2-(imidazo[1,2-a]pyridin-8-ylmethoxy)-3-methoxybenzaldehyde
16		2-(imidazo[1,2-a]pyridin-8-ylmethoxy)-5-methylbenzaldehyde
17		5-methoxy-2-(pyrrolo[1,2-a]pyrazin-4-ylmethoxy)benzaldehyde
18		2-(imidazo[1,5-a]pyridin-6-ylmethoxy)-4-methoxybenzaldehyde
19		2-(imidazo[1,5-a]pyridin-5-ylmethoxy)-5-methoxybenzaldehyde






20		3-formyl-4-(imidazo[1,5-a]pyridin-5-ylmethoxy)benzonitrile
21		2-((1H-pyrrolo[2,3-b]pyridin-4-yl)methoxy)-5-methoxybenzaldehyde
22		5-ethyl-2-(imidazo[1,2-a]pyridin-8-ylmethoxy)benzaldehyde
23		5-methoxy-2-((1-methyl-1H-indazol-4-yl)methoxy)benzaldehyde
24		5-methoxy-2-((8-methylimidazo[1,2-a]pyridin-2-yl)methoxy)benzaldehyde






25		2-((1H-indazol-4-yl)methoxy)-5-methoxybenzaldehyde
26		2-((1H-pyrrolo[2,3-b]pyridin-4-yl)methoxy)-5-methoxybenzaldehyde
27		3-formyl-4-(imidazo[1,2-a]pyridin-8-ylmethoxy)benzonitrile
28		5-methoxy-2-(pyrrolo[1,2-a]pyrazin-6-ylmethoxy)benzaldehyde
29		6-((2-formyl-4-methoxyphenoxy)methyl)pyrrolo[1,2-a]pyrazine-7-carbonitrile







30		6-((2-formyl-4-methoxyphenoxy)methyl)pyrrolo[1,2-a]pyrazine-7-carboxamide
31		2-((1H-pyrazolo[3,4-b]pyridin-4-yl)methoxy)-5-methoxybenzaldehyde
32		5-methoxy-2-(pyrazolo[1,5-a]pyrazin-3-ylmethoxy)benzaldehyde
33		5-methoxy-2-(pyrrolo[1,2-a]pyrazin-6-ylmethoxy)benzaldehyde
34		2-(imidazo[1,5-a]pyridin-6-ylmethoxy)-5-methoxybenzaldehyde






35		3-(3-formyl-4-(imidazo[1,2-a]pyridin-8-ylmethoxy)benzonitrile
36		3-(imidazo[1,2-a]pyridin-8-ylmethyl)-1,3-dihydroisobenzofuran-1-ol
37		2-(2-(3-(5-methoxybenzaldehyde)oxy)imidazo[1,2-a]pyridin-5-ylmethoxy)-5-methoxybenzaldehyde
38		N-(2-(3-(5-methoxybenzaldehyde)oxy)imidazo[1,2-a]pyridin-5-yl)-2-(3-(5-methoxybenzaldehyde)oxy)imidazo[1,2-a]pyridin-5-ylmethoxy)-5-methoxybenzaldehyde
39		N-(2-(3-(5-methoxybenzaldehyde)oxy)imidazo[1,2-a]pyridin-5-yl)-2-(3-(5-methoxybenzaldehyde)oxy)imidazo[1,2-a]pyridin-5-ylmethoxy)-5-methoxybenzaldehyde
40		2-(3-(5-methoxybenzaldehyde)oxy)imidazo[1,2-a]pyridin-5-yl)-2-(3-(5-methoxybenzaldehyde)oxy)imidazo[1,2-a]pyridin-5-ylmethoxy)-5-methoxybenzaldehyde






41		5-methoxy-2-(pyridin-3-ylmethoxy)benzaldehyde
42		4-((2-formyl-3-hydroxyphenoxy)methyl)benzoic acid
43		2-hydroxy-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde
44		2-((3-(2H-tetrazol-5-yl)benzyl)oxy)-6-hydroxybenzaldehyde
45		2-((4-(2H-tetrazol-5-yl)benzyl)oxy)-6-hydroxybenzaldehyde







46		methyl 4-((2-formylphenoxy)methyl)benzoate
47		4-((2-formylphenoxy)methyl)benzoic acid
48		methyl 3-((2-formylphenoxy)methyl)benzoate
49		2-bromo-3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde
50		2-hydroxy-6-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde







51		2-hydroxy-6-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde
52		2-fluoro-6-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde
53		2-fluoro-6-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde
54		2-fluoro-6-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)benzaldehyde
55		1-(2-formyl-3-hydroxyphenethyl)piperidine-4-carboxylic acid






1		4-(pyridin-3-ylmethoxy)nicotinaldehyde
2		3-(pyridin-3-ylmethoxy)isonicotinaldehyde
3		2-(imidazo[1,2-a]pyridin-8-ylmethoxy)nicotinaldehyde
4		3-(imidazo[1,2-a]pyridin-8-ylmethoxy)picolinaldehyde
5		5-(imidazo[1,2-a]pyridin-8-ylmethoxy)-2-methoxyisonicotinaldehyde
6		3-(imidazo[1,2-a]pyridin-8-ylmethoxy)isonicotinaldehyde






7		3-((imidazo[1,5-a]pyridin-8-ylmethoxy)isonicotinaldehyde
8		2-methoxy-5-(pyrazolo[1,5-a]pyrazin-3-ylmethoxy)isonicotinaldehyde
9		8-((3-formylpyridin-2-yl)oxy)methylimidazo[1,2-a]pyridine-6-carboxamide
10		8-((4-formyl-6-methoxypyridin-3-yl)oxy)methylimidazo[1,2-a]pyridine-6-carboxamide
11		5-(imidazo[1,2-a]pyridin-8-ylmethoxy)-2-oxo-1,2-dihydropyridine-4-carbaldehyde






12		2-(2-(imidazo[1,2-a]pyridin-8-yl)ethyl)nicotinaldehyde
13		5-(2-(imidazo[1,2-a]pyridin-8-yl)ethyl)-2-methoxyisonicotinaldehyde
14		5-((1H-pyrazolo[3,4-b]pyridin-4-yl)methoxy)-2-methoxyisonicotinaldehyde
15		3-((4-formyl-6-methoxypyridin-3-yl)oxy)methylpyrazolo[1,5-a]pyrazine-2-carboxamide
16		5-((2-(1H-pyrazol-5-yl)pyrazolo[1,5-a]pyrazin-3-yl)methoxy)-2-methoxyisonicotinaldehyde






17		2-(imidazo[1,2-a]pyridin-2-ylmethoxy)nicotinaldehyde
18		2-methoxy-5-((4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-3-yl)methoxy)isonicotinaldehyde
19		2-(imidazo[1,2-a]pyridin-8-ylmethoxy)nicotinaldehyde
20		5-(imidazo[1,2-a]pyridin-8-ylmethoxy)-2-methylisonicotinaldehyde
21		3-((1H-pyrrolo[2,3-b]pyridin-4-yl)methoxy)isonicotinaldehyde
22		3-(imidazo[1,2-a]pyridin-8-ylmethoxy)isonicotinaldehyde






23		3-((pyrrolo[1,2-a]pyrazin-6-yl)methoxy)isonicotinaldehyde
24		6-((4-formylpyridin-3-yloxy)methyl)pyrrolo[1,2-a]pyrazine-7-carbonitrile
25		6-((4-formylpyridin-3-yloxy)methyl)pyrrolo[1,2-a]pyrazine-7-carboxamide
26		3-((1H-pyrazolo[3,4-b]pyridin-4-yl)methoxy)isonicotinaldehyde
27		3-(pyrazolo[1,5-a]pyrazin-3-yl)methoxy)isonicotinaldehyde
28		2-methoxy-5-((6-oxo-1,6-dihydropyridin-3-yl)methoxy)isonicotinaldehyde





29		2-methoxy-5-((2-oxo-1,2-dihydropyridin-4-yl)methoxy)isonicotinaldehyde
30		2-methoxy-5-(oxazol-5-ylmethoxy)isonicotinaldehyde
31		5-((1H-imidazol-5-yl)methoxy)-2-methoxyisonicotinaldehyde
32		5-((1H-imidazol-2-yl)methoxy)-2-methoxyisonicotinaldehyde
33		5-((4H-1,2,4-triazol-3-yl)methoxy)-2-methoxyisonicotinaldehyde





34		5-((1H-tetrazol-5-yl)methoxy)-2-methoxyisonicotinaldehyde
35		5-((1H-pyrazol-5-yl)methoxy)-2-methoxyisonicotinaldehyde
36		5-((1H-pyrazol-4-yl)methoxy)-2-methoxyisonicotinaldehyde
37		2-methoxy-5-(oxazol-4-ylmethoxy)isonicotinaldehyde
38		2-methoxy-5-((2-methylpyridin-3-yl)methoxy)isonicotinaldehyde






39		2-methoxy-5-((4-methylpyridin-3-yl)methoxy)isonicotinaldehyde
40		2-methoxy-5-((6-(trifluoromethyl)pyridin-3-yl)methoxy)isonicotinaldehyde
41		2-methoxy-5-((6-methylpyridin-3-yl)methoxy)isonicotinaldehyde
42		2-methoxy-5-(pyridin-3-yl)methoxy)isonicotinaldehyde
43		2-methoxy-5-((5-methylpyridin-3-yl)methoxy)isonicotinaldehyde

44		5-(isoquinolin-1-ylmethoxy)-2-methoxyisonicotinaldehyde
45		2-methoxy-5-(quinolin-2-ylmethoxy)isonicotinaldehyde
46		2-methoxy-5-(pyridin-4-ylmethoxy)isonicotinaldehyde
47		2-methoxy-5-((3-methylpyridin-4-yl)methoxy)isonicotinaldehyde
48		5-((3-bromopyridin-4-yl)methoxy)-2-methoxyisonicotinaldehyde





49		3-(imidazo[1,2-a]pyridin-8-ylmethoxy)-6-methylpicolinaldehyde
50		(5-(imidazo[1,2-a]pyridin-5-ylmethoxy)-2-methoxypyridin-4-yl)(methoxy)methanol
51		N-(4-formylpyridin-3-yl)imidazo[1,2-a]pyridine-8-carboxamide
52		2-methoxy-5-((6-(trifluoromethyl)imidazo[1,2-a]pyridin-2-yl)methoxy)isonicotinaldehyde
53		methyl 2-((4-formyl-6-methoxypyridin-3-yloxy)methyl)imidazo[1,2-a]pyridine-8-carboxylate





54		2-methoxy-5-((1-methyl-2-oxo-1,2-dihydropyridin-4-yl)methoxy)isonicotinaldehyde
55		5-((3-bromoimidazo[1,2-a]pyridin-2-yl)methoxy)-2-methoxyisonicotinaldehyde
56		5-((6-bromoimidazo[1,2-a]pyridin-2-yl)methoxy)-2-methoxyisonicotinaldehyde
57		5-((8-bromoimidazo[1,2-a]pyridin-2-yl)methoxy)-2-methoxyisonicotinaldehyde





58		2-methoxy-5-((3-methyl-[1,2,4]triazolo[4,3-a]pyridin-8-yl)methoxy)isonicotinaldehyde
59		5-((3-(1H-pyrazol-5-yl)imidazo[1,2-a]pyridin-2-yl)methoxy)-2-methoxyisonicotinaldehyde
60		5-((6-(1H-pyrazol-3-yl)imidazo[1,2-a]pyridin-2-yl)methoxy)-2-methoxyisonicotinaldehyde
61		2-methoxy-5-((8-(1-methyl-1H-pyrazol-5-yl)imidazo[1,2-a]pyridin-2-yl)methoxy)isonicotinaldehyde





62		5-((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)picolinonitrile
63		5-((2-bromopyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
64		3-((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)picolinonitrile
65		5-((2-(1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
66		5-((5-bromopyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde






67		methyl 2-((4-(1,3-dioxolan-2-yl)-6-methoxypyridin-3-yl)oxy)methyl)imidazo[1,2-a]pyridine-8-carboxylate
68		2-((4-(1,3-dioxolan-2-yl)-6-methoxypyridin-3-yl)oxy)methyl)imidazo[1,2-a]pyridine-8-carboxamide
69		2-((4-(1,3-dioxolan-2-yl)-6-methoxypyridin-3-yl)oxy)methyl)-N-methylimidazo[1,2-a]pyridine-8-carboxamide
70		5-(5-(1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde






71		5-((4-(1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
72		2-((4-(dihydroxymethyl)-6-methoxypyridin-3-yloxy)methyl)-N-methylimidazo[1,2-a]pyridine-8-carboxamide
73		2-((4-(dihydroxymethyl)-6-methoxypyridin-3-yloxy)methyl)imidazo[1,2-a]pyridine-8-carboxamide
74		2-methoxy-5-((5-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde






75		2-methoxy-5-((5-(1-methyl-1H-pyrazol-3-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
76		5-((5-(1H-pyrazol-4-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
77		2-methoxy-5-((5-(1-methyl-1H-pyrazol-4-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
78		methyl 5-((4-formyl-6-methoxypyridin-3-yloxy)methyl)nicotinate






79		5-((4-formyl-6-methoxypyridin-3-yloxy)methyl)nicotinic acid
80		2-methoxy-5-(quinolin-3-ylmethoxy)isonicotinaldehyde
81		6-methyl-3-(quinolin-3-ylmethoxy)picolinaldehyde
82		5-(isoquinolin-7-ylmethoxy)-2-methoxyisonicotinaldehyde





83		3-(isoquinolin-7-ylmethoxy)-6-methylpicolinaldehyde
84		2-methoxy-5-((1-methyl-1H-indazol-4-yl)methoxy)isonicotinaldehyde
85		6-methyl-3-((1-methyl-1H-indazol-4-yl)methoxy)picolinaldehyde
86		tert-butyl 4-((2-formyl-6-methylpyridin-3-yloxy)methyl)-1H-indazole-1-carboxylate






87		5-((1H-indazol-4-yl)methoxy)-2-methoxyisonicotinaldehyde
88		3-((1H-indazol-4-yl)methoxy)-6-methylpicolinaldehyde
89		6-methoxy-3-((1-methyl-1H-indazol-6-yl)methoxy)picolinaldehyde
90		2-methoxy-5-((1-methyl-1H-indazol-7-yl)methoxy)isonicotinaldehyde
91		6-methyl-3-((1-methyl-1H-indazol-6-yl)methoxy)picolinaldehyde






92		6-methyl-3-((1-methyl-1H-indazol-7-yl)methoxy)picolinaldehyde
93		3-(isoquinolin-1-ylmethoxy)-6-methylpicolinaldehyde
94		6-methyl-3-(quinolin-2-ylmethoxy)picolinaldehyde
95		5-((4-(1H-pyrazol-4-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
96		5-((6-bromoimidazo[1,2-a]pyridin-8-yl)methoxy)-2-methoxyisonicotinaldehyde





97		8-((4-formyl-6-methoxypyridin-3-yloxy)methyl)imidazo[1,2-a]pyridine-6-carbonitrile
98		5-((4-formyl-6-methoxypyridin-3-yloxy)methyl)nicotinonitrile
99		3-(benzo[d]oxazol-4-ylmethoxy)-6-methylpicolinaldehyde
100		8-((4-formyl-6-methoxypyridin-3-yloxy)methyl)imidazo[1,2-a]pyridine-6-carboxamide
101		5-((4-formyl-6-methoxypyridin-3-yloxy)methyl)nicotinamide






102		5-((6-(1H-pyrazol-4-yl)imidazo[1,2-a]pyridin-8-yl)methoxy)-2-methoxyisonicotinaldehyde
103		5-(benzo[d]oxazol-4-ylmethoxy)-2-methoxyisonicotinaldehyde
104		5-((6-(1H-pyrazol-5-yl)imidazo[1,2-a]pyridin-8-yl)methoxy)-2-methoxyisonicotinaldehyde
105		5-((1,5-naphthyridin-4-yl)methoxy)-2-methoxyisonicotinaldehyde
106		3-((1,5-naphthyridin-4-yl)methoxy)-6-methylpicolinaldehyde






107		5-((1H-indazol-5-yl)methoxy)-2-methoxyisonicotinaldehyde
108		6-methyl-3-((1-methyl-1H-indazol-5-yl)methoxy)picolinaldehyde
109		3-((3-chloro-1-methyl-1H-indazol-5-yl)methoxy)-6-methylpicolinaldehyde
110		2-methoxy-5-((1-methyl-1H-indazol-5-yl)methoxy)isonicotinaldehyde

111		5-((3-chloro-1-methyl-1H-indazol-5-yl)methoxy)-2-methoxyisonicotinaldehyde
112		N-(4-formyl-6-methoxypyridin-3-yl)imidazo[1,2-a]pyridine-8-carboxamide
113		3-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-4-yl)methoxy)-6-methylpicolinaldehyde
114		5-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-4-yl)methoxy)-2-methoxyisonicotinaldehyde
115		3-((4-formyl-6-methoxypyridin-3-yl)oxy)methylpicolinamide






116		5-((2-chloroquinolin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
117		5-((2-(1H-pyrazol-5-yl)quinolin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
118		2-methoxy-5-(quinoxalin-2-yl)methoxy)isonicotinaldehyde
119		6-methyl-3-(quinolin-5-yl)methoxy)picolinaldehyde
120		2-methoxy-5-(quinolin-5-yl)methoxy)isonicotinaldehyde






121		6-methyl-3-((1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)methoxy)picolinaldehyde
122		2-methoxy-5-((1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)methoxy)isonicotinaldehyde
123		5-((7-(1H-pyrazol-3-yl)imidazo[1,5-a]pyridin-8-yl)methoxy)-2-methoxyisonicotinaldehyde
124		5-((5-(2H-tetrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde






125		5-((6-(2H-tetrazol-5-yl)imidazo[1,2-a]pyridin-8-yl)methoxy)-2-methoxyisonicotinaldehyde
126		ethyl 2-(5-(imidazo[1,2-a]pyridin-8-yl)methoxy)-2-methoxypyridin-4-yl)thiazolidine-4-carboxylate
127		2-methoxy-5-((2-(1-methyl-1H-pyrazol-4-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
128		5-((2-(1H-pyrazol-4-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
129		2-methoxy-5-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde






130		2-methoxy-5-((2-(1-methyl-1H-pyrazol-3-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
131		5-((2-(2H-tetrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
132		2-methoxy-5-((2-(4-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
133		5-((3-(1H-pyrazol-5-yl)isoquinolin-4-yl)methoxy)-2-methoxyisonicotinaldehyde
134		5-((2-(1H-pyrazol-1-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde






135		3-((2-(1H-pyrazol-1-yl)pyridin-3-yl)methoxy)-6-methylpicolinaldehyde
136		6-methyl-3-(pyridin-3-ylmethoxy)picolinaldehyde
137		methyl 8-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)imidazo[1,2-a]pyridine-6-carboxylate
138		methyl 2-bromo-8-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)imidazo[1,2-a]pyridine-6-carboxylate
139		3-(imidazo[1,5-a]pyridin-8-ylmethoxy)-6-methylpicolinaldehyde






140		5-(imidazo[1,5-a]pyridin-8-ylmethoxy)-2-methoxyisonicotinaldehyde
141		(5-(methoxycarbonyl)pyridin-3-yl)methyl 5-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)nicotinate
142		5-((2-(1,4-dimethyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
143		5-((2-(1,5-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
144		2-hydroxyethyl 5-(((4-(1,3-dioxolan-2-yl)-6-methoxypyridin-3-yl)oxy)methyl)nicotinate






145		methyl 5-(((4-(1,3-dioxolan-2-yl)-6-methoxypyridin-3-yl)oxy)methyl)nicotinate
146		methyl 5-(((4-(bis(2-hydroxyethoxy)methyl)-6-methoxypyridin-3-yl)oxy)methyl)nicotinate
147		5-((2-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
148		5-((2-(1,3-dimethyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
149		5-((2-(1-ethyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde






150		5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
151		2-methoxy-5-((2-(3-methyl-1H-pyrazol-1-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
152		5-(((4-(1,3-dioxolan-2-yl)-6-methoxypyridin-3-yl)oxy)methyl)nicotinic acid
153		(E)-2-methoxy-5-((2-(4-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde oxime
154		(E)-2-methoxy-5-(pyridin-3-yl)methoxy)isonicotinaldehyde oxime






155		2-(5-(imidazo[1,2-a]pyridin-8-ylmethoxy)-2-methoxypyridin-4-yl)thiazolidine
156		1-(2-(5-(imidazo[1,2-a]pyridin-8-ylmethoxy)-2-methoxypyridin-4-yl)thiazolidin-3-yl)ethanone
157		5-((2-(4-(1H-pyrazol-3-yl)piperazin-1-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
158		2-(difluoromethoxy)-5-(imidazo[1,2-a]pyridin-8-ylmethoxy)isonicotinaldehyde
159		2-methoxy-5-((2-phenylpyridin-3-yl)methoxy)isonicotinaldehyde






160		5-((3-(1-isopropyl-1H-pyrazol-5-yl)pyridin-4-yl)methoxy)-2-methoxyisonicotinaldehyde
161		5-([2,3'-bipyridin]-3-ylmethoxy)-2-methoxyisonicotinaldehyde
162		2-methoxy-5-((2-(o-tolyl)pyridin-3-yl)methoxy)isonicotinaldehyde
163		2-methoxy-5-((2'-methoxy-[2,3'-bipyridin]-3-yl)methoxy)isonicotinaldehyde
164		methyl 4-(((2-formylpyridin-3-yl)oxy)methyl)benzoate






165		4-(((2-formyl-6-methylpyridin-3-yl)oxy)methyl)benzoic acid
166		4-(((2-formylpyridin-3-yl)oxy)methyl)benzoic acid
167		methyl 3-(((4-formylpyridin-3-yl)oxy)methyl)benzoate
168		methyl 3-(((2-formyl-6-methylpyridin-3-yl)oxy)methyl)benzoate
169		3-(((4-formylpyridin-3-yl)oxy)methyl)benzoic acid






170		3-(((2-formyl-6-methylpyridin-3-yl)oxy)methyl)benzoic acid
171		3-(((2-formylpyridin-3-yl)oxy)methyl)benzoic acid
172		2-methoxy-5-((2-(1-(2-methoxyethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
173		2-methoxy-5-((2-(1-propyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
174		2-methoxy-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde






175		5-((2-(1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
176		3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)picolinaldehyde
177		3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-6-methylpicolinaldehyde
178		2-(difluoromethoxy)-5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
179		5-(imidazo[1,2-a]pyridin-8-yl)methoxy)-2-(2-methoxyethoxy)isonicotinaldehyde






180		5-(2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy-2-(2-methoxyethoxy)isonicotinaldehyde
181		5-(2-(1-isopropyl-1H-pyrazol-5-yl)pyrazin-2-yl)methoxy-2-methoxyisonicotinaldehyde
182		3-(4-formyl-6-methoxypyridin-3-yloxy)methylpicolinate
183		5-(2-(2-hydroxypropan-2-yl)pyridin-3-yl)methoxy-2-methoxyisonicotinaldehyde
184		2-(2-methoxyethoxy)-5-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde

185		2-(2-methoxyethoxy)-5-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)nicotinaldehyde
186		3-hydroxy-5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
187		3-(benzyloxy)-5-hydroxyisonicotinaldehyde
188		3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-5-methoxyisonicotinaldehyde
189		5-((2-(2-isopropyl-2H-1,2,4-triazol-3-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde



190		5-((2-(1-isopropyl-4-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
191		5-((2-(1-(2-hydroxyethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
192		2,2,2-trifluoroacetic acid : 6-(((4-formylpyridin-3-yl)oxy)methyl)picolinic acid (1:1)
193		2-methoxy-5-((2-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
194		5-((2-(4-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-oxo-1,2-dihydropyridine-4-carbaldehyde

195		5-((2-(1-cyclobutyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
196		5-((2-(1-cyclohexyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
197		5-((2-(1-(cyclohexylmethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
198		5-((2-(1-cyclopentyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde
199		2-(5-(3-(4-formyl-6-methoxypyridin-3-yl)oxy)methyl)pyridin-2-yl)-1H-pyrazol-1-yl)acetic acid

200		methyl 3-(5-(3-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)pyridin-2-yl)-1H-pyrazol-1-yl)propanoate
201		3-(3-(3-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)pyridin-2-yl)-1H-pyrazol-1-yl)propanoic acid
202		3-(5-(3-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)pyridin-2-yl)-1H-pyrazol-1-yl)propanoic acid
203		3-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)benzoic acid
204		6-(((4-formylpyridin-3-yl)oxy)methyl)nicotinonitrile 2,2,2-trifluoroacetate

205		6-(((4-formylpyridin-3-yl)oxy)methyl)nicotinic acid hydrochloride
206		2,2,2-trifluoroacetic acid : 6-(((4-formylpyridin-3-yl)oxy)methyl)-N-(methylsulfonyl)nicotinamide (2:1)
207		2-(2-methoxyethoxy)-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
208		2-methoxy-5-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
209		2-(2-methoxyethoxy)-5-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde

210		2-methyl-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
211		2-methyl-5-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
212		3-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
213		3-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
214		3-chloro-5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
215		3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-5-methylisonicotinaldehyde

216		3-chloro-5-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde
217		3-methyl-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde

The compound is selected from 5-hydroxy-2-(2-methoxyethoxy)isonicotinaldehyde (Compound 218), 5-hydroxy-2-(2-methoxyethoxy)nicotinaldehyde (Compound 219), 5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-oxo-1,2-dihydropyridine-4-carbaldehyde (Compound 220), 5-((2-(4-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-oxo-1,2-dihydropyridine-4-carbaldehyde (Compound 221), or a tautomer or pharmaceutically acceptable salt thereof.

5-(imidazo[1,2-a]pyridin-8-ylmethoxy)-2-methoxyisonicotinaldehyde,
 2-methoxy-5-((5-methylpyridin-3-yl)methoxy)isonicotinaldehyde,
 5-(isoquinolin-1-ylmethoxy)-2-methoxyisonicotinaldehyde,
 2-methoxy-5-(quinolin-2-ylmethoxy)isonicotinaldehyde,
 2-methoxy-5-(pyridin-4-ylmethoxy)isonicotinaldehyde,
 3-(imidazo[1,2-a]pyridin-8-ylmethoxy)-6-methylpicolinaldehyde,
 methyl 2-((4-formyl-6-methoxypyridin-3-yloxy)methyl)imidazo[1,2-a]pyridine-8-carboxylate,
 2-methoxy-5-((3-methyl-[1,2,4]triazolo[4,3-a]pyridin-8-yl)methoxy)isonicotinaldehyde,
 5-((2-bromopyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
 5-((2-(1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
 5-((5-bromopyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
 2-methoxy-5-((5-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
 5-((4-formyl-6-methoxypyridin-3-yloxy)methyl)nicotinic acid,
 2-methoxy-5-(quinolin-3-ylmethoxy)isonicotinaldehyde,

2-methoxy-5-((1-methyl-1H-indazol-4-yl)methoxy)isonicotinaldehyde,
tert-butyl 4-((2-formyl-6-methylpyridin-3-yloxy)methyl)-1H-indazole-1-carboxylate,
6-methyl-3-((1-methyl-1H-indazol-6-yl)methoxy)picolinaldehyde,
6-methyl-3-((1-methyl-1H-indazol-7-yl)methoxy)picolinaldehyde,
3-(isoquinolin-1-ylmethoxy)-6-methylpicolinaldehyde,
5-(benzo[d]oxazol-4-ylmethoxy)-2-methoxyisonicotinaldehyde,
3-((1,5-naphthyridin-4-yl)methoxy)-6-methylpicolinaldehyde,
6-methyl-3-((1-methyl-1H-indazol-5-yl)methoxy)picolinaldehyde,
6-methyl-3-(quinolin-5-ylmethoxy)picolinaldehyde,
2-methoxy-5-(quinolin-5-ylmethoxy)isonicotinaldehyde,
2-methoxy-5-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
2-methoxy-5-((2-(1-methyl-1H-pyrazol-3-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
5-((2-(2H-tetrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
2-methoxy-5-((2-(4-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
5-((3-(1H-pyrazol-5-yl)isoquinolin-4-yl)methoxy)-2-methoxyisonicotinaldehyde,
5-((2-(1H-pyrazol-1-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
5-(imidazo[1,5-a]pyridin-8-ylmethoxy)-2-methoxyisonicotinaldehyde,
5-((2-(1,5-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
5-((2-(1-ethyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
2-(difluoromethoxy)-5-(imidazo[1,2-a]pyridin-8-ylmethoxy)isonicotinaldehyde,
2-methoxy-5-((2-phenylpyridin-3-yl)methoxy)isonicotinaldehyde,
5-((3-(1-isopropyl-1H-pyrazol-5-yl)pyridin-4-yl)methoxy)-2-methoxyisonicotinaldehyde,
5-([2,3'-bipyridin]-3-ylmethoxy)-2-methoxyisonicotinaldehyde,
2-methoxy-5-((2-(o-tolyl)pyridin-3-yl)methoxy)isonicotinaldehyde,
2-methoxy-5-((2'-methoxy-[2,3'-bipyridin]-3-yl)methoxy)isonicotinaldehyde,
4-(((2-formyl-6-methylpyridin-3-yl)oxy)methyl)benzoic acid,
4-(((2-formylpyridin-3-yl)oxy)methyl)benzoic acid,
methyl 3-(((4-formylpyridin-3-yl)oxy)methyl)benzoate,
methyl 3-(((2-formyl-6-methylpyridin-3-yl)oxy)methyl)benzoate,
3-(((4-formylpyridin-3-yl)oxy)methyl)benzoic acid,
3-(((2-formyl-6-methylpyridin-3-yl)oxy)methyl)benzoic acid,

3-(((2-formylpyridin-3-yl)oxy)methyl)benzoic acid,
2-methoxy-5-((2-(1-(2-methoxyethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
2-methoxy-5-((2-(1-propyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
2-methoxy-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
5-((2-(1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)picolinaldehyde,
3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-6-methylpicolinaldehyde,
2-(difluoromethoxy)-5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
5-(imidazo[1,2-a]pyridin-8-ylmethoxy)-2-(2-methoxyethoxy)isonicotinaldehyde,
5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-(2-methoxyethoxy)isonicotinaldehyde,
5-((3-(1-isopropyl-1H-pyrazol-5-yl)pyrazin-2-yl)methoxy)-2-methoxyisonicotinaldehyde,
3-((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)picolinate,
5-((2-(2-hydroxypropan-2-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
2-(2-methoxyethoxy)-5-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
2-(2-methoxyethoxy)-5-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)nicotinaldehyde,
3-hydroxy-5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
3-(benzyloxy)-5-hydroxyisonicotinaldehyde,
3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-5-methoxyisonicotinaldehyde,
5-((2-(2-isopropyl-2H-1,2,4-triazol-3-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
5-((2-(1-isopropyl-4-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
5-((2-(1-(2-hydroxyethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,
6-(((4-formylpyridin-3-yl)oxy)methyl)picolinic acid,
2,2,2-trifluoroacetic acid : 6-(((4-formylpyridin-3-yl)oxy)methyl)picolinic acid (1:1),
2-methoxy-5-((2-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,

5-((2-(4-methyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-oxo-1,2-dihydropyridine-4-carbaldehyde,

5-((2-(1-cyclobutyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,

5-((2-(1-cyclohexyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,

5-((2-(1-(cyclohexylmethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,

5-((2-(1-cyclopentyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-2-methoxyisonicotinaldehyde,

2-(5-(3-((4-formyl-6-methoxypyridin-3-yloxy)methyl)pyridin-2-yl)-1H-pyrazol-1-yl)acetic acid,

methyl 3-(5-(3-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)pyridin-2-yl)-1H-pyrazol-1-yl)propanoate,

3-(3-(3-((4-formyl-6-methoxypyridin-3-yloxy)methyl)pyridin-2-yl)-1H-pyrazol-1-yl)propanoic acid,

3-(5-(3-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)pyridin-2-yl)-1H-pyrazol-1-yl)propanoic acid,

3-(((4-formyl-6-methoxypyridin-3-yl)oxy)methyl)benzoic acid,

6-(((4-formylpyridin-3-yl)oxy)methyl)nicotinonitrile 2,2,2-trifluoroacetate,

6-(((4-formylpyridin-3-yl)oxy)methyl)nicotinic acid,

6-(((4-formylpyridin-3-yl)oxy)methyl)nicotinic acid hydrochloride,

6-(((4-formylpyridin-3-yl)oxy)methyl)-N-(methylsulfonyl)nicotinamide,

2,2,2-trifluoroacetic acid : 6-(((4-formylpyridin-3-yl)oxy)methyl)-N-(methylsulfonyl)nicotinamide (2:1), 2-(2-methoxyethoxy)-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,

2-methoxy-5-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,

2-(2-methoxyethoxy)-5-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,

2-methyl-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,

2-methyl-5-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,

3-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,


3-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,

3-chloro-5-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)isonicotinaldehyde,
3-((2-(1-isopropyl-1H-pyrazol-5-yl)pyridin-3-yl)methoxy)-5-methylisonicotinaldehyde,
3-chloro-5-((2-(1-(3,3,3-trifluoropropyl)-1H-pyrazol-5-yl)pyridin-3-
yl)methoxy)isonicotinaldehyde, and
3-methyl-5-((2-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)pyridin-3-
yl)methoxy)isonicotinaldehyde,
or a tautomer or pharmaceutically acceptable salt thereof.

CLAIMS

What is claimed is:

1. A compound of formula (I):



(I)

or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein L¹⁰ is optionally substituted methylene or, preferably, a bond;


ring A is C₆-C₁₀ aryl, a C₃-C₈ cycloalkyl, a 5-10 membered heteroaryl or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, heteroaryl, cycloalkyl, or heterocycle is optionally substituted with 1-4: halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, and/or C₃-C₁₀ cycloalkyl, wherein the C₁-C₆ alkyl is optionally substituted with 1-5 halo, C₁-C₆ alkoxy, and/or C₃-C₁₀ cycloalkyl; or

ring A is C₆-C₁₀ aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, or heteroaryl is optionally substituted with 1-4 C₁-C₆ alkyl and/or C₁-C₆ alkoxy groups;

ring B is a 5-10 membered heteroaryl or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the

heteroaryl and the heterocycle is optionally substituted with 1-4: halo, C_1 - C_6 alkyl and/or $-CO-C_1-C_6$ alkyl, or

ring B is:



wherein ring B' including the $-N-CO-$ moiety is a 5-6 membered heterocycle containing up to 3 heteroatoms selected from nitrogen, oxygen, and sulfur and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted with 1-4 C_1 - C_6 alkyl groups;

 is a single or a double bond;

each X and Y is independently $(CR^{20}R^{21})_e$, O, S, SO, SO_2 , or NR^{20} ; e is 1 to 4, preferably 1; each R^{20} and R^{21} independently is hydrogen or C_1 - C_3 alkyl optionally substituted with 1-3 halo, OH, or C_1 - C_6 alkoxy, or $CR^{20}R^{21}$ is $C=O$, provided that if one of X and Y is O, S, SO, SO_2 , then the other is not CO, and X and Y are both not heteroatoms or oxidized forms thereof;

ring C is C_6 - C_{10} aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, $-OR^1$, C_1 - C_6 alkyl, $-COOR^1$, and/or C_1 - C_6 alkoxy, wherein the C_1 - C_6 alkyl is optionally substituted with 1-5 halo, C_1 - C_6 alkoxy and/or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S; or

ring C is C_6 - C_{10} aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, $-OR^1$, C_1 - C_6 alkyl, $-COOR^1$, NR^5R^6 ,


R^1 is a hydrogen, C_1 - C_6 alkyl or a prodrug moiety; wherein the alkyl is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring

heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, which is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein the heteroaryl is optionally substituted with C₁-C₆ alkyl;

R⁵ and R⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl or -COOR³;

R³ is hydrogen or optionally substituted C₁-C₆ alkyl;

V¹ and V² independently are C₁-C₆ alkoxy; or V¹ and V² together with the carbon atom they are attached to form a ring of formula:



wherein each V³ and V⁴ are independently O, S, or NH, provided that when one of V³ and V⁴ is S, the other is NH, and provided that V³ and V⁴ are both not NH; q is 1 or 2; each V⁵ is independently C₁-C₆ alkyl optionally substituted with 1-3 OH groups, or V⁵ is CO₂R⁶⁰, where each R⁶⁰ independently is C₁-C₆ alkyl or hydrogen; t is 0, 1, 2, or 4; or CV¹V² is C=V, wherein V is O, NOR⁸⁰, or NNR⁸¹R⁸²;

R⁸⁰ is optionally substituted C₁-C₆ alkyl;

R⁸¹ and R⁸² independently are selected from the group consisting of hydrogen, optionally substituted C₁-C₆ alkyl, COR⁸³, or CO₂R⁸⁴;

R⁸³ is hydrogen or optionally substituted C₁-C₆ alkyl; and

R⁸⁴ is optionally substituted C₁-C₆ alkyl;

with the proviso that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;


then ring A excludes optionally substituted 5-10 membered heteroaryl;

and provided that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 5-10 membered heteroaryl;


then ring A is not optionally substituted 4-10 membered heterocycle.

2. A compound of formula (X-I):




(X-I)

or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein ring A is phenyl optionally substituted with 1-3 halo and/or C₁-C₆ alkoxy, or is a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, optionally substituted, or is



wherein R⁷ is C₁-C₆ alkyl, optionally substituted with 3-5 fluoro groups, or is C₃-C₆ cycloalkyl;

ring B is selected from the group consisting of



wherein R⁸ is C₁-C₆ alkyl, -CO-C₁-C₆ alkyl or a prodrug moiety and wherein the pyridyl ring is optionally substituted with a halo or an NR²⁵(CH₂)₂N(R²⁵)₂ group where each R²⁵ is independently hydrogen or C₁-C₆ alkyl;


X is O, S, SO, or SO₂;

 is a single or a double bond;

ring C is phenyl or a 6 membered nitrogen-containing heteroaryl, each of which is optionally substituted with 1-4: halo, oxo, $-OR^1$, C_1-C_6 alkyl, $-COOR^1$, and/or C_1-C_6 alkoxy, wherein the C_1-C_6 alkyl is optionally substituted with 1-5 halo, C_1-C_6 alkoxy and/or 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S; and

each R^1 is hydrogen or a prodrug moiety R;

V^1 and V^2 independently are C_1-C_6 alkoxy; or V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:



wherein each V^3 and V^4 are independently O, S, or NH, provided that when one of V^3 and V^4 is S, the other is NH, and provided that V^3 and V^4 are both not NH; q is 1 or 2; each V^5 is independently C_1-C_6 alkyl or CO_2R^{60} , where each R^{60} independently is C_1-C_6 alkyl or hydrogen; t is 1, 2, or 4; or CV^1V^2 is $C=V$, wherein V is O, NOR^{80} , or $NNR^{81}R^{82}$;

wherein R^{80} is optionally substituted C_1-C_6 alkyl;

R^{81} and R^{82} independently are selected from the group consisting of hydrogen, optionally substituted C_1-C_6 alkyl, COR^{83} , or CO_2R^{84} ;

R^{83} is hydrogen or optionally substituted C_1-C_6 alkyl;

R^{84} is optionally substituted C_1-C_6 alkyl;

provided that when ring C is C_6-C_{10} aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;


then ring A excludes optionally substituted 5-10 membered heteroaryl;

and provided that when ring C is C_6-C_{10} aryl;

and ring B is optionally substituted 5-10 membered heteroaryl;


then ring A is not optionally substituted 4-10 membered heterocycle.

3. The compound of claim 2, wherein V^1 and V^2 independently are C_1 - C_6 alkoxy; or V^1 and V^2 together with the carbon atom they are attached to form a ring of formula:




wherein each V^3 and V^4 are independently O, S, or NH, provided that when one or V^3 and V^4 is S the other is NH, and provided that V^3 and V^4 are both not NH; q is 1 or 2; each V^5 is independently C_1 - C_6 alkyl or CO_2R^{60} , where each R^{60} independently is C_1 - C_6 alkyl or hydrogen; t is 1, 2, or 4; or CV^1V^2 is $C=V$, wherein V is O, and wherein the remaining variables are defined as in claim.

4. The compound of claim 3, of formula:




wherein the remaining variables are defined as in claim 2.


5. A compound of claim 3 or 4 selected from formulas (IIA), (IIB) and (IIC):



IIA



IIB



IIC

wherein

R^9 is hydrogen, $-OR^1$, C_1-C_6 alkoxy optionally substituted with 1-3 C_1-C_6 alkoxy or 4-10 membered heterocycle containing up to 5 ring heteroatoms selected from N, O, S or oxidized forms thereof;

R^{10} is hydrogen, halo, hydroxy, or C_1-C_6 alkoxy;


R^{11} is hydrogen or C_1-C_6 alkyl; and

R^{12} is $-OR^1$;









wherein R^1 is hydrogen or the prodrug moiety R.

6. The compound of claim 4, wherein ring A is phenyl substituted with 1-3 halo or C_1-C_6 alkoxy, or C_3-C_8 heterocyclyl containing 1-3 heteroatoms, wherein the heterocycle is optionally substituted with 1-3 halo.

7. The compound of any one of claims 3-6, wherein




is selected from the group consisting of:




and

8. The compound of any one of claims 3-6, wherein




is



9. A compound of claim 1 wherein:

ring A is C₆-C₁₀ aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, or heteroaryl is optionally substituted with 1-4: C₁-C₆ alkyl and/or C₁-C₆ alkoxy;

ring B is:



wherein ring B' including the -N-CO- moiety is a 5-6 membered heterocycle containing up to 3 heteroatoms selected from nitrogen, oxygen, and sulfur and oxidized forms of N and S, wherein each of the heteroaryl and the heterocycle is optionally substituted with 1-4 C₁-C₆ alkyl groups;

each X and Y is independently CR²⁰R²¹, O, S, SO, SO₂, or NR²⁰; each R²⁰ and R²¹ independently is hydrogen or C₁-C₃ alkyl optionally substituted with 1-3 halo, OH, or C₁-C₆ alkoxy, or CR²⁰R²¹ is C=O, provided that if one of X and Y is O, S, SO, SO₂, then the other is not CO, and X and Y are both not heteroatoms or oxidized forms thereof;

ring C is C₆-C₁₀ aryl or a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, each of which is optionally substituted with 1-4: halo, oxo, -OR¹, C₁-C₆ alkyl, -COOR⁵, NR⁵R⁶,


R¹ is a hydrogen, C₁-C₆ alkyl or a prodrug moiety; wherein the alkyl is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring

heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, which is optionally substituted with a 5-10 membered heteroaryl containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein the heteroaryl is optionally substituted with C₁-C₆ alkyl;

R⁵ and R⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl or -COOR³;

R³ is hydrogen or optionally substituted C₁-C₆ alkyl;

V¹ and V² independently are C₁-C₆ alkoxy; or V¹ and V² together with the carbon atom they are attached to form a ring of formula:



wherein each V³ and V⁴ are independently O, S, or NH, provided that when one of V³ and V⁴ is S, the other is NH, and provided that V³ and V⁴ are both not NH; q is 1 or 2; each V⁵ is independently C₁-C₆ alkyl optionally substituted with 1-3 OH groups, or V⁵ is CO₂R⁶⁰, where each R⁶⁰ independently is C₁-C₆ alkyl or hydrogen; t is 0, 1, 2, or 4; or CV¹V² is C=V, wherein V is O, NOR⁸⁰, or NNR⁸¹R⁸²;

R⁸⁰ is optionally substituted C₁-C₆ alkyl;

R⁸¹ and R⁸² independently are selected from the group consisting of hydrogen, optionally substituted C₁-C₆ alkyl, COR⁸³, or CO₂R⁸⁴;

R⁸³ is hydrogen or optionally substituted C₁-C₆ alkyl; and

R⁸⁴ is optionally substituted C₁-C₆ alkyl;

with the proviso that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 4-10 membered heterocyclyl;

then ring A excludes optionally substituted 5-10 membered heteroaryl;


and provided that when ring C is C₆-C₁₀ aryl;

and ring B is optionally substituted 5-10 membered heteroaryl;

then ring A is not optionally substituted 4-10 membered heterocycle.


10. The compound of claim 9, wherein CV^1V^2 is $C=V$, wherein V is O, and wherein the remaining variables are defined as in claim 10.

11. The compound of claim 10, of formula:



or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein the remaining variables are defined as in claim 9.

12. A compound of claim 10 or 11 of formula (VI) or (VII):



or a tautomer thereof, or a pharmaceutically acceptable salt of each thereof, wherein ring A is C_6 - C_{10} aryl, or a 5-10 membered heteroaryl, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S,

wherein each of the aryl, or heteroaryl is optionally substituted with 1-4 C₁-C₆ alkyl;

ring B is C₆-C₁₀ aryl, C₃-C₈ cycloalkyl, a 5-10 membered heteroaryl containing up to 5 ring heteroatoms or a 4-10 membered heterocycle containing up to 5 ring heteroatoms, wherein the heteroatom is selected from the group consisting of O, N, S, and oxidized forms of N and S, wherein each of the aryl, heteroaryl, cycloalkyl or heterocycle is optionally substituted with 1-4: halo, C₁-C₆ alkyl, or C₁-C₆ alkoxy, wherein the C₁-C₆ alkyl is optionally substituted with 1-5 halo, C₁-C₆ alkoxy, and/or C₃-C₁₀ cycloalkyl;


R⁴ is halo, oxo, -OR¹⁸, C₁-C₆ alkyl, C₁-C₆ alkoxy, -COOR⁵, and/or NR⁵R⁶;

R¹⁸ is hydrogen, substituted C₁-C₆ alkyl, or a prodrug moiety R;


R⁵ and R⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl or -COOR³; and
















R³ is hydrogen, provided that the COOR³ is not joined to a nitrogen atom, or optionally substituted C₁-C₆ alkyl.











13. The compound of claim 12, wherein ring B is selected from the group consisting of:





14. A compound of claim 1 selected from the group consisting of:
















or a prodrug thereof, or a pharmaceutically acceptable salt of each thereof.

15. A compound of claim 1 selected from the group consisting of:





or N oxides thereof, or a pharmaceutically acceptable salt of each thereof.

16. A composition comprising a compound of any one of claims 3-8 and 10-15, and at least one pharmaceutically acceptable excipient.

17. A method for increasing oxygen affinity of hemoglobin S in a subject, the method comprising administering to a subject in need thereof a therapeutically effective amount of a compound of any one of claims 3-8 and 10-15 or the composition of claim 16.

18. A method for treating oxygen deficiency associated with sickle cell anemia, the method comprising administering to a subject in need thereof a therapeutically effective amount of a compound of any one of claims 3-8 and 10-15 or the composition of claim 16.