



MONOCLONAL ANTIIDIOTYPIC ANTIBODIES (AB2) AND THEIR USES.

Field of the Invention.

The present invention is related to the field of Immunology and in particular to the selection and use of antidiotypic monoclonal antibodies (AB2) IgG type with the main characteristic of being highly connected to the idiotypic network and recognition of B and T human lymphocytes, which are major participants in the immune response.

Accordingly the objective of this invention is to provide antidiotypic monoclonal antibodies IgG type connected to the immune network, able to interact with T and B lymphocytes and able to exert an immunoregulatory effect.

Description of the Prior Art.

The immune response is mainly based on the antigen stimulation of some determined B lymphocyte clones bearing in the surface specific receptors for this antigen. T cells have the ability of producing positive and negative signals as a way of regulation of this process. However the immune system regulation mechanisms are very complex as was postulated by Jerne (Jerne NK (1974) Ann Immunol. Inst. Pasteur 125, C:373:389) : the immune system can be "seen" as an organized structure of antibody molecules and interconnected lymphocytes by idiotypic directed interactions.

Idiotypes in antibodies are defined as epitopes in the variable regions in such a way each antibody presents particular idiotypical determinants and among them the antigen- antibody recognition region is included. The theory of the idiotypic network emerges from the demonstrated fact (Urbain .J (1976) Ann Immunol. 1237 C; 357) that the

idiotypic determinants in the immunoglobulin molecule (AB1) are recognized by antidiotypic antibodies (AB2) obtained against it.

The binding of an idiotypic determinant with its corresponding antidiotypic occurs in a dynamic state that could be perturbed during the course of immune response.

The theory of the idiotypic network proposed by Jerne in the seventies has implicit that the immune repertoire contains and recognize all the possible antigens. Ignoring the dicotomy between self and non self antigens, thus the existence of autoreactive clones of lymphocytes occurs in a physiological mode.

Studies performed in the last ten years have demonstrated the presence of natural autoreactive antibodies in normal subjects (Avrameas.S., (1991) Immunol.. Today, 132C: 231: 236, Antin et al (1986), J Immunol. 136,505:510, Lundkvist (1989), PNAS USA 86:5074-5078, Longtenberg .T (1989)J Exp Med, 170:1347.1335)).

These experimental facts have prompted the concept that considers the immune system as a highly connected network of specific receptors ontogenically organized through the internal references (self recognition) philogenetically selected for the maintenance of the homeostatic physiological functions, and more than the normal defense against infections In this sense recognition of the self is normal and tolerance to the self and memory are not any more considered clone properties as postulates by Burnet in his clone selection theory (Burnet F.M. (1956) London Cambridge University Press) but they are emerging properties of the immune network with an integral point of view (Coutinho. A

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED	2
---------------------------------	---

(1984) Immunol. Rev. 79:151-168, Coutinho. A (1989) Immunol. Rev 110:63-87).

This new concept in relation to the immune system have had its impact in understanding the basic mechanisms of autoimmune diseases and also in the new therapy approach Zanetti. M (1985) Immunol. Today 6: 299-302; Kaveri et al (1991) Clin Exp. Immunol. 86:192-198; Kazatchine. D.M (1994) Immunol. Rev 139:79-107.

An amazing therapeutic effect with different immunoglobulin compounds have observed in the wide variety of systemic and organ specific autoimmune diseases that is useful for the clinical validation of this new concept. (Kazatchine et al (1993) Autoimmunity: physiology and disease Wiley-Liss, Inc., New York).

IgG Immunoglobulin preparations are obtained from blood pools of healthy donors representing the autoreactive physiological network of antibodies. The reactivity analysis in these preparations have demonstrated the existence of antibodies interacting with idiotypes expressed in the autoantibodies and autoantibodies per se (Rossi, et al (1988) Clin Exp. Immunol. 74: 311-316; Rossi et al (1989) J Immunol. 143:4104-4109. Both types of reactivity are in high concentration in the total number of IgG molecules that are more connected than other molecules inside the pool (Dietrich et al (1992) Eur J Immunol. 22:1701-1706). In this way the immunoregulatory effect of these preparations justify the design of new therapeutic strategies addressed to restores the connectivity and diversity in the autoimmune patient. Through this clinical value the concept of emergency of autoimmune disease has been validated, through the fail in

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

3

the regulatory network in the variable regions of the connected antibodies.

In previous research a panel of monoclonal antibodies against tumor associated gangliosides was obtained, the antigen, classified as T cell independent self antigen (EPO: No:0 657 471 A1) These Mabs are IgM type bearing idiotypes able to activate the whole idiotypic network and thus generating an antibody response in absence of the antigen, i.e. they are recurrent idiotypes. These antibodies have been previously described as recognizing specifically the NeucAc GM2 ganglioside (Alfonso. M et al (1995) Hibridoma 14(3): 209-216).

The present invention provides novelty in the generation and selection of a panel of antidiotypic monoclonal antibodies (AB2) against the original IgM antibody recognizing ganglioside.

From this panel a further characterization was performed to one of the antibodies, named B7 showing it was an IgG2a with medium affinity for the antigen (Kd 10 E-8) and alpha type even though this type, it was able to induce an antibody (AB1') independent response in a singeneic model (Balb/C).

Surprising properties have been shown in the antidiotypic antibody B7 studies: High idiotypic connectivity with immunoglobulins of different species human B and T lymphocyte recognition and the immunoregulatory effect on previously activated B and T lymphocytes.

Up to now there isn't any therapy proposing this property in antiidiotype antibodies for the treatment of autoimmune, and infectious diseases and cancer.

The novelty of the present invention consists in the selection of the antidiotypic monoclonal antibodies IgG type

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

4

highly connected to the immune network through their variable regions able to interact with human B and T lymphocytes and their immunoregulatory effect.

Detailed Description of the Invention.

GENERATION AND PRODUCTION OF MONOCLONAL ANTIDIOTYPIC ANTIBODIES HIGHLY CONNECTED TO THE IMMUNE NETWORK.

For the generation and production of monoclonal antidiotypic antibodies, the Mab against ganglioside is used as immunogen with an immunization procedure is performed as described elsewhere (European Patent Application 657 471 A1).

Mice with high titers of antidiotypic antibodies in serum are reimmunized and three days afterwards the spleen is removed and splenocytes are obtained. Antibody producing cells can be obtained from any other organ if preferred.

These cells are fused with myeloma cells, cell fusion may be performed according to any of the known methods and thus hybridomas are obtained. This method is described in detail in any of the procedures for this purpose.

The antibodies produced are assayed through any immunoassay method preferably an immunoenzymatic method, in which the supernatant are assessed for the presence of specific recognition of the proper antigen. The antibody- antibody binding is detected through a second antibody labeled with an enzyme.

The supernatants are selected for their capacity of blocking the antiganglioside Mab (E1) binding to the antigen (Neu Ac GM2) the supernatants in adequate concentrations are incubated and then the E1 Mab with the antigen.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

5

Once the desired hybridoma has been selected and cloned at least twice by limiting dilution, the resulting Mab can be produced by several culture techniques.

One method can be inoculating the proper amount of cells in the peritoneal cavity of mice producing ascitic fluid in which the Mab is produced in high concentration.

The antidiotypic Mabs obtained recognize different Mabs anti gangliosides related or not related and also are characterized by not blocking the binding of the anti ganglioside Mab to the corresponding antigen.

PROCEDURE FOR THE EVALUATION OF THE IDIOTYPIC CONNECTIVITY IN THE ANTIDIOTYPIC ANTIBODIES (IGG TYPE).

The evaluation of the idiotypic connectivity of antidiotypic monoclonal antibodies of IgG type obtained against the monoclonal antibody antiganglioside E1 recognizing the Neuc Ac GM2 ganglioside is performed using an indirect Immunoenzimatic assay in PVC plates. Plates are coated with specific antigangliosides Mabs and non specific Mabs of murine origin or any other species also human origin Mabs are used for this purpose and IgG pool. The wells are blocked during 1 hour approximately at room temperature, with PBS and Bovine serum albumin (1%) Followed by the addition of the antidiotypic monoclonal antibodies in a concentration range of 250 ug to 10 ug/mL during two hours in a humid chamber at 37°C . The plate is washed at least 4 times (PBS) and the anti mouse Fc labeled with alkaline Phosphatases conjugate is added during 4 hours. The substrate is added after that time and the plate is read at OD 405nm.

Negative controls are included, i.e. other antidiotypic monoclonal antibodies not connected to the idiotypic network.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

press page

highly connected to the immune network through their variable regions able to interact with human B and T lymphocytes and their immunoregulatory effect.

Detailed Description of the Invention.

GENERATION AND PRODUCTION OF MONOCLONAL ANTIDIOTYPIC ANTIBODIES HIGHLY CONNECTED TO THE IMMUNE NETWORK.

For the generation and production of monoclonal antidiotypic antibodies, the Mab against ganglioside is used as immunogen with an immunization procedure is performed as described elsewhere (European Patent Application 657 471 A1).

Mice with high titers of antidiotypic antibodies in serum are reimmunized and three days afterwards the spleen is removed and splenocytes are obtained. Antibody producing cells can be obtained from any other organ if preferred.

These cells are fused with myeloma cells, cell fusion may be performed according to any of the known methods and thus hybridomas are obtained. This method is described in detail in any of the procedures for this purpose.

The antibodies produced are assayed through any immunoassay method preferably an immunoenzymatic method, in which the supernatant are assessed for the presence of specific recognition of the proper antigen. The antibody- antibody binding is detected through a second antibody labeled with an enzyme.

The supernatants are selected for their capacity of blocking the antiganglioside Mab (E1) binding to the antigen (Neu Ac GM2) the supernatants in adequate concentrations are incubated and then the E1 Mab with the antigen.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

7

2 x 10⁶ cells

Once the desired hybridoma has been selected and cloned at least twice by limiting dilution, the resulting Mab can be produced by several culture techniques.

One method can be inoculating the proper amount of cells in the peritoneal cavity of mice producing ascitic fluid in which the Mab is produced in high concentration.


The antidiotypic Mabs obtained recognize different Mabs anti gangliosides related or not related and also are characterized by not blocking the binding of the anti ganglioside Mab to the corresponding antigen.

PROCEDURE FOR THE EVALUATION OF THE IDIOTYPIC CONNECTIVITY IN THE ANTIDIOTYPIC ANTIBODIES (IGG TYPE).

The evaluation of the idiotypic connectivity of antidiotypic monoclonal antibodies of IgG type obtained against the monoclonal antibody antiganglioside E1 recognizing the Neuc Ac GM2 ganglioside is performed using an indirect Immunoenzimatic assay in PVC plates. Plates are coated with specific antigangliosides Mabs and non specific Mabs of murine origin or any other species also human origin Mabs are used for this purpose and IgG pool. The wells are blocked during 1 hour approximately at room temperature, with PBS and Bovine serum albumin (1%) Followed by the addition of the antidiotypic monoclonal antibodies in a concentration range of 250 ug to 10 ug/mL during two hours in a humid chamber at 37°C . The plate is washed at least 4 times (PBS) and the anti mouse Fc labeled with alkaline Phosphatases conjugate is added during 4 hours. The substrate is added after that time and the plate is read at OD 405nm.

Negative controls are included, i.e. other antidiotypic monoclonal antibodies not connected to the idiotypic network.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

The antidiotypic monoclonal antibodies considered connected recognize the different antibodies coming from different species and their corresponding fragments and also recognize the Human pool IgG obtained from human healthy donors, indicating the high connectivity of these Mabs recognizing the idiotypes in the variable regions of immunoglobulins of other species.

PROCEDURE FOR THE EVALUATION OF DIFFERENT SPECIES B AND T LYMPHOCYTE RECOGNITION BY THE ANTIDIOTYPIC MABS (AB2) IGG TYPE IDIOTYPICALLY CONNECTED.

The evaluation of the recognition of B and T lymphocytes from different species (mouse, rat monkey and human) by the antidiotypic Mabs(AB2) IgG type idiotypically connected is performed by an Indirect Immunofluorescence assay in a Flow cytometer.


B and T lymphocytes from different species are isolated by the Ficoll gradient technique from peripheral blood and other lymphoid organs

(spleen, lymph nodes and peritoneum) according to the species studied. The cell ring is isolated and the cell counting is done with a Neubauer chamber.

The cell suspension (200 uL) is incubated in a humid chamber during 30 minutes together with the antidiotypic Mab (AB2). Positive controls are added (ior t3 or ior t4) in established conditions and a negative control is also included. A proper conjugate that could be anti mouse FITC or RPE is added and incubated during 30 min. and after that time the sample is measured in the Flow cytometer.

PROCEDURE FOR THE EVALUATION OF THE BIOLOGICAL EFFECT OF ANTIDIOTYPIC MABS (AB2) IGG TYPE IDIOTIPICALLY CONNECTED ON THE B AND T LYMPHOCYTES IN VITRO.

This Sheet Has Been Replaced



For the evaluation of the biological effect of antidiotypic Mabs(AB2) IgG type idiotipically connected on the B and T lymphocytes of any mammal species a cell proliferation assay is performed by 3H-Thimidine incorporation.

The Mononuclear peripheral cells are isolated through a Ficoll gradient technique. The cells in amount of 100000 per well are incubated with the proper medium with 10 % serum in microplates . Lymphocyte activators as PHA in a concentration that can be 5 ug/ mL, Con A in the same concentration and PWM in the same concentration are added to the incubation in the presence of different concentrations of the antidiotypic Mabs (300 ug to 15 ug/ mL).

Controls are included in the assay , activators only, or lymphocyte cells only. Cultures are kept during 72 hours in a 5 % CO2 humid atmosphere at 37 oC.

3 H-Thymidine is added (1 uCi, 23 uCi/ mmol) to each well 6 hours before the culture is arrested. Cells are harvested and 3 H-Thymidine is measured by a beta counter (LKB Sweden).

The highly connected antidiotypic mabs are able to suppress or enhance the stimulation of previously activated B and T lymphocytes of any species. The antidiotypic Mabs not connected are not able to exert this effect.

PROCEDURE FOR THE EVALUATION OF THE BIOLOGICAL EFFECT OF ANTIDIOTYPIC MABS (AB2) IGG TYPE IDIOTIPICALLY CONNECTED ON THE B AND T LYMPHOCYTES IN VIVO ON DIFFERENT ANIMAL MODELS FOR AUTOIMMUNE DISEASES, INFECTIOUS DISEASES AND MALIGNANT NEOPLASM'S.

The in vivo biological effect of Antidiotypic Mabs (AB2) IgG type idiotipically connected on the B and T lymphocytes for autoimmune diseases, infectious diseases and malignant

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

neoplasm's, was measured using different animals experimental models. The experimental model MRL/lpr/lpr mouse was used to evaluate B7 Mab effects in the autoimmune diseases. This animal develops the Disseminated Lupus Erythematosus (DSL).

MRL/lpr/lpr mice are intraperitoneally injected with high doses of the B7 antidiotypic Mab according to the following scheme: 1 mg/ mouse, three times a week, initiating the treatment 24 hours after the birth of the animal and maintaining it during 4 weeks.

Also a low dose regime is evaluated with 100 ug of the antidiotypic Mab at the time of the animal birth.

The therapeutic effect is measured through several aspects: the delayed in time of the clinical signs appearance (skin lesions lymph nodes); through the increase of animal survival and also through the variation of any biological parameter considered as indicator (Proteinuria, anti DNA autoantibodies). A control group of non treated animals was used to establish the comparison.

The evaluation of antitumor effect was performed using an experimental animal model. Mice (C57BL/C) with xenografted lung tumor cells (RL-67) were treated with different doses of the antidiotypic Mab according to the following scheme: 1 mg/ mouse, three times a week.

The therapeutic effect is measured through the comparison of survival curves of treated animals versus the control.

The biological effect of the antidiotypic Mabs in infectious diseases was assessed using Balb/C mice previously infected with a clone of Plasmodium Chabaudi.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

(Conquy-Adib. M et al (1993) Mol Immunology 30 (2):119-127). After three days of the inoculation the mice were treated with the following therapeutic scheme:

1) A total dose of 800 ug injected in successive doses of 300, 100, 200, and 200 ug intraperitoneally during one week.

2) A high total doses of 3 mg was administered in 4 weekly injections (two of 1 mg and followed by two of 500 ug) a control group received 0.2 mL of saline solution in the same scheme.

Capacity for delaying the clinical signs appearance is evaluated comparing the control versus the treated group.

PROCEDURE FOR HUMANIZATION AND REDUCE IMMUNOGENICITY OF THE MURINE B7 MONOCLONAL ANTIBODY.

B7 Mab was modified in order to decrease its immunogenicity with a procedure (according European patent bulletin # 1996/10 under publication no 0699755) which simultaneously reduces immunogenicity of the rodent monoclonal antibody while preserving its binding properties in their entirety. Since the antigenicity of an immunoglobulin is dependent on the presence of T-cell antigenic peptides onto their sequence, the immunogenicity of a xenogenic or allogenic antibody could be reduced by replacing the residues included onto the T-cell antigenic sequences which differ from those usually found in another mammalian species antibodies. Of course, the replacement of residues do not include those involved in to the canonical structures or in the Vernier zone. This judicious replacement of residues have no effect on the structural determinants or on the interdomain contacts, thus, ligand binding properties should be unaffected as a

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

consequence of alterations which are limited to the variable region framework residues.

(1)- Analysis of homology of variable regions.

The present procedure makes use of the available sequence data for human antibody variable domains compiled by Kabat et al., "Sequences of proteins of Immunological Interest" Fifth edition., Bethesda, Maryland; National Inst. of Health, 1994.

In the first step the variable domains of the murine B7 heavy or light chain are compared with those corresponding variable domains of the human sequences.

The comparison is made by an automated-computerized method (PC-DOS HIBIO PROSIS 06-00, Hitachi.). The most homologous human variable regions are then compared residue for residue to the corresponding murine regions. This will also define the human subgroup to which each mouse sequence most closely resembles.

(2)- Prediction of T-epitopes.

In the second step, the two homologous variable region sequences, mouse and human are analyzed for prediction of T-antigenic sequences.

The algorithm AMPHI (Bersofsky et al., The Journal of Immunology 138: 2213-2229, (1987)) predicts α Helical sequences. The algorithm SOHHA predicts the strip of helix hydrophobicity (Elliott et al., J. Immunol. 138: 2949-2952, (1987)). These algorithms predict T-cell presented fragments of antigenic proteins.

(3) Analysis for immunogenicity reduction.

Those residues in the mouse framework which differ from its human counterpart are replaced by the residues present

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

in the human counterpart. This switching occurs only with those residues which are at the T-antigenic sequences.

Finally, replacement of those residues responsible for the canonical structures or those involved in the Vernier zone could have a significant effect on the tertiary structure. Hence, they can not be included in the replacement. Additional information about the influence of the proposed replacements on tertiary structure or the binding site, could be obtained from a molecular model of the variable regions.

(4) The method for constructing and expressing the altered antibody.

The following procedures are used to prepare recombinant DNA sequences which incorporate the CDRs of the murine mAb, both light and heavy chains, into human, frameworks that can be used to transfect mammalian cells for the expression of recombinant antibody less immunogenic and with the antigen specificity of the animal monoclonal antibody:

a.-) mutagenesis and assembly of variable region domains including CDRs and FRs regions. The PCR-mutagenesis method (Kamman et al., Nucleic Acids Res. 17: 5404-5409, (1989)) is preferably used to introduce the changes at different positions.


b.-) preparation of an expression vector including one variable region and the corresponding human constant region which upon transfection into cells results in the secretion of protein sufficient for affinity and specificity determinations.

c.-) co-transfection of heavy and light chain expression vectors in appropriate cell lines.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



After about 2 weeks, the cell supernatants are analyzed by ELISA for human IgG production. The samples are then analyzed by any method for human IgG capable for binding to specific antigens.

Example 1: Generation and production of monoclonal antidiotypic antibodies (AB2) highly connected to the immune network against the E1 Mab obtained in the syngeneic model.

The Mab against ganglioside (E1 Mab) is used as immunogen in Balb/C mice (6-8 weeks old) with an immunization procedure performed as described elsewhere (EP No:0 667 471 A1).

Mice with high titers of antidiotypic antibodies in serum are reimmunized and three days afterwards the spleen is removed and splenocytes are obtained.

These cells are fused with myeloma cells, fusion is performed according to the method described by Kohler and Milstein with some minor modifications (Nature (1975) 256:495-497) and hybridomas are obtained. Murine splenocytes are fused with the non secretor murine myeloma cells P3/X63Ag66.5.3 in a rate 10:1 in 0.5 mL of fusion media containing PEG

(3000-5000).

The antibodies produced are assayed through any immunoassay method, in which the supernatants are assessed for the presence of specific recognition of the proper antigen. The antibody-antibody binding is detected through a second antibody labeled with an enzyme.

The supernatants are selected for their capacity of blocking the antiganglioside Mab (E1) binding to the antigen (Neu Ac GM2) the supernatants in adequate concentrations are incubated and then the E1 Mab with the antigen.

The antidiotypic Mabs are selected if they recognized different Mabs anti gangliosides related or not related (Figure 1) and also they are characterized by not blocking the binding of the anti ganglioside Mab to the corresponding antigen. The hybrid 4B7 was cloned twice and the resultant Mab was the antidiotypic Mab B7 an IgG2a class.


Example 2: Procedure for the evaluation of the idiotypic connectivity in the antidiotypic antibodies (IgG type).

The evaluation of the idiotypic connectivity of antidiotypic monoclonal antibodies of IgG type obtained against the monoclonal antibody E1 that recognize the Neuc Ac GM2 ganglioside is performed using an indirect Immunoenzimatic assay in PVC plates (96 wells, High Binding Costar). Plates are coated with specific antigangliosides Mabs including the one used as immunogen the E1 Mab, and non specific Mabs in carbonate -bicarbonate buffer. The wells are blocked during 1 hour at room temperature, with PBS and Bovine serum albumin (1%) Followed by the addition of the antidiotypic monoclonal antibodies in a concentration range of 250 ug to 10 ug/mL during two hours in a humid chamber at 37°C . The plate is washed at least 4 times (PBS) and the anti mouse Fc labeled with alkaline Phosphatase conjugate is added during 4 hours. The substrate is added after that time and the plate is read at OD 405 nm.

Negative controls are included, i.e. other antidiotypic monoclonal antibodies not connected to the idiotypic network.

The supernatants of different hydrides producing antidiotypic Mabs recognized the variable regions of the antiganglioside related Mabs, recognized non related IgM of Balb /C origin Mabs, and also neonatal Mabs. (Figure 1).

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

16

The antidiotypic monoclonal antibody B7 IgG2a type showed a high idiotypic connectivity recognizing the idiotypes in the variable regions of the antiganglioside related Mabs, non related IgM of Balb/C origin Mabs, and also neonatal Mabs with an affinity of 10^{-8} M. (Figure 2).

Example 3: Characterization of human myeloma immunoglobulin (IgG) idiotypes recognition and in the therapeutic IgG preparation by the antidiotypic Mabs idiotypically connected.

The capacity of recognition of the human myeloma immunoglobulin (IgG) idiotypes (variable regions) by the B7 Mab was evaluated .

For this purpose an Indirect Immunoassay is used (96 wells, High Binding Costar). Plates are coated with different IgG fractions or their fragments

(obtained according Colligan. E.J et al, Eds (1995) Current Protocols in Immunology). The wells are coated with myeloma proteins in carbonate-bicarbonate buffer and blocked during 1 hour at room temperature, with PBS and Bovine serum albumin (1%) Followed by the addition of the antidiotypic monoclonal antibodies in a concentration range of 380 ug to 11 ug/mL during two hours in a humid chamber at 37°C . The plate is washed at least 4 times (PBS) and the anti goat Fc region labeled with alkaline Phosphates is added and incubated during 4 hours. The substrate is added after that time and the plate is read at OD 405 nm.


The B7 antidiotypic monoclonal antibody recognized the variable regions

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

2



(idiotypes) present in the human myeloma immunoglobulins (Figure 3). Other murine Mabs not idiotipically connected did not show any reactivity with this IgG fractions (Figure 4).

The supernatants of other hydrides producing high connected antidiotypic Mabs recognized the variable regions of the IgG human myeloma proteins. The recognition profile was different for each IgG human myeloma protein. (Figure 5).

The B7 antidiotypic monoclonal antibody also recognized the variable regions (idiotypes) present in the human myeloma immunoglobulins fragments (Figure 6).

Moreover the recognition specificity was evaluated through a competence assay for the recognition of the human myeloma IgG by the B7 Mab and different concentrations of the therapeutic human IgG the experiment was performed as followed:

Different concentrations of the therapeutic human IgG were incubated overnight with a fixed concentration of the B7 Mab (95 ug/mL) after that time this mixture was incubated with the human myeloma IgG using the immunoassay already described in this example.

Increasing concentrations of therapeutic human IgG inhibits the B7 Mab recognition for the human myeloma IgG idiotypes. (Figure 7).

The fact that this B7 Mab of murine origin is able to recognize the variable regions of a human myeloma IgG and the therapeutic human IgG show us the high idiotipic connectivity and suggest that the zones recognized in the variable regions are highly conserved among species.

Example 4: Evaluation of human B and T Lymphocyte recognition by the antidiotypic Mabs (AB2) IgG type idiotipically connected.

The evaluation of human B and T lymphocytes recognition by the antidiotypic Mabs(AB2) IgG type idiotipically connected is performed by an Indirect Immunofluorescence assay in a Flow cytometer.


B and T lymphocytes are isolated by the Ficoll gradient technique from peripheral blood. The cell ring is isolated and the cell counting is done with a Neubauer chamber. Concentration and cell viability were determined.

The cell suspension (200 uL) adjusted to 10^7 cells is incubated in a humid chamber during 30 min. together with the antidiotypic Mab (AB2) a standard curve is added in the concentration range of 250 to 10 ug/mL. Positive controls are added (ior t3 or ior t4) in established conditions and a negative control is also included (ior r3). After incubation the wells with the cell suspension are washed with PBS BSA Azide and centrifuging eliminating the supernatant and centrifuge again four times. An anti mouse FITC conjugate is added and incubated during 30 min. in humid chamber and after that the sample is washed and centrifuged again four times, finally the cell suspension is resuspended in 100 uL of PBS and the sample is measured in the Flow cytometer.

Figure (8A) shows the recognition analysis of human B and T lymphocytes by the B7 Mab in the Flow Cytometer. Figure (8B) shows the negative control (ior r3 Mab, a non idiotipically connected Mab).

The B7 Mab was able to recognize 53 % of human lymphocytes from a normal donor (Peripheral blood lymphocytes) Therefore this Mab of idiotypic reactivity is able to recognize molecular structures present in T and B human lymphocytes.

This Sheet Has Been Replaced



Example 5: Evaluation of the biological effect of B7 antidiotypic Mab(AB2) IgG type idiotipically connected on the B and T lymphocytes in vitro.

For the evaluation of the biological effect of B7 antidiotypic Mabs (AB2) IgG type idiotipically connected on the B and T lymphocytes of any mammal species a cell proliferation assay is performed by 3H-Thymidine incorporation.

The peripheral blood lymphocyte cells (PBL) are isolated through a Ficoll gradient technique. The cells in amount of 100000 per well are incubated RPMI 1640 medium with 10 % serum in microplates . Lymphocyte activators as PHA in a concentration that can be 5 ug/ mL, Con A in the same concentration and PWM in the same concentration are added to the incubation in the presence of different concentrations of the antidiotypic Mabs (300 ug to 15 ug/ mL). Controls are included in the assay, activators only, or lymphocyte cells only. Cultures are kept during 72 hours in a 5 % CO2 humid atmosphere at 37°C.

3 H-Thymidine is added (1 uCi, 23 uCi/ mmol) to each well 6 hours before the culture is arrested. Cells are harvested and 3 H-Thymidine is measured by a beta counter (LKB Sweden).

The B7 highly connected antidiotypic mab was able to suppress the stimulation of previously activated B and T lymphocytes (Figure 9 A and B). This suppresser effect of the mytogenic stimulation was also produced by therapeutic human IgG (Figure 10 A and B) The antidiotypic Mabs not connected are not able to exert this effect.

The fact that this B7 Mab exerts a biological suppresser effect in previously activated B and T human lymphocytes

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

20



demonstrates the immunoregulator capacity of this highly connected antidiotypic MAB.

Example 6: Evaluation of the biological effect of antidiotypic Mabs

(AB2) IgG type idiotypically connected on the B and T lymphocytes in vivo on different animal models for autoimmune diseases, infectious diseases and malignant neoplasm's.

The in vivo biological effect of antidiotypic Mabs (AB2) IgG type idiotypically connected on the B and T lymphocytes for autoimmune diseases, infectious diseases and malignant neoplasm's, was measured using different animals experimental models. The experimental model MRL/lpr/lpr mouse was used to evaluated B7 Mab effects in the autoimmune diseases. This animal develops the Disseminated Lupus Erythematosus (DSL).

MRL/lpr/lpr mice are intraperitoneally injected with high doses of the B7 antidiotypic Mab according to the following scheme: 1 mg/ mouse, three times a week, initiating the treatment 24 hours after the birth of the animal and maintaining it during 4 weeks.

Also a low dose regime is evaluated with 100 ug of the antidiotypic Mab at the time of the animal birth.

The therapeutic effect is measured through several aspects: the delayed in time of the clinical signs appearance (skin lesions lymph nodes); through the increase of animal survival and also through the variation of any biological parameter considered as indicator (Proteinuria, anti DNA autoantibodies). A control group of non treated animals was used to establish the comparison.

The evaluation of antitumor effect was performed using an experimental animal model. Mice (C57/BL/C) with xenografted



lung tumor cells (RL-67) were treated with different doses of the antidiotypic Mab according to the following shim: 1 mg/mouse, three times a week.

The therapeutic effect is measured through the comparison of survival curves of treated animals versus the control.

The biological effect of the antidiotypic Mabs in Infectious diseases was assessed using Balb/C mice previously infected with a clone of Plasmodium Chabaudi (Conquy-Adib.M et al (1993) Mol Immunology 30(2):119-127) After three days of the inoculation the mice were treated with the following therapeutic scheme:

1) A total dose of 800 ug injected in successive doses of 300, 100, 200, and 200 ug intraperitoneally during one week.

2) A high total doses of 3 mg was administered in 4 weekly injections (two of 1 mg and followed by two of 500 ug) a control group received 0.2 mL of saline solution in the same scheme.

Capacity for delaying the clinical signs appearance is evaluated comparing the control versus the treated group.

Example 7: Pharmaceutical composition containing antidiotypic monoclonal antibodies.

A therapeutic composition was obtained containing an effective amount of the antidiotypic monoclonal antibodies highly idiotypically connected and combinations of them, in a pharmaceutical vehicle composed by Dibasic Sodium phosphate 9.0 mg, Monobasic Sodium Phosphate 2.55 mg and Sodium Chloride 43.00 mg, polysorbato 80 1 mg, WFI 5.00 mL . This pharmaceutical composition is to be used in autoimmune diseases, infectious diseases and cancer.

Example 8: Murine Variable region of B7 monoclonal antibody DNA sequencing.

Cytoplasmic RNA was extracted from about 10^6 B7 hybridoma cells as described by Faloro et al (Faloro, J. et al. Methods in Enzimology 65:718-749, (1989)).

The cDNA synthesis reaction consisted of 5 ug RNA, 50 mM Tris- HCl, pH 7.5, 75 mM KCl, 10 mM DTT, 3 mM $MgCl_2$, 25 pmol CG2AFOR primer (5'GGAAGCTTAGACCGATGGGGCCTGTTGTTTGTG 3') for heavy chain variable region or CK2FOR (5'GGAAGCTTGAAGATGGATACAGTTGGTGCAGC 3') for light chain variable region, 250 uM each of dATP, dTTP, dCTP, dGTP, 15 u ribonuclease inhibitor (RNA guard, Pharmacy) in a total volume of 50ul. Samples were heated at $70^{\circ}C$, for 10 min. and slowly cooled to $37^{\circ}C$ over a period of 30 min. Then, 100units MMLV reverse transcriptase (BRL) were added and the incubation at $37^{\circ}C$ continued for 1 hour.

The VH and VK cDNAs were amplified using the PCR as described by Orlandi et al (Orlandi, R. et al. Proc. Natl.Acad.Sci.USA 86:3833- 3837, (1989)). For PCR amplification of VH, DNA/primer mixtures consisted of 5 ul cDNA, 25 pmoles CG2A FOR (5'GGAAGCTTAGACCGATGGGGCCTGTTGTTTGTG3')

and VH1 BACK primers (5'AGGT(G/C)(A/C)A(A/G)CTGCAG(G/C)AGTC(A/T)GG 3'.

For PCR amplification of VK, DNA/primers mixtures consisted of 5 ul cDNA, 25 pmoles of CK2 FOR (5'GGAAGCTTGAAGATGGATACAGTTGGTGCAGC 3') and

VK10BACK (5'TTGAATTCCAGTGATGTTTGTGATGACCCA 3') primers. To these mixtures were added 2.5mM each of dATP,dCTP,dTTP,and dGTP, 5 ul constituents of 10X buffer thermolase and 1 unit of Thermolase(ABI) in a final volume of 50 ul. Samples were subjected to 25 thermal cycles at



94°C, 30sec; 50°C, 30sec; 72°C, 1 min.; and a last incubation for 5 min. at 72°C. Amplified VH cDNA were purified on Prep. A Gene purification kit (BioRad).

The purified VH and were cloned into M13 vector. Clones were sequenced by the dideoxy method using T7 DNA Pol (Pharmacy). See figure 11.

Example 9: Modification of the variable domain sequences heavy chain of B7 murine monoclonal antibody to humanize the predicted T-cell antigenic sequences.

The variable region sequences of heavy of B7 were analyzed for T-cell antigenic sequences. It was made by using the computer algorithm AMPHI, which predict segments of the sequences 11 amino acids in length with an amphipatic helix structure, that is have one side hydrophobic and one side hydrophilic which bind to MHC II molecules.

On to the variable domain sequence of the heavy chain were predicted 3 segments which are: (It is used Kabat's numbering.).

1. FR 1 between amino acids 8-20.
2. FR 1 , CDR 1 and FR2 between amino acids 28-54.
3. FR3, CDR3 and FR4 between amino acids 88-109.

The Figure 11 shows the sequences corresponding to heavy chain.

This murine sequence is compared with the immunoglobulin sequences included in the Gene Bank and EMBL database. The most homologous human variable region sequence is determined and also the human subgroup to which the murine sequence most closely resembles is defined. In this case the human sequence founded was an TH9 Kabat's subgroup 1.

This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

Both variable region sequences, human and murine are then compared residue by residue and are selected those residues at FR regions which are not involved in the Venire zone or with the canonical structures. Therefore they could be changed by those residues at the same position onto the human sequence.

For the heavy chain of murine B7 we propose 13 replacements:

1. PRO at position 9 by ALA.
2. LEU at position 11 by VAL.
3. VAL at position 12 by LYS.
4. THR at position 16 by ALA.
5. ILE at position 20 by VAL.
6. MET at position 37 by VAL.
7. GLU at position 38 by ARG.
8. SER at position 40 by ALA.
9. HIS at position 41 by PRO.
10. LYS at position 43 by GLN.
11. SER at position 44 by GLY.
12. ILE at position 48 by MET.
13. SER at position 109 by VAL.

This Sheet Has Been Replaced



Example 10: Construction of mutant heavy chain variable region of B7 by PCR mutagenesis.

The changes in the amino acids of mutant heavy chain variable region were constructed using PCR mutagenesis (Kammann, M. et al. Proc. Nat. Aca. Sci. USA, 86, 4220-4224, (1989)).

Briefly: Two amplification by PCR: the reaction mixture was: 0.5 ul the VH supernatant of single strand DNA cloned in M13, 25 pmoles mutagenic oligo 1 or 2, 25 pmoles mutagenic oligo 3 or 4 primers (See below the primers

THIS SHEET HAS BEEN
REPLACED

sequences). To these mixtures were added 2.5mM each of dATP, dCTP, dTTP, and dGTP, 5 ul constituents of 10X Vent Polymerase buffer (NEB) and 1 unit of Vent DNA Polymerase (NEB) in a final volume of 50 ul. Samples were subjected to 12-15 thermal cycles at 94°C, 30 sec.; 50°C, 30 sec.; 75°C, 1 min.; and a last incubation for 5 min. at 75°C. The products of both PCRs are joined in a second PCR using the outside primers only (3 and 4). Amplified VH DNA were purified on Prep. A Gene purification kit (Bio Rad).

For the changes in the positions 9, 11, 12, 16 and 20 the primers used, were:

Primer1:

5'CTTGCAGGATACCTTGACTGAAGCCCCAGGCTTCTTCACCTCAGCTCC 3'.

Primer 3: 5'GTAAAACGACGGCCAGT 3'.

These primers are combined in one PCR.

Primer

2: 5'GGAGCTGAGGTGAAGAAGCCTGGGGCTTCAGTCAAGGTATCCTGCAAG3'.
Primer 4: 5'ACTGGCCGTCGTTTTAC 3'.

These primers are combine in one PCR.

Then, the products of both PCRs are combined in one PCR using 3 and 4 primers.

For the changes in the position 37, 38, 40, 41, 43, 44 and 48, the primers designed were:

Primer1:

5'AACACCTCCCATCCACTCAAGGCCCTGTCCAGGGGCCTGCCTCACCCAGTGCAT
GGT 3'.

Primer 3: 5'GTAAAACGACGGCCAGT 3'.

These primers are combined in one PCR.

Primer2:

5'ATGCACTGGGTGAGGCAGGCCCTGGACAGGGCCTTGAGTGGATGGGAGGT3'.

Primer 4: 5'ACTGGCCGTCGTTTTAC 3'.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

26

These primers are combined in one PCR.

Then, the products of both PCRs are combined in one PCR using 3 and 4 primers.

After mutagenesis VH genes were cloned in expression vectors (pSVgpt) yielding the plasmids B7-mut VH-pSVgpt.

BRIEF DESCRIPTION OF THE FIGURES.

Figure 1: Show the recognition pattern of the high connected antidiotypic hibridomas with different monoclonals antibodies.

Figure 2: Show the binding of the B7 Mab to different monoclonal antibodies.

Figure 3: Show the recognition of different human IgG mieloma proteins and the human intravenous IgG by the B7 Mab.

Figure 4: Show the binding of different murine Mabs by the human mieloma protein #2.

Figure 5: Show the recognition pattern of the IgG high connected antidiotypics Mabs by differents IgG human mieloma proteins.

Figure 6: Show the binding of undigested IgG human mieloma and their Fab'2 by the B7 Mabs.

Figure 7: Show the competitive inhibition by the intravenous human IgG of the B7 Mab binding to the human IgG mieloma protein.

Figure 8: Show the flow cytometric analysis to the peripheral blood human lymphocytes to different normal human donors.

A- Binding of B7 Mab to the human lymphocytes of different normal human donors (B7 Mab at 50 ug/ml).

B- Positive control (CD3 Mab).

C- Negative control (R3 Mab).

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Figure 9: Show the effect of the B7 Mab on human lymphocytes proliferative responses to CONA, PWM and PHA.

A- Human lymphocytes + CONA and PWM.

B- Human lymphocytes +PHA.

Figure 10: Show the effect of the human intravenous IgG on human lymphocyte proliferate responses to CONA and PHA.

Results are expressed as the mean cpm of triplicate values from three independent experiments.

A- iv IgG alone +iv IgG and CONA.

B- iv IgG+PHA.

Figure 11: Analysis for the modification by way of humanization of the variable regions of the heavy chain of antibody B7.

A: Sequence of the variable region of the murine B7 monoclonal antibody.

B: Sequence of the variable region of the most homologous human immunoglobulin.

C: Sequence of the modified variable region of B7.


Shading: predicted T-cell antigenic sequences.

Underlined amino acids residues: amino acids involved in tertiary structure.


Bold font: complementarity determining regions.

Amino acids residues in boxes: replacements which are proposed.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



CLAIMS.

1.- Antidiotypic monoclonal antibodies (AB2) IgG type highly connected to the idiotypic network which are useful as immunoregulators of the immune response.


2.- Antidiotypic monoclonal antibodies according to claim 1 which is the murine monoclonal antibody B7 obtained from the hybridoma of the same name (Deposit Number E.C.A.C.C. 94113324) as well as any humanized variant obtained from it.

3.- Monoclonal antibodies according to claims 1 and 2 which is a humanized variant obtained from the monoclonal antibody B7 which has a variable region of its heavy chain of sequence:

GLN VAL GLN LEU GLN GLN SER GLY ALA GLU VAL LYS
LYS PRO GLY ALA SER VAL LYS VAL SER CYS LYS THR
SER GLY TYR THR PHE THR GLU TYR THR MET HIS TRP
VAL ARG GLN ALA PRO GLY GLN GLY LEU GLU TRP MET
GLY GLY VAL SER PRO ASN ASN GLY GLY ALA SER TYR
ASN GLN LYS PHE LYS GLY LYS ALA THR LEU THR VAL
ASP LYS SER SER ASN THR ALA TYR MET GLU LEU ARG
SER LEU THR SER ASP ASP SER ALA VAL TYR TYR CYS
ALA ARG ARG LEU GLY ARG GLY TYR ASP LEU ALA SER
TYR TRP GLY GLN GLY THR LEU VAL LEU SER LEU GLN

4.- Use of the monoclonal antibodies of claim 1 to 3 for the manufacture of a medicament useful in the treatment of autoimmune diseases.

5.- Use of the monoclonal antibodies of claim 1 to 3 for the manufacture of a medicament useful in the treatment of infectious diseases..



6.- Use of the monoclonal antibodies of claim 1 to 3 for the manufacture of a medicament useful in the treatment of cancer.


7.- Use of the monoclonal antibodies of claim 1 to 3 for the manufacture of a reactive useful for the "in vivo" and "in vitro" diagnostic of autoimmune and infectious diseases and cancer.

8.- Pharmaceutical composition for the treatment of autoimmune diseases which contains any of the monoclonal antibodies of claims 1 to 3.

9.- Pharmaceutical composition for the treatment of infectious diseases which contains any of the monoclonal antibodies of claims 1 to 3.

10.- Pharmaceutical composition for the treatment of cancer which contains any of the monoclonal antibodies of claims 1 to 3.

This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

Claim 11:

11. Reactive useful for diagnostic of autoimmune and infectious diseases and cancer which contains any of the monoclonal antibodies of claims 1 to 3.

This Sheet Has Been Replaced



* C L M S *

THIS SHEET HAS BEEN
REPLACED

ABSTRACT

Monoclonal Antiidiotypic antibodies IgG type connected to the immune network, able to interact with T and B lymphocytes and able to exert an immunoregulatory effect, immunostimulation or immunosuppression that can be used in immunotherapy of autoimmune diseases, infectious diseases and cancer.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

02 FEB 1989

1**MONOCLONAL ANTIIDIOTYPIC ANTIBODIES (AB2) AND THEIR USES.****Field of the Invention.**

The present invention is related to the field of Immunology and in particular to the selection and use of antidiotypic monoclonal antibodies (AB2) IgG type with the main characteristic of being highly connected to the idiotypic network and recognition of B and T human lymphocytes, which are major participants in the immune response.

Accordingly the objective of this invention is to provide antidiotypic monoclonal antibodies IgG type connected to the immune network, able to interact with T and B lymphocytes and able to exert an immunoregulatory effect.

Description of the Prior Art.

The immune response is mainly based on the antigen stimulation of some determined B lymphocyte clones bearing in the surface specific receptors for this antigen. T cells have the ability of producing positive and negative signals as a way of regulation of this process. However the immune system regulation mechanisms are very complex as was postulated by Jerne (Jerne NK (1974) Ann Immunol. Inst. Pasteur 125, C:373:389) : the immune system can be "seen" as an organized structure of antibody molecules and interconnected lymphocytes by idiotypic directed interactions.

Idiotypes in antibodies are defined as epitopes in the variable regions in such a way each antibody presents particular idiotypical determinants and among them the antigen- antibody recognition region is included. The theory of the idiotypic network emerges from the demonstrated fact (Urbain .J (1976) Ann Immunol. 1237 C; 357) that the

02 FEB 1989

2

idiotypic determinants in the immunoglobulin molecule (AB1) are recognized by antidiotypic antibodies (AB2) obtained against it.

The binding of an idiotypic determinant with its corresponding antidiotypic occurs in a dynamic state that could be perturbed during the course of immune response.

The theory of the idiotypic network proposed by Jerne in the seventies has implicit that the immune repertoire contains and recognize all the possible antigens. Ignoring the dicotomy between self and non self antigens, thus the existence of autoreactive clones of lymphocytes occurs in a physiological mode.

Studies performed in the last ten years have demonstrated the presence of natural autoreactive antibodies in normal subjects (Avrameas.S., (1991) Immunol.. Today, 132C: 231: 236, Antin et al (1986), J Immunol. 136,505:510, Lundkvist (1989), PNAS USA 86:5074-5078, Longtenberg .T (1989)J Exp Med, 170:1347.1335)).

These experimental facts have prompted the concept that considers the immune system as a highly connected network of specific receptors ontogenically organized through the internal references (self recognition) philogenetically selected for the maintenance of the homeostatic physiological functions, and more than the normal defense against infections In this sense recognition of the self is normal and tolerance to the self and memory are not any more considered clone properties as postulates by Burnet in his clone selection theory (Burnet F.M. (1956) London Cambridge University Press) but they are emerging properties of the immune network with an integral point of view (Coutinho. A

02 FEB 1989

(1984) Immunol. Rev. 79:151-168, Coutinho. A (1989) Immunol. Rev 110:63-87).

This new concept in relation to the immune system have had its impact in understanding the basic mechanisms of autoimmune diseases and also in the new therapy approach Zanetti. M (1985) Immunol. Today 6: 299-302; Kaveri et al (1991) Clin Exp. Immunol. 86:192-198; Kazatchine. D.M (1994) Immunol. Rev 139:79-107.

An amazing therapeutic effect with different immunoglobulin compounds have observed in the wide variety of systemic and organ specific autoimmune diseases that is useful for the clinical validation of this new concept. (Kazatchine et al (1993) Autoimmunity: physiology and disease Wiley-Liss, Inc., New York).

IgG Immunoglobulin preparations are obtained from blood pools of healthy donors representing the autoreactive physiological network of antibodies. The reactivity analysis in these preparations have demonstrated the existence of antibodies interacting with idiotypes expressed in the autoantibodies and autoantibodies per se (Rossi, et al (1988) Clin Exp. Immunol. 74: 311-316; Rossi et al (1989) J Immunol. 143:4104-4109. Both types of reactivity are in high concentration in the total number of IgG molecules that are more connected than other molecules inside the pool (Dietrich et al (1992) Eur J Immunol. 22:1701-1706). In this way the immunoregulatory effect of these preparations justify the design of new therapeutic strategies addressed to restores the connectivity and diversity in the autoimmune patient. Through this clinical value the concept of emergency of autoimmune disease has been validated, through the fail in

02 FEB 1990

the regulatory network in the variable regions of the connected antibodies.

In previous research a panel of monoclonal antibodies against tumor associated gangliosides was obtained, the antigen, classified as T cell independent self antigen (EPO: No:0 657 471 A1) These Mabs are IgM type bearing idiotypes able to activate the whole idiotypic network and thus generating an antibody response in absence of the antigen, i.e. they are recurrent idiotypes. These antibodies have been previously described as recognizing specifically the NeucAc GM2 ganglioside (Alfonso. M et al (1995) Hibridoma 14(3): 209-216).

The present invention provides novelty in the generation and selection of a panel of antidiotypic monoclonal antibodies (AB2) against the original IgM antibody recognizing ganglioside.

From this panel a further characterization was performed to one of the antibodies, named B7 showing it was an IgG2a with medium affinity for the antigen (Kd 10 E-8) and alpha type even though this type, it was able to induce an antibody (AB1') independent response in a singeneic model (Balb/C).

Surprising properties have been shown in the antidiotypic antibody B7 studies: High idiotypic connectivity with immunoglobulins of different species human B and T lymphocyte recognition and the immunoregulatory effect on previously activated B and T lymphocytes.

Up to know there isn't any therapy proposing this property in antidiotype antibodies for the treatment of autoimmune, and infectious diseases and cancer.

The novelty of the present invention consists in the selection of the antidiotypic monoclonal antibodies IgG type

02 FEB 1989

5

highly connected to the immune network through their variable regions able to interact with human B and T lymphocytes and their immunoregulatory effect.

Detailed Description of the Invention.

GENERATION AND PRODUCTION OF MONOCLONAL ANTIDIOTYPIC ANTIBODIES HIGHLY CONNECTED TO THE IMMUNE NETWORK.

For the generation and production of monoclonal antidiotypic antibodies, the Mab against ganglioside is used as immunogen with an immunization procedure is performed as described elsewhere (European Patent Application 657 471 A1).

Mice with high titers of antidiotypic antibodies in serum are reimmunized and three days afterwards the spleen is removed and splenocytes are obtained. Antibody producing cells can be obtained from any other organ if preferred.

These cells are fused with myeloma cells, cell fusion may be performed according to any of the known methods and thus hybridomas are obtained. This method is described in detail in any of the procedures for this purpose.

The antibodies produced are assayed through any immunoassay method preferably an immunoenzymatic method, in which the supernatant are assessed for the presence of specific recognition of the proper antigen. The antibody- antibody binding is detected through a second antibody labeled with an enzyme.

The supernatants are selected for their capacity of blocking the antiganglioside Mab (E1) binding to the antigen (Neu Ac GM2) the supernatants in adequate concentrations are incubated and then the E1 Mab with the antigen.

02 FEB 1999

Once the desired hybridoma has been selected and cloned at least twice by limiting dilution, the resulting Mab can be produced by several culture techniques.

One method can be inoculating the proper amount of cells in the peritoneal cavity of mice producing ascitic fluid in which the Mab is produced in high concentration.

The antidiotypic Mabs obtained recognize different Mabs anti gangliosides related or not related and also are characterized by not blocking the binding of the anti ganglioside Mab to the corresponding antigen.

PROCEDURE FOR THE EVALUATION OF THE IDIOTYPIC CONNECTIVITY IN THE ANTIDIOTYPIC ANTIBODIES (IGG TYPE).

The evaluation of the idiotypic connectivity of antidiotypic monoclonal antibodies of IgG type obtained against the monoclonal antibody antiganglioside E1 recognizing the Neuc Ac GM2 ganglioside is performed using an indirect Immunoenzimatic assay in PVC plates. Plates are coated with specific antigangliosides Mabs and non specific Mabs of murine origin or any other species also human origin Mabs are used for this purpose and IgG pool. The wells are blocked during 1 hour approximately at room temperature, with PBS and Bovine serum albumin (1%) Followed by the addition of the antidiotypic monoclonal antibodies in a concentration range of 250 ug to 10 ug/mL during two hours in a humid chamber at 37°C . The plate is washed at least 4 times (PBS) and the anti mouse Fc labeled with alkaline Phosphatases conjugate is added during 4 hours. The substrate is added after that time and the plate is read at OD 405nm.

Negative controls are included, i.e. other antidiotypic monoclonal antibodies not connected to the idiotypic network.

highly connected to the immune network through their variable regions able to interact with human B and T lymphocytes and their immunoregulatory effect.

Detailed Description of the Invention.

GENERATION AND PRODUCTION OF MONOCLONAL ANTIDIOTYPIC ANTIBODIES HIGHLY CONNECTED TO THE IMMUNE NETWORK.

For the generation and production of monoclonal antidiotypic antibodies, the Mab against ganglioside is used as immunogen with an immunization procedure is performed as described elsewhere (European Patent Application 657 471 A1).

Mice with high titers of antidiotypic antibodies in serum are reimmunized and three days afterwards the spleen is removed and splenocytes are obtained. Antibody producing cells can be obtained from any other organ if preferred.

These cells are fused with myeloma cells, cell fusion may be performed according to any of the known methods and thus hybridomas are obtained. This method is described in detail in any of the procedures for this purpose.

The antibodies produced are assayed through any immunoassay method preferably an immunoenzymatic method, in which the supernatant are assessed for the presence of specific recognition of the proper antigen. The antibody- antibody binding is detected through a second antibody labeled with an enzyme.

The supernatants are selected for their capacity of blocking the antiganglioside Mab (E1) binding to the antigen (Neu Ac GM2) the supernatants in adequate concentrations are incubated and then the E1 Mab with the antigen.

02 FEB 1993

Once the desired hybridoma has been selected and cloned at least twice by limiting dilution, the resulting Mab can be produced by several culture techniques.

One method can be inoculating the proper amount of cells in the peritoneal cavity of mice producing ascitic fluid in which the Mab is produced in high concentration.

The antidiotypic Mabs obtained recognize different Mabs anti gangliosides related or not related and also are characterized by not blocking the binding of the anti ganglioside Mab to the corresponding antigen.

PROCEDURE FOR THE EVALUATION OF THE IDIOTYPIC CONNECTIVITY IN THE ANTIDIOTYPIC ANTIBODIES (IGG TYPE).

The evaluation of the idiotypic connectivity of antidiotypic monoclonal antibodies of IgG type obtained against the monoclonal antibody antiganglioside E1 recognizing the Neuc Ac GM2 ganglioside is performed using an indirect Immunoenzimatic assay in PVC plates. Plates are coated with specific antigangliosides Mabs and non specific Mabs of murine origin or any other species also human origin Mabs are used for this purpose and IgG pool. The wells are blocked during 1 hour approximately at room temperature, with PBS and Bovine serum albumin (1%) Followed by the addition of the antidiotypic monoclonal antibodies in a concentration range of 250 ug to 10 ug/mL during two hours in a humid chamber at 37°C . The plate is washed at least 4 times (PBS) and the anti mouse Fc labeled with alkaline Phosphatases conjugate is added during 4 hours. The substrate is added after that time and the plate is read at OD 405nm.

Negative controls are included, i.e. other antidiotypic monoclonal antibodies not connected to the idiotypic network.

02 FEB 1999

The antidiotypic monoclonal antibodies considered connected recognize the different antibodies coming from different species and their corresponding fragments and also recognize the Human pool IgG obtained from human healthy donors, indicating the high connectivity of these Mabs recognizing the idiotypes in the variable regions of immunoglobulins of other species.

PROCEDURE FOR THE EVALUATION OF DIFFERENT SPECIES B AND T LYMPHOCYTE RECOGNITION BY THE ANTIDIOTYPIC MABS (AB2) IGG TYPE IDIOTYPICALLY CONNECTED.

The evaluation of the recognition of B and T lymphocytes from different species (mouse, rat monkey and human) by the antidiotypic Mabs (AB2) IgG type idiotypically connected is performed by an Indirect Immunofluorescence assay in a Flow cytometer.

B and T lymphocytes from different species are isolated by the Ficoll gradient technique from peripheral blood and other lymphoid organs

(spleen, lymph nodes and peritoneum) according to the species studied. The cell ring is isolated and the cell counting is done with a Neubauer chamber.

The cell suspension (200 uL) is incubated in a humid chamber during 30 minutes together with the antidiotypic Mab (AB2). Positive controls are added (ior t3 or ior t4) in established conditions and a negative control is also included. A proper conjugate that could be anti mouse FITC or RPE is added and incubated during 30 min. and after that time the sample is measured in the Flow cytometer.

PROCEDURE FOR THE EVALUATION OF THE BIOLOGICAL EFFECT OF ANTIDIOTYPIC MABS (AB2) IGG TYPE IDIOTYPICALLY CONNECTED ON THE B AND T LYMPHOCYTES IN VITRO.

02 FEB 1993

For the evaluation of the biological effect of antidiotypic Mabs(AB2) IgG type idiotipically connected on the B and T lymphocytes of any mammal species a cell proliferation assay is performed by 3H-Thimidine incorporation.

The Mononuclear peripheral cells are isolated through a Ficoll gradient technique. The cells in amount of 100000 per well are incubated with the proper medium with 10 % serum in microplates . Lymphocyte activators as PHA in a concentration that can be 5 ug/ mL, Con A in the same concentration and PWM in the same concentration are added to the incubation in the presence of different concentrations of the antidiotypic Mabs (300 ug to 15 ug/ mL).

Controls are included in the assay , activators only, or lymphocyte cells only. Cultures are kept during 72 hours in a 5 % CO2 humid atmosphere at 37 oC.

3 H-Thymidine is added (1 uCi, 23 uCi/ mmol) to each well 6 hours before the culture is arrested. Cells are harvested and 3 H-Thymidine is measured by a beta counter (LKB Sweden) .

The highly connected antidiotypic mabs are able to suppress or enhance the stimulation of previously activated B and T lymphocytes of any species. The antidiotypic Mabs not connected are not able to exert this effect.

PROCEDURE FOR THE EVALUATION OF THE BIOLOGICAL EFFECT OF ANTIDIOTYPIC MABS (AB2) IGG TYPE IDIOTIPICALLY CONNECTED ON THE B AND T LYMPHOCYTES IN VIVO ON DIFFERENT ANIMAL MODELS FOR AUTOIMMUNE DISEASES, INFECTIOUS DISEASES AND MALIGNANT NEOPLASM'S.

The in vivo biological effect of Antidiotypic Mabs (AB2) IgG type idiotipically connected on the B and T lymphocytes for autoimmune diseases, infectious diseases and malignant

neoplasm's, was measured using different animals experimental models. The experimental model MRL/lpr/lpr mouse was used to evaluate B7 Mab effects in the autoimmune diseases. This animal develops the Disseminated Lupus Erythematosus (DSL).

MRL/lpr/lpr mice are intraperitoneally injected with high doses of the B7 antidiotypic Mab according to the following scheme: 1 mg/ mouse, three times a week, initiating the treatment 24 hours after the birth of the animal and maintaining it during 4 weeks.

Also a low dose regime is evaluated with 100 ug of the antidiotypic Mab at the time of the animal birth.

The therapeutic effect is measured through several aspects: the delayed in time of the clinical signs appearance (skin lesions lymph nodes); through the increase of animal survival and also through the variation of any biological parameter considered as indicator (Proteinuria, anti DNA autoantibodies). A control group of non treated animals was used to establish the comparison.

The evaluation of antitumor effect was performed using an experimental animal model. Mice (C57BL/C) with xenografted lung tumor cells (RL-67) were treated with different doses of the antidiotypic Mab according to the following scheme: 1 mg/ mouse, three times a week.

The therapeutic effect is measured through the comparison of survival curves of treated animals versus the control.

The biological effect of the antidiotypic Mabs in infectious diseases was assessed using Balb/C mice previously infected with a clone of Plasmodium Chabaudi.

02 FEB 1993

(Conquy-Adib. M et al (1993) Mol Immunology 30 (2):119-127). After three days of the inoculation the mice were treated with the following therapeutic scheme:

1) A total dose of 800 ug injected in successive doses of 300, 100, 200, and 200 ug intraperitoneally during one week.

2) A high total doses of 3 mg was administered in 4 weekly injections (two of 1 mg and followed by two of 500 ug) a control group received 0.2 mL of saline solution in the same scheme.

Capacity for delaying the clinical signs appearance is evaluated comparing the control versus the treated group.

PROCEDURE FOR HUMANIZATION AND REDUCE IMMUNOGENICITY OF THE MURINE B7 MONOCLONAL ANTIBODY.

B7 Mab was modified in order to decrease its immunogenicity with a procedure (according European patent bulletin # 1996/10 under publication no 0699755) which simultaneously reduces immunogenicity of the rodent monoclonal antibody while preserving its binding properties in their entirety. Since the antigenicity of an immunoglobulin is dependent on the presence of T-cell antigenic peptides onto their sequence, the immunogenicity of a xenogenic or allogenic antibody could be reduced by replacing the residues included onto the T-cell antigenic sequences which differ from those usually found in another mammalian species antibodies. Of course, the replacement of residues do not include those involved in to the canonical structures or in the Vernier zone. This judicious replacement of residues have no effect on the structural determinants or on the interdomain contacts, thus, ligand binding properties should be unaffected as a

consequence of alterations which are limited to the variable region framework residues.

(1)- Analysis of homology of variable regions.

The present procedure makes use of the available sequence data for human antibody variable domains compiled by Kabat et al., "Sequences of proteins of Immunological Interest" Fifth edition., Bethesda, Maryland; National Inst. of Health, 1994.

In the first step the variable domains of the murine B7 heavy or light chain are compared with those corresponding variable domains of the human sequences.

The comparison is made by an automated-computerized method (PC-DOS HIBIO PROSIS 06-00, Hitachi.). The most homologous human variable regions are then compared residue for residue to the corresponding murine regions. This will also define the human subgroup to which each mouse sequence most closely resembles.

(2)- Prediction of T-epitopes.

In the second step, the two homologous variable region sequences, mouse and human are analyzed for prediction of T-antigenic sequences.

The algorithm AMPHI (Bersofsky et al., The Journal of Immunology 138: 2213-2229, (1987)) predicts α Helical sequences. The algorithm SOHHA predicts the strip of helix hydrophobicity (Elliott et al., J. Immunol. 138: 2949-2952, (1987)). These algorithms predict T-cell presented fragments of antigenic proteins.

(3) Analysis for immunogenicity reduction.

Those residues in the mouse framework which differ from its human counterpart are replaced by the residues present

in the human counterpart. This switching occurs only with those residues which are at the T-antigenic sequences.

Finally, replacement of those residues responsible for the canonical structures or those involved in the Vernier zone could have a significant effect on the tertiary structure. Hence, they can not be included in the replacement. Additional information about the influence of the proposed replacements on tertiary structure or the binding site, could be obtained from a molecular model of the variable regions.

(4) The method for constructing and expressing the altered antibody.

The following procedures are used to prepare recombinant DNA sequences which incorporate the CDRs of the murine mAb, both light and heavy chains, into human, frameworks that can be used to transfect mammalian cells for the expression of recombinant antibody less immunogenic and with the antigen specificity of the animal monoclonal antibody:

a.-) mutagenesis and assembly of variable region domains including CDRs and FRs regions. The PCR-mutagenesis method (Kamman et al., Nucleic Acids Res. 17: 5404-5409, (1989)) is preferably used to introduce the changes at different positions.

b.-) preparation of an expression vector including one variable region and the corresponding human constant region which upon transfection into cells results in the secretion of protein sufficient for affinity and specificity determinations.

c.-) co-transfection of heavy and light chain expression vectors in appropriate cell lines.

02 FEB 1983

15

After about 2 weeks, the cell supernatants are analyzed by ELISA for human IgG production. The samples are then analyzed by any method for human IgG capable for binding to specific antigens.

Example 1: Generation and production of monoclonal antidiotypic antibodies (AB2) highly connected to the immune network against the E1 Mab obtained in the syngeneic model.

The Mab against ganglioside (E1 Mab) is used as immunogen in Balb/C mice (6-8 weeks old) with an immunization procedure performed as described elsewhere (EP No:0 667 471 A1).

Mice with high titers of antidiotypic antibodies in serum are reimmunized and three days afterwards the spleen is removed and splenocytes are obtained.

These cells are fused with myeloma cells, fusion is performed according to the method described by Kohler and Milstein with some minor modifications (Nature (1975) 256:495-497) and hybridomas are obtained. Murine splenocytes are fused with the non secretor murine myeloma cells P3/X63Ag66.5.3 in a rate 10:1 in 0.5 mL of fusion media containing PEG

(3000-5000).

The antibodies produced are assayed through any immunoassay method, in which the supernatants are assessed for the presence of specific recognition of the proper antigen. The antibody-antibody binding is detected through a second antibody labeled with an enzyme.

The supernatants are selected for their capacity of blocking the antiganglioside Mab (E1) binding to the antigen (Neu Ac GM2) the supernatants in adequate concentrations are incubated and then the E1 Mab with the antigen.

The antidiotypic Mabs are selected if they recognized different Mabs anti gangliosides related or not related (Figure 1) and also they are characterized by not blocking the binding of the anti ganglioside Mab to the corresponding antigen. The hybrid 4B7 was cloned twice and the resultant Mab was the antidiotypic Mab B7 an IgG2a class.

Example 2: Procedure for the evaluation of the idiotypic connectivity in the antidiotypic antibodies (IgG type).

The evaluation of the idiotypic connectivity of antidiotypic monoclonal antibodies of IgG type obtained against the monoclonal antibody E1 that recognize the Neuc Ac GM2 ganglioside is performed using an indirect Immunoenzimatic assay in PVC plates (96 wells, High Binding Costar). Plates are coated with specific antigangliosides Mabs including the one used as immunogen the E1 Mab, and non specific Mabs in carbonate -bicarbonate buffer. The wells are blocked during 1 hour at room temperature, with PBS and Bovine serum albumin (1%) Followed by the addition of the antidiotypic monoclonal antibodies in a concentration range of 250 ug to 10 ug/mL during two hours in a humid chamber at 37°C . The plate is washed at least 4 times (PBS) and the anti mouse Fc labeled with alkaline Phosphatase conjugate is added during 4 hours. The substrate is added after that time and the plate is read at OD 405 nm.

Negative controls are included, i.e. other antidiotypic monoclonal antibodies not connected to the idiotypic network.

The supernatants of different hydrides producing antidiotypic Mabs recognized the variable regions of the antiganglioside related Mabs, recognized non related IgM of Balb /C origin Mabs, and also neonatal Mabs. (Figure 1).

02 FEB 1999

17

The antidiotypic monoclonal antibody B7 IgG2a type showed a high idiotypic connectivity recognizing the idiotypes in the variable regions of the antiganglioside related Mabs, non related IgM of Balb/C origin Mabs, and also neonatal Mabs with an affinity of 10^{-8} M. (Figure 2).

Example 3: Characterization of human myeloma immunoglobulin (IgG) idiotypes recognition and in the therapeutic IgG preparation by the antidiotypic Mabs idiotypically connected.

The capacity of recognition of the human myeloma immunoglobulin (IgG) idiotypes (variable regions) by the B7 Mab was evaluated .

For this purpose an Indirect Immunoassay is used (96 wells, High Binding Costar). Plates are coated with different IgG fractions or their fragments

(obtained according Colligan. E.J et al, Eds (1995) Current Protocols in Immunology). The wells are coated with myeloma proteins in carbonate-bicarbonate buffer and blocked during 1 hour at room temperature, with PBS and Bovine serum albumin (1%) Followed by the addition of the antidiotypic monoclonal antibodies in a concentration range of 380 ug to 11 ug/mL during two hours in a humid chamber at 37°C . The plate is washed at least 4 times (PBS) and the anti goat Fc region labeled with alkaline Phosphates is added and incubated during 4 hours. The substrate is added after that time and the plate is read at OD 405 nm.

The B7 antidiotypic monoclonal antibody recognized the variable regions

02 FEB 1969

(idiotypes) present in the human myeloma immunoglobulins (Figure 3). Other murine Mabs not idiotipically connected did not show any reactivity with this IgG fractions (Figure 4).

The supernatants of other hydrides producing high connected antidiotypic Mabs recognized the variable regions of the IgG human myeloma proteins. The recognition profile was different for each IgG human myeloma protein. (Figure 5).

The B7 antidiotypic monoclonal antibody also recognized the variable regions (idiotypes) present in the human myeloma immunoglobulins fragments (Figure 6).

Moreover the recognition specificity was evaluated through a competence assay for the recognition of the human myeloma IgG by the B7 Mab and different concentrations of the therapeutic human IgG the experiment was performed as followed:

Different concentrations of the therapeutic human IgG were incubated overnight with a fixed concentration of the B7 Mab (95 ug/mL) after that time this mixture was incubated with the human myeloma IgG using the immunoassay already described in this example.

Increasing concentrations of therapeutic human IgG inhibits the B7 Mab recognition for the human myeloma IgG idiotypes. (Figure 7).

The fact that this B7 Mab of murine origin is able to recognize the variable regions of a human myeloma IgG and the therapeutic human IgG show us the high idiotipic connectivity and suggest that the zones recognized in the variable regions are highly conserved among species.

Example 4: Evaluation of human B and T Lymphocyte recognition by the antidiotypic Mabs (AB2) IgG type idiotipically connected.

02 FEB 1990

19

The evaluation of human B and T lymphocytes recognition by the antidiotypic Mabs(AB2) IgG type idiotipically connected is performed by an Indirect Immunofluorescence assay in a Flow cytometer.

B and T lymphocytes are isolated by the Ficoll gradient technique from peripheral blood. The cell ring is isolated and the cell counting is done with a Neubauer chamber. Concentration and cell viability were determined.

The cell suspension (200 uL) adjusted to 10^7 cells is incubated in a humid chamber during 30 min. together with the antidiotypic Mab (AB2) a standard curve is added in the concentration range of 250 to 10 ug/mL. Positive controls are added (ior t3 or ior t4) in established conditions and a negative control is also included (ior r3). After incubation the wells with the cell suspension are washed with PBS BSA Azide and centrifuging eliminating the supernatant and centrifuge again four times. An anti mouse FITC conjugate is added and incubated during 30 min. in humid chamber and after that the sample is washed and centrifuged again four times, finally the cell suspension is resuspended in 100 uL of PBS and the sample is measured in the Flow cytometer.

Figure (8A) shows the recognition analysis of human B and T lymphocytes by the B7 Mab in the Flow Cytometer. Figure (8B) shows the negative control (ior r3 Mab, a non idiotipically connected Mab).

The B7 Mab was able to recognize 53 % of human lymphocytes from a normal donor (Peripheral blood lymphocytes) Therefore this Mab of idiotypic reactivity is able to recognize molecular structures present in T and B human lymphocytes.

Example 5: Evaluation of the biological effect of B7 antidiotypic Mab(AB2) IgG type idiotipically connected on the B and T lymphocytes in vitro.

For the evaluation of the biological effect of B7 antidiotypic Mabs (AB2) IgG type idiotipically connected on the B and T lymphocytes of any mammal species a cell proliferation assay is performed by 3H-Thymidine incorporation.

The peripheral blood lymphocyte cells (PBL) are isolated through a Ficoll gradient technique. The cells in amount of 100000 per well are incubated RPMI 1640 medium with 10 % serum in microplates . Lymphocyte activators as PHA in a concentration that can be 5 ug/ mL, Con A in the same concentration and PWM in the same concentration are added to the incubation in the presence of different concentrations of the antidiotypic Mabs (300 ug to 15 ug/ mL). Controls are included in the assay, activators only, or lymphocyte cells only. Cultures are kept during 72 hours in a 5 % CO2 humid atmosphere at 37°C.

3 H-Thymidine is added (1 uCi, 23 uCi/ mmol) to each well 6 hours before the culture is arrested. Cells are harvested and 3 H-Thymidine is measured by a beta counter (LKB Sweden).

The B7 highly connected antidiotypic mab was able to suppress the stimulation of previously activated B and T lymphocytes (Figure 9 A and B). This suppresser effect of the mytogenic stimulation was also produced by therapeutic human IgG (Figure 10 A and B) The antidiotypic Mabs not connected are not able to exert this effect.

The fact that this B7 Mab exerts a biological suppresser effect in previously activated B and T human lymphocytes

demonstrates the immunoregulator capacity of this highly connected antidiotypic MAB.

Example 6: Evaluation of the biological effect of antidiotypic Mabs

(AB2) IgG type idiotypically connected on the B and T lymphocytes in vivo on different animal models for autoimmune diseases, infectious diseases and malignant neoplasm's.

The in vivo biological effect of antidiotypic Mabs (AB2) IgG type idiotypically connected on the B and T lymphocytes for autoimmune diseases, infectious diseases and malignant neoplasm's, was measured using different animals experimental models. The experimental model MRL/lpr/lpr mouse was used to evaluated B7 Mab effects in the autoimmune diseases. This animal develops the Disseminated Lupus Erythematosus (DSL).

MRL/lpr/lpr mice are intraperitoneally injected with high doses of the B7 antidiotypic Mab according to the following scheme: 1 mg/ mouse, three times a week, initiating the treatment 24 hours after the birth of the animal and maintaining it during 4 weeks.

Also a low dose regime is evaluated with 100 ug of the antidiotypic Mab at the time of the animal birth.

The therapeutic effect is measured through several aspects: the delayed in time of the clinical signs appearance (skin lesions lymph nodes); through the increase of animal survival and also through the variation of any biological parameter considered as indicator (Proteinuria, anti DNA autoantibodies). A control group of non treated animals was used to establish the comparison.

The evaluation of antitumor effect was performed using an experimental animal model. Mice (C57/BL/C) with xenografted

02 FEB 1993

lung tumor cells (RL-67) were treated with different doses of the antidiotypic Mab according to the following shim: 1 mg/mouse, three times a week.

The therapeutic effect is measured through the comparison of survival curves of treated animals versus the control.

The biological effect of the antidiotypic Mabs in Infectious diseases was assessed using Balb/C mice previously infected with a clone of Plasmodium Chabaudi (Conquy-Adib.M et al (1993) Mol Immunology 30(2):119-127) After three days of the inoculation the mice were treated with the following therapeutic scheme:

1) A total dose of 800 ug injected in successive doses of 300, 100, 200, and 200 ug intraperitoneally during one week.

2) A high total doses of 3 mg was administered in 4 weekly injections (two of 1 mg and followed by two of 500 ug) a control group received 0.2 mL of saline solution in the same scheme.

Capacity for delaying the clinical signs appearance is evaluated comparing the control versus the treated group.

Example 7: Pharmaceutical composition containing antidiotypic monoclonal antibodies.

A therapeutic composition was obtained containing an effective amount of the antidiotypic monoclonal antibodies highly idiotypically connected and combinations of them, in a pharmaceutical vehicle composed by Dibasic Sodium phosphate 9.0 mg, Monobasic Sodium Phosphate 2.55 mg and Sodium Chloride 43.00 mg, polisorbato 80 1 mg, WFI 5.00 mL . This pharmaceutical composition is to be used in autoimmune diseases, infectious diseases and cancer.

Example 8: Murine Variable region of B7 monoclonal antibody DNA sequencing.

REPLACEMENT SHEET

02 FEB 1999

Cytoplasmic RNA was extracted from about 10^6 B7 hybridoma cells as described by Faloro et al (Faloro, J. et al. Methods in Enzimology 65:718-749, (1989)).

The cDNA synthesis reaction consisted of 5 ug RNA, 50 mM Tris- HCl, pH 7.5, 75 mM KCl, 10 mM DTT, 3 mM $MgCl_2$, 25 pmol CG2AFOR primer (5'GGAAGCTTAGACCGATGGGGCCTGTTGTTTGG 3') for heavy chain variable region or CK2FOR (5'GGAAGCTTGAAGATGGATACAGTTGGTGCAGC 3') for light chain variable region, 250 uM each of dATP, dTTP, dCTP, dGTP, 15 u ribonuclease inhibitor (RNA guard, Pharmacy) in a total volume of 50ul. Samples were heated at $70^{\circ}C$, for 10 min. and slowly cooled to $37^{\circ}C$ over a period of 30 min. Then, 100units MMLV reverse transcriptase (BRL) were added and the incubation at $37^{\circ}C$ continued for 1 hour.

The VH and VK cDNAs were amplified using the PCR as described by Orlandi et al (Orlandi, R. et al. Proc. Natl.Acad.Sci.USA 86:3833- 3837, (1989)). For PCR amplification of VH, DNA/primer mixtures consisted of 5 ul cDNA, 25 pmoles CG2A FOR (5'GGAAGCTTAGACCGATGGGGCCTGTTGTTTGG3')

and VH1 BACK primers (5'AGGT(G/C) (A/C)A(A/G)CTGCAG(G/C)AGTC(A/T)GG 3'.

For PCR amplification of VK, DNA/primers mixtures consisted of 5 ul cDNA, 25 pmoles of CK2 FOR (5'GGAAGCTTGAAGATGGATACAGTTGGTGCAGC 3') and

VK10BACK (5'TTGAATTCCAGTGATGTTTGGATGACCCA 3') primers. To these mixtures were added 2.5mM each of dATP, dCTP, dTTP, and dGTP, 5 ul constituents of 10X buffer thermolase and 1 unit of Thermolase (IBI) in a final volume of 50 ul. Samples were subjected to 25 thermal cycles at

02 FEB 1998

94⁰C, 30sec; 50⁰C, 30sec; 72⁰C, 1 min.; and a last incubation for 5 min. at 72⁰C. Amplified VH cDNA were purified on Prep. A Gene purification kit (BioRad).

The purified VH and were cloned into M13 vector. Clones were sequenced by the dideoxy method using T7 DNA Pol (Pharmacy). See figure 11.

Example 9: Modification of the variable domain sequences heavy chain of B7 murine monoclonal antibody to humanize the predicted T-cell antigenic sequences.


The variable region sequences of heavy of B7 were analyzed for T-cell antigenic sequences. It was made by using the computer algorithm AMPHI, which predict segments of the sequences 11 amino acids in length with an amphipatic helix structure, that is have one side hydrophobic and one side hydrophilic which bind to MHC II molecules.

On to the variable domain sequence of the heavy chain were predicted 3 segments which are: (It is used Kabat's numbering.).

1. FR 1 between amino acids 8-20.
2. FR 1 , CDR 1 and FR2 between amino acids 28-54.
3. FR3, CDR3 and FR4 between amino acids 88-109.

The Figure 11 shows the sequences corresponding to heavy chain.

This murine sequence is compared with the immunoglobulin sequences included in the Gene Bank and EMBL database. The most homologous human variable region sequence is determined and also the human subgroup to which the murine sequence most closely resembles is defined. In this case the human sequence founded was an TH9 Kabat's subgroup 1.



Both variable region sequences, human and murine are then compared residue by residue and are selected those residues at FR regions which are not involved in the Venire zone or with the canonical structures. Therefore they could be changed by those residues at the same position onto the human sequence.


For the heavy chain of murine B7 we propose 13 replacements:

1. PRO at position 9 by ALA.
2. LEU at position 11 by VAL.
3. VAL at position 12 by LYS.
4. THR at position 16 by ALA.
5. ILE at position 20 by VAL.
6. MET at position 37 by VAL.
7. GLU at position 38 by ARG.
8. SER at position 40 by ALA.
9. HIS at position 41 by PRO.
10. LYS at position 43 by GLN.
11. SER at position 44 by GLY.
12. ILE at position 48 by MET.
13. SER at position 109 by VAL.

Example 10: Construction of mutant heavy chain variable region of B7 by PCR mutagenesis.

The changes in the amino acids of mutant heavy chain variable region were constructed using PCR mutagenesis (Kammann, M. et al. Proc. Nat. Aca. Sci. USA, 86, 4220-4224, (1989)).

Briefly: Two amplification by PCR: the reaction mixture was: 0.5 ul the VH supernatant of single strand DNA cloned in M13, 25 pmoles mutagenic oligo 1 or 2, 25 pmoles mutagenic oligo 3 or 4 primers (See below the primers



sequences). To these mixtures were added 2.5mM each of dATP, dCTP, dTTP, and dGTP, 5 ul constituents of 10X Vent Polymerase buffer (NEB) and 1 unit of Vent DNA Polymerase (NEB) in a final volume of 50 ul. Samples were subjected to 12-15 thermal cycles at 94°C, 30 sec.; 50°C, 30 sec.; 75°C, 1 min.; and a last incubation for 5 min. at 75°C. The products of both PCRs are joined in a second PCR using the outside primers only (3 and 4). Amplified VH DNA were purified on Prep. A Gene purification kit (Bio Rad).

For the changes in the positions 9, 11, 12, 16 and 20 the primers used, were:

Primer1:

5'CTTGCAAGGATACCTTGACTGAAGCCCCAGGCTTCTTCACCTCAGCTCC 3'.

Primer 3: 5'GTAAAACGACGGCCAGT 3'.

These primers are combined in one PCR.

Primer

2: 5'GGAGCTGAGGTGAAGAAGCCTGGGGCTTCAGTCAAGGTATCCTGCAAG3'. Pri

mer 4: 5'ACTGGCCGTCGTTTTAC 3'.

These primers are combine in one PCR.

Then, the products of both PCRs are combined in one PCR using 3 and 4 primers.

For the changes in the position 37, 38, 40, 41, 43, 44 and 48, the primers designed were:

Primer1:

5'AACACCTCCCATCCACTCAAGGCCCTGTCCAGGGCCTGCCTCACCCAGTGCAT
GGT 3'.

Primer 3: 5'GTAAAACGACGGCCAGT 3'.

These primers are combined in one PCR.

Primer2:

5'ATGCACTGGGTGAGGCAGGCCCTGGACAGGGCCTTGAGTGGATGGGAGGT3'.

Primer 4: 5'ACTGGCCGTCGTTTTAC 3'.

02 FEB 1999

27

These primers are combined in one PCR.

Then, the products of both PCRs are combined in one PCR using 3 and 4 primers.

After mutagenesis VH genes were cloned in expression vectors (pSVgpt) yielding the plasmids B7-mut VH-pSVgpt.

BRIEF DESCRIPTION OF THE FIGURES.

Figure 1: Show the recognition pattern of the high connected antidiotypic hibridomas with different monoclonals antibodies.

Figure 2: Show the binding of the B7 Mab to different monoclonal antibodies..

Figure 3: Show the recognition of different human IgG mieloma proteins and the human intravenous IgG by the B7 Mab.

Figure 4: Show the binding of different murine Mabs by the human mieloma protein #2.

Figure 5: Show the recognition pattern of the IgG high connected antidiotypics Mabs by differents IgG human mieloma proteins.

Figure 6: Show the binding of undigested IgG human mieloma and their Fab'2 by the B7 Mabs.

Figure 7: Show the competitive inhibition by the intravenous human IgG of the B7 Mab binding to the human IgG mieloma protein.

Figure 8: Show the flow cytometric analysis to the peripheral blood human lymphocytes to different normal human donors.

A- Binding of B7 Mab to the human lymphocytes of different normal human donors (B7 Mab at 50 ug/ml).

B- Positive control (CD3 Mab).

C- Negative control (R3 Mab).

02 FEB 1980

Figure 9: Show the effect of the B7 Mab on human lymphocytes proliferative responses to CONA, PWM and PHA.

A- Human lymphocytes + CONA and PWM.

B- Human lymphocytes +PHA.

Figure 10: Show the effect of the human intravenous IgG on human lymphocyte proliferate responses to CONA and PHA.

Results are expressed as the mean cpm of triplicate values from three independent experiments.

A- iv IgG alone +iv IgG and CONA.

B- iv IgG+PHA.

Figure 11: Analysis for the modification by way of humanization of the variable regions of the heavy chain of antibody B7.

A: Sequence of the variable region of the murine B7 monoclonal antibody.

B: Sequence of the variable region of the most homologous human immunoglobulin.

C: Sequence of the modified variable region of B7.

Shading: predicted T-cell antigenic sequences.

Underlined amino acids residues: amino acids involved in tertiary structure.

Bold font: complementarity determining regions.

Amino acids residues in boxes: replacements which are proposed.

CLAIMS.

02 FEB 1993

1.- Anti-idiotypic monoclonal antibody (AB2) IgG type highly connected to the idiotypic network which is a humanized variant obtained from the monoclonal antibody B7 produced by the hybridoma of the same name, accession number E.C.A.C.C. 94113324 which is useful as immunoregulator of the immune response and the variable region of its heavy chain has the sequence:

GLN VAL GLN LEU GLN GLN SER GLY ALA GLU VAL LYS
LYS PRO GLY ALA SER VAL LYS VAL SER CYS LYS THR
SER GLY TYR THR PHE THR GLU TYR THR MET HIS TRP
VAL ARG GLN ALA PRO GLY GLN GLY LEU GLU TRP MET
GLY GLY VAL SER PRO ASN ASN GLY GLY ALA SER TYR
ASN GLN LYS PHE LYS GLY LYS ALA THR LEU THR VAL
ASP LYS SER SER ASN THR ALA TYR MET GLU LEU ARG
SER LEU THR SER ASP ASP SER ALA VAL TYR TYR CYS
ALA ARG ARG LEU GLY ARG GLY TYR ASP LEU ALA SER
TYR TRP GLY GLN GLY THR LEU VAL LEU SER LEU GLN

2.- Use of the monoclonal antibody of claim 1 for the manufacture of a medicament useful in the treatment of autoimmune diseases.

3.- Use of the monoclonal antibody of claim 1 for the manufacture of a medicament useful in the treatment of infectious diseases.

4.- Use of the monoclonal antibody of claim 1 for the manufacture of a medicament useful in the treatment of cancer.

5.- Use of the monoclonal antibody of claim 1 for the manufacture of a reagent useful for the "in vivo" and "in vitro" diagnosis of autoimmune and infectious diseases and cancer.



* C L H S *

02 FEB 1993

6.- Pharmaceutical composition for the treatment of autoimmune diseases which contains the monoclonal antibody of claim 1.


7.- Pharmaceutical composition for the treatment of infectious diseases which contains the monoclonal antibody of claim 1.

8.- Pharmaceutical composition for the treatment of cancer which contains the monoclonal antibody of claim 1.

9. Reagent for diagnosis of autoimmune diseases, infectious diseases and cancer which contains the monoclonal antibody of claim 1.



* C L M S *



ABSTRACT

Monoclonal Antiidiotypic antibodies IgG type connected to the immune network, able to interact with T and B lymphocytes and able to exert an immunoregulatory effect, immunostimulation or immunosuppression that can be used in immunotherapy of autoimmune diseases, infectious diseases and cancer.

11 AUG 1993

This Sheet Has Been Replaced



CLAIMS.

1.- Anti-idiotypic monoclonal antibody (AB2) IgG type highly connected to the idiotypic network characterized by it is a humanized variant obtained from the monoclonal antibody B7 obtained from the hybridoma of the same name, Deposit Number E.C.A.C.C. 94113324, and which are useful as immunoregulators of the immune response.

2.- The monoclonal antibody of claim 1 characterized by it has a variable region of its heavy chain of sequence:

GLN VAL GLN LEU GLN GLN SER GLY ALA GLU VAL LYS
LYS PRO GLY ALA SER VAL LYS VAL SER CYS LYS THR
SER GLY TYR THR PHE THR GLU TYR THR MET HIS TRP
VAL ARG GLN ALA PRO GLY GLN GLY LEU GLU TRP MET
GLY GLY VAL SER PRO ASN ASN GLY GLY ALA SER TYR
ASN GLN LYS PHE LYS GLY LYS ALA THR LEU THR VAL
ASP LYS SER SER ASN THR ALA TYR MET GLU LEU ARG
SER LEU THR SER ASP ASP SER ALA VAL TYR TYR CYS
ALA ARG ARG LEU GLY ARG GLY TYR ASP LEU ALA SER
TYR TRP GLY GLN GLY THR LEU VAL LEU SER LEU GLN

3.- Use of the monoclonal antibody of claims 1 and 2 for the manufacture of a medicament useful in the treatment of autoimmune diseases.

4.- Use of the monoclonal antibody of claims 1 and 2 for the manufacture of a medicament useful in the treatment of infectious diseases.



* C L M S *

THIS SHEET HAS BEEN
REPLACED

11 AUG 1988

- 5.- Use of the monoclonal antibody of claims 1 and 2 for the manufacture of a medicament useful in the treatment of cancer.
6. - Use of the monoclonal antibody of claims 1 and 2 for the manufacture of a reagent useful for the "in vivo" and "in vitro" diagnosis of autoimmune and infectious diseases and cancer.
- 7.- Pharmaceutical composition for the treatment of autoimmune diseases which contains the monoclonal antibody of claims 1 and 2.
- 8.- Pharmaceutical composition for the treatment of infectious diseases which contains the monoclonal antibody of claims 1 and 2.
- 9.- Pharmaceutical composition for the treatment of cancer which contains the monoclonal antibody of claims 1 and 2.

This Sheet Has Been Replaced

* C L M S *

THIS SHEET HAS BEEN
REPLACED