

Description

This invention relates to novel hair rinse compositions and more particularly, to hair rinse compositions which comprise a minor proportion of cationic surface active agents and a major proportion of oils, but free of hydrophilic nonionic surface active agents. Such hair rinse compositions have good stability and high rinsing performance.

Various troubles after hair washing have been heretofore removed by rinsing the hair with hair rinse compositions which comprise 1 to 5 wt% (hereinafter referred to simply as %) of quaternary ammonium salts such as distearyldimethylammonium chloride as the effective component thereof.

A hair rinse composition comprising a cationic surface active agent and a higher alcohol emulsified in water is described in FR—A—2 357 240.

A hair rinse composition comprising monoglyceride instead of a higher alcohol is known from FR—A—2 345 142.

The hair rinse is used to impart to hair flexibility, smoothness and antistaticity, but use of quaternary ammonium salts alone is not sufficient to give satisfactory performance as to flexibility and smoothness. In order to avoid these drawbacks, it is used to further add oils such as higher alcohols, glycerides, and liquid paraffins. Since, however, quaternary ammonium salts have no ability of stably emulsifying or dispersing oils in amounts sufficient to show their performances, attempts were made to add nonionic active agents of higher hydrophilicity so as to stabilize the dispersion system. Highly hydrophilic nonionic active agents have the action of considerably lowering the rinsing performance. Hair rinses which comprise quaternary ammonium salts, oils, and nonionic surface active agents of high hydrophilicity do not show satisfactory rinsing performances. On the other hand, cationic surface active agents are able to stably emulsify several or larger times of oils but it was difficult to obtain hair rinse compositions with excellent rinsing performances using such cationic active agents.

Thus, known hair rinse compositions are not necessarily satisfactory with respect to rinsing performance, i.e. of imparting flexibility, smoothness, and antistaticity to the hair. There is a demand for develop hair rinses compositions of better performance.


We have made intensive studies to overcome the drawbacks of the prior art hair rinses and, as a result, found that hair rinse compositions of good rinsing performance can be obtained by first preparing an emulsion of high concentration from a small amount of a cationic surface active agent and a specific type of oil, and diluting the emulsion with water whereby the oil can be stably emulsified in large amounts without use of any nonionic surface active agent of high hydrophilicity.

According to the present invention, there is provided a hair rinse composition which comprises

(A) 0.05 to 0.5 wt.-% of cationic surface active agent, and

(B) a higher alcohol having a linear or branched alkyl or alkenyl group having from 12 to 26 carbon atoms, or a glycerine mono fatty acid ester having a melting point not lower than 45°C, and (B) is contained in an amount of 3 to 15 times by weight to ingredient (A) and said composition is produced by diluting a highly concentrated emulsified product containing 0.4 to 34 wt.-% of ingredient (A) and (B) in total with 1 to 20 times by weight to the emulsified product of water.

The cationic surface active agent which is ingredient (A) of the invention is not limited to any specific ones. All cationic surface active agents which are ordinarily used in hair rinse agents may be used in the practice of this invention. Preferable quaternary ammonium salts are those having the following formula (I):



in which one or two of R_1 , R_2 , R_3 , and R_4 independently represent a linear or branched alkyl or hydroxyalkyl group having from 8 to 22 carbon atoms, the others independently represent an alkyl group having from 1 to 3 carbon atoms, hydroxyalkyl group, benzyl group, or polyoxyethylene group in which the total number of added moles is not larger than 10, and X represents a halogen atom or alkylsulfate group having 1 to 2 carbon atoms.

Of the quaternary ammonium salts, the following compounds are more preferable:

distearyldimethylammonium chloride,
stearyltrimethylammonium methosulfate,
stearyltrimethylammonium chloride,
stearyldimethylbenzylammonium chloride,
docosyltrimethylammonium methosulfate,

docosyltrimethylammonium chloride,
 docosyldimethylbenzylammonium chloride,
 didocosyldimethylammonium chloride,
 lauryldiethylbenzylammonium chloride,
 lauryltrimethylammonium bromide,
 distearyl methylhydroxymethyl chloride,
 cetyltrimethylammonium chloride,
 N-stearyl-N,N,N-tri(polyoxyethylene)ammonium chloride (3 moles added in total),
 cetyltriethylammonium bromide and
 stearyldimethylammonium chloride.

The ingredient (A) is generally added in an amount of from 0.05 to 0.5 wt% of the total composition. Outside the range, a satisfactory rinsing performance cannot be achieved.

The higher alcohols which are ingredient (B) are alcohols having a linear or branched alkyl or alkenyl group containing from 12 to 26 carbon atoms. Preferable examples include cetostearyl alcohol (mixture of cetyl alcohol and stearylalcohol), cetyl alcohol, arachidic alcohol, behenyl alcohol, calnabil alcohol, and ceryl alcohol.

The glycerine monofatty acid esters are represented by the following general formula (II)




in which R_5 represents a linear or branched hydrocarbon group having from 11 to 23 carbon atoms. Preferable examples are glycerides composed mainly of oleic acid monoglyceride, palmitic acid monoglyceride, isostearic acid monoglyceride, stearic acid monoglyceride, and behenic acid glyceride. Commercially available products called monoglycerides usually contain, aside from monoesters, large amounts of diesters, triesters and unreacted glycerine and are not suitable for the purpose of the invention. In order to yield good rinsing performance, it is necessary that not lower than 80% of glycerine monofatty acid esters indicated above be contained. The ingredient (B) used in the present invention should have a melting point not lower than 45°C. The esters whose melting temperature is lower than 45°C are more oleophilic and cannot give good feeling to the touch.

The amount of ingredient (B) is preferably in the range from 3 to 15 times as large as the ingredient (A). Less amounts do not show any appreciable rinsing performance, whereas larger amount are unfavorable in that stable emulsification is not accomplished.

The water-soluble polymer compounds used as the ingredient (C) are, for example, cellulose derivatives, polyvinyl alcohol, polyvinylpyrrolidone, sodium polyacrylate, carboxyvinyl polymer and polyethyleneimine.

Of these, there are preferred cellulose derivatives of the formula (III)



in which R_6 s independently represent hydrogen, $-(\text{CH}_2\text{CH}_2\text{O})_m-\text{H}$, $-(\text{CH}_2\text{CH}_2\text{CH}_2\text{O})_m-\text{H}$ (wherein m is an integer of from 1 to 5), $-\text{CH}_3$, $-\text{C}_2\text{H}_5$, or $-\text{COONa}$, and n is an integer of from 5 to 5000. Most preferably, hydroxyethylcellulose and hydroxymethylcellulose are used.

The ingredient (C) does not show its performance when used in amounts less than 0.1%, and any further performance cannot be expected when the amounts exceed 5%. Accordingly, the amount ranging from 0.1 to 5% is preferable.

The stable hair rinse composition of the invention cannot be obtained merely by emulsifying ingredients (A) and (B), or ingredients (A), (B) and (C) in the defined ranges of amounts of water. In order to obtain the hair rinse composition of the invention, 0.1 to 9% of ingredient (A) and ingredient (B) in an amount of 3 to 15 times by weight as large as the ingredient (A) are added to water in such a way that the total amount of both ingredients reaches 0.4 to 34%, followed by heating under agitation to prepare a highly concentrated emulsion. Subsequently, the emulsion was diluted with 1 to 20 times by weight, preferably 2 to 15 times by weight, as large as water so that the content of ingredient (A) is within the above-defined range (0.05 to 0.5%). Ingredient (C) is preferably used after dispersion in water. The hair

rinse composition may further comprise humectants, perfumes and colorants which are ordinary used for these purposes, if necessary.

The thus obtained hair rinse compositions of the invention are stable under either low temperature or high temperature conditions, with very good rinsing performance.

The present invention is illustrated by way of examples, which should not be construed as limiting the present invention thereto. In examples, the following test methods were used.

(1) Appearance

A sample was placed in a 100 ml transparent glass container and its appearance was visually observed. It will be noted that if a sample has bubbles therein, it was subjected to centrifugal separation for defoaming.

O = homogeneous; separation or coagulation is not observed.

X = heterogeneous; separation and/or coagulation is observed.

(2) High temperature stability

A sample was placed in a transparent glass container, tightly sealed, and kept in a thermostatic chamber of 50°C for one month. Thereafter, phase separation of the sample was visually observed.

O = homogeneous; separation or coagulation is not observed.

X = heterogeneous; separation and/or coagulation is observed.

(3) Low temperature stability

A sample was placed in a transparent glass container, tightly sealed, and kept in a thermostatic chamber of -5°C for one month. Thereafter, separation or solidification were visually observed.

O = homogeneous; separation or coagulation is not observed.

X = heterogeneous; separation and/or coagulation is observed.

(4) Emulsified state

A highly concentrated emulsion which was cooled to room temperature was observed visually or through a microscope. Moreover, 5 g of the highly concentrated emulsion was placed in 100 ml of water and agitated at a normal temperature for 1 hour, after which the dispersed state of the resulting emulsion was observed.

O = Uniform in particle size and uniformly dispersed in water

X = Non-uniform in particle size and not uniformly dispersed in water.

(5) Rinsing performance and organoleptic evaluation

Five hundreds ml of 1:50 dilutions of the respective hair rinses and a commercially available hair rinse were each used to treat hair therewith, followed by rinsing twice with warm water and air drying. The hair was organoleptically evaluated with respect to the softness, smoothness and combing ease. In comparison with the commercially available hair rinse, the hair rinses of the invention are evaluated as follows.

⊕ = excellent,

O = better

Δ = equal, and

X = poorer.

Example 1

Compositions of the formulations indicated in Table 1 were prepared and used to determine their appearance, stability, and rinsing performance.

Preparation of Composition:

To water (1) heated at 70°C were added ingredients (A) and (B) heated to and dissolved at the same temperature as indicated above, followed by agitating for emulsification and cooling to room temperature while agitating. After the cooling, water (2) was added to the mixture while agitating. After the cooling, water (2) was added to the mixture while agitating, thereby obtaining the respective compositions.

Results: as shown in Table 1.

Table 1

Composition (%)	<u>Product of Invention</u>			<u>Comparative Product</u>		
	1	2	3	1	2	3
Ingredient (A) Stearyltrimethylammonium chloride	0.5	0.2	0.05	0.8	0.5	0.05
Ingredient (B) Behenyl alcohol	2.1	2.5	0.15	1.0	1.0	1.0
Water (1)	5.73	7.3	4.80	3.2	8.5	3.95
Ratio of Ingredient (B) and Ingredient (A) [(B)/(A)]	4.20	12.5	3.0	1.25	2.0	20.0
Emulsified state	0	0	0	0	0	x
Water (2) (water for dilution)	91.67	90.0	95.0	95.0	90.0	95.0
Appearance	0	0	0	0	0	x
Stability High temperature stability	0	0	0	0	0	x
Low temperature stability	0	0	0	0	0	x
Rinsing performance	⊙	⊙	△	x	x	x

Example 2

To water (1) heated at 70°C were added stearyltrimethylammonium chloride (ingredient A), behenyl alcohol (ingredient B), and propylene glycol, which were heated to and dissolved at the same temperature as indicated above, followed by agitating for emulsification and cooling to room temperature under agitation. The products of the invention were obtained by further adding water (2) to the mixture under agitation, thereby giving dilutions. The results are shown in Table 2.

5

10

15

20

25

30

35

40

45

50

55

60

65

Table 2

Composition (%)	Comparative Products								Products of Invention							
	4	5	6	7	8	4	5	6	7	8	9					
Stearyltrimethylammonium chloride	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2					
Behenyl alcohol	0.2	1.0	2.0	3.0	0.2	1.0	2.0	3.0	1.0	2.0	3.0					
Propylene glycol	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5					
Water (1)	99.1	98.4	97.3	96.3	32.8	31.6	30.6	29.6	8.3	7.3	6.3					
Water (2)	0	0	0	0	66.3	66.7	66.7	66.7	90	90	90					
Appearance	X	X	X	X	X	0	0	0	0	0	0					
High temperature stability	X	X	X	X	X	0	0	0	0	0	0					
Low temperature stability	X	X	X	X	X	0	0	0	0	0	0					

Example 3

To water (1) heated at 70°C (with comparative products, a water-soluble polymer was dispersed in water) were added ingredients (A) and (B) and propylene glycol heated to and dissolved at the same temperature as indicated above, followed by agitating for emulsification and cooling to room temperature under agitation. With the products of the invention, water (2) in which a water-soluble polymer was dispersed was added for dilution to the cooled emulsion, thereby obtaining compositions of the invention. The results are as shown in Table 3.

Table 3

Composition (%)	Comparative Products			Products of Invention			
	9	10	11	10	11	12	13
Ingredient (A)							
Stearyltrimethylammonium chloride	0.2	0.2		0.2	0.2		0.2
Distearyldimethylammonium chloride			0.2			0.2	0.2
Ingredient (B)							
Cetostearyl alcohol	2.0			2.0			2.0
Cetyl alcohol		2.0			2.0		
Stearic acid monoglyceride			2.0			2.0	
Water-soluble Polymer							
Hydroxyethyl cellulose				0.4	0.4	0.4	0.4
Hydroxymethyl cellulose	0.4	0.4	0.4				
Propylene glycol	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Water (1)	96.4	96.4	96.4	16.8	16.8	16.8	15.4
Water (2) (water for dilution)	0	0	0	79.6	79.6	79.6	80.8
Appearance	x	x	x	0	0	0	0
Stability							
High temperature stability	x	x	x	0	0	0	0
Low temperature stability	x	x	x	0	0	0	0

Example 4

4. Rinse compositions were prepared in the same manner as in Example 3. The results are shown in Table

5

10

15

20

25

30

35

40

45

50

55

60

65

Table 4

Ingredient A	Compara- tive Product 12	Inven- tion Product 14	Inven- tion Product 15	Compara- tive Product 13	Inven- tion Product 16	Inven- tion Product 17	Compara- tive Product 14
Stearyltrimethylammonium Chloride	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Ingredient (B)							
Cetostearyl Alcohol	2.0	2.0	2.0				
Cetyl Alcohol	-	-	-	2.0	2.0	-	-
Behenyl alcohol						2.0	2.0
Water-soluble Polymer							
Hydroxyethyl Cellulose				0.2	0.2	0.2	0.2
Hydroxymethyl Cellulose	0.2	0.2	0.2				
Propylene Glycol	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Water (1)	97	30.3	17.0	97	17.0	17.0	3.6
Water (2) (water for dilution)	0	66.7	80.0	0	80.0	80.0	93.4
Appearance	x	0	0	x	0	0	x
Stability							
High Temperature Stability	0	0	0	0	0	0	0
Low Temperature Stability	0	0	0	0	0	0	0
Organoleptic Evaluation							
Softness	x	0	⊙	x	0	⊙	x
Smoothness	△	⊙	⊙	△	⊙	⊙	△
Ease in Combing	x	⊙	⊙	x	⊙	⊙	x

Example 5

Hair rinse compositions of the following formulations were prepared and 500 ml of a 1:50 dilution of each composition was used to treat hair therewith, followed by rinsing twice with warm water and air drying. The compositions were evaluated by 20 panelists using the Sheffe's paired comparison by five ranks. The results are shown in Table 5.

Product No. 1 of Invention:

(1) Monostearyltrimethylammonium chloride	0.2(%)
(2) Behenyl alcohol	1.0
(3) Propylene glycol	0.5
(4) Water (1)	8.3
(5) Water (2)	90.0

To (4) heated to 70°C was added a mixture of (1) to (3) heated to and dissolved at the same temperature as indicated above, followed by agitating for emulsification and cooling to room temperature while agitating. After the cooling, (5) was added to the mixture at a normal temperature under agitation to obtain a composition.

Product No. 2 of Invention:

(1) Monostearyltrimethylammonium chloride	0.2(%)
(2) Behenyl alcohol	1.0
(3) Propylene glycol	0.5
(4) Water (1)	18.3
(5) Water (2)	80.0

To (4) heated to 70°C was added a mixture of (1) to (3) heated to and dissolved at the same temperature as indicated above, followed by agitating for emulsification and adding cooled (5) while agitating, thereby obtaining a composition.

Comparative Product No. 15:

(1) Monostearyltrimethylammonium chloride	0.2(%)
(2) Behenyl alcohol	1.0
(3) Propylene glycol	0.5
(4) Water	98.3

To (4) heated to 70°C was added a mixture of (1) to (3) heated to and dissolved at the same temperature as indicated above, followed by cooling to room temperature while agitating, thereby obtaining a composition.

Comparative Product No. 16:

(1) Monostearyltrimethylammonium chloride	0.2(%)
(2) Behenyl alcohol	1.0
(3) Propylene glycol	0.5
(4) Polyoxyethylene hardened castor oil 60EO	0.5
(5) Water	97.8

To a mixture of (1), (2), (3) and (4) which was heated to and dissolved at 70°C was gradually added (5), which was heated to the same temperature level as indicated above, followed by emulsification and cooling to room temperature with agitation, thereby obtaining a composition.

Table 5

Item	Composition	Good	Fair	Almost the Same	Fair	Good	Composition
Softness Smoothness Ease in Combing	Inventive Product 18	6	7	5	2	0	Comparative Product 15
		6	9	3	2	0	
		7	9	3	1	0	
Softness Smoothness Ease in Combing	Inventive Product 19	2	11	5	2	0	Comparative Product 15
		2	11	6	1	0	
		3	12	6	2	0	
Softness Smoothness Ease in Combing	Inventive Product 18	7	8	3	2	0	Comparative Product 16
		7	10	2	1	0	
		7	9	4	0	0	




Claims

1. A hair rinse composition which comprises

(A) 0.05 to 0.5 wt.-% of cationic surface active agent, and

(B) a higher alcohol having a linear or branched alkyl or alkenyl group having from 12 to 26 carbon atoms, or a glycerine mono fatty acid ester having a melting point not lower than 45°C, and (B) is contained in an amount of 3 to 15 times by weight to ingredient (A) and said composition is produced by diluting a highly concentrated emulsified product containing 0.4 to 34 wt.-% of ingredient (A) and (B) in total with 1 to 20 times by weight to the emulsified product of water.

2. The hair rinse composition according to Claim 1 or 2, wherein said cationic active agent is one or more of compounds represented by the general formula (I)



in which one or two of R_1 , R_2 , R_3 and R_4 independently represent a linear or branched alkyl or hydroxyalkyl group having from 8 to 22 carbon atoms, the rest independently represent an alkyl group having from 1 to 3 carbon atoms, hydroxyalkyl group, benzyl group, or polyoxyethylene group in which the total number of ethylene oxide added is not larger than 10 moles per mole of the amine, and X represents a halogen atom or alkylsulfate group having 1 to 2 carbon atoms.

3. The hair rinse composition according to Claim 1 or 2, wherein said higher alcohol has a linear or branched alkyl or alkenyl group having from 12 to 26 carbon atoms.

4. The hair rinse composition according to Claim 1 or 2, wherein said glycerine monofatty acid ester is represented by the following general formula (II)



(II)

in which R_s represents a saturated or unsaturated linear or branched hydrocarbon group having from 11 to 23 carbon atoms.

5. A hair rinse composition comprising (A) 0.05 to 0.5 wt% of a cationic active agent, (B) a higher alcohol, or a glycerine monofatty acid ester having a melting point not lower than 45°C and contained in an amount of 3 to 15 times by weight as large as ingredient (A), and (C) 0.1 to 5.0 wt% of a water-soluble polymer.

6. A method for preparing a hair rinse composition comprising

(A) 0.05 to 0.5 wt% of cationic surface active agent, and

(B) a higher alcohol or a glycerine monofatty acid ester having a melting point not lower than 45°C and contained in an amount of 3 to 15 times by weight of the ingredient (A), characterized in that a highly concentrated emulsified product containing 0.4 to 34 wt% of combined ingredients (A) and (B) is produced and subsequently diluted with water in an amount of from 1 to 20 times by weight of the emulsified product.

Patentansprüche

1. Haarspülmittelezusammensetzung, umfassend


(A) 0,05 bis 0,5 Gew.-% kationisches, oberflächenaktives Mittel und

(B) einen höheren Alkohol mit einer linearen oder verzweigten Alkyl- oder Alkenylgruppe mit 12 bis 26 Kohlenstoffatomen oder einen Glycerinmonofettsäureester mit einem Schmelzpunkt nicht niedriger als 45°C, wobei (B) in einer Menge des 3- bis 15-fachen nach Gewicht bezüglich Bestandteil (A) enthalten ist und die Zusammensetzung hergestellt wird, indem man ein hochkonzentriertes emulgiertes Produkt, enthaltend 0,4 bis 34 Gew.-% von Bestandteil (A) und (B) zusammen, verdünnt mit dem 1- bis 20-fachen nach Gewicht, bezogen auf das emulgierte Produkt, an Wasser.

2. Haarspülmittelezusammensetzung gemäß Anspruch 1, wobei das kationische aktive Mittel eine oder mehrere von Verbindungen der allgemeinen Formel (I) ist



EP 0 134 998 B1



in der ein oder zwei von R_1 , R_2 , R_3 und R_4 unabhängig eine lineare oder verzweigte Alkyl- oder Hydroxyalkylgruppe mit 8 bis 22 Kohlenstoffatomen bedeuten, die restlichen unabhängig eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen, Hydroxyalkylgruppe, Benzylgruppe oder Polyoxyethylengruppe, in der die Gesamtzahl des addierten Ethylenoxyds nicht größer als 10 Mol pro Mol des Ammins ist, bedeuten und X ein Halogenatom oder eine Alkylsulfatgruppe mit 1 bis 2 Kohlenstoffatomen bedeutet.

3. Haarspülmittelzusammensetzung gemäß Anspruch 1 oder 2, wobei der höhere Alkohol eine lineare oder verzweigte Alkyl- oder Alkenylgruppe mit 12 bis 26 Kohlenstoffatomen aufweist.

4. Haarspülmittelzusammensetzung gemäß Anspruch 1 oder 2, wobei der Glycerinmonofettsäureester durch die folgende Formel (II) dargestellt ist



in der R_s eine gesättigte oder ungesättigte, lineare oder verzweigte Kohlenwasserstoffgruppe mit 11 bis 23 Kohlenstoffatomen bedeutet.

5. Haarspülmittelzusammensetzung umfassend (A) 0,05 bis 0,5 Gew.-% eines kationisch aktiven Mittels, (B) einen höheren Alkohol oder einen Glycerinmonofettsäureester mit einem Schmelzpunkt nicht niedriger als 45°C und in einer Menge vom 3- bis 15-fachen nach Gewicht, bezogen auf den Bestandteil (A), enthalten und (C) 0,1 bis 5,0 Gew.-% eines wasserlöslichen Polymeren.

6. Verfahren zur Herstellung einer Haarspülmittelzusammensetzung, umfassend (A) 0,05 bis 0,5 Gew.-% kationisch oberflächenaktives Mittel und (B) einen höheren Alkohol oder einen Glycerinmonofettsäureester mit einem Schmelzpunkt nicht niedriger als 45°C und in einer Menge vom 3- bis 15-fachen nach Gewicht des Bestandteils (A) enthalten, dadurch gekennzeichnet, daß man ein hochkonzentriertes emulgiertes Produkt, enthaltend 0,4 bis 34 Gew.-% der kombinierten Bestandteile (A) und (B), herstellt und anschließend mit Wasser verdünnt in einer Menge vom 1- bis 20-fachen nach Gewicht des emulgierten Produkts.

Revendications


1. Composition de rinçage capillaire qui comprend:

(A) 0,05 à 0,5% en poids d'agent tensio-actif cationique; et

(B) un alcool supérieur ayant un groupe alkyle ou alcényle, linéaire ou ramifié, ayant de 12 à 26 atomes de carbone, ou un ester de monoacide gras et de glycérol ayant un point de fusion non inférieur à 45°C.

et (B) est présent en une quantité de 3 à 15 fois en poids celle de l'ingrédient (A) et ladite composition est obtenue par dilution d'un produit émulsifié très concentré contenant 0,4 à 34% en poids d'ingrédient (A) et (B) au total avec une quantité d'eau représentant 1 à 20 fois en poids la quantité dudit produit émulsifié.

2. Composition de rinçage capillaire selon la revendication 1, dans laquelle ledit agent tension-actif cationique est constitué par un ou plusieurs des composés représentés par la formule générale (I):



dans laquelle:

un ou deux parmi R_1 , R_2 , R_3 et R_4 représentent indépendamment un groupe alkyle ou hydroxyalkyle, linéaire ou ramifié, ayant de 8 à 22 atomes de carbone,

les autres représentant indépendamment un groupe alkyle ayant de 1 à 3 atomes de carbone, un groupe hydroxyalkyle, un groupe benzyle, ou un groupe polyoxyéthylène dans lequel le nombre total de moles d'oxyde d'éthylène ajoutées n'est pas supérieur à 10 moles par mole de l'amine; et


X représente un atome d'halogène ou un groupe alkylsulfate ayant 1 ou 2 atomes de carbone.



EP 0 134 998 B1

3. Composition de rinçage capillaire selon la revendication 1 ou 2, dans laquelle ledit alcool supérieur a un groupe alkyle ou alcényle, linéaire ou ramifié, ayant 12 à 26 atomes de carbone.

4. Composition de rinçage capillaire selon la revendication 1 ou 2, dans laquelle ledit ester de monoacide gras et de glycérol est représenté par la formule générale (II):



(II)



dans laquelle R_5 représente un groupe hydrocarboné, linéaire ou ramifié, saturé ou insaturé, ayant de 11 à 23 atomes de carbone.

5. Composition de rinçage capillaire comprenant:

(A) 0,05 à 0,5% en poids d'agent tensio-actif cationique;

(B) un alcool supérieur, ou un ester de monoacide gras et de glycérol ayant un point de fusion non inférieur à 45°C, et présent en une quantité de 3 à 15 fois en poids celle de l'ingrédient (A); et

(C) 0,1 à 5,0% en poids d'un polymère soluble dans l'eau.

6. Procédé de préparation d'une composition de rinçage capillaire comprenant:

(A) 0,5 à 0,5% en poids d'agent tensio-actif cationique; et

(B) un alcool supérieur ou un ester de monoacide gras et de glycérol, ayant un point de fusion non inférieur à 45°C et contenu en une quantité de 3 à 15 fois en poids celle de l'ingrédient (A), caractérisé par le fait qu'un produit émulsifié très concentré contenant 0,4 à 34% en poids d'ingrédients (A) et (B) combinés est obtenu et dilué par la suite avec de l'eau en une quantité allant de 1 à 20 fois en poids celle du produit émulsifié.