

"STYRYL-SUBSTITUTED HETEROARYL COMPOUNDS WHICH INHIBIT EGF RECEPTOR
TYROSINE KINASE"

✓ 1 35
✓ 3 40

FIELD OF THE INVENTION

This invention relates to the inhibition of cell proliferation. More specifically, this invention relates to the use of styryl-substituted monocyclic and bicyclic heteroaryl compounds in inhibiting cell proliferation, including compounds which are useful protein tyrosine kinase (PTK) inhibitors.

Normal cellular reproduction is believed to be triggered by the exposure of the cellular substrate to one or more growth factors, examples of which are insulin, epidermal growth factor (EGF) and platelet-derived growth factor (PDGF). Such growth factors are typically specific for corresponding growth factor receptors which are imbedded in and which penetrate through the cellular membrane. The initiation of cellular reproduction is believed to occur when a growth factor binds to the corresponding receptor on the external surface of the cellular membrane. This

This Sheet Has Been Replaced



SUBSTITUTE SHEET

growth factor-receptor binding alters the chemical characteristics of that portion of the receptor which exists within the cell and which functions as an enzyme to catalyze phosphorylation of either an intracellular
5 substrate or the receptor itself, the latter being referred to as autophosphorylation. Examples of such phosphorylation enzymes include tyrosine kinases, which catalyze phosphorylation of tyrosine amino acid residues of substrate proteins.


10 Many diseased states are characterized by the uncontrolled reproduction of cells. These diseased states involve a variety of cell types and include disorders such as leukemia, cancer, psoriasis, atherosclerosis and restenosis injuries. The
15 inhibition of tyrosine kinase is believed to have utility in the control of uncontrolled cellular reproduction, i.e., cellular proliferative disorders.

Initiation of autophosphorylation, i.e., phosphorylation of the growth factor receptor itself,
20 and of the phosphorylation of a host of intracellular substrates are some of the biochemical events which are involved in mitogenesis and cell proliferation. Autophosphorylation of the insulin receptor and phosphorylation of substrate proteins by other
25 receptors are the earliest identifiable biochemical hormonal responses.

Elimination of the protein tyrosine kinase (PTK) activity of the insulin receptor and of the epidermal growth factor (EGF) receptor by site-directed
30 mutagenesis of the cellular genetic material which is responsible for generation of insulin and EGF results

SUBSTITUTE SHEET

This Sheet Has Been Replaced



in the complete elimination of the receptors' biological activity. This is not particularly desirable because insulin is needed by the body to perform other biological functions which are not related to cell proliferation. Accordingly, compounds which inhibit the PTK portion of the EGF receptor at concentrations less than the concentrations needed to inhibit the PTK portion of the insulin receptor could provide valuable agents for selective treatment of cell proliferation disorders.

REPORTED DEVELOPMENTS

U.S. Patent Nos. 4,678,793 and 4,826,984 disclose pharmaceutical compositions including styryl 4,4-dimethyl (bicyclic heteroaryl) compounds as active agents for treating cancer, psoriasis, acne, etc. U.S. Patent No. 4,769,384 discloses pharmaceutical compositions including styryl benzimidazole compounds as active agents for treating ulcers of the stomach and duodenum.

It has been reported that the most potent inhibitors of EGF receptors inhibit EGF-induced proliferation of A431/clone 15 cells with little or no effect on the proliferation of such cells when induced by other growth factors. It has been reported also that erbstatin inhibits the autophosphorylation of the EGF receptor in membranes of A431 cells. Low concentrations of erbstatin are required to inhibit EGF receptor autophosphorylation, whereas much higher concentrations of erbstatin are required to inhibit cyclic adenosine 3',5'-monophosphate (cAMP)-dependent protein kinase.

This Sheet Has Been Replaced



SUMMARY OF THE INVENTION


In accordance with the present invention, there is provided a method of inhibiting cell proliferation in a patient suffering from such disorder comprising the administration to the patient of a styryl-substituted heteroaryl compound exhibiting protein tyrosine kinase inhibition activity wherein the heteroaryl group is a monocyclic ring with 1 or 2 heteroatoms, or a bicyclic ring with 1 to about 4 heteroatoms, said compound optionally substituted or polysubstituted, with the proviso that when said ring is polysubstituted, the substituents do not have a common point of attachment to said ring.

Another aspect of the present invention relates to novel compounds which are those compounds of the aforementioned type wherein no substituent on the heteroaryl group is a carboxy group or an ester group.


Still another aspect of the present invention relates to pharmaceutical compositions comprising, in admixture with a pharmaceutically acceptable carrier, a pharmaceutically-effective amount of a novel compound of the aforementioned type.

With respect to the method aspects of this invention, the compounds described by Formula I below constitute a class of the aforementioned styryl-substituted heteroaryl compounds for use in the practice of the present invention:

This Sheet Has Been Replaced



5



I

wherein:

R₁ is alkyl, -H, -CN, -OH, -COOR, -CONRR or -CSNRR;

R is alkyl, -H or aralkyl;


5 R₂ is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms or 1 or 2 N-oxide groups, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms or 1 to about 4 N-oxide groups, said ring optionally substituted with one to about three R₉ groups, said R₉ substituents having no common points of attachment to said ring;

15 R₃ is alkyl, -H, -CN, -OH, -COOR, -CONRR, -CSNRR or -CH₂CN;

R₄, R₅, R₆, R₇ and R₈ are each independently alkyl, -H, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic or heteroaryl;


20 each R₉ is independently alkyl, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic or heteroaryl

This Sheet Has Been Replaced



SUBSTITUTE SHEET

6



; and

R_3 and R_7 together may be $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$ or, starting from R_3 , $-\text{CONH}-$; or

a pharmaceutically acceptable salt thereof.

5 Also in accordance with the present invention, novel compounds within the scope of the compound and pharmaceutical composition aspects of the present invention are described by Formula I above wherein:

R_1 is alkyl, $-\text{CN}$, $-\text{COOR}$, $-\text{CONRR}$ or $-\text{CSNRR}$;

10 R is alkyl, $-\text{H}$ or aralkyl;


R_2 is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common points of attachment;

15

R_3 is alkyl, $-\text{H}$, $-\text{COOR}$, $-\text{CONRR}$, $-\text{CSNRR}$ or $-\text{CH}_2\text{CN}$;

20 R_4 and R_6 are each independently alkyl, $-\text{H}$, $-\text{CN}$, halo, $-\text{OR}$, $-\text{CHO}$, $-\text{COOH}$, $-\text{NRR}$, $-\text{NO}_2$, $-\text{NHCOCH}_3$, $-\text{SR}$, $-\text{CF}_3$,

This Sheet Has Been Replaced



INSTITUTE SHEET


7

-CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic or heteroaryl;

R₅, R₇ and R₈ are each independently alkyl, -H, -CN, -OR, -CHO, -COOH, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic or heteroaryl;

with the provisos that at least two of R₄, R₅, R₆, R₇ and R₈ are not -H, and R₄, R₅ or R₆ cannot be -OR when R₇ or R₈ is -OR; and

each R₉ is independently alkyl, -CN, halo, -OR, -CHO, -NRR, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic heteroaryl or



: and

R₃ and R₇ together may be -CH₂CH₂-, -CH₂CH₂CH₂- or, starting from R₃, -CONH-.


In an alternate embodiment of the present invention, novel compounds within the scope of the compound and pharmaceutical composition aspects of the present invention are described by Formula II below.

This Sheet Has Been Replaced



SUBSTITUTE SHEET

8



II

wherein:

R is alkyl, -H or aralkyl;

5 R_2 is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common points of attachment;

10

R_4 and R_6 are each independently branched alkyl of about 3 to about 6 carbon atoms, -CN, halo, -SR, -CF₃, heterocyclic, or heteroaryl;

15 R_5 , R_7 and R_8 are each independently alkyl, -H, -CN, -OR, -CHO, -COOH, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic or heteroaryl;


each R_9 is independently alkyl, -CN, halo, -OR, -CHO, -NRR, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, heteroaryl or

20

SUBSTITUTE SHEET

This Sheet Has Been Replaced





Compounds within the scope of the present invention have also a specific affinity toward the substrate site of the tyrosine kinase domain of EGF receptors, inhibit EGF receptor kinase more than they inhibit PDGF receptor kinase and also effectively inhibit EGF-dependent autophosphorylation of the receptor.

DETAILED DESCRIPTION OF THE INVENTION

As employed above and throughout the disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

"Alkyl" means a saturated aliphatic hydrocarbon which may be either straight- or branch-chained containing from about 1 to about 6 carbon atoms.

"Lower alkyl" means an alkyl group as above, having 1 to about 4 carbon atoms which may be straight- or branch-chained such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl.

"Alkoxy" means an alkyl-oxy group in which "alkyl" is as previously described. Lower alkoxy groups are preferred. Exemplary groups include methoxy, ethoxy, n-propoxy, i-propoxy and n-butoxy.

This Sheet Has Been Replaced



SUBSTITUTE SHEET

"Aryl" means an unsaturated or partially unsaturated ring system. Preferred aryl groups are pyridyl and indolyl.

5 "Heterocyclic" means an unsaturated ring system containing 1 or 2 N, O or S heteroatoms. Preferred heterocyclic groups are morpholine, piperidine and piperazine.

10 "Acyl" means an organic radical derived from an organic acid, a carboxylic acid, by the removal of its acid hydroxyl group. Preferred acyl groups are lower alkyl carboxylic acid groups such as acetyl and propionyl. Benzoyl is also preferred.

"Halo" means a halogen. Preferred halogens include chloride, bromide and fluoride.

15 Preferred aralkyl groups are benzyl and phenethyl.

It is believed that therapeutically useful PTK inhibiting compounds should be competitive with the substrate of EGF receptor tyrosine kinase (EGFRK) and not with adenosine triphosphate (ATP). The PTK
20 inhibitors quercetin and genistein, which compete with ATP, inhibit other protein kinases and as a result are highly cytotoxic. As a test of selectivity, compounds which inhibit EGFRK better than they inhibit insulin receptor kinase (IRK) and/or PDGF receptor kinase are
25 of considerable value.


It is theorized that solubility of the compounds of the present invention both in water and in mildly hydrophobic solvents will enhance the probability that



11

they traverse the cell membrane. Various insoluble compounds, however, have exhibited significant EGFRK inhibition in in vitro testing.

5 In accordance with the method aspects of the present invention, a preferred class of styryl-substituted heteroaryl compounds for use in the practice of the present invention are described by Formula II below.



II

10 wherein:

R is alkyl, -H or aralkyl;

15 R₂ is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms or 1 or 2 N-oxide groups, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms or 1 to about 4 N-oxide groups, said ring optionally substituted with one to about three R₉ groups, said R₉ substituents having
20 no common points of attachment to said ring;

R₄, R₅, R₆, R₇ and R₈ are each independently alkyl, -H,

This Sheet Has Been Replaced




SUBSTITUTE SHEET

12

-CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, or heteroaryl;

each R₂ is independently alkyl, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, heteroaryl or



A preferred class of compounds useful in the practice of the present invention include those described by Formula I where:

R₁ is -CN, -COOR, -CONRR or -CSNRR;

R is lower alkyl, -H or aralkyl;

R₂ is a 6-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or a 9- or 10-membered bicyclic aryl ring including 1-4 N, O or S atoms, said ring optionally substituted with one to about three R₃ groups, said R₃ substituents having no common points of attachment to said ring;

R₃ is -H;

This Sheet Has Been Replaced


SUBSTITUTE SHEET

13

R_4 , R_5 , R_6 , R_7 and R_8 are each independently lower alkyl, -H, lower alkoxy or -OH, with the provisos that at least two of R_4 , R_5 , R_6 , R_7 and R_8 are not -H, and R_4 , R_5 or R_6 cannot be lower alkoxy when R_7 or R_8 is lower alkoxy;


R_4 and R_6 are also each independently halo;

each R_9 is independently lower alkyl, halo, lower alkoxy, -OH or



a pharmaceutically acceptable salt thereof.


More preferred compounds for use in the practice of this invention include those of Formulae III and IV below:



III

This Sheet Has Been Replaced

14



IV

where

R_1 is -CN, -COOR, -CONRR or -CSNRR;

R is lower alkyl, -H or aralkyl;

5 R_2 is a 6-membered monocyclic aryl ring including
1 or 2 N, O or S atoms, or a 9- or 10-
membered bicyclic aryl ring including 1-4 N,
O or S atoms, said ring optionally
substituted with one to about three R_9
10 groups, said R_9 substituents having no common
points of attachment to said ring;


R_3 is -H;

R_4 , R_5 , R_6 , R_7 and R_8 are each independently lower alkyl,
-H, lower alkoxy or -OH, with the provisos that at
15 least two of R_4 , R_5 , R_6 , R_7 and R_8 are not -H, and
 R_4 , R_5 or R_6 cannot be lower alkoxy when R_7 or R_8 is
lower alkoxy;

R_4 and R_6 are also each independently halo;


each R_9 is independently lower alkyl, halo, lower
20 alkoxy, -OH or

This Sheet Has Been Replaced




REPLACEMENT SHEET

15



R_{10} is -H, alkyl, aralkyl or



a pharmaceutically acceptable salt thereof.

5 Even more preferred compounds are described by Formulae III and IV where:

R_1 is -CN, -COOR or -CONRR;

10 R_4 , R_5 , R_6 , R_7 and R_8 are independently lower alkyl, -H, lower alkoxy or -OH, with the provisos that at least two of R_4 , R_5 , R_6 , R_7 and R_8 are not -H, and R_4 , R_5 or R_6 cannot be lower alkoxy when R_7 or R_8 is lower alkoxy;

R_4 and R_6 are also each independently halo;

This Sheet Has Been Replaced

16

R is lower alkyl or -H; and

there are no R_9 substituents.

The most preferred compounds are described by Formula II where R_1 is -CN, R_3 , R_7 and R_8 are each
5 independently -H, and R_4 and R_6 are each independently alkyl, halo, -OR or -CF₃.


In an alternate embodiment of the present invention, a preferred class of compounds are described by Formula II above wherein

10 R_2 is a 6-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or a 9- or 10-membered bicyclic aryl ring including 1-4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common
15 points of attachment to said ring;

R_4 and R_6 are each independently branched lower alkyl of about 3 to about 4 carbon atoms or halo;

R_5 , R_7 and R_8 are each independently -H lower alkyl of about 2 to about 4 carbon atoms or halo; and

20 each R_9 is independently lower alkyl, halo, lower alkoxy, -OH or




This Sheet Has Been Replaced



SUBSTITUTE SHEET

In an alternate embodiment of the present invention, more preferred compounds for use in the practice of the present invention are described by Formula V below.



where R_4 , R_5 , R_6 , R_7 , R_8 and R_9 are as described immediately above.

In an alternate embodiment of the present invention, even more preferred compounds are described by Formula V where R_4 and R_6 are each independently halo, isopropyl, isobutyl, sec-butyl or t-butyl; R_5 , R_7 and R_8 are each independently -H, halo, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl or t-butyl; and there are no R_9 substituents.

The most preferred compounds in an alternate embodiment of the present invention are described by Formula V where R_4 and R_6 are each independently halo, and R_5 , R_7 and R_8 are each independently -H.

This Sheet Has Been Replaced

Compounds of this invention may be useful in the form of the free base, in the form of salts and as a hydrate. All forms are within the scope of the invention. Acid addition salts may be formed and are simply a more convenient form for use; and in practice, use of the salt form inherently amounts to use of the base form. The acids which can be used to prepare the acid addition salts include preferably those which produce, when combined with the free base, pharmaceutically acceptable salts, that is, salts whose anions are non-toxic to the animal organism in pharmaceutical doses of the salts, so that the beneficial properties inherent in the free base are not vitiated by side effects ascribable to the anions. Although pharmaceutically acceptable salts of said basic compound are preferred, all acid addition salts are useful as sources of the free base form even if the particular salt per se is desired only as an intermediate product as, for example, when the salt is formed only for purposes of purification and identification, or when it is used as an intermediate in preparing a pharmaceutically acceptable salt by ion exchange procedures.

Pharmaceutically acceptable salts within the scope of the invention include those derived from the following acids: mineral acids such as hydrochloric acid, sulfuric acid, phosphoric acid and sulfamic acid; and organic acids such as acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, cyclohexylsulfamic acid, quinic acid, and the like.

This Sheet Has Been Replaced




The corresponding acid addition salts comprise the following: hydrochloride, sulfate, phosphate, sulfamate, acetate, citrate, lactate, tartarate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate, cyclohexylsulfamate and quinate, respectively.

The acid addition salts of the compounds of this invention are prepared either by dissolving the free base in aqueous or aqueous-alcohol solution or other suitable solvents containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and acid in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.


Compounds useful in the pharmaceutical composition and method aspects of this invention can be prepared by known methods, for example, Knoevenagel condensation reactions such as those disclosed in U.S. Patent No. 3,149,148.

Compounds of this invention may be prepared by the following reaction sequence:



Knoevenagel condensation of a substituted benzaldehyde in a polar media with an active methylene

This Sheet Has Been Replaced



compound of the formula $R_1CH_2R_2$ in the presence of ammonia or amines such as piperidine and raised heat results in the products of this invention. When substitution of the R_3 group is desired, the
5 corresponding ketone starting material is used. Reaction temperatures in the range of 25°C to reflux and reaction times vary depending on the materials being used in the condensation.

Compounds of this invention are either
10 commercially available, known in the literature or can be made by known procedures. For example, U.S. Patent No. 4,600,712 discloses fungicides of Formula I where, for example, R_1 is cyano, R_2 is pyridyl and R_3 and R_7 or R_8 are chloro. U.S. Patent Nos. 3,337,565 and
15 3,337,568 disclose compounds which interfere with carbohydrate metabolism of Formula I where, for example, R_1 is cyano or hydroxy, R_2 is pyridyl and R_3 is hydroxy. U.S. Patent No. 3,196,158 discloses adrenal cortex inhibitors of Formula I where, for example, R_1
20 is cyano, R_2 is pyridyl and R_7 or R_8 are halo. U.S. Patent No. 3,157,663 discloses adrenal cortex inhibitors where, for example, R_1 is cyano, R_2 is pyridyl and R_3 , R_7 or R_8 are amino or nitro groups. Buu-Hoi et al., Journal of the Chemical Society (C),
25 pp. 2069-70 (1969) disclose the conversion of 1,2-diarylacrylonitriles to the corresponding 3-arylcoumarins, wherein the 2-aryl group bears an ortho-alkoxy-substituent. Although the foregoing
30 publications disclose some compounds of the type that can be used in accordance with the method aspects of the present invention, they do not disclose the use of such compounds for inhibiting cell proliferation.

This Sheet Has Been Replaced



Various R, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ substituents on the phenyl and heterocyclic ring or chain can be present in the starting compound or added after formation of the condensation product by methods known in the art for substitution or conversion on one group to another. If the substituents themselves are reactive, then the substituents can themselves be protected according to the techniques known in the art. A variety of protecting groups known in the art, may be employed. Examples of many of these possible groups may be found in "Protective Groups in Organic Synthesis" by T. W. Green, John Wiley and Sons, 1981. For example, nitro groups can be added to the aromatic ring by nitration and the nitro group converted to other groups, such as amino by reduction, and halo by diazotization of the amino group and replacement of the diazo group. Acyl groups can be substituted onto the aryl groups by Friedel-Crafts acylation. The acyl groups can then be transformed to the corresponding alkyl groups by various methods, including the Wolff-Kishner reduction and Clemmenson reduction. Amino groups can be alkylated to form mono- and di-alkylamino groups; and mercapto and hydroxy groups can be alkylated to form corresponding ethers. Primary alcohols can be oxidized by oxidizing agents known in the art to form carboxylic acids or aldehydes, and secondary alcohols can be oxidized to form ketones. Tertiary amino groups can be converted to the corresponding N-oxides by oxidizing agents known in the art, for example, hydrogen peroxide and peracids. Thus, substitution or alteration reactions can be employed to provide a variety of substituents throughout the molecule of the starting material, intermediates, or the final product.

This Sheet Has Been Replaced



SUBSTITUTE SHEET

Compounds within the scope of this invention exhibit significant activity as protein tyrosine kinase inhibitors and possess therapeutic value as cellular antiproliferative agents for the treatment of certain conditions including psoriasis, atherosclerosis and restenosis injuries. It is expected that the invention will be particularly applicable to the treatment of atherosclerosis. With regard to the treatment of some conditions, for example, atherosclerosis, certain people may be identified as being at high risk, for example, due to genetic, environmental or historical factors. Compounds within the scope of the present invention can be used in preventing or delaying the occurrence or reoccurrence of such conditions or otherwise treating the condition.

The compounds of the present invention can be administered to a mammalian host in a variety of forms adapted to the chosen route of administration, i.e., orally, or parenterally. Parenteral administration in this respect includes administration by the following routes: intravenous, intramuscular, subcutaneous, intraocular, intrasynovial, transepithelial including transdermal, ophthalmic, sublingual and buccal; topically including ophthalmic, dermal, ocular, rectal and nasal inhalation via insufflation and aerosol and rectal systemic.

The active compound may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsules, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral

This Sheet Has Been Replaced



SUBSTITUTE SHEET

therapeutic administration, the active compound may be incorporated with excipient and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like.

5 Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 6% of the weight of the unit. The amount of active compound
10 in such therapeutically useful compositions is such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit form contains between about 1 and 1000 mg of active
15 compound.

The tablets, troches, pills, capsules and the like may also contain the following: A binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent
20 such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring. When the
25 dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or
30 capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring

This Sheet Has Been Replaced



SUBSTITUTE SHEET

such as cherry or orange flavor. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound
5 may be incorporated into sustained-release preparations and formulations.

The active compound may also be administered parenterally or intraperitoneally. Solutions of the active compound as a free base or pharmacologically
10 acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersion can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these
15 preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation
20 of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It may be stable under the conditions of manufacture and storage and must be preserved against the contaminating action
25 of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures
30 thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required

This Sheet Has Been Replaced


SUBSTITUTE SHEET

particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

The therapeutic compounds of this invention may be administered to a mammal alone or in combination with pharmaceutically acceptable carriers, as noted above, the proportion of which is determined by the solubility and chemical nature of the compound, chosen route of administration and standard pharmaceutical practice.

This Sheet Has Been Replaced



SUBSTITUTE SHEET

The dosage of the present therapeutic agents which will be most suitable for prophylaxis or treatment will vary with the form of administration, the particular compound chosen and the physiological characteristics of the particular patient under treatment. Generally, small dosages will be used initially and if necessary, will be increased by small increments until the optimum effect under the circumstances is reached. The therapeutic human dosage, based on physiological studies using rats, will generally be from about 0.01 mg to about 100 mg/kg of body weight per day or from about 0.4 mg to about 10 g or and higher although it may be administered in several different dosage units from once to several times a day. Oral administration requires higher dosages.

EXAMPLES


Embodiments of the present invention and comparative examples are described in the following non-limiting examples which include a description of pharmacological test procedures believed to correlate to therapeutic activity in humans and other animals. Examples 1 and 2 below are illustrative of compounds within the scope of the present invention. In examples 1A and 1B, R_1 is cyano, R_2 is pyridyl and R_4 and R_6 are chloro. In example 2, R_2 is pyridyl, R_3 is cyano and R_4 and R_6 are chloro.

Example 1A

trans-2-(3-Pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile

To a stirred solution of 400 g (2.3 mole) 3,5-dichlorobenzaldehyde in 11.4 liters absolute ethanol were added 283.5 g (2.4 mole) 3-

This Sheet Has Been Replaced



pyridylacetonitrile and 720.9 g of K_2CO_3 . Within 2 minutes of the K_2CO_3 addition, a solid precipitate formed. The reaction mixture was stirred about 2.5 hours. Water (22.9 liters) was added to the stirred
5 reaction mixture. After stirring for 1 hour, the mixture was filtered and the filter cake was washed with water (10 liters), dried and recrystallized from isopropanol. Five hundred thirty four g (85% yield) of
10 *trans*-2-(3-pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile were obtained, m.p. 150-151°C.

Example 1B

cis-2-(3-pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile


A stirred solution of *trans*-2-(3-pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile (20 g), for example,
15 as prepared in Example 1A, in 3 liters acetonitrile was irradiated with a sunlamp (275 W) for 24 hours. The reaction mixture was stirred an additional 24 hours and concentrated *in vacuo*. The resulting residue was
20 purified by flash chromatography on silica gel, eluting sequentially with hexanes, 5% EtOAc/hexanes, 10% EtOAc/hexanes, and then 15% EtOAc/hexanes. The resulting yellow solid was recrystallized from 15% EtOAc/heptane to give 8 g (74% yield) of the title
25 compound, m.p. 114-117°C.

Example 2

2-(3,5-Dichlorophenyl)-3-(3-pyridyl)-2-propenenitrile

To a stirred solution of 3 g (17.1 mmole) of 3,5-dichlorobenzaldehyde in 200 ml absolute ethanol
30 were added 1.83 ml (17.1 mmole) of 3-pyridyl-acetonitrile and 1 equivalent of K_2CO_3 (2.36 g). The reaction flask was equipped with a reflux condenser and

This Sheet Has Been Replaced



SUBSTITUTE SHEET

the stirred reaction mixture was refluxed for 2 hours, filtered and concentrated. The resulting residue was purified by flash chromatography on silica gel, eluting with 4:1 hexane/ethyl acetate to give the title
5 compound as a white solid, m.p. 139-141°C. An amount of trans-2-(3-pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile was also obtained from the reaction mixture.

Compounds of this invention are subjected to
10 various biological tests, the results of which correlate to useful cellular antiproliferative activity. These tests are useful in determining EGF receptor kinase, PDGF receptor kinase and insulin receptor kinase inhibition activities of the compounds
15 disclosed herein.

EGF-Receptor Purification

EGF-receptor purification is based on the procedure of Yarden and Schlessinger. A431 cells are grown in 80 cm² bottles to confluency (2 x 10⁷ cells per
20 bottle). The cells are washed twice with PBS and harvested with PBS containing 1.0 mmol EDTA (1 hour at 37°C), and centrifuged at 600g for 10 minutes. The cells are solubilized in 1 ml per 2 x 10⁷ cells of cold solubilization buffer (50 mmol Hepes buffer, pH 7.6, 1%
25 Triton X-100, 150 mmol NaCl, 5 mmol EGTA, 1 mmol PMSF, 50 µg/ml aprotinin, 25 mmol benzamidine, 5 µg/ml leupeptin, and 10 µg/ml soybean trypsin inhibitor) for 20 minutes at 4°C. After centrifugation at 100000g for 30 minutes, the supernatant is loaded onto a WGA-
30 agarose column (100 µl of packed resin per 2 x 10⁷ cells) and shaken for 2 hours at 4°C. The unabsorbed material is removed and the resin washed twice with HTN

This Sheet Has Been Replaced


SUBSTITUTE SHEET

buffer (50 mmol Hepes, pH 7.6, 0.1% Triton X-100, 150 mmol NaCl), twice with HTN buffer containing 1 M NaCl, and twice with HTNG buffer (50 mmol Hepes, pH 7.6, 0.1% Triton X-100, 150 mmol NaCl, and 10% glycerol). The
5 EGF receptor is eluted batchwise with HTNG buffer containing 0.5 M N-acetyl-D-glucosamine (200 μ l per 2×10^7 cells). The eluted material is stored in aliquots at -70°C and diluted before use with TMTNG
10 buffer (50 mmol Tris-Mes buffer, pH 7.6, 0.1% Triton X-100, 150 mmol NaCl, 10% glycerol).

EGFR Kinase Catalyzed Phosphorylation
of Poly(GAT) and its Inhibition

WGA-purified EGFR (0.25 $\mu\text{g}/\text{assay}$) is preactivated with EGF (0.85 μM) in 50 mmol Tris-Mes buffer, pH 7.6
15 for 20 minutes at 4°C . The assay is initiated by addition of a mixture which contains $\text{Mg}(\text{Ac})_2$ (60 mmol), $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ (125 μM , 2-5 $\mu\text{Ci}/\text{assay}$), poly(GAT) (0.0625 mg/ml, 0.125 mg/ml, 0.25 mg/ml), and six concentrations of inhibitor in duplicates. The temperature of the
20 assay is 22°C and the production of phosphorylated copolymer is found to be linear up to 20 minutes. The PTK inhibitors tested are solubilized in water or a mixture of ethanol and water such that the final concentration of ethanol does not exceed 4% in the
25 assay. Up to 4% ethanol in the assay has no effect on the EGFR kinase activity. The concentration of EGF in the assay is 300 nM in a final volume of 40 μl . After 5, 10 or 20 minutes, aliquots of 25 μl are applied onto Whatman 3-mm paper cuttings, which are then soaked in
30 cold 10% TCA containing 0.01 M sodium pyrophosphate. After being washed overnight at 4°C , the paper cuttings are dried and counted, measuring ^{32}P Cerenkov radiation. Concentration dependence on poly(GAT) was Michaelian

This Sheet Has Been Replaced



SUBSTITUTE SHEET

with a $K_m = 0.076 \pm 0.007$ mg/ml or 0.069 ± 0.007 mmol if calculated per Glu₆Ala₃Tyr(GAT) unit. The EGF response for the poly(GAT) phosphorylation is graphed. The K_m for ATP in the assay was found to 2.9 μ M.

5 Time Dependence of EGF-Receptor Autophosphorylation

WGA-purified EGF receptor from A431 cells (0.5 μ g/assay) is activated with EGF (800 nM) for 20 minutes at 4°C. The reaction is initiated by the addition of Mg(Ac)₂ (60 mmol), Tris-Mes buffer, pH 7.6 (50 mmol),
10 and [³²P]ATP (20 μ M, 5 μ Ci/assay). The reaction is conducted at either 4 or 15°C and terminated by addition of sodium dodecyl sulfate (SDS) sample buffer (10% glycerol, 50 mmol Tris, pH 6.8, 5% β -mercapto-ethanol, and 3% (SDS). The samples are run on a 8% SDS
15 polyacrylamide gel (SDS-PAGE) (prepared from 30% acrylamide and 0.8% bis-(acrylamide) and contained 0.375 M Tris, pH 8.8, 0.1% SDS, 0.05% TEMED, and 0.46% ammonium persulfate). The gel is dried and autoradiography performed with Agfa Curix RP2 X-ray
20 film. The relevant radioactive bands are cut and counted in the Cerenkov mode. The fast phase of autophosphorylation continues for another 10 minutes. The extent of phosphorylation completed in the first 10-s at 15°C comprises 1/3 of the total
25 autophosphorylation signal and probably reflects the phosphorylation of the first site on the receptor. The 10-s interval is therefore chosen for use in subsequent autophosphorylation experiments.

ATP and EGF Dependence of Autophosphorylation

30 WGA-purified EGF receptor from A431 cells (0.5 μ g/assay) is activated with EGF (0.85 μ M) for 20 minutes at 4°C. The assay is performed at 15°C and initiated

This Sheet Has Been Replaced



SUBSTITUTE SHEET

by addition of $\text{Mg}(\text{Ac})_2$ (60 mmol), Tris-Mes buffer, pH 7.6 (50 mmol), $[\text{}^{32}\text{P}]\text{ATP}$ (carrier free, 5 $\mu\text{Ci}/\text{assay}$), and increasing concentrations of nonradioactive ATP. The assay is terminated after 10-s by addition of SDS sample buffer. The samples are run on a 6% SDS polyacrylamide gel. The gel is dried and autoradiographed as described above. The relevant radioactive bands are cut and counted in the Cerenkov mode. the K_m for ATP determined in this fashion is found to be 7.2 μM . With use of the 10-s assay protocol, the EGF concentration dependence of EGFRK autophosphorylation is determined.

Inhibition of Copoly(Glu₄Tyr) Phosphorylation by Insulin-Receptor Kinase (InsRK)

Rat liver membranes are prepared from the livers of 6-week-old rats as described by Cuatrecasas. WGA-purified insulin receptor is prepared according to Zick et al. WGA-purified rat liver InsRK (1.25 μg) is preincubated with or without 330 nM insulin in 50 mmol Tris-Mes buffer, pH 7.6, for 30 minutes at 22°C. The assay is performed at 22°C and initiated by addition of a mixture which contains $\text{Mg}(\text{Ac})_2$ (60 mmol), NaVO_3 (40 μM), $[\gamma\text{}^{32}\text{P}]\text{ATP}$ (125 μM , 3-5 $\mu\text{Ci}/\text{assay}$), and poly(GT) [poly(Glu₄Tyr)] at three concentrations: whenever an inhibitor is tested, it is added at the proper concentration. The final concentration of insulin in the assay is 125 nM. The total volume of the assay is 40 μl . After 20 minutes, aliquots of 30 μl are applied on Whatman 3-mm paper and soaked in cold 10% TCA, containing 0.01 M sodium pyrophosphate. After being washed overnight, the papers are dried and counted, measuring Cerenkov radiation. The InsRK-catalyzed

This Sheet Has Been Replaced



phosphorylation of poly(GT) obeys Michaelis-Menten kinetics.

Inhibition of EGFR Autophosphorylation

A431 cells were grown to confluence on human
5 fibronectin coated tissue culture dishes. After
washing 2 times with ice-cold PBS, cells were lysed by
the addition of 500 μ l/dish of lysis buffer (50 mmol
Hepes, pH 7.5, 150 mmol NaCl, 1.5 mmol $MgCl_2$, 1 mmol
EGTA, 10% glycerol, 1% triton X-100, 1 mmol PMSF, 1
10 mg/ml aprotinin, 1 mg/ml leupeptin) and incubating 5
minutes at 4°C. After EGF stimulation (500 μ g/ml 10
minutes at 37°C) immunoprecipitation was performed with
anti EGF-R (Ab 108) and the autophosphorylation
reaction (50 μ l aliquots, 3 μ Ci [γ - 32 P]ATP) sample was
15 carried out in the presence of 2 or 10 μ M of compound,
for 2 minutes at 4°C. The reaction was stopped by
adding hot electrophoresis sample buffer. SDS-PAGE
analysis (7.5% els) was followed by autoradiography and
the reaction was quantitated by densitometry scanning
20 of the x-ray films. In order to test the compounds for
selective inhibition, the procedure is repeated using
PDGF stimulation in place of EGF stimulation. "IC₅₀,"
as used below, refers to the concentration of inhibitor
(μ M) at which the rate of autophosphorylation is
25 halved, compared with media containing no inhibitor.
Preferably, the IC₅₀ values for EGF will be lower than
for PDGF, indicating a high degree of EGF specificity
of the inhibitory compounds. The results of this test
are summarized in Table I below.

This Sheet Has Been Replaced




Table I

Example	IC ₅₀ (μM)	
	EGF	PDGF
2	0.1	20
3	0.5	6
5	0.4	10
6	20	>50
7	0.4	30

These results show that the compounds of the present invention inhibit EGF receptor kinase better than they inhibit PDGF receptor kinase.

Inhibition of Cell Proliferation as Measured by Inhibition of DNA Synthesis

Cells were seeded at 1×10^5 cells per well in 24-well Costar dishes pre-coated with human fibronectin (by incubating for 30 minutes at room temperature with 10 μg/0.5 ml/well). The cells were grown to confluence for 2 days. The medium was changed to DMEM containing 0.5 calf serum for 36-48 hours and the cells were then incubated with PDGF, EGF (Toyobo, New York, NY) (20 ng/ml) or serum (10% calf serum, FCS) and different concentrations of the inhibitory compounds. [³H]thymidine, (NEN, Boston, MA) was added 16-24 hours later at 0.5 μCi/ml for 2 hours. TCA precipitable material was quantitated by scintillation counting (C). Results of this assay are summarized in Table II below. "IC₅₀," as used below, refers to the concentration of inhibitor (nM) at which [³H]thymidine incorporation is halved, compared with media containing no inhibitor. As FCS contains a broad range of growth factors, the IC₅₀ values for PDGF should be lower than for FCS, indicating that the compounds do not act as general inhibitors.

This Sheet Has Been Replaced




TABLE II

<u>Example</u>	<u>IC₅₀ - PDGF</u>	<u>IC₅₀ - FCS</u>
2	30±17	100
3	14±10	27±5% @ 1 µM
5 4	9±2	500
6	17	15% @ 1 µM

These results indicate that the compounds of the invention do not inhibit a broad range of growth factor receptors.


10 Cell Culture

Cells termed HER 14 and K721A (=DK) were prepared by transfecting N1H3T3 cells (clone 2.2) (From C. Fryling, NCI, NIH), which lack endogenous EGF-receptors, with cDNA constructs of wild-type EGF-receptor or mutant EGF-receptor lacking tyrosine kinase activity (in which Lys 721 at the ATP-binding site was replaced by an Ala residue, respectively). All cells were grown in DMEM with 10% calf serum (Hyclone, Logan, Utah).

20 The results obtained by the above experimental methods evidence the useful protein tyrosine kinase inhibition properties of the compounds within the scope of the present invention.

Comparative Examples

25 The following comparative examples illustrate the improved cell proliferation inhibition of compounds of the present invention in comparison to compounds of the prior art. In comparative Example C-1, the preparation of a prior art compound, where R₁ is cyano, R₂ is
30 pyridyl and R₃ and R₄ are chloro is described. The inhibitory activities of the compounds of Examples 1A



and 2 are compared to the inhibitory activity of the compound prepared in comparative Example C-1.

Example C-1

2-(3-pyridyl)-3-(2,4-dichlorophenyl)-2-propenenitrile

5 To a stirred solution of 2,4-dichlorobenzaldehyde (5.0 g, 29 mmole) and 3-pyridylacetonitrile (3.05 ml, 29 mmole) in 150 ml absolute ethenol were added 3.95 g K_2CO_3 . After stirring for about 15 minutes, a white solid precipitated out of the reaction mixture. The
10 reaction mixture was stirred overnight and filtered. The filter cake was washed with absolute ethanol and dissolved in CH_2Cl_2 . The CH_2Cl_2 solution was filtered and concentrated in vacuo. The resulting white solid was recrystallized from 3:2 hexane/ethyl acetate to
15 give 6.0 g (76%) of the title compound.

The inhibition of EGFR autophosphorylation by the compounds of Examples 1A and 2 was compared to the prior art compound of Example C-1. The EGFR autophosphorylation inhibition was evaluated using the
20 same procedure as described hereinbefore. The results are reported in Table III below.

TABLE III


	<u>Example</u>	<u>IC₅₀ (μM)</u>
	1A	2
25	2	2-6
	C-1	≥50

These results demonstrate that the compounds of Examples 1A and 2 are more potent inhibitors of EGF receptor kinase than C-1.

This Sheet Has Been Replaced


CLAIMS

1. A method of inhibiting cell proliferation in a patient suffering from such disorder comprising the administration to the patient of a pharmaceutically effective amount of a pharmaceutical composition containing, in admixture with a pharmaceutically acceptable carrier, a compound, or a pharmaceutically acceptable salt thereof, of the formula



10 wherein:

R is alkyl, -H or aralkyl;

R₂ is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms or 1 or 2 N-oxide groups, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms or 1 to about 4 N-oxide groups, said ring optionally substituted with one to about three R₉ groups, said R₉ substituents having no common points of attachment to said ring;

R₄, R₅, R₆, R₇ and R₈ are each independently alkyl, -H,

This Sheet Has Been Replaced




SUBSTITUTE SHEET




37

-CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, or heteroaryl;

5 each R₉ is independently alkyl, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, heteroaryl or



2. A compound of the formula



10

wherein:

R is alkyl, -H or aralkyl;

15 R₂ is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or an about 8- to about 12-membered bicyclic

This Sheet Has Been Replaced



SUBSTITUTE SHEET




38

aryl ring including 1 to about 4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common points of attachment;

R_4 and R_6 are each independently branched alkyl of about 3 to about 6 carbon atoms, -CN, halo, -SR, -CF₃, heterocyclic, or heteroaryl;

R_5 , R_7 and R_8 are each independently alkyl, -H, -CN, -OR, -CHO, -COOH, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic or heteroaryl;

each R_9 is independently alkyl, -CN, halo, -OR, -CHO, -NRR, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, heteroaryl or



15

3. A pharmaceutical composition for inhibiting cell proliferation in a patient suffering from such disorder comprising, in admixture with a pharmaceutically acceptable carrier, a pharmaceutically effective amount of a compound according to Claim 2.

20

This Sheet Has Been Replaced

SUBSTITUTE SHEET




4. A compound according to Claim 2 wherein:

5 R_2 is a 6-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or a 9- or 10-membered bicyclic aryl ring including 1-4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common points of attachment to said ring;


R_4 and R_6 are each independently branched lower alkyl of about 3 to about 4 carbon atoms or halo;

10 R_5 , R_7 and R_8 are each independently -H lower alkyl of about 2 to about 4 carbon atoms or halo; and

each R_9 is independently lower alkyl, halo, lower alkoxy, -OH or



15 5. A compound according to Claim 4 of the formula



This Sheet Has Been Replaced





6. A compound according to Claim 5 wherein:

R_4 and R_6 are each independently halo, isopropyl, isobutyl, *sec*-butyl or *t*-butyl;

5 R_5 , R_7 and R_8 are each independently -H, halo, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl or *t*-butyl; and

there are no R_9 substituents.

7. A compound according to Claim 6 which is 2-(3,5-dichlorophenyl)-3-(3-pyridyl)-2-propenenitrile.

10 8. A method for inhibiting cell proliferation in a patient suffering from such disorder comprising administering to said patient a pharmaceutically effective amount of a compound according to Claim 2.

15 9. A method for the treatment of psoriasis in a patient suffering from such disorder comprising administering to said patient a composition according to Claim 3.

20 10. A method for the treatment of atherosclerosis in a patient suffering from such disorder comprising administering to said patient a composition according to Claim 3.

9608075-9

02 APR 1996

This Sheet Has Been Replaced



SUBSTITUTE SHEET

28 FEB 1997

✓
✓
✓1
36 40
41 -✓
✓
✓1
36 40
41 -

1

12/2/97


STYRYL-SUBSTITUTED MONOCYCLIC AND BICYCLIC HETEROARYL
COMPOUNDS WHICH INHIBIT EGF RECEPTOR TYROSINE KINASE


FIELD OF THE INVENTION

This invention relates to the inhibition of cell proliferation. More specifically, this invention relates to the use of styryl-substituted monocyclic and bicyclic heteroaryl compounds in inhibiting cell proliferation, including compounds which are useful protein tyrosine kinase (PTK) inhibitors.

Normal cellular reproduction is believed to be triggered by the exposure of the cellular substrate to one or more growth factors, examples of which are insulin, epidermal growth factor (EGF) and platelet-derived growth factor (PDGF). Such growth factors are typically specific for corresponding growth factor receptors which are imbedded in and which penetrate through the cellular membrane. The initiation of cellular reproduction is believed to occur when a growth factor binds to the corresponding receptor on the external surface of the cellular membrane. This

This Sheet Has Been Replaced





growth factor-receptor binding alters the chemical characteristics of that portion of the receptor which exists within the cell and which functions as an enzyme to catalyze phosphorylation of either an intracellular substrate or the receptor itself, the latter being referred to as autophosphorylation. Examples of such phosphorylation enzymes include tyrosine kinases, which catalyze phosphorylation of tyrosine amino acid residues of substrate proteins.

Many diseased states are characterized by the uncontrolled reproduction of cells. These diseased states involve a variety of cell types and include disorders such as leukemia, cancer, psoriasis, atherosclerosis and restenosis injuries. The inhibition of tyrosine kinase is believed to have utility in the control of uncontrolled cellular reproduction, i.e., cellular proliferative disorders.

Initiation of autophosphorylation, i.e., phosphorylation of the growth factor receptor itself, and of the phosphorylation of a host of intracellular substrates are some of the biochemical events which are involved in mitogenesis and cell proliferation. Autophosphorylation of the insulin receptor and phosphorylation of substrate proteins by other receptors are the earliest identifiable biochemical hormonal responses.

Elimination of the protein tyrosine kinase (PTK) activity of the insulin receptor and of the epidermal growth factor (EGF) receptor by site-directed mutagenesis of the cellular genetic material which is responsible for generation of insulin and EGF results


28 FEB 1997

in the complete elimination of the receptors' biological activity. This is not particularly desirable because insulin is needed by the body to perform other biological functions which are not related to cell proliferation. Accordingly, compounds which inhibit the PTK portion of the EGF receptor at concentrations less than the concentrations needed to inhibit the PTK portion of the insulin receptor could provide valuable agents for selective treatment of cell proliferation disorders.

REPORTED DEVELOPMENTS

U.S. Patent Nos. 4,678,793 and 4,826,984 disclose pharmaceutical compositions including styryl 4,4-dimethyl (bicyclic heteroaryl) compounds as active agents for treating cancer, psoriasis, acne, etc. U.S. Patent No. 4,769,384 discloses pharmaceutical compositions including styryl benzimidazole compounds as active agents for treating ulcers of the stomach and duodenum.

It has been reported that the most potent inhibitors of EGF receptors inhibit EGF-induced proliferation of A431/clone 15 cells with little or no effect on the proliferation of such cells when induced by other growth factors. It has been reported also that erbstatin inhibits the autophosphorylation of the EGF receptor in membranes of A431 cells. Low concentrations of erbstatin are required to inhibit EGF receptor autophosphorylation, whereas much higher concentrations of erbstatin are required to inhibit cyclic adenosine 3',5'-monophosphate (cAMP)-dependent protein kinase.



SUMMARY OF THE INVENTION


In accordance with the present invention, there is provided a method of inhibiting cell proliferation in a patient suffering from such disorder comprising the administration to the patient of a styryl-substituted heteroaryl compound exhibiting protein tyrosine kinase inhibition activity wherein the heteroaryl group is a monocyclic ring with 1 or 2 heteroatoms, or a bicyclic ring with 1 to about 4 heteroatoms, said compound optionally substituted or polysubstituted, with the proviso that when said ring is polysubstituted, the substituents do not have a common point of attachment to said ring.

Another aspect of the present invention relates to novel compounds which are those compounds of the aforementioned type wherein no substituent on the heteroaryl group is a carboxy group or an ester group.

Still another aspect of the present invention relates to pharmaceutical compositions comprising, in admixture with a pharmaceutically acceptable carrier, a pharmaceutically-effective amount of a novel compound of the aforementioned type.

With respect to the method aspects of this invention, the compounds described by Formula I below constitute a class of the aforementioned styryl-substituted heteroaryl compounds for use in the practice of the present invention:

73 FEB 1987



I

wherein:

R₁ is alkyl, -H, -CN, -OH, -COOR, -CONRR or -CSNRR;

R is alkyl, -H or aralkyl;


R₂ is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms or 1 or 2 N-oxide groups, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms or 1 to about 4 N-oxide groups, said ring optionally substituted with one to about three R₃ groups, said R₃ substituents having no common points of attachment to said ring;

R₃ is alkyl, -H, -CN, -OH, -COOR, -CONRR, -CSNRR or -CH₂CN;

R₄, R₅, R₆, R₇ and R₈ are each independently alkyl, -H, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic or heteroaryl;

each R₉ is independently alkyl, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic or heteroaryl

28 FEB 1967



; and

R_3 and R_7 together may be $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$ or, starting from R_3 , $-\text{CONH}-$; or

a pharmaceutically acceptable salt thereof.

Also in accordance with the present invention, novel compounds within the scope of the compound and pharmaceutical composition aspects of the present invention are described by Formula I above wherein:

R_1 is alkyl, $-\text{CN}$, $-\text{COOR}$, $-\text{CONRR}$ or $-\text{CSNRR}$;

R is alkyl, $-\text{H}$ or aralkyl;

R_2 is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common points of attachment;

R_3 is alkyl, $-\text{H}$, $-\text{COOR}$, $-\text{CONRR}$, $-\text{CSNRR}$ or $-\text{CH}_2\text{CN}$;


R_4 and R_6 are each independently alkyl, $-\text{H}$, $-\text{CN}$, halo, $-\text{OR}$, $-\text{CHO}$, $-\text{COOH}$, $-\text{NRR}$, $-\text{NO}_2$, $-\text{NHCOCH}_3$, $-\text{SR}$, $-\text{CF}_3$,

$-\text{CH}=\text{CHCOOH}$, $-\text{NHCO}(\text{CH}_2)_2\text{COOH}$, heterocyclic or heteroaryl;

R_5 , R_7 and R_8 are each independently alkyl, $-\text{H}$, $-\text{CN}$, $-\text{OR}$, $-\text{CHO}$, $-\text{COOH}$, $-\text{NHCOCH}_3$, $-\text{SR}$, $-\text{CF}_3$, $-\text{CH}=\text{CHCOOH}$, $-\text{NHCO}(\text{CH}_2)_2\text{COOH}$, heterocyclic or heteroaryl;

with the provisos that at least two of R_4 , R_5 , R_6 , R_7 and R_8 are not $-\text{H}$, and R_4 , R_5 or R_6 cannot be $-\text{OR}$ when R_7 or R_8 is $-\text{OR}$; and


each R_9 is independently alkyl, $-\text{CN}$, halo, $-\text{OR}$, $-\text{CHO}$, $-\text{NRR}$, $-\text{NO}_2$, $-\text{NHCOCH}_3$, $-\text{SR}$, $-\text{CF}_3$, $-\text{CH}=\text{CHCOOH}$, $-\text{NHCO}(\text{CH}_2)_2\text{COOH}$, heterocyclic heteroaryl or



R_3 and R_7 together may be $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$ or, starting from R_3 , $-\text{CONH}-$.

In an alternate embodiment of the present invention, novel compounds within the scope of the compound and pharmaceutical composition aspects of the present invention are described by Formula II below.

8



II

wherein:

R is alkyl, $-H$ or aralkyl;


R_2 is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common points of attachment;

R_4 and R_6 are each independently branched alkyl of about 3 to about 6 carbon atoms, $-CN$, halo, $-SR$, $-CF_3$, heterocyclic, or heteroaryl;

R_5 , R_7 and R_8 are each independently alkyl, $-H$, $-CN$, $-OR$, $-CHO$, $-COOH$, $-NHCOCH_3$, $-SR$, $-CF_3$, $-CH=CHCOOH$, $-NHCO(CH_2)_2COOH$, heterocyclic or heteroaryl;

each R_9 is independently alkyl, $-CN$, halo, $-OR$, $-CHO$, $-NRR$, $-NO_2$, $-NHCOCH_3$, $-SR$, $-CF_3$, $-CH=CHCOOH$, $-NHCO(CH_2)_2COOH$, heterocyclic, heteroaryl or

28 FEB 1997



Compounds within the scope of the present invention have also a specific affinity toward the substrate site of the tyrosine kinase domain of EGF receptors, inhibit EGF receptor kinase more than they inhibit PDGF receptor kinase and also effectively inhibit EGF-dependent autophosphorylation of the receptor.

DETAILED DESCRIPTION OF THE INVENTION

As employed above and throughout the disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

"Alkyl" means a saturated aliphatic hydrocarbon which may be either straight- or branch-chained containing from about 1 to about 6 carbon atoms.

"Lower alkyl" means an alkyl group as above, having 1 to about 4 carbon atoms which may be straight- or branch-chained such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl.

"Alkoxy" means an alkyl-oxy group in which "alkyl" is as previously described. Lower alkoxy groups are preferred. Exemplary groups include methoxy, ethoxy, n-propoxy, i-propoxy and n-butoxy.

28 FEB 1997

"Aryl" means an unsaturated or partially unsaturated ring system. Preferred aryl groups are pyridyl and indolyl.

"Heterocyclic" means an unsaturated ring system containing 1 or 2 N, O or S heteroatoms. Preferred heterocyclic groups are morpholine, piperidine and piperazine.

"Acyl" means an organic radical derived from an organic acid, a carboxylic acid, by the removal of its acid hydroxyl group. Preferred acyl groups are lower alkyl carboxylic acid groups such as acetyl and propionyl. Benzoyl is also preferred.

"Halo" means a halogen. Preferred halogens include chloride, bromide and fluoride.

Preferred aralkyl groups are benzyl and phenethyl.


It is believed that therapeutically useful PTK inhibiting compounds should be competitive with the substrate of EGF receptor tyrosine kinase (EGFRK) and not with adenosine triphosphate (ATP). The PTK inhibitors quercetin and genistein, which compete with ATP, inhibit other protein kinases and as a result are highly cytotoxic. As a test of selectivity, compounds which inhibit EGFRK better than they inhibit insulin receptor kinase (IRK) and/or PDGF receptor kinase are of considerable value.

It is theorized that solubility of the compounds of the present invention both in water and in mildly hydrophobic solvents will enhance the probability that

20 FEB 1977

they traverse the cell membrane. Various insoluble compounds, however, have exhibited significant EGFRK inhibition in in vitro testing.

In accordance with the method aspects of the present invention, a preferred class of styryl-substituted heteroaryl compounds for use in the practice of the present invention are described by Formula II below.



II

wherein:

R is alkyl, -H or aralkyl;

R₂ is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms or 1 or 2 N-oxide groups, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms or 1 to about 4 N-oxide groups, said ring optionally substituted with one to about three R₉ groups, said R₉ substituents having no common points of attachment to said ring;


R₄, R₅, R₆, R₇ and R₈ are each independently alkyl, -H,

28 FEB 1977

12

-CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, or heteroaryl;

each R_i is independently alkyl, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, heteroaryl or



A preferred class of compounds useful in the practice of the present invention include those described by Formula I where:

R₁ is -CN, -COOR, -CONRR or -CSNRR;

R is lower alkyl, -H or aralkyl;


R₂ is a 6-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or a 9- or 10-membered bicyclic aryl ring including 1-4 N, O or S atoms, said ring optionally substituted with one to about three R₃ groups, said R₃ substituents having no common points of attachment to said ring;

R₃ is -H;

R_4 , R_5 , R_6 , R_7 and R_8 are each independently lower alkyl, -H, lower alkoxy or -OH, with the provisos that at least two of R_4 , R_5 , R_6 , R_7 and R_8 are not -H, and R_4 , R_5 or R_6 cannot be lower alkoxy when R_7 or R_8 is lower alkoxy;


R_4 and R_6 are also each independently halo;

each R_i is independently lower alkyl, halo, lower alkoxy, -OH or




a pharmaceutically acceptable salt thereof.

More preferred compounds for use in the practice of this invention include those of Formulae III and IV below:



III

28 FEB 1997



IV

where

R₁ is -CN, -COOR, -CONRR or -CSNRR;

R is lower alkyl, -H or aralkyl;


R₂ is a 6-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or a 9- or 10-membered bicyclic aryl ring including 1-4 N, O or S atoms, said ring optionally substituted with one to about three R₉ groups, said R₉ substituents having no common points of attachment to said ring;

R₃ is -H;

R₄, R₅, R₆, R₇ and R₈ are each independently lower alkyl, -H, lower alkoxy or -OH, with the provisos that at least two of R₄, R₅, R₆, R₇ and R₈ are not -H, and R₄, R₅ or R₆ cannot be lower alkoxy when R₇ or R₈ is lower alkoxy;


R₄ and R₆ are also each independently halo;

each R₉ is independently lower alkyl, halo, lower alkoxy, -OH or



; and

R_{10} is -H, alkyl, aralkyl or



; or

a pharmaceutically acceptable salt thereof.

Even more preferred compounds are described by Formulae III and IV where:

R_1 is -CN, -COOR or -CONRR;

R_4 , R_5 , R_6 , R_7 and R_8 are independently lower alkyl, -H, lower alkoxy or -OH, with the provisos that at least two of R_4 , R_5 , R_6 , R_7 and R_8 are not -H, and R_4 , R_5 or R_6 cannot be lower alkoxy when R_7 or R_8 is lower alkoxy;

R_4 and R_6 are also each independently halo;

28 FEB 1971

R is lower alkyl or -H; and

there are no R_9 substituents.

The most preferred compounds are described by Formula II where R_1 is -CN, R_5 , R_7 and R_8 are each independently -H, and R_4 and R_6 are each independently alkyl, halo, -OR or -CF₃.


In an alternate embodiment of the present invention, a preferred class of compounds are described by Formula II above wherein

R_2 is a 6-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or a 9- or 10-membered bicyclic aryl ring including 1-4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common points of attachment to said ring;

R_4 and R_6 are each independently branched lower alkyl of about 3 to about 4 carbon atoms or halo;


R_5 , R_7 and R_8 are each independently -H lower alkyl of about 2 to about 4 carbon atoms or halo; and

each R_9 is independently lower alkyl, halo, lower alkoxy, -OH or



23 FEB 1997

In an alternate embodiment of the present invention, more preferred compounds for use in the practice of the present invention are described by Formula V below.



V

where R₄, R₅, R₆, R₇, R₈ and R₉ are as described immediately above.

In an alternate embodiment of the present invention, even more preferred compounds are described by Formula V where R₄ and R₆ are each independently halo, isopropyl, isobutyl, *sec*-butyl or *t*-butyl; R₅, R₇ and R₈ are each independently -H, halo, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl or *t*-butyl; and there are no R₉ substituents.

The most preferred compounds in an alternate embodiment of the present invention are described by Formula V where R₄ and R₆ are each independently halo, and R₅, R₇ and R₈ are each independently -H.

28 FEB 1997

Compounds of this invention may be useful in the form of the free base, in the form of salts and as a hydrate. All forms are within the scope of the invention. Acid addition salts may be formed and are simply a more convenient form for use; and in practice, use of the salt form inherently amounts to use of the base form. The acids which can be used to prepare the acid addition salts include preferably those which produce, when combined with the free base, pharmaceutically acceptable salts, that is, salts whose anions are non-toxic to the animal organism in pharmaceutical doses of the salts, so that the beneficial properties inherent in the free base are not vitiated by side effects ascribable to the anions. Although pharmaceutically acceptable salts of said basic compound are preferred, all acid addition salts are useful as sources of the free base form even if the particular salt per se is desired only as an intermediate product as, for example, when the salt is formed only for purposes of purification and identification, or when it is used as an intermediate in preparing a pharmaceutically acceptable salt by ion exchange procedures.


Pharmaceutically acceptable salts within the scope of the invention include those derived from the following acids: mineral acids such as hydrochloric acid, sulfuric acid, phosphoric acid and sulfamic acid; and organic acids such as acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, cyclohexylsulfamic acid, quinic acid, and the like.

The corresponding acid addition salts comprise the following: hydrochloride, sulfate, phosphate, sulfamate, acetate, citrate, lactate, tartarate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate, cyclohexylsulfamate and quinate, respectively.

The acid addition salts of the compounds of this invention are prepared either by dissolving the free base in aqueous or aqueous-alcohol solution or other suitable solvents containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and acid in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.

Compounds useful in the pharmaceutical composition and method aspects of this invention can be prepared by known methods, for example, Knoevenagel condensation reactions such as those disclosed in U.S. Patent No. 3,149,148.

Compounds of this invention may be prepared by the following reaction sequence:



Knoevenagel condensation of a substituted benzaldehyde in a polar media with an active methylene ,


28 FEB 1967

compound of the formula $R_1CH_2R_2$ in the presence of ammonia or amines such as piperidine and raised heat results in the products of this invention. When substitution of the R_3 group is desired, the corresponding ketone starting material is used. Reaction temperatures in the range of 25°C to reflux and reaction times vary depending on the materials being used in the condensation.

Compounds of this invention are either commercially available, known in the literature or can be made by known procedures. For example, U.S. Patent No. 4,600,712 discloses fungicides of Formula I where, for example, R_1 is cyano, R_2 is pyridyl and R_3 and R_7 or R_8 are chloro. U.S. Patent Nos. 3,337,565 and 3,337,568 disclose compounds which interfere with carbohydrate metabolism of Formula I where, for example, R_1 is cyano or hydroxy, R_2 is pyridyl and R_3 is hydroxy. U.S. Patent No. 3,196,158 discloses adrenal cortex inhibitors of Formula I where, for example, R_1 is cyano, R_2 is pyridyl and R_7 or R_8 are halo. U.S. Patent No. 3,157,663 discloses adrenal cortex inhibitors where, for example, R_1 is cyano, R_2 is pyridyl and R_3 , R_7 or R_8 are amino or nitro groups. Buu-Hoi et al., Journal of the Chemical Society (C), pp. 2069-70 (1969) disclose the conversion of 1,2-diarylacrylonitriles to the corresponding 3-arylcoumarins, wherein the 2-aryl group bears an ortho-alkoxy-substituent. Although the foregoing publications disclose some compounds of the type that can be used in accordance with the method aspects of the present invention, they do not disclose the use of such compounds for inhibiting cell proliferation.

28 FEB 1997

Various R, R₁, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ substituents on the phenyl and heterocyclic ring or chain can be present in the starting compound or added after formation of the condensation product by methods known in the art for substitution or conversion on one group to another. If the substituents themselves are reactive, then the substituents can themselves be protected according to the techniques known in the art. A variety of protecting groups known in the art, may be employed. Examples of many of these possible groups may be found in "Protective Groups in Organic Synthesis" by T. W. Green, John Wiley and Sons, 1981. For example, nitro groups can be added to the aromatic ring by nitration and the nitro group converted to other groups, such as amino by reduction, and halo by diazotization of the amino group and replacement of the diazo group. Acyl groups can be substituted onto the aryl groups by Friedel-Crafts acylation. The acyl groups can then be transformed to the corresponding alkyl groups by various methods, including the Wolff-Kishner reduction and Clemmenson reduction. Amino groups can be alkylated to form mono- and di-alkylamino groups; and mercapto and hydroxy groups can be alkylated to form corresponding ethers. Primary alcohols can be oxidized by oxidizing agents known in the art to form carboxylic acids or aldehydes, and secondary alcohols can be oxidized to form ketones. Tertiary amino groups can be converted to the corresponding N-oxides by oxidizing agents known in the art, for example, hydrogen peroxide and peracids. Thus, substitution or alteration reactions can be employed to provide a variety of substituents throughout the molecule of the starting material, intermediates, or the final product.



Compounds within the scope of this invention exhibit significant activity as protein tyrosine kinase inhibitors and possess therapeutic value as cellular antiproliferative agents for the treatment of certain conditions including psoriasis, atherosclerosis and restenosis injuries. It is expected that the invention will be particularly applicable to the treatment of atherosclerosis. With regard to the treatment of some conditions, for example, atherosclerosis, certain people may be identified as being at high risk, for example, due to genetic, environmental or historical factors. Compounds within the scope of the present invention can be used in preventing or delaying the occurrence or reoccurrence of such conditions or otherwise treating the condition.

The compounds of the present invention can be administered to a mammalian host in a variety of forms adapted to the chosen route of administration, i.e., orally, or parenterally. Parenteral administration in this respect includes administration by the following routes: intravenous, intramuscular, subcutaneous, intraocular, intrasynovial, transepithelial including transdermal, ophthalmic, sublingual and buccal; topically including ophthalmic, dermal, ocular, rectal and nasal inhalation via insufflation and aerosol and rectal systemic.

The active compound may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsules, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral

28 FEB 1967

therapeutic administration, the active compound may be incorporated with excipient and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 6% of the weight of the unit. The amount of active compound in such therapeutically useful compositions is such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit form contains between about 1 and 1000 mg of active compound.

The tablets, troches, pills, capsules and the like may also contain the following: A binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring

28 FEB 1957

such as cherry or orange flavor. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and formulations.

The active compound may also be administered parenterally or intraperitoneally. Solutions of the active compound as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersion can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.


The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It may be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required

28 FEB 1967

particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

The therapeutic compounds of this invention may be administered to a mammal alone or in combination with pharmaceutically acceptable carriers, as noted above, the proportion of which is determined by the solubility and chemical nature of the compound, chosen route of administration and standard pharmaceutical practice.



The dosage of the present therapeutic agents which will be most suitable for prophylaxis or treatment will vary with the form of administration, the particular compound chosen and the physiological characteristics of the particular patient under treatment. Generally, small dosages will be used initially and if necessary, will be increased by small increments until the optimum effect under the circumstances is reached. The therapeutic human dosage, based on physiological studies using rats, will generally be from about 0.01 mg to about 100 mg/kg of body weight per day or from about 0.4 mg to about 10 g or and higher although it may be administered in several different dosage units from once to several times a day. Oral administration requires higher dosages.

EXAMPLES

Embodiments of the present invention and comparative examples are described in the following non-limiting examples which include a description of pharmacological test procedures believed to correlate to therapeutic activity in humans and other animals. Examples 1 and 2 below are illustrative of compounds within the scope of the present invention. In examples 1A and 1B, R_1 is cyano, R_2 is pyridyl and R_4 and R_6 are chloro. In example 2, R_2 is pyridyl, R_3 is cyano and R_4 and R_6 are chloro.

Example 1A

trans-2-(3-Pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile

To a stirred solution of 400 g (2.3 mole) 3,5-dichlorobenzaldehyde in 11.4 liters absolute ethanol were added 283.5 g (2.4 mole) 3-

28 FEB 1977

27

pyridylacetonitrile and 720.9 g of K_2CO_3 . Within 2 minutes of the K_2CO_3 addition, a solid precipitate formed. The reaction mixture was stirred about 2.5 hours. Water (22.9 liters) was added to the stirred reaction mixture. After stirring for 1 hour, the mixture was filtered and the filter cake was washed with water (10 liters), dried and recrystallized from isopropanol. Five hundred thirty four g (85% yield) of *trans*-2-(3-pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile were obtained, m.p. 150-151°C.

Example 1B

cis-2-(3-pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile

A stirred solution of *trans*-2-(3-pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile (20 g), for example, as prepared in Example 1A, in 3 liters acetonitrile was irradiated with a sunlamp (275 W) for 24 hours. The reaction mixture was stirred an additional 24 hours and concentrated *in vacuo*. The resulting residue was purified by flash chromatography on silica gel, eluting sequentially with hexanes, 5% EtOAc/hexanes, 10% EtOAc/hexanes, and then 15% EtOAc/hexanes. The resulting yellow solid was recrystallized from 15% EtOAc/heptane to give 8 g (74% yield) of the title compound, m.p. 114-117°C.

Example 2

2-(3,5-Dichlorophenyl)-3-(3-pyridyl)-2-propenenitrile

To a stirred solution of 3 g (17.1 mmole) of 3,5-dichlorobenzaldehyde in 200 ml absolute ethanol were added 1.83 ml (17.1 mmole) of 3-pyridyl-acetonitrile and 1 equivalent of K_2CO_3 (2.36 g). The reaction flask was equipped with a reflux condenser and


28 FEB 1997

the stirred reaction mixture was refluxed for 2 hours, filtered and concentrated. The resulting residue was purified by flash chromatography on silica gel, eluting with 4:1 hexane/ethyl acetate to give the title compound as a white solid, m.p. 139-141°C. An amount of *trans*-2-(3-pyridyl)-3-(3,5-dichlorophenyl)-2-propenenitrile was also obtained from the reaction mixture.

Compounds of this invention are subjected to various biological tests, the results of which correlate to useful cellular antiproliferative activity. These tests are useful in determining EGF receptor kinase, PDGF receptor kinase and insulin receptor kinase inhibition activities of the compounds disclosed herein.

EGF-Receptor Purification

EGF-receptor purification is based on the procedure of Yarden and Schlessinger. A431 cells are grown in 80 cm² bottles to confluency (2×10^7 cells per bottle). The cells are washed twice with PBS and harvested with PBS containing 1.0 mmol EDTA (1 hour at 37°C), and centrifuged at 600g for 10 minutes. The cells are solubilized in 1 ml per 2×10^7 cells of cold solubilization buffer (50 mmol Hepes buffer, pH 7.6, 1% Triton X-100, 150 mmol NaCl, 5 mmol EGTA, 1 mmol PMSF, 50 µg/ml aprotinin, 25 mmol benzamidine, 5 µg/ml leupeptic, and 10 µg/ml soybean trypsin inhibitor) for 20 minutes at 4°C. After centrifugation at 100000g for 30 minutes, the supernatant is loaded onto a WGA-agarose column (100 µl of packed resin per 2×10^7 cells) and shaken for 2 hours at 4°C. The unabsorbed material is removed and the resin washed twice with HTN



buffer (50 mmol Hepes, pH 7.6, 0.1% Triton X-100, 150 mmol NaCl), twice with HTN buffer containing 1 M NaCl, and twice with HTNG buffer (50 mmol Hepes, pH 7.6, 0.1% Triton X-100, 150 mmol NaCl, and 10% glycerol). The EGF receptor is eluted batchwise with HTNG buffer containing 0.5 M N-acetyl-D-glucosamine (200 μ l per 2×10^7 cells). The eluted material is stored in aliquots at -70°C and diluted before use with TMTNG buffer (50 mmol Tris-Mes buffer, pH 7.6, 0.1% Triton X-100, 150 mmol NaCl, 10% glycerol).

EGFR Kinase Catalyzed Phosphorylation of Poly(GAT) and its Inhibition

WGA-purified EGFR (0.25 $\mu\text{g}/\text{assay}$) is preactivated with EGF (0.85 μM) in 50 mmol Tris-Mes buffer, pH 7.6 for 20 minutes at 4°C . The assay is initiated by addition of a mixture which contains $\text{Mg}(\text{Ac})_2$ (60 mmol), $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ (125 μM , 2-5 $\mu\text{Ci}/\text{assay}$), poly(GAT) (0.0625 mg/ml, 0.125 mg/ml, 0.25 mg/ml), and six concentrations of inhibitor in duplicates. The temperature of the assay is 22°C and the production of phosphorylated copolymer is found to be linear up to 20 minutes. The PTK inhibitors tested are solubilized in water or a mixture of ethanol and water such that the final concentration of ethanol does not exceed 4% in the assay. Up to 4% ethanol in the assay has no effect on the EGFR kinase activity. The concentration of EGF in the assay is 300 nM in a final volume of 40 μl . After 5, 10 or 20 minutes, aliquots of 25 μl are applied onto Whatman 3-mm paper cuttings, which are then soaked in cold 10% TCA containing 0.01 M sodium pyrophosphate. After being washed overnight at 4°C , the paper cuttings are dried and counted, measuring ^{32}P Cerenkov radiation. Concentration dependence on poly(GAT) was Michaelian

with a $K_m = 0.076 \pm 0.007$ mg/ml or 0.069 ± 0.007 mmol if calculated per Glu₆Ala₃Tyr(GAT) unit. The EGF response for the poly(GAT) phosphorylation is graphed. The K_m for ATP in the assay was found to 2.9 μ M.

Time Dependence of EGF-Receptor Autophosphorylation

WGA-purified EGF receptor from A431 cells (0.5 μ g/assay) is activated with EGF (800 nM) for 20 minutes at 4°C. The reaction is initiated by the addition of Mg(Ac)₂ (60 mmol), Tris-Mes buffer, pH 7.6 (50 mmol), and [³²P]ATP (20 μ M, 5 μ Ci/assay). The reaction is conducted at either 4 or 15°C and terminated by addition of sodium dodecyl sulfate (SDS) sample buffer (10% glycerol, 50 mmol Tris, pH 6.8, 5% β -mercapto-ethanol, and 3% (SDS). The samples are run on a 8% SDS polyacrylamide gel (SDS-PAGE) (prepared from 30% acrylamide and 0.8% bis-(acrylamide) and contained 0.375 M Tris, pH 8.8, 0.1% SDS, 0.05% TEMED, and 0.46% ammonium persulfate). The gel is dried and autoradiography performed with Agfa Curix RP2 X-ray film. The relevant radioactive bands are cut and counted in the Cerenkov mode. The fast phase of autophosphorylation continues for another 10 minutes. The extent of phosphorylation completed in the first 10-s at 15°C comprises 1/3 of the total autophosphorylation signal and probably reflects the phosphorylation of the first site on the receptor. The 10-s interval is therefore chosen for use in subsequent autophosphorylation experiments.

ATP and EGF Dependence of Autophosphorylation

WGA-purified EGF receptor from A431 cells (0.5 μ g/assay) is activated with EGF (0.85 μ M) for 20 minutes at 4°C. The assay is performed at 15°C and initiated

by addition of $\text{Mg}(\text{Ac})_2$ (60 mmol), Tris-Mes buffer, pH 7.6 (50 mmol), $[\text{}^{32}\text{P}]\text{ATP}$ (carrier free, 5 $\mu\text{Ci}/\text{assay}$), and increasing concentrations of nonradioactive ATP. The assay is terminated after 10-s by addition of SDS sample buffer. The samples are run on a 6% SDS polyacrylamide gel. The gel is dried and autoradiographed as described above. The relevant radioactive bands are cut and counted in the Cerenkov mode. the K_m for ATP determined in this fashion is found to be 7.2 μM . With use of the 10-s assay protocol, the EGF concentration dependence of EGFRK autophosphorylation is determined.

Inhibition of Copoly(Glu₄Tyr) Phosphorylation by Insulin-Receptor Kinase (InsRK)

Rat liver membranes are prepared from the livers of 6-week-old rats as described by Cuatrecasas. WGA-purified insulin receptor is prepared according to Zick et al. WGA-purified rat liver InsRK (1.25 μg) is preincubated with or without 330 nM insulin in 50 mmol Tris-Mes buffer, pH 7.6, for 30 minutes at 22°C. The assay is performed at 22°C and initiated by addition of a mixture which contains $\text{Mg}(\text{Ac})_2$ (60 mmol), NaVO_3 (40 μM), $[\gamma\text{}^{32}\text{P}]\text{ATP}$ (125 μM , 3-5 $\mu\text{Ci}/\text{assay}$), and poly(GT) [poly(Glu₄Tyr)] at three concentrations: whenever an inhibitor is tested, it is added at the proper concentration. The final concentration of insulin in the assay is 125 nM. The total volume of the assay is 40 μl . After 20 minutes, aliquots of 30 μl are applied on Whatman 3-mm paper and soaked in cold 10% TCA, containing 0.01 M sodium pyrophosphate. After being washed overnight, the papers are dried and counted, measuring Cerenkov radiation. The InsRk-catalyzed

phosphorylation of poly(GT) obeys Michaelis-Menten kinetics.

Inhibition of EGFR Autophosphorylation

A431 cells were grown to confluence on human fibronectin coated tissue culture dishes. After washing 2 times with ice-cold PBS, cells were lysed by the addition of 500 μ l/dish of lysis buffer (50 mmol Hepes, pH 7.5, 150 mmol NaCl, 1.5 mmol $MgCl_2$, 1 mmol EGTA, 10% glycerol, 1% triton X-100, 1 mmol PMSF, 1 mg/ml aprotinin, 1 mg/ml leupeptin) and incubating 5 minutes at 4°C. After EGF stimulation (500 μ g/ml 10 minutes at 37°C) immunoprecipitation was performed with anti EGF-R (Ab 108) and the autophosphorylation reaction (50 μ l aliquots, 3 μ Ci [γ - 32 P]ATP) sample was carried out in the presence of 2 or 10 μ M of compound, for 2 minutes at 4°C. The reaction was stopped by adding hot electrophoresis sample buffer. SDS-PAGE analysis (7.5% els) was followed by autoradiography and the reaction was quantitated by densitometry scanning of the x-ray films. In order to test the compounds for selective inhibition, the procedure is repeated using PDGF stimulation in place of EGF stimulation. "IC₅₀," as used below, refers to the concentration of inhibitor (μ M) at which the rate of autophosphorylation is halved, compared with media containing no inhibitor. Preferably, the IC₅₀ values for EGF will be lower than for PDGF, indicating a high degree of EGF specificity of the inhibitory compounds. The results of this test are summarized in Table I below.




Table I

Example	IC ₅₀ (μM)	
	EGF	PDGF
2	0.1	20
3	0.5	6
5	0.4	10
6	20	>50
7	0.4	30

These results show that the compounds of the present invention inhibit EGF receptor kinase better than they inhibit PDGF receptor kinase.

Inhibition of Cell Proliferation as
Measured by Inhibition of DNA Synthesis

Cells were seeded at 1×10^5 cells per well in 24-well Costar dishes pre-coated with human fibronectin (by incubating for 30 minutes at room temperature with 10 μg/0.5 ml/well). The cells were grown to confluence for 2 days. The medium was changed to DMEM containing 0.5 calf serum for 36-48 hours and the cells were then incubated with PDGF, EGF (Toyobo, New York, NY) (20 ng/ml) or serum (10% calf serum, FCS) and different concentrations of the inhibitory compounds.

[³H]thymidine, (NEN, Boston, MA) was added 16-24 hours later at 0.5 μCi/ml for 2 hours. TCA precipitable material was quantitated by scintillation counting (C). Results of this assay are summarized in Table II below.

"IC₅₀," as used below, refers to the concentration of inhibitor (nM) at which [³H]thymidine incorporation is halved, compared with media containing no inhibitor. As FCS contains a broad range of growth factors, the IC₅₀ values for PDGF should be lower than for FCS, indicating that the compounds do not act as general inhibitors.

TABLE II

<u>Example</u>	<u>IC₅₀ - PDGF</u>	<u>IC₅₀ - FCS</u>
2	30±17	100
3	14±10	27±5% @ 1 μM
4	9±2	500
6	17	15% @ 1 μM

These results indicate that the compounds of the invention do not inhibit a broad range of growth factor receptors.

Cell Culture

Cells termed HER 14 and K721A (=DK) were prepared by transfecting N1H3T3 cells (clone 2.2) (From C. Fryling, NCI, NIH), which lack endogenous EGF-receptors, with cDNA constructs of wild-type EGF-receptor or mutant EGF-receptor lacking tyrosine kinase activity (in which Lys 721 at the ATP-binding site was replaced by an Ala residue, respectively). All cells were grown in DMEM with 10% calf serum (Hyclone, Logan, Utah).

The results obtained by the above experimental methods evidence the useful protein tyrosine kinase inhibition properties of the compounds within the scope of the present invention.

Comparative Examples

The following comparative examples illustrate the improved cell proliferation inhibition of compounds of the present invention in comparison to compounds of the prior art. In comparative Example C-1, the preparation of a prior art compound, where R₁ is cyano, R₂ is pyridyl and R₃ and R₄ are chloro is described. The inhibitory activities of the compounds of Examples 1A

and 2 are compared to the inhibitory activity of the compound prepared in comparative Example C-1.

Example C-1

2-(3-pyridyl)-3-(2,4-dichlorophenyl)-2-propenenitrile


To a stirred solution of 2,4-dichlorobenzaldehyde (5.0 g, 29 mmole) and 3-pyridylacetonitrile (3.05 ml, 29 mmole) in 150 ml absolute ethenol were added 3.95 g K_2CO_3 . After stirring for about 15 minutes, a white solid precipitated out of the reaction mixture. The reaction mixture was stirred overnight and filtered. The filter cake was washed with absolute ethanol and dissolved in CH_2Cl_2 . The CH_2Cl_2 solution was filtered and concentrated in vacuo. The resulting white solid was recrystallized from 3:2 hexane/ethyl acetate to give 6.0 g (76%) of the title compound.

The inhibition of EGFR autophosphorylation by the compounds of Examples 1A and 2 was compared to the prior art compound of Example C-1. The EGFR autophosphorylation inhibition was evaluated using the same procedure as described hereinbefore. The results are reported in Table III below.

TABLE III

<u>Example</u>	<u>IC₅₀ (μM)</u>
1A	2
2	2-6
C-1	≥50


These results demonstrate that the compounds of Examples 1A and 2 are more potent inhibitors of EGF receptor kinase than C-1.





CLAIMS

1. A method of inhibiting cell proliferation in a patient suffering from such disorder comprising the administration to the patient of a pharmaceutically effective amount of a pharmaceutical composition containing, in admixture with a pharmaceutically acceptable carrier, a compound, or a pharmaceutically acceptable salt thereof, of the formula



wherein:

R is alkyl, -H or aralkyl;

R₂ is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms or 1 or 2 N-oxide groups, or an about 8- to about 12-membered bicyclic aryl ring including 1 to about 4 N, O or S atoms or 1 to about 4 N-oxide groups, said ring optionally substituted with one to about three R₉ groups, said R₉ substituents having no common points of attachment to said ring;

R₄, R₅, R₆, R₇ and R₈ are each independently alkyl, -H,

This Sheet Has Been Replaced




08 FEB 1997




-CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, or heteroaryl;

each R₉ is independently alkyl, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, heteroaryl or



2. A compound of the formula



wherein:

R is alkyl, -H or aralkyl;

R₂ is an about 5- to about 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or an about 8- to about 12-membered bicyclic ,






aryl ring including 1 to about 4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common points of attachment;

R_4 and R_6 are each independently branched alkyl of about 3 to about 6 carbon atoms, -CN, halo, -SR, -CF₃, heterocyclic, or heteroaryl;


R_5 , R_7 and R_8 are each independently alkyl, -H, -CN, -OR, -CHO, -COOH, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic or heteroaryl;

each R_9 is independently alkyl, -CN, halo, -OR, -CHO, -NRR, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, heteroaryl or



3. A pharmaceutical composition for inhibiting cell proliferation in a patient suffering from such disorder comprising, in admixture with a pharmaceutically acceptable carrier, a pharmaceutically effective amount of a compound according to Claim 2.

This Sheet Has Been Replaced






4. A compound according to Claim 2 wherein:

R_2 is a 6-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or a 9- or 10-membered bicyclic aryl ring including 1-4 N, O or S atoms, said ring optionally substituted with one to about three R_9 groups, said R_9 substituents having no common points of attachment to said ring;


R_4 and R_6 are each independently branched lower alkyl of about 3 to about 4 carbon atoms or halo;

R_5 , R_7 and R_8 are each independently -H lower alkyl of about 2 to about 4 carbon atoms or halo; and


each R_9 is independently lower alkyl, halo, lower alkoxy, -OH or



5. A compound according to Claim 4 of the formula



This Sheet Has Been Replaced





6. A compound according to Claim 5 wherein:

R_4 and R_6 are each independently halo, isopropyl, isobutyl, sec-butyl or *t*-butyl;

R_5 , R_7 and R_8 are each independently -H, halo, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, sec-butyl or *t*-butyl; and

there are no R_9 substituents.

7. A compound according to Claim 6 which is 2-(3,5-dichlorophenyl)-3-(3-pyridyl)-2-propenenitrile.

8. A method for inhibiting cell proliferation in a patient suffering from such disorder comprising administering to said patient a pharmaceutically effective amount of a compound according to Claim 2.

9. A method for the treatment of psoriasis in a patient suffering from such disorder comprising administering to said patient a composition according to Claim 3.

10. A method for the treatment of atherosclerosis in a patient suffering from such disorder comprising administering to said patient a composition according to Claim 3.

This Sheet Has Been Replaced




20 FEB 1

STYRYL-SUBSTITUTED MONOCYCLIC AND BICYCLIC HETEROARYL
COMPOUNDS WHICH INHIBIT EGF RECEPTOR TYROSINE KINASE

ABSTRACT OF THE DISCLOSURE


Methods of inhibiting cell proliferation in a patient suffering from such disorder comprising the use of a styryl-substituted heteroaryl compound wherein the heteroaryl group is a monocyclic ring with 1 or 2 heteroatoms, or a bicyclic ring with 1 to about 4 heteroatoms, said compound optionally substituted or polysubstituted, with the proviso that when said ring is polysubstituted, the substituents do not have a common point of attachment to said ring, and those compounds wherein no substituent on the heteroaryl group is a carboxy group or an ester group, and pharmaceutical compositions comprising such compounds.

This Sheet Has Been Replaced



✓	1	-
✓	36	40
✓	41	-

2/10/94



1

STYRYL-SUBSTITUTED HETEROARYL COMPOUNDS WHICH INHIBIT
EGF RECEPTOR TYROSINE KINASE

FIELD OF THE INVENTION



This invention relates to the inhibition of cell proliferation. More specifically, this invention relates to the use of styryl-substituted monocyclic and bicyclic heteroaryl compounds in inhibiting cell proliferation, including compounds which are useful protein tyrosine kinase (PTK) inhibitors.

Normal cellular reproduction is believed to be triggered by the exposure of the cellular substrate to one or more growth factors, examples of which are insulin, epidermal growth factor (EGF) and platelet-derived growth factor (PDGF). Such growth factors are typically specific for corresponding growth factor receptors which are imbedded in and which penetrate through the cellular membrane. The initiation of cellular reproduction is believed to occur when a growth factor binds to the corresponding receptor on the external surface of the cellular membrane. This



Claims

1. A compound of the formula



wherein:

R is alkyl, -H or aralkyl;

R₂ is a 5- to 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms, or an 8- to 12-membered bicyclic aryl ring including 1 to 4 N, O or S atoms, said ring optionally substituted with one to three R₉ groups, said R₉ substituents having no common points of attachment to said ring;

R₄ and R₆ are each independently branched alkyl of 3 to 6 carbon atoms, -CN, halo, -SR, -CF₃, heterocyclic, or heteroaryl;


R₅, R₇ and R₈ are each independently alkyl, -H, -CN, -OR, -CHO, -COOH, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, or heteroaryl;

with the provisos that at least two of R₄, R₅, R₆, R₇ and R₈ are not -H, and R₄, R₅ or R₆ cannot be -OR when R₇ or R₈ is -OR; and

each R₉ is independently alkyl, -CN, halo, -OR, -CHO, -NRR, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH,



-NHCO(CH₂)₂COOH, heterocyclic, heteroaryl or



and;

R₃ and R₇ together may be -CH₂CH₂-, -CH₂CH₂CH₂- or, starting from R₃, -CONH-.


2. A compound according to Claim 1 wherein:

R₂ is a 6-membered monocyclic acyl ring including 1 or 2 N, O or S atoms, or a 9- to 10-membered bicyclic aryl ring including 1 to 4 N, O or S atoms, said ring optionally substituted with one to three R₉ groups, said R₉ substituents having no common points of attachment to said ring;


R₄ and R₅ are each independently branched alkyl of 3 to 4 carbon atoms or halo;


R₃, R₇ and R₈ are each independently -H, alkyl of 2 to 4 carbon atoms or halo; and

each R₉ is independently alkyl, halo, alkoxy, -OH or



3. A compound according to Claim 2 of the formula





in which R_4 , R_5 , R_6 , R_7 , R_8 , R_9 are as defined in claim 2.

4. A compound according to Claim 3 wherein:

R_4 and R_6 are each independently halo, isopropyl, isobutyl, sec-butyl or t-butyl;


R_5 , R_7 and R_8 are each independently -H, halo, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl or t-butyl; and

there are no R_9 substituents.

5. A compound according to Claim 1 which is 2-(3,5-dichlorophenyl)-3-(3-pyridyl)-2-propenenitrile.

6. A pharmaceutical composition comprising in admixture with a pharmaceutically acceptable carrier, a pharmaceutically effective amount of a compound according to Claim 1.

7. Use of a compound of formula



wherein:


R is alkyl, -H or aralkyl;



R_2 is a 5- to 7-membered monocyclic aryl ring including 1 or 2 N, O or S atoms or 1 or 2 N-oxide groups, or an 8- to 12-membered bicyclic ring including 1 to 4 N, O or S atoms or 1 to 4 N-oxide groups, said ring optionally substituted with one to three R_3 groups, said R_3 substituents having no common points of attachment to said ring;

R_4 , R_5 , R_6 , R_7 and R_8 are each independently alkyl, -H, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, or heteroaryl;

each R_9 is independently alkyl, -CN, halo, -OR, -CHO, -COOH, -NRR or an N-oxide thereof, -NO₂, -NHCOCH₃, -SR, -CF₃, -CH=CHCOOH, -NHCO(CH₂)₂COOH, heterocyclic, heteroaryl or




or a pharmaceutically acceptable salt thereof, for the preparation of a pharmaceutical composition for inhibiting cell proliferation in a patient suffering from such disorder.

8. Use of a compound according to claim 1 for the preparation of a pharmaceutical composition for inhibiting cell proliferation in a patient suffering from such disorder.

9. Use of a compound according of claim 1 for the preparation of a pharmaceutical composition for treating psoriasis in a patient suffering from such disorder.



10. Use of a compound according of claim 1 for the preparation of a pharmaceutical composition for treating atherosclerosis in a patient suffering from such disorder.



- 41 -

STYRYL-SUBSTITUTED HETEROARYL COMPOUNDS WHICH
INHIBIT EGF RECEPTOR TYROSINE KINASEABSTRACT OF THE DISCLOSURE

Methods of inhibiting cell proliferation in a patient suffering from such disorder comprising the use of a styryl-substituted heteroaryl compound wherein the heteroaryl group is a monocyclic ring with 1 or 2 heteroatoms, or a bicyclic ring with 1 to about 4 heteroatoms, said compound optionally substituted or polysubstituted, with the proviso that when said ring is polysubstituted, the substituents do not have a common point of attachment to said ring, and those compounds wherein no substituent on the heteroaryl group is a carboxy group or an ester group, and pharmaceutical compositions comprising such compounds.

