

HERBICIDAL COMPOSITIONS COMPRISING 4-AMINO-3-CHLORO-5-FLUORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL)PYRIDINE-2-CARBOXYLIC ACID OR A DERIVATIVE THEREOF AND CERTAIN PS II INHIBITORS

Priority Claim

5 This application claims the benefit of United States provisional patent application number 61/675,089 filed on July 24, 2012, and United States patent application serial number 13/840,488, filed March 15, 2013, the disclosure of each of which is incorporated herein by reference in its entirety.

Field

10 Provided herein are herbicidal compositions comprising and methods for controlling undesirable vegetation utilizing (a) 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid or an agriculturally acceptable ester or salt thereof and (b) a photosystem (PS) II inhibitor selected from the group consisting of: atrazine, bentazon-sodium, bromoxynil, chlorotoluron, cyanazine, diuron, hexazinone, ioxynil,
15 isoproturon, linuron, methibenzuron, metribuzin, propanil, pyridate, siduron, simazine, simetryne, tebuthiuron and terbuthylazine, or derivative, *e.g.*, salt or ester thereof.


Background

 The protection of crops from weeds and other vegetation which inhibit crop growth is a constantly recurring problem in agriculture. To help combat this problem, researchers in
20 the field of synthetic chemistry have produced an extensive variety of chemicals and chemical formulations effective in the control of such unwanted growth. Chemical herbicides of many types have been disclosed in the literature and a large number are in commercial use. However, there remains a need for compositions and methods that are effective in controlling undesirable vegetation.

25 Summary

 Several embodiments are recited below. In the embodiments, the ratio of compound (a) to compound (b) can be expressed in units of weight to weight (g to g), gae/ha to gae/ha or gae/ha to gai/ha.

1. A synergistic herbicidal composition comprising a herbicidally effective amount of
30 (a) a compound of the formula (I)



or an agriculturally acceptable salt or ester thereof and (b) atrazine, bentazon, bentazon-sodium, bromoxynil, chlorotoluron, cyanazine, diuron, hexazinone, ioxynil, isoproturon, linuron, methibenzuron, metribuzin, propanil, pyridate, siduron, simazine, simetryne, 5 tebuthiuron or terbuthylazine or agriculturally acceptable salt or ester thereof.

2. The composition of embodiment 1, wherein (a) is a C₁₋₄ alkyl or benzyl ester of compound (I).

3. The composition of embodiment 1, wherein (a) is a C₁₋₄ alkyl ester of compound (I).

4. The composition of embodiment 2, wherein (a) is a benzyl ester of compound (I).

10 5. The composition of embodiment 1, wherein (a) is the compound of formula (I), which is the carboxylic acid.

6. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is atrazine.

7. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an 15 agriculturally acceptable benzyl ester and (b) is bentazon or bentazon-sodium.

8. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is bromoxynil or bromoxynil octanoate.

9. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is chlorotoluron.

20 10. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is cyanazine.

11. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is diuron.

12. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an 25 agriculturally acceptable benzyl ester and (b) is hexazinone.

13. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is ioxynil.

14. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is isoproturon.
15. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is linuron.
- 5 16. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is methibenzuron.
17. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is metribuzin.
18. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an
10 agriculturally acceptable benzyl ester and (b) is propanil.
19. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is pyridate.
20. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is siduron.
- 15 21. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is simazine.
22. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is simetryne.
23. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an
20 agriculturally acceptable benzyl ester and (b) is tebuthiuron.
24. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is terbuthylazine.
25. The composition of any of embodiments 1-24, further comprising a herbicide safener.
26. The composition of embodiment 2, wherein the weight ratio of compound of formula
25 (I) or agriculturally acceptable salt or ester thereof to atrazine or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:2240 to about 6:1, about 1:1120 to about 3:1, about 1:256 to about 1:16, about 6:1, about 1:7, about 1:9, about 1:16, about 1:17.5, about 1:32, about 1:35, about 1:64, about 1:70, about 1:128, about 1:140, about 1:256, and about 1:2240, or within any range defined between any pair of
30 the foregoing values.
27. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to bentazon, bentazon-sodium, or

agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:1120 to about 3:1, about 1:560 to about 2:1, about 1:256 to about 1.4:1, about 1:96 to about 1:3.4, about 3:1, about 1:3.4, about 1:6, about 1:7, about 1:12, about 1:14, about 1:16, about 1:20, about 1:24, about 1:26, about 1:32, about 1:40, about 1:48, about
5 1:52.5, about 1:96, about 1:256, and about 1:1120, or within any range defined between any pair of the foregoing values.

28. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to bromoxynil or agriculturally acceptable salt or ester thereof is selected from the group of ranges of ratios and ratios
10 consisting of: from about 1:280 to about 12:1, about 1:140 to about 6:1, about 1:70 to about 3:1, about 1:35 to about 2:1, about 1:17.5 to about 1:1, about 12:1, about 1:1, about 1:2, about 1:4, about 1:4.4, about 1:8, about 1:12.5, about 1:17.5, and about 1:280, or within any range defined between any pair of the foregoing values.

29. The composition of embodiment 2, wherein the weight ratio of the compound of
15 formula (I) or agriculturally acceptable salt or ester thereof to chlorotoluron or agriculturally acceptable salt or ester thereof is from about 1:1750 to about 1.5:1.

30. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to cyanazine or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of:
20 about 1:2650 to about 6:1, about 1:1325 to about 3:1, about 1:662 to about 2:1, about 1:110 to about 1:5.3, about 6:1, about 1:5.3, about 1:8.6, about 1:13.8, about 1:17.2, about 1:25, about 1:27.5, about 1:34.4, about 1:110, about 1:662, about 1:1325, and about 1:2650, or within any range defined between any pair of the foregoing values.

31. The composition of embodiment 2, wherein the weight ratio of the compound of
25 formula (I) or agriculturally acceptable salt or ester thereof to diuron or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:3600 to about 6:1, about 1:1800 to about 3:1, about 1:900 to about 1:2, about 1:70 to about 1:9, about 6:1, about 1:9, about 1:17.5, about 1:35, about 1:70, about 1:900, about 1:1800, and about 1:3600, or within any range defined between any pair of the foregoing
30 values.

32. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to hexazinone or agriculturally

acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:2500 to about 12:1, about 1:1250 to about 2:1, about 1:625 to about 1:1, about 1:156 to about 1:20, about 12:1, about 1:20, about 1:39, about 1:78, about 1:156, and about 1:2500, or within any range defined between any pair of the foregoing values.

5 33. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to ioxynil or agriculturally acceptable salt or ester thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:200 to about 12:1, about 1:100 to about 1:1, about 1:37.5 to about 1:9, about 12:1, about 1:9.4, about 1:19, about 1:37.5, and about 1:200, or within any range
10 defined between any pair of the foregoing values.

34. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to isoproturon or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:750 to about 12:1, about 1:375 to about 6:1, about 1:600 to about 1:37.5, about 1:300
15 to about 1:75, about 1:750, about 1:600, about 1:375, about 1:300, about 1:150, about 1:75, about 1:37.5, about 1:1, about 6:1, and about 12:1.

35. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to linuron or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of:
20 from about 1:2250 to about 6:1, about 1:1125 to about 3:1, about 1:562 to about 1:1, about 1:39 to about 1:19, about 6:1, about 3:1, about 1:19.4, about 1:38.8, about 1:562, about 1:1125, and about 1:2250, or within any range defined between any pair of the foregoing values.

36. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to methibenzuron or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of:
25 about 1:1400 to about 4:1, about 1:700 to about 2:1, and about 1:350 to about 1:1.

37. The composition embodiment 2 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to metribuzin or agriculturally acceptable
30 salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:2800 to about 6:1, about 1:1400 to about 3:1, about 1:700 to about 2:1, about 1:168 to about 1.3:1, about 1:26 to about 1:5, about 6:1, about 3:1, about 2:1, about 1:3, about 1:5, about

1:6.7, about 1:10, about 1:13, about 1:20, about 1:26, about 1:168, about 1:700, about 1:1400, and about 1:2800, or within any range defined between any pair of the foregoing values.

38. The composition of embodiment 2 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to propanil or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:2800 to about 1:1, about 1:1400 to about 1:4, about 1:767 to about 1:24, about 1:767 to about 1:12, about 1:384 to about 1:48, about 1:1, about 1:4, about 1:12, about 1:24, about 1:48, about 1:96, about 1:173, about 1:192, about 1:210, about 1:384, about 1:560, about 1:767, about 1:1400, and about 1:2800, or within any range defined between any pair of the foregoing values.

39. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to pyridate or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:800 to about 12:1, about 1:800 to about 6:1, about 1:800 to about 3:1, about 1:800 to about 1:1, about 1:400 to about 6:1, about 1:400 to about 3:1, about 1:400 to about 1:1, and about 1:200 to about 1:16.

40. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to siduron or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:4500 to about 2:1, about 1:2250 to about 1:1, about 1:1125 to about 1:5, and about 1:625 to about 1:50.

41. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to simazine or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:4250 to about 2:1, about 1:2125 to about 1:1, about 1:560 to about 1:8, about 1:280 to about 1:17.5, about 1:140 to about 1:17.5, about 2:1, about 1:8, about 1:17.5, about 1:35, about 1:70, about 1:140, about 1:280, about 1:560, about 1:2125, and about 1:4250, or within any range defined between any pair of the foregoing values.

42. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to simetryne or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:1000 to about 12:1, about 1:500 to about 6:1, about 1:250 to about 1:1, about 1:56 to

about 1:3.5, about 12:1, about 6:1, about 1.3:5, about 1:1, about 1:7, about 1:14, about 1:28, about 1:56, about 1:250, about 1:500, and about 1:1000, or within any range defined between any pair of the foregoing values.

43. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to tebuthiuron or agriculturally acceptable salt thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:2240 to about 2:1, about 1:1120 to about 1:1, about 1:840 to about 1:6, about 1:210 to about 1:6, about 2:1, about 1:1, about 1:2, about 1:3, about 1:6, about 1:9, about 1:13, about 1:17.5, about 1:26, about 1:35, about 1:52.5, about 1:70, about 1:105, about 1:140, about 1:210, about 1:840, and about 1:2240, or within any range defined between any pair of the foregoing values.


44. The composition of embodiment 2, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to terbuthylazine or agriculturally acceptable thereof is selected from the group of ranges of ratios and ratios consisting of: about 1:1500 to about 2.4:1, about 1:750 to about 1:1, about 1:375 to about 1:3, about 1:200 to about 1:10, about 1:62.5 to about 1:31, about 2.4:1, about 1:31.3, about 1:62.5, and about 1:1500, or within any range defined between any pair of the foregoing values.

45. The composition of any of embodiments 1-44, further comprising an agriculturally acceptable adjuvant or carrier.

46. The composition of any of embodiments 1-44 which is synergistic as determined by the Colby equation.

47. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation the composition of any of embodiments 1-46.

48. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation a herbicidally effective amount of (a) a compound of the formula (I)



or an agriculturally acceptable salt or ester thereof and (b) atrazine, bentazon, bentazon-sodium, bromoxynil, chlorotoluron, cyanazine, diuron, hexazinone, ioxynil, isoproturon, linuron, methibenzuron, metribuzin, propanil, pyridate, siduron, simazine, simetryne, 5 tebuthiuron or terbuthylazine or agriculturally acceptable salt or ester thereof.

49. The method of embodiment 47 or 48, wherein the method is practiced in at least one member of the group consisting of direct-seeded, water-seeded and transplanted rice, cereals, wheat, barley, oats, rye, sorghum, corn/maize, sugarcane, sunflower, oilseed rape, canola, sugar beet, soybean, cotton, pineapple, pastures, grasslands, rangelands, fallowland, turf, tree 10 and vine orchards, aquatics, industrial vegetation management (IVM) or rights of way (ROW).

50. The method of embodiment 47 or embodiment 48, wherein the undesirable vegetation is immature.

51. The method of embodiment 47 or 48, wherein the (a) and (b) are applied to water.

15 52. The method of embodiment 51, wherein the water is part of a flooded rice paddy.

53. The method of embodiment 47 or 48, wherein the (a) and (b) are applied pre-emergently to the weed or the crop.

54. The method of embodiment 47 or 48, wherein the (a) and (b) are applied post-emergently to the weed or the crop.

20 55. The method of embodiment 47 or 48, wherein the undesirable vegetation is controlled in glyphosate-, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitor-, glufosinate-, glutamine synthetase inhibitor-, dicamba-, phenoxy auxin-, pyridyloxy auxin-, synthetic auxin-, auxin transport inhibitor-, aryloxyphenoxypropionate-, cyclohexanedione-, phenylpyrazoline-, acetyl CoA carboxylase (ACCase) inhibitor-, imidazolinone-, 25 sulfonylurea-, pyrimidinylthiobenzoate-, triazolopyrimidine-, sulfonylaminocarbonyltriazolinone-, acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitors-, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitor-,

phytoene desaturase inhibitor-, carotenoid biosynthesis inhibitor-, protoporphyrinogen oxidase (PPO) inhibitor-, cellulose biosynthesis inhibitor-, mitosis inhibitor-, microtubule inhibitor-, very long chain fatty acid inhibitor-, fatty acid and lipid biosynthesis inhibitor-, photosystem I inhibitor-, photosystem II inhibitor-, triazine-, or bromoxynil- tolerant crops.

5 56. The method of embodiment 55, wherein the tolerant crop possesses multiple or stacked traits conferring resistance or tolerance to multiple herbicides or multiple modes of action.

57. The method of embodiment 47 or 48, wherein the undesirable vegetation comprises a herbicide resistant or tolerant weed.

10 58. The method of embodiment 57, wherein the resistant or tolerant weed is a biotype with resistance or tolerance to multiple herbicides, multiple chemical classes, multiple herbicide modes-of-action or via multiple resistance mechanisms.

59. The method of embodiment 57, wherein the resistant or tolerant weed is a biotype resistant or tolerant to acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS)
 15 inhibitors, photosystem II inhibitors, acetyl CoA carboxylase (ACCase) inhibitors, synthetic auxins, auxin transport inhibitors, photosystem I inhibitors, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, microtubule assembly inhibitors, fatty acid and lipid synthesis inhibitors, protoporphyrinogen oxidase (PPO) inhibitors, carotenoid biosynthesis inhibitors, very long chain fatty acid (VLCFA) inhibitors, phytoene desaturase (PDS)
 20 inhibitors, glutamine synthetase inhibitors, 4-hydroxyphenyl-pyruvate-dioxygenase (HPPD) inhibitors, mitosis inhibitors, cellulose biosynthesis inhibitors, herbicides with multiple modes-of-action, quinclorac, arylaminopropionic acids, difenzoquat, endothall, or organoarsenicals.

60. A methods of controlling undesirable vegetation including the step of applying a
 25 herbicidally effective amount of at least one mixture according to the twenty-sixth embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of atrazine selected from the group of rates and ranges of rates consisting of, about: 50, 140, 280, 560, 1120, 2240, and 4400 or within any range defined between any pair of the foregoing values.

61. Methods according to either of the twenty-sixth or sixtieth embodiments wherein the controlled plant is at least one plant selected from the group consisting of: BRAPP, CYPPIR, ECHCG, ECHCO, and XANST, still other embodiments include controlling plants from the genera consisting of: *Brachiaria* or *Urochloa*, *Echinochloa*, *Bolboschoenus*, and *Xanthium*.

5 62. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the twenty-seventh embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of bentazon or bentazon-sodium selected from the group of rates and ranges of rates consisting of, about: 60, 105, 120, 210, 420, 840, 1120, and 2240, or within any range
10 defined between any pair of the foregoing values.

63. Methods according to either of the twenty-seventh or sixty-second embodiments wherein the controlled plant is at least one plant selected from the group consisting of: BRSNW, COMBE, ECHCG, ECHOR, IPOHE, LEFCH, SCPMA, and SETFA, still other
15 embodiments include controlling plants from the genera consisting of: *Brassica*, *Commelina*, *Echinochloa*, *Ipomoea*, *Leptochloa*, *Bolboschoenus* or *Schoenoplectus*, and *Xanthium*.

64. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the twenty-eighth
embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of bromoxynil selected from the group of rates and ranges of rates consisting of,
20 about: 25, 35, 70, 140, 280, or 560 within any range defined between any pair of the foregoing values.

65. Methods according to either of the twenty-eighth or sixty-fourth embodiments wherein the controlled plant is at least one plant selected from the group consisting of:

AMARE, ECHCG, IPOHE, and SETFA, still other embodiments include controlling plants from the genera consisting of: *Amaranthus*, *Echinochloa*, *Ipomoea*, and *Setaria*.

66. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the twenty-ninth embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of chlorotoluron selected from the group of rates and ranges of rates consisting of, about: 200, 400, 800, 1600, or 3500 within any range defined between any pair of the foregoing values.

67. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the thirtieth embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of cyanazine selected from the group of rates and ranges of rates consisting of, about: 50, 170, 275, 440, 880, 1325, 2650, or 5300 within any range defined between any pair of the foregoing values.

68. Methods according to either of the thirtieth or sixty-seventh embodiments wherein the controlled plant is at least one plant selected from the group consisting of: BRAPP and IPOHE, still other embodiments include controlling plants from the genera consisting of: *Brachiaria* or *Urochloa* and *Ipomoea*.

69. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the thirty-first embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of diuron selected from the group of rates and ranges of rates consisting of, about: 50, 100, 140, 280, 560, 1120, 2240, 4480, or 7200 within any range defined between any pair of the foregoing values.

70. Methods according to either of the thirty-first or sixty-ninth embodiments wherein the controlled plant is at least one plant selected from the group consisting of: ABUTH, BRAPP, ECHCG, ECHCO, SETFA, and SETVI, still other embodiments include controlling plants from the genera consisting of: *Abutilon*, *Brachiaria* or *Urochloa*, *Echinochloa*, and *Setaria*.

5 71. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the thirty-second embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of hexazinone selected from the group of rates and ranges of rates consisting of, about: 25, 50, 100, 200, 400, 625, 1250, 2500, or 5000 within any range defined between any pair of
10 the foregoing values.

72. Methods according to either of the thirty-second or seventy-first embodiments wherein the controlled plant is at least one plant selected from the group consisting of: ECHCG, ECHCO, and IPOHE, still other embodiments include controlling plants from the genera consisting of: *Echinochloa* and *Ipomoea*.

15 73. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the thirty-third embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of ioxynil selected from the group of rates and ranges of rates consisting of, about: 25, 50, 100, 200, 300, or 400 within any range defined between any pair of the foregoing values.

20 74. Methods according to either of the thirty-third or seventy-third embodiments wherein the controlled plant is at least one plant selected from the group consisting of: IPOHE, still other embodiments include controlling plants from the genera consisting of: *Ipomoea*.

75. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the thirty-fourth

embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of isoproturon selected from the group of rates and ranges of rates consisting of, about: 25, 50, 100, 200, 400, 800, or 1500 within any range defined between any pair of the foregoing values.

- 5 76. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the thirty-fifth embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of linuron selected from the group of rates and ranges of rates consisting of, about: 50, 100, 200, 400, 620, 1125, 2250, or 4500 within any range defined between any pair of the
10 foregoing values.

77. Methods according to either of the thirty-fifth or seventy-sixth embodiments wherein the controlled plant is at least one plant selected from the group consisting of: BRAPP, still other embodiments include controlling plants from the genera consisting of: *Brachiaria* or *Urochloa*.

- 15 78. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the thirty-sixth embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of methibenzuron selected from the group of rates and ranges of rates consisting of, about: 35, 70, 140, 280, 560, 700, 1400, or 2800 within any range defined between any pair of the
20 foregoing values.

79. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the thirty-seventh embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of metribuzin selected from the group of rates and ranges of rates consisting of, about:

25, 50, 100, 210, 420, 840, 1680, 3360, or 5600 within any range defined between any pair of the foregoing values.

80. Methods according to either of the thirty-seventh or seventy-ninth embodiments wherein the controlled plant is at least one plant selected from the group consisting of:

5 CHEAL, KCHSC, LAMPU, SASKR, SETFA, SINAR, still other embodiments include controlling plants from the genera consisting of: *Chenopodium*, *Kochia*, *Lamium*, *Salsola*, *Setaria*, and *Sinapsis*.

81. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the thirty-eighth

10 embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of propanil selected from the group of rates and ranges of rates consisting of, about: 60, 120, 420, 840, 1680, 3360, or 5600 within any range defined between any pair of the foregoing values.

82. Methods according to either of the thirty-eighth or eighty-first embodiments wherein
15 the controlled plant is at least one plant selected from the group consisting of: COMBE, ECHCG, ECHCO, IPOHE, and SCPMA, still other embodiments include controlling plants from the genera consisting of: *Commelina*, *Echinochloa*, *Ipomoea*, and *Bolboschoenus* or *Schoenoplectus*.

83. Methods of controlling undesirable vegetation comprising the step of applying a

20 herbicidally effective amount of at least one mixture according to the thirty-ninth embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of pyridate selected from the group of rates and ranges of rates consisting of, about: 25, 50, 100, 200, 400, 800, or 1600 within any range defined between any pair of the foregoing values.

84. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the fortieth embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of siduron selected from the group of rates and ranges of rates consisting of, about: 70, 140,
5 280, 560, 1120, 2240, 4480, or 9000 within any range defined between any pair of the foregoing values.

85. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the forty-first embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of
10 simazine selected from the group of rates and ranges of rates consisting of, about: 70, 140, 280, 560, 1120, 2240, 4480, or 8500 within any range defined between any pair of the foregoing values.

86. Methods according to either of the forty-first or eighty-fifth embodiments wherein the controlled plant is at least one plant selected from the group consisting of: BRAPP, ECHCG,
15 ECHCO, CYPPIR, and XANST, still other embodiments include controlling plants from the genera consisting of: *Brachiaria* or *Urochloa*, *Echinochloa*, *Cyperus*, and *Xanthium*.

87. Methods of controlling undesirable vegetation comprising the step of applying a herbicidally effective amount of at least one mixture according to the forty-second embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or
20 gae/ha of simetryne selected from the group of rates and ranges of rates consisting of, about: 25, 50, 112.5, 225, 450, 900, 1350, and 2000 within any range defined between any pair of the foregoing values.

88. Methods according to either of the forty-second or eighty-seventh embodiments wherein the controlled plant is at least one plant selected from the group consisting of:

CYPIR, CYPRO, ECHCO, ECHOR, FIMMI, and LEFCH, still other embodiments include controlling plants from the genera consisting of: *Cyperus*, *Echinochloa*, *Fimbristylis*, and *Leptochloa*.

89. Methods of controlling undesirable vegetation comprising the step of applying a
5 herbicidally effective amount of at least one mixture according to the forty-third embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of tebuthiuron selected from the group of rates and ranges of rates consisting of, about: 70, 140, 210, 280, 420, 560, 840, 1680, 3360, and 4480 within any range defined between any pair of the foregoing values.

10 90. Methods according to either of the forty-third or eighty-ninth embodiments wherein the controlled plant is at least one plant selected from the group consisting of: BRSNN, CIRAR, CYPES, IPOHE, and XANST, still other embodiments include controlling plants from the genera consisting of: *Brassica*, *Cirsium*, *Cyperus*, *Ipomoea*, and *Xanthium*.

91. Methods of controlling undesirable vegetation comprising the step of applying a
15 herbicidally effective amount of at least one mixture according to the forty-fourth embodiment wherein the amount of the mixture is applied at a rate, expressed in gai/ha or gae/ha of terbuthylazine selected from the group of rates and ranges of rates consisting of, about: 60, 125, 250, 500, 1000, 1500, and 3000 within any range defined between any pair of the foregoing values.

20 93. Methods according to either of the forty-fourth or ninety-first embodiments wherein the controlled plant is at least one plant selected from the group consisting of: XANST, still other embodiments include controlling plants from the genera consisting of: *Xanthium*.

94. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is atrazine.

95. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is bentazon-sodium.
96. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is bromoxynil.
- 5 97. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is chlorotoluron.
98. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is cyanazine.
99. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an
10 agriculturally acceptable benzyl ester and (b) is diuron.
100. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is hexazinone.
101. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is ioxynil.
- 15 102. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is isoproturon.
103. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is linuron.
104. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an
20 agriculturally acceptable benzyl ester and (b) is methibenzuron.
105. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is metribuzin.
106. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is propanil.

107. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is pyridate.
108. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is siduron.
- 5 109. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is simazine.
110. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is simetryne.
111. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an
10 agriculturally acceptable benzyl ester and (b) is tebuthiuron.
112. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is terbuthylazine.
113. The composition of embodiment 1, wherein the weight ratio of compound of formula (I) or agriculturally acceptable salt or ester thereof to atrazine or agriculturally acceptable salt
15 or ester thereof is from about 1:2240 to about 6:1.
114. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to bentazon or agriculturally acceptable salt thereof is from about 1:1120 to about 3:1.
115. The composition of embodiment 1, wherein the weight ratio of the compound of
20 formula (I) or agriculturally acceptable salt or ester thereof to bromoxynil or agriculturally acceptable salt thereof is from about 1:280 to about 12:1.
116. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to chlorotoluron or agriculturally acceptable salt or ester thereof is from about 1:1750 to about 1.5:1.

117. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to cyanazine or agriculturally acceptable salt thereof is from about 1:2650 to about 6:1.

118. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to diuron or agriculturally acceptable thereof is from about 1:3600 to about 6:1.

119. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to hexazinone or agriculturally acceptable salt thereof is from about 1:2500 to about 12:1.

120. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to ioxynil or agriculturally acceptable salt thereof is from about 1:200 to about 12:1.

121. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to isoproturon or agriculturally acceptable salt thereof is from about 1:750 to about 12:1.

122. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to linuron or agriculturally acceptable salt thereof is from about 1:2250 to about 6:1.

123. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to methibenzuron or agriculturally acceptable salt thereof is from about 1:1400 to about 4:1.

124. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to metribuzin or agriculturally acceptable salt thereof is from about 1:2800 to about 6:1.

125. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to propanil or agriculturally acceptable salt thereof is from about 1:2800 to about 1:1.

126. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to pyridate or agriculturally acceptable salt thereof is from about 1:800 to about 12:1.

127. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to siduron or agriculturally acceptable salt thereof is from about 1:4500 to about 2:1.

128. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to simazine or agriculturally acceptable salt thereof is from about 1:4250 to about 2:1.


129. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to simetryne or agriculturally acceptable salt thereof is from about 1:1000 to about 12:1.

130. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to tebuthiuron or agriculturally acceptable salt thereof is from about 1:2240 to about 2:1.

131. The composition of embodiment 1, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to terbuthylazine or agriculturally acceptable thereof is from about 1:1500 to about 3:1.

132. The composition of any of embodiments 94-131 which is synergistic as determined by the Colby equation.

Provided herein are herbicidal compositions comprising and methods of controlling undesirable vegetation utilizing a herbicidally effective amount of (a) a compound of the formula (I)




- 5 or an agriculturally acceptable salt or ester of thereof, and (b) a PS II inhibitor selected from the group consisting of: atrazine, bentazon, bentazon-sodium, bromoxynil, bromoxynil octanoate, chlorotoluron, cyanazine, diuron, hexazinone, ioxynil, isoproturon, linuron, methibenzuron, metribuzin, propanil, pyridate, siduron, simazine, simetryne, tebuthiuron and terbuthylazine, or derivative, *e.g.*, salt or ester thereof. The compositions
- 10 may also contain an agriculturally acceptable adjuvant or carrier.

Detailed Description


DEFINITIONS

As used herein, the compound of formula (I) has the following structure:




- 15 The compound of formula (I) can be identified by the name 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropyridine-2-carboxylic acid and has been described in U.S. Patent 7,314,849 (B2), which is incorporated herein by reference in its entirety. Exemplary uses of the compound of the formula (I) include controlling undesirable vegetation, including grass, broadleaf and sedge weeds, in multiple non-crop and cropping situations.
- 20 Atrazine is 6-chloro-*N*-ethyl-*N'*-(1-methylethyl)-1,3,5-triazine-2,4-diamine. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of atrazine include its use for pre-emergence and post-emergence control of broadleaf

and grass weeds, *e.g.*, in corn, sorghum, turf, sugar cane and other crops. Atrazine possesses the following structure:




5 Bentazon is 3-(1-methylethyl)-1*H*-2,1,3-benzothiadiazin-4(3*H*)-one 2,2-dioxide. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. It is also known as bentazone. Bentazon has been used as its sodium salt, bentazon sodium. Exemplary uses of bentazon include its use for control of broadleaf and sedge weeds in broadleaf and grass crops. Bentazon possesses the following structure:



10


Bromoxynil is 3,5-dibromo-4-hydroxybenzonitrile. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Bromoxynil has been used as its octanoate ester, bromoxynil octanoate. Exemplary uses of bromoxynil include its use for post-emergence control of broad-leaved weeds, *e.g.*, in cereals, ryegrass-seed crops, turf, maize, and sorghum. It possesses the following structure:

15




Chlorotoluron is *N'*-(3-chloro-4-methylphenyl)-*N,N*-dimethylurea. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of chlorotoluron include its use for control of broad-leaved and grass weeds, *e.g.*, in winter cereals. It possesses the following structure:


20




Cyanazine is 2-[[4-chloro-6-(ethylamino)-1,3,5-triazin-2-yl]amino]-2-methylpropanenitrile. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of cyanazine include its use for pre-emergent control of weeds in beans, maize and peas and post-emergent control of weeds in early barley and wheat. It possesses the following structure:




Diuron is *N'*-(3,4-dichlorophenyl)-*N,N*-dimethylurea. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of diuron include its use for pre-emergence and post-emergence control of broadleaf and grass weeds in, *e.g.*, corn, sorghum, sugar cane, citrus and other crops. Diuron possesses the following structure:




Hexazinone is 3-cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4(1*H*,3*H*)-dione. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of hexazinone include its use for post-emergent control of annual, biennial and perennial weeds in alfalfa, pineapples, sugar cane and conifers. It possesses the following structure:




Ioxynil is 4-hydroxy-3,5-diiodobenzonitrile. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of ioxynil include its use for post-emergence control of annual broad-leaved weeds in cereals, onions, leeks, garlic, shallots, flax, sugar cane, forage grasses, lawns and newly-sown turf. It possesses the following structure:




Isoproturon is *N,N*-dimethyl-*N'*-[4-(1-methylethyl)phenyl]urea. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of isoproturon include its use for pre- and post-emergence control of annual grasses and annual
 5 broad-leaved weeds, *e.g.*, in spring and winter wheat, spring and winter barley, winter rye and triticale. It possesses the following structure:



Linuron is *N'*-(3,4-dichlorophenyl)-*N*-methoxy-*N*-methylurea. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of linuron
 10 include its use for pre- and post-emergence control of annual grass and broad-leaved weeds, as well as seedling perennial weeds in asparagus, artichokes, carrots, parsley, fennel, parsnips, herbs and spices, celery, celeriac, onions, leeks, garlic, potatoes, peas, field beans, soya beans, cereals, maize, sorghum, cotton, flax, sunflowers, sugar cane, ornamentals, established vines, bananas, cassava, coffee, tea, rice, peanuts, ornamental trees and shrubs,
 15 and other crops. It possesses the following structure:




Methibenzuron is *N*-2-benzothiazolyl-*N,N'*-dimethylurea. Methibenzuron is also known as methabenzthiazuron. The herbicidal activity methibenzuron is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of methibenzuron include its use
 20 for broad-spectrum control of broad-leaved weeds and grasses, *e.g.*, in cereals, legumes, maize, garlic and onions. It possesses the following structure:




Metribuzin is 4-amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009.


Exemplary uses of metribuzin include its use for pre-emergence and post-emergence control of broadleaf weeds, *e.g.*, in soybeans, potatoes, corn, vegetables and other crops. Metribuzin possesses the following structure:




Propanil is *N*-(3,4-dichlorophenyl)propanamide. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of propanil include its use for post-emergence control of broadleaf and grass weeds in, *e.g.*, rice. It possesses the following structure:




Pyridate is *O*-(6-chloro-3-phenyl-4-pyridazinyl) *S*-octyl carbonothioate. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of pyridate include its use for post-emergence control of annual broad-leaved weeds and grass weeds, *e.g.*, in maize, sweet corn, oilseed rape, cereals, rice, peanuts and vegetables. It possesses the following structure:




Siduron is *N*-(2-methylcyclohexyl)-*N'*-phenylurea. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of siduron include its use for pre-emergence control of *Digitaria* spp. and annual grass weeds in, *e.g.*, turf farms, grass seed production and established turf. It possesses the following structure:




Simazine is 6-chloro-*N,N'*-diethyl-1,3,5-triazine-2,4-diamine. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of simazine include its use for control of germinating annual grasses and broad-leaved weeds in, *e.g.*, pome fruit, stone fruit, bush and cane fruit, citrus fruit, vines, strawberries, nuts, olives,
 5 pineapples, field beans, French beans, pea, maize, sweet corn, asparagus, hops, alfalfa, lupins, oilseed rape, artichokes, sugar cane, cocoa, coffee, rubber, oil palms, tea, turf and ornamentals. It possesses the following structure:



Simetryne is *N,N'*-diethyl-6-(methylthio)-1,3,5-triazine-2,4-diamine. Simetryne is
 10 simetryn. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of simetryn include its use for control of broad-leaved weeds, *e.g.*, in rice. It possesses the following structure:




Tebuthiuron is *N*-[5-(1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]-*N,N'*-dimethylurea. Its
 15 herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of tebuthiuron include its use for control of herbaceous and woody plants, annual weeds, and many perennial grass and undersirable woody plants in pastures and rangeland, and control of grass and broadleaved weeds in sugar cane. It possesses the following structure:



Terbuthylazine is 6-chloro-*N*-(1,1-dimethylethyl)-*N'*-ethyl-1,3,5-triazine-2,4-diamine. Its herbicidal activity is described in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of terbuthylazine include its use for pre and post-emergence weed control, *e.g.*, in maize, sorghum, vines, fruit trees, citrus, coffee, oil palm, cocoa, olives, potatoes,

peas, beans, sugarcane, rubber, and forestry (tree nurseries and new plantings). It possesses the following structure:



As used herein, herbicide means a compound, *e.g.*, active ingredient that kills,
5 controls or otherwise adversely modifies the growth of plants.

As used herein, a herbicidally effective or vegetation controlling amount is an amount of active ingredient which causes an adversely modifying effect to the vegetation *e.g.*, causing deviations from natural development, killing, effecting regulation, causing desiccation, causing retardation, and the like.


10 As used herein, controlling undesirable vegetation means preventing, reducing, killing, or otherwise adversely modifying the development of plants and vegetation.

Described herein are methods of controlling undesirable vegetation through the application of certain herbicide combinations or compositions. Methods of application include, but are not limited to applications to the vegetation or locus thereof, *e.g.*, application to the area adjacent
15 to the vegetation, as well as pre-emergence, post-emergence, foliar (broadcast, directed, banded, spot, mechanical, over-the-top, or rescue), and in-water applications (emerged and submerged vegetation, broadcast, spot, mechanical, water-injected, granular broadcast, granular spot, shaker bottle, or stream spray) via hand, backpack, machine, tractor, or aerial (airplane and helicopter) application methods.

20 As used herein, plants and vegetation include, but are not limited to, germinant seeds, emerging seedlings, plants emerging from vegetative propagules, immature vegetation, and established vegetation.

As used herein, agriculturally acceptable salts and esters refer to salts and esters that exhibit herbicidal activity, or that are or can be converted in plants, water, or soil to the
25 referenced herbicide. Exemplary agriculturally acceptable esters are those that are or can be hydrolyzed, oxidized, metabolized, or otherwise converted, *e.g.*, in plants, water, or soil, to the corresponding carboxylic acid which, depending on the pH, may be in the dissociated or undissociated form.

Exemplary salts include those derived from alkali or alkaline earth metals and those derived from ammonia and amines. Exemplary cations include sodium, potassium, magnesium, and aminium cations of the formula:




5 wherein R^1 , R^2 , R^3 and R^4 each, independently represents hydrogen or C_1 - C_{12} alkyl, C_3 - C_{12} alkenyl or C_3 - C_{12} alkynyl, each of which is optionally substituted by one or more hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio or phenyl groups, provided that R^1 , R^2 , R^3 and R^4 are sterically compatible. Additionally, any two of R^1 , R^2 , R^3 and R^4 together may represent an aliphatic difunctional moiety containing one to twelve carbon atoms and up to two oxygen or sulfur
10 atoms. Salts can be prepared by treatment with a metal hydroxide, such as sodium hydroxide, with an amine, such as ammonia, trimethylamine, diethanolamine, 2-methylthiopropylamine, bisallylamine, 2-butoxyethylamine, morpholine, cyclododecylamine, or benzylamine or with a tetraalkylammonium hydroxide, such as tetramethylammonium hydroxide or choline hydroxide.

15 Exemplary esters include those derived from C_1 - C_{12} alkyl, C_3 - C_{12} alkenyl, C_3 - C_{12} alkynyl or C_7 - C_{10} aryl-substituted alkyl alcohols, such as methyl alcohol, isopropyl alcohol, 1-butanol, 2-ethylhexanol, butoxyethanol, methoxypropanol, allyl alcohol, propargyl alcohol, cyclohexanol or unsubstituted or substituted benzyl alcohols. Benzyl alcohols may be substituted with from 1-3 substituents independently selected from halogen, C_1 - C_4 alkyl or
20 C_1 - C_4 alkoxy. Esters can be prepared by coupling of the acids with the alcohol using any number of suitable activating agents such as those used for peptide couplings such as dicyclohexylcarbodiimide (DCC) or carbonyl diimidazole (CDI); by reacting the acids with alkylating agents such as alkylhalides or alkylsulfonates in the presence of a base such as triethylamine or lithium carbonate; by reacting the corresponding acid chloride of an acid
25 with an appropriate alcohol; by reacting the corresponding acid with an appropriate alcohol in the presence of an acid catalyst or by transesterification.

COMPOSITIONS AND METHODS

Provided herein are herbicidal compositions comprising a herbicidally effective amount of (a) a compound of the formula (I)



or an agriculturally acceptable salt or ester of thereof, and (b) a PS II inhibitor selected from the group consisting of atrazine, bentazon, bentazon-sodium, bromoxynil, bromoxynil octanoate, chlorotoluron, cyanazine, diuron, hexazinone, ioxynil, isoproturon, 5 linuron, methibenzuron, metribuzin, propanil, pyridate, siduron, simazine, simetryne, tebuthiuron and terbuthylazine, or a salt or ester thereof.

Provided herein are also methods of controlling undesirable vegetation comprising contacting the vegetation or the locus thereof, *i.e.*, the area adjacent to the undesirable vegetation, with or applying to the soil or water to prevent the emergence or growth of 10 vegetation a herbicidally effective amount of the compound of formula (I) or agriculturally acceptable salt or ester thereof and (b) a PS II inhibitor selected from the group consisting of atrazine, bentazon, bentazon-sodium, bromoxynil, bromoxynil octanoate, chlorotoluron, cyanazine, diuron, hexazinone, ioxynil, isoproturon, linuron, methibenzuron, metribuzin, propanil, pyridate, siduron, simazine, simetryne, tebuthiuron and terbuthylazine, or a salt or 15 ester thereof. In certain embodiments, the methods employ the compositions described herein.

Furthermore, in some embodiments, the combination of compound (I) or agriculturally acceptable salt or ester thereof and a PS II inhibitor or an agriculturally acceptable salt or ester thereof exhibits synergism, *e.g.*, the herbicidal active ingredients are 20 more effective in combination than when applied individually. The *Herbicide Handbook* of the Weed Science Society of America, Ninth Edition, 2007, p. 429 notes that “‘synergism’ [is] an interaction of two or more factors such that the effect when combined is greater than the predicted effect based on the response of each factor applied separately.” In certain embodiments, the compositions exhibit synergy as determined by the Colby’s equation. 25 Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.

In certain embodiments of the compositions and methods described herein, the compound of formula (I), *i.e.*, the carboxylic acid, is employed. In certain embodiments, a carboxylate salt of the compound of formula (I) is employed. In certain embodiments, an aralkyl or alkyl ester is employed. In certain embodiments, a benzyl, substituted benzyl, or C₁₋₄ alkyl, *e.g.*, n-butyl ester is employed. In certain embodiments, the benzyl ester is employed.

In some embodiments, the compound of formula (I) or salt or ester thereof and a PS II inhibitor, or an agriculturally acceptable salt or ester thereof are formulated in one composition, tank mixed, applied simultaneously, or applied sequentially.

Herbicidal activity is exhibited by the compounds when they are applied directly to the plant or to the locus of the plant at any stage of growth. The effect observed depends upon the plant species to be controlled, the stage of growth of the plant, the application parameters of dilution and spray drop size, the particle size of solid components, the environmental conditions at the time of use, the specific compound employed, the specific adjuvants and carriers employed, the soil type, and the like, as well as the amount of chemical applied. These and other factors can be adjusted to promote non-selective or selective herbicidal action. In some embodiments, the compositions described herein are applied as a post-emergence application, pre-emergence application, or in-water application to flooded paddy rice or water bodies (*e.g.*, ponds, lakes and streams), to relatively immature undesirable vegetation to achieve the maximum control of weeds.

In some embodiments, the compositions and methods provided herein are utilized to control weeds in crops, including but not limited to direct-seeded, water-seeded and transplanted rice, cereals, wheat, barley, oats, rye, sorghum, corn/maize, sugarcane, sunflower, oilseed rape, canola, sugarbeet, soybean, cotton, pineapple, pastures, grasslands, rangelands, fallowland, turf, tree and vine orchards, aquatics, plantation crops, vegetables, industrial vegetation management (IVM) and rights-of-way (ROW).

In certain embodiments, the compositions and methods provided herein are utilized to control weeds in rice. In certain embodiments, the rice is direct-seeded, water-seeded, or transplanted rice.

The compositions and methods described herein may be used to control undesirable vegetation on glyphosate-tolerant-, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitor-tolerant-, glufosinate-tolerant-, glutamine synthetase inhibitor-tolerant-, dicamba-

tolerant-, phenoxy auxin-tolerant-, pyridyloxy auxin-tolerant-, auxin-tolerant-, auxin transport inhibitor-tolerant-, aryloxyphenoxypropionate-tolerant-, cyclohexanedione-tolerant-, phenylpyrazoline-tolerant-, acetyl CoA carboxylase (ACCase) inhibitor-tolerant-, imidazolinone-tolerant-, sulfonyleurea-tolerant-, pyrimidinylthiobenzoate-tolerant-,
5 triazolopyrimidine-tolerant-, sulfonyleaminocarbonyltriazolinone-tolerant-, acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitor-tolerant-, 4-hydroxyphenylpyruvate dioxygenase (HPPD) inhibitor -tolerant-, phytoene desaturase inhibitor-tolerant-, carotenoid biosynthesis inhibitor-tolerant-, protoporphyrinogen oxidase (PPO) inhibitor -tolerant-, cellulose biosynthesis inhibitor-tolerant-, mitosis inhibitor-tolerant-, microtubule
10 inhibitor-tolerant-, very long chain fatty acid inhibitor-tolerant-, fatty acid and lipid biosynthesis inhibitor-tolerant-, photosystem I inhibitor-tolerant-, photosystem II inhibitor-tolerant-, triazine-tolerant-, and bromoxynil-tolerant- crops (such as, but not limited to, soybean, cotton, canola/oilseed rape, rice, cereals, corn, sorghum, sunflower, sugar beet, sugarcane, turf, etc.), for example, in conjunction with glyphosate, EPSP synthase inhibitors,
15 glufosinate, glutamine synthase inhibitors, dicamba, phenoxy auxins, pyridyloxy auxins, synthetic auxins, auxin transport inhibitors, aryloxyphenoxypropionates, ACCase inhibitors, cyclohexanediones, phenylpyrazolines, imidazolinones, sulfonyleureas, pyrimidinylthiobenzoates, triazolopyrimidines, sulfonyleaminocarbonyltriazolinones, ALS or AHAS inhibitors, HPPD inhibitors, phytoene desaturase inhibitors, carotenoid biosynthesis
20 inhibitors, PPO inhibitors, cellulose biosynthesis inhibitors, mitosis inhibitors, microtubule inhibitors, very long chain fatty acid inhibitors, fatty acid and lipid biosynthesis inhibitors, photosystem I inhibitors, photosystem II inhibitors, triazines, and bromoxynil. The compositions and methods may be used in controlling undesirable vegetation in crops possessing multiple or stacked traits conferring tolerance to multiple chemistries and/or
25 inhibitors of multiple modes of action.

In some embodiments, the compound of formula (I) or salt or ester thereof and complementary herbicide or salt or ester thereof are used in combination with herbicides that are selective for the crop being treated and which complement the spectrum of weeds controlled by these compounds at the application rate employed. In some embodiments, the
30 compositions described herein and other complementary herbicides are applied at the same time, either as a combination formulation or as a tank mix, or sequentially.

The compositions and methods may be used in controlling undesirable vegetation in crops possessing agronomic stress tolerance (including but not limited to drought, cold, heat, salt, water, nutrient, fertility, pH), pest tolerance (including but not limited to insects, fungi and pathogens) and crop improvement traits (including but not limited to yield; protein, carbohydrate, or oil content; protein, carbohydrate, or oil composition; plant stature and plant architecture).

The compositions and methods provided herein are utilized to control undesirable vegetation. Undesirable vegetation includes, but is not limited to, undesirable vegetation that occurs in rice, cereals, wheat, barley, oats, rye, sorghum, corn/maize, sugarcane, sunflower, oilseed rape, canola, sugar beet, soybean, cotton, pineapple, pastures, grasslands, rangelands, fallowland, turf, tree and vine orchards, aquatics, plantation crops, vegetables, industrial vegetation management (IVM) and rights of way (ROW).

In some embodiments, the methods provided herein are utilized to control undesirable vegetation in rice. In certain embodiments, the undesirable vegetation is *Brachiaria platyphylla* (Groseb.) Nash or *Urochloa platyphylla* (Nash) R.D. Webster (broadleaf signalgrass, BRAPP), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa* species (ECHSS), *Echinochloa crus-galli* (L.) P. Beauv. (barnyardgrass, ECHCG), *Echinochloa crus-pavonis* (Kunth) Schult. (gulf cockspur, ECHCV), *Echinochloa colonum* (L.) LINK (junglerice, ECHCO), *Echinochloa oryzoides* (Ard.) Fritsch (early watergrass, ECHOR), *Echinochloa oryzicola* (Vasinger) Vasinger (late watergrass, ECHPH), *Echinochloa phyllopogon* (Stapf) Koso-Pol. (rice barnyardgrass, ECHPH), *Echinochloa polystachya* (Kunth) Hitchc. (creeping river grass, ECHPO), *Ischaemum rugosum* Salisb. (saramollagrass, ISCRU), *Leptochloa chinensis* (L.) Nees (Chinese sprangletop, LEFCH), *Leptochloa fascicularis* (Lam.) Gray (bearded sprangletop, LEFFA), *Leptochloa panicoides* (Presl.) Hitchc. (Amazon sprangletop, LEFPA), *Oryza* species (red and weedy rice, ORYSS), *Panicum dichotomiflorum* (L.) Michx. (fall panicum, PANDI), *Paspalum dilatatum* Poir. (dallisgrass, PASDI), *Rottboellia cochinchinensis* (Lour.) W.D. Clayton (itchgrass, ROOEX), *Cyperus* species (CYPSS), *Cyperus difformis* L. (smallflower flatsedge, CYPDI), *Cyperus dubius* Rottb. (MAPDU), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus iria* L. (rice flatsedge, CYPPIR), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Cyperus serotinus* Rottb./C.B. Clarke (tidalmarsh flatsedge, CYPSE), *Eleocharis* species (ELOSS), *Fimbristylis miliacea* (L.) Vahl (globe fringerush, FIMMI), *Schoenoplectus* species (SCPSS),

- Schoenoplectus juncooides* Roxb. (Japanese bulrush, SCPJU), *Bolboschoenus maritimus* (L.) Palla or *Schoenoplectus maritimus* L. Lye (sea clubrush, SCPMA), *Schoenoplectus mucronatus* L. (ricefield bulrush, SCPMU), *Aeschynomene* species, (jointvetch, AESSS), *Alternanthera philoxeroides* (Mart.) Griseb. (alligatorweed, ALRPH), *Alisma plantago-*
- 5 *aquatica* L. (common waterplantain, ALSPA), *Amaranthus* species, (pigweeds and amaranths, AMASS), *Ammannia coccinea* Rottb. (redstem, AMMCO), *Commelina benghalensis* L. (Benghal dayflower, COMBE), *Eclipta alba* (L.) Hassk. (American false daisy, ECLAL), *Heteranthera limosa* (SW.) Willd./Vahl (ducksalad, HETLI), *Heteranthera reniformis* R. & P. (roundleaf mudplantain, HETRE), *Ipomoea* species (morningglories,
- 10 IPOSS), *Ipomoea hederacea* (L.) Jacq. (ivyleaf morningglory, IPOHE), *Lindernia dubia* (L.) Pennell (low false pimpernel, LIDDU), *Ludwigia* species (LUDSS), *Ludwigia linifolia* Poir. (southeastern primrose-willow, LUDLI), *Ludwigia octovalvis* (Jacq.) Raven (longfruited primrose-willow, LUDOC), *Monochoria korsakowii* Regel & Maack (monochoria, MOOKA), *Monochoria vaginalis* (Burm. F.) C. Presl ex Kuhth, (monochoria, MOOVA),
- 15 *Murdannia nudiflora* (L.) Brenan (doveweed, MUDNU), *Polygonum pensylvanicum* L., (Pennsylvania smartweed, POLPY), *Polygonum persicaria* L. (ladysthumb, POLPE), *Polygonum hydropiperoides* Michx. (POLHP, mild smartweed), *Rotala indica* (Willd.) Koehne (Indian toothcup, ROTIN), *Sagittaria* species, (arrowhead, SAGSS), *Sesbania exaltata* (Raf.) Cory/Rydb. Ex Hill (hemp sesbania, SEBEX), or *Sphenoclea zeylanica*
- 20 Gaertn. (gooseweed, SPDZE).

In some embodiments, the methods provided herein are utilized to control undesirable vegetation in cereals. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Apera spica-venti* (L.) Beauv. (windgrass, APESV), *Avena fatua* L. (wild oat, AVEFA), *Bromus tectorum* L. (downy brome, BROTE),

25 *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Phalaris minor* Retz. (littleseed canarygrass, PHAMI), *Poa annua* L. (annual bluegrass, POANN), *Setaria pumila* (Poir.) Roemer & J.A. Schultes (yellow foxtail, SETLU), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Amaranthus retroflexus* L. (redroot pigweed, AMARE), *Brassica* species (BRSSS), *Chenopodium album* L. (common lambsquarters, CHEAL), *Cirsium arvense* (L.) Scop.

30 (Canada thistle, CIRAR), *Galium aparine* L. (catchweed bedstraw, GALAP), *Kochia scoparia* (L.) Schrad. (kochia, KCHSC), *Lamium purpureum* L. (purple deadnettle, LAMPU), *Matricaria recutita* L. (wild chamomile, MATCH), *Matricaria matricarioides*

(Less.) Porter (pineappleweed, MATMT), *Papaver rhoeas* L. (common poppy, PAPRH), *Polygonum convolvulus* L. (wild buckwheat, POLCO), *Salsola tragus* L. (Russian thistle, SASKR), *Sinapis* species (SINSS), *Sinapis arvensis* L. (wild mustard, SINAR), *Stellaria media* (L.) Vill. (common chickweed, STEME), *Veronica persica* Poir. (Persian speedwell, VERPE), *Viola arvensis* Murr. (field violet, VIOAR), or *Viola tricolor* L. (wild violet, VIOTR).

In some embodiments, the methods provided herein are utilized to control undesirable vegetation in range and pasture, fallowland, IVM and ROW. In certain embodiments, the undesirable vegetation is *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Cassia obtusifolia* (sickle pod, CASOB), *Centaurea maculosa* auct. non Lam. (spotted knapweed, CENMA), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Convolvulus arvensis* L. (field bindweed, CONAR), *Daucus carota* L. (wild carrot, DAUCA), *Euphorbia esula* L. (leafy spurge, EPHEs), *Lactuca serriola* L./Torn. (prickly lettuce, LACSE), *Plantago lanceolata* L. (buckhorn plantain, PLALA), *Rumex obtusifolius* L. (broadleaf dock, RUMOB), *Sida spinosa* L. (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Sonchus arvensis* L. (perennial sowthistle, SONAR), *Solidago* species (goldenrod, SOOSS), *Taraxacum officinale* G.H. Weber ex Wiggers (dandelion, TAROF), *Trifolium repens* L. (white clover, TRFRE), or *Urtica dioica* L. (common nettle, URTDI).

In some embodiments, the methods provided herein are utilized to control undesirable vegetation found in row crops, tree and vine crops, and perennial crops. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Avena fatua* L. (wild oat, AVEFA), *Brachiaria decumbens* Stapf. or *Urochloa decumbens* (Stapf) R.D. Webster (Surinam grass, BRADC), *Brachiaria brizantha* (Hochst. ex A. Rich.) Stapf. or *Urochloa brizantha* (Hochst. ex A. Rich.) R.D. (beard grass, BRABR), *Brachiaria platyphylla* (Groseb.) Nash or *Urochloa platyphylla* (Nash) R.D. Webster (broadleaf signalgrass, BRAPP), *Brachiaria plantaginea* (Link) Hitchc. or *Urochloa plantaginea* (Link) R.D. Webster (alexandergrass, BRAPL), *Cenchrus echinatus* L. (southern sandbur, CENEC), *Digitaria horizontalis* Willd. (Jamaican crabgrass, DIGHO), *Digitaria insularis* (L.) Mez ex Ekman (sourgrass, TRCIN), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa crus-galli* (L.) P. Beauv. (barnyardgrass, ECHCG), *Echinochloa colonum* (L.) Link (junglerice, ECHCO), *Eleusine indica* (L.) Gaertn. (goosegrass, ELEIN), *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Panicum*

dichotomiflorum Michx. (fall panicum, PANDI), *Panicum miliaceum* L. (wild-proso millet, PANMI), *Setaria faberi* Herrm. (giant foxtail, SETFA), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Sorghum halepense* (L.) Pers. (Johnsongrass, SORHA), *Sorghum bicolor* (L.) Moench ssp. *Arundinaceum* (shattercane, SORVU), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Abutilon theophrasti* Medik. (velvetleaf, ABUTH), *Amaranthus* species (pigweeds and amaranths, AMASS), *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Ambrosia psilostachya* DC. (western ragweed, AMBPS), *Ambrosia trifida* L. (giant ragweed, AMBTR), *Anoda cristata* (L.) Schlecht. (spurred anoda, ANVCR), *Asclepias syriaca* L. (common milkweed, ASCSY),
 5 *Bidens pilosa* L. (hairy beggarticks, BIDPI), *Borreria* species (BOISS), *Borreria alata* (Aubl.) DC. or *Spermacoce alata* Aubl. (broadleaf buttonweed, BOILF), *Spermacoce latifolia* (broadleaved button weed, BOILF), *Chenopodium album* L. (common lambsquarters, CHEAL), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Commelina benghalensis* L. (tropical spiderwort, COMBE), *Datura stramonium* L. (jimsonweed, DATST), *Daucus*
 10 *carota* L. (wild carrot, DAUCA), *Euphorbia heterophylla* L. (wild poinsettia, EPHHL), *Euphorbia hirta* L. or *Chamaesyce hirta* (L.) Millsp. (garden spurge, EPHHI), *Euphorbia dentata* Michx. (toothed spurge, EPHDE), *Erigeron bonariensis* L. or *Conyza bonariensis* (L.) Cronq. (hairy fleabane, ERIBO), *Erigeron canadensis* L. or *Conyza canadensis* (L.) Cronq. (Canadian fleabane, ERICA), *Conyza sumatrensis* (Retz.) E. H. Walker (tall fleabane,
 15 ERIFL), *Helianthus annuus* L. (common sunflower, HELAN), *Jacquemontia tamnifolia* (L.) Griseb. (smallflower morningglory, IAQTA), *Ipomoea hederacea* (L.) Jacq. (ivy leaf morningglory, IPOHE), *Ipomoea lacunosa* L. (white morningglory, IPOLA), *Lactuca serriola* L./Torn. (prickly lettuce, LACSE), *Portulaca oleracea* L. (common purslane, POROL), *Richardia* species (pusley, RCHSS), *Sida* species (sida, SIDSS), *Sida spinosa* L.
 20 (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Solanum ptychanthum* Dunal (eastern black nightshade, SOLPT), *Tridax procumbens* L. (coat buttons, TRQPR), or *Xanthium strumarium* L. (common cocklebur, XANST).

In some embodiments, the methods provided herein are utilized to control undesirable vegetation in turf. In certain embodiments, the undesirable vegetation is *Bellis perennis* L.
 30 (English daisy, BELPE), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus* species (CYPSS), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Diodia virginiana* L. (Virginia buttonweed, DIQVI), *Euphorbia* species (spurge, EPHSS), *Glechoma hederacea* L.

(ground ivy, GLEHE), *Hydrocotyle umbellata* L. (dollarweed, HYDUM), *Kyllinga* species (kyllinga, KYLSS), *Lamium amplexicaule* L. (henbit, LAMAM), *Murdannia nudiflora* (L.) Brenan (doveweed, MUDNU), *Oxalis* species (woodsorrel, OXASS), *Plantago major* L. (broadleaf plantain, PLAMA), *Plantago lanceolata* L. (buckhorn/narrowleaf plantain, PLALA), *Phyllanthus urinaria* L. (chamberbitter, PYLTE), *Rumex obtusifolius* L. (broadleaf dock, RUMOB), *Stachys floridana* Shuttlew. (Florida betony, STAF), *Stellaria media* (L.) Vill. (common chickweed, STEME), *Taraxacum officinale* G.H. Weber ex Wiggers (dandelion, TAROF), *Trifolium repens* L. (white clover, TRFRE), or *Viola* species (wild violet, VIOSS).

In some embodiments, the compositions and methods provided herein are utilized to control undesirable vegetation consisting of grass, broadleaf and sedge weeds. In certain embodiments, the compositions and methods provided herein are utilized to control undesirable vegetation including *Abutilon*, *Amaranthus*, *Brachiaria* or *Urochloa*, *Brassica*, *Chenopodium*, *Commelina*, *Cyperus*, *Echinochloa*, *Fimbristylis*, *Galium*, *Ipomoea*, *Kochia*, *Lamium*, *Leptochloa*, *Salsola*, *Setaria*, *Sinapis*, *Veronica*, *Xanthium*, *Bolboschoenus*, *Schoenoplectus*, *Sinapis*, *Veronica*, and *Xanthium* species.

In some embodiments, the combination of compound (I) or agriculturally acceptable ester or salt thereof and a PS II inhibitor or an agriculturally acceptable salt or ester thereof is used to control *Abutilon theophrasti* Medik. (velvetleaf, ABUTH), *Amaranthus retroflexus* L.

(redroot pigweed, AMARE), *Brachiaria platyphylla* (Groseb.) Nash or *Urochloa platyphylla* (Nash) R.D. Webster (broadleaf signalgrass, BRAPP), *Brassica napus* L. (oilseed rape, BRSNW), *Chenopodium album* L. (common lambsquarters, CHEAL), *Commelina benghalensis* L. (Benghal dayflower, COMBE), *Cyperus iria* L. (rice flatsedge, CYPRI), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Echinochloa crus-galli* (L.) Beauv.

(barnyardgrass, ECHCG), *Echinochloa colona* (L.) Link (junglerice, ECHCO), *Echinochloa oryzoides* (Ard.) Fritsch (early watergrass, ECHOR), *Fimbristylis miliacea* (L.) Vahl (globe fringerush, FIMMI), *Galium aparine* L. (catchweed bedstraw, GALAP), *Ipomoea hederacea* Jacq. (ivy leaf morningglory, IPOHE), *Kochia scoparia* (L.) Schrad. (kochia, KCHSC), *Lamium purpureum* L. (purple deadnettle, LAMPU), *Leptochloa chinensis* (L.) Nees

(Chinese sprangletop, LEFCH), *Bolboschoenus maritimus* (L.) Palla or *Schoenoplectus maritimus* (L.) Lye (sea clubrush, SCPMA), *Salsola tragus* L. (Russian thistle, SASKR), *Setaria faberi* Herrm. (giant foxtail, SETFA), *Setaria viridis* (L.) Beauv. (green foxtail,

SETVI), *Sinapis arvensis* L. (wild mustard, SINAR), and *Veronica persica* Poir. (Persian speedwell, VERPE), and *Xanthium strumarium* L. (common cocklebur, XANST).

The compounds of formula I or agriculturally acceptable salt or ester thereof may be used to control herbicide resistant or tolerant weeds. The methods employing the combination of a compound of formula I or agriculturally acceptable salt or ester thereof and the compositions described herein may also be employed to control herbicide resistant or tolerant weeds. Exemplary resistant or tolerant weeds include, but are not limited to, biotypes resistant or tolerant to acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitors (*e.g.*, imidazolinones, sulfonylureas, pyrimidinylthiobenzoates, triazolopyrimidines, sulfonylaminocarbonyltriazolinones), photosystem II inhibitors (*e.g.*, phenylcarbamates, pyridazinones, triazines, triazinones, uracils, amides, ureas, benzothiadiazinones, nitriles, phenylpyridazines), acetyl CoA carboxylase (ACCase) inhibitors (*e.g.*, aryloxyphenoxypropionates, cyclohexanediones, phenylpyrazolines), synthetic auxins, (*e.g.*, benzoic acids, phenoxycarboxylic acids, pyridine carboxylic acids, quinoline carboxylic acids), auxin transport inhibitors (*e.g.*, phthalamates, semicarbazones), photosystem I inhibitors (*e.g.*, bipyridyliums), 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, (*e.g.*, glyphosate), glutamine synthetase inhibitors (*e.g.*, glufosinate, bialafos), microtubule assembly inhibitors, (*e.g.*, benzamides, benzoic acids, dinitroanilines, phosphoramidates, pyridines), mitosis inhibitors (*e.g.*, carbamates), very long chain fatty acid (VLCFA) inhibitors (*e.g.*, acetamides, chloroacetamides, oxyacetamides, tetrazolinones), fatty acid and lipid synthesis inhibitors (*e.g.*, phosphorodithioates, thiocarbamates, benzofuranes, chlorocarbonic acids), protoporphyrinogen oxidase (PPO) inhibitors (*e.g.*, diphenylethers, N-phenylphthalimides, oxadiazoles, oxazolidinediones, phenylpyrazoles, pyrimidindiones, thiadiazoles, triazolinones), carotenoid biosynthesis inhibitors (*e.g.*, clomazone, amitrole, aclonifen), phytoene desaturase (PDS) inhibitors (*e.g.*, amides, anilidex, furanones, phenoxybutan-amides, pyridiazinones, pyridines), 4-hydroxyphenyl-pyruvate-dioxygenase (HPPD) inhibitors (*e.g.*, callistemones, isoxazoles, pyrazoles, triketones), cellulose biosynthesis inhibitors (*e.g.*, nitriles, benzamides, quinclorac, triazolocarbaxamides), herbicides with multiple modes-of-action such as quinclorac, and unclassified herbicides such as arylaminopropionic acids, difenzoquat, endothall, and organoarsenicals. Exemplary resistant or tolerant weeds include, but are not limited to, biotypes with resistance or tolerance to multiple herbicides, biotypes with resistance or

tolerance to multiple chemical classes, biotypes with resistance or tolerance to multiple herbicide modes-of-action, and biotypes with multiple resistance or tolerance mechanisms (*e.g.*, target site resistance or metabolic resistance).

In some embodiments, an agriculturally acceptable ester or salt of bentazon is employed in the methods or compositions described herein. In certain embodiments, the sodium salt of bentazon is employed.

In some embodiments, an agriculturally acceptable ester or salt of ioxynil is employed in the methods or compositions described herein. In certain embodiments, the sodium salt of ioxynil is employed. In certain embodiments, the potassium salt of ioxynil is employed. In certain embodiments, the octanoate ester of ioxynil is employed.

In some embodiments, an agriculturally acceptable ester or salt of bromoxynil is employed in the methods or compositions described herein. In certain embodiments, the potassium salt of bromoxynil is employed. In certain embodiments, the butyl ester of bromoxynil is employed. In certain embodiments, the heptyl ester of bromoxynil is employed. In certain embodiments, the octanoate ester of bromoxynil is employed.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with atrazine or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to atrazine or salt thereof is within the range from about 1:2200 to about 6:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to atrazine or salt thereof is within the range from 1:509 to about 1:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to atrazine or salt thereof is within the range from about 1:512 to about 1:8. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to atrazine or salt thereof is within the range from about 1:256 to about 1:16. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to atrazine or salt thereof is within the range from about 1:256 to about 1:9. In certain embodiments, the compositions provided herein comprise the compound of formula (I) or its benzyl ester and atrazine or salt thereof. In one embodiment, the composition comprises the compound of formula (I) and atrazine or salt thereof, wherein the weight ratio of the compound of formula (I) to atrazine or salt thereof is about 1:256 to about 1:32. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and

atrazine or salt thereof, wherein the weight ratio of the benzyl ester of the compound of formula (I) to atrazine or salt thereof is about 1:130 to about 1:16. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and atrazine or salt thereof, wherein the weight ratio of the benzyl ester of the compound of formula (I) to atrazine or salt thereof is about 1:140 to about 1:17.5. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 52 grams active ingredient per hectare (g ai/ha) to about 4700 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 60 grams active ingredient per hectare (g ai/ha) to about 2300 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and atrazine or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the atrazine or salt thereof is applied at a rate from about 50 g ai/ha to about 4400 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the atrazine or salt thereof is applied at a rate from about 140 g ai/ha to about 2240 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha. In some embodiments, the atrazine or salt thereof is applied at a rate from about 280 g ai/ha to about 1120 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its benzyl ester and atrazine or salt thereof. In one embodiment, the methods utilize the compound of formula (I) and atrazine or salt thereof, wherein the compound of formula (I) is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 8.75 g ae/ha, and atrazine or salt thereof is applied at a rate of about 280 g ai/ha to about 1120 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and atrazine or salt thereof, wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 17.5 g ae/ha, and atrazine or salt thereof is

applied at a rate of about 280 g ai/ha to about 560 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and atrazine or salt thereof, wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 8 g acid equivalent per hectare (g ae/ha) to about 32 g ae/ha, and atrazine or salt thereof is applied at a rate of about 560 g ai/ha to about 1120 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with atrazine or salt thereof are used to control BRAPP, CYPIR, ECHCG, XANST or ECHCO.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with bentazon or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to bentazon or salt thereof is within the range from about 1:1120 to about 3:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to bentazon or salt thereof is within the range from about 1:256 to about 1:1.2. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to bentazon or salt thereof is within the range from about 1:192 to about 1:3. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to is within the range from about 1:96 to about 1:6. In certain embodiments, the compositions provided herein comprise the compound of formula (I) or its benzyl or *n*-butyl ester and bentazon-sodium. In one embodiment, the composition comprises the compound of formula (I) and bentazon-sodium, wherein the weight ratio of the compound of formula (I) to bentazon-sodium is about 1:96 to about 1:13. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and bentazon-sodium, wherein the weight ratio of the benzyl ester of the compound of formula (I) to bentazon-sodium is about 1:96 to about 1:3.4. In one embodiment, the composition comprises the *n*-butyl ester of the compound of formula (I) and bentazon-sodium, wherein the weight ratio of the *n*-butyl ester of the compound of formula (I) to bentazon-sodium is about 1:48 to about 1:16. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and bentazon-sodium, wherein the weight ratio of the benzyl ester of the compound of formula (I) to bentazon-sodium is about 1:13.7 to about 1:3.4. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition

described herein. In some embodiments, the composition is applied at an application rate from about 107 grams active ingredient per hectare (g ai/ha) to about 2540 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 110 grams active ingredient per hectare (g ai/ha) to about 1205 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and bentazon or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the bentazon or salt thereof is applied at a rate from about 105 g ai/ha to about 2240 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the bentazon or salt thereof is applied at a rate from about 50 g ai/ha to about 2240 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g acid equivalent per hectare (g ae/ha) to about 140 g ae/ha. In some embodiments, the bentazon or salt thereof is applied at a rate from about 60 g ai/ha to about 1120 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 84.4 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its benzyl or *n*-butyl ester and bentazon-sodium. In one embodiment, the methods utilize the compound of formula (I) and bentazon-sodium, wherein the compound of formula (I) is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 84.8 g ae/ha, and bentazon-sodium is applied at a rate of about 105 g ai/ha to about 1120 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and bentazon-sodium, wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 70 g ae/ha, and bentazon-sodium is applied at a rate of about 105 g ai/ha to about 1120 g ai/ha. In one embodiment, the methods utilize the *n*-butyl ester of the compound of formula (I) and bentazon-sodium, wherein the *n*-butyl ester of the compound of formula (I) is applied at a rate of about 17.5 g acid equivalent per hectare (g ae/ha) to about 70 g ae/ha, and bentazon-sodium is applied at a rate of about 840 g ai/ha to about 1120 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and bentazon-sodium, wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 8.75 g acid equivalent per hectare (g ae/ha) to about 17.5 g

ae/ha, and bentazon-sodium is applied at a rate of about 60 g ai/ha to about 120 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with bentazon or salt thereof are used to control BRSNW, COMBE, ECHCG, ECHOR, IPOHE, LEFCH, SCPMA, or SETFA.

5 In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with bromoxynil or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to bromoxynil or salt thereof is within the range from about 1:280 to about 12:1. In certain embodiments, the weight ratio of the
10 compound of formula (I) or salt or ester thereof to bromoxynil or salt thereof is within the range from about 1:254 to about 2:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to bromoxynil or salt thereof is within the range from about 1:18 to about 1:4. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and bromoxynil or salt thereof. With
15 respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 27 grams active ingredient per hectare (g ai/ha) to about 860 g ai/ha based on the total amount of active ingredients in the
20 composition. In certain embodiments, the composition is applied at an application rate from about 50 grams active ingredient per hectare (g ai/ha) to about 610 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or
25 salt or ester thereof and bromoxynil or salt thereof or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the bromoxynil or salt thereof is applied at a rate from about 25 g ai/ha to about 560 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In certain
30 embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and bromoxynil or salt thereof can be use to control AMARE, ECHCG, IPOHE, or SETFA.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with chlorotoluron or

salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to chlorotoluron or salt thereof is within the range from about 1:1750 to about 1.5:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to chlorotoluron or salt thereof is within the range from about 1:909 to about 1:4. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and chlorotoluron or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 202 grams active ingredient per hectare (g ai/ha) to about 3800 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 203 grams active ingredient per hectare (g ai/ha) to about 2500 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and chlorotoluron or salt thereof or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the chlorotoluron or salt thereof is applied at a rate from about 200 g ai/ha to about 3500 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and chlorotoluron or salt thereof.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with cyanazine or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to cyanazine or salt thereof is within the range from about 1:2650 to about 6:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to cyanazine or salt thereof is within the range from about 1:1600 to about 1:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to cyanazine or salt thereof is within the range from about 1:110 to about 1:5.3. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and cyanazine or salt thereof. With

respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 52 grams active ingredient per hectare (g ai/ha) to about 5600 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 53 grams active ingredient per hectare (g ai/ha) to about 3550 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and cyanazine or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the cyanazine or salt thereof is applied at a rate from about 50 g ai/ha to about 5300 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the cyanazine or salt thereof is applied at a rate from about 170 g ai/ha to about 880 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 8 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and cyanazine or salt thereof for the control of BRAPP or IPOHE.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with diuron or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to diuron or salt thereof is within the range from about 1:3600 to about 6:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to diuron or salt thereof is within the range from about 1:509 to about 1:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to diuron or salt thereof is within the range from about 1:70 to about 1:9. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and diuron or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 52 grams active ingredient per hectare (g ai/ha) to about 7500 g ai/ha based on the

total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 55 grams active ingredient per hectare (g ai/ha) to about 2300 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable
5 vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and diuron or salt thereof or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the diuron or salt thereof is applied at a rate from about 50 g ai/ha to about 7200 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to
10 about 300 g ae/ha. In some embodiments, the diuron or salt thereof is applied at a rate from about 280 g ai/ha to about 1,120 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 16 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and diuron or salt thereof for the control of ABUTH, BRAPP, ECHCG, ECHCO, SETFA or
15 SETVI.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with hexazinone or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to hexazinone or salt thereof is within the
20 range from about 1:2500 to about 12:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to hexazinone or salt thereof is within the range from about 1:1360 to about 2:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to hexazinone or salt thereof is within the range from about 1:156 to about 1:19. In certain embodiments, the compositions comprise
25 the compound of formula (I) or its benzyl or n-butyl ester and hexazinone or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 27 grams active ingredient per
30 hectare (g ai/ha) to about 5300 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 28 grams active ingredient per hectare (g ai/ha) to about 3050 g ai/ha based on the total

amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and hexazinone or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the hexazinone or salt thereof is applied at a rate from about 25 g ai/ha to about 5000 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the hexazinone or salt thereof is applied at a rate from about 625 g ai/ha to about 1,250 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 8 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and hexazinone or salt thereof for the control of ECHCO, ECHCG or IPOHE.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with ioxynil or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to ioxynil or salt thereof is within the range from about 1:200 to about 12:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to ioxynil or salt thereof is within the range from about 1:136 to about 6:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to ioxynil or salt thereof is within the range from about 1:38 to about 1:9. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and ioxynil or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 27 grams active ingredient per hectare (g ai/ha) to about 700 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 28 grams active ingredient per hectare (g ai/ha) to about 450 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and ioxynil or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a

composition described herein. In some embodiments, the ioxynil or salt thereof is applied at a rate from about 25 g ai/ha to about 400 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the ioxynil or salt thereof is applied at a rate from about 300 g ai/ha to about 400 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 8 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and ioxynil or salt thereof for the control of IPOHE.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with isoproturon or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to isoproturon or salt thereof is within the range from about 1:750 to about 12:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to isoproturon or salt thereof is within the range from about 1:600 to about 1:37.5. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to isoproturon or salt thereof is within the range from about 1:455 to about 2:1. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and isoproturon or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 27 grams active ingredient per hectare (g ai/ha) to about 1800 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 28 grams active ingredient per hectare (g ai/ha) to about 1050 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and isoproturon or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the isoproturon or salt thereof is applied at a rate from about 25 g ai/ha to about 1500 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the isoproturon or salt

thereof is applied at a rate from about 375 g ai/ha to about 1500 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2.5 g ae/ha to about 10 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and isoproturon or salt thereof for the control of AMARE, SINAR, GALAP,
5 LAMPU or VERPE.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with linuron or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to linuron or salt thereof is within the range
10 from about 1:2250 to about 6:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to linuron or salt thereof is within the range from about 1:39 to about 1:19. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to linuron or salt thereof is within the range from about 1:1364 to about 1:1. In certain embodiments, the compositions comprise the compound of formula (I) or its
15 benzyl or n-butyl ester and linuron or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 52 grams active ingredient per hectare (g ai/ha) to about 4800 g ai/ha based on the
20 total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 53 grams active ingredient per hectare (g ai/ha) to about 3050 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or
25 growth of vegetation with a compound of formula (I) or salt or ester thereof and linuron or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the linuron or salt thereof is applied at a rate from about 50 g ai/ha to about 4500 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the linuron or salt thereof is applied at a rate from about 50 g
30 ai/ha to about 620 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 16 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods

utilize the compound of formula (I) or its benzyl or n-butyl ester and linuron or salt thereof for the control of BRAPP.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with methibenzuron or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to methibenzuron or salt thereof is within the range from about 1:1400 to about 4:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to methibenzuron or salt thereof is within the range from about 1:909 to about 1:1. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and methibenzuron or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 72 grams active ingredient per hectare (g ai/ha) to about 3100 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 73 grams active ingredient per hectare (g ai/ha) to about 2050 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and methibenzuron or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the methibenzuron or salt thereof is applied at a rate from about 70 g ai/ha to about 2800 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and methibenzuron or salt thereof.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with metribuzin or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to metribuzin or salt thereof is within the range from about 1:2800 to about 6:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to metribuzin or salt thereof is within the

range from about 1:509 to about 1:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to metribuzin or salt thereof is within the range from about 1:168 to about 1:1. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and metribuzin or salt thereof. With
5 respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 52 grams active ingredient per hectare (g ai/ha) to about 5900 g ai/ha based on the total amount of active ingredients in the
10 composition. In certain embodiments, the composition is applied at an application rate from about 55 grams active ingredient per hectare (g ai/ha) to about 2300 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or
15 salt or ester thereof and metribuzin or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the metribuzin or salt thereof is applied at a rate from about 50 g ai/ha to about 5600 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the metribuzin or salt thereof is applied at a rate from about 25 g ai/ha to about 420 g ai/ha and the compound of
20 formula (I) or salt or ester thereof is applied at a rate from about 2.5 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and metribuzin or salt thereof for the control of SETFA, KCHSC, LAMPU, SASKR, CHEAL or SINAR.

In certain embodiments of the compositions and methods described herein, the
25 compound of formula (I) or salt or ester thereof is used in combination with propanil or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to propanil or salt thereof is within the range from about 1:2800 to about 1:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to propanil or salt thereof is within the range from about
30 1:767 to about 1:12. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to propanil or salt thereof is within the range from about 1:772 to about 1:12. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester

thereof to propanil or salt thereof is within the range from about 1:384 to about 1:24. In certain embodiments, the compositions provided herein comprise the compound of formula (I) or its benzyl or *n*-butyl ester and propanil. In one embodiment, the composition comprises the compound of formula (I) and propanil, wherein the weight ratio of the compound of formula (I) to propanil is about 1:384 to about 1:48. In one embodiment, the composition comprises the compound of formula (I) and propanil, wherein the weight ratio of the compound of formula (I) to propanil is about 1:384 to about 1:48. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and propanil, wherein the weight ratio of the benzyl ester of the compound of formula (I) to propanil is about 1:767 to about 1:24. In one embodiment, the composition comprises the *n*-butyl ester of the compound of formula (I) and propanil, wherein the weight ratio of the *n*-butyl ester of the compound of formula (I) to propanil is about 1:210. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 422 grams active ingredient per hectare (g ai/ha) to about 5900 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 425 grams active ingredient per hectare (g ai/ha) to about 3400 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and propanil or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the propanil or salt thereof is applied at a rate from about 420 g ai/ha to about 5600 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the propanil or salt thereof is applied at a rate from about 210 g ai/ha to about 6720 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g acid equivalent per hectare (g ae/ha) to about 70 g ae/ha. In some embodiments, the propanil or salt thereof is applied at a rate from about 420 g ai/ha to about 3360 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 4 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha. In some embodiments, the propanil or salt thereof is applied at a rate from about 420 g ai/ha to about 3360 g ai/ha and

the compound of formula (I) of salt or ester thereof is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its benzyl or *n*-butyl ester and propanil. In one embodiment, the methods utilize the compound of formula (I) and propanil, wherein the compound of formula (I) is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha, and propanil is applied at a rate of about 420 g ai/ha to about 3360 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and propanil, wherein the benzyl ester of the compound of formula (I) is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha, and propanil is applied at a rate of about 420 g ai/ha to about 3360 g ai/ha. In one embodiment, the methods utilize the *n*-butyl ester of the compound of formula (I) and propanil, wherein the *n*-butyl ester of the compound of formula (I) is applied at a rate of about 16 g acid equivalent per hectare (g ae/ha), and propanil is applied at a rate of about 3360 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with propanil are used to control ECHCG, ECHCO, COMBE, IPOHE, or SCPMA.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with pyridate. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pyridate is within the range from about 1:800 to about 12:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pyridate is within the range from about 1:545 to about 6:1. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or *n*-butyl ester and pyridate. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 27 grams active ingredient per hectare (g ai/ha) to about 1800 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 28 grams active ingredient per hectare (g ai/ha) to about 1350 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or

applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and pyridate, *e.g.*, sequentially or simultaneously. In some embodiments, the pyridate is applied at a rate from about 25 g ai/ha to about 1600 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and pyridate.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with siduron or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to siduron or salt thereof is within the range from about 1:4500 to about 2:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to siduron or salt thereof is within the range from about 1:2728 to about 1:3. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and siduron or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 142 grams active ingredient per hectare (g ai/ha) to about 9300 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 143 grams active ingredient per hectare (g ai/ha) to about 6050 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and siduron or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the siduron or salt thereof is applied at a rate from about 140 g ai/ha to about 9000 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and siduron or salt thereof.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with simazine or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the

compound of formula (I) or salt or ester thereof to simazine or salt thereof is within the range from about 1:4250 to about 2:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to simazine or salt thereof is within the range from about 1:2728 to about 1:3. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to simazine or salt thereof is within the range from about 1:280 to about 1:17.5. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and simazine or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 142 grams active ingredient per hectare (g ai/ha) to about 8800 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 143 grams active ingredient per hectare (g ai/ha) to about 6050 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 568 grams active ingredient per hectare (g ai/ha) to about 2272 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and simazine or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the simazine or salt thereof is applied at a rate from about 140 g ai/ha to about 8500 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the simazine or salt thereof is applied at a rate from about 560 g ai/ha to about 2240 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 8 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and simazine or salt thereof for the control of BRAPP, ECHCG, ECHCO, CYPIR or XANST.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with simetryne or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to simetryne or salt thereof is within the

range from about 1:1000 to about 12:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to simetryne or salt thereof is within the range from about 1:682 to about 6:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to simetryne or salt thereof is within the

5 range from about 1:56 to about 1:3.5. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and simetryne or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments,

10 the composition is applied at an application rate from about 27 grams active ingredient per hectare (g ai/ha) to about 2300 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 28 grams active ingredient per hectare (g ai/ha) to about 1650 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods

15 comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and simetryne or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the simetryne or salt thereof is applied at a rate from about 25 g ai/ha to about 2000 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate

20 from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the simetryne or salt thereof is applied at a rate from about 110 g ai/ha to about 450 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 8 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and simetryne or salt thereof for the control of ECHCO, ECHOR, CYPPIR,

25 CYPPO, FIMMI or LEFCH.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with tebuthiuron or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to tebuthiuron or salt thereof is within the

30 range from about 1:2240 to about 2:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to tebuthiuron or salt thereof is within the range from about 1:1818 to about 1:1. In certain embodiments, the weight ratio of the

compound of formula (I) or salt or ester thereof to tebuthiuron or salt thereof is within the range from about 1:420 to about 1:6.5. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and tebuthiuron or salt thereof. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 142 grams active ingredient per hectare (g ai/ha) to about 4780 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 143 grams active ingredient per hectare (g ai/ha) to about 4100 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and tebuthiuron or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the tebuthiuron or salt thereof is applied at a rate from about 140 g ai/ha to about 4480 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the tebuthiuron or salt thereof is applied at a rate from about 140 g ai/ha to about 3360 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 4 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and tebuthiuron or salt thereof for the control of BRSNN, CIRAR, CYPES, ECHCG, IPOHE, or XANST.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with terbuthylazine or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to terbuthylazine or salt thereof is within the range from about 1:1500 to about 3:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to terbuthylazine or salt thereof is within the range from about 1:1018 to about 1:2.5. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to terbuthylazine or salt thereof is within the range from about 1:62 to about 1:31. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and terbuthylazine or salt thereof.

With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate from about 127 grams active ingredient per hectare (g ai/ha) to about 3300 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate from about 128 grams active ingredient per hectare (g ai/ha) to about 2290 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and terbuthylazine or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the terbuthylazine or salt thereof is applied at a rate from about 125 g ai/ha to about 3000 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g ae/ha to about 300 g ae/ha. In some embodiments, the terbuthylazine or salt thereof is applied at a rate from about 125 g ai/ha to about 1000 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 16 g ae/ha to about 32 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and terbuthylazine or salt thereof for the control of XANST.

The components of the mixtures described herein can be applied either separately or as part of a multipart herbicidal system.

The mixtures described herein can be applied in conjunction with one or more other herbicides to control a wider variety of undesirable vegetation. When used in conjunction with other herbicides, the composition can be formulated with the other herbicide or herbicides, tank mixed with the other herbicide or herbicides or applied sequentially with the other herbicide or herbicides. Some of the herbicides that can be employed in conjunction with the compositions and methods described herein include, but are not limited to: 4-CPA; 4-CPB; 4-CPP; 2,4-D; 2,4-D choline salt, 2,4-D esters and amines, 2,4-DB; 3,4-DA; 3,4-DB; 2,4-DEB; 2,4-DEP; 3,4-DP; 2,3,6-TBA; 2,4,5-T; 2,4,5-TB; acetochlor, acifluorfen, aclonifen, acrolein, alachlor, allidochlor, alloxydim, allyl alcohol, alorac, ametrifone, ametryn, amibuzin, amicarbazone, amidosulfuron, aminocyclopyrachlor, aminopyralid, amiprofos-methyl, amitrole, ammonium sulfamate, anilofos, anisuron, asulam, atraton, azafenidin, azimsulfuron, aziprotryne, barban, BCPC, beflubutamid, benazolin, bencarbazon, 57

benfluralin, benfuresate, bensulfuron-methyl, bensulide, benthocarb, benzadox,
 benzfendizone, benzipram, benzobicyclon, benzofenap, benzofluor, benzoylprop,
 benzthiazuron, bialaphos, bicyclopypyrone, bifenox, bilanafos, bispyribac-sodium, borax,
 bromacil, bromobonil, bromobutide, bromofenoxim, brompyrazon, butachlor, butafenacil,
 5 butamifos, butenachlor, buthidazole, buthiuron, butralin, butroxydim, buturon, butylate,
 cacodylic acid, cafenstrole, calcium chlorate, calcium cyanamide, cambendichlor,
 carbasulam, carbetamide, carboxazole chlorprocarb, carfentrazone-ethyl, CDEA, CEPC,
 chlomethoxyfen, chloramben, chloranocryl, chlorazifop, chlorazine, chlorbromuron,
 chlorbufam, chloreturon, chlorfenac, chlorfenprop, chlorflurazole, chlorflurenol, chloridazon,
 10 chlorimuron, chlornitrofen, chloropon, chloroxuron, chloroxynil, chlorpropham,
 chlorsulfuron, chlorthal, chlorthiamid, cinidon-ethyl, cinmethylin, cinosulfuron, cisanilide,
 clethodim, cliodinate, clodinafop-propargyl, clofop, clomazone, clomeprop, cloprop,
 cloproxydim, clopyralid, cloransulam-methyl, CMA, copper sulfate, CPMF, CPPC,
 credazine, cresol, cumyluron, cyanatryn, cyclopyrimorate, cycloate, cyclosulfamuron,
 15 cycloxydim, cycluron, cyhalofop-butyl, cyperquat, cyprazine, cyprazole, cypromid,
 daimuron, dalapon, dazomet, delachlor, desmedipham, desmetryn, di-allate, dicamba,
 dichlobenil, dichloralurea, dichlormate, dichlorprop, dichlorprop-P, diclofop-methyl,
 diclosulam, diethamquat, diethatyl, difenopenten, difenoxuron, difenzoquat, diflufenican,
 diflufenzopyr, dimefuron, dimepiperate, dimethachlor, dimethametryn, dimethenamid,
 20 dimethenamid-P, dimexano, dimidazon, dinitramine, dinofenate, dinoprop, dinosam, dinoseb,
 dinoterb, diphenamid, dipropetryn, diquat, disul, dithiopyr, DMPA, DNOC, DSMA, EBEP,
 eglinazine, endothal, epronaz, EPTC, erbon, esprocarb, ethalfluralin, ethbenzamide,
 ethametsulfuron, ethidimuron, ethiolate, ethobenzamid, etobenzamid, ethofumesate,
 ethoxyfen, ethoxysulfuron, etinofen, etnipromid, etobenzanid, EXD, fenasulam, fenoprop,
 25 fenoxaprop, fenoxaprop-P-ethyl, fenoxaprop-P-ethyl + isoxadifen-ethyl, fenoxasulfone,
 fenteracol, fenthiafop, fentrazamide, fenuron, ferrous sulfate, flamprop, flamprop-M,
 flazasulfuron, florasulam, fluazifop, fluazifop-P-butyl, fluazolate, flucarbazone,
 flucetosulfuron, fluchloralin, flufenacet, flufenican, flufenpyr-ethyl, flumetsulam, flumezin,
 flumiclorac-pentyl, flumioxazin, flumipropyn, fluometuron, fluorodifen, fluoroglycofen,
 30 fluoromidine, fluoronitrofen, fluothiuron, flupoxam, flupropacil, flupropanate,
 flupyrsulfuron, fluridone, flurochloridone, fluroxypyr, fluroxypyr-meptyl, flurtamone,
 fluthiacet, fomesafen, foramsulfuron, fosamine, fumiclorac, furyloxyfen, glufosinate,

glufosinate-ammonium, glufosinate-P-ammonium, glyphosate, halauxifen, halauxifen-methyl, halosafen, halosulfuron-methyl, haloxydine, haloxyfop-methyl, haloxyfop-P-methyl, hexachloroacetone, hexaflurate, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazosulfuron, imazethapyr, indanofan, indaziflam, iodobonil, iodomethane, iodosulfuron, iodosulfuron-ethyl-sodium, iofensulfuron, ipazine, ipfencarbazon, iprymidam, isocarbamid, isocil, isomethiozin, isonoruron, isopolinate, isopropalin, isouron, isoxaben, isoxachlortole, isoxaflutole, isoxapyrifop, karbutilate, ketospiradox, lactofen, lenacil, MAA, MAMA, MCPA esters and amines, MCPA-thioethyl, MCPB, mecoprop, mecoprop-P, medinoterb, mefenacet, mefluidide, mesoprazine, mesosulfuron, mesotrione, metam, metamifop, metamitron, metazachlor, metazosulfuron, metflurazon, methabenzthiazuron, methalpropalin, methazole, methiobencarb, methiozin, methiuron, methometon, methoprotrotryne, methyl bromide, methyl isothiocyanate, methyldymron, metobenzuron, metobromuron, metolachlor, metosulam, metoxuron, metsulfuron, metsulfuron-methyl, molinate, monalide, monisouron, monochloroacetic acid, monolinuron, monuron, morfamquat, MSMA, naproanilide, napropamide, naptalam, neburon, nicosulfuron, nipyraclufen, nitratin, nitrofen, nitrofluorfen, norflurazon, noruron, OCH, orbencarb, *ortho*-dichlorobenzene, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon, oxapyrazon, oxasulfuron, oxaziclomefone, oxyfluorfen, paraflufen-ethyl, parafluron, paraquat, pebulate, pelargonic acid, pendimethalin, penoxsulam, pentachlorophenol, pentanochlor, pentoxazone, perfluidone, pethoxamid, phenisopham, phenmedipham, phenmedipham-ethyl, phenobenzuron, phenylmercury acetate, picloram, picolinafen, pinoxaden, piperophos, potassium arsenite, potassium azide, potassium cyanate, pretilachlor, primisulfuron-methyl, procyazine, prodiamine, profluazol, profluralin, profoxydim, proglinazine, prohexadione-calcium, prometon, prometryn, pronamide, propachlor, propaquizafop, propazine, propham, propisochlor, propoxycarbazon, propyrisulfuron, propyzamide, prosulfalin, prosulfocarb, prosulfuron, proxan, prynachlor, pydanon, pyraclonil, pyraflufen-ethyl, pyrasulfotole, pyrazogyl, pyrazolynate, pyrazosulfuron-ethyl, pyrazoxyfen, pyribenzoxim, pyributicarb, pyriclor, pyridafol, pyriftalid, pyriminobac, pyrimisulfan, pyriothiac-sodium, pyroxasulfone, pyroxulam, quinclorac, quinmerac, quinclamine, quinonamid, quizalofop, quizalofop-P-ethyl, rhodethanil, rimsulfuron, saflufenacil, S-metolachlor, sebuthylazine, sebumeton, sethoxydim, simeton, SMA, sodium arsenite, sodium azide, sodium chlorate, sulcotrione, sulfallate, sulfentrazone, sulfometuron, sulfosate, sulfosulfuron, sulfuric acid, sulglycapin,

swep, SYN-523, TCA, tebutam, tefuryltrione, tembotrione, tepraloxym, terbacil, terbucarb, terbutryn, tetrafluron, thenylchlor, thiazafluron, thiazopyr, thidiazimin, thidiazuron, thiencarbazone-methyl, thifensulfuron, thifensulfuron-methyl, thiobencarb, tiocarbamil, tioclorim, topramezone, tralkoxydim, triafamone, tri-allate, triasulfuron, triaziflam, tribenuron, tribenuron-methyl, tricamba, triclopyr choline salt, triclopyr esters and salts, tridiphane, trietazine, trifloxysulfuron, trifluralin, triflusulfuron, trifop, trifopside, trihydroxytriazine, trimeturon, tripropindan, tritac tritosulfuron, vernolate, xylachlor and salts, esters, optically active isomers and mixtures thereof.

The compositions and methods described herein, can further be used in conjunction with glyphosate, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, glufosinate, glutamine synthetase inhibitors, dicamba, phenoxy auxins, pyridyloxy auxins, synthetic auxins, auxin transport inhibitors, aryloxyphenoxypropionates, cyclohexanediones, phenylpyrazolines, acetyl CoA carboxylase (ACCase) inhibitors, imidazolinones, sulfonylureas, pyrimidinylthiobenzoates, triazolopyrimidines, sulfonylaminocarbonyltriazolinones, acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitors, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitors, phytoene desaturase inhibitors, carotenoid biosynthesis inhibitors, protoporphyrinogen oxidase (PPO) inhibitors, cellulose biosynthesis inhibitors, mitosis inhibitors, microtubule inhibitors, very long chain fatty acid inhibitors, fatty acid and lipid biosynthesis inhibitors, photosystem I inhibitors, photosystem II inhibitors, triazines, and bromoxynil on glyphosate-tolerant, EPSP synthase inhibitor-tolerant, glufosinate-tolerant, glutamine synthetase inhibitor-tolerant, dicamba-tolerant, phenoxy auxin-tolerant, pyridyloxy auxin-tolerant, auxin-tolerant, auxin transport inhibitor-tolerant, aryloxyphenoxypropionate-tolerant, cyclohexanedione-tolerant, phenylpyrazoline-tolerant, ACCase-tolerant, imidazolinone-tolerant, sulfonylurea-tolerant, pyrimidinylthiobenzoate-tolerant, triazolopyrimidine-tolerant, sulfonylaminocarbonyltriazolinone-tolerant, ALS- or AHAS-tolerant, HPPD-tolerant, phytoene desaturase inhibitor-tolerant, carotenoid biosynthesis inhibitor tolerant, PPO-tolerant, cellulose biosynthesis inhibitor-tolerant, mitosis inhibitor-tolerant, microtubule inhibitor-tolerant, very long chain fatty acid inhibitor-tolerant, fatty acid and lipid biosynthesis inhibitor-tolerant, photosystem I inhibitor-tolerant, photosystem II inhibitor-tolerant, triazine-tolerant, and bromoxynil-tolerant crops, and crops possessing multiple or stacked traits conferring tolerance to multiple chemistries and/or multiple modes of action via

single and/or multiple resistance mechanisms. In some embodiments, the compound of formula (I) or salt or ester thereof and complementary herbicide or salt or ester thereof are used in combination with herbicides that are selective for the crop being treated and which complement the spectrum of weeds controlled by these compounds at the application rate employed. In some embodiments, the compositions described herein and other complementary herbicides are applied at the same time, either as a combination formulation, as a tank mix, or as a sequential application.

In some embodiments, the compositions described herein are employed in combination with one or more herbicide safeners, such as AD-67 (MON 4660), benoxacor, benthicarb, brassinolide, cloquintocet (mexyl), cyometrinil, daimuron, dichlormid, dicyclonon, dimepiperate, disulfoton, fenchlorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, harpin proteins, isoxadifen-ethyl, jiecaowan, jiecaoxi, mefenpyr-diethyl, mephenate, naphthalic anhydride (NA), oxabetrinil, R29148 and *N*-phenyl-sulfonylbenzoic acid amides, to enhance their selectivity. In some embodiments, the safeners are employed in rice, cereal, corn, or maize settings. In some embodiments, the safener is cloquintocet or an ester or salt thereof. In certain embodiments, cloquintocet is utilized to antagonize harmful effects of the compositions on rice and cereals. In some embodiments, the safener is cloquintocet (mexyl).

In some embodiments, the compositions described herein are employed in combination with one or more plant growth regulators, such as 2,3,5-tri-iodobenzoic acid, IAA, IBA, naphthaleneacetamide, α -naphthaleneacetic acids, benzyladenine, 4-hydroxyphenethyl alcohol, kinetin, zeatin, endothal, ethephon, pentachlorophenol, thidiazuron, tribufos, aviglycine, gibberellins, gibberellic acid, abscisic acid, ancymidol, fosamine, glyphosine, isopyrimol, jasmonic acid, maleic hydrazide, mepiquat, 2,3,5-tri-iodobenzoic acid, morphactins, dichlorflurenol, flurprimidol, mefluidide, paclobutrazol, tetcyclacis, uniconazole, brassinolide, brassinolide-ethyl, cycloheximide, ethylene, methasulfocarb, prohexadione, triapenthenol and trinexapac.

In some embodiments, the plant growth regulators are employed in one or more crops or settings, such as rice, cereal crops, corn, maize, broadleaf crops, oilseed rape/canola, turf, pineapple, sugarcane, sunflower, pastures, grasslands, rangelands, fallowland, turf, tree and vine orchards, plantation crops, vegetables, and non-crop (ornamentals) settings. In some embodiments, the plant growth regulator is mixed with the compound of formula (I), or

mixed with the compound of formula (I) and PSII inhibitors to cause a preferentially advantageous effect on plants.

In some embodiments, compositions provided herein further comprise at least one agriculturally acceptable adjuvant or carrier. Suitable adjuvants or carriers should not be phytotoxic to valuable crops, particularly at the concentrations employed in applying the compositions for selective weed control in the presence of crops, and should not react chemically with herbicidal components or other composition ingredients. Such mixtures can be designed for application directly to weeds or their locus or can be concentrates or formulations that are normally diluted with additional carriers and adjuvants before application. They can be solids, such as, for example, dusts, granules, water-dispersible granules, or wettable powders, or liquids, such as, for example, emulsifiable concentrates, solutions, emulsions or suspensions. They can also be provided as a pre-mix or tank mixed.

Suitable agricultural adjuvants and carriers include, but are not limited to, crop oil concentrate; nonylphenol ethoxylate; benzylcocoalkyldimethyl quaternary ammonium salt; blend of petroleum hydrocarbon, alkyl esters, organic acid, and anionic surfactant; C₉-C₁₁ alkylpolyglycoside; phosphated alcohol ethoxylate; natural primary alcohol (C₁₂-C₁₆) ethoxylate; di-*sec*-butylphenol EO-PO block copolymer; polysiloxane-methyl cap; nonylphenol ethoxylate + urea ammonium nitrate; emulsified methylated seed oil; tridecyl alcohol (synthetic) ethoxylate (8EO); tallow amine ethoxylate (15 EO); PEG(400) dioleate-99.

Liquid carriers that can be employed include water and organic solvents. The organic solvents include, but are not limited to, petroleum fractions or hydrocarbons such as mineral oil, aromatic solvents, paraffinic oils, and the like; vegetable oils such as soybean oil, rapeseed oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; esters of the above vegetable oils; esters of monoalcohols or dihydric, trihydric, or other lower polyalcohols (4-6 hydroxy containing), such as 2-ethyl hexyl stearate, *n*-butyl oleate, isopropyl myristate, propylene glycol dioleate, di-octyl succinate, di-butyl adipate, di-octyl phthalate and the like; esters of mono, di and polycarboxylic acids and the like. Specific organic solvents include, but are not limited to toluene, xylene, petroleum naphtha, crop oil, acetone, methyl ethyl ketone, cyclohexanone, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol monomethyl ether and diethylene glycol

monomethyl ether, methyl alcohol, ethyl alcohol, isopropyl alcohol, amyl alcohol, ethylene glycol, propylene glycol, glycerine, *N*-methyl-2-pyrrolidinone, *N,N*-dimethyl alkylamides, dimethyl sulfoxide, liquid fertilizers and the like. In certain embodiments, water is the carrier for the dilution of concentrates.

5 Suitable solid carriers include but are not limited to talc, pyrophyllite clay, silica, attapulgus clay, kaolin clay, kieselguhr, chalk, diatomaceous earth, lime, calcium carbonate, bentonite clay, Fuller's earth, cottonseed hulls, wheat flour, soybean flour, pumice, wood flour, walnut shell flour, lignin, cellulose, and the like.

10 In some embodiments, the compositions described herein further comprise one or more surface-active agents. In some embodiments, such surface-active agents are employed in both solid and liquid compositions, and in certain embodiments those designed to be diluted with carrier before application. The surface-active agents can be anionic, cationic or nonionic in character and can be employed as emulsifying agents, wetting agents, suspending agents, or for other purposes. Surfactants which may also be used in the present formulations
15 are described, *inter alia*, in "McCutcheon's Detergents and Emulsifiers Annual," MC Publishing Corp., Ridgewood, New Jersey, 1998 and in "Encyclopedia of Surfactants," Vol. I-III, Chemical Publishing Co., New York, 1980-81. Surface-active agents include, but are not limited to salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene
20 oxide addition products, such as nonylphenol-C₁₈ ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C₁₆ ethoxylate; soaps, such as sodium stearate; alkyl-naphthalene-sulfonate salts, such as sodium dibutyl-naphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride; polyethylene
25 glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; salts of mono and dialkyl phosphate esters; vegetable or seed oils such as soybean oil, rapeseed/canola oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; and esters of the above vegetable oils, and in certain embodiments,
30 methyl esters.

In some embodiments, these materials, such as vegetable or seed oils and their esters, can be used interchangeably as an agricultural adjuvant, as a liquid carrier or as a surface active agent.

Other exemplary additives for use in the compositions provided herein include but are not limited to compatibilizing agents, antifoam agents, sequestering agents, neutralizing agents and buffers, corrosion inhibitors, dyes, odorants, spreading agents, penetration aids, sticking agents, dispersing agents, thickening agents, freezing point depressants, antimicrobial agents, and the like. The compositions may also contain other compatible components, for example, other herbicides, plant growth regulants, fungicides, insecticides, and the like and can be formulated with liquid fertilizers or solid, particulate fertilizer carriers such as ammonium nitrate, urea and the like.

In some embodiments, the concentration of the active ingredients in the compositions described herein is from 0.0005 to 98 percent by weight. In some embodiments, the concentration is from 0.0006 to 90 percent by weight. In compositions designed to be employed as concentrates, the active ingredients, in certain embodiments, are present in a concentration from 0.1 to 98 weight percent, and in certain embodiments 0.2 to 90 weight percent. Such compositions are, in certain embodiments, diluted with an inert carrier, such as water, before application. The diluted compositions usually applied to weeds or the locus of weeds contain, in certain embodiments, 0.0052 to 25.0 weight percent active ingredient and in certain embodiments contain 0.01 to 15 weight percent active ingredient.

The present compositions can be applied to weeds or their locus by the use of conventional ground or aerial dusters, sprayers, and granule applicators, by addition to irrigation or paddy water, and by other conventional means known to those skilled in the art.

The described embodiments and following examples are for illustrative purposes and are not intended to limit the scope of the claims. Other modifications, uses, or combinations with respect to the compositions described herein will be apparent to a person of ordinary skill in the art without departing from the spirit and scope of the claimed subject matter.

Examples

Results in Examples I, II, III, IV, and V are greenhouse trial results.


Example I. Evaluation of Postemergence Foliar-Applied Herbicidal Mixtures for Weed Control in Direct Seeded Rice

- 5 Seeds or nutlets of the desired test plant species were planted in a soil matrix prepared by mixing a loam or sandy loam soil (*e.g.*, 28.6 percent silt, 18.8 percent clay, and 52.6 percent sand, with a pH of about 5.8 and an organic matter content of about 1.8 percent) and calcareous grit in an 80 to 20 ratio. The soil matrix was contained in plastic pots with a volume of 1 quart and a surface area of 83.6 square centimeters (cm²). When required to
- 10 ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 8-22 days in a greenhouse with an approximate 14 h photoperiod which was maintained at about 29° C during the day and 26° C during the night. Nutrients (Peters Excel[®] 15-5-15 5-Ca 2-Mg and iron chelate) were applied in the irrigation solution as needed and water was added on a regular basis.
- 15 Supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the first through fourth true leaf stage.



Treatments consisted of the acid or esters of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxy-phenyl) pyridine-2-carboxylic acid (Compound A), each formulated as an

20 SC (suspension concentrate), and various herbicidal components alone and in combination. Forms of compound A were applied on an acid equivalent basis.

Forms of compound A (compound of formula I) tested include:



Compound A Acid

Compound A *n*-Butyl Ester

Compound A Benzyl Ester

5

Other herbicidal components were applied on an active ingredient basis and included Photosystem II (PSII)-inhibiting herbicides propanil formulated as Stam[®] M4 EC, bentazon-sodium formulated as Basagran[®], atrazine formulated as Atrazine 90 WDG, bromoxynil octanoate ester formulated as Buctril[®], simetryne (technical grade material), simazine formulated as Princep[®] Caliber 90, ioxynil (technical grade material), cyanazine (technical grade material), and hexazinone (technical grade material).

Treatment requirements were calculated based upon the rates being tested, the concentration of active ingredient or acid equivalent in the formulation, and a 12 mL application volume at a rate of 187 L/ha.

For treatments comprised of formulated compounds, measured amounts of compounds were placed individually in 25 mL glass vials and diluted in a volume of 1.25% (v/v) Agri-Dex[®] crop oil concentrated to obtain 12X stock solutions. If a test compound did not dissolve readily, the mixture was warmed and/or sonicated. Application solutions were prepared by adding an appropriate amount of each stock solution (*e.g.*, 1 mL) and diluted to the appropriate final concentrations with the addition of 10 mL of an aqueous mixture of

1.25% (v/v) crop oil concentrate so that the final spray solutions contained 1.25 $\pm$ 0.05% (v/v) crop oil concentrate.

For treatments comprised of technical compounds, weighed amounts can be placed individually in 25 mL glass vials and dissolved in a volume of 97:3 v/v acetone/DMSO to obtain 12X stock solutions. If a test compound does not dissolve readily, the mixture can be warmed and/or sonicated. Application solutions can be prepared by adding an appropriate amount of each stock solution (*e.g.*, 1 mL) and diluted to the appropriate final concentrations with the addition of 10 mL of an aqueous mixture of 1.5% (v/v) crop oil concentrate so that the final spray solutions contain 1.25% (v/v) crop oil concentrate. When technical materials are used, the concentrated stock solutions can be added to the spray solutions so that the final acetone and DMSO concentrations of the application solutions are 16.2% and 0.5%, respectively.

For treatments comprised of formulated and technical compounds, weighed amounts of the technical materials were placed individually in 25 mL glass vials and dissolved in a volume of 97:3 v/v acetone/DMSO to obtain 12X stock solutions, and measured amounts of the formulated compounds were placed individually in 25 mL glass vials and diluted in a volume of 1.5% (v/v) crop oil concentrate or water to obtain 12X stock solutions. If a test compound did not dissolve readily, the mixture was warmed and/or sonicated. Application solutions were prepared by adding an appropriate amount of each stock solution (*eg*, 1 mL) and diluted to the appropriate final concentrations with the addition of an appropriate amount of an aqueous mixture of 1.5% (v/v) crop oil concentrate so that the final spray solutions contained 1.25% (v/v) crop oil concentrate. As required, additional water and/or 97:3 v/v acetone/DMSO was added to individual application solutions so that the final acetone and DMSO concentrations of the application solutions being compared were 8.1% and 0.25%, respectively.

All stock solutions and applications solutions were visually inspected for compound compatibility prior to application. Spray solutions were applied to the plant material with an overhead Mandel track sprayer equipped with a 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 m² at a spray height of 18 to 20 inches (46 to 50 cm) above average plant canopy height. Control plants were sprayed in the same manner with the solvent blank.

The treated plants and control plants were placed in a greenhouse as described above and watered by sub-irrigation to prevent wash-off of the test compounds. After approximately 3 weeks, the condition of the test plants as compared with that of the untreated plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

The compounds tested, application rates employed, plant species tested, and results are given in Tables 1-14.

Table 1. Synergistic Activity of Foliar-Applied Compound A Acid and Propanil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Propanil	Visual Weed Control (%) - 21 DAA					
		ECHCG		ECHCO		IPOHE	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp	Obs	Exp
4.38	0	33	-	43	-	15	-
8.75	0	55	-	63	-	35	-
0	420	10	-	10	-	0	-
0	840	10	-	25	-	0	-
0	1680	40	-	25	-	0	-
4.38	420	90	39	75	48	25	15
8.75	420	95	60	85	66	50	35
4.38	840	90	39	75	57	50	15
8.75	840	95	60	90	72	65	35
4.38	1680	95	60	90	57	60	15
8.75	1680	95	73	100	72	55	35

Compound A Acid	Propanil	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
19.4	0	15	-
0	3360	40	-
19.4	3360	73	49

Table 2. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Propanil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Propanil	Visual Weed Control (%) - 21 DAA					
		ECHCG		ECHCO		IPOHE	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp	Obs	Exp
4.38	0	48	-	60	-	0	-
8.75	0	55	-	68	-	15	-
17.5	0	85	-	83	-	38	-
0	420	10	-	10	-	0	-
0	840	10	-	25	-	0	-
0	1680	40	-	25	-	0	-
4.38	420	60	53	75	64	15	0
8.75	420	90	60	90	71	50	15
17.5	420	95	87	95	84	60	38
4.38	840	90	53	85	70	20	0
8.75	840	95	60	80	76	35	15
17.5	840	95	87	99	87	65	38
4.38	1680	70	69	100	70	50	0
8.75	1680	90	73	100	76	50	15
17.5	1680	95	91	100	87	65	38

5

Compound A Benzyl Ester	Propanil	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
16	0	18	-
0	3360	40	-
16	3360	85	51

Compound A Benzyl Ester	Propanil	Visual Weed Control (%) - 20 DAA	
		COMBE	
g ae/ha	g ai/ha	Obs	Exp
6	0	80	-
24	0	0	-
0	3360	10	-
6	3360	80	80
24	3360	95	82

Table 3. Synergistic Activity of Foliar-Applied Compound A n-Butyl Ester and Propanil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A n-Butyl Ester	Propanil	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
16	0	15	-
0	3360	40	-
16	3360	70	49

- 5 Table 4. Synergistic Activity of Foliar-Applied Compound A Acid and Bentazon-sodium Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Bentazon- sodium	Visual Weed Control (%) - 21 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	15	-
0	105	0	-
0	210	0	-
0	420	10	-
4.38	105	30	15
4.38	210	60	15
4.38	420	50	24

Compound A Acid	Bentazon- sodium	Visual Weed Control (%) - 20 DAA	
--------------------	---------------------	--	--

		IPOHE	
g ac/ha	g acha	Obs	Exp
21.2	0	35	-
42.4	0	55	-
0	840	13	-
21.2	840	73	43
42.4	840	80	61

Table 5. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Bentazon-sodium Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Bentazon-sodium	Visual Weed Control (%) - 21 DAA	
		IPOHE	
g ac/ha	g ai/ha	Obs	Exp
4.38	0	0	-
8.75	0	15	-
17.5	0	38	-
0	105	0	-
0	210	0	-
0	420	10	-
4.38	105	40	0
8.75	105	65	15
17.5	105	60	38
4.38	210	30	0
8.75	210	40	15
17.5	210	65	38
4.38	420	30	10
8.75	420	60	24
17.5	420	65	44

5

Compound A Benzyl Ester	Bentazon-sodium	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ac/ha	g acha	Obs	Exp
17.5	0	38	-
35	0	55	-
0	840	13	-
17.5	840	60	45
35	840	78	61

Compound A Benzyl Ester	Bentazon sodium	Visual Weed Control (%) - 20 DAA	
		COMBE	
g ae/ha	g ai/ha	Obs	Exp
6	0	70	-
24	0	80	-
0	280	10	-
6	280	75	73
24	280	95	82

Table 6. Synergistic Activity of Foliar-Applied Compound A n-Butyl Ester and Bentazon-
5 sodium Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A n-Butyl Ester	Bentazon- sodium	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ae/ha	g aecha	Obs	Exp
17.5	0	48	-
0	840	13	-
17.5	840	85	54

Table 7. Synergistic Activity of Foliar-Applied Compound A Acid and Atrazine Herbicidal
10 Compositions on Control of Weeds Common to Rice Cropping Systems.

Compound A Acid	Atrazine	Visual Weed Control (%) - 22 DAA	
		BRAPP	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	45	-
8.75	0	70	-
0	280	0	-
0	560	0	-
4.38	280	60	45
8.75	280	85	70
4.38	560	70	45

8.75	560	75	70
------	-----	----	----

Compound A Acid	Atrazine	Visual Weed Control (%) - 22 DAA					
		ECHCG		ECHCO		CYPPIR	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp	Obs	Exp
4.38	0	20	-	20	-	15	-
0	280	0	-	10	-	0	-
0	560	20	-	10	-	100	-
0	1120	70	-	60	-	80	-
4.38	280	50	20	50	28	100	15
4.38	560	80	36	70	28	100	100
4.38	1120	100	76	99	68	100	83

Table 8. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Atrazine Herbicidal Compositions on Control of Weeds Common to Rice Cropping Systems.

Compound A Benzyl Ester	Atrazine	Visual Weed Control (%) - 22 DAA	
		BRAPP	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	55	-
8.75	0	70	-
17.5	0	75	-
0	280	0	-
0	560	0	-
4.38	280	55	55
8.75	280	80	70
17.5	280	85	75
4.38	560	65	55
8.75	560	85	70
17.5	560	95	75

5 Table 9. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Bromoxynil Herbicidal Compositions on Control of Weeds Common to Rice Cropping Systems.

Compound A Benzyl Ester	Bromoxynil Octanoate Ester	Visual Weed Control (%) - 22 DAA	
		ECHCG	
g ae/ha	g ai/ha	Obs	Exp
8	0	35	-
16	0	65	-
32	0	80	-
0	140	0	-
8	140	50	35
16	140	80	65
32	140	90	80

10 Table 10. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Simetryne Herbicidal Compositions on Control of Weeds Common to Rice Cropping Systems.

Compound A Benzyl Ester	Simetryne	Visual Weed Control (%) - 21 DAA
-------------------------------	-----------	--

		ECHCO	
g ae/ha	g ai/ha	Obs	Exp
8	0	75	-
16	0	85	-
32	0	95	-
0	450	45	-
8	450	100	86
16	450	95	92
32	450	100	97

Compound A Benzyl Ester	Simetryne	Visual Weed Control (%) - 22 DAA	
		LEFCH	
g ae/ha	g ai/ha	Obs	Exp
8	0	35	-
16	0	40	-
0	225	40	-
8	225	80	61
16	225	70	64

Compound A Benzyl Ester	Simetryne	Visual Weed Control (%) - 22 DAA	
		CYPIR	
g ae/ha	g ai/ha	Obs	Exp
8	0	10	-
16	0	60	-
0	112.5	0	-
0	225	0	-
0	450	30	-
8	112.5	100	10
16	112.5	90	60
8	225	60	10
16	225	100	60
8	450	100	37
16	450	100	72

Table 11. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Simazine

5 Herbicidal Compositions on Control of Weeds Common to Rice Cropping Systems.

Compound A Benzyl Ester	Simazine	Visual Weed Control (%) - 20 DAA							
		BRAPP		ECHCG		ECHCO		CYPIR	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
8	0	55	-	70	-	65	-	70	-
16	0	90	-	90	-	85	-	90	-
32	0	90	-	90	-	95	-	90	-
0	560	0	-	0	-	0	-	0	-
0	1120	0	-	0	-	0	-	0	-
0	2240	0	-	0	-	0	-	30	-
8	560	80	55	95	70	80	65	100	70
16	560	90	90	95	90	85	85	85	90
32	560	95	90	99	90	95	95	100	90
8	1120	80	55	85	70	75	65	100	70
16	1120	80	90	100	90	85	85	100	90
32	1120	95	90	99	90	95	95	100	90
8	2240	60	55	95	70	80	65	100	79
16	2240	95	90	95	90	90	85	100	93
32	2240	95	90	90	90	99	95	100	93

Table 12. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Cyanazine Herbicidal Compositions on Control of Weeds Common to Rice Cropping Systems.

Compound A Benzyl Ester	Cyanazine	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
8	0	20	-
16	0	40	-
32	0	55	-
0	275	50	-
8	275	80	60
16	275	90	70
32	275	80	78

5

Table 13. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Ioxynil Herbicidal Compositions on Control of Weeds Common to Rice Cropping Systems.

Compound A Benzyl Ester	Ioxynil	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
8	0	0	-
16	0	15	-
32	0	55	-
0	300	50	-
8	300	50	50
16	300	90	58
32	300	90	78

Table 14. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Hexazinone Herbicidal Compositions on Control of Weeds Common to Rice Cropping Systems.

Compound A Benzyl Ester	Hexazinone	Visual Weed Control (%) - 21 DAA	
		ECHCO	
g ae/ha	g ai/ha	Obs	Exp
8	0	50	-
16	0	65	-
32	0	85	-
0	625	25	-
0	1250	55	-
8	625	100	63
16	625	100	74
32	625	100	89
8	1250	90	78
16	1250	100	84
32	1250	100	93

5

Compound A Benzyl Ester	Hexazinone	Visual Weed Control (%) - 21 DAA			
		ECHCG		IPOHE	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp
8	0	45	-	10	-
16	0	85	-	25	-
32	0	90	-	60	-
0	625	20	-	80	-

8	625	85	56	90	82
16	625	90	88	99	85
32	625	90	92	100	92

BRAPP	<i>Brachiaria platyphylla</i> (Griseb.) Nash	signalgrass, broadleaf
COMBE	<i>Commelina benghalensis</i> L.	dayflower, Benghal
CYPIR	<i>Cyperus iria</i> L.	flatsedge, rice
5 CYPES	<i>Cyperus esculentus</i> L.	nutsedge, yellow
ECHCG	<i>Echinochloa crusgalli</i> (L.) Beauv.	barnyardgrass
ECHCO	<i>Echinochloa colona</i> (L.) Link	junglerice
IPOHE	<i>Ipomoea hederacea</i> Jacq.	morningglory, ivyleaf
LEFCH	<i>Leptochloa chinensis</i> (L.) Nees	sprangletop, Chinese

10 g ae/ha = grams acid equivalent per hectare

g ai/ha = grams active ingredient per hectare

Obs = observed value

Exp = expected value as calculated by Colby's equation

DAA = days after application


15

Example II. Evaluation of In-Water Applied Herbicidal Mixtures for Weed Control in Transplanted Paddy Rice.


Weed seeds or nutlets of the desired test plant species were planted in puddled soil (mud) prepared by mixing a shredded, non-sterilized mineral soil (50.5 percent silt, 25.5 percent clay, and 24 percent sand, with a pH of about 7.6 and an organic matter content of about 2.9 percent) and water at a 1:1 volumetric ratio. The prepared mud was dispensed in 365 mL aliquots into 16-ounce (oz.) non-perforated plastic pots with a surface area of 86.59 square centimeters (cm²) leaving a headspace of 3 centimeters (cm) in each pot. Mud was allowed to dry overnight prior to planting or transplanting. Rice seeds were planted in Sun Gro MetroMix[®] 306 planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of about 30 percent, in plastic plug trays. Seedlings at the second or third leaf stage of growth were transplanted into 840 mL of mud contained in 32-oz. non-perforated plastic pots with a surface area of 86.59 cm² 4 days prior to herbicide application. The paddy was created by filling the headspace of the pots with 2.5 to 3 cm of water. When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or


physical treatment was applied. The plants were grown for 4-22 days in a greenhouse with an approximate 14 h photoperiod which was maintained at about 29°C during the day and 26°C during the night. Nutrients were added as Osmocote® (19:6:12, N:P:K + minor nutrients) at 2 g per 16-oz. pot and 4 g per 32-oz. pot. Water was added on a regular basis to maintain the paddy flood, and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the first through fourth true leaf stage.

Treatments consisted of the acid or esters of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (compound A) each formulated as an SC (suspension concentrate) and various herbicidal components alone and in combination. Forms of compound A were applied on an acid equivalent basis. Forms of compound A (compound of formula I) tested include:



Compound A Acid

Compound A *n*-Butyl Ester



Compound A Benzyl Ester

Other herbicidal components were applied on an active ingredient basis and included
5 photosystem II (PSII)-inhibits propanil formulated as Stam[®] 4M EC bentazon-sodium
formulated as Basagran[®], and simetryne (technical grade material).

Treatment requirements for each compound or herbicidal component were calculated
based upon the rates being tested, the concentration of active ingredient or acid equivalent in
the formulation, an application volume of 2 mL per component per pot, and an application
10 area of 86.59 cm² per pot.

For formulated compounds, a measured amount was placed in an individual 100 or
200 mL glass vial and was dissolved in a volume of 1.25% (v/v) Agri-Dex[®] crop oil
concentrate to obtain application solutions. If the test compound did not dissolve readily, the
mixture was warmed and/or sonicated.

15 For technical grade compounds, a weighed amount was placed in an individual 100 to
200 mL glass vial and was dissolved in a volume of acetone to obtain concentrated stock
solutions. If the test compound did not dissolve readily, the mixture was warmed and/or
sonicated. The concentrated stock solutions obtained were diluted with an equivalent volume
of an aqueous mixture containing 2.5% (v/v) crop oil concentrate so that the final application
20 solutions contained 1.25% (v/v) crop oil concentrate.

Applications were made by injecting with a pipetter appropriate amounts of the
application solutions, individually and sequentially, into the aqueous layer of the paddy.
Control plants were treated in the same manner with the solvent blank. Applications were
made so that all treated plant material received the same concentrations of acetone and crop
25 oil concentrate.

The treated plants and control plants were placed in a greenhouse as described above
and water was added as needed to maintain a paddy flood. After approximately 3 weeks the

condition of the test plants as compared with that of the untreated plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury or growth inhibition and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

Some of the compounds tested, application rates employed, plant species tested, and results are given in Tables 15-20.

Table 15. Synergistic Activity of In-Water Applications of Compound A Acid and Propanil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Propanil	Visual Weed Control (%) - 21 DAA	
		SCPMA	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	0	-
17.5	0	0	-
35	0	0	-
0	1680	50	-
0	3360	90	-
8.75	1680	50	50
17.5	1680	100	50
35	1680	100	50
8.75	3360	100	90
17.5	3360	100	90
35	3360	100	90

Table 16. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Propanil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Propanil	Visual Weed Control (%) - 21 DAA	
		SCPMA	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	0	-
17.5	0	0	-
35	0	0	-
0	1680	50	-
0	3360	90	-
8.75	1680	99	50
17.5	1680	100	50
35	1680	70	50
8.75	3360	100	90
17.5	3360	100	90
35	3360	100	90

- 5 Table 17. Synergistic Activity of In-Water Applications of Compound A Acid and Bentazon-sodium Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Bentazon-sodium	Visual Weed Control (%) - 21 DAA			
		ECHCG		LEFCH	
g ae/ha	g ae/ha	Obs	Exp	Obs	Exp
8.75	0	0	-	0	-
17.5	0	0	-	0	-
0	420	0	-	0	-
0	840	0	-	0	-
8.75	420	30	0	20	0
17.5	420	20	0	30	0
8.75	840	10	0	30	0
17.5	840	25	0	40	0

Compound A Acid	Bentazon-sodium	Visual Weed Control (%) - 20 DAA			
		LEFCH		SCPMA	
g ae/ha	g ae/ha	Obs	Exp	Obs	Exp
42.4	0	15	-	0	-
84.8	0	55	-	0	-

0	1120	0	-	0	-
42.4	1120	38	15	80	0
84.8	1120	63	55	100	0

Table 18. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Bentazon-sodium Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Bentazon-sodium	Visual Weed Control (%) - 21 DAA	
		LEFCH	
g ae/ha	g ae/ha	Obs	Exp
8.75	0	20	-
17.5	0	30	-
35	0	55	-
0	420	0	-
0	840	0	-
8.75	420	30	20
17.5	420	50	30
35	420	70	55
8.75	840	50	20
17.5	840	65	30
35	840	50	55

5

Compound A Benzyl Ester	Bentazon-sodium	Visual Weed Control (%) - 20 DAA	
		LEFCH	
g ae/ha	g ae/ha	Obs	Exp
70	0	80	-
0	1120	0	-
70	1120	93	80
Compound A Benzyl Ester	Bentazon-sodium	Visual Weed Control (%) - 20 DAA	
		SCPMA	
g ae/ha	g ae/ha	Obs	Exp
35	0	0	-
70	0	0	-
0	1120	0	-

35	1120	100	0
70	1120	43	0

Table 19. Synergistic Activity of In-Water Applications of Compound A n-Butyl Ester and Bentazon-sodium Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A n-Butyl Ester	Bentazon- sodium	Visual Weed Control (%) - 20 DAA	
		ECHOR	
g ae/ha	g ae/ha	Obs	Exp
70	0	40	-
0	1120	0	-
70	1120	60	40

5

Compound A n-Butyl Ester	Bentazon- sodium	Visual Weed Control (%) - 20 DAA	
		SCPMA	
g ae/ha	g ae/ha	Obs	Exp
35	0	0	-
70	0	0	-
0	1120	0	-
35	1120	100	0
70	1120	50	0

Table 20. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Simetryne Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Simetryne	Visual Weed Control (%) - 19 DAA			
		ECHOR		FIMMI	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp
8	0	10	-	0	-
16	0	20	-	0	-
32	0	25	-	85	-
0	112.5	0	-	45	-
0	225	10	-	90	-
8	112.5	20	10	100	45
16	112.5	50	20	100	45

32	112.5	20	25	100	92
8	225	50	19	100	90
16	225	50	28	100	90
32	225	100	33	100	99

Compound A Benzyl Ester	Simetryne	Visual Weed Control (%) - 19 DAA	
		CYPRO	
g ae/ha	g ai/ha	Obs	Exp
8	0	50	-
16	0	85	-
32	0	90	-
0	112.5	0	-
0	225	0	-
0	450	0	-
8	112.5	90	50
16	112.5	100	85
32	112.5	100	90
8	225	80	50
16	225	85	85
32	225	100	90
8	450	85	50
16	450	100	85
32	450	100	90

CYPRO *Cyperus rotundus* L.

nutsedge, purple

ECHCG *Echinochloa crusgalli* (L.) Beauv.

barnyardgrass

5 ECHOR *Echinochloa oryzoides* (Ard.) Fritsch

watergrass, early

FIMMI *Fimbristylis miliacea* (L.) Vahl

fringerush, globe

LEFCH *Leptochloa chinensis* (L.) Nees

sprangletop, Chinese

SCPMA *Schoenoplectus maritimus* (L.) Lye

clubrush, sea

g ae/ha = grams acid equivalent per hectare

10 g ai/ha = grams active ingredient per hectare

Obs = observed value

Exp = expected value as calculated by Colby's equation


DAA = days after application

Example III. Evaluation of Postemergence Herbicidal Activity of Mixtures in Cereal Crops in the Greenhouse.

Seeds of the desired test plant species were planted in Sun Gro MetroMix[®] 306 planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of about 30 percent, in plastic pots with a surface area of 103.2 square centimeters (cm²). When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 7-36 days in a greenhouse with an approximate 14 hour photoperiod which was maintained at about 18° C during the day and about 17° C during the night. Nutrients and water were added on a regular basis and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the second or third true leaf stage.

Treatments consisted of the benzyl ester of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (Compound A), formulated as an SC, a second cereal herbicide alone and then both in combination.

Forms of compound A (compound of formula I) tested include:



Compound A Benzyl Ester

Other herbicidal components were applied on an active ingredient basis and included PSII-inhibiting herbicides.

Measured aliquots of benzyl ester of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (Compound A) were placed in 25 milliliter (mL) glass vials and diluted in a volume of 1.25% (v/v) Agri-Dex[®] crop oil concentrate to obtain stock solutions. Compound requirements are based upon a 12 mL application volume at a rate of 187 liters per hectare (L/ha). Spray solutions of the second cereal herbicide and experimental compound mixtures were prepared by adding the stock solutions to the appropriate amount of dilution solution to form 12 mL spray solution with active ingredients

in two- and three-way combinations. Formulated compounds were applied to the plant material with an overhead Mandel track sprayer equipped with an 8002E nozzle calibrated to deliver 187 L/ha over an application area of 0.503 square meters (m²) at a spray height of 18 inches (46 cm) above average plant canopy. Control plants were sprayed in the same manner with the solvent blank.

The treated plants and control plants were placed in a greenhouse as described above and watered by sub-irrigation to prevent wash-off of the test compounds. After 20-22 days, the condition of the test plants as compared with that of the control plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

The compounds tested, application rates employed, plant species tested, and results are given in Tables 21-24.

Table 21. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Metribuzin Herbicidal Compositions on Weed Control in a Cereals Cropping System.

Compound A Benzyl Ester	Metribuzin	Visual Weed Control (%) - 21 DAA					
		SASKR		KCHSC		LAMPU	
g ai/ha	g ai/ha	Obs	Exp	Obs	Exp	Obs	Exp
2.5	0	15	-	20	-	80	-
5	0	50	-	41	-	86	-
10	0	67	-	62	-	94	-
0	25	8	-	0	-	3	-
0	50	8	-	8	-	15	-

0	100	67	-	88	-	23	-
2.5	25	25	21	45	20	90	81
2.5	50	20	21	35	26	89	83
5	25	38	54	58	41	91	86
5	50	63	54	43	45	98	88
5	100	100	83	100	93	99	89
10	100	97	89	100	96	100	95

Compound A Benzyl Ester	Metribuzin	Visual Weed Control (%) - 21 DAA			
		CHEAL		SINAR	
g ai/ha	g ai/ha	Obs	Exp	Obs	Exp
2.5	0	50	-	70	-
5	0	58	-	73	-
0	25	0	-	8	-
0	50	55	-	15	-
2.5	25	68	50	90	72
2.5	50	70	78	95	75
5	25	78	58	93	75
5	50	83	81	95	77

Table 22. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Isoproturon Herbicidal Compositions on Weed Control in a Cereals Cropping System.

5

Compound A Benzyl Ester	Isoproturon	Visual Weed Control (%) - 21 DAA			
		AMARE		SINAR	
g ai/ha	g ai/ha	Obs	Exp	Obs	Exp
2.5	0	20	-	70	-
5	0	38	-	73	-
0	375	3	-	13	-
0	750	10	-	40	-
2.5	375	43	22	84	74
2.5	750	60	28	88	82
5	375	55	39	89	76
5	750	70	44	84	84

Compound A Benzyl Ester	Isoproturon	Visual Weed Control (%) - 21 DAA			
-------------------------------	-------------	----------------------------------	--	--	--

		GALAP		VERPE	
g ai/ha	g ai/ha	Obs	Exp	Obs	Exp
5	0	66	-	17	-
10	0	77	-	23	-
0	375	5	-	8	-
0	750	5	-	10	-
0	1500	27	-	13	-
5	375	68	68	18	23
5	750	80	68	28	25
5	1500	99	75	60	28
10	1500	97	83	70	34

Compound A Benzyl Ester	Isoproturon	Visual Weed Control (%) - 21 DAA	
		LAMP	
g ai/ha	g ai/ha	Obs	Exp
2.5	0	80	-
5	0	86	-
10	0	94	-
0	375	3	-
0	750	3	-
0	1500	13	-
2.5	375	83	81
2.5	750	89	81
5	375	89	86
5	750	91	86
5	1500	97	88
10	1500	97	95

Table 23. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Bentazon-sodium Herbicidal Compositions on Weed Control in a Cereals Cropping System.

Compound A Benzyl Ester	Bentazon-Sodium	Visual Weed Control (%) - 21 DAA	
		IPOHE	
g ai/ha	g ai/ha	Obs	Exp
17.5	0	33	-
0	60	7	-
0	120	10	-

17.5	60	67	38
17.5	120	67	40

Compound A Benzyl Ester	Bentazon- Sodium	Visual Weed Control (%) - 21 DAA			
		SETFA		BRSNW	
g ai/ha	g ai/ha	Obs	Exp	Obs	Exp
8.75	0	13	-	0	-
0	120	0	-	0	-
8.75	120	63	13	28	0

Table 24. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Bromoxynil Herbicidal Compositions on Weed Control in a Cereals Cropping System.

Compound A Benzyl Ester	Bromoxynil	Visual Weed Control (%) - 21 DAA			
		AMARE		IPOHE	
g ai/ha	g ai/ha	Obs	Exp	Obs	Exp
8.75	0	57	-	33	-
17.5	0	85	-	33	-
0	35	3	-	50	-
0	70	17	-	35	-
8.75	35	75	58	42	67
8.75	70	88	64	63	57
17.5	35	98	86	60	67
17.5	70	99	88	75	57

5

Compound A Benzyl Ester	Bromoxynil	Visual Weed Control (%) - 21 DAA	
		SETFA	
g ai/ha	g ai/ha	Obs	Exp
8.75	0	13	-
0	70	0	-
8.75	70	63	13

	AMARE	<i>Amaranthus retroflexus</i> L.	pigweed, redroot
	BRSNW	<i>Brassica napus</i> L.	rape, oilseed (winter)
	CHEAL	<i>Chenopodium album</i> L.	lambsquarters, common
	GALAP	<i>Galium aparine</i> L.	bedstraw, catchweed
5	IPOHE	<i>Ipomoea hederacea</i> Jacq.	morningglory, ivyleaf
	KCHSC	<i>Kochia scoparia</i> (L.) Schrad.	kochia
	LAMPU	<i>Lamium purpureum</i> L.	deadnettle, purple
	SASKR	<i>Salsola tragus</i> L.	thistle, Russian
	SETFA	<i>Setaria faberi</i> Herrm.	foxtail, giant
10	SINAR	<i>Sinapis arvensis</i> L.	mustard, wild
	VERPE	<i>Veronica persica</i> Poir.	Speedwell, Persian

g ae/ha = grams acid equivalent per hectare

g ai/ha = grams active ingredient per hectare

Obs = observed value

15 Exp = expected value as calculated by Colby's equation

DAA = days after application

Example IV. Evaluation of Pre-emergence Soil-Applied Herbicidal Mixtures for Weed

Control

20

Seeds or nutlets of the desired test plant species were planted in a soil matrix prepared by mixing a loam soil (32 percent silt, 23 percent clay, and 45 percent sand, with a pH of about 6.5 and an organic matter content of about 1.9 percent) and calcareous grit in an 80 to 20 ratio. The soil matrix was contained in plastic pots with a volume of 1 quart and a surface area of 83.6 square centimeters (cm²).


25

Treatments consisted of the benzyl ester of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (compound A) formulated as an SC (suspension concentrate) and various herbicidal components alone and in combination.

Forms of compound A were applied on an acid equivalent basis.

30

Forms of compound A (compound of formula I) tested include:



Compound A Benzyl Ester

Other herbicidal components were applied on an acid equivalent or active ingredient basis and included the PSII-inhibiting herbicides, atrazine formulated as Atrazine[®] 90WDG, metribuzin formulated as Metribuzin 75DF, simazine formulated as Princep Caliber 90, tebuthiuron formulated as Spike 80DF, cyanazine (technical grade material), hexazinone (technical grade material), linuron (technical grade material), terbuthylazine (technical grade material), and diuron formulated as Diuron 4L .

Treatment requirements were calculated based upon the rates being tested, the concentration of active ingredient or acid equivalent in the formulation, and a 12 mL application volume at a rate of 187 L/ha.

For treatments comprised of formulated compounds, measured amounts of compounds were placed individually in 25 mL glass vials and diluted in a volume of 1.25% (v/v) Agri-Dex[®] crop oil concentrate (COC) to obtain 12X stock solutions. If a test compound did not dissolve readily, the mixture was warmed and/or sonicated. Application solutions were prepared by adding an appropriate amount of each stock solution (e.g., 1 mL) and diluted to the appropriate final concentrations with the addition of 10 mL of an aqueous mixture of 1.25% (v/v) COC so that the final spray solutions contained 1.25% (v/v) COC.

For treatments comprised of technical compounds, weighed amounts can be placed individually in 25 mL glass vials and dissolved in a volume of 97:3 (v/v) acetone/DMSO to obtain 12X stock solutions. If a test compound does not dissolve readily, the mixture can be warmed and/or sonicated. Application solutions can be prepared by adding an appropriate amount of each stock solution (e.g., 1 mL) and diluted to the appropriate final concentrations with the addition of 10 mL of an aqueous mixture of 1.5% (v/v) COC so that the final spray solutions contain 1.25% (v/v) COC. When technical materials are used, the concentrated

stock solutions can be added to the spray solutions so that the final acetone and DMSO concentrations of the application solutions are 16.2% and 0.5%, respectively.

For treatments comprised of formulated and technical compounds, weighed amounts of the technical materials can be placed individually in 25 mL glass vials and dissolved in a volume of 97:3 (v/v) acetone/DMSO to obtain 12X stock solutions, and measured amounts of the formulated compounds can be placed individually in 25 mL glass vials and diluted in a volume of 1.5% (v/v) COC or water to obtain 12X stock solutions. If a test compound does not dissolve readily, the mixture can be warmed and/or sonicated. Application solutions can be prepared by adding an appropriate amount of each stock solution (*e.g.*, 1 mL) and diluted to the appropriate final concentrations with the addition of an appropriate amount of an aqueous mixture of 1.5% (v/v) COC so that the final spray solutions contain 1.25% (v/v) COC. As required, additional water and/or 97:3 (v/v) acetone/DMSO can be added to individual application solutions so that the final acetone and DMSO concentrations of the application solutions being compared are 8.1% and 0.25%, respectively.

All stock solutions and applications solutions were visually inspected for compound compatibility prior to application. Spray solutions were applied to the soil with an overhead Mandel track sprayer equipped with a 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 m² at a spray height of 18 inches (46 cm) above average pot height. Control pots were sprayed in the same manner with the solvent blank.

The treated and control pots were placed in a greenhouse and top watered as needed. When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The pots were maintained in a greenhouse with an approximate 14 hr photoperiod which was maintained at about 29° C during the day and 26° C during the night. Nutrients (Peters[®] Excel 15-5-15 5-Ca 2-Mg) were applied in the irrigation solution as needed and water was added on a regular basis. Supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. After approximately 4 weeks, the condition of the test plants as compared with that of the untreated plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury or growth inhibition and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

Some of the compounds tested, application rates employed, plant species tested, and results are given in Tables 25-32.

Table 25. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Atrazine Herbicidal Compositions on Weed Control

Compound A Benzyl Ester	Atrazine	Visual Weed Control (%) - 32 DAA	
		XANST	
g ae/ha	g ai/ha	Obs	Exp
8	0	33	-
16	0	35	-
32	0	65	-
0	560	68	-
0	1120	40	-
8	560	95	78
16	560	100	79
32	560	75	89
8	1120	100	60
16	1120	100	61
32	1120	100	79

Table 26. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Metribuzin Herbicidal Compositions on Weed Control

Compound A Benzyl Ester	Metribuzin	Visual Weed Control (%) - 27 DAA	
		SETFA	
g ae/ha	g ai/ha	Obs	Exp
16	0	8	-
32	0	15	-
0	210	43	-
0	420	83	-
16	210	100	48
32	210	100	52
16	420	95	85
32	420	100	86

5 Table 27. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Simazine Herbicidal Compositions on Weed Control

Compound A Benzyl Ester	Simazine	Visual Weed Control (%) - 33 DAA	
		XANST	
g ae/ha	g ai/ha	Obs	Exp
16	0	38	-
32	0	35	-
0	1120	70	-
0	2240	92	-
16	1120	77	82
32	1120	95	81
16	2240	100	95
32	2240	100	95

10 Table 28. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Tebuthiuron Herbicidal Compositions on Weed Control

Compound A Benzyl Ester	Tebuthiuron	Visual Weed Control (%) - 33 DAA
		IPOHE

g ae/ha	g ai/ha	Obs	Exp
16	0	0	-
32	0	5	-
0	420	87	-
0	840	90	-
16	420	93	87
32	420	93	87
16	840	100	90
32	840	100	91

Compound A Benzyl Ester	Tebuthiuron	Visual Weed Control (%) - 29 DAA	
		XANST	
g ae/ha	g ai/ha	Obs	Exp
16	0	0	-
32	0	5	-
0	210	40	-
0	3360	95	-
16	210	63	40
32	210	55	43
16	3360	100	95
32	3360	100	95

Table 29. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Cyanazine Herbicidal Compositions on Weed Control

5

Compound A Benzyl Ester	Cyanazine	Visual Weed Control (%) - 27 DAA	
		BRAPP	
g ae/ha	g ai/ha	Obs	Exp
32	0	0	-
0	170	0	-
0	440	15	-
0	880	10	-
32	170	10	0
32	440	28	15
32	880	40	10

Table 30. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Linuron Herbicidal Compositions on Weed Control

Compound A Benzyl Ester	Linuron	Visual Weed Control (%) - 27 DAA	
		BRAPP	
g ae/ha	g ai/ha	Obs	Exp
16	0	0	-
32	0	0	-
0	620	0	-
16	620	15	0
32	620	20	0

Table 31. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Terbutylazine Herbicidal Compositions on Weed Control

5

Compound A Benzyl Ester	Terbutylazine	Visual Weed Control (%) - 27 DAA	
		XANST	
g ae/ha	g ai/ha	Obs	Exp
16	0	10	-
32	0	25	-
0	1000	45	-
16	1000	83	51
32	1000	75	59

Table 32. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Diuron Herbicidal Compositions on Weed Control

Compound A Benzyl Ester	Diuron	Visual Weed Control (%) - 29 DAA	
		ABUTH	
g ae/ha	g ai/ha	Obs	Exp
16	0	65	-
32	0	93	-
0	280	30	-
0	560	70	-
0	1120	90	-
16	280	90	76
32	280	100	95
16	560	100	90

32	560	100	98
16	1120	100	97
32	1120	100	99

Compound A Benzyl Ester	Diuron	Visual Weed Control (%) - 28 DAA			
		BRAPP		ECHCG	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp
16	0	0	-	0	-
32	0	5	-	0	-
0	280	0	-	25	-
0	560	35	-	65	-
16	280	45	0	35	25
32	280	25	5	23	25
16	560	100	35	80	65
32	560	100	38	98	65

Compound A Benzyl Ester	Diuron	Visual Weed Control (%) - 29 DAA					
		ECHCO		SETFA		SETVI	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp	Obs	Exp
16	0	0	-	0	-	10	-
32	0	5	-	0	-	10	-
0	280	15	-	5	-	0	-
16	280	60	15	13	5	48	10
32	280	80	19	30	5	25	10

ABUTH	<i>Abutilon theophrasti</i> Medik.	velvetleaf
5 BRAPP	<i>Brachiaria platyphylla</i> (Groseb.) Nash or <i>Urochloa platyphylla</i> (Nash) R.D. Webster	signalgrass, broadleaf
ECHCG	<i>Echinochloa crus-galli</i> (L.) P. Beauv.	barnyardgrass
ECHCO	<i>Echinochloa colonum</i> (L.) Link	junclerice
IPOHE	<i>Ipomoea hederacea</i> (L.) Jacq.	morningglory, ivyleaf
10 SETFA	<i>Setaria faberi</i> Herrm.	foxtail, giant
SETVI	<i>Setaria viridis</i> (L.) Beauv.	foxtail, green
XANST	<i>Xanthium strumarium</i> L.	cocklebur, common

g ae/ha = grams acid equivalent per hectare

g ai/ha = grams active ingredient per hectare

Obs = observed value

Exp = expected value as calculated by Colby's equation

DAA = days after application

5

Example V. Evaluation of Postemergence Foliar-Applied Herbicidal Mixtures for General Weed Control

10 Seeds or root cuttings of the desired test plant species were planted in Sun Gro MetroMix[®] 306 planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of about 30 percent, in plastic pots with a surface area of 126.6 square centimeters (cm²). When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 9-30 days in a greenhouse with an approximate 14 hour photoperiod which was maintained at
15 approximately 28° C during the day and 24° C during the night. Nutrients and water were added on a regular basis and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the BBCH11 to BBCH14 leaf stage.

Treatments consisted of the benzyl ester of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-
20 fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (Compound A), formulated as a SC and a tebuthiuron containing herbicide (Spike[®] 80DF). A measured aliquot of Compound A was placed in 25 milliliter (mL) glass vial and diluted in a volume of distilled water or 1.25% volume/volume (v/v) Agri-dex crop oil concentrated to obtain stock solutions. Compound requirements are based upon a 12 mL application volume at a rate of 187 liters per hectare
25 (L/ha). Spray solutions of the glyphosate-containing herbicide and experimental compound mixtures were prepared by adding the stock solutions to the appropriate amount of dilution solution to form 12 mL spray solution with active ingredients in two-way combinations. Formulated compounds were applied to the plant material with an overhead Mandel track sprayer equipped with 8002E nozzles calibrated to deliver 187 L/ha over an application area
30 of 0.503 square meters (m²) at a spray height of 18 inches (43 cm) above average plant canopy. Control plants were sprayed in the same manner with the solvent blank.

The treated plants and control plants were placed in a greenhouse as described above and watered by sub-irrigation to prevent wash-off of the test compounds. After

approximately 14 days, the condition of the test plants, as compared with that of the control plants, was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

Some of the compounds tested, application rates employed, plant species tested, and results are given in Table 33.

Table 33. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Spike 80DF (Tebuthiuron) Herbicidal Compositions formulated in 1.25% (v/v) crop oil concentrate in water on General Weed Control.


Application Rate (g/ha)		ECHCG		CIRAR		BRSNN		CYPES	
Compound A Benzyl Ester	Spike	Obs	Exp	Obs	Exp	Obs	Exp	Obs	Exp
4	0	30	-	35	-	10	-	0	-
8	0	50	-	40	-	15	-	65	-
16	0	60	-	50	-	30	-	85	-
0	140	0	-	25	-	10	-	0	-
0	280	0	-	30	-	20	-	0	-
0	560	0	-	55	-	15	-	0	-
4	140	75	30	55	51	30	19	85	0
8	140	85	50	75	54	65	23	80	65
16	140	95	60	95	71	70	37	90	85
4	280	80	30	85	55	65	28	80	0
8	280	80	50	90	58	80	32	85	65
16	280	90	60	95	73	95	44	95	85
4	560	85	30	90	62	80	23	60	0
8	560	80	50	95	65	90	44	75	65

16	560	90	60	100	77	100	41	100	85
----	-----	----	----	-----	----	-----	----	-----	----


	ECHCG	<i>Echinochloa crus-galli</i> L.	barnyardgrass
	CIRAR	<i>Cirsium arvense</i> (L.) Scop.	thistle, Canada
	BRSNN	<i>Brassica napus</i> L.	rape, oilseed
5	CYPES	<i>Cyperus esculentus</i> L.	nutsedge, yellow

WHAT IS CLAIMED IS:

1. A synergistic herbicidal composition comprising a herbicidally effective amount of
(a) a compound of the formula (I)



- 5 or an agriculturally acceptable salt or ester thereof and (b) at least one compound, or an agriculturally acceptable salt, carboxylic acid, carboxylate salt, or ester thereof, selected from the group consisting of atrazine, bentazon, bentazon-sodium, bromoxynil, chlorotoluron, cyanazine, diuron, hexazinone, ioxynil, isoproturon, linuron, methibenzuron, metribuzin, propanil, pyridate, siduron, simazine, simetryne, tebuthiuron, and terbuthylazine.
- 10 2. The composition of claim 1, wherein (a) is the compound of formula (I), a C₁₋₄ alkyl ester of the compound of formula (I), or a benzyl ester of the compound of formula (I).
3. The composition of any of claims 1-2, further comprising an agriculturally acceptable adjuvant or carrier.
4. The composition of any of claims 1-3, further comprising a herbicide safener.
- 15 5. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation a herbicidally effective amount of (a) a compound of the formula (I)



- or an agriculturally acceptable salt or ester thereof and (b) is at least one compound, or an agriculturally acceptable salt, carboxylic acid, carboxylate salt, or ester thereof, selected from the group consisting of: atrazine, bentazon, bentazon-sodium, bromoxynil, chlorotoluron,
- 20

cyanazine, diuron, hexazinone, ioxynil, isoproturon, linuron, methibenzuron, metribuzin, propanil, pyridate, siduron, simazine, simetryne, tebuthiuron, and terbuthylazine.

6. The method of claim 5, wherein (a) is the compound of formula (I), a C₁₋₄ alkyl ester of the compound of formula (I), or a benzyl ester of the compound of formula (I).

5 7. The method of any of claims 5-6, wherein the undesirable vegetation is controlled in direct-seeded, water-seeded and transplanted rice, cereals, wheat, barley, oats, rye, sorghum, corn/maize, sugarcane, sunflower, oilseed rape, canola, sugar beet, soybean, cotton, pineapple, pastures, grasslands, rangelands, fallowland, turf, tree and vine orchards, aquatics, industrial vegetation management (IVM) or rights of way (ROW).

10 8. The method of any of claims 5-7, wherein the (a) and (b) are applied pre-emergently to the weed or the crop.

9. The method of any of claims 5-8, wherein the undesirable vegetation is controlled in glyphosate-, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitor-, glufosinate-, glutamine synthetase inhibitor-, dicamba-, phenoxy auxin-, pyridyloxy auxin-, synthetic
15 auxin-, auxin transport inhibitor-, aryloxyphenoxypropionate-, cyclohexanedione-, phenylpyrazoline-, acetyl CoA carboxylase (ACCase) inhibitor-, imidazolinone-, sulfonyleurea-, pyrimidinylthiobenzoate-, triazolopyrimidine-, sulfonaminocarbonyltriazolinone-, acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitors-, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitor-,
20 phytoene desaturase inhibitor-, carotenoid biosynthesis inhibitor-, protoporphyrinogen oxidase (PPO) inhibitor-, cellulose biosynthesis inhibitor-, mitosis inhibitor-, microtubule inhibitor-, very long chain fatty acid inhibitor-, fatty acid and lipid biosynthesis inhibitor-, photosystem I inhibitor-, photosystem II inhibitor-, triazine-, or bromoxynil- tolerant crops.

10. The method of claim 9, wherein the tolerant crop possesses multiple or stacked traits
25 conferring tolerance to multiple herbicides or multiple modes of action.

11. The method of any of claims 5-10, wherein the undesirable vegetation comprises a herbicide resistant or tolerant weed.