

- 1 -

3-AMINO-PYRROLO[3,4-C]PYRAZOLE-5 (1H, 4H, 6H) CARBALDEHYDE
DERIVATIVES AS PKC INHIBITORS

This application claims the benefit of U. S. Provisional Application No. 60/888,749 filed February 7, 2007, U.S Provisional Application No. 60/989,086 filed November 19, 2007, and U. S. Provisional Application No. 61/020,965 filed January 14, 2008.

FIELD OF THE INVENTION

The present invention relates to novel compounds, to pharmaceutical compositions comprising the compounds, as well as to the use of the compounds in medicine and for the preparation of a medicament which acts on the human protein kinase C enzyme, and in particular the beta II isoform (pkc β II).

BACKGROUND OF THE INVENTION

Protein kinase C (PKC) is a superfamily of lipid-activated Ser/Thr kinases involved in multiple signal transduction pathways. There are thirteen PKC-isoforms that have been identified and are classified according to their regulation by cellular signaling molecules such as diacylglycerol, phospholipids, and calcium. The protein kinase C isozymes, alpha, beta (two splice variants PKC β I and PKC β II) and gamma, require membrane phospholipids, calcium and diacylglycerolphorbol esters for full activation. The delta, epsilon, eta, and theta forms of PKC are calcium-independent in their mode of activation. The zeta and lambda forms of PKC are independent of both calcium and diacylglycerol and are believed to require only membrane phospholipids for their activation.

The tissue-specific expression and activation of PKC-isoforms suggests that individual PKC-isoforms might be potential therapeutic targets. For diabetes, activation of PKC-beta has been demonstrated in tissues of diabetic animals and has been implicated in the development of microvascular abnormalities related to the hyperglycemic state. Genetic polymorphisms have been identified in the 5'-flanking upstream region of the PKC β gene in Japanese patients with type II diabetes. This PKC β genetic variation was associated with a significant increase in the susceptibility to develop diabetic vascular complications and macrovascular diseases such as coronary heart disease.


- 2 -

In a large case-control study at the Joslin Diabetes Center, additional polymorphisms were identified in the PKC β promoter region that had an association with type I diabetes mellitus (duration <24 years) and a greater risk for development of diabetic nephropathy. Administration of PKC β inhibitors such as ruboxistaurin mesylate (LY333531, Lilly) in diabetic animal models, was shown to prevent or ameliorate the hemodynamic changes and vascular damage associated with diabetic nephropathy, diabetic peripheral neuropathy, and diabetic retinopathy. Way, K.J. et al, Diabet.Med. 18: 945-959 (2001); Vinik, A., Expert Opin. Investig.Drugs 14: 1547-1559 (2005). Together with additional data from phase II and phase III clinical studies of ruboxistaurin mesylate for treatment of diabetes and diabetic microvascular complications, there is a building body of evidence to support the rationale that PKC β can function as a molecular target for diabetic complications and for the development of selective- PKC β inhibitors as potential therapeutic agents.

The compounds of the present invention are protein kinase C beta II inhibitors, and are therefore believed to be useful in the treatment of conditions associated with diabetes mellitus and its complications, cancer, ischemia, inflammation, central nervous system disorders, cardiovascular disease and dermatological disease.

SUMMARY OF THE INVENTION

The present invention is directed to a compound or pharmaceutically acceptable salt of Formula A,



(A)

wherein

X is C-R¹¹ or N, wherein R¹¹ is H, halo, OH, C₁-C₃alkyl, CF₃, or CN;

- 3 -

A and B are independently C or N;

R^1 , R^2 and R^3 are each independently selected from H, R^a -O- R^b , C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-(R^d)_m$ -(C_3 - C_{12} cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m$ -(C_1 - C_6 perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m$ -C(O) R^a , $-(R^d)_m$ -C(O)OR a , $-(R^d)_m$ -C(O)NR a R b , $-(R^d)_m$ -OR a , $-(R^d)_m$ -OC(O) R^a , $-(R^d)_m$ -OC(O)NR a R b , $-(R^d)_m$ -O-S(O) R^a , $-(R^d)_m$ -OS(O) $_2$ R a , $-(R^d)_m$ -OS(O) $_2$ NR a R b , $-(R^d)_m$ -OS(O)NR a R b , $-(R^d)_m$ -NO $_2$, $-(R^d)_m$ -NR a R b , $-(R^d)_m$ -N(R^a)C(O)R b , $-(R^d)_m$ -N(R^a)C(O)OR b , $-(R^d)_m$ -N(R^c)C(O)NR a R b , $-(R^d)_m$ -N(R^a)S(O) $_2$ R b , $-(R^d)_m$ -N(R^a)S(O)R b , $-(R^d)_m$ -SR a , $-(R^d)_m$ -S(O) R^a , $-(R^d)_m$ -S(O) $_2$ R a , $-(R^d)_m$ -S(O)NR a R b , $-(R^d)_m$ -S(O) $_2$ NR a R b , $-(R^d)_m$ -O-(R^e) $_m$ -NR a R b or $-(R^d)_m$ -NR a -(R^e)-OR b ; wherein R^2 and R^3 may together optionally cyclize to form a saturated or unsaturated 3-7 membered heterocyclyl fused to the 6-membered N-containing heteroaryl to which they are attached; and wherein any of the said alkyl, alkenyl, alkynyl, R^a , R^b , R^c , R^d , R^e , C_3 - C_{12} cycloalkyl, phenyl or 3-15 membered heterocyclyl, may independently be further optionally substituted by 0-3 R^{12} groups;

R^4 and R^5 are each independently selected from H, R^a -O- R^b , C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-(R^d)_m$ -(C_3 - C_{12} cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m$ -(C_1 - C_6 perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m$ -C(O) R^a , $-(R^d)_m$ -C(O)OR a , $-(R^d)_m$ -C(O)NR a R b , $-(R^d)_m$ -OR a , $-(R^d)_m$ -OC(O) R^a , $-(R^d)_m$ -OC(O)NR a R b , $-(R^d)_m$ -O-S(O) R^a , $-(R^d)_m$ -OS(O) $_2$ R a , $-(R^d)_m$ -OS(O) $_2$ NR a R b , $-(R^d)_m$ -OS(O)NR a R b , $-(R^d)_m$ -NO $_2$, $-(R^d)_m$ -NR a R b , $-(R^d)_m$ -N(R^a)C(O)R b , $-(R^d)_m$ -N(R^a)C(O)OR b , $-(R^d)_m$ -N(R^c)C(O)NR a R b , $-(R^d)_m$ -N(R^a)S(O) $_2$ R b , $-(R^d)_m$ -N(R^a)S(O)R b , $-(R^d)_m$ -SR a , $-(R^d)_m$ -S(O) R^a , $-(R^d)_m$ -S(O) $_2$ R a , $-(R^d)_m$ -S(O)NR a R b , $-(R^d)_m$ -S(O) $_2$ NR a R b , $-(R^d)_m$ -O-(R^e) $_m$ -NR a R b or $-(R^d)_m$ -NR a -(R^e)-OR b ; wherein any of the said alkyl, alkenyl, alkynyl, R^a , R^b , R^c , R^d , R^e , C_3 - C_{12} cycloalkyl, aryl or 3-15 membered heterocyclyl are independently optionally further substituted by 0-3 R^{12} groups,

R^6 and R^7 are each independently H, R^a -O- R^b , C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-(R^d)_m$ -(C_3 - C_{12} cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m$ -(C_1 - C_6 perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m$ -C(O) R^a , $-(R^d)_m$ -C(O)OR a , $-(R^d)_m$ -C(O)NR a R b , $-(R^d)_m$ -OR a , $-(R^d)_m$ -OC(O) R^a , $-(R^d)_m$ -OC(O)NR a R b , $-(R^d)_m$ -O-S(O) R^a , $-(R^d)_m$ -OS(O) $_2$ R a , $-(R^d)_m$ -OS(O) $_2$ NR a R b , $-(R^d)_m$ -OS(O)NR a R b , $-(R^d)_m$ -NO $_2$, $-(R^d)_m$ -NR a R b , $-(R^d)_m$ -N(R^a)C(O)R b , $-(R^d)_m$ -N(R^a)C(O)OR b , $-(R^d)_m$ -N(R^c)C(O)NR a R b , $-(R^d)_m$ -N(R^a)S(O) $_2$ R b , $-(R^d)_m$ -N(R^a)S(O)R b , $-(R^d)_m$ -SR a , $-(R^d)_m$ -S(O) R^a , $-(R^d)_m$ -S(O) $_2$ R a , $-(R^d)_m$ -S(O)NR a R b , $-(R^d)_m$ -S(O) $_2$ NR a R b , $-(R^d)_m$ -O-(R^e) $_m$ -NR a R b or $-(R^d)_m$ -NR a -(R^e)-OR b ; wherein

- 4 -

R^6 and R^7 may together optionally cyclize to form a C₃-C₇ cycloalkyl and wherein any of the said alkyl, alkenyl, alkynyl, R^a , R^b , R^c , R^d , R^e , C₃-C₁₂ cycloalkyl, aryl or 3-15 membered heterocyclyl are independently optionally further substituted by 0-3 R^{12} groups;

R^8 is H, R^a -O- R^b , C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, $-(R^d)_m$ -(C₃-C₁₂ cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m$ -(C₁-C₆ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m$ -C(O) R^a , $-(R^d)_m$ -C(O)OR^a, $-(R^d)_m$ -C(O)NR^aR^b, $-(R^d)_m$ -OR^a, $-(R^d)_m$ -OC(O) R^a , $-(R^d)_m$ -OC(O)NR^aR^b, $-(R^d)_m$ -O-S(O) R^a , $-(R^d)_m$ -OS(O)₂ R^a , $-(R^d)_m$ -OS(O)₂NR^aR^b, $-(R^d)_m$ -OS(O)NR^aR^b, $-(R^d)_m$ -NO₂, $-(R^d)_m$ -NR^aR^b, $-(R^d)_m$ -N(R^a)C(O) R^b , $-(R^d)_m$ -N(R^a)C(O)OR^b, $-(R^d)_m$ -N(R^c)C(O)NR^aR^b, $-(R^d)_m$ -N(R^a)S(O)₂ R^b , $-(R^d)_m$ -N(R^a)S(O) R^b , $-(R^d)_m$ -SR^a, $-(R^d)_m$ -S(O) R^a , $-(R^d)_m$ -S(O)₂ R^a , $-(R^d)_m$ -S(O)NR^aR^b, $-(R^d)_m$ -S(O)₂NR^aR^b, $-(R^d)_m$ -O-(R^e)_m-NR^aR^b or $-(R^d)_m$ -NR^a-(R^e)-OR^b; and wherein any of the said alkyl, alkenyl, alkynyl, R^a , R^b , R^c , R^d , R^e , C₃-C₁₂ cycloalkyl, phenyl, or 3-15 membered heterocyclyl are independently optionally further substituted by 1-3 groups selected from -F, C₁-C₃ alkyl, C₁-C₃ perfluoroalkyl, hydroxyl, C₁-C₆alkoxyl, or oxo;

R^9 and R^{10} are each independently C₁-C₂ alkyl or can together cyclize to form a cyclopropyl or cyclobutyl;

each R^{12} is independently H, R^a -O- R^b , C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, $-(R^d)_m$ -(C₃-C₁₂ cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m$ -(C₁-C₆ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m$ -C(O) R^a , $-(R^d)_m$ -C(O)OR^a, $-(R^d)_m$ -C(O)NR^aR^b, $-(R^d)_m$ -OR^a, $-(R^d)_m$ -OC(O) R^a , $-(R^d)_m$ -OC(O)NR^aR^b, $-(R^d)_m$ -O-S(O) R^a , $-(R^d)_m$ -OS(O)₂ R^a , $-(R^d)_m$ -OS(O)₂NR^aR^b, $-(R^d)_m$ -OS(O)NR^aR^b, $-(R^d)_m$ -NO₂, $-(R^d)_m$ -NR^aR^b, $-(R^d)_m$ -N(R^a)C(O) R^b , $-(R^d)_m$ -N(R^a)C(O)OR^b, $-(R^d)_m$ -N(R^c)C(O)NR^aR^b, $-(R^d)_m$ -N(R^a)S(O)₂ R^b , $-(R^d)_m$ -N(R^a)S(O) R^b , $-(R^d)_m$ -SR^a, $-(R^d)_m$ -S(O) R^a , $-(R^d)_m$ -S(O)₂ R^a , $-(R^d)_m$ -S(O)NR^aR^b, $-(R^d)_m$ -S(O)₂NR^aR^b, $-(R^d)_m$ -O-(R^e)_m-NR^aR^b or $-(R^d)_m$ -NR^a-(R^e)-OR^b; and wherein any of the said alkyl, alkenyl, alkynyl, R^a , R^b , R^c , R^d , R^e , C₃-C₁₂ cycloalkyl, phenyl, or 3-15 membered heterocyclyl, are independently optionally further substituted by 1-3 groups selected from -F, C₁-C₃ alkyl, C₁-C₃ perfluoroalkyl, hydroxyl, C₁-C₆alkoxyl or oxo;

each R^a , R^b and R^c is independently selected from H, C₁-C₈ alkyl, C₂-C₈ alkenyl, $-(R^d)_m$ -(C₃-C₈ cycloalkyl), $-(R^d)_m$ -(C₃-C₈ cycloalkenyl), C₂-C₈ alkynyl, $-(R^d)_m$ -phenyl, or $-(R^d)_m$ -(3-7 membered heterocyclyl), and each R^a , R^b and R^c is independently

- 5 -

optionally further substituted by 1-3 groups selected from halide, hydroxyl, -CN, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxy and C₁-C₆ alkylamino; or, when connected to the same nitrogen, R^a and R^b may together optionally form a 3-7 membered heterocyclyl, which may optionally be further substituted by 0-3 groups selected from halide, hydroxyl, -CN, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxy or C₁-C₆ alkylamino;

each R^d and R^e is independently-(C₁-C₃ alkylene)-, -(C₂-C₅ alkenylene)-, or-(C₂-C₅ alkynylene)-;

and each m is independently 0 or 1;

with the proviso that when X is N, R⁶ and R⁷ are not both H, and that when X is C-R¹¹, R⁶ and R⁷ are both H.

In one embodiment of the invention, R⁹ and R¹⁰ are both methyl.

In another embodiment of the invention, X is N and R⁶ and R⁷ are each independently H or C₁-C₆alkyl but are not both H.

In one embodiment of the invention, A is N and B is C. In an alternative embodiment, A is C and B is N.

In another embodiment of the invention, R⁶ and R⁷ are both methyl. In an alternative embodiment, R⁶ is H and R⁷ is methyl.

In one embodiment of the invention, R⁴ is R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)₂R^a, -(R^d)_m-OS(O)₂NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO₂, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)₂R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)₂R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)₂NR^aR^b, -(R^d)_m-O--(R^e)_m-NR^aR^b or -(R^d)_m-NR^a--(R^e)_m-OR^b; wherein the said R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ cycloalkyl, aryl, 3-15 membered heterocyclyl, are independently optionally further substituted by 0-3 R¹² groups.

In a further embodiment of the invention, R⁴ is methyl.

In another embodiment of the invention, R¹ is R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-


- 6 -

$S(O)R^a, -(R^d)_m-OS(O)_2R^a, -(R^d)_m-OS(O)_2NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO_2, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)_2R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)_2R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)_2NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b$ or $-(R^d)_m-NR^a-(R^e)-OR^b$; wherein the said- $R^a, R^b, R^c, R^d, R^e, C_3-C_{12}$ cycloalkyl, aryl, the said 3-15 membered heterocyclyl, are independently optionally further substituted by 0-3 R^{12} groups.

In a further embodiment of the invention, R^1 is $-(R^d)_m-OR^a, C_1-C_8$ alkyl, or $-(R^d)_m-NR^aR^b$. In another further embodiment of the invention, R^8 is R^a-O-R^b, C_1-C_8 alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, $-(R^d)_m-(C_3-C_{12}$ cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m-(C_1-C_6$ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m-CN, -(R^d)_m-OR^a$, or $-(R^d)_m-NR^aR^b$.

In another embodiment of the invention, each R^d and R^e is independently an $-(C_1-C_3$ alkylene).

The present invention is further directed to a compound or pharmaceutically acceptable salt of Formula B,



(B)

wherein

X is $C-R^{11}$ or N , wherein R^{11} is $H, halo, OH, C_1-C_3$ alkyl, CF_3 , or CN ;

A and B are independently C or N ;

R^1 is R^a-O-R^b, C_1-C_8 alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, $-(R^d)_m-(C_3-C_{12}$ cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m-(C_1-C_6$ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-$

- 7 -

OS(O)₂R^a, -(R^d)_m-OS(O)₂NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO₂, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)₂R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)₂R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)₂NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b or -(R^d)_m-NR^a-(R^e)-OR^b; and wherein any of the said alkyl, alkenyl, alkynyl, R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ cycloalkyl, phenyl or 3-15 membered heterocyclyl, may independently be further optionally substituted by 0-3 R¹² groups;

R² and R³ are each independently selected from H, R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)₂R^a, -(R^d)_m-OS(O)₂NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO₂, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)₂R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)₂R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)₂NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b or -(R^d)_m-NR^a-(R^e)-OR^b; wherein R² and R³ may together optionally cyclize to form a saturated or unsaturated 3-7 membered heterocyclyl fused to the 6-membered N-containing heteroaryl to which they are attached; and wherein any of the said alkyl, alkenyl, alkynyl, R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ cycloalkyl, phenyl or 3-15 membered heterocyclyl, may independently be further optionally substituted by 0-3 R¹² groups;

R⁴ and R⁵ are each independently selected from H, R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)₂R^a, -(R^d)_m-OS(O)₂NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO₂, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)₂R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)₂R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)₂NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b or -(R^d)_m-NR^a-(R^e)-OR^b; wherein any of the said alkyl, alkenyl, alkynyl, R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ cycloalkyl, aryl or 3-15 membered heterocyclyl are independently optionally further substituted by 0-3 R¹² groups,

R⁶ is H, R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆

perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m$ -C(O)R^a, $-(R^d)_m$ -C(O)OR^a, $-(R^d)_m$ -C(O)NR^aR^b, $-(R^d)_m$ -OR^a, $-(R^d)_m$ -OC(O)R^a, $-(R^d)_m$ -OC(O)NR^aR^b, $-(R^d)_m$ -O-S(O)R^a, $-(R^d)_m$ -OS(O)₂R^a, $-(R^d)_m$ -OS(O)₂NR^aR^b, $-(R^d)_m$ -OS(O)NR^aR^b, $-(R^d)_m$ -NO₂, $-(R^d)_m$ -NR^aR^b, $-(R^d)_m$ -N(R^a)C(O)R^b, $-(R^d)_m$ -N(R^a)C(O)OR^b, $-(R^d)_m$ -N(R^c)C(O)NR^aR^b, $-(R^d)_m$ -N(R^a)S(O)₂R^b, $-(R^d)_m$ -N(R^a)S(O)R^b, $-(R^d)_m$ -SR^a, $-(R^d)_m$ -S(O)R^a, $-(R^d)_m$ -S(O)₂R^a, $-(R^d)_m$ -S(O)NR^aR^b, $-(R^d)_m$ -S(O)₂NR^aR^b, $-(R^d)_m$ -O-(R^e)_m-NR^aR^b or $-(R^d)_m$ -NR^a-(R^e)-OR^b; and wherein any of the said alkyl, alkenyl, alkynyl, R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ cycloalkyl, phenyl, or 3-15 membered heterocyclyl are independently optionally further substituted by 1-3 groups selected from -F, C₁-C₃ alkyl, C₁-C₃ perfluoroalkyl, hydroxyl, C₁-C₆alkoxyl, or oxo;

R⁹ and R¹⁰ are each independently C₁-C₂ alkyl or can together cyclize to form a cyclopropyl or cyclobutyl;

each R¹² is independently H, R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, $-(R^d)_m$ -(C₃-C₁₂ cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m$ -(C₁-C₆ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m$ -C(O)R^a, $-(R^d)_m$ -C(O)OR^a, $-(R^d)_m$ -C(O)NR^aR^b, $-(R^d)_m$ -OR^a, $-(R^d)_m$ -OC(O)R^a, $-(R^d)_m$ -OC(O)NR^aR^b, $-(R^d)_m$ -O-S(O)R^a, $-(R^d)_m$ -OS(O)₂R^a, $-(R^d)_m$ -OS(O)₂NR^aR^b, $-(R^d)_m$ -OS(O)NR^aR^b, $-(R^d)_m$ -NO₂, $-(R^d)_m$ -NR^aR^b, $-(R^d)_m$ -N(R^a)C(O)R^b, $-(R^d)_m$ -N(R^a)C(O)OR^b, $-(R^d)_m$ -N(R^c)C(O)NR^aR^b, $-(R^d)_m$ -N(R^a)S(O)₂R^b, $-(R^d)_m$ -N(R^a)S(O)R^b, $-(R^d)_m$ -SR^a, $-(R^d)_m$ -S(O)R^a, $-(R^d)_m$ -S(O)₂R^a, $-(R^d)_m$ -S(O)NR^aR^b, $-(R^d)_m$ -S(O)₂NR^aR^b, $-(R^d)_m$ -O--(R^e)_m-NR^aR^b or $-(R^d)_m$ -NR^a-(R^e)-OR^b; and wherein any of the said alkyl, alkenyl, alkynyl, R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ cycloalkyl, phenyl, or 3-15 membered heterocyclyl, are independently optionally further substituted by 1-3 groups selected from -F, C₁-C₃ alkyl, C₁-C₃ perfluoroalkyl, hydroxyl, C₁-C₆alkoxyl or oxo;

each R^a, R^b and R^c is independently selected from H, C₁-C₈ alkyl, C₂-C₈ alkenyl, $-(R^d)_m$ -(C₃-C₈ cycloalkyl), $-(R^d)_m$ -(C₃-C₈ cycloalkenyl), C₂-C₈ alkynyl, $-(R^d)_m$ -phenyl, or $-(R^d)_m$ -(3-7 membered heterocyclyl), and each R^a, R^b and R^c is independently optionally further substituted by 1-3 groups selected from halide, hydroxyl, -CN, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxyl and C₁-C₆ alkylamino; or, when connected to the same nitrogen, R^a and R^b may together optionally form a 3-7 membered heterocyclyl, which may optionally be further substituted by 0-3 groups selected from halide, hydroxyl, -CN, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxyl or C₁-C₆ alkylamino;

each R^d and R^e is independently-(C₁-C₃ alkylene)-, -(C₂-C₅ alkenylene)-, or-(C₂-C₅ alkynylene)-;

and each m is independently 0 or 1.

In one embodiment of the invention of Formula (B), A is N and B is C.

In a further embodiment of Formula (B), wherein R⁹ and R¹⁰ are both methyl.

In another embodiment of Formula (B), R⁴ is-(R^d)_m-OR^a, C₁-C₈ alkyl, C₂-C₈ alkenyl or C₂-C₈ alkynyl. In a further embodiment, R⁴ is methyl.

In another embodiment of Formula (B), R¹ is -(R^d)_m-OR^a, C₁-C₈ alkyl, or-(R^d)_m-NR^aR^b.

In another embodiment of Formula (B), each R^d and R^e is independently an-(C₁-C₃ alkylene)-.

The present invention is directed to compounds or pharmaceutically acceptable salts selected from the group consisting of:

N⁴-(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²-ethyl-5-fluoropyrimidine-2,4-diamine ,
 N⁴-(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro-N²,N²-dimethylpyrimidine-2,4-diamine,
 N²-cyclopropyl-N⁴-(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoropyrimidine-2,4-diamine
 N⁴-(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro-N²-methylpyrimidine-2,4-diamine,
 N⁴-(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro-N²-isopropylpyrimidine-2,4-diamine ,
 N⁴-(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²-ethylpyrimidine-2,4-diamine,
 N⁴-(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²,N²-dimethylpyrimidine-2,4-diamine,
 5-([(8S)-6,8-dimethyl-6,9-diazaspiro[4.5]dec-9-yl]carbonyl)-N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
 N⁴-(5-([(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl]carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²-ethyl-5-fluoropyrimidine-2,4-diamine,

*N*⁴-(5-{[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-yl)piperazin-1-yl]carbonyl}-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-yl)-*N*²-ethyl-5-fluoropyrimidine-2,4-diamine,

*N*²-ethyl-5-fluoro-*N*⁴-(5-{[(2*S*,5*R*)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl}-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-yl)pyrimidine-2,4-diamine acetate salt,

*N*⁴-(6,6-dimethyl-5-{[(2*S*,5*R*)-2,4,5-trimethylpiperazin-1-yl]carbonyl}-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-yl)-*N*²-ethyl-5-fluoropyrimidine-2,4-diamine,

4-[(6,6-dimethyl-5-{[(2*S*,5*R*)-2,4,5-trimethylpiperazin-1-yl]carbonyl}-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-yl)amino]pyrimidine-2-carbonitrile,

N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-{[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl}-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(2-ethyl-5-fluoropyrimidin-4-yl)-5-{[(2*S*,5*R*)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl}-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

2-[(5*S*)-4-{[3-[(2-ethyl-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl}-1,5-dimethylpiperazin-2-yl)ethanol,

5-{[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazin-1-yl]carbonyl}-*N*-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-5-{[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl}-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(5-fluoro-2-propylpyrimidin-4-yl)-6,6-dimethyl-5-{[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl}-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(5-fluoro-2-isopropylpyrimidin-4-yl)-6,6-dimethyl-5-{[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl}-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-5-{[(2*S*,5*R*)-2,4,5-trimethylpiperazin-1-yl]carbonyl}-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

5-{[(3*S*,8*aS*)-3,8*a*-dimethylhexahydropyrrolo[1,2-*a*]pyrazin-2(1*H*)-yl]carbonyl}-*N*-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-5-([(3*S*,8*aS*)-3-methylhexahydropyrrolo[1,2-*a*]pyrazin-2(1*H*)-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-5-([(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]-6,6-dimethyl-5-([(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

5-([(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazin-1-yl]carbonyl)-*N*-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

5-([(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-yl)piperazin-1-yl]carbonyl)-*N*-(4-methoxypyrimidin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

5-([(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-yl)piperazin-1-yl]carbonyl)-6,6-dimethyl-*N*-(4-methylpyrimidin-2-yl)-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

5-([(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-yl)piperazin-1-yl]carbonyl)-6,6-dimethyl-*N*-[4-(trifluoromethyl)pyrimidin-2-yl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

5-([(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazin-1-yl]carbonyl)-6,6-dimethyl-*N*-(4-methylpyrimidin-2-yl)-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-([(2*S*,5*R*)-2,4,5-trimethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[4-ethyl(2*S*,5*R*)-2,5-dimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-([(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-([(2*S*,5*R*)-4-(2-methoxyethyl)-2,5-dimethylpiperazin-1-yl]carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-([(2*S*,5*R*)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-5-[[[(2*S*,5*R*)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-6,6-dimethyl-5-[[[(2*S*,5*R*)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

N-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-6,6-dimethyl-5-[[[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

5-[[[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazin-1-yl]carbonyl]-*N*-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

5-[[[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-yl)piperazin-1-yl]carbonyl]-*N*-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine ,

2-((5*S*)-4-[[3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl]-1,5-dimethylpiperazin-2-yl)ethanol ,

2-((5*S*)-4-[[3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl]-1,5-dimethylpiperazin-2-yl)ethanol ,

5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-*N*-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

5-[[[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazin-1-yl]carbonyl]-*N*-[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

2-((5*S*)-4-[[3-[[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl]-1,5-dimethylpiperazin-2-yl)ethanol.

The present invention is further directed to a pharmaceutical composition comprising an effective amount of a compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention is further directed to a method of treating diabetes mellitus and its complications, cancer, ischemia, inflammation, central nervous system disorders, cardiovascular disease, Alzheimer's disease and dermatological disease, virus diseases, inflammatory disorders, or diseases in which the liver is a target organ, the

method comprising administering to a mammal an effective amount of a compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof.

The present invention is further directed to a method of treating diabetic mellitus and its complications, the method comprising administering to a mammal an effective amount of a compound according to any of the preceding claims. In a further embodiment of this invention, the complications comprise diabetic retinopathy (including macular edema), nephropathy and neuropathy.

DEFINITIONS

As used herein, the terms "comprising" and "including" are used in their open, non-limiting sense. As used herein, the terms "C₁-C₈" or "C₂-C₈" and so forth, refer to moieties having 1 to 8 or 2 to 8 carbon atoms, respectively.

The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight or branched moieties. Exemplary alkyl moieties have carbon atoms in the range of 1 to 8 carbon atoms, 1 to 6 carbon atoms or 1 to 4 carbon atoms.

The term "alkenyl", as used herein, unless otherwise indicated, includes alkyl moieties having at least one carbon-carbon double bond wherein alkyl is as defined above and including E and Z isomers of said alkenyl moiety.

The term "alkynyl", as used herein, unless otherwise indicated, includes alkyl moieties having at least one carbon-carbon triple bond wherein alkyl is as defined above.

The term "alkoxyl", as used herein, unless otherwise indicated, includes O-alkyl groups wherein alkyl is as defined above.

The term "hydroxyl", as used herein, unless otherwise indicated, includes -OH.

The term "amino", as used herein, unless otherwise indicated, is intended to include the -NH₂ radical, and any substitutions of the N atom.

The terms "halogen" and "halo", as used herein, unless otherwise indicated, represent chlorine, fluorine, bromine or iodine.

The term "trifluoromethyl", as used herein, unless otherwise indicated, is meant to represent a -CF₃ group.


The term "perfluoroalkyl", as used herein, is meant to represent an alkyl group in which all hydrogens attached to the carbons have been replaced by fluorine, such as CF_3 , $\text{CF}_2\text{-CF}_3$, $\text{C}(\text{CF}_2)(\text{CF}_2)$ and so on.

The term "trifluoromethoxy", as used herein, unless otherwise indicated, is meant to represent a $-\text{OCF}_3$ group.

The term "cyano", as used herein, unless otherwise indicated, is meant to represent a $-\text{CN}$ group.

The term " CH_2Cl_2 ", as used herein, unless otherwise indicated, is meant to represent dichloromethane.

The term " $\text{C}_3\text{-C}_{12}$ cycloalkyl" or " $\text{C}_5\text{-C}_8$ cycloalkyl", as used herein, unless otherwise indicated, refers to a non-aromatic, saturated or partially saturated, monocyclic or fused, spiro or unfused bicyclic or tricyclic hydrocarbon referred to herein containing a total of from 3 to 12 carbon atoms, or 5-8 ring carbon atoms, respectively. Exemplary cycloalkyls include rings having from 3-10 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and adamantyl. Illustrative examples of cycloalkyl are derived from, but not limited to, the following:




The term "aryl", as used herein, unless otherwise indicated, includes an organic radical derived from an aromatic hydrocarbon by removal of one hydrogen, such as phenyl or naphthyl.

The term "(3-15)-membered heterocycl", "(3-7)-membered heterocycl", "(6-10)-membered heterocycl", or "(4 to 10)-membered heterocycl", as used herein, unless otherwise indicated, includes aromatic and non-aromatic heterocyclic groups containing one to four heteroatoms each selected from O, S and N, wherein each heterocyclic group has from 3-15, 3-7, 6-10, or 4 to 10 atoms, respectively, in its ring system, and with the proviso that the ring of said group does not contain two adjacent O or S atoms. Non-aromatic heterocyclic groups include groups having only 3 atoms in their ring

system, but aromatic heterocyclic groups must have at least 5 atoms in their ring system. The heterocyclic groups include benzo-fused ring systems. An example of a 3 membered heterocyclic group is aziridine, an example of a 4 membered heterocyclic group is azetidiny (derived from azetidine). An example of a 5 membered heterocyclic group is thiazolyl, an example of a 7 membered ring is azepiny, and an example of a 10 membered heterocyclic group is quinolinyl. Examples of non-aromatic heterocyclic groups are pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidino, morpholino, thiomorpholino, thioxanyl, piperaziny, azetidiny, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepiny, diazepiny, thiazepiny, 1,2,3,6-tetrahydropyridiny, 2-pyrroliny, 3-pyrroliny, indoliny, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazoliny, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidiny, imidazoliny, imidazolidiny, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl, 3H-indolyl and quinoliziny. Heterocycles include monocyclic and polycyclic aromatic ring structures, with "(5-12)-membered heteroaryls" referring to those that are heterocycles having 5 to 12 atoms in their ring system(s). Examples of "(5-12)-membered heteroaryls" are pyridiny, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyraziny, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrroly, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnoliny, indazolyl, indoliziny, phthalaziny, pyridaziny, triaziny, isoindolyl, pteridinyl, puriny, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridiny, and furopyridiny. The foregoing groups, as derived from the groups listed above, may be C-attached or N-attached where such is possible. For instance, a group derived from pyrrole may be pyrrol-1-yl (N-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole may be imidazol-1-yl (N-attached) or imidazol-3-yl (C-attached). The above-mentioned heterocyclic groups may be optionally substituted on any ring carbon, sulfur, or nitrogen atom(s) by one to two oxo, per ring. An example of a heterocyclic group wherein 2 ring carbon atoms are substituted with oxo moieties is 1,1-dioxo-thiomorpholinyl. Other illustrative examples of 4 to 10 membered heterocyclic are derived from, but not limited to, the following:

- 16 -



The term "(12-15)-membered heterocyclyl", as used herein, unless otherwise indicated, includes aromatic and non-aromatic heterocyclic groups that are in a partially fused or spirocyclic configuration and which contain at least one N and optionally additional 1 to 5 heteroatoms each selected from O, S and N, wherein the heterocyclic group has from 12 to 15 atoms, respectively, in its system, and with the proviso that any ring of said group does not contain two adjacent O or S atoms. The heterocyclic groups include tricyclic fused ring and spirocyclic systems. An example of a 13-membered tricyclic heterocyclic group is 3,4-dihydropyrazino[1,2-a]benzimidazole and an example of a 15-membered spirocyclic heterocyclic group is 3,4-dihydro-1'H-spirochromene.

Unless otherwise indicated, the term "oxo" refers to =O.

A "solvate" is intended to mean a pharmaceutically acceptable solvate form of a specified compound that retains the biological effectiveness of such compound. Examples of solvates include compounds of the invention in combination with water, isopropanol, ethanol, methanol, DMSO (dimethylsulfoxide), ethyl acetate, acetic acid, or ethanolamine.

The phrase "pharmaceutically acceptable salt(s)", as used herein, unless otherwise indicated, includes salts of acidic or basic groups which may be present in the


compounds of formula (A) or formula (B). The compounds of formula (A) or formula (B) that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds of formula (A) or formula (B) are those that form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edislyate, estolate, esylate, ethylsuccinate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodode, and valerate salts.

The term "treating", as used herein, unless otherwise indicated, means reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "treatment", as used herein, unless otherwise indicated, refers to the act of treating as "treating" is defined immediately above.


The phrase "therapeutically effective amount", as used herein, refers to that amount of drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, system, animal, or human that is being sought by a researcher, veterinarian, medical doctor or other.

The term "substituted" means that the specified group or moiety bears one or more substituents. The term "unsubstituted" means that the specified group bears no substituents. The term "optionally substituted" means that the specified group is unsubstituted or substituted by one or more substituents.

In accordance with convention, in some structural formula herein, the carbon atoms and their bound hydrogen atoms are not explicitly depicted e.g.,  represents

a methyl group,  represents an ethyl group,  represents a cyclopentyl group, etc. Moreover, the depiction of any cyclic group (aryl, heterocyclic or cycloalkyl)

- 18 -



with a bond that is not directly attached to a ring atom, e.g., indicates that the point of attachment may be on any available ring atom of the cyclic group.

Certain compounds of formula (A) or formula (B) may have asymmetric centers and therefore exist in different enantiomeric forms. All optical isomers and stereoisomers of the compounds of formula (A) or formula (B), and mixtures thereof, are considered to be within the scope of the invention. With respect to the compounds of formula (A) or formula (B), the invention includes the use of a racemate, one or more enantiomeric forms, one or more diastereomeric forms, or mixtures thereof. The compounds of formula (A) or formula (B) may also exist as tautomers. This invention relates to the use of all such tautomers and mixtures thereof.

Certain functional groups contained within the compounds of the present invention can be substituted for bioisosteric groups, that is, groups which have similar spatial or electronic requirements to the parent group, but exhibit differing or improved physicochemical or other properties. Suitable examples are well known to those of skill in the art, and include, but are not limited to moieties described in Patini et al., Chem. Rev, 1996, 96, 3147-3176 and references cited therein.

The subject invention also includes isotopically-labelled compounds, which are identical to those recited in formula (A) or formula (B), but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Compounds of the present invention and pharmaceutically acceptable salts or solvates of said compounds which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labelled compounds of the present invention, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e., ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage

requirements and, hence, may be preferred in some circumstances. Isotopically labeled compounds of formula (A) or formula (B) of this invention thereof can generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples below, by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

The term "mmol", as used herein, unless otherwise indicated, is intended to mean millimole. The term "equiv", as used herein, unless otherwise indicated, is intended to mean equivalent. The term "mL", as used herein, unless otherwise indicated, is intended to mean milliliter. The term "U", as used herein, unless otherwise indicated, is intended to mean units. The term "mm" as used herein, unless otherwise indicated, is intended to mean millimeter. The term "g", as used herein, unless otherwise indicated, is intended to mean gram. The term "kg", as used herein, unless otherwise indicated, is intended to mean kilogram. The term "h", as used herein, unless otherwise indicated, is intended to mean hour. The term "min", as used herein, unless otherwise indicated, is intended to mean minute. The term " μ L", as used herein, unless otherwise indicated, is intended to mean microliter. The term " μ M", as used herein, unless otherwise indicated, is intended to mean micromolar. The term " μ m", as used herein, unless otherwise indicated, is intended to mean micrometer. The term "M", as used herein, unless otherwise indicated, is intended to mean molar. The term "N", as used herein, unless otherwise indicated, is intended to mean normal. The term "nm", as used herein, unless otherwise indicated, is intended to mean nanometer. The term "nM", as used herein, unless otherwise indicated, is intended to mean nanoMolar. The term "amu", as used herein, unless otherwise indicated, is intended to mean atomic mass unit. The term " $^{\circ}$ C", as used herein, unless otherwise indicated, is intended to mean Celsius. The term "m/z", as used herein, unless otherwise indicated, is intended to mean, mass/charge ratio. The term "wt/wt", as used herein, unless otherwise indicated, is intended to mean weight/weight. The term "v/v", as used herein, unless otherwise indicated, is intended to mean volume/volume. The term "mL/min", as used herein, unless otherwise indicated, is intended to mean milliliter/minute. The term "UV", as used herein, unless otherwise indicated, is intended to mean ultraviolet. The term "APCI-MS", as used herein, unless otherwise indicated, is intended to mean atmospheric pressure chemical ionization mass spectroscopy. The term "HPLC", as used herein, unless otherwise indicated, is intended to mean high performance liquid chromatograph. The chromatography was

performed at a temperature of about 20 °C, unless otherwise indicated. The term "LC", as used herein, unless otherwise indicated, is intended to mean liquid chromatograph. The term "LCMS", as used herein, unless otherwise indicated, is intended to mean liquid chromatography mass spectroscopy. The term "TLC", as used herein, unless otherwise indicated, is intended to mean thin layer chromatography. The term "SFC", as used herein, unless otherwise indicated, is intended to mean supercritical fluid chromatography. The term "sat" as used herein, unless otherwise indicated, is intended to mean saturated. The term "aq" as used herein, is intended to mean aqueous. The term "ELSD" as used herein, unless otherwise indicated, is intended to mean evaporative light scattering detection. The term "MS", as used herein, unless otherwise indicated, is intended to mean mass spectroscopy. The term "HRMS (ESI)", as used herein, unless otherwise indicated, is intended to mean high-resolution mass spectrometry (electrospray ionization). The term "Anal.", as used herein, unless otherwise indicated, is intended to mean analytical. The term "Calcd", as used herein, unless otherwise indicated, is intended to mean calculated. The term "N/A", as used herein, unless otherwise indicated, is intended to mean not tested. The term "RT", as used herein, unless otherwise indicated, is intended to mean room temperature. The term "Mth.", as used herein, unless otherwise indicated, is intended to mean Method. The term "Celite®", as used herein, unless otherwise indicated, is intended to mean a white solid diatomite filter agent commercially available from World Minerals located in Los Angeles, California USA. The term "Eg.", as used herein, unless otherwise indicated, is intended to mean example.

Terms such as $-(CR^3R^4)_t$ or $-(CR^{10}R^{11})_v$, for example, are used, R^3 , R^4 , R^{10} and R^{11} may vary with each iteration of t or v above 1. For instance, where t or v is 2 the terms $-(CR^3R^4)_v$ or $-(CR^{10}R^{11})_t$ may equal $-CH_2CH_2-$, or $-CH(CH_3)C(CH_2CH_3)(CH_2CH_2CH_3)-$, or any number of similar moieties falling within the scope of the definitions of R^3 , R^4 , R^{10} and R^{11} .

The term " K_i ", as used herein, unless otherwise indicated, is intended to mean values of enzyme inhibition constant. The term " K_i app", as used herein, unless otherwise indicated, is intended to mean K_i apparent. The term " IC_{50} ", as used herein, unless otherwise indicated, is intended to mean concentrations required for at least 50% enzyme inhibition.

Other aspects, advantages, and features of the invention will become apparent from the detailed description below.


DETAILED DESCRIPTION AND EMBODIMENTS OF THE INVENTION

The following reaction Schemes illustrate the preparation of the compounds of the present invention. Unless otherwise indicated, R¹ through R¹² and R^a through R^e in the reaction schemes and the discussion that follows are as defined above.

Detailed Description

Compounds of formulas I-III can be made following the synthetic routes in Scheme 1 and Scheme 2. In Scheme 1 and Scheme 2 and the descriptions following, "BOC", "Boc" or "boc" means N-*tert*-butoxycarbonyl, DCM means CH₂Cl₂, DIPEA (also known as Hunig's base) means diisopropyl ethyl amine, DMA means dimethyl amine, "DMF" means dimethyl formamide, "DMSO" means dimethylsulfoxide, Et means -CH₂CH₃, "MTBE" means methyl *t*-butyl ether, NMP means 1-methyl-2-pyrrolidinone, TEA means triethyl amine, TFA means trifluoro acetic acid, THF means tetrahydrofuran. While schemes 1 and 2 and the description refer to compound I, schemes 1 and 2 and the description are equally applicable to compounds II and III.

Scheme 1




Scheme 1 illustrates the synthesis of the intermediate I(A) used to make compounds of formula I. The amino group of the substituted amino acid I(A) is alkylated to give compound I(B). This can typically be done by treating compound I(A) with an alkylating agent in the presence of a base. An activated electrophilic double

bond moiety is a commonly used alkylating reagent. A typical reaction condition of alkylating I(A) with an activated electrophilic double bond moiety is to treat I(A) with the activated double bond moiety in the presence of a strong base. Subsequent aqueous work up affords compound I(B). The amino group of compound I(B) is then protected with a boc group to give compound I(C). This can typically be done by treating compound I(B) with Boc agent in the presence of a base. A typical condition is to treat compound I(B) with (Boc)₂O in the presence of Me₄NOH in MeCN as a solvent. The carboxylic acid group of compound I(C) is then converted into a methyl ester of compound I(D). A typical condition of converting the carboxylic acid group into the methyl ester group is to treat I(C) with methyl iodide in DMF in the presence of a base. Compound I(D) then undergoes an intramolecular aldol condensation to give compound I(E). This can typically be done by treating compound I(D) with a strong base in an aprotic solvent. A typical condition is to treat compound I(D) with t-BuOK in toluene. Subsequent aqueous workup gives compound I(E). Compound I(E) then undergoes a 2+3 cyclization with a hydrazine moiety to form compound I(F). A typical condition of the cyclization is to reflux compound I(E) with hydrazine and acetic acid in EtOH. The free base pyrazole nitrogen of compound I(F) is then acylated to give compound I(G). A typical condition of the acylation is to treat compound I(F) with chloro ethyl carbonate in THF.

More detailed synthetic conditions to compound I(G) of Scheme 1 can be found in U.S. Patent Application Publication No. 2003/0171357 and PCT Publication WO 02/12242.

- 23 -

Scheme 2



Scheme 2 illustrates two routes through which compounds of formula **I** can be made from intermediate **I(G)**. In the first route of Scheme 2, compound **I(G)** undergoes a nucleophilic reaction with an R^a electrophile moiety. This nucleophilic reaction can be an acylation, alkylation, sulfonylation, reductive amination or one of many other reactions that an amine functionality carries out. A typical acylation reaction condition is to treat compound **I(G)** with an acylating agent such as R^a -COCl, in the presence of a base such as 2 equivalents of DIPEA, in a solvent such as dichloromethane. The reaction mixture is stirred at between 0°C and room temperature for 12 hours. Subsequent aqueous workup gives compound **II(A)**. The Boc group on the pyrrole nitrogen of compound **II(A)** is then removed to give compound **II(B)**. This can typically

- 24 -

be done by treating **II(A)** with a strong acid. A typical reaction condition is to treat compound **II(A)** with 4N HCl in dioxane and DCM. Subsequent aqueous workup affords compound **II(B)**. The pyrrole NH of compound **II(B)** is then converted to the chloroformate **II(C)**. This can typically be done using phosgene, triphosgene, or some equivalent. A typical reaction condition is to treat **II(B)** with 2 equivalents of triphosgene in DCM at 0°C for four hours. Subsequent mild basic workup with saturated NaHCO₃ and purification gives compound **II(C)**. Compound **II(C)** is then treated with an R^a nucleophile moiety. The nucleophile can be an alcohol, amine or one of any other functionalities that can react with the electrophile **II(C)**. A typical reaction involves treating **II(C)** with a nucleophile such as 1.5 equivalents of an alcohol in the presence of 2 equivalents of base such as K₂CO₃ in a solvent such as DME. The reaction is heated to 80°C for eight hours and the solvent removed. Alternatively, **II(C)** can be treated with 1.5 equivalents of an amine in the presence of 1 equivalent of base such as DIPEA in a solvent such as THF. Subsequent work up in a protic solvent such as methanol in the presence of base such as TEA followed by purification give compound of formula **I**.

Alternatively, compound **II(B)** can then undergoes a nucleophilic reaction with an (COR^b) electrophile to give compound **II(D)**. The nucleophilic reaction carried out for this transformation can be an alkylation, acylation, sulfonylation, reductive amination. An acylation reaction of **II(B)** to give **II(D)** is carried out by treating compound **II(B)** with an acylating reagent in the presence of base. A typical reaction condition is to mix compound **II(B)** with excess of base, such as DIPEA in DCM and adding the resulting solution to an isocyanate at 0°C. The reaction is stirred for 2 hours and subsequent aqueous workup gives compound **II(D)**. The ethyl ester protecting group on the pyrazole nitrogen of compound **II(D)** is removed to give compound of formula **I**. This can typically be done by treating compound **II(D)** with a base. A typical reaction condition is to reflux compound **II(D)** in dioxane and DCM in the presence of 2-3 equivalents of LiOH. Subsequent aqueous workup affords compound of formula **I**.

In the second route of Scheme 2, the Boc group on the pyrrole nitrogen is removed to give compound **III(A)**. The reaction can typically be carried out by treating compound **I(G)** with a strong acid. A typical reaction condition is to treat compound **I(G)** with 4N HCl in dioxane and DCM. Subsequent aqueous workup affords compound **III(A)**. Compound **III(A)** can then undergo a nucleophilic reaction with an (COR^b) electrophile to give compound **III(B)**. Because the -NH₂ group attached to the pyrazole

in compound **III(A)** is less reactive than the pyrrole nitrogen of **III(A)**, the transformation of **III(A)** to **III(B)** can be carried out without protecting the pyrazole -NH_2 group of compound **III(A)**. The nucleophilic reaction carried out for this transformation can be an alkylation, acylation, sulfonylation, reductive amination. Relative mild reaction conditions are preferred to achieve the reaction selectivity. An acylation reaction of **III(A)** to give **III(B)** is carried out by treating compound **III(A)** with an acylating reagent in the presence of base. A typical reaction condition is to mix compound **III(A)** with excess of base, such as DIPEA in DCM and adding the resulting solution to an isocyanate at 0°C . The reaction mixture is held at 0°C for about two hours and subsequent aqueous workup gives compound **III(B)**.

Compound **III(B)** then undergoes a nucleophilic reaction with an R^a electrophile moiety. This nucleophilic reaction can be an acylation, alkylation, sulfonylation, reductive amination or one of many other reactions that an amine functionality carries out. A typical acylation reaction condition is to treat compound **III(B)** with an acylating agent such as $\text{R}^a\text{-NCO}$ in the presence of a base such as 2 equivalents of DIPEA in a solvent such as dichloromethane for 2 hours. Alternatively, **III(B)** can be treated with an acylating agent such as $\text{R}^a\text{-COOR}$, where R is an activating group such as p-nitrophenyl, in the presence of a base such as 2 equivalents of DIPEA, in a solvent such as 1,2-dichloroethane. Subsequent aqueous workup gives compound **III(C)**. The ethyl ester protecting group on the pyrazole nitrogen of compound **III(C)** is removed typically with a base to give the free base compound **I**. A typical reaction condition is to mix compound **III(C)** with TEA in a protic solvent such as methanol followed by purification to give compound of formula **I**.

Alternatively, the ethyl ester protecting group on the pyrazole nitrogen of compound **III(B)** is removed to give the free base compound **III(D)**. This can typically be done by treating compound **III(B)** with a base. A typical reaction condition is to reflux compound **III(B)** in dioxane and DCM in the presence of 2-3 equivalents of LiOH. Subsequent aqueous workup affords compound **III(D)**. Compound **III(D)** then undergoes a nucleophilic reaction with an R^a electrophile moiety. This nucleophilic reaction can be an acylation, alkylation, sulfonylation, reductive amination or one of many other reactions that an amine functionality carries out. A typical acylation reaction condition is to treating compound **III(D)** with an acylating agent such as $\text{R}^a\text{-COCl}$, in the presence of a base such as 2 equivalents of DIPEA in a solvent such as

dichloromethane. The reaction mixture is stirred for four hours and subsequent aqueous workup and purification gives compound of formula I.

Any of the above compounds of formula (A) and (B) as defined above can be converted into another analogous compound by standard chemical manipulations. All starting materials, reagents, and solvents are commercially available and are known to those of skill in the art unless otherwise stated. These chemical manipulations are known to those skilled in the art and include (a) removal of a protecting group by methods outlined in T. W. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, 2nd Ed., John Wiley and Sons, New York, 1991; (b) displacement of a leaving group (halide, mesylate, tosylate, etc) with a primary or secondary amine, thiol or alcohol to form a secondary or tertiary amine, thioether or ether, respectively; (c) treatment of primary and secondary amines with an isocyanate, acid chloride (or other activated carboxylic acid derivative), alkyl/aryl chloroformate or sulfonyl chloride to provide the corresponding urea, amide, carbamate or sulfonamide; (d) reductive amination of a primary or secondary amine using an aldehyde.

The compounds of the present invention may have asymmetric carbon atoms. Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods known to those skilled in the art, for example, by chromatography or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixtures into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., alcohol), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. All such isomers, including diastereomeric mixtures and pure enantiomers are considered as part of the invention.

The compounds of formula (A) or formula (B) that are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate the compound of formula (A) or formula (B) from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the

chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained. The desired acid salt can also be precipitated from a solution of the free base in an organic solvent by adding to the solution an appropriate mineral or organic acid.

Those compounds of formula (A) or formula (B) that are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particularly, the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the acidic compounds of formula (A) or formula (B). Such non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium, calcium, and magnesium, etc. These salts can easily be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure completeness of reaction and maximum yields of the desired final product.

The compounds of the present invention are inhibitors of protein kinase C and preferably selectively inhibit beta-1, beta-2 and optionally alpha isozymes of protein kinase C. With respect to the beta-2 isozyme in particular, the compounds of the present invention have K_i values of less than 100 nM.

As an inhibitor of protein kinase C the compounds are useful in the treatment of conditions in which protein kinase C has demonstrated a role in the pathology. Conditions recognized in the art include: diabetes mellitus and its complications, cancer, ischemia, inflammation, central nervous system disorders, cardiovascular disease, Alzheimer's disease and dermatological disease.

Protein kinase C has been linked to several different aspects of diabetes. Excessive activity of protein kinase C has been linked to insulin signaling defects and

therefore to the insulin resistance seen in Type II diabetes. Karasik, A. et al. J. Biol. Chem. 265: 10226-10231 (1990); Chen, K.S. et al. Trans.Assoc.Am.Physicians 104: 206-212 (1991); Chin, J.E. et al J. Biol. Chem. 268: 6338-6347 (1993). In addition, studies have demonstrated a marked increase in protein kinase C activity in tissues known to be susceptible to diabetic complications when exposed to hyperglycemic conditions. Lee, T.S. et al., J. Clin. Invest. 83: 90-94 (1989); Lee, T.S. et al. Proc. Natl. Acad.Sci USA 86: 5141-5145 (1989); Craven, P.A. and DeRubertis, F.R. J. Clin. Invest. 83: 1667-1675 (1989); Wolf, B.A. J. Clin. Invest. 87: 1643-1648 (1991).

Protein kinase C activity has long been associated with cell growth, tumor promotion and cancer. Rotenberg, S.A. and Weinstein, I.B. Biochem. Mol.Aspects Sel. Cancer 1: 25-73 (1991). Ahamd et al., Molecular Pharmacology: 43, 858-862 (1993). It is known that inhibitors of protein kinase C inhibitors are effective in preventing tumor growth in animals. Meyer, T. et al. Int. J. Cancer 43: 851-856 (1989); Akinagaka, S. et al. Cancer Res. 51: 4888-4892 (1991). More recently, the protein kinase C β inhibitor, Enzastaurin (LY317615.HCl) was shown to have a direct tumor effect by inducement of apoptosis and suppression of proliferating cultured tumor cell, in particular on human glioblastoma and colon carcinoma. Graff et al, Cancer Res.16: 7462- 7469 (2005). The compounds of the present invention also act as multidrug reversal (MDR) agents making them effective compounds when administered in conjunction with other chemotherapeutic agents.

Protein kinase C inhibitors have been shown to block inflammatory responses such as neutrophil oxidative burst, CD3 down-regulation in T-lymphocytes, and phorbol-induced paw edema. Towemy, B. et al. Biochem. Biophys. Res. Commun. 171: 1087-1092 (1999); Mulqueen, M.J. et al. Agents Actions 37: 85-89 (1992). Accordingly, as inhibitors of PKC, the present compounds are useful in treating inflammation.

Protein kinase C activity plays a central role in the functioning of the central nervous system. Huang, K.P. Trends Neurosci. 12: 425-432 (1989). In addition, protein kinase C inhibitors have been shown to prevent the damage seen in focal and central ischemic brain injury and brain edema. Hara, H. et al. J. Cereb. Blood Flow Metab. 10: 646-653(1990); Shibata, S. et al. Brain Res. 594: 290-294 (1992). Protein kinase C has also been determined to be implicated in Alzheimer's disease. Shimohama, S. et al. Neurology 43: 1407-1413 (1993). Accordingly, the compounds of the present invention are useful in treating Alzheimer's disease and ischemic brain injury.

Protein kinase C activity also plays an important role in cardiovascular disease. Increased protein kinase C activity in the vasculature has been shown to cause increased vasoconstriction and hypertension. A known protein kinase C inhibitor prevented this increase. Bilder, G.E. et al. J. Pharmacol. Exp. Ther. 252: 526-430 (1990). Because protein kinase C inhibitors demonstrate inhibition of the neutrophil oxidative burst, protein kinase C inhibitors are also useful in treating cardiovascular ischemia and improving cardiac function following ischemia. Muid, R.E. et al. FEBS Lett. 293: 169-172 (1990); Sonoki, H. et al. Kokyu-To Junkan 37: 669-674 (1989). The role of protein kinase C in platelet function has also been investigated and as shown elevated protein kinase C levels being correlated with increased response to agonists. Bastyr III, E.J. and Lu, J. Diabetes 42: (Suppl. 1) 97A (1993). PKC has been implicated in the biochemical pathway in the platelet-activity factor modulation of microvascular permeability. Kobayashi et al., Amer. Pjus. Soc. H1214-H1220 (1994). Potent protein kinase C inhibitors have been demonstrated to affect agonist-induced aggregation in platelets. Toullec, D. et al. J. Biol. Chem. 266: 15771-15781 (1991). Protein kinase C inhibitors also block agonist-induced smooth muscle cell proliferation. Matsumoto, H. and Sasaki, Y. Biochem. Biophys. Res. Commun. 158: 105-109 (1989). Therefore, the present compounds are useful in treating cardiovascular disease, atherosclerosis and restenosis.

Abnormal activity of protein kinase C has also been linked to dermatological disorders such as psoriasis. Horn, F. et al. J. Invest. Dermatol. 88: 220-222 (1987); Raynaud, F. and Evain-Brion, D. Br. J. Dermatol. 124: 542-546 (1991). Psoriasis is characterized by abnormal proliferation of keratinocytes. Known protein kinase C inhibitors have been shown to inhibit keratinocyte proliferation in a manner that parallels their potency as PKC inhibitors. Hegemann, L. et al. Aarch. Dermatol. Res. 283: 456-460 (1991); Bollag, W.B. et al. J. Invest. Dermatol. 100: 240-246 (1993). Accordingly, PKC inhibitors are useful in treating psoriasis.

The compounds of the invention are also isozyme-selective. The compounds preferentially inhibit protein kinase C beta-1 and beta-2 isozyme and optionally the alpha isozyme, over the remaining protein kinase C isozymes, i.e., gamma, delta, epsilon, zeta nad eta. Accordingly, compounds of the present invention inhibit beta-1 and beta-2 isozymes of protein kinase C, and optionally the alpha isozyme at much lower concentrations with minimal inhibition of the other PKC isozymes.

The compounds of the present invention are particularly useful in treating those disease states in which protein kinase C isozyme beta-1, beta-2, and optionally alpha, are associated. For example, the elevated blood glucose levels found in diabetes leads to an isozyme-specific elevation of the beta-2 isozyme in vascular tissues. Proc. Natl Acad. Sci. USA 89: 11059-11065 (1992). A diabetes-linked elevation of the beta isozyme in human platelets has been correlated with their altered response to agonists. Bastyr III, E.J. and Lu, J. Diabetes 42: (Suppl 1) 97A (1993). The human vitamin D receptor has been shown to be selectively phosphorylated by protein kinase C beta. This phosphorylation has been linked to alterations in the functioning of the receptor. Hsieh et al, Proc. Natl. Acad. Sci. USA 88: 931509319 (1991); Hsieh et al. J. Biol. Chem. 268: 15118-15126 (1993). In addition, recent work has shown that the beta-2 isozyme is responsible for erythroleukemia cell proliferation while the alpha isozyme is involved in megakaryocyte differentiation in these same cells. Murray et al., J. Biol. Chem. 268: 15847-15853 (1993).

In addition to the beta-1 and beta-2 isozymes discussed above, the protein kinase C alpha isozyme has been shown to have potential in the treatment of nephropathy: a PKC-alpha knockout mouse having STZ-induced diabetes showed improved nephropathy. Menne et al, Diabetes 53: 2101-2109 (2005). PKC alpha was implicated in heart contractility, Braz et al. Nature Medicine 10: 248-254 (2004); and also in the regulation of Akt activation and eNOS phosphorylation in endothelial cells. Partovian & Simons, Cellular Signalling 16: 951-957 (2004).

ASSAY

Protein Kinase C beta 2 (PKC β II) catalyzes the production of ADP from ATP that accompanies the phosphoryl transfer to the PKC Pseudosubstrate peptide (A-> S, RFARKGSLRQKNV). This transfer is coupled to the oxidation of β -NADH through the activities of Pyruvate Kinase (PK) and Lactate Dehydrogenase (LDH). β -NADH conversion to NAD⁺ is monitored by the decrease in absorbance at 340 nm ($\epsilon = 6.22 \text{ cm}^{-1}\text{mM}^{-1}$) using a Molecular Devices SPECTRA max PLUS spectrophotometer.

A typical assay was carried out on a 96-well, clear microtiter plate in a Molecular Devices spectrophotometer for 20 minutes at 30°C in 0.1 mL of assay buffer containing 50 mM HEPES, pH 7.4, 5 nM PKC, 23 units of pyruvate kinase, 33 units of lactate

- 31 -

dehydrogenase, 0.15 mM peptide, 0.1 mM ATP, 1 mM DTT, 4 mM PEP, 8 mM MgCl₂, 0.3 mM NADH, 60 mM CaCl₂, 10 mg/mL PS, 50 ng/mL PMA, 7.5% DMSO and from about 10,000 nM to 0.169 nM compound inhibitor. Stock solutions of 3-sn-phosphatidyl-L-serine (PS) and phorbol-12-myristate-13-acetate (PMA) were sonicated for 30 seconds just prior to addition to assay buffer and assays were initiated by the addition of 100 μM ATP.

Steady-state kinetic parameters for the *bi-bi* kinase reaction were determined at saturating phospho-acceptor peptide substrate concentration (0.15 mM) by fitting initial velocity data to the Michaelis-Menten equation,

$$v = V_{\max}[S]/(K_M + [S])$$

where v is the measured initial velocity, $V_{\max}$ is the maximal enzyme velocity, $[S]$ is the ATP substrate concentration, and K_M is the Michaelis constant for ATP. Enzyme turnover values (k_{cat}) were calculated according to $k_{\text{cat}} = V_{\max}/[E]$, where $[E]$ is the total enzyme concentration. Enzyme inhibition constants (apparent K_i values) were determined by fitting initial velocities at variable inhibitor concentrations to a model for ATP competitive inhibition based on the Morrison equation). Morrison, J.F., Biochim. Biophys Acta 185: 269-286 (1969).

PHARMACEUTICAL COMPOSITIONS/FORMULATIONS, DOSAGING AND MODES OF ADMINISTRATION

Methods of preparing various pharmaceutical compositions with a specific amount of active compound are known, or will be apparent, to those skilled in this art. In addition, those of ordinary skill in the art are familiar with formulation and administration techniques. Such topics would be discussed, e.g. in Goodman and Gilman's The Pharmaceutical Basis of Therapeutics, current ed., Pergamon Press; and Remington's Pharmaceutical Sciences, current ed., Mack Publishing, Co., Easton, PA. These techniques can be employed in appropriate aspects and embodiments of the methods and compositions described herein. The following examples are provided for illustrative purposes only and are not meant to serve as limitations of the present invention.

The compounds of formula (A) and (B) may be provided in suitable topical, oral and parenteral pharmaceutical formulations for use in the treatment of PKCβII mediated diseases. The compounds of the present invention may be administered orally as

tablets or capsules, as oily or aqueous suspensions, lozenges, troches, powders, granules, emulsions, syrups or elixirs. The compositions for oral use may include one or more agents for flavoring, sweetening, coloring and preserving in order to produce pharmaceutically elegant and palatable preparations. Tablets may contain pharmaceutically acceptable excipients as an aid in the manufacture of such tablets. As is conventional in the art these tablets may be coated with a pharmaceutically acceptable enteric coating, such as glyceryl monostearate or glyceryl distearate, to delay disintegration and absorption in the gastrointestinal tract to provide a sustained action over a longer period.

Formulations for oral use may be in the form of hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin. They may also be in the form of soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, such as peanut oil, liquid paraffin or olive oil.

Aqueous suspensions normally contain active ingredients in admixture with excipients suitable for the manufacture of an aqueous suspension. Such excipients may be a suspending agent, such as sodium carboxymethyl cellulose, methyl cellulose, hydroxypropylmethyl cellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; a dispersing or wetting agent that may be a naturally occurring phosphatide such as lecithin, a condensation product of ethylene oxide and a long chain fatty acid, for example polyoxyethylene stearate, a condensation product of ethylene oxide and a long chain aliphatic alcohol such as heptadecaethylenoxycetanol, a condensation product of ethylene oxide and a partial ester derived from a fatty acid and hexitol such as polyoxyethylene sorbitol monooleate or a fatty acid hexitol anhydrides such as polyoxyethylene sorbitan monooleate.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to known methods using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation may also be formulated as a suspension in a non toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringers solution and isotonic sodium chloride solution. For this purpose any bland fixed oil may be employed including synthetic mono- or

diglycerides. In addition fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of formula (A) and (B) may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at about 25 Celsius but liquid at rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter and other glycerides.

For topical use preparations, for example, creams, ointments, jellies solutions, or suspensions, containing the compounds of the present invention are employed.

The compounds of formula (A) and (B) may also be administered in the form of liposome delivery systems such as small unilamellar vesicles, large unilamellar vesicles and multimellar vesicles. Liposomes can be formed from a variety of phospholipides, such as cholesterol, stearylamine or phosphatidylcholines.

Dosage levels of the compounds of the present invention are of the order of about 0.5 mg/kg body weight to about 100 mg/kg body weight. A preferred dosage rate is between about 30 mg/kg body weight to about 100 mg/kg body weight. It will be understood, however, that the specific dose level for any particular patient will depend upon a number of factors including the activity of the particular compound being administered, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy. To enhance the therapeutic activity of the present compounds they may be administered concomitantly with other orally active antidiabetic compounds such as the sulfonylureas, for example, tolbutamide and the like.

For administration to the eye, a compound of the present invention is delivered in a pharmaceutically acceptable ophthalmic vehicle such that the compound is maintained in contact with the ocular surface for a sufficient time period to allow the compound to penetrate the cornea and/or sclera and internal regions of the eye, including, for example, the anterior chamber, posterior chamber, vitreous body, aqueous humor, vitreous humor, cornea, iris/ciliary body, lens, choroid/retina and sclera. The pharmaceutically acceptable ophthalmic vehicle may be an ointment, vegetable oil, or an encapsulating material. A compound of the invention may also be injected directly into the vitreous humor or aqueous humor.

Further, a compound may be also be administered by well known, acceptable methods, such as subtenon and/or subconjunctival injections. As is well known in the ophthalmic art, the macula is comprised primarily of retinal cones and is the region of maximum visual acuity in the retina. A Tenon's capsule or Tenon's membrane is disposed on the sclera. A conjunctiva covers a short area of the globe of the eye posterior to the limbus (the bulbar conjunctiva) and folds up (the upper cul-de-sac) or down (the lower cul-de-sac) to cover the inner areas of the upper eyelid and lower eyelid, respectively. The conjunctiva is disposed on top of Tenon's capsule. The sclera and Tenon's capsule define the exterior surface of the globe of the eye. For treatment of age related macular degeneration (ARMD), choroid neovascularization, retinopathies (such as diabetic retinopathy (including macular edema), retinopathy of prematurity), retinitis, uveitis, cystoid macular edema (CME), glaucoma, and other diseases or conditions of the posterior segment of the eye, it is preferable to dispose a depot of a specific quantity of an ophthalmically acceptable pharmaceutically active agent directly on the outer surface of the sclera and below Tenon's capsule. In addition, in cases of ARMD and CME it is most preferable to dispose the depot directly on the outer surface of the sclera, below Tenon's capsule, and generally above the macula.

The compounds may be formulated as a depot preparation. Such long-acting formulations may be administered by implantation (for example, subcutaneously or intramuscularly) intramuscular injection or by the above mentioned subtenon or intravitreal injection. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

Within particularly preferred embodiments of the invention, the compounds may be prepared for topical administration in saline (combined with any of the preservatives and antimicrobial agents commonly used in ocular preparations), and administered in eyedrop form. The solution or suspension may be prepared in its pure form and administered several times daily. Alternatively, the present compositions, prepared as described above, may also be administered directly to the cornea.

Within preferred embodiments, the composition is prepared with a muco-adhesive polymer which binds to cornea. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example, as an

emulsion in an acceptable oil) or ion-exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for hydrophobic compounds is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The cosolvent system may be a VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) contains VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g. polyvinyl pyrrolidone; and other sugars or polysaccharides may be substituted for dextrose.

Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid- or gel-phase carriers or excipients. Examples of such carriers or excipients include calcium carbonate, calcium phosphate, sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Some of the compounds of the invention may be provided as salts with pharmaceutically compatible counter ions. Pharmaceutically compatible salts may be

formed with many acids, including hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free-base forms.

The preparation of preferred compounds of the present invention is described in detail in the following examples, but the artisan will recognize that the chemical reactions described may be readily adapted to prepare a number of other compounds of the invention. For example, the synthesis of non-exemplified compounds according to the invention may be successfully performed by modifications apparent to those skilled in the art, e.g., by appropriately protecting interfering groups, by changing to other suitable reagents known in the art, or by making routine modifications of reaction conditions. Alternatively, other reactions disclosed herein or known in the art will be recognized as having applicability for preparing other compounds of the invention.


EXAMPLES

The examples and preparations provided below further illustrate and exemplify the compounds of the present invention and methods of preparing such compounds. It is to be understood that the scope of the present invention is not limited in any way by the scope of the following examples and preparations. In the following examples molecules with a single chiral center, unless otherwise noted, exist as a racemic mixture. Those molecules with two or more chiral centers, unless otherwise noted, exist as a racemic mixture of diastereomers. Single enantiomers/diastereomers may be obtained by methods known to those skilled in the art.

The structures of the compounds are confirmed by either elemental analysis or NMR, where peaks assigned to the characteristic protons in the title compound are presented where appropriate. ^1H NMR shift (δ_{H}) are given in parts per million (ppm) down field from an internal reference standard.

The invention will now be described in reference to the following examples. These examples are not to be regarded as limiting the scope of the present invention, but shall only serve in an illustrative manner. Table 1 provides a full listing of the compounds of the present invention and includes relevant ^1H NMR data and K_i values as available.

EXAMPLE A1: N4-(6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N2-ethyl-5-fluoropyrimidine-2,4-diamine



Intermediate A1(i): To a 0 °C solution of 5-tert-butyl 1-ethyl 3-amino-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-1,5-dicarboxylate (16.2 g, 49.9 mmol) in THF (100 mL) was added NaH (2.4 g, 59.9 mmol) in 3 portions. The reaction was stirred for 15 min in an ice bath, then ethyl chloroformate (6.5 g, 59.9 mmol) was added over 10 min. The reaction was warmed to room temperature and stirred for 16h, then quenched with NH₄Cl (sat) and extracted with EtOAc (2 x 50 mL). The combined extracts were washed with brine then dried (MgSO₄) filtered and concentrated to give the desired compound **A1(i)** (19.8 g, 99%). Mass spectrum: Calcd for C₁₈H₂₉N₄O₆ (M+H): 397. Found 397.

Intermediate A1(ii): Ethyl 5-(chlorocarbonyl)-3-[(ethoxycarbonyl)amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate.

To a solution of **A1(i)** (19.8 g, 49.9 mmol) in dioxane (20 mL) was added HCl (60 mL, 4M in dioxane). The reaction was stirred at room temperature for 3h then concentrated and dried under vacuum. The HCl salt of ethyl 3-[(ethoxycarbonyl)amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate was taken up in CH₂Cl₂ (60 mL). DIPEA (16.1 g, 125 mmol) was added and the reaction mixture was cooled in an ice bath. Phosgene (30 mL, 20% in toluene) was added slowly then the reaction was

warmed to room temperature and run overnight. The reaction was concentrated then taken up in EtOAc (100 mL) and water (100 mL). The aqueous phase was extracted with EtOAc (2 x 25 mL) then the combined organic extracts were washed with brine, dried (MgSO_4), filtered and concentrated. The crude material was purified by silica gel column chromatography using CH_2Cl_2 -2% 7N NH_3/MeOH in CH_2Cl_2 to give the title compound A1(ii) as a white solid (9.58 g, 54%). Mass spectrum: Calcd for $\text{C}_{14}\text{H}_{20}\text{ClN}_4\text{O}_5$ (M+H): 359. Found 359.

Intermediate A1(iii): Ethyl 3-[(ethoxycarbonyl)amino]-6,6-dimethyl-5-[[[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate. To a sealed tube was added (5S)-1,2,2,5-tetramethylpiperazine (2.8 g, 20 mmol), DIPEA (7.63 g, 59.1 mmol) and THF (40 mL) followed by A1(ii) (7.1 g, 20 mmol). The tube was sealed and placed in an oil bath at 80 °C and heated for 16 h. The reaction was cooled to room temperature then concentrated and purified by silica gel column chromatography using CH_2Cl_2 -3% (7N NH_3/MeOH) in CH_2Cl_2 to give the title compound A1(iii) (4.62 g, 51%). Mass spectrum: Calcd for $\text{C}_{22}\text{H}_{37}\text{N}_6\text{O}_5$ (M+H): 465. Found 465.

Intermediate A1(iv): 6,6-dimethyl-5-[[[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

To a microwave vial was added A1(iii) (4.6 g, 10 mmol), MeOH (45 mL) and LiOH (6.4 g, 40 mmol). The reaction was heated at 110 °C in the microwave for 20 min. The crude reaction mixture was concentrated and taken up in THF (75 mL). The insoluble material was filtered off and the filtrate was concentrated to give the title compound A1(iv) (2.3 g, 72%). ^1H NMR (300 MHz, CDCl_3) ppm 1.02 (s, 3 H), 1.08- 1.18 (m, 6 H), 1.66 (s, 3 H), 1.76 (s, 3 H), 2.22 (s, 3 H), 2.25- 2.41 (m, 1 H), 2.59- 2.70 (m, 1 H), 2.70- 2.91 (m, 2 H), 3.42- 3.57 (m, 1 H), 4.28- 4.51 (m, 2 H). Mass spectrum: Calcd for $\text{C}_{16}\text{H}_{29}\text{N}_6\text{O}$ (M+H): 321. Found 321.

Intermediate A1(v): N-(2-chloro-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

To a solution of A1(iv) (1.92 g, 5.9 mmol) and 2,4-dichloro-5-fluoropyrimidine (1.0 g, 5.9 mmol) in DMSO (8 mL) was added potassium dihydrogenphosphate (0.82 g, 5.9 mmol) followed by H_3PO_4 (0.12 g, 1.2 mmol). The reaction was placed in a 90 °C oil bath and heated for 20h. The crude reaction was cooled to room temperature then poured into ice cold NaHCO_3 (30 mL) and extracted with EtOAc (2 x 30 mL). The combined organic extracts were washed with brine then dried (MgSO_4), filtered and concentrated. The

- 39 -


crude solid was triturated with EtOAc then the filtrate was purified by silica gel column chromatography using 1%-4% 7N NH₃/MeOH in CH₂Cl₂ to provide the title compound A1(v) as a white solid (1.4 g, 53%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.90 (s, 3 H), 1.00 (s, 3 H), 1.06 (d, *J*=6.4 Hz, 3 H), 1.58 (s, 3 H), 1.69 (s, 3 H), 2.08 (s, 3 H), 2.14-2.25 (m, 1 H), 2.55- 2.64 (m, 2 H), 2.73- 2.89 (m, 1 H), 3.38- 3.51 (m, 1 H), 4.51- 4.72 (m, 2 H), 7.95- 8.45 (m, 1 H), 10.68 (s, 1 H), 11.81- 12.75 (m, 1 H). Mass spectrum: Calcd for C₂₀H₂₉ClN₈O (M+H): 451. Found 451.

Example A1: N4-(6,6-dimethyl-5-[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-yl)-N2-ethyl-5-fluoropyrimidine-2,4-diamine. To a microwave vial was added A1(v) (200 mg, 0.44 mmol) and ethylamine (5 mL, 2M in MeOH). The vial was heated to 150 °C in the microwave for 1.5h. The crude sample was concentrated then purified by silica gel column chromatography using 1%-4% 7N NH₃/MeOH in CH₂Cl₂ to give the desired product A1 as a yellow solid (161 mg, 79%). See Table 1 below for NMR data.

EXAMPLES A2-A9:

Examples A2 through A9 were prepared using methods analogous to Example A1 above. See Table 1 below for names and NMR data

EXAMPLE A10: 5-[(8*S*)-6,8-dimethyl-6,9-diazaspiro[4.5]dec-9-yl]carbonyl)-*N*-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine



Intermediate A10(i): 5-[(8*S*)-6,8-dimethyl-6,9-diazaspiro[4.5]dec-9-yl]carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine

Using the same procedure as in the preparation of intermediate A1(iv) above, the titled intermediate was purified by flash chromatography eluting with methylene chloride containing 5% 7N NH₃ in methanol to a yellow solid (506 mg, 90%). For the synthesis of spiro cyclopentyl piperazine, please refer to the general synthesis of piperazines. ¹H

NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.13 (3 H, d, $J=6.32$ Hz), 1.36 – 1.41 (1 H, m), 1.52- 1.64 (8 H, m), 1.66 (3 H, s), 1.77 (3 H, s), 2.20- 2.22 (3 H, m), 2.23- 2.29 (1 H, m), 2.60 (1 H, dd, $J=11.75, 3.66$ Hz), 2.78- 2.89 (2 H, m), 3.49 (2 H, s), 3.52 – 3.56 (1 H, m), 3.64 (1 H, s), 4.33- 4.44 (2 H, m).

Intermediate A10(ii): 4-Chloro-5-fluoro-2-methylpyrimidine

Sodium hydride (60%, 5.0g, 125 mmol) was washed with hexane to remove the mineral oil and dried, then suspended in THF (50 mL) and cooled to 0°C. Ethyl fluoroacetate (13.30 g, 125 mmol) and ethyl formate (15.14 mL, 187 mmol) were mixed together and added to the stirring suspension. The reaction was slowly warmed to ambient temperature and stirred 3 days. The solvent was removed. A mixture of acetamidine hydrochloride (11.81 g, 125 mmol), sodium ethoxide (8.86 g, 125 mmol), and ethanol (60 mL) were added to the reaction followed by refluxing overnight. The ethanol was removed under reduced pressure. The residue was dissolved in a minimum of water and acidified to pH = 6 with concentrated HCl. The crude products were then extracted by salting out from the aqueous phase and washing exhaustively with 4:1

CHCl₃/isopropanol. The combined organic phases were dried (MgSO₄) and evaporated. The crude solid was purified by silica gel chromatography eluting with 5-90%

EtOAc/hexane to give a white solid (0.95 g, 6%). $R_f = 0.08$ (75% EtOAc/hexane). ¹H

NMR (400MHz, DMSO-*d*₆): δ 2.25 (d, $J = 1.0$ Hz, 3H), 7.93 (d, $J = 3.8$ Hz, 1H), 12.95 (br, 1H). LCMS 129. 2-Methyl-5-fluorouracil (1.04g, 7.21 mmol) and *N,N*-

dimethylaniline (1.80 mL) were heated in POCl₃ at 110°C for 90 minutes. After cooling, the reaction was added carefully to ice. The product was extracted with diethylether.


The ether layer was washed with sequentially with 2N HCl, water, and brine followed by drying (MgSO₄). The ether was carefully removed under reduced pressure to give a volatile liquid (0.39g, 34%) which was used without further purification. $R_f = 0.26$ (10% EtOAc/hexane). ¹H NMR (400MHz, DMSO-*d*₆): δ 3.91 (s, 3H), 8.79 (s, 1H).

Example A10: 5-[[[(8S)-6,8-dimethyl-6,9-diazaspiro[4.5]dec-9-yl]carbonyl]-N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine

Using the same procedure as described above for intermediate A1(v), title compound A10 was purified to a white powder (49.7 mg, 24%). See Table 1 below for NMR data.

- 41 -

EXAMPLE A11: N^4 -(5-[(*(2S,5R)*-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl]carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)- N^2 -ethyl-5-fluoropyrimidine-2,4-diamine



Intermediates A11(ii) and A11(iii): (*(2S,5R)*)-1-benzyl-2,5-dimethylpiperazine and A11(iii) (*(2R,5S)*)-1-benzyl-2,5-dimethylpiperazine

Compound A11(i) (800 g, 7 mol) was dissolved in 12.8 L of EtOH. A solution of benzyl bromide (1 kg, 5.8 mol) in 1.6 L of EtOH was added dropwise at room temperature. The reaction mixture was stirred at room temperature overnight. TLC (CH₂Cl₂: MeOH = 10:1) showed the reaction was complete. The solvent was removed in vacuum. The residue was diluted with 32 L of water and the resulting mixture was stirred at room temperature for 30 minutes, then filtered. The filtrate was extracted with CH₂Cl₂ (8 L×3). The organic phase was dried over Na₂SO₄ and concentrated to dryness to give a mixture of compound A11(ii) and A11(iii) (800 g, 56%) as brown oil.

Intermediate A11(iv) (*(2S,5R)*)-1-benzyl-2,5-dimethylpiperazine):

To a stirred solution of compound A11(ii) and A11(iii) (300 g, 1.48 mol) in MeOH (1 L) was added a solution of L-(+)-tartaric acid (444 g, 2.96 mol) in MeOH (2 L).

- 42 -

After the mixture was stirred at room temperature for 20 minutes, then left standing at 0 °C for 24 hours. The solid formed was collected by filtration, which was re-crystallized from MeOH (2 L) to give a solid. The solid was dissolved in 1.6 L of water. The resulting mixture was basified to pH = 9-10 with saturated aqueous Na₂CO₃. Then the mixture was extracted with CH₂Cl₂ (1.5 L×2). The organic phase was dried over Na₂SO₄ and concentrated to dryness to give A11(iv) (122 g, 81%) as yellow liquid. ¹H NMR (400 MHz, CDCl₃) δ 7.21-7.13(m, 5H), 4.02(d, 1H), 3.00(d, 1H), 2.82-2.51(m, 4H), 2.15(m, 1H), 1.57(m, 2H), 1.05(d, 3H), 0.85(d, 3H).

Intermediate A11(v): (2*S*,5*R*)-1-benzyl-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazine.

To a microwave vial was added A11(iv) (7.50g, 36.7 mmol), (bromomethyl)tetrahydropyran (6.57g, 36.7 mmol), triethylamine (12.8 mL, 91.8 mmol), and MeOH (9 mL). The reaction mixture was heated to 150 °C for 2 hours in the microwave, at which time the resulting solid was triturated with MeOH, filtered, and dried to give the desired product as a white solid A11(v) (6.1 g, 55%). Mass Spectrum: Calcd for C₁₉H₃₁N₂O (M+H): 303. Found: 303.


Intermediate A11(vi): (2*R*,5*S*)-2,5-dimethyl-1-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazine.

To a solution of A11(v) (6.10 g, 20.0 mmol) in MeOH (200 mL) was added 10% palladium on carbon (0.600 g, 0.570 mmol). The suspension was evacuated and backfilled with hydrogen (x 3) and allowed to stir for 15 h under hydrogen. The suspension was filtered through celite, the filter cake washed with CH₂Cl₂, and the filtrate concentrated to give the desired product as a colorless oil A11(vi) (4.3 g, 88%). Mass Spectrum: Calcd. for C₁₂H₂₅N₂O (M+H): 213. Found: 213.

Example A11: *N*⁴-(5-([(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazin-1-yl]carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-yl)-*N*²-ethyl-5-fluoropyrimidine-2,4-diamine.

The title compound was prepared using methods analogous to Example A1 except that A11(vi) was substituted in place of (5*S*)-1,2,2,5-tetramethylpiperazine. See Table 1 below for NMR data.

EXAMPLE AA1: 5-[[[(2S,5R)-4-ethyl-2,5-dimethylpiperazin-1-yl]carbonyl]-N-(5-fluoro-2-methoxypyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine




A solution of 5-[[[(2S,5R)-4-ethyl-2,5-dimethylpiperazin-1-yl]carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine (289 mg, 0.9 mmol) and 4-chloro-5-fluoro-2-methoxypyrimidine (257 mg, 2 eq) in 5 mL of 50% acetic acid in water was heated in a microwave for 30 min at 80°C. Purification as described in Example A1 afforded the title compound **AA1** as a white powder (13.1 mg, 3%). See Table 1 below for NMR data.

EXAMPLES AA2 – AA5:

Examples **AA2** through **AA5** were prepared using methods analogous to Example **AA1**. See Table 1 below for names and NMR data.

EXAMPLE B1: N²-ethyl-5-fluoro-N⁴-(5-[[[(2S,5R)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl]pyrimidine-2,4-diamine acetate salt



Intermediate B1(i): Tert-butyl 3-amino-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carboxylate.

To a slurry of 5-tert-butyl 1-ethyl 3-amino-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-1,5-dicarboxylate (25.00 g, 77.1 mmol) in MeOH (50 mL) was added LiOH (1.92 g, 77.1 mmol). The reaction was stirred at room temperature for 2h then concentrated. The crude reaction mixture was taken up in EtOAc (50 mL) then washed with NaHCO₃ (20 mL) and water (20 mL). The organic layer was dried (MgSO₄), filtered and concentrated to give an orange solid which was triturated with ACN then filtered and rinsed with ACN (50 mL) to give the title compound B1(i) as a white solid (14.8 g, 76%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.40- 1.46 (m, 9 H), 1.47- 1.54 (m, 6 H), 4.03- 4.17 (m, 2 H), 4.95 (br. s., 1 H), 11.15 (s, 1 H). Mass spectrum: Calcd for C₁₂H₂₁N₄O₂ (M+H): 253. Found 253.

Intermediate B1(ii): tert-butyl 3-[(2-chloro-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carboxylate.

To a solution of B1(i) (8.01 g, 31.7 mmol) and 2,4-dichloro-5-fluoro pyrimidine (5.30 g, 31.7 mmol) in DMSO (40 mL) was added potassium dihydrogenphosphate (4.32 g, 31.7 mmol) followed by H₃PO₄ (0.62 g, 6.4 mmol). The reaction was placed in a 95 °C oil bath and heated for 20h. The crude reaction was cooled to room temperature then poured into ice cold NaHCO₃ (sat, 100 mL) followed by addition of EtOAc (75 mL). The resulting mixture was filtered to give the desired compound B1(ii) as a white solid (6.1 g, 50%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.35- 1.50 (m, 9 H), 1.56- 1.65 (m, 6 H), 4.37- 4.64 (m, 2 H), 8.09- 8.42 (m, 1 H), 10.70 (br. s., 1 H), 12.53 (br. s., 1 H). Mass spectrum: Calcd for C₁₆H₂₁ClFN₆O₂ (M+H): 383. Found 383.

Intermediate B1(iii): tert-butyl 3-[[2-(ethylamino)-5-fluoropyrimidin-4-yl]amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carboxylate

To a sealed tube was added B1(ii) (7.64 g, 20.0 mmol) and ethylamine (40.0 mL, 2M in MeOH). The sealed tube was heated in an oil bath at 150 °C for 16 h. The reaction was allowed to cool to room temperature and the resulting solid was filtered and rinsed with cold MeOH to give the title compound B1(iii) as a white solid (6.1 g, 78%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.10 (t, *J*=7.2 Hz, 3 H), 1.39- 1.48 (m, 9 H), 1.53- 1.61 (m, 6 H), 3.18- 3.27 (m, 2 H), 4.23- 4.37 (m, 2 H), 7.06 (br. s., 1 H), 7.93 (s, 1 H), 10.10 (br. s., 1 H), 12.45 (br. s., 1 H). Mass spectrum: Calcd for C₁₈H₂₇FN₇O₂ (M+H): 392. Found 392.

Intermediate B1(iv): 5-tert-butyl 1-ethyl 3-[[2-(ethylamino)-5-fluoropyrimidin-4-yl]amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-1,5-dicarboxylate


A solution of B1(iii) (5.14 g, 13.1 mmol) in THF (60 mL) and DIPEA (4.24 g, 13.1 mmol) was cooled in an ice bath then ethylchloroformate (1.42 g, 13.1 mmol) was added dropwise. The reaction mixture was slowly warmed to room temperature and stirred for 5h, then quenched with water (50 mL) and extracted with EtOAc (2x 100 mL). The combined extracts were washed with brine (50 mL) then dried (MgSO₄), filtered and concentrated. The crude material was purified by silica gel column chromatography using 10-40% EtOAc in CH₂Cl₂ to provide the title compound B1(iv) as a white foam (5.8 g, 95%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.01- 1.11 (m, 3 H), 1.33 (t, *J*=7.2 Hz, 3 H), 1.38- 1.49 (m, 9 H), 1.70- 1.83 (m, 6 H), 3.13- 3.27 (m, 2 H), 4.28- 4.50 (m, 4 H), 6.54- 6.86 (m, 1 H), 7.94 (m, 1 H), 10.06- 10.41 (m, 1 H). Mass spectrum: Calcd for C₂₁H₃₁FN₇O₄ (M+H): 464. Found 464.

Intermediate B1(v): Ethyl 5-(chlorocarbonyl)-3-[[2-(ethylamino)-5-fluoropyrimidin-4-yl]amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate

To a suspension of B1(iv) (4.43 g, 9.6 mmol) in dioxane (25 mL) was added HCl (20 mL, 4M in dioxane). The reaction was stirred at room temperature for 2 h then concentrated. To a solution of the HCl salt of ethyl 3-[[2-(ethylamino)-5-fluoropyrimidin-4-yl]amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate (1.27 g, 2.7 mmol) in CH₂Cl₂ (40 mL) was added DIPEA (1.57 g, 12.1). The reaction was cooled to -78 °C and triphosgene (0.48 g, 0.6 mmol) was added in a solution of CH₂Cl₂ (10 mL) with an addition funnel over 15 min. The reaction was quenched at -78 °C with water then warmed to room temperature. The mixture was made to pH 8-9 with NaHCO₃ and extracted with CH₂Cl₂ (2 x 15 mL). The combined extracts were washed with brine (15 mL) then dried, filtered and concentrated. The crude material was purified by silica gel column chromatography using 0-3% 7N NH₃/MeOH in CH₂Cl₂ to give the desired compound B1(v) as a white foam (489 mg, 36%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.07 (t, *J*=7.2 Hz, 3 H), 1.33 (t, *J*=7.2 Hz, 3 H), 1.78- 1.84 (m, 6 H), 3.12- 3.28 (m, 2 H), 4.42 (q, *J*=7.2 Hz, 2 H), 4.78 (s, 2 H), 6.69- 6.97 (m, 1 H), 7.77- 8.08 (m, 1 H), 10.41 (s, 1 H). Mass spectrum: Calcd for C₁₇H₂₂ClFN₇O₃ (M+H): 426. Found 426.

Preparation of Side-chain B1(c):

- 46 -

**Intermediate B1(a): (2S,5R)-1-benzyl-2,5-dimethylpiperazine**

To a solution of starting material (3.00 g, 9.85 mmol) in CH₂Cl₂ (100 mL) was added 4 N HCl in 1,4-dioxane (20 mL). The solution was stirred for 1 hour, at which time the volatiles were removed *in vacuo* to yield the desired product as a white solid in quantitative yield. Mass Spectrum: Calcd for C₁₃H₂₁N₂ (M+H): 205. Found: 205.

Intermediate B1(b): (2S,5R)-1-benzyl-4-(3-methoxypropyl)-2,5-dimethylpiperazine

To a microwave vial was added starting material (1.60g, 5.77 mmol), 1-bromo-3-methoxypropane (3.09 g, 20.2 mmol), triethylamine (6.03 mL, 43.3 mmol), THF (6 mL), and MeOH (6 mL). The suspension was heated at 150 °C in the microwave for 2 hours. The cooled solution was diluted with EtOAc (20 mL) and the organic layer washed with NaHCO₃ (sat., aq.) (3 x 25 mL) and brine (1 x 25 mL). The organic layer was dried over MgSO₄ and concentrated to give the desired product as brown oil (1.5 g, 92%). Mass Spectrum: Calcd for C₁₇H₂₉N₂O (M+H): 277. Found: 277.

Intermediate B1(c): (2R,5S)-1-(3-methoxypropyl)-2,5-dimethylpiperazine

To a solution of starting material (1.50 g, 5.43 mmol) in MeOH (50 mL) was added 10% palladium on carbon (0.150 g, 1.41 mmol). The suspension was evacuated and backfilled with hydrogen (x 3) and allowed to stir for 15 h under hydrogen. The suspension was filtered through Celite, the filter cake washed with CH₂Cl₂, and the filtrate concentrated to give the desired product as a brown foam (0.83 g, 82%). Mass Spectrum: Calcd. for C₁₀H₂₃N₂O (M+H): 187. Found: 187.

Example B1: N²-ethyl-5-fluoro-N⁴-(5-[[[(2S,5R)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl]pyrimidine-2,4-diamine acetate salt

To a sealed tube was added B1(v) (393 mg, 0.92 mmol) in a solution of THF (10 mL) followed by DIPEA (537 mg, 4.2 mmol) and B1(c) (172 mg, 0.92 mmol). The reaction was placed in an 85 °C oil bath and heated for 16 h. The crude reaction was concentrated then taken up in MeOH (5 mL) and Et₃N (5 mL) then stirred for an additional 16 h. Preparative HPLC using 5-50% ACN/H₂O (0.1% AcOH) provided the title compound B1 as a white solid (185 mg, 36%). See Table 1 below for NMR data.

EXAMPLES B2 – B5:

Examples B2 to B5 were prepared using methods analogous to Example B1 above. See Table 1 below for names and NMR data.

EXAMPLE B6: 4-[(6,6-dimethyl-5-[[[(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl]amino]pyrimidine-2-carbonitrile


Example B6 was prepared using methods analogous to Example B1 above except that (2R, 5S)-1,2,5-trimethylpiperazine hydrochloride was substituted in place of intermediate B1(b) and 4-chloropyrimidine-2-carbonitrile was substituted in place of 2,4-dichloro-5-fluoropyrimidine during the preparation of intermediate B1(ii). See Table 1 below for NMR data.

EXAMPLE B7: N-(5-fluoro-2-morpholin-4-ylpyrimidin-4-yl)-6,6-dimethyl-5-[[[(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

Example B7 was prepared using methods analogous to Example B1 above except that (2R,5S)-1,2,5-trimethylpiperazine hydrochloride was substituted in place of intermediate B1(b) and morpholine was substituted in place of ethylamine during the preparation of intermediate B1(iii). See Table 1 below for NMR data.

EXAMPLE C1: N²-ethyl-5-fluoro-N⁴-(5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)pyrimidine-2,4-diamine

- 48 -




Intermediate C1(i): Ethyl 3-[[2-(ethylamino)-5-fluoropyrimidin-4-yl]amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate.

To a suspension of 5-*tert*-butyl 1-ethyl 3-[[2-(ethylamino)-5-fluoropyrimidin-4-yl]amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-1,5-dicarboxylate (4.43 g, 9.6 mmol) in dioxane (25 mL) was added HCl (20 mL, 4M in dioxane). The reaction was stirred at room temperature for 2 h then concentrated to give the title compound **C1(i)** as the tri HCl salt (3.8 g, 84%). Mass spectrum: Calcd for $C_{16}H_{23}FN_7O_2$ (M+H): 364. Found 364.

Example C1: N²-ethyl-5-fluoro-N⁴-{5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl}pyrimidine-2,4-diamine

To a sealed tube was added **C1(i)** (400 mg, 0.85 mmol) in a solution of THF (10 mL) followed by DIPEA (492 mg, 3.8 mmol) and 4-fluoro-1-methylpiperidine-4-carbonyl chloride (228 mg, 1.3 mmol). The reaction was heated in an oil bath at 80 °C for 16 h then concentrated and taken up in MeOH (3 mL) and Et₃N (5 mL) and stirred for an addition 10 h. The crude reaction was concentrated then purified by silica gel column chromatography using 1-3 % 7N NH₃/MeOH in CH₂Cl₂ to give the title compound **C1** as a yellow solid (175 mg, 47%). See Table 1 below for NMR data.

EXAMPLE D1: N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine




Intermediate D1(i): N-(5-fluoro-2-vinylpyrimidin-4-yl)-6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.


To a septum capped sealed tube was added N-(2-chloro-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine (109 mg, 0.242 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (122 mg, 0.725 mmol), Na₂CO₃ (77 mg, 0.73 mmol) and dichloro[1,1'-

bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane adduct (40 mg, 0.048 mmol) followed by DME (2.5 mL) and water (0.5 mL). The reaction mixture was purged with argon for 2 min. then placed in a preheated 100 °C oil bath and stirred for 16 h. The crude reaction was concentrated, taken up in EtOAc (15 mL) and washed with water (10 mL). The EtOAc solution was dried (MgSO₄), filtered and concentrated. Purification by silica gel column chromatography using 1-3 % 7N NH₃/MeOH in CH₂Cl₂ provided the title compound D1(i) as a yellow solid (45 mg, 42%). ¹H NMR (300 MHz, CDCl₃) δ ppm 0.99 (s, 3 H), 1.05 (s, 3 H), 1.18 (d, *J*=6.4 Hz, 3 H), 1.72 (s, 3 H), 1.83 (s, 3 H), 2.19 (s, 3 H), 2.25- 2.36 (m, 1 H), 2.63- 2.75 (m, 1 H), 2.76- 2.95 (m, 2 H), 3.52- 3.73 (m, 1 H), 4.57- 4.81 (m, 2 H), 5.65 (m, 1 H), 6.40 (m, 1 H), 6.76 (m, 1 H), 8.13- 8.31 (m, 1 H), 8.53 (s, 1 H). Mass spectrum: Calcd for C₂₂H₃₂FN₈O (M+H): 443. Found 443.

Example D1: N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine. To a nitrogen purged flask containing D1(i) (54 mg, 0.12 mmol) was added Pd/C (6.5 mg, 0.006 mmol) and MeOH (3 mL). A H₂ balloon was applied and the reaction was stirred for 16h. The crude mixture was poured over a bed of Celite and rinsed with MeOH (30 mL) then concentrated and purified by column chromatography using 1-3% 7N NH₃/MeOH in CH₂Cl₂ to provide the title compound D1 as a pale yellow solid (33 mg, 61%). See Table 1 below for NMR data.

EXAMPLE E1: N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[(2*S*,5*R*)-2,4,5-trimethylpiperazin-1-yl]carbonyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine acetate salt





Intermediate E1(i): Tert-butyl 3-[(5-fluoro-2-vinylpyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carboxylate

To a septum capped sealed tube was added *tert*-butyl 3-[(2-chloro-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazole-5(1*H*)-carboxylate (2.00 g, 5.22 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2 dioxaborolane (2.41 g, 15.7 mmol), Na₂CO₃ (1.66 g, 15.7 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane adduct (850 mg, 1.04 mmol) followed by DME (50 mL) and water (12 mL). The reaction mixture was purged with argon for 2 min. then placed in a preheated 100 °C oil bath and stirred for 16 h. The crude reaction was concentrated then taken up in EtOAc (100 mL) and washed with water (50 mL). The EtOAc solution was dried (MgSO₄), filtered and concentrated. Purification was performed by silica gel column chromatography using 1-10 % 7N NH₃/MeOH in CH₂Cl₂ to provided the desired compound E1(i) as an orange solid (1.8 g, 91%). ¹H NMR (300 MHz, CDCl₃) δ ppm 1.45- 1.58 (m, 9 H), 1.65- 1.78 (m, 6 H), 4.45- 4.71 (m, 2 H), 5.58- 5.71 (m, 1 H), 6.34- 6.46 (m, 1 H), 6.68- 6.82 (m, 1 H), 7.99 (s, 1 H), 8.12- 8.32 (m, 1 H). Mass spectrum: Calcd for C₁₈H₂₄FN₆O₂ (M+H): 375. Found 375.

Intermediate E1(ii): tert-butyl 3-[(2-ethyl-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carboxylate.

To a nitrogen purged flask containing E1(i) (1.71 g, 4.50 mmol) was added Pd/C (241 mg, 0.277 mmol) and MeOH (30 mL). A H₂ balloon was applied and the reaction was stirred for 16h. The crude mixture was poured over a bed of Celite and rinsed with MeOH (75 mL) then concentrated. The dark brown solid was triturated with Et₂O (35 mL) to give the title compound E1(ii) as an off-white solid (1.02 g, 60%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.21 (t, *J*=7.5 Hz, 3 H), 1.40- 1.49 (m, 9 H), 1.52- 1.65 (m, 6 H), 2.60 - 2.87 (m, 2 H), 4.37 - 4.68 (m, 2 H), 8.01 - 8.55 (m, 1 H), 9.75 - 10.42 (m, 1 H), 11.60 - 12.54 (m, 1 H). Mass spectrum: Calcd for C₁₈H₂₆FN₆O₂ (M+H): 377. Found 377.

Intermediate E1(iii): 5-tert-butyl 1-ethyl 3-[(2-ethyl-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-1,5-dicarboxylate

A solution of E1(ii) (1.00 g, 2.66 mmol) in THF (40 mL) and DIPEA (858 mg, 6.64 mmol) was cooled in an ice bath. Ethyl chloroformate (317 mg, 2.92 mmol) was added and the reaction was warmed to room temperature and stirred for 16 h. The reaction was quenched with water (50 mL) and extracted with EtOAc (2 x 100 mL). The combined extracts were washed with brine (50 mL) then dried (MgSO₄), filtered and concentrated. Purification was performed by silica gel column chromatography using 0-40% EtOAc in CH₂Cl₂ to give the title compound E1(iii) as a yellow solid (879 mg, 74%). ¹H NMR (300 MHz, CDCl₃) δ ppm 1.28- 1.34 (m, 3 H), 1.47 (t, *J*=7.2 Hz, 3 H), 1.50- 1.56 (m, 9 H), 1.79- 1.90 (m, 6 H), 2.75- 2.87 (m, 2 H), 4.45- 4.56 (m, 2 H), 4.74- 4.77 (m, 2 H), 7.65 (s, 1 H), 8.06- 8.31 (m, 1 H). Mass spectrum: Calcd for C₂₁H₃₀FN₆O₄ (M+H): 449. Found 449.

Intermediate E1(iv): Ethyl 5-(chlorocarbonyl)-3-[(2-ethyl-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate

To a suspension of E1(iii) (879 mg, 1.96 mmol) in dioxane (5 mL) was added HCl (10 mL, 4M in dioxane). The reaction was stirred at room temperature for 2 h then concentrated. The di HCl salt was taken up in CH₂Cl₂ (40 mL) and DIPEA (1.27 g, 9.8 mmol) was added then the reaction was cooled to -78 °C and triphosgene (0.41 g, 1.37 mmol) in a solution of CH₂Cl₂ (10 mL) was added slowly with an addition funnel over 15 min. The reaction was quenched at -78°C with water then warmed to room temperature. The mixture was made to pH 8-9 with NaHCO₃ and extracted with CH₂Cl₂ (2 x 15 mL). The combined extracts were washed with brine (15 mL) then dried, filtered and concentrated. The crude material was purified by silica gel column chromatography using 0-3% 7N NH₃/MeOH in CH₂Cl₂ to give the title compound E1(iv) as a white solid (244 mg, 31%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.19- 1.30 (m, 3 H), 1.32- 1.42 (m, 3 H), 1.69 (s, 6 H), 2.67- 2.95 (m, 2 H), 4.47 (q, *J*=7.0 Hz, 2 H), 4.66- 5.51 (m, 2 H), 8.27- 8.76 (m, 1 H), 9.73- 10.36 (m, 1 H). Mass spectrum: Calcd for C₁₇H₂₁ClFN₆O₃ (M+H): 411. Found 411.

Example E1: N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[(2*S*,5*R*)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine acetate salt


To a sealed tube was added E1(iv) (200 mg, 0.487 mmol), DIPEA (283 mg, 2.19 mmol) and THF (10 mL). The tube was placed in a 90 °C oil bath and heated for 16h. The reaction was concentrated then taken up in MeOH (5 mL) and Et₃N (5 mL) and stirred at

room temperature for an additional 16h. Preparative HPLC was performed using 5-50% ACN (0.1% AcOH) to give the title compound E1 as the acetate salt (61 mg, 26%). See Table 1 below for NMR data.

EXAMPLES E2 and E3:

Examples E2 and E3 were prepared using methods analogous to Example E1 above. See Table 1 below for names and NMR data.

EXAMPLE E4: 5-{[(2S,5R)-2,5-dimethyl-4-(3,3,3-trifluoropropyl)piperazin-1-yl]carbonyl}-N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine



Intermediate E4(i): (2S,5R)-1-benzyl-2,5-dimethyl-4-(3,3,3-trifluoropropyl)piperazine

To a solution of (2S,5R)-1-benzyl-2,5-dimethylpiperazine (2.00 g, 7.21 mmol) in THF (10 mL), MeOH (10 mL), and TEA (1.0 mL) was added 3,3,3-trifluoropropionaldehyde (1.62 g, 14.4 mmol) and AcOH (0.826 mL, 14.4 mmol) followed by sodium cyanoborohydride (0.907 g, 14.4 mmol). The reaction was allowed to stir for 2 h, at which time the reaction was quenched with water (15 mL) then made basic with NaHCO₃ and extracted with EtOAc. The organic layer was washed with brine, dried and concentrated *in vacuo* to yield a white powder (2.0 g, 90%). Mass Spectrum: Calcd for C₁₆H₂₄N₂F₃ (M+H): 301. Found: 301.

Intermediate E4(ii): (2R,5S)-2,5-dimethyl-1-(3,3,3-trifluoropropyl)piperazine

To a solution of E4(i) (1.90 g, 6.30 mmol) in MeOH (20 mL) was added Pd(OH)₂ (1.00g, 7.10 mmol). The suspension was evacuated/backfilled with hydrogen (x 3) and allowed to stir for 15 h under hydrogen. The suspension was filtered through Celite, the filter

Example E4 was prepared using methods analogous to Example E1 above, except that intermediate E4(ii) was substituted in place of (2R,5S)-1,2,5-trimethylpiperazine hydrochloride. See Table 1 below for NMR data.

Side-chain E5(ii)

E5(i) $\xrightarrow[\text{K}_2\text{CO}_3, \text{CH}_3\text{CN}]{\text{BnBr}}$ **E5(ii)**


E5(iii) $\xrightarrow[\text{THF, H}_2\text{O}]{\text{Boc}_2\text{O, Na}_2\text{CO}_3}$ **E5(iv)** $\xrightarrow[\text{HATU, DMF}]{\text{E5(ii)}}$ **E5(v)** $\xrightarrow[\text{dioxane}]{\text{HCl}}$ **E5(vi)**

E5(vi) $\xrightarrow{\text{NaHCO}_3}$ **E5(vii)** $\xrightarrow[\text{THF}]{\text{NaH, CH}_3\text{I}}$ **E5(viii)** $\xrightarrow[\text{THF}]{\text{LiAlH}_4}$ **E5(x)**

E5(x) $\xrightarrow{\text{SFC}}$ **E5(x)-isomer A** $\xrightarrow[\text{MeOH}]{\text{H}_2, 10\% \text{ Pd-C}}$ **E5(xi)-Isomer A**

E5(x) $\xrightarrow{\text{SFC}}$ **E5(x)-isomer B** $\xrightarrow[\text{MeOH}]{\text{H}_2, 10\% \text{ Pd-C}}$ **E5(xi)-Isomer B**

Absolute stereochemistry not assigned



Side-chain E5(ii) ((S)-methyl 2-(benzylamino)propanoate):

To a mixture of compound E5(i) (27.7 g, 0.2 mol) in CH₃CN (350 mL) at 0° was added K₂CO₃ in portion, after which BnBr (34.2 g, 0.2 mol) was added dropwise over a period of 1 hour. After addition, the mixture was stirred at room temperature for 2 hours. TLC (petrol ether/ ethyl acetate=4/1) suggested that the reaction was complete. The reaction was quenched by water (500 mL), and extracted with ethyl acetate (400 mL×3). The combined organic layers were dried over Na₂SO₄. Filtration and concentration gave the crude, which was purified by column chromatography (eluted with petrol ether/ ethyl acetate=50/1 to 4/1) to give compound E5(ii) (15 g, 39%) as a colourless liquid. ¹H NMR CDCl₃ 1.2 (d, 3H), 1.9 (bs, 1H), 3.3 (q, 1H), 3.6 (d, 1H), 3.8 (d, 1H), 3.7 (s, 3H), 7.1-7.3 (m 5H)

Intermediate E5(iv) ((R)-2-(tert-butoxycarbonyl)-4-methoxy-4-oxobutanoic acid):

To a stirred solution of compound E5(iii) (73.5 g, 0.5 mol) in THF (500 mL) and water (500 mL) was added Na₂CO₃ (106 g, 1 mol), followed by Boc₂O (120 g, 0.55 mol). Then the mixture was stirred at room temperature overnight. TLC (dichloromethane/methanol=10/1) suggested that the reaction was complete. The reaction mixture was concentrated, and the residue was dissolved in water (200 mL), extracted with ethyl acetate (500 mL×2). The aqueous layer was separated, and acidified with 1N HCl to pH=5, extracted with ethyl acetate (500 mL×20). The combined organic layers were dried over Na₂SO₄. Filtration and concentration gave the compound E5(iv) (75 g, 61%) as an oil.

¹H NMR CDCl₃ 1.4 (s, 9H), 2.8 (dd, 1H), 3.0 (dd, 1H), 3.6 (s, 3H), 4.6 (m, 1H), 5.5 (d, 1H), 9.3 (br, 1H)

Intermediate E5(v) ((R)-methyl 4-(benzyl((R)-1-methoxy-1-oxopropan-2-yl)amino)-3-(tert-butoxycarbonyl)-4-oxobutanoate):

To a stirred solution of compound E5(iv) (117.5 g, 0.475 mol) in DMF (1.5 L) was added NMM (80.1 g, 0.792 mol), HATU (150.6 g, 0.396 mol) in turn. The mixture was stirred at 0° for 30 minutes, then compound 9 (76.5 g, 0.396 mol) was added dropwise. The

solution was stirred at room temperature overnight. TLC (petrol ether/ ethyl acetate=1/1) suggested that the reaction was complete. The reaction solution was poured into water (1 L), and extracted with ethyl acetate (1 L×3). The combined organic layers was washed with 1 N HCl (200 mL×2), aqueous NaHCO₃ (200 mL×2), brine (400 mL), dried over Na₂SO₄, and concentrated to give compound E5(v) (152 g, 91%) as an oil, which was used the next step without further purification.

Intermediate E5(vi) ((R)-2-((R)-2-amino-N-benzyl-4-methoxy-4-oxobutanamido)propanoic acid):]

To a stirred solution of compound E5(v) (152 g, 0.362 mol) in dioxane (200 mL) was added HCl (g) in dioxane (1 L). The solution was stirred at room temperature overnight. The solution was concentrated to give compound E5(vi) (100 g, 94%) as an oil, which was used the next step without further purification.

Intermediate E5(vii) (methyl 2-((2R,5S)-4-benzyl-5-methyl-3,6-dioxopiperazin-2-yl)acetate):

Compound E5(vi) (110 g, 0.307 mol) was dissolved in dichloromethane/water (1 L/500 mL). With stirring, aq. NaHCO₃ was added dropwise till pH=9, then the solution was stirred at room temperature for one hour. The dichloromethane layer was separated, and the aqueous layer was extracted with dichloromethane (300 mL×2). The combined organic layers were dried over Na₂SO₄. Filtration and evaporation gave compound E5(vii) (95 g, 99%) as an oil. ¹H NMR CDCl₃ 1.3-1.4 (d, 3H), 2.7-2.8 (m, 1H), 3.0-3.1 (m, 1H), 3.6 (s, 3H), 3.8 (m, 1H), 4.0 (d, 1H), 4.4 (m, 1H), 5.2 (d, 1H), 7.1-7.3 (m, 5H), 7.7 (br, 1H)

Intermediate E5(viii) (methyl 2-((2R,5S)-4-benzyl-1,5-dimethyl-3,6-dioxopiperazin-2-yl)acetate):

To a solution of compound E5(vii) (1.45 g, 5 mmol) in THF (30 mL) was added NaH (0.24 g, 5 mmol) in portion at 0°C, The mixture was stirred for 20 minutes, then CH₃I (0.85 g, 6mmol) in THF (10 ml) was added dropwise. Then the mixture was stirred at room temperature for 5 hours. TLC (petrol ether/ ethyl acetate=1/1) suggested that the reaction was complete. The reaction was quenched by water (20 mL), and extracted with ethyl acetate (30 mL×2). The combined organic layers were dried over Na₂SO₄. Filtration and concentration gave the crude, which was purified by column chromatography (eluted with petrol ether/ ethyl acetate=5/1) to give compound E5(viii)

- 56 -

(0.7 g, 46%) as an oil. ^1H NMR CDCl_3 1.4 (m, 3H), 2.9 (s, 3H), 3.0-3.2 (m, 1H), 3.5-3.7 (s, 3H), 3.8-4.1 (m, 2H), 4.2-4.4 (m, 1H), 5.1-5.3 (m, 1H), 7.1-7.3 (m 5H)

Intermediate E5(ix) (2-((5S)-4-benzyl-1,5-dimethylpiperazin-2-yl)ethanol):

To a solution of compound E5(ix) (12 g, 0.04 mmol) in THF (250 mL) was added LiAlH_4 (7.6 g, 0.2 mol) in portion at 0°C . After addition, the reaction mixture was heated to reflux for 36 hours. TLC (dichloromethane/methanol=10/1) suggested that the reaction was complete. The reaction was quenched by water (5 mL), and the mixture was filtered, the cake was washed with ethyl acetate several times. The filtrate was washed with Na_2SO_4 . Filtration and concentration gave the crude, which was purified by column chromatography (eluted with dichloromethane/methanol=100/1 to 10/1) to give racemic compound E5(ix) (8.3 g, 84%) as an oil.

Intermediate E5(x) – Isomers A and B: 2-((2R,5S)-4-benzyl-1,5-dimethylpiperazin-2-yl)ethanol and 2-((2S,5S)-4-benzyl-1,5-dimethylpiperazin-2-yl)ethanol

The racemic compound E5(ix) was separated by SFC using AS-H (0.46 x 2.5 cm x $5\mu\text{m}$) column with 5% MeOH (0.025% DEA) and 95% CO_2 as mobile phase to give compounds E5(x) - Isomer A and B (5.2 g and 1.9 g respectively). E5(x)-Isomer A: ^1H NMR CDCl_3 1.0 (d, 3H), 1.4 (m, 1H), 1.7 (m, 1H), 2.0 (m, 2H), 2.1 (m, 1H), 2.2 (s, 3H), 2.4 (m, 1H), 2.6 (dd, 1H), 2.7 (dd, 1H), 3.0 (d, 1H), 3.4 (m, 1H), 3.6 (m, 1H), 4.0 (d, 1H), 4.2 (br, 1H), 7.1-7.3 (m, 5H). E5(x)-Isomer B: ^1H NMR CDCl_3 1.0 (d, 3H), 1.6-1.8 (m, 2H), 2.2-2.7 (m, 9H), 3.2 (d, 1H), 3.4 (m, 1H), 3.6 (m, 1H), 3.8 (d, 1H), 4.8 (br, 1H), 7.1-7.3 (m, 5H).

Intermediates E5(xi) – Isomers A and B: 2-((2R,5S)-1,5-dimethylpiperazin-2-yl)ethanol and 2-((2S,5S)-1,5-dimethylpiperazin-2-yl)ethanol


The mixture of compound E5(x)-Isomer A (5.2 g, 0.021 mol) and Pd/C (0.5 g) in MeOH (40 mL) or the mixture of compound E5(x)-Isomer B (1.9 g, 7.7 mmol) and Pd/C (0.2 g) in MeOH (40 mL) under 50 psi of H_2 was stirred at room temperature overnight. TLC (dichloromethane/methanol=10/1) suggested that the reaction was complete. The reaction mixture was filtered and the filtrate was concentrated to give E5(xi)-Isomer A (3.1 g, 94%) as an offwhite solid or E5(xi)-Isomer B (1.2 g, 93%) as an offwhite solid. E5(xi)-Isomer A: ^1H NMR CDCl_3 1.0 (d, 3H) 1.8 (m, 1H), 2.0 (m, 1H), 2.4-3.1 (m, 9H), 3.8 (m, 2H). E5(xi)-Isomer B: ^1H NMR CDCl_3 1.0 (d, 3H), 1.5 (m, 1H), 1.8 (t, 1H), 2.0-2.3 (m, 2H), 2.4 (s, 3H), 2.7-3.0 (m, 4H), 3.7 (m, 1H), 3.9 (m, 1H).

Example E5: 2-((5S)-4-[[3-[(2-ethyl-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl]-1,5-dimethylpiperazin-2-yl)ethanol

- 57 -

The title compound was prepared using a method analogous to Example E1 above except that E5(xi)-Isomer A was substituted in place of ((2*R*,5*S*)-1,2,5-trimethylpiperazine. See Table 1 below for NMR data.

EXAMPLE F1: 5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-*N*-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine



Intermediate F1(i): Ethyl 3-[(ethoxycarbonyl)amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-*c*]pyrazole-1(4*H*)-carboxylate.

To a solution of 3-Ethoxycarbonylamino-6,6-dimethyl-4,6-dihydro-pyrrolo[3,4-*c*]pyrazole-1,5-dicarboxylic acid 5-tert-butyl ester 1-ethyl ester (5.69 g, 14.4 mmol) in CH₂Cl₂ (10 mL) was added 4M HCl in 1,4-dioxane (20 mL). The solution was stirred for 1 h, at which time the volatiles were removed *in vacuo* to yield the desired product F1(i) as a white solid (4.8 g, 91%). Mass Spectrum: Calcd for C₁₃H₂₁N₄O₄ (M+H): 297. Found: 297.

Intermediate F1(ii): Ethyl 3-[(ethoxycarbonyl)amino]-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-*c*]pyrazole-1(4*H*)-carboxylate.

To a pressure vessel was added F1(i) 4.80 g, 16.2 mmol), 4-fluoro-1-methylpiperidine-4-carbonyl chloride (2.45 g, 13.6 mmol), diisopropylethylamine (9.06 mL, 52.0 mmol), and THF (300 mL). The suspension was heated at 80 °C for 15 h. The volatiles were

removed *in vacuo* to yield the desired product F1(ii) as a brown foam (5.6 g, 98%).

Mass Spectrum: Calcd for $C_{20}H_{31}N_5O_5F$ (M+H): 440. Found: 440.

Intermediate F1(iii): 5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

To a microwave vial was added F1(ii) (9.80 g, 22.0 mmol) in a solution of MeOH (35 mL) and LiOH (2.14 g, 89.2 mmol). The reaction was heated at 110 °C in the microwave for 2 h. The reaction mixture was concentrated *in vacuo* to give the desired product F1(iii) as a tan foam (4.4 g, 67%). Mass Spectrum: Calcd for $C_{14}H_{23}N_5OF$ (M+H): 296. Found: 296.

Intermediate F1(iv): N-(2-chloro-5-fluoropyrimidin-4-yl)-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.


To a solution of F1(iii) (1.40 g, 4.74 mmol) and 2,4-dichloro-5-fluoro pyrimidine (0.791 g, 4.74 mmol) in DMSO (10 mL) was added potassium dihydrogenphosphate (0.645 g, 4.74 mmol) followed by H_3PO_4 (0.0929 g, 0.948 mmol). The reaction heat to 95 °C for 15 h. The crude reaction mixture was cooled to 22 °C then poured into ice cold $NaHCO_3$ (sat., aq.) (100 mL). The aqueous layer was extracted with EtOAc (2 x 50 mL) and the combined organic layers washed with brine (2 x 50 mL), dried over magnesium sulfate, and concentrated to give the desired product F1(iv) as a brown solid (0.633 g, 31%). Mass Spectrum: Calcd for $C_{18}H_{23}N_7OF_2Cl$ (M+H): 426. Found: 426.

Example F1: 5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

To a pressure vessel was added F1(iv) (0.225 g, 0.528 mmol), trimethylboroxine (1.47 mL, 10.6 mmol), cesium carbonate (3.44 g, 10.6 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane adduct (0.0647 g, 0.0792 mmol), H_2O (0.8 mL), and THF (8.0 mL). The suspension was purged with argon for 2 min, then heat at 100 °C for 15 h. The cooled solution was filtered to remove undissolved solids, concentrated, and redissolved in MeOH. Preparative HPLC using 20-60% ACN/ H_2O (0.1% AcOH) provided the desired product F1 as a white solid (0.020 g, 9.3%). See Table 1 below for NMR data.

EXAMPLE G1: N-(2-ethyl-5-fluoropyrimidin-4-yl)-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

- 59 -



Intermediate G1(i): 5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-N-(5-fluoro-2-vinylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.


To a pressure vessel was added starting material [3-(2-Chloro-5-fluoro-pyrimidin-4-ylamino)-6,6-dimethyl-4,6-dihydro-1H-pyrrolo[3,4-c]pyrazol-5-yl]-(4-fluoro-1-methylpiperidin-4-yl)-methanone (0.375 g, 0.881 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (0.407 g, 2.64 mmol), Na₂CO₃ (0.280 g, 2.64 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane adduct (0.144 g, 0.176 mmol) followed by DME (10 mL) and water (2 mL). The reaction mixture was purged with argon for 2 min, then heat to 100 °C for 15 h. The volatiles were concentrated *in vacuo* and the residue redissolved in EtOAc (15 mL). The organic layer was washed with water (10 mL), dried over MgSO₄, filtered, and concentrated to give the desired product **G1(i)** as a brown foam (0.35 g, 95%). Mass Spectrum: Calcd for C₂₀H₂₆N₇OF₂ (M+H): 418. Found: 418.

Example G1: N-(2-ethyl-5-fluoropyrimidin-4-yl)-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

To a solution of starting material (0.413 g, 0.989 mmol) in MeOH (10 mL) was added 10% palladium on carbon (0.0400 g, 0.380 mmol). The suspension was evacuated/backfilled with hydrogen for 3 times and allowed to stir for 15 h under hydrogen. The suspension was filtered through Celite, the filter cake washed with CH₂Cl₂, and the filtrate concentrated. Preparative HPLC provided the desired product **G1** as a white solid (0.018 g, 4.4%). See Table 1 below for NMR data.

- 60 -

EXAMPLE H1: *N*-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-5-[(3*S*,8*aS*)-3-methylhexahydropyrrolo[1,2-*a*]pyrazin-2(1*H*)-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine



Intermediate H1(i): tert-butyl 3-[(5-fluoro-2-methylpyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5-carboxylate.

To a pressure vessel was added 3-(2-Chloro-5-fluoro-pyrimidin-4-ylamino)-6,6-dimethyl-4,6-dihydro-1H-pyrrolo[3,4-c]pyrazole-5-carboxylic acid tert-butyl ester (1.50 g, 3.90 mmol), trimethylboroxine (10.9 mL, 78.4 mmol), cesium carbonate (25.5 g, 78.4 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane adduct (0.480 g, 0.588 mmol), H₂O (6 mL), and THF (60 mL). The suspension was purged with argon for 2 min, then heat at 100 °C for 15 h. The cooled solution was filtered to remove undissolved solids. Purification by silica gel column chromatography using 0-10% ammoniated methanol in CH₂Cl₂ provided the desired product **H1(i)** as a brown solid (0.80 g, 56%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.41 (s, 9 H) 1.57 (s, 6 H) 2.38 (s, 3 H) 4.43- 4.58 (m, 2 H) 8.12 (s, 1 H) 10.02 (s, 1 H) 12.34 (s, 1 H). Mass Spectrum: Calcd for C₁₇H₂₄N₆O₂F (M+H): 363. Found: 363.

Intermediate H1(ii): 5-tert-butyl 1-ethyl 3-[(5-fluoro-2-methylpyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-1,5-dicarboxylate.

To a 0 °C solution of H1(i) (0.800 g, 2.21 mmol) in THF (10 mL) was added NaH (60% dispersion in mineral oil, 0.124g, 3.09 mmol). The solution was stirred at 0 °C for 10 min, at which time ethylchloroformate (0.421 mL, 4.41 mmol) was added. The solution was warmed to 22 °C and stirred for 15 h. The reaction was quenched with NH₄Cl (sat., aq.) (10 mL) and the aqueous layer extracted with EtOAc (3 x 30 mL). The combined organic layers were washed with brine (1 x 20 mL), dried over MgSO₄ and concentrated to give the desired product H1(ii) as a brown solid (0.92 g, 96%). Mass Spectrum: Calcd for C₂₀H₂₈N₆O₄F (M+H): 435. Found: 435.

Intermediate H1(iii): Ethyl 5-(chlorocarbonyl)-3-[(5-fluoro-2-methylpyrimidin-4-yl)amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate.

To a solution of H1(ii) (0.919 g, 2.12 mmol) in CH₂Cl₂ (5 mL) was added 4M HCl in dioxane (10 mL). The solution was stirred for 1 h, concentrated *in vacuo*, and redissolved in CH₂Cl₂ (40 mL) and diisopropylethylamine (1.67 mL, 9.58 mmol). The solution was cooled to -78 °C and triphosgene (0.569 g, 1.92 mmol) in CH₂Cl₂ (10 mL) was added dropwise over 30 min. The reaction was quenched with H₂O (10 mL), warmed to 22 °C, and made pH 8-9 with NaHCO₃. The aqueous layer was extracted with CH₂Cl₂ (2x 30 mL) and the combined organic layers washed with brine (1 x 30 mL), dried over MgSO₄, and concentrated to give the desired product H1(iii) as a brown solid (0.49 g, 46%). Mass Spectrum: Calcd for C₁₆H₁₉ClN₆O₄F (M+H): 397. Found: 397.


Example H1: N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-5-[(3*S*,8*aS*)-3-methylhexahydropyrrolo[1,2-*a*]pyrazin-2(1*H*)-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine.

To a pressure vessel was added H1(iii) (0.244 g, 0.615 mmol), (3*S*,8*aS*)-3-methyloctahydropyrrolo[1,2-*a*]pyrazine (0.0862 g, 0.615 mmol), diisopropylethylamine (0.428 mL, 2.46 mmol), and THF (5 mL). The suspension was stirred at 90 °C for 2 h. Volatiles were removed *in vacuo* and the residue redissolved in MeOH (10 mL). Triethylamine (5 mL) was added and the solution stirred for 15 h. Preparative HPLC using 20-60% ACN/H₂O (0.1% AcOH) provided the desired product H1 as a white solid (0.12 g, 38%).

EXAMPLES H2 – H7:

Examples H2 through H7 were prepared using methods analogous to Example H1 above. See Table 1 below for names and NMR data.

EXAMPLE H8: 4-[(2*R*,5*S*)-4-{[3-[(5-fluoro-2-methylpyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl}-2,5-dimethylpiperazin-1-yl)methyl]tetrahydro-2*H*-pyran-4-ol



Intermediate H8(i): 1,6-dioxaspiro[2.5]octane

A solution of 4-methylenetetrahydro-2*H*-pyran (1.00 g, 10.2 mmol) in CH₂Cl₂ (30 mL) was placed in an ice bath then *meta*-chloroperoxybenzoic acid (2.46 g, 14.3 mmol) was added in three portions. The reaction was slowly warmed to RT and stirred for 3h then quenched with 10% NaOH(aq) (10 mL) and extracted with CH₂Cl₂ (2 x 15 mL). The combined extracts were dried (MgSO₄), filtered and concentrated to provide intermediate H8(i) as a clear oil (607 mg, 52%).

Intermediate H8(ii) (4-[(2*S*,5*R*)-4-benzyl-2,5-dimethylpiperazin-1-yl)methyl]-tetrahydro-2*H*-pyran-4-ol):

To a microwave vial was added H8(i) (259 mg, 2.3 mmol) and (2*S*,5*R*)-1-benzyl-2,5-dimethylpiperazine (464 mg, 2.3 mmol) and 5 mL of MeOH. The vial was heated to 150°C for 2h in the microwave. The crude reaction was concentrated to provide intermediate H8(ii) (723 mg, 100%)

Intermediate H8(iii) 4-[(2*R*,5*S*)-2,5-dimethylpiperazin-1-yl)methyl]tetrahydro-2*H*-pyran-4-ol

To a solution of H8(ii) (723 mg, 2.3 mmol) in MeOH (15 mL) was added Pd/C (72 mg, 0.07 mmol). The reaction was subject evacuation-backfill (3x) with H₂ gas

- 63 -

then run under an H₂ atmosphere overnight. The completed reaction mixture was filtered through a bed of Celite, rinsed with CH₂Cl₂ and MeOH then concentrated to give the title compound (500 mg, 97%) as a yellow-orange semi solid H8(iii).


Example H8: 4-(((2*R*,5*S*)-4-([3-[(5-fluoro-2-methylpyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl)-2,5-dimethylpiperazin-1-yl)methyl]tetrahydro-2*H*-pyran-4-ol.

The title compound was prepared using methods analogous to Example H1 above. See Table 1 below for NMR data.

EXAMPLES H9 –H10:

Examples H9 and H10 were prepared using methods analogous to Example H1 above, except E5(xi)-Isomers A and B, respectively, were substituted in place of (3*S*,8*aS*)-3-methyloctahydropyrrolo[1,2-*a*]pyrazine. See Table 1 below for names and NMR data.

EXAMPLE I1: N-(5-fluoro-2-propylpyrimidin-4-yl)-6,6-dimethyl-5-[[[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine.



To a pressure vessel was added [3-(2-Chloro-5-fluoro-pyrimidin-4-ylamino)-6,6-dimethyl-4,6-dihydro-1*H*-pyrrolo[3,4-*c*]pyrazol-5-yl]-((*S*)-2,4,5,5-tetramethyl-piperazin-1-yl)-


methanone (0.150 g, 0.330 mmol), 2-allyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.12 g, 6.66 mmol), cesium carbonate (0.543 g, 1.66 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane adduct (0.0408 g, 0.0498 mmol), H₂O (0.3 mL), and THF (3 mL). The suspension was purged with argon for 2 min, then stirred at 100 °C for 15 h. The cooled solution was filtered to remove undissolved solids and the volatiles removed *in vacuo*. The residue was redissolved in MeOH (5 mL) and 10% palladium on carbon (0.0400 g, 0.380 mmol) was added. The

suspension was evacuated and backfilled with hydrogen for 3 times and allowed to stir for 15 h under hydrogen. The suspension was filtered through Celite, the filter cake washed with CH_2Cl_2 , and the filtrate concentrated. Preparative HPLC provided the desired product **11** as a white solid (0.025 g, 14%).

EXAMPLE 12:

Example **12** was prepared using methods analogous to Example **11** above. See Table 1 below for name and NMR data.

EXAMPLES J1-J8:



Intermediate J(i): tert-butyl 3-amino-6,6-dimethyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carboxylate.

To a suspension of tert-butyl 3-amino-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carboxylate (8.5 g, 33.9 mmol) and diisopropyl ethylamine (18 mL, 3.0 equiv) in

dichloromethene (200 mL) was added 2-(trimethylsilyl)ethoxymethyl chloride (6.0 mL, 1.0 equiv) dropwise at 0 °C under nitrogen. The mixture was been stirred at 0 °C under nitrogen for two hours then warmed to room temperature and stirred over night. The reaction mixture was concentrated and purified by column chromatography to give the title compound J(i) as a white solid (2.27 g, 18%). ¹H NMR (400 MHz, METHANOL-*d*₄): δ ppm 0.81- 0.97 (m, 2 H) 1.45- 1.59 (m, 9 H) 1.72 (d, *J*=5.29 Hz, 13 H) 3.53- 3.67 (m, 2 H) 4.26 (d, *J*=7.55 Hz, 2 H) 5.17 (s, 2 H). 2D-NOESY NMR showed proton at 5.17 ppm is correlated with proton at 1.72ppm.

Intermediate J(ii): *tert*-butyl 3-[(3-fluoro-6-methylpyridin-2-yl)amino]-6,6-dimethyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-4,6-dihydropyrrolo[3,4-*c*]pyrazole-5(1*H*)-carboxylate.

To a flask equipped with a condenser was added J(i) (4.7g, 12.3 mmol), Cesium Carbonate (8.0g, 25mmol), 2-bromo-3-fluoro-picoline (2.6 g, 14mmol), Pd₂(dba)₃ (600mg, 0.6 mmol, 10 mol% Pd), Xantphos (700mg, 1.2 mmol, 10 mol%) and 1,4 dioxane (150 mL). The reaction mixture was stirred for 15 min under nitrogen atmosphere. While the mixture was being stirred, degassed water (50) mL was added. The mixture was heated at 100°C for 16 hours under nitrogen atmosphere and then evaporated 4/5 volumn solvent. Water was introduced and the mixture extracted with ethyl acetate (3x100 mL). The combined organics were washed with water and saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered, concentrated and purified by column chromatography to give the title compound J(ii) as a white solid (2.9 g, 48%).

Intermediate J(iii): *tert*-butyl 3-[(3-fluoro-6-methylpyridin-2-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazole-5(1*H*)-carboxylate.

To a solution of J(ii) (2.9 g, 5.9mmol) in THF (100 mL) was added TBAF (7.7g, 29.5 mmol, 5 equiv). The reaction mixture was heated at 70 °C for 16 hours and then evaporated 4/5 volumn solvent. Water was introduced and the mixture extracted with ethyl acetate twice. The combined organics were washed with water and saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered, concentrated and purified by column chromatography to give the title compound J(iii) as a white solid (1.6 g, 76%). LCMS (API-ES, M+H⁺): 362.

Intermediate J(iv): Ethyl acetate- *tert*-butyl 3-[(3-fluoro-6-methylpyridin-2-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazole-5(1*H*)-carboxylate.

A solution of carboethoxy chloride (0.58g, 5.35 mmol, 1.2 equiv.) in THF (5 mL) was added slowly to a suspension mixture of J(iii) (1.6g, 4.46mmol) and DIPEA (3.9 mL, 5

- 66 -

equiv) in THF (80 mL) at 0 °C under nitrogen. The reaction was kept at 0 ° for two hours and then warmed to room temperature and stirred over night. The reaction mixture was concentrated and purified by column chromatography to give the title compound J(iv) as a white solid (627 mg, 35%).


Intermediate J(v): Ethyl acetate- N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

To a clear solution of J(iv) (627 mg, 1.45 mmol) in Dioxane (10 mL), 4M HCl in dioxane (10.8 mL, 30equiv) was added. The reaction mixture was stirred at room temperature for 6 hours. Evaporated solvent and volatile, dried in vacuum to give the title compound J(v) as HCl salt.

Intermediate J(vi) (applicable to Examples J3-J6): Ethyl acetate- 3-[(3-fluoro-6-methylpyridin-2-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carbonyl chloride.

To a clear solution of triphosgene (472 mg, 1.59 mmol) in DCM, DIPEA (1.26 mL, 7.23 mmol) was added dropwise (white smoke) under N₂ at 0 °C. The suspension of the crude product J(v) (489 mg, 1.27 mmol) in DCM (20 mL) was added dropwise. The reaction mixture was warmed to room temperature slowly and stirred at room temperature under nitrogen over night. Evaporated solvent and volatile, dried in vacuum to give the title compound J(vi) as HCl salt. See Table 1 below for NMR data.

EXAMPLE J1: 5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine




A solution of J(v) (68 mg, 0.17 mmol), 4-fluoro-1-methylpiperidine-4-carbonyl chloride (74.2 mg, 2 equiv), and DIPEA (0.18mL) in THF (10mL) was heated to 70°C for 16 hrs. THF was concentrated. The reaction mixture was dissolved in CH₃OH (10mL) and Et₃N (5mL) then stirred at room temperature for 16 hrs. The residue was purified by HPLC (0.1% HOAc as a buffer) to give the title compound J1 as a solid (80 mg, 20 % yield). See Table 1 below for NMR data.

EXAMPLES J2-J8:

Examples J2 through J8 were prepared using methods analogous to Example J1 above, with Examples J3 through J8 being further derived from Intermediate J(vi) as described in the schematic above. See Table 1 below for names and NMR data.

EXAMPLE J9: N-[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]-6,6-dimethyl-5-[[[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine




Example J9 was prepared using methods analogous to Examples J1 through J8, except substituting A1(iv) in place of J(i) as shown above. See Table 1 below for NMR data.

EXAMPLES J10-J16:

Examples J10 through J16 were prepared using methods analogous to Example J9 above. See Table 1 below for names and NMR data.

EXAMPLE K1: N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[[(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine



Intermediate K1(i): tert-butyl 3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carboxylate.

A fresh solution of sodium ethoxide was prepared by dissolving 7g of sodium metal in 140ml of anhydrous ethanol. Twelve 20ml Biotage microwave vials, equipped with magnetic stir bars, were charged with 1.2g of the chloride (total 14.4g, 38mmol). To each vial was added 10ml of the aforementioned sodium ethoxide solution followed by 5ml of anhydrous ethanol. The vials were sealed, and processed in a batchwise fashion using a Biotage Initiator microwave. Each reaction was heated to 160°C for 10 minutes. After being cooled, the reaction vials were opened (the reaction was shown to be complete by LC-MS), and the contents combined. Each vial was rinsed with 5ml of ethanol, and 10ml of water (a solid precipitate deposits at the bottom of the vial), and the washings combined with the original contents of the vial. The ethanol was removed *in vacuo*, and the aqueous extracted with ethyl acetate (3 x 150ml). The organic extracts were dried over sodium sulfate, and the solvent removed *in vacuo* to afford the product as a slightly yellow solid (14.1g, 96%). This was sufficiently pure to be used directly in the next step.

Note: Thermal heating to reflux for an extended period of time (18 hours) did not effect the transformation. Use of a sealed tube was not attempted. Microwave heating at 140°C for 10 minutes lead to a 60% conversion by LC-MS.

Intermediate K1(ii): 5-tert-butyl 1-ethyl 3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-1,5-dicarboxylate.

To a solution of K1(i) (175 mg, 0.44 mmol) in methylene chloride was added triethyl amine (0.07 mL, 1.2 eq) followed by ethyl chloroformate (0.04 mL, 1.0 eq). After stirring at room temperature for 12 hours, the solvent was removed *in vacuo* and the residue was partitioned between methylene chloride and water. The organic layer was separated then concentrated to a yellow solid K1(ii) which was carried on without further purification.

Intermediate K1(iii): Ethyl 5-(chlorocarbonyl)-3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate.

To a solution of K1(ii) (170 mg, 0.36 mmol) in 6 mL of dry dioxane was added 1 mL of 4M HCl in dioxane. After stirring at room temperature for 48 hours, the solvent was removed *in vacuo* and the white solid residue was dried for several hours under high vacuum and carried on without further purification. The reaction with triphosgene was

carried out using the same procedure as in the preparation of intermediate E1(iv) ¹H NMR (400 MHz, CHLOROFORM-D) δ ppm 1.38- 1.43 (3 H, m), 1.49- 1.54 (3 H, m), 1.78- 1.86 (6 H, m), 4.32- 4.42 (2 H, m), 4.61 (2 H, qd, $J=7.13$, 1.26 Hz), 5.17 (2 H, d, $J=59.42$ Hz), 8.08 (1 H, t, $J=2.39$ Hz), 10.27 (1 H, d, $J=9.57$ Hz). LCMS (M+H)⁺ 427.1

Example K1: N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-([(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

To a solution of K1 (iii) (70 mg, 0.16 mmol) in tetrahydrofuran (4 mL) was added (2R,5S)-1,2,5-trimethylpiperazine hydrochloride (108 mg, 4 equiv) and diisopropylethyl amine (0.15 mL, 6.8 equiv). The resulting mixture was heated in a sealed tube at 80 °C for 16 hours and then allowed to cool to ambient temperature. To the crude reaction mixture was added one sodium hydroxide pellet dissolved in 2 mL methanol and stirring was continued at room temperature for one hour. The solvent was removed *in vacuo* and the residue was purified by reverse-phase HPLC using a gradient of 15% to 35% acetonitrile in water (0.1% acetic acid modifier) ramped over 25 minutes. Overnight lyophilization afforded the title compound K1 as a white powder (37 mg, 44%). See Table 1 below for NMR data.

EXAMPLES K2 – K19:


Examples K2 through K19 were prepared using methods analogous to Example K1 above. See Table 1 below for names and NMR data.

EXAMPLE K20: N-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-([(3S,8aS)-3-isopropylhexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl]carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine

Example K20 was prepared using methods analogous to Example K1 above, except that E4(ii) was substituted in place of (2R, 5S)-1,2,5-trimethylpiperazine hydrochloride. See Table 1 below for NMR data.

EXAMPLE K21: 5-([(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]carbonyl)-N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine

- 70 -



Intermediate K21(i): (2S,5R)-1-benzyl-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazine.

To a mixture of (2S,5R)-1-benzyl-2,5-dimethylpiperazine (11.0 g, 53.8 mmol) and of tetrahydro-4H-pyran-4-one (5.39 g, 53.8 mmol) at 0°C was added Ti(IV) isopropoxide (19.2 mL, 67.3 mmol). The resulting mixture was stirred at rt overnight and an orange suspension was obtained. To the suspension was added EtOH (25 mL) and NaCNBH₃ (4.27 g, 64.6 mmol, 1.20 eq). The resulting mixture was stirred at rt for 24 h. The reaction mixture was then quenched with water (5.0 mL) and a yellow solid was generated. The suspension was diluted with EtOAc (400 mL) and filtered. The filtrate was concentrated under reduced pressure. The residue was again diluted with EtOAc (500 mL), dried over Na₂SO₄, and filtered. The filtrate was collected and concentrated under reduced pressure to afford a yellow oil. The oil was diluted in THF (400 mL) and PS-isocyanate was added to remove secondary amine. The suspension was filtered and the filtrate concentrated. The residue was purified by column chromatography using 50:5:1 CHCl₃/MeOH/Et₃N to afford the desired compound as yellow oil (13.0g, 84%)

Intermediate K21(ii): (2R,5S)-2,5-dimethyl-1-(tetrahydro-2H-pyran-4-yl)piperazine. (2S,5R)-1-benzyl-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazine (13.0 g, 45.1 mmol) was dissolved in MeOH (500 mL).

The solution was added into a Shaker container, which was previously charged with wet 10 % Pd/C. Hydrogenation was carried out in shaker under 40 psi Hydrogen pressure. The reaction was complete after 12 h. The reaction mixture was filtered through celite and the filtrate was concentrated to afford the desired compound (9.0g, 100%). ¹H NMR (300 MHz, CHLOROFORM-d) δ ppm 1.02 (d, J=3.58 Hz, 3 H), 1.04 (d, J=3.39 Hz, 3 H), 1.44 - 1.54 (m, 2 H), 1.58 (dd, J=8.57, 3.67 Hz, 2 H), 1.78 - 1.98 (m, 2 H), 2.44 - 2.67 (m, 2 H), 2.73 - 2.85 (m, 2 H), 2.85 - 2.94 (m, 1 H), 3.00 - 3.15 (m, 1 H), 3.25 - 3.37 (m, 1 H), 3.37 - 3.49 (m, 1 H), 4.03 (dd, J=6.97, 5.46 Hz, 2 H).

Example K21: 5-[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-yl)piperazin-1-yl]carbonyl)-*N*-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine

The title compound was prepared using methods analogous to Example K1 above. See Table 1 below for NMR data.

EXAMPLE K22: 4-[(2*R*,5*S*)-4-{[3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl}-2,5-dimethylpiperazin-1-yl)methyl]tetrahydro-2*H*-pyran-4-ol


Example K22 was prepared using methods analogous to Example K1 above, except H8(iii) was substituted in place of (2*R*, 5*S*)-1,2,5-trimethylpiperazine. See Table 1 below for NMR data.

EXAMPLES K23-K26 :

Examples K23 and K24 were prepared using the method described in Example K1 above, except that E5(xi)-Isomers A and B, respectively, were substituted in place of (2*R*,5*S*)-1,2,5-trimethylpiperazine. Examples K25 and K26 were also prepared using methods analogous to Example K1 above, except that E5(xi)-Isomers A and B, respectively, were substituted in place of (2*R*,5*S*)-1,2,5-trimethylpiperazine and sodium methoxide was substituted in place of sodium ethoxide. See Table 1 below for names and NMR data.

EXAMPLE L1: *N*-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine

- 72 -




To a solution of ethyl 3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate dihydrochloride (120 mg, 0.27 mmol) in 15 mL THF containing diisopropylethyl amine (0.23 mL, 4 eq) was added a 5 mL THF suspension of 4-fluoro-1-methylpiperidine-4-carbonyl chloride (128 mg, 2.6 eq). After heating for 5 hours at 65°C, the reaction was concentrated to dryness and carried on without further purification. To a solution of ethyl 3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-c]pyrazole-1(4H)-carboxylate in methanol was added 2 mL of a 10% sodium hydroxide solution in methanol. After removing the solvent *in vacuo*, the residue was purified as in Example K1 above. See Table 1 below for NMR data.

EXAMPLES L2 – L4:

Examples L2 through L4 were prepared using methods analogous to Example L1 above. See Table 1 below for names and NMR data.

EXAMPLE L5: N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[1-(3,3,3-trifluoropropyl)piperidin-4-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine



Intermediate L5(i): ethyl 1-(3,3,3-trifluoropropyl)piperidine-4-carboxylate.

- 73 -

To a solution of ethyl isonipecotate (3.00 g, 19.1 mmol) in THF (20 mL) and MeOH (20 mL) was added 3,3,3-trifluoropropionaldehyde (2.14 g, 19.1 mmol) and AcOH (1.09 mL, 19.1 mmol) followed by sodium cyanoborohydride (1.20 g, 19.1 mmol). The reaction was allowed to stir for 10 min., at which time the reaction was quenched with water (15 mL) then made basic with NaHCO₃ and extracted with EtOAc. The organic layer was washed with brine, dried and concentrated *in vacuo* to yield a white powder (3.3 g, 68%). Mass Spectrum: Calcd for C₁₁H₁₉NF₃O₂ (M+H): 254. Found: 254.

Intermediate L5(ii): 1-(3,3,3-trifluoropropyl)piperidine-4-carboxylic acid.

To a solution of ethyl 1-(3,3,3-trifluoropropyl)piperidine-4-carboxylate (3.30 g, 13.0 mmol) in THF (40 mL) and MeOH (40 mL) was added LiOH-H₂O (1.64 g, 39.1 mmol).

The reaction was allowed to stir for 15 h at which time the solution was filtered to remove undissolved solids and the filtrate concentrated. The resulting material was diluted with water (10 mL), made to pH 5 with HCl (conc), then concentrated *in vacuo* to yield a white solid (2.9 g, 150%). Mass Spectrum: Calcd for C₉H₁₅NF₃O₂ (M+H): 226. Found: 226.


Intermediate L5(iii): 1-(3,3,3-trifluoropropyl)piperidine-4-carbonyl chloride.

A suspension of 1-(3,3,3-trifluoropropyl)piperidine-4-carboxylic acid (4.40 g, 20.0 mmol) in thionyl chloride (20 mL) was heat to 80°C for 15 h. The solution was cooled and concentrated *in vacuo* to yield a brown foam (3.0 g, 68%).

Example L5: N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[1-(3,3,3-trifluoropropyl)piperidin-4-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine
The title compound L5 was prepared using methods analogous for the preparation of example L1 above, except 4-fluoro-1-methylpiperidine-4-carbonyl chloride was substituted with intermediate L5(iii). See Table 1 below for NMR data.

EXAMPLE L6: N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

Intermediate L6(i): (1-(tetrahydro-2H-pyran-4-yl)piperidine-4-carbonyl chloride




Intermediate L6(i) was prepared using methods analogous to intermediate L5 above, except tetrahydro-2H-pyran-4-carbaldehyde was substituted in place of 3,3,3-trifluoropropionaldehyde.

Example L6: N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine

The title compound L6 was prepared using methods analogous to L1 above, except 4-fluoro-1-methylpiperidine-4-carbonyl chloride was substituted with intermediate L6(i).

See Table 1 below for NMR data.

EXAMPLE M1: N-(4-ethoxypyrimidin-2-yl)-6,6-dimethyl-5-[[1-(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine



Intermediate M1(i): 3-[(4-ethoxypyrimidin-2-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1H)-carbonyl chloride.

To a -78°C solution of N-(4-ethoxypyrimidin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine hydrochloride (0.300 g, 1.09 mmol) in CH_2Cl_2 (15 mL) and diisopropylethylamine (0.952 mL, 5.47 mmol) was added triphosgene (0.227 g, 0.766 mmol) in CH_2Cl_2 (5 mL) dropwise over 10 min. The reaction was quenched with H_2O (10 mL), warmed to 22°C , and made pH 8-9 with NaHCO_3 . The aqueous layer was extracted with CH_2Cl_2 (2x 30 mL) and the combined organic layers washed with brine (1 x 30 mL), dried over MgSO_4 , and concentrated to give the desired product M1(i) as a brown solid (0.35 g, 95%). Mass Spectrum: Calcd for $\text{C}_{14}\text{H}_{18}\text{ClN}_6\text{O}_2$ (M+H): 337. Found: 337.

Example M1: N-(4-ethoxypyrimidin-2-yl)-6,6-dimethyl-5-[[1-(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.

To a solution of M1(i) (0.350 g, 1.04 mmol) in THF (4mL) was added (2R,5S)-1,2,5-trimethylpiperazine hydrochloride (0.133 g, 1.04 mmol) and DIPEA (0.905 mL, 5.20 mmol). Solution was heat to 90°C in a sealed tube overnight. The cooled solution was

filtered to remove undissolved solids, concentrated, and redissolved in DMSO.


Preparative HPLC provided the desired product **M1** as a white solid (0.030 g, 6.7%).

See Table 1 below for NMR data.

EXAMPLE M2: N-(4-ethoxypyrimidin-2-yl)-5-[[[(2S,5R)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine

Example **M2** was prepared using methods analogous to Example **M1** above. See Table 1 below for NMR data.

EXAMPLE N1: 5-[[[(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl]carbonyl]-N-[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine.



Intermediate N1(i): *tert*-butyl 3-[[[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1*H*)-carboxylate.

To a solution of *tert*-butyl 3-amino-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazole-5(1*H*)-carboxylate (3.00 g, 11.6 mmol) and 4-chloro-5-fluoro-2-(methoxymethyl)pyrimidine (2.06 g, 11.6 mmol) in DMSO (8 mL) was added potassium dihydrogenphosphate (1.58 g, 2.3 mmol) followed by H_3PO_4 (0.29 g, 2.3 mmol). The reaction was placed in a 90 °C oil bath and heated for 20h. The crude reaction was cooled to room temperature then poured into ice cold NaHCO_3 (60 mL) and extracted with EtOAc (2 x 30 mL). The combined organic extracts were washed with brine then dried (MgSO_4), filtered and concentrated. The crude solid was triturated with EtOAc to provide the title compound **N1(i)** as a white solid (3.2 g, 70%). ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ ppm 1.44 (s, 9 H), 1.59

- 76 -

(s, 3 H), 1.61 (s, 3 H), 3.34 (s, 3 H), 4.29 - 4.44 (m, 2 H), 4.45 - 4.64 (m, 2 H), 8.32 (d, $J=3.20$ Hz, 1 H), 10.25 (br. s., 1 H), 12.24 (br. s., 1 H). Mass Spectrum: Calcd for $C_{18}H_{25}FN_6O_3$ (M+H): 393. Found: 393.

Intermediate N1(ii): 5-*tert*-butyl 1-ethyl 3-[[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazole-1,5-dicarboxylate.

A solution of N1(i) (2.90 g, 13.1 mmol) in THF (60 mL) and DIPEA (2.39 g, 18.0 mmol) was cooled in an ice bath then ethylchloroformate (0.89 g, 8.1 mmol) was added dropwise. The reaction mixture was slowly warmed to room temperature and stirred for 5h, then quenched with water (50 mL) and extracted with EtOAc (2x 100 mL). The combined extracts were washed with brine (50 mL) then dried ($MgSO_4$), filtered and concentrated to provide the title compound N1(ii) as a yellow foam (3.3 g, 96%). Mass Spectrum: Calcd for $C_{21}H_{29}FN_6O_5$ (M+H): 465. Found: 465.

Intermediate N1(iii): 3-[[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazole-5(1*H*)-carbonyl chloride.

To a solution of N1(ii) (3.40 g, 7.3 mmol) in dioxane (10 mL) was added HCl (20 mL, 4M in dioxane). The reaction was stirred at room temperature for 3h then concentrated and dried under vacuum. The HCl salt of ethyl 3-[[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]amino]-6,6-dimethyl-5,6-dihydropyrrolo[3,4-*c*]pyrazole-1(4*H*)-carboxylate was taken up in CH_2Cl_2 (150 mL). DIPEA (4.26 g, 5.1 mmol) was added and the reaction mixture was cooled in a dry ice/acetone bath. Triphosgene (1.52 g, 32.9 mmol) was added drop wise in a solution of CH_2Cl_2 (50 mL). The cold reaction was quenched with water (100 mL) and warmed to RT then extracted with CH_2Cl_2 (2x 100 mL). The combined organic extracts were washed with brine, dried ($MgSO_4$), filtered and concentrated. The crude material was triturated with ether to give the title compound N1(iii) as a white solid (1.55 g, 50%). Mass spectrum: Calcd for $C_{17}H_{20}FCIN_6O_4$ (M+H): 427. Found 427.

Example N1: 5-[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazin-1-yl]carbonyl]-*N*-[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine.

To a sealed tube was added N1(iii) (420 mg, 0.98 mmol) in a solution of THF (10 mL) followed by DIPEA (445 mg, 3.4 mmol) and (2*R*,5*S*)-2,5-dimethyl-1-(tetrahydro-2*H*-

- 77 -

pyran-4-ylmethyl)piperazine (209 mg, 0.98 mmol). The reaction was placed in an 85 °C oil bath and heated for 16 h. The crude reaction was concentrated then taken up in MeOH (5 mL) and Et₃N (5 mL) then stirred for an addition 16 h. The crude reaction was concentrated then taken up in DCM (10 mL) and NaHCO₃ (sat) (20 mL) and the solid was filtered, rinsed CH₂Cl₂ (10 mL) and water (10 mL) and dried to give the title compound N1 as a white solid (245 mg, 47%). See Table 1 below for NMR data.


EXAMPLES N2 AND N3:






Examples N2 and N3 were prepared using methods analogous to Example N1 above, except E5(xi)-Isomers A and B, respectively, were substituted in place of 2R,5S)-2,5-dimethyl-1-(tetrahydro-2H-pyran-4-ylmethyl)piperazine. See Table 1 below for names and NMR data.






The following Table 1 depicts further Ki app, structure, nomenclature, and NMR data of further embodiments of the invention. Unless otherwise mentioned, compounds in Table 1 were synthesized starting from commercially available materials or by known methods using routine modifications of the above described examples.






Table 1





While the invention has been illustrated by reference to specific embodiments, those skilled in the art will recognize that variations and modifications may be made through routine experimentation and practice of the invention. Thus, the invention is intended not to be limited by the foregoing description, but to be defined by the appended claims and their equivalents. The foregoing detailed description and examples have been given for clarity of understanding only.





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
A1	2.69	 <p>N4-(6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl))-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N2-ethyl-5-fluoropyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO-d ₆ mixture of rotamers) δ ppm 0.93 (s, 3 H), 0.98-1.07 (m, 6 H), 1.08-1.20 (m, 3 H), 1.55 (s, 3 H), 1.64 (s, 3 H), 2.08 (s, 3 H), 2.14-2.29 (m, 1 H), 2.52-2.64 (m, 2 H), 2.77-2.94 (m, 1 H), 3.23 (s, 3 H), 4.32-4.53 (m, 2 H), 6.49 (br. s., 0.6 H), 7.09 (br. s., 0.4 H), 7.67-8.11 (m, 1 H), 9.42 (br. s., 0.6 H), 10.14 (br. s., 0.4 H), 12.28 (br. s., 0.6 H), 12.44 (br. s., 0.4 H).	442






Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
A2	8.75	 <p>N⁴-(6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro-N²,N²-dimethylpyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆ mixture of rotamers) δ ppm 0.92 (s, 3 H), 0.96-1.08 (m, 6 H), 1.55 (s, 3 H), 1.65 (s, 3 H), 2.08 (s, 3 H), 2.11-2.24 (m, 1 H), 2.53-2.60 (m, 2 H), 2.71-2.84 (m, 1 H), 3.03 (s, 6 H), 3.16-3.28 (m, 1 H), 4.15-4.60 (m, 2 H), 7.84-8.13 (m, 1 H), 9.40 (br. s., 0.4 H), 9.88 (br. s., 0.6 H), 12.11 (br. s., 0.6 H), 12.36 (br. s., 0.4 H).	460
A3	17.9	 <p>N²-(cyclopropylmethyl)-N⁴-(6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoropyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.18 (m, 2 H), 0.35-0.51 (m, 2 H), 0.93 (s, 3 H), 0.96-1.08 (m, 7 H), 1.56 (s, 3 H), 1.65 (s, 3 H), 2.08 (s, 3 H), 2.13-2.25 (m, 1 H), 2.51-2.63 (m, 3 H), 2.84 (d, <i>J</i> =11.9 Hz, 1 H), 3.00-3.19 (m, 2 H), 4.33-4.53 (m, 2 H), 6.35-7.53 (m, 1 H), 7.94 (s, 1 H), 12.51 (br. s., 1 H).	486
A4	37.6	 <p>N⁴-(6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro-N²-isobutylpyrimidine-2,4-diamine acetate salt</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.85-0.94 (m, 9 H), 1.02 (s, 3 H), 1.04 (d, <i>J</i> =6.4 Hz, 3 H), 1.55 (s, 3 H), 1.62-1.69 (m, 3 H), 1.71-1.87 (m, 1 H), 1.90 (s, 3 H), 2.08 (s, 3 H), 2.13-2.24 (m, 1 H), 2.52-2.62 (m, 2 H), 2.83 (m, 1 H), 2.92-3.15 (m, 2 H), 4.27-4.59 (m, 2 H), 7.03 (br. s., 1 H), 7.83-8.01 (m, 1 H), 10.05 (br. s., 1 H).	488
A5	3.95	 <p>N²-cyclopropyl-N⁴-(6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoropyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆ mixture of rotamers) δ ppm 0.32-0.49 (m, 2 H), 0.54-0.75 (m, 2 H), 0.93 (s, 3 H), 0.97-1.12 (m, 6 H), 1.54 (s, 3 H), 1.63 (s, 3 H), 2.08 (s, 3 H), 2.12-2.25 (m, 1 H), 2.53-2.68 (m, 3 H), 2.76-2.90 (m, 1 H), 3.20-3.32 (m, 1 H), 4.15-4.63 (m, 2 H), 6.81 (br. s., 0.3 H), 7.37 (br. s., 0.7 H), 7.82-8.11 (m, 1 H), 9.47 (br. s., 0.3 H), 10.26 (br. s., 0.7 H), 12.29 (br. s., 0.3 H), 12.45 (br. s., 0.7 H).	472
A6	10.8	 <p>N⁴-(6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro-N²-methylpyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆ mixture of rotamers) δ ppm 0.93 (s, 3 H), 0.97-1.10 (m, 6 H), 1.54 (s, 3 H), 1.64 (s, 3 H), 2.08 (s, 3 H), 2.13-2.26 (m, 1 H), 2.51-2.61 (m, 2 H), 2.64-2.93 (m, 4 H), 3.16-3.32 (m, 1 H), 4.29-4.60 (m, 2 H), 6.55 (br. s., 0.3 H), 7.07 (br. s., 0.7 H), 7.76-8.12 (m, 1 H), 9.46 (br. s., 0.3 H), 10.15 (br. s., 0.7 H), 12.29 (br. s., 0.3 H), 12.43 (br. s., 0.7 H).	446






Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
A7	3.81	 <p>N⁴-(6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro-N²-isopropylpyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆ mixture of rotamers) δ ppm 0.93 (s, 3 H) 1.00- 1.08 (m, 6 H) 1.10- 1.16 (m, 6 H) 1.54 (s, 3 H) 1.64 (s, 3 H) 2.09 (s, 3 H) 2.14- 2.26 (m, 1 H) 2.51- 2.62 (m, 2 H) 2.78- 2.95 (m, 1 H) 3.36- 3.45 (m, 1 H) 3.90 (s, 1 H) 4.28- 4.60 (m, 2 H) 6.26 (br. s., 0.3 H), 7.05 (br. s., 0.7 H), 7.76- 8.11 (m, 1 H), 9.42 (br. s., 0.3 H), 10.19 (br. s., 0.7 H), 12.29 (br. s., 0.3 H), 12.50 (br. s., 0.7 H).	474
A8	3.39	 <p>N⁴-(6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²-ethylpyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆ mixture of rotamers) δ ppm 0.93 (s, 3 H), 0.99- 1.06 (m, 6 H), 1.08- 1.18 (m, 3 H), 1.55 (s, 3 H), 1.64 (s, 3 H), 2.09 (s, 3 H), 2.13- 2.24 (m, 1 H), 2.52- 2.58 (m, 2 H), 2.84 (m, 1 H), 3.22- 3.30 (m, 2 H), 3.37- 3.41 (m, 1 H), 4.30- 4.55 (m, 2 H), 5.97 (s, 1 H), 6.31 (br. s., 0.4 H), 7.10 (br. s., 0.6 H), 7.67- 7.98 (m, 1 H), 9.29 (br. s., 0.4 H), 9.84 (br. s., 0.6 H), 12.13 (br. s., 0.4 H), 12.47 (br. s., 0.6 H).	442
A9	9.73	 <p>N⁴-(6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²,N²-dimethylpyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆ mixture of rotamers) δ ppm 0.93 (s, 3 H), 0.96- 1.04 (m, 6 H), 1.56 (s, 3 H), 1.67 (s, 3 H), 2.08 (s, 3 H), 2.12- 2.23 (m, 1 H), 2.52- 2.59 (m, 2 H), 2.80 (m, 1 H), 3.06 (s, 6 H), 3.18- 3.29 (m, 1 H), 4.30- 4.53 (m, 2 H), 5.97 (br. s., 0.5 H), 6.38 (br. s., 0.5 H), 7.78- 8.03 (m, 1 H), 9.24 (br. s., 0.5 H), 9.65 (br. s., 0.5 H), 12.03- 12.26 (m, 1 H).	442
A10	1.01	 <p>5-(((8S)-6,8-dimethyl-6,9-diazaspiro[4.5]dec-9-yl)carbonyl)-N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm 0.94- 1.12 (5 H, m), 1.43 (3 H, m), 1.57 (3 H, s), 1.68 (3 H, s), 1.90 (3 H, s), 2.09 (3 H, s), 2.45 (3 H, s), 2.66 (1 H, m), 2.81 (1 H, m), 4.60 (2 H, m), 8.22 (1 H, s), 10.10 (1 H, s).	457
A11	<10	 <p>N⁴-(5-(((2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl)carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²-ethyl-5-fluoropyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 0.91 - 1.05 (m, 6 H) 1.05 - 1.28 (m, 6 H) 1.47 - 1.61 (m, 4 H) 1.61 - 1.78 (m, 5 H) 1.90 (s, 3 H) 2.43 (br. s., 2 H) 2.69 - 2.94 (m, 2 H) 3.01 - 3.16 (m, 2 H) 3.17 - 3.29 (m, 3 H) 3.74 - 3.93 (m, 2 H) 4.39 - 4.58 (m, 2 H) 7.95 (br. s., 1 H) 10.06 (br. s., 1 H) 12.29 (br. s., 1 H)	530





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
A12	<10	 <p><i>N</i>⁴-((5-(((2<i>S</i>,5<i>R</i>)-2,5-dimethyl-4-(tetrahydro-2<i>H</i>-pyran-4-yl)piperazin-1-yl)carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-yl)-<i>N</i>²-ethyl-5-fluoropyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 0.92 - 1.06 (m, 6 H) 1.06 - 1.17 (m, 3 H) 1.37 - 1.50 (m, 1 H) 1.53 (br. s., 1 H) 1.56 (s, 3 H) 1.61 (br. s., 1 H) 1.65 (s, 3 H) 2.03 - 2.23 (m, 1 H) 2.68 - 2.80 (m, 2 H) 2.80 - 2.99 (m, 2 H) 3.02 - 3.16 (m, 2 H) 3.16 - 3.31 (m, 6 H) 3.81 - 3.97 (m, 2 H) 4.38 - 4.55 (m, 1 H) 7.95 (d, <i>J</i> =3.01 Hz, 1 H)	516
AA1	22.9	 <p>5-(((2<i>S</i>,5<i>R</i>)-4-ethyl-2,5-dimethylpiperazin-1-yl)carbonyl)-<i>N</i>-(5-fluoro-2-methoxypyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.98 (6 H, s), 1.58 (3 H, s), 1.67 (3 H, s), 1.90 (3 H, s), 2.77 (2 H, bm), 3.05 (2 H, bm), 3.32 (8 H, s), 3.82 (3 H, s), 4.59-4.73 (2 H, m), 8.13 (1 H, s), 10.19 (1 H, s), 11.95 (1 H, s).	447
AA2	18.9	 <p>5-(((2<i>S</i>,5<i>R</i>)-4-ethyl-2,5-dimethylpiperazin-1-yl)carbonyl)-<i>N</i>-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.98 (8 H, d, <i>J</i> =4.04 Hz), 1.59 (3 H, s), 1.68 (3 H, s), 1.91 (1 H, s), 2.38 (2 H, s), 2.47 (3 H, s), 2.74 (1 H, s), 3.05 (2 H, s), 3.32 (18 H, s), 4.70 (2 H, s), 8.21 (1 H, s).	431
AA3	14.8	 <p>5-(((2<i>S</i>,5<i>R</i>)-4-ethyl-2,5-dimethylpiperazin-1-yl)carbonyl)-<i>N</i>-(5-fluoro-2,6-dimethylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.91 - 0.99 (10 H, m), 1.58 (3 H, s), 1.68 (3 H, s), 1.90 (3 H, s), 1.93 (1 H, d, <i>J</i> =10.86 Hz), 2.26 - 2.30 (3 H, m), 2.31 - 2.40 (3 H, m), 2.43 (3 H, s), 2.66 - 2.78 (2 H, m), 3.04 (2 H, d, <i>J</i> =8.34 Hz), 4.63 - 4.74 (2 H, m).	445
AA4	147	 <p>2(<i>S</i>),5(<i>S</i>)-[[dimethyl-4-methylpiperazin-1-yl]carbonyl)-<i>N</i>-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, CDCl ₃) δ ppm 1.09 (d, <i>J</i> =8 Hz, 3 H), 1.33 (d, <i>J</i> =8 Hz, 3 H), 1.72 (s, 3 H), 1.76 (s, 3 H), 2.21 (m, 1H), 2.27 (s, 3H), 2.45 (m, 1 H), 2.57 (s, 3H), 2.66 - 2.69 (m, 1 H), 2.92-2.98 (m, 1 H), 3.18-3.22 (m, 1H), 3.86 (m, 1H), 4.60 (d, <i>J</i> =16 Hz, 1 H), 4.69 (d, <i>J</i> =16 Hz, 1 H), 8.11 (s, 1H)	417





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
AA5	29.1	 <p>[3-(5-fluoro-2-methyl-pyrimidin-4-ylamino)-6,6-dimethyl-4,6-dihydro-1H-pyrrolo[3,4-c]pyrazol-5-yl]-[4-(3-hydroxy-propyl)-2,5-dimethyl-piperazin-1-yl]-methanone</p>	¹ H NMR (400 MHz, CD ₃ OD) δ ppm 1.13 (d, <i>J</i> =8 Hz, 3 H), 1.21 (d, <i>J</i> =8 Hz, 3 H), 1.71 (s, 3 H), 1.79 (s, 3 H), 1.81 (m, 2H), 2.43 (m, 1H), 2.56 (s, 3H), 2.71 (m, 2 H), 2.76 (m, 1 H), 3.12 (m, 1 H), 3.19 (m, 1H), 3.26 (m, 2H), 3.65 (m, 2H), 4.81 (m, 2 H), 8.11 (s, 1H)	461
B1	5.09	 <p>N²-ethyl-5-fluoro-N⁴-((5-(((2S,5R)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl)carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)pyrimidine-2,4-diamine acetate salt</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δppm 0.93- 1.01 (m, 6 H), 1.11 (t, <i>J</i> =7.2 Hz, 3 H), 1.56 (s, 3 H), 1.58- 1.63 (m, 2 H), 1.65 (s, 3 H), 1.85- 1.96 (m, 2 H), 2.13- 2.26 (m, 1 H), 2.32- 2.43 (m, 2 H), 2.60- 2.72 (m, 1 H), 2.73- 2.81 (m, 1 H), 2.98- 3.14 (m, 2 H), 3.22- 3.28 (m, 2 H), 3.29- 3.37 (m, 4 H), 4.40- 4.55 (m, 2 H), 6.94 (br. s., 1 H), 7.65- 8.09 (m, 1 H), 10.09 (br. s., 1 H), 12.33 (br. s., 1 H).	504
B2	24.5	 <p>N⁴-(6,6-dimethyl-5-(((3S,8aS)-3-methylhexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²-ethyl-5-fluoropyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆ mixture of rotamers) δ ppm 1.02- 1.16 (m, 3 H), 1.21 (m, 3 H), 1.30 (m, 1 H), 1.58 (m, 6 H), 1.67- 1.89 (m, 3 H), 1.87- 2.03 (m, 1 H), 2.11- 2.27 (m, 1 H), 2.63- 2.85 (m, 2 H), 2.85- 3.00 (m, 1 H), 3.23 (m, 2 H), 3.30- 3.44 (m, 2 H), 3.74- 3.98 (m, 1 H), 4.20- 4.56 (m, 2 H), 6.46 (br. s., 0.3 H), 7.12 (br. s., 0.7 H), 7.77- 8.11 (m, 1 H), 9.46 (br. s., 0.3 H), 10.18 (br. s., 0.7 H), 12.28 (br. s., 0.3 H), 12.42 (br. s., 0.7 H).	458
B3	2.03	 <p>N⁴-(6,6-dimethyl-5-(((2S,5R)-2,4,5-trimethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²-ethyl-5-fluoropyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.93 (d, <i>J</i> =5.65 Hz, 3 H) 0.97 (d, <i>J</i> =6.03 Hz, 3 H) 1.12 (t, <i>J</i> =6.88 Hz, 3 H) 1.57 (s, 3 H) 1.66 (s, 3 H) 1.76- 1.88 (m, 1 H) 2.00- 2.10 (m, 1 H) 2.14 (s, 3 H) 2.24- 2.38 (m, 1 H) 2.69 (d, <i>J</i> =9.80 Hz, 1 H) 2.92- 3.16 (m, 2 H) 3.18- 3.31 (m, 2 H) 4.35- 4.60 (m, 2 H) 7.09 (br. s., 1 H) 7.96 (s, 1 H) 10.24 (br. s., 1 H) 12.47 (br. s., 1 H).	446.4






Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
B4	13.5	 <p><i>N</i>²-ethyl-5-fluoro-<i>N</i>⁴-((2<i>S</i>,5<i>R</i>)-4-(2-methoxyethyl)-2,5-dimethylpiperazin-1-yl)carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-yl)pyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.92- 1.01 (m, 6 H), 1.11 (t, <i>J</i> =7.2 Hz, 3 H), 1.56 (s, 3 H), 1.65 (s, 3 H), 1.98- 2.09 (m, 1 H), 2.29- 2.48 (m, 3 H), 2.73- 2.87 (m, 2 H), 2.97- 3.05 (m, 1 H), 3.09 (d, <i>J</i> =9.8 Hz, 1 H), 3.17- 3.29 (m, 5 H), 3.40 (t, <i>J</i> =5.9 Hz, 2 H), 4.39- 4.57 (m, 2 H), 6.93 (br. s., 1 H), 7.95- 8.01 (m, 1 H), 10.08 (br. s., 1 H), 12.18 (br. s., 1 H).	490.4
B5	29.8	 <p><i>N</i>⁴-((2<i>S</i>,5<i>R</i>)-2,5-dimethyl-4-(3,3,3-trifluoropropyl)piperazin-1-yl)carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-yl)-<i>N</i>²-ethyl-5-fluoropyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 0.91- 1.05 (m, 6 H) 1.06- 1.18 (m, 3 H) 1.57 (s, 3 H) 1.65 (s, 3 H) 1.89- 2.04 (m, 1 H) 2.32- 2.48 (m, 6 H) 2.74- 2.90 (m, 2 H) 3.02- 3.15 (m, 2 H) 3.16- 3.28 (m, 3 H) 4.47 (s, 2 H) 7.85- 8.00 (m, 1 H) 10.26 (bs, 1H).	528
B6	2.43	 <p>4-((6,6-dimethyl-5-((2<i>S</i>,5<i>R</i>)-2,4,5-trimethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-yl)amino]pyrimidine-2-carbonitrile</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 0.91- 1.05 (m, 3 H) 1.14- 1.29 (m, 6 H) 1.60 (s, 3 H) 1.69 (s, 3 H) 2.18 (s, 3 H) 2.27- 2.42 (m, 1 H) 2.64- 2.78 (m, 1 H) 2.96- 3.11 (m, 2 H) 3.16 (s, 1 H) 4.59- 4.72 (m, 2 H) 6.96 - 7.00 (m, 1H) 8.30- 8.40 (m, 1 H)	410
B7	20.1	 <p><i>N</i>-(5-fluoro-2-morpholin-4-ylpyrimidin-4-yl)-6,6-dimethyl-5-((2<i>S</i>,5<i>R</i>)-2,4,5-trimethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 0.89- 1.02 (m, 6 H) 1.59 (s, 3 H) 1.66 (s, 3 H) 1.78- 1.89 (m, 1 H) 1.95- 2.08 (m, 1 H) 2.13 (s, 3 H) 2.23- 2.36 (m, 1 H) 2.64- 2.76 (m, 1 H) 2.90- 3.05 (m, 2 H) 3.50- 3.57 (m, 4 H) 3.57- 3.68 (m, 4 H) 4.30- 4.41 (m, 1 H) 4.42- 4.52 (m, 1 H) 8.02 (s, 1 H) 9.76 (bs, 1H) 12.22 (bs, 1H).	488






Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
C1	33.4	 <p>N²-ethyl-5-fluoro-N⁴-{5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl}pyrimidine-2,4-diamine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 1.11 (m, 3 H), 1.66 (s, 6 H), 1.88- 2.15 (m, 6 H), 2.17 (s, 3 H), 2.59- 2.73 (m, 2 H), 3.12- 3.29 (m, 2 H), 4.65- 4.85 (m, 2 H), 6.22 (br. s., 0.2 H), 7.09 (br. s., 0.8 H), 7.97 (s, 1 H), 9.57 (br. s., 0.2 H), 10.34 (br. s., 0.8 H), 12.35 (br. s., 0.2 H), 12.54 (br. s., 0.8 H).	435
D1	5.43	 <p>N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆ mixture of isomers) δ ppm 0.90 (br. s., 3 H), 0.96 (br. s., 3 H), 1.06 (d, <i>J</i> =6.2 Hz, 3 H), 1.22 (t, <i>J</i> =7.5 Hz, 3 H), 1.57 (s, 3 H), 1.69 (s, 3 H), 2.08 (s, 3 H), 2.13- 2.25 (m, 1 H), 2.53- 2.66 (m, 2 H), 2.66- 2.78 (m, 2 H), 2.77- 2.91 (m, 1 H), 3.37- 3.51 (m, 1 H), 4.37- 4.86 (m, 2 H), 8.11- 8.43 (m, 1 H), 9.99 (br. s., 0.6 H), 10.20 (br. s., 0.4 H), 11.96 (br. s., 0.4 H), 12.40 (br. s., 0.6 H).	445
E1	10.5	 <p>N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[[(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine acetate salt</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.85- 0.99 (m, 6 H), 1.20- 1.31 (m, 3 H), 1.59 (s, 3 H), 1.68 (s, 3 H), 1.73- 1.88 (m, 1 H), 1.90 (s, 3 H), 1.96- 2.07 (m, 1 H), 2.10- 2.18 (m, 3 H), 2.25- 2.39 (m, 1 H), 2.64- 2.82 (m, 3 H), 2.89- 3.06 (m, 3 H), 4.53- 4.91 (m, 2 H), 8.24 (s, 1 H), 10.15 (s, 1 H).	431
E2	1.64	 <p>N-(2-ethyl-5-fluoropyrimidin-4-yl)-5-[[[(2S,5R)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.86- 0.92 (m, 1 H), 0.92- 0.99 (m, 6 H), 1.18- 1.30 (m, 3 H), 1.55- 1.65 (m, 5 H), 1.68 (s, 3 H), 1.83- 1.98 (m, 1 H), 2.10- 2.26 (m, 1 H), 2.30- 2.44 (m, 2 H), 2.62- 2.83 (m, 4 H), 2.95- 3.12 (m, 2 H), 3.19- 3.25 (m, 3 H), 3.27- 3.33 (m, 1 H), 4.47- 4.98 (m, 2 H), 8.25 (s, 1 H), 10.06 (s, 1 H), 11.71- 12.69 (m, 1 H).	489
E3	13.1	 <p>N-(2-ethoxypyrimidin-4-yl)-6,6-dimethyl-5-[[[(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.96 (dd, <i>J</i> =11.75, 5.94 Hz, 6 H), 1.31 (q, <i>J</i> =6.74 Hz, 3 H), 1.55- 1.61 (m, 3 H), 1.67 (s, 3 H), 1.88 (s, 3 H), 2.10- 2.14 (m, 1 H), 2.16 (s, 3 H), 2.33 (t, <i>J</i> =10.61 Hz, 1 H), 2.67- 2.74 (m, 1 H), 2.95- 3.04 (m, 2 H), 3.30 (s, 1 H), 3.35 (d, <i>J</i> =5.05 Hz, 2 H), 4.25- 4.36 (m, 2 H), 4.61- 4.71 (m, 2 H), 8.03 (d, <i>J</i> =5.56 Hz, 1 H).	429





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
E4	31.5	 <p>5-(((2S,5R)-2,5-dimethyl-4-(3,3,3-trifluoropropyl)piperazin-1-yl)carbonyl)-N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 0.93-1.04 (m, 6 H) 1.20- 1.29 (m, 3 H) 1.59 (s, 3 H) 1.69 (s, 3 H) 1.91- 2.02 (m, 1 H) 2.35- 2.48 (m, 3 H) 2.70- 2.78 (m, 1 H) 2.78- 2.91 (m, 2 H) 2.99- 3.12 (m, 1 H) 3.15- 3.23 (m, 4 H) 4.06- 4.18 (m, 2 H) 4.64- 4.79 (m, 2 H) 8.26 (s, 1 H).	513
E5	<10	 <p>2-((5S)-4-[[3-[(2-ethyl-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl]-1,5-dimethylpiperazin-2-yl)ethanol</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.93 (d, <i>J</i> =6.03 Hz, 3 H), 1.25 (t, <i>J</i> =7.63 Hz, 3 H), 1.37 - 1.53 (m, 1 H), 1.59 (s, 3 H), 1.64 - 1.77 (m, 4 H), 1.80 - 1.90 (m, 2 H), 2.16 (s, 3 H), 2.36 - 2.47 (m, 1 H), 2.66 - 2.86 (m, 3 H), 2.93 - 3.14 (m, 2 H), 3.37 - 3.51 (m, 2 H), 4.58 - 4.87 (m, 2 H), 8.26 (s, 1 H), 10.11 (s, 1 H), 12.41 (br. s., 1 H).	461
F1	39.9	 <p>5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 1.67 (s, 6 H) 1.88 (s, 3 H) 1.96- 1.98 (m, 2 H) 2.06- 2.18 (m, 2 H) 2.43 (s, 3 H) 2.62- 2.73 (m, 2 H) 3.17- 3.19 (m, 4 H) 4.98- 4.99 (m, 2 H) 8.19- 8.21 (m, 1 H).	406
G1	25.4	 <p>N-(2-ethyl-5-fluoropyrimidin-4-yl)-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 1.15- 1.27 (m, 3 H) 1.68 (s, 6 H) 1.75 (s, 1 H) 1.87- 2.01 (m, 2 H) 2.02- 2.15 (m, 3 H) 2.18 (s, 3 H) 2.61- 2.75 (m, 4 H) 3.16 (s, 2 H) 4.91- 5.01 (m, 2 H) 8.22 (s, 1 H).	420
H1	39.8	 <p>N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-5-[[[(3S,8aS)-3-methylhexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 1.11- 1.23 (m, 3 H) 1.24- 1.38 (m, 1 H) 1.54 (s, 3 H) 1.61 (s, 3 H) 1.66- 1.82 (m, 3 H) 1.90 (s, 3 H) 1.91- 2.00 (m, 1 H) 2.14- 2.24 (m, 1 H) 2.44 (s, 3 H) 2.70- 2.84 (m, 2 H) 2.86- 3.00 (m, 1 H) 3.78- 3.93 (m, 1 H) 4.42- 4.55 (m, 1 H) 4.58- 4.72 (m, 1 H) 8.16- 8.22 (m, 1 H), 10.1 (bs, 1H).	429





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
H2	18.1	 <p><i>N</i>-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-5-(((2<i>S</i>,5<i>R</i>)-2,4,5-trimethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 0.89-0.99 (m, 6 H) 1.55 (s, 3 H) 1.68 (s, 3 H) 1.73-1.85 (m, 1 H) 1.89 (s, 1 H) 1.97-2.09 (m, 1 H) 2.15 (s, 3 H) 2.25-2.39 (m, 1 H) 2.47 (s, 3 H) 2.63-2.76 (m, 1 H) 2.92-3.06 (m, 2 H) 4.63-4.78 (m, 2 H) 8.17-8.23 (m, 1 H) 10.21 (s, 1 H).	417
H3	7.21	 <p><i>N</i>-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-5-(((2<i>S</i>)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.91 (3 H, s), 0.95-1.02 (3 H, m), 1.06 (3 H, d, <i>J</i> =6.06 Hz), 1.57 (2 H, s), 1.68 (2 H, s), 1.91 (3 H, s), 2.09 (2 H, s), 2.20 (1 H, s), 2.45 (2 H, s), 2.61 (1 H, s), 2.81 (1 H, s), 3.33 (9 H, s), 4.60 (1 H, d, <i>J</i> =7.58 Hz), 8.22 (1 H, s).	
H4	86	 <p>5-(((3<i>S</i>)-3-ethyl-4-methylpiperazin-1-yl)carbonyl)-<i>N</i>-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.81 (t, <i>J</i> =7.45 Hz, 3 H), 1.37 (dt, <i>J</i> =14.46, 7.29 Hz, 1 H), 1.55 (td, <i>J</i> =7.14, 2.65 Hz, 1 H), 1.57-1.65 (m, 7 H), 1.86-1.94 (m, 1 H), 2.08-2.14 (m, 1 H), 2.16 (s, 4 H), 2.46 (s, 3 H), 2.66-2.74 (m, 1 H), 2.76-2.83 (m, 1 H), 3.21-3.28 (m, 2 H), 4.61 (s, 2 H), 8.20 (s, 1 H)	417
H5	50.8	 <p>5-(((3<i>R</i>)-3-ethyl-4-methylpiperazin-1-yl)carbonyl)-<i>N</i>-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.80 (t, <i>J</i> =7.45 Hz, 3 H), 1.37 (dt, <i>J</i> =14.40, 7.20 Hz, 1 H), 1.56 (dd, <i>J</i> =14.15, 7.58 Hz, 1 H), 1.62 (s, 6 H), 1.87-1.93 (m, 4 H), 2.08-2.14 (m, 1 H), 2.16 (s, 3 H), 2.46 (s, 3 H), 2.54 (d, <i>J</i> =10.36 Hz, 1 H), 2.71 (d, <i>J</i> =11.37 Hz, 1 H), 3.25 (t, <i>J</i> =14.02 Hz, 2 H), 3.33 (s, 4 H), 4.61 (s, 2 H), 8.21 (d, <i>J</i> =3.28 Hz, 1 H)	417





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
H6	14.9	 <p>5-[[[(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)methyl]piperazin-1-yl]carbonyl]-N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 0.91 - 1.01 (m, 6 H) 1.01 - 1.20 (m, 3 H) 1.50 (br. s., 1 H) 1.54 - 1.62 (m, 4 H) 1.68 (s, 6 H) 1.80 - 1.89 (m, 2 H) 2.46 (s, 4 H) 2.53 - 2.55 (m, 1 H) 2.76 - 2.90 (m, 1 H) 3.02 - 3.17 (m, 2 H) 3.20 - 3.30 (m, 2 H) 3.75 - 3.90 (m, 2 H) 4.62 - 4.78 (m, 2 H) 8.21 (d, <i>J</i> =3.39 Hz, 1 H) 10.12 (br. s., 1 H) 12.19 (br. s., 1 H)	501
H7	22.0	 <p>5-[[[(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]carbonyl]-N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 0.92 - 1.10 (m, 6 H) 1.13 - 1.25 (m, 1 H) 1.50 - 1.62 (m, 4 H) 1.68 (s, 3 H) 2.09 (br. s., 2 H) 2.47 (br. s., 3 H) 2.53 - 2.57 (m, 2 H) 2.65 - 2.95 (m, 4 H) 3.00 - 3.17 (m, 4 H) 3.80 - 3.96 (m, 2 H) 4.63 - 4.77 (m, 2 H) 8.20 (br. s., 1 H) 10.12 (br. s., 1 H)	487
H8	157	 <p>4-[[[(2R,5S)-4-[[3-[(5-fluoro-2-methylpyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl]-2,5-dimethylpiperazin-1-yl)methyl]tetrahydro-2H-pyran-4-ol</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.95 (d, <i>J</i> =5.27 Hz, 3 H), 1.00 (d, <i>J</i> =6.03 Hz, 3 H), 1.23 - 1.45 (m, 2 H), 1.49 - 1.64 (m, 5 H), 1.68 (s, 3 H), 2.01 - 2.17 (m, 2 H), 2.32 - 2.61 (m, 6 H), 2.94 - 3.18 (m, 2 H), 3.19 - 3.28 (m, 1 H), 3.49 - 3.69 (m, 4 H), 4.09 (br. s., 1 H), 4.67 (s, 2 H), 8.21 (d, <i>J</i> =3.58 Hz, 1 H), 10.15 (s, 1 H), 12.20 (br. s., 1 H).	517
H9	58.2	 <p>2-((5S)-4-[[3-[(5-fluoro-2-methylpyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl]-1,5-dimethylpiperazin-2-yl)ethanol</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.93 (d, <i>J</i> =6.03 Hz, 3 H), 1.35 - 1.50 (m, 1 H), 1.59 (s, 3 H), 1.64 - 1.76 (m, 4 H), 1.78 - 1.96 (m, 2 H), 2.14 (s, 3 H), 2.35 - 2.47 (m, 1 H), 2.41 - 2.49 (m, 3 H), 2.71 (dd, <i>J</i> =11.11, 2.26 Hz, 1 H), 2.92 - 3.02 (m, 1 H), 3.03 - 3.11 (m, 1 H), 3.36 - 3.49 (m, 2 H), 4.56 - 4.85 (m, 2 H), 8.21 (s, 1 H), 10.16 (br. s., 1 H).	447






Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
H10	180	 <p>2-((5S)-4-([3-[(5-fluoro-2-methylpyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl)-1,5-dimethylpiperazin-2-yl)ethanol</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 1.16 (d, <i>J</i> =6.40 Hz, 3 H), 1.38 - 1.53 (m, 1 H), 1.59 (s, 3 H), 1.64 (s, 3 H), 1.68 - 1.83 (m, 1 H), 1.93 - 2.04 (m, 1 H), 2.13 (s, 3 H), 2.18 - 2.29 (m, 1 H), 2.45 (s, 3 H), 2.53 - 2.62 (m, 1 H), 2.74 - 2.91 (m, 1 H), 2.99 - 3.14 (m, <i>J</i> =12.53, 2.17 Hz, 1 H), 3.37 - 3.50 (m, 2 H), 3.61 - 3.81 (m, 1 H), 4.36 - 4.80 (m, 2 H), 8.21 (d, <i>J</i> =3.77 Hz, 1 H), 10.09 (s, 1 H)	447
I1	1.73	 <p>N-(5-fluoro-2-propylpyrimidin-4-yl)-6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 0.90 (s, 3 H) 0.93- 0.94 (m, 2H) 0.97 (s, 3 H) 1.05- 1.07 (m, 3 H) 1.57 (s, 3 H) 1.63 (s, 3 H) 1.73- 1.78 (m, 2H) 1.90 (s, 2 H) 2.08 (s, 3 H) 2.54- 2.63 (m, 3 H) 2.65- 2.72 (m, 2 H) 2.74- 2.90 (m, 2 H) 4.53- 4.66 (m, 2 H) 8.24 (s, 1 H) 10.06 (s, 1 H).	459
I2	3.67	 <p>N-(5-fluoro-2-isopropylpyrimidin-4-yl)-6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 1.03- 1.05 (m, 3 H) 1.21- 1.22 (m, 3 H) 1.23- 1.24 (m, 6H) 1.27- 1.28 (m, 3H), 1.32- 1.34 (m, 3H) 1.58- 1.62 (m, 3 H) 1.69- 1.70 (m, 3 H) 2.63- 2.75 (m, 4 H) 2.98- 3.03 (m, 1 H) 3.17- 3.18 (m, 2 H) 4.49- 4.53 (m, 1 H) 4.69- 4.81 (m, 1 H) 8.28- 8.32 (m, 1 H) 10.10 (s, 1 H)	459
J1	21.2	 <p>5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆): δ ppm 1.23 (s, 3 H) 1.66 (s, 6 H) 1.92- 2.18 (m, 6 H) 2.20 (s, 3 H) 2.35 (s, 3 H) 2.64- 2.76 (m, 2 H) 4.95 (d, <i>J</i> =4.03 Hz, 2 H) 6.61 (d, <i>J</i> =6.55 Hz, 1 H) 7.34- 7.47 (m, 1 H) 9.31 (br, 1 H) 11.65 (br, 1 H)	405
J2	59.9	 <p>5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-N-(6-methylpyridin-2-yl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (MEOD, 400 MHz): δ ppm 1.77 (6H, s), 2.11-2.21 (2H, m), 2.25-2.42 (2H, m), 2.44 (3H, s) 2.52 (3H, s) 2.62- 2.73 (m, 2H) 3.03-3.08 (2H, m), 4.99 (2H, d, <i>J</i> = 4.5 Hz), 6.6 (1H, d, <i>J</i> = 8.3 Hz), 6.67 (1H, d, <i>J</i> = 7.3 Hz), 7.42-7.51 (1H, m).	387





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
J3	2.63	 <p><i>N</i>-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-5-(((2<i>S</i>,5<i>R</i>)-2,4,5-trimethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆): δ ppm 0.90-0.97 (m, 6 H) 1.57 (s, 3 H) 1.66 (s, 3 H) 1.80 (t, <i>J</i> =10.70 Hz, 1 H) 1.96-2.08 (m, 1 H) 2.13 (s, 3 H) 2.28-2.36 (m, 1 H) 2.40 (s, 3 H) 2.65-2.75 (m, 1 H) 2.92-3.06 (m, 2 H) 4.61-4.78 (m, 2 H) 6.58-6.70 (m, 1 H) 7.33-7.49 (m, 1 H) 9.36 (br, 1 H) 11.83 (br, 1 H)	416
J4	8.64	 <p>5-(((3<i>S</i>,8<i>aS</i>)-3,8<i>a</i>-dimethylhexahydropyrrolo[1,2-<i>a</i>]pyrazin-2(1<i>H</i>)-yl)carbonyl)-<i>N</i>-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆): δ ppm 0.86 (s, 3 H) 1.07 (d, <i>J</i> =6.55 Hz, 3 H) 1.39-1.51 (m, 2 H) 1.53 (s, 3 H) 1.60-1.76 (m, 5 H) 2.33-2.45 (m, 4 H) 2.57-2.69 (m, 2 H) 2.71-2.78 (m, 1 H) 3.02-3.14 (m, 2 H) 3.80-3.90 (m, 1 H) 4.40 (d, <i>J</i> =12.59 Hz, 1 H) 4.68 (d, <i>J</i> =12.59 Hz, 1 H) 6.62 (dd, <i>J</i> =8.06, 2.52 Hz, 1 H) 7.40 (dd, <i>J</i> =11.33, 8.06 Hz, 1 H) 9.27 (br, 1 H) 11.85 (br, 1 H)	
J5	6.43	 <p><i>N</i>-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-5-(((3<i>S</i>,8<i>aS</i>)-3-methylhexahydropyrrolo[1,2-<i>a</i>]pyrazin-2(1<i>H</i>)-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 1.19 (d, <i>J</i> =6.80 Hz, 3 H) 1.22-1.35 (m, 1 H) 1.53-1.84 (m, 11 H) 2.19 (dd, <i>J</i> =10.32, 3.53 Hz, 1 H) 2.57-2.82 (m, 3 H) 2.88-2.96 (m, 1 H) 3.79-3.90 (m, <i>J</i> =6.55 Hz, 1 H) 4.46 (d, <i>J</i> =12.84 Hz, 1 H) 4.63 (d, <i>J</i> =12.84 Hz, 1 H) 6.61 (dd, <i>J</i> =8.31, 2.64 Hz, 1 H) 7.40 (dd, <i>J</i> =10.83, 8.31 Hz, 1 H) 9.25 (br, 1 H) 11.73 (br, 1 H)	428
J6	1.17	 <p><i>N</i>-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-5-(((2<i>S</i>)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.90 (s, 3 H) 0.98 (s, 3 H) 1.05 (d, <i>J</i> =6.29 Hz, 3 H) 1.55 (s, 3 H) 1.66 (s, 3 H) 2.08 (s, 3 H) 2.15-2.25 (m, 1 H) 2.37 (s, 3 H) 2.53-2.62 (m, 2 H) 2.74-2.84 (m, 1 H) 3.36-3.45 (m, 1 H) 4.51-4.66 (m, 2 H) 6.62 (dd, <i>J</i> =7.81, 2.27 Hz, 1 H) 7.41 (dd, <i>J</i> =10.95, 7.81 Hz, 1 H) 9.30 (br, 1 H) 11.86 (br, 1 H)	430
J7	111	 <p>5-(((3<i>S</i>,8<i>aS</i>)-3,8<i>a</i>-dimethylhexahydropyrrolo[1,2-<i>a</i>]pyrazin-2(1<i>H</i>)-yl)carbonyl)-6,6-dimethyl-<i>N</i>-(6-methylpyridin-2-yl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, CHLOROFORM- <i>d</i>) δ ppm 1.12-1.22 (m, 6 H) 1.62-1.85 (m, 8 H) 1.86-1.99 (m, 2 H) 2.48 (s, 3 H) 2.53-2.66 (m, 1 H) 2.70-2.82 (m, 1 H) 3.02-3.12 (m, 1 H) 3.13-3.31 (m, 3 H) 3.97-4.10 (m, 1 H) 4.37 (d, <i>J</i> =12.09 Hz, 1 H) 4.53 (d, <i>J</i> =12.09 Hz, 1 H) 6.58 (d, <i>J</i> =8.31 Hz, 1 H) 6.67 (d, <i>J</i> =7.30 Hz, 1 H) 7.44-7.48 (m, 1 H)	424





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
J8	32	 <p>6,6-dimethyl-N-(6-methylpyridin-2-yl)-5-([(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, CHLOROFORM- <i>d</i>) δ ppm 1.04 (d, <i>J</i> =6.04 Hz, 3 H) 1.08 (d, <i>J</i> =6.29 Hz, 3 H) 1.73 (s, 3 H) 1.81 (s, 3 H) 2.19- 2.34 (m, 4 H) 2.49 (s, 3 H) 2.60- 2.71 (m, 1 H) 2.81 (dd, <i>J</i> =11.46, 2.90 Hz, 1 H) 2.91- 2.99 (m, 2 H) 3.22- 3.31 (m, 1 H) 4.48 (d, <i>J</i> =13.09 Hz, 1 H) 4.63 (d, <i>J</i> =13.09 Hz, 1 H) 6.53 (d, <i>J</i> =8.31 Hz, 1 H) 6.69 (d, <i>J</i> =7.55 Hz, 1 H) 7.46 (t, <i>J</i> =7.81 Hz, 1 H)	398
J9	2.1	 <p>N-[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]-6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.90 (s, 3 H), 0.93- 1.00 (m, 3 H), 1.07 (d, <i>J</i> =6.32 Hz, 3 H), 1.56 (s, 3 H), 1.68 (s, 3 H), 1.88 (s, 3 H), 2.05- 2.10 (m, 3 H), 2.19 (dd, <i>J</i> =11.62, 6.32 Hz, 1 H), 2.54- 2.62 (m, 2 H), 2.83 (d, <i>J</i> =12.13 Hz, 1 H), 3.41 (s, 2 H), 4.39 (s, 2 H), 4.53- 4.65 (m, 2 H), 8.31 (s, 1 H)	461
J10	<10	 <p>5-([(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl]carbonyl)-N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 0.89 - 1.02 (m, 6 H) 1.24 (t, <i>J</i> =7.63 Hz, 3 H) 1.46 - 1.61 (m, 1 H) 1.57 (s, 3 H) 1.61 - 1.69 (m, 5 H) 1.71 (s, 3 H) 1.78 - 1.94 (m, 2 H) 1.86 (br. s., 1 H) 2.67 - 2.79 (m, 3 H) 2.79 - 2.89 (m, 1 H) 2.99 - 3.13 (m, 2 H) 3.05 (br. s., 1 H) 3.74 - 3.89 (m, 2 H) 3.82 (br. s., 1 H) 4.59 - 4.84 (m, 2 H) 8.20 (br. s., 1 H)	515
J11	<10	 <p>5-([(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]carbonyl)-N-(4-methoxypyrimidin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 0.92 - 1.04 (m, 6 H) 1.34 - 1.42 (m, 1 H) 1.45 - 1.53 (m, 2 H) 1.53 - 1.60 (m, 5 H) 1.64 (s, 3 H) 2.07 - 2.16 (m, 1 H) 2.70 - 2.78 (m, 2 H) 2.78 - 2.88 (m, 1 H) 3.04 - 3.13 (m, 2 H) 3.16 (s, 3 H) 3.83 - 3.93 (m, 5 H) 4.06 - 4.15 (m, 1 H) 4.43 - 4.60 (m, 2 H) 6.31 (br. s., 1 H) 8.16 (br. s., 1 H)	485





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
J12	<10	 <p>5-[[[(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]carbonyl]-6,6-dimethyl-N-(4-methylpyrimidin-2-yl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 0.95 - 1.01 (m, 6 H) 1.32 - 1.43 (m, 1 H) 1.52 - 1.60 (m, 6 H) 1.65 (s, 3 H) 2.05 - 2.15 (m, 1 H) 2.37 (s, 3 H) 2.69 - 2.78 (m, 2 H) 2.80 - 2.90 (m, 1 H) 3.02 - 3.12 (m, 3 H) 3.14 - 3.19 (m, 4 H) 3.83 - 3.91 (m, 2 H) 4.04 - 4.15 (m, 1 H) 4.56 - 4.68 (m, 1 H) 6.72 (br. s., 1 H) 8.26 (br. s., 1 H)	469
J13	<10	 <p>5-[[[(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]carbonyl]-6,6-dimethyl-N-[4-(trifluoromethyl)pyrimidin-2-yl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 0.91 - 1.14 (m, 7 H) 1.52 - 1.63 (m, 6 H) 1.64 - 1.75 (m, 5 H) 2.04 - 2.28 (m, 1 H) 2.70 - 2.90 (m, 2 H) 2.99 - 3.20 (m, 3 H) 3.81 - 3.97 (m, 3 H) 4.60 (br. s., 2 H) 7.24 (br. s., 1 H) 8.75 (br. s., 1 H) 10.49 (br. s., 1 H) 12.35 (br. s., 1 H)	523
J14	19.2	 <p>5-[[[(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl]carbonyl]-N-(4-methoxypyrimidin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.95 (d, <i>J</i> =5.84 Hz, 3 H), 1.00 (d, <i>J</i> =6.03 Hz, 3 H), 1.05 - 1.26 (m, 2 H), 1.49 - 1.73 (m, 10 H), 1.83 - 1.98 (m, 2 H), 2.31 - 2.45 (m, 2 H), 2.81 (d, <i>J</i> =8.67 Hz, 1 H), 3.00 - 3.21 (m, 4 H), 3.82 (dd, <i>J</i> =9.61, 1.70 Hz, 2 H), 3.88 (s, 3 H), 4.51 (s, 2 H), 6.30 (d, <i>J</i> =5.46 Hz, 1 H), 8.15 (d, <i>J</i> =5.65 Hz, 1 H), 9.86 (br. s., 1 H)	499
J15	<10	 <p>5-[[[(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl]carbonyl]-6,6-dimethyl-N-(4-methylpyrimidin-2-yl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.95 (d, <i>J</i> =5.46 Hz, 3 H), 0.99 (d, <i>J</i> =5.84 Hz, 3 H), 1.04 - 1.19 (m, 2 H), 1.46 - 1.73 (m, 10 H), 1.80 - 1.95 (m, 2 H), 2.37 (s, 3 H), 2.39 - 2.47 (m, 2 H), 2.76 - 2.87 (m, 1 H), 3.00 - 3.14 (m, 2 H), 3.19 - 3.30 (m, 2 H), 3.82 (d, <i>J</i> =8.67 Hz, 2 H), 4.61 (s, 2 H), 6.72 (s, 1 H), 8.27 (s, 1 H), 9.88 (br. s., 1 H), 11.90 (br. s., <i>J</i> =1.51 Hz, 1 H).	483





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
J16	11.1	 <p><i>N</i>-(4,6-dimethylpyrimidin-2-yl)-5-([(2<i>S</i>,5<i>R</i>)-2,5-dimethyl-4-(tetrahydro-2<i>H</i>-pyran-4-ylmethyl)piperazin-1-yl]carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.94 (d, <i>J</i> =5.84 Hz, 3 H), 0.98 (d, <i>J</i> =6.03 Hz, 3 H), 1.01 - 1.22 (m, 2 H), 1.44 - 1.75 (m, 10 H), 1.78 - 1.90 (m, 2 H), 2.30 (s, 6 H), 2.37 - 2.46 (m, 2 H), 2.82 (d, <i>J</i> =10.74 Hz, 1 H), 2.99 - 3.13 (m, 2 H), 3.18 - 3.30 (m, 2 H), 3.82 (d, <i>J</i> =7.54 Hz, 2 H), 4.65 (s, 2 H), 6.61 (s, 1 H), 9.68 (br. s., 1 H), 11.90 (br. s., 1 H). Elemental Analysis: Calcd for C ₂₆ H ₄₀ N ₈ O ₂ •1.47 H ₂ O•0.66 CH ₃ COOH C 58.31, H 8.16, N 19.91	497
K1	1.12	 <p><i>N</i>-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-([(2<i>S</i>,5<i>R</i>)-2,4,5-trimethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (DMSO- <i>d</i> ₆ , 400 MHz): δ ppm 0.95 (5 H, dd, <i>J</i> =11.49, 6.19 Hz), 1.31 (2 H, t, <i>J</i> =7.07 Hz), 1.58 (2 H, s), 1.68 (2 H, s), 1.82- 1.88 (1 H, m), 1.90 (2 H, s), 2.08- 2.13 (1 H, m), 2.15 (2 H, s), 2.30- 2.37 (1 H, m), 2.66- 2.73 (1 H, m), 2.95- 3.03 (2 H, m), 3.34 (7 H, s), 4.23- 4.33 (2 H, m), 4.59- 4.69 (1 H, m), 8.12 (1 H, d, <i>J</i> =2.53 Hz).	447
K2	2.11	 <p><i>N</i>-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[(4-ethyl(2<i>S</i>,5<i>R</i>)-2,5-dimethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.94-0.98 (m, 9 H) 1.28 (t, <i>J</i> = 8 H) 1.57 (s, 3 H) 1.66 (s, 3 H) 1.95 (t, <i>J</i> = 8, 1 H) 2.28- 2.32 (m, 1 H) 2.37-2.41 (m, 2 H) 2.67- 2.77 (m, 2 H) 3.01-3.08 (m, 2 H) 4.21-4.29 (m, 2H) 4.61 (b, 2 H) 8.1 (m, 1 H) 10.1 (br, 1 H) 12.40 (br, 1 H).	461
K3	0.912	 <p><i>N</i>-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-([(2<i>S</i>)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ ppm 0.92 (3 H, s), 0.99 (3 H, s), 1.05 (3 H, d, <i>J</i> =6.32 Hz), 1.30 (3 H, t, <i>J</i> =7.07 Hz), 1.57 (3 H, s), 1.67 (3 H, s), 1.89 (1.4 H, s), 2.09 (3 H, s), 2.19 (1 H, dd, <i>J</i> =11.49, 6.69 Hz), 2.57 (2 H, d, <i>J</i> =12.13 Hz), 2.81 (1 H, d, <i>J</i> =11.87 Hz), 3.31 (24 H, s), 4.24 (2 H, q, <i>J</i> =6.99 Hz), 4.59 (2 H, m), 8.10 (1 H, s).	461






Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
K4	3.33	 <p><i>N</i>-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-((2<i>S</i>,5<i>R</i>)-4-(2-methoxyethyl)-2,5-dimethylpiperazin-1-yl)carbonyl-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>		491.4
K5	0.683	 <p><i>N</i>-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-((2<i>S</i>,5<i>R</i>)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl)carbonyl-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.95 (d, <i>J</i> =5.09 Hz, 3 H) 0.98 (d, <i>J</i> =6.22 Hz, 3 H) 1.29 (t, <i>J</i> =6.97 Hz, 3 H) 1.58 (s, 3 H) 1.59- 1.64 (m, 2 H) 1.67 (s, 3 H) 1.88- 2.01 (m, 1 H) 2.11- 2.26 (m, 1 H) 2.43 (d, <i>J</i> =6.59 Hz, 2 H) 2.58- 2.73 (m, 1 H) 2.73- 2.84 (m, 1 H) 2.98- 3.16 (m, 2 H) 3.21 (s, 3 H) 3.29- 3.36 (m, 2 H) 4.16- 4.35 (m, 2 H) 4.53- 4.73 (m, 2 H) 8.11 (s, 1 H) 10.09 (br. s., 1 H) 12.36 (br. s., 1 H)	505.4
K6	0.376	 <p><i>N</i>-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-5-((2<i>S</i>,5<i>R</i>)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl)carbonyl-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.91 (d, <i>J</i> =5.46 Hz, 3 H) 0.97 (d, <i>J</i> =6.03 Hz, 3 H) 1.58 (s, 3 H) 1.60- 1.65 (m, 2 H) 1.68 (s, 3 H) 1.88- 2.02 (m, 1 H) 2.09- 2.27 (m, 1 H) 2.32- 2.46 (m, 2 H) 2.57- 2.78 (m, 2 H) 2.96- 3.16 (m, 2 H) 3.21 (s, 3 H) 3.29- 3.39 (m, 2 H) 4.45- 4.68 (m, 2 H) 4.78- 5.06 (m, 2 H) 8.18 (s, 1 H) 10.32 (br. s., 1 H) 12.49 (br. s., 1 H)	559.4
K7	0.165	 <p><i>N</i>-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-6,6-dimethyl-5-((2<i>S</i>,5<i>R</i>)-2,4,5-trimethylpiperazin-1-yl)carbonyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.89- 0.96 (m, 6 H), 1.58 (s, 3 H), 1.68 (s, 3 H), 1.79- 1.92 (m, 1 H), 1.98- 2.09 (m, 1 H), 2.12 (s, 3 H), 2.24- 2.39 (m, 1 H), 2.60- 2.70 (m, 1 H), 2.90- 3.05 (m, 2 H), 4.49- 4.74 (m, 2 H), 4.79- 5.06 (m, 2 H), 8.18 (s, 1 H), 10.31 (br. s., 1 H), 12.47 (br. s., 1 H)	501.4
K8	0.181	 <p><i>N</i>-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-6,6-dimethyl-5-((2<i>S</i>)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl-1,4,5,6-tetrahydropyrrolo[3,4-<i>c</i>]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.88 (s, 3 H), 0.95 (s, 3 H), 1.03 (d, <i>J</i> =6.2 Hz, 3 H), 1.57 (s, 3 H), 1.68 (s, 3 H), 2.06 (s, 3 H), 2.12- 2.24 (m, 1 H), 2.50- 2.62 (m, 2 H), 2.82 (d, <i>J</i> =11.7 Hz, 1 H), 3.20- 3.30 (m, 1 H), 4.47- 4.66 (m, 2 H), 4.78- 5.01 (m, 2 H), 8.16 (br. s., 1 H), 10.34 (br. s., 1 H), 12.49 (br. s., 1 H)	515.4





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
K9	12.8	 <p>N-[5-fluoro-2-(3-methoxypropoxy)pyrimidin-4-yl]-6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 0.91 (s, 3 H) 0.98 (s, 3 H) 1.01- 1.11 (m, 3 H) 1.57 (s, 3 H) 1.67 (s, 3 H) 1.91- 1.98 (m, 2 H) 2.08 (s, 3 H) 2.15- 2.24 (m, 1 H) 2.53- 2.61 (m, 3 H) 2.76- 2.86 (m, 1 H) 3.16 (s, 1 H) 3.22 (s, 3 H) 3.40- 3.49 (m, 3 H) 4.18- 4.32 (m, 2 H) 4.52- 4.66 (m, 2 H) 8.08- 8.19 (m, 1 H).	505
K10	28.4	 <p>6,6-dimethyl-5-[(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 0.91- 1.04 (m, 6 H) 1.58 (s, 3 H) 1.68 (s, 3 H) 1.92- 1.99 (m, 2 H) 2.03- 2.13 (m, 1 H) 2.15 (s, 3 H) 2.27- 2.40 (m, 1 H) 2.65- 2.76 (m, 1 H) 2.93- 3.07 (m, 2 H) 3.22 (s, 3 H) 3.39- 3.49 (m, 4 H) 4.18- 4.34 (m, 2 H) 4.54- 4.69 (m, 2 H) 8.12 (d, 1 H) 10.14 (s, 1 H).	491
K11	42.7	 <p>N-[5-fluoro-2-(2-methoxyethoxy)pyrimidin-4-yl]-6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 0.91 (s, 3 H) 0.98 (s, 3 H) 1.02- 1.09 (m, 3 H) 1.57 (s, 3 H) 1.68 (s, 3 H) 2.08 (s, 3 H) 2.16- 2.28 (m, 1 H) 2.53- 2.60 (m, 2 H) 2.77- 2.86 (m, 1 H) 3.16 (s, 2 H) 3.27 (s, 3 H) 3.57- 3.70 (m, 2 H) 4.11 (s, 1 H) 4.27- 4.38 (m, 2 H) 4.50- 4.65 (m, 2 H) 8.12 (s, 1 H)	491
K12	15.2	 <p>N-[5-fluoro-2-(2-methoxyethoxy)pyrimidin-4-yl]-6,6-dimethyl-5-[(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 0.92- 1.04 (m, 6 H) 1.58 (s, 3 H) 1.68 (s, 3 H) 1.82- 1.90 (m, 2 H) 2.04- 2.13 (m, 1 H) 2.15 (s, 3 H) 2.29- 2.40 (m, 1 H) 2.65- 2.75 (m, 1 H) 2.94- 3.06 (m, 2 H) 3.27 (s, 3 H) 3.55- 3.68 (m, 2 H) 4.27- 4.41 (m, 2 H) 4.53- 4.67 (m, 2 H) 8.13 (s, 1 H) 10.16 (s, 1 H)	477





Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
K13	26.5	 <p>2(S),5(S)-{[dimethyl-4-methylpiperazin-1-yl]carbonyl}-N-(5-fluoro-2-ethoxypyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, CDCl ₃) δ ppm 1.18 (b, 3 H), 1.38 (t, J=8 Hz, 3 H), 1.43 (b, 3 H), 1.73 (s, 3 H), 1.74 (s, 3 H), 2.36 (b, 3H), 2.57 (m, 1H), 2.75 (m, 1 H), 3.00 (m, 1 H), 3.29 (m, 1 H), 3.88 (m, 1H), 4.34 (q, J = 8 Hz, 2H), 4.63 (d, J=16 Hz, 1 H), 4.69 (d, J=16 Hz, 1 H), 7.96 (s, 1H), 8.41 (b, 1H).	447
K14	71.7	 <p>[3-(2-Ethoxy-5-fluoro-pyrimidin-4-yl-amino)-6,6-dimethyl-4,6-dihydro-1H-pyrrolo[3,4-c]pyrazol-5-yl]-(R)-hexahydro-pyrrolo[1,2-a]pyrazin-2-yl-methanone</p>	¹ H NMR (400 MHz, ACETONITRILE- d ₃) δ ppm 1.28 (t, J=8 Hz, 3 H), 1.60 (s, 3H), 1.61 (s, 3H), 1.65 (m, 3 H), 1.92 (m, 1 H), 2.03 (m, 1 H), 2.14 (m, 1 H), 2.43 (m, 1 H), 2.76 (m, 1H), 2.90 (m, 1H), 2.96 (m, 1H), 3.40 (m, 2H), 3.52 (m, 1H), 4.23 (q, J=8 Hz, 2 H), 4.51(d, J = 12 Hz, 1 H), 4.61 (d, J = 12 Hz, 1 H), 8.10 (s, 1 H), 10.14 (s, 1H), 12.25 (b, 1 H)	445
K15	11.2	 <p>5-[[[(3S,8aS)-3,8a-dimethylhexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl]carbonyl]-N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, CHLOROFORM-d) δ ppm 1.15- 1.25 (m, 6 H) 1.42 (t, J=7.05 Hz, 3 H) 1.63- 1.87 (m, 8 H) 1.88- 1.99 (m, 2 H) 2.55- 2.66 (m, 1 H) 2.69- 2.79 (m, 1 H) 3.08- 3.39 (m, 4 H) 4.05- 4.18 (m, 1 H) 4.37 (q, J=7.05 Hz, 2 H) 4.53 (d, J=12.84 Hz, 1 H) 4.78 (d, J=12.84 Hz, 1 H) 8.00 (d, J=2.77 Hz, 1 H) 8.66 (brs, 1 H)	473
K16	38.4	 <p>5-[[[(3S)-3,4-dimethylpiperazin-1-yl]carbonyl]-N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, DMSO- d ₆) δ ppm 0.97 (d, J=6.06 Hz, 2 H), 1.30 (t, J=7.07 Hz, 3 H), 1.61 (s, 5 H), 2.06 (td, J=6.32, 3.28 Hz, 1 H), 2.12- 2.17 (m, 1 H), 2.18 (s, 2 H), 2.43 (dd, J=12.25, 9.98 Hz, 1 H), 2.69 (d, J=11.12 Hz, 1 H), 2.76- 2.85 (m, 1 H), 3.20- 3.29 (m, 1 H), 3.32 (s, 7 H), 4.27 (q, J=6.99 Hz, 2 H), 4.59 (s, 2 H), 8.10 (s, 1 H)	456

Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
K17	44.4	 <p>5-[(3R)-3,4-dimethylpiperazin-1-yl]carbonyl)-N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, ACETONITRILE- <i>d</i> ₃) δ ppm 1.17- 1.26 (m, 3 H), 1.35 (t, <i>J</i> =6.95 Hz, 3 H), 1.67 (d, <i>J</i> =3.03 Hz, 6 H), 2.11 (dt, <i>J</i> =4.99, 2.43 Hz, 5 H), 2.53 (s, 3 H), 2.71 (s, 1 H), 2.86 (s, 1 H), 3.43 (d, <i>J</i> =11.12 Hz, 2 H), 4.32 (q, <i>J</i> =7.07 Hz, 2 H), 4.64 (s, 2 H), 7.99 (d, <i>J</i> =2.78 Hz, 1 H), 8.42 (s, 1 H). Calcd for C ₂₀ H ₂₉ FN ₆ O (<i>M</i> + <i>H</i>): 317.	433
K18	25.3	 <p>5-[(2S,5R)-2,5-dimethyl-4-(3,3,3-trifluoropropyl)piperazin-1-yl]carbonyl)-N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 0.94-1.07 (m, 6 H) 1.23- 1.35 (m, 3 H) 1.58 (s, 3 H) 1.67 (s, 3 H) 1.95- 2.08 (m, 1 H) 2.33- 2.48 (m, 6 H) 2.76- 2.91 (m, 2 H) 3.03- 3.18 (m, 2 H) 4.18- 4.32 (m, 2 H) 4.62 (s, 2 H) 8.09 (s, 1 H) 10.10 (s, 1 H)	529
K19	<10	 <p>5-[(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl]carbonyl)-N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 0.89 - 0.97 (m, 3 H) 0.97 - 1.05 (m, 3 H) 1.05 - 1.20 (m, 3 H) 1.21 - 1.35 (m, 3 H) 1.51 (br. s., 1 H) 1.57 (s, 3 H) 1.67 (s, 5 H) 1.94 (br. s., 1 H) 2.37 - 2.47 (m, 3 H) 2.54 (d, <i>J</i> =0.75 Hz, 1 H) 2.76 - 2.89 (m, 1 H) 3.03 - 3.12 (m, 1 H) 3.12 - 3.21 (m, 1 H) 3.21 - 3.29 (m, 2 H) 3.76 - 3.90 (m, 2 H) 4.18 - 4.36 (m, 2 H) 4.53 - 4.74 (m, 2 H) 8.08 - 8.22 (m, 1 H) 10.14 (br. s., 1 H)	531
K20	26	 <p>N-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-[(3S,8aS)-3-isopropylhexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl]carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (400 MHz, CHLOROFORM- <i>d</i>) δ ppm 0.83- 1.05 (m, 7 H) 1.30- 1.51 (m, 4 H) 1.61- 1.95 (m, 10 H) 2.21- 2.46 (m, 2 H) 2.92- 3.11 (m, 2 H) 3.14- 3.32 (m, 1 H) 3.49- 3.57 (m, 1 H) 3.59- 3.76 (m, 1 H) 4.36 (q, <i>J</i> =6.97 Hz, 2 H) 4.54-4.60 (m, 1 H) 4.75- 4.92 (m, 1 H) 8.00 (d, <i>J</i> =2.52 Hz, 1 H)	487

Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
K21	<10	 <p>5-[(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]carbonyl-N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 0.92 - 1.08 (m, 6 H) 1.28 (t, J=6.97 Hz, 3 H) 1.32 - 1.44 (m, 1 H) 1.46 - 1.55 (m, 2 H) 1.57 (s, 3 H) 1.67 (s, 3 H) 2.08 - 2.21 (m, 1 H) 2.51 - 2.58 (m, 1 H) 2.53 (s, 1 H) 2.68 - 2.86 (m, 3 H) 3.04 - 3.15 (m, 1 H) 3.15 - 3.26 (m, 2 H) 3.26 - 3.48 (m, 3 H) 3.77 - 3.95 (m, 2 H) 4.16 - 4.34 (m, 2 H) 4.51 - 4.70 (m, 1 H) 8.10 (d, J=3.01 Hz, 1 H) 10.15 (br. s., 1 H)	517
K22	15.3	 <p>4-[(2R,5S)-4-[[3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl]-2,5-dimethylpiperazin-1-yl)methyl]tetrahydro-2H-pyran-4-ol</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) □ppm 0.96 (d, J=4.52 Hz, 3 H), 1.04 (d, J=5.84 Hz, 3 H), 1.22 - 1.44 (m, 5 H), 1.51 - 1.74 (m, 8 H), 2.06 - 2.24 (m, 2 H), 2.36 - 2.47 (m, 1 H), 2.52 - 2.68 (m, 2 H), 2.99 - 3.09 (m, 1 H), 3.11 - 3.22 (m, 1 H), 3.52 - 3.70 (m, 4 H), 4.09 (br. s., 1 H), 4.19 - 4.35 (m, 2 H), 4.50 - 4.73 (m, 2 H), 8.12 (s, 1 H), 10.14 (s, 1 H), 12.17 (br. s., 1 H).	547
K23	<10	 <p>2-[(5S)-4-[[3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl]-1,5-dimethylpiperazin-2-yl]ethanol</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.95 (d, J=6.03 Hz, 3 H), 1.30 (t, J=6.97 Hz, 3 H), 1.42 - 1.54 (m, 1 H), 1.58 (s, 3 H), 1.64 - 1.77 (m, 4 H), 1.83 - 1.99 (m, 1 H), 2.09 - 2.27 (m, 4 H), 2.51 - 2.59 (m, 1 H), 2.66 - 2.79 (m, 1 H), 2.94 - 3.13 (m, 2 H), 3.37 - 3.51 (m, 2 H), 4.27 (q, J=6.84 Hz, 2 H), 4.52 - 4.72 (m, 2 H), 8.12 (s, 1 H), 10.13 (s, 1 H), 12.42 (br. s., 1 H).	477
K24	<10	 <p>2-[(5S)-4-[[3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl]-1,5-dimethylpiperazin-2-yl]ethanol</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 1.19 (d, J=6.59 Hz, 3 H), 1.29 (t, J=7.06 Hz, 3 H), 1.39 - 1.55 (m, 1 H), 1.59 (s, 3 H), 1.63 (s, 3 H), 1.68 - 1.84 (m, 1 H), 1.97 - 2.10 (m, 1 H), 2.15 (s, 3 H), 2.23 - 2.36 (m, 1 H), 2.53 - 2.61 (m, 1 H), 2.70 - 2.89 (m, 1 H), 3.04 - 3.15 (m, 1 H), 3.39 - 3.48 (m, 2 H), 3.62 - 3.76 (m, 1 H), 4.26 (q, J=6.97 Hz, 2 H), 4.45 - 4.67 (m, 2 H), 8.11 (d, J=2.83 Hz, 1 H), 10.12 (s, 1 H).	477


Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
K25	36.5	 <p>2-((5S)-4-((3-((5-fluoro-2-methoxypyrimidin-4-yl)amino)-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl)carbonyl)-1,5-dimethylpiperazin-2-yl)ethanol</p>	¹ H NMR (300 MHz, CHLOROFORM- <i>d</i>) δ ppm 1.10 (d, <i>J</i> =6.03 Hz, 3 H), 1.74 (s, 3 H), 1.81 (s, 3 H), 1.88 - 2.03 (m, 1 H), 2.32 (t, <i>J</i> =11.40 Hz, 1 H), 2.48 (s, 3 H), 2.76 - 2.89 (m, 1 H), 2.88 - 3.07 (m, 2 H), 3.08 - 3.23 (m, 1 H), 3.28 - 3.43 (m, 2 H), 3.66 - 3.78 (m, 2 H), 3.82 - 3.93 (m, 1 H), 3.97 (s, 3 H), 4.62 - 4.84 (m, 2 H), 7.99 (d, <i>J</i> =2.64 Hz, 1 H), 8.92 (br. s., 1 H).	463
K26	54.7	 <p>2-((5S)-4-((3-((5-fluoro-2-methoxypyrimidin-4-yl)amino)-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl)carbonyl)-1,5-dimethylpiperazin-2-yl)ethanol</p>	¹ H NMR (300 MHz, CHLOROFORM- <i>d</i>) δ ppm 1.22 - 1.33 (m, 1 H), 1.38 (d, <i>J</i> =6.59 Hz, 3 H), 1.76 (d, <i>J</i> =2.64 Hz, 7 H), 2.00 - 2.23 (m, 2 H), 2.45 (s, 3 H), 2.50 - 2.63 (m, 2 H), 2.73 - 2.87 (m, 1 H), 3.14 - 3.44 (m, 2 H), 3.68 - 3.88 (m, 2 H), 3.95 (s, 4 H), 4.74 (s, 2 H), 7.99 (d, <i>J</i> =2.64 Hz, 1 H)	463
L1	16.8	 <p><i>N</i>-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ ppm 1.28 (2 H, t, <i>J</i> =7.07 Hz), 1.68 (4 H, s), 1.91 (2 H, s), 1.91 - 1.95 (1 H, m), 2.13 (2 H, d, <i>J</i> =9.09 Hz), 2.19 (2 H, s), 2.68 (1 H, d, <i>J</i> =5.31 Hz), 3.32 (8 H, s), 4.22 (2 H, q, <i>J</i> =7.07 Hz), 4.90 (1 H, d, <i>J</i> =3.28 Hz), 8.12 (1 H, s), 10.18 (1 H, s).	436
L2	168	 <p><i>N</i>-[5-fluoro-2-(2-methoxyethoxy)pyrimidin-4-yl]-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 1.68 (s, 6 H), 1.84 - 2.09 (m, 4 H), 2.10 - 2.16 (m, 2 H), 2.18 (s, 3 H), 2.60 - 2.74 (m, 2 H), 3.28 (s, 3 H), 3.55 - 3.66 (m, 2 H), 4.22 - 4.38 (m, 2 H), 4.79 - 4.96 (m, 2 H), 8.12 (s, 1 H), 10.14 (br. s., 1 H), 12.46 (br. s., 1 H)	466.4
L3	58.6	 <p><i>N</i>-[5-fluoro-2-(3-methoxypropoxy)pyrimidin-4-yl]-5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 1.68 (s, 6 H), 1.83 - 1.97 (m, 4 H), 1.97 - 2.09 (m, 2 H), 2.09 - 2.15 (m, 2 H), 2.17 (s, 3 H), 2.59 - 2.71 (m, 2 H), 3.23 (s, 3 H), 3.45 (t, <i>J</i> =6.3 Hz, 2 H), 4.21 (t, <i>J</i> =6.3 Hz, 2 H), 4.83 - 4.98 (m, 2 H), 8.11 (s, 1 H), 10.15 (br. s., 1 H), 12.47 (br. s., 1 H).	480.4

Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
L4	1.71	 <p>5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-N-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 1.68 (s, 6 H) 1.80- 1.90 (m, 2 H) 1.92- 2.11 (m, 2 H) 2.10- 2.16 (m, <i>J</i> =8.48 Hz, 2 H) 2.18 (s, 3 H) 2.58- 2.71 (m, 2 H) 4.78- 5.00 (m, 4 H) 8.13- 8.23 (m, <i>J</i> =3.01 Hz, 1 H) 8.29 (br. s., 2 H)	490.4
L5	129	 <p>N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[1-(3,3,3-trifluoropropyl)piperidin-4-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) 1.26- 1.35 (m, 3 H) 1.56- 1.71 (m, 12 H) 1.92- 2.02 (m, 2 H) 2.32- 2.46 (m, 3 H) 2.88- 3.02 (m, 3 H) 4.20- 4.30 (m, 2 H) 4.79 (s, 2 H) 8.04- 8.17 (m, 1 H) 10.18 (s, 1 H)	500
L6	181	 <p>N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 1.24- 1.36 (m, 3 H) 1.38- 1.55 (m, 2 H) 1.57- 1.69 (m, 10 H) 2.03- 2.22 (m, 2 H) 2.26- 2.45 (m, 2 H) 2.83- 3.00 (m, 2 H) 3.12- 3.27 (m, 4 H) 3.79- 3.96 (m, 2 H) 4.25 (q, <i>J</i> =6.8 Hz, 2 H) 4.68- 4.89 (m, 2 H) 8.09 (s, 1 H) 10.17 (br. s., 1 H) 12.41 (br. s., 1 H)	488.4
M1	4.24	 <p>N-(4-ethoxypyrimidin-2-yl)-6,6-dimethyl-5-[[[(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, CDCl ₃ - <i>d</i>) δ ppm 1.09 (m, 6 H) 1.42 (m, 3 H) 1.73 (s, 3 H) 1.80 (s, 3 H) 2.34 (s, 3 H) 2.71 (m, 2 H) 2.83 (m, 2 H) 2.98 (m, 2 H) 3.30 (bs, 2 H) 4.38 (m, 2 H) 4.44- 4.67 (m, 2 H) 6.27 (m, 1 H) 8.16 (m, 1 H)	429

Ex. No.	Ki app (nM)	Structure IUPAC Name	¹ H NMR	MS (m/z)
M2	16.1	 <p>N-(4-ethoxypyrimidin-2-yl)-5-(((2S,5R)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl)carbonyl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.97 (m, 6 H) 1.32 (m, 3 H) 1.56 (s, 3 H) 1.64 (s, 3 H) 1.85- 2.00 (m, 2 H) 2.12- 2.28 (m, 1 H) 2.31- 2.45 (m, 2 H) 2.58- 2.70 (m, 1 H) 2.70- 2.86 (m, 2 H) 2.96- 3.11 (m, 2 H) 3.21 (m, 5 H) 4.26- 4.42 (m, 2 H) 4.42- 4.58 (m, 2 H) 6.20- 6.35 (m, 1 H) 8.14 (s, 1 H) 9.96 (bs, 1H) 11.96 (bs, 1H)	487
N1	<10	 <p>5-(((2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl)carbonyl)-N-[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine</p>	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ ppm 0.94 (d, <i>J</i> =5.27 Hz, 3 H), 0.99 (d, <i>J</i> =5.84 Hz, 3 H), 1.03 - 1.22 (m, 2 H), 1.45 - 1.78 (m, 10 H), 1.82 - 1.98 (m, 2 H), 2.33 - 2.47 (m, 2 H), 2.83 (d, <i>J</i> =9.04 Hz, 1 H), 3.07 (d, <i>J</i> =9.04 Hz, 2 H), 3.20 - 3.30 (m, 2 H), 3.82 (d, <i>J</i> =10.17 Hz, 2 H), 4.42 (s, 2 H), 4.58 - 4.76 (m, 2 H), 8.33 (s, 1 H), 10.28 (br. s., 1 H), 12.36 (br. s., 1 H).	531
N2	<10	 <p>2-(((5S)-4-([3-([5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]amino)-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl)-1,5-dimethylpiperazin-2-yl)ethanol</p>	¹ H NMR (300 MHz, CHLOROFORM- <i>d</i>) δ ppm 1.11 (d, <i>J</i> =5.84 Hz, 3 H), 1.74 (s, 3 H), 1.81 (s, 3 H), 1.85 - 1.99 (m, 2 H), 2.55 (s, 3 H), 2.92 - 3.09 (m, 3 H), 3.10 - 3.24 (m, 1 H), 3.32 - 3.46 (m, 1 H), 3.51 (s, 3 H), 3.68 - 3.80 (m, 1 H), 3.80 - 3.95 (m, 1 H), 4.56 (s, 2 H), 4.66 - 4.88 (m, 3 H), 8.19 (d, <i>J</i> =2.83 Hz, 1 H), 9.24 (br. s., 1 H)	477
N3	15.6	 <p>((5S)-4-([3-([5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]amino)-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl)-1,5-dimethylpiperazin-2-yl)ethanol</p>	¹ H NMR (300 MHz, CHLOROFORM- <i>d</i>) δ ppm 1.30 - 1.38 (m, 3 H), 1.74 (s, 3 H), 1.77 (s, 3 H), 1.81 - 1.90 (m, 1 H), 1.94 - 2.03 (m, 1 H), 2.45 (s, 3 H), 2.49 - 2.63 (m, 2 H), 2.75 - 2.86 (m, 1 H), 3.14 - 3.29 (m, 1 H), 3.29 - 3.39 (m, 1 H), 3.51 (s, 3 H), 3.72 - 3.85 (m, 2 H), 3.86 - 4.00 (m, 1 H), 4.56 (s, 2 H), 4.59 - 4.73 (m, 2 H), 8.23 (d, <i>J</i> =2.83 Hz, 1 H), 9.17 (br. s., 1 H)	477

We claim:

1. A compound of Formula A,



wherein

X is C-R¹¹ or N, wherein R¹¹ is H, halo, OH, C₁-C₃alkyl, CF₃, or CN;

A and B are independently C or N;

R¹, R² and R³ are each independently selected from H, -R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ saturated or partially unsaturated cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)₂R^a, -(R^d)_m-OS(O)₂NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO₂, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)₂R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)₂R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)₂NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b or -(R^d)_m-NR^a-(R^e)-OR^b; wherein R² and R³ may together optionally cyclize to form a saturated or unsaturated 3-7 membered heterocyclyl fused to the 6-membered N-containing heteroaryl to which they are attached; and wherein any of the said alkyl, alkenyl, alkynyl, R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ saturated or partially unsaturated cycloalkyl, phenyl or 3-15 membered heterocyclyl, may independently be further optionally substituted by 0-3 R¹² groups;

R⁴ and R⁵ are each independently selected from H, -R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ saturated or partially unsaturated cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-

$C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)_2R^a, -(R^d)_m-OS(O)_2NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO_2, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)_2R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)_2R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)_2NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b$ or $-(R^d)_m-NR^a-(R^e)-OR^b$; wherein any of the said alkyl, alkenyl, alkynyl, R^a , R^b , R^c , R^d , R^e , C_3 - C_{12} saturated or partially unsaturated cycloalkyl, aryl or 3-15 membered heterocyclyl are independently optionally further substituted by 0-3 R^{12} groups,

R^6 and R^7 are each independently H, $-R^a-O-R^b$, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-(R^d)_m-(C_3-C_{12}$ saturated or partially unsaturated cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m-(C_1-C_6$ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)_2R^a, -(R^d)_m-OS(O)_2NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO_2, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)_2R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)_2R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)_2NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b$ or $-(R^d)_m-NR^a-(R^e)-OR^b$; wherein R^6 and R^7 may together optionally cyclize to form a C_3 - C_7 saturated or partially unsaturated cycloalkyl and wherein any of the said alkyl, alkenyl, alkynyl, R^a , R^b , R^c , R^d , R^e , C_3 - C_{12} saturated or partially unsaturated cycloalkyl, aryl or 3-15 membered heterocyclyl are independently optionally further substituted by 0-3 R^{12} groups;

R^8 is H, $-R^a-O-R^b$, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-(R^d)_m-(C_3-C_{12}$ saturated or partially unsaturated cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m-(C_1-C_6$ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)_2R^a, -(R^d)_m-OS(O)_2NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO_2, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)_2R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)_2R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)_2NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b$ or $-(R^d)_m-NR^a-(R^e)-OR^b$; and wherein any of the said alkyl, alkenyl, alkynyl, R^a , R^b , R^c , R^d , R^e , C_3 - C_{12} saturated or partially unsaturated cycloalkyl, phenyl, or 3-15 membered heterocyclyl are independently optionally further

substituted by 1-3 groups selected from -F, C₁-C₃ alkyl, C₁-C₃ perfluoroalkyl, hydroxyl, C₁-C₆alkoxyl, or oxo;

R⁹ and R¹⁰ are each independently C₁-C₂ alkyl or can together cyclize to form a cyclopropyl or cyclobutyl;

each R¹² is independently H, -R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ saturated or partially unsaturated cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)₂R^a, -(R^d)_m-OS(O)₂NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO₂, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)₂R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)₂R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)₂NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b or -(R^d)_m-NR^a-(R^e)_m-OR^b; and wherein any of the said alkyl, alkenyl, alkynyl, R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ saturated or partially unsaturated cycloalkyl, phenyl, or 3-15 membered heterocyclyl, are independently optionally further substituted by 1-3 groups selected from -F, C₁-C₃ alkyl, C₁-C₃ perfluoroalkyl, hydroxyl, C₁-C₆alkoxyl or oxo;

each R^a, R^b and R^c is independently selected from H, C₁-C₈ alkyl, C₂-C₈ alkenyl, -(R^d)_m-(C₃-C₈ saturated or partially unsaturated cycloalkyl), -(R^d)_m-(C₃-C₈ cycloalkenyl), C₂-C₈ alkynyl, -(R^d)_m-phenyl, or -(R^d)_m-(3-7 membered heterocyclyl), and each R^a, R^b and R^c is independently optionally further substituted by 1-3 groups selected from halide, hydroxyl, -CN, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxyl and C₁-C₆ alkylamino; or, when connected to the same nitrogen, R^a and R^b may together optionally form a 3-7 membered heterocyclyl, which may optionally be further substituted by 0-3 groups selected from halide, hydroxyl, -CN, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxyl or C₁-C₆ alkylamino;

each R^d and R^e is independently -(C₁-C₃ alkylene)-, -(C₂-C₅ alkenylene)-, or -(C₂-C₅ alkynylene)-;

and each m is independently 0 or 1;

with the proviso that when X is N, R⁶ and R⁷ are not both H, and that when X is C-R¹¹, R⁶ and R⁷ are both H;

or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, , wherein R^9 and R^{10} are both methyl.

3. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein X is N and R^6 and R^7 are each independently H or C₁-C₆alkyl but are not both H.

4. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein A is N and B is C.

5. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein A is C and B is N.

6. The compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein R^6 and R^7 are both methyl.

7. The compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein R^6 is H and R^7 is methyl.

8. The compound according to any of the preceding claims, or pharmaceutically acceptable salt thereof, wherein R^4 is $-R^a-O-R^b$, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, $-(R^d)_m-(C_3-C_{12}$ saturated or partially unsaturated cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m-(C_1-C_6$ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m-C(O)R^a$, $-(R^d)_m-C(O)OR^a$, $-(R^d)_m-C(O)NR^aR^b$, $-(R^d)_m-OR^a$, $-(R^d)_m-OC(O)R^a$, $-(R^d)_m-OC(O)NR^aR^b$, $-(R^d)_m-O-S(O)R^a$, $-(R^d)_m-OS(O)_2R^a$, $-(R^d)_m-OS(O)_2NR^aR^b$, $-(R^d)_m-OS(O)NR^aR^b$, $-(R^d)_m-NO_2$, $-(R^d)_m-NR^aR^b$, $-(R^d)_m-N(R^a)C(O)R^b$, $-(R^d)_m-N(R^a)C(O)OR^b$, $-(R^d)_m-N(R^c)C(O)NR^aR^b$, $-(R^d)_m-N(R^a)S(O)_2R^b$, $-(R^d)_m-N(R^a)S(O)R^b$, $-(R^d)_m-SR^a$, $-(R^d)_m-S(O)R^a$, $-(R^d)_m-S(O)_2R^a$, $-(R^d)_m-S(O)NR^aR^b$, $-(R^d)_m-S(O)_2NR^aR^b$, $-(R^d)_m-O-(R^e)_m-NR^aR^b$ or $-(R^d)_m-NR^a-(R^e)-OR^b$; wherein the

said R^a , R^b , R^c , R^d , R^e , C_3 - C_{12} saturated or partially unsaturated cycloalkyl, aryl, 3-15 membered heterocyclyl, are independently optionally further substituted by 0-3 R^{12} groups.

9. The compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein R^4 is methyl.


10. The compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein R^1 is $-R^a-O-R^b$, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-(R^d)_m$ -(C_3 - C_{12} saturated or partially unsaturated cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m$ -(C_1 - C_6 perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m$ - $C(O)R^a$, $-(R^d)_m$ - $C(O)OR^a$, $-(R^d)_m$ - $C(O)NR^aR^b$, $-(R^d)_m$ - OR^a , $-(R^d)_m$ - $OC(O)R^a$, $-(R^d)_m$ - $OC(O)NR^aR^b$, $-(R^d)_m$ - $O-S(O)R^a$, $-(R^d)_m$ - $OS(O)_2R^a$, $-(R^d)_m$ - $OS(O)_2NR^aR^b$, $-(R^d)_m$ - $OS(O)NR^aR^b$, $-(R^d)_m$ - NO_2 , $-(R^d)_m$ - NR^aR^b , $-(R^d)_m$ - $N(R^a)C(O)R^b$, $-(R^d)_m$ - $N(R^a)C(O)OR^b$, $-(R^d)_m$ - $N(R^c)C(O)NR^aR^b$, $-(R^d)_m$ - $N(R^a)S(O)_2R^b$, $-(R^d)_m$ - $N(R^a)S(O)R^b$, $-(R^d)_m$ - SR^a , $-(R^d)_m$ - $S(O)R^a$, $-(R^d)_m$ - $S(O)_2R^a$, $-(R^d)_m$ - $S(O)NR^aR^b$, $-(R^d)_m$ - $S(O)_2NR^aR^b$, $-(R^d)_m$ - $O-(R^e)_m$ - NR^aR^b or $-(R^d)_m$ - NR^a - $-(R^e)_m$ - OR^b ; wherein the said- R^a , R^b , R^c , R^d , R^e , C_3 - C_{12} saturated or partially unsaturated cycloalkyl, aryl, the said 3-15 membered heterocyclyl, are independently optionally further substituted by 0-3 R^{12} groups.

11. The compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein R^1 is $-(R^d)_m$ - OR^a , C_1 - C_8 alkyl, or $-(R^d)_m$ - NR^aR^b .

12. The compound according to any of the preceding claims, or pharmaceutically acceptable salt thereof, wherein R^8 is $-R^a-O-R^b$, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-(R^d)_m$ -(C_3 - C_{12} saturated or partially unsaturated cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m$ -(C_1 - C_6 perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m$ -CN, $-(R^d)_m$ - OR^a , or $-(R^d)_m$ - NR^aR^b .

13. The compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein each R^d and R^e is independently an $-(C_1-C_3$ alkylene).

14. A compound of Formula B,



wherein

X is C-R¹¹ or N, wherein R¹¹ is H, halo, OH, C₁-C₃alkyl, CF₃, or CN;

A and B are independently C or N;

R¹ is -R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ saturated or partially unsaturated cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)₂R^a, -(R^d)_m-OS(O)₂NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO₂, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)₂R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)₂R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)₂NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b or -(R^d)_m-NR^a-(R^e)-OR^b; and wherein any of the said alkyl, alkenyl, alkynyl, R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ saturated or partially unsaturated cycloalkyl, phenyl or 3-15 membered heterocyclyl, may independently be further optionally substituted by 0-3 R¹² groups;

R² and R³ are each independently selected from H, -R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ saturated or partially unsaturated cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)₂R^a, -(R^d)_m-OS(O)₂NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO₂, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-

$N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)_2R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)_2R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)_2NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b$ or $-(R^d)_m-NR^a-(R^e)-OR^b$; wherein R^2 and R^3 may together optionally cyclize to form a saturated or unsaturated 3-7 membered heterocyclyl fused to the 6-membered N-containing heteroaryl to which they are attached; and wherein any of the said alkyl, alkenyl, alkynyl, $R^a, R^b, R^c, R^d, R^e, C_3-C_{12}$ saturated or partially unsaturated cycloalkyl, phenyl or 3-15 membered heterocyclyl, may independently be further optionally substituted by 0-3 R^{12} groups;

R^4 and R^5 are each independently selected from H, $-R^a-O-R^b, C_1-C_8$ alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, $-(R^d)_m-(C_3-C_{12}$ saturated or partially unsaturated cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m-(C_1-C_6$ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, (R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)_2R^a, -(R^d)_m-OS(O)_2NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO_2, (R^d)_m-NR^aR^b, (R^d)_m-N(R^a)C(O)R^b, (R^d)_m-N(R^a)C(O)OR^b, (R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)_2R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)_2R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)_2NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b$ or $-(R^d)_m-NR^a-(R^e)-OR^b$; wherein any of the said alkyl, alkenyl, alkynyl, $R^a, R^b, R^c, R^d, R^e, C_3-C_{12}$ saturated or partially unsaturated cycloalkyl, aryl or 3-15 membered heterocyclyl are independently optionally further substituted by 0-3 R^{12} groups,

R^8 is H, $-R^a-O-R^b, C_1-C_8$ alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, $-(R^d)_m-(C_3-C_{12}$ saturated or partially unsaturated cycloalkyl), $-(R^d)_m$ -phenyl, $-(R^d)_m$ -(3-15 membered heterocyclyl), $-(R^d)_m-(C_1-C_6$ perfluoroalkyl), $-(R^d)_m$ -halide, $-(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, (R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, (R^d)_m-OC(O)R^a, (R^d)_m-OC(O)NR^aR^b, (R^d)_m-O-S(O)R^a, (R^d)_m-OS(O)_2R^a, (R^d)_m-OS(O)_2NR^aR^b, (R^d)_m-OS(O)NR^aR^b, (R^d)_m-NO_2, (R^d)_m-NR^aR^b, (R^d)_m-N(R^a)C(O)R^b, (R^d)_m-N(R^a)C(O)OR^b, (R^d)_m-N(R^c)C(O)NR^aR^b, (R^d)_m-N(R^a)S(O)_2R^b, ((R^d)_m-N(R^a)S(O)R^b, (R^d)_m-SR^a, (R^d)_m-S(O)R^a, (R^d)_m-S(O)_2R^a, (R^d)_m-S(O)NR^aR^b, (R^d)_m-S(O)_2NR^aR^b, (R^d)_m-O-(R^e)_m-NR^aR^b$ or $-(R^d)_m-NR^a-(R^e)-OR^b$; and wherein any of the said alkyl, alkenyl, alkynyl, $R^a, R^b, R^c, R^d, R^e, C_3-C_{12}$ saturated or partially unsaturated cycloalkyl, phenyl, or 3-15 membered heterocyclyl are independently optionally further

substituted by 1-3 groups selected from -F, C₁-C₃ alkyl, C₁-C₃ perfluoroalkyl, hydroxyl, C₁-C₆alkoxyl, or oxo;

R⁹ and R¹⁰ are each independently C₁-C₂ alkyl or can together cyclize to form a cyclopropyl or cyclobutyl;

each R¹² is independently H, -R^a-O-R^b, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, -(R^d)_m-(C₃-C₁₂ saturated or partially unsaturated cycloalkyl), -(R^d)_m-phenyl, -(R^d)_m-(3-15 membered heterocyclyl), -(R^d)_m-(C₁-C₆ perfluoroalkyl), -(R^d)_m-halide, -(R^d)_m-CN, -(R^d)_m-C(O)R^a, -(R^d)_m-C(O)OR^a, -(R^d)_m-C(O)NR^aR^b, -(R^d)_m-OR^a, -(R^d)_m-OC(O)R^a, -(R^d)_m-OC(O)NR^aR^b, -(R^d)_m-O-S(O)R^a, -(R^d)_m-OS(O)₂R^a, -(R^d)_m-OS(O)₂NR^aR^b, -(R^d)_m-OS(O)NR^aR^b, -(R^d)_m-NO₂, -(R^d)_m-NR^aR^b, -(R^d)_m-N(R^a)C(O)R^b, -(R^d)_m-N(R^a)C(O)OR^b, -(R^d)_m-N(R^c)C(O)NR^aR^b, -(R^d)_m-N(R^a)S(O)₂R^b, -(R^d)_m-N(R^a)S(O)R^b, -(R^d)_m-SR^a, -(R^d)_m-S(O)R^a, -(R^d)_m-S(O)₂R^a, -(R^d)_m-S(O)NR^aR^b, -(R^d)_m-S(O)₂NR^aR^b, -(R^d)_m-O-(R^e)_m-NR^aR^b or -(R^d)_m-NR^a-(R^e)_m-OR^b; and wherein any of the said alkyl, alkenyl, alkynyl, R^a, R^b, R^c, R^d, R^e, C₃-C₁₂ saturated or partially unsaturated cycloalkyl, phenyl, or 3-15 membered heterocyclyl, are independently optionally further substituted by 1-3 groups selected from -F, C₁-C₃ alkyl, C₁-C₃ perfluoroalkyl, hydroxyl, C₁-C₆alkoxyl or oxo;

each R^a, R^b and R^c is independently selected from H, C₁-C₈ alkyl, C₂-C₈ alkenyl, -(R^d)_m-(C₃-C₈ saturated or partially unsaturated cycloalkyl), -(R^d)_m-(C₃-C₈ cycloalkenyl), C₂-C₈ alkynyl, -(R^d)_m-phenyl, or -(R^d)_m-(3-7 membered heterocyclyl), and each R^a, R^b and R^c is independently optionally further substituted by 1-3 groups selected from halide, hydroxyl, -CN, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxyl and C₁-C₆ alkylamino; or, when connected to the same nitrogen, R^a and R^b may together optionally form a 3-7 membered heterocyclyl, which may optionally be further substituted by 0-3 groups selected from halide, hydroxyl, -CN, C₁-C₆ alkyl, C₁-C₆ perfluoroalkyl, C₁-C₆ alkoxyl or C₁-C₆ alkylamino;

each R^d and R^e is independently-(C₁-C₃ alkylene)-, -(C₂-C₅ alkenylene)-, or-(C₂-C₅ alkynylene)-;

and each m is independently 0 or 1,

or a pharmaceutically acceptable salt thereof.

15. The compound according to claim 14, or a pharmaceutically acceptable salt thereof, wherein A is N and B is C.

16. The compound according to claims 14 or 15, or a pharmaceutically acceptable salt thereof, wherein R⁹ and R¹⁰ are both methyl.

17. The compound according to any of claims 14, 15, or 16, or a pharmaceutically acceptable salt thereof, wherein R⁴ is-(R^d)_m-OR^a, C₁-C₈ alkyl, C₂-C₈ alkenyl or C₂-C₈ alkynyl.

18. The compound according to any of claims 14, 15, 16 or 17, or a pharmaceutically acceptable salt thereof, wherein R⁴ is methyl.

19. The compound according to any of claims 14, 15, 16, 17 or 18, or a pharmaceutically acceptable salt thereof, wherein R¹ is -(R^d)_m-OR^a, C₁-C₈ alkyl, or-(R^d)_m-NR^aR^b.

20. The compound according to any of claims 14, 15, 16, 17, 18 or 19, or a pharmaceutically acceptable salt thereof, wherein each R^d and R^e is independently an-(C₁-C₃ alkylene)-.

21. A compound selected from the group consisting of:

N⁴-(6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-N²-ethyl-5-fluoropyrimidine-2,4-diamine,
N⁴-(6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro-N²,N²-dimethylpyrimidine-2,4-diamine,
N²-cyclopropyl-N⁴-(6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoropyrimidine-2,4-diamine
N⁴-(6,6-dimethyl-5-[(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro-N²-methylpyrimidine-2,4-diamine,

N^4 -(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl))-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)-5-fluoro- N^2 -isopropylpyrimidine-2,4-diamine ,
 N^4 -(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl))-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)- N^2 -ethylpyrimidine-2,4-diamine,
 N^4 -(6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl))-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)- N^2,N^2 -dimethylpyrimidine-2,4-diamine,
5-([(8S)-6,8-dimethyl-6,9-diazaspiro[4.5]dec-9-yl]carbonyl)- N -(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
 N^4 -(5-([(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl]carbonyl))-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)- N^2 -ethyl-5-fluoropyrimidine-2,4-diamine,
 N^4 -(5-([(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl]carbonyl))-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)- N^2 -ethyl-5-fluoropyrimidine-2,4-diamine,
 N^2 -ethyl-5-fluoro- N^4 -(5-([(2S,5R)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl))-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)pyrimidine-2,4-diamine acetate salt,
 N^4 -(6,6-dimethyl-5-([(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl))-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl)- N^2 -ethyl-5-fluoropyrimidine-2,4-diamine,
4-[(6,6-dimethyl-5-([(2S,5R)-2,4,5-trimethylpiperazin-1-yl]carbonyl))-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl]amino]pyrimidine-2-carbonitrile,
 N -(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-([(2S)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl))-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
 N -(2-ethyl-5-fluoropyrimidin-4-yl)-5-([(2S,5R)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl))-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
2-((5S)-4-[(3-[(2-ethyl-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl]carbonyl))-1,5-dimethylpiperazin-2-yl)ethanol,
5-([(2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl]carbonyl)- N -(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,

N-(5-fluoro-2-methylpyrimidin-4-yl)-6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
N-(5-fluoro-2-propylpyrimidin-4-yl)-6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
N-(5-fluoro-2-isopropylpyrimidin-4-yl)-6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-5-(((2S,5R)-2,4,5-trimethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
5-(((3S,8aS)-3,8a-dimethylhexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)carbonyl)-N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-5-(((3S,8aS)-3-methylhexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
N-(3-fluoro-6-methylpyridin-2-yl)-6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
N-[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]-6,6-dimethyl-5-(((2S)-2,4,5,5-tetramethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
5-(((2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl)carbonyl)-N-(2-ethyl-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
5-(((2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl)carbonyl)-N-(4-methoxypyrimidin-2-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
5-(((2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl)carbonyl)-6,6-dimethyl-N-(4-methylpyrimidin-2-yl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
5-(((2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl)carbonyl)-6,6-dimethyl-N-[4-(trifluoromethyl)pyrimidin-2-yl]-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
5-(((2S,5R)-2,5-dimethyl-4-(tetrahydro-2H-pyran-4-ylmethyl)piperazin-1-yl)carbonyl)-6,6-dimethyl-N-(4-methylpyrimidin-2-yl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,
N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-(((2S,5R)-2,4,5-trimethylpiperazin-1-yl)carbonyl)-1,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-amine,

N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[4-ethyl(2*S*,5*R*)-2,5-dimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
N-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-5-[[[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
N-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-[[[(2*S*,5*R*)-4-(2-methoxyethyl)-2,5-dimethylpiperazin-1-yl]carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
N-(2-ethoxy-5-fluoropyrimidin-4-yl)-5-[[[(2*S*,5*R*)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
N-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-5-[[[(2*S*,5*R*)-4-(3-methoxypropyl)-2,5-dimethylpiperazin-1-yl]carbonyl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
N-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-6,6-dimethyl-5-[[[(2*S*,5*R*)-2,4,5-trimethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
N-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-6,6-dimethyl-5-[[[(2*S*)-2,4,5,5-tetramethylpiperazin-1-yl]carbonyl]-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
5-[[[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazin-1-yl]carbonyl]-*N*-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
5-[[[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-yl)piperazin-1-yl]carbonyl]-*N*-(2-ethoxy-5-fluoropyrimidin-4-yl)-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
2-((5*S*)-4-{[3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl}-1,5-dimethylpiperazin-2-yl)ethanol ,
2-((5*S*)-4-{[3-[(2-ethoxy-5-fluoropyrimidin-4-yl)amino]-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl}-1,5-dimethylpiperazin-2-yl)ethanol ,
5-[(4-fluoro-1-methylpiperidin-4-yl)carbonyl]-*N*-[5-fluoro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,
5-[[[(2*S*,5*R*)-2,5-dimethyl-4-(tetrahydro-2*H*-pyran-4-ylmethyl)piperazin-1-yl]carbonyl]-*N*-[5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]-6,6-dimethyl-1,4,5,6-tetrahydropyrrolo[3,4-*c*]pyrazol-3-amine,

2-((5*S*)-4-([3-([5-fluoro-2-(methoxymethyl)pyrimidin-4-yl]amino)-6,6-dimethyl-4,6-dihydropyrrolo[3,4-*c*]pyrazol-5(1*H*)-yl]carbonyl)-1,5-dimethylpiperazin-2-yl)ethanol; or a pharmaceutically acceptable salt thereof.

22. A pharmaceutical composition comprising an effective amount of a compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

23. A compound according to any of the preceding claims, or a pharmaceutically acceptable salt thereof for use in the treatment of diabetes mellitus and its complications, cancer, ischemia, inflammation, central nervous system disorders, cardiovascular disease, Alzheimer's disease and dermatological disease, virus diseases, inflammatory disorders, or diseases in which the liver is a target organ.

24. A compound or a pharmaceutically acceptable salt thereof, according to any of the preceding claims for use in the treatment of diabetic mellitus and its complications.

25. A compound or a pharmaceutically acceptable salt thereof, according to any of the preceding claims for use in the treatment of diabetic mellitus and its complications, wherein the complications comprise diabetic retinopathy, nephropathy and neuropathy.