


1/9

FIGURE 1


HCT-116 PD study: I-115

PK Parameters of K-I-115 in Female Ncr Nude Mice
Bearing HCT 116 Tumor Xenografts
Following a Single Subcutaneous Administration at 30
mg/kg.

Sample	AUC last (nM*hr)	Cmax (nM)
Plasma	26271	27138

2/9

FIGURE 2


HCT-116 PD study: 10 mg/kg I-216

PK Parameters of I-216-HCl in Female Ncr Nude Mice Bearing HCT 116
Tumor Xenografts
Following a Single Subcutaneous Administration at 10 mg/kg.

Sample	AUC last (nM*hr)	Cmax (nM)
Plasma	24523	10413

3/9

FIGURE 3


HCT-116 PD study: 30 mg/kg I-216

PK Parameters of I-216·HCl in Female Ncr Nude Mice Bearing HCT 116
Tumor Xenografts
Following a Single Subcutaneous Administration at 30 mg/kg.

Sample	AUC _{last} (nM*hr)	C _{max} (nM)
Plasma	54800	24200


4/9

FIGURE 4



5/9


FIGURE 5



6/9


FIGURE 6

TGA




7/9

FIGURE 7



8/9

FIGURE 8



9/9

FIGURE 9

