


**FIG. 1A**




FIG. 1B

Hyaluronic Acid

DMEM/10%FCS

RD-CLMSC-CM




FIG. 2A





FIG. 2B



	Average total cell counts	Standard deviation	% increase over control
DMEM/10% FCS (Control)	1079	45	
FSF-CM	1248	58	16%
H-CLMSC-CM	2293	345	112%
RD-CLMSC-CM	2301	344	113%

FIG. 3




FIG. 4




FIG. 5



FIG. 6A



FIG. 6A (continued)



FIG. 6A (continued)




FIG. 6A (continued)




FIG.6A (continued)

BEFORE



AFTER 3 WEEKS



AFTER 6 WEEKS



FIG. 6B

BEFORE



AFTER



FIG. 6B (continued)




FIG. 6C




FIG. 6C (continued)




FIG. 6C (continued)



FIG. 6C (continued)




FIG. 6C (continued)




FIG. 6C (continued)



FIG. 6C (continued)



FIG. 6C (continued)




FIG. 6C (continued)




FIG. 6C (continued)





FIG. 6C (continued)

BEFORE



AFTER



FIG. 6D

BEFORE



AFTER



FIG. 6E

BEFORE



5 MONTHS AFTER




FIG. 6F



FIG. 6G



FIG. 6G (continued)



FIG. 6G (continued)



FIG. 6G (continued)




FIG. 6G (continued)



FIG. 6G (continued)



FIG. 6G (continued)



FIG. 6G (continued)



FIG. 6G (continued)



FIG. 6G (continued)



FIG. 6G (continued)



FIG. 6G (continued)



FIG. 6G (continued)



FIG. 6G (continued)



FIG. 7




FIG. 8





FIG. 8 (continued)



S/No	Sample ID	Conc of TNF-alpha (pg/ml)
1	CALECIM 2 x Dil with UV Induced NHEK	16.19
2	CALECIM 4 x Dil with UV Induced NHEK	42.69
3	CALECIM 8 x Dil with UV Induced NHEK	116.15
4	CALECIM 16 x Dil with UV Induced NHEK	281.92
5	CALECIM 32 x Dil with UV Induced NHEK	285.77
6	CALECIM 64 x Dil with UV Induced NHEK	346.15
7	Vitamin D (Experimental Positive Control)	30.62
8	Non-UV Induced NHEK (Baseline Control)	1.62
9	UV-Induced NHEK (no CALECIM added)	485.13

FIG. 9

Fig. 10

Lonza

www.lonza.com
 U.S. Scientific Support: 800-521-0390
 scientific.support@lonza.com
 EU/ROW Scientific Support: +49-221-99199-400
 scientific.support.eu@lonza.com
 Document # TS-12-604-3 08/11
 © 2011 Lonza Walkersville, Inc.

Dulbecco's modified eagle medium (DMEM)

Product use

Dulbecco's modified eagle medium was developed in 1969 and is a modification of basal medium eagle (BME) that differs from BME and MEM by the following characteristics:

- Vitamins 4X greater than MEM. Vitamins and amino acids greater than BME
- Types and quantities of amino acids greater than MEM and BME
- Iron (ferric nitrate)

Description

- 12-604 With 4.5 g/L glucose, with L-glutamine
 12-614 With 4.5 g/L glucose, without L-glutamine
 12-707 With 1.0 g/L glucose, without L-glutamine
 12-708 With 1.0 g/L glucose, and 25 mM HEPES buffer, without L-glutamine
 12-709 With 4.5 g/L glucose, and 25 mM HEPES buffer, without L-glutamine
 12-733 With 4.5 g/L glucose, without L-glutamine, without sodium pyruvate
 12-741 With 4.5 g/L glucose, with L-glutamine, without sodium pyruvate
 12-914 With 4.5 g/L glucose, without L-glutamine, screened to support hybridoma growth
 12-917 With 4.5 g/L glucose, without L-glutamine or phenol red
 15-604 With 4.5 g/L glucose, with L-glutamine, without sodium pyruvate, powder
 15-614 With 4.5 g/L glucose, without L-glutamine or sodium pyruvate, powder
 (Powder formulations require the addition of 49.3 mL of NaHCO₃ 7.5% solution or 3.70 g/L of NaHCO₃ powder)

Quick reference chart

	Glucose	L-glutamine	Phenol red	HEPES buffer	Sodium pyruvate
12-604	4.5 g/L	+	+	-	+
12-614	4.5 g/L	-	+	-	+
12-707	1.0 g/L	-	+	-	+
12-708	1.0 g/L	-	+	+	+
12-709	4.5 g/L	-	+	+	+
12-733	4.5 g/L	-	+	-	-
12-741	4.5 g/L	+	+	-	-
12-914	4.5 g/L	-	+	-	+
12-917	4.5 g/L	-	-	-	+

Fig. 10 continued

Lonza**Specifications**

	Sterility	pH	Osmolality (mOsm)	Cell growth promotion (% of control)	Endotoxin (EU/ml)
12-604	Neg.	7.0-7.4	324-352	≥75%	FIO
12-614	Neg.	7.0-7.4	324-352	≥75%	FIO
12-707	Neg.	7.0-7.4	306-346	≥75%	FIO
12-708	Neg.	7.03- 7.27	300-326	≥75%	FIO
12-709	Neg.	7.0-7.4	321-351	≥75%	FIO
12-733	Neg.	7.0-7.4	318-360	≥75%	FIO
12-741	Neg.	7.0-7.4	318-360	≥75%	FIO
12-914	Neg.	7.0-7.4	324-352	≥65% **	FIO
12-917	Neg.	7.0-7.4	329-343	≥75%	FIO
15-604 ***	-	5.5-7.0	235-265	≥75%	≤1.0
15-614 ***	-	5.5-7.0	235-265	≥75%	≤1.0

FIO – for information only
 ** Result of functional test
 *** Moisture content #1.0

Storage

2°C to 8°C

Product use statement

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. Not approved for human or veterinary use, for application to humans or animals, or for use in clinical or *in vitro* procedures.

Ordering information

Catalog number	Description	Size
12-604F	DMEM with 4.5 g/L glucose, with L-glutamine	500 ml
12-604Q	DMEM with 4.5 g/L glucose, with L-glutamine	1 L
12-614F	DMEM with 4.5 g/L glucose, without L-glutamine	500 ml
12-614Q	DMEM with 4.5 g/L glucose, without L-glutamine	1 L
12-707F	DMEM with 1.0 g/L glucose, without L-glutamine	500 ml
12-708F	DMEM with 1.0 g/L glucose, and 25 mM HEPES buffer, without L-glutamine	500 ml
12-708F	DMEM with 4.5 g/L glucose, and 25 mM HEPES buffer, without L-glutamine	500 ml
12-733F	DMEM with 4.5 g/L glucose, without L-glutamine, without sodium pyruvate	500 ml
12-733Q	DMEM with 4.5 g/L glucose, without L-glutamine, without sodium pyruvate	1 L
12-741F	DMEM with 4.5 g/L glucose, with L-glutamine, without sodium pyruvate	500 ml
12-914F	DMEM with 4.5 g/L glucose, without L-glutamine, screened to support hybridoma growth.	500 ml
12-917F	DMEM with 4.5 g/L glucose, without L-glutamine or phenol red.	500 ml
15-604D	DMEM with 4.5 g/L glucose, with L-glutamine, without sodium pyruvate, powder	1 x 10L
15-604F	DMEM with 4.5 g/L glucose, without L-glutamine or sodium pyruvate, powder	1 X 50L
15-614D	DMEM with 4.5 g/L glucose, without L-glutamine or sodium pyruvate, powder	1 x 10L

Fig. 11



Lonza Walkersville, Inc.
www.lonza.com
biotechserv@lonza.com
Tech Service: 800-521-0380
Document # TS 12-615-2 11/08
Walkersville, MD 21793-0127 USA
© 2008 Lonza Walkersville, Inc.

Ham’s F12 Medium

Product Use

Ham's F12 Medium is a nutrient mixture designed to cultivate a wide variety of mammalian and hybridoma cells when used with serum in combination with hormones and transferrin.

Specifications

		Osmolality (mOsm)
Sterility	pH	
Negative	7.07-7.40	286-305
Cell Growth Generation	Endotoxin (EU/ml)	
≥75% of control	FIO	

Storage
2°C to 8°C

Product Use Statement

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. Not approved for human or veterinary use, for application to humans or animals, or for use in clinical or *in vitro* procedures.

Ordering Information

Catalog Number	Description	Size
12-615F	Ham's F12 Medium with L- glutamine	500 ml

Fig. 12



www.lonza.com
U.S. Scientific Support: 800-521-0390
scientific.support@lonza.com
EU/ROW Scientific Support: +49-221-99199-400
scientific.support.eu@lonza.com
Document # TS-12-719-3 08/11
© 2011 Lonza Walkersville, Inc.

DMEM: F12 (1:1) medium

Product use

DMEM combined with Hams F12 has been extensively used to demonstrate the effect of various hormones and growth factors on target tissues.

Description

- 12-719 with L-glutamine, 15 mM HEPES, and 3.151 g/L glucose
- 15-719 powder – with L-glutamine, 15 mM HEPES, and 3.151 g/L glucose. Powdered medium requires the addition of 16.0 ml/L of NaHCO₃, 7.5% solution or 1.2 g/L NaHCO₃ powder
- 04-687 with 3.151 g/L glucose, with L-glutamine, without HEPES

Specifications

	Sterility	pH	Osmolality (mOsm)
12-719	Negative	7.0-7.4	286-356
15-719*	Negative	4.5-6.5	260-290
04-687	Negative	FIO	FIO
	Cell growth promotion	Endotoxin (EU/ml)	Moisture
12-719	≥ 75% of control	FIO	--
15-719*	≥ 75% of control	≤ 1 EU/ml	≤ 2%
04-687	--	FIO	--

* pH, osmolality, endotoxin, and moisture are done without NaHCO₃; NaHCO₃ is added for cell growth test.

Storage

2°C to 8°C

Product use statement

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. Not approved for human or veterinary use, for application to humans or animals, or for use in clinical or *in vitro* procedures.

Ordering information

Catalog number	Description	Size
12-719F	DMEM: F12 with L-glutamine, 15 mM HEPES, and 3.151 g/L glucose	500 ml
12-719Q	DMEM: F12 with L-glutamine, 15 mM HEPES, and 3.151 g/L glucose	1 L
15-719D	DMEM: F12 powder – with L-glutamine, 15 mM HEPES, and 3.151 g/L glucose. Powdered medium requires the addition of 16.0 ml/L of NaHCO ₃ , 7.5% solution or 1.2 g/L NaHCO ₃ powder	1 x 10L
04-687Q	DMEM: F12 with 3.151 g/L glucose, with L-glutamine, without HEPES	1 L

Fig. 13

Cascade Biologics™
invitrogen cell culture



Medium 171, Medium 171PRF, and MEGS

Medium 171

M-171-500

500 ml

Medium 171PRF (Phenol Red-Free)

M-171PRF-500

500 ml

Product Description

Medium 171 and Medium 171PRF are sterile, liquid tissue culture media intended for use as one component in a complete culture environment for the growth of normal human mammary epithelial cells. Medium 171 is a basal medium containing essential and non-essential amino acids, vitamins, other organic compounds, trace minerals, and inorganic salts. Medium 171PRF is a phenol red-free version of Medium 171. These media do not contain antibiotics, antimycotics, hormones, growth factors, or proteins. These media are HEPES and bicarbonate buffered and are designed for use in an incubator with an atmosphere of 5% CO₂/95% air. To support plating and long-term proliferation of normal human mammary epithelial cells, these media must be supplemented with Mammary Epithelial Growth Supplement (MEGS, cat. no S-015-5).

Intended Use

Medium 171 is intended for use in the routine culture of normal human mammary epithelial cells. Medium 171PRF is intended for use by investigators who wish to culture normal human mammary epithelial cells in the absence of phenol red. When supplemented with MEGS, these media will support the plating and proliferation of normal human mammary epithelial cells at densities between 2.5×10^3 cells/cm² and 8×10^4 cells/cm². **This product is for research use only. Not for use in animals, humans, or diagnostic procedures.**

Caution: If handled improperly, some components of this product may present a health hazard. Take appropriate precautions when handling this product, including the wearing of protective clothing and eyewear. Dispose of properly.

Storage and Stability

Medium 171 and Medium 171PRF are stored at 4° C in our facility and are shipped at ambient temperature. Upon receipt, these media should be stored at 4° C and should not be frozen. **Protect from light.** Several components of these tissue culture media are light-labile, and we recommend that the media not be exposed to light for lengthy periods of time. If the media are warmed prior to use, do not exceed 37° C. When stored in the dark at 4° C, the product is stable until the expiration date on the label.

Preparation of Supplemented Medium 171

1. Thaw one bottle of MEGS. Take one bottle of Medium from cold storage. Make sure that the caps of the vessels are tight.
2. Gently swirl the bottle of supplement. Avoid splashing the supplement into the cap of the bottle or causing the supplement to foam.
3. Wipe the outside of the containers with a disinfecting solution such as 70% ethanol or isopropanol.
4. Using sterile technique in a laminar flow culture hood, transfer the entire contents of the bottle of supplement to the bottle of Medium.
5. Tightly cap the bottle of supplemented medium and swirl the contents to ensure a homogeneous solution. Avoid causing the medium to foam.

Storage and Stability of Supplemented Medium 171

Once Medium 171 or Medium 171PRF has been supplemented with MEGS, the supplemented medium should be stored in the dark at 4° C and should not be frozen. When stored in the dark at 4° C, the supplemented medium is stable for 1 month.

Selected References

The Medium 171 formulation is based on medium MCDB 170, with modifications.

Hammond SL, Ham RG, Stampfer MR; PNAS 81:5435-5439, 1984

For research use only.

Life Technologies Corporation • 5791 Van Allen Way • Carlsbad • CA 92008 • Tel: 800.855.6286 • www.invitrogen.com • E-mail: tech_support@invitrogen.com

MAN0001585

Revised: 30 May 2009

Fig. 13 continued

Cascade Biologics™
invitrogen cell culture



MEGS Mammary Epithelial Growth Supplement

Cat. no. S-015-5
5 ml

Product Description

Mammary Epithelial Growth Supplement (MEGS) is a sterile, concentrated (100X) solution intended for use as one component in a complete culture environment for the growth of normal human mammary epithelial cells. Each 5 ml bottle of MEGS contains all of the growth factors, hormones, and tissue extracts necessary for the culture of normal human mammary epithelial cells and is the correct amount of supplement for a 500 ml bottle of Medium 171 or Medium 171PRF. MEGS is an ionically-balanced supplement containing bovine pituitary extract (BPE), bovine insulin, hydrocortisone and recombinant human epidermal growth factor. When a 500 ml bottle of Medium 171 or Medium 171PRF is supplemented with MEGS, the final concentrations of the components in the supplemented medium are: BPE, 0.4% v/v; bovine insulin, 5 µg/ml; hydrocortisone, 0.5 µg/ml; and recombinant human epidermal growth factor, 3 ng/ml.

Intended Use

MEGS is intended for use in conjunction with Medium 171 or Medium 171PRF for the routine serum-free culture of normal human mammary epithelial cells. ***This product is for research use only. Not for use in animals, humans, or diagnostic procedures.***

Caution: If handled improperly, some components of this product may present a health hazard. Take appropriate precautions when handling this product, including the wearing of protective clothing and eyewear. Dispose of properly.

Limited Use Label License No. 5: Invitrogen Technology

The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. The buyer may transfer information or materials made through the use of this product to a scientific collaborator, provided that such transfer is not for any Commercial Purpose, and that such collaborator agrees in writing (a) not to transfer such materials to any third party, and (b) to use such transferred materials and/or information solely for research and not for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For products that are subject to multiple limited use label licenses, the terms of the most restrictive limited use label license shall control. Life Technologies Corporation will not assert a claim against the buyer of infringement of patents owned or controlled by Life Technologies Corporation which cover this product based upon the manufacture, use or sale of a therapeutic, clinical diagnostic, vaccine or prophylactic product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. If the purchaser is not willing to accept the limitations of this limited use statement, Life Technologies is willing to accept return of the product with a full refund. For information on purchasing a license to this product for purposes other than research, contact Licensing Department, Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, California 92008. Phone (760) 603-7200. Fax (760) 602-6500. Email: outlicensing@invitrogen.com.

©2009 Life Technologies Corporation. All rights reserved.

For research use only. Not intended for any animal or human therapeutic or diagnostic use.

For research use only.

Life Technologies Corporation • 5791 Van Allen Way • Carlsbad • CA 92008 • Tel: 800 855 6286 • www.invitrogen.com • E-mail: tech_support@invitrogen.com

MAN0001585

Revised: 30 May 2009

Fig. 14***PTT6 Medium ingredients list***

Medium composition List	Company Name	Catalogue Number
-------------------------	--------------	------------------

Basic media

DMEM (also referred to herein as PTT6-basal medium)	Lonza	12-604F
DMEM/F12	Lonza	12-719F
M171	Life technologies	M171500

Serum

Fetal Bovine Serum	GE Healthcare	A15-151
--------------------	---------------	---------

Antibiotic

Penicillin-Streptomycin-Amphotericin B	Lonza	17-745E
--	-------	---------

Supplements

Adenine (optional)	Sigma	A8626-25G
Hydrocortisone (optional)	Sigma	H-0888
Epidermal Growth Factor	Millipore	GF-144
T3 (3,3',5-Triiodo-L-thyronine sodium salt) (optional)	Sigma	200-223-5
Recombinant Human Insulin AOF	Life Technologies	A11382U

Fig. 15




Fig. 16




Fig. 17A




Fig. 17B




Fig. 17C




Fig. 17D




Fig. 18A




Fig. 18B





Fig. 18C



Week	wk.1		wk.4		wk.6		wk.7		wk.10		wk.12		p-value
Values	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	
Interocular distance (mm)	129	0.37	129	0.21	128	0.29	124	0.26	115	0.14	114	0.16	<0.05
TV Ratio	2485	2724	2008	2158	2185	2388	1509	2477	2327	3110	2584	3154	n.s.
Follicular Units (FU) per sqmm	9288	3839	8561	2565	8878	4385	9188	3544	8312	2845	8868	3128	<0.05
Average hairs per FU	123	0.11	124	0.13	122	0.10	124	0.14	125	0.10	122	0.10	n.s.
Average hair width (micron)	5435	811	5416	875	5823	848	5419	858	5520	882	5538	749	n.s.
Terminal hair count per sqmm	80738	4887	9478	3907	10044	5255	10218	4512	11885	4824	12109	4455	<0.05
Total hair count per sqmm TOTAL	11500	4917	10450	3869	10382	5288	11488	4638	12935	3842	13162	4885	<0.05
Sum of terminal hair width (micron) per sqmm	821132	312785	548012	244103	594538	331071	528865	281705	890044	264275	717608	298296	<0.05
Sum of total hair width (micron) per sqmm	648283	308274	573271	238707	615888	327753	619485	280032	720250	248978	743109	287395	<0.05

Fig. 19 (Table 6)

Measure	wk 1		wk 4		wk 6		wk 7		wk 10		wk 12		p-value
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	
Subsistence with basic needs	1.4	0.7	1.7	0.82	1.6	0.82	2.5	0.71	2.4	0.7	2.4	0.84	<.001
Subsistence with basic security	1.2	0.62	1.4	0.78	1.3	0.67	2.3	0.57	2.2	0.42	2.3	0.67	<.001
Subsistence with basic shelter	1.8	0.79	1.9	0.74	1.6	0.62	2.9	0.46	2.5	0.46	2.3	0.67	<.05
Subsistence with basic quality	2.6	0.97	2.5	0.85	2.7	0.82	2.7	0.67	2.6	0.84	2.8	0.79	n.s.
Subsistence with health	.	.	2.22	0.97	1.6	0.82	2	0.67	2.8	0.83	2.9	0.57	n.s.
Confidence	1.8	0.79	1.8	0.79	2.7	0.67	2.2	0.66	2.3	0.46	2.7	0.67	<.001
Friends & family not making basic quality	2.5	0.53	2.5	0.53	2.9	0.57	2.7	0.46	2.7	0.46	2.9	0.52	n.s.
Thinking of how basic looks like	2.5	0.97	2.4	1.03	2.4	0.97	2.4	1.17	2.2	1.04	2.3	0.88	n.s.
Feeling basic	3.1	0.74	3.1	0.74	2.9	0.57	2.8	0.63	2.9	0.74	2.9	0.74	n.s.
How safe basic	3.2	1.03	3.2	1.14	2.4	1.07	2.3	0.96	1.9	0.99	1.9	0.99	<.01
Relatives quality of life	3.2	1.03	3.6	0.97	2.5	1.18	2.3	1.06	2.3	1.1	2.5	1.08	n.s.
Relatives quality of life	3.2	1.03	3.1	0.98	1.9	0.88	2.3	0.96	1.9	0.74	1.8	0.79	<.001

Fig. 20 (Table 7)