

HIGH-AFFINITY AND AGGREGATIVELY STABLE ANTIBODIES ON THE BASIS OF VARIABLE DOMAINS VL AND A DERIVATIVE VHH

Introduction

Antibodies, also referred to as immunoglobulins (Ig), are soluble blood or interstitial fluid glycoproteins that play a key role in humoral immunity of the Vertebrata. Antibodies are produced by B-cells in response to alien biological and chemical substances (antigens) of various structure for the purpose of their neutralization. Due to high specificity and high affinity of binding with a certain antigen, and ability to produce antibodies against the unlimited antigen repertoire, antibodies and their derivatives are one of the most important reagents to be used in both fundamental and applied medical researches.

Classical antibodies [1, 2] are large multimeric proteins (IgG ~150 kDa), which comprise two identical heavy H-chains (one variable VH domain, three constant domains CH1, CH2 and CH3, and a hinged region between CH1 and CH2 domains) and two identical light L-chains (consisting of the variable domain VL and constant domain CL). A four-chain molecule has non-covalent and covalent (disulfide) bonds connecting the chains. Papain protease can be used to break down an antibody molecule into two fragments: Fab (Fragment antigen binding) and Fc (Fragment crystallizable). Therefore, one region of the molecule (Fab) defines its antigen-related specificity, and another region (Fc) exercises the effector functions targeted at antigen elimination [3, 4]. CH1 and CH2 domains of H-chain are divided by a hinge region that assures mobility of Fab-region and interaction of IgG molecule with Ig effector receptors located on the cells. CH2 domain contains the regions binding both FcY receptors that mediate cell activation (ADCC and ADCP) and complement system molecules (CDC). In addition, this domain contains the site that is an attach point for carbohydrates of all immunoglobulin isotypes. CH3-domain largely determines the stability of IgG dimer and interacts with FcRn-receptor on the cell surface establishing pharmacokinetic properties of antibodies as well as their metabolism and distribution in the body. The

combination of complementarity determining regions (CDR) of the variable domain of the heavy chain (VH) and the variable domain of the light chain (VL) forms an antigen-binding fragment, while the framework regions of the variable domains and the constant domains are not directly involved in antigen recognition. A minimized Fab-derivative for classical antibodies is a single-chain structure in which variable domains of the heavy and light chains are connected with a linker sequence (scFv).

Finding significant amounts of specific non-classical antibodies of simplified structure in the blood of Camelidae animals (camels, llamas, vicunas) was a valuable discovery [5]. Such antibodies (heavy chain antibody, HCAb) consist of a dimeric single shortened heavy chain (without CH1 domain) with no light chain at all. Antigen-binding fragment of HCAb is formed by only one heavy chain variable domain (VHH), which is connected through the hinged region to Fc-domain. VHH is often called a single-domain antibody, “nanobody”, “mini-antibody” or “nano-antibody”. It appeared that in addition to small size (12-15 kDa), such isolated mono-domain structure has a number of advantages as compared to classical IgG antibodies, namely aggregation, chemical and thermal stability. VHH antibodies can be successfully cloned and expressed in bacterial and yeast cells. With such properties, these antibodies were developed in therapeutic field by Ablynx Company and in laboratory and industrial chromatography (CaptureSelect affinity products).

Non-classical antibodies (HCAb) including a dimer of just a single Ig heavy chain were first discovered by electrophoretic analysis of immunoglobulins in the serum obtained from various representatives of Camelidae bloodline [5]. The relative fraction of HCAb varies from about 15–25% (of all IgG) in llamas and vicunas to about 60–80% in camels [6].

It is assumed that non-classical antibodies (HCAb), at least in case of Camelidae, resulted from relatively recent evolution of the genes of classical antibodies. Two heavy chain constant domains, CH2 and CH3, in case of HCAb and classical antibodies are highly conserved. In HCAb there is no domain

corresponding to the first constant domain CH1 of classical antibodies. The genome of the one-humped camel (Dromedary) contains a cluster of about fifty VH- and forty VHH-generative genes followed by multiple genes of D-segments, J-segments and genes encoding the constant regions ($C\mu$, $C\gamma$, $C\epsilon$ and $C\alpha$). It is clear that some of $C\gamma$ -genes serve to form HCAb (mutations result in the loss of CH1-domain), and other serve to form classical antibodies (with sustained CH1-domain). The same genes of D- and J-segments may randomly connect to either one of VH-genes or one of VHH-genes. This indicates that VH- and VHH-genes are located in the same gene locus [7-10].

The organization of variable domains of non-classical antibodies (VHH) and variable domains of classical antibodies (VH) is very similar, as human VH-domains of IgG3 subclass have most evident homology to VH and VHH of Camelidae. In both cases, V-domains comprise four conservative framework regions FR surrounding three hypervariable complementary-determining regions (CDR). In addition, in both cases a 3-D structure typical of immunoglobulin V-domain of two β -layers is formed, one of which comprises 4 amino acid sequences and another comprises 5 amino acid sequences [11, 12]. All three hypervariable regions in this structure form a cluster on one side of V-domain, where they participate in antigen recognition and are located in the loops connecting β -structures. However, there are several important distinctions related to the functioning of single-domain VHH. Thus, CDR1 and CDR3 of VHH are significantly enlarged. Complementary-determining regions of VHH often contain cysteine residues in two sections at once (usually in CDR1 and CDR3, less frequently – in CDR2 and CDR3). The studies of VHH crystal structures have shown that these cysteine residues form disulfide bonds and provide additional stability to the loop structure of these antibodies [12]. The strongest and the most reproducible distinguishing feature of VHH is represented by four substitutes of hydrophobic amino acid residues with hydrophilic ones in the second framework region (Val37Phe, Gly44Glu, Leu45Arg, Trp47Gly, according to the numbering of Kabat et al [13]). This framework region of VH-domain is highly conservative,

enriched with hydrophobic amino acid residues and is essential for linking to the variable domain of the light chain VL. VHH-domain differs greatly in this aspect: substitutes of hydrophobic amino acids with hydrophilic makes the association of VHH and VL impossible. These substitutes also explain high solubility of VHH (nano-antibody), when it is obtained as a recombinant protein [14].

It appears that the repertoires of possible paratopes (antigen-binding parts of an antibody) of HCAb and classical antibodies may be significantly different. Since these two antibody types co-exist in the same organism, it can be assumed that they do not compete but are complementary to each other. For example, it was repeatedly noted that both antibody types could occur simultaneously, exclusively or in different ratios with regard to various epitopes of the antigen material upon immunization of the very same animal. Despite the suspected lower variety of paratopes possible for single-domain antibodies, as compared to classical two-domain antibodies, many publications have clearly demonstrated that HCAb can be obtained against the most diverse epitopes of a rather wide range of antigens [15]. Apparently, this is due to enlarged CDR1 and CDR3 regions. We also should note a surprisingly large (as compared to V-domains of classical antibodies) number of somatic hypermutations in VHH that are likely to accumulate in the course of affine maturation of the antibody during immunization [16]. X-ray diffraction analysis revealed that antigen-binding loop regions of VHH are able to form structures unusual for classical V-domains [12, 16]. While in case of VH- and VL-domains of classical antibodies, all six CDRs contribute almost the same to antigen binding, in case of VHH, CDR3 is usually the most important for the formation of a paratope. It has been shown that CDR3 in VHH (but not in VH or VL) is capable of forming uncommonly long finger-structures that can deepen into the antigen structure and, in particular, detect active enzyme centers [12]. Small size of the antigen-binding region of VHH and its ability to form unusual emerging paratopes explain how HCAb capable of epitopes inaccessible for classical antibodies can be obtained (for example, production of antibodies that efficiently inhibit enzymes) [17].

With all its high potential of specificity, unique, as compared to classical IgG antibodies, the therapeutic use of a single-domain VHH is sometimes limited due to its rapid elimination from the organism. There are several solutions designed to improve the pharmacokinetics of VHH structures, including chemical conjugation with PEG and covalent binding to polypeptides mediating the reduced clearance from the blood, such as Human serum albumin (HSA) or Fc-fragment classical human antibody as fusion proteins that possess a half-life of up to three weeks [18,19,20]. Small peptides attached to VHH by genetic engineering methods and capable of high-affinity non-covalent interaction with said components (HSA and IgG) in human blood have been successfully used [21]. However, the technological efficiency and immunogenicity of these approaches remain questionable, and the feasibility of using thereof in either clinical or earlier study phases is now on trial.

In addition, the greatest limitation of using antibodies as medicinal agents for the treatment of various diseases is due to their aggregation and chemical stability, affinity and immunogenicity. Since the majority of monoclonal antibodies are obtained on the basis of mice, the regular use of such antibodies in humans causes development of immune response to antibody therapy (for example, allergic reactions). These types of immune response finally result at least in lack of efficiency, and in potential anaphylactic reactions at the worst. On the other hand, aggregation or chemically unstable therapeutic antibodies reduce therapeutic properties of the drug product over time and may increase its immunogenicity upon administration to human patients.

In reference with the above, it is important to develop VHH-based antibodies with improved (in comparison to previously known antibodies) functional and therapeutic features, particularly increased aggregation, chemical and thermal stability and improved affinity, which would at the same time be easily and simply produced, including on industrial scale.

Background of the invention

The background of the invention includes various antibody structures containing VHH domain.

PCT/EP2008/066368 publication describes antibodies that comprise separate variable domains linked with Fc-fragment. Nano-bodies can be used as variable domains with Fc obtained from IgE antibodies. The said domain and Fc fragments can be connected through a linker located in the hinge region.

Patent application US 2009/0202979 disclosed antibodies comprising complete VHH antibodies or parts thereof directly connected to the constant regions of human antibodies.

In addition, there are amino acid substitutes known to affect the physical-chemical and biological properties of antibodies.

For example, application US 20110028348 describes the antibody heavy chain variable domains wherein amino acid substitutes were introduced in positions 35, 45, 47, 93- 100 and 100a to improve the hydrophilic properties of the antibody obtained.

Now, the methods have been developed to optimize the structure of isolated VHH and VH mono-domains in order to reduce the immunogenicity and improve the aggregation stability thereof.

Thus, Vincke at al. [22] have found that Glu-49→Gly and Arg-50→Leu substitutes in specific amino acids result in a separate domain that is more stable, though less soluble. Other substitutes in the framework region FR-2 Phe-42→Val and Gly/Ala-52→Trp are crucial for antibody affinity to the antigen due to re-orientation of H3-loop, raising dissociation constant 6-10-fold ($6.85 \cdot 10^{-3}$ 1/s). Phe-42→Val substitute caused the reduced stability of the antibodies obtained. The substitute of Gly-49 and Leu-50 in VH-sequence resulted in lower stability of the domain, while Glu-49 and Arg-50 humanization in VHH allows obtaining stable variable domains.

It is also known from literature, that in the presence of short HCDR3 regions neutralizing the shading effect of the conformation of classical VHH, and upon introduction of VH-characteristic Trp-47→Gly-47 substitute as well as Tyr-37→Val-37, Glu-44→Gly-44 and Arg-45→Leu-45, the isolated VHH domains can regenerate the ability to bind with VL domain [24].

The relationship between the increased aggregation stability of therapeutic antibodies of classical IgG structure and the reduced immunogenicity thereof was demonstrated in multiple studies and summarized in the review by Hermeling et al., 2004 [25]. Yet they did not reveal any antibodies comprising derivative VHH domains, but linked to the variable domains of the light chains within full-size human IgG.

Therefore, there is a need for development of a new format of antibodies that would have improved stability and affinity, good expression and low immunogenicity.

Besides, no approaches were earlier described with regard to the development of such molecules that would be easy to obtain, have improved aggregation stability, increased affinity and high expression level in the mammal cell culture.

With reference to the above, this invention is the first to describe antibodies comprising VHH-derivatives that are able to bind to variable domains of the light chains of the full-size human IgG, which results in the formation of a structure that is similar to the natural one (and, hence, having low immunogenicity), but has improved aggregation stability, increased affinity, and a structure of a therapeutic monoclonal antibody.

Figures

Figure 1. Scheme for development and optimization of an aggregation-resistant VHH-based antibody.

Figure 2, 3, 4. SDS gel electrophoresis of VHHIgG1 antibodies comprising various amino acid substitutes.

Figure 5. Scheme of the synthesis of llama combinatorial Fab libraries.

Figure 6. Phagemid for cloning of phage display Fab libraries.

Figure 7. Amino acid sequences for 3 different VHH mutants.

Figure 8. Sensograms for examining the comparative kinetic characteristics of VHHFab clones for binding with IL-17A.

Figure 9. Amino acid sequences for 3 different VHH mutants.

Figure 10, 11. Aggregation stability of three VHH mutants.

Figure 12, 13. Affinity of three VHH mutants.

Figure 14. Cell test for suppression of IL-6 production by inhibiting IL-17A with VHHIgG1VK4B11, mut1VHHIgG1 and mut4VHHIgG1 antagonists.

Figure 15, 16. Isolation of chimeric variants of mut4 VHHFab comprising human light chains.

Figure 17. Cell test for suppression of IL-6 production by inhibiting IL-17A with mut4 VHHIgG1VK3c8, mut4 VHHIgG1VK1A7 and mut4 VHHIgG1VK4E12 antagonists.

Figure 18, 19. Protein electrophoresis under denaturant conditions performed for products comprising antibodies with mutations in positions 44 and 45 of m4 VHHc8.

Figure 20. Comparative cell test for suppression of IL-6 production by inhibiting IL-17A with m4VHHc8 antagonists comprising mutations in FR2 at positions 44 and 45.

Figure 21. Diagram showing the stability and functional properties of different mutations at positions 44 and 45 in FR2 of m4VHHc8 that determine the interaction of the light and heavy chain variable domains.

Figure 22. Determination of kinetic parameters of BCD109 interaction with IL-17A of various origins.

Figure 23. Results for kinetic parameters of BCD109 interaction with IL17A of various origins.

Figure 24. Chromatogram of BCD109 antibodies obtained before and after the thermal stress.

Definitions

"Monoclonal antibody" as used herein relates to an antibody obtained from llama, chimeric antibody, humanized antibody or fully human antibody, unless otherwise is stated in the present application. Monoclonal antibodies, according to the invention, can be produced using, for example, recombinant technology, phage display technology, synthetic technology or combinations of these or other technologies well known from the prior art.

"Monoclonal antibody" refers to an antibody obtained from a single copy or a clone including, for example, any eukaryotic, prokaryotic or phage clone, rather than to production method thereof. "Monoclonal antibody" can be an intact antibody (with full or full-length Fc-region), actually intact antibody, an antibody part or fragment comprising an antigen-binding region, for example, Fab-fragment, Fab'-fragment or F(ab')₂-fragment from llama or chimeric, humanized or human antibody. "Fab"-fragment comprises a variable light chain domain and a constant light chain domain as well as a variable heavy chain domain and the first constant heavy chain domain (CH1). "F(ab')₂" antibody fragment contains a pair of Fab-fragments, which are mostly covalently bound by hinged cysteine residues at C-terminal regions. Other chemical bonds between antibody fragments are also well known from the prior art.

In addition, "monoclonal antibody" as used herein can be a single-chain Fv that can be obtained by binding DNA encoding VHH and VL with a linker sequence. As long as the protein keeps its ability of specific or preferable binding to the target (for example, epitope or antigen), it is covered by the term "antibody". Antibodies can be either glycosylated or not and are within the framework of the invention.

The term “derivative” or antibody “variant”, as used herein, refers to a molecule, the amino acid sequence of which differs from the parental sequence by adding, deletion and/or substitute of one or more amino acid residues in the sequence of parental antibody. In the preferred embodiment, an antibody contains at least one (for example, from one to about ten, preferably 2, 3, 4, 5, 6, 7 or 8) amino acid substitutes in FR- or CDR-regions of the parental antibody. This application defines the identity or homology regarding the sequence of a variant antibody as the percentage of amino acid residues in a variant antibody sequence that are identical to residues in parental antibody after aligning the sequences and, if needed, introducing gaps in order to achieve the maximum percentage identical sequence.

An antibody derivative (from parental one) keeps its ability to bind the same antigen or, preferably, epitope with which the parental antibody binds, or, preferably, exhibits at least one property or biological activity exceeding that of the parental antibody. For example, the antibody preferably has a better aggregation stability, more strong affinity, improved pharmacokinetics or increased ability to inhibit the antigen biological activity, as compared to parental antibody.

The term "VHH-derivative", as used herein, refers to the derivatives of VHH antibodies, the amino acid sequence of which differs from the sequence of parental VHH antibody by substitute of one or more amino acid residues in the sequence of parental antibody. In the preferred embodiment, VHH antibody contains at least one (for example, from one to about ten preferably 2, 3, 4, 5, 6, 7 or 8) amino acid substitutes in FR- or CDR-regions of the parental antibody.

An antibody derivative keeps its ability to bind the same antigen or, preferably, epitope with which the parental antibody binds, or, preferably, exhibits at least one property or biological activity exceeding that of the parental antibody. For example, the antibody preferably has a better aggregation stability, stronger affinity, improved pharmacokinetics or increased ability to inhibit the antigen biological activity, as compared to parental antibody.

“Parental VHH antibody” or “initial VHH antibody”, or “wild VHH antibody” as used herein refers to VHH antibody isolated from an immunized or non-immunized Camelidae animal encoded with amino acid sequence that is used to produce a VHH variant. Parental antibody can have a framework sequence originating from Camelidae (with respect to VHH variable domain), but preferably the frame sequence of the light chain variable domain is of completely or substantially human origin.

"Parental" or "initial", or "wild" antibody, as used herein, refers to an antibody encoded with amino acid sequence that is used to produce a variant. Parental antibody can have a framework sequence originating from Camelidae (with respect to VHH variable domain), but preferably the frame sequence of the light chain variable domain is of completely or substantially human origin.

As used herein, the term “specifically binds” refers to such a situation in which one party involved in the process of specific binding does not significantly bind molecules different from the ones of its specific binding partner (partners). This term also applies if, for example, an antigen-binding site of the antibody according to the invention is specific for particular epitope that is carried by a number of antigens; in this case, the specific antibody with an antigen-binding domain will be able to bind specifically with various epitope-carrying antigens. Thus, the monoclonal antibody, according to the invention, specifically binds to human IL-17 (IL-17A), while it does not specifically bind human IL-17B, IL-17C, IL-17D or IL-17E. Moreover, a monoclonal antibody of the invention specifically binds human IL-17 and IL-17 of cynomolgus monkey, but it does not specifically bind neither rat IL-17 nor murine IL-17.

As used herein, the term “preferably binds” refers to such a situation in which an antibody binds a specific antigen at least by 20% more, preferably by about 50%, or 2-fold, 20-fold, 50-fold or 100-fold more than it binds any other antigen, as measured according to the procedures known from the prior art (for example, competitive ELISA or K_D measurements obtained with Octet apparatus). Antibody can preferably bind one epitope within an antigen but not bind another

epitope of the same antigen. Thus, an antibody of the invention preferably binds human IL-17, but not rabbit IL-17.

As used herein, the term “epitope” refers to the molecule part that can be recognized by and bind an antibody via one or several antigen-binding sites of an antibody. Epitopes often comprise chemically surface-active groups of molecules such as amino acids or sugar side chains, and have specific 3-D structural characteristics. “Inhibiting epitope” and/or “neutralizing epitope” means an epitope that in the context of an intact antigen molecule and while binding with an antibody specific to the said epitope, causes *in vivo* or *in vitro* loss or reduction of activity of the molecule or organism that contains the molecule.

As used herein, the term “epitope” also refers to a polypeptide fragment, having antigenic and/or immunogenic activity in animals, preferably in mammals such as mice and humans. The term “antigenic epitope” as used herein is a polypeptide fragment which can specifically bind the antibody and can be detected by any technique well known from the prior art (for example, by means of standard immunoassay). Antigen epitopes are not necessary immunogenic, but they can possess immunogenicity. “Immunogenic epitope”, as used herein, is defined as a polypeptide fragment that evokes an antibody response in animals, as determined by any method of the prior art. “Nonlinear epitope” or “conformational epitope” contains nonadjacent polypeptides (amino acids) within the antigen protein, which binds with epitope-specific antibody.

The words "functional activity" or "functional characteristics" or the terms "biological activity" or "activity" referring to an antibody, according to the invention, are interchangeable as used herein, including, but not limited to: epitope/antigen affinity and specificity; ability to neutralize or to be an antagonist to IL-17 *in vivo* or *in vitro*; IC₅₀; antibody stability and *in vivo* immunogenicity of the antibody. Other biological properties or antibody characteristics identified from the prior art include, for example, cross-reactivity (i.e. reaction with non-human homologs of the target peptide or with other proteins or targets) and ability to retain high levels of protein expression in mammal cells. The aforementioned

properties or characteristics may be observed, measured or evaluated using procedures recognized in the prior art, including but not limited to ELISA, competitive ELISA, Octet analysis, neutralization assay *in vitro* or *in vivo* without limitation, receptor binding, production and/or release of cytokine or growth factor, signal transduction and immune histochemical study of tissue sections obtained from various sources including humans, primates or any other source.

The population of “monoclonal antibodies” as used herein refers to homogenous or essentially homogeneous antibody population (i.e. at least 85, 90, 91, 92, 93, 94, 95, or 96%, but more preferably no less than about 97 or 98%, or further preferably, at least 99% of antibodies in the population will compete for the same antigen/epitope in ELISA, or further preferably, antibodies are identical in their amino acid sequence).

A full-size antibody existing in nature is represented by immunoglobulin molecule comprising four polypeptide chains (two heavy full length H chains of about 50-70 kDa, and two light full length L chains of about 25 kDa) linked via disulfide bonds. Amino-terminal part of each chain comprises a variable domain of about 100-110 or more amino acids that are responsible for binding an antigen. Carboxyl-terminal domain of each chain determines the constant region that is mostly responsible for the effector function. Light chains are classified as kappa and lambda and have specific constant regions. Each light chain consists of a variable N-terminal light chain region (hereafter referred to as VL or VK) and a constant light chain region consisting of a single domain (CL or CK). Heavy chains are classified as γ , δ , α , μ and ϵ and determine antibody isotype, such as IgG, IgM, IgE, IgA and IgD respectively; some of them can be additionally divided into sub-classes (isotypes) such as IgG1, IgG2, IgG3, IgG4, IgA1. Each heavy chain type is characterized by specific constant region. Each heavy chain comprises a variable N-terminal region (hereafter referred to as VH) and a constant heavy chain region. Constant heavy chain region consists of three domains (CH1, CH2 and CH3) for IgG, IgD and IgA, and of 4 domains (CH1, CH2, CH3 and CH4) for IgM and IgE. Variable domains VH, VHH and VL can also be divided

into the so-called hypervariable regions (complementarity determining regions, CDR) interspersing with more conservative framework regions (FR). Each variable domain comprises three CDRs and FRs located in the following order from N-terminus to C-terminus: FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4.

In this application, 3 heavy chain CDRs are referred to as "HCDR1, HCDR2 and HCDR3", while 3 light chain CDR are referred to as "LCDR1, LCDR2 и LCDR3". CDRs contain the majority of amino acid residues specifically interacting with an antigen. CDR-residues are numbered and positioned in compliance with Kabat Numbering Scheme.

The term “antigen” refers to an antigen target against which an antibody can be reactive; it is used herein in the same way as specialists use it in this technical field, including, but not limited to, polypeptides, peptides, polysaccharides, glycoproteins, polynucleotides (for example, DNA), or chemical antigens, receptors or interleukins. Interleukins can include interleukins of various groups, such as interleukin 1 (alfa and beta), interleukin 2, interleukin 3, interleukin 4, interleukin 5, interleukin 6, interleukin 7, interleukin 8, interleukin 9, interleukin 10, interleukin 11, interleukin 17, interleukin 18 and interleukin 33.

The term “antigen” can also be used to describe the material that is used for immunization of animals (for example, llama) with the purpose of production of antibodies, according to the invention. In this context, “antigen” can have a broader meaning and may cover purified forms of an antigen as well as non-purified or not fully isolated, or purified antigen products such as cells, cell lysates, or supernatants, cell fractions, for example, cell membranes etc. with added haptens conjugated with a protein-carrier. Antigen used for immunization does not necessary mean an antigen structurally identical to an antigen target to which, finally, an antibody of the invention is able to bind. Usually, antigen used for immunization is a downsized version of an antigen target, for example, a fragment comprising an immunogenic epitope. More details about antigens used for immunization are described in literature and may be familiar to a specialist in this technical field.

Variable regions of each light/heavy chain pair form antigen-binding sites of the antibody. Thus, an intact IgG antibody has two binding sites. Except for bi-functional or bi-specific antibodies, two binding sites are identical. According to this application, “antigen-binding region” or “antigen-binding site”, or “antigen-binding domain”, are interchangeable, as used herein, and refer to an antibody molecule fragment comprising amino acid residues interacting with an antigen and giving the antibody its specificity and affinity to an antigen. This antibody fragment includes frame amino acid residues necessary for maintaining the proper conformation of the antigen-binding residues.

Preferably, CDR of VHH antigen-binding region or the entire antigen-binding region of the antibodies covered by the invention fully originate from Camelidae bloodline or are substantially of Camelidae origin, and comprise specific changed amino acid residues, for example, residues substituted with various amino acid residues (for example, refer to Table 6) in order to improve the particular properties of an antibody (for example, K_D , k_{off} , IC_{50}). Preferably, the antibody framework regions, in accordance with the invention, are of Camelidae origin or of human origin, or substantially of human origin (at least by 80, 85, 90, 95, 96, 97, 98 or 99% human origin), and comply with Kabat numbering.

“Antibody fragment” may represent an antibody fragment or antibody fragment that has the activity of a full-size antibody. The said antibody fragment may be represented by F(ab')₂, F(ab)₂, Fab', Fab Fv and scFv.

"Interleukin 17", also referred to as "IL-17" or "IL-17A", is a 20-30 kD homo-dimeric glycoprotein. The gene of human IL-17A encodes the protein consisting of 155 amino acids and having a 19 amino acid signal sequence and 136 amino acid mature segment. Amino acid sequence of human IL-17A is by 80%, 63% and 58% similar to amino acid sequences of a rabbit, mouse and rat, respectively. Amino acid sequence of human IL-17A is by 97% identical to IL-17A of cynomolgus monkey.

The term "antibody", when applied to anti-IL-17 monoclonal antibody of the invention (hereafter referred to as an "antibody of the invention"), as used herein, means a monoclonal antibody.

As used herein, the terms "inhibit" or "neutralize" regarding the activity of an antibody of the invention shall mean the ability to block, prevent, restrict, slow down, stop, reduce or reverse significantly, for example, the development or severity of inhibition subject, including but not limited to biological activity (such as activity of IL-17) or property, disease or condition. Binding the antibody, according to the invention, with IL-17 results in inhibition or neutralization of IL-17 activity preferably of at least 20, 30, 40, 50, 60, 70, 80, 90, 95% or higher.

The term "separated" or "isolated" with regard to nucleic acids or protein products (such as an antibody) refers to the nucleic acid molecule or protein molecule that is identified and separated from at least one of contaminating substances to which it is usually combined in the natural source. Preferably, an "isolated antibody" is an antibody that substantially contains no other antibodies that have particular antigenic specificity (for example, pharmaceutical compositions of the invention contain an isolated antibody that specifically binds IL-17 and substantially contain no antibodies that specifically bind antigens different from IL-17).

The term "Kabat numbering scheme" or "numbering according to Kabat" as used herein refers to a system of numbering amino acid residues that are more variable (i.e. hypervariable) than other amino acid residues in variable regions of heavy and light antibody chains (Kabat et al. Ann. NY Acad. Sci., 190:382-93 (1971); Kabat et al. Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242 (1991)).

Polynucleotide is "functionally bound", if it has functional links to other polynucleotides. For example, promoter or enhancer is functionally bound to coding sequence if it affects the sequence transcription. Polypeptide is

“functionally bound” to another polypeptide if polynucleotides coding thereof are functionally bound, preferably if they are located in the same open reading frame.

The term “DNA structure”, as used herein, refers to DNA or its fragment coding an antibody of the invention. Generally, DNA or its fragment that encodes an antibody (for example, an antibody of the invention) is functionally (operably) bound, within an open reading frame, to at least one other DNA fragment that encodes an additional polypeptide (for example, domain or region of a receptor for another cytokine, such as IL-2-receptor), and then is inserted into the appropriate expressing vector. Normally, DNA structures are formed in such a way that several DNA fragments encoding certain antibody sites are functionally associated within a reading frame to obtain a solid structure that encodes either an entire antibody or its functional fragment. For example, DNA structure would encode an antibody from N-terminus to C-terminus. Such antibodies can be expressed, isolated and evaluated regarding their activity.

The term “vector”, “plasmid” refers to nucleic acids obtained synthetically and via biotechnology containing a certain sequence of functional elements known in this field. Certain vectors can autonomously replicate in host cells to which they were introduced, while other vectors can be integrated into host cell genome and replicate together with the host genome. Moreover, some vectors can mediate the expression of genes to which they are functionally bound. In this application such vectors are called “recombinant expression vectors” (or “expression vectors”); exemplary vectors are well known from the prior art.

As used herein, the terms “cell”, “host cell”, “cell line” and “cell culture” are interchangeable and refer to an individual cell or cell culture that is a recipient of any isolated polynucleotide, according to the invention, or any recombinant vector (recombinant vectors) that contains the sequence of an antibody of the invention. Host cells involve offspring obtained from an individual host cell; offspring may not necessary be completely identical (regarding the morphology or full DNA complement) to the original host cell due to natural, accidental or intended mutations and/or variations. A host cell includes cells that were transformed,

transduced or infected with recombinant vector, or a monoclonal antibody that expresses a polynucleotide covered by the invention or its heavy or light chain. Host cell that contains a recombinant vector covered by the invention (either incorporated into host chromosomes or not) can also be called “recombinant host cell”. Preferable host cells to be used in the invention are CHO cells (for example, ATCC CRL-9096), NS0 cells, SP2/0 cells, COS cells (ATCC, for example, CRL-1650, CRL-1651) and HeLa (ATCC CCL-2). Additional host cells to be used in the invention include plant cells, yeast cells, other mammalian cells and prokaryotic cells.

The term “Specific binding” between an antibody and an antigen target (antigen) refers to immunological specificity. Antibody can specifically bind an antigen target if it binds an antigen epitope stronger than other antigen epitopes. Specific binding does not exclude cross-reactivity with other antigens that carry similar antigen epitopes.

VL domains in antibodies of the invention can be of either VL lambda type or VL kappa type. The term “VL domain” covers both VL lambda and VL kappa isotypes that contain one or more amino acid substitutes, insertions or deletions.

The term “pharmaceutical composition” covers the formulation and/or composition containing a therapeutically efficient amount of the antibody, according to the invention, and adjuvants (excipients, diluents, filling materials, solvents and other adjuvants, such as).

The term “use” or “treatment” applies to the ability of using an antibody of the invention or a pharmaceutical composition containing thereof to treat, alleviate the course of the disease, expedite the remission or reduce the recurrence rate for the disease or disorders mediated by receptors with which an antibody of the invention can bind.

Description of the invention

The present invention proposes humanized monoclonal antibodies, preferably of IgG type having an increased affinity and improved aggregation stability, wherein variable domains are represented by a combination of VHH-derivative with a variable domain of the light chain V_L .

In one embodiment, VHH-derivative of the antibody, according to the invention, may comprise amino acid substitutes at positions $44X_245X_3$, with $X_2 = G, A, V, S, T$; and $X_3 = A, V, T, H$; or combinations thereof (44 and 45 designate the positions of amino acid substitutes). Hereinafter the position of amino acid substitute is indicated using Kabat numbering scheme (<http://www.bioinf.org.uk/abs/>).

Another embodiment involves an antibody of the invention that has improved aggregation stability of VHH-derivative, as compared to the initial IgG antibody comprising VHH isolated from an immunized animal, wherein an immunized animal can be from Camelidae.

Another embodiment of the invention involves an antibody that comprises V_{HH} -derivative that is a variable domain of the heavy chain of an antibody isolated from an immunized animal of Camelidae family. Herein, V_{HH} -derivative can have additional amino acid substitutes typical of humans at any positions, except for the following substitutes at positions 44 and 45:

- a) $44X_2$, wherein $X_2 = G, A, V, S, T$;
 - b) $45X_3$, wherein $X_3 = A, V, T, H$;
- or combinations thereof.

Another embodiment involves an antibody of the invention that comprises V_{HH} -derivative that can be represented by a heavy chain variable domain isolated from non-immunized animal of Camelidae family. Herein, V_{HH} -derivative can have additional amino acid substitutes typical of humans at any positions, except for the following substitutes at positions 44 and 45:

- a) $44X_2$, wherein $X_2 = G, A, V, S, T$;

- b) $45X_3$, wherein $X_3 = A, V, T, H$;
or combinations thereof.

Another embodiment involves an antibody of the invention that comprises a light chain variable domain V_L that is a derivative of a human antibody. In additional embodiment of the invention, the light chain variable domain V_L is a humanized fragment of an animal antibody.

Another embodiment involves an antibody of the invention that comprises a V_{HH} -derivative that contains cysteine-44 (Kabat numbering scheme), and a light chain variable domain V_L that contains cysteine-100 (Kabat numbering scheme).

Another embodiment of the invention involves an antibody of any of the following isotypes: IgG1, IgG2, IgG3 or IgG4.

Another embodiment involves an antibody of the invention that comprises a non-native modified Fc as a part of IgG.

Another embodiment involves an antibody of the invention that has such aggregation stability under which the content of aggregates increases by not more than 5% of their initial content in the solution, when used in concentrations over 10 mg/ml and stored for >6 months at a temperature of 4°C. In additional embodiment of the invention, an antibody has such aggregation stability under which the content of aggregates increases by not more than 5% of their initial content in the solution, when used in concentrations over 10 mg/ml and stored for >2 weeks at a temperature of 37°C. Another additional embodiment of the invention involves an antibody that has such aggregation stability under which the content of aggregates increases by not more than 5% of their initial content in the solution, when used in concentrations over 10 mg/ml and stored for >48h at a temperature of 42°C. One more additional embodiment of the invention involves an antibody that has such aggregation stability under which the content of aggregates increases by not more than 5% of their initial content in the solution, when used in concentrations over 10 mg/ml and stored for > 6 h at a temperature of 50°C.

One embodiment involves an antibody with dissociation constant $K_D \leq 10^{-9}$ M. Another embodiment involves an antibody of the invention that has an

antibody-antigen interaction association constant $k_{on}(1/Ms) \geq 10^5$ 1/Ms. One more embodiment of the invention involves an antibody that has a antigen-antibody dissociation constant $k_{dis}(1/s) \leq 10^{-4}$ 1/s.

In one of the implementation variants, the invention presents an antibody fragment. This antibody fragment can be represented by a light chain, heavy chain, variable domains of the light and/or heavy chain that are a part of the antibody sequence, which includes a bio-specific antibody variant. Another embodiment of the invention involves an antibody fragment that is represented by a light chain, heavy chain, variable domains of the light and/or heavy chain that are parts of Fab. The antibody fragment of the invention may be represented by a light chain, heavy chain, variable domains of the light and/or heavy chain that are parts of scFv.

In addition, the invention suggests the methods of production of antibodies based on the invention. The said method of antibody production can involve phases selected from the following: directed mutagenesis, display methods, genetic engineering, biochemistry and high-performance biotechnology methods well known from the prior art, which can include the methods for directed mutagenesis in different positions of VHH domain of Camelidae antibodies.

In addition, the invention provides a DNA structure encoding antibodies, according to the invention, or fragments thereof, and an expression vector comprising one or several DNA structures of the invention.

Moreover, the invention suggests a cell line comprising the said expression vector or DNA structure.

In addition, the invention suggests a method of production of a humanized monoclonal antibody or fragment thereof, which involves the cell line culture in a culture medium under the conditions sufficient to obtain the said antibody or fragment thereof, followed by isolation and purification of the obtained antibody or its active fragment.

Moreover, the invention suggests a pharmaceutical composition that contains an antibody or fragment thereof in combination with one or several pharmaceutically suitable excipients, diluents or carriers. Details of the techniques

to produce the composition are described in special biotechnology guidelines, for example in [25].

Another embodiment of the invention involves an antibody or fragment thereof within an active molecule that specifically binds to human IL-17A, wherein the said antibody or fragment thereof comprises a derivative of Camelidae heavy chain variable domain (V_{HH}) that comprises 3 hypervariable regions HCDR1, HCDR2 and HCDR3, wherein:

HCDR1 comprising the amino acid sequence of SEQ ID NO: 1:

G-T-F-A-T-X32-X33-X34 - X35 (numbering according to Kabat index), wherein X32 is an amino acid selected from the group comprising S, N, K, R, E, W, Q, D, A, V and F;

X33 is an amino acid selected from the group comprising P and S;

X34 is an amino acid selected from the group comprising M and I;

X35 is an amino acid selected from the group comprising G, N, S, A, L, I, R, V and Q;

HCDR2 comprises amino acid sequence of SEQ ID NO: 2:

X50-I-S-X52-X52a-S-G-X55-D-R-I-Y-A-D-S-V-K-G, wherein

X50 is an amino acid selected from the group comprising A, G and L;

X52 is an amino acid selected from the group comprising D and E;

X52a is an amino acid selected from the group comprising P and A;

X55 is an amino acid selected from the group comprising G, S, T, L, R, D, E, K, A and W;

HCDR3 comprises amino acid sequence of SEQ ID NO: 3:

C-A- X94-X95- X96- X97-F-X99-X100-X100a- X100b - X100c - X100d - X100e - X100f – D- Y – D-S, wherein

X94 is an amino acid selected from the group comprising K, S, T, V, D and G;

X95 is an amino acid selected from the group comprising R and K;

X96 is an amino acid selected from the group comprising G, R, Y, H, D, W and K;

X97 is an amino acid selected from the group comprising R, A, V, S, L and H;

X99 is an amino acid selected from the group comprising D, E, G, A, R, V, K and Q;

X100 is an amino acid selected from the group comprising G, S and N;

X100a is an amino acid selected from the group comprising G, T, P, V, R, N and K;

X100b is an amino acid selected from the group comprising V, S, T, L, Y, A, H, G and I;

X100c is an amino acid selected from the group comprising Y, W and S;

X100d is an amino acid selected from the group comprising R, V, L, Y, A, W, K, G, Q and I;

X100e is an amino acid selected from the group comprising T, L, A and S;

X100f is an amino acid selected from the group comprising T, L, G, P, N, A, Q, F, I and D;

b) A variable domain of the light chain (V_L) of a human antibody or a variable domain of the light chain of a humanized antibody.

An alternative embodiment involves an antibody or a fragment thereof that specifically binds human IL-17A and comprises a derivative of the heavy chain variable domain (V_{HH}) comprising 3 hypervariable regions HCDR1, HCDR2 and HCDR3 wherein:

HCDR1 comprising an amino acid sequence of G-T-F-A-T-S-P-M-G (SEQ ID NO: 4);

HCDR2, comprising an amino acid sequence of A-I-S-P-S-G-G- D-R-I-Y-A-D-S-V-K-G (SEQ ID NO: 5);

HCDR3 comprising an amino acid sequence of C-A-V-R-R-R-F-D-G-T-S-Y-Y-T-G-D-Y-D-S (SEQ ID NO: 6);

b) A variable domain of the light chain (V_L) of a human antibody or a variable domain of the light chain of a humanized antibody.

An alternative embodiment involves an antibody that specifically binds human IL-17A and comprises a derivative of the heavy chain variable domain (V_{HH}) comprising an amino acid sequence of SEQ ID NO: 7 and a variable domain

of the light chain (V_L) of a human antibody or a variable domain of the light chain of a humanized antibody.

An alternative embodiment involves an antibody or a fragment thereof that specifically binds human IL-17A and comprises V_{HH} -derivative, wherein the said variable domain comprises an amino acid sequence at least 90% identical to SEQ ID NO: 7.

An alternative embodiment involves an antibody that specifically binds human IL-17A and contains a V_{HH} -derivative comprising an amino acid sequence of SEQ ID NO: 7; and a variable domain of the light chain (V_L) of a human antibody comprising an amino acid sequence of SEQ ID NO: 8.

An alternative embodiment involves an antibody or a fragment thereof that specifically binds human IL-17A and comprises V_{HH} -derivative and a variable region of the light chain (V_L) of a human antibody, wherein the said variable domains comprise an amino acid sequence at least 90% identical to SEQ ID NO: 8.

An alternative embodiment involves an antibody that specifically binds human IL-17A and contains a heavy chain comprising SEQ ID NO: 9 amino acid sequence, and a variable domain of a human antibody light chain (V_L) comprising SEQ ID NO: 10 sequence. Another embodiment involves an antibody that specifically binds human IL-17A and contains a heavy chain and a light chain, wherein the said chains comprise amino acid sequences at least 90% identical to SEQ ID NO: 9 and/or SEQ ID NO:10.

An alternative embodiment involves an antibody of the invention that specifically binds human IL-17A, wherein the said antibody has a binding affinity to human IL-17A characterized by K_D of $\leq 10^{-10}$ M. Another embodiment involves an antibody that specifically binds human IL-17A, wherein the kinetic association constant $k_{on}(1/Ms)$ for human IL-17A is at least 10^5 1/Ms. Another embodiment involves an antibody that specifically binds human IL-17A, wherein the kinetic dissociation constant $k_{dis}(1/s)$ for human IL-17A is not more than 10^{-5} 1/s. An alternative embodiment involves an antibody that specifically binds human IL-17A

and inhibits the activity of human IL-17A by no less than 50% with respect to any parameter examined by any specific activity testing.

An alternative embodiment involves an antibody that specifically binds human IL-17A, wherein the said antibody is produced by mammalian, yeast or bacterial cells.

An alternative embodiment involves an antibody that specifically binds human IL-17A and contains one or more additional amino acid substitutes in Fc-region, as compared to natural Fc, wherein the said substitutes improve physical-chemical and pharmacokinetic properties of an antibody, as compared to the antibody with natural Fc, and do not result in loss of antibody's ability to bind IL-17A.

An alternative embodiment suggests a DNA structure encoding an antibody that specifically binds human IL-17A. Moreover, the invention suggests an expression vector comprising one or more DNA structures encoding an antibody that specifically binds human IL-17A. In addition, a host cell was suggested comprising a vector for obtaining an antibody that specifically binds human IL-17A.

In addition, the invention suggests a method for the production of an antibody that specifically binds human IL-17A, based on culturing host cells comprising a DNA structure in a culture medium under conditions suitable to obtain the said antibody or a fragment thereof, and further isolation and purification of the said antibody or active fragment thereof.

Moreover, a pharmaceutical composition was suggested comprising an antibody that specifically binds human IL-17A, and one or more pharmaceutically suitable excipients, diluents or carriers. The said composition can comprise further active pharmaceutical ingredients selected from TNF- α inhibitors. The said composition can be used in treatment of an IL-17A-mediated disease or disorder. An IL-17A-mediated disease or disorder is selected from: rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus,

Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroadenitis, asthma, allergic disorders, psoriasis, dermatitis, systemic sclerosis, graft-versus-host disease, graft rejection, acute or chronic immune disease associated with organ grafting, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki disease, Graves' disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's disease, Henoch-Schonlein purpura, microscopic renal vasculitis, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infections, parasitic diseases, acquired immune deficiency syndrome, acute transverse myelitis, Huntington chorea, Parkinson disease, Alzheimer disease, blood stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure, myocardial infarction, Addison disease, polyglandular autoimmune syndrome type I and type II, Schmidt's syndrome, acute respiratory distress syndrome, alopecia, alopecia areata, seronegative arthropathy, arthropathy, Reiter's syndrome, psoriatic arthropathy, arthropathy associated with ulcerative colitis, enteropathic synovitis, arthropathy associated with Chlamydia, Yersinia and Salmonella, spondyloarthropathy, atheromatosis disease/coronary sclerosis, atopic allergy, autoimmune bullous disease, pemphigus, pemphigus foliaceus, pemphigoid, linear IgA diseases, autoimmune hemolytic anemia, Coombs positive hemolytic anemia, acquired pernicious anemia, juvenile pernicious anemia, Myalgic encephalomyelitis/chronic fatigue syndrome, chronic active hepatic inflammation, cranial giant arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, acquired immune deficiency syndrome (AIDS), AIDS-associated diseases, hepatitis B, hepatitis C, common variable immunodeficiency (common variable hypogammaglobulinemia), dilated cardiomyopathy, female sterility, ovarian insufficiency, premature ovarian failure, pulmonary fibrosis, cryptogenic fibrosing alveolitis, post inflammatory interstitial lung pathologies, interstitial pneumonitis, connective tissue disease associated with interstitial lung disease, mixed connective tissue disease associated with interstitial lung disease, systemic scleroderma associated with interstitial lung disease, rheumatoid arthritis

associated with interstitial lung disease, systemic lupus erythematosus associated with lung disease, dermatomyositis/polymyositis associated with lung disease, Sjogren disease associated with lung disease, ankylosing spondylitis associated with lung disease, diffuse pulmonary vasculitis, hemosiderosis associated with lung disease, drug-induced interstitial lung disease, fibrosis, radiation-induced fibrosis, obliterating bronchiolitis, chronic eosinophilic pneumonia, lung disease with lymphocyte infiltration, post infectious interstitial lung pathologies, gouty arthritis, autoimmune hepatitis, autoimmune hepatitis type I (classic autoimmune or lupoid hepatitis), autoimmune hepatitis type II (associated with anti-LKM antibody), autoimmune hypoglycemia, type B insulin resistance with acanthokeratoderma, hypoparathyroidism, acute graft-associated immune disease, chronic graft-associated immune disease, osteoarthritis, primary sclerosing cholangitis, type I psoriasis, type II psoriasis, idiopathic leukopenia, autoimmune neutropenia, NOS-kidney diseases, glomerulonephritis, microscopic renal polyangiitis, Lyme disease, discoid lupus erythematosus, idiopathic or NOS-male sterility, antisperm immunity, multiple sclerosis (all types), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue disease, Goodpasture syndrome, pulmonary manifestations of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease, systemic scleroderma, Sjogren's Syndrome, Takayasu disease/arthritis, autoimmune thrombocytopenia, idiopathic thrombocytopenia, autoimmune thyroid disorders, hyperthyroid, autoimmune hypothyroidism (Hashimoto disease), atrophic autoimmune hypothyroidism, primary myxedema, phacogenic uveitis, primary vasculitis, vitiligo, acute hepatic disease, chronic hepatic disease, alcoholic cirrhosis, alcohol-induced liver damage, cholestasis, idiosyncratic hepatic disease, drug-induced hepatitis, nonalcoholic steatohepatitis, allergies and asthma, group B streptococcal infection (GBS), mental disorders (including depressions and schizophrenia), Th1- and Th2-mediated disease, acute and chronic pain (various forms), malignancies such as lung cancer, breast cancer, stomach cancer, bladder cancer, colorectal cancer, pancreatic cancer, ovarian cancer, prostate cancer and hematopoietic

malignancies (leukemia and lymphomas), abetalipoproteinaemia, acrocyanosis, acute and chronic infections and infestations, acute leukemia, acute lymphoblastic leukemia, acute myeloid leukemia, acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinoma, atrial ectopics, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic rhinitis, allograft rejection, alpha-I antitrypsin deficiency, lateral amyotrophic sclerosis, anemia, angina, anterior horn cell degeneration, anti-CD3 therapy, antiphospholipid syndrome, hypersensitivity reactions against receptors, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, coronary sclerosis, arteriovenous fistula, ataxia, atrial fibrillation (constant or paroxysmal), atrial flutter, atrioventricular block, B-cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt lymphoma, burns, cardiac arrhythmia, myocardial stunning syndrome, cardiac tumor, cardiomyopathy, inflammatory response to bypass, cartilage graft rejection, brain cortex degeneration, cerebellar disorder, chaotic or multifocal atrial tachycardia, chemotherapy-induced disorders, chronic myelocytic leukemia (CML), chronic alcohol addiction, chronic inflammatory pathologies, chronic lymphatic leukemia (CLL), chronic obstructive pulmonary disease, chronic salicylate intoxication, rectocolic carcinoma, congestive cardiac failure, conjunctivitis, contact dermatitis, pulmonary heart, coronary artery disease, Creutzfeldt-Jakob Disease, culture-negative sepsis, cystic fibrosis, cytokine therapy-induced disorders, boxer's encephalopathy, demyelinating diseases, dengue hemorrhagic fever, dermatitis, dermatological conditions, diabetes, diabetes mellitus, diabetes-related atherosclerotic vascular disease, diffuse Lewy body disease, congestive dilated cardiomyopathy, basal ganglia disease, Down's syndrome in middle age, motor disorders induced by CNS dopamine blockers, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, epiglottiditis, Epstein-Barr viral infection, erythralgia, extrapyramidal and cerebellar symptoms, familial hemophagocytic lymphohistiocytosis, fetal thymus graft rejection, Friedreich's ataxia, peripheral artery disease, fungal sepsis, gas

phlegmon, gastric ulcer, glomerulonephritis, any organ or tissue graft rejection, gram negative sepsis, gram positive sepsis, granulomas due to intracellular organisms, hairy-cell leukemia, Hallervorden-Spatz disease, Hashimoto's thyroiditis, hay fever, heart transplant rejection, hemochromatosis, hemodialysis, hemolytic uremic syndrome / thrombotic thrombocytopenic purpura, bleeding, hepatitis (A), bundle branch arrhythmia, HIV-infections/HIV-neuropathies, Hodgkin disease, hyperkinetic motor disorders, hypersensitivity reactions, hypersensitivity-associated pneumonitis, hypertension, hypokinetic motor disorders, examination of hypothalamo-pituitary-adrenal axis, idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody-mediated cytotoxicity, asthenia, infantile muscular atrophy, aortal inflammation, influenza virus A, exposure to ionizing radiation, iridocyclitis /uveitis/optic neuritis, ischaemia/reperfusion-induced disorders, ischaemic stroke, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, renal transplant rejection, legionellosis, leishmaniasis, leprosy, corticospinal damage, lipoedema, liver transplant rejection, lymphoedema, malaria, malignant lymphoma, malignant histiocytosis, malignant melanoma, meningitis, meningococemia, metabolic/idiopathic diseases, migraine, multiple system mitochondrial disorders, mixed connective-tissue disease, monoclonal gammopathy, multiple myeloma, multiple system degeneration (Mencel Dejerine-Thomas Shi-Drager and Machado-Joseph), myasthenia gravis, intracellular Mycobacterium avium, Mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial ischemic disease, nasopharyngeal cancer, neonatal chronic lung disease, nephritis, nephrotic, neurodegenerative disorders, neurogenic muscular atrophy I, neutropenic fever, non-Hodgkin's lymphomas, abdominal aortic branch occlusion, arterial occlusive disease, OKT3® therapy, orchitis/epididymitis, orchitis/vasectomy reversal operations, organomegaly, osteoporosis, pancreatic graft rejection, pancreatic carcinoma, paraneoplastic disease /tumor-related hypercalcemia, parathyroid graft rejection, pelvic inflammatory disease, perennial rhinitis, pericardial disease, peripheral atherosclerosis (atherosclerotic) disease, peripheral vascular disease, peritonitis,

pernicious anemia, Pneumocystis carinii pneumonia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal plasma-proliferative disorder and skin changes), postperfusion syndrome, pump head syndrome, post-cardiotomy post-infarction syndrome, preeclampsia, progressive supranuclear paralysis, primary pulmonary hypertension, radiation therapy, Raynaud's phenomenon and disease, Raynoud's disease, Refsum's disease, regular narrow QRS tachycardia, renal vascular hypertension, reperfusion injury, restrictive cardiomyopathy, sarcoma, scleroderma disease, senile chorea, senile dementia with Lewy bodies, seronegative arthritis, shock, sickle cell disease, skin allograft rejection, skin changes, small intestinal graft rejection, solid tumors, specific arrhythmias, spinal ataxia, spinocerebellar degradations, streptococcal myositis, cerebellar structural damage, subacute sclerosing panencephalitis, syncope, cardiovascular syphilis, systemic anaphylaxis, a comprehensive systemic inflammatory response syndrome, systemic-onset juvenile rheumatoid arthritis, T cells or FAB ALL, telangiectasia, thrombosis obliterans, thrombocytopenia, toxicity, grafting, trauma/bleeding, hypersensitivity reactions type III, hypersensitivity reactions type IV, unstable angina, uremia, urinary sepsis, urticaria, valvular heart disease, varicose veins, vasculitis, venous diseases, venous thrombosis, ventricular fibrillation, viral and fungal infections, viral encephalitis / aseptic meningitis, viral hemophagocytic syndrome, Wernicke-Korsakoff syndrome, Wilson's disease, heterograft rejection for any organ or tissue, acute coronary syndrome, acute idiopathic polyneuritis, acute inflammatory demyelinating radicular neuropathy, acute ischemia, adult-onset Stills disease, alopecia areata, anaphylaxis, antiphospholipid antibody syndrome, aplastic anemia, coronary sclerosis, atopic eczema, atopic dermatitis, autoimmune dermatitis, autoimmune disorder associated with streptococcus infection, autoimmune enteropathy, autoimmune hearing loss, autoimmune lymphoproliferative syndrome (ALPS), autoimmune myocarditis, autoimmune premature ovarian failure, blepharitis, bronchiectasis, bullous pemphigoid, cardiovascular disease, catastrophic antiphospholipid syndrome, celiac disease, cervical spondylosis,

chronic ischemia, cicatrical pemphigoid, clinically isolated syndrome (cis) with a risk of multiple sclerosis, conjunctivitis, childhood-onset mental disorders, chronic obstructive pulmonary disease (COPD), dacryocystitis, dermatomyositis, diabetic retinopathy, diabetes mellitus, herniated disk, prolapse of intervertebral disc, drug-induced immune haemolytic anaemia, endocarditis, endometreosis, entophthalmia, episcleritis, erythema multiform, severe erythema multiform, gestational pemphigoid, Guillain-Barre syndrome (GBS), hay fever, Hughes syndrome, idiopathic Parkinson's disease, idiopathic interstitial pneumonia, IgE-mediated allergy, autoimmune hemolytic anemia, inclusion body myositis, infectious ocular inflammatory disease, inflammatory demyelinating diseases, inflammatory heart disease, inflammatory kidney disease, idiopathic pulmonary fibrosis/ usual interstitial pneumonia, iritis, keratitis, keratoconjunctivitis sicca, Kussmaul disease or Kussmaul-Meier Disease, Landry palsy, Langerhans' cell histiocytosis, marbled skin, macular degeneration, microscopic polyangiitis, Bechterew disease, motor neuron disease, mucosal pemphigoid, multiple organ failure, myasthenia gravis, spinal cord dysplasia syndrome, myocarditis, nerve root disorders, neuropathy, non-A, non-B hepatitis, optic neuritis, osteolysis, ovarian cancer, oligoarticular juvenile rheumatoid arthritis, peripheral arterial occlusive disease, peripheral vascular disease, peripheral artery disease (PAD), phlebitis, polyarteritis nodosa, polychondritis, polymyalgia rheumatica, poliosis, polyarticular juvenile idiopathic arthritis, multiple endocrine deficiency, polymyositis, polymyalgia rheumatica (PMR), post pump syndrome, primary parkinsonism, prostate and rectal cancer and hematopoietic malignancies (leukemia and lymphoma), prostatitis, pure red-cell aplasia, primary adrenal insufficiency, relapsing neuromyelitis optica, restenosis, rheumatic heart disease, SAPHO (synovitis, acne, pustulosis, hyperostosis and osteitis), scleroderma, secondary amyloidosis, shock lung, scleritis, ischias, secondary adrenal insufficiency, silicon-associated connective tissue disease, Sneddon-Wilkinson disease, ankylosing spondylitis, Stevens–Johnson syndrome, systemic inflammation response syndrome, cranial arteritis, Toxoplasma rhinitis, toxic epidermal necrolysis, transverse myelitis, TRAPS (tumor necrosis factor

receptor-associated periodic syndrome), allergic reactions type I, diabetes type II, urticaria, usual interstitial pneumonia, vasculitis, vernal conjunctivitis, viral retinitis, Vogt-Koyanagi-Harada syndrome (VKH syndrome), wet macular degeneration, wound healing, and Yersinia- or Salmonella-associated arthropathy.

A pharmaceutical composition comprising an antibody that specifically binds human IL-17A can be administered in a therapeutically effective amount to treat an IL-17A-mediated disease or disorder.

An invention suggests a method of treating IL-17A-mediated disease or disorder using an antibody that specifically binds human IL-17A. The treatment method can involve further administration of TNF- α inhibitors.

The following examples illustrate the present invention, yet they are not intended to limit the present invention to those examples per se.

The description of the present application includes references to all information sources.

Examples

Example 1

The scheme for production of VHH-based antibodies

Figure 1 demonstrates a scheme for production and optimization of VHH-based antibody.

Example 2

Production of recombinant antigens and antibodies in suspension mammal cell culture

Antibodies and antigens were generated in established cell line obtained from Chinese hamster ovary cells (CHO-K1) according to the published protocols [26; 27]. Cells constitutively expressing the gene of EBNA1 protein (Epstein-Barr virus nuclear antigen 1) were used. Suspension cultivation was conducted in flasks on orbital shaker using serum-free media from Life Technologies Corporation, in

accordance with manufacturer's guidelines. For transient expression, cells in concentration of 2×10^6 /ml were transfected by means of linear polyethyleneimine (PEI MAX, Polysciences). DNA/PEI ratio was 1:3-1:10. 5-7 days after transfection, the cell culture was centrifuged under 2000g for 20 min and filtered through 0.22 μ m filter. Target proteins from culture liquid were isolated by affine HPLC.

Recombinant IL-17A protein containing 6 His amino acids in C-terminal region was isolated and purified from culture liquid with Profinity IMAC Ni-charged resin (Bio-Rad). Prior to purification procedures, NiCl_2 was added to culture liquid to reach the concentration of 1mM. Then 5 ml of Profinity IMAC Ni-charged was added to culture liquid and mixed on a shaker for 1 h at a room temperature. Sorbent was transferred to 5 ml Thermo scientific Polypropylene columns and washed with 5 column volumes of PBS to remove non-specifically bound components. The bound antigen was eluted with 0.3 M imidazole (pH 8) and 150mM NaCl. Then the protein was dialyzed into PBS (pH 7.4) by means of SnakeSkin Dialysis Tubing technique, filtered (0.22 μ m), transferred into tubes and stored at -70°C . The purity of the protein obtained was evaluated by SDS-PAGE (Figure 2, 3, 4).

The tested and control IgG1 antibodies were purified on 1 ml Hi Trap rProteinA FF column (GE Healthcare) in accordance with the aforementioned procedure for IL-17A-Fc. The purity of the protein obtained was evaluated by SDS-PAGE (Figure 2, 3, 4).

Example 3

Llama immunization with human IL-17A and generation of Fab-library of phage-displayed llama antibodies

Lama Glama animal was immunized 5 times in succession by means of subcutaneous administration of antigen material mixed with an equal volume of full (first injection) or partial (all injections except for the first one) Freund's adjuvant. A mixture of recombinant proteins (0.2 mg of each protein per injection)

one of which was human IL-17A (Kit from R&D Systems) was used as an antigen. The second injection (immunization stage) was performed 3 weeks after the first one; three more immunizations were performed with a 2-week interval. Blood samples (50 ml) were collected 5 days after each injection starting from the third one.

The selected llama blood was diluted two times with PBS solution containing 1 mM EDTA. Then 35 ml of diluted blood was layered over 15 ml of Histopaque® -1077 medium (Sigma, density of 1.077 g/ml) and centrifuged for 20 min under 800g. Mononuclear cells (lymphocytes and monocytes) were selected from plasma/Histopaque medium interphase zone and washed with PBS containing 1 mM EDTA.

The total amount of RNA from mononuclear llama cells was isolated using RNeasy Mini Kit in accordance with the protocol (QIAGEN). RNA concentration assay was performed using Nanovue (GE Healthcare); the quality of isolated RNA was tested by means of 1.5% agarose gel electrophoresis.

The reverse transcription reaction was conducted using MMLV RT kit (Evrogen), according to the recommended protocol, with MMuLV reverse transcriptase and random hexamer primers.

The reverse transcription products were used as a matrix in a two-stage polymerase chain reaction to obtain the genes of variant domains flanked with restriction sites; the reaction was performed using oligonucleotide kit and records of the authors [27; 28; 29]. Further, genes encoding variable domains of the light and heavy chains were put together in one fragment by means of sequential reactions of restriction, ligation and amplification as shown at Figure 5. Heavy chain genes were attached separately to kappa and lambda light chain genes. In this case, the estimated count of matrix molecules in all reactions was no less than 10^{11} . The DNA product obtained (VL-CK-VH) was treated with NheI /Eco91I restriction enzymes and ligated into original phagemid pH 5. Phagemid structure is presented at Figure 6. Ligation products were transformed into SS320 electrocompetent cells prepared in accordance with protocols [30]. The repertoire

of structured kappa and lambda Fab libraries was 5.1×10^8 and 3.7×10^8 , respectively. The product of native phage-display library was prepared in accordance with the procedure described above [31].

Example 4

Selection of Fab-libraries of phage-display antibodies

Specific anti-IL17A phage-display Fab-antibodies were selected from a phage Fab-display library using a recombinant human IL-17A (a kit from R&D Systems); a series of selection cycles was performed, as described above [27; 31; 32]. In order to perform the selection process by panning method, human IL-17A in 50 mM carbonate buffer (pH 9.5) was adsorbed overnight at 4°C on the surface of HighSorb tubes (Nunc). Further, tubes were washed with PBS (pH 7.4) and then blocked with solution containing PBS (pH 7.4) – fat-free milk (0.5% weight/volume) for 1 hour. Then, 2-4 ml of phage solution (10^{12} phage particles per ml) in PBS (pH 7.4) – fat free milk (0.5% w/vol) was transferred to the tube with the antigen, and the system was incubated for 1 h under stirring. Unbound phages were removed by a series of washing cycles with PBS (pH 7.4) - Tween 20 (0.1% vol./vol.). The number of washing cycles was increased from the first round to the third one by 20-30-40 times, respectively. Phage particles that remained bound were eluted with 100 mM Gly-HCl solution (pH 2.5) during 15 min under stirring, and then neutralized with 1 M TRIS-HCl (pH 7.6). *E. coli* TG1 bacteria were infected with the phages obtained; further, then the phages were isolated and used in the next cycle.

After the second and the third selection round, ELISA performed for the polyclonal phage product has shown significant enrichment. Pooled clones enriched with human Fab were re-cloned to expression plasmid LL comprising myctag and His6 tag on C-terminus of CH1gene of the heavy chain.

Example 5

Analysis of Fab specific binding with human IL-17A

ELISA was used to measure the binding of Fab-fragments under research with human IL-17A. Fab with published AIN457 sequence (Novartis) was used as a positive control. ELISA plate wells (Nunc ImmunoMaxisorp) were coated with 50 μ l (0.5 μ g/ml in 1X coating carbonate buffer) IL-17A-Fc, hermetically closed and incubated overnight at 4°C. All further stages were conducted in accordance with the standard ELISA protocol with high-performance systems such as GenetixQ-pix2xt (Molecular Devices) and Tecan Freedom EVO 200 (Tecan). Non-specific binding was blocked by adding the blocking buffer BB (200 μ l 0.5% fat-free milk in PBS). Plates were incubated on a shaker for 1 h at a room temperature. After washing with PBS-Tween, each cell was coated with 50 μ l of test Fab-containing cell supernatant mixed with equal volume of BB. Plates were incubated on a shaker for 1 hour at a room temperature; further, each plate well was washed 5 times with PBS-Tween buffer. After washing, each well was coated (50 μ l/well) with anti-human Fab HRP-conjugated secondary antibody (Pierce-ThermoScientific) in PBS-Tween (1:5000). Plates were transferred to rotation shaker (50 min at room temperature) and then washed 5 times with PBS-Tween buffer as described above. Colorimetric signal was obtained by adding TMB (100 μ l/well) until saturation (the average of 3-5 min); further development was blocked by adding stop solution (100 μ l/well, 10% sulfuric acid). Absorbance was measured at 450 nm using the appropriate Tecan-Sunrise plate reader (Tecan). Antibody binding was proportional to the signal produced. Those clones for which the color signal exceeded the baseline signal by more than 5 times were tested in competitive ELISA in order to reveal antagonistic Fab blocking the interaction between IL-17A ligand and receptor.

Example 6

Competitive ELISA of blocking the interaction of IL17A ligand and IL17R receptor

Competitive ELISA technique was used to test the antagonistic capacity of previously selected specific Fab against human IL-17A. Fab with published AIN457 sequence (Novartis) was used as a positive antagonist control. ELISA plates wells (Nunc Immuno Maxisorp) were covered with 50 μ l/well IL-17RA-Fc receptor (R&D Systems; 1 μ g/ml solution in 1X coating carbonate buffer) and incubated overnight at 4°C. All further stages were performed in accordance with standard ELISA protocols with high-performance systems such as GenetixQ-pix2xt (Molecular Devices) and Tecan Freedom EVO 200 (Tecan). Non-specific binding was suppressed by adding the blocking buffer BB (200 μ l 0.5% fat-free milk in PBS). Plates were incubated for 1 hour on a shaker at a room temperature.

At the same time, 50 μ l of test Fab-containing cell supernatant in non-binding 96-well plate was mixed with 50 μ l of IL-17A-His6-Flag (0.4 μ g/ml in 1% milk diluted with PBS-Tween). The plate was incubated for 1 hour at 37°C on a shaker under 500 rpm.

After the plate containing IL-17RA-Fc receptor was washed from BB solution, it was coated with the reaction mixture of Fab and IL-17A-His6-Flag in the amount of 90 μ l per well. Plates were incubated and shaken for 45 min at a room temperature, and each well was washed 5 times with PBS-Tween buffer. Further, 50 μ l /well of 1 μ g/ml anti-FLAG murine M2 antibody (Sigma) was added, and plates were incubated for 45 min at a room temperature. After incubation, each plate well was washed 5 times with PBS-Tween, then coated with 50 μ l/well of antimurine-IgG HRP-conjugated secondary antibody (Pierce-ThermoScientific) 1:5000 diluted with PBS-Tween. The plates were incubated on rotation shaker for 45 min at a room temperature and washed 5 times with PBS-Tween, as mentioned above. Colorimetric signal was obtained by adding TMB (100 μ l/well) until saturation (the average of 3-5 min); further development was blocked by adding stop solution (100 μ l/well, 10% sulfuric acid). The absorbance

was measured at 450 nm using an appropriate Tecan-Sunrise plate reader (Tecan). The antibody binding was proportional to the color signal produced.

Those clones that demonstrated suppression at the level corresponding to that of control Fab antibody AIN457 were marked as positive and used in further tests. Genes of the variable domains of positive clones were subject to sequencing in accordance with standard protocols on *Applied Biosystems 3130 Genetic Analyzer (Applied Biosystems)* followed by the appropriate analysis. Clones comprising 3 VHHFab variable domains were selected for further studies (Figure 7). In addition, it was found that 3VHHFab clone is represented in combination with 23 light chain domains of various sequences, three of which are shown at Figure 5B. This indicates its relative structural resistance and the fact that this is VHH domain and not the light chain that contributes to IL-17A binding.

Example 7

Comparative k_{off} (k_{diss})-screening of anti-IL-17A VHHFab candidates

The comparative k_{off} screening for anti-IL-17A Fab-candidates was performed using Pall Forte Bio Octet Red 96 system. Anti-FABCH1 biosensors were rehydrated for 30 min in a working buffer comprising 10 mM PBS (pH 7.2-7.4), 0.1% Tween-20 and 0.1% BSA. 10x working buffer was added to test the samples of *E.coli* supernatants up to 1x of the final concentration. Then anti-FABCH1 biosensors were steeped into *E.coli* supernatants containing Fab-fragments of candidate antibodies and incubated for 12 hours at a temperature of 4°C. Sensors coated with Fab-fragments were transferred to wells with working buffer, and a baseline was registered (60 sec). Then sensors were transferred to wells with analyte solution (IL-17A, 30 µg/ml) to achieve the antigen-antibody association (300 sec). After that, sensors were returned into wells with working buffer for further dissociation (300 sec). The used sensors were subject to regeneration after each test: they were placed three times into regenerating buffer (Gly-HCl, pH 1.7) and then were applicable for use in further experiments. The

curves obtained were analyzed with Octet Data Analysis (version 7.0) according to the standard procedure with 1:1 interaction model.

The results obtained for k_{off} -screening of anti-IL-17A Fab candidates are presented at Figure 5C and Table 1. Specific and high-affine binding of all unique VHHFab with human IL-17A was demonstrated, wherein 3VHHFab showed very fast k_{on} and very slow $k_{\text{dis}}(1/c)$ that was beyond the detection limit of the device.

Table 1. VHHFab affinity to human IL-17A

	K_D (M)	$k_{\text{on}}(1/\text{Mc})$	$k_{\text{dis}}(1/c)$
1VHH	2.485E-08	2.498E04	6.206E-04
2VHH	1.352E-08	1.628E05	2.201E-03
3VHHVK4B11	-	-	<2.272E-06

Thus, 3VHHVK4B11 candidate was selected for further investigation based on the analytical results obtained.

Example 8

Generation of VHHIgG1 antibodies with mutations in FR1 and FR2 of VHH variable domain

Genes of the variable domain of the light and heavy VHH chains of 3VHHVK4B11 candidate were cloned in pEE-HcnpEE-Lc plasmid for joint transient expression in CHO-EBNA cells as described in Example 3. Further, substitutes at positions 44 and 45 (Kabat numbering scheme) were introduced by means of oligonucleotide-directed mutagenesis using PfuUltraHS polymerase (Stratagene) in accordance with Protocol [Q5® Site-Directed Mutagenesis Kit (NEB)] and procedure described in [34]. Plasmid pEE-3VHH was used as a matrix. PCR products were fractioned on low-melting agarose and purified on appropriate columns. After ligation, DNA was transformed into *E.coli*. Upon selection of mutant clones with correct sequences, plasmids with mutations in 3VHH were co-transfected with pEE-LcVK4B11(refer to Figure 9 and Table 2).

Table 2. Samples of 3VHHFabVK4B11 mutant clones

Name	Mutations as compared to initial 3VHHFabVK4B11
mut1	E44G, R45L
mut2	S11L, E44G, R45L
mut3	R45L
mut4	S11L, E44G

Example 9

Comparative analysis of kinetic parameters of VHHFab antibodies comprising mutations in FR1 and FR2 of VHH variable domain

The comparative k_{off} screening for anti- IL-17A VHHFab candidates was performed according to the standard protocol with Pall Forte Bio Octet Red 96 system (refer to Example 8). A significant reduction of k_{off} was found for mut1, mut2 and mut4, and less significant reduction for mut3, as compared to natural 3VHHFab (Figure 10, 11).

Example 10

The Comparative analysis of thermal-aggregation characteristics of full-size 3VHHIgG1VK4B11 antibodies comprising mutations in FR1 and FR2 of VHH variable domain

The comparative analysis of aggregation characteristics for anti-IL-17A VHHIgG1 candidates was performed using the following procedure. The preparation of VHHIgG1 antibody (10 mg/ml) in PBS buffer was heated for 6 hours at 50°C. The aggregation induced by thermal stress was evaluated by means of size-exclusion HPLC. The test was performed on 1100 HPLC System (Agilent) using Tosoh TSK-Gel G3000SWXL column, 7.8 mm x 30 cm, Cat. No. 08541 with Tosoh TSKgel Guard SWXL pre-column, 6.0 mm x 4.0 cm (particles of 7 μm , Cat. No. 08543). Isocratic elution with mobile phase containing 50 mM sodium phosphate buffer and 0.3 M NaCl (pH 7.0) was performed under 0.5 ml/min flow rate with detection at 214 nm and 280 nm wavelengths. The antibody

samples were diluted with PBS (pH 7.5) to the concentration of ~1 mg/ml. The injection volume was 10 μ l. Gel filtration standard mixture (Bio-Rad, Cat. No. 151-1901) was used to calibrate the column prior to the test.

Chromatograms presented at Figure 6C and summary Table 3 demonstrate that all mutants aggregate to various extents under thermal stress, wherein the minimum stability was observed for the natural variant and the maximum stability was revealed for mut1 comprising E44G+R45L substitutes typical of classical VH structure.

In addition, the comparative study of thermal stability of the obtained preparations was performed using Thermofluor procedure (also referred to as Thermal shift assay) that determines the protein melting point measuring changes in the fluorescence of a specific dye SYPROOrange that binds to hydrophobic surfaces of the denatured protein [35]. StepOneReal-TimePCRSytem (Applied Biosystems) apparatus and recommended protocol were used to study the mutant products. The study results are shown in Table 3. They rather correlate to the results obtained under thermal stress test, which confirm the stability of mut1 and mut4 in comparison to the initial variant 3VHHIgG1VK4B11.

Table 3. Results of thermal stability study of mutants

Product	Monomer percentage (%) before thermal stress	Monomer percentage (%) after thermal stress	Difference $\sim \Delta\%$	Melting point (Thermofluor), $^{\circ}\text{C}$
Wild 3VHHIgG1VK4B11	60	13	47	46 $\pm$ 1
mut1	95	95	0	64 $\pm$ 1
mut2	65	13	52	42 $\pm$ 1
mut3	40	32	8	ND
mut4	89	80	9	53 $\pm$ 1

Example 11

Cell test of anti-IL17A 3VHHIgG1VK4B11 mutants blocking the ability of IL-17A to induce IL-6 production

The ability of IL-17 to induce the production of IL-6 by HT1080 human cells (ATCC:CCL-121) was used to analyze the neutralizing activity of VHHIgG1 candidates mut1 and mut4 regarding human recombinant IL-17A. Cells were grown on DMEM culture medium with added 10% inactivated fetal serum, gentamycin and glutamine. Aliquots of 5×10^4 cells per well were seeded in 96-well culture plates. Cells were allowed for attachment for 5 hours. The mixture of 40 ng/ml recombinant IL-17 and 20 ng/ml TNF- α was incubated with VHHIgG1 dilutions for 1 hour at 37°C. Then cytokine/antibody mixture was added to the cells and left overnight. The production of IL-6 by HT1080 cell culture was proportional to the amount of IL-17 added. The amount of the released IL-6 in cell supernatant samples was evaluated by ELISA technique using DuoSet ELISA Development System Human IL6 (RD System, Cat. No. DY206). The results of evaluation of the antagonistic properties of VHHIgG1 candidates are presented at Figure 7 in comparison with AIN457 (anti-IL-17A antibody by Novartis). The IC₅₀ value for mut1 was almost 30 times higher than that for the natural variant, while mut4 variant almost completely maintained its inhibiting capacity. The value of IC₅₀ for mut4 candidate was 30 ± 10 pM. Following the results of the present study and the overall physical-chemical and biological characteristics, the said candidate was selected for further development and optimization.

Example 12

Engineering of mut4VHHFab antibodies comprising human light chains

The total RNA of B-lymphocytes collected from 55 human donors was isolated with RNeasy Mini Kit, in accordance with the appropriate protocol (QIAGEN). RNA concentration was measured with Nanovue kit (GE Healthcare), and the quality of isolated RNA was tested by 1.5% agarose gel electrophoresis.

The reverse transcription reaction was held with MMLV RT kit (Evrogen), according to the recommended protocol, with MMuLV reverse transcriptase and random hexamer primers.

The reverse transcription products were used as a matrix in a two-stage polymerase chain reaction to obtain the genes of variable domains flanked with restriction sites; the reaction was performed using oligonucleotide kit and protocols described in [27]. Chimeric Fab specific against IL-17A were generated according to the procedure described in WO93/06213 based on phagemid pH5, as specified above. Genes encoding variable domains of the human light chains and genes encoding the variable domain of mut4VHH were put together in one fragment by means of sequential reactions of restriction, ligation and amplification as shown at Figure 8. In this case, the estimated count of matrix molecules in all reactions was no less than 10^{12} . The DNA product obtained (VL-CK-VH) was treated with NheI /Eco91I restriction enzymes and ligated into original phagemid pH 5. The ligation products were transformed into SS320 electrocompetent cells prepared in accordance with protocols described in [30]. The repertoire of chimeric mut4 VHH Fab-library based on mixed kappa and lambda human chains was $2.8 \cdot 10^9$ transformants. Chimeric phage-displayed Fab-libraries were prepared in accordance with the procedure described above [27].

The selection of phage-displayed chimeric Fab-libraries was performed under conditions described above (refer to Example 5), except for the additional incubation of IL17A-bound phage antibodies at a temperature of 37°C within 12 hours in the presence of 20 µg/ml of IL-17A dissolved.

After the second selection round at IL-17A, significant enrichment of the library was observed. The obtained pooled clones of enriched chimeric mut4 VHH Fab-libraries were used in IL-17A screening, according to the standard protocol (refer to Example 6). The final positive clones were subject to sequencing. Figure 8B shows four sequences of the variable domains of human light chains VK1A7, VK3cl8, VK3cl18 and VK4clE12 from high-affinity mut4 VHHFab. These sequences belong to different kappa human germ lines VK1, VK3 and VK4

comprising the minimum number of somatic mutations. It should be noted that human sequence VK4c1E12 shows high homology to the previously selected llama light chain VK4B1, yet it has its specific distinctions.

Example 13

Comparative analysis of thermal-aggregation characteristics of full-size mut4VHHIgG1 antibodies comprising various light chain variants

The genes of the variable domain of the heavy VHH chains of mut4VHH candidate were cloned in pEE-Hc plasmid, and genes of variable domains of human light chains VK1A7, VK3c18, VK3c18, VK3A4 and VK4E12 were cloned in pEE-Lc plasmid for the joint transient expression in CHO-EBNA cells, as described in Example 3. Further, the obtained antibodies were exposed to thermal stress and their aggregation profile was studied as shown in Example 9. The results obtained are presented in a summary Table 4.

Table 4. Thermal stability of mut4 VHHIgG1 full-size antibodies

Table 4. Thermal stability of mut4 VHHIgG1 full-size antibodiesSa mple name	Structural bloodline of the light chain	% Monomer before the thermal stress	% Monomer after the thermal stress	Differen ce ~Δ%	Formation of visible aggregate s during the purificatio n	Melting point (Thermofl uor)
mut4VHHIg G1VK3c8	VK3	96	82	14	+	58C
mut4VHHIg G1VK3c18	VK3	91	81	10	++	61C

mut4VHHIg G1VK3A4	VK3	71	69	2	+	61C
mut4VHHIg G1VK1A7	VK1	95	88	7	-	63C
mut4VHHIg G1VK4E12	VK4	90	49	41	++	59C

Furthermore, the comparative thermal stability study of the products obtained was conducted using Thermofluor procedure similar to that described in Example 11. Based on the data obtained, the conclusion can be made that the selected mut4VHH pairs comprising human light chains were more stable than the natural variant comprising VVK4B11 llama light chain. Thus, in the comparative study the best aggregation stability parameters were demonstrated by mut4 VHHIgG1VK1A7 and mut4 VHHIgG1VK3cl8 combinations.

Example 14

Cell test for blocking the ability of IL-17A to induce IL-6 production using mut4 VHHIgG1VK3c8 and mut4 VHHIgG1VK1A7

The ability of IL-17 to induce the production of IL-6 by human HT1080 cells (ATCC:CCL-121) was used to analyze the neutralizing capacity of VHHIgG1 candidates mut4 VHHIgG1VK3c8 and mut4 VHHIgG1VK1A7 regarding human recombinant IL-17 (refer to Example 12). Figure 17 shows the results obtained for the blocking test. It should be noted that mut4 VHHIgG1VK1A7 variant (characterized by maximum stability) demonstrated a significant drop of IC₅₀ value, while mut4 VHHIgG1VK3c8 variant with medium stability exceeded by several times the inhibition observed for more stable mut4 VHHIgG1VK1A7 variant.

Example 15

Engineering of mut4 VHHIgG1 full-size antibodies comprising various mutations at positions 44 and 45 in FR2 of VHH variable domain, and comparative analysis of aggregation and functional characteristics thereof

For the purpose of further improvement of the aggregation stability, the method of oligonucleotide-directed mutagenesis was used to introduce substitutes at positions 44 and 45 in FR2 region of VHH for mut4 VHHIgG1VK3c8 candidate (hereafter referred to as m4VHHc8). The study did not include variants with aromatic, aliphatic or positive amino acids at position 44, as they are potentially immunogenic and structurally forbidden. Mutants were transiently expressed as described in Example 3 (Figure 15, 16). Though the amount of products for different mutants varies from one experiment to another, it should be noted that yields within the same experiment were of comparable size for all substitutes. The obtained mutant variants were exposed to thermal stress. Table 5 demonstrates the results obtained for different mutants. It can be seen from the results that 44G45G and 44G45D are less stable mutants the stability of which is lower than that of the initial 44G45R combination within m4VHHc8. Such variants as 44G45N, 44V45T, 44D45T, 44T45T and 44D45V mainly preserved their stability. All other combinations increased the aggregation stability of m4VHHc8, and the most stable were the variants comprising aromatic or aliphatic groups at position 45. However, the variants comprising small hydrophilic and hydrophobic amino acids at both positions, such as 44G45V, 44G45T, 44V45V, 44A45V, 44T45V, 44A45T and 44S45T, have also demonstrated significant stability.

Table 5. Results of transient expression and thermal stress study of mutants comprising substitutes at positions 44 and 45 in FR2 region of m4VHHc8.

No.	Mutant at position 44+45 of FR2 region of m4VHHc8		Monome r.%, initial*	Monome r.%, heated*	~Δ%	Production yield for the product**, (No. of experiment) ≈, mg/L	IC50, picomole** *as defined by the cell IL6 test
1	44G	45R	96	81	15	41 (1)	50
2	44G	45F	98	98	0	22 (3)	1100
3	44G	45W	97	97	0	57 (2)	1200
4	44G	45Y	97	97	0	33 (2)	650
5	44G	45P	98	98	0	41 (1)	800
6	44G	45I	97	97	0	20 (3)	950
7	44G	45 L	97	97	0	25 (3)	1000
8	44G	45G	96	50	46	21 (3)	ND
9	44G	45N	90	77	13	26 (2)	ND
10	44G	45S	93	85	8	45 (2)	100
11	44G	45D	90	40	50	23 (3)	ND
12	44G	45A	93	89	4	27 (3)	100
13	44G	45H	93	93	2	25 (3)	150
14	44G	45V	93	93	0	35 (1)	250
15	44G	45Q	91	84	7	36 (1)	ND
16	44G	45T	93	93	0	46 (1)	100
17	44V	45T	95	84	11	64 (4)	ND
18	44D	45T	95	83	12	72 (4)	ND
19	44T	45T	96	86	10	75 (4)	150
20	44A	45T	95	93	2	74 (4)	150

21	44S	45T	96	94	2	63 (4)	150
22	44V	45V	95	95	0	71 (4)	150
23	44D	45V	96	88	12	28 (4)	150
24	44A	45V	95	95	0	72 (4)	100
25	44T	45V	96	95	1	73 (4)	150
26	AIN457		95	95	0	0	1100

* The Error in different batches is up to 5%

** Varies significantly depending on the experiment; more correct comparison is provided for products within the same batch

*** Error in different batches is up to 50 %

In order to evaluate the neutralizing activity in the cell test as described in Example 12, experiments were performed with various mutants (Figure 20). The test results are shown in Table 5. It was shown that the small amino acids at VHH FR2 positions 44 and 45 demonstrated high antagonistic activity, while the large ones (aliphatic and aromatic) introduced at position 45 resulted in 10-fold reduction of IC_{50} . Figure 21 presents a diagram of stability and functional properties of mutants comprising various amino acid substitutes that determine interaction of the variable domain of the heavy and light chains.

Example 16

Scanning mutagenesis of CDRs antiIL17A 3VHH domain

Mutations of individual positions of candidate's CDRs were inserted by means of NNK randomization technique [26] with Q5® Site-Directed Mutagenesis Kit (NEB), in accordance with the protocol. Plasmid pLL-Fab was used as a matrix. PCR products were fractioned on low-melting agarose and purified at the columns. After ligation, DNA was transformed into *E.coli* expression strain BL21gold (Stratagene). The individual clones obtained were grown by Fab expression in 96-well plates, as described above. Supernatants containing mutant

Fab arms were analyzed by ELISA under the conditions described above and using high-performance Genetix Q-pix2xt and Tecan Freedom EVO200 systems. The concentration of immobilized IL-17A was 0.2 µg/ml. Bound Fab arms were stained with 1:5000 diluted Goat anti-Human IgG (Fab')₂ (HRP) conjugate (Pierce) and TMB+H₂O₂/ H₂SO₄ dye; absorption was measured at 450 nm wavelength.

The results of scanning mutagenesis are presented in Table 6. The Table shows substitutes within CDR corresponding to ≤30% reduction of mutant Fab/human IL-17A binding signal, when compared to the wild type sequence. Thus, such individual mutants or any combinations thereof are the field of the invention.

Table 6. Scanning mutagenesis results

Mutation position	Positive amino acids in mutants
HCDR3	
V94	S, T, A, K, D, G
R95	K
R96	Y, H, W, K, D, G
R97	A, L, M, S, H, V
F98	-
D99	E, G, A, R, V, K, Q
G100	N, S
T100a	G, P, V, R, S, N, K
S100b	V, M, T, L, T, A, H, G, I, C
Y100c	W, S
Y100d	R, L, W, K, A, G, Q, I, V
T100e	A, L, S
G100f	A, L, T, P, N, Q, F, I, D
D107	-
HCDR2	

A50	G, L
S52	-
P52a	A
S53	-
G54	-
G55	S, R, P, D, I, T, E, K , A, L
D56	-
R57	-
I58	-
HCDR1	
S32	N, K, R, E, W, M, Q, D, F, V, L, A
P33	S
M34	I
G35	L, A, I, S, R, V, N, Q, M

Thus, screening was performed for BCD109 antibody CDR amino acid positions that are tolerant to amino acid substitutes. It was demonstrated that the present set of amino acid substitutes does not significantly change the antibody affinity to human IL-17A. The said combination of substitutes can be used to improve various properties of the candidate.

Example 17

Engineering of BCD109 candidate and evaluation of its affinity to IL-17A of various origins

BCD109 antibody was obtained from m4VHHc8 variant comprising 44G45T substitutes (described in Example 16) by introducing humanizing mutations Q5V and R89V to VHH that do not change either the stability or IC₅₀ (no data provided), and three additional mutations in CH2 domain FcIgG1,

232Y/234T/236E, intended to improve the antibody pharmacokinetics. The antibody has been transiently expressed.

The affinity of BCD109 binding to human, monkey and rat IL-17A was studied at OctetRed 96 system (ForteBio). BCD109 was non-specifically immobilized on the surface of amine reactive second-generation sensors (AR2G) according to the standard protocol described in the manufacturer's manual. The test was conducted at a temperature of 30°C and using PBS with 0.1% Tween-20 and 0.1% BSA as a working buffer.

Human, monkey and rat IL-17A were titrated with the working buffer with concentration from 126 nM to 2 nM with an increment of 2.

Binding curves (after subtracting a reference signal) were analyzed with Octet Data Analysis software (Version 7.0), in accordance with the standard procedure and 1:1 interaction model. The results are presented at Figure 22.

It can be seen that BCD109 binds to human and monkey IL-17A with picomolar affinity (Figure 23). Moreover, the candidate does not interact with rat IL-17A (no curves are presented).

Example 18

Determination of aggregation stability of BCD109 under thermal stress

BCD109 preparation of 10 mg/ml in PBS was heated for 6 hours at a temperature of 50°C in accordance with the protocol described in Example 9.

Results represented at Figure 24 show that BCD109 antibody remains stable under thermal stress: the aggregate content was less than 5%.

Example 19

Production of pharmaceutical composition comprising the antibody of the present invention

BCD109 antibody in concentration of 50 mg/ml was dissolved in the required amount of water for injection and pH was brought to 5.5 with citric acid. Solution was filtered (filtration sterilization) and sealed into ampoules.

The product obtained was stable for 6 months with no sedimentation.

Alternatively, BCD109 was dissolved in water containing mannitol as a lyoprotectant, filtered (filtration sterilization) and then subject to freeze-drying. The powder obtained was dispensed to sterile vials. Vials were corked with rubber stoppers and sealed with aluminum caps. The antibody product can be recovered from lyophilizate with water for injection.

References:

1. Porter R.R. 1973. Structural studies of immunoglobulins. *Science*. **180**, 713–716.
2. Padlan E.A. 1994. Anatomy of the antibody molecule. *Mol.Immunol.* **31**, 169–217.
3. Dwek R.A., Sutton B.J., Perkins S.J., Rademacher T.W.1984. Structure–function relationships in immunoglobulins. *Biochem. Soc. Symcl.* **49**, 123–136.
4. Burton D.R. 1985. Immunoglobulin G: functional sites. *Mol. Immunol.* **22**, 161–206.
5. Hamers Casterman C., Atarhouch T., Muyldermans S., Robinson G., Hamers C., Bajyana Songa E., Bendahman N., Hamers R. 1993. Naturally occurring antibodies devoid of light chains. *Nature*.**363**, 446–448.
6. S.V. Tillib *Molecular Biology*, 2011, 45, No.1, 77–85. Camel nanlantibodies – efficient instrument for investigations, diagnostics and therapy.
7. Nguyen V.K., Hamers R., Wyns L., Muyldermans S.1999. Loss of splice consensus signal is responsible for the removal of the entire CH1 domain of the functional camel IGG2A heavy–chain antibodies. *Mol. Immunol.* 36, 515–524.
8. Woolven B.CL., Frenken L., van der Logt CL., Nicholls CL.J. 1999. The structure of the llama heavy chain constant genes reveals a mechanism for heavy–chain antibody formation. *Immunogenetics*.50, 98–101.
9. Nguyen V.K., Hamers R., Wyns L., Muyldermans S.2000. Camel heavy–chain antibodies: diverse germline VHH and specific mechanisms enlarge the antigen–binding repertoire. *EMBO J.* 19, 921–931.
10. De Genst E., Saerens D., Muyldermans S., Conrath K. 2006. Antibody repertoire development in camelids. *Develocl. Comcl. Immunol.* 30, 187–198.
11. Muyldermans S., Cambillau C., Wyns L. 2001. Recognition of antigens by single–domain antibody fragments: the superfluous luxury of paired domains. *TIBS*. 26, 230–235.

12. De Genst E., Silence K., Decanniere K., Loris R., Kinne J., Muyldermans S. 2006. Molecular basis for the preferential cleft recognition by dromedary heavy chain antibodies. *Proc. Natl. Acad. Sci. USA*. **103**, 4586–4591.
13. Kabat E., Wu T.T., Perry H.M., Gottesman K.S., Foeller C. 1991. Sequence of proteins of immunological interest. *US Public Health Services*. NIH, Bethesda, MD, Publication no. 91–3242.
14. Nguyen V.K., Desmyter A., Muyldermans S. 2001. Functional heavy-chain antibodies in camelidae. *Adv. Immunol.* **79**, 261–296.
15. Muyldermans S., Baral T.N., Retamozzo V.C., et al. 2009. Camelid immunoglobulins and nanobody technology. *Vet. Immunol. Immunopathol.* **128** (1–3), 178–183.
16. de Genst E., Silence K., Decanniere K., Loris R., Kinne J., Wyns L., Muyldermans S. 2005. Strong *in vivo* maturation compensates for structurally restricted H3 loops in antibody repertoires. *J. Biol. Chem.* **280**, 14114–14121.
17. Lauwereys M., Ghahroudi M., Desmyter A., Kinne J., Holzer W., De Genst E., Wyns L., Muyldermans S. 1998. Potent enzyme inhibitors derived from dromedary heavy chain antibodies. *EMBO J.* **17**, 3512–3520.
18. Kontermann RE. 2009 Strategies to extend plasma half-lives of recombinant antibodies. *BioDrugs*.; 23(2):93-109.
19. Ken Coppieters, Torsten Dreier, Karen Silence, Hans de Haard, Marc Lauwereys, Peter Casteels, Els Beirnaert, Heidi Jonckheere, 2006 Formatted anti-tumor necrosis factor alpha VHH proteins derived from camelids show superior potency and targeting to inflamed joints in a murine model of collagen-induced arthritis *Arthritis & Rheumatology*; 54(6):1856-66.
20. Gabrielle Richard, Ashley J Meyers, Michael D McLean, Mehdi Arbabi-Ghahroudi, Roger MacKenzie, J Christopher Hall 2013. *In Vivo* Neutralization of α -Cobratoxin with High-Affinity Llama Single-Domain Antibodies (V_HHs) and a V_HH-Fc Antibody. *PLoS One* 22;8(7):e69495.
21. Jan Terje Andersen, Maria Gonzalez-Pajuelo, Stian Foss, Ole J. B. Landsverk, Débora Pinto, Alexander Szyroki, Hans J. de Haard, Michael Saunders, Peter

Vanlandschoot Inger Sandlie 2012, Selection of Nanobodies that Target Human Neonatal Fc Receptor. *Scientific Reports*, 3, 1118.

22. Vincke C¹, Loris R, Saerens D, Martinez-Rodriguez S, Muyldermans S, Conrath K. 2009 General strategy to humanize a camelid single-domain antibody and identification of a universal humanized nanobody scaffold. *J Biol Chem*. Jan 30; 284(5): 3273-84.

23. Barthelemy PA, Raab H, Appleton BA, Bond CJ, Wu P, Wiesmann C, Sidhu SS. 2008 Comprehensive analysis of the factors contributing to the stability and solubility of autonomous human VH domains. *J Biol Chem*. Feb 8;283(6):3639-54.

24. Conrath K, Vincke C, Stijlemans B, Schymkowitz J, Decanniere K, Wyns L, Muyldermans S, Loris R. Antigen binding and solubility effects upon the veneering of a camel VHH in framework-2 to mimic a VH. *J Mol Biol*. 2005 Jul 1; 350 (1): 112-25.

25. Hermeling S, Crommelin DJ, Schellekens H, Jiskoot W (2004) Structure–immunogenicity relationships of therapeutic proteins. *Pharm Res* 21:897–903

26. Handbook of pharmaceutical manufacturing formulations. Volume 6. Sterile products / Sarfaraz K. Niazi, 2004.

27. J. de Haard H., van Neer N., Anneke Reurs, E. Hufton S., Roovers R., Henderikx CL., de Bruïne A., Arends J.-W. and Hoogenboom H. A Large Non-immunized Human Fab Fragment Phage Library That Permits Rapid Isolation and Kinetic Analysis of High Affinity Antibodies. *J Biol Chem*. 1999 Jun 25 274(26):18218-30.

28. WO 2010/001251.

29. Koch-Nolte F., Reyelt J., Schöbnow B., Schwarz N., Scheuplein F., Rothenburg S., Haag F., Alzogaray V., Cauerhff A. and Goldbaum F. Single domain antibodies from llama effectively and specifically block T cell ecto-ADP-ribosyltransferase ART2.2 in vivo. *FASEB J.*, 2007 Nov; 21(13):3490-8.

30. Sidhu S., Lowman H., Cunningham B., Wells J. Phage display for selection of novel binding peptides. *J. Methods Enzymol*. 2000; 328: 333-63.

31. Marks J., Hoogenboom H., Bonnert T., McCafferty J., Griffiths A., Winter G. By-passing immunization: Human antibodies from V-gene libraries displayed on phage. *J Mol Biol.* 1991 Dec 5; 222(3):581-97.
32. Vaughan T., Williams A., Pritchard K., Osbourn J., Pope A., Earnshaw J., McCafferty J., Hodits R., Wilton J. and Johnson K. Human Antibodies with Sub-nanomolar Affinities Isolated from a Large Non-immunized Phage Display Library. *Nat Biotechnol.* 1996 Mar; 14 (3): 309-14.
34. *Environ Microbiol.* 2006 Feb;72(2):1141-7;
MAbs. 2012 May-Jun; 4(3): 341-8.
35. Cummings M., Farnum M., and Nelen M. Universal Screening Methods and Applications of ThermoFluor®. *J Biomol Screen* 2006 11(7): 854-863.

CLAIMS:

1. A humanized monoclonal IgG antibody wherein variable domains are represented by a combination of a V_{HH} -derivative with a variable domain of the light chain V_L .

2. The antibody according to claim 1, characterized by increased aggregation stability of V_{HH} -derivative compared to the native VHH antibody isolated from an immunized animal from Camelidae family.

3. The antibody according to claim 1, wherein the V_{HH} -derivative comprises the following amino acid substitutes at positions 44X₂45X₃:

a) wherein 44X₂ = G, A, V, S, T;

b) wherein 45X₃ = A, V, T, H;

or combinations thereof.

4. The antibody according to claim 3, is characterized by higher aggregation stability of the V_{HH} -derivative, as compared to the native VHH antibody isolated from an immunized animal of Camelidae family.

5. The antibody according to claim 1, wherein the V_{HH} -derivative is represented by a variable domain of the heavy chain isolated from an immunized animal of Camelidae family and comprises additional amino acid substitutes typical of humans and located at any positions, except for the following substitutes at positions 44X₂45X₃:

c) wherein 44X₂ = G, A, V, S, T;

d) wherein 45X₃ = A, V, T, H;

or combinations thereof.

6. The antibody according to claim 1, wherein the V_{HH} -derivative is a variable domain of the heavy chain isolated from a non-immunized animal of Camelidae family, and comprises additional amino acid substitutes typical of humans and located at any positions, except for the following substitutes at positions 44X₂45X₃:

e) wherein 44X₂ = G, A, V, S, T;

f) wherein 45X₃ = A, V, T, H;

or combinations thereof.

7. The antibody according to claim 1, wherein a variable domain of the light chain V_L is a human antibody derivative.

8. The antibody according to claim 1, wherein a variable domain of the light chain V_L is a humanized fragment of an antibody of animal, preferably mammalian.

9. The antibody according to claim 1, wherein the V_{HH} -derivative that contains cysteine-44 (Kabat numbering scheme), and the light chain variable domain V_L contains cysteine-100 (Kabat numbering scheme).

10. The antibody according to claim 1, wherein the said antibody is one of any of the following isotypes: IgG1, IgG2, IgG3 or IgG4.

11. The antibody according to claim 10, comprising a non-native modified Fc as a part of IgG.

12. The antibody according to claim 1, is characterized by such aggregation stability that when it is used in concentrations over 10 mg/ml and stored for >6 months at a temperature of $T=4^{\circ}\text{C}$, the content of aggregates increases by not more than 5% of their initial content in the solution.

13. The antibody according to claim 1, is characterized by such aggregation stability that when it is used in concentrations over 10 mg/ml and stored for >2 weeks at a temperature of 37⁰C, the content of aggregates increases by not more than 5% of their initial content in the solution.

14. The antibody according to claim 1, is characterized by such aggregation stability that when it is used in concentrations over 10 mg/ml and stored for >48h at a temperature of 42⁰C the content of aggregates increases by not more than 5% of their initial content in the solution.

15. The antibody according to claim 1, is characterized by such aggregation stability that when it is used in concentrations over 10 mg/ml and stored for > 6 h at a temperature of 50⁰C the content of aggregates increases by not more than 5% of their initial content in the solution.

16. The antibody according to claim 1, wherein dissociation constant K_D is 10⁻⁹ M and lower.

17. The antibody according to claim 1, wherein the antibody-antigen interaction kinetic association constant k_{on} (1/Ms) is at least 10⁵ 1/Ms and more.

18. The antibody according to claim 1, wherein the antibody-antigen interaction kinetic dissociation constant (1/s) is 10⁻⁴ 1/s and lower.

19. The antibody according to claims 1-18 that specifically binds to human IL-17A, comprising

a) a derivative of Camelidae heavy chain variable domain (V_{HH}) that comprises 3 hypervariable regions HCDR1, HCDR2 and HCDR3, wherein:

HCDR1 comprising the amino acid sequence of SEQ ID NO: 1:

G-T-F-A-T-X32-X33-X34 - X35, wherein

X32 is an amino acid selected from the group comprising S, N, K, R, E, W, Q, D, A, V and F;

X33 is an amino acid selected from the group comprising P or S;

X34 is an amino acid selected from the group comprising M and I;

X35 is an amino acid selected from the group comprising G, N, S, A, L, I, R, V and Q;

HCDR2 comprising the amino acid sequence of SEQ ID NO: 2:

X50-I-X52-X52a-S-G-X55-D-R-I-Y-A-D-S-V-K-G, wherein

X50 is an amino acid selected from the group comprising A, G and L;

X52 is an amino acid selected from the group comprising S, D and E;

X52a is an amino acid selected from the group comprising P and A;

X55 is an amino acid selected from the group comprising G, S, T, L, R, D, E, K, A and W;

HCDR3 comprising the amino acid sequence of SEQ ID NO: 3:

C-A- X94-X95- X96- X97-F-X99-X100-X100a- X100b - X100c - X100d - X100e - X100f – D- Y – D-S, wherein

X94 is an amino acid selected from the group comprising K, S, T, V, D and G;

X95 is an amino acid selected from the group comprising R and K;

X96 is an amino acid selected from the group comprising G, R, Y, H, D, W and K;

X97 is an amino acid selected from the group comprising R, A, V, S, L and H;

X99 is an amino acid selected from the group comprising D, E, G, A, R, V, K and Q;

X100 is an amino acid selected from the group comprising G, S and N;

X100a is an amino acid selected from the group comprising G, T, P, V, R, N and K;

X100b is an amino acid selected from the group comprising V, S, T, L, Y, A, H, G and I;

X100c is an amino acid selected from the group comprising Y, W and S;

X100d is an amino acid selected from the group comprising R, V, L, Y, A, W, K, G, Q and I;

X100e is an amino acid selected from the group comprising T, L, A and S;

X100f is an amino acid selected from the group comprising T, L, G, P, N, A, Q, F, I and D;

b) a variable domain of the light chain (V_L) of a human antibody or a variable domain of the light chain of a humanized antibody.

20. The antibody according to claims 1-18 that specifically binds to human IL-17A, comprising:

a) A derivative of the heavy chain variable domain (V_{HH}) comprising 3 hypervariable regions HCDR1, HCDR2 и HCDR3, wherein:

HCDR1 comprising the amino acid sequence of G-T-F-A-T-S-P-M-G (SEQ ID NO: 4);

HCDR2 comprising the amino acid sequence of A-I-S-P-S-G-G- D-R-I-Y-A-D-S-V-K-G (SEQ ID NO: 5);

HCDR3 comprising the amino acid sequence of C-A-V-R-R-R-F-D-G-T-S-Y-Y-T-G-D-Y-D-S (SEQ ID NO: 6);

b) A variable domain of the light chain (V_L) of a human antibody or a variable domain of the light chain of a humanized antibody.

21. The antibody according to claim 20 that specifically binds to human IL-17A, comprising:

a) A derivative of the heavy chain variable domain (V_{HH}) comprising an amino acid sequence

QVQLVQSGGGLVQAGGSLRLSCAASGGTFATSPMGWLRQAPGKGT
EFVAAISPSGGDRIYADSVKGRFTISRDNAGYFIYLMNSLKPEDTAVYYC
AVRRRFDGTSYYTGDYDSWGQGTLVTVSS (SEQ ID NO: 7);

b) A variable domain of the light chain (V_L) of a human antibody or a variable domain of the light chain of a humanized antibody.

22. The antibody according to claim 21 that specifically binds to human IL-17A, comprising V_{HH} -derivative, wherein said variable domain comprises an amino acid sequence at least 90% identical to SEQ ID NO: 7.

23. The antibody according to claim 22 that specifically binds to human IL-17A, comprising:

a) The V_{HH} -derivative comprising the amino acid sequence of SEQ ID NO: 7;

b) The variable domain of the light chain (V_L) of a human antibody comprising the amino acid sequence of

EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLL
IYDASSRATGIPDRFSGSGGTDFTLISRLEPEDFAVYYCQQYSYSPVTFGQ
GTKVEIKR (SEQ ID NO: 8).

24. The antibody according to claim 23 that specifically binds human IL-17A and comprises V_{HH} -derivative and a variable region of the light chain (V_L) of a human antibody, wherein said variable domains comprise an amino acid sequence at least 90% identical to SEQ ID NO: 8.

25. The antibody according to claims 1-18 that specifically binds human IL-17A and comprises:

a) The heavy chain comprising the following amino acid sequence:

QVQLVQSGGGLVQAGGSLRLSCAASGGTFATSPMGWLRQAPGKGT
EFVAAISPSGGDRIYADSVKGRFTISRDNAGYFIYLMNSLKPEDTAVYYC
AVRRRFDGTSYYTGDYDSWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGT
AALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPS
SSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVF
LFPPKPKDTLYITREPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKP
REEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG
QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY
KTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL
SLSPGK (SEQ ID NO: 9).

b) The variable domain of a human antibody light chain (V_L) comprising the following amino acid sequence:

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRL
IYDASSRATGIPDRFSGSGGTDFTLISRLEPEDFAVYYCQQYSYSPVTFGQ
GTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVD
NALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGL
SSPVTKSFNRGEC (SEQ ID NO: 10).

26. The antibody according to claim 25 that specifically binds human IL-17A and contains the heavy chain and the light chain, wherein said chains comprise amino acid sequences at least 90% identical to SEQ ID NO: 9 and/or SEQ ID NO:10.

27. The antibody according to claim 19, wherein said antibody has the binding affinity to human IL-17A characterized with K_D of 10^{-10} M and lower.

28. The antibody according to claim 19, wherein the kinetic association constant $k_{on}(1/Ms)$ for human IL-17A is not less than 10^5 1/Ms.

29. The antibody according to claim 19, wherein the kinetic dissociation constant $k_{\text{dis}}(1/\text{s})$ for human IL-17A is not more than $10^{-5} 1/\text{s}$.

30. The antibody according to claim 19, wherein said antibody inhibits the activity of human IL-17A by no less than 50%.

31. The antibody according to claim 19, wherein said antibody is produced by mammalian, yeast or bacterial cells.

32. The antibody according to claim 19 comprising one or more additional amino acid substitutions in Fc-region compared to the native Fc, wherein said substitutions improve physical-chemical and pharmacokinetic properties of an antibody, as compared to an antibody with the native Fc, and do not result in the loss of antibody's ability to bind IL-17A.

33. A fragment of an antibody according to claims 1- 32.

34. The fragment of an antibody according to claim 33, wherein said fragment is represented by a light chain, heavy chain, and variable domains of the light and/or heavy chain that are a part of an antibody sequence, which includes a bio-specific antibody variant.

35. The fragment of an antibody according to claim 34, wherein said fragment is represented by a light chain, heavy chain, and variable domains of the light and/or heavy chain that are parts of Fab.

36. The fragment of an antibody according to claim 34, wherein said fragment is represented by a light chain, heavy chain, and variable domains of the light and/or heavy chain that are parts of scFv.

37. Methods of production of antibodies according to any of claims 1 – 32 or fragments thereof according to claims 33-36, involving phases selected from the following: directed mutagenesis, display methods, genetic engineering, biochemistry and high-performance biotechnology methods well known for the art, which can include the methods for directed mutagenesis at different positions of the VHH domain of Camelidae antibodies.

38. A DNA construct encoding the antibody according to any of claims 1 - 32 or the antibody fragment according to any of claims 33-36.

39. An expression vector comprising one or several DNA constructs according to claim 38.

40. A cell line comprising the expression vector according to claim 39 or DNA construct according to claim 38.

41. A method of production of the humanized monoclonal antibody according to claims 1- 32 or the antibody fragment according to claims 33-36, which involves incubation of a cell line according to claim 40 in a culture medium under the conditions sufficient to obtain said antibody or fragment thereof, followed by the isolation and purification of obtained antibody or its active fragment.

42. A host cell comprising the vector according to claim 39.

43. A method of production of the antibody according to any of claims 1-32 or the antibody fragment according to claims 33-36, which involves incubation of the host cell according to claim 42 in a culture medium under the conditions

sufficient to obtain said antibody or fragment thereof, followed by isolation and purification of obtained antibody or its active fragment.

44. A pharmaceutical composition comprising the antibody according to any of claims 1-32 or fragments thereof according to claims 33-36, in combination with one or several pharmaceutically suitable excipients, diluents or carriers.

45. The pharmaceutical composition according to claim 44, further comprising active pharmaceutical ingredients selected from TNF- α inhibitors.

46. The pharmaceutical composition according to claim 44, to be used in the therapy of IL-17A-mediated disease or disorder.

47. The pharmaceutical composition according to claim 46, wherein an IL-17A-mediated disease or disorder is selected from: rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroidenitis, asthma, allergic disorders, psoriasis, dermatitis, systemic sclerosis, graft-versus-host disease, graft rejection, acute or chronic immune disease associated with organ grafting, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki disease, Graves' disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's disease, Henoch-Schonlein purpura, microscopic polyangiitis with renal involvement, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infections, invasions, acquired immune deficiency syndrome, acute transverse myelitis, Huntington chorea, Parkinson disease, Alzheimer disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure, myocardial infarction, Addison disease, polyglandular autoimmune syndrome type I and type II, Schmidt's syndrome, acute respiratory distress syndrome, alopecia, alopecia

areata, seronegative arthropathy, arthropathy, Reiter's syndrome, psoriatic arthropathy, arthropathy associated with ulcerative colitis, enteropathic synovitis, arthropathy associated with Chlamydia, Yersinia and Salmonella, spondyloarthropathy, atheromatosis disease/coronary sclerosis, atopic allergy, autoimmune bullous disease, pemphigus, pemphigus foliaceus, pemphigoid, linear IgA diseases, autoimmune hemolytic anemia, Coombs positive hemolytic anemia, acquired pernicious anemia, juvenile pernicious anemia, Myalgic encephalomyelitis/chronic fatigue syndrome, chronic active hepatic inflammation, cranial giant arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, acquired immune deficiency syndrome (AIDS), AIDS-associated diseases, hepatitis B, hepatitis C, common variable immunodeficiency (common variable hypogammaglobulinemia), dilated cardiomyopathy, female sterility, ovarian insufficiency, Premature ovarian failure, pulmonary fibrosis, cryptogenic fibrosing alveolitis, post inflammatory interstitial lung pathologies, interstitial pneumonitis, connective tissue disease associated with interstitial lung disease, mixed connective tissue disease associated with interstitial lung disease, systemic scleroderma associated with interstitial lung disease, rheumatoid arthritis associated with interstitial lung disease, systemic lupus erythematosus associated with lung disease, dermatomyositis/polymyositis associated with lung disease, Sjogren disease associated with lung disease, ankylosing spondylitis associated with lung disease, diffuse pulmonary vasculitis, hemosiderosis associated with lung disease, drug-induced interstitial lung disease, fibrosis, radiation-induced fibrosis, obliterating bronchiolitis, chronic eosinophilic pneumonia, lung disease with lymphocyte infiltration, post infectious interstitial lung pathologies, gouty arthritis, autoimmune hepatitis, autoimmune hepatitis type I (classic autoimmune or lupoid hepatitis), autoimmune hepatitis type II (associated with anti-LKM antibody), autoimmune hypoglycemia, type B insulin resistance with acanthokeratoderma, hypoparathyroidism, acute graft-associated immune disease, chronic graft-associated immune disease, osteoarthritis, primary sclerosing cholangitis, type I psoriasis, type II psoriasis, idiopathic leukopenia, autoimmune

neutropenia, NOS-kidney diseases, glomerulonephritis, microscopic renal polyangiitis, Lyme disease, discoid lupus erythematosus, idiopathic of NOS-male sterility, antisperm immunity, multiple sclerosis (all types), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue disease, Goodpasture syndrome, pulmonary manifestations of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease, systemic scleroderma, Sjogren's Syndrome, Takayasu disease/arthritis, autoimmune thrombocytopenia, idiopathic thrombocytopenia, autoimmune thyroid disorders, hyperthyroid, autoimmune hypothyroidism (Hashimoto disease), atrophic autoimmune hypothyroidism, primary myxedema, phacogenic uveitis, primary vasculitis, vitiligo, acute hepatic disease, chronic hepatic disease, alcoholic cirrhosis, alcohol-induced liver damage, cholestasis, idiosyncratic hepatic disease, drug-induced hepatitis, nonalcoholic steatohepatitis, allergies and asthma, group B streptococcal infection (GBS), mental disorders (including depressions and schizophrenia), Th1- and Th2-mediated disease, acute and chronic pain (various forms), malignancies such as lung cancer, breast cancer, stomach cancer, bladder cancer, colorectal cancer, pancreatic cancer, ovarian cancer, prostate cancer and hematopoietic malignancies (leukemia and lymphomas), abetalipoproteinaemia, acrocyanosis, acute and chronic infections and infestations, acute leukemia, acute lymphoblastic leukemia, acute myeloid leukemia, acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinoma, atrial ectopics, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic rhinitis, allograft rejection, alpha-I antitrypsin deficiency, lateral amyotrophic sclerosis, anemia, angina, anterior horn cell degeneration, anti-CD3 therapy, antiphospholipid syndrome, hypersensitivity reactions against receptors, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, coronary sclerosis, arteriovenous fistula, ataxia, atrial fibrillation (constant or paroxysmal), atrial flutter, atrioventricular block, B-cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt lymphoma, burns, cardiac arrhythmia, myocardial stunning syndrome, cardiac

tumor, cardiomyopathy, inflammatory response to bypass, cartilage graft rejection, brain cortex degeneration, cerebellar disorder, chaotic or multifocal atrial tachycardia, chemotherapy-induced disorders, chronic myelocytic leukemia (CML), chronic alcohol addiction, chronic inflammatory pathologies, chronic lymphatic leukemia (CLL), chronic obstructive pulmonary disease, chronic salicylate intoxication, rectocolic carcinoma, congestive cardiac failure, conjunctivitis, contact dermatitis, pulmonary heart, coronary artery disease, Creutzfeldt-Jakob Disease, culture-negative sepsis, cystic fibrosis, cytokine therapy-induced disorders, boxer's encephalopathy, demyelinating disease, dengue hemorrhagic fever, dermatitis, dermatological conditions, diabetes, diabetes mellitus, diabetes-related atherosclerotic vascular disease, diffuse Lewy body disease, congestive dilated cardiomyopathy, basal ganglia disease, Down's syndrome in middle age, motor disorders induced by CNS dopamine blockers, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, epiglottiditis, Epstein-Barr viral infection, erythralgia, extrapyramidal and cerebellar symptoms, familial hemophagocytic lymphohistiocytosis, fetal thymus graft rejection, Friedreich's ataxia, peripheral artery disease, fungal sepsis, gas phlegmon, gastric ulcer, glomerulonephritis, any organ or tissue graft rejection, gram negative sepsis, gram positive sepsis, granulomas due to intracellular organisms, hairy-cell leukemia, Hallervorden-Spatz disease, Hashimoto's thyroiditis, hay fever, heart transplant rejection, hemochromatosis, hemodialysis, hemolytic uremic syndrome / thrombotic thrombocytopenic purpura, bleeding, hepatitis (A), bundle branch arrhythmia, HIV-infections/HIV-neuropathies, Hodgkin disease, hyperkinetic motor disorders, hypersensitivity reactions, hypersensitivity-associated pneumonitis, hypertension, hypokinetic motor disorders, examination of hypothalamo-pituitary-adrenal axis, idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody-mediated cytotoxicity, asthenia, infantile muscular atrophy, aortal inflammation, influenza virus A, exposure to ionizing radiation, iridocyclitis /uveitis/optic neuritis, ischaemia/reperfusion-induced disorders, ischaemic stroke, juvenile rheumatoid arthritis, juvenile spinal

muscular atrophy, Kaposi's sarcoma, renal transplant rejection, legionellosis, leishmaniasis, leprosy, corticospinal damage, lipoedema, liver transplant rejection, lymphoedema, malaria, malignant lymphoma, malignant histiocytosis, malignant melanoma, meningitis, meningococemia, metabolic/idiopathic diseases, migraine, multiple system mitochondrial disorders, mixed connective-tissue disease, monoclonal gammopathy, multiple myeloma, multiple system degeneration (Mencel Dejerine-Thomas Shi-Drager and Machado-Joseph), myasthenia gravis, intracellular Mycobacterium avium, Mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial ischemic disease, nasopharyngeal cancer, neonatal chronic lung disease, nephritis, nephrotic, neurodegenerative disorders, neurogenic muscular atrophy I, neutropenic fever, non-Hodgkin's lymphomas, abdominal aortic branch occlusion, arterial occlusive disease, OKT3® treatment, orchitis/epididymitis, orchitis/vasectomy reversal operations, organomegaly, osteoporosis, pancreatic graft rejection, pancreatic carcinoma, paraneoplastic disease /tumor-related hypercalcemia, parathyroid graft rejection, pelvic inflammatory disease, perennial rhinitis, pericardial disease, peripheral atherosclerosis (atherosclerotic) disease, peripheral vascular disease, peritonitis, pernicious anemia, Pneumocystis carinii pneumonia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal plasma-proliferative disorder and skin changes), postperfusion syndrome, pump head syndrome, post-cardiotomy post-infarction syndrome, preeclampsia, progressive supranuclear paralysis, primary pulmonary hypertension, radiation therapy, Raynaud's phenomenon and disease, Raynoud's disease, Refsum's disease, regular narrow QRS tachycardia, renal vascular hypertension, reperfusion injury, restrictive cardiomyopathy, sarcoma, scleroderma disease, senile chorea, Dementia with Lewy bodies, seronegative arthritis, shock, sickle cell disease, skin allograft rejection, skin changes, small intestinal graft rejection, solid tumors, specific arrhythmias, spinal ataxia, spinocerebellar degradations, streptococcal myositis, cerebellar structural damage, subacute sclerosing panencephalitis, syncope, cardiovascular syphilis, systemic anaphylaxis, a comprehensive systemic

inflammatory response syndrome, systemic-onset juvenile rheumatoid arthritis, T cells or FAB ALL, telangiectasia, thrombosis obliterans, thrombocytopenia, toxicity, grafting, trauma/bleeding, hypersensitivity reactions type III, hypersensitivity reactions type IV, unstable angina, uremia, urinary sepsis, urticaria, valvular heart disease, varicose veins, vasculitis, venous diseases, venous thrombosis, ventricular fibrillation, viral and fungal infections, viral encephalitis / aseptic meningitis, viral hemophagocytic syndrome, Wernicke-Korsakoff syndrome, Wilson's disease, heterograft rejection for any organ or tissue, acute coronary syndrome, acute idiopathic polyneuritis, acute inflammatory demyelinating radicular neuropathy, acute ischemia, adult-onset Stills disease, alopecia areata, anaphylaxis, antiphospholipid antibody syndrome, aplastic anemia, coronary sclerosis, atopic eczema, atopic dermatitis, autoimmune dermatitis, autoimmune disorder associated with streptococcus infection, autoimmune enteropathy, autoimmune hearing loss, autoimmune lymphoproliferative syndrome (ALPS), autoimmune myocarditis, autoimmune premature ovarian failure, blepharitis, bronchiectasis, bullous pemphigoid, cardiovascular disease, catastrophic antiphospholipid syndrome, celiac disease, cervical spondylosis, chronic ischemia, cicatricial pemphigoid, clinically isolated syndrome (cis) with the risk for multiple sclerosis, conjunctivitis, childhood-onset mental disorders, chronic obstructive pulmonary disease (COPD), dacryocystitis, dermatomyositis, diabetic retinopathy, diabetes mellitus, herniated disk, prolapse of intervertebral disc, drug-induced immune haemolytic anaemia, endocarditis, endometreosis, entophthalmia, episcleritis, erythema multiform, severe erythema multiform, gestational pemphigoid , Guillain-Barre syndrome (GBS), hay fever, Hughes syndrome, idiopathic Parkinson's disease, idiopathic interstitial pneumonia, IgE-mediated allergy, autoimmune hemolytic anemia, inclusion body myositis, infectious ocular inflammatory disease, inflammatory demyelinating disease, inflammatory heart disease, inflammatory kidney disease, idiopathic pulmonary fibrosis/ usual interstitial pneumonia, iritis, keratitis, keratoconjunctivitis sicca, Kussmaul disease or Kussmaul-Meier Disease, Landry palsy, Langerhans' cell

histiocytosis, marbled skin, macular degeneration, microscopic polyangiitis, Bechterew disease, motor neuron disease, mucosal pemphigoid, multiple organ failure, myasthenia gravis, spinal cord dysplasia syndrome, myocarditis, nerve root disorders, neuropathy, non-A, non-B hepatitis, optic neuritis, osteolysis, ovarian cancer, oligoarticular JIA, peripheral arterial occlusive disease, peripheral vascular disease, peripheral artery disease (PAD), phlebitis, polyarteritis nodosa, polychondritis, polymyalgia rheumatica, poliosis, polyarticular juvenile idiopathic arthritis, multiple endocrine deficiency, polymyositis, polymyalgia rheumatica (PMR), post pump syndrome, primary parkinsonism, prostate and rectal cancer and hematopoietic malignancies (leukemia and lymphoma), prostatitis, pure red-cell aplasia, primary adrenal insufficiency, relapsing neuromyelitis optica, restenosis, rheumatic heart disease, SAPHO (synovitis, acne, pustulosis, hyperostosis and osteitis), scleroderma, secondary amyloidosis, shock lung, scleritis, ischias, secondary adrenal insufficiency, silicon-associated connective tissue disease, Sneddon-Wilkinson disease, ankylosing spondylitis, Stevens–Johnson syndrome, systemic inflammation response syndrome, cranial arteritis, Toxoplasma rhinitis, toxic epidermal necrolysis, transverse myelitis, TRAPS (tumor necrosis factor receptor-associated periodic syndrome), allergic reactions type I, diabetes type II, urticaria, usual interstitial pneumonia, vasculitis, vernal conjunctivitis, viral retinitis, Vogt-Koyanagi-Harada syndrome (VKH syndrome), wet macular degeneration, wound healing, and Yersinia- or Salmonella-associated arthropathy.

48. A method for the treatment of IL-17A-mediated disease or disorder, characterized by administration of a therapeutically effective amount of the isolated antibody according to claims 1-32 or fragment thereof according to claims 33-36 to a patient who needs such a treatment.

49. The method for the treatment according to claim 48, wherein an IL-17A-mediated disease or disorder is selected from: rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis, reactive

arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroadenitis, asthma, allergic disorders, psoriasis, dermatitis, systemic sclerosis, graft-versus-host disease, graft rejection, acute or chronic immune disease associated with organ grafting, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki disease, Graves' disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's disease, Henoch-Schonlein purpura, microscopic polyangiitis with renal involvement, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infections, invasions, acquired immune deficiency syndrome, acute transverse myelitis, Huntington chorea, Parkinson disease, Alzheimer disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure, myocardial infarction, Addison disease, polyglandular autoimmune syndrome type I and type II, Schmidt's syndrome, acute respiratory distress syndrome, alopecia, alopecia areata, seronegative arthropathy, arthropathy, Reiter's syndrome, psoriatic arthropathy, arthropathy associated with ulcerative colitis, enteropathic synovitis, arthropathy associated with Chlamydia, Yersinia and Salmonella, spondyloarthropathy, atheromatosis disease/coronary sclerosis, atopic allergy, autoimmune bullous disease, pemphigus, pemphigus foliaceus, pemphigoid, linear IgA diseases, autoimmune hemolytic anemia, Coombs positive hemolytic anemia, acquired pernicious anemia, juvenile pernicious anemia, Myalgic encephalomyelitis/chronic fatigue syndrome, chronic active hepatic inflammation, cranial giant arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, acquired immune deficiency syndrome (AIDS), AIDS-associated diseases, hepatitis B, hepatitis C, common variable immunodeficiency (common variable hypogammaglobulinemia), dilated cardiomyopathy, female sterility, ovarian insufficiency, Premature ovarian failure, pulmonary fibrosis, cryptogenic fibrosing alveolitis, post inflammatory interstitial lung pathologies, interstitial pneumonitis, connective tissue disease associated with interstitial lung disease, mixed connective tissue disease associated with interstitial lung disease, systemic scleroderma associated with interstitial lung

disease, rheumatoid arthritis associated with interstitial lung disease, systemic lupus erythematosus associated with lung disease, dermatomyositis/polymyositis associated with lung disease, Sjogren disease associated with lung disease, ankylosing spondylitis associated with lung disease, diffuse pulmonary vasculitis, hemosiderosis associated with lung disease, drug-induced interstitial lung disease, fibrosis, radiation-induced fibrosis, obliterating bronchiolitis, chronic eosinophilic pneumonia, lung disease with lymphocyte infiltration, post infectious interstitial lung pathologies, gouty arthritis, autoimmune hepatitis, autoimmune hepatitis type I (classic autoimmune or lupoid hepatitis), autoimmune hepatitis type II (associated with anti-LKM antibody), autoimmune hypoglycemia, type B insulin resistance with acanthokeratoderma, hypoparathyroidism, acute graft-associated immune disease, chronic graft-associated immune disease, osteoarthritis, primary sclerosing cholangitis, type I psoriasis, type II psoriasis, idiopathic leukopenia, autoimmune neutropenia, NOS-kidney diseases, glomerulonephritis, microscopic renal polyangiitis, Lyme disease, discoid lupus erythematosus, idiopathic of NOS-male sterility, antisperm immunity, multiple sclerosis (all types), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue disease, Goodpasture syndrome, pulmonary manifestations of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease, systemic scleroderma, Sjogren's Syndrome, Takayasu disease/arthritis, autoimmune thrombocytopenia, idiopathic thrombocytopenia, autoimmune thyroid disorders, hyperthyroid, autoimmune hypothyroidism (Hashimoto disease), atrophic autoimmune hypothyroidism, primary myxedema, phacogenic uveitis, primary vasculitis, vitiligo, acute hepatic disease, chronic hepatic disease, alcoholic cirrhosis, alcohol-induced liver damage, cholestasis, idiosyncratic hepatic disease, drug-induced hepatitis, nonalcoholic steatohepatitis, allergies and asthma, group B streptococcal infection (GBS), mental disorders (including depressions and schizophrenia), Th1- and Th2-mediated disease, acute and chronic pain (various forms), malignancies such as lung cancer, breast cancer, stomach cancer, bladder cancer, colorectal cancer, pancreatic cancer, ovarian cancer, prostate cancer and hematopoietic

malignancies (leukemia and lymphomas), abetalipoproteinaemia, acrocyanosis, acute and chronic infections and infestations, acute leukemia, acute lymphoblastic leukemia, acute myeloid leukemia, acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinoma, atrial ectopics, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic rhinitis, allograft rejection, alpha-I antitrypsin deficiency, lateral amyotrophic sclerosis, anemia, angina, anterior horn cell degeneration, anti-CD3 therapy, antiphospholipid syndrome, hypersensitivity reactions against receptors, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, coronary sclerosis, arteriovenous fistula, ataxia, atrial fibrillation (constant or paroxysmal), atrial flutter, atrioventricular block, B-cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt lymphoma, burns, cardiac arrhythmia, myocardial stunning syndrome, cardiac tumor, cardiomyopathy, inflammatory response to bypass, cartilage graft rejection, brain cortex degeneration, cerebellar disorder, chaotic or multifocal atrial tachycardia, chemotherapy-induced disorders, chronic myelocytic leukemia (CML), chronic alcohol addiction, chronic inflammatory pathologies, chronic lymphatic leukemia (CLL), chronic obstructive pulmonary disease, chronic salicylate intoxication, rectocolic carcinoma, congestive cardiac failure, conjunctivitis, contact dermatitis, pulmonary heart, coronary artery disease, Creutzfeldt-Jakob Disease, culture-negative sepsis, cystic fibrosis, cytokine therapy-induced disorders, boxer's encephalopathy, demyelinating disease, dengue hemorrhagic fever, dermatitis, dermatological conditions, diabetes, diabetes mellitus, diabetes-related atherosclerotic vascular disease, diffuse Lewy body disease, congestive dilated cardiomyopathy, basal ganglia disease, Down's syndrome in middle age, motor disorders induced by CNS dopamine blockers, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, epiglottiditis, Epstein-Barr viral infection, erythralgia, extrapyramidal and cerebellar symptoms, familial hemophagocytic lymphohistiocytosis, fetal thymus graft rejection, Friedreich's ataxia, peripheral artery disease, fungal sepsis, gas

phlegmon, gastric ulcer, glomerulonephritis, any organ or tissue graft rejection, gram negative sepsis, gram positive sepsis, granulomas due to intracellular organisms, hairy-cell leukemia, Hallervorden-Spatz disease, Hashimoto's thyroiditis, hay fever, heart transplant rejection, hemochromatosis, hemodialysis, hemolytic uremic syndrome / thrombotic thrombocytopenic purpura, bleeding, hepatitis (A), bundle branch arrhythmia, HIV-infections/HIV-neuropathies, Hodgkin disease, hyperkinetic motor disorders, hypersensitivity reactions, hypersensitivity-associated pneumonitis, hypertension, hypokinetic motor disorders, examination of hypothalamo-pituitary-adrenal axis, idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody-mediated cytotoxicity, asthenia, infantile muscular atrophy, aortal inflammation, influenza virus A, exposure to ionizing radiation, iridocyclitis /uveitis/optic neuritis, ischaemia/reperfusion-induced disorders, ischaemic stroke, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, renal transplant rejection, legionellosis, leishmaniasis, leprosy, corticospinal damage, lipoedema, liver transplant rejection, lymphoedema, malaria, malignant lymphoma, malignant histiocytosis, malignant melanoma, meningitis, meningococemia, metabolic/idiopathic diseases, migraine, multiple system mitochondrial disorders, mixed connective-tissue disease, monoclonal gammopathy, multiple myeloma, multiple system degeneration (Mencel Dejerine-Thomas Shi-Drager and Machado-Joseph), myasthenia gravis, intracellular Mycobacterium avium, Mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial ischemic disease, nasopharyngeal cancer, neonatal chronic lung disease, nephritis, nephrotic, neurodegenerative disorders, neurogenic muscular atrophy I, neutropenic fever, non-Hodgkin's lymphomas, abdominal aortic branch occlusion, arterial occlusive disease, OKT3® treatment, orchitis/epididymitis, orchitis/vasectomy reversal operations, organomegaly, osteoporosis, pancreatic graft rejection, pancreatic carcinoma, paraneoplastic disease /tumor-related hypercalcemia, parathyroid graft rejection, pelvic inflammatory disease, perennial rhinitis, pericardial disease, peripheral atherosclerosis (atherosclerotic) disease, peripheral vascular disease, peritonitis,

pernicious anemia, Pneumocystis carinii pneumonia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal plasma-proliferative disorder and skin changes), postperfusion syndrome, pump head syndrome, post-cardiotomy post-infarction syndrome, preeclampsia, progressive supranuclear paralysis, primary pulmonary hypertension, radiation therapy, Raynaud's phenomenon and disease, Raynoud's disease, Refsum's disease, regular narrow QRS tachycardia, renal vascular hypertension, reperfusion injury, restrictive cardiomyopathy, sarcoma, scleroderma disease, senile chorea, Dementia with Lewy bodies, seronegative arthritis, shock, sickle cell disease, skin allograft rejection, skin changes, small intestinal graft rejection, solid tumors, specific arrhythmias, spinal ataxia, spinocerebellar degradations, streptococcal myositis, cerebellar structural damage, subacute sclerosing panencephalitis, syncope, cardiovascular syphilis, systemic anaphylaxis, a comprehensive systemic inflammatory response syndrome, systemic-onset juvenile rheumatoid arthritis, T cells or FAB ALL, telangiectasia, thrombosis obliterans, thrombocytopenia, toxicity, grafting, trauma/bleeding, hypersensitivity reactions type III, hypersensitivity reactions type IV, unstable angina, uremia, urinary sepsis, urticaria, valvular heart disease, varicose veins, vasculitis, venous diseases, venous thrombosis, ventricular fibrillation, viral and fungal infections, viral encephalitis / aseptic meningitis, viral hemophagocytic syndrome, Wernicke-Korsakoff syndrome, Wilson's disease, heterograft rejection for any organ or tissue, acute coronary syndrome, acute idiopathic polyneuritis, acute inflammatory demyelinating radicular neuropathy, acute ischemia, adult-onset Stills disease, alopecia areata, anaphylaxis, antiphospholipid antibody syndrome, aplastic anemia, coronary sclerosis, atopic eczema, atopic dermatitis, autoimmune dermatitis, autoimmune disorder associated with streptococcus infection, autoimmune enteropathy, autoimmune hearing loss, autoimmune lymphoproliferative syndrome (ALPS), autoimmune myocarditis, autoimmune premature ovarian failure, blepharitis, bronchiectasis, bullous pemphigoid, cardiovascular disease, catastrophic antiphospholipid syndrome, celiac disease, cervical spondylosis,

chronic ischemia, cicatrical pemphigoid, clinically isolated syndrome (cis) with the risk for multiple sclerosis, conjunctivitis, childhood-onset mental disorders, chronic obstructive pulmonary disease (COPD), dacryocystitis, dermatomyositis, diabetic retinopathy, diabetes mellitus, herniated disk, prolapse of intervertebral disc, drug-induced immune haemolytic anaemia, endocarditis, endometreosis, entophthalmia, episcleritis, erythema multiform, severe erythema multiform, gestational pemphigoid , Guillain-Barre syndrome (GBS), hay fever, Hughes syndrome, idiopathic Parkinson's disease, idiopathic interstitial pneumonia, IgE-mediated allergy, autoimmune hemolytic anemia, inclusion body myositis, infectious ocular inflammatory disease, inflammatory demyelinating disease, inflammatory heart disease, inflammatory kidney disease, idiopathic pulmonary fibrosis/ usual interstitial pneumonia, iritis, keratitis, keratoconjunctivitis sicca, Kussmaul disease or Kussmaul-Meier Disease, Landry palsy, Langerhans' cell histiocytosis, marbled skin, macular degeneration, microscopic polyangiitis, Bechterew disease, motor neuron disease, mucosal pemphigoid, multiple organ failure, myasthenia gravis, spinal cord dysplasia syndrome, myocarditis, nerve root disorders, neuropathy, non-A, non-B hepatitis, optic neuritis, osteolysis, ovarian cancer, oligoarticular JIA, peripheral arterial occlusive disease, peripheral vascular disease, peripheral artery disease (PAD), phlebitis, polyarteritis nodosa, polychondritis, polymyalgia rheumatica, poliosis, polyarticular juvenile idiopathic arthritis, multiple endocrine deficiency, polymyositis, polymyalgia rheumatica (PMR), post pump syndrome, primary parkinsonism, prostate and rectal cancer and hematopoietic malignancies (leukemia and lymphoma), prostatitis, pure red-cell aplasia, primary adrenal insufficiency, relapsing neuromyelitis optica, restenosis, rheumatic heart disease, SAPHO (synovitis, acne, pustulosis, hyperostosis and osteitis), scleroderma, secondary amyloidosis, shock lung, scleritis, ischias, secondary adrenal insufficiency, silicon-associated connective tissue disease, Sneddon-Wilkinson disease, ankylosing spondylitis, Stevens–Johnson syndrome, systemic inflammation response syndrome, cranial arteritis, Toxoplasma rhinitis, toxic epidermal necrolysis, transverse myelitis, TRAPS (tumor necrosis factor

receptor-associated periodic syndrome), allergic reactions type I, diabetes type II, urticaria, usual interstitial pneumonia, vasculitis, vernal conjunctivitis, viral retinitis, Vogt-Koyanagi-Harada syndrome (VKH syndrome), wet macular degeneration, wound healing, and Yersinia- or Salmonella-associated arthropathy.

50. The method for treatment according to 49, further comprising the administration of TNF- α inhibitors.