


SUBSTITUTED AMINOTETRALINES

The present invention relates to novel substituted (5-amino-5,6,7,8-tetrahydro-naphthalene-1-yloxy)-acetic acids, their manufacture, pharmaceutical compositions
5 containing them and their use as CRTH2 antagonists.

Prostaglandin D₂ (PGD₂) is the major prostanoid produced by activated mast cells and has been implicated in the pathogenesis of allergic diseases such as allergic asthma and atopic dermatitis. Chemoattractant Receptor-homologous molecule
10 expressed on T-helper type cells (CRTH2) is one of the prostaglandin D₂ receptors and is expressed on the effector cells involved in allergic inflammation such as T helper type 2 (Th2) cells, eosinophils, and basophils (Nagata *et al.*, *FEBS Lett* 459: 195-199, 1999). It has been shown to mediate PGD₂-stimulated chemotaxis of Th2 cells, eosinophils, and basophils (Hirai *et al.*, *J Exp Med* 193: 255-261, 2001).
15 Moreover, CRTH2 mediates the respiratory burst and degranulation of eosinophils (Gervais *et al.*, *J Allergy Clin Immunol* 108: 982-988, 2001), induces the production of proinflammatory cytokines in Th2 cells (Xue *et al.*, *J Immunol* 175: 6531-6536), and enhances the release of histamine from basophils (Yoshimura-Uchiyama *et al.*, *Clin Exp Allergy* 34:1283-1290). Sequence variants of the gene encoding CRTH2,
20 which differentially influence its mRNA stability, are shown to be associated with asthma (Huang *et al.*, *Hum Mol Genet* 13, 2691-2697, 2004). Increased numbers of circulating T cells expressing CRTH2 have also been correlated with severity of atopic dermatitis (Cosmi *et al.*, *Eur J Immunol* 30, 2972-2979, 2000). These findings suggest that CRTH2 plays a proinflammatory role in allergic diseases. Therefore,
25 antagonists of CRTH2 are believed to be useful for treating disorders such as asthma, allergic inflammation, COPD, and atopic dermatitis.

- 2 -

The invention is concerned with the compounds of formula I:



- 5 and pharmaceutically acceptable salts and esters thereof, wherein R^1 - R^4 are defined in the detailed description and claims. In addition, the present invention relates to methods of manufacturing and using the compounds of formula I as well as pharmaceutical compositions containing such compounds. The compounds of formula I are antagonists at the CRTH2 receptor and may be useful in treating
- 10 diseases and disorders associated with that receptor such as asthma.

Unless otherwise indicated, the following specific terms and phrases used in the description and claims are defined as follows:

- 15 The term “moiety” refers to an atom or group of chemically bonded atoms that is attached to another atom or molecule by one or more chemical bonds thereby forming part of a molecule. For example, the variables R^1 , R^2 , R^3 and R^4 of formula I refer to moieties that are attached to the core structure of formula I by a covalent bond.

20

In reference to a particular moiety with one or more hydrogen atoms, the term “substituted” refers to the fact that at least one of the hydrogen atoms of that moiety is replaced by another substituent or moiety. For example, the term “lower alkyl substituted by halogen” refers to the fact that one or more hydrogen atoms of a lower

- 3 -

alkyl (as defined below) is replaced by one or more halogen atoms (e.g., trifluoromethyl, difluoromethyl, fluoromethyl, chloromethyl, etc.).

The term "optionally substituted" refers to the fact that one or more hydrogen atoms
5 of a moiety (with one or more hydrogen atoms) can be, but does not necessarily have to be, substituted with another substituent.

The term "alkyl" refers to an aliphatic straight-chain or branched-chain saturated hydrocarbon moiety having 1 to 20 carbon atoms. In particular embodiments the
10 alkyl has 1 to 10 carbon atoms.

The term "lower alkyl" refers to an alkyl moiety having 1 to 7 carbon atoms. In particular embodiments the lower alkyl has 1 to 4 carbon atoms and in other particular embodiments the lower alkyl has 1 to 3 carbon atoms. Examples of lower
15 alkyls include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and *tert*-butyl.

The term "lower cycloalkyl" refers to a saturated or partly unsaturated non-aromatic hydrocarbon ring moiety having 3 to 7 carbon atoms bonded together to form a ring structure. Examples of cycloalkyls include cyclopropyl, cyclobutyl, cyclopentyl,
20 cyclohexyl and cycloheptyl.

The term "lower alkoxy" refers to the moiety -O-R, wherein R is lower alkyl as defined previously. Examples of lower alkoxy moieties include methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, isobutoxy, sec-butoxy and *tert*-butoxy.
25

The term "lower alkanoyl" refers to the moiety -C(O)-R, wherein R is lower alkyl as defined previously. An example of a lower alkanoyl is acetyl.

The term "lower alkylsulfanyl" refers to the moiety -S-R, wherein R is lower alkyl as defined previously. Examples of lower alkylsulfanyls include methylsulfanyl and ethylsulfanyl.
30

- 4 -

The term "lower cycloalkylsulfanyl" refers to the moiety -S-R, wherein R is lower cycloalkyl as defined previously. Examples of lower cycloalkylsulfanyls include cyclopropylsulfanyl, cyclobutylsulfanyl and cyclopentylsulfanyl.

- 5 The term "lower alkylsulfinyl" refers to the moiety -S(O)-R, wherein R is lower alkyl as defined previously. Examples of lower alkylsulfinyls include methylsulfinyl and ethylsulfinyl.

- 10 The term "lower cycloalkylsulfinyl" refers to the moiety -S(O)-R, wherein R is lower cycloalkyl as defined previously. Examples of lower cycloalkylsulfinyls include cyclopropylsulfinyl, cyclobutylsulfinyl and cyclopentylsulfinyl.

- 15 The term "lower alkylsulfonyl" refers to the moiety -S(O)₂-R, wherein R is lower alkyl as defined previously. Examples of lower alkylsulfonyls include methylsulfonyl and ethylsulfonyl.

- The term "lower cycloalkylsulfonyl" refers to the moiety -S(O)₂-R, wherein R is lower cycloalkyl as defined previously. Examples of lower cycloalkylsulfonyls include cyclopropylsulfonyl, cyclobutylsulfonyl and cyclopentylsulfonyl.

20

The term "halogen" refers to a moiety of fluoro, chloro, bromo or iodo.

- 25 Unless otherwise indicated, the term "hydrogen" or "hydro" refers to the moiety of a hydrogen atom (-H) and not H₂.

25

Unless otherwise indicated, the term "a compound of the formula" or "a compound of formula" or "compounds of the formula" or "compounds of formula" refers to any compound selected from the genus of compounds as defined by the formula (including any pharmaceutically acceptable salt or ester of any such compound).

30

The term "pharmaceutically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. Salts may be formed with inorganic acids such

- 5 -

as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, preferably hydrochloric acid, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, salicylic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, *p*-toluenesulfonic acid, *N*-acetylcystein and the like. In addition, salts may be prepared by the addition of an inorganic base or an organic base to the free acid. Salts derived from an inorganic base include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, and magnesium salts and the like. Salts derived from organic bases include, but are not limited to salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, lysine, arginine, *N*-ethylpiperidine, piperidine, polyamine resins and the like.

15

The compounds of the present invention can be present in the form of pharmaceutically acceptable salts. The compounds of the present invention can also be present in the form of pharmaceutically acceptable esters (i.e., the methyl and ethyl esters of the acids of formula I to be used as prodrugs). The compounds of the present invention can also be solvated, i.e. hydrated. The solvation can be effected in the course of the manufacturing process or can take place i.e. as a consequence of hygroscopic properties of an initially anhydrous compound of formula I (hydration).

20

Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers." Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers." Diastereomers are stereoisomers with opposite configuration at one or more chiral centers which are not enantiomers. Stereoisomers bearing one or more asymmetric centers that are non-superimposable mirror images of each other are termed "enantiomers." When a compound has an asymmetric center, for example, if a carbon atom is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be

25

30

- 6 -

characterized by the absolute configuration of its asymmetric center or centers and is described by the R- and S-sequencing rules of Cahn, Ingold and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral
5 compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture".

The term "a therapeutically effective amount" of a compound means an amount of compound that is effective to prevent, alleviate or ameliorate symptoms of disease or
10 prolong the survival of the subject being treated. Determination of a therapeutically effective amount is within the skill in the art. The therapeutically effective amount or dosage of a compound according to this invention can vary within wide limits and may be determined in a manner known in the art. Such dosage will be adjusted to the individual requirements in each particular case including the specific
15 compound(s) being administered, the route of administration, the condition being treated, as well as the patient being treated. In general, in the case of oral or parenteral administration to adult humans weighing approximately 70 Kg, a daily dosage of about 0.1 mg to about 5,000 mg, 1 mg to about 1,000 mg, or 1 mg to 100 mg may be appropriate, although the lower and upper limits may be exceeded when
20 indicated. The daily dosage can be administered as a single dose or in divided doses, or for parenteral administration, it may be given as continuous infusion.

The term "pharmaceutically acceptable carrier" is intended to include any and all material compatible with pharmaceutical administration including solvents, dispersion
25 media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and other materials and compounds compatible with pharmaceutical administration. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions of the invention is contemplated. Supplementary active compounds can also be
30 incorporated into the compositions.

Useful pharmaceutical carriers for the preparation of the compositions hereof, can be solids, liquids or gases; thus, the compositions can take the form of tablets, pills,

- 7 -


capsules, suppositories, powders, enterically coated or other protected formulations (e.g. binding on ion-exchange resins or packaging in lipid-protein vesicles), sustained release formulations, solutions, suspensions, elixirs, aerosols, and the like. The carrier can be selected from the various oils including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, and the like. Water, saline, aqueous dextrose, and glycols are preferred liquid carriers, particularly (when isotonic with the blood) for injectable solutions. For example, formulations for intravenous administration comprise sterile aqueous solutions of the active ingredient(s) which are prepared by dissolving solid active ingredient(s) in water to produce an aqueous solution, and rendering the solution sterile. Suitable pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, talc, gelatin, malt, rice, flour, chalk, silica, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk, glycerol, propylene glycol, water, ethanol, and the like. The compositions may be subjected to conventional pharmaceutical additives such as preservatives, stabilizing agents, wetting or emulsifying agents, salts for adjusting osmotic pressure, buffers and the like. Suitable pharmaceutical carriers and their formulation are described in Remington's Pharmaceutical Sciences by E. W. Martin. Such compositions will, in any event, contain an effective amount of the active compound together with a suitable carrier so as to prepare the proper dosage form for proper administration to the recipient.

In the practice of the method of the present invention, an effective amount of any one of the compounds of this invention or a combination of any of the compounds of this invention or a pharmaceutically acceptable salt or ester thereof, is administered via any of the usual and acceptable methods known in the art, either singly or in combination. The compounds or compositions can thus be administered orally (e.g., buccal cavity), sublingually, parenterally (e.g., intramuscularly, intravenously, or subcutaneously), rectally (e.g., by suppositories or washings), transdermally (e.g., skin electroporation) or by inhalation (e.g., by aerosol), and in the form of solid, liquid or gaseous dosages, including tablets and suspensions. The administration can be conducted in a single unit dosage form with continuous therapy or in a single dose therapy ad libitum. The therapeutic composition can also be in the form of an oil

- 8 -

emulsion or dispersion in conjunction with a lipophilic salt such as pamoic acid, or in the form of a biodegradable sustained-release composition for subcutaneous or intramuscular administration.

5 In detail, the present invention relates to the compounds of formula I:



and pharmaceutically acceptable salts and esters thereof, wherein:

10

R^1 is hydrogen, halogen, methoxy, or phenyl; and R^2 is hydrogen, halogen, or alkyl; with the proviso that R^1 and R^2 are not both hydrogen;

R^3 is hydrogen or methyl; and

15

R^4 is phenyl or pyridine substituted by one or two substituents independently selected from the group consisting of:


- (1) halogen;
- (2) lower alkyl optionally substituted by halogen,
- (3) lower alkanoyl;
- (4) lower alkoxy;
- (5) lower alkylsulfanyl, lower alkylsulfinyl, or lower alkylsulfonyl
- (6) lower cycloalkylsulfanyl, lower cycloalkylsulfinyl, or lower cycloalkylsulfonyl; and

20

- 9 -


(7) phenyl or pyridine, wherein said phenyl or pyridine is optionally substituted by lower alkyl, lower alkylsulfanyl, lower alkylsulfonyl, lower cycloalkylsulfanyl, or lower cycloalkylsulfonyl.

- 5 Unless indicated otherwise, the genus of formula I and any subgenera thereof encompass all possible stereoisomers (i.e., (*R*)-enantiomers and (*S*)-enantiomers) as well as racemic and scalemic mixtures thereof. In one embodiment of the invention, the compounds of formula I are (*R*)-enantiomers or pharmaceutically acceptable salts or esters thereof as depicted in the following subgeneric structural
- 10 formula IA for the (*R*)-enantiomers of formula I:



wherein R¹-R⁴ are as defined previously.

- In another embodiment, the compounds of formula I are (*S*)-enantiomers or
- 15 pharmaceutically acceptable salts or esters thereof as depicted in the following subgeneric structural formula IB for the (*S*)-enantiomers of formula I:



wherein R^1 - R^4 are as defined previously.

5 In another embodiment the present invention is directed to a composition comprising a mixture (racemic or otherwise) of the (*R*)-enantiomers and (*S*)-enantiomers of a compound of formula I.

10 In a more particular embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^3 is hydrogen.

15 In a more particular embodiment the present invention is directed to the compounds of formula IA or pharmaceutically acceptable salts or esters thereof, wherein R^3 is hydrogen.

In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^3 is methyl.

20 In a more particular embodiment the present invention is directed to the compounds of formula IA or pharmaceutically acceptable salts or esters thereof, wherein R^3 is methyl.

25 In one embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^1 is hydrogen, fluoro, chloro, bromo, methoxy, or phenyl and R^2 is hydrogen, fluoro, chloro, methyl, ethyl, propyl, or isopropyl; with the proviso that R^1 and R^2 are not both hydrogen.

30 In a more particular embodiment the present invention is directed to the compounds of formula IA or pharmaceutically acceptable salts or esters thereof, wherein R^1 is hydrogen, fluoro, chloro, bromo, methoxy, or phenyl and R^2 is hydrogen, fluoro, chloro, methyl, ethyl, propyl, or isopropyl; with the proviso that R^1 and R^2 are not both hydrogen.

- 11 -

In a more particular embodiment the present invention is directed to the compounds of formula IA or pharmaceutically acceptable salts or esters thereof, wherein R^1 is hydrogen, fluoro, chloro, bromo, methoxy, or phenyl and R^2 is hydrogen or fluoro; with the proviso that R^1 and R^2 are not both hydrogen.

5

In one embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein at least one of R^1 or R^2 is halogen.

- 10 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^1 is fluoro or chloro.

- In another embodiment the present invention is directed to the compounds of
15 formula I or pharmaceutically acceptable salts or esters thereof, wherein R^1 is hydrogen and R^2 is fluoro.

- In another embodiment the present invention is directed to the compounds of
formula I or pharmaceutically acceptable salts or esters thereof, wherein R^1 is
20 hydrogen and R^2 is chloro.

- In another embodiment the present invention is directed to the compounds of
formula I or pharmaceutically acceptable salts or esters thereof, wherein R^1 is
hydrogen and R^2 is methyl.

25

In another embodiment the present invention is directed to the compounds of
formula I or pharmaceutically acceptable salts or esters thereof, wherein R^1 is
hydrogen and R^2 is ethyl.

- 30 In another embodiment the present invention is directed to the compounds of
formula I or pharmaceutically acceptable salts or esters thereof, wherein R^1 is
hydrogen and R^2 is propyl.

- 12 -

In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is hydrogen and R² is isopropyl.

- 5 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is fluoro and R² is hydrogen.

- 10 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is fluoro and R² is fluoro.

- 15 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is fluoro and R² is chloro.

- 20 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is fluoro and R² is methyl.

- In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is fluoro and R² is ethyl.

- 25 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is fluoro and R² is propyl.

- 30 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is fluoro and R² is isopropyl.

- 13 -

In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is chloro and R² is hydrogen.

- 5 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is chloro and R² is fluoro.

- 10 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is chloro and R² is chloro.

- 15 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is chloro and R² is methyl.

- 20 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is chloro and R² is ethyl.

- In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is chloro and R² is propyl.

- 25 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is chloro and R² is isopropyl.

- 30 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is bromo and R² is hydrogen.

- 14 -

In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is bromo and R² is fluoro.

- 5 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is bromo and R² is chloro.

- 10 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is bromo and R² is methyl.

- 15 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is bromo and R² is ethyl.

- 20 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is bromo and R² is propyl.

- In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is bromo and R² is isopropyl.

- 25 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is methoxy and R² is hydrogen.

- 30 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is methoxy and R² is fluoro.

- 15 -

In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is methoxy and R² is chloro.

- 5 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is methoxy and R² is methyl.

- 10 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is methoxy and R² is ethyl.

- 15 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is methoxy and R² is propyl.

- 20 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is methoxy and R² is isopropyl.

- In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is phenyl and R² is hydrogen.

- 25 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is phenyl and R² is fluoro.

- 30 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is phenyl and R² is chloro.

- 16 -

In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is phenyl and R² is methyl.

- 5 In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is phenyl and R² is ethyl.

- In another embodiment the present invention is directed to the compounds of
10 formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is phenyl and R² is propyl.

- In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R¹ is phenyl
15 and R² is isopropyl.

- In another particular embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is phenyl substituted by one or two substituents as defined previously.
20

- In another embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is pyridine substituted by one or two substituents as defined previously.

- 25 In a more particular embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is phenyl substituted by one or two substituents independently selected from the group consisting of:

- (1) halogen; and
30 (2) lower alkyl optionally substituted by halogen.

In a more specific embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is phenyl

- 17 -

substituted by one or two substituents independently selected from the group consisting of:

- (1) fluoro;
- (2) chloro;
- 5 (3) bromo;
- (4) lower alkyl
- (5) lower alkanoyl;
- (6) lower alkylsulfanyl, lower alkylsulfinyl, or lower alkylsulfonyl;
- (7) lower cycloalkylsulfanyl, lower cycloalkylsulfinyl, or lower
- 10 cycloalkylsulfonyl;
- (8) trifluoromethyl, difluoromethyl, or fluoromethyl; and
- (9) 1,1-difluoroethyl.
- (10) phenyl optionally substituted by lower alkyl or lower alkylsulfonyl; and
- (11) pyridine optionally substituted by lower alkyl or lower alkylsulfonyl.

15

In a more specific embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is phenyl substituted by one or two substituents independently selected from the group consisting of:

- 20 (1) fluoro;
- (2) chloro;
- (3) bromo;
- (4) methyl;
- (5) ethyl;
- 25 (6) propyl or isopropyl;
- (7) butyl, *sec*-butyl, or *tert*-butyl;
- (8) trifluoromethyl, difluoromethyl, or fluoromethyl; and
- (9) 1,1-difluoroethyl.

30 In a more specific embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is phenyl substituted by one or two substituents independently selected from the group consisting of:

- 18 -

- (1) fluoro;
- (2) chloro;
- (3) bromo; and
- (4) trifluoromethyl.

5

In one embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is pyridine substituted by one or two substituents independently selected from the group consisting of:

- 10 (1) halogen;
- (2) lower alkyl optionally substituted by halogen; and
- (3) phenyl or pyridine, wherein said phenyl or pyridine is optionally substituted by lower alkyl.

- 15 In a more specific embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is pyridine substituted by one or two substituents independently selected from the group consisting of:

- (1) fluoro;
- 20 (2) chloro;
- (3) bromo;
- (4) lower alkyl
- (5) lower alkanoyl;
- (6) lower alkylsulfanyl, lower alkylsulfinyl, or lower alkylsulfonyl;
- 25 (7) lower cycloalkylsulfanyl, lower cycloalkylsulfinyl, or lower cycloalkylsulfonyl;
- (8) trifluoromethyl, difluoromethyl, or fluoromethyl; and
- (9) 1,1-difluoroethyl.
- (10) phenyl optionally substituted by lower alkyl or lower alkylsulfonyl; and
- 30 (11) pyridine optionally substituted by lower alkyl or lower alkylsulfonyl.

In a more specific embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is

- 19 -

pyridine substituted by one or two substituents independently selected from the group consisting of:

- (1) fluoro;
- (2) chloro;
- 5 (3) bromo;
- (4) methyl;
- (5) ethyl;
- (6) propyl or isopropyl;
- (7) butyl, *sec*-butyl, or *tert*-butyl;
- 10 (8) trifluoromethyl, difluoromethyl, or fluoromethyl;
- (9) 1,1-difluoroethyl;
- (10) phenyl optionally substituted by lower alkyl; and
- (11) pyridine optionally substituted by lower alkyl.

- 15 In a more specific embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is pyridine substituted by one substituent independently selected from the group consisting of:

- (1) phenyl optionally substituted by lower alkyl; and
- 20 (2) pyridine optionally substituted by lower alkyl.


In a more specific embodiment the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is pyridine substituted by phenyl wherein said phenyl is substituted by a lower alkyl moiety. In a more particular embodiment said phenyl is substituted by an isopropyl moiety.

25

In particular embodiments, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is phenyl substituted by one or two substituents at positions 2, 3, 4, 5, or 6 where such positions are as indicated below in formula IC:

30

- 20 -



In one embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is phenyl substituted by one substituent at the 2 position and another substituent at the 4 or 5 position (as indicated in formula IC).

In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is phenyl substituted by one substituent at the 2 position and another substituent at the 5 position (as indicated in formula IC).

In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is phenyl substituted by one substituent at the 2 or 6 position and another substituent at the 4 position (as indicated in formula IC).

In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is phenyl substituted by one substituent at the 3 position and another substituent at the 5 position (as indicated in formula IC).

- 21 -

In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is phenyl substituted by one substituent at the 3 or 5 position and another substituent at the 4 position (as indicated in formula IC).

5

In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^1 is fluoro or chloro; R^2 is hydrogen or fluoro; and R^4 is phenyl substituted at positions 3 and 5 (as indicated in formula IC) with substituents independently selected from the group consisting of fluoro, bromo, or trifluoromethyl.


10

In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^1 is fluoro or chloro; R^2 is hydrogen or fluoro; and R^4 is phenyl substituted at positions 3 and 5 (as indicated in formula IC) with trifluoromethyl.

15

In other particular embodiments, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R^4 is pyridine substituted by one or two substituents at positions, 4, 5, or 6 where such positions are as indicated below in formula ID:

20



- 22 -

In one embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is pyridine substituted by one substituent at the 6 position and another substituent at the 4 position (as indicated in formula ID).

5

In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is pyridine substituted by one substituent at the 4 position and another substituent at the 5 position (as indicated in formula ID).

10

In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is pyridine substituted by one substituent at the 5 position (as indicated in formula ID).

15 In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is pyridine substituted by one substituent at the 5 position and another substituent at the 6 position (as indicated in formula ID).

20 In another embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is pyridine substituted by one substituent at the 6 position (as indicated in formula ID).

In another embodiment, the present invention is directed to the compounds of
25 formula I or pharmaceutically acceptable salts or esters thereof, wherein R⁴ is pyridine substituted by phenyl at the 6 position (as indicated in formula ID) wherein said phenyl is itself optionally substituted by lower alkyl.

In a more specific embodiment, the present invention is directed to a compound of
30 formula I selected from the group consisting of:

[(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;

- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-isopropyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 5 [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-ethyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 10 [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 15 5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [3-Chloro-5-(2,5-dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 20 5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [3-Chloro-5-(2,5-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 25 [3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [3-Chloro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 30 Tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;

- 24 -

- [(*R*)-3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 5 [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [3-Chloro-2-fluoro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 10 [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-methoxy-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-phenyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 15 [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [2,3-Difluoro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [3-Fluoro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 20 [5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 25 [5-(2,4-Dichloro-benzenesulfonylamino)-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(2,4-Dichloro-benzenesulfonylamino)-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(2,5-Dichloro-benzenesulfonylamino)-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 30 [5-(2,5-Dichloro-benzenesulfonylamino)-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;

- {2,3-Difluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Fluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 5 {3-Chloro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- [(*R*)-3-Bromo-5-(3-bromo-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-3-Chloro-2-fluoro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 10 [(*R*)-5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 15 [(*R*)-3-Chloro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-3-Chloro-5-(2,5-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 20 {3-Chloro-2-fluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- [5-(2,5-Dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 25 [5-(2,4-Dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- {2-Fluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- [(*R*)-3-Bromo-5-(2,5-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 30 [(*R*)-3-Bromo-5-(2,4-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;

- 26 -

[5-(3,5-Bis-methanesulfonyl-benzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid; and
any pharmaceutically acceptable salt or ester thereof.

- 5 In another specific embodiment, the present invention is directed to a compound of formula I selected from the group consisting of:

- [5-(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 10 {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-ethyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 15 {(R)-5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Chloro-5-[(2,4-dichloro-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 20 {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Chloro-5-[(2,5-dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 25 {3-Chloro-5-[(2,5-dichloro-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Chloro-5-[(2,4-dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 30 {5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;

- 27 -

- {3-Chloro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Chloro-2-fluoro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 5 {5-[(2,4-Dichloro-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 10 {5-[(2,4-Dichloro-benzenesulfonyl)-methyl-amino]-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(2,5-Dichloro-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 15 {5-[(2,5-Dichloro-benzenesulfonyl)-methyl-amino]-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Fluoro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 20 {2,3-Difluoro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- (2,3-Difluoro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid;
- 25 (3-Fluoro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid;
- (3-Chloro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid;
- (3-Chloro-2-fluoro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid;
- 30 {5-[(2,5-Dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;

- 28 -

- {5-[(2,4-Dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
(2-Fluoro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid; and
5 any pharmaceutically acceptable salt or ester thereof.

In another particular embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof except for [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-isopropyl-5,6,7,8-tetrahydro-
10 naphthalen-1-yloxy]-acetic acid; and any pharmaceutically acceptable salt or ester thereof.

In another particular embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof except for {5-[(3,5-
15 Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid; and any pharmaceutically acceptable salt or ester thereof.

In another particular embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof except for {5-[(3,5-
20 Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-ethyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid; and any pharmaceutically acceptable salt or ester thereof.

In another particular embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof except for {5-[(3,5-
25 Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid; and any pharmaceutically acceptable salt or ester thereof.

30

In another particular embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof except for {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-methyl-5,6,7,8-

- 29 -

tetrahydro-naphthalen-1-yloxy}-acetic acid; and any pharmaceutically acceptable salt or ester thereof.

In another particular embodiment, the present invention is directed to the compounds
5 of formula I or pharmaceutically acceptable salts or esters thereof except for {3-Chloro-5-[(2,5-dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid; and any pharmaceutically acceptable salt or ester thereof.

10 In another particular embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof except for {3-Chloro-5-[(2,4-dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid; and any pharmaceutically acceptable salt or ester thereof.

15

In another particular embodiment, the present invention is directed to the compounds of formula I or pharmaceutically acceptable salts or esters thereof except for {5-[(2,4-dichloro-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid; and any pharmaceutically acceptable salt or ester thereof.

20


In a further embodiment the invention provides, the use of a compound of formula I for the preparation of a medicament for the treatment of severe asthma or chronic obstructive pulmonary disease.

25

The compounds of the present invention can be prepared by any conventional means. Suitable processes for synthesizing these compounds are provided in the examples. Generally, compounds of formula I can be prepared according to the
30 schemes illustrated below.

- 30 -

Scheme 1



Compounds of interest **I** and **IA** with substituents on the tetrahydro-naphthalene ring can be prepared according to Scheme 1. Starting with phenols **III** and succinic anhydride (**IV**), esterification followed by Fries rearrangement gives the butyric acid derivatives **VI**. Clemmensen reduction of keto acids **VI** yields compounds of structures **VII**. Methylation followed by ester hydrolysis converts phenols **VII** to the methoxy acid derivatives **IX**, which can form naphthalenones **X** via intramolecular Friedel-Crafts reactions. Demethylation of compounds of structure **X** gives 5-hydroxy-naphthalenones **XI**, which undergo nucleophilic substitution with *tert*-butyl bromoacetate (**XII**) under basic conditions to generate the ethers **XIII**. Reductive

- 31 -

amination reactions of intermediates **XIII** afford the corresponding amino derivatives **XV**. Sulfonylation of amino compounds **XV** with sulfonyl chlorides **XVI** affords the corresponding sulfonamides of structure **XVII**. N-Methylation of the sulfonamides **XVII** gives the corresponding derivatives **XIX**. Ester hydrolysis of either **XVII** or **XIX** produces compounds of interest **I**, which can be resolved to the optically pure (R)-enantiomers **IA** by chiral HPLC.

In the first step of the sequence, nucleophilic substitution reactions of phenols **III** and succinic anhydride (**IV**) give the phenol esters **V**. Subsequent Fries rearrangement of intermediates **V** to yield the *ortho* keto acids **VI** can be carried out in the presence of a strong Lewis acid catalyst such as aluminum chloride, boron trifluoride, or titanium tetrachloride under anhydrous conditions, either neat, or in a solvent such as tetrahydrofuran, dichloromethane, 1,1,2,2-tetrachloroethane, or dichloroethane, at a temperature between room temperature and reflux temperature for several hours.

Conversion of keto acids **VI** to derivatives **VII** can be accomplished under Clemmensen reduction conditions. The reaction is typically carried out using zinc amalgam and hydrochloric acid with or without an auxiliary solvent such as toluene or ether, at a temperature between room temperature and reflux temperature for 3 days.

Methylation of phenols **VII** to compounds **VIII** can be achieved using methods that are well known to someone skilled in the art. For example, dimethyl sulfate can be used in presence of a base such as potassium carbonate or potassium hydroxide in a solvent such as acetone, acetonitrile, 1,4-dioxane, toluene, water or mixtures thereof, at a temperature between room temperature and reflux temperature, preferably reflux temperature, for several hours. Alternatively, methyl iodide can be used as a methylating agent under similar reaction conditions.

For the specific case of compound **VIII** where R1 is bromo and R2 is hydrogen, the key intermediate **VIII** can be prepared as described in U.S. Patent No. 7,226,951 (WO 2005058798) which is hereby incorporated by reference in its entirety.

- 32 -

Hydrolysis of methyl ester **VIII** to give the corresponding acid derivatives **IX** can be carried out in the presence of an aqueous inorganic base such as sodium hydroxide or potassium hydroxide, in an inert solvent such as 1,4-dioxane or tetrahydrofuran, at room temperature for several hours.

5

The cyclization of acid derivatives **IX** to form naphthalenones **X** can be accomplished in two steps. The first step is to convert the acids **IX** to the corresponding acyl chloride derivatives using methods that are well known to someone skilled in the art. The acids can be treated with thionyl chloride or certain phosphorus chloride

10 reagents such as phosphorus trichloride or phosphorus pentachloride at a temperature between room temperature and reflux temperature for several hours. The second step consists of intramolecular Friedel-Crafts acylation of the resulting acyl chloride intermediates in the presence of a strong Lewis acid such as aluminum chloride, under anhydrous conditions. The reactions can be carried out either neat
15 (no solvent), or in a solvent such as tetrahydrofuran, dichloromethane, 1,1,2,2-tetrachloroethane, benzene or 1,2-dichloroethane, at a temperature between room temperature and 80 degrees for several hours. Alternatively, naphthalenones **X** can be obtained by treating compounds **IX** with phosphorus oxychloride at an elevated temperature in a solvent such as 1,2-dichloroethane, tetrahydrofuran, or 1,1,2,2-
20 tetrachloroethane for several hours.

Demethylation of the methoxy compounds **X** to give the phenol derivatives **XI** can be carried out in the presence of aluminum chloride in a solvent such as
25 dichloromethane, acetonitrile, toluene, or *p*-xylene at a temperature between room temperature and reflux temperature for several hours.

Nucleophilic substitution reactions of 8-hydroxy-naphthalenones **X** with *tert*-butyl bromoacetate (**XII**) to give compounds **XIII** can be accomplished using methods that are well known to someone skilled in the art. The reactions are typically carried out
30 in the presence of a carbonate base (e.g. cesium carbonate, potassium carbonate, or the like) or potassium hydroxide in an aprotic solvent such as acetonitrile, *N,N*-dimethylformamide, or dimethyl sulfoxide, at a temperature between 50 and 100 °C for several hours.

Transformation of the naphthalenones **XIII** to the amine derivatives **XV** can be achieved via reductive amination. The reactions can be carried out in stepwise fashion by treating ketones **XIII** with an amine such as ammonium acetate (**XIV**) or ammonia to generate the corresponding imines, which can then be isolated and reduced with a suitable reducing agent (e.g. sodium borohydride). It is also possible to carry out the same reaction sequence in one pot, with the imine formation and reduction occurring concurrently with reducing agents such as sodium cyanoborohydride (NaBH_3CN) or sodium triacetoxyborohydride ($\text{NaBH}(\text{OCOCH}_3)_3$). The reaction is typically performed in a solvent such as methanol or tetrahydrofuran, at a temperature between room temperature and reflux temperature for several hours.

Sulfonylation of the amines **XV** with sulfonyl chlorides **XVI** to give sulfonamides **XVII** can be easily accomplished using methods well known to someone skilled in the art. The reaction is typically carried out in the presence of a base such as diisopropylethylamine, triethylamine, pyridine, or dimethyl-pyridin-4-yl-amine in a suitable inert solvent such as dichloromethane, acetonitrile, 1,4-dioxane, tetrahydrofuran or mixtures thereof, at 0 °C to room temperature for 16 hours.

A large number of sulfonyl chlorides are commercially available. The preparation of non-commercial sulfonyl chlorides is well known to someone skilled in the art, through a variety of methods [for examples, see a) Gilbert, E.E. *Sulfonation and Related Reactions*; Wiley: NY, **1965**; b) Gilbert, E.E. *Synthesis*, **1969**, 1, 6; and c) the procedure described for the preparation of 5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl chloride from 2-chloro-5-(3-isopropyl-phenyl)-pyridine, below].

N-Methylation of compounds **XVII** to produce the derivatives **XIX** can be achieved by treating compounds **XVII** with methyl iodide (**XVIII**) in the presence of a weak base such as potassium carbonate or sodium carbonate, in an inert solvent such as *N,N*-dimethylformamide, acetonitrile, or tetrahydrofuran, at 65 degrees for 5 hours.

- 34 -


Hydrolysis of esters **XVII** or **XIX** gives the corresponding acids **I**. The reaction can be carried out in the presence of an aqueous inorganic base such as lithium hydroxide, sodium hydroxide, or potassium hydroxide, in an inert solvent such as 1,4-dioxane or tetrahydrofuran, at room temperature for several hours.

5

The compounds of interest **IA**, in enantiomerically pure (R)- form, can be obtained through a resolution of the racemic mixtures **I** using chiral chromatography.

10

Alternatively, the (R)- enantiomers **IA** can be generated by following the exact same route as described in Scheme 1, this time employing the enantiomerically pure form of intermediates **XV**, which can be obtained by chiral resolution of racemic **XV**.



15

For the specific case where R_1 is chloro and R_2 is hydrogen, an alternative synthesis may be utilized to access the key intermediate **IXa** (prior to intramolecular Friedel-Crafts cyclization), as outlined in Scheme 2. A Sonogashira coupling between 2-bromo-5-chloroanisole (**XX**), and 3-butyn-1-ol (**XXI**) provides the acetylene intermediate **XXII**, which can be reduced by hydrogenation to furnish the intermediate alcohol **XXIII**. Oxidation of the alcohol **XXIII** to the corresponding carboxylic acid provides the key intermediate **IXa** (wherein R_1 is chloro and R_2 is

20

- 35 -

hydrogen). The intermediate **IXa** can then be converted to compounds of interest **I** wherein R1 is chloro and R2 is hydrogen, as previously described above (in Scheme 1).


- 5 The Sonogashira coupling reaction between 2-bromo-5-chloroanisole (**XX**) and 3-butyn-1-ol (**XXI**) to give the aryl-substituted acetylene **XXII** can be achieved in the presence of a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0) or bis(triphenylphosphine)palladium(II) chloride, and a copper(I) catalyst such as copper(I) iodide. The reaction can be carried out in an inert solvent such as
10 pyrrolidine, at a temperature between room temperature and 80 °C for several hours.

The reduction of the acetylene **XXII** to the corresponding saturated alcohol **XXIII** can be achieved by hydrogenation in the presence of PtO₂ under 50 psi hydrogen atmosphere. The reaction can be carried out in a solvent such as ethyl acetate, at
15 room temperature, for several hours.

The oxidation of alcohol **XXIII** to the corresponding acid **IXa** (wherein R1 is chloro and R2 is hydrogen) can be accomplished using methods well known to someone skilled in the art, by utilizing a variety of oxidizing agents. For example, H₅IO₆/CrO₃
20 can be used as the oxidant. The reaction can be carried out in a suitable solvent mixture, such as acetonitrile and water, at 0 °C.

- 36 -

Scheme 3



- For the specific case where R_1 is bromo and R_2 is hydrogen, an alternative
- 5 synthesis may be utilized to access the key intermediate **XIIIb** (prior to reductive amination), as outlined in Scheme 3. A non-regioselective Pd-catalyzed hydroxylation of the commercially available dibromo derivative **XXIV** provides the phenol derivative **Xlb** in a single step. Two other products are also observed in the reaction, namely the dihydroxyl derivative **XXV**, and the undesired regioisomer **XXVI**.
- 10 The intermediate **Xlb** can then be separated and converted to esters **XIIIb** by treatment with *tert*-butyl bromoacetate (**XII**), as previously described. The compounds of interest **I** or **IA**, wherein R_1 is bromo and R_2 is hydrogen, are then prepared from **XIIIb** in a manner analogous to the one previously described above (in Scheme 1).

15

The intermediate **XIIIb** (prepared via the reaction of **Xlb** with *tert*-butyl bromoacetate (**XII**), as described in Scheme 1) would undergo a Suzuki cross-coupling reaction

- 37 -

with phenylboronic acid to provide the phenyl-substituted derivative **XIIIC**. A similar sequence of reactions as described in Scheme 1 would then provide the compounds of interest **I** or **IA**, wherein R1 is phenyl and R2 is hydrogen.


- 5 The Pd-catalyzed hydroxylation of 5,7-dibromo-3,4-dihydro-2H-naphthalen-1-one (**XXIV**) to produce the phenol derivative **XIb** can be achieved using the methodology described by Buchwald *et. al* (*J. Am. Chem. Soc.* **2006**, 128, 10694). The reaction can be carried out with potassium hydroxide, using catalytic amounts of tris(dibenzylideneacetone)dipalladium (0), and di-*tert*-butyl-(2',4',6'-triisopropyl-
10 biphenyl-2-yl)-phosphane, in a mixture of water and dioxane, at 100 °C.

- The conversion of **XIb** to **XIIb** can be accomplished as described above (Scheme 1). The Pd-catalyzed Suzuki coupling of (3-bromo-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (**XIIb**) with phenylboronic acid is then carried out
15 in the presence of a Pd catalyst, such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium, and a base such as cesium carbonate, in a solvent such as dimethoxyethane, at 95 °C, for several hours.

- The intermediate **XIIc** can then be converted to the compounds of interest **I** or **IA**
20 wherein R1 is phenyl and R2 is hydrogen, as previously described above (in Scheme 1).

- 38 -

Scheme 4



For the specific case where R_1 is methoxy and R_2 is hydrogen, an alternative synthesis may be utilized to access the key intermediate **XIIId** (prior to reductive amination), as outlined in Scheme 4. The dihydroxyl derivative **XXV** (prepared as described in Scheme 3) can be sulfonlated to form the corresponding bis-triflate derivative **XXVII**, which is then converted to the mono-triflate **XXVIII** via selective cleavage of the trifluoromethylsulfonyl group at the 7-position. Conversion of **XXVIII** to the *tert*-butyl ester **XXIX** can be achieved as previously described in Scheme 1, by treatment with *tert*-butyl bromoacetate. The removal of the remaining triflate at the 4-position, followed by methylation of the resulting phenol will produce the key intermediate **XIIId**, which is then subsequently transformed into the compounds of interest **I** or **IA** wherein R_1 is methoxy and R_2 is hydrogen, as previously described in Scheme 1.

The sulfonylation of the dihydroxy compound **XXV** to the bis-triflate derivative **XXVII** can be accomplished according to procedures well known to someone skilled in the art. For example, the reaction can be carried out using a sulfonylating agents such as trifluoromethanesulfonic anhydride, and a base such as diisopropylethylamine, in an inert solvent such as dichloromethane, at a temperature near 0 °C.

The cleavage of the triflate at the 7-position to produce the hydroxyl derivative **XXVIII** can be accomplished using a base such as cesium carbonate, in a solvent such as dimethoxyethane, at 80 °C, or the reflux temperature.

The nucleophilic substitution reaction between **XXVIII** and *tert*-butyl bromoacetate (**XII**) to give the *tert*-butyl ester **XXIX** can be accomplished as described previously. The reaction is typically carried out in the presence of a carbonate base (such as cesium carbonate, potassium carbonate, and the like), in an aprotic solvent such as acetonitrile, *N,N*-dimethylformamide, or dimethyl sulfoxide, at a temperature between room temperature and 100 °C for several hours.

The cleavage of the remaining triflate functionality in **XXIX** to provide the hydroxyl derivative **XXX** may be accomplished by treatment with tetrabutylammonium fluoride, in a solvent such as tetrahydrofuran, at room temperature, for several hours.

The O-methylation of phenol **XXX**, to provide the methoxy derivative **XIIId**, can be carried out using methods that are well known to someone skilled in the art. For example, the reaction can be performed using a methylating agent such as iodomethane or dimethyl sulfate, in the presence of a base such as potassium carbonate, in a solvent such as acetone, acetonitrile, 1,4-dioxane, or *N,N*-dimethylformamide, or mixtures thereof, at a temperature between room temperature and reflux temperature (preferably the reflux temperature), for several hours.

The intermediate **XIIId** can then be converted to the compounds of interest **I** or **IA** wherein R1 is methoxy and R2 is hydrogen, as previously described above (in Scheme 1).

EXAMPLES

Intermediates and final compounds were purified by either flash chromatography
5 and/or preparative HPLC (high performance liquid chromatography). Flash
chromatography was performed using (1) the Biotage SP1™ system and the Quad
12/25 Cartridge module from Biotage AB) or (2) the ISCO CombiFlash®
chromatography instrument (from Teledyne Isco, Inc.); unless otherwise noted. The
silica gel brand and pore size utilized were: (1) KP-SIL™ 60 Å, particle size: 40-60
10 micron (from Biotage AB); (2) Silica Gel CAS registry No: 63231-67-4, particle size:
47-60 micron; or (3) ZCX from Qingdao Haiyang Chemical Co., Ltd, pore size: 200-
300 mesh or 300-400 mesh. Preparative HPLC was performed on a reversed
phase column using an Xbridge™ Prep C₁₈ (5 µm, OBD™ 30 × 100 mm) column
(from Waters Corporation), a SunFire™ Prep C₁₈ (5 µm, OBD™ 30 × 100 mm)
15 column (from Waters Corporation), or a Varian Pursuit® C-18 column 20 X 150 mm
(from Varian, Inc.).

Mass spectrometry (MS) or high resolution mass spectrometry (HRMS) was
performed using a Waters® ZQ™ 4000 (from Waters Corporation), a Waters®
20 Alliance® 2795-ZQ™ 2000 (from Waters Corporation), a Waters® Quattro micro™ API
(from Waters Corporation), or an MDS Sciex™ API-2000™n API (from MDS Inc.).
Mass spectra data generally only indicates the parent ions unless otherwise stated.
MS or HRMS data is provided for a particular intermediate or compound where
indicated.

25
Nuclear magnetic resonance spectroscopy (NMR) was performed using a Varian®
Mercury300 NMR spectrometer (for the HNMR spectrum acquired at 300 MHz) and
a Varian® Inova400 NMR spectrometer (for the HNMR spectrum acquired at 400
MHz) both from Varian Inc. NMR data is provided for a particular intermediate or
30 compound where indicated.

- 41 -

The microwave assisted reactions were carried out in a Biotage Initiator™ Sixty (or its early models) (from Biotage AB) or by a CEM Discover® model (with gas addition accessory) (from CEM Corporation).


- 5 Chiral separation was performed by supercritical fluid chromatography (SFC) using a Multigram® III instrument (from Thar Technologies, Inc.).

All reactions involving air-sensitive reagents were performed under an inert atmosphere. Reagents were used as received from commercial suppliers unless
10 otherwise noted.


PREPARATION OF KEY INTERMEDIATES

Preparation of 4-(4-Chloro-2-methoxy-phenyl)-butyric acid (IXa)-Prepared as described in Scheme 2

15



4-(4-Chloro-2-methoxy-phenyl)-but-3-yn-1-ol




- 20 To a stirred solution of 2-bromo-5-chloroanisole (2 g, 9 mmol), Pd(PPh₃)₄ (520 mg, 0.45 mmol), and CuI (400 mg) in pyrrolidine (20 mL) was added a solution of 3-butyn-1-ol (1.26 g, 18 mmol) in pyrrolidine (30 mL). The reaction mixture was stirred at 80 °C for 6 h, and then concentrated *in vacuo*. The resulting residue was extracted with ethyl acetate and the organic phase was washed with 1N
25 hydrochloric acid, followed by water, then brine, and dried over sodium sulfate.

- 42 -


Upon removal of the solvent under reduced pressure, the crude product was subjected to column chromatography on silica gel (gradient elution with 0%-40% EtOAc/hexane) to give 4-(4-chloro-2-methoxy-phenyl)-but-3-yn-1-ol as a colorless oil (1.46 g, 76.7%). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.29 (d, J = 8.2 Hz, 1 H), 6.88 (dd, J = 8.2, 1.6 Hz, 1 H), 6.85 (d, J = 1.6 Hz, 1 H), 3.87 (s, 3 H), 3.82 (t, J = 6.2 Hz, 2 H), 2.73 (t, J = 6.2 Hz, 2 H), 1.92 (br. s, 1 H).

4-(4-Chloro-2-methoxy-phenyl)-butan-1-ol



- 10 A solution of 4-(4-chloro-2-methoxy-phenyl)-but-3-yn-1-ol (2.2 g, 10.4 mmol) and PtO₂ (200 mg) in ethyl acetate (15 mL) was hydrogenated under 50 psi hydrogen atmosphere until no more hydrogen uptake was observed. The reaction mixture was then filtered and concentrated *in vacuo*. The crude product was subjected to column chromatography on silica gel (gradient elution with 0%-40% EtOAc/hexane) to give
- 15 4-(4-chloro-2-methoxy-phenyl)-butan-1-ol as a colorless oil (2 g, 89.7 %). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.03 (d, J = 8.2 Hz, 1 H), 6.86 (dd, J = 8.2, 1.6 Hz, 1 H), 6.81 (d, J = 1.6 Hz, 1 H), 3.81 (s, 3 H), 3.66 (t, J = 6.0 Hz, 2 H), 2.49 - 2.65 (m, 2 H), 1.50 - 1.72 (m, 5 H).

20 4-(4-Chloro-2-methoxy-phenyl)-butyric acid (IXa)




- A stock solution of H₅IO₆/CrO₃ was prepared by dissolving H₅IO₆ (11.4 g, 50 mmol) and CrO₃ (23 mg, 1.2 mol %) in wet acetonitrile (0.75 v % water) to a volume of 114 mL (complete dissolution typically required 1-2 hours). The H₅IO₆/CrO₃ solution (8.7
- 25 mL) was then added to a solution of 4-(4-chloro-2-methoxy-phenyl)-butan-1-ol (330 mg, 1.54 mmol) in wet acetonitrile (10 mL, 0.75 v % water) dropwise over 30-60 minutes while maintaining the reaction temperature at 0-5 °C. The resulting reaction

- 43 -

mixture was stirred at 0 °C for 0.5 h, when completion of the reaction was confirmed by LC/MS. The reaction was quenched by the addition of an aqueous solution of Na₂HPO₄ (0.60 g in 10 mL H₂O). Ethyl acetate was added and the organic layer was separated and washed with water and then brine. The organic layer was then dried and concentrated to give the crude carboxylic acid, which was purified by flash chromatography on silica gel (gradient elution with 25%-50% EtOAc/hexane) to give 4-(4-chloro-2-methoxy-phenyl)-butyric acid as a white solid (200 mg, 56.9%). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 11.64 (br. s, 1 H), 7.12 (d, J = 8.2 Hz, 1 H), 7.00 (s, 1 H), 6.92 (d, J = 8.2 Hz, 1 H), 3.79 (s, 3 H), 2.51 - 2.57 (m, 2 H), 2.18 (t, J = 7.4 Hz, 2 H), 1.72 (quin, J = 7.4 Hz, 2 H).

**Preparation of 7-Bromo-5-hydroxy-3,4-dihydro-2H-naphthalen-1-one (XIb)-
Prepared as described in Scheme 3**



15


To a mixture of tris(dibenzylideneacetone)dipalladium(0) (210 mg, 0.23 mmol), di-*tert*-butyl-(2',4',6'-triisopropyl-biphenyl-2-yl)-phosphane (403 mg, 0.92 mmol), and potassium hydroxide (1.67 g, 25.32 mmol) were added degassed water (5.25 mL) and a solution of 5,7-dibromo-3,4-dihydro-2H-naphthalen-1-one (3.5 g, 11.51 mmol) in dioxane (7.0 mL) at room temperature under nitrogen. The resulting suspension was heated to 100 °C for 3.5 h. The reaction mixture was then cooled to room temperature, diluted with water (~10 mL) and acidified with 1N HCl solution. The resulting mixture was extracted with ethyl acetate (2 x 100 mL) and the combined organic extracts were washed with brine (200 mL), then dried over anhydrous magnesium sulfate, filtered, and concentrated *in vacuo*. The crude product was purified by chromatography, using an ISCO (120 g) column (gradient elution with 5-30% ethyl acetate in hexanes) to obtain 7-bromo-5-hydroxy-3,4-dihydro-2H-naphthalen-1-one (0.45 g, 16%) as a white solid: ES(-)-HRMS *m/e* calcd. (calculated) for C₁₀H₉O₂Br (M-H)⁻ 238.9713, obsd. (observed) 238.9712. Two other

25

- 44 -

side products were also isolated and characterized as 5-bromo-7-hydroxy-3,4-dihydro-2*H*-naphthalen-1-one (0.3 g, 10%) and 5,7-dihydroxy-3,4-dihydro-2*H*-naphthalen-1-one (0.191 g, 7%).


5 **Preparation of (3-bromo-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (XIIIb)-Prepared as described in Scheme 3**



To a suspension of 7-bromo-5-hydroxy-3,4-dihydro-2*H*-naphthalen-1-one (**XIb**, 158
10 mg, 0.65 mmol) and cesium carbonate (431 mg, 1.31 mmol) in acetonitrile (5 mL)
was added *tert*-butyl bromoacetate (**XII**, 261 mg, 1.31 mmol) at room temperature
under nitrogen. The resulting suspension was stirred for 15 h at room temperature
and then concentrated under vacuum. The residue was diluted with water (10 mL)
and ethyl acetate (20 mL). The aqueous layer was separated and extracted with
15 ethyl acetate (20 mL), and the combined organic extracts were washed with brine
solution (50 mL), then dried over anhydrous magnesium sulfate, filtered, and
concentrated *in vacuo* to give the crude product. Chromatography using an ISCO
(40 g) column, (gradient elution with 5-25% ethyl acetate in hexanes) provided (3-
bromo-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (150
20 mg, 64%) as a white solid: ES(+)-HRMS *m/e* calcd. for C₁₆H₁₉O₄Br (M+Na)⁺
377.0359, obsd. 377.0358.


Preparation of (5-oxo-3-phenyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (XIIIc)-Prepared as described in Scheme 3

- 45 -




To a mixture of (3-bromo-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (**XIIIb**, 78 mg, 0.22 mmol), phenylboronic acid (55.3 mg, 0.44 mmol),
 5 [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (24.1 mg, 0.033 mmol), and cesium carbonate (144.8 mg, 0.44 mmol) was added dimethoxyethane (2 mL) at room temperature under nitrogen. The resulting brown reaction mixture was heated at 95 °C for 15 h. Then, the reaction mixture was cooled to room temperature, and diluted with water (20 mL) and ethyl acetate (20 mL). The aqueous layer was
 10 separated and extracted with ethyl acetate (20 mL), and the combined organic extracts were washed with water (50 mL) and brine solution (50 mL), then dried over anhydrous magnesium sulfate, filtered, and concentrated under vacuum to provide the colored residue. Chromatography using an ISCO (12 g) column, (gradient elution with 2-25% ethyl acetate in hexanes) afforded (5-oxo-3-phenyl-5,6,7,8-
 15 tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (54 mg, 70%) as a white solid: ES(+)-HRMS *m/e* calcd. for C₂₂H₂₄O₄ (M+Na)⁺ 375.1567, obsd. 375.1567.

Preparation of (3-Methoxy-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (XIIIId**)-Prepared as described in Scheme 4**




Trifluoro-methanesulfonic acid 8-oxo-4-trifluoromethanesulfonyloxy-5,6,7,8-tetrahydro-naphthalen-2-yl ester (XXVII)



- 5 To a suspension of 5,7-dihydroxy-3,4-dihydro-2*H*-naphthalen-1-one (**XXV**, prepared as described in Scheme 3 above, 320 mg, 1.79 mmol) in dichloromethane (25 mL) was added diisopropylethylamine (925 mg, 7.16 mmol) at -5 to 0°C. To the resulting solution was then added trifluoromethanesulfonic anhydride (1.06 g, 3.76 mmol) dropwise over 5 minutes at this temperature. The resulting brown solution was
- 10 stirred for 45 minutes at ~0-5°C and then warmed to room temperature and stirred for another 15 minutes. The reaction mixture was then diluted with dichloromethane (50 mL) and water (70 mL). The aqueous layer was separated and extracted with dichloromethane (25 mL). The combined organic extracts were washed with brine (100 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated under
- 15 vacuum. The crude product was dissolved in dichloromethane (10 mL) and diluted with hexanes (50 mL). The brown solution was stored in the refrigerator for 15 h. The resulting solids were collected by filtration and washed with hexanes. After drying in the air, trifluoro-methanesulfonic acid 8-oxo-4-trifluoromethanesulfonyloxy-5,6,7,8-tetrahydro-naphthalen-2-yl ester (0.548 g, 69%) was obtained as a light brown solid:
- 20 EI(+)-HRMS *m/e* calcd. for C₁₂H₈F₆O₇S₂ (M⁺) 442.9616, obsd. 442.9618.


Trifluoro-methanesulfonic acid 4-hydroxy-8-oxo-5,6,7,8-tetrahydro-naphthalen-2-yl ester (XXVIII)

- 47 -



To a mixture of trifluoro-methanesulfonic acid 8-oxo-4-trifluoromethanesulfonyloxy-5,6,7,8-tetrahydro-naphthalen-2-yl ester (**XXVII**, 542 mg, 1.23 mmol) and cesium carbonate (599 mg, 1.84 mmol) was added dimethoxyethane (11 mL) at room temperature. The resulting brown suspension was then heated at 80°C and stirred for 3 h. The reaction mixture was cooled to room temperature and diluted with saturated ammonium chloride solution (50 mL), and then extracted with ethyl acetate (2 x 50 mL). The combined organic extracts were washed with brine (100 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated under vacuum. The crude product was dissolved in dichloromethane (15 mL) and diluted with hexanes (100 mL), then heated to remove the dichloromethane. The cloudy solution was then stored in the refrigerator for 15 h. The resulting solids were collected by filtration and washed with hexanes. After drying in the air, trifluoro-methanesulfonic acid 4-hydroxy-8-oxo-5,6,7,8-tetrahydro-naphthalen-2-yl ester (182 mg, 51%) was obtained as a dark brown solid: ES(-)-HRMS m/e calcd. for C₁₁H₉F₃O₅S (M-H)⁻ 309.0050, obsd. 309.0049.


(5-Oxo-3-trifluoromethanesulfonyloxy-5,6,7,8-tetrahydro-naphthalen-1-yl oxy)-acetic acid *tert*-butyl ester (XXIX**)**



- 48 -

To a suspension of trifluoro-methanesulfonic acid 4-hydroxy-8-oxo-5,6,7,8-tetrahydro-naphthalen-2-yl ester (**XXVIII**, 177 mg, 0.57 mmol) and cesium carbonate (372 mg, 1.14 mmol) in acetonitrile (5 mL) was added *tert*-butyl bromoacetate (167 mg, 0.86 mmol) at room temperature under nitrogen. The resulting suspension was stirred for 15 h at room temperature, and then concentrated under vacuum. The residue was diluted with water (10 mL) and ethyl acetate (20 mL). The aqueous layer was separated and extracted with ethyl acetate (20 mL). The combined organic extracts were washed with brine (50 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated *in vacuo*. The crude product was purified by ISCO (40 g) column chromatography, (gradient elution with 5-25% ethyl acetate in hexanes) to provide (5-oxo-3-trifluoromethanesulfonyloxy-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (143 mg, 59%) as a white solid: ES(+)-HRMS *m/e* calcd. for C₁₇H₁₉F₃O₇S (M+H)⁺ 425.0877, obsd. 425.0876.

15 **(3-Hydroxy-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (**XXX**)**




To a solution of (5-oxo-3-trifluoromethanesulfonyloxy-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (**XXIX**, 140 mg, 0.33 mmol) in tetrahydrofuran (5 mL) was added a 1.0 M solution of tetrabutylammonium fluoride (660 μ L, 0.66 mmol) in tetrahydrofuran at room temperature under nitrogen. The resulting light yellow solution was stirred for 5 h at room temperature, and then diluted with water (10 mL) and 1.0N hydrochloric acid (5 mL). The mixture was extracted with ethyl acetate (2 x 35 mL). The combined organic extracts were washed with brine (50 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated *in vacuo*. The crude product was purified by ISCO (40 g) column chromatography (gradient elution with 5-

- 49 -

50% ethyl acetate in hexanes) to provide (3-hydroxy-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (82 mg, 85%) as a yellow solid: ES(+)-HRMS *m/e* calcd. for C₁₆H₂₀O₅ (M+H)⁺ 293.1384, obsd. 293.1383.

5 **(3-Methoxy-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (XIIId)**




To a suspension of (3-hydroxy-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (**XXX**, 78 mg, 0.27 mmol) and potassium carbonate (111 mg, 0.8 mmol) in acetone (4 mL) and DMF (2 mL) was added iodomethane (114 mg, 0.8 mmol) at room temperature under nitrogen. The resulting red-brown suspension was heated at reflux for 2 days. The reaction mixture was then cooled to room temperature and diluted with water (10 mL) and brine (10 mL). The resulting mixture was extracted with ethyl acetate (2 x 35 mL). The combined organic extracts were washed with brine (50 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated *in vacuo*. The crude product was purified by using ISCO (40 g) column chromatography (gradient elution with 5-40% ethyl acetate in hexanes) to obtain (3-methoxy-5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (76 mg, 94%) as a light yellow solid: ES(+)-HRMS *m/e* calcd. for C₁₇H₂₂O₅ (M+H)⁺ 307.1540, obsd. 307.1540.


Preparation of sulfonyl chlorides:

Preparation of 5-(3-Isopropyl-phenyl)-pyridine-2-sulfonyl chloride

- 50 -




3-Isopropyl-phenylboronic acid



- 5 *n*-BuLi (7.7 mL, 14.0 mmol) was added dropwise to a stirred solution of 1-bromo-3-isopropylbenzene (2.4 g, 12.1 mmol) in dry THF (40 mL) over 30 min at -70°C under nitrogen. The reaction mixture was degassed for 15 min, and triisopropylborate (2.63 g, 14.0 mmol) was added at the same temperature. The reaction mixture was gradually warmed to 0°C during 90 min and stirred for an
- 10 additional 30 min at 0°C . The reaction mixture was then cooled to -20°C , and an aqueous solution of 2N HCl (20 mL) was added slowly. THF was distilled off and the reaction mixture was extracted with ethyl acetate (2 x 20 mL). The combined organic layers were dried over Na_2SO_4 , and concentrated *in vacuo* to obtain the crude product (1.85 g, 93.5%), which was used for the next step without further purification.

15

2-Chloro-5-(3-isopropylphenyl)-pyridine




- To a stirred solution of 5-bromo-2-chloropyridine (1.27 g, 6.63 mmol) in toluene (30 mL) was added an ethanolic solution (10 mL) of 3-isopropylphenylboronic acid (1.8 g, 11.28 mmol) followed by water (10 mL) and potassium carbonate (2.76 g, 20.0 mmol). The resulting mixture was degassed for 15 min (by purging with argon). At
- 20 this time, $\text{Pd}(\text{PPh}_3)_4$ (0.613 g, 0.53 mmol) was added under argon and the reaction

- 51 -


mixture was heated at 80 °C for 4 h. The reaction mixture was then cooled to room temperature and extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with water (2 x 10 mL), dried over Na₂SO₄ and concentrated *in vacuo* to obtain the crude product, which was purified over silica gel (Biotage column chromatography, 2% ethyl acetate in hexane) to afford 0.900 g (59%) of 2-chloro-5-(3-isopropyl-phenyl)-pyridine.

2-Benzylsulfanyl-5-(3-isopropyl-phenyl)-pyridine



A solution of phenyl-methanethiol (0.8 mL, 5.84 mmol) in DMF (5 mL) was treated with NaH (0.140 g, 5.84 mmol) at room temperature under argon. After 15 min of stirring, the reaction mixture was treated with a solution of 2-chloro-5-(3-isopropyl-phenyl)-pyridine (0.900 g, 3.90 mmol) in DMF (15 mL). After 2 h of additional stirring at room temperature (reaction was monitored by TLC), the reaction mixture was quenched with ice cold water (20 mL), and extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with water (2 x 10 mL), dried over Na₂SO₄, and concentrated *in vacuo* to obtain a crude product, which was purified over silica gel (Biotage column chromatography, gradient elution using 0-1% ethyl acetate in hexane) to furnish 2-benzylsulfanyl-5-(3-isopropyl-phenyl)-pyridine (0.680 g, 54.71%).


5-(3-Isopropyl-phenyl)-pyridine-2-sulfonyl chloride



- 52 -


Chlorine gas was bubbled through a stirred solution of 2-benzylsulfanyl-5-(3-isopropyl-phenyl)-pyridine (0.680 g, 2.13 mmol) in carbon tetrachloride-water mixture (5:1; 30 mL) at 0 °C for 15 min. After 15 min of additional stirring, argon was bubbled through the reaction mixture, which was then quenched with ice (~ 10 mL) and
5 extracted with dichloromethane (2 x 20 mL). The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuo* to obtain 5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl chloride (0.700 g, 71.5%), which was used without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.92 (s, 1 H), 8.72 (d, J = 8.0 Hz, 1 H), 8.13 (d, J = 8.0 Hz, 1 H), 7.74 (s, 1 H), 7.66 (d, J = 7.3 Hz, 1 H), 7.48 (t, J = 7.3 Hz, 1 H), 7.40
10 (d, J = 7.3 Hz, 1 H), 2.92 - 3.06 (m, 1 H), 1.27 (d, J = 6.8 Hz, 6 H).

Preparation of 3-fluoro-5-trifluoromethyl-benzenesulfonyl chloride



A mixture of 3-fluoro-5-trifluoromethyl-phenylamine (9.7 g, 54 mmol) in trifluoroacetic acid (100 mL) was cooled at 0 °C. To the mixture was slowly added concentrated
15 hydrochloric acid (10 mL), followed by a solution of sodium nitrite (4.7 g, 68 mmol) in water (5 mL) dropwise over 20 minutes. The mixture was stirred for another 10 minutes at 0 °C, and then poured into a stirred mixture of acetic acid (120 mL), sulfurous acid (0.94 N aqueous sulfur dioxide solution, 120 mL), copper(II) chloride
20 (9.2 g, 93 mmol) and copper(I) chloride (100 mg, 0.74 mmol) at 0 °C. The resulting reaction mixture was allowed to warm to room temperature and stirred for 15 hours. Water (200 mL) was added, and the resulting mixture was extracted with ethyl acetate (100 mL x 3). The combined organic layers were dried over sodium sulfate,
25 filtered through a glass funnel and concentrated *in vacuo*. The residue was purified by column chromatography (20% ethyl acetate in petroleum ether) to afford 3-fluoro-5-trifluoromethyl-benzenesulfonyl chloride (3.7 g, 26 %) as a white solid (reference: Cherney, R.J. *et al.*, *J. Med. Chem.* 46 (2003) 1811). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.15 (s, 1 H); 7.97-7.99 (d, J = 4.0 Hz, 1 H); 7.74-7.76 (d, J = 4.0 Hz, 1 H).

Preparation of 3-methoxy-5-trifluoromethyl-benzenesulfonyl chloride




3-methoxy-5-trifluoromethyl-phenylamine (10 g, 54 mmol) was added to trifluoroacetic acid (100 mL) in a 250 mL flask, and the mixture was cooled to 0°C. Concentrated hydrochloric acid (10 mL) was then added slowly to the reaction mixture, followed by the dropwise addition of a solution of sodium nitrite (4.7 g, 68 mmol) in water (5 mL) over 20 min. The mixture was stirred for another 10 minutes at 0 °C, and then poured into a stirred mixture of acetic acid (120 mL), sulfurous acid (0.94 N aqueous sulfur dioxide solution, 120 mL, 113 mmol), copper(II) chloride (9.2 g, 68 mmol) and copper(I) chloride (100 mg, 1 mmol) at 0 °C. The reaction mixture was allowed to warm to room temperature and stirred for 15 hours, and then treated with water (200 mL). The aqueous layer was extracted with ethyl acetate (100 mL × 3). The combined organic layers were dried over sodium sulfate, filtered through a glass funnel and concentrated *in vacuo*. The residue was purified by column chromatography (20% ethyl acetate in petroleum ether) to afford 3-methoxy-5-trifluoromethyl-benzenesulfonyl chloride (3.9 g, 27%) as a white solid [reference: Cherney, R.J. *et al.*, *J. Med. Chem.* 46 (2003) 1811]. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.89 (s, 1 H); 7.70 (s, 1 H); 7.50 (s, 1 H); 4.00 (s, 3 H).


PREPARATION OF COMPOUNDS OF INTEREST

EXAMPLE 1-1 (Prepared as described in Scheme 1)


[(R)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid



- 54 -

4-(4-Chloro-3-fluoro-2-hydroxy-phenyl)-4-oxo-butyrac acid

To a stirred and cooled solution of 2-fluoro-3-chloro-phenol (21.9 g, 0.15 mol) and succinic anhydride (15.3 g, 0.15 mol) in 1,1,2,2-tetrachloroethane (135 mL) was added anhydrous aluminum chloride (38.7 g, 0.29 mol) portion-wise. The reaction mixture was stirred at room temperature for 3 hours and then heated at 130 °C for 2 hours. After being cooled to room temperature, to the reaction mixture were added concentrated hydrochloric acid and ice alternatively in small portions. The resulting mixture was extracted with ethyl acetate (75 mL x 3). The combined organic layers were evaporated. The solid residue was basified by slow addition of an aqueous solution of 2N sodium hydroxide solution under cooling conditions, and then extracted with ethyl acetate (30 mL x 3). The aqueous layer was acidified by dropwise addition of concentrated hydrochloric acid to a pH of about 2, and then extracted with ethyl acetate (50 mL x 3). The combined organic layers were dried over sodium sulfate, filtered, and concentrated *in vacuo*. Column chromatography (silica gel, 100-120 mesh, 15% ethyl acetate in hexane) gave 4-(4-chloro-3-fluoro-2-hydroxy-phenyl)-4-oxo-butyrac acid (7.0 g, 19%). ¹H NMR (400 MHz, DMSO-d₆) δ ppm 2.59 (t, J= 6.26 Hz, 2H), 3.31 (t, J=6.28 Hz, 2H), 7.15 (d, J=6.56 Hz, 1H), 7.74 (d, J=6.56 Hz, 1H), 11.90-12.31 (m, 2H).


4-(4-Chloro-3-fluoro-2-hydroxy-phenyl)-butyrac acid

A mixture of zinc pieces (11.4 g), mercuric chloride (3.0 g), concentrated hydrochloric acid (4.6 mL) and water (140 mL) was stirred at room temperature for 5 minutes. The solvent was then decanted off. To the solid, were added concentrated hydrochloric acid (41 mL), water (18.6 mL), toluene (23 mL) and 4-(4-chloro-3-fluoro-2-hydroxy-phenyl)-4-oxo-butyrac acid (8.6 g, 35 mmol). The resulting mixture was

- 55 -


heated at reflux for 2 days, with the addition of concentrated hydrochloric acid in 12 hour intervals. The reaction mixture was cooled and extracted with ether (50 mL x 3). The collected organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo* to give pure 4-(4-chloro-3-fluoro-2-hydroxy-phenyl)-butyric acid (6.0 g, 74%), which was used in the next step without further purification. ¹H NMR (400 MHz, DMSO-d₆) δ ppm 1.74 (dt, 2H), 2.20 (t, J=7.40 Hz, 2H), 2.57 (t, J=7.56 Hz, 2H), 6.87-6.92 (m, 2H), 9.90 (br s, 1H), 12.03 (br s, 1H).

4-(4-Chloro-3-fluoro-2-methoxy-phenyl)-butyric acid methyl Ester



To a stirred solution of 4-(4-chloro-3-fluoro-2-hydroxy-phenyl)-butyric acid (6.0 g, 0.026 mol) in acetone (20 mL) was added anhydrous potassium carbonate (9.7 g, 0.07 mol) at room temperature under nitrogen. After 10 minutes, dimethyl sulfate (5.49 mL, 0.058 mol) was added and the resulting mixture was stirred for an additional 3-4 hours. The solvent was removed *in vacuo* and the residue was extracted with dichloromethane (25 mL x 3). The combined organic layers were washed with water (15 mL x 2), dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo* to give 4-(4-chloro-3-fluoro-2-methoxy-phenyl)-butyric acid methyl ester (5.6 g, 83%), which was used in the next step without further purification. MS calcd. for C₁₂H₁₄ClFO₃ 260, obsd. 261 [(M+H)⁺].


4-(4-Chloro-3-fluoro-2-methoxy-phenyl)-butyric acid




To a stirred solution of 4-(4-chloro-3-fluoro-2-methoxy-phenyl)-butyric acid methyl ester (5.6 g, 0.022 mol) in tetrahydrofuran (25 mL) was added a solution of lithium hydroxide (2.7 g, 0.067 mol) in water-methanol (2:1; 20 mL) at room temperature. The reaction mixture was stirred for 4 hours at room temperature. The solvents were

- 56 -

removed under reduced pressure. The residue was diluted with water (25 mL) and then acidified by dropwise addition of aqueous 1N hydrochloric acid under cooling conditions. The resulting solution was extracted with ethyl acetate (30 mL x 3). The combined organic layers were washed with water (30 mL), dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo* to give 4-(3,4-dichloro-2-methoxy-phenyl)-butyric acid (4.7 g, 87%), which was used in the next step without further purification. MS calcd. for C₁₁H₁₂ClFO₃ 246, obsd. 247 [(M+H)⁺].

7-Chloro-6-fluoro-5-methoxy-3,4-dihydro-2H-naphthalen-1-one

To a stirred solution of 4-(3,4-dichloro-2-methoxy-phenyl)-butyric acid (1.03 g, 8.10 mmol) in benzene (100 mL) were added phosphorus pentachloride (2.70 g, 12.96 mmol) and aluminum chloride (3.24 g, 24.30 mmol) at room temperature. After the reaction mixture was stirred for 4 hours, it was poured into ice water with continuous stirring, and then extracted with ethyl acetate (50 mL x 3). The collected organic layers were subsequently washed with saturated sodium bicarbonate solution (20 mL) and water (20 mL), dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo*. Column chromatography (silica gel, 100-200 mesh, 10% ethyl acetate in hexane) gave 7-chloro-6-fluoro-5-methoxy-3,4-dihydro-2H-naphthalen-1-one (1.04 g, 58%). ¹H-NMR (400 MHz, CDCl₃) δ ppm 2.10 (dt, 2H), 2.61 (t, J=6.58 Hz, 2H), 2.90 (t, J=6.10 Hz, 2H), 3.93 (s, 3H), 7.83 (s, 1H).


7-Chloro-6-fluoro-5-hydroxy-3,4-dihydro-2H-naphthalen-1-one

To a stirred solution of 7-chloro-6-fluoro-5-methoxy-3,4-dihydro-2H-naphthalen-1-one (1.04 g, 4.7 mmol) in xylene (35 mL) was added anhydrous aluminum chloride (1.93 g, 14.5 mmol) at room temperature. The reaction mixture was then heated at

- 57 -

reflux for 1 hour. After being cooled to room temperature, the reaction mixture was slowly poured into ice water with continuous stirring. The resulting solution was extracted with ethyl acetate (50 mL x 3). The collected organic layers were then washed with saturated sodium bicarbonate solution and water, dried with anhydrous sodium sulfate, filtered, and concentrated *in vacuo* to give the crude residue, which was washed with ether to afford 7-chloro-6-fluoro-5-hydroxy-3,4-dihydro-2*H*-naphthalen-1-one (0.90 g, 89%), which was used in the next step without further purification. MS calcd. for C₁₀H₈ClFO₂ 214, obsd. 215 [(M+H)⁺].


10 **(3-Chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester**



To a stirred solution of 7-chloro-6-fluoro-5-hydroxy-3,4-dihydro-2*H*-naphthalen-1-one (0.90 g, 4.18 mmol) in dry *N,N*-dimethylformamide (10 mL) was added potassium carbonate (1.16 g, 8.36 mmol) and *tert*-butyl bromoacetate (1.22 g, 6.27 mmol) at room temperature and the resulting mixture was stirred for 2 hours, then extracted with ethyl acetate (25 mL x 3), washed with water (25 mL x 3), dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo* to afford (3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (1.32 g, 96%), which was used in the next step without further purification. MS calcd. for C₁₆H₁₈ClFO₄ 328, obsd. 329 [(M+H)⁺].


(5-Amino-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester

- 58 -



To a stirred and ice cold solution of (3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (1.32 g, 4.01 mmol) and ammonium acetate (4.7 g, 61.0 mmol) in methanol (21 mL) was added sodium cyanoborohydride (504 mg, 8.02 mmol) under nitrogen. After 10 minutes of stirring at room temperature, the reaction mixture was heated at reflux for 4 hours. The solvent was removed under reduced pressure and to the residue was added saturated sodium bicarbonate solution (10 mL). The resulting solution was extracted with dichloromethane (25 mL x 3). The organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo* to afford (5-amino-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (1.26 g, 98%), which is used in the next step without further purification.

[5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid *tert*-butyl ester




To a solution of (5-amino-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid *tert*-butyl ester (125.0 mg, 0.38 mmol) and 3,5-di-(trifluoromethyl)benzene-1-sulfonyl chloride (0.117 g, 0.38 mmol) in dry tetrahydrofuran (5 mL) was added diisopropylethylamine (0.2 mL, 0.94 mmol) at 0 °C under nitrogen. After 2 hours of stirring at room temperature, the reaction mixture was concentrated *in vacuo*. The residue was partitioned between ethyl acetate and water. The collected organic layers were dried over anhydrous sodium sulfate,

- 59 -

filtered, and concentrated under reduced pressure. Column chromatography (silica gel, 100-200 mesh, 5% ethyl acetate in hexane) gave [5-(3,5-bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid *tert*-butyl ester (163.2 mg, 71%). MS cald. for C₂₄H₂₃ClF₇NO₅S 605, obsd. 604 [(M+H)⁺].

[(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)- 3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid



To a stirred solution of [5-(3,5-bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid *tert*-butyl ester (163.4 mg, 0.27 mmol) in tetrahydrofuran (4 mL) were added an aqueous (2 mL) solution of lithium hydroxide (0.036 g, 0.87 mmol) and methanol (1 mL) at room temperature and the resulting reaction mixture was stirred for 2 hours. The solvent was removed under reduced pressure. The residue was diluted with water (15 mL) and acidified with dilute hydrochloric acid, which resulted in the formation of a precipitate. The resulting mixture was extracted with ethyl acetate (15 mL x 3). The combined organic layers was washed with water (15 mL x 3), dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo*. Column chromatography (silica gel, 100-200 mesh, 50% ethyl acetate in hexane) gave [5-(3,5-bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid (130.4 mg, 88%). MS cald. for C₂₀H₁₅ClF₇NO₅S 549, obsd. 548 (ESI⁺) [(M-H)⁺].

Chiral separation by super fluid chromatography (SFC) (Thar Technologies, Inc.'s Multigram[®] II instrument AD column, 100 bar back pressure, 35 °C oven, 10% methanol in carbon dioxide at 70 ml/min) gave (*R*)-[5-(3,5-bis-trifluoromethyl-

- 60 -

benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid (58 mg). ^1H NMR (400 MHz, DMSO- d_6) δ ppm 13.06 (br. s, 1 H), 8.60 (d, $J = 7.9$ Hz, 1 H), 8.54 (s, 1 H), 8.45 (s, 2 H), 6.63 (d, $J = 6.8$ Hz, 1 H), 4.67 (s, 2 H), 4.46 - 4.57 (m, 1 H), 2.57 - 2.75 (m, 2 H), 1.70 - 1.83 (m, 1 H), 1.60 (m, 3 H); MS

5 cald. for $\text{C}_{20}\text{H}_{15}\text{ClF}_7\text{NO}_5\text{S}$ 549, obsd. 548 (ESI^+) $[(\text{M}-\text{H})^+]$.

EXAMPLES 1-2 to 1-52

The following examples 1-2 to 1-52 were prepared in an analogous manner to example 1-1 starting with commercially available substituted phenols and aryl sulfonyl chlorides. Where appropriate, the examples were prepared from the






10 corresponding key intermediates **IXa**, **XIb**, or **XIIIc**, as described above. In cases where the example is depicted as a single enantiomer, a chiral separation of enantiomers was performed on the racemic final product, as described above.

Example No.	Systematic Name	^1H NMR (400 MHz, DMSO- d_6) δ ppm	MS (M-H) ⁻	Structure
1-2	[5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-isopropyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	(CDCl_3) 8.39 (s, 2 H), 8.12 (s, 1 H), 7.06 (d, $J = 8.1$ Hz, 1 H), 6.80 (d, $J = 8.1$ Hz, 1 H), 4.88 (d, $J = 8.1$ Hz, 1 H), 4.52 - 4.62 (m, 1 H), 4.41 (s, 2 H), 3.13 - 3.30 (m, 1 H), 2.80 (ddd, $J = 17.2, 5.3, 5.1$ Hz, 1 H), 2.56 - 2.67 (m, 1 H), 1.73 - 1.91 (m, 5 H), 1.20 (d, $J = 7.1$ Hz, 6 H)	538	
1-3	[5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	12.90 (br. s, 1 H), 8.51 (br. s, 1 H), 8.50 (d, $J = 8.3$ Hz, 1 H), 8.44 (s, 2 H), 6.91 (d, $J = 8.0$ Hz, 1 H), 6.65 (d, $J = 8.0$ Hz, 1 H), 4.50 (br. s, 1 H), 4.29 (s, 2 H), 2.54 - 2.74 (m, 2 H), 2.17 (s, 3 H), 1.65 - 1.86 (m, 1 H), 1.56 (br. s, 3 H)	510	

1-4	[5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-ethyl-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	12.88 (br. s, 1 H), 8.48 - 8.52 (m, 2 H), 8.45 (s, 2 H), 6.94 (d, J = 7.8 Hz, 1 H), 6.69 (d, J = 7.8 Hz, 1 H), 4.47 - 4.54 (m, 1 H), 4.28 (s, 2 H), 2.52 - 2.72 (m, 4 H), 1.66 - 1.82 (m, 1 H), 1.50 - 1.67 (m, 3 H), 1.10 (t, J=7.3 Hz, 3 H)	524	
1-5	[5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-methyl-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	8.50 (s, 1 H), 8.42 (s, 2 H), 6.56 (s, 1 H), 4.42 - 4.53 (m, 1 H), 4.30 (s, 2 H), 2.54 - 2.70 (m, 2 H), 2.17 (s, 3 H), 1.68 - 1.80 (m, 1 H), 1.56 - 1.68 (m, 3 H)	544	
1-6	[5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.07 (br. s, 1 H), 8.63 (d, J = 8.1 Hz, 1 H), 8.54 (s, 1 H), 8.46 (s, 2 H), 6.78 (d, J = 1.6 Hz, 1 H), 6.37 (d, J = 1.6 Hz, 1 H), 4.71 (s, 2 H), 4.44 - 4.63 (m, 1 H), 2.54 - 2.70 (m, 2 H), 1.69 - 1.85 (m, 1 H), 1.60 (m, 3 H)	530	
1-7	[5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-chloro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.04 (br. s, 1 H), 8.62 (d, J = 8.3 Hz, 1 H), 8.53 (s, 1 H), 8.46 (s, 2 H), 6.79 (s, 1 H), 6.38 (s, 1 H), 4.73 (s, 2 H), 4.45 - 4.60 (m, 1 H), 2.42 - 2.49 (m, 2 H), 1.70 - 1.83 (m, 1 H), 1.53 - 1.70 (m, 3 H)	530	
1-8	[(R)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.08 (br. s, 1 H), 8.62 (d, J = 8.1 Hz, 1 H), 8.54 (s, 1 H), 8.46 (s, 2 H), 6.79 (d, J = 1.4 Hz, 1 H), 6.38 (d, J = 1.4 Hz, 1 H), 4.71 (s, 2 H), 4.47 - 4.58 (m, 1 H), 2.40 - 2.61 (m, 2 H), 1.70 - 1.84 (m, 1 H), 1.55 - 1.71 (m, 3 H)	530	

1-9	[5-(3,5-Bis-trifluoromethyl-benzenesulfonyl-amino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	13.07 (br. s, 1 H), 8.61 (d, J = 8.1 Hz, 1 H), 8.54 (s, 1 H), 8.45 (s, 2 H), 6.63 (d, J = 6.8 Hz, 1 H), 4.68 (d, J = 2.0 Hz, 2 H), 4.49 - 4.57 (m, 1 H), 2.54 - 2.73 (m, 2 H), 1.70 - 1.84 (m, 1 H), 1.52 - 1.69 (m, 3 H)	548	
1-10	[3-Chloro-5-(2,5-dichloro-benzenesulfonyl-amino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	13.06 (br. s, 1 H), 8.71 (d, J = 8.6 Hz, 1 H), 8.00 (d, J = 2.2 Hz, 1 H), 7.68 - 7.85 (m, 2 H), 6.92 (d, J = 6.6 Hz, 1 H), 4.68 (s, 2 H), 4.29 - 4.47 (m, 1 H), 2.58 - 2.69 (m, 2 H), 1.83 (br. s, 1 H), 1.62 (br. s, 3 H)	480	
1-11	[3-Chloro-5-(2,4-dichloro-benzenesulfonyl-amino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	13.05 (br. s, 1 H), 8.61 (d, J = 9.3 Hz, 1 H), 8.06 (d, J = 8.8 Hz, 1 H), 7.93 (d, J = 2.0 Hz, 1 H), 7.67 (dd, J = 8.8, 2.0 Hz, 1 H), 6.80 (s, 1 H), 6.69 (s, 1 H), 4.73 (s, 2 H), 4.26 - 4.38 (m, 1 H), 2.44-2.50 (m, 2 H), 1.77 - 1.89 (m, 1 H), 1.53 - 1.67 (m, 3 H)	462	
1-12	[5-(3,5-Bis-trifluoromethyl-benzenesulfonyl-amino)-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	13.08 (s, 1 H), 8.63 (d, J = 8.1 Hz, 1 H), 8.55 (s, 1 H), 8.46 (s, 2 H), 6.81 (s, 1 H), 4.51 - 4.61 (m, 1 H), 4.48 (s, 2 H), 2.54 - 2.81 (m, 2 H), 1.71 - 1.85 (m, 1 H), 1.54 - 1.71 (m, 3 H)		
1-13	[3-Chloro-5-(2,5-dichloro-benzenesulfonyl-amino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	12.88 (br. s, 1 H), 8.55 (s, 1 H), 8.52 (s, 2 H), 7.10 (dd, J = 11.7, 8.7 Hz, 1 H), 6.84 (dd, J = 8.7, 5.1 Hz, 1 H), 5.22 - 5.30 (m, 1 H), 4.64 (s, 2 H), 2.78 - 2.90 (m, 1 H), 2.57 (s, 3 H), 2.43 - 2.49 (m, 1 H), 1.80 (br. s, 1 H), 1.50 - 1.76 (m, 2 H), 1.46 (br. s, 1 H)	462	

1-14	[3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.03 (br. s, 1 H), 8.62 (d, J = 9.0 Hz, 1 H), 8.05 (d, J = 8.4 Hz, 1 H), 7.93 (d, J = 2.2 Hz, 1 H), 7.67 (dd, J = 8.4, 2.2 Hz, 1 H), 6.91 (d, J = 6.6 Hz, 1 H), 4.69 (s, 2 H), 4.24 - 4.38 (m, 1 H), 2.63 (br. s, 2 H), 1.83 (br. s, 1 H), 1.52 - 1.75 (m, 3 H)	480	
1-15	[5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.09 (br. s, 1 H), 8.53 (d, J = 8.3 Hz, 1 H), 8.38 (s, 1 H), 8.31 (s, 1 H), 8.14 (s, 1 H), 6.67 (d, J = 6.8 Hz, 1 H), 4.68 (s, 2 H), 4.44 - 4.52 (m, 1 H), 2.57 - 2.72 (m, 2 H), 1.70 - 1.80 (m, 1 H), 1.59 (br. s, 3 H)	560/562	
1-16	[3-Chloro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.05 (br. s, 1 H), 8.54 (d, J = 8.1 Hz, 1 H), 8.13 (d, J = 9.0 Hz, 1 H), 8.05 (d, J = 7.3 Hz, 1 H), 8.02 (s, 1 H), 6.80 (d, J = 1.4 Hz, 1 H), 6.46 (d, J = 1.4 Hz, 1 H), 4.74 (s, 2 H), 4.44 - 4.52 (m, 1 H), 2.54 - 2.60 (m, 1 H), 2.42 - 2.48 (m, 1 H), 1.70 - 1.84 (m, 1 H), 1.47 - 1.70 (m, 3 H)	480/482	
1-17	[5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.05 (br. s, 1 H), 8.54 (d, J = 8.6 Hz, 1 H), 8.38 (s, 1 H), 8.32 (s, 1 H), 8.15 (s, 1 H), 6.80 (d, J = 1.6 Hz, 1 H), 6.43 (d, J = 1.6 Hz, 1 H), 4.74 (s, 2 H), 4.41 - 4.53 (m, 1 H), 2.53 - 2.60 (m, 1 H), 2.40 - 2.49 (m, 1 H), 1.75 (br. s, 1 H), 1.54 - 1.71 (m, 3 H)	540/542	






1-18	[(<i>R</i>)-3-Chloro-5-(2,4-dichlorobenzenesulfonylamino)-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	8.61 (br. s, 1 H), 8.06 (d, <i>J</i> = 8.5 Hz, 1 H), 7.93 (d, <i>J</i> = 2.0 Hz, 1 H), 7.67 (dd, <i>J</i> = 8.5, 2.0 Hz, 1 H), 6.75 (s, 1 H), 6.67 (s, 1 H), 4.64 (s, 2 H), 4.31 (br. s, 1 H), 2.45 - 2.50 (m, 2 H), 1.75 - 1.93 (m, 1 H), 1.51 - 1.66 (m, 3 H)	462	
1-19	[5-(3,5-Bis-trifluoromethylbenzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	(300 MHz, DMSO- <i>d</i> ₆) 13.07 (br. s, 1 H), 8.63 (d, <i>J</i> = 8.2 Hz, 1 H), 8.54 (s, 1 H), 8.46 (s, 2 H), 6.89 (s, 1 H), 6.49 (s, 1 H), 4.73 (s, 2 H), 4.46-4.60 (m, 1 H), 2.33-2.47 (m, 2 H), 1.52-1.82 (m, 4 H)	597 [#]	
1-20	[(<i>R</i>)-5-(3,5-Bis-trifluoromethylbenzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid			
1-21	[3-Chloro-2-fluoro-5-(3-fluoro-5-trifluoromethylbenzenesulfonylamino)-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.36 (br. s, 1 H), 8.52 (d, <i>J</i> = 8.3 Hz, 1 H), 8.13 (d, <i>J</i> = 8.8 Hz, 1 H), 8.04 (d, <i>J</i> = 7.8 Hz, 1 H), 8.01 (s, 1 H), 6.70 (d, <i>J</i> = 6.8 Hz, 1 H), 4.68 (s, 2 H), 4.41 - 4.53 (m, 1 H), 2.57 - 2.73 (m, 2 H), 1.69 - 1.82 (m, 1 H), 1.54 - 1.69 (m, 3 H)	498/500	
1-22	[5-(3,5-Bis-trifluoromethylbenzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	12.92 (br. s, 1 H), 8.53 (d, <i>J</i> = 7.8 Hz, 1 H), 8.51 (s, 1 H), 8.45 (s, 2 H), 7.00 (dd, <i>J</i> = 11.7, 8.7 Hz, 1 H), 6.76 (dd, <i>J</i> = 8.7, 5.1 Hz, 1 H), 4.62 (s, 2 H), 4.47 - 4.57 (m, 1 H), 2.56 - 2.81 (m, 2 H), 1.66 - 1.83 (m, 1 H), 1.48 - 1.66 (m, 3 H)		

1-23*	[(<i>R</i>)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-methoxy-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	14.31 (br. s, 1 H), 8.50-8.56 (m, 2 H), 8.48 (s, 2 H), 6.25 (d, J = 1.8 Hz, 1 H), 5.92 (d, J = 1.8 Hz, 1 H), 4.55 (br. s, 2 H), 4.44-4.52 (m, 1 H), 3.45 (s, 3 H), 2.80 (br. s, 1 H), 2.36-2.46 (m, 1 H), 1.68-1.82 (m, 1 H), 1.62 (br. s, 3 H)		
1-24	[5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-phenyl-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	(300 MHz, DMSO-d ₆) 12.99 (br. s, 1 H), 8.62 (d, J = 7.5 Hz, 1 H), 8.47 (s, 3 H), 7.28-7.48 (m, 3 H), 7.24 (d, J = 7.5 Hz, 2 H), 6.92 (s, 1 H), 6.4 (s, 1 H), 4.79 (s, 2 H), 4.64 (br. s, 1 H), 2.63-2.76 (m, 1 H), 2.5-2.57 (m, 1 H), 1.74 (br. s, 4 H)	596 [#]	
1-25	[(<i>R</i>)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-chloro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid			
1-26	[2,3-Difluoro-5-(3-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	12.91 (br. s, 1 H), 8.52 (d, J = 8.3 Hz, 1 H), 8.12 (d, J = 7.8 Hz, 1 H), 8.04 (d, J = 7.8 Hz, 1 H), 8.01 (s, 1 H), 6.64 (dd, J = 11.2, 7.8 Hz, 1 H), 4.71 (d, J = 2.0 Hz, 2 H), 4.33 - 4.56 (m, 1 H), 2.53 - 2.72 (m, 2 H), 1.67 - 1.82 (m, 1 H), 1.47 - 1.67 (m, 3 H)		
1-27	[3-Fluoro-5-(3-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	12.69 (br. s, 1 H), 8.52 (d, J = 8.3 Hz, 1 H), 8.12 (d, J = 8.3 Hz, 1 H), 8.00 - 8.08 (m, 2 H), 6.66 (d, J = 10.3 Hz, 1 H), 6.31 (d, J = 8.8 Hz, 1 H), 4.70 (s, 2 H), 4.39 - 4.51 (m, 1 H), 2.51 - 2.58 (m, 2 H), 1.75 (br. s, 1 H), 1.47 - 1.69 (m, 3 H)		

1-28	[5-(3-Bromo-5-trifluoromethylbenzenesulfonylamino)-3-fluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.02 (br. s, 1 H), 8.52 (d, J = 8.3 Hz, 1 H), 8.37 (s, 1 H), 8.32 (s, 1 H), 8.16 (s, 1 H), 6.66 (d, J = 11.2 Hz, 1 H), 6.30 (d, J = 8.8 Hz, 1 H), 4.71 (s, 2 H), 4.39 - 4.52 (m, 1 H), 2.41 - 2.48 (m, 2 H), 1.75 (br. s, 1 H), 1.58 (br. s, 3 H)	
1-29	[5-(3-Bromo-5-trifluoromethylbenzenesulfonylamino)-2,3-difluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.06 (br. s, 1 H), 8.52 (d, J = 7.8 Hz, 1 H), 8.38 (s, 1 H), 8.31 (s, 1 H), 8.15 (s, 1 H), 6.63 (dd, J = 11.2, 7.3 Hz, 1 H), 4.72 (s, 2 H), 4.29 - 4.51 (m, 1 H), 2.55 - 2.65 (m, 2 H), 1.69 - 1.82 (m, 1 H), 1.49 - 1.69 (m, 3 H)	
1-30	[5-(2,4-Dichlorobenzenesulfonylamino)-2,3-difluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.05 (br. s, 1 H), 8.62 (d, J = 8.8 Hz, 1 H), 8.05 (d, J = 8.5 Hz, 1 H), 7.94 (d, J = 2.0 Hz, 1 H), 7.66 (dd, J = 8.5, 2.0 Hz, 1 H), 6.83 (dd, J = 11.5, 7.6 Hz, 1 H), 4.72 (s, 2 H), 4.19 - 4.35 (m, 1 H), 2.60 (br. s, 2 H), 1.83 (br. s, 1 H), 1.50 - 1.67 (m, 3 H)	
1-31	[5-(2,4-Dichlorobenzenesulfonylamino)-3-fluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.05 (br. s, 1 H), 8.60 (d, J = 9.3 Hz, 1 H), 8.06 (d, J = 8.3 Hz, 1 H), 7.93 (d, J = 2.0 Hz, 1 H), 7.66 (dd, J = 8.3, 2.0 Hz, 1 H), 6.67 (dd, J = 11.0, 2.2 Hz, 1 H), 6.53 (dd, J = 10.0, 2.2 Hz, 1 H), 4.72 (s, 2 H), 4.25 - 4.37 (m, 1 H), 2.50 (s, 2 H), 1.74 - 1.92 (m, 1 H), 1.49 - 1.71 (m, 3 H)	

1-32	[5-(2,5-Dichloro-benzenesulfonylamino)-2,3-difluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.04 (br. s, 1 H), 8.69 (d, J = 9.3 Hz, 1 H), 8.01 (s, 1 H), 7.73 - 7.83 (m, 2 H), 6.80 (dd, J = 11.2, 7.3 Hz, 1 H), 4.72 (s, 2 H), 4.28 - 4.42 (m, 1 H), 2.60 (br. s, 2 H), 1.75 - 1.89 (m, 1 H), 1.49 - 1.71 (m, 3 H)	
1-33	[5-(2,5-Dichloro-benzenesulfonylamino)-3-fluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.03 (br. s, 1 H), 8.67 (d, J = 8.8 Hz, 1 H), 8.02 (s, 1 H), 7.77 (br. s, 2 H), 6.67 (d, J = 10.8 Hz, 1 H), 6.51 (d, J = 9.8 Hz, 1 H), 4.71 (s, 2 H), 4.26 - 4.44 (m, 1 H), 2.50 (br. s, 2 H), 1.85 (br. s, 1 H), 1.63 (br. s, 3 H)	
1-34	{2,3-Difluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydronaphthalen-1-yloxy}-acetic acid	12.82 (br. s, 1 H), 9.10 (s, 1 H), 8.45 (d, J = 8.3 Hz, 1 H), 8.39 (dd, J = 8.2, 1.7 Hz, 1 H), 8.06 (d, J = 8.2 Hz, 1 H), 7.70 (s, 1 H), 7.64 (d, J = 7.3 Hz, 1 H), 7.48 (t, J = 7.3 Hz, 1 H), 7.39 (d, J = 7.3 Hz, 1 H), 6.91 (dd, J = 12.0, 7.6 Hz, 1 H), 4.71 (s, 2 H), 4.34 - 4.61 (m, 1 H), 2.85 - 3.10 (m, 1 H), 2.57 - 2.66 (m, 2 H), 1.62 (br. s, 4 H), 1.28 (d, J = 6.8 Hz, 6 H)	
1-35	{3-Fluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydronaphthalen-1-yloxy}-acetic acid	13.04 (br. s, 1 H), 9.10 (d, J = 2.0 Hz, 1 H), 8.42 (d, J = 8.8 Hz, 1 H), 8.38 (dd, J = 8.0, 2.0 Hz, 1 H), 8.06 (d, J = 8.0 Hz, 1 H), 7.70 (s, 1 H), 7.64 (d, J = 7.8 Hz, 1 H), 7.48 (t, J = 7.8 Hz, 1 H), 7.39 (d, J = 7.8 Hz, 1 H), 6.65 (dd, J = 11.2, 2.0 Hz, 1 H), 6.61 (d, J = 10.3 Hz, 1 H), 4.71 (s, 2 H), 4.44 - 4.55 (m, 1 H), 2.91 - 3.07 (m, 1 H), 2.50 (br. s, 2 H), 1.53 - 1.90 (m, 4 H), 1.28 (d, J = 7.3 Hz, 6 H)	

1-36	{3-Chloro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.04 (br. s, 1 H), 9.09 (s, 1 H), 8.45 (d, J = 8.8 Hz, 1 H), 8.38 (d, J = 7.8 Hz, 1 H), 8.06 (d, J = 7.8 Hz, 1 H), 7.69 (s, 1 H), 7.64 (d, J = 7.3 Hz, 1 H), 7.48 (t, J = 7.3 Hz, 1 H), 7.38 (d, J = 7.3 Hz, 1 H), 6.75 (br. s, 1 H), 6.73 (s, 1 H), 4.67 (br. s, 2 H), 4.49 (br. s, 1 H), 2.88 - 3.09 (m, 1 H), 2.51 - 2.65 (m, 2 H), 1.54 - 1.90 (m, 4 H), 1.27 (d, J = 6.8 Hz, 6 H)		
1-37	[(R)-3-Bromo-5-(3-bromo-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	13.14 (br. s, 1 H), 8.61 (d, J = 8.5 Hz, 1 H), 8.44 (s, 1 H), 8.38 (s, 1 H), 8.21 (s, 1 H), 6.97 (s, 1 H), 6.61 (s, 1 H), 4.80 (s, 2 H), 4.47-4.64 (m, 1 H), 2.58-2.76 (m, 2 H), 1.56-1.91 (m, 4 H)	583	
1-38	[(R)-3-Chloro-2-fluoro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	13.01 (br. s, 1 H), 8.54 (d, J = 7.7 Hz, 1 H), 8.13 (d, J = 8.1 Hz, 1 H), 8.04 (d, J = 8.0 Hz, 1 H), 8.01 (s, 1 H), 6.69 (d, J = 6.8 Hz, 1 H), 4.67 (s, 2 H), 4.42 - 4.52 (m, 1 H), 2.57 - 2.77 (m, 2 H), 1.68 - 1.85 (m, 1 H), 1.49 - 1.70 (m, 3 H)	498	
1-39	[(R)-5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	12.96 (br. s, 1 H), 8.53 (d, J = 6.2 Hz, 1 H), 8.38 (s, 1 H), 8.31 (s, 1 H), 8.14 (s, 1 H), 6.66 (d, J = 6.8 Hz, 1 H), 4.66 (s, 2 H), 4.40 - 4.54 (m, 1 H), 2.56 - 2.75 (m, 2 H), 1.71 - 1.85 (m, 1 H), 1.51 - 1.70 (m, 3 H)	558	

1-40	[(<i>R</i>)-3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.08 (br. s, 1 H), 8.63 (d, <i>J</i> = 9.0 Hz, 1 H), 8.05 (d, <i>J</i> = 8.5 Hz, 1 H), 7.93 (d, <i>J</i> = 2.0 Hz, 1 H), 7.67 (dd, <i>J</i> = 8.5, 2.0 Hz, 1 H), 6.90 (d, <i>J</i> = 6.8 Hz, 1 H), 4.68 (s, 2 H), 4.21 - 4.41 (m, 1 H), 2.59 - 2.70 (m, 2 H), 1.84 (br. s, 1 H), 1.48 - 1.69 (m, 3 H)	480	
1-41	[(<i>R</i>)-3-Chloro-5-(3-fluoro-5-trifluoromethylbenzenesulfonylamino)-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.34 (br. s, 1 H), 8.55 (d, <i>J</i> = 8.2 Hz, 1 H), 8.13 (d, <i>J</i> = 8.2 Hz, 1 H), 8.05 (d, <i>J</i> = 7.7 Hz, 1 H), 8.02 (s, 1 H), 6.79 (d, <i>J</i> = 1.3 Hz, 1 H), 6.46 (s, 1 H), 4.72 (s, 2 H), 4.43 - 4.52 (m, 1 H), 2.39 - 2.51 (m, 2 H), 1.75 (br. s, 1 H), 1.48 - 1.70 (m, 3 H)	480	
1-42	[(<i>R</i>)-5-(3-Bromo-5-trifluoromethylbenzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	12.97 (br. s, 1 H), 8.54 (d, <i>J</i> = 7.9 Hz, 1 H), 8.38 (s, 1 H), 8.32 (s, 1 H), 8.15 (s, 1 H), 6.79 (d, <i>J</i> = 1.3 Hz, 1 H), 6.43 (d, <i>J</i> = 1.3 Hz, 1 H), 4.71 (s, 2 H), 4.42 - 4.53 (m, 1 H), 2.40 - 2.50 (m, 2 H), 1.69 - 1.83 (m, 1 H), 1.49 - 1.72 (m, 3 H)	540	
1-43	[(<i>R</i>)-3-Chloro-5-(2,5-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydronaphthalen-1-yloxy]-acetic acid	13.14 (br. s, 1 H), 8.70 (d, <i>J</i> = 9.0 Hz, 1 H), 8.03 (d, <i>J</i> = 2.2 Hz, 1 H), 7.79 (dd, <i>J</i> = 8.5, 2.2 Hz, 1 H), 7.76 (d, <i>J</i> = 8.5 Hz, 1 H), 6.79 (d, <i>J</i> = 1.5 Hz, 1 H), 6.72 (d, <i>J</i> = 1.5 Hz, 1 H), 4.72 (s, 2 H), 4.38 (br. s, 1 H), 2.46 - 2.49 (m, 2 H), 1.76 - 1.96 (m, 1 H), 1.50 - 1.74 (m, 3 H)	462	
1-44	{3-Chloro-2-fluoro-5-[5-(3-isopropylphenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydronaphthalen-1-yloxy}-acetic acid			

1-45	[5-(2,5-Dichloro-benzenesulfonyl-amino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	12.93 (br. s, 1 H), 8.60 (d, J = 9.0 Hz, 1 H), 8.00 (d, J = 1.8 Hz, 1 H), 7.77 (dd, J = 8.5, 1.8 Hz, 1 H), 7.74 (d, J = 8.5 Hz, 1 H), 7.06 (dd, J = 11.6, 8.7 Hz, 1 H), 6.91 (dd, J = 8.7, 5.1 Hz, 1 H), 4.61 (s, 2 H), 4.32 - 4.44 (m, 1 H), 2.67 (br. s, 2 H), 1.75 - 1.88 (m, 1 H), 1.53 - 1.66 (m, 3 H)	
1-46	[5-(2,4-Dichloro-benzenesulfonyl-amino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	12.94 (br. s, 1 H), 8.52 (d, J = 9.0 Hz, 1 H), 8.05 (d, J = 8.5 Hz, 1 H), 7.91 (d, J = 2.1 Hz, 1 H), 7.65 (dd, J = 8.5, 2.1 Hz, 1 H), 7.07 (dd, J = 11.7, 8.8 Hz, 1 H), 6.94 (dd, J = 8.8, 5.0 Hz, 1 H), 4.61 (s, 2 H), 4.22 - 4.39 (m, 1 H), 2.62 - 2.74 (m, 2 H), 1.83 (br. s, 1 H), 1.58 (br. s, 3 H)	
1-47	{2-Fluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid		
1-48	[(R)-3-Bromo-5-(2,5-dichloro-benzenesulfonyl-amino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	(300 MHz, DMSO-d6) 8.69 (br. s, 1 H), 8.00 (d, J = 2.1 Hz, 1 H), 7.77 (dd, J = 8.8, 2.1 Hz, 1 H), 7.73 (d, J = 8.8 Hz, 1 H), 6.74 (br. s, 1 H), 6.73 (br. s, 1 H), 4.38 (s, 2 H), 4.29-4.36 (m, 1 H), 2.36-2.48 (m, 2 H), 1.70-1.93 (m, 1 H), 1.46-1.70 (m, 3 H).	

- 71 -

1-49	[(<i>R</i>)-3-Bromo-5-(2,4-dichlorobenzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	13.00 (br. s, 1 H), 8.61 (d, <i>J</i> = 9.1 Hz, 1 H), 8.04 (d, <i>J</i> = 8.8 Hz, 1 H), 7.9 (s, 1 H), 7.65 (d, <i>J</i> = 8.5 Hz, 1 H), 6.86 (s, 1 H), 6.76 (s, 1 H), 4.67 (s, 2 H), 4.28 (br. s, 1 H), 2.45 (br. s, 2 H), 1.83 (br. s, 1 H), 1.58 (br. s, 3 H)	505	
1-50	[5-(3,5-Bis-methanesulfonylbenzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	13.18 (br. s, 1 H), 8.65-8.85 (m, 4 H), 6.96 (br. s, 1 H), 6.76 (s, 1 H), 4.79 (br. s, 2 H), 4.57-4.70 (m, 1 H), 3.51 (s, 6 H), 2.50-2.55 (m, 2 H), 1.54-1.91 (m, 4 H)	593	
1-51	[(<i>R</i>)-3-Bromo-5-(3-methoxy-5-trifluoromethylbenzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	(300 MHz, CDCl ₃) 7.74 (s, 1 H), 7.59 (s, 1 H), 7.35 (s, 1 H), 6.67 (d, <i>J</i> = 1.8 Hz, 2 H), 5.38 (d, <i>J</i> = 8.5 Hz, 1 H), 4.61 (s, 2 H), 4.30 - 4.50 (m, 1 H), 3.91 (s, 3 H), 2.42 - 2.77 (m, 2 H), 1.76 (br. s, 4 H)	560 [#]	
1-52	[(<i>R</i>)-3-Bromo-5-(3-fluoro-5-trifluoromethylbenzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid	(300 MHz) 13.04 (br. s, 1 H), 8.55 (d, <i>J</i> = 8.2 Hz, 1 H), 8.13 (d, <i>J</i> = 8.2 Hz, 1 H), 8.05 (d, <i>J</i> = 7.8 Hz, 1 H), 8.02 (s, 1 H), 6.90 (s, 1 H), 6.57 (s, 1 H), 4.74 (s, 2 H), 4.39 - 4.56 (m, 1 H), 2.38 - 2.48 (m, 2 H), 1.70 - 1.86 (m, 1 H), 1.47 - 1.70 (m, 3 H)	524	


[#] (M+Na)⁺

* Resolved using chiral chromatography using Thar Technologies, Inc.'s Multigram[®] III instrument, Daicel[®] OD column 3x25 cm, 25% methanol


5

EXAMPLE 2-1

{5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid




[5-(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid tert-butyl ester



- 5 To a cooled solution of [5-(3,5-bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid tert-butyl ester (example 1-1, 9th step) (169.4 mg, 0.28 mmol) in *N,N*-dimethylformamide (2.5 mL) was added potassium carbonate (0.059 g, 0.42 mmol) and methyl iodide (0.02 mL, 0.31 mmol). The reaction mixture was stirred for 2 hours at room temperature and then extracted
- 10 with ethyl acetate (15 mL x 3). The collected organic layers were washed with water (15 mL x 3), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to give [5-(3,5-bis-trifluoromethyl-benzenesulfonyl)-methyl-amino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid tert-butyl ester (156.0 mg, 90%), which was used in the next step without further purification.

15

{5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid



Starting with [5-(3,5-bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid tert-butyl ester, using an analogous method to example 1-1, [5-(3,5-bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid was obtained (69%). ¹H NMR (400 MHz, DMSO-d₆) δ ppm 13.05 (br. s, 1 H), 8.58 (s, 1 H), 8.56 (s, 2 H), 6.72 (d, J = 6.8 Hz, 1 H), 5.19 - 5.32 (m, 1 H), 4.71 (d, J = 1.7 Hz, 2 H), 2.60 (s, 3 H), 2.54 - 2.88 (m, 2 H), 1.77 - 1.92 (m, 1 H), 1.56 - 1.76 (m, 2 H), 1.51 (m, 1 H); MS calcd. for C₂₁H₁₉ClF₆NO₅S 563, obsd. 562 (M-H⁺).

10

EXAMPLES 2-2 to 2-33

The following examples 2-2 to 2-33 were prepared in an analogous manner to example 2-1. In cases where the example is depicted as a single enantiomer, a chiral separation of enantiomers was performed on the racemic final product, as described above.

15

Example No.	Systematic Name	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm	MS (M-H) ⁻	Structure
2-2	{5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	12.89 (br. s, 1 H), 8.55 (s, 1 H), 8.52 (s, 2 H), 7.01 (d, J = 7.7 Hz, 1 H), 6.75 (d, J = 7.7 Hz, 1 H), 5.17 - 5.32 (m, 1 H), 4.33 (s, 2 H), 2.70 - 2.82 (m, 1 H), 2.58 - 2.64 (m, 1 H), 2.56 (s, 3 H), 2.19 (s, 3 H), 1.73 - 1.88 (m, 1 H), 1.46 - 1.72 (m, 3 H)	524	




2-3	{5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-ethyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	12.86 (br. s, 1 H), 8.55 (s, 1 H), 8.51 (s, 2 H), 7.05 (d, J = 7.8 Hz, 1 H), 6.80 (d, J = 7.8 Hz, 1 H), 5.21 - 5.29 (m, 1 H), 4.23 - 4.39 (m, 2 H), 2.57 (s, 3 H), 2.52 - 2.81 (m, 4 H), 1.72 - 1.88 (m, 1 H), 1.43 - 1.73 (m, 3 H), 1.12 (t, J=7.3 Hz, 3 H)	538	
2-4	{5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.09 (br. s, 1 H), 8.57 (s, 1 H), 8.55 (s, 2 H), 6.83 (d, J = 2.0 Hz, 1 H), 6.49 (s, 1 H), 5.24 - 5.32 (m, 1 H), 4.74 (dd, J = 18.3, 16.4 Hz, 2 H), 2.60 (s, 3 H), 2.54 - 2.73 (m, 2 H), 1.79 - 1.90 (m, 1 H), 1.59 - 1.77 (m, 2 H), 1.52 (m, 1 H)	544	
2-5	{(R)-5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid		544	
2-6	{3-Chloro-5-[(2,4-dichloro-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	12.33 - 13.99 (m, 1 H), 8.10 (d, J = 8.5 Hz, 1 H), 7.99 (d, J = 2.0 Hz, 1 H), 7.68 (dd, J = 8.5, 2.0 Hz, 1 H), 6.83 (s, 1 H), 6.78 (s, 1 H), 4.94 (dd, J = 10.5, 5.6 Hz, 1 H), 4.72 (s, 2 H), 2.66 - 2.76 (m, 1 H), 2.64 (s, 3 H), 2.54 (br. s, 1 H), 2.28 - 2.44 (m, 1 H), 1.54 - 1.97 (m, 3 H)	476/478	
2-7	{5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.05 (br. s, 1 H), 8.57 (s, 1 H), 8.55 (s, 2 H), 6.90 (s, 1 H), 5.25 - 5.32 (m, 1 H), 4.51 (s, 2 H), 2.76 - 2.87 (m, 1 H), 2.62 (s, 3 H), 2.52 - 2.60 (m, 1 H), 1.75 - 1.89 (m, 1 H), 1.58 - 1.75 (m, 2 H), 1.53 (br. s., 1 H)		

2-8	{5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	12.96 (br. s, 1 H), 8.57 (s, 1 H), 8.54 (s, 2 H), 6.69 (s, 1 H), 5.21 - 5.30 (m, 1 H), 4.36 (s, 2 H), 2.68 (d, J = 11.7 Hz, 1 H), 2.60 (s, 3 H), 2.40 - 2.46 (m, 1 H), 2.22 (s, 3 H), 1.81 (br. s, 1 H), 1.45 - 1.73 (m, 3 H)		
2-9	{3-Chloro-5-[(2,5-dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.13 (br. s, 1 H), 8.08 (t, J = 1.5 Hz, 1 H), 7.81 - 7.83 (m, 2 H), 6.93 (d, J = 6.6 Hz, 1 H), 4.96 (dd, J = 10.3, 5.4 Hz, 1 H), 4.67 (s, 2 H), 2.83 (m, 1 H), 2.68 (s, 3 H), 2.54 (br. s, 1 H), 1.82 - 1.96 (m, 1 H), 1.72 - 1.82 (m, 1 H), 1.54 - 1.74 (m, 2 H)	496/49 8	
2-10	{3-Chloro-5-[(2,5-dichloro-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	8.06 (s, 1 H), 7.65 - 7.89 (m, 2 H), 6.76 (d, J = 2.0 Hz, 1 H), 6.73 (br. s, 1 H), 4.94 (dd, J = 10.3, 5.9 Hz, 1 H), 4.67 (s, 2 H), 2.64 (s, 3 H), 2.61 (br. s, 1 H), 2.29 - 2.44 (m, 1 H), 1.84 - 1.93 (m, 1 H), 1.71 - 1.84 (m, 1 H), 1.51 - 1.71 (m, 2 H)	476/47 8	
2-11	{3-Chloro-5-[(2,4-dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.07 (br. s, 1 H), 8.10 (d, J = 8.5 Hz, 1 H), 8.00 (d, J = 1.5 Hz, 1 H), 7.69 (dd, J = 8.5, 1.5 Hz, 1 H), 6.96 (d, J = 6.6 Hz, 1 H), 4.82 - 4.98 (m, 1 H), 4.68 (br. s, 2 H), 2.71 - 2.92 (m, 1 H), 2.65 (s, 3 H), 2.57 - 2.62 (m, 1 H), 1.85 - 1.98 (m, 1 H), 1.53 - 1.83 (m, 3 H)	496/49 8	


2-12	{5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-fluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy}-acetic acid	13.00 (br. s, 1 H), 8.43 (br. s, 1 H), 8.42 (br. s, 1 H), 8.21 (s, 1 H), 6.72 (d, J = 6.8 Hz, 1 H), 5.14 - 5.24 (m, 1 H), 4.69 (s, 2 H), 2.75 - 2.90 (m, 1 H), 2.59 (s, 3 H), 2.42 - 2.46 (m, 1 H), 1.83 (br. s, 1 H), 1.56 - 1.76 (m, 2 H), 1.45 - 1.56 (m, 1 H)	574/57 6	
2-13	{5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydronaphthalen-1-yloxy}-acetic acid	13.05 (br. s, 1 H), 8.42 (d, J = 6.4 Hz, 2 H), 8.20 (s, 1 H), 6.83 (s, 1 H), 6.50 (s, 1 H), 5.16 - 5.25 (m, 1 H), 4.73 (s, 2 H), 2.68 (d, J = 16.1 Hz, 1 H), 2.58 (s, 3 H), 2.36 (dd, J = 18.8, 10.5 Hz, 1 H), 1.85 (br. s, 1 H), 1.59 - 1.77 (m, 2 H), 1.45 - 1.59 (m, 1 H)	554/55 6	
2-14	{3-Chloro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydronaphthalen-1-yloxy}-acetic acid	13.09 (br. s, 1 H), 8.14 - 8.23 (m, 2 H), 8.06 (s, 1 H), 6.83 (s, 1 H), 6.51 (s, 1 H), 5.15 - 5.23 (m, 1 H), 4.73 (s, 2 H), 2.68 (d, J = 16.1 Hz, 1 H), 2.59 (s, 3 H), 2.36 (dd, J = 17.4, 10.5 Hz, 1 H), 1.84 (br. s, 1 H), 1.59 - 1.75 (m, 2 H), 1.45 - 1.58 (m, 1 H)	494/49 6	
2-15	{3-Chloro-2-fluoro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydronaphthalen-1-yloxy}-acetic acid	13.02 (br. s, 1 H), 8.19 (t, J = 9.8 Hz, 2 H), 8.06 (s, 1 H), 6.73 (d, J = 6.4 Hz, 1 H), 5.07 - 5.22 (m, 1 H), 4.70 (s, 2 H), 2.82 (d, J = 17.1 Hz, 1 H), 2.59 (s, 3 H), 2.42 - 2.48 (m, 1 H), 1.83 (br. s, 1 H), 1.55 - 1.76 (m, 2 H), 1.43 - 1.55 (m, 1 H)	512/51 4	
2-16	{5-[(2,4-Dichloro-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydronaphthalen-1-yloxy}-acetic acid	13.03 (br. s, 1 H), 8.10 (d, J = 8.5 Hz, 1 H), 7.99 (d, J = 2.0 Hz, 1 H), 7.68 (dd, J = 8.5, 2.0 Hz, 1 H), 6.70 (d, J = 11.7 Hz, 1 H), 6.56 (d, J = 10.3 Hz, 1 H), 4.88 - 4.98 (m, 1 H), 4.71 (s, 2 H), 2.63 (s, 3 H), 2.59 - 2.74 (m, 1 H), 2.26 - 2.44 (m, 1 H), 1.89 (br. s, 1 H), 1.53 - 1.85 (m, 3 H)	460/46 2	

2-17	{5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	12.88 (br. s, 1 H), 8.55 (s, 1 H), 8.52 (s, 2 H), 7.10 (dd, J = 11.7, 8.7 Hz, 1 H), 6.84 (dd, J = 8.7, 5.1 Hz, 1 H), 5.22 - 5.30 (m, 1 H), 4.64 (s, 2 H), 2.78 - 2.90 (m, 1 H), 2.57 (s, 3 H), 2.43 - 2.49 (m, 1 H), 1.80 (br. s, 1 H), 1.50 - 1.76 (m, 2 H), 1.46 (br. s, 1 H)	
2-18	{5-[(2,4-Dichloro-benzenesulfonyl)-methyl-amino]-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.04 (br. s, 1 H), 8.10 (d, J = 8.5 Hz, 1 H), 8.00 (d, J = 2.0 Hz, 1 H), 7.68 (dd, J = 8.5, 2.0 Hz, 1 H), 6.84 (dd, J = 11.5, 7.6 Hz, 1 H), 4.91 (dd, J = 9.0, 6.1 Hz, 1 H), 4.74 (s, 2 H), 2.81 (d, J = 18.1 Hz, 1 H), 2.64 (s, 3 H), 2.41 - 2.48 (m, 1 H), 1.50 - 1.95 (m, 4 H)	
2-19	{5-[(2,5-Dichloro-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.03 (br. s, 1 H), 8.08 (s, 1 H), 7.81 (s, 2 H), 6.70 (d, J = 10.3 Hz, 1 H), 6.53 (d, J = 10.3 Hz, 1 H), 4.99 (dd, J = 10.0, 5.1 Hz, 1 H), 4.72 (s, 2 H), 2.66 - 2.76 (m, 1 H), 2.66 (s, 3 H), 2.29 - 2.44 (m, 1 H), 1.56 - 1.95 (m, 4 H)	
2-20	{5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.04 (br. s, 1 H), 8.43 (s, 1 H), 8.41 (s, 1 H), 8.21 (s, 1 H), 6.66 (dd, J = 11.0, 7.6 Hz, 1 H), 5.10 - 5.23 (m, 1 H), 4.74 (s, 2 H), 2.78 (d, J = 17.6 Hz, 1 H), 2.59 (s, 3 H), 2.36 - 2.45 (m, 1 H), 1.82 (br. s, 1 H), 1.37 - 1.76 (m, 3 H)	
2-21	{5-[(2,5-Dichloro-benzenesulfonyl)-methyl-amino]-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.54 (br. s, 1 H), 8.08 (s, 1 H), 7.82 (s, 2 H), 6.80 (dd, J = 11.5, 7.6 Hz, 1 H), 4.96 (dd, J = 10.0, 5.6 Hz, 1 H), 4.73 (s, 2 H), 2.74 - 2.86 (m, 1 H), 2.67 (s, 3 H), 2.39 - 2.47 (m, 1 H), 1.54 - 1.93 (m, 4 H)	

2-22	{5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.05 (br. s, 1 H), 8.43 (br. s, 1 H), 8.41 (br. s, 1 H), 8.20 (s, 1 H), 6.70 (dd, J = 10.8, 1.5 Hz, 1 H), 6.35 (dd, J = 9.8, 1.5 Hz, 1 H), 5.12 - 5.27 (m, 1 H), 4.72 (s, 2 H), 2.67 (d, J = 17.6 Hz, 1 H), 2.58 (s, 3 H), 2.27 - 2.43 (m, 1 H), 1.84 (br. s, 1 H), 1.56 - 1.77 (m, 2 H), 1.52 (br. s, 1 H)		
2-23	{3-Fluoro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.05 (br. s, 1 H), 8.20 (d, J = 7.8 Hz, 1 H), 8.17 (d, J = 8.8 Hz, 1 H), 8.06 (s, 1 H), 6.70 (dd, J = 10.8, 2.0 Hz, 1 H), 6.36 (d, J = 9.8 Hz, 1 H), 5.13 - 5.23 (m, 1 H), 4.73 (s, 2 H), 2.67 (d, J = 16.1 Hz, 1 H), 2.59 (s, 3 H), 2.26 - 2.43 (m, 1 H), 1.84 (br. s, 1 H), 1.57 - 1.76 (m, 2 H), 1.44 - 1.57 (m, 1 H)		
2-24	{2,3-Difluoro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.05 (br. s, 1 H), 8.20 (d, J = 7.8 Hz, 1 H), 8.16 (d, J = 8.8 Hz, 1 H), 8.06 (s, 1 H), 6.66 (dd, J = 11.5, 7.6 Hz, 1 H), 5.07 - 5.20 (m, 1 H), 4.74 (s, 2 H), 2.78 (d, J = 16.6 Hz, 1 H), 2.59 (s, 3 H), 2.39 - 2.48 (m, 1 H), 1.82 (br. s, 1 H), 1.54 - 1.75 (m, 2 H), 1.42 - 1.53 (m, 1 H)		
2-25	(2,3-Difluoro-5-[(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	13.04 (br. s, 1 H), 9.14 (s, 1 H), 8.41 (d, J = 8.2 Hz, 1 H), 8.08 (d, J = 8.2 Hz, 1 H), 7.71 (br. s, 1 H), 7.65 (d, J = 7.1 Hz, 1 H), 7.48 (t, J = 7.1 Hz, 1 H), 7.39 (d, J = 7.1 Hz, 1 H), 6.86 (dd, J = 10.9, 7.7 Hz, 1 H), 5.07 (br. s, 1 H), 4.74 (br. s, 2 H), 2.89 - 3.08 (m, 1 H), 2.81 (d, J = 17.1 Hz, 1 H), 2.65 (s, 3 H), 2.40 - 2.48 (m, 1 H), 1.88 (br. s, 1 H), 1.73 (br. s, 3 H), 1.28 (d, J = 6.6 Hz, 6 H)		

2-26	(3-Fluoro-5-([5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid	13.06 (br. s, 1 H), 9.14 (s, 1 H), 8.41 (d, J = 8.4 Hz, 1 H), 8.08 (d, J = 8.4 Hz, 1 H), 7.71 (s, 1 H), 7.65 (d, J = 7.8 Hz, 1 H), 7.48 (t, J = 7.8 Hz, 1 H), 7.39 (d, J = 7.8 Hz, 1 H), 6.70 (d, J = 10.5 Hz, 1 H), 6.56 (d, J = 10.0 Hz, 1 H), 5.01 - 5.13 (m, 1 H), 4.72 (s, 2 H), 2.93 - 3.07 (m, 1 H), 2.67 - 2.75 (m, 1 H), 2.65 (s, 3 H), 2.30 - 2.45 (m, 1 H), 1.89 (br. s, 1 H), 1.55 - 1.81 (m, 3 H), 1.28 (d, 6 H)	
2-27	(3-Chloro-5-([5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid	12.88 (br. s, 1 H), 9.14 (d, J = 2.0 Hz, 1 H), 8.41 (dd, J = 8.2, 2.0 Hz, 1 H), 8.09 (d, J = 8.2 Hz, 1 H), 7.70 (s, 1 H), 7.65 (d, J = 8.1 Hz, 1 H), 7.44 - 7.51 (m, 1 H), 7.39 (d, J = 7.6 Hz, 1 H), 6.82 (d, J = 1.2 Hz, 1 H), 6.71 (s, 1 H), 4.98 - 5.15 (m, 1 H), 4.73 (s, 2 H), 2.90 - 3.07 (m, 1 H), 2.68 - 2.75 (m, 1 H), 2.66 (s, 3 H), 2.31 - 2.46 (m, 1 H), 1.88 (br. s., 1 H), 1.55 - 1.80 (m, 3 H), 1.27 (d, 6 H)	
2-28	(3-Chloro-2-fluoro-5-([5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid	13.02 (s, 1 H), 9.14 (br. s, 1 H), 8.41 (d, J = 8.8 Hz, 1 H), 8.09 (d, J = 7.3 Hz, 1 H), 7.70 (br. s, 1 H), 7.61 - 7.68 (m, 1 H), 7.43 - 7.53 (m, 1 H), 7.40 (br. s, 1 H), 6.87 - 6.98 (m, 1 H), 5.07 (br. s, 1 H), 4.70 (br. s, 2 H), 3.00 (br. s, 1 H), 2.75 - 2.90 (m, 1 H), 2.66 (br. s, 3 H), 2.41 - 2.48 (m, 1 H), 1.89 (br. s, 1 H), 1.54 - 1.81 (m, 3 H), 1.27 (d, J = 5.9 Hz, 6 H)	

2-29	{5-[(2,5-Dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	12.92 (br. s, 1 H), 8.07 (s, 1 H), 7.80 (s, 2 H), 7.12 (dd, J = 11.7, 8.6 Hz, 1 H), 6.91 (dd, J = 8.6, 5.0 Hz, 1 H), 4.94 - 5.04 (m, 1 H), 4.64 (s, 2 H), 2.83 - 2.94 (m, 1 H), 2.64 (s, 3 H), 2.52 - 2.60 (m, 1 H), 1.84 - 1.93 (m, 1 H), 1.68 (br. s, 3 H)		
2-30	{5-[(2,4-Dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	12.93 (br. s, 1 H), 8.09 (d, J = 8.2 Hz, 1 H), 7.98 (s, 1 H), 7.68 (d, J = 8.2 Hz, 1 H), 7.13 (dd, J = 11.2, 8.8 Hz, 1 H), 6.94 (dd, J = 8.8, 4.9 Hz, 1 H), 4.85 - 5.04 (m, 1 H), 4.64 (s, 2 H), 2.79 - 2.96 (m, 1 H), 2.61 (s, 3 H), 2.53 - 2.58 (m, 1 H), 1.82 - 1.94 (m, 1 H), 1.74 (br. s, 3 H)		
2-31	(2-Fluoro-5-{[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino}-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid	12.93 (br. s, 1 H), 9.14 (d, J = 2.1 Hz, 1 H), 8.40 (dd, J = 8.2, 2.1 Hz, 1 H), 8.07 (d, J = 8.2 Hz, 1 H), 7.72 (s, 1 H), 7.66 (d, J = 7.8 Hz, 1 H), 7.48 (t, J = 7.8 Hz, 1 H), 7.39 (d, J = 7.8 Hz, 1 H), 7.13 (dd, J = 12.0, 8.8 Hz, 1 H), 7.00 (dd, J = 8.8, 5.3 Hz, 1 H), 5.00 - 5.19 (m, 1 H), 4.65 (s, 2 H), 2.95 - 3.07 (m, 1 H), 2.79 - 2.95 (m, 1 H), 2.63 (s, 3 H), 2.52 - 2.61 (m, 1 H), 1.81 - 1.96 (m, 1 H), 1.51 - 1.77 (m, 3 H), 1.28 (d, J = 6.8 Hz, 6 H)		
2-32	{{(R)-3-Bromo-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid	(300 MHz, CDCl ₃) 7.95 (s, 1 H), 7.79 (d, J = 7.5 Hz, 1 H), 7.59 (d, J = 7.8 Hz, 1 H), 6.87 (s, 1 H), 6.72 (s, 1 H), 5.07 - 5.23 (m, 1 H), 4.66 (s, 2 H), 2.81 (d, J = 17.8 Hz, 1 H), 2.64 (s, 3 H), 2.34 - 2.51 (m, 1 H), 1.89 - 2.02 (m, 1 H), 1.65 (d, J = 7.2 Hz, 3 H)	562 [#]	

2-33	{{(R)-3-Bromo-5-[(3-methoxy-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydronaphthalen-1-yloxy}-acetic acid	(300 MHz, CDCl ₃) 7.71 (s, 1 H), 7.55 (s, 1 H), 7.35 (s, 1 H), 6.89 (s, 1 H), 6.71 (s, 1 H), 5.01 - 5.25 (m, 1 H), 4.64 (s, 2 H), 3.93 (s, 3 H), 2.80 (d, J = 16.3 Hz, 1 H), 2.62 (s, 3 H), 2.30 - 2.53 (m, 1 H), 1.83 - 2.02 (m, 1 H), 1.53 - 1.80 (m, 3 H)	574 [#]	
------	--	--	------------------	---

[#] (M+Na)⁺

ACTIVITY AND USE OF THE COMPOUNDS

The compounds of formula I possess valuable pharmacological properties. It has

- 5 been found that said compounds are antagonists at the CRTH2 receptor and may be useful in treating diseases and disorders associated with that receptor such as asthma. The activity of the present compounds as CRTH2 receptor antagonists is demonstrated by the following biological assays.

10 Human CRTH2 Receptor Binding Assay

A whole cell receptor binding assay using [³H]ramatroban as the competing radioactive ligand was employed to evaluate the compound binding activity to human CRTH2. The radioactive ligand [³H]ramatroban was synthesized according to Sugimoto *et. al.* (*Eur. J. Pharmacol.* 524, 30 - 37, 2005) to a specific activity of 42

15 Ci/mmol.

A cell line stably expressing human CRTH2 was established by transfecting CHO-K1 cells with two mammalian expression vectors that harbored human CRTH2 and G-alpha16 cDNAs, respectively, using FuGene[®] 6 transfection reagent (from Roche).

- 20 Stable clones expressing CRTH2 were selected by staining each clone with BM16 (BD Pharmingen[™] from BD Biosciences, a division of Becton, Dickinson and Company), which is a rat monoclonal antibody to human CRTH2. The cells were maintained as monolayer cultures in Ham's F-12 medium containing 10% fetal bovine serum, 100 units/mL penicillin, 100 mg/mL streptomycin, 2 mM glutamine, 0.5
- 25 mg/mL G418 (geneticin) for CRTH2, and 0.2 mg/mL hygromycin-B (for G-alpha 16). For whole cell receptor binding assay, the monolayer cells were rinsed once with

- 82 -

PBS (phosphate buffered saline), dissociated using diammonium ethylenediaminetetraacetate (Versene™ EDTA from the Dow Chemical Company), and suspended in PBS containing 10 mM MgCl₂ and 0.06% BSA (bovine serum albumin) at 1.5×10^6 cells/mL.

5

The binding reactions (0.2 mL) were performed in 96-well plates at room temperature in PBS containing 1.5×10^5 cells, 10 mM MgCl₂, 0.06% BSA, 20 nM [³H]ramatroban, and test compound at various concentrations. After 1 hour of binding reactions, the cells were harvested on GF™/B filter microplates (microtiter plates with embedded glass fiber from PerkinElmer, Inc.) and washed 5 times with PBS using a Filtermate™ Harvester (a cell harvester that harvests and washes cells from microplates from PerkinElmer, Inc.). The radioactivities bound to the cells were determined using a microplate scintillation counter (TopCount® NXT, from PerkinElmer, Inc.) after adding 50 µL of Microscint™ 20 scintillation fluid (from PerkinElmer, Inc.) to each well of the filter plates. The radioactivity from non-specific binding was determined by replacing compound with 10 µM of 15(R)-15-methyl PGD₂ (from Cayman Chemical Company) in the reaction mixtures. The radioactivity bound to the cells in the absence of compound (total binding) was determined by replacing compound with 0.25% of DMSO (dimethyl sulfoxide) in the reaction mixture.

Specific binding data were obtained by subtracting the radioactivity of non-specific binding from each binding data.

The IC₅₀ value is defined as the concentration of the tested compound that is required for 50% inhibition of total specific binding. In order to calculate the IC₅₀ value, the percent inhibition data were determined for 7 concentrations for each compound. The percent inhibition for a compound at each concentration was calculated according to the following formula, $[1 - (\text{specific binding in the presence of compound}) / (\text{total specific binding})] \times 100$. The IC₅₀ value was then obtained by fitting the percent inhibition data to a sigmoidal dose-response (4 parameter logistic) model in the XLfit® software Excel add-in program [from ID Business Solutions Ltd., model 205, where $F(x) = (A + (B - A) / (1 + ((C/x)^D)))$].

The acid compounds of the foregoing examples were tested using the above Human CRTH2 Receptor Binding Assay. The results of the assay showed that all of these compounds have binding activity exhibiting IC₅₀ values ranging from 0.0024 µM to 1.1561 µM. For instance, the following table shows the specific IC₅₀ values for these

5 compounds:

Example No.	Human CRTH2 Binding IC ₅₀ (µM)
Example 1-1	0.0024
Example 1-2	0.1950
Example 1-3	0.1470
Example 1-4	0.0640
Example 1-5	0.0284
Example 1-6	0.0058
Example 1-7	0.2920
Example 1-8	0.0035
Example 1-9	0.0058
Example 1-10	0.0114
Example 1-11	0.0035
Example 1-12	0.0261
Example 1-13	0.0071
Example 1-14	0.0077
Example 1-15	0.008
Example 1-16	0.0038
Example 1-17	0.0036
Example 1-18	0.0028
Example 1-19	0.0029
Example 1-20	0.0025
Example 1-21	0.0028
Example 1-22	0.0337
Example 1-23	0.0036

- 84 -

Example 1-24	0.0037
Example 1-25	0.0959
Example 1-26	0.0422
Example 1-27	0.0401
Example 1-28	0.0042
Example 1-29	0.0182
Example 1-30	0.0461
Example 1-31	0.1544
Example 1-32	0.2686
Example 1-33	0.081
Example 1-34	0.0086
Example 1-35	0.007
Example 1-36	0.0045
Example 1-37	0.0028
Example 1-38	0.0034
Example 1-39	0.0026
Example 1-40	0.0063
Example 1-41	0.0034
Example 1-42	0.0032
Example 1-43	0.0109
Example 1-44	0.0164
Example 2-1	0.0080
Example 2-2	0.3110
Example 2-3	0.1770
Example 2-4	0.0083
Example 2-5	0.0059
Example 2-6	0.0929
Example 2-7	0.3333
Example 2-8	0.6379
Example 2-9	1.1561

- 85 -

Example 2-10	0.8594
Example 2-11	0.5357
Example 2-12	0.0123
Example 2-13	0.0062
Example 2-14	0.0112
Example 2-15	0.0141
Example 2-16	0.9810
Example 2-17	0.1990
Example 2-18	2.5240
Example 2-19	2.6428
Example 2-20	0.2884
Example 2-21	2.1016
Example 2-22	0.0054
Example 2-23	0.1983
Example 2-24	0.3799

Calcium Flux Assay Using Fluorometric Imaging Plate Reader

Cell Culture Conditions:

CHO-K1 cells previously transfected with G-alpha 16 were subsequently transfected
5 with the human CRTH2 receptor and the neomycin resistance gene. Following
selection in 800 µg/mL G418 (geneticin), individual clones were assayed for their
receptor expression based on staining with an anti human CRTH2 IgG, followed by
assaying for their response to 13,14-dihydro-15-keto Prostaglandin D₂ (DK-PDG₂)
(ligand) in the Ca²⁺ Flux assay. Positive clones were then cloned by limiting dilution
10 cloning. The transfected cells were cultured in Ham's F-12 medium supplemented
with 10% fetal bovine serum, 2 mM glutamine , 100 U/mL penicillin/100 µg/mL
streptomycin, 200 µg/mL hygromycin B, and 800 µg/mL G418 (geneticin). Cells
were harvested with trypsin-EDTA (trypsin-ethylenediaminetetraaceticacid) and
counted using ViaCount[®] reagent (from Guava Technologies, Inc. which contains
15 two DNA-binding dyes that enable the reagent user to distinguish between viable
and non-viable cells). The cell suspension volume was adjusted to 2.5 x10⁵ cells
/mL with complete growth media. Aliquots of 50 µL were dispensed into BD

- 86 -

Falcon™ 384 well black/clear microplates (from BD Biosciences, a division of Becton, Dickinson and Company) and the microplates were placed in a 37 °C CO₂ incubator overnight. The following day, the microplates were used in the assay.

5 ***Dye Loading and Assay:***

- Loading Buffer containing dye (from the FLIPR® Calcium 3 Assay Kit from Molecular Devices, a division of MDS Analytical Technologies and MDS Inc.) was prepared by dissolving the contents of one bottle into 200 mL Hank's Balanced Salt Solution containing 20mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid) and
- 10 2.5 mM probenecid. Growth media was removed from the cell plates and 25 µL of Hank's Balanced Salt Solution (HBSS) containing 20 mM HEPES, 0.05% BSA and 2.5 mM probenecid was added to each well followed by 25 µL of diluted dye using a Multidrop dispenser. The plates were then incubated for 1 hour at 37 °C.
- 15 During the incubation, test compound plates were prepared by adding 90 µL of HBSS/20 mM HEPES/0.005% BSA buffer to the 2 µL of serial diluted compounds. To prepare serial diluted compounds, 20 mM stocks of compounds were dissolved in 100% DMSO. The compound dilution plate was set up as follows: well # 1 received 5 µL of compound plus 10 µL of DMSO. Wells 2-10 received 10 µL of DMSO. 5 µL
- 20 was mixed and transferred from well #1 into well #2. 1:3 serial dilutions were continued out 10 steps. 2 µL of diluted compound was transferred into duplicate wells of a 384 well "assay plate" and then 90 µL of buffer was added.

- After incubation, both the cell and "assay plate" plates were brought to the
- 25 fluorometric imaging plate reader (FLIPR®) and 20 µL of the diluted compounds were transferred to the cell plates by the FLIPR®. Plates were then incubated for 1 hour at room temperature. After the hour incubation, plates were returned to the FLIPR® and 20 µL of 4.5X concentrated ligand was added to the cell plates. During the assay, fluorescence readings were taken simultaneously from all 384 wells of the cell
- 30 plate every 1.5 seconds. Five readings were taken to establish a stable baseline, then 20 µL of sample was rapidly (30 µL/sec) and simultaneously added to each well of the cell plate. The fluorescence was continuously monitored before, during and after sample addition for a total elapsed time of 100 seconds. Responses (increase

- 87 -

in peak fluorescence) in each well following agonist addition were determined. The initial fluorescence reading from each well, prior to ligand stimulation, was used as a zero baseline value for the data from that well. The responses were expressed as % inhibition of the buffer control. The IC₅₀ value, defined as the concentration of a compound that was required for 50% inhibition of the buffer control, was calculated by fitting the percent inhibition data for 10 concentrations to a sigmoidal dose-response (4 parameter logistic) model using Genedata Screener[®] Condoseo software program [from Genedata AG, model 205, where $F(x) = (A + (B - A) / (1 + ((C/x)^D)))$].

Representative compounds tested in the binding assay were tested using the above FLIPR[®] assay (examples 1-1 to 1-17, 1-24, 1-25, 2-1 to 2-14). The results of the FLIPR[®] assay showed that all of the representative compounds tested in this assay have activity exhibiting IC₅₀ values ranging from 0.0001 µM to 3.46 µM.

DK-PGD₂-induced IL-13 production assay in Th2 cells

Inhibition of 13,14-dihydro-15-keto Prostaglandin D₂ (DK-PGD₂)-induced IL-13 production in T helper type 2 (Th2) cells was applied to evaluate compound cellular potency.

Cultures of Th2 cells were established from blood of healthy human volunteers according to the following procedure. Peripheral blood mononuclear cells (PBMC) were first isolated from 50 mL of fresh blood by Ficoll-Hypaque density gradient centrifugation, followed by CD4⁺ cell purification using a CD4⁺ T Cell Isolation Kit II (from Miltenyi Biotec Inc.). The CD4⁺ T cells were then differentiated to Th2 cells by culturing the cells in X-VIVO 15[®] medium (from Cambrex BioScience Walkersville Inc.) containing 10% human AB serum (serum of blood type AB from Invitrogen Corporation), 50 U/mL of recombinant human interleukin-2 (rhIL-2) (from PeproTech Inc.) and 100 ng/mL of recombinant human interleukin-4 (rhIL-4) (from PeproTech Inc.) for 7 days. The Th2 cells were isolated using a CD294 (CRTH2) MicroBead Kit (from Miltenyi Biotec Inc.) and amplified in X-VIVO 15[®] medium containing 10% human AB serum and 50 U/mL of rhIL-2 for 2 to 5 weeks. In general, 70% to 80% of the Th2 cells used in the assay are CRTH2-positive when analyzed by fluorescence-

- 88 -

activated cell sorting using the BM16 antibody (as previously described) conjugated to phycoerythrin (PE).

To determine cellular inhibitory potency, compounds at various concentrations were
5 incubated with 2.5×10^4 Th2 cells and 500 nM DK-PGD₂ for 4 hrs at 37 °C in 200 µL
of X-VIVO 15[®] medium containing 10% human AB serum. IL-13 production to the
medium was detected by ELISA (enzyme-linked immunosorbent assay) using an
“Instant ELISA™” kit (from Bender MedSystems Inc.) according to the procedure
suggested by the vendor. The spontaneous production of IL-13 by Th2 cells was
10 determined in the absence of DK-PGD₂ stimulation and the value was subtracted
from that in the presence of each compound for percent inhibition and IC₅₀
calculations.

The percent inhibition of interleukin 13 (IL-13) production for a compound at various
15 concentrations was calculated according to the following formula, $[1 - (\text{IL-13 production in the presence of compound}) / (\text{IL-13 production in the presence of 0.15\% DMSO})] \times 100$. The IC₅₀ value, defined as the concentration of a compound that is
required for 50% inhibition of IL-13 production, was calculated by fitting the percent
inhibition data for 7 concentrations to a sigmoidal dose-response (4 parameter
20 logistic) model in the XLfit[®] software Excel add-in program [ID Business Solutions
Ltd., model 205, where $F(x) = (A + (B - A) / (1 + ((C/x)^D)))$].

Representative compounds tested in the binding assay were tested using the
foregoing DK-PGD₂-induced IL-13 production assay (examples 1-1 to 1-18, 1-20 to
25 1-25, 1-37 to 1-43, 2-01 to 2-17). The results of the DK-PGD₂-induced IL-13
production assay showed that, with the exception of examples 1-2, 2-2, 2-3, 2-7 to 2-
9, 2-11, and 2-16 (which exhibited IC₅₀ values greater than 10), the compounds
tested in this assay exhibited activity in inhibiting IL-13 production, with IC₅₀ values
ranging from 0.004 µM to 5.982 µM.

30

Thus, the compounds of the present invention are useful since the compounds
tested show some activity in at least one of the above three assays (i.e., binding at

- 89 -

the CRTH2 receptor), and therefore may be useful as antagonists in treating diseases and disorders associated with this receptor such as asthma.

In one embodiment, the present invention relates to a method for the treatment
5 and/or prevention of diseases and disorders which are associated with the
modulation of CRTH2 receptors, which method comprises administering a
therapeutically effective amount of a compound of formula I to a human being or
animal. A method for the treatment and/or prevention of an inflammatory or allergic
disease or disorder is preferred. Such diseases or disorders may include (but are
10 not limited to) asthma, chronic obstructive pulmonary disease (COPD), allergic
inflammation, and atopic dermatitis.

The present invention is also directed to the administration of a therapeutically
effective amount of a compound of formula I in combination or association with other
15 drugs or active agents for the treatment of inflammatory or allergic diseases and
disorders. In one embodiment, the present invention relates to a method for the
treatment and/or prevention of such diseases or disorders comprising administering
to a human or animal simultaneously, sequentially, or separately, a therapeutically
effective amount of a compound of formula I and another drug or active agent (such
20 as another anti-inflammatory or anti-allergic drug or agent). These other drugs or
active agents may have the same, similar, or a completely different mode of action.
Suitable other drugs or active agents may include, but are not limited to: Beta2-
adrenergic agonists such as albuterol or salmeterol; corticosteroids such as
dexamethasone or fluticasone; antihistamines such as loratidine; leukotriene
25 antagonists such as montelukast or zafirlukast; anti-IgE antibody therapies such as
omalizumab; anti-infectives such as fusidic acid (particularly for the treatment of
atopic dermatitis); anti-fungals such as clotrimazole (particularly for the treatment of
atopic dermatitis); immunosuppressants such as tacrolimus and pimecrolimus; other
antagonists of PGD₂ acting at other receptors such as DP antagonists; inhibitors of
30 phosphodiesterase type 4 such as cilomilast; drugs that modulate cytokine production
such as inhibitors of TNF- α converting enzyme (TACE); drugs that modulate the
activity of Th2 cytokines IL-4 and IL-5 such as blocking monoclonal antibodies and

- 90 -


soluble receptors; PPAR-gamma agonists such as rosiglitazone; and 5-lipoxygenase inhibitors such as zileuton.

Unless stated to the contrary, all compounds in the examples were prepared and
5 characterized as described. All patents and publications cited herein are hereby
incorporated by reference in their entirety.

- 91 -

CLAIMS

1. A compound of formula I:



or a pharmaceutically acceptable salt or ester thereof, wherein:

R¹ is hydrogen, halogen, methoxy, or phenyl; and R² is hydrogen, fluoro, chloro, or alkyl; with the proviso that R¹ and R² are not both hydrogen;

R³ is hydrogen or methyl; and

R⁴ is phenyl or pyridine substituted by one or two substituents independently selected from the group consisting of:

- (1) halogen;
- (2) lower alkyl optionally substituted by halogen,
- (3) lower alkanoyl;
- (4) lower alkoxy;
- (5) lower alkylsulfanyl, lower alkylsulfinyl, or lower alkylsulfonyl
- (6) lower cycloalkylsulfanyl, lower cycloalkylsulfinyl, or lower cycloalkylsulfonyl; and

- 92 -

(7) phenyl or pyridine, wherein said phenyl or pyridine is optionally substituted by lower alkyl, lower alkylsulfanyl, lower alkylsulfonyl, lower cycloalkylsulfanyl, or lower cycloalkylsulfonyl.

- 5 2. A compound of claim 1 wherein R³ is hydrogen.
3. A compound according to claim 1 or 2 wherein R³ is methyl.
4. A compound according to any one of claims 1 to 3 wherein R⁴ is phenyl
- 10 substituted by two substituents independently selected from the group consisting of:
(1) halogen, and (2) lower alkyl optionally substituted by halogen.
5. A compound according to any one of claims 1 to 3 wherein R⁴ is phenyl substituted
by two substituents independently selected from the group consisting of:
- 15 (1) fluoro;
 (2) chloro;
 (3) bromo;
 (4) lower alkyl
 (5) lower alkanoyl;
- 20 (6) lower alkylsulfanyl, lower alkylsulfinyl, or lower alkylsulfonyl;
 (7) lower cycloalkylsulfanyl, lower cycloalkylsulfinyl, or lower
 cycloalkylsulfonyl;
 (8) trifluoromethyl, difluoromethyl, or fluoromethyl; and
 (9) 1,1-difluoroethyl.
- 25 6. A compound according to any one of claims 1 to 5 wherein R⁴ is phenyl
substituted by two substituents independently selected from the group consisting of:
- (1) fluoro;
 (2) chloro;
- 30 (3) bromo;
 (4) methyl;
 (5) ethyl;
 (6) propyl or isopropyl;

- 93 -

- (7) butyl, *sec*-butyl, or *tert*-butyl;
- (8) trifluoromethyl, difluoromethyl, or fluoromethyl; and
- (9) 1,1-difluoroethyl.

- 5 7. A compound according to any one of claims 1 to 6 wherein R⁴ is phenyl substituted by two substituents independently selected from the group consisting of:
- (1) fluoro;
 - (2) chloro;
 - (3) bromo; and
 - 10 (4) trifluoromethyl.

8. A compound according to any one of claims 1 to 3 wherein R⁴ is pyridine substituted by one substituent selected from the group consisting of:
- 15 (1) halogen;
- (2) lower alkyl optionally substituted by halogen; and
- (3) phenyl or pyridine, wherein said phenyl or pyridine is optionally substituted by lower alkyl.

- 20 9. A compound according to any one of claims 1 to 3 wherein R⁴ is pyridine substituted by phenyl wherein said phenyl is optionally substituted by lower alkyl.

10. A compound according to any one of claims 1 to 9 wherein R¹ is hydrogen, fluoro, chloro, bromo, methoxy, or phenyl and R² is hydrogen, fluoro, chloro, methyl,
- 25 ethyl, propyl, or isopropyl; with the proviso that R¹ and R² are not both hydrogen.

11. A compound according to any one of claims 1 to 10 wherein R¹ is hydrogen, fluoro, chloro, or bromo and R² is hydrogen, fluoro, chloro, or methyl, ethyl, or isopropyl; with the proviso that R¹ and R² are not both hydrogen.


- 30 12. A compound according to any one of claims 1 to 11 wherein at least one of R¹ or R² is fluoro, chloro, or bromo.

- 94 -

13. A compound according to any one of claims 1 to 12 wherein R¹ is fluoro or chloro.

14. A compound of claim 1 wherein R⁴ is phenyl substituted by two substituents at positions 3 and 5 on the phenyl ring, where such positions are indicated below in formula IC, independently selected from the group consisting of (1) halogen, (2) lower alkyl optionally substituted by halogen, (3) lower alkanoyl; (4) lower alkoxy; (5) lower alkylsulfanyl, lower alkylsulfinyl, or lower alkylsulfonyl and (6) lower cycloalkylsulfanyl, lower cycloalkylsulfinyl, or lower cycloalkylsulfonyl:

10




wherein R¹- R³ are as defined in claim 1.

15 15. A compound of claim 14 wherein R¹ is fluoro or chloro; R² is hydrogen or fluoro; R³ is hydrogen; and the phenyl ring is substituted at positions 3 and 5 as indicated in formula IC with substituents independently selected from the group consisting of fluoro, bromo, or trifluoromethyl.

20 16. A compound of claim 14 wherein R¹ is fluoro or chloro; R² is hydrogen or fluoro; R³ is hydrogen; and the phenyl ring is substituted at positions 3 and 5 with trifluoromethyl.

17. A compound of claim 1 which is an (*R*)-enantiomer as depicted in formula IA:



5 wherein R¹- R⁴ are as defined in claim 1.

18. A compound of claim 1 selected from the group consisting of:

- 10 [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-isopropyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 15 [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-ethyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 20 [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;

- 96 -

- 5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
[3-Chloro-5-(2,5-dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
5 [3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
[3-Chloro-5-(2,5-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
10 [3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
[5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
15 [3-Chloro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-Tetrahydro-naphthalen-1-yloxy]-acetic acid;
[5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
[(*R*)-3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
20 [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
[(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
25 [3-Chloro-2-fluoro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid; and
[5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.

- 30 19. A compound of claim 1 selected from the group consisting of:

[(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-methoxy-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;

- 97 -

- [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-phenyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 5 [2,3-Difluoro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [3-Fluoro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 10 [5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(2,4-Dichloro-benzenesulfonylamino)-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 15 [5-(2,4-Dichloro-benzenesulfonylamino)-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(2,5-Dichloro-benzenesulfonylamino)-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 20 [5-(2,5-Dichloro-benzenesulfonylamino)-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- {2,3-Difluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Fluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 25 {3-Chloro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- [(*R*)-3-Bromo-5-(3-bromo-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-3-Chloro-2-fluoro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 30 [(*R*)-5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;

- 98 -

- [(*R*)-3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-3-Chloro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 5 [(*R*)-5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-3-Chloro-5-(2,5-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- {3-Chloro-2-fluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 10 [5-(2,5-Dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [5-(2,4-Dichloro-benzenesulfonylamino)-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- 15 {2-Fluoro-5-[5-(3-isopropyl-phenyl)-pyridine-2-sulfonylamino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- [(*R*)-3-Bromo-5-(2,5-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- [(*R*)-3-Bromo-5-(2,4-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid; and
- 20 [5-(3,5-Bis-methanesulfonyl-benzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.

20. A compound of claim 1 selected from the group consisting of:

- 25 [5-(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid;
- {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 30 {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-ethyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;

- {(R)-5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Chloro-5-[(2,4-dichloro-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 5 {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-methyl-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Chloro-5-[(2,5-dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 10 {3-Chloro-5-[(2,5-dichloro-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Chloro-5-[(2,4-dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 15 {5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {3-Chloro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 20 {3-Chloro-2-fluoro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(2,4-Dichloro-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 25 {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(2,4-Dichloro-benzenesulfonyl)-methyl-amino]-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(2,5-Dichloro-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 30 {5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;

- 100 -

- {5-[(2,5-Dichloro-benzenesulfonyl)-methyl-amino]-2,3-difluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(3-Bromo-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- 5 {3-Fluoro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {2,3-Difluoro-5-[(3-fluoro-5-trifluoromethyl-benzenesulfonyl)-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- (2,3-Difluoro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid;
- 10 (3-Fluoro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid;
- (3-Chloro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid;
- 15 (3-Chloro-2-fluoro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid;
- {5-[(2,5-Dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid;
- {5-[(2,4-Dichloro-benzenesulfonyl)-methyl-amino]-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid; and
- 20 (2-Fluoro-5-[[5-(3-isopropyl-phenyl)-pyridine-2-sulfonyl]-methyl-amino]-5,6,7,8-tetrahydro-naphthalen-1-yloxy)-acetic acid.

21. A pharmaceutically acceptable salt of a compound according to any one of
25 claims 18 to 20.

22. A pharmaceutically acceptable ester of a compound according to any one of
claims 18 to 20.

30 23. A compound of claim 1 which is [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-2,3-dichloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.

- 101 -

24. A compound of claim 1 which is [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
25. A compound of claim 1 which is [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
26. A compound of claim 1 which is [5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
27. A compound of claim 1 which is [5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
28. A compound of claim 1 which is [3-Chloro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-Tetrahydro-naphthalen-1-yloxy]-acetic acid.
29. A compound of claim 1 which is [5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
30. A compound of claim 1 which is [(*R*)-3-Chloro-5-(2,4-dichloro-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
31. A compound of claim 1 which is [(*R*)-5-(3,5-Bis-trifluoromethyl-benzenesulfonylamino)-3-bromo-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
32. A compound of claim 1 which is [5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
33. A compound of claim 1 which is [(*R*)-3-Bromo-5-(3-bromo-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.

- 102 -

34. A compound of claim 1 which is [(*R*)-5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-2-fluoro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
- 5 35. A compound of claim 1 which is [(*R*)-3-Chloro-5-(3-fluoro-5-trifluoromethyl-benzenesulfonylamino)-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
36. A compound of claim 1 which is [(*R*)-5-(3-Bromo-5-trifluoromethyl-benzenesulfonylamino)-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy]-acetic acid.
- 10 37. A compound of claim 1 which is {5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid.
38. A compound of claim 1 which is {(*R*)-5-[(3,5-Bis-trifluoromethyl-benzenesulfonyl)-methyl-amino]-3-chloro-5,6,7,8-tetrahydro-naphthalen-1-yloxy}-acetic acid.
- 15 39. A pharmaceutical composition comprising a therapeutically effective amount of a compound according any one of claims 1 to 42 and a pharmaceutically acceptable carrier.
- 20 40. A compound according to any one of claims 1 to 38 for use as therapeutic active substances.
- 25 41. A compound according to any one of claims 1 to 38 for the treatment and/or prophylaxis of diseases treatable by CRTH2 receptors antagonist.
42. Use of a compound according to any one of claims 1 to 38 for the preparation of medicaments for the therapeutic and/or prophylactic treatment of diseases treatable
- 30 by a CRTH2 receptors antagonist.

- 103 -

43. The use according to claim 42, wherein the disease is asthma, chronic obstructive pulmonary disease, allergic inflammation, allergic rhinitis or atopic dermatitis.

- 5 44. The invention as hereinbefore defined, particularly with reference to the new compounds, intermediates, medicaments, uses and processes.