

INSECT REPELLENT

Field of the Invention

[0001]

The present invention relates to an insect repellent having insect repellent activity.

Background of the Invention

[0002]

Most hematophagous insects cause infections among humans and animals via transmission of a pathogen, or cause skin inflammation. In particular, mosquitoes carry serious diseases, such as Dengue fever, Zika fever, yellow fever, encephalitis, and malaria, to humans, and are thus hygienically very harmful insects.

[0003]

Such insects each have an excellent chemoreception system including a heat sensor to sense the body temperature of an animal, a taste receptor to sense taste, an olfactory receptor to sense volatile substances such as body odor, and a carbon dioxide receptor to sense carbon dioxide, which is a highly-volatile substance, and exhibit a wide variety of behaviors. For example, female mosquitoes before oviposition are known to trace carbon dioxide exhaled from an animal and the body odor to approach the animal for the blood, detect the targeted animal by sensing the body temperature with the heat sensor, and suck the blood.

[0004]

In recent years, a means to repel such insects has been devised which causes a change to the chemoreception system of the insect to disable the cognitive sense of the insect, and substances acting on the olfactory receptor, such as N,N-diethyl-3-methylbenzamide (DEET) and p-menthane-3,8-diol (PMD), have been used as a repellent.

However, DEET is disadvantageous in that it has uncomfortable odor, and use of it for infants and individuals with sensitive skin is restricted because of the high skin permeability, and the duration time is limited to 2 to 3 hours.

[0005]

While natural essential oils such as lemon eucalyptus oil, lemongrass oil, orange oil, and cassia oil are reported, for example, to have insect-repelling effect (e.g., Patent Literature 1), however, the repelling effect is not necessarily satisfactory, and thus insufficient for practical use.

Patent Literature

[0006]

[Patent Literature 1] JP-A-2002-173407

Summary of the Invention

[0007]

The present invention relates to the following 1) to 15).

1) An insect repellent comprising, as an active ingredient, at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde.

2) A thermoreception inhibitor for an insect, comprising, as an active ingredient, at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde.

3) An insect repellent comprising, as active ingredients, the compound defined in 1) and a fragrance compound having carbon dioxide response-inhibiting action.

4) A method for repelling an insect, the method comprising applying at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde onto the skin of a mammal.

5) A method for repelling an insect, the method comprising applying the compound defined in 4) and a fragrance compound having carbon dioxide response-inhibiting action onto the skin of a mammal.

6) A method for repelling an insect, the method comprising allowing at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-

inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde to attach or be held on an apparel article directly or indirectly contacting a skin.

7) A method for repelling an insect, the method comprising allowing the compound defined in 6) and a fragrance compound having carbon dioxide response-inhibiting action to attach or be held on an apparel article directly or indirectly contacting a skin.

8) A method for repelling an insect, the method comprising spreading at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde in a defined space.

9) A method for repelling an insect, the method comprising spreading the compound defined in 8) and a fragrance compound having carbon dioxide response-inhibiting action in a defined space.

10) Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for producing an insect repellent.

11) Use of the compound defined in 10) and a fragrance compound having carbon dioxide response-inhibiting action for producing an insect repellent.

12) Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for producing a thermoreception inhibitor for an insect.

13) Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for repelling an insect.

14) Use of the compound defined in 13) and a fragrance compound having carbon dioxide response-inhibiting action for repelling an insect.

15) Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for inhibiting the thermoreception of an insect.

Detailed Description of the Invention

[0008]

The present invention relates to providing an insect repellent having excellent repellent activity and being highly safe.

[0009]

The present inventors searched for a material capable of inhibiting the thermal response of an insect, and found that specific fragrance compounds have excellent thermoreception-inhibiting effect and are useful for an insect repellent, and further found that use of the compound in combination with a fragrance compound having carbon dioxide response-inhibiting action can inhibit the seeking behavior of an insect for humans, and is thus more useful for an insect repellent.

[0010]

The present invention can provide an insect repellent which exhibits excellent repelling effect by inhibition of the thermoreception response of an insect on the basis of a mechanism different from those of common repellents such as DEET and PMD, and is highly safe.

[0011]

Each of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde,

γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde in the present invention (hereinafter, also referred to as "the compound according to the present invention") is a compound known as a fragrance, and commercially available as follows.

- 3-methyl-5-propyl-2-cyclohexen-1-one (GRAVENONE): Symrise AG

- octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde ("SCENTENAL" (registered trademark)): Firmenich SA

- phenoxyacetaldehyde: International Flavors & Fragrances (IFF) Inc.

- γ -octalactone: Sigma-Aldrich Co., LLC.

- 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one ("FLOREX" (registered trademark)): Firmenich SA

- octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one (PLICATONE): Firmenich SA

- 2,3,3-trimethyl-2H-inden-1-one ("SAFRALEINE" (registered trademark)): Givaudan SA

- 4-methyl-5-thiazoleethanol: Sigma-Aldrich Co., LLC.

- 7-methoxy-3,7-dimethyloctanal: Innospec

- 3-methyl-1-phenyl-3-pentanol: Tokyo Chemical Industry Co., Ltd.

· 2-phenylpropionaldehyde dimethyl acetal: Sigma-Aldrich Co., LLC.

· octahydro-2H-1-benzopyran-2-one (octahydrocoumarin): Takasago International Corporation, International Flavors & Fragrances (IFF) Inc.

· 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde ("safranal"): Givaudan SA

· 6-methylquinoline: Tokyo Chemical Industry Co., Ltd.

One of these compounds according to the present invention may be used singly, or two or more thereof may be used in a mixture.

From the viewpoint of insect-repelling effect and feeling of use, for example, preferred compounds among the compounds according to the present invention are octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one (PLICATONE): Firmenich SA, 3-methyl-5-propyl-2-cyclohexen-1-one (GRAVENONE), and 6-methylquinoline.

[0012]

The compound according to the present invention may be used in combination with a fragrance compound having carbon dioxide response-inhibiting action.

Here, "carbon dioxide response-inhibiting" refers to inhibiting the seeking behavior of an insect for a carbon dioxide source by disabling or reducing the response of the insect to carbon dioxide reception to control the cognitive sense to carbon dioxide. The carbon dioxide response of an insect can be measured through

investigation of a nerve firing pattern of the sensory hair with electrodes inserted therein in response to a carbon dioxide stimulus by using known single sensillum recording (Nature 461, 277-281 (10 September 2009)).

Examples of fragrance compounds having carbon dioxide response-inhibiting action include compounds listed in Reference Example below, specifically, 2-methoxy-4-propylphenol, 2-isopropyl-4-methylthiazol, isocyclocitral, 9-decen-1-ol, 1-octen-3-ol, 1-decanol, ethyl 2-ethylhexanoate, trans-2-hexenyl acetate, ethyl 2-cyclohexylpropionate, ethyl phenylacetate, 2,4,6-trimethyl-3-cyclohexene-1-methanol, 1-octanol, 2,4-dimethylcyclohex-3-ene-1-methanol, o-tert-butylcyclohexyl acetate, 2-methylbutyric acid, benzyl propionate, 3-methyl-2-pentylcyclopent-2-en-1-one, and cis-3-hexenyl acetate. Among them, 2,4-dimethylcyclohex-3-ene-1-methanol (floralol), 1-octanol, cis-3-hexenyl acetate, etc., are preferred, and 2,4-dimethylcyclohex-3-ene-1-methanol is more preferred.

[0013]

The insect intended in the present invention is not limited and may be any insect having a thermoreception response system, and the insect may be hematophagous or non-hematophagous. Example of such insects include mosquitoes such as northern house mosquitoes, *Culex tritaeniorhynchus*, yellow fever mosquitoes, Asian tiger mosquitoes, and Chinese malaria mosquitoes; flies such as houseflies, false stable flies, little house flies, blow

flies, flesh flies, seed-corn flies, onion flies, fruit flies, vinegar flies, moth flies, and tsetse flies; horseflies; black flies; stable flies; biting midges; and chironomids. Among them, mosquitos, flies, black flies, and stable flies are preferred.

[0014]

As demonstrated in Examples later, the compound according to the present invention inhibits the thermoreception system of an Asian tiger mosquito. Thus, the compound according to the present invention can serve as an insect repellent or serve for inhibition of the thermoreception of an insect, and can be used for producing an insect repellent or thermoreception inhibitor. In other words, the compound according to the present invention can be used for repelling an insect and inhibiting the thermoreception. In addition, use of the compound according to the present invention in combination with a fragrance compound having carbon dioxide response-inhibiting action effectively inhibits the seeking behavior of an Asian tiger mosquito for humans, and thus such combined use provides a more excellent insect repellent and is useful for repelling an insect, especially, repelling a mosquito.

Here, "repelling an insect" refers to preventing an insect from approaching to an object or forcing an insect to avoid an object, and "inhibiting the thermoreception of an insect" refers to inhibiting the seeking behavior of an insect for a heat source by disabling or reducing

the thermoreception response of the insect to inhibit the cognitive sense to the heat source, and the concept differs from having insecticidal ability to exterminate an insect.

For determination of thermoreception, insects are released into a cage with which a heat source such as a plate heater is provided and the number of insects attracted to the heat source is counted.

[0015]

The insect repellent or thermoreception inhibitor according to the present invention can repel an insect, for example, through applying it onto the skin of a mammal such as a human, allowing it to attach or be held on an apparel article directly or indirectly contacting the skin, such as a cloth and an accessory, or spreading (including evaporating and volatilizing) it in a defined space such as a room interior and a car interior, and can be appropriately blended with additional components such as a solvent and an additive for formulation into a composition to prepare a formulation. Alternatively, the insect repellent or thermoreception inhibitor according to the present invention may be used for a material for imparting insect repellent activity to an external preparation for cutaneous administration, a cleaning agent, a skin cosmetic, a hair cosmetic, a textile softener, a fabric treatment agent, or the like.

[0016]

The amount of the compound according to the present invention to be blended in the above composition is preferably 0.001% by mass or more, more preferably 0.01% by mass or more, even more preferably 0.1% by mass or more and preferably 80% by mass or less, more preferably 40% by mass or less, even more preferably 10% by mass or less, or preferably from 0.001 to 80% by mass, more preferably 0.01 to 40% by mass, even more preferably from 0.01 to 10% by mass, and further more preferably from 0.1 to 10% by mass, relative to the total amount of the composition, although it varies depending on the dosage form.

[0017]

In the case that the compound according to the present invention is used in combination with a fragrance compound having carbon dioxide response-inhibiting action, the content of the fragrance compound in the composition is preferably 0.001% by mass or more, more preferably 0.01% by mass or more, even more preferably 0.1% by mass or more and preferably 80% by mass or less, more preferably 40% by mass or less, even more preferably 10 mass% or less, or preferably from 0.001 to 80% by mass, more preferably from 0.01 to 40% by mass, even more preferably from 0.01 to 10% by mass, and further more preferably from 0.1 to 10% by mass, relative to the total amount of the compound.

[0018]

The type of a solvent or additive may be appropriately selected in accordance with the mode of use or intended use of the insect repellent or thermoreception inhibitor, and examples of the solvent include lower alcohols such as methanol, ethanol, and propanol; and water.

Examples of the additive include additives commonly used for various chemical products and cosmetics, specifically, surfactants, organic solvents, oily components, moisturizing agents, powders, solubilizers, thickeners, resins, cleaning agents, preservatives, UV absorbers, inorganic substances, fragrances, pigments, deodorants, essential oils, pharmaceutical agents, plant extracts, and other repellents.

[0019]

The dosage form of the insect repellent or thermoreception inhibitor according to the present invention can be appropriately set in accordance with the mode of use, and may be any of a liquid, a cream, a lotion, an emulsion, a gel, an ointment, a powder, a granule, a spray, and an aerosol spray, etc.

[0020]

In the case that the insect repellent or thermoreception inhibitor is spread in a defined space, it is suitable to employ a method in which the insect repellent or thermoreception inhibitor supported on a carrier or dissolved in a solvent is packed in a container with a liquid-absorbing wick and heated or

evaporated through air blow, or a method in which the insect repellent or thermoreception inhibitor supported on a carrier is evaporated into the space at normal temperature. In this case, examples of carriers to support the insect repellent or thermoreception inhibitor include natural fibers such as pulp, cotton, wool, linen, and silk; synthetic fibers such as polypropylene, polyethylene, polyamide, polyethylene terephthalate, polybutylene terephthalate, polysulfone, rayon, methacrylic resin, and glass fiber; and porous materials such as zeolite, talc, white carbon, diatomaceous earth, lime, silica gel, and activated carbon.

[0021]

Examples of the method for allowing the insect repellent or thermoreception inhibitor according to the present invention to attach or be held on an apparel article include, in addition to a method in which the insect repellent or thermoreception inhibitor is applied onto an intended article, a method in which the insect repellent or thermoreception inhibitor is blended in a textile softener and allowed to attach or be held in treatment with the softener, and a method in which a fabric base material which has been treated with the insect repellent or thermoreception inhibitor in advance to impart insect repellent activity is used for production of an intended article.

[0022]

In relation to the above-described embodiments, the following modes are disclosed in the present invention.

<1> An insect repellent comprising, as an active ingredient, at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde.

<2> A thermoreception inhibitor for an insect, comprising, as an active ingredient, at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde.

<3> An insect repellent comprising, as active ingredients, the compound defined in <1> and a fragrance compound having carbon dioxide response-inhibiting action.

<4> The insect repellent according to <3>, wherein the fragrance compound having carbon dioxide response-inhibiting action is at least one selected from the group consisting of 2,4-dimethylcyclohex-3-ene-1-methanol, 1-octanol, and cis-3-hexenyl acetate.

<5> The insect repellent according to <3> or <4>, wherein the compound defined in <1> is octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one or 3-methyl-5-propyl-2-cyclohexen-1-one.

<6> The insect repellent according to <1>, <3>, <4>, or <5> or the thermoreception inhibitor for an insect according to <2>, wherein the insect is at least one selected from the group consisting of mosquitoes, flies, black flies, and stable flies.

<7> The insect repellent or thermoreception inhibitor for an insect according to <6>, wherein the insect is a mosquito.

<8> A method for repelling an insect, comprising applying at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-

inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde onto the skin of a mammal.

<9> A method for repelling an insect, comprising applying the compound defined in <8> and a fragrance compound having carbon dioxide response-inhibiting action onto the skin of a mammal.

<10> The method for repelling an insect according to <9>, wherein the fragrance compound having carbon dioxide response-inhibiting action is at least one selected from the group consisting of 2,4-dimethylcyclohex-3-ene-1-methanol, 1-octanol, and cis-3-hexenyl acetate.

<11> The method for repelling an insect according to <9> or <10>, wherein the compound defined in <8> is octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one or 3-methyl-5-propyl-2-cyclohexen-1-one.

[0023]

<12> Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-

phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for producing an insect repellent.

<13> Use of the compound defined in <12> and a fragrance compound having carbon dioxide response-inhibiting action for producing an insect repellent.

<14> Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for producing a thermoreception inhibitor for an insect.

<15> Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-

phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for repelling an insect.

<16> Use of the compound defined in <15> and a fragrance compound having carbon dioxide response-inhibiting action for repelling an insect.

<17> Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for inhibiting the thermoreception of an insect.

<18> The insect repellent or thermoreception inhibitor for an insect according to any one of <1> to <6>, wherein the content of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-

dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, or 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde in the composition is preferably 0.001% by mass or more, more preferably 0.01% by mass or more, even more preferably 0.1% by mass or more and preferably 80% by mass or less, more preferably 40% by mass or less, even more preferably 10% by mass or less, or preferably from 0.001 to 80% by mass, more preferably from 0.01 to 40% by mass, even more preferably from 0.01 to 10% by mass, and further more preferably from 0.1 to 10% by mass.

<19> The insect repellent according to any one of <3> to <5>, wherein the content of the fragrance compound having carbon dioxide response-inhibiting action in the composition is preferably 0.001% by mass or more, more preferably 0.01% by mass or more, even more preferably 0.1% by mass or more and preferably 80% by mass or less, more preferably 40% by mass or less, even more preferably 10% by mass or less, or preferably from 0.001 to 80% by mass, more preferably from 0.01 to 40% by mass, even more preferably from 0.01 to 10% by mass, and further more preferably from 0.1 to 10% by mass, relative to the total amount of the composition.

<20> A method for repelling an insect, comprising attaching or holding at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-

indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde on an apparel article directly or indirectly contacting the skin.

<21> A method for repelling an insect, comprising spreading at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde in a defined space.

<22> A method for repelling an insect, comprising attaching or holding the compound defined in <20> and a fragrance compound having carbon dioxide response-inhibiting action on an apparel article directly or indirectly contacting the skin.

<23> A method for repelling an insect, comprising spreading the compound defined in <21> and a fragrance compound having carbon dioxide response-inhibiting action in a defined space such as a room interior or a car interior.

<24> In <13>, <16>, <22>, or <23>, the fragrance compound is at least one selected from the group consisting of 2,4-dimethylcyclohex-3-ene-1-methanol, 1-octanol, and cis-3-hexenyl acetate.

<25> In <13>, <16>, <22>, <23>, or <24>, the compound defined in <12>, the compound defined in <15>, the compound defined in <20>, or the compound defined in <21> is octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one or 3-methyl-5-propyl-2-cyclohexen-1-one.

Examples

[0024]

Example 1: Evaluation of thermoreception-inhibiting effect

(1) Preparation of Asian tiger mosquitoes

Asian tiger mosquitoes grown from eggs purchased from Sumika Technoservice Corporation were used. A transparent plastic pan was filled with water to a depth of approximately 1 cm, and a filter paper on which the eggs had been deposited was put therein. The larvae were fed with tropical fish feed (TetraMin) every day. After approximately 1 week, the pupae were collected with a dropper, and transferred into a 20 mL plastic cup, and

the cup was placed in a netted cage. The adults were fed with 10% by mass sucrose contained in a 25 mL plastic tube. The males and females were kept together in one cage and allowed to mate for 5 days after eclosion. After 5 days of captivity, the adults were collected with an insect aspirator and anesthetized on ice for 5 minutes, and visually sorted into males and females to collect only the females.

[0025]

(2) Evaluation of inhibitory effect on seeking behavior for heat source

Approximately 150 to 200 female Asian tiger mosquitoes were transferred in a plastic cage (30 × 30 × 30 cm). To monitor the behavior of the female mosquitoes from the top of the plastic cage, the top surface of the cage was replaced with an acrylic plate. A plate heater (multi-panel heater 16W for small animals, manufactured by Vivaria) as a heat source was provided at a corner of the cage from the outside of the mesh with an area of 5 × 3 cm exposed. To take a photograph of the heat source from the top surface in experiment, a video camera (manufactured by Sony Corporation) was disposed above the top surface. A sheet for application of a sample was fixed to the heat source with a double-sided adhesive tape.

The concentration of a fragrance listed in Table 1 was adjusted to 0.1% with ethanol to prepare a sample, and 50 μ L of the sample or ethanol as a solvent was

administered in experiment. To activate the female mosquitoes, 1% of carbon dioxide was administered from the top for 1 second, and then the heat source was brought into contact with the cage. After 30 seconds of contact, a photograph of the heat source was taken to count the number of mosquitoes attracted. The sheet used for the heat source was discarded and replaced with another sheet every time the assay was completed. In view of the influence of a sample on the mosquitoes, the number of evaluations performed for one cage was limited to (7 samples + 1 control) at a maximum. The inhibitory rate relative to the number of landings with application of ethanol solvent only was calculated for one set of assays (7 samples + 1 control) to determine the inhibitory effect by using the following equation.

(Equation 1)

$$\text{Inhibitory rate (\%)} = 100 \times (1 - [\text{number of landings with application of sample}] / [\text{number of landings with application of ethanol}])$$

[0026]

Table 2 shows the results. The compounds according to the present invention are considered to inhibit the seeking behavior of an Asian tiger mosquito for a heat source to inhibit the thermoreception response.

[0027]

[Table 1]

Name	Distributor
3-Methyl-5-propyl-2-cyclohexen-1-one (GRAVENONE)	SYMRISE AG
Octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde ("SCENTENAL" (registered trademark))	FIRMENICH SA
Phenoxyacetaldehyde (50% benzyl alcohol solution)	IFF INC.
γ -Octalactone	SIGMA-ALDRICH CO., LLC.
9-Ethylidene-3-oxatricyclo(6.2.1.0 ^{2,7})undecan-4-one ("FLOREX" (registered trademark))	FIRMENICH SA
Octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one (PLICATONE)	FIRMENICH SA
2,3,3-Trimethyl-2H-inden-1-one ("SAFRALEINE" (registered trademark))	GIVAUDAN SA
4-Methyl-5-thiazoleethanol	SIGMA-ALDRICH CO., LLC.
7-Methoxy-3,7-dimethyloctanal	INNOSPEC
3-Methyl-1-phenyl-3-pentanol	Tokyo Chemical Industry Co., Ltd.
2-Phenylpropionaldehyde dimethyl acetal	SIGMA-ALDRICH CO., LLC.
Octahydro-2H-1-benzopyran-2-one (octahydrocoumarin)	Takasago International Corporation,
2,6,6-Trimethylcyclohexa-1,3-diene-1-carbaldehyde ("safranal")	GIVAUDAN SA
6-Methylquinoline	Tokyo Chemical Industry Co., Ltd.

[0028]

[Table 2]

Sample	Inhibitory rate for seeking for heat source (%)
3-Methyl-5-propyl-2-cyclohexen-1-one	97.2
Octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde	89.2
Phenoxyacetaldehyde	89.0
γ -Octalactone	87.4
9-Ethylidene-3-oxatricyclo(6.2.1.0 ^{2,7})undecan-4-one	85.9
Octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one	81.3
2,3,3-Trimethyl-2H-inden-1-one	76.2
4-Methyl-5-thiazoleethanol	74.5
7-Methoxy-3,7-dimethyloctanal	72.4
3-Methyl-1-phenyl-3-pentanol	69.3
2-Phenylpropionaldehyde dimethyl acetal	66.8
Octahydro-2H-1-benzopyran-2-one	93.3%
2,6,6-Trimethylcyclohexa-1,3-diene-1-carbaldehyde	91.7%
6-Methylquinoline	64.7%

[0029]

Example 2: Evaluation of inhibition of seeking behavior for human arm

Adult Asian tiger mosquitoes consisting of 220 females and 80 males each at 5 to 10 day old were purchased from Sumika Technoservice Corporation, and released in a 30 cm × 30 cm × 30 cm plastic cage (bug dorm).

The concentration of each of 2,4-dimethylcyclohex-3-ene-1-methanol ("floralol", IFF Inc.) and octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one ("PLICATONE", Firmenich SA) as evaluation samples was adjusted to 0.4%

with ethanol. A sample containing only ethanol was used as a control.

Each of the evaluation samples and control sample was sprayed 14 times in total (0.6 g) to both sides of a hand wearing a 100% cotton glove doubly.

After application of the sample, the gloved hand was left to stand for 1 minute to volatilize ethanol, and the gloved hand with the sample applied thereto was then put in the plastic cage, and the number of mosquitoes which landed on the glove was counted every 1 minute.

The numbers of landings after 1 minute, 2 minutes, and 3 minutes, and the sum total of them were counted, and the landing rate relative to the number of landings with application of the control sample as 100% was calculated. This experiment was repeated three times. Table 3 shows the results. It was demonstrated that use of the compound according to the present invention in combination with floralol (2,4-dimethylcyclohex-3-ene-1-methanol) more effectively inhibits the seeking behavior of a mosquito for the human arm.

[0030]

[Table 3]

Sample	Number of landings				Total landing rate (%)
	1 min	2 min	3 min	total	
Floralol	3	6	7	16	71.1
PLICATONE	2	4	6	12	53.3
Floralol + PLICATONE	2	1	3	6	26.7
Control (average of three runs)	6.5	6	10	22.5	100.0

[0031]

Reference Example: Evaluation of carbon dioxide response-inhibiting effect

(1) Preparation of Asian tiger mosquitoes

Asian tiger mosquitoes grown from eggs purchased from Sumika Technoservice Corporation were used. A transparent plastic pan was filled with water to a depth of approximately 1 cm, and a filter paper on which the eggs had been deposited was put therein. The larvae were fed with tropical fish feed (TetraMin) every day. After approximately 1 week, the pupae were collected with a dropper, and transferred into a 20 mL plastic cup, and the cup was placed in a netted cage. The adults were fed with 10 mass% sucrose contained in a 25 mL plastic tube. The males and females were kept together in one cage and allowed to mate for 5 days after eclosion. After 5 days of captivity, the adults were collected with an insect aspirator and anesthetized on ice for 5 minutes, and visually sorted into males and females to collect only the females. The wings and legs were removed from each female mosquito under anesthesia, and the female mosquito was pushed onto a double-sided adhesive tape pasted on a microscope slide with the back on the tape and thus fixed thereto, and the antennae and maxillary palpi were positioned so as to achieve the accessibility of electrodes.

[0032]

(2) Preparation of electrodes for recording nerve response

A 1 mol/L aqueous solution of potassium hydroxide was injected into a 50 mL syringe, and the syringe was horizontally fixed to a magnet stand with a clamp. A tungsten wire fixed to an electrode holder was horizontally inserted into the syringe, and polished by electrolysis under microscopic observation. A current was supplied from an AC power source to the syringe through the electrode holder via crocodile clips.

[0033]

(3) Recording of nerve response

The Asian tiger mosquito prepared on a microscope slide was set under a microscope, and positioned so that the antennae were at the center of the view through a $\times 10$ objective lens. A reference electrode was inserted in the eye of the insect. A pipette for providing an olfactory stimulus was placed near the maxillary palpi, and the objective lens was replaced with a $\times 100$ objective lens. The recording electrode was brought close to the sensory hair, and carefully inserted into the sensory hair with a micromanipulator. The peg-like shape of the carbon dioxide-sensitive sensory hair of an Asian tiger mosquito can be confirmed under microscopic observation. After insertion of the electrode into the sensory hair, the spontaneous nerve firing pattern was investigated. Carbon dioxide the concentration of which had been adjusted to 0.1% was fed to an olfactory stimulation

apparatus via a silicone tube to stimulate the maxillary palpi for 1 second, and the carbon dioxide response was investigated.

[0034]

(4) Evaluation of carbon dioxide response-inhibiting effect

The concentration of each of the fragrances listed in Table 4 was adjusted to 1% with paraffin oil, and the resultant was used as a sample. On a filter paper (3 mm × 50 mm) 15 µL of the sample was dropped, and the filter paper was inserted in a Pasteur pipette. The Pasteur pipette was connected to a stimulation apparatus via a plastic tube, and olfactory stimulation was performed for 1 second while the response to carbon dioxide in the air was monitored, and thus the carbon dioxide response-inhibiting effect was evaluated. The number of nerve firings in 1 second after stimulation with paraffin oil as a solvent and the number of nerve firings in 1 second after stimulation with each sample were counted, and the ratio was used for evaluation of the response-inhibiting effect (N = 1).

(Equation 2)

$$\text{Inhibitory rate (\%)} = 100 \times ([\text{number of firings after stimulation with paraffin oil} - \text{number of firings after stimulation with sample}] / [\text{number of firings after stimulation with paraffin oil}])$$

Table 5 shows the results.

[0035]

[Table 4]

Name	Distributor
2-Methoxy-4-propylphenol	SIGMA-ALDRICH CO., LLC.
2-Isopropyl-4-methylthiazol	SIGMA-ALDRICH CO., LLC.
Isocyclocitral	GIVAUDAN SA
9-Decen-1-ol	SIGMA-ALDRICH CO., LLC.
1-Octen-3-ol	SIGMA-ALDRICH CO., LLC.
1-Decanol	SIGMA-ALDRICH CO., LLC.
Ethyl 2-ethylhexanoate ("IROTYL" (registered trademark))	Kao Corporation
trans-2-Hexenyl acetate	SIGMA-ALDRICH CO., LLC.
Ethyl 2-cyclohexylpropionate ("POIRENATE" (registered trademark))	Kao Corporation
Ethyl phenylacetate	Tokyo Chemical Industry Co., Ltd.
2,4,6-Trimethyl-3-cyclohexene-1-methanol	IFF INC.
1-Octanol	Kao Corporation
2,4-Dimethylcyclohex-3-ene-1-methanol (floralol)	IFF INC.
o-tert-Butylcyclohexyl acetate	Kao Corporation
2-Methylbutyric acid	Tokyo Chemical Industry Co., Ltd.
Benzyl propionate	Tokyo Chemical Industry Co., Ltd.
cis-3-Hexenyl acetate	SIGMA-ALDRICH CO., LLC.

[0036]

[Table 5]

Sample	Inhibition of carbon dioxide response (%)
2-Methoxy-4-propylphenol	95
2-Isopropyl-4-methylthiazol	71
Isocyclocitral	61
9-Decen-1-ol	56
1-Octen-3-ol	55
1-Decanol	54
Ethyl 2-ethylhexanoate	51
trans-2-Hexenyl acetate	48
Ethyl 2-cyclohexylpropionate	48
Ethyl phenylacetate	79
2,4,6-Trimethyl-3-cyclohexene-1-methanol	61
1-Octanol	97
2,4-Dimethylcyclohex-3-ene-1-methanol	67
o-tert-Butylcyclohexyl acetate	87
2-Methylbutyric acid	75
Benzyl propionate	97
3-Methyl-2-pentylcyclopent-2-en-1-one	68
cis-3-Hexenyl acetate	50

Claims

[Claim 1]

An insect repellent comprising, as an active ingredient, at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde.

[Claim 2]

A thermoreception inhibitor for an insect, comprising, as an active ingredient, at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-

benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde.

[Claim 3]

An insect repellent comprising, as active ingredients, the compound defined in claim 1 and a fragrance compound having carbon dioxide response-inhibiting action.

[Claim 4]

The insect repellent according to claim 3, wherein the fragrance compound having carbon dioxide response-inhibiting action is at least one selected from the group consisting of 2,4-dimethylcyclohex-3-ene-1-methanol, 1-octanol, and cis-3-hexenyl acetate.

[Claim 5]

The insect repellent according to claim 3 or 4, wherein the compound defined in claim 1 is octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one or 3-methyl-5-propyl-2-cyclohexen-1-one.

[Claim 6]

The insect repellent according to claim 1, 3, 4, or 5 or the thermoreception inhibitor for an insect according to claim 2, wherein the insect is at least one selected from the group consisting of mosquitoes, flies, black flies, and stable flies.

[Claim 7]

The insect repellent or thermoreception inhibitor for an insect according to claim 6, wherein the insect is a mosquito.

[Claim 8]

A method for repelling an insect, comprising applying at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde onto the skin of a mammal.

[Claim 9]

A method for repelling an insect, comprising applying the compound defined in claim 8 and a fragrance compound having carbon dioxide response-inhibiting action onto the skin of a mammal.

[Claim 10]

The method for repelling an insect according to claim 9, wherein the fragrance compound having carbon dioxide response-inhibiting action is at least one selected from the group consisting of 2,4-dimethylcyclohex-3-ene-1-methanol, 1-octanol, and cis-3-hexenyl acetate.

[Claim 11]

The method for repelling an insect according to claim 9 or 10, wherein the compound defined in claim 8 is octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one or 3-methyl-5-propyl-2-cyclohexen-1-one.

[Claim 12]

A method for repelling an insect, comprising attaching or holding at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde on an apparel article directly or indirectly contacting the skin.

[Claim 13]

A method for repelling an insect, comprising attaching or holding the compound defined in claim 12 and a fragrance compound having carbon dioxide response-inhibiting action on an apparel article directly or indirectly contacting the skin.

[Claim 14]

The method for repelling an insect according to claim 13, wherein the fragrance compound having carbon

dioxide response-inhibiting action is at least one selected from the group consisting of 2,4-dimethylcyclohex-3-ene-1-methanol, 1-octanol, and cis-3-hexenyl acetate.

[Claim 15]

The method for repelling an insect according to claim 13 or 14, wherein the compound defined in claim 12 is octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one or 3-methyl-5-propyl-2-cyclohexen-1-one.

[Claim 16]

A method for repelling an insect, comprising spreading at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde in a defined space.

[Claim 17]

A method for repelling an insect, comprising spreading the compound defined in claim 16 and a fragrance compound having carbon dioxide response-inhibiting action in a defined space.

[Claim 18]

The method for repelling an insect according to claim 17, wherein the fragrance compound having carbon dioxide response-inhibiting action is at least one selected from the group consisting of 2,4-dimethylcyclohex-3-ene-1-methanol, 1-octanol, and cis-3-hexenyl acetate.

[Claim 19]

The method for repelling an insect according to claim 17 or 18, wherein the compound defined in claim 16 is octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one or 3-methyl-5-propyl-2-cyclohexen-1-one.

[Claim 20]

Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for producing an insect repellent.

[Claim 21]

Use of the compound defined in claim 20 and a fragrance compound having carbon dioxide response-inhibiting action for producing an insect repellent.

[Claim 22]

Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for producing a thermoreception inhibitor for an insect.

[Claim 23]

Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-

benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for repelling an insect.

[Claim 24]

Use of the compound defined in claim 23 and a fragrance compound having carbon dioxide response-inhibiting action for repelling an insect.

[Claim 25]

Use of at least one compound selected from the group consisting of octahydro-7-methyl-1,4-methanonaphthalen-6(2H)-one, 3-methyl-5-propyl-2-cyclohexen-1-one, octahydro-5-methoxy-4,7-methano-1H-indene-2-carboxaldehyde, 6-methylquinoline, phenoxyacetaldehyde, γ -octalactone, 9-ethylidene-3-oxatricyclo(6.2.1.0^{2,7})undecan-4-one, 2,3,3-trimethyl-2H-inden-1-one, 4-methyl-5-thiazoleethanol, 7-methoxy-3,7-dimethyloctanal, 3-methyl-1-phenyl-3-pentanol, 2-phenylpropionaldehyde dimethyl acetal, octahydro-2H-1-benzopyran-2-one, and 2,6,6-trimethylcyclohexa-1,3-diene-1-carbaldehyde for inhibiting thermoreception of an insect.