



(51) International Patent Classification:
C07D 217/26 (2006.01)

(21) International Application Number:
PCT/US2013/050538

(22) International Filing Date:
15 July 2013 (15.07.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/672,191 16 July 2012 (16.07.2012) US

(71) Applicant: **FIBROGEN, INC.** [US/US]; 409 Illinois St.,
San Francisco, California 94158 (US).

(72) Inventors: **THOMPSON, Michael D.**; c/o FibroGen, Inc.,
409 Illinois St., San Francisco, California 94158 (US).
PARK, Jung Min; c/o FibroGen, Inc., 409 Illinois St., San
Francisco, California 94158 (US). **AREND, Michael P.**;
c/o FibroGen, Inc., 409 Illinois St., San Francisco, Califor-
nia 94158 (US).

(74) Agents: **TANNER, Lorna L.** et al.; Sheppard Mullin
Richter & Hampton LLP, 379 Lytton Avenue, Palo Alto,
CA 94301-1479 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: PROCESS FOR MAKING ISOQUINOLINE COMPOUNDS

(57) Abstract: The present invention relates to methods for making isoquinoline compounds and the intermediate compounds achieved thereby. Such compounds can be used to prepare compounds and compositions capable of decreasing HIF hydroxylase enzyme activity, thereby increasing the stability and/or activity of hypoxia inducible factor (HIF).

