

Description**Background of the invention**

In the commercial production of chemiluminescent light devices which are used as emergency light sources or for aesthetic purposes etc, the prime concern is that the light be strong enough for a particularly desired duration. The usual criteria is that the duration of the light range from about 5 to about 12 hours without becoming so dim so as to lose its effectiveness.

At the other end of the spectrum is the need for light of a high intensity and a short duration. Usually the requirement is for an intensity of at least output 100 foot lamberts and a duration of from about 5 to 15 minutes. Light of this type is most commonly required for emergency or military uses such as, for example, flares. Explosive devices account for most of the production of these systems, however, because of the explosion, they, being pyroforic, sometimes cause undesirable side-effects such as brush fires, etc. It has therefore become apparent that there is a need for non-pyroforic devices which emit a strong light for a short duration and if such could be devised, a long-felt need would be satisfied.

Summary of the invention

A novel activator composition and a novel oxalate composition have now been found, which compositions, which admixed together, for a chemiluminescent light of high intensity and short duration. The activator composition comprises a mixture of two salicylate catalysts in critical concentrations whereas the oxalate composition comprises a critical admixture of an oxalate and a fluorescer, also in critical concentrations.

Description of the invention including preferred embodiments

The novel compositions of the present invention are useful in the production of chemical light devices of the kind comprising a liquid chemiluminescent mixture in a transparent or translucent container. They may be used in chemical lightsticks of the kind comprising an outer flexible sealed tube containing one of the novel compositions hereof separated by frangible means such as a glass vial or the like inside the tube and containing the other of the novel compositions hereof. By bending the tube, the frangible means is broken causing its composition to admix with the other composition in the tube and thereby form a chemiluminescent mixture which emits light through the walls of the tube. Typically, one of the compositions is an enhancer composition composed of a catalyst, hydrogen peroxide and a solvent while the other composition is in oxalate composition comprising an oxalate, a fluorescer and a solvent.

The instant invention resides in a novel activator composition and a novel oxalate composition which, when admixed, form a chemiluminescent light of high light intensity and short duration. By "high light intensity", as used herein, is meant a light intensity of at least about 100 foot lamberts and by "short duration", as used herein, is meant from about 5 to about 15 minutes.

The novel activator composition of the present invention which produces chemiluminescent light of high light intensity and short duration when admixed with the novel oxalate composition of the present invention, comprises a solvent solution of hydrogen peroxide and a mixture of from 0.01% to 0.08%, preferably from 0.03% to 0.06% by weight, based on the total weight of the activator composition, of sodium salicylate and from 0.01% to 0.08%, preferably from 0.03% to 0.06% by weight, same basis, of sodium 5-bromosalicylate, the ratio of sodium salicylate to sodium 5-bromosalicylate ranging from 9:1 to 1:9, respectively, preferably 7:3—3:7, respectively.

Useful solvents for the catalyst components of the activator composition are well known in the art and can comprise any of those set forth in any one or more of the following U.S. Patent Nos. 3749679; 3391068; 3391069; 3974368; 3816316; 3557233; 3597362; 3775336; 3888786, which patents are incorporated herein by reference.

Preferred solvents are esters, aromatic hydrocarbons and chlorinated hydrocarbons. Most preferred are the dialkyl phthalates wherein the alkyl group contains 1—12 carbon atoms, inclusive. Dimethyl phthalate is exemplary. Mixture of dimethyl phthalate and t-butyl alcohol are even more preferred.

The novel oxalate composition of the present invention which are adapted to produce chemiluminescent light of high light intensity and short duration when admixed with the novel activator compositions hereof comprise a solvent solution of a mixture of from 5% to 15%, by weight, based on the total weight of the oxalate composition, of an oxalate and from 0.16% to 0.36%, by weight, same basis, of a fluorescer, preferably from 0.20% to 0.32%.

Any of the above-disclosed solvents may also be used as the solvent for the oxalate solution. Dibutyl phthalate is preferred. Any oxalate known in the art may be used such as those disclosed in U.S. Patent Nos. 3749679 and 3816316. Substituted carbalkoxyphenyl oxalates are preferred and bis(2,4,5-trichloro-6-carbopentoxyphenyl)oxalate is most preferred.

The fluorescers useful in the oxalate

B¹) The oxalate composition of Example 1 contains:

88.625%—dibutyl phthalate

11.113%—bis(2,4,5-trichloro-6-carbopentoxyphenyl)oxalate (CPPO)

0.262%—1-chloro-9,10-bis(phenylethynyl)anthracene. (CBPEA)

A²) The activator composition of Example 2 contains:

Same as Example 1.

B²) The oxalate composition of Example 2 contains:

Same as Example 1, except that CBPEA replaced by rubrene.

A³) The activator composition of Example 3 contains:

82.04%—dimethyl phthalate

12.64%—t-butyl alcohol

5.24%—90% H₂O₂

0.04%—sodium salicylate

0.04%—sodium 5-bromosalicylate

B³) The oxalate composition of Example 3 contains:

91.14%—dibutyl phthalate

8.60%—CPPO

0.26%—rubrene

compositions of the present invention are well known in the art and any such fluorescer may be employed. Exemplary fluorscers include rubrene, i.e., 5,6,11,12-tetraphenylnaphthalene; 9,10-diphenylanthracene; 1-chloro-9,10-bis(phenylethynyl)anthracene; 9,10-bis(phenylethynyl)anthracene; 9,10-bis(4-methoxyphenyl)-2-chloroanthracene, alone or especially in admixture with N,N'-bis(2,5-di-t-butylphenyl)-3,4,9,10-perylenedicarboximide; mixtures of the above fluorscers, and the like.

The following examples are set forth for purposes of illustration only and are not to be construed as limitations on the present invention except as set forth in the appended claims. All parts and percentages are by weight unless otherwise specified.

Examples 1—5

In the following examples, all measurements are made using conventional six (6) inch lightsticks containing a glass ampule of the activator composition floating in the oxalate composition structure of which is known in the art.

In Table I, the light output (LOP) in foot lambert (approx. $10^3/\pi$ cd/cm²) units is given for compositions of the present invention (Examples 3, 4 and 5) in comparison to a standard red lightstick known in the art (Example 1) and a lightstick containing rubrene in the oxalate component and a standard activator solution (Example 2).

TABLE I

	Time (min.)	Example 1	Example 2	Example 3	Example 4	Example 5
5	On activation	176	305	4970	2765	1464
	1	18	38	291	266	234
10	2	15	29	237	254	226
	3	14	25	228	255	224
15	4	13	23	227	258	229
	5	13	21	232	260	224
	6	12	21	291	254	224
20	7	12	20	253	246	221
	8	12	19	248	226	214
	9	12	19	223	212	208
25	10	12	18	184	186	206
	11	—	18	137	154	196
30	12	—	18	88	126	188
	13	—	17	52	98	184
	14	—	17	28	72	181
35	15	—	16	14	52	172
	16	—	—	7	24	—

A¹⁾ The activator composition of Example 1 contains:

81.445%—dimethylphthalate

13.319%—*t*-butyl alcohol

5.227%—90% H₂O₂

0.009%—sodium salicylate

A⁴⁾ The activator compositions of Example 4 contains:

Same as Example 3.

B⁴⁾ The oxalate composition of Example 4 contains:

Same as Example 3 except the rubrene is replaced by a mixture of 10% rubrene and 90% CBPEA.

A⁵⁾ The activator composition of Example 5 contains:

Same as Example 3.

B⁵⁾ The oxalate composition of Example 5 contains:

Same as Example 3 except that rubrene replaced by CBPEA.

Example 6

When the procedure of Example 5 is again followed except that the CBPEA is replaced by 9,10-bis(phenylethynyl)anthracene (BPEA), similar results are achieved.

Example 7

Replacement of the CBPEA of Example 5 by diphenyl anthracene achieves similar results.

Example 8

The procedure of Example 3 is again followed except that the ratio of sodium salicylate to sodium 5-bromosalicylate is 9:1, respectively. Again a high light output of short duration is effected.

Example 9

Following the procedure of Example 8 except that the catalyst ratio is 1:9, respectively, equivalent results accrue.

5 Claims

1. An activator composition adapted to produce a chemiluminescent light of high light output for a short duration when admixed with an oxalate solution comprising a mixture of from 5% to 15%, by weight, based on the weight of the oxalate solution, of an oxalate and from 0.16% to 0.36%, by weight, same basis, of a fluorescer, comprising a solution of hydrogen peroxide and a mixture of from 0.01% to 0.08%, by weight, based on the total weight of said composition, of sodium salicylate and from 0.01% to 0.08%, by weight, same basis, of sodium 5-bromosalicylate, the ratio of sodium salicylate to sodium 5-bromosalicylate ranging from 9:1 to 1:9 respectively.
2. The composition of Claim 1 wherein the concentration of sodium salicylate ranges from 0.03% to 0.06%, by weight, and the concentration of sodium 5-bromosalicylate ranges from 0.03% to 0.06%, by weight.
3. An oxalate composition adapted to produce a chemiluminescent light of high light output for a short duration when admixed with an activator solution comprising hydrogen peroxide and a mixture of from 0.01% to 0.08%, by weight, based on the weight of said solution, of sodium salicylate and from 0.01% to 0.08%, by weight same basis, of sodium 5-bromosalicylate, the ratio of sodium salicylate to sodium 5-bromosalicylate, ranging from 9:1 to 1:9, respectively, comprising a solution of a mixture of from 5% to 15%, by weight, based on the weight of said composition, of an oxalate and from 0.16% to 0.36%, by weight, same basis, of a fluorescer.
4. A composition according to Claim 3 wherein said fluorescer is rubrene.
5. A composition according to Claim 3 wherein said fluorescer is a mixture of rubrene and 1-chloro-9,10-bis(phenylethynyl)anthracene.
6. A composition according to Claim 3, wherein said fluorescer is 9,10-bis(phenylethynyl)anthracene.
7. A composition according to Claim 3, wherein said fluorescer is a mixture of diphenyl anthracene and a soluble perylene dye.
8. A composition according to Claim 3 wherein said oxalate is bis(2,4,5-trichloro-6-carbopentoxyphenyl)oxalate.
9. A method for the production of chemiluminescent light of high light output and short duration which comprises adding to an oxalate solution comprising a mixture of from 5% to 15%, by weight, based on the weight of the oxalate solution, of an oxalate and from 0.16% to 0.36%, by weight, same basis, of a fluorescer, an activator composition comprising a solution of hydrogen peroxide and a mixture of from 0.01% to 0.08%, by weight, based on the total weight of the composition, of sodium salicylate and from 0.01% to 0.08%, by weight, same basis, of sodium 5-bromosalicylate, the ratio of sodium salicylate to sodium 5-bromosalicylate ranging from 9:1 to 1:9, respectively.
10. A method according to Claim 9 wherein the concentration of sodium salicylate ranges from 0.03% to 0.06%, by weight, and the concentration of sodium 5-bromosalicylate ranges from 0.03% to 0.06%, by weight.
11. A method according to Claim 10 wherein said fluorescer is rubrene.
12. A method according to Claim 10, wherein said fluorescer is a mixture of rubrene and 1-chloro-9,10-bis(phenylethynyl)anthracene.
13. A method according to Claim 10 wherein said fluorescer is 9,10-bis(phenylethynyl)anthracene.
14. A method according to Claim 10 wherein said fluorescer is a mixture of 9,10-diphenylanthracene and a soluble perylene dye.
15. A method according to Claim 10, wherein said oxalate is bis(2,4,5-trichloro-6-carbopentoxyphenyl)oxalate.

50 Patentansprüche

1. Aktivatorzusammensetzung für die Erzeugung eines Chemilumineszenz-Lichts mit hoher Lichtausgangsleistung während einer kurzen Dauer, wenn sie mit einer Oxalatlösung vermischt wird, die ein Gemisch von 5 bis 15 Gew.-%, bezogen auf das Gewicht der Oxalatlösung, eines Oxalats und von 0,16 bis 0,36 Gew.-%, bezogen auf die gleiche Basis, eines Leuchtstoffs umfaßt, umfassend eine Lösung von Wasserstoffperoxid und ein Gemisch von 0,01 bis 0,08 Gew.-%, bezogen auf das Gesamtgewicht der Zusammensetzung, Natriumsalicylat und von 0,01 bis 0,08 Gew.-%, bezogen auf die gleiche Basis, Natrium-5-bromosalicylat, wobei das Verhältnis von Natriumsalicylat zu Natrium-5-bromosalicylat im Bereich von 9:1 bis 1:9 liegt.
2. Zusammensetzung gemäß Anspruch 1, wobei die Konzentration von Natriumsalicylat im Bereich von 0,03 bis 0,06 Gew.-% liegt und die Konzentration von Natrium-5-bromosalicylat im Bereich von 0,03 bis 0,06 Gew.-% liegt.
3. Oxalatzusammensetzung für die Erzeugung eines Chemilumineszenz-Lichts mit hoher Lichtausgangsleistung während einer kurzen Dauer, wenn sie mit einer Aktivatorlösung vermischt wird, die



Example 9

Following the procedure of Example 8 except that the catalyst ratio is 1:9, respectively, equivalent results accrue.

5 Claims

1. An activator composition adapted to produce a chemiluminescent light of high light output for a short duration when admixed with an oxalate solution comprising a mixture of from 5% to 15%, by weight, based on the weight of the oxalate solution, of an oxalate and from 0.16% to 0.36%, by weight, same basis, of a fluorescer, comprising a solution of hydrogen peroxide and a mixture of from 0.01% to 0.08%, by weight, based on the total weight of said composition, of sodium salicylate and from 0.01% to 0.08%, by weight, same basis, of sodium 5-bromosalicylate, the ratio of sodium salicylate to sodium 5-bromosalicylate ranging from 9:1 to 1:9 respectively.

2. The composition of Claim 1 wherein the concentration of sodium salicylate ranges from 0.03% to 0.06%, by weight, and the concentration of sodium 5-bromosalicylate ranges from 0.03% to 0.06%, by weight.

3. An oxalate composition adapted to produce a chemiluminescent light of high light output for a short duration when admixed with an activator solution comprising hydrogen peroxide and a mixture of from 0.01% to 0.08%, by weight, based on the weight of said solution, of sodium salicylate and from 0.01% to 0.08%, by weight same basis, of sodium 5-bromosalicylate, the ratio of sodium salicylate to sodium 5-bromosalicylate, ranging from 9:1 to 1:9, respectively, comprising a solution of a mixture of from 5% to 15%, by weight, based on the weight of said composition, of an oxalate and from 0.16% to 0.36%, by weight, same basis, of a fluorescer.

4. A composition according to Claim 3 wherein said fluorescer is rubrene.

5. A composition according to Claim 3 wherein said fluorescer is a mixture of rubrene and 1-chloro-9,10-bis(phenylethynyl)anthracene.

6. A composition according to Claim 3, wherein said fluorescer is 9,10-bis(phenylethynyl)anthracene.

7. A composition according to Claim 3, wherein said fluorescer is a mixture of diphenyl anthracene and a soluble perylene dye.

8. A composition according to Claim 3 wherein said oxalate is bis(2,4,5-trichloro-6-carbopentoxyphenyl)oxalate.

9. A method for the production of chemiluminescent light of high light output and short duration which comprises adding to an oxalate solution comprising a mixture of from 5% to 15%, by weight, based on the weight of the oxalate solution, of an oxalate and from 0.16% to 0.36%, by weight, same basis, of a fluorescer, an activator composition comprising a solution of hydrogen peroxide and a mixture of from 0.01% to 0.08%, by weight, based on the total weight of the composition, of sodium salicylate and from 0.01% to 0.08%, by weight, same basis, of sodium 5-bromosalicylate, the ratio of sodium salicylate to sodium 5-bromosalicylate ranging from 9:1 to 1:9, respectively.

10. A method according to Claim 9 wherein the concentration of sodium salicylate ranges from 0.03% to 0.06%, by weight, and the concentration of sodium 5-bromosalicylate ranges from 0.03% to 0.06%, by weight.

11. A method according to Claim 10 wherein said fluorescer is rubrene.

12. A method according to Claim 10, wherein said fluorescer is a mixture of rubrene and 1-chloro-9,10-bis(phenylethynyl)anthracene.

13. A method according to Claim 10 wherein said fluorescer is 9,10-bis(phenylethynyl)anthracene.

14. A method according to Claim 10 wherein said fluorescer is a mixture of 9,10-diphenylanthracene and a soluble perylene dye.

15. A method according to Claim 10, wherein said oxalate is bis(2,4,5-trichloro-6-carbopentoxyphenyl)oxalate.

Patentansprüche

1. Aktivatorzusammensetzung für die Erzeugung eines Chemilumineszenz-Lichts mit hoher Lichtausgangsleistung während einer kurzen Dauer, wenn sie mit einer Oxalatlösung vermischt wird, die ein Gemisch von 5 bis 15 Gew.-%, bezogen auf das Gewicht der Oxalatlösung, eines Oxalats und von 0,16 bis 0,36 Gew.-%, bezogen auf die gleiche Basis, eines Leuchtstoffs umfaßt, umfassend eine Lösung von Wasserstoffperoxid und ein Gemisch von 0,01 bis 0,08 Gew.-%, bezogen auf das Gesamtgewicht der Zusammensetzung, Natriumsalicylat und von 0,01 bis 0,08 Gew.-%, bezogen auf die gleiche Basis, Natrium-5-bromosalicylat, wobei das Verhältnis von Natriumsalicylat zu Natrium-5-bromosalicylat im Bereich von 9:1 bis 1:9 liegt.

2. Zusammensetzung gemäß Anspruch 1, wobei die Konzentration von Natriumsalicylat im Bereich von 0,03 bis 0,06 Gew.-% liegt und die Konzentration von Natrium-5-bromosalicylat im Bereich von 0,03 bis 0,06 Gew.-% liegt.

3. Oxalatzusammensetzung für die Erzeugung eines Chemilumineszenz-Lichts mit hoher Lichtausgangsleistung während einer kurzen Dauer, wenn sie mit einer Aktivatorlösung vermischt wird, die



Wasserstoffperoxid und eine Mischung von 0,01 bis 0,08 Gew.-%, bezogen auf das Gewicht der Lösung, Natriumsalicylat und von 0,01 bis 0,08 Gew.-%, bezogen auf die gleiche Basis, Natrium-5-bromsalicylat umfaßt, wobei das Verhältnis von Natriumsalicylat zu Natrium-5-bromsalicylat im Bereich von 9:1 bis 1:9 liegt, umfassend eine Lösung von einem Gemisch von 5 bis 15 Gew.-%, bezogen auf das Gewicht der Zusammensetzung eines Oxalats, und von 0,16 bis 0,36 Gew.-%, bezogen auf die gleiche Basis, eines Leuchtstoffs.

4. Zusammensetzung gemäß Anspruch 3, wobei der Leuchstoff Rubren ist.

5. Zusammensetzung gemäß Anspruch 3, wobei der Leuchstoff ein Gemisch von Rubren und 1-Chlor-9,10-bis(phenylethynyl)anthracen ist.

6. Zusammensetzung gemäß Anspruch 3, wobei der Leuchstoff 9,10-Bis-(phenylethynyl)-anthracen ist.

7. Zusammensetzung gemäß Anspruch 3, wobei der Leuchstoff ein Gemisch von Diphenylanthracen und einem löslichen Perylen-Farbstoff ist.

8. Zusammensetzung gemäß Anspruch 3, wobei das Oxalat Bis-(2,4,5-trichlor-6-carbopentoxyphenyl)-oxalat ist.

9. Verfahren zur Erzeugung von Chemilumineszenz-Licht mit hoher Lichtausgangsleistung und kurzer Dauer, wobei man zu einer Oxalatlösung, umfassend ein Gemisch von 5 bis 15 Gew.-%, bezogen auf das Gewicht der Oxalatlösung, eines Oxalats und von 0,16 bis 0,36 Gew.-%, bezogen auf die gleiche Basis, eines Leuchtstoffs, eine Aktivatorzusammensetzung gibt, welche eine Lösung von Wasserstoffperoxid und ein Gemisch von 0,01 bis 0,08 Gew.-%, bezogen auf das Gesamtgewicht der Zusammensetzung, Natriumsalicylat und von 0,01 bis 0,08 Gew.-%, bezogen auf die gleiche Basis, Natrium-5-bromsalicylat umfaßt, wobei das Verhältnis von Natriumsalicylat zu Natrium-5-bromsalicylat im Bereich von 9:1 bis 1:9 liegt.

10. Verfahren gemäß Anspruch 9, wobei die Konzentration von Natriumsalicylat im Bereich von 0,03 bis 0,06 Gew.-% liegt und die Konzentration von Natrium-5-bromsalicylat im Bereich von 0,03 bis 0,06 Gew.-% liegt.

11. Verfahren gemäß Anspruch 10, wobei der Leuchstoff Rubren ist.

12. Verfahren gemäß Anspruch 10, wobei der Leuchstoff ein Gemisch von Rubren und 1-Chlor-9,10-bis-(phenylethynyl)-anthracen ist.

13. Verfahren gemäß Anspruch 10, wobei der Leuchstoff 9,10-Bis-(phenylethynyl)-anthracen ist.

14. Verfahren gemäß Anspruch 10, wobei der Leuchstoff ein Gemisch von 9,10-Diphenylanthracen und einem löslichen Perylen-Farbstoff ist.

15. Verfahren gemäß Anspruch 10, wobei das Oxalat Bis-(2,4,5-trichlor-6-carbopentoxyphenyl)-oxalat ist.

Revendications

1. Une composition activatrice adaptée à produire une lumière chimioluminescente de forte intensité lumineuse pendant une brève durée lorsqu'elle est mélangée à une solution à base d'oxalate comprenant un mélange de 5 à 15% en poids, par rapport au poids de la solution à base d'oxalate, d'un oxalate et de 0,16 à 0,36% en poids, sur la même base, d'un agent de fluorescence, comprenant une solution de peroxyde d'hydrogène et un mélange de 0,01 à 0,08% en poids, par rapport au poids total de ladite composition, de salicylate de sodium et de 0,01 à 0,08% en poids, sur la même base, de 5-bromosalicylate de sodium le rapport du salicylate de sodium au 5-bromosalicylate de sodium se situant de 9:1 à 1:9, respectivement.

2. La composition de la revendication 1, dans laquelle la concentration de salicylate de sodium est de 0,03 à 0,06% en poids, et la concentration de 5-bromosalicylate de sodium est de 0,03 à 0,06% en poids.

3. Une composition à base d'oxalate adaptée à produire une lumière chimioluminescente de forte intensité lumineuse pendant une brève durée lorsqu'elle est mélangée à une solution activatrice comprenant du peroxyde d'hydrogène et un mélange de 0,01 à 0,08% en poids, par rapport au poids de ladite solution, de salicylate de sodium et de 0,01 à 0,08% en poids, sur la même base, de 5-bromosalicylate de sodium, le rapport du salicylate de sodium au 5-bromosalicylate de sodium se situant de 9:1 à 1:9, respectivement, comprenant une solution d'un mélange de 5 à 15% en poids, par rapport au poids de ladite composition, d'un oxalate et de 0,16 à 0,36% en poids, sur la même base, d'un agent de fluorescence.

4. Une composition selon la revendication 3 dans laquelle ledit agent de fluorescence est le rubrène.

5. Une composition selon la revendication 3 dans laquelle ledit agent de fluorescence est un mélange de rubrène et de 1-chloro-9,10-bis(phenylethynyl)anthracène.

6. Une composition selon la revendication 3 dans laquelle ledit agent de fluorescence est le 9,10-bis(phenylethynyl)anthracène.

7. Une composition selon la revendication 3 dans laquelle ledit agent de fluorescence est un mélange de diphenyl anthracène et d'un colorant perylène soluble.

8. Une composition selon la revendication 3 dans laquelle ledit oxalate est l'oxalate de bis(2,4,5-trichloro-6-carbopentoxyphényle).

9. Un procédé pour la production de lumière chimioluminescente de forte intensité lumineuse et de brève durée qui comprend l'addition à une solution à base d'oxalate comprenant un mélange de 5 à 15% en poids, par rapport au poids de la solution à base d'oxalate, d'un oxalate et de 0,16 à 0,36% en poids, sur



la même base, d'un agent de fluorescence, d'une composition activatrice comprenant une solution de peroxyde d'hydrogène et un mélange de 0,01 à 0,08% en poids, par rapport au poids total de la composition, de salicylate de sodium et de 0,01 à 0,08% en poids, sur la même base, de 5-bromosalicylate de sodium, le rapport du salicylate de sodium au 5-bromosalicylate de sodium se situant de 9:1 à 1:9, respectivement.

10. Un procédé selon la revendication 9 dans lequel la concentration du salicylate de sodium se situe de 0,03 à 0,06% en poids, et la concentration du 5-bromosalicylate de sodium se situe de 0,03 à 0,06% en poids.

11. Un procédé selon la revendication 10 dans lequel ledit agent de fluorescence est le rubrène.

12. Un procédé selon la revendication 10 dans lequel ledit agent de fluorescence est un mélange de rubrène et de 1-chloro-9,10-bis(phényléthynyl)anthracène.

13. Un procédé selon la revendication 10 dans lequel ledit agent de fluorescence est le 9,10-bis(phényléthynyl)anthracène.

14. Un procédé selon la revendication 10 dans lequel ledit agent de fluorescence est un mélange de 9,10-diphénylanthracène et d'un colorant pérylène soluble.

15. Un procédé selon la revendication 10 dans lequel ledit oxalate est l'oxalate de bis(2,4,5-trichloro-6-carbopentoxyphényle).