


1/5

Figure 1: X-Ray Powder Diffraction Pattern of Form A of 8-[6-(3-dimethylaminopropoxy)pyridin-3-yl]-3-methyl-1-(oxan-4-yl)imidazo[5,4-c]quinolin-2-one


5



2/5


Figure 2: DSC Thermogram of Form A of 8-[6-(3-dimethylaminopropoxy)pyridin-3-yl]-3-methyl-1-(oxan-4-yl)imidazo[5,4-c]quinolin-2-one

5




3/5

Figure 3: Tumour Growth Inhibition in the Mouse Xenograft Model by 8-[6-(3-Dimethylaminopropoxy)pyridin-3-yl]-1-(*cis*-3-methoxycyclobutyl)-3-methylimidazo[4,5-
 5 c]quinolin-2-one (Example 2) in Combination with Irinotecan



4/5

Figure 4: Tumour Growth Inhibition in the Mouse Xenograft Model by
 Dimethylaminopropoxy)pyridin-3-yl]-3-methyl-1-(oxan-4-yl)imidazo[5,4-c]quinolin-2-
 one (Example 1) in Combination with Irinotecan



5/5

Figure 5: Tumour Growth Inhibition in the Mouse Xenograft Model by 8-[6-(3-Dimethylaminopropoxy)pyridin-3-yl]-3-methyl-1-(oxan-4-yl)imidazo[5,4-c]quinolin-2-one (Example 1) in Combination with Olaparib

