

KIF18A INHIBITORS

[0001] The invention relates to the field of pharmaceutical agents and, more specifically, is directed to compounds and compositions useful for modulating KIF18A, and to uses and methods for managing cell proliferation and for treating cancer.

BACKGROUND OF THE INVENTION

[0002] Cancer is one of the most widespread diseases afflicting mankind and a major cause of death worldwide. In an effort to find an effective treatment or a cure for one or more of the many different cancers, over the last couple of decades, numerous groups have invested a tremendous amount of time, effort and financial resources. However, to date, of the available cancer treatments and therapies, only a few offer any considerable degree of success.

[0003] Cancer is often characterized by unregulated cell proliferation. Damage to one or more genes, responsible for the cellular pathways, which control progress of proliferation through the cell cycle and centrosome cycle, can cause the loss of normal regulation of cell proliferation. These deregulated genes can code for various tumor suppressor or oncogene proteins, which participate in a cascade of events, leading to unchecked cell-cycling progression and cell proliferation. Various kinase and kinesin proteins have been identified, which play key roles in cell cycle and mitotic regulation and progression of normal dividing cells and cancer cells.

[0004] Kinesins are molecular motors that play important roles in cell division and intracellular vesicle and organelle transport. Mitotic kinesin plays roles in several aspects of spindle assembly, chromosome segregation, centrosome separation and dynamics (reviewed in O. Rath and F. Kozielski, *Nature Review Cancer*, 12:527-39, 2012). Human kinesins are categorized into 14 subfamilies based on sequence homology within the so-called “motor domain”, this domains ATPase activity drives unidirectional movement along microtubules (MTs). The non-motor domain of these proteins is responsible for cargo attachment; a “cargo” can include any one of a variety of different membranous organelles, signal transduction scaffolding systems, and chromosomes. Kinesins use the energy of ATP hydrolysis to move cargo along polarized microtubules. Thus, kinesins are often called “plus-end” or “minus-end” directed motors.

[0005] *KIF18A* gene belongs to Kinesin-8 subfamily and is a plus-end-directed motor. KIF18A is believed to influence dynamics at the plus end of kinetochore microtubules to control correct chromosome positioning and spindle tension. Depletion of human KIF18A leads to longer spindles, increased chromosome oscillation at metaphase, and activation of the mitotic spindle assembly checkpoint in HeLa cervical cancer cells (MI Mayr et al, *Current Biology* 17, 488–98, 2007). KIF18A appears to be viable target for the treatment of cancer. KIF18A is overexpressed in various types of cancers, including but not limited to colon, breast, lung, pancreas, prostate, bladder, head, neck, cervix, and ovarian cancers. Further, genetic deletion or knockdown, or inhibition of KIF18A effects mitotic spindle apparatus in cancer cell lines. Particularly, inhibition of KIF18A has been found to induce mitotic cell arrest, a known vulnerability that can promote cell death in mitosis via apoptosis, mitotic catastrophe, or multipolarity


driven lethality or death after mitotic slippage in interphase. Accordingly, there has been a strong interest in finding inhibitors of KIF18A proteins.

[0006] Thus, the inhibition of KIF18A ATPase activity is a promising approach for the development of novel anti-cancer agents.

SUMMARY

[0007] The present disclosure provides a new class of compounds useful for modulating KIF18A protein alone or in a bound complex with microtubules for treating KIF18A-mediated conditions and/or diseases, including cancer, inflammation, or ciliopathologies.

[0007a] In one aspect, the present disclosure refers to a compound of formula I:



or any pharmaceutically-acceptable salt thereof, wherein:

X¹ is N or -CR⁶;

R¹ is -CN, or a group -Z-R¹² wherein Z is -C₀₋₄alk-, -NR¹¹-, -NR¹¹SO₂-, -SO₂NR¹¹-, -NR¹¹-S(=O)(=NH)-, -S(=O)(=NH)-, -S-, -S(=O)-, -SO₂-, C₀₋₄alk-O-, -(C=O)-, -(C=O)NR¹¹-, -C=N(OH)-, or -NR¹¹(C=O); or

the group -Z-R¹² is -N=S(=O)-(R¹²)₂, wherein the two R¹² pair can alternatively combine with the sulfur atom attached to each of them to form a saturated or partially-saturated 3-, 4-, 5-, or 6-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S;

R² is halo or a group -Y-R¹³, wherein Y is -C₀₋₄alk-, -NH-C₀₋₄alkylene-, -N(CH₃)-C₀₋₄alkylene-, -C(=O)NR^a(C₁₋₄alk), -O-C₀₋₄alk-, S, S=O, S(=O)₂, -SO₂NR¹³, or -S(=O)(=NH)-;

R³ is H, C₁₋₄alk, or C₁₋₄haloalk;

R⁴ is H, halo, R^{4a} or R^{4b};


R⁵ is H, halo, C₁₋₈alk, or C₁₋₄haloalk;

R⁶ is H, halo, C₁₋₈alk, C₁₋₄haloalk, -O-C₁₋₈alk, or -O-R^{6a}; wherein R^{6a} is a saturated or partially-saturated 3-, 4-, 5-, or 6-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S;

R⁷ is H, halo, C₁₋₈alk, or C₁₋₄haloalk;

R⁸ is H, halo, C₁₋₈alk, C₁₋₄haloalk, -OH, -O-R^{8a}, or -O-R^{8b};

R⁹ is H, halo, C₁₋₈alk, or C₁₋₄haloalk;



R^x is represented by the following formula:

Each of R^{10a}, R^{10b}, R^{10c}, R^{10d}, R^{10e}, R^{10f}, R^{10g}, R^{10h}, R¹⁰ⁱ, and R^{10j} is H, halo, R^{10k}, or R^{10l},
or alternatively, each of R^{10a} and R^{10b} pair, R^{10c} and R^{10d} pair, R^{10e} and R^{10f} pair, R^{10g} and R^{10h} pair, or R¹⁰ⁱ and R^{10j} pair, independently, can combine with the carbon atom attached to each of them to form a saturated or partially-saturated 3-, 4-, 5-, 6-membered monocyclic ring spiro to the R^x ring; wherein said 3-, 4-, 5-, 6-membered monocyclic ring contains 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S, and further wherein said 3-, 4-, 5-, 6-membered monocyclic ring is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, C₁₋₆alk, C₁₋₄haloalk, -OR^a, -OC₁₋₄haloalk, CN, -NR^aR^a, or oxo;

R¹¹ is H, R^{11a}, or R^{11b};

R¹² is H, R^{12a}, or R^{12b};

R¹³ is R^{13a} or R^{13b};

R^{4a}, R^{8a}, R^{10k}, R^{11a}, R^{12a}, and R^{13a} is independently, at each instance, selected from the group consisting of a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic or 5-, 6-, 7-, 8-, 9-, 10-, 11-, or 12-membered bicyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S, which is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, C₁₋₆alk, C₁₋₄haloalk, -OR^a, -OC₁₋₄haloalk, CN, -C(=O)R^b, -C(=O)OR^a, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a, -NR^aC₂₋₆alkOR^a, -NR^aC₂₋₆alkO R^a,

-C₁₋₆alkNR^aR^a, -C₁₋₆alkOR^a, -C₁₋₆alkN(R^a)C(=O)R^b, -C₁₋₆alkOC(=O)R^b, -C₁₋₆alkC(=O)NR^aR^a, -C₁₋₆alkC(=O)OR^a, R¹⁴, and oxo;

R^{4b}, R^{8b}, R^{10l}, R^{11b}, R^{12b}, and R^{13b} is independently, at each instance, selected from the group consisting of C₁₋₆alk substituted by 0, 1, 2, 3, 4, or 5 group(s) selected from F, Cl, Br, -OR^a, -OC₁₋₄haloalk, or CN;

R¹⁴ is independently, at each instance, selected from the group consisting of a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic or 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, or 12-membered bicyclic ring containing 0, 1, 2 or 3 N atoms and 0 or 1 atoms selected from O and S, which is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, C₁₋₆alk, C₁₋₄haloalk, -OR^a,

-OC₁₋₄haloalk, CN, -C(=O)R^b, -C(=O)OR^a, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a, -NR^aC₂₋₆alkOR^a, -C₁₋₆alkNR^aR^a, -C₁₋₆alkOR^a, -C₁₋₆alkN(R^a)C(=O)R^b, -C₁₋₆alkOC(=O)R^b, -C₁₋₆alkC(=O)NR^aR^a, -C₁₋₆alkC(=O)OR^a, and oxo;

R^a is independently, at each instance, H or R^b; and

R^b is independently, at each instance, C₁₋₆alk, phenyl, or benzyl, wherein the C₁₋₆alk is being substituted by 0, 1, 2 or 3 substituents selected from halo, -OH, -OC₁₋₄alk, -NH₂, -NHC₁₋₄alk, -OC(=O)C₁₋₄alk, or -N(C₁₋₄alk)C₁₋₄alk; and the phenyl or benzyl is being substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄alk, C₁₋₃haloalk, -OH, -OC₁₋₄alk, -NH₂, -NHC₁₋₄alk, -OC(=O)C₁₋₄alk, or -N(C₁₋₄alk)C₁₋₄alk.

[0007b] In another aspect, the present disclosure refers to a pharmaceutical composition comprising the compound according to the disclosure, or the pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable diluent or carrier.

[0007c] In another aspect, the present disclosure refers to a compound in accordance with the disclosure, or the pharmaceutical composition according to the disclosure in the preparation of a medicament for treating a condition, wherein said condition is cancer selected from the group consisting of (a) a solid or hematologically derived tumor selected from cancer of the bladder, endometrial, lung squamous cell, breast, colon, kidney, liver, lung, small cell lung cancer, esophagus, gall-bladder, brain, head and neck, ovary, pancreas, stomach, cervix, thyroid, prostate and skin, (b) a hematopoietic tumor of lymphoid lineage selected from leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-

cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma and Burkett's lymphoma, (c) a hematopoietic tumor of myeloid lineage selected from acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia (d) a tumor of mesenchymal origin selected from fibrosarcoma and rhabdomyosarcoma, (e) a tumor of the central and peripheral nervous system selected from astrocytoma, neuroblastoma, glioma and schwannoma, and (f) a melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderma pigmentosum, keratocanthoma, thyroid follicular cancer or Kaposi's sarcoma.

[0007d] In another aspect, the present disclosure refers to a use of the compound in accordance with the disclosure, or the pharmaceutically acceptable salt thereof, or the composition according to the disclosure in the manufacture of a medicament for reducing the size of a solid tumor in a subject.


[0007e] In another aspect, the present disclosure refers to a use of the compound in accordance with the disclosure, or the pharmaceutically acceptable salt thereof, or the composition according to the disclosure in the manufacture of a medicament for treating a cell proliferation disorder in a subject.

[0007f] In another aspect, the present disclosure refers to a compound, or pharmaceutically acceptable salts thereof, in accordance with the disclosure, or the pharmaceutically acceptable salt thereof, or the composition according to the disclosure for use in therapy.

[0008] The compounds provided by the disclosure have MT-based KIF18A modulatory activity and, in particular, KIF18A inhibitory activity. To this end, the disclosure also provides the use of these compounds, as well as pharmaceutically acceptable salts thereof, in the preparation and manufacture of a pharmaceutical composition or medicament for therapeutic, prophylactic, acute or chronic treatment of KIF18A mediated diseases and disorders, including without limitation, cancer. Thus, the compounds of the disclosure are

useful in the manufacture of anti-cancer medicaments. The disclosure also provides processes for making compounds of Formula I, as well as intermediates useful in such processes.

[0009] In example 1, the present disclosure provides a compound of Formula (I), A compound of formula I:



or any pharmaceutically-acceptable salt thereof, wherein:

X¹ is N or -CR⁶;

R¹ is -CN, or a group -Z-R¹² wherein Z is -C₀₋₄alk-, -NR¹¹-, -NR¹¹SO₂-, -SO₂NR¹¹-, -NR¹¹-S(=O)(=NH), -S(=O)(=NH)-, -S-, -S(=O)-, -SO₂-, C₀₋₄alk-O-, -(C=O)-, -(C=O)NR¹¹-, -C=N(OH)-, or -NR¹¹(C=O); or

the group -Z-R¹² is -N=S(=O)-(R¹²)₂, wherein the two R¹² pair can alternatively combine with the sulfur atom attached to each of them to form a saturated or partially-saturated 3-, 4-, 5-, or 6-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S;

R² is halo or a group -Y-R¹³, wherein Y is -C₀₋₄alk-, -N(C₀₋₁alk)-C₀₋₄alk-, -C(=O)NR^aR^a(C₁₋₄alk), -O-C₀₋₄alk-, S, S=O, S(=O)₂-, -SO₂NR¹³, or -S(=O)(=NH)-;

R³ is H, C₁₋₄alk, or C₁₋₄haloalk;

R⁴ is H, halo, R^{4a} or R^{4b};


R⁵ is H, halo, C₁₋₈alk, or C₁₋₄haloalk;

R⁶ is H, halo, C₁₋₈alk, C₁₋₄haloalk, -O-C₁₋₈alk, or -O-R^{6a}; wherein R^{6a} is a saturated or partially-saturated 3-, 4-, 5-, or 6-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S;

R⁷ is H, halo, C₁₋₈alk, or C₁₋₄haloalk;

R⁸ is H, halo, C₁₋₈alk, C₁₋₄haloalk, -OH, -O-R^{8a}, or -O-R^{8b};

R⁹ is H, halo, C₁₋₈alk, or C₁₋₄haloalk;



Each of R^{10a} , R^{10b} , R^{10c} , R^{10d} , R^{10e} , R^{10f} , R^{10g} , R^{10h} , R^{10i} , and R^{10j} is H, halo, R^{10k} , or R^{10l} ;

or alternatively, each of R^{10a} and R^{10b} pair, R^{10c} and R^{10d} pair, R^{10e} and R^{10f} pair, R^{10g} and R^{10h} pair, or R^{10i} and R^{10j} pair, independently, can combine with the carbon atom attached to each of them to form a saturated or partially-saturated 3-, 4-, 5-, 6-membered monocyclic ring spiro to the R^x ring; wherein said 3-, 4-, 5-, 6-membered monocyclic ring contains 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S, and further wherein said 3-, 4-, 5-, 6-membered monocyclic ring is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, $C_{1-6}alk$, $C_{1-4}haloalk$, $-OR^a$, $-OC_{1-4}haloalk$, CN, $-NR^aR^a$, or oxo;

R^{11} is H, R^{11a} , or R^{11b} ;

R^{12} is H, R^{12a} , or R^{12b} ;

R^{13} is R^{13a} or R^{13b} ;

R^{4a} , R^{8a} , R^{10k} , R^{11a} , R^{12a} , and R^{13a} is independently, at each instance, selected from the group consisting of a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic or 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, or 12-membered bicyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S, which is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, $C_{1-6}alk$, $C_{1-4}haloalk$, $-OR^a$, $-OC_{1-4}haloalk$, CN, $-C(=O)R^b$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}alkNR^aR^a$, $-OC_{2-6}alkOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkNR^aR^a$, $-NR^aC_{2-6}alkOR^a$, $-C_{1-6}alkNR^aR^a$, $-C_{1-6}alkOR^a$, $-C_{1-6}alkN(R^a)C(=O)R^b$, $-C_{1-6}alkOC(=O)R^b$, $-C_{1-6}alkC(=O)NR^aR^a$, $-C_{1-6}alkC(=O)OR^a$, R^{14} , and oxo;

R^{4b} , R^{8b} , R^{10l} , R^{11b} , R^{12b} , and R^{13b} is independently, at each instance, selected from the group consisting of $C_{1-6}alk$ substituted by 0, 1, 2, 3, 4, or 5 group(s) selected from F, Cl, Br, $-OR^a$, $-OC_{1-4}haloalk$, or CN;


R^{14} is independently, at each instance, selected from the group consisting of a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic or 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, or 12-membered bicyclic ring containing 0, 1, 2 or 3 N atoms and 0 or 1 atoms selected from O and S, which is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, $C_{1-6}alk$, $C_{1-4}haloalk$, $-OR^a$,

-OC₁₋₄haloalk, CN, -C(=O)R^b, -C(=O)OR^a, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a, -NR^aC₂₋₆alkOR^a, -C₁₋₆alkNR^aR^a, -C₁₋₆alkOR^a, -C₁₋₆alkN(R^a)C(=O)R^b, -C₁₋₆alkOC(=O)R^b, -C₁₋₆alkC(=O)NR^aR^a, -C₁₋₆alkC(=O)OR^a, and oxo;


R^a is independently, at each instance, H or R^b; and

R^b is independently, at each instance, C₁₋₆alk, phenyl, or benzyl, wherein the C₁₋₆alk is being substituted by 0, 1, 2 or 3 substituents selected from halo, -OH, -OC₁₋₄alk, -NH₂, -NHC₁₋₄alk, -OC(=O)C₁₋₄alk, or -N(C₁₋₄alk)C₁₋₄alk; and the phenyl or benzyl is being substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄alk, C₁₋₃haloalk, -OH, -OC₁₋₄alk, -NH₂, -NHC₁₋₄alk, -OC(=O)C₁₋₄alk, or -N(C₁₋₄alk)C₁₋₄alk.

[0010] In example 2, the present disclosure provides compounds wherein X¹ is N; having the formula (Ia):



[0011] In example 3, the present disclosure provides compounds wherein X¹ is -CR⁶; having the formula (Ib):




[0012] In example 4, the present disclosure provides compounds wherein R³ is H or methyl. Preferably, R³ is H.

[0013] In example 5, the present disclosure provides compounds wherein each of R^{10c}, R^{10d}, R^{10e}, R^{10f}, R^{10g}, R^{10h}, R¹⁰ⁱ, and R^{10j} is H, halo, C₁₋₆alk, or C₁₋₄haloalk; and each of R^{10a} and R^{10b} pair combine with the carbon atom attached to each of them form a saturated 3-, 4-, or 5-membered monocyclic ring spiro to the R^x ring; wherein said ring contains 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S.


[0014] In example 6, the present disclosure provides compounds wherein each of R^{10c} , R^{10d} , R^{10e} , R^{10f} , R^{10g} , R^{10h} , R^{10i} , and R^{10j} is H, methyl, or ethyl; and each of R^{10a} and R^{10b} pair combine with the carbon atom attached to each of them form a cyclopropyl, cyclobutyl, or cyclopentyl ring spiro to the R^x ring.

[0015] In example 7, the present disclosure provides compounds in accordance with examples 1-6, or




pharmaceutically acceptable salts thereof, wherein the group

is selected from:



[0016] In example 8, the present disclosure provides compounds in accordance with examples 1-7, or




pharmaceutically acceptable salts thereof, wherein the group

is

[0017] In example 9, the present disclosure provides compounds in accordance with examples 1-8, or pharmaceutically acceptable salts thereof, wherein R^1 is $-\text{CN}$, or a group $-\text{Z}-\text{R}^{12}$, wherein Z is a bond, $-\text{NH}-$, $-\text{NH}\text{SO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{S}(=\text{O})(=\text{NH})-$, $-\text{S}-$, $-\text{S}(=\text{O})-$, $-\text{SO}_2-$, $-(\text{C}=\text{O})-$, $-(\text{C}=\text{O})\text{NH}-$, or $-\text{NH}(\text{C}=\text{O})-$; and R^{12} is selected from:

(a) H;

(b) cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, oxiranyl, oxetanyl, tetrahydrofuranyl, azetidiny, imidazolyl, morpholiny, pyrrolidiny, piperaziny,



; wherein each said ring is substituted by 0, 1, 2 or 3 group(s) selected from wherein each ring is substituted by 0, 1, 2 or 3 OH, F, methyl, $-\text{CH}_2\text{OH}$, $-\text{C}(=\text{O})\text{OCH}_3$, $-\text{C}(=\text{O})\text{OC}(\text{CH}_3)_3$, NH_2 , CN, and oxo; or

(c) C₁₋₆alk substituted by 0, 1, 2 or 3 OH, F, -C(=O)OCH₃, -NH₂, -NH(CH₃), or -N(CH₃)₂.


[0018] In example 10, the present disclosure provides compounds in accordance with examples 1-9, or pharmaceutically acceptable salts thereof, wherein R¹ is -CN, or a group -Z-R¹², wherein Z is absent, -NH-, -NHSO₂-, -SO₂NH-, -S(=O)(=NH)-, -S-, -S(=O)-, -SO₂-, -(C=O)-, -(C=O)NH-, or -NH(C=O)-; and

(a) R¹² is H;

(b) R¹² is oxetanyl, cyclopropyl; or

(c) R¹² is C₁₋₆alk substituted by 0, 1, 2 or 3 OH group(s).

[0019] In example 11, the present disclosure provides compounds in accordance with examples 1-10, or pharmaceutically acceptable salts thereof, wherein the group -Z-R¹² is -N=S(=O)-(R¹²)₂, wherein the two R¹² pair can alternatively combine with the sulfur atom attached to each of them to form a saturated or partially-saturated 3-, 4-, 5-, or 6-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S; which is selected from:



[0020] In example 12, the present disclosure provides compounds in accordance with examples 1-11, or pharmaceutically acceptable salts thereof, wherein R¹ is a group -Z-R¹², wherein Z is -NHSO₂- or -SO₂NH-; and R¹² is oxetanyl, cyclopropyl, or R¹² is C₁₋₆alk substituted by 0, 1, 2 or 3 OH group(s).

[0021] In example 13, the present disclosure provides compounds in accordance with examples 1-12, or pharmaceutically acceptable salts thereof, wherein R¹ is a group -Z-R¹², wherein Z is -NHSO₂- and R¹² is -CH₂-CH₂-OH.


[0022] In example 14, the present disclosure provides compounds in accordance with examples 1-13, or pharmaceutically acceptable salts thereof, wherein R² is halo or a group -Y-R¹³, wherein Y is a bond, -NH-, -NH-(CH₂)₀₋₄-, or -O-(CH₂)₀₋₄-; and R¹³ is a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic or 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, or 12-membered bicyclic ring containing 0, 1, 2 or 3 N atoms and 0 or 1 atoms selected from O and S, which is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, C₁₋₆alk, C₁₋₄haloalk, -OH, -OC₁₋₄haloalk, CN, R¹⁴, and oxo; or

R¹³ is C₁₋₆alk substituted by 0, 1, 2, 3, 4, or 5 group(s) selected from F, Cl, Br, -OH, -OC₁₋₄haloalk, or CN.


[0023] In example 15, the present disclosure provides compounds in accordance with examples 1-14, or pharmaceutically acceptable salts thereof, wherein R² is a saturated 5- or 6-membered monocyclic ring wherein each said ring contains 0, 1, or 2 N atoms and 0 or 1 O atom, and wherein each said ring is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, C₁₋₆alk, C₁₋₄haloalk, -OH, -OC₁₋₄haloalk, CN, R¹⁴, and oxo.

[0024] In example 16, the present disclosure provides compounds in accordance with examples 1-15, or pharmaceutically acceptable salts thereof, wherein R² is (a) halo; (b) a group -Y-R¹³, wherein Y is a bond;

and R¹³ is morpholinyl, piperidinyl, azetidiny, pyrrolidinyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, piperazinyl, tetrahydrofuranyl,



; wherein each said ring is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, methyl, CF₃, -OH, -OCHF₂, CN, and oxo; or (c) a group -Y-R¹³, wherein Y is NH, -O-, -O-(CH₂)-, -O-(CH₂)-(CH₂)-, or


-O-(CH₂)-(CH₂)-(CH₂)-, and wherein R¹³ is ; or R¹³ is C₁₋₆alk substituted by 0, 1, 2, 3, 4, or 5 group(s) selected from F, Cl, Br, methyl, CF₃, -OH, or CN.

[0025] In example 17, the present disclosure provides compounds in accordance with examples 1-16, or pharmaceutically acceptable salts thereof, wherein R² is morpholinyl or piperidinyl substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, methyl, CF₃, -OH, -OCHF₂, CN, or oxo

[0026] In example 18, the present disclosure provides compounds in accordance with examples 1-17, or pharmaceutically acceptable salts thereof, wherein R² is morpholinyl substituted by 1, 2 or 3 methyl group(s).

[0027] In example 19, the present disclosure provides compounds in accordance with examples 1-18, or pharmaceutically acceptable salts thereof, wherein R² is piperidinyl substituted by 1, 2 or 3 fluoro group(s).

[0028] In example 20, the present disclosure provides compounds in accordance with examples 1-19, or pharmaceutically acceptable salts thereof, wherein R² is



[0029] In example 21, the present disclosure provides compounds in accordance with examples 1-20, or pharmaceutically acceptable salts thereof, wherein Z is a bond, -NH-, -NH₂SO₂-, -SO₂NH-, -N=S(=O)<(R^a)₂ (wherein each R¹¹ is independently selected from the group consisting of H, methyl, or isopropyl), -S(=O)(=NH)-, -S-, -S(=O)-, -SO₂-, -(C=O)-, -(C=O)NH-, or -NH(C=O)-.

[0030] In example 22, the present disclosure provides compounds in accordance with examples 1-21, or pharmaceutically acceptable salts thereof, wherein R¹² is selected from (a) H; (b) C₁₋₆alk substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, -OH, -OCH₃, or cyclopropyl; or (c) a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0 or 1 atoms selected from O and S, which is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, C₁₋₆alk, C₁₋₄haloalk, -C₁₋₆alkOH, -OH, -OCH₃, -NH₂, or oxo.

[0031] In example 23, the present disclosure provides compounds in accordance with examples 1-22, or pharmaceutically acceptable salts thereof, wherein R¹² is selected from cyclopropyl, cyclobutyl, cyclopentyl, oxetanyl, azetidiny, tetrahydrofuranyl, or 1,3,4-oxathiazinanyl.

[0032] In example 24, the present disclosure provides compounds in accordance with examples 1-23, or pharmaceutically acceptable salts thereof, wherein R⁴ is selected from (a) H; (b) C₁₋₆alk substituted by 0, 1, 2 or 3 OH group(s); or (c) cyclopropyl.

[0033] In example 25, the present disclosure provides compounds in accordance with examples 1-24, or pharmaceutically acceptable salts thereof, wherein R⁴ is methyl.

[0034] In example 26, the present disclosure provides compounds in accordance with examples 1-25, or pharmaceutically acceptable salts thereof, wherein R⁵ is H.

[0035] In example 27, the present disclosure provides compounds in accordance with examples 1-26, or pharmaceutically acceptable salts thereof, wherein R⁶ is H or F.

[0036] In example 28, the present disclosure provides compounds in accordance with examples 1-27, or pharmaceutically acceptable salts thereof, wherein R⁷ is H or F.

[0037] In example 29, the present disclosure provides compounds in accordance with examples 1-28, or pharmaceutically acceptable salts thereof, wherein R⁸ is H.

[0038] In example 30, the present disclosure provides compounds in accordance with examples 1-29, or pharmaceutically acceptable salts thereof, wherein R⁹ is H.

[0039] In example 31, the present disclosure provides a compound, or pharmaceutically acceptable salts thereof, selected from:

Ex. #	Chemical Structure	Name
1		<i>N</i> -(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
1-7		<i>N</i> -(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
2		<i>N</i> -(2-(2-Hydroxypropan-2-yl)pyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
2-4		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
2-8		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
3		(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
4		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

Ex. #	Chemical Structure	Name
5-1		(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
5-2		(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
6-7		N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(ethylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
7		N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
8-1		(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
8-2		(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
10-1		(R)-N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

Ex. #	Chemical Structure	Name
10-2		(S)-N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
12		N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
13-1		(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
13-2		(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
14		N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxy-2-methylpropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
15		N-(2-(4,4-Difluorocyclohexyl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
16-1		(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

Ex. #	Chemical Structure	Name
16-2		(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
17		N-(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(N-(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
18-1		2-(6-Azaspiro[2.5]octan-6-yl)-4-(R-cyclopropylsulfonimidoyl)-N-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide
18-2		2-(6-Azaspiro[2.5]octan-6-yl)-4-(S-cyclopropylsulfonimidoyl)-N-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide
20		(N¹-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)terephthalamide
21		4-(Azetidin-3-ylsulfonyl)-N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

Ex. #	Chemical Structure	Name
22		N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylazetidin-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

or any pharmaceutically-acceptable salt thereof.

[0040] In sub-example 31a, the present disclosure provides *N*-(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0041] In sub-example 31b, the present disclosure provides *N*-(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0042] In sub-example 31c, the present disclosure provides *N*-(2-(2-Hydroxypropan-2-yl)pyrimidin-4-yl)-4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0043] In sub-example 31d, the present disclosure provides *N*-(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0044] In sub-example 31e, the present disclosure provides *N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0045] In sub-example 31f, the present disclosure provides (*R*)-4-((2-Hydroxyethyl)sulfonamido)-*N*-(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0046] In sub-example 31g, the present disclosure provides *N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0047] In sub-example 31h, the present disclosure provides (*R*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0048] In sub-example 31i, the present disclosure provides (*S*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0049] In sub-example 31j, the present disclosure provides *N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(ethylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0050] In sub-example 31k, the present disclosure provides *N*-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0051] In sub-example 31l, the present disclosure provides (*R*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0052] In sub-example 31m, the present disclosure provides (*S*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0053] In sub-example 31n, the present disclosure provides (*R*)-*N*-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0054] In sub-example 31o, the present disclosure provides (*S*)-*N*-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0055] In sub-example 31p, the present disclosure provides *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0056] In sub-example 31q, the present disclosure provides (*S*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0057] In sub-example 31r, the present disclosure provides (*R*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0058] In sub-example 31s, the present disclosure provides *N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxy-2-methylpropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0059] In sub-example 31t, the present disclosure provides *N*-(2-(4,4-Difluorocyclohexyl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0060] In sub-example 31u, the present disclosure provides (*R*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0061] In sub-example 31v, the present disclosure provides (*S*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0062] In sub-example 31w, the present disclosure provides *N*-(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(*N*-(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0063] In sub-example 31x, the present disclosure provides 2-(6-Azaspiro[2.5]octan-6-yl)-4-(*R*-cyclopropylsulfonimidoyl)-*N*-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide, or the pharmaceutically acceptable salt thereof.

[0064] In sub-example 31y, the present disclosure provides 2-(6-Azaspiro[2.5]octan-6-yl)-4-(*S*-cyclopropylsulfonimidoyl)-*N*-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide, or the pharmaceutically acceptable salt thereof.

[0065] In sub-example 31z, the present disclosure provides (*N*¹-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)terephthalamide, or the pharmaceutically acceptable salt thereof.

[0066] In sub-example 31aa, the present disclosure provides 4-(Azetidin-3-ylsulfonyl)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0067] In sub-example 31ab, the present disclosure provides *N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylazetidin-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide, or the pharmaceutically acceptable salt thereof.

[0068] In example 32, the present disclosure provides pharmaceutical compositions comprising a compound, or pharmaceutically acceptable salts thereof, in accordance with any one of examples 1 to 31, and a pharmaceutically acceptable diluent or carrier.

[0069] In example 33, the present disclosure provides a method of treating a condition that may be treated with KIF18a inhibitors, the method comprising administering to a patient in need thereof a therapeutically effective amount of the compound in accordance with examples 1 to 31, or the composition according to example 31.

[0070] In example 34, the present disclosure provides the method of example 33, wherein said condition is cancer selected from the group consisting of (a) a solid or hematologically derived tumor selected from cancer of the cancer of the bladder, endometrial, lung squamous cell, breast, colon, kidney, liver, lung, small cell lung cancer, esophagus, gall-bladder, brain, head and neck, ovary, pancreas, stomach, cervix, thyroid, prostate and skin, (b) a hematopoietic tumor of lymphoid lineage selected from leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma and Burkett's lymphoma, (c) a hematopoietic tumor of myeloid lineage selected from acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia (d) a tumor of mesenchymal origin selected from fibrosarcoma and rhabdomyosarcoma, (e) a tumor of the central and peripheral nervous system selected from astrocytoma, neuroblastoma, glioma and schwannoma, or (f) a melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderoma pigmentosum, keratocanthoma, thyroid follicular cancer or Kaposi's sarcoma.

[0071] In a sub-example 34a, the present disclosure provides the method of example 33, wherein said condition is cancer selected from the group consisting of melanoma, prostate cancer, cervical cancer, breast cancer, colon cancer, sarcoma, or leukemia. See: Zhang C. et. al., “Kif18A is involved in human breast carcinogenesis”, *Carcinogenesis*, 2010 Sep;31(9):1676-84. doi: 10.1093/carcin/bgq134. Epub 2010 Jul 1. See also: (1) <https://www.proteinatlas.org/ENSG00000121621-KIF18A/pathology>; (2) Nagahara, M. et. al., “Kinesin 18A expression: clinical relevance to colorectal cancer progression”, *Int. J. Cancer*: 129, 2543–2552 (2011) VC 2011 UIC; and (3) Yu, Y. et. al., “The Role of Kinesin Family Proteins in Tumorigenesis and Progression - Potential Biomarkers and Molecular Targets for Cancer Therapy”, *Cancer* 2010;116:5150–60. VC 2010 American Cancer Society. In a sub-example 34b, the present disclosure provides the method of example 33, wherein said condition is any one of the cancer specified in examples (a), (b), (c), (d), (e), or (f).

[0072] In example 35, the present disclosure provides a method of reducing the size of a solid tumor in a subject, the method comprising administering to the subject in need thereof a therapeutically effective amount of the compound in accordance with any one of examples 1 to 31, or the composition according to example 32.

[0073] In example 36, the present disclosure provides a method of treating a cell proliferation disorder in a subject, the method comprising administering to the subject in need thereof a therapeutically effective amount of the compound in accordance with any one of examples 1 to 31, or the composition according to example 32.

[0074] In example 37, the present disclosure provides a method of inhibiting KIF18A in a cell, comprising contacting the cell with a compound, or pharmaceutically acceptable salts thereof, in accordance with any one of examples 1 to 31, or the composition according to example 32.

[0075] In example 38, the disclosure provides use of the compound or the pharmaceutically acceptable salt of said compound according to any one of examples 1 to 31, or the pharmaceutical composition according to example 32, in the preparation of a medicament for treating a condition that may be treated with KIF18a inhibitors.

[0076] In example 39, the disclosure provides use of examples 38, wherein the condition is cancer selected from the group consisting of (a) a solid or hematologically derived tumor selected from cancer of the cancer of the bladder, endometrial, lung squamous cell, breast, colon, kidney, liver, lung, small cell lung cancer, esophagus, gall-bladder, brain, head and neck, ovary, pancreas, stomach, cervix, thyroid, prostate and skin, (b) a hematopoietic tumor of lymphoid lineage selected from leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma and Burkett's lymphoma, (c) a hematopoietic tumor of myeloid lineage selected from acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia (d) a tumor of mesenchymal origin selected from fibrosarcoma and rhabdomyosarcoma, (e) a tumor of the central and peripheral nervous system selected from astrocytoma, neuroblastoma, glioma and

schwannoma, or (f) a melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderoma pigmentosum, keratocanthoma, thyroid follicular cancer or Kaposi's sarcoma.

[0077] In a sub-example 39a, the present disclosure provides the use of example 33, wherein said condition is any one of the cancer specified in examples (a), (b), (c), (d), (e), or (f).

[0078] In example 40, the disclosure provides use of the compound or the pharmaceutically acceptable salt of said compound according to any one of examples 1 to 31, or the pharmaceutical composition according to example 32, in the preparation of a medicament for reducing the size of a solid tumor in a subject.

[0079] In example 41, the disclosure provides use of the compound or the pharmaceutically acceptable salt of said compound according to any one of examples 1 to 31, or the pharmaceutical composition according to example 32, in the preparation of a medicament for treating a cell proliferation disorder in a subject.

[0080] In example 42, the disclosure provides use of the compound or the pharmaceutically acceptable salt of said compound according to any one of examples 1 to 31, or the pharmaceutical composition according to example 32, in the preparation of a medicament for inhibiting KIF18A in a cell.

[0081] In example 43, the disclosure provides a method of preparing a compound of Formula (I) as described herein.

[0082] In example 44, the disclosure provides an intermediate compound used in the method of preparing a compound of Formula (I) as described herein.

[0083] It is to be understood that the above reference to any example is intended to include any and all its sub-example thereof. For example, reference to example 31 includes reference to sub-examples 31a-31ab.

[0084] The present disclosure includes all pharmaceutically acceptable isotopically-labelled compounds of the present disclosure wherein one or more atoms are replaced by atoms having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number which predominates in nature.

[0085] Examples of isotopes suitable for inclusion in the compounds of the disclosure include, but are not limited to, isotopes of hydrogen, such as ^2H and ^3H , carbon, such as ^{11}C , ^{13}C and ^{14}C , chlorine, such as ^{38}Cl , fluorine, such as ^{18}F , iodine, such as ^{123}I and ^{125}I , nitrogen, such as ^{13}N and ^{15}N , oxygen, such as ^{15}O , ^{17}O and ^{18}O , phosphorus, such as ^{32}P , and sulphur, such as ^{35}S .

[0086] Certain isotopically-labelled compounds of the present disclosure, for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, i.e. ^3H , and carbon-14, i.e. ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

[0087] Substitution with heavier isotopes such as deuterium, i.e. ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased in vivo half-life or reduced dosage requirements, and hence may be preferred in some circumstances.

[0088] Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy.


[0089] Isotopically-labeled compounds of the present disclosure can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples and Preparations using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

[0090] Pharmaceutically acceptable solvates in accordance with the disclosure include those wherein the solvent of crystallization may be isotopically substituted, e.g. D₂O, d₆-acetone, d₆-DMSO.

[0091] Specific examples of the present disclosure include the compounds exemplified in the Examples below and their pharmaceutically acceptable salts, complexes, solvates, polymorphs, stereoisomers, metabolites, prodrugs, and other derivatives thereof.

[0092] Unless otherwise specified, the following definitions apply to terms found in the specification and claims:

[0093] “C_{α-β}alk” means an alkyl group comprising a minimum of α and a maximum of β carbon atoms in a branched or linear relationship or any combination of the three, wherein α and β represent integers. The alkyl groups described in this section may also contain one or two double or triple bonds. A designation of C₀alk indicates a direct bond. Examples of C₁₋₆alkyl include, but are not limited to the following:




[0094] "Benzo group", alone or in combination, means the divalent radical C₆H₄=, one representation of which is -CH=CH-CH=CH-, that when vicinally attached to another ring forms a benzene-like ring--for example tetrahydronaphthylene, indole and the like.

[0095] The terms “oxo” and “thioxo” represent the groups =O (as in carbonyl) and =S (as in thiocarbonyl), respectively.


[0096] “Halo” or “halogen” means a halogen atom selected from F, Cl, Br and I.

[0097] “C_{α-β}haloalk” means an alk group, as described above, wherein any number --at least one-- of the hydrogen atoms attached to the alk chain are replaced by F, Cl, Br or I.


[0098] The group N(R^a)R^a and the like include substituents where the two R^a groups together form a ring, optionally including a N, O or S atom, and include groups such as:




[0099] The group N(C_{α-β}alk) C_{α-β}alk, wherein α and β are as defined above, include substituents where the two C_{α-β}alk groups together form a ring, optionally including a N, O or S atom, and include groups






[0100] “Bicyclic ring” means a group that features two joined rings. A bicyclic ring can be carbocyclic (all of the ring atoms are carbons), or heterocyclic (the rings atoms consist, for example, 1, 2 or 3 heteroatoms, such as N, O, or S, in addition to carbon atoms). The two rings can both be aliphatic (*e.g.* decalin and norbornane), or can be aromatic (*e.g.* naphthalene), or a combination of aliphatic and aromatic (*e.g.* tetralin). Bicyclic rings include (a) spirocyclic compounds, wherein the two rings share only one single atom, the spiro atom, which is usually a quaternary carbon. Examples of spirocyclic compound include, but are not limited to:



[0101] (b) fused bicyclic compounds, wherein two rings share two adjacent atoms. In other words, the rings share one covalent bond, *i.e.* the bridgehead atoms are directly connected (*e.g.* α -thujene and decalin). Examples of fused bicyclic rings include, but are not limited to:



[illegible]



[0105] "Pharmaceutically-acceptable salt" means a salt prepared by conventional means, and are well known by those skilled in the art. The "pharmacologically acceptable salts" include basic salts of inorganic and organic acids, including but not limited to hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, malic acid, acetic acid, oxalic acid, tartaric acid, citric acid, lactic acid, fumaric acid, succinic acid, maleic acid, salicylic acid, benzoic acid, phenylacetic acid, mandelic acid and the like. When compounds of the disclosure include an acidic function such as a carboxy group, then suitable pharmaceutically acceptable cation pairs for the carboxy group are well known to those skilled in the art and include alkaline, alkaline earth, ammonium, quaternary ammonium cations and the like. For additional examples of "pharmacologically acceptable salts," *see infra* and Berge et al., J. Pharm. Sci. 66:1 (1977).

[0106] "Saturated, partially-saturated or unsaturated" includes substituents saturated with hydrogens, substituents completely unsaturated with hydrogens and substituents partially saturated with hydrogens.

[0107] "Leaving group" generally refers to groups readily displaceable by a nucleophile, such as an amine, a thiol or an alcohol nucleophile. Such leaving groups are well known in the art. Examples of such leaving groups include, but are not limited to, N-hydroxysuccinimide, N-hydroxybenzotriazole, halides, triflates, tosylates and the like. Preferred leaving groups are indicated herein where appropriate.

[0108] "Protecting group" generally refers to groups well known in the art which are used to prevent selected reactive groups, such as carboxy, amino, hydroxy, mercapto and the like, from undergoing undesired reactions, such as nucleophilic, electrophilic, oxidation, reduction and the like. Preferred protecting groups are indicated herein where appropriate. Examples of amino protecting groups include, but are not limited to, aralkyl, substituted aralkyl, cycloalkenylalkyl and substituted cycloalkenyl alkyl, allyl, substituted allyl, acyl, alkoxycarbonyl, aralkoxycarbonyl, silyl and the like. Examples of aralkyl


include, but are not limited to, benzyl, ortho-methylbenzyl, trityl and benzhydryl, which can be optionally substituted with halogen, alkyl, alkoxy, hydroxy, nitro, acylamino, acyl and the like, and salts, such as phosphonium and ammonium salts. Examples of aryl groups include phenyl, naphthyl, indanyl, anthracenyl, 9-(9-phenylfluorenyl), phenanthrenyl, durenyl and the like. Examples of cycloalkenylalkyl or substituted cycloalkenylalkyl radicals, preferably have 6-10 carbon atoms, include, but are not limited to, cyclohexenyl methyl and the like. Suitable acyl, alkoxycarbonyl and aralkoxycarbonyl groups include benzyloxycarbonyl, t-butoxycarbonyl, iso-butoxycarbonyl, benzoyl, substituted benzoyl, butyryl, acetyl, trifluoroacetyl, trichloro acetyl, phthaloyl and the like. A mixture of protecting groups can be used to protect the same amino group, such as a primary amino group can be protected by both an aralkyl group and an aralkoxycarbonyl group. Amino protecting groups can also form a heterocyclic ring with the nitrogen to which they are attached, for example, 1,2-bis(methylene)benzene, phthalimidyl, succinimidyl, maleimidyl and the like and where these heterocyclic groups can further include adjoining aryl and cycloalkyl rings. In addition, the heterocyclic groups can be mono-, di- or tri-substituted, such as nitrophthalimidyl. Amino groups may also be protected against undesired reactions, such as oxidation, through the formation of an addition salt, such as hydrochloride, toluenesulfonic acid, trifluoroacetic acid and the like. Many of the amino protecting groups are also suitable for protecting carboxy, hydroxy and mercapto groups. For example, aralkyl groups. Alkyl groups are also suitable groups for protecting hydroxy and mercapto groups, such as tert-butyl.

[0109] Silyl protecting groups are silicon atoms optionally substituted by one or more alkyl, aryl and aralkyl groups. Suitable silyl protecting groups include, but are not limited to, trimethylsilyl, triethylsilyl, triisopropylsilyl, tert-butyl dimethylsilyl, dimethylphenylsilyl, 1,2-bis(dimethylsilyl)benzene, 1,2-bis(dimethylsilyl)ethane and diphenylmethylsilyl. Silylation of an amino groups provide mono- or di-silylamino groups. Silylation of aminoalcohol compounds can lead to a N,N,O-trisilyl derivative. Removal of the silyl function from a silyl ether function is readily accomplished by treatment with, for example, a metal hydroxide or ammonium fluoride reagent, either as a discrete reaction step or in situ during a reaction with the alcohol group. Suitable silylating agents are, for example, trimethylsilyl chloride, tert-butyl-dimethylsilyl chloride, phenyldimethylsilyl chloride, diphenylmethyl silyl chloride or their combination products with imidazole or DMF. Methods for silylation of amines and removal of silyl protecting groups are well known to those skilled in the art. Methods of preparation of these amine derivatives from corresponding amino acids, amino acid amides or amino acid esters are also well known to those skilled in the art of organic chemistry including amino acid/amino acid ester or aminoalcohol chemistry.

[0110] Protecting groups are removed under conditions which will not affect the remaining portion of the molecule. These methods are well known in the art and include acid hydrolysis, hydrogenolysis and the like. A preferred method involves removal of a protecting group, such as removal of a benzyloxycarbonyl group by hydrogenolysis utilizing palladium on carbon in a suitable solvent system such as an alcohol, acetic acid, and the like or mixtures thereof. A t-butoxycarbonyl protecting group can be removed

utilizing an inorganic or organic acid, such as HCl or trifluoroacetic acid, in a suitable solvent system, such as dioxane or methylene chloride. The resulting amino salt can readily be neutralized to yield the free amine. Carboxy protecting group, such as methyl, ethyl, benzyl, tert-butyl, 4-methoxyphenylmethyl and the like, can be removed under hydrolysis and hydrogenolysis conditions well known to those skilled in the art.

[0111] It should be noted that compounds of the disclosure may contain groups that may exist in tautomeric forms, such as cyclic and acyclic amidine and guanidine groups, heteroatom substituted heteroaryl groups ($Y' = O, S, NR$), and the like, which are illustrated in the following examples:



[0112] and though one form is named, described, displayed and/or claimed herein, all the tautomeric forms are intended to be inherently included in such name, description, display and/or claim.

[0113] Prodrugs of the compounds of this disclosure are also contemplated by this disclosure. A prodrug is an active or inactive compound that is modified chemically through in vivo physiological action, such as hydrolysis, metabolism and the like, into a compound of this disclosure following administration of the prodrug to a patient. The suitability and techniques involved in making and using prodrugs are well known by those skilled in the art. For a general discussion of prodrugs involving esters see Svensson and Tunek *Drug Metabolism Reviews* 165 (1988) and Bundgaard *Design of Prodrugs*, Elsevier (1985). Examples of a masked carboxylate anion include a variety of esters, such as alkyl (for example, methyl, ethyl), cycloalkyl (for example, cyclohexyl), aralkyl (for example, benzyl, p-methoxybenzyl), and alkylcarbonyloxyalkyl (for example, pivaloyloxymethyl). Amines have been masked as arylcarbonyloxymethyl substituted derivatives which are cleaved by esterases in vivo releasing the free drug and formaldehyde (Bunggaard *J. Med. Chem.* 2503 (1989)). Also, drugs containing an acidic NH group, such as imidazole, imide, indole and the like, have been masked with N-acyloxymethyl groups (Bundgaard *Design of Prodrugs*, Elsevier (1985)). Hydroxy groups have been masked as esters and ethers. EP 039,051 (Sloan and Little, 4/11/81) discloses Mannich-base hydroxamic acid prodrugs, their preparation and use.

[0114] The specification and claims contain listing of species using the language “selected from . . . and . . .” and “is . . . or . . .” (sometimes referred to as Markush groups). When this language is used in this application, unless otherwise stated it is meant to include the group as a whole, or any single members thereof, or any subgroups thereof. The use of this language is merely for shorthand purposes and is not meant in any way to limit the removal of individual elements or subgroups as needed.

[0115] PHARMACEUTICAL COMPOSITIONS, DOSING, AND ROUTES OF ADMINISTRATION

[0116] Also provided herein are pharmaceutical compositions that includes a compound as disclosed herein, together with a pharmaceutically acceptable excipient, such as, for example, a diluent or carrier. Compounds and pharmaceutical compositions suitable for use in the present disclosure include those wherein the compound can be administered in an effective amount to achieve its intended purpose. Administration of the compound described in more detail below.

[0117] Suitable pharmaceutical formulations can be determined by the skilled artisan depending on the route of administration and the desired dosage. See, e.g., Remington’s Pharmaceutical Sciences, 1435-712 (18th ed., Mack Publishing Co, Easton, Pennsylvania, 1990). Formulations may influence the physical state, stability, rate of in vivo release and rate of in vivo clearance of the administered agents. Depending on the route of administration, a suitable dose may be calculated according to body weight, body surface areas or organ size. Further refinement of the calculations necessary to determine the appropriate treatment dose is routinely made by those of ordinary skill in the art without undue experimentation, especially in light of the dosage information and assays disclosed herein as well as the pharmacokinetic data obtainable through animal or human clinical trials.

[0118] The phrases “pharmaceutically acceptable” or “pharmacologically acceptable” refer to molecular entities and compositions that do not produce adverse, allergic, or other untoward reactions when administered to an animal or a human. As used herein, “pharmaceutically acceptable e” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such excipients for pharmaceutically active substances is well

known in the art. Except insofar as any conventional media or agent is incompatible with the therapeutic compositions, its use in therapeutic compositions is contemplated. Supplementary active ingredients also can be incorporated into the compositions. In exemplary embodiments, the formulation may comprise corn syrup solids, high-oleic safflower oil, coconut oil, soy oil, L-leucine, calcium phosphate tribasic, L-tyrosine, L-proline, L-lysine acetate, DATEM (an emulsifier), L-glutamine, L-valine, potassium phosphate dibasic, L-isoleucine, L-arginine, L-alanine, glycine, L-asparagine monohydrate, L-serine, potassium citrate, L-threonine, sodium citrate, magnesium chloride, L-histidine, L-methionine, ascorbic acid, calcium carbonate, L-glutamic acid, L-cystine dihydrochloride, L-tryptophan, L-aspartic acid, choline chloride, taurine, m-inositol, ferrous sulfate, ascorbyl palmitate, zinc sulfate, L-carnitine, alpha-tocopheryl acetate, sodium chloride, niacinamide, mixed tocopherols, calcium pantothenate, cupric sulfate, thiamine chloride hydrochloride, vitamin A palmitate, manganese sulfate, riboflavin, pyridoxine hydrochloride, folic acid, beta-carotene, potassium iodide, phylloquinone, biotin, sodium selenate, chromium chloride, sodium molybdate, vitamin D3 and cyanocobalamin.

[0119] The compound can be present in a pharmaceutical composition as a pharmaceutically acceptable salt. As used herein, “pharmaceutically acceptable salts” include, for example base addition salts and acid addition salts.

[0120] Pharmaceutically acceptable base addition salts may be formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible. Examples of metals used as cations are sodium, potassium, magnesium, ammonium, calcium, or ferric, and the like. Examples of suitable amines include isopropylamine, trimethylamine, histidine, N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine.

[0121] Pharmaceutically acceptable acid addition salts include inorganic or organic acid salts. Examples of suitable acid salts include the hydrochlorides, formates, acetates, citrates, salicylates, nitrates, phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include, for example, formic, acetic, citric, oxalic, tartaric, or mandelic acids, hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, trifluoroacetic acid (TFA), propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane 1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene 2-

sulfonic acid, naphthalene 1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose 6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid.

[0122] Pharmaceutical compositions containing the compounds disclosed herein can be manufactured in a conventional manner, e.g., by conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen.

[0123] For oral administration, suitable compositions can be formulated readily by combining a compound disclosed herein with pharmaceutically acceptable excipients such as carriers well known in the art. Such excipients and carriers enable the present compounds to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by adding a compound as disclosed herein with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, for example, fillers and cellulose preparations. If desired, disintegrating agents can be added. Pharmaceutically acceptable ingredients are well known for the various types of formulation and may be for example binders (e.g., natural or synthetic polymers), lubricants, surfactants, sweetening and flavoring agents, coating materials, preservatives, dyes, thickeners, adjuvants, antimicrobial agents, antioxidants and carriers for the various formulation types.

[0124] When a therapeutically effective amount of a compound disclosed herein is administered orally, the composition typically is in the form of a solid (e.g., tablet, capsule, pill, powder, or troche) or a liquid formulation (e.g., aqueous suspension, solution, elixir, or syrup).

[0125] When administered in tablet form, the composition can additionally contain a functional solid and/or solid carrier, such as a gelatin or an adjuvant. The tablet, capsule, and powder can contain about 1 to about 95% compound, and preferably from about 15 to about 90% compound.

[0126] When administered in liquid or suspension form, a functional liquid and/or a liquid carrier such as water, petroleum, or oils of animal or plant origin can be added. The liquid form of the composition can further contain physiological saline solution, sugar alcohol solutions, dextrose or other saccharide solutions, or glycols. When administered in liquid or suspension form, the composition can contain about 0.5 to about 90% by weight of a compound disclosed herein, and preferably about 1 to about 50% of a compound disclosed herein. In one embodiment contemplated, the liquid carrier is non-aqueous or substantially non-aqueous. For administration in liquid form, the composition may be supplied as a rapidly-dissolving solid formulation for dissolution or suspension immediately prior to administration.

[0127] When a therapeutically effective amount of a compound disclosed herein is administered by intravenous, cutaneous, or subcutaneous injection, the composition is in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred composition for intravenous, cutaneous, or subcutaneous injection typically contains, in addition to a

compound disclosed herein, an isotonic vehicle. Such compositions may be prepared for administration as solutions of free base or pharmacologically acceptable salts in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions also can be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations can optionally contain a preservative to prevent the growth of microorganisms.

[0128] Injectable compositions can include sterile aqueous solutions, suspensions, or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions, suspensions, or dispersions. In all embodiments the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must resist the contaminating action of microorganisms, such as bacteria and fungi, by optional inclusion of a preservative. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. In one embodiment contemplated, the carrier is non-aqueous or substantially non-aqueous. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size of the compound in the embodiment of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many embodiments, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0129] Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the embodiment of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0130] Slow release or sustained release formulations may also be prepared in order to achieve a controlled release of the active compound in contact with the body fluids in the GI tract, and to provide a substantially constant and effective level of the active compound in the blood plasma. For example, release can be controlled by one or more of dissolution, diffusion, and ion-exchange. In addition, the slow release approach may enhance absorption via saturable or limiting pathways within the GI tract. For example, the compound may be embedded for this purpose in a polymer matrix of a biological degradable polymer, a water-soluble polymer or a mixture of both, and optionally suitable surfactants. Embedding can mean in this context the incorporation of micro-particles in a matrix of polymers. Controlled release

formulations are also obtained through encapsulation of dispersed micro-particles or emulsified micro-droplets via known dispersion or emulsion coating technologies.

[0131] For administration by inhalation, compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant. In the embodiment of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin, for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0132] The compounds disclosed herein can be formulated for parenteral administration by injection (e.g., by bolus injection or continuous infusion). Formulations for injection can be presented in unit dosage form (e.g., in ampules or in multidose containers), with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing, and/or dispersing agents.

[0133] Pharmaceutical formulations for parenteral administration include aqueous solutions of the compounds in water-soluble form. Additionally, suspensions of the compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils or synthetic fatty acid esters. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension. Optionally, the suspension also can contain suitable stabilizers or agents that increase the solubility of the compounds and allow for the preparation of highly concentrated solutions. Alternatively, a present composition can be in powder form for constitution with a suitable vehicle (e.g., sterile pyrogen-free water) before use.

[0134] Compounds disclosed herein also can be formulated in rectal compositions, such as suppositories or retention enemas (e.g., containing conventional suppository bases). In addition to the formulations described previously, the compounds also can be formulated as a depot preparation. Such long-acting formulations can be administered by implantation (e.g., subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0135] In particular, a compound disclosed herein can be administered orally, buccally, or sublingually in the form of tablets containing excipients, such as starch or lactose, or in capsules or ovules, either alone or in admixture with excipients, or in the form of elixirs or suspensions containing flavoring or coloring agents. Such liquid preparations can be prepared with pharmaceutically acceptable additives, such as suspending agents. A compound also can be injected parenterally, for example, intravenously, intramuscularly, subcutaneously, or intracoronarily. For parenteral administration, the compound is best used in the form of a sterile aqueous solution which can contain other substances, for example, salts, or sugar alcohols, such as mannitol, or glucose, to make the solution isotonic with blood.

[0136] For veterinary use, a compound disclosed herein is administered as a suitably acceptable formulation in accordance with normal veterinary practice. The veterinarian can readily determine the dosing regimen and route of administration that is most appropriate for a particular animal.

[0137] In some embodiments, all the necessary components for the treatment of KIF18A-related disorder using a compound as disclosed herein either alone or in combination with another agent or intervention traditionally used for the treatment of such disease may be packaged into a kit. Specifically, the present invention provides a kit for use in the therapeutic intervention of the disease comprising a packaged set of medicaments that include the compound disclosed herein as well as buffers and other components for preparing deliverable forms of said medicaments, and/or devices for delivering such medicaments, and/or any agents that are used in combination therapy with the compound disclosed herein, and/or instructions for the treatment of the disease packaged with the medicaments. The instructions may be fixed in any tangible medium, such as printed paper, or a computer readable magnetic or optical medium, or instructions to reference a remote computer data source such as a world wide web page accessible via the internet.

[0138] A "therapeutically effective amount" means an amount effective to treat or to prevent development of, or to alleviate the existing symptoms of, the subject being treated. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. Generally, a "therapeutically effective dose" refers to that amount of the compound that results in achieving the desired effect. For example, in one preferred embodiment, a therapeutically effective amount of a compound disclosed herein decreases KIF18A activity by at least 5%, compared to control, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, or at least 90%.

[0139] The amount of compound administered can be dependent on the subject being treated, on the subject's age, health, sex, and weight, the kind of concurrent treatment (if any), severity of the affliction, the nature of the effect desired, the manner and frequency of treatment, and the judgment of the prescribing physician. The frequency of dosing also can be dependent on pharmacodynamic effects on arterial oxygen pressures. While individual needs vary, determination of optimal ranges of effective amounts of the compound is within the skill of the art. Such doses may be administered in a single dose or it may be divided into multiple doses.

[0140] The terms "cancer" and "cancerous" when used herein refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include, without limitation, carcinoma, lymphoma, sarcoma, blastoma and leukemia. More particular examples of such cancers include squamous cell carcinoma, lung cancer, pancreatic cancer, cervical cancer, bladder cancer, hepatoma, breast cancer, colon carcinoma, and head and neck cancer, ovarian cancer, and endometrial cancer. While the term "cancer" as used herein is not limited to any one specific form of the disease, it is believed that the methods of the invention will be particularly effective for

cancers which are found to be accompanied by unregulated levels of KIF18A or dependent on KIF18A for proper chromosome segregation and survival in the mammal.

[0141] The terms "treat", "treating" and "treatment" as used herein refer to therapy, including without limitation, curative therapy, prophylactic therapy, and preventative therapy. Prophylactic treatment generally constitutes either preventing the onset of disorders altogether or delaying the onset of a pre-clinically evident stage of disorders in individuals.

[0142] The term "patient", "subject", or "mammal" as used herein refers to any "patient", "subject", or "mammal" including humans, cows, horses, dogs and cats. In one embodiment of the invention, the mammal is a human.

[0143] The term "comprising" is meant to be open ended, including the indicated component(s) but not excluding other elements.

[0144] The terms "Formula I" include any sub formulas, such as (Ia), (Ib), (Ic), (Id), etc.

[0145] METHODS OF USING KIF18A INHIBITORS

[0146] The present disclosure provides compounds having MT-based KIF18A modulatory activity in general, and inhibitory activity in particular. In one embodiment of the invention, there is provided a method of modulating KIF18A protein in a subject, the method comprising administering to the subject an effective dosage amount of a compound of Formulas I. As such, the compounds of the invention may be used to treat cellular proliferation disorders, including uncontrolled cell growth, aberrant cell cycle regulation, centrosome abnormalities (structural and or numeric, fragmentation). Other diseases or disorders associated with the accumulation of extra centrosomes (>2) include human papillomavirus (HPV) infection, including HPV-associated neoplasias. The compounds are also useful for cilia-related diseases as well as ablating haploid germ cell population which could be used as a male contraceptive.

[0147] In addition, compounds of the invention are useful for, but not limited to, the prevention or treatment of cancer and other KIF18A-mediated diseases or disorders. For example, compounds of the invention would be useful for the treatment of various solid and hematologically derived tumors, such as carcinomas, including, without limitation, cancer of the bladder, breast, colon, kidney, liver, lung (including squamous cell and small cell lung cancer), esophagus, gall-bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin (including squamous cell carcinoma); hematopoietic tumors of lymphoid lineage (including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma and Burkett's lymphoma); hematopoietic tumors of myeloid lineage (including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia); tumors of mesenchymal origin (including fibrosarcoma and rhabdomyosarcoma, and other sarcomas, e.g. soft tissue and bone); tumors of the central and peripheral nervous system (including astrocytoma, neuroblastoma, glioma and schwannomas); and other tumors (including melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderma pigmentosum, keratocanthoma, thyroid follicular cancer and Kaposi's sarcoma).

[0148] The compounds of the invention are also useful in the treatment of cancer related indications such as solid tumors, sarcomas (especially Ewing's sarcoma and osteosarcoma), retinoblastoma,

rhabdomyosarcomas, neuroblastoma, hematopoietic malignancies, including leukemia and lymphoma, tumor-induced pleural or pericardial effusions, and malignant ascites.

[0149] Based on the ability to modulate kinesin impacting angiogenesis, the compounds of the invention are also useful in treatment and therapy of proliferative diseases. Particularly, these compounds can be used for the treatment of an inflammatory disease, especially of manifestations at the locomotor apparatus, such as various inflammatory rheumatoid diseases, especially chronic polyarthritis including rheumatoid arthritis, juvenile arthritis or psoriasis arthropathy; paraneoplastic syndrome or tumor-induced inflammatory diseases, turbid effusions, collagenosis, such as systemic Lupus erythematosus, polymyositis, dermatomyositis, systemic scleroderma or mixed collagenosis; postinfectious arthritis (where no living pathogenic organism can be found at or in the affected part of the body), seronegative spondylarthritis, such as spondylitis ankylosans; vasculitis, sarcoidosis, or arthrosis; or further any combinations thereof.

[0150] The compounds of the invention can also be used as active agents against such disease states as arthritis, atherosclerosis, psoriasis, hemangiomas, myocardial angiogenesis, coronary and cerebral collaterals, ischemic limb angiogenesis, wound healing, peptic ulcer Helicobacter related diseases, fractures, cat scratch fever, rubeosis, neovascular glaucoma and retinopathies such as those associated with diabetic retinopathy or macular degeneration. In addition, some of these compounds can be used as active agents against solid tumors, malignant ascites, hematopoietic cancers and hyperproliferative disorders such as thyroid hyperplasia (especially Grave's disease), and cysts (such as hypervascularity of ovarian stroma, characteristic of polycystic ovarian syndrome (Stein-Leventhal syndrome)) since such diseases require a proliferation of blood vessel cells for growth and/or metastasis.

[0151] Besides being useful for human treatment, these compounds are useful for veterinary treatment of companion animals, exotic animals and farm animals, including mammals, rodents, and the like. For example, animals including horses, dogs, and cats may be treated with compounds provided by the invention.

[0152] COMBINATIONS

[0153] While the compounds of the invention can be dosed or administered as the sole active pharmaceutical agent, they can also be used in combination with one or more compounds of the invention or in conjunction with other agents. When administered as a combination, the therapeutic agents can be formulated as separate compositions that are administered simultaneously or sequentially at different times, or the therapeutic agents can be given as a single composition.

[0154] The phrase "co-therapy" (or "combination-therapy"), in defining use of a compound of the present invention and another pharmaceutical agent, is intended to embrace administration of each agent in a sequential manner in a regimen that will provide beneficial effects of the drug combination, and is intended as well to embrace co-administration of these agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of these active agents or in multiple, separate capsules for each agent.

[0155] Specifically, the administration of compounds of the present invention may be in conjunction with additional therapies known to those skilled in the art in the prevention or treatment of cancer, such as with radiation therapy, small molecule targeted agents (e.g. PARP inhibitors, kinase inhibitors), therapeutic antibodies (e.g. naked and drug-conjugate) immunotherapy antibodies (checkpoint inhibitors, bi-specific T-cell engagers) with neoplastic or cytotoxic agents.

[0156] If formulated as a fixed dose, such combination products employ the compounds of this invention within the accepted dosage ranges. Compounds of Formula I may also be administered sequentially with known anticancer or cytotoxic agents when a combination formulation is inappropriate. The invention is not limited in the sequence of administration; compounds of the invention may be administered either prior to, simultaneous with or after administration of the known anticancer or cytotoxic agent.

[0157] There are large numbers of anticancer agents available in commercial use, in clinical evaluation and in pre-clinical development, which would be selected for treatment of neoplasia by combination drug chemotherapy. Such agents fall into several major categories such as antibiotic-type agents, alkylating and alkylating-like agents, antimitotic agents, targeted small molecule agents, antimetabolite agents, hormonal agents, immunological agents, anti-angiogenic agents, interferon-type agents and a category of miscellaneous agents.

[0158] The present disclosure also provides methods for combination therapies in which an agent known to modulate other pathways, or other components of the same pathway, or even overlapping sets of target enzymes are used in combination with a compound of the present disclosure, or a pharmaceutically acceptable salt thereof. In one aspect, such therapy includes but is not limited to the combination of one or more compounds of the disclosure with chemotherapeutic agents, therapeutic antibodies, targeted small molecule agents, and radiation treatment, to provide a synergistic or additive therapeutic effect.

[0159] Many chemotherapeutics are presently known in the art and can be used in combination with the compounds of the disclosure. In some embodiments, the chemotherapeutic is selected from the group consisting of antimitotic agents, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, angiogenesis inhibitors, and anti-androgens. Non-limiting examples are chemotherapeutic agents, cytotoxic agents, and non-peptide small molecules such as Gleevec® (Imatinib Mesylate), Kypolis® (carfilzomib), Velcade® (bortezomib), Casodex (bicalutamide), Iressa® (gefitinib), and Adriamycin as well as a host of chemotherapeutic agents. Non-limiting examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXANTM); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins,

actinomycin, anthramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin, CasodexTM, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo- L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, androgens such as calusterone, dromostanolone propionate, epitio stanol, mepitio stanol, testolactone; anti-adrenals such as aminogluthethimide, mitotane, trilostane; folic acid replenisher such as frolic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diazi quone; elfomithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK; razoxane; sizofiran; spirogermanium; tenuazonic acid; triazi quone; 2,2',2"-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxanes, e.g. paclitaxel and docetaxel, Nab-paclitaxel; retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[0160] Also included as suitable chemotherapeutic cell conditioners are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, (NolvadexTM), raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY 117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin, oxaliplatin, carboplatin; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vinblastine vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; topotecan; camptothecin-11 (CPT-11); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO),

[0161] Where desired, the compounds or pharmaceutical composition of the present disclosure can be used in combination with commonly prescribed anti-cancer drugs such as Herceptin®, Avastin®, Erbitux®, Rituxan®, Taxol®, Abraxane, Arimidex®, Taxotere®, ABVD, AVICINE, Abagovomab, Acridine carboxamide, Adecatumumab, 17-N-Allylamino-17-demethoxygeldanamycin, Alpharadin, Alvocidib, 3-Aminopyridine-2-carboxaldehyde thiosemicarbazone, Amonafide, Anthracenedione, Anti-CD22 immunotoxins, Antineoplastic, Antitumorigenic herbs, Apazi quone, Atiprimod, Azathioprine, Belotecan, Bendamustine, BIBW 2992, Biricodar, Brostallicin, Bryostatin, Buthionine sulfoximine, CBV (chemotherapy), Calyculin, cell-cycle nonspecific antineoplastic agents, Dichloroacetic acid, Discodermolide, Elsamitrucin, Enocitabine, Etophilone, Eribulin, Everolimus, Exatecan, Exisulind, Ferruginol, Forodesine, Fosfestrol, ICE chemotherapy regimen, IT-101, Imexon, Imiquimod,

Indolocarbazole, Irofulven, Laniquidar, Larotaxel, Lenalidomide, Lucanthone, Lurtotecan, Mafosfamide, Mitozolomide, Nafoxidine, Nedaplatin, Olaparib, Talazoparib, Niraparib, Ortataxel, PAC-1, Pawpaw, Pixantrone, Proteasome inhibitor, Rebeccamycin, Resiquimod, Rubitecan, SN-38, Salinosporamide A, Sapacitabine, Stanford V, Swainsonine, Talaporfin, Tariquidar, Tegafur-uracil, Temodar, Tesetaxel, Triplatin tetranitrate, Tris(2-chloroethyl)amine, Troxacitabine, Uramustine, Vadimezan, Vinflunine, ZD6126 or Zosuquidar, CDK4/6 inhibitors (Palbociclib, Ibrance; Ribociclib, Kisqali; Abemaciclib, Verzenio).

[0162] This disclosure further relates to a method for using the compounds or pharmaceutical compositions provided herein, in combination with radiation therapy for inhibiting abnormal cell growth or treating the hyperproliferative disorder in the mammal. Techniques for administering radiation therapy are known in the art, and these techniques can be used in the combination therapy described herein. The administration of the compound of the disclosure in this combination therapy can be determined as described herein.

[0163] Radiation therapy can be administered through one of several methods, or a combination of methods, including without limitation external-beam therapy, internal radiation therapy, implant radiation, stereotactic radiosurgery, systemic radiation therapy, radiotherapy and permanent or temporary interstitial brachytherapy. The term “brachytherapy,” as used herein, refers to radiation therapy delivered by a spatially confined radioactive material inserted into the body at or near a tumor or other proliferative tissue disease site. The term is intended without limitation to include exposure to radioactive isotopes (e.g. At-211, I-131, I-125, Y-90, Re-186, Re-188, Sm-153, Bi-212, P-32, and radioactive isotopes of Lu). Suitable radiation sources for use as a cell conditioner of the present disclosure include both solids and liquids. By way of non-limiting example, the radiation source can be a radionuclide, such as I-125, I-131, Yb-169, Ir-192 as a solid source, I-125 as a solid source, or other radionuclides that emit photons, beta particles, gamma radiation, or other therapeutic rays. The radioactive material can also be a fluid made from any solution of radionuclide(s), e.g., a solution of I-125 or I-131, or a radioactive fluid can be produced using a slurry of a suitable fluid containing small particles of solid radionuclides, such as Au-198, Y-90. Moreover, the radionuclide(s) can be embodied in a gel or radioactive micro spheres.

[0164] The compounds or pharmaceutical compositions of the disclosure can be used in combination with an amount of one or more substances selected from anti-angiogenesis agents, signal transduction inhibitors, antiproliferative agents, glycolysis inhibitors, or autophagy inhibitors.

[0165] Anti-angiogenesis agents, such as MMP-2 (matrix-metalloproteinase 2) inhibitors, MMP-9 (matrix-metalloprotenase 9) inhibitors, and COX-11 (cyclooxygenase 11) inhibitors, can be used in conjunction with a compound of the disclosure and pharmaceutical compositions described herein. Anti-angiogenesis agents include, for example, rapamycin, temsirolimus (CCI-779), everolimus (RAD001), sorafenib, sunitinib, and bevacizumab. Examples of useful COX-II inhibitors include alecoxib, valdecoxib, and rofecoxib. Examples of useful matrix metalloproteinase inhibitors are described in WO 96/33172 WO 96/27583 European Patent Publication EP0818442, European Patent Publication EP1004578, WO 98/07697, WO 98/03516, WO 98/34918, WO 98/34915, WO 98/33768, WO 98/30566,

European Patent Publication 606046, European Patent Publication 931 788, WO 90/05719, WO 99/52910, WO 99/52889, WO 99/29667, WO1999007675, European Patent Publication EP1786785, European Patent Publication No. EP1181017, United States Publication No. US20090012085, United States Publication US5863 949, United States Publication US5861 510, and European Patent Publication EP0780386, all of which are incorporated herein in their entireties by reference. Preferred MMP-2 and MMP-9 inhibitors are those that have little or no activity inhibiting MMP-1. More preferred, are those that selectively inhibit MMP-2 and/or AMP-9 relative to the other matrix- metalloproteinases (i. e., MAP-1, MMP-3, MMP-4, MMP-5, MMP-6, MMP- 7, MMP- 8, MMP-10, MMP-11, MMP-12, and MMP-13). Some specific examples of MMP inhibitors useful in the disclosure are AG-3340, RO 32-3555, and RS 13-0830.

[0166] The present compounds may also be used in co-therapies with other anti-neoplastic agents, such as acemannan, aclarubicin, aldesleukin, alemtuzumab, alitretinoin, altretamine, amifostine, aminolevulinic acid, amrubicin, amsacrine, anagrelide, anastrozole, ANCER, aneastim, ARGLABIN, arsenic trioxide, BAM 002 (Novelos), bexarotene, bicalutamide, broxuridine, capecitabine, celmoleukin, cetorelix, cladribine, clotrimazole, cytarabine ocfosfate, DA 3030 (Dong-A), daclizumab, denileukin diftitox, deslorelin, dexrazoxane, dilazep, docetaxel, docosanol, doxercalciferol, doxifluridine, doxorubicin, bromocriptine, carmustine, cytarabine, fluorouracil, HIT diclofenac, interferon alfa, daunorubicin, doxorubicin, tretinoin, edelfosine, edrecolomab, eflornithine, emitfur, epirubicin, epoetin beta, etoposide phosphate, exemestane, exisulind, fadrozole, filgrastim, finasteride, fludarabine phosphate, formestane, fotemustine, gallium nitrate, gemcitabine, gemtuzumab zogamicin, gimeracil/oteracil/tegafur combination, glycopine, goserelin, heptaplatin, human chorionic gonadotropin, human fetal alpha fetoprotein, ibandronic acid, idarubicin, (imi)quimod, interferon alfa, interferon alfa, natural, interferon alfa-2, interferon alfa-2a, interferon alfa-2b, interferon alfa-N1, interferon alfa-n3, interferon alfacon-1, interferon alpha, natural, interferon beta, interferon beta-1a, interferon beta-1b, interferon gamma, natural interferon gamma-1a, interferon gamma-1b, interleukin-1 beta, iobenguane, irinotecan, irsogladine, lanreotide, LC 9018 (Yakult), leflunomide, lenograstim, lentinan sulfate, letrozole, leukocyte alpha interferon, leuporelin, levamisole + fluorouracil, liarozole, lobaplatin, lonidamine, lovastatin, masoprocol, melarsoprol, metoclopramide, mifepristone, miltefosine, mirimostim, mismatched double stranded RNA, mitoguazone, mitolactol, mitoxantrone, molgramostim, nafarelin, naloxone + pentazocine, nartograstim, nedaplatin, nilutamide, noscapine, novel erythropoiesis stimulating protein, NSC 631570 octreotide, oprelvekin, osaterone, oxaliplatin, paclitaxel, pamidronic acid, pegaspargase, peginterferon alfa-2b, pentosan polysulfate sodium, pentostatin, picibanil, pirarubicin, rabbit antithymocyte polyclonal antibody, polyethylene glycol interferon alfa-2a, porfimer sodium, raloxifene, raltitrexed, rasburiembodiment, rhenium Re 186 etidronate, RII retinamide, rituximab, romurtide, samarium (153 Sm) lexicidronam, sargramostim, sizofiran, sobuzoxane, sonermin, strontium-89 chloride, suramin, tasonermin, tazarotene, tegafur, temoporfin, temozolomide, teniposide, tetrachlorodecaoxide, thalidomide, thymalfasin, thyrotropin alfa, topotecan, toremifene, tositumomab-iodine 131, trastuzumab, treosulfan, tretinoin, trilostane, trimetrexate, triptorelin, tumor necrosis factor alpha, natural, ubenimex, bladder

cancer vaccine, Maruyama vaccine, melanoma lysate vaccine, valrubicin, verteporfin, vinorelbine, VIRULIZIN, zinostatin stimalamer, or zoledronic acid; abarelix; AE 941 (Aeterna), ambamustine, antisense oligonucleotide, bcl-2 (Genta), APC 8015 (Dendreon), cetuximab, decitabine, dexamino-glutethimide, diaziquone, EL 532 (Elan), EM 800 (Endorecherche), eniluracil, etanidazole, fenretinide, filgrastim SD01 (Amgen), fulvestrant, galocitabine, gastrin 17 immunogen, HLA-B7 gene therapy (Vical), granulocyte macrophage colony stimulating factor, histamine dihydrochloride, ibritumomab tiuxetan, ilomastat, IM 862 (Cytran), interleukin-2, iproxifene, LDI 200 (Milkhaus), leridistim, lintuzumab, CA 125 MAb (Biomira), cancer MAb (Japan Pharmaceutical Development), HER-2 and Fc MAb (Medarex), idiotypic 105AD7 MAb (CRC Technology), idiotypic CEA MAb (Trilex), LYM-1-iodine 131 MAb (Techniclone), polymorphic epithelial mucin-tyttrium 90 MAb (Antisoma), marimastat, menogaril, mitumomab, motexafin gadolinium, MX 6 (Galderma), nelarabine, nolatrexed, P 30 protein, pegvisomant, pemetrexed, porfiromycin, prinomastat, RL 0903 (Shire), rubitecan, satraplatin, sodium phenylacetate, sparfosic acid, SRL 172 (SR Pharma), SU 5416 (SUGEN), TA 077 (Tanabe), tetrathiomolybdate, thaliblastine, thrombopoietin, tin ethyl etiopurpurin, tirapazamine, cancer vaccine (Biomira), melanoma vaccine (New York University), melanoma vaccine (Sloan Kettering Institute), melanoma oncolysate vaccine (New York Medical College), viral melanoma cell lysates vaccine (Royal Newcastle Hospital), or valspodar.

[0167] The compounds of the invention may further be used with VEGFR inhibitors. Other compounds described in the following patents and patent applications can be used in combination therapy: US 6,258,812, US 2003/0105091, WO 01/37820, US 6,235,764, WO 01/32651, US 6,630,500, US 6,515,004, US 6,713,485, US 5,521,184, US 5,770,599, US 5,747,498, WO 02/68406, WO 02/66470, WO 02/55501, WO 04/05279, WO 04/07481, WO 04/07458, WO 04/09784, WO 02/59110, WO 99/45009, WO 00/59509, WO 99/61422, US 5,990,141, WO 00/12089, and WO 00/02871.

[0168] In some embodiments, the combination comprises a composition of the present invention in combination with at least one anti-angiogenic agent. Agents are inclusive of, but not limited to, in vitro synthetically prepared chemical compositions, antibodies, antigen binding regions, radionuclides, and combinations and conjugates thereof. An agent can be an agonist, antagonist, allosteric modulator, toxin or, more generally, may act to inhibit or stimulate its target (e.g., receptor or enzyme activation or inhibition), and thereby promote cell death or arrest cell growth.

[0169] Exemplary anti-angiogenic agents include ERBITUX™ (IMC-C225), KDR (kinase domain receptor) inhibitory agents (e.g., antibodies and antigen binding regions that specifically bind to the kinase domain receptor), anti-VEGF agents (e.g., antibodies or antigen binding regions that specifically bind VEGF, or soluble VEGF receptors or a ligand binding region thereof) such as AVASTIN™ or VEGF-TRAP™, and anti-VEGF receptor agents (e.g., antibodies or antigen binding regions that specifically bind thereto), EGFR inhibitory agents (e.g., antibodies or antigen binding regions that specifically bind thereto) such as Vectibix (panitumumab), IRESSA™ (gefitinib), TARCEVA™ (erlotinib), anti-Ang1 and anti-Ang2 agents (e.g., antibodies or antigen binding regions specifically binding thereto or to their receptors, e.g., Tie2/Tek), and anti-Tie2 kinase inhibitory agents (e.g., antibodies or antigen binding regions that

specifically bind thereto). The pharmaceutical compositions of the present invention can also include one or more agents (e.g., antibodies, antigen binding regions, or soluble receptors) that specifically bind and inhibit the activity of growth factors, such as antagonists of hepatocyte growth factor (HGF, also known as Scatter Factor), and antibodies or antigen binding regions that specifically bind its receptor "c-met".

[0170] Other anti-angiogenic agents include Campath, IL-8, B-FGF, Tek antagonists (Ceretti et al., U.S. Publication No. 2003/0162712; U.S. Patent No. 6,413,932), anti-TWEAK agents (e.g., specifically binding antibodies or antigen binding regions, or soluble TWEAK receptor antagonists; see, Wiley, U.S. Patent No. 6,727,225), ADAM distintegrin domain to antagonize the binding of integrin to its ligands (Fanslow et al., U.S. Publication No. 2002/0042368), specifically binding anti-ephrin receptor and/or anti-ephrin antibodies or antigen binding regions (U.S. Patent Nos. 5,981,245; 5,728,813; 5,969,110; 6,596,852; 6,232,447; 6,057,124 and patent family members thereof), and anti-PDGF-BB antagonists (e.g., specifically binding antibodies or antigen binding regions) as well as antibodies or antigen binding regions specifically binding to PDGF-BB ligands, and PDGFR kinase inhibitory agents (e.g., antibodies or antigen binding regions that specifically bind thereto).

[0171] Additional anti-angiogenic/anti-tumor agents include: SD-7784 (Pfizer, USA); cilengitide (Merck KGaA, Germany, EPO 770622); pegaptanib octasodium, (Gilead Sciences, USA); Alphastatin, (BioActa, UK); M-PGA, (Celgene, USA, US 5712291); ilomastat, (Arriva, USA, US 5892112); emaxanib, (Pfizer, USA, US 5792783); vatalanib, (Novartis, Switzerland); 2-methoxyestradiol, (EntreMed, USA); TLC ELL-12, (Elan, Ireland); anecortave acetate, (Alcon, USA); alpha-D148 Mab, (Amgen, USA); CEP-7055, (Cephalon, USA); anti-Vn Mab, (Crucell, Netherlands) DAC:antiangiogenic, (ConjuChem, Canada); Angiocidin, (InKine Pharmaceutical, USA); KM-2550, (Kyowa Hakko, Japan); SU-0879, (Pfizer, USA); CGP-79787, (Novartis, Switzerland, EP 970070); ARGENT technology, (Ariad, USA); YIGSR-Stealth, (Johnson & Johnson, USA); fibrinogen-E fragment, (BioActa, UK); angiogenesis inhibitor, (Trigen, UK); TBC-1635, (Encysive Pharmaceuticals, USA); SC-236, (Pfizer, USA); ABT-567, (Abbott, USA); Metastatin, (EntreMed, USA); angiogenesis inhibitor, (Tripep, Sweden); maspin, (Sosei, Japan); 2-methoxyestradiol, (Oncology Sciences Corporation, USA); ER-68203-00, (IVAX, USA); Benefin, (Lane Labs, USA); Tz-93, (Tsumura, Japan); TAN-1120, (Takeda, Japan); FR-111142, (Fujisawa, Japan, JP 02233610); platelet factor 4, (RepliGen, USA, EP 407122); vascular endothelial growth factor antagonist, (Boreon, Denmark); bevacizumab (pINN), (Genentech, USA); angiogenesis inhibitors, (SUGEN, USA); XL 784, (Exelixis, USA); XL 647, (Exelixis, USA); MAb, alpha5beta3 integrin, second generation, (Applied Molecular Evolution, USA and MedImmune, USA); gene therapy, retinopathy, (Oxford BioMedica, UK); enzastaurin hydrochloride (USAN), (Lilly, USA); CEP 7055, (Cephalon, USA and Sanofi-Synthelabo, France); BC 1, (Genoa Institute of Cancer Research, Italy); angiogenesis inhibitor, (Alchemia, Australia); VEGF antagonist, (Regeneron, USA); rBPI 21 and BPI-derived antiangiogenic, (XOMA, USA); PI 88, (Progen, Australia); cilengitide (pINN), (Merck KGaA, German; Munich Technical University, Germany, Scripps Clinic and Research Foundation, USA); cetuximab (INN), (Aventis, France); AVE 8062, (Ajinomoto, Japan); AS 1404, (Cancer Research Laboratory, New Zealand); SG 292, (Telios, USA); Endostatin, (Boston Childrens Hospital, USA); ATN 161, (Attenuon,

USA); ANGIOSTATIN, (Boston Childrens Hospital, USA); 2-methoxyestradiol, (Boston Childrens Hospital, USA); ZD 6474, (AstraZeneca, UK); ZD 6126, (Angiogene Pharmaceuticals, UK); PPI 2458, (Praecis, USA); AZD 9935, (AstraZeneca, UK); AZD 2171, (AstraZeneca, UK); vatalanib (pINN), (Novartis, Switzerland and Schering AG, Germany); tissue factor pathway inhibitors, (EntreMed, USA); pegaptanib (Pinn), (Gilead Sciences, USA); xanthorrhizol, (Yonsei University, South Korea); vaccine, gene-based, VEGF-2, (Scripps Clinic and Research Foundation, USA); SPV5.2, (Supratek, Canada); SDX 103, (University of California at San Diego, USA); PX 478, (ProlX, USA); METASTATIN, (EntreMed, USA); troponin I, (Harvard University, USA); SU 6668, (SUGEN, USA); OXI 4503, (OXiGENE, USA); o-guanidines, (Dimensional Pharmaceuticals, USA); motuporamine C, (British Columbia University, Canada); CDP 791, (Celltech Group, UK); atiprimod (pINN), (GlaxoSmithKline, UK); E 7820, (Eisai, Japan); CYC 381, (Harvard University, USA); AE 941, (Aeterna, Canada); vaccine, angiogenesis, (EntreMed, USA); urokinase plasminogen activator inhibitor, (Dendreon, USA); oglufanide (pINN), (Melmotte, USA); HIF-1alfa inhibitors, (Xenova, UK); CEP 5214, (Cephalon, USA); BAY RES 2622, (Bayer, Germany); Angiocidin, (InKine, USA); A6, (Angstrom, USA); KR 31372, (Korea Research Institute of Chemical Technology, South Korea); GW 2286, (GlaxoSmithKline, UK); EHT 0101, (ExonHit, France); CP 868596, (Pfizer, USA); CP 564959, (OSI, USA); CP 547632, (Pfizer, USA); 786034, (GlaxoSmithKline, UK); KRN 633, (Kirin Brewery, Japan); drug delivery system, intraocular, 2-methoxyestradiol, (EntreMed, USA); angonex, (Maastricht University, Netherlands, and Minnesota University, USA); ABT 510, (Abbott, USA); AAL 993, (Novartis, Switzerland); VEGI, (ProteomTech, USA); tumor necrosis factor-alpha inhibitors, (National Institute on Aging, USA); SU 11248, (Pfizer, USA and SUGEN USA); ABT 518, (Abbott, USA); YH16, (Yantai Rongchang, China); S-3APG , (Boston Childrens Hospital, USA and EntreMed, USA); MAb, KDR, (ImClone Systems, USA); MAb, alpha5 beta1, (Protein Design, USA); KDR kinase inhibitor, (Celltech Group, UK, and Johnson & Johnson, USA); GFB 116, (South Florida University, USA and Yale University, USA); CS 706, (Sankyo, Japan); combretastatin A4 prodrug, (Arizona State University, USA); chondroitinase AC, (IBEX, Canada); BAY RES 2690, (Bayer, Germany); AGM 1470, (Harvard University, USA, Takeda, Japan, and TAP, USA); AG 13925, (Agouron, USA); Tetrathiomolybdate, (University of Michigan, USA); GCS 100, (Wayne State University, USA) CV 247, (Ivy Medical, UK); CKD 732, (Chong Kun Dang, South Korea); MAb, vascular endothelium growth factor, (Xenova, UK); irsogladine (INN), (Nippon Shinyaku, Japan); RG 13577, (Aventis, France); WX 360, (Wilex, Germany); squalamine (pINN), (Genaera, USA); RPI 4610, (Sirna, USA); cancer therapy, (Marinova, Australia); heparanase inhibitors, (InSight, Israel); KL 3106, (Kolon, South Korea); Honokiol, (Emory University, USA); ZK CDK, (Schering AG, Germany); ZK Angio, (Schering AG, Germany); ZK 229561, (Novartis, Switzerland, and Schering AG, Germany); XMP 300, (XOMA, USA); VGA 1102, (Taisho, Japan); VEGF receptor modulators, (Pharmacopeia, USA); VE-cadherin-2 antagonists , (ImClone Systems, USA); Vasostatin, (National Institutes of Health, USA);vaccine, Flk-1, (ImClone Systems, USA); TZ 93, (Tsumura, Japan); TumStatin, (Beth Israel Hospital, USA); truncated soluble FLT 1 (vascular endothelial growth factor receptor 1), (Merck & Co,

USA); Tie-2 ligands, (Regeneron, USA); and, thrombospondin 1 inhibitor, (Allegheny Health, Education and Research Foundation, USA).

[0172] Autophagy inhibitors include, but are not limited to chloroquine, 3- methyladenine, hydroxychloroquine (Plaquenil™), bafilomycin A1, 5-amino-4- imidazole carboxamide riboside (AICAR), okadaic acid, autophagy-suppressive algal toxins which inhibit protein phosphatases of type 2A or type 1, analogues of cAMP, and drugs which elevate cAMP levels such as adenosine, LY204002, N6- mercaptopurine riboside, and vinblastine. In addition, antisense or siRNA that inhibits expression of proteins including but not limited to ATG5 (which are implicated in autophagy), may also be used.

[0173] Additional pharmaceutically active compounds/agents that can be used in the treatment of cancers and that can be used in combination with one or more compound of the present invention include: epoetin alfa; darbepoetin alfa; panitumumab; pegfilgrastim; palifermin; filgrastim; denosumab; ancestim; AMG 102; AMG 386; AMG 479; AMG 655; AMG 745; AMG 951; and AMG 706, or a pharmaceutically acceptable salt thereof.

[0174] In certain embodiments, a composition provided herein is conjointly administered with a chemotherapeutic agent . Suitable chemotherapeutic agents may include, natural products such as vinca alkaloids (e.g., vinblastine, vincristine, and vinorelbine), paclitaxel, epidipodophyllotoxins (e.g., etoposide and teniposide), antibiotics (e.g., dactinomycin (actinomycin D), daunorubicin, doxorubicin, and idarubicin), anthracyclines, mitoxantrone, bleomycins, plicamycin (mithramycin), mitomycin, enzymes (e.g., L-asparaginase which systemically metabolizes L-asparagine and deprives cells which do not have the capacity to synthesize their own asparagine), antiplatelet agents, antiproliferative/antimitotic alkylating agents such as nitrogen mustards (e.g., mechlorethamine, cyclophosphamide and analogs, melphalan, and chlorambucil), ethylenimines and methylmelamines (e.g., hexaamethylmelamine and thiotepa), CDK inhibitors (e.g., seliciclib, UCN-01, P1446A-05, PD-0332991, dinaciclib, P27-00, AT-7519, RGB286638, and SCH727965), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine (BCNU) and analogs, and streptozocin), trazenes-dacarbazine (DTIC), antiproliferative/antimitotic antimetabolites such as folic acid analogs (e.g., methotrexate), pyrimidine analogs (e.g., fluorouracil, floxuridine, and cytarabine), purine analogs and related inhibitors (e.g., mercaptopurine, thioguanine, pentostatin and 2-chlorodeoxyadenosine), aromatase inhibitors (e.g., anastrozole, exemestane, and letrozole), and platinum coordination complexes (e.g., cisplatin and carboplatin), procarbazine, hydroxyurea, mitotane, aminoglutethimide, histone deacetylase (HDAC) inhibitors (e.g., trichostatin, sodium butyrate, apicidan, suberoyl anilide hydroamic acid, vorinostat, LBH 589, romidepsin, ACY-1215, and panobinostat), mTor inhibitors (e.g., temsirolimus, everolimus, ridaforolimus, and sirolimus), KSP(Eg5) inhibitors (e.g., Array 520), DNA binding agents (e.g., Zalypsis), PI3K delta inhibitor (e.g., GS-1101 and TGR-1202), PI3K delta and gamma inhibitor (e.g., CAL-130), multi-kinase inhibitor (e.g., TG02 and sorafenib), hormones (e.g., estrogen) and hormone agonists such as leutinizing hormone releasing hormone (LHRH) agonists (e.g., goserelin, leuprolide and triptorelin), BAFF-neutralizing antibody (e.g., LY2127399), IKK inhibitors, p38MAPK inhibitors, anti-IL-6 (e.g., CNT0328), telomerase inhibitors (e.g., GRN 163L), aurora kinase inhibitors (e.g., MLN8237, AMG 900, AZD-1152), cell

surface monoclonal antibodies (e.g., anti-CD38 (HUMAX-CD38), anti-CS1 (e.g., elotuzumab), HSP90 inhibitors (e.g., 17 AAG and KOS 953), P13K / Akt inhibitors (e.g., perifosine), Akt inhibitor (e.g., GSK-2141795), PKC inhibitors (e.g., enzastaurin), FTIs (e.g., Zarnestra™), anti-CD138 (e.g., BT062), Torc1/2 specific kinase inhibitor (e.g., INK128), kinase inhibitor (e.g., GS-1101), ER/UPR targeting agent (e.g., MKC-3946), cFMS inhibitor (e.g., ARRY-382), JAK1/2 inhibitor (e.g., CYT387), PARP inhibitor (e.g., olaparib, Talazoparib, Niraparib veliparib (ABT-888)), BCL-2 antagonist. Other chemotherapeutic agents may include mechlorethamine, camptothecin, ifosfamide, tamoxifen, raloxifene, gemcitabine, navelbine, sorafenib, or any analog or derivative variant of the foregoing.

[0175] The compounds of the present invention may also be used in combination with radiation therapy, hormone therapy, surgery and immunotherapy, which therapies are well known to those skilled in the art.

[0176] In certain embodiments, a pharmaceutical composition provided herein is conjointly administered with a steroid. Suitable steroids may include, but are not limited to, 21-acetoxypregnenolone, alclometasone, algestone, amcinonide, beclomethasone, betamethasone, budesonide, chloroprednisone, clobetasol, clocortolone, cloprednol, corticosterone, cortisone, cortivazol, deflazacort, desonide, desoximetasone, dexamethasone, diflorasone, diflucortolone, difuprednate, enoxolone, fluazacort, flucoronide, flumethasone, flunisolide, fluocinolone acetonide, fluocinonide, fluocortin butyl, fluocortolone, fluorometholone, fluperolone acetate, fluprednidene acetate, fluprednisolone, flurandrenolide, fluticasone propionate, formocortal, halcinonide, halobetasol propionate, halometasone, hydrocortisone, loteprednol etabonate, mazipredone, medrysone, meprednisone, methylprednisolone, mometasone furoate, paramethasone, prednicarbate, prednisolone, prednisolone 25-diethylaminoacetate, prednisolone sodium phosphate, prednisone, prednival, prednylidene, rimexolone, tixocortol, triamcinolone, triamcinolone acetonide, triamcinolone benetonide, triamcinolone hexacetonide, and salts and/or derivatives thereof. In a particular embodiment, the compounds of the present invention can also be used in combination with additional pharmaceutically active agents that treat nausea. Examples of agents that can be used to treat nausea include: dronabinol; granisetron; metoclopramide; ondansetron; and prochlorperazine; or a pharmaceutically acceptable salt thereof.

[0177] The compounds or pharmaceutical compositions of the disclosure can also be used in combination with an amount of one or more substances selected from EGFR inhibitors, MEK inhibitors, PI3K inhibitors, AKT inhibitors, TOR inhibitors, and immune therapies, including anti-PD-1, anti-PDL-1, anti-CTLA4, anti-LAG1, and anti-OX40 agents, GITR agonists, CAR-T cells, and BiTEs.

[0178] EGFR inhibitors include, but are not limited to, small molecule antagonists, antibody inhibitors, or specific antisense nucleotide or siRNA. Useful antibody inhibitors of EGFR include cetuximab (Erbix), panitumumab (Vectibix), zalutumumab, nimotuzumab, and matuzumab. Small molecule antagonists of EGFR include gefitinib, erlotinib (Tarceva), and most recently, lapatinib (TykerB). See e.g., Yan L, et. al., Pharmacogenetics and Pharmacogenomics In Oncology Therapeutic Antibody Development, BioTechniques 2005; 39(4): 565-8, and Paez J G, et. al., EGFR Mutations In Lung Cancer Correlation With Clinical Response To Gefitinib Therapy, Science 2004; 304(5676): 1497-500.

[0179] Non-limiting examples of small molecule EGFR inhibitors include any of the EGFR inhibitors described in the following patent publications, and all pharmaceutically acceptable salts and solvates of said EGFR inhibitors: European Patent Application EP 520722, published Dec. 30, 1992; European Patent Application EP 566226, published Oct. 20, 1993; PCT International Publication WO 96/33980, published Oct. 31, 1996; U.S. Pat. No. 5,747,498, issued May 5, 1998; PCT International Publication WO 96/30347, published Oct. 3, 1996; European Patent Application EP 787772, published Aug. 6, 1997; PCT International Publication WO 97/30034, published Aug. 21, 1997; PCT International Publication WO 97/30044, published Aug. 21, 1997; PCT International Publication WO 97/38994, published Oct. 23, 1997; PCT International Publication WO 97/49688, published Dec. 31, 1997; European Patent Application EP 837063, published Apr. 22, 1998; PCT International Publication WO 98/02434, published Jan. 22, 1998; PCT International Publication WO 97/38983, published Oct. 23, 1997; PCT International Publication WO 95/19774, published Jul. 27, 1995; PCT International Publication WO 95/19970, published Jul. 27, 1995; PCT International Publication WO 97/13771, published Apr. 17, 1997; PCT International Publication WO 98/02437, published Jan. 22, 1998; PCT International Publication WO 98/02438, published Jan. 22, 1998; PCT International Publication WO 97/32881, published Sep. 12, 1997; German Application DE 19629652, published Jan. 29, 1998; PCT International Publication WO 98/33798, published Aug. 6, 1998; PCT International Publication WO 97/32880, published Sep. 12, 1997; PCT International Publication WO 97/32880 published Sep. 12, 1997; European Patent Application EP 682027, published Nov. 15, 1995; PCT International Publication WO 97/02266, published Jan. 23, 1997; PCT International Publication WO 97/27199, published Jul. 31, 1997; PCT International Publication WO 98/07726, published Feb. 26, 1998; PCT International Publication WO 97/34895, published Sep. 25, 1997; PCT International Publication WO 96/31510', published Oct. 10, 1996; PCT International Publication WO 98/14449, published Apr. 9, 1998; PCT International Publication WO 98/14450, published Apr. 9, 1998; PCT International Publication WO 98/14451, published Apr. 9, 1998; PCT International Publication WO 95/09847, published Apr. 13, 1995; PCT International Publication WO 97/19065, published May 29, 1997; PCT International Publication WO 98/17662, published Apr. 30, 1998; U.S. Pat. No. 5,789,427, issued Aug. 4, 1998; U.S. Pat. No. 5,650,415, issued Jul. 22, 1997; U.S. Pat. No. 5,656,643, issued Aug. 12, 1997; PCT International Publication WO 99/35146, published Jul. 15, 1999; PCT International Publication WO 99/35132, published Jul. 15, 1999; PCT International Publication WO 99/07701, published Feb. 18, 1999; and PCT International Publication WO 92/20642 published Nov. 26, 1992. Additional non-limiting examples of small molecule EGFR inhibitors include any of the EGFR inhibitors described in Traxler, P., 1998, Exp. Opin. Ther. Patents 8(12):1599-1625.

[0180] Antibody-based EGFR inhibitors include any anti-EGFR antibody or antibody fragment that can partially or completely block EGFR activation by its natural ligand. Non-limiting examples of antibody-based EGFR inhibitors include those described in Modjtahedi, H., et al., 1993, Br. J. Cancer 67:247-253; Teramoto, T., et al., 1996, Cancer 77:639-645; Goldstein et al., 1995, Clin. Cancer Res. 1:1311-1318; Huang, S. M., et al., 1999, Cancer Res. 59:1236-1243. Thus, the EGFR inhibitor can be monoclonal antibody Mab E7.6.3 (Yang, 1999 supra), or

Mab C225 (ATCC Accession No. HB-8508), or an antibody or antibody fragment having the binding specificity thereof.

[0181] MEK inhibitors include, but are not limited to, CI-1040, AZD6244, PD318088, PD98059, PD334581, RDEA119, ARRY-142886, ARRY-438162, and PD-325901.

[0182] PI3K inhibitors include, but are not limited to, wortmannin, 17-hydroxywortmannin analogs described in WO 06/044453, 4-[2-(1H-Indazol-4-yl)-6-[[4-(methylsulfonyl)piperazin-1-yl]methyl]thieno[3,2-d]pyrimidin-4-yl]morpholine (also known as GDC 0941 and described in PCT Publication Nos. WO 09/036,082 and WO 09/055,730), 2-Methyl-2-[4-[3-methyl-2-oxo-8-(quinolin-3-yl)-2,3-dihydroimidazo[4,5-c]quinolin-1-yl]phenyl]propionitrile (also known as BEZ 235 or NVP-BEZ 235, and described in PCT Publication No. WO 06/122806), (S)-1-(4-((2-(2-aminopyrimidin-5-yl)-7-methyl-4-morpholinothieno[3,2-d]pyrimidin-6-yl)methyl)piperazin-1-yl)-2-hydroxypropan-1-one (described in PCT Publication No. WO 2008/070740), LY294002 (2-(4-Morpholinyl)-8-phenyl-4H-1-benzopyran-4-one available from Axon Medchem), PI 103 hydrochloride (3-[4-(4-morpholinyl)pyrido-[3',2':4,5]furo[3,2-d]pyrimidin-2-yl]phenol hydrochloride available from Axon Medchem), PIK 75 (N'-[(1E)-(6-bromoimidazo[1,2-a]pyridin-3-yl)methylene]-N,2-dimethyl-5-nitrobenzenesulfono-hydrazide hydrochloride available from Axon Medchem), PIK 90 (N-(7,8-dimethoxy-2,3-dihydro-imidazo[1,2-c]quinazolin-5-yl)-nicotinamide available from Axon Medchem), GDC-0941 bismesylate (2-(1H-Indazol-4-yl)-6-(4-methanesulfonyl-piperazin-1-ylmethyl)-4-morpholin-4-yl-thieno[3,2-d]pyrimidine bismesylate available from Axon Medchem), AS-252424 (5-[1-[5-(4-Fluoro-2-hydroxy-phenyl)-furan-2-yl]-meth-(Z)-ylidene]-thiazolidine-2,4-dione available from Axon Medchem), and TGX-221 (7-Methyl-2-(4-morpholinyl)-9-[1-(phenylamino)ethyl]-4H-pyrido-[1,2-a]pyrimidin-4-one available from Axon Medchem), XL-765, and XL-147. Other PI3K inhibitors include demethoxyviridin, perifosine, CAL101, PX-866, BEZ235, SF1126, INK1117, IPI-145, BKM120, XL147, XL765, Palomid 529, GSK1059615, ZSTK474, PWT33597, IC87114, TG100-115, CAL263, PI-103, GNE-477, CUDC-907, and AEZS-136.

[0183] AKT inhibitors include, but are not limited to, Akt-1-1 (inhibits Akt1) (Barnett et al. (2005) Biochem. J., 385 (Pt. 2), 399-408); Akt-1-1,2 (inhibits Akt1 and 2) (Barnett et al. (2005) Biochem. J. 385 (Pt. 2), 399-408); API-59CJ-Ome (e.g., Jin et al. (2004) Br. J. Cancer 91, 1808-12); 1-H-imidazo[4,5-c]pyridinyl compounds (e.g., WO05011700); indole-3-carbinol and derivatives thereof (e.g., U.S. Pat. No. 6,656,963; Sarkar and Li (2004) J Nutr. 134(12 Suppl), 3493S-3498S); perifosine (e.g., interferes with Akt membrane localization; Dasmahapatra et al. (2004) Clin. Cancer Res. 10(15), 5242-52, 2004); phosphatidylinositol ether lipid analogues (e.g., Gills and Dennis (2004) Expert. Opin. Investig. Drugs 13, 787-97); and triciribine (TCN or API-2 or NCI identifier: NSC 154020; Yang et al. (2004) Cancer Res. 64, 4394-9).

[0184] TOR inhibitors include, but are not limited to, inhibitors include AP-23573, CCI-779, everolimus, RAD-001, rapamycin, temsirolimus, ATP-competitive TORC1/TORC2 inhibitors, including PI-103, PP242, PP30 and Torin 1. Other TOR inhibitors in FKBP12 enhancer; rapamycins and derivatives thereof, including: CCI-779 (temsirolimus), RAD001 (Everolimus; WO 9409010) and AP23573; rapalogs, e.g. as disclosed in WO 98/02441 and WO 01/14387, e.g. AP23573, AP23464, or AP23841; 40-

(2-hydroxyethyl)rapamycin, 40-[3-hydroxy(hydroxymethyl)methylpropanoate]-rapamycin (also called CC1779), 40-epi-(tetrazolyt)-rapamycin (also called ABT578), 32-deoxorapamycin, 16-pentynyloxy-32(S)-dihydrorapamycin, and other derivatives disclosed in WO 05005434; derivatives disclosed in U.S. Pat. No. 5,258,389, WO 94/090101, WO 92/05179, U.S. Pat. No. 5,118,677, U.S. Pat. No. 5,118,678, U.S. Pat. No. 5,100,883, U.S. Pat. No. 5,151,413, U.S. Pat. No. 5,120,842, WO 93/111130, WO 94/02136, WO 94/02485, WO 95/14023, WO 94/02136, WO 95/16691, WO 96/41807, WO 96/41807 and U.S. Pat. No. 5,256,790; phosphorus-containing rapamycin derivatives (e.g., WO 05016252); 4H-1-benzopyran-4-one derivatives (e.g., U.S. Provisional Application No. 60/528,340).

[0185] Immune therapies include, but are not limited to, anti-PD-1 agents, anti-PDL-1 agents, anti-CTLA-4 agents, anti-LAG1 agents, and anti-OX40 agents. Exemplary anti-PD-1 antibodies and methods for their use are described by Goldberg et al., Blood 110(1):186-192 (2007), Thompson et al., Clin. Cancer Res. 13(6):1757-1761 (2007), and Korman et al., International Application No. PCT/JP2006/309606 (publication no. WO 2006/121168 A1), each of which are expressly incorporated by reference herein. include: Yervoy™ (ipilimumab) or Tremelimumab (to CTLA-4), galiximab (to B7.1), BMS-936558 (to PD-1), MK-3475 (to PD-1), AMP224 (to B7DC), BMS-936559 (to B7-H1), MPDL3280A (to B7-H1), MEDI-570 (to ICOS), AMG557 (to B7H2), MGA271 (to B7H3), IMP321 (to LAG-3), BMS-663513 (to CD137), PF-05082566 (to CD137), CDX-1127 (to CD27), anti-OX40 (Providence Health Services), huMAbOX40L (to OX40L), Atacicept (to TACI), CP-870893 (to CD40), Lucatumumab (to CD40), Dacetuzumab (to CD40), Muromonab-CD3 (to CD3), Ipilimumab (to CTLA-4). Immune therapies also include genetically engineered T-cells (e.g., CAR-T cells) and bispecific antibodies (e.g., BiTEs).

[0186] GITR agonists include, but are not limited to, GITR fusion proteins and anti-GITR antibodies (e.g., bivalent anti-GITR antibodies), such as, a GITR fusion protein described in U.S. Pat. No. 6,111,090box.c, European Patent No.: 090505B1, U.S. Pat. No. 8,586,023, PCT Publication Nos.: WO 2010/003118 and 2011/090754, or an anti-GITR antibody described, e.g., in U.S. Pat. No. 7,025,962, European Patent No.: 1947183B1, U.S. Pat. No. 7,812,135, U.S. Pat. No. 8,388,967, U.S. Pat. No. 8,591,886, European Patent No.: EP 1866339, PCT Publication No.: WO 2011/028683, PCT Publication No.: WO 2013/039954, PCT Publication No.: WO2005/007190, PCT Publication No.: WO 2007/133822, PCT Publication No.: WO2005/055808, PCT Publication No.: WO 99/40196, PCT Publication No.: WO 2001/03720, PCT Publication No.: WO99/20758, PCT Publication No.: WO2006/083289, PCT Publication No.: WO 2005/115451, U.S. Pat. No. 7,618,632, and PCT Publication No.: WO 2011/051726.

[0187] The compounds described herein can be used in combination with the agents disclosed herein or other suitable agents, depending on the condition being treated. Hence, in some embodiments the one or more compounds of the disclosure will be co-administered with other agents as described above. When used in combination therapy, the compounds described herein are administered with the second agent simultaneously or separately. This administration in combination can include simultaneous administration of the two agents in the same dosage form, simultaneous administration in separate dosage forms, and separate administration. That is, a compound described herein and any of the agents described above can

be formulated together in the same dosage form and administered simultaneously. Alternatively, a compound of the disclosure and any of the agents described above can be simultaneously administered, wherein both the agents are present in separate formulations. In another alternative, a compound of the present disclosure can be administered just followed by and any of the agents described above, or vice versa. In some embodiments of the separate administration protocol, a compound of the disclosure and any of the agents described above are administered a few minutes apart, or a few hours apart, or a few days apart.

[0188] As one aspect of the present invention contemplates the treatment of the disease/conditions with a combination of pharmaceutically active compounds that may be administered separately, the invention further relates to combining separate pharmaceutical compositions in kit form. The kit comprises two separate pharmaceutical compositions: a compound of the present invention, and a second pharmaceutical compound. The kit comprises a container for containing the separate compositions such as a divided bottle or a divided foil packet. Additional examples of containers include syringes, boxes, and bags. In some embodiments, the kit comprises directions for the use of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral and parenteral), are administered at different dosage intervals, or when titration of the individual components of the combination is desired by the prescribing health care professional.

EXPERIMENTAL

[0189] Abbreviations: The following abbreviations may be used herein:

ACN or MeCN	Acetonitrile
AcOH	acetic acid
aq or aq.	Aqueous
BOC or Boc	<i>tert</i> -butoxycarbonyl
cataCXium® A	rac-((3 <i>R</i> ,5 <i>R</i> ,7 <i>R</i>)-adamantan-1-yl)((3 <i>S</i> ,5 <i>S</i> ,7 <i>S</i>)-adamantan-1-yl)(butyl)phosphane
DABSO	1,4-diazabicyclo[2.2.2]octane bis(sulfur dioxide) adduct
DCE	1,2-dichloroethane
DCM	Dichloromethane
DMAP	4-dimethylaminopyridine
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethyl sulfoxide
Dppf, DPPF or dppf	1,1'-bis(diphenylphosphino)ferrocene
dppp	1,3-bis(diphenylphosphino)propane
eq or eq. or equiv.	Equivalent
ESI or ES	electrospray ionization
Et	Ethyl
Et ₂ O	diethyl ether
EtOH	ethyl alcohol
EtOAc	ethyl acetate
g	Grams
h	Hour
HATU	1-[bis(dimethylamino)methylene]-1 <i>H</i> -1,2,3-triazolo[4,5- <i>b</i>]pyridinium 3-oxid hexafluorophosphate
HPLC	high pressure liquid chromatography
IPA	Isopropanol
iPr	Isopropyl

iPr ₂ NEt or DIPEA	<i>N</i> -ethyl diisopropylamine (Hünig's base)
KOAc	potassium acetate
LDA	lithium diisopropylamide
LC MS, LCMS, LC-MS or LC/MS	liquid chromatography mass spectroscopy
LG	leaving group (e.g., halogen, mesylate, triflate)
m/z	mass divided by charge
Me	Methyl
MeOH	Methanol
Met	metal species for cross-coupling (e.g., MgX, ZnX, SnR ₃ , SiR ₃ , B(OR) ₂)
mg	milligrams
min	minutes
mL	milliliters
MS	mass spectra
MsCl	methanesulfonyl chloride
MTBE	<i>tert</i> -butyl methyl ether
NMP	1-methyl-2-pyrrolidine
<i>n</i> -BuLi	<i>N</i> -butyllithium
NMR	nuclear magnetic resonance
Pd ₂ (dba) ₃	tris(dibenzylideneacetone)dipalladium(0)
Pd(dppf)Cl ₂ ·DCM	[1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane
Pd(PPh ₃) ₄	tetrakis(triphenylphosphine)palladium(0)
Ph	phenyl
Phen	1,10-phenanthroline
PR or PG or Prot. group	protecting group
rbf	round-bottom flask
RP-HPLC	reverse phase high pressure liquid chromatography
RT or rt	room temperature
sat. or satd.	saturated
SFC	supercritical fluid chromatography
SPhos Pd G3 or SPhos G3	(2-Dicyclohexylphosphino-2',6'-dimethoxybiphenyl) [2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate
TBAF	tetra- <i>n</i> -butylammonium fluoride
<i>t</i> -BuOH	<i>tert</i> -butanol
TEA or Et ₃ N	trimethylamine
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TLC	thin layer chromatography
T ₃ P	2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide
UV	ultraviolet
Xantphos	4,5-bis(diphenylphosphino)-9,9-dimethylxanthene
XtalFluor-M	difluoro(morpholino)sulfonium tetrafluoroborate


[0190] Unless otherwise noted, all materials were obtained from commercial suppliers and used without further purification. All parts are by weight and temperatures are in degrees centigrade unless otherwise indicated. All microwave assisted reactions were conducted with a Smith Synthesizer™ from Biotage™. All compounds showed NMR spectra consistent with their assigned structures. Melting points were determined on a Buchi apparatus and are uncorrected. Mass spectral data was determined by electrospray ionization technique. All examples were purified to >90% purity as determined by high-performance liquid chromatography. Unless otherwise stated, reactions were run at room temperature.

[0191] In synthesizing compounds of the present invention, it may be desirable to use certain leaving groups. The term “leaving groups” (“LG”) generally refer to groups that are displaceable by a nucleophile. Such leaving groups are known in the art. Examples of leaving groups include, but are not limited to, halides (e.g., I, Br, F, Cl), sulfonates (e.g., mesylate, tosylate), sulfides (e.g., SCH₃), N-hydroxysuccinimide, N-hydroxybenzotriazole, and the like. Examples of nucleophiles include, but are not limited to, amines, thiols, alcohols, Grignard reagents, anionic species (e.g., alkoxides, amides, carbanions) and the like.

[0192] The examples presented below illustrate specific embodiments of the present invention. These examples are meant to be representative and are not intended to limit the scope of the claims in any manner.

[0193] It is noted that when a percent (%) is used with regard to a liquid, it is a percent by volume with respect to the solution. When used with a solid, it is the percent with regard to the solid composition. Materials obtained from commercial suppliers were typically used without further purification. Reactions involving air or moisture sensitive reagents were typically performed under a nitrogen or argon atmosphere. Purity was measured using high performance liquid chromatography (HPLC) system with UV detection at 254 nm and 215 nm (System A: Agilent Zorbax Eclipse XDB-C8 4.6 x 150 mm, 5 μm, 5 to 100% CH₃CN in H₂O with 0.1% TFA for 15 min at 1.5 mL/min; System B: Zorbax SB-C8, 4.6 x 75 mm, 10 to 90% CH₃CN in H₂O with 0.1% formic acid for 12 min at 1.0 mL/min) (Agilent Technologies, Santa Clara, CA). Silica gel chromatography was generally performed with prepacked silica gel cartridges (Biotage, Uppsala, Sweden or Teledyne-Isco, Lincoln, NE). ¹H NMR spectra were recorded on a Bruker AV-400 (400 MHz) spectrometer (Bruker Corporation, Madison, WI) or a Varian (Agilent Technologies, Santa Clara, CA) 400 MHz spectrometer at ambient temperature. All observed protons are reported as parts per million (ppm) downfield from tetramethylsilane (TMS) or other internal reference in the appropriate solvent indicated. Data are reported as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), coupling constants, and number of protons. Low-resolution mass spectral (MS) data were determined on an Agilent 1100 Series (Agilent Technologies, Santa Clara, CA) LC/MS with UV detection at 254 nm and 215 nm and a low resonance electrospray mode (ESI).

[0194] For the purpose of clarity in this general synthesis section, Compounds of Formula (I) as defined in the summary of the inventions can be schematically drawn to contain Ring Ar¹ and Ring Ar² as follows:



; wherein the group -NR³-(C=O)- is a linker, Ring Ar¹ is located to the left of the linker, and ring Ar² is located to the right of linker.

[0195] Generally, compounds of Formula (I), can be synthesized via three general steps as follows:

[0196] Step 1: Preparation of Ring Ar¹ compound.

[0197] Step 2: Preparation of Ring Ar² compound.


[0198] Step 3: Coupling of Ring Ar¹ compound to Ring Ar² compound

[0199] The generic Schemes A-E below are meant to provide guidance to the ordinarily skilled synthetic chemist, who will readily appreciate that the solvent, concentration, reagent, protecting group, order of synthetic steps, time, temperature, and the like can be modified as necessary, well within the skill and judgment of the ordinarily skilled artisan.

SCHEME A


[0200] According to Scheme A, in one embodiment a compound of Formula (I) as disclosed herein can be synthesized as follows:

[0201] Step 1a: Preparation of Ring Ar¹ compound:




[0202] Step 1a: Preparation of Ring Ar¹ compound: Compound A-1, wherein W¹ is a halogen, for example fluoro or chloro, can be reacted with an R² group containing agent in the presence of a suitable base, in a suitable organic solvent such as NMP, dioxane, acetonitrile, tetrahydrofuran, DMF, methylene chloride, and the like, to form compound A-2. Compound A-1 is commercially available or can be synthesized by known methods by those skilled in the art. Examples of compound A-1 include, but are not limited to, 2-chloropyrimidin-4-amine, 2-chloro-6-methylpyrimidin-4-amine, 2-fluoro-6-methylpyridin-4-amine, 2-chloropyridin-4-amine, 2-chloro-6-methylpyridin-4-amine, 2-chloro-6-ethylpyrimidin-4-amine, or 2-chloro-6-cyclopropylpyrimidin-4-amine. Examples of R² reagents include, but are not limited to (1) (R)-2-methylmorpholine, (2) 4,4-difluoropiperidine hydrochloride, (3) 3,3-difluoroazetidine hydrochloride, or (4) 3,3,3-trifluoropropan-1-ol. Examples of bases include, but are not limited to diisopropylethyl amine, potassium carbonate, or sodium hydride.

[0203] Step 1b: Preparation of Ring Ar¹ compound:




[0204] Alternatively, compound A-1 as defined in step 1a, can be converted to compound A-2, as defined in step 1a, via Suzuki cross coupling reaction with a suitable organoboron R² reagent (R²-BY₂, wherein Y is an organic functional group) such as 2-(4,4-difluorocyclohex-1-en-1-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane or 2-(4-fluorocyclopent-1-en-1-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane, and a suitable palladium catalyst and a base, such as PdCl₂(dppf)-DCM adduct and potassium phosphate tribasic. This step is followed by a reduction with a suitable palladium catalyst and a hydrogen source, such as Pd/C in the presence of hydrogen gas, to form compound A-2. This alternative Suzuki reaction can be used when the group R² is linked to the Ar¹ ring via a carbon-carbon bond.

[0205] Step 2a: Preparation of Ring Ar² compound:



[0206] In Step 2a, Compound **A-3**, wherein each of W^2 and W^3 is independently a halogen, for example fluoro, chloro, bromo, or iodo, can be reacted with an R^X reagent, such as (1) 6-azaspiro[2.5]octane hydrochloride, (2) 4,4-dimethylpiperidine hydrochloride, (3) 3,4,4-trimethylpiperidine hydrochloride, (4) 4-methyl-6-azaspiro[2.5]octane hydrochloride, or (5) 7-azaspiro[3.5]nonane hydrochloride, in a suitable organic solvent such as NMP, acetonitrile, tetrahydrofuran, DMF, methylene chloride, DMSO, and the like, to form Compound **A-4**.

[0207] Step 3a: Coupling of Ring Ar^1 compound to Ring Ar^2 compound followed by introduction of R^1 :




[0208] In Step 3a, compound **A-4**, which was obtained from Step 2a, can be reacted with an activating agent such as acid chloride $(COCl)_2$ or $SOCl_2$, in a suitable organic solvent such as tetrahydrofuran, methylene chloride, and the like, to form an acid chloride derivative, which can then react with compound **A-2** to form compound **A-5**. Alternatively, compound **A-2** can be directly coupled with compound **A-4**, which was obtained from Step 2a, in a suitable organic solvent such as acetonitrile, tetrahydrofuran, DMF, methylene chloride, and the like, in the presence of a coupling reagent, such as N, N'-diisopropylcarbodiimide, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide, benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, carbonyl diimidazole, and polyphosphonic anhydride. Those ordinary skilled synthetic chemists will readily understand that other coupling agents can be used. Further manipulation of halogen group W^3 by transformation reactions such as, metal-catalyzed sulfoamidation, sulfonation, or sulfonylation, in a suitable organic solvent such as DMSO, acetonitrile, tetrahydrofuran, DMF, and the like, in the presence of a metal catalyst and an R^1 reagent, such as (1) 1-methylcyclopropane-1-sulfonamide, (2) 3-methyloxetan-3-amine, (3) tert-butyl 3-mercaptoazetidine-1-carboxylate, (4) ethyl 2-

sulfamoylpropanoate, (5) 2-hydroxypropane-1-sulfonamide, (6) 2-hydroxyethane-1-sulfonamide, (7) ethyl iodoacetate, (8) 2-mercaptopropan-1-ol, (9) 2-mercapto-2-methylpropan-1-ol, (10) 2-aminoethan-1-ol, or (11) cyclopropanethiol can be used to form Compound (I). Those ordinary skilled chemists will readily understand that coupling reaction such as shown in Step 3a can be performed under various known conditions.

SCHEME B


[0209] Step 1a or 1b: Preparation of Ring Ar¹ compound: see SCHEME A above

[0210] Step 2b: Preparation of Ring Ar² compound:



[0211] Scheme B provides an alternative method for formation of compounds of Formula (I) as disclosed herein. After Step 1a or Step 1b as described in Scheme A, the group R¹ can alternatively be introduced to Ring Ar² in Step 2b rather than in Step 3a as in Scheme A. According to Step 2b, compound **B-1**, wherein each of W⁴ and W⁵ is independently a halogen, for example fluoro, chloro, bromo, or iodo, can be reacted with an appropriate carboxylic acid protecting group (PG₁ reagent), such as methyl iodide in the presence of a base such as potassium carbonate, to form a methyl ester, or other appropriate protecting group to form other ester such as benzyl ester, in a suitable organic solvent such as NMP, acetonitrile, tetrahydrofuran, DMF, methylene chloride and the like, to form compound **B-2**, wherein each of W⁴ and W⁵ are as defined in compound **B-1**. Compound **B-2** can then be reacted with an R^x reagent, such as 6-azaspiro[2.5]octane, in a suitable organic solvent such as NMP, acetonitrile, tetrahydrofuran, DMF, methylene chloride, DMSO, and the like, to form compound **B-3**, wherein W⁵ is as defined in compound **B-1**. Compound **B-3** can then be reacted with an R¹ reagent by a transformation reaction such as, metal-catalyzed sulfoamidation, sulfonation, or sulfonylation, in a suitable organic solvent such as DMSO, acetonitrile, tetrahydrofuran, DMF, and the like, in the presence of a metal catalyst, such as copper iodide, Pd₂(dba)₃ to form compound **B-4**, which can then be reacted further with an appropriate carboxylic acid deprotecting agent to form compound **B-5**. Appropriate carboxylic acid protecting groups and deprotection agents are known to those skilled in the art, e.g., as discussed in Greene's Protective Groups in Organic Synthesis.

[0212] Step 3b: Coupling of Ring Ar¹ compound to Ring Ar² compound:




[0213] Step 3b is similar to the coupling reaction as described above in Step 3a.

SCHEME C

[0214] Scheme C provides yet another alternative method for formation of compounds of Formula (I) as disclosed herein. According to Scheme C, Step 1a can be performed as described in Scheme A, followed by Step 2b as described in Scheme B.

[0215] Step 3c: Coupling of Ring Ar^1 compound to Ring Ar^2 compound:




[0216] In Step 3c, compound **A-1a**, which is compound **A-1** of Scheme A, wherein X^1 is N and W^1 is a halogen, for example fluoro or chloro, can be reacted with compound **B-5** obtained from step 2b of scheme B, in the presence of an activating agents under conditions similar to step 3a and 3b above, to form compound **C-1**, wherein W^1 is as defined in compound **A-1a**, which then can be reacted with an R^2 group containing agent, such as (1) (*R*)-2-methylmorpholine, (2) 4,4-difluoropiperidine hydrochloride, (3) 3,3-difluoroazetidine hydrochloride, or (4) 3,3,3-trifluoropropan-1-ol, in the presence of a suitable base, such as diisopropylethyl amine, potassium carbonate, or sodium hydride, in a suitable organic solvent such as NMP, dioxane, acetonitrile, tetrahydrofuran, DMF, methylene chloride, and the like, to form compound of formula (I).

SCHEME D

[0217] Scheme D provides yet another alternative method for formation of compounds of Formula (I) as disclosed herein. According to Scheme D, Step 1a or 1b can be performed as described in Scheme A, followed by Step 2b as described in Scheme B.

[0218] Step 3d: Coupling of Ring Ar^1 compound to Ring Ar^2 compound:





[0219] In Step 3d, compound **A-1a**, which is compound **A-1** of Scheme A, wherein X^1 is N and W^1 is a halogen, for example fluoro or chloro, can be reacted with compound **B-5** obtained from step 2a of scheme A, in the presence of a activating agent under conditions similar to step 3a and 3b above, to form compound **D-1**, wherein W^1 is as defined in compound **A-1a** and W^3 is as defined in compound **B-5**, which then can be reacted with an R^2 group containing agent, such as (1) (*R*)-2-methylmorpholine, (2) 4,4-difluoropiperidine hydrochloride, (3) 3,3-difluoroazetidine hydrochloride, or (4) 3,3,3-trifluoropropan-1-ol, optionally in the presence of a suitable base, such as diisopropylethyl amine, potassium carbonate, or sodium hydride, in a suitable organic solvent such as NMP, dioxane, acetonitrile, tetrahydrofuran, DMF, methylene chloride, and the like, to form compound **A-5a**, which is compound **A-5**, wherein X^1 is N and W^3 is as defined in compound **B-5**, which can then be reacted with an R^1 group containing agent by a transformation reaction such as, metal-catalyzed sulfoamidation, sulfination, or sulfonylation, in a suitable organic solvent such as DMSO, acetonitrile, tetrahydrofuran, DMF, and the like, in the presence of a metal catalyst, to form compound formula (I).

SCHEME E

[0220] Scheme E provides yet another alternative method for formation of compounds of Formula (I) as disclosed herein. According to Scheme E, Step 1a or 1b can be performed as described in Scheme A to prepare compound **A-2**. Compound **E-1**, wherein W^6 is a halogen, for example fluoro or chloro, which includes but is not limited to 2-fluoro-4-nitrobenzoic acid, 2,5-difluoro-4-nitrobenzoic acid, or 2,6-difluoro-4-nitrobenzoic acid, is commercially available or can be synthesized according to known methods by those skilled in the art.

[0221] Step 3e: Coupling of Ring Ar^1 compound to Ring Ar^2 compound






[0222] In step 3e, compound A-2 can be reacted with compound E-1, the presence of an activation reagent under conditions similar to step 3a and 3b above, to form compound E-2, which then can be reacted with an R^x reagent in a similar way as described in Step 2a to form compound E-3. The nitro group on compound E-4 can then be converted into an amino group by reacting with a reducing agent, which includes but is not limited to palladium on carbon and hydrogen gas, to form compound E-4, which can then be reacted with an R^1 reagent, such as (1) 1-methylcyclopropane-1-sulfonamide, (2) 3-methyloxetan-3-amine, (3) tert-butyl 3-mercaptoazetidine-1-carboxylate, (4) ethyl 2-sulfamoylpropanoate, (5) 2-hydroxypropane-1-sulfonamide, (6) 2-hydroxyethane-1-sulfonamide, (7) ethyl iodoacetate, (8) 2-mercaptopropan-1-ol, (9) 2-mercapto-2-methylpropan-1-ol, (10) 2-aminoethan-1-ol, or (11) cyclopropanethiol, by a transformation reaction such as, metal-catalyzed sulfoamidation, sulfination, or sulfonylation, in a suitable organic solvent such as DMSO, acetonitrile, tetrahydrofuran, DMF, and the like, in the presence of a metal catalyst, to form compound (I).

[0223] EXAMPLES

[0224] PREPARATION OF SYNTHETIC INTERMEDIATES

[0225] RING AR¹ INTERMEDIATES:


[0226] Intermediate 1: (R)-2-(2-Methylmorpholino)pyrimidin-4-amine



[0227] A mixture of 2-chloropyrimidin-4-amine (60.0 g, 463 mmol, Combi-Blocks, San Diego, CA), (R)-2-methylmorpholine (65.6 g, 648 mmol, Wuxi Apptec) and DIPEA (243 mL, 1389 mmol) in NMP (600 mL) was taken in an autoclave and heated at 150 °C for 36 h. The reaction mixture was cooled to room temperature and quenched with water (1 L) and extracted with ethyl acetate (3 x 500 mL). The organic layer was washed with brine solution (500 mL), dried (Na_2SO_4), filtered and concentrated under reduced pressure to give the crude material as a tan oil. The crude material was adsorbed onto a plug of silica gel


and purified by column chromatography over silica gel (60-120 mesh), eluting with a gradient of 50% to 100% ethyl acetate in hexanes to give a yellow solid. This solid was further triturated with hexanes (300 mL), filtered and dried under vacuum to give the title compound (70 g, 78% yield) as a pale yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 7.75 (d, $J = 5.6$ Hz, 1H), 6.42 (s, 2H), 5.75 (d, $J = 5.6$ Hz, 1H), 4.26 – 4.49 (m, 2H), 3.83 (ddd, $J = 11.4, 3.6, 1.4$ Hz, 1H), 3.31 – 3.50 (m, 2H), 2.78 (ddd, $J = 13.2, 11.8, 3.5$ Hz, 1H), 2.42 – 2.48 (m, 1H), 1.11 (d, $J = 6.2$ Hz, 3H). m/z (ESI): 195.2 ($\text{M}+\text{H}$) $^+$.

[0228] Intermediate 2: (R)-6-Methyl-2-(2-methylmorpholino)pyrimidin-4-amine



[0229] A mixture of 2-chloro-6-methylpyrimidin-4-amine (30.0 g, 209 mmol, Combi-Blocks, San Diego, CA), (R)-2-methylmorpholine (40.3 g, 293 mmol, Wuxi Apptec, PR China), and DIPEA (109 mL, 627 mmol) was taken in an autoclave (600 mL) and heated at 150 °C for 12 h. The reaction mixture was quenched with water (500 mL) and extracted with ethyl acetate (2 x 1500 mL). The organic layer was washed with brine solution (500 mL), dried (Na_2SO_4), filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography over silica gel (60-120 mesh) using 50% ethyl acetate in hexanes as an eluent to give the title compound (25.0 g, 57% yield) as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 6.28 (s, 2H), 5.62 (s, 1H), 4.45 – 4.31 (m, 2H), 3.83 (ddd, $J = 11.4, 3.5, 1.3$ Hz, 1H), 3.42 (ddt, $J = 14.4, 9.7, 2.8$ Hz, 2H), 2.78 – 2.70 (m, 1H), 2.43 (dd, $J = 13.0, 10.3$ Hz, 1H), 2.05 (s, 3H), 1.11 (d, $J = 6.2$ Hz, 3H). m/z (ESI): 209.2 ($\text{M}+\text{H}$) $^+$.


[0230] Intermediate 3: 2-(4,4-Difluoropiperidin-1-yl)pyrimidin-4-amine



[0231] A glass microwave reaction vessel was successively charged with 2-chloro-4-aminopyrimidine (1.0 g, 7.7 mmol, Combi-Blocks, San Diego, CA), 4,4-difluoropiperidine hydrochloride (1.82 g, 11.58 mmol, Combi-Blocks, San Diego, CA) and DIPEA (4.04 mL, 23.2 mmol) in NMP (12 mL). The reaction mixture was stirred and heated in a microwave at 200 °C for 1 h. Reaction mixture was diluted with water (50 mL), extracted with ethyl acetate (30 x 2 mL). The organic extract was washed with brine (30 x mL), dried over Na_2SO_4 , concentrated in vacuo to give the crude material as a brownish sticky liquid. The crude material was absorbed onto a plug of silica gel and purified by flash chromatography through a Redi-Sep pre-packed silica gel column (40 g), eluting with 0 % to 100 % ethyl acetate in heptane to provide the title

compound as an off-white solid (6.02 g, 91 %). ^1H NMR (300 MHz, $\text{DMSO-}d_6$) δ ppm 7.21 (d, $J = 5.6$ Hz, 1H), 6.46 (br s, 2H), 5.77 (d, $J = 5.6$ Hz, 1H), 3.80 (t, $J = 5.6$ Hz, 4H), 1.85-1.90 (m, 4H). m/z (ESI): 215.2 ($\text{M}+\text{H}$) $^+$.


[0232] Intermediate 4: 2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-amine



Intermediate 4

[0233] A mixture of 2-chloro-6-methylpyrimidin-4-amine (46 g, 320 mmol, Combi-Blocks, San Diego, CA), 4,4-difluoropiperidine hydrochloride (76 g, 481 mmol, Combi-Blocks, San Diego, CA) and DIPEA (166 mL, 961 mmol) in NMP (460 mL, 10.00 mL/g) was taken in an autoclave (1 L) and heated at 180 °C for 30 h. The reaction mixture was cooled to room temperature and quenched with water (500 mL), extracted with ethyl acetate (2 x 1000 mL). The organic layer was washed with brine (500 mL), dried (Na_2SO_4), filtered and concentrated under reduced pressure. The crude material was adsorbed onto a plug of silica gel and purified by column chromatography over silica gel (60-120 mesh), eluting with 50% to 100% ethyl acetate in hexanes as an eluent to give the product. This was re-dissolved in ethyl acetate (500 mL), washed with water (2 x 500 mL). The organic layer was dried (Na_2SO_4), filtered and concentrated under reduced pressure. The yellow solid was once again suspended in hexanes (400 mL) and stirred for 30 min. The slurry was filtered, washed with hexanes (100 mL), dried under vacuum to provide the title compound (58 g, 79% yield) as a pale yellow solid. ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ ppm 6.33 (s, 2H), 5.63 (s, 1H), 3.80 – 3.78 (dd, $J = 6.8, 4.7$ Hz, 4H), 2.06 (s, 3H), 1.95 – 1.85 (tt, $J = 14.2, 5.7$ Hz, 4H). m/z (ESI): 229.2 ($\text{M}+\text{H}$) $^+$.

[0234] Intermediate 5: 2-(3,3-Difluoroazetidin-1-yl)pyrimidin-4-amine




Intermediate 5

[0235] A mixture of 2-chloro-4-aminopyrimidine (5.0 g, 38.6 mmol, Combi-Blocks, San Diego, CA), 3,3-difluoroazetidine hydrochloride (7.50 g, 57.9 mmol, Combi-Blocks, San Diego, CA) and potassium carbonate (5.33 g, 38.6 mmol) in dioxane (25 mL) was heated at 95 °C for 16 h. The reaction mixture was cooled to room temperature and the suspension was filtered. The crude material was absorbed onto a plug of silica gel and purified by flash chromatography through a Redi-Sep pre-packed silica gel column (40 g), eluting with 10 % MeOH in DCM to provide the title compound as light brown solid (6.1 g, 85 %). ^1H

NMR (300 MHz, DMSO- d_6) δ ppm 8.29-8.66 (m, 2H), 7.81 (d, J = 7.05 Hz, 1H), 6.22 (d, J = 7.26 Hz, 1H), 4.61 (t, J = 12.23 Hz, 4H). m/z (ESI): 187.2 (M+H)⁺.


[0236] Intermediate 6: 2-(4,4-Difluorocyclohexyl)-6-methylpyrimidin-4-amine



[0237] Step 1: To a solution of 2-chloro-6-methylpyrimidin-4-amine (70.0 g, 488 mmol, Combi-Blocks, San Diego, CA) in 1,4-dioxane (560 mL) and water (210 mL) were added 2-(4,4-difluorocyclohex-1-en-1-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (119 g, 488 mmol, Combi-Blocks, San Diego, CA) and potassium phosphate tribasic (310 g, 1463 mmol). The reaction mixture was degassed and purged with nitrogen for 5 min. PdCl₂(dppf)-DCM adduct (39.8 g, 48.8 mmol) was added to the reaction mixture and stirred at 100 °C for 16 h. The dark heterogeneous mixture was filtered through a CELITE® bed and the filter cake was washed with ethyl acetate (2 x 1000 mL). The filtrate was washed with 1N NaOH solution (300 mL), followed by water (500 mL). The organic layer was dried (Na₂SO₄), filtered and concentrated under reduced pressure. The crude residue was adsorbed onto a plug of silica gel and purified by column chromatography over silica gel (60-120 mesh), eluting with a gradient of 50% to 60% ethyl acetate in hexanes, to give 2-(4,4-difluorocyclohex-1-en-1-yl)-6-methylpyrimidin-4-amine (70 g, 64% yield) as a pale yellow solid. ¹H NMR (400 MHz, DMSO- d_6) δ ppm 6.85 (br s, 1H), 6.59 (br s, 2H), 6.13 (s, 1H), 2.62 – 2.80 (m, 4H), 2.19 (s, 3H), 2.04 – 2.17 (m, 2H). m/z (ESI): 226.2 (M+H)⁺.

[0238] Step 2: To a solution of 2-(4,4-difluorocyclohex-1-en-1-yl)-6-methylpyrimidin-4-amine (70.0 g, 311 mmol) in EtOH (700 mL) was added 10% Pd on carbon (33.1 g, 155 mmol) under nitrogen. The reaction mixture was stirred at room temperature under hydrogen pressure (1 atm) for 16 h. The reaction mixture was filtered through a CELITE® bed and washed with a mixture of ethyl acetate and ethanol (1:1, 500 mL). The filtrate was concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel (60-120 mesh) using 50% ethyl acetate in hexanes to give the title compound (58.5 g, 83% yield) as an off white solid. ¹H NMR (400 MHz, DMSO- d_6) δ ppm 6.61 (s, 2H), 6.09 (s, 1H), 2.56 – 2.67 (m, 1H), 2.16 (s, 3H), 1.72 – 2.10 (m, 8H). m/z (ESI): 228.1 (M+H)⁺.


[0239] Intermediate 7: 2-Methyl-6-(3,3,3-trifluoropropoxy)pyridin-4-amine




[0240] To a solution of 3,3,3-trifluoropropan-1-ol (1.99 g, 17.44 mmol, Combi-Blocks) in 30 mL of THF at 0 °C was added sodium hydride (60% wt. in mineral oil, 0.79 g, 19.82 mmol). The mixture was stirred at RT for 30 min then treated with 2-fluoro-6-methylpyridin-4-amine (1.00 g, 7.93 mmol, AstaTech Inc).

The mixture was heated at 65 °C in an oil bath for 5 h. It was cooled to RT, quenched with water (10 mL) and extracted with EtOAc (2 x 50 mL). The organic solution was dried over Na₂SO₄, filtered and concentrated in vacuo to give the crude material as a yellow oil. The crude material was absorbed onto a plug of silica gel and purified on a silica gel column (15% to 30% EtOAc in heptane) to give 2-methyl-6-(3,3,3-trifluoropropoxy)pyridin-4-amine (0.47 g, 2.13 mmol, 27% yield) as a light-yellow oil. *m/z* (ESI): 221.1 (M+H)⁺.

[0241] Table 1: The intermediate below was prepared following a similar procedure for Intermediate 7:



Int. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
7-1		6-Methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-amine	222.0

[0242] Intermediate 8: 2-(4,4-Difluoropiperidin-1-yl)pyridin-4-amine




[0243] A mixture of 2-chloropyridin-4-amine (2.00 g, 15.56 mmol, Combi Blocks), DIPEA (6.03 g, 46.70 mmol, Sigma-Aldrich) and 4,4-difluoropiperidine (2.45 g, 20.22 mmol, Enamine) in NMP (6 mL) was heated in a microwave at 200 °C for 6 h. The reaction mixture was diluted with water (20 mL) and extracted with EtOAc (2 x 50 mL). The organic extracts were concentrated. The residue was purified by silica gel chromatography (20% to 70% EtOAc in heptane) to provide 2-(4,4-difluoropiperidin-1-yl)pyridin-4-amine (2.91 g, 13.65 mmol, 88% yield) as a light yellow solid. *m/z* (ESI): (M+H)⁺ 214.1.

[0244] Table 2: Intermediates below were prepared following a similar procedure as described for Intermediate 8

Int. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
8-1		2-(4,4-Difluoropiperidin-1-yl)-6-methylpyridin-4-amine	228.2
8-2		2-(4,4-Difluoropiperidin-1-yl)-6-ethylpyrimidin-4-amine	243.1

Int. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
8-3		6-Cyclopropyl-2-(4,4-difluoropiperidin-1-yl)pyrimidin-4-amine	255.1

[0245] Intermediate 9: 2-(4,4-Difluoropiperidin-1-yl)-3-fluoro-6-methylpyridin-4-amine



[0246] Step 1: To a solution of diisopropylamine (5.85 mL, 41.0 mmol) in tetrahydrofuran (50 mL) was added *n*-butyllithium (2M solution in hexanes, 20.52 mL, 41.0 mmol) dropwise at -60 °C. The reaction mixture was slowly warmed to 0 °C and was stirred at the same temperature for 45 minutes. In another round bottom flask, to a solution of 2-bromo-3-fluoro-6-methylpyridine (3.9 g, 20.52 mmol) in tetrahydrofuran (50 mL) was added the above prepared LDA solution dropwise at -78 °C. The reaction resultant reaction mixture was stirred for 45 minutes at the same temperature and then iodine (10.42 g, 41.0 mmol) in THF (40 mL) was added dropwise. The reaction mixture was stirred at same temperature for 1h. After completion of the reaction, it was quenched with a saturated solution of ammonium chloride and was extracted in ethyl acetate. The organic layer was washed with sodium thiosulfate solution, water, and brine, dried over Na₂SO₄, filtered and concentrated to afford 2-bromo-3-fluoro-4-iodo-6-methylpyridine (6 g, 18.99 mmol, 93 % yield) as yellow solid. The product was taken for next step without further purification. ¹H NMR (400 MHz, Chloroform-*d*) δ ppm 7.53 (d, *J* = 3.8 Hz, 1 H), 2.52 (s, 3 H). m/z (ESI): 315.8, 317.8 (M+H)⁺.


[0247] Step 2: A mixture of 2-bromo-3-fluoro-4-iodo-6-methylpyridine (1.5 g, 4.75 mmol), (4-methoxyphenyl)methanamine (0.782 g, 5.70 mmol), cesium carbonate (4.64 g, 14.24 mmol), Xantphos (0.549 g, 0.950 mmol) and Pd₂(dba)₃ (0.065 g, 0.071 mmol) in 1,4-dioxane (30 mL) was stirred at ambient temperature for 16 h. Then the reaction mixture was filtered through a plug of CELITE® and filtrate was diluted with EtOAc. The resulting solution was washed with water, brine, dried over Na₂SO₄, filtered and concentrated. The concentrate was purified by flash column chromatography using a gradient of 0 - 18 % ethyl acetate in petroleum ether to afford 2-bromo-3-fluoro-*N*-(4-methoxybenzyl)-6-methylpyridin-4-amine (0.7 g, 2.15 mmol, 45 % yield) as pale-yellow solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 7.38 (s,


1 H), 7.21 – 7.31 (m, 2 H), 6.84 – 6.95 (m, 2 H), 6.52 (d, $J = 6.0$ Hz, 1 H), 4.32 (d, $J = 6.3$ Hz, 2 H), 3.72 (s, 3 H), 2.20 (s, 3 H). m/z (ESI): 325.0, 327.0 (M+H)⁺.

[0248] Step 3: A mixture of 2-bromo-3-fluoro-*N*-(4-methoxybenzyl)-6-methylpyridin-4-amine (0.7 g, 2.153 mmol), 4,4-difluoropiperidine hydrochloride (0.407 g, 2.58 mmol), cesium carbonate (2.81 g, 8.61 mmol), Xantphos (0.249 g, 0.431 mmol) and Pd₂(dba)₃ (0.030 g, 0.032 mmol) in 1,4-dioxane (15 mL) was stirred in a sealed tube at 100 °C for 16 h. Then the reaction mixture was filtered through a plug of CELITE® and filtrate was diluted with EtOAc. The resulting solution was washed with water, brine, dried over Na₂SO₄, filtered, and concentrated. The concentrate was purified by flash column chromatography using a gradient of 0 - 6 % ethyl acetate in petroleum ether to afford 2-(4,4-difluoropiperidin-1-yl)-3-fluoro-*N*-(4-methoxybenzyl)-6-methylpyridin-4-amine (0.67 g, 1.83 mmol, 85 % yield) as pale-yellow solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 7.19 – 7.29 (m, 2 H), 6.83 – 6.93 (m, 2 H), 6.75 (m, 1H), 6.13 (d, $J = 5.4$ Hz, 1 H), 4.26 (d, $J = 6.3$ Hz, 2 H), 3.72 (s, 3 H), 3.40 (d, $J = 11.5$ Hz, 4 H), 1.92 – 2.14 (m, 7 H). m/z (ESI): 366.1 (M+H)⁺.

[0249] Step 4: To a solution of 2-(4,4-difluoropiperidin-1-yl)-3-fluoro-*N*-(4-methoxybenzyl)-6-methylpyridin-4-amine (0.3 g, 0.821 mmol) in dichloromethane (3 mL) were added anisole (0.179 mL, 1.642 mmol) and TFA (1.5 mL, 19.47 mmol) at ambient temperature and the reaction mixture was stirred at 50 °C for 2.5 h. Then the reaction mixture quenched with water and pH was adjusted to 8 with 10 % sodium bicarbonate solution before it was extracted with ethyl acetate. The organic layer was washed with water and brine, dried over Na₂SO₄, filtered and concentrated. The concentrate was purified by flash column chromatography using a gradient of 20 % ethyl acetate in petroleum ether to afford 2-(4,4-difluoropiperidin-1-yl)-3-fluoro-6-methylpyridin-4-amine (0.17 g, 0.69 mmol, 84% yield) as yellow oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 6.15 (d, $J = 5.5$ Hz, 1 H), 5.79 (s, 2 H), 3.40 (t, $J = 5.6$ Hz, 4 H), 2.13 (s, 3 H), 2.02 (tt, $J = 14.2, 5.6$ Hz, 4 H). m/z (ESI): 246.2 (M+H)⁺.

[0250] Intermediate 10: 2-(6-Amino-2-(4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)propan-2-ol





[0251] Step 1: To a solution of methyl 2,6-dichloropyrimidine-4-carboxylate (20.00 g, 97 mmol), in tetrahydrofuran (200 mL) were added (3,5-dimethoxyphenyl)methanamine (19.39 g, 116 mmol) and DIPEA (33.7 mL, 193 mmol) at 0 °C. Then the reaction mixture was stirred at ambient temperature for 16 h before it was quenched with water and was extracted with EtOAc. The organic layer was washed with brine, and dried over Na₂SO₄, filtered and concentrated. The concentrate was triturated with DCM and hexane to afford methyl 2-chloro-6-((3,5-dimethoxybenzyl)amino)pyrimidine-4-carboxylate (19.5 g, 57.7 mmol, 59.8 % yield) as off white solid. ¹H NMR(400 MHz, DMSO-*d*₆) δ ppm 8.52 (t, *J* = 5.5 Hz, 1 H), 7.16 (d, *J* = 8.9 Hz, 2 H), 6.59 (d, *J* = 2.4 Hz, 1 H), 6.50 (dd, *J* = 8.4, 2.4 Hz, 1 H), 4.40 (d, *J* = 5.4 Hz, 2 H), 3.82 (d, *J* = 14.5 Hz, 6 H), 3.75 (d, *J* = 5.8 Hz, 3 H). *m/z* (ESI): 338.1 (M+H)⁺.

[0252] Step 2: A solution of methyl 2-chloro-6-((2,4-dimethoxybenzyl)amino)pyrimidine-4-carboxylate (3 g, 8.88 mmol), 4,4-difluoropiperidine hydrochloride (2.10 g, 13.32 mmol) and DIPEA (3.44 g, 26.6 mmol) in DMF (30 mL) was stirred in a sealed tube at 90 °C for 16 h. Then the reaction mixture was quenched with water and was extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The crude material was purified by flash column chromatography using a gradient of 50% to 60% ethyl acetate in petroleum ether to provide methyl 2-(4,4-difluoropiperidin-1-yl)-6-((2,4-dimethoxybenzyl)amino)pyrimidine-4-carboxylate (2.6 g, 6.15 mmol, 69.3 % yield) as pale yellow solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 7.74 (s, 1 H), 7.13 (d, *J* = 8.3 Hz, 1 H), 6.55 (d, *J* = 2.4 Hz, 1 H), 6.47 (dd, *J* = 8.3, 2.4 Hz, 2 H), 4.39 (s, 2 H), 3.82 (s, 3 H), 3.79 (s, 3 H), 3.73 (s, 3 H), 3.33 (m, 4 H), 1.85 – 2.00 (m, 4 H).

[0253] Step 3: To a solution of methyl 2-(4,4-difluoropiperidin-1-yl)-6-((2,4-dimethoxybenzyl)amino)pyrimidine-4-carboxylate (1 g, 2.367 mmol) in DCM (10 mL) was dropwise added sulfuric acid (0.126 mL, 2.37 mmol) at 0 °C. Then the mixture was allowed to warm to RT and the progress of the reaction was monitored by TLC. Upon completion of starting material, the reaction mixture was quenched with ice-water and pH was adjusted to 9 by using 10% NaHCO₃ solution. Then, the reaction mixture was extracted with ethyl acetate and the organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated to provide methyl 6-amino-2-(4,4-difluoropiperidin-1-yl)pyrimidine-4-carboxylate (0.45 g, 1.653 mmol, 69.8 % yield) as off-white solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 6.90 (br s, 2 H), 6.39 (s, 1 H), 3.85 – 3.83 (m, 4 H), 3.79 (s, 3 H), 1.99-1.35 (m, 4 H). *m/z* (ESI): 273.1 (M+H)⁺.


[0254] Step 4: To a solution of methyl 6-amino-2-(4,4-difluoropiperidin-1-yl)pyrimidine-4-carboxylate (0.45 g, 1.653 mmol) in tetrahydrofuran (5 mL) was added methyl magnesium bromide (2.0 M diethyl ether) (2.07 mL, 4.13 mmol) at 0 °C and was stirred at room temperature for 2 h. Upon consumption of

starting material, the reaction mixture was quenched with ice water and was extracted with ethyl acetate. The organic layer was washed with brine, dried over Na_2SO_4 , filtered and concentrated. The concentrate was purified by flash column chromatography using a gradient of 0% - 90% ethyl acetate in petroleum ether to provide 2-(6-amino-2-(4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)propan-2-ol (0.35 g, 1.28 mmol, 78 % yield) as pale yellow solid. m/z (ESI): 273.1 ($\text{M}+\text{H}$)⁺.

[0255] Table 3: The intermediate below was prepared following a similar procedure as described for intermediate 10

Int. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
10-1		(<i>R</i>)-2-(6-Amino-2-(2-methylmorpholino)pyrimidin-4-yl)propan-2-ol	252.2

[0256] Intermediate 11: 2-(4-Amino-6-methylpyrimidin-2-yl)propan-2-ol



[0257] Step 1: A mixture of 2,4-dichloro-6-methylpyrimidine (3.0 g, 18.40 mmol, Aldrich, St. Louis, MO, USA), bis(4-methoxybenzyl)-amine (7.10 g, 27.6 mmol, Combi-Blocks Inc., San Diego, CA, USA), and potassium carbonate (7.63 g, 55.2 mmol, Aldrich, St. Louis, MO, USA) in tetrahydrofuran (100 mL) was stirred at room temperature for 72 h. Then, the mixture was diluted with water (50 mL) and was then extracted with EtOAc (2 x 100 mL). The combined organic extracts were then dried over MgSO_4 and concentrated in vacuo. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc/heptane) provided 2-chloro-*N,N*-bis(4-methoxybenzyl)-6-methylpyrimidin-4-amine (3.11 g, 8.10 mmol, 44.0 % yield) as an off white solid. ^1H NMR ($\text{DMSO}-d_6$) δ ppm 7.16 (br d, $J=6.2$ Hz, 4H), 6.89 (d, $J=8.7$ Hz, 4H), 6.60 (s, 1H), 4.39-4.84 (m, 4H), 3.73 (s, 6H), 2.20 (s, 3H). m/z (ESI): 384.2 ($\text{M}+\text{H}$)⁺.

[0258] Step 2: A mixture of 2-chloro-*N,N*-bis(4-methoxybenzyl)-6-methylpyrimidin-4-amine (1.0 g, 2.61 mmol), 1,3-bis(diphenylphosphino)propane (64.5 mg, 0.156 mmol, Aldrich, St. Louis, MO, USA), diethyl oxalate (0.529 mL, 3.91 mmol, Aldrich, St. Louis, MO, USA), trans-dichlorobis(triphenylphosphine)palladium (ii) (54.9 mg, 0.078 mmol, Strem Chemicals Inc., Newburyport, MA, USA), and 4-(dimethylamino) pyridine (477 mg, 3.91 mmol, Aldrich, St. Louis, MO, USA) in ethanol (0.5 mL) was subjected to a microwave irradiation at 140 °C for 20 min. Then, the mixture was diluted with water (50


mL) and was then extracted with EtOAc (2 x 50 mL). The combined organic extracts were then dried over MgSO₄ and concentrated in vacuo. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc/heptane) provided ethyl 4-(bis(4-methoxybenzyl)amino)-6-methylpyrimidine-2-carboxylate (274 mg, 0.650 mmol, 24.95 % yield) as a light yellow solid. ¹H NMR (Methanol-d₄) δ ppm 7.29 (br s, 4H), 6.97 (d, *J*=8.5 Hz, 4H), 6.68 (s, 1H), 4.78-4.93 (m, 4H), 4.54 (q, *J*=7.2 Hz, 2H), 3.88 (s, 6H), 2.43 (s, 3H), 1.53 (t, *J*=7.0 Hz, 3H). *m/z* (ESI): 422.1 (M+H)⁺.

[0259] Step 3: To a solution of ethyl 4-(bis(4-methoxybenzyl)amino)-6-methylpyrimidine-2-carboxylate (396 mg, 0.940 mmol) in 2-methyltetrahydrofuran (7 mL) at 0 °C under N₂ was added methylmagnesium bromide, 3.4M in 2-methyltetrahydrofuran (0.829 mL, 2.82 mmol, Aldrich, St. Louis, MO, USA) dropwise. After addition, the mixture was then stirred at 0°C for 3.5 hours. Then, the mixture was quenched with saturated NH₄Cl (10 mL) and was then extracted with EtOAc (2 x 50 mL). The combined organic extracts were then dried over MgSO₄ and concentrated in vacuo. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc/heptane) provided 2-(4-(bis(4-methoxybenzyl)amino)-6-methylpyrimidin-2-yl)propan-2-ol (307 mg, 0.753 mmol, 80 % yield) as a yellow solid. *m/z* (ESI): 408.2 (M+H)⁺.

[0260] Step 4: A solution of 2-(4-(bis(4-methoxybenzyl)amino)-6-methylpyrimidin-2-yl)propan-2-ol (300 mg, 0.736 mmol) in trifluoroacetic acid (10 mL, Aldrich, St. Louis, MO, USA) was subjected to a microwave irradiation at 110 °C for 30 min. Then, the mixture was concentrated under reduced pressure. The crude was then dissolved in DCM (10 mL) and was then quenched with saturated Na₂CO₃ (15 mL). The mixture was then extracted with EtOAc (2 x 50 mL). The combined organic extracts were then dried over MgSO₄ and concentrated in vacuo. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc: EtOH (3:1)/heptane) provided 2-(4-amino-6-methylpyrimidin-2-yl)propan-2-ol (123 mg) as a light yellow solid. *m/z* (ESI): 168.2 (M+H)⁺.

[0261] PREPARATION OF RING AR² INTERMEDIATES:

[0262] Intermediate 12: 4-Iodo-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid



Intermediate 12


[0263] To a solution of 2-fluoro-4-iodobenzoic acid (300 g, 1.13 mol, Combi-Blocks, San Diego, CA) in DMSO (2.10 L) was added 6-azaspiro[2.5]octane hydrochloride (216 g, 1.47 mol, Wuxi AppTec) at 20°C. Then K₂CO₃ (468 g, 3.38 mol) was added and the reaction solution was stirred at 140°C for 48 hours under N₂. The reaction solution was slowly poured into ice water (4.20 L), then extracted with hexanes (2.00 L x 3). The water phase was separated and adjusted to pH = 6 with HCl (2.00 mol/L, aq). Solid was precipitated out and collected. The solid was washed with water (700 mL x 3) and filtered. The moist solid was spread out on a large watch glass and dried in the air at 25°C for 72 hours. 4-Iodo-2-(6-


azaspiro[2.5]octan-6-yl)benzoic acid (280 g, 777 mmol, 68.9% yield) was obtained as a light yellow solid. 400 MHz DMSO- d_6 δ ppm 8.07 (s, 1H), 7.76 - 7.66 (m, 2H), 3.10 (t, J = 5.2 Hz, 4H), 1.55 (br s, 4H), 0.41 (s, 4H).

[0264] Table 4: Intermediates below were prepared following a similar procedure for Int. 12:

Int. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
12-1		4-Bromo-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid	310.2/312.2
12-2		Methyl 4-bromo-2-(6-azaspiro[2.5]octan-6-yl)benzoate	324.0/326.0
12-3		4-Iodo-2-(7-azaspiro[3.5]nonan-7-yl)benzoic acid	372.0
12-4		2-(4,4-Dimethylpiperidin-1-yl)-4-iodobenzoic acid	360.0

[0265] Intermediate 13: 4-(Methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoic






[0266] Step 1: To a solution of 2-fluoro-4-(methylsulfonyl)benzoic acid (90.0 g, 412.1 mmol) in *N,N*-dimethylformamide (1.0 L) were added benzyl bromide (78.1 g, 454.0 mmol) and sodium carbonate (52.5 g, 495 mmol) at 0 °C. The reaction mixture was stirred for 12 h at room temperature. The reaction mixture was quenched with water (1 L) and extracted with MTBE (3 x 1 L). The combined organic layer was washed with brine (1 L), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography over silica gel using 0 to 30% ethyl acetate in hexanes as an eluent to give benzyl 2-fluoro-4-(methylsulfonyl)benzoate (100 g, 79% yield) as a white solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 8.16 (dd, *J* = 8.2, 6.9 Hz, 1H), 7.98 – 7.86 (m, 2H), 7.48 – 7.31 (m, 5H), 5.40 (s, 2H), 3.33 (s, 3H).

[0267] Step 2: To a solution of benzyl 2-fluoro-4-(methylsulfonyl)benzoate (55g, 178 mmol) in dimethyl sulfoxide (550 mL) was added DIPEA (57.6 g, 446 mmol) followed by 6-azaspiro[2.5]octane (29.8 g, 268 mmol) and the reaction mixture was stirred at 100 °C for 24 h. The reaction mixture was quenched with water (1 L) and extracted with MTBE (3 x 1 L). The combined organic layer was washed with brine solution (1 L), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel (230-400 mesh) using 0 to 10% ethyl acetate in hexanes as an eluent to give benzyl 4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoate (55 g, 77 % yield) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.76 (d, *J* = 8.0 Hz, 1H), 7.52 – 7.45 (m, 4H), 7.43 – 7.35 (m, 3H), 5.35 (s, 2H), 3.25 (s, 3H), 3.05 (t, *J* = 5.3 Hz, 4H), 1.36 (t, *J* = 5.3 Hz, 4H), 0.30 (s, 4H). *m/z* (ESI): 400.1 (M+H)⁺.

[0268] Step 3: To a solution of benzyl 4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoate (65 g, 163 mmol) in tetrahydrofuran (108 mL) and methanol (36 mL) was added 1N aqueous sodium hydroxide solution (407 mL, 407 mmol) and the reaction mixture was stirred for 12 h at 60 °C. The reaction mixture was concentrated under reduced pressure to remove THF and methanol. The remaining aqueous solution was acidified to pH ~2 with 1.5 N HCl solution. The precipitated solid was filtered, washed with water (200 mL) followed by hexanes (200 mL), dried under vacuum for 12 h to give 4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid (42 g, 83% yield) as an off white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 16.13 (s, 1H), 8.04 (d, *J* = 8.0 Hz, 1H), 7.99 (s, 1H), 7.75 (d, *J* = 8.0 Hz, 1H), 3.29 (s, 3H), 3.17 (b s, 4H), 1.55 (b s, 4H), 0.41 (s, 4H). *m/z* (ESI): 310.1 (M+H)⁺.

[0269] Intermediate 14: 4-((1-Methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid



Intermediate 14

[0270] Step 1: To a solution of 4-bromo-2-fluorobenzoic acid (50.0 g, 228 mmol, F Chemicals, China) in DMF (500 mL) was added sodium carbonate (31.5 g, 297 mmol) followed by benzyl bromide (43.0 g, 251 mmol) at 0 °C and the reaction mixture was stirred for 24 h at room temperature. The reaction mixture was quenched with water (1000 mL) and extracted with ethyl acetate (3 x 2000 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography over silica gel using 10% ethyl acetate in hexanes as an eluent to give benzyl 4-bromo-2-fluorobenzoate (65 g, 92% yield) as a colorless viscous oil. ¹H NMR (400 MHz, Chloroform-*d*) δ ppm 7.91 (t, *J* = 8.4 Hz, 1H), 7.47 (dt, *J* = 6.0, 1.5 Hz, 2H), 7.43 (dd, *J* = 6.8, 1.8 Hz, 1H), 7.41 (q, *J* = 1.5 Hz, 1H), 7.39 – 7.35 (m, 1H), 7.02 (dd, *J* = 8.7, 2.1 Hz, 1H), 6.79 (s, 1H), 5.38 (s, 2H). *m/z* (ESI): 310.2 (MH)⁺.


[0271] Step 2: To a solution of benzyl 4-bromo-2-fluorobenzoate (60.0 g, 194 mmol) in DMSO (200 mL) were added 6-azaspiro[2.5]octane (32.4, 291 mmol, Wuxi Appec) in DMSO (200 mL) and DIPEA (30 g, 291 mmol) and stirred at 100 °C for 12 h. The reaction mixture was quenched with water (2000 mL) and extracted with ethyl acetate (3 x 2000 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography over silica gel using 10% ethyl acetate in hexanes to give benzyl 4-bromo-2-(6-azaspiro[2.5]octan-6-yl)benzoate (70 g, 90% yield) as a yellow oil. ¹H NMR (400 MHz, Chloroform-*d*) δ ppm 7.60 (dd, *J* = 8.4, 1.9 Hz, 1H), 7.50 – 7.45 (m, 2H), 7.49 – 7.28 (m, 3H), 7.20 (d, *J* = 1.9 Hz, 1H), 7.07 (dd, *J* = 8.3, 1.9 Hz, 1H), 5.36 (s, 2H), 3.12 – 3.02 (m, 4H), 1.47 (t, *J* = 5.3 Hz, 4H), 0.33 (d, *J* = 1.8 Hz, 4H). *m/z* (ESI): 398.1, 400.1 (M+H)⁺.

[0272] Step 3: To a 250-mL sealed tube were added benzyl 4-bromo-2-(6-azaspiro[2.5]octan-6-yl)benzoate (9 g, 22.48 mmol), 1-methylcyclopropane-1-sulfonamide (3.95 g, 29.2 mmol, Combi-Blocks, San Diego, CA) and K₂CO₃ (6.21 g, 45.0 mmol) in 1,4-dioxane (90 mL) and the reaction was degassed and purged with nitrogen for 5 min. To this reaction mixture was added Xantphos (1.301 g, 2.248 mmol) followed by Pd₂(dba)₃ (1.03 g, 1.12 mmol) and the sealed tube was closed and stirred at 110 °C for 18 h. The reaction mixture was quenched with water (250 mL) and extracted with ethyl acetate (2 x 150 mL).

The organic layer was washed with water (100 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography over silica gel eluting with a gradient of 0% to 15% EtOAc in hexanes to give benzyl 4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzoate (6.1 g, 59% yield) as an orange oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 10.06 (s, 1H), 7.62 (d, *J* = 8.5 Hz, 1H), 7.48 – 7.31 (m, 5H), 6.98 (s, 1H), 6.81 (d, *J* = 8.5 Hz, 1H), 5.27 (s, 2H), 2.92 (t, *J* = 4.96 Hz, 4H), 1.40 – 1.30 (m, 7H), 1.16 (dd, *J* = 6.4, 4.7 Hz, 2H), 0.81 (dd, *J* = 6.4, 4.7 Hz, 2H), 0.28 (s, 4H). *m/z* (ESI): 455.2 (M+H)⁺.

[0273] Step 4: To a solution of benzyl 4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzoate (2.1 g, 4.62 mmol) in methanol (20 mL) and ethyl acetate (10 mL) was added 10% Pd-C (1.05 g, 50% wt/wt) under nitrogen atmosphere. The reaction mixture was degassed and stirring under hydrogen pressure (1 atm, balloon pressure) for 4 h. The reaction mixture was filtered through a CELITE® bed and washed with methanol (20 mL). The filtrate was concentrated under reduced pressure. The residue was triturated with Et₂O (50 mL) to give 4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid (1.2 g, 71% yield) as an off white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.20 (s, 1H), 10.33 (s, 1H), 7.95 (d, *J* = 8.6 Hz, 1H), 7.41 (s, 1H), 7.20 (d, *J* = 8.6 Hz, 1H), 2.99 (s, 4H), 1.56 (s, 4H), 1.39 (s, 3H), 1.18 (t, *J* = 4.8 Hz, 2H), 0.83 (t, *J* = 4.7 Hz, 2H), 0.42 (s, 4H). *m/z* (ESI): 363.2 (M-H)⁺.

[0274] Intermediate 15: 4-(*N*-(3-Methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid




[0275] Step 1: To a solution of 3-methyl-3-oxetanamine hydrochloride (5.50 g, 44.5 mmol) and *N,N*-diisopropylethylamine (23.26 mL, 134 mmol) in DCM (200 mL) at 0 °C, methyl 4-(chlorosulfonyl)-2-fluorobenzoate (12.37 g, 49.0 mmol) was added and the mixture stirred from 0 °C to room temperature for 1 h. The mixture was diluted with 1.0 N HCl (200 mL) and extracted with dichloromethane (150 mL x 2). The combined organic layers were washed with brine, dried with anhydrous sodium sulfate, filtered and concentrated under reduced pressure to afford crude product. The crude product was purified with Biotage SNAP 100 g column eluting with 0-30% 3:1 EtOAc-EtOH in heptane to afford methyl 2-fluoro-4-(*N*-(3-

methyloxetan-3-yl)sulfamoyl)benzoate (13.59 g, 44.8 mmol, 100 % yield) as a white solid. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 8.67 (s, 1H), 8.11 (t, $J = 7.37$ Hz, 1H), 7.76-7.82 (m, 1H), 7.69-7.76 (m, 1H), 4.56 (d, $J = 6.23$ Hz, 2H), 4.18 (d, $J = 6.75$ Hz, 2H), 3.90 (s, 3H), 1.42 (s, 3H).

[0276] Step 2: A mixture of *N,N*-diisopropylethylamine (16.23 mL, 93 mmol), 6-azaspiro[2.5]octane (6.22 g, 55.9 mmol), and methyl 2-fluoro-4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)benzoate (14.13 g, 46.6 mmol) in anhydrous 1,4-dioxane was stirred at 100 °C for 20 h. The mixture was cooled down to room temperature, quenched with water and extracted with ethyl acetate. The combined organic phases were washed with brine, dried and evaporated to dryness under reduced pressure. The crude product was purified using the Biotage SNAP 340 g column eluting with 0-40% 3:1 EtOAc-EtOH in heptane to get methyl 4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoate (14.15 g, 35.9 mmol, 77 % yield) as an off-white solid. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 8.42 (s, 1H), 7.72 (d, $J = 8.04$ Hz, 1H), 7.47 (d, $J = 1.56$ Hz, 1H), 7.36 (dd, $J = 1.82, 8.04$ Hz, 1H), 4.55 (d, $J = 5.97$ Hz, 2H), 4.14 (d, $J = 6.49$ Hz, 2H), 3.85 (s, 3H), 3.02-3.09 (m, 4H), 1.44-1.50 (m, 4H), 1.42 (s, 3H), 0.35 (s, 4H).

[0277] Step 3: A mixture of methyl 4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoate (14.15 g, 35.9 mmol) and lithium hydroxide monohydrate (22.58 g, 538 mmol) in THF-water-MeOH (1:1:1, 300 mL) was stirred at room temperature overnight. The mixture was concentrated under reduced pressure to partially remove the organic solvent. The solution was acidified with 2N HCl to reach pH < 3. The precipitated solid was filtered and dried in air to give 4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid (9.94 g, 26.1 mmol, 72.8 % yield) as a white solid. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 8.51 (s, 1H), 8.04 (d, $J = 8.04$ Hz, 1H), 7.89 (d, $J = 1.30$ Hz, 1H), 7.66 (dd, $J = 1.69, 8.17$ Hz, 1H), 4.55 (d, $J = 6.23$ Hz, 2H), 4.09-4.17 (m, 2H), 3.06-3.19 (m, 4H), 1.56 (t, $J = 5.19$ Hz, 4H), 1.40 (s, 3H), 0.36-0.46 (s, 4H).

[0278] Intermediate 16: 4-((1-(*tert*-Butoxycarbonyl)azetidin-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid



[0279] Step 1: A mixture of methyl 4-bromo-2-(6-azaspiro[2.5]octan-6-yl)benzoate (10.50 g, 32.4 mmol, Int. 12-2), DIPEA (8.37 mL, 64.8 mmol), Xantphos (1.874 g, 3.24 mmol) and $\text{Pd}_2(\text{dba})_3$ (2.97 g, 3.24


mmol) in 1,4-dioxane was bubbled with argon flow, then was added *tert*-butyl 3-mercaptoazetidine-1-carboxylate (7.66 mL, 40.5 mmol). The mixture was stirred at 100 °C for 18 h. The mixture was cooled down to room temperature, concentrated and purified through a Biotage SNAP eluting with a gradient of 0% to 25% 3:1 EtOAc-EtOH in heptane to provide *tert*-butyl 3-((4-(methoxycarbonyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)thio)azetidine-1-carboxylate (13.85 g, 32.0 mmol, 99 % yield) as a light-yellow sticky solid. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 7.55 (d, *J* = 8.04 Hz, 1H), 6.79 (s, 1H), 6.76 (d, *J* = 8.14 Hz, 1H), 4.34-4.43 (m, 2H), 4.27-4.34 (m, 1H), 3.79 (s, 3H), 3.70 (dd, *J* = 4.67, 8.56 Hz, 2H), 2.97-3.04 (m, 4H), 1.41-1.51 (m, 4H), 1.38 (s, 9H), 0.29-0.37 (m, 4H).

[0280] Step2: To a solution of *tert*-butyl 3-((4-(methoxycarbonyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)thio)azetidine-1-carboxylate (13.85 g, 32.0 mmol) in 1,4-dioxane (300 mL) was added Oxone monopersulfate (39.4 g, 64.0 mmol) in 150 mL water. The mixture was stirred at room temperature for 5 h and added 150 mL of ethyl acetate and 150 mL of water and the mixture was stirred for 10 minutes. The organic layer was separated, the aqueous layer was extracted with ethyl acetate. The combined organics were washed with brine, dried, filtered and concentrated. The crude material was purified through a Biotage SNAP 340 g column eluting with a gradient of 0% to 25% EtOAc-EtOH (3:1) in heptane to give *tert*-butyl 3-((4-(methoxycarbonyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfonyl)azetidine-1-carboxylate (12.77 g, 27.5 mmol, 86 % yield) as an off-white solid. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 7.75 (d, *J* = 8.04 Hz, 1H), 7.44-7.50 (m, 2 H), 4.48-4.55 (m, 1 H), 4.09 (br s, 2 H), 3.97-4.02 (m, 2H), 3.86 (s, 3H), 3.04-3.17 (m, 4 H), 1.42-1.51 (m, 4 H), 1.38 (s, 9 H), 0.35 (s, 4 H).

[0281] Step 3: A mixture of *tert*-butyl 3-((4-(methoxycarbonyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfonyl) azetidine-1-carboxylate (12.77 g, 27.5 mmol) and lithium hydroxide monohydrate (11.53 g, 275 mmol) in THF-water-MeOH (1:1:1, 230 mL) was stirred at room temperature for 15 h. The mixture was concentrated under reduced pressure to remove some organic solvent. The solution was acidified with 2N HCl to pH < 3. The precipitated solid was filtered and dried to give 4-((1-(*tert*-butoxycarbonyl)azetidin-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid (10.6 g, 23.53 mmol, 86 % yield) as an off-white solid. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 8.03 (d, *J* = 8.30 Hz, 1H), 7.93 (d, *J* = 1.82 Hz, 1H), 7.72 (dd, *J* = 1.69, 8.17 Hz, 1H), 4.48-4.60 (m, 1H), 4.10 (br. s., 2H), 3.99-4.06 (m, 3H), 3.14-3.22 (m, 4H), 1.49-1.59 (m, 4H), 1.38 (s, 9H), 0.41 (s, 4H).


[0282] INTERMEDIATES COMPOUNDS HAVING AR¹ AND AR² RINGS

[0283] Intermediate 17: *N*-(2-Chloro-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0284] 2,4,6-Tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (50 wt% solution in EtOAc, 12.50 mL, 21.00 mmol) and triethylamine (2.93 mL, 21.00 mmol) were added to a suspension of 4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid (3.0 g, 8.40 mmol, Int. 12) and 2-chloro-6-methylpyrimidin-4-amine (1.45 g, 10.08 mmol, Aurum Pharmtech Inc.) in DCE (20 mL). The mixture was heated to 85 °C for 24 h, then cooled to room temperature. Water (10 mL) was added, the layers were separated, and the aqueous layer was extracted with DCM (1x 10 mL). The combined organic extracts were dried over anhydrous MgSO₄, filtered, and concentrated in vacuo to give a solid. The solid was suspended in 1:1 EtOAc/heptane and filtered to provide *N*-(2-chloro-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (3.61 g, 7.48 mmol, 89% yield) as an off white solid. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 13.53 (br s, 1 H) 8.11 (s, 1 H) 7.92 (d, *J* = 8.29 Hz, 1 H) 7.70 (s, 1 H) 7.65 - 7.69 (m, 1 H) 3.05 (t, *J* = 5.39 Hz, 4 H) 2.53 (s, 3 H) 1.61 - 1.88 (m, 4 H) 0.44 (s, 4 H). *m/z* (ESI): 483.0 (M+H)⁺.

[0285] Intermediate 18: (*R*)-4-Bromo-*N*-(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide




[0286] Step 1: To a 100-mL round-bottomed flask was added 4-bromo-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid (0.9 g, 2.90 mmol, Int. 12-1), pyridine (0.657 mL, 8.12 mmol) and perfluorophenyl 2,2,2-trifluoroacetate (0.717 g, 3.77 mmol) in DCM (8 mL). The resultant mixture was stirred at rt for 16 h and the solvent was removed under vacuum to get crude product which was used for the next step without purification. *m/z* (ESI): 476 and 478 (M+1).

[0287] Step 2: To a 50-mL round-bottomed flask was added (*R*)-2-methyl-4-(4-methylpyrimidin-2-yl)morpholine (0.223 g, 1.15 mmol, Int. 2) dissolved in *N,N*-dimethylformamide (6 mL) followed by sodium hydride (0.084 g, 2.1 mmol) at rt under nitrogen atmosphere. The reaction mixture was stirred for 10 min and then it was treated with perfluorophenyl 4-bromo-2-(6-azaspiro[2.5]octan-6-yl)benzoate (0.5 g, 1.050 mmol) at RT under nitrogen atmosphere. The resultant reaction mixture was stirred at rt for additional 2 h. Reaction mixture was extracted with DCM (2 x 30 mL), separated, dried over anhydrous sodium sulphate and evaporated to dryness to get crude material. It was then absorbed onto a plug of silica gel and purified by silica gel flash column chromatography eluting with 30% to 50 % EtOAc/hexanes, to provide the title compound (0.20 g, 0.40 mmol, 38 % yield) as white solid. ¹H NMR (300 MHz, DMSO-


d_6) δ ppm 13.19 (bs, 1H), 7.99 (d, $J = 8.4$ Hz, 1H), 7.69 (s, 1H), 7.55 (d, $J = 8.4$ Hz, 1H), 7.36 (s, 1H), 4.45 (t, $J = 12$ Hz, 2H), 3.88 (m, 1H), 3.52-3.48 (m, 2H) 3.10-2.90 (m, 6H), 2.30 (s, 3H), 1.71-1.62 (m, 4H), 1.14 (d, $J = 6$ Hz, 3H), 0.36 (s, 4H). m/z (ESI): 500 and 502 (M+1).

[0288] Intermediate 19: *N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0289] 4-Iodo-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid (150.0 g, 420 mmol, Int. 12) was suspended in dichloromethane (1000 mL) under argon. Catalytic DMF (1.0 mL) was added followed by dropwise addition of a solution of thionyl chloride (54.6 g, 28 mL, 459 mmol, Sigma-Aldrich Corporation) in dichloromethane (500 mL) over 10 minutes. After stirring at ambient temperature for 30 minutes, the mixture was evaporated to dryness under reduced pressure. The crude was azeotroped with toluene (2 x 300 mL) and suspended in dichloromethane (300 mL) under argon. Tribasic potassium phosphate (267 g, 1.26 mol, Sigma-Aldrich Corporation) was added followed by a solution of 2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-amine (100 g, 438 mmol, Int. 4) and *N,N*-diisopropylethylamine (200 mL, 1.14 mol, Sigma-Aldrich Corporation) in DCM (300 mL, added over 5 minutes). The yellow mixture was stirred at ambient temperature for 3 hours then evaporated to dryness under reduced pressure. The crude solids were suspended in dichloromethane (1 L) and stirred for 10 minutes. The mixture was filtered through a frit and the solids washed with additional dichloromethane (2 x 100 mL). The solids were discarded, and the filtrate was evaporated to dryness under reduced pressure. The crude residue was suspended in acetonitrile (750 mL) and stirred at ambient temperature for 15 minutes. The suspension was filtered through a glass frit and the solids washed with additional acetonitrile (75 mL). The solids were dried under a stream of nitrogen to give *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (186 g, 328 mmol, 78 % yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 13.38 (br s, 1 H) 7.72 - 7.87 (m, 3 H) 7.39 (s, 1 H) 3.91 (br s, 4 H) 2.99 - 3.06 (m, 4 H) 2.32 (s, 3 H) 1.92 - 2.07 (m, 4 H) 1.62 - 1.85 (m, 4 H) 0.38 (s, 4 H). m/z (ESI): 568.0 (M-H) $^+$.


[0290] Intermediate 20: 4-Bromo-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-fluoro-6-(6-azaspiro[2.5]octan-6-yl)benzamide



[0291] Step 1: To a solution of 4-bromo-2,6-difluorobenzoic acid (3.0 g, 12.7 mmol, Apollo Scientific Ltd.) in THF (50 mL) was oxalyl chloride (1.7 mL, 19.0 mmol) followed by 1 drop of DMF. The mixture was stirred for 1 h then the solvent was removed in vacuo to give a solid that was taken directly to the next stage without further characterization. The solid was dissolved in DCM (50 mL) and anhydrous pyridine (4.31 mL, 50.6 mmol) followed by 2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-amine (2.89 g, 12.7 mmol, Int. 4) were added and the mixture was stirred for 16 h at room temperature. EtOAc (200 mL) was added and the mixture was washed with saturated NH_4Cl (1x), water (1x), brine (1x), dried over anhydrous MgSO_4 , filtered and concentrated in vacuo to give an oil. The oil was purified by silica gel chromatography, eluting with 0 to 40% EtOAc/heptane, to provide 4-bromo-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2,6-difluorobenzamide (1.36 g, 3.04 mmol, 24 % yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 11.24 (s, 1 H) 7.64 (d, $J=7.05$ Hz, 2 H) 7.19 - 7.37 (m, 1 H) 3.86 (br s, 4 H) 2.32 (s, 3 H) 1.98 (br d, $J=11.40$ Hz, 4 H). m/z (ESI): 447.0, 449.0 ($\text{M}+\text{H}$) $^+$.

[0292] Step 2: A mixture of 4-bromo-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2,6-difluorobenzamide (0.65 g, 1.45 mmol), 6-azaspiro[2.5]octane (0.18 g, 1.60 mmol, Wuxi App Tech), and DIPEA (0.31 mL, 1.74 mmol) in DMSO (2.5 mL) was heated to 100 °C for 8 h then cooled to room temperature. Water was added, and the resulting suspension was filtered and the solid obtained was dried. This solid was purified by silica gel chromatography, eluting with 0 to 15% EtOAc/heptane, to provide 4-bromo-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-fluoro-6-(6-azaspiro[2.5]octan-6-yl)benzamide (0.23 g, 0.43 mmol, 28.2% yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 10.86 (br s, 1 H) 7.30 (br s, 1 H) 7.21 (br d, $J=9.12$ Hz, 1 H) 7.10 (br s, 1 H) 3.88 (br s, 4 H) 3.05 (br s, 4 H) 2.32 (br s, 3 H) 1.77 - 2.04 (m, 4 H) 1.36 (br s, 4 H) 0.27 (s, 4 H). ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ ppm -94.88 (s, 1 F) -113.60 (s, 1 F). m/z (ESI): 538.2, 540.2 ($\text{M}+\text{H}$) $^+$.

[0293] Intermediate 21: 4-Amino-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0294] Step 1: Triethylamine (3.11 mL, 22.2 mmol) and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (50 wt% in EtOAc, 13.2 mL, 22.2 mmol) were added to a solution of 2,5-difluoro-4-nitrobenzenecarboxylic acid (1.5 g, 7.39 mmol, Combi-Blocks Inc.) and 2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-amine (1.69 g, 7.39 mmol, Int. 4) in DCE (15 mL) and the mixture was heated to 85 °C for 2 h then cooled to room temperature. Water (15 mL) was added, the resulting biphasic mixture was separated, and the organic layer was dried over anhydrous MgSO_4 , filtered, and concentrated in vacuo to give a solid. The solid was suspended in DCM (15 mL), filtered, and dried to provide *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2,5-difluoro-4-nitrobenzamide (3.05g, 4.55 mmol, 62% yield) as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 11.15 (s, 1 H) 8.27 (dd, $J=8.81, 5.91$ Hz, 1 H) 7.99 (dd, $J=10.57, 5.39$ Hz, 1 H) 7.23 (br s, 1 H) 3.85 (br s, 3 H) 2.33 (s, 4 H) 1.89 - 2.05 (m, 4 H). ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ ppm -95.11 (s, 1 F) -123.35 (s, 1 F) -123.40 (s, 1 F). m/z (ESI): 414.2 ($\text{M}+\text{H}$) $^+$.

[0295] Step 2: A mixture of *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2,5-difluoro-4-nitrobenzamide (1.0 g, 2.42 mmol) and palladium (10 wt. % on activated carbon, 0.45 g, 0.42 mmol) was placed under argon atmosphere and then EtOH (15 mL) was added, followed by ammonium formate (0.76 g, 12.1 mmol). The mixture was stirred at 75 °C for 10 min, then cooled to room temperature. The palladium was filtered off using CELITE® and the filtrate was concentrated in vacuo. The resulting residue was dissolved in EtOAc (10 mL) and the solution was washed with water (2 x 10 mL), brine (1 x 10 mL), dried over anhydrous MgSO_4 , filtered, and concentrated in vacuo to provide 4-amino-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2,5-difluorobenzamide (0.93 g, 2.19 mmol, 91% yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 9.68 - 9.80 (m, 1 H) 7.40 (dd, $J=11.61, 6.84$ Hz, 1 H) 7.28 (s, 1 H) 6.54 (dd, $J=13.48, 7.26$ Hz, 1 H) 6.26 (s, 2 H) 3.83 - 3.90 (m, 4 H) 2.30 (s, 3 H) 1.92 -

2.05 (m, 4 H). ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ ppm -95.07 (s, 1 F) -116.10 (s, 1 F) -140.24 (s, 1 F). m/z (ESI): 384.2 (M+H) $^+$.

[0296] Step 3: A mixture of 4-amino-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2,5-difluorobenzamide (0.80 g, 2.09 mmol), 6-azaspiro[2.5]octane (0.70 g, 6.26 mmol, Wuxi App Tech), and DIPEA (1.1 mL, 6.26 mmol) in NMP (4 mL) was heated to 200 °C in a microwave reactor for 4 h. Water (4 mL) was added and the resulting suspension was stirred for 30 min, filtered, and the collected solid was dried to give a white solid. This solid was dissolved in DCM, fused to silica gel and purified by silica gel chromatography, eluting with 0 to 50% EtOAc/heptane, to provide 4-amino-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-2-(6-azaspiro[2.5]octan-6-yl)benzamide (702 mg, 1.48 mmol, 70.9% yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 13.77 (s, 1 H) 7.64 (d, J = 12.85 Hz, 1 H) 7.38 (s, 1 H) 6.83 (d, J = 8.09 Hz, 1 H) 6.04 (s, 2 H) 3.90 (br t, J = 5.39 Hz, 4 H) 2.91 (br s, 4 H) 2.29 (s, 3 H) 1.88 - 2.04 (m, 4 H) 0.37 (s, 4 H). (Note: 4 protons not observed). ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ ppm -94.73 (s, 1 F) -138.31 (s, 1 F). m/z (ESI): 475.2 (M+H) $^+$.


[0297] Table 5: Intermediates 21-1 to 21-20 were prepared following similar procedures for Int. 17 to 21:

Int. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
21-1		(<i>R</i>)-4-Bromo- <i>N</i> -(2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	486.1/488.1
21-2		(<i>R</i>)-4-Iodo- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	548.2
21-3		4-Bromo- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	520.1/522.1
21-4		4-Bromo- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-(2-hydroxypropan-2-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	564.1/566.1
21-5		4-Bromo- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-ethylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	534.2/536.2

Int. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
21-6		4-Bromo- <i>N</i> -(6-cyclopropyl-2-(4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	546.1/548.1
21-7		4-Bromo-5-chloro- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	553.0/555.0
21-8		4-Bromo- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-methyl-2-(6-azaspiro[2.5]octan-6-yl)benzamide	534.2/536.2
21-9		(<i>R</i>)-4-Amino-5-fluoro- <i>N</i> -(2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	441.2
21-10		(<i>R</i>)-4-Amino-5-fluoro- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	455.2
21-11		4-Amino- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-methoxy-2-(6-azaspiro[2.5]octan-6-yl)benzamide	487.4
21-12		4-Bromo- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-3-fluoro-6-methylpyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	537.1/539.1
21-13		4-Bromo- <i>N</i> -(2-(4,4-difluorocyclohexyl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	519.1/521.1
21-14		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide	553.1

Int. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
21-15		4-Bromo- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	519.1/521.1
21-16		4-Iodo- <i>N</i> -(6-methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	561.0
21-17		4-Bromo- <i>N</i> -(2-chloro-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	435.0/437.0
21-18		<i>N</i> -(2-Chloro-6-methylpyrimidin-4-yl)-4-nitro-2-(6-azaspiro[2.5]octan-6-yl)benzamide	402.2
21-19		<i>N</i> -(2-Chloro-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	434.4
21-20		<i>N</i> -(2-Chloro-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	506.1

[0298] Intermediate 22: Ethyl 2-sulfamoylpropanoate




[0299] Step 1: To a solution of *N,N*-bis(4-methoxybenzyl)ethanesulfonamide (200.0 g, 572.0 mmol) in tetrahydrofuran (4000 mL) was added *n*BuLi (1.6 M in hexane, 608.0 mL, 973.0 mmol) at -78 °C slowly and stirred for 30 min. Ethyl carbonochloridate (92.0 mL, 973.0 mmol) in THF (50 mL) was added to the reaction mixture and stirred at -78 °C for 1 h. The reaction mixture was quenched with HCl (1.5 N, 3000 mL) and extracted with EtOAc (2 x 3000 mL). The organic extract was dried over sodium sulphate,

filtered and concentrated in reduced pressure to give the crude material of ethyl 2-(*N,N*-bis(4-methoxybenzyl)sulfamoyl)propanoate (250.0 g, 60% pure) as yellow oil. The ^1H -NMR showed desired peaks and proceeded to next step without any purification.

[0300] Step 2: To solution of ethyl 2-(*N,N*-bis(4-methoxybenzyl)sulfamoyl)propanoate (600.0 g, 1.4 mol) in trifluoroacetic acid (2.50 L, 32.45 mol) was added anisole (500.0 mL, 4.57 mol) and stirred at room temperature for 3 h. The reaction mixture was concentrated under reduced pressure, quenched with 10% cold aqueous NaHCO_3 solution (3 L) and extracted with EtOAc (2 x 3 L). The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography over silica gel using 25% ethyl acetate in hexanes to give a pale yellow solid (168 g) which was dissolved in DCM (1 L) and precipitated by the addition of hexanes (3000 mL). The solid was filtered, dried under vacuum to give the title compound (109.0 g, 42% yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 7.14 (s, 2H), 4.15 (q, $J = 7.1$ Hz, 2H), 3.98 (q, $J = 7.0$ Hz, 1H), 1.45 (d, $J = 7.0$ Hz, 3H), 1.21 (t, $J = 7.1$ Hz, 3H). m/z (ESI): 180.1 ($\text{M}-\text{H}$) $^+$.

[0301] Intermediate 23: 2-Hydroxypropane-1-sulfonamide




[0302] Step 1: Methanesulfonyl chloride (1.73 mL, 22.3 mmol) was added dropwise over 5 min to a 0 °C solution of bis(4-methoxybenzyl)amine (5.0 g, 19.4 mmol, Combi-Blocks Inc.) and triethylamine (8.12 mL, 58.3 mmol) in DCM (40 mL). The mixture was then stirred for 2 h at room temperature and then 1N HCl (50 mL) was added. The layers were separated, and the organic layer was washed with brine (1 x 50 mL), dried over anhydrous MgSO_4 , filtered, and then concentrated in vacuo to give a brown oil. The oil was dissolved in MeOH (50 mL) and partially concentrated in vacuo until a thick suspension formed. The suspension was stirred for 30 min, filtered, and the collected solid was dried in vacuo to provide *N,N*-bis(4-methoxybenzyl)methanesulfonamide (5.11 g, 15.2 mmol, 78% yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 7.19 (d, $J = 8.50$ Hz, 4 H) 6.90 (d, $J = 8.50$ Hz, 4 H) 4.19 (s, 4 H) 3.75 (s, 6 H) 2.89 (s, 3 H).

[0303] Step 2: *n*-Butyllithium (4.10 mL, 6.56 mmol) was added dropwise to a solution of *N,N*-bis(4-methoxybenzyl)methanesulfonamide (2.0 g, 5.96 mmol) in THF (15 mL) at -78 °C. The mixture was stirred for 10 min before acetaldehyde (0.37 mL, 6.56 mmol) was added dropwise. This mixture was stirred at -78 °C for 5 min, and then the -78 °C bath was replaced with a 0 °C bath. The mixture was stirred for 15 min and then the reaction was quenched with saturated NH_4Cl . EtOAc was added, the resulting biphasic mixture was separated, and the organic layer was dried over anhydrous MgSO_4 , filtered,

and concentrated in vacuo to give an oil. The oil was purified by silica gel chromatography, eluting with 0 to 70% EtOAc/heptane gradient, to provide 2-hydroxy-*N,N*-bis(4-methoxybenzyl)propane-1-sulfonamide (1.84 g, 4.85 mmol, 81% yield) as an oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.16 (d, *J*=8.50 Hz, 4 H) 6.80 - 6.92 (m, 4 H) 4.99 (d, *J*=5.18 Hz, 1 H) 4.15 - 4.28 (m, 4 H) 4.01 - 4.13 (m, 1H) 3.74 (s, 6 H) 3.16 (dd, *J*=13.99, 6.53 Hz, 1 H) 3.04 (dd, *J*=13.89, 5.39 Hz, 1 H) 1.12 - 1.27 (m, 3 H). *m/z* (ESI): 402.2 (M+Na)⁺.

[0304] Step 3: A mixture of 2-hydroxy-*N,N*-bis(4-methoxybenzyl)propane-1-sulfonamide (1.84 g, 4.85 mmol) and anisole (1.06 mL, 9.70 mmol) in TFA (10 mL) was stirred at room temperature for 2 h, then the volatiles were removed in vacuo. The resulting oil was purified by silica gel chromatography, eluting with 0 to 100% EtOAc/heptane gradient, to provide 2-hydroxypropane-1-sulfonamide (592 mg, 4.25 mmol, 88% yield) as a colorless oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 6.70 (s, 2 H) 4.02 - 4.13 (m, 1 H) 3.50 - 3.25 (br s, 1H) 3.07 - 3.15 (m, 1 H) 2.98 - 3.06 (m, 1 H) 1.20 (d, *J*=6.22 Hz, 3 H).

[0305] Example 1: *N*-(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide




[0306] A mixture of *N*-(2-chloro-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (200 mg, 0.46 mmol, Int. 21-19), 2-amino-2-methyl-1-propanol (180 uL, 1.80 mmol, Sigma-Aldrich, St Louis, MO), and DIPEA (200 uL, 1.14 mmol) in NMP (1 mL) was heated at 130 °C for 60 h. The reaction mixture was cooled to room temperature and quenched with water (10 mL), extracted with ethyl acetate (2 x 10 mL). The organic layer was washed with brine (500 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The crude material was adsorbed onto a plug of silica gel and purified by column chromatography over silica gel (60-120 mesh), eluting with 0% to 70% ethyl acetate in heptane to give the title compound (111 mg, 49%) as an off white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 11.65-11.96 (m, 1H), 8.07-8.25 (m, 1H), 7.82-7.87 (m, 1H), 7.75-7.81 (m, 1H), 7.32-7.41 (m, 1H), 6.13-6.24 (m, 1H), 4.73-4.91 (m, 1H), 3.42-3.54 (m, 2H), 3.30-3.33 (m, 3H), 3.01-3.16 (m, 4H), 2.18-2.28 (m, 3H), 1.55-1.75 (m, 4H), 1.25-1.42 (m, 6H), 0.27-0.40 (m, 4H). *m/z* (ESI): 487.4 (M+H)⁺.

[0307] Table 6: Examples 1-1 to 1-7 were prepared following a similar procedure as described for Example 1:

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
1-1		(<i>R</i>)- <i>N</i> -(2-((1-Hydroxypropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	474.0
1-2		(<i>S</i>)- <i>N</i> -(2-((1-Hydroxypropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	474.0
1-3		<i>N</i> -(2-((2-Hydroxyethyl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	460.4
1-4		(<i>R</i>)- <i>N</i> -(2-((2-Hydroxypropyl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	474.4
1-5		(<i>S</i>)- <i>N</i> -(2-((2-Hydroxypropyl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	474.4
1-6		<i>N</i> -(6-Methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	584.2
1-7		<i>N</i> -(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.4

[0308] Example 2: *N*-(2-(2-Hydroxypropan-2-yl)pyrimidin-4-yl)-4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide




[0309] To a solution of 4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid (150 mg, 0.394 mmol, Int. 15) and 2-(4-aminopyrimidin-2-yl)propan-2-ol (91 mg, 0.591 mmol, AstaTech, Bristol, PA, USA) in dichloromethane (2.6 mL) at 0 °C was added 1-propanephosphonic anhydride (50% in ethyl acetate, 0.469 mL, 0.789 mmol, Aldrich) followed by DIPEA (0.207 mL, 1.18 mmol, Aldrich). The resulting mixture was then stirred at room temperature overnight. Then, the mixture was then diluted with saturated NaHCO₃ (2 mL) followed by saturated NH₄Cl (7 mL). The mixture was then extracted with EtOAc (2 x 15 mL). The combined organic extracts were then dried over MgSO₄ and concentrated. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc/heptane) provided *N*-(2-(2-hydroxypropan-2-yl)pyrimidin-4-yl)-4-(*N*-(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (45 mg, 0.087 mmol) as a light yellow solid. ¹H NMR (DMSO-*d*₆) δ ppm 13.52 (br s, 1H), 8.76 (d, *J* = 5.6 Hz, 1H), 8.54 (br d, *J* = 3.5 Hz, 1H), 8.24 (d, *J* = 8.1 Hz, 1H), 8.08 (br d, *J* = 5.4 Hz, 1H), 7.86 (d, *J* = 1.0 Hz, 1H), 7.74 (dd, *J* = 8.1, 1.5 Hz, 1H), 4.96 (s, 1H), 4.55 (d, *J* = 6.0 Hz, 2H), 4.13 (d, *J* = 6.4 Hz, 2H), 3.08 (br t, *J* = 5.2 Hz, 4H), 1.70 (br s, 4H), 1.51 (s, 6H), 1.41 (s, 3H), 0.38 (s, 4H). *m/z* (ESI): 516.2 (M+H)⁺.

Table 7: Examples 2-1 to 2-8 were prepared following a similar procedure as described for example 2:

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
2-1		<i>N</i> -(2-(2-Hydroxypropan-2-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	530.2
2-2		(<i>R</i>)- <i>N</i> -(2-(2-Methylmorpholino)pyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	557.2
2-3		(<i>R</i>)- <i>N</i> -(6-Methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	571.2

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
2-4		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	576.2
2-5		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	591.2
2-6		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	520.0
2-7		(<i>R</i>)- <i>N</i> -(2-(2-Methylmorpholino)pyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	486.1
2-8		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	575.4


[0310] Example 3: (*R*)-4-((2-Hydroxyethyl)sulfonamido)-*N*-(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0311] Tribasic potassium phosphate (9.89 g, 46.6 mmol, Sigma-Aldrich), copper(I) iodide (0.710 g, 3.73 mmol, Sigma-Aldrich), 2-hydroxyethane-1-sulfonamide (1.166 g, 9.32 mmol, Wuxi Aptec, China), sarcosine (0.830 g, 9.32 mmol), and (*R*)-4-iodo-*N*-(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (5.1 g, 9.32 mmol, Int. 21-2) were combined in a three neck flask under argon. Dry, degassed DMF (20 mL) was added and the mixture heated to 110°C with overhead stirring for 45 minutes. The reaction was cooled to ambient temperature and saturated ammonium

chloride (75 mL), water (200 mL) and ethyl acetate (200 mL) were added. The phases were mixed and separated and the organic dried with brine (75 mL) before evaporating to dryness under reduced pressure. The crude solids were stirred in boiling ethanol (15 mL) for 10 minutes then cooled to ambient temperature and filtered through a sintered glass frit. The solids were dried on the frit then suspended in water (75 mL) and heated to 80°C. After 10 minutes, the mixture was cooled to ambient temperature and filtered through a sintered glass frit. The solids were dried under a stream of nitrogen to give the title compound (3.3 g, 6.06 mmol, 65.0 % yield) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.23 (bs, 1H), 10.26 (bs, 1H), 8.05 (m, 1H), 7.37 (s, 1H), 7.26 (s, 1H), 7.13 (s, 1H), 4.95 (bs, 1H), 4.50-4.42 (m, 2H), 3.84 (m, 1H), 3.76-3.74 (m, 2H), 3.51-3.45 (m, 2H), 3.00-2.82 (m, 6H), 2.61-2.58 (m, 2H), 2.31 (s, 3H), 1.91-1.65 (m, 4H), 1.17 (d, *J* = 6.0 Hz, 3H), 0.39 (s, 4H). *m/z* (ESI): 545.2 (M+H)⁺.


[0312] Example 4: *N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0313] A mixture of 2-hydroxyethane-1-sulfonamide (1.28 g, 10.3 mmol, Wuxi AppTec), copper(I) iodide (0.49 g, 2.56 mmol), potassium phosphate tribasic (5.44 g, 25.6 mmol), and Sarcosine (0.48 g, 5.13 mmol) in a 100 mL round bottom flask was placed under argon atmosphere. Anhydrous DMF (20 mL) was added and the mixture was warmed to 50 °C for 5 minutes. *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (2.91 g, 5.13 mmol, Int. 19) was added as a solid and the mixture was heated to 100 °C and stirred for 2 h, then cooled to room temperature. EtOAc (20 mL) and water (20 mL) were added, the resulting biphasic mixture was separated, and the aqueous layer was extracted with EtOAc (3x). The combined organic extracts were then washed with water (2x), 9:1 NH₄Cl/NH₄OH (aq), brine, dried over anhydrous MgSO₄, filtered, and concentrated in vacuo to give an oil. The oil was purified by silica gel chromatography, eluting with 0 to 50% EtOAc/heptane gradient, then 50% EtOAc/heptane isocratic elution, to provide an off-white solid. This solid was suspended in methanol, filtered, and dried to give a white solid. This solid was then suspended in water, stirred for 24 h, filtered, and dried in vacuo to provide *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (1.55 g, 2.75 mmol, 54% yield) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.37 (s, 1 H) 10.03 - 10.52 (m, 1 H) 8.06 (d, *J* = 8.71 Hz, 1 H) 7.41 (s, 1 H) 7.28 (d, *J* = 1.87 Hz, 1 H) 7.15 (dd, *J* = 8.71, 1.87 Hz, 1 H) 4.73 - 5.14 (m, 1 H) 3.92 (br t, *J* = 5.39 Hz, 4 H) 3.77 (t, *J* = 6.43 Hz, 2 H) 3.34 - 3.40 (m, 2 H) 2.98 (br t, *J* = 4.56 Hz, 4 H) 2.32 (s, 3 H) 1.93 - 2.07 (m, 4 H) 1.58 - 1.85 (m, 4 H) 0.40 (s, 4 H). ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ ppm -94.74 (s, 1 F). *m/z* (ESI): 565.2 (M+H)⁺.

[0314] Examples 5-1 and 5-2: (*R*)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide and (*S*)-*N*-(2-(4,4-

di(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0315] Step 1: A mixture of ethyl 2-sulfamoylpropanoate (1.44 g, 7.93 mmol, Int. 22), copper(I) iodide (0.503 g, 2.64 mmol, Strem), sarcosine (0.47 g, 5.29 mmol, Sigma-Aldrich Corporation), and potassium phosphate (4.49 g, 21.2 mmol) in DMF (15 mL) was placed under argon atmosphere and warmed to 50 °C for 5 min. *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (3.0 g, 5.29 mmol, Int. 19) was added and the mixture was heated to 100 °C for 3 h, then cooled to room temperature. EtOAc (50 mL), IPA (5 mL) and water (50 mL) were added and the mixture was stirred vigorously for 5 min. The resulting biphasic mixture was transferred to a separatory funnel and the layers were separated. The aqueous layer was extracted with EtOAc (2 x 20 mL) and the combined extracts were then washed with water (2 x 50 mL), 9:1 NH₄Cl/NH₄OH (1 x 50 mL), dried over anhydrous MgSO₄, filtered, and concentrated in vacuo to give an oil. The crude oil was purified by silica gel chromatography using a Redi-Sep pre-packed silica gel column (80 g), eluting with 0 to 50% EtOAc/heptane gradient, to provide ethyl 2-(*N*-(4-((2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)carbamoyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfamoyl)propanoate (2.76 g, 4.45 mmol, 84 % yield) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.35 (s, 1 H) 10.69 (br s, 1 H) 8.07 (d, *J*=8.71 Hz, 1 H) 7.40 (s, 1 H) 7.31 (d, *J*=1.87 Hz, 1 H) 7.17 (dd, *J*= 8.60, 1.97 Hz, 1 H) 4.06 (qd, *J*=7.08, 4.87 Hz, 2 H) 3.92 (br t, *J*=5.49 Hz, 4 H) 2.98 (br t, *J*=4.77 Hz, 4 H) 2.32 (s, 3 H) 1.85 - 2.06 (m, 5 H) 1.73 (br s, 4 H) 1.48 (d, *J*=6.84 Hz, 3 H) 1.14 (t, *J*=7.05 Hz, 3 H) 0.39 (s, 4 H). ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ ppm -94.75 (s, 1 F). *m/z* (ESI): 621.2 (M+H)⁺.

[0316] Step 2: To a 250 mL round bottom flask was added ethyl 2-(*N*-(4-((2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)carbamoyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfamoyl)propanoate (10.39 g, 16.74 mmol) and lithium borohydride solution, (2.0M in THF, 16.7 mL, 33.5 mmol, Sigma-Aldrich Corporation) in THF (100 mL). Methanol (4.29 mL, 134 mmol) was added slowly over 5 min and the resulting solution was stirred at room temperature for 30 min. 1 N HCl (20 mL) was slowly added followed by EtOAc (20 mL) and the resulting biphasic mixture was transferred to a separatory funnel and the phases were separated. The aqueous layer was extracted with EtOAc (1 x 25 mL) and the combined

extracts were washed with saturated NaHCO₃ (1 x 50 mL), brine (1 x 50 mL), dried over anhydrous MgSO₄, filtered, and concentrated to give 8.9 g racemic mixture. This material was separated by preparative SFC using a Chiral Tech AD column (250 X 30 mm, 5mm) with a mobile phase of 85% Liquid CO₂ and 15% MeOH with 0.2% TEA using a flowrate of 150 mL/min to give:

[0317] Example 5-1: (R)-N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. First eluting peak (3.50 g, 6.05 mmol, 36.1 % yield, >99%ee). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.36 (s, 1 H) 8.05 (d, *J* = 8.50 Hz, 1 H) 7.40 (s, 1 H) 7.31 (d, *J* = 1.87 Hz, 1 H) 7.17 (dd, *J* = 8.71, 2.07 Hz, 1H) 3.88 - 3.97 (m, 4 H) 3.84 (dd, *J* = 10.99, 4.35 Hz, 1 H) 3.37 - 3.54 (m, 1 H) 3.25 - 3.30 (m, 1 H) 2.97 (br t, *J* = 4.77 Hz, 4 H) 2.32 (s, 3 H) 1.84 - 2.06 (m, 4 H) 1.57 - 1.84 (br s, 4 H) 1.30 (d, *J* = 6.84 Hz, 3 H) 0.39 (s, 4 H). 2 exchangeable protons not observed. ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ ppm -94.74 (s, 1 F). *m/z* (ESI): 579.2 (M+H)⁺.

[0318] Example 5-2: (S)-N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. Second eluting peak (2.66 g, 4.60 mmol, 27.5 % yield, 98.9%ee). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.35 (s, 1 H) 8.05 (d, *J* = 8.50 Hz, 1 H) 7.40 (s, 1 H) 7.31 (d, *J* = 2.07 Hz, 1 H) 7.17 (dd, *J* = 8.60, 1.97 Hz, 1H) 3.88 - 3.97 (m, 4 H) 3.84 (dd, *J* = 10.99, 4.35 Hz, 1 H) 3.50 (dd, *J* = 10.99, 7.46 Hz, 1 H) 3.25 - 3.32 (m, 1 H) 2.97 (br t, *J* = 4.77 Hz, 4 H) 2.31 (s, 3H) 1.83 - 2.06 (m, 4 H) 1.73 (br s, 4 H) 1.30 (d, *J* = 6.84 Hz, 3 H) 0.39 (s, 4 H). 2 exchangeable protons not observed. ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ ppm -94.75 (s, 1 F). *m/z* (ESI): 579.2 (M+H)⁺. The stereochemistry was arbitrarily assigned.

Table 8: Examples 6-1 to 6-40 were prepared following a similar procedure for Examples 3 to 5-2:

Ex.#	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
6-1		4- ((Fluoromethyl)sulfonamido)- N-(2-(4-fluoropiperidin-1-yl)- 6-methylpyrimidin-4-yl)-2-(6- azaspiro[2.5]octan-6- yl)benzamide	535.3
6-2		N-(2-(4,4-Difluoropiperidin-1- yl)-6-methylpyrimidin-4-yl)-4- ((fluoromethyl)sulfonamido)- 2-(6-azaspiro[2.5]octan-6- yl)benzamide	553.3
6-3		(R)-4- ((Fluoromethyl)sulfonamido)- N-(6-methyl-2-(2- methylmorpholino)pyrimidin- 4-yl)-2-(6-azaspiro[2.5]octan- 6-yl)benzamide	533.2

Ex.#	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
6-4		(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- N-(2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	531.2
6-5		(<i>R</i>)-N-(6-Methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-4-(methylsulfonamido)- 2-(6-azaspiro[2.5]octan-6-yl)benzamide	515.2
6-6		(<i>R</i>)-4-(Ethylsulfonamido)-N-(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	529.2
6-7		N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(ethylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	549.2
6-8		(<i>R</i>)-4-((3-Hydroxypropyl)sulfonamido)- N-(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.2
6-9		N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(3,3-dioxido-1,3,4-oxathiazinan-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	577.2
6-10		4-(Cyclopentanesulfonamido)- N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	589.3
6-11		4-(Cyclobutanesulfonamido)- N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	575.2

Ex.#	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
6-12		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(oxetane-3-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	577.2
6-13		4-(Cyclobutanesulfonamido)- <i>N</i> -(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	547.2
6-14		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1,1-dimethylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	577.3
6-15		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((3-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	579.2
6-16		4-(Cyclopropanesulfonamido)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	561.2
6-17		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-methoxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	579.9
6-18		4-((Cyclopropylmethyl)sulfonamido)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	575.2
6-19		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(((3-hydroxyoxetan-3-yl)methyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	607.2


Ex.#	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
6-20		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1,1-dimethylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	593.2
6-21		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(methylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	535.2
6-22		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	563.2
6-23-1		4-(((<i>S</i>)-2-Hydroxy-1-methylethyl)sulfonamido)- <i>N</i> -(6-methyl-2-((<i>R</i>)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.2
6-23-2		4-(((<i>R</i>)-2-Hydroxy-1-methylethyl)sulfonamido)- <i>N</i> -(6-methyl-2-((<i>R</i>)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.2
6-24-1		4-(((<i>S</i>)-2-Hydroxy-1-methylethyl)sulfonamido)- <i>N</i> -(2-((<i>R</i>)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	545.2
6-24-2		4-(((<i>R</i>)-2-hydroxy-1-methylethyl)sulfonamido)- <i>N</i> -(2-((<i>R</i>)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	545.2
6-25-1		(<i>S</i>)- <i>N</i> -(2-(4-fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	561.2

Ex.#	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
6-25-2		(<i>R</i>)- <i>N</i> -(2-(4-Fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	561.2
6-26-1		(<i>S</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	565.2
6-26-2		(<i>R</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	565.2
6-27-1		4-(((<i>R</i>)-2-Hydroxypropyl)sulfonamido)- <i>N</i> -(6-methyl-2-((<i>R</i>)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.2
6-27-2		4-(((<i>S</i>)-2-Hydroxypropyl)sulfonamido)- <i>N</i> -(6-methyl-2-((<i>R</i>)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.2
6-28-1		(<i>R</i>)-4-((2-Hydroxypropyl)sulfonamido)- <i>N</i> -(6-methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	572.2
6-28-2		(<i>S</i>)-4-((2-Hydroxypropyl)sulfonamido)- <i>N</i> -(6-methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	572.2
6-29		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	550.2

Ex.#	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
6-30		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyridin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	564.2
6-31-1		(<i>R</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-(1-hydroxyethyl)cyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	605.2
6-31-2		(<i>S</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-(1-hydroxyethyl)cyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	605.2
6-32-1		(<i>R</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	579.1
6-32-2		(<i>S</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	579.1
6-33-1		4-(((<i>S</i>)-2-Hydroxypropyl)sulfonamido)- <i>N</i> -(2-((<i>R</i>)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	545.2
6-33-2		4-(((<i>R</i>)-2-hydroxypropyl)sulfonamido)- <i>N</i> -(2-((<i>R</i>)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	545.2
6-34		5-Chloro- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	599.2

Ex.#	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
6-35		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-5-methyl-2-(6-azaspiro[2.5]octan-6-yl)benzamide	579.2
6-36		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-5-methoxy-2-(6-azaspiro[2.5]octan-6-yl)benzamide	595.2
6-37		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-(2-hydroxypropan-2-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	609.1
6-38		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-ethylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	579.2
6-39		<i>N</i> -(6-Cyclopropyl-2-(4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	591.2
6-40		(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-(2-hydroxypropan-2-yl)-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	589.3

[0319] Example 7: *N*-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0320] Step 1: A solution of *N*-(2-chloro-6-methylpyrimidin-4-yl)-4-nitro-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.3 g, 0.747 mmol, Int. 21-18), 3,3-difluoroazetidine hydrochloride (0.145 g, 1.120 mmol, Combi-Blocks) and DIPEA (0.261 mL, 1.49 mmol) in DMF (0.5 mL) and ethanol (1 mL) was heated at 80 °C for 4 h. Then the reaction mixture was quenched with water (50 mL) and extracted with ethyl acetate (3 × 50 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered, and concentrated. The concentrate was purified by flash column chromatography eluting with 30% to 50% ethyl acetate in petroleum ether to provide *N*-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-nitro-2-(6-azaspiro[2.5]octan-6-yl)benzamide (180 mg, 0.393 mmol, 52.6 % yield) as yellow solid. ¹H NMR (400 MHz, Chloroform-*d*) δ ppm 8.47 (d, *J* = 8.7 Hz, 1 H), 8.25 (d, *J* = 2.2 Hz, 1 H), 8.17 (dd, *J* = 8.7, 2.2 Hz, 1 H), 7.68 (s, 1 H), 3.79 – 3.66 (m, 4 H), 3.17 (t, *J* = 5.4 Hz, 4 H), 2.60 (s, 3 H), 1.28 (s, 4 H), 0.50 (s, 4 H). *m/z* (ESI): 459.2 (M+H)⁺.


[0321] Step 2: To a solution of *N*-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-nitro-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.18 g, 0.39 mmol) in ethanol (8 mL) and water (8 mL) were added iron powder (0.066 g, 1.18 mmol) and ammonium chloride (0.063 g, 1.18 mmol). Then the mixture was heated at 90 °C for 3 h before it was filtered through a bed of CELITE® and was washed with ethyl acetate (3 × 100 mL). The filtrate was washed with brine, dried over Na₂SO₄, filtered, and concentrated to provide 4-amino-*N*-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.12 g, 0.280 mmol, 71.3 % yield) as pale yellow solid. It was used directly for the next step without further purification. *m/z* (ESI): 429.2 (M+H)⁺.

[0322] Step 3: To a solution of 4-amino-*N*-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.120 g, 0.280 mmol) in DCM (5 mL) were added Et₃N (0.078 mL, 0.560 mmol) and methyl 2-(chlorosulfonyl)acetate (0.058 g, 0.336 mmol, Combi-blocks) at 0 °C. The mixture was stirred at room temperature for 4 h before it was quenched with water (50 mL) and was extracted with DCM (3 × 50 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated to afford methyl 2-(*N*-(4-((2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)carbamoyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfamoyl)acetate (140 mg, 0.25 mmol, 89 % yield) as pale yellow solid. It was used for the next step without further purification.

m/z (ESI): 565.2 (M+H)⁺.

[0323] Step 4: A solution of methyl 2-(*N*-(4-((2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)carbamoyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfamoyl)acetate (130 mg, 0.230 mmol) in THF (5 mL) was treated with LiBH₄ (230 μL, 0.460 mmol) at -30 °C. The reaction mixture was stirred for 30 min at 0 °C before it was quenched with a saturated aqueous solution of NH₄Cl (50 mL) at 0 °C and was extracted with EtOAc (2 × 10 mL). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered and concentrated. The concentrate was purified by flash column chromatography eluting with a gradient of 20% to 100% EtOAc in hexanes to provide *N*-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (60 mg, 0.112 mmol, 48.6 % yield) as white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.55 (s, 1 H), 10.28 (s, 1 H), 8.05 (d, *J*=8.7 Hz, 1 H), 7.51 (s, 1 H), 7.28 (d, *J*=2.2 Hz, 1 H), 7.14 (dd, *J*=8.6, 2.2 Hz, 1 H), 4.95 (s, 1 H), 4.45 (d, *J*=12.3 Hz, 4 H), 3.76 (t, *J*=6.4 Hz, 2 H), 3.64 (s, 1 H), 3.57 (s, 1 H), 2.97 (t, *J*=5.4 Hz, 4 H), 2.34 (s, 3 H), 1.74 (br s, 4 H), 0.40 (s, 4 H). *m/z* (ESI): 537.2 (M+H)⁺.

[0324] Examples 8-1 and 8-2: (*R*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide and (*S*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0325] Step 1: Methyl 2-(chlorosulfonyl)propanoate (177 mg, 0.95 mmol, Enamine) was added to a solution of 4-amino-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.41 g, 0.86 mmol, Int. 21) and triethylamine (0.18 mL, 1.30 mmol) in DCM (3 mL) at 0 °C. The mixture was stirred for 1 h then the mixture was concentrated in vacuo and then purified by silica gel chromatography using a Redi-Sep pre-packed silica gel column (12 g), eluting with 25% EtOAc/heptane gradient, to provide methyl 2-(*N*-(4-((2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)carbamoyl)-2-fluoro-5-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfamoyl)propanoate (0.54 g, 0.48 mmol, 54.8% yield) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.71 (s, 1 H) 10.66 (br s, 1 H) 7.88 (d, *J*=11.61 Hz, 1 H) 7.60 (d, *J*=7.26 Hz, 1 H) 7.39 (s, 1 H) 4.38 (d, *J*=7.05 Hz, 1 H) 3.88 - 3.96 (m, 4 H) 3.57 (s, 3 H) 2.98 (br t, *J*=4.56 Hz, 4 H) 2.33 (s, 3 H) 1.92 - 2.07 (m, 4 H) 1.57 - 1.91 (m, 4 H) 1.52 (d, *J*=6.84 Hz, 3 H) 0.40 (s, 4 H). ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ ppm -94.77 (s, 1 F) -126.39 (s, 1 F). *m/z* (ESI): 625.2 (M+H)⁺.

[0326] Step 2: Methanol (0.10 mL, 2.52 mmol) was added dropwise to a THF (2.5 mL) solution of methyl 2-(*N*-(4-((2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)carbamoyl)-2-fluoro-5-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfamoyl)propanoate (192 mg, 0.31 mmol) and lithium borohydride (2.0 M in THF, 0.35 mL, 0.69 mmol). The mixture was stirred for 30 minutes and then aqueous NH₄Cl was added. The product was extracted into EtOAc (2x), and the combined extracts were dried over anhydrous MgSO₄, filtered, and concentrated in vacuo to give racemic product as a solid. This material was separated by preparative SFC using a OD column (250 x 21 mm, 5 mm) and OD column (150 X 21 mm, 5 mm) with a mobile phase of 90% Liquid CO₂ and 10% EtOH/0.2% triethylamine using a flowrate of 80 mL/min to give:


[0327] Example 8-1: (*R*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. First eluting peak (94 mg, 0.16 mmol, 33.2% yield, >99% ee). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.74 (s, 1 H) 7.82 (d, *J*=11.82 Hz, 1 H) 7.58 (d, *J*=7.26 Hz, 1 H) 7.39 (s, 1 H) 3.77 - 3.96 (m, 6 H) 3.37 - 3.54 (m, 1 H) 3.22 - 3.30 (m, 1 H) 2.88 - 3.05 (m, 4 H) 2.32 (s, 3 H) 1.92 - 2.05 (m, 4 H) 1.73 (br s, 4 H) 1.31 (d, *J*=6.84 Hz, 3 H) 0.40 (s, 4 H). ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ ppm -94.77 (s, 1 F) -127.36 (s, 1 F). *m/z* (ESI): 597.2 (M+H)⁺.

[0328] Example 8-2: (*S*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. Second eluting peak (102 mg, 0.17 mmol, 36.1% yield, 98.4% ee). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.74 (s, 1 H) 7.82 (d, *J*=11.82 Hz, 1 H) 7.58 (d, *J*=7.26 Hz, 1 H) 7.39 (s, 1 H) 3.77 - 3.96 (m, 6 H) 3.37 - 3.54 (m, 1 H) 3.22 - 3.30 (m, 1 H) 2.88 - 3.05 (m, 4 H) 2.32 (s, 3 H) 1.92 - 2.05 (m, 4 H) 1.73 (br s, 4 H) 1.31 (d, *J*=6.84 Hz, 3 H) 0.40 (s, 4 H). ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ ppm -94.76 (s, 1 F) -127.73 (s, 1 F). *m/z* (ESI): 597.2 (M+H)⁺. The stereochemistry was arbitrarily assigned.

[0329] Table 9: Examples 9-1 to 9-2 were prepared following similar procedure for Examples 8-1 and 8-2;

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
9-1		(<i>R</i>)-5-Fluoro-4-((2-hydroxyethyl)sulfonamido)-N-(2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	549.2
9-2		(<i>R</i>)-5-Fluoro-4-((2-hydroxyethyl)sulfonamido)-N-(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	563.2

[0330] Examples 10-1 and 10-2: (*R*)-*N*-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide and (*S*)-*N*-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0331] Step 1: A mixture of *N*-(2-chloro-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (2.0 g, 4.14 mmol, Int. 17), 3,3-difluoroazetidine hydrochloride (1.07 g, 8.29 mmol, Combi-Blocks Inc.), and potassium carbonate (1.72 g, 12.4 mmol, Combi-Blocks, Inc.) in NMP (10 mL) was heated to 90 °C for 24 h. The mixture was cooled to room temperature, EtOAc (10 mL) was added, and then the mixture was washed with water (1 x 10 mL), 1N HCl (1 x 10 mL) and brine (1 x 10 mL). The mixture was then dried over anhydrous MgSO₄, filtered, and concentrated in vacuo to give a solid. The solid was then suspended in MeOH and filtered, then dried in vacuo to provide *N*-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (1.16 g, 2.15 mmol, 51.9% yield) as a light brown solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.53 (br s, 1 H) 7.79 - 7.86 (m, 2 H) 7.73 - 7.77 (m, 1 H) 7.49 (s, 1 H) 4.43 (t, *J*=12.44 Hz, 4 H) 3.02 (br t, *J*= 5.08 Hz, 4 H) 2.31 - 2.38 (m, 3

H) 1.65 - 1.82 (m, 4 H) 0.39 (s, 4 H). ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ ppm -99.09 (s, 1 F). m/z (ESI): 540.0 (M+H) $^+$.

[0332] Step 2: A mixture of 2-hydroxypropane-1-sulfonamide (206 mg, 1.48 mmol, Int. 23), copper(I) iodide (71 mg, 0.37 mmol), Sarcosine (66 mg, 0.74 mmol), and potassium phosphate (787 mg, 3.71 mmol) was placed under argon atmosphere, taken up in anhydrous DMF (3 mL), and warmed to 50 °C for 5 min. *N*-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.40 g, 0.74 mmol) was added in one portion and the mixture was heated to 100 °C for 2.5 h, then cooled to room temperature. Water was added, and the product was extracted into EtOAc (2x). The combined extracts were washed with water (2x), 9:1 saturated $\text{NH}_4\text{Cl}/\text{NH}_4\text{OH}$ (1x), dried over anhydrous MgSO_4 , filtered, and concentrated in vacuo to give racemic product as an oil. This material was separated by preparative SFC using an IF column (250 x 301 mm, 5 mm) with a mobile phase of 75% Liquid CO_2 and 25% MeOH using a flowrate of 130 mL/min to give:

[0333] Example 10-1: (R)-N-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. First eluting peak (88 mg, 0.16 mmol, 21.6% yield, >99%ee). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 13.54 (s, 1 H) 8.05 (d, J = 8.71 Hz, 1 H) 7.51 (s, 1 H) 7.27 (d, J = 1.87 Hz, 1 H) 7.14 (dd, J = 8.71, 1.87 Hz, 1H) 4.44 (t, J = 12.44 Hz, 4 H) 4.07 - 4.15 (m, 1 H) 3.22 - 3.29 (m, 2 H) 2.97 (br t, J = 4.87 Hz, 4 H) 2.34 (s, 3 H) 1.74 (br s, 4 H) 1.19 (d, J = 6.22 Hz, 3H) 0.40 (s, 4 H). 2 exchangeable protons not observed. ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ ppm -99.08 (s, 1 F). m/z (ESI): 551 (M+H) $^+$.

[0334] Example 10-2: (S)-N-(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. Second eluting peak (89 mg, 0.162 mmol, 21.8% yield, 99%ee). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 13.55 (s, 1 H) 8.05 (d, J = 8.71 Hz, 1 H) 7.51 (s, 1 H) 7.26 (d, J = 1.87 Hz, 1 H) 7.14 (dd, J = 8.71, 1.87 Hz, 1H) 4.43 (t, J = 12.44 Hz, 4 H) 4.11 (d, J = 6.01 Hz, 1 H) 3.21 - 3.31 (m, 2 H) 2.97 (br t, J = 4.87 Hz, 4 H) 2.34 (s, 3 H) 1.53 - 2.01 (m, 4 H) 1.19 (d, J = 6.43 Hz, 3 H) 0.40 (s, 4 H). 2 exchangeable protons not observed. ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ ppm -99.09 (s, 1 F). m/z (ESI): 551 (M+H) $^+$. The stereochemistry was arbitrarily assigned.

[0335] Table 10: Examples 11-1 to 11-83 were prepared following similar procedures for Example 10:

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
11-1		(S)-4-((2-Hydroxyethyl)sulfonamido)-N-(2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	531.2
11-2		4-((2-Hydroxyethyl)sulfonamido)-N-(2-(2-isopropyl-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	488.1

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
11-3		<i>N</i> -(2-Cyclopropyl-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	486.1
11-4		<i>N</i> -(2-Cyclobutoxy-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	516.2
11-5		<i>N</i> -(2-(3-Fluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	519.2
11-6		<i>N</i> -(2-(3-Fluoroazetidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	505.1
11-7		<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	523.1
11-8		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	543.2
11-9		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(6-oxa-1-azaspiro[3.3]heptan-1-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	543.2
11-10		<i>N</i> -(2-(3-(Difluoromethoxy)azetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	567.2

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
11-11		<i>N</i> -(2-(1,1-Difluoro-5-azaspiro[2.3]hexan-5-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	563.2
11-12		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(3-(trifluoromethyl)azetidin-1-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	569.2
11-13		<i>N</i> -(2-(3-Cyanoazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	526.2
11-14		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-(3-(2-hydroxypropan-2-yl)azetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.2
11-15		<i>N</i> -(2-(3-Hydroxy-3-methylazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	531.2
11-16		<i>N</i> -(2-(3-Cyclopropyl-3-hydroxyazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	557.2
11-17		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(5-azaspiro[2.3]hexan-5-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	527.2
11-18		<i>N</i> -(2-(3-Hydroxy-3-(trifluoromethyl)azetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	585.2

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
11-19		<i>N</i> -(2-(Azetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	501.2
11-20		<i>N</i> -(2-(3-Hydroxyazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	517.2
11-21		(<i>S</i>)- <i>N</i> -(2-(3-(1-Hydroxyethyl)azetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	545.2
11-22		4-(Cyclopropanesulfonamido)- <i>N</i> -(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	533.2
11-23		<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-(oxetane-3-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	549.2
11-24		<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-(methylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	507.1
11-25		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(5-azaspiro[2.4]heptan-5-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	541.2
11-26		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	515.2

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
11-27		(<i>R</i>)- <i>N</i> -(2-(3-Fluoro-3-methylpyrrolidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	547.3
11-28		<i>N</i> -(2-(2-Azabicyclo[3.1.0]hexan-2-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	527.2
11-29		<i>N</i> -(2-((1 <i>R</i> ,5 <i>S</i>)-3-Azabicyclo[3.1.0]hexan-3-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	527.2
11-30		<i>N</i> -(2-(3,3-Difluoropyrrolidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	551.2
11-31		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(6-azaspiro[2.5]octan-6-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	555.2
11-32		<i>N</i> -(2-((1 <i>R</i>)-1-Hydroxy-3-azabicyclo[3.1.0]hexan-3-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	543.2
11-33		<i>N</i> -(2-(4-Cyanopiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	554.2
11-34		(<i>R</i>)- <i>N</i> -(2-(3-Cyanopiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	554.2

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
11-35		(<i>S</i>)- <i>N</i> -(2-(3-Cyanopiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	554.2
11-36		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(piperidin-1-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	529.3
11-37		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-(4-hydroxypiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	545.2
11-38		<i>N</i> -(2-(4-Hydroxy-4-methylpiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.2
11-39		<i>N</i> -(2-(3,3-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	565.2
11-40		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(7-azaspiro[3.5]nonan-7-yl)benzamide	579.2
11-41		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(4,4-dimethylpiperidin-1-yl)-4-((2-hydroxyethyl)sulfonamido)benzamide	567.2
11-42		<i>N</i> -(2-((1 <i>R</i>)-1-Hydroxy-3-azabicyclo[4.1.0]heptan-3-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	557.2

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
11-43		<i>N</i> -(2-((1 <i>R</i> ,6 <i>R</i>)-3-Azabicyclo[4.1.0]heptan-3-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	541.2
11-44		<i>N</i> -(2-((1 <i>S</i> ,6 <i>R</i>)-2-Azabicyclo[4.1.0]heptan-2-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	541.2
11-45		(<i>S</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	545.2
11-46		<i>N</i> -(2-(2,2-Dimethylmorpholino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.3
11-47		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(4-oxa-7-azaspiro[2.5]octan-7-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	557.3
11-48		<i>N</i> -(2-((3 <i>S</i> ,5 <i>R</i>)-3,5-Dimethylmorpholino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.3
11-49		<i>N</i> -(2-((3 <i>S</i> ,5 <i>S</i>)-3,5-Dimethylmorpholino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.3
11-50		<i>N</i> -(2-((3 <i>R</i> ,5 <i>R</i>)-3,5-Dimethylmorpholino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	559.3

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
11-51		<i>N</i> -(2-(4-Fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	547.2
11-52		<i>N</i> -(2-(4-Fluoro-4-methylpiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	561.3
11-53		(<i>S</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-(3-hydroxypiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	545.2
11-54		(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-(3-hydroxypiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	545.2
11-55		(<i>R</i>)- <i>N</i> -(2-(3-Fluoro-3-methylpiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	561.2
11-56		<i>N</i> -(2-(4-Fluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	533.2
11-57		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	551.2
11-58		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)-4-((2-hydroxyethyl)sulfonamido)benzamide	583.2


Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
11-59		4-((2-Hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)-N-(2-(3,3,3-trifluoropropoxy)pyridin-4-yl)benzamide	543.2
11-60		4-((2-Hydroxyethyl)sulfonamido)-N-(2-methyl-6-(3,3,3-trifluoropropoxy)pyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	557.2
11-61		4-((2-Hydroxyethyl)sulfonamido)-N-(6-methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	558.2
11-62		<i>N</i> -(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	533.2
11-63		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-3-fluoropyridin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	568.2
11-64		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-3-fluoro-6-methylpyridin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	582.2
11-65		(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-(2-methylmorpholino)pyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	530.2
11-66		(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-methyl-6-(2-methylmorpholino)pyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	544.3

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
11-67		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	583.4
11-68		<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-(hydroxymethyl)cyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	563.2
11-69		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-(hydroxymethyl)cyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	591.2
11-70		(<i>S</i>)- <i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)-4-((tetrahydrofuran)-3-sulfonamido)benzamide	563.2
11-71		(<i>R</i>)- <i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)-4-((tetrahydrofuran)-3-sulfonamido)benzamide	563.2
11-72		(<i>S</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)-4-((tetrahydrofuran)-3-sulfonamido)benzamide	591.3
11-73		(<i>R</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)-4-((tetrahydrofuran)-3-sulfonamido)benzamide	591.3
11-74		<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	547.2

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
11-75		<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	535.2
11-76		(<i>R</i>)- <i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	551.2
11-77		(<i>S</i>)- <i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	551.2
11-78		4-((Cyclopropylmethyl)sulfonamido)- <i>N</i> -(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	547.2
11-79		4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-isopropoxy-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	504.2
11-80		(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-((tetrahydrofuran-3-yl)oxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	532.2
11-81		<i>N</i> -(2-(4-Fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(methylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	517.2
11-82		4-(Ethylsulfonamido)- <i>N</i> -(2-(4-fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	531.2

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
11-83		<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1,1-dimethylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	549.2


[0336] Example 12: *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0337] Step 1: A mixture of *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (500 mg, 0.881 mmol, Int. 19), triphenylphosphine (34.7 mg, 0.132 mmol, Aldrich St. Louis, MO USA), 1,10-phenanthroline (23.82 mg, 0.132 mmol, Aldrich St. Louis, MO USA), palladium(II) acetate (9.89 mg, 0.044 mmol, Strem Chemicals, Inc. Newburyport, MA USA), sodium formate (132 mg, 1.939 mmol, Thermo Fisher Scientific, Grand Island, NY USA), and tetrabutylammonium bromide (426 mg, 1.322 mmol, Aldrich St. Louis, MO USA) in dimethyl sulfoxide (3 mL) under N₂ was stirred at 70 °C for 45 min. Then, the mixture was cooled to room temperature and ethyl iodoacetate (0.157 mL, 1.322 mmol, Aldrich St. Louis, MO USA) was added. The mixture was then stirred at room temperature for 10 min. Then, the mixture was diluted with water (20 mL) and was then extracted with EtOAc (2 x 40 mL). The combined organic extracts were then dried over MgSO₄ and concentrated. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc/heptane) provided ethyl 2-((4-((2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)carbamoyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfonyl)acetate (330 mg, 0.558 mmol) as a light yellow solid. ¹H NMR (DMSO-*d*₆) δ 13.14 (br s, 1H), 8.27 (br d, *J* = 8.1 Hz, 1H), 7.97 (s, 1H), 7.84 (br d, *J* = 7.7 Hz, 1H), 7.40 (s, 1H), 4.80 (s, 2H), 4.05 (q, *J* = 7.2 Hz, 2H), 3.92 (br s, 4H), 3.03-3.13 (m, 4H), 2.34 (s, 3H), 1.90-2.07 (m, 4H), 1.71 (br s, 4H), 1.07 (t, *J* = 7.0 Hz, 3H), 0.39 (s, 4H). m/z (ESI): 592.3 (M+H)⁺.

[0338] Step 2: To a solution of ethyl 2-((4-((2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)carbamoyl)-3-(6-azaspiro[2.5]octan-6-yl)phenyl)sulfonyl)acetate (320 mg, 0.541 mmol) in 2-methyltetrahydrofuran (3.5 mL) under N₂ at 0 °C was added lithium borohydride solution (2.0M in tetrahydrofuran 0.541 mL, 1.082 mmol, Aldrich St. Louis, MO USA) dropwise. After addition, the mixture was stirred at room temperature overnight. Then, the mixture was quenched with saturated NH₄Cl (18 mL) and was stirred at room temperature for 15 min. The mixture was then extracted with EtOAc (2 x 30 mL). The combined organic extracts were then dried over MgSO₄ and concentrated. Chromatographic purification of the residue (silica gel, 0% - 100% EtOAc/heptane) provided *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (60 mg, 0.109 mmol, 20% yield) provided *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (60 mg, 0.109 mmol) as a white solid. ¹H NMR (DMSO-*d*₆) δ ppm 13.17 (br s, 1H), 8.25 (br d, *J* = 8.3 Hz, 1H), 7.93 (br s, 1H), 7.82 (br d, *J* = 8.3 Hz, 1H), 7.40 (br s, 1H), 4.91 (br t, *J* = 5.0 Hz, 1H), 3.92 (br s, 4H), 3.68-3.77 (m, 2H), 3.52-3.60 (m, 2H), 3.09 (br s, 4H), 2.34 (s, 3H), 1.87-2.08 (m, 4H), 1.70 (br d, *J* = 1.0 Hz, 4H), 0.39 (s, 4H). ¹⁹F NMR (DMSO-*d*₆) δ ppm -94.76 (s, 2F). *m/z* (ESI): 550.1 (M+H)⁺.

[0339] Examples 13-1 and 13-2: (*S*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide and (*R*)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0340] Step 1: *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (1.08 g, 1.90 mmol, Int. 19), 2-mercapto-1-propan-1-ol (0.48 g, 5.21 mmol, Enamine), and potassium carbonate (0.237 mL, 3.91 mmol) in 4 mL DMSO were heated at 90 °C


in a sealed vial for 4h. The mixture was cooled to RT, added 50 mL ethyl acetate and 10 mL brine. The organic layer was separated, washed with brine, dried, and evaporated. The resulting product was adsorbed onto a plug of silica gel and purified by silica gel chromatography (0-30% of EtOAc in heptane) to give *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)thio)-2-(6-azaspiro[2.5]octan-6-yl)benzamide as a yellow solid. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 0.39 - 0.44 (m, 4 H) 1.37 - 1.43 (m, 3 H) 1.55 - 1.60 (m, 4 H) 1.96 - 2.04 (m, 4 H) 2.35 - 2.41 (m, 3 H) 3.01 - 3.11 (m, 4 H) 3.46 - 3.78 (m, 3 H) 3.96 - 4.05 (m, 4 H) 7.28 - 7.36 (m, 2 H) 7.48 - 7.53 (m, 1 H) 8.12 - 8.32 (m, 1 H) 13.01 - 13.37 (m, 1 H). *m/z* (ESI): 531.4 (M+H)⁺.

[0341] Step 2: To *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)thio)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.62 g, 1.17 mmol) in 15 mL THF was added oxone(r) monopersulfate compound (0.72 g, 1.17 mmol) in 5 mL water. After stirring for 1.5 h, LCMS showed a mixture of sulfone and sulfoxide formed. Additional 0.4 g of oxone in 3 mL water was added. After stirring for additional 2 h, EtOAc (50 mL) and brine (20 mL) were added to the reaction mixture and the organic layer was taken, washed with brine, dried, and evaporated. The crude product was purified via preparative SFC using an (S,S) Whelk-01 (250 X 21 mm, 5mm) column with a mobile phase of 60% Liquid CO₂ and 40% MeOH, flowrate of 80 mL/min. to give *N*-(5-chloro-2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (337 mg, 0.58 mmol, 50% yield). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 0.30 - 0.57 (m, 4 H) 1.27 - 1.37 (m, 3 H) 1.59 - 1.72 (m, 4 H) 1.94 - 2.10 (m, 4 H) 2.40 - 2.54 (m, 3 H) 2.55 - 2.97 (m, 1 H) 3.05 - 3.23 (m, 4 H) 3.28 - 3.41 (m, 1 H) 3.84 - 4.06 (m, 6 H) 7.63 - 7.86 (m, 2 H) 8.13 - 8.37 (m, 1 H) 11.08 - 11.59 (m, 1 H). *m/z* (ESI): 598.3 (M+H)⁺. This racemic mixture was separated by preparative SFC using an OD (250 x 21 mm, 5 mm) with a mobile phase of 85% Liquid CO₂ and 15% iPrOH and a flowrate of 90 mL/min to generate:

[0342] Example 13-1: (S)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. First eluting peak (85 mg, ee > 99%). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 12.74 - 13.01 (m, 1 H), 8.36 - 8.54 (m, 1 H), 7.69 - 8.03 (m, 2 H), 7.41 - 7.58 (m, 1 H), 3.83 - 4.06 (m, 6 H), 3.25 - 3.43 (m, 1 H), 3.03 - 3.17 (m, 4 H), 2.50 - 2.81 (m, 1 H), 2.31 - 2.42 (m, 3 H), 1.93 - 2.10 (m, 4 H), 1.61 - 1.90 (m, 4 H), 1.28 - 1.36 (m, 3 H), 0.35 - 0.50 (m, 4 H). *m/z* (ESI): 563.2 (M+H)⁺.

[0343] Example 13-2: (R)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. Second eluting peak (84 mg, ee 97%). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 12.71 - 13.05 (m, 1 H), 8.39 - 8.56 (m, 1 H), 7.74 - 8.03 (m, 2 H), 7.38 - 7.54 (m, 1 H), 3.81 - 4.04 (m, 6 H), 3.26 - 3.39 (m, 1 H), 3.03 - 3.19 (m, 4 H), 2.48 - 2.83 (m, 1 H), 2.33 - 2.41 (m, 3 H), 1.92 - 2.10 (m, 4 H), 1.60 - 1.90 (m, 4 H), 1.28 - 1.33 (m, 3 H), 0.36 - 0.46 (m, 4 H). *m/z* (ESI): 563.2 (M+H)⁺. The stereochemistry was arbitrary assigned.

[0344] Example 14: *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxy-2-methylpropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0345] Step 1: *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.55 g, 0.967 mmol, Int. 19), 4,5-bis(diphenylphosphino)-9,9-dimethyl-xanthene (0.034 g, 0.058 mmol), tris(dibenzylideneacetone)dipalladium (0) chloroform adduct (0.035 g, 0.034 mmol), DIPEA (0.4 mL, 2.29 mmol), and 2-mercapto-2-methylpropan-1-ol (0.134 g, 1.29 mmol) in 3 mL dioxane were passed through N₂ for 5 min in a sealed tube. The mixture was heated at 90 °C for 3h and cooled to RT. The resulting crude product was adsorbed onto a plug of silica gel and purified by silica gel chromatography (0-7% EtOAc in DCM) to give *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxy-2-methylpropan-2-yl)thio)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 0.38 - 0.47 (m, 4 H) 1.25 - 1.31 (m, 6 H) 1.56 - 1.56 (m, 4 H) 1.94 - 2.04 (m, 4 H) 2.35 - 2.41 (m, 3 H) 3.03 - 3.12 (m, 4 H) 3.31 - 3.38 (m, 2 H) 3.95 - 4.04 (m, 4 H) 7.41 - 7.47 (m, 2 H) 7.48 - 7.52 (m, 1 H) 8.14 - 8.32 (m, 1 H) 12.96 - 13.41 (m, 1 H). 4H overlapped with water peak. *m/z* (ESI): 546.2 (M+H)⁺.


[0346] Step 2: To *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxy-2-methylpropan-2-yl)thio)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.36 g, 0.66 mmol) in THF (15 mL) cooled to 0 °C was added oxone(r) monopersulfate compound (0.52 g, 0.85 mmol) in water (5 mL). The mixture was stirred for 2.5 h from 0 °C to RT. Additional 0.35 g oxone was added. After 1h lcms showed the sulfoxide was almost consumed. Ethyl acetate (40 mL) and brine (20 mL) were added and organic layer was separated, dried and evaporated. The crude mixture was purified via preparative SFC using an (*S,S*) Whelk-01 (250 x 21 mm, 5mm) with a mobile phase of 60% Liquid CO₂ and 40% MeOH using a flowrate of 80 mL/min to give *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxy-2-methylpropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 8.15 - 8.73 (m, 1 H), 7.64 - 7.90 (m, 2 H), 7.41 - 7.52 (m, 1 H), 3.94 - 4.07 (m, 4 H), 3.70 - 3.83 (m, 2 H), 3.04 - 3.20 (m, 4 H), 2.34 - 2.47 (m, 3 H), 1.96 - 2.12 (m, 4 H), 1.47 - 1.95 (m, 4 H), 1.30 - 1.41 (m, 6 H), 0.37 - 0.52 (m, 4 H). ¹⁹F NMR (376 MHz, CHLOROFORM-*d*) δ ppm -96.68 (br s, 1 F). *m/z* (ESI): 578.2 (M+H)⁺.

Table 11: Examples 14-1 to 14-8 were prepared following similar procedures for Examples 12 to 14:

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
14-1		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(oxetan-3-ylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	562.3
14-2		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(ethylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	534.2
14-3		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(isopropylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	548.2
14-4		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-2-methylpropyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	578.2
14-5		4-(Cyclopropylsulfonyl)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	546.2
14-6		4-(<i>tert</i> -Butylsulfonyl)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	562.3
14-7		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(((3 <i>R</i> ,4 <i>R</i>)-4-hydroxytetrahydrofuran-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	592.2
14-8		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(((3 <i>S</i> ,4 <i>S</i>)-4-hydroxytetrahydrofuran-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	592.2


Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
		azaspiro[2.5]octan-6-yl)benzamide	
14-9		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(((1R,2R)-2-hydroxycyclopentyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	590.2

[0347] Example 15: *N*-(2-(4,4-Difluorocyclohexyl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0348] A mixture of 4-bromo-*N*-(2-(4,4-difluorocyclohexyl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.055 g, 0.106 mmol, Int. 21-13), 2-hydroxyethane-1-sulfonamide (0.020 g, 0.159 mmol, Wuxi), potassium phosphate tribasic (0.045 g, 0.212 mmol), copper(I) iodide (0.020 g, 0.106 mmol) and (1*R*,2*R*)-*N,N'*-dimethyl-1,2-cyclohexanediamine (7.53 mg, 0.053 mmol, Combi-Blocks) in DMF (1.5mL) was heated at 90 °C for 16 h. Then the reaction mixture was filtered through a pad of CELITE® pad and filtrate was diluted with EtOAc. The resulting solution was washed with water and brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by reverse phase HPLC using a gradient of 60 % ACN in water (0.1% TFA) to afford *N*-(2-(4,4-difluorocyclohexyl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.025 g, 0.044 mmol, 42 % yield) as white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.70 (s, 1 H), 8.04 (d, *J* = 8.8 Hz, 1 H), 7.92 (s, 1 H), 7.25 (d, *J* = 2.2 Hz, 1 H), 7.16 – 7.08 (m, 1 H), 3.75 (t, *J* = 6.3 Hz, 2 H), 3.03 – 2.85 (m, 6 H), 2.44 (d, *J* = 4.5 Hz, 5 H), 2.05 (s, 5 H), 1.92 (d, *J* = 11.7 Hz, 4 H), 1.72 (s, 4 H), 0.38 (s, 4 H). m/z (ESI): 564.1 (M+H)⁺.

[0349] Example 16-1 and 16-2: (*S*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide and (*R*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0350] Step 1: To a solution of 1-(benzyloxy)-3-((tert-butyldimethylsilyl)oxy)propane-2-sulfonamide (0.803 g, 2.23 mmol) in THF was added tetrabutylammonium fluoride solution (2.75 mL, 2.75 mmol, 1M in THF) at RT. The reaction was stirred for 1 hour and then concentrated under reduced pressure. The resulting material was used immediately in the next step.

[0351] Step 2: To a pressure relief vial charged with the sulfonamide from the previous step was added copper(I) iodide (0.196 g, 1.03 mmol), methyl glycine (0.128 g, 1.440 mmol), potassium phosphate, tribasic (0.934 g, 4.40 mmol), and *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.975 g, 1.72 mmol, Int. 19). The vial was sealed and evacuated / backfilled with nitrogen and then DMF (7 mL) was added. The cap was replaced, and the reaction was stirred in a pre-heated 100 °C oil bath for 16 h. The reaction mixture was partitioned between satd. NH₄Cl:NH₄OH (9:1) and EtOAc. The organic phase was separated, washed with brine, and concentrated in vacuo. The material was purified by silica gel chromatography (20-100% EtOAc in heptane) to afford

4-((2-(benzyloxy)-1-(hydroxymethyl)ethyl)sulfonamido)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.635 g, 0.927 mmol, 54.0 % yield). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 13.11 (br dd, *J*=4.66, 2.38 Hz, 1 H) 8.14 (d, *J*=8.50 Hz, 1 H) 7.49 (br s, 1 H) 7.31 - 7.44 (m, 5 H) 7.13 (d, *J*=1.87 Hz, 1 H) 6.83 (dd, *J*=8.60, 2.18 Hz, 2 H) 4.49 - 4.62 (m, 2 H) 4.08 (dt, *J*=11.77, 5.83 Hz, 1 H) 3.94 - 4.03 (m, 5 H) 3.88 - 3.93 (m, 1 H) 3.95 (br s, 1 H) 3.45 - 3.55 (m, 1 H) 2.96 (br t, *J*=4.98 Hz, 4 H) 2.29 - 2.49 (m, 4 H) 1.95 - 2.07 (m, 4 H) 1.57 (br s, 4 H) 1.18 - 1.35 (m, 2 H) 0.85 - 0.91 (m, 1 H) 0.40 (s, 4 H). *m/z* (ESI, +ve ion): 683.8 (M+H)⁺.

[0352] Step 3: To a solution of xtalfleur-m (0.351 g, 1.446 mmol) in DCM (6 mL) in a brine/ice bath was added triethylamine trihydrofluoride (0.262 mL, 1.61 mmol) followed by a solution of 4-((2-(benzyloxy)-1-(hydroxymethyl)ethyl)sulfonamido)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.55 g, 0.803 mmol) in DCM (10 mL) dropwise via addition funnel. The reaction was gradually warmed to RT and stirred for 16h. The reaction was quenched with saturated sodium bicarbonate (aq) and diluted with water and DCM. The organic phase was separated, washed with brine, dried over magnesium sulfate, and concentrated in vacuo. The crude material was purified by silica gel chromatography (20-100% EtOAc in heptane) to afford a 1:0.8 mixture of rac- 4-((2-(benzyloxy)-1-(fluoromethyl)ethyl)sulfonamido)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide and 4-((1-((benzyloxy)methyl)vinyl)sulfonamido)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (265 mg). The mixture was carried forward without further purification.


[0353] Step 4: To a suspension of 265 mg of the product mixture from the previous step and palladium hydroxide on carbon (0.135 g, 0.193 mmol), in EtOH (20 mL) was added and AcOH (0.033 mL, 0.579 mmol). The reaction was hydrogenated under 55 psi hydrogen at room temperature for 24 hours. After flushing the reaction with Nitrogen, an additional portion of palladium hydroxide on carbon (0.135 mL, 0.193 mmol) was added followed by additional AcOH (0.033 mL, 0.579 mmol). The reaction vessel was flushed with N₂ before replacing the atmosphere with 55 psi hydrogen. The reaction was continued at room temperature for an additional 48 hours. The reaction was flushed with nitrogen then filtered over CELITE® and the filtrate was concentrated in vacuo. The racemic mixture was purified by preparative SFC using an IE (250 x 21 mm, 5 mm) with a mobile phase of 80% Liquid CO₂ and 20% MeOH (flowrate of 80 mL/min) to give:

[0354] Example 16-1: (*S*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. First eluting peak ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 13.35 (s, 1 H), 10.21 - 10.84 (m, 1 H), 8.06 (d, *J* = 8.50 Hz, 1 H) 7.40 (s, 1 H), 7.29 (d, *J*=2.07 Hz, 1 H), 7.16 (dd, *J*=8.50, 2.07 Hz, 1 H), 5.12 - 5.38 (m, 1 H), 4.88 - 4.97 (m, 1 H) 4.71 - 4.84 (m, 1 H), 3.85 - 4.01 (m, 5 H), 3.74 (dd, *J* = 11.30, 7.98 Hz, 1 H), 3.51 - 3.63 (m, 1 H) 2.98 (br t, *J* = 4.77 Hz, 4 H), 2.32 (s, 3 H), 1.93 - 2.07 (m, 4 H), 1.51 - 1.91 (m, 4 H), 0.40 (s, 4 H). *m/z* (ESI, +ve ion): 597.2 (M+H)⁺.

[0355] Example 16-2: (*R*)-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide. Second eluting peak. ¹H

NMR (400 MHz, DMSO- d_6) δ ppm 13.35 (s, 1 H) 10.32 - 10.70 (m, 1 H) 8.05 (d, J = 8.71 Hz, 1 H) 7.40 (s, 1 H) 7.29 (d, J = 2.07 Hz, 1 H) 7.16 (dd, J = 8.71, 2.07 Hz, 1 H) 5.14 - 5.40 (m, 1 H) 4.85 - 4.97 (m, 1 H) 4.73 - 4.84 (m, 1 H) 3.87 - 3.98 (m, 5 H), 3.74 (dd, J = 11.09, 7.98 Hz, 1 H) 3.48. m/z (ESI, +ve ion): 597.2 ($M+H$)⁺. The stereochemistry was arbitrarily assigned

[0356] Example 17: *N*-(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(*N*-(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide




[0357] A mixture of *N*-(2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (212 mg, 0.384 mmol, Int. 21-14), 1,4-diazabicyclo[2.2.2]octane bis(sulfur dioxide) adduct (DABSO) (55 mg, 0.23 mmol, Sigma-Aldrich), diacetoxypalladium (13 mg, 0.06 mmol, Strem), rac-((3*R*,5*R*,7*R*)-adamantan-1-yl)((3*S*,5*S*,7*S*)-adamantan-1-yl)(butyl)phosphane (cataCXium® A) (28 mg, 0.08 mmol, Strem), and triethylamine (107 μ L, 0.77 mmol) in IPA (3 mL) in a glass tube was degassed for 3 min. The tube was sealed then heated at 85 °C in an oil bath for 3 h. The heterogeneous mixture was cooled to RT and treated with 2-aminoethan-1-ol (47 mg, 0.77 mmol, Sigma-Aldrich) followed by sodium hypochlorite solution (10% wt., 571 mg, 0.77 mmol, Sigma-Aldrich) and stirred at RT for 18 h. The mixture was treated with 2-aminoethan-1-ol (23 mg) followed by sodium hypochlorite solution (10% wt., 275 mg) then stirred at RT for 5 h. EtOAc (20 mL) and water (5 mL) were added to the heterogeneous mixture and the insoluble solid was filtered off. The filter cake was washed with water (2 x 2 mL) followed by EtOAc (2 x 4 mL). The organic solution was taken and concentrated under vacuum. The residue was purified by silica gel chromatography (10% to 60% EtOAc in heptane) to give *N*-(2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)-4-(*N*-(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (100 mg, 0.18 mmol, 47% yield) as an off-white solid. ¹H NMR (400 MHz, METHANOL- d_4) δ ppm 7.99-8.14 (m, 2H), 7.79 (s, 1H), 7.68 (d, J = 7.88 Hz, 1H), 7.47 (s, 1H), 7.01 (d, J = 4.77 Hz, 1H), 3.76 (t, J = 5.29 Hz, 4H), 3.58 (t, J = 5.91 Hz, 2H), 3.16 (t, J = 5.08 Hz, 4H), 3.02 (t, J = 5.80 Hz, 2H), 1.98-2.11 (m, 4H), 1.62 (s, 4H), 0.42 (s, 4H). m/z (ESI): ($M+H$)⁺ 550.1.

[0358] Table 12: Examples 17-1 to 17-8 were prepared following similar procedures for Example 17:

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
17-1		4-(<i>N</i> -(2-Hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)- <i>N</i> -(2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)benzamide	544.2

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
17-2		4-(<i>N</i> -(2-Hydroxyethyl)sulfamoyl)- <i>N</i> -(6-methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	558.2
17-3		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	565.2
17-4		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyrimidin-4-yl)-4-(<i>N</i> -(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	551.2
17-5		<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	537.2
17-6		(<i>R</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(1-hydroxypropan-2-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	579.4
17-7		(<i>S</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(1-hydroxypropan-2-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	579.4
17-8		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(2-hydroxy-2-methylpropyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	593.4

[0359] Examples 18-1 and 18-2: 2-(6-Azaspiro[2.5]octan-6-yl)-4-(*R*-cyclopropylsulfonimidoyl)-*N*-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide and 2-(6-azaspiro[2.5]octan-6-yl)-4-(*S*-cyclopropylsulfonimidoyl)-*N*-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide



[0360] Step 1: Into a 20 mL microwave vessel were placed *N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-iodo-2-(6-azaspiro[2.5]octan-6-yl)benzamide (1.00 g, 1.762 mmol, Int. 19), tris (dibenzylideneacetone) dipalladium (0) (0.161 g, 0.176 mmol) and 4,5-bis(diphenylphosphino)-9,9-dimethyl-xanthene (0.102 g, 0.176 mmol) followed by 1,4-dioxane (10 mL). The resulting mixture was stirred and purged with nitrogen for 5 min before 1,1'-dimethyltriethylamine (0.616 mL, 3.52 mmol) was added under nitrogen followed by cyclopropanethiol (0.142 mL, 1.939 mmol). The vessel was sealed and subjected to microwave condition (10 h, 90 °C). The crude mixture was directly loaded onto a silica gel precolumn and subjected to combi-flash column chromatography on a 40-g ISCO gold column eluting with MeOH/DCM (5 min at 0% and 25 min from 0 to 6%) twice to give 4-(cyclopropylthio)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.92 g, 1.791 mmol, 102 % yield) as an off-white solid. ¹H NMR (400 MHz, DICHLOROMETHANE-*d*₂) δ ppm 13.33 (s, 1H), 8.15 (d, *J*=8.29 Hz, 1H), 7.48 (s, 1H), 7.22-7.35 (m, 2H), 3.91-4.09 (m, 4H), 3.06 (br t, *J*=5.18 Hz, 4H), 2.35 (s, 3H), 2.17-2.28 (m, 1H), 1.62-2.10 (m, 6H), 1.52 (s, 2H), 1.13-1.21 (m, 2H), 0.68-0.76 (m, 2H), 0.40 (s, 4H). *m/z* (ESI): 514.1 (M+H)⁺.

[0361] Step 2: To a stirred solution of 4-(cyclopropylthio)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.89 g, 1.733 mmol) and ammonium carbonate (0.250 g, 2.60 mmol) in MeOH (4.5 mL) and dichloromethane (9.0 mL) was added (acetyloxy)(phenyl)-iodanyl acetate (1.284 g, 3.99 mmol) in one portion as a solid. The resulting mixture was stirred in open air at rt for 18 h. The resulting mixture was directly loaded onto silica gel precolumn (25 g) and subjected to combi-flash column chromatography on a 40-g ISCO gold column eluting with MeOH/DCM (3 min at 0% and 25 min from 0 to 14%) to give a racemic mixture of 4-(cyclopropanesulfonimidoyl)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.95 g, 1.744 mmol, 101 % yield) as an off-white solid. The enantiomers were separated via preparative SFC using a Regis (S,S) Whelk-01 (250 X 21 mm, 5mm) with a mobile phase of 50% Liquid CO₂ and 50% MeOH using a flow rate of 60 mL/min to generate:

[0362] Example 18-1: 2-(6-Azaspiro[2.5]octan-6-yl)-4-(*R*-cyclopropylsulfonimidoyl)-*N*-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide. First eluting peak, ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 13.20 (br d, *J*=3.73 Hz, 1H), 8.44 (d, *J*=8.29 Hz, 1H), 7.96 (d, *J*=1.45 Hz, 1H), 7.87 (dd, *J*=1.66, 8.29 Hz, 1H), 7.52 (s, 1H), 4.03 (br s, 4H), 3.14 (t, *J*=5.29 Hz, 4H), 2.53-2.63 (m, 1H), 2.44 (br s, 3H), 1.95-2.10 (m, 4H), 1.53-1.89 (m, 5H), 1.45 (tdd, *J*=5.08, 6.92, 10.29 Hz, 1H), 1.20-1.30 (m, 1H), 1.07-1.17 (m, 1H), 0.93-1.03 (m, 1H), 0.44 (s, 4H). *m/z* (ESI): 545.2 (M+H)⁺.


[0363] Example 18-2: 2-(6-Azaspiro[2.5]octan-6-yl)-4-(*R*-cyclopropylsulfonimidoyl)-*N*-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide. Second eluting peak. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 13.20 (br d, *J*=3.73 Hz, 1H), 8.44 (d, *J*=8.29 Hz, 1H), 7.96 (d, *J*=1.45 Hz, 1H), 7.87 (dd, *J*=1.66, 8.29 Hz, 1H), 7.52 (s, 1H), 4.03 (br s, 4H), 3.14 (t, *J*=5.29 Hz, 4H), 2.53-2.63 (m, 1H), 2.44 (br s, 3H), 1.95-2.10 (m, 4H), 1.53-1.89 (m, 5H), 1.45 (tdd, *J*=5.08, 6.92, 10.29 Hz, 1H), 1.20-1.30 (m, 1H), 1.07-1.17 (m, 1H), 0.93-1.03 (m, 1H), 0.44 (s, 4H). *m/z* (ESI): 545.2 (**M+H**)⁺. The stereochemistry assignments were arbitrary.

[0364] Table 13: Examples 19-1 to 19-9 were prepared following the procedure described for Examples 18-1 and 18-2:

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) <i>m/z</i>
19-1		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(oxetane-3-sulfonimidoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	561.2
19-2		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(methylsulfonimidoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	519.2
19-3		2-(6-Azaspiro[2.5]octan-6-yl)- <i>N</i> -(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)-4-(<i>S</i> -ethylsulfonimidoyl)benzamide	533.2
19-4		2-(6-Azaspiro[2.5]octan-6-yl)- <i>N</i> -(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)-4-(<i>R</i> -ethylsulfonimidoyl)benzamide	533.2
19-5		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(propan-2-ylsulfonimidoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	547.3

Ex. #	Chemical Structure	Name	LRMS: (ESI + ve ion) m/z
19-6		4-(Cyclopropanesulfonylimidoyl)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	530.0
19-7		2-(6-Azaspiro[2.5]octan-6-yl)-4-(<i>S</i> -cyclopropylsulfonylimidoyl)- <i>N</i> -(6-methyl-2-((2 <i>R</i>)-2-methyl-4-morpholinyl)-4-pyrimidinyl)benzamide	525.2
19-8		2-(6-Azaspiro[2.5]octan-6-yl)-4-(<i>R</i> -cyclopropylsulfonylimidoyl)- <i>N</i> -(6-methyl-2-((2 <i>R</i>)-2-methyl-4-morpholinyl)-4-pyrimidinyl)benzamide	525.2
19-9		4-(Cyclopropanesulfonylimidoyl)- <i>N</i> -(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	517.3

[0365] Example 20: (*N*¹-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)terephthalamide




[0366] Step 1: To a solution of 4-bromo-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (250 mg, 0.48 mmol, Int. 21-3) in DMF (2.5 mL) was added $\text{Pd}(\text{PPh}_3)_4$

(6 mg, 4.8 μ mol) and $\text{Zn}(\text{CN})_2$ (113 mg, 0.961 mmol) and the reaction mixture was stirred at 100 °C for 16 h. Then the reaction mixture was diluted with EtOAc and filtered through a CELITE® bed. The filtrate was washed with water and brine, dried over Na_2SO_4 , filtered, concentrated and purified by flash column chromatography using a gradient of 30% EtOAc in petroleum ether to provide 4-cyano-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (160 mg, 0.343 mmol, 71.4 % yield) as an off white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 13.20 (s, 1 H), 8.18 (d, $J=8.1$ Hz, 1 H), 8.03 (s, 1 H), 7.79 (d, $J=8.1$ Hz, 1 H), 7.39 (s, 1 H), 4.00 – 3.80 (m, 4 H), 3.07 (t, $J=5.2$ Hz, 4 H), 2.34 (s, 3 H), 1.99 (tt, $J=13.3, 5.7$ Hz, 4 H), 1.69 (s, 4 H), 0.38 (s, 4 H). m/z (ESI): 467.2 ($\text{M}+\text{H}$) $^+$.

Step 2: To a solution of 4-cyano-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (110 mg, 0.236 mmol) in dimethyl sulfoxide (2 mL) were added K_2CO_3 (6.52 mg, 0.047 mmol) and H_2O_2 (103 μL , 1.179 mmol) at 0 °C and the reaction mixture was stirred for 1 h before it was quenched with water and was extracted with EtOAc (2 x 20 mL). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered, concentrated and purified by flash column chromatography using a gradient of 10% methanol in dichloromethane to provide *N* 1 -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)terephthalamide (108 mg, 0.223 mmol, 95 % yield) as off white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 13.75 (s, 1 H), 8.26 (m, 1 H), 8.17 (m, 1 H), 8.00 (s, 1 H), 7.89 – 7.78 (m, 1 H), 7.61 (s, 1 H), 7.42 (s, 1 H), 3.94 (d, $J=5.8$ Hz, 4 H), 3.06 (d, $J=6.0$ Hz, 4 H), 2.34 (d, $J=2.3$ Hz, 3 H), 2.01 (q, $J=8.5, 8.1$ Hz, 4 H), 1.74 (s, 4 H), 0.41 (d, $J=2.3$ Hz, 4 H). m/z (ESI): 485.2 ($\text{M}+\text{H}$) $^+$.


[0367] Example 21: 4-(Azetidin-3-ylsulfonyl)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0368] To a solution of 4-((1-(*tert*-butoxycarbonyl)azetidin-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzoic acid (2.54 g, 5.64 mmol, Int. 16) and HATU (3.22 g, 8.46 mmol, ChemPep) in DMF (35 mL), was added 2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-amine (1.93 g, 8.46 mmol, Int. 4) and DIPEA (2.46 mL, 14.09 mmol). The mixture was stirred at room temperature for 18 h. The mixture was diluted with saturated Na_2CO_3 and EtOAc. The organic was separated and washed with Na_2CO_3 , water and brine, dried over Na_2SO_4 and concentrated in vacuo. The crude was purified by silica gel chromatography: 0-30%-60% EtOAc in heptane. m/z (ESI): 661.3 ($\text{M}+\text{H}$) $^+$. The residue was treated with DCM (8 mL) and TFA (4 mL) at room temperature for 30 min and concentrated in vacuo. The solid obtained was suspended in EtOAc and washed with 1N NaOH solution and the mixture was extracted

with EtOAc. The organic extracts were washed with water and brine, dried over Na₂SO₄ and concentrated in vacuo. The crude was purified by silica gel chromatography: 0-20% MeOH in DCM with 2% NH₄OH to give the title compound as a white solid. ¹H NMR (400 MHz, METHANOL-d₄) δ ppm 8.35 (d, *J*=8.09 Hz, 1H), 7.90 (d, *J*=1.24 Hz, 1H), 7.81 (dd, *J*=1.66, 8.09 Hz, 1H), 7.46 (s, 1H), 4.46-4.60 (m, 1H), 3.97-4.06 (m, 6H), 3.76-3.86 (m, 2H), 3.10-3.21 (m, 4H), 2.37 (s, 3H), 1.93-2.01 (m, 4H), 1.73-1.88 (m, 4H), 0.45 (s, 4H). *m/z* (ESI): 561.2 (M+H)⁺.

[0369] Example 22: 4-*N*-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylazetidin-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide



[0370] To a mixture of 4-(azetidin-3-ylsulfonyl)-*N*-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide (0.045 g, 0.080 mmol, ex. 21), MeOH (1 mL), and formaldehyde with 10-15% MeOH (0.016 g, 0.482 mmol, Fisher) was added AcOH (0.037 mL, 0.642 mmol, Aldrich), followed by sodium triacetoxyborohydride (0.204 g, 0.963 mmol, Aldrich). The mixture was stirred at room temperature for 18 h and concentrated in vacuo. The acid was neutralized with 1N NaOH solution and the mixture was extracted with EtOAc. The organic phase was washed with water and brine, dried over Na₂SO₄ and concentrated in vacuo. Purification by silica gel chromatography: 0-100% EtOAc/EtOH (3/1) in heptane gave the title compound as a white solid. ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 12.82 (br s, 1H), 8.44 (d, *J*=8.29 Hz, 1H), 7.82 (d, *J*=1.66 Hz, 1H), 7.73 (dd, *J*=1.76, 8.19 Hz, 1H), 7.46 (s, 1H), 3.97-4.11 (m, 5H), 3.60-3.90 (m, 4H), 3.12 (t, *J*=5.29 Hz, 4H), 2.44-2.57 (m, 3H), 2.39 (s, 3H), 1.68-2.06 (m, 8H), 0.43 (s, 4H). *m/z* (ESI): 575.3 (M+H)⁺.

[0371] BIOLOGICAL EXAMPLES

[0372] The following assays were used in testing the exemplary compounds of the invention. Data for those examples tested in accordance with the procedures described below are presented in Table A below.

[0373] KIF18A Enzyme Assay: Microtubule-stimulated ATPase activity assay is used to measure KIF18A enzyme activity after treatment with compound. Compounds were 2-fold serially diluted in DMSO (Sigma Inc) over 22-point concentration range. Recombinant human KIF18A (1-467 His-tagged) protein was expressed using a baculovirus system and purified by affinity chromatography by Amgen Inc. Concentrations of KIF18A protein, microtubules (MT), and ATP in the reaction were optimized for standardized homogenous enzyme assay using ADP-Glo™ Kinase/ATPase Assay Kit (Promega Inc). The assay measures ADP formed from the ATPase reaction. Prepare reaction buffer [(15 mM Tris, pH 7.5 (Teknova Inc), 10 mM MgCl₂ (JT Baker Inc), 0.01% Pluronic F-68 (Life Technologies Inc), 1 μM Taxol (Cytoskeleton Inc), and 30 μg/mL pig microtubules (Cytoskeleton Inc)]. Add compound and KIF18A

protein (30 nM) to prepared reaction buffer and incubated for 15 minutes at room temperature, next add ATP (at K_m , 75 μ M) to the reaction mixture and incubated for an additional 15 minutes at room temperature. Mix 5 μ l of ADP-GloTM Reagent and 2.5 μ l of the reaction mixture and incubate for 40 minutes at room temperature. Add 10 μ l ADP-GloTM Detection Reagent and incubate for 40 minutes at room temperature. Read luminescence using EnVision microplate reader with ultra-luminescence module (Perkin Elmer Inc). Concentration-response curve-fitting and IC_{50} determination was performed using Genedata Screener Software (Standard 15.0.1, Genedata Inc) with a four-parameter logistic regression fit model.

[0374] Table A provides data for compounds exemplified in the present application and priority document thereof, as representative compounds of the present invention, as follows: compound name and biological data. (IC_{50} in μ M, where available. Ex. # refers to Example No.)

[0375] TABLE A: BIOLOGICAL DATA

Ex. #	Compound Name	KIF18A ATPase IC_{50} (μ M)
1	<i>N</i> -(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.186
1-1	(<i>R</i>)- <i>N</i> -(2-((1-Hydroxypropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.216
1-2	(<i>S</i>)- <i>N</i> -(2-((1-Hydroxypropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.491
1-3	<i>N</i> -(2-((2-Hydroxyethyl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.849
1-4	(<i>R</i>)- <i>N</i> -(2-((2-Hydroxypropyl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.580
1-5	(<i>S</i>)- <i>N</i> -(2-((2-Hydroxypropyl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.451
1-6	<i>N</i> -(6-Methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.032
1-7	<i>N</i> -(2-((1-hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.168
2	<i>N</i> -(2-(2-Hydroxypropan-2-yl)pyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.143
2-1	<i>N</i> -(2-(2-Hydroxypropan-2-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.065
2-2	(<i>R</i>)- <i>N</i> -(2-(2-methylmorpholino)pyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.068
2-3	(<i>R</i>)- <i>N</i> -(6-Methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.115
2-4	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.029
2-5	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.103
2-6	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.260
2-7	(<i>R</i>)- <i>N</i> -(2-(2-Methylmorpholino)pyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.240

Ex. #	Compound Name	KIF18A ATPase IC ₅₀ (μM)
2-8	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.713
3	(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.057
4	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.071
5-1	(<i>R</i>)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.062
5-2	(<i>S</i>)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.070
6-1	4-((Fluoromethyl)sulfonamido)- <i>N</i> -(2-(4-fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.062
6-2	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((fluoromethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.042
6-3	(<i>R</i>)-4-((Fluoromethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.079
6-4	(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.032
6-5	(<i>R</i>)- <i>N</i> -(6-Methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-4-(methylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.120
6-6	(<i>R</i>)-4-(Ethylsulfonamido)- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.067
6-7	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(ethylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.122
6-8	(<i>R</i>)-4-((3-Hydroxypropyl)sulfonamido)- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.107
6-9	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(3,3-dioxido-1,3,4-oxathiazinan-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.246
6-10	4-(Cyclopentanesulfonamido)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	2.42
6-11	4-(Cyclobutanesulfonamido)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.127
6-12	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(oxetane-3-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.040
6-13	4-(Cyclobutanesulfonamido)- <i>N</i> -(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.052
6-14	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1,1-dimethylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.413
6-15	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((3-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.086
6-16	4-(Cyclopropanesulfonamido)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.052
6-17	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-methoxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.032

Ex. #	Compound Name	KIF18A ATPase IC ₅₀ (μM)
6-18	4-((Cyclopropylmethyl)sulfonamido)-N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.085
6-19	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(((3-hydroxyoxetan-3-yl)methyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.039
6-20	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1,1-dimethylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.049
6-21	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(methylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.039
6-22	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.151
6-23-1	4-(((S)-2-Hydroxy-1-methylethyl)sulfonamido)-N-(6-methyl-2-((R)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.039
6-23-2	4-(((R)-2-Hydroxy-1-methylethyl)sulfonamido)-N-(6-methyl-2-((R)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.0373
6-24-1	4-(((S)-2-Hydroxy-1-methylethyl)sulfonamido)-N-(2-((R)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.0174
6-24-2	4-(((R)-2-hydroxy-1-methylethyl)sulfonamido)-N-(2-((R)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.028
6-25-1	(S)-N-(2-(4-fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.047
6-25-2	(R)-N-(2-(4-Fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.041
6-26-1	(S)-N-(2-(4,4-Difluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.031
6-26-2	(R)-N-(2-(4,4-Difluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.021
6-27-1	4-(((R)-2-Hydroxypropyl)sulfonamido)-N-(6-methyl-2-((R)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.162
6-27-2	4-(((S)-2-Hydroxypropyl)sulfonamido)-N-(6-methyl-2-((R)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.151
6-28-1	(R)-4-((2-Hydroxypropyl)sulfonamido)-N-(6-methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.068
6-28-2	(S)-4-((2-Hydroxypropyl)sulfonamido)-N-(6-methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.071
6-29	N-(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.017
6-30	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyridin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.0384
6-31-1	(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-(1-hydroxyethyl)cyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.324
6-31-2	(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-(1-hydroxyethyl)cyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.281

Ex. #	Compound Name	KIF18A ATPase IC ₅₀ (μM)
6-32-1	(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.112
6-32-2	(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.141
6-33-1	4-(((S)-2-Hydroxypropyl)sulfonamido)-N-(2-((R)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.115
6-33-2	4-(((R)-2-hydroxypropyl)sulfonamido)-N-(2-((R)-2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.086
6-34	5-Chloro-N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.0922
6-35	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-5-methyl-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.144
6-36	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-5-methoxy-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.114
6-37	N-(2-(4,4-Difluoropiperidin-1-yl)-6-(2-hydroxypropan-2-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.125
6-38	N-(2-(4,4-Difluoropiperidin-1-yl)-6-ethylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.060
6-39	N-(6-Cyclopropyl-2-(4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.225
6-40	(R)-4-((2-Hydroxyethyl)sulfonamido)-N-(6-(2-hydroxypropan-2-yl)-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.161
7	N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.023
8-1	(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.091
8-2	(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.073
9-1	(R)-5-Fluoro-4-((2-hydroxyethyl)sulfonamido)-N-(2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.078
9-2	(R)-5-Fluoro-4-((2-hydroxyethyl)sulfonamido)-N-(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.158
10-1	(R)-N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.083
10-2	(S)-N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.073
11-1	(S)-4-((2-Hydroxyethyl)sulfonamido)-N-(2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.037
11-2	4-((2-Hydroxyethyl)sulfonamido)-N-(2-isopropyl-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.027
11-3	N-(2-Cyclopropyl-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.049

Ex. #	Compound Name	KIF18A ATPase IC ₅₀ (μM)
11-4	<i>N</i> -(2-Cyclobutoxy-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.018
11-5	<i>N</i> -(2-(3-Fluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.032
11-6	<i>N</i> -(2-(3-Fluoroazetidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.068
11-7	<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.023
11-8	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.640
11-9	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(6-oxa-1-azaspiro[3.3]heptan-1-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.128
11-10	<i>N</i> -(2-(3-(Difluoromethoxy)azetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.192
11-12	<i>N</i> -(2-(1,1-Difluoro-5-azaspiro[2.3]hexan-5-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.093
11-13	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(3-(trifluoromethyl)azetidin-1-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.067
11-14	<i>N</i> -(2-(3-Cyanoazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.172
11-15	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-(3-(2-hydroxypropan-2-yl)azetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.935
11-16	<i>N</i> -(2-(3-Hydroxy-3-methylazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	1.01
11-17	<i>N</i> -(2-(3-Cyclopropyl-3-hydroxyazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	1.09
11-18	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(5-azaspiro[2.3]hexan-5-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.064
11-19	<i>N</i> -(2-(3-Hydroxy-3-(trifluoromethyl)azetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.205
11-20	<i>N</i> -(2-(Azetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.0483
11-21	(<i>S</i>)- <i>N</i> -(2-(3-(1-Hydroxyethyl)azetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.704
11-22	4-(Cyclopropanesulfonamido)- <i>N</i> -(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.788
11-23	<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-(oxetane-3-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.0489
11-24	<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-(methylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.0551
11-25	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(5-azaspiro[2.4]heptan-5-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.0176
11-26	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.0716
11-27	(<i>R</i>)- <i>N</i> -(2-(3-Fluoro-3-methylpyrrolidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.020
11-28	<i>N</i> -(2-(2-Azabicyclo[3.1.0]hexan-2-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.059

Ex. #	Compound Name	KIF18A ATPase IC ₅₀ (μM)
11-29	<i>N</i> -(2-((1 <i>R</i> ,5 <i>S</i>)-3-Azabicyclo[3.1.0]hexan-3-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.020
11-30	<i>N</i> -(2-(3,3-Difluoropyrrolidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.033
11-31	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(6-azaspiro[2.5]octan-6-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.146
11-32	<i>N</i> -(2-((1 <i>R</i>)-1-Hydroxy-3-azabicyclo[3.1.0]hexan-3-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.193
11-33	<i>N</i> -(2-(4-Cyanopiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.088
11-34	(<i>R</i>)- <i>N</i> -(2-(3-Cyanopiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.092
11-35	(<i>S</i>)- <i>N</i> -(2-(3-Cyanopiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.043
11-36	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(piperidin-1-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.082
11-37	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-(4-hydroxypiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.144
11-38	<i>N</i> -(2-(4-Hydroxy-4-methylpiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.065
11-39	<i>N</i> -(2-(3,3-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.045
11-40	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(7-azaspiro[3.5]nonan-7-yl)benzamide	0.109
11-41	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(4,4-dimethylpiperidin-1-yl)-4-((2-hydroxyethyl)sulfonamido)benzamide	0.127
11-42	<i>N</i> -(2-((1 <i>R</i>)-1-Hydroxy-3-azabicyclo[4.1.0]heptan-3-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.034
11-43	<i>N</i> -(2-((1 <i>R</i> ,6 <i>R</i>)-3-Azabicyclo[4.1.0]heptan-3-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.035
11-44	<i>N</i> -(2-((1 <i>S</i> ,6 <i>R</i>)-2-Azabicyclo[4.1.0]heptan-2-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.033
11-45	(<i>S</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.035
11-46	<i>N</i> -(2-(2,2-Dimethylmorpholino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.150
11-47	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(4-oxa-7-azaspiro[2.5]octan-7-yl)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.095
11-48	<i>N</i> -(2-((3 <i>S</i> ,5 <i>R</i>)-3,5-Dimethylmorpholino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.095
11-49	<i>N</i> -(2-((3 <i>S</i> ,5 <i>S</i>)-3,5-Dimethylmorpholino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.245
11-50	<i>N</i> -(2-((3 <i>R</i> ,5 <i>R</i>)-3,5-Dimethylmorpholino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.187
11-51	<i>N</i> -(2-(4-Fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.020
11-52	<i>N</i> -(2-(4-Fluoro-4-methylpiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.063

Ex. #	Compound Name	KIF18A ATPase IC ₅₀ (μM)
11-53	(S)-4-((2-Hydroxyethyl)sulfonamido)-N-(2-(3-hydroxypiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.105
11-54	(R)-4-((2-Hydroxyethyl)sulfonamido)-N-(2-(3-hydroxypiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.074
11-55	(R)-N-(2-(3-Fluoro-3-methylpiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.041
11-56	N-(2-(4-Fluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.022
11-57	N-(2-(4,4-Difluoropiperidin-1-yl)pyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.022
11-58	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-fluoro-4-((2-hydroxyethyl)sulfonamido)-6-(6-azaspiro[2.5]octan-6-yl)benzamide	0.071
11-59	4-((2-Hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)-N-(2-(3,3,3-trifluoropropoxy)pyridin-4-yl)benzamide	0.058
11-60	4-((2-Hydroxyethyl)sulfonamido)-N-(2-methyl-6-(3,3,3-trifluoropropoxy)pyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.084
11-61	4-((2-Hydroxyethyl)sulfonamido)-N-(6-methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.070
11-62	N-(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.018
11-63	N-(2-(4,4-Difluoropiperidin-1-yl)-3-fluoropyridin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.022
11-64	N-(2-(4,4-Difluoropiperidin-1-yl)-3-fluoro-6-methylpyridin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.054
11-65	(R)-4-((2-Hydroxyethyl)sulfonamido)-N-(2-(2-methylmorpholino)pyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.151
11-66	(R)-4-((2-Hydroxyethyl)sulfonamido)-N-(2-methyl-6-(2-methylmorpholino)pyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.148
11-67	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.034
11-68	N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-(hydroxymethyl)cyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.053
11-69	N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-(hydroxymethyl)cyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.099
11-70	(S)-N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)-4-((tetrahydrofuran)-3-sulfonamido)benzamide	0.072
11-71	(R)-N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)-4-((tetrahydrofuran)-3-sulfonamido)benzamide	0.042
11-72	(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)-4-((tetrahydrofuran)-3-sulfonamido)benzamide	0.135
11-73	(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)-4-((tetrahydrofuran)-3-sulfonamido)benzamide	0.079
11-74	N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.094
11-75	N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.056
11-76	(R)-N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.020

Ex. #	Compound Name	KIF18A ATPase IC ₅₀ (μM)
11-77	(<i>S</i>)- <i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.034
11-78	4-((Cyclopropylmethyl)sulfonamido)- <i>N</i> -(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.125
11-79	4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(2-isopropoxy-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.057
11-80	(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-((tetrahydrofuran-3-yl)oxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.117
11-81	<i>N</i> -(2-(4-Fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(methylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.060
11-82	4-(Ethylsulfonamido)- <i>N</i> -(2-(4-fluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.051
11-83	<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1,1-dimethylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.098
12	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.076
13-1	(<i>S</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.057
13-2	(<i>R</i>)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.064
14	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxy-2-methylpropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.122
14-1	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(oxetan-3-ylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.182
14-2	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(ethylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.108
14-3	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(isopropylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.128
14-4	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-2-methylpropyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.806
14-5	4-(Cyclopropylsulfonyl)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.826
14-6	4-(<i>tert</i> -Butylsulfonyl)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	1.88
14-7	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(((3 <i>R</i> ,4 <i>R</i>)-4-hydroxytetrahydrofuran-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.094
14-8	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(((3 <i>S</i> ,4 <i>S</i>)-4-hydroxytetrahydrofuran-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.097
14-9	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(((1 <i>R</i> ,2 <i>R</i>)-2-hydroxycyclopentyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.195
15	<i>N</i> -(2-(4,4-Difluorocyclohexyl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.017
16-1	(<i>R</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.089
16-2	(<i>S</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.096

Ex. #	Compound Name	KIF18A ATPase IC ₅₀ (μM)
17	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(<i>N</i> -(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.023
17-1	4-(<i>N</i> -(2-Hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)- <i>N</i> -(2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)benzamide	0.027
17-2	4-(<i>N</i> -(2-Hydroxyethyl)sulfamoyl)- <i>N</i> -(6-methyl-2-(3,3,3-trifluoropropoxy)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.042
17-3	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.063
17-4	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyrimidin-4-yl)-4-(<i>N</i> -(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.023
17-5	<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.133
17-6	(<i>R</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(1-hydroxypropan-2-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.127
17-7	(<i>S</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(1-hydroxypropan-2-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.147
17-8	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(2-hydroxy-2-methylpropyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.286
18-1	2-(6-Azaspiro[2.5]octan-6-yl)-4-(<i>R</i> -cyclopropylsulfonimidoyl)- <i>N</i> -(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide	0.064
18-2	2-(6-azaspiro[2.5]octan-6-yl)-4-(<i>S</i> -cyclopropylsulfonimidoyl)- <i>N</i> -(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide	0.057
19-1	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(oxetane-3-sulfonimidoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.163
19-2	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(<i>S</i> -methylsulfonimidoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.438
19-3	2-(6-Azaspiro[2.5]octan-6-yl)- <i>N</i> -(2-(4,4-difluoro-1-pieridinyl)-6-methyl-4-pyrimidinyl)-4-(<i>S</i> -ethylsulfonimidoyl)benzamide	0.078
19-4	2-(6-Azaspiro[2.5]octan-6-yl)- <i>N</i> -(2-(4,4-difluoro-1-pieridinyl)-6-methyl-4-pyrimidinyl)-4-(<i>R</i> -ethylsulfonimidoyl)benzamide	0.035
19-5	<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(propan-2-ylsulfonimidoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.175
19-6	4-(Cyclopropanesulfonimidoyl)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.042
19-7	2-(6-Azaspiro[2.5]octan-6-yl)-4-(<i>S</i> -cyclopropylsulfonimidoyl)- <i>N</i> -(6-methy-2-((2 <i>R</i>)-2-methyl-4-moprpholinyl)-4-pyrimidinyl)benzamide	0.195
19-8	2-(6-Azaspiro[2.5]octan-6-yl)-4-(<i>R</i> -cyclopropylsulfonimidoyl)- <i>N</i> -(6-methy-2-((2 <i>R</i>)-2-methyl-4-moprpholinyl)-4-pyrimidinyl)benzamide	0.078
19-9	4-(Cyclopropanesulfonimidoyl)- <i>N</i> -(2-(3,3-difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.069
20	(<i>N</i> ¹ -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)terephthalamide	0.119
21	4-(Azetidin-3-ylsulfonyl)- <i>N</i> -(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.040
22	4- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylazetidin-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide	0.406


[0376] The foregoing invention has been described in some detail by way of illustration and example, for purposes of clarity and understanding. Those skilled in the art understand that changes and modifications may be practiced within the scope of the appended claims. Therefore, it is to be understood that the above

description is intended to be illustrative and not restrictive. The scope of the invention should, therefore, be determined not with reference to the above description, but should instead be determined with reference to the following appended claims, along with the full scope of equivalents to which such claims are entitled.

[0377] All patents, patent applications and publications cited herein are hereby incorporated by reference in their entirety for all purposes to the same extent as if each individual patent, patent application or publication were so individually denoted.

CLAIMS

1. A compound of formula I:



or any pharmaceutically-acceptable salt thereof, wherein:

X^1 is N or $-CR^6$;

R^1 is $-CN$, or a group $-Z-R^{12}$ wherein Z is $-C_{0-4}alk-$, $-NR^{11}-$, $-NR^{11}SO_2-$, $-SO_2NR^{11}-$, $-NR^{11}-S(=O)(=NH)-$, $-S(=O)(=NH)-$, $-S-$, $-S(=O)-$, $-SO_2-$, $C_{0-4}alk-O-$, $-(C=O)-$, $-(C=O)NR^{11}-$, $-C=N(OH)-$, or $-NR^{11}(C=O)$; or

the group $-Z-R^{12}$ is $-N=S(=O)-(R^{12})_2$, wherein the two R^{12} pair can alternatively combine with the sulfur atom attached to each of them to form a saturated or partially-saturated 3-, 4-, 5-, or 6-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S;

R^2 is halo or a group $-Y-R^{13}$, wherein Y is $-C_{0-4}alk-$, $-NH-C_{0-4}alkylene-$, $-N(CH_3)-C_{0-4}alkylene-$, $-C(=O)NR^a(C_{1-4}alk)$, $-O-C_{0-4}alk-$, S, $S=O$, $S(=O)_2$, $-SO_2NR^{13}$, or $-S(=O)(=NH)-$;

R^3 is H, $C_{1-4}alk$, or $C_{1-4}haloalk$;

R^4 is H, halo, R^{4a} or R^{4b} ;


R^5 is H, halo, $C_{1-8}alk$, or $C_{1-4}haloalk$;

R^6 is H, halo, $C_{1-8}alk$, $C_{1-4}haloalk$, $-O-C_{1-8}alk$, or $-O-R^{6a}$; wherein R^{6a} is a saturated or partially-saturated 3-, 4-, 5-, or 6-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S;

R^7 is H, halo, $C_{1-8}alk$, or $C_{1-4}haloalk$;

R^8 is H, halo, $C_{1-8}alk$, $C_{1-4}haloalk$, $-OH$, $-O-R^{8a}$, or $-O-R^{8b}$;

R^9 is H, halo, $C_{1-8}alk$, or $C_{1-4}haloalk$;



R^x is the group represented by the following formula:

Each of R^{10a} , R^{10b} , R^{10c} , R^{10d} , R^{10e} , R^{10f} , R^{10g} , R^{10h} , R^{10i} , and R^{10j} is H, halo, R^{10k} , or R^{10l} ;

or alternatively, each of R^{10a} and R^{10b} pair, R^{10c} and R^{10d} pair, R^{10e} and R^{10f} pair, R^{10g} and R^{10h} pair, or R^{10i} and R^{10j} pair, independently, can combine with the carbon atom attached to each of them to form a saturated or partially-saturated 3-, 4-, 5-, 6-membered monocyclic ring spiro to the R^x ring; wherein said 3-, 4-, 5-, 6-membered monocyclic ring contains 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S, and further wherein said 3-, 4-, 5-, 6-membered monocyclic ring is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, $C_{1-6}alk$, $C_{1-4}haloalk$, $-OR^a$, $-OC_{1-4}haloalk$, CN, $-NR^aR^a$, or oxo;

R^{11} is H, R^{11a} , or R^{11b} ;

R^{12} is H, R^{12a} , or R^{12b} ;

R^{13} is R^{13a} or R^{13b} ;

R^{4a} , R^{8a} , R^{10k} , R^{11a} , R^{12a} , and R^{13a} is independently, at each instance, selected from the group consisting of a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic or 5-, 6-, 7-, 8-, 9-, 10-, 11-, or 12-membered bicyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S, which is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, $C_{1-6}alk$,

$C_{1-4}haloalk$, $-OR^a$, $-OC_{1-4}haloalk$, CN, $-C(=O)R^b$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}alkNR^aR^a$, $-OC_{2-6}alkOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkNR^aR^a$, $-NR^aC_{2-6}alkOR^a$, $-C_{1-6}alkNR^aR^a$, $-C_{1-6}alkOR^a$, $-C_{1-6}alkN(R^a)C(=O)R^b$, $-C_{1-6}alkOC(=O)R^b$, $-C_{1-6}alkC(=O)NR^aR^a$, $-C_{1-6}alkC(=O)OR^a$, R^{14} , and oxo;

R^{4b} , R^{8b} , R^{10l} , R^{11b} , R^{12b} , and R^{13b} is independently, at each instance, selected from the group consisting of $C_{1-6}alk$ substituted by 0, 1, 2, 3, 4, or 5 group(s) selected from F, Cl, Br, $-OR^a$, $-OC_{1-4}haloalk$, or CN;

R^{14} is independently, at each instance, selected from the group consisting of a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic or 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, or 12-membered bicyclic ring containing 0, 1, 2 or 3 N atoms and 0 or 1 atoms selected from O and S, which is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, $C_{1-6}alk$, $C_{1-4}haloalk$, $-OR^a$,

$-OC_{1-4}haloalk$,

CN, $-C(=O)R^b$, $-C(=O)OR^a$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$,

$-OC_{2-6}alkNR^aR^a$, $-OC_{2-6}alkOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$,

$-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$,


$-NR^aC_{2-6}alkNR^aR^a$, $-NR^aC_{2-6}alkOR^a$, $-C_{1-6}alkNR^aR^a$, $-C_{1-6}alkOR^a$, $-C_{1-6}alkN(R^a)C(=O)R^b$,

$-C_{1-6}alkOC(=O)R^b$, $-C_{1-6}alkC(=O)NR^aR^a$, $-C_{1-6}alkC(=O)OR^a$, and oxo;


R^a is independently, at each instance, H or R^b ; and

R^b is independently, at each instance, $C_{1-6}alk$, phenyl, or benzyl, wherein the $C_{1-6}alk$ is being substituted by 0, 1, 2 or 3 substituents selected from halo, -OH, $-OC_{1-4}alk$, $-NH_2$, $-NHC_{1-4}alk$, $-OC(=O)C_{1-4}alk$, or $-N(C_{1-4}alk)C_{1-4}alk$; and the phenyl or benzyl is being substituted by 0, 1, 2 or 3 substituents selected from halo, $C_{1-4}alk$, $C_{1-3}haloalk$, -OH, $-OC_{1-4}alk$, $-NH_2$, $-NHC_{1-4}alk$, $-OC(=O)C_{1-4}alk$, or $-N(C_{1-4}alk)C_{1-4}alk$.

2. The compound of claim 1, or the pharmaceutically-acceptable salt thereof, wherein X^1 is N; having the formula (Ia):



3. The compound of claim 1, or the pharmaceutically-acceptable salt thereof, wherein X^1 is $-CR^6$; having the formula (Ib):




4. The compound of claim 1, or the pharmaceutically-acceptable salt thereof, wherein R^3 is H or methyl.

5. The compound of any one of claims 1 to 4, or the pharmaceutically-acceptable salt thereof, wherein each of R^{10c} , R^{10d} , R^{10e} , R^{10f} , R^{10g} , R^{10h} , R^{10i} , and R^{10j} is H, halo, C_{1-6} alk, or C_{1-4} haloalk; and each of R^{10a} and R^{10b} pair combine with the carbon atom attached to each of them form a saturated 3-, 4-, or 5-membered monocyclic ring spiro to the R^x ring; wherein said ring contains 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S.


6. The compound of any one of claims 1 to 5, or the pharmaceutically-acceptable salt thereof, wherein each of R^{10c} , R^{10d} , R^{10e} , R^{10f} , R^{10g} , R^{10h} , R^{10i} , and R^{10j} is H, methyl, or ethyl; and each of R^{10a} and R^{10b} pair combine with the carbon atom attached to each of them form a cyclopropyl, cyclobutyl, or cyclopentyl ring spiro to the R^x ring.

7. The compound of any one of claims 1 to 6, or the pharmaceutically-




acceptable salt thereof, wherein the group

is selected from:



8. The compound of any one of claims 1 to 7, or the pharmaceutically-



acceptable salt thereof, wherein the group


is

9. The compound of any one of claims 1 to 8, or the pharmaceutically-acceptable salt thereof, wherein R^1 is -CN, or a group -Z- R^{12} , wherein Z is a bond, -NH-, -NH SO_2 -, - SO_2 NH-, -S(=O)(=NH)-, -S-, -S(=O)-, - SO_2 -, -(C=O)-, -(C=O)NH-, or -NH(C=O)-; and

R^{12} is selected from:

(a) H;

(b) cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, oxiranyl, oxetanyl, tetrahydrofuranyl, azetidiny, imidazolyl, morpholiny, pyrrolidiny, piperaziny,



or 3 group(s) selected from wherein each ring is substituted by 0, 1, 2 or 3 OH, F,

methyl, $-\text{CH}_2\text{OH}$,

$-\text{C}(=\text{O})\text{OC}(\text{CH}_3)_3$, and CN ; or

(c) C_{1-6}alk substituted by 0, 1, 2 or 3 OH, or F.

10. The compound of any one of claims 1 to 9, or the pharmaceutically-acceptable salt thereof, wherein R^1 is $-\text{CN}$, or a group $-\text{Z}-\text{R}^{12}$, wherein Z is a bond, $-\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$,


$-\text{S}(=\text{O})(=\text{NH})-$, $-\text{S}-$, $-\text{S}(=\text{O})-$, $-\text{SO}_2-$, $-(\text{C}=\text{O})-$, $-(\text{C}=\text{O})\text{NH}-$, or $-\text{NH}(\text{C}=\text{O})-$; and

(a) R^{12} is H;

(b) R^{12} is oxetanyl, cyclopropyl; or

(c) R^{12} is C_{1-6}alk substituted by 0, 1, 2 or 3 OH group(s).

11. The compound of any one of claims 1 to 8, or the pharmaceutically-acceptable salt thereof, wherein the group $-\text{Z}-\text{R}^{12}$ is $-\text{N}=\text{S}(=\text{O})-(\text{R}^{12})_2$, wherein the two R^{12} pair can alternatively combine with the sulfur atom attached to each of them to form a saturated or partially-saturated 3-, 4-, 5-, or 6-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0, 1, or 2 atoms selected from O and S; which is selected from:



12. The compound of any one of claims 1 to 10, or the pharmaceutically-acceptable salt thereof, wherein R^1 is a group $-\text{Z}-\text{R}^{12}$, wherein Z is $-\text{NHSO}_2-$ or $-\text{SO}_2\text{NH}-$; and R^{12} is oxetanyl, cyclopropyl, or R^{12} is C_{1-6}alk substituted by 0, 1, 2 or 3 OH group(s).

13. The compound of any one of claims 1 to 10 or 12, or the pharmaceutically-acceptable salt thereof, wherein R^1 is a group $-\text{Z}-\text{R}^{12}$, wherein Z is $-\text{NHSO}_2-$ and R^{12} is $-\text{CH}_2-\text{CH}_2-\text{OH}$.

14. The compound of any one of claims 1 to 13, or the pharmaceutically-acceptable salt thereof, wherein R^2 is halo or a group $-\text{Y}-\text{R}^{13}$, wherein Y is a bond, $-\text{NH}-$, $-\text{NH}(\text{CH}_2)_{0-4}-$, or $-\text{O}(\text{CH}_2)_{0-4}-$; and

R^{13} is a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic or

4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, or 12-membered bicyclic ring containing 0, 1, 2 or 3 N atoms and 0 or 1 atoms selected from O and S, which is substituted by 0, 1, 2 or 3


group(s) selected from F, Cl, Br, C₁₋₆alk, C₁₋₄haloalk, -OH, -OC₁₋₄haloalk, CN, R¹⁴, and oxo; or


R¹³ is C₁₋₆alk substituted by 0, 1, 2, 3, 4, or 5 group(s) selected from F, Cl, Br, -OH, -OC₁₋₄haloalk, or CN.

15. The compound of any one of claims 1 to 14, or the pharmaceutically-acceptable salt thereof, wherein R² is a saturated 5- or 6-membered monocyclic ring wherein each said ring contains 0, 1, or 2 N atoms and 0 or 1 O atom, and wherein each said ring is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, C₁₋₆alk, C₁₋₄haloalk, -OH, -OC₁₋₄haloalk, CN, R¹⁴, and oxo.

16. The compound of any one of claims 1 to 15, or the pharmaceutically-acceptable salt thereof, wherein

R² is (a) halo; (b) a group -Y-R¹³, wherein Y is a bond; and R¹³ is morpholinyl, piperidinyl, azetidiny, pyrrolidinyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, piperazinyl, tetrahydrofuranyl,





each said ring is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, -OH, -OCHF₂, CN, and oxo; or

(c) a group -Y-R¹³, wherein Y is NH, -O-, -O-(CH₂)-, -O-(CH₂)-(CH₂)-, or


-O-(CH₂)-(CH₂)-(CH₂)-, and wherein R¹³ is ; or R¹³ is C₁₋₆alk substituted by 0, 1, 2, 3, 4, or 5 group(s) selected from F, Cl, Br, -OH, or CN.

17. The compound of any one of claims 1 to 16, or the pharmaceutically-acceptable salt thereof, wherein R² is morpholinyl or piperidinyl substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, methyl, CF₃, -OH, -OCHF₂, CN, or oxo.

18. The compound of any one of claims 1 to 17, or the pharmaceutically-acceptable salt thereof, of any one of claims 1-4, or the pharmaceutically-acceptable salt thereof, wherein R² is morpholinyl substituted by 1, 2 or 3 methyl group(s).

19. The compound of any one of claims 1 to 17, or the pharmaceutically-acceptable salt thereof, wherein R² is piperidinyl substituted by 1, 2 or 3 fluoro group(s).

20. The compound of any one of claims 1 to 16, or the pharmaceutically-acceptable salt thereof, wherein R² is



21. The compound of any one of claims 1 to 13, or the pharmaceutically-acceptable salt thereof, wherein Z is a bond, -NH-, -NHSO₂-, -SO₂NH-, -S(=O)(=NH)-, -S-, -S(=O)-, -SO₂-, -(C=O)-, -(C=O)NH-, or -NH(C=O)-.

22. The compound of any one of claims 1 to 13 or 21, or the pharmaceutically-acceptable salt thereof, wherein R¹² is selected from (a) H; (b) C₁₋₆alk substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, -OH, or -OCH₃; or (c) a saturated, partially-saturated or unsaturated 3-, 4-, 5-, 6-, or 7-membered monocyclic ring containing 0, 1, 2 or 3 N atoms and 0 or 1 atoms selected from O and

S, which is substituted by 0, 1, 2 or 3 group(s) selected from F, Cl, Br, C₁₋₆alk, C₁₋₄haloalk, -C₁₋₆alkOH, -OH, -OCH₃, -NH₂, or oxo.

23. The compound of any one of claims 1 to 13 or 21 to 22, or the pharmaceutically-acceptable salt thereof, wherein R¹² is selected from cyclopropyl, cyclobutyl, cyclopentyl, oxetanyl, azetidiny, tetrahydrofuranyl, or 1,3,4-oxathiazinanyl.

24. The compound of any one of claims 1 to 23, or the pharmaceutically-acceptable salt thereof, wherein R⁴ is selected from (a) H; (b) C₁₋₆alk substituted by 0, 1, 2 or 3 OH group(s); or (c) cyclopropyl.

25. The compound of any one of claims 1 to 24, or the pharmaceutically-acceptable salt thereof, wherein R⁴ is H or methyl.

26. The compound of any one of claims 1 to 25, or the pharmaceutically-acceptable salt thereof, wherein R⁵ is H.


27. The compound of any one of claims 1 or 3 to 26, or the pharmaceutically-acceptable salt thereof, wherein R⁶ is H or F.

28. The compound of any one of claims 1 to 27, or the pharmaceutically-acceptable salt thereof, wherein R⁷ is H or F.






29. The compound of any one of claims 1 to 28, or the pharmaceutically-acceptable salt thereof, wherein R⁸ is H.

30. The compound of any one of claims 1 to 29, or the pharmaceutically-acceptable salt thereof, wherein R⁹ is H.

31. The compound of any one of claims 1 to 30, selected from:

Ex. #	Chemical Structure	Name
1		<i>N</i> -(2-((1-Hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(methylsulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

Ex. #	Chemical Structure	Name
1-7		<i>N</i> -(2-((1-hydroxy-2-methylpropan-2-yl)amino)-6-methylpyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
2		<i>N</i> -(2-(2-Hydroxypropan-2-yl)pyrimidin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
2-4		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(<i>N</i> -(3-methyloxetan-3-yl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
2-8		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylcyclopropane)-1-sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
3		(<i>R</i>)-4-((2-Hydroxyethyl)sulfonamido)- <i>N</i> -(6-methyl-2-(2-methylmorpholino)pyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

Ex. #	Chemical Structure	Name
4		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
5-1		(<i>R</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
5-2		(<i>S</i>)- <i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
6-7		<i>N</i> -(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-(ethylsulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
7		<i>N</i> -(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

Ex. #	Chemical Structure	Name
8-1		(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
8-2		(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
10-1		(R)-N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
10-2		(S)-N-(2-(3,3-Difluoroazetidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxypropyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
12		N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

Ex. #	Chemical Structure	Name
13-1		(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
13-2		(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxypropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
14		N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-hydroxy-2-methylpropan-2-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
15		N-(2-(4,4-Difluorocyclohexyl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
16-1		(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

Ex. #	Chemical Structure	Name
16-2		(S)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-fluoro-1-(hydroxymethyl)ethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
17		N-(2-(4,4-Difluoropiperidin-1-yl)pyridin-4-yl)-4-(N-(2-hydroxyethyl)sulfamoyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
18-1		2-(6-Azaspiro[2.5]octan-6-yl)-4-(R-cyclopropylsulfonimidoyl)-N-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide
18-2		2-(6-Azaspiro[2.5]octan-6-yl)-4-(S-cyclopropylsulfonimidoyl)-N-(2-(4,4-difluoro-1-piperidinyl)-6-methyl-4-pyrimidinyl)benzamide
20		(N¹-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)terephthalamide

Ex. #	Chemical Structure	Name
21		4-(Azetidin-3-ylsulfonyl)-N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
22		N-(2-(4,4-difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((1-methylazetidin-3-yl)sulfonyl)-2-(6-azaspiro[2.5]octan-6-yl)benzamide

; or any pharmaceutically-acceptable salt thereof.

32. A pharmaceutical composition comprising the compound according to any one of claims 1 to 31, or the pharmaceutically acceptable salt thereof, and a pharmaceutically-acceptable diluent or carrier.

33. Use of a compound in accordance with any one of claims 1 to 31, or the pharmaceutical composition according to claim 32 in the preparation of a medicament for treating a condition, wherein said condition is cancer selected from the group consisting of (a) a solid or hematologically derived tumor selected from cancer of the cancer of the bladder, endometrial, lung squamous cell, breast, colon, kidney, liver, lung, small cell lung cancer, esophagus, gall-bladder, brain, head and neck, ovary, pancreas, stomach, cervix, thyroid, prostate and skin, (b) a hematopoietic tumor of lymphoid lineage selected from leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma and Burkett's lymphoma, (c) a hematopoietic tumor of myeloid lineage selected from acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia (d) a tumor of mesenchymal origin selected from fibrosarcoma and rhabdomyosarcoma, (e) a tumor of the central and peripheral nervous system selected from astrocytoma, neuroblastoma, glioma and schwannoma, and (f) a melanoma, seminoma,


teratocarcinoma, osteosarcoma, xenoderoma pigmentosum, keratoctanthoma, thyroid follicular cancer or Kaposi's sarcoma.

34. Use of the compound in accordance with any one of claims 1 to 31, or the pharmaceutically acceptable salt thereof, or the composition according to claim 32 in the manufacture of a medicament for reducing the size of a solid tumor in a subject.

35. Use of the compound in accordance with any one of claims 1 to 31, or the pharmaceutically acceptable salt thereof, or the composition according to claim 32 in the manufacture of a medicament for treating a cell proliferation disorder in a subject.


36. A compound, or pharmaceutically acceptable salts thereof, in accordance with any one of claims 1 to 31, or the pharmaceutically acceptable salt thereof, or the composition according to claim 32 for use in therapy.

37. The compound of claim 1, which is:

4		N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
---	--	--


or the pharmaceutically acceptable salt thereof.

38. The compound of claim 1, which is:

8-1		(R)-N-(2-(4,4-Difluoropiperidin-1-yl)-6-methylpyrimidin-4-yl)-5-fluoro-4-((2-hydroxy-1-methylethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
-----	---	--

or the pharmaceutically acceptable salt thereof.

39. The compound of claim 1, which is:

15		N-(2-(4,4-Difluorocyclohexyl)-6-methylpyrimidin-4-yl)-4-((2-hydroxyethyl)sulfonamido)-2-(6-azaspiro[2.5]octan-6-yl)benzamide
----	---	--

or the pharmaceutically acceptable salt thereof.