

MICROORGANISMS AND METHODS FOR THE CO-PRODUCTION OF ISOPROPANOL WITH PRIMARY ALCOHOLS, DIOLS AND ACIDS

BACKGROUND OF THE INVENTION

The present invention relates generally to biosynthetic processes, and more specifically to
5 organisms having n-propanol and isopropanol, 1,4-butanediol and isopropanol, 1,3-butanediol
and isopropanol or methacrylic and isopropanol biosynthetic capability.

Isopropanol (IPA) is a colorless, flammable liquid that mixes completely with most solvents,
including water. The largest use for IPA is as a solvent, including its well known yet small use
as "rubbing alcohol," which is a mixture of IPA and water. As a solvent, IPA is found in many
10 everyday products such as paints, lacquers, thinners, inks, adhesives, general-purpose cleaners,
disinfectants, cosmetics, toiletries, de-icers, and pharmaceuticals. Low-grade IPA is also used in
motor oils. The second largest use is as a chemical intermediate for the production of
isopropylamines, isopropylethers, and isopropyl esters. Isopropanol can potentially be
dehydrated to form propylene, a polymer precursor with an annual market of more than 2
15 million metric tons.

Current global production capacity of isopropanol (IPA) is approximately 6 B lb/yr, with
approximately 74% of global IPA capacity concentrated in the US, Europe, and Japan.
Isopropanol is manufactured by two petrochemical routes. The predominant process entails the
hydration of propylene either with or without sulfuric acid catalysis. Secondly, IPA is
20 produced via hydrogenation of acetone, which is a by-product formed in the production of
phenol and propylene oxide. High-priced propylene is currently driving costs up and margins
down throughout the chemical industry motivating the need for an expanded range of low cost
feedstocks.

n-Propanol can be potentially used as a gasoline substitute. It is currently used as a multi-
25 purpose solvent in the pharmaceutical industry, for surface coatings and in ink formulations. It
is used as a building block for resins and esters, propyl amines and halides. It is also used for
packaging and food contact applications. Global production of n-propanol in 2005 was more
than 140,000 metric tonnes.

n-Propanol is manufactured by the catalytic hydrogenation of propionaldehyde.
30 Propionaldehyde is itself produced via the oxo process, by hydroformylation of ethylene using
carbon monoxide and hydrogen in the presence of a catalyst such as cobalt octacarbonyl or a

rhodium complex. It is formed naturally in small amounts in many fermentation processes. For example, microbial production of very small quantities of n-propanol has been detected from certain species of *Clostridium* via threonine catabolism and from yeast in beer fermentation. No existing microorganism has been reported to produce 1-propanol from sugars in significant amounts.

- 1,4-Butanediol (14-BDO) is a polymer intermediate and industrial solvent with a global market of about 3 billion lb/year. BDO is currently produced from petrochemical precursors, primarily acetylene, maleic anhydride, and propylene oxide. For example, acetylene is reacted with 2 molecules of formaldehyde in the Reppe synthesis reaction (Kroschwitz and Grant, *Encyclopedia of Chem. Tech.*, John Wiley and Sons, Inc., New York (1999)), followed by catalytic hydrogenation to form 1,4-butanediol. Downstream, 14-BDO can be further transformed; for example, by oxidation to gamma-butyrolactone, which can be further converted to pyrrolidone and N-methyl-pyrrolidone, or hydrogenolysis to tetrahydrofuran. These compounds have varied uses as polymer intermediates, solvents, and additives, and have a combined market of nearly 2 billion lb/year. 1,3-Butanediol (13-BDO) is a four carbon diol commonly used as an organic solvent for food flavoring agents. It is also used as a co-monomer for polyurethane and polyester resins and is widely employed as a hypoglycaemic agent. Optically active 13-BDO is a useful starting material for the synthesis of biologically active compounds and liquid crystals. A substantial commercial use of 1,3-butanediol is subsequent dehydration to afford 1,3-butadiene (Ichikawa, *J. Mol. Catalysis*. 256:106-112 (2006)), a 25 billion lb/yr petrochemical used to manufacture synthetic rubbers (e.g., tires), latex, and resins. 13-BDO is traditionally produced from acetylene via its hydration. The resulting acetaldehyde is then converted to 3-hydroxybutyraldehyde which is subsequently reduced to form 1,3-BDO. In more recent years, acetylene has been replaced by ethylene as a source of acetaldehyde.
- Methylacrylic acid (MAA) is a key precursor of methyl methacrylate (MMA), a chemical intermediate with a global demand in excess of 4.5 billion pounds per year, much of which is converted to polyacrylates. The conventional process for synthesizing methyl methacrylate (i.e., the acetone cyanohydrin route) involves the conversion of hydrogen cyanide (HCN) and acetone to acetone cyanohydrin which then undergoes acid assisted hydrolysis and esterification with methanol to give MAA. Difficulties in handling potentially deadly HCN along with the high costs of byproduct disposal (1.2 tons of ammonium bisulfate are formed per ton of MAA) have sparked a great deal of research aimed at cleaner and more economical processes. As a starting material, MAA can easily be converted into MMA via esterification with methanol. No existing microorganism has been reported to produce MAA from sugars in significant amounts.

Microbial organisms and methods for effectively co-producing commercial quantities of n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol are described herein and include related advantages.

SUMMARY OF THE INVENTION

5 The invention provides non-naturally occurring microbial organisms having an n-propanol pathway and an isopropanol pathway. In one aspect, the embodiments disclosed herein relate to a non-naturally occurring microbial organism that includes a microbial organism having an n-propanol and an isopropanol pathway, where the n-propanol pathway includes at least one
10 exogenous nucleic acid encoding an n-propanol pathway enzyme expressed in a sufficient amount to produce n-propanol and where the isopropanol pathway includes at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol. In one aspect, the n-propanol pathway includes a propionaldehyde dehydrogenase, a propanol dehydrogenase, a propionyl-CoA:phosphate
15 propanoyltransferase, a propionyl-CoA hydrolase, a propionyl-CoA transferase, a propionyl-CoA synthetase, a propionate kinase, a propionate reductase or a propionyl phosphate reductase and the isopropanol pathway includes an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

In another embodiment, the invention provides a non-naturally occurring microbial organism
20 that includes a microbial organism having an n-propanol and an isopropanol pathway, where the n-propanol pathway includes a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol and where the isopropanol pathway includes a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol. In one aspect, the first set
25 encodes n-propanol pathway enzymes including a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a
30 propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase. In another aspect, the second set encodes isopropanol pathway enzymes including an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase, an

acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In another aspect, the invention provides a non-naturally occurring microbial organism having a first set of exogenous nucleic acids encoding n-propanol pathway enzymes and a second set of exogenous nucleic acids encoding isopropanol pathway enzymes, where the first set encodes a PEP carboxykinase or a PEP carboxylase; a malate dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA synthetase; a methylmalonyl-CoA mutase; a methylmalonyl-CoA decarboxylase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase and a propionyl phosphate reductase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the second set encodes a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In another aspect, the invention provides a non-naturally occurring microbial organism having a first set of exogenous nucleic acids encoding n-propanol pathway enzymes and a second set of exogenous nucleic acids encoding isopropanol pathway enzymes, where the first set encodes a PEP carboxykinase or a PEP carboxylase; a threonine deaminase; and a 2-oxobutanoate decarboxylase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the second set encodes a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-

CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In another aspect, the invention provides a non-naturally occurring microbial organism having a first set of exogenous nucleic acids encoding n-propanol pathway enzymes and a second set of exogenous nucleic acids encoding isopropanol pathway enzymes, where the first set encodes a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA carboxylase; a malonyl-CoA reductase; a malonate semialdehyde reductase; propionyl-CoA synthase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase and the second set encodes an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In another aspect, the invention provides a non-naturally occurring microbial organism having a first set of exogenous nucleic acids encoding n-propanol pathway enzymes and a second set of exogenous nucleic acids encoding isopropanol pathway enzymes, where the first set encodes a lactate dehydrogenase; a lactate-CoA transferase; a lactyl-CoA dehydratase; acryloyl CoA reductase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase and the second set encodes a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In another embodiment, the invention provides a non-naturally occurring microbial organism having an n-propanol pathway, the n-propanol pathway including at least one exogenous nucleic acid encoding an n-propanol pathway enzyme expressed in a sufficient amount to produce n-propanol. In one aspect the n-propanol pathway includes a propionaldehyde dehydrogenase, a
5 propanol dehydrogenase, a propionyl-CoA:phosphate propanoyltransferase, a propionyl-CoA hydrolase, a propionyl-CoA transferase, a propionyl-CoA synthetase, a propionate kinase, a propionate reductase or a propionyl phosphate reductase.

In another embodiment, the invention provides a non-naturally occurring microbial organism having an n-propanol pathway, the n-propanol pathway including a set of exogenous nucleic
10 acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the set of exogenous nucleic acids encoding a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl
15 phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase.

In still other aspects, embodiments disclosed herein relate to a method for producing n-propanol and isopropanol that includes culturing the aforementioned non-naturally occurring microbial
20 organisms. In still other aspect, embodiments disclosed herein relate to a method for producing n-propanol that includes culturing the aforementioned non-naturally occurring microbial organisms.

In one embodiment, the invention provides non-naturally occurring microbial organisms having an isopropanol pathway and a 1,4-butanediol (14-BDO) pathway, a 1,3-butanediol (13-BDO)
25 pathway or a methylacrylic acid (MAA) pathway. In one aspect, the embodiments disclosed herein relate to a non-naturally occurring microbial organism that includes a microbial organism having a 1,4-butanediol and an isopropanol pathway, where the 1,4-butanediol pathway includes at least one exogenous nucleic acid encoding a 1,4-butanediol pathway enzyme expressed in a sufficient amount to produce 1,4-butanediol and where the isopropanol pathway includes at least
30 one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol. In one aspect, the embodiments disclosed herein relate to a non-naturally occurring microbial organism that includes a microbial organism having a 1,3-butanediol and an isopropanol pathway, where the 1,3-butanediol pathway includes at least one

exogenous nucleic acid encoding a 1,3-butanediol pathway enzyme expressed in a sufficient amount to produce 1,3-butanediol and where the isopropanol pathway includes at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol. In one aspect, the embodiments disclosed herein relate to a

5 non-naturally occurring microbial organism that includes a microbial organism having a methylacrylic acid and an isopropanol pathway, where the methylacrylic acid pathway includes at least one exogenous nucleic acid encoding a methylacrylic acid pathway enzyme expressed in a sufficient amount to produce methylacrylic acid and where the isopropanol pathway includes at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a

10 sufficient amount to produce isopropanol.

In one embodiment, the isopropanol pathway comprises an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

In one embodiment, the 14-BDO pathway comprises a succinyl-CoA reductase, a succinate

15 reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde-forming), a 4-hydroxybutyraldehyde reductase, a 4-hydroxybutyrate reductase; a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, 4-hydroxybutyryl-phosphate reductase, or a 4-hydroxybutyryl-CoA reductase (alcohol-forming).

20 In one embodiment, the 13-BDO pathway comprises a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA dehydratase, a crotonase, a 3-hydroxybutyryl-CoA reductase (aldehyde forming), a 3-hydroxybutyraldehyde reductase, a 3-hydroxybutyryl-CoA

25 reductase (alcohol-forming), a 3-hydroxybutyryl-CoA transferase, a 3-hydroxybutyryl-CoA synthetase, a 3-hydroxybutyryl-CoA hydrolase, or a 3-hydroxybutyrate reductase.

In one embodiment, the MAA pathway comprises a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-

30 hydroxybutyrylase, a 4-hydroxybutyryl-CoA mutase, a 3-hydroxyisobutyryl-CoA dehydratase, a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase, a methacrylyl-CoA hydrolase, a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase, a 3-hydroxyisobutyryl-CoA hydrolase, a 3-hydroxyisobutyrate dehydratase, a methylmalonyl-CoA

mutase, a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase, a methylmalonyl-CoA hydrolase, a methylmalonate reductase, a methylmalonyl-CoA reductase (aldehyde forming), a 3-hydroxyisobutyrate dehydrogenase, a methylmalonyl-CoA reductase (alcohol forming) or a 3-hydroxyisobutyrate dehydratase.

- 5 In one embodiment, the invention provides a non-naturally occurring microbial organism that includes a microbial organism having an 14-BDO and an isopropanol pathway, where the 14-BDO pathway includes a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO and where the isopropanol pathway includes a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol.

- In one embodiment, the invention provides a non-naturally occurring microbial organism that includes a microbial organism having an 13-BDO and an isopropanol pathway, where the 13-BDO pathway includes a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO and where the isopropanol pathway includes a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol.

- In one embodiment, the invention provides a non-naturally occurring microbial organism that includes a microbial organism having an methylacrylic acid and an isopropanol pathway, where the methylacrylic acid pathway includes a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid and where the isopropanol pathway includes a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol.

- It is understood that methylacrylic acid pathways passing through a 3-hydroxyisobutyrate intermediate can be applied for 3-hydroxyisobutyrate production as opposed to methylacrylic acid production if the downstream enzyme, that is, a dehydratase, is omitted (see Figures 7 and 8). In this case, the non-naturally occurring organism would produce 3-hydroxyisobutyrate instead of methylacrylic acid. The non-naturally occurring organism could alternatively produce a mixture of 3-hydroxyisobutyrate and methylacrylic acid. The maximum molar yields of ATP and product will be unchanged regardless of whether methylacrylic acid or 3-hydroxyisobutyrate is produced.

It is further understood that, if desired, 3-hydroxyisobutyric acid expressed by a microbial organism of the invention can be chemically converted to methylacrylic acid. For example, 3-hydroxyisobutyric acid, or β -hydroxyisobutyric acid, can be dehydrated to form methylacrylic acid as described, for example, in U.S. Patent No. 7,186,856.

- 5 In still other aspects, embodiments disclosed herein relate to a method for producing 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol that includes culturing the aforementioned non-naturally occurring microbial organisms.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows an exemplary pathway for co-production of n-propanol and isopropanol from
 10 glucose. Abbreviations: Glc - glucose, PEP - phosphoenolpyruvate, PYR - pyruvate, FOR - formate, ACCOA - acetyl-CoA, AACOA - acetoacetyl-CoA, ACAC- acetoacetate, AC - acetone, PPOH-2 - isopropanol, OAA - oxaloacetate, MAL - malate, FUM - fumarate, SUCC - succinate, SUCCOA - succinyl- CoA, MMCOA- methylmalonyl-CoA, PPCOA - propionyl-CoA, PPA – propionate, PPAL - propionaldehyde, PPPi - propionyl phosphate, PPOH-1 – n-propanol.
 15

Figure 2 shows an exemplary pathway for co-production of n-propanol and isopropanol from glucose. Abbreviations: Glc - glucose, PEP - phosphoenolpyruvate, PYR - pyruvate, FOR - formate, ACCOA - acetyl-CoA, AACOA - acetoacetyl-CoA, ACAC- acetoacetate, AC - acetone, PPOH-2 - isopropanol, OAA - oxaloacetate, THR - threonine, 2-OBUT - 2-oxobutanoate, PPCOA - propionyl-CoA, PPA – propionate, PPAL- propionaldehyde, PPPi - propionyl phosphate, PPOH-1 – n-propanol.
 20

Figure 3 shows an exemplary pathway for co-production of n-propanol and isopropanol from glucose. Abbreviations: Glc - glucose, PEP - phosphoenolpyruvate, PYR - pyruvate, FOR - formate, ACCOA - acetyl-CoA, AACOA - acetoacetyl-CoA, ACAC- acetoacetate, AC - acetone, PPOH-2 - isopropanol, MALCOA – malonyl-CoA, MALAL- malonate semialdehyde, 3HP- 3-hydroxypropionate, PPCOA - propionyl-CoA, PPA – propionate, PPAL- propionaldehyde, PPPi - propionyl phosphate, PPOH-1 – n-propanol.
 25

Figure 4 shows an exemplary pathway for co-production of n-propanol and isopropanol from glucose. Abbreviations: Glc - glucose, PEP - phosphoenolpyruvate, PYR - pyruvate, FOR - formate, ACCOA - acetyl-CoA, AACOA - acetoacetyl-CoA, ACAC- acetoacetate, AC - acetone, PPOH-2 – isopropanol, LAC- D-lactate, LACCOA- lactoyl-CoA, ACRYLCOA-
 30

acryloyl-CoA, PPCoA - propionyl-CoA, PPA – propionate, PPAL - propionaldehyde, PPPi - propionyl phosphate, PPOH-1 – n-propanol.

- Figure 5 shows an exemplary pathway for coproduction of 1,4-BDO and isopropanol from glucose. Abbreviations: Glc - glucose, PEP - phosphoenolpyruvate, PYR - pyruvate, FOR - formate, ACCoA - acetyl-CoA, AACoA - acetoacetyl-CoA, ACAC- acetoacetate, AC - acetone, PPOH-2 - isopropanol, OAA - oxaloacetate, MAL - malate, FUM - fumarate, SUCC - succinate, SUCCoA - succinyl- CoA, SUCSAL – succinic semialdehyde, 4-HB – 4-hydroxybutyrate, 4-HBCoA – 4-hydroxybutyryl-CoA, 4-HBALD – 4-hydroxybutyraldehyde, 1,4-BDO – 1,4-butanediol, 4-HBP – 4-hydroxybutyryl-phosphate.
- Figure 6 shows an exemplary pathway for coproduction of 1,3-BDO and isopropanol from glucose. Abbreviations: Glc - glucose, PEP - phosphoenolpyruvate, PYR - pyruvate, FOR - formate, ACCoA - acetyl-CoA, AACoA - acetoacetyl-CoA, ACAC- acetoacetate, AC - acetone, PPOH-2 - isopropanol, OAA - oxaloacetate, MAL - malate, FUM - fumarate, SUCC - succinate, SUCCoA - succinyl- CoA, SUCSAL – succinic semialdehyde, 3-HB – 3-hydroxybutyrate, 4-HB – 4-hydroxybutyrate, 4-HBCoA – 4-hydroxybutyryl-CoA, CRTCoA – crotonyl-CoA, 3-HBCoA – 3-hydroxybutyryl-CoA, 3-HBALD – 3-hydroxybutyraldehyde, 1,3-BDO – 1,3-butanediol.

- Figure 7 shows an exemplary pathway for coproduction of methacrylic acid and isopropanol from glucose. Abbreviations: Glc - glucose, PEP - phosphoenolpyruvate, PYR - pyruvate, FOR - formate, ACCoA - acetyl-CoA, AACoA - acetoacetyl-CoA, ACAC- acetoacetate, AC - acetone, PPOH-2 - isopropanol, OAA - oxaloacetate, MACoA – methacrylyl-CoA, MAL - malate, FUM - fumarate, SUCC - succinate, SUCCoA - succinyl- CoA, SUCSAL – succinic semialdehyde, 4-HB – 4-hydroxybutyrate, 4-HBCoA – 4-hydroxybutyryl-CoA, 3-HIBCoA – 3-hydroxyisobutyryl-CoA, 3-HIB – 3-hydroxyisobutyrate, MAA – methacrylic acid.

- Figure 8 shows an exemplary pathway for coproduction of methacrylic acid and isopropanol from glucose. Abbreviations: Glc - glucose, PEP - phosphoenolpyruvate, PYR - pyruvate, FOR - formate, ACCoA - acetyl-CoA, AACoA - acetoacetyl-CoA, ACAC- acetoacetate, AC - acetone, PPOH-2 - isopropanol, OAA - oxaloacetate, MAL - malate, FUM - fumarate, SUCC - succinate, SUCCoA - succinyl- CoA, MM – methylmalonate, MMCoA- methylmalonyl-CoA, MMSA – methylmalonate semialdehyde, 3-HIB – 3-hydroxyisobutyrate, MAA – methacrylic acid.

DETAILED DESCRIPTION THE INVENTION

Embodiments of the present invention provide non-naturally occurring microbial organisms having redox-balanced anaerobic pathways for co-production of n-propanol and isopropanol from 3 phosphoenolpyruvate (PEP) molecules as exemplified in Figures 1-4. Some advantages of this co-production strategy include: (1) the co-production affords the maximum theoretical yield of n-propanol and isopropanol at 1.33 moles total/mole of glucose; and (2) the pathway for co-production is completely redox balanced and has a net positive yield of ATP. This facilitates a completely anaerobic production of the C3 alcohols as opposed to culturing microbial organisms having the isopropanol pathway alone, which requires aeration for regeneration of NAD.

Embodiments of the present invention also provide non-naturally occurring microbial organisms that can co-produce n-propanol and isopropanol from renewable resources as shown in Figures 1-4. Specifically, the organisms include all enzymes utilized in the co-production of n-propanol and isopropanol from acetyl-CoA and propionyl-CoA. Formate can be converted to carbon dioxide by a formate dehydrogenase that provides an additional reducing equivalent that can be used for n-propanol and isopropanol syntheses. Additionally, reducing equivalents can be obtained from other steps in the pathway, such as, the glycolysis pathway during conversion of glucose to phosphoenolpyruvate, pyruvate dehydrogenase or pyruvate ferredoxin oxidoreductase during conversion of pyruvate to acetyl-CoA, or 2-oxobutanoate dehydrogenase during conversion of 2-oxobutanoate to propionyl-CoA.

Embodiments of the present invention also provide non-naturally occurring microbial organisms that can produce n-propanol via propionyl-CoA. This conversion is carried out by two different enzymes: an aldehyde and alcohol dehydrogenase or in one step by a bifunctional aldehyde/alcohol dehydrogenase. Alternatively, propionyl-CoA can be converted into propionyl phosphate and then transformed into propionaldehyde by an acyl phosphate reductase. Alternatively, propionyl-CoA can be converted to propionate then to propionyl phosphate by a propionyl-CoA hydrolase, transferase, or synthetase and a propionate dinase, respectively. Alternatively, propionate can be converted to propionaldehyde by a propionate reductase. Pathways for production of propionyl-CoA are exemplified in Figures 1-4. In one embodiment, the pathway for production of propionyl-CoA proceeds via oxaloacetate as exemplified in Figure 1. Oxaloacetate is converted to propionyl-CoA by means of the reductive TCA cycle, a methylmutase, a decarboxylase, and a decarboxylase. An epimerase may be required to convert the (R) stereoisomer of methylmalonyl-CoA to the (S) configuration. In another embodiment,

the pathway for production of propionyl-CoA proceeds via threonine as exemplified in Figure 2. Oxaloacetate is converted into threonine by the native threonine pathway engineered for high yields. It is then deaminated to form 2-oxobutanoate and subsequently converted into propionyl-CoA. In one alternative, 2-oxobutanoate is converted to propionaldehyde by a decarboxylase, which is then reduced to n-propanol by a propanol dehydrogenase. In yet another embodiment, the pathway for production of propionyl-CoA proceeds via malonyl-CoA as exemplified in Figure 3. Acetyl-CoA is carboxylated to form malonyl-CoA. This is then reduced to malonate semialdehyde, and subsequently transformed into 3-hydroxypropionate (3HP). 3HP is converted into propionyl-CoA via propionyl-CoA synthase. In yet another embodiment, the pathway for production of propionyl-CoA proceeds via lactate as exemplified in Figure 4. Pyruvate is reduced to form lactate which is then activated to form lactoyl-CoA. The lactoyl-CoA is dehydrated to form acryloyl-CoA and then reduced to generate propionyl-CoA.

Embodiments of the present invention also provide non-naturally occurring microbial organisms that can produce isopropanol via acetyl-CoA. Isopropanol production is achieved via conversion of acetyl-CoA by an acetoacetyl-CoA thiolase, an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase,, an acetoacetate decarboxylase, and an isopropanol dehydrogenase as exemplified in Figures 1-4. In one embodiment the pathway for production of acetyl-CoA from glucose proceeds via phosphoenolpyruvate (PEP). Glucose is converted into PEP by the native glycolysis pathway of the microbial organism. PEP is converted to pyruvate by pyruvate kinase and then to acetyl-CoA by pyruvate dehydrogenase or pyruvate ferredoxin oxidoreductase. Alternatively, pyruvate is converted to acetyl-CoA and formate by pyruvate formate lyase. The formate is then converted to carbon dioxide and hydrogen by a formate dehydrogenase.

Embodiments of the present invention provide alternate methods for coproduction of isopropanol with the compounds 14-BDO, 13-BDO and MAA. The production of isopropanol proceeds via acetyl-CoA as described above. Alone this route is not redox-balanced and thus requires aeration to achieve high isopropanol yields. Embodiments described herein use this route and combine it with pathways for synthesizing the coproducts 1,4-butanediol (14-BDO), 1,3-butanediol (13-BDO) and methylacrylic acid (MAA). Coproduction routes are redox-balanced under anaerobic conditions as opposed to the requirement of oxygen if isopropanol is produced solely through acetone. Coproduction also provides related advantages, such as, the ease of separating isopropanol from other fermentation products due to its low boiling point (82°C) relative to 14-BDO (230°C), 13-BDO (203°C) and MAA (163°C) and the coproduction

using any of the microbial organisms described herein provides that maximum theoretical yield of the carbon from glucose is afforded.

Embodiments of the present invention provide non-naturally occurring microbial organisms that can produce 14-BDO via succinyl-CoA or in some aspects via succinate. For production of 14-BDO, succinyl-CoA is converted to succinic semialdehyde by a succinyl-CoA reductase. Alternatively, succinate can be converted to succinic semialdehyde by a succinate reductase. Next, succinic semialdehyde is reduced to 4-hydroxybutyrate by 4-hydroxybutyrate dehydrogenase. Activation of 4-HB to its acyl-CoA is catalyzed by a CoA transferase or synthetase. Alternatively, 4-HB can be converted into 4-hydroxybutyryl-phosphate and subsequently transformed into 4-HB-CoA by a phosphotrans-4-hydroxybutyrylase. 4-HB-CoA is then converted to 14-BDO by either a bifunctional CoA-dependent aldehyde/alcohol dehydrogenase, or by two separate enzymes with aldehyde and alcohol dehydrogenase activity. Yet another alternative that bypasses the 4-HB-CoA intermediate is direct reduction of 4-HB to 4-hydroxybutyrylaldehyde by a carboxylic acid reductase. 4-Hydroxybutyrylaldehyde is subsequently reduced to 14-BDO by an alcohol dehydrogenase. Yet another route that bypasses 4-HB-CoA entails reducing 4-hydroxybutyryl-phosphate to 4-hydroxybutyraldehyde by a phosphate reductase.

Embodiments of the present invention provide non-naturally occurring microbial organisms that can produce 13-BDO via succinyl-CoA or in some aspects via succinate. Production of 13-BDO also proceeds through 4-hydroxybutyryl-CoA, formed as described above. In this route, 4-hydroxybutyryl-CoA is dehydrated and isomerized to form crotonyl-CoA. The dehydration and vinylisomerisation reactions are catalyzed by a bifunctional enzyme, 4-hydroxybutyryl-CoA dehydratase. Crotonyl-CoA is then hydrated to 3-hydroxybutyryl-CoA. Removal of the CoA moiety and concurrent reduction yields 3-hydroxybutyraldehyde. Alternatively, 3-hydroxybutyryl-CoA is converted to 3-hydroxybutyrate by a 3-hydroxybutyryl-CoA transferase, hydrolase, or synthetase and then reduced by a 3-hydroxybutyrate reductase to yield 3-hydroxybutyraldehyde. Finally reduction of the aldehyde by 3-hydroxybutyraldehyde reductase yields 13-BDO.

Embodiments of the present invention provide non-naturally occurring microbial organisms that can produce MAA via two alternative routes. The first route proceeds through 4-hydroxybutyryl-CoA, formed as described above. 4-Hydroxybutyryl-CoA is converted to 3-hydroxyisobutyryl-CoA by a methyl mutase. The CoA moiety of 3-Hydroxyisobutyryl-CoA is then removed by a CoA transferase, hydrolase or synthetase. Finally, dehydration of the 3-

hydroxy group yields MAA. Alternatively, 3-hydroxyisobutyryl-CoA is converted to methacrylyl-CoA by a 3-hydroxyisobutyryl-CoA dehydratase and then the CoA moiety is removed by a CoA transferase, hydrolase or synthetase to yield MAA. In the alternate MAA production route, succinyl-CoA is converted to methylmalonyl-CoA by methylmalonyl-CoA mutase. An epimerase may be required to convert the (R) stereoisomer of methylmalonyl-CoA to the (S) configuration. A CoA-dependent aldehyde dehydrogenase then converts methylmalonyl-CoA to methylmalonate semialdehyde. Alternatively, the CoA moiety of (R)-methylmalonyl-CoA or (S)-methylmalonyl-CoA is removed by a CoA transferase, hydrolase or synthetase to form methylmalonate, which is then converted to the semialdehyde by a reductase.

Reduction of the aldehyde to 3-hydroxyisobutyrate, followed by dehydration, yields MAA. Alternately, methylmalonyl-CoA is converted to 3-hydroxyisobutyrate by an alcohol-forming CoA reductase.

Embodiments of the present invention provide non-naturally occurring microbial organisms having pathways for production of succinyl-CoA as exemplified in Figures 5-8. In one embodiment, the pathway for production of succinyl-CoA proceeds via oxaloacetate. Oxaloacetate is converted to succinyl-CoA by means of the reductive TCA cycle, including a malate dehydrogenase, a fumerase, a fumarate reductase and a succinyl-CoA transferase or alternatively a succinyl-CoA synthetase.

Engineering these pathways into a microorganism involves cloning an appropriate set of genes encoding a set of enzymes into a production host described herein, optimizing fermentation conditions, and assaying product formation following fermentation. To engineer a production host for the production of n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol, one or more exogenous DNA sequence(s) can be expressed in a microorganism. In addition, the microorganism can have endogenous gene(s) functionally disrupted, deleted or overexpressed. The metabolic modifications disclosed herein enable the production of n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol using renewable feedstock.

In some embodiments, the invention provides non-naturally occurring microbial organisms that include at least one exogenous nucleic acid that encode an n-propanol pathway enzyme expressed in a sufficient amount to produce n-propanol.

In another embodiment, the invention provides non-naturally occurring microbial organisms that include at least one exogenous nucleic acid that encode an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol.

In still other embodiments, the invention provides methods for co-producing n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol. Such methods involve culturing the microbial organisms described herein.

As used herein, the term “non-naturally occurring” when used in reference to a microbial
5 organism or microorganism of the invention is intended to mean that the microbial organism has at least one genetic alteration not normally found in a naturally occurring strain of the referenced species, including wild-type strains of the referenced species. Genetic alterations include, for example, modifications introducing expressible nucleic acids encoding metabolic polypeptides, other nucleic acid additions, nucleic acid deletions and/or other functional disruption of the
10 microbial organism’s genetic material. Such modifications include, for example, coding regions and functional fragments thereof, for heterologous, homologous or both heterologous and homologous polypeptides for the referenced species. Additional modifications include, for example, non-coding regulatory regions in which the modifications alter expression of a gene or operon. Exemplary metabolic polypeptides include enzymes or proteins within an n-propanol,
15 an isopropanol, a 14-BDO, a 13-BDO and/or MAA biosynthetic pathways.

A metabolic modification refers to a biochemical reaction that is altered from its naturally occurring state. Therefore, non-naturally occurring microorganisms can have genetic modifications to nucleic acids encoding metabolic polypeptides or, functional fragments thereof. Exemplary metabolic modifications are disclosed herein.

As used herein, the term “isolated” when used in reference to a microbial organism are intended
20 to mean an organism that is substantially free of at least one component as the referenced microbial organism is found in nature. The term includes a microbial organism that is removed from some or all components as it is found in its natural environment. The term also includes a microbial organism that is removed from some or all components as the microbial organism is
25 found in non-naturally occurring environments. Therefore, an isolated microbial organism is partly or completely separated from other substances as it is found in nature or as it is grown, stored or subsisted in non-naturally occurring environments. Specific examples of isolated microbial organisms include partially pure microbes, substantially pure microbes and microbes cultured in a medium that is non-naturally occurring.

As used herein, the terms “microbial,” “microbial organism” or “microorganism” is intended to
30 mean any organism that exists as a microscopic cell that is included within the domains of archaea, bacteria or eukarya. Therefore, the term is intended to encompass prokaryotic or eukaryotic cells or organisms having a microscopic size and includes bacteria, archaea and

eubacteria of all species as well as eukaryotic microorganisms such as yeast and fungi. The term also includes cell cultures of any species that can be cultured for the production of a biochemical.

As used herein, "n-propanol" is intended to mean a primary alcohol with the molecular formula of C_3H_8O and a molecular mass of 60.1 g/mol. N-propanol is also known in the art as 1-propanol, 1-propyl alcohol, n-propyl alcohol, propan-1-ol, or simply propanol. N-propanol is an isomer of isopropanol.

As used herein, "isopropanol" is intended to mean a secondary alcohol, with the molecular formula of C_3H_8O and a molecular mass of 60.1 g/mol, wherein the alcohol carbon is attached to two other carbons. This attachment is sometimes shown as $(CH_3)_2CHOH$. Isopropanol is also known in the art as propan-2-ol, 2-propanol or the abbreviation IPA. Isopropanol is an isomer of n-propanol.

As used herein, the term "1,4-butanediol" is intended to mean an alcohol derivative of the alkane butane, carrying two hydroxyl groups which has the chemical formula $C_4H_{10}O_2$ and a molecular mass of 90.12 g/mol. The chemical compound 1,4-butanediol also is known in the art as 1,4-BDO and is a chemical intermediate or precursor for a family of compounds commonly referred to as the BDO family of compounds.

As used herein, the term "1,3-butanediol" is intended to mean one of four stable isomers of butanediol having the chemical formula $C_4H_{10}O_2$ and a molecular mass of 90.12 g/mol. The chemical compound 1,3-butanediol is known in the art as 13-BDO or β -butane glycol and is also a chemical intermediate or precursor for a family of compounds commonly referred to as the BDO family of compounds.

As used herein, "methacrylic acid," having the chemical formula $CH_2=C(CH_3)CO_2$ (also known as methacrylic acid and IUPAC name 2-methyl-2-propenoic acid), is the acid form of methylacrylate, and it is understood that methacrylic acid and methylacrylate can be used interchangeably throughout to refer to the compound in any of its neutral or ionized forms, including any salt forms thereof. It is understood by those skilled in the art that the specific form will depend on the pH. Similarly, 3-hydroxyisobutyrate and 3-hydroxyisobutyric acid can be used interchangeably throughout to refer to the compound in any of its neutral or ionized forms, including any salt forms thereof.

As used herein, the term “CoA” or “coenzyme A” is intended to mean an organic cofactor or prosthetic group (nonprotein portion of an enzyme) whose presence is required for the activity of many enzymes (the apoenzyme) to form an active enzyme system. Coenzyme A functions in certain condensing enzymes, acts in acetyl or other acyl group transfer and in fatty acid synthesis and oxidation, pyruvate oxidation and in other acetylation.

As used herein, the term “substantially anaerobic” when used in reference to a culture or growth condition is intended to mean that the amount of oxygen is less than about 10% of saturation for dissolved oxygen in liquid media. The term also is intended to include sealed chambers of liquid or solid medium maintained with an atmosphere of less than about 1% oxygen.

- “Exogenous” as it is used herein is intended to mean that the referenced molecule or the referenced activity is introduced into the host microbial organism. The molecule can be introduced, for example, by introduction of an encoding nucleic acid into the host genetic material such as by integration into a host chromosome or as non-chromosomal genetic material such as a plasmid. Therefore, the term as it is used in reference to expression of an encoding nucleic acid refers to introduction of the encoding nucleic acid in an expressible form into the microbial organism. When used in reference to a biosynthetic activity, the term refers to an activity that is introduced into the host reference organism. The source can be, for example, a homologous or heterologous encoding nucleic acid that expresses the referenced activity following introduction into the host microbial organism. Therefore, the term “endogenous” refers to a referenced molecule or activity that is present in the host. Similarly, the term when used in reference to expression of an encoding nucleic acid refers to expression of an encoding nucleic acid contained within the microbial organism. The term “heterologous” refers to a molecule or activity derived from a source other than the referenced species whereas “homologous” refers to a molecule or activity derived from the host microbial organism. Accordingly, exogenous expression of an encoding nucleic acid of the invention can utilize either or both a heterologous or homologous encoding nucleic acid.

- The non-naturally occurring microbial organisms of the invention can contain stable genetic alterations, which refers to microorganisms that can be cultured for greater than five generations without loss of the alteration. Generally, stable genetic alterations include modifications that persist greater than 10 generations, particularly stable modifications will persist more than about 25 generations, and more particularly, stable genetic modifications will be greater than 50 generations, including indefinitely.

Those skilled in the art will understand that the genetic alterations, including metabolic modifications exemplified herein, are described with reference to a suitable host organism such as *E. coli* and their corresponding metabolic reactions or a suitable source organism for desired genetic material such as genes for a desired metabolic pathway. However, given the complete
5 genome sequencing of a wide variety of organisms and the high level of skill in the area of genomics, those skilled in the art will readily be able to apply the teachings and guidance provided herein to essentially all other organisms. For example, the *E. coli* metabolic alterations exemplified herein can readily be applied to other species by incorporating the same or analogous encoding nucleic acid from species other than the referenced species. Such genetic
10 alterations include, for example, genetic alterations of species homologs, in general, and in particular, orthologs, paralogs or nonorthologous gene displacements.

An ortholog is a gene or genes that are related by vertical descent and are responsible for substantially the same or identical functions in different organisms. For example, mouse epoxide hydrolase and human epoxide hydrolase can be considered orthologs for the biological
15 function of hydrolysis of epoxides. Genes are related by vertical descent when, for example, they share sequence similarity of sufficient amount to indicate they are homologous, or related by evolution from a common ancestor. Genes can also be considered orthologs if they share three-dimensional structure but not necessarily sequence similarity, of a sufficient amount to indicate that they have evolved from a common ancestor to the extent that the primary sequence
20 similarity is not identifiable. Genes that are orthologous can encode proteins with sequence similarity of about 25% to 100% amino acid sequence identity. Genes encoding proteins sharing an amino acid similarity less than 25% can also be considered to have arisen by vertical descent if their three-dimensional structure also shows similarities. Members of the serine protease family of enzymes, including tissue plasminogen activator and elastase, are considered to have
25 arisen by vertical descent from a common ancestor.

Orthologs include genes or their encoded gene products that through, for example, evolution, have diverged in structure or overall activity. For example, where one species encodes a gene product exhibiting two functions and where such functions have been separated into distinct genes in a second species, the three genes and their corresponding products are considered to be
30 orthologs. For the production of a biochemical product, those skilled in the art will understand that the orthologous gene harboring the metabolic activity to be introduced or disrupted is to be chosen for construction of the non-naturally occurring microorganism. An example of orthologs exhibiting separable activities is where distinct activities have been separated into distinct gene products between two or more species or within a single species. A specific example is the

separation of elastase proteolysis and plasminogen proteolysis, two types of serine protease activity, into distinct molecules as plasminogen activator and elastase. A second example is the separation of mycoplasma 5'-3' exonuclease and Drosophila DNA polymerase III activity. The DNA polymerase from the first species can be considered an ortholog to either or both of the
5 exonuclease or the polymerase from the second species and vice versa.

In contrast, paralogs are homologs related by, for example, duplication followed by evolutionary divergence and have similar or common, but not identical functions. Paralogs can originate or derive from, for example, the same species or from a different species. For example, microsomal epoxide hydrolase (epoxide hydrolase I) and soluble epoxide hydrolase (epoxide
10 hydrolase II) can be considered paralogs because they represent two distinct enzymes, co-evolved from a common ancestor, that catalyze distinct reactions and have distinct functions in the same species. Paralogs are proteins from the same species with significant sequence similarity to each other suggesting that they are homologous, or related through co-evolution from a common ancestor. Groups of paralogous protein families include HipA homologs,
15 luciferase genes, peptidases, and others.

A nonorthologous gene displacement is a nonorthologous gene from one species that can substitute for a referenced gene function in a different species. Substitution includes, for example, being able to perform substantially the same or a similar function in the species of origin compared to the referenced function in the different species. Although generally, a
20 nonorthologous gene displacement will be identifiable as structurally related to a known gene encoding the referenced function, less structurally related but functionally similar genes and their corresponding gene products nevertheless will still fall within the meaning of the term as it is used herein. Functional similarity requires, for example, at least some structural similarity in the active site or binding region of a nonorthologous gene product compared to a gene encoding
25 the function sought to be substituted. Therefore, a nonorthologous gene includes, for example, a paralog or an unrelated gene.

Therefore, in identifying and constructing the non-naturally occurring microbial organisms of the invention having n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol biosynthetic capability, those skilled in the art will
30 understand with applying the teaching and guidance provided herein to a particular species that the identification of metabolic modifications can include identification and inclusion or inactivation of orthologs. To the extent that paralogs and/or nonorthologous gene displacements are present in the referenced microorganism that encode an enzyme catalyzing a similar or

substantially similar metabolic reaction, those skilled in the art also can utilize these evolutionally related genes.

Orthologs, paralog and nonorthologous gene displacements can be determined by methods well known to those skilled in the art. For example, inspection of nucleic acid or amino acid sequences for two polypeptides will reveal sequence identity and similarities between the compared sequences. Based on such similarities, one skilled in the art can determine if the similarity is sufficiently high to indicate the proteins are related through evolution from a common ancestor. Algorithms well known to those skilled in the art, such as Align, BLAST, Clustal W and others compare and determine a raw sequence similarity or identity, and also determine the presence or significance of gaps in the sequence which can be assigned a weight or score. Such algorithms also are known in the art and are similarly applicable for determining nucleotide sequence similarity or identity. Parameters for sufficient similarity to determine relatedness are computed based on well known methods for calculating statistical similarity, or the chance of finding a similar match in a random polypeptide, and the significance of the match determined. A computer comparison of two or more sequences can, if desired, also be optimized visually by those skilled in the art. Related gene products or proteins can be expected to have a high similarity, for example, 25% to 100% sequence identity. Proteins that are unrelated can have an identity which is essentially the same as would be expected to occur by chance, if a database of sufficient size is scanned (about 5%). Sequences between 5% and 24% may or may not represent sufficient homology to conclude that the compared sequences are related. Additional statistical analysis to determine the significance of such matches given the size of the data set can be carried out to determine the relevance of these sequences.

Exemplary parameters for determining relatedness of two or more sequences using the BLAST algorithm, for example, can be as set forth below. Briefly, amino acid sequence alignments can be performed using BLASTP version 2.0.8 (Jan-05-1999) and the following parameters: Matrix: 0 BLOSUM62; gap open: 11; gap extension: 1; x_dropoff: 50; expect: 10.0; wordsize: 3; filter: on. Nucleic acid sequence alignments can be performed using BLASTN version 2.0.6 (Sept-16-1998) and the following parameters: Match: 1; mismatch: -2; gap open: 5; gap extension: 2; x_dropoff: 50; expect: 10.0; wordsize: 11; filter: off. Those skilled in the art will know what modifications can be made to the above parameters to either increase or decrease the stringency of the comparison, for example, and determine the relatedness of two or more sequences.

In one embodiment, the invention provides a non-naturally occurring microbial organism, including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway having at least one exogenous nucleic acid encoding an n-propanol pathway enzyme expressed in a sufficient amount to produce n-propanol, the n-propanol pathway including a propionaldehyde dehydrogenase, a propanol dehydrogenase, a propionyl-CoA:phosphate propanoyltransferase, a propionyl-CoA hydrolase, a propionyl-CoA transferase, a propionyl-CoA synthetase, a propionate kinase, a propionate reductase or a propionyl phosphate reductase, the isopropanol pathway comprising at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, the isopropanol pathway including an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

In a further aspect of the above embodiment, the microbial organism has an acetyl-CoA pathway having at least one exogenous nucleic acid encoding an acetyl-CoA pathway enzyme expressed in a sufficient amount to produce acetyl-CoA, the acetyl-CoA pathway including a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, or a formate dehydrogenase.

In further embodiment, the microbial organism has a propionyl-CoA pathway having at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, the propionyl-CoA pathway including a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a methylmalonyl-CoA mutase, a methylmalonyl-CoA epimerase or a methylmalonyl-CoA decarboxylase. In a further aspect, the propionyl-CoA pathway includes a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

In another further embodiment, the microbial organism has a propionyl-CoA pathway having at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, the propionyl-CoA pathway including a PEP carboxykinase, a PEP carboxylase, a threonine deaminase, or a 2-oxobutanoate dehydrogenase. In a further aspect, the n-propanol pathway includes 2-oxobutanoate decarboxylase.

In another further embodiment, the microbial organism has a propionyl-CoA pathway having at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, the propionyl-CoA pathway including an acetyl-

CoA carboxylase, a malonyl-CoA reductase, a malonate semialdehyde reductase or propionyl-CoA synthase.

In another further embodiment, the microbial organism has a propionyl-CoA pathway having at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a
5 sufficient amount to produce propionyl-CoA, the propionyl-CoA pathway including a lactate dehydrogenase, a lactate-CoA transferase, a lactyl-CoA dehydratase or acryloyl CoA reductase.

In yet another embodiment, the invention provides a non-naturally occurring microbial organism, including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway having a first set of exogenous nucleic acids encoding n-
10 propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids encoding a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase
15 and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the isopropanol pathway having a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase
20 or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In a further aspect of the above embodiment, the microbial organism has an acetyl-CoA pathway having a third set of exogenous nucleic acids encoding acetyl-CoA pathway enzymes expressed in a sufficient amount to produce acetyl-CoA, the third set of exogenous nucleic acids encoding
25 a pyruvate kinase; and a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase.

In another further embodiment, the microbial organism has a propionyl-CoA pathway having a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a
30 sufficient amount to produce propionyl-CoA, the third set of exogenous nucleic acids encoding a PEP carboxykinase or a PEP carboxylase; a malate dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA synthetase; a methylmalonyl-CoA mutase; and a methylmalonyl-CoA decarboxylase. In a further aspect, the third set of

exogenous nucleic acids further encodes a methylmalonyl-CoA epimerase, a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

- In another further embodiment, the microbial organism has a propionyl-CoA pathway having a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA, said third set of exogenous nucleic acids encoding a PEP carboxykinase or a PEP carboxylase; a threonine deaminase; and a 2-oxobutanoate dehydrogenase. In a further aspect, the third set of exogenous nucleic acids further encodes a methylmalonyl-CoA decarboxylase or a pyruvate carboxylase. In yet another aspect, the second set of exogenous nucleic acids further encodes a 2-oxobutanoate decarboxylase.
- 10 In another further embodiment, the microbial organism has a propionyl-CoA pathway having a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA, the third set of exogenous nucleic acids encoding an acetyl-CoA carboxylase; a malonyl-CoA reductase; a malonate semialdehyde reductase; and propionyl-CoA synthase.
- 15 In another further embodiment, the microbial organism has a propionyl-CoA pathway having a third set of exogenous nucleic acids encoding a lactate dehydrogenase; a lactate-CoA transferase; a lactyl-CoA dehydratase; and acryloyl CoA reductase.

- In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids encoding a PEP carboxykinase or a PEP carboxylase; a malate dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA synthetase; a methylmalonyl-CoA mutase; a methylmalonyl-CoA decarboxylase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase and a propionyl phosphate reductase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a

pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids encoding a PEP carboxykinase or a PEP carboxylase; a threonine deaminase; and a 2-oxobutanoate decarboxylase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; a 2-oxobutanoate dehydrogenase, a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase. In a further aspect, the second set of exogenous nucleic acids further encodes a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids encoding a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA carboxylase; a malonyl-CoA reductase; a malonate semialdehyde reductase; propionyl-CoA synthase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate

propanoyltransferase, a propionyl phosphate reductase and propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway including a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids encoding a lactate dehydrogenase; a lactate-CoA transferase; a lactyl-CoA dehydratase; acryloyl CoA reductase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway, the n-propanol pathway comprising at least one exogenous nucleic acid encoding an n-propanol pathway enzyme expressed in a sufficient amount to produce n-propanol, the n-propanol pathway including a propionaldehyde dehydrogenase, a propanol dehydrogenase, a propionyl-CoA:phosphate

propanoyltransferase, a propionyl-CoA hydrolase, a propionyl-CoA transferase, a propionyl-CoA synthetase, a propionate kinase, a propionate reductase, or a propionyl phosphate reductase.

In another embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway, the n-propanol pathway comprising a set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the set of exogenous nucleic acids encoding a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase.

In a further aspect of the above embodiment, the non-naturally occurring microbial organism having an n-propanol pathway also has a propionyl-CoA pathway including exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA as exemplified herein. For example, in some aspects the exogenous nucleic acids encode a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a methylmalonyl-CoA mutase or a methylmalonyl-CoA decarboxylase. In another aspect, the exogenous nucleic acids further encode a methylmalonyl-CoA epimerase. Additionally, in yet another aspect of the above embodiment, the non-naturally occurring microbial organism having an n-propanol pathway can have a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, wherein the first set of exogenous nucleic acids encode a PEP carboxykinase or a PEP carboxylase; a malate dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA synthetase; a methylmalonyl-CoA mutase; a methylmalonyl-CoA epimerase, a methylmalonyl-CoA decarboxylase; a propionaldehyde dehydrogenase and a propanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism, including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway having at least one exogenous nucleic acid encoding an 14-BDO pathway enzyme expressed in a sufficient amount to produce 14-BDO, the 14-BDO pathway including a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA

reductase (aldehyde-forming), a 4-hydroxybutyraldehyde reductase, a 4-hydroxybutyrate reductase; a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-phosphate reductase or a 4-hydroxybutyryl-CoA reductase (alcohol-forming), the isopropanol pathway including at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, the isopropanol pathway including an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism, including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway having at least one exogenous nucleic acid encoding an 13-BDO pathway enzyme expressed in a sufficient amount to produce 13-BDO, the 13-BDO pathway including a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA dehydratase, a crotonase, a 3-hydroxybutyryl-CoA reductase (aldehyde forming), a 3-hydroxybutyraldehyde reductase, a 3-hydroxybutyryl-CoA transferase, a 3-hydroxybutyryl-CoA synthetase, a 3-hydroxybutyryl-CoA hydrolase, or a 3-hydroxybutyrate reductase, or a 3-hydroxybutyryl-CoA reductase (alcohol-forming), the isopropanol pathway including at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, the isopropanol pathway including an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism, including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway having at least one exogenous nucleic acid encoding an MAA pathway enzyme expressed in a sufficient amount to produce MAA, the MAA pathway including a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA mutase, a 3-hydroxyisobutyryl-CoA dehydratase, a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase, a methacrylyl-CoA hydrolase, a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase, a 3-hydroxyisobutyryl-CoA hydrolase, a 3-hydroxyisobutyrate dehydratase, a methylmalonyl-CoA mutase, a methylmalonyl-CoA epimerase, a methylmalonyl-CoA transferase, a methylmalonyl-

CoA synthetase, a methylmalonyl-CoA hydrolase, a methylmalonate reductase, a methylmalonyl-CoA reductase (aldehyde forming), a 3-hydroxyisobutyrate dehydrogenase, a methylmalonyl-CoA reductase (alcohol forming) or a 3-hydroxyisobutyrate dehydratase, the isopropanol pathway including at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, the isopropanol pathway including an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

10 In a further aspect of the above embodiments, the microbial organism has an acetyl-CoA pathway having at least one exogenous nucleic acid encoding an acetyl-CoA pathway enzyme expressed in a sufficient amount to produce acetyl-CoA, the acetyl-CoA pathway including a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, or a formate dehydrogenase.

15 In further aspect of the above embodiments, the microbial organism has a succinyl-CoA pathway having at least one exogenous nucleic acid encoding a succinyl-CoA pathway enzyme expressed in a sufficient amount to produce succinyl-CoA, the succinyl-CoA pathway including a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase or a succinyl-CoA synthetase. In a further aspect, the succinyl-CoA pathway includes a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

20 In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate reductase; and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a 4-hydroxybutyryl-phosphate reductase; and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-

CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate reductase; and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a 4-hydroxybutyryl-phosphate reductase; and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the

second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of

exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- 5 In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-
- 10 hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3 hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient
- 15 amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the
- 20 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a
- 25 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- 30 In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic

- acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a
- 5 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.
- 10 In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a
- 15 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase
- 20 or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway
- 25 enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding
- 30 isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3 hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce

isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4 hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3 hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a

crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA

pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-
5 hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an
10 acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes
15 expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4 hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase, and the isopropanol pathway comprising a second
20 set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism
25 including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-
30 hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an

acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4 hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-

hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a methylmalonyl-CoA mutase; a methylmalonyl-CoA reductase (aldehyde forming); a 3-hydroxyisobutyrate dehydrogenase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a methylmalonyl-CoA mutase; a methylmalonyl-CoA epimerase; a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase, or a methylmalonyl-CoA hydrolase; a methylmalonate reductase; a 3-hydroxyisobutyrate dehydrogenase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids

encoding a methylmalonyl-CoA mutase; a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase or a methylmalonyl-CoA hydrolase; a methylmalonate reductase; a 3-hydroxyisobutyrate dehydrogenase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a methylmalonyl-CoA mutase; a methylmalonyl-CoA epimerase; a methylmalonyl-CoA reductase (alcohol forming); and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In a further aspect of the above embodiments, the microbial organism has an acetyl-CoA pathway having a third set of exogenous nucleic acids encoding acetyl-CoA pathway enzymes expressed in a sufficient amount to produce acetyl-CoA, the third set of exogenous nucleic acids encoding a pyruvate kinase; and a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase.

In another further embodiment, the microbial organism has a succinyl-CoA pathway having a third set of exogenous nucleic acids encoding succinyl-CoA pathway enzymes expressed in a sufficient amount to produce succinyl-CoA, the third set of exogenous nucleic acids encoding a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase and a succinyl-CoA synthetase. In a further aspect, the third set of exogenous nucleic acids further encodes a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a pyruvate carboxylase, a methylmalonyl-CoA carboxytransferase, a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde-forming), a 4-hydroxybutyraldehyde reductase, a 4-hydroxybutyrate reductase; a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-phosphate reductase, a 4-hydroxybutyryl-CoA reductase (alcohol-forming), and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, a formate dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase, and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a pyruvate carboxylase, a methylmalonyl-CoA carboxytransferase, a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA dehydratase, a crotonase, a 3-hydroxybutyryl-CoA reductase (aldehyde forming), a 3-hydroxybutyraldehyde reductase, a 3-hydroxybutyryl-CoA transferase, a 3-hydroxybutyryl-CoA synthetase, a 3-hydroxybutyryl-CoA hydrolase, a 3-hydroxybutyrate reductase, and a 3-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of

exogenous nucleic acids encoding a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, a formate dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate
5 decarboxylase, and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids
10 encoding a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a pyruvate carboxylase, a methylmalonyl-CoA carboxytransferase, a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-
15 hydroxybutyrylase, a 4-hydroxybutyryl-CoA mutase, a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase, a 3-hydroxyisobutyryl-CoA hydrolase, 3-hydroxyisobutyryl-CoA dehydratase, methacrylyl-CoA transferase, methacrylyl-CoA synthetase, methacrylyl-CoA hydrolase and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol
20 pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, a formate dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate
25 decarboxylase, and an isopropanol dehydrogenase.

In one embodiment, the invention provides a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids
30 encoding a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a pyruvate carboxylase, a methylmalonyl-CoA carboxytransferase, a methylmalonyl-CoA mutase, a methylmalonyl-CoA epimerase, a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase, a methylmalonyl-CoA hydrolase, a methylmalonate reductase, a methylmalonyl-

- CoA reductase (aldehyde forming), a 3-hydroxyisobutyrate dehydrogenase, a methylmalonyl-CoA reductase (alcohol forming) and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous
- 5 nucleic acids encoding a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, a formate dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase, and an isopropanol dehydrogenase.
- 10 In a further aspect of each of the above embodiments, the exogenous nucleic acid is a heterologous nucleic acid.

In a further aspect of each of the above embodiments, the non-naturally occurring microbial organism is in a substantially anaerobic culture medium.

- In an additional embodiment, the invention provides a non-naturally occurring microbial
- 15 organism having an n-propanol and isopropanol pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product selected from the group consisting of phosphoenolpyruvate to oxaloacetate, oxaloacetate to malate, malate to fumarate, fumarate to succinate, succinate to succinyl-CoA, succinyl-CoA to (R)-methylmalonyl-CoA, (R)-
- 20 methylmalonyl-CoA to (S)-methylmalonyl-CoA, (S)-methylmalonyl-CoA to propionyl-CoA, propionyl-CoA to propionaldehyde, propionaldehyde to n-propanol, propionyl-CoA to propionyl phosphate, propionyl-CoA to propionate, propionate to propionyl phosphate, propionate to propionaldehyde, propionyl phosphate to propionaldehyde, phosphoenolpyruvate to pyruvate, pyruvate to oxaloacetate, pyruvate to acetyl-CoA, pyruvate to acetyl-CoA and formate, formate
- 25 to CO₂, 2 acetyl-CoA substrates to 1 acetoacetyl-CoA product, acetoacetyl-CoA to acetoacetate, acetoacetate to acetone, acetone to isopropanol. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of the substrate to the product can be readily determined by one skilled in the art based on the
- 30 teachings herein. Thus, the invention provides a non-naturally occurring microbial organism containing at least one exogenous nucleic acid encoding an enzyme or protein, where the enzyme or protein converts the substrates and products of an n-propanol and isopropanol pathway, such as that shown in Figure 1.

- In an additional embodiment, the invention provides a non-naturally occurring microbial organism having an n-propanol and isopropanol pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product selected from the group consisting of
- 5 phosphoenolpyruvate to oxaloacetate, oxaloacetate to threonine, threonine to 2-oxobutanoate, 2-oxobutanoate to propionyl-CoA, propionyl-CoA to propionaldehyde, propionaldehyde to n-propanol, 2-oxobutanoate to propionaldehyde, propionyl-CoA to propionyl phosphate, propionyl-CoA to propionate, propionate to propionyl phosphate, propionate to propionaldehyde, propionyl phosphate to propionaldehyde, phosphoenolpyruvate to pyruvate,
- 10 pyruvate to oxaloacetate, pyruvate to acetyl-CoA, pyruvate to acetyl-CoA and formate, formate to CO₂, 2 acetyl-CoA substrates to 1 acetoacetyl-CoA product, acetoacetyl-CoA to acetoacetate, acetoacetate to acetone, acetone to isopropanol. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of
- 15 the substrate to the product can be readily determined by one skilled in the art based on the teachings herein. Thus, the invention provides a non-naturally occurring microbial organism containing at least one exogenous nucleic acid encoding an enzyme or protein, where the enzyme or protein converts the substrates and products of an n-propanol and isopropanol pathway, such as that shown in Figure 2.
- 20 In an additional embodiment, the invention provides a non-naturally occurring microbial organism having an n-propanol and isopropanol pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product selected from the group consisting of
- 25 phosphoenolpyruvate to pyruvate, pyruvate to acetyl-CoA, pyruvate to acetyl-CoA and formate, formate to CO₂, acetyl-CoA to malonyl-CoA, malonyl-CoA to malonate semialdehyde, malonate semialdehyde to 3-hydroxypropionate, 3-hydroxypropionate to propionyl-CoA, propionyl-CoA to propionaldehyde, propionaldehyde to n-propanol, propionyl-CoA to propionyl phosphate, propionyl-CoA to propionate, propionate to propionyl phosphate, propionate to propionaldehyde, propionyl phosphate to propionaldehyde, 2 acetyl-CoA substrates to 1
- 30 acetoacetyl-CoA product, acetoacetyl-CoA to acetoacetate, acetoacetate to acetone, acetone to isopropanol. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of the substrate to the product can be readily determined by one skilled in the art based on the teachings herein. Thus, the invention

provides a non-naturally occurring microbial organism containing at least one exogenous nucleic acid encoding an enzyme or protein, where the enzyme or protein converts the substrates and products of an n-propanol and isopropanol pathway, such as that shown in Figure 3.

In an additional embodiment, the invention provides a non-naturally occurring microbial organism having an n-propanol and isopropanol pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product selected from the group consisting of pyruvate to D-lactate, D-lactate to lactoyl-CoA, lactoyl-CoA to acryloyl-CoA, acryloyl-CoA to propionyl-CoA, propionyl-CoA to propionaldehyde, propionaldehyde to n-propanol, propionyl-CoA to propionyl phosphate, propionyl-CoA to propionate, propionate to propionyl phosphate, propionate to propionaldehyde, propionyl phosphate to propionaldehyde, pyruvate to acetyl-CoA, pyruvate to acetyl-CoA and formate, formate to CO₂, 2 acetyl-CoA substrates to 1 acetoacetyl-CoA product, acetoacetyl-CoA to acetoacetate, acetoacetate to acetone, acetone to isopropanol. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of the substrate to the product can be readily determined by one skilled in the art based on the teachings herein. Thus, the invention provides a non-naturally occurring microbial organism containing at least one exogenous nucleic acid encoding an enzyme or protein, where the enzyme or protein converts the substrates and products of an n-propanol and isopropanol pathway, such as that shown in Figure 4.

In an additional embodiment, the invention provides a non-naturally occurring microbial organism having an n-propanol pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product selected from the group consisting of propionyl-CoA to propionaldehyde, propionaldehyde to n-propanol, propionyl-CoA to propionyl phosphate, propionyl-CoA to propionate, propionate to propionyl phosphate, propionate to propionaldehyde, and propionyl phosphate to propionaldehyde. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of the substrate to the product can be readily determined by one skilled in the art based on the teachings herein. Thus, the invention provides a non-naturally occurring microbial organism containing at least one exogenous nucleic acid encoding an enzyme or protein, where

the enzyme or protein converts the substrates and products of an n-propanol pathway, such as that shown in Figures 1-4.

In an additional embodiment, the invention provides a non-naturally occurring microbial organism having an 14-BDO and an isopropanol pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product selected from the group consisting of phosphoenolpyruvate to oxaloacetate, oxaloacetate to malate, malate to fumarate, fumarate to succinate, succinate to succinyl-CoA, succinyl-CoA to succinic semialdehyde, succinic semialdehyde to 4-hydroxybutyrate, 4-hydroxybutyrate to 4-hydroxybutyryl-CoA, 4-hydroxybutyryl-CoA to 4-hydroxybutyraldehyde, 4-hydroxybutyraldehyde to 14-BDO, succinate to succinic semialdehyde, 4-hydroxybutyrate to 4-hydroxybutyraldehyde, 4-hydroxybutyrate to 4-hydroxybutyryl-phosphate, 4-hydroxybutyryl-phosphate to 4-hydroxybutyryl-CoA, 4-hydroxybutyryl-phosphate to 4-hydroxybutyraldehyde, 4-hydroxybutyryl-CoA to 14-BDO, propionyl-CoA to propionyl phosphate, propionyl phosphate to propionaldehyde, phosphoenolpyruvate to pyruvate, pyruvate to oxaloacetate, pyruvate to acetyl-CoA, pyruvate to acetyl-CoA and formate, formate to CO₂, 2 acetyl-CoA substrates to 1 acetoacetyl-CoA product, acetoacetyl-CoA to acetoacetate, acetoacetate to acetone, acetone to isopropanol. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of the substrate to the product can be readily determined by one skilled in the art based on the teachings herein. Thus, the invention provides a non-naturally occurring microbial organism containing at least one exogenous nucleic acid encoding an enzyme or protein, where the enzyme or protein converts the substrates and products of an n-propanol and isopropanol pathway, such as that shown in Figure 5.

In an additional embodiment, the invention provides a non-naturally occurring microbial organism having an 13-BDO and an isopropanol pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product selected from the group consisting of phosphoenolpyruvate to oxaloacetate, oxaloacetate to malate, malate to fumarate, fumarate to succinate, succinate to succinyl-CoA, succinyl-CoA to succinic semialdehyde, succinic semialdehyde to 4-hydroxybutyrate, 4-hydroxybutyrate to 4-hydroxybutyryl-CoA, succinate to succinic semialdehyde, 4-hydroxybutyrate to 4-hydroxybutyryl-phosphate, 4-hydroxybutyryl-phosphate to 4-hydroxybutyryl-CoA, 4-hydroxybutyryl-CoA to crotonyl-CoA, crotonyl-CoA to 3-hydroxybutyryl-CoA, 3-hydroxybutyryl-CoA to 3-hydroxybutyraldehyde, 3-hydroxybutyryl-

CoA to 3-hydroxybutyrate, 3-hydroxybutyrate to 3-hydroxybutyraldehyde, 3-hydroxybutyraldehyde to 13-BDO, 3-hydroxybutyryl-CoA to 13-BDO, propionyl-CoA to propionyl phosphate, propionyl phosphate to propionaldehyde, phosphoenolpyruvate to pyruvate, pyruvate to oxaloacetate, pyruvate to acetyl-CoA, pyruvate to acetyl-CoA and formate, formate to CO₂, 2 acetyl-CoA substrates to 1 acetoacetyl-CoA product, acetoacetyl-CoA to acetoacetate, acetoacetate to acetone, acetone to isopropanol. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of the substrate to the product can be readily determined by one skilled in the art based on the teachings herein. Thus, the invention provides a non-naturally occurring microbial organism containing at least one exogenous nucleic acid encoding an enzyme or protein, where the enzyme or protein converts the substrates and products of an n-propanol and isopropanol pathway, such as that shown in Figure 6.

In an additional embodiment, the invention provides a non-naturally occurring microbial organism having an MAA and an isopropanol pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product selected from the group consisting of phosphoenolpyruvate to oxaloacetate, oxaloacetate to malate, malate to fumarate, fumarate to succinate, succinate to succinyl-CoA, succinyl-CoA to succinic semialdehyde, succinic semialdehyde to 4-hydroxybutyrate, 4-hydroxybutyrate to 4-hydroxybutyryl-CoA, succinate to succinic semialdehyde, 4-hydroxybutyrate to 4-hydroxybutyryl-phosphate, 4-hydroxybutyryl-phosphate to 4-hydroxybutyryl-CoA, 4-hydroxybutyryl-CoA to 3-hydroxyisobutyryl-CoA, 3-hydroxyisobutyryl-CoA to 3-hydroxyisobutyrate, 3-hydroxyisobutyryl-CoA to methacrylyl-CoA, methacrylyl-CoA to MAA, 3-hydroxyisobutyrate to MAA, succinyl-CoA to (R)-methylmalonyl-CoA, (R)-methylmalonyl-CoA to (S)-methylmalonyl-CoA, (S)-methylmalonyl-CoA to methylmalonate semialdehyde, (S)-methylmalonyl-CoA to 3-hydroxyisobutyrate, methylmalonate semialdehyde to 3-hydroxyisobutyrate, propionyl-CoA to propionyl phosphate, propionyl phosphate to propionaldehyde, phosphoenolpyruvate to pyruvate, pyruvate to oxaloacetate, pyruvate to acetyl-CoA, pyruvate to acetyl-CoA and formate, formate to CO₂, 2 acetyl-CoA substrates to 1 acetoacetyl-CoA product, acetoacetyl-CoA to acetoacetate, acetoacetate to acetone, acetone to isopropanol. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of the substrate to the product can be readily determined by one skilled in the art based on the

teachings herein. Thus, the invention provides a non-naturally occurring microbial organism containing at least one exogenous nucleic acid encoding an enzyme or protein, where the enzyme or protein converts the substrates and products of an n-propanol and isopropanol pathway, such as that shown in Figure 7 and 8.

- 5 While generally described herein as a microbial organism that contains an n-propanol and an isopropanol, a 14-BDO and an isopropanol, a 13-BDO and an isopropanol or a MAA and an isopropanol pathway, it is understood that the invention additionally provides a non-naturally occurring microbial organism comprising at least one exogenous nucleic acid encoding an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or MAA pathway enzyme expressed in a
- 10 sufficient amount to produce an intermediate of an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or MAA pathway. For example, as disclosed herein, an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or MAA pathway is exemplified in Figures 1-8. Therefore, in addition to a microbial organism containing an n-propanol and an isopropanol, a 14-BDO and an isopropanol, a 13-BDO and an isopropanol or a MAA and an isopropanol pathway that produces
- 15 n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol, the invention additionally provides a non-naturally occurring microbial organism comprising at least one exogenous nucleic acid encoding an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or MAA pathway enzyme, where the microbial organism produces an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or MAA pathway intermediate, for
- 20 example, acetone, methylmalonyl-CoA, propionyl phosphate, 2-oxobutanoate, 3-hydroxypropionate, lactoyl-CoA, 4-hydroxybutyrate, 4-hydroxybutyryl-phosphate, crotonyl-CoA, succinyl-CoA, succinic semialdehyde or 3-hydroxyisobutyryl-CoA.

- It is understood that any of the pathways disclosed herein, as described in the Examples and exemplified in the Figures, including the pathways of Figures 1-8, can be utilized to generate a
- 25 non-naturally occurring microbial organism that produces any pathway intermediate or product, as desired. As disclosed herein, such a microbial organism that produces an intermediate can be used in combination with another microbial organism expressing downstream pathway enzymes to produce a desired product. However, it is understood that a non-naturally occurring microbial organism that produces an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or MAA
- 30 intermediate can be utilized to produce the intermediate as a desired product.

The invention is described herein with general reference to the metabolic reaction, reactant or product thereof, or with specific reference to one or more nucleic acids or genes encoding an enzyme associated with or catalyzing, or a protein associated with, the referenced metabolic

reaction, reactant or product. Unless otherwise expressly stated herein, those skilled in the art will understand that reference to a reaction also constitutes reference to the reactants and products of the reaction. Similarly, unless otherwise expressly stated herein, reference to a reactant or product also references the reaction, and reference to any of these metabolic constituents also references the gene or genes encoding the enzymes that catalyze or proteins involved in the referenced reaction, reactant or product. Likewise, given the well known fields of metabolic biochemistry, enzymology and genomics, reference herein to a gene or encoding nucleic acid also constitutes a reference to the corresponding encoded enzyme and the reaction it catalyzes or a protein associated with the reaction as well as the reactants and products of the reaction.

The non-naturally occurring microbial organisms of the invention can be produced by introducing expressible nucleic acids encoding one or more of the enzymes or proteins participating in one or more n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthetic pathways. Depending on the host microbial organism chosen for biosynthesis, nucleic acids for some or all of a particular n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthetic pathway can be expressed. For example, if a chosen host is deficient in one or more enzymes or proteins for a desired biosynthetic pathway, then expressible nucleic acids for the deficient enzyme(s) or protein(s) are introduced into the host for subsequent exogenous expression. Alternatively, if the chosen host exhibits endogenous expression of some pathway genes, but is deficient in others, then an encoding nucleic acid is needed for the deficient enzyme(s) or protein(s) to achieve n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthesis. Thus, a non-naturally occurring microbial organism of the invention can be produced by introducing exogenous enzyme or protein activities to obtain a desired biosynthetic pathway or a desired biosynthetic pathway can be obtained by introducing one or more exogenous enzyme or protein activities that, together with one or more endogenous enzymes or proteins, produces a desired product such as n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA.

Depending on the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthetic pathway constituents of a selected host microbial organism, the non-naturally occurring microbial organisms of the invention will include at least one exogenously expressed n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway-encoding nucleic acid and up to all encoding nucleic acids for one or more n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthetic pathways. For example, n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthesis can be established in a host deficient in a pathway enzyme or protein through

exogenous expression of the corresponding encoding nucleic acid. In a host deficient in all enzymes or proteins of an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway, exogenous expression of all enzyme or proteins in the pathway can be included, although it is understood that all enzymes or proteins of a pathway can be expressed even if the host contains at least one of the pathway enzymes or proteins. For example, exogenous expression of all enzymes or proteins in a pathway for production of n-propanol and isopropanol can be included, such as a PEP carboxykinase or a PEP carboxylase; a malate dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA synthetase; a methylmalonyl-CoA mutase; a methylmalonyl-CoA epimerase; a methylmalonyl-CoA decarboxylase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase and a propionyl phosphate reductase, a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase, as exemplified in Figure 1.

Given the teachings and guidance provided herein, those skilled in the art will understand that the number of encoding nucleic acids to introduce in an expressible form will, at least, parallel the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway deficiencies of the selected host microbial organism. Therefore, a non-naturally occurring microbial organism of the invention can have one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty or twenty one, up to all nucleic acids encoding the enzymes or proteins constituting an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA biosynthetic pathway disclosed herein. In some embodiments, the non-naturally occurring microbial organisms also can include other genetic modifications that facilitate or optimize n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthesis or that confer other useful functions onto the host microbial organism. One such other functionality can include, for example, augmentation of the synthesis of one or more of the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway precursors such as phosphoenolpyruvate or pyruvate.

Generally, a host microbial organism is selected such that it produces the precursor of an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway, either as a naturally produced molecule or as an engineered product that either provides *de novo* production of a desired precursor or increased production of a precursor naturally produced by the host

microbial organism. For example, phosphoenolpyruvate and pyruvate are produced naturally in a host organism such as *E. coli*. A host organism can be engineered to increase production of a precursor, as disclosed herein. In addition, a microbial organism that has been engineered to produce a desired precursor can be used as a host organism and further engineered to express enzymes or proteins of an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway.

In some embodiments, a non-naturally occurring microbial organism of the invention is generated from a host that contains the enzymatic capability to synthesize n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA. In this specific embodiment it can be useful to increase the synthesis or accumulation of an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway product to, for example, drive n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway reactions toward n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA production. Increased synthesis or accumulation can be accomplished by, for example, overexpression of nucleic acids encoding one or more of the above-described n-propanol and/or isopropanol pathway enzymes or proteins. Over expression of the enzyme or enzymes and/or protein or proteins of the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway can occur, for example, through exogenous expression of the endogenous gene or genes, or through exogenous expression of the heterologous gene or genes. Therefore, naturally occurring organisms can be readily generated to be non-naturally occurring microbial organisms of the invention, for example, producing n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA, through overexpression of one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty or twenty one, that is, up to all nucleic acids encoding n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthetic pathway enzymes or proteins. In addition, a non-naturally occurring organism can be generated by mutagenesis of an endogenous gene that results in an increase in activity of an enzyme in the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthetic pathway.

In particularly useful embodiments, exogenous expression of the encoding nucleic acids is employed. Exogenous expression confers the ability to custom tailor the expression and/or regulatory elements to the host and application to achieve a desired expression level that is controlled by the user. However, endogenous expression also can be utilized in other embodiments such as by removing a negative regulatory effector or induction of the gene's promoter when linked to an inducible promoter or other regulatory element. Thus, an endogenous gene having a naturally occurring inducible promoter can be up-regulated by providing the appropriate inducing agent, or the regulatory region of an endogenous gene can be

engineered to incorporate an inducible regulatory element, thereby allowing the regulation of increased expression of an endogenous gene at a desired time. Similarly, an inducible promoter can be included as a regulatory element for an exogenous gene introduced into a non-naturally occurring microbial organism.

- 5 It is understood that, in methods of the invention, any of the one or more exogenous nucleic acids can be introduced into a microbial organism to produce a non-naturally occurring microbial organism of the invention. The nucleic acids can be introduced so as to confer, for example, an n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol biosynthetic pathway onto the microbial organism. Alternatively,
- 10 encoding nucleic acids can be introduced to produce an intermediate microbial organism having the biosynthetic capability to catalyze some of the required reactions to confer n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthetic capability. For example, a non-naturally occurring microbial organism having an n-propanol and an isopropanol, a 14-BDO and an isopropanol, a 13-BDO and an isopropanol or a MAA and an isopropanol biosynthetic
- 15 pathway can comprise at least two exogenous nucleic acids encoding desired enzymes or proteins, such as the combination of propionaldehyde dehydrogenase and isopropanol dehydrogenase, or alternatively propionyl-CoA synthase and acetyl-CoA acetyl thiolase, or alternatively lactate dehydrogenase and acetyl-CoA thiolase, or alternatively a succinyl-CoA reductase and 4-hydroxybutyryl-CoA reductase (alcohol-forming), or alternatively crotonase and
- 20 acetoacetate decarboxylase, or alternatively 4-hydroxybutyrate kinase and phosphotrans-4-hydroxybutyrylase or alternatively methylmalonyl-CoA reductase (alcohol forming) and pyruvate kinase and the like. Thus, it is understood that any combination of two or more enzymes or proteins of a biosynthetic pathway can be included in a non-naturally occurring microbial organism of the invention. Similarly, it is understood that any combination of three or
- 25 more enzymes or proteins of a biosynthetic pathway can be included in a non-naturally occurring microbial organism of the invention, for example, PEP carboxykinase, acetyl-CoA acetyl thiolase and propanol dehydrogenase, or alternatively pyruvate kinase, acetoacetate decarboxylase and 2-oxobutanoate dehydrogenase, or alternatively propionyl-CoA:phosphate propanoyltransferase, propionyl phosphate reductase and isopropanol dehydrogenase, or
- 30 alternatively lactate-CoA transferase and lactyl-CoA dehydratase and pyruvate formate lyase, or alternatively succinyl-CoA dehydrogenase, 4-hydroxybutyrate reductase and 4-hydroxybutyraldehyde reductase, or alternatively crotonase, PEP carboxylase and acetoacetate decarboxylase, or alternatively 3-hydroxyisobutyryl-CoA synthetase, fumarase and isopropanol dehydrogenase, or alternatively acetyl-CoA acetyl thiolase, acetoacetate decarboxylase and

5 methylmalonyl-CoA reductase (alcohol forming) and so forth, as desired, so long as the combination of enzymes and/or proteins of the desired biosynthetic pathway results in production of the corresponding desired product. Similarly, any combination of four or more enzymes or proteins of a biosynthetic pathway as disclosed herein, for example, pyruvate
10 carboxylase, malate dehydrogenase, methylmalonyl-CoA epimerase and acetoacetyl-CoA hydrolase, or alternatively acetyl-CoA acetyl thiolase, isopropanol dehydrogenase, propionaldehyde dehydrogenase and propanol dehydrogenase, or alternatively acetyl-CoA carboxylase, malonyl-CoA reductase, malonate semialdehyde and acetoacetate decarboxylase, or alternatively, acryloyl CoA reductase, acetoacetyl-CoA transferase, acetoacetate
15 decarboxylase, and isopropanol dehydrogenase, or alternatively succinyl-CoA dehydrogenase, 4-hydroxybutyrate dehydrogenase, 4-hydroxybutyryl-CoA transferase, and isopropanol dehydrogenase, or alternatively succinate reductase, 3-hydroxyisobutyryl-CoA synthetase, 3-hydroxyisobutyrate dehydratase and pyruvate ferredoxin oxidoreductase, or alternatively acetyl-CoA acetyl thiolase, acetoacetyl-CoA transferase, methylmalonyl-CoA mutase and
20 hydroxyisobutyrate dehydratase, can be included in a non-naturally occurring microbial organism of the invention, as desired, so long as the combination of enzymes and/or proteins of the desired biosynthetic pathway results in production of the corresponding desired product.

It is understood that when more than one exogenous nucleic acid is included in a microbial organism that the more than one exogenous nucleic acids refers to the referenced encoding
25 nucleic acid or biosynthetic activity, as discussed above. It is further understood, as disclosed herein, that more than one exogenous nucleic acids can be introduced into the host microbial organism on separate nucleic acid molecules, on polycistronic nucleic acid molecules, or a combination thereof, and still be considered as more than one exogenous nucleic acid. For example, as disclosed herein a microbial organism can be engineered to express two or more
30 exogenous nucleic acids encoding a desired pathway enzyme or protein. In the case where two exogenous nucleic acids encoding a desired activity are introduced into a host microbial organism, it is understood that the two exogenous nucleic acids can be introduced as a single nucleic acid, for example, on a single plasmid, on separate plasmids, can be integrated into the host chromosome at a single site or multiple sites, and still be considered as two exogenous
35 nucleic acids. Similarly, it is understood that more than two exogenous nucleic acids can be introduced into a host organism in any desired combination, for example, on a single plasmid, on separate plasmids, can be integrated into the host chromosome at a single site or multiple sites, and still be considered as two or more exogenous nucleic acids, for example three exogenous nucleic acids. Thus, the number of referenced exogenous nucleic acids or

biosynthetic activities refers to the number of encoding nucleic acids or the number of biosynthetic activities, not the number of separate nucleic acids introduced into the host organism.

In addition to the biosynthesis of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA as described herein, the non-naturally occurring microbial organisms and methods of the invention also can be utilized in various combinations with each other and with other microbial organisms and methods well known in the art to achieve product biosynthesis by other routes. For example, one alternative to produce n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA other than use of the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA producers is through addition of another microbial organism capable of converting an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway intermediate to n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA. One such procedure includes, for example, the fermentation of a microbial organism that produces an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway intermediate. The n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol pathway intermediate can then be used as a substrate for a second microbial organism that converts the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway intermediate to n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA. The n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway intermediate can be added directly to another culture of the second organism or the original culture of the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway intermediate producers can be depleted of these microbial organisms by, for example, cell separation, and then subsequent addition of the second organism to the fermentation broth can be utilized to produce the final product without intermediate purification steps.

In other embodiments, the non-naturally occurring microbial organisms and methods of the invention can be assembled in a wide variety of subpathways to achieve biosynthesis of, for example, n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol. In these embodiments, biosynthetic pathways for a desired product of the invention can be segregated into different microbial organisms, and the different microbial organisms can be co-cultured to produce the final product. In such a biosynthetic scheme, the product of one microbial organism is the substrate for a second microbial organism until the final product is synthesized. For example, the biosynthesis of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA can be accomplished by constructing a microbial organism that contains biosynthetic pathways for conversion of one pathway intermediate to another pathway intermediate or the product. Alternatively, n-propanol, isopropanol, 14-BDO, 13-BDO and/or

MAA also can be biosynthetically produced from microbial organisms through co-culture or co-fermentation using two organisms in the same vessel, where the first microbial organism produces a propionyl-CoA, succinyl-CoA and/or an acetyl-CoA intermediate and the second microbial organism converts the intermediate(s) to n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA.

Given the teachings and guidance provided herein, those skilled in the art will understand that a wide variety of combinations and permutations exist for the non-naturally occurring microbial organisms and methods of the invention together with other microbial organisms, with the co-culture of other non-naturally occurring microbial organisms having subpathways and with combinations of other chemical and/or biochemical procedures well known in the art to produce n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA.

Sources of encoding nucleic acids for an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway enzyme or protein can include, for example, any species where the encoded gene product is capable of catalyzing the referenced reaction. Such species include both prokaryotic and eukaryotic organisms including, but not limited to, bacteria, including archaea and eubacteria, and eukaryotes, including yeast, plant, insect, animal, and mammal, including human. Exemplary species for such sources include, for example, *Escherichia coli*, *Acetobacter pasteurians*, *Acidans brierleyi*, *Acinetobacter baylyi*, *Acinetobacter calcoaceticus*, *Acinetobacter* sp. Strain M-1, *Actinobacillus succinogenes*, *Anaerobiospirillum succiniciproducens*, *Anaerostipes caccae* DSM 14662, *Arabidopsis thaliana*, *Bacillus cereus* ATCC 14579, *Bacillus subtilis*, *Bacillus subtilis* subsp. *subtilis* str. 168, *Bos taurus*, *Bradyrhizobium japonicum* USDA110, *Caenorhabditis elegans*, *Campylobacter jejuni*, *Chlamydomonas reinhardtii*, *Chloroflexus aurantiacus*, *Clostridium acetobutylicum*, *Clostridium acetobutylicum* ATCC 824, *Clostridium beijerinckii*, *Clostridium botulinum* C str. Eklund, *Clostridium kluyveri*, *Clostridium kluyveri* DSM 555, *Clostridium novyi*-NT, *Clostridium propionicum*, *Clostridium saccharobutylicum*, *Clostridium saccharoperbutylacetonicum*, *Corynebacterium glutamicum*, *Desulfovibrio africanus*, *Erythrobacter* sp. NAP1, *Escherichia coli* K12, *Escherichia coli* K12 str. MG1655, *Escherichia coli* O157:H7, *Geobacillus thermoglucosidarius* M10EXG, *Haemophilus influenza*, *Helicobacter pylori*, *Homo sapiens*, *Klebsiella pneumonia* MGH78578, *Kluyveromyces lactis*, *Lactobacillus casei*, *Lactobacillus plantarum* WCFS1, *Lactococcus lactis*, *Leuconostoc mesenteroides*, *Mannheimia succiniciproducens*, marine gamma proteobacterium HTCC2080, *Mesorhizobium loti*, *Metallosphaera sedula*, *Methylobacterium extorquens*, *Moorella thermoacetica*, *Mycobacterium smegmatis*, *Mycobacterium tuberculosis*, *Oryctolagus cuniculus*,

Plasmodium ovale, *Porphyromonas gingivalis*, *Propionibacterium acnes*, *Propionibacterium fredenreichii* sp. *shermanii*, *Propionibacterium freudenreichii*, *Propionigenium modestum*,
Pseudomonas aeruginosa, *Pseudomonas aeruginosa* PAO1, *Pseudomonas fluorescens*,
Pseudomonas putida, *Pseudomonas putida* E23, *Pseudomonas putida* KT2440, *Pseudomonas*
5 *sp*, *Pseudomonas stutzeri*, *Ralstonia eutropha*, *Ralstonia eutropha* H16, *Rattus norvegicus*,
Rhodobacter spaeroides, *Rhodoferrax ferrireducens* DSM 15236, *Rhodospirillum rubrum*,
Roseiflexus castenholzii, *Saccharomyces cerevisiae*, *Salmonella enterica*, *Salmonella*
typhimurium, *Shigella flexneri*, *Simmondsia chinensis*, *Streptococcus mutans*, *Sulfolobus*
acidocaldarius, *Sulfolobus solfataricus*, *Sulfolobus tokodaii*, *Syntrophobacter fumaroxidans*,
10 *Thermococcus litoralis*, *Thermotoga maritime*, *Thermus thermophilus*, *Trichomonas vaginalis*
G3, *Trypanosoma brucei*, *Veillonella parvula*, *Yersinia frederiksenii*, *Zymomonas mobilis*,
Bacillus megaterium, *butyrate-producing bacterium* L2-50, *Clostridium aminobutyricum*,
Geobacillus thermoglucosidasius, *Mycobacterium bovis* BCG, *Nocardia farcinica* IFM 10152,
Nocardia iowensis (sp. NRRL 5646), *Penicillium chrysogenum*, *Porphyromonas gingivalis*
15 *ATCC 33277*, *Pseudomonas mendocina*, *Streptomyces griseus subsp. griseus* NBRC 13350 as
well as other exemplary species disclosed herein are available as source organisms for
corresponding genes. However, with the complete genome sequence available for now more
than 550 species (with more than half of these available on public databases such as the NCBI),
including 395 microorganism genomes and a variety of yeast, fungi, plant, and mammalian
20 genomes, the identification of genes encoding the requisite n-propanol, isopropanol, 14-BDO,
13-BDO and/or MAA biosynthetic activity for one or more genes in related or distant species,
including for example, homologues, orthologs, paralogs and nonorthologous gene displacements
of known genes, and the interchange of genetic alterations between organisms is routine and
well known in the art. Accordingly, the metabolic alterations allowing biosynthesis of n-
25 propanol, isopropanol, 14-BDO, 13-BDO and/or MAA described herein with reference to a
particular organism such as *E. coli* can be readily applied to other microorganisms, including
prokaryotic and eukaryotic organisms alike. Given the teachings and guidance provided herein,
those skilled in the art will know that a metabolic alteration exemplified in one organism can be
applied equally to other organisms.

30 In some instances, such as when an alternative n-propanol, isopropanol, 14-BDO, 13-BDO
and/or MAA biosynthetic pathway exists in an unrelated species, n-propanol, isopropanol, 14-
BDO, 13-BDO and/or MAA biosynthesis can be conferred onto the host species by, for
example, exogenous expression of a paralog or paralogs from the unrelated species that
catalyzes a similar, yet non-identical metabolic reaction to replace the referenced reaction.

Because certain differences among metabolic networks exist between different organisms, those skilled in the art will understand that the actual gene usage between different organisms may differ. However, given the teachings and guidance provided herein, those skilled in the art also will understand that the teachings and methods of the invention can be applied to all microbial organisms using the cognate metabolic alterations to those exemplified herein to construct a microbial organism in a species of interest that will synthesize n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA.

Host microbial organisms can be selected from, and the non-naturally occurring microbial organisms generated in, for example, bacteria, yeast, fungus or any of a variety of other microorganisms applicable to fermentation processes. Exemplary bacteria include species selected from *Escherichia coli*, *Klebsiella oxytoca*, *Anaerobiospirillum succiniciproducens*, *Actinobacillus succinogenes*, *Mannheimia succiniciproducens*, *Rhizobium etli*, *Bacillus subtilis*, *Corynebacterium glutamicum*, *Gluconobacter oxydans*, *Zymomonas mobilis*, *Lactococcus lactis*, *Lactobacillus plantarum*, *Streptomyces coelicolor*, *Clostridium acetobutylicum*, *Pseudomonas fluorescens*, and *Pseudomonas putida*. Exemplary yeasts or fungi include species selected from *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces marxianus*, *Aspergillus terreus*, *Aspergillus niger* and *Pichia pastoris*. *E. coli* is a particularly useful host organism since it is a well characterized microbial organism suitable for genetic engineering. Other particularly useful host organisms include yeast such as *Saccharomyces cerevisiae*. Other particularly useful host organisms include microbial organisms which naturally produce sufficient quantities of propionyl-CoA and/or acetyl-CoA for co-production of n-propanol and isopropanol. Examples of such organisms include, but are not limited to, *Clostridium propionicum*, *Escherichia coli* and *Propionibacterium freudenreichii* subsp. *shermanii*.

Methods for constructing and testing the expression levels of a non-naturally occurring n-propanol-, isopropanol-, 14-BDO-, 13-BDO- and/or MAA-producing host can be performed, for example, by recombinant and detection methods well known in the art. Such methods can be found described in, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Third Ed., Cold Spring Harbor Laboratory, New York (2001); and Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, MD (1999).

Exogenous nucleic acid sequences involved in a pathway for production of n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol can be introduced stably or transiently into a host cell using techniques well known in the art

including, but not limited to, conjugation, electroporation, chemical transformation, transduction, transfection, and ultrasound transformation. For exogenous expression in *E. coli* or other prokaryotic cells, some nucleic acid sequences in the genes or cDNAs of eukaryotic nucleic acids can encode targeting signals such as an N-terminal mitochondrial or other targeting signal, which can be removed before transformation into prokaryotic host cells, if desired. For example, removal of a mitochondrial leader sequence led to increased expression in *E. coli* (Hoffmeister et al., *J. Biol. Chem.* 280:4329-4338 (2005)). For exogenous expression in yeast or other eukaryotic cells, genes can be expressed in the cytosol without the addition of leader sequence, or can be targeted to mitochondrion or other organelles, or targeted for secretion, by the addition of a suitable targeting sequence such as a mitochondrial targeting or secretion signal suitable for the host cells. Thus, it is understood that appropriate modifications to a nucleic acid sequence to remove or include a targeting sequence can be incorporated into an exogenous nucleic acid sequence to impart desirable properties. Furthermore, genes can be subjected to codon optimization with techniques well known in the art to achieve optimized expression of the proteins.

An expression vector or vectors can be constructed to include one or more n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA biosynthetic pathway encoding nucleic acids as exemplified herein operably linked to expression control sequences functional in the host organism. Expression vectors applicable for use in the microbial host organisms of the invention include, for example, plasmids, phage vectors, viral vectors, episomes and artificial chromosomes, including vectors and selection sequences or markers operable for stable integration into a host chromosome. Additionally, the expression vectors can include one or more selectable marker genes and appropriate expression control sequences. Selectable marker genes also can be included that, for example, provide resistance to antibiotics or toxins, complement auxotrophic deficiencies, or supply critical nutrients not in the culture media. Expression control sequences can include constitutive and inducible promoters, transcription enhancers, transcription terminators, and the like which are well known in the art. When two or more exogenous encoding nucleic acids are to be co-expressed, both nucleic acids can be inserted, for example, into a single expression vector or in separate expression vectors. For single vector expression, the encoding nucleic acids can be operationally linked to one common expression control sequence or linked to different expression control sequences, such as one inducible promoter and one constitutive promoter. The transformation of exogenous nucleic acid sequences involved in a metabolic or synthetic pathway can be confirmed using methods well known in the art. Such methods include, for example, nucleic acid analysis such as

Northern blots or polymerase chain reaction (PCR) amplification of mRNA, or immunoblotting for expression of gene products, or other suitable analytical methods to test the expression of an introduced nucleic acid sequence or its corresponding gene product. It is understood by those skilled in the art that the exogenous nucleic acid is expressed in a sufficient amount to produce the desired product, and it is further understood that expression levels can be optimized to obtain sufficient expression using methods well known in the art and as disclosed herein.

Directed evolution is a powerful approach that involves the introduction of mutations targeted to a specific gene in order to improve and/or alter the properties of an enzyme. Improved and/or altered enzymes can be identified through the development and implementation of sensitive high-throughput screening assays that allow the automated screening of many enzyme variants (e.g., $>10^4$). Iterative rounds of mutagenesis and screening typically are performed to afford an enzyme with optimized properties. Computational algorithms that can help to identify areas of the gene for mutagenesis also have been developed and can significantly reduce the number of enzyme variants that need to be generated and screened.

Numerous directed evolution technologies have been developed (for reviews, see Hibbert et al., *Biomol. Eng* 22:11-19 (2005); Huisman and Lalonde, *In Biocatalysis in the pharmaceutical and biotechnology industries* pgs. 717-742 (2007), Patel (ed.), CRC Press; Otten and Quax. *Biomol.Eng* 22:1-9 (2005).; and Sen et al., *Appl Biochem. Biotechnol* 143:212-223 (2007)) to be effective at creating diverse variant libraries and these methods have been successfully applied to the improvement of a wide range of properties across many enzyme classes.

Enzyme characteristics that have been improved and/or altered by directed evolution technologies include, for example, selectivity/specificity – for conversion of non-natural substrates; temperature stability – for robust high temperature processing; pH stability – for bioprocessing under lower or higher pH conditions; substrate or product tolerance – so that high product titers can be achieved; binding (K_m) – broadens substrate binding to include non-natural substrates; inhibition (K_i) – to remove inhibition by products, substrates, or key intermediates; activity (k_{cat}) – increases enzymatic reaction rates to achieve desired flux; expression levels – increases protein yields and overall pathway flux; oxygen stability – for operation of air sensitive enzymes under aerobic conditions; and anaerobic activity – for operation of an aerobic enzyme in the absence of oxygen.

The following exemplary methods have been developed for the mutagenesis and diversification of genes to target desired properties of specific enzymes. Any of these can be used to alter/optimize activity of a decarboxylase enzyme.

EpPCR (Pritchard et al., *J Theor.Biol* 234:497-509 (2005)) introduces random point mutations by reducing the fidelity of DNA polymerase in PCR reactions by the addition of Mn^{2+} ions, by biasing dNTP concentrations, or by other conditional variations. The five step cloning process to confine the mutagenesis to the target gene of interest involves: 1) error-prone PCR
5 amplification of the gene of interest; 2) restriction enzyme digestion; 3) gel purification of the desired DNA fragment; 4) ligation into a vector; 5) transformation of the gene variants into a suitable host and screening of the library for improved performance. This method can generate multiple mutations in a single gene simultaneously, which can be useful. A high number of mutants can be generated by EpPCR, so a high-throughput screening assay or a selection method
10 (especially using robotics) is useful to identify those with desirable characteristics.

Error-prone Rolling Circle Amplification (epRCA) (Fujii et al., *Nucleic Acids Res* 32:e145 (2004); and Fujii et al., *Nat.Protoc.* 1:2493-2497 (2006)) has many of the same elements as epPCR except a whole circular plasmid is used as the template and random 6-mers with
15 exonuclease resistant thiophosphate linkages on the last 2 nucleotides are used to amplify the plasmid followed by transformation into cells in which the plasmid is re-circularized at tandem repeats. Adjusting the Mn^{2+} concentration can vary the mutation rate somewhat. This technique uses a simple error-prone, single-step method to create a full copy of the plasmid with 3 - 4 mutations/kbp. No restriction enzyme digestion or specific primers are required. Additionally,
this method is typically available as a kit.

20 DNA or Family Shuffling (Stemmer, *Proc Natl Acad Sci U.S.A.* 91:10747-10751 (1994); and Stemmer, *Nature* 370:389-391 (1994)) typically involves digestion of two or more variant genes with nucleases such as Dnase I or EndoV to generate a pool of random fragments that are reassembled by cycles of annealing and extension in the presence of DNA polymerase to create a library of chimeric genes. Fragments prime each other and recombination occurs when one
25 copy primes another copy (template switch). This method can be used with >1kbp DNA sequences. In addition to mutational recombinants created by fragment reassembly, this method introduces point mutations in the extension steps at a rate similar to error-prone PCR. The method can be used to remove deleterious, random and neutral mutations that might confer antigenicity.

30 Staggered Extension (StEP) (Zhao et al., *Nat.Biotechnol* 16:258-261 (1998)) entails template priming followed by repeated cycles of 2 step PCR with denaturation and very short duration of annealing/extension (as short as 5 sec). Growing fragments anneal to different templates and extend further, which is repeated until full-length sequences are made. Template switching

means most resulting fragments have multiple parents. Combinations of low-fidelity polymerases (Taq and Mutazyme) reduce error-prone biases because of opposite mutational spectra.

In Random Priming Recombination (RPR) random sequence primers are used to generate many short DNA fragments complementary to different segments of the template. (Shao et al., *Nucleic Acids Res* 26:681-683 (1998)) Base misincorporation and mispriming via epPCR give point mutations. Short DNA fragments prime one another based on homology and are recombined and reassembled into full-length by repeated thermocycling. Removal of templates prior to this step assures low parental recombinants. This method, like most others, can be performed over multiple iterations to evolve distinct properties. This technology avoids sequence bias, is independent of gene length, and requires very little parent DNA for the application.

In Heteroduplex Recombination linearized plasmid DNA is used to form heteroduplexes that are repaired by mismatch repair. (Volkov et al, *Nucleic Acids Res* 27:e18 (1999); and Volkov et al., *Methods Enzymol.* 328:456-463 (2000)) The mismatch repair step is at least somewhat mutagenic. Heteroduplexes transform more efficiently than linear homoduplexes. This method is suitable for large genes and whole operons.

Random Chimeragenesis on Transient Templates (RACHITT) (Coco et al., *Nat.Biotechnol* 19:354-359 (2001)) employs Dnase I fragmentation and size fractionation of ssDNA. Homologous fragments are hybridized in the absence of polymerase to a complementary ssDNA scaffold. Any overlapping unhybridized fragment ends are trimmed down by an exonuclease. Gaps between fragments are filled in, and then ligated to give a pool of full-length diverse strands hybridized to the scaffold (that contains U to preclude amplification). The scaffold then is destroyed and is replaced by a new strand complementary to the diverse strand by PCR amplification. The method involves one strand (scaffold) that is from only one parent while the priming fragments derive from other genes; the parent scaffold is selected against. Thus, no reannealing with parental fragments occurs. Overlapping fragments are trimmed with an exonuclease. Otherwise, this is conceptually similar to DNA shuffling and StEP. Therefore, there should be no siblings, few inactives, and no unshuffled parentals. This technique has advantages in that few or no parental genes are created and many more crossovers can result relative to standard DNA shuffling.

Recombined Extension on Truncated templates (RETT) entails template switching of unidirectionally growing strands from primers in the presence of unidirectional ssDNA

fragments used as a pool of templates. (Lee et al., *J.Molec.Catalysis* 26:119-129 (2003)) No DNA endonucleases are used. Unidirectional ssDNA is made by DNA polymerase with random primers or serial deletion with exonuclease. Unidirectional ssDNA are only templates and not primers. Random priming and exonucleases don't introduce sequence bias as true of enzymatic cleavage of DNA shuffling/RACHITT. RETT can be easier to optimize than StEP because it uses normal PCR conditions instead of very short extensions. Recombination occurs as a component of the PCR steps--no direct shuffling. This method can also be more random than StEP due to the absence of pauses.

In Degenerate Oligonucleotide Gene Shuffling (DOGS) degenerate primers are used to control recombination between molecules; (Bergquist and Gibbs, *Methods Mol.Biol* 352:191-204 (2007); Bergquist et al., *Biomol.Eng* 22:63-72 (2005); Gibbs et al., *Gene* 271:13-20 (2001)) this can be used to control the tendency of other methods such as DNA shuffling to regenerate parental genes. This method can be combined with random mutagenesis (epPCR) of selected gene segments. This can be a good method to block the reformation of parental sequences. No endonucleases are needed. By adjusting input concentrations of segments made, one can bias towards a desired backbone. This method allows DNA shuffling from unrelated parents without restriction enzyme digests and allows a choice of random mutagenesis methods.

Incremental Truncation for the Creation of Hybrid Enzymes (ITCHY) creates a combinatorial library with 1 base pair deletions of a gene or gene fragment of interest. (Ostermeier et al., *Proc Natl Acad Sci U.S.A.* 96:3562-3567 (1999); and Ostermeier et al., *Nat.Biotechnol* 17:1205-1209 (1999)) Truncations are introduced in opposite direction on pieces of 2 different genes. These are ligated together and the fusions are cloned. This technique does not require homology between the 2 parental genes. When ITCHY is combined with DNA shuffling, the system is called SCRATCHY (see below). A major advantage of both is no need for homology between parental genes; for example, functional fusions between an *E. coli* and a human gene were created via ITCHY. When ITCHY libraries are made, all possible crossovers are captured.

Thio-Incremental Truncation for the Creation of Hybrid Enzymes (THIO-ITCHY) is similar to ITCHY except that phosphothioate dNTPs are used to generate truncations. (Lutz et al., *Nucleic Acids Res* 29:E16 (2001)) Relative to ITCHY, THIO-ITCHY can be easier to optimize, provide more reproducibility, and adjustability.

SCRATCHY combines two methods for recombining genes, ITCHY and DNA shuffling. (Lutz et al., *Proc Natl Acad Sci U.S.A.* 98:11248-11253 (2001)) SCRATCHY combines the best features of ITCHY and DNA shuffling. First, ITCHY is used to create a comprehensive set of

fusions between fragments of genes in a DNA homology-independent fashion. This artificial family is then subjected to a DNA-shuffling step to augment the number of crossovers.

Computational predictions can be used in optimization. SCRATCHY is more effective than DNA shuffling when sequence identity is below 80%.

- 5 In Random Drift Mutagenesis (RNDM) mutations made via epPCR followed by screening/selection for those retaining usable activity. (Bergquist et al., *Biomol.Eng* 22:63-72 (2005)) Then, these are used in DOGS to generate recombinants with fusions between multiple active mutants or between active mutants and some other desirable parent. Designed to promote isolation of neutral mutations; its purpose is to screen for retained catalytic activity whether or
- 10 not this activity is higher or lower than in the original gene. RNDM is usable in high throughput assays when screening is capable of detecting activity above background. RNDM has been used as a front end to DOGS in generating diversity. The technique imposes a requirement for activity prior to shuffling or other subsequent steps; neutral drift libraries are indicated to result in higher/quicker improvements in activity from smaller libraries. Though published using
- 15 epPCR, this could be applied to other large-scale mutagenesis methods.

- Sequence Saturation Mutagenesis (SeSaM) is a random mutagenesis method that: 1) generates pool of random length fragments using random incorporation of a phosphothioate nucleotide and cleavage; this pool is used as a template to 2) extend in the presence of “universal” bases such as inosine; 3) replication of a inosine-containing complement gives random base incorporation and,
- 20 consequently, mutagenesis. (Wong et al., *Biotechnol J* 3:74-82 (2008); Wong et al., *Nucleic Acids Res* 32:e26 (2004); and Wong et al., *Anal.Biochem.* 341:187-189 (2005)) Using this technique it can be possible to generate a large library of mutants within 2 –3 days using simple methods. This technique is non-directed in comparison to the mutational bias of DNA polymerases. Differences in this approach make this technique complementary (or an
- 25 alternative) to epPCR.

- In Synthetic Shuffling, overlapping oligonucleotides are designed to encode “all genetic diversity in targets” and allow a very high diversity for the shuffled progeny. (Ness et al., *Nat.Biotechnol* 20:1251-1255 (2002)) In this technique, one can design the fragments to be shuffled. This aids in increasing the resulting diversity of the progeny. One can design
- 30 sequence/codon biases to make more distantly related sequences recombine at rates approaching those observed with more closely related sequences. Additionally, the technique does not require physically possessing the template genes.

- Nucleotide Exchange and Excision Technology Next exploits a combination of dUTP incorporation followed by treatment with uracil DNA glycosylase and then piperidine to perform endpoint DNA fragmentation. (Muller et al., *Nucleic Acids Res* 33:e117 (2005)) The gene is reassembled using internal PCR primer extension with proofreading polymerase. The sizes for shuffling are directly controllable using varying dUPT::dTTP ratios. This is an endpoint reaction using simple methods for uracil incorporation and cleavage. Other nucleotide analogs, such as 8-oxo-guanine, can be used with this method. Additionally, the technique works well with very short fragments (86 bp) and has a low error rate. The chemical cleavage of DNA used in this technique results in very few unshuffled clones.
- 10 In Sequence Homology-Independent Protein Recombination (SHIPREC) a linker is used to facilitate fusion between two distantly/unrelated genes. Nuclease treatment is used to generate a range of chimeras between the two genes. These fusions result in libraries of single-crossover hybrids. (Sieber et al., *Nat.Biotechnol* 19:456-460 (2001)) This produces a limited type of shuffling and a separate process is required for mutagenesis. In addition, since no homology is
- 15 needed this technique can create a library of chimeras with varying fractions of each of the two unrelated parent genes. SHIPREC was tested with a heme-binding domain of a bacterial CP450 fused to N-terminal regions of a mammalian CP450; this produced mammalian activity in a more soluble enzyme.
- In Gene Site Saturation MutagenesisTM (GSSMTM) the starting materials are a supercoiled
- 20 dsDNA plasmid containing an insert and two primers which are degenerate at the desired site of mutations. (Kretz et al., *Methods Enzymol.* 388:3-11 (2004)) Primers carrying the mutation of interest, anneal to the same sequence on opposite strands of DNA. The mutation is typically in the middle of the primer and flanked on each side by ~20 nucleotides of correct sequence. The sequence in the primer is NNN or NNK (coding) and MNN (noncoding) (N = all 4, K = G, T, M
- 25 = A, C). After extension, DpnI is used to digest dam-methylated DNA to eliminate the wild-type template. This technique explores all possible amino acid substitutions at a given locus (i.e., one codon). The technique facilitates the generation of all possible replacements at a single-site with no nonsense codons and results in equal to near-equal representation of most possible alleles. This technique does not require prior knowledge of the structure, mechanism,
- 30 or domains of the target enzyme. If followed by shuffling or Gene Reassembly, this technology creates a diverse library of recombinants containing all possible combinations of single-site up-mutations. The utility of this technology combination has been demonstrated for the successful evolution of over 50 different enzymes, and also for more than one property in a given enzyme.

Combinatorial Cassette Mutagenesis (CCM) involves the use of short oligonucleotide cassettes to replace limited regions with a large number of possible amino acid sequence alterations. (Reidhaar-Olson et al. *Methods Enzymol.* 208:564-586 (1991); and Reidhaar-Olson et al. *Science* 241:53-57 (1988)) Simultaneous substitutions at two or three sites are possible using
5 this technique. Additionally, the method tests a large multiplicity of possible sequence changes at a limited range of sites. This technique has been used to explore the information content of the lambda repressor DNA-binding domain.

Combinatorial Multiple Cassette Mutagenesis (CMCM) is essentially similar to CCM except it is employed as part of a larger program: 1) Use of epPCR at high mutation rate to 2) ID hot spots
10 and hot regions and then 3) extension by CMCM to cover a defined region of protein sequence space. (Reetz, M. T., S. Wilensek, D. Zha, and K. E. Jaeger, 2001, Directed Evolution of an Enantioselective Enzyme through Combinatorial Multiple-Cassette Mutagenesis. *Angew.Chem.Int.Ed Engl.* 40:3589-3591.) As with CCM, this method can test virtually all possible alterations over a target region. If used along with methods to create random mutations
15 and shuffled genes, it provides an excellent means of generating diverse, shuffled proteins. This approach was successful in increasing, by 51-fold, the enantioselectivity of an enzyme.

In the Mutator Strains technique conditional *ts* mutator plasmids allow increases of 20- to 4000-
X in random and natural mutation frequency during selection and block accumulation of deleterious mutations when selection is not required. (Selifonova et al., *Appl Environ Microbiol*
20 67:3645-3649 (2001)) This technology is based on a plasmid-derived *mutD5* gene, which encodes a mutant subunit of DNA polymerase III. This subunit binds to endogenous DNA polymerase III and compromises the proofreading ability of polymerase III in any strain that harbors the plasmid. A broad-spectrum of base substitutions and frameshift mutations occur. In order for effective use, the mutator plasmid should be removed once the desired phenotype is
25 achieved; this is accomplished through a temperature sensitive origin of replication, which allows for plasmid curing at 41°C. It should be noted that mutator strains have been explored for quite some time (e.g., see Low et al., *J. Mol. Biol.* 260:359-3680 (1996)). In this technique very high spontaneous mutation rates are observed. The conditional property minimizes non-desired background mutations. This technology could be combined with adaptive evolution to
30 enhance mutagenesis rates and more rapidly achieve desired phenotypes.

"Look-Through Mutagenesis (LTM) is a multidimensional mutagenesis method that assesses and optimizes combinatorial mutations of selected amino acids." (Rajpal et al., *Proc Natl Acad Sci U.S.A.* 102:8466-8471 (2005)) Rather than saturating each site with all possible amino acid

changes, a set of nine is chosen to cover the range of amino acid R-group chemistry. Fewer changes per site allows multiple sites to be subjected to this type of mutagenesis. A >800-fold increase in binding affinity for an antibody from low nanomolar to picomolar has been achieved through this method. This is a rational approach to minimize the number of random

- 5 combinations and can increase the ability to find improved traits by greatly decreasing the numbers of clones to be screened. This has been applied to antibody engineering, specifically to increase the binding affinity and/or reduce dissociation. The technique can be combined with either screens or selections.

- 10 Gene Reassembly is a DNA shuffling method that can be applied to multiple genes at one time or to creating a large library of chimeras (multiple mutations) of a single gene. (Tunable GeneReassembly™ (TGR™) Technology supplied by Verenum Corporation) Typically this technology is used in combination with ultra-high-throughput screening to query the represented sequence space for desired improvements. This technique allows multiple gene recombination independent of homology. The exact number and position of cross-over events can be pre-
- 15 determined using fragments designed via bioinformatic analysis. This technology leads to a very high level of diversity with virtually no parental gene reformation and a low level of inactive genes. Combined with GSSM™, a large range of mutations can be tested for improved activity. The method allows “blending” and “fine tuning” of DNA shuffling, e.g. codon usage can be optimized.

- 20 *In Silico* Protein Design Automation (PDA) is an optimization algorithm that anchors the structurally defined protein backbone possessing a particular fold, and searches sequence space for amino acid substitutions that can stabilize the fold and overall protein energetics. (Hayes et al., *Proc Natl Acad Sci U.S.A.* 99:15926-15931 (2002)) This technology uses *in silico* structure-based entropy predictions in order to search for structural tolerance toward protein amino acid
- 25 variations. Statistical mechanics is applied to calculate coupling interactions at each position. Structural tolerance toward amino acid substitution is a measure of coupling. Ultimately, this technology is designed to yield desired modifications of protein properties while maintaining the integrity of structural characteristics. The method computationally assesses and allows filtering of a very large number of possible sequence variants (10^{50}). The choice of sequence variants to
- 30 test is related to predictions based on the most favorable thermodynamics. Ostensibly only stability or properties that are linked to stability can be effectively addressed with this technology. The method has been successfully used in some therapeutic proteins, especially in engineering immunoglobulins. *In silico* predictions avoid testing extraordinarily large numbers of potential variants. Predictions based on existing three-dimensional structures are more likely

to succeed than predictions based on hypothetical structures. This technology can readily predict and allow targeted screening of multiple simultaneous mutations, something not possible with purely experimental technologies due to exponential increases in numbers.

Iterative Saturation Mutagenesis (ISM) involves: 1) use knowledge of structure/function to
5 choose a likely site for enzyme improvement; 2) saturation mutagenesis at chosen site using Stratagene QuikChange (or other suitable means); 3) screen/select for desired properties; and 4) with improved clone(s), start over at another site and continue repeating. (Reetz et al., *Nat. Protoc.* 2:891-903 (2007); and Reetz et al., *Angew. Chem. Int. Ed Engl.* 45:7745-7751 (2006))
This is a proven methodology, which assures all possible replacements at a given position are
10 made for screening/selection.

Any of the aforementioned methods for mutagenesis can be used alone or in any combination. Additionally, any one or combination of the directed evolution methods can be used in conjunction with adaptive evolution techniques.

In one embodiment, the invention provides a method for producing n-propanol and isopropanol
15 that includes culturing a non-naturally occurring microbial organism, including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway having at least one exogenous nucleic acid encoding an n-propanol pathway enzyme expressed in a sufficient amount to produce n-propanol, the n-propanol pathway including a propionaldehyde dehydrogenase, a propanol dehydrogenase, a propionyl-CoA:phosphate
20 propanoyltransferase, a propionyl-CoA hydrolase, a propionyl-CoA transferase, a propionyl-CoA synthetase, a propionate kinase, a propionate reductase or a propionyl phosphate reductase,, the isopropanol pathway comprising at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, the isopropanol pathway including an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase,
25 an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

In a further aspect of the above embodiment, the method includes a microbial organism having an acetyl-CoA pathway having at least one exogenous nucleic acid encoding an acetyl-CoA pathway enzyme expressed in a sufficient amount to produce acetyl-CoA, the acetyl-CoA
30 pathway including a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, or a formate dehydrogenase.

- In further embodiment, the method includes a microbial organism having a propionyl-CoA pathway having at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, the propionyl-CoA pathway including a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a methylmalonyl-CoA mutase, a methylmalonyl-CoA epimerase or a methylmalonyl-CoA decarboxylase. In a further aspect, the propionyl-CoA pathway includes a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.
- In another further embodiment, the method includes a microbial organism having a propionyl-CoA pathway having at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, the propionyl-CoA pathway including a PEP carboxykinase, a PEP carboxylase, a threonine deaminase, or a 2-oxobutanoate dehydrogenase. In a further aspect, the n-propanol pathway includes 2-oxobutanoate decarboxylase.
- In another further embodiment, the method includes a microbial organism having a propionyl-CoA pathway having at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, the propionyl-CoA pathway including an acetyl-CoA carboxylase, a malonyl-CoA reductase, a malonate semialdehyde reductase or propionyl-CoA synthase.
- In another further embodiment, the method includes a microbial organism having a propionyl-CoA pathway having at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, the propionyl-CoA pathway including a lactate dehydrogenase, a lactate-CoA transferase, a lactyl-CoA dehydratase or acryloyl CoA reductase.
- In yet another embodiment, the invention provides a method for producing n-propanol and isopropanol that includes culturing a non-naturally occurring microbial organism, including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway having a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids encoding a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol

dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the isopropanol pathway having a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In a further aspect of the above embodiment, the method includes a microbial organism having an acetyl-CoA pathway having a third set of exogenous nucleic acids encoding acetyl-CoA pathway enzymes expressed in a sufficient amount to produce acetyl-CoA, the third set of exogenous nucleic acids encoding a pyruvate kinase; and a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase.

In another further embodiment, the method includes a microbial organism having a propionyl-CoA pathway having a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA, the third set of exogenous nucleic acids encoding a PEP carboxykinase or a PEP carboxylase; a malate dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA synthetase; a methylmalonyl-CoA mutase; and a methylmalonyl-CoA decarboxylase. In a further aspect, the third set of exogenous nucleic acids further encodes a methylmalonyl-CoA epimerase or a pyruvate carboxylase.

In another further embodiment, the method includes a microbial organism having a propionyl-CoA pathway having a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA, said third set of exogenous nucleic acids encoding a PEP carboxykinase or a PEP carboxylase; a threonine deaminase; and a 2-oxobutanoate dehydrogenase. In a further aspect, the third set of exogenous nucleic acids further encodes a methylmalonyl-CoA decarboxylase or a pyruvate carboxylase. In yet another aspect, the second set of exogenous nucleic acids further encodes a 2-oxobutanoate decarboxylase.

In another further embodiment, the method includes a microbial organism having a propionyl-CoA pathway having a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA, the third set of exogenous

nucleic acids encoding an acetyl-CoA carboxylase; a malonyl-CoA reductase; a malonate semialdehyde reductase; and propionyl-CoA synthase.

In another further embodiment, the method includes a microbial organism having a propionyl-CoA pathway having a third set of exogenous nucleic acids encoding a lactate dehydrogenase; a
5 lactate-CoA transferase; a lactyl-CoA dehydratase; and acryloyl CoA reductase.

In one embodiment, the invention provides a method for producing n-propanol and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes
10 expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids encoding a PEP carboxykinase or a PEP carboxylase; a malate dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA synthetase; a methylmalonyl-CoA mutase; a methylmalonyl-CoA decarboxylase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase and a propionyl
15 phosphate reductase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway
20 enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase;
25 and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing n-propanol and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes
30 expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids encoding a PEP carboxykinase or a PEP carboxylase; a threonine deaminase; and a 2-oxobutanoate decarboxylase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a 2-oxobutanoate

- dehydrogenase, a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a 2-oxobutanoate
- 5 dehydrogenase, a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase,, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin
- 10 oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase. In a further aspect, the second set of exogenous nucleic acids further encodes a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.
- 15 In one embodiment, the invention provides a method for producing n-propanol and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids
- 20 encoding a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA carboxylase; a malonyl-CoA reductase; a malonate semialdehyde reductase; propionyl-CoA synthase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate
- 25 reductase and propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol
- 30 pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing n-propanol and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway and an isopropanol pathway, the n-propanol pathway including a first set of exogenous nucleic acids encoding n-propanol pathway enzymes
5 expressed in a sufficient amount to produce n-propanol, the first set of exogenous nucleic acids encoding a lactate dehydrogenase; a lactate-CoA transferase; a lactyl-CoA dehydratase; acryloyl CoA reductase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a
10 propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous
15 nucleic acids encoding a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 In one embodiment, the invention provides a method for producing n-propanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an n-propanol pathway, the n-propanol pathway comprising at least one exogenous nucleic acid encoding an n-propanol pathway enzyme expressed in a sufficient amount to produce n-propanol, the n-propanol pathway including a propionaldehyde dehydrogenase, a propanol
25 dehydrogenase, a propionyl-CoA:phosphate propanoyltransferase, a propionyl-CoA hydrolase, a propionyl-CoA transferase, a propionyl-CoA synthetase, a propionate kinase, a propionate reductase, or a propionyl phosphate reductase.

In another embodiment, the invention provides a method for producing n-propanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having
30 an n-propanol pathway, the n-propanol pathway comprising a set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, the set of exogenous nucleic acids encoding a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA

transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase.

In a further aspect of the above embodiment, the method for producing an propanol includes
5 culturing the non-naturally occurring microbial organism having an n-propanol pathway that
also has a propionyl-CoA pathway including exogenous nucleic acids encoding propionyl-CoA
pathway enzymes expressed in a sufficient amount to produce propionyl-CoA as exemplified
herein. For example, in some aspects the exogenous nucleic acids encode a PEP carboxykinase,
a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA
10 transferase, a succinyl-CoA synthetase, a methylmalonyl-CoA mutase, or a methylmalonyl-CoA
decarboxylase. In another aspect, the exogenous nucleic acids further encode a methylmalonyl-
CoA epimerase. Additionally, in yet another aspect of the above embodiment, the method for
producing an propanol includes culturing the non-naturally occurring microbial organism having
an n-propanol pathway that has a first set of exogenous nucleic acids encoding n-propanol
15 pathway enzymes expressed in a sufficient amount to produce n-propanol, wherein the first set
of exogenous nucleic acids encode a PEP carboxykinase or a PEP carboxylase; a malate
dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA
synthetase; a methylmalonyl-CoA mutase; a methylmalonyl-CoA epimerase; a methylmalonyl-
CoA decarboxylase; a propionaldehyde dehydrogenase and a propanol dehydrogenase.

20 In one embodiment, the invention provides a method for producing 14-BDO and isopropanol
that includes culturing a non-naturally occurring microbial organism, including a microbial
organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway having
at least one exogenous nucleic acid encoding an 14-BDO pathway enzyme expressed in a
sufficient amount to produce 14-BDO, the 14-BDO pathway including a succinyl-CoA
25 reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA
transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde-
forming), a 4-hydroxybutyraldehyde reductase, a 4-hydroxybutyrate reductase; a 4-
hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-phosphate
reductase or a 4-hydroxybutyryl-CoA reductase (alcohol-forming), the isopropanol pathway
30 including at least one exogenous nucleic acid encoding an isopropanol pathway enzyme
expressed in a sufficient amount to produce isopropanol, the isopropanol pathway including an
acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an
acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism, including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway having at least one exogenous nucleic acid encoding an 13-BDO pathway enzyme expressed in a

5 sufficient amount to produce 13-BDO, the 13-BDO pathway including a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA dehydratase, a crotonase, a 3-hydroxybutyryl-CoA reductase (aldehyde forming), a 3-hydroxybutyraldehyde reductase, a 3-hydroxybutyryl-CoA

10 transferase, a 3-hydroxybutyryl-CoA synthetase, a 3-hydroxybutyryl-CoA hydrolase, or a 3-hydroxybutyrate reductase, or a 3-hydroxybutyryl-CoA reductase (alcohol-forming), the isopropanol pathway including at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, the isopropanol pathway including an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an

15 acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism, including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway having at least one

20 exogenous nucleic acid encoding an MAA pathway enzyme expressed in a sufficient amount to produce MAA, the MAA pathway including a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA mutase, a 3-hydroxyisobutyryl-CoA dehydratase, a methacrylyl-CoA

25 transferase, a methacrylyl-CoA synthetase, a methacrylyl-CoA hydrolase, a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase, a 3-hydroxyisobutyryl-CoA hydrolase, a 3-hydroxyisobutyrate dehydratase, a methylmalonyl-CoA mutase, a methylmalonyl-CoA epimerase, a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase, a methylmalonyl-CoA hydrolase, a methylmalonate reductase, a

30 methylmalonyl-CoA reductase (aldehyde forming), a 3-hydroxyisobutyrate dehydrogenase, a methylmalonyl-CoA reductase (alcohol forming) or a 3-hydroxyisobutyrate dehydratase, the isopropanol pathway including at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, the isopropanol pathway including an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an

acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase or an isopropanol dehydrogenase.

In a further aspect of the above embodiments, the microbial organism has an acetyl-CoA pathway having at least one exogenous nucleic acid encoding an acetyl-CoA pathway enzyme
5 expressed in a sufficient amount to produce acetyl-CoA, the acetyl-CoA pathway including a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, or a formate dehydrogenase.

In further aspect of the above embodiments, the microbial organism has a succinyl-CoA pathway having at least one exogenous nucleic acid encoding a succinyl-CoA pathway enzyme
10 expressed in a sufficient amount to produce succinyl-CoA, the succinyl-CoA pathway including a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase or a succinyl-CoA synthetase. In a further aspect, the succinyl-CoA pathway includes a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

15 In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a
20 succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic
25 acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial
30 organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate reductase;

and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a 4-hydroxybutyryl-phosphate reductase; and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed

in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding
5 isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol
10 that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA
15 transferase or a 4-hydroxybutyryl-CoA synthetase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming); and a 4-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase
20 or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway
25 including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set
30 of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed
5 in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate reductase; and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-
10 CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway
15 including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of
20 exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol
25 that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a 4-
30 hydroxybutyryl-phosphate reductase; and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase

or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial
5 organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming),
10 and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

15 In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a
20 succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-
25 CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial
30 organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a

crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3 hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.
- In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a

crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3 hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed
5 in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding
10 isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol
15 that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4 hydroxybutyrate kinase; a
20 phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3 hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the
25 second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial
30 organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a crotonase; and a 3-hydroxybutyryl-CoA reductase

(alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism

having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4 hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce

isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to

- produce MAA, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase; and an isopropanol dehydrogenase.
- 10 In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4 hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a methylmalonyl-CoA mutase; a methylmalonyl-CoA reductase (aldehyde forming); a 3-hydroxyisobutyrate dehydrogenase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a methylmalonyl-CoA mutase; a methylmalonyl-CoA epimerase; a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase, or a methylmalonyl-CoA hydrolase; a methylmalonate reductase; a 3-hydroxyisobutyrate dehydrogenase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a methylmalonyl-CoA mutase; a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase or a methylmalonyl-CoA hydrolase; a methylmalonate reductase; a 3-hydroxyisobutyrate dehydrogenase; and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids a methylmalonyl-CoA mutase; a methylmalonyl-CoA reductase (alcohol forming); and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding an acetyl-CoA acetyl thiolase; an acetoacetyl-

CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

In a further aspect of the above embodiments, the microbial organism has an acetyl-CoA pathway having a third set of exogenous nucleic acids encoding acetyl-CoA pathway enzymes expressed in a sufficient amount to produce acetyl-CoA, the third set of exogenous nucleic acids encoding a pyruvate kinase; and a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase.

In another further embodiment, the microbial organism has a succinyl-CoA pathway having a third set of exogenous nucleic acids encoding succinyl-CoA pathway enzymes expressed in a sufficient amount to produce succinyl-CoA, the third set of exogenous nucleic acids encoding a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase and a succinyl-CoA synthetase. In a further aspect, the third set of exogenous nucleic acids further encodes a methylmalonyl-CoA epimerase, a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

In one embodiment, the invention provides a method for producing 14-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an 14-BDO pathway and an isopropanol pathway, the 14-BDO pathway including a first set of exogenous nucleic acids encoding 14-BDO pathway enzymes expressed in a sufficient amount to produce 14-BDO, the first set of exogenous nucleic acids encoding a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a pyruvate carboxylase, a methylmalonyl-CoA carboxytransferase, a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde-forming), a 4-hydroxybutyraldehyde reductase, a 4-hydroxybutyrate reductase; a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-phosphate reductase, a 4-hydroxybutyryl-CoA reductase (alcohol-forming), and a 4-hydroxybutyraldehyde reductase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, a formate dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-

CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase, and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing 13-BDO and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial
 5 organism having an 13-BDO pathway and an isopropanol pathway, the 13-BDO pathway including a first set of exogenous nucleic acids encoding 13-BDO pathway enzymes expressed in a sufficient amount to produce 13-BDO, the first set of exogenous nucleic acids encoding PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a pyruvate carboxylase, a
 10 methylmalonyl-CoA carboxytransferase, a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA dehydratase, a crotonase, a 3-hydroxybutyryl-CoA reductase (aldehyde forming), a 3-hydroxybutyraldehyde reductase, a 3-hydroxybutyryl-CoA transferase, a 3-
 15 hydroxybutyryl-CoA synthetase, a 3-hydroxybutyryl-CoA hydrolase, a 3-hydroxybutyrate reductase, and a 3-hydroxybutyryl-CoA reductase (alcohol-forming), and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin
 20 oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, a formate dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase, and an isopropanol dehydrogenase.

In one embodiment, the invention provides a method for producing MAA and isopropanol that
 25 includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA
 30 transferase, a succinyl-CoA synthetase, a pyruvate carboxylase, a methylmalonyl-CoA carboxytransferase, a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA mutase, a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase, a 3-

- hydroxyisobutyryl-CoA hydrolase, 3-hydroxyisobutyryl-CoA dehydratase, methacrylyl-CoA transferase, methacrylyl-CoA synthetase, methacrylyl-CoA hydrolase and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient
- 5 amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, a formate dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase, and an isopropanol dehydrogenase.
- 10 In one embodiment, the invention provides a method for producing MAA and isopropanol that includes culturing a non-naturally occurring microbial organism including a microbial organism having an MAA pathway and an isopropanol pathway, the MAA pathway including a first set of exogenous nucleic acids encoding MAA pathway enzymes expressed in a sufficient amount to produce MAA, the first set of exogenous nucleic acids encoding a PEP carboxykinase, a PEP
- 15 carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a pyruvate carboxylase, a methylmalonyl-CoA carboxytransferase, a methylmalonyl-CoA mutase, a methylmalonyl-CoA epimerase, a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase, a methylmalonyl-CoA hydrolase, a methylmalonate reductase, a methylmalonyl-CoA reductase (aldehyde forming), a
- 20 3-hydroxyisobutyrate dehydrogenase, a methylmalonyl-CoA reductase (alcohol forming) and a 3-hydroxyisobutyrate dehydratase, and the isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, the second set of exogenous nucleic acids encoding a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate
- 25 lyase, a pyruvate formate lyase activating enzyme, a formate dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, an acetoacetate decarboxylase, and an isopropanol dehydrogenase.

In a further aspect of each of the above embodiments, the exogenous nucleic acid is a heterologous nucleic acid.

- 30 In a further aspect of each of the above embodiments, the conditions include substantially anaerobic culture conditions.

Suitable purification and/or assays to test for the production of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA can be performed using well known methods. Suitable replicates

such as triplicate cultures can be grown for each engineered strain to be tested. For example, product and byproduct formation in the engineered production host can be monitored. The final product and intermediates, and other organic compounds, can be analyzed by methods such as HPLC (High Performance Liquid Chromatography), GC-MS (Gas Chromatography-Mass Spectroscopy) and LC-MS (Liquid Chromatography-Mass Spectroscopy) or other suitable analytical methods using routine procedures well known in the art. The release of product in the fermentation broth can also be tested with the culture supernatant. Byproducts and residual glucose can be quantified by HPLC using, for example, a refractive index detector for glucose and alcohols, and a UV detector for organic acids (Lin et al., *Biotechnol. Bioeng.* 90:775-779 (2005)), or other suitable assay and detection methods well known in the art. The individual enzyme or protein activities from the exogenous DNA sequences can also be assayed using methods well known in the art. Various alcohols can be quantified by gas chromatography by using a flame ionization detector as described in Atsumi et al. *Metab Eng* (2007) and Hanai et al. *Appl Environ Microbiol* 73:7814-7818 (2007).

The n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA can be separated from other components in the culture using a variety of methods well known in the art. Such separation methods include, for example, extraction procedures as well as methods that include continuous liquid-liquid extraction, pervaporation, membrane filtration, membrane separation, reverse osmosis, electrodialysis, distillation, crystallization, centrifugation, extractive filtration, ion exchange chromatography, size exclusion chromatography, adsorption chromatography, and ultrafiltration. All of the above methods are well known in the art.

Any of the non-naturally occurring microbial organisms described herein can be cultured to produce and/or secrete the biosynthetic products of the invention. For example, the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA producers can be cultured for the biosynthetic production of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA.

For the production of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA, the recombinant strains are cultured in a medium with a carbon source and other essential nutrients. It is highly desirable to maintain anaerobic conditions in the fermenter to reduce the cost of the overall process. Such conditions can be obtained, for example, by first sparging the medium with nitrogen and then sealing the flasks with a septum and crimp-cap. For strains where growth is not observed anaerobically, microaerobic conditions can be applied by perforating the septum with a small hole for limited aeration. Exemplary anaerobic conditions have been described previously and are well-known in the art. Exemplary aerobic and anaerobic conditions are

described, for example, in U.S. publication 2009/0047719, filed August 10, 2007.

Fermentations can be performed in a batch, fed-batch or continuous manner, as disclosed herein.


If desired, the pH of the medium can be maintained at a desired pH, in particular neutral pH, such as a pH of around 7 by addition of a base, such as NaOH or other bases, or acid, as needed to maintain the culture medium at a desirable pH. The growth rate can be determined by measuring optical density using a spectrophotometer (600 nm), and the glucose uptake rate by monitoring carbon source depletion over time.

The growth medium can include, for example, any carbohydrate source which can supply a source of carbon to the non-naturally occurring microorganism. Such sources include, for example, sugars such as glucose, xylose, arabinose, galactose, mannose, fructose, sucrose and starch. Other sources of carbohydrate include, for example, renewable feedstocks and biomass. Exemplary types of biomasses that can be used as feedstocks in the methods of the invention include cellulosic biomass, hemicellulosic biomass and lignin feedstocks or portions of feedstocks. Such biomass feedstocks contain, for example, carbohydrate substrates useful as carbon sources such as glucose, xylose, arabinose, galactose, mannose, fructose and starch. Given the teachings and guidance provided herein, those skilled in the art will understand that renewable feedstocks and biomass other than those exemplified above also can be used for culturing the microbial organisms of the invention for the production of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA.

In addition to renewable feedstocks such as those exemplified above, the n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol microbial organisms of the invention also can be modified for growth on syngas as its source of carbon. In this specific embodiment, one or more proteins or enzymes are expressed in the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA producing organisms to provide a metabolic pathway for utilization of syngas or other gaseous carbon source.

Synthesis gas, also known as syngas or producer gas, is the major product of gasification of coal and of carbonaceous materials such as biomass materials, including agricultural crops and residues. Syngas is a mixture primarily of H₂ and CO and can be obtained from the gasification of any organic feedstock, including but not limited to coal, coal oil, natural gas, biomass, and waste organic matter. Gasification is generally carried out under a high fuel to oxygen ratio. Although largely H₂ and CO, syngas can also include CO₂ and other gases in smaller quantities. Thus, synthesis gas provides a cost effective source of gaseous carbon such as CO and, additionally, CO₂.

The Wood-Ljungdahl pathway catalyzes the conversion of CO and H₂ to acetyl-CoA and other products such as acetate. Organisms capable of utilizing CO and syngas also generally have the capability of utilizing CO₂ and CO₂/H₂ mixtures through the same basic set of enzymes and transformations encompassed by the Wood-Ljungdahl pathway. H₂-dependent conversion of CO₂ to acetate by microorganisms was recognized long before it was revealed that CO also could be used by the same organisms and that the same pathways were involved. Many acetogens have been shown to grow in the presence of CO₂ and produce compounds such as acetate as long as hydrogen is present to supply the necessary reducing equivalents (see for example, Drake, *Acetogenesis*, pp. 3-60 Chapman and Hall, New York, (1994)). This can be summarized by the following equation:



Hence, non-naturally occurring microorganisms possessing the Wood-Ljungdahl pathway can utilize CO₂ and H₂ mixtures as well for the production of acetyl-CoA and other desired products.

The Wood-Ljungdahl pathway is well known in the art and consists of 12 reactions which can be separated into two branches: (1) methyl branch and (2) carbonyl branch. The methyl branch converts syngas to methyl-tetrahydrofolate (methyl-THF) whereas the carbonyl branch converts methyl-THF to acetyl-CoA. The reactions in the methyl branch are catalyzed in order by the following enzymes or proteins: ferredoxin oxidoreductase, formate dehydrogenase, formyltetrahydrofolate synthetase, methenyltetrahydrofolate cyclodehydratase, methylenetetrahydrofolate dehydrogenase and methylenetetrahydrofolate reductase. The reactions in the carbonyl branch are catalyzed in order by the following enzymes or proteins: methyltetrahydrofolate:corrinoid protein methyltransferase (for example, AcsE), corrinoid iron-sulfur protein, nickel-protein assembly protein (for example, AcsF), ferredoxin, acetyl-CoA synthase, carbon monoxide dehydrogenase and nickel-protein assembly protein (for example, CooC). Following the teachings and guidance provided herein for introducing a sufficient number of encoding nucleic acids to generate an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway, those skilled in the art will understand that the same engineering design also can be performed with respect to introducing at least the nucleic acids encoding the Wood-Ljungdahl enzymes or proteins absent in the host organism. Therefore, introduction of one or more encoding nucleic acids into the microbial organisms of the invention such that the modified organism contains the complete Wood-Ljungdahl pathway will confer syngas utilization ability.

Additionally, the reductive (reverse) tricarboxylic acid cycle is and/or hydrogenase activities can also be used for the conversion of CO, CO₂ and/or H₂ to acetyl-CoA and other products such as acetate. Organisms capable of fixing carbon via the reductive TCA pathway can utilize one or more of the following enzymes: ATP citrate-lyase, citrate lyase, aconitase, isocitrate

5 dehydrogenase, alpha-ketoglutarate:ferredoxin oxidoreductase, succinyl-CoA synthetase, succinyl-CoA transferase, fumarate reductase, fumarase, malate dehydrogenase, NAD(P)H:ferredoxin oxidoreductase, carbon monoxide dehydrogenase, and hydrogenase. Specifically, the reducing equivalents extracted from CO and/or H₂ by carbon monoxide dehydrogenase and hydrogenase are utilized to fix CO₂ via the reductive TCA cycle into acetyl-

10 CoA or acetate. Acetate can be converted to acetyl-CoA by enzymes such as acetyl-CoA transferase, acetate kinase/phosphotransacetylase, and acetyl-CoA synthetase. Acetyl-CoA can be converted to an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA precursors, glyceraldehyde-3-phosphate, phosphoenolpyruvate, and pyruvate, by pyruvate:ferredoxin oxidoreductase and the enzymes of gluconeogenesis. Following the teachings and guidance

15 provided herein for introducing a sufficient number of encoding nucleic acids to generate an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway, those skilled in the art will understand that the same engineering design also can be performed with respect to introducing at least the nucleic acids encoding the reductive TCA pathway enzymes or proteins absent in the host organism. Therefore, introduction of one or more encoding nucleic acids into

20 the microbial organisms of the invention such that the modified organism contains the complete reductive TCA pathway will confer syngas utilization ability.

Accordingly, given the teachings and guidance provided herein, those skilled in the art will understand that a non-naturally occurring microbial organism can be produced that secretes the biosynthesized compounds of the invention when grown on a carbon source such as a

25 carbohydrate. Such compounds include, for example, n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA and any of the intermediate metabolites in the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway. All that is required is to engineer in one or more of the required enzyme or protein activities to achieve biosynthesis of the desired compound or intermediate including, for example, inclusion of some or all of the n-propanol, isopropanol, 14-

30 BDO, 13-BDO and/or MAA biosynthetic pathways. Accordingly, the invention provides a non-naturally occurring microbial organism that produces and/or secretes n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA when grown on a carbohydrate or other carbon source and produces and/or secretes any of the intermediate metabolites shown in the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA pathway when grown on a carbohydrate or other

carbon source. The n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol producing microbial organisms of the invention can initiate synthesis from an intermediate, for example, succinyl-CoA, propionyl-CoA and/or acetyl-CoA.

- 5 The non-naturally occurring microbial organisms of the invention are constructed using methods well known in the art as exemplified herein to exogenously express at least one nucleic acid encoding an n-propanol, an isopropanol, a 14-BDO, a 13-BDO and/or a MAA pathway enzyme or protein in sufficient amounts to produce n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA. It is understood that the microbial organisms of the invention are cultured under
- 10 conditions sufficient to produce n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA. Following the teachings and guidance provided herein, the non-naturally occurring microbial organisms of the invention can achieve biosynthesis of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA resulting in intracellular concentrations between about 0.1-200 mM or more. Generally, the intracellular concentration of n-propanol, isopropanol, 14-BDO, 13-BDO and/or
- 15 MAA is between about 3-150 mM, particularly between about 5-125 mM and more particularly between about 8-100 mM, including about 10 mM, 20 mM, 50 mM, 80 mM, or more. Intracellular concentrations between and above each of these exemplary ranges also can be achieved from the non-naturally occurring microbial organisms of the invention.

- In some embodiments, culture conditions include anaerobic or substantially anaerobic growth or
- 20 maintenance conditions. Exemplary anaerobic conditions have been described previously and are well known in the art. Exemplary anaerobic conditions for fermentation processes are described herein and are described, for example, in U.S. publication 2009/0047719, filed August 10, 2007. Any of these conditions can be employed with the non-naturally occurring microbial organisms as well as other anaerobic conditions well known in the art. Under such anaerobic
- 25 conditions, the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA producers can synthesize n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA at intracellular concentrations of 5-10 mM or more as well as all other concentrations exemplified herein. It is understood that, even though the above description refers to intracellular concentrations, n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA producing microbial organisms can
- 30 produce n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA intracellularly and/or secrete the product into the culture medium.

The culture conditions can include, for example, liquid culture procedures as well as fermentation and other large scale culture procedures. As described herein, particularly useful

yields of the biosynthetic products of the invention can be obtained under anaerobic or substantially anaerobic culture conditions.

As described herein, one exemplary growth condition for achieving biosynthesis of n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol
5 includes anaerobic culture or fermentation conditions. In certain embodiments, the non-naturally occurring microbial organisms of the invention can be sustained, cultured or fermented under anaerobic or substantially anaerobic conditions. Briefly, anaerobic conditions refers to an environment devoid of oxygen. Substantially anaerobic conditions include, for example, a culture, batch fermentation or continuous fermentation such that the dissolved oxygen
10 concentration in the medium remains between 0 and 10% of saturation. Substantially anaerobic conditions also includes growing or resting cells in liquid medium or on solid agar inside a sealed chamber maintained with an atmosphere of less than 1% oxygen. The percent of oxygen can be maintained by, for example, sparging the culture with an N₂/CO₂ mixture or other suitable non-oxygen gas or gases.

15 The culture conditions described herein can be scaled up and grown continuously for manufacturing of n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol. Exemplary growth procedures include, for example, fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or continuous fermentation and continuous separation. All of these processes are well known in
20 the art. Fermentation procedures are particularly useful for the biosynthetic production of commercial quantities of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA. Generally, and as with non-continuous culture procedures, the continuous and/or near-continuous production of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA will include culturing a non-naturally occurring n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and
25 isopropanol or MAA and isopropanol producing organism of the invention in sufficient nutrients and medium to sustain and/or nearly sustain growth in an exponential phase. Continuous culture under such conditions can be include, for example, growth for 1 day, 2, 3, 4, 5, 6 or 7 days or more. Additionally, continuous culture can include longer time periods of 1 week, 2, 3, 4 or 5 or more weeks and up to several months. Alternatively, organisms of the invention can be cultured
30 for hours, if suitable for a particular application. It is to be understood that the continuous and/or near-continuous culture conditions also can include all time intervals in between these exemplary periods. It is further understood that the time of culturing the microbial organism of the invention is for a sufficient period of time to produce a sufficient amount of product for a desired purpose.

Fermentation procedures are well known in the art. Briefly, fermentation for the biosynthetic production of n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol can be utilized in, for example, fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or continuous fermentation and continuous separation. Examples of batch and continuous fermentation procedures are well known in the art.

In addition to the above fermentation procedures using the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA producers of the invention for continuous production of substantial quantities of n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol, the n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA producers also can be, for example, simultaneously subjected to chemical synthesis procedures to convert the product to other compounds or the product can be separated from the fermentation culture and sequentially subjected to chemical conversion to convert the product to other compounds, if desired.

In addition to the culturing and fermentation conditions described herein, growth condition for achieving biosynthesis of n-propanol and isopropanol, 14-BDO and isopropanol, 13-BDO and isopropanol or MAA and isopropanol can include the addition of an osmoprotectant to the culturing conditions. In certain embodiments, the non-naturally occurring microbial organisms of the invention can be sustained, cultured or fermented as described herein in the presence of an osmoprotectant. Briefly, an osmoprotectant means a compound that acts as an osmolyte and helps a microbial organism as described herein survive osmotic stress. Osmoprotectants include, but are not limited to, betaines, amino acids, and the sugar trehalose. Non-limiting examples of such are glycine betaine, praline betaine, dimethylthetin, dimethylsulfoniopropionate, 3-dimethylsulfonio-2-methylpropionate, pipercolic acid, dimethylsulfonioacetate, choline, L-carnitine and ectoine. In one aspect, the osmoprotectant is glycine betaine. It is understood to one of ordinary skill in the art that the amount and type of osmoprotectant suitable for protecting a microbial organism described herein from osmotic stress will depend on the microbial organism used. The amount of osmoprotectant in the culturing conditions can be, for example, no more than about 0.1 mM, no more than about 0.5 mM, no more than about 1.0 mM, no more than about 1.5 mM, no more than about 2.0 mM, no more than about 2.5 mM, no more than about 3.0 mM, no more than about 5.0 mM, no more than about 7.0 mM, no more than about 10mM, no more than about 50mM, no more than about 100mM or no more than about 500mM.

To generate better producers, metabolic modeling can be utilized to optimize growth conditions. Modeling can also be used to design gene knockouts that additionally optimize utilization of the pathway (see, for example, U.S. patent publications US 2002/0012939, US 2003/0224363, US 2004/0029149, US 2004/0072723, US 2003/0059792, US 2002/0168654 and US 2004/0009466, and U.S. Patent No. 7,127,379). Modeling analysis allows reliable predictions of the effects on cell growth of shifting the metabolism towards more efficient production of n-propanol, isopropanol, 14-BDO, 13-BDO and/or MAA.

One computational method for identifying and designing metabolic alterations favoring biosynthesis of a desired product is the OptKnock computational framework (Burgard et al., *Biotechnol. Bioeng.* 84:647-657 (2003)). OptKnock is a metabolic modeling and simulation program that suggests gene deletion or disruption strategies that result in genetically stable microorganisms which overproduce the target product. Specifically, the framework examines the complete metabolic and/or biochemical network of a microorganism in order to suggest genetic manipulations that force the desired biochemical to become an obligatory byproduct of cell growth. By coupling biochemical production with cell growth through strategically placed gene deletions or other functional gene disruption, the growth selection pressures imposed on the engineered strains after long periods of time in a bioreactor lead to improvements in performance as a result of the compulsory growth-coupled biochemical production. Lastly, when gene deletions are constructed there is a negligible possibility of the designed strains reverting to their wild-type states because the genes selected by OptKnock are to be completely removed from the genome. Therefore, this computational methodology can be used to either identify alternative pathways that lead to biosynthesis of a desired product or used in connection with the non-naturally occurring microbial organisms for further optimization of biosynthesis of a desired product.

Briefly, OptKnock is a term used herein to refer to a computational method and system for modeling cellular metabolism. The OptKnock program relates to a framework of models and methods that incorporate particular constraints into flux balance analysis (FBA) models. These constraints include, for example, qualitative kinetic information, qualitative regulatory information, and/or DNA microarray experimental data. OptKnock also computes solutions to various metabolic problems by, for example, tightening the flux boundaries derived through flux balance models and subsequently probing the performance limits of metabolic networks in the presence of gene additions or deletions. OptKnock computational framework allows the construction of model formulations that allow an effective query of the performance limits of metabolic networks and provides methods for solving the resulting mixed-integer linear

programming problems. The metabolic modeling and simulation methods referred to herein as OptKnock are described in, for example, U.S. publication 2002/0168654, filed January 10, 2002, in International Patent No. PCT/US02/00660, filed January 10, 2002, and U.S. publication 2009/0047719, filed August 10, 2007.

- 5 Another computational method for identifying and designing metabolic alterations favoring biosynthetic production of a product is a metabolic modeling and simulation system termed SimPheny®. This computational method and system is described in, for example, U.S. publication 2003/0233218, filed June 14, 2002, and in International Patent Application No. PCT/US03/18838, filed June 13, 2003. SimPheny® is a computational system that can be used
10 to produce a network model *in silico* and to simulate the flux of mass, energy or charge through the chemical reactions of a biological system to define a solution space that contains any and all possible functionalities of the chemical reactions in the system, thereby determining a range of allowed activities for the biological system. This approach is referred to as constraints-based modeling because the solution space is defined by constraints such as the known stoichiometry
15 of the included reactions as well as reaction thermodynamic and capacity constraints associated with maximum fluxes through reactions. The space defined by these constraints can be interrogated to determine the phenotypic capabilities and behavior of the biological system or of its biochemical components.

- These computational approaches are consistent with biological realities because biological
20 systems are flexible and can reach the same result in many different ways. Biological systems are designed through evolutionary mechanisms that have been restricted by fundamental constraints that all living systems must face. Therefore, constraints-based modeling strategy embraces these general realities. Further, the ability to continuously impose further restrictions on a network model via the tightening of constraints results in a reduction in the size of the
25 solution space, thereby enhancing the precision with which physiological performance or phenotype can be predicted.

- Given the teachings and guidance provided herein, those skilled in the art will be able to apply various computational frameworks for metabolic modeling and simulation to design and implement biosynthesis of a desired compound in host microbial organisms. Such metabolic
30 modeling and simulation methods include, for example, the computational systems exemplified above as SimPheny® and OptKnock. For illustration of the invention, some methods are described herein with reference to the OptKnock computation framework for modeling and simulation. Those skilled in the art will know how to apply the identification, design and

implementation of the metabolic alterations using OptKnock to any of such other metabolic modeling and simulation computational frameworks and methods well known in the art.

The methods described above will provide one set of metabolic reactions to disrupt. Elimination of each reaction within the set or metabolic modification can result in a desired product as an
5 obligatory product during the growth phase of the organism. Because the reactions are known, a solution to the bilevel OptKnock problem also will provide the associated gene or genes encoding one or more enzymes that catalyze each reaction within the set of reactions. Identification of a set of reactions and their corresponding genes encoding the enzymes participating in each reaction is generally an automated process, accomplished through
10 correlation of the reactions with a reaction database having a relationship between enzymes and encoding genes.

Once identified, the set of reactions that are to be disrupted in order to achieve production of a desired product are implemented in the target cell or organism by functional disruption of at least one gene encoding each metabolic reaction within the set. One particularly useful means to
15 achieve functional disruption of the reaction set is by deletion of each encoding gene. However, in some instances, it can be beneficial to disrupt the reaction by other genetic aberrations including, for example, mutation, deletion of regulatory regions such as promoters or cis binding sites for regulatory factors, or by truncation of the coding sequence at any of a number of locations. These latter aberrations, resulting in less than total deletion of the gene set can be
20 useful, for example, when rapid assessments of the coupling of a product are desired or when genetic reversion is less likely to occur.

To identify additional productive solutions to the above described bilevel OptKnock problem which lead to further sets of reactions to disrupt or metabolic modifications that can result in the biosynthesis, including growth-coupled biosynthesis of a desired product, an optimization
25 method, termed integer cuts, can be implemented. This method proceeds by iteratively solving the OptKnock problem exemplified above with the incorporation of an additional constraint referred to as an integer cut at each iteration. Integer cut constraints effectively prevent the solution procedure from choosing the exact same set of reactions identified in any previous iteration that obligatorily couples product biosynthesis to growth. For example, if a previously
30 identified growth-coupled metabolic modification specifies reactions 1, 2, and 3 for disruption, then the following constraint prevents the same reactions from being simultaneously considered in subsequent solutions. The integer cut method is well known in the art and can be found described in, for example, Burgard et al., *Biotechnol. Prog.* 17:791-797 (2001). As with all

methods described herein with reference to their use in combination with the OptKnock computational framework for metabolic modeling and simulation, the integer cut method of reducing redundancy in iterative computational analysis also can be applied with other computational frameworks well known in the art including, for example, SimPheny®.

- 5 The methods exemplified herein allow the construction of cells and organisms that biosynthetically produce a desired product, including the obligatory coupling of production of a target biochemical product to growth of the cell or organism engineered to harbor the identified genetic alterations. Therefore, the computational methods described herein allow the identification and implementation of metabolic modifications that are identified by an *in silico* method selected from OptKnock or SimPheny®. The set of metabolic modifications can include, for example, addition of one or more biosynthetic pathway enzymes and/or functional disruption of one or more metabolic reactions including, for example, disruption by gene deletion.

As discussed above, the OptKnock methodology was developed on the premise that mutant microbial networks can be evolved towards their computationally predicted maximum-growth phenotypes when subjected to long periods of growth selection. In other words, the approach leverages an organism's ability to self-optimize under selective pressures. The OptKnock framework allows for the exhaustive enumeration of gene deletion combinations that force a coupling between biochemical production and cell growth based on network stoichiometry. The identification of optimal gene/reaction knockouts requires the solution of a bilevel optimization problem that chooses the set of active reactions such that an optimal growth solution for the resulting network overproduces the biochemical of interest (Burgard et al., *Biotechnol. Bioeng.* 84:647-657 (2003)).

An *in silico* stoichiometric model of *E. coli* metabolism can be employed to identify essential genes for metabolic pathways as exemplified previously and described in, for example, U.S. patent publications US 2002/0012939, US 2003/0224363, US 2004/0029149, US 2004/0072723, US 2003/0059792, US 2002/0168654 and US 2004/0009466, and in U.S. Patent No. 7,127,379. As disclosed herein, the OptKnock mathematical framework can be applied to pinpoint gene deletions leading to the growth-coupled production of a desired product. Further, the solution of the bilevel OptKnock problem provides only one set of deletions. To enumerate all meaningful solutions, that is, all sets of knockouts leading to growth-coupled production formation, an optimization technique, termed integer cuts, can be implemented. This entails iteratively solving

the OptKnock problem with the incorporation of an additional constraint referred to as an integer cut at each iteration, as discussed above.

It is understood that modifications which do not substantially affect the activity of the various embodiments of this invention are also provided within the definition of the invention provided herein. Accordingly, the following examples are intended to illustrate but not limit the present invention.

EXAMPLE I

Pathways for Co-production of n-Propanol and Isopropanol from Glucose

This example describes exemplary pathways for co-production of n-propanol and isopropanol.

Novel pathways for co-producing n-propanol and isopropanol and related products are described herein. This invention provides four alternate methods for co-production of n-propanol and isopropanol. The production of isopropanol in *E. coli* has been described previously (Hanai et al., *Appl Environ Microbiol* 73:7814-7818 (2007)). Briefly, acetyl CoA is converted into acetoacetyl CoA, transformed into acetoacetate, decarboxylated to form acetone and then reduced to form isopropanol (Figures 1-4). The microbial organisms and methods described herein combine this known route with four novel pathways for synthesizing n-propanol. This co-production will provide completely redox balanced routes for production of the C3 alcohols, i.e. n-propanol and isopropanol, allowing for anaerobic production as opposed to the requirement of oxygen if isopropanol is produced solely via acetone as described by Hanai et al., *supra*. One advantage to the co-production of n-propanol and isopropanol using any of the pathways described herein is that the maximum theoretical yield of the C3 alcohols is afforded:



Furthermore, all of these pathways have a net positive yield of ATP.

Production of isopropanol utilizing acetyl-CoA

Isopropanol production is achieved via conversion of acetyl-CoA by an acetoacetyl-CoA thiolase, an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase,, an acetoacetate decarboxylase, and an isopropanol dehydrogenase as exemplified in Figures 1-4. Isopropanol production has been described for recombinant *E. coli* following expression of two heterologous genes from *C. acetobutylicum* (*thl* and *adc* encoding acetoacetyl-CoA thiolase and acetoacetate decarboxylase, respectively) and one from *C. beijerinckii* (*adh*

encoding a secondary alcohol dehydrogenase), along with the increased expression of the native *atoA* and *atoD* genes which encode acetoacetyl-CoA:acetate:CoA transferase activity (Hanai et al., *Appl Environ Microbiol* 73:7814-7818 (2007)). The conversion of acetoacetyl-CoA to acetoacetate can alternately be catalyzed by an enzyme with acetoacetyl-CoA hydrolase or acetoacetyl-CoA synthetase activities.

Acetoacetyl-CoA thiolase

Acetoacetyl-CoA thiolase (also known as acetyl-CoA acetyltransferase) converts two molecules of acetyl-CoA into one molecule each of acetoacetyl-CoA and CoA. Exemplary acetoacetyl-CoA thiolase enzymes include the gene products of *atoB* from *E. coli* (Martin et al., *Nat. Biotechnol* 21:796-802 (2003)), *thlA* and *thlB* from *C. acetobutylicum* (Hanai et al., *Appl Environ Microbiol* 73:7814-7818 (2007); Winzer et al., *J. Mol. Microbiol Biotechnol* 2:531-541 (2000), and *ERG10* from *S. cerevisiae* Hiser et al., *J. Biol. Chem.* 269:31383-31389 (1994)). These genes/proteins are identified below in Table 1.

Table 1.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>AtoB</i>	NP_416728	16130161	<i>Escherichia coli</i>
<i>ThlA</i>	NP_349476.1	15896127	<i>Clostridium acetobutylicum</i>
<i>ThlB</i>	NP_149242.1	15004782	<i>Clostridium acetobutylicum</i>
<i>ERG10</i>	NP_015297	6325229	<i>Saccharomyces cerevisiae</i>

Acetoacetyl-CoA transferase

Acetoacetyl-CoA transferase catalyzes the conversion of acetoacetyl-CoA to acetoacetate while transferring the CoA moiety to a CoA acceptor molecule. Many transferases have broad specificity and thus may utilize CoA acceptors as diverse as acetate, succinate, propionate, butyrate, 2-methylacetoacetate, 3-ketohexanoate, 3-ketopentanoate, valerate, crotonate, 3-mercaptopropionate, propionate, vinylacetate, butyrate, among others.

Acetoacetyl-CoA:acetate:CoA transferase converts acetoacetyl-CoA and acetate to acetoacetate and acetyl-CoA. Exemplary enzymes include the gene products of *atoAD* from *E. coli* (Hanai et al., *Appl Environ Microbiol* 73:7814-7818 (2007)), *ctfAB* from *C. acetobutylicum* (Jojima et al., *Appl Microbiol Biotechnol* 77:1219-1224 (2008), and *ctfAB* from *Clostridium saccharoperbutylacetonicum* (Kosaka et al., *Biosci. Biotechnol Biochem.* 71:58-68 (2007)) are shown below in Table 2. A succinyl-CoA:3-ketoacid CoA transferase (SCOT) can also catalyze the conversion of the 3-ketoacyl-CoA, acetoacetyl-CoA, to the 3-ketoacid, acetoacetate. As opposed to acetoacetyl-CoA:acetate:CoA transferase, SCOT employs succinate as the CoA

acceptor instead of acetate. Exemplary succinyl-CoA:3-ketoacid-CoA transferases are present in *Helicobacter pylori* (Corthesy-Theulaz et al., *J Biol Chem* 272:25659-25667 (1997)), *Bacillus subtilis* (Stols et al., *Protein Expr Purif* 53:396-403 (2007)), and *Homo sapiens* (Fukao et al., *Genomics* 68:144-151 (2000); Tanaka et al., *Mol Hum Reprod* 8:16-23 (2002)). Yet another

5 transferase capable of this conversion is butyryl-CoA:acetoacetate CoA-transferase. Exemplary enzymes can be found in *Fusobacterium nucleatum* (Barker et al., *J Bacteriol* 152(1):201-7 (1982)), *Clostridium* SB4 (Barker et al., *J Biol Chem* 253(4):1219-25 (1978)), and *Clostridium acetobutylicum* (Wiesenborn et al., *Appl Environ Microbiol* 55(2):323-9 (1989)). Although

specific gene sequences were not provided for butyryl-CoA:acetoacetate CoA-transferase in

10 these references, the genes *FN0272* and *FN0273* have been annotated as a butyrate-acetoacetate CoA-transferase (Kapatral et al., *J Bact* 184(7) 2005-2018 (2002)). Homologs in *Fusobacterium nucleatum* such as *FN1857* and *FN1856* also likely have the desired acetoacetyl-CoA transferase activity. *FN1857* and *FN1856* are located adjacent to many other genes involved in lysine

fermentation and are thus very likely to encode an acetoacetate:butyrate CoA transferase

15 (Kreimeyer, et al., *J Biol Chem* 282 (10) 7191-7197 (2007)). Additional candidates from *Porphyromonas gingivalis* and *Thermoanaerobacter tengcongensis* can be identified in a similar fashion (Kreimeyer, et al., *J Biol Chem* 282 (10) 7191-7197 (2007)). These genes/proteins are identified below in Table 2.

Table 2.

20

Gene	GenBank ID	GI Number	Organism
<i>AtoA</i>	NP_416726.1	2492994	<i>Escherichia coli</i>
<i>AtoD</i>	NP_416725.1	2492990	<i>Escherichia coli</i>
<i>CtfA</i>	NP_149326.1	15004866	<i>Clostridium acetobutylicum</i>
<i>CtfB</i>	NP_149327.1	15004867	<i>Clostridium acetobutylicum</i>
<i>CtfA</i>	AAP42564.1	31075384	<i>Clostridium saccharoperbutylacetonicum</i>
<i>CtfB</i>	AAP42565.1	31075385	<i>Clostridium saccharoperbutylacetonicum</i>
<i>HPAG1_0676</i>	YP_627417	108563101	<i>Helicobacter pylori</i>
<i>HPAG1_0677</i>	YP_627418	108563102	<i>Helicobacter pylori</i>
<i>ScoA</i>	NP_391778	16080950	<i>Bacillus subtilis</i>
<i>ScoB</i>	NP_391777	16080949	<i>Bacillus subtilis</i>
<i>OXCT1</i>	NP_000427	4557817	<i>Homo sapiens</i>
<i>OXCT2</i>	NP_071403	11545841	<i>Homo sapiens</i>
<i>FN0272</i>	NP_603179.1	19703617	<i>Fusobacterium nucleatum</i>
<i>FN0273</i>	NP_603180.1	19703618	<i>Fusobacterium nucleatum</i>
<i>FN1857</i>	NP_602657.1	19705162	<i>Fusobacterium nucleatum</i>
<i>FN1856</i>	NP_602656.1	19705161	<i>Fusobacterium nucleatum</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>PG1066</i>	NP_905281.1	34540802	<i>Porphyromonas gingivalis</i> W83
<i>PG1075</i>	NP_905290.1	34540811	<i>Porphyromonas gingivalis</i> W83
<i>TTE0720</i>	NP_622378.1	20807207	<i>Thermoanaerobacter tengcongensis</i> MB4
<i>TTE0721</i>	NP_622379.1	20807208	<i>Thermoanaerobacter tengcongensis</i> MB4

Acetoacetyl-CoA synthetase

A CoA synthetase can also catalyze the removal of the CoA moiety from acetoacetyl-CoA. One candidate enzyme, ADP-forming acetyl-CoA synthetase (ACD, EC 6.2.1.13), couples the conversion of acyl-CoA esters to their corresponding acids with the concurrent synthesis of

5 ATP. Several enzymes with broad substrate specificities have been described in the literature. ACD I from *Archaeoglobus fulgidus*, encoded by AF1211, was shown to operate on a variety of linear and branched-chain substrates including acetyl-CoA, propionyl-CoA, butyryl-CoA, acetate, propionate, butyrate, isobutyryate, isovalerate, succinate, fumarate, phenylacetate, indoleacetate (Musfeldt et al., *J Bacteriol* 184:636-644 (2002)). The enzyme from *Haloarcula*

10 *marismortui* (annotated as a succinyl-CoA synthetase) accepts propionate, butyrate, and branched-chain acids (isovalerate and isobutyrate) as substrates, and was shown to operate in the forward and reverse directions (Brasen et al., *Arch Microbiol* 182:277-287 (2004)). The ACD encoded by PAE3250 from hyperthermophilic crenarchaeon *Pyrobaculum aerophilum* showed the broadest substrate range of all characterized ACDs, reacting with acetyl-CoA, isobutyryl-

15 CoA (preferred substrate) and phenylacetyl-CoA (Brasen et al., *supra*). The enzymes from *A. fulgidus*, *H. marismortui* and *P. aerophilum* have all been cloned, functionally expressed, and characterized in *E. coli* (Musfeldt et al., *supra*; Brasen et al., *supra*). These genes/proteins are identified below in Table 3.

Table 3.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>AF1211</i>	NP_070039.1	11498810	<i>Archaeoglobus fulgidus</i> DSM 4304
<i>scs</i>	YP_135572.1	55377722	<i>Haloarcula marismortui</i> ATCC 43049
<i>PAE3250</i>	NP_560604.1	18313937	<i>Pyrobaculum aerophilum</i> str. IM2

20 Another candidate CoA synthetase is succinyl-CoA synthetase. The *sucCD* genes of *E. coli* form a succinyl-CoA synthetase complex which naturally catalyzes the formation of succinyl-CoA from succinate with the concomitant consumption of one ATP, a reaction which is reversible *in*

vivo (Buck et al., *Biochem.* 24:6245-6252 (1985)). These genes/proteins are identified below in Table 4.

Table 4.

<u>Protein</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>sucC</i>	NP_415256.1	16128703	<i>Escherichia coli</i>
<i>sucD</i>	AAC73823.1	1786949	<i>Escherichia coli</i>

Additional exemplary CoA-ligases include the rat dicarboxylate-CoA ligase for which the
 5 sequence is yet uncharacterized (Vamecq et al., *Biochemical Journal* 230:683-693 (1985)),
 either of the two characterized phenylacetate-CoA ligases from *P. chrysogenum* (Lamas-
 Maceiras et al., *Biochem. J.* 395:147-155 (2005); Wang et al., *Biochem Biophys Res Commun*
 360(2):453-458 (2007)), the phenylacetate-CoA ligase from *Pseudomonas putida* (Martinez-
 Blanco et al., *J. Biol. Chem.* 265:7084-7090 (1990)), and the 6-carboxyhexanoate-CoA ligase
 10 from *Bacillus subtilis* (Boweret al., *J. Bacteriol.* 178(14):4122-4130 (1996)). Additional
 candidate enzymes are acetoacetyl-CoA synthetases from *Mus musculus* (Hasegawa et al.,
Biochim Biophys Acta 1779:414-419 (2008)) and *Homo sapiens* (Ohgami et al., *Biochem*
Pharmacol 65:989-994 (2003)) which naturally catalyze the ATP-dependant conversion of
 acetoacetate into acetoacetyl-CoA. These genes/proteins are identified below in Table 5.

15

Table 5.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>phl</i>	CAJ15517.1	77019264	<i>Penicillium chrysogenum</i>
<i>phlB</i>	ABS19624.1	152002983	<i>Penicillium chrysogenum</i>
<i>paaF</i>	AAC24333.2	22711873	<i>Pseudomonas putida</i>
<i>bioW</i>	NP_390902.2	50812281	<i>Bacillus subtilis</i>
<i>AACS</i>	NP_084486.1	21313520	<i>Mus musculus</i>
<i>AACS</i>	NP_076417.2	31982927	<i>Homo sapiens</i>

Acetoacetyl-CoA hydrolase

Acetoacetyl-CoA can also be converted to acetoacetate by a CoA hydrolase. Acetoacetyl-CoA
 hydrolase enzyme candidates include acyl-CoA hydrolase, 3-hydroxyisobutyryl-CoA hydrolase,
 acetyl-CoA hydrolase, and dicarboxylic acid thioesterase. A short-chain acyl-CoA hydrolase in
 20 rat liver mitochondria was found to accept acetoacetyl-CoA as a substrate; however, the gene
 associated with this enzyme has not been identified to date (Svensson et al. *Eur. J. Biochem.*,
 239:526-531 (1996)).

3-Hydroxyisobutyryl-CoA hydrolase efficiently catalyzes the conversion of 3-
 hydroxyisobutyryl-CoA to 3-hydroxyisobutyrate during valine degradation (Shimomura et al., *J*

Biol Chem. 269:14248-14253 (1994)). Genes encoding this enzyme include *hibch* of *Rattus norvegicus* (Shimomura et al., *supra*; Shimomura et al., *Methods Enzymol.* 324:229-240 (2000)) and *Homo sapiens* (Shimomura et al., *supra*). The *H. sapiens* enzyme also accepts 3-hydroxybutyryl-CoA and 3-hydroxypropionyl-CoA as substrates (Shimomura et al., *supra*).

- 5 Candidate genes by sequence homology include *hibch* of *Saccharomyces cerevisiae* and *BC_2292* of *Bacillus cereus*. These genes/proteins are identified below in Table 6.

Table 6.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>hibch</i>	Q5XIE6.2	146324906	<i>Rattus norvegicus</i>
<i>hibch</i>	Q6NVY1.2	146324905	<i>Homo sapiens</i>
<i>hibch</i>	P28817.2	2506374	<i>Saccharomyces cerevisiae</i>
<i>BC_2292</i>	AP09256	29895975	<i>Bacillus cereus</i>

- Several eukaryotic acetyl-CoA hydrolases (EC 3.1.2.1) have broad substrate specificity and thus represent suitable candidate enzymes. For example, the enzyme from *Rattus norvegicus* brain (Robinson et al., *Res. Commun.* 71:959-965 (1976)) can react with butyryl-CoA, hexanoyl-CoA and malonyl-CoA. Though its sequence has not been reported, the enzyme from the mitochondrion of the pea leaf also has a broad substrate specificity, with demonstrated activity on acetyl-CoA, propionyl-CoA, butyryl-CoA, palmitoyl-CoA, oleoyl-CoA, succinyl-CoA, and crotonyl-CoA (Zeihner et al., *Plant. Physiol.* 94:20-27 (1990)). The acetyl-CoA hydrolase, *ACH1*, from *S. cerevisiae* represents another candidate hydrolase (Buu et al., *J. Biol. Chem.* 278:17203-17209 (2003)). These genes/proteins are identified below in Table 7.

Table 7.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>acot12</i>	NP_570103.1	18543355	<i>Rattus norvegicus</i>
<i>ACH1</i>	NP_009538	6319456	<i>Saccharomyces cerevisiae</i>

- Another candidate hydrolase is the human dicarboxylic acid thioesterase, *acot8*, which exhibits activity on glutaryl-CoA, adipyl-CoA, suberyl-CoA, sebacyl-CoA, and dodecanedioyl-CoA (Westin et al., *J Biol. Chem.* 280:38125-38132 (2005)) and the closest *E. coli* homolog, *tesB*, which can also hydrolyze a broad range of CoA thioesters (Naggert et al., *J Biol. Chem.* 266:11044-11050 (1991)). A similar enzyme has also been characterized in the rat liver (Deana et al., *Biochem. Int.* 26:767-773 (1992)). Other potential *E. coli* thioester hydrolases include the gene products of *tesA* (Bonner et al., *Chem.* 247:3123-3133 (1972)), *ybgC* (Kuznetsova et al., *FEMS Microbiol Rev* 29:263-279 (2005); and (Zhuang et al., *FEBS Lett.* 516:161-163 (2002)),

paal (Song et al., *J Biol. Chem.* 281:11028-11038 (2006)), and *ybdB* (Leduc et al., *J Bacteriol.* 189:7112-7126 (2007)). These genes/proteins are identified below in Table 8.

Table 8.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>tesB</i>	NP_414986	16128437	<i>Escherichia coli</i>
<i>acot8</i>	CAA15502	3191970	<i>Homo sapiens</i>
<i>acot8</i>	NP_570112	51036669	<i>Rattus norvegicus</i>
<i>tesA</i>	NP_415027	16128478	<i>Escherichia coli</i>
<i>ybgC</i>	NP_415264	16128711	<i>Escherichia coli</i>
<i>paal</i>	NP_415914	16129357	<i>Escherichia coli</i>
<i>ybdB</i>	NP_415129	16128580	<i>Escherichia coli</i>

- 5 Yet another candidate hydrolase is the glutaconate CoA-transferase from *Acidaminococcus fermentans*. This enzyme was transformed by site-directed mutagenesis into an acyl-CoA hydrolase with activity on glutaryl-CoA, acetyl-CoA and 3-butenoyl-CoA (Mack et al., *FEBS Lett.* 405:209-212 (1997)). This suggests that the enzymes encoding succinyl-CoA:3-ketoacid-CoA transferases and acetoacetyl-CoA:acetyl-CoA transferases may also serve as candidates for
- 10 this reaction step but would require certain mutations to change their function. These genes/proteins are identified below in Table 9.

Table 9.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>gctA</i>	CAA57199	559392	<i>Acidaminococcus fermentans</i>
<i>gctB</i>	CAA57200	559393	<i>Acidaminococcus fermentans</i>

Acetoacetate decarboxylase

- Acetoacetate decarboxylase converts acetoacetate into carbon dioxide and acetone. Exemplary
- 15 acetoacetate decarboxylase enzymes are encoded by the gene products of *adc* from *C. acetobutylicum* (Petersen and Bennett, *Appl Environ. Microbiol* 56:3491-3498 (1990)) and *adc* from *Clostridium saccharoperbutylacetonicum* (Kosaka et al., *Biosci. Biotechnol Biochem.* 71:58-68 (2007)). The enzyme from *C. beijerinckii* can be inferred from sequence similarity. These genes/proteins are identified below in Table 10.

Table 10.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Adc</i>	NP_149328.1	15004868	<i>Clostridium acetobutylicum</i>
<i>Adc</i>	AAP42566.1	31075386	<i>Clostridium saccharoperbutylacetonicum</i>
<i>Adc</i>	YP_001310906.1	150018652	<i>Clostridium beijerinckii</i>

Isopropanol dehydrogenase

- 5 The final step in the isopropanol synthesis pathway involves the reduction of acetone to isopropanol. Exemplary alcohol dehydrogenase enzymes capable of this transformation include *adh* from *C. beijerinckii* (Hanai et al., *Appl Environ Microbiol* 73:7814-7818 (2007); Jojima et al., *Appl Microbiol Biotechnol* 77:1219-1224 (2008)) and *adh* from *Thermoanaerobacter brockii* (Hanai et al., *Appl Environ Microbiol* 73:7814-7818 (2007); Peretz et al., *Anaerobe* 3:259-270 (1997)). Additional characterized enzymes include alcohol dehydrogenases from *Ralstonia eutropha* (formerly *Alcaligenes eutrophus*) (Steinbuchel and Schlegel et al., *Eur.J.Biochem.* 141:555-564 (1984)) and *Phytomonas* species (Uttaro and Opperdoes et al., *Mol.Biochem.Parasitol.* 85:213-219 (1997)). These genes/proteins are identified below in Table 11.

Table 11.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>sadh</i>	CAD36475	21615553	<i>Rhodococcus rubber</i>
<i>adhA</i>	AAC25556	3288810	<i>Pyrococcus furiosus</i>
<i>Adh</i>	P14941.1	113443	<i>Thermoanaerobacter brockii</i>
<i>Adh</i>	AAA23199.2	60592974	<i>Clostridium beijerinckii</i>

Production of n-propanol utilizing propionyl-CoA

- The pathways described herein for production of n-propanol utilize reduction of propionyl-CoA into propionaldehyde by a CoA-dependent aldehyde dehydrogenase that is then reduced further to form n-propanol (Figures 1-4). This conversion is carried out by two different enzymes: an aldehyde and alcohol dehydrogenase or in one step by a bifunctional aldehyde/alcohol dehydrogenase. Alternatively, propionyl CoA can be converted into propionyl phosphate and then transformed into propionaldehyde by an acyl phosphate reductase.

Propionaldehyde Dehydrogenase and Propanol Dehydrogenase

The conversion of propionyl-CoA to propanol is catalyzed by either a bifunctional enzyme that has both the CoA-dependent aldehyde dehydrogenase and the alcohol dehydrogenase activities or by two different enzymes with the aldehyde and alcohol dehydrogenase activities.

- 5 Exemplary two-step oxidoreductases that convert an acyl-CoA to alcohol include those that transform substrates such as acetyl-CoA to ethanol (e.g., *adhE* from *E. coli*) (Kessler, *FEBS Lett.* 281:59-63 (1991)) and butyryl-CoA to butanol (e.g. *adhE2* from *C. acetobutylicum*) (Fontaine et al., *J. Bacteriol.* 184:821-830 (2002)). In addition to reducing acetyl-CoA to ethanol, the enzyme encoded by *adhE* in *Leuconostoc mesenteroides* has been shown to oxidize
- 10 the branched chain compound isobutyraldehyde to isobutyryl-CoA (Kazahaya, *Microbiol.* 18:43-55 (1972); and Koo et al., *Biotechnol Lett.* 27:505-510 (2005)). These genes/proteins are identified below in Table 12.

Table 12.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>adhE</i>	NP_415757.1	16129202	<i>Escherichia coli</i>
<i>adhE2</i>	AAK09379.1	12958626	<i>Clostridium acetobutylicum</i>
<i>adhE</i>	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>

- 15 Another exemplary enzyme can convert malonyl-CoA to 3-HP. An NADPH-dependent enzyme with this activity has been characterized in *Chloroflexus aurantiacus* where it participates in the 3-hydroxypropionate cycle (Hugler, *J. Bacteriol.* 184:2404-2410 (2002); and Strauss, *Eur. J. Biochem.* 215:633-643 (1993)). This enzyme, with a mass of 300 kDa, is highly substrate-specific and shows little sequence similarity to other known oxidoreductases (Hugler, *J.*
- 20 *Bacteriol.* 184:2404-2410 (2002)). No enzymes in other organisms have been shown to catalyze this specific reaction; however there is bioinformatic evidence that other organisms may have similar pathways (Klatt, *Environ. Microbiol.* 9:2067-2078 (2007)). Enzyme candidates in other organisms including *Roseiflexus castenholzii*, *Erythrobacter sp. NAP1* and marine gamma proteobacterium HTCC2080 can be inferred by sequence similarity. These genes/proteins are
- 25 identified below in Table 13.

Table 13.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mcr</i>	AAS20429.1	42561982	<i>Chloroflexus aurantiacus</i>
<i>Rcas_2929</i>	YP_001433009.1	156742880	<i>Roseiflexus castenholzii</i>
<i>NAPI_02720</i>	ZP_01039179.1	85708113	<i>Erythrobacter sp. NAPI</i>
MGP2080_00535	ZP_01626393.1	119504313	marine gamma proteobacterium HTCC2080

Longer chain acyl-CoA molecules can be reduced by enzymes such as the jojoba (*Simmondsia chinensis*) FAR which encodes an alcohol-forming fatty acyl-CoA reductase. Its overexpression in *E. coli* resulted in FAR activity and the accumulation of fatty alcohol (Metz, *Plant Physiology* 122:635-644 (2000)). These genes/proteins are identified below in Table 14.

Table 14.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>FAR</i>	AAD38039.1	5020215	<i>Simmondsia chinensis</i>

Several acyl-CoA dehydrogenases are capable of reducing an acyl-CoA to its corresponding aldehyde. Exemplary genes that encode such enzymes include the *Acinetobacter calcoaceticus* *acrI* encoding a fatty acyl-CoA reductase, (Reiser, *Journal of Bacteriology* 179:2969-2975 (1997)) the *Acinetobacter sp. M-1* fatty acyl-CoA reductase, (Ishige et al., *Appl. Environ. Microbiol.* 68:1192-1195 (2002)) and a CoA- and NADP- dependent succinate semialdehyde dehydrogenase encoded by the *sucD* gene in *Clostridium kluyveri* (Sohling, *J. Bacteriol.* 178:871-880 (1996)). *SucD* of *P. gingivalis* is another succinate semialdehyde dehydrogenase (Takahashi, *J. Bacteriol* 182:4704-4710 (2000)). The enzyme acylating acetaldehyde dehydrogenase in *Pseudomonas sp.*, encoded by *bphG*, is yet another candidate as it has been demonstrated to oxidize and acylate acetaldehyde, propionaldehyde, butyraldehyde, isobutyraldehyde and formaldehyde (Powlowski, *J. Bacteriol.* 175:377-385 (1993)). In addition to reducing acetyl-CoA to ethanol, the enzyme encoded by *adhE* in *Leuconostoc mesenteroides* has been shown to oxidize the branched chain compound isobutyraldehyde to isobutyryl-CoA (Kazahaya, *J. Gen. Appl. Microbiol.* 18:43-55 (1972); and Koo et al., *Biotechnol Lett.* 27:505-510 (2005)). These genes/proteins are identified below in Table 15.

Table 15.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>acrI</i>	YP_047869.1	50086359	<i>Acinetobacter calcoaceticus</i>
<i>acrI</i>	AAC45217	1684886	<i>Acinetobacter baylyi</i>
<i>acrI</i>	BAB85476.1	18857901	<i>Acinetobacter sp. Strain M-1</i>
<i>sucD</i>	P38947.1	172046062	<i>Clostridium kluyveri</i>
<i>sucD</i>	NP_904963.1	34540484	<i>Porphyromonas gingivalis</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>bphG</i>	BAA03892.1	425213	<i>Pseudomonas sp</i>
<i>adhE</i>	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>

An additional enzyme type that converts an acyl-CoA to its corresponding aldehyde is malonyl-CoA reductase which transforms malonyl-CoA to malonic semialdehyde. Malonyl-CoA reductase is a key enzyme in autotrophic carbon fixation via the 3-hydroxypropionate cycle in thermoacidophilic archaeal bacteria (Berg, *Science* 318:1782-1786 (2007); and Thauer, *Science* 318:1732-1733 (2007)). The enzyme utilizes NADPH as a cofactor and has been characterized in *Metallosphaera* and *Sulfolobus spp.* (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006); and Hugler, *J. Bacteriol.* 184:2404-2410 (2002)). The enzyme is encoded by *Msed_0709* in *Metallosphaera sedula* (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006); and Berg, *Science* 318:1782-1786 (2007)). A gene encoding a malonyl-CoA reductase from *Sulfolobus tokodaii* was cloned and heterologously expressed in *E. coli* (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006)). This enzyme has also been shown to catalyze the conversion of methylmalonyl-CoA to its corresponding aldehyde (WO2007141208 (2007)). Although the aldehyde dehydrogenase functionality of these enzymes is similar to the bifunctional dehydrogenase from *Chloroflexus aurantiacus*, there is little sequence similarity. Both malonyl-CoA reductase enzyme candidates have high sequence similarity to aspartate-semialdehyde dehydrogenase, an enzyme catalyzing the reduction and concurrent dephosphorylation of aspartyl-4-phosphate to aspartate semialdehyde. Additional gene candidates can be found by sequence homology to proteins in other organisms including *Sulfolobus solfataricus* and *Sulfolobus acidocaldarius* and have been listed below. Yet another candidate for CoA-acylating aldehyde dehydrogenase is the *ald* gene from *Clostridium beijerinckii* (Toth, *Appl. Environ. Microbiol.* 65:4973-4980 (1999)). This enzyme has been reported to reduce acetyl-CoA and butyryl-CoA to their corresponding aldehydes. This gene is very similar to *eutE* that encodes acetaldehyde dehydrogenase of *Salmonella typhimurium* and *E. coli* (Toth, *Appl. Environ. Microbiol.* 65:4973-4980 (1999)). These genes/proteins are identified below in Table 16.

25

Table 16.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Msed_0709</i>	YP_001190808.1	146303492	<i>Metallosphaera sedula</i>
<i>mcr</i>	NP_378167.1	15922498	<i>Sulfolobus tokodaii</i>
<i>asd-2</i>	NP_343563.1	15898958	<i>Sulfolobus solfataricus</i>
<i>Saci_2370</i>	YP_256941.1	70608071	<i>Sulfolobus acidocaldarius</i>
<i>Ald</i>	AAT66436	49473535	<i>Clostridium beijerinckii</i>
<i>eutE</i>	AAA80209	687645	<i>Salmonella typhimurium</i>
<i>eutE</i>	P77445	2498347	<i>Escherichia coli</i>

Exemplary genes encoding enzymes that catalyze the conversion of an aldehyde to alcohol (i.e., alcohol dehydrogenase or equivalently aldehyde reductase) include *alrA* encoding a medium-chain alcohol dehydrogenase for C2-C14, (Tani, *Appl. Environ. Microbiol.* 66:5231-5235 (2000)) *ADH2* from *Saccharomyces cerevisiae*, (Atsumi, *Nature* 451:86-89 (2008)) *yqhD* from *E. coli* which has preference for molecules longer than C3, (Sulzenbacher et al., *Journal of Molecular Biology* 342:489-502 (2004)) and *bdh* I and *bdh* II from *C. acetobutylicum* which converts butyraldehyde into butanol (Walter, *Journal of Bacteriology* 174: 7149-7158 (1992)). The gene product of *yqhD* catalyzes the reduction of acetaldehyde, malondialdehyde, propionaldehyde, butyraldehyde, and acrolein using NADPH as the cofactor (Perez, *J. Biol. Chem.* 283:7346-7353 (2008)). *ADH1* from *Zymomonas mobilis* has been demonstrated to have activity on a number of aldehydes including formaldehyde, acetaldehyde, propionaldehyde, butyraldehyde, and acrolein (Kinoshita, *Appl. Microbiol. Biotechnol.* 22:249-254 (1985)). These genes/proteins are identified below in Table 17.

Table 17.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>alrA</i>	BAB12273.1	9967138	<i>Acinetobacter</i> sp. Strain M-1
<i>ADH2</i>	NP_014032.1	6323961	<i>Saccharomyces cerevisiae</i>
<i>yqhD</i>	NP_417484.1	16130909	<i>Escherichia coli</i>
<i>bdh</i> I	NP_349892.1	15896543	<i>Clostridium acetobutylicum</i>
<i>bdh</i> II	NP_349891.1	15896542	<i>Clostridium acetobutylicum</i>
<i>adhA</i>	YP_162971.1	56552132	<i>Zymomonas mobilis</i>

Enzymes exhibiting 3-hydroxybutyraldehyde reductase activity (EC 1.1.1.61) also fall into this category. Such enzymes have been characterized in *Ralstonia eutropha*, (Bravo J. *Forensic Sci.* 49:379-387 (2004)) *Clostridium kluyveri* (Wolff, *Protein Expr. Purif.* 6:206-212 (1995)) and *Arabidopsis thaliana* (Breitkreuz et al., *J. Biol. Chem.* 278:41552-41556 (2003)). Yet another gene candidate is the alcohol dehydrogenase *adhI* from *Geobacillus thermoglucosidasius* (Jeon et al., *J. Biotechnol.* 135:127-133 (2008)). These genes/proteins are identified below in Table 18.

Table 18.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>4hbd</i>	YP_726053.1	113867564	<i>Ralstonia eutropha</i> H16
<i>4hbd</i>	L21902.1	146348486	<i>Clostridium kluyveri</i> DSM 555
<i>4hbd</i>	Q94B07	75249805	<i>Arabidopsis thaliana</i>
<i>adhI</i>	AAR91477.1	40795502	<i>Geobacillus thermoglucosidasius</i> M10EXG

Another exemplary enzyme is 3-hydroxyisobutyrate dehydrogenase which catalyzes the reversible oxidation of 3-hydroxyisobutyrate to methylmalonate semialdehyde. This enzyme participates in valine, leucine and isoleucine degradation and has been identified in bacteria, eukaryotes, and mammals. The enzyme encoded by P84067 from *Thermus thermophilus* HB8 has been structurally characterized (Lokanath et al., *J Mol Biol* 352:905-917 (2005)). The reversibility of the human 3-hydroxyisobutyrate dehydrogenase was demonstrated using isotopically-labeled substrate (Manning, *Biochem J* 231:481-484 (1985)). Additional genes encoding this enzyme include *3hidh* in *Homo sapiens* (Hawes et al., *Methods Enzymol.* 324:218-228 (2000)) and *Oryctolagus cuniculus*, (Hawes et al., *Methods Enzymol.* 324:218-228 (2000); and Chowdhury, *Biosci. Biotechnol Biochem.* 60:2043-2047 (1996)) (*mmsb* in *Pseudomonas aeruginosa*, and *dhat* in *Pseudomonas putida* (Aberhart, *J Chem. Soc.* 6:1404-1406 (1979); Chowdhury, *Biosci. Biotechnol Biochem.* 60:2043-2047 (1996) and Chowdhury, *Biosci. Biotechnol Biochem.* 67:438-441 (2003)). These genes/proteins are identified below in Table 19.

Table 19.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>P84067</i>	P84067	75345323	<i>Thermus thermophilus</i>
<i>mmsb</i>	P28811.1	127211	<i>Pseudomonas aeruginosa</i>
<i>dhat</i>	Q59477.1	2842618	<i>Pseudomonas putida</i>
<i>3hidh</i>	P31937.2	12643395	<i>Homo sapiens</i>
<i>3hidh</i>	P32185.1	416872	<i>Oryctolagus cuniculus</i>

Propionyl-CoA:phosphate propanoyltransferase

The conversion of propanoyl-CoA to propanoyl phosphate can be catalyzed by a phosphate transferase. Among the phosphate acetyltransferases (EC 2.3.1.8), several enzymes including those from *Bacillus subtilis*, (Rado, *Biochem. Biophys. Acta* 321:114-125 (1973)) *Clostridium kluyveri*, (Stadtman, *Methods Enzymol* 1:596-599 (1955)) and *Thermotoga maritima* (Bock, *J Bacteriol.* 181:1861-1867 (1999)) have been shown to have activity on propionyl-CoA. Therefore, the genes coding for these phosphate acetyltransferases as well as *Escherichia coli* *pta* gene will be utilized to catalyze this step. These genes/proteins are identified below in Table 20.

Table 20.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pta</i>	P39646	730415	<i>Bacillus subtilis</i>
<i>pta</i>	A5N801	146346896	<i>Clostridium kluyveri</i>
<i>pta</i>	Q9X0L4	6685776	<i>Thermotoga maritima</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pta</i>	P0A9M8	71152910	<i>Escherichia coli</i> K12

Propionyl phosphate reductase

The conversion of propanoyl phosphate to propionaldehyde is catalyzed by the propionyl phosphate reductase. Even though such direct conversion has not been demonstrated yet, similar transformations were well documented including glyceraldehyde-3-phosphate dehydrogenase and aspartate-semialdehyde dehydrogenase. The following genes encoding glyceraldehyde-3-phosphate dehydrogenase and aspartate-semialdehyde dehydrogenase will be considered for catalyzing this step. These genes/proteins are identified below in Table 21.

Table 21.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>asd</i>	NP_417891	16131307	<i>Escherichia coli</i> K12
<i>gapA</i>	NP_785996	28379104	<i>Lactobacillus plantarum</i> WCFS1
<i>gapA</i>	NP_416293	71159358	<i>Escherichia coli</i> K12
<i>gapA</i>	NP_347346	15893997	<i>Clostridium acetobutylicum</i> ATCC 824
<i>gapN</i>	NP_350239	15896890	<i>Clostridium acetobutylicum</i> ATCC 824

10 Propionyl-CoA hydrolase

Propionyl-CoA can be converted to propionate by a CoA hydrolase, synthetase or transferase. The hydrolysis of propionyl-CoA to propionate occurs in organic acid degradation pathways that proceed through the intermediate 2-oxobutanoate. This reaction is catalyzed by acyl-CoA hydrolase enzymes (EC 3.1.2.18). Propionyl-CoA is the preferred substrate of the short chain acyl-CoA hydrolase found in rat liver mitochondria (Alexson et al., *Biochim Biophys. Acta.*, 1105(1):13-9 (1989)). This enzyme has been characterized but the sequence encoding the gene is not yet identified (Garras et al., *Biochim. Biophys. Acta.*, 1255:154-160 (1995)). Another enzyme exhibiting CoA hydrolase activity on propionyl-CoA is found in the mitochondrion of the pea leaf. Though its sequence has not been reported, this enzyme has a broad substrate specificity, with demonstrated activity on acetyl-CoA, propionyl-CoA, butyryl-CoA, palmitoyl-CoA, oleoyl-CoA, succinyl-CoA, and crotonyl-CoA (Zeiger et al., *Plant. Physiol.* 94:20-27 (1990)). Additional propionyl-CoA hydrolase candidates include 3-hydroxyisobutyryl-CoA hydrolase, acetyl-CoA hydrolase, and dicarboxylic acid thioesterase.

3-Hydroxyisobutyryl-CoA hydrolase efficiently catalyzes the conversion of 3-hydroxyisobutyryl-CoA to 3-hydroxyisobutyrate during valine degradation (Shimomura et al., *J Biol Chem.* 269:14248-14253 (1994)). Genes encoding this enzyme include *hibch* of *Rattus norvegicus* (Shimomura et al., *supra*; Shimomura et al., *Methods Enzymol.* 324:229-240 (2000)) and *Homo sapiens* (Shimomura et al., *supra*). The *H. sapiens* enzyme also accepts 3-hydroxybutyryl-CoA and 3-hydroxypropionyl-CoA as substrates (Shimomura et al., *supra*). Candidate genes by sequence homology include *hibch* of *Saccharomyces cerevisiae* and *BC_2292* of *Bacillus cereus*. These genes/proteins are identified below in Table 22.

Table 22.

Gene	GenBank ID	GI Number	Organism
<i>hibch</i>	Q5XIE6.2	146324906	<i>Rattus norvegicus</i>
<i>hibch</i>	Q6NVY1.2	146324905	<i>Homo sapiens</i>
<i>hibch</i>	P28817.2	2506374	<i>Saccharomyces cerevisiae</i>
<i>BC_2292</i>	AP09256	29895975	<i>Bacillus cereus</i>

Several eukaryotic acetyl-CoA hydrolases (EC 3.1.2.1) have broad substrate specificity and thus represent suitable candidate enzymes. For example, the enzyme from *Rattus norvegicus* brain (Robinson et al., *Res. Commun.* 71:959-965 (1976)) can react with butyryl-CoA, hexanoyl-CoA and malonyl-CoA. The acetyl-CoA hydrolase, *ACH1*, from *S. cerevisiae* represents another candidate hydrolase (Buu et al., *J. Biol. Chem.* 278:17203-17209 (2003)). These genes/proteins are identified below in Table 23.

Table 23.

Gene	GenBank ID	GI Number	Organism
<i>acot12</i>	NP_570103.1	18543355	<i>Rattus norvegicus</i>
<i>ACH1</i>	NP_009538	6319456	<i>Saccharomyces cerevisiae</i>

Another candidate hydrolase is the human dicarboxylic acid thioesterase, *acot8*, which exhibits activity on glutaryl-CoA, adipyl-CoA, suberyl-CoA, sebacyl-CoA, and dodecanedioyl-CoA (Westin et al., *J Biol. Chem.* 280:38125-38132 (2005)) and the closest *E. coli* homolog, *tesB*, which can also hydrolyze a broad range of CoA thioesters (Naggert et al., *J Biol. Chem.* 266:11044-11050 (1991)). A similar enzyme has also been characterized in the rat liver (Deana et al., *Biochem. Int.* 26:767-773 (1992)). Other potential *E. coli* thioester hydrolases include the gene products of *tesA* (Bonner et al., *Chem.* 247:3123-3133 (1972)), *ybgC* (Kuznetsova et al., *FEMS Microbiol Rev* 29:263-279 (2005); and (Zhuang et al., *FEBS Lett.* 516:161-163 (2002)),

paal (Song et al., *J Biol. Chem.* 281:11028-11038 (2006)), and *ybdB* (Leduc et al., *J Bacteriol.* 189:7112-7126 (2007)). These genes/proteins are identified below in Table 24.

Table 24.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>tesB</i>	NP_414986	16128437	<i>Escherichia coli</i>
<i>acot8</i>	CAA15502	3191970	<i>Homo sapiens</i>
<i>acot8</i>	NP_570112	51036669	<i>Rattus norvegicus</i>
<i>tesA</i>	NP_415027	16128478	<i>Escherichia coli</i>
<i>ybgC</i>	NP_415264	16128711	<i>Escherichia coli</i>
<i>paal</i>	NP_415914	16129357	<i>Escherichia coli</i>
<i>ybdB</i>	NP_415129	16128580	<i>Escherichia coli</i>

Yet another candidate hydrolase is the glutaconate CoA-transferase from *Acidaminococcus fermentans*. This enzyme was transformed by site-directed mutagenesis into an acyl-CoA hydrolase with activity on glutaryl-CoA, acetyl-CoA and 3-butenoyl-CoA (Mack et al., *FEBS Lett.* 405:209-212 (1997)). This suggests that the enzymes encoding succinyl-CoA:3-ketoacid-CoA transferases and acetoacetyl-CoA:acetyl-CoA transferases may also serve as candidates for this reaction step but would require certain mutations to change their function. These genes/proteins are identified below in Table 25.

Table 25.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>gctA</i>	CAA57199	559392	<i>Acidaminococcus fermentans</i>
<i>gctB</i>	CAA57200	559393	<i>Acidaminococcus fermentans</i>

Propionyl-CoA synthetase

A CoA synthetase can also catalyze the removal of the CoA moiety from propionyl-CoA. One candidate enzyme, ADP-forming acetyl-CoA synthetase (ACD, EC 6.2.1.13), couples the conversion of acyl-CoA esters to their corresponding acids with the concurrent synthesis of ATP. Several enzymes with broad substrate specificities have been described in the literature. ACD I from *Archaeoglobus fulgidus*, encoded by AF1211, was shown to operate on a variety of linear and branched-chain substrates including acetyl-CoA, propionyl-CoA, butyryl-CoA, acetate, propionate, butyrate, isobutyryate, isovalerate, succinate, fumarate, phenylacetate, indoleacetate (Musfeldt et al., *J Bacteriol* 184:636-644 (2002)). The enzyme from *Haloarcula marismortui* (annotated as a succinyl-CoA synthetase) accepts propionate, butyrate, and branched-chain acids (isovalerate and isobutyrate) as substrates, and was shown to operate in the forward and reverse directions (Brasen et al., *Arch Microbiol* 182:277-287 (2004)). The ACD encoded by PAE3250 from hyperthermophilic crenarchaeon *Pyrobaculum aerophilum* showed

the broadest substrate range of all characterized ACDs, reacting with acetyl-CoA, isobutyryl-CoA (preferred substrate) and phenylacetyl-CoA (Brasen et al., *supra*). The enzymes from *A. fulgidus*, *H. marismortui* and *P. aerophilum* have all been cloned, functionally expressed, and characterized in *E. coli* (Musfeldt et al., *supra*; Brasen et al., *supra*). These genes/proteins are

5 identified below in Table 26.

Table 26.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>AF1211</i>	NP_070039.1	11498810	<i>Archaeoglobus fulgidus</i> DSM 4304
<i>scs</i>	YP_135572.1	55377722	<i>Haloarcula marismortui</i> ATCC 43049
<i>PAE3250</i>	NP_560604.1	18313937	<i>Pyrobaculum aerophilum</i> str. IM2

Another candidate CoA synthetase is succinyl-CoA synthetase. The *sucCD* genes of *E. coli* form a succinyl-CoA synthetase complex which naturally catalyzes the formation of succinyl-CoA from succinate with the concomitant consumption of one ATP, a reaction which is reversible *in*

10 *vivo* (Buck et al., *Biochem. 24*:6245-6252 (1985)). These genes/proteins are identified below in Table 27.

Table 27.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>sucC</i>	NP_415256.1	16128703	<i>Escherichia coli</i>
<i>sucD</i>	AAC73823.1	1786949	<i>Escherichia coli</i>

Additional exemplary CoA-ligases include the rat dicarboxylate-CoA ligase for which the sequence is yet uncharacterized (Vamecq et al., *Biochemical Journal* 230:683-693 (1985)),

15 either of the two characterized phenylacetate-CoA ligases from *P. chrysogenum* (Lamas-Maceiras et al., *Biochem. J.* 395:147-155 (2005); Wang et al., *Biochem Biophys Res Commun* 360(2):453-458 (2007)), the phenylacetate-CoA ligase from *Pseudomonas putida* (Martinez-Blanco et al., *J. Biol. Chem.* 265:7084-7090 (1990)), and the 6-carboxyhexanoate-CoA ligase from *Bacillus subtilis* (Bower et al., *J. Bacteriol.* 178(14):4122-4130 (1996)). Additional

20 candidate enzymes are acetoacetyl-CoA synthetases from *Mus musculus* (Hasegawa et al., *Biochim Biophys Acta* 1779:414-419 (2008)) and *Homo sapiens* (Ohgami et al., *Biochem Pharmacol* 65:989-994 (2003)) which naturally catalyze the ATP-dependant conversion of acetoacetate into acetoacetyl-CoA. These genes/proteins are identified below in Table 28.

Table 28.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>phl</i>	CAJ15517.1	77019264	<i>Penicillium chrysogenum</i>
<i>phlB</i>	ABS19624.1	152002983	<i>Penicillium chrysogenum</i>
<i>paaF</i>	AAC24333.2	22711873	<i>Pseudomonas putida</i>
<i>bioW</i>	NP_390902.2	50812281	<i>Bacillus subtilis</i>
AACS	NP_084486.1	21313520	<i>Mus musculus</i>
AACS	NP_076417.2	31982927	<i>Homo sapiens</i>

Propionyl-CoA transferase

Propionyl-CoA transferase catalyzes the conversion of propionyl-CoA to propionate while transferring the CoA moiety to a CoA acceptor molecule. Many transferases have broad
5 specificity and thus may utilize CoA acceptors as diverse as acetate, succinate, propionate, butyrate, 2-methylacetoacetate, 3-ketohexanoate, 3-ketopentanoate, valerate, crotonate, 3-mercaptopropionate, propionate, vinylacetate, butyrate, among others.

Several genes have been identified that have propionyl-CoA transferase activity. The enzyme from *Roseburia* sp. A2-183 was shown to have butyryl-CoA:acetate:CoA transferase and
10 propionyl-CoA:acetate:CoA transferase activity (Charrier et al., *Microbiology* 152, 179-185 (2006)). Close homologs can be found in, for example, *Roseburia intestinalis* L1-82, *Roseburia inulinivorans* DSM 16841, *Eubacterium rectale* ATCC 33656. Another enzyme with propionyl-CoA transferase activity can be found in *Clostridium propionicum* (Selmer et al., *Eur J Biochem* 269, 372-380 (2002)). This enzyme can use acetate, (R)-lactate, (S)-lactate, acrylate, and
15 butyrate as the CoA acceptor (Selmer et al., *Eur J Biochem* 269, 372-380 (2002); Schweiger and Buckel, *FEBS Letters*, 171(1) 79-84 (1984)). Close homologs can be found in, for example, *Clostridium novyi* NT, *Clostridium beijerinckii* NCIMB 8052, and *Clostridium botulinum* C str. Eklund. *YgfH* encodes a propionyl CoA:succinate CoA transferase in *E. coli* (Haller et al., *Biochemistry*, 39(16) 4622-4629). Close homologs can be found in, for example, *Citrobacter youngae* ATCC 29220, *Salmonella enterica* subsp. *arizonae* serovar, and *Yersinia intermedia*
20 ATCC 29909. These genes/proteins are identified below in Table 29.

Table 29.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Ach1</i>	AAX19660.1	60396828	<i>Roseburia</i> sp. A2-183
<i>ROSINTL182_07121</i>	ZP_04743841.2	257413684	<i>Roseburia intestinalis</i> L1-82
<i>ROSEINA2194_03642</i>	ZP_03755203.1	225377982	<i>Roseburia inulinivorans</i> DSM 16841
<i>EUBREC_3075</i>	YP_002938937.1	238925420	<i>Eubacterium rectale</i> ATCC

			33656
<i>pct</i>	CAB77207.1	7242549	<i>Clostridium propionicum</i>
<i>NT01CX_2372</i>	YP_878445.1	118444712	<i>Clostridium novyi NT</i>
<i>Cbei_4543</i>	YP_001311608.1	150019354	<i>Clostridium beijerinckii NCIMB 8052</i>
<i>CBC_A0889</i>	ZP_02621218.1	168186583	<i>Clostridium botulinum C str. Eklund</i>
<i>ygfH</i>	NP_417395.1	16130821	<i>Escherichia coli str. K-12 substr. MG1655</i>
<i>CIT292_04485</i>	ZP_03838384.1	227334728	<i>Citrobacter youngae ATCC 29220</i>
<i>SARI_04582</i>	YP_001573497.1	161506385	<i>Salmonella enterica subsp. arizonae serovar</i>
<i>yinte0001_14430</i>	ZP_04635364.1	238791727	<i>Yersinia intermedia ATCC 29909</i>

An additional candidate enzyme is the two-unit enzyme encoded by *pcaI* and *pcaJ* in *Pseudomonas*, which has been shown to have 3-oxoadipyl-CoA/succinate transferase activity (Kaschabek et al., *supra*). Similar enzymes based on homology exist in *Acinetobacter* sp. ADP1 (Kowalchuk et al., *Gene* 146:23-30 (1994)) and *Streptomyces coelicolor*. Additional exemplary succinyl-CoA:3-oxoacid-CoA transferases are present in *Helicobacter pylori* (Corthesy-Theulaz et al., *J.Biol.Chem.* 272:25659-25667 (1997)) and *Bacillus subtilis* (Stols et al., *Protein.Expr.Purif.* 53:396-403 (2007)). These genes/proteins are identified below in Table 30.

Table 30.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pcaI</i>	AAN69545.1	24985644	<i>Pseudomonas putida</i>
<i>pcaJ</i>	NP_746082.1	26990657	<i>Pseudomonas putida</i>
<i>pcaI</i>	YP_046368.1	50084858	<i>Acinetobacter</i> sp. ADP1
<i>pcaJ</i>	AAC37147.1	141776	<i>Acinetobacter</i> sp. ADP1
<i>pcaI</i>	NP_630776.1	21224997	<i>Streptomyces coelicolor</i>
<i>pcaJ</i>	NP_630775.1	21224996	<i>Streptomyces coelicolor</i>
<i>HPAG1_0676</i>	YP_627417	108563101	<i>Helicobacter pylori</i>
<i>HPAG1_0677</i>	YP_627418	108563102	<i>Helicobacter pylori</i>
<i>ScoA</i>	NP_391778	16080950	<i>Bacillus subtilis</i>
<i>ScoB</i>	NP_391777	16080949	<i>Bacillus subtilis</i>

A CoA transferase that can utilize acetate as the CoA acceptor is acetoacetyl-CoA transferase, encoded by the *E. coli* *atoA* (alpha subunit) and *atoD* (beta subunit) genes (Vanderwinkel et al., *Biochem.Biophys.Res Commun.* 33:902-908 (1968); Korolev et al., *Acta Crystallogr.D Biol Crystallogr.* 58:2116-2121 (2002)). This enzyme has also been shown to transfer the CoA moiety to acetate from a variety of branched and linear acyl-CoA substrates, including isobutyrate (Matthies et al., *Appl Environ Microbiol* 58:1435-1439 (1992)), valerate (Vanderwinkel et al., *supra*) and butanoate (Vanderwinkel et al., *supra*). Similar enzymes exist

in *Corynebacterium glutamicum* ATCC 13032 (Duncan et al., *Appl Environ Microbiol* 68:5186-5190 (2002)), *Clostridium acetobutylicum* (Cary et al., *Appl Environ Microbiol* 56:1576-1583 (1990)), and *Clostridium saccharoperbutylacetonicum* (Kosaka et al., *Biosci. Biotechnol Biochem.* 71:58-68 (2007)). These genes/proteins are identified below in Table 31.

5

Table 31.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>atoA</i>	P76459.1	2492994	<i>Escherichia coli K12</i>
<i>atoD</i>	P76458.1	2492990	<i>Escherichia coli K12</i>
<i>actA</i>	YP_226809.1	62391407	<i>Corynebacterium glutamicum</i> ATCC 13032
<i>cg0592</i>	YP_224801.1	62389399	<i>Corynebacterium glutamicum</i> ATCC 13032
<i>ctfA</i>	NP_149326.1	15004866	<i>Clostridium acetobutylicum</i>
<i>ctfB</i>	NP_149327.1	15004867	<i>Clostridium acetobutylicum</i>
<i>ctfA</i>	AAP42564.1	31075384	<i>Clostridium saccharoperbutylacetonicum</i>
<i>ctfB</i>	AAP42565.1	31075385	<i>Clostridium saccharoperbutylacetonicum</i>

The above enzymes may also exhibit the desired activities on propionyl-CoA. Additional exemplary transferase candidates are catalyzed by the gene products of *cat1*, *cat2*, and *cat3* of *Clostridium kluyveri* which have been shown to exhibit succinyl-CoA, 4-hydroxybutyryl-CoA, and butyryl-CoA transferase activity, respectively (Seedorf et al., *supra*; Sohling et al., *Eur. J Biochem.* 212:121-127 (1993); Sohling et al., *J Bacteriol.* 178:871-880 (1996)). Similar CoA transferase activities are also present in *Trichomonas vaginalis* (van Grinsven et al., *J. Biol. Chem.* 283:1411-1418 (2008)) and *Trypanosoma brucei* (Riviere et al., *J. Biol. Chem.* 279:45337-45346 (2004)). These genes/proteins are identified below in Table 32.

10

Table 32.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>cat1</i>	P38946.1	729048	<i>Clostridium kluyveri</i>
<i>cat2</i>	P38942.2	172046066	<i>Clostridium kluyveri</i>
<i>cat3</i>	EDK35586.1	146349050	<i>Clostridium kluyveri</i>
<i>TVAG_395550</i>	XP_001330176	123975034	<i>Trichomonas vaginalis G3</i>
<i>Tb11.02.0290</i>	XP_828352	71754875	<i>Trypanosoma brucei</i>

15

The glutaconate-CoA-transferase (EC 2.8.3.12) enzyme from anaerobic bacterium *Acidaminococcus fermentans* reacts with diacid glutaconyl-CoA and 3-butenoyl-CoA (Mack et al., *FEBS Lett.* 405:209-212 (1997)). The genes encoding this enzyme are *gctA* and *gctB*. This enzyme has reduced but detectable activity with other CoA derivatives including glutaryl-CoA,

2-hydroxyglutaryl-CoA, adipyl-CoA and acrylyl-CoA (Buckel et al., *Eur.J.Biochem.* 118:315-321 (1981)). The enzyme has been cloned and expressed in *E. coli* (Mack et al., *Eur.J.Biochem.* 226:41-51 (1994)). These genes/proteins are identified below in Table 33.

Table 33.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>gctA</i>	CAA57199.1	559392	<i>Acidaminococcus fermentans</i>
<i>gctB</i>	CAA57200.1	559393	<i>Acidaminococcus fermentans</i>

5 Propionate kinase

Propionate is activated to propionyl-phosphate by an enzyme with propionate kinase activity. Butyrate kinase (EC 2.7.2.7) carries out the reversible conversion of butyryl-phosphate to butyrate during acidogenesis in *C. acetobutylicum* (Cary et al., *Appl. Environ. Microbiol* 56:1576-1583 (1990)). This enzyme is encoded by either of the two *buk* gene products (Huang et al., *J Mol. Microbiol Biotechnol* 2:33-38 (2000)). This enzyme was shown to accept propionate, isobutanoate and valerate as alternate substrates (Hartmanis, *J. Biol. Chem.*, 262(2):617-21 (1987)). Other butyrate kinase enzymes are found in *C. butyricum* and *C. tetanomorphum* (Twarog et al., *J Bacteriol.* 86:112-117 (1963)). These enzymes also accept propionate, isobutanoate and valerate as secondary substrates. Related enzyme isobutyrate kinase from *Thermotoga maritima* has also been expressed in *E. coli* and crystallized (Diao et al., *E. Biol. Crystallogr.* 59:1100-1102 (2003); and Diao et al., *J Bacteriol.* 191:2521-2529 (2009)). Aspartokinase catalyzes the ATP-dependent phosphorylation of aspartate and participates in the synthesis of several amino acids. The aspartokinase III enzyme in *E. coli*, encoded by *lysC*, has a broad substrate range and the catalytic residues involved in substrate specificity have been elucidated (Keng et al., *Arch. Biochem. Biophys.* 335:73-81 (1996)). Two additional kinases in *E. coli* are also good candidates: acetate kinase and gamma-glutamyl kinase. The *E. coli* acetate kinase, encoded by *ackA* (Skarstedt et al., *J. Biol. Chem.* 251:6775-6783 (1976)), phosphorylates propionate in addition to acetate (Hesslinger et al., *Mol. Microbiol* 27:477-492 (1998)). The *E. coli* gamma-glutamyl kinase, encoded by *proB* (Smith et al., *J. Bacteriol.* 157:545-551 (1984)), phosphorylates the gamma carbonic acid group of glutamate. These genes/proteins are identified below in Table 34.

Table 34.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>buk1</i>	NP_349675	15896326	<i>Clostridium acetobutylicum</i>
<i>buk2</i>	Q97III1	20137415	<i>Clostridium acetobutylicum</i>
<i>buk2</i>	Q9X278.1	6685256	<i>Thermotoga maritima</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>lysC</i>	NP_418448.1	16131850	<i>Escherichia coli</i>
<i>ackA</i>	NP_416799.1	16130231	<i>Escherichia coli</i>
<i>proB</i>	NP_414777.1	16128228	<i>Escherichia coli</i>

Propionate reductase

The reduction of propionate to propionic semialdehyde is catalyzed by a carboxylic acid reductase. Exemplary enzyme candidates for succinate reductase and 4-hydroxybutyrate reductase enzyme, described below, are also applicable here.

5

EXAMPLE II

Pathways for Production of Acetyl-CoA from Glucose

Further to Example I, the pathway for production of acetyl-CoA from glucose proceeds via phosphoenolpyruvate (PEP) (Figures 1-4). Glucose is converted into PEP by the native glycolysis pathway of the microbial organism. PEP is converted to pyruvate by pyruvate kinase and then to acetyl-CoA by pyruvate dehydrogenase or pyruvate ferredoxin oxidoreductase. Alternatively, pyruvate is converted to acetyl-CoA and formate by pyruvate formate lyase. Formate is then converted to carbon dioxide by a formate dehydrogenase that also produces NADH. The acetyl-CoA produced by these pathways are then utilized for production of isopropanol as described in Example I or utilized for production of both n-propanol and isopropanol as described in Example V below (Figure 3).

15

Pyruvate Dehydrogenase

The pyruvate dehydrogenase complex, catalyzing the conversion of pyruvate to acetyl-CoA, has been extensively studied. The *S. cerevisiae* complex consists of an E2 (*LAT1*) core that binds E1 (*PDA1*, *PDB1*), E3 (*LPDI*), and Protein X (*PDX1*) components (Pronk, *Yeast* 12:1607-1633 (1996)). In the *E. coli* enzyme, specific residues in the E1 component are responsible for substrate specificity (Bisswanger, *J. Biol Chem.* 256:815-822 (1981); Bremer, *Eur. J Biochem.* 8:535-540 (1969) and Gong et al., *J. Biol Chem.* 275:13645-13653 (2000)). Engineering efforts have improved the *E. coli* PDH enzyme activity under anaerobic conditions (Kim, *J. Bacteriol* 190:3851-3858 (2008); Kim, *Appl. Environ. Microbiol.* 73:1766-1771 (2007) and Zhou, *Biotechnol. Lett.* 30:335-342 (2008)). In contrast to the *E. coli* PDH, the *B. subtilis* complex is active and required for growth under anaerobic conditions (Nakano, *J. Bacteriol* 179:6749-6755 (1997)). The *Klebsiella pneumoniae* PDH, characterized during growth on glycerol, is also active under anaerobic conditions (Menzel, *J. Biotechnol.* 56:135-142 (1997)). Crystal

20

25

structures of the enzyme complex from bovine kidney (Zhou, *Proc. Natl. Acad. Sci. U.S.A.* 98:14802-14807 (2001)) and the E2 catalytic domain from *Azotobacter vinelandii* are available (Mattevi et al., *Science*. 255:1544-1550 (1992)). Some mammalian PDH enzymes complexes can react on alternate substrates such as 2-oxobutanoate (Paxton, *J Bacteriol.* 179:5684-5692 (1997)). These genes/proteins are identified below in Table 35.

Table 35.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>LAT1</i>	NP_014328	6324258	<i>Saccharomyces cerevisiae</i>
<i>PDA1</i>	NP_011105	37362644	<i>Saccharomyces cerevisiae</i>
<i>PDB1</i>	NP_009780	6319698	<i>Saccharomyces cerevisiae</i>
<i>LPD1</i>	NP_116635	14318501	<i>Saccharomyces cerevisiae</i>
<i>PDX1</i>	NP_011709	6321632	<i>Saccharomyces cerevisiae</i>
<i>aceE</i>	NP_414656.1	16128107	<i>Escherichia coli str. K12 substr. MG1655</i>
<i>aceF</i>	NP_414657.1	16128108	<i>Escherichia coli str. K12 substr. MG1655</i>
<i>lpd</i>	NP_414658.1	16128109	<i>Escherichia coli str. K12 substr. MG1655</i>
<i>pdhA</i>	P21881.1	3123238	<i>Bacillus subtilis</i>
<i>pdhB</i>	P21882.1	129068	<i>Bacillus subtilis</i>
<i>pdhC</i>	P21883.2	129054	<i>Bacillus subtilis</i>
<i>pdhD</i>	P21880.1	118672	<i>Bacillus subtilis</i>
<i>aceE</i>	YP_001333808.1	152968699	<i>Klebsiella pneumonia MGH78578</i>
<i>aceF</i>	YP_001333809.1	152968700	<i>Klebsiella pneumonia MGH78578</i>
<i>lpdA</i>	YP_001333810.1	152968701	<i>Klebsiella pneumonia MGH78578</i>
<i>Pdha1</i>	NP_001004072.2	124430510	<i>Rattus norvegicus</i>
<i>Pdha2</i>	NP_446446.1	16758900	<i>Rattus norvegicus</i>
<i>Dlat</i>	NP_112287.1	78365255	<i>Rattus norvegicus</i>
<i>Dld</i>	NP_955417.1	40786469	<i>Rattus norvegicus</i>

Pyruvate Ferredoxin Oxidoreductase

Pyruvate ferredoxin oxidoreductase (PFOR) catalyzes the oxidation of pyruvate to form acetyl-CoA. The PFOR from *Desulfovibrio africanus* has been cloned and expressed in *E. coli* resulting in an active recombinant enzyme that was stable for several days in the presence of oxygen (Pieulle, *J Bacteriol* 179:5684-5692 (1997)). Oxygen stability is relatively uncommon in PFORs and is believed to be conferred by a 60 residue extension in the polypeptide chain of the *D. africanus* enzyme. The *M. thermoacetica* PFOR is also well characterized (Menon, *Biochemistry* 36:8484-8494 (1997)) and was even shown to have high activity in the direction of pyruvate synthesis during autotrophic growth (Furdui, *J Biol Chem.* 275:28494-28499 (2000)). Further, *E. coli* possesses an uncharacterized open reading frame, *ydbK*, that encodes a protein that is 51% identical to the *M. thermoacetica* PFOR. Evidence for pyruvate oxidoreductase

activity in *E. coli* has been described (Blaschkowski, *Eur. J Biochem.* 123:563-569 (1982)).

Several additional PFOR enzymes are described in the following review (Ragsdale, *Chem. Rev.* 103:2333-2346 (2003)). Finally, flavodoxin reductases (e.g., *fqrB* from *Helicobacter pylori* or *Campylobacter jejuni*) (St Maurice et al., *J. Bacteriol.* 189:4764-4773 (2007)) or Rnf-type

- 5 proteins (Seedorf et al., *Proc. Natl. Acad. Sci. U.S.A.* 105:2128-2133 (2008); and Herrmann, *J. Bacteriol* 190:784-791 (2008)) provide a means to generate NADH or NADPH from the reduced ferredoxin generated by PFOR. These genes/proteins are identified below in Table 36.

Table 36.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>por</i>	CAA70873.1	1770208	<i>Desulfovibrio africanus</i>
<i>por</i>	YP_428946.1	83588937	<i>Moorella thermoacetica</i>
<i>ydbK</i>	NP_415896.1	16129339	<i>Escherichia coli</i>
<i>fqrB</i>	NP_207955.1	15645778	<i>Helicobacter pylori</i>
<i>fqrB</i>	YP_001482096.1	157414840	<i>Campylobacter jejuni</i>
<i>RnfC</i>	EDK33306.1	146346770	<i>Clostridium kluyveri</i>
<i>RnfD</i>	EDK33307.1	146346771	<i>Clostridium kluyveri</i>
<i>RnfG</i>	EDK33308.1	146346772	<i>Clostridium kluyveri</i>
<i>RnfE</i>	EDK33309.1	146346773	<i>Clostridium kluyveri</i>
<i>RnfA</i>	EDK33310.1	146346774	<i>Clostridium kluyveri</i>
<i>RnfB</i>	EDK33311.1	146346775	<i>Clostridium kluyveri</i>

10 Pyruvate Formate Lyase

Pyruvate formate lyase is an enzyme that catalyzes the conversion of pyruvate and CoA into acetyl-CoA and formate. Pyruvate formate lyase is a common enzyme in prokaryotic organisms that is used to help modulate anaerobic redox balance. Exemplary enzymes can be found in *Escherichia coli* (Knappe, *FEMS. Microbiol Rev.* 6:383-398 (1990)), *Lactococcus lactis*

- 15 (Melchiorson, *Appl Microbiol Biotechnol* 58:338-344(2002)), and *Streptococcus mutans*. (Takahashi-Abbe, *Oral. Microbiol Immunol.* 18:293-297 (2003)). A mitochondrial pyruvate formate lyase has also been identified in the eukaryote, *Chlamydomonas reinhardtii*. (Hemschemeier, *Eukaryot. Cell* 7:518-526 (2008); and Atteia, *J. Biol. Chem.* 281:9909-9918 (2008)). These genes/proteins are identified below in Table 37.

Table 37.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pflB</i>	NP_415423	16128870	<i>Escherichia coli</i>
<i>pfl</i>	CAA03993	2407931	<i>Lactococcus lactis</i>
<i>pfl</i>	BAA09085	1129082	<i>Streptococcus mutans</i>
<i>PFLI</i>	EDP09457	158283707	<i>Chlamydomonas reinhardtii</i>

Formate Hydrogen Lyase

A formate hydrogen lyase enzyme can be employed to convert formate to carbon dioxide and hydrogen. An exemplary formate hydrogen lyase enzyme can be found in *Escherichia coli*. The *E. coli* formate hydrogen lyase consists of hydrogenase 3 and formate dehydrogenase-H

- 5 (Maeda, *Appl Microbiol Biotechnol* 77:879-890 (2007)). It is activated by the gene product of *fhlA* (Maeda, *Appl Microbiol Biotechnol* 77:879-890 (2007)). The addition of the trace elements, selenium, nickel and molybdenum, to a fermentation broth has been shown to enhance formate hydrogen lyase activity (Soini, *Microb. Cell Fact.* 7:26 (2008)). These genes/proteins are identified below in Table 38.

10

Table 38.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
Hydrogenase 3:			
<i>hycD</i>	NP_417202	16130629	<i>Escherichia coli</i>
<i>hycC</i>	NP_417203	16130630	<i>Escherichia coli</i>
<i>hycF</i>	NP_417200	16130627	<i>Escherichia coli</i>
<i>hycG</i>	NP_417199	16130626	<i>Escherichia coli</i>
<i>hycB</i>	NP_417204	16130631	<i>Escherichia coli</i>
<i>hycE</i>	NP_417201	16130628	<i>Escherichia coli</i>
Formate dehydrogenase-H:			
<i>fdhF</i>	NP_418503	16131905	<i>Escherichia coli</i>
Activator:			
<i>fhlA</i>	NP_417211	16130638	<i>Escherichia coli</i>

A formate hydrogen lyase enzyme also exists in the hyperthermophilic archaeon, *Thermococcus litoralis* (Takacs et al., *BMC. Microbiol* 8:88 (2008)). These genes/proteins are identified below in Table 39.

Table 39.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mhyC</i>	ABW05543	157954626	<i>Thermococcus litoralis</i>
<i>mhyD</i>	ABW05544	157954627	<i>Thermococcus litoralis</i>
<i>mhyE</i>	ABW05545	157954628	<i>Thermococcus litoralis</i>
<i>myhF</i>	ABW05546	157954629	<i>Thermococcus litoralis</i>
<i>myhG</i>	ABW05547	157954630	<i>Thermococcus litoralis</i>
<i>myhH</i>	ABW05548	157954631	<i>Thermococcus litoralis</i>
<i>fdhA</i>	AAB94932	2746736	<i>Thermococcus litoralis</i>
<i>fdhB</i>	AAB94931	157954625	<i>Thermococcus litoralis</i>

- 15 Additional formate hydrogen lyase systems have been found in *Salmonella typhimurium*, *Klebsiella pneumoniae*, *Rhodospirillum rubrum*, *Methanobacterium formicicum* (Vardar-Schara, *Microbial Biotechnology* 1:107-125 (2008)).

Formate Dehydrogenase

Formate dehydrogenase activity is present in both *E. coli* and *Saccharomyces cerevisiae* among other organisms. *S. cerevisiae* contains two formate dehydrogenases, *FDH1* and *FDH2*, that catalyze the oxidation of formate to CO₂. (Overkamp et al., *Yeast* 19:509-520 (2002)) In *Moorella thermoacetica*, the loci, *Moth_2312* and *Moth_2313*, are actually one gene that is responsible for encoding the alpha subunit of formate dehydrogenase while the beta subunit is encoded by *Moth_2314* (Pierce et al., *Environ. Microbiol* (2008); Andreesen, *J. Bacteriol.* 116:867-873 (1973); Li, *J. Bacteriol* 92:405-412 (1966) and Yamamoto, *J. Biol. Chem.* 258:1826-1832 (1983)) Another set of genes encoding formate dehydrogenase activity is encoded by *Sfum_2703* through *Sfum_2706* in *Syntrophobacter fumaroxidans* (Reda, *Proc. Natl. Acad. Sci. U.S.A.* 105:10654-10658 (2008); and de Bok et al., *Eur. J. Biochem.* 270:2476-2485 (2003)). Similar to their *M. thermoacetica* counterparts, *Sfum_2705* and *Sfum_2706* are actually one gene. *E. coli* contains multiple formate dehydrogenases. These genes/proteins are identified below in Table 40.

Table 40.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>FDH1</i>	NP_015033	6324964	<i>Saccharomyces cerevisiae</i>
<i>FDH2</i>	Q08987	88909613	<i>Saccharomyces cerevisiae</i>
<i>Moth_2312</i>	YP_431142	148283121	<i>Moorella thermoacetica</i>
<i>Moth_2313</i>	YP_431143	83591134	<i>Moorella thermoacetica</i>
<i>Moth_2314</i>	YP_431144	83591135	<i>Moorella thermoacetica</i>
<i>Sfum_2703</i>	YP_846816.1	116750129	<i>Syntrophobacter fumaroxidans</i>
<i>Sfum_2704</i>	YP_846817.1	116750130	<i>Syntrophobacter fumaroxidans</i>
<i>Sfum_2705</i>	YP_846818.1	116750131	<i>Syntrophobacter fumaroxidans</i>
<i>Sfum_2706</i>	YP_846819.1	116750132	<i>Syntrophobacter fumaroxidans</i>
<i>fdnG, H, I</i>	NP_415991-993.1	16129433 16129434 16129435	<i>Escherichia coli</i>
<i>fdoG, H, I</i>	NP_418330,29,28.1	16131734 16131733 16131732	<i>Escherichia coli</i>

EXAMPLE III

Pathways for Production of Propionyl-CoA from Glucose Utilizing the Reductive TCA Cycle

Further to Examples I and II, the pathway for production of propionyl-CoA proceeds via oxaloacetate (Figure 1). PEP is converted into oxaloacetate either via PEP carboxykinase or PEP carboxylase. Alternatively, PEP is converted first to pyruvate by pyruvate kinase and then

to oxaloacetate by methylmalonyl-CoA carboxytransferase or pyruvate carboxylase.

Oxaloacetate is converted to propionyl-CoA by means of the reductive TCA cycle, a methylmutase, a decarboxylase, an epimerase and a decarboxylase.

PEP Carboxykinase

- 5 Although the net conversion of phosphoenolpyruvate to oxaloacetate is redox-neutral, the mechanism of this conversion is important to the overall energetics of the co- production pathway. The most desirable enzyme for the conversion of PEP to oxaloacetate is PEP carboxykinase which simultaneously forms an ATP while carboxylating PEP. In most organisms, however, PEP carboxykinase serves a gluconeogenic function and converts
- 10 oxaloacetate to PEP at the expense of one ATP. *S. cerevisiae* is one such organism whose native PEP carboxykinase, *PCK1*, serves a gluconeogenic role (Valdes-Hevia, *FEBS. Lett.* 258:313-316 (1989)). *E. coli* is another such organism, as the role of PEP carboxykinase in producing oxaloacetate is believed to be minor when compared to PEP carboxylase, which does not form ATP, possibly due to the higher K_m for bicarbonate of PEP carboxykinase (Kim, *Appl Environ*
- 15 *Microbiol* 70:1238-1241 (2004)). Nevertheless, activity of the native *E. coli* PEP carboxykinase from PEP towards oxaloacetate has been recently demonstrated in *ppc* mutants of *E. coli* K-12 (Kwon, *Journal of Microbiology and Biotechnology* 16:1448-1452 (2006)). These strains exhibited no growth defects and had increased succinate production at high NaHCO_3 concentrations. In some organisms, particularly rumen bacteria, PEP carboxykinase is quite
- 20 efficient in producing oxaloacetate from PEP and generating ATP. Examples of PEP carboxykinase genes that have been cloned into *E. coli* include those from *Mannheimia succiniciproducens* (Lee, *Biotechnol. Bioprocess Eng.* 7:95-99 (2002)), *Anaerobiospirillum succiniciproducens* (Laivenieks, *Appl Environ Microbiol* 63:2273-2280 (1997)), and *Actinobacillus succinogenes* (Kim, *Appl Environ Microbiol* 70:1238-1241 (2004)). Internal
- 25 experiments have also found that the PEP carboxykinase enzyme encoded by *Haemophilus influenza* is highly efficient at forming oxaloacetate from PEP. These genes/proteins are identified below in Table 41.

Table 41.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>PCK1</i>	NP_013023	6322950	<i>Saccharomyces cerevisiae</i>
<i>pck</i>	NP_417862.1	16131280	<i>Escherichia coli</i>
<i>pckA</i>	YP_089485.1	52426348	<i>Mannheimia succiniciproducens</i>
<i>pckA</i>	O09460.1	3122621	<i>Anaerobiospirillum succiniciproducens</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pckA</i>	Q6W6X5	75440571	<i>Actinobacillus succinogenes</i>
<i>pckA</i>	P43923.1	1172573	<i>Haemophilus influenza</i>

These sequences and sequences for subsequent enzymes listed in this report can be used to identify homologue proteins in GenBank or other databases through sequence similarity searches (e.g. BLASTp). The resulting homologue proteins and their corresponding gene sequences provide additional DNA sequences for transformation into the host organism of choice.

PEP Carboxylase

PEP carboxylase represents an alternative enzyme for the formation of oxaloacetate from PEP. Since the enzyme does not generate ATP upon decarboxylating oxaloacetate, its utilization decreases the maximum ATP yield of the production pathway and represents a less favorable alternative for converting oxaloacetate to PEP. Nevertheless, the maximum theoretical C3 alcohols yield of 1.33 mol/mol will remain unchanged if PEP carboxylase is utilized to convert PEP to oxaloacetate. *S. cerevisiae* does not naturally encode a PEP carboxylase, but exemplary organisms that possess genes that encode PEP carboxylase include *E. coli* (Kai, Arch. Biochem. Biophys. 414:170-179 (2003)), *Methylobacterium extorquens* AM1 (Arps, J. Bacteriol. 175:3776-3783 (1993)), and *Corynebacterium glutamicum* (Eikmanns, Mol. Gen. Genet. 218:330-339 (1989)). These genes/proteins are identified below in Table 42.

Table 42.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ppc</i>	NP_418391	16131794	<i>Escherichia coli</i>
<i>ppcA</i>	AAB58883	28572162	<i>Methylobacterium extorquens</i>
<i>ppc</i>	ABB53270	80973080	<i>Corynebacterium glutamicum</i>

Pyruvate Kinase and Methylmalonyl-CoA Carboxyltransferase

An additional energetically efficient route to oxaloacetate from PEP requires two enzymatic activities: pyruvate kinase and methylmalonyl-CoA carboxytransferase. Pyruvate kinase catalyzes the ATP-generating conversion of PEP to pyruvate and is encoded by the *PYK1* (Burke, J. Biol. Chem. 258:2193-2201 (1983)) and *PYK2* (Boles et al., J. Bacteriol. 179:2987-2993 (1997)) genes in *S. cerevisiae*. In *E. coli*, this activity is catalyzed by the gene product of *pykF* and *pykA*. Methylmalonyl-CoA carboxytransferase catalyzes the conversion of pyruvate to oxaloacetate. Importantly, this reaction also simultaneously catalyzes the conversion of (S)-methylmalonyl-CoA to propionyl-CoA (see Figures 1 and 2). An exemplary methylmalonyl-CoA carboxytransferase which is comprised of 1.3S, 5S, and 12S subunits can be found in

Propionibacterium freudenreichii (Thornton et al., *J. Bacteriol* 175:5301-5308 (1993)). These genes/proteins are identified below in Table 43.

Table 43.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>PYK1</i>	NP_009362	6319279	<i>Saccharomyces cerevisiae</i>
<i>PYK2</i>	NP_014992	6324923	<i>Saccharomyces cerevisiae</i>
<i>pykF</i>	NP_416191.1	16129632	<i>Escherichia coli</i>
<i>pykA</i>	NP_416368.1	16129807	<i>Escherichia coli</i>
1.3S subunit	P02904	114847	<i>Propionibacterium freudenreichii</i>
5S subunit	Q70AC7	62901478	<i>Propionibacterium freudenreichii</i>
12S subunit	Q8GBW6	62901481	<i>Propionibacterium freudenreichii</i>

Pyruvate Kinase and Pyruvate Carboxylase

- 5 A combination of enzymes can convert PEP to oxaloacetate with a stoichiometry identical to that of PEP carboxylase. These enzymes are encoded by pyruvate kinase, *PYK1* (Burke, *J. Biol. Chem.* 258:2193-2201 (1983)) or *PYK2* (Boles et al., *J. Bacteriol*, 179:2987-2993 (1997)) and pyruvate carboxylase, *PYC1* (Walker, *Biochem. Biophys. Res. Commun.* 176:1210-1217 (1991)) or *PYC2* (Walker, *Biochem. Biophys. Res. Commun.* 176:1210-1217 (1991)). The latter
- 10 genes/proteins are identified below in Table 44.

Table 44.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>PYC1</i>	NP_011453	6321376	<i>Saccharomyces cerevisiae</i>
<i>PYC2</i>	NP_009777	6319695	<i>Saccharomyces cerevisiae</i>
<i>Pyc</i>	YP_890857.1	118470447	<i>Mycobacterium smegmatis</i>

Malate Dehydrogenase, Fumarase, Fumarate Reductase

- Oxaloacetate can be converted to succinate by malate dehydrogenase, fumarase and fumarate reductase when the TCA cycle is operating in the reductive cycle. *S. cerevisiae* possesses three
- 15 copies of malate dehydrogenase, *MDH1* (McAlister-Henn, *J. Bacteriol* 169:5157-5166 (1987)) *MDH2* (Minard, *Mol. Cell. Biol.* 11:370-380 (1991); and Gibson, *J. Biol. Chem.* 278:25628-25636 (2003)), and *MDH3* (Steffan, *J. Biol. Chem.* 267:24708-24715 (1992)), which localize to the mitochondrion, cytosol, and peroxisome, respectively. *S. cerevisiae* contains one copy of a fumarase-encoding gene, *FUM1*, whose product localizes to both the cytosol and mitochondrion
- 20 (Sass, *J. Biol. Chem.* 278:45109-45116 (2003)). Fumarate reductase is encoded by two soluble enzymes, FRDS1 (Enomoto, *DNA. Res.* 3:263-267 (1996)) and FRDS2 (Muratsubaki, *Arch. Biochem. Biophys.* 352:175-181 (1998)), which localize to the cytosol and promitochondrion,

respectively, and are required for anaerobic growth on glucose (Arikawa, *Microbiol Lett.* 165:111-116 (1998)). *E. coli* is known to have an active malate dehydrogenase. It has three fumarases encoded by *fumA*, *B* and *C*, each one of which is active under different conditions of oxygen availability. The fumarate reductase in *E. coli* is composed of four subunits. These

5 genes/proteins are identified below in Table 45.

Table 45.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>MDH1</i>	NP_012838	6322765	<i>Saccharomyces cerevisiae</i>
<i>MDH2</i>	NP_014515	116006499	<i>Saccharomyces cerevisiae</i>
<i>MDH3</i>	NP_010205	6320125	<i>Saccharomyces cerevisiae</i>
<i>FUM1</i>	NP_015061	6324993	<i>Saccharomyces cerevisiae</i>
<i>FRDS1</i>	P32614	418423	<i>Saccharomyces cerevisiae</i>
<i>FRDS2</i>	NP_012585	6322511	<i>Saccharomyces cerevisiae</i>
<i>frdA</i>	NP_418578.1	16131979	<i>Escherichia coli</i>
<i>frdB</i>	NP_418577.1	16131978	<i>Escherichia coli</i>
<i>frdC</i>	NP_418576.1	16131977	<i>Escherichia coli</i>
<i>frdD</i>	NP_418475.1	16131877	<i>Escherichia coli</i>
<i>Mdh</i>	NP_417703.1	16131126	<i>Escherichia coli</i>
<i>FumA</i>	NP_416129.1	16129570	<i>Escherichia coli</i>
<i>FumB</i>	NP_418546.1	16131948	<i>Escherichia coli</i>
<i>FumC</i>	NP_416128.1	16129569	<i>Escherichia coli</i>

Succinyl-CoA Transferase

Succinyl-CoA transferase catalyzes the conversion of succinyl-CoA to succinate while transferring the CoA moiety to a CoA acceptor molecule. Many transferases have broad
 10 specificity and thus may utilize CoA acceptors as diverse as acetate, succinate, propionate, butyrate, 2-methylacetoacetate, 3-ketohexanoate, 3-ketopentanoate, valerate, crotonate, 3-mercaptopropionate, propionate, vinylacetate, butyrate, among others.

The conversion of succinate to succinyl-CoA is ideally carried by a transferase which does not require the direct consumption of an ATP or GTP. This type of reaction is common in a number
 15 of organisms. Perhaps the top candidate enzyme for this reaction step is succinyl-CoA:3-ketoacid-CoA transferase. This enzyme converts succinate to succinyl-CoA while converting a 3-ketoacyl-CoA to a 3-ketoacid. Exemplary succinyl-CoA:3-ketoacid-CoA transferases are present in *Helicobacter pylori* (Corthesy-Theulaz et al., *J. Biol. Chem.* 272:25659-25667 (1997)), *Bacillus subtilis* (Stols et al., *Protein. Expr. Purif.* 53:396-403 (2007)), and *Homo sapiens* (Fukao et al., *Genomics*, 68:144-151 (2000); and Tanaka, *Mol. Hum. Reprod.* 8:16-23
 20 (2002)). These genes/proteins are identified below in Table 46.

Table 46.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>HPAG1_0676</i>	YP_627417	108563101	<i>Helicobacter pylori</i>
<i>HPAG1_0677</i>	YP_627418	108563102	<i>Helicobacter pylori</i>
<i>ScoA</i>	NP_391778	16080950	<i>Bacillus subtilis</i>
<i>ScoB</i>	NP_391777	16080949	<i>Bacillus subtilis</i>
<i>OXCT1</i>	NP_000427	4557817	<i>Homo sapiens</i>
<i>OXCT2</i>	NP_071403	11545841	<i>Homo sapiens</i>

The conversion of succinate to succinyl-CoA can also be catalyzed by succinyl-CoA: Acetyl-CoA transferase. The gene product of *catI* of *Clostridium kluyveri* has been shown to exhibit succinyl-CoA: acetyl-CoA transferase activity (Sohling, *J Bacteriol.* 178:871-880 (1996)). In addition, the activity is present in *Trichomonas vaginalis* (van Grinsven et al., *J. Biol. Chem.* 283:1411-1418 (2008)) and *Trypanosoma brucei* (Riviere et al., *J. Biol. Chem.* 279:45337-45346 (2004)). These genes/proteins are identified below in Table 47.

Table 47.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>catI</i>	P38946.1	729048	<i>Clostridium kluyveri</i>
<i>TVAG_395550</i>	XP_001330176	123975034	<i>Trichomonas vaginalis</i> G3
<i>Tb11.02.0290</i>	XP_828352	71754875	<i>Trypanosoma brucei</i>

Yet another possible CoA acceptor is benzylsuccinate. Succinyl-CoA:(R)-Benzylsuccinate CoA-Transferase functions as part of an anaerobic degradation pathway for toluene in organisms such as *Thauera aromatica* (Leutwein and Heider, *J. Bact.* 183(14) 4288-4295 (2001)). Homologs can be found in *Azoarcus* sp. T, *Aromatoleum aromaticum* EbN1, and *Geobacter metallireducens* GS-15. These genes/proteins are identified below in Table 48.

Table 48.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>bbsE</i>	AAF89840	9622535	<i>Thauera aromatica</i>
<i>bbsF</i>	AAF89841	9622536	<i>Thauera aromatica</i>
<i>bbsE</i>	AAU45405.1	52421824	<i>Azoarcus</i> sp. T
<i>bbsF</i>	AAU45406.1	52421825	<i>Azoarcus</i> sp. T
<i>bbsE</i>	YP_158075.1	56476486	<i>Aromatoleum aromaticum</i> EbN1
<i>bbsF</i>	YP_158074.1	56476485	<i>Aromatoleum aromaticum</i> EbN1
<i>Gmet_1521</i>	YP_384480.1	78222733	<i>Geobacter metallireducens</i> GS-15
<i>Gmet_1522</i>	YP_384481.1	78222734	<i>Geobacter metallireducens</i> GS-15

Finally, *ygfH* encodes a propionyl CoA:succinate CoA transferase in *E. coli* (Haller et al., *Biochemistry*, 39(16) 4622-4629). Close homologs can be found in, for example, *Citrobacter*

youngae ATCC 29220, *Salmonella enterica* subsp. *arizonae* serovar, and *Yersinia intermedia* ATCC 29909. These genes/proteins are identified below in Table 49.

Table 49.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ygfH</i>	NP_417395.1	16130821	<i>Escherichia coli</i> str. K-12 substr. MG1655
<i>CIT292_04485</i>	ZP_03838384.1	227334728	<i>Citrobacter youngae</i> ATCC 29220
<i>SARI_04582</i>	YP_001573497.1	161506385	<i>Salmonella enterica</i> subsp. <i>arizonae</i> serovar
<i>yinte0001_14430</i>	ZP_04635364.1	238791727	<i>Yersinia intermedia</i> ATCC 29909

Succinyl-CoA Synthetase

- 5 The product of the *LSC1* and *LSC2* genes of *S. cerevisiae* and the *sucC* and *sucD* genes of *E. coli* naturally form a succinyl-CoA synthetase complex that catalyzes the formation of succinyl-CoA from succinate with the concomitant consumption of one ATP, a reaction which is reversible *in vivo* (Przybyla-Zawilask et al., *Eur. J. Biochem.* 258(2):736-743 (1998) and Buck et al., *J. Gen. Microbiol.* 132(6):1753-1762 (1986)). These genes/proteins are identified below in
- 10 Table 50.

Table 50.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>LSC1</i>	NP_014785	6324716	<i>Saccharomyces cerevisiae</i>
<i>LSC2</i>	NP_011760	6321683	<i>Saccharomyces cerevisiae</i>
<i>sucC</i>	NP_415256.1	16128703	<i>Escherichia coli</i>
<i>sucD</i>	AAC73823.1	1786949	<i>Escherichia coli</i>

Methylmalonyl-CoA Mutase

- Succinyl-CoA can be converted into (R)-methylmalonyl-CoA by methylmalonyl-CoA mutase (MCM). In *E. coli*, the reversible adenosylcobalamin-dependant mutase participates in a three-
- 15 step pathway leading to the conversion of succinate to propionate (Haller, *Biochemistry* 39:4622-9 (2000)). MCM is encoded by genes *scpA* in *Escherichia coli* (Haller, *Biochemistry* 39: 4622-4629 (2000); and Bobik, *Anal. Bioanal. Chem.* 375:344-349 (2003)) and *mutA* in *Homo sapiens* (Padovani, *Biochemistry* 45:9300-9306 (2006)). In several other organisms
- 20 MCM contains alpha and beta subunits and is encoded by two genes. Exemplary gene candidates encoding the two-subunit protein are *Propionibacterium fredenreichii* sp. *shermani* *mutA* and *mutB* (Korotkova, *J Biol Chem.* 279:13652-13658 (2004)) and *Methylobacterium*

extorquens mcmA and *mcmB* (Korotkova, *J Biol Chem.* 279:13652-13658 (2004)). These genes/proteins are identified below in Table 51.

Table 51.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>scpA</i>	NP_417392.1	16130818	<i>Escherichia coli K12</i>
<i>mutA</i>	P22033.3	67469281	<i>Homo sapiens</i>
<i>mutA</i>	P11652.3	127549	<i>Propionibacterium fredenreichii sp. shermanii</i>
<i>mutB</i>	P11653.3	127550	<i>Propionibacterium fredenreichii sp. shermanii</i>
<i>mcmA</i>	Q84FZ1	75486201	<i>Methylobacterium extorquens</i>
<i>mcmB</i>	Q6TMA2	75493131	<i>Methylobacterium extorquens</i>

Additional enzyme candidates identified based on high homology to the *E. coli scpA* gene product are identified below in Table 52.

Table 52.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>sbm</i>	NP_838397.1	30064226	<i>Shigella flexneri</i>
<i>SARI_04585</i>	ABX24358.1	160867735	<i>Salmonella enterica</i>
<i>YfreA_01000861</i>	ZP_00830776.1	77975240	<i>Yersinia frederiksenii</i>

There further exists evidence that genes adjacent to the methylmalonyl-CoA mutase catalytic genes are also required for maximum activity. For example, it has been demonstrated that the *meaB* gene from *M. extorquens* forms a complex with methylmalonyl-CoA mutase, stimulates *in vitro* mutase activity, and possibly protects it from irreversible inactivation (Korotkova, *J Biol Chem.* 279:13652-13658 (2004)). The *M. extorquens meaB* gene product is highly similar to the product of the *E. coli argK* gene (BLASTp: 45% identity, e-value: 4e-67) which is adjacent to *scpA* on the chromosome. No sequence for a *meaB* homolog in *P. freudenreichii* is catalogued in GenBank. However, the *Propionibacterium acnes* KPA171202 gene product, YP_055310.1, is 51% identical to the *M. extorquens meaB* protein and its gene is also adjacent to the methylmalonyl-CoA mutase gene on the chromosome. These genes/proteins are identified below in Table 53.

Table 53.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>argK</i>	AAC75955.1	1789285	<i>Escherichia coli K12</i>
KPA171202	YP_055310.1	50842083	<i>Propionibacterium acnes</i>
<i>meaB</i>	2QM8_B	158430328	<i>Methylobacterium extorquens</i>

Methylmalonyl-CoA Epimerase

Methylmalonyl-CoA epimerase (MMCE) is the enzyme that interconverts (R)-methylmalonyl-CoA and (S)-methylmalonyl-CoA. MMCE is an essential enzyme in the breakdown of odd-numbered fatty acids and of the amino acids valine, isoleucine, and methionine. Methylmalonyl-CoA epimerase is present in organisms such as *Bacillus subtilis* (YqjC) (Haller, *Biochemistry*. 39:4622-4629 (2000)), *Homo sapiens* (YqjC) (Fuller, *Biochem. J* 213:643-650 (1983)), *Rattus norvegicus* (Mcee) (Bobik, *J Biol Chem*. 276:37194-37198 (2001)), *Propionibacterium shermanii* (AF454511) (Haller, *Biochemistry* 39:4622-9 (2000); McCarthy, *Structure* 9:637-46 (2001) and (Fuller, *Biochem. J* 213:643-650 (1983)) and *Caenorhabditis elegans* (mmce) (Kuhnl et al., *FEBS J* 272:1465-1477 (2005)). The additional gene candidate, AE016877 in *Bacillus cereus*, has high sequence homology to the other characterized enzymes. MMCE activity is required if the employed methylmalonyl-CoA decarboxylase or methylmalonyl-CoA carboxytransferase requires the (S) stereoisomer of methylmalonyl-CoA. These genes/proteins are identified below in Table 54.

Table 54.

Gene	GenBank ID	GI Number	Organism
<i>YqjC</i>	NP_390273	255767522	<i>Bacillus subtilis</i>
<i>MCEE</i>	Q96PE7.1	50401130	<i>Homo sapiens</i>
<i>Mcee_predicted</i>	NP_001099811.1	157821869	<i>Rattus norvegicus</i>
<i>AF454511</i>	AAL57846.1	18042135	<i>Propionibacterium fredenreichii</i> sp. <i>shermanii</i>
<i>mmce</i>	AAT92095.1	51011368	<i>Caenorhabditis elegans</i>
<i>AE016877</i>	AAP08811.1	29895524	<i>Bacillus cereus</i> ATCC 14579

15 Methylmalonyl-CoA Decarboxylase

Methylmalonyl-CoA decarboxylase, is a biotin-independent enzyme that catalyzes the conversion of methylmalonyl-CoA to propionyl-CoA in *E. coli* (Benning, *Biochemistry*. 39:4630-4639 (2000); and Haller, *Biochemistry*. 39:4622-4629 (2000)). The stereo specificity of the *E. coli* enzyme was not reported, but the enzyme in *Propionigenium modestum* (Bott et al., *Eur. J. Biochem*. 250:590-599 (1997)) and *Veillonella parvula* (Huder, *J. Biol. Chem*. 268:24564-24571 (1993)) catalyzes the decarboxylation of the (S)-stereoisomer of methylmalonyl-CoA (Hoffmann, *FEBS. Lett*. 220:121-125 (1987)). The enzymes from *P. modestum* and *V. parvula* are comprised of multiple subunits that not only decarboxylate (S)-methylmalonyl-CoA, but also create a pump that transports sodium ions across the cell membrane as a means to generate energy. These genes/proteins are identified below in Table 55.

Table 55.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>YgfG</i>	NP_417394	90111512	<i>Escherichia coli</i>
<i>mmdA</i>	CAA05137	2706398	<i>Propionigenium modestum</i>
<i>mmdD</i>	CAA05138	2706399	<i>Propionigenium modestum</i>
<i>mmdC</i>	CAA05139	2706400	<i>Propionigenium modestum</i>
<i>mmdB</i>	CAA05140	2706401	<i>Propionigenium modestum</i>
<i>mmdA</i>	CAA80872	415915	<i>Veillonella parvula</i>
<i>mmdC</i>	CAA80873	415916	<i>Veillonella parvula</i>
<i>mmdE</i>	CAA80874	415917	<i>Veillonella parvula</i>
<i>mmdD</i>	CAA80875	415918	<i>Veillonella parvula</i>
<i>mmdB</i>	CAA80876	415919	<i>Veillonella parvula</i>

EXAMPLE IV**Pathways for Production of Propionyl-CoA from Glucose via Threonine**

- 5 Further to Examples I and II, the pathway for production of propionyl-CoA via threonine is exemplified in Figure 2. PEP is converted into oxaloacetate either via PEP carboxykinase or PEP carboxylase as described in Example III. Alternatively, PEP is converted first to pyruvate by pyruvate kinase and then to oxaloacetate by methylmalonyl-CoA carboxytransferase or pyruvate carboxylase as described in Example III. Oxaloacetate is converted into threonine by
- 10 the native threonine pathway engineered for high yields. It is then deaminated to form 2-oxobutanoate and subsequently converted into propionyl-CoA. In one alternative, 2-oxobutanoate is converted to propionaldehyde by a decarboxylase, which is then reduced to n-propanol by a propanol dehydrogenase.

Threonine Deaminase

- 15 The conversion of threonine to 2-oxobutanoate (or 2-ketobutyrate) can be accomplished by a threonine deaminase. It is encoded by one or more genes selected from *ilvA* (Calhoun et al., *J. Biol. Chem.* 248(10):3511-6, (1973)) and *tdcB* (Umbarger et al., *J. Bacteriol.* 73(1):105-12, (1957); Datta et al., *Proc. Natl. Acad. Sci. U S A* 84(2): 393-7(1987)). *Rhodospirillum rubrum* represents an additional exemplary organism containing threonine deaminase (Feldberg et al.,
- 20 *Eur. J. Biochem.* 21(3): 438-46 (1971); U.S. Patent 5,958,745). Details for exemplary enzymes for carrying out this transformation are shown below. These genes/proteins are identified below in Table 56.

Table 56.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ilvA</i>	AAC77492	1790207	<i>Escherichia coli</i>
<i>tdcB</i>	AAC76152	1789505	<i>Escherichia coli</i>
<i>Rru_A2877</i>	YP_427961.1	83594209	<i>Rhodospirillum rubrum</i>
<i>Rru_A0647</i>	YP_425738.1	83591986	<i>Rhodospirillum rubrum</i>

2-Oxobutanoate Dehydrogenase

2-oxobutanoate (2-ketobutyrate) can be converted to propionyl-CoA via a pyruvate formate lyase and a pyruvate formate lyase activating enzyme. The pyruvate formate lyase is encoded by gene selected from *pflB* and *tdcE*, while the pyruvate formate lyase activating enzyme is encoded by a *pflA* gene. Details for these exemplary genes for carrying out this transformation are already listed.

Alternatively, 2-oxobutanoate can be converted to propionyl-CoA by means of pyruvate dehydrogenase, pyruvate ferredoxin oxidoreductase (PFOR), or any other enzyme with 2-ketoacid dehydrogenase functionality. Such enzymes are also capable of converting pyruvate to acetyl-CoA. Exemplary pyruvate dehydrogenase enzymes are present in *E. coli* (Bisswanger, H., *J. Biol. Chem.* 256:815-822 (1981); Bremer, J., *Eur.J. Biochem.* 8:535-540 (1969); Gong et al., *J. Biol. Chem.* 275:13645-13653 (2000)), *B. subtilis* (Nakano et al., *J.Bacteriol.* 179:6749-6755 (1997)), *K. pneumonia* (Menzel et al., *J.Biotechnol.* 56:135-142 (1997)), *R. norvegicus* (Paxton et al., *Biochem. J.* 234:295-303 (1986)), for example. Exemplary gene information is provided below. These genes/proteins are identified below in Table 57.

Table 57.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>aceE</i>	NP_414656.1	16128107	<i>Escherichia coli str. K12 substr. MG1655</i>
<i>aceF</i>	NP_414657.1	16128108	<i>Escherichia coli str. K12 substr. MG1655</i>
<i>lpd</i>	NP_414658.1	16128109	<i>Escherichia coli str. K12 substr. MG1655</i>
<i>pdhA</i>	P21881.1	3123238	<i>Bacillus subtilis</i>
<i>pdhB</i>	P21882.1	129068	<i>Bacillus subtilis</i>
<i>pdhC</i>	P21883.2	129054	<i>Bacillus subtilis</i>
<i>pdhD</i>	P21880.1	118672	<i>Bacillus subtilis</i>
<i>aceE</i>	YP_001333808.1	152968699	<i>Klebsiella pneumonia MGH78578</i>
<i>aceF</i>	YP_001333809.1	152968700	<i>Klebsiella pneumonia MGH78578</i>
<i>lpdA</i>	YP_001333810.1	152968701	<i>Klebsiella pneumonia MGH78578</i>
<i>Pdha1</i>	NP_001004072.2	124430510	<i>Rattus norvegicus</i>
<i>Pdha2</i>	NP_446446.1	16758900	<i>Rattus norvegicus</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Dlat</i>	NP_112287.1	78365255	<i>Rattus norvegicus</i>
<i>Dld</i>	NP_955417.1	40786469	<i>Rattus norvegicus</i>

Exemplary PFOR enzymes include, for example, the enzyme from *Desulfovibrio africanus* which has been cloned and expressed in *E. coli*, resulting in an active recombinant enzyme that was stable for several days in the presence of oxygen (Pieulle et al., J. Bacteriol. 179:5684-5692 (1997)). Oxygen stability is relatively uncommon in PFORs and is reported to be conferred by a 60 residue extension in the polypeptide chain of the *D. africanus* enzyme. The *M. thermoacetica* PFOR is also well characterized (Menon et al. Biochemistry 36:8484-8494 (1997)) and was shown to have high activity in the direction of pyruvate synthesis during autotrophic growth (Furdui et al. J. Biol. Chem. 275:28494-28499 (2000)). Further, *E. coli* possesses an uncharacterized open reading frame, *ydbK*, that encodes a protein that is 51% identical to the *M. thermoacetica* PFOR. Evidence for pyruvate oxidoreductase activity in *E. coli* has been described (Blaschkowski et al., Eur. J. Biochem. 123:563-569 (1982)). The protein sequences of these exemplary PFOR enzymes can be identified by the following GenBank accession and/or GI numbers as shown below. Several additional PFOR enzymes have been described (Ragsdale, Chem. Rev. 103:2333-2346 (2003)). These genes/proteins are identified below in Table 58.

Table 58.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
Por	CAA70873.1	1770208	<i>Desulfovibrio africanus</i>
Por	YP_428946.1	83588937	<i>Moorella thermoacetica</i>
YdbK	NP_415896.1	16129339	<i>Escherichia coli</i>

Additional routes for producing propionyl-CoA are disclosed in U.S. Patent 5,958,745 which is incorporated by reference herein in its entirety. One such route involves converting 2-ketobutyrate to propionate by pyruvate oxidase, and converting propionate to propionyl-CoA via an acyl-CoA synthetase.

2-Oxobutanoate Decarboxylase

A keto acid decarboxylase can catalyze the conversion of 2-oxobutanoate to propionaldehyde. Several 2-keto acid decarboxylases have been identified. Enzyme candidates for this step are pyruvate decarboxylase (EC 4.1.1.1), benzoylformate decarboxylase (4.1.1.7), alpha-ketoglutarate decarboxylase (EC 4.1.1.71), branched-chain alpha-keto-acid decarboxylase (4.1.1.72), and indolepyruvate decarboxylase (EC 4.1.1.74). These classes of decarboxylases are

NADH-independent, they utilize thiamine diphosphate as a cofactor, and the interaction of the substrate with the enzyme-bound cofactor is thought to be the rate-limiting step for enzyme activation (Hubner, *Eur. J Biochem.* 92:175-181 (1978)). Pyruvate decarboxylase and benzoylformate decarboxylase have broad substrate ranges for diverse keto-acids and have been characterized in structural detail. Fewer alpha-ketoglutarate and branched-chain alpha-ketoacid decarboxylases have been characterized experimentally; however these enzymes also appear to decarboxylate a variety of keto-acid substrates.

Pyruvate decarboxylase (PDC), also termed keto-acid decarboxylase, is a key enzyme in alcoholic fermentation, catalyzing the decarboxylation of pyruvate to acetaldehyde. The enzyme from *Saccharomyces cerevisiae* has a broad substrate range for aliphatic 2-keto acids including 2-ketobutyrate, 2-ketovalerate, 3-hydroxypyruvate and 2-phenylpyruvate (22). The PDC from *Zymomonas mobilis*, encoded by *pdh*, has been a subject of directed engineering studies that altered the affinity for different substrates (Siegert et al., *Protein Eng Des Sel* 18:345-357 (2005)). The PDC from *Saccharomyces cerevisiae* has also been extensively studied, engineered for altered activity, and functionally expressed in *E. coli* (Li, *Biochemistry*. 38:10004-10012 (1999); ter Schure, *Appl. Environ. Microbiol.* 64:1303-1307 (1998) and Killenberg-Jabs, *Eur. J. Biochem.* 268:1698-1704 (2001)). The crystal structure of this enzyme is available (Killenberg-Jabs, *Eur. J. Biochem.* 268:1698-1704 (2001)). Other well-characterized PDC candidates include the enzymes from *Acetobacter pasteurians* (Chandra, *Arch. Microbiol.* 176:443-451 (2001)) and *Kluyveromyces lactis* (Krieger, *Eur. J. Biochem.* 269:3256-3263 (2002)). These genes/proteins are identified below in Table 59.

Table 59.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pdh</i>	P06672.1	118391	<i>Zymomonas mobilis</i>
<i>pdh1</i>	P06169	30923172	<i>Saccharomyces cerevisiae</i>
<i>pdh</i>	Q8L388	20385191	<i>Acetobacter pasteurians</i>
<i>pdh1</i>	Q12629	52788279	<i>Kluyveromyces lactis</i>

Like PDC, benzoylformate decarboxylase has a broad substrate range and has been the target of enzyme engineering studies. The enzyme from *Pseudomonas putida* has been extensively studied and crystal structures of this enzyme are available (Polovnikova et al, *Biochemistry* 42:1820-1830 (2003); and Hasson et al., *Biochemistry* 37:9918-9930 (1998)). Site-directed mutagenesis of two residues in the active site of the *Pseudomonas putida* enzyme altered the affinity (Km) of naturally and non-naturally occurring substrates (Siegert et al., *Protein Eng Des Sel* 18:345-357 (2005)). The properties of this enzyme have been further modified by directed engineering (Lingen et al., *Chembiochem.* 4:721-726 (2003); and Lingen, *Protein Eng* 15:585-

593 (2002)). The enzyme from *Pseudomonas aeruginosa*, encoded by *mdlC*, has also been characterized experimentally (Barrowman, *FEMS Microbiology Letters* 34:57-60 (1986)). Additional gene candidates from *Pseudomonas stutzeri*, *Pseudomonas fluorescens* and other organisms can be inferred by sequence homology or identified using a growth selection system developed in *Pseudomonas putida* (Henning et al., *Appl. Environ. Microbiol.* 72:7510-7517 (2006)). These genes/proteins are identified below in Table 60.

Table 60.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mdlC</i>	P20906.2	3915757	<i>Pseudomonas putida</i>
<i>mdlC</i>	Q9HUR2.1	81539678	<i>Pseudomonas aeruginosa</i>
<i>dpgB</i>	ABN80423.1	126202187	<i>Pseudomonas stutzeri</i>
<i>ilvB-1</i>	YP_260581.1	70730840	<i>Pseudomonas fluorescens</i>

A third enzyme capable of decarboxylating 2-oxoacids is alpha-ketoglutarate decarboxylase (KGD). The substrate range of this class of enzymes has not been studied to date. The KDC from *Mycobacterium tuberculosis* (Tian, *Proc Natl Acad Sci U S A* 102:10670-10675 (2005)) has been cloned and functionally expressed in other internal projects at Genomatica. However, it is not an ideal candidate for strain engineering because it is large (~130 kD) and GC-rich. KDC enzyme activity has been detected in several species of Rhizobia including *Bradyrhizobium japonicum* and *Mesorhizobium loti* (Green, *J Bacteriol.* 182:2838-2844 (2000)). Although the KDC-encoding gene(s) have not been isolated in these organisms, the genome sequences are available and several genes in each genome are annotated as putative KDCs. A KDC from *Euglena gracilis* has also been characterized but the gene associated with this activity has not been identified to date (Shigeoka, *Arch. Biochem. Biophys.* 288:22-28 (1991)). The first twenty amino acids starting from the N-terminus were sequenced MTYKAPVKDVKFLLDKVFKV (Shigeoka, *Arch. Biochem. Biophys.* 288:22-28 (1991)). The gene could be identified by testing candidate genes containing this N-terminal sequence for KDC activity. These genes/proteins are identified below in Table 61.

Table 61.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>kgd</i>	O50463.4	160395583	<i>Mycobacterium tuberculosis</i>
<i>kgd</i>	NP_767092.1	27375563	<i>Bradyrhizobium japonicum</i> <i>USDA110</i>
<i>kgd</i>	NP_105204.1	13473636	<i>Mesorhizobium loti</i>

A fourth candidate enzyme for catalyzing this step is branched chain alpha-ketoacid decarboxylase (BCKA). This class of enzyme has been shown to act on a variety of compounds varying in chain length from 3 to 6 carbons (Oku, *J Biol Chem.* 263:18386-18396 (1988); and

Smit et al., *Appl Environ Microbiol* 71:303-311 (2005)). The enzyme in *Lactococcus lactis* has been characterized on a variety of branched and linear substrates including 2-oxobutanoate, 2-oxohexanoate, 2-oxopentanoate, 3-methyl-2-oxobutanoate, 4-methyl-2-oxobutanoate and isocaproate (Smit et al., *Appl Environ Microbiol* 71:303-311 (2005)). The enzyme has been

5 structurally characterized (Berthold et al., *Acta Crystallogr. D Biol Crystallogr.* 63:1217-1224 (2007)). Sequence alignments between the *Lactococcus lactis* enzyme and the pyruvate decarboxylase of *Zymomonas mobilis* indicate that the catalytic and substrate recognition residues are nearly identical (Siegert et al., *Protein Eng Des Sel* 18:345-357 (2005)), so this enzyme would be a promising candidate for directed engineering. Decarboxylation of alpha-

10 ketoglutarate by a BCKA was detected in *Bacillus subtilis*; however, this activity was low (5%) relative to activity on other branched-chain substrates (Oku, *J Biol Chem.* 263:18386-18396 (1988)) and the gene encoding this enzyme has not been identified to date. Additional BCKA gene candidates can be identified by homology to the *Lactococcus lactis* protein sequence. Many of the high-scoring BLASTp hits to this enzyme are annotated as indolepyruvate

15 decarboxylases (EC 4.1.1.74). Indolepyruvate decarboxylase (IPDA) is an enzyme that catalyzes the decarboxylation of indolepyruvate to indoleacetaldehyde in plants and plant bacteria. This gene/protein is identified below in Table 62.

Table 62.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>kdcA</i>	AAS49166.1	44921617	<i>Lactococcus lactis</i>

Recombinant branched chain alpha-keto acid decarboxylase enzymes derived from the E1

20 subunits of the mitochondrial branched-chain keto acid dehydrogenase complex from *Homo sapiens* and *Bos taurus* have been cloned and functionally expressed in *E. coli* (Wynn, *J. Biol. Chem.* 267:12400-12403 (1992); Davie, *J. Biol. Chem.* 267:16601-16606 (1992) and Wynn et al., *J. Biol. Chem.* 267:1881-1887 (1992)). In these studies, the authors found that co-expression of chaperonins GroEL and GroES enhanced the specific activity of the decarboxylase by 500-

25 fold (Wynn, *J. Biol. Chem.* 267:12400-12403 (1992)). These enzymes are composed of two alpha and two beta subunits. These genes/proteins are identified below in Table 63.

Table 63.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>BCKDHB</i>	NP_898871.1	34101272	<i>Homo sapiens</i>
<i>BCKDHA</i>	NP_000700.1	11386135	<i>Homo sapiens</i>
<i>BCKDHB</i>	P21839	115502434	<i>Bos taurus</i>
<i>BCKDHA</i>	P11178	129030	<i>Bos taurus</i>

EXAMPLE V

Pathways for Production of Propionyl-CoA from Glucose via Malonyl-CoA

Further to Examples I and II, the pathway for production of propionyl-CoA via malonyl-CoA is exemplified in Figure 3. Acetyl CoA is carboxylated to form malonyl-CoA. This is then
 5 reduced to malonate semialdehyde, and subsequently transformed into 3-hydroxypropionate (3HP). 3HP is converted into propionyl-CoA via propionyl-CoA synthase.

Acetyl-CoA Carboxylase

The multisubunit acetyl-CoA carboxylase complex (ACC), broadly conserved among bacteria, catalyzes the ATP-dependent formation of malonyl-CoA by acetyl-CoA and bicarbonate. This
 10 reaction serves as the first committed step in fatty acid biosynthesis, and the enzyme has been targeted in efforts to develop antibacterial drugs and inhibitors in *E. coli* (Freiberg et al., *J. Biol. Chem.* 279: 26066-26073 (2004)), yeast (Zhang, *Proc. Natl. Acad. Sci. U S A.* 101:5910-5915 (2004)), *Bacillus subtilis* (Freiberg et al., *J. Biol. Chem.* 279:26066-26073 (2004)) and other organisms (Barber, *Biochim. Biophys. Acta* 1733:1-28 (2005)). In *E. coli* and many other
 15 bacteria, ACC is composed of four subunits encoded by *accA*, *accB*, *accC* and *accD* (Choi-Rhee, *J. Biol. Chem.* 278:30806-30812 (2003)). Expression of two subunits, *accB* and *accC*, is autoregulated by the gene product of *accB* (James, *J. Biol. Chem.* 279:2520-2527 (2004)). In yeast, the enzyme is encoded by two genes, *hfa1* and *acc1*. The gene *bpl1*, encoding a biotin:apoprotein ligase, is required for enzyme function.

20 Autotrophic members of the archaeal taxonomic group *Sulfolobales* exhibit high levels of acetyl-CoA carboxylase activity in the context of the 3-hydroxypropionate cycle (Chuakrut, *J. Bacteriol.* 185:938-947 (2003); and Hugler, *Eur. J. Biochem.* 270:736-744 (2003)). In *Metallosphaera sedula*, the acyl-CoA carboxylase holoenzyme is a multimer composed of subunits encoded by three genes: *Msed_0148* (biotin/lipoyl attachment), *Msed_0147* (biotin
 25 carboxylase), and *Msed_1375* (carboxyl transferase). The enzyme has been purified and characterized and was found to be bifunctional, reacting with acetyl-CoA and propionyl-CoA (Hugler, *Eur. J. Biochem.* 270:736-744 (2003)). A bifunctional archaeal acetyl-CoA carboxylase enzyme from *Acidanus brierleyi*, encoded by three genes, has been cloned into *E. coli* and characterized (Chuakrut, *J. Bacteriol.* 185:938-947 (2003)). The sequences of *A. brierleyi* acyl-
 30 CoA carboxylase genes and flanking regions were submitted to the DNA Data Bank of Japan (DDBJ) under accession no. AB088419. Although these archaeal enzymes exhibit high activity it

should be noted that the optimum temperature is 65°C (Chuakrut, *J. Bacteriol.* 185:938-947 (2003)). These genes/proteins are identified below in Table 64.

Table 64.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>accA</i>	NP_414727	16128178	<i>Escherichia coli</i> K12 str. MG1655
<i>accB</i>	NP_417721	16131143	<i>Escherichia coli</i> K12 str. MG1655
<i>accC</i>	NP_417722	16131144	<i>Escherichia coli</i> K12 str. MG1655
<i>accD</i>	NP_416819	16130251	<i>Escherichia coli</i> K12 str. MG1655
<i>accA</i>	NP_390798.1	16079972	<i>Bacillus subtilis</i> subsp. <i>subtilis</i> str. 168
<i>accB</i>	NP_390315.1	16079491	<i>Bacillus subtilis</i> subsp. <i>subtilis</i> str. 168
<i>accC</i>	NP_390314.1	16079490	<i>Bacillus subtilis</i> subsp. <i>subtilis</i> str. 168
<i>accD</i>	NP_390799.1	16079973	<i>Bacillus subtilis</i> subsp. <i>subtilis</i> str. 168
<i>bpl1</i>	NP_010140.1	6320060	<i>Saccharomyces cerevisiae</i>
<i>hfa1</i>	NP_013934.1	6323863	<i>Saccharomyces cerevisiae</i>
<i>acc1</i>	NP_014413.1	6324343	<i>Saccharomyces cerevisiae</i>
<i>accB Msed_0148</i>	Q8J2Z3	74499802	<i>Metallosphaera sedula</i>
<i>accC Msed_0147</i>	Q8J2Z4	74499032	<i>Metallosphaera sedula</i>
<i>pccB Msed_1375</i>	Q8J2Z5	74499033	<i>Metallosphaera sedula</i>
<i>accB</i>	BAC55868.1	27877098	<i>Acidanus brierleyi</i>
<i>accC</i>	BAC55867.1	27877097	<i>Acidanus brierleyi</i>
<i>pccB</i>	BAC55869.1	27877099	<i>Acidanus brierleyi</i>

Malonyl-CoA Reductase and Malonate Semialdehyde Reductase

- 5 The reduction of malonyl-CoA to 3-HP can be accomplished by a bifunctional malonyl-CoA reductase with aldehyde dehydrogenase and alcohol dehydrogenase functionality. An NADPH-dependent enzyme with this activity has been characterized in *Chloroflexus aurantiacus* where it participates in the 3-hydroxypropionate cycle (Hugler, *J. Bacteriol.* 184:2404-2410 (2002); and Strauss, *Eur. J. Biochem.* 215:633-643 (1993)). This enzyme, with a mass of 300 kDa, is highly
- 10 substrate-specific and shows little sequence similarity to other known oxidoreductases (Hugler, *J. Bacteriol.* 184:2404-2410 (2002)). No enzymes in other organisms have been shown to catalyze this specific reaction; however there is bioinformatic evidence that other organisms may have similar pathways (Klatt, *Environ. Microbiol.* 9:2067-2078 (2007)). Enzyme candidates in other organisms including *Roseiflexus castenholzii*, *Erythrobacter* sp. *NAP1* and marine gamma
- 15 proteobacterium HTCC2080 can be inferred by sequence similarity. These genes/proteins are identified below in Table 65.

Table 65.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mcr</i>	AAS20429.1	42561982	<i>Chloroflexus aurantiacus</i>
<i>Rcas_2929</i>	YP_001433009.1	156742880	<i>Roseiflexus castenholzii</i>
<i>NAP1_02720</i>	ZP_01039179.1	85708113	<i>Erythrobacter sp. NAP1</i>
MGP2080_00535	ZP_01626393.1	119504313	marine gamma proteobacterium HTCC2080

Alternatively, the reduction of malonyl-CoA to 3-HP can be catalyzed by two separate enzymes: a CoA-acylating aldehyde dehydrogenase and a primary alcohol dehydrogenase. By this route, malonyl-CoA is first reduced to malonate semialdehyde (MSA) by malonate-semialdehyde dehydrogenase or malonyl-CoA reductase. MSA is subsequently converted to 3-HP by 3-HP-dehydrogenase.

Malonyl-CoA reductase is a key enzyme in autotrophic carbon fixation via the 3-hydroxypropionate cycle in thermoacidophilic archaeal bacteria (Berg, *Science*. 318:1782-1786 (2007); and Thauer, *Science*. 318:1732-1733 (2007)). The enzyme utilizes NADPH as a cofactor and has been characterized in *Metallosphaera* and *Sulfolobus spp* (Alber et al, *J. Bacteriol.* 188:8551-8559 (2006); and Hugler, *J. Bacteriol.* 184:2404-2410 (2002)). The enzyme encoded by *Msed_0709* in *Metallosphaera sedula* is known to convert malonyl-CoA to malonic semialdehyde and operate in the direction of interest (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006); and (Berg, *Science*. 318:1782-1786 (2007)). A gene encoding a malonyl-CoA reductase from *Sulfolobus tokodaii* was cloned and heterologously expressed in *E. coli* (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006). Although the aldehyde dehydrogenase functionality of these enzymes is similar to the bifunctional dehydrogenase from *Chloroflexus aurantiacus*, there is little sequence similarity. Both malonyl-CoA reductase enzyme candidates have high sequence similarity to aspartate-semialdehyde dehydrogenase, an enzyme catalyzing the reduction and concurrent dephosphorylation of aspartyl-4-phosphate to aspartate semialdehyde. Additional gene candidates can be found by sequence homology to proteins in other organisms including *Sulfolobus solfataricus* and *Sulfolobus acidocaldarius*. These genes/proteins are identified below in Table 66.

Table 66.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Msed_0709</i>	YP_001190808.1	146303492	<i>Metallosphaera sedula</i>
<i>mcr</i>	NP_378167.1	15922498	<i>Sulfolobus tokodaii</i>
<i>asd-2</i>	NP_343563.1	15898958	<i>Sulfolobus solfataricus</i>
<i>Saci_2370</i>	YP_256941.1	70608071	<i>Sulfolobus acidocaldarius</i>

The subsequent conversion of malonic semialdehyde to 3-HP can be accomplished by an enzyme with 3-HP dehydrogenase activity. Three enzymes are known to catalyze this

- conversion: NADH-dependent 3-hydroxypropionate dehydrogenase, NADPH-dependent malonate semialdehyde reductase, and NADH-dependent 3-hydroxyisobutyrate dehydrogenases. An NADH-dependent 3-hydroxypropionate dehydrogenase is thought to participate in beta-alanine biosynthesis pathways from propionate in bacteria and plants (Rathinasabapathi, *Journal of Plant Pathology* 159:671-674 (2002); and Stadtman, A. *J. Am. Chem. Soc.* 77:5765-5766 (1955)). This enzyme has not been associated with a gene in any organism to date. NADPH-dependent malonate semialdehyde reductase catalyzes the reverse reaction in autotrophic CO₂-fixing bacteria. Although the enzyme activity has been detected in *Metallosphaera sedula*, the identity of the gene is not known (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006)).
- Several 3-hydroxyisobutyrate dehydrogenase enzymes have also been shown to convert malonic semialdehyde to 3-HP. Three gene candidates exhibiting this activity are *mmsB* from *Pseudomonas aeruginosa* PAO1 (Gokam et al., U.S. Patent 7,393,676 (2008)). *mmsB* from *Pseudomonas putida* KT2440 (Liao, U.S. Patent Publication 2005-0221466 (2005) and *mmsB* from *Pseudomonas putida* E23 (Chowdhury, *Biosci. Biotechnol. Biochem.* 60:2043-2047 (1996)). The protein from *Pseudomonas putida* E23 has been characterized and functionally expressed in *E. coli*; however, its activity on 3-HP was relatively low (Chowdhury, *Biosci. Biotechnol. Biochem.* 60:2043-2047 (1996)). An enzyme with 3-hydroxybutyrate dehydrogenase activity in *Alcaligenes faecalis* M3A has also been identified (Liao, U.S. Patent Publication 2005-0221466 (2005); and Liao, U.S. Patent Publication 2005-0221466 (2005)).
- Additional gene candidates from other organisms including *Rhodobacter spaeroides* can be inferred by sequence similarity. These genes/proteins are identified below in Table 67.

Table 67.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mmsB</i>	AAA25892.1	151363	<i>Pseudomonas aeruginosa</i>
<i>mmsB</i>	NP_252259.1	15598765	<i>Pseudomonas aeruginosa</i> PAO1
<i>mmsB</i>	NP_746775.1	26991350	<i>Pseudomonas putida</i> KT2440
<i>mmsB</i>	JC7926	60729613	<i>Pseudomonas putida</i> E23
<i>orfB1</i>	AAL26884	16588720	<i>Rhodobacter spaeroides</i>

- Enzymes exhibiting a 4-hydroxybutyrate activity (EC 1.1.1.61) may also be able to convert malonic semialdehyde to 3-HP, as the chemical transformation is very similar. Such enzymes have been characterized in *Ralstonia eutropha* (Bravo, *J. Forensic Sci.* 49:379-387 (2004)), *Clostridium kluyveri* (Wolff, *Protein Expr. Purif.* 6:206-212 (1995)) and *Arabidopsis thaliana* (Breitkreuz et al., *J. Biol. Chem.* 278:41552-41556 (2003)). Activity of these enzymes on malonic semialdehyde has not been demonstrated experimentally to date. However, since these

enzymes have been studied in other internal projects at Genomatica they could easily be tested for 3-HP dehydrogenase activity. These genes/proteins are identified below in Table 68.

Table 68.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>4hbd</i>	YP_726053.1	113867564	<i>Ralstonia eutropha</i> H16
<i>4hbd</i>	L21902.1	146348486	<i>Clostridium kluyveri</i> DSM 555
<i>4hbd</i>	Q94B07	75249805	<i>Arabidopsis thaliana</i>

Propionyl-CoA Synthase

- 5 The conversion of 3-hydroxypropionate (3HP) to propionyl-CoA is accomplished by a propionyl-CoA synthase. This step is known to be catalyzed by a single fusion protein of 201 KDa in *Chloroflexus aurantiacus* (Alber, *J Biol. Chem.* 277:12137-12143 (2002)). The protein is comprised of a CoA ligase, an enoyl-CoA hydratase and an enoyl-CoA reductase. The enzyme has been purified 30-fold to near homogeneity and has a very large native molecular
- 10 mass between 500 and 800 kDa. In thermoacidophilic *Metallosphaera sedula* (and members of the *Sulfolobaceae* family), this function is catalyzed by three different enzymes, a 3-hydroxypropionyl-CoA synthetase that activates 3HP to its CoA ester, a 3-hydroxypropionyl-CoA dehydratase that converts 3-HP-CoA to acryloyl-CoA followed by the reduction of the latter to form propionyl-CoA. A 3-HP-CoA synthetase had been reported (Alber, *J Bacteriol.*
- 15 190:1383-1389 (2008)). The gene encoding the protein has been sequenced and gene encoding a homologous protein identified in the genome of *Sulfolobus tokodaii*; similar genes were found in *S. solfataricus* and *S. acidocaldarius*. The gene was heterologously expressed in *Escherichia coli*. These genes/proteins are identified below in Table 69.

Table 69.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Msed_1456</i>	YP_001191537	146304221	<i>M. sedula</i>
<i>ST0783</i>	NP_376686	15921017	<i>S. tokodaii</i>
<i>acsA-10</i>	NP_344510	15899905	<i>S. solfataricus</i>
<i>Saci_1184</i>	YP_255824	70606954	<i>S. acidocaldarius</i>
<i>pcs</i>	AAL47820	29126583	<i>C. aurantiacus</i>

- 20 Recently, 3-hydroxypropionyl-CoA dehydratase and acryloyl-CoA reductase were purified from *M. sedula* (Teufel, *J Bacteriol.* 191:4572-4581 (2009)), the coding genes were identified from the genome of *M. sedula* and other members of the *Sulfolobales*, and recombinant enzymes were produced as a proof of function. It was concluded that the genes coding for 3-hydroxypropionyl-CoA dehydratase and acryloyl-CoA reductase are not clustered on the *Metallosphaera* or the

Sulfolobus genome. Comparison of the respective domains of propionyl-CoA synthase in these two organisms has revealed that the enzyme(s) catalyzing the conversion of 3HP to propionyl-CoA has evolved independently in these two phyla. The GenBank accession and/or GI numbers for the 3-HP-CoA dehydratase from *M. sedula* are identified below in Table 70.

5

Table 70.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Msed_2001</i>	YP_001192065.1	146304749	<i>M. sedula</i>

The GenBank IDs for acryloyl-CoA reductases are identified below in Table 71.

Table 71.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Msed_1426</i>	YP_001191508.1	146304192	<i>M. sedula</i>
<i>ST0480</i>	NP_376364	15920695	<i>S. tokodaii</i>

Other gene candidates encoding these two enzymes can be obtained by sequence homology searches.

10

EXAMPLE VI

Pathways for Production of Propionyl-CoA from Glucose via Lactate

Further to Examples I and II, the pathway for production of propionyl-CoA via lactate is exemplified in Figure 4. This pathway presents yet another redox balanced route for the formation of propionyl-CoA. Pyruvate is reduced to form lactate which is then activated to form lactoyl-CoA. The lactoyl-CoA is dehydrated to form acryloyl-CoA and then reduced to generate propionyl-CoA.

15

Lactate dehydrogenase

The conversion of pyruvate to lactate is catalyzed by lactate dehydrogenase (EC 1.1.1.27). Many lactate dehydrogenases have been described in detail (Garvie, *Microbiol Rev* 44:106-139 (1980)). The fermentative lactate dehydrogenase of *Escherichia coli* will be the first candidate to be overexpressed for converting pyruvate to lactate (Bunch, *Microbiology* 143 (Pt 1), 187-195 (1997)). Other lactate dehydrogenase candidates will be utilized for this step including those with low K_m for pyruvate that favors the formation of lactate, such as lactate dehydrogenase from: *Lactobacillus casei* (Gordon, *Eur. J Biochem.* 67:543-555 (1976)), *Plasmodium falciparum* (Brown et al., *Biochemistry* 43:6219-6229 (2004)), and *Thermotoga*

25

maritime (Auerbach et al., *Structure*. 6:769-781 (1998)). These genes/proteins are identified below in Table 72.

Table 72.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ldh</i>	P52643	1730102	<i>Escherichia coli</i>
<i>ldh</i>	P00343	126063	<i>Lactobacillus casei</i>
<i>ldh</i>	Q6JH32	74911026	<i>Plasmodium ovale</i>
<i>ldh</i>	P16115	547837	<i>Thermotoga maritima</i>

5 Lactate-CoA transferase

The activation of lactate to lactoyl-CoA can be catalyzed by lactate-CoA transferase activity associated with propionate CoA-transferase (EC 2.8.3.1). *Clostridium propionicum* ferments alanine via the nonrandomising pathway with acryloyl-CoA as characteristic intermediate. In this pathway, lactate is activated to lactoyl-CoA by the enzyme propionate:acetyl-CoA CoA-transferase (EC 2.8.3.1, or propionate CoA-transferase) using propionyl-CoA or acetyl-CoA as a coenzyme A donor (Schweiger, *FEBS Lett.* 171:79-84 (1984)). The enzyme exhibited rather broad substrate specificities for monocarboxylic acids including acrylate, propionate and butyrate whereas dicarboxylic acids were not used. Gene coding for this enzyme was cloned (Selmer, *Eur. J Biochem.* 269:372-380 (2002)). Other propionate CoA-transferase can be candidates for this step include homologues of *Clostridium propionicum* propionate CoA-transferase. These genes/proteins are identified below in Table 73.

Table 73.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pct</i>	Q9L3F7	75416255	<i>Clostridium propionicum</i>
<i>pct</i>	YP_002270763.1	209397911	<i>Escherichia coli</i> O157:H7
<i>pct</i>	Q220N6	122479931	<i>Rhodospirillum rubrum</i> DSM 15236
<i>pct</i>	Q46MA6	123621528	<i>Ralstonia eutropha</i>

Lactoyl-CoA dehydratase

The dehydration of lactoyl-CoA to acryloyl-CoA is catalyzed by lactoyl-CoA dehydratase (EC 4.2.1.54). *Clostridium propionicum* ferments alanine via the nonrandomising pathway with acryloyl-CoA as characteristic intermediate (Schweiger, *FEBS Lett.* 171:79-84 (1984)). In this pathway, lactoyl-CoA is dehydrated to acryloyl-CoA by the lactoyl-CoA dehydratase (Hofmeister, *Eur. J Biochem.* 206:547-552 (1992)). Cloning of the propionate CoA-transferase

also identified a second ORF (*lcdB*) likely encoding one subunit of the lactoyl-CoA dehydratase required in the pathway. The *lcdB* is similar to the 2-hydroxyglutaryl-CoA dehydratase β subunit. Homologues of *lcdB* will be tested for their activity in this step. These genes/proteins are identified below in Table 74.

5

Table 74.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
CBC_A0885	ZP_02621214	168186579	<i>Clostridium botulinum</i> C str. Eklund
CBC_A0886	ZP_02621215	168186580	<i>Clostridium botulinum</i> C str. Eklund
<i>hgdB</i>	YP_878441	118444181	<i>Clostridium novyi</i> -NT
<i>hgdA</i>	YP_878442	118444701	<i>Clostridium novyi</i> -NT

Acryloyl-CoA reductase

The conversion of acryloyl-CoA to propionyl-CoA is catalyzed by the acryloyl-CoA reductase. In alanine-fermenting *Clostridium propionicum*, acryloyl-CoA reductase catalyses the irreversible NADH-dependent formation of propionyl-CoA from acryloyl-CoA. The enzyme has been purified and the N-termini of the subunits of the enzyme have been determined (Hetzel et al., *Eur. J Biochem.* 270:902-910 (2003)). The N-terminus of the dimeric propionyl-CoA dehydrogenase subunit is similar to those of butyryl-CoA dehydrogenases from several *Clostridia* and related anaerobes (up to 55% sequence identity). The N-termini of the β and γ subunits share 40% and 35% sequence identities with those of the A and B subunits of the electron-transferring flavoprotein (ETF) from *Megasphaera elsdenii*, respectively, and up to 60% with those of putative ETFs from other anaerobes. Since the complete genome sequence of *Clostridium propionicum* is not available, the N-terminus of the propionyl-CoA dehydrogenase subunit “MDFKLTKTQVLQQWLFAEFAGIGIKPIAE” (SEQ ID NO.) was used in similarity search and resulted in the following homologues of the propionyl-CoA dehydrogenase for their activities in this step. These genes/proteins are identified below in Table 75.

20

Table 75.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>bcdA</i>	CAQ53135	188027001	<i>Clostridium saccharobutylicum</i>
<i>Cbei_2035</i>	ABR34203	149903370	<i>Clostridium beijerinckii</i>
<i>ANACAC_00471</i>	EDR98937	167654808	<i>Anaerostipes caccae</i> DSM 14662

Additionally, a tri-functional propionyl-CoA synthase (*pcs*) gene was identified from the phototrophic green non-sulfur eubacterium *Chloroflexus aurantiacus* (Alber, *J Biol. Chem.* 277:12137-12143 (2002)). The propionyl-CoA synthase is a natural fusion protein of 201 kDa consisting of a CoA ligase, an enoyl-CoA hydratase, and an enoyl-CoA reductase. The enzyme

25

catalyzes the conversion from 3-hydroxypropionate to 3-hydroxypropionyl-CoA to acryloyl-CoA then to propionyl-CoA. This enzyme can be utilized in whole or in part for its enoyl-CoA reductase activity. The gene/protein is identified below in Table 76.

Table 76.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pcs</i>	AAL47820	29126583	<i>Chloroflexus aurantiacus</i>

5


EXAMPLE VII

Pathways for Co-production of 1,4-butanediol (1,4-BDO) and Isopropanol from Glucose

This example describes exemplary pathways for co-production of 1,4-butanediol (1,4-BDO) and isopropanol.

- 10 Novel pathways for co-producing 1,4-butanediol (1,4-BDO) and isopropanol and related products are described herein. In the 1,4-butanediol (1,4-BDO) and isopropanol co-production pathway of Figure 5, central metabolism intermediates are first channeled into succinyl-CoA. For formation of succinyl-CoA, phosphoenolpyruvate (PEP) is converted into oxaloacetate either via PEP carboxykinase or PEP carboxylase. Alternatively, PEP is converted first to
- 15 pyruvate by pyruvate kinase and then to oxaloacetate by methylmalonyl-CoA carboxytransferase or pyruvate carboxylase. Oxaloacetate is then converted to succinyl-CoA by means of the reductive TCA cycle. Succinyl-CoA is then converted to succinic semialdehyde by a CoA-dependent aldehyde dehydrogenase. Alternatively, succinate can be converted to succinic semialdehyde by a succinate reductase. Next, succinic semialdehyde is reduced to 4-
- 20 hydroxybutyrate by 4-hydroxybutyrate dehydrogenase. Activation of 4-HB to its acyl-CoA is catalyzed by a CoA transferase or synthetase. Alternatively, 4-HB can be converted into 4-hydroxybutyryl-phosphate and subsequently transformed into 4-HB-CoA by a phosphotrans-4-hydroxybutyrylase. 4-HB-CoA is then converted to 1,4-BDO by either a bifunctional CoA-dependent aldehyde/alcohol dehydrogenase, or by two separate enzymes with aldehyde and
- 25 alcohol dehydrogenase activity. Yet another alternative that bypasses the 4-HB-CoA intermediate is direct reduction of 4-HB to 4-hydroxybutyrylaldehyde by a carboxylic acid reductase. 4-Hydroxybutyrylaldehyde is subsequently reduced to 1,4-BDO by an alcohol dehydrogenase. Yet another route that bypasses the CoA intermediate is reduction of 4-hydroxybutyryl-phosphate to 4-hydroxybutyrylaldehyde by a phosphate reductase. Pathways for
- 30 production of isopropanol proceed as described above in Examples I and II.

The maximum theoretical yield of a 14-BDO and isopropanol producing organism is 0.77 moles isopropanol and 0.46 moles 14-BDO per mole glucose consumed (0.26 g/g IPA and 0.23 g/g 14-BDO), per the following equation:



5


EXAMPLE VIII

Pathways for Co-production of 1,3-butanediol (1,3-BDO) and Isopropanol from Glucose

This example describes exemplary pathways for co-production of 1,3-butanediol (13-BDO) and isopropanol.

- 10 Novel pathways for co-producing 1,3-butanediol (13-BDO) and isopropanol and related products are described herein. The coproduction route to 1,3-butanediol (13-BDO) and isopropanol, shown in Figure 6, also proceeds through 4-hydroxybutyryl-CoA, formed as described in Example VI. In this route, 4-hydroxybutyryl-CoA is dehydrated and isomerized to form crotonyl-CoA. The dehydration and vinylisomerisation reactions are catalyzed by a
- 15 bifunctional enzyme, 4-hydroxybutyryl-CoA dehydratase. Crotonyl-CoA is then hydrated to 3-hydroxybutyryl-CoA. Removal of the CoA moiety and concurrent reduction yields 3-hydroxybutyraldehyde. Finally reduction of the aldehyde by 3-hydroxybutyraldehyde reductase yields 13-BDO. Alternately, 3-hydroxybutyryl-CoA can be converted to 13-BDO directly by a 3-hydroxybutyryl-CoA reductase (alcohol forming). Several other alternate routes are possible
- 20 in this pathway. Succinate can be converted to succinic semialdehyde by a carboxylic acid reductase, bypassing the formation of succinyl-CoA. 4-HB can be phosphorylated to 4-HB-phosphate by a kinase, then subsequently converted to 4-HB-CoA. Finally 3-hydroxybutyryl-CoA can be de-acylated by a CoA hydrolase, transferase or synthetase, then subsequently reduced to 3-hydroxybutyraldehyde by a carboxylic acid reductase.
- 25 Pathways for production of isopropanol proceed as described above in Examples I and II.

The maximum theoretical yield of 13-BDO and isopropanol via this pathway is 0.77 moles isopropanol and 0.46 moles 13-BDO per mole glucose consumed (0.26 g/g IPA and 0.23 g/g 13-BDO), per the following equation:



EXAMPLE IX

Pathways for Co-production of methylacrylic acid (MAA) and Isopropanol from Glucose

This example describes exemplary pathways for co-production of methylacrylic acid (MAA) and isopropanol.

- 5 Novel pathways for co-producing methylacrylic acid (MAA) and isopropanol and related products are described herein. Two coproduction routes to methylacrylic acid (MAA) are shown in Figures 7 and 8. The route shown in Figure 7 proceeds through 4-hydroxybutyryl-CoA, formed as described previously. 4-Hydroxybutyryl-CoA is converted to 3-hydroxyisobutyryl-CoA by a methyl mutase. The CoA moiety of 3-Hydroxyisobutyryl-CoA is then removed by a
 10 CoA transferase, hydrolase or synthetase. Finally, dehydration of the 3-hydroxy group yields MAA. Several of the key steps in this route can be bypassed by alternate routes. Succinate, for example, can be directly converted to succinic semialdehyde by a succinate reductase, bypassing the formation of succinyl-CoA. The conversion of 4-HB to 4-HB-CoA can proceed through the intermediate 4-hydroxybutyrylphosphate, via the enzymes 4-hydroxybutyrate kinase and
 15 phosphotrans-4-hydroxybutyrylase. 3-HIBCOA can be converted to MAA via the intermediate methacrylyl-CoA. Pathways for production of isopropanol proceed as described above in Examples I and II.

- In the alternate MAA coproduction route shown in Figure 8, succinyl-CoA is formed through the reductive TCA cycle, then converted to methylmalonyl-CoA by methylmalonyl-CoA
 20 mutase. An epimerase may be required to convert the (R) stereoisomer of methylmalonyl-CoA to the (S) configuration. A CoA-dependent aldehyde dehydrogenase then converts methylmalonyl-CoA to methylmalonate semialdehyde. Reduction of the aldehyde to 3-hydroxyisobutyrate, followed by dehydration, yields MAA. Alternately, methylmalonyl-CoA is converted to 3-hydroxyisobutyrate by an alcohol-forming CoA reductase. In yet another
 25 alternate route, methylmalonyl-CoA is converted to methylmalonate by a CoA hydrolase, transferase or synthetase. Methylmalonate is subsequently converted to methylmalonate semialdehyde by a carboxylic acid reductase. Methylmalonate semialdehyde is converted to MAA as described previously. Pathways for production of isopropanol proceed as described above in Examples I and II.
- 30 Both MAA coproduction pathways achieve yields 0.67 moles each of isopropanol and MAA per mole glucose utilized (0.22 g/g isopropanol and 0.32 g/g MAA) per the equation:



EXAMPLE X

Enzyme Classification System for Production of Isopronaol and 1,4-Butanediol (1,4-BDO), 1,3-Butanediol (1,3-BDO) or Methylacrylic acid (MAA)

5 This example describes the enzyme classification system for the exemplary pathways described in Examples VII and IX for production of 1,4-butanediol (1,4-BDO), 1,3-butanediol (1,3-BDO) or methylacrylic acid (MAA). Exemplary enzymes for production of isopropanol from acetyl-CoA are described in Example I and exemplary enzymes for production acetyl-CoA from glucose are described in Example II.

10 PEP Carboxykinase

Although the net conversion of phosphoenolpyruvate to oxaloacetate is redox-neutral, the mechanism of this conversion is important to the overall energetics of the co- production pathway. The most desirable enzyme for the conversion of PEP to oxaloacetate is PEP carboxykinase which simultaneously forms an ATP while carboxylating PEP. In most
 15 organisms, however, PEP carboxykinase serves a gluconeogenic function and converts oxaloacetate to PEP at the expense of one ATP. *S. cerevisiae* is one such organism whose native PEP carboxykinase, *PCK1*, serves a gluconeogenic role (Valdes-Hevia et al., *FEBS. Lett.* 258:313-316 (1989)). *E. coli* is another such organism, as the role of PEP carboxykinase in producing oxaloacetate is believed to be minor when compared to PEP carboxylase, which does
 20 not form ATP, possibly due to the higher K_m for bicarbonate of PEP carboxykinase (Kim, et al., *Appl Environ Microbiol* 70:1238-1241 (2004)). Nevertheless, activity of the native *E. coli* PEP carboxykinase from PEP towards oxaloacetate has been recently demonstrated in *ppc* mutants of *E. coli* K-12 (Kwon et al., *Journal of Microbiology and Biotechnology* 16:1448-1452 (2006)). These strains exhibited no growth defects and had increased succinate production at high
 25 NaHCO_3 concentrations. In some organisms, particularly rumen bacteria, PEP carboxykinase is quite efficient in producing oxaloacetate from PEP and generating ATP. Examples of PEP carboxykinase genes that have been cloned into *E. coli* include those from *Mannheimia succiniciproducens* (Lee et al., *Gene. Biotechnol. Bioprocess Eng.* 7:95-99 (2002)), *Anaerobiospirillum succiniciproducens* (Laivenieks et al., *Appl Environ Microbiol* 63:2273-
 30 2280 (1997)), and *Actinobacillus succinogenes* (Kim et al., *Appl Environ Microbiol* 70:1238-1241 (2004)). Internal experiments have also found that the PEP carboxykinase enzyme

encoded by *Haemophilus influenza* is highly efficient at forming oxaloacetate from PEP. These genes/proteins are identified below in Table 77.

Table 77.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>PCK1</i>	NP_013023	6322950	<i>Saccharomyces cerevisiae</i>
<i>pck</i>	NP_417862.1	16131280	<i>Escherichia coli</i>
<i>pckA</i>	YP_089485.1	52426348	<i>Mannheimia succiniciproducens</i>
<i>pckA</i>	O09460.1	3122621	<i>Anaerobiospirillum succiniciproducens</i>
<i>pckA</i>	Q6W6X5	75440571	<i>Actinobacillus succinogenes</i>
<i>pckA</i>	P43923.1	1172573	<i>Haemophilus influenza</i>

These sequences and sequences for subsequent enzymes listed in this report can be used to

- 5 identify homologue proteins in GenBank or other databases through sequence similarity searches (e.g. BLASTp). The resulting homologue proteins and their corresponding gene sequences provide additional DNA sequences for transformation into the host organism of our choice.

PEP Carboxylase

- 10 PEP carboxylase represents an alternative enzyme for the formation of oxaloacetate from PEP. *S. cerevisiae* does not naturally encode a PEP carboxylase, but exemplary organisms that possess genes that encode PEP carboxylase include *E. coli* (Kai et al., *Arch. BioChem. Biophys.* 414:170-179 (2003)), *Methylobacterium extorquens* AM1 (Arps et al., *J. Bacteriol.* 175:3776-3783 (1993)), and *Corynebacterium glutamicum* (Eikmanns et al., *Mol. Gen. Genet.* 218:330-15 339 (1989)). These genes/proteins are identified below in Table 78.

Table 78.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ppc</i>	NP_418391	16131794	<i>Escherichia coli</i>
<i>ppcA</i>	AAB58883	28572162	<i>Methylobacterium extorquens</i>
<i>ppc</i>	ABB53270	80973080	<i>Corynebacterium glutamicum</i>

Pyruvate Kinase and Methylmalonyl-CoA Carboxytransferase

- 20 An additional energetically efficient route to oxaloacetate from PEP requires two enzymatic activities: pyruvate kinase and methylmalonyl-CoA carboxytransferase. Pyruvate kinase catalyzes the ATP-generating conversion of PEP to pyruvate and is encoded by the *PYK1*

(Burke et al., *J. Biol. Chem.* 258:2193-2201 (1983)) and *PYK2* (Boles et al., *J. Bacteriol.* 179:2987-2993 (1997)) genes in *S. cerevisiae*. In *E. coli*, this activity is catalyzed by the gene product of *pykF* and *pykA*. Methylmalonyl-CoA carboxytransferase catalyzes the conversion of pyruvate to oxaloacetate. Importantly, this reaction also simultaneously catalyzes the conversion of (S)-methylmalonyl-CoA to propionyl-CoA (see Figures 1 and 2). An exemplary methylmalonyl-CoA carboxytransferase which is comprised of 1.3S, 5S, and 12S subunits can be found in *Propionibacterium freudenreichii* (Thornton et al., *J. Bacteriol.* 175:5301-5308 (1993)). These genes/proteins are identified below in Table 79.

Table 79.

Gene	GenBank ID	GI Number	Organism
<i>PYK1</i>	NP_009362	6319279	<i>Saccharomyces cerevisiae</i>
<i>PYK2</i>	NP_014992	6324923	<i>Saccharomyces cerevisiae</i>
<i>pykF</i>	NP_416191.1	16129632	<i>Escherichia coli</i>
<i>pykA</i>	NP_416368.1	16129807	<i>Escherichia coli</i>
1.3S subunit	P02904	114847	<i>Propionibacterium freudenreichii</i>
5S subunit	Q70AC7	62901478	<i>Propionibacterium freudenreichii</i>
12S subunit	Q8GBW6	62901481	<i>Propionibacterium freudenreichii</i>

Pyruvate Kinase and Pyruvate Carboxylase

A combination of enzymes can convert PEP to oxaloacetate with a stoichiometry identical to that of PEP carboxylase. These enzymes are encoded by pyruvate kinase, *PYK1* (Burke et al., *J. Biol. Chem.* 258:2193-2201 (1983)) or *PYK2* (Boles et al., *J. Bacteriol.* 179:2987-2993 (1997)), and pyruvate carboxylase, *PYC1* (Walker et al., *BioChem. Biophys. Res. Commun.* 176:1210-1217 (1991)) or *PYC2* (224). Some candidates for pyruvate carboxylase function are identified below in Table 80.

Table 80.

Gene	GenBank ID	GI Number	Organism
<i>PYC1</i>	NP_011453	6321376	<i>Saccharomyces cerevisiae</i>
<i>PYC2</i>	NP_009777	6319695	<i>Saccharomyces cerevisiae</i>
<i>Pyc</i>	YP_890857.1	118470447	<i>Mycobacterium smegmatis</i>

Malate Dehydrogenase, Fumarase, Fumarate Reductase

Oxaloacetate can be converted to succinate by malate dehydrogenase, fumarase and fumarate reductase when the TCA cycle is operating in the reductive cycle. *S. cerevisiae* possesses three

copies of malate dehydrogenase, *MDH1* (McAlister-Henn et al., *J. Bacteriol.* 169:5157-5166 (1987)), *MDH2* (Gibson *J. Biol. Chem.* 278:25628-25636 (2003); and Minard et al., *Mol. Cell Biol.* 11:370-380 (1991)), and *MDH3* (Steffan et al., *J. Biol. Chem.* 267:24708-24715 (1992)), which localize to the mitochondrion, cytosol, and peroxisome, respectively. *S. cerevisiae*

5 contains one copy of a fumarase-encoding gene, *FUM1*, whose product localizes to both the cytosol and mitochondrion (Sass et al., *J. Biol. Chem.* 278:45109-45116 (2003)). Fumarate reductase is encoded by two soluble enzymes, *FRDS1* (Enomoto et al., *DNA. Res.* 3:263-267 (1996)) and *FRDS2* (Muratsubaki et al., *Arch. BioChem. Biophys.* 352:175-181 (1998)), which

10 localize to the cytosol and promitochondrion, respectively, and are required for anaerobic growth on glucose (Arikawa et al., *Microbiol Lett.* 165:111-116 (1998)). *E. coli* is known to have an active malate dehydrogenase. It has three fumarases encoded by *fumA*, *B* and *C*, each one of which is active under different conditions of oxygen availability. The fumarate reductase in *E. coli* is composed of four subunits. These genes/proteins are identified below in Table 81.

Table 81.

Gene	GenBank ID	GI Number	Organism
<i>MDH1</i>	NP_012838	6322765	<i>Saccharomyces cerevisiae</i>
<i>MDH2</i>	NP_014515	116006499	<i>Saccharomyces cerevisiae</i>
<i>MDH3</i>	NP_010205	6320125	<i>Saccharomyces cerevisiae</i>
<i>FUM1</i>	NP_015061	6324993	<i>Saccharomyces cerevisiae</i>
<i>FRDS1</i>	P32614	418423	<i>Saccharomyces cerevisiae</i>
<i>FRDS2</i>	NP_012585	6322511	<i>Saccharomyces cerevisiae</i>
<i>frdA</i>	NP_418578.1	16131979	<i>Escherichia coli</i>
<i>frdB</i>	NP_418577.1	16131978	<i>Escherichia coli</i>
<i>frdC</i>	NP_418576.1	16131977	<i>Escherichia coli</i>
<i>frdD</i>	NP_418475.1	16131877	<i>Escherichia coli</i>
<i>Mdh</i>	NP_417703.1	16131126	<i>Escherichia coli</i>
<i>FumA</i>	NP_416129.1	16129570	<i>Escherichia coli</i>
<i>FumB</i>	NP_418546.1	16131948	<i>Escherichia coli</i>
<i>FumC</i>	NP_416128.1	16129569	<i>Escherichia coli</i>

15

Succinyl-CoA Transferase

Succinyl-CoA transferase catalyzes the conversion of succinyl-CoA to succinate while transferring the CoA moiety to a CoA acceptor molecule. Many transferases have broad specificity and thus may utilize CoA acceptors as diverse as acetate, succinate, propionate,

20 butyrate, 2-methylacetoacetate, 3-ketohexanoate, 3-ketopentanoate, valerate, crotonate, 3-mercaptopropionate, propionate, vinylacetate, butyrate, among others.

The conversion of succinate to succinyl-CoA is ideally carried by a transferase which does not require the direct consumption of an ATP or GTP. This type of reaction is common in a number of organisms. Perhaps the top candidate enzyme for this reaction step is succinyl-CoA:3-ketoacid-CoA transferase. This enzyme converts succinate to succinyl-CoA while converting a 3-ketoacyl-CoA to a 3-ketoacid. Exemplary succinyl-CoA:3-ketoacid-CoA transferases are present in *Helicobacter pylori* (Corthesy-Theulaz et al., *J. Biol. Chem.* 272:25659-25667 (1997)), *Bacillus subtilis* (Stols et al., *Protein. Expr. Purif.* 53:396-403 (2007)), and *Homo sapiens* (Fukao et al., *Genomics.* 68:144-151 (2000); and Tanaka et al., *Mol. Hum. Reprod.* 8:16-23 (2002)). These genes/proteins are identified below in Table 82.

10

Table 82.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>HPAG1_0676</i>	YP_627417	108563101	<i>Helicobacter pylori</i>
<i>HPAG1_0677</i>	YP_627418	108563102	<i>Helicobacter pylori</i>
<i>ScoA</i>	NP_391778	16080950	<i>Bacillus subtilis</i>
<i>ScoB</i>	NP_391777	16080949	<i>Bacillus subtilis</i>
<i>OXCT1</i>	NP_000427	4557817	<i>Homo sapiens</i>
<i>OXCT2</i>	NP_071403	11545841	<i>Homo sapiens</i>

The conversion of succinate to succinyl-CoA can also be catalyzed by succinyl-CoA: Acetyl-CoA transferase. The gene product of *catI* of *Clostridium kluyveri* has been shown to exhibit succinyl-CoA: acetyl-CoA transferase activity (Sohling et al., *J. Bacteriol.* 178:871-880 (1996)).

15 In addition, the activity is present in *Trichomonas vaginalis* (van Grinsven et al., *J. Biol. Chem.* 283:1411-1418 (2008)) and *Trypanosoma brucei* (Riviere et al., *J. Biol. Chem.* 279:45337-45346 (2004)). These genes/proteins are identified below in Table 83.

Table 83.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>catI</i>	P38946.1	729048	<i>Clostridium kluyveri</i>
<i>TVAG_395550</i>	XP_001330176	123975034	<i>Trichomonas vaginalis</i> G3
<i>Tb11.02.0290</i>	XP_828352	71754875	<i>Trypanosoma brucei</i>

Yet another possible CoA acceptor is benzylsuccinate. Succinyl-CoA:(R)-Benzylsuccinate CoA-Transferase functions as part of an anaerobic degradation pathway for toluene in organisms such as *Thauera aromatica* (Leutwein and Heider, *J. Bact.* 183(14) 4288-4295 (2001)). Homologs can be found in *Azoarcus* sp. T, *Aromatoleum aromaticum* EbN1, and *Geobacter metallireducens* GS-15. These genes/proteins are identified below in Table 84.

20

Table 84.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>bbsE</i>	AAF89840	9622535	<i>Thauera aromatica</i>
<i>bbsf</i>	AAF89841	9622536	<i>Thauera aromatica</i>
<i>bbsE</i>	AAU45405.1	52421824	<i>Azoarcus sp. T</i>
<i>bbsF</i>	AAU45406.1	52421825	<i>Azoarcus sp. T</i>
<i>bbsE</i>	YP_158075.1	56476486	<i>Aromatoleum aromaticum EbN1</i>
<i>bbsF</i>	YP_158074.1	56476485	<i>Aromatoleum aromaticum EbN1</i>
<i>Gmet_1521</i>	YP_384480.1	78222733	<i>Geobacter metallireducens GS-15</i>
<i>Gmet_1522</i>	YP_384481.1	78222734	<i>Geobacter metallireducens GS-15</i>

Finally, *ygfH* encodes a propionyl CoA:succinate CoA transferase in *E. coli* (Haller et al., *Biochemistry*, 39(16) 4622-4629). Close homologs can be found in, for example, *Citrobacter youngae* ATCC 29220, *Salmonella enterica* subsp. *arizonae* serovar, and *Yersinia intermedia*

5 ATCC 29909. These genes/proteins are identified below in Table 85.

Table 85.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ygfH</i>	NP_417395.1	16130821	<i>Escherichia coli str. K-12 substr. MG1655</i>
<i>CIT292_04485</i>	ZP_03838384.1	227334728	<i>Citrobacter youngae</i> ATCC 29220
<i>SARI_04582</i>	YP_001573497.1	161506385	<i>Salmonella enterica</i> subsp. <i>arizonae</i> serovar
<i>yinte0001_14430</i>	ZP_04635364.1	238791727	<i>Yersinia intermedia</i> ATCC 29909

Succinyl-CoA Synthetase

The product of the *LSC1* and *LSC2* genes of *S. cerevisiae* and the *sucC* and *sucD* genes of *E. coli* naturally form a succinyl-CoA synthetase complex that catalyzes the formation of succinyl-

10 CoA from succinate with the concomitant consumption of one ATP, a reaction which is reversible *in vivo* (Bravo et al., *J. Forensic Sci.* 49:379-387 (2004)). These genes/proteins are identified below in Table 86.

Table 86.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>LSC1</i>	NP_014785	6324716	<i>Saccharomyces cerevisiae</i>
<i>LSC2</i>	NP_011760	6321683	<i>Saccharomyces cerevisiae</i>
<i>sucC</i>	NP_415256.1	16128703	<i>Escherichia coli</i>
<i>sucD</i>	AAC73823.1	1786949	<i>Escherichia coli</i>

Pyruvate Formate Lyase

Pyruvate formate lyase is an enzyme that catalyzes the conversion of pyruvate and CoA into acetyl-CoA and formate. Pyruvate formate lyase is a common enzyme in prokaryotic organisms that is used to help modulate anaerobic redox balance. Exemplary enzymes can be found in

5 *Escherichia coli* (Knappe et al., *FEMS. Microbiol Rev.* 6:383-398 (1990)), *Lactococcus lactis* (Melchiorson et al., *Appl Microbiol Biotechnol* 58:338-344 (2002)), and *Streptococcus mutans* (Takahashi-Abbe et al., *Oral. Microbiol Immunol.* 18:293-297 (2003)). A mitochondrial pyruvate formate lyase has also been identified in the eukaryote, *Chlamydomonas reinhardtii* (Atteia et al., *J. Biol. Chem.* 281:9909-9918 (2006); and Hemschemeier et al., *Eukaryot. Cell*

10 7:518-526 (2008)). These genes/proteins are identified below in Table 87.

Table 87.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pflB</i>	NP_415423	16128870	<i>Escherichia coli</i>
<i>pfl</i>	CAA03993	2407931	<i>Lactococcus lactis</i>
<i>pfl</i>	BAA09085	1129082	<i>Streptococcus mutans</i>
<i>PFL1</i>	EDP09457	158283707	<i>Chlamydomonas reinhardtii</i>

Formate Hydrogen Lyase

A formate hydrogen lyase enzyme can be employed to convert formate to carbon dioxide and hydrogen. An exemplary formate hydrogen lyase enzyme can be found in *Escherichia coli*. The

15 *E. coli* formate hydrogen lyase consists of hydrogenase 3 and formate dehydrogenase-H (Maeda et al., *Appl Microbiol Biotechnol* 77:879-890 (2007)). It is activated by the gene product of *fhlA* (Maeda et al., *Appl Microbiol Biotechnol* 77:879-890 (2007)). The addition of the trace elements, selenium, nickel and molybdenum, to a fermentation broth has been shown to enhance formate hydrogen lyase activity (Soini, et al., *Microb. Cell Fact.* 7:26 (2008)). These

20 genes/proteins are identified below in Table 88.

Table 88.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
Hydrogenase 3:			
<i>hycD</i>	NP_417202	16130629	<i>Escherichia coli</i>
<i>hycC</i>	NP_417203	16130630	<i>Escherichia coli</i>
<i>hycF</i>	NP_417200	16130627	<i>Escherichia coli</i>
<i>hycG</i>	NP_417199	16130626	<i>Escherichia coli</i>
<i>hycB</i>	NP_417204	16130631	<i>Escherichia coli</i>
<i>hycE</i>	NP_417201	16130628	<i>Escherichia coli</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
Formate dehydrogenase-H:			
<i>fdhF</i>	NP_418503	16131905	<i>Escherichia coli</i>
Activator:			
<i>fhlA</i>	NP_417211	16130638	<i>Escherichia coli</i>

A formate hydrogen lyase enzyme also exists in the hyperthermophilic archaeon, *Thermococcus litoralis* (Takacs et al., *Microbiol* 8:88 2008)). These genes/proteins are identified below in Table 89.

Table 89.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mhyC</i>	ABW05543	157954626	<i>Thermococcus litoralis</i>
<i>mhyD</i>	ABW05544	157954627	<i>Thermococcus litoralis</i>
<i>mhyE</i>	ABW05545	157954628	<i>Thermococcus litoralis</i>
<i>myhF</i>	ABW05546	157954629	<i>Thermococcus litoralis</i>
<i>myhG</i>	ABW05547	157954630	<i>Thermococcus litoralis</i>
<i>myhH</i>	ABW05548	157954631	<i>Thermococcus litoralis</i>
<i>fdhA</i>	AAB94932	2746736	<i>Thermococcus litoralis</i>
<i>fdhB</i>	AAB94931	157954625	<i>Thermococcus litoralis</i>

- 5 Additional formate hydrogen lyase systems have been found in *Salmonella typhimurium*, *Klebsiella pneumoniae*, *Rhodospirillum rubrum*, *Methanobacterium formicicum* (Vardar-Schara et al., *Microbial Biotechnology* 1:107-125)).

Formate Dehydrogenase

- Formate dehydrogenase activity is present in both *E. coli* and *Saccharomyces cerevisiae* among other organisms. *S. cerevisiae* contains two formate dehydrogenases, *FDH1* and *FDH2*, that catalyze the oxidation of formate to CO₂ (Overkamp et al., *Yeast* 19:509-520 (2002)). In *Moorella thermoacetica*, the loci, Moth_2312 and Moth_2313, are actually one gene that is responsible for encoding the alpha subunit of formate dehydrogenase while the beta subunit is encoded by Moth_2314 (Andreesen et al., *J. Bacteriol.* 116:867-873 (1973); Li et al., *J. Bacteriol.* 92:405-412 (1966); Pierce et al., *Environ. Microbiol* (2008) and Yamamoto et al., *J. Biol. Chem.* 258:1826-1832 (1983)). Another set of genes encoding formate dehydrogenase activity is encoded by Sfum_2703 through Sfum_2706 in *Syntrophobacter fumaroxidans* (de Bok, et al., *Eur. J. BioChem.* 270:2476-2485 (2003); and Reda et al., *Proc. Natl. Acad. Sci. U S A.* 105:10654-10658 (2008)). Similar to their *M. thermoacetica* counterparts, Sfum_2705 and Sfum_2706 are actually one gene. *E. coli* contains multiple formate dehydrogenases. These genes/proteins are identified below in Table 90.

Table 90.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>FDH1</i>	NP_015033	6324964	<i>Saccharomyces cerevisiae</i>
<i>FDH2</i>	Q08987	88909613	<i>Saccharomyces cerevisiae</i>
<i>Moth_2312</i>	YP_431142	148283121	<i>Moorella thermoacetica</i>
<i>Moth_2313</i>	YP_431143	83591134	<i>Moorella thermoacetica</i>
<i>Moth_2314</i>	YP_431144	83591135	<i>Moorella thermoacetica</i>
<i>Sfum_2703</i>	YP_846816.1	116750129	<i>Syntrophobacter fumaroxidans</i>
<i>Sfum_2704</i>	YP_846817.1	116750130	<i>Syntrophobacter fumaroxidans</i>
<i>Sfum_2705</i>	YP_846818.1	116750131	<i>Syntrophobacter fumaroxidans</i>
<i>Sfum_2706</i>	YP_846819.1	116750132	<i>Syntrophobacter fumaroxidans</i>
<i>fdnG, H, I</i>	NP_415991-993.1	16129433 16129434 16129435	<i>Escherichia coli</i>
<i>fdoG, H, I</i>	NP_418330,29,28.1	16131734 16131733 16131732	<i>Escherichia coli</i>

Pyruvate Dehydrogenase

The pyruvate dehydrogenase complex, catalyzing the conversion of pyruvate to acetyl-CoA, has been extensively studied. The *S. cerevisiae* complex consists of an E2 (*LAT1*) core that binds E1 (*PDA1*, *PDB1*), E3 (*LPD1*), and Protein X (*PDX1*) components (Pronk et al., *Yeast* 12:1607-1633 (1996)). In the *E. coli* enzyme, specific residues in the E1 component are responsible for substrate specificity (Bisswanger *J Biol Chem.* 256:815-822. (1981); Bremer *J BioChem.* 8:535-540 (1969) and Gong et al., *J Biol Chem.* 275:13645-13653 (2000)). Engineering efforts have improved the *E. coli* PDH enzyme activity under anaerobic conditions (Kim et al., *Appl. Environ. Microbiol.* 73:1766-1771 (2007); Kim et al., *J. Bacteriol.* 190:3851-3858 (2008) and Zhou et al., *Biotechnol. Lett.* 30:335-342 (2008)). In contrast to the *E. coli* PDH, the *B. subtilis* complex is active and required for growth under anaerobic conditions (Nakano et al., *J. Bacteriol.* 179:6749-6755 (1997)). The *Klebsiella pneumoniae* PDH, characterized during growth on glycerol, is also active under anaerobic conditions (Menzel et al., *J. Biotechnol.* 56:135-142 (1997)). Crystal structures of the enzyme complex from bovine kidney (Zhou et al., *Proc. Natl. Acad. Sci. U. S. A* 98:14802-14807 (2001)) and the E2 catalytic domain from *Azotobacter vinelandii* are available (Mattevi et al., *Science.* 255:1544-1550 (1992)). Some maMAAlian PDH enzymes complexes can react on alternate substrates such as 2-oxobutanoate (Paxton et al., *BioChem. J.* 234:295-303 (1986)). These genes/proteins are identified below in Table 91.

Table 91.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>LAT1</i>	NP_014328	6324258	<i>Saccharomyces cerevisiae</i>
<i>PDA1</i>	NP_011105	37362644	<i>Saccharomyces cerevisiae</i>
<i>PDB1</i>	NP_009780	6319698	<i>Saccharomyces cerevisiae</i>
<i>LPD1</i>	NP_116635	14318501	<i>Saccharomyces cerevisiae</i>
<i>PDX1</i>	NP_011709	6321632	<i>Saccharomyces cerevisiae</i>
<i>aceE</i>	NP_414656.1	16128107	<i>Escherichia coli str. K12 substr. MG1655</i>
<i>aceF</i>	NP_414657.1	16128108	<i>Escherichia coli str. K12 substr. MG1655</i>
<i>Lpd</i>	NP_414658.1	16128109	<i>Escherichia coli str. K12 substr. MG1655</i>
<i>pdhA</i>	P21881.1	3123238	<i>Bacillus subtilis</i>
<i>pdhB</i>	P21882.1	129068	<i>Bacillus subtilis</i>
<i>pdhC</i>	P21883.2	129054	<i>Bacillus subtilis</i>
<i>pdhD</i>	P21880.1	118672	<i>Bacillus subtilis</i>
<i>aceE</i>	YP_001333808.1	152968699	<i>Klebsiella pneumonia MGH78578</i>
<i>aceF</i>	YP_001333809.1	152968700	<i>Klebsiella pneumonia MGH78578</i>
<i>lpdA</i>	YP_001333810.1	152968701	<i>Klebsiella pneumonia MGH78578</i>
<i>Pdha1</i>	NP_001004072.2	124430510	<i>Rattus norvegicus</i>
<i>Pdha2</i>	NP_446446.1	16758900	<i>Rattus norvegicus</i>
<i>Dlat</i>	NP_112287.1	78365255	<i>Rattus norvegicus</i>
<i>Dld</i>	NP_955417.1	40786469	<i>Rattus norvegicus</i>

Pyruvate Ferredoxin Oxidoreductase

Pyruvate ferredoxin oxidoreductase (PFOR) catalyzes the oxidation of pyruvate to form acetyl-CoA. The PFOR from *Desulfovibrio africanus* has been cloned and expressed in *E. coli* resulting in an active recombinant enzyme that was stable for several days in the presence of oxygen (Pieulle et al., *J Bacteriol.* 179:5684-5692 (1997)). Oxygen stability is relatively uncommon in PFORs and is believed to be conferred by a 60 residue extension in the polypeptide chain of the *D. africanus* enzyme. The *M. thermoacetica* PFOR is also well characterized (Menon et al., *BioChemistry* 36:8484-8494 (1997)) and was even shown to have high activity in the direction of pyruvate synthesis during autotrophic growth (Furdui et al., *J Biol Chem.* 275:28494-28499 (2000)). Further, *E. coli* possesses an uncharacterized open reading frame, *ydbK*, encoding a protein that is 51% identical to the *M. thermoacetica* PFOR. Evidence for pyruvate oxidoreductase activity in *E. coli* has been described (Blaschkowski et al., *J BioChem.* 123:563-569 (1982)). Several additional PFOR enzymes are described in the following review (Ragsdale, *Chem. Rev.* 103:2333-2346 (2003)). Finally, flavodoxin reductases (e.g., *fqrB* from *Helicobacter pylori* or *Campylobacter jejuni* (St Maurice et al., *J. Bacteriol.*

189:4764-4773 (2007)) or Rnf-type proteins (Herrmann et al., *J. Bacteriol.* 190:784-791 (2008); and Seedorf et al., *Proc. Natl. Acad. Sci. U S A.* 105:2128-2133 (2008)) provide a means to generate NADH or NADPH from the reduced ferredoxin generated by PFOR. These genes/proteins are identified below in Table 92.

5

Table 92.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>Por</i>	CAA70873.1	1770208	<i>Desulfovibrio africanus</i>
<i>Por</i>	YP_428946.1	83588937	<i>Moorella thermoacetica</i>
<i>ydbK</i>	NP_415896.1	16129339	<i>Escherichia coli</i>
<i>fqrB</i>	NP_207955.1	15645778	<i>Helicobacter pylori</i>
<i>fqrB</i>	YP_001482096.1	157414840	<i>Campylobacter jejuni</i>
<i>RnfC</i>	EDK33306.1	146346770	<i>Clostridium kluyveri</i>
<i>RnfD</i>	EDK33307.1	146346771	<i>Clostridium kluyveri</i>
<i>RnfG</i>	EDK33308.1	146346772	<i>Clostridium kluyveri</i>
<i>RnfE</i>	EDK33309.1	146346773	<i>Clostridium kluyveri</i>
<i>RnfA</i>	EDK33310.1	146346774	<i>Clostridium kluyveri</i>
<i>RnfB</i>	EDK33311.1	146346775	<i>Clostridium kluyveri</i>

Succinic semialdehyde dehydrogenase (CoA-dependent)

Succinic semialdehyde dehydrogenase (CoA-dependent), also referred to as succinyl-CoA reductase, is a CoA- and NAD(P)H- dependent oxidoreductase that reduces succinyl-CoA to its corresponding aldehyde. Exemplary enzymes are encoded by the *sucD* gene in *Clostridium kluyveri* (Sohling et al., *J Bacteriol* 178:871-80 (1996); and Sohling et al., *J Bacteriol.* 178:871-880 (1996)) and the *sucD* gene of *P. gingivalis* (Takahashi et al., *J. Bacteriol.* 182:4704-4710 (2000)). Other enzymes that catalyze similar reactions are the fatty acyl-CoA reductases of *Acinetobacter calcoaceticus* (Reiser et al., *Journal of Bacteriology* 179:2969-2975 (2007)) and *Acinetobacter sp. M-1* (Ishige et al., *Appl. Environ. Microbiol.* 68:1192-1195 (2002)), and the acylating acetaldehyde dehydrogenase in *Pseudomonas sp.*, which has been demonstrated to oxidize and acylate acetaldehyde, propionaldehyde, butyraldehyde, isobutyraldehyde and formaldehyde (Powlowski et al., *J Bacteriol.* 175:377-385 (1993)). In addition to reducing acetyl-CoA to ethanol, the enzyme encoded by *adhE* in *Leuconostoc mesenteroides* has been shown to oxidize the branched chain compound isobutyraldehyde to isobutyryl-CoA (Koo et al., *Biotechnol Lett.* 27:505-510 (2005)). These genes/proteins are identified below in Table 93.

Table 93.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>sucD</i>	P38947.1	172046062	<i>Clostridium kluyveri</i>
<i>sucD</i>	NP_904963.1	34540484	<i>Porphyromonas gingivalis</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>acrI</i>	YP_047869.1	50086359	<i>Acinetobacter calcoaceticus</i>
<i>acrI</i>	AAC45217	1684886	<i>Acinetobacter baylyi</i>
<i>acrI</i>	BAB85476.1	18857901	<i>Acinetobacter</i> sp. Strain M-1
<i>bphG</i>	BAA03892.1	425213	<i>Pseudomonas</i> sp
<i>adhE</i>	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>

4-Hydroxybutyrate dehydrogenase

4-Hydroxybutyrate dehydrogenase catalyzes the NAD(P)H dependent reduction of succinic semialdehyde to 4-HB. Enzymes exhibiting this activity are found in *Ralstonia eutropha* (Bravo et al., *J. Forensic Sci.* 49:379-387 (2004)), *Clostridium kluyveri* (Wolff et al., *Protein Expr. Purif.* 6:206-212 (1995)) and *Arabidopsis thaliana* (Breitkreuz et al., *J. Biol. Chem.* 278:41552-41556 (2003)). Yet another gene is the alcohol dehydrogenase *adhI* from *Geobacillus thermoglucosidasius* (Jeon et al., *J Biotechnol* 135:127-133 (2008)). These genes/proteins are identified below in Table 94.

Table 94.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>4hbd</i>	YP_726053.1	113867564	<i>Ralstonia eutropha</i> H16
<i>4hbd</i>	EDK35022.1	146348486	<i>Clostridium kluyveri</i>
<i>4hbd</i>	Q94B07	75249805	<i>Arabidopsis thaliana</i>
<i>adhI</i>	AAR91477.1	40795502	<i>Geobacillus thermoglucosidasius</i>

4-Hydroxybutyryl-CoA transferase

The conversion of 4-HB to 4-hydroxybutyryl-CoA is catalyzed by an enzyme with 4-hydroxybutyryl-CoA transferase activity. Candidate enzymes include the gene products of *catI*, *cat2*, and *cat3* of *Clostridium kluyveri* which have been shown to exhibit succinyl-CoA, 4-hydroxybutyryl-CoA, and butyryl-CoA transferase activity, respectively (Gerhardt et al., *Arch. Microbiol* 174:189-199 (2000); Arikawa et al., *Microbiol Lett.* 165:111-116 (1998) and Sohling et al., *J Bacteriol.* 178:871-880 (1996)). Similar CoA transferase activities are also present in *Trichomonas vaginalis* (van Grinsven et al., *J. Biol. Chem.* 283:1411-1418 (2008)) and *Trypanosoma brucei* (Riviere et al., *J. Biol. Chem.* 279:45337-45346 (2004)). The *atoA* and *atoD* genes of *E. coli* encode an acetoacetyl-CoA transferase with a broad substrate range (Sramek et al., *Arch. BioChem. Biophys.* 171:14-26 (1975)). This enzyme has been shown to transfer a CoA moiety from acetyl-CoA to a variety of branched and linear substrates including isobutyrate (Matthies et al., *Appl Environ. Microbiol* 58:1435-1439 (1992)), valerate

(Vanderwinkel et al., *BioChem. Biophys. Res. Commun.* 33:902-908 (1968)) and butanoate (Vanderwinkel et al., *BioChem. Biophys. Res. Commun.* 33:902-908 (1968)). These genes/proteins are identified below in Table 95.

Table 95.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>cat1</i>	P38946.1	729048	<i>Clostridium kluyveri</i>
<i>cat2</i>	P38942.2	172046066	<i>Clostridium kluyveri</i>
<i>cat3</i>	EDK35586.1	146349050	<i>Clostridium kluyveri</i>
<i>TVAG_395550</i>	XP_001330176	123975034	<i>Trichomonas vaginalis G3</i>
<i>Tb11.02.0290</i>	XP_828352	71754875	<i>Trypanosoma brucei</i>
<i>atoA</i>	P76459.1	2492994	<i>Escherichia coli</i>
<i>atoD</i>	P76458.1	2492990	<i>Escherichia coli</i>

5

4-Hydroxybutyryl-CoA synthetase

The conversion of 4-HB to 4-hydroxybutyryl-CoA can also be catalyzed by a CoA acid-thiol ligase, also known as a CoA synthetase. Enzymes catalyzing this exact transformation have not been characterized to date; however, several enzymes with broad substrate specificities have been described in the literature. An exemplary candidate is the enzyme encoded by *sucCD* in *E. coli*, which naturally catalyzes the formation of succinyl-CoA from succinate with the concomitant consumption of one ATP, a reaction which is reversible *in vivo* (Buck et al., *BioChemistry* 24:6245-6252 (1985)). Additional CoA-ligase candidates include the ADP-forming phenylacetate-CoA ligases from *P. chrysogenum* (Lamas-Maceiras et al., *BioChem. J* 395:147-155 (2006); and Wang et al., *BioChem. Biophys. Res. Commun.* 360:453-458 (2007)) and the pimeloyl-CoA ligase from *Pseudomonas mendocina*. The AMP-forming enzyme from *Pseudomonas mendocina*, cloned into *E. coli*, was shown to accept the alternate substrates hexanedioate and nonanedioate (Binieda et al., *BioChem. J* 340 (Pt 3):793-801 (1999)). These genes/proteins are identified below in Table 96. CoA synthetase enzyme candidates identified for acetoacetyl-CoA synthetase, succinyl-CoA synthetase, propionyl-CoA synthetase, 3-hydroxybutyryl-CoA synthetase, 3-hydroxyisobutyryl-CoA synthetase, methylmalonyl-CoA synthetase and methacrylyl-CoA synthase are also applicable here.

20

Table 96.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>sucC</i>	NP_415256.1	16128703	<i>Escherichia coli</i>
<i>sucD</i>	AAC73823.1	1786949	<i>Escherichia coli</i>
<i>phl</i>	CAJ15517.1	77019264	<i>Penicillium chrysogenum</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>pauA</i>	NP_249708.1	15596214	<i>Pseudomonas mendocina</i>

4-Hydroxybutyryl-CoA reductase (aldehyde forming)

4-Hydroxybutyryl-CoA reductase catalyzes the NAD(P)H dependent reduction of 4-hydroxybutyryl-CoA to 4-hydroxybutyraldehyde. Enzymes that exhibit this activity include succinate semialdehyde dehydrogenase enzymes encoded by the *sucD* gene in *Clostridium kluyveri* (Sohling et al., *J Bacteriol* 178:871-80 (1996); and Sohling et al., *J Bacteriol.* 178:871-880 (1996)) and *sucD* of *P. gingivalis* (Takahashi et al., *J. Bacteriol.* 182:4704-4710 (2000)). Butyraldehyde dehydrogenase enzymes, found in solventogenic organisms such as *Clostridium saccharoperbutylacetonicum* (Kosaka et al., *Biosci. Biotechnol BioChem.* 71:58-68 (2007)), catalyzes a similar reaction: conversion of butyryl-CoA to butyraldehyde. The enzyme acylating acetaldehyde dehydrogenase in *Pseudomonas sp*, encoded by *bphG*, is yet another candidate as it has been demonstrated to oxidize and acylate acetaldehyde, propionaldehyde, butyraldehyde, isobutyraldehyde and formaldehyde (Powlowski et al., *J Bacteriol.* 175:377-385 (1993)). Fatty acyl-CoA reductase enzymes from *Acinetobacter calcoaceticus* (Reiser et al., *Journal of Bacteriology* 179:2969-2975 (1997)) and the *Acinetobacter sp. M-1* (Ishige, et al., *Appl. Environ. Microbiol.* 68:1192-1195 (2002)) catalyze similar reactions. In addition to reducing acetyl-CoA to ethanol, the enzyme encoded by *adhE* in *Leuconostoc mesenteroides* has been shown to oxidize the branched chain compound isobutyraldehyde to isobutyryl-CoA (Koo et al., *Biotechnol Lett.* 27:505-510 (2005)). These genes/proteins are identified below in Table 97.

Table 97.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>sucD</i>	P38947.1	172046062	<i>Clostridium kluyveri</i>
<i>sucD</i>	NP_904963.1	34540484	<i>Porphyromonas gingivalis</i>
<i>bld</i>	AAP42563.1	31075383	<i>Clostridium saccharoperbutylacetonicum</i>
<i>bphG</i>	BAA03892.1	425213	<i>Pseudomonas sp</i>
<i>acrI</i>	YP_047869.1	50086359	<i>Acinetobacter calcoaceticus</i>
<i>acrI</i>	AAC45217	1684886	<i>Acinetobacter baylyi</i>
<i>acrI</i>	BAB85476.1	18857901	<i>Acinetobacter sp. Strain M-1</i>
<i>adhE</i>	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>

4-hydroxybutyraldehyde reductase

The conversion of 4-hydroxybutyrylaldehyde to 14-BDO is catalyzed by an alcohol dehydrogenase. Several native dehydrogenases in *E. coli* such as *yqhD* (Sulzenbacher et al., *Journal of Molecular Biology* 342:489-502 (2004)) exhibit broad substrate specificity and are able to catalyze this reaction. The gene product of *yqhD* catalyzes the reduction of acetaldehyde, malondialdehyde, propionaldehyde, butyraldehyde, and acrolein using NADPH as the cofactor (Perez et al., *J Biol. Chem.* 283:7346-7353 (2008); and Perez et al., *J Biol. Chem.* 283:7346-7353 (2008)). Additional enzyme candidates that catalyze the conversion of an aldehyde to alcohol include *alrA* encoding a medium-chain alcohol dehydrogenase for C2-C14 (Tani et al., *Appl. Environ. Microbiol.* 66:5231-5235 (2000)), ADH2 from *Saccharomyces cerevisiae* (Atsumi et al., *Nature* 451:86-89 (2008)) and *bdh* I and *bdh* II from *C. acetobutylicum* which converts butyraldehyde into butanol (Walter et al., *Journal of Bacteriology* 174:7149-7158 (1992)). The *adhA* gene product from *Zymomonas mobilis* has been demonstrated to have activity on a number of aldehydes including formaldehyde, acetaldehyde, propionaldehyde, butyraldehyde, and acrolein (Kinoshita et al., *Appl Microbiol Biotechnol* 22:249-254 (1985)). These genes/proteins are identified below in Table 98.

Table 98.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>yqhD</i>	NP_417484.1	16130909	<i>Escherichia coli</i>
<i>alrA</i>	BAB12273.1	9967138	<i>Acinetobacter</i> sp. Strain M-1
<i>ADH2</i>	NP_014032.1	6323961	<i>Saccharomyces cerevisiae</i>
<i>bdh</i> I	NP_349892.1	15896543	<i>Clostridium acetobutylicum</i>
<i>bdh</i> II	NP_349891.1	15896542	<i>Clostridium acetobutylicum</i>
<i>adhA</i>	YP_162971.1	56552132	<i>Zymomonas mobilis</i>

4-Hydroxybutyryl-CoA reductase (alcohol forming)

The conversion of 4-hydroxybutyryl-CoA to 14-BDO can also be catalyzed by a bifunctional oxidoreductase with aldehyde dehydrogenase and alcohol dehydrogenase capabilities. For example, the *adheE2* gene product from *Clostridium acetobutylicum* converts butyryl-CoA to butanol (Fontaine et al., *J. Bacteriol.* 184:821-830 (2002)). This enzyme also accepts 4-hydroxybutyryl-CoA as a substrate. Additional bifunctional alcohol-forming reductase enzymes include the gene products of *adhE* in *Leuconostoc mesenteroides* (Kazahaya et al., *J. Gen. Appl. Microbiol.* 18:43-55 (1972); and Koo et al., *Biotechnol Lett.* 27:505-510 (2005)) and FAR from *Simmondsia chinensis* (Metz et al., *Plant Physiology* 122:635-644 (2000)). Another exemplary

enzyme is the NADPH-dependent malonyl-CoA reductase in *Chloroflexus aurantiacus* encoded by *mcr* (Hugler et al., *J. Bacteriol.* 184:2404-2410 (2002); and Strauss et al., *Eur. J. BioChem.* 215:633-643 (1993)). These genes/proteins are identified below in Table 99.

Table 99.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>adhE2</i>	AAK09379.1	12958626	<i>Clostridium acetobutylicum</i>
<i>adhE</i>	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>
<i>FAR</i>	AAD38039.1	5020215	<i>Simmondsia chinensis</i>
<i>mcr</i>	AAS20429.1	42561982	<i>Chloroflexus aurantiacus</i>

5

4-Hydroxybutyrate phosphotransferase (aka. kinase)

4-Hydroxybutyrate phosphotransferase, also known as 4-hydroxybutyrate kinase, transforms 4-HB to 4-hydroxybutyryl phosphate with concurrent hydrolysis of one ATP. Candidate enzymes for catalyzing these transformations include butyrate kinase, aspartokinase, acetate kinase and gaMAA-glutamyl kinase. Butyrate kinase (EC 2.7.2.7) enzymes carry out the reversible conversion of butyryl-phosphate to butyrate during acidogenesis in *C. acetobutylicum* (Cary et al., *Appl. Environ. Microbiol* 56:1576-1583 (1990)). This enzyme is encoded by either of the two *buk* gene products (Huang et al., *J Mol. Microbiol Biotechnol* 2:33-38 (2000)). Other butyrate kinase enzymes are found in *C. butyricum* and *C. tetanomorphum* (TWAROG et al., *J Bacteriol.* 86:112-117 (1963)). Related enzyme isobutyrate kinase from *Thermotoga maritima* has also been expressed in *E. coli* and crystallized (Diao et al., *D. Biol. Crystallogr.* 59:1100-1102 (2003); and Diao et al., *J Bacteriol.* 191:2521-2529 (2009)). Aspartokinase catalyzes the ATP-dependent phosphorylation of aspartate and participates in the synthesis of several amino acids. The aspartokinase III enzyme in *E. coli*, encoded by *lysC*, has a broad substrate range and the catalytic residues involved in substrate specificity have been elucidated (Keng et al., *Arch. BioChem. Biophys.* 335:73-81 (1996)). Two additional kinases in *E. coli* are also good candidates: acetate kinase and gaMAA-glutamyl kinase. The *E. coli* acetate kinase, encoded by *ackA* (Skarstedt et al., *J. Biol. Chem.* 251:6775-6783 (1976)), phosphorylates propionate in addition to acetate (Hesslinger et al., *Mol. Microbiol* 27:477-492 (1998)). The *E. coli* gaMAA-glutamyl kinase, encoded by *proB* (Smith et al., *J. Bacteriol.* 157:545-551 (1984)), phosphorylates the gaMAA carbonic acid group of glutamate. These genes/proteins are identified below in Table 100.

25

Table 100.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>buk1</i>	NP_349675	15896326	<i>Clostridium acetobutylicum</i>
<i>buk2</i>	Q97II1	20137415	<i>Clostridium acetobutylicum</i>
<i>buk2</i>	Q9X278.1	6685256	<i>Thermotoga maritima</i>
<i>lysC</i>	NP_418448.1	16131850	<i>Escherichia coli</i>
<i>ackA</i>	NP_416799.1	16130231	<i>Escherichia coli</i>
<i>proB</i>	NP_414777.1	16128228	<i>Escherichia coli</i>

Phosphotrans-4-hydroxybutyrylase

Phosphotrans-4-hydroxybutyrylase exchanges the phosphate moiety of 4-hydroxybutyryl-phosphate for a CoA moiety, forming 4-hydroxybutyryl-CoA. A candidate enzyme for this transformation is phosphotransbutyrylase (EC 2.3.1.19) an enzyme that reversibly converts butyryl-CoA into butyryl-phosphate. This enzyme is encoded by *ptb* genes found in *C. acetobutylicum* (Walter et al., *Gene* 134:107-111 (1993); and Wiesenborn et al., *Appl Environ. Microbiol* 55:317-322 (1989)), butyrate-producing bacterium L2-50 (Louis et al., *J. Bacteriol.* 186:2099-2106 (2004)) and *Bacillus megaterium* (Vazquez et al., *Curr. Microbiol* 42:345-349 (2001)). These genes/proteins are identified below in Table 101.

Table 101.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ptb</i>	NP_349676	34540484	<i>Clostridium acetobutylicum</i>
<i>ptb</i>	AAR19757.1	38425288	butyrate-producing bacterium L2-50
<i>ptb</i>	CAC07932.1	10046659	<i>Bacillus megaterium</i>

4-Hydroxybutyryl-phosphate reductase

The reduction of 4-hydroxybutyryl-phosphate to its corresponding aldehyde is catalyzed by phosphate reductase. This reaction is not catalyzed by known enzymes, but a similar reaction is catalyzed by aspartate semialdehyde dehydrogenase (ASD, EC 1.2.1.11): the NADPH-dependent reduction of 4-aspartyl phosphate to aspartate-4-semialdehyde. ASD participates in amino acid biosynthesis and recently has been studied as an antimicrobial target (Hadfield et al., *Biochemistry* 40:14475-14483 (2001)). The *E. coli* ASD structure has been solved (Hadfield et al., *J Mol. Biol.* 289:991-1002 (1999)) and the enzyme has been shown to accept the alternate substrate beta-3-methylaspartyl phosphate (Shames et al., *J Biol. Chem.* 259:15331-15339 (1984)). The *Haemophilus influenzae* enzyme has been the subject of enzyme engineering studies to alter substrate binding affinities at the active site (Blanco et al., *Acta Crystallogr. D. Biol. Crystallogr.* 60:1388-1395 (2004); and Blanco et al., *Acta Crystallogr. D. Biol.*

Crystallogr. 60:1808-1815 (2004)). Other ASD candidates are found in *Mycobacterium tuberculosis* (Shafiani et al., *J Appl Microbiol* 98:832-838 (2005)), *Methanococcus jannaschii* (Faehnle et al., *J Mol. Biol.* 353:1055-1068 (2005)), and the infectious microorganisms *Vibrio cholera* and *Helicobacter pylori* (Moore et al., *Protein Expr. Purif.* 25:189-194 (2002)). A related enzyme candidate is acetylglutamylphosphate reductase (EC 1.2.1.38), an enzyme that naturally reduces acetylglutamylphosphate to acetylglutamate-5-semialdehyde, found in *S. cerevisiae* (Pauwels et al., *Eur. J Biochem.* 270:1014-1024 (2003)), *B. subtilis* (O'Reilly and Devine, *Microbiology* 140 (Pt 5):1023-1025 (1994)) and other organisms. These genes/proteins are identified below in Table 102.

10

Table 102

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>asd</i>	NP_417891.1	16131307	<i>Escherichia coli</i>
<i>asd</i>	YP_248335.1	68249223	<i>Haemophilus influenzae</i>
<i>asd</i>	AAB49996	1899206	<i>Mycobacterium tuberculosis</i>
<i>VC2036</i>	NP_231670	15642038	<i>Vibrio cholera</i>
<i>asd</i>	YP_002301787.1	210135348	<i>Helicobacter pylori</i>
<i>ARG5,6</i>	NP_010992.1	6320913	<i>Saccharomyces cerevisiae</i>
<i>argC</i>	NP_389001.1	16078184	<i>Bacillus subtilis</i>

Other exemplary phosphate reductase enzymes include glyceraldehyde 3-phosphate dehydrogenase which converts glyceraldehyde-3-phosphate into D-glycerate 1,3-bisphosphate (e.g., *E. coli gapA* (Branlant et al., *Eur. J. Biochem.* 150:61-66 (1985))), N-acetyl-gamma-glutamyl-phosphate reductase which converts N-acetyl-L-glutamate-5-semialdehyde into N-acetyl-L-glutamyl-5-phosphate (e.g., *E. coli argC* (Parsot et al. *Gene*, 68: 275-283 (1988))), and glutamate-5-semialdehyde dehydrogenase which converts L-glutamate-5-semialdehyde into L-glutamyl-5-phosphate (e.g., *E. coli proA* (Smith et al., *J. Bacteriol.*, 157:545-551 (1984))). Genes encoding glutamate-5-semialdehyde dehydrogenase enzymes from *Salmonella typhimurium* (Mahan et al., *J. Bacteriol.*, 156: 1249-1262 (1983)) and *Campylobacter jejuni* (Louie et al., *Mol. Gen. Genet.*, 240:29-35 (1993)) were cloned and expressed in *E. coli*. These genes/proteins are identified below in Table 103.

20

Table 103.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>gapA</i>	P0A9B2.2	71159358	<i>Escherichia coli</i>
<i>argC</i>	NP_418393.1	16131796	<i>Escherichia coli</i>
<i>proA</i>	NP_414778.1	16128229	<i>Escherichia coli</i>
<i>proA</i>	NP_459319.1	16763704	<i>Salmonella typhimurium</i>
<i>proA</i>	P53000.2	9087222	<i>Campylobacter jejuni</i>

Succinate reductase and 4-Hydroxybutyrate reductase

The direct reduction of succinate to succinic semialdehyde or 4-HB to 4-hydroxybutyraldehyde can be catalyzed by a carboxylic acid reductase. The carboxylic acid reductase of *Nocardia iowensis*, known equivalently as aryl-aldehyde dehydrogenase, catalyzes the magnesium, ATP and NADPH-dependent reduction of carboxylic acids to their corresponding aldehydes (Venkitasubramanian et al., J Biol. Chem. 282:478-485 (2007)) and is capable of catalyzing the conversion of 4-hydroxybutyrate to 4-hydroxybutanal. This enzyme, encoded by *car*, was cloned and functionally expressed in *E. coli* (Venkitasubramanian et al., J Biol. Chem. 282:478-485 (2007)). Expression of the *npt* gene product improved activity of the enzyme via post-transcriptional modification. The *npt* gene encodes a specific phosphopantetheine transferase (PPTase) that converts the inactive apo-enzyme to the active holo-enzyme. The natural substrate of this enzyme is vanillic acid and the enzyme exhibits broad acceptance of aromatic and aliphatic substrates (Venkitasubramanian et al. "Biocatalytic Reduction of Carboxylic Acids: Mechanism and Applications" Chapter 15 in Biocatalysis in the Pharmaceutical and Biotechnology Industries, ed. R.N. Patel, CRC Press LLC, Boca Raton, FL. (2006)). These genes/proteins are identified below in Table 104.

Table 104.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>car</i>	AAR91681.1	40796035	<i>Nocardia iowensis</i> (sp. NRRL 5646)
<i>npt</i>	ABI83656.1	114848891	<i>Nocardia iowensis</i> (sp. NRRL 5646)

Additional *car* and *npt* genes can be identified based on sequence homology. Non-limiting examples of proteins encoded by these genes are shown in Table 105.

Table 105.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>fadD9</i>	YP_978699.1	121638475	<i>Mycobacterium bovis</i> BCG
<i>BCG_2812c</i>	YP_978898.1	121638674	<i>Mycobacterium bovis</i> BCG
<i>nfa20150</i>	YP_118225.1	54023983	<i>Nocardia farcinica</i> IFM 10152
<i>nfa40540</i>	YP_120266.1	54026024	<i>Nocardia farcinica</i> IFM 10152
<i>SGR_6790</i>	YP_001828302.1	182440583	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350
<i>SGR_665</i>	YP_001822177.1	182434458	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350
<i>MSMEG_2956</i>	YP_887275.1	YP_887275.1	<i>Mycobacterium smegmatis</i> MC2 155
<i>MSMEG_5739</i>	YP_889972.1	118469671	<i>Mycobacterium smegmatis</i> MC2 155

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>MSMEG_2648</i>	YP_886985.1	118471293	<i>Mycobacterium smegmatis</i> MC2 155
<i>MAP1040c</i>	NP_959974.1	41407138	<i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> K-10
<i>MAP2899c</i>	NP_961833.1	41408997	<i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> K-10
<i>MMAR_2117</i>	YP_001850422.1	183982131	<i>Mycobacterium marinum</i> M
<i>MMAR_2936</i>	YP_001851230.1	183982939	<i>Mycobacterium marinum</i> M
<i>MMAR_1916</i>	YP_001850220.1	183981929	<i>Mycobacterium marinum</i> M
<i>TpauDRAFT_33060</i>	ZP_04027864.1	227980601	<i>Tsukamurella paurometabola</i> DSM 20162
<i>TpauDRAFT_20920</i>	ZP_04026660.1	227979396	<i>Tsukamurella paurometabola</i> DSM 20162
<i>CPCC7001_1320</i>	ZP_05045132.1	254431429	<i>Cyanobium PCC7001</i>
<i>DDBDRAFT_0187729</i>	XP_636931.1	66806417	<i>Dictyostelium discoideum</i> AX4

An additional enzyme candidate found in *Streptomyces griseus* is encoded by the *griC* and *griD* genes. This enzyme is believed to convert 3-amino-4-hydroxybenzoic acid to 3-amino-4-hydroxybenzaldehyde as deletion of either *griC* or *griD* led to accumulation of extracellular 3-acetylamin-4-hydroxybenzoic acid, a shunt product of 3-amino-4-hydroxybenzoic acid metabolism (Suzuki, et al., *J. Antibiot.* 60(6):380-387 (2007)). Co-expression of *griC* and *griD* with SGR_665, an enzyme similar in sequence to the *Nocardia iowensis npt*, may be beneficial. These genes/proteins are identified below in Table 106.

Table 106.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>griC</i>	YP_001825755.1	182438036	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350
<i>griD</i>	YP_001825756.1	182438037	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350

An enzyme with similar characteristics, alpha-aminoadipate reductase (AAR, EC 1.2.1.31), participates in lysine biosynthesis pathways in some fungal species. This enzyme naturally reduces alpha-aminoadipate to alpha-aminoadipate semialdehyde. The carboxyl group is first activated through the ATP-dependent formation of an adenylate that is then reduced by NAD(P)H to yield the aldehyde and AMP. Like CAR, this enzyme utilizes magnesium and requires activation by a PPTase. Enzyme candidates for AAR and its corresponding PPTase are found in *Saccharomyces cerevisiae* (Morris et al., *Gene* 98:141-145 (1991)), *Candida albicans* (Guo et al., *Mol. Genet. Genomics* 269:271-279 (2003)), and *Schizosaccharomyces pombe* (Ford et al., *Curr. Genet.* 28:131-137 (1995)). The AAR from *S. pombe* exhibited significant activity when expressed in *E. coli* (Guo et al., *Yeast* 21:1279-1288 (2004)). The AAR from *Penicillium chrysogenum* accepts S-carboxymethyl-L-cysteine as an alternate substrate, but did not react

with adipate, L-glutamate or diaminopimelate (Hijarrubia et al., *J Biol. Chem* 278:8250-8256 (2003)). The gene encoding the *P. chrysogenum* PPTase has not been identified to date and no high-confidence hits were identified by sequence comparison homology searching. These genes/proteins are identified below in Table 107.

5

Table 107.

Gene	GenBank ID	GI Number	Organism
<i>LYS2</i>	AAA34747.1	171867	<i>Saccharomyces cerevisiae</i>
<i>LYS5</i>	P50113.1	1708896	<i>Saccharomyces cerevisiae</i>
<i>LYS2</i>	AAC02241.1	2853226	<i>Candida albicans</i>
<i>LYS5</i>	AAO26020.1	28136195	<i>Candida albicans</i>
<i>Lys1p</i>	P40976.3	13124791	<i>Schizosaccharomyces pombe</i>
<i>Lys7p</i>	Q10474.1	1723561	<i>Schizosaccharomyces pombe</i>
<i>Lys2</i>	CAA74300.1	3282044	<i>Penicillium chrysogenum</i>

4-Hydroxybutyryl-CoA dehydratase

4-Hydroxybutyryl-CoA dehydratase catalyzes the reversible conversion of 4-hydroxybutyryl-CoA to crotonyl-CoA. This enzyme possesses an intrinsic vinylacetyl-CoA Δ -isomerase activity, shifting the double bond from the 3,4 position to the 2,3 position (Scherf et al., *Eur. J BioChem.* 215:421-429 (1993); and Scherf et al., *Arch. Microbiol* 161:239-245 (1994)). 4-Hydroxybutyryl-CoA dehydratase enzymes from *C. aminobutyricum* and *C. kluyveri* were purified, characterized, and sequenced at the N-terminus (Scherf et al., *Eur. J BioChem.* 215:421-429 (1993); and Scherf et al., *Arch. Microbiol* 161:239-245 (1994)). The *C. kluyveri* enzyme, encoded by *abfD*, was cloned, sequenced and expressed in *E. coli* (Gerhardt et al., *Arch. Microbiol* 174:189-199 (2000)). The *abfD* gene product from *Porphyromonas gingivalis* ATCC 33277 is closely related by sequence homology to the Clostridial gene products. These genes/proteins are identified below in Table 108.

10

15

Table 108.

Gene	GenBank ID	GI Number	Organism
<i>abfD</i>	YP_001396399.1	153955634	<i>Clostridium kluyveri</i> DSM 555
<i>abfD</i>	P55792	84028213	<i>Clostridium aminobutyricum</i>
<i>abfD</i>	YP_001928843	188994591	<i>Porphyromonas gingivalis</i> ATCC 33277

Crotonase

3-Hydroxybutyryl-CoA dehydratase (EC 4.2.1.55), also called crotonase, is an enoyl-CoA hydratase that reversibly dehydrates 3-hydroxyisobutyryl-CoA to form crotonyl-CoA. Crotonase enzymes are required for n-butanol formation in some organisms, particularly

20

Clostridial species, and also comprise one step of the 3-hydroxypropionate/4-hydroxybutyrate cycle in thermoacidophilic Archaea of the genera *Sulfolobus*, *Acidianus*, and *Metallosphaera*. Exemplary genes encoding crotonase enzymes can be found in *C. acetobutylicum* (Atsumi et al., *Metab Eng* 10:305-311 (2008); and Boynton et al., *J Bacteriol.* 178:3015-3024 (1996)), *C. kluyveri* (Hillmer et al., *FEBS Lett.* 21:351-354 (1972)), and *Metallosphaera sedula* (Berg et al., *Science.* 318:1782-1786 (2007)) though the sequence of the latter gene is not known. The enoyl-CoA hydratase of *Pseudomonas putida*, encoded by *ech*, catalyzes the conversion of crotonyl-CoA to 3-hydroxybutyryl-CoA (Roberts et al., *Arch. Microbiol* 117:99-108 (1978)). Additional enoyl-CoA hydratase candidates are *phaA* and *phaB*, of *P. putida*, and *paaA* and *paaB* from *P. fluorescens* (Olivera et al., *Proc. Natl. Acad. Sci U. S. A* 95:6419-6424 (1998)). Lastly, a number of *Escherichia coli* genes have been shown to demonstrate enoyl-CoA hydratase functionality including *maoC* (Park et al., *J Bacteriol.* 185:5391-5397 (2003)), *paaF* (Ismail et al., *Eur. J BioChem.* 270:3047-3054 (2003); Park et al., *Appl. BioChem. Biotechnol* 113-116:335-346 (2004) and Park et al., *Biotechnol Bioeng* 86:681-686 (2004)) and *paaG* (Ismail et al., *Eur. J BioChem.* 270:3047-3054 (2003); Park et al., *Appl. BioChem. Biotechnol* 113-116:335-346 (2004) and Park et al., *Biotechnol Bioeng* 86:681-686 (2004)). These genes/proteins are identified below in Table 109.

Table 109.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>crt</i>	NP_349318.1	15895969	<i>Clostridium acetobutylicum</i>
<i>crtI</i>	YP_001393856.1	153953091	<i>Clostridium kluyveri</i>
<i>ech</i>	NP_745498.1	26990073	<i>Pseudomonas putida</i>
<i>paaA</i>	NP_745427.1	26990002	<i>Pseudomonas putida</i>
<i>paaB</i>	NP_745426.1	26990001	<i>Pseudomonas putida</i>
<i>phaA</i>	ABF82233.1	106636093	<i>Pseudomonas fluorescens</i>
<i>phaB</i>	ABF82234.1	106636094	<i>Pseudomonas fluorescens</i>
<i>maoC</i>	NP_415905.1	16129348	<i>Escherichia coli</i>
<i>paaF</i>	NP_415911.1	16129354	<i>Escherichia coli</i>
<i>paaG</i>	NP_415912.1	16129355	<i>Escherichia coli</i>

3-Hydroxybutyryl-CoA reductase (aldehyde forming)

3-Hydroxybutyryl-CoA dehydrogenase catalyzes the NAD(P)H dependent reduction of 3-hydroxybutyryl-CoA to 3-hydroxybutyraldehyde. An enzyme catalyzing this transformation has not been identified to date. An exemplary CoA-acylating aldehyde dehydrogenase is the *ald* gene from *Clostridium beijerinckii* (Toth et al., *Appl Environ. Microbiol* 65:4973-4980 (1999)). This enzyme has been reported to reduce acetyl-CoA and butyryl-CoA to their corresponding aldehydes. Another enzyme that converts an acyl-CoA to its corresponding aldehyde is malonyl-CoA reductase which transforms malonyl-CoA to malonic semialdehyde. Malonyl-

CoA reductase is a key enzyme in autotrophic carbon fixation via the 3-hydroxypropionate cycle in thermoacidophilic archaeal bacteria (Berg et al., *Science*. 318:1782-1786 (2007); and Thauer, *Science*. 318:1732-1733 (2007)). The enzyme utilizes NADPH as a cofactor and has been characterized in *Metallosphaera* and *Sulfolobus spp* (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006); and Hugler et al., *J. Bacteriol.* 184:2404-2410 (2002)). The enzyme is encoded by *Msed_0709* in *Metallosphaera sedula* (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006); Berg et al., *Science*. 318:1782-1786 (2007)). A gene encoding a malonyl-CoA reductase from *Sulfolobus tokodaii* was cloned and heterologously expressed in *E. coli* (Alber et al., *J. Bacteriol.* 188:8551-8559 (2006)). This enzyme has also been shown to catalyze the conversion of methylmalonyl-CoA to its corresponding aldehyde (WO/2007/141208). Aldehyde dehydrogenase enzyme candidates for converting 4-hydroxybutyryl-CoA to 4-hydroxybutyraldehyde, described above, are also applicable here. These genes/proteins are identified below in Table 110.

Table 110.

Gene	GenBank ID	GI Number	Organism
<i>Ald</i>	AAT66436	49473535	<i>Clostridium beijerinckii</i>
<i>Msed_0709</i>	YP_001190808.1	146303492	<i>Metallosphaera sedula</i>
<i>mcr</i>	NP_378167.1	15922498	<i>Sulfolobus tokodaii</i>

15 3-Hydroxybutyraldehyde reductase

An enzyme with 3-hydroxybutyraldehyde reductase activity is required to convert 3-hydroxybutyraldehyde to 1,3-butanediol. Exemplary genes encoding enzymes that catalyze the conversion of an aldehyde to alcohol (i.e., alcohol dehydrogenase or equivalently aldehyde reductase) include *alrA* encoding a medium-chain alcohol dehydrogenase for C2-C14 (Tani et al., *Appl. Environ. Microbiol.* 66:5231-5235 (2000)), ADH2 from *Saccharomyces cerevisiae* (Atsumi et al., *Nature* 451:86-89 (2008)), *yqhD* from *E. coli* which has preference for molecules longer than C(3) (Sulzenbacher et al., *Journal of Molecular Biology* 342:489-502 (2004)), and *bdh I* and *bdh II* from *C. acetobutylicum* which converts butyraldehyde into butanol (Walter et al., *Journal of Bacteriology* 174:7149-7158 (1992)). The gene product of *yqhD* catalyzes the reduction of acetaldehyde, malondialdehyde, propionaldehyde, butyraldehyde, and acrolein using NADPH as the cofactor (Perez et al., *J Biol. Chem.* 283:7346-7353 (2008); and Perez et al., *J Biol. Chem.* 283:7346-7353 (2008)). The *adhA* gene product from *Zymomonas mobilis* has been demonstrated to have activity on a number of aldehydes including formaldehyde, acetaldehyde, propionaldehyde, butyraldehyde, and acrolein (Kinoshita et al., *Appl Microbiol Biotechnol* 22:249-254 (1985)). These genes/proteins are identified below in Table 111.

Table 111.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>alrA</i>	BAB12273.1	9967138	<i>Acinetobacter</i> sp. Strain M-1
<i>ADH2</i>	NP_014032.1	6323961	<i>Saccharomyces cerevisiae</i>
<i>yqhD</i>	NP_417484.1	16130909	<i>Escherichia coli</i>
<i>bdh I</i>	NP_349892.1	15896543	<i>Clostridium acetobutylicum</i>
<i>bdh II</i>	NP_349891.1	15896542	<i>Clostridium acetobutylicum</i>
<i>adhA</i>	YP_162971.1	56552132	<i>Zymomonas mobilis</i>

Additional candidates include 4-hydroxybutyrate dehydrogenase and 3-hydroxyisobutyrate dehydrogenase enzymes. 4-Hydroxybutyrate dehydrogenase enzymes naturally convert 4-hydroxybutyraldehyde to 4-HB and have been characterized in *Ralstonia eutropha* (Bravo et al.,
5 *J. Forensic Sci.* 49:379-387 (2004)), *Clostridium kluyveri* (Wolff et al., *Protein Expr. Purif.* 6:206-212 (1995)) and *Arabidopsis thaliana* (Breitkreuz, et al., *J. Biol. Chem.* 278:41552-41556 (2003)). 3-Hydroxyisobutyrate dehydrogenase enzyme candidates include *mmsB* from *Pseudomonas aeruginosa* PAO1 (Gokam et al., *United States patent* 7393676 (2008)), *mmsB* from *Pseudomonas putida* KT2440 (118) and *mmsB* from *Pseudomonas putida* E23
10 (Chowdhury, et al., *Biosci. Biotechnol. BioChem.* 60:2043-2047 (1996)). These genes/proteins are identified below in Table 112.

Table 112.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>4hbd</i>	YP_726053.1	113867564	<i>Ralstonia eutropha</i> H16
<i>4hbd</i>	EDK35022.1	146348486	<i>Clostridium kluyveri</i>
<i>4hbd</i>	Q94B07	75249805	<i>Arabidopsis thaliana</i>
<i>mmsB</i>	NP_252259.1	15598765	<i>Pseudomonas aeruginosa</i> PAO1
<i>mmsB</i>	NP_746775.1	26991350	<i>Pseudomonas putida</i> KT2440
<i>mmsB</i>	JC7926	60729613	<i>Pseudomonas putida</i> E23

3-Hydroxybutyryl-CoA reductase (alcohol forming)

15 A bifunctional oxidoreductase is required for the direct conversion of 3-hydroxybutyryl-CoA to 1,3-butanediol. Exemplary enzymes that convert an acyl-CoA to alcohol include those that transform substrates such as acetyl-CoA to ethanol (e.g., *adhE* from *E. coli* (Kessler et al., *FEBS. Lett.* 281:59-63 (1991))), butyryl-CoA to butanol (e.g. *adhE2* from *C. acetobutylicum* (Fontaine et al., *J. Bacteriol.* 184:821-830 (2002))) and 4-hydroxybutyryl-CoA to 1,4-butanediol
20 (see candidates in previous section). The jojoba (*Simmondsia chinensis*) FAR encodes an alcohol-forming fatty acyl-CoA reductase. This gene was cloned and overexpressed in *E. coli*, resulting in FAR activity and the accumulation of fatty alcohol (Metz et al., *Plant Physiology* 122:635-644 (2000)). Another exemplary enzyme convert malonyl-CoA to 3-

hydroxypropionate. An NADPH-dependent enzyme with this activity has characterized in *Chloroflexus aurantiacus* where it participates in the 3-hydroxypropionate cycle (Hugler et al., *J. Bacteriol.* 184:2404-2410 (2002); and Strauss et al., *Eur. J. BioChem.* 215:633-643 (1993)).

This enzyme, with a mass of 300 kDa, is highly substrate-specific and shows little sequence
5 similarity to other known oxidoreductases (Hugler et al., *J. Bacteriol.* 184:2404-2410 (2002)).

These genes/proteins are identified below in Table 113.

Table 113.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>adhE</i>	NP_415757.1	16129202	<i>Escherichia coli</i>
<i>adhE2</i>	AAK09379.1	12958626	<i>Clostridium acetobutylicum</i>
<i>adhE</i>	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>
<i>FAR</i>	AAD38039.1	5020215	<i>Simmondsia chinensis</i>
<i>mcr</i>	AAS20429.1	42561982	<i>Chloroflexus aurantiacus</i>

3-Hydroxybutyryl-CoA transferase

10 The conversion of 3-hydroxybutyryl-CoA to 3-hydroxybutyrate (3-HB) is catalyzed by a CoA transferase, hydrolase or synthetase. A CoA transferase enzyme catalyzing this specific transformation has not been identified to date. The *E. coli* enzyme acyl-CoA:acetate-CoA transferase, also known as acetate-CoA transferase (EC 2.8.3.8), has been shown to transfer the CoA moiety to acetate from a variety of branched and linear acyl-CoA substrates, including
15 isobutyrate (Matthies et al., *Appl Environ Microbiol* 58:1435-1439 (1992)), valerate (Vanderwinkel et al., *BioChem. Biophys. Res Commun.* 33:902-908 (1968)) and butanoate (Vanderwinkel et al., *BioChem. Biophys. Res Commun.* 33:902-908 (1968)). This enzyme is encoded by *atoA* (alpha subunit) and *atoD* (beta subunit) in *E. coli* sp. K12 (Korolev et al., *D Biol Crystallogr.* 58:2116-2121 (2002); and Vanderwinkel et al., *BioChem. Biophys. Res*
20 *Commun.* 33:902-908 (1968)) and *actA* and *cg0592* in *Corynebacterium glutamicum* ATCC 13032 (Duncan et al., *Appl Environ Microbiol* 68:5186-5190 (2002)). Similar enzymes exist in *Corynebacterium glutamicum* ATCC 13032 (Eikmanns et al., *Mol. Gen. Genet.* 218:330-339 (1989)), *Clostridium acetobutylicum* (Cary et al., *Appl. Environ. Microbiol* 56:1576-1583 (1990); and Wiesenborn et al., *Appl. Environ. Microbiol* 55:323-329 (1989)), and *Clostridium*
25 *saccharoperbutylacetonicum* (Kosaka et al., *Biosci. Biotechnol BioChem.* 71:58-68 (2007)). These genes/proteins are identified below in Table 114.

Table 114.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>atoA</i>	P76459.1	2492994	<i>Escherichia coli</i>
<i>atoD</i>	P76458.1	2492990	<i>Escherichia coli</i>
<i>actA</i>	YP_226809.1	62391407	<i>Corynebacterium glutamicum</i>
<i>cg0592</i>	YP_224801.1	62389399	<i>Corynebacterium glutamicum</i>
<i>ctfA</i>	NP_149326.1	15004866	<i>Clostridium acetobutylicum</i>
<i>ctfB</i>	NP_149327	15004867	<i>Clostridium acetobutylicum</i>
<i>ctfA</i>	AAP42564.1	31075384	<i>Clostridium saccharoperbutylacetonicum</i>
<i>ctfB</i>	AAP42565.1	31075385	<i>Clostridium saccharoperbutylacetonicum</i>

CoA transferase gene candidates described for propionyl-CoA transferase, methylmalonyl-CoA transferase, acetoacetyl-CoA transferase, methacrylyl-CoA transferase, 3-hydroxyisobutyryl-

- 5 CoA transferase, 4-hydroxybutyryl-CoA transferase and succinyl-CoA transferase are also applicable here.

3-Hydroxybutyryl-CoA synthetase

- 3-Hydroxybutyryl-CoA can also be converted to 3-HB by a CoA synthetase (also known as
 10 ligase or synthase). A candidate ATP synthase is ADP-forming acetyl-CoA synthetase (ACD, EC 6.2.1.13), an enzyme that couples the conversion of acyl-CoA esters to their corresponding acids with the concurrent synthesis of ATP. Although this enzyme has not been shown to react with 3-hydroxybutyryl-CoA as a substrate, several enzymes with broad substrate specificities have been described in the literature. ACD I from *Archaeoglobus fulgidus*, encoded by *AF1211*,
 15 was shown to operate on a variety of linear and branched-chain substrates including isobutyrate, isopentanoate, and fumarate (Musfeldt et al., *J Bacteriol.* 184:636-644 (2002)). A second reversible ACD in *Archaeoglobus fulgidus*, encoded by *AF1983*, was also shown to have a broad substrate range with high activity on cyclic compounds phenylacetate and indoleacetate (Musfeldt, et al., *J Bacteriol.* 184:636-644 (2002)). The enzyme from *Haloarcula marismortui*
 20 (annotated as a succinyl-CoA synthetase) accepts propionate, butyrate, and branched-chain acids (isovalerate and isobutyrate) as substrates, and was shown to operate in the forward and reverse directions (Brasen et al., *Arch. Microbiol* 182:277-287 (2004)). The ACD encoded by *PAE3250* from hyperthermophilic crenarchaeon *Pyrobaculum aerophilum* showed the broadest substrate range of all characterized ACDs, reacting with acetyl-CoA, isobutyryl-CoA (preferred substrate)
 25 and phenylacetyl-CoA (Brasen et al., *Arch. Microbiol* 182:277-287 (2004)). However, directed evolution or engineering may be necessary for this enzyme to operate at the physiological temperature of the host organism. The enzymes from *A. fulgidus*, *H. marismortui* and *P.*

aerophilum have all been cloned, functionally expressed, and characterized in *E. coli* (Brasen et al., *Arch. Microbiol* 182:277-287 (2004); and Musfeldt et al., *J Bacteriol.* 184:636-644 (2002)). An additional candidate is the enzyme encoded by *sucCD* in *E. coli*, which naturally catalyzes the formation of succinyl-CoA from succinate with the concomitant consumption of one ATP, a reaction which is reversible *in vivo* (Buck et al., *BioChemistry* 24:6245-6252 (1984)). These genes/proteins are identified below in Table 115.

Table 115.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>AF1211</i>	NP_070039.1	11498810	<i>Archaeoglobus fulgidus</i> DSM 4304
<i>AF1983</i>	NP_070807.1	11499565	<i>Archaeoglobus fulgidus</i> DSM 4304
<i>scs</i>	YP_135572.1	55377722	<i>Haloarcula marismortui</i>
<i>PAE3250</i>	NP_560604.1	18313937	<i>Pyrobaculum aerophilum</i> str. IM2
<i>sucC</i>	NP_415256.1	16128703	<i>Escherichia coli</i>
<i>sucD</i>	AAC73823.1	1786949	<i>Escherichia coli</i>

CoA synthetase gene candidates described for propionyl-CoA synthetase, methylmalonyl-CoA synthetase, methacrylyl-CoA synthetase, acetoacetyl-CoA synthetase, 3-hydroxyisobutyryl-CoA synthetase, 4-hydroxybutyryl-CoA synthetase and succinyl-CoA synthetase are also applicable here.

3-Hydroxybutyryl-CoA hydrolase

A 3-hydroxybutyryl-CoA hydrolase is required to convert 3-hydroxybutyryl-CoA to 3-HB. The enzyme 3-hydroxyisobutyryl-CoA hydrolase (EC 3.1.2.4) catalyzes a related transformation: the hydrolysis of 3-hydroxyisobutyryl-CoA. The 3-hydroxyisobutyryl-CoA hydrolase from *Homo sapiens* also accepts 3-hydroxybutyryl-CoA as a substrate (Shimomura et al., *Methods Enzymol.* 324:229-240 (2000)). This enzyme has also been characterized in *Rattus norvegicus* (Shimomura et al., *J Biol Chem.* 269:14248-14253 (1994); and Shimomura et al., *Methods Enzymol.* 324:229-240 (2000)). Candidate genes by sequence homology include *hibch* of *Saccharomyces cerevisiae* and *BC_2292* of *Bacillus cereus*. These proteins are identified below in Table 116. Additional CoA hydrolase enzyme candidates identified for propionyl-CoA hydrolase, methylmalonyl-CoA hydrolase, methacrylyl-CoA hydrolase, acetoacetyl-CoA hydrolase and 3-hydroxyisobutyryl-CoA are also applicable here. These genes/proteins are identified below in Table 116.

Table 116.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>hibch</i>	Q5XIE6.2	146324906	<i>Rattus norvegicus</i>
<i>hibch</i>	Q6NVY1.2	146324905	<i>Homo sapiens</i>
<i>hibch</i>	P28817.2	2506374	<i>Saccharomyces cerevisiae</i>
<i>BC_2292</i>	AP09256	29895975	<i>Bacillus cereus</i>

3-Hydroxybutyrate reductase

The reduction of 3-hydroxybutyrate to 3-hydroxybutyraldehyde is catalyzed by a carboxylic acid reductase. Exemplary enzyme candidates for succinate reductase and 4-hydroxybutyrate reductase enzymes are also applicable here.

4-Hydroxybutyryl-CoA mutase

The conversion of 4HB-CoA to 3-hydroxyisobutyryl-CoA is catalyzed by a methylmutase. Such a conversion has yet to be demonstrated experimentally. However, two methylmutases (i.e., isobutyryl-CoA mutase and methylmalonyl-CoA mutase) that catalyze similar reactions are promising candidates given the structural similarity of their corresponding substrates.

Methylmalonyl-CoA mutase (MCM) is a cobalamin-dependent enzyme that naturally converts succinyl-CoA to methylmalonyl-CoA. In *E. coli*, the reversible adenosylcobalamin-dependant mutase participates in a three-step pathway leading to the conversion of succinate to propionate (Haller et al., *BioChemistry* 39:4622-9 (2000)). MCM is encoded by genes *scpA* in *Escherichia coli* (Bobik et al., *Anal. Bioanal. Chem.* 375:344-349 (2003); and Haller et al., *BioChemistry* 39:4622-4629 (2000)) and *mutA* in *Homo sapiens* (Padovani et al., *BioChemistry* 45:9300-9306 (2006)). In several other organisms MCM contains alpha and beta subunits and is encoded by two genes. Exemplary gene candidates encoding the two-subunit protein are *Propionibacterium freudenreichii* sp. *shermanii* *mutA* and *mutB* (Korotkova et al., *J Biol Chem.* 279:13652-13658 (2004)) and *Methylobacterium extorquens* *mcmA* and *mcmB* (Korotkova et al., *J Biol Chem.* 279:13652-13658 (2004)). These genes/proteins are identified below in Table 117.

Table 117.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>scpA</i>	NP_417392.1	16130818	<i>Escherichia coli</i> K12
<i>mutA</i>	P22033.3	67469281	<i>Homo sapiens</i>
<i>mutA</i>	P11652.3	127549	<i>Propionibacterium freudenreichii</i> sp. <i>shermanii</i>

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>mutB</i>	P11653.3	127550	<i>Propionibacterium freudenreichii</i> <i>sp. shermanii</i>
<i>mcmA</i>	Q84FZ1	75486201	<i>Methylobacterium extorquens</i>
<i>mcmB</i>	Q6TMA2	75493131	<i>Methylobacterium extorquens</i>

Additional enzyme candidates identified based on high homology to the *E. coli* *spcA* gene product include those identified below in Table 118.

Table 118.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>sbm</i>	NP_838397.1	30064226	<i>Shigella flexneri</i>
<i>SARI_04585</i>	ABX24358.1	160867735	<i>Salmonella enterica</i>
<i>YfreA_01000861</i>	ZP_00830776.1	77975240	<i>Yersinia frederiksenii</i>

- 5 There further exists evidence that genes adjacent to the methylmalonyl-CoA mutase catalytic genes are also required for maximum activity. For example, it has been demonstrated that the *meaB* gene from *M. extorquens* forms a complex with methylmalonyl-CoA mutase, stimulates *in vitro* mutase activity, and possibly protects it from irreversible inactivation (Korotkova et al., *J Biol Chem.* 279:13652-13658 (2004)). The *M. extorquens* *meaB* gene product is highly similar
- 10 to the product of the *E. coli* *argK* gene (BLASTp: 45% identity, e-value: 4e-67) which is adjacent to *scpA* on the chromosome. No sequence for a *meaB* homolog in *P. freudenreichii* is catalogued in GenBank. However, the *Propionibacterium acnes* KPA171202 gene product at the locus *PPA0597* is 51% identical to the *M. extorquens* *meaB* protein and its gene is also adjacent to the methylmalonyl-CoA mutase gene on the chromosome. These genes/proteins are identified
- 15 below in Table 119.

Table 119.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>argK</i>	AAC75955.1	1789285	<i>Escherichia coli</i> K12
<i>PPA0597</i>	YP_055310.1	50842083	<i>Propionibacterium acnes</i>
KPA171202	2QM8_B	158430328	<i>Methylobacterium extorquens</i>

- Alternatively, isobutyryl-CoA mutase (ICM) could catalyze the proposed transformation. ICM is a cobalamin-dependent methylmutase in the MCM family that reversibly rearranges the carbon backbone of butyryl-CoA into isobutyryl-CoA (Figure 7B of Ratnatilleke, *J Biol Chem.* 274:31679-31685 (1999)). A recent study of a novel ICM in *Methylibium petroleiphilum*, along with previous work, provides evidence that changing a single amino acid near the active site alters the substrate specificity of the enzyme (Ratnatilleke et al., *J Biol Chem.* 274:31679-31685 (1999); and Rohwerder et al., *Appl Environ Microbiol* 72:4128-4135 (2006)). This implies that
- 20

if a native enzyme is unable to catalyze the conversion of 4HB-CoA to 3HIB-CoA, the enzyme could undergo rational engineering. Exemplary ICM genes encoding homodimeric enzymes include *icmA* in *Streptomyces coelicolor* A3 (2) and *Mpe_B0541* in *Methylibium petroleiphilum* PM1 (Ratnatilleke et al., *J Biol Chem.* 274:31679-31685 (1999); and Rohwerder et al., *Appl Environ Microbiol* 72:4128-4135 (2006)). Genes encoding heterodimeric enzymes include *icm* and *icmB* in *Streptomyces cinnamomensis* (Ratnatilleke, et al., *J Biol Chem.* 274:31679-31685 (1999); Vrijbloed et al., *J Bacteriol.* 181:5600-5605 (1999) and Zerbe-Burkhardt et al., *J Biol Chem.* 273:6508-6517 (1998)). Enzymes encoded by *icmA* and *icmB* genes in *Streptomyces avermitilis* MA-4680 show high sequence similarity to known ICMs. These genes/proteins are identified below in Table 120.

Table 120.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>icmA</i>	CAB40912.1	4585853	<i>Streptomyces coelicolor</i> A3(2)
<i>Mpe_B0541</i>	YP_001023546.1	124263076	<i>Methylibium petroleiphilum</i> PM1
<i>icm</i>	AAC08713.1	3002492	<i>Streptomyces cinnamomensis</i>
<i>icmB</i>	CAB59633.1	6137077	<i>Streptomyces cinnamomensis</i>
<i>icmA</i>	NP_824008.1	29829374	<i>Streptomyces avermitilis</i>
<i>icmB</i>	NP_824637.1	29830003	<i>Streptomyces avermitilis</i>

3-Hydroxyisobutyryl-CoA transferase

The next step in this pathway entails the conversion of 3-hydroxyisobutyryl-CoA into 3-hydroxyisobutyrate (3-HIB) by a CoA transferase. An enzyme catalyzing this specific transformation has not been identified to date. The *E. coli* enzyme acyl-CoA:acetate-CoA transferase, also known as acetate-CoA transferase (EC 2.8.3.8), has been shown to transfer the CoA moiety to acetate from a variety of branched and linear acyl-CoA substrates, including isobutyrate (Matthies et al., *Appl Environ Microbiol* 58:1435-1439 (1992)), valerate (Vanderwinkel et al., *BioChem. Biophys. Res Commun.* 33:902-908 (1968)) and butanoate (Vanderwinkel et al., *BioChem. Biophys. Res Commun.* 33:902-908 (1968)). This enzyme is encoded by *atoA* (alpha subunit) and *atoD* (beta subunit) in *E. coli* sp. K12 (Korolev et al., *D Biol Crystallogr.* 58:2116-2121 (2002); and Vanderwinkel et al., *BioChem. Biophys. Res Commun.* 33:902-908 (1968)) and *actA* and *cg0592* in *Corynebacterium glutamicum* ATCC 13032 (Duncan et al., *Appl Environ Microbiol* 68:5186-5190 (2002)). Similar enzymes exist in *Corynebacterium glutamicum* ATCC 13032 (Eikmanns et al., *Mol. Gen. Genet.* 218:330-339 (1989)), *Clostridium acetobutylicum* (Cary et al., *Appl. Environ. Microbiol* 56:1576-1583 (1990); and Wiesenborn et al., *Appl. Environ. Microbiol* 55:323-329 (1989)), and *Clostridium saccharoperbutylacetonicum* (Kosaka et al., *Biosci. Biotechnol BioChem.* 71:58-68 (2007)). 4-

Hydroxybutyryl-CoA transferase enzyme candidates, described previously, are also applicable here. These genes/proteins are identified below in Table 121.

Table 121.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>atoA</i>	P76459.1	2492994	<i>Escherichia coli</i>
<i>atoD</i>	P76458.1	2492990	<i>Escherichia coli</i>
<i>actA</i>	YP_226809.1	62391407	<i>Corynebacterium glutamicum</i>
<i>cg0592</i>	YP_224801.1	62389399	<i>Corynebacterium glutamicum</i>
<i>ctfA</i>	NP_149326.1	15004866	<i>Clostridium acetobutylicum</i>
<i>ctfB</i>	NP_149327	15004867	<i>Clostridium acetobutylicum</i>
<i>ctfA</i>	AAP42564.1	31075384	<i>Clostridium saccharoperbutylacetonicum</i>
<i>ctfB</i>	AAP42565.1	31075385	<i>Clostridium saccharoperbutylacetonicum</i>

5 3-Hydroxyisobutyryl-CoA synthetase

3-Hydroxyisobutyryl-CoA can also be converted to 3-HIB by a CoA synthetase (also known as ligase or synthase). A candidate ATP synthase is ADP-forming acetyl-CoA synthetase (ACD, EC 6.2.1.13), an enzyme that couples the conversion of acyl-CoA esters to their corresponding acids with the concurrent synthesis of ATP. Although this enzyme has not been shown to react with 3-hydroxyisobutyryl-CoA as a substrate, several enzymes with broad substrate specificities have been described in the literature. ACD I from *Archaeoglobus fulgidus*, encoded by *AF1211*, was shown to operate on a variety of linear and branched-chain substrates including isobutyrate, isopentanoate, and fumarate (Musfeldt et al., *J Bacteriol.* 184:636-644 (2002)). A second reversible ACD in *Archaeoglobus fulgidus*, encoded by *AF1983*, was also shown to have a broad substrate range with high activity on cyclic compounds phenylacetate and indoleacetate (Musfeldt, et al., *J Bacteriol.* 184:636-644 (2002)). The enzyme from *Haloarcula marismortui* (annotated as a succinyl-CoA synthetase) accepts propionate, butyrate, and branched-chain acids (isovalerate and isobutyrate) as substrates, and was shown to operate in the forward and reverse directions (Brasen et al., *Arch. Microbiol* 182:277-287 (2004)). The ACD encoded by *PAE3250* from hyperthermophilic crenarchaeon *Pyrobaculum aerophilum* showed the broadest substrate range of all characterized ACDs, reacting with acetyl-CoA, isobutyryl-CoA (preferred substrate) and phenylacetyl-CoA (Brasen et al., *Arch. Microbiol* 182:277-287 (2004)). However, directed evolution or engineering may be necessary for this enzyme to operate at the physiological temperature of the host organism. The enzymes from *A. fulgidus*, *H. marismortui* and *P. aerophilum* have all been cloned, functionally expressed, and characterized in *E. coli* (Brasen et al., *Arch. Microbiol* 182:277-287 (2004); and Musfeldt et al., *J Bacteriol.* 184:636-644 (2002)).

An additional candidate is the enzyme encoded by *sucCD* in *E. coli*, which naturally catalyzes the formation of succinyl-CoA from succinate with the concomitant consumption of one ATP, a reaction which is reversible *in vivo* (Buck et al., *BioChemistry* 24:6245-6252 (1984)). These genes/proteins are identified below in Table 122.

5

Table 122.

Gene	GenBank ID	GI Number	Organism
<i>AF1211</i>	NP_070039.1	11498810	<i>Archaeoglobus fulgidus</i> DSM 4304
<i>AF1983</i>	NP_070807.1	11499565	<i>Archaeoglobus fulgidus</i> DSM 4304
<i>scs</i>	YP_135572.1	55377722	<i>Haloarcula marismortui</i>
<i>PAE3250</i>	NP_560604.1	18313937	<i>Pyrobaculum aerophilum</i> str. IM2
<i>sucC</i>	NP_415256.1	16128703	<i>Escherichia coli</i>
<i>sucD</i>	AAC73823.1	1786949	<i>Escherichia coli</i>

3-Hydroxyisobutyryl-CoA hydrolase

The enzyme 3-hydroxyisobutyryl-CoA hydrolase selectively converts 3-hydroxyisobutyryl-CoA to 3-HIB during valine degradation (Shimomura et al., *J Biol Chem* 269:14248-53 (1994)).

- 10 Genes encoding this enzyme were described previously. 3-Hydroxybutyryl-CoA hydrolase and propionyl-CoA gene candidates, described previously, are also applicable here.

3-Hydroxyisobutyrate dehydratase

- The dehydration of 3-hydroxyisobutyrate to methylacrylic acid is catalyzed by an enzyme with 3-hydroxyisobutyrate dehydratase activity. No direct evidence for this specific enzymatic
- 15 transformation has been identified. However, most dehydratases catalyze the alpha, beta-elimination of water which involves activation of the alpha-hydrogen by an electron-withdrawing carbonyl, carboxylate, or CoA-thiol ester group and removal of the hydroxyl group from the beta-position (Buckel et al., *J Bacteriol.* 117:1248-1260 (1974); and Martins, et al., *Proc Natl Acad Sci U S A* 101:15645-9 (2004)). This is the exact type of transformation
- 20 proposed for the final step in the methylacrylic acid pathway. The proposed transformation is highly similar to the 2-(hydroxymethyl)glutarate dehydratase of *Eubacterium barkeri* (Figure 3A). This enzyme has been studied in the context of nicotinate catabolism and is encoded by *hmd* (Alhapel et al., *Proc Natl Acad Sci U S A* 103:12341-6 (2006)). An enzyme with similar functionality in *E. barkeri* is dimethylmaleate hydratase, a reversible Fe^{2+} -dependent and
- 25 oxygen-sensitive enzyme in the aconitase family that hydrates dimethylmaleate to form (2R,3S)-2,3-dimethylmalate. This enzyme is encoded by *dmdAB* (Alhapel et al., *Proc Natl Acad Sci U S A* 103:12341-6 (2006); and Kollmann-Koch et al., *Hoppe Seylers. Z. Physiol Chem.* 365:847-857 (1984)). These genes/proteins are identified below in Table 123.

Table 123.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>hmd</i>	ABC88407.1	86278275	<i>Eubacterium barkeri</i>
<i>dmdA</i>	ABC88408	86278276	<i>Eubacterium barkeri</i>
<i>dmdB</i>	ABC88409.1	86278277	<i>Eubacterium barkeri</i>

An additional enzyme candidate is 2-methylmalate dehydratase, also called citramalate hydrolyase, a reversible hydrolyase that catalyzes the alpha, beta elimination of water from citramalate to form mesaconate. This enzyme has been studied in *Methanocaldococcus jannaschii* in the context of the pyruvate pathway to 2-oxobutanoate, where it has been shown to have a broad substrate specificity (Drevland et al., *J Bacteriol.* 189:4391-4400 (2007)). This enzyme activity was also detected in *Clostridium tetanomorphum*, *Morganella morganii*, *Citrobacter amalonaticus* where it is thought to participate in glutamate degradation (Kato et al., *Arch. Microbiol* 168:457-463 (1997)). The *M. jannaschii* protein sequence does not bear significant homology to genes in these organisms. This genes/proteins is identified below in Table 124.

Table 124.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>leuD</i>	Q58673.1	3122345	<i>Methanocaldococcus jannaschii</i>

Fumarate hydratase enzymes, which naturally catalyze the dehydration of malate to fumarate, represent an additional set of candidates. Although the ability of fumarate hydratase to react on 3-hydroxyisobutyrate as a substrate has not been described, a wealth of structural information is available for this enzyme and other researchers have successfully engineered the enzyme to alter activity, inhibition and localization (Weaver, *D Biol Crystallogr.* 61:1395-1401 (2005)). *E. coli* has three fumarases: FumA, FumB, and FumC that are regulated by growth conditions. FumB is oxygen sensitive and only active under anaerobic conditions. FumA is active under microanaerobic conditions, and FumC is the only active enzyme in aerobic growth (Guest et al., *J Gen Microbiol* 131:2971-2984 (1985); Tseng et al., *J Bacteriol* 183:461-467 (2001) and Woods et al., *Biochim Biophys Acta* 954:14-26 (1988)). Additional enzyme candidates are found in *Campylobacter jejuni* (Smith et al., *Int. J BioChem. Cell Biol* 31:961-975 (1999)), *Thermus thermophilus* (Mizobata et al., *Arch. BioChem. Biophys.* 355:49-55 (1998)) and *Rattus norvegicus* (Kobayashi et al., *J BioChem.* 89:1923-1931 (1981)). The *MmcBC* fumarase from *Pelotomaculum thermopropionicum* is another class of fumarase with two subunits (Shimoyama

et al., *FEMS Microbiol Lett* 270:207-213 (2007)). These genes/proteins are identified below in Table 125.

Table 125.

Gene	GenBank ID	GI Number	Organism
<i>fumA</i>	P0AC33	81175318	<i>Escherichia coli K12</i>
<i>fumB</i>	P14407	33112655	<i>Escherichia coli K12</i>
<i>fumC</i>	P05042.1	120601	<i>Escherichia coli K12</i>
<i>fumC</i>	O69294.1	9789756	<i>Campylobacter jejuni</i>
<i>fumC</i>	P84127	75427690	<i>Thermus thermophilus</i>
<i>fumH</i>	P14408.1	120605	<i>Rattus norvegicus</i>
<i>MmcB</i>	YP_001211906	147677691	<i>Pelotomaculum thermopropionicum</i>
<i>MmcC</i>	YP_001211907	147677692	<i>Pelotomaculum thermopropionicum</i>

5 3-Hydroxyisobutyryl-CoA dehydratase

Dehydration of 3-hydroxyisobutyryl-CoA by a CoA dehydratase yields methacrylyl-CoA. Enoyl-CoA hydratases (EC 4.2.1.17) catalyze the dehydration of 3-hydroxyacyl-CoA substrates (Agnihotri and Liu., *J. Bacteriol.* 188:8551-8559(2003); Conrad et al., *J. Bacteriol.* 118:103-111 (1974); and Roberts et al., *Arch. Microbiol* 117:99-108 (1978)). The enoyl-CoA hydratase (ECH) found in bovine liver accepts a variety of substrates including methacrylyl-CoA, 2- and 3-methyl-crotonoyl-CoA, acryloyl-CoA and 1-carboxycyclohexenoyl-CoA (Agnihotri et al., *Bioorg Med Chem.*, 11(1):9-20 (2003)). A recombinant bovine liver ECH enzyme has been overexpressed in *E. coli* and found to have similar catalytic properties (Dakoji et al., *J Am Chem Soc.*, 123:9749 (2001)). The enoyl-CoA hydratase of *Pseudomonas putida*, encoded by *ech*, catalyzes the conversion of 3-hydroxybutyryl-CoA to crotonoyl-CoA (Roberts et al., *Arch. Microbiol* 117:99-108 (1978)). Additional enoyl-CoA hydratase candidates are *phaA* and *phaB*, of *P. putida*, and *paaA* and *paaB* from *P. fluorescens* (Olivera et al., *Proc. Natl. Acad. Sci U. S. A* 95:6419-6424 (1998)). The gene product of *pimF* in *Rhodopseudomonas palustris* is predicted to encode an enoyl-CoA hydratase that participates in pimeloyl-CoA degradation (Harrison and Harwood, *Microbiology* 151:727-736 (2005)). Lastly, a number of *Escherichia coli* genes have been shown to demonstrate enoyl-CoA hydratase functionality including *maoC* (Park and Lee, *J. Bacteriol.* 185:5391-5397 (2003)), *paaF* (Ismail et al., *J Biochem.* 270:3047-3054 (2003); Park and Lee, *Appl. Biochem. Biotechnol* 113-116:335-346 (2004); and Park and Yup, *Biotechnol Bioeng* 86:681-686 (2004)) and *paaG* (Ismail et al., *J Biochem.* 270:3047-3054(2003); Park and Lee, *Appl. Biochem. Biotechnol* 113-116:335-346 (2004); and Park and

Yup, *Biotechnol Bioeng* 86:681-686 (2004)). These genes/proteins are identified below in Table 126.

Table 126.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>ECHS1</i>	NP_001020377.2	70778822	<i>Bos taurus</i>
<i>ech</i>	NP_745498.1	26990073	<i>Pseudomonas putida</i>
<i>paaA</i>	NP_745427.1	26990002	<i>Pseudomonas putida</i>
<i>paaB</i>	NP_745426.1	26990001	<i>Pseudomonas putida</i>
<i>phaA</i>	ABF82233.1	106636093	<i>Pseudomonas fluorescens</i>
<i>phaB</i>	ABF82234.1	106636094	<i>Pseudomonas fluorescens</i>
<i>pimF</i>	CAE29158	39650635	<i>Rhodopseudomonas palustris</i>
<i>maoC</i>	NP_415905.1	16129348	<i>Escherichia coli</i>
<i>paaF</i>	NP_415911.1	16129354	<i>Escherichia coli</i>
<i>paaG</i>	NP_415912.1	16129355	<i>Escherichia coli</i>

Another exemplary enzyme candidate for catalyzing this reaction is crotonase. Gene candidates
 5 for this enzyme are described above. Alternatively, the *E. coli* gene products of *fadA* and *fadB* encode a multienzyme complex involved in fatty acid oxidation that exhibits enoyl-CoA hydratase activity (Nakahigashi and Inokuchi, *Nucleic Acids Res.* 18:4937 (1990); Yang, *J. Bacteriol.* 173:7405-7406 (1991); and Yang et al., *Biochemistry* 30:6788-6795 (1991)). Knocking out a negative regulator encoded by *fadR* can be utilized to activate the *fadB* gene
 10 product (Sato et al., *J Biosci. Bioeng* 103:38-44 (2007)). The *fadI* and *fadJ* genes encode similar functions and are naturally expressed under anaerobic conditions (Campbell et al., *Mol. Microbiol* 47:793-805 (2003)). These genes/proteins are identified below in Table 127.

Table 127.

<u>Gene</u>	<u>GenBank ID</u>	<u>GI Number</u>	<u>Organism</u>
<i>fadA</i>	YP_026272.1	49176430	<i>Escherichia coli</i>
<i>fadB</i>	NP_418288.1	16131692	<i>Escherichia coli</i>
<i>fadI</i>	NP_416844.1	16130275	<i>Escherichia coli</i>
<i>fadJ</i>	NP_416843.1	16130274	<i>Escherichia coli</i>
<i>fadR</i>	NP_415705.1	16129150	<i>Escherichia coli</i>

15 Methacrylyl-CoA hydrolase

Conversion of methacrylyl-CoA to MAA is catalyzed by a CoA transferase, synthetase or hydrolase. CoA hydrolase gene candidates described for propionyl-CoA hydrolase, methylmalonyl-CoA hydrolase, acetoacetyl-CoA hydrolase, 3-hydroxybutyryl-CoA hydrolase and 3-hydroxyisobutyryl-CoA hydrolase are also applicable here.

Methacrylyl-CoA transferase

Conversion of methacrylyl-CoA to MAA is catalyzed by a CoA transferase, synthetase or hydrolase. CoA transferase gene candidates described for propionyl-CoA transferase, methylmalonyl-CoA transferase, acetoacetyl-CoA transferase, 3-hydroxybutyryl-CoA transferase, 3-hydroxyisobutyryl-CoA transferase, 4-hydroxybutyryl-CoA transferase and succinyl-CoA transferase are applicable here.

Methacrylyl-CoA synthetase

Conversion of methacrylyl-CoA to MAA is catalyzed by a CoA transferase, synthetase or hydrolase. CoA synthetase gene candidates described for propionyl-CoA synthetase, methylmalonyl-CoA synthetase, acetoacetyl-CoA synthetase, 3-hydroxybutyryl-CoA synthetase, 3-hydroxyisobutyryl-CoA synthetase, 4-hydroxybutyryl-CoA synthetase and succinyl-CoA synthetase are applicable here.

Methylmalonyl-CoA hydrolase

Methylmalonyl-CoA is converted to methylmalonate by methylmalonyl-CoA hydrolase (EC 3.1.2.17). This enzyme, isolated from *Rattus norvegicus* liver, is also active on malonyl-CoA and propionyl-CoA as alternative substrates (Kovachy et al., *J. Biol. Chem.*, 258: 11415-11421 (1983)). The gene associated with this enzyme is not known. Other CoA hydrolase enzyme candidates for propionyl-CoA hydrolase, methacrylyl-CoA hydrolase, acetoacetyl-CoA hydrolase, 3-hydroxybutyryl-CoA hydrolase and 3-hydroxyisobutyryl-CoA hydrolase, described in previous sections, are applicable here.

Methylmalonyl-CoA transferase

Alternately, methylmalonyl-CoA is converted to methylmalonate by a CoA transferase. CoA transferase gene candidates described for propionyl-CoA transferase, methacrylyl-CoA transferase, acetoacetyl-CoA transferase, 3-hydroxybutyryl-CoA transferase, 3-hydroxyisobutyryl-CoA transferase, 4-hydroxybutyryl-CoA transferase and succinyl-CoA transferase are also applicable here.

Methylmalonyl-CoA synthetase

Yet another enzyme that forms methylmalonate from methylmalonyl-CoA is methylmalonyl-CoA synthetase. CoA synthetase gene candidates described for propionyl-CoA synthetase, methacrylyl-CoA synthetase, acetoacetyl-CoA synthetase, 3-hydroxybutyryl-CoA synthetase, 3-

hydroxyisobutyryl-CoA synthetase, 4-hydroxybutyryl-CoA synthetase and succinyl-CoA synthetase are applicable here.

Methylmalonate reductase

5 The reduction of methylmalonate to methylmalonate semialdehyde is catalyzed by a carboxylic acid reductase. Exemplary enzyme candidates for succinate reductase and 4-hydroxybutyrate reductase enzymes are also applicable here.

10 Throughout this application various publications have been referenced. The disclosures of these publications in their entireties, including GenBank and GI number publications, are hereby incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains. Although the invention has been described with reference to the examples provided above, it should be understood that various modifications can be made without departing from the spirit of the invention.

What is claimed is:

1. A non-naturally occurring microbial organism, comprising a microbial organism having an n-propanol pathway and an isopropanol pathway, said n-propanol pathway comprising at least one exogenous nucleic acid encoding an n-propanol pathway enzyme expressed in a sufficient amount to produce n-propanol, said n-propanol pathway comprising:
5 a propanol dehydrogenase, a propionaldehyde dehydrogenase, a propionyl-CoA:phosphate propanoyltransferase, a propionyl-CoA hydrolase, a propionyl-CoA transferase, a propionyl-CoA synthetase, a propionate kinase, a propionate reductase or a propionyl phosphate reductase,
10 said isopropanol pathway comprising at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, said isopropanol pathway comprising:
an isopropanol dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, or an acetoacetate decarboxylase.
15
2. The non-naturally occurring microbial organism of claim 1, further comprising an acetyl-CoA pathway comprising at least one exogenous nucleic acid encoding an acetyl-CoA pathway enzyme expressed in a sufficient amount to produce acetyl-CoA, said
20 acetyl-CoA pathway comprising:
a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, or a formate dehydrogenase.
3. The non-naturally occurring microbial organism of claim 1, further comprising a
25 propionyl-CoA pathway comprising at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, said propionyl-CoA pathway comprising:
a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase, a succinyl-CoA synthetase, a
30 methylmalonyl-CoA mutase, a methylmalonyl-CoA epimerase or a methylmalonyl-CoA decarboxylase.

4. The non-naturally occurring microbial organism of claim 3, wherein said propionyl-CoA pathway further comprises a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.
5. The non-naturally occurring microbial organism of claim 1, further comprising a propionyl-CoA pathway comprising at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, said propionyl-CoA pathway comprising:
- a PEP carboxykinase, a PEP carboxylase, a threonine deaminase, or a 2-oxobutanoate dehydrogenase.
6. The non-naturally occurring microbial organism of claim 5, wherein said n-propanol pathway further comprises 2-oxobutanoate decarboxylase.
7. The non-naturally occurring microbial organism of claim 1, further comprising a propionyl-CoA pathway comprising at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, said propionyl-CoA pathway comprising:
- an acetyl-CoA carboxylase, a malonyl-CoA reductase, a malonate semialdehyde reductase or propionyl-CoA synthase.
8. The non-naturally occurring microbial organism of claim 1, further comprising a propionyl-CoA pathway comprising at least one exogenous nucleic acid encoding a propionyl-CoA pathway enzyme expressed in a sufficient amount to produce propionyl-CoA, said propionyl-CoA pathway comprising:
- a lactate dehydrogenase, a lactate-CoA transferase, a lactyl-CoA dehydratase or acryloyl CoA reductase.
9. A non-naturally occurring microbial organism, comprising a microbial organism having an n-propanol pathway and an isopropanol pathway, said n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, said first set of exogenous nucleic acids encoding:
- a propionaldehyde dehydrogenase and a propanol dehydrogenase; or

a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or

a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or

a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

10. The non-naturally occurring microbial organism of claim 9, further comprising an acetyl-CoA pathway comprising a third set of exogenous nucleic acids encoding acetyl-CoA pathway enzymes expressed in a sufficient amount to produce acetyl-CoA, said third set of exogenous nucleic acids encoding:

a pyruvate kinase; and a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase.

11. The non-naturally occurring microbial organism of claim 9, further comprising a propionyl-CoA pathway comprising a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA, said third set of exogenous nucleic acids encoding:

a PEP carboxykinase or a PEP carboxylase; a malate dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA synthetase; a methylmalonyl-CoA mutase; and a methylmalonyl-CoA decarboxylase.

12. The non-naturally occurring microbial organism of claim 11, wherein said third set of exogenous nucleic acids further encodes a methylmalonyl-CoA epimerase, a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

13. The non-naturally occurring microbial organism of claim 9, further comprising a propionyl-CoA pathway comprising a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA, said third set of exogenous nucleic acids encoding:

5 a PEP carboxykinase or a PEP carboxylase; a threonine deaminase; and a 2-oxobutanoate dehydrogenase.

14. The non-naturally occurring microbial organism of claim 13, wherein said third set of exogenous nucleic acids further encodes a or a methylmalonyl-CoA carboxytransferase or a pyruvate carboxylase.

10 15. The non-naturally occurring microbial organism of claim 13, wherein said second set of exogenous nucleic acids further encode a 2-oxobutanoate decarboxylase.

16. The non-naturally occurring microbial organism of claim 9, further comprising a propionyl-CoA pathway comprising a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA, said third set of exogenous nucleic acids encoding:

15 an acetyl-CoA carboxylase; a malonyl-CoA reductase; a malonate semialdehyde reductase; and propionyl-CoA synthase.

17. The non-naturally occurring microbial organism of claim 9, further comprising a propionyl-CoA pathway comprising a third set of exogenous nucleic acids encoding propionyl-CoA pathway enzymes expressed in a sufficient amount to produce propionyl-CoA, said third set of exogenous nucleic acids encoding:

20 a lactate dehydrogenase; a lactate-CoA transferase; a lactyl-CoA dehydratase; and acryloyl CoA reductase.

18. A non-naturally occurring microbial organism, comprising a microbial organism having an n-propanol pathway and an isopropanol pathway, said n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, said first set of exogenous nucleic acids encoding:

25 a PEP carboxykinase or a PEP carboxylase; a malate dehydrogenase; a fumarase; a fumarate reductase; a succinyl-CoA transferase or a succinyl-CoA synthetase; a

30

methyalmalonyl-CoA mutase; a methyalmalonyl-CoA decarboxylase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase and a propionyl phosphate reductase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

19. The non-naturally occurring microbial organism of claim 18, wherein said first set of exogenous nucleic acids further encodes a methyalmalonyl-CoA epimerase, a pyruvate carboxylase or a methyalmalonyl-CoA carboxytransferase.

20. A non-naturally occurring microbial organism, comprising a microbial organism having an n-propanol pathway and an isopropanol pathway, said n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, said first set of exogenous nucleic acids encoding:

a PEP carboxykinase or a PEP carboxylase; a threonine deaminase; and a 2-oxobutanoate decarboxylase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a 2-oxobutanoate dehydrogenase, a propionyl-CoA

hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol,

5 said second set of exogenous nucleic acids encoding:

a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

10

21. The non-naturally occurring microbial organism of claim 20, wherein said second set of exogenous nucleic acids further encodes a methylmalonyl-CoA decarboxylase or a pyruvate carboxylase.

22. A non-naturally occurring microbial organism, comprising a microbial organism having an n-propanol pathway and an isopropanol pathway, said n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, said first set of exogenous nucleic acids encoding:

15

a pyruvate kinase; a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA carboxylase; a malonyl-CoA reductase; a malonate semialdehyde reductase; propionyl-CoA synthase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase,

20

25

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

30

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

23. A non-naturally occurring microbial organism, comprising a microbial organism having an n-propanol pathway and an isopropanol pathway, said n-propanol pathway comprising a first set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, said first set of exogenous nucleic acids encoding:
- a lactate dehydrogenase; a lactate-CoA transferase; a lactyl-CoA dehydratase; acryloyl CoA reductase; and a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase,
- said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:
- a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase; an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

24. The non-naturally occurring microbial organism of any one of claims 1-23, wherein said exogenous nucleic acid is a heterologous nucleic acid.

25. The non-naturally occurring microbial organism of any one of claims 1-23, wherein said non-naturally occurring microbial organism is in a substantially anaerobic culture medium.

26. A method for producing n-propanol and isopropanol, comprising culturing a non-naturally occurring microbial organism of any one of claims 1-23 under conditions and for a sufficient period of time to produce n-propanol and isopropanol.

27. The method of claim 26, wherein said conditions comprise substantially anaerobic culture conditions.

28. The method of claim 26, wherein said exogenous nucleic acid is a heterologous nucleic acid.

5 29. A non-naturally occurring microbial organism, comprising a microbial organism having an n-propanol pathway, said n-propanol pathway comprising at least one exogenous nucleic acid encoding an n-propanol pathway enzyme expressed in a sufficient amount to produce n-propanol, said n-propanol pathway comprising:

10 a propanol dehydrogenase, a propionaldehyde dehydrogenase, a propionyl-CoA:phosphate propanoyltransferase a propionyl-CoA hydrolase, a propionyl-CoA transferase, a propionyl-CoA synthetase, a propionate kinase, a propionate reductase, or a propionyl phosphate reductase.

15 30. A non-naturally occurring microbial organism, comprising a microbial organism having an n-propanol pathway, said n-propanol pathway comprising a set of exogenous nucleic acids encoding n-propanol pathway enzymes expressed in a sufficient amount to produce n-propanol, said set of exogenous nucleic acids encoding:

20 a propionaldehyde dehydrogenase and a propanol dehydrogenase; or a propionyl-CoA:phosphate propanoyltransferase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate kinase, a propionyl phosphate reductase and a propanol dehydrogenase; or a propionyl-CoA hydrolase or a propionyl-CoA transferase or a propionyl-CoA synthetase, a propionate reductase and a propanol dehydrogenase.

31. The non-naturally occurring microbial organism of claim 29 or 30, wherein said exogenous nucleic acid is a heterologous nucleic acid.

25 32. The non-naturally occurring microbial organism of claim 29 or 30, wherein said non-naturally occurring microbial organism is in a substantially anaerobic culture medium.

30 33. A method for producing n-propanol, comprising culturing a non-naturally occurring microbial organism of claim 29 or 30 under conditions and for a sufficient period of time to produce n-propanol.

34. The method of claim 33, wherein said conditions comprise substantially anaerobic culture conditions.

35. The method of claim 33, wherein said exogenous nucleic acid is a heterologous nucleic acid.

5 36. A non-naturally occurring microbial organism, comprising a microbial organism having a 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising at least one exogenous nucleic acid encoding a 1,4-butanediol pathway enzyme expressed in a sufficient amount to produce 1,4-butanediol, said 1,4-butanediol pathway comprising:

10 a 4-hydroxybutyraldehyde reductase, a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde-forming), 4-hydroxybutyryl-phosphate reductase, a 4-hydroxybutyrate reductase; a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, or a 4-hydroxybutyryl-CoA reductase (alcohol-forming);

15 said isopropanol pathway comprising at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, said isopropanol pathway comprising:

20 an isopropanol dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, or an acetoacetate decarboxylase.

37. A non-naturally occurring microbial organism, comprising a microbial organism having a 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising at least one exogenous nucleic acid encoding a 1,3-butanediol pathway enzyme expressed in a sufficient amount to produce 1,3-butanediol, said 1,3-butanediol pathway comprising:

25 a 3-hydroxybutyraldehyde reductase, a succinyl-CoA reductase, a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA dehydratase, a crotonase, a 3-hydroxybutyryl-CoA transferase, a 3-hydroxybutyryl-CoA synthetase, a 3-

hydroxybutyryl-CoA hydrolase, a 3-hydroxybutyrate reductase, a 3-hydroxybutyryl-CoA reductase (aldehyde forming), or a 3-hydroxybutyryl-CoA reductase (alcohol-forming);

said isopropanol pathway comprising at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol,

5 said isopropanol pathway comprising:

an isopropanol dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, or an acetoacetate decarboxylase.

38. A non-naturally occurring microbial organism, comprising a microbial organism
10 having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising at least one exogenous nucleic acid encoding a methylacrylic acid pathway enzyme expressed in a sufficient amount to produce methylacrylic acid, said methylacrylic acid pathway comprising:

a 4-hydroxybutyryl-CoA mutase, a succinyl-CoA reductase, a succinate reductase, a 4-
15 hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 3-hydroxyisobutyryl-CoA dehydratase, a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase, a methacrylyl-CoA hydrolase, a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase, a 3-
20 hydroxyisobutyryl-CoA hydrolase, a 3-hydroxyisobutyrate dehydratase, a methylmalonyl-CoA mutase, a methylmalonyl-CoA epimerase, a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase, a methylmalonyl-CoA hydrolase, a methylmalonate reductase, a methylmalonyl-CoA reductase (aldehyde forming), a 3-hydroxyisobutyrate dehydrogenase, a methylmalonyl-CoA reductase (alcohol forming)
25 or a 3-hydroxyisobutyrate dehydratase;

said isopropanol pathway comprising at least one exogenous nucleic acid encoding an isopropanol pathway enzyme expressed in a sufficient amount to produce isopropanol, said isopropanol pathway comprising:

an isopropanol dehydrogenase, an acetyl-CoA acetyl thiolase, an acetoacetyl-CoA
30 transferase, an acetoacetyl-CoA hydrolase, an acetoacetyl-CoA synthetase, or an acetoacetate decarboxylase.

39. The non-naturally occurring microbial organism of any one of claims 36-38, further comprising an acetyl-CoA pathway comprising at least one exogenous nucleic acid encoding an acetyl-CoA pathway enzyme expressed in a sufficient amount to produce acetyl-CoA, said acetyl-CoA pathway comprising:

5 a pyruvate kinase, a pyruvate dehydrogenase, a pyruvate ferredoxin oxidoreductase, a pyruvate formate lyase, a pyruvate formate lyase activating enzyme, or a formate dehydrogenase.

40. The non-naturally occurring microbial organism of any one of claims 36-38, further comprising a succinyl-CoA pathway comprising at least one exogenous nucleic acid encoding a succinyl-CoA pathway enzyme expressed in a sufficient amount to produce succinyl-CoA, said succinyl-CoA pathway comprising:

10 a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase or a succinyl-CoA synthetase.

41. The non-naturally occurring microbial organism of claim 40, wherein said succinyl-CoA pathway further comprises a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

42. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase,

25 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

30

43. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:
- a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate reductase; and a 4-hydroxybutyraldehyde reductase,
- said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:
- an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.
44. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:
- a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase,
- said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:
- an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.
45. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol

pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a 4-hydroxybutyryl-phosphate reductase; and a 4-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

46. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming),

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

47. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming),

5 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

10 48. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

15 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol,
20 said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

25 49. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

30 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate reductase; and a 4-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

5 an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

10 50. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA reductase (aldehyde-forming); and a 4-hydroxybutyraldehyde reductase,

15 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

20 an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

25 51. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a 4-hydroxybutyryl-phosphate reductase; and a 4-hydroxybutyraldehyde reductase,

30 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

52. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

10 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming),

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

15 an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

53. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,4-butanediol pathway and an isopropanol pathway, said 1,4-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,4-butanediol pathway enzymes expressed in a sufficient amount to produce 1,4-butanediol, said first set of exogenous nucleic acids encoding:

20 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; and a 4-hydroxybutyryl-CoA reductase (alcohol-forming),

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

54. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

55. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

5 56. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

10 a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol,
15 said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 57. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

25 a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol,
30 said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

5 58. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

10 a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming),

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

15 an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 59. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

25 a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming),

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

5 60. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

10 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol,
15 said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 61. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

25 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol,
30 said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

5 62. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

10 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA reductase (aldehyde forming); and a 3-hydroxybutyraldehyde reductase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol,
15 said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 63. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:

25 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; a 3-hydroxybutyryl-CoA transferase or a 3-hydroxybutyryl-CoA synthetase or a 3-hydroxybutyryl-CoA hydrolase; a 3-hydroxybutyrate reductase; and a 3-hydroxybutyraldehyde reductase,

30 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- 5 64. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:
- 10 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming),
- said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:
- 15 an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- 20 65. A non-naturally occurring microbial organism, comprising a microbial organism having an 1,3-butanediol pathway and an isopropanol pathway, said 1,3-butanediol pathway comprising a first set of exogenous nucleic acids encoding 1,3-butanediol pathway enzymes expressed in a sufficient amount to produce 1,3-butanediol, said first set of exogenous nucleic acids encoding:
- 25 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA dehydratase; a crotonase; and a 3-hydroxybutyryl-CoA reductase (alcohol-forming),
- said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

5 66. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

10 a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol,
15 said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 67. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

25 a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol,
30 said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

5 68. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

10 a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase,

15 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 69. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

25 a succinyl-CoA reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase,

30 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

5 70. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

10 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase,

15 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 71. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

25 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase,

30 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- 5 72. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:
- 10 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA transferase, a 3-hydroxyisobutyryl-CoA synthetase or a 3-hydroxyisobutyryl-CoA hydrolase; and a 3-hydroxyisobutyrate dehydratase,
- 15 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:
- an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

- 20 73. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:
- 25 a succinate reductase; a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA mutase; a 3-hydroxyisobutyryl-CoA dehydratase; and a methacrylyl-CoA transferase, a methacrylyl-CoA synthetase or a methacrylyl-CoA hydrolase,
- 30 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

5 74. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

10 a methylmalonyl-CoA mutase; a methylmalonyl-CoA reductase (aldehyde forming); a 3-hydroxyisobutyrate dehydrogenase; and a 3-hydroxyisobutyrate dehydratase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

15 an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 75. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

25 a methylmalonyl-CoA mutase; a methylmalonyl-CoA epimerase; a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase or a methylmalonyl-CoA hydrolase; a methylmalonate reductase; a 3-hydroxyisobutyrate dehydrogenase; and a 3-hydroxyisobutyrate dehydratase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

5 76. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

10 a methylmalonyl-CoA mutase; a methylmalonyl-CoA transferase, a methylmalonyl-CoA synthetase or a methylmalonyl-CoA hydrolase; a methylmalonate reductase; a 3-hydroxyisobutyrate dehydrogenase; and a 3-hydroxyisobutyrate dehydratase,

said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

15 an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

20 77. A non-naturally occurring microbial organism, comprising a microbial organism having a methylacrylic acid pathway and an isopropanol pathway, said methylacrylic acid pathway comprising a first set of exogenous nucleic acids encoding methylacrylic acid pathway enzymes expressed in a sufficient amount to produce methylacrylic acid, said first set of exogenous nucleic acids encoding:

a methylmalonyl-CoA mutase; a methylmalonyl-CoA reductase (alcohol forming); and a 3-hydroxyisobutyrate dehydratase,

25 said isopropanol pathway comprising a second set of exogenous nucleic acids encoding isopropanol pathway enzymes expressed in a sufficient amount to produce isopropanol, said second set of exogenous nucleic acids encoding:

30 an acetyl-CoA acetyl thiolase; an acetoacetyl-CoA transferase or an acetoacetyl-CoA hydrolase or an acetoacetyl-CoA synthetase; an acetoacetate decarboxylase; and an isopropanol dehydrogenase.

78. The non-naturally occurring microbial organism of any one of claims 42-77, further comprising an acetyl-CoA pathway comprising a third set of exogenous nucleic acids encoding acetyl-CoA pathway enzymes expressed in a sufficient amount to produce acetyl-CoA, said third set of exogenous nucleic acids encoding:

5 a pyruvate kinase; and a pyruvate dehydrogenase or a pyruvate ferredoxin oxidoreductase; or a pyruvate formate lyase, a pyruvate formate lyase activating enzyme and a formate dehydrogenase.

79. The non-naturally occurring microbial organism of any one of claims 42-77, further comprising a succinyl-CoA pathway comprising a third set of exogenous nucleic acids encoding succinyl-CoA pathway enzymes expressed in a sufficient amount to
10 produce succinyl-CoA, said third set of exogenous nucleic acids encoding:

a PEP carboxykinase, a PEP carboxylase, a malate dehydrogenase, a fumarase, a fumarate reductase, a succinyl-CoA transferase and a succinyl-CoA synthetase.

80. The non-naturally occurring microbial organism of claim 79, wherein said third
15 set of exogenous nucleic acids further encodes a pyruvate carboxylase or a methylmalonyl-CoA carboxytransferase.

81. The non-naturally occurring microbial organism of any one of claims 36-38 or 42-77, wherein said exogenous nucleic acid is a heterologous nucleic acid.

82. The non-naturally occurring microbial organism of any one of claims 36-38 or
20 42-77, wherein said non-naturally occurring microbial organism is in a substantially anaerobic culture medium.

83. A method for producing 1,4-butanediol and isopropanol, comprising culturing a non-naturally occurring microbial organism of any one of claims 36, or 42-53 under conditions and for a sufficient period of time to produce 1,4-butanediol and isopropanol.

84. The method of claim 83, wherein said conditions comprise substantially
25 anaerobic culture conditions.

85. The method of claim 83, wherein said exogenous nucleic acid is a heterologous nucleic acid.

86. A method for producing 1,3-butanediol and isopropanol, comprising culturing a non-naturally occurring microbial organism of any one of claims 37, or 54-65 under conditions and for a sufficient period of time to produce 1,3-butanediol and isopropanol.

5 87. The method of claim 86, wherein said conditions comprise substantially anaerobic culture conditions.

88. The method of claim 86, wherein said exogenous nucleic acid is a heterologous nucleic acid.

10 89. A method for producing methylacrylic acid and isopropanol, comprising culturing a non-naturally occurring microbial organism of any one of claims 38, or 67-77 under conditions and for a sufficient period of time to produce methylacrylic acid and isopropanol.

90. The method of claim 89, wherein said conditions comprise substantially anaerobic culture conditions.

15 91. The method of claim 89, wherein said exogenous nucleic acid is a heterologous nucleic acid.