

<u>Figure 1</u>: Effect of 8-cyclopropyl-3-[2-(3-fluorophenyl)ethyl]-7,8-dihydro-3*H*-[1,3]oxazino[6,5-*g*][1,2,3]benzotriazine-4,9-dione (3, 10, 30 mg/kg, i.p.) in the marble burying test in NMRI mice.



Values: mean ± sem, n = 6 mice / group.


<u>Figure 2</u>: Effect of citalopram on its own (1, 3 and 10 mg/kg, i.p.) (A) and of the coadministration of 8-cyclopropyl-3-[2-(3-fluorophenyl)ethyl]-7,8-dihydro-3*H*-[1,3]oxazino[6,5-*g*][1,2,3]benzotriazine-4,9-dione (30 mg/kg, i.p.; compound (I)) and citalopram (1, 3 and 10 mg/kg, i.p.) (B) in the marble burying test in NMRI mice



Values: mean ± sem, n = 6 mice / group

** and *** : $p < 0.01$ and $p < 0.001$, respectively, *vs.* control group.

<u>Figure 3</u>: Effect of paroxetine on its own (1, 3 and 10 mg/kg i.p.) (A) and of the coadministration of 8-cyclopropyl-3-[2-(3-fluorophenyl)ethyl]-7,8-dihydro-3*H*-[1,3]oxazino[6,5-*g*][1,2,3]benzotriazine-4,9-dione (30 mg/kg, i.p.; compound (I)) and paroxetine (1, 3 and 10 mg/kg, i.p.) (B) in the marble burying test in NMRI mice



Values: mean ± sem, n = 6 mice / group

*, ** and *** : $p < 0.05$, $p < 0.01$ and $p < 0.001$, respectively, *vs.* control group.