PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 :		(11) International Publication Number: WO 00/32189
A61K 31/415	A1	(43) International Publication Date: 8 June 2000 (08.06.00)

(21) International Application Number: PCT/US99/28411

(22) International Filing Date: 30 November 1999 (30.11.99)

(30) Priority Data:
60/110,333 30 November 1998 (30.11.98) US

(71) Applicant *(for all designated States except US)*: G. D. SEARLE & CO. [US/US]; 5200 Old Orchard Road, Skokie, IL 60077 (US).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: GAO, Danchen [CN/US]; 1446 S. Prairie Avenue, Chicago, IL 60605 (US). HLINAK, Anthony, J. [US/US]; 1803 Longmeadow Drive, Lindenhurst, IL 60046 (US). MAZHARY, Ahmad, M. [US/US]; 1461 Lancaster Lane, Algonquin, IL 60102 (US). TRUELOVE, James, E. [US/US]; 1006 Regency Lane, Libertyville, IL 60048 (US). VAUGHN, Margaret, B., Woodhull [US/US]; 764 Rosewood, Winnetka, IL 60093 (US).

(74) Agent: STREIT, Richard, J.; Ladas & Parry, 224 S. Michigan Avenue, Chicago, IL 60604 (US).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.

(54) Title: CELECOXIB COMPOSITIONS

(57) Abstract

Pharmaceutical compositions are provided comprising one or more orally deliverable dose units, each comprising particulate celecoxib in an amount of about 10 mg to about 1000 mg in intimate mixture with one or more pharmaceutically acceptable excipients. The compositions are useful in treatment or prophylaxis of cyclooxygenase–2 mediated conditions and disorders.