




FIG.1

Brain and Bone Uptakes in Normal Mice




FIG.2




FIG.3




FIG.4

Compound	K_i (nM $\pm$ SEM)
<u>2</u>	<u>2.5</u> $\pm$ 0.4
<u>5</u>	<u>150</u> $\pm$ 30
<u>6</u>	<u>10</u> $\pm$ 3.3
<u>8</u>	<u>91.2</u> $\pm$ 8.7
<u>9</u>	<u>2.2</u> $\pm$ 0.2
<u>11a</u>	<u>6.8</u> $\pm$ 1.4
<u>11b</u>	<u>4.5</u> $\pm$ 0.9
<u>11e</u>	<u>14.2</u> $\pm$ 0.9
<u>13a</u>	<u>7.5</u> $\pm$ 0.8
<u>13b</u>	<u>9.0</u> $\pm$ 1.0
<u>13e</u>	<u>21</u> $\pm$ 8.0
<u>14a</u>	<u>3.6</u> $\pm$ 0.8
<u>14b</u>	<u>5.0</u> $\pm$ 1.6
<u>14e</u>	<u>6.8</u> $\pm$ 0.8
<u>16a</u>	<u>7.5</u> $\pm$ 1.5
<u>16b</u>	<u>8.5</u> $\pm$ 2.5
<u>16e</u>	<u>15.5</u> $\pm$ 0.5

Potencies (K_i) of compounds for displacement
of I-125-IMPY binding to amyloid plaques
in AD brain homogenates

FIG.5




FIG.6




FIG.7




FIG.8




FIG.9