

Breast Fibroadenoma Susceptibility Mutations And Use Thereof

Technical Field

5 The present disclosure relates to a method of assaying the risk or susceptibility of fibroadenoma development in a human subject. The disclosed method can be or at least form part of the diagnosis to confirm the occurrence of fibroadenoma in the subject. More particularly, the disclosed method sets out to detect one or more mutations residing within the *MED12* gene of the subject that presence of these mutations has shown significant
10 association with the occurrence of fibroadenomas in a human subject.

Background

Fibroadenomas (FAs) are benign breast tumors that represent the most frequently occurring breast tumors in women under the age of 30 years^{1,2}. Often observed in adolescent girls and
15 young adult women, fibroadenomas are clinically known to be hormone-dependent and to fluctuate in size according to periods of pregnancy and menopause³. A study of 265,402 women in China reported a fibroadenoma incidence of 241 per 100,000 among women under 35 years and 165 per 100,000 among women of age 35-39 years⁴. Histologically, fibroadenomas comprise an admixture of stromal and epithelial cells⁵. Although benign,
20 fibroadenomas are reported to be associated with an approximately two-fold increase in risk of developing invasive breast carcinoma in 20 years⁶. The diagnosis of fibroadenoma is typically achieved by biopsy, and patients with larger lesions are often subjected to surgery which can incur cost, anxiety and in rare cases, procedure-related complications⁷. At present, little is known about the genetic abnormalities that underlie fibroadenoma particularly when
25 compared to breast carcinoma, where much recent progress has been made in the characterization of its mutational landscape^{8,9}. For example, previous targeted mutational screens of TP53 in fibroadenomas have been equivocal. One study reported that one out of eight (12.5%) fibroadenomas exhibited a non-silent TP53 mutation¹⁰, whereas another study reported no somatic TP53 mutations in fibroadenomas from women who remained unaffected
30 by breast cancer after an average follow-up of ten years¹¹. A single PIK3CA mutation has also been reported from a screen of ten fibroadenoma tumors¹². Based upon the findings of these studies, it appears that certain genetic makeups may inevitably predispose an individual

to greater risk of FAs development. Therefore, early identification of such genetic attributes or methods allowing one to discover the likelihood of genetic deficiencies is greatly desired.

Summary

- 5 The present disclosure aims to provide a method of assaying the risk of breast fibroadenomas occurrence in a human subject, preferably a female subject, through genotyping a specific allele or gene of the subject.

Still, an object of the present disclosure is to bring forth a method capable of serving as
10 confirming diagnosis or forming at least part of the confirming diagnosis towards the occurrence of fibroadenoma in a human subject.

A further object of the present disclosure is to offer a method of assaying susceptibility and/or confirming diagnosis of fibroadenoma development in a human subject by detecting
15 one or more mutations located in the *MED12* gene using any genotyping approaches known in the art.

Another object of the present disclosure is to provide a method of detecting mutations resulting particularly in non-synonymous substitution in the encoded mediator complex
20 subunit 12 (*MED12*). The mutations to be detected are associated with higher risk of fibroadenomas occurrence in a female subject.

At least one of the preceding objects is met, in whole or in part, by the present invention, in which one of the embodiments of the present invention involves a method of assaying
25 susceptibility and/or confirming diagnosis of breast fibroadenomas development in a human subject. The method essentially comprises the steps of performing a nucleic acid-based assay to analyze an isolated polynucleotide encoding at least exon 2 of *MED12* gene from a sample acquired from the human subject; and regarding the human subject with greater susceptibility and/or confirming diagnosis of breast fibroadenomas development by detecting a mutation in
30 the isolated polynucleotide. Preferably, the mutation is a splice site mutation located at position -8 of exon 2 of the *MED12* gene, a missense mutation located at codon 44 of cDNA of the *MED12* gene or a missense mutation located at codon 36 of cDNA of the *MED12* gene.

In several preferred embodiments, the sample comprises stromal tissues which may acquire the mutation through one or more somatic events progressively acquired in the subject.

- 5 Some embodiments of the disclosed method preferably detect the missense mutation, which is located at nucleotide position 107 of codon 36 cDNA of the *MED12* gene.

For a number of embodiments, the missense mutation is located at position 130 and/or 131 of codon 44 cDNA of the *MED12* gene. More preferably, the missense mutation results in
 10 p.G44A, p.G44C, p.G44D, p.G44R, p.G44S, or p.G44V in a polypeptide translated from the *MED12* gene.

In several preferred embodiments, the disclosed method may include additional steps of detecting at least one mutation located at *PIK3CA* and/or *TP53* gene of the subject upon
 15 detecting a mutation in the isolated polynucleotide encoding at least exon 2 of *MED12* gene; and regarding developed fibroadenoma in the subject as benign state in the absence of detectable mutation located at *PIK3CA* and/or *TP53* gene.

Brief Description of The Drawings

20 Fig. 1 is a schematic diagram showing the distribution of *MED12* exon 2 mutations in which the top panel shows regions of deletions, the middle panel and bottom respectively show nucleotide changes associated with point mutations and the corresponding codon alterations;

25 Fig. 2 shows results of Genomic DNA Sanger sequencing of *MED12* variants in eight fresh frozen FAs and their matched whole-blood;

Fig. 3 shows results of Complementary DNA (cDNA) Sanger sequencing of *MED12* variants in eight fresh frozen FAs and their matched whole-blood that variant
 30 peaks were unambiguous except for Sample002, possibly due to RNA degradation.;

Fig. 4 (a) is Hematoxylin and eosin (H&E) stained section of Sample006 with the epithelial compartments marked in green and (b) shows respective Sanger sequencing results of *MED12* bulk tissue, epithelial and stromal compartments, revealing that p.G44D mutations in *MED12* are exclusive to the stromal compartment;

Fig. 5 (a) is a heat map showing differential activation of gene sets associated with breast cancer and estrogen signaling associated with *MED12* alterations in FA and unsupervised clustering of gene sets with significantly differential activation scores as determined by GSVA, and (b) is a GSEA enrichment plot that genes are rank-ordered according to fold-change between mutant *MED12* and wild-type *MED12* FA samples (bottom panel), with Genes upregulated > 4x in UL being indicated as black bars in the middle panel.;

Fig. 6 is a GSEA enrichment plot against genes upregulated 2x in UL instead of 4x, as shown in Fig.5b; and

Fig. 7 shows (a) cDNA sequence of *MED12*, as in Seq ID No.15, and (b) amino acid sequence of *MED12* peptide, as in Seq ID No.16.

Detailed Description

The present invention may be embodied in other specific forms without departing from its structures, methods, or other essential characteristics as broadly described herein and claimed hereinafter. The described embodiments are to be considered in all respects only as illustrative, and not restrictive. The scope of the invention is, therefore, indicated by the appended claims, rather than by the description provided hereinafter. All changes that come within the meaning and range of equivalency of the claims are to be embraced within their scope.

Unless specified otherwise, the terms "comprising" and "comprise" as used herein, and grammatical variants thereof, are intended to represent "open" or "inclusive" language such that they include recited elements but also permit inclusion of additional, un-recited elements.

- 5 As used herein, the phrase "in embodiments" means in some embodiments but not necessarily in all embodiments.

As used herein, the terms "approximately" or "about", in the context of concentrations of components, conditions, other measurement values, etc., means +/- 5% of the stated value, or
10 +/- 4% of the stated value, or +/- 3% of the stated value, or +/- 2% of the stated value, or +/- 1% of the stated value, or +/- 0.5% of the stated value, or +/- 0% of the stated value.

The term "polynucleotide" or "nucleic acid" as used herein designates mRNA, RNA, cRNA, cDNA or DNA. The term typically refers to oligonucleotides greater than 30 nucleotide
15 residues in length.

The term "primer" used herein throughout the specification refers to an oligonucleotide which, when paired with a strand of DNA, is capable of initiating the synthesis of a primer extension product in the presence of a suitable polymerizing agent. The primer is preferably single-
20 stranded for maximum efficiency in amplification but can alternatively be double-stranded. A primer must be sufficiently long to prime the synthesis of extension products in the presence of the polymerization agent. Primers can be "substantially complementary" to the sequence on the template to which it is designed to hybridize and serve as a site for the initiation of synthesis. By "substantially complementary", it is meant that the primer is sufficiently
25 complementary to hybridize with a target polynucleotide. Preferably, the primer contains no mismatches with the template to which it is designed to hybridize but this is not essential. For example, non-complementary nucleotide residues can be attached to the 5' end of the primer, with the remainder of the primer sequence being complementary to the template. Alternatively, non-complementary nucleotide residues or a stretch of non-complementary
30 nucleotide residues can be interspersed into a primer, provided that the primer sequence has sufficient complementarity with the sequence of the template to hybridize therewith and thereby form a template for synthesis of the extension product of the primer.

The term "gene" as used herein may refer to a DNA sequence with functional significance. It can be a native nucleic acid sequence, or a recombinant nucleic acid sequences derived from natural source or synthetic construct. The term "gene" may also be used to refer to, for
5 example and without limitation, a cDNA and/or an mRNA encoded by or derived from, directly or indirectly, genomic DNA sequence.

One aspect of the present disclosure refers to a method of assaying susceptibility and/or confirming diagnosis of breast fibroadenomas development in a human subject, preferably a
10 female human subject. Essentially, the method of assaying comprises the steps of performing a nucleic acid-based assay to analyze an isolated polynucleotide encoding at least exon 2 of *MED12* gene from a sample acquired from the human subject; and regarding the human subject with greater susceptibility and/or confirming diagnosis of breast fibroadenomas development by detecting a mutation in the isolated polynucleotide. The sample applicable
15 for the disclosed method can be any biological sample having the extractable or accessible genetic materials suspected to have carrying the mutations of interested, either acquired or constitutional, and detectable in the nucleic acid-based assay. The sample can be, but not limited to, biopsy tissue or blood sample of the subject. More preferably, the sample or the biopsy tissue comprises stromal tissues, which is found by the inventors of the present
20 disclosure being prone to adversely affected, likely owing to dysregulated extracellular matrix organization, in a significant extent by the mutations of interested located at exon 2 of the *MED12* gene. The sample may be subjected to pre-treatment to isolate the preferred tissue type prior to extracting the genetic material for analysis.

25 Further, the polynucleotide to be reacted or analyzed in the nucleic-acid based assay can be directly or indirectly derived from the genetic materials extracted or obtained from the sample of the subject. Polynucleotides can be acquired directly from the sample source, but not limited to, by digesting or cutting the targeted gene segment utilizing restriction enzymes recognizing the specific restriction site located adjacent to the interested portion. On the other
30 hand, the polynucleotides can be amplicons generated and duplicated from the extracted genetic materials through any known PCR or the like approaches. These amplicons are

further subjected to the analysis of the nucleic acid-based assay to identify the possible mutations resulting in the occurrence of fibroadenomas.

According to several preferred embodiments, the nucleic acid-based assay can be performed to identify and/or detect the mutations comprises sequencing the polynucleotide. More specifically, the sequencing approach implementable in the present disclosure to effect the detection can be Sanger sequencing and/or ultra-deep targeted amplicon sequencing which is effective and capable of catering highly precise and reliable result in identifying the interested mutations in exon 2 of the *MED12* gene. Primer pairs Seq ID No. 1 and Seq ID No. 2 listed in Table 1 below are one embodiment of the primers usable in the present disclosure to realize the sequencing process on exon 2 of the *MED12* gene. The sequencing process shall provide reliable reading about the sequence of the polynucleotide that substantial outcome can be inferred thereby regarding the tested subject, at least the in the sample, whether the FA-associated mutation are carried. Furthermore, Seq ID No.3-14 are sequences of the primers pairs can be used to perform the ultra-deep targeted amplicon sequencing. The details of the Sanger sequencing and/or ultra-deep targeted amplicon sequencing utilizing the listed primer pairs are further elaborated in the examples incorporated hereafter. It is important for other skilled artisans to appreciate the fact that the disclosed method can be conducted utilizing other known sequencing equivalent or non-equivalent procedures or approaches to detect presence of the interested mutation in the analyzed polynucleotides and such modification shall not depart from the scope of the present disclosure. Other known processes implementable to identify or assist in identifying these mutations can be any one of, but not limited to, temperature gradient gel electrophoresis, capillary electrophoresis, amplification-refractory mutation system-polymerase chain reaction (ARMS-PCR), dynamic allele-specific hybridization (DASH), target capture for next generation sequencing (NGS), high-density oligonucleotide SNP arrays or Restriction fragment length polymorphism (RFLP).

Table 1 Primers for sequencing

Primer pairs of Sanger sequencing	Sequences (5' to 3')
Seq ID No. 1	TGTTCTACACGGAACCCTCCTC
Seq ID No. 2	CTGGGCAAATGCCAATGAGAT

Primer pairs for ultra-deep sequencing	Sequences (5' to 3')
Seq ID No. 3	TTCTCCTGCCCTACTCTCCAC
Seq ID No. 4	CAGGCTGGTTATTGAAACCTTG
Seq ID No. 5	CCCTAAGGAAAAAACAATAAACGC
Seq ID No. 6	CTGCCATGCTCATCCCCAGA
Seq ID No. 7	CTTGTTCTTCTTTTCTCCTGCC
Seq ID No. 8	GTTTACATTCAAGGCCGTCAG
Seq ID No. 9	CAACTAAACGCCGCTTTCCTG
Seq ID No. 10	AAGCTGACGTTCTTGGAAGTGC
Seq ID No. 11	GCTTTCCTGCCTCAGGATGAACT
Seq ID No. 12	CCTTGGCAGGATTGAAGCTGAC
Seq ID No. 13	GATGAACTGACGGCCTTGAATGTA
Seq ID No. 14	CCTGGCAGAGTTGTCTCACCTTG

Pursuant to the preferred embodiments, the disclosed method targets to analyze and/or identify multiple potential mutations residing in exon 2 the *MED12* gene concurrently. One of the interesting mutations is a splice site mutation located at position -8 of exon 2 of the *MED12* gene. More specifically, the splice site mutation is an intronic T>A substitution located 8bp upstream of exon 2 of the *MED12* gene in the genomic DNA. This splice site mutation results in an aberrant splice acceptor site further leading to retention of the last six bases of the *MED12* gene intron 1 in the mRNA transcribed thereof.

Another mutation to be identified by the disclosed method, in a number of embodiments, is a missense mutation located at codon 36 of cDNA of the *MED12* gene. Preferably, the missense mutation is located at position 107 of codon 36 cDNA of the *MED12* gene causing a non-synonymous substitution of the encoded amino acid thereof. More preferably, the disclosed method seeks to detect presence of any mutation resulting in any one of p.L36R or p.L36P. Correspondingly, the equivalent mutations positioned on the cDNA result in the non-synonymous substitution of p.L36R and p.L36P are respectively c.107T>G and c.107T>A. Considering degeneracy of the codon involved, other mutations may result in similar synonymous substitution of the involved amino acids, p.L36R and p.L36P, besides c.107T>G and c.107T>A.

According to other preferred embodiments, the disclosed method also aims to identify a missense mutation located at codon 44 of cDNA of the *MED12* gene. Specifically, the

missense mutation is located at position 130 and/or 131 of codon 44 cDNA of the *MED12* gene giving rise to non-synonymous substitution of encoded amino acids such as p.G44A, p.G44C, p.G44D, p.G44R, p.G44S, or p.G44V in a polypeptide translated from the *MED12* gene.

5

It is important to note that inventors of the present disclosure found that the aforesaid mutations may subsequently upregulate or activate other genes associated with extracellular matrix organization, estrogen signaling, and TGF β and Wnt signaling. Up-regulation or uncontrolled activation of these genes or gene products shall hence promote development of FA in the subject.

10

In several preferred embodiments, the disclosed method may include additional steps of detecting at least one mutation located at *PIK3CA* and/or *TP53* gene of the subject upon detecting a mutation in the isolated polynucleotide encoding at least exon 2 of *MED12* gene; and regarding developed fibroadenoma in the subject as benign state in the absence of detectable mutation located at *PIK3CA* and/or *TP53* gene.

15

Another aspect of the present disclosure may include use of Seq. ID No. 1 and 2 in the preparation of a platform for nucleic-acid based assay for assaying susceptibility and/or confirming diagnosis of breast fibroadenomas development in a female human subject.

20

Likewise, in further aspect, the present disclosure may include use of Seq. ID No. 3 and 4, Seq. ID No. 5 and 6, Seq. ID No. 7 and 8, Seq. ID No. 9 and 10, Seq. ID No. 11 and 12, and/or Seq. ID No. 13 and 14 in preparation of a platform for nucleic-acid based assay for assaying susceptibility and/or confirming diagnosis of breast fibroadenomas development in a female human subject. Preferably, the use of Seq. ID No. 3 and 4, Seq. ID No. 5 and 6, Seq. ID No. 7 and 8, Seq. ID No. 9 and 10, Seq. ID No. 11 and 12, and/or Seq. ID No. 13 and 14 in the mentioned platform facilitates or materializes identification of a missense mutation located at codon 44 of cDNA of *MED12* gene or a missense mutation located at codon 36 of cDNA of *MED12* gene. Presence of at least one of these mutations in the breast tissue, more preferably stromal cells of the breast tissue, has been shown to associate with greater risk in developing FA in relation to those with the wild type allele by the present disclosure.

25
30

The following example is intended to further illustrate the invention, without any intent for the invention to be limited to the specific embodiments described therein.

5 Example 1

A total of 98 fibroadenoma tumors were included in this study, of which 12 were from fresh frozen tumors and a further 86 from archival FFPE (formalin-fixed paraffin-embedded) samples. Tumors and whole-blood were obtained from patients undergoing surgical excision of fibroadenoma or from the SingHealth Tissue Repository, with signed informed consent.
10 Archival samples were obtained from the Department of Pathology of Singapore General Hospital. Clinicopathological information for subjects (age and tumor size) was reviewed retrospectively.

Genomic DNA (gDNA) from fresh frozen tissue was extracted and purified using the Qiagen
15 Blood and Cell Culture DNA kit. In the case of FFPE samples, the Qiagen FFPE DNA kit was used on freshly sectioned FFPE tissue. Genomic DNA yield and quality were determined using PicogreenTM fluorometric analysis as well as visual inspection of agarose gel electrophoresis images.

20 Example 2

Native genomic DNA was fragmented with the *Covaris*TM S2 (Covaris) system using recommended settings. Sequencing adaptor ligation was performed using the Truseq Paired-End Genomic DNA kit (Illumina). For enrichment of coding sequences, the present disclosure used the SureSelectXTTM Human All Exon v3 (50 Mb) kit (Agilent Technologies)
25 according to manufacturer's recommended protocol. Exome-enriched libraries were then sequenced on Illumina's HiSeq 2000 sequencing platform to generate 76 bp paired-end reads. Bioinformatics analysis, comprising of sequence alignment, variant calling and identification of candidate somatic variants was performed as described in previous work²⁸. For point mutations, at least 10 variant reads in the tumor and a total read depth of 10 in the normal
30 sample was required. In the case of Indels, a support of at least 20 variant reads amounting to at least 10% of total reads was required. Indels overlapping simple repeat regions were also discarded. All remaining candidate variants were visually inspected in a genome browser to

identify missed probable germline variants or those in regions of anomalous alignment. The variant calling pipeline missed the p.Glu33_Asp34insProGln aberrant splice site variant in Sample004 as it was not in the exome capture kit manufacturer's target region file. It was later identified from a systematic visual inspection of *MED12* in a genome browser as it was the only gene recurrently mutated in multiple samples. All candidate somatic variants were confirmed by Sanger sequencing.

The present disclosure used the following PCR primer pair to identify mutations in *MED12* exon 2; forward primer: TGTTCTACACGGAACCCTCCTC, reverse primer: CTGGGCAAATGCCAATGAGAT, Tm: 54.6 °C, 56.3 °C, product length: 373 bp. PCR amplification was conducted using neat DNA and Platinum™ Taq Polymerase (Life Technologies). PCR cycling regime included one cycle at 95 °C for 10 min, 35 cycles at 95 °C for 30 s, 58 °C for 30 s and 72 °C for 1 min, and one cycle at 72 °C for 10 min. BigDye Terminator v.3.1 kit (Applied Biosystems) was used for bi-directional sequencing on generated PCR amplicons and products were fractionated employing ABI PRISM 3730 Genetic Analyzer (Applied Biosystems). Sequencing traces were aligned to reference sequences using Lasergene 10.1 (DNASTAR) and were visually analyzed.

Example 3

For sensitive detection of low-frequency variants in *MED12* exon 2, the present disclosure further used ultra-deep targeted amplicon sequencing. Six PCR amplicons were designed and tiled across exon 2 of *MED12* using primers pairs listed in Table 2 below.

Table 2. PCR amplicon sequencing primers used in ultra-deep targeted amplicon sequencing of *MED12* exon 2

Gene	Region	Primer Name	Primer sequence (5' -> 3')	Melting Point	PCR product size (bp)
MED12	Exon2	MD12-ex2-2F	TTCTCCTGCCCTACTCTCCCA	56.2	125
		MD12-ex2-2R	CAGGCTGGTTATTGAAACCTTG	53.2	
		MD12-ex2-3F	CCCTAAGGAAAAAACAATA	55.9	115
		MD12-ex2-3R	CTGCCATGCTCATCCCCAGA	58.1	

	MD12-ex2-1F	CTTGTCCTTCTTTCTCCTGC C	55.3	117
	MD12-ex2-1R	GTTTTACATTCAAGGCCGTCA G	53.2	
	MD12-ex2-4F	CAACTAAACGCCGCTTTCCTG	56.4	119
	MD12-ex2-4R	AAGCTGACGTTCTTGGAAGTGA C	58.2	
	MD12-ex2-5F	GCTTTCCTGCCTCAGGATGAA CT	57.6	121
	MD12-ex2-5R	CCTTGGCAGGATTGAAGCTGA C	57.5	
	MD12-ex2-6F	GATGAAGTACGGCCTTGAAT GTA	57	124
	MD12-ex2-6R	CCTGGCAGAGTTGTCTCACCT TG	57.6	

The present disclosure then used Fluidigm's Access Array System to generate and pool the amplicons according to manufacturer's instructions. For each sample, 50 ng of genomic DNA was used as template. Sequencing library preparation of the pooled amplicons was performed using the TruSeq HT DNA Sample Preparation Kit (Illumina) according to manufacturer's instructions. Sequencing was performed on the Illumina MiSeq next-generation sequencing platform for 150 cycles using the MiSeq Reagent kit v3.

Bioinformatics analysis of sequencing reads was performed as follows. Briefly, undetermined ('N') base calls at the ends of reads were trimmed. Following this, the 5' end of each read was trimmed by 25 bases to eliminate the possibility of primer inclusion. The Burrows-Wheeler Alignment²⁶ (BWA) tool (0.6.2) was used to align the resulting reads to the reference human genome (hg19). For more sensitive detection of insertions and deletions (indels), the present disclosure also ran a separate alignment process using modified settings (o=2, e=30, d=30, O=0, E=0, L=0). Indels were identified through manual inspection, whereas automated detection of point mutations was performed using the samtools²⁷ (0.1.18) *mpileup* tool. Variant calls were restricted to regions covered by amplicons generated from primers pairs provided in the Table 1. Variant allele frequencies were calculated for each position in the targeted region, and those that exceeded a threshold of 5% were considered candidate variants. In order to minimize the possibility of PCR-induced artifacts, variants were only considered valid if present in at least two amplicons. Candidate variants had at least 21,620 sequencing reads overlapping them, with an average coverage of 184,526 reads.

- To ascertain the sensitivity of our assay, positive control samples containing spiked-in validated mutant *MED12* at allele frequencies (15%, 10%, 5%, 3%) were generated via serial dilution. The present disclosure accurately detected variants in positive control samples at allele frequencies down to 3%. The present disclosure also calculated alternate (non-reference) allele frequencies across all positions in our target region in order to estimate the likelihood of error from sequencing and alignment artifacts. The mean alternate allele frequency was 0.281% with a standard deviation of 1.09%. Thus, our detection threshold of 5% exceeds four standard deviations from the estimated background error rate.
- To identify genes with recurrent somatic mutations across multiple samples, the present disclosure first sequenced the exomes of eight fresh frozen FA tumors together with matched whole-blood to a mean coverage of 124x, with an average of 87% of bases covered by at least 20 reads in each sample.

Table 3. Summary of whole-exome sequencing of FA tumors and matched normal tissue (whole-blood). Highlighted samples contain somatic *MED12* exon 2 mutations.

No.	Sample	Sample Type	Bases in Target Region	Reads Mapped to Target Region*	Ave. Depth Per Targeted Base	Targeted Bases with Depth at Least 1X (%)	Targeted Bases with Depth at Least 20X (%)	Candidate somatic mutations
1	Sample002N	Normal	51,860,012	117,269,831	130	95.6	88	1
	Sample002T	Tumor	51,860,012	164,371,773	184	95.9	90	
2	Sample004N	Normal	51,860,012	114,347,376	127	95.6	88	19
	Sample004T	Tumor	51,860,012	114,347,376	106	95.5	86	
3	Sample006N	Normal	51,860,012	98,767,446	111	95.4	86	7
	Sample006T	Tumor	51,860,012	161,158,144	155	95.8	90	
4	Sample007N	Normal	51,860,012	116,511,803	130	95.4	88	7
	Sample007T	Tumor	51,860,012	113,165,060	126	95.5	88	
5	Sample009N	Normal	51,860,012	83,736,104	93	95.5	86	7
	Sample009T	Tumor	51,860,012	114,378,362	128	95.6	88	
6	Sample010N	Normal	51,860,012	119,910,977	136	95.5	87	4
	Sample010T	Tumor	51,860,012	98,563,020	110	95.4	86	
7	Sample011N	Normal	51,860,012	112,204,764	125	95.5	88	1
	Sample011T	Tumor	51,860,012	94,612,966	105	95.3	86	
8	Sample012N	Normal	51,860,012	84,410,000	95	95.3	85	5
	Sample012T	Tumor	51,860,012	114,280,813	128	95.5	88	
Average				113,877,238	124	96	87	6

Consistent FA being a benign tumor, samples had an average of only seven somatic mutations. Almost all genes were found to be mutated only once. These included tumor suppressors such as *NFI* and *RBI*. The only gene that was recurrently mutated was *MED12* (mediator complex subunit 12), which is a member of the Mediator Complex, a multiprotein complex that is widely involved in transcriptional regulation of gene expression. Four out of the eight FA samples sequenced (50%) contained somatic mutations in exon 2 of *MED12* as presented in Table 4.

Table 4 List of candidate somatic mutations identified from whole-exome sequencing of eight FAs.

No	Gene Symbol	Sample	Transcript ID	Nucleotide (genomic)	Nucleotide (cDNA)	Amino acid (protein)	Mutation type
1	MED12	Sample002	CCDS43970.1	g.chrX:70339254 G>A	c.131 G>A	p.G44D	Missense
2	MED12	Sample006	CCDS43970.1	g.chrX:70339254 G>A	c.131 G>A	p.G44D	Missense
3	MED12	Sample007	CCDS43970.1	g.chrX:70339254 G>A	c.131 G>A	p.G44D	Missense
4	MED12	Sample004	CCDS43970.1	g.chrX:70339215 T>A	c.100-8 T>A	Splice site	Splice site
5	ANK2	Sample004	CCDS3702.1	g.chr4:114277287 G>T	c.84+29032 G>T	p.V2505L	Missense
6	C1orf173	Sample007	CCDS30755.1	g.chr1:75037715 G>T	c.3679 G>T	p.V1227L	Missense
7	C22orf23	Sample004	CCDS13962.1	g.chr22:38340198 delCCTT	c.638 delCCTT	Frameshift	Indel
8	CC2D1A	Sample006	CCDS42512.1	g.chr19:14034557 delCT	c.1873 delCT	Frameshift	Indel
9	CHD6	Sample012	CCDS13317.1	g.chr20:40033303 G>A	c.8078 G>A	p.P2693L	Missense
10	CKAP5	Sample012	CCDS31477.1	g.chr11:46799825 A>G	c.2612 A>G	p.D871G	Missense
11	CREBBP	Sample004	CCDS45399.1	g.chr16:3786805 delCT	c.4292 delCT	Frameshift	Indel
12	DNAH11	Sample009	NM_001277115	g.chr7:21781777 T>A	c.8178 T>A	p.V2716D	Missense
13	FGB	Sample004	CCDS3786.1	g.chr4:155487155 G>T	c.306+4 G>T	Splice site	Splice site
14	FRMD4A	Sample004	CCDS7101.1	g.chr10:13804618 T>A	c.441+6 T>A	Splice site	Splice site
15	GRIN3B	Sample012	CCDS32861.1	g.chr19:1003331 C>T	c.629 C>T	p.T210M	Missense
16	ISL1	Sample009	CCDS43314.1	g.chr5:50685533 C>T	c.532 C>T	p.P178S	Missense
17	IST1	Sample004	CCDS10905.1	g.chr16:71956504 C>T	c.680 C>T	p.T227M	Missense
18	KCNG4	Sample006	CCDS10945.1	g.chr16:84255957 C>T	c.1426 C>T	p.R476C	Missense
19	KIAA1211L	Sample004	CCDS42720.1	g.chr2:99454665 C>A	c.156 C>A	p.S52R	Missense
20	KRTAP1-3	Sample006	CCDS42323.1	g.chr17:39190785 delCT	c.289 delCT	Frameshift	Indel
21	LAMB4	Sample010	CCDS34732.1	g.chr7:107735743 C>G	c.1400 C>G	p.T467S	Missense
22	LPA	Sample006	CCDS43523.1	g.chr6:160998309	c.4289+6078	p.P1497L	Missense

Table 4 List of candidate somatic mutations identified from whole-exome sequencing of eight FAs.

No	Gene Symbol	Sample	Transcript ID	Nucleotide (genomic)	Nucleotide (cDNA)	Amino acid (protein)	Mutation type
				C>T	C>T		
23	LRRC10	Sample012	CCDS31856.1	g.chr12:70004273 G>A	c.346 G>A	p.E116K	Missense
24	LRRC42	Sample012	CCDS585.1	g.chr1:54432042 C>G	c.1001 C>G	p.A334G	Missense
25	LRRTM3	Sample011	CCDS7270.1	g.chr10:68686900 delT	c.226 delT	Frameshift	Indel
26	MAGEE1	Sample009	CCDS14433.1	g.chrX:75650516 T>G	c.2193 T>G	p.Y731X	Missense
27	MAPT	Sample006	CCDS45715.1	g.chr17:44055797 G>A	c.364 G>A	p.V122M	Missense
28	MYO9A	Sample007	CCDS10239.1	g.chr15:72170501 G>A	c.5811 G>A	p.M1937I	Missense
29	NF1	Sample009	CCDS42292.1	g.chr17:29560073 A>T	c.3550 A>T	p.T1184S	Missense
30	NODAL	Sample004	CCDS7304.1	g.chr10:72195115 C>T	c.818 C>T	p.A273V	Missense
31	NOTCH2	Sample004	CCDS908.1	g.chr1:120491681 delTT	c.2548 delTT	Frameshift	Indel
32	NUMA1	Sample004	CCDS31633.1	g.chr11:71724080 G>T	c.4469 G>T	p.R1490L	Missense
33	PCLO	Sample010	CCDS47630.1	g.chr7:82580690 C>T	c.9214 C>T	p.P3072S	Missense
34	PGAP1	Sample004	CCDS2318.1	g.chr2:197791238 delCTC	c.103 delCTC	In-frame	Indel
35	POM121L12	Sample004	CCDS43584.1	g.chr7:53103758 C>T	c.394 C>T	p.R132W	Missense
36	POTEA	Sample007	NM_001002920	g.chr8:43211931 G>T	c.1295 G>T	p.A464S	Missense
37	PRAF2	Sample009	CCDS14317.1	g.chrX:48929554 G>C	c.511 G>C	p.G171R	Missense
38	PSME4	Sample009	CCDS33197.2	g.chr2:54158971 G>A	c.1316+1 G>A	Splice site	Splice site
39	RARA	Sample007	CCDS11366.1	g.chr17:38510626 C>T	c.880 C>T	p.R294W	Missense
40	RB1	Sample004	CCDS31973.1	g.chr13:48881465 G>T	c.187 G>T	p.K63X	Missense
41	RB1	Sample004	CCDS31973.1	g.chr13:48937094 A>T	c.861+1 A>T	Splice site	Splice site
42	ROS1	Sample004	CCDS5116.1	g.chr6:117609731 A>T	c.6968 A>T	p.Y2323F	Missense
43	SAAL1	Sample006	CCDS31439.1	g.chr11:18112008 A>G	c.446 A>G	p.D149G	Missense
44	SCN10A	Sample004	CCDS33736.1	g.chr3:38783906 T>C	c.1982 T>C	p.L661P	Missense
45	SEMA4F	Sample007	CCDS1955.1	g.chr2:74902152 G>T	c.1139 G>T	p.R380I	Missense
46	SHROOM4	Sample007	CCDS35277.1	g.chrX:50350882 C>T	c.3260 C>T	p.T1087I	Missense
47	SIAH3	Sample009	CCDS41883.1	g.chr13:46357894 C>T	c.434 C>T	p.A145V	Missense
48	SYNE4	Sample004	NM_001039876	g.chr19:36494181 C>A	c.1362 C>A	p.T365N	Missense
49	TNFAIP3	Sample010	CCDS5187.1	g.chr6:138196931 T>C	c.593 T>C	p.V198A	Missense
50	TRPM1	Sample010	CCDS58347.1	g.chr15:31323296 G>A	c.3068 G>A	p.R1023H	Missense

Example 4

In order to further ascertain the prevalence of *MED12* exon 2 mutations in FA, the present disclosure performed ultra-deep targeted amplicon sequencing of *MED12* exon 2 in 90 additional FA samples (4 fresh frozen tissue samples and 86 archival samples). This confirmed a strikingly high *MED12* exon 2 mutation frequency in FA of 59%. Frequency of the various detected mutation is summarized in Table 5 and Fig.1.

Table 5 A tabular summary of *MED12* exon 2 mutations in Fa in comparison with corresponding mutation frequencies in UL are indicated where applicable (FA = fibroadenoma, UL = uterine leiomyoma, ins = insertion, del = deletion, fs = frameshift).

Type	cDNA	Protein	# mutated out of 98 samples in FA (%)	# mutated out of 225 samples in UL ¹³ (%)
Missense	c.131G>C	p.G44A	1 (1.1)	11 (5.0)
	c.130G>T	p.G44C	2 (2.2)	7 (3.1)
	c.131G>A	p.G44D	20 (20.4)	47 (20.9)
	c.130G>C	p.G44R	3 (3.3)	16 (7.1)
	c.130G>A	p.G44S	12 (13.3)	17 (7.6)
	c.131G>T	p.G44V	3 (3.3)	12 (5.3)
	c.128A>C	p.Q43P	1 (1.1)	3 (1.3)
	c.107T>G	p.L36R	3 (3.3)	11 (5.0)
	c.107T>A	p.L36P	1 (1.0)	0 (0.0)
Splice Site	Exon2 (-8 T>A)	p.E33_D34insPQ	4 (4.1)	10 (4.4)
Deletions	intronic -23bp c.100_101del2	p.D34fs	1 (1.1)	0 (0.0)
	c.134_151del18	p.F45_V51>F	1 (1.1)	0 (0.0)
	c.130_147del18	p.G44_P49	1 (1.1)	0 (0.0)
	c.120_149del30	p.N40_A50>N	2 (2.2)	0 (0.0)
	c.118_132del15	p.N40_G44	1 (1.1)	0 (0.0)
	c.118_135del18	p.N40_F45	2 (2.2)	0 (0.0)
Total			58 (59.2)	-

Out of the 98 FA samples sequenced, 41 (42%) had point mutations in codon 44 (20 p.G44D, 12 p.G44S, 3 p.G44R, 3 p.G44V, 2 p.G44C, 1 p.G44A). A single point mutation (1.1%) was also found in codon 43 (p.Q43P) and four (4.1%) in codon 36 (3 p.L36R, 1 p.L36P).

Additionally, seven (7.8%) samples were found to have insertions or deletions that were expected to preserve the reading frame, and one (1.1%) further sample harbored a frameshift deletion. The present disclosure also identified four samples with an intronic T>A substitution 8 bp upstream of exon 2 that resulted in an aberrant splice acceptor site, causing the last six bases of intron 1 to be retained¹³. Several lines of evidence indicate the *MED12* exon 2 mutations are somatic. The present disclosure performed Sanger sequencing on eight *MED12* mutant fresh-frozen samples with available whole-blood and confirmed that all eight mutations were somatic as indicated in Fig.2. All but one point mutations and 25% (2/8) of deletions detected in our archival samples for which there was no matched whole-blood were found to have COSMIC¹⁴ (Catalog of Somatic Mutations in Cancer) entries with reference to the Table 6, none were classified as germline variants in dbSNP¹⁵ and the 1000 Genomes Project¹⁶, and an examination of our in-house database of germline variants from a predominantly East Asian cohort of 470 subjects revealed no variants in *MED12* exon 2.

Table 6 Mutations detected in ultra-deep targeted amplicon sequencing of *MED12* exon 2 in 98 FA samples.

No .	Sample ID	Total Reads	Variant Reads	Variant allele Frequency (%)	Amino acid change	cDNA change	COSMIC	Tissue type
19	Sample03_5	130736	6222	5	p.G44V	c.131G>T	COSM131597	FFPE
20	Sample03_6	80052	8714	11	p.G44R	c.130G>C	COSM131592	FFPE
21	Sample03_7	273272	22424	8.21	p.N40_A50>N	c.120_149del30		FFPE
22	Sample03_8	159978	11466	7.17	p.G44_P49	c.130_147del18		FFPE
23	Sample03_9	369350	63858	17.29	p.G44D	c.131G>A	COSM131596	FFPE
24	Sample04_1	348710	69038	19.8	p.G44S	c.130G>A	COSM131594	FFPE
25	Sample04_2	102994	7788	7.56	p.D34fs	intronic - 23bp c.100_101del2	COSM1235330	FFPE
26	Sample04_4	106860	10694	10.01	p.G44D	c.131G>A	COSM131596	FFPE
27	Sample04_5	509920	142790	28	p.G44S	c.130G>A	COSM131594	FFPE
28	Sample04_6	202270	36546	18.07	p.G44D	c.131G>A	COSM131596	FFPE
29	Sample04_7	146698	19732	13.45	p.G44R	c.130G>C	COSM131592	FFPE

Table 6 Mutations detected in ultra-deep targeted amplicon sequencing of *MED12* exon 2 in 98 FA samples.

No .	Sample ID	Total Reads	Variant Reads	Variant allele Frequency (%)	Amino acid change	cDNA change	COSMIC	Tissue type
30	Sample048	176580	32634	18.48	p.G44S	c.130G>A	COSM131594	FFPE
31	Sample049	193484	21046	10.88	p.G44D	c.131G>A	COSM131596	FFPE
32	Sample050	108602	17060	15.7	p.G44V	c.131G>T	COSM131597	FFPE
33	Sample051	243910	29634	12.15	p.G44D	c.131G>A	COSM131596	FFPE
34	Sample053	87288	13832	15.85	p.G44D	c.131G>A	COSM131596	FFPE
35	Sample054	289626	69788	24.1	p.G44R	c.130G>C	COSM131592	FFPE
36	Sample055	142914	16800	11.76	p.L36R	c.107T>G	COSM131590	FFPE
37	Sample056	292544	34640	11.8	p.G44A	c.131G>C	COSM131595	FFPE
38	Sample057	246888	52504	21.27	p.G44C	c.130G>T	COSM131593	FFPE
39	Sample058	82052	11428	13.93	p.G44D	c.131G>A	COSM131596	FFPE
40	Sample060	189936	21498	11.32	p.F45_V51>F	c.134_151 del18		FFPE
41	Sample065	72026	9184	12.75	p.G44D	c.131G>A	COSM131596	FFPE
42	Sample066	23772	3504	14.74	p.G44D	c.131G>A	COSM131596	FFPE
43	Sample067	207170	35934	17.35	p.G44D	c.131G>A	COSM131596	FFPE
44	Sample068	21620	1274	5.89	p.G44D	c.131G>A	COSM131596	FFPE
45	Sample069	26076	2810	10.78	p.G44D	c.131G>A	COSM131596	FFPE
46	Sample070	42560	4516	10.61	p.G44S	c.130G>A	COSM131594	FFPE
47	Sample074	54226	7328	13.51	p.G44S	c.130G>A	COSM131594	FFPE
48	Sample077	268636	62138	23.13	p.G44S	c.130G>A	COSM131594	FFPE
49	Sample078	42170	5700	13.5	p.G44S	c.130G>A	COSM131594	FFPE
50	Sample080	29112	2618	8	p.G44S	c.130G>A	COSM131594	FFPE
51	Sample085	60604	4474	7.38	p.G44S	c.130G>A	COSM131594	FFPE
52	Sample086	81188	6800	8.38	p.G44S	c.130G>A	COSM131594	FFPE
53	Sample087	101244	7140	7.05	p.N40_F45	c.118_135 del18		FFPE
54	Sample090	100526	114548	11.39	p.G44C	c.130G>T	COSM131593	FFPE
55	Sample091	43602	8344	19.14	p.L36P	c.107T>C		FFPE

Table 6 Mutations detected in ultra-deep targeted amplicon sequencing of *MED12* exon 2 in 98 FA samples.

No .	Sample ID	Total Reads	Variant Reads	Variant allele Frequency (%)	Amino acid change	cDNA change	COSMIC	Tissue type
56	Sample09_2	25034	3034	12.12	p.E33_D34 insPQ	Exon2 (-8 T>A)	COSM1316_18	FFPE
57	Sample09_5	42452	7530	17.74	p.E33_D34 insPQ	Exon2 (-8 T>A)	COSM1316_18	FFPE
58	Sample09_6	89982	19106	21.23	p.G44D	c.131G>A	COSM1315_96	FFPE
-	288_PC3	49790	1730	3.59	p.G44S	c.130G>A	COSM1315_94	Spike-in Control
-	287_PC5	222990	13576	6.1	p.G44S	c.130G>A	COSM1315_94	Spike-in Control
-	286_PC1_0	167234	18412	11.01	p.G44S	c.130G>A	COSM1315_94	Spike-in Control
-	285_PC1_5	185356	28162	15.2	p.G44S	c.130G>A	COSM1315_94	Spike-in Control

Example 5

The *MED12* gene lies on chromosome X, and in females, one copy is normally silenced by epigenetic inactivation¹⁷. To confirm that mutant *MED12* transcripts are expressed, and are not suppressed by X-inactivation, the present disclosure performed Sanger sequencing on complementary DNA (cDNA) generated by reverse-transcribing messenger RNA (mRNA) from eight fresh frozen samples that were determined to harbor *MED12* exon 2 mutations by targeted amplicon sequencing. Particularly, the present disclosure sequenced the cDNA of seven *MED12*-mutant samples with available fresh frozen tissue. The present disclosure converted 100 ng of RNA to cDNA with SuperScript III First-Strand Synthesis SuperMix from Invitrogen according to manufacturer's recommended protocol. The present disclosure performed PCR and sequenced the *MED12* region between exon 1 and 3 with primers from Mäkinen et al¹³; forward primer: CTTCCGGATCTTGAGCTACG, reverse primer: GATCTTGGCAGGATTGAAGC, product length: 199 bp. PCR amplification, sequencing and fractionation was performed as described above for Sanger sequencing of genomic DNA.

The present disclosure were able to unambiguously identify the correct *MED12* mutations in the cDNA of all but one sample, as illustrated in Fig.3, indicating that mutant *MED12* is indeed transcribed.

5 Example 6

Due to the biphasic nature of FA and relatively low variant allele frequencies observed in *MED12* mutations (14.1%), it was suspected that *MED12* mutations may be present in either the epithelial or stromal compartments. To confirm this, the present disclosure performed LCM (laser capture microdissection) on one sample (Sample006) and Sanger sequenced the
10 individual compartments.

Briefly, fresh frozen tissue from Sample006 was embedded in Optimal Cutting Temperature (OCT) compound (Tissue-Tek, Sakura Finetek), and sections (8µm thick) were cut in a Microtome-cryostat (Leica), mounted onto Arcturus® PEN membrane glass slides (Life
15 Technologies), and then stored at -80 °C till required. Slides were dehydrated & stained with Arcturus® Histogene® following manufacturer's recommendations. The stained slide was loaded onto the laser capture microscope stage (ArcturusXT™ Laser Capture Microdissection (LCM) System). A Capsure™ Macro LCM cap (Life Technologies) was then placed automatically over the chosen area of the tissue. Once the cells of interest that
20 were highlighted by the software were verified by the user, the machine automatically dissected out the highlighted cells of interest using a near infrared laser or UV pulse that transferred them onto the Capsure™ Macro LCM Cap.

The DNA was extracted directly from LCM caps using Qiagen FFPE DNA Tissue kit
25 following manufacturer's protocol with the following modifications. Each sample cap was incubated with the lysis buffer (ATL & Proteinase K) in a 500 µl microcentrifuge at 60 °C for 5 hrs & enzyme deactivation at 90 °C for 10 minutes. The eluted DNA was used directly for PCR & BigDye® sequencing.

30 Results show that *MED12* mutations are only found in the stromal compartment, and that epithelial portions of the FA tumor contained only wild-type as in Fig.4. Frequent *MED12* exon 2 somatic mutations have hitherto been found only in uterine leiomyoma (UL)¹³. The

point mutations found in FA are remarkably similar to that of UL both in location and variant codon preference as indicated in Table 5. Both tumors are dominated by frequent codon 44 missense mutations (42% in FA and 49% in UL, $p = 0.28$, two-tailed Fisher's exact test). Codon 36 missense mutations were the second-most frequent in both tumors and occurred at similar frequencies (4.1% in FA and 5% in UL, $p = 1.00$, two-tailed Fisher's exact test). The present disclosure also observed codon 43 mutations and intronic T>A aberrant splice acceptor site mutation previously observed in UL. Altogether, every single point mutation in MED12 exon 2 detected in UL was also detected in FA. Additionally, both tumors also share a preference for in-frame deletions. These observations suggest that FAs and ULs may have a common underlying genetic basis.

Example 7

Total RNA was extracted from 10 fresh frozen fibroadenoma tumors using Trizol (Invitrogen) and purified using the RNeasy mini kit (Qiagen). 10 μ g of purified total RNA was then labelled according to standard Affymetrix protocol and then hybridized to Affymetrix GeneChip Human Genome U133 Plus 2.0 microarrays. Scanning of the microarrays was performed using the Affymetrix GeneChip Scanner 7G. CEL files were loaded into the R statistical environment (version 2.15.2) using the *simpleaffy* package³¹ and preprocessed using the robust multi-array average (RMA) algorithm³² with quantile normalization. Mapping of Affymetrix probe sets to genes was performed using the BrainArray custom CDF³³ (chip definition file) version 17. Differentially expressed genes between mutant *MED12* and wild-type *MED12* samples were identified based on empirical Bayes moderated t-statistics calculated using the *limma* package³⁴. A list of genes differentially expressed over 1.5 fold in either direction and with a p-value less than 0.05 is presented in Table 7. P-values were not significant after adjusting for multiple hypotheses due to the limited sample size. The microarray data has been deposited in the Gene Expression Omnibus³⁵ (GEO accession ID: GSE55594).

Table 7 Differentially expressed genes between mutant and wild-type *MED12* fibroadenoma samples.

Gene Symbol	log2 fold-change	t-statistic	p-value
MMP13	3.823	3.113	0.010

Table 7 Differentially expressed genes between mutant and wild-type *MED12* fibroadenoma samples.

Gene Symbol	log2 fold-change	t-statistic	p-value
TAT	2.764	4.569	0.001
RFX6	2.034	3.469	0.006
ERP27	1.936	3.693	0.004
CYP4X1	1.880	2.341	0.040
SUSD5	1.874	2.560	0.027
IL13RA2	1.826	2.838	0.017
C12orf69	1.675	2.371	0.038
KCNK15	1.607	3.345	0.007
CPA3	1.388	2.624	0.024
SOWAHA	1.359	2.587	0.026
ENTPD1	1.347	2.323	0.041
ADRA2A	1.334	3.579	0.005
C1orf64	1.266	3.642	0.004
FSIP1	1.246	2.278	0.045
REEP1	1.242	2.561	0.027
RHOH	1.238	2.397	0.036
TTC39A	1.232	3.907	0.003
RERGL	1.015	3.002	0.013
TUBB2B	1.008	2.736	0.020
ITGA8	1.007	2.634	0.024
FAM70A	1.004	2.421	0.035
SLC19A2	0.967	3.469	0.006
LTBP2	0.957	2.344	0.040
HEPH	0.925	2.549	0.028
SYTL4	0.910	3.022	0.012
NRIP3	0.908	2.398	0.036
ZNF552	0.895	2.736	0.020
PREX1	0.883	3.389	0.006
TTC36	0.865	2.739	0.020
MLPH	0.863	3.054	0.012
AZGP1	0.861	2.634	0.024
LOC100507165	0.854	2.882	0.016
TUBB2A	0.827	2.635	0.024
GEM	0.826	3.464	0.006
ECM2	0.824	3.441	0.006
TSPAN2	0.811	2.321	0.042
HOMER1	0.810	2.809	0.018
C11orf96	0.805	2.951	0.014
CASC1	0.791	5.394	0.000

Table 7 Differentially expressed genes between mutant and wild-type *MED12* fibroadenoma samples.

Gene Symbol	log2 fold-change	t-statistic	p-value
FOXA1	0.776	2.481	0.031
CSRP2	0.762	2.289	0.044
KIAA1467	0.751	3.039	0.012
TSPAN7	0.748	3.837	0.003
LOC729970	0.739	5.084	0.000
ACP5	0.719	2.632	0.024
RNF175	0.710	2.606	0.025
LYPD6	0.708	2.926	0.014
FGFR1OP	0.702	2.763	0.019
MUC1	0.678	2.906	0.015
C10orf116	0.675	2.704	0.021
GPR160	0.663	2.387	0.037
FRK	0.657	2.273	0.045
FJX1	0.656	2.255	0.047
ECI2	0.645	3.662	0.004
STK17A	0.642	3.037	0.012
SNX10	0.636	2.965	0.013
TPBG	0.631	2.939	0.014
CDC42EP3	0.624	3.780	0.003
C7orf10	0.613	2.583	0.026
RAB38	0.608	2.268	0.046
MYRIP	0.608	2.675	0.022
WWP1	0.603	2.413	0.035
HS3ST1	0.594	2.337	0.040
SLC25A16	0.592	2.741	0.020
LOC100506100	0.586	3.252	0.008
MFSD4	-0.586	-2.410	0.036
PIK3R1	-0.586	-2.563	0.027
PPAP2B	-0.598	-2.482	0.031
EMILIN3	-0.601	-2.455	0.033
HLF	-0.605	-3.216	0.009
ARHGAP26	-0.612	-2.611	0.025
IGSF5	-0.613	-2.310	0.042
HOXA13	-0.620	-2.603	0.025
PRKX	-0.626	-2.635	0.024
PPP3CA	-0.627	-2.645	0.024
DDR2	-0.638	-2.400	0.036
FCRLB	-0.640	-3.497	0.005
UNC80	-0.655	-2.471	0.032

Table 7 Differentially expressed genes between mutant and wild-type *MED12* fibroadenoma samples.

Gene Symbol	log2 fold-change	t-statistic	p-value
RHOV	-0.657	-2.554	0.028
ZFHX4	-0.664	-2.437	0.034
CD44	-0.700	-2.860	0.016
FUT9	-0.700	-2.256	0.046
TPTE2P6	-0.703	-2.450	0.033
CXCR4	-0.705	-2.345	0.040
EYA4	-0.723	-2.551	0.028
HSD17B1	-0.730	-2.409	0.036
FAM19A5	-0.745	-2.928	0.014
C1orf51	-0.757	-2.361	0.039
ITIH5	-0.761	-2.684	0.022
FZD7	-0.775	-3.074	0.011
CPE	-0.796	-2.772	0.019
OOEP	-0.844	-3.102	0.011
FAM13A	-0.860	-3.198	0.009
MNX1	-0.863	-4.462	0.001
EMR2	-0.867	-4.237	0.002
ITGA6	-0.878	-2.298	0.043
MFAP3L	-0.881	-2.450	0.033
MAGEL2	-0.883	-2.400	0.036
EBF3	-0.890	-2.720	0.021
ADAMTSL3	-0.933	-2.340	0.040
LOC158434	-0.980	-2.483	0.031
ST8SIA1	-1.003	-2.898	0.015
PDE9A	-1.051	-2.601	0.026
UG0898H09	-1.051	-2.624	0.024
SLIT2	-1.179	-2.460	0.033
SLC12A2	-1.222	-2.555	0.028
CXCL1	-1.278	-2.222	0.049
RYR3	-1.358	-2.419	0.035
FGF10	-1.380	-2.932	0.014
DDX43	-1.384	-4.831	0.001
MFAP5	-1.391	-2.294	0.044
NRK	-1.731	-4.317	0.001
SOX8	-1.741	-2.500	0.030
PTH2R	-1.762	-2.346	0.040
GPC3	-1.958	-3.516	0.005
ZFPM2	-1.962	-2.284	0.044
LOC100652994	-1.994	-2.425	0.035

Table 7 Differentially expressed genes between mutant and wild-type *MED12* fibroadenoma samples.

Gene Symbol	log2 fold-change	t-statistic	p-value
LTF	-1.996	-2.588	0.026

Example 8

To characterize transcriptional changes associated with aberrant *MED12*, the present disclosure generated and compared the gene expression profiles of six *MED12* mutated fibroadenoma samples against four *MED12* wild-type fibroadenomas. Due to the limited sample size and fibroepithelial nature of fibroadenomas, the present disclosure used GSEA¹⁹ (Gene Set Enrichment Analysis) in order to identify potentially dysregulated pathways. Genes were rank-ordered by fold-change between *MED12*-mutant and wild-type fibroadenomas and subjected to GSEA against MSigDB¹⁹ (Molecular Signatures Database) curated (c2) gene sets. Particularly, the present disclosure integrated our gene expression data with publicly-available gene expression data of UL tumors (GEO accession ID: GSE30673). Lists of genes upregulated two-fold and four-fold were obtained by calculating fold-change of averages between mutant *MED12* (n = 8) and wild-type UL samples (n = 2). To calculate if the overlap between genes upregulated in *MED12*-mutant fibroadenoma and *MED12*-mutant UL is significant, the present disclosure used the Gene Set Enrichment Analysis (GSEA) tool¹⁹. Briefly, genes in the fibroadenoma dataset were ranked-ordered according to log fold-change. The GSEA algorithm then examines where genes upregulated in UL fall in the rank-ordered list, and generates an enrichment score corresponding to how enriched a gene set is in either extreme end of the rank-ordered list as can be seen in Fig. 5b. Random, size-matched gene sets are then used to generate an empirical p-value. Similarly, GSEA analysis was also performed on our fibroadenoma microarray dataset against the MSigDB c2 (curated) gene sets¹⁹, which are derived from publications, canonical pathways and expert knowledge.

25

A list of candidate mutant *MED12* target genes was obtained from the core-enriched genes (Fig. 6) in the GSEA analysis of our FA microarray data against upregulated genes in the UL dataset. Core-enriched genes are defined as those in the leading edge subset of the gene set (i.e. those that contributed most to the enrichment score). These genes were then used as

input in the MSigDB web site ‘Compute Overlaps’ tool (accessed on 10th February 2014, see URLs). Two classes of gene sets were used in the analysis; c2 (curated) and c5 (gene ontology). Given a gene list, the tool uses the hypergeometric test to compare it against gene sets to determine if the overlap exceeds chance. Gene sets with FDR³³ (false discovery rate) q-values < 0.05 were considered to be significantly over-represented with members of the input gene list.

In order to study relative pathway activity on the level of individual samples, the present disclosure used the Gene Set Variation Analysis (GSVA) method³⁶. Using a non-parametric approach, GSVA transforms a gene by sample matrix into a gene set by sample matrix, facilitating the identification of differential activation of functionally related genes. Empirical Bayes moderated t-statistics³⁴ were then calculated and gene sets with p-values < 0.05 were considered to have significantly differential activity between mutant *MED12* and wild-type *MED12* samples. GSVA was performed on two groups of gene sets. MSigDB c2 gene sets associated with breast cancer and estrogen signalling were considered, as shown in Fig.5b. Unsupervised clustering of samples and gene sets in heatmaps was performed using the *gplots* package in R using a Euclidean distance metric and complete-linkage clustering.

Among others, genes upregulated in *MED12*-mutant fibroadenomas are associated with ER+ breast cancers, estrogen stimulus in ER+ breast cancer cells, extracellular matrix (ECM) regulation and TGF β signalling as revealed in Table 8 and 9. As the top GSEA results suggested an association between *MED12* mutations and activated estrogen signalling, the present disclosure performed GSVA (Gene Set Variation Analysis) on our microarrays to detect differential pathway activity between samples.

Table 8. Top 50 enriched MSigDB curated (c2) gene sets for genes upregulated in *MED12* mutant FA. Gene sets of interest are highlighted. ES: Enrichment Score, NES: Normalized Enrichment Score, FDR: False Discovery Rate

NAME	SIZE	ES	NES	FDR
DOANE_BREAST_CANCER_ESR1_UP	103	0.801	2.826	0.000
SMID_BREAST_CANCER_RELAPSE_IN_BRAIN_DN	70	0.754	2.531	0.000
SMID_BREAST_CANCER_RELAPSE_IN_BONE_UP	86	0.724	2.526	0.000
SMID_BREAST_CANCER_BASAL_DN	597	0.579	2.508	0.000
LIEN_BREAST_CARCINOMA_METAPLASTIC_VS_DUC	94	0.705	2.472	0.000

TAL_DN				
SMID_BREAST_CANCER_LUMINAL_B_UP	153	0.638	2.419	0.000
YANG_BREAST_CANCER_ESR1_UP	33	0.821	2.413	0.000
VANTVEER_BREAST_CANCER_ESR1_UP	132	0.642	2.373	0.000
NAGASHIMA_EGF_SIGNALING_UP	50	0.738	2.310	0.000
MASSARWEH_RESPONSE_TO ESTRADIOL	51	0.697	2.196	0.000
NAGASHIMA_NRG1_SIGNALING_UP	157	0.579	2.185	0.000
REACTOME_DEGRADATION_OF_THE_EXTRACELLULAR_MATRIX	26	0.765	2.140	0.000
CHIBA_RESPONSE_TO_TSA_UP	49	0.685	2.135	0.000
POOLA_INVASIVE_BREAST_CANCER_DN	123	0.580	2.124	0.001
CHARAFE_BREAST_CANCER_LUMINAL_VS_BASAL_UP	314	0.509	2.088	0.003
DORN_ADENOVIRUS_INFECTION_48HR_DN	34	0.698	2.075	0.003
COWLING_MYCN_TARGETS	36	0.692	2.074	0.003
PID_UPA_UPAR_PATHWAY	38	0.699	2.072	0.003
AMIT_SERUM_RESPONSE_60_MCF10A	53	0.643	2.061	0.003
VANTVEER_BREAST_CANCER_METASTASIS_UP	43	0.660	2.039	0.005
WANG_TNF_TARGETS	23	0.747	2.017	0.007
AMIT_EGF_RESPONSE_40_HELA	38	0.672	2.003	0.009
PLASARI_TGFB1_TARGETS_1HR_UP	30	0.693	2.002	0.009
LIM_MAMMARY_LUMINAL_MATURE_UP	106	0.563	2.000	0.009
DORN_ADENOVIRUS_INFECTION_32HR_DN	33	0.670	1.992	0.010
FARMER_BREAST_CANCER_BASAL_VS_LUMINAL	291	0.487	1.985	0.011
SMID_BREAST_CANCER_LUMINAL_A_UP	81	0.578	1.981	0.011
MASSARWEH_TAMOXIFEN_RESISTANCE_DN	201	0.507	1.977	0.011
NIELSEN_LEIOMYOSARCOMA_CNN1_UP	18	0.765	1.971	0.013
AMIT_EGF_RESPONSE_40_MCF10A	18	0.768	1.954	0.016
FRASOR_RESPONSE_TO ESTRADIOL_UP	35	0.654	1.952	0.016
REACTOME_EXTRACELLULAR_MATRIX_ORGANIZATION	83	0.554	1.950	0.016
YANG_BREAST_CANCER_ESR1_BULK_UP	19	0.741	1.948	0.016
WATTEL_AUTONOMOUS_THYROID_ADENOMA_DN	50	0.611	1.948	0.016
PHONG_TNF_TARGETS_UP	60	0.589	1.948	0.015
SU_THYMUS	19	0.755	1.946	0.015
DUTERTRE ESTRADIOL_RESPONSE_24HR_UP	290	0.474	1.941	0.016
WILSON_PROTEASES_AT_TUMOR_BONE_INTERFACE_UP	21	0.731	1.933	0.018
ROSTY_CERVICAL_CANCER_PROLIFERATION_CLUSTER	127	0.521	1.932	0.017
PLASARI_TGFB1_TARGETS_10HR_UP	182	0.503	1.932	0.017
MCMURRAY_TP53_HRAS_COOPERATION_RESPONSE_DN	61	0.586	1.920	0.020
UZONYI_RESPONSE_TO_LEUKOTRIENE_AND_THRO	34	0.654	1.915	0.021

MBIN				
DIRMEIER_LMP1_RESPONSE_EARLY	61	0.591	1.911	0.022
DAZARD_UV_RESPONSE_CLUSTER_G4	17	0.746	1.909	0.022
JAZAERI_BREAST_CANCER_BRCA1_VS_BRCA2_DN	37	0.641	1.904	0.023
TIAN_TNF_SIGNALING_NOT_VIA_NFKB	21	0.716	1.897	0.025
CREIGHTON_ENDOCRINE_THERAPY_RESISTANCE_4	245	0.468	1.893	0.026
WANG_RESPONSE_TO_FORSKOLIN_UP	21	0.696	1.872	0.033
TRAYNOR_RETT_SYNDROM_UP	41	0.608	1.869	0.034
KORKOLA_TERATOMA	34	0.635	1.869	0.033

Table 9. Top 50 enriched MSigDB curated (c2) gene sets for genes downregulated in *MED12* mutant FA. Gene sets of interest are highlighted. ES: Enrichment Score, NES: Normalized Enrichment Score, FDR: False Discovery Rate

NAME	SIZE	ES	NES	FDR
LIM_MAMMARY_LUMINAL_PROGENITOR_UP	53	-0.718	-2.468	0.000
SMID_BREAST_CANCER_RELAPSE_IN_BONE_DN	283	-0.517	-2.293	0.001
DOANE_BREAST_CANCER_ESR1_DN	46	-0.685	-2.239	0.001
SMID_BREAST_CANCER_LUMINAL_B_DN	500	-0.459	-2.113	0.010
SMID_BREAST_CANCER_BASAL_UP	581	-0.452	-2.093	0.014
ONDER_CDH1_TARGETS_3_DN	50	-0.611	-2.052	0.023
YANG_BREAST_CANCER_ESR1_DN	24	-0.696	-1.988	0.048
REACTOME_LATENT_INFECTION_OF_HOMO_SAPIENS_WITH_MYCOBACTERIUM_TUBERCULOSIS	30	-0.654	-1.948	0.074
CHIBA_RESPONSE_TO_TSA_DN	21	-0.699	-1.917	0.101
KEGG_GLYCOPHINGOLIPID_BIOSYNTHESIS_LACTO_AND_NEOLACTO_SERIES	26	-0.649	-1.891	0.126
KEGG_LONG_TERM_POTENTIATION	64	-0.532	-1.880	0.132
YANG_BREAST_CANCER_ESR1_BULK_DN	19	-0.679	-1.867	0.142
CHIARADONNA_NEOPLASTIC_TRANSFORMATION_CDC25_UP	110	-0.481	-1.842	0.177
BIOCARTA_CXCR4_PATHWAY	22	-0.654	-1.838	0.172
PID_A6B1_A6B4_INTEGRIN_PATHWAY	44	-0.565	-1.837	0.162
BIOCARTA_IL7_PATHWAY	17	-0.699	-1.816	0.194
LEE_LIVER_CANCER_DENA_UP	58	-0.517	-1.815	0.184
ROY_WOUND_BLOOD_VESSEL_DN	20	-0.649	-1.815	0.175
LIM_MAMMARY_LUMINAL_MATURE_DN	89	-0.481	-1.799	0.199
REACTOME_INTERACTION_BETWEEN_L1_AND_ANKYRINS	20	-0.654	-1.790	0.207
REACTOME_ACTIVATED_POINT_MUTANTS_OF_FGF_R2	16	-0.666	-1.787	0.203
KEGG_AXON_GUIDANCE	124	-0.453	-1.786	0.196
NAKAYAMA_SOFT_TISSUE_TUMORS_PCA2_DN	75	-0.486	-1.773	0.215
KEGG_ALZHEIMERS_DISEASE	142	-0.440	-1.762	0.230
REACTOME_TRAFFICKING_OF_AMPA_RECEPTORS	26	-0.601	-1.753	0.241

PID_NCADHERINPATHWAY	30	-0.585	-1.751	0.238
CHEN_LVAD_SUPPORT_OF_FAILING_HEART_UP	93	-0.465	-1.750	0.231
KEGG_SMALL_CELL_LUNG_CANCER	79	-0.481	-1.746	0.231
NIELSEN_SCHWANNOMA_UP	15	-0.683	-1.744	0.228
SUZUKI_RESPONSE_TO_TSA_AND_DECITABINE_1A	19	-0.629	-1.740	0.228
CHEMELLO_SOLEUS_VS_EDL_MYOFIBERS_DN	19	-0.637	-1.737	0.229
JOHNSTONE_PARVB_TARGETS_1_DN	43	-0.533	-1.731	0.235
PID_INTEGRIN1_PATHWAY	63	-0.489	-1.731	0.228
TIEN_INTESTINE_PROBIOTICS_6HR_UP	39	-0.542	-1.726	0.232
REACTOME_SIGNALING_BY_INSULIN_RECEPTOR	98	-0.446	-1.724	0.229
REACTOME_NEPHRIN_INTERACTIONS	19	-0.641	-1.723	0.225
GUILLAUMOND_KLF10_TARGETS_DN	24	-0.608	-1.718	0.231
JAEGER_METASTASIS_DN	234	-0.399	-1.716	0.230
REACTOME_UNBLOCKING_OF_NMDA_RECEPTOR_G LUTAMATE_BINDING_AND_ACTIVATION	15	-0.685	-1.708	0.242
MAHADEVAN_RESPONSE_TO_MP470_UP	19	-0.606	-1.699	0.255
REACTOME_FGFR_LIGAND_BINDING_AND_ACTIVA TION	22	-0.610	-1.694	0.260
LEE_LIVER_CANCER_E2F1_UP	57	-0.489	-1.693	0.256
CHIARADONNA_NEOPLASTIC_TRANSFORMATION_ KRAS_UP	112	-0.444	-1.693	0.251
ONDER_CDH1_TARGETS_1_DN	146	-0.423	-1.691	0.248
VANTVEER_BREAST_CANCER_ESR1_DN	206	-0.408	-1.691	0.243
REACTOME_PI3K_CASCADE	62	-0.487	-1.686	0.249
PID_IL2_STAT5PATHWAY	30	-0.567	-1.685	0.247
KEGG_AMYOTROPHIC_LATERAL_SCLEROSIS_ALS	47	-0.503	-1.683	0.245
YANG_MUC2_TARGETS_DUODENUM_6MO_DN	19	-0.609	-1.677	0.255
KEGG_GLIOMA	63	-0.477	-1.676	0.252

Given the similarity of the *MED12* mutation spectrum in FAs and ULs, the present disclosure hypothesized the integration of FA and UL molecular data might allow further pinpointing of genes and pathways. Indeed, GSEA on our FA dataset against a previously-published set of genes upregulated in *MED12*-mutated ULs revealed a strong similarity of upregulated genes in *MED12*-mutated FAs and ULs. Specifically, genes upregulated in *MED12*-mutant FA samples were significantly enriched for genes upregulated over two-fold in *MED12*-mutant UL (enrichment score =0.61, p=0) as shown in Fig.2, with enrichment becoming even more profound when only genes upregulated four-fold were considered (enrichment score = 0.81, p = 0), with reference to Fig. 5b. Analysis of core-enriched genes (i.e. genes commonly upregulated in both FA and UL with mutant *MED12*) revealed, as in Table 10 below, that

they were over-represented with genes associated with extracellular matrix (ECM) organization, estrogen signalling, as well as TGF β and Wnt signalling.

Table 10. MSigDB curated (c2) gene sets significantly overlapping with candidate

Gene Set Name	Gene Set	# Genes in	FDR q-
RIGGI_EWING_SARCOMA_PROGENITOR_UP	430	17	0.00E+00
VECCHI_GASTRIC_CANCER_ADVANCED_VS_EARLY	175	9	2.20E-08
SCHUETZ_BREAST_CANCER_DUCTAL_INVASIVE_U	351	11	2.20E-08
TURASHVILI_BREAST_LOBULAR_CARCINOMA_VS_	74	7	4.50E-08
WONG_ADULT_TISSUE_STEM_MODULE	721	13	1.54E-07
BENPORATH_SUZ12_TARGETS	1038	14	1.01E-06
BENPORATH_EED_TARGETS	1062	14	1.15E-06
CHANDRAN_METASTASIS_DN	306	9	1.15E-06
BENPORATH_ES_WITH_H3K27ME3	1118	14	1.74E-06
YANG_BCL3_TARGETS_UP	364	9	4.16E-06
MARTORIATI_MDM4_TARGETS_NEUROEPITHELIUM	164	7	4.50E-06
CHIANG_LIVER_CANCER_SUBCLASS_CTNNB1_DN	170	7	5.29E-06
MCBRYAN_PUBERTAL_BREAST_4_5WK_UP	271	8	5.74E-06
SMID_BREAST_CANCER_BASAL_DN	701	11	6.56E-06
POOLA_INVASIVE_BREAST_CANCER_UP	288	8	7.97E-06
BOQUEST_STEM_CELL_CULTURED_VS_FRESH_UP	425	9	9.83E-06
PLASARI_TGFB1_TARGETS_10HR_UP	199	7	1.10E-05
GOZGIT_ESR1_TARGETS_DN	781	11	1.52E-05
CROMER_TUMORIGENESIS_UP	63	5	1.64E-05
TURASHVILI_BREAST_LOBULAR_CARCINOMA_VS_	69	5	2.39E-05
ZWANG_TRANSIENTLY_UP_BY_2ND_EGF_PULSE_O	1725	15	2.39E-05
AFFAR_YY1_TARGETS_DN	234	7	2.57E-05
SABATES_COLORECTAL_ADENOMA_UP	141	6	2.60E-05
SENESE_HDAC3_TARGETS_UP	501	9	2.65E-05
REACTOME_DEGRADATION_OF_THE_EXTRACELLU	29	4	2.81E-05
LEE_NEURAL_CREST_STEM_CELL_UP	146	6	2.81E-05
MEISSNER_BRAIN_HCP_WITH_H3K4ME3_AND_H3K2	1069	12	2.81E-05
MCLACHLAN_DENTAL_CARIES_UP	253	7	3.43E-05
MIYAGAWA_TARGETS_OF_EWSR1_ETS_FUSIONS_U	259	7	3.88E-05
SANA_TNF_SIGNALING_UP	83	5	4.18E-05
HAN_SATB1_TARGETS_UP	395	8	4.27E-05
REACTOME_SIGNALING_BY_GPCR	920	11	4.27E-05
SERVITJA_ISLET_HNF1A_TARGETS_UP	163	6	4.27E-05
GHANDHI_BYSTANDER_IRRADIATION_UP	86	5	4.41E-05
BENPORATH_SOX2_TARGETS	734	10	4.47E-05
REACTOME_GPCR_LIGAND_BINDING	408	8	4.75E-05
CHICAS_RB1_TARGETS_CONFLUENT	567	9	4.84E-05
WANG_SMARCE1_TARGETS_UP	280	7	5.01E-05
NUYTTEN_NIPPI_TARGETS_UP	769	11	6.12E-05
SCHAEFFER_PROSTATE_DEVELOPMENT_48HR_DN	428	8	6.12E-05
BROWNE_HCMV_INFECTION_48HR_UP	180	6	6.15E-05
GAUSSMANN_MLL_AF4_FUSION_TARGETS_E_UP	97	5	6.52E-05
KANG_IMMORTALIZED_BY_TERT_DN	102	5	8.17E-05
COWLING_MYCN_TARGETS	43	4	8.26E-05
JAEGER_METASTASIS_UP	44	4	8.87E-05
NUYTTEN_EZH2_TARGETS_UP	1037	11	9.87E-05

Table 10. MSigDB curated (c2) gene sets significantly overlapping with candidate

Gene Set Name	Gene Set	# Genes in	FDR q-
REACTOME_GASTRIN_CREB_SIGNALLING_PATHWA	205	6	1.15E-04
BENPORATH_PRC2_TARGETS	652	9	1.16E-04
MCCLUNG_DELTA_FOSB_TARGETS_2WK	48	4	1.16E-04
RIZKI_TUMOR_INVASIVENESS_3D_UP	210	6	1.24E-04
RODWELL_AGING_KIDNEY_UP	487	8	1.26E-04
MCBRYAN_PUBERTAL_BREAST_3_4WK_UP	214	6	1.33E-04
RODWELL_AGING_KIDNEY_NO_BLOOD_UP	222	6	1.61E-04
YAMASHITA_METHYLATED_IN_PROSTATE_CANCE	57	4	2.12E-04
ZHOU_INFLAMMATORY_RESPONSE_FIMA_UP	544	8	2.65E-04
ANASTASSIOU_CANCER_MESENCHYMAL_TRANSITI	64	5	3.25E-04
DOUGLAS_BMI1_TARGETS_UP	566	8	3.41E-04
KUMAR_TARGETS_OF_MLL_AF9_FUSION	405	7	3.78E-04
CUI_TCF21_TARGETS_2_UP	428	7	5.33E-04
CORRE_MULTIPLE_MYELOMA_UP	74	4	5.34E-04
NAKAYAMA_SOFT_TISSUE_TUMORS_PCA1_DN	74	4	5.34E-04
WANG_MLL_TARGETS	289	6	6.18E-04
DELYS_THYROID_CANCER_UP	443	7	6.18E-04
BENPORATH_OCT4_TARGETS	290	6	6.18E-04
SABATES_COLORECTAL_ADENOMA_DN	291	6	6.21E-04
WONG_ENDMETRIUM_CANCER_DN	82	4	7.43E-04
SMID_BREAST_CANCER_BASAL_UP	648	8	7.76E-04
ONDER_CDH1_TARGETS_2_DN	464	7	7.80E-04
KEGG_ECM_RECEPTOR_INTERACTION	84	4	7.82E-04
KEGG_TGF_BETA_SIGNALING_PATHWAY	86	4	8.47E-04
REACTOME_EXTRACELLULAR_MATRIX_ORGANIZA	87	4	8.54E-04
SASSON_RESPONSE_TO_GONADOTROPHINS_DN	87	4	8.54E-04
SMID_BREAST_CANCER_RELAPSE_IN_BONE_DN	315	6	8.54E-04
REACTOME_G_ALPHA_Q_SIGNALLING_EVENTS	184	5	8.54E-04
BYSTRYKH_HEMATOPOIESIS_STEM_CELL_QTL_TR	882	9	8.54E-04
SASSON_RESPONSE_TO_FORSKOLIN_DN	88	4	8.54E-04
ABE_VEGFA_TARGETS_30MIN	29	3	9.07E-04
SCHAEFFER_PROSTATE_DEVELOPMENT_48HR_UP	487	7	9.28E-04
RIGGI_EWING_SARCOMA_PROGENITOR_DN	191	5	9.59E-04
STAEGE_EWING_FAMILY_TUMOR	33	3	1.30E-03
RICKMAN_HEAD_AND_NECK_CANCER_A	100	4	1.33E-03
PLASARI_TGFB1_TARGETS_1HR_UP	34	3	1.39E-03
PEREZ_TP63_TARGETS	355	6	1.48E-03
LI_CISPLATIN_RESISTANCE_DN	35	3	1.48E-03
WIERENGA_STAT5A_TARGETS_UP	217	5	1.64E-03
IZADPANAH_STEM_CELL_ADIPOSE_VS_BONE_DN	108	4	1.67E-03
WESTON_VEGFA_TARGETS	108	4	1.67E-03
CUI_TCF21_TARGETS_UP	37	3	1.67E-03
BLALOCK_ALZHEIMERS_DISEASE_DN	1237	10	1.73E-03
GHANDHI_DIRECT_IRRADIATION_UP	110	4	1.74E-03
LABBE_TARGETS_OF_TGFB1_AND_WNT3A_UP	111	4	1.78E-03
SMID_BREAST_CANCER_LUMINAL_B_DN	564	7	2.00E-03
SCHAEFFER_PROSTATE_DEVELOPMENT_12HR_UP	116	4	2.07E-03
CERVERA_SDHB_TARGETS_1_UP	118	4	2.17E-03
MARKEY_RB1_CHRONIC_LOF_DN	118	4	2.17E-03
PID_THROMBIN_PAR1_PATHWAY	43	3	2.42E-03
YOSHIMURA_MAPK8_TARGETS_UP	1305	10	2.46E-03

Table 10. MSigDB curated (c2) gene sets significantly overlapping with candidate

Gene Set Name	Gene Set	# Genes in	FDR q-
HOOI_ST7_TARGETS_DN	123	4	2.46E-03
PLASARI_TGFB1_TARGETS_10HR_DN	244	5	2.46E-03
MARTINEZ_RB1_AND_TP53_TARGETS_DN	591	7	2.47E-03

Moreover, previous study reported that one out of eight (12.5%) fibroadenomas exhibited a non-silent TP53 mutation¹⁰, whereas another study reported no somatic TP53 mutations in fibroadenomas from women who remained unaffected by breast cancer after an average follow-up of ten years¹¹. A single PIK3CA mutation has also been reported from a screen of ten fibroadenoma tumors¹². It is likely that those cases that harbour PIK3CA and TP53 mutations may actually indicate more aggressive phylloides tumors³⁷ (subtype of fibroepithelial tumors) rather than the true benign fibroadenomas. Therefore the presence of a single genetic alteration, in the absence of others such as P53 or PIK3CA, may be a more accurate biomarker for benign fibroadenoma. Therefore, early identification of such genetic attributes or methods allowing one to discover the likelihood of genetic deficiencies is greatly desired.

Candidate aberrant *MED12* target genes were also enriched for genes downregulated in liver cancer with activated beta-catenin (*CTNNB1*). As *MED12* plays a vital role in transducing Wnt/beta-catenin signaling²⁰, this observation is consistent with *MED12* mutations resulting in aberrant beta-catenin signalling, which is involved in regulating focal adhesion. Accordingly, GO (gene ontology) analysis showed that genes upregulated in mutant *MED12* samples were over-represented with those expressed in the extracellular region as shown in Table 11.

Table 11. MSigDB GO (c5) gene sets significantly overlapping with candidate mutant *MED12* target genes.

Gene Set Name	Gene Set Size (K)	# Genes in Overlap (k)	FDR q-value
EXTRACELLULAR_REGION_PART	338	13	1.79E-11
EXTRACELLULAR_REGION	447	13	3.12E-10
EXTRACELLULAR_SPACE	245	9	1.35E-07
MULTICELLULAR_ORGANISMAL_DEVELOPMENT	1049	13	5.28E-06
PROTEINACEOUS_EXTRACELLULAR_MATRIX	98	5	1.59E-04

EXTRACELLULAR_MATRIX	100	5	1.59E-04
ANATOMICAL_STRUCTURE_DEVELOPMENT	1013	11	1.59E-04
REGULATION_OF_BIOLOGICAL_QUALITY	419	7	1.05E-03
INTEGRAL_TO_MEMBRANE	1330	11	1.44E-03
SYSTEM_DEVELOPMENT	861	9	1.44E-03
INTRINSIC_TO_MEMBRANE	1348	11	1.44E-03
METALLOENDOPEPTIDASE_ACTIVITY	27	3	1.44E-03
CELL_FRACTION	493	7	1.85E-03
REGULATION_OF_SIGNAL_TRANSDUCTION	222	5	3.42E-03
ORGAN_DEVELOPMENT	571	7	4.09E-03
TRANSMEMBRANE_RECEPTOR_ACTIVITY	418	6	5.89E-03
METALLOPEPTIDASE_ACTIVITY	50	3	6.26E-03
MEMBRANE_PART	1670	11	6.26E-03
HYDROLASE_ACTIVITY_ACTING_ON_ESTER_BONDS	269	5	6.26E-03
MEMBRANE	1994	12	6.46E-03
SOLUBLE_FRACTION	161	4	1.01E-02
RESPONSE_TO_EXTERNAL_STIMULUS	312	5	1.08E-02
AXON	12	2	1.08E-02
INTEGRAL_TO_PLASMA_MEMBRANE	977	8	1.18E-02
INTRINSIC_TO_PLASMA_MEMBRANE	991	8	1.25E-02
PROTEOLYSIS	191	4	1.56E-02
HOMEOSTATIC_PROCESS	207	4	2.02E-02
RECEPTOR_ACTIVITY	583	6	2.02E-02
CATION_BINDING	213	4	2.11E-02
CELLULAR_PROTEIN_METABOLIC_PROCESS	1117	8	2.25E-02
MUSCLE_DEVELOPMENT	93	3	2.25E-02
CELLULAR_MACROMOLECULE_METABOLIC_PROCESS	1131	8	2.25E-02
PLASMA_MEMBRANE	1426	9	2.25E-02
CELL_MIGRATION	96	3	2.25E-02
NEURON_PROJECTION	21	2	2.25E-02
PLASMA_MEMBRANE_PART	1158	8	2.43E-02
CELL_SURFACE_RECEPTOR_LINKED_SIGNAL_TRANSDUCTION_GO_0007166	641	6	2.49E-02
REGULATION_OF_G_PROTEIN_COUPLED_RECEPTOR_PROTEIN_SIGNALING_PATHWAY	23	2	2.49E-02
CELLULAR_CATION_HOMEOSTASIS	106	3	2.66E-02
CATION_HOMEOSTASIS	109	3	2.81E-02

PROTEIN_METABOLIC_PROCESS	1231	8	3.17E-02
ENDOPEPTIDASE_ACTIVITY	117	3	3.29E-02
ION_BINDING	273	4	3.60E-02
ION_HOMEOSTASIS	129	3	4.16E-02
SIGNAL_TRANSDUCTION	1634	9	4.37E-02
RHODOPSIN_LIKE_RECEPTOR_ACTIVITY	134	3	4.44E-02
ENZYME_LINKED_RECEPTOR_PROTEIN_SIGNALING_PATHWAY	140	3	4.92E-02

It is crucial to note that the present disclosure is implementable to detect constitutional mutations regarding *MED12* gene mutation, more particularly mutations located in exon 2 of *MED12* gene, despite the subjects experimented in the foregoing examples may have obtained these mutations as somatic mutations. It is to be understood also that the present invention may be embodied in other specific forms and is not limited to the sole embodiment described above. However modification and equivalents of the disclosed concepts such as those which readily occur to one skilled in the art are intended to be included within the scope of the claims which are appended thereto

REFERENCES

1. Krishnamurthy, S., Ashfaq, R., Shin, H. J. C. & Sneige, N. Distinction of phyllodes tumor from fibroadenoma. *Cancer Cytopathol.* **90**, 342–349 (2000).
2. Fine, R. E. *et al.* Low-risk palpable breast masses removed using a vacuum-assisted handheld device. *Am. J. Surg.* **186**, 362–367 (2003).
3. Bernardes, J. R. M., Jr, Seixas, M. T., Lima, G. R., Marinho, L. C. & Gebrim, L. H. The effect of tamoxifen on PCNA expression in fibroadenomas. *Breast J.* **9**, 302–306 (2003).
4. Coriaty Nelson, Z., Ray, R. M., Gao, D. L. & Thomas, D. B. Risk factors for fibroadenoma in a cohort of female textile workers in Shanghai, China. *Am. J. Epidemiol.* **156**, 599–605 (2002).
5. Noguchi, S., Motomura, K., Inaji, H., Imaoka, S. & Koyama, H. Clonal Analysis of Fibroadenoma and Phyllodes Tumor of the Breast. *Cancer Res.* **53**, 4071–4074 (1993).
6. Dupont, W. D. *et al.* Long-term risk of breast cancer in women with fibroadenoma. *N. Engl. J. Med.* **331**, 10–15 (1994).
7. Liu, X. F. *et al.* A clinical study on the resection of breast fibroadenoma using two types of incision. *Scand. J. Surg. SJS Off. Organ Finn. Surg. Soc. Scand. Surg. Soc.* **100**, 147–152 (2011).

8. The Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature* **490**, 61–70 (2012).
9. Stephens, P. J. *et al.* The landscape of cancer genes and mutational processes in breast cancer. *Nature* **486**, 400–404 (2012).
- 5 10. Millikan, R. *et al.* p53 mutations in benign breast tissue. *J. Clin. Oncol.* **13**, 2293–2300 (1995).
11. Franco, N., Picard, S.-F., Mege, F., Arnould, L. & Lizard-Nacol, S. Absence of Genetic Abnormalities in Fibroadenomas of the Breast Determined at p53 Gene Mutations and Microsatellite Alterations. *Cancer Res.* **61**, 7955–7958 (2001).
12. Vorkas, P. A. *et al.* PIK3CA Hotspot Mutation Scanning by a Novel and Highly Sensitive High-Resolution Small Amplicon Melting Analysis Method. *J. Mol. Diagn. JMD* **12**, 697–704 (2010).
- 10 13. Vogelstein, B. *et al.* Cancer Genome Landscapes. *Science* **339**, 1546–1558 (2013).
14. Mäkinen, N. *et al.* MED12, the mediator complex subunit 12 gene, is mutated at high frequency in uterine leiomyomas. *Science* **334**, 252–255 (2011).
15. Forbes, S. A. *et al.* COSMIC: mining complete cancer genomes in the Catalogue of Somatic Mutations in Cancer. *Nucleic Acids Res.* **39**, D945–D950 (2011).
- 15 16. Sherry, S. T. *et al.* dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res.* **29**, 308–311 (2001).
17. Consortium, T. 1000 G. P. An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56–65 (2012).
- 20 18. Harper, P. S. Mary Lyon and the hypothesis of random X chromosome inactivation. *Hum. Genet.* **130**, 169–174 (2011).
19. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–15550 (2005).
- 25 20. Barbieri, C. E. *et al.* Exome sequencing identifies recurrent SPOP, FOXA1 and MED12 mutations in prostate cancer. *Nat. Genet.* **44**, 685–689 (2012).
21. Assié, G. *et al.* Integrated genomic characterization of adrenocortical carcinoma. *Nat. Genet.* (2014). doi:10.1038/ng.2953
22. Network, T. C. G. A. R. Integrated genomic analyses of ovarian carcinoma. *Nature* **474**, 609–615 (2011).
- 30 23. Je, E. M., Kim, M. R., Min, K. O., Yoo, N. J. & Lee, S. H. Mutational analysis of MED12 exon 2 in uterine leiomyoma and other common tumors. *Int. J. Cancer* **131**, E1044–E1047 (2012).
24. Kämpjärvi, K. *et al.* Somatic MED12 mutations in uterine leiomyosarcoma and colorectal cancer. *Br. J. Cancer* **107**, 1761–1765 (2012).

25. Zhu, B. T. & Conney, A. H. Functional role of estrogen metabolism in target cells: review and perspectives. *Carcinogenesis* **19**, 1–27 (1998).
26. Kang, Y. K., Guermah, M., Yuan, C.-X. & Roeder, R. G. The TRAP/Mediator coactivator complex interacts directly with estrogen receptors α and β through the TRAP220 subunit and directly enhances estrogen receptor function in vitro. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 2642–2647 (2002).
27. Mäkinen, N., Vahteristo, P., Bützow, R., Sjöberg, J. & Aaltonen, L. A. Exomic landscape of MED12 mutation-negative and -positive uterine leiomyomas. *Int. J. Cancer J. Int. Cancer* **134**, 1008–1012 (2014).
28. Chan-on, W. *et al.* Exome sequencing identifies distinct mutational patterns in liver fluke-related and non-infection-related bile duct cancers. *Nat. Genet.* **45**, 1474–1478 (2013).
29. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinforma. Oxf. Engl.* **25**, 1754–1760 (2009).
30. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* **25**, 2078–2079 (2009).
31. Wilson, C. L. & Miller, C. J. Simpleaffy: a BioConductor package for Affymetrix Quality Control and data analysis. *Bioinforma. Oxf. Engl.* **21**, 3683–3685 (2005).
32. Irizarry, R. A. *et al.* Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostat. Oxf. Engl.* **4**, 249–264 (2003).
33. Dai, M. *et al.* Evolving gene/transcript definitions significantly alter the interpretation of GeneChip data. *Nucleic Acids Res.* **33**, e175–e175 (2005).
34. Smyth, G. K. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat. Appl. Genet. Mol. Biol.* **3**, Article3 (2004).
35. Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res.* **30**, 207–210 (2002).
36. Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray and RNA-Seq data. *BMC Bioinformatics* **14**, 7 (2013).
37. Jardim, D.L.F., Coney, A., & Subbiah, V. Comprehensive characterization of malignant phyllodes tumor by whole genomic and proteomic analysis: biological implications for targeted therapy opportunities. *Orphanet Journal of Rare Diseases* **8**:112 (2013).

Claims

1. A method of assaying susceptibility and/or confirming diagnosis of breast
fibroadenomas development in a human subject comprising:
performing a nucleic acid-based assay to analyze an isolated polynucleotide encoding
at least exon 2 of *MED12* gene from a sample acquired from the human subject; and
regarding the human subject with greater susceptibility and/or confirming diagnosis of
breast fibroadenomas development by detecting a mutation in the isolated
polynucleotide, wherein the mutation is a splice site mutation located at position -8 of
exon 2 of the *MED12* gene, a missense mutation located at codon 44 of cDNA of the
MED12 gene or a missense mutation located at codon 36 of cDNA of the *MED12*
gene, wherein cDNA sequence of the *MED12* gene is set forth in SEQ ID No. 15.
2. The method of claim 1, wherein the missense mutation is located at position 107 of
codon 36 cDNA of the *MED12* gene.
3. The method of claim 1, wherein the missense mutation is located at position 130
and/or 131 of codon 44 cDNA of the *MED12* gene
4. The method of claim 1, wherein the missense mutation results in p.G44A, p.G44C,
p.G44D, p.G44R, p.G44S, or p.G44V in a polypeptide translated from the *MED12*
gene.
5. The method of claim 1, wherein the performing a nucleic acid-based assay comprises
sequencing the polynucleotide.
6. The method of claim 1, wherein the sample comprises stromal tissues.

Breast Fibroadenoma Susceptibility Mutations And Use Thereof

5

Abstract

The present disclosure provides a method of assaying susceptibility and/or confirming diagnosis of breast fibroadenomas development in a human subject. Preferably, the method comprises the steps of performing a nucleic acid-based assay to analyze an isolated
10 polynucleotide encoding at least exon 2 of *MED12* gene from a sample acquired from the human subject; and regarding the human subject with greater susceptibility and/or confirming diagnosis of breast fibroadenomas development by detecting a mutation in the isolated polynucleotide. The mutation can be a splice site mutation located at position -8 of exon 2 of the *MED12* gene, a missense mutation located at codon 44 of cDNA of the *MED12* gene or a
15 missense mutation located at codon 36 of cDNA of the *MED12* gene.

(Most illustrated by Figure 1)





Fig.1


gDNA-Tumour

ATGTAAACAAAGTTTCAATAACCAGC



Sample001

ATGTAAACAAAGTTTCAATAACCAGC




Sample002

ATGTAAACAAAGTTTCAATAACCAGC




Sample003

ATGTAAACAAAGTTTCAATAACCAGC




Sample006

ATGTAAACAAAGTTTCAATAACCAGC




Sample007

ATGTAAACAAAGTTTCAATAACCAGC




Sample008

CTAAACGCCGCTTTCCGCCTCAGGATGAAC



Sample004


CTAAACGCCGCTTTCCGCCTCAGGATGAAC




Sample005

gDNA-Normal


ATGTAAACAAAGTTTCAATAACCAGC



ATGTAAACAAAGTTTCAATAACCAGC




ATGTAAACAAAGTTTCAATAACCAGC




ATGTAAACAAAGTTTCAATAACCAGC




ATGTAAACAAAGTTTCAATAACCAGC



ATGTAAACAAAGTTTCAATAACCAGC



CTAAACGCCGCTTTCCGCCTCAGGATGAAC



CTAAACGCCGCTTTCCGCCTCAGGATGAAC




Fig.2

cDNA




Fig.3

a



b




Fig.4




Fig.5




Fig.6

SEQ ID NO. 15

5

10

15

20

25

30

35

40

45

50

55

```

1 attgtccgat ggttcccggc gtacctcggc ttccctcggt agtttccggc aatggtcgag
61 agtttctaac gtgccccctt gttgtctctc ggccgccgtc ctctcaacca cgcgccccct
121 ttccggctcc ctctccccct tcccggtccc ccagtcagcc tggccctgct ggtgcctccg
181 gcgctacggg ctgggcaaga tggcgccctt cgggatcttg agctacgaac accggccccct
241 gaagcggccg cggctggggc ctcccgatgt ttaccctcag gaccccaaac agaaggagga
301 tgaactgacg gccttgaatg taaaacaagg tttcaataac cagcctgctg tctctgggga
361 tgagcatggc agtgccaaga acgtcagctt caatcctgcc aagatcagtt ccaacttcag
421 cagcattatt gcagagaaat tacgttgtaa tacccttcct gacactggtc gcaggaagcc
481 ccaagtgaac cagaaggata acttctggct ggtgactgca cgatcccaga gtgccattaa
541 cacttggttc actgacttgg ctggcaccaa gccactcacg caactagcca aaaaggtccc
601 cattttcagt aagaaggaag aggtgtttgg gtacttagcc aaatacacag tgcctgtgat
661 gcgggctgcc tggctcatta agatgacctg tgcctactat gcagcaatct ctgagacca
721 ggttaagaag agacatggtg accctttcat ggaatggact cagatcatca ccaagtactt
781 atgggagcag ttacagaaga tggctgaata ctaccggcca gggcctgcag gaagtggggg
841 ctgtgggtcc acgatagggc ccttgcccca tgatgtagag gtggcaatcc ggcagtggga
901 ttacaccgag aagctggcca tgttcatgtt tcaggatgga atgctggaca gacatgagtt
961 cctgacctgg gtgcttgagt gttttgagaa gatccgccct ggagaggatg aattgcttaa
1021 actgctgctg cctctgcttc tccgatactc tggggaattt gttcagtcgt catacctgtc
1081 ccgcccggct gcctacttct gtacacggag actggccctg cagctggatg tgtgtagcag
1141 tcactcatct catgttatat ctgctcagtc aacaagcacg ctaccacca cccctgctcc
1201 tcagccccc aactagcaga caccctcgac tccctttagt gacctgtta tgtgccctca
1261 gcaccggccc ctgggttttg gcctcagctg tatcctacag accatcctcc tgtgctgtcc
1321 tagtgccttg gtttggcact actcactgac tgatagcaga attaagaccg gctcaccact
1381 tgaccacttg cctattgccc cgtccaacct gcccatgcca gagggtaaca gtgccttcac
1441 tcagcaggtc cgtgcaaagt tgcgggagat cgagcagcag atcaaggagc ggggacaggc
1501 agttgaagtt cgctggctct tcgataaatg ccaggaagct actgcaggct tcaccattgg
1561 acgggtactt catacttttg aagtgtgga cagccatagt tttgaacgct ctgacttcag
1621 caactctctt gactcccttt gtaaccgaat ctttggattg ggacctagca aggatgggca
1681 tgagatctcc tcagatgatg atgctgtggt gtcattgcta tgtgaatggg ctgtcagctg
1741 caagcgttct ggtcggcatc gtgctatggt ggtagccaag ctctggaga agagacaggc
1801 ggagattgag gctgagcgtt gtggagaatc agaagccgca gatgagaagg gttccatcgc
1861 ctctggetcc ctttctgctc ccagtgtctc cattttccag gatgtcctcc tgcagtttct
1921 ggatacacag gctcccatgc tgacggaccc tcgaagttag agtgagcggg tgggaattctt
1981 taacttagta ctgctgttct gtgaactgat tcgacatgat gttttctccc acaacatgta
2041 tacttgcact ctcatctccc gaggggacct tgcttttga gccctggtc cccggcctcc
2101 ctctcccttt gatgatcctg ccgatgacct agagcacaag gaggctgaag gcagcagcag
2161 cagcaagctg gaagatccag ggctctcaga atctatggac attgacccta gttccagtgt
2221 tctctttgag gacatggaga agcctgattt ctcatgttcc tcccacta tgcctgtga
2281 ggggaagggc agtccatccc ctgagaagcc agatgtcgag aaggaggatga agccccacc
2341 caaggagaag attgaaggga cccttggggg tctttacgac cagccacgac acgtgcagta
2401 cgccacccat tttcccatcc ccaggagga gtcatgcagc catgagtgca accagcgggt
2461 ggtcgtactg tttgggtgg gaaagcagcg agatgatgcc cgccatgcca tcaagaaaat
2521 caccaaggat atcttgaagg ttctgaaccg caaagggaca gcagaaactg accagcttgc
2581 tctattgtg cctctgaatc ctggagacct gacattctta ggtggggagg atgggcagaa
2641 gcggcgacgc aaccggcctg aagccttccc cactgctgaa gatatctttg ctaagttcca
2701 gcacctttca cattatgacc aacaccaggc cacggctcag gtctcccgga atgttctgga
2761 gcagatcacg agctttgccc ttggcatgtc ataccacttg cctctggtgc agcatgtgca
2821 gttcatcttc gacctcatgg aatattcact cagcatcagt ggctcatcg actttgccat
2881 tcagctgctg aatgaactga gtgtagttga ggctgagctg cttctcaaat cctcgatct
2941 ggtgggcagc tacactacta gcctgtgcct gtgcatcgtg gctgtcctgc ggcactatca
3001 tgctgcctc atcctcaacc aggaccagat ggcacaggct tttgaggggc tgtgtggcgt

```

	3061	cgtgaagcat	gggatgaacc	ggtccgatgg	ctcctctgca	gagcgcgtgta	tccttgctta
	3121	tctctatgat	ctgtacacct	cctgtagcca	tttaaagaac	aaatttgggg	agctcttcag
	3181	cgacttttgc	tcaaagggtga	agaacacccat	ctactgcaac	gtggagccat	cggaatcaaa
	3241	tatgcgctgg	gcacctgagt	tcatgatcga	cactctagag	aacctgcag	ctcacacctt
5	3301	cacctacacg	gggctaggca	agagtcttag	tgagaaccct	gctaaccgct	acagctttgt
	3361	ctgcaatgcc	cttatgcacg	tctgtgtggg	gcaccatgat	cccgataggg	tgaatgacat
	3421	cgcaatcctg	tgtgcagagc	tgaccggcta	ttgcaagtca	ctgagtgcag	aatggctagg
	3481	agtgccttaag	gccttgtgct	gctcctctaa	caatggcact	tgtggtttca	acgatctcct
10	3541	ctgcaatggt	gatgtcagtg	acctatcttt	tcatgactcg	ctggctactt	ttgttgccat
	3601	cctcatcgct	cggcagtggt	tgctcctgga	agatctgatt	cgctgtgctg	ccatcccttc
	3661	actccttaat	gctgcttgta	gtgaacagga	ctctgagcca	ggggcccggc	ttacctgccg
	3721	catcctcctt	caccttttca	agacaccgca	gctcaatcct	tgccagctcg	atggaaacaa
	3781	gcctacagta	ggaatccgct	cctcctgcga	ccgccacctg	ctggctgcct	cccagaaccg
15	3841	catcgtggat	ggagccgtgt	ttgctgttct	caaggctgtg	tttgtacttg	gggatgcgga
	3901	actgaaaggt	tcaggcttca	ctgtgacagg	aggaacagaa	gaacttccag	aggaggaggg
	3961	aggaggtggc	agtgggtggc	ggaggcaggg	tggccgcaac	atctctgtgg	agacagccag
	4021	tctggatgtc	tatgccaagt	acgtgctgcg	cagcatctgc	caacaggaat	gggtaggtag
	4081	acgttgcctt	aagtctctgt	gtgaggacag	caatgacctg	caagaccag	tgttgagttag
	4141	tgcccaggcg	cagcgccctca	tgcaagctcat	ttgctatcca	catcgactgc	tggacaatga
20	4201	ggatggggaa	aacccccagc	ggcagcgcct	aaagcgcatt	ctccagaact	tggaccagtg
	4261	gaccatgcgc	cagtcttcct	tggagctgca	gctcatgatc	aagcagaccc	ctaacaatga
	4321	gatgaactcc	ctcttgagga	acatcgccaa	ggccacaatc	gagggttttc	aacagtcagc
	4381	agagacaggg	tcatcttctg	gaagtactgc	aagcaacatg	cccagcagca	gcaagaccaa
25	4441	gcctgtgctc	agctctctag	agcgcctctg	tgtatggctg	gtggccccc	tcattgctaa
	4501	actgcccacc	tcagtccagg	gacatgtgtt	aaaggctgct	ggggaagaat	tggagaaggg
	4561	tcagcacctg	ggttcctcct	cacgcaaaga	acgtgatcga	caaaagcaga	agagcatgtc
	4621	cctattgagc	cagcagccct	tcttatcgct	ggtgctaaca	tgtctgaaag	ggcaggatga
	4681	acaacgcgag	ggactcctta	cctccctcta	cagccagggtg	caccagattg	tgaataattg
30	4741	gcgagatgac	cagtacttag	atgattgcaa	accaaagcag	cttatgcatg	aggcactcaa
	4801	actgcggctc	aacctggtgg	ggggcatgtt	tgacacgggtg	cagcgcagca	cccagcagac
	4861	cacggagtgg	gccatgctcc	tcctggagat	catcatcagc	ggcactgtcg	acatgcagtc
	4921	caacaatgag	ctcttcaacta	ctgtgttgga	catgctgagc	gtgctcatca	atgggacatt
	4981	ggctgcagac	atgtctagca	tctcgcaagg	tagcatggag	gaaaacaagc	gtgcatacat
35	5041	gaacctggcg	aagaagttgc	agaaggagtt	gggggagcgc	cagtcagaca	gtctggaaaa
	5101	ggttcgccag	ctgctgccac	tgcccaagca	gacccgagat	gtcatcacgt	gtgagccaca
	5161	gggtccctt	atcgatacca	agggcaacaa	gattgctggc	ttcgattcca	tcttcaagaa
	5221	ggagggtcta	caggttttcca	ccaaacagaa	gatctcgccc	tgggatcttt	ttgaggggtt
	5281	gaagccgtca	gcaccactct	cttggggctg	gtttggaaca	gtccgagtgg	accggcgagt
40	5341	ggctcgagga	gaggagcagc	agcggttgct	gctctaccac	acacacctga	ggccccggcc
	5401	ccgcgcctat	tacctggagc	cactgccact	gccccagaa	gatgaggagc	cgctgtctcc
	5461	tacctgcta	gagcctgaga	aaaaggctcc	agagccccc	aaaactgaca	aaccgggggc
	5521	tgtccaccc	agtactgagg	aacgcaagaa	gaagtccacc	aagggcaaga	aacgcagcca
	5581	gccagctacc	aagacagagg	actatggaat	gggcccggt	cggagcggcc	cttatggtgt
45	5641	gacagtgcct	ccggacctcc	tgcaccaccc	aaaccctggt	tctataacac	accttaacta
	5701	caggcaaggc	tccataggcc	tgtacaccca	gaaccagcca	ctacctgcag	gtggccctcg
	5761	tgtggaccca	taccgtcctg	tgcgcttacc	aatgcagaag	ctgcccaccc	gaccaactta
	5821	ccctggagtg	ctgcccacaa	ccatgactgg	cgtcatgggt	ttagaaccct	cctcttataa
	5881	gacctctgtg	taccggcagc	agcaacctgc	ggtgccccaa	ggacagcgcc	ttcgccaaca
50	5941	gctccagcag	agtcaaggca	tgttgggaca	gtcatctgtc	catcagatga	ctcccagctc
	6001	ttcctacggt	ttgcagactt	cccagggtca	tactccttat	gtttctcatg	tgggattgca
	6061	gcaacacaca	ggccctgcag	gtaccatggt	gcccccagc	tactccagcc	agccttacca
	6121	gagcaccac	ccttctacca	atcctactct	tgtagatcct	accgcgccc	tgcacagcg
	6181	gccagtggt	tatgtgcacc	agcaggcccc	cacctatgga	catgactga	cctccactca
55	6241	aaggttttca	caccagacac	tgcagcagac	acccatgata	agtaccatga	ctccaatgag
	6301	tgcccagggc	gtccaggcag	gcgtccgttc	aacagccatc	ctacctgagc	agcagcagca
	6361	gcagcaacag	cagcaacagc	aacagcagca	gcagcagcaa	cagcaacagc	agcagcagca
	6421	gcagcagtac	cacatccggc	agcagcagca	gcagcagatc	ctgcggcagc	agcagcaaca
	6481	gcaacagcag	cagcagcagc	agcagcaaca	gcaacagcag	cagcagcaac	agcaacaaca

5 6541 gcaacaccag cagcaacagc agcaacaggc ggctcctccc caaccccagc cccagtccca
 6601 gccccagttc cagcgccagg ggcttcagca gacccagcag cagcaacaga cagcagcttt
 6661 ggtccggcaa cttcaacaac agctctctaa taccagcca cagccagta ccaacatatt
 6721 tggacgctac tgagccacct ggaggaactg cttgtgcact ggatgtggcc ccaccctttc
 6781 ctcttaattc ccaatcccat tcctgggcta gcaccagtag tgggtggggc cctccctca
 6841 ggctccattt ttaataagtt tttagtatatt ttgttaatgt gaggcattga gctgttgggt
 6901 tttgtatatt atttatatag agaccccaga gctgttgac ccaatacaca gagcttcttt
 6961 gcaaagggaa aaaaaaaaaa aaaaa

Fig.7a

SEQ ID NO. 16

15 MAAFGILSYEHRPLKRPRLGPPDVYPQDPKQKEDELTAENVKQGFNNQPAVSGDEHGSANKVSNPAKISSNFSS
 IIAEKLRCNTLPDTGRRKPQVNQKDNFWLVARSQSAINTWFTDLATGKPLTQLAKKVP IFSKKEEVFGYLAKYT
 VPVMRAAWLIKMTCAYYAAISETKVKKRHVDPFMEWTQIIITKYLWEQLQKMAEYYRPGPAGSGGCGSTIGPLPHD
 VEVAIRQWDYTEKLAMFMFQDGM LDRHEFLTWVLECFEKIRPGEDELLKLLPLLLRYSGEFVQSAYLSRRLAYF
 CTRRLALQLDGVSSSHSHVISAQSTSTLPTTPAPQPPTSSTPSTPFSDDL MCPQHRPLVFGLSCILQTIILLCCPS
 20 ALVWHYSLTDSRIKTGSPLDHLP IAPSNLPMPEGNSAFTQQVRAKLREIEQQIKERGQAVEVRWSFDKCEATAG
 FTIGRVLHTLEVLD SHSFERSDFSNSLDSL CNRIFGLGPSKD GHEISSDDDAVVSLLCEWAVSCKRSGRHRAMVV
 AKLLEKRQAEIEAERCGESEAADEKGSIASGSL SAPSAP IFQDVLLQFLDTQAPMLTDPRSESERVEFFNLVLLF
 CELIRHDVFSHNMYTCTLISRGLAFGAPGPRPPSPFDDPADDPEHKEAEGSSSSKLEDPGLSEMDIDPSSSVL
 FEDMEKPDFSLFPTMPCEGKGSPSPEKPDVEKEVKPPPEKIEGLGVLYDQPRHVQYATHFPIQEESSHEC
 25 NQRLVVLFGVGKQRDDARHAIKKITKDILKVLNRKGTAE TDQLAPIVPLNPGDLTFLGGEDGQKRRNRPEAFPT
 AEDIFAKFQHL SHYDQHQVTAQVSRNVLEQITSFALGMSYHLPLVQHVFIFDLMEYSLSISGLIDFAIQLNEL
 SVVEAELLKSSDLVGSYTTSLCLCIVAVLRHYHACLILNQDQMAQVFEGLCGVVKHGMNRS DGSSAERCILAYL
 YDLYTSCSHLKNKFGE LFSDFCSKVKN TIYCNVEPSESNMRAWPEFMIDTLENPAHTFTYTGLGKSLSEN PANR
 YSFVCNALMHVCVGHHD PDRVNDIAILCAELTGYCKSLSAEWLGV LKALCCSSNNGTCGFNDLLCNVDVSDLSFH
 30 DSLATFVAIILARQCLLLEDLIRCAAIPSLNNAACSEQDSEPGARLT CRILLHLFKTPQLNPCQSDGNKPTVGIR
 SSCDRHLLAASQNRIVDGA VFAVLKAVFVLGDAELKGS GFTVTGGTEELPEEEGGGGSGRRQGRNISVETASL
 DVYAKYVLR SICQEWVGERCLKSLCEDSN DLQDPVLSSAQARLMQLICYPHRLLDNEDGENPQRQRIKRILQN
 LDQWTRQSSLELQLMIKQTPNNEMNSLLENIAKATIEVFQSSAETGSSSGSTASNMPSSSKTKPVLSSSLERSGV
 WLVAPLIAKLPTSVQGHVLKAAGEELEKGQHLGSSSRKERDRQKQKMSLLSQQPFLSLVLTCLKGQDEQREGLL
 35 TSLYSQVHQIVNNWRDDQYLD DCKPKQLMHEALKRLNLVGGMFDTVQRSTQQTTEWAMLLLEIIISGTVMQSN
 NELFTTVLDMLSVLINGTLAADMSSISQGSMEENKRAYMNLAKKLQKELGERQSDSLEKVRQLPLPKQTRDVIT
 CEPQGS LIDTKGNKIAGFDSIFKKEGLQVSTKQKISPWDLFEG LKPSAPLSWGWF GTVRVDRRVARGEEQQRLLL
 YHTHLRPRPRAYYLEPLPLPPEDEEPPAP TLLPEKKAPEPPKTDKPGAAPPSTEERKKKSTKGKKRSQPATKTE
 DYGMGPRSGPYGVTVPPDLLHHPNPGSITHLNYRQGSIGLYTQNP LPAGGPRVDPYRPVRLPMQKLPTRPTYP
 40 GVLPTTMTGVMGLEPSSYKTSVYRQQQPAVPQGQRLRQQLQQSQGMLGQSSVHQMT PSSSYGLQTSQGYTPYVSH
 VGLQQHTGPAGTMVPPSYSSQPYQSTHPSTNPTLVDPTRHLQQRPSGYVHQQAPTYGHGLTSTQRFHQTLQQTP
 MISTMTPM SAQGVQAGVRSTAILPEQQQQQQQQQQQQQQQQQQQQQQYHIRQQQQQQQILRQQQQQQQQQQQ
 QQQQQQQQQQQQQQHQQQQQQQAAPPQPPQSQPQFQRQGLQQTQQQQQTAA LVRQLQQQLSNTQPQPSTNIFG
 RY

Fig.7b