

DESCRIPTION

Title of Invention: NOVEL PROSTAGLANDIN DERIVATIVE

[Technical Field]

[0001]

5 The present invention relates to a novel prostaglandin derivative having an alkynyl group on the ω -chain of a prostaglandin, a pharmaceutically acceptable salt thereof, or a cyclodextrin clathrate compound thereof, particularly, a novel prostaglandin derivative having a double bond at the 2-position
10 of a prostaglandin and an alkynyl group on the ω -chain. In addition, the present invention relates to a medicament containing at least one of the prostaglandin derivative, a pharmaceutically acceptable salt thereof, and a cyclodextrin clathrate compound thereof as an active ingredient,
15 particularly a medicament for the prophylaxis or treatment of a blood flow disorder.

[Background Art]

[0002]

Peripheral arterial occlusive disease is a disease in
20 which the artery is constricted or occluded due to arteriosclerosis and thrombus formation, the periphery, particularly the lower limb, falls into an ischemic state, and exhibits symptoms such as cryaesthesia, intermittent claudication, pain, and ulcer or necrosis of lower limb.
25 Improvement of the blood flow to the ischemic part is important for improving the lower limb symptoms, and treatments aiming at resuming blood circulation by drugs or physical methods are performed. Drugs having vasodilating action and platelet aggregation inhibiting action are used for drug therapy.

30 [0003]

Peripheral vascular disease also manifests as atherosclerotic stenosis of the renal artery and can lead to renal ischemia and kidney dysfunction. Chronic diabetes can also lead to atherosclerosis and vascular complications
35 involving large vessel, arteriole and capillary. Diabetic

patients are at high risk of developing foot ulcer due to long-term complications of, for example, neuropathy and ischemia.

[0004]

Spinal canal stenosis is a disease in which the spinal
5 canal is narrowed due to hypertrophy degeneration of the spine
and the ligamentum flavum that constitute the spinal canal and
protrusion of the intervertebral disc, and neural tissues such
as the nerve root and cauda equina are compressed and various
symptoms are exhibited. Spinal canal stenosis is classified
10 into extensive spinal canal stenosis, thoracic spinal canal
stenosis, lumbar spinal canal stenosis and the like according
to the narrow portion of the spinal canal. The symptoms
thereof include lumbago, pain in the upper limb or lower limb,
numbness and the like due to nerve compression. In particular,
15 when the cauda equina is injured, lumbago, lower limb pain,
numbness, weakness become severe during walking, and this
symptom is called intermittent claudication.

[0005]

Natural prostaglandins (hereinafter prostaglandin is to
20 be indicated as PG) are a group of bioactive substances
synthesized in vivo, and regulate cellular functions of each
tissue of the body as a topical hormone having various
physiological activities. In particular, PGE₁s, which are one
kind of natural PGs, have, for example, a vasodilating action,
25 an angiogenesis action, a platelet aggregation suppressive
action, and an epithelial regeneration promoting action. They
are used as an antiplatelet agent, an agent for improving
peripheral blood flow disorders and the like in the drug
therapy of the above-mentioned diseases. While PGEs may be
30 applicable to other indications, natural PGEs are extremely
unstable chemically and metabolically. Thus, the development
of PGE derivatives which are more stable and effective, and
cause fewer side effects has been extensively studied.

[0006]

35 A PG derivative having a double bond at the 2-position of

PG and a production method thereof are reported in the following patent documents 1 - 5 and non-patent documents 1 - 2. In addition, a PG derivative having an alkynyl group in the ω -chain of PG and a production method thereof are reported in the following patent documents 6 - 7.

[Document List]

[Patent documents]

[0007]

Patent document 1: JP-A-50-71649

10 Patent document 2: JP-A-50-116452

Patent document 3: JP-A-52-85151

Patent document 4: JP-A-53-149954

Patent document 5: JP-A-55-100360

Patent document 6: JP-A-51-131860

15 Patent document 7: JP-A-54-12352

[Non-patent documents]

[0008]

Non-patent document 1: Ann. Acad. N. Y. Sci., 1971, vol. 180, p. 181.

20 Non-patent document 2: Prostaglandins, 1974, vol. 8, p. 341.

[Summary of Invention]

[Problems to be Solved by the Invention]

[0009]

25 An object of the present invention is to provide a blood flow improving agent that can be administered orally, causes fewer side effects, shows higher safety, and is superior in the effectiveness and pharmacokinetic property, especially, a compound useful as a therapeutic agent for blood flow disorder and pain associated with spinal canal stenosis. Among the compounds described in the aforementioned documents, particularly, (2E)-7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,5S)-3-hydroxy-5-methylnon-1-en-1-yl)-5-oxocyclopentyl)hept-2-enoic acid (limaprost) is a compound showing superior efficacy, and
30
35 blood flow improving agents containing limaprost as an active

ingredient are being used for the treatment of pain and
cryaesthesia associated with thromboangiitis obliterans and
spinal canal stenosis in the actual medical field. However,
limaprost has side effects on the gastrointestinal tract, and
5 there is still room for improvement in the blood flow
increasing action and sustainability of drug efficacy. In
particular, the side effect of limaprost that has a high smooth
muscle contraction action and causes diarrhea is left as a
problem to be solved. Therefore, the development of a PGE1
10 derivative having physiological activities similar to those of
natural type, causing low side effects and having high
sustainability has been intensively studied in and out.
[0010]

However, a PG compound having a double bond at the 2-
15 position of PG and an alkynyl group in the ω -chain has not been
reported. Furthermore, synthetic examples, properties,
physiological activities and the like of a PGE compound having
a double bond at the 2-position of PG and a triple bond at the
18-position have not been reported.

20 In addition, synthetic examples, properties,
physiological activities and the like of a PGE compound having
a triple bond at the 18-position of PG and a cycloalkyl group
on the end of the triple bond have not been reported at all.
[Means of Solving the Problems]

25 [0011]


The present inventors have synthesized novel PG compounds
and conducted studies in an attempt to clarify the properties
and physiological activities. As a result, they have found
that a compound represented by the following formula (1)
30 (hereinafter sometimes to be indicated as compound (1)) or a
pharmaceutically acceptable salt thereof (hereinafter compound
(1) and a pharmaceutically acceptable salt thereof are
sometimes to be generically indicated as "the compound of the
present invention"), or cyclodextrin clathrate compounds
35 thereof have superior properties and pharmacological action,

and further that they are extremely superior compounds as blood flow improving agents, especially therapeutic drugs for blood flow disorder and pain associated with spinal canal stenosis, which resulted in the completion of the present invention.

5 Therefore, the present invention provides the following.

[1] A compound represented by the following formula (1) or a pharmaceutically acceptable salt thereof;

[0012]



10 [0013]

wherein

R^A is $-\text{CH}_2-\text{CZ}^1\text{Z}^2(\text{COX})$ or $-\text{CH}=\text{CZ}^1(\text{COX})$, Z^1 and Z^2 are each independently a hydrogen atom or a fluorine atom, X is OR^4 or NR^5R^6 , R^4 is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms or a substituted alkyl group having 1 - 6 carbon atoms, R^5 is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms or a substituted alkyl group having 1 - 6 carbon atoms, R^6 is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms, a substituted alkyl group having 1 - 6 carbon atoms, an alkylsulfonyl group having 1 - 6 carbon atoms, a substituted alkylsulfonyl group having 1 - 6 carbon atoms, an arylsulfonyl group having 6 - 10 carbon atoms or a substituted arylsulfonyl group having 6 - 10 carbon atoms;

Y is CH_2 , S or O ;

25 A-B is a carbon-carbon single bond, a carbon-carbon double bond or a carbon-carbon triple bond;

a hydroxy group bonded by a wavy line is a hydroxy group having α -configuration, β -configuration or a mixed configuration of α -configuration and β -configuration;

30 R^1 and R^2 are each independently a hydrogen atom, an

alkyl group having 1 - 3 carbon atoms or a substituted alkyl group having 1 - 3 carbon atoms;

n is an integer of 0 - 2;

R³ is an alkyl group having 1 - 4 carbon atoms, a substituted alkyl group having 1 - 4 carbon atoms, a cycloalkyl group having 3 - 6 carbon atoms, a substituted cycloalkyl group having 3 - 6 carbon atoms, an aryl group having 6 - 10 carbon atoms or a substituted aryl group having 6 - 10 carbon atoms;

excluding 7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4RS)-3-hydroxy-4-methyloct-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoic acid;

[2] the compound or a pharmaceutically acceptable salt thereof of [1], wherein Y is CH₂;

[3] the compound or a pharmaceutically acceptable salt thereof of [1] or [2], wherein R^A is -CH=CZ¹(COX);

[4] the compound or a pharmaceutically acceptable salt thereof of [1] or [2], wherein R^A is -CH₂-CZ¹Z²(COX);

[5] the compound or a pharmaceutically acceptable salt thereof of any of [1] - [4], wherein n is 1;

[6] the compound or a pharmaceutically acceptable salt thereof of any of [1] - [5], wherein A-B is a carbon-carbon double bond;

[7] the compound of or a pharmaceutically acceptable salt thereof any of [1] - [6], wherein R¹ is a hydrogen atom and R² is an alkyl group having 1 - 3 carbon atoms;

[8] the compound or a pharmaceutically acceptable salt thereof of any of [1] - [7], wherein R³ is an alkyl group having 2 - 3 carbon atoms or a cycloalkyl group having 3 - 5 carbon atoms;

[9] the compound or a pharmaceutically acceptable salt thereof of any of [1] - [8], wherein R³ is a cycloalkyl group having 3 - 5 carbon atoms;

[10] the compound or a pharmaceutically acceptable salt thereof of any of [1] - [9], wherein X is OR⁴;

[11] a cyclodextrin clathrate compound of the compound of any of [1] - [10] or a pharmaceutically acceptable salt thereof;

[12] a medicament comprising the compound or a pharmaceutically acceptable salt thereof of any of [1] - [10], or the cyclodextrin clathrate compound of [11] as an active ingredient;

5 [13] the medicament of [12], which is a prophylactic or therapeutic agent for a blood flow disorder;

[14] the medicament of [13], wherein the blood flow disorder is a blood flow disorder of nerve;

[15] the medicament of [14], wherein the blood flow disorder of
10 nerve is a blood flow disorder associated with spinal canal stenosis;

[16] the medicament of [13], wherein the blood flow disorder is a blood flow disorder of peripheral artery, skin or brain;

[17] the medicament of [16], wherein the blood flow disorder of
15 peripheral artery is a blood flow disorder associated with chronic arterial occlusion or pulmonary hypertension;

[18] the medicament of [16], wherein the blood flow disorder of skin is a blood flow disorder associated with pressure ulcer;

[19] the medicament of [16], wherein the blood flow disorder of
20 brain is a blood flow disorder associated with suppression of recurrence after cerebral infarction;

[20] a method for prophylaxis or treatment of a blood flow disorder, comprising administering an effective amount of the compound or a pharmaceutically acceptable salt thereof of any
25 of [1] - [10], or the cyclodextrin clathrate compound of [11] to a patient in need thereof;

[21] the method of [20], wherein the blood flow disorder is a blood flow disorder of nerve;

[22] the method of [21], wherein the blood flow disorder of
30 nerve is a blood flow disorder associated with spinal canal stenosis;

[23] the method of [20], wherein the blood flow disorder is a blood flow disorder of peripheral artery, skin or brain;

[24] the method of [23], wherein the blood flow disorder of
35 peripheral artery is a blood flow disorder associated with

chronic arterial occlusion or pulmonary hypertension;
[25] the method of [23], wherein the blood flow disorder of
skin is a blood flow disorder associated with pressure ulcer;
[26] the method of [23], wherein the blood flow disorder of
5 brain is a blood flow disorder associated with suppression of
recurrence after cerebral infarction;
[27] the compound or a pharmaceutically acceptable salt thereof
of any of [1] - [10], or the cyclodextrin clathrate compound of
[11] for use in the treatment of a blood flow disorder;
10 [28] the compound or a pharmaceutically acceptable salt thereof,
or the cyclodextrin clathrate compound for use of [27], wherein
the blood flow disorder is a blood flow disorder of nerve;
[29] the compound or a pharmaceutically acceptable salt thereof,
or the cyclodextrin clathrate compound for use of [28], wherein
15 the blood flow disorder of nerve is a blood flow disorder
associated with spinal canal stenosis;
[30] the compound or a pharmaceutically acceptable salt thereof,
or the cyclodextrin clathrate compound for use of [27], wherein
the blood flow disorder is a blood flow disorder of peripheral
20 artery, skin or brain;
[31] the compound or a pharmaceutically acceptable salt thereof,
or the cyclodextrin clathrate compound for use of [30], wherein
the blood flow disorder of peripheral artery is a blood flow
disorder associated with chronic arterial occlusion or
25 pulmonary hypertension;
[32] the compound or a pharmaceutically acceptable salt thereof,
or the cyclodextrin clathrate compound for use of [30], wherein
the blood flow disorder of skin is a blood flow disorder
associated with pressure ulcer;
30 [33] the compound or a pharmaceutically acceptable salt thereof,
or the cyclodextrin clathrate compound for use of [30], wherein
the blood flow disorder of brain is a blood flow disorder
associated with suppression of recurrence after cerebral
infarction;
35 [34] use of the compound or a pharmaceutically acceptable salt

thereof of any of [1] - [10], or the cyclodextrin clathrate compound of [11] in the manufacture of a prophylactic or therapeutic agent for a blood flow disorder;

[35] the use of [34], wherein the blood flow disorder is a
5 blood flow disorder of nerve;

[36] the use of [35], wherein the blood flow disorder of nerve is a blood flow disorder associated with spinal canal stenosis;

[37] the use of [34], wherein the blood flow disorder is a blood flow disorder of peripheral artery, skin or brain;

10 [38] the use of [37], wherein the blood flow disorder of peripheral artery is a blood flow disorder associated with chronic arterial occlusion or pulmonary hypertension;

[39] the use of [37], wherein the blood flow disorder of skin is a blood flow disorder associated with pressure ulcer;

15 [40] the use of [37], wherein the blood flow disorder of brain is a blood flow disorder associated with suppression of recurrence after cerebral infarction and the like.

[Effect of the Invention]

20 [0014]

The compound of the present invention has a very potent suppressive action on platelet aggregation in human and rat, and a superior blood flow increasing action when intravenously administered to rat in a cauda equine blood flow test. Thus,
25 the compound is useful as a prophylactic or therapeutic agent for blood flow disorders.

In addition, the compound of the present invention is less susceptible to oxidative metabolism by PG dehydrogenase and has high metabolic stability since an alkyl group
30 (particularly a methyl group) is bonded to the 16-position carbon atom adjacent to the 15-position hydroxy group of the PG skeleton. Consequently, the compound has a long half-life in blood and can maintain the effective blood concentration for a long time as compared to natural type PG compounds. The
35 improved metabolic stability can markedly improve

bioavailability of a drug containing the compound of the present invention.

[Description of Embodiments]

[0015]

5 The definition of each symbol in the formula (1) is described in detail below.

For naming of the compounds in the present specification, the IUPAC nomenclature is used in principle. To indicate the position of the PG skeleton, numerical numbering based on the
10 prostanoic acid skeleton may sometimes be indicated as appropriate. In the present specification, a group in which a hydrogen atom of the alkyl group is substituted is also indicated as a substituted alkyl group. Similar indication is adopted for other groups as well. The "lower" means that the
15 carbon number is 1 - 6, and 1 - 4 is preferable.

[0016]

The "alkyl group" may be a linear chain or a branched chain. The alkyl group is preferably an alkyl group having 1 - 6 carbon atoms, particularly preferably an alkyl group having 1 -
20 - 4 carbon atoms. Examples of the alkyl group include methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, sec-butyl group, tert-butyl group, pentyl group, hexyl group.

[0017]

25 The "alkenyl group" may be a linear chain or a branched chain. The alkenyl group is preferably an alkenyl group having 2 - 6 carbon atoms. Examples of the alkenyl group include vinyl group, allyl group, 1-propenyl group, isopropenyl group, 3-butenyl group, 3-pentenyl group, 4-hexenyl group and the like.

30 [0018]

The "alkynyl group" may be a linear chain or a branched chain. The alkynyl group is preferably an alkynyl group having 2 - 6 carbon atoms. Examples of the alkynyl group include ethynyl group, 1-propynyl group, 2-propynyl group, 3-butynyl
35 group, 3-pentynyl group, 4-hexynyl group and the like.

[0019]

The "alkoxy group" is a group in which an oxygen atom is bonded to the terminal carbon atom of the alkyl group. The alkoxy group may be a linear chain or a branched chain. The
5 alkoxy group is preferably an alkoxy group having 1 - 6 carbon atoms, and particularly preferably an alkoxy group having 1 - 4 carbon atoms. Examples of the alkoxy group include methoxy group, ethoxy group, propoxy group, butoxy group and the like.

[0020]

10 The "halogen atom" is a fluorine atom, a chlorine atom, a bromine atom, or an iodine atom.

[0021]

The "aryl group" is an aromatic hydrocarbon group, preferably an aryl group having 6 - 18 carbon atoms,
15 particularly preferably an aryl group having 6 - 10 carbon atoms. Examples of the aryl group include phenyl group, naphthyl group, anthryl group and the like, and phenyl group is particularly preferable.

[0022]

20 The "aralkyl group" is an alkyl group to which an aryl group is bonded. The aryl group of the aralkyl group is preferably an aryl group having 6 - 10 carbon atoms, particularly preferably a phenyl group. The alkyl group of the aralkyl group is preferably an alkyl group having 1 - 4 carbon
25 atoms. Examples of the aralkyl group include benzyl group, benzhydryl group, trityl group, phenylethyl group and the like.

[0023]

The "cycloalkyl group" is a cyclic alkyl group having a ring with three or more members, preferably a 3- to 6-membered
30 cycloalkyl group. Examples of the cycloalkyl group include cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group, cycloheptyl group and the like.

[0024]

35 The "alkylsulfonyl group" is a monovalent group in which

-S(O)₂- is bonded to the terminal carbon atom of the alkyl group. The carbon number of the alkyl group of the alkylsulfonyl group is preferably 1 to 6. Examples of the alkylsulfonyl group include methanesulfonyl group,
5 ethanesulfonyl group, propanesulfonyl group, butanesulfonyl group, pentanesulfonyl group, hexanesulfonyl group and the like.
[0025]

The "arylsulfonyl group" is a monovalent group in which -S(O)₂- is bonded to the binding end of the aryl group. As the
10 aryl group of the arylsulfonyl group, a group having 6 - 10 carbon atoms is preferable, and a group similar to the aforementioned aryl group is preferable. Examples of the arylsulfonyl group include an arylsulfonyl group having 6 - 10 carbon atoms such as benzenesulfonyl group, 1-
15 naphthalenesulfonyl group, and 2-naphthalenesulfonyl group, and benzenesulfonyl group is preferable.
[0026]

The "acyl group" is a monovalent group obtained by removing a hydroxy group from the carboxy group of carboxylic
20 acid. As the carboxylic acid, carboxylic acid having 1 - 10 carbon atoms is preferable, formic acid, saturated aliphatic carboxylic acid (e.g., alkylcarboxylic acid such as acetic acid, propionic acid, and butyric acid), unsaturated aliphatic carboxylic acid (e.g., alkenylcarboxylic acid such as acrylic
25 acid), carbocyclic carboxylic acid, heterocyclic carboxylic acid and the like can be mentioned.

Examples of the carbocyclic carboxylic acid include saturated cyclic aliphatic carboxylic acid (e.g., cycloalkylcarboxylic acid such as cyclopropylcarboxylic acid,
30 cyclopentylcarboxylic acid, and cyclohexylcarboxylic acid), unsaturated cyclic aliphatic carboxylic acid (e.g., cycloalkenylcarboxylic acid such as cyclohexenylcarboxylic acid) and aromatic carboxylic acid. The aromatic carboxylic acid is a compound in which the aforementioned aryl group is
35 bonded to a carboxy group and, for example, arylcarboxylic acid

such as benzoic acid and naphthylcarboxylic acid can be mentioned. The heterocyclic carboxylic acid is a compound in which a heterocyclic group is bonded to the carboxy group.

[0027]

5 The "heterocyclic group" is preferably a 3- to 10-membered monovalent saturated or unsaturated heterocyclic group containing at least one hetero atom selected from the group consisting of a nitrogen atom, an oxygen atom and a sulfur atom. Examples of the 3- to 10-membered saturated or unsaturated
10 heterocycle include aziridine, azetidine, pyrrole, pyrroline, pyrrolidine, imidazole, imidazoline, imidazolidine, pyrazole, pyrazoline, pyrazolidine, pyridine, piperidine, pyrazine, piperazine, pyrimidine, pyridazine, thiazole, thiazoline, thiazolidine, isothiazole, isothiazoline, isothiazolidine,
15 oxazole, oxazoline, oxazolidine, isoxazole, isoxazoline, isoxazolidine, morpholine, azepane, azepine and the like.

[0028]

 In the present specification, substitution of one or more hydrogen atoms in a group with a halogen atom is called
20 halogenation. For example, a halogenated alkoxy group is a group in which one or more hydrogen atoms of an alkoxy group is/are substituted by a halogen atom. The same applies to other groups. As the halogen atom, a fluorine atom or a chlorine atom is preferable.

25 A preferable example of the halogenated group having a halogenated alkyl group in its structure is a halogenated lower alkyl group, and fluoromethyl group, difluoromethyl group, trifluoromethyl group, trifluoroethyl group, pentafluoroethyl group, chloromethyl group, bromomethyl group and the like can
30 be mentioned.

[0029]

 The "alkoxyalkyl group" is an alkyl group in which one of the hydrogen atoms in the alkyl group is substituted by an alkoxy group. The alkoxy group in the alkoxyalkyl group is
35 preferably an alkoxy group having 1 - 4 carbon atoms. The

alkyl group in the alkoxyalkyl group is preferably an alkyl group having 1 - 4 carbon atoms. The carbon number of the alkoxyalkyl group is preferably 2 to 6, more preferably 2 to 4. Examples of the alkoxyalkyl group include methoxymethyl group, 5 ethoxymethyl group, propoxymethyl group, ethoxymethyl group, ethoxyethyl group and the like.

[0030]

The "hydroxyalkyl group" is a group in which one or more hydrogen atoms in the alkyl group is/are substituted by a 10 hydroxy group. The carbon number of the hydroxyalkyl group is preferably 2 to 6. The number of hydroxy groups of the hydroxyalkyl group is preferably one. Examples of the hydroxyalkyl group include 2-hydroxyethyl group, 3-hydroxypropyl group, 2-hydroxypropyl group, 4-hydroxybutyl 15 group and the like.

[0031]

The "alkoxycarbonyl group" is a group in which the aforementioned alkoxy group is bonded the carbonyl group. The alkoxy carbonyl group is preferably an alkoxy carbonyl group in 20 which an alkoxy group having 1 - 6 carbon atoms is bonded to a carbonyl group. Examples of the alkoxy carbonyl group include methoxycarbonyl group, ethoxycarbonyl group, n-propoxycarbonyl group, isopropoxycarbonyl group, n-butoxycarbonyl group, t-butoxycarbonyl group, amyloxycarbonyl group and the like.

25 [0032]

The substituted alkyl group is a group in which one or more hydrogen atoms of the alkyl group is/are substituted by a substituent(s). Examples of the substituent include halogen atom, hydroxy group, alkoxy group, halogenated alkoxy group, 30 acyl group, acyloxy group, halogenated acyloxy group, alkylsulfonyloxy group, halogenated alkylsulfonyloxy group, arylsulfonyloxy group, arylsulfonyloxy group substituted halogenated alkyl group, alkoxy carbonyloxy group, amino group, substituted amino group, aryl group, heterocyclic group and the 35 like, with preference given to halogen atom, hydroxy group or

alkoxy group.

[0033]

The substituted cycloalkyl group is a group in which one or more hydrogen atoms in the cycloalkyl group is/are substituted by a substituent(s). Examples of the substituent include alkyl group, halogenated alkyl group, halogen atom, hydroxy group, alkoxy group, halogenated alkoxy group, acyloxy group, halogenated acyloxy group, alkylsulfonyloxy group, halogenated alkylsulfonyloxy group, arylsulfonyloxy group, halogenated arylsulfonyloxy group, arylsulfonyloxy group substituted by alkyl, alkoxycarbonyloxy group and the like, with preference given to alkyl group, halogen atom or alkoxy group.

Examples of the substituted cycloalkyl group include 2-methylcyclopropyl group, 2,2-dimethylcyclopropyl group, 2-fluorocyclopropyl group, 2-methylcyclobutyl group, 2,2-dimethylcyclobutyl group, 2-methylcyclopentyl group, 2-methylcyclohexyl group and the like.

[0034]

The substituted aryl group is a group in which one or more hydrogen atoms of the aryl group is/are substituted by a substituent(s). The substituent of the substituted aryl group is halogen atom, alkyl group, halogenated alkyl group, hydroxy group, alkoxy group, halogenated alkoxy group, alkylenedioxy group having 1 - 3 carbon atoms (e.g., methylenedioxy group and ethylenedioxy group) or the like, with preference given to halogen atom, alkyl group, halogenated alkyl group, or alkoxy group.

Examples of the substituted aryl group include mono-halophenyl group (e.g., 1-, 2- or 3-chlorophenyl group, 1-, 2- or 3-fluorophenyl group, and 1-, 2- or 3-bromophenyl group), (halogenated alkyl)phenyl group (e.g., 1-, 2- or 3-trifluoromethylphenyl group), and alkoxyphenyl group (e.g., 1-, 2- or 3-methoxyphenyl group and 1-, 2- or 3-ethoxyphenyl group).

[0035]

The substituted alkylsulfonyl group is a group in which one or more hydrogen atoms of the alkylsulfonyl group is/are substituted by a substituent(s). As the substituent of the substituted alkylsulfonyl group, a halogen atom is preferable.

5 Examples of the substituted alkylsulfonyl group include trifluoromethanesulfonyl group, pentafluoroethanesulfonyl group and the like.

[0036]

The substituted arylsulfonyl group is a group in which
10 one or more hydrogen atoms of the arylsulfonyl group is/are substituted by a substituent(s). The substituent of the substituted arylsulfonyl group is preferably halogen atom, alkyl group, halogenated alkyl group, or alkoxy group. Examples of the substituted arylsulfonyl group include
15 monohalogenated phenylsulfonyl group (e.g., 1-, 2- or 3-chlorobenzenesulfonyl group, 1-, 2- or 3-fluorobenzenesulfonyl group, and 1-, 2- or 3-bromobenzenesulfonyl group), (alkyl)phenylsulfonyl group (e.g., p-toluenesulfonyl group), (halogenated alkyl)phenylsulfonyl group (e.g.,
20 trifluoromethylbenzenesulfonyl group) and (alkoxy)phenyl group (e.g., methoxybenzenesulfonyl group and ethoxybenzenesulfonyl group).

[0037]

The substituted amino group is a group in which one or
25 two hydrogen atoms of the amino group is/are substituted. When two hydrogen atoms of the amino group are substituted by substituents, the substituents may be the same or different. Examples of the substituent of the substituted amino group include the amino-protecting groups described in
30 "Greene's/Protective/Groups/in/Organic/Synthesis 4th Edition" (P. G. M. Wuts, T. W. Greene, J. Wiley & Sons, 2007) and the like. Specifically, alkyl group having 1 - 6 carbon atoms, alkoxy group having 1 - 6 carbon atoms, alkenyl group having 2 - 6 carbon atoms, alkynyl group having 2 - 6 carbon atoms, aryl
35 group having 6 - 10 carbon atoms, monovalent heterocyclic group,

aralkyl group having 7 - 14 carbon atoms, benzhydryl group, trityl group, acyl group, tri-organosilyl group and the like can be mentioned. When the aforementioned substituent is a group containing a hydrogen atom, one or more of the hydrogen
5 atoms may be further substituted by a substituent(s) (hereinafter to be indicated as substituent 2). As the substituent 2, halogen atom, alkyl group having 1 - 6 carbon atoms, alkoxy group having 1 - 6 carbon atoms, alkoxycarbonyl group having 1 - 6 carbon atoms, cyano group, nitro group and
10 the like can be mentioned.



Examples of the substituent of the substituted amino group include methyl group, ethyl group, isopropyl group, benzyl group, phenyl group, pyridyl group, methoxy group, acetyl group, trifluoroacetyl group, pivaloyl group, benzoyl
15 group, naphthoyl group, tert-butoxycarbonyl group, 2,2,2-trichloroethoxycarbonyl group, benzyloxycarbonyl group, 9-fluorenylmethyloxycarbonyl group, benzhydryl group, trityl group, phthaloyl group, allyloxycarbonyl group, methanesulfonyl group, trifluoromethanesulfonyl group, p-toluenesulfonyl group,
20 o-nitrobenzenesulfonyl group, trimethylsilylethoxycarbonyl group, dimethylcarbamoyl group and the like.

In the compound of the present invention, when the substituent is a group selected from alkyl group, halogenated alkyl group, alkoxy group and halogenated alkoxy group, or a
25 group having the selected group as a partial structure, the selected group is preferably a lower group.

[0038]

In compound (1) of the present invention, R^A in the α -chain is $-\text{CH}_2-\text{CZ}^1\text{Z}^2(\text{COX})$ or $-\text{CH}=\text{CZ}^1(\text{COX})$. That is, compound (1)
30 is a compound represented by the following formula ($\alpha 1$) or the following formula ($\alpha 2$).

[0039]







[0040]

Both Z^1 and Z^2 in compound (α1) are preferably hydrogen atoms or fluorine atoms.

5 The configuration of the group bonded to the double bond of the $CH=CZ^1(COX)$ moiety of compound (α2) may be E or Z. When Z^1 is a hydrogen atom, E is preferable, and when Z^1 is a fluorine atom, Z is preferable.

Furthermore, compound (α1) is preferably compound (α1-1) or compound (α1-2), and compound (α2) is preferably compound
 10 (α2-1) or (α2-2).

[0041]



[0042]

In the aforementioned formulas, X is OR⁴ or NR⁵R⁶, preferably OR⁴. R⁴ of OR⁴ is preferably a hydrogen atom, an
 5 alkyl group having 1 - 4 carbon atoms or a substituted alkyl group having 1 - 4 carbon atoms. As the substituent, hydroxy group, acyl group, acyloxy group, alkoxy carbonyloxy group, substituted amino group, aryl group, heterocyclic group and the like can be mentioned, and acyl group and heterocyclic group
 10 are particularly preferable.

A compound in which X is OR⁴ and R⁴ is an alkyl group

having 1 - 6 carbon atoms or a substituted alkyl group having 1 - 6 carbon atoms can also be led from a compound in which the COX moiety (i.e., CO₂R⁴ moiety) has a structure that produces a prodrug of carboxylic acid.

5 Specific examples of R⁴ include hydrogen atom, methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, isobutyl group, sec-butyl group, tert-butyl group, benzyl group, pivaloylmethyl group, (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl group, 2-hydroxyethyl group, 2-(dimethylamino)ethyl
10 group, 2-(morpholino)ethyl group, 4-pyridylmethyl group, pivaloyloxymethyl group, 1-[(1-cyclohexyloxy)carbonyl]oxy)ethyl group and the like. Preferred are hydrogen atom, alkyl group having 1 - 4 carbon atoms (e.g., methyl group and ethyl group), pivaloylmethyl group, and (5-
15 methyl-2-oxo-1,3-dioxol-4-yl)methyl group, and particularly preferred are hydrogen atom, methyl group and ethyl group.

 R⁵ when X is NR⁵R⁶ is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms or a substituted alkyl group having 1 - 6
- 6 carbon atoms. Examples of the alkyl group having 1 - 6
20 carbon atoms include methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, isobutyl group, sec-butyl group, tert-butyl group and the like. R⁵ is preferably a hydrogen atom, a methyl group, an ethyl group or the like.

[0043]

25 R⁶ is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms, a substituted alkyl group having 1 - 6 carbon atoms, an alkylsulfonyl group having 1 - 6 carbon atoms, a substituted alkylsulfonyl group having 1 - 6 carbon atoms, an arylsulfonyl group having 6 - 10 carbon atoms or a substituted arylsulfonyl
30 group having 6 - 10 carbon atoms.

 When R⁶ is an alkyl group having 1 - 6 carbon atoms or a substituted alkyl group having 1 - 6 carbon atoms, examples thereof include lower alkyl group such as methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, isobutyl
35 group, sec-butyl group, and tert-butyl group, hydroxyalkyl

group such as 2-hydroxyethyl group, 3-hydroxypropyl group, 2-hydroxypropyl group, and 2,3-dihydroxypropyl group, and the like.

When R^6 is an alkylsulfonyl group having 1 - 6 carbon atoms or a substituted alkylsulfonyl group having 1 - 6 carbon atoms, examples thereof include methanesulfonyl group, ethanesulfonyl group and the like. Examples of the arylsulfonyl group having 6 - 10 carbon atoms or a substituted arylsulfonyl group having 6 - 10 carbon atoms include phenylsulfonyl group, p-toluenesulfonyl group and the like.

R^6 is preferably a hydrogen atom, a methyl group, an ethyl group, a hydroxyethyl group, a methanesulfonyl group, an ethanesulfonyl group or the like.

The structure of the moiety expressed by A-B shows that the A-B structure is a carbon-carbon single bond (i.e., $-\text{CH}_2-\text{CH}_2-$), a carbon-carbon double bond (i.e., $-\text{CH}=\text{CH}-$), or a carbon-carbon triple bond (i.e., $-\text{C}\equiv\text{C}-$). The structure of the A-B moiety is preferably a carbon-carbon single bond or a carbon-carbon double bond, and the configuration of the group bonded to the double bond is preferably E.

The hydroxy group bonded by a wavy line shows that the orientation of the bond of the hydroxy group is α -configuration (binding downward on the paper surface), β -configuration (binding upward on the paper surface) or a mixed configuration of α -configuration and β -configuration. The configuration of the hydroxy group is preferably α -configuration, or a mixed configuration of α -configuration and β -configuration, particularly preferably α -configuration.

Y is CH_2 , S or O, preferably CH_2 or S, particularly preferably CH_2 .

In the structure shown by $(\text{CH}_2)_n$, n is the number of methylene units. When n is 0, the methylene unit is absent. In this case, the carbon atom to which R^1 and R^2 are bonded, and the carbon atom of the triple bond are directly bonded to each other. n is preferably 1 or 2, particularly preferably 1.

[0044]

R¹ and R² are each independently a hydrogen atom, an alkyl group having 1 - 3 carbon atoms or a substituted alkyl group having 1 - 3 carbon atoms. It is preferable that one of
5 R¹ and R² is a hydrogen atom and the other is an alkyl group having 1 - 2 carbon atoms (preferably a methyl group), or R¹ and R² are alkyl groups having 1 - 2 carbon atoms (preferably a methyl group). It is more preferable that R¹ is a hydrogen atom and R² is an alkyl group having 1 - 2 carbon atoms, and
10 particularly preferably, R¹ is a hydrogen atom and R² is a methyl group.

[0045]


R³ is an alkyl group having 1 - 4 carbon atoms, a substituted alkyl group having 1 - 4 carbon atoms, a cycloalkyl
15 group having 3 - 6 carbon atoms, a substituted cycloalkyl group having 3 - 6 carbon atoms, an aryl group having 6 - 10 carbon atoms or a substituted aryl group having 6 - 10 carbon atoms. It is preferably an alkyl group having 1 - 4 carbon atoms or a cycloalkyl group having 3 - 6 carbon atoms, particularly
20 preferably an alkyl group having 2 - 4 carbon atoms or a cycloalkyl group having 3 - 6 carbon atoms, further preferably an alkyl group having 2 - 3 carbon atoms or a cycloalkyl group having 3 - 5 carbon atoms, especially preferably a cycloalkyl group having 3 - 5 carbon atoms. Specific examples of R³
25 include ethyl group, n-propyl group, isopropyl group, cyclopropyl group, cyclobutyl group, cyclopentyl group, phenyl group and the like. Ethyl group, cyclopropyl group or cyclobutyl group is preferable, and cyclopropyl group is particularly preferable.

30 [0046]

In compound (1) of the present invention, a compound wherein ω -chain has the following structure is preferable.

[Compound (ω -1)]

[0047]



[0048]

The symbols in the formula (ω-1) have the same meanings as in the formula (1).


5 It is preferable that Y is CH₂ or S, R¹ is a hydrogen atom, R² is an alkyl group having 1 - 2 carbon atoms, R³ is an alkyl group having 1 - 4 carbon atoms (preferably an alkyl group having 2 - 3 carbon atoms) or a cycloalkyl group having 3 - 6 carbon atoms (preferably a cycloalkyl group having 3 - 5
10 carbon atoms), X is OR⁴, R⁴ is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms or a substituted alkyl group having 1 - 6 carbon atoms. Note that 7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4RS)-3-hydroxy-4-methyloct-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoic acid is excluded from compound (ω-1).
15 Furthermore, the substituent of the substituted alkyl group having 1 - 6 carbon atoms is preferably a group selected from hydroxy group, acyl group, acyloxy group, alkoxycarbonyloxy group, substituted amino group, aryl group and heterocyclic group.

20 [0049]

As compound (1) of the present invention, a compound wherein the structure of α-chain and ω-chain is the following structure is preferable from the aspects of pharmacological activity and property.

25 [Compound (ω-10)]

[0050]



[0051]

The symbols in the formula (ω-10) have the same meanings as in the formula (1). R³⁰ is an alkyl group having 1 - 4
 5 carbon atoms (preferably an alkyl group having 2 - 3 carbon atoms), or a cycloalkyl group having 3 - 6 carbon atoms (preferably a cycloalkyl group having 3 - 5 carbon atoms).


[0052]

It is preferable that R¹ is a hydrogen atom, R² is an
 10 alkyl group having 1 - 2 carbon atoms, X is OR⁴, and R⁴ is a hydrogen atom or an alkyl group having 1 - 4 carbon atoms.

[0053]

[Compound (ω-11)]

[0054]



15

[0055]

The symbols in the formula (ω-11) have the same meanings as in the formula (1). R³⁰ is an alkyl group having 1 - 4
 20 carbon atoms (preferably an alkyl group having 2 - 3 carbon atoms), or a cycloalkyl group having 3 - 6 carbon atoms (preferably a cycloalkyl group having 3 - 5 carbon atoms).

[0056]

It is preferable that R¹ is a hydrogen atom, R² is an
 25 alkyl group having 1 - 2 carbon atoms, X is OR⁴, and R⁴ is a hydrogen atom or 1 - 4 carbon. Both Z¹ and Z² are preferably

hydrogen atoms or fluorine atoms.


[0057]

Preferable compounds (1) are shown below.

[0058]

5 [Compound (1)-A]

[0059]




[0060]

In the formula (1)-A, the configuration of a hydroxy
10 group bonded by a wavy line is α -configuration, β -configuration,
or a mixed configuration of α -configuration and β -configuration,
Y is CH_2 , S or O, R^1 is preferably a hydrogen atom, R^2 is
preferably an alkyl group having 1 - 2 carbon atoms, R^3 is
preferably an alkyl group having 1 - 4 carbon atoms or a
15 cycloalkyl group having 3 - 6 carbon atoms, R^A is $-\text{CH}_2-$
 $\text{CZ}^1\text{Z}^2(\text{COX})$ or $-\text{CH}=\text{CZ}^1(\text{COX})$, Z^1 and Z^2 are each independently a
hydrogen atom or
a fluorine atom, X is preferably OR^4 , and R^4 is preferably a
hydrogen atom, an alkyl group having 1 - 6 carbon atoms, or an
20 alkyl group having 1 - 6 carbon atoms substituted by a
substituent(s) selected from the group consisting of a hydroxy
group, an acyl group, an acyloxy group, an alkoxy carbonyloxy
group, a substituted amino group, an aryl group and a
heterocyclic group. When R^A is $-\text{CH}=\text{CZ}^1(\text{COX})$, the configuration
25 of the group bonded to the double bond may be E or Z, when Z^1
is a hydrogen atom, E is preferable, and when Z^1 is a fluorine
atom, Z is preferable. Note that 7-((1R,2R,3R)-3-hydroxy-2-
((1E,3S,4RS)-3-hydroxy-4-methyloct-1-en-6-yn-1-yl)-5-
oxocyclopentyl)heptanoic acid is excluded.

30 [0061]

[Compound (1)-B]

[0062]




[0063]

5 In the formula (1)-B, the configuration of a hydroxy group bonded by a wavy line is α -configuration, β -configuration, or a mixed configuration of α -configuration and β -configuration, Y is preferably CH₂ or S, R¹ is preferably a hydrogen atom, R² is preferably an alkyl group having 1 - 2 carbon atoms, R³ is
10 preferably an alkyl group having 2 - 3 carbon atoms or a cycloalkyl group having 3 - 5 carbon atoms, R^A is -CH₂-CZ¹Z²(COX) or -CH=CZ¹(COX), Z¹ and Z² are each independently a hydrogen atom or a fluorine atom, X is preferably OR⁴, and R⁴ is preferably a hydrogen atom, an alkyl group having 1 - 6
15 carbon atoms, or an alkyl group having 1 - 6 carbon atoms substituted by a substituent(s) selected from the group consisting of a hydroxy group, an acyl group, an acyloxy group, an alkoxy carbonyloxy group, a substituted amino group, an aryl group and a heterocyclic group. When R^A is -CH=CZ¹(COX), the
20 configuration of the group bonded to the double bond may be E or Z, when Z¹ is a hydrogen atom, E is preferable, and when Z¹ is a fluorine atom, Z is preferable.

[0064]

[Compound (1)-C]

25 [0065]




[0066]

In the formula (1)-C, the configuration of a hydroxy group bonded by a wavy line is α -configuration, β -configuration, or a mixed configuration of α -configuration and β -configuration, R^1 is preferably a hydrogen atom, R^2 is preferably an alkyl group having 1 - 2 carbon atoms, R^3 is preferably an alkyl group having 2 - 3 carbon atoms, Z^1 is a hydrogen atom or a fluorine atom, X is preferably OR^4 , and R^4 is preferably a hydrogen atom or an alkyl group having 1 - 4 carbon atoms.

[0067]

[Compound (1)-D]

[0068]




[0069]

In the formula (1)-D, the configuration of a hydroxy group bonded by a wavy line is α -configuration, β -configuration, or a mixed configuration of α -configuration and β -configuration, R^1 is preferably a hydrogen atom, R^2 is preferably an alkyl group having 1 - 2 carbon atoms, R^3 is preferably a cycloalkyl group having 3 - 5 carbon atoms, Z^1 is a hydrogen atom or a fluorine atom, X is preferably OR^4 , and R^4 is preferably a hydrogen atom or an alkyl group having 1 - 4 carbon atoms.

[0070]

[Compound (1)-E]

[0071]




[0072]

In the formula (1)-E, the configuration of a hydroxy
5 group bonded by a wavy line is α -configuration, β -configuration,
or a mixed configuration of α -configuration and β -configuration,
 R^1 is preferably a hydrogen atom, R^2 is preferably an alkyl
group having 1 - 2 carbon atoms, R^3 is preferably an alkyl
group having 2 - 3 carbon atoms, Z^1 and Z^2 are each
10 independently a hydrogen atom or a fluorine atom, X is
preferably OR^4 , and R^4 is preferably a hydrogen atom or an
alkyl group having 1 - 4 carbon atoms.

[0073]

[Compound (1)-F]

15 [0074]



[0075]

In the formula (1)-F, the configuration of a hydroxy
group bonded by a wavy line is α -configuration, β -configuration,
20 or a mixed configuration of α -configuration and β -configuration,
 R^1 is preferably a hydrogen atom, R^2 is preferably an alkyl
group having 1 - 2 carbon atoms, R^3 is preferably a cycloalkyl
group having 3 - 5 carbon atoms, Z^1 and Z^2 are each
independently a hydrogen atom or a fluorine atom, X is
25 preferably OR^4 , and R^4 is preferably a hydrogen atom or an

alkyl group having 1 - 4 carbon atoms.

[0076]

Particularly preferable examples of compound (1) include the following compounds.

- 5 Methyl 4-((2-((1R,2R,3R)-3-hydroxy-2-((1E,3RS,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)ethyl)thio)butanoate [compound (1)-1],
methyl (2E)-4-((2-((1R,2R,3R)-3-hydroxy-2-((1E,3RS,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)ethyl)thio)but-2-enoate [compound (1)-2],
10 methyl ((1R,2R,3R)-3-hydroxy-2-((1E,3RS,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoate [compound (1)-3],
(5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoate [compound (1)-4],
15 3,3-dimethyl-2-oxobutyl 7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoate [compound (1)-5],
20 7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoic acid [compound (1)-6],
(2E)-7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)hept-2-enoic acid
25 [compound (1)-7],
2,2-difluoro-7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoic acid [compound (1)-8],
7-((1R,2R,3R)-2-((1E,3S,4S)-7-cyclopropyl-3-hydroxy-4-methylhept-1-en-6-yn-1-yl)-3-hydroxy-5-oxocyclopentyl)heptanoic acid [compound (1)-9], and
30 (2E)-7-((1R,2R,3R)-2-((1E,3S,4S)-7-cyclopropyl-3-hydroxy-4-methylhept-1-en-6-yn-1-yl)-3-hydroxy-5-oxocyclopentyl)hept-2-enoic acid [compound (1)-10].

35 [0077]

The compound of the present invention may be a pharmaceutically acceptable salt of compound (1). The salt is a salt of a carboxylic acid moiety with a basic substance (nontoxic inorganic base or nontoxic organic base), and has a structure in which the carboxylic acid moiety is -COO^- and the hydrogen atom of the carboxylic acid is a cation.

The pharmaceutically acceptable salt is, for example, a salt derived from a nontoxic inorganic base or a salt derived from a nontoxic organic base, and a salt derived from a nontoxic inorganic base is preferable.

Examples of the base derived from an inorganic salt include sodium salt, potassium salt, calcium salt, magnesium salt, zinc salt, aluminum salt, ammonium salt and the like, as well as lithium salt, copper salt, ferric salt, ferrous salt, manganese salt, manganous salt and the like, and sodium salt, potassium salt, calcium salt, magnesium salt and ammonium salt are preferable, and sodium salt and potassium salt are particularly preferable.

Examples of the salt derived from an organic base include salts with primary amine, secondary amine, tertiary amine, substituted amine of these (including naturally present substituted amine), cyclic amine, basic amino acid, basic ion exchange resin and the like. Examples of the organic amine and amino acid include isopropylamine, diethylamine, triethylamine, trimethylamine, tripropylamine, ethylenediamine, N,N' -dibenzylethylenediamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, morpholine, N -ethyl-morpholine, piperazine, piperidine, N -ethylpiperidine, betaine, caffeine, choline, glucamine, glucosamine, histidine, hydrabamine, methylglucamine, lysine, arginine, polyamine resin, procaine, purine, theobromine and the like.

Compound (1) or a pharmaceutically acceptable salt thereof may take the form of a hydrate or solvate.

[0078]

Compound (1) of the present invention may be derived from

a pharmaceutically acceptable prodrug of compound (1). The pharmaceutically acceptable prodrug is a compound having a group converted to a hydroxy group, a carboxy group or the like by solvolysis or under physiological conditions. Examples of
5 the group forming a prodrug include the groups described in Prog. Med., 1985, vol. 5, pp. 2157-2161, and "IYAKUJIN no KAIHATSU (Development of Pharmaceuticals)", Vol. 7, Design of Molecules, pp. 163-198, Published by HIROKAWA SHOTEN (1990).
[0079]

10 A prodrug of compound (1) is preferably a compound converted to a compound by a reaction due to an enzyme, gastric acid and the like in the body.

As the compound, the following compound (1-1) - compound (1-4) can be mentioned.

15 Compound (1-1) wherein the hydroxy group in compound (1) is acylated, alkylated, phosphorylated or borated. Specifically, a compound wherein the hydroxy group in compound (1) is acetylated, propanoylated, butanoylated, pivaloylated, oleylated, palmitoylated, succinylated, fumarylated, alanylated,
20 or dimethylaminomethylcarbonylated.

Compound (1-2) wherein the carboxy group in compound (1) is esterified or amidated. Specifically, a compound wherein the carboxy group in compound (1) is phenyl-esterified or phthalidyl-esterified, and the like.

25 Compound (1-3) wherein the carboxy group in compound (1) is substituted by a hydroxymethyl group.

Compound (1-4) wherein the carbonyl group of the 5-membered ring moiety in compound (1) is enol-esterified.

These compound (1-1) - compound (1-4) can be produced by
30 a known method. Compound (1-1) - compound (1-4) may be a hydrate or a non-hydrate.

[0080]

Compound (1) of the present invention or a pharmaceutically acceptable salt thereof may take the form of a
35 clathrate compound of cyclodextrin. As the cyclodextrin, α -

cyclodextrin, β -cyclodextrin, γ -cyclodextrin, sulfobutylether-
 β -cyclodextrin, hydroxypropyl- β -cyclodextrin, or a mixture of
one or more kinds selected from these is used. As the
cyclodextrin, α -cyclodextrin or β -cyclodextrin is preferable,
5 and α -cyclodextrin is particularly preferable.

As the production method of a cyclodextrin clathrate
compound, methods generally known widely as emulsion method,
saturated aqueous solution method, kneading method, mixing and
pulverizing method and the like can be used. Specifically, it
10 can be produced using compound (1) or a pharmaceutically
acceptable salt thereof, and cyclodextrin and by the method
described in JP-B-50-3362, JP-B-2-31404 or JP-B-61-52146. By
formulating compound (1) as a cyclodextrin clathrate compound,
properties such as stability and water solubility increase, and
15 properties preferable as pharmaceutical products can be
imparted.

[0081]

(Production method of the compound of the present invention)

The compound of the present invention can be synthesized
20 by a general PG synthesis method. The concept of the synthetic
route is shown in the following formulas.


For example, a compound wherein the -A-B- moiety is a
double bond can be synthesized using a compound in which α -
chain is introduced into Corey lactone as a starting material,
25 by reducing the 15-position carbonyl group of the α, β -
unsaturated carbonyl moiety in the ω -chain (compound of the
following formula wherein W is H) introduced by a Horner-
Wadsworth-Emmons reaction, introducing a double bond into the
 α -chain when desired, oxidizing the 9-position and deprotecting
30 the hydroxy-protecting group.

A PG derivative having fluorine at the 2-position can be
synthesized by, for example, synthesizing the starting material
by introducing α -chain having a difluoro unit into Corey
lactone, and performing a reaction similar to the above-
35 mentioned reaction.

A compound wherein the -A-B- moiety is a single bond can be synthesized by performing 1,4-reduction of the α,β -unsaturated carbonyl moiety in the ω -chain (W in the following formula is H) introduced by a Horner-Wadsworth-Emmons reaction using Stryker's reagent ($[(\text{Ph}_3\text{P})\text{CuH}]_6$), and a reducing agent
 5 such as sodium dithionite.

A compound wherein the -A-B- moiety is a triple bond can be synthesized by reducing, in the presence of a bromination agent such as N-bromosuccinimide and 1,3-dibromo-5,5-
 10 dimethylhydantoin, the 15-position carbonyl group of the α -bromo- α,β -unsaturated carbonyl moiety in the ω -chain (W in the following formula is a bromine atom) introduced by a Horner-Wadsworth-Emmons reaction, and performing a dehydrobromination reaction using sodium hydroxide, potassium tert-butoxide, or a
 15 base such as cesium acetate/18-crown-6.

[0082]



[0083]

In the formulas, R^7 , R^8 and R^9 are each independently a

hydroxy-protecting group, W is a hydrogen atom or a bromine atom, and each of other symbols is as defined above.

[0084]


As the hydroxy-protecting group for $R^7 - R^9$, the hydroxy-protecting groups described in "Greene's/Protective/Groups/in/Organic/Synthesis 4th Edition" (by P. G. M. Wuts, T. W. Greene, J. Wiley & Sons, 2007), pp. 16-430 and the like can be used. Specifically, acyl group, tri-organosilyl group, alkoxyalkyl group, a monovalent group having a cyclic ether structure and the like can be mentioned. As the acyl group, acetyl group, benzoyl group, chloroacetyl group, dichloroacetyl group, trichloroacetyl group, trifluoroacetyl group and the like are preferable. As the tri-organosilyl group, a group in which three of alkyl group, aryl group, aralkyl group and alkoxy group are bonded to the silicon atom is preferable. Specifically, for example, tert-butyldimethylsilyl group, tert-butyldiphenylsilyl group, trimethylsilyl group, triethylsilyl group, triphenylsilyl group, triisopropylsilyl group and the like are preferable. As the alkoxyalkyl group, methoxymethyl group, benzyloxymethyl group, tert-butoxymethyl group, 2-methoxyethoxymethyl group, 1-ethoxyethyl group, 1-methyl-1-methoxyethyl group and the like are preferable. As the monovalent group having a cyclic ether structure, tetrahydropyranyl group, tetrahydrofuranyl group and the like are preferable. Acetyl group, benzoyl group, tetrahydropyranyl group, tert-butyldimethylsilyl group, tert-butyldiphenylsilyl group and the like are particularly preferable.

The hydroxy-protecting groups can be deprotected easily by a conventional method. Specifically, for example, they can be deprotected by the methods described in "Greene's/Protective/Groups/in/Organic/Synthesis 4th Edition" (by P. G. M. Wuts, T. W. Greene, J. Wiley & Sons, 2007), pp. 16-430 and the like.

[0085]

A PG derivative in which the 5-position (Y) of the α -chain is a hetero atom can be synthesized as shown in the following by using Corey lactone that constructed ω -chain as a starting material, reducing the lactone moiety to give diol, introducing α -chain, oxidizing the 9-position, and deprotecting other hydroxy-protecting groups.

[0086]



[0087]

10 In the formulas, each symbol is as defined above.

[0088]

Representative examples of the production method of compound (1) can be shown by the following formulas.


In the following formulas, the following abbreviations
15 are used.

Me: methyl

Ac: acetyl

THP: tetrahydro-2H-pyran-2-yl

[0089]



[0090]

5 The starting material in the aforementioned formulas is aldehyde (A-1) described in JP-B-60-36422 and the like, and α,β -unsaturated ketone (A-3) is obtained from aldehyde (A-1) and phosphonate (A-2) by the Horner-Wadsworth-Emmons reaction. After reduction of the 15-position carbonyl group, deprotection

10 is performed to give diol (A-4). After protection of the hydroxy group, hydrolysis is performed to give carboxylic acid (A-5). The unprotected hydroxy group on the 5-membered ring is oxidized and deprotected, whereby compound (A-6) of the present invention (compound (1)-9) can be synthesized.

15 [0091]

After protection of two hydroxy groups of compound (A-4), deacetylation is performed to give alcohol (A-7). The α -position of the ester is phenylselenenylated, an elimination reaction with hydrogen peroxide water is performed and ester

20 hydrolysis is performed to give α,β -unsaturated carboxylic acid (A-8). The unprotected hydroxy group on the 5-membered ring is

oxidized and deprotected, whereby compound (A-9) of the present invention (compound (1)-10) can be obtained.

For esterification of the carboxy group of compound (A-6) or (A-9) of the present invention, a general method, for example, the method described in "Shinjikken Kagaku Koza (Courses in Experimental Chemistry) 14 synthesis and reaction of organic compound (II)" (Maruzen Company, Limited) can be used. For example, methods such as esterification by condensation with alcohols and phenols, esterification by an O-alkylating agent, esterification using alkenes and alkynes, esterification by dialkyl sulfate and halogenated hydrocarbons and the like are used. For conversion to acylamides and sulfonamides, for example, the method of Tithereley et al. (J. Chem. Soc., 1904, vol. 85, p. 1673), the method of Lynch et al. (Can. J. Chem., 1972, vol. 50, p. 2143), the method of Davidson et al. (J. Am. Chem. Soc., 1958, vol. 80, p. 376) and the like can be adopted. In addition, a method including converting carboxylic acid to acid halide or active ester, then a condensation reaction directly with amides or sulfonamides, or a reaction with amines to once convert to amides, followed by acylation or sulfonylation and the like is used.

[0092]

The compound of the present invention has an asymmetric carbon in the structure thereof. Thus, various stereoisomers and optical isomers are present. The present invention encompasses all stereoisomers, optical isomers, and mixtures thereof.

[0093]

The compound of the present invention is a derivative less susceptible to metabolism in vivo and having improved stability. Compound (1) has an alkyl group at the 16-position carbon atom, and therefore, the hydroxy group at the 15-position of the PG skeleton is less susceptible to metabolism by PG dehydrogenase and shows high metabolic stability. Thus, it has a long half-life in blood and can maintain effective

blood concentration for a long time as compared to natural type PG. Having improved metabolic stability, the compound can improve bioavailability of drugs.

[0094]

5 The compound of the present invention has an alkynyl group in the ω -chain and shows very potent actions such as platelet aggregation suppressive activity. In particular, a derivative having a cycloalkyl group on the end of the alkynyl group is stable and exhibits potent activity.

10 [0095]

 The compound of the present invention and a cyclodextrin clathrate compound thereof are useful as medicaments, and can show a superior effect as a drug for circulatory diseases, central nervous system diseases, inflammatory diseases, pain
15 and the like. Specifically, the circulatory diseases include peripheral circulation disorder, Buerger's disease, Raynaud's disease, angina pectoris, myocardial infarction, cardiac failure, pulmonary hypertension, pulmonary obliteration, diabetes, cerebral infarction, cerebral thrombus, deafness,
20 Meniere's disease and the like. The central nervous system diseases include sleeplessness, anxiety, depression, schizophrenia, dementia and the like. The inflammatory diseases include acute hepatitis, chronic hepatitis, cirrhosis, cholecystitis, cholangitis, acute pancreatitis, chronic
25 pancreatitis, chronic peritonitis, acute peritonitis, cystitis, acute nephritis, chronic nephritis, encephalitis, polyneuritis, meningitis, myelitis, arthritis, rheumatoid arthritis, bronchitis, pneumonia, pleurisy, phlebitis, pericarditis, rhinitis, pharyngitis, labyrinthitis, otitis externa and the
30 like.

[0096]

 A medicament containing the compound of the present invention or a cyclodextrin clathrate compound thereof (hereinafter to be indicated as "the medicament of the present
35 invention") is particularly useful as a medicament for the

prophylaxis or treatment of blood flow disorders. The medicament of the present invention may be a medicament containing the compound of the present invention and optionally containing a pharmaceutically acceptable carrier and other
5 therapeutic components.

[0097]

When the medicament of the present invention is administered to patients as an agent for the prophylaxis or treatment of the aforementioned diseases, the daily dose varies
10 depending on the age and body weight of patients, pathology and severity in patients, and the like. It is generally 0.00001 mg to 1 mg, preferably 0.0001 mg to 0.3 mg, as compound (1), which is desirably administered in one to several portions. For example, 0.0001 mg to 0.3 mg, particularly 0.001 mg to 0.1 mg,
15 is preferable for oral administration. Meanwhile, 0.00001 mg to 0.1 mg, particularly 0.0001 mg to 0.03 mg, is preferable for intravenous administration. The dose can be appropriately increased or decreased according to the disease or pathology, and continuous drip administration is sometimes desirable in
20 the case of injection.

[0098]

The medicament of the present invention can be administered to the body by oral administration or parenteral administration (e.g., blood vessel (intravenous and
25 intraarterial) administration, intrarectal administration, and intraarticular administration). Examples of the administration form include oral administration forms such as tablet (including orally disintegrating tablet and sublingual tablet), film (including oral films), capsule (including soft capsule
30 and microcapsule), granule, powder, troche, and syrup, and parenteral administration forms such as liquid injection such as solution, emulsion, and suspension (e.g., subcutaneous injection, intravenous injection, intramuscular injection, intraperitoneal injection, and drip infusion), suppository
35 (e.g., rectal suppository and vaginal suppository), eye drop,

nasal drop, external preparation (e.g., transdermal preparation such as adhesive preparation, ointment, and cream), and pulmonary preparation (inhalant etc.). These preparations may be controlled-release preparations such as rapid release
5 preparation and sustained-release preparation.

[0099]

The preparation in the aforementioned administration form can be produced by adding additives necessary for preparations such as general carrier, excipient, lubricant, disintegrant,
10 binder, and stabilizer to the medicament of the present invention and formulating the resulting mixture by a conventional method. For example, when the preparation is a tablet, granule, powder or the like, it can be produced using any pharmaceutical carriers suitable for producing a solid
15 preparation, for example, excipient, lubricant, disintegrant, and binder.

[0100]

Examples of the excipient include lactose, glucose, fructose, maltose, isomerized lactose, reduced lactose, lactose
20 crystal, saccharose, D-mannitol, erythritol, maltitol, xylitol, palatinose, trehalose, sorbitol, cornstarch, potato starch, wheat starch, rice starch, crystalline cellulose, dextrans (e.g., dextran, dextran 40, and dextran 70), pullulan, dextrin, talc, silicic anhydride, anhydrous calcium phosphate,
25 precipitated calcium carbonate, calcium silicate and the like.

[0101]

Examples of the lubricant include magnesium stearate, calcium stearate, sucrose fatty acid ester, sodium stearyl fumarate, stearic acid, talc, polyethylene glycol, beeswax and
30 the like.

[0102]

Examples of the disintegrant include starch, carboxymethylcellulose, hydroxypropylcellulose, carmellose, carmellose calcium, sodium carboxymethyl starch, croscarmellose
35 sodium, hydroxypropylstarch and the like.

[0103]

Examples of the binder include crystalline cellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, hydroxyethylcellulose, methylcellulose, sodium
5 carboxymethylcellulose, partially pregelatinized starch, pregelatinized starch, sodium alginate, pullulan, gum arabic, dextrin, polyvinylpyrrolidone, poly(vinyl alcohol), gelatin and the like.

[0104]

10 A preparation in the aforementioned administration form may contain inert diluent, lubricant, stabilizer, solubilizing agent, corrigent and the like as necessary. Examples of the inert diluent include calcium carbonate, sodium carbonate, lactose, calcium phosphate, sodium phosphate and the like.
15 Examples of the lubricant include magnesium stearate, stearic acid, colloidal silica, talc and the like. Examples of the stabilizer include ascorbic acid, sodium sulfite, glycine, L-aspartic acid, tocopheryl acetate, β -cyclodextrin, fumaric acid and the like. Examples of the solubilizing agent include
20 polyethylene glycol, propylene glycol, glutamic acid, aspartic acid and the like. Examples of the corrigent include citric acid, ascorbic acid, lactic acid, acetic acid, tartaric acid, malic acid, aspartame, acesulfame potassium, thaumatin, saccharin sodium, dipotassium glycyrrhetinate, sodium glutamate,
25 sodium 5'-inosinate, sodium 5'-guanylate and the like.

[0105]

These tablets may be uncoated, or coated by a known technique to delay disintegration and absorption in the gastrointestinal tract, thereby ensuring a sustained action for
30 a longer time. For example, a time delay material, for example, glyceryl monostearate or glyceryl distearate may be used.

[0106]

The medicament of the present invention may be provided as a hard gelatin capsule containing a mixture with an inert
35 solid diluent, for example, calcium carbonate, sodium carbonate,

lactose, calcium phosphate, sodium phosphate or kaolin. In addition, the medicament of the present invention may be provided as a soft gelatin capsule containing a mixture with water-miscible solvent, for example, alcohols such as propylene glycol, polyethylene glycol, glycerol, and ethanol and an oil medium.

[0107]

Examples of the oil medium include fatty acid ester, liquid paraffin, vegetable oil such as peanut oil and olive oil and the like. Fatty acid ester is a compound in which a carboxy group of fatty acid forms an ester bond with alcohol and, specifically, glyceride optionally having a saturated or unsaturated branched chain, fatty acid ester composed of an ester compound of fatty acid and monohydric alcohol and the like can be mentioned. Preferable fatty acid is medium-chain or long-chain fatty acid having 6 - 24 carbon atoms, and caproic acid, caprylic acid, capric acid, lauric acid, myristic acid, palmitic acid, palmitoleic acid, stearic acid, oleic acid, linoleic acid, linolenic acid, ricinoleic acid, arachidonic acid and the like can be mentioned. Preferable alcohols include monohydric alcohols having 1 - 6 carbon atoms, polyols which are polyhydric alcohols such as glycerol, polyethylene glycol, and propylene glycol and the like. Preferable fatty acid esters include glyceride optionally having saturated or unsaturated branch, glycerol fatty acid ester, and propyleneglycol fatty acid ester. In addition, a mixture of two or more kinds of glycerides can also be used. The mixtures of glycerides include a mixture of caprylic acid triglyceride and capric acid triglyceride, vegetable oil such as castor oil, corn oil, olive oil, sesame oil, rape seed oil, salad oil, cottonseed oil, camellia oil, peanut oil, palm oil, and sunflower oil. Examples of the fatty acid ester composed of an ester compound of fatty acid and monohydric alcohol include isopropyl myristate, isopropyl palmitate, ethyl linoleate, ethyl oleate and the like.

[0108]

The major component of the coating film of the capsule is not particularly limited and, for example, various known components such as gelatin, carrageenan, pectin, pullulan, 5 sodium alginate, starch, hypromellose, and hydroxypropylcellulose can be mentioned.

[0109]

Preferable examples of the plasticizer used for the production of the coating film of soft capsules include 10 polyhydric alcohol such as glycerol, ethylene glycol, polyethylene glycol, propylene glycol, and polypropylene glycol, sugar alcohol such as sorbitol, mannitol, xylitol, maltitol, cornstarch-derived sugar alcohol solution, and reduced maltose syrup, and the like. Two or more kinds of these plasticizers 15 may be used in combination.

[0110]

When the preparation is liquid such as syrup, solution, emulsion, and suspension, for example, it can be produced by appropriately selecting stabilizer, suspending agent, corrigent, 20 aromatic, soothing agent and the like. When injection is produced, it can be aseptically prepared by dissolving the active ingredient together with a pH adjuster such as hydrochloric acid, sodium hydroxide, sodium lactate, acetic acid, sodium hydrogen phosphate, and sodium dihydrogen 25 phosphate; an isotonicity agent such as sodium chloride, glucose, D-sorbitol, glycerol, and D-mannitol; a surfactant such as stearylethanolamine, sodium lauryl sulfate, lauryl aminopropionate, lecithin, benzalkonium chloride, and glycerol monostearate; a suspending agent such as poly(vinyl alcohol), 30 polyvinylpyrrolidone, sodium carboxymethylcellulose, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, and hydroxypropylcellulose; and a soothing agent such as benzyl alcohol in distilled water for injection. It may also be prepared using an inert non-aqueous diluent such as propylene 35 glycol, polyethylene glycol, olive oil, ethanol, and

polysorbate 80. Furthermore, injection to be dissolved when in use may be prepared by adding mannitol, dextrin, cyclodextrin, gelatin and the like and freeze-drying the resulting mixture in vacuum. For stabilization and improving lesion reachability, a liposome preparation may also be prepared by a known method and used as an injection. What is called a lipoid preparation (fat emulsion) in which the compound of the present invention is dissolved in fine fat emulsion particles can also be prepared, and it can be produced by mixing an oil component such as fatty acid glycerides; phospholipid such as egg-yolk lecithin and soybean lecithin; and additives such as emulsion adjuvant, stabilizer, polymer substance, and isotonicity agent as necessary.

[0111]

In addition, an intrarectal administration preparation may be prepared by using a base for suppository such as cacao butter, fatty acid triglyceride, fatty acid diglyceride, fatty acid monoglyceride, and polyethylene glycol. Furthermore, an intrarectally injecting ointment can also be prepared by adjusting to a suitable viscosity with a water-soluble base such as polyethylene glycol, polypropylene glycol, glycerol, and glycerolgelatin, an oily base such as white petrolatum, hard fat, paraffin, liquid paraffin, plastibase, lanolin, and purified lanolin, and the like.

[0112]

The medicament of the present invention can be administered topically to the skin or mucosa, i.e., transdermally or transmucosally. Examples of the general dosage form for this object include gel, hydrogel, lotion, solution, cream, ointment, dusting powder, dressing agent, foam preparation, film, skin patch, wafer, implant, sponge, fiber, bandage, microemulsion and the like. Examples of the general carrier include alcohol, water, mineral oil, liquid paraffin, white petrolatum, glycerol, polyethylene glycol, propylene glycol and the like.

[0113]

The medicament can be directly administered to the eye or ear, in the form of an isotonic, pH controlled eye drop of a suspension of fine powder or solution in sterile saline. Other dosage forms suitable for ocular or aural administration include ointment, biodegradable implant, non-biodegradable implant, wafer, lens, microparticles such as liposome, and the like. Crosslinked polyacrylic acid, poly(vinyl alcohol), hyaluronic acid, cellulose polymers such as hydroxypropylmethyl cellulose, hydroxyethyl cellulose, and methyl cellulose, or polymers such as complex polysaccharides such as gellan gum can be mixed with a preservative such as benzalkonium chloride.

[0114]

To produce a pulmonary administration preparation, the medicament of the present invention is dissolved or dispersed in a conventional spray and filled in a pressure resistant container. It can also be pulmonarily administered as a liposome preparation by a known method.

[0115]

In order to use the medicament of the present invention in any of the aforementioned administration forms and improve solubility, dissolution rate, bioavailability and/or stability thereof, the medicament can be further mixed with cyclodextrin and an appropriate derivative thereof or a soluble polymer unit of polyethylene glycol-containing polymer and the like. For example, a mixture, a complex etc. of these have been confirmed to be generally useful for most dosage forms and administration routes. Both inclusion and non-inclusion complexes can be used. As an alternative method for directly forming a complex with a drug, cyclodextrin can also be used as an auxiliary additive, i.e., carrier, excipient or solubilizer. For these objects, α -cyclodextrin, β -cyclodextrin, hydroxypropyl- β -cyclodextrin, γ -cyclodextrin and the like are generally used.

[0116]

The compound of the present invention may be a

pharmaceutically acceptable salt of a compound represented by the formula (1). As the salt, those mentioned above can be recited.

[0117]

5 Other therapeutic components (hereinafter to be also referred simply to as "other drugs") to be contained in the medicament of the present invention are not particularly limited as long as they are drugs that supplement and/or enhance the prophylactic and/or therapeutic effect of the
10 compound of the present invention on blood flow disorders (e.g., spinal canal stenosis) and the like, drugs that can improve pharmacokinetics and absorption of the compound of the present invention and reduce the dose of the compound of the present invention, drugs that reduce the side effects of the compound
15 of the present invention and the like.

[0118]

Examples of other drugs for supplementing and/or enhancing the prophylactic and/or therapeutic effect of the compound of the present invention on blood flow disorders (e.g.,
20 spinal canal stenosis) and the like include, but are not limited to, prostaglandins, prostaglandin derivative preparations, vasodilators, anti-platelet drugs (platelet aggregation inhibitors), analgesic drugs, vitamins, muscle relaxants, antidepressants and the like.

25 [0119]

Examples of the prostaglandins include PG receptor agonist, PG receptor antagonist and the like. Examples of the PG receptor include PGE receptor (EP1, EP2, EP3, and EP4), PGD receptor (DP and CRTH2), PGF receptor (FP), PGI receptor (IP),
30 TX receptor (TP) and the like. Examples of the prostaglandin derivative preparation include limaprost, beraprost and the like.

[0120]

Examples of the vasodilator include angiotensin
35 converting enzyme inhibitor (e.g., captopril, enalapril, and

delapril), angiotensin II antagonist (e.g., candesartan
cilexetil, losartan, eprosartan, valsartan, telmisartan,
irbesartan, olmesartan medoxomil, tasosartan, 1-{{[2'-(2,5-
dihydro-5-oxo-4H-1,2,4-oxadiazol-3-yl)biphenyl-4-yl]methyl}-2-
5 ethoxy-1H-benzimidazole-7-carboxylic acid), calcium channel
blocker (e.g., manidipine, nifedipine, nicardipine, amlodipine,
and efonidipine), potassium channel opener (e.g., levcromakalim,
L-27152, AL0671, and NIP-121), clonidine and the like.
[0121]

10 Examples of the anti-platelet drug (platelet aggregation
inhibitor) include clopidogrel sulfate, ticlopidine
hydrochloride, cilostazol, ethyl icosapentate, sarpogrelate
hydrochloride and the like.
[0122]

15 Examples of the analgesic drug include acetaminophen,
non-steroidal antiinflammatory drugs, neuropathic pain
relieving drugs, pyrazolone drugs and the like. For example,
sodium salicylate, aspirin, aspirin-dialuminate formulation,
diflunisal, indomethacin, suprofen, ufenamate,
20 dimethylisopropylazulene, bufexamac, felbinac, diclofenac,
tolmetin sodium, clinoril, fenbufen, nabumetone, proglumetacin,
indomethacin farnesyl, acemetacin, proglumetacin maleate,
amfenac sodium, mofezolac, etodolac, ibuprofen, ibuprofen
piconol, naproxen, flurbiprofen, flurbiprofen axetil,
25 ketoprofen, fenoprofen calcium, tiaprofen, oxaprozin,
pranoprofen, loxoprofen sodium, alminoprofen, zaltoprofen,
mefenamic acid, mefenamic acid aluminum, tolfenamic acid,
floctafenine, ketophenylbutazone, oxyphenbutazone, piroxicam,
tenoxicam, ampiroxicam, epirizole, tiaramide hydrochloride,
30 tinoridine hydrochloride, emorfazone, neurotropin, pregabalin,
sulpyrine, migrenin, pyrine cold remedies, dimetotiazine
mesylate, simetride combination drug, non-pyrine cold remedies
and the like can be mentioned.
[0123]

35 Examples of the vitamin include mecobalamin and the like.

[0124]

Examples of the muscle relaxant include tolperisone hydrochloride, chlorzoxazone, chlormezanone, methocarbamol, phenprobamate, pridinol mesylate, chlorphenesin, baclofen
5 carbamate, eperisone hydrochloride, afloqualone, tizanidine hydrochloride, alcuronium chloride, suxamethonium chloride, tubocurarine chloride, dantrolene sodium, pancuronium bromide, vecuronium bromide and the like.

[0125]

10 Examples of the antidepressant include tricyclic or tetracyclic antidepressant, SSRI (selective serotonin reuptake inhibitor), SNRI (serotonin-noradrenaline reuptake inhibitor) and the like. Examples of the tricyclic antidepressant include imipramine hydrochloride, desipramine hydrochloride,
15 clomipramine hydrochloride, trimipramine maleate, amitriptyline hydrochloride, nortriptyline hydrochloride, lofepramine hydrochloride, amoxapine, dosulepin hydrochloride and the like, and examples of the tetracyclic antidepressant include maprotiline, mianserin and the like. Examples of SSRI include
20 paroxetine hydrochloride hydrate and the like, and examples of SNRI include milnacipran hydrochloride, duloxetine hydrochloride and the like.

[0126]

The weight ratio of the compound of the present invention
25 to other drugs is not particularly limited, and any two or more kinds of other drugs may be administered in combination.

[0127]

Preferable embodiments of the present invention are the following inventions.

- 30 <1> Compound (1) or a pharmaceutically acceptable salt thereof;
<2> the compound or a pharmaceutically acceptable salt thereof of the above-mentioned <1>, wherein Y is CH₂ or S;
<3> the compound or a pharmaceutically acceptable salt thereof of the above-mentioned <2>, wherein Y is CH₂;
35 <4> the compound or a pharmaceutically acceptable salt thereof

- of any of the above-mentioned <1> - <3>, wherein R^A is -
 $CH=CZ^1(COX)$;
- <5> the compound or a pharmaceutically acceptable salt thereof
of any of the above-mentioned <1> - <3>, wherein R^A is $-CH_2-$
5 $CZ^1Z^2(COX)$;
- <6> the compound or a pharmaceutically acceptable salt thereof
of any of the above-mentioned <1> - <5>, wherein n is 1;
- <7> the compound or a pharmaceutically acceptable salt thereof
of any of the above-mentioned <1> - <6>, wherein A-B is a
10 carbon-carbon double bond;
- <8> the compound or a pharmaceutically acceptable salt thereof
of any of the above-mentioned <1> - <7>, wherein Z^1 and Z^2 are
hydrogen atoms;
- <9> the compound or a pharmaceutically acceptable salt thereof
15 of any of the above-mentioned <1> - <8>, wherein R^1 is a
hydrogen atom and R^2 is an alkyl group having 1 - 3 carbon
atoms;
- <10> the compound or a pharmaceutically acceptable salt thereof
of the above-mentioned <9>, wherein R^2 is a methyl group;
- 20 <11> the compound or a pharmaceutically acceptable salt thereof
of any of the above-mentioned <1> - <10>, wherein R^3 is an
alkyl group having 1 - 4 carbon atoms or a cycloalkyl group
having 3 - 6 carbon atoms;
- <12> the compound or a pharmaceutically acceptable salt thereof
25 of the above-mentioned <11>, wherein R^3 is an alkyl group
having 2 - 4 carbon atoms or a cycloalkyl group having 3 - 6
carbon atoms;
- <13> the compound or a pharmaceutically acceptable salt thereof
of the above-mentioned <12>, wherein R^3 is a cycloalkyl group
30 having 3 - 6 carbon atoms;
- <14> the compound or a pharmaceutically acceptable salt thereof
of the above-mentioned <12>, wherein R^3 is an alkyl group
having 2 - 3 carbon atoms or a cycloalkyl group having 3 - 5
carbon atoms;
- 35 <15> the compound or a pharmaceutically acceptable salt thereof

of the above-mentioned <14>, wherein R³ is a cycloalkyl group having 3 - 5 carbon atoms;

<16> the compound or a pharmaceutically acceptable salt thereof of any of the above-mentioned <1> - <15>, wherein X is OR⁴;

5 <17> the compound or a pharmaceutically acceptable salt thereof of the above-mentioned <16>, wherein R⁴ is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms or a substituted alkyl group having 1 - 6 carbon atoms;

<18> the compound or a pharmaceutically acceptable salt thereof
10 of the above-mentioned <17>, wherein R⁴ is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms, or an alkyl group having 1 - 6 carbon atoms substituted by a substituent(s) selected from the group consisting of a hydroxy group, an acyl group, an acyloxy group, an alkoxycarbonyloxy group, a substituted amino
15 group, an aryl group and a heterocyclic group;

<19> the compound or a pharmaceutically acceptable salt thereof of the above-mentioned <17>, wherein R⁴ is a hydrogen atom, an alkyl group having 1 - 4 carbon atoms or a substituted alkyl group having 1 - 4 carbon atoms;

20 <20> the compound or a pharmaceutically acceptable salt thereof of the above-mentioned <19>, wherein R⁴ is a hydrogen atom, an alkyl group having 1 - 4 carbon atoms, or an alkyl group having 1 - 4 carbon atoms substituted by a substituent(s) selected from the group consisting of a hydroxy group, an acyl group, an
25 acyloxy group, an alkoxycarbonyloxy group, a substituted amino group, an aryl group and a heterocyclic group;

<21> the compound or a pharmaceutically acceptable salt thereof of the above-mentioned <20>, wherein R⁴ is a hydrogen atom, an alkyl group having 1 - 4 carbon atoms, a pivaloylmethyl group
30 or a (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl group;

<22> a cyclodextrin clathrate compound or a pharmaceutically acceptable salt thereof of the compound of any of the above-mentioned <1> - <21>;

<23> a medicament comprising the compound or a pharmaceutically
35 acceptable salt thereof of any of the above-mentioned <1> -

<21>, or the cyclodextrin clathrate compound of the above-mentioned <22> as an active ingredient;

<24> the medicament of the above-mentioned <23>, which is a prophylactic or therapeutic agent for a blood flow disorder;

5 <25> the medicament of the above-mentioned <24>, wherein the blood flow disorder is a blood flow disorder of nerve;

<26> the medicament of the above-mentioned <25>, wherein the blood flow disorder of nerve is a blood flow disorder associated with spinal canal stenosis;

10 <27> the medicament of the above-mentioned <24>, wherein the blood flow disorder is a blood flow disorder of peripheral artery, skin or brain;

<28> the medicament of the above-mentioned <27>, wherein the blood flow disorder of peripheral artery is a blood flow

15 disorder associated with chronic arterial occlusion or pulmonary hypertension;

<29> the medicament of the above-mentioned <27>, wherein the blood flow disorder of skin is a blood flow disorder associated with pressure ulcer;

20 <30> the medicament of the above-mentioned <27>, wherein the blood flow disorder of brain is a blood flow disorder associated with suppression of recurrence after cerebral infarction;

<31> the compound or a pharmaceutically acceptable salt thereof

25 of any of the above-mentioned <1> - <21>, or the cyclodextrin clathrate compound of the above-mentioned <22> for use in the prophylaxis or treatment of a blood flow disorder;

<32> the compound or a pharmaceutically acceptable salt thereof, or the cyclodextrin clathrate compound for use of the above-

30 mentioned <31>, wherein the blood flow disorder is a blood flow disorder of nerve;

<33> the compound or a pharmaceutically acceptable salt thereof, or the cyclodextrin clathrate compound for use of the above-mentioned <32>, wherein the blood flow disorder of nerve is a

35 blood flow disorder associated with spinal canal stenosis;

<34> the compound or a pharmaceutically acceptable salt thereof,
or the cyclodextrin clathrate compound for use of the above-
mentioned <31>, wherein the blood flow disorder is a blood flow
disorder of peripheral artery, skin or brain;

5 <35> the compound or a pharmaceutically acceptable salt thereof,
or the cyclodextrin clathrate compound for use of the above-
mentioned <34>, wherein the blood flow disorder of peripheral
artery is a blood flow disorder associated with chronic
arterial occlusion or pulmonary hypertension;

10 <36> the compound or a pharmaceutically acceptable salt thereof,
or the cyclodextrin clathrate compound for use of the above-
mentioned <34>, wherein the blood flow disorder of skin is a
blood flow disorder associated with pressure ulcer;

<37> the compound or a pharmaceutically acceptable salt thereof,
15 or the cyclodextrin clathrate compound for use of the above-
mentioned <34>, wherein the blood flow disorder of brain is a
blood flow disorder associated with suppression of recurrence
after cerebral infarction;

<38> a method for prophylaxis or treatment of a blood flow
20 disorder in a mammal, comprising administering an effective
amount of the compound or a pharmaceutically acceptable salt
thereof of any of the above-mentioned <1> - <21>, or the
cyclodextrin clathrate compound of the above-mentioned <22> to
the mammal;

25 <39> the method of the above-mentioned <38>, wherein the blood
flow disorder is a blood flow disorder of nerve;

<40> the method of the above-mentioned <39>, wherein the blood
flow disorder of nerve is a blood flow disorder associated with
spinal canal stenosis;

30 <41> the method of the above-mentioned <38>, wherein the blood
flow disorder is a blood flow disorder of peripheral artery,
skin or brain;

<42> the method of the above-mentioned <41>, wherein the blood
flow disorder of peripheral artery is a blood flow disorder
35 associated with chronic arterial occlusion or pulmonary

hypertension;

<43> the method of the above-mentioned <41>, wherein the blood flow disorder of skin is a blood flow disorder associated with pressure ulcer;

5 <44> the method of the above-mentioned <41>, wherein the blood flow disorder of brain is a blood flow disorder associated with suppression of recurrence after cerebral infarction;

<45> use of the compound or a pharmaceutically acceptable salt thereof of any of the above-mentioned <1> - <21>, or the
10 cyclodextrin clathrate compound of the above-mentioned <22> in the manufacture of a prophylactic or therapeutic agent for a blood flow disorder;

<46> the use of the above-mentioned <45>, wherein the blood flow disorder is a blood flow disorder of nerve;

15 <47> the use of the above-mentioned <46>, wherein the blood flow disorder of nerve is a blood flow disorder associated with spinal canal stenosis;

<48> the use of the above-mentioned <45>, wherein the blood flow disorder is a blood flow disorder of peripheral artery,
20 skin or brain;

<49> the use of the above-mentioned <48>, wherein the blood flow disorder of peripheral artery is a blood flow disorder associated with chronic arterial occlusion or pulmonary hypertension;

25 <50> the use of the above-mentioned <48>, wherein the blood flow disorder of skin is a blood flow disorder associated with pressure ulcer; and

<51> the use of the above-mentioned <48>, wherein the blood flow disorder of brain is a blood flow disorder associated with
30 suppression of recurrence after cerebral infarction.

[0128]

The present invention is explained in more detail in the following by referring to Reference Examples, Examples, Preparation Examples and Experimental Examples. The present
35 invention is not limited by these Examples and the like and may

be changed within the scope of the present invention.

Unless otherwise specified, the apparatuses, reagents and the like used in the present Examples and the like are easily obtained or prepared according to the methods generally
5 conducted in the pertinent technical field, or commercially available.

In the following Reference Examples and Examples, the "room temperature" is generally about 10°C to about 30°C. Unless otherwise specified, the ratios of mixed solvents are
10 volume mixing ratios. % shows mol/mol% for yield and wt% for others.

The following abbreviations are used in the Reference Examples and Examples.

Me: methyl

15 Et: ethyl

Pr: propyl

Bu: butyl

Ph: phenyl

Ac: acetyl

20 THP: tetrahydro-2H-pyran-2-yl

TBS: tert-butyldimethylsilyl

KHMDS: potassium bistrimethylsilylamide

DBU: 1,8-diazabicyclo[5.4.0]undec-7-ene

DMAP: 4-dimethylaminopyridine

25 Py: pyridine

DMSO: dimethyl sulfoxide

DME: 1,2-dimethoxyethane


p-TsOH: p-toluenesulfonic acid

aq: aqueous solution

30 [0129]

Reference Example 1: methyl (S)-5-cyclopropyl-2-methylpent-4-ynoate

[0130]



[0131]

Lithium chloride (4.24 g) and zinc powder (9.15 g) were dried under reduced pressure, and tetrahydrofuran (THF) (100 mL), 1,2-dibromoethane (0.433 mL) and trimethylchlorosilane (0.127 mL) were added thereto at room temperature. To the mixture was added dropwise a solution of methyl (R)-3-iodo-2-methylpropionate (22.8 g) in THF (30 mL), and the mixture was stirred at 40°C for 1.5 hr to prepare an organozinc reagent.

10 In another reaction container were added lithium chloride (7.63 g), copper(I) cyanide (8.06 g) and THF (90 mL), and the mixture was stirred for 1 hr. The mixture was cooled to -10°C, and the aforementioned organo zinc reagent was added dropwise thereto. The reaction mixture was stirred at -10°C for 10 min, cooled to

15 -78°C, and a solution of 2-(bromoethynyl)cyclopropane (14.5 g) in THF (50 mL) was added dropwise. The mixture was stirred at the same temperature for 15 hr and poured into an aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water, and


20 the insoluble material was filtered off and washed with ethyl acetate. The filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (diethyl ether:hexane = 1:30 - 1:5) to give the title compound (9.93 g). Yield: 66%.

25 ¹H NMR (400 MHz, CDCl₃) δ 3.69 (s, 3H), 2.59 (dq, J = 6.8, 7.2 Hz, 1H), 2.46 (ddd, J = 2.0, 6.0, 16.4 Hz, 1H), 2.29 (ddd, J = 2.0, 7.8, 16.4 Hz, 1H), 1.23 (d, J = 7.2 Hz, 3H), 1.22 (m, 1H), 0.71 (m, 2H), 0.60 (m, 2H).

[0132]

30 Reference Example 2: dimethyl (S)-(+)-(6-cyclopropyl-3-methyl-2-oxohex-5-yn-1-yl)phosphonate

[0133]




THF (25 mL) was added to dimethyl methylphosphonate (4.34 mL), and 1.6 M n-butyllithium (23.7 mL) was added dropwise thereto at -78°C . The reaction mixture was stirred at -78°C for 1 hr, a solution of methyl (S)-5-cyclopropyl-2-methylpent-4-ynoate obtained in Reference Example 1 in THF (10 mL) was added thereto, and the mixture was stirred at the same temperature for 4 hr. To the reaction mixture was added an aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate:hexane = 1:3 - ethyl acetate alone) to give the title compound (2.36 g). Yield: 61%.

^1H NMR (400 MHz, CDCl_3) δ 3.81 (s, 3H), 3.78 (s, 3H), 3.20 (ddd, $J = 14.4, 22.8, 28.4$ Hz, 2H), 2.91 (q, $J = 6.8$ Hz, 1H), 2.33 (dddd, $J = 2.0, 6.8, 16.8, 44.4$ Hz, 2H), 1.18 (d, $J = 7.2$ Hz, 3H), 1.18 (m, 1H), 0.71 (m, 2H), 0.60 (m, 2H).

[0135]

Reference Example 3: methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-hydroxymethyl-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate

[0136]



To a suspension of 4-(carboxybutyl)triphenylphosphonium bromide (14.06 g) in THF (86 mL) was added 1M potassium

bistrimethylsilylamide (64 mL), and the mixture was stirred for 1 hr. The mixture was cooled to -78°C, and a solution of (3aR,4S,5R,6aS)-4-(((tert-butyltrimethylsilyl)oxy)methyl)hexahydro-5-((tetrahydro-2H-pyran-2-yl)oxy)-2H-cyclopenta[b]furan-2-ol (3.92 g) in THF (50 mL) was added thereto. The mixture was stirred at the same temperature for 30 min, heated to room temperature, and stirred overnight. Water was added to the reaction mixture, and the mixture was extracted with tert-butylmethylether, acidified with disodium citrate, and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure, and acetone (390 mL) was added thereto. Diisopropylethylamine (9.16 mL), iodomethane (2.95 mL) and 1,8-diazabicyclo[5.4.0]undec-7-ene (7.85 mL) were added thereto at 0°C. The mixture was stirred at room temperature for 3.5 hr, saturated aqueous sodium hydrogen carbonate was added thereto, and the mixture was extracted with ethyl acetate. After concentration under reduced pressure, the residue was purified by silica gel column chromatography to give a viscous oil (3.21 g). Methanol (220 mL) was added to the oil (3.13 g), and the mixture was stirred using 5% Pd/C (1.13 g) under a hydrogen atmosphere for 40 min. The catalyst was filtered off, and the filtrate was concentrated under reduced pressure. To the residue were added pyridine (14 mL), acetic anhydride (14 mL), and 4-dimethylaminopyridine (170 mg), and the mixture was stirred for 1 hr. Water was added to the reaction mixture, and the mixture was extracted with a hexane-ethyl acetate mixed solvent. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. THF (32 mL) was added to the residue, and the mixture was ice-cooled. 1M tetrabutylammonium fluoride (6.7 mL) was added thereto, and the mixture was stirred at room temperature for 3 days. The reaction mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (hexane:ethyl acetate = 3:1 - 1:3) to


give the title compound (2.60 g).

^1H NMR (300 MHz, CDCl_3) δ 5.07 (m, 1H), 4.71 (m, 0.55H), 4.54 (m, 0.45H), 4.15-4.04 (m, 1H), 4.04-3.88 (m, 1H), 3.88-3.73 (m, 2H), 3.66 (s, 3H), 3.60-3.48 (m, 2H), 2.29 (m, 2H),
5 2.04 (t, $J = 7.5$ Hz, 2H), 1.95-1.15 (m, 20H).

[0138]

Reference Example 4: methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-formyl-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate

[0139]




Methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-hydroxymethyl-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate (3.0 g) was dissolved in ethyl acetate (48 mL), and N,N-diisopropylethylamine (7.6 mL) was added thereto under ice-cooling. To the mixture was added a solution of SO_3 -pyridine (3.6 g) in dimethylsulfoxide (24 mL), and the mixture was stirred for 30 min under ice-cooling. The reaction mixture was poured into ethyl acetate (50 mL) and water (50 mL),
20 partitioned by adding aqueous sodium hydrogen carbonate, and the organic layer was concentrated under reduced pressure. Hexane was added to the residue, and the mixture was washed successively with an aqueous copper sulfate solution, saturated brine and water. The organic layer was dried over anhydrous
25 sodium sulfate and concentrated under reduced pressure to give the title compound (2.75 g).

^1H NMR (300 MHz, CDCl_3) δ 9.77 (dd, $J = 16.1, 3.2$ Hz, 1H), 5.13 (m, 1H), 4.70-4.50 (m, 1H), 4.41 (m, 1H), 3.81 (m, 2H), 3.66 (s, 3H), 3.46 (m, 2H), 3.40-2.92 (m, 0.46H), 2.88-2.78 (m, 0.54H), 2.28 (t, $J = 7.5$ Hz, 2H), 2.50-2.20 (m, 1H), 2.06 (s, 3H), 1.98-1.52 (m, 16H).

[0141]

Reference Example 5: methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-
 ((1E,4S)-7-cyclopropyl-4-methyl-3-oxohept-1-en-6-yn-1-yl)-3-
 ((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate
 [0142]



To sodium hydride (60%, dispersion in oil) (575 mg) was added 1,2-dimethoxyethane (126 mL), and a solution of dimethyl (S)-(+)-(6-cyclopropyl-3-methyl-2-oxohex-5-yn-1-yl)phosphonate
 10 (3.41 g) synthesized in Reference Example 2 in 1,2-dimethoxyethane (20 mL) was added dropwise thereto under ice-cooling. The mixture was stirred under ice-cooling for 1 hr, a solution of methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-formyl-3-
 ((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate (2.75 g)
 15 synthesized in Reference Example 4 in 1,2-dimethoxyethane (40 mL) was added dropwise thereto, and the mixture was stirred at the same temperature for 3 hr. To the reaction mixture were added ethyl acetate (290 mL) and saturated brine (290 mL), the mixture was diluted with water to carry out a liquid separation,
 20 and the aqueous layer was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (hexane:ethyl acetate = 86:14 - ethyl acetate alone) to give the title
 25 compound (3.43 g) as a colorless oil. Yield: 61% (2 step yield from methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-hydroxymethyl-3-
 ((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate obtained in Reference Example 3).

¹H NMR (400 MHz, CDCl₃) δ 6.80 (m, 1H), 6.28 (m, 1H), 5.12
 30 (brt, J = 5.7 Hz, 1H), 4.54 (dt, J = 12.8, 3.0 Hz, 1H), 4.12-3.96 (m, 1H), 3.83-3.58 (m, 2H), 3.65 (s, 3H), 3.50-3.38 (m,


1H), 3.14-2.83 (m, 1H), 2.75-2.16 (m, 4H), 2.06 (s, 3H), 1.95-1.05 (m, 20H), 1.18 (d, J = 7.2 Hz, 3H), 0.75-0.54 (m, 4H).

[0144]

Reference Example 6: methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-

5 ((1E,3S,4S)-7-cyclopropyl-3-hydroxy-4-methylhept-1-en-6-yn-1-yl)-3-hydroxycyclopentyl)heptanoate

[0145]



[0146]

10 Methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-((1E,4S)-7-cyclopropyl-4-methyl-3-oxohept-1-en-6-yn-1-yl)-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate (2.47 g) synthesized in Reference Example 5 was dissolved in methanol (42 mL), cerium chloride heptahydrate (1.74 g) was added thereto, and
15 the mixture was cooled to -40°C. To the mixture was added sodium borohydride (90.2 mg), and the mixture was stirred at the same temperature for 75 min. The reaction mixture was diluted with ethyl acetate, partitioned by adding saturated aqueous sodium hydrogen carbonate and water, and the aqueous
20 layer was extracted with ethyl acetate. The organic layers were combined, washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (2.45 g) as a crude product.

¹H NMR (400 MHz, CDCl₃) δ 5.65-5.45 (m, 2H), 5.09 (brs, 25 1H), 4.60 (m, 1H), 4.14-3.98 (m, 1H), 3.92-3.78 (m, 2H), 3.66 (s, 3H), 2.60-2.30 (m, 1H), 2.28 (t, J = 7.6 Hz, 2H), 2.30-2.13 (m, 2H), 2.04 (s, 3H), 1.85-1.10 (m, 23H), 0.96 (m, 3H), 0.72-0.58 (m, 4H).

[0147]

30 The aforementioned crude product (2.45 g) was dissolved in THF (9.3 mL), a 65% aqueous acetic acid solution (93 mL) was added thereto, and the mixture was stirred at 45°C for 2 hr.

The reaction mixture was allowed to cool to room temperature, water (100 mL) was added thereto, and the mixture was extracted with an ethyl acetate-hexane (1:1) mixed solvent. The organic layers were combined, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane:ethyl acetate = 67:33 - ethyl acetate alone) to give the title compound (723 mg) and a stereoisomer thereof (1.23 g).

^1H NMR (400 MHz, CDCl_3)


The title compound: δ 5.58 (dd, $J = 15.2, 7.4$ Hz, 1H), 5.45 (dd, $J = 15.2, 9.2$ Hz, 1H), 5.14 (brt, $J = 4.6$ Hz, 1H), 3.97 (t, $J = 7.4$ Hz, 1H), 3.87 (q, $J = 8.5$ Hz, 1H), 3.66 (s, 3H), 2.72-2.56 (m, 2H), 2.50 (dq, $J = 8.5, 6.5$ Hz, 1H), 2.28 (t, $J = 7.6$ Hz, 2H), 2.30-2.17 (m, 2H), 2.05 (s, 3H), 1.80-1.49 (m, 6H), 1.49-1.00 (m, 9H), 0.93 (d, $J = 6.5$ Hz, 3H), 0.75-0.68 (m, 2 H), 0.62-0.57 (m, 2H).

Isomer of the title compound: δ 5.61 (dd, $J = 15.2, 5.6$ Hz, 1H), 5.50 (dd, $J = 15.2, 8.6$ Hz, 1H), 5.15 (brt, $J = 4.6$ Hz, 1H), 4.18 (t, $J = 4.8$ Hz, 1H), 3.90 (m, 1H), 3.65 (s, 3H), 2.48 (dq, $J = 8.6, 6.4$ Hz, 1H), 2.28 (t, $J = 7.6$ Hz, 2H), 2.05 (s, 3H), 1.86-1.17 (m, 10H), 1.49-1.00 (m, 9H), 0.96 (d, $J = 6.4$ Hz, 3H), 0.72-0.68 (m, 2H), 0.62-0.58 (m, 2H).

[0148]

Reference Example 7: 7-((1R,2R,3R,5S)-2-((1E,3S,4S)-7-cyclopropyl-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)hept-1-en-6-yn-1-yl)-5-hydroxy-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoic acid

[0149]



[0150]

Methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-((1E,3S,4S)-7-

cyclopropyl-3-hydroxy-4-methylhept-1-en-6-yn-1-yl)-3-hydroxycyclopentyl)heptanoate (67.6 mg) synthesized in Reference Example 6 was dissolved in methylene chloride (1 mL), 3,4-dihydro-2H-pyran (41.3 μ L) and p-toluenesulfonic acid monohydrate (2.9 mg) were added thereto at 0°C, and the mixture was stirred at the same temperature for 30 min. To the reaction mixture were added diethylether (10 mL) and saturated aqueous sodium hydrogen carbonate (10 mL), the mixture was partitioned, and the aqueous layer was extracted with diethylether. The organic layers were combined, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give acetoxy-bis tetrahydropyranyl (THP) ether form (117.1 mg) as a crude product.

^1H NMR (400 MHz, CDCl_3) δ 5.56-5.26 (m, 2H), 5.10 (m, 1H), 4.81-4.58 (m, 2H), 4.20-3.80 (m, 6H), 3.65 (s, 3H), 3.48-3.40 (m, 2H), 2.60-2.31 (m, 1H), 2.28 (t, J = 7.6 Hz, 2H), 2.31-2.10 (m, 3H), 2.04 (s, 3H), 1.90-1.10 (m, 22H), 0.93 (m, 3H), 0.70-0.68 (m, 2H), 0.62-0.58 (m, 2H).
[0151]


The crude product (117 mg) of acetoxy-bis THP ether form obtained above was dissolved in methanol (0.7 mL), a 1N aqueous sodium hydroxide solution (0.45 mL) was added thereto at 0°C, and the mixture was stirred at room temperature for 26 hr. To the reaction mixture was added an aqueous citric acid solution, and the mixture was extracted with ethyl acetate. The organic layers were combined, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (hexane:ethyl acetate = 1:1 - 1:3) to give the title compound (70.9 mg).

^1H NMR (400 MHz, CDCl_3) δ 5.55-5.25 (m, 2H), 4.71 (m, 2H), 4.15-3.84 (m, 6H), 3.47 (m, 2H), 2.50-2.35 (m, 1H), 2.34 (t, J = 7.2 Hz, 2H), 2.25-2.10 (m, 2H), 1.97-1.20 (m, 28H), 0.94 (m, 3H), 0.71-0.68 (m, 2H), 0.60-0.58 (m, 2H).
[0152]

Reference Example 8: methyl 7-((1R,2R,3R,5S)-2-((1E,3S,4S)-7-

cyclopropyl-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)hept-1-en-6-yn-1-yl)-5-hydroxy-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate

[0153]




To acetoxy-bis THP ether form synthesized in the same manner as in Reference Example 7 (682.4 mg) were added methanol (6 mL) and potassium carbonate (382.8 mg), and the mixture was stirred at 40°C for 2 hr. The reaction mixture was allowed to cool to room temperature, diluted with ethyl acetate, neutralized with acetic acid, and partitioned by adding saturated aqueous sodium hydrogen carbonate. The organic layer was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (hexane:ethyl acetate = 84:16 - ethyl acetate alone) to give the title compound (566.6 mg).

¹H NMR (400 MHz, CDCl₃) δ 5.58-5.42 (m, 1.5H), 5.28-5.18 (m, 0.5H), 4.77-4.60 (m, 2H), 4.18-4.08 (m, 2H), 3.90-3.75 (m, 3H), 3.66 (s, 3H), 3.51-3.38 (m, 2H), 2.50-2.00 (m, 5H), 2.29 (t, J = 7.4 Hz, 2H), 2.00-1.40 (m, 26H), 0.92 (dddd, J = 17.2, 17.2, 6.8, 6.8 Hz, 3H), 0.74-0.68 (m, 2H), 0.62-0.58 (m, 2H).

[0155]

25 Reference Example 9: (2E)-7-((1R,2R,3R,5S)-2-((1E,3S,4S)-7-cyclopropyl-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)hept-1-en-6-yn-1-yl)-5-hydroxy-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)hept-2-enoic acid

[0156]



To methyl 7-((1R,2R,3R,5S)-2-((1E,3S,4S)-7-cyclopropyl-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)hept-1-en-6-yn-1-yl)-5-hydroxy-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate (364.4 mg) synthesized in Reference Example 8 was added pyridine (2 mL), and the mixture was cooled to 0°C. Trimethylchlorosilane (0.12 mL) was added thereto, and the mixture was stirred at the same temperature for 30 min. The reaction mixture was diluted with a mixed solvent of hexane and ethyl acetate (10:1), partitioned by adding saturated aqueous sodium hydrogen carbonate, and the organic layer was washed successively with water and saturated brine. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give silylether form (366.1 mg) as a crude product.

The crude product (366.1 mg) of silylether form was dissolved in THF (2.5 mL), a THF solution (1.41 mL) of 1.12 M lithium diisopropylamide was added thereto at -78°C, and the mixture was stirred at the same temperature for 30 min. To the mixture was added dropwise a solution of diphenyldiselenide (412.9 mg) in THF (1 mL) at -78°C, and the mixture was stirred at the same temperature for 45 min and at 0°C for 1 hr. The reaction mixture was diluted with ethyl acetate, partitioned by adding saturated ammonium chloride, and the aqueous layer was extracted with ethyl acetate. The organic layers were combined, washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give Selenylated form (763 mg) as a crude product.

To the crude product (763 mg) of the Selenylated form were added ethyl acetate (4 mL), THF (2 mL) and sodium hydrogen carbonate (179.2 mg), and the mixture was stirred. To the


mixture was added 30% hydrogen peroxide water (0.21 mL), and
 the mixture was stirred at 35°C for 1 hr. The reaction mixture
 was diluted with ethyl acetate, saturated aqueous sodium
 hydrogen carbonate was added thereto, and the mixture was
 5 extracted with ethyl acetate. The organic layers were combined,
 washed with saturated brine, dried over anhydrous sodium
 sulfate, and concentrated under reduced pressure. The residue
 was dissolved in methanol (5 mL), a 1N aqueous sodium hydroxide
 solution (1.9 mL) was added thereto at 0°C, and the mixture was
 10 stirred at room temperature for 43 hr. The reaction mixture
 was diluted with ethyl acetate, neutralized with a saturated
 aqueous disodium citrate solution, and extracted with ethyl
 acetate. The extract was dried over anhydrous sodium sulfate,
 concentrated under reduced pressure, and purified by silica gel
 15 column chromatography (hexane:ethyl acetate = 3:1 - ethyl
 acetate - ethyl acetate:methanol = 20:1) to give the title
 compound (169.8 mg).

¹H NMR (400 MHz, CDCl₃) δ 7.00-6.87 (m, 1H), 5.81 (d, J =
 14.8 Hz, 1H), 5.63-5.41 (m, 2H), 5.30-5.18 (m, 1H), 4.82-4.60
 20 (m, 2H), 4.20-3.52 (m, 6H), 3.72 (s, 3H), 3.53-3.39 (m, 2H),
 2.59-1.00 (m, 25H), 0.92 (m, 3H), 0.74-0.63 (m, 2H), 0.60-0.52
 (m, 2H).

[0158]

Reference Example 10: methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-
 25 ((1E,4S)-4-methyl-3-oxonon-1-en-6-yn-1-yl)-3-((tetrahydro-2H-
 pyran-2-yl)oxy)cyclopentyl)heptanoate

[0159]



[0160]

30 The title compound was synthesized in the same manner as
 in Reference Example 5 from dimethyl (S)-3-methyl-2-oxooct-5-


yn-1-yl)phosphonate and methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-formyl-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate synthesized in Reference Example 4.

¹H NMR (300 MHz, CDCl₃) δ 6.76 (m, 1H), 6.29 (m, 1H),
 5 5.13 (m, 1H), 4.58 (m, 1H), 4.12-3.44 (m, 3H), 3.66 (s, 3H),
 2.89 (m, 1H), 2.74-2.10 (m, 5H), 2.07 (s, 3H), 1.99-1.08 (m,
 21H), 1.20 (d, J = 7.2 Hz, 3H), 1.10 (t, J = 7.4 Hz, 3H).

[0161]

Reference Example 11: methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-
 10 hydroxymethyl-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)-
 2,2-difluoroheptanoate

[0162]



[0163]

15 The title compound was synthesized in the same manner as
 in Reference Example 3 from (3aR,4S,5R,6aS)-4-(((tert-
 butyldimethylsilyl)oxy)methyl)hexahydro-5-((tetrahydro-2H-
 pyran-2-yl)oxy)-2H-cyclopenta[b]furan-2-ol by using 4-(1-
 carboxy-2,2-difluorobutyl)triphenylphosphonium bromide instead
 20 of 4-(carboxybutyl)triphenylphosphonium bromide.


¹H NMR (300 MHz, CDCl₃) δ 5.64 (dt, J = 5.2, 5.2, 2.0 Hz,
 1H), 4.70 (dd, J = 4.0, 0.8 Hz, 0.5H), 4.53 (dd, J = 5.1, 2.4
 Hz, 0.5H), 4.08 (dddd, J = 8.4, 6.6, 4.8, 4.4 Hz, 0.5H), 4.53
 (dd, J = 8.0, 6.2, 4.4, 4.4 Hz, 0.5H), 3.96-3.71 (m, 2H), 3.85
 25 (s, 3H), 3.58-3.43 (m, 2H), 3.00 (brs, 1H), 2.29 (m, 2H), 2.12-
 1.15 (m, 18H), 2.03 (d, J = 2.4 Hz, 3H). ¹⁹F NMR (282.65 MHz,
 CD₃OD) δ -106.26 (t, J = 17.3 Hz, 2F).

[0164]

Reference Example 12: 2-((1S,2R,3R,5S)-5-hydroxy-2-
 30 ((1E,3RS,4S)-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)non-1-
 en-6-yn-1-yl)-3-((tetrahydro-2H-pyran-2-

yl)oxy)cyclopentyl)ethanol

[0165]



[0166]


5 (3aR,4R,5R,6aS)-hexahydro-5-hydroxy-4-((1E,3RS,4S)-3-hydroxy-4-methyl-1-nonen-6-ynyl)-2H-cyclopenta[b]furan-2-one (3.50 g) was dissolved in methylene chloride (61 mL), 2,3-dihydropyran (3.3 mL) and p-toluenesulfonic acid monohydrate (24.5 mg) were added thereto under ice-cooling, and the mixture
10 was stirred at room temperature for 30 min. The reaction mixture was poured into saturated aqueous sodium hydrogen carbonate, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure, and the residue was
15 purified by silica gel column chromatography (hexane:ethyl acetate = 85/15 - ethyl acetate) to give bis THP ether form (7.26 g). The bis THP ether form (994 mg) was dissolved in THF (4 mL) and added to a suspension of lithium aluminum hydride (75 mg) in THF (4 mL) under ice-cooling. The mixture was
20 stirred at the same temperature for 40 min. The reaction mixture was diluted with diethylether, an aqueous sodium sulfate solution was added thereto, and the mixture was filtered through celite and washed with ethyl acetate. The filtrate was concentrated under reduced pressure to give the
25 title compound (953 mg) as a viscous oil.

^1H NMR (300 MHz, CDCl_3) δ 5.60-5.26 (m, 2H), 4.77-4.65 (m, 2H), 4.30-3.60 (m, 7H), 3.48 (m, 2H), 2.20-1.40 (m, 25H), 1.13 (t, $J = 6.9$ Hz, 3H), 0.96 (m, 3H).

[0167]

30 Reference Example 13: methyl 4-((2-((1R,2R,3R,5S)-5-hydroxy-2-((1E,3RS,4S)-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)non-1-

en-6-yn-1-yl)-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)ethyl)thio)butanoate
[0168]



5 [0169]

2-((1S,2R,3R,5S)-5-hydroxy-2-((1E,3RS,4S)-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)non-1-en-6-yn-1-yl)-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)ethanol (122 mg) synthesized in Reference Example 12 was dissolved in THF (2 mL),
10 diisopropylethylamine (91 μ L) and methanesulfonyl chloride (30 μ L) were added thereto at -20°C , and the mixture was stirred for 1 hr. To the reaction mixture were added diisopropylethylamine (44 μ L) and methanesulfonyl chloride (10 μ L), and the mixture was stirred at -10°C for 1 hr.
15 Diisopropylamine (91 μ L) and chlorotrimethylsilane (40 μ L) were added thereto, and the mixture was stirred for 1 hr. Furthermore, a solution of S-potassium thioacetate (92 mg) in N,N-dimethylformamide (2.5 mL) was added thereto at -10°C , potassium carbonate (211 mg) was added thereto, and the mixture
20 was stirred at room temperature overnight. Water was added to the reaction mixture, the mixture was extracted with a mixed solvent of ethyl acetate and hexane (1:1), and the organic layer was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The
25 residue was dissolved in methanol (1 mL), methyl 4-iodobutanoate (72 mg) and potassium carbonate (86 mg) were added thereto, and the mixture was stirred at room temperature for 1.5 hr. Thereafter, a saturated aqueous ammonium chloride solution was added thereto, the mixture was extracted with
30 tert-butylmethylether, dried over anhydrous sodium sulfate, and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (hexane:ethyl

acetate = 4:1 - 1:1) to give the title compound (71 mg).

^1H NMR (300 MHz, CDCl_3) δ 5.55 (m, 2H), 4.68 (m, 2H),
4.30-3.70 (m, 5H), 3.67 (s, 3H), 3.46 (m, 2H), 2.60-2.40 (m,
4H), 2.23 (m, 2H), 2.10-1.40 (m, 26H), 1.13 (t, $J = 6.9$ Hz, 3H),
5 0.96 (m, 3H).

[0170]

Example 1: methyl 4-((2-((1R,2R,3R)-3-hydroxy-2-((1E,3RS,4S)-3-
hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-
oxocyclopentyl)ethyl)thio)butanoate [compound (1)-1]

10 [0171]



[0172]

To methyl 4-((2-((1R,2R,3R,5S)-5-hydroxy-2-((1E,3RS,4S)-
4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)non-1-en-6-yn-1-yl)-
15 3-((tetrahydro-2H-pyran-2-
yl)oxy)cyclopentyl)ethyl)thio)butanoate (70 mg) synthesized in
Reference Example 13 were added ethyl acetate (0.8 mL) and N,N-
diisopropylethylamine (0.15 mL), a solution of SO_3 -pyridine (63
mg) in dimethylsulfoxide (0.67 mL) was slowly added thereto at
20 0°C , and the mixture was stirred for 20 min. Thereafter, the
mixture was diluted with ethyl acetate, and partitioned by
adding water. The organic layer was washed with 1N
hydrochloric acid and saturated aqueous sodium hydrogen
carbonate, dried over anhydrous sodium sulfate, and
25 concentrated under reduced pressure, and the residue was
purified by silica gel column chromatography (hexane:ethyl
acetate = 93/7 - 20/80) to give the corresponding 5-oxo-bis THP
ether form (52 mg). To the 5-oxo-bis THP ether form (52 mg)
were added THF (0.2 mL) and 65% acetic acid (3.6 mL), and the
30 mixture was stirred at 45°C for 1 hr, diluted with water, and
extracted with a mixed solvent of hexane and ethyl acetate
(1:1). The organic layer was dried over anhydrous sodium


sulfate and dried under reduced pressure, and the residue was purified by silica gel column chromatography (hexane:ethyl acetate = 50:50 - 35:65) to give the title compound (29 mg).

¹H NMR (300 MHz, CDCl₃) δ 5.64 (dddd, J = 15.1, 15.1, 8.0, 6.9 Hz, 2H), 4.12-4.03 (m, 2H), 3.70 (s, 1H), 3.68 (s, 2H), 2.76 (dd, J = 7.5, 6.6 Hz, 1H), 2.62 (dt, J = 7.2, 3.2 Hz, 2H), 2.51 (t, J = 7.4 Hz, 2H), 2.43 (t, J = 7.4 Hz, 2H), 2.39-2.10 (m, 7H), 2.10-1.61 (m, 7H), 1.12 (t, J = 7.5 Hz, 3H), 0.97 (d, J = 6.6 Hz, 3H).

[0173]

Example 2: methyl (2E)-4-((2-((1R,2R,3R)-3-hydroxy-2-((1E,3RS,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)ethyl)thio)but-2-enoate [compound (1)-2]

[0174]



[0175]


The title compound was obtained by synthesis in the same manner as in Reference Example 13 and Example 1 by using methyl 4-bromo-2-butenate instead of methyl 4-iodobutanoate from 2-((1S,2R,3R,5S)-5-hydroxy-2-((1E,3RS,4S)-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)non-1-en-6-yn-1-yl)-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)ethanol synthesized in Reference Example 12.

¹H NMR (300 MHz, CDCl₃) δ 6.86 (ddd, J = 15.6, 7.6, 7.6 Hz, 1H), 6.07 (dt, J = 15.6, 9.5 Hz, 1H), 5.92-5.56 (m, 2H), 4.30-4.00 (m, 2H), 3.75 (s, 3H), 3.60-3.00 (m, 2H), 2.90-2.70 (m, 2H), 2.70-2.08 (m, 9H), 2.05-1.64 (m, 4H), 1.13 (t, J = 7.2 Hz, 3H), 1.02-0.95 (m, 3H).

[0176]

Example 3: methyl ((1R,2R,3R)-3-hydroxy-2-((1E,3RS,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoate [compound (1)-3]

[0177]



[0178]

Methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-((1E,4S)-4-methyl-3-oxonon-1-en-6-yn-1-yl)-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate (44 mg) synthesized in Reference Example 10 was dissolved in methanol (0.5 mL), cerium chloride heptahydrate (9.5 mg) and sodium borohydride (1.9 mg) were added thereto under ice-cooling, and the mixture was stirred at room temperature for 30 min. The reaction mixture was diluted with ethyl acetate, partitioned by adding a saturated aqueous ammonium chloride solution, and the aqueous layer was extracted with ethyl acetate. The organic layers were combined, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was dissolved in methylene chloride (1 mL), 3,4-dihydro-2H-pyran (10.5 μ L) and p-toluenesulfonic acid monohydrate (0.73 mg) were added thereto at 0°C, and the mixture was stirred at the same temperature for 1 hr. The reaction mixture was partitioned by adding ethyl acetate and saturated aqueous sodium hydrogen carbonate, and the aqueous layer was extracted with ethyl acetate. The organic layers were combined, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. To the residue were added methanol (0.5 mL) and potassium carbonate (10.6 mg), and the mixture was stirred at room temperature for 16 hr. The reaction mixture was partitioned by adding ethyl acetate and a saturated aqueous ammonium chloride solution, and the aqueous layer was extracted with ethyl acetate, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane:ethyl acetate = 84:16 - 50:50) to give methyl 7-((1R,2R,3R,5S)-2-((1E,3RS,4S)-4-methyl-3-((tetrahydro-


2H-pyran-2-yl)oxy)non-1-en-6-yn-1-yl)-5-hydroxy-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate (33 mg). The obtained compound was treated in the same manner as in Example 1 to give the title compound.

5 ^1H NMR (300 MHz, CDCl_3) δ 5.75-5.55 (m, 2H), 4.22-3.98 (m, 2H), 3.66 (s, 3H), 2.75 (dd, $J = 18.4, 7.2$ Hz, 1H), 2.43-1.28 (m, 22H), 1.12 (t, $J = 7.3$ Hz, 3H) 0.97 (d, $J = 6.9$ Hz, 3H).

[0179]

Example 4: 7-((1R,2R,3R)-2-((1E,3S,4S)-7-cyclopropyl-3-hydroxy-10 4-methylhept-1-en-6-yn-1-yl)-3-hydroxy-5-oxocyclopentyl)heptanoic acid [compound (1)-9]

[0180]



[0181]

15 7-((1R,2R,3R,5S)-2-((1E,3S,4S)-7-cyclopropyl-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)hept-1-en-6-yn-1-yl)-5-hydroxy-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoic acid (70.9 mg) synthesized in Reference Example 7 was dissolved in acetone, and 2.67 M Jones reagent (70.8 μL) was added dropwise thereto at -20°C . The mixture was stirred at the same temperature for 30 min. Then, the Jones reagent (7.1 μL) was added thereto, and the mixture was further stirred for 10 min. The reaction mixture was partitioned by adding 2-propanol (0.15 mL), diethyl ether (10 mL) and water (6 mL). The aqueous layer 25 was extracted with diethylether, and the organic layers were combined, washed successively with water and brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to give oxo-bis THP ether form (53.7 mg) as a crude product.

30 The obtained oxo-bis THP ether form as a crude product (53.7 mg) was dissolved in THF (0.252 mL), a 65% aqueous acetic


acid solution (2.52 mL) was added thereto, and the mixture was stirred at 45°C for 2 hr and at room temperature for 1 hr. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed
 5 with saturated brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane:ethyl acetate = 1:2 - ethyl acetate alone - ethyl acetate:methanol = 20:1) to give the title compound (29.5 mg).

10 ^1H NMR (400 MHz, CDCl_3) δ 5.69 (dd, J = 15.2, 7.6 Hz, 1H), 5.60 (dd, J = 15.2, 8.8, 1H), 4.13-4.03 (m, 2H), 2.75 (dd, J = 18.4, 7.2 Hz, 1H), 2.43-1.20 (m, 22 H), 0.94 (d, J = 6.4 Hz, 3H), 0.75-0.71 (m, 2H), 0.63-0.59 (m, 2H).

[0182]

15 Example 5: (2E)-7-((1R,2R,3R)-2-((1E,3S,4S)-7-cyclopropyl-3-hydroxy-4-methylhept-1-en-6-yn-1-yl)-3-hydroxy-5-oxocyclopentyl)hept-2-enoic acid [compound (1)-10]

[0183]



20 [0184]

The title compound was synthesized by Jones oxidation followed by deTHP reaction in the same manner as in Example 4 by using (2E)-7-((1R,2R,3R,5S)-2-((1E,3S,4S)-7-cyclopropyl-4-methyl-3-((tetrahydro-2H-pyran-2-yl)oxy)hept-1-en-6-yn-1-yl)-5-
 25 hydroxy-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)hept-2-enoic acid synthesized in Reference Example 9. Yield: 66.4%.


^1H NMR (400 MHz, CDCl_3) δ 6.98 (dt, J = 15.6, 6.9 Hz, 1H), 5.98 (brs, 2H), 5.77 (d, J = 15.6 Hz, 1H), 5.58 (dddd, J = 15.3, 15.3, 7.4, 6.8 Hz, 2H), 4.05-3.93 (m, 2H), 2.71 (dd, J = 18.2, 7.4 Hz, 1H), 2.40-1.00 (m, 16H), 0.89 (d, J = 6.8 Hz, 3H),
 30 0.71-0.67 (m, 2H), 0.59-0.55 (m, 2H).

[0185]

Example 6: 7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoic acid

[compound (1)-6]

[0186]



[0187]


The title compound was synthesized in the same manner as in Reference Example 6, Reference Example 7 and Example 4 from methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-((1E,4S)-4-methyl-3-oxonon-1-en-6-yn-1-yl)-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate synthesized in Reference Example 10.

¹H NMR (400 MHz, CDCl₃) δ 5.65 (dddd, J = 15.3, 15.3, 8.3, 7.2 Hz, 2H), 4.12-4.01 (m, 2H), 2.74 (dd, J = 18.0, 7.2 Hz, 1H), 2.45-1.96 (m, 11H), 1.79 (m, 1H), 1.50-1.40 (m, 4H), 1.40-1.30 (m, 7H), 1.13 (t, J = 6.9 Hz, 3H), 0.96 (d, J = 6.9 Hz, 3H).

[0188]

Example 7: (2E)-7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)hept-2-enoic acid [compound (1)-7]

[0189]



[0190]

The title compound was synthesized in the same manner as in Reference Example 6, Reference Example 7, Reference Example 8, Reference Example 9 and Example 5 from methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-((1E,4S)-4-methyl-3-oxonon-1-en-6-yn-1-yl)-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)heptanoate synthesized in Reference Example


10.

^1H NMR (300 MHz, CDCl_3) δ 7.01 (dt, $J = 15.4, 7.0$ Hz, 1H),
5.80 (d, $J = 15.4$ Hz, 1H), 5.62 (dddd, $J = 15.3, 15.3, 8.2, 7.7$
Hz, 2H), 4.08–3.80 (m, 2H), 2.73 (dd, $J = 18.3, 7.5$ Hz, 1H),
5 2.46–1.70 (m, 11H), 1.87–1.30 (m, 8H), 1.14 (t, $J = 7.4$ Hz, 3H),
0.94 (t, $J = 6.9$ Hz, 3H).

[0191]

Example 8: (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 7-
((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-
10 6-yn-1-yl)-5-oxocyclopentyl)heptanoate [compound (1)-4]

[0192]



[0193]

The oxo-bis THP ether form (250 mg) obtained as an
15 intermediate in Example 6 in the same manner as in Example 4
was dissolved in acetone (4.5 mL), and 4-chloromethyl-5-methyl-
1,3-dioxol-2-one (20.8 mg) was added dropwise thereto at 0°C .
Sodium iodide (68.4 mg) and 1,8-diazabicyclo[5.4.0]undec-7-ene
(82.2 mg) were added thereto, and the mixture was stirred at
20 the same temperature for 21 hr. To the reaction mixture were
added diethylether and saturated aqueous sodium hydrogen
carbonate, and the aqueous layer was extracted with
diethylether, dried over anhydrous sodium sulfate, and
concentrated under reduced pressure. The residue was purified
25 by silica gel column chromatography (hexane alone -
hexane:ethyl acetate = 16:84 - ethyl acetate alone - ethyl
acetate:methanol = 98:2) to give ester form (278 mg). Then,
THF (0.95 mL) and a 65% aqueous acetic acid solution (9.5 mL)
were added to the obtained ester form (278 mg), and the mixture
30 was stirred at 45°C for 1 hr. Water was added to the reaction
mixture, and the mixture was extracted with ethyl acetate. The
organic layer was washed with saturated brine, dried over


anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane:ethyl acetate = 3:1 - 1:5 - ethyl acetate alone - ethyl acetate:methanol = 50:1) to give the
5 title compound (168.2 mg).

^1H NMR (300 MHz, CDCl_3) δ 5.64 (dddd, J = 15.2, 15.2, 8.4, 7.1 Hz, 2H), 4.83 (s, 2H), 4.13-3.99 (m, 2H), 2.74 (dd, J = 17.7, 7.5 Hz, 1H), 2.46-1.94 (m, 11H), 1.85-1.19 (m, 14H), 1.13 (d, J = 7.4 Hz, 3H), 0.95 (d, J = 4.2 Hz, 3H).

10 [0194]

Example 9: 3,3-dimethyl-2-oxobutyl 7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoate [compound (1)-5]

[0195]



[0196]

The oxo-bis THP ether form (200.9 mg) obtained as an intermediate in Example 6 in the same manner as in Example 4 was dissolved in acetone (3.7 mL), and 1-chloropinacolone (28.2
20 mg), sodium iodide (14 mg) and 1,8-diazabicyclo[5.4.0]undec-7-ene (16.6 mg) were added thereto at 0°C, and the mixture was stirred at the same temperature for 17 hr. To the reaction mixture were added diethylether and saturated aqueous sodium hydrogen carbonate, and the aqueous layer was extracted with
25 diethylether. The organic layer was dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane alone - hexane:ethyl acetate = 16:84 - ethyl acetate alone - ethyl acetate:methanol = 95:5) to give ester form
30 (163.6 mg). Then, THF (0.95 mL) and a 65% aqueous acetic acid solution (9.5 mL) were added to the obtained ester form (21.6


mg), and the mixture was stirred at 45°C for 2.5 hr. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane:ethyl acetate = 3:1 - ethyl acetate alone) to give the title compound (116.6 mg).

¹H NMR (300 MHz, CDCl₃) δ 5.76-5.57 (m, 2H), 4.88 (s, 2H), 4.18-4.00 (m, 2H), 2.82-2.68 (m, 1H), 2.46-1.95 (m, 10H), 1.85-1.17 (m, 12H), 1.21 (d, J = 4.7 Hz, 9H), 1.14 (dt, J = 7.4, 4.7 Hz, 3H), 0.97 (dd, J = 6.9, 4.5 Hz, 3H).

[0197]

Example 10: 2,2-difluoro-7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4S)-3-hydroxy-4-methylnon-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoic acid [compound (1)-8]

[0198]



[0199]

The title compound was synthesized in the same manner as in Reference Example 4 - Reference Example 7 and Example 4 from methyl 7-((1R,2R,3R,5S)-5-acetoxy-2-hydroxymethyl-3-((tetrahydro-2H-pyran-2-yl)oxy)cyclopentyl)-2,2-difluoroheptanoate synthesized in Reference Example 11.

¹H NMR (400 MHz, CDCl₃) δ 5.60 (dddd, J = 15.2, 15.2, 8.4, 7.4 Hz, 2H), 4.03 (dd, J = 17.2, 8.4 Hz, 1H), 3.97 (t, J = 7.6 Hz, 1H), 2.86-2.65 (m, 1H), 2.51-1.88 (m, 9H), 1.74 (m, 1H), 1.72-1.17 (m, 11H), 1.12 (t, J = 7.4 Hz, 3H), 0.94 (d, J = 6.8 Hz, 3H). ¹⁹F NMR (282.65 MHz, CD₃OD) δ -106.42 (t, J = 15.4 Hz, 2F).

[0200]

Preparation Example 1: Production of tablet

1) compound (1) 0.003 g

2) lactose	50 g
3) cornstarch	15 g
4) calcium carboxymethylcellulose	44 g
5) magnesium stearate	1 g
5 1000 tablets	total 110.003 g

The total amount of 1), 2), and 3) and 30 g of 4) are kneaded with water, vacuum dried and sieved. The sieved powder is mixed with 14 g of 4) and 1 g of 5), and the mixture is punched by a tableting machine. In this manner, 1000 tablets
 10 containing 3 μ g of the Example compound per tablet are obtained.
 [0201]

Preparation Example 2: Production of capsule

1) compound (1)	0.003 g
2) fine powder cellulose	10 g
15 3) lactose	19 g
4) magnesium stearate	1 g
	total 30.003 g

1), 2), 3) and 4) are mixed and filled in gelatin capsules to give 1000 capsules containing 3 μ g of the Example
 20 compound per capsule.
 [0202]

Experimental Example 1: Evaluation of human platelet aggregation inhibitory action in vitro

(1) Preparation of human platelet-rich plasma

25 Using 3.8 w/v% sodium citrate as an anticoagulant, blood samples were collected from human brachial vein at a ratio of sodium citrate solution: blood (volume ratio) = 1:9, and an upper layer of platelet-rich plasma (PRP) was collected by centrifugation (120 $\times$ g, 10 min).. Platelet poor plasma (PPP)
 30 was prepared from PRP-free blood by centrifugation (1700 $\times$ g, 10 min).. The number of platelets in PRP was measured and adjusted to 25 $\times$ 10⁴/ μ L by diluting with PPP.

[0203]

(2) Measurement of platelet aggregation ability

35 The platelet aggregation ability was measured by a

platelet aggregometer (PRP313 M, IMI Co., Ltd.) according to the method of Born (Nature, 1962, vol. 194, pp.927-929). PRP (190 μ L) was placed in a glass cuvette and mounted on the platelet aggregometer. Then, 5 μ L of the compound of the present invention, limaprost (positive control substance) or vehicle control (0.25 vol% dimethyl sulfoxide [DMSO]) was added as a test substance, and the mixture was incubated at 37°C for 10 min. ADP (LMS Co., Ltd.) (5 μ L) as a platelet aggregation agent was added at 3 μ mol/L in a final concentration to cause aggregation, and the maximum response of the aggregation was measured.

[0204]

(3) Evaluation of inhibitory action on platelet aggregation

The aggregation inhibition ratio of a test substance was determined by defining the maximum aggregation ratio of control as 100%, and IC₅₀ value was determined by using "Dx calculation (linear fitting): non-logit transformation" of SAS System Version 9.1.3 (Stat Preclinica Client Ver. 1.1 SAS Institute Japan, Ltd.). Furthermore, the relative activity (%) of each test substance was calculated by defining the activity of IC₅₀ value of limaprost as 100.

The results are shown in Table 1. As shown in Table 1, the compound of the present invention exhibited a very potent activity in the human platelet aggregation inhibitory action test in vitro, and some of them exhibited activity not less than 10-fold that of limaprost.

[0205]

[Table 1]

Test substance	Relative activity (%)
compound (1)-4	643
compound (1)-5	161
compound (1)-6	923
compound (1)-7	1319
compound (1)-9	1200
compound (1)-10	1481
limaprost	100

[0206]

5 Experimental Example 2: Evaluation of inhibitory action on rat platelet aggregation in vitro

(1) Preparation of rat platelet-rich plasma

In the same manner as in the above-mentioned Experimental Example 1(1), blood samples were collected from the abdominal
10 artery of 8- to 9-week-old male SD rats. PRP was diluted with PPP and the number of platelet was adjusted to $63.2 \times 10^4/\mu\text{L}$.

[0207]

(2) Measurement of platelet aggregation ability

In the same manner as in the above-mentioned Experimental
15 Example 1(2) except that the final concentration of ADP was 2 $\mu\text{mol/L}$, the compound of the present invention, prostaglandin E1 (PGE1:positive control substance) or vehicle control was added as a test substance, and the maximum response of aggregation was measured.

20 [0208]

(3) Evaluation of platelet aggregation inhibitory action

In the same manner as in the above-mentioned Experimental Example 1(3), IC_{50} values were determined from the aggregation inhibition ratio of the test substance, and the relative
25 activity (%) of each test substance was calculated by defining the activity of IC_{50} value of PGE1 as 100.

The results are shown in Table 2. The compound of the present invention showed a potent activity in the rat platelet aggregation inhibitory action test in vitro. All of compounds (1)-1 to (1)-3 as test substances are diastereomeric mixtures of R form and S form of the stereochemistry of the hydroxy group on the ω -chain. Therefore, when only the effective active form (one of the diastereomers) is separated from the mixture and the above-mentioned test is performed, the activity may be almost doubled.

10 [0209]

[Table 2]

Test substance	Relative activity (%)
compound (1)-1	169
compound (1)-2	62
compound (1)-3	2175
compound (1)-8	817
PGE1	100

[0210]

Experimental Example 3: Evaluation of persistent inhibitory action on guinea pig platelet aggregation ex vivo

(1) Measurement of IC₅₀ value

As a test substance, a single oral dose of the compound of the present invention (3 doses of 79, 132 and 263 nmol/kg), limaprost (positive control substance: 3 doses of 79, 263 and 789 nmol/kg) or vehicle control (water for injection containing 0.26 vol% DMSO) was administered at 10 mL/kg to 4- to 10-week-old male Hartley guinea pigs. Blood samples were collected from the abdominal artery 4 hr later, platelet aggregation ability was measured, and the IC₅₀ value was calculated from the aggregation inhibitory activity (aggregation was induced by 0.5 μ mol/L ADP) in the same manner as in the above-mentioned Experimental Example 1(3).

[0211]

(2) Administration of test substance and preparation of platelet-rich plasma

A single oral dose showing the IC_{50} value determined in the above-mentioned test (1) of compound (1)-7 (168 nmol/kg), compound (1)-9 (178 nmol/kg), compound (1)-10 (187 nmol/kg) or limaprost (266 nmol/kg) was administered. Blood samples were collected from the abdominal artery of the guinea pig at 1, 2, 4, 6, 10, 16 and 24 hr later. PRP was diluted with PPP and the number of platelet was adjusted to about $60 \times 10^4/\mu L$ in the same manner as in the above-mentioned Experimental Example 1(1) (n=3 - 4).

[0212]

(3) Measurement of platelet aggregation

In the same manner as in the above-mentioned Experimental Example 1(2), the platelet aggregation ability was measured according to the method of Born. PRP (190 μL) was placed in a glass cuvette, mounted on the platelet aggregometer and incubated at 37°C for 10 min. ADP (10 μL) as platelet aggregation agent was added at 0.5 $\mu mol/L$ in a final concentration to cause aggregation, and the maximum reaction of the aggregation was measured.

[0213]

(4) Evaluation of persistent inhibitory action on platelet aggregation

The maximum aggregation inhibition ratio of the test substance based on the maximum aggregation inhibition ratio of the control group as 0% was determined from the maximum coagulation ratio of the control group and the maximum coagulation ratio of the test substance when IC_{50} values were calculated. Furthermore, in a graph showing time course changes of the maximum aggregation inhibition ratio, wherein the vertical axis shows the maximum aggregation inhibition ratio (%) and the horizontal axis shows time (hr) after administration of test substance, the area of the part surrounded by the line connecting the points of the maximum

aggregation inhibition ratios plotted at each measurement time and the horizontal axis was determined as AUC. In addition, the assumed time at which the maximum aggregation inhibition ratio became 0% on the graph was determined as the time at which inhibition of aggregation disappeared (inhibition 0 hr).
[0214]

The results are shown in Table 3. As shown in Table 3, in the guinea pig platelet aggregation inhibitory action test ex vivo, the compounds of the present invention showed higher AUC values from 1 hr after administration to inhibition 0 hr (AUC: 1-inhibition 0 hr) than limaprost did. It was confirmed that the compounds of the present invention had a more potent platelet aggregation inhibitory action than limaprost did. When the maximum aggregation inhibition ratio after 6 hr from the administration as an index of persistent activity was evaluated, the compounds of the present invention showed higher AUC values from 6 hr after administration to inhibition 0 hr (AUC: 6-inhibition 0 hr) and a longer time up to the disappearance of inhibitory of aggregation (inhibition 0 hr) in comparison with limaprost. It was therefore confirmed that the compounds of the present invention are superior to limaprost in the persistent efficacy.

[0215]

[Table 3]

Test substance	Inhibition 0 hr (hr)	AUC:1-inhibition 0 hr (% relative to limaprost)	AUC:6-inhibition 0 hr (% relative to limaprost)
compound (1)-7	>24	713 (202)	526 (359)
compound (1)-9	19.7	627 (178)	401 (273)
compound (1)-10	>24	719 (204)	508 (346)
limaprost	12.1	353 (100)	147 (100)

25

[0216]

Experimental Example 4: Evaluation of increasing effect on rat cauda equine blood flow

(1) Administration of test substance

The compound of the present invention or limaprost
5 (positive control substance) as a test substance was injected at 0.263, 0.789 and 2.63 nmol/kg in principle to Slc:Wistar male rats (body weight: about 200 - 250 g) from the tail vein at a flow rate of 0.2 mL/min for 10 min with an infusion pump (model 11, Harvard Apparatus). A vehicle (1 vol% DMSO-
10 physiological saline) was similarly injected to the control group (5 - 6 mice in each group).

[0217]

(2) Measurement of cauda equine blood flow

The rats were fixed in a prone position under halothane
15 anesthesia and kept warm by a heater (Nippon Medical & Chemical Instruments Co., Ltd.). The lower back of the rats was shaved, dissected to expose the spine, the vertebral arch of the fifth lumbar vertebra was excised and cauda equina was exposed. The cauda equine blood flow was measured by a laser Doppler blood
20 flow meter (TBF-LN1, Unique Medical Co., Ltd.) via a non-contact type probe (LP-NC special type, Unique Medical Co., Ltd.) fixed perpendicularly to a position of the exposed cauda equina upper by about 1 mm from before administration of each test substance until 60 min after the administration. The open
25 wound was filled with liquid paraffin to prevent drying of the exposed cauda equina.

[0218]

(3) Data analysis

The obtained data was taken into a personal computer via
30 PowerLab8/30 (registered trade mark, ML870, ADInstruments) every 1/400 second, and analyzed using LabChart (registered trade mark, ADInstruments). The analysis time points were before administration (administration 0 min), and 3, 5, 8, 10, 13, 15, 18, 20, 30, 40, 50 and 60 min after administration, or
35 before administration (administration 0 min), and 3, 5, 10, 15,

20, 30, 40, 50 and 60 min after administration. The mean value for 10 seconds at each time point was taken as the measurement value, and the change rate (%) relative to the value before administration was obtained by the following equation.

5 The rate of change (%) = (measurement value at each time point after administration - measurement value before administration)/measurement value before administration x 100

Furthermore, difference in the rate of change (%) relative to the control group at each time point [Δ change rate (%)] was calculated by the following equation, and the maximum value of the Δ change rate was taken as the maximum change rate. Δ change rate (%) = change rate of each individual at each time point - mean of change rate of control group

From the maximum change rate at each dose of the test substance, the dose at which the cauda equine blood flow increases by 20% (ED₂₀ value) was determined by linear regression equation, and the relative activity (%) of each test substance was calculated based on the activity of ED₂₀ value of limaprost as 100.

20 The results are shown in Table 4. The compounds of the present invention exhibited a potent activity in the rat cauda equine blood flow test.

[0219]

[Table 4]

Test substance	Relative activity (%)
compound (1)-4	147
compound (1)-6	383
compound (1)-7	375
compound (1)-9	311
compound (1)-10	550
limaprost	100

25

[Industrial Applicability]

[0220]


The prostaglandin derivative of the present invention is useful as an active ingredient of a medicament. A medicament containing the prostaglandin derivative of the present invention as an active ingredient is useful as a medicament for
5 the prophylaxis or treatment of blood flow disorders, particularly, a medicament for the prophylaxis or treatment of a blood flow disorder associated with spinal canal stenosis or chronic arterial occlusion.

[0221]

10 This application is based on a patent application No. 2016-094196 filed in Japan, the contents of which are incorporated in full herein.

CLAIMS

1. A compound represented by the following formula (1) or a pharmaceutically acceptable salt thereof;



wherein

R^A is $-\text{CH}_2-\text{CZ}^1\text{Z}^2(\text{COX})$ or $-\text{CH}=\text{CZ}^1(\text{COX})$, Z^1 and Z^2 are each independently a hydrogen atom or a fluorine atom, X is OR^4 or NR^5R^6 , R^4 is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms or a substituted alkyl group having 1 - 6 carbon atoms, R^5 is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms or a substituted alkyl group having 1 - 6 carbon atoms, R^6 is a hydrogen atom, an alkyl group having 1 - 6 carbon atoms, a substituted alkyl group having 1 - 6 carbon atoms, an alkylsulfonyl group having 1 - 6 carbon atoms, a substituted alkylsulfonyl group having 1 - 6 carbon atoms, an arylsulfonyl group having 6 - 10 carbon atoms or a substituted arylsulfonyl group having 6 - 10 carbon atoms;

Y is CH_2 , S or O ;

A-B is a carbon-carbon single bond, a carbon-carbon double bond or a carbon-carbon triple bond;

a hydroxy group bonded by a wavy line is a hydroxy group having α -configuration, β -configuration or a mixed configuration of α -configuration and β -configuration;

R^1 and R^2 are each independently a hydrogen atom, an alkyl group having 1 - 3 carbon atoms or a substituted alkyl group having 1 - 3 carbon atoms;

n is an integer of 0 - 2;

R^3 is an alkyl group having 1 - 4 carbon atoms, a substituted alkyl group having 1 - 4 carbon atoms, a cycloalkyl

group having 3 - 6 carbon atoms, a substituted cycloalkyl group having 3 - 6 carbon atoms, an aryl group having 6 - 10 carbon atoms or a substituted aryl group having 6 - 10 carbon atoms; excluding 7-((1R,2R,3R)-3-hydroxy-2-((1E,3S,4RS)-3-hydroxy-4-methyloct-1-en-6-yn-1-yl)-5-oxocyclopentyl)heptanoic acid.

2. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein Y is CH₂.

10

3. The compound or a pharmaceutically acceptable salt thereof according to claim 1 or 2, wherein R^A is -CH=CZ¹(COX).

4. The compound or a pharmaceutically acceptable salt thereof according to claim 1 or 2, wherein R^A is -CH₂-CZ¹Z²(COX).

15

5. The compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 4, wherein n is 1.

20 6. The compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 5, wherein A-B is a carbon-carbon double bond.

7. The compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 6, wherein R¹ is a hydrogen atom and R² is an alkyl group having 1 - 3 carbon atoms.

25

8. The compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 7, wherein R³ is an alkyl group having 2 - 3 carbon atoms or a cycloalkyl group having 3 - 5 carbon atoms.

30

9. The compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 8, wherein R³ is a cycloalkyl group having 3 - 5 carbon atoms.

35

10. The compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 9, wherein X is OR⁴.
- 5 11. A cyclodextrin clathrate compound of the compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 10.
12. A medicament comprising the compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 10,
10 or the cyclodextrin clathrate compound according to claim 11 as an active ingredient.
13. The medicament according to claim 12, which is a
15 prophylactic or therapeutic agent for a blood flow disorder.
14. The medicament according to claim 13, wherein the blood flow disorder is a blood flow disorder of nerve.
- 20 15. The medicament according to claim 14, wherein the blood flow disorder of nerve is a blood flow disorder associated with spinal canal stenosis.
16. The medicament according to claim 13, wherein the blood
25 flow disorder is a blood flow disorder of peripheral artery, skin or brain.
17. The medicament according to claim 16, wherein the blood flow disorder of peripheral artery is a blood flow disorder
30 associated with chronic arterial occlusion or pulmonary hypertension.
18. The medicament according to claim 16, wherein the blood flow disorder of skin is a blood flow disorder associated with
35 pressure ulcer.

19. The medicament according to claim 16, wherein the blood flow disorder of brain is a blood flow disorder associated with suppression of recurrence after cerebral infarction.

5