
What is claimed is:


1. A metal-ligand complex of Formula I:



Formula I

wherein M is titanium, zirconium, or hafnium; R is hydrogen or a (C_1 to C_4) alkyl; R^1 is a (C_1 to C_4) alkyl ; any one or two of R^2, R^3, R^4, R^5 is independently a (C_1 to C_{20}) alkyl and the three or two of R^2, R^3, R^4, R^5 is H; and each X is independently a halide, a (C_1 to C_{20}) alkyl, a (C_7 to C_{20}) aralkyl, a (C_1 to C_6) alkyl-substituted (C_6 to C_{12}) aryl, a (C_1 to C_6) alkyl-substituted benzyl, or a silicon-containing alkyl.

2. The metal-ligand complex of Formula I, wherein R^1 is a C_1 alkyl, R^2 is hydrogen, R^3 is a C_1 alkyl, R^4 is hydrogen, R^5 is hydrogen, the metal-ligand complex represented by Formula II:



Formula II

wherein R is hydrogen or a (C_1 to C_2) alkyl, M is titanium, zirconium, or hafnium, and where each X is independently a halide, a (C_1 to C_{20}) alkyl, a (C_7 to C_{20}) aralkyl, a (C_1 to C_6) alkyl-substituted (C_6 to C_{12}) aryl, a (C_1 to C_6) alkyl-substituted benzyl, or a silicon-containing alkyl.

3. The metal-ligand complex of claim 1 or claim 2, wherein R is hydrogen.

4. The metal-ligand complex of claim 1 or claim 2, wherein R is a (C_1 to C_2) alkyl.

5. The metal-ligand complex of claim 1 or claim 2, wherein R is methyl.

6. The metal-ligand complex of claim 1, wherein R is ethyl.

7. The metal-ligand complex of claim 1 or claim 2, wherein M is Zr.

8. The metal-ligand complex of claim 1 or claim 2, wherein each X is independently Cl, a (C_1 to C_3) alkyl, or benzyl.

9. The metal-ligand complex of claim 1 or claim 2 selected from:

metal-ligand complex of Formula I, wherein R is hydrogen, M is zirconium, and each X is Cl;

metal-ligand complex of Formula I, wherein R is methyl, M is zirconium, and each X is Cl;


metal-ligand complex of Formula I, wherein R is ethyl, M is zirconium, and each X is Cl;

metal-ligand complex of Formula I, wherein R is hydrogen, M is zirconium, and each X is methyl;

metal-ligand complex of Formula I, wherein R is methyl, M is zirconium, and each X is methyl; and

metal-ligand complex of Formula I, wherein R is ethyl, M is zirconium, and each X is methyl.

10. A bimodal catalyst system comprising the metal-ligand complex of claim 1 or claim 2 and a bis(2-(pentamethylphenylamido)ethyl)-amine zirconium complex of Formula III:



Formula III

wherein M is Zr and each R^1 independently is Cl, Br, a (C_1 to C_{20}) alkyl, a (C_1 to C_6) alkyl-substituted (C_6-C_{12}) aryl, benzyl, or a (C_1 to C_6) alkyl-substituted benzyl.

11. A method of making a polymerization catalyst, the method comprising contacting, under activating conditions, the metal-ligand complex of claim 1 or claim 2 or the bimodal catalyst system of claim 10 with an activator so as to activate the metal-ligand complex or the bimodal catalyst system, thereby making the polymerization catalyst.

12. The method of claim 11 wherein the activator is methylaluminoxane.

13. A polymerization catalyst made by the method of claim 11.

14. A method of making a polyolefin polymer, the method comprising:

contacting, under polymerization conditions, an olefin with the polymerization catalyst of claim 13 so as to polymerize the olefin, thereby making a polyolefin polymer.

15. A polyolefin polymer made by the method of claim 14.