

METHOD OF DECOMPOSING ORGANOPHOSPHORUS COMPOUNDS

RELATED APPLICATIONS

This application claims the benefit of the filing date of U.S. Provisional Patent Application No. 60/453,762, filed on March 12, 2003, the contents of which are incorporated herein by reference in their entirety.




FIELD OF THE INVENTION

This invention relates to methods of decomposing organophosphorus compounds. The invention more particularly relates to metal ion and metal species catalysis of an alcoholysis reaction which converts toxic organophosphorus compounds into non-toxic compounds. The invention further relates to lanthanum ion catalyzed degradation of chemical warfare agents, insecticides and pesticides.

BACKGROUND OF THE INVENTION

The Chemical Weapons Convention was adopted by the Conference on Disarmament in Geneva on September 3, 1992, entered into force on April 29, 1997, and calls for a prohibition of the development, production, stockpiling and use of chemical weapons and for their destruction under universally applied international control. Eliminating the hazard of chemical warfare agents is desirable both in storage sites and on the battlefield. Decontamination of battlefields requires speed and ease of application of decontaminant. Surfaces involved pose a challenge for decontamination techniques since some surfaces absorb such agents, making decontamination difficult. Examples of surfaces that could be involved include those of tanks, ships, aircraft, weapons, electronic devices, ground, protective clothing and human skin. The decontaminants should not be corrosive, so that surfaces are not damaged during/following decontamination. An optimum solvent of a decontaminating method should provide ease of application, solubility of the chemical warfare agent, non-corrosiveness, and minimal environmental contamination. Since the establishment of the Convention, considerable effort has been directed toward methods of facilitating the controlled decomposition of organophosphorus compounds.

Aqueous decontamination systems, such as hydrolysis systems, have been used in the past, most notably for nerve agents, particularly for the G-agents tabun (GA), sarin (GB), soman (GD) and GF. However, hydrolysis reactions are not suitable for all chemical warfare nerve agents such as V-agents VX (S-2-(diisopropylamino)ethyl O-ethyl methylphosphonothiolate) and Russian-VX (S-2-(diethylamino)ethyl O-isobutyl methylphosphonothiolate), whose decontamination chemistries are very similar to one another (Yang, 1999). The V-agents are about 1000-fold less reactive with hydroxide than the G-agents (due to their poor solubility in water under basic conditions), and they produce product mixtures containing the hydrolytically stable, but toxic, thioic acid byproduct.



Although some chemical warfare agents are water soluble, they may be applied in combination with a polymer so that, being thickened, they adhere well to surfaces. These "thickened" agents are only minimally soluble in water. In the case of decomposition using

a hydrolysis reaction, products in which a phosphorus-sulfur bond is preserved are common; these are toxic in their own right and are relatively resistant to further reaction. Another disadvantage of an aqueous decontamination system is that hydrolysis reactions are not catalytic, and therefore require stoichiometric amounts of reagents. Furthermore, commonly used aqueous methods, due to their alkaline pH, are not suitable for decontamination of human skin. Yet another disadvantage of aqueous decontamination methods is the caustic wastewater produced as an end product, which poses a challenge for disposal.

Historically, decontamination of chemical warfare agents has been effected using hydrolysis or oxidation using bleach or alkali salts. Bleach is corrosive to skin, rubber, and metal surfaces and is ineffective in cold weather conditions. Alkali salts require excess hydroxide ion in order for the reaction to go to completion rapidly, thus resulting in a caustic product. Non-catalytic methanolysis of V-agents has been studied, wherein the reaction of VX with alkoxide leads primarily to a displacement of the SR^- group (Yang *et al.*, 1997).


Transition metal ions and lanthanide series ions and certain mono- and dinuclear complexes thereof are known to promote hydrolysis of neutral phosphate and/or phosphonate esters. However, the available literature on the hydrolysis of phosphothiolate ($\text{P}=\text{S}$) esters and phosphothiolates is quite sparse with only the softer ions such as Cu^{2+} , Hg^{2+} and Pd^{2+} showing significant catalysis. The lack of examples may be due to reduced activity of $\text{P}=\text{S}$ esters, their poor aqueous solubility and the fact that anionic hydrolytic products bind to the metal ions thereby inhibiting further catalysis.

There is a need for a viable catalytic decontamination method which is inexpensive, has high catalyst turnover, and occurs at relatively neutral pH and ambient temperature, and most importantly, proceeds rapidly, e.g. $t_{1/2} < 1$ min.

BRIEF STATEMENT OF THE INVENTION

According to one aspect of the invention there is provided a method for decomposing an organophosphorus compound comprising subjecting said organophosphorus compound to an alcoholysis reaction in a medium comprising non-radioactive metal ions and at least a trace amount of alkoxide ions, wherein, through said alcoholysis reaction, said organophosphorus compound is decomposed.

In one embodiment of the invention, said organophosphorus compound has the following formula (10):



(10)

where:

J is O or S;

X, G, Z are the same or different and are selected from the group consisting of Q, OQ, QA, OA, F, Cl, Br, I, QS, SQ and C≡N;

Q is hydrogen or a substituted or unsubstituted branched, straight-chain or cyclic alkyl group having 1-100 carbon atoms; and

A is a substituted or unsubstituted aryl group selected from the group consisting of phenyl, biphenyl, benzyl, pyridyl, naphthyl, polynuclear aromatic, and aromatic and non-aromatic heterocyclic;

wherein, when X, G, Z are the same,

(i) X, G, Z are not Q; or

(ii) Q is not H; and

wherein said substituents are selected from the group consisting of Cl, Br, I, F, nitro, nitroso, Q, alkenyl, OQ, carboxyalkyl, acyl, SO₃H, SO₃Q, S=O(Q), S(=O)₂Q, amino, alkylamino (NHQ), arylamino (NHA), alkylaryl amino, dialkylamino and diarylamino.

In some embodiments, said medium is a solution further comprising a solvent selected from the group consisting of methanol, substituted and unsubstituted primary, secondary and tertiary alcohols, alkoxyalkanol, aminoalkanol, and combinations thereof.

In a preferred embodiment, said organophosphorus compound has at least one phosphorus atom double bonded to an oxygen or a sulfur atom.

In another embodiment, said medium further comprises a non-inhibitory buffering agent.

In yet another embodiment said buffering agent is selected from the group consisting of anilines, N-alkylanilines, N,N-dialkylanilines, N-alkylmorpholines, N-alkylimidazoles, 2,6-dialkylpyridines, primary, secondary and tertiary amines, trialkylamines, and combinations thereof.

In another embodiment, said medium is a solution further comprising a solvent selected from the group consisting of methanol, ethanol, n-propanol, *iso*-propanol, n-butanol, 2-butanol, methoxyethanol, and combinations thereof.

In further embodiments, said solution further comprises a solvent selected from the group consisting of nitriles, esters, ketones, amines, ethers, hydrocarbons, substituted hydrocarbons, unsubstituted hydrocarbons, chlorinated hydrocarbons, and combinations thereof.

In further embodiments, said medium further comprises alkoxide ions in addition to said at least a trace amount of alkoxide ions.

In further embodiments, the concentration of said alkoxide ions is about 0.1 to about 2 equivalents of the concentration of the metal ions.

In further embodiments, the concentration of said alkoxide ions is about 1 to about 1.5 equivalents of the concentration of the metal ions.

In further embodiments, said medium is prepared by combining a metal salt and an alkoxide salt with at least one of alcohol, alkoxyalkanol and aminoalkanol.

In further embodiments, said metal ions are selected from the group consisting of lanthanide series metal ions, transition metal ions, and combinations thereof.

In further embodiments, said metal ions are selected from the group consisting of lanthanide series metal ions, copper, platinum, palladium, zinc, nickel, yttrium, scandium ions, and combinations thereof.

In further embodiments, said metal ions are selected from the group consisting of Cu^{2+} , Pt^{2+} , Pd^{2+} , Zn^{2+} , Y^{3+} , Sc^{3+} , Ce^{3+} , La^{3+} , Pr^{3+} , Nd^{3+} , Sm^{3+} , Eu^{3+} , Gd^{3+} , Tb^{3+} , Dy^{3+} , Ho^{3+} , Er^{3+} , Tm^{3+} , Yb^{3+} , and combinations thereof.

In further embodiments, said metal ions are lanthanide series metal ions.

In further embodiments, said lanthanide series metal ions are selected from the group consisting of Ce^{3+} , La^{3+} , Pr^{3+} , Nd^{3+} , Sm^{3+} , Eu^{3+} , Gd^{3+} , Tb^{3+} , Dy^{3+} , Ho^{3+} , Er^{3+} , Tm^{3+} , Yb^{3+} , and combinations thereof.

In further embodiments, said metal ions are selected from the group consisting of Cu^{2+} , Pt^{2+} , Pd^{2+} , Zn^{2+} , and combinations thereof.

In further embodiments, said metal ions are selected from the group consisting of Y^{3+} , Sc^{3+} , and combinations thereof.

In further embodiments, said metal ion is La^{3+} .

In further embodiments, said organophosphorus compound is a pesticide.

In further embodiments, said organophosphorus compound is an insecticide.

In further embodiments, said organophosphorus compound is paraoxon.

In further embodiments, said organophosphorus compound is a chemical warfare agent.

In further embodiments, said organophosphorus compound is a G-agent.

In further embodiments, said organophosphorus compound is selected from the group consisting of VX and Russian-VX.

In further embodiments, said organophosphorus compound is a nerve agent.

In further embodiments, said chemical warfare agent is combined with a polymer.

In further embodiments, said medium further comprises one or more ligands.

In further embodiments, said ligand is selected from the group consisting of 2,2'-bipyridyl, 1,10-phenanthryl, 2,9-dimethylphenanthryl, crown ether, and 1,5,9-triazacyclododecyl.

In further embodiments, said ligand further comprises solid support material.

In further embodiments, said solid support material is selected from a polymer, silicate, aluminate, and combinations thereof.

In further embodiments, said medium is a solid.

In further embodiments, said medium is a solution.

In further embodiments, said solution is disposed on an applicator.

In further embodiments, the concentration of said alkoxide ions is about 0.5 to about 1.5 equivalents of the concentration of the metal ions.

In another broad aspect, the invention provides a kit for decomposing an organophosphorus compound comprising a substantially non-aqueous medium for an

alcoholysis reaction, said medium comprising non-radioactive metal ions and at least a trace amount of alkoxide ions.

In a first embodiment, said medium is contained in an ampule.

In a second embodiment, the kit comprises an applicator bearing the medium, said applicator being adapted so that the medium is applied to the organophosphorus compound and the compound decomposes.

In some embodiments, the kit further comprises written instructions for use.

BRIEF DESCRIPTION OF THE DRAWINGS

For a better understanding of the invention and to show more clearly how it may be carried into effect, reference will now be made by way of example to the accompanying drawings, which illustrate aspects and features according to preferred embodiments of the present invention, and in which:

Figure 1A shows a proposed mechanism for catalysis by a lanthanum methoxide dimer of the methanolysis of an aryl phosphate.

Figure 1B shows a proposed mechanism for catalysis by a zinc methoxide complex of the methanolysis of an aryl phosphate.

Figure 1C shows the reaction scheme for Cu:[12]aneN₃ catalyzing the methanolysis of fenitrothion.

Figure 2 shows a plot of k_{obs} vs. concentration of La(OTf)₃ for the La³⁺-catalyzed methanolysis of paraoxon (2.04×10^{-5} M) at 25 °C, where

■, ^spH 8.96;

○, ^spH 8.23; and

●, ^spH 7.72.

Figure 3 shows a plot of the $\log k_2^{\text{obs}}$ (M⁻¹s⁻¹) vs. ^spH for La³⁺-catalyzed methanolysis of paraoxon at 25 °C. The dotted line through the data was computed on the basis of a fit of the k_{obs} data to equation 3, the two dominant forms being La₂(OCH₃)₂ and La₂(OCH₃)₃.

Figure 4 shows a speciation diagram for the distribution of La₂(OCH₃)_n forms in methanol, $n = 1-5$, as a function of ^spH, calculated for [La(OTf)₃] = 2×10^{-3} M. Data

represented as (●) correspond to second order rate constants (k_2^{obs}) for La^{3+} -catalyzed methanolysis of paraoxon presented in Table 13.

Figure 5 shows a plot of the predicted k_2^{obs} vs. pH rate profile for La^{3+} -catalyzed methanolysis of paraoxon (-----) based on the kinetic contributions of $\text{La}_2(\text{OCH}_3)_1$, (.....); $\text{La}_2(\text{OCH}_3)_2$ (solid line) and $\text{La}_2(\text{OCH}_3)_3$, (-·-·-) computed from the $k_2^{2:1}$, $k_2^{2:2}$ and $k_2^{2:3}$ rate constants (Table 14), and their speciation (Figure 4); data points (■) are experimental k_2^{obs} rate constants from Table 13.

Figure 6 shows the effect of copper triflate (in the presence of equimolar ligand and 0.5 equivalents of methoxide) on the rate of methanolysis of fenitrothion as a plot of the k_{obs} vs. total concentration of $\text{Cu}(\text{OTf})_2$ for the methanolysis of fenitrothion catalyzed by various species at $T = 25^\circ\text{C}$ and $[\text{OCH}_3^-]/[\text{Cu}^{2+}]_t = 0.5$, when ligand is used, $[\text{Cu}^{2+}]_t = [\text{Ligand}]$, where

●, $\{\text{Cu}^{2+}:\text{no ligand}:(\text{OCH}_3^-)\}$;

◆, $\{\text{Cu}^{2+}:\text{phen}:(\text{OCH}_3^-)\}$; and

■, $\{\text{Cu}^{2+}:\text{bpy}:(\text{OCH}_3^-)\}$.

Figure 7 shows the effect of $\text{Cu}^{2+}:[12]\text{aneN}_3:(\text{OCH}_3^-)$ (copper triflate in the presence of equimolar ligand and 0.5 equivalents of methoxide) on the rate of methanolysis of paraoxon (●) and fenitrothion (■) as a plot of the k_{obs} vs. total concentration of $\text{Cu}(\text{OTf})_2$ conducted at $T = 25^\circ\text{C}$.

Figure 8 shows the effect of methoxide ion concentration on the rate of Zn^{2+} -catalyzed methanolysis of paraoxon as plots of k_{obs} vs added NaOCH_3 for the methanolysis of paraoxon in the presence of 1 mM $\text{Zn}(\text{ClO}_4)_2$, where:

●, no added ligand;

◇, 1 mM phen;

■, 1mM diMephen; and

□, 1 mM $[12]\text{aneN}_3$

(lines through the data drawn as visual aid only).

Figure 9A shows the catalyzed methanolysis of fenitrothion as a plot of k_{obs} vs. concentration of zinc ion ($\text{Zn}(\text{OTf})_2$) alone, and in the presence of equimolar ligand at constant $[(\text{OCH}_3^-)]/[\text{Zn}^{2+}]_{\text{total}}$ ratios, where:

●, no ligand, $[(\text{OCH}_3^-)]/[\text{Zn}^{2+}]_{\text{total}} = 0.3$;

O, phen, $[(\text{OCH}_3)]/[\text{Zn}^{2+}]_{\text{total}} = 0.5$; and

■, diMephen, $[(\text{OCH}_3)]/[\text{Zn}^{2+}]_{\text{total}} = 1.0$.

Figure 9B shows the catalyzed methanolysis of paraoxon as a plot of k_{obs} vs. concentration of zinc ion ($\text{Zn}(\text{OTf})_2$) alone and in the presence of equimolar ligand at constant $[(\text{OCH}_3)]/[\text{Zn}^{2+}]_{\text{total}}$ ratios, where:

●, no ligand, $[(\text{OCH}_3)]/[\text{Zn}^{2+}]_{\text{total}} = 0.3$;

O, phen, $[(\text{OCH}_3)]/[\text{Zn}^{2+}]_{\text{total}} = 0.5$; and

■, diMephen, $[(\text{OCH}_3)]/[\text{Zn}^{2+}]_{\text{total}} = 1.0$.

Figure 10 shows the disappearance of paraoxon (●) and appearance of diethyl methyl phosphate (■) product over time for a methanolysis reaction in the presence of zinc ion, methoxide, and ligand in deuterated methanol in a plot of relative signal integration of the reagent and product ^{31}P NMR signals for a system containing 15 mM paraoxon, 1 mM $\text{Zn}(\text{OTf})_2$, 1 mM NaOCH_3 and 1 mM diMephen at $T = 25^\circ\text{C}$.

Figure 11 shows the effect of increasing concentration of methoxide on the rate of Zn^{2+} -catalyzed methanolysis of paraoxon in a plot of the pseudo-first order rate constants (k_{obs}) for methanolysis of paraoxon in the presence of 1 mM $\text{Zn}(\text{OTf})_2$ and absence of added ligand as a function of added NaOCH_3 .

Figure 12 shows the effect of zinc ion concentration on the rate of Zn^{2+} -catalyzed methanolysis of paraoxon as plots of the k_{obs} for the methanolysis of fenitrothion (●), paraoxon (O) and p-nitrophenyl acetate (■) vs. $[\text{Zn}(\text{ClO}_4)_2]$ at a constant

$[\text{Zn}^{2+}:(\text{OCH}_3)]/[\text{Zn}^{2+}]_{\text{total}}$ ratio of 0.3, $T = 25^\circ\text{C}$. Lines through the data are calculated on the basis of fits to equation (6).

Figure 13A shows the effect of $\text{Zn}^{2+}:[12]\text{aneN}_3$ on the rate of Zn^{2+} -catalyzed methanolysis of paraoxon as a plot of k_{obs} for methanolysis of paraoxon as a function of $[\text{Zn}(\text{OTf})_2]_{\text{total}}$ containing equimolar $[12]\text{aneN}_3$ and NaOCH_3 , $T = 25^\circ\text{C}$. Right axis gives $[\text{Zn}^{2+}:[12]\text{aneN}_3:(\text{OCH}_3)]$ determined by Hyperquad™ fitting of titration data. The arrows are presented as a visual aid to connect the various species concentrations with the kinetic rate constant.

Figure 13B shows the effect of $\text{Zn}^{2+}:\text{phen}$ on the rate of Zn^{2+} -catalyzed methanolysis of paraoxon as a plot of k_{obs} for methanolysis of paraoxon as a function of $[\text{Zn}(\text{OTf})_2]_{\text{total}}$

containing equimolar phen and NaOCH_3 , $T = 25^\circ\text{C}$. Right axis gives $[\text{Zn}^{2+}:\text{phen}:(\text{OCH}_3)]$ determined by Hyperquad™ fitting of titration data. The arrows are presented as a visual aid to connect the various species concentrations with the kinetic rate constant.

Figure 14 shows the titration profiles obtained by potentiometric titration of 2mM $\text{Zn}(\text{OTf})_2$ with no added ligand (●), with 2 mM phen (◆), with 2 mM diMephen (■), with 2 mM $[\text{12}] \text{aneN}_3$ (□) and with 1.2 mM added HClO_4 . Lines through the titration curves with phen and $[\text{12}] \text{aneN}_3$ were derived from Hyperquad™ fitting of the data.

DETAILED DESCRIPTION OF THE INVENTION

According to a broad aspect of the invention there is provided a method of decomposing an organophosphorus compound by combining the organophosphorus compound with a substantially non-aqueous medium comprising alcohol, alkoxyalkanol or aminoalkanol, metal ions and at least a trace amount of alkoxide ions. When so combined the organophosphorus compound undergoes an alcoholysis reaction and forms a less toxic or non-toxic compound.

More particularly, the invention provides a method of increasing the rate of decomposition of an organophosphorus compound by combining the compound with a catalytic species formed in a substantially non-aqueous medium comprising metal ions; alcohol, alkoxyalkanol or aminoalkanol; and alkoxide ions. In some embodiments, the medium is a solution.

As used herein, the term “alcohol” means a compound which comprises an R-OH group, for example, methanol, primary alcohols, and substituted or unsubstituted secondary alcohols, tertiary alcohols, alkoxyalkanol, aminoalkanol, or a mixture thereof.

As used herein, “substantially non-aqueous medium” means an organic solvent, solution, mixture or polymer. As it is very difficult to obtain anhydrous alcohol, a person of ordinary skill in the art would recognize that trace amounts of water may be present. For example, absolute ethanol is much less common than 95% ethanol. However, the amount of alcohol present in a medium or solution according to the invention should not have so much water present as to inhibit the alcoholysis reaction, nor should a substantial amount of hydrolysis occur.

As used herein, the term “organophosphorus compound” includes compounds which comprise a phosphorus atom doubly bonded to an oxygen or a sulfur atom. In preferred

embodiments such organophosphorus compounds are deleterious to biological systems, for example, a compound may be an acetylcholine esterase inhibitor, a pesticide or a chemical warfare agent.

As used herein, the term "decomposing an organophosphorus compound" refers to rendering a deleterious organophosphorus compound into a less toxic or non-toxic form.

Decomposition of an organophosphorus compound according to the invention may be carried out in solution form, or in solid form. Examples of such decomposition include, applying catalyst as a solution directly to a solid chemical warfare agent or pesticide. Such a solution would be for example, an appropriately buffered alcoholic, alkoxyalkanolic or aminoalkanolic solution comprising metal ions and alkoxide ions, in which one or more catalytic species forms spontaneously, which may be applied to a surface which has been contacted with an organophosphorus agent.

As used herein, the term "catalytic species" means a molecule or molecules, comprising metal ions and alkoxide ions, whose presence in an alcoholic, alkoxyalkanolic or aminoalkanolic solvent containing an organophosphorus compound increases the rate of alcoholysis of the organophosphorus compound relative to its rate of alcoholysis in the solvent without the catalytic species.

As used herein, the term "appropriately buffered" means that the ^spH of a solution is controlled by adding non-inhibitory buffering agents, or by adding about 0.1 to about 2.0 equivalents of alkoxide ion per equivalent of metal ion.

As used herein, the term " ^spH " is used to indicate pH in a non-aqueous solution (Bosch *et al.*, 1999, Rived *et al.*, 1998, Bosch *et al.*, 1996). One skilled in the art will recognize that if a measuring electrode is calibrated with aqueous buffers and used to measure pH of an aqueous solution, the term ^wpH is used. If the electrode is calibrated in water and the 'pH' of a neat methanol solution is then measured, the term ^wpH is used, and if the latter reading is made, and a correction factor of 2.24 (in the case of methanol) is added, then the term ^spH is used.

As used herein, the term "non-inhibitory agent or compound" means that the agent or compound does not substantially diminish the rate of a catalyzed reaction when compared to the rate of the reaction in the absence thereof.

As used herein, the term "inhibitory agent or compound" means that the agent or compound does substantially diminish the rate of a catalyzed reaction when compared to the rate of the reaction in the absence thereof.

As used herein, the term "metal species" means a metal in an oxidation state of zero to 9.

As used herein, the term "mononuclear" or "monomeric" means a species comprising one metal atom.


In an embodiment of the invention, the catalytic species is a metal alkoxide species of the stoichiometry $\{M^{n+}(\text{OR})_m L_g\}_s$ where M is a metal selected from lanthanide series metals or transition metals; n is the charge on the metal which may be 1 to 9, most preferably 2 to 4; OR is alkoxide; m is the number of associated alkoxide ions and may be 1, 2, ..., $n-1$, n , $n+1$, $n+2$, ..., $n+6$, most preferably 1 to $n-1$; s is 1 to 100; L is ligand; g is the number of ligands complexed to the metal ion, and may be 0 to 9; where g is greater than 1, the ligands may be the same or different. Examples of this embodiment include the lanthanum dimer $\{\text{La}^{3+}(\text{OMe})\}_2$ and copper monomer $\{\text{Cu}^{2+}(\text{OMe})\text{L}\}$.

The inventors contemplate an embodiment wherein the oxidation state of the metal atom is zero. For example, it is well known in the art that transition metals having an oxidation state of zero may be reactive and may form complexes. Copper is an example of such a metal, and it is expected that Cu^0 may catalyze alcoholysis of organophosphorus compounds according to the invention.

As used herein, the term "ligand" means a species containing a donor atom or atoms that has a non-bonding lone pair or pairs of electrons which are donated to a metal centre to form one or more metal-ligand coordination bonds. In this way, ligands bond to coordination sites on a metal and thereby limit dimerization and prevent further oligomerization of the metal species, thus allowing a greater number of active mononuclear species to be present than is the case in the absence of ligand or ligands.

As used herein, the term " $\{M^{n+}:L:\text{OR}\}$ " (which differs from the above described system, $\{M^{n+}(\text{OR})_m L_g\}_s$, by the use of the symbol ":" between constituents of the brace "{ }") is used when no stoichiometry is defined for a system comprising metal ions (M^{n+}), ligand (L), and alkoxide (OR). This technique is meant to encompass any and all catalytically active stoichiometries thereof including but not limited to dimers, trimers and longer oligomers, monoalkoxides, dialkoxides, polyalkoxides, etc.

In another embodiment of the invention, the catalytic species has the general formula 20:



where Z^1 and Z^2 are the same or different non-radioactive lanthanide, copper, platinum or palladium ions;

R^1 , R^2 , R^3 and R^4 are each independently alkyl groups selected from a branched, cyclic or straight-chain hydrocarbon containing 1-12 carbon atoms, preferably 1-4 carbon atoms;

p is a number from 1-6; and

m and q are each independently zero or 1 or more, preferably 1-5, such that the dimer has a net charge of zero.

In another embodiment of the invention, the catalytic species has the general formula 20:

where Z^1 and Z^2 are the same or different non-radioactive lanthanide series metal ions, copper, platinum or palladium ions;

R^1 , R^2 , R^3 and R^4 are each independently alkyl groups selected from a branched, cyclic or straight-chain hydrocarbon containing 1-12 carbon atoms, preferably 1-4 carbon atoms;

p is a number from 1-6; and

m and q are each independently zero or 1 or more, preferably 1-5, such that the dimer has a net charge of zero.

In another embodiment of the invention, the catalytic species has the general formula 20:

where Z^1 and Z^2 are the same or different non-radioactive lanthanide series metal ions, and/or transition metal ions;

R^1 , R^2 , R^3 and R^4 are each independently alkyl groups selected from a branched, cyclic or straight-chain hydrocarbon containing 1-12 carbon atoms, preferably 1-4 carbon atoms;

p is a number from 0-6; and

m and q are each independently zero or 1 or more, preferably 1-5, such that the dimer has a net positive charge.

In another embodiment of the invention, the catalytic species has the general formula 20:


where Z^1 and Z^2 are the same or different non-radioactive lanthanide series metal ions, and/or transition metal ions;

R^1 , R^2 , R^3 and R^4 are each independently alkyl groups selected from a branched, cyclic or straight-chain hydrocarbon containing 1-12 carbon atoms, preferably 1-4 carbon atoms;

p is a number from 1-6; and

m and q are each independently zero or 1 or more, preferably 1-5, such that the dimer has a net positive charge.

In another embodiment of the invention, the catalytic species has the general formula 30:



where Z^1 is a non-radioactive lanthanide, copper, platinum or palladium ion;

R^2 and R^3 are each independently alkyl groups selected from a branched, cyclic or straight-chain hydrocarbon containing 1-12 carbon atoms, preferably 1-4 carbon atoms;

a is a number from 1-3; and

b is zero or 1 or more, such that the catalytic species has a net charge of zero.

In another embodiment of the invention, the catalytic species has the general formula 30:

where Z^1 is a non-radioactive lanthanide series metal ion or a transition metal ion;

R^2 and R^3 are each independently alkyl groups selected from a branched, cyclic or straight-chain hydrocarbon containing 1-12 carbon atoms, preferably 1-4 carbon atoms;


a is a number from 1-3; and

b is zero or 1 or more, such that the catalytic species has a net positive charge.

Another embodiment of the invention, the catalytic species has the general formula 30:

where Z^1 is a non-radioactive lanthanide series metal ion or a transition metal ion;
 R^2 and R^3 are each independently alkyl groups selected from a branched, cyclic or straight-chain hydrocarbon containing 1-12 carbon atoms, preferably 1-4 carbon atoms;
 a is a number from 1-3; and
 b is zero or 1 or more, such that the catalytic species has a net positive charge;
 wherein unoccupied coordination sites on the metal may be occupied by one or more ligands.

In another embodiment of the invention, the catalytic species has the general formula 40:



where Z^1 , Z^2 and Z^3 are the same or different non-radioactive lanthanide, copper, platinum or palladium ions;

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are each independently alkyl groups selected from a branched, cyclic or straight-chain hydrocarbon containing 1-12 carbon atoms, preferably 1-4 carbon atoms;

p is a number from 1-4;

m , d , q and t are each independently zero or 1 or more, preferably 1-5, such that the oligomer has a net charge of zero; and

r is a number from 0 to 100, or in the case of polymeric material may be greater than 100.

In yet another embodiment of the invention, the catalytic species has the general formula 40:

where Z^1 , Z^2 and Z^3 are the same or different non-radioactive lanthanide series metal ions, or transition metal ions or combinations thereof;

$R^1, R^2, R^3, R^4, R^5, R^6$ and R^7 are each independently alkyl groups selected from a branched, cyclic or straight-chain hydrocarbon containing 1-12 carbon atoms, preferably 1-4 carbon atoms;

p is a number from 1-4;

m, d, q and t are each independently zero or 1 or more, preferably 1-5, such that the oligomer has a net positive charge; and

r is a number from 0-100, or in the case of polymeric material may be greater than 100.

The alcoholic solution comprises a primary, secondary or tertiary alcohol, an alkoxyalkanol, an aminoalkanol, or a mixture thereof. In one embodiment, a non-inhibitory buffering agent is added to the solution to maintain the s_pH at the optimum range of s_pH , for example in the case of La^{3+} in methanol, s_pH 7 to 11 (see Figure 3). Examples of non-inhibitory buffering agents include: anilines; N-alkylanilines; N,N-dialkylanilines; N-alkylmorpholines; N-alkylimidazoles; 2,6-dialkylpyridines; primary, secondary and tertiary amines such as trialkylamines; and their various derivatives.

In another embodiment, non-inhibitory buffering agents are not added, but additional alkoxide ion is added in the form of an alkoxide salt to obtain metal ions and alkoxide ions in a metal:alkoxide ratio of about 1:0.01 to about 1:2, for some embodiments preferably about 1:1 to about 1:1.5, for other embodiments preferably about 1:0.5 to about 1:1.5. A person skilled in the art will recognize that an alcoholic solution contains trace amounts of alkoxide ions. This concept is analogous to water containing a trace amount of hydrogen ions and hydroxide ions, thus water of pH 7 contains, by definition, $[H^+] = 1 \times 10^{-7} M$ and $[OH^-] = 1 \times 10^{-7} M$. For this reason, when alkoxide salts are added according to this embodiment of the invention, they are referred to as "additional" alkoxide ions. Suitable non-inhibitory cations for the alkoxide salts include monovalent ions such as, for example, Na^+ , K^+ , Cs^+ , Rb^+ , NR_4^+ and $NR'R''R'''R''''^+$ (where R' , R'' , R''' , and R'''' may be the same or different and may be hydrogen or substituted or unsubstituted alkyl or aryl groups) and divalent ions such as the alkali earth metals, and combinations thereof. In some instances such ions may prolong the life of a catalyst by bonding to and, for example, precipitating, an inhibitory product of organophosphorus decomposition, an example of which is Ca^{2+} bonding to fluoride.

To obtain the metal ions, metal salts are added to the solution. Preferably, the metal ion is a non-radioactive lanthanide series metal ion. Suitable lanthanide series metal ions include, for example, Ce^{3+} , La^{3+} , Pr^{3+} , Nd^{3+} , Sm^{3+} , Eu^{3+} , Gd^{3+} , Tb^{3+} , Dy^{3+} , Ho^{3+} , Er^{3+} , Tm^{3+} and Yb^{3+} and combinations thereof or complexes thereof. Suitable non-lanthanide series metal ions include, for example, divalent transition metal ions such as, for example, Cu^{2+} , Pd^{2+} , Pt^{2+} ,

Zn^{2+} , and trivalent transition metal ions such as, for example, Sc^{3+} and Y^{3+} , as well as combinations thereof or complexes thereof, including combinations/complexes of those with non-radioactive lanthanide series metal ions. While La^{3+} ($\text{pK}_{\text{a}1} = 7.8$) has good catalytic efficacy from pH 7.3 to 10.3, other metal ions which have lower pK_{a} values (for example Ho^{3+} and Eu^{3+} have $\text{pK}_{\text{a}1}$ values of 6.6, while Yb^{3+} has a $\text{pK}_{\text{a}1}$ value of 5.3, Gibson *et al.* 2003) may be efficacious at lower pH .

An embodiment of the invention is a catalytic system comprising mixtures of metal ions, for example, mixtures of lanthanide series metal ions which would be active between the wide pH range of 5 to 11. Lanthanide series metal ions and alkoxide may form several species in solution, an example of which, species forming from La^{3+} and methoxide is shown in the figures. In the case of La^{3+} , a dimer containing 1 to 3 alkoxides is a particularly active catalyst for the degradation of organophosphorus compounds. In the case of non-lanthanide series metal ions, such as, for example Zn^{2+} and Cu^{2+} , a mononuclear complex containing alkoxides is an active catalyst for the degradation of organophosphorus compounds.

In some embodiments, the invention provides limiting of dimerization and prevention of further oligomerization by addition of ligand such as, for example, bidentate and tridentate ligands. By coordination at one or more sites on a metal, a ligand limits dimerization and prevents further oligomerization of a metal species, thus allowing a greater number of active mononuclear species than is the case in the absence of ligand. Although not meant to be limiting, examples of such ligands are 2,2'-bipyridyl ("bpy"), 1,10-phenanthryl ("phen"), 2,9-dimethylphenanthryl ("diMephen") and 1,5,9-triazacyclododecyl ("12]aneN₃"), crown ether, and their substituted forms. Such ligands may be attached via linkages to solid support structures such as polymers, silicates or aluminates to provide solid catalysts for the alcoholysis of organophosphorus compounds which are decomposed according to the invention. The point of attachment of the metal:ligand:alkoxide complex to the solid support is preferably at the 3 or 4 position in the case of bipyridyl or the 3, 4 or 5 position in the case of phenanthrolines using linking procedures and connecting spacers which are known in the art. In the case of aza ligands, such as, for example, 12]aneN₃, the point of attachment of the complex to the solid support would preferably be on one of the nitrogens of the macrocycle, using methods and connecting spacers known in the art. Such attachment to solid supports offers advantages in that the solid catalysts may be conveniently recovered from the reaction media by filtration or decantation. In an embodiment of the invention wherein ligands are attached to solid support structures, organophosphorus compounds may be decomposed by running a solution through a column such as a chromatography column. In another embodiment of the invention wherein ligands are attached to solid support

structures, organophosphorus compounds may be decomposed by contact with a polymer comprising metal species and alkoxide ions.


Suitable anions of the metal salts are non-inhibitory or substantially non-inhibitory and include, for example, ClO_4^- , BF_4^- , BR_4^- , I^- , Br^- , CF_3SO_3^- (also referred to herein as "triflate" or "OTf") and combinations thereof. Preferred anions are ClO_4^- and CF_3SO_3^- . In the case of BF_4^- , a solvent other than methanol is preferred.

The solution comprises solvents, wherein preferred solvents are alcohols, including primary and secondary alcohols such as methanol, ethanol, n-propanol, *iso*-propanol, n-butanol, 2-butanol and methoxyethanol, and combinations thereof. Most preferably the solution is all alcohol or all alkoxyalkanol or all aminoalkanol; however, combinations with non-aqueous non-inhibitory solvents can also be used, including, for example, nitriles, ketones, amines, ethers, hydrocarbons including chlorinated hydrocarbons and esters. In the case of esters, it is preferable that the alkoxy group is the same as the conjugate base of the solvent alcohol. In some embodiments, esters may cause side reactions which may be inhibitory.

Initial studies have been undertaken in methanol since methanol is closest to water in terms of structure and chemical properties and is readily available. However, methanol is less desirable than other solvents due to its toxicity and its relatively low boiling point of 64.7 °C which makes it volatile and prone to evaporation from open vessels. For these reasons, use of higher alcohols such as ethanol, n-propanol and *iso*-propanol has been explored (see Examples 1 and 2). Ethanol, n-propanol and *iso*-propanol are substantially less volatile (boiling points 78, 97.2 and 82.5 °C respectively), are less toxic, and have better solubilizing characteristics for hydrophilic substrates. The higher boiling points mean that these solvents are more amenable to field conditions since there would conveniently be less evaporation and thus less solvent would be lost to the atmosphere.

Other preferred solvents include n-butanol and 2-butanol since they have higher boiling points than the lower alcohols.

In accordance with the invention, the metal ion species catalyzes an alcoholysis reaction of an organophosphorus compound or a mixture of organophosphorus compounds represented by the following general formula (10):



(10)

where P is phosphorus;

J is O (oxygen) or S (sulfur);

X, G, Z are the same or different and are selected from the group consisting of Q, OQ, QA, OA, F (fluoride), Cl (chloride), Br (bromide), I (iodide), QS, SQ and C≡N;

where Q is hydrogen or a substituted or unsubstituted branched, straight-chain or cyclic alkyl group consisting of 1-100 carbon atoms; wherein when X, G, Z are the same, X, G, Z are not Q, and when X, G, Z are the same Q is not H;

A is a mono-, di-, or poly-substituted or unsubstituted aryl group selected from phenyl, biphenyl, benzyl, pyridine, naphthyl, polynuclear aromatics, and 5- and 6-membered aromatic and non-aromatic heterocycles;

wherein each said substituent is selected from Cl, Br, I, F, nitro, nitroso, Q, alkenyl, OQ, carboxyalkyl, acyl, SO₃H, SO₃Q, S=O(Q), S(=O)₂Q, amino, alkylamino (NHQ), arylamino (NHA), alkylarylamino, dialkylamino and diarylamino.

Most preferably, the phosphorus atom of figure 10 has at least one good leaving group attached. For this reason, organophosphorus compounds which are decomposed according to the invention do not have three alkyl groups, nor three hydrogens, nor three hydroxyl groups attached. One skilled in the art will recognize that a "good leaving group" is a substituent with an unshared electron pair that readily departs from the substrate in a nucleophilic substitution reaction. The best leaving groups are those that become either a relatively stable anion or a neutral molecule when they depart, because they cause a stabilization of the transition state. Also, leaving groups that become weak bases when they depart are good leaving groups. Good leaving groups include halogens, alkanesulfonates, alkyl sulfates, and p-toluenesulfonates.


As used herein, the term "heterocycle" means a substituted or unsubstituted 5- or 6-membered aromatic or non-aromatic hydrocarbon ring containing one or more O, S or N atoms, or polynuclear aromatic heterocycle containing one or more N, O, or S atoms.

An advantage of the decomposition method of the invention is that the solvent, being hydrophobic, relative to water, permits good solubility of organophosphorus agents such as


VX, Russian-VX, tabun (GA), soman (GD), sarin (GB), GF, hydrophobic polymers, insecticides and pesticides.

Another advantage of the invention is that it provides a non-aqueous solution and reaction products that can be easily and safely disposed of by incineration. It will thus be appreciated that the decontamination method of the invention can be used for a broad range of chemical warfare agents, or mixtures of such agents, or blends of such agents with polymers, as well as other toxic compounds such as insecticides, pesticides and related organophosphorus agents in general.


A further advantage of the invention is that destruction of organophosphorus agents occurs with or without the addition of heat. An ambient temperature reaction is cost-efficient for large scale destruction of stockpiled organophosphorus material such as chemical weapons, insecticides or pesticides. The catalyst species can catalyze the alcoholysis over the full temperature range between the freezing and boiling points of the solvents or mixture of solvents used.



Paraoxon



O,O'-diethyl-S-p-nitrophenylphosphothioate



Fenitrothion

The G-type and V-type classes of chemical warfare agents are too toxic to be handled without specialized facilities and are often modeled by simulants such as, for the G-agents: paraoxon and *p*-nitrophenyl diphenyl phosphate, and for the V-agents: O,S-dialkyl- or O,S-arylalkyl-phosphonothioates or S-alkyl-phosphinothioates or S-aryl-phosphinothioates (Yang, 1999). We have used three such simulants and report herein, degradation of paraoxon as a model of G-agents, degradation of O,O'-diethyl-S-p-nitrophenylphosphorothioate as a model of V-agents, and degradation of fenitrothion as a model of (P=S)-containing pesticides. Structures for these model compounds are shown below. These three compounds were chosen because each possess a chromophore which makes the UV-vis kinetics simpler to study with low concentrations of materials. It is expected that this invention has wide applicability for other organophosphorus compounds

including chemical warfare agents and other pesticides such as, for example, parathion and malathion.

In our studies, which are detailed in the following examples, we have: confirmed the degradation of paraoxon, O,O'-diethyl-S-p-nitrophenylphosphorothioate and fenitrothion when placed in an alcoholic solution of metal ions and at least a trace amount of alkoxide ions; determined the rate of the decomposition of paraoxon in a methanol solution containing La^{3+} and additional methoxide ions; characterized stoichiometry and proposed a structure of active $\{\text{La}^{3+}(\text{OCH}_3)\}_2$ dimers; studied catalyzed alcoholysis in the presence of ligand and determined that faster rates are possible in some such systems relative to catalysis in the absence of ligand; and confirmed the complete destruction of paraoxon and O,O'-diethyl-S-p-nitrophenylphosphorothioate relative to catalyst in $\{\text{La}^{3+}:\text{OMe}\}$, $\{\text{Cu}^{2+}:\text{OMe}\}$, and $\{\text{Zn}^{2+}:\text{OMe}\}$ systems thus confirming the true catalytic nature of this method.

The data presented in the following examples support the following conclusions:

Destruction of Paraoxon (Model G Agent): A preferred embodiment for methanolysis of paraoxon is a $\{\text{La}^{3+}:\text{OCH}_3\}$ system according to the invention. The procedure involves preparation of a 2 mM $\text{La}(\text{OTf})_3$ methanolic solution, containing equimolar NaOCH_3 which affords a 10^9 -fold acceleration of the methanolysis of paraoxon relative to the background reaction at the same pH in the absence of catalyst ($t_{1/2} \sim 20$ sec). A second preferred embodiment for the methanolysis of paraoxon is a $\{\text{Zn}^{2+}:\text{diMephen}:\text{OMe}\}$ system. This system affords accelerations of up to 1.8×10^6 -fold for the methanolysis of paraoxon and has broader applicability than La^{3+} as Zn^{2+} also catalyzes the decomposition of fenitrothion.

Destruction of O,O'-diethyl-S-p-nitrophenylphosphorothioate (Model V Agent):

A preferred embodiment for methanolysis of O,O'-diethyl-S-p-nitrophenylphosphorothioate is a $\{\text{Cu}^{2+}:\text{OCH}_3:[12]\text{andN}_3\}$ system. A second preferred embodiment for the methanolysis of O,O'-diethyl-S-p-nitrophenylphosphorothioate is methanolic solution of $\{\text{Zn}^{2+}:\text{diMephen}:\text{OCH}_3\}$. A third preferred embodiment for the methanolysis of O,O'-diethyl-S-p-nitrophenylphosphorothioate is a methanolic solution of $\{\text{La}^{3+}:\text{OCH}_3\}$.

Destruction of Fenitrothion (Model Pesticide): A preferred embodiment for methanolysis of fenitrothion is a $\{\text{Cu}^{2+}:[12]\text{aneN}_3:\text{OCH}_3\}$ system according to the invention. The procedure involves preparation of a 2 mM $\text{Cu}(\text{OTf})_2$ methanolic solution containing 0.5 equivalents of $\text{N}(\text{Bu})_4\text{OCH}_3$ and 1 equivalent of $[12]\text{aneN}_3$ which catalyzes the methanolysis of fenitrothion with a $t_{1/2}$ of ~ 58 sec accounting for a 1.7×10^9 -fold acceleration of the

reaction at near neutral pH (8.75). A second preferred embodiment for the methanolysis of fenitrothion is a $\{Zn^{2+}:diMephen:OCH_3\}$ system. This system affords accelerations of 13×10^6 -fold for the methanolysis of fenitrothion at 2 mM each of $Zn(OTf)_2$, ligand diMephen and $NaOCH_3$ and exhibits broad applicability as it also catalyzes the decomposition of paraoxon. Fenitrothion decomposition is not appreciably accelerated in the presence of a La^{3+} system according to the invention. This points out the importance of matching the relative hard/soft characteristics of catalyst and substrate, and suggests that softer metal ions such as Cu^{2+} and Pd^{2+} could show enhanced catalytic activity toward the methanolysis of sulfur-containing phosphorus species.

Destruction of a Suspected Organophosphorus Compound of Unknown

Structure: A preferred embodiment of the invention for catalyzed alcoholysis of an unknown agent which is suspected to be an organophosphorus compound, is a mixture of $\{M^{3+}:OCH_3\}$ and $\{M^{2+}:L:OCH_3\}$ in an alcohol solution. Examples of such a mixture include $\{La^{3+}:OCH_3\}$ and $\{Cu^{2+}:[12]aneN_3:OCH_3\}$; and $\{La^{3+}:OCH_3\}$ and $\{Zn^{2+}:diMephen:OCH_3\}$. Although such a M^{2+} system is less reactive toward paraoxon than the M^{3+} system; unlike M^{3+} , the M^{2+} system does catalyze alcoholysis of fenitrothion. This mixture produces an effective method for destruction of both $P=S$ pesticides and $P=O$ chemical warfare agents.

The invention also provides a kit for decomposing an organophosphorus compound comprising a substantially non-aqueous medium for an alcoholysis reaction, said medium comprising non-radioactive metal ions and at least a trace amount of alkoxide ions. The kit may include a container, e.g., an ampule, which is opened so that the medium can be applied to the organophosphorus compound. Alternatively, the kit may include an applicator bearing the medium, wherein the applicator is adapted so that the medium is applied to the organophosphorus compound and the compound consequently decomposes. The applicator may comprise a moist cloth, i.e., a cloth bearing a solution according to the invention. The applicator may be a sprayer which sprays medium according to the invention on the organophosphorus compound. In some embodiments, the kit comprises written instructions for use to decompose an organophosphorus compound.

The following examples further illustrate the present invention and are not intended to be limiting in any respect. All scientific and patent publications cited herein are hereby incorporated by reference in their entirety.

Examples

Examples 5 to 8 provide a summary of the La^{3+} ion catalyzed alcoholysis of paraoxon. Example 10 is a prophetic example of an La^{3+} ion catalyzed alcoholysis of VX. Due to the fact that the dimeric lanthanum methoxide catalyst is stable in solution, and the reaction takes place at room temperature and at neutral pH (neutral ^spH in methanol is ~ 8.4), we expect that this reaction is amenable to scale-up and to use in the field.

In the examples, methanol (99.8 % anhydrous), sodium methoxide (0.5 M solution in methanol), $\text{La}(\text{CF}_3\text{SO}_3)_3$ and paraoxon were purchased from Sigma-Aldrich (St. Louis, Missouri) and used without any further purification. HClO_4 (70% aqueous solution) was purchased from BDH (Dorset, England). ^1H NMR and ^{31}P NMR spectra were determined at 400 MHz and 161.97 MHz. ^{31}P NMR spectra were referenced to an external standard of 70% phosphoric acid in water, and up-field chemical shifts are negative.

In the examples, the CH_3OH_2^+ concentration was determined using a Radiometer Vit 90 Autotitrator, equipped with a Radiometer GK2322 combination (glass/calomel) electrode calibrated with Fisher Certified Standard aqueous buffers (pH = 4.00 and 10.00) as described in recent papers (Neverov *et al* 2000; Neverov *et al.*, 2001(a); Neverov *et al.*, 2001(b); Neverov *et al.*, 2001(c); Brown *et al.*, 2002; Tsang *et al.*, 2003). Values of ^spH were calculated by adding a correction constant of 2.24 to the experimental meter reading as reported by Bosch *et al.*, 1999.

The $^s\text{pK}_a$ values of buffers used in the examples were obtained from the literature or measured at half neutralization of the bases with 70% HClO_4 in MeOH.

Example 1. M^{n+} -Catalyzed Ethanolysis Of Paraoxon And Fenitrothion: Reaction Conditions And Rates

The ethanolysis of fenitrothion and paraoxon was studied in ethanol using various metal ions with varying amounts of added base. These reactions were followed by UV-vis spectroscopy by observing the rate of disappearance of a starting material signal or the rate of appearance of a product signal such as 4-nitrophenol in the case of paraoxon or 3-methyl-4-nitrophenol in the case of fenitrothion. Reaction conditions and the catalyzed reaction's rate constants are summarized in Table 1.

Table 1 -- Maximum pseudo-first order kinetic rate constants for the ethanolysis of fenitrothion and paraoxon catalyzed by metal ions (0.001M) in the presence of optimum amount of base (max k_{obs}) and at equimolar amount (k_{obs} 1:1 OCH_3/M^x ratio), $T = 25^\circ\text{C}$.

Metals ^a	Paraoxon		Fenitrothion
	$10^4 \text{ Max } k_{\text{obs}}, \text{s}^{-1}$	$10^4 k_{\text{obs}}, \text{s}^{-1} \text{ }^b$	$10^4 k_{\text{obs}}, \text{s}^{-1} \text{ }^b$
Lanthanides			
La ³⁺	544.15 (1:1)	544.15	No catalysis
Pr ³⁺	253.24 (1:1)	253.24	No catalysis
Nd ³⁺	247.59 (1:1)	247.59	No catalysis
Gd ³⁺	220.14 (1:1)	220.14	No catalysis
Sm ³⁺	185.88 (1:1)	185.88	No catalysis
Eu ³⁺	160.0(1:1)	160	No catalysis
Tb ³⁺	146.34 (1:1)	146.34	No catalysis
Ho ³⁺	99.72 (1:1)	99.72	No catalysis
Dy ³⁺	63.65 (1:1)	63.65	No catalysis
Er ³⁺	62.61 (1:1)	62.61	No catalysis
Tm ³⁺	49.34 (1:1)	49.34	No catalysis
Transition Metals			
Zn ²⁺	48.22 (1:0:5)	37.28	5.42
Y ³⁺	32.56 (1:1)	32.56	No catalysis
Co ²⁺	25.70 (1:0:5)	Catalysis, rate unknown ^c	Catalysis, rate unknown ^c
Yb ³⁺	25.73(1:1)	25.73	No catalysis
Ni ²⁺	23.63 (1:0:5)	12.18	No catalysis
Cu ²⁺	No catalysis	No catalysis	Catalysis, rate unknown ^c
Sc ³⁺	No catalysis	No catalysis	No catalysis

^a Introduced as commercially available triflate salts and used as received

^b 0.001 M in each of M^{n+} salt and added NaOCH_3

^c Product formation was observed by final UV-vis spectra, but determination of exact value of the rate constant was not possible due to high absorbance of the solutions.

Example 2. La³⁺ And Zn²⁺-Catalyzed Solvolysis of Paraoxon in Propanols:

Kinetics and NMR Studies

The solvolysis of paraoxon was studied in two alcohols that are less polar than methanol, namely 1-propanol and 2-propanol. In the case of 1-propanol, kinetics were monitored by UV-vis spectroscopic techniques following the appearance of the product of the solvolysis, 4-nitrophenol, at $\lambda=335$ nanometers. For example, at a concentration of $\text{La}(\text{OTf})_3 = 0.5\text{mM}$ = concentration of NaOCH_3 , in the absence of any ligand, catalyzed solvolysis of paraoxon proceeded with a pseudo-first order rate constant of $2.1 \times 10^{-4} \text{ s}^{-1}$. At a concentration of $\text{Zn}(\text{OTf})_2 = 0.5\text{mM}$ = concentration of NaOCH_3 , in the presence of equimolar diMephen, the catalyzed solvolysis of paraoxon proceeded with a pseudo-first rate constant of $1.93 \times 10^{-4} \text{ s}^{-1}$.

The true catalytic nature of the system was demonstrated in the following Nuclear Magnetic Resonance (NMR) studies. To 2.5 mL of a solution of 1-propanol containing 5% methanol, and 0.5mM each of $\text{Zn}(\text{OTf})_2$, diMephen and NaOMe was added 8.3 μL of paraoxon so that the latter's total concentration was 15.4 mM. The alcoholic solution was then incubated at room temperature for 72 hours after which the ^{31}P NMR spectrum was recorded. This spectrum showed complete disappearance of the paraoxon starting material and complete formation of diethyl methyl phosphate (product of reaction with methanol) ($\delta = -0.3\text{ppm}$) and diethyl 1-propyl phosphate (product of reaction with 1-propanol) ($\delta = -1.23\text{ppm}$). This indicates true catalysis with more than 30 turnovers in 72 hr. The solvents were removed, and the residues dissolved in deuterated methanol- d_4 and the ^1H NMR spectra were recorded showing the presence of the products: 4-nitrophenol, diethyl methyl phosphate and diethyl 1-propyl phosphates. Similarly, an NMR study was done such that 2.5 mL of 2-propanol containing 5% methanol, 0.5mM each of $\text{Zn}(\text{OTf})_2$, diMephen and NaOMe was added 8.3 μL of paraoxon so that the latter's total concentration was 15.4 mM. The alcoholic solution was then incubated at room temperature for 72 hours after which the ^{31}P NMR spectrum was recorded. This spectrum showed complete disappearance of the paraoxon starting material and complete formation of diethyl methyl phosphate (product of reaction with methanol) ($\delta = -0.3\text{ppm}$) and diethyl 2-propyl phosphate (product of reaction with 2-propanol) ($\delta = -2.4\text{ppm}$) was observed and formation of the products 4-nitrophenol, diethyl methyl phosphate and diethyl 2-propyl phosphate were confirmed by ^1H NMR.

The ratio of the two phosphate products from each of the propanol solvents was determined from their ^{31}P NMR spectra and were found to be:


MeOH reaction product : Propanol reaction product

1-propanol reaction	1 : 2.8
2-propanol reaction	2.2 : 1.

These ratios show that if the medium for catalysis according to the invention is a mixture of alcohol, alkoxyalkanol and aminoalkanol, the reaction will select for the least hindered one. This factor may determine what an "effective amount" of methanol will be for a given system.

Example 3. La^{3+} -Catalyzed Methanolysis of Paraoxon: Experimental Details

Paraoxon, when placed in an appropriately buffered methanol solution containing La^{3+} ions held in a pH region between 7 and 11, underwent rapid methanolysis at ambient temperature to produce diethyl methyl phosphate and p-nitrophenol. A detailed reaction scheme is given in Scheme 1.

Scheme 1

To two mL of dry methanol at ambient temperature was added N-ethylmorpholine (25.5 μL or 23 mg) half neutralized with 11.4 M HClO_4 (8.6 μL) so that the final total buffer concentration was 0.1 M. To this was added 16.0 mg of paraoxon. The ^{31}P NMR spectrum showed a single signal at δ -6.35 ppm. To the resulting mixture was added 12.9 mg of $\text{La}(\text{O}_3\text{SCF}_3)_3$ and 40 μL of 0.5 M NaOCH_3 in methanol solution. At this point the concentration of paraoxon was 0.057 M and that of $\text{La}(\text{O}_3\text{SCF}_3)_3$ was 0.011 M and the measured pH of the methanol solution was 8.75, essentially neutrality. This solution was allowed to stand for 10 minutes, after which time the ^{31}P NMR spectrum indicated complete disappearance of the paraoxon signal and the appearance of a new signal at δ 0.733 ppm corresponding to diethyl methyl phosphate. The ^1H NMR spectrum indicated complete disappearance of the starting material and full release of free p-nitrophenol.

Example 4. La^{3+} -Catalyzed Methanolysis of G-agent: A Prophetic Example

To 200 mL of methanol is added 2.55 mL of N-ethylmorpholine (2.3 g) and 0.86 mL of 11.4 M HClO_4 to bring the total buffer concentration to 0.1 M. To this solution is added 1.29 g of $\text{La}(\text{O}_3\text{SCF}_3)_3$ and 4 mL of a 0.5 M solution of NaOCH_3 in methanol.

To the above solution is added 2 g of the G-agent Sarin (0.016 moles, 0.08 M) and the solution is allowed to stand at ambient temperature for 15 minutes. It is expected that analysis of the resulting solution would indicate substantially complete disappearance of Sarin. This reaction may be inhibited by F^- in which case Ca^{2+} may be added to the reaction solution to precipitate this inhibitory product.

Example 5. La^{3+} -Catalyzed Methanolysis of Paraaxon: Kinetics

The kinetics of the alcoholysis degradation reaction have been thoroughly investigated using the pesticide paraoxon. For methanolysis with dimeric lanthanum catalysts at 25°C, as little as 10^{-3} M of the catalytic specie(s) promotes the methanolysis reaction by $\sim 10^9$ -fold relative to the background reaction at a neutral ^spH of ~ 8.5 . The uncatalyzed methoxide-promoted reaction of paraoxon proceeds with the second order rate constant, $k_2^{\text{OCH}_3}$ of $0.011 \text{ M}^{-1}\text{s}^{-1}$ determined from concentrations of NaOCH_3 between 1×10^{-2} M and 4×10^{-2} M. Methanolysis of paraoxon is markedly accelerated in the presence of La^{3+} with an observed second order rate constant, k_2^{obs} of $\sim 17.5 \text{ M}^{-1}\text{s}^{-1}$ at the near neutral ^spH of 8.23. Assuming that the methoxide reaction persists at ^spH 8.23, the acceleration afforded to the methanolysis of paraoxon at that ^spH by a 2×10^{-3} M solution of $\text{La}(\text{O}_3\text{SCF}_3)_3$ is 1.1×10^9 -fold giving a half-life time of 20 seconds. The acceleration is 2.3×10^9 -fold at ^spH 7.72 and 2.7×10^8 -fold at ^spH 8.96.

UV kinetics of the methanolysis of paraoxon were monitored at 25 °C by observing the rate of loss of paraoxon at 268 nm or by the rate of appearance of *p*-nitrophenol at 313 nm or 328 nm at a concentration of paraoxon = 2.04×10^{-5} M using an OLIS®-modified Cary 17 UV-vis spectrophotometer. The concentration of $\text{La}(\text{O}_3\text{SCF}_3)_3$ was varied from 8×10^{-6} M to 4.8×10^{-3} M. All reactions were followed to at least three half-times and found to exhibit good pseudo-first order rate behavior. The pseudo-first order rate constants (k_{obs}) were evaluated by fitting the Absorbance vs. time traces to a standard exponential model.

The kinetics were determined under buffered conditions. Buffers were prepared from N, N-dimethylaniline ($^s\text{pK}_a = 5.00$), 2,6-lutidine ($^s\text{pK}_a = 6.70$), N-methylimidazole ($^s\text{pK}_a = 7.60$), N-ethylmorpholine ($^s\text{pK}_a = 8.60$) and triethylamine ($^s\text{pK}_a = 10.78$). Due to the fact that added counterions can ion-pair with La^{3+} ions and affect its speciation in solution, ionic strength was controlled through neutralization of the buffer and not by added salts. The total concentration of buffer varied between $7 \times 10^{-3} \text{ M}$ and $3 \times 10^{-2} \text{ M}$, and the buffers were partially neutralized with 70 % HClO_4 to keep the concentration of ClO_4^- at a low but constant value of $5 \times 10^{-3} \text{ M}$ which leads to a reasonably constant ionic strength in solution. With the concentration of $\text{La}^{3+} > 5 \times 10^{-4} \text{ M}$ at $^s\text{pH} > 7.0$, the metal ion was partially neutralized by adding an appropriate amount of NaOMe to help control the ^spH at the desired value. ^spH measurements were performed before and after each experiment and in all cases the values were consistent to within 0.1 units.

Shown in Figure 2 are three representative plots of the pseudo-first order rate constants (k_{obs}) for methanolysis of paraoxon as a function of added concentration of $\text{La}(\text{O}_3\text{SCF}_3)_3$ at ^spH 7.72, 8.23 and 8.96. (For original k_{obs} vs. concentration of La^{3+} kinetic data see Tables 2-12).

Table 2 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5} \text{ M}$) at 25°C ; ^spH 5.15 [dimethylaniline buffer] = $1.00 \times 10^{-2} \text{ M}$, $\lambda = 328 \text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{ M}$	$k_{\text{obs}}, \text{ s}^{-1}$
4.00E-05	3.11E-07
6.00E-05	5.46E-07
8.00E-05	4.90E-07
2.00E-04	1.17E-05
4.00E-04	2.46E-05
6.00E-04	3.78E-05
8.00E-04	5.34E-05
1.00E-03	6.13E-05
1.20E-03	7.72E-05

Table 3 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5}\text{M}$) at 25°C ; $\text{pH } 5.58$ [dimethylaniline buffer] = $2.00 \times 10^{-2}\text{M}$, $\lambda = 328\text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{M}$	$k_{\text{obs}}, \text{s}^{-1}$
4.00E-05	5.37E-06
6.00E-05	6.23E-06
8.00E-05	5.63E-06
2.00E-04	8.33E-06
4.00E-04	4.28E-05
6.00E-04	6.93E-05
8.00E-04	9.48E-05
1.00E-03	1.05E-04
1.20E-03	1.26E-04

Table 4 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5}\text{M}$) at 25°C ; $\text{pH } 5.82$ [dimethylaniline buffer] = $2.93 \times 10^{-2}\text{M}$, $\lambda = 328\text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{M}$	$k_{\text{obs}}, \text{s}^{-1}$
4.00E-05	1.15E-06
6.00E-05	1.71E-06
8.00E-05	2.52E-06
2.00E-04	3.13E-05
4.00E-04	7.11E-05
6.00E-04	1.15E-04
8.00E-04	1.92E-04
1.00E-03	2.17E-04
1.20E-03	3.07E-04

Table 5 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5} \text{ M}$) at 25°C ; $\text{pH } 6.69$ [2,6-Lutidine buffer] = $6.61 \times 10^{-3} \text{ M}$, $\lambda = 313 \text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{ M}$	$k_{\text{obs}}, \text{ s}^{-1}$
4.00E-05	1.18E-05
6.00E-05	3.13E-05
8.00E-05	4.43E-05
2.00E-04	1.21E-04
4.00E-04	3.04E-04
6.00E-04	5.24E-04
8.00E-04	8.00E-04
1.00E-03	9.31E-04
1.20E-03	1.18E-03

Table 6 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5} \text{ M}$) at 25°C , $\text{pH } 7.10$ [2,6-Lutidine buffer] = $1.00 \times 10^{-2} \text{ M}$, $\lambda = 313 \text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{ M}$	$k_{\text{obs}}, \text{ s}^{-1}$
4.00E-05	2.58E-05
6.00E-05	4.86E-05
8.00E-05	6.68E-05
2.00E-04	2.62E-04
4.00E-04	7.22E-04
6.00E-04	1.26E-03
8.00E-04	1.88E-03
1.00E-03	2.14E-03
1.20E-03	2.67E-03

Table 7 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5} \text{ M}$) at 25°C ; $\text{pH } 7.30$ [*N*-methylimidazole buffer] = $6.67 \times 10^{-3} \text{ M}$, $\lambda = 268 \text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{ M}$	$k_{\text{obs}}, \text{ s}^{-1}$
8.00E-06	3.83E-05
2.00E-05	1.50E-05
8.00E-05	7.95E-05
2.00E-04	7.17E-04
4.00E-04	1.58E-03
8.00E-04	3.97E-03
1.60E-03	8.45E-03
3.20E-03	1.70E-02
4.80E-03	2.28E-02

Table 8 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5} \text{ M}$) at 25°C ; $\text{pH } 7.72$ [*N*-methylimidazole buffer] = $1.00 \times 10^{-2} \text{ M}$, $\lambda = 268 \text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{ M}$	$k_{\text{obs}}, \text{ s}^{-1}$
2.00E-05	2.83E-06
8.00E-05	1.18E-04
2.00E-04	9.30E-04
4.00E-04	3.49E-03
6.00E-04	6.10E-03
8.00E-04	8.46E-03
1.20E-03	1.22E-02
1.60E-03	1.51E-02

Table 9 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5} \text{ M}$) at 25°C ; $\text{pH } 8.23$ [*N*-methylimidazole buffer] = $2.00 \times 10^{-2} \text{ M}$, $\lambda = 268 \text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{ M}$	$k_{\text{obs}}, \text{ s}^{-1}$
4.00E-05	5.08E-05
6.00E-05	9.74E-05
8.00E-05	1.63E-04
2.00E-04	1.94E-03
4.00E-04	5.65E-03
6.00E-04	1.01E-02
8.00E-04	1.26E-02
1.00E-03	1.66E-02
1.20E-03	1.98E-02

Table 10 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5} \text{ M}$) at 25°C ; $\text{pH } 8.96$ [*N*-ethylmorpholine buffer] = $2.00 \times 10^{-2} \text{ M}$, $\lambda = 268 \text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{ M}$	$k_{\text{obs}}, \text{ s}^{-1}$
4.00E-05	8.50E-05
6.00E-05	2.03E-04
8.00E-05	3.75E-04
2.00E-04	2.70E-03
4.00E-04	8.25E-03
6.00E-04	1.38E-02
8.00E-04	1.76E-02
1.00E-03	2.14E-02
1.20E-03	2.65E-02

Table 11 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5}\text{M}$) at 25°C ; ^spH 10.34 [triethylamine buffer] = $6.67 \times 10^{-3}\text{M}$, $\lambda = 268\text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{M}$	$k_{\text{obs}}, \text{s}^{-1}$
4.00E-05	1.75E-04
6.00E-05	4.52E-04
8.00E-05	1.43E-03
2.00E-04	4.75E-03
4.00E-04	8.08E-03
6.00E-04	1.10E-02
8.00E-04	1.28E-02
1.00E-03	1.42E-02
1.20E-03	1.66E-02

Table 12 -- Observed pseudo-first order rate constants for La^{3+} catalyzed methanolysis of paraoxon ($2.04 \times 10^{-5}\text{M}$) at 25°C ; ^spH 10.97 [triethylamine buffer] = $1.00 \times 10^{-2}\text{M}$, $\lambda = 268\text{ nm}$.

$\text{La}(\text{O}_3\text{SCF}_3)_3, \text{M}$	$k_{\text{obs}}, \text{s}^{-1}$
4.00E-05	1.60E-04
6.00E-05	3.98E-04
8.00E-05	5.21E-04
2.00E-04	3.49E-03
4.00E-04	5.42E-03
6.00E-04	6.23E-03
8.00E-04	7.57E-03
1.00E-03	8.17E-03
1.20E-03	9.15E-03

As was observed in our earlier studies of the La^{3+} -catalyzed methanolysis of esters (Neverov *et al.*, 2001) and acetyl imidazole, (Neverov *et al.*, 2000 & Neverov *et al.*, 2001) these plots exhibit two domains, a nonlinear one at low concentration of La^{3+} suggestive of a second order behavior in La^{3+} , followed by a linear domain at higher concentration of La^{3+} . Following the approach we have used before, (Neverov *et al.*, 2001, Neverov *et al.*, 2000 & Neverov *et al.*, 2001) we use the linear portion of these plots to calculate the observed

second order rate constants (k_2^{obs}) for La^{3+} -catalyzed methanolysis of paraoxon at the various ^spH values. These are tabulated in Table 13 and graphically presented in Figure 3 as a $\log k_2^{\text{obs}}$ vs. ^spH plot which is seen to have a skewed bell-shape, maximizing at $^s\text{pH} \sim 9$.

Table 13 -- Observed second order rate constants for La^{3+} catalyzed methanolysis of paraoxon at various ^spH values, $T = 25^\circ\text{C}$.

^spH	$k_2^{\text{obs}}, \text{M}^{-1} \text{s}^{-1} \text{ }^a$
5.15	0.065 ± 0.002
5.58	0.11 ± 0.01
5.82	0.28 ± 0.02
6.69	1.07 ± 0.04
7.10	2.4 ± 0.1
7.30	5.6 ± 0.1
7.72	11.3 ± 0.5
8.23	17.5 ± 0.5
8.96	23.2 ± 0.9
10.34	11.4 ± 0.8
10.97	5.4 ± 0.4


^a k_2 determined from slope of the k_{obs} vs. $[\text{La}^{3+}]_{\text{total}}$ plots at higher $[\text{La}^{3+}]$ at each ^spH .

Example 6. La^{3+} Catalyst Species: Stoichiometries

As shown in Figure 3, the reactivity of the catalytic species increases with increasing ^spH up to ~ 9.0 . This fact seems to indicate the involvement of at least one methoxide, although the general shape of the plot suggests the catalytic involvement of more than one species. Since the second order k_2^{obs} values for the La^{3+} -catalyzed reactions in the neutral ^spH region are some 1000- to 2300-fold larger than the methoxide $k_2^{\text{OCH}_3}$, the role of the metal ion is not to simply decrease the $^s\text{pK}_a$ of any bound CH_3OH molecules that act as

nucleophiles. This points to a dual role for the metal, such as acting as a Lewis acid and as a source of the nucleophile.

Detailed mechanistic evaluation of kinetic data requires additional information such as the stoichiometries and concentrations of various La^{3+} -containing species that are formed as a function of both ^spH and concentration of La^{3+} . A study of the potentiometric titration of La^{3+} was performed under various conditions, with the concentration of $\text{La}(\text{O}_3\text{SCF}_3)_3$ from $1 \times 10^{-3} \text{ M}$ to $3 \times 10^{-3} \text{ M}$, which is within the concentration range where the kinetic plots of k_{obs} vs. concentration of La^{3+} in this study are linear. The potentiometric titration data were successfully analyzed with the computer program Hyperquad™ (Gans *et al.*, 1996) through fits to the dimer model presented in equation(1) where n assumes values of 1-5, to give the various stability constants (sK_n) that are defined in equation(2). On the basis of the five computed stability constants, $\log ^sK_{1-5} = 11.66 \pm 0.04, 20.86 \pm 0.07, 27.52 \pm 0.09, 34.56 \pm 0.20$ and 39.32 ± 0.26 , we constructed the speciation diagram shown in Figure 4 which presents the distribution of the various $\text{La}_2(\text{OCH}_3)_n$ forms as a function of ^spH at $[\text{La}(\text{O}_3\text{SCF}_3)_3]_{\text{total}} = 2 \times 10^{-3} \text{ M}$.



$$^sK_n = [\text{La}^{3+}_2(\text{OCH}_3)_n] / [\text{La}^{3+}]^2 [\text{OCH}_3^-]^n \quad (2)$$

Also included on Figure 4 as data points (•) are the k_2^{obs} data for La^{3+} -catalyzed methanolysis of paraoxon which predominantly coincide with the ^spH distribution of $\text{La}^{3+}_2(\text{OCH}_3)_2$ but with an indication that higher order species such as $\text{La}^{3+}_2(\text{OCH}_3)_3$ and/or $\text{La}^{3+}_2(\text{OCH}_3)_4$ have some activity. To determine the activities for the various $\text{La}^{3+}_2(\text{OCH}_3)_n$ we analyzed the k_2^{obs} data as a linear combination of individual rate constants (equation(3)).

$$k_2^{\text{obs}} = (k_2^{2:1}[\text{La}^{3+}_2(\text{OCH}_3)_1] + k_2^{2:2}[\text{La}^{3+}_2(\text{OCH}_3)_2] + \dots \\ \dots k_2^{2:n}[\text{La}^{3+}_2(\text{OCH}_3)_n]) / [\text{La}^{3+}(\text{O}_3\text{SCF}_3)_3]_{\text{total}} \quad (3)$$

where $k_2^{2:1}, k_2^{2:2}, \dots, k_2^{2:n}$ are the second order rate constants for the methanolysis of paraoxon promoted by the various dimeric forms. Given in Table 14 are the best-fit rate constants produced by fitting under various assumptions.

Table 14 -- Computed second order rate constants for various dimeric forms $\text{La}_2(\text{OCH}_3)_n$, catalyzing the methanolysis of paraoxon, as determined from fits of k_2^{obs} data in Table 13 to equation(3), $[\text{La}(\text{O}_3\text{SCF}_3)_3]_{\text{total}} = 2 \times 10^{-3} \text{ M}$, $T=25^\circ \text{C}$.

Fit #	$K_2^{2:1} (\text{M}^{-1}\text{s}^{-1})$	$k_2^{2:2} (\text{M}^{-1}\text{s}^{-1})$	$k_2^{2:3} (\text{M}^{-1}\text{s}^{-1})$	$k_2^{2:4} (\text{M}^{-1}\text{s}^{-1})$	R^2
1 ^a	15.9 ± 3.2	49.8 ± 2.2	67.2 ± 36.0	8.8 ± 11.2	0.9976
2 ^b	18.4 ± 5.4	47.2 ± 2.4	110.4 ± 11.8	-	0.9861
3 ^c	-	51.4 ± 2.8	103.4 ± 17	-	0.9664

^a Including all dimeric forms except $\text{La}_2(\text{OCH}_3)_0$ and $\text{La}_2(\text{OCH}_3)_6$. Computed value of $k_2^{2:5} = (-3.4 \pm 10.8) \text{ M}^{-1}\text{s}^{-1}$.

^b Computed without the involvement of $k_2^{2:4}$ and $k_2^{2:5}$.

^c Computed without the involvement of $k_2^{2:1}$, $k_2^{2:4}$ and $k_2^{2:5}$.

We have analyzed the titration data to determine speciation for a total La^{3+} concentration of $2 \times 10^{-3} \text{ M}$ which is in the general concentration range where the kinetic behavior of the methanolysis of paraoxon is linearly dependent on concentration of La^{3+} , and thus largely controlled by dimeric species. In Figure 5 are presented kinetic plots for all three species ($\text{La}^{3+}_2(\text{OCH}_3)_1$, $\text{La}^{3+}_2(\text{OCH}_3)_2$ and $\text{La}^{3+}_2(\text{OCH}_3)_3$) based on their second order rate constants for catalyzed methanolysis of paraoxon, and their concentrations as a function of ^spH . Their combined reactivities as a function of ^spH give the predicted $\log k_2^{\text{obs}}$ vs. ^spH profile shown as the dashed line on Figure 5. The computed line is also presented in the plot in Figure 2 of $\log k_2^{\text{obs}}$ vs. ^spH . Included on Figure 5 as data points (■) are the actual experimentally-determined values which fit on the computed profile with remarkable fidelity, strongly indicating that these three species are responsible for the observed activity. At ^spH values below 9, the $\text{La}^{3+}_2(\text{OCH}_3)_2$ complex accounts for essentially all the activity, while at ^spH 10 and above, the dominantly active form is $\text{La}^{3+}_2(\text{OCH}_3)_4$.

Through joint consideration of the k_{obs} vs. concentration of La^{3+} kinetics and a detailed analysis of the potentiometric titration data for La^{3+} in methanol, we have determined that the dominant species in solution are dimers of the general formula $\text{La}_2(\text{OCH}_3)_n$ where $n = 1-5$, and three of these dimers, $\text{La}^{3+}_2(\text{OCH}_3)_1$, $\text{La}^{3+}_2(\text{OCH}_3)_2$ and $\text{La}^{3+}_2(\text{OCH}_3)_3$, account for all the catalytic activity with $\text{La}^{3+}_2(\text{OCH}_3)_2$ being the most important form at $^s\text{pH} < 9$.

The ^spH dependence of the metal ion is such that several complexes are present with their individual concentrations maximized at different ^spH values. It is only through complementary analyses of the kinetic and potentiometric titration data that one can satisfactorily explain the kinetic behavior of complex mixtures having several ^spH dependent forms.

Through a series of detailed potentiometric titrations of the $\{\text{La}^{3+}:\text{OMe}\}$ system in methanol, and through studies of the kinetics of methanolysis of paraoxon as a function of La^{3+} concentration and ^spH , it has been determined that in this $\{\text{La}^{3+}:\text{OMe}:\text{paraoxon}\}$ system there are two dominant stoichiometries of catalysts, $\text{La}_2(\text{OCH}_3)_2$ with a proposed structure of a bis-methoxy bridged dimer between ^spH 8 and 10 (maximum concentration of ~80% at ^spH 8.9), and $\text{La}_2(\text{OCH}_3)_3$ with a proposed structure of tris-methoxy bridged dimer between ^spH 9 and 11 (maximum concentration of ~25% at ^spH 10). Above a total $[\text{La}^{3+}]$ of about 2×10^{-4} M, these species form spontaneously in solution without any requirement for added ligands, so that in the millimolar concentration range, dimer formation is essentially complete.

Given that we know the dominantly active forms are $\text{La}^{3+}_2(\text{OCH}_3)_2$ and $\text{La}^{3+}_2(\text{OCH}_3)_3$, we can derive a kinetic expression (equation 4) which gives values of $k_2^{2:2} = 51.4 \pm 2.8$ and $k_2^{2:3} = 103 \pm 17 \text{ M}^{-1}\text{s}^{-1}$ for the second order rate constants for methanolysis of paraoxon catalyzed by the bis-methoxy dimer and the tris-methoxy dimer respectively (Table 14).

$$k_2^{\text{obs}} = k_2^{2:2}[\text{La}^{3+}_2(\text{OCH}_3)_2] + k_2^{2:3}[\text{La}^{3+}_2(\text{OCH}_3)_3] \quad (4)$$

The net effect of this is that a solution containing 2×10^{-3} M of $\text{La}(\text{OTf})_3$, generating 1×10^{-3} M of total dimer, will catalyze the methanolysis of paraoxon with $t_{1/2}$ values of 30s, 20s, 15s and 30s at respective ^spH values of 7.7, 8.2, 9.0 and 10.3. By way of reference, at ^spH 7.7 the methoxide background rate constant is $(0.011 \text{ M}^{-1}\text{s}^{-1} \times 10^{-9} \text{ M } [\text{OCH}_3^-]) = 1.1 \times 10^{-11} \text{ s}^{-1}$, corresponding to a $t_{1/2}$ of 1994 years, so that the acceleration afforded by the La^{3+} catalyst is some two billion-fold at that ^spH .

Example 7. La^{3+} Catalysis: Proposed Mechanism


We have shown above that La^{3+} in methanol is a remarkably effective catalyst for the decomposition of paraoxon and that there are three forms of dimeric species which have maximal activities at different ^spH values. Of these, the highest activity is attributed to

$\text{La}^{3+}_2(\text{OCH}_3)_2$ operating most effectively in the neutral pH region between 7.7 and 9.2 (neutral pH in methanol is 8.4). Given in Figure 1A is a proposed mechanism by which $\text{La}^{3+}_2(\text{OCH}_3)_2$, as a bis methoxy bridged dimer, promotes the methanolysis of paraoxon. Although none of our k_{obs} vs. $[\text{La}^{3+}]$ kinetics profiles shows saturation behavior indicative of formation of a strong complex between paraoxon and La^{3+} , given the well-known coordinating ability of trialkyl phosphates to lanthanide series metal ions and actinide series metal ions, a first step probably involves transient formation of a $\{\text{paraoxon}:\text{La}^{3+}_2(\text{OCH}_3)_2\}$ complex. Since it is unlikely that the bridged methoxy is sufficiently nucleophilic to attack the coordinated phosphate, in the proposed mechanism, one of the $\text{La}^{3+}\text{-OCH}_3\text{-La}^{3+}$ bridges opens to reveal a singly coordinated $\{\text{La}^{3+}:\text{OCH}_3\}$ adjacent to a Lewis acid coordinated phosphate which then undergoes intramolecular nucleophilic addition followed by ejection of the *p*-nitrophenoxy leaving group. $\text{La}^{3+}_2(\text{OCH}_3)_2$ is regenerated from the final product by a simple deprotonation of one of the methanols of solvation and dissociation of the phosphate product, $(\text{EtO})_2\text{P}(\text{O})\text{OCH}_3$.

Example 8. M^{n+} -Catalyzed Methanolysis of O,O'-diethyl-S-*p*-nitrophenylphosphorothioate: Experimental Details, Kinetics and NMR Studies

O,O'- diethyl-S-*p*-nitrophenyl phosphorothiolate, when placed in an appropriately buffered methanol solution containing La^{3+} and OCH_3 ions held in the pH region between 7 and 11, underwent rapid methanolysis at ambient temperature to produce diethyl methyl phosphate and 5-mercapto-2-nitrobenzene. A detailed reaction scheme is given in Scheme 2 and reaction conditions are detailed below.

Scheme 2.



To 4.9 mL of anhydrous methanol at ambient temperature was added N-ethylmorpholine (63.8 μL or 57.7 mg) half neutralized with 11.4 M HClO_4 (21.5 μL), so that the final total buffer concentration was 0.1M in 4.95mL solution. The measured ^spH of the buffer solution was 8.89. To 0.8 mL of this buffer and 0.2 mL deuterated methanol was added 8.8 mg of O,O-diethyl-S-p-nitrophenyl phosphorothiolate. The ^{31}P NMR spectrum of this solution showed a single signal at δ 22.39 ppm. Following NMR analysis, a 10 μL aliquot of a lanthanum ion/sodium methoxide/methanol solution was added which had been prepared by dissolving 16.4 mg $\text{La}(\text{O}_3\text{SCF}_3)_3$ in 56.9 μL of 0.5 M sodium methoxide methanol solution. At this point, the concentrations in the NMR tube were: 0.030 M phosphorothiolate, 0.1 M N-ethylmorpholine, 0.01M sodium methoxide and 0.0098 M $\text{La}(\text{O}_3\text{SCF}_3)_3$. The ^{31}P NMR spectrum, obtained 103 sec after addition of the aliquot indicated complete disappearance of the phosphorothiolate signal and the appearance of a new signal at δ 3.57 ppm, attributable to diethyl methyl phosphate in the presence of 0.0098 M La^{3+} .

The absorbance of a 0.5 mL solution of methanol containing 1 mM of $\text{Cu}(\text{OTf})_2$, 1 mM of $[\text{12}] \text{aneN}_3$, 0.5 mM of NaOCH_3 and 0.5 mM of O,O'-diethyl-S-p-nitrophenylphosphorothioate was monitored at 280 nm as a function of time. The reaction exhibited first order kinetics with $k_{\text{obs}} = 4.3 \times 10^{-2} \text{ s}^{-1}$ ($t_{1/2} = 16 \text{ sec}$) corresponding to a 8.3×10^7 -fold acceleration over the background reaction at $^s\text{pH} = 8.41$.

The absorbance of a 2.5 mL solution of methanol containing 1 mM of $\text{Zn}(\text{OTf})_2$, 1 mM of $[\text{12}] \text{aneN}_3$, 0.5 mM of NaOCH_3 and 0.5 mM of O,O'-diethyl-S-p-nitrophenylphosphorothioate was monitored at 280 nm as a function of time. The reaction exhibited first order kinetics with $k_{\text{obs}} = 4.1 \times 10^{-4} \text{ s}^{-1}$ ($t_{1/2} = 28 \text{ min}$) corresponding to a 4.1×10^5 -fold acceleration over the background reaction at $^s\text{pH} = 8.70$.

Example 9. La^{3+} -Catalyzed Methanolysis of VX: A Prophetic Example

To 200 mL of methanol is added 2.55 mL of N-ethylmorpholine (2.3 g) and 0.86 mL of 11.4 M HClO_4 to bring the total buffer concentration to 0.1 M. To this solution is added 1.29 g of $\text{La}(\text{O}_3\text{SCF}_3)_3$ and 4 mL of a 0.5 M solution of NaOCH_3 in methanol.


To the above solution is added 2 g of VX (8.33×10^{-3} moles, 0.041 M) and the solution is allowed to stand at ambient temperature for 15 minutes. It is expected that analysis of the resulting solution would indicate substantially complete disappearance of VX.

Example 10. M^{2+} -Catalyzed Methanolysis of Fenitrothion

The activity of this system may be increased by adding equimolar amounts of bi- or tri-dentate ligands to complex $Zn^{2+}(\text{OCH}_3)$ and limit oligomerization of $Zn^{2+}(\text{OCH}_3)_2$ in solution. The systems studied herein used methoxide and the ligands phen, diMephen and [12]aneN₃. The active forms of the metal ions at neutral $s_p\text{H}$ are $Zn^{2+}(\text{OCH}_3)$ with no added ligand and $\{Zn^{2+}:L:(\text{OCH}_3)\}$ when ligand (L) is present. In the case of phen ligand, decreasing the oligomerization does not prevent the formation $Zn^{2+}(\text{OCH}_3)$ dimers since the bulk of the material is now present as $\{LZn^{2+}(\text{OCH}_3)_2Zn^{2+}L\}$ which is not catalytically active, but is in equilibrium with an active mononuclear form. The propensity to form the latter inactive dimers can be reduced either by increasing the steric interaction (ligand diMephen) or by changing the coordination number (ligand [12]aneN₃) in which cases the overall activity of the catalytic system increases. In the case of ligand diMephen, the dimerization is definitely reduced but the binding to the metal ion is not as strong as in the case of phen or [12]aneN₃, which means that there is some free Zn^{2+} in solution under the concentrations and $s_p\text{H}$ region where the catalyst is active.

A reaction scheme is given below (Scheme 3) for the methanolysis of fenitrothion where M^{2+} is a transition metal ion, most preferably Zn^{2+} or Cu^{2+} . In a preferred embodiment a ligand is present, preferably a bidentate or tridentate ligand, most preferably [12]aneN₃ for Cu^{2+} and diMephen or [12]aneN₃ for Zn^{2+} .

Scheme 3



As seen in Figures 6 and 7, $Cu^{2+}:(\text{OCH}_3)$ at 25°C either alone or in the presence of

equimolar [12]aneN₃, bpy or phen shows both great catalytic efficacy and specificity toward the P=S derivatives.

Apparently matching the hard/soft characteristics of the metal ion and the substrate is important in designing an effective catalytic system for P=S substrates. With due consideration for matching the hard/soft characteristics of the substrate and the metal ion, dramatic rate and selectivity can be achieved in the methanolysis of P=O vs. P=S phosphates.

Example 11. Zn²⁺-Catalyzed Methanolysis of Paraoxon and Fenitrothion


The methanolyses of paraoxon and fenitrothion were investigated as a function of added Zn(OTf)₂ or Zn(ClO₄)₂ in methanol at 25 °C either alone, or in the presence of equimolar concentration of ligands: phen, diMephen and [12]aneN₃. The catalysis requires the presence of methoxide, and when studied as a function of added [NaOCH₃], the rate constants (*k*_{obs}) for methanolysis with Zn²⁺ alone or in the presence of equimolar phen or diMephen, maximize at different [OCH₃]/[Zn²⁺]_{total} ratios of 0.3, 0.5 and 1.0 respectively. Plots of *k*_{obs} vs. [Zn²⁺]_t either alone or in the presence of equimolar ligands phen and diMephen at the [OCH₃]/[Zn²⁺]_{total} ratios corresponding to the rate maxima are curved and show a square root dependence on [Zn²⁺]_t. In the cases of phen and diMephen, this is explained as resulting from formation of a non-active dimer, formulated as a bis-μ-methoxide bridged form (L:Zn²⁺(⁻OCH₃)₂Zn²⁺:L) in equilibrium with an active mononuclear form, L:Zn²⁺(⁻OCH₃). In the case of the Zn²⁺: [12]aneN₃ system, no dimeric forms are present as can be judged by the strict linearity of the plots of *k*_{obs} vs. [Zn²⁺]_t in the presence of equimolar [12]aneN₃ and ⁻OCH₃. Analysis of the potentiometric titration curves for Zn²⁺ alone and in the presence of the ligands allows calculation of the speciation of the various Zn²⁺ forms and shows that the binding to ligands phen and [12]aneN₃ is very strong, while the binding to ligand diMephen is weaker. This {Zn²⁺: [12]aneN₃: ⁻OMe} system exhibits excellent turnover of the methanolysis of paraoxon when the substrate is in excess. A mechanism for the catalyzed reactions is proposed (see Figure 1B) which involves a dual role for the metal ion as a Lewis acid and source of nucleophilic Zn²⁺-bound ⁻OCH₃.

Example 12. Zn²⁺-Catalyzed Methanolysis of Paraoxon and Fenitrothion and p-nitrophenyl acetate

A second set of methanolysis experiments was performed with three substrates, namely paraoxon, fenitrothion and p-nitrophenyl acetate, as a function of total added

$[\text{Zn}(\text{ClO}_4)_2]$ maintaining the $[(\text{OCH}_3)]/[\text{Zn}^{2+}]_{\text{total}}$ ratio at 0.3 with added NaOCH_3 . The three plots shown in Figure 12 all exhibit a similar curvature independent of the nature of the substrate. The curvature thus cannot be due to substrate binding and is modeled according to the overall process given in equation(5) where an active mononuclear form (assumed to be $[\text{Zn}(\text{OCH}_3)]^+$ is in equilibrium with a non-active dimer. Given in equation (6) is the appropriate kinetic expression based on equation(5) which includes a possible methoxide dependent term ($k_{\text{background}}$) which is present for the most reactive substrate (p-nitrophenyl acetate) but not important for the phosphate triesters. This expression shows a square-root dependence on the $[\text{M}^{2+}]_{\text{total}}$. Shown in Figure 9A and 9B are the concentration dependencies for the methanolysis of fenitrothion (Figure 9A) and paraoxon (Figure 9B) catalyzed by Zn^{2+} alone and in the presence of ligands phen and diMephen where the ratio of $[(\text{OCH}_3)]/[\text{Zn}^{2+}]_{\text{total}}$ is kept at a constant value (i.e. 0.3 for Zn^{2+} alone, 0.5 for phen, and 1.0 for diMephen).

These plots are also curved, not due to a saturation binding of the phosphorus triesters to the metal, but due to the monomer:dimer equilibrium given in equation(5). The lines through the Figure 9A, 9B data are derived on the basis of NLLSQ fits to equation(6) and yield the kinetic constants given in Table 16. As shown in Figure 13A, the kinetic dependence in the presence of ligand [12]aneN₃ is substantially linear and shows no evidence of monomer:dimer equilibrium.



$$k_{\text{obs}} = \left\{ k_m K_{\text{dis}} \left(\sqrt{1 + \frac{8[\text{M}^{2+}]_{\text{total}}}{K_{\text{dis}}}} - 1 \right) / 4 + k_{\text{background}} \right\} \quad (6)$$

Example 13. Zn^{2+} -Catalyst Stoichiometry

Potentiometric titration of $\text{Zn}(\text{OTf})_2$ solutions of varying concentrations (0.5-2 mM) in anhydrous methanol were performed in the absence and presence of equimolar amounts of ligands phen, diMephen and [12]aneN₃ in order to determine the speciation of the Zn^{2+} ions under conditions similar to those of the kinetic experiments.

Independent titrations of 1 mM solutions of each ligand were performed and the

resulting data were analyzed using Hyperquad™ 2000 fitting routine providing the s_pK_a values for the last acid dissociation step, of 5.63 ± 0.01 for phen- H^+ , 6.43 ± 0.01 for diMephen- H^+ and > 13 for [12]aneN₃- H^+ respectively.

The potentiometric titration curve of Zn(OTf)₂ presented in Figure 14 shows the consumption of two equivalents of methoxide occurring in one rather steep step. In the presence of ligands phen, diMephen and [12]aneN₃, the titration curve changes due to the formation of complexes. To analyze these titration data, a number of different dissociation schemes were attempted and the final adopted ones were selected based on goodness of fit to the titration profiles along with due consideration of the various species suggested by the kinetic studies.

The case of the ligand triazacrown ether [12]aneN₃ is the simplest to analyze since we have no evidence supporting the presence of any species containing more than one Zn²⁺ ion. This fact, coupled with the high s_pK_2 of [12]aneN₃- H^+ , allows one to define the relevant species in solution as [12]aneN₃- H^+ , Zn²⁺: [12]aneN₃, Zn²⁺: [12]aneN₃: (OCH₃) and Zn²⁺: [12]aneN₃: (OCH₃)₂, which, when fit via the Hyperquad™ 2000 program, produces a theoretical titration curve (Figure 14) which is in excellent agreement with the observed curve. The best fit formation constants for [12]aneN₃- H^+ , Zn²⁺: [12]aneN₃, Zn²⁺: [12]aneN₃: (OCH₃) and Zn²⁺: [12]aneN₃: 2(OCH₃) are given in Table 15. The Zn²⁺ speciation diagram constructed from these constants (not shown) indicates that in the s_pH region used in our kinetic studies, greater than 95% of the total Zn²⁺ is present as Zn²⁺: [12]aneN₃: (OCH₃). Shown in Figure 13A is a plot of the pseudo-first order rate constants for the methanolysis of paraoxon in the presence of Zn(OTf)₂ with a right hand axis depicting the [Zn²⁺: [12]aneN₃: (OCH₃)] as function of total [Zn(OTf)₂]. The very good correlations between the kinetic data and the speciation data strongly supports Zn²⁺: [12]aneN₃: (OCH₃) as the catalytically active component, with a derived second order rate constant of 50.4 M⁻¹ min⁻¹ for the methanolysis of paraoxon.

Potentiometric titration of an equimolar mixture of Zn(OTf)₂ and phen in the presence of 0.6 equivalents of perchloric acid showed that all the added H⁺ was released in the strong acid region below s_pH 3 with one additional step consuming a single equivalent of methoxide around s_pH 10. The former indicates strong binding between Zn²⁺ and phen even at s_pH = 3, but does not allow us to determine an exact value of the Zn²⁺: phen binding constant other than to set a lower limit for its formation constant of 10¹⁰ M⁻¹ which was used as a fixed value in all subsequent fittings. In the higher s_pH region where the kinetic

experiments were performed, we employed a model where the Zn^{2+} exists predominantly as $\{\text{Zn}^{2+}:\text{phen}:(\text{OCH}_3)_2\}$ and $\text{Zn}^{2+}:\text{phen}:(\text{OCH}_3)_2$, both of these being inferred by the kinetic data. Hyperquad™ 2000 fitting of the full titration profile using the previously determined stability constants for phen-H^+ and $\text{Zn}^{2+}:\text{phen}$, produces a good fit and provides respective stability constants for $\{\text{Zn}^{2+}:\text{phen}:(\text{OCH}_3)_2\}$ and $\text{Zn}^{2+}:\text{phen}:(\text{OCH}_3)_2$ given in Table 15.

In the catalysis of methanolysis of paraoxon and fenitrothion by $\{\text{Zn}^{2+}:\text{OMe}\}$, either alone or in the presence of complexing ligands, two things are clear: first, Zn^{2+} species are appreciably soluble in solution at all pH values and all concentrations employed; and second, equilibria consisting of dimeric species in equilibrium with a kinetically active mononuclear species are formed in the case of Zn^{2+} , $\{\text{Zn}^{2+}:\text{phen}\}$ and $\{\text{Zn}^{2+}:\text{diMephen}\}$, but not in the case of $\{\text{Zn}^{2+}:[12]\text{aneN}_3\}$ where only the kinetically active mononuclear form is present. High solubility of Zn^{2+} has been found with triflate and perchlorate counterions. These anions are preferred for their relative kinetic inertness since they give the highest rates for catalyzed reactions relative to other anions such as bromide, chloride or acetate. Methanolysis of paraoxon, catalyzed by 1 mM $\text{Zn}(\text{OTf})_2$ with 0.3 equivalent of added NaOCH_3 is relatively unaffected by the addition of up to 5 mM NaOTf or NaClO_4 , but is significantly inhibited by the addition of 1 mM NaCl , NaBr or $\text{Na}(\text{O}_2\text{CCH}_3)$.

The ability of the Zn^{2+} species to methanolyze both the P=O and P=S species with second-order rate constants 50-to 1000-fold larger than the corresponding second-order rate constants for methoxide attack alone may be due to the bifunctional nature of the catalyst and partly due to the reduced dielectric constant of the medium and its reduced solvation of metal ions relative to water.

Preparatively useful forms of catalysts can be generated by the addition of known amounts of ligand, $\text{Zn}(\text{OTf})_2$ and methoxide. In the case of a solution comprising 2 mM $\text{Zn}(\text{OTf})_2$, 2 mM diMephen ligand and 2 mM NaOCH_3 which generates a pH of ~ 9.5 , methanolysis of paraoxon is accelerated 1.8×10^6 -fold and methanolysis of fenitrothion is accelerated 13×10^6 -fold. Likewise, a solution comprising 1 mM of $\text{Zn}(\text{OTf})_2$, 1 mM $[12]\text{aneN}_3$ ligand and 0.5 mM NaOCH_3 generates a pH of 9.3 and methanolysis of paraoxon is accelerated 1.7×10^6 -fold.

Unlike the dimeric form of La^{3+} , which are effective for methanolyzing paraoxon, dimeric forms of Zn^{2+} are not as effective as its monomers.

Example 14. Zn^{2+} -Catalyzed Methanolysis of Paraoxon and Fenitrothion: Kinetic and Potentiometric Studies

The kinetics for Zn^{2+} -catalyzed methanolysis of paraoxon and fenitrothion fall into two distinct classes depending on what ligand is coordinated to the metal ion and how much methoxide is added. Without any ligand, as shown in Figure 11, the k_{obs} for methanolysis of paraoxon in the presence of 1mM $\text{Zn}(\text{OTf})_2$ is maximized between 0.1 and 0.4 mM added NaOCH_3 . There is an initially very strong dependence on the concentration of methoxide, the slope of which for the first 0.05 equationadded yields a second order rate constant of $\sim 34 \text{ M}^{-1} \text{ min}^{-1}$ for methanolysis of paraoxon. Undoubtedly this methoxide is coordinated to Zn^{2+} to establish the $\{\text{Zn}(\text{OCH}_3)\}_2^{2+} \rightleftharpoons 2 \{\text{Zn}(\text{OCH}_3)\}^+$ equilibrium but as additional methoxide is added, the overall rate drops significantly suggesting formation of inactive species having a $[\text{OCH}_3]/[\text{Zn}^{2+}]$ greater than 1. This agrees with a potentiometric titration of Zn^{2+} in methanol which displayed a steeper-than-normal consumption of 2 methoxides in an apparent single event having a midpoint of $\sim {}^s\text{pK}_a$ 9.8 which, when analyzed via HyperquadTM fitting to a model containing only the mononuclear species $\text{Zn}^{2+}(\text{OCH}_3^-)$ and $\text{Zn}^{2+}(\text{OCH}_3^-)_2$, gives apparent ${}^s\text{pK}_{a1}$ and ${}^s\text{pK}_{a2}$ values of 10.66 and 8.94. While our original fitting (Gibson, *et al.*, 2003) did not include dimer and oligomer formation, the fact that the second apparent ${}^s\text{pK}_a$ is lower than the first indicates some cooperative effect facilitating addition of a second methoxide per Zn^{2+} ion before the first addition is stoichiometrically complete. This fact limits the amount of any forms having a methoxide/ Zn^{2+} stoichiometry of 1 and shifts the maximum of the kinetic plot in Figure 11 to the left. Species where the methoxide/ Zn^{2+} ratio >1 probably exist in solution as oligomers of $\{\text{Zn}^{2+}(\text{OCH}_3)_{1.5,2}\}_n$ held together with methoxide bridges. Added bi- or tridentate ligands could, in principle, disrupt this arrangement by capping one face of the Zn favouring the formation of dimers and monomers of stoichiometry $\{\text{Zn}^{2+}:\text{L}(\text{OCH}_3)\}_2$, $\text{Zn}^{2+}:\text{L}(\text{OCH}_3)(\text{HOCH}_3)$ or $\text{Zn}^{2+}:\text{L}(\text{OCH}_3)_2$ depending on the methoxide/ Zn^{2+} ratio. Indeed, as shown in Figure 8, ligands phen, diMephen and [12]aneN₃ modify the kinetic behaviour in important ways depending on whether the methoxide/ Zn^{2+} ratio is less than or greater than 1.

Example 15. Zn^{2+} -Catalyzed Methanolysis of Paraoxon: NMR Studies of Catalytic Turnover

A ${}^{31}\text{P}$ NMR experiment was performed to determine a turnover rate for the methanolysis of paraoxon using $\text{Zn}^{2+}:\text{diMephen}:\text{OCH}_3$.

To 0.6 mL of dry methanol (with 20% of CD₃OD as an NMR lock signal) containing 1 mM each of Zn(OTf)₂, diMephen and NaOCH₃ at ambient temperature was added 2.54 mg of paraoxon. At this point the concentration of paraoxon was 15 mM and that of Zn²⁺: diMephen: ⁻OCH₃ was taken as 1.0 mM with the measured ^spH of the methanol solution being 8.75, close to neutrality (8.34). The ³¹P NMR spectrum of the solution was monitored periodically over ~160 minutes at which time it indicated complete disappearance of the paraoxon signal which had been at δ -6.35 ppm and complete appearance of a new signal at δ 0.733 ppm corresponding to the product diethyl methyl phosphate. The ¹H NMR spectrum was obtained after 150 min and it confirmed the complete disappearance of the starting material and full release of the product *p*-nitrophenol.

The ³¹P NMR spectrum of a solution containing 15 mM paraoxon and 1 mM in each of Zn(OTf)₂, NaOCH₃ and ligand diMephen was continuously monitored at ambient temperature over a period of ~160 minutes. The spectra were summed each 15 minutes to produce the time profile given in Figure 10 which displays the disappearance of paraoxon and the appearance of a new signal at δ 0.733 ppm attributed to diethyl methyl phosphate. Fitting of these two time profiles to a first order expression gave an average pseudo-first order rate constant of $(4.5 \pm 0.1) \times 10^{-4} \text{ s}^{-1}$ over 15 turnovers ($t_{1/2} = 25 \text{ min}$), thus showing the true catalytic nature of the system.

Example 16. Zn²⁺-Catalyzed Methanolysis of Paraaxon and Fenitrothion: Kinetics

As shown by the various formation constants given in Table 15, phen binds very tightly to Zn²⁺ at all values in methanol. According to potentiometric titration data, the major species in the ^spH domain surrounding $0 < [\text{methoxide}]/[\text{Zn}^{2+}]_t < 1$ is the dimer {Zn²⁺:phen:(⁻OCH₃)₂} which is in equilibrium with a small amount of kinetically active monomer, {Zn²⁺phen(⁻OCH₃)}. Under conditions where the $[\text{methoxide}]/[\text{Zn}^{2+}]_t = 0.5$, a plot of k_{obs} for catalyzed methanolysis of paraoxon vs. $[\text{Zn}^{2+}]_{\text{total}}$ (see Figure 14B) follows the square root dependence of equation (6) that corresponds to the process presented in equation (5) with the derived kinetic parameters being given in Table 16. The same general phenomenon is seen with ligand diMephen although its binding to Zn²⁺ is weaker than phen (as is known to be the case in water) such that at any given ^spH, only about 85% of the Zn²⁺ is bound to diMephen.

Table 15 -- Formation constants for various species determined by potentiometric titration.

Equilibrium	$\text{Log } {}^s K$ L = phen	$\text{Log } {}^s K$ L = diMephen	$\text{Log } {}^s K$ L = [12]aneN ₃
$[\text{L-H}^+]/[\text{L}][\text{H}^+]$	5.63	6.43	14.92
$[\text{ZnL}]/[\text{L}][\text{Zn}]$	10	4.25	10.11
$[\text{Zn}_2\text{L}_2(\text{OMe})_2]/[\text{L}]^2[\text{Zn}]^2[\text{OMe}]^2$	36.33	28.05	
$[\text{ZnL}(\text{OMe})_2]/[\text{L}][\text{Zn}^{2+}][\text{OMe}]^2$	20.58		21.67
$[\text{ZnL}(\text{OMe})]/[\text{L}][\text{Zn}][\text{OMe}]$			17.79

Table 16 -- Kinetic constants for the methanolysis of fenitrothion and paraoxon catalyzed by Zn^{2+} in the absence and presence of ligands phen, diMephen, [12]aneN₃, at T = 25 °C.

		Paraoxon	Fenitrothion
Catalyst	$K_{\text{dis}} (\text{mM})^a$	$k_m (\text{M}^{-1} \text{min}^{-1})^a$	$k_m (\text{M}^{-1} \text{min}^{-1})^a$
OCH_3^-	-	0.66	0.043 ± 0.001
Zn^{2+} ^b	< 0.005	72.5 ± 1.5	11.2 ± 0.4
$\{\text{Zn}^{2+}:\text{phen}\}^c$	< 0.005	124 ± 2.5	19.0 ± 0.6
$\{\text{Zn}^{2+}:\text{phen}:2(\text{OCH}_3)^-\}^d$	-	29.5 ± 0.7	2.7 ± 0.1
$\text{Zn}^{2+}:\text{diMephen}^e$	0.6 ± 0.2	101 ± 1	48.0 ± 0.7
$\text{Zn}^{2+}:[12]\text{aneN}_3:(\text{OCH}_3)^-f$	-	50.8 ± 0.8	2.9 ± 0.1
$\{2\text{La}^{3+}:2(\text{OCH}_3)^-\}^g$	-	2830 ± 140	No catalysis

^a Dimer dissociation constant (K_{dis}) and conditional second order rate constant (k_m) for monomer defined as in equation(5); "-" means non-applicable since there is no observable dimerization under the specific conditions.

^b Based on NLLSQ fits of k_{obs} vs. $[\text{Zn}^{2+}]_{\text{total}}$ data to equation(6) at $[\text{methoxide}]/[\text{Zn}^{2+}]_{\text{total}}$ ratio of 0.3

^c Based on NLLSQ fits of K_{obs} vs. $[\text{Zn}^{2+}:\text{phen}]_{\text{total}}$ data to equation(6) at $[\text{methoxide}]/[\text{Zn}^{2+}]_{\text{t}}$ ratio of 0.5

^d Based on linear fits of K_{obs} vs. $[\text{Zn}^{2+}:\text{phen}]_{\text{total}}$ data to equation(6) at $[\text{methoxide}]/[\text{Zn}^{2+}]_{\text{t}}$ ratio of 2.0

^e Based on NLLSQ fits of K_{obs} vs. $[\text{Zn}^{2+}:\text{diMephen}]_{\text{total}}$ data to equation(6) at $[\text{methoxide}]/[\text{Zn}^{2+}]_{\text{total}}$ ratio of 1.0

^f Based on linear fits of K_{obs} vs. $[\text{Zn}^{2+}:[12]\text{aneN}_3:(\text{OCH}_3)^-]_{\text{t}}$ data at $[\text{methoxide}] = [\text{Zn}^{2+}]_{\text{total}} = [[12]\text{aneN}_3]$.

^g From reference Tsang *et al.*, 2003

As shown in Figure 8 for the methanolysis of paraoxon, the $\text{Zn}^{2+}:\text{phen}$ and $\text{Zn}^{2+}:\text{diMephen}$ systems behave differently in the $1 < [\text{methoxide}]/[\text{Zn}^{2+}]_{\text{total}} < 2$ domains with the overall activity increasing and decreasing respectively. Because of the weak binding inherent in the $\text{Zn}^{2+}:\text{diMephen}$ system, the additional methoxide probably displaces the ligand from the $\{\text{Zn}^{2+}:\text{diMephen}:(\text{OCH}_3)\}_{1,2}$ forms to generate uncomplexed diMephen and $\{\text{Zn}(\text{OCH}_3)_2\}_n$ oligomers which are not active. However, because of the far stronger binding of phen to Zn^{2+} , the additional methoxide breaks apart the $\{\text{Zn}^{2+}:\text{phen}:(\text{OCH}_3)\}_2$ dimer as shown in Figure 1B to form $\text{Zn}^{2+}:\text{phen}:(\text{OCH}_3)_2$. The presence of $\text{Zn}^{2+}:\text{phen}:(\text{OCH}_3)_2$ and its catalytic viability is respectively confirmed by the potentiometric titration data and by the fact that a plot of k_{obs} for methanolysis of both substrates vs. $[\text{Zn}^{2+}]_t$ under conditions where the $[\text{Zn}^{2+}]:\text{phen}:\text{methoxide}$ ratio is 1:1:2 gives a straight line with a slope of $k_m = 29.5 \text{ M}^{-1} \text{ min}^{-1}$ for the methanolysis of paraoxon and $k_m = 2.7 \text{ M}^{-1} \text{ s}^{-1}$ for the methanolysis of fenitrothion.

The $\text{Zn}^{2+}:[12]\text{aneN}_3:\text{OCH}_3^-$ system is a simple one because of very strong binding and the lack of formation dimers $\{\text{Zn}^{2+}:[12]\text{aneN}_3:(\text{OCH}_3)\}_2$ under employed conditions. In methanol, the $\text{M}^{2+}\text{-L}$ binding constant is large ($\log {}^s K = 10.11$), ensuring that there is essentially no free ligand in solution, and the ${}^s \text{p}K_a$ for ionization of the complex $\text{Zn}^{2+}:[12]\text{aneN}_3:\text{HOCH}_3$ is 9.1. The k_{obs} vs. $[\text{Zn}^{2+}]_{\text{total}}$ plot shown in Figure 13A is a straight line consistent with $(\text{Zn}^{2+}:[12]\text{aneN}_3:(\text{OCH}_3))$ being the active catalyst and predominant form.

Example 17. Cu^{2+} -Catalyzed Methanolysis of Paraoxon and Fenitrothion: Kinetic Studies

In the absence of metal ions, uncatalyzed attack of methoxide on paraoxon is some 15 times faster than on fenitrothion, but in the presence of all $\{\text{Cu}^{2+}:(\text{OCH}_3)\}$ species are more effective for fenitrothion than paraoxon. This can be quantified by the relative selectivity parameter given in Table 18 which compares the relative reactivity of the metal-coordinated methoxide reaction relative to free methoxide attack for $\text{P}=\text{S}$ and $\text{P}=\text{O}$ substrates. Relative Selectivity parameters clearly correlate with the hard/soft properties of the metal ion. The "hard" ion La^{3+} exhibits exclusive selectivity for the $\text{P}=\text{O}$ substrate (relative selectivity parameter ~ 0), while the softer Zn^{2+} ion shows almost equal affinity for $\text{P}=\text{O}$ and $\text{P}=\text{S}$ substrates (relative selectivity parameter ~ 1). Of the three ions, Cu^{2+} is softest, and exhibits very high selectivities for the $\text{P}=\text{S}$ substrates with relative selectivity parameter values from ~ 55 -340 with the highest values exhibited in the case of the aromatic ligands. The best combination of selectivity and overall high catalytic activity is achieved with $\{[12]\text{aneN}_3:\text{Cu}^{2+}:(\text{OCH}_3)\}$ perhaps due to reduced dimerization. All the Cu^{2+} -catalyzed

reactions proceed with computed second order rate constants larger than those for the uncatalyzed attack of methoxide on paraoxon or fenitrothion which indicates that there is a dual role for the metal ion. As in other M^{n+} -promoted hydrolytic and methanolytic reactions, the metal ion is reasonably proposed to deliver a M^{n+} -coordinated OH^- or CH_3O^- and act as a Lewis acid to polarize a $P=S$ or $P=O$ unit, which provides both rate and selectivity enhancement. There is a 17,000-fold enhancement of attack of $[12]aneN_3:Cu^{2+}:(^-OCH_3)$ on fenitrothion vs. attack of free $^-OCH_3$ even though the latter is $\sim 10^8$ -fold more basic. This represents the largest acceleration reported for metal-catalyzed phosphoryl transfer reactions to solvent. Through turnover experiments, it has been demonstrated that this is a truly catalytic system which, at millimolar concentration can provide 1.7×10^9 -fold acceleration of the methanolysis of fenitrothion at neutral s_pH and ambient temperature.

In the presence of ligand $[12]aneN_3$, the kinetic plots, k_{obs} vs. $[Cu(OTf)_2]_{total}$ (see Figure 7), for methanolysis of paraoxon and methanolysis of fenitrothion are strictly linear which is indicative of complete formation of a mononuclear catalyst of the structure: $[12]aneN_3:Cu^{2+}:(^-OCH_3)$. The second order rate constants, k_m , for paraoxon and for fenitrothion were evaluated as the gradients of the linear plots, these values being given in Table 18.

Table 18 -- Kinetic constants for the methanolysis of paraoxon and fenitrothion catalyzed by Cu^{2+} in the absence and presence of ligands $[12]aneN_3$, bpy and phen at $T = 25^\circ C$.

			Paraoxon	Fenitrothion	
Catalyst	s_pH at 0.5 eq of base	K_{dis} (mM) ^a	k_m ($M^{-1}s^{-1}$) ^a	k_m ($M^{-1}s^{-1}$) ^a	Relative selectivity ^b
$^-OCH_3$		N.A	1.1×10^{-2}	$(7.2 \pm 0.2) \times 10^{-4}$	1
$Cu^{2+}:(^-OCH_3)^c$	6.86 ± 0.2	< 0.005	0.22 ± 0.02	0.79 ± 0.03	55
$Cu^{2+}:bpy:(^-OCH_3)^d$	7.8 ± 0.2	< 0.005	< 0.2	4.48 ± 0.12	342
$Cu^{2+}:phen:(^-OCH_3)^e$	7.45 ± 0.2	< 0.005	< 0.2	2.44 ± 0.06	186
$Cu^{2+}:[12]aneN_3:(^-OCH_3)^f$	8.75 ± 0.1	-	2.76 ± 0.17	12.2 ± 0.4	67
$Zn^{2+}:[12]aneN_3:(^-OCH_3)^g$	9.3	-	0.85 ± 0.01	$(4.8 \pm 0.2) \times 10^{-2}$	0.86
$La^{3+}_2(^-OCH_3)_2^h$		-	47.2 ± 2.3	No catalysis	~ 0

^a Dimer dissociation constant (K_{dis}) and conditional second order rate constant (k_m) for reaction with monomer defined as in text. "-" means non-applicable since there is no observable dimerization under the specific conditions. The K_{dis} of < 0.005 indicates very strong dimerization and is quoted as an upper limit based on an iterative fitting procedure which provided the lowest standard deviations.

^b Defined as $(k_m/k_{OCH_3})^{fenitrothion} / (k_m/k_{OCH_3})^{paraoxon}$

- ^c Based on NLLSQ fits of k_{obs} vs. $[\text{Cu}^{2+}]_{\text{total}}$ data to equation(6) at $[\text{methoxide}]/[\text{Cu}^{2+}]_{\text{total}}$ ratio of 0.5
- ^d Based on NLLSQ fits of k_{obs} vs. $[\text{bpy}:\text{Cu}^{2+}]_{\text{total}}$ data to equation(6) at $[\text{methoxide}]/[\text{Cu}^{2+}]_{\text{total}}$ ratio of 0.5
- ^e Based on NLLSQ fits of k_{obs} vs. $[\text{phen}:\text{Cu}^{2+}]_{\text{total}}$ data to equation(6) at $[\text{methoxide}]/[\text{Cu}^{2+}]_{\text{t}}$ ratio of 0.5
- ^f Based on linear fits of k_{obs} vs. $[\text{Cu}^{2+}:[12]\text{aneN}_3:(\text{OCH}_3)]_{\text{total}}$ data at $\text{methoxide}/[\text{Cu}^{2+}]_{\text{t}}$ ratio of 0.5.
- ^g From reference Desloges, *et al.* 2004.
- ^h From reference Tsang *et al.*, 2003.

The kinetics of methanolysis were monitored at 25°C in anhydrous methanol by observing the rate of appearance of p-nitrophenol or 3-methyl-4-nitrophenol between 312 and 335 nm at $[\text{paraoxon}]$ or $[\text{fenitrothion}] = 4$ to 12×10^{-5} M under pseudo-first order conditions of excess $\text{Cu}(\text{OTf})_2$ (0.2 to 5.0×10^{-3} M). All reactions were followed to at least three half times and found to exhibit good pseudo-first order rate behavior and the first order rate constants (k_{obs}) were evaluated by fitting the Abs. vs. time traces to a standard exponential model. The kinetics were all determined under self-buffered conditions where the ^spH was controlled by a constant $\text{Cu}^{2+}/\text{Cu}^{2+}(\text{OCH}_3)$ ratio and in the cases with ligands $[12]\text{aneN}_3$, bpy and phen, these were added in amounts equivalent to the $[\text{Cu}^{2+}]_{\text{total}}$. Under these conditions the observed ^spH values correspond to the apparent $^s\text{pK}_a$ value for ionization of the $\{\text{Cu}^{2+}:\text{L}:(\text{HOCH}_3)\} \rightleftharpoons \{\text{Cu}^{2+}:\text{L}:(\text{OCH}_3)\} + {}^+\text{H}_2\text{OCH}_3$ system.

As shown in Figures 6 and 7 the overall behaviour portrayed in the k_{obs} vs. $[\text{Cu}^{2+}]$ plots falls into two categories depending on the nature of the ligand employed. In the absence of any ligand, or in the presence of equimolar bpy or phen, the Figure 6 plots are non-linear and indicative of a square-root dependence which can be fit via a standard Non-Linear Least Squares (NLLSQ) treatment to equation (6) derived on the following assumptions: all the ligand is bound to Cu^{2+} ; an active (rate constant k_m) mononuclear species $\{\text{Cu}^{2+}:\text{L}:(\text{OCH}_3)\}$ is in rapid equilibrium (dissociation constant K_{dis}) with an inactive dimer (equation4) and $k_{\text{background}}$ is negligible since it is undetectable. How good the fit of the lines is may be seen by examining the computed lines through the Figure 6 data and the best fit constants are given in Table 18. Also in Table 18 are the measured ^spH values over the entire $[\text{Cu}^{2+}]$ range under the self-buffering conditions which deviate by an acceptable 0.2 or less units. In the case of paraoxon, the catalyzed reactions were sufficiently slow that we have placed upper limits on the rate and equilibrium constants.

A system comprising 2 mM $\text{Cu}(\text{OTf})_2$, along with 0.5 equivalent of $\text{N}(\text{Bu})_4\text{OCH}_3$ and 1 equivalent of $[12]\text{aneN}_3$ catalyzes the methanolysis of fenitrothion with a $t_{1/2}$ of ~58 sec

accounting for a 1.7×10^9 -fold acceleration of the reaction relative to the background reaction at a near neutral pH of 8.75. In this system the concentration of catalyst is in excess over the concentration of fenitrothion.

A turnover experiment with substrate in excess of catalyst was conducted using 0.4 mM $Cu(OTf)_2$ along with equimolar $[12]aneN_3$ and 0.5 equivalent of NBu_4OCH_3 . The methanolysis of 2mM fenitrothion was monitored by UV/vis at $T = 25.0^\circ C$ and showed 10 turnovers relative to the active catalyst ($0.2\text{ mM } Cu^{2+}:[12]aneN_3:(^-\text{OCH}_3)$) within 100 min.

Although this invention is described in detail with reference to preferred embodiments thereof, these embodiments are offered to illustrate but not to limit the invention. It is possible to make other embodiments that employ the principles of the invention and that fall within its spirit and scope as defined by the claims appended hereto.

REFERENCES

- Bosch, E.; Rived, F.; Roses, M.; Sales, J., "Hammett-Taft and Drago Models in the Prediction of Acidity Constant Values of Neutral and Cationic Acids in Methanol" *J. Chem. Soc., Perkin Trans. 2*, **1999**, 1953.
- Bosch, E.; Bou, P.; Allemann, H.; Rosés, M. "Retention of Ionizable Compounds on HPLC. pH Scale in Methanol-Water and the pK and pH Values of Buffers" *Anal. Chem.* **1996**, 3651
- Brown, R. S.; Neverov, A. A., "Acyl and Phosphoryl Transfer to Methanol Promoted by Metal Ions" *J. Chem. Soc. Perkin 2* **2002**, 1039.
- Brown, R. S.; Zamkane, M., "Hydrolysis of Neutral Phosphate and Phosphonate Esters Catalysed by Co^{2+} -Chelates of Tris-Imidazolyl Phosphines" *Inorg. Chim. Acta.* **1985**, 108, 201.
- Desloges, W.; Neverov, A. A.; Brown, R.S., "Zinc²⁺-Catalyzed Methanolysis of Phosphate Triesters: a Process for Catalytic Degradation of the Organophosphorus Pesticides Paraoxon and Fenitrothion" *Inorg. Chem.* **2004**, submitted.
- Gans, P.; Sabatini, A.; Vacca, A., "Investigation of Equilibria in Solution. Determination of Equilibrium Constants with the HYPERQUAD Suite of Programs" *Talanta.* **1996** 43, 1739.
- Gibson, G.; Neverov, A. A.; Brown, R.S., "Potentiometric Titration of Metal Ions in Methanol" *Can. J. Chem.* **2003**, 81, 495.
- Neverov, A. A.; Brown, R. S., "Catalysis of the Methanolysis of Acetylimidazole by Lanthanum Triflate" *Can. J. Chem.* **2000**, 78, 1247.
- Neverov, A. A.; Brown, R. S., "La³⁺-Catalyzed Methanolysis of Phosphate Diesters. Remarkable Rate Acceleration of the Methanolysis of Diphenyl Phosphate, Methyl *p*-Nitrophenyl Phosphate, and Bis(*p*-nitrophenyl) Phosphate" *Inorg. Chem.* **2001(a)**, 40, 3588.
- Neverov, A. A.; McDonald, T.; Gibson, G.; Brown, R. S., "Catalysis of Transesterification Reactions by Lanthanides — Unprecedented Acceleration of Methanolysis of Aryl and Alkyl Esters Promoted by $\text{La}(\text{OTf})_3$ at Neutral pH and Ambient Temperatures" *Can. J. Chem.* **2001(b)**, 79, 1704.
- Neverov, A. A.; Montoya-Pelaez, P.J.; Brown, R. S., "Catalysis of the Methanolysis of Activated Amides by Divalent and Trivalent Metal Ions. The Effect of Zn^{2+} , Co^{2+} , and La^{3+} on

the Methanolysis of Acetylmidazole and Its $(\text{NH}_3)_5\text{Co}^{\text{III}}$ Complex" *J. Am. Chem. Soc.* **2001**(c), 123, 210.

Rived, F.; Rosés, M.; Bosch, E., "Dissociation Constants of Neutral and Charged Acids in Methyl Alcohol. The Acid Strength Resolution" *Anal. Chim. Acta* **1998**, 374, 309.

Tsang, J.; Neverov, A. A.; Brown, R. S., "Billion-Fold Acceleration of the Methanolysis of Paraoxon Promoted by $\text{La}(\text{OTf})_3$ in Methanol" *J. Am. Chem. Soc.* **2003**, 125, 7602.


Yang, Y.-C.; Berg, F. J.; Szafraniec, L. L.; Beaudry, W. T.; Bunton, C. A.; Kumar, A., "Peroxyhydrolysis of Nerve Agent VX and Model Compounds and Related Nucleophilic Reactions" *J. Chem. Soc. Perkin Trans. 2* **1997**, 607.

Yang, Y.-C., "Chemical Detoxification of Nerve Agent VX" *Acc. Chem. Res.* **1999**, 32, 109-115.

We claim:

1. A method for decomposing an organophosphorus compound comprising subjecting said organophosphorus compound to an alcoholysis reaction in a medium comprising non-radioactive metal ions and at least a trace amount of alkoxide ions, wherein, through said alcoholysis reaction, said organophosphorus compound is decomposed.

2. The method of claim 1, wherein said organophosphorus compound has the following formula (10):



(10)

where:

J is O or S;

X, G, Z are the same or different and are selected from the group consisting of Q, OQ, QA, OA, F, Cl, Br, I, QS, SQ and C≡N;

Q is hydrogen or a substituted or unsubstituted branched, straight-chain or cyclic alkyl group having 1-100 carbon atoms; and

A is a substituted or unsubstituted aryl group selected from the group consisting of phenyl, biphenyl, benzyl, pyridyl, naphthyl, polynuclear aromatic, and aromatic and non-aromatic heterocyclic;

wherein, when X, G, Z are the same,

(i) X, G, Z are not Q; or

(ii) Q is not H; and

wherein said substituents are selected from the group consisting of Cl, Br, I, F, nitro, nitroso, Q, alkenyl, OQ, carboxyalkyl, acyl, SO₃H, SO₃Q, S=O(Q), S(=O)₂Q, amino, alkylamino (NHQ), arylamino (NHA), alkylaryl amino, dialkylamino and diarylamino.

3. The method of claim 1, wherein said medium is a solution further comprising a solvent selected from the group consisting of methanol, substituted and unsubstituted primary, secondary and tertiary alcohols, alkoxyalkanol, aminoalkanol, and combinations thereof.

4. The method of claim 1, wherein said organophosphorus compound has at least one phosphorus atom double bonded to an oxygen or a sulfur atom.

5. The method of claim 1, wherein said medium further comprises a non-inhibitory buffering agent.

6. The method of claim 5, wherein said buffering agent is selected from the group consisting of anilines, N-alkylanilines, N,N-dialkylanilines, N-alkylmorpholines, N-alkylimidazoles, 2,6-dialkylpyridines, primary, secondary and tertiary amines, trialkylamines, and combinations thereof.

7. The method of claim 1, wherein said medium is a solution further comprising a solvent selected from the group consisting of methanol, ethanol, n-propanol, *iso*-propanol, n-butanol, 2-butanol, methoxyethanol, and combinations thereof.

8. The method of claim 7, wherein said solution further comprises a solvent selected from the group consisting of nitriles, esters, ketones, amines, ethers, hydrocarbons, substituted hydrocarbons, unsubstituted hydrocarbons, chlorinated hydrocarbons, and combinations thereof.

9. The method of claim 1, wherein said medium further comprises alkoxide ions in addition to said at least a trace amount of alkoxide ions.

10. The method of claim 9, wherein the concentration of said alkoxide ions is about 0.1 to about 2 equivalents of the concentration of the metal ions.

11. The method of claim 9, wherein the concentration of said alkoxide ions is about 1 to about 1.5 equivalents of the concentration of the metal ions.

12. The method of claim 1, wherein said medium is prepared by combining a metal salt and an alkoxide salt with at least one of alcohol, alkoxyalkanol and aminoalkanol.

13. The method of claim 1, wherein said metal ions are selected from the group consisting of lanthanide series metal ions, transition metal ions, and combinations thereof.

14. The method of claim 1, wherein said metal ions are selected from the group consisting of lanthanide series metal ions, copper, platinum, palladium, zinc, nickel, yttrium, scandium ions, and combinations thereof.

15. The method of claim 1, wherein said metal ions are selected from the group consisting of Cu^{2+} , Pt^{2+} , Pd^{2+} , Zn^{2+} , Y^{3+} , Sc^{3+} , Ce^{3+} , La^{3+} , Pr^{3+} , Nd^{3+} , Sm^{3+} , Eu^{3+} , Gd^{3+} , Tb^{3+} , Dy^{3+} , Ho^{3+} , Er^{3+} , Tm^{3+} , Yb^{3+} , and combinations thereof.

16. The method of claim 1, wherein said metal ions are lanthanide series metal ions.

17. The method of claim 16, wherein said lanthanide series metal ions are selected from the group consisting of Ce^{3+} , La^{3+} , Pr^{3+} , Nd^{3+} , Sm^{3+} , Eu^{3+} , Gd^{3+} , Tb^{3+} , Dy^{3+} , Ho^{3+} , Er^{3+} , Tm^{3+} , Yb^{3+} , and combinations thereof.
18. The method of claim 1, wherein said metal ions are selected from the group consisting of Cu^{2+} , Pt^{2+} , Pd^{2+} , Zn^{2+} , and combinations thereof.
19. The method of claim 1, wherein said metal ions are selected from the group consisting of Y^{3+} , Sc^{3+} , and combinations thereof.
20. The method of claim 1, wherein said metal ion is La^{3+} .
21. The method of claim 1, wherein said organophosphorus compound is a pesticide.
22. The method of claim 1, wherein said organophosphorus compound is an insecticide.
23. The method of claim 1, wherein said organophosphorus compound is paraoxon.
24. The method of claim 1, wherein said organophosphorus compound is a chemical warfare agent.
25. The method of claim 24, wherein said organophosphorus compound is a G-agent.
26. The method of claim 24, wherein said organophosphorus compound is selected from the group consisting of VX and Russian-VX.
27. The method of claim 24, wherein said organophosphorus compound is a nerve agent.
28. The method of claim 24, wherein said chemical warfare agent is combined with a polymer.
29. The method of claim 1, wherein said medium further comprises one or more ligands.
30. The method of claim 29, wherein said ligand is selected from the group consisting of 2,2'-bipyridyl, 1,10-phenanthryl, 2,9-dimethylphenanthryl, crown ether, and 1,5,9-triazacyclododecyl.
31. The method of claim 29, wherein said ligand further comprises solid support material.

32. The method of claim 31, wherein said solid support material is selected from a polymer, silicate, aluminate, and combinations thereof.

33. The method of claim 1, wherein said medium is a solid.

34. The method of claim 1, wherein said medium is a solution

35. . The method of claim 34, wherein said solution is disposed on an applicator.

36. The method of claim 9, wherein the concentration of said alkoxide ions is about 0.01 to about 2 equivalents of the concentration of the metal ions.

37. The method of claim 9, wherein the concentration of said alkoxide ions is about 0.5 to about 1.5 equivalents of the concentration of the metal ions.

38. A kit for decomposing an organophosphorus compound comprising a substantially non-aqueous medium for an alcoholysis reaction, said medium comprising non-radioactive metal ions and at least a trace amount of alkoxide ions.

39. The kit of claim 38, wherein said medium is contained in an ampule.

40. The kit of claim 38, comprising an applicator bearing the medium, said applicator being adapted so that the medium is applied to the organophosphorus compound and the compound decomposes.

41. The kit of claim 38, further comprising written instructions for use.