

POLYMORPH FORMS OF

(2S)-(4E)-N-METHYL-5-(3-ISOPROPOXYPYRIDIN)YL]-4-PENTEN-2-AMINE FOR THE
TREATMENT OF CENTRAL NERVOUS SYSTEM DISORDERS

This application claims benefit of U.S. Provisional Patent Application No.
5 60/746,821, filed May 9, 2006.

Field of the Invention

The present invention relates to novel polymorph forms of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine p-hydroxybenzoate, as well
10 as pharmaceutical compositions including the polymorph forms. The present invention also relates to methods for treating a wide variety of conditions and disorders, and particularly conditions and disorders associated with dysfunction of the central and autonomic nervous systems, using the novel polymorph forms.

15

Background of the Invention

In the formulation of drug compositions, it is important for the drug substance to be in a form in which it can be conveniently handled and processed. This is of importance, not only from the point of view of obtaining a commercially-viable manufacturing process, but also from the point of view of subsequent manufacture of
20 pharmaceutical compositions comprising the active compound.

Further, in the manufacture of drug compositions, it is important that a reliable, reproducible and constant plasma concentration profile of drug is provided following administration to a patient.

Chemical stability, solid state stability, and "shelf life" of the active
25 ingredients are also very important factors. The drug substance, and compositions containing it, should preferably be capable of being effectively stored over appreciable periods of time, without exhibiting a significant change in the active component's physico-chemical characteristics (e.g. its chemical composition, density, hygroscopicity and solubility).

30 Moreover, it is also important to be able to provide drugs in a form, that are as chemically pure as possible.

The skilled person will appreciate that, typically, if a drug can be readily obtained in a stable form, such as a stable crystalline form, advantages may be

provided, in terms of ease of handling, ease of preparation of suitable pharmaceutical compositions, and a more reliable solubility profile.

The compound (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine is known to provide benefits in the area of the treatment and/or prevention of central nervous system disorders. The compound, its synthesis, and its use in methods of medical treatment, is described, for example, in PCT WO 99/65876 to Caldwell et al. and in US application 11/270,018, the contents of which are hereby incorporated by reference in their entirety.

10

Summary of the Invention

The present invention relates to two polymorphic forms of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine p-hydroxybenzoate (hereinafter referred to as Polymorphs A and B), methods of preparing the polymorph forms, and pharmaceutical compositions including the polymorph forms. Polymorphs A and B of the p-hydroxybenzoate salt of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine are specifically disclosed.

The present invention also relates to methods for treating and/or preventing a wide variety of conditions or disorders, and particularly those disorders characterized by dysfunction of nicotinic cholinergic neurotransmission including disorders involving neuromodulation of neurotransmitter release, such as dopamine release. The present invention also relates to methods for treating and/or preventing disorders, such as central nervous system (CNS) disorders, which are characterized by an alteration in normal neurotransmitter release, and also for treating certain conditions (e.g., alleviating pain). The methods involve administering to a subject an effective amount of the novel polymorph forms, or pharmaceutical compositions including such polymorph forms.

The polymorph forms can be provided in the form of a pharmaceutical composition that includes an effective amount of the one of the polymorph forms or mixtures thereof, as described herein. The pharmaceutical compositions incorporating the polymorph form(s) of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine p-hydroxybenzoate, which, when employed in effective amounts, interacts with relevant nicotinic receptor sites of a subject, and hence has acts as a therapeutic agent to treat and prevent a wide variety of conditions and disorders,

particularly those disorders characterized by an alteration in normal neurotransmitter release.

The pharmaceutical compositions are believed to be safe and effective with regard to prevention and treatment of a wide variety of conditions and disorders as described below.

The foregoing and other aspects of the present invention are explained in detail in the detailed description and examples set forth below.

Detailed Description

Two polymorph forms of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine p-hydroxybenzoate, pharmaceutical compositions, methods of preparing these forms, and methods of treatment and/or prevention using these forms, are described in detail below.

According to a further aspect of the invention there is provided a salt of the invention in substantially crystalline form. Although we have found that it is possible to produce salt forms of the invention in forms which are greater than 80% crystalline, by "substantially crystalline" we include greater than 20%, preferably greater than 30%, and more preferably greater than 40% (e.g. greater than any of 50, 60, 70, 80 or 90%) crystalline.


According to a further aspect of the invention there is also provided a salt forms of the invention in partially crystalline form. By "partially crystalline" we include 5% or between 5% and 20% crystalline. The degree (%) of crystallinity may be determined by the skilled person using X-ray powder diffraction (XRPD). Other techniques, such as solid state NMR, FT-IR, Raman spectroscopy, differential scanning calorimetry (DSC) and microcalorimetry, may also be used.

The term "stability" as defined herein includes chemical stability and solid state stability. By "chemical stability", we include that it may be possible to store salt forms of the invention in an isolated form, or in the form of a composition in which it is provided in admixture with pharmaceutically acceptable carriers, diluents or adjuvants (e.g. in an oral dosage form, such as a tablet, capsule etc.), under normal storage conditions, with an insignificant degree of chemical degradation or decomposition. By "solid state stability", we include that it may be possible to store salt forms of the invention in an isolated solid form, or in the form of a solid

composition in which it is provided in admixture with pharmaceutically acceptable carriers, diluents or adjuvants (e.g. in an oral dosage form, such as a tablet, capsule etc.), under normal storage conditions, with an insignificant degree of solid state transformation (e.g. crystallisation, recrystallisation, solid state phase transition, hydration, dehydration, solvation or desolvation). Examples of “normal storage conditions” include temperatures of between minus 80 and plus 50°C (preferably between 0 and 40°C and more preferably room temperatures, such as 15 to 30°C), pressures of between 0.1 and 2 bars (preferably at atmospheric pressure), relative humidities of between 5 and 95% (preferably 10 to 60%), and/or exposure to 460 lux of UV/visible light, for prolonged periods (i.e. greater than or equal to six months). Under such conditions, salt forms of the invention may be found to be less than 15%, more preferably less than 10%, and especially less than 5%, chemically degraded/decomposed, or solid state transformed, as appropriate. The skilled person will appreciate that the above-mentioned upper and lower limits for temperature, pressure and relative humidity represent extremes of normal storage conditions, and that certain combinations of these extremes will not be experienced during normal storage (e.g. a temperature of 50°C and a pressure of 0.1 bar).

I. (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine

The compounds described herein are polymorphs of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine p-hydroxybenzoate, which has the formula:



25

The manner in which the (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine p-hydroxybenzoate, in its polymorphic forms, can be prepared can vary. Approaches for preparing (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-

penten-2-amine are described, for example, in PCT WO 99/65876 to Caldwell et al. and in US application 11/270,018, pertinent portions of which are summarized below.

One synthetic approach involves a convergent synthesis, in which the side chain, (2S)-N-methyl-N-(tert-butoxycarbonyl)-4-penten-2-amine is coupled with the
5 3-substituted 5-halo-substituted pyridine, 5-bromo-3-isopropoxypyridine, under Heck reaction conditions, followed by removal of the tert-butoxycarbonyl protecting group. Typically, the types of procedures set forth in W. C. Frank et al., *J. Org. Chem.* **43**: 2947 (1978) and N. J. Malek et al., *J. Org. Chem.* **47**: 5395 (1982) involving a palladium-catalyzed coupling of an olefin and an aromatic halide are used. The
10 required (2S)-N-methyl-N-(tert-butoxycarbonyl)-4-penten-2-amine can be synthesized as follows: (i) (2R)-4-Penten-2-ol can be prepared from (R)-(+)-propylene oxide, according to procedures set forth in A. Kalivretenos, J. K. Stille, and L. S. Hegedus, *J. Org. Chem.* **56**: 2883 (1991), and then treated with p-toluenesulfonyl chloride in pyridine to yield (2R)-4-penten-2-ol p-toluenesulfonate. (ii) The resulting tosylate can
15 be heated with 20 molar equivalents of methylamine (as a 40% aqueous solution) in dimethylformamide to yield (2S)-N-methyl-4-penten-2-amine. (iii) The resulting amine can be allowed to react with di-tert-butyl dicarbonate in ether to yield the side chain, (2S)-N-methyl-N-(tert-butoxycarbonyl)-4-penten-2-amine.

The halo-substituted pyridine, (e.g., 5-bromo-3-isopropoxypyridine) can be
20 synthesized by two different routes. In one preparation, 3,5-dibromopyridine is heated at 140°C for 14 hours with 2 molar equivalents of potassium isopropoxide in dry isopropanol in the presence of copper powder (5%, w/w of the 3,5-dibromopyridine) in a sealed glass tube to yield 5-bromo-3-isopropoxypyridine. A second preparation of 5-bromo-3-isopropoxypyridine from 5-bromonicotinic acid can be performed as
25 follows: (i) 5-Bromonicotinic acid is converted to 5-bromonicotinamide by treatment with thionyl chloride, followed by reaction of the intermediate acid chloride with aqueous ammonia. (ii) The resulting 5-bromonicotinamide, previously described by C. V. Greco et al., *J. Heterocyclic Chem.* **7(4)**: 761 (1970), is subjected to Hofmann degradation by treatment with sodium hydroxide and a 70% solution of calcium
30 hypochlorite. (iii) The resulting 3-amino-5-bromopyridine, previously described by C. V. Greco et al., *J. Heterocyclic Chem.* **7(4)**: 761 (1970), can be converted to 5-bromo-3-isopropoxypyridine by diazotization with isoamyl nitrite under acidic conditions, followed by treatment of the intermediate diazonium salt with isopropanol to yield 5-

bromo-3-isopropoxypyridine. The palladium-catalyzed coupling of 5-bromo-3-isopropoxypyridine and (2S)-N-methyl-N-(tert-butoxycarbonyl)-4-penten-2-amine is carried out in acetonitrile-triethylamine (2:1, v,v) using a catalyst consisting of 1 mole % palladium(II) acetate and 4 mole % tri-o-tolylphosphine. The reaction can be
5 carried out by heating the components at 80°C for 20 hours to yield (2S)-(4E)-N-methyl-N-(tert-butoxycarbonyl)-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine. Removal of the tert-butoxycarbonyl protecting group can be accomplished by treatment with 30 molar equivalents of trifluoroacetic acid in anisole at 0°C to afford (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine.

10

II. p-Hydroxybenzoate Salt of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine

Typically the p-hydroxybenzoate salt of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine is prepared by reacting about one equivalent
15 of p-hydroxybenzoic acid with (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine in some suitable solvent, such as methanol, ethanol, isopropyl alcohol, acetone, ethyl acetate, or acetonitrile. Sometimes water is present in the reaction mixture. These procedures are described in detail in US application 11/270,018.

20

III. Polymorphs of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine - hydroxybenzoate salt

Polymorphs A and B of the p-hydroxybenzoate salt of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine are disclosed.

25 The polymorph forms of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine p-hydroxybenzoate are prepared by recrystallizing the salt in various solvents. Representative procedures are given in the Examples. A variety of solvents may be utilized in the formation of polymorphs A and B. Appropriate choice of solvent(s) for recrystallization provides either polymorph A or B in relatively pure
30 form (i.e., greater than 95% polymorph A or greater than 95% polymorph B). The invention thus provides both polymorph A and polymorph B in relatively pure form (greater than 95% polymorphic purity).

Depending upon the manner by which the polymorphs described herein are formed, the polymorph can have crystal structures that occlude solvents that are present during polymorph formation. Thus, the polymorph can occur as hydrates and other solvates of varying stoichiometry of solvent relative to the (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin-2-yl)]-4-penten-2-amine p-hydroxybenzoate. Representative solvents that can be used to prepare and/or recrystallize the polymorphs include, without limitation, ethanol, methanol, isopropyl alcohol, acetone, ethyl acetate, and/or acetonitrile.

A series of solid-state techniques can be used to evaluate two polymorphic forms of the p-hydroxybenzoate salt of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin-2-yl)]-4-penten-2-amine. X-ray Powder Diffraction (XRPD) was used to evaluate the unique crystalline phase and structural characteristics of the polymorphs and mixtures. Differential Scanning Calorimetry (DSC) may also be performed.

One aspect of the invention relates to a mixture of polymorph A and polymorph B. In one embodiment the ratio of polymorph A to B is 40 to 60. In another embodiment the ratio is 30 to 70. In a further embodiment the ratio is 20 to 80. In yet another embodiment the ratio is 10 to 90. In yet another embodiment the ratio is 5 to 95. Likewise, in one embodiment the ratio of polymorph B to A is 40 to 60. In another embodiment the ratio is 30 to 70. In a further embodiment the ratio is 20 to 80. In yet another embodiment the ratio is 10 to 90. In yet another embodiment the ratio is 5 to 95. Thus, the invention also provides mixtures of polymorphs A and B in ratios ranging from 5:95 to 95:5.

IV. Methods of Treatment

The salt forms described herein can be used in methods for preventing and/or treating a condition or disorder in a subject susceptible to such a condition or disorder. For example, an effective amount of a compound effective for providing some degree of prevention of the progression of a CNS disorder (i.e., provide protective effects), amelioration of the symptoms of a CNS disorder, and amelioration of the reoccurrence of a CNS disorder, can be administered to a patient in need thereof.

The compounds can be used to treat and/or prevent those types of conditions and disorders for which other types of nicotinic compounds have been proposed as

therapeutics. See, for example, Williams et al. DN&P 7(4):205-227 (1994), Arneric et al., CNS Drug Rev. 1(1):1-26 (1995), Arneric et al., Exp. Opin. Invest. Drugs 5(1):79-100 (1996), Bencherif et al., JPET 279:1413 (1996), Lippiello et al., JPET 279:1422 (1996), Damaj et al., Neuroscience (1997), Holladay et al., J. Med. Chem 40(28): 4169-4194 (1997), Bannon et al., Science 279: 77-80 (1998), PCT WO 94/08992, PCT WO 96/31475, and U.S. Patent Nos. 5,583,140 to Bencherif et al., 5,597,919 to Dull et al., and 5,604,231 to Smith et al the disclosures of which are incorporated herein by reference in their entirety.

The compounds modulate nicotinic receptors in the patient's brain. As such, such compounds have the ability to express nicotinic pharmacology, and in particular, to act as nicotinic partial agonists.

Receptor binding constants provide a measure of the ability of the compound to bind to half of the relevant receptor sites of certain brain cells of the patient. See, Cheng, et al., Biochem. Pharmacol. 22:3099 (1973). The (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine compound used to prepare the salt forms has extremely high affinity for the relevant receptors, with a binding affinity in the low nM range.

The compounds have the ability to demonstrate a nicotinic function by effectively modulating neurotransmitter secretion from neurons. As such, such compounds have the ability to affect relevant the release of acetylcholine, dopamine, and other neurotransmitters by neurons.

The compounds, when employed in effective amounts in accordance with the methods described herein, are selective to certain relevant nicotinic receptors, but do not cause significant activation of receptors associated with undesirable side effects at concentrations at least greater than those required for modulation of CNS neuronal activity. The selectivity of the compounds against those ganglia-type receptors responsible for cardiovascular side effects is demonstrated by a lack of the ability of those compounds to activate nicotinic function of adrenal chromaffin tissue at concentrations greater than those required for modulation of CNS neuronal activity.

Thus, administration of the compounds provides a therapeutic window in which treatment of certain CNS disorders is provided, and certain side effects are avoided. That is, an effective dose of the compound is sufficient to provide the desired effects upon the CNS, but is insufficient (i.e., is not at a high enough level) to

provide undesirable side effects. The pharmaceutical compositions are useful in the treatment of a variety of CNS disorders, including neurodegenerative disorders, neuropsychiatric disorders, neurologic disorders, and addictions. The pharmaceutical compositions can be used to treat cognitive deficits (age-related and otherwise),
5 attentional disorders and dementias (including those due to infectious agents or metabolic disturbances); to provide neuroprotection; to treat convulsions and multiple cerebral infarcts; to treat mood disorders, compulsions and addictive behaviors; to provide analgesia; and to control inflammation (such as mediated by cytokines and nuclear factor kappa B) and treat inflammatory disorders. Among the disorders,
10 diseases and conditions, that pharmaceutical compositions of the present invention can be used to treat, are: age-associated memory impairment, mild cognitive impairment, pre-senile dementia (early onset Alzheimer's disease), senile dementia (dementia of the Alzheimer's type), Lewy body dementia, vascular dementia, Alzheimer's disease, stroke, AIDS dementia complex, attention deficit disorder,
15 attention deficit hyperactivity disorder, dyslexia, schizophrenia, schizophreniform disorder, schizoaffective disorder, Parkinsonism including Parkinson's disease, Pick's disease, Huntington's chorea, tardive dyskinesia, hyperkinesia, progressive supranuclear palsy, Creutzfeld-Jakob disease, multiple sclerosis, amyotrophic lateral sclerosis, epilepsy, depression, panic disorders, bipolar disorders, generalized anxiety
20 disorder, obsessive compulsive disorder, rage outbursts, Tourette's syndrome, autism, drug and alcohol addiction, tobacco addiction, obesity, acute pain, neuropathic pain, inflammatory pain, ulcerative colitis, irritable bowel syndrome, cachexia, osteoarthritis, psoriasis, rheumatoid arthritis, endotoxaemia, sepsis, asthma, atherosclerosis and idiopathic pulmonary fibrosis.

25 Thus, the present invention relates to the salt forms mentioned above for use in therapy. The present invention further relates to the use of said salt forms, in the manufacture of a medicament for treatment of a central nervous system disorder. Also provided is a method for treatment of a central nervous system disorder, comprising administering to a mammal in need of such treatment, a therapeutically effective
30 amount of the salt forms of the present invention. Further provided is a method for treatment of disorders selected from the group consisting of age-associated memory impairment, mild cognitive impairment, pre-senile dementia (early onset Alzheimer's disease), senile dementia (dementia of the Alzheimer's type), Lewy body dementia,

vascular dementia, Alzheimer's disease, stroke, AIDS dementia complex, attention deficit disorder, attention deficit hyperactivity disorder, dyslexia, schizophrenia, schizophreniform disorder, and schizoaffective disorder. Even further provided is a method for treatment of disorders selected from the group consisting of the treatment
5 of mild to moderate dementia of the Alzheimer's type, attention deficit disorder, mild cognitive impairment and age associated memory impairment.

V. Pharmaceutical Compositions

According to one embodiment of the present invention there is provided a
10 pharmaceutical composition comprising as active ingredient a therapeutically effective amount of salt forms of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine, in association with one or more pharmaceutically acceptable diluents, excipients and/or inert carriers.

The manner in which the compositions are administered can vary. The
15 compounds can be administered by inhalation (e.g., in the form of an aerosol either nasally or using delivery articles of the type set forth in U.S. Patent No. 4,922,901 to Brooks et al., the disclosure of which is incorporated herein in its entirety); topically (e.g., in lotion form); orally (e.g., in liquid form within a solvent such as an aqueous or non-aqueous liquid; intravenously (e.g., within a dextrose or saline solution); as an
20 infusion or injection (e.g., as a suspension or as an emulsion in a pharmaceutically acceptable liquid or mixture of liquids); intrathecally; intracerebroventricularly; or transdermally (e.g., using a transdermal patch or by powder injection). Although it is possible to administer the compositions in the form of a bulk active chemical, it is preferred to present each compound in the form of a pharmaceutical composition for
25 efficient and effective administration. Exemplary methods for administering such compounds will be apparent to the skilled artisan. For example, the compositions can be administered in the form of a tablet, a hard gelatin capsule or as a time release capsule. The administration of the pharmaceutical compositions described herein can be intermittent, or at a gradual, continuous, constant or controlled rate to a warm-blooded animal, (e.g., a mammal such as a mouse, rat, cat, rabbit, dog, pig, cow, or
30 monkey); but advantageously is preferably administered to a human being. In addition, the time of day and the number of times per day that the pharmaceutical composition is administered can vary. Administration preferably is such that the active

ingredients of the pharmaceutical composition interact with receptor sites within the body of the subject that effect the functioning of the CNS. More specifically, in treating a CNS disorder administration preferably is such so as to optimize the effect upon those relevant receptor subtypes, which have an effect upon the functioning of the CNS, while minimizing the effects upon muscle-type receptor subtypes.

The appropriate dose of the compound is that amount effective to prevent occurrence of the symptoms of the disorder or to treat some symptoms of the disorder from which the patient suffers. By "effective amount", "therapeutic amount" or "effective dose" is meant that amount sufficient to elicit the desired pharmacological or therapeutic effects, thus resulting in effective prevention or treatment of the disorder. Thus, when treating a CNS disorder, an effective amount of compound is an amount sufficient to pass across the blood-brain barrier of the subject, to bind to relevant receptor sites in the brain of the subject, and to modulate the activity of relevant nicotinic receptor subtypes (e.g., modulate neurotransmitter secretion, thus resulting in effective prevention or treatment of the disorder). Prevention of the disorder is manifested by delaying the onset of the symptoms of the disorder. Treatment of the disorder is manifested by a decrease in the symptoms associated with the disorder or an amelioration of the reoccurrence of the symptoms of the disorder.

The effective dose can vary, depending upon factors such as the condition of the patient, the severity of the symptoms of the disorder, and the manner in which the pharmaceutical composition is administered. For human patients, the effective dose of typical compounds generally requires administering the compound in an amount sufficient to modulate relevant receptors to effect neurotransmitter (e.g., dopamine) release but the amount should be insufficient to induce effects on skeletal muscles and ganglia to any significant degree. The effective dose of compounds will of course differ from patient to patient but in general includes amounts starting where CNS effects or other desired therapeutic effects occur, but below the amount where muscular and ganglionic effects are observed.

Typically, the effective dose of compounds may require administering the compound in an amount of less than 5 mg/kg of patient weight. Often, the compounds may be administered in an amount from less than about 1 mg/kg patient weight to less than about 100 µg/kg of patient weight, and occasionally between from 10 µg/kg to less than 100 µg/kg of patient weight. The foregoing effective doses

typically represent that amount administered as a single dose, or as one or more doses administered over a 24 hour period.

For human patients, the effective dose of the compounds may require administering the compound in an amount of at least about 1, but not more than about 5 1000, often not more than about 500 mg/ 24 hr./ patient.

The compounds also can be administered in composition compositions that incorporate other ingredients, such as those types of ingredients that are useful in formulating a diagnostic composition. Compositions useful as diagnostics can be employed as set forth in U.S. Patent Nos. 5,853,696 to Elmalch et al. and 5,969,144 to 10 London et al., the contents of which are hereby incorporated by reference.

The compounds can also be formulated and/or administered in combination with other therapeutic compounds, such as those used in the treatment and or prevention of CNS disorders.

US Provisional Application Serial Nos. 60/746,808 and 60/746,821 are 15 incorporated herein by reference in their entirety.

VI: Experimentals

Example 1: Analysis of the Two Polymorphic Forms

20 The Polymorphs A and B were generated as described in Example 4.

A. Methods

X-ray powder diffraction analyses were performed using a PANalytical X'Pert Pro MPD diffractometer for 96 minutes from 1 to 60° 2θ with and without internal standard reference. The 2θ angles were corrected with regard to the standard values 25 whereafter calculation into d-values (distance values) was done. The d-values may vary in the range ±2 on the last given decimal place. The sample preparation was performed according to standard methods, for example those described in Giacovazzo, C. et al (1995), Fundamentals of Crystallography, Oxford University Press; Jenkins, R. and Snyder, R. L. (1996), Introduction to X-Ray Powder Diffractometry, John 30 Wiley & Sons, New York; Bunn, C. W. (1948), Chemical Crystallography, Clarendon Press, London or Klug, H. P. & Alexander, L. E. (1974), X-ray Diffraction Procedures, John Wiley and Sons, New York.

The term “stability” as defined herein includes chemical stability and solid state stability.

By “chemical stability”, we include that it may be possible to store salt forms of the invention in an isolated form, or in the form of a formulation in which it is provided in admixture with pharmaceutically acceptable carriers, diluents or adjuvants (e.g. in an oral dosage form, such as a tablet, capsule etc.), under normal storage conditions, with an insignificant degree of chemical degradation or decomposition.

By “solid state stability”, we include that it may be possible to store salt forms of the invention in an isolated solid form, or in the form of a solid formulation in which it is provided in admixture with pharmaceutically acceptable carriers, diluents or adjuvants (e.g. in an oral dosage form, such as a tablet, capsule etc.), under normal storage conditions, with an insignificant degree of solid state transformation (e.g. crystallisation, recrystallisation, solid state phase transition, hydration, dehydration, solvation or desolvation).

Examples of “normal storage conditions” include temperatures of between minus 80 and plus 50°C (preferably between 0 and 40°C and more preferably room temperatures, such as 15 to 30°C), pressures of between 0.1 and 2 bars (preferably at atmospheric pressure), relative humidities of between 5 and 95% (preferably 10 to 60%), and/or exposure to 460 lux of UV/visible light, for prolonged periods (i.e. greater than or equal to six months). Under such conditions, salt forms of the invention may be found to be less than 15%, more preferably less than 10%, and especially less than 5%, chemically degraded/decomposed, or solid state transformed, as appropriate. The skilled person will appreciate that the above-mentioned upper and lower limits for temperature, pressure and relative humidity represent extremes of normal storage conditions, and that certain combinations of these extremes will not be experienced during normal storage (e.g. a temperature of 50°C and a pressure of 0.1 bar).

B. Results

Characterization of Polymorph A

A sample of pure Polymorph A was provided and tested “As-Is” by XRPD and shown in table 1.

Characterization of Polymorph B

A sample of pure Polymorph B was provided and tested "As-Is" by XRPD and shown in table 2.

The diffractability and preferred orientation of the two polymorphic forms were studied to determine if a method could be developed to minimize preferred orientation without increasing the amorphous content or altering the polymorphic form. This study involved attempts to randomize the sample presentation such that during data collection the powder pattern intensities remain somewhat constant. The crystals must be reduced to mimic the size of the other samples.

10 Table 1. Selected Unique X-Ray Powder Peaks for Polymorphs A.

Corrected 2 θ	distance (Å)	Rel intensity
6.4	13.9	vw
7.6	11.7	vs
8.2	10.7	w
10.2	8.6	vw
14.7	6.0	vw
15.2	5.8	s
16.4	5.4	vw
17.7	5.00	vw
18.1	4.89	vw
19.0	4.67	w
19.7	4.51	m
21.9	4.05	vw
22.2	4.01	vw
22.6	3.93	vw
22.8	3.89	vw
23.5	3.79	m
23.8	3.73	vw
24.2	3.68	vw
27.2	3.28	w
28.5	3.13	vw

30.6	2.92	m
34.9	2.57	vw
36.6	2.46	vw

Definitions used:

% Relative intensity*	Definition
100 - 50	vs (very strong)
50-10	s (strong)
10-2	m (medium)
2-0.7	w (weak)
0.7 >	vw (very weak)

An embodiment of the invention is 2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine p-hydroxybenzoate which exhibits at least the following characteristic X-ray powder diffraction peaks (expressed in degrees 2 θ):

- (1) 7.6 and 15.2 and 19.7, or
- (2) 6.4, 7.6, 15.2 and 19.7, or
- (3) 6.4, 7.6, 15.2, 19.7 and 30.6, or
- (4) 6.4, 7.6, 8.2, 10.2, 15.2, 19.7 and 30.6.

*The relative intensities are derived from diffractograms measured with variable slits.

Table 2. Selected Unique X-Ray Powder Peaks for Polymorphs B.

Corrected 2 θ	distance (Å)	Rel intensity
7.7	11.4	s
8.0	11.0	vs

9.6	9.2	m
10.5	8.4	vw
15.5	5.7	m
16.1	5.5	s
16.9	5.2	m
18.4	4.82	w
19.0	4.66	w
21.1	4.20	vw
21.9	4.05	vw
22.9	3.87	vw
23.5	3.79	vw
23.7	3.75	vw
27.0	3.30	vw

Definitions used:

% Relative intensity*	Definition
100 - 70	vs (very strong)
70 - 40	s (strong)
40-10	m (medium)
10-5	w (weak)
5 >	vw (very weak)

*The relative intensities are derived from diffractograms measured with variable slits.

An embodiment of the invention is (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine p-hydroxybenzoate which exhibits at least
 5 the following characteristic X-ray powder diffraction peaks (expressed in degrees 2 θ):

- 1) 8.0, 9.6 and 16.1, or
- 2) 8.0, 9.6, 16.1 and 16.9, or
- 3) 8.0, 9.6, 16.1, 16.9 and 18.4, or
- 4) 8.0, 9.6, 10.5, 16.1, 16.9 and 18.4.

10

Example 2: Determination of Binding to Relevant Receptor Sites

Binding of the compounds to relevant receptor sites was determined in accordance with the techniques described in U.S. Patent No. 6,953,855 to Mazurov et al. Inhibition constants (K_i values), reported in nM, were calculated from the IC_{50} values using the method of Cheng et al., Biochem, Pharmacol. 22:3099 (1973). Low
5 binding constants indicate that the compounds of the present invention exhibit good high affinity binding to certain CNS nicotinic receptors.

Example 3: Synthesis of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine p-hydroxybenzoate

10 (2S)-(4E)-N-Methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine p-hydroxybenzoate was prepared according to the procedures described in US application 11/270,018.

Example 4: Preparation of Polymorphs A and B

15 Polymorph A:

A 10 g sample of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine p-hydroxybenzoate was dissolved in refluxing acetone (81 ml) and 0.94 ml water. The solution was cooled to 48 °C and seeded with 100 mg of said compound. The resulting slurry was allowed to crystallize for 50 min at 48 °C. This was then
20 cooled down to 0°C during 3 hr and left overnight at 0°C. The recrystallized material was collected by vacuum filtration, washed with 40 ml acetone and dried in a vacuum oven at 60 °C, for 24 hr to give polymorph A.

Polymorph B:

25 A 10 g sample of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxypyridin)yl]-4-penten-2-amine p-hydroxybenzoate was dissolved in refluxing acetone (81 ml) and 0.94 ml water. The solution was cooled to 48 °C and seeded with 100 mg of said compound, slurried in 1 ml acetone, and additionally washed with 1 ml acetone. The resulting slurry was allowed to crystallize for 50 min at 48 °C. This was then cooled down to
30 0°C during 3 hr and left overnight at 0°C. The recrystallized material was collected by vacuum filtration, washed with 40 ml acetone and dried in a vacuum oven at 60 °C, for 72 hr to give polymorph B. This salt form was also dried up to 1 week in this last step.

The foregoing is illustrative of the present invention and is not to be construed as limiting thereof. The invention is defined by the following claims, with equivalents of the claims to be included therein.

Claims:

1. A crystalline polymorph A of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine p-hydroxybenzoate, exhibiting an X-ray powder diffraction pattern having characteristic peaks expressed in degrees ($2\theta^\circ$) at around 7.6 and 15.2 and 19.7.
2. A crystalline polymorph B of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine p-hydroxybenzoate, exhibiting an X-ray powder diffraction pattern having characteristic peaks expressed in degrees ($2\theta^\circ$) at around 8.0, 9.6 and 16.1.
3. A mixture of polymorphs comprising the polymorphs A of claim 1 and B of claim 2 in ratios of 5:95 to 95:5.
4. A polymorph of (2S)-(4E)-N-methyl-5-[3-(5-isopropoxy-pyridin)yl]-4-penten-2-amine p-hydroxybenzoate, wherein the polymorph is isolated by recrystallization from ethanol, methanol, isopropyl alcohol, acetone, ethyl acetate and/or acetonitrile.
5. The crystalline polymorph A having an X-ray powder diffraction pattern substantially the same as that shown in table 1.
6. The crystalline polymorph B having an X-ray powder diffraction pattern substantially the same as that shown in table 2.
7. The polymorph according to any one of claims 1 to 6, for use in therapy.
8. Use of the polymorph according to any one of claims 1 to 6, in the manufacture of a medicament for treatment of central nervous system disorders.

9. A method for treatment of central nervous system disorders, comprising administering to a mammal, in need of such treatment, a therapeutically effective amount of the polymorph according to any one of claims 1 to 6.

5 10. The method of Claim 9, wherein the disorder is selected from the group consisting of age-associated memory impairment, mild cognitive impairment, pre-senile dementia (early onset Alzheimer's disease), senile dementia (dementia of the Alzheimer's type), Lewy body dementia, vascular dementia, Alzheimer's disease, stroke, AIDS dementia complex, attention deficit disorder, attention deficit
10 hyperactivity disorder, dyslexia, schizophrenia, schizophreniform disorder and schizoaffective disorder.

 11. The method of Claim 9, wherein the disorders selected from the group consisting of the treatment of mild to moderate dementia of the Alzheimer's type,
15 attention deficit disorder, mild cognitive impairment and age associated memory impairment.

 12. A pharmaceutical composition comprising as active ingredient a therapeutically effective amount of the polymorph according to any one of claims 1 to
20 6, in association with one or more pharmaceutically acceptable diluents, excipients and/or inert carriers.

 13. The pharmaceutical composition according to claim 12 for treatment of central nervous system disorders.
25

 14. An agent for treatment of central nervous system disorders, which comprises as active ingredient a polymorph according to any one of claims 1 to 6.