

Substituted quinoline-2-carboxamides, their preparation and their use as pharmaceuticals, and intermediates

- 5 The invention relates to substituted quinoline-2-carboxylic acid amides, to their preparation and their use as inhibitors of prolyl 4-hydroxylase, and to their use as pharmaceuticals for the treatment of fibrotic diseases, as well as to intermediates in their
10 preparation.

- Compounds which inhibit the enzymes prolyl and lysyl hydroxylase have the effect of very selective inhibition of collagen biosynthesis by influencing collagen-specific hydroxylation reactions. In the course thereof, protein-
15 bound proline or lysine is hydroxylated by the enzymes prolyl or lysyl hydroxylase. If this reaction is suppressed by inhibitors, a nonfunctional hypohydroxylated collagen molecule which can be released by the cells into the extracellular space only in a small amount is formed.
- 20 The hypohydroxylated collagen furthermore cannot be incorporated into the collagen matrix and is very readily degraded proteolytically. As a consequence of these effects, the amount of collagen deposited extracellularly is reduced overall.

- 25 Inhibitors of prolyl hydroxylase are therefore suitable substances in the treatment of diseases in which the deposition of collagens contributes decisively to the clinical picture. These include, inter alia, fibroses of the lung, liver and skin (scleroderma and scarring after
30 burns, injuries and surgical operations) and atherosclerosis.

It is known that the enzyme prolyl hydroxylase is inhibited effectively by pyridine-2,4- and -2,5-

dicarboxylic acid (K. Majamaa et al., Eur. J. Biochem. 138 (1984) 239-245). However, these compounds are active as inhibitors in the cell culture only at very high concentrations (Tschank, G et al., Biochem. J. 238 (1987) 625 to 633).

Prodrugs of pyridine-2,4(5)-dicarboxylates are also known. These are described in EP-A-0 590 520 and EP-A-0 562 512.

N-Oxalylglycines as inhibitors of prolyl 4-hydroxylase are known from J. Med. Chem. 1992, 35, 2652 to 2658 (Cunliffe et al.), and EP-A-0 457 163 (Baader et al.).

3-Hydroxypyridine-2-carboxylic acid N-(carboxymethyl)-amide is known from G. Yolles et al. in: Bull. Soc. Chim. Fr. 1965, 8, 2252 to 2259.


Hydroxyisoquinoline- and hydroxycinnolinecarboxylic acid glycydamides are known from Biochem. Soc. Trans. 1991, 19, 812 to 815 (Franklin et al.).

EP-A-0 661 269 describes substituted heterocyclic carboxylic acid amides and their use as inhibitors of prolyl 4-hydroxylase and as inhibitors of collagen biosynthesis.

The object was to search for even more active inhibitors of prolyl hydroxylase and for other inhibitors of collagen biosynthesis.

It has now been found that a selection of the compounds included in EP-A-0 661 269, that is to say the quinoline-2-carboxylic acid amides having an OH function in the ortho position relative to the amide function, show a surprisingly high inhibition of prolyl 4-hydroxylase in cell cultures.

The compounds according to the invention correspond to the formula (I)



in which

- A is (C₁-C₄)-alkylene, which is optionally substituted by one or two substituents from the series consisting of halogen, cyano, nitro, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)Hal_g, preferably (C₁-C₈)-fluoroalkoxy, (C₁-C₈)-fluoroalkenyloxy, (C₁-C₈)-fluoroalkynyloxy, -OCF₂Cl or -O-CF₂-CHFCl, (C₁-C₆)-alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl, phenyl, benzyl, phenoxy, benzyl-oxy, anilino, N-methylanilino, phenylmercapto, phenylsulfonyl, phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl and N,N-di-(C₁-C₄)-alkylsulfamoyl, or
- by a (C₆-C₁₂)-aryloxy, (C₇-C₁₁)-aralkyloxy, (C₆-C₁₂)-aryl or (C₇-C₁₁)-aralkyl radical, which carries, in the aryl part, 1, 2, 3, 4 or 5 identical or different substituents from the series consisting of halogen, cyano, nitro, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)Hal_g, -OCF₂Cl, -O-CF₂-CHFCl, (C₁-C₆)-alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl and N,N-di-(C₁-C₄)-alkylsulfamoyl, or

by one or more substituents R^x of the α -C atom of an α -amino acid, it being possible for the naturally occurring L-amino acids and their D isomers to be used;

- 5 B is an acid grouping from the series comprising
 $-\text{CO}_2\text{H}$, $-\text{CONHCOR}'''$, $-\text{CONHSOR}'''$, $\text{CONHSO}_2\text{R}'''$, $-\text{NHSO}_2\text{CF}_3$,
 tetrazolyl, imidazolyl and 3-hydroxyisoxazolyl, in
 which R''' is aryl, heteroaryl, $(\text{C}_3\text{-C}_7)$ -cycloalkyl or
 $(\text{C}_1\text{-C}_4)$ -alkyl, optionally monosubstituted by
 10 $(\text{C}_6\text{-C}_{12})$ -aryl, heteroaryl, OH, SH, $(\text{C}_1\text{-C}_4)$ -alkyl,
 $(\text{C}_1\text{-C}_4)$ -alkoxy, $(\text{C}_1\text{-C}_4)$ -thioalkyl, -sulfinyl or
 -sulfonyl, CF_3 , Cl, Br, F, I, NO_2 , $-\text{COOH}$, $(\text{C}_2\text{-C}_9)$ -
 alkoxycarbonyl, NH_2 , mono- or di- $(\text{C}_1\text{-C}_4)$ -amino
 or $(\text{C}_1\text{-C}_4)$ -perfluoroalkyl,
- 15 R^1 , R^2 , R^3 , R^4 and R^5 are identical or different and are
 hydrogen, hydroxyl, halogen, cyano, trifluoromethyl,
 nitro, carboxyl, $(\text{C}_1\text{-C}_{20})$ -alkyl, $(\text{C}_3\text{-C}_8)$ -cycloalkyl,
 $(\text{C}_3\text{-C}_8)$ -cycloalkyl- $(\text{C}_1\text{-C}_{12})$ -alkyl, $(\text{C}_3\text{-C}_8)$ -cyclo-
 alkoxy, $(\text{C}_3\text{-C}_8)$ -cycloalkyl- $(\text{C}_1\text{-C}_{12})$ -alkoxy, $(\text{C}_3\text{-C}_8)$ -
 20 cycloalkyloxy- $(\text{C}_1\text{-C}_{12})$ -alkyl, $(\text{C}_3\text{-C}_8)$ -cycloalkyloxy-
 $(\text{C}_1\text{-C}_{12})$ -alkoxy, $(\text{C}_3\text{-C}_8)$ -cycloalkyl- $(\text{C}_1\text{-C}_8)$ -alkyl-
 $(\text{C}_1\text{-C}_6)$ -alkoxy, $(\text{C}_3\text{-C}_8)$ -cycloalkyl- $(\text{C}_1\text{-C}_8)$ -alkoxy-
 $(\text{C}_1\text{-C}_6)$ -alkyl, $(\text{C}_3\text{-C}_8)$ -cycloalkyloxy- $(\text{C}_1\text{-C}_8)$ -alkoxy-
 $(\text{C}_1\text{-C}_6)$ -alkyl, $(\text{C}_3\text{-C}_8)$ -cycloalkoxy- $(\text{C}_1\text{-C}_8)$ -alkoxy-
 25 $(\text{C}_1\text{-C}_8)$ -alkoxy, $(\text{C}_6\text{-C}_{12})$ -aryl, $(\text{C}_7\text{-C}_{16})$ -aralkyl,
 $(\text{C}_7\text{-C}_{16})$ -aralkenyl, $(\text{C}_7\text{-C}_{16})$ -aralkynyl, $(\text{C}_2\text{-C}_{20})$ -
 alkenyl, $(\text{C}_2\text{-C}_{20})$ -alkynyl, $(\text{C}_1\text{-C}_{20})$ -alkoxy, $(\text{C}_2\text{-C}_{20})$ -
 alkenyloxy, $(\text{C}_2\text{-C}_{20})$ -alkynyloxy, retinyloxy,
 $(\text{C}_1\text{-C}_{20})$ -alkoxy- $(\text{C}_1\text{-C}_{12})$ -alkyl, $(\text{C}_1\text{-C}_{12})$ -alkoxy-
 30 $(\text{C}_1\text{-C}_{12})$ -alkoxy, $(\text{C}_1\text{-C}_{12})$ -alkoxy- $(\text{C}_1\text{-C}_8)$ -alkoxy-
 $(\text{C}_1\text{-C}_8)$ -alkyl, $(\text{C}_6\text{-C}_{12})$ -aryloxy, $(\text{C}_7\text{-C}_{16})$ -aralkyloxy,
 $(\text{C}_6\text{-C}_{12})$ -aryloxy- $(\text{C}_1\text{-C}_6)$ -alkoxy, $(\text{C}_7\text{-C}_{16})$ -aralkoxy-
 $(\text{C}_1\text{-C}_6)$ -alkoxy, $(\text{C}_1\text{-C}_{16})$ -hydroxyalkyl, $(\text{C}_6\text{-C}_{16})$ -
 aryloxy- $(\text{C}_1\text{-C}_8)$ -alkyl, $(\text{C}_7\text{-C}_{16})$ -aralkoxy- $(\text{C}_1\text{-C}_8)$ -
 35 alkyl, $(\text{C}_6\text{-C}_{12})$ -aryloxy- $(\text{C}_1\text{-C}_8)$ -alkoxy- $(\text{C}_1\text{-C}_6)$ -alkyl,
 $(\text{C}_7\text{-C}_{12})$ -aralkyloxy- $(\text{C}_1\text{-C}_8)$ -alkoxy- $(\text{C}_1\text{-C}_6)$ -alkyl,
 $(\text{C}_2\text{-C}_{20})$ -alkenyloxy- $(\text{C}_1\text{-C}_6)$ -alkyl, $(\text{C}_2\text{-C}_{20})$ -

alkynyloxy-(C₁-C₆)-alkyl, retinyloxy-(C₁-C₆)-alkyl,
-O-[CH₂]_xC_fH_(2f+1-g)Hal_g, (C₁-C₆)-chlorofluoroalkoxy,
such as -OCF₂Cl, -OCF₂-CHFCl,

5 (C₁-C₂₀)-alkylcarbonyl, (C₃-C₈)-cycloalkylcarbonyl,
(C₆-C₁₂)-arylcarbonyl, (C₇-C₁₆)-aralkylcarbonyl,
cinnamoyl, (C₂-C₂₀)-alkenylcarbonyl, (C₂-C₂₀)-
alkynylcarbonyl,

10 (C₂-C₂₀)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-
alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-
aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl,
(C₂-C₂₀)-alkenyloxycarbonyl, retinyloxycarbonyl,
(C₂-C₂₀)-alkynyloxycarbonyl, (C₆-C₁₂)-aryloxy-
(C₁-C₆)-alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-
alkoxycarbonyl, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxy-
15 carbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxycarbonyl,


(C₁-C₁₂)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl-
carbonyloxy, (C₆-C₁₂)-arylcarbonyloxy, (C₇-C₁₆)-
aralkylcarbonyloxy, cinnamoyloxy, (C₂-C₁₂)-alkenyl-
carbonyloxy, (C₂-C₁₂)-alkynylcarbonyloxy,

20 (C₁-C₁₂)-alkoxycarbonyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-
alkoxycarbonyloxy, (C₆-C₁₂)-aryloxycarbonyloxy,
(C₇-C₁₆)-aralkyloxycarbonyloxy, (C₃-C₈)-cycloalkoxy-
carbonyloxy, (C₂-C₁₂)-alkenyloxycarbonyloxy,
(C₂-C₁₂)-alkynyloxycarbonyloxy,

25 carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-
(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-
carbamoyl, N,N-dicyclo-(C₃-C₈)-alkylcarbamoyl,
N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl,
N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)carbamoyl,
30 N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-
alkyl)carbamoyl, N-(+)-dehydroabietylcarbamoyl,
N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl,
N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkyl-
carbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcarbamoyl,

N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl,
 N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl,
 N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl,
 N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyl,
 5 N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-
 alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-
 aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-
 ((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyl,
 10 CON(CH₂)_h, in which a CH₂ group can be replaced by
 O, S, N-(C₁-C₈)-alkylimino, N-(C₃-C₈)-cycloalkyl-
 imino, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkylimino, N-
 (C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or N-
 (C₁-C₁₂)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7,
 or a carbamoyl radical of the formula J

15



in which

R^x is the substituent of an L- or D-α-amino acid,
 s is 1, 2, 3, 4 or 5 and
 20 T is OH, OR or NR^{*}R^{**}, in which

R^{*}, R^{**} and R^{***} are identical or different and are
 hydrogen, (C₆-C₁₂)-aryl, (C₇-C₁₁)-aralkyl,
 (C₁-C₈)-alkyl, (C₃-C₈)-cycloalkyl, (+)-dehydro-
 abietyl, (C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-
 25 aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-
 (C₁-C₈)-alkyl, (C₁-C₁₀)-alkanoyl, optionally
 substituted (C₇-C₁₆)-aralkanoyl or optionally
 substituted (C₆-C₁₂)-aroyl, or

30 R^{*} and R^{**} together are -[CH₂]_h, in which one CH₂
 group can be replaced by O, S, SO, SO₂,

5 N-acylamino, N-(C₁-C₁₀)-alkoxycarbonylimino, N-(C₁-C₈)-alkylimino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or N-(C₁-C₄)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7,

or further wherein R¹, R², R³, R⁴ and R⁵ are identical or different and are:
 carbamoyloxy, N-(C₁-C₁₂)-alkylcarbamoyloxy, N,N-di-(C₁-C₁₂)-alkylcarbamoyloxy, N-(C₃-C₈)-cycloalkylcarbamoyloxy, N-(C₆-C₁₂)-arylcarbamoyloxy, N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-((C₁-C₁₀)-alkyl)carbamoyloxy, N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy)-(C₁-C₁₀)-alkyl)carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy)-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyloxy,

NR^YR^Z, wherein R^Y and R^Z are independently selected from (C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyl, (C₃-C₁₂)-alkenyl, (C₃-C₁₂)-alkynyl, (C₆-C₁₂)-aryl, (C₇-C₁₁)-aralkyl, (C₁-C₁₂)-alkoxy, (C₇-C₁₂)-aralkoxy,

(C₁-C₁₂)-alkanoyl, (C₃-C₈)-cycloalkanoyl, (C₆-C₁₂)-aroyl, (C₇-C₁₆)-aralkanoyl, and further wherein R^Y and R^Z together are -[CH₂]_h-, wherein one CH₂-group can be replaced by O, S, N-(C₁-C₄)-alkylcarbonylimino or N-(C₁-C₄)-alkoxycarbonylimino, and h is 3 to 6,

(C₁-C₁₂)-alkanoylamino-(C₁-C₈)-alkyl, (C₃-C₈)-cycloalkanoylamino-(C₁-C₈)-alkyl, (C₆-C₁₂)-aroylamino-(C₁-C₈)-alkyl, (C₇-C₁₆)-aralkanoylamino-(C₁-C₈)-alkyl, amino-(C₁-C₁₀)-alkyl, N-(C₁-C₁₀)-alkylamino-

- (C₁-C₁₀)-alkyl, N,N-di(C₁-C₁₀)-alkylamino-(C₁-C₁₀)-alkyl, (C₃-C₈)-cycloalkylamino-(C₁-C₁₀)-alkyl, (C₁-C₂₀)-alkylmercapto, (C₁-C₂₀)-alkylsulfinyl, (C₁-C₂₀)-alkylsulfonyl, (C₆-C₁₂)-arylmercapto, (C₆-C₁₂)-arylsulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₆)-aralkylmercapto, (C₇-C₁₆)-aralkylsulfinyl, (C₇-C₁₆)-aralkylsulfonyl, (C₁-C₁₂)-alkylmercapto-(C₁-C₆)-alkyl, (C₁-C₁₂)-alkylsulfinyl-(C₁-C₆)-alkyl, (C₁-C₁₂)-alkylsulfonyl-(C₁-C₆)-alkyl, (C₆-C₁₂)-arylmercapto-(C₁-C₆)-alkyl, (C₆-C₁₂)-arylsulfinyl-(C₁-C₆)-alkyl, (C₆-C₁₂)-arylsulfonyl-(C₁-C₆)-alkyl, (C₇-C₁₆)-aralkylmercapto-(C₁-C₆)-alkyl, (C₇-C₁₆)-aralkylsulfinyl-(C₁-C₆)-alkyl, (C₇-C₁₆)-aralkylsulfonyl-(C₁-C₆)-alkyl,
- 15 sulfamoyl, N-(C₁-C₁₀)-alkylsulfamoyl, N,N-di-(C₁-C₁₀)-alkylsulfamoyl, (C₃-C₈)-cycloalkylsulfamoyl, N-(C₆-C₁₂)-arylsulfamoyl, N-(C₇-C₁₆)-aralkylsulfamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylsulfamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylsulfamoyl,
- 20 (C₁-C₁₀)-alkyl-sulfonamido, N-((C₁-C₁₀)-alkyl)-(C₁-C₁₀)-alkylsulfonamido, (C₇-C₁₆)-aralkyl-sulfonamido, or N-(C₁-C₁₀)-alkyl-(C₇-C₁₆)-aralkyl-sulfonamido,
- 25 where the radicals which contain an aryl radical can in turn be substituted on the aryl by 1 to 5 identical or different radicals from the series comprising:
- hydroxyl, halogen, cyano, trifluoromethyl, nitro, carboxyl, (C₁-C₁₆)-alkyl, (C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkoxy, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl, (C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy, (C₁-C₆)-alkyl, (C₃-C₈)-cycloalkyloxy-(C₁-C₈)-alkoxy, (C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₈)-alkoxy, (C₁-C₈)-alkoxy, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, (C₂-C₁₆)-alkenyl, (C₂-C₁₂)-alkynyl, (C₁-C₁₆)-alkoxy,

- (C₁-C₁₆)-alkenyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl,
 (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-
 (C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy,
 (C₇-C₁₆)-aralkyloxy, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy,
 5 (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxy, (C₁-C₈)-hydroxy-
 alkyl, (C₆-C₁₆)-aryloxy-(C₁-C₈)-alkyl, (C₇-C₁₆)-
 aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-
 alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₂)-aralkyloxy-(C₁-C₈)-
 alkoxy-(C₁-C₆)-alkyl, -O-[CH₂]_xC_FH_(2f+1-g)F_g, -OCF₂Cl,
 10 -OCF₂-CHFCl,
- (C₁-C₁₂)-alkylcarbonyl, (C₃-C₈)-cycloalkylcarbonyl,
 (C₆-C₁₂)-arylcabonyl, (C₇-C₁₆)-aralkylcarbonyl,
- (C₁-C₁₂)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-
 alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-
 15 aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl,
 (C₂-C₁₂)-alkenyloxycarbonyl, (C₂-C₁₂)-alkynyloxy-
 carbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl,
 (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-
 cycloalkyl-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cyclo-
 20 alkoxy-(C₁-C₆)-alkoxycarbonyl,
- (C₁-C₁₂)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl-
 carbonyloxy, (C₆-C₁₂)-arylcabonyloxy, (C₇-C₁₆)-
 aralkylcarbonyloxy, cinnamoyloxy, (C₂-C₁₂)-alkenyl-
 carbonyloxy, (C₂-C₁₂)-alkynylcarbonyloxy,
- 25 (C₁-C₁₂)-alkoxycarbonyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-
 alkoxycarbonyloxy, (C₆-C₁₂)-aryloxycarbonyloxy,
 (C₇-C₁₆)-aralkyloxycarbonyloxy, (C₃-C₈)-cycloalkoxy-
 carbonyloxy, (C₂-C₁₂)-alkenyloxycarbonyloxy,
 (C₂-C₁₂)-alkynyloxycarbonyloxy,
- 30 carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-
 (C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-
 carbamoyl, N,N-dicyclo-(C₃-C₈)-alkylcarbamoyl,
 N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl,
 N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)carbamoyl,

- N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-
 alkyl) carbamoyl, N-(+)-dehydroabietylcarbamoyl,
 N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl,
 N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkylcarb-
 5 amoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcarbamoyl,
 N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl,
 N-((C₁-C₁₆)-alkoxy-(C₁-C₁₀)-alkyl) carbamoyl,
 N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl) carbamoyl,
 N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl) carbamoyl,
 10 N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-
 alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-
 aryloxy-(C₁-C₁₀)-alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-N-
 ((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl) carbamoyl,
 CON(CH₂)_h, in which one CH₂ group can be replaced by
 15 O, S, N-(C₁-C₈)-alkylimino, N-(C₃-C₈)-cycloalkyl-
 imino, N-(C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylimino,
 N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or
 N-(C₁-C₄)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7,

 carbamoyloxy, N-(C₁-C₁₂)-alkylcarbamoyloxy, N,N-di-
 20 (C₁-C₁₂)-alkylcarbamoyloxy, N-(C₃-C₈)-cycloalkyl-
 carbamoyloxy, N-(C₆-C₁₆)-arylcarbamoyloxy,
 N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-
 (C₆-C₁₂)-arylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-
 (C₇-C₁₆)-aralkylcarbamoyloxy, N-((C₁-C₁₀)-
 25 alkyl) carbamoyloxy, N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-
 alkyl) carbamoyloxy, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-
 alkyl) carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-
 alkoxy-(C₁-C₁₀)-alkyl) carbamoyloxy, N-(C₁-C₁₀)-
 alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl) car-
 30 bamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-
 (C₁-C₁₀)-alkyl) carbamoyloxy,

 amino, (C₁-C₁₂)-alkylamino, di-(C₁-C₁₂)-alkylamino,
 (C₃-C₈)-cycloalkylamino, (C₃-C₁₂)-alkenylamino,
 (C₃-C₁₂)-alkynylamino, N-(C₆-C₁₂)-arylamino,
 35 N-(C₇-C₁₁)-aralkylamino, N-alkyl-aralkylamino,
 N-alkyl-arylamino, (C₁-C₁₂)-alkoxyamino, (C₁-C₁₂)-
 alkoxy-N-(C₁-C₁₀)-alkylamino,

5 (C₁-C₁₂)-alkanoylamino, (C₃-C₈)-cycloalkanoylamino,
(C₆-C₁₂)-aroylamino, (C₇-C₁₆)-aralkanoylamino,
(C₁-C₁₂)-alkanoyl-N-(C₁-C₁₀)-alkylamino, (C₃-C₈)-
cycloalkanoyl-N-(C₁-C₁₀)-alkylamino, (C₆-C₁₂)-aroyl-
N-(C₁-C₁₀)-alkylamino, (C₇-C₁₁)-aralkanoyl-N-(C₁-C₁₀)-
alkylamino,

10 (C₁-C₁₂)-alkanoylamino-(C₁-C₈)-alkyl, (C₃-C₈)-cyclo-
alkanoylamino-(C₁-C₈)-alkyl, (C₆-C₁₂)-aroylamino-
(C₁-C₈)-alkyl, (C₇-C₁₆)-aralkanoylamino-(C₁-C₈)-
alkyl, amino-(C₁-C₁₀)-alkyl, N-(C₁-C₁₀)-alkylamino-
(C₁-C₁₀)-alkyl, N,N-di-(C₁-C₁₀)-alkylamino-(C₁-C₁₀)-
alkyl, (C₃-C₈)-cycloalkylamino-(C₁-C₁₀)-alkyl,

15 (C₁-C₁₂)-alkylmercapto, (C₁-C₁₂)-alkylsulfinyl,
(C₁-C₁₂)-alkylsulfonyl, (C₆-C₁₆)-arylmercapto,
(C₆-C₁₆)-arylsulfinyl, (C₆-C₁₆)-arylsulfonyl,
(C₇-C₁₆)-aralkylmercapto, (C₇-C₁₆)-aralkylsulfinyl
and (C₇-C₁₆)-aralkylsulfonyl, or optionally carries
up to 3 of said radical substituents for said aryl
and two adjacent carbon atoms of an arylalkoxy
20 radical, recited in the definition of R¹-R⁵ together
carry a chain -[CH₂]- and/or -CH=CH-CH=CH, where one
CH₂-group of the chain is optionally replaced by O,
S, SO, SO₂ or NR',

25 R¹ and R², R² and R³, R³ and R⁴, or R⁴ and R⁵ form a chain
[CH₂]_o, in which one or two CH₂ groups of the chain,
which is saturated or unsaturated with a C=C double
bond, are optionally replaced by O, S, SO, SO₂ or
NR', in which o is 3, 4 or 5, and

30 R' is hydrogen, (C₆-C₁₂)-aryl, (C₁-C₈)-alkyl, (C₁-C₈)-
alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkoxy-(C₁-C₈)-
alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkyl, (C₁-C₁₀)-
alkanoyl, optionally substituted (C₇-C₁₆)-aralkanoyl
or optionally substituted (C₆-C₁₂)-aroyl, and

m is 0 or 1,

- f is 1 to 8,
g is 0 or 1 to $(2f + 1)$,
x is 0 to 3 and
h is 3 to 6, including the physiologically active
5 salts.

Aryl is understood as meaning, in particular, phenyl and naphthyl which are unsubstituted or preferably substituted as described, heteroaryl is understood as meaning, in particular, pyridyl, picolyl or thienylmethyl which
10 are unsubstituted or substituted as described, cycloalkyl is preferably understood as meaning cyclohexyl and halogen is understood as meaning, in particular, fluorine, chlorine and bromine.

The invention furthermore relates to salts of the compounds of the formula I.
15

The salt formation with basic reagents can take place on one or two acid groups of the compounds of the formula I, i.e. on the radicals B, R^1 , R^2 , R^3 , R^4 and R^5 and/or on the acid phenolic OH group, in particular on the radicals
20 B and the phenolic OH group.

Reagents which are used are, for example, alcoholates, hydroxides, carbonates, bicarbonates, hydrogen phosphates and/or metal-organyls of the alkali metal and alkaline earth metal elements, the elements of main groups 3 and
25 4 of the periodic table and the elements of the transition metals,

amines, optionally mono- to trisubstituted by (C_1-C_8) -hydroxyalkyl, (C_1-C_4) -alkoxy- (C_1-C_8) -alkyl, phenyl, benzyl or (C_1-C_8) -alkyl, which can be mono- to trisubstituted by
30 hydroxyl or (C_1-C_4) -alkoxy,

for example Tromethane (Tris buffer), 2-aminoethanol, 3-aminopropanol, hydroxylamine, dimethylhydroxylamine, 2-methoxy-ethylamine and 3-ethoxypropylamine, and

basic amino acids and derivatives, such as amino acid esters, histidine, arginine and lysine and derivatives thereof, and

medicaments which contain a basic group, such as, for example, amiloride, verapamil and beta-blockers.

The invention furthermore relates to the compounds according to the formula I for use as medicaments.

Compounds of the formula I which are of particular importance are those in which

10 A is (C_1-C_3) -alkylene, which is optionally mono-substituted by halogen, cyano, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -hydroxyalkyl, (C_1-C_6) -alkoxy or $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$ or.

15 A is $-CHR^x-$, in which R^x is one of the substituents of the α -C atom of an α -amino acid, in particular of a naturally occurring L-amino acid or its D isomer,

B is $-CO_2H$,

20 R^1 and R^5 are identical or different and are hydrogen, (C_1-C_{12}) -alkyl, (C_1-C_{12}) -alkenyl, chlorine, fluorine, bromine, trifluoromethyl, (C_1-C_{12}) -alkylsulfonyl, (C_1-C_{12}) -alkylsulfinyl, phenylsulfonyl or phenylsulfinyl; where phenyl is optionally substituted by fluorine, chlorine or (C_1-C_5) -alkoxy, (C_1-C_{10}) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$, $(C-C)$ -alkoxy-
25 (C_1-C_6) -alkoxy or (C_3-C_8) -cycloalkoxy,

30 R^2 , R^3 and R^4 are identical or different and are hydrogen, hydroxyl, (C_1-C_{20}) -alkyl, (C_2-C_{20}) -alkenyl, (C_2-C_{20}) -alkynyl, (C_1-C_{20}) -alkoxy, (C_2-C_{20}) -alkenyl-oxy, (C_2-C_{20}) -alkynyloxy, retinyloxy, (C_1-C_{20}) -alkoxy- (C_1-C_8) -alkyl, (C_2-C_{20}) -alkenyloxy- (C_1-C_8) -alkyl, retinyloxy- (C_1-C_6) -alkyl, (C_2-C_{20}) -alkynyloxy- (C_1-C_8) -alkyl, halogen, cyano, trifluoromethyl,

- (C₁-C₁₂)-hydroxyalkyl, (C₁-C₂₀)-alkanoyl, (C₇-C₁₆)-
aralkanoyl, (C₆-C₁₂)-aroyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-
aralkyl, -O-[CH₂]_x-C_FH_(2f+1-g)F_g, (C₁-C₁₈)-alkyl-
mercapto, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkyl-
5 sulfonyl, (C₆-C₁₂)-arylmercapto, (C₆-C₁₂)-aryl-
sulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₂)-aralkyl-
mercapto, (C₇-C₁₂)-aralkylsulfinyl, (C₇-C₁₂)-aralkyl-
sulfonyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy,
carboxyl, (C₁-C₂₀)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-
10 (C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl,
(C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxy-
carbonyl, (C₂-C₂₀)-alkenyloxycarbonyl, retinyloxy-
carbonyl, (C₂-C₂₀)-alkynyloxycarbonyl, (C₃-C₈)-cyclo-
alkyl-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkoxy-
15 (C₁-C₆)-alkoxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-
alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-
alkoxycarbonyl,
- (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-
(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy-
20 (C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy, (C₃-C₈)-cyclo-
alkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-
cycloalkyloxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy,
(C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cyclo-
alkyloxy-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-
25 alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₆)-
alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-
alkoxy-(C₁-C₆)-alkyl, NR^YR^Z, substituted (C₆-C₁₂)-
aryloxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkoxy-(C₁-C₆)-
alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl,
30 (C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl,
(C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy or (C₇-C₁₁)-aralkoxy-
(C₁-C₆)-alkoxy,
- carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-
(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-
35 carbamoyl, N,N-dicyclo(C₃-C₈)-alkylcarbamoyl,
N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl,
N-(C₃-C₈)-cycloalkyl-(C₁-C₁₆)-alkyl)carbamoyl,

N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-
 alkyl)carbamoyl, N-(+)-dehydroabietylcarbamoyl,
 N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl,
 N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkyl-
 5 carbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcarbamoyl,
 N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl,
 N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl,
 N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl,
 N-((C₇-C₁₆)-aralkyloxy-(C₁-C₆)-alkyl)carbamoyl,
 10 N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-
 alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-
 aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-
 ((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyl, or
 CON(CH₂)_h, in which one CH₂ group can be replaced by
 15 O, S, N-(C₁-C₁₆)-alkylimino, N-(C₃-C₈)-cycloalkyl-
 imino, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkylimino,
 N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or
 N-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7,
 20 where an aromatic radical carries 1, 2, 3, 4 or 5
 identical or different substituents from the series
 comprising hydrogen, halogen, cyano, nitro,
 hydroxyl, trifluoromethyl, (C₁-C₁₆)-alkyl, (C₂-C₁₆)-
 alkenyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₁₆)-alkoxy,
 (C₁-C₁₆)-alkenyloxy, -O-[CH₂]_xC_fH_(2f+1-g)F_g, -OCF₂Cl,
 25 -O-CF₂-CHFCl, (C₁-C₆)-alkylmercapto, (C₁-C₆)-
 alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-
 alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-
 (C₁-C₄)-alkylcarbamoyl, N,N-di-(C₁-C₄)-
 alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy, (C₃-C₈)-
 30 cycloalkylcarbamoyl, phenyl, benzyl, phenoxy,
 benzyloxy, phenylmercapto, phenylsulfonyl,
 phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl
 and N,N-di-(C₁-C₄)-alkylsulfamoyl, or optionally
 carries up to 3 of the abovementioned identical or
 35 different substituents and two adjacent carbon atoms
 of the aralkyloxy radical recited in the definition
 of R¹-R⁵ together carry a chain -[CH₂]- and/or -
 CH=CH-CH=CH-, where one CH₂ group of the chain is
 optionally replaced by O, S, SO, SO₂ or NR',

R^Y and R^Z are identical or different and are hydrogen,
 (C₆-C₁₂)-aryl, (C₁-C₁₀)-alkyl, (C₃-C₁₀)-cycloalkyl,
 (C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkoxy-(C₁-
 5 C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkyl, (C₁-C₁₀)-
 alkanoyl, optionally substituted (C₇-C₁₆)-
 aralkanoyl, or optionally substituted (C₆-C₁₂)-
 aroyl, or

R^Y and R^Z together are $-[CH_2]_h-$, wherein one CH group can
 be replaced by O, S, N-(C₁-C₄)-alkanoylimino or
 10 N-(C₁-C₄)-alkoxycarbonylimino,

m is 0 or 1,
 f is 1 to 8,
 g is 0 or 1 to (2f + 1),
 h is 3 to 6 and
 15 x is 0 to 3, including the physiologically active
 salts.

Particularly preferred compounds of the formula I are
 those in which

m is 0,
 20 A is a -CH₂- group, which can be substituted by a
 methyl group,
 B is -CO₂H,
 R¹ is hydrogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, chlorine
 or bromine,
 25 R⁵ is hydrogen, fluorine, chlorine or methyl and
 R², R³ and R⁴ are identical or different and are hydro-
 gen, (C₁-C₁₈)-alkyl, (C₂-C₁₈)-alkenyl, (C₂-C₁₈)-
 alkynyl, phenyl, chlorine, fluorine, bromine,
 hydroxyl, trifluoromethyl, (C₁-C₁₈)-alkylsulfinyl,
 30 (C₁-C₁₈)-alkylsulfonyl, phenylsulfinyl, phenyl-
 sulfonyl, naphthylsulfinyl, naphthylsulfonyl,
 (C₁-C₁₈)-alkoxy, (C₃-C₈)-cycloalkoxy, (C₁-C₈)-alkoxy-
 (C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, phenyl-
 (C₁-C₄)-alkoxy, phenoxy, (C₁-C₁₂)-alkanoyl, phenyl-
 35 (C₁-C₄)-alkanoyl or benzoyl, where, in substituents
 with a phenyl or naphthyl ring, this optionally

carries up to 5 identical or different substituents from the series comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₆)-alkylsulfonyl, including the physiologically active salts.

Especially preferred compounds of the formula I are those in which

- m is 0,
A is a -CH₂- group,
B is -CO₂H,
R¹ and R⁵ are hydrogen and
R², R³ and R⁴ are identical or different and are hydrogen, (C₁-C₁₈)-alkyl, (C₁-C₁₈)-alkenyl, phenyl, chlorine, fluorine, bromine, trifluoromethyl, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, (C₁-C₁₈)-alkoxy, (C₃-C₈)-cycloalkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, phenyl-(C₁-C₄)-alkoxy, phenoxy, (C₁-C₁₂)-alkanoyl, phenyl-(C₁-C₄)-alkanoyl or benzoyl, where, in substituents having a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g and (C₁-C₆)-alkylsulfonyl, including the physiologically active salts.

Particularly preferred compounds of the formula I are those in which

- m is 0,
A is a -CH₂- group,
R¹ and R⁵ are hydrogen,
one of the substituents R², R³ or R⁴ is hydrogen and the other two are identical or different and are hydrogen, (C₁-C₁₆)-alkyl, fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₁₆)-alkylsulfonyl, phenylsulfonyl, (C₁-C₁₆)-

alkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1g)F_g, benzyloxy or phenoxy, where, in substituents which contain a phenyl ring, this optionally carries up to 3 substituents from the series comprising fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₄)-alkylsulfonyl, including the physiologically active salts.

Particularly preferred compounds of the formula I are those in which

- 10 m is 0,
A is a -CH₂- group,
R¹, R² and R⁵ are hydrogen and
R³ and R⁴ are identical or different and are hydrogen, (C₁-C₁₆)-alkyl, fluorine, chlorine, trifluoromethyl, (C₁-C₁₆)-alkylsulfonyl, phenylsulfonyl, naphthylsulfonyl, (C₁-C₁₆)-alkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, benzyloxy or phenoxy, where, in substituents which contain a phenyl ring, this optionally carries up to 3 substituents from the series comprising fluorine, chlorine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g and (C₁-C₄)-alkylsulfonyl, including the physiologically active salts.

Particularly preferred compounds of the formula I are furthermore those in which

- 25 m is 0,
A is a -CH₂- group,
R¹, R², R³ and R⁵ are hydrogen and
R⁴ is hydrogen, (C₁-C₁₂)-alkyl, fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₁₂)-alkylsulfonyl, phenylsulfonyl, naphthylsulfonyl, (C₁-C₁₂)-alkoxy, (C₁-C₆)-alkoxy-(C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, benzyloxy or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this is optionally monosubstituted by fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₄)-alkylsulfonyl, including the physiologically active salts.

Particularly preferred compounds of the formula I are, in addition, those in which

m is 0,

A is a $-\text{CH}_2-$ group,

5 R^1 , R^3 , R^4 and R^5 are hydrogen and

R^2 is hydrogen, $(\text{C}_1-\text{C}_{12})$ -alkyl, fluorine, chlorine, bromine, trifluoromethyl, $(\text{C}_1-\text{C}_{12})$ -alkylsulfonyl, phenylsulfonyl, naphthylsulfonyl, $(\text{C}_1-\text{C}_{12})$ -alkoxy, (C_1-C_6) -alkoxy- (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$, benzyloxy or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this is optionally monosubstituted by fluorine, chlorine, bromine, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ or (C_1-C_4) -alkylsulfonyl, including the physiologically active

15 salts.

Particularly preferred compounds of the formula I are furthermore those in which

m is 0,

A is a $-\text{CH}_2-$ group,

20 R^1 , R^2 and R^5 are hydrogen,

R^4 is hydrogen or chlorine and

R^3 is hydrogen, fluorine, chlorine, $(\text{C}_1-\text{C}_{12})$ -alkoxy, (C_1-C_4) -alkoxy- (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1g)}\text{F}_g$, phenylsulfonyl, naphthylsulfonyl or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this optionally carries up to 3 identical or different substituents, preferably one substituent, from the series comprising fluorine, chlorine, bromine, trifluoromethyl, trifluoromethoxy, (C_1-C_6) -alkyl and (C_1-C_6) -alkoxy, including the physiologically active salts.

Particularly preferred compounds are furthermore those of the formula I in which

m is 0,

35 A is a $-\text{CH}_2-$ group,

R^1 , R^2 , R^4 and R^5 are hydrogen and

R^3 is hydrogen, fluorine, chlorine, $(\text{C}_1-\text{C}_{12})$ -alkoxy,

(C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, phenylsulfonyl, naphthylsulfonyl or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this optionally carries up to 3 identical or different substituents, preferably one substituent, from the series comprising fluorine, chlorine, bromine, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-alkyl and (C₁-C₆)-alkoxy, including the physiologically active salts.

10 The invention furthermore relates to prodrugs to the compounds of the formula (I) which have the effect of inhibiting collagen biosynthesis in vivo owing to liberation of compounds of the formula I or salts thereof.

15 Finally, the invention also relates to prodrugs which have the effect of inhibiting prolyl 4-hydroxylase in vivo owing to liberation of compounds of the formula I or salts thereof.

Prodrug groupings are chemical groups which, in vivo,

- 20 - can be converted into the carboxylate group of the compounds of the formula I and/or
- can be split off from the amide N atom and/or
- can be converted into a quinoline ring.

The prodrug groups possible are known to the expert.

25 The following prodrug groupings are mentioned in particular:

for the carboxylate group ester, amide, hydroxymethyl and aldehyde groups and derivatives thereof, and for the quinoline N atom N-oxides and N-alkyl derivatives.

30 The invention relates to the use of compounds of the formula I and the physiologically tolerated salts for inhibition of collagen biosynthesis.

The invention relates to the use of compounds of the general formula I and the physiologically tolerated salts for inhibition of prolyl 4-hydroxylase.

5 The invention furthermore relates to the use of compounds of the formula I and the physiologically tolerated salts for the preparation of a pharmaceutical (medicament) against fibrotic diseases.

10 The invention furthermore relates to the use of compounds of the formula I and the physiologically tolerated salts for the preparation of a medicament against fibrotic diseases of the liver, the lungs and the skin.

Finally, the invention relates to the compounds of the formula I for use as medicaments.

15 The invention particularly relates to the compounds of the formula I for use as fibrosuppressants.

The invention furthermore relates to a process for the preparation of compounds of the formula I.

The compounds of the formula I in which

A - B is $-(CH_2)_{1-4}-CO_2H$ and

20 m is 0 are prepared by

1.i1.) reacting quinoline-2-carboxylic acids of the formula II ($R^{11} = H$) with the amino esters of the formula III to give the amide esters of the formula IV, or

25 1.i2.) reacting quinoline-2-carboxylic acid esters of the formula II ($R^{11} = \text{lower alkyl}$) under the conditions of aminolysis to give the compounds of the formula IV;



- 1.ii) liberating the compounds of the formulae I and V from the esters of the formulae IV and VI; and
- 1.iii) obtaining the compounds of the formulae I and VI from the compounds of the formulae V and IV by splitting off the hydroxyl-protective group R^{10} , and if appropriate
- 1.iv) oxidizing the compounds of the formula I, IV, V or VI to give compounds of the formulae Ia, IVa, Va or VIa.
- 10 Suitable protecting groups (PG = Protecting Groups) such as are familiar to the expert are, in particular, benzyl, p-methoxybenzyl, 3,4-dimethoxybenzyl, alkyl, methoxymethyl (MOM), methylthio, benzyloxymethyl (BOM), t-butyloxymethyl, 2-methoxyethoxymethyl (MEM) and
- 15 tetrahydropyranyl (THP).

Further protecting groups and the conditions under which they are split off (conversion of compounds of the formula V into compounds of the formula I) are described by Theodoro W. Greene, Peter G.M. Wuts, in Protective

20 Groups in Organic Synthesis, Second Edition 1991, John Wiley, Chapter 2 and 3, pages 10 to 174.

Equation 1.1. $R^{10} = \text{PG}$ (Protecting Group)

$A - B = (\text{CH}_2)_{1-4} - \text{CO}_2\text{H}$




R¹¹ = H, (C₁-C₈)-alkyl or benzyl

R¹² = H, (C₁-C₈)-alkyl or benzyl

5 If the 3-OH function in the quinoline-2-carboxylic acids of the formula II a is present without a protecting group, these compounds can be reacted directly with the amino acid esters of the formula III to give the compounds of the formula VI.

Equation 1.2., $R^{10} = H$



To prepare the quinoline N-oxides of the formula I (= Ia), the compounds of the formulae I, IV, V or VI are converted into their corresponding N-oxides of the formulae Ia, IVa, Va or VIa, and these are further treated analogously to equation 1.1.

Suitable processes for amide formation (reaction 1.ii.) are the methods of carboxyl activation and the condensation reactions known from peptide chemistry.

Reagents which can be used for activation of the carboxylic acids are the substances known to the expert, such as thionyl chloride, oxalyl chloride, pivaloyl chloride, chloroformic acid ester derivatives or N,N'-carbonyldiimidazole. The activated derivatives of the compounds of the formula II are reacted in situ, after their preparation, with the amide derivatives of the formula III.


A suitable condensing agent is, for example, the combination of N,N'-dicyclohexylcarbodiimide, 1-hydroxy-1H-benzotriazole and N-ethylmorpholine.

Suitable solvents are methylene chloride, carbon tetrachloride, butyl acetate, ethyl acetate, toluene, tetrahydrofuran, dimethoxyethane, 1,4-dioxane, acetonitrile, N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulfoxide, nitromethane and/or

pyridine.

To prepare the compounds of the formula II, appropriately substituted quinolines having a protected hydroxyl function in the 3-position and an oxidizable group G in the 2-position are subjected to oxidation reactions (equation 2.1.).

Equation 2.1., G = oxidizable group, for example CH_3 , CHO , CH_2OH or CO-CH_3



Variously substituted 3-hydroxy derivatives of the quinoline-2-carboxylic acids of the formula II are obtained in accordance with equation 2.2.

2.i) Substituted 2-nitrobenzoic acids of the formula VIII are treated by the customary methods of carboxylic acid activation and reacted with salts of acetylacetone to give the compounds of the formula IX.


2.ii) The compounds of the formula IX are cyclized with a base to give the 2-acetyl-3-hydroxyquinolines of the formulae Xa/b.

2.iii) The compounds of the formula Xa are reacted with a reagent $\text{R}^{10}\text{-X}$ to give the hydroxyl-protected compounds of the formula XI.

2.iv) The compounds of the formula XI are oxidized to

give the compounds of the formula II, preferably under the conditions of the haloform reaction.

Equation 2.2. $R^1 = H$, $R^{10} = PG$ (Protecting Group)



The preparation of 3-(2-nitrobenzoyl)acetylacetone (formula IX; $R^2-R^5 = H$) is known, J. Prakt. Chem. 1987, 329, page 1063.

The reaction of the compounds of the formula IX with aqueous potassium hydroxide solution to give the compounds of the formula Xa in which R^2 , R^4 and R^5 are hydrogen and R^3 is hydrogen, methyl or chlorine is known from G. Tennant et al., J. Chem. Soc. Chem. Comm. 1975, 782.

It has now been found that the substituents R^2 , R^3 , R^4 and R^5 can be varied according to the invention to achieve the object described.

5 It has furthermore been found that the compounds of the formula XI can be oxidized in a mild manner under the conditions of the haloform reaction (aqueous alkali metal or alkaline earth metal hydroxide/bromine) to give the compounds of the formula II (IIb).

10 2-Nitrobenzoic acids of the formula VIII are commercially obtainable or known from the literature, or can be prepared by various synthesis methods - depending on the desired substitution pattern.

The compounds of the formulae II, IIa and IIb (intermediate products) are novel, with the following exceptions:
15 Compounds of the formula II in which - if R^1 to R^5 and R^{11} are hydrogen - R^{10} is benzyl or methyl,
(or compounds of the formula IIb in which - if R^2 to R^5 are hydrogen - R^{10} is benzyl or methyl) and

20 compounds of the formula II in which - if R^1 to R^5 are hydrogen and R^{11} is ethyl - R^{10} is hydrogen, benzyl or p-toluenesulfonyl and

compounds of the formula IIa in which - if R^1 , R^2 , R^4 and R^5 are hydrogen - R^3 is hydrogen or methoxy.

25 3-Benzyloxyquinoline-2-carboxylic acid is known from J. Am. Chem. Soc., Volume 115, No. 24, 1993, pages 11624-11625.

3-Hydroxyquinoline-2-carboxylic acid and methyl 3-methoxyquinoline-2-carboxylate are described in J. Antibiot. 1987, 40 (7), pages 953-960, and the former
30 is also described in Chem. Pharm. Bull. 1967, 15 (5), pages 663-669.

3-Hydroxy-6-methoxyquinoline-2-carboxylic acid is described in J. Am. Chem. Soc., Vol. 103, No. 5, 1981, pages 1241 - 1243.

3-Methoxyquinoline-2-carboxylic acid is known from J. Am. Chem. Soc. 1960, pages 3371-3377.

Ethyl 3-hydroxyquinoline-2-carboxylate and ethyl 3-(p-toluenesulfonyloxy)quinoline-2-carboxylate are known from J. Chem. Soc. C. 1967, 4, pages 256-261.

Intermediate products of the formulae Xa and Xb are furthermore novel, with the following exceptions:
Compounds of the formula Xa in which - if R^2 , R^4 and R^5 are hydrogen - R^3 is hydrogen, methyl or chlorine. These known compounds of the formula Xa are described in J. Chem. Soc. Chem. Comm. 1975, pages 782 to 783.

The invention therefore also relates to the following intermediate products for the preparation of the compounds of the formula I:

Intermediate products of the formula II in which R^1 to R^5 have the meanings which apply to the compounds of the formula I,

R^{10} is H or an HO-protecting group (Protecting Group PG) and

R^{11} is hydrogen, (C_1-C_8) -alkyl or benzyl,


with the exception of compounds of the formula II in which R^{10} is benzyl or methyl if R^1 to R^5 and R^{11} are hydrogen, and those compounds from the formula II in which R^{10} is hydrogen, benzyl or p-toluenesulfonyl if R^1 to R^5 are hydrogen and R^{11} is ethyl.

Intermediate products of the formula Xa in which R^2 to R^5 have the meanings which apply to the compounds of the formula I,

with the exception of compounds of the formula Xa in which R^3 is hydrogen, methyl or chlorine if R^2 , R^4 and R^5 are hydrogen.

Alkoxy- and benzyloxy-substituted quinoline systems of the present invention are prepared by alkylation of phenolic OH group(s) of 2-nitrobenzoic acids or derivatives thereof in which, after alkylation, the carboxylate group is introduced by oxidation of methyl, acetyl, hydroxymethyl or aldehyde groups or obtained by hydrolysis of an ester group, cf., for example, equation 3.1.

Equation 3.1.




$X^1 = \text{Hal, tosylate or mesylate}$

$R = (C_1-C_{12})\text{-alkyl, benzyl or } (C_1-C_6)\text{-alkoxy-}(C_1-C_6)\text{alkyl}$

6-Aryloxyquinoline and 6-arylsulfonyl derivatives are accessible by nucleophilic aromatic substitution on 4-halo-2-nitrotoluenes XV and subsequent oxidation to give the corresponding 5-aryloxy (5-arylsulfonyl)-2-nitrobenzoic acids XVII, cf. equation 3.2..

Equation 3.2.



$X^2 = \text{F, Cl, OSO}_2\text{CF}_3$



$Y = \text{O, S}$

$Z = \text{O, SO}_2$

Another possibility is provided by nitration of appropriately substituted toluenes, benzoic acids, benzaldehydes

or acetophenones. This method is preferably used if the substituents present direct the nitration exclusively or predominantly into the 2 position, cf., for example, equation 3.3..

5 Equation 3.3.



The compounds of the formula XIII and XVI were reacted with an oxidizing agent, preferably with KMnO_4 in an aqueous alkaline medium, to give the 2-nitrobenzoic acid derivatives.

- 10 The compounds of the formula I are inhibitors of prolyl 4-hydroxylase. The inhibition of this enzyme was determined as described by Kaule and Günzler in *Annal. Biochem.* 184, 291 to 297 (1990).

15 The compounds of the formula I according to the invention furthermore have valuable pharmacological properties and show, in particular, an antifibrotic activity.

20 The antifibrotic action can be determined in the model of fibrosis of the liver induced by carbon tetrachloride. For this, rats are treated twice weekly with CCl_4 (1 ml/kg), dissolved in olive oil. The test substance is administered daily, if appropriate even twice daily, perorally or intraperitoneally - dissolved in a suitable tolerated solvent. The extent of the fibrosis of the liver is determined histologically and the content of

collagen in the liver is analyzed by hydroxyproline determination - as described by Kivirikko et al. (Anal. Biochem. 19, 249 et seq. (1967)). The activity of the fibrogenesis can be determined by radioimmunological determination of collagen fragments and procollagen peptides in the serum. The compounds according to the invention are active in this model in a concentration of 1 to 100 mg/kg.

The activity of the fibrogenesis can be determined by radioimmunological assay of the N-terminal propeptide of type III collagen or of the N- or C-terminal crosslinking domain of type IV collagen (7s collagen or type IV collagen NC₁) in the serum.

For this purpose, the hydroxyproline, procollagen III peptide, 7s collagen and type IV collagen NC concentrations in the liver of

- a) untreated rats (control)
 - b) rats to which carbon tetrachloride was administered (CCl₄ control)
 - c) rats to which first CCl₄ and then a compound according to the invention was administered
- were measured (this test method is described by Rouiller, C., experimental toxic injury of the liver; in The Liver, C. Rouiller, Volume 2, 5. 335 to 476, New York, Academic Press, 1964).

An activity of the compounds according to the invention can furthermore be detected in the following systems.

Inhibition of hepatic prolyl 4-hydroxylase in vivo:

This model is used for detection of acute inhibition of prolyl 4-hydroxylase in vivo. For this, the test substance and the corresponding vehicle are administered (intraperitoneally, intravenously, perorally) to rats of both sexes (healthy and with induced fibrosis of the liver) and, after administration of the substance, ¹⁴C-L-

proline (250 μ Ci/kg of body weight) is administered intraperitoneally. 14 C-L-Proline (250 μ Ci/kg of body weight) is then administered again intraperitoneally. Finally, the animals are exsanguinated under pento-
5 barbitol anesthesia and the liver is removed. The hepatic collagen was purified by pepsin digestion and fractional ammonium sulfate precipitation in accordance with published protocols (Ref. 1, 2). The purified liver collagen was hydrolyzed and the content of 14 C-hydroxyproline and
10 14 C-proline was determined by amino acid analysis by means of ion exchange chromatography. Inhibition of prolyl 4-hydroxylase can be seen from a reduction in the quotient 14 C-hydroxyproline/[14 C-hydroxyproline + 14 C-proline]. 2,2'-Dipyridyl is used as the reference
15 substance. (Ref. 1: Chojkier, M. 1986. Hepatocyte collagen production in vivo in normal rats. J. Clin. Invest. 78: 333-339 and Ref. 2: Ogata I. et al. 1991. Minor contribution of hepatocytes to collagen production in normal and early fibrotic livers. Hepatology 14: 361-367).

20 Inhibition of prolyl 4-hydroxylase in cell cultures:

The following cell types are used for testing prolyl 4-hydroxylase inhibitors in cell cultures:
Normal human dermal fibroblasts (NHDF), rat liver epithelial cells (Ref. 1) and primary fat storing cells
25 from the rat liver (Ref. 2). For this, the cells are cultured in the presence of inhibitors. At the same time, the collagen newly synthesized during this period is labeled metabolically by 4- 3 H-L-proline and 14 C-proline. The influence of the test substances on the degree of
30 hydroxylation of the collagen is then determined in accordance with the method of Chojkier et al. (Ref. 3). 2,2'-Dipyridyl is employed as the reference substance. (1.: Schrode, W., Mecke, D., Gebhard, R. 1990. Induction of glutamine synthetase in periportal hepatocytes by co-
35 cultivation with a liver epithelial cell line. Eur. J. Cell. Biol. 53: 35-41, 2. Blomhoff, R., Berg T. 1990. Isolation and cultivation of rat liver stellate cells.

Methods Enzymol. 190: 59-71 and 3.: Chojkier, M.
Peterkofsky, B. Bateman J. 1980. A new method for
determining the extent of proline hydroxylation by
measuring changes in the ration of [4-³H]:[¹⁴C] proline in
5 collagenase digests. Anal. Biochem. 108: 385-393).

The compounds of the formula I can be used as medicaments
in the form of pharmaceutical preparations which comprise
them, if appropriate with tolerated pharmaceutical
excipients. The compounds can be used as medicines, for
10 example in the form of pharmaceutical preparations, which
comprise these compounds as a mixture with a pharmaceuti-
cal, organic or inorganic excipient suitable for enteral,
percutaneous or parenteral administration, such as, for
example, water, gum arabic, gelatin, lactose, starch,
15 magnesium stearate, talc, vegetable oils, polyalkylene
glycols, petroleum jelly and the like.

For this purpose, they can be administered orally in
doses of 0.1 to 25 mg/kg/day, preferably 1 to 5 mg/kg/
day, or parenterally in doses of 0.01 to 5 mg/kg/day,
20 preferably 0.01 to 2.5 mg/kg/day, in particular 0.5 to
1.0 mg/kg/day. In severe cases, the dosage can also be
increased. However, lower doses are also sufficient in
many cases. These data relate to an adult weighing about
75 kg.

25 In the examples described below, the compounds of the
formula I according to the invention are understood as
meaning substituted quinoline-2-carboxylic acid N-
(carboxymethyl)amides (quinoline-2-carboxylic acid
glycinamides).

30 The term substituted N-((quinolin-2-yl)-carbonyl)-
glycines is used in parallel.

Example 1

3-Hydroxyquinoline-2-carboxylic acid N-(carboxymethyl)-
amide

35 a) 3-(2-Nitrobenzoyl)acetylacetone was obtained from

acetylacetone and 2-nitrobenzoyl chloride, melting point 69°C; cf. J. Prakt. Chem. 1987, 329, page 1063, 29 % yield.

5 b) 2-Acetyl-3-hydroxyquinoline was obtained from the product a) under basic conditions (KOH/water, Smiles rearrangement), melting point 105°C; cf. J. Chem. Soc. Chem. Comm. 1975, 782; 53 % yield.

10 c) 2-Acetyl-3-benzyloxyquinoline was obtained from the product b) with benzyl bromide (potash/acetone), 52 % yield.

¹H-NMR (CDCl₃): δ = 2.89 (2, 3 H), 5.25 (s, 2 H), 7.38 (m, 3 H), 7.58 (m, 5 H), 7.70 (m, 1 H), 8.08 (m, 1 H).

15 d) 3-Benzyloxyquinoline-2-carboxylic acid was obtained from the product c) with potassium hypochlorite (dioxane/water), oily crude product, 47 % yield,

¹H-NMR (CDCl₃): δ = 5.40 (s, 2 H), 7.40 (m, 3 H), 7.63 (m, 4 H), 7.75 (m, 2 H), 8.07 (m, 1 H).

20 e) 3-Benzyloxyquinoline-2-carboxylic acid N-((benzyloxycarbonyl)methyl)amide was obtained from the product d) with triethylamine/ethyl chloroformate (mixed anhydride method) and glycine benzyl ester tosylate, oily crude product, 64 % yield,

25 ¹H-NMR (CDCl₃): δ = 4.40 (d, 2 H), 5.25 (s, 2 H), 5.35 (s, 2 H), 7.10 to 7.75 (m, 14 H), 8.10 (m, 1H), 7.28 (t, 1 H).

f) The title compound was obtained by hydrogenating the product e) in methanol with Pd/C (10 %) in a duck-shaped shaking vessel, melting point 191°C (from aqueous hydrochloric acid), 40 % yield.

30 Example 2

N-((3-Hydroxy-6-methoxyquinolin-2-yl)carbonyl)glycine

a) 3-(5-Methoxy-2-nitrobenzoyl) acetylacetone

28 g of crude product was obtained from 19.75 g of 5-methoxy-2-nitrobenzoic acid with oxalyl chloride and Mg acetylacetonide analogously to Example 4c).

b) 2-Acetyl-3-hydroxy-6-methylquinoline

5 28 g (0.1 mol) of the above crude product were heated under reflux in 250 ml of 20 % strength aqueous KOH solution for 30 minutes. The mixture was worked up as under Example 4d), the crude product was chromatographed over silica gel with n-heptane/ethyl acetate (1:1) and
10 5.1 g of product was crystallized from corresponding fractions with diisopropyl ether, melting point 124-126°C.

c) 2-Acetyl-3-benzyloxy-6-methoxyquinoline, from the above compound by alkylation with benzyl bromide, melting
15 point 100-102°C (from diisopropyl ether).

d) 3-Benzyloxy-6-methoxyquinoline-2-carboxylic acid

1.2 g (4 mmol) of the above acetyl compound, dissolved in 5 ml of 1,4-dioxane, were added dropwise to a solution of 1.6 g of NaOH (40 mmol) in 8 ml of water, to which 0.6 ml
20 (12 mmol) of bromine had been added at 5-10°C. After a little diethyl ether had been added and the organic phase had been separated off, a solution of 0.4 g of sodium disulfite in 6 ml of water was added to the alkaline solution, the mixture was concentrated in vacuo, the
25 residue was taken up in 8 ml of water, the pH was brought to 2 with concentrated HCl, while cooling, and the product which had crystallized was filtered off with suction and dried, 0.83 g, melting point 272-275°C (sintering 90°C).

30 e) N-((3-Benzyloxy-6-methoxyquinolin-2-yl)carbonyl)-glycine 1-butyl ester

0.83 g (2.7 mmol) of the above carboxylic acid were

reacted with 0.82 g (2.7 mmol) of glycine butyl ester tosylate, 1.1 ml (8.1 mmol) of NEM, 0.405 g (3 mmol) of HOBT and 1.14 g (2.7 mmol) of CMC in 200 ml of anhydrous methylene chloride analogously to Example 4g). The crude
5 product was chromatographed over silica gel with ethyl acetate. 0.8 g of product was crystallized from corresponding fractions with diisopropyl ether, melting point 90-92°C.

f) N-((3-Benzylloxy-6-methoxyquinolin-2-yl)carbonyl)-
10 glycine

0.5 g (1.2 mmol) of the above ester was hydrolyzed in 1.5 N methanolic sodium hydroxide solution, 0.45 g of product, melting point 118°C (sintering at 105°C, from aqueous hydrochloric acid).

15 g) The title compound was obtained by hydrogenating 0.4 g (1.1 mmol) of the above benzyl compound in methanol/tetrahydrofuran (1:1) with Pd/C (10 %) in a duck-shaped shaking vessel. After the catalyst had been filtered off with suction and the filtrate concentrated
20 in vacuo, the residue was crystallized with diethyl ether, 0.16 g, melting point 258-260°C.

Example 3

N-((6-(1-Hexyloxy)-3-hydroxyquinolin-2-yl)carbonyl)-
glycine

25 Example 4

N-((6-(1-Butyloxy)-3-hydroxyquinolin-2-yl)carbonyl)-
glycine

a) 5-(1-Butyloxy)-2-nitrobenzyl alcohol

5.1 g (30 mmol) of 5-hydroxy-2-nitrobenzyl alcohol were
30 stirred with 2.5 g (18 mmol) of potassium carbonate and 25 ml of N,N-dimethylacetamide at 70°-80°C for 30 minutes. After cooling to 20°C, 3.9 ml (36 mmol) of

1-butylbromide were added dropwise and the mixture was stirred at 90°C for 2 hours. After cooling, the reaction mixture was concentrated in vacuo, the residue was treated with water and the pH was brought to 7 with 2 N HCl. The oil obtained crystallizes after about 30 minutes; the product was filtered off with suction, washed with water and dried; 6 g, melting point 47-49°C.

b) 5-(1-Butyloxy)-2-nitrobenzoic acid

0.5 g (10 mmol) of potassium hydroxide was dissolved in 40 ml of water, 2.2 g (10 mmol) of the above alcohol were added at 20°C, while stirring, the mixture was heated to 70-80°C and 3.16 g (20 mmol) of potassium permanganate were added in portions. After the mixture had been stirred at 60-70°C for 30 minutes, it was filtered hot with suction, the residue on the filter was washed with hot water, the filtrate was concentrated to 20 ml and half-concentrated hydrochloric acid was added down to a pH of 1-2. The oil which separates off crystallized after 1 hour. The product was filtered off with suction and washed with water, 2 g, melting point 82-84°C.

c) 3-(5-(1-Butyloxy)-2-nitrobenzoyl)acetylacetone

1. 5-(1-Butyloxy)-2-nitrobenzoyl chloride

3 drops of N,N-dimethylformamide and, dropwise, 7 ml of oxalyl chloride (in 12 ml of tetrahydrofuran) were added to 12 g (50 mmol) of the above benzoic acid in 90 ml of anhydrous tetrahydrofuran at 5°C.

The mixture was allowed to warm to room temperature and was stirred for 1 hour and concentrated in vacuo, and the residue was taken up in 25 ml of anhydrous toluene.

2. Magnesium acetylacetonide

1.23 g (50 mmol) of magnesium chips were heated under reflux with a crystal of iodine in 20 ml of anhydrous ethanol and 2.5 ml of xylene for 8 hours. The mixture was then concentrated in vacuo, the residue was taken up in

100 ml of anhydrous toluene, 5.2 ml (50 mmol) of freshly distilled acetylacetone were added dropwise at 20°C and the mixture was stirred for 30 minutes.

5 This solution of magnesium acetylacetonide in toluene was cooled to 0°C and the solution, prepared under 1., of the acid chloride was added dropwise, while stirring. The mixture was stirred at 0°C for 1 hour and left to stand overnight at 20°C, 200 ml of ice-water and concentrated hydrochloric acid to pH 1 were added, the mixture was
10 stirred for 30 minutes and the organic phase was washed with water and dried to give, after concentration in vacuo, 14 g of oily crude product.

d) 2-Acetyl-6-(1-butyloxy)-3-hydroxyquinoline

14 g (43.6 mmol) of the above product were heated under
15 reflux in 150 ml of 20 % strength aqueous KOH solution for 30 minutes. After cooling, the pH was brought to 1-2 with half-concentrated hydrochloric acid, while stirring, the mixture was extracted three times with methylene chloride, the organic phase was dried and concentrated in
20 vacuo and the residue was chromatographed over silica gel with n-heptane/ethyl acetate (3:1). 2.5 g of product were crystallized from corresponding fractions with petroleum ether, melting point 71-73°C.

e) 2-Acetyl-3-benzyloxy-6-(1-butyloxy)quinoline

25 2.5 g (about 10 mmol) of the above product were reacted with 1.38 g (10 mmol) of potassium carbonate and 1.2 ml (10 mmol) of benzyl bromide in 30 ml of N,N-dimethylacetamide (1 hour at 75-80°C). 2.2 g of product were obtained, melting point 91-93°C (from diisopropyl ether).

30 f) 3-Benzyloxy-6-(1-butyloxy)quinoline-2-carboxylic acid (haloform oxidation)

0.45 ml (9 mmol) of bromine was added to a solution of

1.2 g (30 mmol) of NaOH in 6 ml of water at 5-10°C, while stirring. A solution of 1.05 g (3 mmol) of the above acetyl compound, dissolved in 5 ml of 1,4-dioxane, was then added dropwise. The mixture was stirred at 20°C for
5 a further hour and extracted with 5 ml of diethyl ether, the organic phase was concentrated, the residue was treated with 2 N aqueous HCl, the mixture was decanted, the residue was dissolved in ethyl acetate, the solution was dried with magnesium sulfate and concentrated and the
10 residue was crystallized with diisopropyl ether. 0.8 g was obtained, melting point 108-110°C.

g) N-((3-Benzoyloxy-6-(1-butyloxy)quinolin-2-yl)-carbonyl)glycine 1-butyl ester

0.8 g (2.28 mmol) of the above quinoline-2-carboxylic
15 acid was dissolved in 200 ml of anhydrous methylene chloride, 0.74 g (2.43 mmol) of glycine 1-butyl ester tosylate, 1.1 ml (8.1 mmol) of N-ethylmorpholine (NEM), 0.405 g (3 mmol) of 1-hydroxybenzotriazole (HOBT) and 1.0 g (2.43 mmol) of N-cyclohexyl-N'-(2-morpholino-
20 ethyl)carbodiimide-methyl-p-toluenesulfonate (CMC) were added in succession and the mixture was stirred at 20°C for 30 hours.

The reaction solution was then shaken with 2 N aqueous HCl, dilute NaOH was added to a precipitate, the mixture
25 was extracted with diethyl ether, the combined organic phases were extracted with aqueous Na bicarbonate solution, dried and concentrated and the residue was chromatographed over silica gel with n-heptane/ethyl acetate (2:1). After corresponding fractions had been
30 concentrated and treated with diisopropyl ether, 0.55 g of product was obtained, melting point 99-101°C.

h) The title compound was obtained by hydrolyzing 0.5 g of the above glycine ester with 50 ml of 1 N methanolic NaOH at 20°C. The mixture was concentrated in vacuo,
35 water was added to the residue, the mixture was extracted

with diethyl ether, the aqueous, oily phase was acidified after addition of tetrahydrofuran, while cooling, and concentrated in vacuo, and the aqueous phase was extracted with methylene chloride, dried and concentrated to give 0.46 g of oily product. 0.3 g of the glycine obtained was dissolved in 75 ml of tetrahydrofuran, Pd/C (10 %) was added and the glycine was hydrogenated in a duck-shaped shaking vessel. The catalyst was filtered off with suction, the filtrate was concentrated and the residue was crystallized with diisopropyl ether. 140 mg of the title compound were obtained as a colorless crystalline substance, melting point 180-182°C.

Example 5

N-((6-Ethyloxy-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 6

N-((3-Hydroxy-6-(1-octyloxy)quinolin-2-yl)carbonyl)-glycine

The title compound was obtained analogously to Example 4:

- a) 2-Nitro-5-(1-octyloxy)benzyl alcohol
- 20 The oil obtained after treatment with 1 N hydrochloric acid slowly crystallized completely, melting point 55-57°C.
- b) 2-Nitro-5-(1-octyloxy)benzoic acid, oily crude product.
- 25 c) 3-(2-Nitro-5-(1-octyloxy)benzoyl)acetylacetone; oily product (after chromatography over silica gel with n-heptane/ethyl acetate (3:1)).
- d) 2-Acetyl-3-hydroxy-6-(1-octyloxy)quinoline, oily product, which solidifies on standing.
- 30 e) 2-Acetyl-3-benzyloxy-6-(1-octyloxy)quinoline; oily product which crystallized completely on standing.

f) 3-Benzylloxy-6-(1-octyloxy)quinoline-2-carboxylic acid; melting point 145°C (with decomposition, from aqueous hydrochloric acid/tetrahydrofuran).

g) N-((3-Benzylloxy-6-(1-octyloxy)quinolin-2-yl)-carbonyl)glycine benzyl ester; melting point 93-95°C (from petroleum ether).

h) The title compound was obtained by hydrogenating the above benzyl compound with Pd/C; melting point 153-155°C (from petroleum ether).

10 Example 7

N-((6-(1-Decyloxy)-3-hydroquinolin-2-yl)carbonyl)glycine

Example 8

N-((3-Hydroxy-6-((2,2,2-trifluoroethyl)oxy)quinolin-2-yl)carbonyl)glycine hydrochloride

15 0.5 g (1.3 mmol) of N-((7-chloro-3-hydroxy-6-(2,2,2-trifluoroethyloxy)quinolin-2-yl)carbonyl)glycine (title compound from Example 31) was dissolved in 26 ml of methanolic formic acid (4.4 % strength), and a further 26 ml of methanolic formic acid were added at 20°C. After
20 addition of 0.5 g of Pd/C (10 %), the mixture was stirred at 20°C for 1 hour, the catalyst was filtered off, the filtrate was concentrated and the residue was crystallized with diisopropyl ether. 250 mg of product were obtained, melting point > 300°C (sintering at 207°C).

25 Example 9

N-((3-Hydroxy-6-((2,2,3,3,3-pentafluoropropyl)oxy)-quinolin-2-yl)carbonyl)glycine

Example 10

30 N-((6-((2,2,3,3,4,4,4-Heptafluorobutyl)oxy)-3-hydroxy-quinolin-2-yl)carbonyl)glycine

Example 11

N-((6-Chloro-3-hydroxyquinolin-2-yl)carbonyl)glycine

a) 2-Acetyl-6-chloro-3-hydroxyquinoline, cf. J. Chem. Soc. Chem. Comm. 1975, 782.

5 b) 2-Acetyl-3-benzyloxy-6-chloroquinoline

5.0 g (22.5 mmol) of the above chloroquinoline were dissolved in 50 ml of N,N-dimethylacetamide, the solution was stirred with 3.4 g (25 mmol) of potassium carbonate at 70-80°C for 30 minutes, 3.1 ml (25 mmol) of benzyl
10 bromide were added dropwise at 20°C and the mixture was stirred at 90°C for 2 hours. After working up, the residue was chromatographed over silica gel with n-heptane/ethyl acetate (4:1) and 4.2 g of product were crystallized from corresponding fractions with di-
15 isopropyl ether, melting point 107-110°C.

c) 3-Benzyloxy-6-chloroquinoline-2-carboxylic acid

2.1 g (6.7 mmol) of the above acetyl compound, dissolved in 7 ml of 1,4-dioxane, were added dropwise to a solution of 1 ml (20 mmol) of bromine in aqueous sodium hydroxide
20 solution (from 2.7 g (6.7 mmol) in 13.5 ml of water) at 0°C. After working up, 1.2 g of product were obtained, melting point 90-93°C (sintering from 80°C, from aqueous hydrochloric acid).

d) N-((3-Benzyloxy-6-chloroquinolin-2-yl)carbonyl)-
25 glycine 1-butyl ester

The product was obtained analogously to Example 4g) from 1.1 g (3.3 mmol) of the above quinolinecarboxylic acid and the coupling reagents NEM, HOBT, CMC and glycine 1-butyl ester tosylate, 0.75 g, melting point 121-123°C
30 (from diisopropyl ether).

e) The title compound was obtained by hydrolyzing 0.3 g

(0.75 mmol) of the above glycine ester in 25 ml of 1 N methanolic sodium hydroxide solution at 20°C for 1 hour. 0.3 g of the resinous Na salt was isolated, 12.5 ml of 48 % strength aqueous HBr were added and the mixture was stirred at a bath temperature of 80-85°C for 1 hour. After the mixture had been concentrated in vacuo, the crystalline residue was extracted with tetrahydrofuran, the extract was concentrated in vacuo and the residue was crystallized with diisopropyl ether. 0.16 g of the title compound was obtained, melting point 224°C (with decomposition).

Example 12

N-((6-Bromo-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 13

N-((3-Hydroxy-6-(phenylsulfonyl)quinolin-2-yl)carbonyl)-glycine

a) 2-Nitro-5-(phenylsulfonyl)toluene

4 g (100 mmol) of NaOH were dissolved in 150 ml of anhydrous ethanol, while stirring and heating, 10.3 ml (100 mmol) of thiophenol were added dropwise at 20°C and, after 10 minutes, 15.5 g (12.2 ml; 100 mmol) of 5-fluoro-2-nitrotoluene, dissolved in 20 ml of ethanol, were added dropwise. After 30 minutes at 20°C, the mixture was heated under reflux for 30 minutes and filtered hot. After trituration, crystallization started. After addition of water, the product was filtered off with suction and dried; 19 g of 2-nitro-5-phenylthiotoluene, melting point 70-71°C. Oxidation was then carried out with m-chloroperbenzoic acid in methylene chloride; 20.5 g of product, melting point 108-110°C (from petroleum ether).

b) 2-Nitro-5-(phenylsulfonyl)benzoic acid

15 g (55 mmol) of 2-nitro-5-phenylsulfonyltoluene were suspended in 210 ml of pyridine/water (1:2), and 79 g

(500 mmol) of potassium permanganate were added in portions, in each case until decoloration occurred, under reflux. The mixture was then filtered hot with suction and the residue was rinsed three times with hot water.

5 Starting material (2.8 g) which had precipitated out in the filtrate was recovered. The second filtrate was concentrated, 200 ml of water were added, the mixture was extracted with diethyl ether and the aqueous phase was brought to pH 1 with half-concentrated aqueous HCl, while

10 cooling. The product which had precipitated out was filtered off with suction and dried; 9 g, melting point 210-212°C.

c) 3-(2-Nitro-5-(phenylsulfonyl)benzoyl)acetylacetone

9.4 g of product were obtained from 9 g (30 mmol) of the

15 above substance with oxalyl chloride and magnesium acetylacetonide analogously to Example 4c), melting point 143-145°C (sintering at 140°C, from diisopropyl ether).

d) 2-Acetyl-3-hydroxy-6-(phenylsulfonyl)quinoline

9.5 g of the above compound were introduced into 95 ml of

20 aqueous potassium hydroxide solution (20 % strength) at 20°C, while stirring, a clear solution forming after 45 minutes. After 90 minutes, the pH was brought to 1 with half-concentrated HCl, while cooling, and the crude product was filtered off with suction. This was chromatog-

25 graphed over silica gel with n-heptane/ethyl acetate (3:1). 1.5 g of product were isolated from corresponding fractions, melting point 179-180°C.

e) 2-Acetyl-3-benzyloxy-6-(phenylsulfonyl)quinoline

1.5 g (4.6 mmol) of the above compound were reacted with

30 0.63 g (4.6 mmol) of potassium carbonate and 0.55 ml (4.6 mmol) of benzyl bromide in 50 ml of diethyl ketone (heating under reflux for 3 hours). 2.0 g of product were obtained, melting point 160-162°C (from water).

f) 3-Benzyloxy-6-(phenylsulfonyl)quinoline-2-carboxylic acid

2.0 g (4.8 mmol) of the above acetyl compound were subjected to the haloform reaction analogously to Example 4 f). 1.4 g of product, which was reacted further, were obtained.

g) N-((3-Benzyloxy-6-(phenylsulfonyl)quinolin-2-yl)-carbonyl)glycine benzyl ester

1.4 g (3.3 mmol) of the above carboxylic acid were reacted with glycine benzyl ester tosylate, N-ethylmorpholine, 1-hydroxy-1H-benzotriazole and CMC analogously to Example 4 g). 1.1 g of product were obtained, melting point 165-167°C (from diethyl ether).

h) The title compound was obtained by hydrogenating 1.1 g (1.9 mmol) of the above benzyl ester in tetrahydrofuran with Pd/C (10 %) in a duck-shaped shaking vessel. 0.5 g of colorless product was obtained, melting point 196-198°C (from diisopropyl ether).

Example 14

N-((6-((4-Fluorophenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 15

N-((6-Benzyloxy-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 16

N-((6-(4-Fluorobenzyloxy)-3-hydroxyquinolin-2-yl)-carbonyl)glycine

Example 17

N-((7-Butyloxy-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 18

N-((7-Benzyloxy-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 19

N-((6-(cis-3-Hexenyl-1-oxy)-3-hydroxyquinolin-2-yl)-carbonyl)glycine

Example 20

5 **N-((6-(trans-3-Hexenyl-1-oxy)-3-hydroxyquinolin-2-yl)-carbonyl)glycine**

Example 21

N-((3-Hydroxy-6-trifluoromethoxyquinolin-2-yl)carbonyl)-glycine

10 a) **2-Nitro-5-trifluoromethoxybenzoic acid**

29.7 ml of nitric acid were added dropwise to 76.5 ml of concentrated sulfuric acid, while cooling and stirring, and 9.0 g (43.7 mmol) of 3-trifluoromethoxybenzoic acid were then added. The reaction mixture was heated at 50-
15 55°C for 1 hour and, after cooling, was poured onto ice, and the residue was filtered off with suction to give, after drying, 8.8 g of product, melting point 90-93°C (sintering at 85°C).

b) **3-(2-Nitro-5-trifluoromethoxybenzoyl)acetylacetone**

20 was obtained from 8.8 g (35 mmol) of the above 2-nitro-5-trifluoromethoxybenzoic acid analogously to Example 4c), 13 g of oily crude product.

c) **2-Acetyl-3-hydroxy-6-trifluoromethoxyquinoline**

11.5 g (34.5 mmol) of the above product were introduced
25 into 105 ml of 20 % strength aqueous potassium hydroxide solution at 20°C, while stirring, and the mixture was heated under reflux for 30 minutes. After cooling, the pH was brought to 1 with half-concentrated aqueous HCl, the precipitate was filtered off with suction and the crude
30 product (11 g) was chromatographed over silica gel with n-heptane/ethyl acetate. Corresponding fractions were

concentrated and crystallized with petroleum ether; 5.1 g, melting point 87-88°C. 0.45 g of the corresponding N-oxide was obtained as a by-product, melting point 190-191°C.

5 d) 2-Acetyl-3-benzyloxy-5-trifluoromethoxyquinoline

5.0 g (18.4 mmol) of the above compound were reacted with 2.2 ml (18.4 mmol) of benzyl bromide in diethyl ketone analogously to Example 4e); 5.6 g of product, melting point 90-92°C (from petroleum ether).

10 e) 3-Benzyloxy-5-trifluoromethoxyquinoline-2-carboxylic acid

2.9 g of product were obtained from 2.7 g (7.5 mmol) of the above compound with 1.15 ml (22.5 mmol) of bromine in sodium hydroxide solution/1,4-dioxane, melting point 286°C (with evolution of gas, from the reaction solution).

f) N-((3-Benzyloxy-5-trifluoromethoxyquinolin-2-yl)-carbonyl)glycine benzyl ester

1.35 g of product were obtained from 1.8 g (5 mmol) of the above quinoline-2-carboxylic acid, 1.7 g (5 mmol) of glycine benzyl ester tosylate, 2.7 ml (20 mmol) of NEM, 0.74 g (5.5 mmol) of HOBT and 2.1 g (5 mmol) of CMC analogously to Example 4 g), melting point 123-124°C (from diisopropyl ether).

25 g) The title compound was obtained by hydrogenating 1.35 g (2.65 mmol) of the above glycine benzyl ester in 100 ml of tetrahydrofuran with Pd/C (10 %) in a duck-shaped shaking vessel. After the catalyst had been filtered off with suction and the filtrate concentrated, 30 the residue was crystallized with petroleum ether to give 0.7 g of product, melting point 192-194°C.

Examples 22, 23 and 24 were obtained from corresponding precursors analogously to Example 21.

Example 22

5 N-((6-((Heptafluoropropyl)oxy)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 23

N-((3-Hydroxy-6-((nonafluorobutyl)oxy)quinolin-2-yl)-carbonyl)glycine

Example 24

10 N-((3-Hydroxy-6-((pentafluoroethyl)oxy)quinolin-2-yl)-carbonyl)glycine

Example 25

N-((6-(3-(Ethyloxy)-1-propyloxy)-3-hydroxyquinolin-2-yl)carbonyl)glycine

15 a) 5-(3-(Ethyloxy)-1-propyloxy)-2-nitro-benzyl alcohol

5.0 g ($\approx$ 30 mmol) of 5-hydroxy-2-nitrobenzyl alcohol were reacted with 7.1 g (about 40 mmol) of 3-ethoxy-1-propyl bromide (prepared from 8.3 g (9.2 ml, 80 mmol) of 3-ethoxy-1-propanol and 15.3 ml (160 mmol) of phosphorus tribromide in methylene chloride) analogously to Example 20 4a); 6.1 g of crude product.

b) 5-(3-(Ethyloxy)-1-propyloxy)-2-nitrobenzoic acid

8.1 g of product were obtained from 10.2 g (40 mmol) of the above alcohol analogously to Example 4b), melting 25 point 109-111° (from aqueous hydrochloric acid).

c) 3-(5-(3-(Ethyloxy)-1-propyloxy)-2-nitrobenzoyl)acetyl-
acetone

10.3 g of oily crude product were obtained from 10.2 g 30 (about 40 mmol) of the above benzoic acid analogously to

Example 4c), after chromatography over silica gel with n-heptane/ethyl acetate (1:1).

d) 2-Acetyl-6-(3-ethyloxy-1-propyloxy)-3-hydroxy-quinoline was obtained by reaction of 10 g of the above
5 compound in 100 ml of 20 % strength aqueous potassium hydroxide solution (40 minutes, 60°C), 2.8 g, melting point 62-64°C (from n-heptane/ethyl acetate (3:1)).

e) 2-Acetyl-3-benzyloxy-6-(3-ethyloxy-1-propyloxy)-quinoline

10 2.8 g (10 mmol) of the above compound were reacted with 1.4 g of potassium carbonate and 1.2 ml (10 mmol) of benzyl bromide in 80 ml of diethyl ketone, 3.8 g of oily product after chromatography over silica gel with n-heptane/ethyl acetate (1:1).

15 f) 3-Benzyloxy-6-(3-ethyloxy-1-propyloxy)quinoline-2-carboxylic acid

3.8 g (10 mmol) of the above acetyl compound were oxidized with aqueous NaOH/bromine/1,4-dioxane (haloform reaction). 3.0 g of product were obtained, melting point
20 117-119°C (from aqueous hydrochloric acid).

g) N-((3-Benzyloxy-6-(3-ethyloxy-1-propyloxy)quinolin-2-yl)carbonyl)glycine benzyl ester

2.7 g of oily product were obtained from 2.3 g (6 mmol) of the above carboxylic acid in 300 ml of anhydrous
25 methylene chloride with 2.0 g (6 mmol) of glycine benzyl ester tosylate, 3.3 ml (24 mmol) of NEM, 1.0 g (7.5 mmol) of HOBT and 2.7 g (6.3 mmol) of CMC analogously to Examples 4g) and 21f), after chromatography over silica gel with ethyl acetate.

30 h) The title compound was obtained by dissolving 2.7 g of the above dibenzyl compound in 100 ml of tetrahydro-

furan and hydrogenating it with Pd/C (10 %) in a duck-shaped shaking vessel. The catalyst was then filtered off with suction, the filtrate was concentrated in vacuo and the residue was crystallized with petroleum ether. 1.3 g of product were obtained, melting point 133-135°C.

Example 26

N-((3-Hydroxy-6-(2-propyloxy)quinolin-2-yl)carbonyl)-glycine

a) 2-Nitro-5-(2-propyloxy)benzyl alcohol

10 10 g (60 mmol) of 5-hydroxy-2-nitrobenzyl alcohol were reacted with 7.0 ml (70 mmol) of isopropyl iodide analogously to Example 4a); after purification over silica gel with n-heptane/ethyl acetate (3:2), 10.3 g of oily crude product.

15 b) 2-Nitro-5-(2-propyloxy)benzoic acid

10.3 g (about 50 mmol) of the above benzyl alcohol were oxidized analogously to Example 4b); 9.4 g of product, melting point 131-133°C (from methylene chloride).

c) 3-(5-(2-Propyloxy)-2-nitrobenzoyl)acetyl acetone

20 12.4 g of product were obtained from 9 g (40 mmol) of the above product (analogously to Example 4c)).

d) 2-Acetyl-3-hydroxy-6-(2-propyloxy)quinoline

25 12.3 g (40 mmol) of the above product were reacted in 150 ml of 20 % strength aqueous potassium hydroxide solution at 50°C for 45 minutes analogously to Example 4d). Chromatography gave 4.3 g of product, melting point 95-97°C (from n-heptane/ethyl acetate (3:1)).

e) 2-Acetyl-3-benzyloxy-6-(2-propyloxy)quinoline

4.2 g (17 mmol) of the above substance were reacted with
1.4 g (10 mmol) of potassium carbonate and 2.9 g (2.0 ml,
17 mmol) of benzyl bromide in 100 ml of diethyl ketone,
5 5.0 g of product, melting point 102-104°C (from petroleum
ether).

f) 3-Benzyloxy-6-(2-propyloxy)quinoline-2-carboxylic
acid

3.4 g (10 mmol) of the above acetyl compound, dissolved
10 in 10 ml of 1,4-dioxane, were added dropwise to a sol-
ution of 4 g of NaOH and 1.6 g (30 mmol) of bromine in
20 ml of water at 0 to 5°C. After a thick slurry had
formed, 50 ml of diethyl ether were added, the mixture
was stirred, 1 g of sodium disulfite in 15 ml of water
15 was then added, the precipitate was filtered off with
suction and dissolved in tetrahydrofuran/water, the
solution was acidified with half-concentrated aqueous
hydrochloric acid, the aqueous phase was extracted with
ethyl acetate and concentrated and the residue was
20 crystallized with diisopropyl ether. 3.0 g of product
were obtained, melting point 140°C (with decomposition).

g) N-((3-Benzyloxy-6-(2-propyloxy)quinolin-2-yl)-
carbonyl)glycine benzyl ester

2.7 g (8 mmol) of the above quinolinecarboxylic acid were
25 reacted with 2.7 g (8 mmol) of glycine benzyl ester
tosylate, 4.4 ml (32 mmol) of NEM, 1.35 g (10 mmol) of
.HOBT and 3.6 g (8.8 mmol) of CMC in 300 ml of anhydrous
methylene chloride at 20°C for 24 hours, analogously to
Example 25g). After working up and chromatography over
30 silica gel with n-heptane/ethyl acetate (3:2), 3.5 g of
oily product were obtained.

h) The title compound was obtained by hydrogenating
3.4 g (7 mmol) of the above dibenzyl compound in 100 ml

of tetrahydrofuran with Pd/C (10 %) in a duck-shaped shaking vessel (uptake of 260 ml of hydrogen). The catalyst was filtered off with suction, the filtrate was concentrated and the residue was crystallized with
5 diisopropyl ether. 1.5 g of the colorless title compound were obtained, melting point 149-151°C.

Example 27

N-((3-Hydroxy-6-phenoxyquinolin-2-yl)carbonyl)glycine

a) 2-Nitro-5-phenoxytoluene

10 14.1 g (150 mmol) of phenol were dissolved in 120 ml of anhydrous N,N-dimethylacetamide, while stirring, 20.6 g (150 mmol) of finely powdered potassium carbonate were added and the mixture was stirred at 70-80°C for 30 minutes. After cooling to 20°C, 23.25 g (18.3 ml,
15 150 mmol) of 5-fluoro-2-nitrotoluene were added dropwise, the mixture was heated at 135-140°C for 2 hours and, after cooling, concentrated in vacuo, the residue was introduced into ice and the crystalline product was filtered off with suction, washed with water and dried,
20 30 g, melting point 43-46°C.

b) 2-Nitro-5-phenoxybenzoic acid

(cf. also Liebigs Ann. 593, 113 (1955))

13 g (56.7 mmol) of the above substance were stirred under reflux with 85 g of potassium permanganate in
25 pyridine/water (1:2) for 4 hours. The manganese dioxide was filtered off hot with suction and rinsed with hot water, the filtrate was concentrated, the residue was taken up in 250 ml of saturated aqueous Na bicarbonate solution, unreacted starting material was filtered off
30 with suction and the filtrate was brought to pH 1 with half-concentrated HCl, 4.3 g, melting point 148-150°C.

c) 3-(2-Nitro-5-phenoxybenzoyl)acetylacetone

12 g of oily crude product were obtained from 15.6 g (60 mmol) of the above benzoic acid with oxalyl chloride and magnesium acetylacetonide analogously to Example 4 c).

5 d) 2-Acetyl-3-hydroxy-6-phenoxyquinoline

10 g (29.3 mmol) of the above compound were reacted in 100 ml of 20 % strength aqueous KOH solution analogously to Example 4 d). After chromatography of the crude product over silica gel with n-heptane/ethyl acetate (3:1), 2.5 g of product were obtained, melting point 131-133°C. 0.55 g of the corresponding N-oxide was obtained as a byproduct, melting point 173-175°C.

e) 2-Acetyl-3-benzyloxy-5-phenoxyquinoline

2.5 g (8.9 mmol) of the above compound were reacted with 1.2 g (8.9 mmol) of potassium carbonate and 1.1 ml (8.9 mmol) of benzyl bromide in 80 ml of diethyl ketone, 3.3 g of product, melting point 116-118°C (from water, sintering at 100°C).

f) 3-Benzyloxy-6-phenoxyquinoline-2-carboxylic acid

20 2.2 g (5.96 mmol) of the above compound were subjected to the haloform reaction analogously to Example 4 f) (12 ml of water, 2.6 g of NaOH (60 mmol), 0.9 ml (18 mmol) of bromine, 10 ml of dioxane), 2.1 g of product, melting point 281°C-283°C.

25 g) N-((3-Benzyloxy-6-phenoxyquinolin-2-yl)carbonyl)-glycine benzyl ester

1.5 g (4 mmol) of the above carboxylic acid were reacted with 1.35 g (4 mmol) of glycine benzyl ester tosylate, 2.2 ml (16 mmol) of NEM, 0.67 g (15 mmol) of HOBT and 30 1.8 g (4.2 mmol) of CMC in 250 ml of anhydrous methylene chloride analogously to Example 4 g). 1.5 g of product

were obtained, melting point 143-145°C (from diisopropyl ether).

h) The title compound was obtained by hydrogenating 1.5 g of the above glycine ester in tetrahydrofuran with Pd/C (10 %) in a duck-shaped shaking vessel. 0.7 g of the title compound was obtained, melting point 225-227°C (from diisopropyl ether).

Example 28

N-((3-Hydroxy-6-(3-methoxyphenoxy)quinolin-2-yl)-carbonyl)glycine

The title compound was obtained analogously to Example 27.

a) 5-((3-Methoxyphenyl)oxy)-2-nitrotoluene

35 g of product were obtained from 18 g (150 mmol) of 3-methoxyphenol and 23.3 g (150 mmol) of 5-fluoro-2-nitrotoluene, melting point 48-50°C (from water).

b) 5-(3-Methoxyphenoxy)-2-nitrobenzoic acid
Melting point 146-151°C (from aqueous hydrochloric acid).

c) 3-(5-((3-Methoxyphenyl)oxy)-2-nitrobenzoyl)acetylacetone
oily product after column chromatography.

d) 2-Acetyl-3-hydroxy-6-((3-methoxyphenyl)oxy)quinoline
4.9 g of product were obtained from 7.5 g (20 mmol) of the above substance after treatment with 75 ml of 20 % strength KOH (90 minutes, 55°C), melting point 102-104°C (from n-heptane/ethyl acetate (3:1)).

e) 2-Acetyl-3-benzyloxy-6-((3-methoxyphenyl)oxy)quinoline
oily product after chromatography over silica gel.

f) 3-Benzylloxy-6-((3-methoxyphenyl)oxy)quinoline-2-carboxylic acid,
Melting point 142°C (with decomposition, from diisopropyl ether).

5 g) N-((3-Benzylloxy-6-((3-methoxyphenyl)oxy)quinolin-2-yl)carbonyl)glycine benzyl ester
Melting point 118-120°C (from petroleum ether).

h) The title compound was obtained analogously to Example 27 h).

10 1.1 g of product were obtained from 2 g of the above substance, melting point 166-167°C (from diisopropyl ether).

Example 29

15 N-((3-Hydroxy-6-((3-trifluoromethylphenyl)oxy)quinolin-2-yl)carbonyl)glycine

The title compound was obtained analogously to Example 27.

a) 2-Nitro-5-((3-trifluoromethylphenyl)oxy)toluene

20 62 g of oily crude product were obtained from 36.5 g (225 mmol) of 3-hydroxybenzotrifluoride and 27.9 g (225 mmol) of 5-fluoro-2-nitrotoluene.

b) 2-Nitro-5-((3-trifluoromethylphenyl)oxy)benzoic acid, melting point 153-155°C (from aqueous hydrochloric acid).

25 c) 3-(2-Nitro-5-((3-trifluoromethylphenyl)oxy)benzoyl)-acetylacetone

15 g of oily product were obtained from 18 g (55 mmol) of the above compound after chromatography with n-heptane/ethyl acetate (4:1).

d) 2-Acetyl-3-hydroxy-6-((3-trifluoromethylphenyl)oxy)-quinoline, melting point 97-99°C (from petroleum ether).

e) 2-Acetyl-3-benzyloxy-6-((3-trifluoromethylphenyl)oxy)quinoline,

5 oily product after chromatography.

f) 3-Benzyloxy-6-((3-trifluoromethylphenyl)oxy)-quinoline-2-carboxylic acid

5.7 g (13 mmol) of the above acetyl compound were oxidized analogously to Example 4 f). 5.7 g of product were
10 obtained from the organic phase without acidification, melting point 150°C (from petroleum ether, foaming).

g) N-((3-Benzyloxy-6-((3-trifluoromethylphenyl)oxy)-quinolin-2-yl)carbonyl)glycine benzyl ester

5.6 g (12.7 mmol) of the above carboxylic acid were
15 reacted with glycine benzyl ester tosylate, NEM, HOBT and CMC analogously to Example 4 g); 3.5 g of product, melting point 103-105°C.

h) The title compound was obtained by hydrogenating
3.4 g of the above benzyl ester in tetrahydrofuran with
20 Pd/C (10 %) in a duck-shaped shaking vessel. 2.2 g of product were isolated, melting point 166-168°C (from petroleum ether).

Example 30

N-((7-Chloro-3-hydroxyquinolin-2-yl)carbonyl)glycine

25 a) 3-(4-Chloro-2-nitrobenzoyl)acetylacetone

45 g of crude product were obtained from 60.5 g (0.3 mol) of 4-chloro-2-nitrobenzoic acid analogously to Example 4 e).

b) 2-Acetyl-7-chloro-3-hydroxyquinoline

18.3 g of product were obtained from 44 g (0.15 mol) of the above compound analogously to Example 4 d), after chromatography over silica gel with n-heptane/ethyl acetate (2:1), melting point 97-99°C (from petroleum ether). 2.6 g of the corresponding N-oxide were furthermore obtained, melting point 191°C (from diisopropyl ether).

c) 2-Acetyl-3-benzyloxy-7-chloroquinoline

7.5 g (34 mmol) of the above substance were reacted with 4.7 g (34 mmol) of potassium carbonate and 4.1 ml (34 mmol) of benzyl bromide in 120 ml of diethyl ketone. Chromatography over silica gel with n-heptane/ethyl acetate (4:1) gave 8 g of product, melting point 76-77°C (from petroleum ether).

d) 3-Benzyloxy-7-chloroquinoline-2-carboxylic acid

4.1 g of product were obtained from 4.7 g (15 mmol) of the above acetyl compound after haloform oxidation (NaOH/water/bromine/dioxane), melting point 140°C (with decomposition, from water).

e) N-((3-Benzyloxy-7-chloroquinolin-2-yl)carbonyl)-glycine ethyl ester

3.5 g of product were obtained from 3.2 g (10 mmol) of the above carboxylic acid, 1.4 g (10 mmol) of glycine ethyl ester hydrochloride, 3.8 ml (30 mmol) of NEM, 1.5 g (11 mmol) of HOBT and 4.3 g (10 mmol) of CMC analogously to Example 4 g), melting point 138-140°C (from diisopropyl ether).

f) N-((3-Benzyloxy-7-chloroquinolin-2-yl)carbonyl)-glycine

2.0 g (5.0 mmol) of the above glycine ethyl ester were hydrolyzed in 150 ml of 1.5 N methanolic NaOH. 1.3 g of

product were obtained, melting point 105-108°C (sintering 70°C, foaming, from diisopropyl ether).

g) The title compound was obtained by introducing 1.2 g (3.2 mmol) of the above benzyl compound into 60 ml of 48 % strength aqueous hydrogen bromide and stirring the mixture at 80°C for 45 minutes. After cooling, the mixture was concentrated in vacuo and the residue was washed three times with 50 ml of tetrahydrofuran each time to give 0.57 g of product, melting point 268-270°C.

10 Example 31

N-((7-Chloro-3-hydroxy-6-(2,2,2-trifluoroethoxy)-quinolin-2-yl)carbonyl)glycine

a) 4,5-Dichloro-2-nitrotoluene

3,4-Dichlorotoluene was nitrated with concentrated sulfuric acid and fuming nitric acid ($d = 1.52$).

b) 4-Chloro-2-nitro-5-(2,2,2-trifluoroethoxy)toluene

40.4 g (360 mmol) of potassium tert-butyrate were added in portions to 60 ml of trifluoroethanol, while stirring and cooling, and the mixture was heated to 80-90°C. After cooling, 14.4 g (70 mmol) of 4,5-dichloro-2-nitrotoluene were added and the mixture was stirred at 105°C for 1 hour. It was concentrated in vacuo and the residue was treated with 300 ml of water to give, after filtration with suction, 12 g of crude product, which were purified over silica gel with n-heptane/ethyl acetate, 10 g of product, melting point 76-79°C.

c) 4-Chloro-2-nitro-5-(2,2,2-trifluoroethoxy)benzoic acid

10 g of the above compound were oxidized in pyridine/water (1:2) with 85 g of potassium permanganate; 8.2 g of product, melting point 153-155°C (from aqueous hydro-

chloric acid).

d) 3-(4-Chloro-2-nitro-5-(2,2,2-trifluoroethoxy)-benzoyl)acetyl acetone

10.3 g of product were obtained from 10 g (34 mmol) of the above acid analogously to Example 4c), melting point 138-140°C (from petroleum ether).

e) 2-Acetyl-7-chloro-3-hydroxy-6-(2,2,2-trifluoroethoxy)quinoline

10.2 g (26.6 mmol) of the above compound were reacted in 100 ml of 10 % strength aqueous KOH at 50-70°C. After acidification with half-concentrated HCl, 10 g of crude product were purified over silica gel with n-heptane/ethyl acetate (4:1); 4.2 g of product, melting point 104-105°C.

15 f) 2-Acetyl-3-benzyloxy-7-chloro-6-(2,2,2-trifluoroethoxy)quinoline

4.2 g (13 mmol) of the above compound were reacted with 1.8 g (13 mmol) of potassium carbonate and 1.6 ml (13 mmol) of benzyl bromide in 100 ml of diethyl ketone, 4.4 g of product, melting point 110-112°C (from water).

g) Sodium 3-benzyloxy-7-chloro-6-(2,2,2-trifluoroethoxy)quinoline-2-carboxylate

3.3 g (8 mmol) of the above acetyl compound were oxidized in aqueous NaOH/bromine/1,4-dioxane. After addition of aqueous sodium pyrosulfite (Na₂S₂O₅) solution, the product which had precipitated out was filtered off with suction to give 3.6 g of the sodium salt, melting point > 325°C.

h) N-((3-Benzyloxy-7-chloro-6-(2,2,2-trifluoroethoxy)quinolin-2-yl)carbonyl)glycine benzyl ester

3.3 g (8 mmol) of the above carboxylic acid were reacted with glycine benzyl ester tosylate analogously to Example 4 g); 2.92 g of product, which was reacted further.

5 i) N-((3-Hydroxy-7-chloro-6-(2,2,2-trifluoroethoxy)-quinolin-2-yl)carbonyl)glycine-tetrahydrofuran complex (0.5 equivalent)

10 1.12 g (2 mmol) of the above benzyl compound were dissolved in tetrahydrofuran and hydrogenated with Pd/C (10 %) in a duck-shaped shaking vessel. 0.58 g of colorless product was obtained, melting point 204-206°C (from petroleum ether), which, according to ¹H-NMR, comprises 0.5 equivalent of tetrahydrofuran.

Example 32

15 N-((7-Chloro-3-hydroxy-6-(2,2,3,3,3-pentafluoropropoxy)quinolin-2-yl)carbonyl)glycine

Example 33

N-((7-Chloro-6-(2,2,3,3,4,4,4-heptafluorobutyloxy)-3-hydroxyquinolin-2-yl)carbonyl)glycine

20 a) 4-Chloro-4-nitro-5-(2,2,3,3,4,4,4-heptafluorobutyloxy)toluene was obtained with heptafluorobutanol analogously to Example 31 b), oily crude product.

b) 4-Chloro-5-(2,2,3,3,4,4,4-heptafluorobutyloxy)-2-nitrobenzoic acid, analogously to Example 31c), melting point 115-117°C (from petroleum ether).

25 c) 3-(4-Chloro-5-(2,2,3,3,4,4,4-heptafluorobutyloxy)-2-nitrobenzoyl)acetylacetone

7.77 g of product were obtained from 7.7 g (19 mmol) of the above acid analogously to Examples 31 d) and 4 c), melting point 80-81°C (from petroleum ether).

30 d) 2-Acetyl-7-chloro-6-(2,2,3,3,4,4,4-heptafluoro-

butyloxy)-3-hydroxyquinoline was obtained from 7.7 g of the above compound analogously to Example 31 e), 2.9 g of oily product after chromatography over silica gel with n-heptane/ethyl acetate (4:1).

- 5 e) 2-Acetyl-3-benzyloxy-7-chloro-6-(2,2,3,3,4,4,4-heptafluorobutyloxy)quinoline

3.8 g of oily product were obtained from 2.9 g (6.9 mmol) of the above compound analogously to Example 31 f).

- 10 f) 3-Benzyloxy-7-chloro-6-(2,2,3,3,4,4,4-heptafluorobutyloxy)quinoline-2-carboxylic acid

3.4 g of product were obtained from 3.6 g (7 mmol) of the above acetyl compound analogously to Example 31 g), melting point 183-185°C (from petroleum ether).

- 15 g) N-((3-Benzyloxy-7-chloro-6-(2,2,3,3,4,4,4-heptafluorobutyloxy)quinolin-2-yl)carbonyl)glycine benzyl ester

- 20 h) 3.3 g (6.5 mmol) of the above carboxylic acid were reacted with glycine benzyl ester tosylate analogously to Example 4 g). After purification of the crude product over silica gel with n-heptane/ethyl acetate (2:1), 3.3 g of oily product were obtained.

- i) 1.32 g (2 mmol) of the above compound were hydrogenated analogously to Example 31 i); 0.72 g of colorless product, melting point 172-173°C (from petroleum ether).

25 Example 34

N-((3-Hydroxy-6-(2-propyl)quinolin-2-yl)carbonyl)glycine

Example 35

N-((3-Hydroxy-5-phenoxyquinolin-2-yl)carbonyl)glycine

Example 36

N-((5-(1,3-Dichlorophenoxy)-3-hydroxyquinolin-2-yl)-carbonyl)glycine

Example 37

- 5 N-((5-(1-Butyloxy)-3-hydroxyquinolin-2-yl)carbonyl)-glycine

Example 38

N-((6-(1-Butyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 39

- 10 N-((6-(1-Hexyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 40

N-((6-(1-Decyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 41

N-((3-Hydroxy-6-(1-octyl)quinolin-2-yl)carbonyl)glycine

- 15 Example 42

N-((3-Hydroxy-6-phenylquinolin-2-yl)carbonyl)glycine

Example 43

N-((3-Hydroxy-6-(2,2,3,3-tetrafluoropropoxy)quinolin-2-yl)carbonyl)glycine

- 20 a) 2-Nitro-5-(2,2,3,3-tetrafluoropropoxy)toluene
28 g of potassium carbonate, 70 ml of tetrafluoropropanol and 26.4 g of 5-fluoro-2-nitrotoluene were reacted; 45 g of oily crude product.

- 25 b) 2-Nitro-5-(2,2,3,3-tetrafluoropropoxy)benzoic acid,
Melting point 140°C (with foaming, from aqueous hydrochloric acid).

- c) 3-(2-Nitro-5-(2,2,3,3-tetrafluoropropoxy)benzoyl)-acetylacetone,

oily product.

d) 2-Acetyl-3-hydroxy-6-(2,2,3,3-tetrafluoropropoxy)-quinoline

13.3 g (35 mmol) of the above substance were stirred in
5 125 ml of 20 % strength aqueous KOH at 50°C for 45 minutes; 7.5 g of product, melting point 119-121°C (from petroleum ether).

e) 2-Acetyl-3-benzyloxy-6-(2,2,3,3-tetrafluoropropoxy)quinoline

10 Melting point 121-123°C (from petroleum ether).

f) 3-Benzyloxy-6-(2,2,3,3-tetrafluoropropoxy)-quinoline-2-carboxylic acid

Melting point 123°C (with decomposition, from petroleum ether).

15 g) N-((3-Benzyloxy-6-(2,2,3,3-tetrafluoropropoxy)-quinolin-2-yl)carbonyl)glycine benzyl ester, melting point 99-101°C (from petroleum ether).

h) The title compound was obtained by hydrogenation of the above benzyl compound, melting point 175-177°C (from
20 petroleum ether).

Example 44

N-((3-Hydroxy-6-(2,2,3,3,4,4,5,5-octafluoropentyloxy)-quinolin-2-yl)carbonyl)glycine

a) 2-Nitro-5-(2,2,3,3,4,4,5,5-octafluoropentyloxy)-toluene prepared from 5-fluoro-2-nitrotoluene and
25 octafluoro-1-pentanol, oil.

b) 2-Nitro-5-(2,2,3,3,4,4,5,5-octafluoropentyloxy)-benzoic acid

Melting point 95-96°C (from petroleum ether).

- c) 3-(2-Nitro-5-(2,2,3,3,4,4,5,5-octafluoropentyloxy)-benzoyl)acetylacetone
oily product.
- d) 2-Acetyl-3-hydroxy-6-(2,2,3,3,4,4,5,5-octafluoropentyloxy)quinoline
5 Melting point 69-71°C (from n-heptane/ethyl acetate (3:1)).
- e) 2-Acetyl-3-benzyloxy-6-(2,2,3,3,4,4,5,5-octafluoropentyloxy)quinoline
- 10 f) 3-Benzyloxy-6-(2,2,3,3,4,4,5,5-octafluoropentyloxy)-quinoline-2-carboxylic acid, resinous product.
- g) N-((3-Benzyloxy-(2,2,3,3,4,4,5,5-octafluoropentyloxy)quinolin-2-yl)carbonyl)-glycine benzyl ester, oil.
- 15 h) The title compound was obtained by hydrogenation, melting point 170-172°C (sintering 115°C, from petroleum ether).

Example 45

- 20 N-((6-((3-Chlorophenyl)oxy)-3-hydroxyquinolin-2-yl)-carbonyl)glycine

Example 46

N-((6-((3-Fluorophenyl)oxy)-3-hydroxyquinolin-2-yl)-carbonyl)glycine

Example 47

- 25 N-((3-Hydroxy-6-(3-(1-propyloxy)phenyloxy)quinolin-2-yl)carbonyl)glycine

Example 48

N-((6-(3-((Cyclohexylamino)carbonyl)phenyloxy)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 49

N-((6-((4-Chlorophenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 50

5 N-((3-Hydroxy-6-((4-(2,2,2-trifluoroethoxy)phenyl)sulfonyl)quinolin-2-yl)carbonyl)glycine

Example 51

N-((3-Hydroxy-6-((4-methoxyphenyl)sulfonyl)quinolin-2-yl)carbonyl)glycine

10 Example 52

N-((3-Hydroxy-6-((4-(1-propyloxy)phenyl)sulfonyl)quinolin-2-yl)carbonyl)glycine

Example 53

15 N-((6-((4-(1-Butyloxy)phenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 54

N-((6-((3-Chlorophenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 55

20 N-((3-Hydroxy-6-((3-methoxyphenyl)sulfonyl)quinolin-2-yl)carbonyl)glycine

Example 56

N-((3-Hydroxy-6-((3-trifluoromethylphenyl)sulfonyl)quinolin-2-yl)carbonyl)glycine

25 Example 57

N-((6-((4-Ethyloxyphenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 58

30 N-((6-((4-Bromophenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 59

N-((6-((4-(3-Ethoxypropyloxy)phenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 60

5 N-((6-((2,5-Dichlorophenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 61

N-((6-((3,4-Dichlorophenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

10 Example 62

N-((6-((3,5-Dichlorophenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 63

15 N-((6-((3,4-Dimethoxyphenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 64

N-((6-((3,5-Bis-trifluoromethylphenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 65

20 N-((3-Hydroxy-6-((4-trifluoromethoxyphenyl)sulfonyl)-quinolin-2-yl)carbonyl)glycine

Example 66

N-((3-Hydroxy-6-((4-trifluoromethylphenyl)sulfonyl)-quinolin-2-yl)carbonyl)glycine

25 Example 67

N-((3-Hydroxy-6-((4-(1-propyl)phenyl)sulfonyl)quinolin-2-yl)carbonyl)glycine

Example 68

30 N-((6-((4-(1-Butyl)phenyl)sulfonyl)-3-hydroxyquinolin-2-yl)carbonyl)glycine

Example 69

**N-((3-Hydroxy-6-(1-naphthylsulfonyl)quinolin-2-yl)-
carbonyl)glycine**

Example 70


**5 N-((3-Hydroxy-6-(2-naphthylsulfonyl)quinolin-2-yl)-
carbonyl)glycine**



HOE 95/F 224 K

Patent claims:

1. A compound of the formula (I)



in which

- 5 A is (C₁-C₄)-alkylene, which is optionally substituted by one or two substituents from the series consisting of halogen, cyano, nitro, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)Hal_g, (C₁-C₆)-alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl, phenyl, benzyl, phenoxy, benzyloxy, anilino, N-methylanilino, phenylmercapto, phenylsulfonyl, phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl and N,N-di-(C₁-C₄)-alkylsulfamoyl, or
- 20 by (C₆-C₁₂)-aryloxy, (C₇-C₁₁)-aralkyloxy, (C₆-C₁₂)-aryl or (C₇-C₁₁)-aralkyl radical, which carries, in the aryl part, 1, 2, 3, 4 or 5 identical or different substituents from the series consisting of halogen, cyano, nitro, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)Hal_g, (C₁-C₆)-alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylcarbonyl,
- 25

THIS SHEET HAS BEEN
REPLACED



* C L M S *

This Sheet Has Been Replaced



- 69 -

5 (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl and N,N-di-(C₁-C₄)-alkylsulfamoyl, or

by one or more substituents R^x of the α-C atom of an α-amino acid, being a naturally occurring L-amino acid or a D isomer thereof;

10 B is an acid grouping from the series comprising -CO₂H, -CONHCOR^{'''}, -CONHSOR^{'''}, CONHSO₂R^{'''}, -NHSO₂CF₃, tetrazolyl, imidazolyl and 3-hydroxyisoxazolyl, in which R^{'''} is aryl, heteroaryl, (C₃-C₇)-cycloalkyl or (C₁-C₄)-alkyl, optionally monosubstituted by (C₆-C₁₂)-aryl, heteroaryl, OH, SH, (C₁-C₄)-alkyl, 15 (C₁-C₄)-alkoxy, (C₁-C₄)-thioalkyl, -sulfinyl or -sulfonyl, CF₃, Cl, Br, F, I, NO₂, -COOH, (C₂-C₅)-alkoxycarbonyl, NH₂, mono- or di-(C₁-C₄-alkyl)-amino or (C₁-C₄)-perfluoroalkyl,


20 R¹, R², R³, R⁴ and R⁵ are identical or different and are hydrogen, hydroxyl, halogen, cyano, trifluoromethyl, nitro, carboxyl, (C₁-C₂₀)-alkyl, (C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkoxy, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl- 25 (C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkyloxy-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, (C₇-C₁₆)-aralkenyl, (C₇-C₁₆)-aralkynyl, (C₂-C₂₀)- 30 alkenyl, (C₂-C₂₀)-alkynyl, (C₁-C₂₀)-alkoxy, (C₂-C₂₀)-alkenyloxy, (C₂-C₂₀)-alkynyloxy, retinyloxy, (C₁-C₂₀)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy- 35 (C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy, (C₇-C₁₆)-aralkoxy-

THIS SHEET HAS BEEN
REPLACED



* C L M S *

This Sheet Has Been Replaced




- 70 -

- (C₁-C₆)-alkoxy, (C₁-C₁₆)-hydroxyalkyl, (C₆-C₁₆)-aryloxy-(C₁-C₈)-alkyl, (C₇-C₁₆)-aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₂)-aralkyloxy-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl,
- 5 (C₂-C₂₀)-alkenyloxy-(C₁-C₆)-alkyl, (C₂-C₂₀)-alkynyloxy-(C₁-C₆)-alkyl, retinyloxy-(C₁-C₆)-alkyl, -O-[CH₂]_xC_fH_(2f+1-g)Hal_g, (C₁-C₆)-chlorofluoroalkoxy,
- (C₁-C₂₀)-alkylcarbonyl, (C₃-C₈)-cycloalkylcarbonyl, (C₆-C₁₂)-arylcarbonyl, (C₇-C₁₆)-aralkylcarbonyl,
- 10 cinnamoyl, (C₂-C₂₀)-alkenylcarbonyl, (C₂-C₂₀)-alkynylcarbonyl,
- (C₂-C₂₀)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl,
- 15 (C₂-C₂₀)-alkenyloxycarbonyl, retinyloxycarbonyl, (C₂-C₂₀)-alkynyloxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxy-carbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxycarbonyl,
- 20 (C₁-C₁₂)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl-carbonyloxy, (C₆-C₁₂)-arylcarbonyloxy, (C₇-C₁₆)-aralkylcarbonyloxy, cinnamoyloxy, (C₂-C₁₂)-alkenyl-carbonyloxy, (C₂-C₁₂)-alkynylcarbonyloxy,
- (C₁-C₁₂)-alkoxycarbonyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyloxy, (C₆-C₁₂)-aryloxycarbonyloxy,
- 25 (C₇-C₁₆)-aralkyloxycarbonyloxy, (C₃-C₈)-cycloalkoxy-carbonyloxy, (C₂-C₁₂)-alkenyloxycarbonyloxy, (C₂-C₁₂)-alkynyloxycarbonyloxy,
- carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-
- 30 (C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-carbamoyl, N,N-dicyclo-(C₃-C₈)-alkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl, N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)carbamoyl, N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-

THIS SHEET HAS BEEN
REPLACED




alkyl)carbamoyl, N-(+)-dehydroabietylcarbamoyl,
 N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl,
 N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-
 aralkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-
 5 arylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-
 aralkylcarbamoyl, N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-
 alkyl)carbamoyl, N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-
 alkyl)carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-
 alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-
 10 (C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-
 ((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl,
 N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-
 alkyl)carbamoyl, CON(CH₂)_h, in which a CH₂ group can
 be replaced by O, S, N-(C₁-C₈)-alkylimino, N-(C₃-C₈)-
 15 cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl-
 imino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino
 or N-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to
 7, or a carbamoyl radical of the formula J



in which
 20 R^x is the substituent of an L- or D-α-amino acid,
 s is 1, 2, 3, 4 or 5 and
 T is OH, OR or NR^{*}R^{**}, in which

25 R^{*}, R^{**} and R^{***} are identical or different and are
 hydrogen, (C₆-C₁₂)-aryl, (C₇-C₁₁)-aralkyl,
 (C₁-C₈)-alkyl, (C₃-C₈)-cycloalkyl, (+)-dehydro-
 abietyl, (C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-
 aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-
 (C₁-C₈)-alkyl, (C₁-C₁₀)-alkanoyl, optionally
 substituted (C₇-C₁₆)-aralkanoyl or optionally



substituted (C₆-C₁₂)-aroyl, or

5 R* and R** together are -[CH₂]_h, in which one CH₂ group can be replaced by O, S, SO, SO₂, N-acylamino, N-(C₁-C₁₀)-alkoxycarbonylimino, N-(C₁-C₈)-alkylimino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or N-(C₁-C₄)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7,

10 or further wherein R¹, R², R³, R⁴ and R⁵ are identical or different and are:

carbamoyloxy, N-(C₁-C₁₂)-alkylcarbamoyloxy, N,N-di-(C₁-C₁₂)-alkylcarbamoyloxy, N-(C₃-C₈)-cycloalkylcarbamoyloxy, N-(C₆-C₁₂)-arylcarbamoyloxy, 15 N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-((C₁-C₁₀)-alkyl)carbamoyloxy, N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, 20 N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy)-(C₁-C₁₀)-alkyl)carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy)-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyloxy,

25 NR^YR^Z, wherein R^Y and R^Z are independently selected from (C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyl, (C₃-C₁₂)-alkenyl, (C₃-C₁₂)-alkynyl, (C₆-C₁₂)-aryl, (C₇-C₁₁)-aralkyl, (C₁-C₁₂)-alkoxy, (C₇-C₁₂)-aralkoxy,


30 (C₁-C₁₂)-alkanoyl, (C₃-C₈)-cycloalkanoyl, (C₆-C₁₂)-aroyl, (C₇-C₁₆)-aralkanoyl,

and further wherein R^Y and R^Z together are -[CH₂]_h-, wherein one CH₂-group can be replaced by O, S, N-(C₁-C₄)-alkylcarbonylimino or N-(C₁-C₄)-alkoxycarbonylimino, and h is 3 to 6,



* C L M S *

This Sheet Has Been Replaced



- 73 -

- (C₁-C₁₂)-alkanoylamino-(C₁-C₈)-alkyl, (C₃-C₈)-
 cycloalkanoylamino-(C₁-C₈)-alkyl, (C₆-C₁₂)-
 aroylamino-(C₁-C₈)-alkyl, (C₇-C₁₆)-aralkanoylamino-
 (C₁-C₈)-alkyl, amino-(C₁-C₁₀)-alkyl, N-(C₁-C₁₀)-
 5 alkylamino-(C₁-C₁₀)-alkyl, N,N-di(C₁-C₁₀)-alkyl-
 amino-(C₁-C₁₀)-alkyl, (C₃-C₈)-cycloalkylamino-
 (C₁-C₁₀)-alkyl, (C₁-C₂₀)-alkylmercapto, (C₁-C₂₀)-
 alkylsulfinyl, (C₁-C₂₀)-alkylsulfonyl, (C₆-C₁₂)-
 arylmercapto, (C₆-C₁₂)-arylsulfinyl, (C₆-C₁₂)-aryl-
 10 sulfonyl, (C₇-C₁₆)-aralkylmercapto, (C₇-C₁₆)-aralkyl-
 sulfinyl, (C₇-C₁₆)-aralkylsulfonyl, (C₁-C₁₂)-alkyl-
 mercapto-(C₁-C₆)-alkyl, (C₁-C₁₂)-alkylsulfinyl-
 (C₁-C₆)-alkyl, (C₁-C₁₂)-alkylsulfonyl-(C₁-C₆)-alkyl,
 (C₆-C₁₂)-arylmercapto-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryl-
 15 sulfinyl-(C₁-C₆)-alkyl, (C₆-C₁₂)-arylsulfonyl-
 (C₁-C₆)-alkyl, (C₇-C₁₆)-aralkylmercapto-(C₁-C₆)-
 alkyl, (C₇-C₁₆)-aralkylsulfinyl-(C₁-C₆)-alkyl,
 (C₇-C₁₆)-aralkylsulfonyl-(C₁-C₆)-alkyl,
 sulfamoyl, N-(C₁-C₁₀)-alkylsulfamoyl, N,N-di-
 20 (C₁-C₁₀)-alkylsulfamoyl, (C₃-C₈)-cycloalkylsulfamoyl,
 N-(C₆-C₁₂)-arylsulfamoyl, N-(C₇-C₁₆)-aralkylsulf-
 amoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylsulfamoyl,
 N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylsulfamoyl,
 (C₁-C₁₀)-alkyl-sulfonamido, N-((C₁-C₁₀)-alkyl)-
 25 (C₁-C₁₀)-alkylsulfonamido, (C₇-C₁₆)-aralkyl-
 sulfonamido, or N-(C₁-C₁₀)-alkyl-(C₇-C₁₆)-aralkyl-
 sulfonamido,
 where the radicals which contain an aryl radical can
 in turn be substituted on the aryl by 1 to 5 ident-
 30 ical or different radicals from the series
 comprising:
 hydroxyl, halogen, cyano, trifluoromethyl, nitro,
 carboxyl, (C₁-C₁₆)-alkyl, (C₃-C₈)-cycloalkyl,
 (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cyclo-
 35 alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkoxy, (C₃-C₈)-
 cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy-
 (C₁-C₁₂)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl-
 (C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy-

THIS SHEET HAS BEEN
REPLACED



* C L M S *

This Sheet Has Been Replaced



- 74 -


- 5 (C₁-C₆)-alkyl, (C₃-C₈)-cycloalkyloxy-(C₁-C₈)-alkoxy-
(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₈)-alkoxy-
(C₁-C₈)-alkoxy, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl,
(C₂-C₁₆)-alkenyl, (C₂-C₁₂)-alkynyl, (C₁-C₁₆)-alkoxy,
(C₁-C₁₆)-alkenyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl,
(C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-
(C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy,
(C₇-C₁₆)-aralkyloxy, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy,
(C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxy, (C₁-C₈)-
10 hydroxyalkyl, (C₆-C₁₆)-aryloxy-(C₁-C₈)-alkyl,
(C₇-C₁₆)-aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-
(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₂)-aralkyloxy-
(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, -O-[CH₂]_xC_FH_(2f+1-g)F_g,
-OCF₂Cl, -OCF₂-CHFCl,
- 15 (C₁-C₁₂)-alkylcarbonyl, (C₃-C₈)-cycloalkylcarbonyl,
(C₆-C₁₂)-arylcarbonyl, (C₇-C₁₆)-aralkylcarbonyl,
- 20 (C₁-C₁₂)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-
alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-
aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl,
(C₂-C₁₂)-alkenyloxycarbonyl, (C₂-C₁₂)-alkynyloxy-
carbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl,
(C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-
cycloalkyl-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cyclo-
alkoxy-(C₁-C₆)-alkoxycarbonyl,
- 25 (C₁-C₁₂)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl-
carbonyloxy, (C₆-C₁₂)-arylcarbonyloxy, (C₇-C₁₆)-
aralkylcarbonyloxy, cinnamoyloxy, (C₂-C₁₂)-alkenyl-
carbonyloxy, (C₂-C₁₂)-alkynylcarbonyloxy,
- 30 (C₁-C₁₂)-alkoxycarbonyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-
alkoxycarbonyloxy, (C₆-C₁₂)-aryloxycarbonyloxy,
(C₇-C₁₆)-aralkyloxycarbonyloxy, (C₃-C₈)-cycloalkoxy-
carbonyloxy, (C₂-C₁₂)-alkenyloxycarbonyloxy,
(C₂-C₁₂)-alkynyloxycarbonyloxy,

THIS SHEET HAS BEEN
REPLACED



* C L M S *

This Sheet Has Been Replaced



- 75 -

- carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkylcarbamoyl, N,N-dicyclo-(C₃-C₈)-alkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl, 5 N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)carbamoyl, N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)carbamoyl, N-(+)-dehydroabietylcarbamoyl, N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl, N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkylcarbamoyl, 10 N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl, N-((C₁-C₁₆)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyl, 15 N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyl, CON(CH₂)_h, in which one CH₂ group can be replaced by 20 O, S, N-(C₁-C₈)-alkylimino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or N-(C₁-C₄)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7,
- carbamoyloxy, N-(C₁-C₁₂)-alkylcarbamoyloxy, N,N-di-(C₁-C₁₂)-alkylcarbamoyloxy, N-(C₃-C₈)-cycloalkylcarbamoyloxy, N-(C₆-C₁₆)-arylcarbamoyloxy, 25 N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-((C₁-C₁₀)-alkyl)carbamoyloxy, N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, 30 N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, 35 N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyloxy,

THIS SHEET HAS BEEN
REPLACED



This Sheet Has Been Replaced



- 76 -

5 amino, (C₁-C₁₂)-alkylamino, di-(C₁-C₁₂)-alkylamino,
(C₃-C₈)-cycloalkylamino, (C₃-C₁₂)-alkenylamino,
(C₃-C₁₂)-alkynylamino, N-(C₆-C₁₂)-arylamino,
N-(C₇-C₁₁)-aralkylamino, N-alkyl-aralkylamino,
N-alkyl-arylamino, (C₁-C₁₂)-alkoxyamino, (C₁-C₁₂)-
alkoxy-N-(C₁-C₁₀)-alkylamino,

10 (C₁-C₁₂)-alkanoylamino, (C₃-C₈)-cycloalkanoylamino,
(C₆-C₁₂)-aroylamino, (C₇-C₁₆)-aralkanoylamino,
(C₁-C₁₂)-alkanoyl-N-(C₁-C₁₀)-alkylamino, (C₃-C₈)-
cycloalkanoyl-N-(C₁-C₁₀)-alkylamino, (C₆-C₁₂)-aroyl-
N-(C₁-C₁₀)-alkylamino, (C₇-C₁₁)-aralkanoyl-N-(C₁-C₁₀)-
alkylamino,

15 (C₁-C₁₂)-alkanoylamino-(C₁-C₈)-alkyl, (C₃-C₈)-cyclo-
alkanoylamino-(C₁-C₈)-alkyl, (C₆-C₁₂)-aroylamino-
(C₁-C₈)-alkyl, (C₇-C₁₆)-aralkanoylamino-(C₁-C₈)-
alkyl, amino-(C₁-C₁₀)-alkyl, N-(C₁-C₁₀)-alkylamino-
(C₁-C₁₀)-alkyl, N,N-di-(C₁-C₁₀)-alkylamino-(C₁-C₁₀)-
alkyl, (C₃-C₈)-cycloalkylamino-(C₁-C₁₀)-alkyl,

20 (C₁-C₁₂)-alkylmercapto, (C₁-C₁₂)-alkylsulfinyl,
(C₁-C₁₂)-alkylsulfonyl, (C₆-C₁₆)-arylmercapto,
(C₆-C₁₆)-arylsulfinyl, (C₆-C₁₆)-arylsulfonyl,
(C₇-C₁₆)-aralkylmercapto, (C₇-C₁₆)-aralkylsulfinyl
and (C₇-C₁₆)-aralkylsulfonyl, or optionally carries
25 up to 3 of said radical substituents for said aryl
and two adjacent carbon atoms of an arylalkoxy,
radical recited in the definition of R¹-R⁵ together
carry a chain -[CH₂]- and/or -CH=CH-CH=CH, where one
CH₂ group of the chain is optionally replaced by O,
S, SO, SO₂ or NR',


30 R¹ and R², R² and R³, R³ and R⁴, or R⁴ and R⁵ form a chain
[CH₂]_o, in which one or two CH₂ groups of the chain,
which is saturated or unsaturated with a C=C double
bond, are optionally replaced by O, S, SO, SO₂ or
NR', in which o is 3, 4 or 5, and

THIS SHEET HAS BEEN
REPLACED



* C L M S *

This Sheet Has Been Replaced



- 77 -

R' is hydrogen, (C₆-C₁₂)-aryl, (C₁-C₈)-alkyl, (C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkyl, (C₁-C₁₀)-alkanoyl, optionally substituted (C₇-C₁₆)-aralkanoyl or optionally substituted (C₆-C₁₂)-aroyl, and

m is 0 or 1,
f is 1 to 8,
g is 0 or 1 to (2f + 1),
x is 0 to 3 and
10 h is 3 to 6, or a physiologically active salt thereof.

2. A compound of the formula I as claimed in claim 1, in which

A is (C₁-C₃)-alkylene, which is optionally mono-substituted by halogen, cyano, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₆)-alkoxy or -O-[CH₂]_x-C_fH_(2f+1-g)F_g or

A is -CHR^x-, in which R^x is one of the substituents of the α-C atom of an α-amino acid, in particular of a naturally occurring L-amino acid or its D isomer,

20 B is -CO₂H,

R¹ and R⁵ are identical or different and are hydrogen, (C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkenyl, chlorine, fluorine, bromine, trifluoromethyl, (C₁-C₁₂)-alkylsulfonyl, (C₁-C₁₂)-alkylsulfinyl, phenylsulfonyl or phenylsulfinyl; where phenyl is optionally substituted by fluorine, chlorine or (C₁-C₅)-alkoxy, (C₁-C₁₀)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, (C₁-C₆)-alkoxy-(C₁-C₆)-alkoxy or (C₃-C₈)-cycloalkoxy,

30 R², R³ and R⁴ are identical or different and are hydrogen, hydroxyl, (C₁-C₂₀)-alkyl, (C₂-C₂₀)-alkenyl, (C₂-C₂₀)-alkynyl, (C₁-C₂₀)-alkoxy, (C₂-C₂₀)-alkenyl-oxy, (C₂-C₂₀)-alkynyloxy, retinyloxy, (C₁-C₂₀)-

THIS SHEET HAS BEEN
REPLACED

THIS SHEET HAS BEEN
REPLACED



- alkoxy-(C₁-C₈)-alkyl, (C₂-C₂₀)-alkenyloxy-(C₁-C₈)-alkyl, retinyloxy-(C₁-C₆)-alkyl, (C₂-C₂₀)-alkynyloxy-(C₁-C₈)-alkyl, halogen, cyano, trifluoromethyl, (C₁-C₁₂)-hydroxyalkyl, (C₁-C₂₀)-alkanoyl, (C₇-C₁₆)-aralkanoyl, (C₆-C₁₂)-aroyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, (C₁-C₁₈)-alkyl-mercapto, (C₁-C₁₈)-alkylsulfanyl, (C₁-C₁₈)-alkyl-sulfonyl, (C₆-C₁₂)-arylmercapto, (C₆-C₁₂)-aryl-sulfanyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₂)-aralkyl-mercapto, (C₇-C₁₂)-aralkylsulfanyl, (C₇-C₁₂)-aralkyl-sulfonyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy, carboxyl, (C₁-C₂₀)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkoxy-carbonyl, (C₂-C₂₀)-alkenyloxy-carbonyl, retinyloxy-carbonyl, (C₂-C₂₀)-alkynyloxy-carbonyl, (C₃-C₈)-cyclo-alkyl-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl,
- (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy, (C₃-C₈)-cyclo-alkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cyclo-alkyloxy-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, NR^xR^z, substituted (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy or (C₇-C₁₁)-aralkoxy-(C₁-C₆)-alkoxy,
- carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-



* C L H S *

This Sheet Has Been Replaced



- 79 -

carbamoyl, N,N-dicyclo(C₃-C₈)-alkylcarbamoyl,
N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl,
N-(C₃-C₈)-cycloalkyl-(C₁-C₁₆)-alkyl) carbamoyl,
N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-
5 alkyl) carbamoyl, N-(+)-dehydroabietylcarbamoyl,
N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl,
N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-
aralkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-
arylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-
10 aralkylcarbamoyl, N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-
alkyl) carbamoyl, N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-
alkyl) carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₆)-
alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-
alkoxy-(C₁-C₁₀)-alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-N-
15 ((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl) carbamoyl,
N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-
alkyl) carbamoyl, or CON(CH₂)_h, in which one CH₂
group can be replaced by O, S, N-(C₁-C₁₆)-alkyl-
imino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cyclo-
20 alkyl-(C₁-C₁₂)-alkylimino, N-(C₆-C₁₂)-arylimino,
N-(C₇-C₁₆)-aralkylimino or N-(C₁-C₁₂)-alkoxy-(C₁-C₆)-
alkylimino and h is 3 to 7, where an aromatic rad-
ical carries 1, 2, 3, 4 or 5 identical or different
substituents from the series comprising of hydrogen,
25 halogen, cyano, nitro, hydroxyl, trifluoromethyl,
(C₁-C₁₆)-alkyl, (C₂-C₁₆)-alkenyl, (C₁-C₆)-hydroxy-
alkyl, (C₁-C₁₆)-alkoxy, (C₁-C₁₆)-alkenyloxy, -O-
[CH₂]_xC_fH_(2f+1-g)F_g, -OCF₂Cl, -O-CF₂-CHFCl, (C₁-C₆)-
alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkyl-
30 sulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxy-
carbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-
di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy,
(C₃-C₈)-cycloalkylcarbamoyl, phenyl, benzyl,
phenoxy, benzyloxy, phenylmercapto, phenylsulfonyl,
35 phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl
and N,N-di-(C₁-C₄)-alkylsulfamoyl, or optionally
carries up to 3 of the abovementioned identical or
different substituents and two adjacent carbon atoms
of the aralkyloxy radical recited in the definition

THIS SHEET HAS BEEN
REPLACED



This Sheet Has Been Replaced



- 80 -

of R^1-R^5 together carry a chain $-[CH_2]-$ and/or $-CH=CH-CH=CH-$, where one CH_2 group of the chain is optionally replaced by O, S, SO, SO_2 or NR^1 ,

5 R^Y and R^Z are identical or different and are hydrogen, (C_6-C_{12}) -aryl, (C_1-C_{10}) -alkyl, (C_3-C_{10}) -cycloalkyl, (C_1-C_8) -alkoxy- (C_1-C_8) -alkyl, (C_7-C_{12}) -aralkoxy- (C_1-C_8) -alkyl, (C_6-C_{12}) -aryloxy- (C_1-C_8) -alkyl, (C_1-C_{10}) -alkanoyl, optionally substituted (C_7-C_{16}) -aralkanoyl, or optionally substituted (C_6-C_{12}) -
10 aroyl, or

R^Y and R^Z together are $-[CH_2]_h-$, wherein one CH group can be replaced by O, S, $N-(C_1-C_4)$ -alkanoylimino or $N-(C_1-C_4)$ -alkoxycarbonylimino,

m is 0 or 1,
15 f is 1 to 8,
g is 0 or 1 to $(2f + 1)$,
h is 3 to 6 and
x is 0 to 3, or a physiologically active salt thereof.

3. A compound of the formula I as claimed in claims 1
20 or 2, in which

m is 0,
A is a $-CH_2-$ group, which can be substituted by a methyl group,
B is $-CO_2H$,
25 R^1 is hydrogen, (C_1-C_4) -alkyl, (C_1-C_4) -alkoxy, chlorine or bromine,
 R^5 is hydrogen, fluorine, chlorine or methyl and
 R^2 , R^3 and R^4 are identical or different and are hydrogen,
30 (C_1-C_{18}) -alkyl, (C_2-C_{18}) -alkenyl, (C_2-C_{18}) -alkynyl, phenyl, chlorine, fluorine, bromine, hydroxyl, trifluoromethyl, (C_1-C_{18}) -alkylsulfinyl, (C_1-C_{18}) -alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, (C_1-C_{18}) -alkoxy, (C_3-C_8) -cycloalkoxy, (C_1-C_8) -alkoxy-

THIS SHEET HAS BEEN
REPLACED



- 81 -

(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_FH_(2f+1-g)F_g, phenyl-
 (C₁-C₄)-alkoxy, phenoxy, (C₁-C₁₂)-alkanoyl, phenyl-
 (C₁-C₄)-alkanoyl or benzoyl, where, in substituents
 with a phenyl or naphthyl ring, this optionally
 carries up to 5 identical or different substituents
 from the series comprising fluorine, chlorine,
 bromine, nitrile, trifluoromethyl, (C₁-C₆)-alkyl,
 (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_FH_(2f+1-g)F_g or (C₁-C₆)-
 alkylsulfonyl,
 or a physiologically active salt thereof.

4. A compound of the formula I as claimed in claims 1
 to 3, in which
 m is 0,
 A is a -CH₂- group,
 B is -CO₂H,
 R¹ and R⁵ are hydrogen and
 R², R³ and R⁴ are identical or different and are hydro-
 gen, (C₁-C₁₈)-alkyl, (C₁-C₁₈)-alkenyl, phenyl, chlor-
 ine, fluorine, bromine, trifluoromethyl, (C₁-C₁₈)-
 alkylsulfinyl, (C₁-C₁₈)-alkylsulfonyl, phenyl-
 sulfinyl, phenylsulfonyl, naphthylsulfinyl,
 naphthylsulfonyl, (C₁-C₁₈)-alkoxy, (C₃-C₈)-cyclo-
 alkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-
 C_FH_(2f+1-g)F_g, phenyl-(C₁-C₄)-alkoxy, phenoxy,
 (C₁-C₁₂)-alkanoyl, phenyl-(C₁-C₄)-alkanoyl or
 benzoyl, where, in substituents having a phenyl or
 naphthyl ring, this optionally carries up to 5
 identical or different substituents from the series
 comprising fluorine, chlorine, bromine, nitrile,
 trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy,
 -O-[CH₂]_x-C_FH_(2f+1-g)F_g and (C₁-C₆)-alkylsulfonyl, or
 a physiologically active salt thereof.


5. A compound of the formula I as claimed in claims 1
 to 4, in which
 m is 0,
 A is a -CH₂- group,
 R¹ and R⁵ are hydrogen,

 THIS SHEET HAS BEEN
 REPLACED



* C L M S *

This Sheet Has Been Replaced



- 82 -

one of the substituents R^2 , R^3 or R^4 is hydrogen and the other two are identical or different and are hydrogen, (C_1-C_{16}) -alkyl, fluorine, chlorine, bromine, trifluoromethyl, (C_1-C_{16}) -alkylsulfonyl, phenylsulfonyl, (C_1-C_{16}) -alkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$, benzyloxy or phenoxy, where, in substituents which contain a phenyl ring, this optionally carries up to 3 substituents from the series comprising fluorine, chlorine, bromine, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$ or (C_1-C_4) -alkylsulfonyl, or a physiologically active salt thereof.

6. A compound of the formula I as claimed in claims 1 to 5, in which

m is 0,

15 A is a $-CH_2-$ group,

R^1 , R^2 and R^5 are hydrogen and

R^3 and R^4 are identical or different and are hydrogen, (C_1-C_{16}) -alkyl, fluorine, chlorine, trifluoromethyl, (C_1-C_{16}) -alkylsulfonyl, phenylsulfonyl, naphthylsulfonyl,

20 (C_1-C_{16}) -alkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$, benzyloxy or phenoxy, where, in substituents which contain a phenyl ring, this optionally carries up to 3 substituents from the series comprising fluorine, chlorine, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$ and (C_1-C_4) -alkylsulfonyl, or a physiologically active salt thereof.

7. A compound of the formula I as claimed in claims 1 to 6, in which

m is 0,

30 A is a $-CH_2-$ group,

R^1 , R^2 , R^3 and R^5 are hydrogen and

R^4 is hydrogen, (C_1-C_{12}) -alkyl, fluorine, chlorine, bromine, trifluoromethyl, (C_1-C_{12}) -alkylsulfonyl, phenylsulfonyl, naphthylsulfonyl, (C_1-C_{12}) -alkoxy, (C_1-C_6) -alkoxy- (C_1-C_6) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$, benzyloxy or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this is optionally monosubstituted by

35

THIS SHEET HAS BEEN
REPLACED



fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₄)-alkylsulfonyl, or a physiologically active salt thereof.

8. A compound of the formula I as claimed in claims 1 to 6, in which

m is 0,

A is a -CH₂- group,

R¹, R³, R⁴ and R⁵ are hydrogen and

R² is hydrogen, (C₁-C₁₂)-alkyl, fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₁₂)-alkylsulfonyl, phenylsulfonyl, naphthylsulfonyl, (C₁-C₁₂)-alkoxy, (C₁-C₆)-alkoxy-(C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, benzyloxy or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this is optionally monosubstituted by fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₄)-alkylsulfonyl, or a physiologically active salt thereof.

9. A compound of the formula I as claimed in claims 1 to 6, in which

m is 0,

A is a -CH₂- group,

R¹, R² and R⁵ are hydrogen,

R⁴ is hydrogen or chlorine and

R³ is hydrogen, fluorine, chlorine, (C₁-C₁₂)-alkoxy, (C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, phenylsulfonyl, naphthylsulfonyl or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this optionally carries up to 3 identical or different substituents, preferably one substituent, from the series comprising fluorine, chlorine, bromine, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-alkyl and (C₁-C₆)-alkoxy, or a physiologically active salt thereof.

10. A compound of the formula I as claimed in claims 1 to 6 and 9, in which

m is 0,

A is a -CH₂- group,

THIS SHEET HAS BEEN
REPLACED



R^1 , R^2 , R^4 and R^5 are hydrogen and
 R^3 is hydrogen, fluorine, chlorine, (C_1-C_{12}) -alkoxy,
 (C_1-C_4) -alkoxy- (C_1-C_6) -alkoxy, $-O-[CH_2]_x-C_FH_{(2f+1-g)}F_g$
phenylsulfonyl, naphthylsulfonyl or phenoxy, where, in
5 substituents which contain a phenyl or naphthyl ring,
this optionally carries up to 3 identical or different
substituents, preferably one substituent, from the series
consisting of fluorine, chlorine, bromine, trifluoro-
methyl, trifluoromethoxy, (C_1-C_6) -alkyl and (C_1-C_6) -
10 alkoxy, or a physiologically active salt thereof.

11. A compound of the formula I as claimed in claims 1
and 2, in which

A is (C_1-C_3) -alkylene, which is optionally mono-
substituted by halogen, cyano, trifluoromethyl,
15 (C_1-C_6) -alkyl, (C_1-C_6) -hydroxyalkyl, (C_1-C_6) -alkoxy
or $-O-[CH_2]_x-C_FH_{(2f+1-g)}F_g$ or

A is $-CHR^x-$, in which R^x is one of the substituents of
the α -C atom of an α -amino acid, in particular of a
naturally occurring L-amino acid or its D isomer,

20 B is $-CO_2H$,

R^1 and R^5 are identical or different and are hydrogen,
 (C_1-C_{12}) -alkyl, (C_2-C_{12}) -alkenyl, chlorine, fluorine,
bromine, trifluoromethyl, (C_1-C_{12}) -alkylsulfonyl,
 (C_1-C_{12}) -alkylsulfinyl, phenylsulfonyl or phenyl-
25 sulfinyl; where phenyl is optionally substituted by
fluorine, chlorine or (C_1-C_5) -alkoxy, (C_1-C_{10}) -
alkoxy, $-O-[CH_2]_x-C_FH_{(2f+1-g)}F_g$, (C_1-C_6) -alkoxy-
 (C_1-C_6) -alkoxy or (C_3-C_8) -cycloalkoxy,

R^2 is hydroxyl, (C_1-C_{20}) -alkyl, (C_2-C_{20}) -alkenyl,
30 (C_2-C_{20}) -alkynyl, (C_1-C_{20}) -alkoxy, (C_2-C_{20}) -alkenyl-
oxy, (C_2-C_{20}) -alkynyloxy, retinyloxy, (C_1-C_{20}) -
alkoxy- (C_1-C_8) -alkyl, (C_2-C_{20}) -alkenyloxy- (C_1-C_8) -
alkyl, retinyloxy- (C_1-C_6) -alkyl, (C_2-C_{20}) -alkynyloxy-

THIS SHEET HAS BEEN
REPLACED



* C L H S *

This Sheet Has Been Replaced



- 85 -

- (C₁-C₈)-alkyl, halogen, cyano, trifluoromethyl,
(C₁-C₁₂)-hydroxyalkyl, (C₁-C₂₀)-alkanoyl, (C₇-C₁₆)-
aralkanoyl, (C₆-C₁₂)-aroyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-
aralkyl, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, (C₁-C₁₈)-alkyl-
5 mercapto, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkyl-
sulfonyl, (C₆-C₁₂)-arylmecapto, (C₆-C₁₂)-aryl-
sulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₂)-aralkyl-
mercapto, (C₇-C₁₂)-aralkylsulfinyl, (C₇-C₁₂)-aralkyl-
sulfonyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy,
10 carboxyl, (C₁-C₂₀)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-
(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl,
(C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxy-
carbonyl, (C₂-C₂₀)-alkenyloxycarbonyl, retinyloxy-
carbonyl, (C₂-C₂₀)-alkynyloxycarbonyl, (C₃-C₈)-cyclo-
15 alkyl-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkoxy-
(C₁-C₆)-alkoxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-
alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-
alkoxycarbonyl,

(C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-
20 (C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy-
(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy, (C₃-C₈)-cyclo-
alkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-
cycloalkyloxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy,
(C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cyclo-
25 alkyloxy-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-
alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₆)-
alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-
alkoxy-(C₁-C₆)-alkyl, NR^yR^z, substituted (C₆-C₁₂)-
aryloxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkoxy-(C₁-C₆)-
30 alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl,
(C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl,
(C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy or (C₇-C₁₁)-aralkoxy-
(C₁-C₆)-alkoxy,


carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-
35 (C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-
carbamoyl, N,N-dicyclo(C₃-C₈)-alkylcarbamoyl,
N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl,

THIS SHEET HAS BEEN
REPLACED



* C L H S *

This Sheet Has Been Replaced



- 86 -


N-(C₃-C₈)-cycloalkyl-(C₁-C₁₆)-alkyl carbamoyl,
N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-
alkyl) carbamoyl, N-(+)-dehydroabietyl carbamoyl,
N-(C₁-C₆)-alkyl-N-(+)-dehydroabietyl carbamoyl,
5 N-(C₆-C₁₂)-aryl carbamoyl, N-(C₇-C₁₆)-aralkyl-
carbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-aryl carbamoyl,
N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkyl carbamoyl,
N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-alkyl) carbamoyl,
N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl) carbamoyl,
10 N-((C₇-C₁₆)-aralkyloxy-(C₁-C₆)-alkyl) carbamoyl,
N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-
alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-
aryloxy-(C₁-C₁₀)-alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-
N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl) carbamoyl, or
15 CON(CH₂)_h, in which one CH₂ group can be replaced by
O, S, N-(C₁-C₁₆)-alkylimino, N-(C₃-C₈)-cycloalkyl-
imino, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkylimino,
N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or
N-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7,
20 where an aromatic radical carries 1, 2, 3, 4 or 5
identical or different substituents from the series
comprising hydrogen, halogen, cyano, nitro,
hydroxyl, trifluoromethyl, (C₁-C₁₆)-alkyl, (C₂-C₁₆)-
alkenyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₁₆)-alkoxy,
25 (C₁-C₁₆)-alkenyloxy, -O-[CH₂]_xC_fH_(2f+1-g)F_g, -OCF₂Cl,
-O-CF₂-CHFCl, (C₁-C₆)-alkylmercapto, (C₁-C₆)-
alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C-C₁)-₆
alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-
(C₁-C₄)-alkyl carbamoyl, N,N-di-(C₁-C₄)-
30 alkyl carbamoyl, (C₁-C₆)-alkylcarbonyloxy, (C₃-C₈)-
cycloalkyl carbamoyl, phenyl, benzyl, phenoxy,
benzyloxy, phenylmercapto, phenylsulfonyl,
phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl
and N,N-di-(C₁-C₄)-alkylsulfamoyl, or optionally
35 carries up to 3 of the abovementioned identical or
different substituents and two adjacent carbon atoms
of the aralkyloxy radical together carry a chain
-[CH₂]- and/or -CH=CH-CH=CH-, where one CH₂ group of

THIS SHEET HAS BEEN
REPLACED



* C L M S *

This Sheet Has Been Replaced



- 87 -

the chain is optionally replaced by O, S, SO, SO₂ or NR',

- R³ is hydroxyl, (C₁-C₂₀)-alkyl, (C₂-C₂₀)-alkenyl, (C₂-C₂₀)-alkynyl, (C₁₁-C₂₀)-alkoxy, (C₇-C₂₀)-alkenyl-oxy, (C₂-C₂₀)-alkynyloxy, retinyloxy, (C₁-C₂₀)-alkoxy-(C₁-C₈)-alkyl, (C₂-C₂₀)-alkenyloxy-(C₁-C₈)-alkyl, retinyloxy-(C₁-C₆)-alkyl-(C₂-C₂₀)-alkynyloxy-(C₁-C₈)-alkyl, fluorine, cyano, trifluoromethyl, (C₁-C₁₂)-hydroxyalkyl, (C₁-C₂₀)-alkanoyl, (C₇-C₁₆)-aralkanoyl, (C₆-C₁₂)-aroyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, (C₁-C₁₈)-alkylmercapto, (C₁-C₁₈)-alkyl-sulfinyl, (C₁-C₁₈)-alkylsulfonyl, (C₆-C₁₂)-aryl-mercapto, (C₆-C₁₂)-arylsulfinyl, (C₇-C₁₂)-aralkyl-mercapto, (C₇-C₁₂)-aralkylsulfinyl, (C₇-C₁₂)-aralkyl-sulfonyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy, apart from benzyloxy, carboxyl, (C₁-C₂₀)-alkoxy-carbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl, (C₂-C₂₀)-alkenyloxycarbonyl, retinyloxycarbonyl, (C₂-C₂₀)-alkynyloxycarbonyl, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyloxycarbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl,
- O-[CH₂]_x-C_fH_(2f+1-g)Hal_g apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy, (C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, NR^YR^Z, substituted (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-

THIS SHEET HAS BEEN REPLACED




- aralkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy or (C₇-C₁₁)-aralkoxy-(C₁-C₆)-alkoxy,
- 5 carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkylcarbamoyl, N,N-dicyclo(C₃-C₈)-alkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₆)-alkyl)carbamoyl,
- 10 N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)carbamoyl, N-(+)-dehydroabietylcarbamoyl, N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl, N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcarbamoyl,
- 15 N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl, N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₆)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl,
- 20 N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyl, or CON(CH₂)_h, in which one CH₂ group can be replaced by O, S, N-(C₁-C₁₆)-alkylimino,
- 25 N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or N-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7, where an aromatic radical carries 1, 2, 3, 4 or 5 identical or different
- 30 substituents from the series comprising hydrogen, halogen, cyano, nitro, hydroxyl, trifluoromethyl, (C₁-C₁₆)-alkyl, (C₂-C₁₆)-alkenyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₁₆)-alkoxy, (C₁-C₁₆)-alkenyloxy, -O-[CH₂]_xC_FH_(2f+1-g)F_g, -OCF₂Cl, -O-CF₂-CHFCl, (C₁-C₆)-alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkylsulfonyl,
- 35 (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy,

THIS SHEET HAS BEEN
REPLACED



* C L H S *

This Sheet Has Been Replaced



- 89 -

(C₃-C₈)-cycloalkylcarbamoyl, phenyl, benzyl, phenoxy, benzyloxy, phenylmercapto, phenylsulfonyl, phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl and N,N-di-(C₁-C₄)-alkylsulfamoyl, or optionally carries up to 3 of the abovementioned identical or different substituents and two adjacent carbon atoms of the aralkyloxy radical together carry a chain -[CH₂]- and/or -CH=CH-CH=CH-, where one CH₂ group of the chain is optionally replaced by O, S, SO, SO₂ or NR', and

R⁴ is hydroxyl, (C₁-C₂₀)-alkyl, (C₂-C₂₀)-alkenyl, (C₂-C₂₀)-alkynyl, (C₁-C₂₀)-alkoxy apart from 1-butoxy, (C₂-C₂₀)-alkenyloxy, (C₂-C₂₀)-alkynyloxy, retinyloxy, (C₁-C₂₀)-alkoxy-(C₁-C₆)-alkyl, (C₂-C₂₀)-alkenyloxy-(C₁-C₈)-alkyl, retinyloxy-(C₁-C₆)-alkyl, (C₂-C₂₀)-alkynyloxy-(C₁-C₈)-alkyl, halogen, cyano, trifluoromethyl, (C₁-C₁₂)-hydroxyalkyl, (C₁-C₂₀)-alkanoyl, (C₇-C₁₆)-aralkanoyl, (C₆-C₁₂)-aroyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, (C₁-C₁₈)-alkylmercapto, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkylsulfonyl, (C₆-C₁₂)-arylmercapto, (C₆-C₁₂)-arylsulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₂)-aralkylmercapto, (C₇-C₁₂)-aralkylsulfinyl, (C₇-C₁₂)-aralkylsulfonyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy apart from benzyloxy, carboxyl, (C₁-C₂₀)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl, (C₂-C₂₀)-alkenyloxycarbonyl, retinyloxycarbonyl, (C₂-C₂₀)-alkynyloxycarbonyl, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy, (C₃-C₈)-cyclo-

THIS SHEET HAS BEEN
REPLACED



- alkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, NR^YR^Z, substituted (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy or (C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy,
- carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkylcarbamoyl, N,N-dicyclo(C₃-C₈)-alkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₆)-alkylcarbamoyl, N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)carbamoyl, N-(+)-dehydroabietylcarbamoyl, N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl, N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl, N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₆)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyl, or CON(CH₂)_h, in which one CH₂ group can be replaced by O, S, N-(C₁-C₁₆)-alkylimino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or N-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7, where an aromatic radical carries 1, 2, 3, 4 or 5 identical or different substituents from the series comprising hydrogen,

THIS SHEET HAS BEEN
REPLACED



* C L M S *

- 91 -

This Sheet Has Been Replaced



5 halogen, cyano, nitro, hydroxyl, trifluoromethyl,
(C₁-C₁₆)-alkyl, (C₂-C₁₆)-alkenyl, (C₁-C₆)-hydroxy-
alkyl, (C₁-C₁₆)-alkoxy, (C₁-C₁₆)-alkenyloxy, -O-
[CH₂]_xC_fH_(2f+1-g)F_g, -O-CF₂Cl, -OCF₂-CHFCl, (C₁-C₆)-
10 alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkyl-
sulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxy-
carbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-
di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy,
15 (C₃-C₈)-cycloalkylcarbamoyl, phenyl, benzyl,
phenoxy, benzyloxy, phenylmercapto, phenylsulfonyl,
phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl
and N,N-di-(C₁-C₄)-alkylsulfamoyl, or optionally
carries up to 3 of the abovementioned identical or
different substituents and two adjacent carbon atoms
of the aralkyloxy radical together carry a chain
- [CH₂]- and/or -CH=CH-CH=CH-, where one CH₂ group of
the chain is optionally replaced by O, S, SO, SO₂ or
NR',

20 R^y and R^z are identical or different and are hydrogen,
(C₆-C₁₂)-aryl, (C₁-C₁₀)-alkyl, (C₃-C₁₀)-cycloalkyl,
(C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkoxy-(C₁-
C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkyl, (C₁-C₁₀)-
alkanoyl, optionally substituted (C₇-C₁₆)-
25 aralkanoyl, or optionally substituted (C₆-C₁₂)-
aroyl, or

R^y and R^z together are -[CH₂]_h-, wherein one CH₂ group can
be replaced by O, S, N-(C₁-C₄)-alkanoylimino or
N-(C₁-C₄)-alkoxycarbonylimino,

30 m is 0 or 1,
f is 1 to 8,
g is 0 or 1 to (2f + 1),
h is 3 to 6 and
x is 0 to 3, or a physiologically active salt thereof.


12. A compound of the formula I as claimed in claims 3
35 and 11, in which

THIS SHEET HAS BEEN
REPLACED



* C L H S *


This Sheet Has Been Replaced



- 92 -

- m is 0,
- A is a $-\text{CH}_2-$ group, which can be substituted by a methyl group,
- B is $-\text{CO}_2\text{H}$,
- 5 R^1 is hydrogen, (C_1-C_4) -alkyl, (C_1-C_4) -alkoxy, chlorine or bromine,
- R^5 is hydrogen, fluorine, chlorine or methyl,
- R^2 is $(\text{C}_1-\text{C}_{18})$ -alkyl, $(\text{C}_2-\text{C}_{18})$ -alkenyl, $(\text{C}_2-\text{C}_{18})$ -alkenyloxy, $(\text{C}_2-\text{C}_{18})$ -alkynyl, phenyl, chlorine, 10 fluorine, bromine, hydroxyl, trifluoromethyl, $(\text{C}_1-\text{C}_{18})$ -alkylsulfinyl, $(\text{C}_1-\text{C}_{18})$ -alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, $(\text{C}_1-\text{C}_{18})$ -alkoxy, (C_3-C_8) -cycloalkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy, 15 $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$, phenyl- (C_1-C_4) -alkoxy, phenoxy, $(\text{C}_1-\text{C}_{12})$ -alkanoyl, phenyl- (C_1-C_4) -alkanoyl or benzoyl, where, in substituents with a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series 20 comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ or (C_1-C_6) -alkylsulfonyl, and
- R^3 is $(\text{C}_1-\text{C}_{18})$ -alkyl, $(\text{C}_2-\text{C}_{18})$ -alkenyl, $(\text{C}_7-\text{C}_{18})$ -alkenyloxy, $(\text{C}_2-\text{C}_{18})$ -alkynyl, phenyl, fluorine, 25 hydroxyl, trifluoromethyl, $(\text{C}_1-\text{C}_{18})$ -alkylsulfinyl, $(\text{C}_1-\text{C}_{18})$ -alkylsulfonyl, phenylsulfinyl, naphthylsulfinyl, naphthylsulfonyl, $(\text{C}_{11}-\text{C}_{18})$ -alkoxy, (C_3-C_8) -cycloalkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy, 30 $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy, phenyl- (C_2-C_4) -alkoxy, phenoxy, $(\text{C}_1-\text{C}_{12})$ -alkanoyl, phenyl- (C_1-C_4) -alkanoyl or benzoyl, where, in substituents with a phenyl or naphthyl ring, this optionally 35 carries up to 5 identical or different substituents from the series comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ or (C_1-C_6) -alkylsulfonyl, and

THIS SHEET HAS BEEN
REPLACED



R^4 is (C_1-C_{18}) -alkyl, (C_2-C_{18}) -alkenyl, (C_2-C_{18}) -alkenyloxy, (C_2-C_{18}) -alkynyl, phenyl, chlorine, fluorine, bromine, hydroxyl, trifluoromethyl, (C_1-C_{18}) -alkylsulfinyl, (C_1-C_{18}) -alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, (C_1-C_{18}) -alkoxy apart from 1-butoxy, (C_3-C_8) -cycloalkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$, phenyl- (C_2-C_4) -alkoxy, phenoxy, (C_1-C_{12}) -alkanoyl, phenyl- (C_1-C_4) -alkanoyl or benzoyl, where, in substituents with a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series consisting of fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$ or (C_1-C_6) -alkylsulfonyl,

or a physiologically active salt thereof.

13. A compound of the formula I as claimed in claims 4 and 12, in which

m is 0,
A is a $-CH_2-$ group,
B is $-CO_2H$,
 R^1 and R^5 are hydrogen and
 R^2 is (C_1-C_{18}) -alkyl, (C_1-C_{18}) -alkenyl, phenyl, chlorine, fluorine, bromine, trifluoromethyl, (C_1-C_{18}) -alkylsulfinyl, (C_1-C_{18}) -alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, (C_1-C_{18}) -alkoxy, (C_3-C_8) -cycloalkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$, where, in substituents having a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series consisting of fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$ and (C_1-C_6) -alkylsulfonyl, and
 R^3 is (C_1-C_{18}) -alkyl, (C_1-C_{18}) -alkenyl, phenyl,



- fluorine, trifluoromethyl, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkylsulfonyl, phenylsulfinyl, naphthylsulfinyl, naphthylsulfonyl, (C₁₁-C₁₈)-alkoxy, (C₃-C₈)-cycloalkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy, phenyl-(C₂-C₄)-alkoxy, phenoxy, (C₁-C₁₂)-alkanoyl, phenyl-(C₁-C₄)-alkanoyl or benzoyl, where, in substituents having a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g and (C₁-C₆)-alkylsulfonyl, and
- 15 R⁴ is (C₁-C₁₈)-alkyl, (C₁-C₁₈)-alkenyl, phenyl, chlorine, fluorine, bromine, trifluoromethyl, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, (C₁-C₁₈)-alkoxy apart from 1-butoxy, (C₃-C₈)-cycloalkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, phenyl-(C₂-C₄)-alkoxy, phenoxy, (C₁-C₁₂)-alkanoyl, phenyl-(C₂-C₄)-alkanoyl or benzoyl, where, in substituents with a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₆)-alkylsulfonyl,
- 30 or a physiologically active salt thereof.

14. A compound of the formula I as claimed in claims 5 and 13, in which

m is 0,

A is a -CH₂- group,

35 R¹ and R⁵ are hydrogen,

one of the substituents R², R³ or R⁴ is hydrogen and the other two are identical or different and are hydrogen,

THIS SHEET HAS BEEN
REPLACED



- (C₁-C₁₆)-alkyl, fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₁₆)-alkylsulfonyl, phenylsulfonyl, (C₁-C₁₆)-alkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1g)F_g, benzyloxy or phenoxy, where, in substituents which contain a phenyl ring, this optionally carries up to 3 substituents from the series comprising fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₄)-alkylsulfonyl, or a physiologically active salt thereof.
- 10 15. A compound of the formula I as claimed in claims 6 and 14, in which
- m is 0,
- A is a -CH₂- group,
- R¹, R² and R⁵ are hydrogen and
- 15 R³ is (C₁-C₁₆)-alkyl, fluorine, trifluoromethyl, (C₁-C₁₆)-alkylsulfonyl, naphthylsulfonyl, (C₁₁-C₁₆)-alkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy or phenoxy, where, in substituents which contain a phenyl ring, this optionally carries up to 3 substituents from the series comprising fluorine, chlorine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g and (C₁-C₄)-alkylsulfonyl, and
- 20
- 25 R⁴ is (C₁-C₁₆)-alkyl, fluorine, chlorine, trifluoromethyl, (C₁-C₁₆)-alkylsulfonyl, phenylsulfonyl, naphthylsulfonyl, (C₁-C₁₆)-alkoxy apart from 1-butoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy or -O-[CH₂]_x-C_fH_(2f+1-g)F_g, where, in substituents which contain a phenyl ring, this optionally carries up to 3 substituents from the series consisting of fluorine, chlorine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g and (C₁-C₄)-alkylsulfonyl, or a physiologically active salt thereof.
- 30
- 35 16. A compound of the formula I as claimed in claims 7 and 15, in which



m is 0,
A is a $-\text{CH}_2-$ group,
 R^1 , R^2 , R^3 and R^5 are hydrogen and
 R^4 is $(\text{C}_1-\text{C}_{12})$ -alkyl, fluorine, chlorine, bromine, tri-
5 fluoromethyl, $(\text{C}_1-\text{C}_{12})$ -alkylsulfonyl, phenylsulfonyl,
naphthylsulfonyl, $(\text{C}_1-\text{C}_{12})$ -alkoxy apart from 1-butoxy,
 (C_1-C_6) -alkoxy- (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$, or
phenoxy, where, in substituents which contain a phenyl or
naphthyl ring, this is optionally monosubstituted by
10 fluorine, chlorine, bromine, trifluoromethyl, (C_1-C_6) -
alkyl, (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ or (C_1-C_4) -
alkylsulfonyl, or a physiologically active salt thereof.

17. A compound of the formula I as claimed in claims 8
and 16, in which
15 m is 0,
A is a $-\text{CH}_2-$ group,
 R^1 , R^3 , R^4 and R^5 are hydrogen and
 R^2 is $(\text{C}_1-\text{C}_{12})$ -alkyl, fluorine, chlorine, bromine, tri-
fluoromethyl, $(\text{C}_1-\text{C}_{12})$ -alkylsulfonyl, phenylsulfonyl,
20 naphthylsulfonyl, $(\text{C}_1-\text{C}_{12})$ -alkoxy, (C_1-C_6) -alkoxy- (C_1-C_6) -
alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$, benzyloxy or phenoxy,
where, in substituents which contain a phenyl or naphthyl
ring, this is optionally monosubstituted by fluorine,
chlorine, bromine, trifluoromethyl, (C_1-C_6) -alkyl,
25 (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ or (C_1-C_4) -alkyl-
sulfonyl, or a physiologically active salt thereof.

18. A compound of the formula I as claimed in claims 9
and 17, in which
m is 0,
30 A is a $-\text{CH}_2-$ group,
 R^1 , R^2 and R^5 are hydrogen,
 R^4 is hydrogen or chlorine and
 R^3 is hydrogen, fluorine, (C_1-C_4) -alkoxy- (C_1-C_6) -alkoxy,
 $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ apart from trifluoroethoxy,
35 pentafluoropropoxy and heptafluorobutoxy, naphthyl-
sulfonyl or phenoxy, where, in substituents which
contain a phenyl or naphthyl ring, this optionally





* C L M S *

- 97 -

carries up to 3 identical or different substituents, preferably one substituent, from the series comprising fluorine, chlorine, bromine, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-alkyl and (C₁-C₆)-alkoxy, or a physiologically active salt thereof.

19. A compound of the formula I as claimed in claims 10 and 18, in which
m is 0,
A is a -CH₂- group,
R¹, R², R⁴ and R⁵ are hydrogen and
R³ is hydrogen, fluorine, (C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy, naphthylsulfonyl or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this optionally carries up to 3 identical or different substituents, preferably one substituent, from the series comprising fluorine, chlorine, bromine, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-alkyl and (C₁-C₆)-alkoxy, or a physiologically active salt thereof.

20. A prodrug to a compound of the formula I as claimed in claims 1 to 19.

21. A process for the preparation of a compound of the formula I as claimed in claims 1 to 19, which comprises

1.i1.) reacting a quinoline-2-carboxylic acid of the formula II (R¹¹ = H) with an amino ester of the formula III to give an amide ester of the formula IV, or

1.i2.) reacting a quinoline-2-carboxylic acid ester of the formula II (R¹¹ = lower alkyl) under the conditions of aminolysis to give a compound of the formula IV; where R¹⁰ is PG (Protecting


This Sheet Has Been Replaced





THIS SHEET HAS BEEN
REPLACED



Group) and A-B is $(CH_2)_{1-4}-CO_2H$



1.ii) liberating a compound of the formula I or V from its ester of the formula IV or VI; and/or




5 1.iii) obtaining a compound of the formula I or VI from a compound of the formula V or IV by splitting off the hydroxyl-protecting group R^{10} , and if appropriate


This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED




V



I




IV





VI

1.iv) oxidizing a compound of the formula I, IV, V or VI to give a compound of the formula Ia, IVa, Va or VIa.



I



Ia

22. A compound as claimed in claims 1 to 19 for use for
5 inhibiting collagen biosynthesis.

23. A compound as claimed in claims 1 to 19 as an
inhibitor of prolyl hydroxylase.

24. A compound as claimed in claims 1 to 19 for use as
a fibrosuppressant.






This Sheet Has Been Replaced



- 100 -

25. A compound as claimed in claims 1 to 19 for the preparation of a medicament against fibrotic diseases.
26. A compound as claimed in claims 1 to 19 for the preparation of a medicament against fibrotic diseases of the liver.
27. A compound as claimed in claims 1 to 19 for the preparation of a medicament against fibrotic diseases of the lung.
28. A compound as claimed in claims 1 to 19 for the preparation of a medicament against fibrotic diseases of the skin.
29. The use of a compound of the formula I as claimed in claims 1 to 19 as an inhibitor of prolyl hydroxylase and as a fibrosuppressant.
30. A medicament comprising a compound of the formula I as claimed in claims 1 to 19.
31. An intermediate product of the formula II



II

- in which
- R^1 to R^5 have the meanings which apply to a compound of the formula I as claimed in claims 1 to 19,
- R^{10} is hydrogen or an HO-protecting group and
- R^{11} is hydrogen, (C_1-C_8) -alkyl or benzyl,


THIS SHEET HAS BEEN
REPLACED



- 101 -

with the exception of compounds of the formula II in which R^{10} is benzyl or methyl if R^1 to R^5 and R^{11} are hydrogen, and those compounds of the formula II in which R^{10} is hydrogen, benzyl or p-toluenesulfonyl if R^1 to R^5 are hydrogen and R^{11} is ethyl and those compounds of formula II in which R^3 is hydrogen or methoxy, if R^1 , R^2 , R^3 , R^4 , R^5 , R^{10} and R^{11} are hydrogen.

32. An intermediate of the formula X



in which

10 m is 0 or 1 and R^2 to R^5 have the meanings which apply to a compound of the formula I as claimed in claims 1 to 19, with the exception of compounds of the formula X in which m is 0, R^3 is hydrogen, methyl or chlorine if R^2 , R^4 and

15 R^5 are hydrogen.

This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

Abstract

Substituted quinoline-2-carboxamides, their preparation and their use as pharmaceuticals, and intermediates

The invention relates to substituted quinoline-2-carboxylic acid amides of the formula I



their preparation and their use, and intermediates which are formed in the preparation of the compounds of the formula I.

The compounds according to the invention are used as inhibitors of prolyl 4-hydroxylase and as pharmaceuticals (medicaments) for treatment of fibrotic diseases.

This Sheet Has Been Replaced



4-2850188


THIS SHEET HAS BEEN
REPLACED



11 APR 1998

PATENT CLAIMS:

1. A compound of the formula I



in which

- A is (C_1-C_3) -alkylene, which is optionally mono-substituted by halogen, cyano, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -hydroxyalkyl, (C_1-C_6) -alkoxy or $-O-[CH_2]_x-C_FH_{(2f+1-g)}F_g$ or
- A is $-CHR^x-$, in which R^x is one of the substituents of the α -C atom of an α -amino acid, in particular of a naturally occurring L-amino acid or its D isomer,
- 10 B is $-CO_2H$,
- 15 R^1 and R^5 are identical or different and are hydrogen, (C_1-C_{12}) -alkyl, (C_2-C_{12}) -alkenyl, chlorine, fluorine, bromine, trifluoromethyl, (C_1-C_{12}) -alkylsulfonyl, (C_1-C_{12}) -alkylsulfinyl, phenylsulfonyl or phenylsulfinyl; where phenyl is optionally substituted by fluorine, chlorine or (C_1-C_5) -alkoxy, (C_1-C_{10}) -alkoxy, $-O-[CH_2]_x-C_FH_{(2f+1-g)}F_g$, (C_1-C_6) -alkoxy- (C_1-C_6) -alkoxy or (C_3-C_8) -cycloalkoxy,
- 20 R^2 is hydroxyl, (C_1-C_{20}) -alkyl, (C_2-C_{20}) -alkenyl, (C_2-C_{20}) -alkynyl, (C_1-C_{20}) -alkoxy, (C_2-C_{20}) -alkenyl-oxy, (C_2-C_{20}) -alkynyloxy, retinyloxy, (C_1-C_{20}) -alkoxy- (C_1-C_8) -alkyl, (C_2-C_{20}) -alkenyl-oxy- (C_1-C_8) -alkyl, retinyloxy- (C_1-C_6) -alkyl, (C_2-C_{20}) -alkynyloxy-



11 APR 1998

-69-

- (C₁-C₈)-alkyl, halogen, cyano, trifluoromethyl, (C₁-C₁₂)-hydroxyalkyl, (C₁-C₂₀)-alkanoyl, (C₇-C₁₆)-aralkanoyl, (C₆-C₁₂)-aroyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, (C₁-C₁₈)-alkyl-mercapto, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkyl-sulfonyl, (C₆-C₁₂)-arylmercapto, (C₆-C₁₂)-aryl-sulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₂)-aralkyl-mercapto, (C₇-C₁₂)-aralkylsulfinyl, (C₇-C₁₂)-aralkyl-sulfonyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy, carboxyl, (C₁-C₂₀)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxcarbonyl, (C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxy-carbonyl, (C₂-C₂₀)-alkenyloxcarbonyl, retinyloxy-carbonyl, (C₂-C₂₀)-alkynyloxcarbonyl, (C₃-C₈)-cyclo-alkyl-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl,
- (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy, (C₃-C₈)-cyclo-alkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cyclo-alkyloxy-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, NR^YR^Z, substituted (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy or (C₇-C₁₁)-aralkoxy-(C₁-C₆)-alkoxy,
- carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-carbamoyl, N,N-dicyclo(C₃-C₈)-alkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl,



11 APR 1998

-70-

5 N-(C₃-C₈)-cycloalkyl-(C₁-C₁₆)-alkyl) carbamoyl,
N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-
alkyl) carbamoyl, N-(+)-dehydroabietyl carbamoyl,
N-(C₁-C₆)-alkyl-N-(+)-dehydroabietyl carbamoyl,
N-(C₆-C₁₂)-aryl carbamoyl, N-(C₇-C₁₆)-aralkyl-
carbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-aryl carbamoyl,
N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkyl carbamoyl,
N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-alkyl) carbamoyl,
N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl) carbamoyl,
10 N-((C₇-C₁₆)-aralkyloxy-(C₁-C₆)-alkyl) carbamoyl,
N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-
alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-
aryloxy-(C₁-C₁₀)-alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-
N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl) carbamoyl, or
15 CON(CH₂)_h, in which one CH₂ group can be replaced by
O, S, N-(C₁-C₁₆)-alkylimino, N-(C₃-C₈)-cycloalkyl-
imino, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkylimino,
N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or
N-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7,
20 where an aromatic radical carries 1, 2, 3, 4 or 5
identical or different substituents from the series
comprising hydrogen, halogen, cyano, nitro,
hydroxyl, trifluoromethyl, (C₁-C₁₆)-alkyl, (C₂-C₁₆)-
alkenyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₁₆)-alkoxy,
25 (C₁-C₁₆)-alkenyloxy, -O-[CH₂]_xC_fH_(2f+1-g)F_g, -OCF₂Cl,
-O-CF₂-CHFCl, (C₁-C₆)-alkylmercapto, (C₁-C₆)-
alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C-C₁)-₆
alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-
(C₁-C₄)-alkyl carbamoyl, N,N-di-(C₁-C₄)-
30 alkyl carbamoyl, (C₁-C₆)-alkylcarbonyloxy, (C₃-C₈)-
cycloalkyl carbamoyl, phenyl, benzyl, phenoxy,
benzyloxy, phenylmercapto, phenylsulfonyl,
phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl
and N,N-di-(C₁-C₄)-alkylsulfamoyl, or optionally
35 carries up to 3 of the abovementioned identical or
different substituents and two adjacent carbon atoms
of the aralkyloxy radical together carry a chain
-[CH₂]- and/or -CH=CH-CH=CH-, where one CH₂ group of



11 APR 1998

-71-

the chain is optionally replaced by O, ~~S, SO, SO₂ or~~
NR',

- R^3 is hydroxyl, (C₁-C₂₀)-alkyl, (C₂-C₂₀)-alkenyl, (C₂-C₂₀)-alkynyl, (C₁₁-C₂₀)-alkoxy, (C₇-C₂₀)-alkenyl-
oxy, (C₂-C₂₀)-alkynyloxy, retinyloxy, (C₁-C₂₀)-
alkoxy-(C₁-C₈)-alkyl, (C₂-C₂₀)-alkenyloxy-(C₁-C₈)-
alkyl, retinyloxy-(C₁-C₆)-alkyl-(C₂-C₂₀)-alkynyloxy-
(C₁-C₈)-alkyl, fluorine, cyano, trifluoromethyl, (C₁-C₁₂)-hydroxyalkyl, (C₁-C₂₀)-alkanoyl, (C₇-C₁₆)-
aralkanoyl, (C₆-C₁₂)-aroyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-
aralkyl, (C₁-C₁₈)-alkylmercapto, (C₁-C₁₈)-alkyl-
sulfinyl, (C₁-C₁₈)-alkylsulfonyl, (C₆-C₁₂)-aryl-
mercapto, (C₆-C₁₂)-arylsulfinyl, (C₇-C₁₂)-aralkyl-
mercapto, (C₇-C₁₂)-aralkylsulfinyl, (C₇-C₁₂)-aralkyl-
sulfonyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy,
apart from benzyloxy, carboxyl, (C₁-C₂₀)-alkoxy-
carbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl,
(C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-aralkoxycarbonyl,
(C₃-C₈)-cycloalkoxycarbonyl, (C₂-C₂₀)-
alkenyloxycarbonyl, retinyloxycarbonyl, (C₂-C₂₀)-
alkynyloxycarbonyl, (C₃-C₈)-cycloalkyl-(C₁-C₆)-
alkyloxycarbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-
alkoxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-
alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-
alkoxycarbonyl,

-O-[CH₂]_x-C_fH_(2f+1-g)Hal_g apart from trifluoroethoxy,
pentafluoropropoxy and heptafluorobutoxy, (C₁-C₁₂)-
alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-
alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy, (C₃-C₈)-
cycloalkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl,
(C₃-C₈)-cycloalkyloxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-
alkoxy, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl,
(C₃-C₈)-cycloalkyloxy-(C₁-C₈)-alkoxy, (C₃-C₈)-cyclo-
alkyl-(C₁-C₁₂)-alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cyclo-
alkyl-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-
cycloalkoxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, NR^YR^Z,
substituted (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-



11 APR 1998

-72-

aralkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy or (C₇-C₁₁)-aralkoxy-(C₁-C₆)-alkoxy,

- 5 carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkylcarbamoyl, N,N-dicyclo(C₃-C₈)-alkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₆)-alkyl carbamoyl,
- 10 N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl) carbamoyl, N-(+)-dehydroabietylcarbamoyl, N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl, N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcarbamoyl,
- 15 N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl, N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-alkyl) carbamoyl, N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl) carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₆)-alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl) carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl) carbamoyl, or CON(CH₂)_h, in which one CH₂ group can be replaced by O, S, N-(C₁-C₁₆)-alkyl-
- 25 imino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or N-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7, where an aromatic radical carries 1, 2, 3, 4 or 5 identical or different
- 30 substituents from the series comprising hydrogen, halogen, cyano, nitro, hydroxyl, trifluoromethyl, (C₁-C₁₆)-alkyl, (C₂-C₁₆)-alkenyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₁₆)-alkoxy, (C₁-C₁₆)-alkenyloxy, -O-[CH₂]_xC_FH_(2f+1-g)F_g, -OCF₂Cl, -O-CF₂-CHFCl, (C₁-C₆)-alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy,
- 35



11 APR 1998

-73-

5 (C₃-C₈)-cycloalkylcarbamoyl, phenyl, benzyl, phenoxy, benzyloxy, phenylmercapto, phenylsulfonyl, phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl and N,N-di-(C₁-C₄)-alkylsulfamoyl, or optionally carries up to 3 of the abovementioned identical or different substituents and two adjacent carbon atoms of the aralkyloxy radical together carry a chain -[CH₂]- and/or -CH=CH-CH=CH-, where one CH₂ group of the chain is optionally replaced by O, S, SO, SO₂ or NR', and

10

R⁴ is hydroxyl, (C₁-C₂₀)-alkyl, (C₂-C₂₀)-alkenyl, (C₂-C₂₀)-alkynyl, (C₁-C₂₀)-alkoxy apart from 1-butoxy, (C₂-C₂₀)-alkenyloxy, (C₂-C₂₀)-alkynyloxy, retinyloxy, (C₁-C₂₀)-alkoxy-(C₁-C₆)-alkyl, (C₂-C₂₀)-alkenyloxy-(C₁-C₈)-alkyl, retinyloxy-(C₁-C₆)-alkyl, (C₂-C₂₀)-alkynyloxy-(C₁-C₈)-alkyl, halogen, cyano, trifluoromethyl, (C₁-C₁₂)-hydroxyalkyl, (C₁-C₂₀)-alkanoyl, (C₇-C₁₆)-aralkanoyl, (C₆-C₁₂)-aroyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, -O-[CH₂]_x-C_FH_(2f+1-g)F_g, (C₁-C₁₈)-alkylmercapto, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkylsulfonyl, (C₆-C₁₂)-arylmercapto, (C₆-C₁₂)-arylsulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₂)-aralkylmercapto, (C₇-C₁₂)-aralkylsulfinyl, (C₇-C₁₂)-aralkylsulfonyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy apart from benzyloxy, carboxyl, (C₁-C₂₀)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl, (C₂-C₂₀)-alkenyloxycarbonyl, retinyloxycarbonyl, (C₂-C₂₀)-alkynyloxycarbonyl, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl,

20

25

30

35 (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy, (C₃-C₈)-cyclo-



11 APR 1998

-74-

- alkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy-(C₁-C₈)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, NR^YR^Z, substituted (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy or (C₇-C₁₁)-aralkyloxy-(C₁-C₆)-alkoxy,
- carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkylcarbamoyl, N,N-dicyclo(C₃-C₈)-alkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₆)-alkylcarbamoyl, N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)carbamoyl, N-(+)-dehydroabietylcarbamoyl, N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl, N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl, N-((C₁-C₁₂)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₆)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyl, or CON(CH₂)_h, in which one CH₂ group can be replaced by O, S, N-(C₁-C₁₆)-alkylimino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino or N-(C₁-C₁₂)-alkoxy-(C₁-C₆)-alkylimino and h is 3 to 7, where an aromatic radical carries 1, 2, 3, 4 or 5 identical or different substituents from the series comprising hydrogen,



11 APR 1998

halogen, cyano, nitro, hydroxyl, trifluoromethyl,
(C₁-C₁₆)-alkyl, (C₂-C₁₆)-alkenyl, (C₁-C₆)-hydroxy-
alkyl, (C₁-C₁₆)-alkoxy, (C₁-C₁₆)-alkenyloxy, -O-
[CH₂]_xC_fH_(2f+1-g)F_g, -O-CF₂Cl, -OCF₂-CHFCl, (C₁-C₆)-
5 alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkyl-
sulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxy-
carbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-
di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy,
10 (C₃-C₈)-cycloalkylcarbamoyl, phenyl, benzyl,
phenoxy, benzyloxy, phenylmercapto, phenylsulfonyl,
phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl
and N,N-di-(C₁-C₄)-alkylsulfamoyl, or optionally
15 carries up to 3 of the abovementioned identical or
different substituents and two adjacent carbon atoms
of the aralkyloxy radical together carry a chain
-[CH₂]- and/or -CH=CH-CH=CH-, where one CH₂ group of
the chain is optionally replaced by O, S, SO, SO₂ or
NR',

R^Y and R^Z are identical or different and are hydrogen,
20 (C₆-C₁₂)-aryl, (C₁-C₁₀)-alkyl, (C₃-C₁₀)-cycloalkyl,
(C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkoxy-(C₁-
C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkyl, (C₁-C₁₀)-
alkanoyl, optionally substituted (C₇-C₁₆)-
aralkanoyl, or optionally substituted (C₆-C₁₂)-
25 aroyl, or

R^Y and R^Z together are -[CH₂]_n-, wherein one CH₂ group can
be replaced by O, S, N-(C₁-C₄)-alkanoylimino or
N-(C₁-C₄)-alkoxycarbonylimino,

m is 0 or 1,
30 f is 1 to 8,
g is 0 or 1 to (2f + 1),
h is 3 to 6 and
x is 0 to 3, or a physiologically active salt thereof.

2. A compound of the formula I as claimed in claim 1,
35 in which



11 APR 1998

-76-

- m is 0,
A is a $-\text{CH}_2-$ group, which can be substituted by a methyl group,
B is $-\text{CO}_2\text{H}$,
5 R^1 is hydrogen, (C_1-C_4) -alkyl, (C_1-C_4) -alkoxy, chlorine or bromine,
 R^5 is hydrogen, fluorine, chlorine or methyl,
 R^2 is $(\text{C}_1-\text{C}_{18})$ -alkyl, $(\text{C}_2-\text{C}_{18})$ -alkenyl, $(\text{C}_2-\text{C}_{18})$ -alkenyloxy, $(\text{C}_2-\text{C}_{18})$ -alkynyl, phenyl, chlorine,
10 fluorine, bromine, hydroxyl, trifluoromethyl, $(\text{C}_1-\text{C}_{18})$ -alkylsulfinyl, $(\text{C}_1-\text{C}_{18})$ -alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, $(\text{C}_1-\text{C}_{18})$ -alkoxy, (C_3-C_8) -cycloalkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy,
15 $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$, phenyl- (C_1-C_4) -alkoxy, phenoxy, $(\text{C}_1-\text{C}_{12})$ -alkanoyl, phenyl- (C_1-C_4) -alkanoyl or benzoyl, where, in substituents with a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series
20 comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ or (C_1-C_6) -alkylsulfonyl, and
 R^3 is $(\text{C}_1-\text{C}_{18})$ -alkyl, $(\text{C}_2-\text{C}_{18})$ -alkenyl, $(\text{C}_7-\text{C}_{18})$ -alkenyloxy, $(\text{C}_2-\text{C}_{18})$ -alkynyl, phenyl, fluorine,
25 hydroxyl, trifluoromethyl, $(\text{C}_1-\text{C}_{18})$ -alkylsulfinyl, $(\text{C}_1-\text{C}_{18})$ -alkylsulfonyl, phenylsulfinyl, naphthylsulfinyl, naphthylsulfonyl, $(\text{C}_{11}-\text{C}_{18})$ -alkoxy, (C_3-C_8) -cycloalkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy,
30 $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy, phenyl- (C_2-C_4) -alkoxy, phenoxy, $(\text{C}_1-\text{C}_{12})$ -alkanoyl, phenyl- (C_1-C_4) -alkanoyl or benzoyl, where, in substituents with a phenyl or naphthyl ring, this optionally
35 carries up to 5 identical or different substituents from the series comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-\text{O}-[\text{CH}_2]_x-\text{C}_f\text{H}_{(2f+1-g)}\text{F}_g$ or (C_1-C_6) -alkylsulfonyl, and



11 APR 1998

-77-

R^4 is (C_1-C_{18}) -alkyl, (C_2-C_{18}) -alkenyl, (C_2-C_{18}) -alkenyloxy, (C_2-C_{18}) -alkynyl, phenyl, chlorine, fluorine, bromine, hydroxyl, trifluoromethyl, (C_1-C_{18}) -alkylsulfinyl, (C_1-C_{18}) -alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, (C_1-C_{18}) -alkoxy apart from 1-butoxy, (C_3-C_8) -cycloalkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$, phenyl- (C_2-C_4) -alkoxy, phenoxy, (C_1-C_{12}) -alkanoyl, phenyl- (C_1-C_4) -alkanoyl or benzoyl, where, in substituents with a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series consisting of fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$ or (C_1-C_6) -alkylsulfonyl,

or a physiologically active salt thereof.

3. A compound of the formula I as claimed in claim 2, in which

m is 0,
A is a $-CH_2-$ group,
B is $-CO_2H$,
 R^1 and R^5 are hydrogen and
 R^2 is (C_1-C_{18}) -alkyl, (C_1-C_{18}) -alkenyl, phenyl, chlorine, fluorine, bromine, trifluoromethyl, (C_1-C_{18}) -alkylsulfinyl, (C_1-C_{18}) -alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, (C_1-C_{18}) -alkoxy, (C_3-C_8) -cycloalkoxy, (C_1-C_8) -alkoxy- (C_1-C_8) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$, where, in substituents having a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series consisting of fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C_1-C_6) -alkyl, (C_1-C_6) -alkoxy, $-O-[CH_2]_x-C_fH_{(2f+1-g)}F_g$ and (C_1-C_6) -alkylsulfonyl, and
 R^3 is (C_1-C_{18}) -alkyl, (C_1-C_{18}) -alkenyl, phenyl,



11 APR 1998

- fluorine, trifluoromethyl, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkylsulfonyl, phenylsulfinyl, naphthylsulfinyl, naphthylsulfonyl, (C₁₁-C₁₈)-alkoxy, (C₃-C₈)-cycloalkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy, phenyl-(C₂-C₄)-alkoxy, phenoxy, (C₁-C₁₂)-alkanoyl, phenyl-(C₁-C₄)-alkanoyl or benzoyl, where, in substituents having a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g and (C₁-C₆)-alkylsulfonyl, and
- 15 R⁴ is (C₁-C₁₈)-alkyl, (C₁-C₁₈)-alkenyl, phenyl, chlorine, fluorine, bromine, trifluoromethyl, (C₁-C₁₈)-alkylsulfinyl, (C₁-C₁₈)-alkylsulfonyl, phenylsulfinyl, phenylsulfonyl, naphthylsulfinyl, naphthylsulfonyl, (C₁-C₁₈)-alkoxy apart from 1-butoxy, (C₃-C₈)-cycloalkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, phenyl-(C₂-C₄)-alkoxy, phenoxy, (C₁-C₁₂)-alkanoyl, phenyl-(C₂-C₄)-alkanoyl or benzoyl, where, in substituents with a phenyl or naphthyl ring, this optionally carries up to 5 identical or different substituents from the series comprising fluorine, chlorine, bromine, nitrile, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₆)-alkylsulfonyl,
- 30 or a physiologically active salt thereof.
4. A compound of the formula I as claimed in claim 3, in which
- m is 0,
- A is a -CH₂- group,
- 35 R¹ and R⁵ are hydrogen,
- one of the substituents R², R³ or R⁴ is hydrogen and the other two are identical or different and are hydrogen,



11 APR 1998

-79-

- (C₁-C₁₆)-alkyl, fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₁₆)-alkylsulfonyl, phenylsulfonyl, (C₁-C₁₆)-alkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1g)F_g, benzyloxy or phenoxy, where, in substituents
5 which contain a phenyl ring, this optionally carries up to 3 substituents from the series comprising fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₄)-alkylsulfonyl, or a physiologically active salt thereof.
- 10 5. A compound of the formula I as claimed in claim 4,
in which
m is 0,
A is a -CH₂- group,
R¹, R² and R⁵ are hydrogen and
15 R³ is (C₁-C₁₆)-alkyl, fluorine, trifluoromethyl, (C₁-C₁₆)-alkylsulfonyl, naphthylsulfonyl, (C₁₁-C₁₆)-alkoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy or phenoxy, where, in
20 substituents which contain a phenyl ring, this optionally carries up to 3 substituents from the series comprising fluorine, chlorine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g and (C₁-C₄)-alkylsulfonyl, and
- 25 R⁴ is (C₁-C₁₆)-alkyl, fluorine, chlorine, trifluoromethyl, (C₁-C₁₆)-alkylsulfonyl, phenylsulfonyl, naphthylsulfonyl, (C₁-C₁₆)-alkoxy apart from 1-butoxy, (C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy or -O-[CH₂]_x-C_fH_(2f+1-g)F_g, where, in substituents which contain a
30 phenyl ring, this optionally carries up to 3 substituents from the series consisting of fluorine, chlorine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g and (C₁-C₄)-alkylsulfonyl, or a physiologically active salt thereof.

- 35 6. A compound of the formula I as claimed in claim 5,
in which



11 APR 1998

-80-

m is 0,
A is a -CH₂- group,
R¹, R², R³ and R⁵ are hydrogen and
R⁴ is (C₁-C₁₂)-alkyl, fluorine, chlorine, bromine, tri-
5 fluoromethyl, (C₁-C₁₂)-alkylsulfonyl, phenylsulfonyl,
naphthylsulfonyl, (C₁-C₁₂)-alkoxy apart from 1-butoxy,
(C₁-C₆)-alkoxy-(C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, or
phenoxy, where, in substituents which contain a phenyl or
naphthyl ring, this is optionally monosubstituted by
10 fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₆)-
alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₄)-
alkylsulfonyl, or a physiologically active salt thereof.

7. A compound of the formula I as claimed in claim 6
, in which
15 m is 0,
A is a -CH₂- group,
R¹, R³, R⁴ and R⁵ are hydrogen and
R² is (C₁-C₁₂)-alkyl, fluorine, chlorine, bromine, tri-
fluoromethyl, (C₁-C₁₂)-alkylsulfonyl, phenylsulfonyl,
20 naphthylsulfonyl, (C₁-C₁₂)-alkoxy, (C₁-C₆)-alkoxy-(C₁-C₆)-
alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g, benzyloxy or phenoxy,
where, in substituents which contain a phenyl or naphthyl
ring, this is optionally monosubstituted by fluorine,
chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl,
25 (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g or (C₁-C₄)-alkyl-
sulfonyl, or a physiologically active salt thereof.

8. A compound of the formula I as claimed in claim 7,
in which
m is 0,
30 A is a -CH₂- group,
R¹, R² and R⁵ are hydrogen,
R⁴ is hydrogen or chlorine and
R³ is hydrogen, fluorine, (C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy,
-O-[CH₂]_x-C_fH_(2f+1-g)F_g apart from trifluoroethoxy,
35 pentafluoropropoxy and heptafluorobutoxy, naphthyl-
sulfonyl or phenoxy, where, in substituents which
contain a phenyl or naphthyl ring, this optionally



THIS SHEET HAS BEEN
REPLACED

REPLACEMENT SHEET

11 APR 1998

-81-

carries up to 3 identical or different substituents, preferably one substituent, from the series comprising fluorine, chlorine, bromine, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-alkyl and (C₁-C₆)-alkoxy, or a physiologically active salt thereof.

5

9. A compound of the formula I as claimed in claim 8, in which

m is 0,

10 A is a -CH₂- group,

R¹, R², R⁴ and R⁵ are hydrogen and

R³ is hydrogen, fluorine, (C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy, naphthyl-sulfonyl or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this optionally carries up to 3 identical or different substituents, preferably one substituent, from the series comprising fluorine, chlorine, bromine, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-alkyl and (C₁-C₆)-alkoxy, or a physiologically active salt thereof.

15

20

10. A prodrug to a compound of the formula I as claimed in claims 1 to 9.

25 11. A process for the preparation of a compound of the formula I as claimed in claims 1 to 9, which comprises

1.i1.) reacting a quinoline-2-carboxylic acid of the formula II (R¹¹ = H) with an amino ester of the formula III to give an amide ester of the formula IV, or

30

1.i2.) reacting a quinoline-2-carboxylic acid ester of the formula II (R¹¹ = lower alkyl) under the conditions of aminolysis to give a compound of the formula IV; where R¹⁰ is PG (Protecting

This Sheet Has Been Replaced






THIS SHEET HAS BEEN
REPLACED

REPLACEMENT SHEET



11 APR 1998

-82-

Group) and A-B is $(\text{CH}_2)_{1-4}\text{-CO}_2\text{H}$



1.ii) liberating a compound of the formula I or V from its ester of the formula IV or VI; and/or



5 1.iii) obtaining a compound of the formula I or VI from a compound of the formula V or IV by splitting off the hydroxyl-protecting group R^{10} , and if appropriate

This Sheet Has Been Replaced







THIS SHEET HAS BEEN
REPLACED


-83-

REPLACEMENT SHEET

11 APR 1998



1.iv) oxidizing a compound of the formula I, IV, V or VI to give a compound of the formula Ia, IVa, Va or VIa.



12. A compound as claimed in claims 1 to 9 for use for inhibiting collagen biosynthesis.

13. A compound as claimed in claims 1 to 9 as an inhibitor of prolyl hydroxylase.

14. A compound as claimed in claims 1 to 9 for use as a fibrosuppressant.

This Sheet Has Been Replaced






THIS SHEET HAS BEEN
REPLACED

REPLACEMENT SHEET

11 APR 1998

-84-

15. A compound as claimed in claims 1 to 9 for the preparation of a medicament against fibrotic diseases.
16. A compound as claimed in claims 1 to 9 for the preparation of a medicament against fibrotic diseases of the liver.
17. A compound as claimed in claims 1 to 9 for the preparation of a medicament against fibrotic diseases of the lung.
18. A compound as claimed in claims 1 to 9 for the preparation of a medicament against fibrotic diseases of the skin.
19. The use of a compound of the formula I as claimed in claims 1 to 9 as an inhibitor of prolyl hydroxylase and as a fibrosuppressant.
20. A medicament comprising a compound of the formula I as claimed in claims 1 to 9.
21. An intermediate product of the formula II



This Sheet Has Been Replaced



II

in which

- R^1 to R^5 have the meanings which apply to a compound of the formula I as claimed in claims 1 to 9,
- R^{10} is hydrogen or an HO-protecting group and
- R^{11} is hydrogen, (C_1-C_8) -alkyl or benzyl,



THIS SHEET HAS BEEN
REPLACED


REPLACEMENT SHEET

11 APR 1998

-85-

with the exception of compounds of the formula II in which R^{10} is benzyl or methyl if R^1 to R^5 and R^{11} are hydrogen, and those compounds of the formula II in which R^{10} is hydrogen, benzyl or p-toluenesulfonyl if R^1 to R^5 are hydrogen and R^{11} is ethyl and those compounds of formula II in which R^3 is hydrogen or methoxy, if R^1 , R^2 , R^3 , R^4 , R^5 , R^{10} and R^{11} are hydrogen.

22. An intermediate of the formula X



in which

10 m is 0 or 1 and R^2 to R^5 have the meanings which apply to a compound of the formula I as claimed in claims 1 to 9, with the exception of compounds of the formula X in which m is 0, R^3 is hydrogen, methyl or chlorine if R^2 , R^4 and

15 R^5 are hydrogen.

This Sheet Has Been Replaced




11 APR 1998

Abstract

Substituted quinoline-2-carboxamides, their preparation and their use as pharmaceuticals, and intermediates

The invention relates to substituted quinoline-2-carboxylic acid amides of the formula I



their preparation and their use, and intermediates which are formed in the preparation of the compounds of the formula I.

The compounds according to the invention are used as inhibitors of prolyl 4-hydroxylase and as pharmaceuticals (medicaments) for treatment of fibrotic diseases.



-81-

carries up to 3 identical or different substituents, preferably one substituent, from the series comprising fluorine, chlorine, bromine, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-alkyl and (C₁-C₆)-alkoxy, or a physiologically active salt thereof.

9. A compound of the formula I as claimed in claim 8, in which
- m is 0,
- 10 A is a -CH₂- group,
- R¹, R², R⁴ and R⁵ are hydrogen and
- R³ is hydrogen, fluorine, (C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)F_g apart from trifluoroethoxy, pentafluoropropoxy and heptafluorobutoxy, naphthyl-sulfonyl or phenoxy, where, in substituents which contain a phenyl or naphthyl ring, this optionally carries up to 3 identical or different substituents, preferably one substituent, from the series comprising fluorine, chlorine, bromine, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-alkyl and (C₁-C₆)-alkoxy, or a physiologically active salt thereof.

10. A process for the preparation of a compound of the formula I as claimed in claims 1 to 9, which comprises
- 25 1.i1.) reacting a quinoline-2-carboxylic acid of the formula II (R¹¹ = H) with an amino ester of the formula III to give an amide ester of the formula IV, or
- 30 1.i2.) reacting a quinoline-2-carboxylic acid ester of the formula II (R¹¹ = lower alkyl) under the conditions of aminolysis to give a compound of the formula IV; where R¹⁰ is PG (Protecting


REPLACEMENT SHEET

19 NOV 1998





-82-

Group) and A-B is $(\text{CH}_2)_{1-4}\text{-CO}_2\text{H}$



1.ii) liberating a compound of the formula I or V from its ester of the formula IV or VI; and/or





5 1.iii) obtaining a compound of the formula I or VI from a compound of the formula V or IV by splitting off the hydroxyl-protecting group R^{10} , and if appropriate

REPLACEMENT SHEET


19 NOV 1998



-83-



1.iv) oxidizing a compound of the formula I, IV, V or VI to give a compound of the formula Ia, IVa, Va or VIa.



11. A compound as claimed in claims 1 to 9 for use for
5 inhibiting collagen biosynthesis.

12. A compound as claimed in claims 1 to 9 as an
inhibitor of prolyl hydroxylase.

13. A compound as claimed in claims 1 to 9 for use as
a fibrosuppressant.

REPLACEMENT SHEET


19 NOV 1998



19 NOV 1998

-84-

14. A compound as claimed in claims 1 to 9 for the preparation of a medicament against fibrotic diseases.
15. A compound as claimed in claims 1 to 9 for the preparation of a medicament against fibrotic diseases of the liver.
16. A compound as claimed in claims 1 to 9 for the preparation of a medicament against fibrotic diseases of the lung.
17. A compound as claimed in claims 1 to 9 for the preparation of a medicament against fibrotic diseases of the skin.
18. The use of a compound of the formula I as claimed in claims 1 to 9 as an inhibitor of prolyl hydroxylase and as a fibrosuppressant.
19. A medicament comprising a compound of the formula I as claimed in claims 1 to 9.
20. An intermediate product of the formula II



II


- in which
- 20 R^1 to R^5 have the meanings which apply to a compound of the formula I as claimed in claims 1 to 9,
- R^{10} is hydrogen or an HO-protecting group and
- R^{11} is hydrogen, (C_1-C_8) -alkyl or benzyl,



-85-

with the exception of compounds of the formula II in which R^{10} is benzyl or methyl if R^1 to R^5 and R^{11} are hydrogen, and those compounds of the formula II in which R^{10} is hydrogen, benzyl or p-toluenesulfonyl if R^1 to R^5 are hydrogen and R^{11} is ethyl and those compounds of formula II in which R^3 is hydrogen or methoxy, if R^1 , R^2 , R^3 , R^4 , R^5 , R^{10} and R^{11} are hydrogen.

21. An intermediate of the formula X



in which

10 m is 0 or 1 and R^2 to R^5 have the meanings which apply to a compound of the formula I as claimed in claims 1 to 9, with the exception of compounds of the formula X in which m is 0, R^3 is hydrogen, methyl or chlorine if R^2 , R^4 and

15 R^5 are hydrogen.

REPLACEMENT SHEET

19 NOV 1998