

Description

The present invention relates to an amphotericin B composition for the treatment of fungal infections, and in particular, to an amphotericin B composition which has both a high LD₅₀ and therapeutic efficacy.

References

1. Holz, R.W., in F.E. Hahn, ed., Antibiotics, vol. 2, Springer-Verlag, N.Y. (1979).
2. Trembley, C., et al, Antimicrob Agents and Chemoth, 26(2):170 (1984).
3. Mehta, R.T., et al, Infection and Immunity, 47(2):429 (1985).
4. Lopez-Berestein, G., et al, Cancer Drug Delivery, 1(1):37 (1983).
5. New, R.R.C., et al, J Antimicrob Chemoth, 8:371 (1981).
6. Graybill, J.R., et al, J Infect Dis, 145:5 (1982).
7. Lopez-Berestein, G., J Infect Dis, 150(2):278 (1984).
8. Trembley, C., et al, Invest Ophthalmol, 26:711 (1985).
9. Lopez-Berestein, G., et al, J Infect Dis, 151(4):704 (1985).
10. Juliano, R., et al, Biology of the Cell, 4(39) (1983).
11. Mehta, R., et al, Biochem Biophys Acta, 770:230 (1984).
12. Hopfer, R.L., et al, Antimicrob Agents and Chemoth, 25(3):387 (1984).
13. Brockerhoff, H. et al, Biochimica biophysica Acta 691:227 (1982).
14. Crowe, L. M., et al, Biochimica Biophysica Acta 769:141 (1984).
15. Remington's Pharmaceutical Sciences, Gennaro, A.R., ed., Mack Publishing Company (1985).3.

Amphotericin B (AMB) is an effective antifungal agent, and at present, is the drug of choice for most serious systemic fungal infections (reference 1). The drug is presently available for human use as a lyophilized powder of AMB and deoxycholate ("Fungizone"). The drug binds strongly to ergosterol, a major sterol component of fungal membranes, forming pores in the membranes which allow leakage of solute molecules. The drug also has a strong binding affinity for cholesterol, a sterol present in most mammalian cell membranes, and is therefore capable of disrupting host cells.

When AMB is administered in free form (i.e., as a reconstituted AMB/deoxycholate complex) side effects resulting from red blood cell disruption are observed initially, followed by more serious cardiotoxicity, CNS and bone-marrow effects. Renal toxicity, resulting from the body's attempt to clear the drug, is also present.

Several studies have shown that AMB toxicity can be reduced by administering the drug in a liposome-bound form (references 2-12). Typically, the LD₅₀ of the drug increases from about 2-3 mg/kg body weight for the free drug up to about 8-15 mg/kg when the drug is administered in liposomal form. One limitation of liposomal formulations, however, is the apparent size instability of amphotericin B/liposomal particles when stored in an aqueous medium. Typically, AMB-containing liposomes which have an initial size distribution between 200-300 nm will spontaneously form large liposomal structures of up to several μ m (microns) on long-term storage in an aqueous medium. Liposomes with sizes greater than 1-2 μ m (microns) are generally more toxic than smaller liposomes when administered parenterally, i.e., into the bloodstream. The toxicity of large liposomes in the bloodstream is related in part to liposome blockage of the alveolar capillaries. There are also indications that relatively large liposomes are more toxic to the liver, presumably due to liposome accumulation in reticuloendothelial cells. Co-owned U.S. patent application for "Amphotericin B Liposome Composition", Serial # 781,395, filed September 27, 1985: (now US-A-4766046), discloses a novel method of preparing and storing AMB liposomes which largely overcome the size-growth problem mentioned above.

An amphotericin B composition formed by complexing AMB with a polyethylene derivative of cholesterol (PEG-cholesterol) has also been proposed (PCT application WO-A-8505030). The formulation increased the LD₅₀ of AMB to 10.0 mg/kg in mice, from 3.8 mg/kg for Fungizone, and was also less cytotoxic in cell culture. It is not known how and whether AMB complexing to PEG-cholesterol affects therapeutic efficacy against fungal infection *in vivo*, nor whether the complex can be stored in a size-stable form.

One object of the invention to provide an AMB composition which has a substantially higher LD₅₀ than AMB formulations reported in the prior art, and also is significantly more effective in treating fungal infections, *in vivo*.

Another object of the invention is to provide an AMB formulation which can be stored in suspension form over a several day period without significant particle size change, and which can be stored long term as a lyophilized preparation.

Yet another object of the invention is provide an improved method for treating fungal infections with AMB.

The invention includes an AMB composition containing particles of AMB and cholesterol sulfate, in a mole ratio of AMB to cholesterol sulfate of between 1:1 to 1:4. When prepared in a suspension form in an aqueous medium, the particles have preferred sizes between 100-400 nm. Osmotic swelling and solute trapping studies indicate that the particles are non-liposomal. A preferred composition, containing AMB and cholesterol sulfate in a mole ratio of about 1:1 shows very little change in particle size when stored in solution form over a several-day period.

The composition is formed, according to one aspect of the invention, by dispersing an aqueous suspension of amphotericin B/cholesterol sulfate particles, in a molar ratio of between 1:1 to 1:4, to optical clarity, where particle sizes are predominantly between 100-200 nm. The suspension is then lyophilized in the presence of a cryoprotectant, such as lactose, before substantial particle size increase occurs. After reconstitution in an aqueous medium, particle sizes are predominantly in the range 200-300 nm.

The composition has an LD₅₀ of greater than 15 mg/kg in mice, compared with 3.2 mg/kg for Fungizone in the same animal model system, and the 1:1 sole ratio composition has an LD₅₀ greater than 30-40 mg/kg. The therapeutic efficacy of the AMB/cholesterol sulfate composition, in treating fungal infections *in vivo*, is comparable to or greater than that observed for free AMB, depending on total AMB dosage level.

The invention also includes a use for treating fungal infections with AMB, with substantially less toxicity and greater efficacy than has been achieved heretofore. The use includes formulating an AMB/cholesterol sulfate aqueous particle suspension of the type described above and administering the suspension in a therapeutically effective amount.

These and other objects and features of the invention will become more fully apparent from the following detailed description of the invention.

Figures 1A-1C are plots of percent survivors, as a function of days following infection, of animals infected systemically with *Candida albicans*, and treated thereafter with bolus-injection doses of 0.5 mg/kg/dose (Figure 1A), 0.125 mg/kg/dose (Figure 1B), or 0.031 mg/kg/dose (Figure 1C) of AMB.

I. Preparation of the Particle Composition

A. Particle Suspension

To prepare the AMB/cholesterol sulfate particle suspension of the invention, AMB and cholesterol sulfate are combined in dry or solution form at a selected mole ration of AMB to cholesterol sulfate between 1:1 to 1:4. In a preferred method, the two components are mixed in dry form, at the selected mole ratio, then dissolved in a suitable solvent, and preferably an alcohol such as methanol. A combined molarity of AMB and cholesterol sulfate in methanol of 50 μ mole/ml, in the mole ratio range between 1:1 and 1:4 AMB to cholesterol, is suitable. For example, in a 1:1 formulation, both AMB and cholesterol sulfate would be present at 25 μ moles/ml.

A cryoprotectant may be added to the AMB/cholesterol sulfate solution, to a final preferred concentration of between 5-15%. The cryoprotectant serves two purposes in later processing steps. First, it provides a crystalline, water-soluble bulking agent on which the AMB and cholesterol sulfate components can form, when the solvent in the mixture is removed. That is, the dried crystals of cryoprotectant increase the surface area of lipid film formed on solvent removal, and this facilitates particle hydration when an aqueous medium is added to dried mixture. Secondly, where the hydrated particles are stored in a lyophilized state, as discussed below, the cryoprotectant may reduce particle damage which can occur on freezing, and thereby reduce size growth of the rehydrated particles. Suitable cryoprotectants are carbohydrates such as trehalose, lactose, maltose, cellobiose, sucrose, glucose, fructose, sorbitol, raffinose, myo-inositol, and glycerol (references 14). A variety of other water-soluble bulking agents, such as maltodextrin or salts may be substituted for the cryoprotectant where particle processing does not involve a freezing step, such as where the particles are dried for storing by spray drying.

The lipid solution of AMB/cholesterol sulfate is dried to a lipid film. As just indicated, the film is preferably formed from a solution containing a bulking agent, yielding dried particles of the agent coated with the lipid mixture. Solvent removal is by vacuum evaporation or under a stream of inert gas, e.g., nitrogen. The dried lipid film may be stored under an inert gas, preferably at 4° C or less.

An aqueous particle suspension is formed by addition of an aqueous medium to the dried lipid mixture. The medium used in Example 1, containing 10 mM Tris-HCl, 0.1 mM EDTA, pH 7.4, is suitable. The amount of medium added is sufficient to produce a final AMB concentration of preferably between 25-100 μ mole/ml.

Initially, the lipid material is crudely suspended by spatula or mechanical agitation until the lipid clumps are released into the aqueous medium, to form a slurry-like mixture. This material is now dispersed to a fine particle size by sonication, homogenization, French press or other high-energy input. The dispersion is carried out until a desired particle size, preferably between 0.1 to 1 μm (micron) is achieved. The suspension may be warmed during dispersion, and should be maintained under an inert atmosphere. In the method described in Example 1, the suspension is sonicated at 45°C to optical clarity. Final particle sizes were between 0.1-0.2 μm (microns). Here it is noted that formulations containing less than about 1 mole cholesterol sulfate per mole AMB do not sonicate to optical clarity, indicating that particle dispersion requires at least a stoichiometric amount of cholesterol sulfate.

The particle suspension may be treated such as by molecular sieve chromatography or dialysis, to remove traces of unincorporated AMB. The dialysis conditions noted in Example 1 are suitable. The final concentration of AMB in the dispersed particle suspension can be determined by diluting an aliquot of the suspension in methanol, and measuring AMB spectrophotometrically at 406 nm. Typical AMB concentrations at various stages of the preparation of the dispersion are given in Table 1 in Example 1 below.

B. Dried Particle Suspension

According to one aspect of the invention, it has been discovered that the AMB/cholesterol sulfate particles of the invention can be stored long-term in dried form without significant increase in particle size on rehydration.

The dried particle formulation can be prepared either by lyophilization or spray drying. In both cases, the drying step is performed before the AMB/cholesterol sulfate particles increase in size substantially. In the lyophilization method, the small-particle suspension is quick frozen and lyophilized at a shelf temperature of preferably 20°C or less, as described in Example 1. The effect of lyophilizing on particle size is seen in Table 2 in Example 2, for each of four formulations having AMB:cholesterol sulfate mole ratios between 1:1 and 1:4. In each case, mean particle sizes increased from 100-200 nm before lyophilization, to between 200-300 after lyophilization and rehydration with water. The stability of the particles, pre and post lyophilization is considered in Section II below.

For spray drying, the particle suspension is dried in a conventional apparatus in which the particles to be dried are sprayed in aerosolized suspension form into a stream of heated air or inert gas, and the aerosolized droplets are dried in the gas stream as they are carried toward a plate collector where the dried liposomes are collected. An exemplary spray dry apparatus is a Buchi 190 Mini Spray Dryer.

The drying temperature is at least 37°C, and preferably between 40-50°C. The temperature of the collection chamber is generally lower than that of the heated air, and typically 37°C. The dried particles are collected and stored in dehydrated form, under an inert atmosphere.

II. Size Stability

This section examines the size stability of AMB:cholesterol particle suspensions under a variety of conditions relating to molar composition of the particles, suspension medium, and storage time.

In a first study, reported in Example 3, AMB:cholesterol sulfate particles having mole ratios of AMB:cholesterol sulfate of 1:1, 1:2, 1:3, and 1:4 were prepared and immediately after dialysis were stored for periods of up to 8 days at 4°C. The results are shown in Table 3 of Example 3. The 1:1 formulation was substantially stable over the 8-day test period, whereas the other formulations showed progressively greater size increases with increasing mole ratios of cholesterol sulfate.

The size stability of the same four formulations after lyophilization and rehydration was similarly studied, also as reported in Example 3. Size stability data for the eight day test is shown in Table 3. Interestingly, there was little difference in size stability among the four formulations, and for each formulation mean particle sizes increased at most about 2 fold over the eight day test period. The combined results from Tables 2 and 4 demonstrate that (a) lyophilized AMB/cholesterol particles can be reconstituted with little increase in mean size and size distribution and (b) the particles in the reconstituted suspension are relatively stable on storage in solution over a several-day period.

The effect of physiological-strength saline and plasma on the size characteristics of the particles was also examined, as reported in Example 4. In a first study, the four post-dialysis AMB/cholesterol sulfate formulations from above were diluted in 0.9% saline, and the particle sizes examined immediately thereafter. As shown in at the top row in Table 5, all of the particles showed a large size increase, although the 1:1 formulation was less aggregated. A similar study on post-lyophilisation particles was also carried out, with the results shown in the top row of Table 6 in Example 4. A comparison of the Table 5 and 6 data

shows that the 1:1 formulation is substantially more size stable in saline after lyophilization than post-dialysis. The other three formulations, having greater cholesterol sulfate mole ratios, showed large size increases in saline both pre and post lyophilization.

A second study was designed to examine AMB/cholesterol sulfate particle size in blood plasma, and the effect of subsequent dilution of the plasma medium with suspending buffer. Initially, each of the four samples (both pre and post lyophilization) were diluted 1:1 with human plasma, then diluted after a few minutes with suspending buffer containing 10% lactose. Size measurements were made immediately after dilution, and again 20 minutes later. The results are shown in the bottom two rows of Tables 5 and 6 in Example 4. Summarizing the data, plasma caused a size increase in all of the formulations. Smallest size increases were seen in the 1:1 formulation, where particle sizes were less than 1 μm (micron) i.e. 1,000 nm. The size increase produced on contact with plasma was at least partially reversible for all formulations except the 1:4 formulation, as evidenced by the a significant reduction in particle size after 20 minutes incubation in dilute form in suspension medium. There was little difference in the size behavior of particles in pre- and post-lyophilization formulations.

The data above demonstrate that the AMB:cholesterol sulfate formulation of the invention can be stored in dried form long term, without significant increase in size, on rehydration, or significant change in size stability in plasma. One significant advantage of the dried particles which was observed was substantially greater size stability on storage in buffer. Within the range of AMB:cholesterol sulfate mole ratios which was examined, the 1:1 formulation, gave greatest size stability and smallest mean particle sizes under the various conditions examined.

III. Particle Characteristics

It has been reported that cholesterol sulfate is capable of forming lipid vesicles or liposomes on extended (several hour) sonication (reference 13). It was therefore of interest to determine whether the AMB μ cholesterol sulfate particles of the present invention are liposomal in form. For these studies, the 1:4 AMB/cholesterol sulfate formulation was selected, since a relatively high ratio of cholesterol sulfate is more likely to form liposomal structures.

One characteristic of liposomes is a continuous lipid bilayer capable of encapsulating water-soluble solute molecules. Many water-soluble molecules, such as sugars and other marker solutes, are readily encapsulated in liposomes by preparing (dispersing) the liposomal lipids in an aqueous medium containing the marker solute. Smaller marker molecules, such as sugars, also tend to pass through lipid bilayer membranes slowly, as evidenced by equilibration of the solute between encapsulated and bulk phase aqueous compartments over a several-hour to several-day solute-exchange period.

To test the ability of AMB/cholesterol sulfate (1:4) particles to encapsulate sucrose, the particles were prepared by dispersion in a medium containing ^{14}C sucrose. After sonication to optical clarity, the particles were separated from the suspending medium by molecular sieve chromatography, using a column sieving material which excludes particles in the size range of the AMB/cholesterol sulfate particles. Details of the test are given in Example 5. Briefly, 95% of the AMB was associated with the particles eluted in the void volume, but no detectable peak of radioactivity was associated with the particles.

Based on this study, it appears that the particles do not form encapsulating (liposomal) structures, or alternatively, that the particles form very leaky structures. The latter explanation is unlikely, since (a) cholesterol tends to decrease permeability in liposomes to small water-soluble permeants, and (b) the pure cholesterol derivative liposomes which have been described (reference 13) have very low permeability. Studies on cholesterol hemisuccinate liposomes also show stable encapsulation of a variety of small water-soluble molecules (PCT Patent application WO-A-8504578)

Another characteristic feature of liposomes is the ability of isotonic liposomes to swell on injection into a hypotonic medium. Here the liposomes are acting as small osmometers in response solute gradients across the bilayer membranes. Isotonic liposome swelling has been observed in liposomes prepared from a variety of cholesterol derivatives, including cholesterol-PEG and cholesterol sulfate (reference 13) and cholesterol hemisuccinate liposomes (PCT patent application WO-A-8504578). Cholesterol-derivative liposomes show the expected increased absorbance when injected into increasingly dilute media, although these liposomes behave less like ideal osmometers than do liposomes formed from conventional phospholipid components.

Each of the above four AMB/cholesterol particle compositions from above (1:1, 1:2, 1:3, and 1:4 mole ratios) was prepared in 10% lactose. Both pre- and post-dialysis particles were tested for osmotic swelling in distilled water, comparing particle size immediately after dilution with particle size 20 minutes after dilution. The results are shown in Table 7 in Example 6. No swelling was observed in any of the particle formulations. The test supports the finding from the encapsulation studies above that the AMB/cholesterol

sulfate particles of the invention do not form closed vesicle structures.

IV. Therapeutic Uses

5 AMB is useful in treating a variety of systemic fungal organisms, including coccidiomycosis, cryptococcosis, systemic moniliasis, histoplasmosis, aspergillosis, rhodoptorulosis, sporotrichosis, phycomycosis, and blastomycosis, and is also effective against some species of *Leishmania* (reference 15). Because of the severe side effects of the drug, in its presently available free form, it is generally administered only to patients with progressive, potentially fatal systemic fungal infections, and the patient must remain hospital-
10 ized during drug treatment for constant monitoring of renal function.

This section describes the decreased toxicity and improved efficacy of AMB in the present formulation, and the potential therefore of wider drug use, particularly for prophylactic use in preventing opportunistic fungal infection in immune-deficient or immune-compromised patients, such as those receiving cancer chemotherapy, immunosuppressive drugs, or radiation therapy.

15

A. AMB/Cholesterol Sulfate Toxicity

According to an important feature of the invention, it has been discovered that the AMB:cholesterol sulfate composition described herein is substantially less toxic than free AMB (Fungizone), as evidenced by a much higher LD₅₀ value. Further, the composition is considerably less toxic than liposomal or lipid-complex forms of AMB which have been described in the prior art, as judged by a comparison with reported LD₅₀ values. Toxicity studies to determine LD₅₀ values for the composition of the invention are detailed in Example 7. An initial test examined the lethal toxicity of Fungazone and 1:4 AMB/cholesterol sulfate particles. Based on the data shown in Table 8, the LD₅₀ of the free AMB (Fungizone) composition is
20 between 1-4 mg/kg animal weight. This value is increased to between 15-25 mg/kg in the AMB/cholesterol sulfate composition. LD₅₀ value is substantially higher than values previously reported for liposomal or lipid-complex AMB formulations described in the prior art.

In additional tests, also reported in Example 8, the lethal toxicity of the four different mole ratio AMB formulation described above, at a dose of up to 40 mg/kg was investigated. As reported in the example, the
30 AMB/CHSO₄ formulation gave an LD₅₀ value of up to 40 mg/kg. The data show that the LD₅₀ for the other three formulations is less than 20 mg/kg, and at least for the 1:4 formulation, therefore between 15-20 mg/kg.

The composition also enjoys the advantage that cholesterol sulfate is a natural cholesterol component found widely in animals. The cholesterol compound has no known toxicity, and is metabolized in the body
35 by cholesterol sulfatase.

Finally, since toxicity is expected to increase with increased particle size, the stable and relatively small particles which can be injected may contribute to reduced toxicity.

B. Efficacy

40

Another important feature of the AMB/cholesterol sulfate composition is significantly enhanced drug efficacy in treating systemic fungal infection. One efficacy study performed in support of the present invention is detailed in Example 9. Here animals infected intravenously with *C. albicans* were treated with Fungizone, at doses between 0.3 and 0.9 mg/kg body weight, or with 1:4 AMB:cholesterol sulfate, at doses between 0.3
45 and 2.0 mg/kg. Drug efficacy was determined by survival at 25 days post-drug administration. The data, presented in Table 11 show that the AMB/cholesterol sulfate composition gives significantly higher survival rates at each dose level between 0.3 and 0.9 mg/kg. At 2.0 mg/kg, all of the animals treated with the formulation of the invention survived.

A second efficacy study, also reported in Example 9, compared percent survival in *C. albicans*-infected
50 mice at dose levels 0.031, 0.125, and 0.5 mg/kg, with each animal receiving two doses on the day of infection, and two doses on the following day. The data are plotted in Figures 1A-1C for control (circles), free AMB (squares), AMB/CHSO₄ (1:4) (triangles), and AMB/CHSO₄ (1:1) (diamonds), at each of the three different dose levels. At the two lower dose levels (Figures 1B and 1C), the two AMB/CHSO₄ formulations were comparable in treatment efficacy to free AMB. At the highest dose level (Figure 1A), the 1:1
55 AMB/CHSO₄ formulation gave 100 percent survival, compared with 60-80% survival for either free AMB or the 1:4 AMB/CHSO₄ formulation.

C. Modes of Administration

The present invention provides a dehydrated AMB composition which, when rehydrated after an extended storage period, forms a suspension of AMB particles having a selected size range less than about 1 micron. Because the particles can be stored in an anhydrous, inert environment, toxicity and lipid and drug breakdown problems related to oxidation and mechanical damage at a gas/liquid interface are minimized. For parenteral use, e.g., intravenous administration, the composition is preferably formed from AMB liposomes having sizes of between 0.1 to 0.4 μm (microns), such as can be prepared by the methods above. The AMB/lipid composition is hydrated typically to a selected AMB concentration between about 50 and 200 mg/ml, and administered at a concentration of between 1 and 5 mg AMB/kg body weight. Administration is carried out before the AMB/CHSO₄ particles increase in size substantially.

Where the drug is given intramuscularly, to provide slow drug release from the site of injection, the composition is preferably rehydrated to a more concentrated form, which can be conveniently localized at an injection site.

From the foregoing, it can be appreciated how various objects and features of the invention are met. The invention provides an AMB formulation which has substantially reduced toxicity with respect to prior art free AMB or AMB lipid formulations, and drug efficacy which is at least as great as reported prior-art AMB formulations, including free AMB. The enhanced therapeutic index of the drug, particularly related to reduced toxicity, allows much broader use of the drug, for example, for prophylactic treatment of immune-compromised patients, and also provides greater therapeutic efficacy, in the treatment of active systemic fungal infections.

The composition is readily prepared, and the cholesterol sulfate component is relatively inexpensive in purified form, and naturally utilized when administered parenterally. The formulation is easily stored in dried form, and when rehydrated, yields a particle suspension with selected small sizes.

The following examples illustrate methods of preparing, characterizing, and using the AMB/cholesterol sulfate composition of the invention. The examples are in no way intended to limit the scope of the invention.

Materials

CHSO₄ (cholesterol 3-sulphate, sodium salt) was obtained from Sigma Chemical Co., St. Louis, MO; AMB (amphotericin-B, Type 1) was donated by E.R. Squibb & Sons, Inc. (Batch No. 20-914-5978-001); All other materials are of reagent grade or superior from commercial sources.

35 Example 1Preparation of AMB/CHSO₄ Particles

AMB and CHSO₄ in dry powder form were weighed out and combined to give one of the four AMB:cholesterol sulfate mole ratios listed in Table 1 below. The amount of AMB and cholesterol sulfate added was sufficient to produce a final AMB plus cholesterol sulfate concentration in the particle suspension of 50 $\mu\text{mole/ml}$.

Dry methanol was added to the AMB/cholesterol sulfate powder to a final AMB concentration of between 0.2-0.6 mg/ml, and the suspension was stirred until all of the powder dissolved. Lactose was added to this solution to produce a 10% (w/v) lactose solution in the final aqueous product. The solution was dried in vacuo, yielding dried lactose particles coated with a lipophilic AMB/cholesterol sulfate film.

A suspending buffer containing 10 mM Tris-HCl, 0.1 mM EDTA pH 7.4, 67 mOsm, was added to the dried mixture in an amount sufficient to produce a final AMB plus cholesterol sulfate concentration of 50 $\mu\text{mole/ml}$. This suspension was sonicated with an Ultrasonic Liquid Processor (Heat Ultrasonics, Inc., Farmingdale, NY), Model W-800, probe sonicator until the suspension becomes optically clear. (This process is facilitated if the suspension is warmed to 45° C in a water bath.) Sonication was performed under nitrogen gas.

The sonicated AMB/CHSO₄ particles were dialyzed to remove traces of unincorporated AMB, using 6000-8000 molecular weight cut-off dialysis tubing. The material was dialyzed against a buffer containing 10 mM Tris-HCl, 0.1 mM EDTA, 10% (w/v) lactose, pH 7.4, 300 mOsm. The clear suspension was dried by rapid freezing in a dry ice/isopropanol mixture and lyophilized overnight at a shelf temperature of -25° C, followed by a further two hours at 25° C (15 SRC-X Lyophilizer; Virtis, Gardiner, NY). Lyophilized samples were reconstituted by addition of an equal volume of water and gentle mixing. Table 1 below shows the

AMB concentrations of the four compositions, at various stages of preparation.

Table 1

AMB Concentration (mg/ml)				
Molar Ratio	Theoretical	Pre-Dialysis	Post-Dialysis	Post-Lyophilization/Rehydration
1:1	23.10	20.66	24.20	21.52
1:2	15.40	10.80	16.46	15.19
1:3	11.55	11.02	12.19	11.60
1:4	9.24	9.49	10.12	8.79

Example 2

Effect of Lyophilization on Particle Size

Particle sizes were determined by dynamic laser-light scattering using a Nicomp Model 200 sizer (Nicomp Instruments Inc., Goleta, CA). Samples were typically diluted to 0.3 μ mole/ml for this measurement using 10 mM Tris/HCl, 0.1 mM EDTA, 10% (w/v) lactose buffer, pH 7.4. The mean particle sizes and standard deviations (S.D.) for the four compositions from Example 1 are given in Table 2 below. As seen, all four compositions have mean particle sizes between 130-180 nm prior to lyophilization, and between 210-280 nm after lyophilization.

Table 2

Particle Diameter (mean $\pm$ S.D. nm)		
Molar Ratio AMB/CHSO ₄	Particles Immediately Post-Dialysis	Particles Post-Lyophilization/Rehydration
1:1	138 $\pm$ 54	268 $\pm$ 137
1:2	148 $\pm$ 66	211 $\pm$ 105
1:3	172 $\pm$ 89	265 $\pm$ 147
1:4	138 $\pm$ 61	274 $\pm$ 151

Example 3

Effect of Storage in Solution on Particle Size

The four samples from Example 1, each containing an AMB plus cholesterol sulfate concentration of about 50 μ mole/ml, were incubated at 4° C for up to eight days. At days 0, 2, 6, and 8, an aliquot of the each suspension was withdrawn, diluted to about 0.3 μ mole/ml, and examined for particle size distribution, as in Example 2. The results are shown in Table 2 below. It is seen that the 1:1 composition is stable to particle size change, whereas the compositions containing higher molar amounts of cholesterol sulfate are progressively less stable on storage.

Table 3

Days of Storage	Particle Diameter (mean $\pm$ S.D. nm) as a Function of AMB/CHSO ₄ / Molar Ratio (Post-Dialysis)			
	1:1	1:2	1:3	1:4
0	138 $\pm$ 54	148 $\pm$ 66	172 $\pm$ 89	138 $\pm$ 61
2	193 $\pm$ 98	336 $\pm$ 194	527 $\pm$ 299	334 $\pm$ 192
6	161 $\pm$ 82	462 $\pm$ 265	679 $\pm$ 377	823 $\pm$ 481
8	179 $\pm$ 95	512 $\pm$ 296	968 $\pm$ 551	1034 $\pm$ 587

A similar stability study was performed on the same compositions after lyophilization and reconstitution in distilled water, as in Example 1. The results, given in Table 4, show (a) relatively small size increases over the eight day test for each of the four compositions, and (b) little effect on size changes of the molar amount of cholesterol sulfate.

Table 4

Days of Storage	Particle Size (mean $\pm$ S.D. nm) as a Function of AMB/CHSO ₄ Molar Ratio (Post-Lyophilization)			
	1:1	1:2	1:3	1:4
0	268 $\pm$ 137	211 $\pm$ 105	265 $\pm$ 147	279 $\pm$ 151
2	304 $\pm$ 153	331 $\pm$ 177	541 $\pm$ 293	554 $\pm$ 303
6	340 $\pm$ 171	418 $\pm$ 224	515 $\pm$ 283	478 $\pm$ 242
8	371 $\pm$ 188	436 $\pm$ 228	570 $\pm$ 314	487 $\pm$ 258

Example 4

Effect of Saline and Plasma on Particle Size

The four samples from Example 1 were diluted to approximately 0.3 μ mole/ml with 0.9% (w/v) saline and their sizes measured as in Example 2. The results are shown at the top line in Table 5 below. For each formulation, saline produced a more than tenfold increase in mean particle size. The size growth of the 1:1 composition was substantially less than for the three compositions with greater amounts of cholesterol sulfate.

The four samples were also diluted 1:1 (v/v) with human plasma and subsequently (within a few minutes of contact with the plasma) diluted with 10 mM Tris/HCl, 0.1 mM EDTA, 10% lactose (w/v) buffer pH 7.4, for sizing. Size measurements, reported in Table 5 below, were made immediately after diluting, and 20 minutes after diluting. The data indicate that the 1:1 formulation is least sensitive to size change, upon contact with plasma, and that for all formulations, incubation in the diluted medium for 20 minutes produced some size decrease.

Table 5

Treatment	Particle Size (mean $\pm$ S.D. nm) as a Function of AMB/CHSO ₄ Molar Ratio (Post-Dialysis)			
	1:1	1:2	1:3	1:4
Dilute in saline	3328 $\pm$ 1990	13209 $\pm$ 8288	6789 $\pm$ 4276	11072 $\pm$ 6618
Mix + plasma, dilute + suspending buffer	728 $\pm$ 334	1200 $\pm$ 628	4017 $\pm$ 1951	1942 $\pm$ 1043
20 minutes later	358 $\pm$ 169	823 $\pm$ 457	1745 $\pm$ 1007	1667 $\pm$ 942

Similar size measurements, after mixing with 0.9 % saline or plasma, were made on AMB/cholesterol sulfate particles after lyophilization and rehydration with distilled water, as in Example 1. The results are shown in Table 6 below. Size changes similar to those observed with pre-lyophilized particles (Table 5 data) were observed.

Table 6

Treatment	Particle Size (mean $\pm$ S.D. nm) as a Function of AMB/CHSO ₄ Molar Ratio (Post-Lyophilization/Hydration)			
	1:1	1:2	1:3	1:4
Dilute in saline	1738 $\pm$ 1007	8405 $\pm$ 5549	13158 $\pm$ 8625	9813 $\pm$ 6265
Mix + plasma, dilute + suspending buffer	955 $\pm$ 446	1030 $\pm$ 550	1766 $\pm$ 976	2467 $\pm$ 1233
20 minutes later	534 $\pm$ 267	776 $\pm$ 378	1147 $\pm$ 643	2661 $\pm$ 1374

Example 5

Particle Encapsulation Studies

The ability of the AMB/cholesterol sulfate particles to encapsulate a radiolabeled marker was examined. CHSO₄/AMB, 4:1 molar ratio particles, were prepared as in Example 1, except that the Tris buffer medium used to suspend the dried AMB/cholesterol sulfate mix contained 1 μ Ci of ¹⁴C-sucrose. The suspension was applied to a Sephadex G50 gel exclusion column equilibrated with 10 mM Tris/HCl, 0.1 mM EDTA, 10% (w/v) lactose buffer, pH 7.4, and the applied material was eluted with the same buffer. The particles were eluted in the void volume, which was monitored by UV absorption at 280 nm. The samples were collected and examined for radioactivity by conventional scintillation counting. It was found that 95% of the AMB was in the particle peak, whereas no detectable peak of ¹⁴C-sucrose occurred in this region.

Example 6

Osmotic Swelling Studies

CHSO₄/AMB formulations containing the four different mole ratios of AMB and cholesterol sulfate were prepared as in Example 1, (in the usual suspension medium containing 10% lactose). These samples are designated as post-dialysis (P.D.) suspensions in the Table 7 below. A portion of each sample (containing 10% lactose) was lyophilized and reconstituted in distilled water, and these samples are designated as lyophilized and reconstituted (L.R.) in the table.

The P.D. and L.R. samples were each diluted to 0.3 μ mole/ml with distilled water, and the size distribution of the particles immediately after dilution in the hypotonic medium, and 20 minutes after dilution was measured as in Example 2. The results are given in Table 7 below. As seen, there is no appreciable swelling, over a 20 minute incubation period, as evidenced by an increase in mean particle size, in any of the samples examined.

Table 7

	Molar Ratio AMB/ChSO ₄	Sample Post-Dialysis (PD) or Post-Lyophilization/Rehydration (LR)	Time (min)	Particle Size (mean $\pm$ S.D. nm)
5	1:1	PD	0 20	109 $\pm$ 42 114 $\pm$ 45
10	1:1	LR	0 20	212 $\pm$ 107 212 $\pm$ 105
	1:2	PD	0 20	126 $\pm$ 59 134 $\pm$ 64
15	1:2	LR	0 20	170 $\pm$ 81 184 $\pm$ 89
	1:3	PD	0 20	124 $\pm$ 58 131 $\pm$ 61
20	1:3	LR	0 20	205 $\pm$ 117 204 $\pm$ 107
	1:4	PD	0 20	161 $\pm$ 86 185 $\pm$ 100
25	1:4	LR	0 20	228 $\pm$ 116 236 $\pm$ 122

Example 7Toxicity (LD₅₀) of the Particle Suspensions

Outbred male Swiss/Webster mice were obtained from Simonsen Labs, Inc. The animals weighed approximately 15-45 grams on the day of treatment and were between 4-8 weeks old. The animals were quarantined for at least three days prior to the study, and only mice that remained healthy during the quarantine period were used. The animals were given food and water ad libitum

In a first study, the animal groups were treated with either Fungizone (Squibb) suspended in sterile saline or a 1:4 AMB/cholesterol sulfate composition prepared as in Example 1. In each case, the AMB concentration was adjusted such that the selected dose of AMB (given in Table 8) could be administered in a final volume of 0.2 ml. Four-eight animals were employed for each dose group. The test material was administered by a single intravenous injection via the lateral tail vein. Each dose was administered over about 1.5 minutes.

The animals were observed for signs of toxicity and death at least three times (1, 2, and 4 hour post treatment) on the day of treatment. During the remaining observation period of five days, the animals were examined daily in the morning and afternoon. The test results, expressed as the ratio of number of survivors on day five:total number of animals treated, are given in Table 8. The LD₅₀ value for Fungizone, calculated by conventional methods, is 3.2 mg/kg. The LD₅₀ for the AMB/cholesterol sulfate composition between 15-20 mg/KG.

Table 8

Treatment
 Number of Survivors on Day 5
 (Post-Injection/Total Animals Injected)

FUNGIZONE

0.5 mg/Kg	8/8
1.0 mg/Kg	8/8
2.0 mg/Kg	8/8
4.0 mg/Kg	2/8
6.0 mg/Kg	0/8
8.0 mg/Kg	0/4

CHSO₄/AMB (4:1)

10 mg/Kg	3/3
15 mg/Kg	3/4
20 mg/Kg	2/5
25 mg/Kg	1/3

In a second toxicity test, mice were treated with 20mg/kg of one of the four AMB/cholesterol sulfate formulations from Example 1, with drug administration and animal monitoring being done as above. The results, presented below in Table 9, show that the 1:1 formulation has an LD₅₀ value higher than 20 mg/kg.

Table 9

Treatment (AMB/CHSO ₄)	Number of Survivors on Day 5 (Post-Injection/Total Animals Injected)
1:1 molar ratio	6/6
1:2 molar ratio	0/5
1:3 molar ratio	1/5
1:4 molar ratio	0/5

A third toxicity test was conducted by an independent laboratory, using male Crl:Cb-I(ICR)BR mice, 18-20 grams, obtained from Charles River Laboratory (Wilmington, MA). AMB/CHSO₄ formulations containing either 1:4 or 1:1 mole ratios of AMB:cholesterol sulfate were prepared as above. Fungizone™ and free AMB were solubilized in pyrogen-free water and sonicated for 10 minutes at room temperature. All drug preparations were diluted to the desired concentration with a suitable dilution buffer.

Uninfected mice received a single dose of one of the AMB preparations by slow bolus, intravenous injection. Doses for the AMB/CHSO₄ cholesterol sulfate formulations were 30, 15, 7.5, 3.75, 1.88 and 0.94 mg AMB/kg body weight. Doses for Fungizone™ and free AMB were 8, 4, 2, 1, 0.5, and 0.25 mg AMB/kg body weight. Mortality was recorded for a period of 5 days following treatment, and the 50% lethal dose (LD₅₀) for each preparation was calculated by standard methods. The results are shown in Table 10. As seen, the LD₅₀ value of the 1:1 and 1:4 AMP/CHSO₄ formulations were several times higher than that of free AMB or Fungizone™, with the 1:1 formulation giving an LD₅₀ value of greater than 30 mg/kg.

Table 10

AMB Form	LD ₅₀ (mg/kg/dose) (95% Conf. Int.)
Free AMB	4.9 (3.7, 6.5)
Fungizone	4.3 (3.1, 6.0)
AMB/CHSO ₄ (4:1)	16.1 (10.9, 23.8)
AMB/CHSO ₄ (1:1)	>30

More recently, the toxicity of the 1:4 and 1:1 AMB/CHSO₄ formulations has been tested by another independent laboratory, in which bolus injections of up to 40 mg AMB/kg body weight were tested. The study found an LD₅₀ value for the 1:1 AMB/CHSO₄ formulation of greater than 40 mg/kg.

Example 8

Efficacy of the AMB/Cholesterol Sulfate Formulation

CrI:CFW(SW)BR mice weighing 20-25 grams were obtained from the Charles River Breeding Laboratories, and were given food and water ad libitum. *C. albicans* strain 30 was grown at 35° C on SDA (Sabourand Dextrose A gas) for 18 hours, and the organism is harvested and diluted with sterile nonpyrogenic saline to yield about 7×10^8 colony forming units in a 0.2 ml volume.

Eight-ten animals are injected in the tail vein each with 0.2 ml of the above *C. albicans* mixture. Two days after the fungal injection, the animals were injected with graded doses of Fungizone or AMB/cholesterol sulfate (1:4) prepared as in Example 1. The AMB preparations were adjusted in concentration so that each animal received a total volume, administered intravenously through the tail vein, of ml. The amount of AMB administered, expressed in terms of mg drug/kg body weight of the animal is given at the left in Table 10. The animals were followed for 25 days post-drug administration. The number of survivors at 25 days per total number of test animals is shown in the table for the two AMB preparations, and a buffer control.

Table 11

Dose	Survivors Post-25 Days/Total Animals Injected		
	Free AMB	CHSO ₄ /AMB = 4:1	Control
0.0 mg/Kg	-	-	0/11
0.3 mg/Kg	2/10	5/10	-
0.6 mg/Kg	1/10	9/10	-
0.9 mg/Kg	3/10	9/10	-
2.0 mg/Kg	-	10/10	-

In a second study, male CrI:CD-1(ICR)BR mice weighing 18-20 gm were infected with 0.1 ml *C. albicans* (10^6 cells/animal) and divided into treatment groups of ten animals/group. Free AMB and 1:1 and 1:4 AMB/CHSO₄ formulations were prepared as in Example 7 and Example 1. Infected mice received two doses of each drug preparation on the day of infection and 2 more doses on the following day, by intravenous route for a total of four doses. The single-dose AMB levels for each test group were 0.5, 0.125, and 0.031 mg AMB/kg body weight/dose. A control group received only the dilution buffer. Mice were observed for mortality for a period of 30 days following infection. The 50% protective dose (PD₅₀) was calculated from the survival data by conventional methods.

Table 12 summarizes the PD₅₀ data obtained for the two AMB/CHSO₄ formulations and free AMB. Both of the AMB/CHSO₄ formulations were as effective as free AMB in the treatment of the fungal infection.

Table 12

Amphotericin B Form	PD50 (mg/kg/dose) (95% Conf. Int.)
Free AMB	0.25 (0.11, 0.55)
AMB/CHSO ₄ (4:1)	0.25 (0.14, 0.45)
AMB/CHSO ₄ (1:1)	0.22 (0.16, 0.29)

The survival curves from the above study are presented in Figures 1A-1C, as discussed above. The curves show, in particular, the greater treatment efficacy over free AMB achievable with the 1:1 AMB/CHSO₄ formulation at higher AMB dose levels.

Although the invention has been described and illustrated with respect to specific embodiments, uses and methods of preparation, it will be appreciated that a variety of changes and modifications may be made without departing from the scope of the invention.

Claims

1. An amphotericin B composition comprising particles of amphotericin B containing cholesterol sulfate in a mole ratio of amphotericin B to cholesterol sulfate of between 1:1 to 1:4.
2. The composition of claim 1, wherein the particles are suspended in an aqueous medium, and have particle sizes between 100-400 nm.
3. The composition of claim 2, which is produced by the steps of:
 - (a) dispersing an aqueous suspension of amphotericin B and cholesterol sulfate, in a molar ratio of between 1:1 to 1:4, to optical clarity,
 - (b) lyophilizing the suspension in the presence of a cryoprotectant before the particle sizes in the suspension increase substantially, and
 - (c) reconstituting the lyophilized material with an aqueous medium.
4. The composition of claim 1, wherein the molar ratio of amphotericin B to cholesterol sulfate is about 1:1.
5. A composition according to any one of claims 1 to 4, for use in the treatment of fungal infections.
6. A method of preparing amphotericin B in a form which has an LD₅₀ greater than 15 mg/kg, and can be reconstituted with an aqueous medium, to particle sizes between 100-400 nm, comprising
 - (a) dispersing an aqueous suspension of amphotericin B and cholesterol sulfate, in a molar ratio of between 1:1 to 1:4, to optical clarity, and
 - (b) lyophilizing the suspension in the presence of a cryoprotectant, before the particles in the suspension increase substantially in size.
7. The method of claim 6, wherein the molar ratio of amphotericin B to cholesterol in the suspension is about 1:1.

Patentansprüche

1. Amphotericin B-Zusammensetzung, umfassend Partikel von Amphotericin B, die Cholesterinsulfat in einem Molverhältnis von Amphotericin B zu Cholesterinsulfat von 1:1 bis 1:4 enthalten.
2. Zusammensetzung nach Anspruch 1, worin die Partikel in einem wäßrigen Medium suspendiert sind und eine Partikelgröße zwischen 100-400 nm aufweisen.
3. Zusammensetzung nach Anspruch 2, die durch die Verfahrensschritte erhalten wurde:
 - (a) Dispergieren einer wäßrigen Suspension von Amphotericin B und Cholesterinsulfat in einem Molverhältnis von 1:1 bis 1:4 bis zur optischen Klärung,
 - (b) Lyophilisieren der Suspension in Gegenwart eines Kälteschutzmittels, ehe die Partikelgröße in der Suspension wesentlich zunimmt und



Table 12

Amphotericin B Form	PD50 (mg/kg/dose) (95% Conf. Int.)
Free AMB	0.25 (0.11, 0.55)
AMB/CHSO ₄ (4:1)	0.25 (0.14, 0.45)
AMB/CHSO ₄ (1:1)	0.22 (0.16, 0.29)

The survival curves from the above study are presented in Figures 1A-1C, as discussed above. The curves show, in particular, the greater treatment efficacy over free AMB achievable with the 1:1 AMB/CHSO₄ formulation at higher AMB dose levels.

Although the invention has been described and illustrated with respect to specific embodiments, uses and methods of preparation, it will be appreciated that a variety of changes and modifications may be made without departing from the scope of the invention.

Claims

1. An amphotericin B composition comprising particles of amphotericin B containing cholesterol sulfate in a mole ratio of amphotericin B to cholesterol sulfate of between 1:1 to 1:4.
2. The composition of claim 1, wherein the particles are suspended in an aqueous medium, and have particle sizes between 100-400 nm.
3. The composition of claim 2, which is produced by the steps of:
 - (a) dispersing an aqueous suspension of amphotericin B and cholesterol sulfate, in a molar ratio of between 1:1 to 1:4, to optical clarity,
 - (b) lyophilizing the suspension in the presence of a cryoprotectant before the particle sizes in the suspension increase substantially, and
 - (c) reconstituting the lyophilized material with an aqueous medium.
4. The composition of claim 1, wherein the molar ratio of amphotericin B to cholesterol sulfate is about 1:1.
5. A composition according to any one of claims 1 to 4, for use in the treatment of fungal infections.
6. A method of preparing amphotericin B in a form which has an LD₅₀ greater than 15 mg/kg, and can be reconstituted with an aqueous medium, to particle sizes between 100-400 nm, comprising
 - (a) dispersing an aqueous suspension of amphotericin B and cholesterol sulfate, in a molar ratio of between 1:1 to 1:4, to optical clarity, and
 - (b) lyophilizing the suspension in the presence of a cryoprotectant, before the particles in the suspension increase substantially in size.
7. The method of claim 6, wherein the molar ratio of amphotericin B to cholesterol in the suspension is about 1:1.

Patentansprüche

1. Amphotericin B-Zusammensetzung, umfassend Partikel von Amphotericin B, die Cholesterinsulfat in einem Molverhältnis von Amphotericin B zu Cholesterinsulfat von 1:1 bis 1:4 enthalten.
2. Zusammensetzung nach Anspruch 1, worin die Partikel in einem wäßrigen Medium suspendiert sind und eine Partikelgröße zwischen 100-400 nm aufweisen.
3. Zusammensetzung nach Anspruch 2, die durch die Verfahrensschritte erhalten wurde:
 - (a) Dispergieren einer wäßrigen Suspension von Amphotericin B und Cholesterinsulfat in einem Molverhältnis von 1:1 bis 1:4 bis zur optischen Klärung,
 - (b) Lyophilisieren der Suspension in Gegenwart eines Kälteschutzmittels, ehe die Partikelgröße in der Suspension wesentlich zunimmt und



(c) Rekonstituieren des lyophilisierten Materials mit einem wäßrigen Medium.

4. Zusammensetzung nach Anspruch 1, worin das Molverhältnis von Amphotericin B und Cholesterinsulfat etwa 1:1 beträgt.
5. Zusammensetzung nach einem der Ansprüche 1 bis 4 zur Verwendung bei der Behandlung von Pilzinfektionen.
6. Verfahren zur Herstellung von Amphotericin B in einer Form, die einen LD₅₀-Wert von größer als 15 mg/kg aufweist, und die mit einem wäßrigen Medium zu Partikeln mit einer Größe zwischen 100-400 nm rekonstituiert werden kann, umfassend
 - (a) Dispergieren einer wäßrigen Suspension von Amphotericin B und Cholesterinsulfat in einem Molverhältnis zwischen 1:1 bis 1:4, bis zur optischen Klärung und
 - (b) Lyophilisieren der Suspension in Gegenwart eines Kälteschutzmittels, ehe die Größe der Partikel in der Suspension wesentlich zunimmt.
7. Verfahren nach Anspruch 6, in dem das Molverhältnis von Amphotericin B zu Cholesterin in der Suspension etwa 1:1 beträgt.

20 Revendications

1. Composition d'amphotéricine B comprenant des particules d'amphotéricine B contenant du sulfate de cholestérol, dans une proportion molaire de l'amphotéricine B au sulfate de cholestérol entre 1:1 et 1:4.
2. Composition selon la revendication 1, dans laquelle les particules sont en suspension dans un milieu aqueux et ont des dimensions entre 100 et 400 nm.
3. Composition selon la revendication 2, qui est produite selon les étapes de :
 - (a) dispersion d'une suspension aqueuse d'amphotéricine B et de sulfate de cholestérol dans une proportion molaire entre 1:1 et 1:4 jusqu'à limpidité optique,
 - (b) lyophilisation de la suspension en présence d'un cryoprotecteur avant que les dimensions des particules dans la suspension n'augmentent sensiblement et
 - (c) reconstitution de la matière lyophilisée avec un milieu aqueux.
4. Composition selon la revendication 1, dans laquelle la proportion molaire de l'amphotéricine B au sulfate de cholestérol est d'environ 1:1.
5. Composition selon l'une quelconque des revendications 1 à 4, pour l'utilisation dans le traitement des infections fongiques.
6. Procédé de préparation d'amphotéricine B, sous une forme ayant une DL₅₀ supérieure à 15 mg/kg et qui peut être reconstituée avec un milieu aqueux, avec des dimensions des particules entre 100 et 400 nm, comprenant
 - (a) la dispersion d'une suspension aqueuse d'amphotéricine B et de sulfate de cholestérol dans une proportion molaire entre 1:1 et 1:4 jusqu'à limpidité optique et
 - (b) la lyophilisation de la suspension en présence d'un cryoprotecteur avant que les dimensions des particules dans la suspension n'augmentent sensiblement.
7. Procédé selon la revendication 6, dans lequel la proportion molaire de l'amphotéricine B au cholestérol dans la suspension est d'environ 1:1.