

CLAIMS

What is claimed is:

1. A method of preparing an abuse deterrent pharmaceutical composition
5 comprising a drug-containing core enclosed by a layer comprising one or more metal oxide materials, the method comprising the sequential steps of:
 - (a) loading the particles comprising the drug into a reactor;
 - (b) applying a vaporous or gaseous metal precursor to the particles in the reactor;
 - (c) performing one or more pump-purge cycles of the reactor using inert gas;
 - 10 (d) applying a vaporous or gaseous oxidant to the particles in the reactor; and
 - (e) performing one or more pump-purge cycles of the reactor using inert gas;wherein the, thereby producing an abuse deterrent pharmaceutical composition comprising a drug containing core enclosed by a layer comprising one or more metal oxide materials.
15
2. The method of claim 1, wherein the sequential steps (b)-(e) are repeated one or more times to increase the total thickness of the layer comprising one or more metal oxide materials that enclose the core.
- 20 3. The method of claim 1, wherein the reactor pressure is allowed to stabilize following step (a), step (b), and/or step (d).
4. The method of claim 1, wherein the reactor contents are agitated prior to and/or during step (b), step (c), and/or step (e).
25
5. The method of claim 1, wherein a subset of vapor or gaseous content is pumped out prior to step (c) and/or step (e).
6. The method of claim 1, wherein the metal oxide layer has a thickness in range
30 of 0.1 nm to 100 nm.
7. The method of claim 1, wherein the particles comprise a drug and one or more pharmaceutically acceptable excipients.

8. The method of claim 1, wherein the particles have a median particle size, on a volume average basis between 0.1 μm and 1000 μm .

9. The method of claim 1, wherein the pharmaceutical composition is removed
5 from the reactor and admixed with a pharmaceutically acceptable diluent or carrier.

10. The method of claim 1, wherein the particles consist essentially of the drug.

11. The method of claim 1, wherein the drug is an opioid.
10

12. An abuse deterrent pharmaceutical composition comprising a drug-containing core enclosed by one or more metal oxide materials, the method comprising the sequential steps of:

(a) loading particles comprising the drug into a reactor;
15 (b) applying a vaporous or gaseous metal precursor to the particles in the reactor;
(c) performing one or more pump-purge cycles of the reactor using inert gas;
(d) applying a vaporous or gaseous oxidant to the particles in the reactor; and
(e) performing one or more pump-purge cycles of the reactor using inert gas;
thereby producing an abuse deterrent pharmaceutical composition comprising a drug
20 containing core enclosed by one layer comprising a metal oxide materials.

13. The composition of claim 12, wherein the sequential steps (b)-(e) are repeated one or more times to increase the total thickness of the one or more metal oxide materials that enclose the core.
25

14. The composition of claim 12, wherein the reactor pressure is allowed to stabilize following step (a), step (b), and/or step (d).

15. The composition of claim 12, wherein the reactor contents are agitated prior
30 and/or during to step (b), step (c), and/or step (e).

16. The composition of claim 12, wherein a subset of vapor or gaseous content is pumped out prior to step (c) and/or step (e).

17. The composition of claim 12, wherein the metal oxide layer has a thickness in range of 0.1 nm to 100 nm.

18. The composition of claim 12, wherein the particles comprise a drug and one or
5 more pharmaceutically acceptable excipients.

19. The composition of claim 12, wherein the particles have a median particle size, on a volume average basis between 0.1 μm and 1000 μm .

20. The composition of claim 12, wherein the pharmaceutical composition is removed from the reactor and admixed with a pharmaceutically acceptable diluent or carrier.

21. The composition of claim 12, wherein the core consists essentially of the drug.

22. The composition of claim 12, wherein the drug is a small molecule, virus particle, polypeptide, polynucleotide, a composition comprising polypeptide and lipid, or a composition comprising polynucleotide and lipid.

23. The method of claim 1, wherein the layer is less than 200 nanometers thick.

20

24. The method of claim 1, wherein the layer is less than 100 nanometers thick.

25. The method of claim 1, wherein the layer is less than 50 nanometers thick.

26. The method of claim 1, wherein the layer is less than 10 nanometers thick.

25

27. The method of claim 1, wherein the particle have a median particle size, on a volume average basis, of less than 10 microns.

28. The method of claim 1, wherein the particle have a median particle size, on a volume average basis, of less than 1 micron

30

29. The method of claim 1, wherein the particle have a median particle size, on a volume average basis, of less than 100 nanometers.