7. The compound according to claim 1, wherein the compound is a (R)(-)-stereoisomer.

8. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of the compound according to claim 1.

9. Use of therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for treating a disorder selected from the group consisting attention deficit hyperactivity disorder (ADHD), cognition impairment, anxiety disorders, generalized anxiety disorder (GAD), panic disorder, bipolar disorder or manic depression or manic-depressive disorder, obsessive compulsive disorder (OCD), posttraumatic stress disorder (PTSD), acute stress disorder, social phobia, simple phobias, pre-menstrual dysphoric disorder (PMDD), social anxiety disorder (SAD), major depressive disorder (MDD), postnatal depression, dysthymia, depression associated with Alzheimer's disease, Parkinson's disease, or psychosis, supranuclear palsy, eating disorders, obesity, anorexia nervosa, bulimia nervosa, binge eating disorder, diabetes, ischemic diseases, pain, substance abuse disorders, chemical dependencies, nicotine addiction, cocaine addiction, amphetamine addiction, alcohol addiction, Lesch-Nyhan syndrome, neurodegenerative diseases, Parkinson's disease, late luteal phase syndrome or narcolepsy, psychiatric symptoms, anger, rejection sensitivity, movement disorders, extrapyramidal syndrome, Tic disorders, restless leg syndrome (RLS), tardive dyskinesia, ~~supranuclear palsy,~~ sleep related eating disorder (SRED), night eating syndrome (NES), stress urinary incontinence (SUI), migraine, neuropathic pain, diabetic neuropathy, lower back pain, fibromyalgia syndrome (FS), osteoarthritis pain, arthritis pain, chronic fatigue syndrome (CFS), sexual dysfunction, premature ejaculation, male impotence, thermoregulatory disorders (e.g., hot flashes associated with menopause), and irritable bowel syndrome (IBS).

10. Use of a compound for the manufacture of a medicament according to claim 9 further comprising:

a therapeutically effective amount of a serotonin 1A receptor antagonist or a pharmaceutically acceptable salt thereof.

7. The compound according to claim 1, wherein the compound is a (R)(-)-stereoisomer.

8. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of the compound according to claim 1.

9. Use of therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for treating a disorder selected from the group consisting of attention deficit hyperactivity disorder (ADHD), cognition impairment, anxiety disorders, generalized anxiety disorder (GAD), panic disorder, bipolar disorder or manic depression or manic-depressive disorder, obsessive compulsive disorder (OCD), posttraumatic stress disorder (PTSD), acute stress disorder, social phobia, simple phobias, pre-menstrual dysphoric disorder (PMDD), social anxiety disorder (SAD), major depressive disorder (MDD), postnatal depression, dysthymia, depression associated with Alzheimer's disease, Parkinson's disease, or psychosis, supranuclear palsy, eating disorders, obesity, anorexia nervosa, bulimia nervosa, binge eating disorder, diabetes, ischemic diseases, pain, substance abuse disorders, chemical dependencies, nicotine addiction, cocaine addiction, amphetamine addiction, alcohol addiction, Lesch-Nyhan syndrome, neurodegenerative diseases, Parkinson's disease, late luteal phase syndrome or narcolepsy, psychiatric symptoms, anger, rejection sensitivity, movement disorders, extrapyramidal syndrome, Tic disorders, restless leg syndrome (RLS), tardive dyskinesia, sleep related eating disorder (SRED), night eating syndrome (NES), stress urinary incontinence (SUI), migraine, neuropathic pain, diabetic neuropathy, lower back pain, fibromyalgia syndrome (FS), osteoarthritis pain, arthritis pain, chronic fatigue syndrome (CFS), sexual dysfunction, premature ejaculation, male impotence, thermoregulatory disorders (e.g., hot flashes associated with menopause), and irritable bowel syndrome (IBS).

10. Use of a compound for the manufacture of a medicament according to claim 9 further comprising:

a therapeutically effective amount of a serotonin 1A receptor antagonist or a pharmaceutically acceptable salt thereof.