

ANDROGEN RECEPTOR MODULATORS AND METHODS FOR THEIR USE

CROSS-REFERENCE TO RELATED APPLICATIONS

[1] This application claims priority to U.S. Provisional Application No. 62/747,209, filed October 18, 2018, U.S. Provisional Application No. 62/803,516, filed February 10, 2019, and U.S. Provisional Application No. 62/857,516, filed June 5, 2019, the disclosures of which are herein incorporated by reference in their entirety for all purposes.

FIELD OF THE INVENTION

[2] The present disclosure generally relates to tricyclic compounds and their use for treatment of various indications. In particular the disclosure relates to tricyclic compounds and their use for treatment of various cancers, for example prostate cancer, including but not limited to, primary/localized prostate cancer (newly diagnosed), locally advanced prostate cancer, recurrent prostate cancer, non-metastatic castration-resistant prostate cancer, metastatic prostate cancer, metastatic castration-resistant prostate cancer (CRPC), and hormone-sensitive prostate cancer. This invention also relates to tricyclic compounds and their use for modulating androgen receptor (AR) activity, including truncated AR.

BACKGROUND OF THE INVENTION

[3] Androgens mediate their effects through the androgen receptor (AR). Androgens play a role in a wide range of developmental and physiological responses and are involved in male sexual differentiation, maintenance of spermatogenesis, and male gonadotropin regulation (R. K. Ross, G. A. Coetzee, C. L. Pearce, J. K. Reichardt, P. Bretsky, L. N. Kolonel, B. E. Henderson, E. Lander, D. Altshuler & G. Daley, *Eur Urol* 35, 355-361 (1999); A. A. Thomson, *Reproduction* 121, 187-195 (2001); N. Tanji, K. Aoki & M. Yokoyama, *Arch Androl* 47, 1-7 (2001)). Several lines of evidence show that androgens are associated with the development of prostate carcinogenesis. Firstly, androgens induce prostatic carcinogenesis in rodent models (R. L. Noble, *Cancer Res* 37, 1929-1933 (1977); R. L. Noble, *Oncology* 34, 138-141 (1977)) and men receiving androgens in the form of anabolic steroids have a higher incidence of prostate cancer (J. T. Roberts & D. M. Essenhight, *Lancet* 2, 742 (1986); J. A. Jackson, J. Waxman & A. M. Spiekerman, *Arch Intern Med* 149, 2365-2366 (1989); P. D. Guinan, W. Sadoughi, H. Alsheik, R. J. Ablin, D. Alrenga & I. M. Bush, *Am J Surg* 131, 599-600 (1976)). Secondly, prostate cancer does not develop if humans or dogs are castrated before puberty (J. D. Wilson & C. Roehrborn, *J Clin Endocrinol Metab* 84, 4324-4331 (1999); G. Wilding,

Cancer Surv 14, 113-130 (1992)). Castration of adult males causes involution of the prostate and apoptosis of prostatic epithelium while eliciting no effect on other male external genitalia (E. M. Bruckheimer & N. Kyprianou, *Cell Tissue Res* 301, 153-162 (2000); J. T. Isaacs, *Prostate* 5, 545-557 (1984)). This dependency on androgens provides the underlying rationale for treating prostate cancer with chemical or surgical castration (androgen ablation), also known as androgen ablation therapy (ABT) or androgen deprivation therapy (ADT).

[4] Androgens also play a role in female diseases such as polycystic ovary syndrome as well as cancers. One example is ovarian cancer where elevated levels of androgens are associated with an increased risk of developing ovarian cancer (K. J. Helzlsouer, A. J. Alberg, G. B. Gordon, C. Longcope, T. L. Bush, S. C. Hoffman & G. W. Comstock, *JAMA* 274, 1926-1930 (1995); R. J. Edmondson, J. M. Monaghan & B. R. Davies, *Br J Cancer* 86, 879-885 (2002)). The AR has been detected in a majority of ovarian cancers (H. A. Risch, *J Natl Cancer Inst* 90, 1774-1786 (1998); B. R. Rao & B. J. Slotman, *Endocr Rev* 12, 14-26 (1991); G. M. Clinton & W. Hua, *Crit Rev Oncol Hematol* 25, 1-9 (1997)), whereas estrogen receptor-alpha (ERa) and the progesterone receptor are detected in less than 50% of ovarian tumors.

[5] The only effective treatment available for advanced prostate cancer is the withdrawal of androgens which are essential for the survival of prostate luminal cells. Androgen ablation therapy causes a temporary reduction in tumor burden concomitant with a decrease in serum prostate-specific antigen (PSA). Unfortunately prostate cancer can eventually grow again in the absence of testicular androgens (castration-resistant disease) (Huber *et al* 1987 *Scand J. Urol Nephrol.* 104, 33-39). Castration-resistant prostate cancer that is still driven by AR is biochemically characterized before the onset of symptoms by a rising titre of serum PSA (Miller *et al* 1992 *J. Urol.* 147, 956-961). Once the disease becomes castration-resistant most patients succumb to their disease within two years.

[6] The AR has distinct functional domains that include the carboxy-terminal ligand-binding domain (LBD), a DNA-binding domain (DBD) comprising two zinc finger motifs, and an N-terminus domain (NTD) that contains two transcriptional activation units (tau1 and tau5) within activation function-1 (AF-1). Binding of androgen (ligand) to the LBD of the AR results in its activation such that the receptor can effectively bind to its specific DNA consensus site, termed the androgen response element (ARE), on the promoter and enhancer regions of "normally" androgen regulated genes, such as PSA, to initiate transcription. The AR can be activated in the absence of androgen by stimulation of the cAMP-dependent protein kinase (PKA) pathway, with interleukin-6 (IL-6) and by various growth factors (Culig *et al* 1994 *Cancer Res.* 54, 5474-5478; Nazareth *et al* 1996 *J. Biol. Chem.* 271, 19900-19907; Sadar

1999 *J. Biol. Chem.* 274, 7777-7783; Ueda *et al* 2002 *A J. Biol. Chem.* 277, 7076-7085; and Ueda *et al* 2002 *B J. Biol. Chem.* 277, 38087-38094). The mechanism of ligand-independent transformation of the AR has been shown to involve: 1) increased nuclear AR protein suggesting nuclear translocation; 2) increased AR/ARE complex formation; and 3) the AR-NTD (Sadar 1999 *J. Biol. Chem.* 274, 7777-7783; Ueda *et al* 2002 *A J. Biol. Chem.* 277, 7076-7085; and Ueda *et al* 2002 *B J. Biol. Chem.* 277, 38087-38094). The AR can be activated in the absence of testicular androgens by alternative signal transduction pathways in castration-resistant disease, which is consistent with the finding that nuclear AR protein is present in secondary prostate cancer tumors (Kim *et al* 2002 *Am. J. Pathol.* 160, 219-226; and van der Kwast *et al* 1991 *Inter. J. Cancer* 48, 189-193).

[7] Clinically available inhibitors of the AR include nonsteroidal antiandrogens such as apalutamide, darolutamide, bicalutamide (Casodex™), nilutamide, flutamide, and enzalutamide. There is also a class of steroidal antiandrogens, such as cyproterone acetate and spironolactone. Both steroidal and non-steroidal antiandrogens target the LBD of the AR and predominantly fail presumably due to poor affinity and mutations that lead to activation of the AR by these same antiandrogens (Taplin, M.E., Bubley, G.J., Kom Y.J., Small E.J., Upton M., Rajeshkumarm B., Balkm S.P., *Cancer Res.*, 59, 2511-2515 (1999)), and constitutively active AR splice variants. Antiandrogens have no effect on the constitutively active AR splice variants that lack the ligand-binding domain (LBD) and are associated with castration-recurrent prostate cancer (Dehm SM, Schmidt LJ, Heemers HV, Vessella RL, Tindall DJ., *Cancer Res* 68, 5469-77, 2008; Guo Z, Yang X, Sun F, Jiang R, Linn DE, Chen H, Chen H, Kong X, Melamed J, Tepper CG, Kung HJ, Brodie AM, Edwards J, Qiu Y., *Cancer Res.* 69, 2305-13, 2009; Hu *et al* 2009 *Cancer Res.* 69, 16-22; Sun *et al* 2010 *J Clin Invest.* 2010 120, 2715-30) and resistant to abiraterone and enzalutamide (Antonarakis *et al.*, *N Engl J Med.* 2014, 371, 1028-38; Scher *et al JAMA Oncol.* 2016 doi: 10.1001). Conventional therapy has concentrated on androgen-dependent activation of the AR through its C-terminal domain.

[8] Other relevant AR antagonists previously reported (*see*, WO 2010/000066, WO 2011/082487; WO 2011/082488; WO 2012/145330; WO 2015/031984; WO 2016/058080; and WO 2016/058082) that bind to full-length AR and/or truncated AR splice variants that have been recently developed or are currently being developed include: AR degraders such as niclosamide (Liu C *et al* 2014), galeterone (Njar *et al* 2015; Yu Z *et al* 2014), and ARV-330/Androgen receptor PROTAC (Neklesa *et al* 2016 *J Clin Oncol* 34 suppl 2S; abstr 267); AR DBD inhibitor VPC-14449 (Dalal K *et al* 2014 *J Biol Chem.* 289(38):26417-29; Li H *et al* 2014 *J Med Chem.* 57(15):6458-67); antiandrogens apalutamide (Clegg NJ *et al* 2012), ODM-

201 (Moilanen AM et al 2015), ODM-204 (Kallio et al *J Clin Oncol* 2016 vol. 34 no. 2_suppl 230), TAS3681 (Minamiguchi et al 2015 *J Clin Oncol* 33, suppl 7; abstr 266); and AR NTD inhibitors 3E10-AR441bsAb (Goicochea NL et al 2015), and sintokamide (Sadar et al 2008; Banuelos et al 2016).

[9] The AR-NTD is also a target for drug development (e.g. WO 2000/001813; Myung *et al. J. Clin. Invest* **2013**, 123, 2948), since the NTD contains Activation-Function-1 (AF-1) which is the essential region required for AR transcriptional activity (Jenster et al 1991. *Mol Endocrinol.* 5, 1396-404). The AR-NTD importantly plays a role in activation of the AR in the absence of androgens (Sadar, M.D. 1999 *J. Biol. Chem.* 274, 7777-7783; Sadar MD *et al* 1999 *Endocr Relat Cancer.* 6, 487-502; Ueda *et al* 2002 *J. Biol. Chem.* 277, 7076-7085; Ueda 2002 *J. Biol. Chem.* 277, 38087-38094; Blaszczyk *et al* 2004 *Clin Cancer Res.* 10, 1860-9; Dehm *et al* 2006 *J Biol Chem.* 28, 27882-93; Gregory *et al* 2004 *J Biol Chem.* 279, 7119-30). The AR-NTD is important in hormonal progression of prostate cancer as shown by application of decoy molecules (Quayle *et al* 2007, *Proc Natl Acad Sci U S A.* 104,1331-1336).

[10] While the crystal structure has been resolved for the AR C-terminus LBD, this has not been the case for the NTD due to its high flexibility and intrinsic disorder in solution (Reid *et al* 2002 *J. Biol. Chem.* 277, 20079-20086) thereby hampering virtual docking drug discovery approaches. Compounds that modulate AR, potentially through interaction with NTD domain, include the bisphenol compounds disclosed in published PCT Nos: WO 2010/000066, WO 2011/082487; WO 2011/082488; WO 2012/145330; WO 2012/139039; WO 2012/145328; WO 2013/028572; WO 2013/028791; WO 2014/179867; WO 2015/031984; WO 2016/058080; WO 2016/058082; WO 2016/112455; WO 2016/141458; WO 2017/177307; WO 2017/210771; and WO 2018/045450, and which are hereby incorporated by reference in their entireties.


[11] Transcriptionally active androgen receptor plays a major role in CRPC in spite of reduced blood levels of androgen (Karantanos, T. *et al Oncogene* **2013**, 32, 5501-5511; Harris, W. P. *et al Nature Clinical Practice Urology*, **2009**, 6, 76-85). AR mechanisms of resistance to ADT include: overexpression of AR (Visakorpi, T. *et al Nature Genetics* **1995**, 9, 401-406; Koivisto, P. *et al Scandinavian Journal of Clinical and Laboratory Investigation Supplementum* **1996**, 226, 57-63); gain-of-function mutations in the AR LBD (Culig Z. *et al Molecular Endocrinology* **1993**, 7, 1541-1550); intratumoral androgen synthesis (Cai, C. *et al Cancer Research* **2011**, 71, 6503-6513); altered expression and function of AR coactivators (Ueda, T. *et al The Journal of Biological Chemistry* **2002**, 277, 38087-38094; Xu J. *et al Nature Reviews Cancer* **2009**, 9, 615-630); aberrant post-translational modifications of AR (Gioeli D.

et al Molecular and Cellular Endocrinology **2012**, 352, 70-78; van der Steen T. *et al International Journal of Molecular Sciences* **2013**, 14, 14833-14859); and expression of AR splice variants (AR-Vs) which lack the ligand-binding domain (LBD) (Karantanos, T. *et al Oncogene* **2013**, 32, 5501-5511; Andersen R. J. *et al Cancer Cell* 2010, 17, 535-546; Myung J. K. *et al The Journal of Clinical Investigation* **2013**, 123, 2948-2960; Sun S. *et al The Journal of Clinical Investigation* **2010**, 120, 2715-2730). Anti-androgens such as bicalutamide and enzalutamide target AR LBD, but have no effect on truncated constitutively active AR-Vs such as AR-V7 (Li Y. *et al Cancer Research* **2013**, 73, 483-489). Expression of AR-V7 is associated with resistance to current hormone therapies (Li Y. *et al Cancer Research* **2013**, 73, 483-489; Antonarakis E. S. *et al The New England Journal of Medicine* **2014**, 371, 1028-1038).

[12] While significant advances have been made in this field, there remains a need for improved treatment for AR-mediated disorders including prostate cancer, especially metastatic castration-resistant prostate cancer. Development of compounds, via unique interactions with AR NTD, would provide patients alternative options and new hope.

SUMMARY OF THE INVENTION

[13] The compounds of the present disclosure are androgen receptor modulators which may be useful in treating various diseases and conditions as disclosed herein. In one embodiment, the present disclosure provides compounds comprising the structure of formula (IIIA):



[14] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

- [15] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;
- [16] C is a 3- to 10-membered ring;
- [17] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;
- [18] Y is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, $-NR^7-$, or $-N(COCH_3)-$;

- [19] W is a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;
- [20] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;
- [21] V is $-CH_2-$ and L is halogen, $-NH_2$, $-CHCl_2$, $-CCl_3$, or $-CF_3$; or
- [22] V is $-CH_2CH_2-$ and L is halogen or $-NH_2$;
- [23] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;
- [24] R^3 is selected from halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;
- [25] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, or C_1 - C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [26] R^7 is hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;
- [27] R^8 and R^9 are each independently hydrogen, halogen, or C_1 - C_3 alkyl;
- [28] R^{8a} and R^{9a} are each independently hydrogen, $-OH$, halogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}COR^{16}$, $-(C_1-C_3 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, or $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [29] R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;
- [30] R^{16} is hydrogen, optionally substituted C_1 - C_3 alkyl, optionally substituted C_2 - C_3 alkenyl, optionally substituted C_2 - C_3 alkynyl, C_3 - C_6 cycloalkyl, or phenyl;
- [31] each m is independently 0, 1, or 2;
- [32] n1 and n2 are each independently 0, 1, or 2;

[33] R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[34] R^{16} is hydrogen, optionally substituted C_1 - C_3 alkyl, optionally substituted C_2 - C_3 alkenyl, optionally substituted C_2 - C_3 alkynyl, C_3 - C_6 cycloalkyl, or phenyl;


[35] each m is independently 0, 1, or 2;

[36] n_1 and n_2 are each independently 0, 1, or 2;

[37] n_3 is 1, 2, 3, 4 or 5; and

[38] t is 0, 1 or 2.

[39] In one embodiment, the present disclosure provides compounds comprising the structure of formula (IVA):



[40] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[41] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[42] C is a 3- to 10-membered ring;

[43] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[44] Y and Z are each independently a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[45] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[46] V is $-CH_2-$ and L is halogen, $-NH_2$, or $-CF_3$; or

[47] V is $-CH_2CH_2-$ and L is halogen or $-NH_2$;

[48] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)- $(C_1$ - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-$

SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶ or optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶;

[49] R³ is selected from halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);

[50] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R⁵ and R⁶ taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[51] R⁷ is H or C₁-C₆ alkyl;

[52] R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R¹⁴ and R¹⁵ taken together form a 3- to 6-membered heterocyclyl;


[53] R¹⁶ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;


[54] n₁ and n₂ are each independently 0, 1, or 2;

[55] n₃ is 1, 2, 3, 4 or 5; and

[56] t is 0, 1 or 2.

[57] In one embodiment of the compound of formula (IVA), C is 5- to 10-membered heteroaryl or aryl. In some embodiments, C is 5- to 7-membered heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member. In some embodiments, C, which is substituted with (R³)_{n3}, is pyrazole, imidazole, oxazole, oxadiazole, oxazolone, isoxazole, thiazole, pyridyl, pyrazine, furan or pyrimidyl. In some embodiments, C, which is substituted





[58] In one embodiment of the compound of formula (IVA), R¹ and R² are each independently Cl, -CN, -CF₃, -OH, methyl, methoxy, or -CONH₂.

[59] In one embodiment of the compound of formula (IVA),

[60] A and B are phenyl;

[61] X is -(CR⁵R⁶)_t;

[62] Y and Z are each -O-;

[63] V is -CH₂- or -CH₂CH₂-;

[64] L is halogen;

[65] R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, or optionally substituted C₁-C₆ alkyl;

[66] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, or C₁-C₃ alkyl; and

[67] R¹⁶ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl.

[68] In one embodiment of the compound of formula (IVA),

[69] A and B are phenyl;

[70] X is -(CR⁵R⁶)_t;

[71] W is -CH₂- or -C(CH₃)H-;

[72] Y and Z are each -O-;

[73] V is -CH₂CH₂-;


[74] L is halogen;

[75] R¹ and R² are each independently hydrogen, halogen, or -CN;

[76] R^5 and R^6 are each independently hydrogen, or C₁-C₃ alkyl; and

[77] R^{16} is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl.

[78] In one embodiment, the present disclosure provides compounds comprising the structure of formula (A-I):



[79] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[80] C is a phenyl or a 5- to 7-membered monocyclic heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member;

[81] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[82] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[83] Z is a bond, $-CH_2-$, $-O-$, or $-NH-$;

[84] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[85] V is $-CH_2-$ and L is halogen, $-NH_2$, or $-CF_3$; or

[86] V is $-CH_2CH_2-$ and L is halogen or $-NH_2$;

[87] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, methyl, or $-CONH_2$;

[88] R^3 is selected from from $-CN$, C₁-C₃ alkoxy, $-CF_3$, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, $-S(C_1-C_3 \text{ alkyl})$, $-SO_2(C_1-C_3 \text{ alkyl})$, $-NH_2$, $-(C_1-C_3 \text{ alkyl})NH_2$, $-NHSO_2CH_3$, $-NHSO_2CF_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1-C_3 \text{ alkyl})_2$, $-CONH(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, or $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$;

[89] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, or C₁-C₃ alkyl;

[90] R^7 is H or C₁-C₆ alkyl;

[91] n_1 and n_2 are each independently 0, 1, or 2;

[92] n_3 is 1, 2, 3, 4 or 5; and

[93] t is 0, 1 or 2.

[94] In one embodiment of the compound of formula (IVA),

[95] In one embodiment of the compound of formula (A-I), at least one R^3 is selected from -CN, C₁-C₃ alkoxy, -CONH₂, -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, or -SO₂CH₃ and the other R^3 , if present, is selected from -CN, -CF₃, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -(C₁-C₃ alkyl)NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl).

[96] In one embodiment of the compound of formula (A-I),

[97] X is a bond or -(CR⁵R⁶)_i;

[98] W is a bond, -CH₂-, or -C(CH₃)H-;


[99] Y is -O-;

[100] Z is -O-;


[101] V is -CH₂- or -CH₂CH₂-; and


[102] L is halogen

[103] In one embodiment, the present disclosure provides compounds comprising the structure of formula (G-II):



[104] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:





[106] X is $-(CR^5R^6)_t-$;

[107] Y is $-O-$;

[108] Z is $-O-$;

[109] W is $-CH_2-$ or $-C(CH_3)H-$;

[110] V is $-CH_2CH_2-$;

[111] L is halogen;

[112] R^1 and R^2 are each independently Cl or $-CN$;

[113] at least one R^3 is selected from $-CN$, C_1-C_3 alkoxy, $-CONH_2$, $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, or $-SO_2CH_3$ and the other R^3 , if present, is selected from $-CN$, $-CF_3$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy, $-S(C_1-C_3 \text{ alkyl})$, $-SO_2(C_1-C_3 \text{ alkyl})$, $-NH_2$, $-(C_1-C_3 \text{ alkyl})NH_2$, $-NHSO_2CH_3$, $-NHSO_2CF_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1-C_3 \text{ alkyl})_2$, $-CONH(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, or $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$;

[114] R^5 and R^6 are each independently hydrogen or methyl;

[115] n_1 and n_2 are each independently 0, 1, or 2;

[116] n_3 is 1 or 2; and

[117] t is 1.

[118] In one embodiment of the compound of formula (G-II), at least one R^3 is selected from $-NHSO_2CH_3$, $-NHSO_2CH_2CH_3$, or $-SO_2CH_3$ and the other R^3 , if present, is selected from $-CN$, C_1-C_3 alkyl, C_1-C_3 alkoxy, $-SO_2(C_1-C_3 \text{ alkyl})$, $-NH_2$, $-(C_1-C_3 \text{ alkyl})NH_2$, $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1-C_3 \text{ alkyl})_2$, $-CONH(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, or $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$.

[119] In one embodiment, the present disclosure provides Compounds A1-A96, A98-A116, A118-A159, A161-A175, and A177-A234, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides Compounds A13, A57, A74, A93, A109, A112, A122, A126, A131, A134, A136, A137, A164, A168, A169, A170, A171, A172, A184, A185, A195, and/or A204, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

[120] In one embodiment, the present disclosure provides a pharmaceutical composition comprising any one of the compound of formula (IIIA), or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, and a pharmaceutically acceptable carrier. In one embodiment, the present disclosure provides a pharmaceutical composition comprising any one of the compound of formula (IVA), or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, and a pharmaceutically acceptable carrier. In one embodiment, the present disclosure provides a pharmaceutical composition comprising any one of the compound of formula (A-I), or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, and a pharmaceutically acceptable carrier. In one embodiment, the present disclosure provides a pharmaceutical composition comprising any one of the compound of formula (G-II), or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, and a pharmaceutically acceptable carrier.

[121] In one embodiment, the present disclosure provides a pharmaceutical composition comprising any one of Compounds A1-A96, A98-A116, A118-A159, A161-A175, and A177-A234, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, and a pharmaceutically acceptable carrier. In one embodiment, the present disclosure provides a pharmaceutical composition comprising any one of Compounds A13, A57, A74, A93, A109, A112, A122, A126, A131, A134, A136, A137, A164, A168, A169, A170, A171, A172, A184, A185, A195, and A204, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, and a pharmaceutically acceptable carrier.

[122] In one embodiment, the present disclosure provides a method for treating cancer, comprising administering any one of the compound of formula (IIIA), (IVA), (A-I), or (G-II), and Compounds A1-A96, A98-A116, A118-A159, A161-A175, or A177-A234, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, to a subject in need thereof. In one embodiment, the cancer is prostate cancer. In one embodiment, the prostate cancer is metastatic castration-resistant prostate cancer. In other embodiments, the prostate cancer expresses full-length androgen receptor or truncated androgen receptor splice variant.

[123] In one embodiment, the present disclosure provides a method for treating cancer, comprising administering any one of Compounds A13, A57, A74, A93, A109, A112, A122, A126, A131, A134, A136, A137, A164, A168, A169, A170, A171, A172, A184, A185, A195, and A204, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[124] Fig. 1 shows a dose-dependent curve of PSA-luciferase activities in response to representative compounds in transiently transfected LNCaP cells treated with synthetic androgen (R1881).

[125] Fig. 2A shows the percent PSA-luciferase activities of representative compounds in LNCaP cells transiently transfected with the PSA(6.1kb)-luciferase reporter and treated without or with R1881. Fig. 2B shows the percent PSA-luciferase activities of representative compounds in LNCaP cells co-transfected with an expression vector for AR-V7 and the PSA-luciferase reporter and treated without or with R1881.

[126] Fig. 3 shows concentration of representative compounds in plasma of male CD-1 mice after a single dose PO (oral) dose.

[127] Fig. 4 shows change in tumor volume in male NCG mice bearing LNCaP tumors after oral administration of representative compounds.

[128] Fig. 5 shows change in % body weight in male NCG mice bearing LNCaP tumors after oral administration of representative compounds.

[129] Fig. 6 shows individual tumor volume change from baseline measured at the end of experiment for oral administration of representative compounds to male NCG mice bearing LNCaP tumors.

[130] Fig. 7 shows concentration dependent effects on cell proliferation of LNCaP, PC3, and LNCaP95 cells treated with Compound A13.

DETAILED DESCRIPTION

[131] All publications, patents and patent applications, including any drawings and appendices therein are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication, patent or patent application, drawing, or appendix was specifically and individually indicated to be incorporated by reference in its entirety for all purposes.

Definitions

[132] While the following terms are believed to be well understood by one of ordinary skill in the art, the following definitions are set forth to facilitate explanation of the presently disclosed subject matter.

[133] Throughout the present specification, the terms “about” and/or “approximately” may be used in conjunction with numerical values and/or ranges. The term “about” is understood to mean those values near to a recited value. Furthermore, the phrases “less than about [a value]”

or “greater than about [a value]” should be understood in view of the definition of the term “about” provided herein. The terms “about” and “approximately” may be used interchangeably.

[134] Throughout the present specification, numerical ranges are provided for certain quantities. It is to be understood that these ranges comprise all subranges therein. Thus, the range “from 50 to 80” includes all possible ranges therein (e.g., 51-79, 52-78, 53-77, 54-76, 55-75, 60-70, etc.). Furthermore, all values within a given range may be an endpoint for the range encompassed thereby (e.g., the range 50-80 includes the ranges with endpoints such as 55-80, 50-75, etc.).

[135] The term “a” or “an” refers to one or more of that entity; for example, “a androgen receptor modulator” refers to one or more androgen receptor modulators or at least one androgen receptor modulator. As such, the terms “a” (or “an”), “one or more” and “at least one” are used interchangeably herein. In addition, reference to “an inhibitor” by the indefinite article “a” or “an” does not exclude the possibility that more than one of the inhibitors is present, unless the context clearly requires that there is one and only one of the inhibitors.

[136] As used herein, the verb “comprise” as is used in this description and in the claims and its conjugations are used in its non-limiting sense to mean that items following the word are included, but items not specifically mentioned are not excluded. The present invention may suitably “comprise”, “consist of”, or “consist essentially of”, the steps, elements, and/or reagents described in the claims.

[137] It is further noted that the claims may be drafted to exclude any optional element. As such, this statement is intended to serve as antecedent basis for use of such exclusive terminology as “solely”, “only” and the like in connection with the recitation of claim elements, or the use of a “negative” limitation.

[138] The term “pharmaceutically acceptable salts” includes both acid and base addition salts. Pharmaceutically acceptable salts include those obtained by reacting the active compound functioning as a base, with an inorganic or organic acid to form a salt, for example, salts of hydrochloric acid, sulfuric acid, phosphoric acid, methanesulfonic acid, camphorsulfonic acid, oxalic acid, maleic acid, succinic acid, citric acid, formic acid, hydrobromic acid, benzoic acid, tartaric acid, fumaric acid, salicylic acid, mandelic acid, carbonic acid, etc. Those skilled in the art will further recognize that acid addition salts may be prepared by reaction of the compounds with the appropriate inorganic or organic acid via any of a number of known methods.

[139] The term “treating” means one or more of relieving, alleviating, delaying, reducing, improving, or managing at least one symptom of a condition in a subject. The term “treating”

may also mean one or more of arresting, delaying the onset (i.e., the period prior to clinical manifestation of the condition) or reducing the risk of developing or worsening a condition.

[140] The compounds of the invention, or their pharmaceutically acceptable salts can contain one or more asymmetric centers and can thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that can be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)- or, as (*D*)- or (*L*)- for amino acids. The present disclosure is meant to include all such possible isomers, as well as their racemic and optically pure forms whether or not they are specifically depicted herein. Optically active (+) and (-), (*R*)- and (*S*)-, or (*D*)- and (*L*)- isomers can be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques, for example, chromatography and fractional crystallization. Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of the racemate (or the racemate of a salt or derivative) using, for example, chiral high pressure liquid chromatography (HPLC). When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both *E* and *Z* geometric isomers. Likewise, all tautomeric forms are also intended to be included.

[141] A "stereoisomer" refers to a compound made up of the same atoms bonded by the same bonds but having different three-dimensional structures, which are not interchangeable. The present disclosure contemplates various stereoisomers and mixtures thereof and includes "enantiomers", which refers to two stereoisomers whose molecules are nonsuperimposable mirror images of one another.

[142] A "tautomer" refers to a proton shift from one atom of a molecule to another atom of the same molecule. The present disclosure includes tautomers of any said compounds.

[143] A "prodrug" refers to a derivative of a compound of the present disclosure that will be converted to the compound *in vivo*. In one embodiment of the present disclosure, a prodrug includes a compound of formula (I), (IA), (IB), (IC), (II), (IIA), (IIIA), (IIB), (III), (IV), (V), (VA), (VI), (A), (A-I), (B)-(D), (E), (E-I)-(E-VII), (F), (G), (G-I), (G-II), (H), and (H-I), having a free hydroxyl group (-OH) that is acetylated (-OCOMe) at one or more positions.

[144] An "effective amount" means the amount of a formulation according to the invention that, when administered to a patient for treating a state, disorder or condition is sufficient to effect such treatment. The "effective amount" will vary depending on the active ingredient, the state, disorder, or condition to be treated and its severity, and the age, weight, physical condition and responsiveness of the mammal to be treated.

[145] The term "therapeutically effective" applied to dose or amount refers to that quantity of a compound or pharmaceutical formulation that is sufficient to result in a desired clinical benefit after administration to a patient in need thereof.

[146] As used herein, a "subject" can be a human, non-human primate, mammal, rat, mouse, cow, horse, pig, sheep, goat, dog, cat and the like. The subject can be suspected of having or at risk for having a cancer, such as prostate cancer, breast cancer, ovarian cancer, salivary gland carcinoma, or endometrial cancer, or suspected of having or at risk for having acne, hirsutism, alopecia, benign prostatic hyperplasia, ovarian cysts, polycystic ovary disease, precocious puberty, spinal and bulbar muscular atrophy, or age-related macular degeneration. Diagnostic methods for various cancers, such as prostate cancer, breast cancer, ovarian cancer, bladder cancer, pancreatic cancer, hepatocellular cancer, salivary gland carcinoma, or endometrial cancer, and diagnostic methods for acne, hirsutism, alopecia, benign prostatic hyperplasia, ovarian cysts, polycystic ovary disease, precocious puberty, spinal and bulbar muscular atrophy, or age-related macular degeneration and the clinical delineation of cancer, such as prostate cancer, breast cancer, ovarian cancer, bladder cancer, pancreatic cancer, hepatocellular cancer, salivary gland carcinoma, or endometrial cancer, diagnoses and the clinical delineation of acne, hirsutism, alopecia, benign prostatic hyperplasia, ovarian cysts, polycystic ovary disease, precocious puberty, spinal and bulbar muscular atrophy, or age-related macular degeneration are known to those of ordinary skill in the art.

[147] "Mammal" includes humans and both domestic animals such as laboratory animals (e.g., mice, rats, monkeys, dogs, etc.) and household pets (e.g., cats, dogs, swine, cattle, sheep, goats, horses, rabbits), and non-domestic animals such as wildlife and the like.

[148] All weight percentages (i.e., "% by weight" and "wt. %" and w/w) referenced herein, unless otherwise indicated, are measured relative to the total weight of the pharmaceutical composition.

[149] As used herein, "substantially" or "substantial" refers to the complete or nearly complete extent or degree of an action, characteristic, property, state, structure, item, or result. For example, an object that is "substantially" enclosed would mean that the object is either completely enclosed or nearly completely enclosed. The exact allowable degree of deviation from absolute completeness may in some cases depend on the specific context. However, generally speaking, the nearness of completion will be so as to have the same overall result as if absolute and total completion were obtained. The use of "substantially" is equally applicable when used in a negative connotation to refer to the complete or near complete lack of action, characteristic, property, state, structure, item, or result. For example, a composition that is

"substantially free of" other active agents would either completely lack other active agents, or so nearly completely lack other active agents that the effect would be the same as if it completely lacked other active agents. In other words, a composition that is "substantially free of" an ingredient or element or another active agent may still contain such an item as long as there is no measurable effect thereof

[150] The terms below, as used herein, have the following meanings, unless indicated otherwise:

[151] "Amino" refers to the -NH_2 radical.

[152] "Cyano" refers to the -CN radical.

[153] "Halo" or "halogen" refers to bromo, chloro, fluoro or iodo radical, including their radioisotopes. " ^{123}I " refers to the radioactive isotope of iodine having atomic mass 123. The compounds of Formula I can comprise at least one ^{123}I moiety. Throughout the present application, where structures depict a ^{123}I moiety at a certain position it is meant that the I moiety at this position is enriched for ^{123}I . In other words, the compounds contain more than the natural abundance of ^{123}I at the indicated position(s). It is not required that the compounds comprise 100% ^{123}I at the indicated positions, provided ^{123}I is present in more than the natural abundance. Typically the ^{123}I isotope is enriched to greater than 50%, greater than 60%, greater than 70%, greater than 80% or greater than 90%, relative to ^{127}I . " ^{18}F " refers to the radioactive isotope of fluorine having atomic mass 18. " ^{19}F " refers to the abundant, non-radioactive fluorine isotope having atomic mass 19. The compounds of Formula I can comprise at least one ^{18}F moiety. Throughout the present application, where structures depict a ^{18}F moiety at a certain position it is meant that the F moiety at this position is enriched for ^{18}F . In other words, the compounds contain more than the natural abundance of ^{18}F at the indicated position(s). It is not required that the compounds comprise 100% ^{18}F at the indicated positions, provided ^{18}F is present in more than the natural abundance. Typically the ^{18}F isotope is enriched to greater than 50%, greater than 60%, greater than 70%, greater than 80% or greater than 90%, relative to ^{19}F .

[154] "Hydroxy" or "hydroxyl" refers to the -OH radical.

[155] "Imino" refers to the =NH substituent.

[156] "Nitro" refers to the -NO_2 radical.

[157] "Oxo" refers to the =O substituent.

[158] "Thioxo" refers to the =S substituent.

[159] "Alkyl" or "alkyl group" refers to a fully saturated, straight or branched hydrocarbon chain radical having from one to twelve carbon atoms, and which is attached to the rest of the

molecule by a single bond. Alkyls comprising any number of carbon atoms from 1 to 12 are included. An alkyl comprising up to 12 carbon atoms is a C₁-C₁₂ alkyl, an alkyl comprising up to 10 carbon atoms is a C₁-C₁₀ alkyl, an alkyl comprising up to 6 carbon atoms is a C₁-C₆ alkyl and an alkyl comprising up to 5 carbon atoms is a C₁-C₅ alkyl. A C₁-C₅ alkyl includes C₅ alkyls, C₄ alkyls, C₃ alkyls, C₂ alkyls and C₁ alkyl (i.e., methyl). A C₁-C₆ alkyl includes all moieties described above for C₁-C₅ alkyls but also includes C₆ alkyls. A C₁-C₁₀ alkyl includes all moieties described above for C₁-C₅ alkyls and C₁-C₆ alkyls, but also includes C₇, C₈, C₉ and C₁₀ alkyls. Similarly, a C₁-C₁₂ alkyl includes all the foregoing moieties, but also includes C₁₁ and C₁₂ alkyls. Non-limiting examples of C₁-C₁₂ alkyl include methyl, ethyl, n-propyl, i-propyl, sec-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, n-pentyl, t-amyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl, n-undecyl, and n-dodecyl. Unless stated otherwise specifically in the specification, an alkyl group can be optionally substituted.

[160] “Alkylene” or “alkylene chain” refers to a fully saturated, straight or branched divalent hydrocarbon chain radical, and having from one to twelve carbon atoms. Non-limiting examples of C₁-C₁₂ alkylene include methylene, ethylene, propylene, n-butylenes, ethenylene, propenylene, n-butenylene, propynylene, n-butylenylene, and the like. The alkylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond. The points of attachment of the alkylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkylene chain can be optionally substituted.

[161] “Alkenyl” or “alkenyl group” refers to a straight or branched hydrocarbon chain radical having from two to twelve carbon atoms, and having one or more carbon-carbon double bonds. Each alkenyl group is attached to the rest of the molecule by a single bond. Alkenyl group comprising any number of carbon atoms from 2 to 12 are included. An alkenyl group comprising up to 12 carbon atoms is a C₂-C₁₂ alkenyl, an alkenyl comprising up to 10 carbon atoms is a C₂-C₁₀ alkenyl, an alkenyl group comprising up to 6 carbon atoms is a C₂-C₆ alkenyl and an alkenyl comprising up to 5 carbon atoms is a C₂-C₅ alkenyl. A C₂-C₅ alkenyl includes C₅ alkenyls, C₄ alkenyls, C₃ alkenyls, and C₂ alkenyls. A C₂-C₆ alkenyl includes all moieties described above for C₂-C₅ alkenyls but also includes C₆ alkenyls. A C₂-C₁₀ alkenyl includes all moieties described above for C₂-C₅ alkenyls and C₂-C₆ alkenyls, but also includes C₇, C₈, C₉ and C₁₀ alkenyls. Similarly, a C₂-C₁₂ alkenyl includes all the foregoing moieties, but also includes C₁₁ and C₁₂ alkenyls. Non-limiting examples of C₂-C₁₂ alkenyl include ethenyl (vinyl), 1-propenyl, 2-propenyl (allyl), iso-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-

hexenyl, 5-hexenyl, 1-heptenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 5-heptenyl, 6-heptenyl, 1-octenyl, 2-octenyl, 3-octenyl, 4-octenyl, 5-octenyl, 6-octenyl, 7-octenyl, 1-nonenyl, 2-nonenyl, 3-nonenyl, 4-nonenyl, 5-nonenyl, 6-nonenyl, 7-nonenyl, 8-nonenyl, 1-decenyl, 2-decenyl, 3-decenyl, 4-decenyl, 5-decenyl, 6-decenyl, 7-decenyl, 8-decenyl, 9-decenyl, 1-undecenyl, 2-undecenyl, 3-undecenyl, 4-undecenyl, 5-undecenyl, 6-undecenyl, 7-undecenyl, 8-undecenyl, 9-undecenyl, 10-undecenyl, 1-dodecenyl, 2-dodecenyl, 3-dodecenyl, 4-dodecenyl, 5-dodecenyl, 6-dodecenyl, 7-dodecenyl, 8-dodecenyl, 9-dodecenyl, 10-dodecenyl, and 11-dodecenyl. Unless stated otherwise specifically in the specification, an alkyl group can be optionally substituted.

[162] “Alkenylene” or “alkenylene chain” refers to a straight or branched divalent hydrocarbon chain radical, having from two to twelve carbon atoms, and having one or more carbon-carbon double bonds. Non-limiting examples of C₂-C₁₂ alkenylene include ethene, propene, butene, and the like. The alkenylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond. The points of attachment of the alkenylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkenylene chain can be optionally substituted.

[163] “Alkynyl” or “alkynyl group” refers to a straight or branched hydrocarbon chain radical having from two to twelve carbon atoms, and having one or more carbon-carbon triple bonds. Each alkynyl group is attached to the rest of the molecule by a single bond. Alkynyl group comprising any number of carbon atoms from 2 to 12 are included. An alkynyl group comprising up to 12 carbon atoms is a C₂-C₁₂ alkynyl, an alkynyl comprising up to 10 carbon atoms is a C₂-C₁₀ alkynyl, an alkynyl group comprising up to 6 carbon atoms is a C₂-C₆ alkynyl and an alkynyl comprising up to 5 carbon atoms is a C₂-C₅ alkynyl. A C₂-C₅ alkynyl includes C₅ alkynyls, C₄ alkynyls, C₃ alkynyls, and C₂ alkynyls. A C₂-C₆ alkynyl includes all moieties described above for C₂-C₅ alkynyls but also includes C₆ alkynyls. A C₂-C₁₀ alkynyl includes all moieties described above for C₂-C₅ alkynyls and C₂-C₆ alkynyls, but also includes C₇, C₈, C₉ and C₁₀ alkynyls. Similarly, a C₂-C₁₂ alkynyl includes all the foregoing moieties, but also includes C₁₁ and C₁₂ alkynyls. Non-limiting examples of C₂-C₁₂ alkenyl include ethynyl, propynyl, butynyl, pentynyl and the like. Unless stated otherwise specifically in the specification, an alkyl group can be optionally substituted.

[164] “Alkynylene” or “alkynylene chain” refers to a straight or branched divalent hydrocarbon chain radical, having from two to twelve carbon atoms, and having one or more carbon-carbon triple bonds. Non-limiting examples of C₂-C₁₂ alkynylene include ethynylene,

propargylene and the like. The alkynylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond. The points of attachment of the alkynylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkynylene chain can be optionally substituted.

[165] “Alkoxy” refers to a radical of the formula $-OR_a$ where R_a is an alkyl, alkenyl or alkynyl radical as defined above containing one to twelve carbon atoms. Unless stated otherwise specifically in the specification, an alkoxy group can be optionally substituted.

[166] “Alkylamino” refers to a radical of the formula $-NHR_a$ or $-NR_aR_a$ where each R_a is, independently, an alkyl, alkenyl or alkynyl radical as defined above containing one to twelve carbon atoms. Unless stated otherwise specifically in the specification, an alkylamino group can be optionally substituted.

[167] “Alkylcarbonyl” refers to the $-C(=O)R_a$ moiety, wherein R_a is an alkyl, alkenyl or alkynyl radical as defined above. A non-limiting example of an alkyl carbonyl is the methyl carbonyl (“acetal”) moiety. Alkylcarbonyl groups can also be referred to as “ C_w-C_z acyl” where w and z depicts the range of the number of carbon in R_a , as defined above. For example, “ C_1-C_{10} acyl” refers to alkylcarbonyl group as defined above, where R_a is C_1-C_{10} alkyl, C_1-C_{10} alkenyl, or C_1-C_{10} alkynyl radical as defined above. Unless stated otherwise specifically in the specification, an alkyl carbonyl group can be optionally substituted.

[168] “Aryl” refers to a hydrocarbon ring system radical comprising hydrogen, 6 to 18 carbon atoms and at least one aromatic ring. For purposes of this invention, the aryl radical can be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which can include fused or bridged ring systems. Aryl radicals include, but are not limited to, aryl radicals derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, fluoranthene, fluorene, as-indacene, s-indacene, indane, indene, naphthalene, phenalene, phenanthrene, pleiadene, pyrene, and triphenylene. Unless stated otherwise specifically in the specification, the term “aryl” is meant to include aryl radicals that are optionally substituted.

[169] “Aralkyl” or “arylalkyl” refers to a radical of the formula $-R_b-R_c$ where R_b is an alkylene group as defined above and R_c is one or more aryl radicals as defined above, for example, benzyl, diphenylmethyl and the like. Unless stated otherwise specifically in the specification, an aralkyl group can be optionally substituted.

[170] “Aralkenyl” or “arylalkenyl” refers to a radical of the formula $-R_b-R_c$ where R_b is an alkenylene group as defined above and R_c is one or more aryl radicals as defined above.

Unless stated otherwise specifically in the specification, an aralkenyl group can be optionally substituted.

[171] “Aralkynyl” or “arylalkynyl” refers to a radical of the formula $-R_b-R_c$ where R_b is an alkynylene group as defined above and R_c is one or more aryl radicals as defined above. Unless stated otherwise specifically in the specification, an aralkynyl group can be optionally substituted.

[172] “Carbocyclyl,” “carbocyclic ring” or “carbocycle” refers to a rings structure, wherein the atoms which form the ring are each carbon. Carbocyclic rings can comprise from 3 to 20 carbon atoms in the ring. Carbocyclic rings include aryls and cycloalkyl, cycloalkenyl and cycloalkynyl as defined herein. Unless stated otherwise specifically in the specification, a carbocyclyl group can be optionally substituted.

[173] “Cycloalkyl” refers to a stable non-aromatic monocyclic or polycyclic fully saturated hydrocarbon radical consisting solely of carbon and hydrogen atoms, which can include fused or bridged ring systems, having from three to twenty carbon atoms, preferably having from three to ten carbon atoms, and which is attached to the rest of the molecule by a single bond. Monocyclic cycloalkyl radicals include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic cycloalkyl radicals include, for example, adamantyl, norbornyl, decalyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like. Unless otherwise stated specifically in the specification, a cycloalkyl group can be optionally substituted.

[174] “Cycloalkenyl” refers to a stable non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, having one or more carbon-carbon double bonds, which can include fused or bridged ring systems, having from three to twenty carbon atoms, preferably having from three to ten carbon atoms, and which is attached to the rest of the molecule by a single bond. Monocyclic cycloalkenyl radicals include, for example, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, and the like. Polycyclic cycloalkenyl radicals include, for example, bicyclo[2.2.1]hept-2-enyl and the like. Unless otherwise stated specifically in the specification, a cycloalkenyl group can be optionally substituted.

[175] “Cycloalkynyl” refers to a stable non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, having one or more carbon-carbon triple bonds, which can include fused or bridged ring systems, having from three to twenty carbon atoms, preferably having from three to ten carbon atoms, and which is attached to the rest of the molecule by a single bond. Monocyclic cycloalkynyl radicals include, for example,

cycloheptynyl, cyclooctynyl, and the like. Unless otherwise stated specifically in the specification, a cycloalkynyl group can be optionally substituted.

[176] “Cycloalkylalkyl” refers to a radical of the formula $-R_b-R_d$ where R_b is an alkylene, alkenylene, or alkynylene group as defined above and R_d is a cycloalkyl, cycloalkenyl, cycloalkynyl radical as defined above. Unless stated otherwise specifically in the specification, a cycloalkylalkyl group can be optionally substituted.

[177] “Haloalkyl” refers to an alkyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, e.g., trifluoromethyl, difluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 1,2-difluoroethyl, 3-bromo-2-fluoropropyl, 1,2-dibromoethyl, and the like. Unless stated otherwise specifically in the specification, a haloalkyl group can be optionally substituted.

[178] “Haloalkenyl” refers to an alkenyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, e.g., 1-fluoropropenyl, 1,1-difluorobutenyl, and the like. Unless stated otherwise specifically in the specification, a haloalkenyl group can be optionally substituted.

[179] “Haloalkynyl” refers to an alkynyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, e.g., 1-fluoropropynyl, 1-fluorobutynyl, and the like. Unless stated otherwise specifically in the specification, a haloalkenyl group can be optionally substituted.

[180] “Heterocyclyl,” “heterocyclic ring” or “heterocycle” refers to a stable 3- to 20-membered non-aromatic, partially aromatic, or aromatic ring radical which consists of two to twelve carbon atoms and from one to six heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. Heterocyclyl or heterocyclic rings include heteroaryls as defined below. Unless stated otherwise specifically in the specification, the heterocyclyl radical can be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which can include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heterocyclyl radical can be optionally oxidized; the nitrogen atom can be optionally quaternized; and the heterocyclyl radical can be partially or fully saturated. Examples of such heterocyclyl radicals include, but are not limited to, dioxolanyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazolyl, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholinyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, quinuclidinyl, thiazolidinyl, tetrahydrofuryl, trithianyl, tetrahydropyranyl, thiomorpholinyl, thiamorpholinyl,

1-oxo-thiomorpholinyl, and 1,1-dioxo-thiomorpholinyl. Unless stated otherwise specifically in the specification, a heterocyclyl group can be optionally substituted.

[181] “Heterocyclalkyl” refers to a radical of the formula $-R_b-R_e$ where R_b is an alkylene group as defined above and R_e is a heterocyclyl radical as defined above. Unless stated otherwise specifically in the specification, a heterocycloalkylalkyl group can be optionally substituted.

[182] “Heterocyclalkenyl” refers to a radical of the formula $-R_b-R_e$ where R_b is an alkenylene group as defined above and R_e is a heterocyclyl radical as defined above. Unless stated otherwise specifically in the specification, a heterocycloalkylalkenyl group can be optionally substituted.

[183] “Heterocyclalkynyl” refers to a radical of the formula $-R_b-R_e$ where R_b is an alkynylene group as defined above and R_e is a heterocyclyl radical as defined above. Unless stated otherwise specifically in the specification, a heterocycloalkylalkynyl group can be optionally substituted.

[184] “N-heterocyclyl” refers to a heterocyclyl radical as defined above containing at least one nitrogen and where the point of attachment of the heterocyclyl radical to the rest of the molecule is through a nitrogen atom in the heterocyclyl radical. Unless stated otherwise specifically in the specification, a N-heterocyclyl group can be optionally substituted.

[185] “Heteroaryl” refers to a 5- to 20-membered ring system radical comprising hydrogen atoms, one to thirteen carbon atoms, one to six heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, and at least one aromatic ring. For purposes of this invention, the heteroaryl radical can be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which can include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heteroaryl radical can be optionally oxidized; the nitrogen atom can be optionally quaternized. Examples include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzothiazolyl, benzindolyl, benzodioxolyl, benzofuranyl, benzooxazolyl, benzothiazolyl, benzothiadiazolyl, benzo[b][1,4]dioxepinyl, 1,4-benzodioxanyl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl, benzofuranonyl, benzothieryl (benzothiophene), benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, carbazolyl, cinnolinyl, dibenzofuranyl, dibenzothiophene, furanyl, furanonyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indoliziny, isoxazolyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 1-oxidopyridinyl, 1-oxidopyrimidinyl, 1-oxidopyrazinyl, 1-oxidopyridazinyl, 1-phenyl-1H-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl,

purinyl, pyrrolyl, pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, quinazolinyl, quinoxalinyl, quinolinyl, quinuclidinyl, isoquinolinyl, tetrahydroquinolinyl, thiazolyl, thiadiazolyl, triazolyl, tetrazolyl, triazinyl, and thiophene (i.e. thienyl). Unless stated otherwise specifically in the specification, a heteroaryl group can be optionally substituted.

[186] “N-heteroaryl” refers to a heteroaryl radical as defined above containing at least one nitrogen and where the point of attachment of the heteroaryl radical to the rest of the molecule is through a nitrogen atom in the heteroaryl radical. Unless stated otherwise specifically in the specification, an N-heteroaryl group can be optionally substituted.

[187] “Heteroarylalkyl” refers to a radical of the formula $-R_b-R_f$ where R_b is an alkylene chain as defined above and R_f is a heteroaryl radical as defined above. Unless stated otherwise specifically in the specification, a heteroarylalkyl group can be optionally substituted.

[188] “Heteroarylalkenyl” refers to a radical of the formula $-R_b-R_f$ where R_b is an alkenylene chain as defined above and R_f is a heteroaryl radical as defined above. Unless stated otherwise specifically in the specification, a heteroarylalkenyl group can be optionally substituted.

[189] “Heteroarylalkynyl” refers to a radical of the formula $-R_b-R_f$ where R_b is an alkynylene chain as defined above and R_f is a heteroaryl radical as defined above. Unless stated otherwise specifically in the specification, a heteroarylalkynyl group can be optionally substituted.

[190] “Ring” refers to a cyclic group which can be fully saturated, partially saturated, or fully unsaturated. A ring can be monocyclic, bicyclic, tricyclic, or tetracyclic. Unless stated otherwise specifically in the specification, a ring can be optionally substituted.

[191] “Thioalkyl” refers to a radical of the formula $-SR_a$ where R_a is an alkyl, alkenyl, or alkynyl radical as defined above containing one to twelve carbon atoms. Unless stated otherwise specifically in the specification, a thioalkyl group can be optionally substituted.

[192] The term “substituted” used herein means any of the above groups (i.e., alkyl, alkylene, alkenyl, alkenylene, alkynyl, alkynylene, alkoxy, alkylamino, alkylcarbonyl, thioalkyl, aryl, aralkyl, carbocyclyl, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, haloalkyl, heterocyclyl, N-heterocyclyl, heterocyclylalkyl, heteroaryl, N-heteroaryl and/or heteroarylalkyl) wherein at least one hydrogen atom is replaced by a bond to a non-hydrogen atoms such as, but not limited to: a halogen atom such as F, Cl, Br, and I; an oxygen atom in groups such as hydroxyl groups, alkoxy groups, and ester groups; a sulfur atom in groups such as thiol groups, thioalkyl groups, sulfone groups, sulfonyl groups, and sulfoxide groups; a nitrogen atom in groups such as amines, amides, alkylamines, dialkylamines, arylamines, alkylarylamines, diarylamines, N-oxides, imides, and enamines; a silicon atom in groups such

as trialkylsilyl groups, dialkylarylsilyl groups, alkyl diarylsilyl groups, and triarylsilyl groups; and other heteroatoms in various other groups.

[193] “Substituted” also means any of the above groups in which one or more hydrogen atoms are replaced by a higher-order bond (e.g., a double- or triple-bond) to a heteroatom such as oxygen in oxo, carbonyl, carboxyl, and ester groups; and nitrogen in groups such as imines, oximes, hydrazones, and nitriles. For example, “substituted” includes any of the above groups in which one or more hydrogen atoms are replaced with $-\text{NR}_g\text{R}_h$, $-\text{NR}_g\text{C}(=\text{O})\text{R}_h$, $-\text{NR}_g\text{C}(=\text{O})\text{NR}_g\text{R}_h$, $-\text{NR}_g\text{C}(=\text{O})\text{OR}_h$, $-\text{NR}_g\text{SO}_2\text{R}_h$, $-\text{OC}(=\text{O})\text{NR}_g\text{R}_h$, $-\text{OR}_g$, $-\text{SR}_g$, $-\text{SOR}_g$, $-\text{SO}_2\text{R}_g$, $-\text{OSO}_2\text{R}_g$, $-\text{SO}_2\text{OR}_g$, $=\text{NSO}_2\text{R}_g$, and $-\text{SO}_2\text{NR}_g\text{R}_h$. “Substituted” also means any of the above groups in which one or more hydrogen atoms are replaced with $-\text{C}(=\text{O})\text{R}_g$, $-\text{C}(=\text{O})\text{OR}_g$, $-\text{C}(=\text{O})\text{NR}_g\text{R}_h$, $-\text{CH}_2\text{SO}_2\text{R}_g$, $-\text{CH}_2\text{SO}_2\text{NR}_g\text{R}_h$. In the foregoing, R_g and R_h are the same or different and independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylamino, thioalkyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, haloalkyl, haloalkenyl, haloalkynyl, heterocyclyl, N-heterocyclyl, heterocyclylalkyl, heteroaryl, N-heteroaryl and/or heteroarylalkyl. “Substituted” further means any of the above groups in which one or more hydrogen atoms are replaced by a bond to an amino, cyano, hydroxyl, imino, nitro, oxo, thioxo, halo, alkyl, alkenyl, alkynyl, alkoxy, alkylamino, thioalkyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, haloalkyl, haloalkenyl, haloalkynyl, heterocyclyl, N-heterocyclyl, heterocyclylalkyl, heteroaryl, N-heteroaryl and/or heteroarylalkyl group. In addition, each of the foregoing substituents can also be optionally substituted with one or more of the above substituents.

[194] As used herein, the symbol “” (hereinafter can be referred to as “a point of attachment bond”) denotes a bond that is a point of attachment between two chemical entities, one of which is depicted as being attached to the point of attachment bond and the other of which is not depicted as being attached to the point of attachment bond. For example, “ $\text{XY}-\text{$ ” indicates that the chemical entity “XY” is bonded to another chemical entity via the point of attachment bond. Furthermore, the specific point of attachment to the non-depicted chemical entity can be specified by inference. For example, the compound CH_3-R^3 , wherein R^3 is H or “ $\text{XY}-\text{$ ” infers that when R^3 is “XY”, the point of attachment bond is the same bond as the bond by which R^3 is depicted as being bonded to CH_3 .


[195] “Fused” refers to any ring structure described herein which is fused to an existing ring structure in the compounds of the invention. When the fused ring is a heterocyclyl ring or a heteroaryl ring, any carbon atom on the existing ring structure which becomes part of the fused heterocyclyl ring or the fused heteroaryl ring can be replaced with a nitrogen atom.

[196] The following description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed inventions, or that any publication specifically or implicitly referenced is prior art.

Compounds of the Present Disclosure

[197] The compound of the present disclosure can be useful for modulating androgen receptor (AR). Further, the compound of the present disclosure can be useful for treating various diseases and conditions including, but not limited to, cancer. In some embodiments, the cancer is prostate cancer or breast cancer.

[198] In one embodiment, the present disclosure provides compounds comprising the structure of formula (I):



[199] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[200] A and B are each independently aryl or heteroaryl;

[201] C is a 3-to 10-membered ring;

[202] X is a bond, $-(CR^5R^6)_t-$, $-O-$, $-C(=O)-$, $-S-$, $-S(=O)-$, $-SO_2-$, $-NR^7-$, $-N(R^7)CO-$, $-CON(R^7)-$, or $-NSO_2R^7-$;

[203] Y and Z are each independently a bond, $-(CR^8R^9)_m-$, $-O-$, $-C(=O)-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;

[204] W and V are each independently a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

[205] L is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;

[206] R^1 and R^2 are each independently hydrogen, halogen, -CN, -CF₃, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[207] R^3 is hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -SR¹⁶, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[208] R^5 and R^6 are each independently hydrogen, halogen, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^5 and R^6 taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[209] R^8 and R^9 are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[210] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -OCO(C₁-C₆ alkyl), -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-

CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^{8a} and R^{8b} taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[211] R⁷, R¹⁰ and R¹⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁷ and R^{8a} taken together form an optionally substituted heterocyclyl;

[212] R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or (R¹¹ and R¹²) or (R¹⁴ and R¹⁵) taken together form an optionally substituted heterocyclyl;


[213] each m is independently 0, 1 or 2;

[214] n₁ and n₂ are each independently 0, 1, 2, 3, or 4;

[215] n₃ is 0, 1, 2, 3, 4 or 5; and

[216] each t is independently 0, 1 or 2.

[217] In one embodiment the present disclosure provides compounds comprising the structure of formula (IA):



[218] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[219] A and B are each independently aryl or heteroaryl;

[220] C is a 3- to 10-membered ring;

[221] X is a bond, -(CR⁵R⁶)_t-, -O-, -C(=O)-, -S-, -S(=O)-, -SO₂-, -NR⁷-, -N(R⁷)CO-, -CON(R⁷)-, or -NSO₂R⁷-;

[222] Y and Z are each independently a bond, -(CR⁸R⁹)_m-, -O-, -C(=O)-, -S-, -S(=O)-, -SO₂-, or -NR⁷-;

[223] W and V are each independently a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

[224] L is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;

[225] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[226] R^3 is hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-SR^{16}$, optionally substituted C_1-C_6 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COOR^{16}$, $-NR^{14}COR^{16}$, $-NR^{14}CONR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[227] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, optionally substituted C_1-C_6 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted C_1-C_6 alkoxy, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^5 and R^6 taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[228] R^8 and R^9 are each independently hydrogen, halogen, or C_1-C_3 alkyl;

[229] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -OCO(C₁-C₆ alkyl), -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^{8a} and R^{8b} taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[230] R^7 , R^{10} and R^{16} are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, C₁-C₆ haloalkyl, C₂-C₆ haloalkenyl, C₂-C₆ haloalkynyl, optionally substituted carbocyclyl, optionally substituted -CO(C₁-C₆ alkyl), -CO(optionally substituted heterocyclyl), optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^7 and R^{8a} taken together form an optionally substituted heterocyclyl;

[231] R^{11} , R^{12} , R^{13} , R^{14} and R^{15} are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted -COO(C₁-C₆ alkyl), optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or (R^{11} and R^{12}) or (R^{14} and R^{15}) taken together form an optionally substituted heterocyclyl;


[232] each m is independently 0, 1 or 2;

[233] n1 and n2 are each independently 0, 1, 2, 3, or 4;

[234] n3 is 0, 1, 2, 3, 4 or 5; and

[235] each t is independently 0, 1 or 2.

[236] In one embodiment the present disclosure provides compounds comprising the structure of formula (IB):



[237] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

- [238] A and B are each independently aryl or heteroaryl;
- [239] C is a 3- to 10-membered ring;
- [240] X is a bond, $-(CR^5R^6)_t-$, $-O-$, $-C(=O)-$, $-S-$, $-S(=O)-$, $-SO_2-$, $-NR^7-$, $-N(R^7)CO-$, $-CON(R^7)-$, or $-NSO_2R^7-$;
- [241] Y is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;
- [242] W is a bond, $-(CR^{8a}R^{9a})_m-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;
- [243] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;
- [244] V is $-CH_2-$, $-CH_2CH_2-$, $-CH(CH_3)CH_2-$, $-CH_2CH(CH_3)-$, or $-CH_2CH_2CH_2-$;
- [245] L is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;
- [246] R^1 and R^2 are each independently hydrogen, deuterium, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)-(C_1 - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} , optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;
- [247] R^3 is hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-SR^{16}$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)-(C_1 - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COOR^{16}$, $-NR^{14}COR^{16}$, $-NR^{14}CONR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} , optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;
- [248] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 alkoxy, $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, -

CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁵ and R⁶ taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[249] R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[250] R^{8a} and R^{9a} are each independently hydrogen, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^{8a} and R^{8b} taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[251] R⁷, R¹⁰ and R¹⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkyl-NH₂, C₁-C₆ haloalkyl, C₂-C₆ haloalkenyl, C₂-C₆ haloalkynyl, optionally substituted carbocyclyl, optionally substituted -CO(C₁-C₆ alkyl), -CO(optionally substituted heterocyclyl), optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁷ and R^{8a} taken together form an optionally substituted heterocyclyl;

[252] R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted -COO(C₁-C₆ alkyl), optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or (R¹¹ and R¹²) or (R¹⁴ and R¹⁵) or (R¹⁴ and R¹⁶) taken together form an optionally substituted heterocyclyl;


[253] each m is independently 0, 1 or 2;

[254] n₁ and n₂ are each independently 0, 1, 2, 3, or 4;

[255] n₃ is 0, 1, 2, 3, 4 or 5; and

[256] each t is independently 0, 1 or 2.

[257] In one embodiment the present disclosure provides compounds comprising the structure of formula (IC)



[258] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[259] A and B are each independently aryl or heteroaryl;

[260] C is a 3- to 10-membered ring;

[261] X is a bond, $-(CR^5R^6)_t$ -, -O-, -C(=O)-, -S-, -S(=O)-, -SO₂-, -NR⁷-, -N(R⁷)CO-, -CON(R⁷)-, or -NSO₂R⁷-;

[262] Y is a bond, $-(CR^8R^9)_m$ -, -O-, -S-, -S(=O)-, -SO₂-, or -NR⁷-;

[263] W is a bond, $-(CR^{8a}R^{9a})_m$ -, -C(=O)-, N(R⁷)CO-, -CONR⁷-, or -NSO₂R⁷-;

[264] Z is a bond, $-(CR^8R^9)_m$ -, -O-, -S-, -S(=O)-, -SO₂-, or -NR⁷-;

[265] V is -CH₂-, -CH₂CH₂-, -CH(CH₃)CH₂-, -CH₂CH(CH₃)-, or -CH₂CH₂CH₂-;

[266] L is hydrogen, halogen, -CF₂R¹⁰-, -CF₃-, -CCl₂R¹⁰-, -CCl₃-, -CN-, -OR¹⁰-, -NR¹¹R¹²-, or -CONR¹¹R¹²-;

[267] R¹ and R² are each independently hydrogen, deuterium, halogen, -CN-, -CF₃-, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$ -, -NR¹³R¹⁴-, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$ -, -NR¹⁴SO₂R¹⁶-, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$ -, -NR¹⁴COR¹⁶-, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$ -, -CONR¹⁴R¹⁵-, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$ -, -SO₂NR¹⁴R¹⁵-, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$ -, optionally substituted -SO₂R¹⁶-, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$ -, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[268] R³ is hydrogen, halogen, oxo, =S, =NR¹⁶-, -CN-, -CF₃-, -OH-, -SR¹⁶-, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$ -, -NR¹³R¹⁴-, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$ -, -NR¹⁴SO₂R¹⁶-, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$ -, -NR¹⁴COR¹⁶-, -NR¹⁴CONR¹⁴R¹⁵-, optionally substituted $-(C_1-C_6 \text{ alkyl})-$

NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[269] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹³R¹⁴, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁵ and R⁶ taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[270] R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[271] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^{8a} and R^{8b} taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[272] R⁷, R¹⁰ and R¹⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkyl-NH₂, C₁-C₆ haloalkyl, C₂-C₆ haloalkenyl, C₂-C₆ haloalkynyl, optionally substituted carbocyclyl, optionally substituted -CO(C₁-C₆ alkyl), -CO(optionally substituted heterocyclyl), optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁷ and R^{8a} taken together form an optionally substituted heterocyclyl;

[273] R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted -COO(C₁-C₆ alkyl), optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted

heteroaryl; or (R¹¹ and R¹²) or (R¹⁴ and R¹⁵) or (R¹⁴ and R¹⁶) taken together form an optionally substituted heterocyclyl;


[274] each m is independently 0, 1 or 2;

[275] n₁ and n₂ are each independently 0, 1, 2, 3, or 4;

[276] n₃ is 0, 1, 2, 3, 4 or 5; and

[277] each t is independently 0, 1 or 2.

[278] In one embodiment, the present disclosure provides compounds comprising the structure of formula (II):



[279] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[280] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[281] C is a 3- to 10-membered ring;

[282] X is a bond, -(CR⁵R⁶)_t-, or -NR⁷-;

[283] Y is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-;

[284] Z is a bond, -(CR⁸R⁹)_m-, -O-, -S-, -S(=O)-, -SO₂-, or -NR⁷-;

[285] W is a bond, -CH₂-, -C(CH₃)H-, -C(=O)-, -NHCO-, -N(C₁-C₃ alkyl)CO-, or -CONH-, or -CON(C₁-C₃ alkyl)-;

[286] V is a bond, -(CR^{8a}R^{9a})_m-, -C(=O)-, -N(R⁷)CO-, -CONR⁷-, or -NSO₂R⁷-;

[287] L is hydrogen, halogen, -CF₂R¹⁰, -CF₃, -CN, -OR¹⁰, -NR¹¹R¹², or -CONR¹¹R¹²;

[288] R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -

SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶ or optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶;

[289] R³ is selected from hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);

[290] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R⁵ and R⁶ taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[291] R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[292] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, or -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[293] R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R¹⁴ and R¹⁵ taken together form a 3- to 6-membered heterocyclyl;

[294] R⁷, R¹⁰ and R¹⁶ are each independently hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl;


[295] each m is independently 0, 1 or 2;

[296] n₁ and n₂ are each independently 0, 1, or 2;

[297] n₃ is 0, 1, 2, 3, 4 or 5; and

[298] t is 0, 1 or 2.

[299] In one embodiment, the present disclosure provides compounds comprising the structure of formula (IIA):



[300] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[301] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[302] C is a 3- to 10-membered ring;

[303] X is a bond, $-(CR^5R^6)_l-$, or $-NR^7-$;

[304] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[305] Z is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;

[306] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-NHCO-$, $-N(C_1-C_3 \text{ alkyl})CO-$, or $-CONH-$, or $-CON(C_1-C_3 \text{ alkyl})-$;

[307] V is $-CH_2-$, $-CH_2CH_2-$, $-CH(CH_3)CH_2-$, $-CH_2CH(CH_3)-$, or $-CH_2CH_2CH_2-$;

[308] L is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;

[309] R^1 and R^2 are each independently hydrogen, deuterium, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;

[310] R^3 is selected from hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;

[311] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, or C_1-C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[312] R^8 and R^9 are each independently hydrogen, halogen, or C_1-C_3 alkyl;

[313] R^{11} , R^{12} , R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1-C_3 alkyl, C_2-C_3 alkenyl, or C_2-C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[314] R^7 , R^{10} and R^{16} are each independently hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl or C_1 - C_6 alkyl- NH_2 ; or R^{14} and R^{16} taken together form a 3- to 6-membered heterocyclyl;


[315] each m is independently 0, 1 or 2;

[316] n_1 and n_2 are each independently 0, 1, or 2;

[317] n_3 is 0, 1, 2, 3, 4 or 5; and

[318] t is 0, 1 or 2.

[319] In one embodiment, the present disclosure provides compounds comprising the structure of formula (IIB):



[320] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[321] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[322] C is a 3- to 10-membered ring;

[323] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[324] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[325] Z is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;

[326] W is a bond, $-CH_2-$, or $-C(CH_3)H-$;

[327] V is a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

[328] L is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;

[329] R^1 and R^2 are each independently hydrogen, deuterium, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, -

SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶ or optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶;

[330] R³ is selected from hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);

[331] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R⁵ and R⁶ taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[332] R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[333] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, or -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[334] R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R¹⁴ and R¹⁵ taken together form a 3- to 6-membered heterocyclyl;

[335] R⁷, R¹⁰ and R¹⁶ are each independently hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl or C₁-C₆ alkyl-NH₂; or R¹⁴ and R¹⁶ taken together form a 3- to 6-membered heterocyclyl;


[336] each m is independently 0, 1 or 2;

[337] n₁ and n₂ are each independently 0, 1, or 2;

[338] n₃ is 1, 2, 3, 4 or 5; and

[339] t is 0, 1 or 2.

[340] In one embodiment, the present disclosure provides compounds comprising the structure of formula (III):



[341] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[342] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[343] C is a 3- to 10-membered ring;

[344] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[345] Y is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;

[346] W is a bond, $-(CR^8R^9)_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

[347] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[348] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[349] L is halogen, $-NH_2$, or $-CF_3$;


[350] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;

[351] R^3 is selected from hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;

[352] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, or C_1-C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

- [353] R^7 is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl;
- [354] R^8 and R^9 are each independently hydrogen, halogen, or C₁-C₃ alkyl;
- [355] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, or -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [356] R^{13} , R^{14} and R^{15} are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;
- [357] R^{16} is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;
- [358] each m is independently 0, 1 or 2;
- [359] n₁ and n₂ are each independently 0, 1, or 2;
- [360] n₃ is 0, 1, 2, 3, 4 or 5; and
- [361] t is 0, 1 or 2.

[362] In one embodiment, the present disclosure provides compounds comprising the structure of formula (IIIA):



[363] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

- [364] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;
- [365] C is a 3- to 10-membered ring;
- [366] X is a bond, -(CR⁵R⁶)_t-, or -NR⁷;
- [367] Y is a bond, -(CR⁸R⁹)_m-, -O-, -S-, -S(=O)-, -SO₂-, -NR⁷-, or -N(COCH₃)₂-;
- [368] W is a bond, -(CR^{8a}R^{9a})_m-, -C(=O)-, -N(R⁷)CO-, -CONR⁷-, or -NSO₂R⁷-;
- [369] Z is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)₂-;
- [370] V is -CH₂- and L is halogen, -NH₂, -CHCl₂, -CCl₃, or -CF₃; or
- [371] V is -CH₂CH₂- and L is halogen or -NH₂;

[372] R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶ or optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶;

[373] R³ is selected from halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);

[374] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R⁵ and R⁶ taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[375] R⁷ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl;

[376] R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[377] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, or -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[378] R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R¹⁴ and R¹⁵ taken together form a 3- to 6-membered heterocyclyl;

[379] R¹⁶ is hydrogen, optionally substituted C₁-C₃ alkyl, optionally substituted C₂-C₃ alkenyl, optionally substituted C₂-C₃ alkynyl, C₃-C₆ cycloalkyl, or phenyl;


[380] each m is independently 0, 1 or 2;

[381] n₁ and n₂ are each independently 0, 1, or 2;

[382] n₃ is 1, 2, 3, 4 or 5; and

[383] t is 0, 1 or 2.

[384] In one embodiment, the present disclosure provides compounds comprising the structure of formula (IV):



[385] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[386] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[387] C is a 3- to 10-membered ring;

[388] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[389] Y and Z are each independently a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[390] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[391] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[392] L is halogen, $-NH_2$, or $-CF_3$;

[393] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;

[394] R^3 is selected from hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;

[395] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, or C_1-C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[396] R^7 is H or C_1 - C_6 alkyl;

[397] R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;


[398] R^{16} is hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl;

[399] n_1 and n_2 are each independently 0, 1, or 2;

[400] n_3 is 0, 1, 2, 3, 4 or 5; and

[401] t is 0, 1 or 2.

[402] In one embodiment, the present disclosure provides compounds comprising the structure of formula (V):



[403] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[404] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[405] C is a 5- to 10-membered heteroaryl or aryl;

[406] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[407] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[408] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[409] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[410] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[411] L is halogen, $-NH_2$, or $-CF_3$;

[412] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, -

SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶ or optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶;

[413] R³ is selected from hydrogen, halogen, oxo, =S, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);

[414] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy; or R⁵ and R⁶ taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[415] R⁷ is H or C₁-C₆ alkyl;

[416] R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl; or R¹⁴ and R¹⁵ taken together form a 3- to 6-membered heterocyclyl;


[417] R¹⁶ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;

[418] n₁ and n₂ are each independently 0, 1, or 2;

[419] n₃ is 0, 1, 2, 3, 4 or 5; and

[420] t is 0, 1 or 2.

[421] In one embodiment the present disclosure provides compounds comprising the structure of formula (VA):



[422] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[423] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[424] C is a 5- to 10-membered heteroaryl or aryl;


[425] X is a bond, -(CR⁵R⁶)_t-, or -NR⁷;

[426] Y is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-;

[427] Z is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-;

- [428] W is a bond, -CH₂-, -C(CH₃)H-, -C(=O)-, -N(R⁷)CO-, or -CONR⁷-;
- [429] V is -CH₂-, -CH₂CH₂-, -CH(CH₃)CH₂-, -CH₂CH(CH₃)-, or -CH₂CH₂CH₂-;
- [430] L is hydrogen, halogen, -OH, -NH₂, or -CF₃;
- [431] R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶ or optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶;
- [432] R³ is selected from hydrogen, halogen, oxo, =S, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -NR¹⁴COOR¹⁶, -NR¹⁴CONR¹⁴R¹⁵, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);
- [433] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, -NH₂, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R⁵ and R⁶ taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [434] R⁷ is H, C₁-C₆ alkyl, -CO(C₁-C₆ alkyl);
- [435] R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or -COO(C₁-C₆ alkyl); or R¹⁴ and R¹⁵ taken together form a 3- to 6-membered heterocyclyl;
- [436] R¹⁶ is hydrogen, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;
- [437] n₁ and n₂ are each independently 0, 1, or 2;
- [438] n₃ is 0, 1, 2, 3, 4 or 5; and
- [439] t is 0, 1 or 2.

[440] In one embodiment, the present disclosure provides compounds comprising the structure of formula (VI):



[441] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[442] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[443] C is a 5- to 10-membered heterocyclyl;

[444] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[445] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[446] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[447] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[448] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[449] L is halogen, $-NH_2$, or $-CF_3$;

[450] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)-(C_1 - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} ;

[451] R^3 is selected from hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1$ - C_3 alkyl), C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1$ - C_3 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1$ - C_3 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $SO_2NR^{14}R^{15}$, $-SO_2(C_1$ - C_3 alkyl), or $-(C_1$ - C_6 alkyl)- $SO_2(C_1$ - C_3 alkyl);

[452] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, or C_1 - C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[453] R^7 is H or C₁-C₆ alkyl;

[454] R^{13} , R^{14} and R^{15} are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;


[455] R^{16} is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;

[456] n_1 and n_2 are each independently 0, 1, or 2;

[457] n_3 is 0, 1, 2, 3, 4 or 5; and

[458] t is 0, 1 or 2.

[459] In one embodiment, the present disclosure provides compounds comprising the structure of formula (A):



[460] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[461] C is a phenyl or a 5- to 7-membered monocyclic heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member;

[462] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[463] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[464] Z is a bond, $-CH_2-$, $-O-$, or $-NH-$;

[465] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[466] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[467] L is halogen, $-NH_2$, or $-CF_3$;

[468] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, methyl, or $-CONH_2$;

[469] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $-CN$, $-CF_3$, $-OH$, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, $-S(C_1-C_3 \text{ alkyl})$, $-SO_2(C_1-C_3 \text{ alkyl})$, $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1-C_3 \text{ alkyl})_2$, $-CONH(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, or $-N(CH_3)CO(C_1-C_3 \text{ alkyl})$;

[470] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, or C₁-C₃ alkyl;


[471] R^7 is H or C₁-C₆ alkyl;

[472] n_1 and n_2 are each independently 0, 1, or 2;

[473] n_3 is 0, 1, 2, 3, 4 or 5; and

[474] t is 0, 1 or 2.

[475] In one embodiment, the present disclosure provides compounds comprising the structure of formula (B)



[476] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[477] C is a 5- to 7-membered saturated or partially saturated monocyclic heterocycle comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member;

[478] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[479] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[480] Z is a bond, $-CH_2-$, $-O-$, or $-NH-$;

[481] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[482] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[483] L is halogen, $-NH_2$, or $-CF_3$;

[484] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, methyl, or $-CONH_2$;

[485] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, $-S(C_1-C_3 \text{ alkyl})$, $-SO_2(C_1-C_3 \text{ alkyl})$, $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1-C_3 \text{ alkyl})_2$, $-CONH(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, or $-N(CH_3)CO(C_1-C_3 \text{ alkyl})$;

[486] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, or C_1-C_3 alkyl;

[487] R^7 is H or C_1-C_6 alkyl;

[488] R^{16} is hydrogen or C_1-C_3 alkyl;

[489] n_1 and n_2 are each independently 0, 1, or 2;

[490] n_3 is 0, 1, 2, 3, 4 or 5; and


[491] t is 0, 1 or 2.

[492] In one embodiment of the compounds of formula (I), (IC), (II), (III), (IIIA), (IV), (V), (VI), (A), or (B), -V-L is -CH₂CH₂Cl, -CH₂CH₂CH₂Cl, -CH₂CH₂NH₂, or -CH₂CH₂CH₂NH₂.

[493] In one embodiment of the compounds of formula (I), (IC), (II), (III), (IIIA), (IV), (V), (VI), (A), or (B), -Y-W- is a bond, -OCH₂-, -OCH₂CH₂-, -OCH(CH₃)-, -NH-, -NHCH₂-, -NHC(=O)-, or -C(=O)NH-.

[494] In one embodiment of the compounds of formula (I), (IC), (II), (III), (IIIA), (IV), (V), (VI), (A), or (B), X is a bond, -CH₂-, -C(CH₃)H-, -C(CH₃)₂-, or -CH₂CH₂-.

[495] In one embodiment, the present disclosure provides compounds comprising the structure of formula (C):



[496] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[497] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[498] X is a bond, (CR⁵R⁶)_n-, or -NR⁷-;

[499] Y is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)_n-;

[500] Z is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)_n-;

[501] W is a bond, -CH₂-, -C(CH₃)H-, -C(=O)-, -N(R⁷)CO-, or -CONR⁷-;

[502] V is -CH₂-, -CH₂CH₂-, or -CH₂CH₂CH₂-;

[503] L is halogen, -NH₂, or -CF₃;

[504] D is -NH or -NR³;

[505] U is each independently O, S, or NR¹⁶;

[506] R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -

SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶ or optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶;

[507] R³ is selected from hydrogen, halogen, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);

[508] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R⁵ and R⁶ taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[509] R⁷ is H or C₁-C₆ alkyl;

[510] R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R¹⁴ and R¹⁵ taken together form a 3- to 6-membered heterocyclyl;


[511] R¹⁶ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;

[512] n₁ and n₂ are each independently 0, 1, or 2;

[513] n₃ is 0, 1, 2, or 3; and

[514] t is 0, 1 or 2.

[515] In one embodiment, the present disclosure provides compounds comprising the structure of formula (D):



[516] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[517] C is a 5- or 6-membered heteroaryl comprising 1 or 2 heteroatoms selected from O, S, or N as a ring member;

[518] X is -(CR³R⁶)_t- or -NR⁷-;


[519] Y is a bond, -CH₂-, -O-, or -NH-;

[520] Z is a bond, -CH₂-, -O-, or -NH-;


[521] W is a bond, -CH₂-, or -C(CH₃)H-;

- [522] V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$;
 [523] L is hydrogen, halogen, $-\text{NH}_2$, or $-\text{CF}_3$;
 [524] R^1 and R^2 are each independently hydrogen, halogen, $-\text{CN}$, $-\text{CF}_3$, methyl, or $-\text{CONH}_2$;
 [525] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $-\text{CN}$, $-\text{CF}_3$, $-\text{OH}$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, $-\text{S}(\text{C}_1$ - C_3 alkyl), $-\text{SO}_2(\text{C}_1$ - C_3 alkyl), $-\text{NHSO}_2\text{CH}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{CH}_2\text{NHSO}_2\text{CH}_3$, $-\text{CH}_2\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{SO}_2\text{NH}_2$, $-\text{CONH}_2$, $-\text{CON}(\text{C}_1$ - C_3 alkyl) $_2$, $-\text{CONH}(\text{C}_1$ - C_3 alkyl), $-\text{NHCO}(\text{C}_1$ - C_3 alkyl), or $-\text{N}(\text{CH}_3)\text{CO}(\text{C}_1$ - C_3 alkyl);
 [526] R^5 and R^6 are each independently hydrogen, halogen, $-\text{OH}$, or C_1 - C_3 alkyl;
 [527] R^7 is H or C_1 - C_6 alkyl;
 [528] n_1 and n_2 are each independently 0, 1, or 2;
 [529] n_3 is 0, 1, 2, 3, 4 or 5; and
 [530] t is 0, 1 or 2.

[531] In one embodiment, the present disclosure provides compounds comprising the structure of formula (E):



[532] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



- [534] X is $-(\text{CR}^5\text{R}^6)_t-$ or $-\text{NR}^7-$;
 [535] Y is a bond, $-\text{CH}_2-$, $-\text{O}-$, or $-\text{NH}-$;
 [536] Z is a bond, $-\text{CH}_2-$, $-\text{O}-$, or $-\text{NH}-$;
 [537] W is a bond, $-\text{CH}_2-$, or $-\text{C}(\text{CH}_3)\text{H}-$;
 [538] V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$;

[539] L is hydrogen or halogen;

[540] R^1 and R^2 are each independently hydrogen, halogen, -CN, -CF₃, methyl, or -CONH₂;

[541] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)CO(C₁-C₃ alkyl);

[542] R^5 and R^6 are each independently hydrogen or C₁-C₃ alkyl;


[543] R^7 is H or C₁-C₆ alkyl;

[544] n_1 and n_2 are each independently 0, 1, or 2;


[545] n_3 is 0, 1, or 2; and

[546] t is 1 or 2.

[547] In one embodiment the present disclosure provides compounds comprising the structure of formula (E-I):



[548] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[550] X is -(CR⁵R⁶)_t- or -NR⁷-;


[551] Y is a bond, -CH₂-, -O-, or -NH-;

[552] Z is a bond, -CH₂-, -O-, or -NH-;


- [553] W is a bond, $-\text{CH}_2-$, or $-\text{C}(\text{CH}_3)\text{H}-$;
- [554] V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, or $-\text{CH}_2\text{CHClCH}_2-$;
- [555] L is hydrogen, $-\text{OH}$, or halogen;
- [556] R^1 and R^2 are each independently hydrogen, halogen, $-\text{CN}$, $-\text{CF}_3$, methyl, or $-\text{CONH}_2$;
- [557] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $-\text{CN}$, $-\text{CF}_3$, $-\text{OH}$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-\text{S}(\text{C}_1$ - C_3 alkyl), $-\text{SO}_2(\text{C}_1$ - C_3 alkyl), $-\text{NH}_2$, $-\text{NHSO}_2\text{CH}_3$, $-\text{NHSO}_2\text{CF}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{CH}_2\text{NHSO}_2\text{CH}_3$, $-\text{CH}_2\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{SO}_2\text{NH}_2$, $-\text{CONH}_2$, $-\text{CON}(\text{C}_1$ - C_3 alkyl) $_2$, $-\text{CONH}(\text{C}_1$ - C_3 alkyl), $-\text{NHCO}(\text{C}_1$ - C_3 alkyl), $-\text{N}(\text{CH}_3)\text{COO}(\text{C}_1$ - C_3 alkyl), $-\text{NHCO}(\text{C}_1$ - C_3 alkyl), or $-\text{N}(\text{CH}_3)\text{COO}(\text{C}_1$ - C_3 alkyl);
- [558] R^5 and R^6 are each independently hydrogen or C_1 - C_3 alkyl;
- [559] R^7 is H or C_1 - C_6 alkyl;
- [560] n_1 and n_2 are each independently 0, 1, or 2;
- [561] n_3 is 0, 1, or 2; and
- [562] t is 1 or 2.


[563] In one embodiment of the compounds of formula (A)-(C) or (E-1), R^3 is selected from hydrogen, F, Cl, Br, I, $-\text{CN}$, $-\text{CF}_3$, $-\text{OH}$, methyl, methoxy, $-\text{S}(\text{C}_1$ - C_3 alkyl), $-\text{SO}_2(\text{C}_1$ - C_3 alkyl), $-\text{NH}_2$, $-\text{NHSO}_2\text{CH}_3$, $-\text{NHSO}_2\text{CF}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{SO}_2\text{NH}_2$, $-\text{CONH}_2$, $-\text{CON}(\text{C}_1$ - C_3 alkyl) $_2$, $-\text{CONH}(\text{C}_1$ - C_3 alkyl), $-\text{NHCO}(\text{C}_1$ - C_3 alkyl), or $-\text{NHCO}(\text{C}_1$ - C_3 alkyl).

[564] In one embodiment, the present disclosure provides compounds comprising the structure of formula (F):



[565] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:





[567] X is $-(CR^5R^6)_t-$ or $-NR^7-$;

[568] Y is $-O-$;

[569] Z is $-O-$;

[570] W is $-CH_2-$ or $-C(CH_3)H-$;

[571] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[572] L is hydrogen or halogen;

[573] R^1 and R^2 are each independently halogen or $-CN$;

[574] R^3 is selected from $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, or $-SO_2CH_3$;

[575] R^5 and R^6 are each independently hydrogen or C_1 - C_3 alkyl;


[576] R^7 is H or C_1 - C_6 alkyl;

[577] n_1 and n_2 are each independently 0, 1, or 2;


[578] n_3 is 0, 1, or 2; and

[579] t is 1 or 2

[580] In one embodiment, the present disclosure provides compounds comprising the structure of formula (G):



[581] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:




[583] X is $-(CR^5R^6)_t-$ or $-NR^7-$;


[584] Y is $-O-$;

- [585] Z is -O-;
- [586] W is -CH₂- or -C(CH₃)H-;
- [587] V is -CH₂- and L is hydrogen;
- [588] or alternatively, V is -CH₂CH₂- or -CH₂CH₂CH₂-, and L is halogen;
- [589] R¹ and R² are each independently Cl or -CN;
- [590] R³ is selected from -NHSO₂CH₃, -N(CH₃)SO₂CH₃, or -SO₂CH₃;
- [591] R⁵ and R⁶ are each independently hydrogen or methyl;
- [592] R⁷ is H or C₁-C₆ alkyl;
- [593] n₁ and n₂ are each independently 0, 1, or 2;
- [594] n₃ is 1, or 2; and
- [595] t is 1.

[596] In one embodiment the present disclosure provides compounds comprising the structure of (G-I)



[597] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



- [599] X is -(CR⁵R⁶)_t;
- [600] Y is -O-;
- [601] Z is -O-;
- [602] W is -CH₂- or -C(CH₃)H-;
- [603] V is -CH₂CH₂- or -CH₂CH₂CH₂-;
- [604] L is halogen;
- [605] R¹ and R² are each independently Cl or -CN;

[606] R^3 is selected from $-\text{NHSO}_2\text{CH}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, or $-\text{SO}_2\text{CH}_3$;


[607] R^5 and R^6 are each independently hydrogen or methyl;

[608] n_1 and n_2 are each independently 0, 1, or 2;


[609] n_3 is 1 or 2; and

[610] t is 1.

[611] In one embodiment, the present disclosure provides compounds comprising the structure of formula (G-II)



[612] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[614] X is $-(\text{CR}^5\text{R}^6)_t-$;

[615] Y is $-\text{O}-$;

[616] Z is $-\text{O}-$;

[617] W is $-\text{CH}_2-$ or $-\text{C}(\text{CH}_3)\text{H}-$;

[618] V is $-\text{CH}_2\text{CH}_2-$;

[619] L is halogen;

[620] R^1 and R^2 are each independently Cl or $-\text{CN}$;

[621] at least one R^3 is selected from -CN, C₁-C₃ alkoxy, -CONH₂, -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, or -SO₂CH₃ and the other R^3 , if present, is selected from -CN, -CF₃, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -(C₁-C₃ alkyl)NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl);


[622] R^5 and R^6 are each independently hydrogen or methyl;

[623] n_1 and n_2 are each independently 0, 1, or 2;


[624] n_3 is 1 or 2; and

[625] t is 1.

[626] In one embodiment, the present disclosure provides compounds comprising the structure of formula (H):



[627] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[629] X is -(CR⁵R⁶)_t- or -NR⁷-;

[630] Y is -O-;

[631] Z is -O-;


[632] W is -CH₂- or -C(CH₃)H-;

[633] V is -CH₂-, -CH₂CH₂- or -CH₂CH₂CH₂-;


[634] L is halogen;

- [635] R^1 and R^2 are each independently Cl or -CN;
 [636] R^5 and R^6 are each independently hydrogen or methyl;
 [637] R^7 is H or C₁-C₆ alkyl;
 [638] n_1 and n_2 are each independently 0, 1, or 2; and
 [639] t is 1.

[640] In one embodiment the present disclosure provides compounds comprising the structure of formula (H-I):




[641] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:




- [643] X is $-(CR^5R^6)_t$;
 [644] Y is -O-;
 [645] Z is -O-;
 [646] W is -CH₂- or -C(CH₃)H-;

- [647] V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2\text{CH}_2-$;
 [648] L is halogen;
 [649] R^1 and R^2 are each independently Cl or $-\text{CN}$;
 [650] R^5 and R^6 are each independently hydrogen or methyl;
 [651] R^7 is H or C_1 - C_6 alkyl;
 [652] n_1 and n_2 are each independently 0, 1, or 2; and
 [653] t is 1.

[654] In one embodiment the present disclosure provides compounds comprising the structure of formula (E-II):



[655] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:




- [657] X is $-(\text{CR}^5\text{R}^6)-$;
 [658] Y is a bond, $-\text{CH}_2-$, $-\text{O}-$, or $-\text{NH}-$;
 [659] Z is a bond, $-\text{CH}_2-$, $-\text{O}-$, or $-\text{NH}-$;
 [660] W is a bond, $-\text{CH}_2-$, or $-\text{C}(\text{CH}_3)\text{H}-$;
 [661] V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$;
 [662] L is halogen;
 [663] R^1 , R^{2A} and R^{2B} are each independently hydrogen, halogen, $-\text{CN}$, $-\text{CF}_3$, methyl, or $-\text{CONH}_2$;
 [664] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $-\text{CN}$, $-\text{CF}_3$, $-\text{OH}$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-\text{S}(\text{C}_1$ - C_3 alkyl), $-\text{SO}_2(\text{C}_1$ - C_3 alkyl), $-\text{NH}_2$, $-\text{NHSO}_2\text{CH}_3$, $-\text{NHSO}_2\text{CF}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{CH}_2\text{NHSO}_2\text{CH}_3$, $-\text{CH}_2\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, -


SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl);

[665] R⁵ and R⁶ are each independently hydrogen or C₁-C₃ alkyl;


[666] n₁ is 0, 1, or 2; and

[667] n₃ is 1 or 2;




[668] wherein when C-R³ is , R^{2A} and R^{2B} are not both Cl.

[669] In one embodiment the present disclosure provides compounds comprising the structure of formula (E-III):



[670] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[672] X is -(CR⁵R⁶)-;

[673] Y is a bond, -CH₂-, -O-, or -NH-;

[674] Z is a bond, -CH₂-, -O-, or -NH-;

[675] W is a bond, -CH₂-, or -C(CH₃)H-;

[676] V is -CH₂-, -CH₂CH₂-, or -CH₂CH₂CH₂-;

[677] L is halogen;


[678] R¹, R^{2A} and R^{2B} are each independently hydrogen, halogen, -CN, -CF₃, methyl, or -CONH₂;


[679] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl);

[680] R^5 and R^6 are each independently hydrogen or C₁-C₃ alkyl;


[681] n_1 is 0, 1, or 2; and

[682] n_3 is 1 or 2;




[683] wherein when C- R^3 is  and one of R^{2A} and R^{2B} is Cl, then the other of R^{2A} and R^{2B} is not -CN.

[684] In one embodiment the present disclosure provides compounds comprising the structure of formula (E-IV):



[685] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[687] X is -(CR⁵R⁶)-;

[688] Y is a bond, -CH₂-, -O-, or -NH-;

[689] Z is a bond, -CH₂-, -O-, or -NH-;

[690] W is a bond, -CH₂-, or -C(CH₃)H-;

[691] V is -CH₂-, -CH₂CH₂-, or -CH₂CH₂CH₂-;

[692] L is halogen;

[693] R^1 , R^{2A} and R^{2B} are each independently hydrogen, halogen, -CN, -CF₃, methyl, or -CONH₂;


[694] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl);

[695] R^5 and R^6 are each independently hydrogen or C₁-C₃ alkyl;


[696] n_1 is 0, 1, or 2; and

[697] n_3 is 1 or 2.

[698] In one embodiment the present disclosure provides compounds comprising the structure of formula (E-V):



[699] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[701] X is -(CR⁵R⁶)-;

[702] Y is a bond, -CH₂-, -O-, or -NH-;

[703] Z is a bond, -CH₂-, -O-, or -NH-;

[704] W is a bond, -CH₂-, or -C(CH₃)H-;

[705] V is -CH₂-, -CH₂CH₂-, or -CH₂CH₂CH₂-;

[706] L is halogen;


[707] R^1 , R^{2A} and R^{2B} are each independently hydrogen, halogen, -CN, -CF₃, methyl, or -CONH₂;

[708] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl);


[709] R^5 and R^6 are each independently hydrogen or C₁-C₃ alkyl;

[710] n1 is 0, 1, or 2; and


[711] n3 is 1 or 2;

[712] wherein when C-R³ is , R^{2A} and R^{2B} are not both Cl.

[713] In one embodiment the present disclosure provides compounds comprising the structure of formula (E-VI):



[714] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[715] C is , or

[716] X is -(CR⁵R⁶)-;

[717] Y is a bond, -CH₂-, -O-, or -NH-;

[718] Z is a bond, -CH₂-, -O-, or -NH-;

[719] W is a bond, -CH₂-, or -C(CH₃)H-;

[720] V is -CH₂-, -CH₂CH₂-, or -CH₂CH₂CH₂-;

[721] L is halogen;


[722] R^1 , R^{2A} and R^{2B} are each independently hydrogen, halogen, -CN, -CF₃, methyl, or -CONH₂;

[723] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl);


[724] R^5 and R^6 are each independently hydrogen or C₁-C₃ alkyl;

[725] n₁ is 0, 1, or 2; and


[726] n₃ is 1 or 2;

[727] wherein when C-R³ is  and one of R^{2A} and R^{2B} is Cl, then the other of R^{2A} and R^{2B} is not -CN.

[728] In one embodiment the present disclosure provides compounds comprising the structure of formula (E-VII):



[729] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[731] X is -(CR⁵R⁶)-;

[732] Y is a bond, -CH₂-, -O-, or -NH-;

[733] Z is a bond, -CH₂-, -O-, or -NH-;

[734] W is a bond, $-\text{CH}_2-$, or $-\text{C}(\text{CH}_3)\text{H}-$;

[735] V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$;

[736] L is halogen;

[737] R^1 , $\text{R}^{2\text{A}}$ and $\text{R}^{2\text{B}}$ are each independently hydrogen, halogen, $-\text{CN}$, $-\text{CF}_3$, methyl, or $-\text{CONH}_2$;

[738] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $-\text{CN}$, $-\text{CF}_3$, $-\text{OH}$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-\text{S}(\text{C}_1$ - C_3 alkyl), $-\text{SO}_2(\text{C}_1$ - C_3 alkyl), $-\text{NH}_2$, $-\text{NHSO}_2\text{CH}_3$, $-\text{NHSO}_2\text{CF}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{CH}_2\text{NHSO}_2\text{CH}_3$, $-\text{CH}_2\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{SO}_2\text{NH}_2$, $-\text{CONH}_2$, $-\text{CON}(\text{C}_1$ - C_3 alkyl) $_2$, $-\text{CONH}(\text{C}_1$ - C_3 alkyl), $-\text{NHCO}(\text{C}_1$ - C_3 alkyl), $-\text{N}(\text{CH}_3)\text{COO}(\text{C}_1$ - C_3 alkyl), $-\text{NHCO}(\text{C}_1$ - C_3 alkyl), or $-\text{N}(\text{CH}_3)\text{COO}(\text{C}_1$ - C_3 alkyl);

[739] R^5 and R^6 are each independently hydrogen or C_1 - C_3 alkyl;


[740] n_1 is 0, 1, or 2; and

[741] n_3 is 1 or 2.

[742] In one embodiment of the compounds of formula (I), (IA), (IB), (IC), (II), (IIA), (IIB), (III), (IIIA), (IV), (V), (VA), or (VI) (denoted as formula “(I)-(VI)”), A and B are each independently 5- or 6-membered aryl or heteroaryl. In one embodiment, A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene. In one embodiment, A and B are each phenyl.

[743] In another embodiment, A has a meta or para connectivity with X and Y. In some embodiments, B has a meta or para connectivity with X and Z.

[744] In one embodiment of the compounds of formula (I)-(VI), A and B are phenyl and has one of the connectivity as shown:




[745] In one embodiment of the compounds of formula (I)-(VA) and (A) (e.g., formula (I), (IA), (IB), (IC), (II), (IIA), (IIB), (III), (IIIA), (IV), (V) (VA), and (A)), C is aryl or heteroaryl. In some embodiments, C is 5- to 10-membered aryl or heteroaryl. In other embodiments, C is



aryl. In some embodiments, C is phenyl or naphthyl. In other embodiments, C is aryl. In some embodiments, C is phenyl.


[746] In one embodiment of the compounds of formula (I)-(VA) and (A), C is heteroaryl. In one embodiment, C monocyclic or bicyclic heteroaryl. In another embodiment, C is monocyclic heteroaryl. In some embodiments, C is 5- or 10-membered heteroaryl. In some embodiments, C is 5- or 6-membered heteroaryl, which is optionally substituted with 1, 2, 3, 4, or 5 R^3 . In some embodiments, C is 5- or 6-membered heteroaryl containing 1, 2, or 3 heteroatoms selected from O, S, or N, wherein the heteroaryl is optionally substituted with 1, 2, 3, 4, or 5 R^3 . In some embodiments, C is 5- or 6-membered heteroaryl containing 1 or 2 heteroatoms selected from O, S, or N, wherein the heteroaryl is optionally substituted with 1, 2, 3, 4, or 5 R^3 .

[747] In one embodiment of the compounds of formula (I)-(VA), (A), or (D), C is pyrazole, imidazole, oxazole, oxadiazole, oxazolone, isoxazole, thiazole, pyridyl, or pyrimidyl, which are each optionally substituted with 1, 2, 3, 4, or 5 R^3 . In one embodiment, C, which is substituted with $(R^3)_{n3}$, is pyrazole, imidazole, oxazole, oxadiazole, oxazolone, isoxazole, thiazole, pyridyl, pyrazine, furan or pyrimidyl. In one embodiment, C is pyrazole, imidazole, oxazole, oxadiazole, oxazolone, isoxazole, thiazole, pyridyl, pyrazine, furan or pyrimidyl, which are each substituted with 1, 2, 3, 4, or 5 R^3 .


[748] In one embodiment of the compounds of formula (I)-(VA), (A), or (D), C is selected from








[750] In one embodiment of the compounds of formula (I)-(VA), (A), or (D), C is



[751] In one embodiment, C is




In one embodiment, C is




[752] In one embodiment of the compounds of formula (I)-(IV), C is heterocyclyl. In one embodiment, C is saturated or partially saturated heterocycle. In some embodiments, C is monocyclic or bicyclic. In some embodiments, C is 5- to 7-membered heterocyclyl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

[753] In one embodiment of the compounds of formula (I)-(VA), (B) and (C), C is imidazolidine, imidazolidine-dione, or dihydrooxazole. In one embodiment, C is selected from




[754] In one embodiment of the compounds of formula (I)-(VA), (B) and (C), C is




; D is $-\text{O}-$, $-\text{NH}-$ or $-\text{NR}^3-$; and U is each independently O, S, or NR^{16} . In one embodiment, D is $-\text{NH}-$ or $-\text{NR}^3-$. In some embodiments, at least one U is O. In other embodiments, each U is O. In some embodiments, at least one R^3 is $-\text{SO}_2\text{CH}_3$, $-\text{NHSO}_2\text{CH}_3$, $-\text{CH}_2\text{NHSO}_2\text{CH}_3$, $-\text{SO}_2\text{NH}_2$, $-\text{CONH}_2$, or $-\text{NHCOCH}_3$.

[755] In one embodiment of the compounds of formula (I)-(VA), C is aryl. In some embodiments, C is phenyl or naphthyl. In one embodiment of the compounds of formula (I)-(VA) or (A), C is phenyl.

[756] In one embodiment compounds of formula (I)-(VI), C is bicyclic heteroaryl or



heterocyclyl. In one embodiment, C is



[757] In one embodiment of the compounds of formula (I), Z is a bond, $-(\text{CR}^8\text{R}^9)_m-$, $-\text{O}-$, $-\text{C}(=\text{O})-$, $-\text{S}-$, $-\text{S}(=\text{O})-$, $-\text{SO}_2-$, or $-\text{NR}^7-$. In one embodiment, Z is $-(\text{CR}^8\text{R}^9)_m-$, $-\text{O}-$, $-\text{C}(=\text{O})-$, $-\text{S}-$, $-\text{S}(=\text{O})-$, $-\text{SO}_2-$, or $-\text{NR}^7-$. In some embodiments, Z is $-\text{C}(=\text{O})-$.

[758] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), Z is a bond, $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{NCH}_3-$, or $-\text{N}(\text{COCH}_3)-$. In one embodiment, Z is $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{NCH}_3-$, or $-\text{N}(\text{COCH}_3)-$. In some embodiments, Z is a bond, $-\text{CH}_2-$, $-\text{O}-$, or $-\text{NCH}_3-$. In some embodiments, Z is a bond, $-\text{CH}_2-$, $-\text{O}-$, or $-\text{NH}-$. In some embodiments of the compounds of formula (I)-(VI) and (A)-(H-I), Z is $-\text{O}-$. As used herein, “compounds of formula (I)-(IV) and (A)-(H-I)” refers to compounds of formula (I), (IA), (IB), (IC), (II), (IIA), (IIIA), (IIB), (IIB), (III), (IV), (IVA), (V), (VA), (VI), (A), (A-I), (B)-(D), (E), (E-I)-(E-VII), (F), (G), (G-I), (G-II), (H), and (H-I).

[759] In one embodiment of the compounds of formula (I), Y is a bond, $-(\text{CR}^8\text{R}^9)_m-$, $-\text{O}-$, $-\text{C}(=\text{O})-$, $-\text{S}-$, $-\text{S}(=\text{O})-$, $-\text{SO}_2-$, or $-\text{NR}^7-$. In one embodiment, Y is $-(\text{CR}^8\text{R}^9)_m-$, $-\text{O}-$, $-\text{C}(=\text{O})-$, $-\text{S}-$, $-\text{S}(=\text{O})-$, $-\text{SO}_2-$, or $-\text{NR}^7-$. In some embodiments, Y is $-\text{C}(=\text{O})-$.

[760] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), Y is a bond, $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{NCH}_3-$, or $-\text{N}(\text{COCH}_3)-$. In one embodiment, Y is $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{NCH}_3-$, or $-\text{N}(\text{COCH}_3)-$. In some embodiments, Y is a bond, $-\text{CH}_2-$, $-\text{O}-$, or $-\text{NH}-$. In some embodiments of the compounds of formula (I)-(VI) and (A)-(H-I), Y is $-\text{O}-$.

[761] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), V is a bond, $-(\text{CR}^{8a}\text{R}^{9a})_m-$, or $-\text{C}(=\text{O})-$. In some embodiments, V is bond, or $-(\text{CR}^{8a}\text{R}^{9a})_m-$.

[762] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), V is $-(\text{CR}^{8a}\text{R}^{9a})_m-$, wherein m is 1, 2, or 3. In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), V is $-(\text{CR}^{8a}\text{R}^{9a})_m-$, wherein R^{8a} and R^{8b} are each independently hydrogen, $-\text{OH}$, halogen, optionally substituted C_1 - C_3 alkyl, optionally substituted C_2 - C_3 alkenyl, optionally substituted C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-\text{NR}^{13}\text{R}^{14}$, optionally substituted $-(\text{C}_1$ - C_3 alkyl)- $\text{NR}^{13}\text{R}^{14}$, $-\text{NR}^{14}\text{COR}^{16}$, optionally substituted $-(\text{C}_1$ - C_3 alkyl)- $\text{NR}^{14}\text{COR}^{16}$, $-\text{CONR}^{14}\text{R}^{15}$, or optionally substituted $-(\text{C}_1$ - C_3 alkyl)- $\text{CONR}^{14}\text{R}^{15}$; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl. In one embodiment, V is $-(\text{CR}^{8a}\text{R}^{9a})_m-$, wherein R^{8a} and R^{8b} are each independently hydrogen, $-\text{OH}$, halogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-\text{NR}^{13}\text{R}^{14}$, $-(\text{C}_1$ - C_3 alkyl)- $\text{NR}^{13}\text{R}^{14}$, $-\text{NR}^{14}\text{COR}^{16}$, $-(\text{C}_1$ - C_3 alkyl)- $\text{NR}^{14}\text{COR}^{16}$, $-\text{CONR}^{14}\text{R}^{15}$, or $-(\text{C}_1$ - C_3 alkyl)- $\text{CONR}^{14}\text{R}^{15}$; or R^{8a} and R^{8b} , on the same carbon atom or on a different carbon atom, taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl.

[763] In one embodiment of the compounds of formula (I)-(IIB) and (VA), V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}(\text{CH}_3)\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_3)-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$.

[764] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$, each optionally substituted with one or more of $-\text{OH}$, halogen, or C_1 - C_3 alkyl. In other embodiments, V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{OH})\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2\text{CH}_2-$. In some embodiments of the compounds of formula (I)-(VI) and (A)-(H-I), V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$. In some embodiments of the compounds of formula (I)-(VI) and (A)-(H-I), V is $-\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2-$.

[765] In some embodiments of the compounds of formula (I)-(VI) and (A)-(D), V is $-\text{CH}_2-$ and L is halogen, $-\text{NH}_2$, or $-\text{CF}_3$; or V is $-\text{CH}_2\text{CH}_2-$ and L is halogen or $-\text{NH}_2$.

[766] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), L is hydrogen, halogen, $-\text{CF}_2\text{H}$, $-\text{CF}_3$, $-\text{CN}$, $-\text{O}(\text{C}_1$ - C_3 alkyl), $-\text{NR}^{11}\text{R}^{12}$, or $-\text{CONR}^{11}\text{R}^{12}$. In one embodiment, L is hydrogen, halogen, $-\text{CF}_2\text{H}$, $-\text{CF}_3$, $-\text{CN}$, $-\text{O}(\text{C}_1$ - C_3 alkyl), $-\text{NH}_2$, $-\text{NH}(\text{C}_1$ - C_3

alkyl), -N(C₁-C₃ alkyl)₂, -CONH₂, -CONH(C₁-C₃ alkyl), or -CON(C₁-C₃ alkyl)₂. In some embodiments, L is hydrogen, halogen, -CF₃, or -NH₂.

[767] In some embodiments of the compounds of formula (IC) and (IIIA), L is halogen, -CCl₃, -CCl₂, -CF₃, or -NH₂. In some embodiments of the compounds of formula (I)-(VI) and (A)-(H-I), L is halogen, -CF₃, or -NH₂. In one embodiment, L is hydrogen or halogen. In one embodiment, L is halogen. In other embodiments, L is Cl, or Br. In one embodiment, L is Cl.

[768] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), W is a bond. In one embodiment, W is -(CR^{8a}R^{9a})_m-, -C(=O)-, -N(R⁷)CO-, -CONR⁷-, or -NSO₂R⁷-. In one embodiment, W is -(CR^{8a}R^{9a})_m-, wherein m is 1, 2, or 3. In some embodiments, W is a bond, -CH₂-, -C(CH₃)H-, -C(=O)-, -N(R⁷)CO-, or -CONR⁷-, wherein R⁷ is H or C₁-C₆ alkyl. In some embodiments, W is a bond, -CH₂-, -C(CH₃)H-, -C(=O)-, -N(R⁷)CO-, or -CONR⁷-, wherein R⁷ is H or C₁-C₃ alkyl. In one embodiment, W is a bond, -CH₂-, -C(CH₃)H-, -C(=O)-, -NHCO-, -N(C₁-C₃ alkyl)CO-, -CONH-, or -CON(C₁-C₃ alkyl)-. In one embodiment of the compounds of formula (I)-(VI) and (A)-(E-VI), W is a bond, -CH₂-, or -C(CH₃)H-. In one embodiment of the compounds of formula (I)-(VI) and (A)-(H-I), W is a -CH₂- or -C(CH₃)H-.

[769] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), -Y-W- is a bond, -OCH₂-, -OCH₂CH₂-, -OCH(CH₃)-, -NH-, -NHCH₂-, -NHC(=O)-, or -C(=O)NH-. In one embodiment of the compounds of formula (I)-(VI) and (A)-(H-I), -Y-W- is -OCH₂-, -OCH₂CH₂-, or -OCH(CH₃)-.

[770] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C),

[771] Z is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-; and

[772] V is -CH₂-, -CH₂CH₂-, or -CH₂CH₂CH₂-.

[773] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C),

[774] Z is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-;

[775] V is -CH₂-, -CH₂CH₂-, or -CH₂CH₂CH₂-; and

[776] L is halogen, -NH₂, or -CF₃.

[777] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), -Z-V-L is -Z-CH₂CH₂Cl, -Z-CH₂CH₂CH₂Cl, -Z-CH₂CH₂NH₂, or -Z-CH₂CH₂CH₂NH₂, wherein Z is a bond, -O-, -NH-, or -N(COCH₃)-. In one embodiment, -Z-V-L is -OCH₃.

[778] In one embodiment of the compounds of formula (I)-(VI) and (A)-(H-I), -Z-V-L is -O-CH₂CH₂Cl or -O-CH₂CH₂CH₂Cl.

[779] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), -V-L is -CH₂CH₂Cl, -CH₂CH₂CH₂Cl, -CH₂CH₂NH₂, or -CH₂CH₂CH₂NH₂. In one embodiment, -V-L is -CH₃.

[780] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), X is a bond, $-(CR^5R^6)_t-$, $-O-$, $-C(=O)-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$. In one embodiment, X is a bond, $-(CR^5R^6)_t-$, $-O-$, $-C(=O)-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$, wherein R^7 is H or C₁-C₆ alkyl. In some embodiments, X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$. In some embodiments, X is a bond or $-(CR^5R^6)_t-$. In some embodiments, X is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(CH_3)_2-$, $-CH_2CH_2-$, $-NH-$, or $-N(C_1-C_6 \text{ alkyl})-$. In some embodiments, X is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(CH_3)_2-$, $-CH_2CH_2-$, $-NH-$, $-N(CH_3)-$, $-N(CH_2CH_3)-$, $-N(iPr)-$, or $-N(tBu)-$. In some embodiments, X is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(CH_3)_2-$, or $-CH_2CH_2-$.

[781] In one embodiment of the compounds of formula (I)-(VI) and (A)-(H-I), X is $-CH_2-$, $-C(CH_3)H-$, or $-C(CH_3)_2-$. In one embodiment, X is $-C(CH_3)_2-$.

[782] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R^1 and R^2 are each independently halogen, $-CN$, $-CF_3$, $-OH$, C₁-C₃ alkyl, C₁-C₃ alkoxy, $-(C_1-C_3 \text{ alkyl})-(C_1-C_3 \text{ alkoxy})$, $-(C_1-C_3 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_3 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})-SO_2R^{16}$, 3- to 7-membered carbocyclyl, 3- to 7-membered heterocyclyl, phenyl, or 5- to 6-membered heteroaryl.

[783] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$, or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$. In one embodiment, R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C₁-C₃ alkyl, C₁-C₃ alkoxy, optionally substituted $-(C_1-C_3 \text{ alkyl})-(C_1-C_3 \text{ alkoxy})$, optionally substituted $-(C_1-C_3 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_3 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2R^{16}$, or optionally substituted $-(C_1-C_3 \text{ alkyl})-SO_2R^{16}$. In one embodiment, R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, C₁-C₃ alkyl, C₁-C₃ alkoxy, $-(C_1-C_3 \text{ alkyl})-(C_1-C_3 \text{ alkoxy})$, $-(C_1-C_3 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_3 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2R^{16}$, or optionally substituted $-(C_1-C_3 \text{ alkyl})-SO_2R^{16}$.


alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂R¹⁶, or -(C₁-C₃ alkyl)-SO₂R¹⁶.

[784] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, C₁-C₃ alkyl, or -CONR¹⁴R¹⁵. In some embodiments, R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, methyl, methoxy, or -CONH₂. In one embodiment, R¹ and R² are each independently hydrogen, Cl, -CN, -CF₃, -OH, methyl, methoxy, or -CONH₂. In one embodiment, R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, or methyl. In one embodiment, R¹ and R² are each independently Cl, -CN, -CF₃, -OH, methyl, methoxy, or -CONH₂. In one embodiment of the compounds of formula (I)-(VI), R¹ and R² are each independently halogen, -CN, -CF₃, -OH, or methyl.


[785] In one embodiment of the compounds of formula (I)-(VI) and (A)-(E-I), R¹ and R² are each halogen, methyl, -CF₃, or -CN. In one embodiment of the compounds of formula (I)-(VI) and (A)-(F), R¹ and R² are each halogen or -CN. In one embodiment of the compounds of formula (I)-(VI) and (A)-(H-I), at least one of R¹ and R² is Cl or -CN. In one embodiment of the compounds of formula (I)-(VI) and (A)-(H-I), at least two of R¹ and R² are each independently Cl or -CN. In one embodiment of the compounds of formula (I)-(VI) and (A)-(H-I), R¹ and R² are each Cl or -CN.

[786] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R¹ and R² are each independently optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In one embodiment, R¹ and R² are each independently 3- to 7-membered carbocyclyl, 3- to 7-membered heterocyclyl, phenyl, or 5- to 6-membered heteroaryl.

[787] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R¹ have one of the connectivity as shown below with respect to X and Y:



[788] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R² have one of the connectivity as shown below with respect to X and Z:



[789] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), n_1 is 0, 1, or 2. In some embodiments, n_1 is 0 or 1. In other embodiments, n_1 is 0. In some embodiments, n_1 is 1. In one embodiment, the sum of n_1 and n_2 is 0, 1, 2, 3, or 4. In some embodiments, the sum of n_1 and n_2 is 1, 2, 3, or 4. In one embodiment, the sum of n_1 and n_2 is 2.

[790] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), n_2 is 0, 1, or 2. In some embodiments, n_2 is 1 or 2. In other embodiments, n_2 is 0. In some embodiments, n_2 is 1. In some embodiments, n_2 is 2.

[791] In some embodiments of the compounds of (I)-(VI), (A), (A-I), (B), and (C), n_3 is 1, 2, 3, 4, or 5. In some embodiments, n_3 is 1, 2, 3, or 4. In one embodiment, n_3 is 1, 2, or 3. In one embodiment, n_3 is 1 or 2.

[792] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R^3 is selected from hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -SR¹⁶, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In some embodiments, R^3 is selected from hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -SR¹⁶, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-

NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, or optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶. In another embodiment, R³ is hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -SR¹⁶, optionally substituted C₁-C₃ alkyl, optionally substituted C₂-C₃ alkenyl, optionally substituted C₂-C₃ alkynyl, optionally substituted C₁-C₃ alkoxy, optionally substituted -(C₁-C₃ alkyl)-(C₁-C₃ alkoxy), optionally substituted -(C₁-C₃ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, or optionally substituted -(C₁-C₃ alkyl)-SO₂R¹⁶.

[793] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R³ is selected from -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, or optionally substituted -SO₂R¹⁶; wherein R¹⁶ is hydrogen, optionally substituted C₁-C₃ alkyl, optionally substituted C₂-C₃ alkenyl, optionally substituted C₂-C₃ alkynyl, C₃-C₆ cycloalkyl, or phenyl. In one embodiment, R³ is selected from -NR¹⁴SO₂R¹⁶, -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, or -SO₂R¹⁶; wherein R¹⁶ is hydrogen, C₁-C₃ alkyl, -(C₁-C₃ alkyl)-NH₂, C₃-C₆ cycloalkyl, or phenyl.

[794] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R³ is selected from hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -(C₁-C₃ alkyl)-(C₁-C₃ alkoxy), -(C₁-C₃ alkyl)-OH, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl). In some embodiments, R³ is selected from hydrogen, F, Cl, Br, I, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, -S(C₁-C₃ alkyl), -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)CO(C₁-C₃ alkyl). In some embodiments, R³ is selected from hydrogen, F, Cl, Br, I, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, -S(C₁-C₃ alkyl), -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)CO(C₁-C₃ alkyl). In one embodiment, R³ is selected from hydrogen, F, Cl, Br, I, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, methyl, -SCH₃, -SO₂CH₃, -NHSO₂CH₃, -NHSO₂CH₂CH₃, -CH₂NHSO₂CH₃, -SO₂NH₂, -

CONH₂, or -NHCOCH₃. In one embodiment, R³ is selected from hydrogen, F, Cl, Br, I, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, methyl, -SCH₃, -SO₂CH₃, -NHSO₂CH₃, -CH₂NHSO₂CH₃, -SO₂NH₂, -CONH₂, or -NHCOCH₃. In some embodiments, R³ is selected from F, Cl, Br, I, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, -S(C₁-C₃ alkyl), -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)CO(C₁-C₃ alkyl). In some embodiments, R³ is selected from F, Cl, Br, I, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, -S(C₁-C₃ alkyl), -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)CO(C₁-C₃ alkyl). In one embodiment, R³ is selected from F, Cl, Br, I, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, methyl, -SCH₃, -SO₂CH₃, -NHSO₂CH₃, -CH₂NHSO₂CH₃, -SO₂NH₂, -CONH₂, or -NHCOCH₃. In another embodiment, R³ is -SO₂CH₃, -NHSO₂CH₃, -CH₂NHSO₂CH₃, -SO₂NH₂, -CONH₂, or -NHCOCH₃. In one embodiment, R³ is oxo, =S, =NR¹⁶, C₁-C₃ alkyl, -SO₂(C₁-C₃ alkyl), or -NHSO₂(C₁-C₃ alkyl). In one embodiment, at least one of R³ is oxo, =S, or =NR¹⁶. In one embodiment, at least one of R³ is oxo, =S, or =NR¹⁶, wherein R¹⁶ is H or C₁-C₃ alkyl.

[795] In one embodiment of the compounds of formula (I)-(VI) and (A)-(E-VII), R³ is selected from hydrogen, F, Cl, Br, I, oxo, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl). In one embodiment of the compounds of formula (I)-(VI) and (A)-(E-VII), R³ is selected from hydrogen, F, Cl, Br, I, oxo, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl).

[796] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R³ on a sp³ carbon is each selected from hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -(C₁-C₃ alkyl)-(C₁-C₃ alkoxy), -(C₁-C₃ alkyl)-OH, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-

CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl). When R³ on a sp³ carbon is oxo, =S, or =NR¹⁶, the carbon becomes sp².

[797] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R³ on a sp² carbon is each selected from hydrogen, halogen, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -(C₁-C₃ alkyl)-(C₁-C₃ alkoxy), -(C₁-C₃ alkyl)-OH, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl).


[798] In one embodiment of the compounds of formula (I)-(VI), (A), (A-I), (B), and (C), R³ on a nitrogen atom is each selected from C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -(C₁-C₃ alkyl)-(C₁-C₃ alkoxy), -(C₁-C₃ alkyl)-OH, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl).

[799] In one embodiment of the compounds of formula (I)-(VI) and (A)-(G-II), at least one R³ is selected from -CN, C₁-C₃ alkoxy, -CONH₂, -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, or -SO₂CH₃ and the other R³, if present, is selected from -CN, -CF₃, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -(C₁-C₃ alkyl)NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl). In one embodiment, at least one R³ is selected from -NHSO₂CH₃, -NHSO₂CH₂CH₃, or -SO₂CH₃ and the other R³, if present, is selected from -CN, C₁-C₃ alkyl, C₁-C₃ alkoxy, -SO₂(C₁-C₃ alkyl), -NH₂, -(C₁-C₃ alkyl)NH₂, -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl).

[800] In one embodiment of the compounds of formula (I)-(VI) and (A)-(E-II), R³ is not hydrogen.

[801] In one embodiment of the compounds of formula (I)-(VI) and (A)-(G-II), at least one R³ is -SO₂CH₃, -NHSO₂CH₃, -NCH₃SO₂CH₃, -NHSO₂CH₂CH₃, or -N(CH₃)SO₂CH₂CH₃. In one embodiment of the compounds of formula (I)-(VI) and (A)-(G-II), at least one R³ is -SO₂CH₃, -NHSO₂CH₃, or -NCH₃SO₂CH₃.

[802] In one embodiment the compounds of formula (I), (IA), (IB), or (IC), R^3 is heterocyclyl.



[803] In one embodiment of the compounds of formula (IB), (IC), (IIA), or (IIB), R^3 is $-NR^{14}SO_2R^{16}$, wherein R^{14} and R^{16} together form a 5 or 6 membered ring including the nitrogen and sulfur atoms.

[804] In one embodiment of the compounds of formula (I), (IA), (IB), or (IC), R^3 is $-NR^{14}SO_2R^{16}$, wherein R^{16} is optionally substituted C_1 - C_6 alkyl. In one embodiment, R^3 is $-NR^{14}SO_2R^{16}$, wherein R^{16} is C_1 - C_6 alkyl optionally substituted with one or more groups selected from halogen, $-CN$, $-CF_3$, $-OH$, C_1 - C_3 alkyl, C_1 - C_3 alkoxy, $-NH_2$, $-NH(C_1$ - C_3 alkyl), $-N(C_1$ - C_3 alkyl) $_2$, $-SCH_3$. In one embodiment, R^3 is $-NR^{14}SO_2R^{16}$, wherein R^{16} is C_1 - C_3 alkyl substituted with $-NH_2$.

[805] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, or C_1 - C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl. In some embodiments, R^5 and R^6 are each independently hydrogen, halogen, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl. In one embodiment, R^5 and R^6 are hydrogen, halogen, $-OH$, or C_1 - C_3 alkyl. In one embodiment, R^5 and R^6 are each independently hydrogen, F, $-OH$, or C_1 - C_3 alkyl. In one embodiment, R^5 and R^6 are each independently, hydrogen, F, $-OH$, or methyl. In one embodiment, R^5 and R^6 are each H. In one embodiment, R^5 and R^6 are each methyl. In one embodiment of the compounds of formula (I)-(VI) and (A)-(H-I), R^5 and R^6 are each H or methyl.

[806] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R^7 is hydrogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In some embodiments, R^7 is hydrogen, optionally substituted C_1 - C_3 alkyl, optionally substituted C_2 - C_3 alkenyl, optionally substituted C_2 - C_3 alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In some embodiments, R^7 is hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl. In some embodiments, R^7 is hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl. In some embodiments,

R⁷ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl. In some embodiments, R⁷ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl. In some embodiments, R⁷ is hydrogen or C₁-C₆ alkyl. In some embodiments, R⁷ is hydrogen or C₁-C₄ alkyl. In some embodiments of the compounds of formula (I)-(VI) and (A)-(H-I), R⁷ is hydrogen or C₁-C₃ alkyl.

[807] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R^{8a} and R^{9a} are each independently hydrogen, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl, or R^{8a} and R^{9a} taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl. In some embodiments, R^{8a} and R^{8b} are each independently hydrogen, -OH, halogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, or -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl. In one embodiment, R^{8a} and R^{9a} are each independently hydrogen, halogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, or -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵. In one embodiment of the compounds of formula (IB), (IC), (III), or (IIIA), R^{8a} and R^{9a} are not -OH. In one embodiment, R^{8a} and R^{9a} are not -OH.

[808] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R⁷ and R^{8a} taken together form an optionally substituted heterocyclyl. In one embodiment, R⁷ and R^{8a} taken together form an optionally substituted 3- to 7-membered heterocycle.

[809] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl.

[810] In one embodiment of the compounds of formula (I)-(IIB), R¹⁰ is hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In some embodiments, R¹⁰ is hydrogen, optionally substituted C₁-C₃ alkyl, optionally substituted C₂-C₃ alkenyl, optionally substituted C₂-C₃ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In some embodiments, R¹⁰ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, carbocyclyl, heterocyclyl, aryl, or

heteroaryl. In some embodiments, R¹⁰ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl. In some embodiments, R¹⁰ is hydrogen or C₁-C₃ alkyl.

[811] In one embodiment of the compounds of formula (I)-(IIB), R¹¹ and R¹² are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In one embodiment, R¹¹ and R¹² are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, or optionally substituted C₂-C₆ alkynyl. In some embodiments, R¹¹ and R¹² are each independently hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl. In some embodiments, R¹¹ and R¹² are each independently hydrogen or C₁-C₃ alkyl.

[812] In one embodiment of the compounds of formula (I)-(IIB), R¹¹ and R¹² taken together form an optionally substituted heterocyclyl. In one embodiment, R¹¹ and R¹² taken together form an optionally substituted 3- to 7-membered heterocyclyl. In other embodiments, R¹¹ and R¹² taken together form 3- to 7-membered heterocyclyl.

[813] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R¹³ and R¹⁴ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In one embodiment, R¹³ and R¹⁴ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, or optionally substituted C₂-C₆ alkynyl. In some embodiments R¹³ and R¹⁴ are each independently hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl. In some embodiments R¹³ and R¹⁴ are each independently hydrogen or C₁-C₃ alkyl.

[814] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R¹⁵ is hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In one embodiment, R¹⁵ is hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, or optionally substituted C₂-C₆ alkynyl. In some embodiments, R¹⁵ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl. In some embodiments, R¹⁵ is hydrogen or C₁-C₃ alkyl.

[815] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R¹⁴ and R¹⁵ taken together form an optionally substituted heterocyclyl. In one embodiment, R¹⁴ and R¹⁵ taken together form an optionally substituted 3- to 7-membered heterocyclyl. In other embodiments, R¹⁴ and R¹⁵ taken together form 3- to 7-membered heterocyclyl.

[816] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), R¹⁶ is hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In some embodiments, R¹⁶ is hydrogen, optionally substituted C₁-C₃ alkyl, optionally substituted C₂-C₃ alkenyl, optionally substituted C₂-C₃ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In some embodiments, R¹⁶ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl. In some embodiments, R¹⁶ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl. In some embodiments, R¹⁶ is hydrogen or C₁-C₃ alkyl.

[817] In one embodiment of the compounds of formula (I)-(IIIA), m is 1 or 2.

[818] In one embodiment of the compounds of formula (I)-(VI) and (A)-(F), t is 1 or 2. In one embodiment of the compounds of formula (I)-(VI) and (A)-(H-I), t is 1.

[819] In one embodiment of the compounds of formula (I)-(VI), (A), (B), and (C), optional substituent is selected from halogen, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -(C₁-C₃ alkyl)-(C₁-C₃ alkoxy), -(C₁-C₃ alkyl)-OH, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl). In another embodiment, the optional substituent is selected from halogen, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₁-C₃ alkoxy, -NH₂, -SCH₃, -SO₂CH₃, -NHSO₂CH₃, -CH₂NHSO₂CH₃, -SO₂NH₂, -CONH₂, or -NHCOCH₃.

[820] In one embodiment of the compounds of formula (I)-(VI), A and B are each monocyclic ring.

[821] In one embodiment of the compounds of formula (I)-(VI), B is phenyl, pyridyl, or pyrimidyl.

[822] In one embodiment of the compounds of formula (I)-(IIIA), Z and V are not both a bond.

[823] In one embodiment of the compounds of formula (I)-(VI), (A)-(C), Y and W are not both a bond.

[824] In one embodiment of the compounds of formula (I)-(VI), C is a 4- to 10-membered ring.

[825] In one embodiment of the compounds of formula (D)-(H-I), X is a bond, -CH₂-, -C(CH₃)H-, -C(CH₃)₂-, or -CH₂CH₂-. In one embodiment, X is -CH₂-, -C(CH₃)H-, or -C(CH₃)₂-. In some embodiments, X is -C(CH₃)₂-.

[826] In one embodiment of the compounds of formula (D)-(H), X is -NR⁷-. In one embodiment, X is -NH-, -N(CH₃)-, -N(CH₂CH₃)-, -N(*i*Pr)-, or -N(*t*Bu)-.

[827] In one embodiment of the compounds of formula (D)-(H-I), Y is -O-. In one embodiment of the compounds of formula (D)-(H-I), Z is -O-. In one embodiment of the compounds of formula (D)-(H-I), Y and Z are both -O-.

[828] In one embodiment of the compounds of formula (D)-(H-I), -V-L is CH₂CH₂Cl, -CH₂CH₂CH₂Cl, or -CH₃. In some embodiments, -V-L is CH₂CH₂Cl or -CH₂CH₂CH₂Cl.

[829] In one embodiment of the compounds of formula (D)-(H-I), n₁ is 0.

[830] In one embodiment of the compounds of formula (D)-(H-I), n₂ is 0, 1, or 2. In some embodiments, n₂ is 2. In some embodiments, n₂ is 2 and R² are each ortho to Z. In other embodiments, n₂ is 2 and R² are each ortho to Z, wherein R² is halogen or -CN.

[831] In one embodiment of the compound of formula (I)-(VI) and (A)-(H-I), the compound can be a stereoisomer. For example, if X is -(CR⁵R⁶)- and R⁵ and R⁶ are different, the carbon attached to R⁵ and R⁶ can be in an S configuration or an R configuration.

[832] In some embodiments of the compounds of formula (I)-(VI) and (A)-(H-I), a hydrogen atom can be replaced with a deuterium atom.

[833] In one embodiment of the compound of formula (I)-(VA), (A), (A-I), or (D)-(H-I), the compound is selected from **Table A** below, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(VA), (A), or (D)-(H-I), the compound is selected from Compounds A3, A5, A7, A13, A17, A18, A22, A23, A24, A25, A28, A30, A31, A32, A34, A35, A38, A40, A41, A42, A45, A49, A52, A53, A54, A56, A57, A58, A62, A63, A64, A65, A68, A73, A74, A75, or A76, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(VA), (A), (A-I), or (D)-(H-I), the compound is selected from Compounds A1, A2, A4, A6, A8, A9, A10, A11, A12, A14, A15, A16, A19, A20, A21, A26, A27, A29, A33, A36, A37, A39, A43, A44, A46, A47, A48, A50, A51, A55, A59, A60, A61, A66, A67, A69, A70, A71, A72, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, A88, A89, A90, A91, A92, A93, A94, A95, A96, or A97, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(VA), (A), (A-I), or (D)-(H-I), the compound is selected from Compounds A98-A186, or a pharmaceutically acceptable salt, tautomer,

stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(VA), (A), (A-I), or (D)-(H-I), the compound is selected from Compounds A187-A211, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(VA), (A), (A-I), or (D)-(H-I), the compound is selected from Compounds A1-A211, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(VA), (A), (A-I), or (D)-(H-I), the compound is selected from Compounds A212-A234, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(VA), (A), (A-I), or (D)-(H-I), the compound is selected from Compounds A1-A234, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(VA), (A), (A-I), or (D)-(H-I), the compound is selected from Compounds A1-A96, A98-A116, A118-A159, A161-A175, and A177-A234, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

[834] In one embodiment of the compound of formula (I)-(VA), (A), (A-I), or (D)-(H-I), the compound is selected from A13, A57, A74, A93, A109, A112, A122, A126, A131, A134, A136, A137, A164, A168, A169, A170, A171, A172, A184, A185, A195, and/or A204, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.








[835] Table A. Compounds








Compound ID	Structure
A1	
A2	
A3	








Compound ID	Structure
A4	
A5	
A6	
A7	
A8	
A9	
A10	
A11	

Compound ID	Structure
A12	
A13	
A14	
A15	
A16	
A17	
A18	
A19	








Compound ID	Structure
A20	
A21	
A22	
A23	
A24	
A25	
A26	








Compound ID	Structure
A27	
A28	
A29	
A30	
A31	
A32	
A33	








Compound ID	Structure
A34	
A35	
A36	
A37	
A38	
A39	
A40	








Compound ID	Structure
A41	
A42	
A43	
A44	
A45	
A46	
A47	

Compound ID	Structure
A48	
A49	
A50	
A51	
A52	
A53	
A54	

Compound ID	Structure
A55	
A56	
A57	
A58	
A59	
A60	
A61	

Compound ID	Structure
A62	
A63	
A64	
A65	
A66	
A67	
A68	








Compound ID	Structure
A69	
A70	
A71	
A72	
A73	
A74	
A75	

Compound ID	Structure
A76	
A77	
A78	
A79	
A80	
A81	
A82	

Compound ID	Structure
A83	
A84	
A85	
A86	
A87	
A88	
A89	
A90	










Compound ID	Structure
A91	
A92	
A93	
A94	
A95	
A96	
A97	
A98	









Compound ID	Structure
A99	Chemical structure A99: A biphenyl derivative. The left phenyl ring is substituted with a sulfonamide group (-SO₂NH-) at the 4-position and a methoxy group (-O-) at the 3-position. The right phenyl ring is substituted with a cyano group (-CN) at the 4-position and a chloroethoxy group (-OCH₂CH₂Cl) at the 3-position. The two rings are connected by a central carbon atom bonded to two methyl groups.
A100	Chemical structure A100: A biphenyl derivative. The left phenyl ring is substituted with a sulfonamide group (-SO₂NH-) at the 4-position and a methoxy group (-O-) at the 3-position. The right phenyl ring is substituted with a cyano group (-CN) at the 4-position and a chloroethoxy group (-OCH₂CH₂Cl) at the 3-position. The two rings are connected by a central carbon atom bonded to two methyl groups.
A101	Chemical structure A101: A biphenyl derivative. The left phenyl ring is substituted with a sulfonamide group (-SO₂NH-) at the 4-position and a methoxy group (-O-) at the 3-position. The right phenyl ring is substituted with a cyano group (-CN) at the 4-position and a chloroethoxy group (-OCH₂CH₂Cl) at the 3-position. The two rings are connected by a central carbon atom bonded to two methyl groups.
A102	Chemical structure A102: A biphenyl derivative. The left phenyl ring is substituted with a sulfonamide group (-SO₂NH-) at the 4-position and a methoxy group (-O-) at the 3-position. The right phenyl ring is substituted with a cyano group (-CN) at the 4-position and a chloroethoxy group (-OCH₂CH₂Cl) at the 3-position. The two rings are connected by a central carbon atom bonded to two methyl groups.
A103	Chemical structure A103: A biphenyl derivative. The left phenyl ring is substituted with a sulfonamide group (-SO₂NH-) at the 4-position and a methoxy group (-O-) at the 3-position. The right phenyl ring is substituted with a cyano group (-CN) at the 4-position and a chloroethoxy group (-OCH₂CH₂Cl) at the 3-position. The two rings are connected by a central carbon atom bonded to two methyl groups.
A104	Chemical structure A104: A biphenyl derivative. The left phenyl ring is substituted with a sulfonamide group (-SO₂NH-) at the 4-position and a methoxy group (-O-) at the 3-position. The right phenyl ring is substituted with a cyano group (-CN) at the 4-position and a chloroethoxy group (-OCH₂CH₂Cl) at the 3-position. The two rings are connected by a central carbon atom bonded to two methyl groups.
A105	Chemical structure A105: A biphenyl derivative. The left phenyl ring is substituted with a sulfonamide group (-SO₂NH-) at the 4-position and a methoxy group (-O-) at the 3-position. The right phenyl ring is substituted with a cyano group (-CN) at the 4-position and a chloroethoxy group (-OCH₂CH₂Cl) at the 3-position. The two rings are connected by a central carbon atom bonded to two methyl groups.

Compound ID	Structure
A106	
A107	
A108	
A109	
A110	
A111	
A112	









Compound ID	Structure
A113	
A114	
A115	
A116	
A117	
A118	
A119	
A120	









Compound ID	Structure
A121	
A122	
A123	
A124	
A125	
A126	
A127	

Compound ID	Structure
A128	
A129	
A130	
A131	
A132	
A133	
A134	
A135	
A136	









Compound ID	Structure
A137	
A138	
A139	
A140	
A141	
A142	
A143	
A144	







Compound ID	Structure
A145	
A146	
A147	
A148	
A149	
A150	
A151	
A152	

Compound ID	Structure
A153	
A154	
A155	
A156	
A157	
A158	
A159	
A160	

Compound ID	Structure
A161	
A162	
A163	
A164	
A165	
A166	
A167	
A168	









Compound ID	Structure
A169	
A170	
A171	
A172	
A173	
A174	
A175	
A176	
A177	

Compound ID	Structure
A178	
A179	
A180	
A181	
A182	
A183	
A184	
A185	










Compound ID	Structure
A186	
A187	
A188	
A189	
A190	
A191	

Compound ID	Structure
A192	
A193	
A194	
A195	
A196	
A197	
A198	
A199	
A200	

Compound ID	Structure
A201	
A202	
A203	
A204	
A205	
A206	
A207	
A208	






Compound ID	Structure
A209	
A210	
A211	
A212	
A213	
A214	
A215	
A216	







Compound ID	Structure
A217	
A218	
A219	
A220	
A221	
A222	
A223	
A224	
A225	

Compound ID	Structure
A226	
A227	
A228	
A229	
A230	
A231	
A232	
A233	
A234	

[836] In one embodiment of the compound of formula (I)-(IV), (VI), (B) or (C), the compound is selected from **Table B** below, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(IV), (VI), (B) or (C), the compound is selected from Compounds B1, B2, B3, or B6 or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(IV), (VI), (B) or (C), the compound is selected from Compounds B4, B5, B7, B8, B9, B10, or B11 or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment of the compound of formula (I)-(IV), (VI), (B) or (C), the compound is selected from Compounds B1-B11 or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

[837] **Table B.** Compounds

Compound ID	Structure
B1	
B2	
B3	
B4	
B5	

Compound ID	Structure
B6	
B7	
B8	
B9	
B10	
B11	

[838] In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound of any one of formula (I), (IA), (IB), (IC), (II), (IIA), (IIIA), (IIB), (III), (IV), (IVA), (V), (VA), (VI), (A), (A-I), (B)-(D), (E), (E-I)-(E-VII), (F), (G), (G-I), (G-II), (H), and (H-I) ("formula (I)-(VI) and (A)-(H-I)") or compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound selected from Compounds A3, A5, A7, A13, A17, A18, A22, A23, A24, A25, A28, A30, A31, A32, A34, A35, A38, A40, A41, A42, A45, A49, A52, A53, A54, A56, A57, A58, A62, A63, A64, A65, A68, A73, A74, A75, A76, B1, B2, B3, or B6 or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound selected from Compounds A1, A2, A4, A6, A8, A9, A10, A11, A12, A14, A15, A16, A19, A20, A21, A26, A27, A29, A33, A36, A37, A39, A43, A44, A46, A47, A48, A50, A51, A55, A59, A60, A61, A66, A67, A69, A70, A71, A72, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, A88, A89, A90, A91, A92, A93, A94, A95, A96, A97, B4, B5, B7, B8, B9, B10, or B11, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound selected from Compounds A98-A186, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound selected from Compounds A187-A211, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound selected from Compounds A212-A234, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound selected from Compounds A1-A96, A98-A116, A118-A159, A161-A175, and A177-A234, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides

a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound selected from Compounds A13, A57, A74, A93, A109, A112, A122, A126, A131, A134, A136, A137, A164, A168, A169, A170, A171, A172, A184, A185, A195, and/or A204, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound selected from Compounds A1-A211 or B1-B11, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound selected from Compounds A1-A234 or B1-B11, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In one embodiment, the present disclosure provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier or a pharmaceutically acceptable excipient and a compound of any one of formula (D)-(H-I), or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

Therapeutic Use

[839] The present compounds find use in any number of methods. For example, in some embodiments the compounds are useful in methods for modulating androgen receptor (AR). Accordingly, in one embodiment, the present disclosure provides the use of any one of the foregoing compounds of formula (I)-(VI) and (A)-(H-I), or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, for modulating androgen receptor (AR) activity. For example in some embodiments, modulating androgen receptor (AR) activity is in a mammalian cell. Modulating androgen receptor (AR) can be in a subject in need thereof (e.g., a mammalian subject) and for treatment of any of the described conditions or diseases.

[840] In one embodiment, the modulating AR is binding to AR. In other embodiments, the modulating AR is inhibiting AR.

[841] In one embodiment, the modulating AR is modulating AR N-terminal domain (NTD). In one embodiment, the modulating AR is binding to AR NTD. In other embodiments, the modulating AR is inhibiting AR NTD. In one embodiment, the modulating AR is modulating AR N-terminal domain (NTD). In some embodiments, modulating the AR is inhibiting transactivation of androgen receptor N-terminal domain (NTD).

[842] In other embodiments, modulating androgen receptor (AR) activity is for treatment of at least one indication selected from the group consisting of: prostate cancer, breast cancer,

ovarian cancer, bladder cancer, pancreatic cancer, hepatocellular cancer, endometrial cancer, salivary gland carcinoma, hair loss, acne, hirsutism, ovarian cysts, polycystic ovary disease, precocious puberty, spinal and bulbar muscular atrophy, age related macular degeneration, and combinations thereof. For example in some embodiments, the indication is prostate cancer. In other embodiments, the prostate cancer is primary/localized prostate cancer, locally advanced prostate cancer, recurrent prostate cancer, non-metastatic castration-resistant prostate cancer, metastatic prostate cancer, advanced prostate cancer, or metastatic castration-resistant prostate cancer (CRPC), or hormone-sensitive prostate cancer. While in other embodiments, the prostate cancer is androgen dependent prostate cancer. In other embodiments, the spinal and bulbar muscular atrophy is Kennedy's disease. In other embodiments, the prostate cancer is primary/localized prostate cancer, locally advanced prostate cancer, recurrent prostate cancer, metastatic prostate cancer, advanced prostate cancer, or metastatic castration-resistant prostate cancer (CRPC), or hormone-sensitive prostate cancer. While in other embodiments, the prostate cancer is androgen dependent prostate cancer. In other embodiments, the spinal and bulbar muscular atrophy is Kennedy's disease.

[843] In one embodiment of the present disclosure, a method of treating a condition associated with cell proliferation in a patient in need thereof is provided, comprising administering a compounds of formula (I)-(VI) and (A)-(H-I), or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, to a subject in need thereof. In one embodiment, the present invention provides a method of treating cancer or tumors. In another embodiment, the present invention provides a method of treating prostate cancer or breast cancer.

[844] In one embodiment of the present disclosure, a method of reducing, inhibiting, or ameliorating proliferation, comprising administering a therapeutically effective amount of a compound of formula (I)-(VI) and (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof is provided. In one embodiment, the reducing, inhibiting, or ameliorating in the method disclosed herein, is *in vivo*. In another embodiment, the reducing, inhibiting, or ameliorating is *in vitro*.

[845] In one embodiment, the cells in the method disclosed herein, are a cancer cells. In one embodiment, the cancer cells are a prostate cancer cells. In one embodiment, the prostate cancer cells are cells of primary/localized prostate cancer (newly diagnosed or early stage), locally advanced prostate cancer, recurrent prostate cancer (*e.g.*, prostate cancer which was not responsive to primary therapy), metastatic prostate cancer, non-metastatic castration-resistant prostate cancer, advanced prostate cancer (*e.g.*, after castration for recurrent prostate cancer), metastatic castration-resistant prostate cancer (CRPC), or hormone-sensitive prostate cancer.

In one embodiment, the prostate cancer cells are cells of primary/localized prostate cancer (newly diagnosed or early stage), locally advanced prostate cancer, recurrent prostate cancer (*e.g.*, prostate cancer which was not responsive to primary therapy), metastatic prostate cancer, advanced prostate cancer (*e.g.*, after castration for recurrent prostate cancer), metastatic castration-resistant prostate cancer (CRPC), or hormone-sensitive prostate cancer. In another embodiment, the prostate cancer cells are cells of a metastatic castration-resistant prostate cancer. In other embodiments, the prostate cancer cells are an androgen-dependent prostate cancer cells or an androgen-independent prostate cancer cells. In one embodiment, the cancer cells are breast cancer cells.

[846] In one embodiment, the condition or disease associated with cell proliferation is cancer. In one embodiment of any one of the methods disclosed herein, the cancer is selected from the group consisting of: prostate cancer, breast cancer, ovarian cancer, endometrial cancer, salivary gland carcinoma, hair loss, acne, hirsutism, ovarian cysts, polycystic ovary disease, precocious puberty, spinal and bulbar muscular atrophy, and age-related macular degeneration. In one embodiment, the condition or disease is prostate cancer. In one embodiment, prostate cancer is selected from primary/localized prostate cancer, locally advanced prostate cancer, recurrent prostate cancer, metastatic prostate cancer, non-metastatic castration-resistant prostate cancer, advanced prostate cancer, metastatic castration-resistant prostate cancer (CRPC), or hormone-sensitive prostate cancer. In one embodiment, prostate cancer is selected from primary/localized prostate cancer, locally advanced prostate cancer, recurrent prostate cancer, metastatic prostate cancer, advanced prostate cancer, metastatic castration-resistant prostate cancer (CRPC), or hormone-sensitive prostate cancer. In another embodiment, the prostate cancer is a metastatic castration-resistant prostate cancer. In some embodiments, the prostate cancer is an androgen-dependent prostate cancer cells or an androgen-independent prostate cancer. In one embodiment, the condition or disease is breast cancer.

[847] In another embodiment of the present disclosure, a method for reducing or preventing tumor growth, comprising contacting tumor cells with a therapeutically effective amount of a compound of (I)-(VI) and (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof is provided.

[848] In one embodiment, reducing or preventing tumor growth includes reduction in tumor volume. In one embodiment, reducing or preventing tumor growth includes complete elimination of tumors. In one embodiment, reducing or preventing tumor growth includes stopping or halting the existing tumor to grow. In one embodiment, reducing or preventing tumor growth includes reduction in the rate of tumor growth. In one embodiment, reducing or

preventing tumor growth includes reduction in the rate of tumor growth such that the rate of tumor growth before treating a patient with the methods disclosed herein (r_1) is faster than the rate of tumor growth after said treatment (r_2) such that $r_1 > r_2$.

[849] In one embodiment, the reducing or preventing in the method disclosed herein is *in vivo*. In another embodiment, the treating is *in vitro*.

[850] In one embodiment, the tumor cell in the method disclosed herein is selected from prostate cancer, breast cancer, ovarian cancer, endometrial cancer, or salivary gland carcinoma. In one embodiment, the tumor cells are prostate cancer tumor cells. In one embodiment, the prostate cancer tumor cells are tumor cells of primary/localized prostate cancer, locally advanced prostate cancer, recurrent prostate cancer, metastatic prostate cancer, non-metastatic castration-resistant prostate cancer, advanced prostate cancer, metastatic castration-resistant prostate cancer (CRPC), or hormone-sensitive prostate cancer. In one embodiment, the prostate cancer tumor cells are tumor cells of primary/localized prostate cancer, locally advanced prostate cancer, recurrent prostate cancer, metastatic prostate cancer, advanced prostate cancer, metastatic castration-resistant prostate cancer (CRPC), or hormone-sensitive prostate cancer. In other embodiments, the prostate cancer is a metastatic castration-resistant prostate cancer. In some embodiments, the prostate cancer is androgen-dependent prostate cancer or androgen-independent prostate cancer. In another embodiment, the tumor cells are breast cancer tumor cells.

[851] In accordance with another embodiment, there is provided a use of the compounds of formula (I)-(VI) and (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof for preparation of a medicament for modulating androgen receptor (AR). In one embodiment, the compounds of the present disclosure are compounds having the structure of formula(I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds provide metabolic stability. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in human liver microsomes (HLMs) of less than about 500 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 450 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 400 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 350 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 300 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 250 $\mu\text{L}/\text{min}/\text{mg}$

protein, less than about 225 $\mu\text{L}/\text{min}/\text{mg}$ protein, or less than about 200 $\mu\text{L}/\text{min}/\text{mg}$ protein. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HLMs of less than about 200 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 190 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 180 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 170 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 160 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 150 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 140 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 130 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 120 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 110 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 100 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 90 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 80 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 70 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 60 $\mu\text{L}/\text{min}/\text{mg}$ protein, or less than about 50 $\mu\text{L}/\text{min}/\text{mg}$ protein. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HLMs of less than about 50 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 48 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 45 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 40 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 35 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 30 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 25 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 20 $\mu\text{L}/\text{min}/\text{mg}$ protein, or less than about 15 $\mu\text{L}/\text{min}/\text{mg}$ protein. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HLMs of less than about 15 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 14 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 13 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 11 $\mu\text{L}/\text{min}/\text{mg}$ protein, or less than about 10 $\mu\text{L}/\text{min}/\text{mg}$ protein.

[852] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HLMs of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-

(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HLMs in the range of about 0.1 $\mu\text{L}/\text{min}/\text{mg}$ protein to about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HLMs of less than 48 $\mu\text{L}/\text{min}/\text{mg}$ protein. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HLMs in the range of about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein to about 48 $\mu\text{L}/\text{min}/\text{mg}$ protein. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HLMs in the range of about 0.1 $\mu\text{L}/\text{min}/\text{mg}$ protein to about 48 $\mu\text{L}/\text{min}/\text{mg}$ protein.

[853] In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM *in vitro* metabolic clearance of less than about 20 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM *in vitro* metabolic clearance of less than about 15 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are compounds having the

structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM *in vitro* metabolic clearance of less than about 11.55 $\mu\text{L}/\text{min}/\text{mg}$ protein.

[854] In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM *in vitro* metabolic clearance of less than about 20 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM *in vitro* metabolic clearance of less than about 15 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM *in vitro* metabolic clearance of less than about 11.55 $\mu\text{L}/\text{min}/\text{mg}$ protein.

[855] In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in mouse liver microsomes (MLMs) of less than about 500 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 450 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 400 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 350 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 300 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 250 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 225 $\mu\text{L}/\text{min}/\text{mg}$ protein, or less than about 200 $\mu\text{L}/\text{min}/\text{mg}$ protein. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MLMs of less than about 200 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 190 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 180 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 170 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 160 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 150 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 140 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 130 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 120 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 110 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 100 $\mu\text{L}/\text{min}/\text{mg}$

protein, less than about 90 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 80 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 70 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 60 $\mu\text{L}/\text{min}/\text{mg}$ protein, or less than about 50 $\mu\text{L}/\text{min}/\text{mg}$ protein. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MLMs of less than about 50 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 48 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 45 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 40 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 35 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 30 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 25 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 20 $\mu\text{L}/\text{min}/\text{mg}$ protein, or less than about 15 $\mu\text{L}/\text{min}/\text{mg}$ protein. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MLMs of less than about 15 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 14 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 13 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein, less than about 11 $\mu\text{L}/\text{min}/\text{mg}$ protein, or less than about 10 $\mu\text{L}/\text{min}/\text{mg}$ protein.

[856] In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MLMs of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MLMs in the range of about 0.1 $\mu\text{L}/\text{min}/\text{mg}$ protein to about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MLMs of less than 48 $\mu\text{L}/\text{min}/\text{mg}$ protein. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MLMs in the range of about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein to about 48 $\mu\text{L}/\text{min}/\text{mg}$ protein. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro*

metabolic clearance in MLMs in the range of about 0.1 $\mu\text{L}/\text{min}/\text{mg}$ protein to about 48 $\mu\text{L}/\text{min}/\text{mg}$ protein.

[857] In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an MLM *in vitro* metabolic clearance of less than about 20 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an MLM *in vitro* metabolic clearance of less than about 15 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an MLM *in vitro* metabolic clearance of less than about 11.55 $\mu\text{L}/\text{min}/\text{mg}$ protein.

[858] In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an MLM *in vitro* metabolic clearance of less than about 20 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an MLM *in vitro* metabolic clearance of less than about 15 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an MLM *in vitro* metabolic clearance of less than about 11.55 $\mu\text{L}/\text{min}/\text{mg}$ protein.

[859] In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in human hepatocytes (HHs) of less than about 500 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 450 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 400 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 350 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 300 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 250 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 225 $\mu\text{L}/\text{min}/\text{million cells}$, or less than about 200 $\mu\text{L}/\text{min}/\text{million cells}$. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HHs of less than about 200 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 190 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 180 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 170 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 160 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 150 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 140 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 130 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 120 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 110 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 100 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 90 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 80 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 70 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 60 $\mu\text{L}/\text{min}/\text{million cells}$, or less than about 50 $\mu\text{L}/\text{min}/\text{million cells}$. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HHs of less than about 50 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 48 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 45 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 40 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 35 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 30 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 25 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 20 $\mu\text{L}/\text{min}/\text{million cells}$, or less than about 15 $\mu\text{L}/\text{min}/\text{million cells}$. In some embodiments, the compounds of the present disclosure have an *in vitro* metabolic clearance in HHs of less than about 15 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 14 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 13 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 12 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 11 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 10 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 9 $\mu\text{L}/\text{min}/\text{million cells}$.

cells, less than about 8 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 8 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 6 $\mu\text{L}/\text{min}/\text{million cells}$, or less than about 5 $\mu\text{L}/\text{min}/\text{million cells}$.

[860] In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in HHs of less than 4 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in HHs in the range of about 0.1 $\mu\text{L}/\text{min}/\text{million cells}$ to about 4 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in HHs of less than 18 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in HHs in the range of about 4 $\mu\text{L}/\text{min}/\text{million cells}$ to about 18 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in HHs in the range of about 0.1 $\mu\text{L}/\text{min}/\text{million cells}$ to about 18 $\mu\text{L}/\text{min}/\text{million cells}$.

[861] In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HHs of less than about 20 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HHs of less than about 10 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HHs of less than about 5 $\mu\text{L}/\text{min}/\text{million}$

cells. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HHs of less than about 3.85 $\mu\text{L}/\text{min}/\text{million cells}$.

[862] In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HHs of less than about 20 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HHs of less than about 10 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in HHs of less than about 5 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in human hepatocytes of less than about 3.85 $\mu\text{L}/\text{min}/\text{million cells}$.

[863] In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in mouse hepatocytes (MHs) of less than about 500 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 450 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 400 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 350 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 300 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 250 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 225 $\mu\text{L}/\text{min}/\text{million cells}$, or less than about 200 $\mu\text{L}/\text{min}/\text{million cells}$. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 200 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 190 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 180 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 170 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 160 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 150 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 140 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 130

$\mu\text{L}/\text{min}/\text{million cells}$, less than about 120 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 110 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 100 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 90 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 80 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 70 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 60 $\mu\text{L}/\text{min}/\text{million cells}$, or less than about 50 $\mu\text{L}/\text{min}/\text{million cells}$. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 50 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 48 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 45 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 40 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 35 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 30 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 25 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 20 $\mu\text{L}/\text{min}/\text{million cells}$, or less than about 15 $\mu\text{L}/\text{min}/\text{million cells}$. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 15 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 14 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 13 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 12 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 11 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 10 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 9 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 8 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 8 $\mu\text{L}/\text{min}/\text{million cells}$, less than about 6 $\mu\text{L}/\text{min}/\text{million cells}$, or less than about 5 $\mu\text{L}/\text{min}/\text{million cells}$.

[864] In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MHs of less than 4 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MHs in the range of about 0.1 $\mu\text{L}/\text{min}/\text{million cells}$ to about 4 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MHs of less than 18 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein

the compounds have an *in vitro* metabolic clearance in MHs in the range of about 4 $\mu\text{L}/\text{min}/\text{million cells}$ to about 18 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MHs in the range of about 0.1 $\mu\text{L}/\text{min}/\text{million cells}$ to about 18 $\mu\text{L}/\text{min}/\text{million cells}$.

[865] In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 20 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 17 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 7 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 3.85 $\mu\text{L}/\text{min}/\text{million cells}$.

[866] In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 20 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 17 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of

less than about 7 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A or B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in MHs of less than about 3.85 $\mu\text{L}/\text{min}/\text{million cells}$.

[867] In another embodiment, the compounds of the present disclosure are any compounds with a combination of *in vitro* metabolic clearance in HLM, MLM, HHs, and MHs as described herein.

[868] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 20 $\mu\text{L}/\text{min}/\text{mg protein}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 15 $\mu\text{L}/\text{min}/\text{mg protein}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg protein}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 11.55 $\mu\text{L}/\text{min}/\text{mg protein}$.

[869] In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 20 $\mu\text{L}/\text{min}/\text{mg protein}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 15 $\mu\text{L}/\text{min}/\text{mg protein}$. In another embodiment, the compounds of

the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein. In another embodiment, the compounds of the present disclosure are any compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 11.55 $\mu\text{L}/\text{min}/\text{mg}$ protein.

[870] In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 20 $\mu\text{L}/\text{min}/\text{million}$ cells. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in mouse hepatocytes of less than about 17 $\mu\text{L}/\text{min}/\text{million}$ cells and an *in vitro* metabolic clearance in human hepatocytes of less than about 10 $\mu\text{L}/\text{min}/\text{million}$ cells. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in mouse hepatocytes of less than about 7 $\mu\text{L}/\text{min}/\text{million}$ cells, and an *in vitro* metabolic clearance in human hepatocytes of less than about 5 $\mu\text{L}/\text{min}/\text{million}$ cells. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 3.85 $\mu\text{L}/\text{min}/\text{million}$ cells.

[871] In another embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 20 $\mu\text{L}/\text{min}/\text{million}$ cells. In another embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt,

tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in mouse hepatocytes of less than about 17 $\mu\text{L}/\text{min}/\text{million cells}$ and an *in vitro* metabolic clearance in human hepatocytes of less than about 10 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in mouse hepatocytes of less than about 7 $\mu\text{L}/\text{min}/\text{million cells}$, and an *in vitro* metabolic clearance in human hepatocytes of less than about 5 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 3.85 $\mu\text{L}/\text{min}/\text{million cells}$.

[872] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 20 $\mu\text{L}/\text{min}/\text{mg protein}$ and an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 20 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 15 $\mu\text{L}/\text{min}/\text{mg protein}$, an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 17 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg protein}$, an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 7 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds s have an HLM and MLM in

vitro metabolic clearance of less than about 11.55 $\mu\text{L}/\text{min}/\text{mg}$ protein, an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 3.85 $\mu\text{L}/\text{min}/\text{million cells}$.

[873] In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an a) HLM or MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein and b) an *in vitro* metabolic clearance in human or mouse hepatocytes of less than about 4 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an a) HLM or MLM *in vitro* metabolic clearance of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein and b) an *in vitro* metabolic clearance in human or mouse hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an a) HLM and MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein and b) an *in vitro* metabolic clearance in human or mouse hepatocytes of less than about 4 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an a) HLM or MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein and b) an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 4 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an a) HLM and MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein and b) an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 4 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or

stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an a) HLM and MLM *in vitro* metabolic clearance of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein and b) an *in vitro* metabolic clearance in human and mouse hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an a) HLM and MLM *in vitro* metabolic clearance of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein and b) an *in vitro* metabolic clearance in human and mouse hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million cells}$.

[874] In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 20 $\mu\text{L}/\text{min}/\text{mg}$ protein and an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 20 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 15 $\mu\text{L}/\text{min}/\text{mg}$ protein, an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 17 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present are any compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 12 $\mu\text{L}/\text{min}/\text{mg}$ protein, an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 7 $\mu\text{L}/\text{min}/\text{million cells}$. In another embodiment, the compounds of the present disclosure are any compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an HLM and MLM *in vitro* metabolic clearance of less than about 11.55 $\mu\text{L}/\text{min}/\text{mg}$ protein, an *in vitro* metabolic clearance in human and mouse hepatocytes of less than about 3.85 $\mu\text{L}/\text{min}/\text{million cells}$.

[875] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human, mouse, rat, monkey or dog microsome greater than about 120 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or

stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human, mouse, rat, monkey or dog microsome greater than about 110 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human, mouse, rat, monkey or dog microsome greater than about 100 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human, mouse, rat, monkey or dog microsome greater than about 90 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human, mouse, rat, monkey or dog microsome greater than about 80 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human, mouse, rat, monkey or dog microsome greater than about 70 minutes.

[876] In another embodiment, the compounds of the present disclosure are any compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human microsomes greater than about 120 min.

[877] In another embodiment, the compounds of the present disclosure are any compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse microsomes greater than about 120 min.

[878] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically

acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 120 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 110 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 100 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 90 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 80 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 70 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 60 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 50 minutes. In some embodiments, the compounds of the present disclosure are compounds

having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 40 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 30 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 20 minutes. In some embodiments, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat microsome greater than about 10 minutes. In one embodiment, the compound is selected from any of the compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof.

[879] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 360 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 350 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 325 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of

formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 310 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 300 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 290 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 250 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 200 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 150 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 100 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically

acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in human hepatocytes greater than about 80 minutes. In one embodiment, the compound is selected from any of the compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof.

[880] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 360 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 350 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 325 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 300 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 250 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 225 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the

compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 200 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 150 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 100 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 50 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mouse hepatocytes greater than about 30 minutes. In one embodiment, the compound is selected from any of the compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof.

[881] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 360 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 350 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt,

tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 325 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 300 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 250 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 225 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 200 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 150 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 100 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in rat hepatocytes greater than about 50

minutes. In one embodiment, the compounds have an *in vitro* half-life in rat hepatocytes greater than about 30 minutes. In one embodiment, the compound is selected from any of the compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof.

[882] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome of greater than 80 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome of greater than 90 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome of greater than 100 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome of greater than 110 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome of greater than 115 minutes. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome of greater than 120 minutes. In one embodiment, the compound is selected from any of the compound of Tables A

and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof. In one embodiment, the mammal microsome is human or mouse microsome.

[883] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome in the range of about 50 min to about 200 min. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome in the range of about 80 min to about 180 min. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome in the range of about 100 min to about 160 min. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome in the range of about 110 min to about 150 min. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome in the range of about 110 min to about 140 min. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life stability in mammal microsome in the range of about 110 min to about 130 min. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or

[884] In one embodiment, the present disclosure provides compounds which demonstrate blocking androgen-induced PSA-luciferase activity (PSA-luciferase assay). In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 4500 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 4000 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 3500 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 3000 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 2500 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables

A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 2000 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 1500 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 1000 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 950 nm. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 900 nM.

[885] In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 890 nM, less than about 880 nM, less than about 870 nM, less than about 860 nM, less than about 850 nM, less than about 840 nM, less than about 830 nM, less than about 820 nM, less than about 810 nM, or less than about 800 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 790 nM, less than about 780 nM, less than about 770 nM, less than about 760 nM, less than about 750 nM, less than about 740 nM, less than about 730 nM, less than about 720 nM, less than about 710 nM, or less than about 700 nM. In one

embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 690 nM, less than about 680 nM, less than about 670 nM, less than about 660 nM, less than about 650 nM, less than about 640 nM, less than about 630 nM, less than about 620 nM, less than about 610 nM, or less than about 600 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 590 nM, less than about 580 nM, less than about 570 nM, less than about 560 nM, less than about 550 nM, less than about 540 nM, less than about 530 nM, less than about 520 nM, less than about 510 nM, or less than about 500 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 500 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds having *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 490 nM, less than about 480 nM, less than about 470 nM, less than about 460 nM, less than about 450 nM, less than about 440 nM, less than about 430 nM, less than about 420 nM, less than about 410 nM, or less than about 400 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 390 nM, less than about 380 nM, less than about 370 nM, less than about 360 nM, less than about 350 nM, less than about 340 nM, less than about 330 nM, less than about 320 nM, less than about 310 nM, or less than about 300 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt,

tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 290 nM, less than about 280 nM, less than about 270 nM, less than about 260 nM, less than about 250 nM, less than about 240 nM, less than about 230 nM, less than about 220 nM, less than about 210 nM, or less than about 200 nM. In one embodiment, the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 200 nM. In one embodiment, the compound is selected from any of the compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof.

[886] In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 80 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 2000 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 90 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 1500 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 100 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 1000 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 105 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 900 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 110 minutes

and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 850 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 115 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 800 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 115 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 750 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 120 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 700 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 120 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 650 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 120 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 600 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 120 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 550 nM. In one embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI)

and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* half-life in mammal microsome of greater than about 120 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 500 nM. In one embodiment, the compound is selected from any of the compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof. In one embodiment, the mammal microsome is human or mouse microsome.

[887] In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 1000 nM and a HLM or MLM *in vitro* metabolic clearance of less than 12 μ L/min/mg protein. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 1000 nM and a HLM and MLM *in vitro* metabolic clearance of less than 12 μ L/min/mg protein. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 1000 nM and an *in vitro* metabolic clearance in human or mouse hepatocytes of less than 4 μ L/min/million cells. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 1000 nM and an *in vitro* metabolic clearance in human and mouse hepatocytes of less than 4 μ L/min/million cells.

[888] In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC₅₀

in a PSA-luciferase assay of less than about 1000 nM; a HLM or MLM *in vitro* metabolic clearance of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein; and an *in vitro* metabolic clearance in human or mouse hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million}$ cells. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay of less than about 800 nM; a HLM or MLM *in vitro* metabolic clearance of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein; and an *in vitro* metabolic clearance in human or mouse hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million}$ cells. In another embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, or the compounds of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay of less than about 700 nM; a HLM or MLM *in vitro* metabolic clearance of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein; and an *in vitro* metabolic clearance in human or mouse hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million}$ cells. In one embodiment, the compound is selected from any of the compound of Tables A and B or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof.

[889] In a specific embodiment, the compounds of the present disclosure are any compound of Tables A and B, or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay of less than about 1000 nM; a HLM or MLM *in vitro* metabolic clearance of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein; and an *in vitro* metabolic clearance in human or mouse hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million}$ cells.

[890] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in HHs of less than 4 $\mu\text{L}/\text{min}/\text{million}$ cells. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in HHs of less than 4 $\mu\text{L}/\text{min}/\text{million}$ cells and *in vitro* IC_{50} in a PSA-luciferase assay of less than about 700 nM. In one embodiment, the compounds have an *in vitro* metabolic clearance in HHs of less than 4 $\mu\text{L}/\text{min}/\text{million}$ cells and *in vitro* IC_{50} in a PSA-luciferase assay of less than about 675 nM. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-

(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* half-life in HHs of less than about 350 minutes. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* half-life in HHs of greater than about 320 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 700 nM. In one embodiment, the compounds have an *in vitro* half-life in HHs of greater than about 350 minutes and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 700 nM.

[891] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MLMs of less than 15 $\mu\text{L}/\text{min}/\text{mg}$ protein. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MLMs of less than 15 $\mu\text{L}/\text{min}/\text{mg}$ protein and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 700 nM. In one embodiment, the compounds have an *in vitro* metabolic clearance in MLMs of less than 15 $\mu\text{L}/\text{min}/\text{mg}$ protein and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 675 nM. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MHs of less than 10 $\mu\text{L}/\text{min}/\text{million cells}$. In one embodiment, the compounds have an *in vitro* metabolic clearance in MHs of less than 8 $\mu\text{L}/\text{min}/\text{million cells}$. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MHs of less than 10 $\mu\text{L}/\text{min}/\text{million cells}$ and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 700 nM. In one embodiment, the compounds have an *in vitro* metabolic clearance in MHs of less than 8 $\mu\text{L}/\text{min}/\text{million cells}$ and *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 675 nM. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* half-life in MLMs of greater than about 100 minutes. In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a

pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* half-life in MHs of greater than about 100 minutes. In one embodiment, the compounds have an *in vitro* half-life in MHs of greater than about 180 minutes.

[892] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* metabolic clearance in MHs in the range of 3 $\mu\text{L}/\text{min}/\text{million cells}$ to about 20 $\mu\text{L}/\text{min}/\text{million cells}$. In one embodiment, the compounds have an *in vitro* metabolic clearance in MHs in the range of 3.2 $\mu\text{L}/\text{min}/\text{million cells}$ to about 18 $\mu\text{L}/\text{min}/\text{million cells}$. In one embodiment, the compounds have an *in vitro* metabolic clearance in MHs in the range of 3.4 $\mu\text{L}/\text{min}/\text{million cells}$ to about 17 $\mu\text{L}/\text{min}/\text{million cells}$.

[893] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay of less than 650 nM. In one embodiment, the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay of less than about 640 nM. In one embodiment, the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay of less than about 635 nM.

[894] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay in the range of 200 nM to about 700 nM. In one embodiment, the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay in the range of about 250 nM to about 700 nM. In one embodiment, the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay in the range of about 300 nM to about 700 nM. In one embodiment, the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay in the range of about 350 nM to about 675 nM.

[895] In a specific embodiment, the compounds of the present disclosure are compounds having the structure of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, and wherein the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay of less than 525 nM. In one embodiment, the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay of less than about 500 nM. In one embodiment, the compounds have an *in vitro* IC_{50} in a PSA-luciferase assay of less than about 450 nM.

Pharmaceutical Compositions and Formulations

[896] The present disclosure also includes pharmaceutical compositions for modulating androgen receptor (AR) in a subject. In one embodiment, a pharmaceutical composition comprises one or more compounds of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt or solvate thereof.

[897] In one embodiment of the present disclosure, a pharmaceutical composition comprises a therapeutically effective amounts of one or more compounds of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt or solvate thereof.

[898] In a specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Table A, or a pharmaceutically acceptable salt or solvate thereof. In one embodiment, a pharmaceutical composition as described herein comprise one or more compounds selected from Table B, or a pharmaceutically acceptable salt or solvate thereof.

[899] In a specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Compounds A3, A5, A7, A13, A17, A18, A22, A23, A24, A25, A28, A30, A31, A32, A34, A35, A38, A40, A41, A42, A45, A49, A52, A53, A54, A56, A57, A58, A62, A63, A64, A65, A68, A73, A74, A75, A76, B1, B2, B3, or B6, or a pharmaceutically acceptable salt or solvate thereof. In a specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Compounds A1, A2, A4, A6, A8, A9, A10, A11, A12, A14, A15, A16, A19, A20, A21, A26, A27, A29, A33, A36, A37, A39, A43, A44, A46, A47, A48, A50, A51, A55, A59, A60, A61, A66, A67, A69, A70, A71, A72, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, A88, A89, A90, A91, A92, A93, A94, A95, A96, A97, B4, B5, B7, B8, B9, B10, or B11, or a pharmaceutically acceptable salt or solvate thereof. In a specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Compounds A98-A186. In a specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Compounds A187-A211. In a specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Compounds A212-A234. In a specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Compounds A1-A96, A98-A116, A118-A159, A161-A175, and A177-A234. In a specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Compounds A13, A57, A74, A93, A109, A112, A122, A126, A131, A134, A136, A137, A164, A168, A169, A170, A171, A172, A184, A185, A195, and/or A204, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof. In a

specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Compounds A1-A211 or B1-B11. In a specific embodiment, a pharmaceutical composition, as described herein, comprises one or more compounds selected from Compounds A1-A234 or B1-B11.

[900] In one embodiment, a pharmaceutical composition, as described herein, comprising one or more compounds of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt or solvate thereof, further comprises one or more additional therapeutically active agents. In one embodiment, one or more additional therapeutically active agents are selected from therapeutics useful for treating cancer, neurological disease, a disorder characterized by abnormal accumulation of α -synuclein, a disorder of an aging process, cardiovascular disease, bacterial infection, viral infection, mitochondrial related disease, mental retardation, deafness, blindness, diabetes, obesity, autoimmune disease, glaucoma, Leber's Hereditary Optic Neuropathy, and rheumatoid arthritis.

[901] In some embodiments, the one or more additional therapeutic agents is a poly (ADP-ribose) polymerase (PARP) inhibitor including but not limited to olaparib, niraparib, rucaparib, talazoparib; an androgen receptor ligand binding domain inhibitor including but not limited to enzalutamide, apalutamide, darolutamide, bicalutamide, nilutamide, flutamide, ODM-204, TAS3681; an inhibitor of CYP17 including but not limited to galeterone, abiraterone, abiraterone acetate; a microtubule inhibitor including but not limited to docetaxel, paclitaxel, cabazitaxel (XRP-6258); a modulator of PD-1 or PD-L1 including but not limited to pembrolizumab, durvalumab, nivolumab, atezolizumab; a gonadotropin releasing hormone agonist including but not limited to cyproterone acetate, leuprolide; a 5- α reductase inhibitor including but not limited to finasteride, dutasteride, turosteride, bexlosteride, izonsteride, FCE 28260, SKF105,111; a vascular endothelial growth factor inhibitor including but not limited to bevacizumab (Avastin); a histone deacetylase inhibitor including but not limited to OSU-HDAC42; an integrin α -v- β -3 inhibitor including but not limited to VITAXIN; a receptor tyrosine kinase inhibitor including but not limited to sunitumib; a phosphoinositide 3-kinase inhibitor including but not limited to alpelisib, buparlisib, idealisib; an anaplastic lymphoma kinase (ALK) inhibitor including but not limited to crizotinib, alectinib; an endothelin receptor A antagonist including but not limited to ZD-4054; an anti-CTLA4 inhibitor including but not limited to MDX-010 (ipilimumab); an heat shock protein 27 (HSP27) inhibitor including but not limited to OGX 427; an androgen receptor degrader including but not limited to ARV-330, ARV-110; a androgen receptor DNA-binding domain inhibitor including but not limited to VPC-14449; a bromodomain and extra-terminal motif

(BET) inhibitor including but not limited to BI-894999, GSK25762, GS-5829; an androgen receptor N-terminal domain inhibitor including but not limited to a sintokamide; an alpha-particle emitting radioactive therapeutic agent including but not limited to radium 233 or a salt thereof; niclosamide; or related compounds thereof; a selective estrogen receptor modulator (SERM) including but not limited to tamoxifen, raloxifene, toremifene, arzoxifene, bazedoxifene, pipindoxifene, lasofoxifene, enclomiphene; a selective estrogen receptor degrader (SERD) including but not limited to fulvestrant, ZB716, OP-1074, elacestrant, AZD9496, GDC0810, GDC0927, GW5638, GW7604; an aromatase inhibitor including but not limited to anastrozole, exemestane, letrozole; selective progesterone receptor modulators (SPRM) including but not limited to mifepristone, lonaprisone, onapristone, asoprisnil, lonaprisnil, ulipristal, telapristone; a glucocorticoid receptor inhibitor including but not limited to mifepristone, COR108297, COR125281, ORIC-101, PT150; CDK4/6 inhibitors including palbociclib, abemaciclib, ribociclib; HER2 receptor antagonist including but not limited to trastuzumab, neratinib; a mammalian target of rapamycin (mTOR) inhibitor including but not limited to everolimus, temsirolimus.

[902] In a further embodiment of the present disclosure, a pharmaceutical composition comprising one or more compounds of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable excipient or adjuvant is provided. The pharmaceutically acceptable excipients and adjuvants are added to the composition or formulation for a variety of purposes. In another embodiment, a pharmaceutical composition comprising one or more compounds of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt or solvate thereof, further comprises a pharmaceutically acceptable carrier. In one embodiment, a pharmaceutically acceptable carrier includes a pharmaceutically acceptable excipient, binder, and/or diluent. In one embodiment, suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylase, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose and polyvinylpyrrolidone.

[903] In certain embodiments, the pharmaceutical compositions of the present disclosure may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the pharmaceutical compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives,

antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the oligonucleotide(s) of the formulation.

[904] For the purposes of this disclosure, the compounds of the present disclosure can be formulated for administration by a variety of means including orally, parenterally, by inhalation spray, topically, or rectally in formulations containing pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used here includes subcutaneous, intravenous, intramuscular, and intraarterial injections with a variety of infusion techniques. Intraarterial and intravenous injection as used herein includes administration through catheters.

[905] The compounds disclosed herein can be formulated in accordance with the routine procedures adapted for desired administration route. Accordingly, the compounds disclosed herein can take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. The compounds disclosed herein can also be formulated as a preparation for implantation or injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (e.g., as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives (e.g., as a sparingly soluble salt). Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use. Suitable formulations for each of these methods of administration can be found, for example, in Remington: The Science and Practice of Pharmacy, A. Gennaro, ed., 20th edition, Lippincott, Williams & Wilkins, Philadelphia, PA.

[906] In certain embodiments, a pharmaceutical composition of the present disclosure is prepared using known techniques, including, but not limited to mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or tableting processes.

[907] In one embodiment, the present disclosure provides a pharmaceutical composition comprising a compound of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt or solvate thereof, as disclosed herein, combined with a pharmaceutically acceptable carrier. In one embodiment, suitable pharmaceutically acceptable carriers include, but are not limited to, inert solid fillers or diluents and sterile aqueous or organic solutions. Pharmaceutically acceptable carriers are well known to those skilled in the art and include, but are not limited to, from about 0.01 to about 0.1 M and preferably 0.05M phosphate buffer or

0.8% saline. Such pharmaceutically acceptable carriers can be aqueous or non-aqueous solutions, suspensions and emulsions. Examples of non-aqueous solvents suitable for use in the present application include, but are not limited to, propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate.

[908] Aqueous carriers suitable for use in the present application include, but are not limited to, water, ethanol, alcoholic/aqueous solutions, glycerol, emulsions or suspensions, including saline and buffered media. Oral carriers can be elixirs, syrups, capsules, tablets and the like.

[909] Liquid carriers suitable for use in the present application can be used in preparing solutions, suspensions, emulsions, syrups, elixirs and pressurized compounds. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fats. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators.

[910] Liquid carriers suitable for use in the present application include, but are not limited to, water (partially containing additives as above, e.g. cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g. glycols) and their derivatives, and oils (e.g. fractionated coconut oil and arachis oil). For parenteral administration, the carrier can also include an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are useful in sterile liquid form comprising compounds for parenteral administration. The liquid carrier for pressurized compounds disclosed herein can be halogenated hydrocarbon or other pharmaceutically acceptable propellant.

[911] Solid carriers suitable for use in the present application include, but are not limited to, inert substances such as lactose, starch, glucose, methyl-cellulose, magnesium stearate, dicalcium phosphate, mannitol and the like. A solid carrier can further include one or more substances acting as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents; it can also be an encapsulating material. In powders, the carrier can be a finely divided solid which is in admixture with the finely divided active compound. In tablets, the active compound is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active compound. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose,

polyvinylpyrrolidone, low melting waxes and ion exchange resins. A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free flowing form such as a powder or granules, optionally mixed with a binder (e.g., povidone, gelatin, hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (e.g., sodium starch glycolate, cross-linked povidone, cross-linked sodium carboxymethyl cellulose) surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropyl methylcellulose in varying proportions to provide the desired release profile. Tablets may optionally be provided with an enteric coating, to provide release in parts of the gut other than the stomach.

[912] Parenteral carriers suitable for use in the present application include, but are not limited to, sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's and fixed oils. Intravenous carriers include fluid and nutrient replenishers, electrolyte replenishers such as those based on Ringer's dextrose and the like. Preservatives and other additives can also be present, such as, for example, antimicrobials, antioxidants, chelating agents, inert gases and the like.

[913] Carriers suitable for use in the present application can be mixed as needed with disintegrants, diluents, granulating agents, lubricants, binders and the like using conventional techniques known in the art. The carriers can also be sterilized using methods that do not deleteriously react with the compounds, as is generally known in the art.

[914] Diluents may be added to the formulations of the present invention. Diluents increase the bulk of a solid pharmaceutical composition and/or combination, and may make a pharmaceutical dosage form containing the composition and/or combination easier for the patient and care giver to handle. Diluents for solid compositions and/or combinations include, for example, microcrystalline cellulose (e.g., AVICEL), microfine cellulose, lactose, starch, pregelatinized starch, calcium carbonate, calcium sulfate, sugar, dextrans, dextrin, dextrose, dibasic calcium phosphate dihydrate, tribasic calcium phosphate, kaolin, magnesium carbonate, magnesium oxide, maltodextrin, mannitol, polymethacrylates (e.g., EUDRAGIT(r)), potassium chloride, powdered cellulose, sodium chloride, sorbitol, and talc.

[915] Additional embodiments relate to the pharmaceutical formulations wherein the formulation is selected from the group consisting of a solid, powder, liquid and a gel. In certain

embodiments, a pharmaceutical composition of the present invention is a solid (e.g., a powder, tablet, a capsule, granulates, and/or aggregates). In certain of such embodiments, a solid pharmaceutical composition comprising one or more ingredients known in the art, including, but not limited to, starches, sugars, diluents, granulating agents, lubricants, binders, and disintegrating agents.

[916] Solid pharmaceutical compositions that are compacted into a dosage form, such as a tablet, may include excipients whose functions include helping to bind the active ingredient and other excipients together after compression. Binders for solid pharmaceutical compositions and/or combinations include acacia, alginic acid, carbomer (e.g., carbopol), carboxymethylcellulose sodium, dextrin, ethyl cellulose, gelatin, guar gum, gum tragacanth, hydrogenated vegetable oil, hydroxyethyl cellulose, hydroxypropyl cellulose (e.g., KLUCEL), hydroxypropyl methyl cellulose (e.g., METHOCEL), liquid glucose, magnesium aluminum silicate, maltodextrin, methylcellulose, polymethacrylates, povidone (e.g., KOLLIDON, PLASDONE), pregelatinized starch, sodium alginate, and starch.

[917] The dissolution rate of a compacted solid pharmaceutical composition in the patient's stomach may be increased by the addition of a disintegrant to the composition and/or combination. Disintegrants include alginic acid, carboxymethylcellulose calcium, carboxymethylcellulose sodium (e.g., AC-DI-SOL and PRIMELLOSE), colloidal silicon dioxide, croscarmellose sodium, crospovidone (e.g., KOLLIDON and POLYPLASDONE), guar gum, magnesium aluminum silicate, methyl cellulose, microcrystalline cellulose, polacrilin potassium, powdered cellulose, pregelatinized starch, sodium alginate, sodium starch glycolate (e.g., EXPLOTAB), potato starch, and starch.

[918] Glidants can be added to improve the flowability of a non-compacted solid composition and/or combination and to improve the accuracy of dosing. Excipients that may function as glidants include colloidal silicon dioxide, magnesium trisilicate, powdered cellulose, starch, talc, and tribasic calcium phosphate.

[919] When a dosage form such as a tablet is made by the compaction of a powdered composition, the composition is subjected to pressure from a punch and dye. Some excipients and active ingredients have a tendency to adhere to the surfaces of the punch and dye, which can cause the product to have pitting and other surface irregularities. A lubricant can be added to the composition and/or combination to reduce adhesion and ease the release of the product from the dye. Lubricants include magnesium stearate, calcium stearate, glyceryl monostearate, glyceryl palmitostearate, hydrogenated castor oil, hydrogenated vegetable oil, mineral oil,

polyethylene glycol, sodium benzoate, sodium lauryl sulfate, sodium stearyl fumarate, stearic acid, talc, and zinc stearate.

[920] Flavoring agents and flavor enhancers make the dosage form more palatable to the patient. Common flavoring agents and flavor enhancers for pharmaceutical products that may be included in the composition and/or combination of the present invention include maltol, vanillin, ethyl vanillin, menthol, citric acid, fumaric acid, ethyl maltol, and tartaric acid.

[921] Solid and liquid compositions may also be dyed using any pharmaceutically acceptable colorant to improve their appearance and/or facilitate patient identification of the product and unit dosage level.

[922] In certain embodiments, a pharmaceutical composition of the present invention is a liquid (e.g., a suspension, elixir and/or solution). In certain of such embodiments, a liquid pharmaceutical composition is prepared using ingredients known in the art, including, but not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents.

[923] Liquid pharmaceutical compositions can be prepared using compounds of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt or solvate thereof, and any other solid excipients where the components are dissolved or suspended in a liquid carrier such as water, vegetable oil, alcohol, polyethylene glycol, propylene glycol, or glycerin.

[924] For example, formulations for parenteral administration can contain as common excipients sterile water or saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes and the like. In particular, biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxyethylene-polyoxypropylene copolymers can be useful excipients to control the release of active compounds. Other potentially useful parenteral delivery systems include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation administration contain as excipients, for example, lactose, or can be aqueous solutions containing, for example, polyoxyethylene-9-arylether, glycocholate and deoxycholate, or oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally. Formulations for parenteral administration can also include glycocholate for buccal administration, methoxysalicylate for rectal administration, or citric acid for vaginal administration.

[925] Liquid pharmaceutical compositions can contain emulsifying agents to disperse uniformly throughout the composition and/or combination an active ingredient or other excipient that is not soluble in the liquid carrier. Emulsifying agents that may be useful in liquid compositions and/or combinations of the present invention include, for example, gelatin, egg

yolk, casein, cholesterol, acacia, tragacanth, chondrus, pectin, methyl cellulose, carbomer, cetostearyl alcohol, and cetyl alcohol.

[926] Liquid pharmaceutical compositions can also contain a viscosity enhancing agent to improve the mouth-feel of the product and/or coat the lining of the gastrointestinal tract. Such agents include acacia, alginic acid bentonite, carbomer, carboxymethylcellulose calcium or sodium, cetostearyl alcohol, methyl cellulose, ethylcellulose, gelatin guar gum, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, maltodextrin, polyvinyl alcohol, povidone, propylene carbonate, propylene glycol alginate, sodium alginate, sodium starch glycolate, starch tragacanth, and xanthan gum.

[927] Sweetening agents such as aspartame, lactose, sorbitol, saccharin, sodium saccharin, sucrose, aspartame, fructose, mannitol, and invert sugar may be added to improve the taste.

[928] Preservatives and chelating agents such as alcohol, sodium benzoate, butylated hydroxyl toluene, butylated hydroxyanisole, and ethylenediamine tetraacetic acid may be added at levels safe for ingestion to improve storage stability.

[929] A liquid composition can also contain a buffer such as guconic acid, lactic acid, citric acid or acetic acid, sodium guconate, sodium lactate, sodium citrate, or sodium acetate. Selection of excipients and the amounts used may be readily determined by the formulation scientist based upon experience and consideration of standard procedures and reference works in the field.

[930] In one embodiment, a pharmaceutical composition is prepared for administration by injection (e.g., intravenous, subcutaneous, intramuscular, etc.). In certain of such embodiments, a pharmaceutical composition comprises a carrier and is formulated in aqueous solution, such as water or physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. In certain embodiments, other ingredients are included (e.g., ingredients that aid in solubility or serve as preservatives). In certain embodiments, injectable suspensions are prepared using appropriate liquid carriers, suspending agents and the like. Certain pharmaceutical compositions for injection are presented in unit dosage form, e.g., in ampoules or in multi-dose containers. Certain pharmaceutical compositions for injection are suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Certain solvents suitable for use in pharmaceutical compositions for injection include, but are not limited to, lipophilic solvents and fatty oils, such as sesame oil, synthetic fatty acid esters, such as ethyl oleate or triglycerides, and liposomes. Aqueous injection suspensions may contain substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or

dextran. Optionally, such suspensions may also contain suitable stabilizers or agents that increase the solubility of the pharmaceutical agents to allow for the preparation of highly concentrated solutions.

[931] The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as a solution in 1,3-butane-diol or prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile fixed oils may conventionally be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables. Formulations for intravenous administration can comprise solutions in sterile isotonic aqueous buffer. Where necessary, the formulations can also include a solubilizing agent and a local anesthetic to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampule or sachet indicating the quantity of active agent. Where the compound is to be administered by infusion, it can be dispensed in a formulation with an infusion bottle containing sterile pharmaceutical grade water, saline or dextrose/water. Where the compound is administered by injection, an ampule of sterile water for injection or saline can be provided so that the ingredients can be mixed prior to administration.

[932] Suitable formulations further include aqueous and non-aqueous sterile injection solutions that can contain antioxidants, buffers, bacteriostats, bactericidal antibiotics and solutes that render the formulation isotonic with the bodily fluids of the intended recipient; and aqueous and non-aqueous sterile suspensions, which can include suspending agents and thickening agents.

[933] In certain embodiments, a pharmaceutical composition of the present invention is formulated as a depot preparation. Certain such depot preparations are typically longer acting than non-depot preparations. In certain embodiments, such preparations are administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. In certain embodiments, depot preparations are prepared using suitable polymeric or hydrophobic materials (for example an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[934] In certain embodiments, a pharmaceutical composition of the present invention comprises a delivery system. Examples of delivery systems include, but are not limited to,

liposomes and emulsions. Certain delivery systems are useful for preparing certain pharmaceutical compositions including those comprising hydrophobic compounds. In certain embodiments, certain organic solvents such as dimethylsulfoxide are used.

[935] In certain embodiments, a pharmaceutical composition of the present invention comprises a co-solvent system. Certain of such co-solvent systems comprise, for example, benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. In certain embodiments, such co-solvent systems are used for hydrophobic compounds. A non-limiting example of such a co-solvent system is the VPD co-solvent system, which is a solution of absolute ethanol comprising 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant Polysorbate 80 and 65% w/v polyethylene glycol 300. The proportions of such co-solvent systems may be varied considerably without significantly altering their solubility and toxicity characteristics. Furthermore, the identity of co-solvent components may be varied: for example, other surfactants may be used instead of Polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g., polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose.

[936] In certain embodiments, a pharmaceutical composition of the present invention comprises a sustained-release system. A non-limiting example of such a sustained-release system is a semi-permeable matrix of solid hydrophobic polymers. In certain embodiments, sustained-release systems may, depending on their chemical nature, release pharmaceutical agents over a period of hours, days, weeks or months.

[937] Appropriate pharmaceutical compositions of the present disclosure can be determined according to any clinically-acceptable route of administration of the composition to the subject. The manner in which the composition is administered is dependent, in part, upon the cause and/or location. One skilled in the art will recognize the advantages of certain routes of administration. The method includes administering an effective amount of the agent or compound (or composition comprising the agent or compound) to achieve a desired biological response, e.g., an amount effective to alleviate, ameliorate, or prevent, in whole or in part, a symptom of a condition to be treated, e.g., oncology and neurology disorders. In various aspects, the route of administration is systemic, e.g., oral or by injection. The agents or compounds, or pharmaceutically acceptable salts or derivatives thereof, are administered orally, nasally, transdermally, pulmonary, inhalationally, buccally, sublingually, intraperitoneally, subcutaneously, intramuscularly, intravenously, rectally, intrapleurally, intrathecally, intraportally, and parenterally. Alternatively or in addition, the route of

administration is local, e.g., topical, intra-tumor and peri-tumor. In some embodiments, the compound is administered orally.

[938] In certain embodiments, a pharmaceutical composition of the present disclosure is prepared for oral administration. In certain of such embodiments, a pharmaceutical composition is formulated by combining one or more agents and pharmaceutically acceptable carriers. Certain of such carriers enable pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a subject. Suitable excipients include, but are not limited to, fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). In certain embodiments, such a mixture is optionally ground and auxiliaries are optionally added. In certain embodiments, pharmaceutical compositions are formed to obtain tablets or dragee cores. In certain embodiments, disintegrating agents (e.g., cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate) are added.

[939] In certain embodiments, dragee cores are provided with coatings. In certain such embodiments, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to tablets or dragee coatings.

[940] In certain embodiments, pharmaceutical compositions for oral administration are push-fit capsules made of gelatin. Certain of such push-fit capsules comprise one or more pharmaceutical agents of the present invention in admixture with one or more filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In certain embodiments, pharmaceutical compositions for oral administration are soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. In certain soft capsules, one or more pharmaceutical agents of the present invention are dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added.

[941] In certain embodiments, pharmaceutical compositions are prepared for buccal administration. Certain of such pharmaceutical compositions are tablets or lozenges formulated in conventional manner.

[942] In certain embodiments, a pharmaceutical composition is prepared for transmucosal administration. In certain of such embodiments penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

[943] In certain embodiments, a pharmaceutical composition is prepared for administration by inhalation. Certain of such pharmaceutical compositions for inhalation are prepared in the form of an aerosol spray in a pressurized pack or a nebulizer. Certain of such pharmaceutical compositions comprise a propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In certain embodiments using a pressurized aerosol, the dosage unit may be determined with a valve that delivers a metered amount. In certain embodiments, capsules and cartridges for use in an inhaler or insufflator may be formulated. Certain of such formulations comprise a powder mixture of a pharmaceutical agent of the invention and a suitable powder base such as lactose or starch.

[944] In other embodiments the compound of the present disclosure are administered by the intravenous route. In further embodiments, the parenteral administration may be provided in a bolus or by infusion.

[945] In certain embodiments, a pharmaceutical composition is prepared for rectal administration, such as a suppository or retention enema. Certain of such pharmaceutical compositions comprise known ingredients, such as cocoa butter and/or other glycerides.

[946] In certain embodiments, a pharmaceutical composition is prepared for topical administration. Certain of such pharmaceutical compositions comprise bland moisturizing bases, such as ointments or creams. Exemplary suitable ointment bases include, but are not limited to, petrolatum, petrolatum plus volatile silicones, and lanolin and water in oil emulsions. Exemplary suitable cream bases include, but are not limited to, cold cream and hydrophilic ointment.

[947] In certain embodiments, the therapeutically effective amount is sufficient to prevent, alleviate or ameliorate symptoms of a disease or to prolong the survival of the subject being treated. Determination of a therapeutically effective amount is well within the capability of those skilled in the art.

[948] In certain embodiments, one or more compounds of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt or solvate thereof are formulated as a prodrug. In certain embodiments, upon in vivo administration, a prodrug is chemically converted to the biologically, pharmaceutically or therapeutically more active form. In certain embodiments, prodrugs are useful because they are easier to administer than the corresponding active form. For example, in certain instances, a prodrug may be more bioavailable (e.g., through oral

administration) than is the corresponding active form. In certain instances, a prodrug may have improved solubility compared to the corresponding active form. In certain embodiments, prodrugs are less water soluble than the corresponding active form. In certain instances, such prodrugs possess superior transmittal across cell membranes, where water solubility is detrimental to mobility. In certain embodiments, a prodrug is an ester. In certain such embodiments, the ester is metabolically hydrolyzed to carboxylic acid upon administration. In certain instances the carboxylic acid containing compound is the corresponding active form. In certain embodiments, a prodrug comprises a short peptide (polyaminoacid) bound to an acid group. In certain of such embodiments, the peptide is cleaved upon administration to form the corresponding active form.

[949] In certain embodiments, a prodrug is produced by modifying a pharmaceutically active compound such that the active compound will be regenerated upon in vivo administration. The prodrug can be designed to alter the metabolic stability or the transport characteristics of a drug, to mask side effects or toxicity, to improve the flavor of a drug or to alter other characteristics or properties of a drug. By virtue of knowledge of pharmacodynamic processes and drug metabolism in vivo, those of skill in this art, once a pharmaceutically active compound is known, can design prodrugs of the compound (see, e.g., Nogrady (1985) *Medicinal Chemistry A Biochemical Approach*, Oxford University Press, New York, pages 388-392).

[950] In various aspects, the amount of the compound of formula (I)-(VI) and/or (A)-(H-I), or a pharmaceutically acceptable salt or solvate thereof, or compounds disclosed in Tables A and B, or a pharmaceutically acceptable salt or solvate thereof, can be administered at about 0.001 mg/kg to about 100 mg/kg body weight (e.g., about 0.01 mg/kg to about 10 mg/kg or about 0.1 mg/kg to about 5 mg/kg).

[951] The concentration of a disclosed compound in a pharmaceutically acceptable mixture will vary depending on several factors, including the dosage of the compound to be administered, the pharmacokinetic characteristics of the compound(s) employed, and the route of administration. The agent may be administered in a single dose or in repeat doses. The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. Treatments may be administered daily or more frequently depending upon a number of factors, including the overall health of a patient, and the formulation and route of administration of the selected compound(s). An ordinarily skilled physician or veterinarian can readily determine

and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

[952] The compounds or pharmaceutical compositions of the present disclosure may be manufactured and/or administered in single or multiple unit dose forms.

[953] Having now generally described the invention, the same will be more readily understood through reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.

EXAMPLES

[954] The disclosure now being generally described, it will be more readily understood by reference to the following examples which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

[955] Synthetic Preparation

[956] The novel compounds of the present invention can be prepared in a variety of ways known to one skilled in the art of organic synthesis. The compounds of the present invention can be synthesized using the methods as hereinafter described below, together with synthetic methods known in the art of synthetic organic chemistry or variations thereon as appreciated by those skilled in the art.

[957] Preparation of compounds can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the selection of appropriate protecting groups can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in Greene and Wuts, *Protective Groups in Organic Synthesis*, 4th. Ed., Wiley & Sons, 2006, as well as in Jerry March, *Advanced Organic Chemistry*, 4th edition, John Wiley & Sons, publisher, New York, 1992 which are incorporated herein by reference in their entirety.

[958] Compounds of the present invention can be prepared by the literature methods cited in the following text. The following schemes depict established, known syntheses of these scaffolds.


[959] The groups and/or the substituents of the compounds of the present invention can be synthesized and attached to these scaffolds by the literature methods cited in the following text. The following schemes depict the known techniques for accomplishing this joinder.

[960] General Synthesis

[961] Compounds of the present invention can be synthesized using the following methods. General reaction conditions are given, and reaction products can be purified by general known methods including crystallization, silica gel chromatography using various organic solvents such as hexane, cyclohexane, ethyl acetate, methanol and the like, preparative high pressure liquid chromatography or preparative reverse phase high pressure liquid chromatography.

[962] **Representative Synthesis**

[963] **Example 1: Synthesis of 5-[[4-[1-[3,5-dichloro-4-(3-chloropropoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-4-methylsulfonyl-oxazole (A3)**



[964] **(4-Methylsulfonyloxazol-5-yl) 4-methylbenzenesulfonate (2):** To a solution of 4-methylsulfonyloxazol-5-ol (1) (0.10 g, 0.56 mmol), TEA (0.11 g, 1.1 mmol) and DMAP (14 mg, 0.11 mol) in DCM (3 mL) was added Tosyl chloride (0.10 g, 0.52 mmol) dropwise at 0 °C. Then the solution was stirred at the same temperature for 1 hour. TLC showed the reaction was completed. The mixture was poured into H₂O (5 mL), extracted with DCM (3 mL x 3). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give (4-methylsulfonyloxazol-5-yl) 4-methylbenzenesulfonate (2) (0.11 g, yield: 56.6 %) as yellow oil. ¹H NMR (400 MHz, CHCl₃-d) δ 7.98 (s, 1 H), 7.93 (d, *J*=8.40 Hz, 2 H), 7.42 (d, *J*=8.80 Hz, 2 H), 4.95 (s, 2 H), 3.22 (s, 3 H).

[965] **Methyl 3,5-dichloro-4-(3-chloropropoxy)benzoate (5):** To a mixture of methyl 3,5-dichloro-4-hydroxybenzoate (3) (100.0 g, 0.45 mol) and Cs₂CO₃ (293.2 g, 0.90 mol) in DMF (1.5 L) was added 1-chloro-3-iodopropane (4) (102.2 g, 0.50 mol), and the mixture was

stirred at 80 °C for 16 hours. TLC showed the reaction was completed. The mixture was cooled down, quenched with water (3.5 L) and extracted with MTBE (1.0 L x 2). The combined organic layers were washed with brine (1 L), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica-gel column chromatography to give methyl 3,5-dichloro-4-(3-chloropropoxy)benzoate (70.0 g, yield: 52.0%) as yellow oil. ¹H NMR (400MHz, CHCl₃-*d*) δ 2.29 (q, *J*=6.03 Hz, 2 H), 3.85 (t, *J*=6.36 Hz, 2 H), 3.91 (s, 3 H), 4.22 (t, *J*=5.70 Hz, 2 H), 7.92 - 8.00 (m, 2 H).


[966] 2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-ol (6): To a solution of 3,5-dichloro-4-(3-chloropropoxy)benzoate (**5**) (70.0 g, 0.24 mol) in THF (500 mL) was added MeMgBr (400 mL, 1.20 mol) at 0 °C under N₂ atmosphere. The mixture was stirred at 20 °C for 16 hours under N₂ atmosphere. TLC showed the reaction was completed. The mixture was poured into H₂O (500 mL), extracted with EtOAc (200 mL x 5), and the combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-ol (44.0 g, yield: 62.9%) as yellow oil. ¹H NMR (400MHz, CHCl₃-*d*) δ 1.56 (s, 6 H), 2.29 (q, *J*=6.06 Hz, 2 H), 3.87 (t, *J*=6.39 Hz, 2 H), 4.13 - 4.19 (m, 2 H), 7.41 (s, 2 H).

[967] 4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenol (7): To a solution of 2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-ol (**6**) (20.0 g, 67.2 mmol) and phenol (7.0 g, 73.9 mmol) in CCl₄ (400 mL) was added BF₃·Et₂O (16.6 mL, 134.2 mmol) dropwise at 0 °C under N₂ atmosphere. The mixture was stirred at 0 °C for 30 min. TLC showed the reaction was completed. The resulting mixture was poured into H₂O (300 mL), extracted with DCM (300 mL x 3). The combined organic layers were washed with brine (500 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenol (18.0 g, yield: 71.7 %) as yellow oil. ¹H NMR (400MHz, CHCl₃-*d*) δ 7.12 (s, 2H), 7.10 - 7.05 (m, 2H), 6.79 - 6.75 (m, 2H), 4.65 (s, 1H), 4.15 (t, *J*=5.7 Hz, 2H), 3.86 (t, *J*=6.4 Hz, 2H), 2.28 (q, *J*=6.1 Hz, 2H), 1.62 (s, 6H).

[968] 5-[[4-[1-[3,5-Dichloro-4-(3-chloropropoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-4-methylsulfonyl-oxazole (A3): To a suspension of 4-[1-[3,5-dichloro-4-(3-chloropropoxy)phenyl]-1-methyl-ethyl]phenol (**7**) (0.135 g, 0.36 mmol) and Cs₂CO₃ (0.197 g, 0.6 mmol) in DMF (3 mL) was added (4-methylsulfonyloxazol-5-yl)methyl 4-methylbenzenesulfonate (**2**) (0.1 g, 0.3 mmol) at 25 °C. The mixture was stirred at 60 °C for 6 hours. LCMS showed the reaction was completed. The resulting mixture was poured into

H₂O (8 mL) and extracted with EtOAc (5 mL x 3). The combined organic layers were washed with brine (5 mL x 2), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by prep-HPLC (HCl) to give 5-[[4-[1-[3,5-dichloro-4-(3-chloropropoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-4-methylsulfonyl-oxazole (**A1**) (37.9 mg, yield:23.6%) as yellow oil. HPLC purity (220 nm): 96.25%. ¹H NMR (400 MHz, CHCl₃-d) δ 7.99 (s, 1H), 7.16-7.10 (m, 4H), 6.94 (d, *J*=8.82 Hz, 2H), 5.42 (s, 2H), 4.15 (t, *J*=5.73 Hz, 2H), 3.86 (t, *J*=6.50 Hz, 2H) 3.18 (s, 3H), 2.28 (quin, *J*=6.17 Hz, 2H), 1.62 (s, 6H). LCMS (M + 23) *m/z*: calcd 533; found 556.

[699] **Example 2:** Synthesis of 4-((4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy) methyl)-1-(methylsulfonyl)-1H-imidazole (**A5**)



[970] **(1-Trityl-1H-imidazol-4-yl)methanol (2):** To a solution of (1H-imidazol-4-yl)methanol (0.5 g, 5.3 mmol) and Et₃N (1.1 g, 10.2 mmol) in DCM (20 mL) was added TrCl (1.5 g, 5.3 mmol) and the mixture was stirred at 50 °C for 4 hours. TLC showed the reaction was completed. The reaction was cooled down, quenched with water (40 mL), extracted with DCM (5 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica-gel column chromatography to give (1-trityl-1H-imidazol-4-yl)methanol (**2**) (1.0 g, yield: 76.8%) as colorless oil. ¹H NMR (400MHz, CHCl₃-d) δ = 7.37-7.44 (m, 9H), 7.29 (d, *J*=1.34 Hz, 2H), 7.10 (dd, *J*=8.01, 1.41 Hz, 6H), 4.87 (t, *J*=5.56 Hz, 1H), 4.33 (d, *J*=5.50 Hz, 2H).


[971] **4-(Chloromethyl)-1-trityl-1H-imidazole (3):** To a solution of (1-tritylimidazol-4-yl)methanol (**2**) (200 mg, 0.6 mmol) in DCM (3 mL) was added SOCl₂ (0.2 mL, 2.9 mmol) dropwise at 0 °C and the mixture was stirred at 25 °C for 10 min. TLC showed the reaction was completed. The reaction was concentrated under reduced pressure to give 4-(chloromethyl)-1-trityl-1H-imidazole (**3**) (150 mg, yield: 71.1 %) as yellow oil.

[972] **4-((4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-1-trityl-1H-imidazole (5):** A mixture of 4-(chloromethyl)-1-trityl-1H-imidazole (**3**) (150 mg, 0.4 mmol), Cs₂CO₃ (272 mg, 0.8 mmol) and 4-[1-[3,5-dichloro-4-(3-chloropropoxy)phenyl]-1-methyl-ethyl]phenol (**4**) (156 mg, 0.4 mmol) in MeCN (5 mL) was stirred at 80 °C for 4 hours. TLC showed the reaction was completed. The resulting mixture was cooled down, poured into H₂O (10 mL), extracted with EtOAc (4 mL x 2), and the combined organic layers were washed with brine (4 mL x 5), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica-gel column chromatography to give 4-((4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-1-trityl-1H-imidazole (**5**) (150 mg, yield: 71.1 %) as white solid. ¹H NMR (400MHz, CHCl₃-d) δ = 7.45 (d, *J*=1.2 Hz, 1H), 7.38-7.30 (m, 9H), 7.19-7.06 (m, 10H), 6.95-6.86 (m, 3H), 4.99 (s, 2H), 4.14-4.10 (m, 2H), 3.86 (t, *J*=6.5 Hz, 2H), 2.28 (quin, *J*=6.1 Hz, 2H), 1.62 (s, 6H).

[973] **4-((4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-1H-imidazole (6):** A solution of 4-((4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-1-trityl-1H-imidazole (**5**) (150 mg, 0.2 mmol) in HCl/EtOAc (4 M, 2 mL) was stirred at 25 °C for 2 hours. LCMS showed the reaction was completed. The reaction was filtered and the filtrate concentrated under reduced pressure to give 4-((4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-1H-imidazole (**6**) (95 mg, yield: 97.2 %) as colorless oil. LCMS (*M* + 1) *m/z*: calcd 452.1; found 452.9.

[974] **4-((4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-1-(methyl sulfonyl)-1H-imidazole (A5):** To a mixture of 4-((4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-1H-imidazole (**6**) (80 mg, 0.2 mmol) and TEA (0.1 mL, 0.5 mmol) in DCM (2 mL) was added methanesulfonyl chloride (41 mg, 0.4 mmol) dropwise at 0 °C, and the mixture was stirred at 25 °C for 2 hours. TLC showed the reaction was completed. The mixture was diluted with water (20 mL), extracted with DCM (5 mL x 3), and the combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (NH₄HCO₃) to give 4-((4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-1-(methyl sulfonyl)-imidazole (**A5**) (16 mg, yield: 16.6 %) as colorless oil. ¹H NMR (400MHz, CHCl₃-d) δ = 7.99 (d, *J*=1.3 Hz, 1H), 7.40 (s, 1H), 7.15-7.12 (m, 4H), 6.95-6.90 (m, 2H), 5.05 (s, 2H), 4.15 (t, *J*=5.7 Hz, 2H), 3.86 (t, *J*=6.4 Hz, 2H), 3.30 (s, 3H), 2.31-2.26 (m, 2H), 1.63 (s, 6H). LCMS (220 nm): 95.2%. LCMS (*M* + 1) *m/z*: calcd 530.1; found 531.0.

[975] **Example 3:** Synthesis of 2-((4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-5-(methylsulfonyl)-1,3,4-oxadiazole (**A7**)




[976] **Methyl 2-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)acetate (3):** To a solution of 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenol (**1**) (1.00 g, 2.6 mmol) and methyl 2-bromoacetate (**2**) (0.49 g, 3.2 mmol) in DMF (10 mL) was added Cs_2CO_3 (1.30 g, 4.0 mmol) at 25 °C. The mixture was stirred at 50 °C for 4 hours. LCMS showed the reaction was completed. The mixture was quenched with water (20 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (20 mL x 5), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give the methyl 2-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)acetate (**3**) (1.00 g, yield: 70 %) as colorless oil. LCMS ($M + 23$) m/z : calcd 444.07; found 467.1.

[977] **2-(4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)acetohydrazide (4):** To a solution of methyl 2-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)acetate (**3**) (1.32 g, 2.9 mmol) in MeOH (10 mL) was added $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$ (0.59 g, 11.8 mmol) at 20°C. The mixture was stirred at 90 °C for 16 hours. TLC showed of the reaction was completed. The mixture was quenched with water (10 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (10 mL x2), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give 2-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)acetohydrazide (**4**) (1.0 g, yield: 75.8%) as colorless oil. ^1H NMR (400 MHz, $\text{CHCl}_3\text{-d}$) δ ppm: 7.73 (br, 1 H), 7.11 - 7.14 (d, $J=8.9$ Hz, 2 H), 7.11 (s, 2 H), 6.84 (d, $J=8.9$ Hz, 2 H), 4.58 (s, 2 H), 4.14 - 4.16 (t, $J=5.2$ Hz, 2 H), 3.94 (br, 1 H), 3.86 (t, $J=6.4$ Hz, 2 H), 2.28 (m, 2 H), 1.62 (s, 6 H).

[978] **3-(1-(4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)ethyl)-5-methylthio)-4H-pyrazole (5):** To a solution of 2-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy) acetohydrazide (**4**) (500 mg, 1.12 mmol) in MeOH (2 mL) was added KOH (75 mg, 1.33 mmol) and CS₂ (76 mg, 2.8 mmol) at 20 °C. The mixture was stirred at 65 °C for 2 hours. Then CH₃I (2 mL) was added to the reaction mixture and stirred at 65 °C for 2 hours. LCMS showed the reaction was completed. The mixture was quenched with water (20 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give 3-(1-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)ethyl)-5-(methylthio)-4H-pyrazole (**5**) (300 mg, yield: 54.5%) as colorless oil. ¹H NMR (400 MHz, CHCl₃-d) δ ppm 7.14 (d, *J*=8.9 Hz, 2 H), 7.11 (s, 2 H), 6.94 (d, *J*=8.9 Hz, 2 H), 5.22 (s, 1 H), 4.15 (t, *J*=5.8 Hz, 2 H), 3.86 (t, *J*=6.4 Hz, 2 H), 2.74 (s, 3 H), 2.28 (t, *J*=5.99 Hz, 2 H), 1.62 (s, 6 H).

[979] **2-((4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-5-(methylsulfonyl)-1,3,4-oxadiazole (A7):** To a solution of 3-(1-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)ethyl)-5-(methylthio)-4H-pyrazole (**5**) (220 mg, 0.49 mmol) in DCM (5 mL) was added m-CPBA (85% purity, 226 mg, 4.03 mmol) at 0 °C. The reaction was stirred at 20 °C for 4 hours. LCMS showed the reaction was completed. The mixture was quenched with saturated aqueous Na₂S₂O₃ (5 mL) and saturated aqueous NaHCO₃ (5 mL), then extracted with DCM (10 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (TFA) to give 2-((4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-5-(methylsulfonyl)-1,3,4-oxadiazole (**A7**) (74 mg, yield: 31.6%) as colorless oil. ¹H NMR (400 MHz, DMSO-d₆) δ ppm 7.14 (d, *J*=8.9 Hz, 2 H), 7.11 (s, 2 H), 6.95 (d, *J*=8.8 Hz, 2 H), 5.35 (s, 2 H), 4.15 (t, *J*=5.73 Hz, 2 H), 3.86 (t, *J*=6.39 Hz, 2 H), 3.50 (s, 3 H), 2.28 (t, *J*=6.06 Hz, 2 H), 1.63 (s, 6 H). LCMS (220 nm): 97%. LCMS M+H⁺ *m/z*: calcd 532.04, found 533.1.

[980] **Example 4: Synthesis of 5-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)-4-(methylsulfonyl)oxazole (A13)**



[981] Ethyl 4-bromooxazole-5-carboxylate (2): To a solution of ethyl oxazole-5-carboxylate (**1**) (100.0 g, 0.71 mol) in THF (500 mL) and DMPU (500 mL) was added LiHMDS (1.0 M, 921 mL, 921 mmol) dropwise at -78 °C under N₂ atmosphere, and the mixture was stirred for 1.5 hours. Then Br₂ (54.5 mL, 1.06 mol) was added dropwise at the same temperature and the resulting mixture was stirred for 0.5 hours. TLC showed the starting material disappeared. The resulting solution was poured into a room temperature saturated aqueous NH₄Cl (1000 mL), extracted with EtOAc (500 mL x 3). The combined organic layers were washed with brine (500 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give ethyl 4-bromooxazole-5-carboxylate (25.0 g, purity: 90 %, yield: 11 %) as yellow oil. ¹H NMR (400MHz, CHCl₃-d): δ 7.94 (s, 1H), 4.43 (q, *J*=7.3 Hz, 2H), 1.42 (t, *J*=7.2 Hz, 3H)

[982] Ethyl 4-(methylthio)oxazole-5-carboxylate (3): A mixture of ethyl 4-bromooxazole-5-carboxylate (**2**) (28.0 g, 127 mmol) and NaSMe (10.8 g, 153 mmol) in EtOH (400 mL) was stirred at 80 °C for 2 hours. LCMS showed the reaction was completed. The resulting solution was cooled down and concentrated under reduced pressure. The residue was diluted with water (300 mL) and extracted with EtOAc (300 mL x 3). The combined organic layers were washed with brine (300 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The

residue was purified by MPLC to give ethyl 4-methylsulfanyloxazole-5-carboxylate (13.0 g, yield: 80%, yield: 43.7 %) as yellow oil. ¹H NMR (400MHz, CHCl₃-d): δ 7.91 (s, 1H), 4.40 (q, *J*=7.2 Hz, 2H), 2.62 (s, 3H), 1.41 (t, *J*=7.2 Hz, 3H).

[983] Ethyl 4-(methylsulfonyl)oxazole-5-carboxylate (4): A solution of ethyl 4-methylsulfanyloxazole-5-carboxylate (3) (2.9 g, 15.5 mmol) and m-CPBA (10.0 g, 46.5 mmol) in DCM (10 mL) was stirred at 25 °C for 16 hours. The reaction was completed detected by TLC. The reaction was quenched by sat. aqueous Na₂S₂O₃ (100 mL) and sat. aqueous NaHCO₃ (100 mL). The mixture was extracted with DCM (60 mL x 3). The combined organic layers were washed with brine (50 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give ethyl 4-methylsulfonyloxazole-5-carboxylate (1.20 g, yield: 35.3 %) as colorless oil. ¹H NMR (400MHz, CHCl₃-d): δ 8.04 (s, 1H), 4.50 (q, *J*=7.1 Hz, 2H), 3.38 (s, 3H), 1.44 (t, *J*=7.2 Hz, 3H).

[984] (4-(Methylsulfonyl)oxazol-5-yl)methanol (5): To a solution of ethyl 4-methylsulfonyloxazole-5-carboxylate (4) (1.2 g, 5.47 mmol) in THF (10 mL) was added LiBH₄ (179 mg, 8.21mmol) at 0 °C. The mixture was stirred at 0 °C for 1 hour. TLC showed the reaction completed. The mixture was quenched with aqueous HCl solution (3M, 2 mL) at 0 °C and extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine (3 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give (4-methylsulfonyloxazol-5-yl)methanol (900 mg, yield: 92.8 %) as colorless oil. ¹H NMR (400MHz, CHCl₃-d): δ = 7.92 (s, 1H), 4.93 (s, 2H), 3.24 (s, 3H).

[985] 5-(Chloromethyl)-4-(methylsulfonyl)oxazole (6): To a solution of ethyl 4-methylsulfonyloxazole-5-carboxylate (5) (700 mg, 3.95 mmol) in DCM (10 mL) and DMF (0.5 mL) was added SOCl₂ (1.44 mL, 19.8 mmol) at 0 °C. The mixture was stirred at 25 °C for 2 hours. TLC showed the reaction completed. The mixture was quenched with H₂O (15 mL), extracted with DCM (10 mL x 3). The combined organic layers were washed with brine (10 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 5-(chloromethyl)-4-methylsulfonyl-oxazole (700 mg, yield: 90.6 %) as yellow oil. ¹H NMR (400MHz, CHCl₃-d) δ = 7.99 (s, 1H), 4.94 (s, 2H), 3.22 (s, 3H).

[986] Methyl 3,5-dichloro-4-(2-chloroethoxy)benzoate (9): A mixture of methyl 3,5-dichloro-4-hydroxy-benzoate (7) (10.0 g, 45.2 mmol), Cs₂CO₃ (36.9 g, 113 mmol) and 1-bromo-2-chloro-ethane (26.0 g, 181 mmol) in DMF (60.0 mL) was stirred at 70 °C for 1 hour. TLC showed the reaction completed. The resulting solution was cooled down and diluted with water (150 mL), extracted with EtOAc (100 mL x 3). The combined organic layers were

washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give methyl 3,5-dichloro-4-(2-chloroethoxy)benzoate (10.0 g, yield: 78.0 %) as a yellow oil. ¹H NMR (400MHz, CHCl₃-d) δ = 8.01-7.97 (m, 2H), 4.35 (t, J=6.2 Hz, 2H), 3.93 (s, 3H), 3.89 (t, J=6.2 Hz, 2H).


[987] 2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-ol (10): To a solution of MeMgBr (3 M, 65.6 mL, 197 mmol) in THF (30 mL) was added methyl 3,5-dichloro-4-(2-chloroethoxy)benzoate (9) (9.3 g, 32.8 mmol) in THF (50 mL) dropwise at 0 °C. The reaction was stirred at 0 °C for 1 hour and TLC showed the reaction completed. The mixture was added to sat. aqueous NH₄Cl solution (300 mL) and extracted with EtOAc (200 mL x 3). The combined organic layers were washed with brine (200 mL), dried over Na₂SO₄, filtered and concentrated to give 2-[3,5-dichloro-4-(2-chloroethoxy)phenyl]propan-2-ol (9.0 g, yield: 96.8 %) as yellow oil. ¹H NMR (400MHz, CHCl₃-d) δ = 7.42 (s, 2H), 4.27 (t, J=6.3 Hz, 2H), 3.87 (t, J=6.3 Hz, 2H), 1.55 (s, 6H).

[988] 4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenol (11): To a solution of 2-[3,5-dichloro-4-(2-chloroethoxy)phenyl]propan-2-ol (9.0 g, 31.7 mmol) (10) and phenol (3.58 g, 38.1 mmol) in CCl₄ (100 mL) was added BF₃·Et₂O (7.83 mL, 63.5 mmol) dropwise at 0 °C under N₂ atmosphere. The mixture was stirred at 0 °C for 1 h. LCMS TLC showed the reaction almost completed. The mixture was added to water (150 mL) and extracted with DCM (100 mL x 3). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, filtered and concentrated. The crude product was purified by reverse phase MPLC (neutral) to give 4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenol (7.0 g, yield: 61.3 %) as yellow oil. ¹H NMR (400MHz, CHLOROFORM-d) δ = 7.13 (s, 2H), 7.10 - 7.04 (m, 2H), 6.79 - 6.74 (m, 2H), 4.81 (br s, 1H), 4.26 (t, J=6.3 Hz, 2H), 3.86 (t, J=6.4 Hz, 2H), 1.62 (s, 6H).

[989] 5-((4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)-4-(methylsulfonyl)oxazole (A13): A solution of 5-(chloromethyl)-4-methylsulfonyl-oxazole (6) (500 mg, 2.56 mmol), 4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenol (11) (919 mg, 2.56 mmol) and Cs₂CO₃ (1.67 g, 5.11 mmol) in DMF (20 mL) was stirred at 25 °C for 2 hours. Then the resulting solution was stirred at 40 °C for 0.5 hr. The reaction was completed detected by TLC. The reaction was quenched with water (50 mL), and the mixture was extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (30 mL x 2), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by MPLC to give 5-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-

ethyl]phenoxy]methyl]-4-methylsulfonyl-oxazole (528 mg, yield: 39.8 %) as white solid. ^1H NMR (400MHz, CHLOROFORM- d) δ = 7.92 (s, 1H), 7.09 - 7.02 (m, 4H), 6.89 - 6.83 (m, 2H), 5.34 (s, 2H), 4.18 (t, J =6.4 Hz, 2H), 3.79 (t, J =6.4 Hz, 2H), 3.11 (s, 3H), 1.54 (s, 6H). MS($M+H^+$) m/z : clcd.517.0; found 518.1, 540.0.

[990] Example 5: Synthesis of N-((3-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl)isoxazol-5-yl)methyl)methanesulfonamide (A22)



[991] 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl

trifluoromethane-sulfonate (2): To a solution of 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenol (1) (4.0 g, 10.7 mmol) and Et_3N (4.5 mL, 54.7 mmol) in DCM (40 mL) was added trifluoromethanesulfonic anhydride (5.4 g, 19.3 mmol) dropwise at 0°C under N_2 atmosphere. The mixture was stirred at the same temperature for 1.5 hrs. TLC showed the reaction was completed. The mixture was poured into H_2O (60 mL), extracted with DCM (20 mL x 3). The combined organic layers were washed with brine (20 mL x 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by MPLC to give 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl trifluoromethanesulfonate (4.0 g, yield: 73.4 %) as yellow oil. ^1H NMR (400MHz, CHCl_3 - d) δ 7.29-7.27 (m, 2H), 7.22-7.19 (m, 2H), 7.11 (s, 2H), 4.17 (t, J =6.3 Hz, 2H), 3.87 (t, J =6.3 Hz, 2H), 3.87 (t, J =6.2 Hz, 2H), 1.66 (s, 6H).

[992] **4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)benzaldehyde (3):** To a solution of 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl trifluoromethanesulfonate (**6**) (4.0 g, 7.9 mmol), Et₃SiH (1.9 mL, 11.9 mmol) and Et₃N (2.2 mL, 15.8 mmol) in DMF (50 mL) was added Pd(dppf)Cl₂ (290 mg, 0.4 mmol) at 20 °C. The reaction was stirred at 60 °C for 16 hours under CO atmosphere at 50 psi. TLC showed the reaction was completed. The resulting mixture was cooled down, poured into H₂O (100 mL), extracted with EtOAc (40 mL x 3). The combined organic layers were washed with brine (50 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)benzaldehyde (2.6 g, yield: 85.2 %) as yellow oil. ¹H NMR (400 MHz, CHCl₃-d) δ 10.02 (s, 1 H), 7.83 (d, *J*=8.8 Hz, 2 H), 7.41 (d, *J*=8.8 Hz, 2 H), 7.13 (s, 2 H), 4.18 (t, *J*=5.8 Hz, 2 H), 3.88 (t, *J*=6.4 Hz, 2 H), 2.31 (m, 2 H), 1.70 (s, 6 H).

[993] **(E)-4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)benzaldehyde oxime (4):** To a solution of 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)benzaldehyde (0.5 g, 1.3 mmol) (**3**) in DCM (5 mL) was added NH₂OH.HCl (117 mg, 1.7 mmol) and TEA (0.5 mL, 3.9 mmol) at 20 °C under N₂ atmosphere. The reaction was stirred at the same temperature for 16 hrs. LCMS showed the reaction was completed. The mixture was poured into H₂O (10 mL) and extracted with DCM (10 mL x 3). The organic layers were washed with brine (10 mL x 2), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)benzaldehyde oxime (**4**) (0.410 g, yield: 78.9 %) as white solid. ¹H NMR (400MHz, CHCl₃-d) δ = 8.13 (s, 1H), 7.52 (d, *J*=8.4 Hz, 2H), 7.37 (s, 1H), 7.23 (d, *J*=8.4 Hz, 2H), 7.12 (s, 2H), 4.15 (t, *J*=5.6 Hz, 2H), 3.86 (t, *J*=6.5 Hz, 2H), 2.28 (quin, *J*=6.1 Hz, 2H), 1.65 (s, 6H). MS(M+H⁺) *m/z*: calcd. 399.0; found 399.8.


[994] **tert-Butyl ((3-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl)isoxazol-5-yl)methyl)carbamate (6):** To a solution of 4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)benzaldehyde oxime (**4**) (320 mg, 0.80 mmol) in DCM (10 mL) was added aqueous NaClO (7.0 %, 1.3 g, 1.2 mmol) dropwise at 0 °C. The reaction was stirred for 5 min. Tert-butyl prop-2-yn-1-ylcarbamate (**5**) (120 mg, 0.80 mmol) and TEA (0.22 mL, 1.6 mmol) were added to the reaction mixture at 0 °C. The mixture was stirred at 0 °C~25 °C for 16 hours. LCMS showed ~50% of desired product and ~10% of starting material **2** remained. The mixture was poured into H₂O (10 mL), extracted with DCM (10 mL x 2). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure and purified by MPLC to give tert-butyl ((3-(4-(2-(3,5-

dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl)isoxazol-5-yl)methyl carbamate (0.22 g, yield: 51 %) as colorless oil. ¹H NMR (400MHz, CHCl₃-d) δ = 7.72 (d, *J*=8.60 Hz, 2 H) 7.31 (d, *J*=8.60 Hz, 2 H) 7.10-7.20 (m, 2 H) 6.48 (s, 1 H) 5.03 (br s, 1 H) 4.48 (br d, *J*=6.17 Hz, 2 H) 4.16 (t, *J*=5.73 Hz, 2 H) 3.86 (t, *J*=6.39 Hz, 2 H) 2.29 (q, *J*=6.06 Hz, 2 H) 1.68 (s, 6 H) 1.47 (s, 9 H). MS(M+H⁺) *m/z*: clcd. 552.1; found 553.2.

[995] **(3-(4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl)isoxazol-5-yl) methanamine hydrochloride (7)**: A solution of tert-butyl ((3-(4-(2-(3,5-dichloro-4-(3-chloropropoxy) phenyl)propan-2-yl)phenyl)isoxazol-5-yl)methyl)carbamate (**6**) (190 mg, 0.34 mmol) in HCl/EtOAc (4 M, 3 mL, 12 mmol) was stirred at 20 °C for 0.5 hour. LCMS showed the reaction was completed. The mixture was concentrated under reduced pressure to give (3-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl)isoxazol-5-yl)methanamine hydrochloride (**7**) (60 mg, yield: 36 %) as brown oil. MS(M+H⁺) *m/z*: clcd. 452.1; found 453.1.

[996] **N-((3-(4-(2-(3,5-Dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl)isoxazol-5-yl)methyl)methanesulfonamide (A22)**: To a solution of [3-[4-[1-[3,5-dichloro-4-(3-chloropropoxy)phenyl]-1-methyl-ethyl]phenyl]isoxazol-5-yl]methanamine (**7**) (60 mg, 0.13 mmol) in DCM (3 mL) was added TEA (40 mg, 0.40 mmol) and MsCl (18 mg, 0.16 mmol) under N₂ atmosphere at 0 °C. The reaction was stirred at 20 °C for 5 hrs. TLC showed the reaction was completed. The mixture was poured into H₂O (5 mL) and extracted with DCM (10 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (TFA) to give N-((3-(4-(2-(3,5-dichloro-4-(3-chloropropoxy)phenyl)propan-2-yl)phenyl)isoxazol-5-yl)methyl) methanesulfonamide (**A22**) (5 mg, yield: 7.11 %) as brown oil. LCMS purity (220 nm): 89.4%. ¹H NMR (400MHz, CHCl₃-d) δ = 7.73 (br d, *J*=7.9 Hz, 2H), 7.31 (br d, *J*=8.2 Hz, 2H), 7.14 (s, 2H), 6.60 (s, 1H), 4.89-4.80 (m, 1H), 4.54 (d, *J*=6.2 Hz, 2H), 4.16 (t, *J*=5.6 Hz, 2H), 3.86 (t, *J*=6.4 Hz, 2H), 2.99 (s, 3H), 2.33 - 2.25 (m, 2H), 1.68 (s, 6H). LCMS(M+H⁺) *m/z*: clcd. 530.0; found 531.0.

[997] **Example 6: Synthesis of N-(tert-Butyl)-3,5-dichloro-4-(2-chloroethoxy)-N-(4-((4-(methyl-sulfonyl)oxazol-5-yl)methoxy)phenyl)aniline (A31)**



[998] tert-Butyl-(4-iodophenoxy)-dimethyl-silane (2): To a solution of 4-iodophenol (**1**) (3.0 g, 13.6 mmol) and imidazole (1.9 g, 27.3 mmol) in DCM (30 mL) was added TBSCl (1.8 g, 12.3 mmol) at 25 °C. Then the mixture was stirred at the same temperature for 16 hours. TLC showed the reaction was completed. The reaction mixture was washed with water (10 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give tert-butyl-(4-iodophenoxy)-dimethyl-silane (**2**) (2.5 g, yield: 54.8%) as colorless oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.57-7.44 (m, 2H), 6.67-6.55 (m, 2H), 1.02-0.93 (m, 9H), 0.19 (s, 6H).

[999] 3,5-Dichloro-4-(2-chloroethoxy)aniline (5): To a solution of 4-amino-2,6-dichlorophenol (**3**) (5.0 g, 28.1 mmol) in DMF (50 mL) was added NaH (60% purity, 1.1 g, 28.1 mmol) in portions at 0 °C under N₂ atmosphere. After stirring at the same temperature for 15 min, 1-bromo-2-chloro-ethane (**4**) (4.4 g, 30.9 mmol) was added and the resulting mixture was stirred at 25 °C for 16 hours. TLC showed the reaction was completed. The resulting mixture was quenched with saturated aqueous NH₄Cl solution (100 mL) and extracted with EtOAc (50 mL x 3). The combined organic layers were washed with brine (50 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give 3,5-dichloro-4-(2-chloroethoxy)aniline (**5**) (2.5 g, yield: 37%) as yellow solid. ¹H NMR (400MHz, CHCl₃-d) δ 6.64-6.55 (m, 2H), 4.18 (t, *J*=6.4 Hz, 2H), 3.84 (t, *J*=6.4 Hz, 2H), 3.69-3.58 (m, 2H).

[1000] N-tert-butyl-3,5-dichloro-4-(2-chloroethoxy)aniline (7): To a solution of 3,5-dichloro-4-(2-chloroethoxy)aniline (**5**) (1.0 g, 4.16 mmol) and tert-butyl 2,2,2-trichloroethanimidate (**6**) (2.27 g, 10.4 mmol) in MeNO₂ (10 mL) was added Copper(II) trifluoromethanesulfonate (1.5 g, 4.16 mmol) under N₂ atmosphere. Then the resulting mixture was stirred at 25 °C for 16 hours. TLC showed the reaction completed. The mixture was

quenched with aqueous NaHCO₃ solution (5 mL) and extracted with EtOAc (5 mL x 3). The combined organic layers were washed with brine (5 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give the N-tert-butyl-3,5-dichloro-4-(2-chloroethoxy)aniline (**7**) (460 mg, yield: 37.3%) as a brown gum. ¹H NMR (400MHz, CHCl₃-d) δ 6.64 (s, 2H), 4.19 (t, *J*=6.4 Hz, 2H), 3.84 (t, *J*=6.4 Hz, 2H), 1.37-1.29 (m, 9H).


[1001] N-tert-Butyl-N-[4-[tert-butyl(dimethyl)silyl]oxyphenyl]-3,5-dichloro-4-(2-chloroethoxy)aniline (8**):** To a mixture of N-tert-butyl 3,5-dichloro-4-(2-chloroethoxy)aniline (**7**) (210 mg, 0.71 mmol), tert-butyl-(4-iodophenoxy)-dimethyl-silane (237 mg, 0.71 mmol) and t-BuONa (162 mg, 1.77 mmol) in toluene (5 mL) was added palladium (I) tritertbutyl phosphine bromide (55 mg, 0.07 mmol) under N₂ atmosphere. Then the resulting mixture was stirred at 110 °C for 16 hours. TLC showed the reaction was completed. The resulting mixture was cooled down and quenched with water (2 mL) and extracted with EtOAc (2 mL x 3). The combined organic layers were washed with brine (3 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give N-tert-butyl-N-[4-[tert-butyl(dimethyl)silyl]oxyphenyl]-3,5-dichloro-4-(2-chloroethoxy)aniline (**8**): (125 mg, yield: 35.1%) as yellow solid. ¹H NMR (400MHz, CHCl₃-d) δ 6.99-6.93 (m, 2H), 6.86-6.77 (m, 2H), 6.70 (s, 2H), 4.18 (t, *J*=6.4 Hz, 2H), 3.83 (t, *J*=6.5 Hz, 2H), 1.37 (s, 9H), 1.04-0.94 (m, 9H), 0.26-0.21 (m, 6H).

[1002] 4-[N-tert-Butyl-3,5-dichloro-4-(2-chloroethoxy)anilino]phenol (9**):** To a solution of N-tert-butyl-N-[4-[tert-butyl(dimethyl)silyl]oxyphenyl]-3,5-dichloro-4-(2-chloroethoxy)aniline (**8**) (150 mg, 0.29 mmol) in THF (5 mL) was added TBAF (1 M, 1.0 mL, 1 mmol). Then the resulting mixture was stirred at 25 °C for 2 hours. TLC showed the reaction was completed. The mixture was quenched with water (5 mL) and extracted with EtOAc (3 mL x 3). The combined organic layers were washed with water (5 mL x 2), dried over Na₂SO₄, filtered and concentrated to give the 4-[N-tert-butyl-3,5-dichloro-4-(2-chloroethoxy)anilino]phenol (**9**) (110 mg, yield: 94.9%) as yellow gum. ¹H NMR (400MHz, CHCl₃-d) δ 7.03-6.93 (m, 2H), 6.87-6.79 (m, 2H), 6.72 (s, 2H), 4.18 (t, *J*=6.4 Hz, 2H), 3.83 (t, *J*=6.4 Hz, 2H), 1.37 (s, 9H).

[1003] N-tert-Butyl-3,5-dichloro-4-(2-chloroethoxy)-N-[4-[(4-methylsulfonyloxazol-5-yl)-methoxy]-phenyl]aniline (A31**):** To a mixture of 4-[N-tert-butyl-3,5-dichloro-4-(2-chloroethoxy)anilino]phenol (**9**) (110 mg, 0.283 mmol) and Cs₂CO₃ (277 mg, 0.85 mmol) in DMF (5 mL) was added 5-(chloromethyl)-4-methylsulfonyl-oxazole (**10**) (83 mg, 0.42 mmol). Then the resulting mixture was stirred at 40 °C for 2 hours. LCMS showed the reaction was

completed. The mixture was cooled down, quenched with water (5 mL) and extracted with EtAOc (5 mL x 3). The combined organic layers were washed with brine (5 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by p-HPLC (TFA) to give the N-tert-butyl-3,5-dichloro-4-(2-chloroethoxy)-N-[4-[(4-methylsulfonyl-oxazol-5-yl)-methoxy]-phenyl]aniline (**A31**) (36.5 mg, yield: 23.5%) as yellow solid. HPLC purity (220 nm): 91.7%. ¹H NMR (400MHz, CHCl₃-d) δ 8.01 (s, 1H), 7.06-7.02 (m, 2H), 6.99-6.94 (m, 2H), 6.73 (s, 2H), 5.41 (s, 2H), 4.17 (t, *J*=6.4 Hz, 2H), 3.82 (t, *J*=6.4 Hz, 2H), 3.20 (s, 3H), 1.35 (s, 9H). LCMS (M + Na⁺) *m/z*: calcd 546.1; found 569.1.

[1004] Example 7: Synthesis of 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)-N-((4-(methylsulfonyl)oxazol-5-yl)methyl)aniline hydrochloride (**A32**)



[1005] 4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)-N-


(diphenylmethylene)aniline (3): To a mixture of 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl trifluoromethanesulfonate (**1**) (5.0 g, 10.2 mmol), diphenylmethanimine (**2**) (1.8 g, 10.2 mmol), Cs₂CO₃ (6.6 g, 20.3 mmol) and BINAP (1.3 g, 5.08 mmol) in 1,4-dioxane (100 mL) was added Pd(OAc)₂ (0.5 g, 2.0 mmol) at 25 °C under N₂ atmosphere. The mixture was stirred at 65 °C for 2 hrs. TLC showed the reaction was completed. The resulting solution was cooled down, quenched with H₂O (100 mL), extracted with EtOAc (30 mL x 2). The combined organic layers were washed with brine (40 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica-gel column chromatography to give 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)-N-(diphenylmethylene)aniline (4.5 g, yield: 84.6 %) as colorless oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.78 - 7.74 (m, 2H), 7.63 - 7.57 (m, 2H), 7.45 -

7.38 (m, 2H), 7.32 - 7.27 (m, 2H), 7.14 - 7.07 (m, 2H), 7.03 - 6.94 (m, 4H), 6.66 (d, $J=8.6$ Hz, 2H), 4.26 (t, $J=6.4$ Hz, 2H), 3.86 (t, $J=6.4$ Hz, 2H), 1.58 (s, 6H).

[1006] 4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)aniline hydrochloride (4): To a solution of 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)-N-(diphenylmethylene)aniline (**3**) (4.5 g, 8.6 mmol) in THF (50 mL) was added aqueous HCl (2 M, 15 mL) and the mixture was stirred at 25 ° C for 15 min. TLC showed the reaction was completed. The mixture was concentrated under reduced pressure and purified by silica-gel column chromatography to give 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)aniline hydrochloride (2.5 g, yield: 81.0%) as yellow solid. ^1H NMR (400MHz, CHCl_3-d) δ 10.54 (br s, 2H), 7.48 (br d, $J=8.6$ Hz, 2H), 7.31 - 7.27 (m, 2H), 7.09 (s, 2H), 4.26 (t, $J=6.4$ Hz, 2H), 3.86 (t, $J=6.3$ Hz, 2H), 1.63 (s, 6H).

[1007] 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)-N-((4-(methylsulfonyl)oxazol-5-yl)methyl)aniline hydrochloride (A32): To a suspension of 5-(chloromethyl)-4-(methylsulfonyl)oxazole (**5**) (200 mg, 0.5 mmol) and Ag_2CO_3 (564 mg, 0.2 mmol) in DMF (2 mL) was added 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)aniline (**4**) (382 mg, 0.1 mmol), and the mixture was stirred at 65 °C for 2 hours. TLC showed the reaction was completed. The resulting mixture was cooled down, poured into H_2O (6 mL), extracted with EtOAc (2 mL x 2). The combined organic layers were washed with brine (4 mL x 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (HCl) to give 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)-N-((4-(methylsulfonyl)oxazol-5-yl)methyl)aniline hydrochloride (**A32**) (20 mg, yield: 3.8 %) as white solid. ^1H NMR (400MHz, CHCl_3-d) δ 7.85 (s, 1H), 7.16-7.09 (m, 4H), 7.04-6.93 (m, 2H), 4.80 (s, 2H), 4.26 (t, $J=6.4$ Hz, 2H), 3.86 (t, $J=6.4$ Hz, 2H), 3.16 (s, 3H), 1.61 (s, 6H). LCMS ($\text{M}+\text{H}^+$) m/z : calcd: 516.0; found 517.0.

[1008] Example 8: Synthesis of 5-(1-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)ethyl)-4-(methylsulfonyl)oxazole (**A35**)



[1009] 4-(Methylthio)oxazole-5-carboxylic acid (2): To a solution of ethyl 4-methylsulfanyloxazole-5-carboxylate (**1**) (1 g, 80% purity, 4.3 mmol) in THF (10 mL) and H₂O (4 mL) was added NaOH (300 mg, 8.6 mmol) at 25 °C, and the mixture was stirred at the same temperature for 2 hours. LCMS showed the reaction was completed. The mixture was acidified to pH = 3 with aqueous HCl solution (3 M, 3 mL) and extracted with EtOAc (15 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 4-(methylthio)oxazole-5-carboxylic acid (**2**) (0.7 g, 80% purity, yield: 82.3%) as white solid. ¹H NMR (400MHz, CHCl₃-d) δ 7.99 (s, 1H), 2.65 (s, 3H). LCMS (M + 1) m/z: calcd: 531.0; found 532.0.

[1010] N-Methoxy-N-methyl-4-(methylthio)oxazole-5-carboxamide (4): To a mixture of 4-(methylthio)oxazole-5-carboxylic acid (**2**) (0.7 g, 4.4 mmol) in THF (5 mL) and DMF (5 mL) was added HATU (2.5 g, 6.6 mmol) at 25 °C under N₂ atmosphere. After stirring at the same temperature for 15 minutes, N-methoxymethanamine hydrochloride (**3**) (0.6 g, 5.7 mmol) and TEA (3.1 mL, 22.0 mmol) were added and the resulting mixture stirred at 25 °C for 2 hours. TLC showed the reaction was completed. The reaction was diluted with H₂O (10 mL), extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine (3 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give N-methoxy-N-methyl-4-(methylthio)oxazole-5-carboxamide (**4**) (0.5 g, yield: 56.2%) as yellow oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.90 (s, 1H), 4.12 (q, *J*=7.1 Hz, 1H), 3.81 (s, 3H), 3.30 (s, 3H), 2.58 (s, 3H).

[1011] 1-(4-(Methylthio)oxazol-5-yl)ethenone (5): To a mixture of N-methoxy-N-methyl-4-(methylthio)oxazole-5-carboxamide (**4**) (0.4 g, 1.9 mmol) in THF (8 mL) was added MeMgBr (3 M, 0.8 mL, 2.4 mmol) at 0 °C under N₂ atmosphere, and the resulting mixture was stirred at the same temperature for 1 hour. TLC showed the reaction was completed. The reaction was quenched with saturated aqueous NH₄Cl (4 mL), extracted with EtOAc (5 mL x 3). The combined organic layers were washed with brine (3 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 1-(4-(methylthio)oxazol-5-yl)ethenone (**5**) (0.4 g, 75% purity, yield: 96.2%) as yellow oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.89 (s, 1H), 2.62 (s, 3H), 2.50 (s, 3H).


[1012] 1-(4-(Methylthio)oxazol-5-yl)ethanol (6): To a mixture of 1-(4-(methylthio)oxazol-5-yl)ethanone (**5**) (0.4 g, 2.3 mmol) in THF (8 mL) was added NaBH₄ (0.1 g, 2.7 mmol) at 0 °C under N₂ atmosphere, and the mixture was stirred at the same temperature for 1 hour. TLC showed the reaction was completed. The reaction was quenched with saturated aqueous NH₄Cl (4 mL), extracted with EtOAc (5 mL x 3). The combined organic layers were washed with brine (3 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 1-(4-(methylthio)oxazol-5-yl)ethanol (**6**) (0.2 g, yield: 54.9 %) as yellow oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.83 (s, 1H), 5.12 (br d, *J*=5.5 Hz, 1H), 2.44 (s, 3H), 1.58 (d, *J*=6.6 Hz, 3H).

[1013] 5-(1-(4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)ethyl)-4-(methylthio)oxazole (8): To a mixture of 1-(4-(methylthio)oxazol-5-yl)ethanol (**6**) (0.2 g, 1.1 mmol), 4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl] phenol (**7**) (0.4 g, 1.1 mmol) and PPh₃ (0.6 g, 2.3 mmol) in THF (2 mL) was added DEAD (0.3 g, 1.7 mmol) at 0 °C under N₂ atmosphere, and the mixture was stirred at 25 °C for 16 hours. TLC showed the reaction was completed. The reaction was quenched with H₂O (5 mL), extracted with EtOAc (6 mL x 3). The combined organic layers were washed with brine (3 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give the 5-(1-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)ethyl)-4-(methylthio)oxazole (**8**) (50 mg, yield: 8.83 %) as yellow oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.85-7.82 (m, 1H), 7.11-7.03 (m, 4H), 6.88 (d, *J*=8.8 Hz, 2H), 5.51 (q, *J*=6.6 Hz, 1H), 4.24 (t, *J*=6.4 Hz, 2H), 3.84 (t, *J*=6.4 Hz, 2H), 2.35 (s, 3H), 1.69 (d, *J*=6.6 Hz, 3H), 1.58 (s, 6H).

[1014] 5-(1-(4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)ethyl)-4-(methylsulfonyl)oxazole (A35): To a mixture of 5-(1-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)ethyl)-4-(methylthio)oxazole (**8**) (50 mg, 0.1 mmol) was added mCPBA (80% purity, 64 mg, 0.3 mmol) in DCM (3 mL) at 25 °C, and the mixture was stirred at the same temperature for 16 hours. LCMS showed the reaction was

completed. The reaction was quenched with H₂O (5 mL), extracted with EtOAc (6 mL x 3). The combined organic layers were washed with brine (3 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by p-HPLC (TFA) 5-(1-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)ethyl)-4-(methylsulfonyl)oxazole (21.7 mg, yield: 40.8%) as white solid. HPLC purity (220 nm): 98.5%. ¹H NMR (400MHz, CHCl₃-d) δ = 7.94 (s, 1H), 7.14 - 7.03 (m, 4H), 6.92 (d, *J*=8.9 Hz, 2H), 6.10 (q, *J*=6.5 Hz, 1H), 4.26 (t, *J*=6.3 Hz, 2H), 3.86 (t, *J*=6.3 Hz, 2H), 3.06 (s, 3H), 1.74 (d, *J*=6.7 Hz, 3H), 1.59 (s, 6H). LCMS (M+H⁺) *m/z*: calcd: 531.0; found 532.0, 534.0.

[1015] Example 9: Synthesis of N-(4-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)oxazol-2-yl)methanesulfonamide (A38)



[1016] (2-Chlorooxazol-4-yl)methanol (2): To a solution of ethyl 2-chlorooxazole-4-carboxylate (**1**) (1.0 g, 5.7 mmol) in DCM (20 mL) was added DIBAL-H (1 M in toluene 8.5 mL, 8.5 mmol) dropwise at 0 °C under N₂ atmosphere. The mixture was stirred at the same temperature for 1 hour. TLC showed the reaction was completed. The mixture was quenched with saturated aqueous NH₄Cl (20 mL), then extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (20 mL x 2), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by MPLC to give (2-chlorooxazol-4-yl)methanol (**2**) (470 mg, yield: 62 %) as brown oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.59 (s, 1H), 4.58 (d, *J*=0.8 Hz, 2H).


[1017] (2-Chlorooxazol-4-yl)methyl methanesulfonate (3): To a mixture of (2-chlorooxazol-4-yl)methanol (**2**) (0.47 g, 3.52 mmol) and TEA (1 mL, 7.04 mol) in DCM (15 mL) was added methanesulfonyl chloride (444 mg, 3.87 mmol) at 0 °C. The mixture was stirred at 0-25 °C for 3 hours. TLC showed the reaction was completed. The mixture was poured into H₂O (30 mL), extracted with DCM (10 mL x 2). The combined organic layers were washed with brine (20

mL x 2), dried over Na₂SO₄, filtered and concentrated under reduce pressure to give (2-chlorooxazol-4-yl)methyl methanesulfonate (**3**) (800 mg, 70% purity) as yellow oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.77 (s, 1H), 5.14 (d, *J*=0.8 Hz, 2H), 3.08 (s, 3H).

[1018] 2-Chloro-4-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)oxazole (5): To a solution of (2-chlorooxazol-4-yl)methyl methanesulfonate (**3**) (0.70 g, 3.31 mmol) in DMF (10 mL) was added 4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenol (**4**) (1.19 g, 3.31 mmol) and Cs₂CO₃ (2.16 g, 6.62 mmol). The resulting mixture was stirred at 25 °C for 10 hours. LCMS showed the reaction was completed. The mixture was poured into H₂O (30 mL) and extracted with EtOAc (30 mL x 3). The combined organic layers were washed with brine (20 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduce pressure. The residue was purified by silica gel column chromatography to give 2-chloro-4-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]oxazole (0.50 g, yield: 31.8 %) as yellow oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.68 (s, 1H), 7.15-7.11 (m, 4H), 6.92-6.87 (m, 2H), 4.97 (d, *J*=0.8 Hz, 2H), 4.26 (t, *J*=6.4 Hz, 2H), 3.86 (t, *J*=6.4 Hz, 2H), 1.63 (s, 6H). LCMS (M+H⁺) *m/z*: calcd: 473.0; found 474.1.

[1019] N-(4-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)oxazol-2-yl)methanesulfonamide (A38): To a solution of 2-chloro-4-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]oxazole (**5**) (10 mg, 0.02mmol) in 1,4-dioxane (0.2 mL) was added methanesulfonamide (2.4 mg, 0.02 mmol), Brettphos Pd G3 (2 mg, w20%) and t-BuONa (3 mg, 0.03 mmol). The mixture was stirred at 80 °C for 10 hours under N₂ atmosphere. LCMS showed 5% desired MS and 90% starting material. The resulting 20 reaction mixtures were cooled down and combined. The mixture was filtered and the filtrate concentrated under reduced pressure. The residue was purified by prep-HPLC (TFA) to give N-[[5-bromo-4-[4-[1-[3,5-dichloro-4-(3-chloropropoxy)phenyl]-1-methyl-ethyl]phenyl]oxazol-2-yl]methyl]methanesulfonamide (2 mg, yield: 1.8 %) as pale yellow solid. LCMS (220 nm): 85.79%. ¹H NMR (400MHz, CHCl₃-d) δ 7.16 (d, *J*=8.8 Hz, 2H), 7.12 (s, 2H), 7.08 (s, 1H), 6.86 (d, *J*=8.8 Hz, 2H), 4.86 (s, 2H), 4.27 (t, *J*=6.4 Hz, 2H), 3.86 (t, *J*=6.4 Hz, 2H), 3.09 (s, 3H), 1.64 (s, 6H). LCMS (M+H⁺) *m/z*: calcd: 532.04; found 533.0.

[1020] Example 10: Synthesis of N-[3-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy] methyl]-1H-pyrazol-4-yl]methanesulfonamide (A40)



[1021] Methyl 4-nitro-1-tetrahydropyran-2-yl-pyrazole-3-carboxylate (3): To a solution of methyl 4-nitro-1H-pyrazole-5-carboxylate (**1**) (2.00 g, 1.17 mmol) in DCM (20 mL) was added 3,4-dihydro-2H-pyran (**2**) (3.11 g, 14.0 mmol) and p-Toluenesulfonic acid (0.20 g, 1.17 mmol) at 20 °C. The mixture was stirred at the same temperature for 4 hours. LCMS showed the reaction was completed. The reaction was poured into water (100 mL) and extracted with DCM (30 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give methyl 4-nitro-1-tetrahydropyran-2-yl-pyrazole-3-carboxylate (**3**) (2.7 g, yield: 90.5 %) as colorless oil. ¹H NMR (400 MHz, CHCl₃-d) δ ppm 8.39 (s, 1 H), 5.44 (dd, *J* = 9.04, 2.87 Hz, 1 H), 4.06-4.15 (m, 1 H), 4.00 (s, 3 H), 3.74 (ddd, *J* = 11.69, 9.04, 4.41 Hz, 1 H), 2.23 (dd, *J* = 12.79, 3.09 Hz, 1 H), 1.86-2.07 (m, 3 H), 1.63-1.77 (m, 4 H), 1.57 (s, 6 H). LCMS (220 nm): 68% Exact Mass 255.09; found 256.0.

[1022] (4-nitro-1-tetrahydropyran-2-yl-pyrazol-3-yl) methanol (4): To a solution of methyl 4-nitro-1-tetrahydropyran-2-yl-pyrazole-3-carboxylate (**3**) (1.00 g, 3.92 mmol) in THF (10 mL) was added DIBALH (7.8 mL, 7.8 mmol, 1.0 mol/L in toluene) at -78 °C under N₂ atmosphere and the mixture was stirred at 20 °C for 4 hours. LCMS showed the reaction was completed. The mixture was quenched with water (20 mL) and extracted with EtOAc (30 mL x 5). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give (4-nitro-1-tetrahydropyran-2-yl-pyrazol-3-yl)methanol (70.0 %, 0.90 g, yield: 70.8 %) as colorless oil. HPLC purity (220 nm): 69%. ¹H NMR (400 MHz, CHCl₃-d) δ ppm 8.39 (s, 1 H), 5.37 (dd, *J* = 9.17, 2.69 Hz, 1 H), 4.93 (s, 2 H), 4.06-4.14 (m, 1 H), 3.70 - 3.77 (m, 1 H), 2.13-2.22 (m, 1 H), 1.87-2.09 (m, 2 H), 1.64-1.77 (m, 4 H). LCMS (M+H⁺) *m/z*: calcd: 227.09; found 250.2.

[1023] 3-(Chloromethyl)-4-nitro-1-tetrahydropyran-2-yl-pyrazole (5): To a solution of (4-nitro-1-tetrahydropyran-2-yl-pyrazol-3-yl) methanol (**4**) (1.3 g, 5.72 mmol) in DCM (13 mL) was added triethylamine (1.2 g, 1.14 mmol) at 20 °C. Methanesulfonyl chloride (0.8 g, 6.87 mmol) was dropwise at 0 °C. The reaction was stirred at 20 °C for 4 hours. TLC showed the reaction was completed. The mixture was quenched with water (30 mL) and extracted with DCM (30 mL x 3). The combined organic layers were washed with brine (40 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give 3-(chloromethyl)-4-nitro-1-tetrahydropyran-2-yl-pyrazole (**5**) (60.0 %, yield: 30.9 %) as colorless oil. ¹H NMR (400 MHz, CHCl₃-d) δ ppm 8.39 (s, 1 H), 5.38 (dd, J = 9.26, 2.65 Hz, 1 H), 4.88 (s, 2 H), 3.75 (m, 2 H), 1.86 (m, 2 H), 1.67-1.77 (m, 4 H).

[1024] 3-[[4-[1-[3,5-Dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-4-nitro-1-tetrahydropyran-2-yl-pyrazole (7): To a solution of 3-(chloromethyl)-4-nitro-1-tetrahydropyran-2-yl-pyrazole (**5**) (0.20 g, 0.814 mmol) in DMF (2 mL) was added CS₂CO₃ (0.4 g, 1.22 mmol) and 4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenol (0.18 g, 0.49 mmol) (**6**) at 20 °C. The mixture was stirred for 2 hours at 60 °C. LCMS showed the reaction was completed. The mixture was quenched with water (10 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine (10 mL x 5), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 3-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-4-nitro-1-tetrahydropyran-2-yl-pyrazole 3-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-4-nitro-1-tetrahydropyran-2-yl-pyrazole (**7**) (50.0% purity, 0.20 g, yield: 21.6 %) as colorless oil. LCMS (M+Na⁺) *m/z*: calcd: 567.1; found 590.2.


[1025] 3-[[4-[1-[3,5-Dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-1-tetrahydropyran-2-yl-pyrazol-4-amine (8): To a solution of 3-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-4-nitro-1-tetrahydropyran-2-yl-pyrazole (**7**) (1.0 g, 1.76 mmol) in THF/H₂O (6 mL/3 mL) was added NH₄Cl (0.282 g, 5.27 mmol) at 20 °C, Zn (0.575 g, 8.79 mmol) was added to the mixture at 0 °C. The mixture was stirred at 20 °C for 8 hours. LCMS showed the reaction was completed. The reaction mixture was quenched with water (10 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by p-HPLC (TFA) to give the 3-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-1-

tetrahydropyran-2-yl-pyrazol-4-amine (8) (60 mg, yield: 6.3 %) as colorless oil. ¹H NMR (400 MHz, CHCl₃-d) δ ppm 7.07-7.17 (m, 4 H) 6.89-6.99 (m, 2 H), 5.23 (d, J = 9.48 Hz, 1 H), 5.10 (s, 2 H), 4.24-4.30 (m, 2 H), 4.05 (d, J = 11.47 Hz, 1 H), 3.83-3.89 (m, 2 H), 3.63-3.72 (m, 1 H), 2.06 (m, 2 H), 2.02 (s, 1 H), 1.62 (s, 6 H). LCMS (M+H⁺) m/z: calcd: 537.14; found 538.2.

[1026] N-[3-[[4-[1-[3,5-Dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-1-tetrahydropyran-2-yl-pyrazol-4-yl]methanesulfonamide (9): To a solution of 3-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-1-tetrahydropyran-2-yl-pyrazol-4-amine (8) (60 mg, 0.111 mmol) in DCM (0.5 mL) was added pyridine (17.6 mg, 0.223 mmol) at 20 °C. Mesyl chloride (0.0129 mL, 0.167 mmol) was added dropwise to the reaction mixture at 0 °C. The mixture was stirred at 20 °C for 4 hours. LCMS showed the reaction was completed. The reaction mixture was quenched with water (5 mL) and extracted with DCM (1 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give N-[3-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-1-tetrahydropyran-2-yl-pyrazol-4-yl]methanesulfonamide (9) (70 mg, yield: 81.5 %) as colorless oil. HPLC purity (220 nm): 79%. LCMS (M+Na⁺) m/z: calcd: 615.1; found 616.2.

[1027] N-[3-[[4-[1-[3,5-Dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-1H-pyrazol-4-yl]methanesulfonamide (A40): A solution of N-[3-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-1-tetrahydropyran-2-yl-pyrazol-4-yl]methanesulfonamide (9) (70 mg, 0.113 mmol) in HCl/EtOAc (4M, 2 mL) was stirred at 20 °C for 2 hours. LCMS showed the reaction was completed. The mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (FA) to give N-[3-[[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-1H-pyrazol-4-yl]methanesulfonamide(A40) (11.6 mg, yield: 18.2 %) as white solid. ¹H NMR (400 MHz, CHCl₃-d) δ ppm 7.71 (s, 1 H), 7.10 - 7.17 (m, 4 H), 6.89 - 6.94 (m, 2 H), 6.22 (s, 1 H), 5.21 (s, 2 H), 4.26 (t, J = 6.39 Hz, 2 H), 3.86 (t, J = 6.28 Hz, 2 H), 2.90 (s, 3 H), 1.62 (s, 6 H). LCMS (M+Na⁺) m/z: calcd: 531.06; found 532.1.

[1028] Example 11: Synthesis of N-(4-((4-(2-(3, 5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy) methyl) pyrimidin-5-yl)methanesulfonamide (A41):



[1029] N-(4-Methylpyrimidin-5-yl)methanesulfonamide (2): To a solution of 4-methylpyrimidin-5-amine (**1**) (3.0 g, 27.5 mmol) and pyridine (10.9 g, 0.137 mol) in DCM (30 mL) was added methanesulfonyl chloride (6.3 g, 55.0 mmol) at 0 °C and the resulting mixture was stirred at 20 °C for 3 hours. TLC showed the reaction was completed. The mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give N-(4-methylpyrimidin-5-yl)methanesulfonamide (**2**) (1.0 g, 19.4%) as yellow solid. ¹H NMR (400 MHz, CHCl₃-*d*) δ ppm 8.98 (s, 1 H), 8.76 (s, 1 H), 6.68 (br s, 1 H), 3.13 (s, 3 H), 2.61 (s, 3 H).

[1030] tert-Butyl (4-methylpyrimidin-5-yl)(methylsulfonyl)carbamate (3): To a solution of N-(4-methylpyrimidin-5-yl) methanesulfonamide (**2**) (1.0 g, 5.34 mmol) and TEA (1.1 g, 10.7 mmol), DMAP (196 mg, 1.60 mmol) in THF (15 mL) was added tert-butoxycarbonyl tert-butyl carbonate (1.75 g, 8.01 mmol) at 0 °C and the resulting mixture was stirred at 50 °C for 3 hours. TLC showed the reaction was completed. The resulting mixture was cooled down and quenched with water (20 mL). The mixture was extracted with EtOAc (10 mL x 5). The combined organic layers were washed with brine (20 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give tert-butyl N-(4-methylpyrimidin-5-yl)-N-methylsulfonyl-carbamate (**3**) (1.0 g, yield: 65.2%) as yellow solid. ¹H NMR (400 MHz, CHCl₃-*d*) δ ppm 9.07 (s, 1 H), 8.46 (s, 1 H), 3.53 (s, 3 H), 2.56 (s, 3 H), 1.49 (s, 9 H).


[1031] tert-Butyl (4-(bromomethyl)pyrimidin-5-yl)(methylsulfonyl)carbamate (4): To a mixture of tert-butyl N-(4-methylpyrimidin-5-yl)-N-methylsulfonyl-carbamate (**3**) (1.0 g, 3.48 mmol) and NBS (929 mg, 5.22 mmol) in CCl₄ (20 mL) was added AIBN (495 mg, 3.48 mmol) under N₂ atmosphere, and the resulting mixture was stirred at 100 °C for 16 hours. LCMS showed the reaction was completed. The mixture was cooled down and quenched with water

(40 mL), and then extracted with DCM (10 mL x 3). The combined organic layers were washed with brine (20 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give tert-butyl N-(bromomethyl)pyrimidin-5-yl)-N-methylsulfonyl-carbamate (**4**) (170 mg, yield: 13.3%) as yellow solid. ¹H NMR (400 MHz, CHCl₃-d) δ ppm 9.20 (br s, 1 H), 8.60 (br s, 1 H), 4.59 (d, *J*=10.64 Hz, 1 H), 4.42 (d, *J*=10.64 Hz, 1 H), 3.57 - 3.60 (m, 3 H), 1.48 - 1.51 (m, 9 H). LCMS (M + H⁺) *m/z*: 367.0; found 368.0.

[1032] tert-Butyl (4-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)pyrimidin-5-yl)(methylsulfonyl)carbamate (6): To a mixture of 4-(1-(3,5-dichloro-4-(2-chloroethoxy)phenyl)-1-methyl-ethyl)phenol (**5**) (134 mg, 0.37 mmol) and Cs₂CO₃ (181 mg, 0.56 mmol) in DMF (5 mL) was added tert-butyl N-(4-(bromomethyl)pyrimidin-5-yl)-N-methylsulfonyl-carbamate (**4**) (170 mg, 0.37 mmol) at 0 °C under N₂ atmosphere, and the resulting mixture was stirred at the same temperature for 3 hours. TLC showed the reaction was completed. The mixture was poured into water (10 mL) and extracted with EtOAc (5 mL x 5). The combined organic layers were washed with brine (10 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give tert-butyl N-(4-((4-(1-(3,5-dichloro-4-(2-chloroethoxy)phenyl)-1-methyl-ethyl)phenoxy)methyl)pyrimidin-5-yl)-N-methylsulfonyl-carbamate (**6**) (50 mg, yield: 20.9%) as light oil. ¹H NMR (400 MHz, CHCl₃-d) δ ppm 9.18 (s, 1 H), 8.62 (s, 1 H), 7.12 (s, 2 H), 7.09 (s, 2 H), 6.85-6.91 (m, 2 H), 5.15-5.30 (m, 2 H), 4.24 (t, *J*=6.28 Hz, 2 H), 3.85 (t, *J*=6.28 Hz, 2 H), 3.52 (s, 3 H), 1.60 (s, 6 H), 1.37 (s, 9 H).

[1033] N-(4-((4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)pyrimidin-5-yl)methanesulfonamide (A41): A mixture of tert-butyl N-(4-((4-(1-(3,5-dichloro-4-(2-chloroethoxy)phenyl)-1-methyl-ethyl)phenoxy)methyl)pyrimidin-5-yl)-N-methylsulfonyl-carbamate (**6**) (50 mg, 0.062 mmol) in DCM (5.0 mL) and TFA (0.5 mL) was stirred at 20 °C for 1 hour. LCMS showed the reaction was completed. The resulting mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (TFA) to give N-(4-((4-(1-(3,5-dichloro-4-(2-chloroethoxy)phenyl)-1-methyl-ethyl)phenoxy)methyl)pyrimidin-5-yl) methanesulfonamide (**A41**) (8 mg, yield: 23.7%) as yellow oil. ¹H NMR (400 MHz, CHCl₃-d) δ ppm 9.01 (d, *J*=4.40 Hz, 2 H), 7.81 (br s, 1 H), 7.16 (d, *J*=8.93 Hz, 2 H), 7.10 (s, 2 H), 6.93 (d, *J*=8.93 Hz, 2 H), 5.36 (s, 2 H), 4.26 (t, *J*=6.36 Hz, 2 H), 3.86 (t, *J*=6.36 Hz, 2 H), 3.03 (s, 3 H), 1.62 (s, 6 H). LCMS (M + H⁺) *m/z*: 545.05; found 546.0. HPLC purity(220 nm): 84.4%.

[1034] Example 12: Synthesis of N-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)-2-(methylsulfonylamido)oxazole-4-carboxamide (**A49**)




[1035] Ethyl 2-(methylsulfonylamido)oxazole-4-carboxylate (2): To a solution of methanesulfonamide (3.3 g, 34.2 mmol) in DMF (30 mL) was added NaH (1.3 g, 60% purity, 34.2 mmol) in portions at 0 °C under N₂ atmosphere. The mixture was stirred at 0 °C for 0.5 hour, and then ethyl 2-chlorooxazole-4-carboxylate (**1**) (3.0 g, 17.1 mmol) was added to the mixture. The resulting mixture was stirred at 60 °C for 10 hours. LCMS showed the reaction was completed. The reaction was quenched with H₂O (10 mL), and directly purified by reversed phase MPLC (TFA) to give ethyl 2-(methanesulfonamido)oxazole-4-carboxylate (**2**) (1.17 g, yield: 29.2%) as colorless oil. ¹H NMR (400MHz, CHCl₃-d) δ 7.61 (s, 1H), 4.39 (q, *J*=7.2 Hz, 2H), 3.12 (s, 3H), 1.37 (t, *J*=7.2 Hz, 3H). LCMS (M+H)⁺ *m/z*: cld 234.03; found 235.1.

[1036] 2-(Methylsulfonylamido)oxazole-4-carboxylic acid (3): To a mixture of ethyl 2-(methanesulfonamido)oxazole-4-carboxylate (**2**) (100 mg, 0.43 mmol) in THF (3 mL) and H₂O (1 mL) was added LiOH·H₂O (72 mg, 1.7 mmol) at 25 °C. The mixture was stirred at the same temperature for 10 hours. LCMS showed the reaction was completed. The mixture was adjusted pH to 4 by aqueous HCl (2 M), and then concentrated under reduce pressure to give 2-(methanesulfonamido)oxazole-4-carboxylic acid (**3**) (90 mg, 65% purity) as yellow solid. ¹H NMR (400MHz, MeOH-d₄) δ 7.94 (s, 1H), 3.08 (s, 3H). LCMS (M-H)⁻ *m/z*: cld 206.0; found 204.9.

[1037] N-(4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)-2-(methylsulfonylamido)oxazole-4-carboxamide (A49): To a solution of 2-(methanesulfonamido)oxazole-4-carboxylic acid (**3**) (60 mg, 0.3 mmol) in DMF (3 mL) was added 4-[1-(3,5-dichloro-4-(2-chloroethoxy)phenyl)-1-methyl-ethyl]aniline (**4**) (104 mg, 0.3 mmol), HATU (133 mg, 0.35 mmol) and TEA (0.12 mL, 0.9 mmol) at 25 °C. The mixture was stirred at the same temperature for 3 hours. LCMS showed the reaction was completed, the mixture

was quenched with H₂O (1 mL), and directly purified by prep-HPLC (TFA), to give N-[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]phenyl]-2-(methanesulfonamido)oxazole-4-carboxamide (**A49**) (23.2 mg, yield: 14.6 %) as white solid. ¹H NMR (400MHz, CHCl₃-d) δ 8.44 (s, 1H), 7.91 (s, 1H), 7.56 (d, *J*=8.8 Hz, 2H), 7.21 (d, *J*=8.8 Hz, 2H), 7.13 (s, 2H), 4.27 (t, *J*=6.4 Hz, 2H), 3.86 (t, *J*=6.4 Hz, 2H), 3.32 (s, 3H), 1.65 (s, 6H). LCMS (M+H⁺) *m/z*: calcd 545.03; found 546.0.

[1038] Example 13: Synthesis of 5-((4-(2-(3,5-dichloro-4-(3,3,3-trifluoropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-4-(methylsulfonyl)oxazole (A54**)**




[1039] 2,6-Dichloro-4-(2-(4-hydroxyphenyl)propan-2-yl)phenol (3): To a solution of 2,6-dichloro-4-[1-(4-hydroxyphenyl)-1-methyl-ethyl]phenol (**1**) (0.2 g, 0.67 mmol) in DMF (2 mL) were added 1,1,1-trifluoro-3-iodo-propane (**2**) (1.5 g, 6.7 mmol) and Cs₂CO₃ (0.66 g, 2.0 mmol). The resulting reaction was stirred at 60°C for 16 hours. TLC showed the reaction was completed. The mixture was cooled down and poured into water (10 mL), extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine (10 mL x 3), dried over Na₂SO₄, filtered and concentrated. The residue was purified by p-TLC to give 4-[1-[3,5-dichloro-4-(3,3,3-trifluoropropoxy)phenyl]-1-methyl-ethyl]phenol (**3**) (80 mg, yield: 30.2%) as colorless oil. ¹H NMR (400 MHz, CHCl₃-d) δ 7.13 (s, 2H), 7.09-7.05 (m, 2H), 6.78-6.75 (m, 2H), 4.71 (s, 1H), 4.21 (t, *J*=6.8 Hz, 2H), 2.78-2.66 (m, 2H), 1.62 (s, 6H).

[1040] 5-((4-(2-(3,5-Dichloro-4-(3,3,3-trifluoropropoxy)phenyl)propan-2-yl)phenoxy)methyl)-4-(methylsulfonyl)oxazole (A54**):** To a mixture of 4-[1-[3,5-dichloro-4-(3,3,3-trifluoro propoxy)phenyl]-1-methyl-ethyl]phenol (**3**) (40 mg, 0.10 mmol) and 5-(chloromethyl)-4-methylsulfonyl-oxazole (**4**) (24 mg, 0.12 mmol) in DMF (0.5 mL) was added Cs₂CO₃ (66 mg, 0.20 mmol) and the mixture was stirred at 25 °C for 16 hours. LCMS showed the reaction was completed. The mixture was poured into water (2 mL) and extracted with EtOAc (5 mL x 3). The combined organic layers were washed with brine (5 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by p-TLC to give 5-[[4-[1-[3,5-dichloro-4-(3,3,3-trifluoropropoxy)phenyl]-1-methyl-ethyl]phenoxy]methyl]-4-methylsulfonylo-xazole (**A54**) (18 mg, yield: 29.9%) as yellow oil. LCMS purity: (220 nm): 93.3%. ¹H NMR (400 MHz, CHCl₃-d) δ 8.00 (s, 1H), 7.16-7.12 (m,

4H), 6.94 (d, $J=8.8$ Hz, 2H), 5.42 (s, 2H), 4.22 (t, $J=6.8$ Hz, 2H), 3.19 (s, 3H), 2.78-2.64 (m, 2H), 1.62 (s, 6H). LCMS (M + NH₄⁺) m/z : calcd 551.1; found 569.0.

[1041] Example 14: Synthesis of 2-(2-chloroethoxy)-5-(2-(3-cyano-4-((4-(methylsulfonyl)oxazol-5-yl)methoxy)phenyl)propan-2-yl)benzonitrile (A63)



[1042] Methyl 3-bromo-4-(2-chloroethoxy)benzoate (2): To a solution of methyl 3-bromo-4-hydroxybenzoate (**1**) (15 g, 65 mmol) in DMF (100 mL) was added 1-bromo-2-chloroethane (11 g, 78 mmol) and Cs₂CO₃ (32 g, 97 mmol) under N₂ atmosphere at 20 °C. Then the reaction mixture was stirred at the same temperature for 16 hours. TLC showed the reaction was completed. The mixture was poured into H₂O (100 mL), extracted with EtOAc (100 mL x 2). The combined organic layers were washed with brine (100 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by MPLC to give methyl 3-bromo-4-(2-chloroethoxy)benzoate (**2**) (13 g, yield: 67%) as white solid. ¹H NMR (400 MHz, CHCl₃-d) δ 8.29-8.21 (m, 1H), 8.02-7.90 (m, 1H), 6.91 (d, $J=8.6$ Hz, 1H), 4.35 (t, $J=6.1$ Hz, 2H), 3.95-3.86 (m, 5H).

[1043] Methyl 4-(2-chloroethoxy)-3-cyanobenzoate (3): To a solution of methyl 3-bromo-4-(2-chloroethoxy)benzoate (**2**) (0.6 g, 2.0 mmol) in DMF (20 mL) was added Zn(CN)₂ (0.72 g, 6.1 mmol), Xantphos (0.1 g, 0.2 mmol) and Pd₂(dba)₃ (0.18 g, 0.2 mmol) under N₂ atmosphere at 20 °C. The reaction mixture was stirred at 130 °C for 3 hours. TLC showed the reaction was completed. Thirteen batches of reaction were combined, and then poured into H₂O (50 mL). The mixture was extracted with EtOAc (80 mL x 3). The combined organic layers were washed with brine (80 mL x 4), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude product was purified by MPLC to give methyl 4-(2-chloroethoxy)-3-cyanobenzoate

(3) (4.60 g, yield: 72.2 %) as brown solid. ^1H NMR(400MHz, $\text{CHCl}_3\text{-d}$) δ 8.27 (s, 1H), 8.22 (dd, $J=2.1, 8.9$ Hz, 1H), 7.02 (d, $J=8.8$ Hz, 1H), 4.41 (s, 2H), 3.92 (s, 3H), 3.92-3.88 (m, 2H).

[1044] 2-(2-Chloroethoxy)-5-(2-hydroxypropan-2-yl)benzonitrile (4): To a solution of MeMgBr (3 M, 24 mL, 72 mmol) in THF (20 mL) was added methyl 4-(2-chloroethoxy)-3-cyanobenzoate (3) (4.3 g, 18 mmol) in THF (40 mL) dropwise under N_2 atmosphere at 0 °C. Then the reaction was stirred at 20 °C for 3 hours. LCMS showed the reaction was completed. The mixture was poured into saturated aqueous NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (20 mL x 2), dried over Na_2SO_4 , filtered, concentrated under reduced pressure. The residue was purified by MPLC to give 2-(2-chloroethoxy)-5-(2-hydroxypropan-2-yl)benzonitrile (4) (2.9 g, yield: 67.4 %) as brown oil. ^1H NMR(400MHz, $\text{CHCl}_3\text{-d}$) δ = 7.71 (d, $J=2.4$ Hz, 1H), 7.67 (dd, $J=2.3, 8.7$ Hz, 1H), 6.98 - 6.92 (m, 1H), 4.34 (t, $J=6.1$ Hz, 2H), 3.87 (t, $J=6.2$ Hz, 2H), 1.57 (s, 6H). LCMS ($\text{M} + \text{H}^+$) m/z : calcd 239.1; found: 240.1.

[1045] 2-(2-Chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (5): To a mixture of 2-(2-chloroethoxy)-5-(2-hydroxypropan-2-yl)benzonitrile (4) (2.9 g, 12.1 mmol) and phenol (1.4 g, 14.5 mmol) in CCl_4 (40 mL) was added $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (3.0 mL, 24 mmol) dropwise under N_2 atmosphere at 0 °C. Then the reaction was stirred at 20 °C for 3 hours. TLC showed the reaction was completed. The mixture was poured into saturated aqueous NaHCO_3 (40 mL), extracted with DCM (20 mL x 2). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by MPLC to give 2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (5) (2.77 g, yield: 72.5%) as brown oil. ^1H NMR(400MHz, $\text{CHCl}_3\text{-d}$) δ 7.46 (s, 1H), 7.39-7.33 (dd, $J=8.8$ Hz, 1H), 7.05 (d, $J=8.6$ Hz, 2H), 6.88-6.83 (d, $J=8.8$ Hz, 1H), 6.80-6.73 (d, $J=8.4$ Hz, 2H), 5.00 (s, 1H), 4.31 (t, $J=6.2$ Hz, 2H), 3.85 (t, $J=6.2$ Hz, 2H), 1.63 (s, 6H).

[1046] 5-(2-(3-Bromo-4-hydroxyphenyl)propan-2-yl)-2-(2-chloroethoxy)benzonitrile (6): To a solution of 2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (5) (1.35 g, 4.27 mmol) in EtOH (30 mL) was added Br_2 (0.22 mL, 4.27 mmol) dropwise under N_2 atmosphere at 0 °C. Then the reaction was stirred at 20 °C for 3 hours, and then at 40 °C for another 6 hours. LCMS showed 23% of starting material and 73% of desired product in reaction mixture. The mixture was poured into saturated aqueous Na_2SO_3 (20 mL) then concentrated under reduced to remove EtOH. The resulting solution was extracted with EtOAc (20 mL x 2). The combined organic layers were washed with brine (20 mL), dried over Na_2SO_4 , filtered, and concentrated under reduced. The residue was purified by MPLC to give 5-(2-(3-bromo-4-hydroxyphenyl)propan-2-yl)-2-(2-chloroethoxy)benzonitrile (6) (1.16

g, yield: 68.7 %) as brown oil. ^1H NMR(400MHz, $\text{CHCl}_3\text{-d}$) δ = 7.45 (d, J =2.4 Hz, 1H), 7.37-7.31 (d, J =8.8 Hz, 1H), 7.30-7.28 (m, 1H), 7.03-6.98 (m, 1H), 6.96-6.92 (m, 1H), 6.87 (d, J =8.8 Hz, 1H), 5.44 (br s, 1H), 4.32 (t, J =6.2 Hz, 2H), 3.86 (t, J =6.2 Hz, 2H), 1.63 (s, 6H). LCMS ($\text{M} + \text{H}^+$) m/z : calcd 393.0; found: 393.8.

[1047] 2-(2-Chloroethoxy)-5-(2-(3-cyano-4-hydroxyphenyl)propan-2-yl)benzonitrile (7):


To a solution of 5-(2-(3-bromo-4-hydroxyphenyl)propan-2-yl)-2-(2-chloroethoxy) benzonitrile (**6**) (0.60 g, 1.52 mmol) in NMP (5 mL) was added CuCN (0.68 g, 7.6 mol) under N_2 atmosphere at 20°C . The reaction was stirred at 180°C for 0.5 hour. TLC showed the reaction was completed. The mixture was poured into H_2O (20 mL), extracted with EtOAc (10 mL x2). The combined organic layers were washed with brine (10 mL x 6), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by MPLC to give 2-(2-chloroethoxy)-5-(2-(3-cyano-4-hydroxyphenyl)propan-2-yl)benzonitrile (**7**) (130 mg, yield: 25.1%) as brown oil. ^1H NMR(400MHz, $\text{CHCl}_3\text{-d}$) δ = 7.44-7.41 (m, 1H), 7.35 (d, J =2.4 Hz, 1H), 7.32 (dd, J =2.5, 8.9 Hz, 1H), 7.23 (dd, J =2.4, 8.8 Hz, 1H), 6.91 (d, J =7.3 Hz, 1H), 6.90 - 6.88 (dd, J =2.5, 8.9 Hz, 1H), 6.10 (br, 1H), 4.37-4.28 (t, J =6.2 Hz, 2H), 3.87 (t, J =6.2 Hz, 2H), 1.64 (s, 6H).

[1048] 2-(2-Chloroethoxy)-5-(2-(3-cyano-4-((4-(methylsulfonyl)oxazol-5-yl)methoxy)phenyl)propan-2-yl) benzonitrile (A63):

To a solution of 2-(2-chloroethoxy)-5-(2-(3-cyano-4-hydroxyphenyl)propan-2-yl)benzonitrile (**7**) (130 mg, 0.38 mmol) in DMF (2 mL) was added 5-(chloromethyl)-4-(methylsulfonyl)oxazole (**G**) (75 mg, 0.38 mol) and Cs_2CO_3 (249 mg, 0.76 mmol) under N_2 atmosphere. The reaction was stirred at 0°C for 3 hours. LCMS showed the reaction was completed. The mixture was diluted with EtOAc (5 mL) and poured into H_2O (5 mL). The aqueous phase was extracted with EtOAc (5 mL x2). The combined organic layers were washed with brine (5 mL x 4), then dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by prep-HPLC(TFA) to give 2-(2-chloroethoxy)-5-(2-(3-cyano-4-((4-(methylsulfonyl)oxazol-5-yl)methoxy)phenyl)propan-2-yl)benzonitrile (**A63**) (53 mg, yield: 27.8 %) as white solid. LCMS purity (220 nm): 91.1%. ^1H NMR(400MHz, $\text{CHCl}_3\text{-d}$) δ = 8.04 (s, 1H), 7.41 (br s, 2H), 7.39 - 7.29 (m, 2H), 7.15 - 7.04 (m, 1H), 6.94 - 6.85 (d, J = 8.9 Hz, 1H), 5.51 (s, 2H), 4.33 (br t, J =6.0 Hz, 2H), 3.87 (br t, J =6.0 Hz, 2H), 3.25 (s, 3H), 1.64 (s, 6H). LCMS ($\text{M} + \text{H}^+$) m/z : calcd 499.1; found 500.1.

[1049] Example 15: Synthesis of N-((2-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)amino)oxazol-5-yl)methyl)methanesulfonamide

(A75)



[1050] Ethyl 2-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)amino)oxazole-5-carboxylate (3): To a solution of 4-[1-[3,5-dichloro-4-(2-chloroethoxy)-phenyl]-1-methyl-ethyl]aniline (**1**) (1.00 g, 2.5 mmol) in DMF (10 mL) was added Ag_2CO_3 (1.05 g, 3.81 mol) and ethyl 2-chlorooxazole-5-carboxylate (**2**) (0.668 g, 3.81 mmol) under N_2 atmosphere at 20 °C. Then the mixture was stirred at 70 °C for 16 hours. TLC showed the reaction was completed. The mixture was cooled down, poured into H_2O (50 mL) and extracted with EtOAc (25 mL x 2). The organic layers were combined and washed with brine (20 mL x 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give ethyl 2-[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]anilino]oxazole-5-carboxylate (**3**) (0.7 g, yield: 56.2%) as brown oil. ^1H NMR (400MHz, $\text{CHCl}_3\text{-d}$) δ 7.62 (s, 1H), 7.46 (s, 1H), 7.43 (d, $J=8.8$ Hz, 2H), 7.20 (d, $J=8.8$ Hz, 2H), 7.14 (s, 2H), 4.37 (q, $J=7.2$ Hz, 2H), 4.26 (t, $J=6.4$ Hz, 2H), 3.86 (t, $J=6.4$ Hz, 2H), 1.64 (s, 6H), 1.38 (t, $J=7.2$ Hz, 3H).


[1051] (2-((4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)amino)oxazol-5-yl)methanol (4): To a solution of ethyl 2-[4-[1-[3,5-dichloro-4-(2-chloroethoxy) phenyl]-1-methyl-ethyl]anilino]oxazole-5-carboxylate (**3**) (0.6 g, 1.2 mmol) in DCM (10 mL) was added DIBAL-H (1 M in toluene, 2.4 mL, 2.4 mmol) under N_2 atmosphere at 0 °C. Then the resulting mixture was stirred at 20 °C for 3 hours. TLC showed the reaction was completed. The mixture was poured into saturated aqueous NH_4Cl (20 mL) and extracted with DCM (10 mL x 3). The organic layers were washed with brine (15 mL x 2), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give 2-[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]anilino]oxazol-5-yl]-methanol (**4**) (410 mg, yield: 74.6%) as brown oil. ^1H NMR (400MHz, $\text{CHCl}_3\text{-d}$) δ 7.38 (d, $J=8.8$ Hz, 2H), 7.17 (d, $J=8.8$

Hz, 2H), 7.14 (s, 2H), 6.81 (s, 1H), 4.60 (s, 2H), 4.26 (t, $J=6.4$ Hz, 2H), 3.86 (t, $J=6.4$ Hz, 2H), 1.63 (s, 6H).

[1052] tert-Butyl ((2-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)-amino)oxazol-5-yl)methyl)(methylsulfonyl)carbamate (5): To a mixture of [2-[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]anilino]oxazol-5-yl]methanol (**4**) (50 mg, 0.1 mmol) and tert-butyl N-methylsulfonylcarbamate (21 mg, 0.1 mmol) in THF (3 mL) was added PPh_3 (58 mg, 0.22 mmol) and DEAD (29 mg, 0.16 mmol). The mixture was stirred at 25 °C for 10 hours under N_2 atmosphere. LCMS showed 30% desired product was detected. The mixture was concentrated and purified by prep-TLC to give tert-butyl N-[[2-[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]anilino]oxazol-5-yl]methyl]-N-methylsulfonyl-carbamate (**5**) (25 mg, yield: 36.0%) as colorless oil. ^1H NMR (400MHz, $\text{CHCl}_3\text{-d}$) δ 7.39 (d, $J=8.8$ Hz, 2H), 7.16 (d, $J=8.8$ Hz, 2H), 7.13 (s, 2H), 6.84 (s, 1H), 4.86 (s, 2H), 4.26 (t, $J=6.4$ Hz, 2H), 3.86 (t, $J=6.4$ Hz, 2H), 3.24 (s, 3H), 1.63 (s, 9H), 1.53 (s, 6H).

[1053] N-((2-((4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)amino)oxazol-5-yl)methyl)methanesulfonamide (A75): A solution of tert-butyl N-[[2-[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]anilino]oxazol-5-yl]methyl]-N-methylsulfonyl-carbamate (**5**) (25 mg, 0.04 mmol) in DCM (2 mL) and TFA (0.2 mL) was stirred at 25 °C for 3 hours. LCMS showed the reaction was completed. The mixture was concentrated and purified by prep-HPLC (TFA) to give N-[[2-[4-[1-[3,5-dichloro-4-(2-chloroethoxy)phenyl]-1-methyl-ethyl]anilino]oxazol-5-yl]methyl]methanesulfonamide (**4.6** mg, yield: 21.9%) as yellow oil. LCMS purity (220 nm): 86%. ^1H NMR (400MHz, $\text{CHCl}_3\text{-d}$) δ 7.36 - 7.33 (m, 2H), 7.25 - 7.22 (m, 2H), 7.11 (s, 2H), 7.02 (s, 1H), 5.18 (s, 1H), 4.36 (s, 2H), 4.27 (t, $J=6.4$ Hz, 2H), 3.87 (t, $J=6.4$ Hz, 2H), 3.00 (s, 3H), 1.64 (s, 6H). LCMS ($\text{M} + \text{H}^+$) m/z : calcd: 531.0; found 531.6.

[1054] Example 16: Synthesis of 3-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzyl)-1,5,5-trimethylimidazolidine-2,4-dione (B2**)**



[1055] Methyl 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzoate (2):

To a solution of 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl trifluoromethanesulfonate (1) (600 mg, 1.7 mmol) and TEA (1 mL, 5.0 mmol) in MeOH (15 mL) was added Pd(dppf)Cl₂ (200 mg, 0.3 mmol). The resulting mixture was stirred at 50 Psi of CO atmosphere and the mixture at 60 °C for 16 hours. TLC showed the reaction was completed. The mixture was poured into H₂O (30 mL), extracted with EtOAc (10 mL x 2). The combined organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-TLC to give methyl 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzoate (2) (300 mg, yield: 44.5%) as yellow oil. ¹H NMR (400 MHz, CHCl₃-d) δ 8.00-7.90 (m, 2H), 7.27-7.23 (m, 2H), 7.09 (s, 2H), 4.28-4.21 (m, 2H), 3.90 (s, 3H), 3.87-3.81 (m, 2H), 1.68-1.62 (m, 6H).

[1056] (4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)methanol (3):


To a solution of methyl 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzoate (2) (300 mg, 0.7 mmol) in THF (3 mL) was added DIBAL-H (1.00 M, 2 mL) at 0 °C and the mixture was stirred at 0-25 °C for 2 hours. TLC showed the reaction was completed. The mixture was poured into H₂O (15 mL), extracted with EtOAc (5 mL x 2). The combined organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give (4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)methanol (3) (170 mg, yield: 60.9%) as yellow oil. ¹H NMR (400 MHz, CHCl₃-d) δ = 7.32 (d, *J* = 8.3 Hz, 2H), 7.24-7.18 (m, 2H), 7.17-7.11 (m, 2H), 4.75-4.63 (m, 2H), 4.31-4.22 (m, 2H), 3.86 (t, *J* = 6.4 Hz, 2H), 1.70-1.61 (m, 6H).

[1057] 1,3-Dichloro-2-(2-chloroethoxy)-5-(2-(4-(chloromethyl)phenyl)propan-2-yl)benzene (4): To a solution of (4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)methanol (3) (40 mg, 0.1 mmol) in DCM (2 mL) was added SOCl₂ (0.1 mL, 1.1 mmol) at 0 °C and the mixture was stirred at 0-25 °C for 2 hours. TLC showed the reaction was completed. The mixture was poured into H₂O (10 mL), extracted with DCM (3 mL x 2). The combined organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 1,3-dichloro-2-(2-chloroethoxy)-5-(2-(4-(chloromethyl)phenyl)propan-2-yl)benzene (4) (40 mg, yield: 95.3%) as yellow oil. ¹H NMR (400 MHz, CHCl₃-d) δ = 7.36-7.30 (m, 2H), 7.20 (d, *J* = 8.4 Hz, 2H), 7.15-7.15 (m, 1H), 7.14 (s, 1H), 4.59 (s, 2H), 4.27 (t, *J* = 6.4 Hz, 2H), 3.86 (t, *J* = 6.4 Hz, 2H), 1.68-1.62 (m, 6H).

[1058] 3-(4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzyl)-1,5,5-trimethylimida zolidine-2,4-dione (B2): To a mixture of 1,5,5-trimethylimidazolidine-2,4-dione (5) (20 mg, 0.2 mmol) and K₂CO₃ (70 mg, 0.5 mmol) in DMF (3 mL) was added 1,3-

dichloro-2-(2-chloroethoxy)-5-(2-(4-(chloromethyl)phenyl)propan-2-yl)benzene (**4**) (50 mg, 0.1 mmol) at 25 °C and the mixture was stirred at the same temperature for 2 hours. LCMS showed the reaction was completed. The mixture was poured into H₂O (10 mL), extracted with EtOAc (5 mL x 2). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (HCl) to give 3-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzyl)-1,5,5-trimethylimidazolidine-2,4-dione (**B2**) (20 mg, yield: 31.8 %) as colorless oil. LCMS purity (220 nm): 96.1%. ¹H NMR (400 MHz, CHCl₃-d) δ = 7.30 - 7.25 (m, 1H), 7.28 - 7.25 (m, 1H), 7.30 - 7.25 (m, 1H), 7.15 - 7.10 (m, 2H), 7.10 - 7.08 (m, 2H), 4.66 - 4.57 (m, 2H), 4.24 (t, *J* = 6.4 Hz, 2H), 3.89 - 3.77 (m, 2H), 2.87 (s, 3H), 1.65 - 1.54 (m, 6H), 1.41 - 1.34 (m, 6H). LCMS (M + H⁺) *m/z*: calcd: 496.1; found 497.1.

[1059] Example 17: Synthesis of 3-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzyl)-5,5-dimethyl-1-(methylsulfonyl)imidazolidine-2,4-dione (B3**)**



[1060] Methyl 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzoate (2**):**

To a solution of 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl trifluoromethanesulfonate (**1**) (600 mg, 1.7 mmol) and TEA (1 mL, 5.0 mmol) in MeOH (15 mL) was added Pd(dppf)Cl₂ (200 mg, 0.3 mmol). The resulting mixture was stirred at 50 Psi of CO atmosphere and the mixture at 60 °C for 16 hours. TLC showed the reaction was completed. The mixture was poured into H₂O (30 mL), extracted with EtOAc (10 mL x 2). The combined organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-TLC to give methyl 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzoate (**2**) (300 mg, yield: 44.5%) as yellow oil. ¹H NMR (400 MHz, CHCl₃-d) δ 8.00-7.90 (m, 2H), 7.27-7.23 (m, 2H), 7.09 (s, 2H), 4.28-4.21 (m, 2H), 3.90 (s, 3H), 3.87-3.81 (m, 2H), 1.68-1.62 (m, 6H).

[1061] (4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)methanol (3**):**

To a solution of methyl 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzoate (**2**)

(300 mg, 0.7 mmol) in THF (3 mL) was added DIBAL-H (1.00 M, 2 mL) at 0 °C and the mixture was stirred at 0-25 °C for 2 hours. TLC showed the reaction was completed. The mixture was poured into H₂O (15 mL), extracted with EtOAc (5 mL x 2). The combined organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give (4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)methanol (**3**) (170 mg, yield: 60.9%) as yellow oil. ¹H NMR (400 MHz, CHCl₃-d) δ = 7.32 (d, *J* = 8.3 Hz, 2H), 7.24-7.18 (m, 2H), 7.17-7.11 (m, 2H), 4.75-4.63 (m, 2H), 4.31-4.22 (m, 2H), 3.86 (t, *J* = 6.4 Hz, 2H), 1.70-1.61 (m, 6H).


[1062] 1,3-Dichloro-2-(2-chloroethoxy)-5-(2-(4-(chloromethyl)phenyl)propan-2-yl)benzene (4): To a solution of (4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenyl)methanol (**3**) (40 mg, 0.1 mmol) in DCM (2 mL) was added SOCl₂ (0.1 mL, 1.1 mmol) at 0 °C and the mixture was stirred at 0-25 °C for 2 hours. TLC showed the reaction was completed. The mixture was poured into H₂O (10 mL), extracted with DCM (3 mL x 2). The combined organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 1,3-dichloro-2-(2-chloroethoxy)-5-(2-(4-(chloromethyl)phenyl)propan-2-yl)benzene (**4**) (40 mg, yield: 95.3%) as yellow oil. ¹H NMR (400 MHz, CHCl₃-d) δ = 7.36-7.30 (m, 2H), 7.20 (d, *J* = 8.4 Hz, 2H), 7.15-7.15 (m, 1H), 7.14 (s, 1H), 4.59 (s, 2H), 4.27 (t, *J* = 6.4 Hz, 2H), 3.86 (t, *J* = 6.4 Hz, 2H), 1.68-1.62 (m, 6H).

[1063] 3-(4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzyl)-5,5-dimethylimidazolidine-2,4-dione (6): To a mixture of 1,3-dichloro-2-(2-chloroethoxy)-5-(2-(4-(chloromethyl)phenyl)propan-2-yl)benzene (**4**) (100 mg, 0.2 mmol) and 5,5-dimethylimidazolidine-2,4-dione (**5**) (30 mg, 0.2 mmol) in DMF (5 mL) was added K₂CO₃ (100 mg, 0.8 mmol) at 25 °C and the mixture was stirred at 25 °C for 16 hours. LCMS showed the reaction was completed. The mixture was poured into H₂O (15 mL), extracted with EtOAc (5 mL x 2). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 3-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzyl)-5,5-dimethylimidazolidine-2,4-dione (**6**) (40 mg, yield: 64.8 %) as yellow oil. LCMS purity (220 nm): 97.7%. ¹H NMR (400MHz, CHCl₃-d) δ = 7.16 - 7.08 (m, 4H), 6.83 (d, *J*=8.8 Hz, 2H), 5.36 (s, 1H), 5.27 (s, 1H), 4.51 (s, 2H), 4.45 (s, 2H), 4.17 - 4.12 (m, 2H), 3.86 (t, *J*=6.4 Hz, 2H), 3.31 (s, 3H), 2.28 (quin, *J*=6.1 Hz, 2H), 1.62 (s, 6H), 1.49 (s, 9H). LCMS (M + H⁺) m/z: calcd: 482.1; found 483.2.

[1064] 3-(4-(2-(3,5-Dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzyl)-5,5-dimethyl-1-(methylsulfonyl)imidazolidine-2,4-dione (B3): To a solution of 3-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzyl)-5,5-dimethylimidazolidine-2,4-

dione (**6**) ((40 mg, 0.1 mmol) in THF (2 mL) was added Mesyl chloride (0.1 mL, 0.2 mmol) and NaH (60.0 %, 6 mg, 0.2 mmol) at 0 °C and the mixture was stirred at 80 °C for 16 hours. TLC showed the reaction was completed. The reaction was quenched with saturated aqueous NH₄Cl (10 mL) and extracted with EtOAc (3 mL x 2). The combined organic layers were washed with brine (3 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (HCl) to give 3-(4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)benzyl)-5,5-dimethyl-1-(methylsulfonyl)imidazolidine-2,4-dione (5 mg, yield: 10.8 %) as yellow oil. LCMS purity (220 nm): 81.8%. ¹H NMR (400 MHz, CHCl₃-d) δ = 7.31 - 7.28 (m, 2H), 7.20 - 7.14 (m, 2H), 7.14 - 7.11 (m, 2H), 4.71 - 4.65 (m, 2H), 4.27 (t, *J* = 6.4 Hz, 2H), 3.86 (t, *J* = 6.4 Hz, 2H), 3.38 (s, 3H), 1.76 - 1.71 (m, 6H), 1.64 (s, 6H). LCMS (M + H⁺) *m/z*: calcd:526, found: 527.


[1065] Example 18: Synthesis of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((4-(methylsulfonyl)oxazol-5-yl) methoxy)phenyl)propan-2-yl)benzonitrile (**A28**)



[1066] To a mixture of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (900 mg, 2.6 mmol) in DMF (10 mL) was added Cs₂CO₃ (167 mg, 5.1 mmol) and 5-(chloromethyl)-4-(methylsulfonyl)oxazole (750 mg, 3.9 mmol) under N₂ atmosphere at 20 °C. The mixture was stirred at 40 °C for 1 h. TLC showed the reaction was completed. The mixture was cooled down to room temperature, poured into H₂O (40 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (20 mL x 4), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by MPLC and then triturated with MTBE: EtOAc = 10 mL: 2 mL to give 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((4-(methylsulfonyl)oxazol-5-yl)ethoxy)phenyl)propan-2-yl)benzonitrile (310 mg, yield: 23.2 %) as white solid. ¹H NMR (400 MHz, CDCl₃) δ = 8.00 (s, 1H), 7.43 (d, *J* = 2.4 Hz, 1H), 7.32 (d, *J* = 2.4 Hz, 1H), 7.13 (d, *J* = 8.4 Hz, 2H), 6.96 (d, *J* = 8.8 Hz, 2H), 5.42 (s, 2H), 4.43 (t, *J* = 6.4 Hz, 2H), 3.88 (t, *J* = 6.4 Hz, 2H), 3.21 (s, 3H), 1.64 (s, 6H). LCMS (220 nm): 97.8%. Exact Mass: 508.1; found 509.1, 511.1.

[1067] Example 19: Synthesis of 2-(2-chloroethoxy)-3-methyl-5-(2-(4-((4-

(methylsulfonyl)oxazol-5-yl)methoxy) phenyl)proan-2-yl)benzonitrile (A29)



[1068] Methyl 4-hydroxy-3-iodo-5-methylbenzoate (2): To a solution of methyl 4-hydroxy-3-methyl-benzoate (3.0 g, 18.0 mmol) in DCM (30 mL) was added NIS (4.0 g, 18.0 mmol) at 0 °C under N₂ atmosphere. The mixture was stirred at 20 °C for 16 hrs. TLC showed the reaction was completed. The mixture was poured into water (20 mL), extracted with DCM (30 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography to give methyl 4-hydroxy-3-iodo-5-methyl-benzoate (3.6 g, yield: 68.3 %) as yellow solid. ¹H NMR (400 MHz, CDCl₃) δ = 8.21 (d, *J*=1.2 Hz, 1 H), 7.80 (s, 1 H), 5.70 (s, 1 H), 3.89 (s, 3 H), 2.34 (s, 3 H).

[1069] Methyl 4-(2-chloroethoxy)-3-iodo-5-methylbenzoate (4): To a mixture of methyl 4-hydroxy-3-iodo-5-methyl-benzoate (3.6 g, 12.3 mmol) and 1-bromo-2-chloro-ethane (1.9 g, 13.6 mmol) in DMF (80 mL) was added Cs₂CO₃ (8.0 g, 24.7 mmol) and the mixture was stirred at 70 °C for 16 hrs. TLC showed the reaction was completed. The mixture was cooled down and poured into water (80 mL), extracted with EtOAc (50 mL x 3). The combined organic layers were washed with brine (50 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography to give methyl 4-(2-chloroethoxy)-3-iodo-5-methyl-benzoate (2.3 g, yield: 52.6 %) as colorless oil. ¹H NMR (400 MHz, CDCl₃) δ = 8.31 (d, *J*=1.6 Hz, 1 H), 7.87 - 7.86 (m, 1 H), 4.17 (t, *J*=6.0 Hz, 2 H), 3.89 - 3.95 (m, 5 H), 2.41 (s, 3 H).

[1070] Methyl 4-(2-chloroethoxy)-3-cyano-5-methylbenzoate (5): To a solution of methyl 4-(2-chloroethoxy)-3-iodo-5-methyl-benzoate (2.1 g, 6.0 mmol) in NMP (20 mL) was added CuCN (637 mg, 7.1 mmol) under N₂ atmosphere and the mixture was stirred at 160 °C for 2 hrs. LCMS showed the reaction was completed. The mixture was cooled down and poured into water (60 mL) and extracted with EtOAc (30 mL x 3). The combined organic layers were washed with brine (30 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography to give methyl 4-(2-

chloroethoxy)-3-cyano-5-methyl-benzoate (0.7 g, yield: 46.6 %) as yellow solid. ¹H NMR (400 MHz, CHCl₃-*d*): δ 8.13 (d, *J*=2.0 Hz, 1 H), 8.09 (d, *J*=1.2 Hz, 1 H), 4.54 (t, *J*=5.6 Hz, 2 H), 3.93 (s, 3 H), 3.89 (t, *J*=5.6 Hz, 2 H), 2.39 (s, 3 H). LCMS (220 nm): 78.1%. Exact Mass: 253.1; found: 254.1.


[1071] 2-(2-chloroethoxy)-5-(2-hydroxypropan-2-yl)-3-methylbenzonitrile (6): To a mixture of methyl 4-(2-chloroethoxy)-3-cyano-5-methyl-benzoate (500 mg, 2.1 mmol) and TEA (1.7 mL, 11.8 mmol) in THF (10 mL) was added MeMgBr (3M, 9.8 mmol, 3.3 mL) dropwise at 0 °C under N₂ atmosphere and the mixture was stirred at 0 °C for 2 hrs. TLC showed the reaction was completed. The mixture was poured into saturated aqueous NH₄Cl (20 mL), extracted with EtOAc (10 mL x 4), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography to give 2-(2-chloroethoxy)-5-(2-hydroxypropan-2-yl)-3-methylbenzonitrile (250 mg, yield: 50.0 %) as yellow oil. ¹H NMR (400 MHz, CDCl₃) δ = 7.55 (s, 2 H), 4.38 (t, *J*=5.6 Hz, 2 H), 3.87 (t, *J*=5.6 Hz, 2 H), 2.36 (s, 3 H), 1.57 (s, 6 H).

[1072] 2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)-3-methylbenzonitrile (7): To a solution of 2-(2-chloroethoxy)-5-(2-hydroxy-2-methyl-ethyl)-3-methyl-benzonitrile (250 mg, 1.0 mmol) and phenol (110 mg, 1.2 mmol) in DCM (3 mL) was added BF₃·Et₂O (0.5 mL, 2 mmol) at -78 °C under N₂ atmosphere and the mixture was stirred at the same temperature for 2 hrs. TLC showed the reaction was completed. The mixture was poured into water (5 mL) and extracted with DCM (5 mL x 3). The combined organic layers were washed with brine (15 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography to give 2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)-3-methylbenzonitrile (220 mg, yield: 67.7 %) as colorless oil. ¹H NMR (400 MHz, CDCl₃) δ = 7.29 (d, *J*= 2.4 Hz, 1H), 7.21 (d, *J*= 2.0 Hz, 1H), 7.06 (d, *J*= 6.4 Hz, 2H), 6.76 (d, *J*= 6.8 Hz, 2H), 4.73 (s, 1H), 4.36 (t, *J*=6.0 Hz, 2 H), 3.86 (t, *J*=6.0 Hz, 2 H), 2.29 (s, 3H), 1.63 (s, 6H).

[1073] 2-(2-chloroethoxy)-3-methyl-5-(2-(4-((4-(methylsulfonyl)oxazol-5-yl)methoxy)phenyl)propan-2-yl)benzonitrile (A29): To a mixture of 2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)-3-methylbenzonitrile (100 mg, 0.3 mmol) and 5-(chloromethyl)-4-methylsulfonyl-oxazole (60 mg, 0.3 mmol) in DMF (3 mL) was added Cs₂CO₃ (200 mg, 0.6 mmol) at 20 °C and the mixture was stirred at 40 °C for 4 hrs. LCMS showed the reaction was completed. The mixture was cooled down, filtered and purified by prep-HPLC (TFA) to give 2-(2-chloroethoxy)-3-methyl-5-(2-(4-((4-(methylsulfonyl)oxazol-5-yl)methoxy)phenyl)propan-2-yl)benzonitrile (26 mg, yield: 17.5%) as yellow oil. ¹H NMR

(400 MHz, MeOD): δ 8.39 (s, 1 H), 7.33 - 7.35 (m, 1 H), 7.31 - 7.32 (m, 1 H), 7.16 (d, $J=8.8$ Hz, 2 H), 6.96 (d, $J=8.8$ Hz, 2 H), 5.41 (s, 2 H), 4.32 (t, $J=5.6$ Hz, 2 H), 3.88 (t, $J=5.4$ Hz, 2 H), 3.18 (s, 3 H), 2.28 (s, 3 H), 1.64 (s, 6 H). LCMS: (220 nm): 98.8 %. Exact Mass: 488.1; found 489.0.

[1074] Example 20: Synthesis of N-(4-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)pyrimidin-2-yl)methanesulfonamide (**A74**)



[1075] 4-(bromomethyl)-2-chloropyrimidine (2): To a mixture of 2-chloro-4-methylpyrimidine (5.0 g, 38.9 mmol) and NBS (10.4 g, 58.3 mmol) in CCl_4 (100 mL) was added AIBN (2.18 g, 19.4 mmol) at 20 °C and the mixture was stirred at 100 °C for 16 hrs. LCMS showed most of the starting material consumed. The mixture was cooled down, added to water (200 mL) and extracted with DCM (100 mL x 5). The combined organic phases were washed with brine (200 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by MPLC to give 4-(bromomethyl)-2-chloropyrimidine (2.0 g, yield: 24.8%) as brown oil. ^1H NMR (400MHz, CDCl_3) δ = 8.65 (d, $J=5.2$ Hz, 1H), 7.45 (d, $J=5.2$ Hz, 1H), 4.43 (s, 2H).


[1076] 2-chloro-4-((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl) pyrimidine (4): To a mixture of 4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenol (0.2 g, 0.6 mmol) and 4-(bromomethyl)-2-chloropyrimidine (127 mg, 0.6 mmol) in MeCN (5 mL) was added K_2CO_3 (115 mg, 0.83 mmol) and the mixture was stirred at 25 °C for 16 hrs. LCMS showed the reaction was completed. The resulting mixture was quenched with saturated aqueous NH_4Cl (5 mL), extracted with EtOAc (5 mL x 3). The combined organic layers were washed with brine (10 mL x 2), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give 2-chloro-4-

((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)pyrimidine (270 mg, yield: 82.9 %) as yellow solid. ^1H NMR (400MHz, CDCl_3) δ = 8.66 (d, J =5.2 Hz, 1H), 7.56 (d, J =5.2 Hz, 1H), 7.16 (d, J =8.8 Hz, 2H), 7.13 (s, 2H), 6.88 (d, J =8.8 Hz, 2H), 5.15 (s, 2H), 4.26 (t, J =6.4 Hz, 2H), 3.86 (t, J =6.4 Hz, 2H), 1.63 (s, 6H).

[1077] N-(4-(((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)pyrimidin-2-yl)methanesulfonamide (A74):

To a mixture of 2-chloro-4-(((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)pyrimidine (200 mg, 0.3 mmol), methanesulfonamide (62.6 mg, 0.66 mmol), Cs_2CO_3 (322 mg, 0.99 mmol) in anhydrous dioxane (8 mL) was added $\text{Pd}(\text{dba})_3$ (30.1 mg, 0.03 mmol) and Xantphos (38.1 mg, 0.06 mmol) and the mixture was stirred at 100 °C for 16 hrs under N_2 atmosphere. LCMS showed the reaction was completed. The mixture was cooled down, poured into water (10 mL), extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by p-HPLC(TFA) to obtain the desired product N-(4-(((4-(2-(3,5-dichloro-4-(2-chloroethoxy)phenyl)propan-2-yl)phenoxy)methyl)pyrimidin-2-yl)methanesulfonamide (61.4 mg, yield: 34.2 %) as light red solid. ^1H NMR (400MHz, CDCl_3) δ = 8.61 (d, J =5.2 Hz, 1H), 7.29 (d, J =5.2 Hz, 1H), 7.16 - 7.11 (m, 4H), 6.88 (d, J =8.8 Hz, 2H), 5.09 (s, 2H), 4.26 (t, J =6.4 Hz, 2H), 3.86 (t, J =6.4 Hz, 2H), 3.47 (s, 3H), 1.63 (s, 6H). LCMS: (220 nm): 98.1%. Exact Mass: 543.1; found 544.

[1078] Example 21: Synthesis of 3-chloro-2-(2-(chloroethoxy)-5-(2-(4-((5-(methylsulfonyl)thiazol-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (A93)



[1079] 2-(((tert-butyl)diphenylsilyl)oxy)methyl)thiazole (2): To a mixture of thiazol-2-ylmethanol (1.0 g, 8.7 mmol) and imidazole (1.2 g, 17.4 mmol) in DCM (50 mL) was added TBDPSCI (2.9 g, 10.4 mmol) in portions at 25 °C. The mixture was stirred at the same

temperature for 2 hrs. TLC showed the reaction was completed. The reaction was quenched with water (80 mL) and extracted with DCM (50 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduce pressure. The residue was purified by MPLC to give 2-(((tert-butyl)diphenylsilyl)oxy)methylthiazole (3.0 g, yield: 97.7 %) as colorless oil. ¹H NMR (400MHz, CDCl₃) δ = 7.73 - 7.70 (m, 4H), 7.48 - 7.45 (m, 1H), 7.44 - 7.38 (m, 6H), 7.31 (d, *J*=3.2 Hz, 1H), 5.02 (s, 2H), 1.14 (s, 9H).

[1080] 2-(((tert-butyl)diphenylsilyl)oxy)methyl-5-(methylthio)thiazole (3): To a solution of 2-(((tert-butyl)diphenylsilyl)oxy)methylthiazole (4.2 g, 11.9 mmol) in THF (80 mL) was added n-BuLi (2.50 M, 9.50 mL) dropwise at -78 °C under N₂ atmosphere. The mixture was stirred at the same temperature for 30 min. Then 1,2-dimethyldisulfane (2.2 g, 23.8 mmol) was added to the mixture dropwise at -78 °C under N₂ atmosphere and stirred for 30 min. TLC showed the reaction was completed. The reaction mixture was poured into saturated aqueous NH₄Cl (100 mL) and extracted with EtOAc (50 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduce pressure to give 2-(((tert-butyl)diphenylsilyl)oxy)methyl-5-(methylthio)thiazole (4.6 g, yield: 96.9 %) as brown oil. ¹H NMR (400MHz, CDCl₃) δ = 7.74 - 7.67 (m, 4H), 7.60 (s, 1H), 7.48 - 7.38 (m, 6H), 4.92 (s, 2H), 2.50 (s, 3H), 1.13 (s, 9H).

[1081] (5-(methylthio)thiazol-2-yl)methanol (4): To a solution of 2-(((tert-butyl)diphenylsilyl)oxy)methyl-5-(methylthio)thiazole (5.4 g, 13.5 mmol) in anhydrous THF (27 mL) was added TBAF (1.00 M, 27.0 mL) dropwise at 20 °C and stirred at the same temperature for 1 hour. TLC showed the reaction was completed. The reaction was quenched with water (50 mL). The aqueous layer was extracted with EtOAc (30 mL x 3). The combined organic layers were washed with brine (30 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduce pressure. The residue was purified by MPLC to give (5-(methylthio)thiazol-2-yl)methanol (1.4 g, yield: 64.3 %) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ = 7.63 (s, 1H), 4.91 (d, *J*=5.6 Hz, 2H), 2.72 (t, *J*=6.0 Hz, 1H), 2.49 (s, 3H).

[1082] 2-(chloromethyl)-5-(methylthio)thiazole (5): To a solution of (5-(methylthio)thiazol-2-yl)methanol (0.4 g, 2.5 mmol) in DCM (5 mL) was added SOCl₂ (0.5 mL, 7.4 mmol) at 20 °C, then DMF (0.1 mL) was added. The mixture was stirred at the same temperature for 1 h. TLC showed the reaction was completed. The reaction was concentrated under reduce pressure. The residue was quenched with water (10 mL) and extracted with DCM (10 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduce pressure to give 2-(chloromethyl)-5-

(methylthio)thiazole (0.4 g, yield: 89.7 %) as brown oil. ¹H NMR (400MHz, CDCl₃) δ = 7.63 (s, 1H), 4.82 (s, 2H), 2.52 (s, 3H).


[1083] 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((5-(methylthio)thiazol-2-

yl)methoxy)phenyl)propan-2-yl)benzonitrile (6): To a mixture of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (100 mg, 0.3 mmol) and 2-(chloromethyl)-5-(methylthio)thiazole (103 mg, 0.6 mmol) in DMF (3 mL) was added Cs₂CO₃ (186 mg, 0.6 mmol) and the mixture was stirred at 20 °C for 2 hrs. LCMS showed the reaction was completed. The resulting mixture was quenched with water (10 mL) and extracted with EtOAc (5 mL x 3). The combined organic layers were washed with brine (5 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduce pressure. The residue was purified by MPLC to give 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((5-(methylthio)thiazol-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (80 mg, yield: 56.8 %) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ = 7.66 (s, 1H), 7.44 (s, 1H), 7.33 (s, 1H), 7.12 (br d, *J*=8.4 Hz, 2H), 6.95 (br d, *J*=8.8 Hz, 2H), 5.30 (s, 2H), 4.42 (t, *J*=6.0 Hz, 2H), 3.88 (t, *J*=6.0 Hz, 2H), 2.50 (s, 3H), 1.65 (s, 6H).

[1084] 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((5-(methylthio)thiazol-2-

yl)methoxy)phenyl)propan-2-yl)benzonitrile (A93): A suspension of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((5-(methylthio)thiazol-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (80 mg, 0.16 mmol) and m-CPBA (80.0 %, 105 mg, 0.49 mmol) in DCM (3 mL) was stirred at 20 °C for 1 h. LCMS showed the reaction was completed. The resulting mixture was poured into saturated aqueous NaHCO₃ (3 mL) and extracted with DCM (3 mL x 3). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered and concentrated under reduce pressure. The residue was purified by prep-HPLC (TFA) to give 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((5-(methylsulfonyl)thiazol-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (64.2 mg, yield: 75.4 %) as white solid. ¹H NMR (400MHz, CDCl₃) δ = 8.31 (s, 1H), 7.45 (d, *J*=2.4 Hz, 1H), 7.32 (d, *J*=2.4 Hz, 1H), 7.16 (d, *J*=8.8 Hz, 2H), 6.95 (d, *J*=8.8 Hz, 2H), 5.40 (s, 2H), 4.43 (t, *J*=6.0 Hz, 2H), 3.88 (t, *J*=6.0 Hz, 2H), 3.25 (s, 3H), 1.66 (s, 6H). LCMS (220 nm): 99.5%. Exact Mass: 524.0; found 525.1, 527.1.

[1085] Example 22: Synthesis of N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)pyrimidin-2-yl)methanesulfonamideN-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)pyrimidin-2-yl)methanesulfonamide (A109)




[1086] 2-chloro-4-(chloromethyl)pyrimidine (2): To a mixture of 2-chloro-4-methylpyrimidine (50.0 g, 398 mmol) and NCS (77.9 g, 583 mmol) in MeCN (250 mL) was added benzoyl benzenecarboperoxoate (28.3 g, 117 mmol) in portions at 20 °C and the mixture was stirred at 100 °C for 16 hrs under N₂ atmosphere. TLC showed most of the starting material consumed and two new spots appeared. The mixture was cooled down to room temperature, poured into water (500 mL) and extracted with EtOAc (200 mL x 3). The organic layers were combined and washed with brine (200 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give 2-chloro-4-(chloromethyl) pyrimidine (22 g, yield: 31.2 %) as yellow oil. ¹H NMR (400 MHz, CDCl₃) δ = 8.69 (d, *J*=5.2 Hz, 1H), 7.54 (d, *J*=5.0 Hz, 1H), 4.61 (s, 2H).

[1087] 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((2-chloropyrimidin-4-yl)methoxy)phenyl)propan-2-yl)benzonitrile (4): To a mixture of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (18.0 g, 51.4 mmol) and 2-chloro-4-(chloromethyl) pyrimidine (10.1 g, 61.7 mmol) in DMF (150 mL) was added Cs₂CO₃ (33.5 g, 103.4 mmol) at 20 °C and the mixture was stirred at the same temperature for 16 hrs. LCMS showed the reaction was completed. The reaction mixture was poured into H₂O (300 mL) and extracted with EtOAc (150 mL x 3). The combined organic layers were washed with brine (150 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((2-chloropyrimidin-4-yl)methoxy)phenyl)propan-2-yl)benzonitrile (15.5 g, yield: 63.3%) as white solid. ¹H NMR (400 MHz, CDCl₃) δ = 8.67 (d, *J* = 5.2 Hz, 1H), 7.56 (d, *J* = 5.2 Hz, 1H), 7.45 (d, *J* = 2.4 Hz, 1H), 7.35 - 7.29 (m, 1H), 7.13 (d, *J* = 8.8 Hz, 2H), 6.90 (d, *J* = 8.8 Hz, 2H), 5.16 (s, 2H), 4.43 (t, *J* = 6.0 Hz, 2H), 3.88 (t, *J* = 6.0 Hz, 2H), 1.65 (s, 6H).

[1088] N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-

yl)phenoxy)methyl) pyrimidin-2-yl)methanesulfonamide (A109): To a mixture of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((2-chloropyrimidin-4-yl)methoxy)phenyl)propan-2-yl)benzonitrile (15.5 g, 32.5 mmol), methane sulfonamide (9.3 g, 97.5 mmol), Cs₂CO₃ (21.2 g, 65.0 mmol) and Xantphos (1.88 g, 3.25 mmol) in 1,4-dioxane (450 mL) was added Pd₂(dba)₃ (3.0 g, 3.3 mmol) at 20 °C and the mixture was stirred at 90 °C for 6 hrs under N₂ atmosphere. LCMS showed the reaction was completed. The mixture was cooled down to room temperature, poured into water (300 mL) and extracted with EtOAc (300 mL x 3). The combined organic layers were washed with brine (300 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give the crude product and then further purified by p-HPLC (TFA) to give N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)pyrimidin-2-yl)methanesulfonamide (5.30 g, yield: 30.1 %) as yellow solid. ¹H NMR (400 MHz, CDCl₃) δ = 10.02 (br s, 1H), 8.69 (d, *J* = 5.2 Hz, 1H), 7.45 (d, *J* = 2.4 Hz, 1H), 7.34 - 7.31 (m, 1H), 7.30 (d, *J* = 5.2 Hz, 1H), 7.13 (d, *J* = 8.8 Hz, 2H), 6.91 (d, *J* = 8.8 Hz, 2H), 5.13 (s, 2H), 4.43 (t, *J* = 6.0 Hz, 2H), 3.88 (t, *J* = 6.0 Hz, 2H), 3.47 (s, 3H), 1.65 (s, 6H). LCMS (220 nm): 99.0%. Exact Mass: 534.09; found 535.1, 537.0.

[1089] **Example 23:** Synthesis of N-(4-(1-(4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)ethyl)oxazol-2-yl)methanesulfonamide (A122)



[1090] **2-Chloro-4-(1-ethoxyvinyl)-5-methylpyrimidine (2):** To a mixture of 2,4-dichloro-5-methylpyrimidine (10.0 g, 61.3 mmol) and tributyl(1-ethoxyvinyl)stannane (22.2 g, 0.0613

mol) in DMF (200 mL) was added Pd(PPh₃)₂Cl₂ (2.15 g, 3.1 mmol) at 20 °C and the mixture was stirred at 100°C for 5 hrs under N₂ atmosphere. TLC showed the reaction was completed. The reaction was quenched by KF solution (30 g, 0.172 mol dissolved 500 mL in water), filtered and extracted with EtOAc (100 mL x 3). The combined organic layers were washed with brine (100 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give 2-chloro-4-(1-ethoxyvinyl)-5-methyl-pyrimidine (6.0 g, yield: 41.5 %) as yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 8.40 (s, 1H), 5.01 (d, *J* = 2.4 Hz, 1H), 4.55 (d, *J* = 2.4 Hz, 1H), 3.94 (q, *J* = 7.2 Hz, 2H), 2.38 (s, 3H), 1.41 (t, *J* = 7.2 Hz, 3H).

[1091] Ethyl 2-chloro-5-methylpyrimidine-4-carboxylate (3): O₃ (gas) was bubbled into a solution of 2-chloro-4-(1-ethoxyvinyl)-5-methyl-pyrimidine (**2**) (purity: 50.0 %, 6.0 g, 0.0151 mol) in DCM (60.0 mL) at -30 °C for 0.5 hour. TLC showed the reaction was completed. O₂ and N₂ was bubbled into the above solution at 10 °C for 0.5 hr. (CH₃)₂S (10.0 mL, 0.0151 mol) was added into the reaction and the mixture was stirred for 0.5 hr at 20 °C. TLC showed the reaction was completed. The mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give ethyl 2-chloro-5-methylpyrimidine-4-carboxylate (1.70 g, yield: 56.1 %) as colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 8.62 (s, 1H), 4.48 (q, *J* = 7.2 Hz, 2H), 2.50 (s, 3H), 1.44 (t, *J* = 7.2 Hz, 3H).

[1092] Ethyl 2-(N-(2,4-dimethoxybenzyl)methylsulfonamido)-5-methylpyrimidine-4-carboxylate (4): To a mixture of ethyl 2-chloro-5-methylpyrimidine-4-carboxylate (**3**) (0.20 g, 0.997 mmol) and N-[(2,4-dimethoxyphenyl)methyl]methanesulfonamide (0.49 g, 1.99 mmol) in DMF (5.0 mL) was added NaH (60.0 %, 0.076 g, 1.99 mmol) at 0 °C. The mixture was stirred at 60°C for 6 hrs. LCMS showed the reaction was completed. The mixture was quenched by saturated aqueous NH₄Cl (10 mL) and extracted with EtOAc (15 mL x 3). The combined organic layers were combined and washed with brine (15 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give ethyl 2-(N-(2,4-dimethoxybenzyl)methylsulfonamido)-5-methylpyrimidine-4-carboxylate (0.180 g, yield: 44.1 %) as colorless oil. ¹H NMR (400 MHz, CDCl₃): δ 8.53 (s, 1H), 7.33-7.29 (m, 1H), 6.42-6.37 (m, 2H), 5.29 (s, 2H), 4.44 (q, *J* = 7.2 Hz, 2H), 3.78 (s, 3H), 3.66 (s, 3H), 3.36 (s, 3H), 2.44 (s, 3H), 1.43 (t, *J* = 7.2 Hz, 3H).

[1093] N-(2,4-Dimethoxybenzyl)-N-(4-(hydroxymethyl)-5-methylpyrimidin-2-yl)methanesulfonamide (5): To a solution of ethyl 2-(N-(2,4-dimethoxybenzyl)methylsulfonamido)-5-methylpyrimidine-4-carboxylate (**4**) (0.18 g, 0.44 mmol) in THF (5 mL) was added LiBH₄ (0.038 g, 1.8 mmol) in portions at 0 °C. The mixture

was stirred at 20°C for 2 hrs. TLC showed the reaction was completed. The mixture was quenched by saturated aqueous NH₄Cl (20 mL), extracted with EtOAc (10 mL x 2). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography to give N-(2,4-dimethoxybenzyl)-N-(4-(hydroxymethyl)-5-methylpyrimidin-2-yl) methanesulfonamide (0.080 g, yield: 50 %) as colorless oil. ¹H NMR (400 MHz, CDCl₃): δ 8.29 (s, 1H), 7.25 (br s, 1H), 6.45-6.41 (m, 2H), 5.31 (s, 2H), 4.63 (d, *J* = 4.8 Hz, 2H), 4.03 (t, *J* = 4.8 Hz, 1H), 3.78 (s, 3H), 3.74 (s, 3H), 3.41 (s, 3H), 2.12 (s, 3H).


[1094] (2-(N-(2,4-Dimethoxybenzyl)methylsulfonamido)-5-methylpyrimidin-4-yl)methyl methanesulfonate (6): To a solution of N-(2,4-dimethoxybenzyl)-N-(4-(hydroxymethyl)-5-methylpyrimidin-2-yl)methanesulfonamide (**5**) (0.10 g, 0.272 mmol) in DCM (2 mL) was added TEA (0.0759 mL, 0.544 mmol) and MsCl (0.047 g, 0.41 mmol) at 0°C. The mixture was stirred at 20°C for 0.5 hr under N₂ atmosphere. LCMS showed the reaction was completed. The reaction was quenched by saturated aqueous NH₄Cl solution (2 mL), extracted with DCM (3 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give (2-(N-(2,4-dimethoxybenzyl)methylsulfonamido)-5-methylpyrimidin-4-yl)methyl methanesulfonate (0.15 g, crude). ¹H NMR (400 MHz, CDCl₃): δ 8.39 (s, 1H), 7.24 (s, 1H), 6.39 (s, 2H), 5.29 - 5.28 (m, 2H), 4.54 (s, 2H), 3.78 (s, 6H), 3.62 (s, 3H), 3.28 (s, 3H), 2.80 (s, 3H).

[1095] N-(4-((4-(2-(3-Chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)-5-methylpyrimidin-2-yl)-N-(2,4-dimethoxybenzyl)methanesulfonamide (7): To a solution of (2-(N-(2,4-dimethoxybenzyl)methylsulfonamido)-5-methylpyrimidin-4-yl)methyl methanesulfonate (**6**) (0.10 g, 0.22 mmol) in DMF (2.0 mL) was added Cs₂CO₃ (0.15 g, 0.45 mmol) and 3-chloro-2-(2-chloroethoxy)-5-[1-(4-hydroxyphenyl)-1-methyl-ethyl]benzonitrile (**Int H**) (0.079 g, 0.22 mmol) at 20°C. The mixture was stirred at 20 °C for 10 hrs. TLC showed the reaction was completed. The mixture was quenched with H₂O (3 mL), extracted with EtOAc (3 mL x 3). The combined organic layers were washed with brine (4 mL x 3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)-5-methylpyrimidin-2-yl)-N-(2,4-dimethoxybenzyl)methanesulfonamide (0.10 g, crude) as yellow oil which was used directly without further purification.

[1096] N-(4-((4-(2-(3-Chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)-5-methylpyrimidin-2-yl)methanesulfonamide (A122): To a solution

of N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)-5-methylpyrimidin-2-yl)-N-(2,4-dimethoxybenzyl)methanesulfonamide (**7**) (0.05 g, crude) in DCM (1 mL) was added TFA (0.0159 mL, 0.214 mmol) and the mixture was stirred at 20 °C for 15 min. LCMS showed the reaction was completed. The mixture was quenched with H₂O (2 mL), extracted with DCM (3 mL x 3). The combined organic layers were washed with brine (2 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (FA) to give N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)-5-methylpyrimidin-2-yl)methanesulfonamide (purify: 100%, 7 mg, yield: 17.8%) as white solid. ¹H NMR (400 MHz, CDCl₃): δ 8.41 (s, 2H), 8.25-8.30 (br, 1H), 7.44 (d, *J* = 2.4 Hz, 1H), 7.31 (d, *J* = 2.4 Hz, 1H), 7.11 (d, *J* = 6.8 Hz, 2H), 6.92 (d, *J* = 6.8 Hz, 2H), 5.12 (s, 2H), 4.42 (t, *J* = 6.1 Hz, 2H), 3.88 (t, *J* = 6.4 Hz, 2H), 3.41 (s, 3H), 2.35 (s, 3H), 1.64 (s, 6H). LCMS: (220 nm): 100%. Exact Mass: 548.1; found 549.2, 551.1.


[1097] Example 24: Synthesis of N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazol-2-yl)methanesulfonamide (**A134**)



[1098] 5-(2-(4-((2-Aminothiazol-4-yl)methoxy)phenyl)propan-2-yl)-3-chloro-2-(2-chloroethoxy)benzonitrile (3**):** To a mixture of 4-(chloromethyl)thiazol-2-aminohydrochloride (**1**) (200 mg, 1.08 mmol) and 3-chloro-2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (**2**) (378 mg, 1.08 mmol) in DMF (5 mL) was added Cs₂CO₃ (1.4 g, 4.32 mol) at 20 °C and the mixture was stirred at 130 °C for 10 hrs. LCMS showed the reaction was completed. The reaction was cooled down, quenched with water (10 mL). The mixture was extracted with DCM (5 mL x 3). The combined organic layers were washed with brine (5 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by HPLC (FA) to obtain 5-(2-(4-((2-aminothiazol-4-yl)methoxy)phenyl)propan-2-yl)-3-chloro-2-(2-chloroethoxy) benzonitrile (**3**) (15.0 mg, yield: 3 %) as yellow solid. ¹H NMR (400MHz, CDCl₃) δ = 7.44 (s, 1H), 7.32 (s, 1H), 7.11 (br d, *J*=6.6 Hz, 2H), 6.93 (br s, 2H), 6.53 (br s, 1H), 5.06 - 4.83 (m, 2H), 4.42 (t, *J*=6.2 Hz, 2H), 3.88 (t, *J*=6.2 Hz, 2H), 1.64 (s, 6H).

[1099] N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazol-2-yl)methanesulfonamide (A134): To a mixture of 5-(2-(4-((2-aminothiazol-4-yl)methoxy)phenyl)propan-2-yl)-3-chloro-2-(2-chloroethoxy)benzonitrile (**3**) (15.0 mg, 0.03 mmol) and TEA (0.01 mL, 0.10 mmol) in DCM (1 mL) was added methanesulfonyl chloride (4.46 mg, 0.03 mmol) and the mixture was stirred at 25 °C for 30 min (two Ms groups are installed in LCMS). Then Cs₂CO₃ (21.1 mg, 0.0649 mol) in H₂O (0.5 mL) was added into the reaction. The mixture was stirred at 50 °C for 10 min. LCMS showed the reaction was completed. The reaction was quenched with water (2 mL) and extracted with DCM (3 mL x 3). The combined organic layers were washed with brine (3 mL x 2), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by prep-HPLC (HCOOH) to obtain N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazol-2-yl) methanesulfonamide (**A134**) (1.8 mg, yield: 10.3 %) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ = 7.44 (d, *J*=2.2 Hz, 1H), 7.32 (d, *J*=2.2 Hz, 1H), 7.13 (d, *J*=8.8 Hz, 2H), 6.90 (d, *J*=8.8 Hz, 2H), 6.49 (s, 1H), 4.98 (s, 2H), 4.43 (t, *J*=6.2 Hz, 2H), 3.88 (t, *J*=6.2 Hz, 2H), 3.01 (s, 3H), 1.65 (s, 6H). LCMS: (220 nm): 94.9%. Exact Mass: 539.1; found 540.0/542.0.

[1100] Example 25: Synthesis of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((4-(methylsulfonyl)thiazol-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (A136)



[1101] (4-Bromothiazol-2-yl)methanol (2): To a solution of 4-bromothiazole-2-carbaldehyde (10.4 g, 0.0542 mol) in MeOH (150 mL) was added NaBH₄ (4.10 g, 0.108 mol) at 0 °C. The mixture was stirred at 25 °C for 2 hrs. TLC showed the reaction was completed and one new spot was observed. The mixture was quenched with water (70 mL), stirred for 0.5 hr and concentrated to remove most of MeOH. The aqueous layer was extracted with

EtOAc (30 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (SiO₂, Petroleum ether/Ethyl acetate=10:1 to 3:1) to give (4-bromothiazol-2-yl)methanol (10.0 g, 0.0515 mol, yield: 95.2 %) as white solid. ¹H NMR (400MHz, CDCl₃) δ 7.23 (s, 1H), 4.96 (d, *J*=6.0 Hz, 2H), 2.79 (t, *J*=6.0 Hz, 1H).

[1102] 4-Bromo-2-(((tert-butyldimethylsilyl)oxy)methyl)thiazole (3): To a solution of (4-bromothiazol-2-yl)methanol (11.7 g, 0.0603 mol) in DCM (120 mL) was added imidazole (8.21 g, 0.121 mol) and tert-butyl-chloro-dimethyl-silane (10.0 g, 0.0663 mol). The mixture was stirred at 25 °C for 1 hr. TLC showed the starting material was consumed and one new spot was observed. The reaction was poured into water (100 mL), extracted with DCM (70 mL x 2). The combined organic layers were washed with brine (70 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica-gel column chromatography to give (4-bromothiazol-2-yl)methoxy-tert-butyl-dimethyl-silane (17.0 g, 0.0551 mol, yield: 91.5%) as colorless oil. ¹H NMR (400MHz, CDCl₃) δ 7.18 (s, 1H), 4.95 (s, 2H), 0.96 (s, 9H), 0.14 (s, 6H).

[1103] 2-(((tert-Butyldimethylsilyl)oxy)methyl)-4-(methylthio)thiazole (4): To a solution of (4-bromothiazol-2-yl)methoxy-tert-butyl-dimethyl-silane (11.5 g, 0.0373 mol) in MTBE (300 mL) was added n-BuLi (2.50 M, 17.9 mL, 0.0448 mol) dropwise at -78 °C under N₂ atmosphere. The mixture was stirred at -78 °C for 30 min. Then MeSSMe (14.3 g, 0.152 mol) was added at -78 °C and stirred at the same temperature for 1 hr. LCMS showed 70% desired MS was detected. The mixture was quenched with sat. NH₄Cl solution (100 mL), extracted with EtOAc (30 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give crude tert-butyl-dimethyl-[(4-methylsulfanylthiazol-2-yl)methoxy]silane (90.0 %, 10.0 g, 0.0327 mol, yield: 87.6 %) as brown oil. ¹H NMR (400MHz, CDCl₃) δ 6.87 (s, 1H), 4.95 (s, 2H), 2.53 (s, 3H), 0.96 (s, 9H), 0.14 (s, 6H).

[1104] (4-(Methylthio)thiazol-2-yl)methanol (5): To a solution of (4-bromothiazol-2-yl)methoxy-tert-butyl-dimethyl-silane (12.0 g, 0.0389 mol) in THF (150 mL) was added TBAF (1.00 M, 46.7 mL, 0.0467 mol) at 0 °C. The mixture was stirred at 20 °C for 1 hr. TLC showed the reaction was completed. The reaction was poured into water (50 mL), extracted with EtOAc (30 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica-gel column chromatography to give (4-methylsulfanylthiazol-2-yl) methanol (3.60 g, 0.0223 mol, yield: 57.4 %) as yellow solid. ¹H NMR (400MHz, CDCl₃) δ 6.89 (s, 1H), 4.94

(br d, $J=2.0$ Hz, 2H), 2.95 (br s, 1H), 2.54 (s, 3H).


[1105] (4-(Methylthio)thiazol-2-yl)methyl methanesulfonate (6): To a mixture of (4-methylsulfanyltiazol-2-yl)methanol (3.40 g, 0.0211 mol) and TEA (0.432 mL, 0.00310 mol) in DCM (50 mL) was added MsCl (1.96 mL, 0.0253 mol) at 0°C. The mixture was stirred at 0 °C for 1hr. TLC showed the starting material was consumed and one new spot was observed. The reaction was poured into water (30 mL), extracted with DCM (15 mL x 3). The combined organic layers were washed with brine (40 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give (4-methylsulfanyltiazol-2-yl)methyl methanesulfonate (5.40 g, 0.0226 mol, crude) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ 6.99 (s, 1H), 5.45 (s, 2H), 3.10 (s, 3H), 2.55 (s, 3H).

[1106] 3-Chloro-2-(2-chloroethoxy)-5-(2-(4-((4-(methylthio)thiazol-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (7): To a suspension of (4-methylsulfanyltiazol-2-yl)methyl methanesulfonate (80.0 %, 3.08 g, 0.0103 mol) and Cs₂CO₃ (5.58 g, 0.0171 mol) in DMF (20 mL) was added 3-chloro-2-(2-chloroethoxy)-5-[1-(4-hydroxyphenyl)-1-methyl-ethyl]benzonitrile (3.00 g, 0.00857 mol) at 20°C. The mixture was stirred at 20 °C for 16 hrs. LCMS showed the reaction was completed. The reaction was poured into H₂O (50 mL), extracted with EtOAc (30 mL x 3). The combined organic layers were washed with brine (30 mL x 3), filtered and concentrated under reduced pressure to give 3-chloro-2-(2-chloroethoxy)-5-[1-methyl-1-[4-[(4-methylsulfanyltiazol-2-yl)methoxy]phenyl]ethyl]benzonitrile (3.20 g, 0.00648 mol, yield: 75.7 %) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ 7.44 (d, $J=2.4$ Hz, 1H), 7.32 (d, $J=2.4$ Hz, 1H), 7.12 (d, $J=8.4$ Hz, 2H), 6.99 - 6.91 (m, 3H), 5.34 (s, 2H), 4.42 (t, $J=6.4$ Hz, 2H), 3.88 (t, $J=6.4$ Hz, 2H), 2.56 (s, 3H), 1.64 (s, 6H).

[1107] 3-Chloro-2-(2-chloroethoxy)-5-(2-(4-((4-(methylsulfonyl)thiazol-2-yl)methoxy)phenyl) propan-2-yl)benzonitrile (A136): To a solution of 3-chloro-2-(2-chloroethoxy)-5-[1-methyl-1-[4-[(4-methylsulfanyltiazol-2-yl)methoxy]phenyl]ethyl]benzonitrile (3.00 g, 0.00608 mol) in DCM (50 mL) was added m-CPBA (80.0 %, 5.25 g, 0.0243 mol) at 0°C. The mixture was stirred at 25°C for 3 hrs. LCMS showed the reaction was completed and 60% desired product was detected. The mixture was quenched with water (30 mL). The mixture was extracted with DCM (20 mL x 3). The combined organic layers were washed with brine (50 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (FA) to give 3-chloro-2-(2-chloroethoxy)-5-[1-methyl-1-[4-[(4-methylsulfonyltiazol-2-yl)methoxy]phenyl]ethyl]benzonitrile (**A136**) (1.71 g, 0.00325 mol, yield: 53.5 %) as white

solid. ^1H NMR (400MHz, CDCl_3) δ 8.22 (s, 1H), 7.45 (d, $J=2.4$ Hz, 1H), 7.32 (d, $J=2.0$ Hz, 1H), 7.15 (d, $J=6.8$ Hz, 2H), 6.95 (d, $J=6.8$ Hz, 2H), 5.39 (s, 2H), 4.43 (t, $J=6.4$ Hz, 2H), 3.88 (t, $J=6.0$ Hz, 2H), 3.23 (s, 3H), 1.66 (s, 6H).

[1108] Example 26: Synthesis of N-(2-((4-(2-(3-Chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazol-4-yl)methanesulfonamide (**A137**)



[1109] Ethyl 2-methylthiazole-4-carboxylate (2): To a mixture of ethyl 2-methylthiazole-4-carboxylate (**1**) (10.0 g, 58.4 mmol) and AIBN (0.959 g, 5.84 mmol) in CCl_4 (80 mL) was added NBS (12.5 g, 70.1 mmol) at 20°C and the mixture was stirred at 80°C for 16 hrs. TLC showed a new spot appeared and trace of starting material was remained. The reaction was poured into H_2O (150 mL) and extracted with DCM (80 mL x 2). The combined organic layers were washed with brine (80 mL), dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The residue was purified by silica gel to give ethyl 2-methylthiazole-4-carboxylate (**2**) (4.6 g, Yield: 31.5%) as yellow oil. ^1H NMR (400MHz, CDCl_3) δ = 8.23 (s, 1H), 4.77 (s, 2H), 4.41 (q, $J = 6.8$ Hz, 2H), 1.42 (t, $J = 7.2$ Hz, 3H).

[1110] Ethyl 2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy) methyl)thiazole-4-carboxylate (3): A solution of 3-chloro-2-(2-chloroethoxy)-5-(1-(4-hydroxyphenyl)-1-methyl-ethyl)benzonitrile (**int H**) (0.5 g, 1.43 mmol), ethyl 2-(bromomethyl) thiazole-4-carboxylate (**2**) (0.428 g, 1.71 mmol) and Cs_2CO_3 (0.93, 2.86 mmol) in DMF (10 mL) was stirred at 20°C for 16 hrs. LCMS showed the reaction was completed. The reaction was poured into H_2O (15 mL) and extracted with EtOAc (15 mL x 3). The combined organic layers were washed with brine (10 mL x 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by silica gel to give ethyl 2-

((4-(1-(3-chloro-4-(2-chloroethoxy)-5-cyano-phenyl)-1-methyl-ethyl)phenoxy)methyl)thiazole-4-carboxylate (**3**) (0.6 g, Yield: 80.9%) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ = 8.27 (s, 1H), 7.44 (d, *J*=2.4 Hz, 1H), 7.32 (d, *J*=2.4 Hz, 1H), 7.12 (d, *J*=8.8 Hz, 2H), 5.41 (s, 2H), 4.44 - 4.09 (m, 4H), 3.88 (t, *J*=6.4 Hz, 2H), 1.65 (s, 1H), 1.43 (t, *J*=7.6 Hz, 3H).

[1111] 2-((4-(2-(3-Chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazole-4-carboxylic acid (4**):**

A solution of ethyl 2-((4-(1-(3-chloro-4-(2-chloroethoxy)-5-cyano-phenyl)-1-methyl-ethyl)phenoxy)methyl)thiazole-4-carboxylate (**3**) (0.5 g, 0.963 mmol) and LiOH.H₂O (0.162 g, 3.85 mmol) in THF (3 mL) and H₂O (3 mL) was stirred at 20°C for 16 hrs. LCMS showed the reaction was completed. The reaction was poured into H₂O (8 mL). The pH of the aqueous layer was adjusted to 3~4 by adding aqueous HCl solution (3M). The aqueous layer was separated and extracted with EtOAc (8 mL x 3). The combined organic layers were washed with brine (8 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazole-4-carboxylic acid (**4**) (0.4 g, yield: 84.6%) as white solid. ¹H NMR (400MHz, CDCl₃) δ = 8.34 (s, 1H), 7.44 (d, *J*=2.4 Hz, 1H), 7.32 (d, *J*=2.4 Hz, 1H), 7.14 (d, *J* = 8.8 Hz, 2H), 6.95 (d, *J* = 8.8 Hz, 2H), 5.41 (s, 2H), 4.43 (t, *J* = 6.0 Hz, 2H), 3.88 (t, *J*=6.4 Hz, 2H), 1.65 (s, 6H).

[1112] tert-Butyl (2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazol-4-yl)carbamate (5**):**

To a solution of 2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazole-4-carboxylic acid (**4**) (0.1 g, 0.204 mmol) and TEA (0.0851 mL, 0.000611 mol) in toluene (1.0 mL) was added t-BuOH (1.0 mL) and DPPA (0.132 mL, 0.611 mmol) under N₂ atmosphere. The mixture was stirred at 90°C for 4 hrs. LCMS showed the starting material was consumed. The mixture was poured into H₂O (6 mL), extracted with DCM (3 mL x 2). And the combined organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel tert-butyl (2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazol-4-yl)carbamate (**5**) (0.11g, yield: 96.1%) as white oil. ¹H NMR (400MHz, CDCl₃) δ = 7.43 (d, *J* = 2.4 Hz, 1H), 7.33 (d, *J*=2.4 Hz, 1H), 7.25 (s, 1H), 7.11 (d, *J*=8.8 Hz, 2H), 6.93 (d, *J*=8.8 Hz, 2H), 5.25 (s, 2H), 4.43 (t, *J*=6.0 Hz, 2H), 3.88 (t, *J*=6.0 Hz, 2H), 1.65 (s, 6H), 1.53 (s, 9H).


[1113] 5-(2-(4-((4-Aminothiazol-2-yl)methoxy)phenyl)propan-2-yl)-3-chloro-2-propoxybenzonitrile (6**):**

To a solution of tert-butyl (2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazol-4-yl)carbamate (**5**) (0.1 g, 0.0178 mmol)

in DCM (1 mL) was added TFA (0.1 mL) and stirred at 25°C for 2 hrs. LCMS showed the reaction was completed. The reaction was poured into saturated aqueous NaHCO₃ solution (4 mL) and extracted with DCM (2 mL x 3). The combined organic layers were washed with brine (2 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 5-(2-(4-((4-aminothiazol-2-yl)methoxy)phenyl)propan-2-yl)-3-chloro-2-propoxybenzonitrile (**6**) (30 mg, Yield: 35.1%) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ = 7.44 (d, *J*=2.0 Hz, 1H), 7.33 (d, *J*=2.4 Hz, 1H), 7.25 (s, 1H), 7.11 (d, *J*=8.8 Hz, 2H), 6.94 (d, *J*=8.8 Hz, 2H), 5.24 (s, 2H), 4.42 (t, *J* = 2.4 Hz, 2H), 3.88 (t, *J* = 2.4 Hz, 2H), 1.65 (s, 6H)

[1114] N-(2-((4-(2-(3-Chloro-4-(2-chloroethoxy))-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazol-4-yl)methanesulfonamide (A137): To a solution of 5-(2-(4-((4-aminothiazol-2-yl)methoxy)phenyl)propan-2-yl)-3-chloro-2-propoxybenzonitrile (**6**) (50 mg, 0.108 mmol) and TEA (21.9 mg, 0.216 mmol) in DCM (2 mL) was added methanesulfonyl chloride (18.6 mg, 0.0649 mmol) dropwisely at 0°C under nitrogen and the mixture was stirred at 25°C for 2 hrs. (two Ms groups were installed). Then the mixture was added Cs₂CO₃ (0.141 g, 0.433 mmol) and H₂O (1 mL) and stirred at 25°C for 2 hrs. LCMS showed the reaction was completed. The reaction was poured into H₂O (4 mL). The aqueous layer was extracted with EtOAc (4 mL x 3). The combined organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by prep-HPLC (HCOOH) to give product N-(2-((4-(2-(3-chloro-4-(2-chloroethoxy))-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazol-4-yl)methanesulfonamide (**A137**) (10 mg, yield: 17.1%) as white solid. ¹H NMR (400MHz, CDCl₃) δ = 7.44 (d, *J*=2.4 Hz, 1H), 7.32 (d, *J*=2.4 Hz, 1H), 7.13 (d, *J*=8.8 Hz, 2H), 7.04 (s, 1H), 7.00 (s, 1H), 6.94 (d, *J*=8.8 Hz, 2H), 5.29 (s, 2H), 4.46 (t, *J* = 6.0 Hz, 2H), 3.88 (t, *J* = 6.0 Hz, 2H), 3.12 (s, 3H), 1.65 (s, 6H). LCMS: (220 nm): 93.78%. Exact Mass: 540.5; found: 540.1/542.1.


[1115] Example 27: Synthesis of 2-((4-(2-(3-chloro-4-(2-chloroethoxy))-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazole-4-carboxamide (**A171**)



[1116] To a solution of 2-((4-(2-(3-chloro-4-(2-chloroethoxy))-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazole-4-carboxylic acid (**4**) (20 mg, 0.0407 mmol), NH₄Cl (5.4 mg,

0.102 mmol), HATU (0.0186 g, 0.0488 mmol) in DCM (1.00 mL) was added TEA (8.2 mg, 0.0814 mmol) at 20 °C. The mixture was stirred at same temperature for 2 hrs. LCMS showed the starting material was consumed. The mixture was poured into H₂O (4 mL) and extracted with DCM (3 mL x 2). The combined organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified prep-HPLC (HCOOH) to give 2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazole-4-carboxamide (**A171**) (9.2 mg, yield: 46.1%) as white solid. ¹H NMR (400MHz, CDCl₃) δ = 8.19 (s, 1H), 7.44 (d, *J*=2.2 Hz, 1H), 7.32 (d, *J*=2.2 Hz, 1H), 7.14 (br d, *J*=8.8 Hz, 2H), 6.95 (d, *J*=8.8 Hz, 2H), 5.60 (br s, 1H), 5.34 (s, 2H), 4.43 (t, *J*=6.2 Hz, 2H), 3.88 (t, *J*=6.2 Hz, 2H), 1.66 (s, 6H). LCMS (220 nm): 99.5%. Exact Mass: 489.0; found 490.

[1117] Example 28: Synthesis of 2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazole-4-carbonitrile (**A172**)



[1118] To a solution 2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazole-4-carboxamide (**A171**) (20 mg, 0.0408 mmol) and pyridine (3.27 mg, 0.0413 mmol) in dioxane (2.0 mL) was added TFAA (17.4 mg, 0.00827 mmol) under N₂ at 0 °C. Then the reaction was stirred at 20 °C for 16 hrs. LCMS showed the reaction was completed. The reaction was poured into H₂O (4 mL). The reaction mixture were separated and extracted with EtOAc (4 mL x 3). The organic layers were washed with brine (4 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by prep-HPLC (HCOOH) to give the product 2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)thiazole-4-carbonitrile (**A172**) (10 mg, yield: 52.1%) as white solid. ¹H NMR (400MHz, CDCl₃) δ = 8.05 (s, 1H), 7.44 (d, *J*=2.2 Hz, 1H), 7.31 (d, *J*=2.2 Hz, 1H), 7.15 (d, *J*=8.8 Hz, 2H), 6.94 (d, *J*=8.8 Hz, 2H), 5.37 (s, 2H), 4.43 (t, *J*=6.2 Hz, 2H), 3.88 (t, *J*=6.2 Hz, 2H), 1.65 (s, 6H). LCMS (220 nm): 100.0%. Exact Mass: 471.0; found 472.0.

[1119] Example 29: Synthesis of N-(4-(aminomethyl)-6-((4-(2-(3-chloro-4-(2-

Synthesis of A164:

Starting material **1** (2-chloro-4-hydroxy-5-(chloromethyl)pyrimidine) is reacted with POCl_3 at 100°C for 3 hrs to form intermediate **2** (2,4-dichloro-5-(chloromethyl)pyrimidine). Intermediate **2** is then reacted with **3** (N-benzyl-2-chloro-1H-indole-3-carboxamide) in DMF at 20°C for 2 hrs to form intermediate **4** (N-benzyl-2-chloro-1H-indole-3-carboxamide derivative). Intermediate **4** is reacted with BF_3K , TEA, and $\text{Pd}(\text{dppf})\text{Cl}_2$ in n-PrOH at 100°C for 2 hrs to form intermediate **5** (N-benzyl-2-chloro-1H-indole-3-carboxamide derivative with a vinyl group).

Intermediate **5** is then reacted with O_3 and Me_2S in DCM at -78 to -20°C for 1 hr to form intermediate **6** (N-benzyl-2-chloro-1H-indole-3-carboxamide derivative with an aldehyde group). Intermediate **6** is then reacted with NaBH_4 in THF at 0°C for 15 min to form intermediate **7** (N-benzyl-2-chloro-1H-indole-3-carboxamide derivative with a primary alcohol group).

Intermediate **7** is then reacted with MsCl and TEA in DCM at 0°C for 20 min to form intermediate **8** (N-benzyl-2-chloro-1H-indole-3-carboxamide derivative with a mesylate group). Intermediate **8** is then reacted with **int H** (4-chloro-2-(chloromethoxy)-3-cyano-5-(4-chlorophenyl)phenol) in the presence of Cs_2CO_3 in DMF at 20 to 40°C for 1.5 hrs to form intermediate **9** (N-benzyl-2-chloro-1H-indole-3-carboxamide derivative with a 4-chlorophenyl group).

Intermediate **9** is then reacted with MsNH_2 , Xantphos, $\text{Pd}(\text{dba})_3$, and Cs_2CO_3 in dioxane at 100°C for 2 hrs to form intermediate **10** (N-benzyl-2-chloro-1H-indole-3-carboxamide derivative with a 4-chlorophenyl group and a mesylate group).

Intermediate **10** is then reacted with $\text{NH}_2\text{NH}_2\cdot\text{H}_2\text{O}$ in DCM/MeOH at 20°C for 1 hr to form the final product **A164** (N-benzyl-2-chloro-1H-indole-3-carboxamide derivative with a 4-chlorophenyl group and a primary amine group).

[1121] 2-((2,6-Dichloropyrimidin-4-yl)methyl)isoindoline-1,3-dione (4): A suspension of 2,4-dichloro-6-(chloromethyl)pyrimidine (**2**) (1.5 g, 7.60 mmol) and potassium 1,3-dioxoisoindolin-2-ide (**3**) (1.55 g, 8.36 mmol) in DMF (20 mL) was stirred at 20°C for 2 hrs. LCMS showed the reaction was completed. The reaction was quenched with water (60 mL)

and extracted with EtOAc (40 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by R-MPLC to obtain 2-((2,6-dichloropyrimidin-4-yl)methyl)isoindoline-1,3-dione (**4**) (1.5 g, yield: 64.1 %) as yellow solid. ¹H NMR (400MHz, DMSO-d₆) δ = 8.00 (s, 1H), 7.96-7.92 (m, 2H), 7.92-7.87 (m, 2H), 4.92 (s, 2H).

[1122] 2-((2-Chloro-6-vinylpyrimidin-4-yl)methyl)isoindoline-1,3-dione (5): A suspension of 2-((2,6-dichloropyrimidin-4-yl)methyl)isoindoline-1,3-dione (**4**) (1.5 g, 4.87 mmol), potassium;trifluoro(vinyl)boranuide (0.587 g, 4.38 mmol), TEA (0.746 mL, 5.36 mmol) and Pd(dppf)Cl₂ (0.249 g, 0.34 mmol) in n-PrOH (80 mL) was stirred at 100°C for 2 hrs under N₂ atmosphere. LCMS showed the reaction was completed. Then the resulting mixture was quenched with saturated aqueous NH₄Cl (50 mL) and extracted with EtOAc (30 mL x3). The combined organic layers were washed with brine (80 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to obtain 2-((2-chloro-6-vinylpyrimidin-4-yl)methyl)isoindoline-1,3-dione (**5**) (~80.0 % in HNMR, 0.8 g) as yellow solid. ¹H NMR (400MHz, DMSO-d₆) δ = 7.96-7.93 (m, 2H), 7.92-7.88 (m, 2H), 7.72 (s, 1H), 6.82-6.73 (m, 1H), 6.55 (dd, *J* = 1.0, 17.4 Hz, 1H), 5.83 (dd, *J* = 1.0, 10.7 Hz, 1H), 4.90 (s, 2H).

[1123] 2-Chloro-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidine-4-carbaldehyde (6): To a solution of 2-((2-chloro-6-vinylpyrimidin-4-yl)methyl)isoindoline-1,3-dione (**5**) (80.0 %, 0.7 g, 1.87 mmol) in DCM (20 mL) was bubbled with O₃ at -78°C for 15 min. Then the solution was bubbled with O₂ for 10 min at same temperature until the reaction turned colorless. Me₂S (2 mL) was added at -78°C and the mixture was stirred at 20°C for 1 hr. TLC showed the reaction was completed. The resulting mixture was quenched with water (20 mL) and extracted with DCM (10 mL x 2). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to give 2-chloro-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidine-4-carbaldehyde (**6**) (470 mg, yield: 83.4 %) as yellow solid. ¹H NMR (400MHz, DMSO-d₆) δ = 9.96 (s, 1H), 7.95 (dd, *J* = 3.2, 5.4 Hz, 2H), 7.82 (dd, *J* = 2.8, 5.6 Hz, 2H), 7.63 (s, 1H), 5.08 (s, 2H).

[1124] 2-((2-Chloro-6-(hydroxymethyl)pyrimidin-4-yl)methyl)isoindoline-1,3-dione (7): To a solution of 2-chloro-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidine-4-carbaldehyde (**6**) (490 mg, 1.62 mmol) in THF (10 mL) was added NaBH₄ (24.7 mg, 0.65 mmol) in water (0.5 mL) dropwise at 0°C. The reaction was stirred at same temperature for 15 min. LCMS showed the reaction was completed. The residue was partitioned between EtOAc (10 mL) and water

(10mL). The aqueous layer was extracted with EtOAc (5 mL x 3). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 2-((2-chloro-6-(hydroxymethyl)pyrimidin-4-yl)methyl)isoindoline-1,3-dione (**7**) (430 mg, yield: 78.5%) as light yellow solid. ¹H NMR (400MHz, DMSO-d₆) δ = 7.98-7.94 (m, 2H), 7.93-7.89 (m, 2H), 7.61 (s, 1H), 5.76 (t, *J*=5.9 Hz, 1H), 4.95 (s, 2H), 4.54 (d, *J*=5.9 Hz, 2H).

[1125] (2-Chloro-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-4-yl)methyl

methanesulfonate (8): To a solution of 2-((2-chloro-6-(hydroxymethyl)pyrimidin-4-yl)methyl)isoindoline-1,3-dione (**7**) (410 mg, 1.35 mmol) and TEA (0.3 mL, 2.03 mmol) in DCM (10 mL) was added MsCl (170 mg, 1.49 mmol) at 0°C. The reaction was stirred at same temperature for 20 min. TLC showed the reaction was completed. The reaction was quenched with saturated aqueous NH₄Cl (10 mL) and extracted with DCM (10 mL x 2). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give (2-chloro-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-4-yl)methylmethanesulfonate (**8**) (500 mg, yield: 97.0 %) as yellow oil, which was used for next step directly. ¹H NMR (400MHz, CDCl₃) δ = 7.94 (dd, *J*=2.8, 5.4 Hz, 2H), 7.83-7.79 (m, 2H), 7.35 (s, 1H), 5.25 (s, 2H), 5.01 (s, 2H), 3.15 (s, 3H).

[1126] 3-Chloro-5-(2-(4-((2-chloro-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-4-yl)methoxy) phenyl)propan-2-yl)-2-(2-chloroethoxy)benzonitrile (9):

A suspension of (2-chloro-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-4-yl)methyl methanesulfonate (**8**) (500 mg, 1.31 mmol), 3-chloro-2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (**int H**) (436 mg, 1.24 mmol) and Cs₂CO₃ (853 mg, 2.62 mmol) in DMF (10 mL) was stirred at 20°C for 16 hrs. Then the mixture was stirred at 40°C for 0.5 hrs. LCMS showed the reaction was completed. The resulting mixture was quenched with saturated aqueous NH₄Cl solution (10 mL) and water (10 mL). The aqueous layer was extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to obtain 3-chloro-5-(2-(4-((2-chloro-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-4-yl)methoxy)phenyl)propan-2-yl)-2-(2-chloroethoxy)benzonitrile (**9**) (300 mg, 36.0 %) as yellow solid. ¹H NMR (400MHz, CDCl₃) δ = 7.92 (dd, *J* = 3.0, 5.4 Hz, 2H), 7.80 (dd, *J* = 3.0, 5.4 Hz, 2H), 7.46-7.43 (m, 2H), 7.33 (d, *J* = 2.4 Hz, 1H), 7.11 (d, *J* = 8.8 Hz, 2H), 6.86 (d, *J* = 8.8 Hz, 2H), 5.11 (s, 2H), 4.99 (s, 2H), 4.43 (t, *J* = 6.2 Hz, 2H), 3.88 (t, *J* = 6.2 Hz, 2H), 1.65 (s, 6H).

[1127] N-(4-((4-(2-(3-Chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-

yl)phenoxy)methyl)-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-


yl)methanesulfonamide (**10**): A suspension of N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)methanesulfonamide (**10**) (180 mg, 0.28 mmol), methanesulfonamide (80.8 mg, 0.85 mmol), Cs₂CO₃ (184 mg, 0.57 mmol), Xantphos (32.8 mg, 0.056 mmol) and Pd₂(dba)₃ (25.9 mg, 0.028 mmol) in dioxane (6 mL) was stirred at 100°C for 2 hrs. TLC showed the reaction was completed. The resulting mixture was cooled down, quenched with saturated aqueous NH₄Cl (5 mL) and water (5 mL). The aqueous layer was extracted with EtOAc (5 mL x 3). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC to obtain N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)methanesulfonamide (**10**) (150 mg, yield: 76.3 %) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ = 7.92 (dd, *J* = 3.2, 5.4 Hz, 2H), 7.79 (dd, *J* = 3.2, 5.4 Hz, 2H), 7.45 (d, *J* = 2.2 Hz, 1H), 7.34 (d, *J* = 2.6 Hz, 1H), 7.25 (s, 1H), 7.12 (d, *J* = 8.8 Hz, 2H), 6.89 (d, *J* = 8.8 Hz, 2H), 5.05 (s, 2H), 4.96 (s, 2H), 4.43 (t, *J* = 6.0 Hz, 2H), 3.88 (t, *J* = 6.0 Hz, 2H), 3.25 (s, 3H), 1.66 (s, 6H).

[1128] N-(4-(Aminomethyl)-6-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-

cyanophenyl)propan-2-yl)phenoxy)methyl)pyrimidin-2-yl)methanesulfonamide

(**A164**): A solution of N-(4-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)-6-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)methanesulfonamide (**10**) (50 mg, 0.072 mmol) and NH₂NH₂·H₂O (80.0 %, 0.0218 mL, 0.36 mmol) in MeOH (1 mL) and DCM (0.5 mL) was stirred at 20°C for 1 hr. LCMS showed the reaction was completed. The resulting mixture was adjusted pH to 4~5 by adding aqueous HCl (1 N) and the residue was purified by prep-HPLC (HCl) to obtain N-(4-(aminomethyl)-6-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)pyrimidin-2-yl)methanesulfonamide (**A164**) (5.54 mg, yield: 13.6 %) as white solid. ¹H NMR (400MHz, DMSO-d₆) δ = 11.52 (s, 1H), 8.45 (s, 3H), 7.63 (s, 1H), 7.58 (s, 1H), 7.39 (s, 1H), 7.20 (d, *J* = 8.4 Hz, 2H), 6.98 (d, *J* = 8.8 Hz, 2H), 5.14 (s, 2H), 4.41 (t, *J* = 4.8 Hz, 2H), 4.22 (s, 2H), 3.95 (t, *J* = 4.8 Hz, 2H), 3.42 (s, 3H), 1.63 (s, 6H). LCMS (220 nm): 99.7%. Exact Mass : 563.12; found 564.1.

[1129] **Example 30:** Synthesis of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((5-(methylsulfonyl)furan-2-yl) methoxy)phenyl)propan-2-yl)benzonitrile (**A168**)



[1130] Methyl 5-(methylsulfonyl)furan-2-carboxylate (2): A suspension of methyl 5-nitrofuran-2-carboxylate (1.0 g, 4.9 mmol), sodium methanesulfonate (115 mg, 9.8 mmol) and CuI (139 mg, 7.3 mmol) in DMSO (30 mL) was stirred at 100°C for 6 hrs under N_2 atmosphere. TLC showed the reaction was completed. The mixture was cooled down. The reaction was diluted with EtOAc (30 mL) and quenched with water (70 mL). The mixture was filtered, and the filtrate was extracted with EtOAc (10 mL). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by MPLC to give methyl 5-(methylsulfonyl)furan-2-carboxylate (0.3 g, yield: 30.1 %) as brown oil. ^1H NMR (400MHz, CDCl_3) δ = 7.25 (d, J =3.4 Hz, 1H), 7.22 (d, J =3.4 Hz, 1H), 3.96 (s, 3H), 3.23 (s, 3H).


[1131] (5-(Methylsulfonyl)furan-2-yl)methanol (3): To a solution of methyl 5-(methylsulfonyl)furan-2-carboxylate (0.3 g, 1.5 mmol) in THF (3 mL) was added LiBH_4 (128 mg, 5.9 mmol) under N_2 atmosphere at 20°C . The mixture was stirred at 50°C for 1 hr. TLC showed the reaction was completed. The mixture was poured into saturated aqueous NH_4Cl (5 mL) and extracted with EtOAc (3 mL x 3). The combined organic layers were washed with brine (3 mL x 2), dried over Na_2SO_4 , filtered, and concentrated under reduced pressure to give (5-(methylsulfonyl)furan-2-yl)methanol (220 mg, yield: 85.0 %) as brown oil. ^1H NMR (400MHz, CDCl_3) δ = 7.14 (d, J =3.4 Hz, 1H), 6.48 (d, J =3.4 Hz, 1H), 4.72 (d, J =6.0 Hz, 2H), 3.17 (s, 3H), 1.97 (t, J =6.4 Hz, 1H).

[1132] (5-(Methylsulfonyl)furan-2-yl)methyl methanesulfonate (4): To a solution of (5-(methylsulfonyl)furan-2-yl)methanol (110 mg, 0.6 mmol) in DCM (2 mL) was added TEA (0.18 g, 1.87 mmol) and MsCl (107 mg, 0.936 mol) at 0°C under N_2 atmosphere. The mixture was stirred at 20°C for 30 min. TLC showed the reaction was completed. The mixture was poured into saturated aqueous NH_4Cl solution (2 mL). The mixture was extracted with DCM (2 mL x 3). The combined organic layers were washed with brine (1 mL x 2), dried over

Na₂SO₄, filtered, and concentrated under reduced pressure to give (5-(methylsulfonyl)furan-2-yl)methyl methanesulfonate (80.0 %, 150 mg, yield: 75.6 %) as light yellow solid. ¹H NMR (400MHz, CDCl₃) δ = 7.17 (d, *J*=3.4 Hz, 1H), 6.70 (d, *J*=3.4 Hz, 1H), 5.25 (s, 2H), 3.19 (s, 3H), 3.07 (s, 3H)

[1133] 3-Chloro-2-(2-chloroethoxy)-5-(2-(4-((5-(methylsulfonyl)furan-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (A168): To a solution of (5-(methylsulfonyl)furan-2-yl)methyl methanesulfonate (150 mg, 0.6 mmol) in DMF (2.00 mL) was added 3-chloro-2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (207 mg, 0.6mmol) and Cs₂CO₃ (384 mg, 1.2 mmol) under N₂ atmosphere at 20°C. The mixture was stirred at 20°C for 16 hrs. LCMS showed the reaction was completed. The mixture was poured into H₂O (5 mL). The mixture was extracted with EtOAc (3 mL x 2). The combined organic layers were washed with brine (2 mL x 4), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by prep-HPLC (FA) to give 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((5-(methylsulfonyl)furan-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (**A168**) (62 mg, yield: 20.7%) as white solid. ¹H NMR (400MHz, CDCl₃) δ = 7.44 (d, *J*=2.4 Hz, 1H), 7.32 (d, *J*=2.4 Hz, 1H), 7.17 (d, *J*=3.2 Hz, 1H), 7.13 (d, *J*=8.8 Hz, 2H), 6.90 (d, *J*=8.8 Hz, 2H), 6.59 (d, *J*=3.2 Hz, 1H), 5.07 (s, 2H), 4.43 (t, *J*=6.0 Hz, 2H), 3.88 (t, *J*=6.0 Hz, 2H), 3.18 (s, 3H), 1.65 (s, 6H). LCMS: (220 nm): 32.43%. Exact Mass: 507.1; found 508.1.

[1134] Example 31: Synthesis of (R)-N-(4-(1-(4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)ethyl)pyrimidin-2-yl)methanesulfonamide (**A169**) and (S)-N-(4-(1-(4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)ethyl)pyrimidin-2-yl)methanesulfonamide (**A170**)



[1135] 2-Chloro-4-(1-ethoxyvinyl)pyrimidine (2): To a solution of 2, 4-dichloropyrimidine (**1**) (20.0 g, 0.134 mol) in DMF (250 mL) was added tributyl(1-ethoxyvinyl)stannane (53.3 g, 0.148 mol), Pd(PPh₃)₄ (7.7 g, 0.0067 mol) and KF (23.4 g, 0.403 mol) at 20°C under N₂. The mixture was stirred at 100°C for 2 hrs under N₂ atmosphere. TLC showed the reaction was completed. The mixture was poured into water (500 mL). The reaction mixture was filtered and the filter cake was washed with 200 mL water, dried in vacuum to give 2-chloro-4-(1-ethoxyvinyl)pyrimidine (**2**) (25.0 g, crude) as yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.79 (d, *J*=5.0 Hz, 1 H), 7.64 (d, *J*=5.0 Hz, 1 H), 5.54 (d, *J*=2.2 Hz, 1 H), 4.72 (d, *J*=2.2 Hz, 1 H), 3.95 (q, *J*=6.8 Hz, 2 H), 1.34 (t, *J*=7.0 Hz, 3 H).

[1136] 1-(2-Chloropyrimidin-4-yl)ethanone (3): To a solution of 2-chloro-4-(1-ethoxyvinyl) pyrimidine (**2**) (25 g, crude) in MeCN (100 mL) was added HCl (10.0 %, 100 mL) at 20°C. The mixture was stirred at 75°C for 1 hour. TLC showed the reaction was completed. The mixture was concentrated to remove MeCN. The mixture was extracted with EtOAc (50 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by MPLC (petroleum ether:EtOAc=10:1 to 5:1) to give 1-(2-chloropyrimidin-4-yl)ethanone (**3**) (11.0 g, over two steps yield: 52.3 %) as white solid. ¹H NMR (400 MHz, CDCl₃) δ ppm 8.79 (d, *J*=4.89 Hz, 1 H), 7.77 (d, *J*=4.89 Hz, 1 H), 2.61-2.68 (m, 3 H).

[1137] 1-(2-Chloropyrimidin-4-yl)ethanol (4): To a solution of 1-(2-chloropyrimidin-4-yl)ethanone (**3**) (11.0 g, 0.0703 mol) in THF (50.0 mL) and H₂O (50.0 mL) was added NaBH₄ (2.92 g, 0.0773 mol) at 0°C. The mixture was stirred for 1 hour at 20°C. TLC showed the

reaction was completed. The mixture was poured into saturated aqueous NH_4Cl (100 mL) and extracted with EtOAc (30 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column (petroleum ether:EtOAc = 10:1 to 5:1) to give 1-(2-chloropyrimidin-4-yl)ethanol (**4**) (5.50 g, yield: 49.4 %) as colorless oil. ^1H NMR (400 MHz, CDCl_3) δ ppm 8.62 (d, $J=5.0$ Hz, 1 H), 7.38 (d, $J=5.0$ Hz, 1 H), 4.84-4.94 (m, 1 H), 2.98 (d, $J=5.0$ Hz, 1 H), 1.55 (d, $J=6.8$ Hz, 2 H), 1.52-1.53 (m, 1 H).

[1138] 1-(2-Chloropyrimidin-4-yl)ethyl methanesulfonate (5): To a solution of 1-(2-chloropyrimidin-4-yl)ethanol (**4**) (14.0 g, 0.0883 mol) and TEA (24.6 mL, 0.177 mol) in DCM (200 mL) was added methanesulfonyl chloride (15.2 g, 0.132 mol) at 0°C . The mixture was stirred at 20°C for 1 hr. LCMS showed the reaction was completed. The mixture was poured into water (400 mL) and extracted with DCM (100 mL x 3). The combined organic layers were washed with brine (200 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give 1-(2-chloropyrimidin-4-yl)ethyl methanesulfonate (**5**) (19.0 g, 0.0803 mol, yield: 90.9 %) as yellow oil.


[1139] 3-Chloro-2-(2-chloroethoxy)-5-(2-(4-(1-(2-chloropyrimidin-4-yl)ethoxy)phenyl)propan-2-yl)benzonitrile (6): To a solution of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (**int H**) (20.0 g, 0.0571 mol) in DMF (250 mL) was added 1-(2-chloropyrimidin-4-yl)ethyl methanesulfonate (**5**) (18.9 g, 0.0799 mol) and Cs_2CO_3 (37.2 g, 0.114 mol) at 20°C . The mixture was stirred at 20°C for 20 hrs and 30°C for 4 hrs. TLC showed the reaction was completed. The mixture was poured into saturated aqueous NH_4Cl solution (200 mL) and water (200 mL). The aqueous layer was extracted with EtOAc (300 mL X 3). The combined organic layers were washed with brine (300 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by MPLC (petroleum ether:EtOAc = 10:1 to 5:1) to give 3-chloro-2-(2-chloroethoxy)-5-(2-(4-(1-(2-chloropyrimidin-4-yl)ethoxy)phenyl)propan-2-yl)benzonitrile (**6**) (14.8 g, 0.0302 mol, yield: 52.8 %) as yellow oil.

^1H NMR (400MHz, CDCl_3) δ = 8.60 (d, $J=5.2$ Hz, 1H), 7.45-7.41 (m, 2H), 7.29 (d, $J=2.4$ Hz, 1H), 7.06 (d, $J=8.8$ Hz, 2H), 6.78 (d, $J=8.8$ Hz, 2H), 5.33 - 5.28 (m, 1H), 4.42 (t, $J=6.2$ Hz, 2H), 3.88 (t, $J=6.2$ Hz, 2H), 1.69 (d, $J=6.6$ Hz, 3H), 1.62 (s, 6H).

[1140] (R)-N-(4-(1-(4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)ethyl)pyrimidin-2-yl)methanesulfonamide (A169) and (S)-N-(4-(1-(4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)ethyl)pyrimidin-2-yl)methanesulfonamide (A170): A suspension of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-(1-

(2-chloropyrimidin-4-yl)ethoxy)phenyl)propan-2-yl)benzonitrile (**6**) (8.00 g, 0.0163 mol), methanesulfonamide (4.65 g, 0.0489 mol), Cs_2CO_3 (10.6 g, 0.0326 mol), Xantphos (1.89 g, 0.00326 mol) and $\text{Pd}_2(\text{dba})_3$ (1.49 g, 0.00163 mol) in dioxane (250 mL) was stirred at 100°C for 2 hrs under N_2 atmosphere. TLC showed the reaction was completed. The resulting mixture was quenched with saturated aqueous NH_4Cl (100 mL) and water (100 mL). The aqueous layer was extracted with EtOAc (100 mL x 3). The combined organic layers were washed with brine (100 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by MPLC to obtain the crude product. The mixture was purified by p-HPLC (FA) and SFC to give (R)-N-(4-(1-(4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)ethyl)pyrimidin-2-yl)methanesulfonamide (1.77 g, yield: 19.8 %, peak 1 in SFC was named as **A169**) ^1H NMR (400MHz, CDCl_3) δ = 8.56 (d, J =5.1 Hz, 1H), 8.35 (br s, 1H), 7.43 (d, J =2.2 Hz, 1H), 7.29 (d, J =2.1 Hz, 1H), 7.17 (d, J =5.1 Hz, 1H), 7.05 (d, J =8.7 Hz, 2H), 6.79 (d, J =8.8 Hz, 2H), 5.23 (q, J =6.7 Hz, 1H), 4.42 (t, J =6.2 Hz, 2H), 3.88 (t, J =6.2 Hz, 2H), 3.49 (s, 3H), 1.67 (d, J =6.6 Hz, 3H), 1.62 (s, 6H). LCMS (220 nm): 98.4%. Exact Mass: 548.1; found 549.1. (S)-N-(4-(1-(4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)ethyl)pyrimidin-2-yl)methanesulfonamide (1.23 g, yield: 13.7 %, peak 2 in SFC was named as **A170**) as light yellow solid. ^1H NMR (400MHz, CDCl_3) δ = 8.69 (br s, 1H), 8.58 (d, J =5.3 Hz, 1H), 7.43 (d, J =1.6 Hz, 1H), 7.29 (d, J =1.6 Hz, 1H), 7.17 (d, J =5.1 Hz, 1H), 7.05 (br d, J =8.6 Hz, 2H), 6.79 (d, J =8.7 Hz, 2H), 5.23 (q, J =6.3 Hz, 1H), 4.42 (t, J =6.1 Hz, 2H), 3.88 (t, J =6.1 Hz, 2H), 3.49 (s, 3H), 1.67 (d, J =6.6 Hz, 3H), 1.62 (s, 6H). LCMS (220 nm): 98.6%. Exact Mass: 548.1; found 549.1.

[1141] Example 32: Synthesis of N-(2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)pyrimidin-4-yl)methanesulfonamide (**A184**)




[1142] 2-(Bromomethyl)-4-chloropyrimidine (2): To a mixture of 4-chloro-2-methylpyrimidine (10.0 g, 7.8 mmol) and AIBN (5.1 g, 31.1 mmol) in CCl₄ (100 mL) was added NBS (16.6 g, 93.3 mmol) at 25°C and the mixture was stirred at 80°C for 16 hrs. LCMS showed the reaction was completed. The reaction mixture was cooled down to room temperature, poured into H₂O (150 mL) and extracted with DCM (80 mL x 2). The organic layers were combined and washed with brine (80 mL x 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give 2-(bromomethyl)-4-chloropyrimidine (1.7 g, yield 10.5%) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ = 8.64 (d, *J* = 5.6 Hz, 1H), 7.29 (d, *J* = 5.2 Hz, 1H), 4.56 (s, 2H).

[1143] 3-Chloro-2-(2-chloroethoxy)-5-(2-(4-((4-chloropyrimidin-2-yl)methoxy)phenyl)propan-2-yl) benzonitrile (4): To a mixture of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl) propan-2-yl)benzonitrile (300 mg, 0.9 mmol) and 2-(bromomethyl)-4-chloropyrimidine (444 mg, 2.1 mmol) in DMF (5 mL) was added NaH (60%, 39 mg, 1.0 mmol) in portions at 0°C under N₂ and the mixture was stirred at 0°C for 4 hrs. LCMS showed the reaction was completed. The reaction was poured into H₂O (10 mL), extracted with EtOAc (8 mL x 3). The combined organic layers were washed with brine (12 mL x 4), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((4-chloropyrimidin-2-yl) methoxy)phenyl)propan-2-yl) benzonitrile (300 mg, Yield: 73.5%) as yellow oil. ¹H NMR (400MHz, CDCl₃) δ = 8.68 (d, *J* = 5.2 Hz, 1H), 7.43 (d, *J* = 2.4 Hz, 1H), 7.33-7.32 (m, 2H), 7.09 (d, *J* = 8.8 Hz, 2H), 6.94 (d, *J* = 9.2 Hz, 2H), 5.27 (s, 2H), 4.42 (t, *J* = 6.4 Hz, 2H), 3.88 (t, *J* = 6.0 Hz, 2H), 1.64 (s, 6H).

[1144] N-(2-((4-(2-(3-Chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl) pyrimidin-4-yl)methanesulfonamide (A184): To a solution of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((4-chloropyrimidin-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (100 mg, 0.2 mmol) in dioxane (2 mL) was added methanesulfonamide (60 mg, 6.3 mmol), Xantphos (24.3 mg, 0.06 mmol), Cs₂CO₃ (137 mg, 0.4 mmol) and Pd₂(dba)₃ (19.2 mg, 0.02 mmol) at 100°C under N₂ atmosphere. The mixture was stirred at 100°C for 5 hrs. LCMS showed the mixture was completed. The reaction was cooled down to room temperature, quenched with water (4 mL) and extracted with DCM (3 mL x 3). The combined organic layers were washed with brine (6 mL x 2), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude was purified by silica gel column chromatography and prep-HPLC (FA) to give N-(2-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-

cyanophenyl)propan-2-yl)phenoxy)methyl)pyrimidin-4-yl) methanesulfonamide (**A184**) (8.4 mg, yield: 7.84%) as white solid. ¹H NMR (400 MHz, DMSO) δ = 8.31-8.36 (m, 1H), 7.63 (d, J = 2.4 Hz, 1H), 7.55 (d, J = 2.4 Hz, 1H), 7.13 (d, J = 8.8 Hz, 1H), 7.13 (d, J = 8.8 Hz, 2H), 7.06 (s, 1H), 6.93 (d, J = 8.8 Hz, 2H), 5.16 (s, 2H), 4.43 (t, J = 6.4 Hz, 2H), 3.88 (t, J = 6.0 Hz, 2H), 3.39 (s, 3H), 1.61 (s, 6H). LCMS (220 nm): 98.6%. Exact Mass: 534.10; found 535.1.

[1145] Example 33: Synthesis of N-(6-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)pyrazin-2-yl)methanesulfonamide (**A185**)




[1146] 3-Chloro-2-(2-chloroethoxy)-5-(2-(4-((6-chloropyrazin-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (3): To a solution of (6-chloropyrazin-2-yl)methanol (**1**) (0.500 g, 3.46 mmol), 3-chloro-2-(2-chloroethoxy)-5-(1-(4-hydroxyphenyl)-1-methyl-ethyl)benzonitrile (**2**) (0.606 g, 1.73 mmol) and triphenylphosphane (1.09 g, 4.15 mmol) in THF (10.0 mL) was added DIAD (0.817 mg, 4.15 mmol) at 0°C and stirred at 25°C for 3 hrs under N₂. TLC showed the reaction was completed. The mixture was poured into water (20 mL) and extracted with EtOAc (15 mL x 3), the combined organic layers were washed with brine (5 mL x 2), dried over Na₂SO₄, filtered and concentrated to give the crude product which was purified by silica gel column chromatography to give 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((6-chloropyrazin-2-yl)methoxy)phenyl)propan-2-yl)benzonitrile (**3**) (purity: 90.0 %, 0.400 g, yield: 21.8%) as colorless oil. ¹H NMR (400MHz, CDCl₃) δ = 8.75 (s, 1H), 8.58 (s, 1H), 7.44 (d, J =2.4 Hz, 1H), 7.33 (d, J =2.0 Hz, 1H), 7.13 (d, J = 8.8 Hz, 2H), 6.93 (d, J = 8.8 Hz, 2H), 5.22 (s, 2H), 4.43 (t, J =6.4 Hz, 2H), 3.88 (t, J =6.0 Hz, 2H), 1.65 (s, 6H).

[1147] N-(6-((4-(2-(3-Chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl) pyrazin-2-yl)methanesulfonamide (A185): To a solution of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((6-chloropyrazin-2-yl)methoxy)phenyl)propan-2-yl) benzonitrile (**3**) (120 mg, 0.25 mmol) in 1,4-Dioxane (3.00 mL) was added methanesulfonamide (71.8 mg, 0.76 mmol), tris(dibenzylideneacetone)dipalladium(0) (14.5 mg, 0.025 mmol), Xantphos (14.6 mg, 0.025 mmol), and Cs₂CO₃ (0.164g, 0.50 mmol) at

20°C under N₂. The mixture was stirred at 100°C for 3 hrs. LCMS showed the reaction was completed. The reaction was poured into water (10 mL) and extracted with EtOAc (3 mL x 3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (FA) to give

N-((6-((4-(2-(3-chloro-4-(2-chloroethoxy)-5-cyanophenyl)propan-2-yl)phenoxy)methyl)pyrazin-2-yl)methanesulfonamide (**A185**) (38 mg, purity: 96.0%, yield: 27.1%) as white solid. ¹H NMR (400 MHz, CDCl₃) δ = 8.58 (s, 1H), 8.48 (s, 1H), 7.45 (d, *J* = 2.4 Hz, 1H), 7.32 (d, *J* = 2.4 Hz, 1H), 7.14 (d, *J* = 8.8 Hz, 2H), 6.95 (d, *J* = 8.8 Hz, 2H), 5.16 (s, 2H), 4.43 (t, *J* = 6.0 Hz, 2H), 3.88 (t, *J* = 6.4 Hz, 2H), 3.33 (s, 3H), 1.66 (s, 6H). LCMS (220 nm): 96.8%. Exact Mass: 534.09; found 535.0.

[1148] Example 34: Synthesis of 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((2-methoxyoxazol-5-yl)methoxy)phenyl)propan-2-yl)benzonitrile (**A112**)



[1149] (2-Methoxyoxazol-5-yl)methanol (2): To a mixture of ethyl 2-chlorooxazole-5-carboxylate (300 mg, 1.7 mmol) in MeOH (5 mL) was added NaOMe (138 mg, 2.6 mmol) at 25°C and the mixture was stirred at 25°C for 2 hrs. Then NaBH₄ (259 mg, 6.8 mmol) was added to the mixture and stirred at 25°C for another 2 hrs. TLC showed the starting material was consumed and one new spot was observed. The mixture was filtered and concentrated under reduced pressure. The residue was purified by silica-gel column chromatography to give (2-methoxyoxazol-5-yl)methanol (110 mg, yield: 49.9 %) as yellow oil. ¹H NMR (400 MHz, CD₃OD) δ = 6.74 (s, 1H), 4.44 (s, 2H), 4.04 (s, 3H).

[1150] 3-Chloro-2-(2-chloroethoxy)-5-(2-(4-((2-methoxyoxazol-5-yl)methoxy)phenyl)propan-2-yl)benzonitrile (A112**):** To a mixture of (2-methoxyoxazol-5-yl)methanol (35 mg, 0.27 mmol), 3-chloro-2-(2-chloroethoxy)-5-(2-(4-hydroxyphenyl)propan-2-yl)benzonitrile (95 mg, 0.27 mmol) and (C₆H₅)₃P (142 mg, 0.5

mmol) in THF (4 mL) was added DIAD (0.1 mL, 0.5 mmol) at 25°C under N₂ atmosphere. The mixture was stirred at 25°C for 2 hrs. LCMS showed the starting material was consumed and one new peak was observed. The mixture was poured into water (15 mL), extracted with EtOAc (4 mL x 2). The combined organic layers were washed with brine (6 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-HPLC (FA) to give 3-chloro-2-(2-chloroethoxy)-5-(2-(4-((2-methoxyoxazol-5-yl)methoxy)phenyl)propan-2-yl)benzonitrile (A112) (9.05 mg, yield: 7.6 %) as yellow oil. ¹H NMR (400 MHz, CDCl₃) δ = 7.43 (d, *J* = 2.4 Hz, 1H), 7.33 (d, *J* = 2.4 Hz, 1H), 7.11 (d, *J* = 8.8 Hz, 2H), 6.91 (d, *J* = 2.4 Hz, 2H), 6.85 (s, 1H), 4.92 (s, 2H), 4.42 (t, *J* = 6.4 Hz, 2H), 4.08 (s, 3H), 3.88 (t, *J* = 6.0 Hz, 2H), 1.65 (s, 6H). LCMS (220 nm): 98.1%. Exact Mass: 460.1; found: 461.2.

[1151] Characterization of representative compounds synthesized are shown in Table C.

[1152] Table C. Characterization of compounds

Compound ID	¹ H NMR and LCMS
A185	¹ H NMR (400 MHz, CDCl ₃): δ = 8.58 (s, 1H), 8.48 (s, 1H), 7.45 (d, <i>J</i> = 2.4 Hz, 1H), 7.32 (d, <i>J</i> = 2.4 Hz, 1H), 7.14 (d, <i>J</i> = 8.8 Hz, 2H), 6.95 (d, <i>J</i> = 8.8 Hz, 2H), 5.16 (s, 2H), 4.43 (t, <i>J</i> = 6.0 Hz, 2H), 3.88 (t, <i>J</i> = 6.4 Hz, 2H), 3.33 (s, 3H), 1.66 (s, 6H). LCMS (220 nm): 96.8%. Exact Mass: 534.09; found 535.0, 537.0.
A189	¹ H NMR (400 MHz, CDCl ₃): δ = 8.67 (s, 1H), 8.53 (br d, <i>J</i> = 6.0 Hz, 1H), 8.02 (br d, <i>J</i> = 6.0 Hz, 1H), 7.87 (d, <i>J</i> = 8.4 Hz, 2H), 7.38 (d, <i>J</i> = 8.4 Hz, 2H), 7.13 (s, 2H), 4.28 (t, <i>J</i> = 6.0 Hz, 2H), 3.87 (t, <i>J</i> = 6.8 Hz, 2H), 3.42 (s, 3H), 1.69 (s, 6H). LCMS (220 nm): 95.2%. Exact Mass: 556.1; found 557.0/559.0.
A190	¹ H NMR (400 MHz, CDCl ₃): δ = 7.51 (br s, 1H), 7.44 (d, <i>J</i> = 2.0 Hz, 1H), 7.31 (d, <i>J</i> = 2.4 Hz, 1H), 7.10 (d, <i>J</i> = 8.8 Hz, 2H), 6.88 (d, <i>J</i> = 8.8 Hz, 2H), 5.04 (s, 2H), 4.84 (s, 2H), 4.43 (t, <i>J</i> = 6.4 Hz, 2H), 4.03 (t, <i>J</i> = 6.0 Hz, 2H), 3.88 (t, <i>J</i> = 2.0 Hz, 2H), 3.45 (s, 3H), 2.95 (t, <i>J</i> = 5.6 Hz, 2H), 1.64 (s, 6H). LCMS (220 nm): 97.8%. Exact Mass: 590.1; found: 591.1/593.1.
A191	¹ H NMR (400MHz, CDCl ₃) δ = 9.34 (br s, 1H), 7.62 (d, <i>J</i> =2.2 Hz, 1H), 7.57 (d, <i>J</i> =2.2 Hz, 1H), 7.18 (d, <i>J</i> =8.8 Hz, 2H), 6.96 (d, <i>J</i> =8.8 Hz, 2H), 5.17 (s, 2H), 4.41 (t, <i>J</i> =7.2 Hz, 2H), 4.34 (s, 2H), 3.95 (t, <i>J</i> =7.2 Hz, 2H), 3.51 - 3.42 (m, 2H), 3.30 (s, 3H), 3.03 (br s, 2H), 1.63 (s, 6H). LCMS (220 nm): 97.6%. Exact Mass: 589.1; found: 590.1/592.2.
A192	¹ H NMR (400MHz, DMSO-d ₆) δ = 7.61 (d, <i>J</i> =2.2 Hz, 1H), 7.56 (d, <i>J</i> =2.2 Hz, 1H), 7.40 (s, 1H), 7.17 (d, <i>J</i> =8.8 Hz, 2H), 6.96 (d, <i>J</i> =8.8 Hz, 2H), 5.14 (s, 2H), 4.39 (t, <i>J</i> =4.8 Hz, 2H), 4.28 (s, 2H), 3.94 (t, <i>J</i> =4.8 Hz, 2H), 1.62 (s, 6H). LCMS (220 nm): 99.7%. Exact Mass: 459.1; found: 460.1/462.1.
A193	¹ H NMR (400 MHz, CDCl ₃) δ = 7.46 (d, <i>J</i> = 2.0 Hz, 1H), 7.31 (d, <i>J</i> = 2.4 Hz, 1H), 7.13 (d, <i>J</i> = 8.4 Hz, 2H), 6.95 - 6.89 (m, 3H), 6.32 (d, <i>J</i> = 4.0 Hz, 1H), 5.00 (s, 2H), 4.43 (t, <i>J</i> = 6.4 Hz, 2H), 3.94 (s, 3H), 3.88 (t, <i>J</i> = 6.4 Hz,

Compound ID	¹ H NMR and LCMS
	2H), 3.14 (s, 3H), 1.66 (s, 6H). LCMS (220 nm): 99.3%. Exact Mass: 520.10; found 521.1, 523.1.
A194	¹ H NMR (400MHz, CDCl ₃) δ = 7.73 (t, <i>J</i> = 7.6 Hz, 1H), 7.45 (d, <i>J</i> = 2.4 Hz, 1H), 7.32 (d, <i>J</i> = 2.4 Hz, 1H), 7.24 (d, <i>J</i> = 7.6 Hz, 1H), 7.14 - 7.10 (m, 3H), 6.92 (d, <i>J</i> = 8.8 Hz, 2H), 5.09 (s, 2H), 4.42 (t, <i>J</i> = 6.0 Hz, 2H), 3.88 (t, <i>J</i> = 6.8 Hz, 2H), 3.22 (s, 3H), 1.65 (s, 6H). LCMS (220 nm): 94.7%. Exact Mass: 533.1; found 534.1/536.1.
A195	¹ H NMR (400MHz, CDCl ₃) δ = 8.17 (d, <i>J</i> = 5.6 Hz, 1H), 7.44 (d, <i>J</i> = 2.4 Hz, 1H), 7.37 (s, 1H), 7.32 (d, <i>J</i> = 2.4 Hz, 1H), 7.12 (d, <i>J</i> = 8.8 Hz, 2H), 6.98 (d, <i>J</i> = 5.6 Hz, 1H), 6.89 (d, <i>J</i> = 8.8 Hz, 2H), 5.08 (s, 2H), 4.43 (t, <i>J</i> = 6.4 Hz, 2H), 3.88 (t, <i>J</i> = 6.4 Hz, 2H), 3.12 (s, 3H), 1.65 (s, 6H). LCMS: (220 nm): 97.5%. Exact Mass: 533.1; found 534.1/536.1.
A196	¹ H NMR (400MHz, CDCl ₃) δ = 8.48 (br s, 1H), 7.44 (d, <i>J</i> = 2.8 Hz, 1H), 7.31 (d, <i>J</i> = 2.4 Hz, 1H), 7.23 (s, 1H), 7.12 (d, <i>J</i> = 8.8 Hz, 2H), 7.09 (br s, 1H), 6.93 (d, <i>J</i> = 8.8 Hz, 2H), 6.79 (br s, 1H), 5.17 (s, 2H), 4.42 (t, <i>J</i> = 6.0 Hz, 2H), 3.88 (t, <i>J</i> = 6.0 Hz, 2H), 3.12 (s, 3H), 1.65 (s, 6H). LCMS (220 nm): 97% Exact Mass: 533; found 534.1/536.1.
A197	¹ H NMR (400MHz, CDCl ₃) δ = 8.51 (br d, <i>J</i> = 4.6 Hz, 2H), 7.45 (s, 1H), 7.32 (s, 1H), 7.13 (br d, <i>J</i> = 8.6 Hz, 2H), 6.94 (br d, <i>J</i> = 8.2 Hz, 2H), 5.21 (s, 2H), 4.43 (t, <i>J</i> = 6.2 Hz, 2H), 3.88 (t, <i>J</i> = 6.2 Hz, 2H), 3.33 (s, 3H), 1.65 (s, 6H). LCMS: (220 nm): 95.8 Exact Mass: 534.1; found 535.1 / 537.1
A198	¹ H NMR (400MHz, CDCl ₃) δ = 8.67 (d, <i>J</i> = 5.0 Hz, 1H), 7.45 (d, <i>J</i> = 2.4 Hz, 1H), 7.34 (d, <i>J</i> = 5.2 Hz, 1H), 7.32 (d, <i>J</i> = 2.4 Hz, 1H), 7.12 (d, <i>J</i> = 8.8 Hz, 2H), 6.90 (d, <i>J</i> = 8.8 Hz, 2H), 5.13 (s, 2H), 4.43 (t, <i>J</i> = 6.2 Hz, 2H), 4.30 - 4.26 (m, 2H), 3.88 (t, <i>J</i> = 6.2 Hz, 2H), 3.36 - 3.31 (m, 2H), 2.34 (quin, <i>J</i> = 6.2 Hz, 2H), 1.85 (td, <i>J</i> = 5.8, 11.7 Hz, 2H), 1.65 (s, 6H). LCMS: (220 nm): 90.5%. Exact Mass: 574.1; found 575.2/9,577.2.
A199	¹ H NMR (400MHz, CDCl ₃) δ = 8.59 (d, <i>J</i> = 5.0 Hz, 1H), 7.44 (s, 1H), 7.33 (s, 1H), 7.21 (d, <i>J</i> = 5.0 Hz, 1H), 7.11 (br d, <i>J</i> = 8.4 Hz, 2H), 6.89 (br d, <i>J</i> = 8.4 Hz, 2H), 5.11 (s, 2H), 4.43 (t, <i>J</i> = 6.4 Hz, 2H), 4.07 (t, <i>J</i> = 6.4 Hz, 2H), 3.88 (t, <i>J</i> = 6.4 Hz, 2H), 3.48 (t, <i>J</i> = 7.2 Hz, 2H), 2.52 (br t, <i>J</i> = 6.8 Hz, 2H), 1.65 (s, 6H). LCMS: (220 nm): 95.2%. Exact Mass: 560.1; found 561.1 / 563.1
A200	¹ H NMR (400 MHz, CDCl ₃) δ = 8.88 (d, <i>J</i> = 5.2 Hz, 1H), 7.82 (d, <i>J</i> = 5.2 Hz, 1H), 7.45 (d, <i>J</i> = 2.4 Hz, 1H), 7.30 (d, <i>J</i> = 2.4 Hz, 1H), 7.14 (d, <i>J</i> = 6.8 Hz, 2H), 6.90 (d, <i>J</i> = 8.8 Hz, 2H), 5.21 (s, 2H), 4.43 (t, <i>J</i> = 6.0 Hz, 2H), 3.88 (t, <i>J</i> = 6.4 Hz, 2H), 1.66 (s, 6H). LCMS (220 nm): 91.1%. Exact Mass: 466.1; found 467.0, 469.0.
A201	¹ H NMR (400MHz, CDCl ₃) δ = 8.80 (brs, 1H), 8.64 (d, <i>J</i> = 5.2 Hz, 1H), 7.45 (d, <i>J</i> = 2.4 Hz, 1H), 7.32 (d, <i>J</i> = 2.4 Hz, 1H), 7.29 (d, <i>J</i> = 5.2 Hz, 1H), 5.11 (s, 2H), 4.43 (t, <i>J</i> = 6.4 Hz, 2H), 3.88 (t, <i>J</i> = 5.6 Hz, 2H), 3.47 (s, 3H), 1.65 (s, 6H). LCMS: (220 nm): 98.53%. Exact Mass: 538.1; found 539.2/541.1
A202	¹ H NMR (400MHz, CDCl ₃) δ = 7.66 (br s, 1H), 7.44 (d, <i>J</i> = 2.4 Hz, 1H), 7.31 (d, <i>J</i> = 2.4 Hz, 1H), 7.10 (d, <i>J</i> = 8.8 Hz, 2H), 6.88 (d, <i>J</i> = 8.8 Hz, 2H), 5.08 (s, 2H), 4.43 (t, <i>J</i> = 6.0 Hz, 2H), 3.88 (t, <i>J</i> = 6.4 Hz, 2H), 3.45 (s, 3H), 3.02 - 2.93 (m, 4H), 2.16 - 2.12 (m, 2H), 1.64 (s, 6H). LCMS (220 nm): 92.0%. Exact Mass: 574.1; found 575.1/577.1
A203	¹ H NMR (400MHz, CDCl ₃) δ = 8.56 (s, 1H), 8.48 (br s, 1H), 7.45 (d, <i>J</i> = 2.4 Hz, 1H), 7.34 (d, <i>J</i> = 1.6 Hz, 1H), 7.16 - 7.11 (m, 2H), 7.10 - 7.05 (m, 2H),

Compound ID	¹ H NMR and LCMS
	5.70 - 5.59 (m, 1H), 4.43 (t, <i>J</i> =6.4 Hz, 2H), 3.89 (t, <i>J</i> =6.4 Hz, 2H), 3.31 (s, 3H), 3.19 - 3.07 (m, 1H), 2.96 - 2.85 (m, 1H), 2.72 - 2.59 (m, 1H), 2.41 - 2.27 (m, 1H), 1.66 (s, 6H) LCMS: (220 nm): 97.7% Exact Mass: 560.1; found 561.1/563.1
A204	¹ H NMR (400MHz, CDCl ₃) δ = 10.14 (br s, 1H), 8.74 (s, 2H), 7.45 (d, <i>J</i> =2.2 Hz, 1H), 7.33 (d, <i>J</i> =2.2 Hz, 1H), 7.14 (d, <i>J</i> =8.8 Hz, 2H), 6.91 (d, <i>J</i> =8.8 Hz, 2H), 5.02 (s, 2H), 4.43 (t, <i>J</i> =6.4 Hz, 2H), 3.88 (t, <i>J</i> =6.4 Hz, 2H), 3.49 (s, 3H), 1.66 (s, 6H) LCMS: (220 nm): 98.6% Exact Mass: 534.1; found 535.0/ 537.0
A205	¹ H NMR (400MHz, CDCl ₃) δ = 8.61 (d, <i>J</i> =5.2 Hz, 1H), 7.27-7.29 (m, 1H), 7.17-7.19 (m, 1H), 7.13- 7.14 (m, 1H), 7.12 (d, <i>J</i> =8.8 Hz, 1H), 6.89 (d, <i>J</i> =8.8 Hz, 1H), 5.10 (s, 2H), 4.48 (t, <i>J</i> =6.0 Hz, 2H), 3.83 (t, <i>J</i> =5.6 Hz, 2H), 3.48 (s, 3H), 1.64 (s, 6H) LCMS: (220 nm): 99.74%. Exact Mass: 518.1; found 519.1/521.1
A206	¹ H NMR (400MHz, CDCl ₃) δ = 8.62 (d, <i>J</i> =5.2 Hz, 1H), 8.60 (brs, 1H), 7.29 (d, <i>J</i> =5.2 Hz, 1H), 7.15 (d, <i>J</i> =8.8 Hz, 1H), 7.02 (d, <i>J</i> =2.0 Hz, 1H), 6.89 (d, <i>J</i> =9.2 Hz, 1H), 6.86 - 6.83 (dd, <i>J</i> =12.4 Hz, 2.0 Hz, 1H), 5.10 (s, 2H), 4.31 (t, <i>J</i> =6.0 Hz, 2H), 3.81 (t, <i>J</i> =6.0 Hz, 2H), 3.47 (s, 3H), 1.63 (s, 6H) LCMS: (220 nm): 99.51%. Exact Mass: 527.1; found 528.0/529.9
A207	¹ H NMR (400MHz, CDCl ₃) δ = 7.63 (d, <i>J</i> =0.8 Hz, 1H), 7.46 (d, <i>J</i> =2.4 Hz, 1H), 7.30 (d, <i>J</i> =2.4 Hz, 1H), 7.16 - 7.07 (m, 3H), 6.89 (d, <i>J</i> =8.8 Hz, 2H), 5.01 - 4.96 (m, 1H), 4.96 (br s, 1H), 4.95 (s, 2H), 4.43 (t, <i>J</i> =6.4 Hz, 2H), 3.88 (t, <i>J</i> =6.4 Hz, 2H), 1.65 (s, 6H) LCMS: (220 nm): 99.2% Exact Mass: 508.1; found 509.0/ 511.0
A208	¹ H NMR (400MHz, CDCl ₃) δ = 7.71 (s, 1H), 7.45 (d, <i>J</i> =2.4 Hz, 1H), 7.32 (d, <i>J</i> =2.2 Hz, 1H), 7.26 (br s, 1H), 7.13 (d, <i>J</i> =8.8 Hz, 2H), 6.89 (d, <i>J</i> =8.8 Hz, 2H), 4.96 (s, 2H), 4.43 (t, <i>J</i> =6.2 Hz, 2H), 3.88 (t, <i>J</i> =6.2 Hz, 2H), 3.18 (s, 3H), 1.66 (s, 6H) LCMS: (220 nm): 99.7% Exact Mass: 507.1; found 508.1/510.1
A209	¹ H NMR (400 MHz, CDCl ₃) δ = 8.53 (d, <i>J</i> = 4.8 Hz, 1H), 7.44 (d, <i>J</i> = 2.4 Hz, 1H), 7.32 (d, <i>J</i> = 2.4 Hz, 1H), 7.21 (d, <i>J</i> = 5.2 Hz, 1H), 7.11 (d, <i>J</i> = 8.8 Hz, 2H), 6.89 (d, <i>J</i> = 8.8 Hz, 2H), 5.09 (s, 2H), 4.42 (t, <i>J</i> = 6.0 Hz, 2H), 3.88 (t, <i>J</i> = 6.0 Hz, 2H), 2.59 (s, 3H), 1.65 (s, 6H).
A210	¹ H NMR (400MHz, CDCl ₃) δ = 7.79 (s, 1H), 7.70 (s, 1H), 7.47 (d, <i>J</i> =1.8 Hz, 1H), 7.34 (d, <i>J</i> =1.8 Hz, 1H), 7.14 (d, <i>J</i> =8.8 Hz, 2H), 6.91 (d, <i>J</i> =8.6 Hz, 2H), 5.07 (s, 2H), 4.44 (t, <i>J</i> =6.2 Hz, 2H), 3.90 (t, <i>J</i> =6.2 Hz, 2H), 3.23 (s, 3H), 1.67 (s, 6H) LCMS: (220 nm): 99.4% Exact Mass: 523.1; found 524.1/526.1
A211	¹ H NMR (400MHz, CDCl ₃) δ = 7.72 (s, 1H), 7.58 (s, 1H), 7.46 (d, <i>J</i> =2.2 Hz, 1H), 7.31 (d, <i>J</i> =2.2 Hz, 1H), 7.12 (d, <i>J</i> =8.8 Hz, 2H), 6.89 (d, <i>J</i> =8.8 Hz, 2H), 5.03 (s, 2H), 4.99 (s, 2H), 4.43 (t, <i>J</i> =6.2 Hz, 2H), 3.88 (t, <i>J</i> =6.2 Hz, 2H), 1.65 (s, 6H). LCMS: (220 nm): 99.4% Exact Mass: 524.0; found 524.9/522.8

[1153] Biological Assays

[1154] Example 35: Activity of Exemplary Compounds in Cellular Assays


[1155] The PSA (6.1kb)-luciferase reporter contains functional AREs (androgen response elements) to which AR binds in response to androgen to induce luciferase activity. LNCaP

cells were transiently transfected with the PSA (6.1 kb)-luciferase reporter for 24 h, and then pretreated with vehicle (DMSO) or indicated concentration of representative compounds for one hour prior to the addition of synthetic androgen, R1881 (1 nM). After 24 h of incubation with R1881, the cells were harvested, and relative luciferase activities were determined (Fig 1). To determine the IC₅₀, treatments were normalized to vehicle control activity induced by R1881 (Table 1A).

[1156] Luciferase Assay: Cells were lysed using Passive Lysis Buffer (Promega) and then collected into V-bottom 96-well tissue culture plates. Lysates were centrifuged at 4°C for 5 minutes at 3000 rpm. To measure luminescence of LNCaP cell lysates the Firefly Luciferase Assay System (Promega) was employed, according to manufacturer's protocol. Relative luminescence units (RLU) in cell lysates were detected for 10 seconds using Promega GloMax-Multi Detection Luminometer (Promega). Values were normalized to protein content. GraphPad Prism graphing software was used to calculate IC₅₀ values.

[1157] Statistical analyses were performed using GraphPad Prism (Version 6.01 for Windows; La Jolla, CA, USA). Comparisons between treatment and control groups were compared using Two-Way ANOVA with post-hoc Dunnett's and Tukey's tests. Differences were considered statistically significant at P values less than 0.05.

[1158] Table 1A shows IC₅₀ ranges of Compounds from Tables A and B from the luciferase assay. Fig. 1 shows dose dependent curve of representative compounds shown in Table 1A. Compound X and EPI-002 have the following structures:



[1159] The PSA-Luc % inhibition IC₅₀ values of the representative compounds are shown in Table 1B.

[1160] LBD-Expressing Reporter Assay: To assess if the representative compounds (Compounds A13, A28, A38, A74, A93, and A109) compete with ligand (agonist) for the ligand-binding domain (LBD) of steroid hormone receptors (which would be off-target), the GeneBLAzer assay was employed (Thermo Fisher). This assay uses cells that stably express a chimera fusion protein comprising the LBD of a steroid hormone receptor fused to the DNA-binding domain (DBD) of GAL4 along with beta-lactamase reporter under transcriptional control of an upstream activator sequence (UAS). When agonist binds to the LBD of the GAL4

(DBD)-steroid hormone (LBD) chimera fusion protein, the protein binds to the UAS, resulting in expression of beta-lactamase. LBDs of androgen receptor (AR), progesterone receptor-beta (PR β), glucocorticoid receptor (GR), estrogen receptor- α (ER α), and estrogen receptor- β (ER β) were tested and IC₅₀s calculated for inhibiting maximal activity attained with agonist. See Table 1C.

[1161] Cell Proliferation Assay: Cell proliferation/viability was measured in LNCaP and PC3 cells with Alamar blue, and proliferation was measured in LN-CaP95 cells with BrdU incorporation. In LNCaP cells, AR specific proliferation is calculated by measuring the difference between control cells treated with or without 0.1 nM R1881. See Table 1D and Fig. 7.

[1162] AR-V7 Transcriptional Activity Assay: Transcriptional activity of AR-V7 was measured in LNCaP cells transiently cotransfected with an expression vector for AR-V7 and the PSA-luciferase reporter and incubated with the representative compounds in the presence or absence of synthetic androgen, R1881. As a control, LNCaP cells were transfected with pcDNA plasmid (absent of AR-V7 insert) with or without R1881. See Figs. 2A-2B.

[1163] The results from this Example supports that the compounds of the present disclosure are selective to AR NTD.

[1164] **Table 1A.** IC₅₀ Ranges of Various Compounds in Androgen-Induced PSA-Luciferase Assay

Compound ID	PSA-Luc IC ₅₀ Range	Compound ID	PSA-Luc IC ₅₀ Range
A1	N/A	A112	***
A2	N/A	A113	*
A3	**	A114	**
A4	N/A	A115	***
A5	**	A116	**
A6	N/A	A117	**
A7	*	A118	**
A8	N/A	A119	***
A9	N/A	A120	*
A10	N/A	A121	**
A11	N/A	A122	***
A12	N/A	A123	***
A13	**	A124	*
A14	N/A	A125	***
A15	N/A	A126	**
A16	N/A	A127	N/A
A17	**	A128	N/A

A18	*	A129	N/A
A19	N/A	A130	**
A20	N/A	A131	**
A21	N/A	A132	**
A22	*	A133	***
A23	*	A134	**
A24	**	A135	***
A25	*	A136	**
A26	N/A	A137	***
A27	*	A138	N/A
A28	***	A139	N/A
A29	***	A140	***
A30	**	A141	***
A31	***	A142	***
A32	*	A143	*
A33	N/A	A144	N/A
A34	**	A145	**
A35	**	A146	N/A
A36	N/A	A147	*
A37	N/A	A148	**
A38	**	A149	***
A39	N/A	A150	N/A
A40	**	A151	**
A41	*	A152	N/A
A42	**	A153	**
A43	N/A	A154	N/A
A44	N/A	A155	***
A45	**	A156	***
A46	N/A	A157	N/A
A47	N/A	A158	N/A
A48	N/A	A159	N/A
A49	*	A160	N/A
A50	**	A161	N/A
A51	N/A	A162	***
A52	*	A163	N/A
A53	*	A164	**
A54	**	A165	N/A
A55	N/A	A166	N/A
A56	**	A167	N/A
A57	***	A168	***
A58	*	A169	***
A59	N/A	A170	**
A60	N/A	A171	**

A61	N/A	A172	***
A62	*	A173	*
A63	**	A174	**
A64	*	A175	N/A
A65	**	A176	N/A
A66	**	A177	N/A
A67	N/A	A178	N/A
A68	***	A179	N/A
A69	N/A	A180	N/A
A70	***	A181	N/A
A71	**	A182	N/A
A72	N/A	A183	N/A
A73	**	A184	*
A74	**	A185	***
A75	**	A186	N/A
A76	**	A187	N/A
A77	**	A188	N/A
A78	**	A189	**
A79	***	A190	**
A80	***	A191	**
A81	***	A192	***
A82	N/A	A193	**
A83	N/A	A194	**
A84	N/A	A195	***
A85	***	A196	**
A86	***	A197	**
A87	**	A198	***
A88	**	A199	***
A89	**	A200	**
A90	**	A201	**
A91	**	A202	**
A92	***	A203	**
A93	***	A204	***
A94	**	A205	**
A95	*	A206	**
A96	*	A207	***
A97	***	A208	***
A98	**	A209	N/A
A99	**	A210	N/A
A100	***	A211	N/A
A101	**	B1	**
A102	***	B2	***
A103	***	B3	**

A104	***	B4	N/A
A105	*	B5	**
A106	***	B6	*
A107	*	B7	*
A108	**	B8	**
A109	**	B9	***
A110	N/A	B10	*
A111	***		

Note: *** represent IC₅₀ < 500 nM, ** represents IC₅₀ in the range of 500-2000 nM, * represents IC₅₀ in the range of ≥ 2000 nM

[1165] Table 1B. IC₅₀ of Representative Compounds in Androgen-Induced PSA-Luciferase Assay

Compound ID	PSA-Luc IC ₅₀ (nM)	n
X	1054	3
A13	592	8
A28	400	5
A38	631	6
A74	658	6
A93	205	4
A109	535	2
EPI-002	9580	2
Enzalutamide	189	8
Bicalutamide	306	2

[1166] Table 1C. IC₅₀ of Representative Compounds in LBD-Expressing Reporter Cell Lines

Compound ID	Steroid Receptor LBD IC ₅₀ (μM)				
	AR	ERα	ERβ	PR	GR
X	9.83	5.82	7.37	4.96	>10
A13	>10	>10	>10	0.97	>10
A28	>10	>10	>10	0.93	>10
A38	>10	>10	>10	5.07	>10
A74	>10	>10	>10	5.41	9.51
A93	>10	>10	>10	5.04	>10
A109	>10	>10	>10	5.61	>10
EPI-002	>10	>10	>10	>10	>10
Enzalutamide	0.54	>10	>10	4.49	>10

[1167] Table 1D. IC₅₀ of Representative Compounds on Cell Proliferation/Viability

Compound ID	Cellular Proliferation/Viability IC ₅₀ (μM)		
	LNCaP	PC-3	LNCaP95
X	3.00	>10	4.00
A13	0.85	>10	4.00

Compound ID	Cellular Proliferation/Viability IC ₅₀ (μM)		
	LNCaP	PC-3	LNCaP95
A28	0.68	>10	6.43
A74	1.16	>10	2.65
A109	0.44	>10	3.78
EPI-002	9.00	>10	~20
Enzalutamide	0.35	>10	>10

[1168] Example 36: Stability Assays

[1169] Microsomal Stability Assay: Microsomal stability assay is a widely used in vitro model to characterize the metabolic conversion by phase I enzymes, such as cytochrome P450 (CYP) enzymes. Since metabolism is known to be highly variable in different species, microsomal stability assay is commonly run in multiple species. Metabolic stability of testing compound can be evaluated using human, rat, mouse, or other animal liver or intestine microsomes to predict intrinsic clearance.

[1170] The assay was carried out in 96-well microtiter plates at 37 °C. Reaction mixtures (25 μL) contained a final concentration of 1 μM test compound, 0.5 mg/mL liver microsomes protein, and 1 mM NADPH and/or 1 mM UDPGA (with alamethicin) in 100 mM potassium phosphate, pH 7.4 buffer with 3 mM MgCl₂. The incubation was done with N=1, but duplicate incubation at each time point can be prepared if necessary. At each of the time points (for example, 0, 15, 30, and 60 minutes), 150 μL of quench solution (100% acetonitrile with 0.1% formic acid) with internal standard was transferred to each well. Besides the zero minute controls, mixtures containing the same components except the NADPH can also be prepared as the negative control. Verapamil was included as a positive control to verify assay performance. Plates were sealed, vortexed, and centrifuged at 4 °C for 15 minutes at 4000 rpm. The supernatant is transferred to fresh plates for LC/MS/MS analysis.

[1171] Summarized conditions: [Compound] = 1 μM, [LM] = 0.5 mg/mL, [NADPH] = 1 mM and/or [UDPGA] = 1 mM, Buffer = 100 mM Potassium Phosphate, pH 7.4 with 3 mM MgCl₂, Time = 0, 15, 30, and 60 min, and Temperature = 37 °C

[1172] All samples were analyzed on LC/MS/MS using an AB Sciex API 4000 instrument, coupled to a Shimadzu LC-20AD LC Pump system. Analytical samples were separated using a Waters Atlantis T3 dC18 reverse phase HPLC column (20 mm x 2.1 mm) at a flow rate of 0.5 mL/min. The mobile phase consisted of 0.1% formic acid in water (solvent A) and 0.1% formic acid in 100% acetonitrile (solvent B).

[1173] The extent of metabolism is calculated as the disappearance of the test compound, compared to the 0-min time incubation. Initial rates are calculated for the compound

concentration and used to determine half-life ($t_{1/2}$) values and subsequently, the intrinsic clearance (Cl_{int}).

[1174] $Cl_{int} = (0.693)(1/t_{1/2} \text{ (min)})(\text{g of liver/kg of body weight})(\text{mL incubation/mg of microsomal protein})(45\text{mg of microsomal protein/g of liver weight})$

[1175] Hepatocyte Stability Assay: The termination of *in vitro* intrinsic clearance (Cl_{int}) for drug candidates in the early discovery stage is common practice in the pharmaceutical industry. The Cl_{int} values of drug candidates can help to confirm whether metabolism is the main clearance pathway when it is compared with the total body clearance *in vivo*. Hepatocytes are among the routinely used *in vitro* systems for measuring clearance. Hepatocytes have been reported to retain most of the phase I and phase II metabolic activities, and therefore, it is a useful tool to estimate the intrinsic clearance for potential drug candidates.

[1176] Metabolic stability of testing compound were be evaluated using human and mouse hepatocytes and can also be use rat, mouse, or other animal hepatocytes to predict intrinsic clearance. Cryopreserved hepatocytes were removed from the liquid nitrogen tank and thawed in a 37 °C water bath. As soon as the cells pulled away from the vial wall, they were decanted into 48 ml of warm HT medium. Cells were centrifuged for four minutes at 420 rpm (50 g). After removing the supernatant, pellet was re-suspended in warm DMEM medium. Cell density was counted by a hemacytometer.

[1177] The assay was carried out in 96-well microtiter plates. Compounds were incubated for 0, 60, 120, and 180 minutes at 37 °C with hepatocytes. Reaction mixtures (50 µL) contained a final concentration of 1 µM test compound, 0.5 million cells/mL hepatocytes in the DMEM medium. At each of the time points (for example, 0, 1, 2, and 3 hours), 200 µL of quench solution (100% acetonitrile with 0.1% formic acid) with internal standard was transferred to each well. Midazolam was included as a positive control to verify assay performance. Plates were sealed and centrifuged at 4 °C for 15 minutes at 4000 rpm. The supernatant is transferred to fresh plates for LC/MS/MS analysis.

[1178] Summarized conditions (customizable): [Compound] = 1 µM, Positive control: midazolam and/or naloxone, [Hepatocyte] = 0.5 million cells/ML, time = 0, 60, 120, and 180 min, $t = 37$ °C.

[1179] All samples were analyzed on LC/MS/MS using an AB Sciex API 4000 instrument, coupled to a Shimadzu LC-20AD LC Pump system. Analytical samples were separated using a Waters Atlantis T3 dC18 reverse phase HPLC column (20 mm x 2.1 mm) at a flow rate of

0.5 mL/min. The mobile phase consisted of 0.1% formic acid in water (solvent A) and 0.1% formic acid in 100% acetonitrile (solvent B).

[1180] The extent of metabolism was calculated as the disappearance of the test compound, compared to the 0-min control reaction incubations. Initial rates were calculated for the compound concentration and used to determine $t_{1/2}$ values and subsequently, the intrinsic clearance, Cl_{int} .

[1181] $Cl_{int} = (0.693)(1/t_{1/2} \text{ (min)})(\text{mL incubation/million cells})$

[1182] Metabolic stability assay results of representative compounds from Tables A and B are listed in Table 2A, below. Metabolic stability assay results of representative compounds from Tables A and B for human hepatocytes are demonstrated in Table 2B. Liver microsome and hepatocyte half life of representative compounds are shown in Tables 2C and 2D.

[1183] Table 2A. Stability Results of Representative Compounds from Tables A and B

Compound ID	MLM +/- NADPH: Cl_{int} ($\mu\text{L}/\text{min}/\text{mg}$ protein) ¹	HLM +/- NADPH: Cl_{int} ($\mu\text{L}/\text{min}/\text{mg}$ protein) ¹	MH +/- NADPH: Cl_{int} ($\mu\text{L}/\text{min}/\text{million cells}$) ²
A3	***	***	NA
A5	NA	***	NA
B1	*	**	NA
A7	NA	*	NA
A13	***	***	**
A17	NA	***	NA
A18	NA	*	NA
A21	NA	***	NA
A22	NA	***	NA
A23	NA	**	NA
A24	NA	***	NA
A25	NA	***	NA
A27	*	***	NA
B2	*	*	NA
B3	***	***	NA
B5	***	***	**
A28	**	***	**
A29	**	**	NA
A30	**	**	NA
A31	NA	**	NA
A32	*	**	NA
A34	**	***	NA
A35	**	***	NA
A38	***	***	***
A40	NA	**	NA
A41	NA	*	NA
A42	***	***	NA
A45	***	***	NA

Compound ID	MLM +/- NADPH: Cl _{int} (μL/min/mg protein) ¹	HLM +/- NADPH: Cl _{int} (μL/min/mg protein) ¹	MH +/- NADPH: Cl _{int} (μL/min/million cells) ²
A49	***	**	NA
A50	NA	**	NA
A52	NA	***	NA
A53	NA	***	NA
A54	NA	***	NA
A56	NA	***	NA
A57	*	**	NA
A58	NA	**	NA
B8	NA	*	NA
A62	NA	***	NA
A63	NA	**	NA
B9	**	**	NA
A64	NA	*	NA
A65	**	**	NA
A66	***	***	NA
A68	***	***	NA
A70	*	**	**
A71	***	**	**
A73	***	***	NA
A74	***	***	NA
A75	NA	***	NA
B6	NA	**	NA
A76	NA	***	NA
B7	***	***	NA
A77	*	**	NA
B10	**	***	NA
A78	***	***	NA
A79	*	***	*
A80	**	***	*
A81	*	**	*
A85	*	***	*
A86	**	**	*
A87	**	***	*
A88	*	*	*
A89	*	**	*
A90	*	***	*
A91	***	***	**
A92	**	**	*
A93	***	***	***
A94	**	**	**
A95	*	**	*
A96	*	*	*
A97	**	**	*
A98	***	***	***
A99	**	***	**

Compound ID	MLM +/- NADPH: Cl _{int} (μL/min/mg protein) ¹	HLM +/- NADPH: Cl _{int} (μL/min/mg protein) ¹	MH +/- NADPH: Cl _{int} (μL/min/million cells) ²
A100	*	*	*
A101	**	*	**
A102	***	**	***
A103	**	**	***
A104	*	**	*
A105	*	**	*
A106	NA	**	NA
A107	NA	**	NA
A108	*	***	*
A109	***	***	***
A110	NA	NA	NA
A111	**	**	**
A112	**	***	**
A113	**	**	*
A114	**	***	*
A115	***	***	*
A116	**	**	**
A117	**	**	*
A118	*	*	**
A119	*	*	**
A120	***	*	**
A121	**	**	**
A122	**	***	**
A123	*	***	**
A124	*	***	*
A125	*	***	*
A126	***	***	***
A127	**	**	**
A128	NA	NA	NA
A129	NA	NA	NA
A130	***	**	***
A131	***	***	***
A132	***	***	***
A133	*	*	*
A134	***	**	***
A135	**	**	*
A136	**	**	**
A137	**	***	***
A138	NA	NA	NA
A139	NA	NA	NA
A140	*	*	*
A141	*	***	*
A142	*	**	*
A143	***	***	***
A144	**	**	*

Compound ID	MLM +/- NADPH: Cl _{int} (μL/min/mg protein) ¹	HLM +/- NADPH: Cl _{int} (μL/min/mg protein) ¹	MH +/- NADPH: Cl _{int} (μL/min/million cells) ²
A145	*	*	*
A146	***	**	*
A147	*	**	*
A148	***	*	***
A149	***	***	*
A150	NA	NA	NA
A151	NA	NA	NA
A152	NA	NA	NA
A153	**	***	**
A154	NA	NA	NA
A155	*	**	**
A156	**	**	*
A157	NA	NA	NA
A158	NA	NA	NA
A159	NA	NA	NA
A160	NA	NA	NA
A161	***	***	NA
A162	*	**	*
A163	NA	NA	NA
A164	***	***	***
A165	NA	NA	NA
A166	NA	NA	NA
A167	NA	NA	NA
A168	***	***	**
A169	***	***	***
A170	***	***	***
A171	***	***	**
A172	***	***	***
A173	***	***	NA
A174	**	***	NA
A175	NA	NA	NA
A176	NA	NA	NA
A177	NA	NA	NA
A178	NA	NA	NA
A179	NA	NA	NA
A180	NA	NA	NA
A181	NA	NA	NA
A182	NA	NA	NA
A183	NA	NA	NA
A184	***	***	***
A185	**	***	**
A186	NA	NA	NA
A187	NA	NA	NA
A188	NA	NA	NA
A189	***	***	NA

Compound ID	MLM +/- NADPH: Cl _{int} (μL/min/mg protein) ¹	HLM +/- NADPH: Cl _{int} (μL/min/mg protein) ¹	MH +/- NADPH: Cl _{int} (μL/min/million cells) ²
A190	*	***	*
A191	NA	NA	NA
A192	**	**	*
A193	*	*	**
A194	***	***	***
A195	***	***	**
A196	***	**	***
A197	***	***	***
A198	**	***	**
A199	*	***	**
A200	***	***	**
A201	***	***	***
A202	*	***	*
A203	NA	NA	NA
A204	***	***	***
A205	NA	NA	NA
A206	***	***	***
A207	***	***	***
A208	***	***	***
A209	NA	NA	NA
A210	NA	NA	NA
A211	NA	NA	NA

¹ For MLM and HLM: *** represent Cl < 12 μL/min/mg protein, ** represents Cl in the range of 12-48 μL/min/mg protein, * represents Cl in the range of ≥ 48 μL/min/mg protein.

² For MH: *** represent Cl < 4 μL/min/million cells, ** represents Cl in the range of 4-18 μL/min/million cells, * represents Cl in the range of ≥ 18 μL/min/million cells.

[1184] Table 2B. Human Hepatocyte Stability Results of Representative Compounds from Tables A and B

	Number of Compounds
Cl < 4 μL/min/million cells	14 compounds
Cl in the range of 4-18 μL/min/million cells	31 compounds
Cl in the range of ≥ 18 μL/min/million cells	36 compounds

[1185] Table 2C. Liver Microsome Half-Life

Compound ID	Liver microsome T _{1/2} (min)				
	Human	Mouse	Rat	Dog	Monkey
A13	>120	>120	64	>120	>120

A28	>120	74	23	>120	73
A38	>120	>120	12	>120	>120
A74	>120	>120	>120	>120	>120
A93	>120	>120	83	>120	101
A109	>120	>120	>120	>120	>120
X	>120	NA	NA	NA	NA
Enzalutamide	>120	>120	>120	>120	>120

[1186] Table 2D. Hepatocytes Half-Life

Compound ID	Hepatocytes T _{1/2} (min)		
	Human	Mouse	Rat
A13	290	203	75
A28	>360	86	31
A38	>360	>360	56
A74	>360	NA	>360
A93	>360	>360	94
A109	>360	>360	>360
X	>360	33	NA
Enzalutamide	>360	NA	NA

[1187] Example 37: *In Vivo* Pharmacokinetic Properties

[1188] The purpose of this study is to determine the pharmacokinetics of the tested compounds in plasma, following oral gavage administration (PO) to male CD-1 mice.

[1189] Animal Husbandry: Animals were group housed during acclimation and the study. The animal room environment were controlled (target conditions: temperature 18 to 26 °C, relative humidity 30 to 70%, 12 hours artificial light and 12 hours dark). Temperature and relative humidity were monitored daily. Animals were fasted at least 12 hours prior to the administration. All animals had access to Certified Rodent Diet ad libitum 4 hours post dosing.

[1190] Dose Formulation: prepared 85-100 uL of a stock solution of representative compounds of the disclosure in DMSO at 50 mM; 1.5% v/v of Tween 80 were added and mixed gently; 95.5% v/v saline was added gently to the organic phase. The solution was mixed slowly by reverse pipetting to get a clear solution.

[1191] Dose Administration: the dose formulations were administered via oral gavage per facility SOPs. The dose volume were determined by the animals' body weight collected on the morning of dosing day

[1192] Sample Collection: blood collection (about 0.05 mL per time point) was performed from saphenous vein of each animal into polypropylene tubes at each timepoints (0, 0.25, 0.5, 1, 2, 4, 8, 12, and 24 hr). All blood samples were transferred into pre-chilled tubes containing 2 µL of K2-EDTA (0.5M) and placed on wet ice until centrifugation. Each collected blood

sample was in the wet-ice before centrifugation. Each collected blood was under centrifugation for 15 minutes at 4 °C and 3000 g for plasma collection. Plasma samples were stored in polypropylene tubes, quickly frozen over dry ice and kept at -70 ± 10 °C until LC/MS/MS analysis.

[1193] Bioanalytical analysis: A LC-MS/MS method for the quantitative determination of tested compound in biological matrix were developed under non-GLP compliance. A calibration curve with at least 7 non-zero calibration standards were applied for the method including LLOQ.

[1194] Plasma concentration versus time data were analyzed by non-compartmental approaches using the Phoenix WinNonlin 6.3 software program. Cl, Vdss, C₀, C_{max}, T_{max}, T_{1/2}, AUC(0-t), AUC(0-inf), MRT(0-t), MRT(0-inf), %F (oral availability) and graphs of plasma concentration versus time profile were reported. Fig. 3 shows the time-concentration curve of representative compounds in mice as measured in plasma.

[1195] Tables 3A shows PK parameters determined from single PO dose of representative compounds in mice (5 mg/kg).

[1196] Table 3A. Summary of PO PK Parameters After Single Dose (5 mg/kg) in Male CD-1 Mice

Compound ID	t _{1/2} (hr)	t _{max} (hr)	C _{max} (ng/mL)	AUC _{last} (hr*ng/mL)	AUC/D (hr*kg*ng/mL/mg)	F (%)
A13	8.3	1.0	1143.3	8309.3	1661.9	85.8
A28	2.8	0.5	2013.3	8425.4	1685.1	50.3
A29	2.5	0.5	2010.0	5186.6	1037.3	60.6
A35	4.0	0.5	818.7	2415.1	483.0	49.6
A38	4.9	3.3	1270.0	12805.1	2561.0	86.4
A66	12.0	0.8	1613.3	11927.2	2385.4	71.4
A74	12.5	2.7	6306.7	72711.4	14542.3	71.9
A93	17.1	0.8	2973.3	32878.6	6575.7	62.8
A109	8.1	2.2	2673.3	30714.8	6143.0	33.6
A122	1.1	0.5	4080.0	6800.0	1360.0	92.4
A126	4.7	1.3	3113.0	24547.0	4909.4	86.0
A131	5.0	1.3	7053.0	76047.0	15209.4	81.0
A136	10.2	1.5	3427.0	42038.0	8407.6	14.0
A170	3.0	3.0	8503.0	80038.0	16007.6	150.0

[1197] **Example 38:** In vivo Activity of Representative Compounds in LNCaP Xenografts Model

[1198] Tumor growth was measured in male NCG mice bearing LNCaP tumors. Castration was performed when tumors reached $\sim 100 \text{ mm}^3$ and dosing (60 mg/kg PO qd) started 1 week after castration. See Fig. 4. Body weight of the mice were captured biweekly in the animals which showed no drug related toxicity. See Fig. 5. Individual tumor volume change from baseline measured at the end of the experiment. See Fig. 6. Data demonstrated in Figs. 4-6 shows that the representative compounds showed activity and induced partial regressions of tumor growth. The $C_{\min}$ at 5 mg/kg PO and extrapolated $C_{\min}$ in efficacy of the representative compounds are shown in Table 4.


[1199] **Table 4.** $C_{\min}$ at 5 mg/kg PO and $C_{\min}$ at 5 mg/kg PO

Compound ID	$C_{\min}$ at 5 mg/kg PO (μM)	$C_{\min}$ at 5 mg/kg PO (μM)
A13	0.23	2.78
A74	2.42	29.1
A109	0.68	8.18
Enzalutamide	3.80	22.8

[1200] Numbered Embodiments

[1201] In some embodiments, the present disclosure relates to the following embodiments.

[1202] Embodiment 1: A compound having a structure of formula (I):



[1203] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1204] A and B are each independently aryl or heteroaryl;

[1205] C is a 3- to 10-membered ring;

[1206] X is a bond, $-(CR^5R^6)_-$, $-O-$, $-C(=O)-$, $-S-$, $-S(=O)-$, $-SO_2-$, $-NR^7-$, $-N(R^7)CO-$, $-CON(R^7)-$, or $-NSO_2R^7-$;

[1207] Y and Z are each independently a bond, $-(CR^8R^9)_m-$, $-O-$, $-C(=O)-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;

[1208] W and V are each independently a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

[1209] L is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;

[1210] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[1211] R^3 is hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-SR^{16}$, optionally substituted C_1-C_6 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-$

C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[1212] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁵ and R⁶ taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[1213] R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[1214] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -OCO(C₁-C₆ alkyl), -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^{8a} and R^{8b} taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[1215] R⁷, R¹⁰ and R¹⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁷ and R^{8a} taken together form an optionally substituted heterocyclyl;

[1216] R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or (R¹¹ and R¹²) or (R¹⁴ and R¹⁵) taken together form an optionally substituted heterocyclyl;

[1217] each m is independently 0, 1 or 2;

[1218] n_1 and n_2 are each independently 0, 1, 2, 3, or 4;

[1219] n_3 is 0, 1, 2, 3, 4 or 5; and

[1220] each t is independently 0, 1 or 2.

[1221] **Embodiment 2:** The compound of **Embodiment 1**, wherein A and B are each independently 5- or 6-membered aryl or heteroaryl.

[1222] **Embodiment 3:** The compound of **Embodiment 1 or 2**, wherein A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene.

[1223] **Embodiment 4:** The compound of any one of **Embodiments 1-3**, wherein A has a meta or para connectivity with X and Y.


[1224] **Embodiment 5:** The compound of any one of **Embodiments 1-4**, wherein B has a meta or para connectivity with X and Z.


[1225] **Embodiment 6:** The compound of any one of **Embodiments 1-5**, wherein C is 5- to 10-membered heteroaryl or aryl.

[1226] **Embodiment 7:** The compound of any one of **Embodiments 1-6**, wherein C is 5- to 7-membered heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

[1227] **Embodiment 8:** The compound of any one of **Embodiments 1-7**, wherein C is pyrazole, imidazole, oxazole, oxadiazole, oxazolone, isoxazole, thiazole, pyridyl, or pyrimidyl.

[1228] **Embodiment 9:** The compound of any one of **Embodiments 1-7**, wherein C, which is






[1229] **Embodiment 10:** The compound of any one of **Embodiments 1-6**, wherein C is phenyl or naphthyl.

[1230] **Embodiment 11:** The compound of any one of **Embodiments 1-5**, wherein C is 5- to 10-membered heterocyclyl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

[1231] **Embodiment 12:** The compound of **Embodiment 11**, wherein



[1233] D is -O-, -NH- or -NR³-; and

[1234] U is each independently O, S, or NR¹⁶.

[1235] **Embodiment 13:** The compound of **Embodiment 11**, wherein C is imidazolidine, imidazolidine-dione, or dihydrooxazole.

[1236] **Embodiment 14:** The compound of any one of **Embodiments 1-13**, wherein Y is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-.

[1237] **Embodiment 15:** The compound of any one of **Embodiments 1-14**, wherein W is a bond, -CH₂-, -C(CH₃)H-, -C(=O)-, -N(R⁷)CO-, or -CONR⁷-, wherein R⁷ is H or C₁-C₃ alkyl.

[1238] **Embodiment 16:** The compound of any one of **Embodiments 1-15**, wherein:

[1239] Z is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-; and

[1240] V is -CH₂-, -CH₂CH₂-, or -CH₂CH₂CH₂-.

[1241] **Embodiment 17:** The compound of any one of **Embodiments 1-16**, wherein:

[1242] Z is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-;

[1243] V is -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-; and

[1244] L is halogen, -NH₂, or -CF₃.

[1245] **Embodiment 18:** The compound of any one of **Embodiments 1-17**, wherein X is a bond or -(CR⁵R⁶)_n.

[1246] **Embodiment 19:** The compound of any one of **Embodiments 1-18**, wherein X is a bond, -CH₂-, -C(CH₃)H-, -C(CH₃)₂-, or -CH₂CH₂-.

[1247] **Embodiment 20:** The compound of any one of **Embodiments 1-19**, wherein R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₁-C₃ alkoxy, -(C₁-C₃ alkyl)-(C₁-C₃ alkoxy), -(C₁-C₃ alkyl)-OH, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂R¹⁶, -(C₁-C₃ alkyl)-SO₂R¹⁶, 3- to 7-membered carbocyclyl, 3- to 7-membered heterocyclyl, phenyl, or 5- to 6-membered heteroaryl.

[1248] **Embodiment 21:** The compound of any one of **Embodiments 1-20**, wherein R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₁-C₃ alkoxy, -(C₁-C₃ alkyl)-(C₁-C₃ alkoxy), -(C₁-C₃ alkyl)-OH, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂R¹⁶, or -(C₁-C₃ alkyl)-SO₂R¹⁶.

[1249] **Embodiment 22:** The compound of any one of **Embodiments 1-21**, wherein R¹ and R² are each independently halogen, -CN, -CF₃, -OH, methyl, methoxy, or -CONH₂.

[1250] **Embodiment 23:** The compound of any one of **Embodiments 1-22**, wherein R¹ and R² are each independently Cl, -CN, -CF₃, -OH, methyl, methoxy, or -CONH₂.

[1251] **Embodiment 24:** The compound of any one of **Embodiments 1-23**, wherein n₁ and n₂ are each independently 0, 1, or 2.

[1252] **Embodiment 25:** The compound of any one of **Embodiments 1-24**, wherein the sum of n₁ and n₂ is 0, 1, 2, 3, or 4.

[1253] **Embodiment 26:** The compound of any one of **Embodiments 1-25**, wherein R³ is selected from hydrogen, halogen, oxo, =S, or =NR¹⁶, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -(C₁-C₃ alkyl)-(C₁-C₃ alkoxy), -(C₁-C₃ alkyl)-OH, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl).

[1254] **Embodiment 27:** The compound of any one of **Embodiments 1-26**, wherein R^3 is selected from hydrogen, F, Cl, Br, I, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, -S(C₁-C₃ alkyl), -SO(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)CO(C₁-C₃ alkyl).

[1255] **Embodiment 28:** The compound of any one of **Embodiments 1-27**, wherein R^3 is selected from hydrogen, F, Cl, Br, I, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, methyl, -SCH₃, -SO₂CH₃, -NHSO₂CH₃, -CH₂NHSO₂CH₃, -SO₂NH₂, -CONH₂, or -NHCOCH₃.


[1256] **Embodiment 29:** The compound of any one of **Embodiments 1-28**, wherein at least one of R^3 is -SO₂CH₃, -NHSO₂CH₃, -CH₂NHSO₂CH₃, -SO₂NH₂, -CONH₂, or -NHCOCH₃.

[1257] **Embodiment 30:** The compound of any one of **Embodiments 1-29**, wherein at least one of R^3 is oxo, =S, =NR¹⁶.

[1258] **Embodiment 31:** The compound of any one of **Embodiments 1-30**, wherein R^{8a} and R^{9a} are each independently hydrogen, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl.

[1259] **Embodiment 32:** The compound of any one of **Embodiments 1-30**, wherein R^{8a} and R^{9a} are not -OH.

[1260] **Embodiment 33:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (II):



[1261] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1262] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

- [1263] C is a 3- to 10-membered ring;
- [1264] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;
- [1265] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;
- [1266] Z is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;
- [1267] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-NHCO-$, $-N(C_1-C_3 \text{ alkyl})CO-$, or $-CONH-$, or $-CON(C_1-C_3 \text{ alkyl})-$;
- [1268] V is a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;
- [1269] L is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;
- [1270] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;
- [1271] R^3 is selected from hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;
- [1272] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, or C_1-C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [1273] R^8 and R^9 are each independently hydrogen, halogen, or C_1-C_3 alkyl;
- [1274] R^{8a} and R^{9a} are each independently hydrogen, $-OH$, halogen, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}COR^{16}$, $-(C_1-C_3 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, or $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [1275] R^{11} , R^{12} , R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1-C_3 alkyl, C_2-C_3 alkenyl, or C_2-C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[1276] R^7 , R^{10} and R^{16} are each independently hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl;


[1277] each m is independently 0, 1 or 2;

[1278] n_1 and n_2 are each independently 0, 1, or 2;

[1279] n_3 is 0, 1, 2, 3, 4 or 5; and

[1280] t is 0, 1 or 2.

[1281] **Embodiment 34:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (III):



[1282] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1283] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[1284] C is a 3- to 10-membered ring;

[1285] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;

[1286] Y is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;

[1287] W is a bond, $-(CR^8R^9)_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

[1288] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[1289] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[1290] L is halogen, $-NH_2$, or $-CF_3$;

[1291] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)-(C_1 - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} ;

[1292] R^3 is selected from hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);

[1293] R^5 and R^6 are each independently hydrogen, halogen, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1294] R^7 is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl;

[1295] R^8 and R^9 are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[1296] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, or -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1297] R^{13} , R^{14} and R^{15} are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[1298] R^{16} is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;

[1299] each m is independently 0, 1 or 2;

[1300] n₁ and n₂ are each independently 0, 1, or 2;


[1301] n₃ is 0, 1, 2, 3, 4 or 5; and

[1302] t is 0, 1 or 2.

[1303] **Embodiment 35:** The compound of **Embodiment 33 or 34**, wherein R^{8a} and R^{9a} are each independently hydrogen, halogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, or -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵.

[1304] **Embodiment 36:** The compound of **Embodiment 33 or 34**, wherein R^{8a} and R^{9a} are not -OH.

[1305] **Embodiment 37:** The compound of **Embodiments 1**, wherein the compound has the structure of formula (IV):



[1306] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1307] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[1308] C is a 3- to 10-membered ring;

[1309] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;

[1310] Y and Z are each independently a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[1311] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[1312] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[1313] L is halogen, $-NH_2$, or $-CF_3$;

[1314] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)-(C_1 - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} ;

[1315] R^3 is selected from hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1$ - C_3 alkyl), C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1$ - C_3 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1$ - C_3 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $SO_2NR^{14}R^{15}$, $-SO_2(C_1$ - C_3 alkyl), or $-(C_1$ - C_6 alkyl)- $SO_2(C_1$ - C_3 alkyl);

[1316] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, or C_1 - C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1317] R^7 is H or C₁-C₆ alkyl;

[1318] R^{13} , R^{14} and R^{15} are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[1319] R^{16} is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;

[1320] n_1 and n_2 are each independently 0, 1, or 2;

[1321] n_3 is 0, 1, 2, 3, 4 or 5; and


[1322] t is 0, 1 or 2.


[1323] **Embodiment 38:** The compound of any one of **Embodiments 33-37**, wherein C is 5- to 10-membered heteroaryl or aryl.

[1324] **Embodiment 39:** The compound of any one of **Embodiments 33-38**, wherein C is 5- to 7-membered heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

[1325] **Embodiment 40:** The compound of any one of **Embodiments 33-39**, wherein C is pyrazole, imidazole, oxazole, oxadiazole, oxazolone, isoxazole, thiazole, pyridyl, or pyrimidyl.

[1326] **Embodiment 41:** The compound of any one of **Embodiments 33-39**, wherein C, which






[1327] **Embodiment 42:** The compound of any one of **Embodiments 33-38**, wherein C is phenyl or naphthyl.

[1328] **Embodiment 43:** The compound of any one of **Embodiments 33-37**, wherein C is 5- to 10-membered heterocyclyl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

[1329] **Embodiment 44:** The compound of **Embodiment 43**, wherein



[1331] D is -O-, -NH- or -NR³-; and

[1332] U is each independently O, S, or NR¹⁶.


[1333] **Embodiment 45:** The compound of **Embodiment 43**, wherein C is imidazolidine, imidazolidine-dione, or dihydrooxazole.

[1334] **Embodiment 46:** The compound of any one of **Embodiments 33-45**, wherein A has a meta or para connectivity with X and Y.

[1335] **Embodiment 47:** The compound of any one of **Embodiments 33-46**, wherein B has a meta or para connectivity with X and Z.

[1336] **Embodiment 48:** The compound of any one of **Embodiments 33-47**, wherein n₁ and n₂ are each independently 0, 1, or 2.

[1337] **Embodiment 49:** The compound of **Embodiments 1**, wherein the compound has the



structure of formula (V):

[1338] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1339] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[1340] C is a 5- to 10-membered heteroaryl or aryl;

[1341] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;

[1342] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[1343] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[1344] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[1345] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[1346] L is halogen, $-NH_2$, or $-CF_3$;

[1347] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;

[1348] R^3 is selected from hydrogen, halogen, oxo, $=S$, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;

[1349] R^5 and R^6 are each independently hydrogen, halogen, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1350] R^7 is H or C₁-C₆ alkyl;

[1351] R^{13} , R^{14} and R^{15} are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[1352] R^{16} is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;

[1353] n_1 and n_2 are each independently 0, 1, or 2;


[1354] n_3 is 0, 1, 2, 3, 4 or 5; and


[1355] t is 0, 1 or 2.

[1356] **Embodiment 50:** The compound of **Embodiment 49**, wherein C is 5- to 7-membered heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

[1357] **Embodiment 51:** The compound of **Embodiment 49 or 50**, wherein C is pyrazole, imidazole, oxazole, oxadiazole, oxazolone, isoxazole, thiazole, pyridyl, or pyrimidyl.

[1358] **Embodiment 52:** The compound of **Embodiment 49**, wherein C, which is optionally





[1359] **Embodiment 53:** The compound of any one of **Embodiments 49-52**, wherein A has a meta or para connectivity with X and Y.

[1360] **Embodiment 54:** The compound of any one of **Embodiments 49-53**, wherein B has a meta or para connectivity with X and Z.


[1361] **Embodiment 55:** The compound of any one of **Embodiments 49-54**, wherein A and B are each phenyl.

[1362] **Embodiment 56:** The compound of any one of **Embodiments 49-55**, wherein $-Z-V-L$ is $-Z-CH_2CH_2Cl$, $-Z-CH_2CH_2CH_2Cl$, $-Z-CH_2CH_2NH_2$, or $-Z-CH_2CH_2CH_2NH_2$, wherein Z is a bond, $-O-$, $-NH-$, or $-N(COCH_3)-$.

[1363] **Embodiment 57:** The compound of any one of **Embodiments 49-56**, wherein $-Y-W-$ is a bond, $-OCH_2-$, $-OCH_2CH_2-$, $-OCH(CH_3)-$, $-NH-$, $-NHCH_2-$, $-NHC(=O)-$, or $-C(=O)NH-$.

[1364] **Embodiment 58:** The compound of any one of **Embodiments 49-57**, wherein X is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(CH_3)_2-$, or $-CH_2CH_2-$.

[1365] **Embodiment 59:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (VI):



[1366] wherein:

[1367] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;


[1368] C is a 5- to 10-membered heterocyclyl;

- [1369] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;
- [1370] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;
- [1371] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;
- [1372] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;
- [1373] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;
- [1374] L is halogen, $-NH_2$, or $-CF_3$;
- [1375] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)- $(C_1$ - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} ;
- [1376] R^3 is selected from hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1$ - C_3 alkyl), C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1$ - C_3 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1$ - C_3 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $SO_2NR^{14}R^{15}$, $-SO_2(C_1$ - C_3 alkyl), or $-(C_1$ - C_6 alkyl)- $SO_2(C_1$ - C_3 alkyl);
- [1377] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, or C_1 - C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [1378] R^7 is H or C_1 - C_6 alkyl;
- [1379] R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl; and
- [1380] R^{16} is hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl;
- [1381] n_1 and n_2 are each independently 0, 1, or 2;
- [1382] n_3 is 0, 1, 2, 3, 4 or 5; and
- [1383] t is 0, 1 or 2.

[1384] **Embodiment 60:** The compound of **Embodiment 57**, wherein C is 5- to 7-membered saturated or partially saturated heterocycle comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

[1385] **Embodiment 61:** The compound of **Embodiment 59 or 60**, wherein C is imidazolidine, imidazolidine-dione, or dihydrooxazole.

[1386] **Embodiment 62:** The compound of any one of **Embodiments 59-61**, wherein C, which is optionally substituted with R^3 , is selected from



[1387] **Embodiment 63:** The compound of any one of **Embodiments 59-62**, wherein A has a meta or para connectivity with X and Y.

[1388] **Embodiment 64:** The compound of any one of **Embodiments 59-63**, wherein B has a meta or para connectivity with X and Z.

[1389] **Embodiment 65:** The compound of any one of **Embodiments 59-64**, wherein A and B are each phenyl.


[1390] **Embodiment 66:** The compound of any one of **Embodiments 59-65**, wherein -V-L is -CH₂CH₂Cl, -CH₂CH₂CH₂Cl, -CH₂CH₂NH₂, or -CH₂CH₂CH₂NH₂, and Z is a bond, -O-, -NH-, or -N(COCH₃)-.

[1391] **Embodiment 67:** The compound of any one of **Embodiments 59-66**, wherein -Y-W- is a bond, -OCH₂-, -OCH₂CH₂-, -OCH(CH₃)-, -NH-, -NHCH₂-, -NHC(=O)-, or -C(=O)NH-.

[1392] **Embodiment 68:** The compound of any one of **Embodiments 59-67**, wherein X is a bond, -CH₂-, -C(CH₃)H-, -C(CH₃)₂-, or -CH₂CH₂-.

[1393] **Embodiment 69:** The compound of any one of **Embodiments 59-68**, R^3 is oxo, =S, =NR¹⁶, C₁-C₃ alkyl, -SO₂(C₁-C₃ alkyl), or -NHSO₂(C₁-C₃ alkyl).

[1394] **Embodiment 70:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (A):




[1395] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1396] C is a phenyl or a 5- to 7-membered monocyclic heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member;

- [1397] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;
- [1398] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;
- [1399] Z is a bond, $-CH_2-$, $-O-$, or $-NH-$;
- [1400] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;
- [1401] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;
- [1402] L is halogen, $-NH_2$, or $-CF_3$;
- [1403] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, methyl, or $-CONH_2$;
- [1404] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $-CN$, $-CF_3$, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, $-S(C_1$ - C_3 alkyl), $-SO_2(C_1$ - C_3 alkyl), $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1$ - C_3 alkyl) $_2$, $-CONH(C_1$ - C_3 alkyl), $-NHCO(C_1$ - C_3 alkyl), or $-N(CH_3)CO(C_1$ - C_3 alkyl);
- [1405] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, or C_1 - C_3 alkyl;
- [1406] R^7 is H or C_1 - C_6 alkyl;
- [1407] n_1 and n_2 are each independently 0, 1, or 2;
- [1408] n_3 is 0, 1, 2, 3, 4 or 5; and
- [1409] t is 0, 1 or 2.

[1410] **Embodiment 71:** The compound of **Embodiment 70**, wherein C is 5- to 7-membered monocyclic heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

[1411] **Embodiment 72:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (B)



[1412] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

- [1413] C is a 5- to 7-membered saturated or partially saturated monocyclic heterocycle comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member;
- [1414] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;


- [1415] Y is a bond, $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{NCH}_3-$, or $-\text{N}(\text{COCH}_3)-$;
- [1416] Z is a bond, $-\text{CH}_2-$, $-\text{O}-$, or $-\text{NH}-$;
- [1417] W is a bond, $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{C}(=\text{O})-$, $-\text{N}(\text{R}^7)\text{CO}-$, or $-\text{CONR}^7-$;
- [1418] V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$;
- [1419] L is halogen, $-\text{NH}_2$, or $-\text{CF}_3$;
- [1420] R^1 and R^2 are each independently hydrogen, halogen, $-\text{CN}$, $-\text{CF}_3$, methyl, or $-\text{CONH}_2$;
- [1421] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $=\text{S}$, $=\text{NR}^{16}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{OH}$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, $-\text{S}(\text{C}_1$ - C_3 alkyl), $-\text{SO}_2(\text{C}_1$ - C_3 alkyl), $-\text{NHSO}_2\text{CH}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{CH}_2\text{NHSO}_2\text{CH}_3$, $-\text{CH}_2\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{SO}_2\text{NH}_2$, $-\text{CONH}_2$, $-\text{CON}(\text{C}_1$ - C_3 alkyl) $_2$, $-\text{CONH}(\text{C}_1$ - C_3 alkyl), $-\text{NHCO}(\text{C}_1$ - C_3 alkyl), or $-\text{N}(\text{CH}_3)\text{CO}(\text{C}_1$ - C_3 alkyl);
- [1422] R^5 and R^6 are each independently hydrogen, halogen, $-\text{OH}$, or C_1 - C_3 alkyl;
- [1423] R^7 is H or C_1 - C_6 alkyl;
- [1424] R^{16} is hydrogen or C_1 - C_3 alkyl;
- [1425] n_1 and n_2 are each independently 0, 1, or 2;
- [1426] n_3 is 0, 1, 2, 3, 4 or 5; and
- [1427] t is 0, 1 or 2.

[1428] **Embodiment 73:** The compound of any one of **Embodiments 70-72**, wherein $-\text{V-L}$ is $-\text{CH}_2\text{CH}_2\text{Cl}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$, $-\text{CH}_2\text{CH}_2\text{NH}_2$, or $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$.

[1429] **Embodiment 74:** The compound of any one of **Embodiments 70-73**, wherein $-\text{Y-W-}$ is a bond, $-\text{OCH}_2-$, $-\text{OCH}_2\text{CH}_2-$, $-\text{OCH}(\text{CH}_3)-$, $-\text{NH}-$, $-\text{NHCH}_2-$, $-\text{NHC}(=\text{O})-$, or $-\text{C}(=\text{O})\text{NH}-$.

[1430] **Embodiment 75:** The compound of any one of **Embodiments 70-74**, wherein X is a bond, $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{C}(\text{CH}_3)_2-$, or $-\text{CH}_2\text{CH}_2-$.


[1431] **Embodiment 76:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (C):



[1432] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

- [1433] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;
- [1434] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;
- [1435] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;
- [1436] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;
- [1437] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7$;
- [1438] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;
- [1439] L is halogen, $-NH_2$, or $-CF_3$;
- [1440] D is $-NH$ or $-NR^3$;
- [1441] U is each independently O, S, or NR^{16} ;
- [1442] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;
- [1443] R^3 is selected from hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;
- [1444] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [1445] R^7 is H or C_1-C_6 alkyl;
- [1446] R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;
- [1447] R^{16} is hydrogen, C_1-C_3 alkyl, C_2-C_3 alkenyl, or C_2-C_3 alkynyl;
- [1448] n_1 and n_2 are each independently 0, 1, or 2;
- [1449] n_3 is 0, 1, 2, or 3; and
- [1450] t is 0, 1 or 2.

[1451] **Embodiment 77:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (D):



[1452] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1453] C is a 5- or 6-membered heteroaryl comprising 1 or 2 heteroatoms selected from O, S, or N as a ring member;

[1454] X is $-(CR^5R^6)_t-$ or $-NR^7-$;

[1455] Y is a bond, $-CH_2-$, $-O-$, or $-NH-$;

[1456] Z is a bond, $-CH_2-$, $-O-$, or $-NH-$;

[1457] W is a bond, $-CH_2-$, or $-C(CH_3)H-$;

[1458] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[1459] L is hydrogen, halogen, $-NH_2$, or $-CF_3$;

[1460] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, methyl, or $-CONH_2$;

[1461] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $-CN$, $-CF_3$, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, $-S(C_1$ - C_3 alkyl), $-SO_2(C_1$ - C_3 alkyl), $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1$ - C_3 alkyl) $_2$, $-CONH(C_1$ - C_3 alkyl), $-NHCO(C_1$ - C_3 alkyl), or $-N(CH_3)CO(C_1$ - C_3 alkyl);

[1462] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, or C_1 - C_3 alkyl;


[1463] R^7 is H or C_1 - C_6 alkyl;

[1464] n_1 and n_2 are each independently 0, 1, or 2;

[1465] n_3 is 0, 1, 2, 3, 4 or 5; and

[1466] t is 0, 1 or 2.


[1467] **Embodiment 78:** The compound of **Embodiments 1**, wherein the compound has the



structure of formula (E)

(E)

[1468] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[1470] X is $-(CR^5R^6)_t-$ or $-NR^7-$;

[1471] Y is a bond, $-CH_2-$, $-O-$, or $-NH-$;

[1472] Z is a bond, $-CH_2-$, $-O-$, or $-NH-$;

[1473] W is a bond, $-CH_2-$, or $-C(CH_3)H-$;

[1474] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[1475] L is hydrogen or halogen;

[1476] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, methyl, or $-CONH_2$;

[1477] R^3 is selected from hydrogen, F, Cl, Br, I, oxo, $-CN$, $-CF_3$, $-OH$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, $-S(C_1-C_3 \text{ alkyl})$, $-SO_2(C_1-C_3 \text{ alkyl})$, $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1-C_3 \text{ alkyl})_2$, $-CONH(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, or $-N(CH_3)CO(C_1-C_3 \text{ alkyl})$;

[1478] R^5 and R^6 are each independently hydrogen or C_1-C_3 alkyl;


[1479] R^7 is H or C_1-C_6 alkyl;

[1480] $n1$ and $n2$ are each independently 0, 1, or 2;


[1481] $n3$ is 0, 1, or 2; and

[1482] t is 1 or 2.

[1483] **Embodiment 79:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (F):



[1484] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[1486] X is $-(CR^5R^6)_t-$ or $-NR^7-$;

[1487] Y is $-O-$;

[1488] Z is $-O-$;

[1489] W is $-CH_2-$ or $-C(CH_3)H-$;

[1490] V is $-CH_2-$, $-CH_2CH_2-$, or $-CH_2CH_2CH_2-$;

[1491] L is hydrogen or halogen;

[1492] R^1 and R^2 are each independently halogen or $-CN$;

[1493] R^3 is selected from $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, or $-SO_2CH_3$;

[1494] R^5 and R^6 are each independently hydrogen or C_1 - C_3 alkyl;


[1495] R^7 is H or C_1 - C_6 alkyl;

[1496] n_1 and n_2 are each independently 0, 1, or 2;


[1497] n_3 is 0, 1, or 2; and

[1498] t is 1 or 2.

[1499] **Embodiment 80:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (G):



[1500] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[1502] X is $-(CR^5R^6)_t-$ or $-NR^7-$;

[1503] Y is $-O-$;

[1504] Z is $-O-$;

[1505] W is $-CH_2-$ or $-C(CH_3)H-$;

[1506] V is $-CH_2-$ and L is hydrogen;

[1507] or alternatively, V is $-CH_2CH_2-$ or $-CH_2CH_2CH_2-$, and L is halogen

[1508] R^1 and R^2 are each independently Cl or $-CN$;

[1509] R^3 is selected from $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, or $-SO_2CH_3$;

[1510] R^5 and R^6 are each independently hydrogen or methyl;


[1511] R^7 is H or C_1-C_6 alkyl;

[1512] n_1 and n_2 are each independently 0, 1, or 2;


[1513] n_3 is 1, or 2; and

[1514] t is 1.

[1515] **Embodiment 81:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (H)



[1516] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[1518] X is $-(\text{CR}^5\text{R}^6)_t-$ or $-\text{NR}^7-$;

[1519] Y is -O-;

[1520] Z is -O-;

[1521] W is $-\text{CH}_2-$ or $-\text{C}(\text{CH}_3)\text{H}-$;

[1522] V is -CH₂-, -CH₂CH₂- or -CH₂CH₂CH₂-;

[1523] L is halogen;

[1524] R¹ and R² are each independently Cl or -CN;

[1525] R⁵ and R⁶ are each independently hydrogen or methyl;

[1526] R⁷ is H or C₁-C₆ alkyl;

[1527] n_1 and n_2 are each independently 0, 1, or 2; and

[1528] t is 1.

[1529] Embodiment 82: The compound of any one of **Embodiments 1-69**, wherein A and B are each monocyclic ring.

[1530] Embodiment 83: The compound of any one of **Embodiments 1-69**, wherein B is phenyl, pyridyl, or pyrimidyl.

[1531] Embodiment 84: The compound of any one of **Embodiments 1-36**, wherein Z and V are not both a bond.

[1532] Embodiment 85: The compound of any one of **Embodiments 1-78**, wherein Y and W are not both a bond.

[1533] **Embodiment 86:** The compound of any one of **Embodiments 1-48**, wherein **C** is a 4- to 10-membered ring.


[1534] **Embodiment 87:** A compound selected from Compounds **A1-A97** or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

[1535] **Embodiment 88:** A compound selected from Compounds **A98-A186** or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

[1536] **Embodiment 89:** A compound selected from Compounds **A187-A211** or a pharmaceutically acceptable salt thereof.

[1537] **Embodiment 90:** A compound selected from Compounds **B1-B11** or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

[1538] **Embodiment 91:** A compound having a structure of formula (IA):



[1539] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1540] **A** and **B** are each independently aryl or heteroaryl;

[1541] **C** is a 3- to 10-membered ring;

[1542] **X** is a bond, $-(CR^5R^6)_l-$, $-O-$, $-C(=O)-$, $-S-$, $-S(=O)-$, $-SO_2-$, $-NR^7-$, $-N(R^7)CO-$, $-CON(R^7)-$, or $-NSO_2R^7-$;

[1543] **Y** and **Z** are each independently a bond, $-(CR^8R^9)_m-$, $-O-$, $-C(=O)-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;

[1544] **W** and **V** are each independently a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

[1545] **L** is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;

[1546] **R**¹ and **R**² are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, -

SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[1547] R³ is hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -SR¹⁶, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COOR¹⁶, -NR¹⁴COR¹⁶, -NR¹⁴CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[1548] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁵ and R⁶ taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[1549] R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[1550] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -OCO(C₁-C₆ alkyl), -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^{8a} and R^{8b} taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[1551] R⁷, R¹⁰ and R¹⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, C₁-C₆

haloalkyl, C₂-C₆ haloalkenyl, C₂-C₆ haloalkynyl, optionally substituted carbocyclyl, optionally substituted -CO(C₁-C₆ alkyl), -CO(optionally substituted heterocyclyl), optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁷ and R^{8a} taken together form an optionally substituted heterocyclyl;

[1552] R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted -COO(C₁-C₆ alkyl), optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or (R¹¹ and R¹²) or (R¹⁴ and R¹⁵) taken together form an optionally substituted heterocyclyl;


[1553] each m is independently 0, 1 or 2;

[1554] n₁ and n₂ are each independently 0, 1, 2, 3, or 4;

[1555] n₃ is 0, 1, 2, 3, 4 or 5; and

[1556] each t is independently 0, 1 or 2.

[1557] **Embodiment 92:** The compound of **Embodiment 91**, wherein the compound has the



structure of formula (VA):

[1558] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1559] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[1560] C is a 5- to 10-membered heteroaryl or aryl;

[1561] X is a bond, -(CR⁵R⁶)_t-, or -NR⁷-;

[1562] Y is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-;

[1563] Z is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-;

[1564] W is a bond, -CH₂-, -C(CH₃)H-, -C(=O)-, -N(R⁷)CO-, or -CONR⁷-;

[1565] V is -CH₂-, -CH₂CH₂-, -CH(CH₃)CH₂-, -CH₂CH(CH₃)-, or -CH₂CH₂CH₂-;

[1566] L is hydrogen, halogen, -OH, -NH₂, or -CF₃;

[1567] R^1 and R^2 are each independently hydrogen, halogen, -CN, -CF₃, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶ or optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶;

[1568] R^3 is selected from hydrogen, halogen, oxo, =S, -CN, -CF₃, -OH, -S(C₁-C₃ alkyl), C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, -(C₁-C₃ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, -NR¹⁴COOR¹⁶, -NR¹⁴CONR¹⁴R¹⁵, -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);

[1569] R^5 and R^6 are each independently hydrogen, halogen, -OH, -NH₂, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1570] R^7 is H, C₁-C₆ alkyl, -CO(C₁-C₆ alkyl);

[1571] R^{13} , R^{14} and R^{15} are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or -COO(C₁-C₆ alkyl); or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;


[1572] R^{16} is hydrogen, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl;


[1573] n_1 and n_2 are each independently 0, 1, or 2;

[1574] n_3 is 0, 1, 2, 3, 4 or 5; and


[1575] t is 0, 1 or 2.

[1576] **Embodiment 93:** The compound of any one of **Embodiments 1-7, 33-39, and 49**,






[1577] **Embodiment 94:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (E-I):



[1578] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[1580] X is $-(\text{CR}^5\text{R}^6)_t-$ or $-\text{NR}^7-$:

[1581] Y is a bond, -CH₂-, -O-, or -NH-;

[1582] Z is a bond, -CH₂-, -O-, or -NH-;

[1583] W is a bond, -CH₂-, or -C(CH₃)H-;

[1584] V is -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, or -CH₂CHClCH₂-;

[1585] L is hydrogen, -OH, or halogen;

[1586] R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, methyl, or -CONH₂;

[1587] R³ is selected from hydrogen, F, Cl, Br, I, oxo, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl);

[1588] R⁵ and R⁶ are each independently hydrogen or C₁-C₃ alkyl;

[1589] R⁷ is H or C₁-C₆ alkyl;


[1590] n1 and n2 are each independently 0, 1, or 2;

[1591] n_3 is 0, 1, or 2; and


[1592] t is 1 or 2.

[1593] Embodiment 95: The compound of **Embodiment 94**, wherein R³ is selected from hydrogen, F, Cl, Br, I, -CN, -CF₃, -OH, methyl, methoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -NHCO(C₁-C₃ alkyl).

[1594] **Embodiment 96:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (G-I):



[1595] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[1597] X is $-(CR^5R^6)_t-$;

[1598] Y is $-O-$;

[1599] Z is $-O-$;

[1600] W is $-\text{CH}_2-$ or $-\text{C}(\text{CH}_3)\text{H}-$;

[1601] V is $-\text{CH}_2\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2\text{CH}_2-$;

[1602] L is halogen

[1603] R^1 and R^2 are each independently Cl or $-\text{CN}$;

[1604] R^3 is selected from $-\text{NHSO}_2\text{CH}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, or $-\text{SO}_2\text{CH}_3$;


[1605] R^5 and R^6 are each independently hydrogen or methyl;

[1606] n_1 and n_2 are each independently 0, 1, or 2;


[1607] n_3 is 1 or 2; and

[1608] t is 1.

[1609] **Embodiment 97:** The compound of **Embodiment 1**, wherein the compound has the structure of formula (H-I):



[1610] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[1612] X is $-(CR^5R^6)_t-$;

[1613] Y is $-O-$;

[1614] Z is $-O-$;

[1615] W is $-CH_2-$ or $-C(CH_3)H-$;

[1616] V is $-CH_2-$, $-CH_2CH_2-$ or $-CH_2CH_2CH_2-$;

[1617] L is halogen;

[1618] R¹ and R² are each independently Cl or $-CN$;


[1619] R⁵ and R⁶ are each independently hydrogen or methyl;

[1620] R⁷ is H or C₁-C₆ alkyl;


[1621] n₁ and n₂ are each independently 0, 1, or 2; and

[1622] t is 1.

[1623] Embodiment 98: The compound of **Embodiment 1**, wherein the compound has the structure of formula (E-II)



[1624] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[1626] X is $-(CR^5R^6)-$;

[1627] Y is a bond, -CH₂-, -O-, or -NH-;

[1628] Z is a bond, -CH₂-, -O-, or -NH-;

[1629] W is a bond, -CH₂-, or -C(CH₃)H-;

[1630] V is -CH₂-, -CH₂CH₂-, or -CH₂CH₂CH₂-;

[1631] L is halogen;


[1632] R^1 , R^{2A} and R^{2B} are each independently hydrogen, halogen, -CN, -CF₃, methyl, or -CONH₂;


[1633] R³ is selected from hydrogen, F, Cl, Br, I, oxo, -CN, -CF₃, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl);

[1634] R⁵ and R⁶ are each independently hydrogen or C₁-C₃ alkyl;


[1635] n_1 is 0, 1, or 2; and

[1636] n3 is 1 or 2;



[1637] wherein when C-R³ is , R^{2A} and R^{2B} are not both Cl.

[1638] **Embodiment 99:** A compound having the structure of formula (IB):



[1639] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1640] A and B are each independently aryl or heteroaryl;

[1641] C is a 3- to 10-membered ring;

[1642] X is a bond, -(CR⁵R⁶)_n-, -O-, -C(=O)-, -S-, -S(=O)-, -SO₂-, -NR⁷-, -N(R⁷)CO-, -CON(R⁷)-, or -NSO₂R⁷-;

[1643] Y is a bond, -(CR⁸R⁹)_m-, -O-, -S-, -S(=O)-, -SO₂-, or -NR⁷-;

[1644] W is a bond, -(CR^{8a}R^{9a})_m-, -N(R⁷)CO-, -CONR⁷-, or -NSO₂R⁷-;

[1645] Z is a bond, -CH₂-, -C(CH₃)H-, -O-, -S-, -NH-, -NCH₃-, or -N(COCH₃)-;

[1646] V is -CH₂-, -CH₂CH₂-, -CH(CH₃)CH₂-, -CH₂CH(CH₃)-, or -CH₂CH₂CH₂-;

[1647] L is hydrogen, halogen, -CF₂R¹⁰, -CF₃, -CN, -OR¹⁰, -NR¹¹R¹², or -CONR¹¹R¹²;

[1648] R¹ and R² are each independently hydrogen, deuterium, halogen, -CN, -CF₃, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[1649] R³ is hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -SR¹⁶, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆

alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COOR¹⁶, -NR¹⁴COR¹⁶, -NR¹⁴CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[1650] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁵ and R⁶ taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[1651] R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[1652] R^{8a} and R^{9a} are each independently hydrogen, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^{8a} and R^{8b} taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[1653] R⁷, R¹⁰ and R¹⁶ are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkyl-NH₂, C₁-C₆ haloalkyl, C₂-C₆ haloalkenyl, C₂-C₆ haloalkynyl, optionally substituted carbocyclyl, optionally substituted -CO(C₁-C₆ alkyl), -CO(optionally substituted heterocyclyl), optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R⁷ and R^{8a} taken together form an optionally substituted heterocyclyl;

[1654] R^{11} , R^{12} , R^{13} , R^{14} and R^{15} are each independently hydrogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted $-\text{COO}(C_1\text{-}C_6 \text{ alkyl})$, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or (R^{11} and R^{12}) or (R^{14} and R^{15}) or (R^{14} and R^{16}) taken together form an optionally substituted heterocyclyl;


[1655] each m is independently 0, 1 or 2;

[1656] n_1 and n_2 are each independently 0, 1, 2, 3, or 4;

[1657] n_3 is 0, 1, 2, 3, 4 or 5; and

[1658] each t is independently 0, 1 or 2.

[1659] **Embodiment 100:** A compound having the structure of formula (IIA):



[1660] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1661] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[1662] C is a 3- to 10-membered ring;

[1663] X is a bond, $-(\text{CR}^5\text{R}^6)_t-$, or $-\text{NR}^7-$;

[1664] Y is a bond, $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{NCH}_3-$, or $-\text{N}(\text{COCH}_3)-$;

[1665] Z is a bond, $-(\text{CR}^8\text{R}^9)_m-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(=\text{O})-$, $-\text{SO}_2-$, or $-\text{NR}^7-$;

[1666] W is a bond, $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{NHCO}-$, $-\text{N}(C_1\text{-}C_3 \text{ alkyl})\text{CO}-$, or $-\text{CONH}-$, or $-\text{CON}(C_1\text{-}C_3 \text{ alkyl})-$;

[1667] V is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}(\text{CH}_3)\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_3)-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$;

[1668] L is hydrogen, halogen, $-\text{CF}_2\text{R}^{10}$, $-\text{CF}_3$, $-\text{CN}$, $-\text{OR}^{10}$, $-\text{NR}^{11}\text{R}^{12}$, or $-\text{CONR}^{11}\text{R}^{12}$;

[1669] R^1 and R^2 are each independently hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{OH}$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1\text{-}C_6 \text{ alkyl})-(C_1\text{-}C_6 \text{ alkoxy})$, optionally substituted $-(C_1\text{-}C_6 \text{ alkyl})-\text{OH}$, $-\text{NR}^{13}\text{R}^{14}$, optionally substituted $-(C_1\text{-}C_6 \text{ alkyl})-\text{NR}^{13}\text{R}^{14}$, $-\text{NR}^{14}\text{SO}_2\text{R}^{16}$, optionally

substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;

[1670] R^3 is selected from hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, $C_2-C_3 \text{ alkynyl}$, $C_1-C_3 \text{ alkoxy}$, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;

[1671] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, $C_2-C_3 \text{ alkynyl}$, or $C_1-C_3 \text{ alkoxy}$; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1672] R^8 and R^9 are each independently hydrogen, halogen, or $C_1-C_3 \text{ alkyl}$;

[1673] R^{11} , R^{12} , R^{13} , R^{14} and R^{15} are each independently hydrogen, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, or $C_2-C_3 \text{ alkynyl}$; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[1674] R^7 , R^{10} and R^{16} are each independently hydrogen, $C_1-C_6 \text{ alkyl}$, $C_2-C_6 \text{ alkenyl}$, $C_2-C_6 \text{ alkynyl}$ or $C_1-C_6 \text{ alkyl-NH}_2$; or R^{14} and R^{16} taken together form a 3- to 6-membered heterocyclyl;


[1675] each m is independently 0, 1 or 2;

[1676] n_1 and n_2 are each independently 0, 1, or 2;

[1677] n_3 is 0, 1, 2, 3, 4 or 5; and

[1678] t is 0, 1 or 2.

[1679] **Embodiment 101:** A compound having the structure of formula (IIB):



[1680] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

- [1681] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;
- [1682] C is a 3- to 10-membered ring;
- [1683] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7-$;
- [1684] Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;
- [1685] Z is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, or $-NR^7-$;
- [1686] W is a bond, $-CH_2-$, or $-C(CH_3)H-$;
- [1687] V is a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;
- [1688] L is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;
- [1689] R^1 and R^2 are each independently hydrogen, deuterium, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)-(C_1 - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} ;
- [1690] R^3 is selected from hydrogen, halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1$ - C_3 alkyl), C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1$ - C_3 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1$ - C_3 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $SO_2NR^{14}R^{15}$, $-SO_2(C_1$ - C_3 alkyl), or $-(C_1$ - C_6 alkyl)- $SO_2(C_1$ - C_3 alkyl);
- [1691] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, or C_1 - C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [1692] R^8 and R^9 are each independently hydrogen, halogen, or C_1 - C_3 alkyl;
- [1693] R^{8a} and R^{9a} are each independently hydrogen, $-OH$, halogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1$ - C_3 alkyl)- $NR^{13}R^{14}$, $-NR^{14}COR^{16}$, $-(C_1$ - C_3 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, or $-(C_1$ - C_3 alkyl)- $CONR^{14}R^{15}$; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;
- [1694] R^{11} , R^{12} , R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[1695] R^7 , R^{10} and R^{16} are each independently hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl or C_1 - C_6 alkyl- NH_2 ; or R^{14} and R^{16} taken together form a 3- to 6-membered heterocyclyl;


[1696] each m is independently 0, 1 or 2;

[1697] n_1 and n_2 are each independently 0, 1, or 2;

[1698] n_3 is 1, 2, 3, 4 or 5; and

[1699] t is 0, 1 or 2.

[1700] **Embodiment 102:** A compound having the structure of formula (IC):



[1701] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1702] A and B are each independently aryl or heteroaryl;

[1703] C is a 3- to 10-membered ring;

[1704] X is a bond, $-(CR^5R^6)_t$ -, $-O$ -, $-C(=O)$ -, $-S$ -, $-S(=O)$ -, $-SO_2$ -, $-NR^7$ -, $-N(R^7)CO$ -, $-CON(R^7)$ -, or $-NSO_2R^7$;

[1705] Y is a bond, $-(CR^8R^9)_m$ -, $-O$ -, $-S$ -, $-S(=O)$ -, $-SO_2$ -, or $-NR^7$;

[1706] W is a bond, $-(CR^8R^9)_m$ -, $-C(=O)$ -, $N(R^7)CO$ -, $-CONR^7$ -, or $-NSO_2R^7$;

[1707] Z is a bond, $-(CR^8R^9)_m$ -, $-O$ -, $-S$ -, $-S(=O)$ -, $-SO_2$ -, or $-NR^7$;

[1708] V is $-CH_2$ -, $-CH_2CH_2$ -, $-CH(CH_3)CH_2$ -, $-CH_2CH(CH_3)$ -, or $-CH_2CH_2CH_2$;

[1709] L is hydrogen, halogen, $-CF_2R^{10}$, $-CF_3$, $-CCl_2R^{10}$, $-CCl_3$, $-CN$, $-OR^{10}$, $-NR^{11}R^{12}$, or $-CONR^{11}R^{12}$;

[1710] R^1 and R^2 are each independently hydrogen, deuterium, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)-(C_1 - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} , optionally substituted carbocyclyl,

optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[1711] R^3 is hydrogen, halogen, oxo, =S, =NR¹⁶, -CN, -CF₃, -OH, -SR¹⁶, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, optionally substituted -(C₁-C₆ alkyl)-(C₁-C₆ alkoxy), optionally substituted -(C₁-C₆ alkyl)-OH, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)NR¹⁴SO₂R¹⁶, -NR¹⁴COOR¹⁶, -NR¹⁴COR¹⁶, -NR¹⁴CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-SO₂NR¹⁴R¹⁵, optionally substituted -SO₂R¹⁶, optionally substituted -(C₁-C₆ alkyl)-SO₂R¹⁶, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl;

[1712] R^5 and R^6 are each independently hydrogen, halogen, -OH, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹³R¹⁴, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^5 and R^6 taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[1713] R^8 and R^9 are each independently hydrogen, halogen, or C₁-C₃ alkyl;

[1714] R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkoxy, -NR¹³R¹⁴, optionally substituted -(C₁-C₆ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, optionally substituted -(C₁-C₆ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, optionally substituted -(C₁-C₆ alkyl)-CONR¹⁴R¹⁵, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^{8a} and R^{8b} taken together form an optionally substituted carbocyclyl or optionally substituted heterocyclyl;

[1715] R^7 , R^{10} and R^{16} are each independently hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ alkyl-NH₂, C₁-C₆ haloalkyl, C₂-C₆ haloalkenyl, C₂-C₆ haloalkynyl, optionally substituted carbocyclyl, optionally substituted -CO(C₁-C₆ alkyl), -

CO(optionally substituted heterocyclyl), optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or R^7 and R^{8a} taken together form an optionally substituted heterocyclyl;

[1716] R^{11} , R^{12} , R^{13} , R^{14} and R^{15} are each independently hydrogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted $-COO(C_1$ - C_6 alkyl), optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; or (R^{11} and R^{12}) or (R^{14} and R^{15}) or (R^{14} and R^{16}) taken together form an optionally substituted heterocyclyl;


[1717] each m is independently 0, 1 or 2;

[1718] n_1 and n_2 are each independently 0, 1, 2, 3, or 4;

[1719] n_3 is 0, 1, 2, 3, 4 or 5; and

[1720] each t is independently 0, 1 or 2.

[1721] **Embodiment 103:** A compound having a structure of formula (IIIA):



[1722] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1723] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[1724] C is a 3- to 10-membered ring;

[1725] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;

[1726] Y is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, $-NR^7-$ or $-N(COCH_3)-$;

[1727] W is a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

[1728] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[1729] V is $-CH_2-$ and L is halogen, $-NH_2$, $-CHCl_2$, $-CCl_3$, or $-CF_3$; or

[1730] V is $-CH_2CH_2-$ and L is halogen or $-NH_2$;

[1731] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally

substituted $-(C_1-C_6 \text{ alkyl})-(C_1-C_6 \text{ alkoxy})$, optionally substituted $-(C_1-C_6 \text{ alkyl})-OH$, $-NR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;

[1732] R^3 is selected from halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, $C_2-C_3 \text{ alkynyl}$, $C_1-C_3 \text{ alkoxy}$, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;

[1733] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, $C_2-C_3 \text{ alkynyl}$, or $C_1-C_3 \text{ alkoxy}$; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1734] R^7 is hydrogen, $C_1-C_6 \text{ alkyl}$, $C_2-C_6 \text{ alkenyl}$, or $C_2-C_6 \text{ alkynyl}$;

[1735] R^8 and R^9 are each independently hydrogen, halogen, or $C_1-C_3 \text{ alkyl}$;

[1736] R^{8a} and R^{9a} are each independently hydrogen, $-OH$, halogen, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, $C_2-C_3 \text{ alkynyl}$, $C_1-C_3 \text{ alkoxy}$, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}COR^{16}$, $-(C_1-C_3 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, or $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1737] R^{13} , R^{14} and R^{15} are each independently hydrogen, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, or $C_2-C_3 \text{ alkynyl}$; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[1738] R^{16} is hydrogen, optionally substituted $C_1-C_3 \text{ alkyl}$, optionally substituted $C_2-C_3 \text{ alkenyl}$, optionally substituted $C_2-C_3 \text{ alkynyl}$, $C_3-C_6 \text{ cycloalkyl}$, or phenyl;


[1739] each m is independently 0, 1 or 2;

[1740] n_1 and n_2 are each independently 0, 1, or 2;


[1741] n_3 is 1, 2, 3, 4 or 5; and

[1742] t is 0, 1 or 2.

[1743] **Embodiment 104:** A compound having a structure of formula (G-II)



[1744] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



[1746] X is $-(CR^3R^6)_t-$;

[1747] Y is $-O-$;

[1748] Z is $-O-$;

[1749] W is $-CH_2-$ or $-C(CH_3)H-$;

[1750] V is $-CH_2CH_2-$;

[1751] L is halogen;

[1752] R^1 and R^2 are each independently Cl or $-CN$;

[1753] at least one R^3 is selected from $-CN$, C_1 - C_3 alkoxy, $-CONH_2$, $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, or $-SO_2CH_3$ and the other R^3 , if present, is selected from $-CN$, $-CF_3$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-S(C_1$ - C_3 alkyl), $-SO_2(C_1$ - C_3 alkyl), $-NH_2$, $-(C_1$ - C_3 alkyl) NH_2 , $-NHSO_2CH_3$, $-NHSO_2CF_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1$ - C_3 alkyl) $_2$, $-CONH(C_1$ - C_3 alkyl), $-NHCO(C_1$ - C_3 alkyl), $-N(CH_3)COO(C_1$ - C_3 alkyl), $-NHCO(C_1$ - C_3 alkyl), or $-N(CH_3)COO(C_1$ - C_3 alkyl);

[1754] R⁵ and R⁶ are each independently hydrogen or methyl;

[1755] n₁ and n₂ are each independently 0, 1, or 2;

[1756] n₃ is 1 or 2; and

[1757] t is 1.

[1758] **Embodiment 105:** The compound of any one of **Embodiments 1-104**, wherein at least one of R³ is -CN, -OCH₃, -SO₂CH₃, -NHSO₂CH₃, -CH₂NHSO₂CH₃, -SO₂NH₂, -CONH₂, or -NHCOCH₃.

[1759] **Embodiment 106:** A compound selected from Compounds **A212-A234** or a pharmaceutically acceptable salt thereof.

[1760] **Embodiment 107:** A pharmaceutical composition comprising a compound of any one of **Embodiments 1-106** and a pharmaceutically acceptable carrier.

[1761] **Embodiment 108:** The pharmaceutical composition of **Embodiment 107**, further comprising one or more additional therapeutic agents.

[1762] **Embodiment 109:** The pharmaceutical composition of **Embodiment 108**, wherein the one or more additional therapeutic agents is for treating prostate cancer, breast cancer, ovarian cancer, bladder cancer, pancreatic cancer, hepatocellular cancer, endometrial cancer, salivary gland carcinoma, hair loss, acne, hirsutism, ovarian cysts, polycystic ovary disease, precocious puberty, spinal and bulbar muscular atrophy, or age-related macular degeneration.

[1763] **Embodiment 110:** The pharmaceutical composition of **Embodiment 108**, wherein the one or more additional therapeutic agents is a poly (ADP-ribose) polymerase (PARP) inhibitor including but not limited to olaparib, niraparib, rucaparib, talazoparib; an androgen receptor ligand binding domain inhibitor including but not limited to enzalutamide, apalutamide, darolutamide, bicalutamide, nilutamide, flutamide, ODM-204, TAS3681; an inhibitor of CYP17 including but not limited to galeterone, abiraterone, abiraterone acetate; a microtubule inhibitor including but not limited to docetaxel, paclitaxel, cabazitaxel (XRP-6258); a modulator of PD-1 or PD-L1 including but not limited to pembrolizumab, durvalumab, nivolumab, atezolizumab; a gonadotropin releasing hormone agonist including but not limited to cyproterone acetate, leuprolide; a 5- α reductase inhibitor including but not limited to finasteride, dutasteride, turosteride, bexlosteride, izonsteride, FCE 28260, SKF105,111; a vascular endothelial growth factor inhibitor including but not limited to bevacizumab (Avastin); a histone deacetylase inhibitor including but not limited to OSU-HDAC42; an integrin α -v- β -3 inhibitor including but not limited to VITAXIN; a receptor tyrosine kinase inhibitor including but not limited to sunitumib; a phosphoinositide 3-kinase inhibitor including but not limited to alpelisib, buparlisib, idealisib; an anaplastic lymphoma kinase

(ALK) inhibitor including but not limited to crizotinib, alectinib; an endothelin receptor A antagonist including but not limited to ZD-4054; an anti-CTLA4 inhibitor including but not limited to MDX-010 (ipilimumab); an heat shock protein 27 (HSP27) inhibitor including but not limited to OGX 427; an androgen receptor degrader including but not limited to ARV-330, ARV-110; a androgen receptor DNA-binding domain inhibitor including but not limited to VPC-14449; a bromodomain and extra-terminal motif (BET) inhibitor including but not limited to BI-894999, GSK25762, GS-5829; an androgen receptor N-terminal domain inhibitor including but not limited to a sintokamide; an alpha-particle emitting radioactive therapeutic agent including but not limited to radium 233 or a salt thereof, niclosamide; or related compounds thereof; a selective estrogen receptor modulator (SERM) including but not limited to tamoxifen, raloxifene, toremifene, arzoxifene, bazedoxifene, pipindoxifene, lasofoxifene, enclomiphene; a selective estrogen receptor degrader (SERD) including but not limited to fulvestrant, ZB716, OP-1074, elacestrant, AZD9496, GDC0810, GDC0927, GW5638, GW7604; an aromatase inhibitor including but not limited to anastrozole, exemestane, letrozole; selective progesterone receptor modulators (SPRM) including but not limited to mifepristone, lonaprisone, onaprisone, asoprisnil, lonaprisnil, ulipristal, telapristone; a glucocorticoid receptor inhibitor including but not limited to mifepristone, COR108297, COR125281, ORIC-101, PT150; CDK4/6 inhibitors including palbociclib, abemaciclib, ribociclib; HER2 receptor antagonist including but not limited to trastuzumab, neratinib; or a mammalian target of rapamycin (mTOR) inhibitor including but not limited to everolimus, temsirolimus.

[1764] Embodiment 111: A method for modulating androgen receptor activity, comprising administering a compound of any one of **Embodiments 1-106**, to a subject in need thereof.

[1765] Embodiment 112: The method of any one of **Embodiment 111**, wherein the modulating androgen receptor activity is for treating a condition or disease selected from prostate cancer, breast cancer, ovarian cancer, bladder cancer, pancreatic cancer, hepatocellular cancer, endometrial cancer, salivary gland carcinoma, hair loss, acne, hirsutism, ovarian cysts, polycystic ovary disease, precocious puberty, spinal and bulbar muscular atrophy, or age-related macular degeneration.

[1766] Embodiment 113: A method for treating cancer, comprising administering a compound of any one of **Embodiments 1-106**, to a subject in need thereof.

[1767] Embodiment 114: The method of **Embodiment 113**, wherein the cancer is selected from prostate cancer, breast cancer, ovarian cancer, bladder cancer, pancreatic cancer, hepatocellular cancer, endometrial cancer, or salivary gland carcinoma.

[1768] **Embodiment 115:** The method of **Embodiment 113**, wherein the cancer is prostate cancer.

[1769] **Embodiment 116:** The method of **Embodiment 115**, wherein the prostate cancer is primary or localized prostate cancer, locally advanced prostate cancer, recurrent prostate cancer, advanced prostate cancer, metastatic prostate cancer, metastatic castration-resistant prostate cancer, and hormone-sensitive prostate cancer. The method of **Embodiment 115**, wherein the prostate cancer is primary or localized prostate cancer, locally advanced prostate cancer, recurrent prostate cancer, advanced prostate cancer, metastatic prostate cancer, non-metastatic castration-resistant prostate cancer, metastatic castration-resistant prostate cancer, and hormone-sensitive prostate cancer.

[1770] **Embodiment 117:** The method of **Embodiment 115**, wherein the prostate cancer is metastatic castration-resistant prostate cancer.

[1771] **Embodiment 118:** The method of **Embodiment 115**, wherein the prostate cancer expresses full-length androgen receptor or truncated androgen receptor splice variant.

[1772] **Embodiment 119:** The compound of any one of **Embodiments 1-106**, wherein the compound has an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 1000 nM.

[1773] **Embodiment 120:** The compound of any one of **Embodiments 1-106**, wherein the compound has an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 900 nM.

[1774] **Embodiment 121:** The compound of any one of **Embodiments 1-106**, wherein the compound has an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 800 nM.

[1775] **Embodiment 122:** The compound of any one of **Embodiments 1-106**, wherein the compound has an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 700 nM.

[1776] **Embodiment 123:** The compound of any one of **Embodiments 1-106**, wherein the compound has an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 600 nM.

[1777] **Embodiment 124:** The compound of any one of **Embodiments 1-106**, wherein the compound has an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 500 nM.

[1778] **Embodiment 125:** The compound of any one of **Embodiments 1-106**, wherein the compound has an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 400 nM.

[1779] **Embodiment 126:** The compound of any one of **Embodiments 1-106**, wherein the compound has an *in vitro* IC₅₀ in a PSA-luciferase assay of less than about 300 nM.

[1780] **Embodiment 127:** The compound of any one of **Embodiments 1-106 or 119-126**, wherein the compound has an *in vitro* metabolic clearance of less than 12 μ L/min/mg protein in mammalian microsomes.

[1781] **Embodiment 128:** The compound of **Embodiment 126**, wherein the mammalian microsomes are human microsomes.

[1782] **Embodiment 129:** The compound of any one of **Embodiments 1-106 or 119-128**, wherein the compound has an *in vitro* metabolic clearance in mammalian hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million cells}$.

[1783] **Embodiment 130:** The compound of **Embodiment 128**, wherein the mammalian microsomes are human microsomes.

[1784] **Embodiment 131:** The compound of any one of **Embodiment 1-106**, wherein the compound has:

[1785] (a) an *in vitro* IC_{50} in a PSA-luciferase assay of less than about 1000 nM;

[1786] (b) an *in vitro* metabolic clearance of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein in mammalian microsomes; and

[1787] (c) *in vitro* metabolic clearance in mammalian hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million cells}$.

[1788] **Embodiment 132:** The compound of any one of **Embodiments 1-106**, wherein the compound has:

[1789] (a) an *in vitro* IC_{50} in a PSA-luciferase assay of less than about 1000 nM;


[1790] (b) an *in vitro* metabolic clearance of less than 12 $\mu\text{L}/\text{min}/\text{mg}$ protein in human microsomes; and

[1791] (c) *in vitro* metabolic clearance in human hepatocytes of less than 4 $\mu\text{L}/\text{min}/\text{million cells}$.

[1792] **Embodiment 133:** The compounds of any one of **Embodiments 119-132**, wherein the PSA-luciferase assay is performed as described in Example 35.

[1793] **Embodiment 134:** The compounds of any one of **Embodiments 119-132**, wherein the metabolic clearance is measured according to Example 36.

[1794] **Embodiment 135:** A compound having the structure of formula (IIIA):



[1795] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1796] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[1797] C is a 3- to 10-membered ring;

[1798] X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;

[1799] Y is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, $-NR^7-$, or $-N(COCH_3)-$;

[1800] W is a bond, $-(CR^8R^9)_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

[1801] Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[1802] V is $-CH_2-$ and L is halogen, $-NH_2$, $-CHCl_2$, $-CCl_3$, or $-CF_3$; or

[1803] V is $-CH_2CH_2-$ and L is halogen or $-NH_2$;

[1804] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)-(C_1 - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} ;

[1805] R^3 is selected from halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1$ - C_3 alkyl), C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1$ - C_3 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1$ - C_3 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $SO_2NR^{14}R^{15}$, $-SO_2(C_1$ - C_3 alkyl), or $-(C_1$ - C_6 alkyl)- $SO_2(C_1$ - C_3 alkyl);

[1806] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, or C_1 - C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1807] R^7 is hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

[1808] R^8 and R^9 are each independently hydrogen, halogen, or C_1 - C_3 alkyl;

[1809] R^{8a} and R^{9a} are each independently hydrogen, $-OH$, halogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1$ - C_3 alkyl)- $NR^{13}R^{14}$, $-NR^{14}COR^{16}$, $-(C_1$ - C_3 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, or $-(C_1$ - C_3 alkyl)- $CONR^{14}R^{15}$; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1810] R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[1811] R^{16} is hydrogen, optionally substituted C_1 - C_3 alkyl, optionally substituted C_2 - C_3 alkenyl, optionally substituted C_2 - C_3 alkynyl, C_3 - C_6 cycloalkyl, or phenyl;


[1812] each m is independently 0, 1, or 2;

[1813] n_1 and n_2 are each independently 0, 1, or 2;

[1814] n_3 is 1, 2, 3, 4 or 5; and

[1815] t is 0, 1 or 2.

[1816] **Embodiment 136:** The compound of embodiment 135, wherein the compound has the structure of formula (IVA):



[1817] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1818] A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

[1819] C is a 3- to 10-membered ring;

[1820] X is a bond, $-(CR^5R^6)_t$, or $-NR^7$;

[1821] Y and Z are each independently a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

[1822] W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

[1823] V is $-CH_2-$ and L is halogen, $-NH_2$, or $-CF_3$; or

[1824] V is $-CH_2CH_2-$ and L is halogen or $-NH_2$;

[1825] R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)- $(C_1$ - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl) $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally

substituted $-(C_1-C_6 \text{ alkyl})-SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$, optionally substituted $-(C_1-C_6 \text{ alkyl})-SO_2R^{16}$;

[1826] R^3 is selected from halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1-C_3 \text{ alkyl})$, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, $C_2-C_3 \text{ alkynyl}$, $C_1-C_3 \text{ alkoxy}$, $-NR^{13}R^{14}$, $-(C_1-C_3 \text{ alkyl})-NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1-C_3 \text{ alkyl})NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1-C_6 \text{ alkyl})-NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1-C_3 \text{ alkyl})-SO_2NR^{14}R^{15}$, $-SO_2(C_1-C_3 \text{ alkyl})$, or $-(C_1-C_6 \text{ alkyl})-SO_2(C_1-C_3 \text{ alkyl})$;

[1827] R^5 and R^6 are each independently hydrogen, halogen, $-OH$, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, $C_2-C_3 \text{ alkynyl}$, or $C_1-C_3 \text{ alkoxy}$; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

[1828] R^7 is H or $C_1-C_6 \text{ alkyl}$;

[1829] R^{13} , R^{14} and R^{15} are each independently hydrogen, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, or $C_2-C_3 \text{ alkynyl}$; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

[1830] R^{16} is hydrogen, $C_1-C_3 \text{ alkyl}$, $C_2-C_3 \text{ alkenyl}$, or $C_2-C_3 \text{ alkynyl}$;

[1831] n_1 and n_2 are each independently 0, 1, or 2;

[1832] n_3 is 1, 2, 3, 4 or 5; and


[1833] t is 0, 1 or 2.


[1834] **Embodiment 137:** The compound of embodiment 135 or 136, wherein C is 5- to 10-membered heteroaryl or aryl

[1835] **Embodiment 138:** The compound of embodiment 137, wherein C is 5- to 7-membered heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

[1836] **Embodiment 139:** The compound of embodiment 138, wherein C, which is substituted with $(R^3)_{n_3}$, is pyrazole, imidazole, oxazole, oxadiazole, oxazolone, isoxazole, thiazole, pyridyl, pyrazine, furan or pyrimidyl.

[1837] **Embodiment 140:** The compound of any one of embodiments 135-139, wherein C, which is substituted with $(R^3)_{n_3}$, is selected from





C₁-C₃ alkyl.

[1838] **Embodiment 141:** The compound of any one of **Embodiments 135-140**, wherein R¹ and R² are each independently Cl, -CN, -CF₃, -OH, methyl, methoxy, or -CONH₂.

[1839] **Embodiment 142:** The compound of any one of **Embodiments 135-141**, wherein:

[1840] A and B are phenyl;

[1841] X is -(CR⁵R⁶)_l-;

[1842] Y and Z are each -O-;

[1843] V is -CH₂- or -CH₂CH₂-;

[1844] L is halogen;

[1845] R¹ and R² are each independently hydrogen, halogen, -CN, -CF₃, -OH, or optionally substituted C₁-C₆ alkyl;

[1846] R⁵ and R⁶ are each independently hydrogen, halogen, -OH, or C₁-C₃ alkyl; and

[1847] R¹⁶ is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl.

[1848] **Embodiment 143:** The compound of any one of **Embodiments 135-142**, wherein:


[1849] R⁵ and R⁶ are each independently hydrogen, or C₁-C₃ alkyl;

[1850] W is -CH₂- or -C(CH₃)H-;

[1851] V is -CH₂CH₂-; and

[1852] R^1 and R^2 are each independently hydrogen, halogen, or $-\text{CN}$.

[1853] **Embodiment 144:** The compound of any one of **Embodiments 135-143**, wherein the compound has the structure of formula (A-I):



[1854] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

[1855] C is a 5- to 7-membered monocyclic heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member;

[1856] X is a bond, $-(\text{CR}^5\text{R}^6)_t-$, or $-\text{NR}^7-$;

[1857] Y is a bond, $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{NCH}_3-$, or $-\text{N}(\text{COCH}_3)-$;

[1858] Z is a bond, $-\text{CH}_2-$, $-\text{O}-$, or $-\text{NH}-$;

[1859] W is a bond, $-\text{CH}_2-$, $-\text{C}(\text{CH}_3)\text{H}-$, $-\text{C}(=\text{O})-$, $-\text{N}(\text{R}^7)\text{CO}-$, or $-\text{CONR}^7-$;

[1860] V is $-\text{CH}_2-$ and L is halogen, $-\text{NH}_2$, or $-\text{CF}_3$; or

[1861] V is $-\text{CH}_2\text{CH}_2-$ and L is halogen or $-\text{NH}_2$;

[1862] R^1 and R^2 are each independently hydrogen, halogen, $-\text{CN}$, $-\text{CF}_3$, methyl, or $-\text{CONH}_2$;

[1863] R^3 is selected from $-\text{CN}$, C_1 - C_3 alkoxy, CF_3 , C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, $-\text{S}(\text{C}_1$ - C_3 alkyl), $-\text{SO}_2(\text{C}_1$ - C_3 alkyl), $-\text{NH}_2$, $-(\text{C}_1$ - C_3 alkyl) NH_2 , $-\text{NHSO}_2\text{CH}_3$, $-\text{NHSO}_2\text{CF}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{NHSO}_2\text{CH}_2\text{CH}_3$, $-\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{NHSO}_2\text{CH}_3$, $-\text{CH}_2\text{N}(\text{CH}_3)\text{SO}_2\text{CH}_3$, $-\text{SO}_2\text{NH}_2$, $-\text{CONH}_2$, $-\text{CON}(\text{C}_1$ - C_3 alkyl) $_2$, $-\text{CONH}(\text{C}_1$ - C_3 alkyl), $-\text{NHCO}(\text{C}_1$ - C_3 alkyl), $-\text{N}(\text{CH}_3)\text{COO}(\text{C}_1$ - C_3 alkyl), $-\text{NHCO}(\text{C}_1$ - C_3 alkyl), or $-\text{N}(\text{CH}_3)\text{COO}(\text{C}_1$ - C_3 alkyl);

[1864] R^5 and R^6 are each independently hydrogen, halogen, $-\text{OH}$, or C_1 - C_3 alkyl;

[1865] R^7 is H or C_1 - C_6 alkyl;

[1866] n_1 and n_2 are each independently 0, 1, or 2;

[1867] n_3 is 1, 2, 3, 4 or 5; and

[1868] t is 0, 1 or 2.

[1869] **Embodiment 145:** The compound of any one of **Embodiments 135-144**, wherein: at least one R^3 is selected from -CN, C₁-C₃ alkoxy, -CONH₂, -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, or -SO₂CH₃ and the other R^3 , if present, is selected from -CN, -CF₃, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -(C₁-C₃ alkyl)NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl).

[1870] **Embodiment 146:** The compound of any one of **Embodiments 135-145**, wherein:

[1871] X is a bond or -(CR⁵R⁶)_i;

[1872] W is a bond, -CH₂-, or -C(CH₃)H-;


[1873] Y is -O-;

[1874] Z is -O-;


[1875] V is -CH₂- or -CH₂CH₂-; and


[1876] L is halogen.

[1877] **Embodiment 147:** The compound of any one of **Embodiments 135-146**, wherein the compound has the structure of formula (G-II):



[1878] or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:





[1880] X is $-(CR^5R^6)_t-$;

[1881] Y is $-O-$;

[1882] Z is $-O-$;

[1883] W is $-CH_2-$ or $-C(CH_3)H-$;

[1884] V is $-CH_2CH_2-$;

[1885] L is halogen;

[1886] R^1 and R^2 are each independently Cl or $-CN$;

[1887] at least one R^3 is selected from $-CN$, C_1 - C_3 alkoxy, $-CONH_2$, $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, or $-SO_2CH_3$ and the other R^3 , if present, is selected from $-CN$, $-CF_3$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-S(C_1$ - C_3 alkyl), $-SO_2(C_1$ - C_3 alkyl), $-NH_2$, $-(C_1$ - C_3 alkyl) NH_2 , $-NHSO_2CH_3$, $-NHSO_2CF_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1$ - C_3 alkyl) $_2$, $-CONH(C_1$ - C_3 alkyl), $-NHCO(C_1$ - C_3 alkyl), $-N(CH_3)COO(C_1$ - C_3 alkyl), $-NHCO(C_1$ - C_3 alkyl), or $-N(CH_3)COO(C_1$ - C_3 alkyl);

[1888] R^5 and R^6 are each independently hydrogen or methyl;

[1889] n_1 and n_2 are each independently 0, 1, or 2;


[1890] n_3 is 1 or 2; and


[1891] t is 1.

[1892] **Embodiment 148:** The compound of any one of **Embodiments 135-147**, wherein: at least one R^3 is selected from $-NHSO_2CH_3$, $-NHSO_2CH_2CH_3$, or $-SO_2CH_3$ and the other R^3 , if present, is selected from $-CN$, C_1 - C_3 alkyl, C_1 - C_3 alkoxy, $-SO_2(C_1$ - C_3 alkyl), $-NH_2$, $-(C_1$ - C_3 alkyl) NH_2 , $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1$ - C_3 alkyl) $_2$, $-CONH(C_1$ - C_3 alkyl), $-NHCO(C_1$ - C_3 alkyl), $-N(CH_3)COO(C_1$ - C_3 alkyl), $-NHCO(C_1$ - C_3 alkyl), or $-N(CH_3)COO(C_1$ - C_3 alkyl).

[1893] **Embodiment 149:** A compound selected from **Compounds A1-A96**, **A98-A116**, **A118-A159**, **A161-A175**, or **A177-A234** or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

[1894] **Embodiment 150:** The compound of **Embodiment 149**, wherein the compound is selected from





, or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof.

[1895] **Embodiment 151:** A pharmaceutical composition comprising a compound of any one of **Embodiments 135-150** and a pharmaceutically acceptable carrier

[1896] **Embodiment 152:** A method for treating cancer, comprising administering the compound, pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug of any one of the compound of **Embodiments 135-150** or any pharmaceutical composition of **Embodiment 151**, to a subject in need thereof.

[1897] **Embodiment 153:** The method of **Embodiment 152**, wherein the cancer is prostate cancer.

[1898] **Embodiment 154:** The method of **Embodiment 153**, wherein the prostate cancer is metastatic castration-resistant prostate cancer.


[1899] **Embodiment 155:** The method of **Embodiment 154**, wherein the prostate cancer expresses full-length androgen receptor or truncated androgen receptor splice variant.

[1900] The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention.

[1901] While the invention has been described in connection with proposed specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth and as follows in the scope of the appended claims.

WHAT IS CLAIMED IS:

1. A compound having the structure of formula (IIIA):



or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

C is a 3- to 10-membered ring;

X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;

Y is a bond, $-(CR^8R^9)_m-$, $-O-$, $-S-$, $-S(=O)-$, $-SO_2-$, $-NR^7-$, or $-N(COCH_3)-$;

W is a bond, $-(CR^{8a}R^{9a})_m-$, $-C(=O)-$, $-N(R^7)CO-$, $-CONR^7-$, or $-NSO_2R^7-$;

Z is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

V is $-CH_2-$ and L is halogen, $-NH_2$, $-CHCl_2$, $-CCl_3$, or $-CF_3$; or

V is $-CH_2CH_2-$ and L is halogen or $-NH_2$;

R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)- $(C_1$ - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$ or optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} ;

R^3 is selected from halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1$ - C_3 alkyl), C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1$ - C_3 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1$ - C_3 alkyl)- $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)-

CONR¹⁴R¹⁵, -SO₂NR¹⁴R¹⁵, -(C₁-C₃ alkyl)-SO₂NR¹⁴R¹⁵, -SO₂(C₁-C₃ alkyl), or -(C₁-C₆ alkyl)-SO₂(C₁-C₃ alkyl);

R⁵ and R⁶ are each independently hydrogen, halogen, -OH, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, or C₁-C₃ alkoxy; or R⁵ and R⁶ taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

R⁷ is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, or C₂-C₆ alkynyl;

R⁸ and R⁹ are each independently hydrogen, halogen, or C₁-C₃ alkyl;

R^{8a} and R^{9a} are each independently hydrogen, -OH, halogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -NR¹³R¹⁴, -(C₁-C₃ alkyl)-NR¹³R¹⁴, -NR¹⁴COR¹⁶, -(C₁-C₃ alkyl)-NR¹⁴COR¹⁶, -CONR¹⁴R¹⁵, or -(C₁-C₃ alkyl)-CONR¹⁴R¹⁵; or R^{8a} and R^{8b} taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

R¹³, R¹⁴ and R¹⁵ are each independently hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl; or R¹⁴ and R¹⁵ taken together form a 3- to 6-membered heterocyclyl;

R¹⁶ is hydrogen, optionally substituted C₁-C₃ alkyl, optionally substituted C₂-C₃ alkenyl, optionally substituted C₂-C₃ alkynyl, C₃-C₆ cycloalkyl, or phenyl;


each m is independently 0, 1, or 2;

n1 and n2 are each independently 0, 1, or 2;

n3 is 1, 2, 3, 4 or 5; and

t is 0, 1 or 2.

2. The compound of claim 1, wherein the compound has the structure of formula (IVA):



or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

A and B are each independently selected from phenyl, pyridyl, pyrimidyl, or thiophene;

C is a 3- to 10-membered ring;

X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;

Y and Z are each independently a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

V is $-CH_2-$ and L is halogen, $-NH_2$, or $-CF_3$; or

V is $-CH_2CH_2-$ and L is halogen or $-NH_2$;

R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, $-OH$, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 alkoxy, optionally substituted $-(C_1$ - C_6 alkyl)-(C_1 - C_6 alkoxy), optionally substituted $-(C_1$ - C_6 alkyl)- OH , $-NR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{13}R^{14}$, optionally substituted $-(C_1$ - C_6 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, optionally substituted $-(C_1$ - C_6 alkyl)- $SO_2NR^{14}R^{15}$, optionally substituted $-SO_2R^{16}$, optionally substituted $-(C_1$ - C_6 alkyl)- SO_2R^{16} ;

R^3 is selected from halogen, oxo, $=S$, $=NR^{16}$, $-CN$, $-CF_3$, $-OH$, $-S(C_1$ - C_3 alkyl), C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_1 - C_3 alkoxy, $-NR^{13}R^{14}$, $-(C_1$ - C_3 alkyl)- $NR^{13}R^{14}$, $-NR^{14}SO_2R^{16}$, $-(C_1$ - C_3 alkyl)- $NR^{14}SO_2R^{16}$, $-NR^{14}COR^{16}$, $-(C_1$ - C_6 alkyl)- $NR^{14}COR^{16}$, $-CONR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $CONR^{14}R^{15}$, $-SO_2NR^{14}R^{15}$, $-(C_1$ - C_3 alkyl)- $SO_2NR^{14}R^{15}$, $-SO_2(C_1$ - C_3 alkyl), or $-(C_1$ - C_6 alkyl)- $SO_2(C_1$ - C_3 alkyl)

R^5 and R^6 are each independently hydrogen, halogen, $-OH$, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, or C_1 - C_3 alkoxy; or R^5 and R^6 taken together form an optionally substituted 3- to 6-membered carbocyclyl or heterocyclyl;

R^7 is H or C_1 - C_6 alkyl;

R^{13} , R^{14} and R^{15} are each independently hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl; or R^{14} and R^{15} taken together form a 3- to 6-membered heterocyclyl;

R^{16} is hydrogen, C_1 - C_3 alkyl, C_2 - C_3 alkenyl, or C_2 - C_3 alkynyl;

n_1 and n_2 are each independently 0, 1, or 2;

n_3 is 1, 2, 3, 4 or 5; and


t is 0, 1 or 2.

3. The compound of claim 2, wherein C is 5- to 10-membered heteroaryl or aryl.


4. The compound of claim 3, wherein C is 5- to 7-membered heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member.

5. The compound of claim 4, wherein C, which is substituted with $(R^3)_{n3}$, is pyrazole, imidazole, oxazole, oxadiazole, oxazolone, isoxazole, thiazole, pyridyl, pyrazine, furan or pyrimidyl.

6. The compound of claim 5, wherein C, which is substituted with $(R^3)_{n3}$, is selected from



7. The compound of claim 6, wherein R^1 and R^2 are each independently Cl, -CN, -CF₃, -OH, methyl, methoxy, or -CONH₂.
8. The compound of claim 2, wherein:
 A and B are phenyl;
 X is -(CR⁵R⁶)_t;
 Y and Z are each -O-;
 V is -CH₂- or -CH₂CH₂-;
 L is halogen;
 R^1 and R^2 are each independently hydrogen, halogen, -CN, -CF₃, -OH, or optionally substituted C₁-C₆ alkyl;
 R^5 and R^6 are each independently hydrogen, halogen, -OH, or C₁-C₃ alkyl; and
 R^{16} is hydrogen, C₁-C₃ alkyl, C₂-C₃ alkenyl, or C₂-C₃ alkynyl.
9. The compound of claim 8, wherein:
 R^5 and R^6 are each independently hydrogen, or C₁-C₃ alkyl;
 W is -CH₂- or -C(CH₃)H-;
 V is -CH₂CH₂-; and
 R^1 and R^2 are each independently hydrogen, halogen, or -CN.
10. The compound of claim 1, wherein the compound has the structure of formula (A-I)



(A-I)

or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:

C is a 5- to 7-membered monocyclic heteroaryl comprising 1, 2, or 3 heteroatoms selected from O, S, or N as a ring member;

X is a bond, $-(CR^5R^6)_t-$, or $-NR^7$;

Y is a bond, $-CH_2-$, $-C(CH_3)H-$, $-O-$, $-S-$, $-NH-$, $-NCH_3-$, or $-N(COCH_3)-$;

Z is a bond, $-CH_2-$, $-O-$, or $-NH-$;

W is a bond, $-CH_2-$, $-C(CH_3)H-$, $-C(=O)-$, $-N(R^7)CO-$, or $-CONR^7-$;

V is $-CH_2-$ and L is halogen, $-NH_2$, or $-CF_3$; or

V is $-CH_2CH_2-$ and L is halogen or $-NH_2$;

R^1 and R^2 are each independently hydrogen, halogen, $-CN$, $-CF_3$, methyl, or $-CONH_2$;

R^3 is selected from $-CN$, C_1-C_3 alkoxy, $-CF_3$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, $-S(C_1-C_3 \text{ alkyl})$, $-SO_2(C_1-C_3 \text{ alkyl})$, $-NH_2$, $-(C_1-C_3 \text{ alkyl})NH_2$, $-NHSO_2CH_3$, $-NHSO_2CF_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1-C_3 \text{ alkyl})_2$, $-CONH(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, or $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$;

R^5 and R^6 are each independently hydrogen, halogen, $-OH$, or C_1-C_3 alkyl;

R^7 is H or C_1-C_6 alkyl;

n_1 and n_2 are each independently 0, 1, or 2;

n_3 is 1, 2, 3, 4 or 5; and

t is 0, 1 or 2.

11. The compound of claim 10, wherein:

at least one R^3 is selected from $-CN$, C_1-C_3 alkoxy, $-CONH_2$, $-NHSO_2CH_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, or $-SO_2CH_3$ and the other R^3 , if present, is selected from $-CN$, $-CF_3$, C_1-C_3 alkyl, C_2-C_3 alkenyl, C_2-C_3 alkynyl, C_1-C_3 alkoxy, $-S(C_1-C_3 \text{ alkyl})$, $-SO_2(C_1-C_3 \text{ alkyl})$, $-NH_2$, $-(C_1-C_3 \text{ alkyl})NH_2$, $-NHSO_2CH_3$, $-NHSO_2CF_3$, $-N(CH_3)SO_2CH_3$, $-NHSO_2CH_2CH_3$, $-N(CH_3)SO_2CH_2CH_3$, $-CH_2NHSO_2CH_3$, $-CH_2N(CH_3)SO_2CH_3$, $-SO_2NH_2$, $-CONH_2$, $-CON(C_1-C_3 \text{ alkyl})_2$, $-CONH(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$, $-NHCO(C_1-C_3 \text{ alkyl})$, or $-N(CH_3)COO(C_1-C_3 \text{ alkyl})$.

12. The compound of claim 10, wherein:

X is a bond or $-(CR^5R^6)_t$;

W is a bond, $-CH_2-$, or $-C(CH_3)H-$;


Y is -O-;

Z is -O-;

V is -CH₂- or -CH₂CH₂-; and


L is halogen.

13. The compound of claim 1, wherein the compound has the structure of formula (G-II)



(G-II)

or a pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug thereof, wherein:



X is -(CR⁵R⁶)_t;

Y is -O-;

Z is -O-;

W is -CH₂- or -C(CH₃)H-;

V is -CH₂CH₂-;

L is halogen;

R¹ and R² are each independently Cl or -CN;

at least one R³ is selected from -CN, C₁-C₃ alkoxy, -CONH₂, -NH₂SO₂CH₃, -N(CH₃)SO₂CH₃, -NH₂SO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, or -SO₂CH₃ and the other R³, if present,

is selected from -CN, -CF₃, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₁-C₃ alkoxy, -S(C₁-C₃ alkyl), -SO₂(C₁-C₃ alkyl), -NH₂, -(C₁-C₃ alkyl)NH₂, -NHSO₂CH₃, -NHSO₂CF₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -CH₂NHSO₂CH₃, -CH₂N(CH₃)SO₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl);

R⁵ and R⁶ are each independently hydrogen or methyl;

n₁ and n₂ are each independently 0, 1, or 2;


n₃ is 1 or 2; and


t is 1.


14. The compound of claim 13, wherein:


at least one R³ is selected from -NHSO₂CH₃, -NHSO₂CH₂CH₃, or -SO₂CH₃ and the other R³, if present, is selected from -CN, C₁-C₃ alkyl, C₁-C₃ alkoxy, -SO₂(C₁-C₃ alkyl), -NH₂, -(C₁-C₃ alkyl)NH₂, -NHSO₂CH₃, -N(CH₃)SO₂CH₃, -NHSO₂CH₂CH₃, -N(CH₃)SO₂CH₂CH₃, -SO₂NH₂, -CONH₂, -CON(C₁-C₃ alkyl)₂, -CONH(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), -N(CH₃)COO(C₁-C₃ alkyl), -NHCO(C₁-C₃ alkyl), or -N(CH₃)COO(C₁-C₃ alkyl).

15. A compound selected from the group consisting of:





















































16. The compound of claim 15, wherein the compound is selected from







, or a pharmaceutically acceptable salt,

tautomer, stereoisomer or prodrug thereof.

17. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

18. A pharmaceutical composition comprising a compound of claim 4 and a pharmaceutically acceptable carrier.

19. A pharmaceutical composition comprising a compound of claim 10 and a pharmaceutically acceptable carrier.

20. A pharmaceutical composition comprising a compound of claim 13 and a pharmaceutically acceptable carrier.

21. A pharmaceutical composition comprising a compound of claim 15 and a pharmaceutically acceptable carrier.

22. A pharmaceutical composition comprising a compound of claim 16 and a pharmaceutically acceptable carrier.


23. A method for treating cancer, comprising administering the compound, pharmaceutically acceptable salt, tautomer, stereoisomer or prodrug of the compound of claim 1, to a subject in need thereof.


24. The method of claim 23, wherein the cancer is prostate cancer.

25. The method of claim 24, wherein the prostate cancer is metastatic castration-resistant prostate cancer.

26. The method of claim 25, wherein the prostate cancer expresses full-length androgen receptor or truncated androgen receptor splice variant.

27. The method of claim 24, wherein the compound is selected from





tautomer, stereoisomer or prodrug thereof.