

METHODS FOR PURIFICATION OF MESSENGER RNA

RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application Serial No. 61/984,503, filed April 25, 2014, the disclosure of which is hereby incorporated by reference.

BACKGROUND

[0002] Messenger RNA therapy is becoming an increasingly important approach for the treatment of a variety of diseases. Messenger RNA therapy involves administration of messenger RNA (mRNA) into a patient in need of the therapy and production of the protein encoded by the mRNA within the patient body. Thus, there is a great need for large scale production of highly pure and safe mRNA product suitable for therapeutic use.

SEQUENCE LISTING

[0003] The present specification makes reference to a Sequence Listing (submitted electronically as a .txt file named "Sequence listing.txt" on April 24, 2015). The .txt file was generated on April 22, 2015 and is 11,184 bytes in size. The entire contents of the Sequence Listing are herein incorporated by reference.

SUMMARY OF THE INVENTION

[0004] The present invention provides improved methods for effective purification of messenger RNA (mRNA), in particular, *in vitro* synthesized mRNA suitable for therapeutic use. The present invention is, in part, based on the surprising discovery that precipitation of mRNA followed by membrane filtration, a highly unusual combination, resulted in unexpectedly successful large scale production of high quality mRNA.

[0005] Thus, in one aspect, the present invention provides methods of purifying messenger RNA (mRNA) including the steps of (a) precipitating mRNA from an impure preparation; (b) subjecting the impure preparation comprising precipitated mRNA to a

purification process involving membrane filtration such that the precipitated mRNA is captured by a membrane; and (c) eluting the captured precipitated mRNA from the membrane by re-solubilizing the mRNA, thereby resulting in a purified mRNA solution. In some embodiments, a purification process involving membrane filtration suitable for the present invention is tangential flow filtration. In some embodiments, a purification process involving membrane filtration suitable for the present invention is direct flow filtration.

[0006] In some embodiments, the step of precipitating mRNA comprises treating the impure preparation with a solution comprising a reagent selected from the group consisting of lithium chloride, sodium chloride, potassium chloride, guanidinium chloride, guanidinium thiocyanate, guanidinium isothiocyanate, ammonium acetate and combinations thereof. In some embodiments, a suitable reagent is guanidinium thiocyanate.

[0007] In some embodiments, a solution suitable for mRNA precipitation comprises guanidinium thiocyanate at a concentration of about 1 M or greater, of about 2 M or greater, of about 3 M or greater, of about 4 M or greater, of about 5 M or greater, of about 6 M or greater, of about 7 M or greater, of about 8 M or greater, of about 9 M or greater, or of about 10 M or greater. In some embodiments, a solution suitable for mRNA precipitation comprises guanidinium thiocyanate at a concentration of about 4M. In some such embodiments, a suitable solution further includes sodium lauryl sarcosyl and/or sodium citrate. For example, in certain embodiments, a solution suitable for mRNA precipitation comprises 4M guanidinium thiocyanate, 0.5% sodium lauryl sarcosyl, and 25 mM sodium citrate. In certain embodiments, a solution suitable for mRNA precipitation comprises 4M guanidinium thiocyanate, and 0.5% sodium lauryl sarcosyl. In certain embodiments, a solution suitable for mRNA precipitation comprises 4M guanidinium thiocyanate, and 25 mM sodium citrate.

[0008] In some embodiments, the step of precipitating mRNA further comprises a step of treating the impure preparation with absolute ethanol.

[0009] In some embodiments, the step of precipitating mRNA further comprises a step of treating the impure preparation with isopropyl alcohol.

[0010] In some embodiments, a membrane suitable for the present invention is made of material selected from the group consisting of polyethersulfone (mPES) (not modified), polyethersulfone (mPES) hollow fiber membrane, polyvinylidene fluoride (PVDF), cellulose

acetate, nitrocellulose, MCE (mixed cellulose esters), ultra-high MW polyethylene (UPE), polyfluorotetraethylene (PTFE), nylon, and combination thereof.

[0011] In some embodiments, a method according to the invention further comprises washing the captured precipitated mRNA before eluting. In some embodiments, the washing step comprises multiple rinse cycles using a wash solution comprising a guanidinium buffer and ethanol, followed by about 70-80% ethanol (e.g., about 70%, 75%, or 80% ethanol). In some embodiments, the multiple rinse cycles suitable for the present invention are at least 5 or more than 5 cycles (e.g., about 5 to 10 cycles or about 5, 6, 7, 8, 9 or 10 cycles).

[0012] In some embodiments, the eluting step comprises re-solubilizing the captured precipitated mRNA with RNase-free water. In some embodiments, the RNase-free water is re-circulated for about 5-10 minutes (e.g., for about 5, 6, 7, 8, 9 or 10 minutes).

[0013] In some embodiments, a method according to the present invention further comprises a step of dialyzing the purified mRNA solution. In some embodiments, the purified mRNA solution is dialyzed with 1mM sodium citrate using a 100 kDa molecular weight cut-off (MWCO) membrane.

[0014] In various embodiments, the present invention may be used to purify mRNA *in vitro* synthesized from an impure preparation containing an *in vitro* mRNA synthesis reaction mixture. In some embodiments, the impure preparation comprises prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis.

[0015] In some embodiments, the purified mRNA solution contains less than about 5% (e.g., less than about 4.5%, 4%, 3.5%, 3%, 2.5%, 2%, 1.5%, 1%, 0.5%, 0.1%) of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis. In certain embodiments, the purified mRNA solution contains less than about 1% (e.g., less than about 0.9%, 0.8%, 0.7%, 0.6%, 0.5%) of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis. In certain embodiments, the purified mRNA solution contains less than about 0.5% (e.g., less than about 0.4%, 0.3%, 0.2%, or 0.1%) of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis. In some embodiments, the purified mRNA solution contains less than about 0.1% of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis. In some embodiments, the purified mRNA solution

is substantially free of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis.

[0016] In some embodiments, the prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis are measured via silver stain, gel electrophoresis, high-performance liquid chromatography (HPLC), ultra-performance liquid chromatography (UPLC), and/or capillary electrophoresis.

[0017] In some embodiments, the prematurely aborted RNA sequences contain less than 15 bases (e.g., less than 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, or 3 bases). In some embodiments, the prematurely aborted RNA sequences contain about 8-15, 8-14, 8-13, 8-12, 8-11, or 8-10 bases.

[0018] In some embodiments, the enzyme reagents used in *in vitro* synthesis comprise T7 RNA polymerase, DNase I, pyrophosphatase, and/or RNase inhibitor.

[0019] In some embodiments, the mRNA is purified at a scale of or greater than 1 gram, 5 gram, 10 gram, 15 gram, 20 gram, 25 gram, 30 gram, 35 gram, 40 gram, 45 gram, 50 gram, 75 gram, 100 gram, 150 gram, 200 gram, 250 gram, 300 gram, 350 gram, 400 gram, 450 gram, 500 gram, 550 gram, 600 gram, 650 gram, 700 gram, 750 gram, 800 gram, 850 gram, 900 gram, 950 gram, 1 kg, 2.5 kg, 5 kg, 7.5 kg, 10 kg, 25 kg, 50 kg, 75 kg, or 100 kg per batch. As shown in the examples below, a batch comprising purified mRNA in the amount of 10 gram or greater (25 gram or more) can be achieved easily with the methods of the invention.

[0020] In some embodiments, the mRNA is purified before a cap and/or tail are added to the mRNA. In some embodiments, the mRNA is purified after a cap and/or tail are added to the mRNA. In some embodiments, the mRNA is purified after a cap is added. In some embodiments, the mRNA is purified both before and after a cap and/or tail are added to the mRNA.

[0021] In some embodiments, the mRNA is or greater than about 1 kb, 1.5 kb, 2 kb, 2.5 kb, 3 kb, 3.5 kb, 4 kb, 4.5 kb, 5 kb, 6 kb, 7 kb, 8 kb, 9 kb, 10 kb, 11 kb, 12 kb, 13 kb, 14 kb, 15 kb, or 20 kb in length.

[0022] In some embodiments, the mRNA comprises one or more modifications to enhance stability. In some embodiments, the one or more modifications comprise modified

nucleotide and/or modified sugar phosphate backbones. In some embodiments, the mRNA is unmodified.

[0023] In some embodiments, the purified mRNA has an integrity of or greater than about 95% (e.g., of or greater than about 96%, 97%, 98%, or 99%). In some embodiments, the purified mRNA has an integrity of or greater than about 98%. In some embodiments, the purified mRNA has an integrity of or greater than about 99%.

[0024] In some embodiments, the present invention provides a method of purifying messenger RNA (mRNA), comprising (a) precipitating mRNA from an impure preparation; (b) subjecting the impure preparation comprising precipitated mRNA to tangential flow filtration such that the precipitated mRNA is captured by a filtration membrane while impurities are discarded through permeation; and (c) eluting the captured precipitated mRNA by re-solubilizing the precipitated mRNA, resulting in a purified mRNA solution.

[0025] In another aspect, the present invention provides a method for manufacturing messenger RNA (mRNA) comprising synthesizing mRNA *in vitro*; and purifying the *in vitro* synthesized mRNA using a method described herein.

[0026] Among other things, the present invention also provides a messenger RNA (mRNA) purified using a method described herein.

[0027] In another aspect, the present invention also provides a batch of purified mRNA comprising 5 gram or more of a single mRNA species suitable for administration to a human subject. In some embodiments, the batch comprises 10 gram or more of a single mRNA species. In some embodiments, the batch comprises 25 gram or more of a single mRNA species. In some embodiments, the batch is substantially free of impurities from an mRNA synthesis process. In some embodiments, the batch is substantially free of prematurely aborted RNA sequences, DNA templates, and/or enzyme reagents used in *in vitro* synthesis of the single mRNA species. In some embodiments, the purified mRNA contains less than about 5% of enzyme reagents used in *in vitro* synthesis. In some embodiments, the purified mRNA has an integrity greater than about 95%.

[0028] In another aspect, the present invention also provides a composition comprising messenger RNA purified using a method as described herein.

.....

[0029] In another aspect, the present invention also provides a composition comprising *in vitro* synthesized messenger RNA, wherein the composition contains less than 1% of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis.

[0030] In another aspect, the present invention also provides a composition comprising *in vitro* synthesized messenger RNA (mRNA), wherein the mRNA has an integrity of or greater than 95%.

[0031] As used in this application, the terms “about” and “approximately” are used as equivalents. Any numerals used in this application with or without about/approximately are meant to cover any normal fluctuations appreciated by one of ordinary skill in the relevant art.

[0032] Other features, objects, and advantages of the present invention are apparent in the detailed description that follows. It should be understood, however, that the detailed description, while indicating embodiments of the present invention, is given by way of illustration only, not limitation. Various changes and modifications within the scope of the invention will become apparent to those skilled in the art from the detailed description.

BRIEF DESCRIPTION OF THE DRAWING

[0033] The following figures are for illustration purposes only and not for limitation.

[0034] **Figure 1** depicts an exemplary process for large-scale purification of mRNA involving load, wash and elute steps. For example, the precipitated mRNA can be loaded to membranes such that the precipitated mRNA may be captured as retentate while soluble impurities as well as insoluble ones less than 0.22 μm are discarded through the permeate. After capture, the solid precipitate is washed with various buffers followed by re-solubilization and elution for a pure messenger RNA product.

[0035] **Figure 2** shows an exemplary silver-stained protein gel of modified cystic fibrosis transmembrane conductance regulator (CFTR) mRNA from an initial 1 gram *in vitro* transcription (IVT) reaction according to provided methods (with three elutions) and control enzymes present in the IVT reaction.

[0036] **Figure 3** shows an exemplary silver-stained protein gel of modified CFTR mRNA from an initial 1.5 gram *in vitro* transcription (IVT) reaction according to provided methods (with six elutions) and control enzymes present in the IVT reaction.

[0037] **Figure 4** shows the length of exemplary mRNA in *in vitro* transcription (IVT) samples of modified CFTR mRNA purified and filtered according to provided methods as shown by agarose gel electrophoresis. Full integrity remained for modified CFTR mRNA post large-scale precipitation.

[0038] **Figure 5** shows an exemplary silver-stained protein gel of argininosuccinate synthetase (ASS1) mRNA from an initial 1 gram *in vitro* transcription (IVT) reaction according to provided methods (with five elutions) and control enzymes present in the IVT reaction.

[0039] **Figure 6** shows an exemplary SYPRO-stained protein gel of argininosuccinate synthetase (ASS1) mRNA from an initial 1 gram *in vitro* transcription (IVT) reaction according to provided methods (with five elutions) and control enzymes present in the IVT reaction.

[0040] **Figure 7** shows the length of exemplary mRNA in *in vitro* transcription (IVT) samples of ASS1 mRNA purified and filtered according to provided methods as shown by agarose gel electrophoresis. Full integrity remained for ASS1 mRNA post large-scale precipitation.

[0041] **Figure 8** shows an exemplary silver-stained protein gel of firefly luciferase (FFL) mRNA from an initial *in vitro* transcription (IVT) reaction according to provided methods (with either a single or double precipitation) and control enzymes present in the IVT reaction.

[0042] **Figure 9** shows an exemplary SYPRO-stained protein gel of firefly luciferase (FFL) mRNA from an initial *in vitro* transcription (IVT) reaction according to provided methods (with either a single or double precipitation) and control enzymes present in the IVT reaction.

[0043] **Figure 10** shows the length of exemplary mRNA in *in vitro* transcription (IVT) samples of firefly luciferase (FFL) mRNA purified and filtered according to provided methods as shown by agarose gel electrophoresis. Full integrity remained for FFL mRNA post large-scale precipitation and re-precipitation.

[0044] **Figure 11** shows an exemplary silver-stained protein gel of argininosuccinate synthetase (ASS1) mRNA from a final capping and tailing reaction (with three elutions) and control enzymes present in the capping and tailing reactions.

[0045] **Figure 12** shows the length of exemplary final ASS1 mRNA purified and filtered according to provided methods as shown by agarose gel electrophoresis. Full integrity remained for ASS1 mRNA post large-scale precipitation and re-precipitation.

[0046] **Figure 13** shows exemplary luminescence observed within cell lysates of FFL mRNA treated HEK293T cells. Cells were harvested 24 hours post-transfection. A comparison of vendor derived mRNA versus spin-column purified mRNA (commercial kit) versus precipitation-TFF purified FFL mRNA translational ability is represented.

[0047] **Figure 14** shows exemplary CFTR protein levels observed within cell lysates of hCFTR mRNA-transfected HEK293T cells. Cells were harvested 24 hours post-transfection. A comparison of TFF purified mRNA versus spin-column (commercial kit) hCFTR mRNA translational ability is represented.

[0048] **Figure 15** shows the length of an exemplary mRNA from an *in vitro* transcription (IVT) sample of argininosuccinate synthetase (ASS1) mRNA purified and filtered according to provided methods. Full integrity remained for ASS1 mRNA post large-scale (5G) precipitation.

[0049] **Figure 16** shows the length of an exemplary mRNA from an *in vitro* transcription (IVT) sample of argininosuccinate synthetase (ASS1) mRNA (pre- and post-capped/tailed) purified and filtered according to provided methods. Full integrity remained for ASS1 mRNA post large-scale (5G) precipitation.

[0050] **Figure 17** shows an exemplary silver-stained protein gel of argininosuccinate synthetase (ASS1) mRNA after final purification (5G) according to provided methods as well as control enzymes present in the reaction.

[0051] **Figure 18** shows exemplary human ASS1 protein production observed within cell lysates of ASS1 mRNA treated HEK293T cells. Cells were harvested 24 hours post-transfection. A comparison of spin-column purified mRNA (commercial kit) versus precipitation-TFF purified ASS1 mRNA (5G) translational ability is represented.

[0052] **Figure 19** shows the length of an exemplary mRNA in *in vitro* transcription (IVT) sample of cystic fibrosis transmembrane conductance regulator (CFTR) mRNA (pre- and post-capped/tailed) purified and filtered according to provided methods as shown by agarose gel electrophoresis. Full integrity remained for CFTR mRNA post large-scale (10G) precipitation.

[0053] **Figure 20** shows an exemplary silver-stained protein gel of cystic fibrosis transmembrane conductance regulator (CFTR) mRNA after final purification (10G) according to provided methods as well as control enzymes present in the reaction.

[0054] **Figure 21** shows exemplary CFTR protein levels observed within cell lysates of hCFTR mRNA-transfected HEK293T cells from a large scale production and purification. Cells were harvested 24 hours post-transfection.

[0055] **Figure 22** shows the length of an exemplary mRNA in *in vitro* transcription (IVT) sample of argininosuccinate synthetase (ASS1) mRNA at 25G scale production. Full integrity remained for ASS1 mRNA post large-scale (25G) isolation.

DEFINITIONS

[0056] In order for the present invention to be more readily understood, certain terms are first defined below. Additional definitions for the following terms and other terms are set forth throughout the specification.

[0057] *Animal*: As used herein, the term “animal” refers to any member of the animal kingdom. In some embodiments, “animal” refers to humans, at any stage of development. In some embodiments, “animal” refers to non-human animals, at any stage of development. In certain embodiments, the non-human animal is a mammal (*e.g.*, a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, and/or a pig). In some embodiments, animals include, but are not limited to, mammals, birds, reptiles, amphibians, fish, insects, and/or worms. In some embodiments, an animal may be a transgenic animal, genetically-engineered animal, and/or a clone.

[0058] *Approximately or about*: As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain embodiments, the term “approximately” or “about” refers to a range of values

that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[0059] *Biologically active:* As used herein, the phrase “biologically active” refers to a characteristic of any agent that has activity in a biological system, and particularly in an organism. For instance, an agent that, when administered to an organism, has a biological effect on that organism, is considered to be biologically active.

[0060] *Expression:* As used herein, “expression” of a nucleic acid sequence refers to translation of an mRNA into a polypeptide, assembly of multiple polypeptides into an intact protein and/or post-translational modification of a polypeptide or fully assembled protein. In this application, the terms “expression” and “production,” and grammatical equivalent, are used inter-changeably.

[0061] *Functional:* As used herein, a “functional” biological molecule is a biological molecule in a form in which it exhibits a property and/or activity by which it is characterized.

[0062] *Improve, increase, or reduce:* As used herein, the terms “improve,” “increase” or “reduce,” or grammatical equivalents, indicate values that are relative to a baseline measurement, such as a measurement in the same individual prior to initiation of the treatment described herein, or a measurement in a control subject (or multiple control subject) in the absence of the treatment described herein. A “control subject” is a subject afflicted with the same form of disease as the subject being treated, who is about the same age as the subject being treated.

[0063] *Impurities:* As used herein, the term “impurities” refers to substances inside a confined amount of liquid, gas, or solid, which differ from the chemical composition of the target material or compound. Impurities are also referred to as contaminants.

[0064] *In Vitro:* As used herein, the term “*in vitro*” refers to events that occur in an artificial environment, *e.g.*, in a test tube or reaction vessel, in cell culture, *etc.*, rather than within a multi-cellular organism.

[0065] *In Vivo:* As used herein, the term “*in vivo*” refers to events that occur within a multi-cellular organism, such as a human and a non-human animal. In the context of cell-based

systems, the term may be used to refer to events that occur within a living cell (as opposed to, for example, *in vitro* systems).

[0066] *Isolated*: As used herein, the term “isolated” refers to a substance and/or entity that has been (1) separated from at least some of the components with which it was associated when initially produced (whether in nature and/or in an experimental setting), and/or (2) produced, prepared, and/or manufactured by the hand of man. Isolated substances and/or entities may be separated from about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% of the other components with which they were initially associated. In some embodiments, isolated agents are about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% pure. As used herein, a substance is “pure” if it is substantially free of other components. As used herein, calculation of percent purity of isolated substances and/or entities should not include excipients (*e.g.*, buffer, solvent, water, *etc.*).

[0067] *messenger RNA (mRNA)*: As used herein, the term “messenger RNA (mRNA)” refers to a polynucleotide that encodes at least one polypeptide. mRNA as used herein encompasses both modified and unmodified RNA. mRNA may contain one or more coding and non-coding regions.

[0068] *mRNA integrity*: As used herein, the term “mRNA integrity” generally refers to the quality of mRNA. In some embodiments, mRNA integrity refers to the percentage of mRNA that is not degraded after a purification process (*e.g.*, tangential flow filtration). mRNA integrity may be determined using methods well known in the art, for example, by RNA agarose gel electrophoresis (*e.g.*, Ausubel et al., John Wiley & Sons, Inc., 1997, Current Protocols in Molecular Biology).

[0069] *Nucleic acid*: As used herein, the term “nucleic acid,” in its broadest sense, refers to any compound and/or substance that is or can be incorporated into a polynucleotide chain. In some embodiments, a nucleic acid is a compound and/or substance that is or can be incorporated into a polynucleotide chain via a phosphodiester linkage. In some embodiments, “nucleic acid” refers to individual nucleic acid residues (*e.g.*, nucleotides and/or nucleosides). In

some embodiments, “nucleic acid” refers to a polynucleotide chain comprising individual nucleic acid residues. In some embodiments, “nucleic acid” encompasses RNA as well as single and/or double-stranded DNA and/or cDNA. Furthermore, the terms “nucleic acid,” “DNA,” “RNA,” and/or similar terms include nucleic acid analogs, *i.e.*, analogs having other than a phosphodiester backbone. For example, the so-called “peptide nucleic acids,” which are known in the art and have peptide bonds instead of phosphodiester bonds in the backbone, are considered within the scope of the present invention. The term “nucleotide sequence encoding an amino acid sequence” includes all nucleotide sequences that are degenerate versions of each other and/or encode the same amino acid sequence. Nucleotide sequences that encode proteins and/or RNA may include introns. Nucleic acids can be purified from natural sources, produced using recombinant expression systems and optionally purified, chemically synthesized, *etc.* Where appropriate, *e.g.*, in the case of chemically synthesized molecules, nucleic acids can comprise nucleoside analogs such as analogs having chemically modified bases or sugars, backbone modifications, *etc.* A nucleic acid sequence is presented in the 5' to 3' direction unless otherwise indicated. In some embodiments, a nucleic acid is or comprises natural nucleosides (*e.g.*, adenosine, thymidine, guanosine, cytidine, uridine, deoxyadenosine, deoxythymidine, deoxyguanosine, and deoxycytidine); nucleoside analogs (*e.g.*, 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3-methyl adenosine, 5-methylcytidine, C-5 propynyl-cytidine, C-5 propynyl-uridine, 2-aminoadenosine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5-propynyl-cytidine, C5-methylcytidine, 2-aminoadenosine, 7-deazaadenosine, 7-deazaguanosine, 8-oxoadenosine, 8-oxoguanosine, O(6)-methylguanine, and 2-thiocytidine); chemically modified bases; biologically modified bases (*e.g.*, methylated bases); intercalated bases; modified sugars (*e.g.*, 2'-fluororibose, ribose, 2'-deoxyribose, arabinose, and hexose); and/or modified phosphate groups (*e.g.*, phosphorothioates and 5'-*N*-phosphoramidite linkages). In some embodiments, the present invention is specifically directed to “unmodified nucleic acids,” meaning nucleic acids (*e.g.*, polynucleotides and residues, including nucleotides and/or nucleosides) that have not been chemically modified in order to facilitate or achieve delivery.

[0070] *Patient:* As used herein, the term “patient” or “subject” refers to any organism to which a provided composition may be administered, *e.g.*, for experimental, diagnostic, prophylactic, cosmetic, and/or therapeutic purposes. Typical patients include animals (*e.g.*,

mammals such as mice, rats, rabbits, non-human primates, and/or humans). In some embodiments, a patient is a human. A human includes pre and post natal forms.

[0071] *Pharmaceutically acceptable*: The term “pharmaceutically acceptable” as used herein, refers to substances that, within the scope of sound medical judgment, are suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0072] *Prematurely aborted RNA sequences*: The term “prematurely aborted RNA sequences”, as used herein, refers to incomplete products of an mRNA synthesis reaction (e.g., an *in vitro* synthesis reaction). For a variety of reasons, RNA polymerases do not always complete transcription of a DNA template; i.e., RNA synthesis terminates prematurely. Possible causes of premature termination of RNA synthesis include quality of the DNA template, polymerase terminator sequences for a particular polymerase present in the template, degraded buffers, temperature, depletion of ribonucleotides, and mRNA secondary structures. Prematurely aborted RNA sequences may be any length that is less than the intended length of the desired transcriptional product. For example, prematurely aborted mRNA sequences may be less than 1000 bases, less than 500 bases, less than 100 bases, less than 50 bases, less than 40 bases, less than 30 bases, less than 20 bases, less than 15 bases, less than 10 bases or fewer.

[0073] *Salt*: As used herein the term “salt” refers to an ionic compound that does or may result from a neutralization reaction between an acid and a base.

[0074] *Subject*: As used herein, the term “subject” refers to a human or any non-human animal (e.g., mouse, rat, rabbit, dog, cat, cattle, swine, sheep, horse or primate). A human includes pre- and post-natal forms. In many embodiments, a subject is a human being. A subject can be a patient, which refers to a human presenting to a medical provider for diagnosis or treatment of a disease. The term “subject” is used herein interchangeably with “individual” or “patient.” A subject can be afflicted with or is susceptible to a disease or disorder but may or may not display symptoms of the disease or disorder.

[0075] *Substantially*: As used herein, the term “substantially” refers to the qualitative condition of exhibiting total or near-total extent or degree of a characteristic or property of interest. One of ordinary skill in the biological arts will understand that biological and chemical

phenomena rarely, if ever, go to completion and/or proceed to completeness or achieve or avoid an absolute result. The term “substantially” is therefore used herein to capture the potential lack of completeness inherent in many biological and chemical phenomena.

[0076] *Substantially free:* As used herein, the term “substantially free” refers to a state in which relatively little or no amount of a substance to be removed (e.g., prematurely aborted RNA sequences) are present. For example, “substantially free of prematurely aborted RNA sequences” means the prematurely aborted RNA sequences are present at a level less than approximately 5%, 4%, 3%, 2%, 1.0%, 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1% or less (w/w) of the impurity. Alternatively, “substantially free of prematurely aborted RNA sequences” means the prematurely aborted RNA sequences are present at a level less than about 100 ng, 90 ng, 80 ng, 70 ng, 60 ng, 50 ng, 40 ng, 30 ng, 20 ng, 10 ng, 1 ng, 500 pg, 100 pg, 50 pg, 10 pg, or less.

DETAILED DESCRIPTION

[0077] The present invention provides, among other things, improved methods for purifying mRNA from an impure preparation (e.g., *in vitro* synthesis reaction mixture) based on a process involving precipitating mRNA followed by membrane filtration (e.g., tangential flow filtration).

[0078] Various aspects of the invention are described in detail in the following sections. The use of sections is not meant to limit the invention. Each section can apply to any aspect of the invention. In this application, the use of “or” means “and/or” unless stated otherwise.

Synthesis of mRNA

[0079] The present invention may be used to purify any mRNA. mRNA is typically thought of as the type of RNA that carries information from DNA to the ribosome. The existence of mRNA is typically very brief and includes processing and translation, followed by degradation. Typically, in eukaryotic organisms, mRNA processing comprises the addition of a “cap” on the N-terminal (5') end, and a “tail” on the C-terminal (3') end. A typical cap is a 7-

methylguanosine cap, which is a guanosine that is linked through a 5'-5'-triphosphate bond to the first transcribed nucleotide. The presence of the cap is important in providing resistance to nucleases found in most eukaryotic cells. The tail is typically a polyadenylation event whereby a polyadenylyl moiety is added to the 3' end of the mRNA molecule. The presence of this "tail" serves to protect the mRNA from exonuclease degradation. Messenger RNA is translated by the ribosomes into a series of amino acids that make up a protein.

[0080] mRNAs according to the present invention may be synthesized according to any of a variety of known methods. For example, mRNAs according to the present invention may be synthesized via *in vitro* transcription (IVT). Briefly, IVT is typically performed with a linear or circular DNA template containing a promoter, a pool of ribonucleotide triphosphates, a buffer system that may include DTT and magnesium ions, and an appropriate RNA polymerase (e.g., T3, T7 or SP6 RNA polymerase), DNase I, pyrophosphatase, and/or RNase inhibitor. The exact conditions will vary according to the specific application. The presence of these reagents is undesirable in the final product according to several embodiments and may thus be referred to as impurities and a preparation containing one or more of these impurities may be referred to as an impure preparation.

[0081] According to various embodiments, the present invention may be used to purify *in vitro* synthesized mRNA of a variety of lengths. In some embodiments, the present invention may be used to purify *in vitro* synthesized mRNA of or greater than about 1 kb, 1.5 kb, 2 kb, 2.5 kb, 3 kb, 3.5 kb, 4 kb, 4.5 kb, 5 kb, 6 kb, 7 kb, 8 kb, 9 kb, 10 kb, 11 kb, 12 kb, 13 kb, 14 kb, 15 kb, or 20 kb in length. In some embodiments, the present invention may be used to purify *in vitro* synthesized mRNA ranging from about 1-20 kb, about 1-15 kb, about 1-10 kb, about 5-20 kb, about 5-15 kb, about 5-12 kb, about 5-10 kb, about 8-20 kb, or about 8-15 kb in length. For example, typical mRNAs may be about 1 kb to about 5 kb in length. More typically, the mRNA will have a length of about 1 kb to about 3 kb. However, in some embodiments, the mRNA in the composition of the invention is much longer (greater than about 20 kb). In some embodiments, the present invention may be used to purify mRNA containing one or more modifications that typically enhance stability. In some embodiments, one or more modifications are selected from modified nucleotide, modified sugar phosphate backbones, 5' and/or 3' untranslated region. In some embodiments, the present invention may be used to purify *in vitro* synthesized mRNA that is unmodified.

[0082] Typically, mRNAs are modified to enhance stability. Modifications of mRNA can include, for example, modifications of the nucleotides of the RNA. An modified mRNA according to the invention can thus include, for example, backbone modifications, sugar modifications or base modifications. In some embodiments, antibody encoding mRNAs (e.g., heavy chain and light chain encoding mRNAs) may be synthesized from naturally occurring nucleotides and/or nucleotide analogues (modified nucleotides) including, but not limited to, purines (adenine (A), guanine (G)) or pyrimidines (thymine (T), cytosine (C), uracil (U)), and as modified nucleotides analogues or derivatives of purines and pyrimidines, such as e.g. 1-methyl-adenine, 2-methyl-adenine, 2-methylthio-N-6-isopentenyl-adenine, N6-methyl-adenine, N6-isopentenyl-adenine, 2-thio-cytosine, 3-methyl-cytosine, 4-acetyl-cytosine, 5-methyl-cytosine, 2,6-diaminopurine, 1-methyl-guanine, 2-methyl-guanine, 2,2-dimethyl-guanine, 7-methyl-guanine, inosine, 1-methyl-inosine, pseudouracil (5-uracil), dihydro-uracil, 2-thio-uracil, 4-thio-uracil, 5-carboxymethylaminomethyl-2-thio-uracil, 5-(carboxyhydroxymethyl)-uracil, 5-fluoro-uracil, 5-bromo-uracil, 5-carboxymethylaminomethyl-uracil, 5-methyl-2-thio-uracil, 5-methyl-uracil, N-uracil-5-oxyacetic acid methyl ester, 5-methylaminomethyl-uracil, 5-methoxyaminomethyl-2-thio-uracil, 5'-methoxycarbonylmethyl-uracil, 5-methoxy-uracil, uracil-5-oxyacetic acid methyl ester, uracil-5-oxyacetic acid (v), 1-methyl-pseudouracil, queosine, .beta.-D-mannosyl-queosine, wybutoxosine, and phosphoramidates, phosphorothioates, peptide nucleotides, methylphosphonates, 7-deazaguanosine, 5-methylcytosine and inosine. The preparation of such analogues is known to a person skilled in the art e.g. from the U.S. Pat. No. 4,373,071, U.S. Pat. No. 4,401,796, U.S. Pat. No. 4,415,732, U.S. Pat. No. 4,458,066, U.S. Pat. No. 4,500,707, U.S. Pat. No. 4,668,777, U.S. Pat. No. 4,973,679, U.S. Pat. No. 5,047,524, U.S. Pat. No. 5,132,418, U.S. Pat. No. 5,153,319, U.S. Pat. Nos. 5,262,530 and 5,700,642, the disclosure of which is included here in its full scope by reference.

[0083] Typically, mRNA synthesis includes the addition of a “cap” on the N-terminal (5') end, and a “tail” on the C-terminal (3') end. The presence of the cap is important in providing resistance to nucleases found in most eukaryotic cells. The presence of a “tail” serves to protect the mRNA from exonuclease degradation.

[0084] Thus, in some embodiments, mRNAs include a 5' cap structure. A 5' cap is typically added as follows: first, an RNA terminal phosphatase removes one of the terminal phosphate groups from the 5' nucleotide, leaving two terminal phosphates; guanosine

triphosphate (GTP) is then added to the terminal phosphates via a guanylyl transferase, producing a 5'5'5 triphosphate linkage; and the 7-nitrogen of guanine is then methylated by a methyltransferase. Examples of cap structures include, but are not limited to, m7G(5')ppp(5'(A,G(5')ppp(5')A and G(5')ppp(5')G.

[0085] While mRNA provided from *in vitro* transcription reactions may be desirable in some embodiments, other sources of mRNA are contemplated as within the scope of the invention including wild-type mRNA produced from bacteria, fungi, plants, and/or animals.

[0086] In some embodiments, mRNAs include a 5' and/or 3' untranslated region. In some embodiments, a 5' untranslated region includes one or more elements that affect an mRNA's stability or translation, for example, an iron responsive element. In some embodiments, a 5' untranslated region may be between about 50 and 500 nucleotides in length.

[0087] In some embodiments, a 3' untranslated region includes one or more of a polyadenylation signal, a binding site for proteins that affect an mRNA's stability of location in a cell, or one or more binding sites for miRNAs. In some embodiments, a 3' untranslated region may be between 50 and 500 nucleotides in length or longer.

[0088] The present invention may be used to purify mRNAs encoding a variety of proteins. Non-limiting examples of purification of mRNAs encoding firefly luciferase, argininosuccinate synthetase, Factor IX, and CFTR, are described in detail in the Examples section.

[0089] Typically, the present invention is used to purify a single mRNA species, i.e. the mRNA preparation to be purified contains mRNA derived from a single gene or a single synthesis or expression construct. In contrast, total mRNA purified from a cell contains multiple mRNA species.

[0090] A purification process according to the present invention may be carried out during or subsequent to synthesis. For example, mRNA may be purified as described herein before a cap and/or tail are added to the mRNA. In some embodiments, the mRNA is purified after a cap and/or tail are added to the mRNA. In some embodiments, the mRNA is purified after a cap is added. In some embodiments, the mRNA is purified both before and after a cap and/or tail are added to the mRNA. In general, a purification step as described herein may be

performed after each step of mRNA synthesis, optionally along with other purification processes, such as dialysis. For example, mRNA may undergo dialysis to remove shortmers after initial synthesis (e.g., with or without a tail) and then be subjected to precipitation and purification as described herein, then after addition of the cap and/or tail, be purified again by precipitation and purification.

Precipitation of mRNA

[0091] According to the present invention, mRNA may be precipitated from an impure preparation, such as an *in vitro* synthesis reaction mixture, using various precipitation methods known in the art. As used herein, the term “precipitation” (or any grammatical equivalent thereof) refers to the formation of a solid in a solution. When used in connection with mRNA, the term “precipitation” refers to the formation of insoluble or solid form of mRNA in a liquid.

[0092] Any and all methods suitable for precipitating mRNA may be used to practice the present invention. Typically, mRNA precipitation involves a denaturing condition. As used herein, the term “denaturing condition” refers to any chemical or physical condition that can cause disruption of native conformation of mRNA. Since the native conformation of a molecule is usually the most water soluble, disrupting the secondary and tertiary structures of a molecule may cause changes in solubility and may result in precipitation of mRNA from solution.

[0093] For example, a suitable method of precipitating mRNA from an impure preparation involves treating the impure preparation with a denaturing reagent such that the mRNA precipitates. Exemplary denaturing reagents suitable for the invention include, but are not limited to, lithium chloride, sodium chloride, potassium chloride, guanidinium chloride, guanidinium thiocyanate, guanidinium isothiocyanate, ammonium acetate and combinations thereof. Suitable reagent may be provided in a solid form or in a solution.

[0094] As a non-limiting example, guanidinium thiocyanate may be used to precipitate mRNA. Typically, guanidinium thiocyanate may be provided in a solution at a concentration of about 1 M or greater, of about 2 M or greater, of about 3 M or greater, of about 4 M or greater, of about 5 M or greater, of about 6 M or greater, of about 7 M or greater, of about 8 M or greater, of

about 9 M or greater, or of about 10 M or greater. In some embodiments, a solution suitable for mRNA precipitation contains guanidinium thiocyanate at a concentration of about 4M.

[0095] In addition to denaturing reagent, a suitable solution for mRNA precipitation may include additional salt, surfactant and/or buffering agent. For example, a suitable solution may further include sodium lauryl sarcosyl and/or sodium citrate. As non-limiting examples, a solution suitable for mRNA precipitation may contain 4M guanidinium thiocyanate, 0.5% sodium lauryl sarcosyl, and 25 mM sodium citrate; or 4M guanidinium thiocyanate, and 0.5% sodium lauryl sarcosyl; or 4M guanidinium thiocyanate, and 25 mM sodium citrate.

[0096] Typically, it is desirable to incubate the impure preparation with one or more denaturing reagents described herein for a period of time at a desired temperature that permits precipitation of substantial amount of mRNA. For example, the mixture of an impure preparation and a denaturing agent may be incubated at room temperature or ambient temperature for a period of time. Typically, a suitable incubation time is a period of or greater than about 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, or 60 minutes. In some embodiments, a suitable incubation time is a period of or less than about 60, 55, 50, 45, 40, 35, 30, 25, 20, 15, 10, 9, 8, 7, 6, or 5 minutes. In some embodiments, the mixture is incubated for about 5 minutes at room temperature. Typically, “room temperature” or “ambient temperature” refers to a temperature with the range of about 20-25 °C, for example, about 20 °C, 21 °C, 22 °C, 23 °C, 24 °C, or 25 °C. In some embodiments, the mixture of an impure preparation and a denaturing agent may also be incubated above room temperature (e.g., about 30-37 °C or in particular, at about 30 °C, 31 °C, 32 °C, 33 °C, 34 °C, 35 °C, 36 °C, or 37 °C) or below room temperature (e.g., about 15-20 °C or in particular, at about 15 °C, 16 °C, 17 °C, 18 °C, 19 °C, or 20 °C). The incubation period may be adjusted based on the incubation temperature. Typically, a higher incubation temperature requires shorter incubation time.

[0097] Alternatively or additionally, a solvent may be used to facilitate mRNA precipitation. Suitable exemplary solvent includes, but is not limited to, isopropyl alcohol, acetone, methyl ethyl ketone, methyl isobutyl ketone, ethanol, methanol, denatonium, and combinations thereof. For example, a solvent (e.g., absolute ethanol) may be added to an impure preparation together with a denaturing reagent or after the addition of a denaturing reagent and the incubation as described herein, to further enhance and/or expedite mRNA

precipitation. Typically, after the addition of a suitable solvent (e.g., absolute ethanol), the mixture may be incubated at room temperature for another period of time. Typically, a suitable period of incubation time is or greater than about 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, or 60 minutes. In some embodiments, a suitable period of incubation is a period of or less than about 60, 55, 50, 45, 40, 35, 30, 25, 20, 15, 10, 9, 8, 7, 6, or 5 minutes. Typically, the mixture is incubated at room temperature for another about 5 minutes. Temperature above or below room may be used with proper adjustment of incubation time. Alternatively, incubation could occur at 4 °C or -20 °C for precipitation.

[0098] Typically, methods described herein result in precipitation of a substantial amount of mRNA from an impure preparation. In some embodiments, methods described herein result in precipitation of at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% of total mRNA from an impure preparation. In some embodiments, methods described herein result in precipitation of substantially 100% of total mRNA from an impure preparation.

Membrane Filtration

[0099] According to the present invention, an impure preparation containing the precipitated mRNA may be subjected to a purification process involving membrane filtration such that the precipitated mRNA is captured or retained by a membrane. Thus, in some embodiments, an impure preparation is subjected to membrane filtration following precipitation without pre-treatment to remove insolubles.

[0100] Various types of membrane filtration may be used to capture or retain precipitated mRNA. Typically, membrane filtration involves separating solids from fluids using one or more interposed permeable membranes. Membrane filtration may also be used to filter particles from a gaseous sample. Generally speaking, there are two major forms of membrane filtration, passive filtration, which proceeds solely due to solution-diffusion, and active filtration, which uses positive pressure or negative pressure (i.e. vacuum) to force the liquid or gas across the membrane.

[0101] An exemplary process involving membrane filtration for purifying mRNA is shown in Figure 1. Typically, such a process involves load, wash and elute steps.

.....

Load

[0102] Typically, the load step involves loading the feed (e.g., an impure preparation containing precipitated mRNA) onto a membrane and force it through by positive or negative pressure, leaving retentate captured or retained on the membrane. As used herein, the term “retentate” refers to any non-permeating solute and/or insoluble that is retained by a membrane. According to the present invention, precipitated mRNA is captured by a membrane as retentate. As used herein, the term “membrane” refers to any porous layer or sheet of material. In this application, the term “membrane” is used inter-changeably with filter.

[0103] In some embodiments, a suitable membrane has a pore size appropriate for capturing or retaining precipitated mRNA, while letting impurities (including soluble impurities and/or insoluble with size less than the pore size) pass through as permeate. In some embodiments, a suitable membrane has an average pore size of or greater than about 0.10 μm , 0.20 μm , 0.22 μm , 0.24 μm , 0.26 μm , 0.28 μm , 0.30 μm , 0.40 μm , 0.5 μm , or 1.0 μm . In a particular embodiment, a suitable membrane has an average pore size of about 0.22 μm . In some embodiments, appropriate pore size for retaining precipitated mRNA may be determined by the nominal molecular weight limits (NMWL) of the precipitated mRNA, also referred to as the molecular weight cut off (MWCO). Typically, a membrane with pore size less than the NMWL or MWCO of the precipitated mRNA is used. In some embodiments, a membrane with pore size two to six (e.g., 2, 3, 4, 5, or 6) times below the NMWL or MWCO of the precipitated mRNA is used. In some embodiments, a suitable membrane for the present invention may have pore size of or greater than about 100 kilodaltons (kDa), 300 kDa, 500 kDa, 1,000 kDa, 1,500 kDa, 2,000 kDa, 2,500 kDa, 3,000 kDa, 3,500 kDa, 4,000 kDa, 4,500 kDa, 5,000 kDa, 5,500 kDa, 6,000 kDa, 6,500 kDa, 7,000 kDa, 7,500 kDa, 8,000 kDa, 8,500 kDa, 9,000 kDa, 9,500 kDa, or 10,000 kDa.

[0104] A suitable membrane for the present invention may be made of any material. Exemplary membrane materials include, but are not limited to, polyethersulfone (mPES) (not modified), polyethersulfone (mPES) hollow fiber membrane, polyvinylidene fluoride (PVDF), cellulose acetate, nitrocellulose, MCE (mixed cellulose esters), ultra-high MW polyethylene (UPE), polyfluorotetraethylene (PTFE), nylon, polysulfone, polyether sulfone, polyacrylonitrile, polypropylene, polyvinyl chloride, and combination thereof.

[0105] A suitable membrane for the present invention may have various surface area. In some embodiments, a suitable membrane has a sufficiently large surface area to facilitate large scale production of mRNA. For example, a suitable membrane may have a surface area of or greater than about 2,000 cm², 2,500 cm², 3,000 cm², 3,500 cm², 4,000 cm², 4,500 cm², 5,000 cm², 7,500 cm², 10,000 cm², 5 m², 10 m², 12 m², 15m², 20m², 24 m², 25 m², 30m², or 50 m².

[0106] Membrane filtration may be performed in various format to capture precipitated mRNA. In some embodiments, membrane filtration is performed as part of tangential flow filtration (TFF).

[0107] Tangential flow filtration (TFF), also referred to as cross-flow filtration, is a type of filtration wherein the material to be filtered is passed tangentially *across* a filter rather than through it. In TFF, undesired permeate passes through the filter, while the desired retentate (e.g., precipitated mRNA) passes along the filter and is captured or retained on the filter or membrane downstream.

[0108] A principal advantage of tangential flow filtration is that non-permeable retentate that may aggregate in and block the filter (sometimes referred to as “filter cake”) during traditional “dead-end” filtration, are instead carried along the surface of the filter. This advantage allows tangential flow filtration to be particularly suitable for large scale purification of precipitated mRNA. In some embodiments, a load of mRNA of or greater than about 1 gram, 10 gram, 50 gram, 100 gram, 200 gram, 300 gram, 400 gram, 500 gram, 600 gram, 700 gram, 800 gram, 900 gram, 1 kg, 5 kg, 10 kg, 50 kg, or 100 kg may be applied per batch.

[0109] At least three process variables that are important in a typical TFF process: the transmembrane pressure, feed rate, and flow rate of the permeate. The transmembrane pressure is the force that drives fluid through the filter, carrying with it permeable molecules. In some embodiments, the transmembrane pressure is between 1 and 30 pounds per square inch (psi), inclusive.

[0110] The feed rate (also known as the crossflow velocity) is the rate of the solution flow through the feed channel and across the filter. The feed rate determines the force that sweeps away molecules that may otherwise clog or foul the filter and thereby restrict filtrate flow. In some embodiments, the feed rate is between 1 and 500 L/minute. In some embodiments, the feed rate is between 50 and 800 mL/minute. In some embodiments, the feed

rate is between 50 and 750 mL/minute. In some embodiments, the feed rate is between 50 and 300 mL/minute. In some embodiments, the feed rate is between 50 and 200 mL/minute. In some embodiments, the feed rate is between 75 and 200 mL/minute. In some embodiments, the feed rate is between 100 and 200 mL/minute. In some embodiments, the feed rate is between 125 and 175 mL/minute. In some embodiments, the feed rate is 130 mL/minute. In some embodiments, the feed rate is between 60 mL/min and 220 mL/min. In some embodiments, the feed rate is 60 mL/min or greater. In some embodiments, the feed rate is 100 mL/min or greater. In some embodiments, the feed rate is 150 mL/min or greater. In some embodiments, the feed rate is 200 mL/min or greater. In some embodiments, the feed rate is 220 mL/min or greater.

[0111] The flow rate of the permeate is the rate at which the permeate is removed from the system. For a constant feed rate, increasing permeate flow rates can increase the pressure across the filter, leading to enhanced filtration rates while also potentially increasing the risk of filter clogging or fouling. The principles, theory, and devices used for TFF are described in Michaels et al., "Tangential Flow Filtration" in Separations Technology, Pharmaceutical and Biotechnology Applications (W. P. Olson, ed., Interpharm Press, Inc., Buffalo Grove, Ill. 1995). See also U.S. Pat. Nos. 5,256,294 and 5,490,937 for a description of high-performance tangential flow filtration (HP-TFF), which represents an improvement to TFF. In some embodiments, the flow rate is between 1 and 100 L/minute. In some embodiments, the flow rate is between 10 and 100 mL/minute. In some embodiments, the flow rate is between 10 and 90 mL/minute. In some embodiments, the flow rate is between 10 and 80 mL/minute. In some embodiments, the flow rate is between 10 and 70 mL/minute. In some embodiments, the flow rate is between 10 and 60 mL/minute. In some embodiments, the flow rate is between 10 and 50 mL/minute. In some embodiments, the flow rate is between 10 and 40 mL/minute. In some embodiments, the flow rate is between 20 and 40 mL/minute. In some embodiments, the flow rate is 30 mL/minute.

[0112] Any combinations of various process variables described herein may be used. In some embodiments, the tangential flow filtration is performed at a feed rate of approximately 100-200 mL/minute (e.g., approximately 100-180 mL/minute, 100-160 mL/minute, 100-140 mL/minute, 110-190 mL/minute, 110-170 mL/minute, or 110-150 mL/minute) and/or a flow rate of approximately 10-50 mL/minute (e.g., approximately 10-40 mL/minute, 10-30 mL/minute, 20-50 mL/minute, or 20-40 mL/minute). In some embodiments, the tangential flow filtration is performed at a feed rate of approximately 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, or

200 mL/minute and/or a flow rate of approximately 10, 20, 30, 40, or 50 mL/minute. In other embodiments, the tangential flow filtration is performed at a feed rate of approximately 500, 750 mL/minute, 1, 2, 3, 4, or 5 L/min and/or a flow rate of approximately 100, 200, 250, 500, 750 mL/minute or 1 L/min.

Wash

[0113] Typically, the captured insoluble mRNA may be washed before eluting to get rid of impurities retained on the membrane. In some embodiments, a wash step comprises multiple rinse cycles using one or more wash solutions. For example, a wash step may be carried out by multiple rinse cycles using a guanidinium buffer and ethanol, followed by 70-80% ethanol (e.g., about 70%, 75%, or 80% ethanol). In certain embodiments, the multiple rinse cycles is more than 2, more than 3, more than 4, more than 5, more than 6, more than 7, more than 8, more than 9, or more than 10 cycles.

Elute

[0114] Typically, captured or retained mRNA may be eluted by re-solubilizing the precipitated mRNA into a solution. For example, captured mRNA may be eluted with RNase-free water. In certain embodiments, eluting the captured mRNA involves recirculating the RNase-free water. For example, the RNase-free water may be circulated for about 5-30 minutes (e.g., about 5-25 minutes, about 5-20 minutes, or about 5-15 minutes). In particular embodiments, the RNase-free water is re-circulated for about 5-10 minutes (e.g., for about 5, 6, 7, 8, 9 or 10 minutes).

[0115] In some embodiments, re-solubilized mRNA may be dialyzed into a desired formulation at a desired concentration. Various formulations may be used for dialysis. In some embodiments, the purified mRNA solution is dialyzed with 1mM sodium citrate. In some embodiments, the purified mRNA solution is dialyzed with sodium acetate, ammonium carbonate, ammonium bicarbonate, pyridinium acetate, pyridinium formate, ammonium acetate, urea, potassium chloride, etc. Depending on the size of mRNA of interest, dialysis membranes with appropriate molecular weight cut-off (MWCO) may be used. For example, suitable dialysis membranes may have a MWCO of about 50 kDa, 60 kDa, 70 kDa, 80 kDa, 90 kDa, 100 kDa, 150 kDa, 200 kDa, 250 kDa, 300 kDa, 350 kDa, 400 kDa, 450 kDa, or 500 kDa.

Characterization of purified mRNA

[0116] A particular advantage provided by the present invention is the ability to purify mRNA, in particular, mRNA synthesized *in vitro*, at a large or commercial scale. For example, *in vitro* synthesized mRNA may be purified at a scale of or greater than about 1 gram, 10 gram, 50 gram, 100 gram, 200 gram, 300 gram, 400 gram, 500 gram, 600 gram, 700 gram, 800 gram, 900 gram, 1 kg, 5 kg, 10 kg, 50 kg, or 100 kg per batch. In one particular embodiment, *in vitro* synthesized mRNA may be purified at a scale of 10 gram per batch. In another particular embodiment, *in vitro* synthesized mRNA may be purified at a scale of 100 gram per batch. In yet another particular embodiment, *in vitro* synthesized mRNA may be purified at a scale of 1 kg per batch.

[0117] In various embodiments, mRNA purified according to the present invention is substantially free of impurities from mRNA synthesis process including, but not limited to, prematurely aborted RNA sequences, DNA templates, and/or enzyme reagents used in *in vitro* synthesis.

[0118] In particular, the present invention removes or eliminates a high degree of prematurely aborted RNA sequences (also known as “shortmers”). In some embodiments, a method according to the invention removes more than about 90%, 95%, 96%, 97%, 98%, 99% or substantially all prematurely aborted RNA sequences. In some embodiments, mRNA purified according to the present invention is substantially free of prematurely aborted RNA sequences. In some embodiments, mRNA purified according to the present invention contains less than about 5% (e.g., less than about 4%, 3%, 2%, or 1%) of prematurely aborted RNA sequences. In some embodiments, mRNA purified according to the present invention contains less than about 1% (e.g., less than about 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, 0.2%, or 0.1%) of prematurely aborted RNA sequences. In some embodiments, mRNA purified according to the present invention contains undetectable prematurely aborted RNA sequences as determined by, e.g., ethidium bromide and/or Coomassie staining. In some embodiments, prematurely aborted RNA sequences comprise less than 15 bases (e.g., less than 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, or 3 bases). In some embodiments, the prematurely aborted RNA sequences contain about 8-15, 8-14, 8-13, 8-12, 8-11, or 8-10 bases.

[0119] In some embodiments, a method according to the present invention removes or eliminates a high degree of enzyme reagents used in *in vitro* synthesis including, but not limited to, T7 RNA polymerase, DNase I, pyrophosphatase, and/or RNase inhibitor. In some embodiments, the present invention is particularly effective to remove T7 RNA polymerase. In some embodiments, a method according to the invention removes more than about 90%, 95%, 96%, 97%, 98%, 99% or substantially all enzyme reagents used in *in vitro* synthesis including. In some embodiments, mRNA purified according to the present invention is substantially free of enzyme reagents used in *in vitro* synthesis including. In some embodiments, mRNA purified according to the present invention contains less than about 5% (e.g., less than about 4%, 3%, 2%, or 1%) of enzyme reagents used in *in vitro* synthesis including. In some embodiments, mRNA purified according to the present invention contains less than about 1% (e.g., less than about 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, 0.2%, or 0.1%) of enzyme reagents used in *in vitro* synthesis including. In some embodiments, mRNA purified according to the present invention contains undetectable enzyme reagents used in *in vitro* synthesis including as determined by, e.g., ethidium bromide and/or Coomassie staining.

[0120] The level of prematurely aborted RNA sequences and/or enzyme reagents in the purified mRNA may be measured using various methods known in the art. For example, the prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis may be measured via silver stain, gel electrophoresis, high-performance liquid chromatography (HPLC), ultra-performance liquid chromatography (UPLC), and/or capillary electrophoresis.

[0121] In various embodiments, mRNA purified using a method described herein maintain high degree of integrity. As used herein, the term “mRNA integrity” generally refers to the quality of mRNA after purification. In some embodiments, mRNA integrity refers to the percentage of mRNA that is not degraded after tangential flow filtration. mRNA integrity may be determined using methods well known in the art, for example, by RNA agarose gel electrophoresis (e.g., Ausubel et al., John Wiley & Sons, Inc., 1997, Current Protocols in Molecular Biology). In some embodiments, mRNA purified according to the present invention has an integrity greater than about 95% (e.g., greater than about 96%, 97%, 98%, 99% or more). In some embodiments, mRNA purified according to the present invention has an integrity greater than 98%. In some embodiments, mRNA purified according to the present invention has an

integrity greater than 99%. In some embodiments, mRNA purified according to the present invention has an integrity of approximately 100%.

Large-scale batch production of mRNA for therapeutic applications

[0122] The present invention addresses an urgent need for the large-scale production of purified mRNAs that have the high degree of purity and integrity required for therapeutic applications. Existing methods are typically small in scale and cannot be scaled up to the extent that is required to make commercial production of mRNA that is suitable for administration to a human subject sufficiently cost-effective. In contrast, the methods of the invention are fully scalable as demonstrated in the examples and allow the cost-effective large-scale production of pharmaceutical-grade mRNA.

[0123] Each batch of purified mRNA produced in accordance with the invention comprises 5 gram or more of a single mRNA species suitable for administration to a human subject. In some embodiments, a single batch comprises 10 gram or more of a single mRNA species. In a particular embodiment, a single batch comprises 25 gram or more of a single mRNA species.

[0124] The method of the invention yields purified mRNA batches that are substantially free of impurities from an mRNA synthesis process. In particular, the batches are substantially free of prematurely aborted RNA sequences, DNA templates, and/or enzyme reagents used in in vitro synthesis of the single mRNA species. For example, a batch of purified mRNA produced in accordance with the invention contains less than about 5% of enzyme reagents used in in vitro synthesis. The purified mRNA in each batch typically has an integrity greater than about 95%.

[0125] The mRNA batches produced in accordance with the methods of the invention can be used for to prepare a therapeutic agent, requiring one or more downstream processing step(s). Typically, each mRNA batch will be formulated, e.g. by encapsulating the mRNA into lipid nanoparticles, liposomes, polymer-based polyplexes, etc. that can be administered to a patient. Typical routes of administration, but not exclusively, involve intravenous or pulmonary delivery of the purified mRNA.

EXAMPLES

Example 1. Generation and Purification of Messenger RNA (mRNA)

Messenger RNA Synthesis

[0126] Firefly Luciferase (FFL), Argininosuccinate Synthetase (ASS1), and human Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) messenger RNA were synthesized by *in vitro* transcription from a plasmid DNA template encoding the gene, which was followed by the addition of a 5' cap structure (Cap 1) (Fechter, P.; Brownlee, G.G. "Recognition of mRNA cap structures by viral and cellular proteins" J. Gen. Virology 2005, 86, 1239-1249) and a 3' poly(A) tail of approximately 200 nucleotides in length as determined by gel electrophoresis. 5' and 3' untranslated regions present in each mRNA product are represented as X and Y, respectively and defined as stated (*vide infra*). The synthesis of the target messenger RNA construct involved a two-step process in which the initial strand consisting of the coding sequence flanked by 5' and 3' untranslated regions is synthesized. This uncapped and untailed construct was purified and processed further via a capping step, followed by a poly-A addition tailing step. A second purification step at the conclusion of the tailing reaction was performed in a similar fashion to the initial purification process.

Exemplary Codon-Optimized Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) mRNA

Construct design:

X₁ - SEQ ID NO: 1 - Y₁

```
AUGCAGCGGUCCCCGCUCGAAAAGGCCAGUGUCGUGUCCAAACUCUUCUUCUCAU
GGACUCGGCCUAUCCUUAGAAAGGGGUUAUCGGCAGAGGCUUGAGUUGUCUGACA
UCUACCAGAUCCCCUCGGUAGAUUCGGCGGAUAACCUCUCGGAGAAGCUCGAACG
GGAAUGGGACCGCGAACUCGCGUCUAAGAAAAACCCGAAGCUCAUCAACGCACUG
AGAAGGUGCUUCUUCUGGCGGUUCAUGUUCUACGGUAUCUUCUUGUAUCUCGGG
GAGGUCACAAAAGCAGUCCAACCCUGUUGUUGGGUCGCAUUAUCGCCUCGUACG
ACCCCGAUAAACAAAGAAGACGGAGCAUCGCGAUUCUACCUCGGGAUCGGACUGUG
UUUGCUUUUCAUCGUCAGAACACUUUUGUUGCAUCCAGCAAUCUUCGGCCUCCAU
CACAUCCGUAUGCAGAUCCGAUCCGCUAUGUUUAGCUUGAUCUACAAAAAGACA
CUGAAACUCUCGUCGCGGGUGUUGGAUAAGAUUCCAUCGGUCAGUUGGUGUCC
CUGCUUAGUAAUAACCUCAACAAAUUCGAUGAGGGACUGGCGCUGGCACAUUUC
GUGUGGAUUGCCCCGUUGCAAGUCGCCCUUUUGAUGGGCCUUAUUUGGGAGCUG
```

.....

UUGCAGGCAUCUGCCUUUUGUGGGCCUGGGAUUUCUGAUUGUGUUGGCAUUGUUU
CAGGCUGGGCUUGGGCGGAUGAUGAUGAAGUAUCGCGACCAGAGAGCGGGUAAA
AUCUCGGAAAGACUCGUCAUCACUUCGGAAAUGAUCGAAAACAUC CAGUCGGUCA
AAGCCUAUUGCUGGGAAGAAGCUAUGGAGAAGAUGAUUGAAAACCUCCGCCAAA
CUGAGCUGAAACUGACCCGCAAGGCGGCGUAUGUCCGGUAUUUCAAUUCGUCAGC
GUUCUUCUUUUCGGGUUCUUCGUUGUCUUUCUCUCGGUUUUGCCUUAUGCCUUG
AUUAAGGGGAUUAUCCUCCGCAAGAUUUUCACCACGAUUUCGUUCUGCAUUGUA
UUGCGCAUGGCAGUGACACGGCAAUUUCCGUGGGCCGUGCAGACAUGGUAUGAC
UCGCUUGGAGCGAUCAACAAA AUCCAAGACUUCUUGCAAAAGCAAGAGUACAAG
ACCCUGGAGUACAAUCUUA CUACUACGAGGAGUAGUAAUGGAGAAUGUGACGGCU
UUUUGGGAAGAGGGUUUUGGAGAACUGUUUGAGAAAGCAAAGCAGAAUAAACAAC
AACCGCAAGACCUCAAAUGGGGACGAUUC CUGUUUUUCUGAACUUCUCCUGC
UCGGAACACCCGUGUUGAAGGACAUCAAUUUCAAGAUUGAGAGGGGACAGCUUC
UCGCGGUAGCGGGAAGCACUGGUGCGGGAAAAACUAGCCUCUUGAUGGUGAUUA
UGGGGGAGCUUGAGCCCAGCGAGGGGAAGAUUAAACACUCCGGGCGUAUCUCAU
UCUGUAGCCAGUUUUCAUUGGAUCAUGCCCGGAACCAUUAAGAGAAACAUCAUUU
UCGGAGUAUCCUAUGAUGAGUACCGAUACAGAUCCGUCAUUAAGGCGUGCCAGU
UGGAAGAGGACAUUUCUAAGUUCGCCGAGAAGGAUAAACAU CGUCUUGGGAGAAG
GGGUAAUUA CAUUGUCGGGAGGGCAGCGAGCGCGGAUCAGCCUCGCGAGAGCGG
UAUACAAAGAU GCAGAUUUGUAUCUGCUUGAUUCACCGUUUGGAUACCUCGACG
UAUUGACAGAAAAAGAAAUCUUCGAGUCGUGCGUGUGUAAACUUAUGGCUAAUA
AGACGAGAAUCCUGGUGACAUCAAAAAUGGAACACCUUAAGAAGGCGGACAAGA
UCCUGAUCCUCCACGAAGGAUCGUCCUACUUUACGGCACUUCUCAGAGUUGCA
AAACUUGCAGCCGGACUUCUCAAGCAAACUCAUGGGGUGUGACUCAUUCGACCAG
UUCAGCGCGGAACGGCGGAACUCGAUCUUGACGGAAACGCUGCACCGAUUCUCGC
UUGAGGGUGAUGCCCCGGUAUCGUGGACCGAGACAAAGAAGCAGUCGUUUAAGC
AGACAGGAGAAUUGGUGAGAAAAAGAAAGAACAGUAUCUUGAAUCCUAUUAACU
CAAUUCGCAAGUUCUCAAUUCGUCCAGAAAACUCCACUGCAGAUGA AUGGAUUG
AAGAGGAUUCGGACGAACCCCUUGGAGCGCAGGCUUAGCCUCGUGCCGGAUUCAGA
GCAAGGGGAGGCCAUUCUUC CCGGAUUUCGGUGAUUUAACCGGACCUACACUU
CAGGCGAGGCGAAGGCAAUCCGUGCUCAACCUCAUGACGCAUUCGGUAAACCAGG
GGCAAAACAUUCACCGCAAACGACGGCCUCAACGAGAAAAGUGUCACUUGCACC
CCAGGCGAAUUGACUGAACUCGACAUCUACAGCCGUAGGCUUUCGCAAGAAACC
GGACUUGAGAU CAGCGAAGAAAUCAAUGAAGAAGAUUUGAAAGAGUGUUUCUUU
GAUGACAUGGAUCAAUCC CAGCGGUGACAACGUGGAACACAUAUCUUGCGUAC
AUCACGGUGCACAAGUCCUUGAUUUUCGUCCUCAUCUGGUGUCUCGUGAUCUUUC
UCGCUGAGGUCGCAGCGUCACUUGUGGUCCUCUGGCUGCUUGGUAAUACGCCCU
GCAAGACAAAGGCAAUUCUACACACUCAAGAAACAAUCCUAUGCCGUGAUUAUC
ACUUCUACAAGCUCGUAAUACGUGUUUACAUCUACGUAGGAGUGGCCGACACUC
UGCUCGCGAUGGGUUUCUUC CAGGACUCCACUCGUUCACACGCUUAUCACUGU
CUCCAAGAUUCUCCACCAUAAGAUGCUUCAUAGCGUACUGCAGGCUC CCAUGUCC
ACCUUGAAUACGCUCAAGGCGGGAGGUUUUUGAAUCGCUUCUCAAAAGAUUU
GCAAUUUUGGAUGACCUUCUGCCCCUGACGAUCUUCGACUUCAUCCAGUUGUUGC
UGAUCGUGAUUUGGGGCUAUUGCAGUAGUCGCUUGCCUCCAGCCUUA CAUUUUUG
UCGCGACCGUUC CCGUGAUCGUGGCGUUUAUCAUGCUGCGGGCCUAUUUCUUGCA
GACGUCACAGCAGCUUAAGCAAACUGGAGUCUGAAGGGAGGUCGCCUAUCUUUAC

GCAUCUUGUGACCAGUUUGAAGGGAUUGUGGACGUUGCGCGCCUUUGGCAGGCA
 GCCCUACUUUGAAACACUGUCCACAAAGCGCUGAAUCUCCAUACGGCAAAUUGG
 UUUUUGUAUUUGAGUACCCUCCGAUGGUUUCAGAUUGCAGAUUGAGAUGAUUUUU
 GUGAUCUUCUUUAUCGCGGUGACUUUUUUCUCCAUCUUGACCACGGGAGAGGGC
 GAGGGACGGGUCGGUAUUUAUCCUGACACUCGCCAUGAACAUUAUGAGCACUUUG
 CAGUGGGCAGUGAACAGCUCGAUUGAUGUGGAUAGCCUGAUGAGGUCCGUUUCG
 AGGGUCUUUAAGUUCAUUCGACAUGCCGACGGAGGGAAAGCCCACAAAAAGUACG
 AAACCCUAUAAGAAUGGGCAAUUGAGUAAGGUAAUGAUCAUCGAGAACAGUCAC
 GUGAAGAAGGAUGACAUCUGGCCUAGCGGGGGUCAGAUGACCGUGAAGGACCUG
 ACGGCAAAAUACACCGAGGGAGGGAAACGCAAUCCUUGAAAACAUCUCGUUCAGCA
 UUAGCCCCGGUCAGCGUGUGGGGUUGCUCGGGAGGACCGGGUCAGGAAAAUCGA
 CGUUGCUGUCGGCCUUCUUGAGACUUCUGAAUACAGAGGGUGAGAUCAGAUCCG
 ACGGCGUUUCGUGGGAUAGCAUACCUUGCAGCAGUGGCGGAAAGCGUUUGGAG
 UAAUCCCCCAAAGGUCUUUAUCUUUAGCGGAACCUUCCGAAAGAAUCUCGAUCC
 UUAUGAACAGUGGUCAGAUAAGAGAUUUGGAAAGUCGCGGACGAGGUUGGCCU
 UCGGAGUGUAAUCGAGCAGUUUCCGGGAAAACUCGACUUUGUCCUUGUAGAUGG
 GGAUGCGUCCUGUCGCAUGGGGCACAAGCAGCUCAUGUGCCUGGCGCGAUCCGUC
 CUCUCUAAAGCGAAAAUUCUUCUCUUGGAUGAACCUUCGGCCCAUCUGGACCCGG
 UAACGUAUCAGAUAUCAGAAGGACACUUAAGCAGGCGUUUGCCGACUGCACGG
 UGAUUCUCUGUGAGCAUCGUAUCGAGGCCAUGCUCGAAUGCCAGCAAUUUCUUG
 UCAUCGAAGAGAAUAAGGUCCGCCAGUACGACUCCAUCCAGAAGCUGCUUAAUGA
 GAGAUAUUGUUCGGCAGGCGAUUUCACCAUCCGAUAGGGUGAAACUUUUUCC
 ACACAGAAAUUCGUCGAAGUGCAAGUCCAAACCGCAGAUUCGCGGCCUUGAAAGAA
 GAGACUGAAGAAGAAGUUCAAGACACGCGUCUUUAA [SEQ ID NO: 1]

Exemplary Codon-Optimized Firefly Luciferase (FFL) mRNA

Construct design:

X₁ - SEQ ID NO: 2 - **Y₁**

AUGGAAGAUGCCAAAAACAUAAGAAGGGCCCAGCGCCAUUCUACCCACUCGAAG
 ACGGGACCGCCGGCGAGCAGCUGCACAAAGCCAUGAAGCGCUACGCCUGGUGCC
 CGGCACCAUCGCCUUUACCGACGCACAUUUCGAGGUGGACAUUACCUACGCCGAG
 UACUUCGAGAUGAGCGUUCGGCUGGCAGAAGCUAUGAAGCGCUAUGGGCUGAAU
 ACAAACCAUCGGAUCGUGGUGUGCAGCGAGAAUAGCUUGCAGUUCUUAUGCCCG
 UGUUGGGUGCCCUGUUAUCGGUGUGGCUGUGGGCCCCAGCUAACGACAUCUACAA
 CGAGCGCGAGCUGCUGAACAGCAUGGGCAUCAGCCAGCCCACCGUCGUUUCGUG
 AGCAAGAAAGGGCUGCAAAAGAUCCUCAACGUGCAAAAGAAGCUACCGAUCAUA
 CAAAAGAUAUCAUCAUGGAUAGCAAGACCGACUACCAAGGGCUUCCAAAGCAUGU
 ACACCUUCGUGACUUCCEAUUUGCCACCCGGCUUCAACGAGUACGACUUCGUGCC
 CGAGAGCUUCGACCGGGACAAAACCAUCGCCUGAUCAUGAACAGUAGUGGCAGU
 ACCGGAUUGCCCAAGGGCGUAGCCCUACCGCACCGCACCGCUUGUGUCCGAUUA
 GUCAUGCCCGCGACCCCAUCUUCGGCAACCAGAUCAUCCCCGACACCGCUAUCCU
 CAGCGUGGUGCCAUUUCACACGGCUUCGGCAUGUUCACCACGCUGGGGCUACUUG
 AUCUGCGGCUUUCGGGUCGUGCUCAUGUACCGCUUCGAGGAGGAGCUAUUCUUGC

GCAGCUUGCAAGACUAUAAGAUUCAAUUCUGCCCUGCUGGUGCCCACACUAUUUAG
 CUUCUUCGCUAAGAGCACUCUCAUCGACAAGUACGACCUAAGCAACUUGCACGAG
 AUCGCCAGCGGGCGGGGCGCCGCUCAGCAAGGAGGUAGGUGAGGCCGUGGCCAAAC
 GCUUCCACCUACCAGGCAUCCGCCAGGGCUACGGCCUGACAGAAACAACCAGCGC
 CAUUCUGAUCACCCCCGAAGGGGACGACAAGCCUGGCGCAGUAGGCAAGGUGGUG
 CCCUUCUUCGAGGCUAAGGUGGUGGACUUGGACACCGGUAAGACACUGGGUGUG
 AACCAGCGCGGGCAGCUGUGCGUCCGUGGCCCAUGAUCAUGAGCGGCUACGUUA
 ACAACCCCGAGGCUACAAACGCUCUCAUCGACAAGGACGGCUGGCUGCACAGCGG
 CGACAUCGCCUACUGGGACGAGGACGAGCACUUCUUCAUCGUGGACCGGCUGAAG
 AGCCUGAUCAAAUACAAGGGCUACCAGGUAGCCCCAGCCGAACUGGAGAGCAUCC
 UGCUGCAACACCCCAACAUCUUCGACGCCGGGGUCCGCCGCCUGCCCCGACGACGA
 UGCCGGCGAGCUGCCCCGCCGCAGUCGUCGUGCUGGAACACGGUAAAACCAUGACC
 GAGAAGGAGAUCGUGGACUAUGUGGCCAGCCAGGUUACAACCGCCAAGAAGCUG
 CGCGGUGGUGUUGUGUUCGUGGACGAGGUGCCUAAAGGACUGACCGGCAAGUUG
 GACGCCCCGCAAGAUCGCGAGAUUCUCAUUAAGGCCAAGAAGGGCGGCAAGAUCG
 CCGUGUAA [SEQ ID NO: 2]

Exemplary Codon-Optimized Human Argininosuccinate Synthetase (ASS1) mRNA

Construct design:

X₁ - SEQ ID NO: 3 - Y₂

AUGAGCAGCAAGGGCAGCGUGGUGCUGGCCUACAGCGGGCGGCCUGGACACCAGCU
 GCAUCCUGGUGUGGCUGAAGGAGCAGGGCUACGACGUGAUCGCCUACCUGGCCAA
 CAUCGGCCAGAAGGAGGACUUCGAGGAGGCCCGCAAGAAGGCCCUAGAAGCUGGGC
 GCCAAGAAGGUGUUCAUCGAGGACGUGAGCCGCGAGUUCGUGGAGGAGUUCAUC
 UGGCCCGCCAUCCAGAGCAGCGCCUGUACGAGGACCGCUACCUGCUGGGCACCA
 GCCUGGCCCCGCCCCUGCAUCGCCCGCAAGCAGGUGGAGAUCGCCCAGCGCGAGGG
 CGCCAAGUACGUGAGCCACGGCGCCACCGGCAAGGGCAACGACCAGGUGCGCUUC
 GAGCUGAGCUGCUACAGCCUGGCCCCCAGAUAAGGUGAUCGCCCCCUGGCGCA
 UGCCCAGAUUCUACAACCGCUUCAAGGGCCGCAACGACCUGAUGGAGUACGCCAA
 GCAGCACGGCAUCCCCAUCCCCGUGACCCCCAAGAACCCUGGAGCAUGGACGAG
 AACCUGAUGCACAUCAGCUACGAGGCCGGCAUCCUGGAGAACCCCAAGAACCAGG
 CCCCCCGGGCCUGUACACCAAGACCCAGGACCCCGCCAAGGCCCCCAACACCCCC
 GACAUCCUGGAGAUUCGAGUUAAGAAGGGCGUGCCCGUGAAGGUGACCAACGUG
 AAGGACGGCACCAACCCACCAGACCAGCCUGGAGCUGUUCAUGUACCUGAACGAGG
 UGGCCGGCAAGCACGGCGUGGGCCGCAUCGACAUCGUGGAGAACCGCUUCAUCGG
 CAUGAAGAGCCGCGGCAUCUACGAGACCCCGCCGGCACCAUCCUGUACCACGCC
 CACCUGGACAUCGAGGCCUUCACCAUGGACCGCGAGGUGCGCAAGAUCAAGCAGG
 GCCUGGGCCUGAAGUUCGCCGAGCUGGUGUACACCGGCUUCUGGCACAGCCCCGA
 GUGCGAGUUCGUGCGCCACUGCAUCGCCAAGAGCCAGGAGCGCGUGGAGGGCAAG
 GUGCAGGUGAGCGUGCUGAAGGGCCAGGUGUACAUCUGGGCCGCGAGAGCCCCC
 UGAGCCUGUACAACGAGGAGCUGGUGAGCAUGAACGUGCAGGGCGACUACGAGC
 CCACCGACGCCACCGGCUUCAUCAACAUCACAGCCUGCGCCUGAAGGAGUACCA
 CCGCCUGCAGAGCAAGGUGACCGCCAAGUGA [SEQ ID NO: 3]

5' and 3' UTR Sequences $X_1 =$

GGACAGAUCGCCUGGAGACGCCAUCCACGCUGUUUUGACCUCCAUAAGAAGACACC
 GGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGAACGCGGAUUCCTCCG
 UGCCAAGAGUGACUCACCGUCCUUGACACG [SEQ ID NO: 4]

 $Y_1 =$

CGGGUGGCAUCCCUGUGACCCCUCCCCAGUGCCUCUCCUGGCCCUGGAAGUUGCC
 ACUCCAGUGCCCACCAGCCUUGUCCUAAUAAAAUUAAGUUGCAUCAAGCU [SEQ
 ID NO: 5]

 $Y_2 =$

GGGUGGCAUCCCUGUGACCCCUCCCCAGUGCCUCUCCUGGCCCUGGAAGUUGCCA
 CUCCAGUGCCCACCAGCCUUGUCCUAAUAAAAUUAAGUUGCAUCAAGCU [SEQ
 ID NO: 6]

Synthesis of mRNA

[0127] In each of the examples below, the synthesis of mRNA was conducted under complete RNase-free conditions. All tubes, vials, pipette tips, pipettes, buffers, etc. were required to be nuclease-free, unless explicitly stated otherwise.

[0128] In the following examples, unless otherwise described, mRNA was synthesized via *in vitro* transcription from a linearized DNA template. To produce the desired mRNA precursor (IVT) construct, a mixture of ~8 mg of linearized DNA, rNTPs (7.25 mM), DTT (10 mM), T7 RNA polymerase, RNase Inhibitor, Pyrophosphatase and reaction buffer (10x, 800mM Hepes (pH 8.0), 20 mM Spermidine, 250 mM MgCl₂, pH 7.7) was prepared with RNase-free water to a final volume of 180 mL. The reaction mixture is incubated at 37°C for a range of time between 20 minutes- 60 minutes. Upon completion, the mixture is treated with DNase I for an additional 15 minutes and quenched accordingly.

Addition of 5' Cap and 3' Tail

[0129] The purified mRNA product from the aforementioned IVT step (and possibly initial TFF filtration as well) was denatured at 65°C for 10 minutes. Separately, portions of GTP (1.0 mM), S-adenosyl methionine, RNase inhibitor, 2'-O-Methyltransferase and guanylyl

.....

transferase are mixed together with reaction buffer (10x, 500 mM Tris-HCl (pH 8.0), 60 mM KCl, 12.5 mM MgCl₂) to a final concentration of 1.6 L. Upon denaturation, the mRNA was cooled on ice and then added to the reaction mixture. The combined solution was incubated for a range of time at 37°C for 25-90 minutes. Upon completion, aliquots of ATP (2.0 mM), PolyA Polymerase and tailing reaction buffer (10x, 500 mM Tris-HCl (pH 8.0), 2.5 M NaCl, 100 mM MgCl₂) were added and the total reaction mixture was further incubated at 37°C for a range of time from 20-45 minutes. Upon completion, the final reaction mixture was quenched and purified accordingly.

Purification of mRNA

Precipitation of mRNA

[0130] Messenger RNA can be precipitated using a variety of methods. For example, the use of lithium chloride, potassium chloride, guanidinium chloride, guanidinium thiocyanate, guanidinium isothiocyanate, ammonium acetate and other salts afford efficient precipitation of mRNA from the reaction mixture.

Tangential Flow Filtration

[0131] In the following examples, unless otherwise described, the tangential flow filtration (TFF) system consisted of a filtration membrane and a peristaltic pump (Spectrum system) with tangential circulation of the fluid across the membrane at a feed rate of ~130 mL/min with a 30 mL/min flow rate for the permeate. The TFF membrane employed was a MidiKros 500 kDa mPES 115cm² (Spectrum Labs). Before use, the filter cartridge was washed with nuclease free water and further cleaned with 0.2 N NaOH. Finally the system was cleaned with nuclease free water until the pH of permeate and retentate reached a pH ~6. Isolation of mRNA via TFF can be accomplished as depicted in Figure 1.

Purification of Firefly Luciferase mRNA

[0132] In a representative example of mRNA purification, a 1 gram batch of firefly luciferase (FFL) mRNA was transcribed via *in vitro* methods to produce the aforementioned intermediate construct with no cap and no polyA tail. This reaction maintained a total volume of

~180 mL and was quenched upon completion by the addition of DNase I (~9.0 KU). The resulting solution was treated with a homogeneous solution of 4 M guanidinium thiocyanate, 0.5% sodium lauryl sarcosyl, and 25 mM sodium citrate (1500 mL), bringing the total volume to 1.68 L. The resultant mixture was kept at ambient temperature for ~ 5 minutes followed by further treatment of 1.1 L of absolute ethanol. The mRNA slowly precipitated and the suspension was kept at ambient temperature for ~ 5 minutes. Upon completion, the entire heterogeneous suspension was pumped through a filtration membrane using tangential flow filtration (TFF).

[0133] In this example, modified polyethersulfone (mPES) hollow fiber membranes were employed with a surface area of 2600 cm². The resulting heterogenous mixture (post-precipitation) was pumped through the filter system at a flow rate of 750 mL/min for approximately 8 min in portions. Upon completion, the resulting captured precipitate was rinsed with the guanidinium buffer/ethanol combined buffer solution followed by 80% ethanol (500 mL, 750 mL/min) and repeated multiple times (>5x). Once completely washed, the solid mRNA distributed across the membrane was treated with RNase-free water (250 mL) and re-circulated over 5-10 minutes to ensure dissolution. This process was repeated until there was no more mRNA recovered. Upon completion, the resulting mRNA solution was further dialyzed with 1 mM sodium citrate (pH 6.4) using a 100 KDa MWCO membrane to remove any residual ethanol and obtained a final mRNA product in the proper storage solution. Final concentration was determined via absorption at 260 nm ($\lambda_{\max}$). Messenger RNA purity was determined via UV absorption (260/280 ratio) as well as protein gels (silver stain, SYPRO stain, coomassie, etc). Messenger RNA integrity was determined via gel electrophoresis as well as *in vitro/in vivo* analysis of protein production.

Purification of ASS1 mRNA

[0134] In a second representative example of mRNA purification, a 1 gram batch of ASS1 mRNA was transcribed via *in vitro* methods to produce the aforementioned intermediate construct with no cap and no polyA tail. This reaction maintained a total volume of ~180 mL and was quenched upon completion by the addition of DNase I (~9.0 KU). The resulting solution was treated with a homogeneous solution of 4 M guanidinium thiocyanate, 0.5% sodium lauryl sarcosyl, and 25 mM sodium citrate (1500 mL), bringing the total volume to 1.68 L. The

resultant mixture was kept at ambient temperature for ~ 5 minutes followed by further treatment of 1.1 L of absolute ethanol. The messenger RNA slowly precipitated and the suspension was kept at ambient temperature for ~ 5 minutes. Upon completion, the entire heterogeneous suspension was pumped through a filtration membrane using tangential flow filtration (TFF) and isolated as described above.

Purification of CFTR mRNA

[0135] In a third representative example of mRNA purification, a 1.5 gram batch of modified CFTR mRNA being transcribed via *in vitro* methods to produce the aforementioned intermediate construct with no cap and no polyA tail. This reaction maintains a total volume of ~270 mL and is quenched upon completion by the addition of DNase I (~13.5 KU). The resulting solution was treated with a homogeneous solution of 4 M guanidinium thiocyanate, 0.5% sodium lauryl sarcosyl, and 25 mM sodium citrate (1500 mL), bringing the total volume to 1.77 L. The resultant mixture was kept at ambient temperature for ~ 5 minutes followed by further treatment of 1.1 L of absolute ethanol. The mRNA slowly precipitated and the suspension was kept at ambient temperature for ~ 5 minutes. Upon completion, the entire heterogeneous suspension was pumped through a filtration membrane using tangential flow filtration (TFF) and isolated as described above.

Example 2. Analysis of Purified mRNA

Testing for Presence of Enzymes in Purified mRNA

SYPRO Stain Gels

[0136] Standard SYPRO-stained protein gels were performed to determine the presence of any residual reagent enzymes present before and after purifications. Gels were run at 200V for 35 minutes.

Silver Stain Gels

[0137] Silver stains of all mRNA batches and fractions were performed using SilverQuest® (Life Technologies, Catalog # LC6070) using the manufacturer's protocol. Briefly, samples were loaded (with and without treatment of RNase) and monitored as compared

to loaded enzyme control lanes. Gels were run at 200 V for 35 minutes. Exposure time was allotted for 8 minutes.

Assessment of mRNA Integrity via Agarose Gel Electrophoresis Assays

[0138] Unless otherwise described, mRNA size and integrity were assessed via gel electrophoresis. Either self-poured 1.0% agarose gel or Invitrogen E-Gel precast 1.2% agarose gels were employed. Messenger RNA was loaded at 1.0-1.5 µg quantities per well. Upon completion, mRNA bands were visualized using ethidium bromide.

In Vitro mRNA Integrity Assays

[0139] Unless otherwise described, *in vitro* transfections of firefly luciferase (FFL), argininosuccinate synthetase (ASS1) mRNA, and CFTR mRNA were performed using HEK293T cells. Transfections of one microgram of each mRNA construct were performed in separate wells using lipofectamine. Cells were harvested at select time points (e.g. 4 hour, 8 hour, etc.) and respective protein production was analyzed. For FFL mRNA, cell lysates were analyzed for luciferase production via bioluminescence assays. For ASS1 mRNA, cell lysates were analyzed for ASS1 production via ELISA assays. For CFTR mRNA, cell lysates were analyzed for CFTR production via western blot procedures

Luciferase Assay

[0140] The bioluminescence assay was conducted using a Promega Luciferase Assay System (Item # E1500). The Luciferase Assay Reagent was prepared by adding 10 mL of Luciferase Assay Buffer to Luciferase Assay Substrate and mix via vortex. 20 µL of homogenate samples were loaded onto a 96-well plate, followed by 20 µL of plate control to each sample. Separately, 120 µL of Luciferase Assay Reagent (prepared as described above) was loaded into each well of a 96-well flat bottomed plate. Each plate was inserted into the appropriate chambers using a Molecular Device Flex Station instrument and the luminescence was measured in relative light units (RLU).

ASS1 ELISA Assay

[0141] Standard ELISA procedures were followed employing mouse anti-ASS1 2D1-2E12 IgG as the capture antibody with rabbit anti-ASS1 #3285 IgG as the secondary (detection)

antibody (Shire Human Genetic Therapies). Horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG was used for activation of the 3,3',5,5'-tetramethylbenzidine (TMB) substrate solution. The reaction was quenched using 2N H₂SO₄ after 20 minutes. Detection was monitored via absorption (450 nm) on a Molecular Device SpectraMax instrument. Untreated mouse serum and organs and human ASS1 protein were used as negative and positive controls, respectively.

CFTR Western Blot Analysis

[0142] Western blots were performed on protein samples obtained using immunoprecipitation methods (Dynabeads G). In general, cells or tissue homogenates were processed and treated with Dynabead G pre-bound to anti-human CFTR antibody (R&D Systems, MAB25031). Detection of human CFTR protein was accomplished using Ab570.

Example 3. Purification Results

[0143] This example demonstrates that firefly luciferase (FFL) mRNA, argininosuccinate synthetase (ASS1) mRNA and cystic fibrosis transmembrane conductance regulator (CFTR) mRNA have been successfully purified using precipitation followed by tangential flow filtration with removal of enzymatic reagents as well as shortmers. Many typical chaotropic conditions were successfully employed to precipitate the mRNA as listed above.

[0144] To demonstrate this success, a large scale production (~1 gram) of modified CFTR mRNA IVT reaction mixture was subjected to 4 M guanidinium buffer solution (described above). The resulting mixture was then treated with absolute ethanol and incubated for five minutes at room temperature. Isolation of the precipitated mRNA was achieved via TFF as described above. Figure 2 represents a silver stained protein gel that shows the resulting mRNA isolated after TFF employing the aforementioned conditions. There is no detectable enzyme present upon completion (T7 polymerase, DNase I, pyrophosphatase, RNase Inhibitor). Lanes 1-3 contained modified CFTR after elution 1 (E1), elution 2 (E2) and elution 3 (E3). Lanes 4-7 contained control enzymes present in the IVT reaction. Figure 3 represents a silver stained protein gel that shows pure mRNA resulting from a larger scale (1.5 gram batch) precipitation and filtration process showing no residual enzyme. Lanes 1-6 contained mRNA

after elutions 1-6. Lanes 7-10 contained control enzymes present in the IVT reaction (T7 polymerase, DNase I, RNase Inhibitor, and pyrophosphatase). Figure 4 demonstrates that the modified CFTR mRNA is fully intact after such purification. It shows the results of agarose (1.0%) gel electrophoresis of modified CFTR mRNA after purification via precipitation and filtration. Lane 1 contained a RiboRuler HR ladder. Lane 2-7 contained modified CFTR mRNA purified via filtration after precipitation.

[0145] This process is widely applicable to multiple different mRNA constructs. For example, a second mRNA construct was synthesized and purified using this method. In this instance, argininosuccinate synthetase (ASS1) mRNA was produced and precipitated using methods described above. The solid precipitate was loaded onto the TFF system and isolated according to the process described. Figure 5 represents a silver-stain analysis of the resulting isolated ASS1 mRNA showing no residual enzyme present (T7 polymerase, DNase I, RNase inhibitor, pyrophosphatase). Lanes 1-5 contained mRNA after elutions 1-5. Lanes 6-9 contained control enzymes present in the IVT reaction. Lane 10 contained RNase A as a control because mRNA samples were pre-treated with RNase A prior to loading. As demonstrated in Figure 5, no enzymes were present after extensive exposure to silver stain development. This was further confirmed using SYPRO staining for residual enzymes as shown in Figure 6. Using this method, one can again demonstrate the purity of the ASS1 mRNA as isolated using this process. Lanes 1-5 contained mRNA after elutions 1-5. Lanes 6-9 contained control enzymes present in the IVT reaction. In addition to this, RNA gel electrophoresis showed that the ASS1 mRNA integrity was maintained with full length ASS1 mRNA fully intact after this process (Figure 7). In Figure 7, Lane 1 contained a RiboRuler HR ladder and lanes 2-6 contained ASS1 mRNA after elutions 1-5.

[0146] To further demonstrate that this purification technique is widely applicable to multiple different mRNA constructs, FFL mRNA was synthesized and purified using this method. The mRNA was precipitated using 4 M guanidinium buffer system and filtered. Figure 8 represents a silver-stain analysis of the resulting isolated FFL mRNA showing no residual enzyme present. Lanes 1-4 contained FFL mRNA purified via a single precipitation (XL1) or a double precipitation (XL2), with lanes 2-4 containing different elutions from the second precipitation recovery. Lanes 6-9 contained control enzymes present in the IVT reaction. As demonstrated in Figure 8, no enzymes were present after extensive exposure to silver stain

development. Further, one can see that a single precipitation appears to be sufficient in removing all of the unwanted residual enzyme. This was further confirmed using SYPRO staining for residual enzymes as shown in Figure 9. Using this method, one can again demonstrate the purity of the FFL mRNA isolated using this process. In Figure 9, lanes 1-4 contained FFL mRNA purified via a single precipitation (XL1) or a double precipitation (XL2), with lanes 2-4 containing different elutions from the second precipitation recovery. Lanes 6-9 contained control enzymes present in the IVT reaction. In addition to this, RNA gel electrophoresis showed that the FFL mRNA integrity was maintained with full length FFL mRNA fully intact after this process (Figure 10). Further, the mRNA shows no difference after multiple re-precipitations. In Figure 10, lane 1 contained a RiboRuler HR ladder and lanes 2-5 contained FFL mRNA purified via a single precipitation (XL1) or a double precipitation (XL2).

[0147] Upon subjecting this isolated mRNA further to afford a capped and tailed final product, TFF methods were employed further to purify the final target mRNA. Figure 11 demonstrates that precipitation, followed by capture and elution using tangential flow filtration, resulted in a successfully pure final mRNA product. In Figure 11, lanes 1-3 contained ASS1 mRNA from a cap/tail reaction purified via a single precipitation (E1-3=elution 1-3). Lanes 4-6 contained key control enzymes present in both reaction steps. Lane 7 contained an RNase I control because mRNA samples were pre-treated with RNase I prior to loading. In addition to this, RNA gel electrophoresis showed that the FFL mRNA integrity was maintained with full length FFL mRNA fully intact after this process (Figure 12). In Figure 12, lanes 1-3 contained capped and tailed ASS1 mRNA.

[0148] While such characterization affords information with respect to purity and mRNA size/integrity, a true measure of mRNA quality further lies within its ability to produce the desired protein. Therefore, a comparison of each of the isolated FFL mRNA constructs (TFF vs. spin-column) was made. Each of the three constructs listed below were transfected into HEK293T cells and the corresponding FFL protein production was assessed via FFL protein activity in the form of FFL luminescence upon exposure to luciferin (*vida supra*). Cells were harvested 24 hours post-transfection.

FFL Constructs:

1. FFL mRNA purchased from an outside vendor

.....

2. FFL mRNA purified via commercial kit
3. FFL mRNA purified via precipitation-TFF method

[0149] A comparison of luminescence output of FFL protein produced from each is represented in Figure 13. The integrity of the TFF-purified FFL mRNA was maintained throughout the precipitation and tangential flow filtration process under the conditions described.

[0150] Further, successful detection of human CFTR protein was achieved from transfection of hCFTR mRNA isolated using the aforementioned process. Figure 14 shows successful production of human CFTR protein after transfection of hCFTR mRNA purified via either commercial kits or the aforementioned TFF-based precipitation method. Visualization of this “C-band” for CFTR protein, which is indicative of a full length, properly trafficked CFTR protein, supports the full integrity and active nature of the isolated mRNA via such conditions. This process is further easily scalable. For example, a separate mRNA construct was synthesized and purified at the 5 gram scale using this method. In this instance, argininosuccinate synthetase (ASS1) mRNA was produced and precipitated using methods described above. The solid precipitate was loaded onto the TFF system and isolated according to the process described.

[0151] Integrity of the manufactured mRNA drug substance was demonstrated using two separate methods. Figure 15 shows the length of an exemplary mRNA from an *in vitro* transcription (IVT) sample of argininosuccinate synthetase (ASS1) mRNA purified and filtered according to provided methods. mRNA length was demonstrated via agarose gel-on-a-chip electrophoresis. Intact and full length mRNA was confirmed using gel electrophoresis (Figure 16) for both the IVT pre-cursor as well as the final capped and tailed construct. Full integrity remained for ASS1 mRNA post large-scale (5G) precipitation.

[0152] As demonstrated previously, Figure 17 shows that precipitation, followed by capture and elution using tangential flow filtration, resulted in a successfully pure final mRNA product. Lane 1 contained ASS1 mRNA from a cap/tail reaction purified via a single precipitation. Lanes 2-8 contained key control enzymes present in both reaction steps. Lane 9 contained an RNase I control because mRNA samples were pre-treated with RNase I prior to loading. Such a silver stain method demonstrates that no residual enzymes are present.

[0153] While such characterization affords information with respect to purity and mRNA size/integrity, a true measure of mRNA quality further lies within its ability to produce the desired protein. Therefore, a comparison of the isolated ASS1 mRNA constructs (TFF vs. spin-column) was made (Figure 18). Each of the constructs listed below were transfected into HEK293T cells and the corresponding ASS1 protein production was assessed via ELISA methods. Cells were harvested ~18 hours post-transfection.

[0154] A further demonstration of scalability was achieved with synthesis and purification of CFTR mRNA at the 10 gram scale using this method. In this instance, cystic fibrosis transmembrane conductance regulator (CFTR) mRNA was produced and precipitated using methods described above. The solid precipitate was loaded onto the TFF system and isolated according to the process described. Integrity of the manufactured mRNA drug substance was demonstrated using agarose gel electrophoresis. Intact and full length mRNA was confirmed (Figure 19) for both the IVT pre-cursor as well as the final capped and tailed construct. Full integrity remained for CFTR mRNA post large-scale (10G) precipitation.

[0155] Again, purity of the manufactured product at the 10 gram scale was demonstrated via silver stain as shown in Figure 20. Lane 1 contains CFTR mRNA from a cap/tail reaction purified via a single precipitation. Lanes 2-8 contained key control enzymes present in both reaction steps. Lane 9 contained an RNase I control because mRNA samples were pre-treated with RNase I prior to loading. Such a silver stain method demonstrates that no residual enzymes are present.

[0156] As described above, such characterization affords information with respect to purity and mRNA size/integrity, however a true measure of mRNA quality further lies within its ability to produce the desired protein. Therefore, various amounts of CFTR mRNA were transfected into HEK293T cells and the corresponding CFTR protein production was assessed via western blot methods (Figure 21). Cells were harvested ~18 hours post-transfection.

[0157] In another embodiment, further demonstration of scalability was achieved with synthesis and purification of ASS1 mRNA at the 25 gram scale using this method. In this instance, argininosuccinate synthetase (ASS1) mRNA was produced and precipitated using methods described above. The solid precipitate was loaded onto the TFF system and isolated according to the process described. Integrity of the manufactured mRNA drug substance was

demonstrated using agarose gel electrophoresis. Intact and full length mRNA was confirmed (Figure 22) for ASS1 mRNA post large-scale (25G) precipitation.

EQUIVALENTS AND SCOPE

[0158] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. The scope of the present invention is not intended to be limited to the above Description, but rather is as set forth in the following claims:

.....

CLAIMS

We claim:

1. A method of purifying messenger RNA (mRNA), comprising
 - (a) precipitating mRNA from an impure preparation; and
 - (b) subjecting the impure preparation comprising precipitated mRNA to a purification process involving membrane filtration such that the precipitated mRNA is captured by a membrane; and
 - (c) eluting the captured precipitated mRNA from the membrane by re-solubilizing the mRNA, thereby resulting in a purified mRNA solution.
2. The method of claim 1, wherein the purification process involving membrane filtration is tangential flow filtration.
3. The method of claim 1, wherein the purification process involving membrane filtration is direct flow filtration.
4. The method of claim 1, 2 or 3, wherein the step of precipitating mRNA comprises treating the impure preparation with a solution comprising a reagent selected from the group consisting of lithium chloride, potassium chloride, guanidinium chloride, guanidinium thiocyanate, guanidinium isothiocyanate, ammonium acetate and combinations thereof.
5. The method of claim 4, wherein the reagent is guanidinium thiocyanate.
6. The method of claim 5, wherein the solution comprises 4M guanidinium thiocyanate, 0.5% sodium lauryl sarcosyl, and 25 mM sodium citrate.
7. The method of claim 5, wherein the solution comprises 4M guanidinium thiocyanate, and 0.5% sodium lauryl sarcosyl.
8. The method of claim 5, wherein the solution comprises 4M guanidinium thiocyanate, and 25 mM sodium citrate.
9. The method of any one of the preceding claims, wherein the step of precipitating mRNA further comprises a step of treating the impure preparation with absolute ethanol.

10. The method of any one of the preceding claims, wherein the membrane is selected from the group consisting of polyethersulfone (mPES) (not modified), polyethersulfone (mPES) hollow fiber membrane, polyvinylidene fluoride (PVDF), cellulose acetate, nitrocellulose, MCE (mixed cellulose esters), ultra-high MW polyethylene (UPE), polyfluorotetraethylene (PTFE), nylon, and combination thereof.
11. The method of any one of the preceding claims, wherein the method further comprises washing the captured precipitated mRNA before eluting.
12. The method of claim 11, wherein the washing step comprises multiple rinse cycles using a wash solution comprising a guanidinium buffer and ethanol, followed by about 70-80% ethanol.
13. The method of claim 12, wherein the multiple rinse cycles are more than 5 cycles.
14. The method of any one of the preceding claims, wherein the eluting step comprises re-solubilizing the captured precipitated mRNA with RNase-free water.
15. The method of claim 14, wherein the RNase-free water is re-circulated for 5-10 minutes.
16. The method of any one of the preceding claims, wherein the method further comprises a step of dialyzing the purified mRNA solution.
17. The method of claim 16, wherein the purified mRNA solution is dialyzed with 1mM sodium citrate using a 100 kDa molecular weight cut-off (MWCO) membrane.
18. The method of any one of the preceding claims, wherein the mRNA is *in vitro* synthesized and the impure preparation comprises an *in vitro* mRNA synthesis reaction mixture.
19. The method of claim 18, wherein the impure preparation comprises prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis.
20. The method of claim 19, wherein the purified mRNA solution contains less than 1% of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis.
21. The method of claim 19, wherein the purified mRNA solution contains less than 0.5% of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis.
22. The method of claim 19, wherein the purified mRNA solution contains less than 0.1% of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis.

23. The method of claim 19, wherein the purified mRNA solution is substantially free of prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis.
24. The method of claim 23, wherein the prematurely aborted RNA sequences and/or enzyme reagents used in *in vitro* synthesis are measured via silver stain, gel electrophoresis, HPLC, UPLC, and/or capillary electrophoresis.
25. The method of any one of claims 19-24, wherein the prematurely aborted RNA sequences comprise less than 15 bases.
26. The method of any one of claims 19-24, wherein the prematurely aborted RNA sequences comprise about 8-12 bases.
27. The method of any one of claims 19-24, wherein the enzyme reagents used in *in vitro* synthesis comprise T7 RNA polymerase, DNase I, pyrophosphatase, and/or RNase inhibitor.
28. The method of any one of the preceding claims, wherein the mRNA is purified at a scale of or greater than 1 gram, 10 gram, 100 gram, 1 kg, 10 kg, or 100 kg per batch.
29. The method of any one of the preceding claims, wherein the mRNA is purified before a cap and tail are added to the mRNA.
30. The method of any one of claims 1-28, wherein the mRNA is purified after a cap and tail are added to the mRNA.
31. The method of any one of claims 1-28, wherein the mRNA is purified after a cap is added.
32. The method of any one of claims 1-28, wherein the mRNA is purified both before and after a cap and/or tail are added to the mRNA.
33. The method of any one of the preceding claims, wherein the mRNA is or greater than about 1 kb, 1.5 kb, 2 kb, 2.5 kb, 3 kb, 3.5 kb, 4 kb, 4.5 kb, 5 kb, 6 kb, 7 kb, 8 kb, 9 kb, 10 kb, 11 kb, 12 kb, 13 kb, 14 kb, 15 kb, or 20 kb in length.
34. The method of any one of the preceding claims, wherein the mRNA comprises one or more modifications to enhance stability.
35. The method of claim 34, wherein the one or more modifications comprises modified nucleotide and/or modified sugar phosphate backbones.

36. The method of any one of claims 1-33, wherein the mRNA is unmodified.
37. The method of any one of the preceding claims, wherein the purified mRNA has an integrity of or greater than 95%.
38. The method of any one of the preceding claims, wherein the purified mRNA has an integrity of or greater than 98%.
39. The method of any one of the preceding claims, wherein the purified mRNA has an integrity of or greater than 99%.
40. A method of purifying messenger RNA (mRNA), comprising
- (a) precipitating mRNA from an impure preparation; and
 - (b) subjecting the impure preparation comprising precipitated mRNA to tangential flow filtration such that the precipitated mRNA is captured by a filtration membrane while impurities are discarded through permeation; and
 - (c) eluting the captured precipitated mRNA by re-solubilizing the precipitated mRNA, resulting in a purified mRNA solution.
41. A method for manufacturing messenger RNA (mRNA) comprising:
- synthesizing mRNA *in vitro*; and
 - purifying the *in vitro* synthesized mRNA using a method according to any one of the preceding claims.
42. A messenger RNA (mRNA) purified using a method according to any one of the preceding claims.
43. A batch of purified mRNA comprising 5 gram or more of a single mRNA species suitable for administration to a human subject.
44. The batch of claim 43, comprising 10 gram or more of a single mRNA species.
45. The batch of claim 43, comprising 25 gram or more of a single mRNA species.
46. The batch according to any one of claims 42-45, which is substantially free of impurities from an mRNA synthesis process.

47. The batch of claim 46, wherein the batch is substantially free of prematurely aborted RNA sequences, DNA templates, and/or enzyme reagents used in in vitro synthesis of the single mRNA species.

48. The batch according to any one of claims 43-47, wherein the purified mRNA contains less than about 5% of enzyme reagents used in in vitro synthesis.

49. The batch according to any one of claims 43-48, wherein the purified mRNA has an integrity greater than about 95%.

50. A composition comprising messenger RNA purified using a method according to any one of the preceding claims.

51. A composition comprising in vitro synthesized messenger RNA, wherein the composition contains less than 1% of prematurely aborted RNA sequences and/or enzyme reagents used in in vitro synthesis.

52. A composition comprising in vitro synthesized messenger RNA (mRNA), wherein the mRNA has an integrity of or greater than 95%.
