

F00–F99, T36–T50, X60–X84: Mental and behavioural disorders
9 710 events




Figure 1a




Figure 1b




Figure 2a

4/24




Figure 2b

5/24




Figure 3a

6/24




Figure 3b

7/24




Figure 4a

8/24




Figure 4b

9/24




Figure 4c

10/24




Figure 5a

11/24




Figure 5b

12/24




Figure 5c

13/24




Figure 5d

14/24




Figure 5e




Figure 5f




Figure 5g

F00–F99, T36–T50, X60–X84: Mental and behavioural disorders




Figure 6

F01–F09: Mental disorders due to known physiological conditions (940 events)






Figure 7a

F01–F09: Mental disorders due to known physiological conditions (940 events)


Men and women separately



Age at blood sampling



Additional adjustments



Short vs. long term risk




Figure 7b

F30-F39: Mood affective disorders (2656 events)





Figure 8a

F30-F39: Mood affective disorders (2656 events)

Men and women separately

- Women
- Men



Age at blood sampling

- 62-70
- 53-62
- 40-53



Additional adjustments

- Sex, assessment center, BMI and smoking status
- Sex and assessment center



Short vs. long term risk

- 3-10 years from blood sampling
- Within 3 years from blood sampling




Figure 8b

20/24

F40–F48: Anxiety, dissociative, stress related, somatoform and other nonpsychotic mental disorders (2266 events)






Figure 9a

F40–F48: Anxiety, dissociative, stress related, somatoform and other nonpsychotic mental disorders


Men and women separately



Age at blood sampling



Additional adjustments



Short vs. long term risk




Figure 9b

F05: Delirium due to known physiological condition (543 events)






Figure 10a

F05: Delirium due to known physiological condition (543 events)


Men and women separately



Age at blood sampling



Additional adjustments



Short vs. long term risk




Figure 10b

22/24

F32: Major depressive disorder, single episode (2616 events)






Figure 11a

F32: Major depressive disorder, single episode (2616 events)


Men and women separately



Age at blood sampling



Additional adjustments



Short vs. long term risk




Figure 11b

F41: Other anxiety disorders (2013 events)






Figure 12a

F41: Other anxiety disorders (2013 events)


Men and women separately



Age at blood sampling



Additional adjustments



Short vs. long term risk




Figure 12b

R41: Other symptoms and signs involving cognitive functions and awareness (923 events)





Figure 13a

R41: Other symptoms and signs involving cognitive functions and awareness (923 events)


Men and women separately

- Women
- Men




Age at blood sampling

- 61-70
- 53-61
- 40-53



Additional adjustments

- Sex, assessment center, BMI and smoking status
- Sex and assessment center



Short vs. long term risk

- 3-10 years from blood sampling
- Within 3 years from blood sampling




Figure 13b