

Antibody-conjugates for targeting of tumours expressing PTK7**Field of the invention**

[0001] The present invention is in the field of bioconjugation. More specifically, the present invention relates to antibody-drug conjugates for targeted treatment of patients with cancer, in particular PTK7-expressing tumours.

Background

[0002] A promising approach for targeted treatment of tumours entails the conjugation of a multitude (2 to 8) of highly toxic payloads to a monoclonal antibody, thereby generating an antibody-drug conjugate (ADCs). ADCs are well known in the art, as for example described by Chari et al., *Angew. Chem. Int. Ed.* **2014**, 53, 3796 and Beck et al., *Nat. Rev. Drug Discov.* **2017**, 16, 315–37. Mechanistically, the antibody is designed to bind with high specificity to tumour-associated receptor that is overexpressed versus healthy tissue. The ADC is thought to internalize into the tumour cell after binding to the receptor, then to release the toxic payload upon degradation of the antibody and/or the linker in the lysosome.

[0003] Current ADCs are commonly prepared by various conjugation technologies (summarized in Figure 1), mostly based on conjugation to cysteine side chains with maleimides or to lysine side chains with activated esters. To generate an ADC based on native cysteines naturally engaged in disulfide bonds, the thiol in the side-chain can be liberated by subjection of the antibody to a suitable reducing agent such as TCEP or DTT, followed by treatment with a maleimide-functionalized linker-drug. The resulting ADC will typically consist of a mixture of positional isomers, in case only the total of eight free thiols are not comprehensively alkylated. Alternatively, to generate a site-specific ADC, an antibody can be generated by mutating one or more amino acids at defined positions in the sequence to cysteines, the side-chain of which can be selectively liberated for alkylation by a reduction-oxidation sequence. Commonly employed cysteines for site-specific conjugation are LC-41C (light chain 41C), HC-41C (heavy chain 41C), LC-80C, HC-118C, HC-265C, HC-140C, LC-149C, LC-124C, LC-180C, HC-190C, HC-160C, LC-183C, HC-290C, LC-205C, HC-220C, HC-239C, HC-442C. An additional cysteine can also be inserted into the sequence, for example HC-i239C. Besides maleimides as alkylating agents, reaction of cysteine side chain with haloacetamide or vinylbenzene derivatives has also been reported. Besides reaction to natural amino acid side chains, specific unnatural (non-canonical) amino acids can also be engineered into the amino acid sequence of an antibody, thereby providing a unique handle for chemical conjugation, such as ketone, acetylene, azide, cyclic alkyne or cyclic alkene, for reaction with oxime, azide, alkyne or tetrazine, respectively. However, a disadvantage of the latter approach is that the natural sequence of the antibody must be re-engineered, which besides being time-consuming and costly, may lead to instability issues.

[0004] Conjugation through the glycan by an oxidation–ligation sequence is known in the art and has for example been described by Hamann et al. (*Bioconjugate Chem.* **2002**, 13, 47–58). Chemoenzymatic conjugation through the glycan is known in the art and has been described for

the use of sialyltransferase by Boons et al, *Angew. Chem. Int. Ed.* **2014**, 53, 7179, and for the use of a mutant galactosyltransferase by Zhu et al, *mAbs* **2014**, 6, 1 and Cook et al, *Bioconjugate Chem.* **2016**, 27, 1789.

[0005] Chemoenzymatic conjugation through the glycan including first trimming of the glycan is known in the art and has been described by van Geel et al, *Bioconjugate Chem.* **2015**, 26, 2233 and is schematically depicted in Figure 2. In short, the monoclonal antibody is treated with an endoglycosidase to trim the glycan down to the core GlcNAc (attached directly to Asn-297), followed by transfer of an azido-modified sugar under the action of a glycosyltransferase. Various structures of UDP-azidosugars are depicted in Figure 3. One particularly suitable combination involves that transfer of GalNAz **2b** (2-azidoacetyl-N-galactosamine) under the action of a mutant galactosyltransferase GalT(Y289L) as disclosed in WO 2007/095506, EP 2911699 B1 and van Geel et al. Another useful combination entails GlcNAz with α -1,3-mannosyl-glycoprotein-2- β -N-acetylglucosaminyltransferase (MGAT1) and α -1,6-mannosyl-glycoprotein-2- β -N-acetylglucosaminyltransferase (MGAT2) and as disclosed in WO2018/126092 and WO2021/248048.

[0006] Various cyclooctynes for application in metal-free click chemistry are known in the art (Figure 4). In particular, various cyclooctynes such as DIBO (I), DBCO/DIBAC (J), s-DIBO (K), BCN (L) and TMTHSI (T) are regularly applied for conjugation to azide.

[0007] A payload for an ADCs is typically a highly cytotoxic molecule, with IC₅₀-value in low nanomolar or picomolar range, in particular low to medium molecular weight compounds (e.g. about 200 to about 2500 Da). Examples of suitable cytotoxin classes for ADCs include anthracyclines, camptothecins, taxanes, tubulysins, enediynes, inhibitory peptides, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, hemiasterlins, BCL-XL inhibitors, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs) and analogues or prodrugs thereof. A representative set of cytotoxic molecules, and/or synthetic derivatives or prodrugs thereof, with suitable attachment point for conjugation to a monoclonal antibody, is depicted in Figure 5.

[0008] Specific examples of anthracyclins suitable of application in ADCs include (but are not limited to) doxorubicin, daunorubicin, nemorubicin and PNU-159,682.

[0009] Specific examples of camptothecins suitable for application in ADCs include (but are not limited to) SN-38, exatecan, exatecan-S, topotecan, silatecan, cositecan, lurtotecan, gimatecan, belotecan, rubitecan, AMDCPT, G-AMDCPT and other synthetic camptothecins the structures of which are depicted in Figure 6. Various novel camptothecins have been disclosed in EP0296597, WO2019/236954, WO2020/200880, WO2020/219287, CN113816969, CN113710277 and US20180200273.

[0010] Specific examples of enediynes suitable for application in ADCs include (but are not limited to) calicheamicin, esperamicins, shishijimicins and namenamycin and other enediynes as summarized by Galm et al., *Chem. Rev.* **2005**, 105, 739-758.

[0011] Specific examples of auristatins suitable for application in ADCs include (but are not limited to) MMAE, MMAF and PF-06380101 and other auristatins as summarized by Maderna et

al., *Mol. Pharmaceutics* **2015**, 12, 1798-1812.

[0012] Protein tyrosine kinase 7 (PTK7) is a highly conserved member of the pseudokinase family of receptor tyrosine kinases, without observable kinase activity across various species. Genetic and biochemical studies have demonstrated a key function for PTK7 in noncanonical Wnt signaling, and PTK7-deficient embryos exhibit severe developmental defects in planar cell polarity. There is also evidence for additional functions of PTK7 in the vascular endothelial growth factor (VEGF), semaphorin/plexin, and canonical Wnt signaling pathways. Oncogenic functions of PTK7 have been documented in colon cancer, lung cancer, breast cancer, ovarian cancer and esophageal cancer, and PTK7 promotes cell survival and resistance to chemotherapy in acute myeloid leukemia.

[0013] Antibodies against PTK7 are known in the art, as disclosed in US9777070B2 (H23 and H24, also known as cofetuzumab), US9,777,070B2, US9,505,845B2/US9,102,738B2 (4D6, 12C6, 12C6A/1F12 and 7C8) and WO2015/168019 (e.g. Hu23 and Hu58).

[0014] ADCs targeting PTK7 are known in the art, such as for example PF-06647020/ABBV-647 (cofetuzumab pelidotin), based on a humanized, anti-PTK7 antibody and an auristatin microtubule inhibitor payload (Aur0101). PF-06647020 was shown to induce prolonged tumor regression in patient-derived, tumor xenograft preclinical models and a tolerable safety profile and preliminary clinical activity in previously treated patients administered every 2 or 3 weeks, with locally advanced/metastatic, PTK7-positive NSCLC, TNBC, and platinum-resistant OvCa.

Summary of the invention

[0015] The inventors have developed antibody-conjugates that are highly suitable for targeting PTK7-expressing cells, in particular tumours. Thus, the antibody-conjugates according to the invention are highly suitable for treating PTK7-positive cancer, especially colon cancer, lung cancer, breast cancer, ovarian cancer and esophageal cancer.

[0016] In a first aspect, the present invention concerns an antibody-conjugate. Related thereto, in a second aspect, the invention concerns a process for preparing the antibody-conjugate according to the invention. In a third aspect, the invention concerns a method for targeting PTK7-expressing cells. Related thereto are the first medical use of the antibody-conjugate according to the invention, as well as the second medical use for the treatment of cancer. In a last aspect, the invention concerns the use of a mode of conjugation for increasing the therapeutic index of an antibody-conjugate in the treatment of PTK7-expressing tumours.

Detailed description

Definitions

[0017] The verb "to comprise", and its conjugations, as used in this description and in the claims is used in its non-limiting sense to mean that items following the word are included, but items not specifically mentioned are not excluded.

[0018] In addition, reference to an element by the indefinite article "a" or "an" does not exclude the possibility that more than one of the element is present, unless the context clearly requires that

there is one and only one of the elements. The indefinite article "a" or "an" thus usually means "at least one".

[0019] A linker is herein defined as a moiety that connects (covalently links) two or more elements of a compound. A linker may comprise one or more spacer moieties. A spacer-moiety is herein defined as a moiety that spaces (i.e. provides distance between) and covalently links together two (or more) parts of a linker. The linker may be part of e.g. a linker-construct, a linker-conjugate, a linker-payload (e.g. linker-drug) or an antibody-conjugate, as defined below.

[0020] A "hydrophilic group" or "polar linker" is herein defined as any molecular structure containing one or more polar functional groups that imparts improved polarity, and therefore improved aqueous solubility, to the molecule it is attached to. Preferred hydrophilic groups are selected from a carboxylic acid group, an alcohol group, an ether group, a polyethylene glycol group, an amino group, an ammonium group, a sulfonate group, a phosphate group, an acyl sulfamide group or a carbamoyl sulfamide group. In addition to higher solubility other effects of the hydrophilic group include improved click conjugation efficiency, and, once incorporated into an antibody-drug conjugate: less aggregation, improved pharmacokinetics resulting in higher efficacy and in vivo tolerability.

[0021] The term "salt thereof" means a compound formed when an acidic proton, typically a proton of an acid, is replaced by a cation, such as a metal cation or an organic cation and the like. Where applicable, the salt is a pharmaceutically acceptable salt, although this is not required for salts that are not intended for administration to a patient. For example, in a salt of a compound the compound may be protonated by an inorganic or organic acid to form a cation, with the conjugate base of the inorganic or organic acid as the anionic component of the salt. The term "pharmaceutically accepted" salt means a salt that is acceptable for administration to a patient, such as a mammal (salts with counterions having acceptable mammalian safety for a given dosage regime). Such salts may be derived from pharmaceutically acceptable inorganic or organic bases and from pharmaceutically acceptable inorganic or organic acids. "Pharmaceutically acceptable salt" refers to pharmaceutically acceptable salts of a compound, which salts are derived from a variety of organic and inorganic counter ions known in the art and include, for example, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, etc., and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, formate, tartrate, besylate, mesylate, acetate, maleate, oxalate, etc.

[0022] The term "enediyne" or "enediyne antibiotic" or "enediyne-containing cytotoxin" refers to any cytotoxin characterized by the presence of a 3-ene-1,5-diyne structural feature as part of a cyclic molecule as known in the art and include neocarzinostatin (NCS), C-1027, kedarcidin (KED), maduropeptin (MDP), N1999A2, the sporolides (SPO), the cyanosporasides (CYA and CYN), and the fijiolides, calicheamicins (CAL), the esperamicins (ESP), dynemicin (DYN), namenamycin, shishijimicin, and uncialamycin (UCM).

[0023] The term "alkylaminosugar" as used herein means a tetrahydropyranyl moiety connected to an alcohol function via its 2-position, thereby forming an acetal function, and further substituted by (at least) one N-alkylamino group in position 3, 4 or 5. "N-alkylamino group" in this context refers

to an amino group having one methyl, ethyl or 2-propyl group.

[0024] The term “click probe” refers to a functional moiety that is capable of undergoing a click reaction, i.e. two compatible click probes mutually undergo a click reaction such that they are covalently linked in the product. Compatible probes for click reactions are known in the art, and preferably include (cyclic) alkynes and azides. In the context of the present invention, click probe Q in the compound according to the invention is capable of reacting with click probe F on the (modified) protein, such that upon the occurrence of a click reaction, a conjugate is formed wherein the protein is conjugated to the compound according to the invention. Herein, F and Q are compatible click probes.

[0025] An “acylsulfamide moiety” is herein defined as a sulfamide moiety ($\text{H}_2\text{NSO}_2\text{NH}_2$) that is N-acylated or N-carbamoylated on one end of the molecule and N-alkylated (mono or bis) at the other end of the molecule. In the context of the present invention, especially in the examples, this group is also referred to as “HS”.

[0026] A “coding sequence” or a sequence “encoding” an expression product, such as a RNA, polypeptide, protein, or enzyme, is a nucleotide sequence that, when expressed, results in the production of that RNA, polypeptide, protein, or enzyme, i.e., the nucleotide sequence encodes an amino acid sequence for that polypeptide, protein or enzyme. A coding sequence for a protein may include a start codon (usually ATG) and a stop codon.

[0027] The term “gene” means a DNA sequence that codes for, or corresponds to, a particular sequence of amino acids which comprises all or part of one or more proteins or enzymes, and may or may not include regulatory DNA sequences, such as promoter sequences, which determine for example the conditions under which the gene is expressed. Some genes, which are not structural genes, may be transcribed from DNA to RNA, but are not translated into an amino acid sequence. Other genes may function as regulators of structural genes or as regulators of DNA transcription. In particular, the term gene may be intended for the genomic sequence encoding a protein, i.e. a sequence comprising regulator, promoter, intron and exon sequences.

[0028] The term “glycoprotein” is herein used in its normal scientific meaning and refers to a protein comprising one or more monosaccharide or oligosaccharide chains (“glycans”) covalently bonded to the protein. A glycan may be attached to a hydroxyl group on the protein (O-linked-glycan), e.g. to the hydroxyl group of serine, threonine, tyrosine, hydroxylysine or hydroxyproline, or to an amide function on the protein (N-glycoprotein), e.g. asparagine or arginine, or to a carbon on the protein (C-glycoprotein), e.g. tryptophan. A glycoprotein may comprise more than one glycan, may comprise a combination of one or more monosaccharide and one or more oligosaccharide glycans, and may comprise a combination of N-linked, O-linked and C-linked glycans. It is estimated that more than 50% of all proteins have some form of glycosylation and therefore qualify as glycoprotein. Examples of glycoproteins include PSMA (prostate-specific membrane antigen), CAL (candida antarctica lipase), gp41, gp120, EPO (erythropoietin), antifreeze protein and antibodies.

[0029] The term “glycan” is herein used in its normal scientific meaning and refers to a monosaccharide or oligosaccharide chain that is linked to a protein. The term glycan thus refers to the carbohydrate-part of a glycoprotein. The glycan is attached to a protein via the C-1 carbon of

one sugar, which may be without further substitution (monosaccharide) or may be further substituted at one or more of its hydroxyl groups (oligosaccharide). A naturally occurring glycan typically comprises 1 to about 10 saccharide moieties. However, when a longer saccharide chain is linked to a protein, said saccharide chain is herein also considered a glycan. A glycan of a glycoprotein may be a monosaccharide. Typically, a monosaccharide glycan of a glycoprotein consists of a single N-acetylglucosamine (GlcNAc), glucose (Glc), mannose (Man) or fucose (Fuc) covalently attached to the protein. A glycan may also be an oligosaccharide. An oligosaccharide chain of a glycoprotein may be linear or branched. In an oligosaccharide, the sugar that is directly attached to the protein is called the core sugar. In an oligosaccharide, a sugar that is not directly attached to the protein and is attached to at least two other sugars is called an internal sugar. In an oligosaccharide, a sugar that is not directly attached to the protein but to a single other sugar, *i.e.* carrying no further sugar substituents at one or more of its other hydroxyl groups, is called the terminal sugar. For the avoidance of doubt, there may exist multiple terminal sugars in an oligosaccharide of a glycoprotein, but only one core sugar. A glycan may be an O-linked glycan, an N-linked glycan or a C-linked glycan. In an O-linked glycan a monosaccharide or oligosaccharide glycan is bonded to an O-atom in an amino acid of the protein, typically via a hydroxyl group of serine (Ser) or threonine (Thr). In an N-linked glycan a monosaccharide or oligosaccharide glycan is bonded to the protein via an N-atom in an amino acid of the protein, typically via an amide nitrogen in the side chain of asparagine (Asn) or arginine (Arg). In a C-linked glycan a monosaccharide or oligosaccharide glycan is bonded to a C-atom in an amino acid of the protein, typically to a C-atom of tryptophan (Trp).

[0030] The term "antibody" (AB) is herein used in its normal scientific meaning. An antibody is a protein generated by the immune system that is capable of recognizing and binding to a specific antigen. An antibody is an example of a glycoprotein. The term antibody herein is used in its broadest sense and specifically includes monoclonal antibodies, polyclonal antibodies, dimers, multimers, multispecific antibodies (e.g. bispecific antibodies), antibody fragments, and double and single chain antibodies. The term "antibody" is herein also meant to include human antibodies, humanized antibodies, chimeric antibodies and antibodies specifically binding cancer antigen. The term "antibody" is meant to include whole antibodies, but also fragments of an antibody, for example an antibody Fab fragment, F(ab')₂, Fv fragment or Fc fragment from a cleaved antibody, a scFv-Fc fragment, a minibody, a diabody or a scFv. Furthermore, the term includes genetically engineered antibodies and derivatives of an antibody. Antibodies, fragments of antibodies and genetically engineered antibodies may be obtained by methods that are known in the art.

[0031] An antibody may be a natural or conventional antibody in which two heavy chains are linked to each other by disulfide bonds and each heavy chain is linked to a light chain by a disulfide bond. There are two types of light chain, lambda (l) and kappa (k). The light chain includes two domains or regions, a variable domain (VL) and a constant domain (CL). The heavy chain includes four domains, a variable domain (VH) and three constant domains (CH1, CH2 and CH3, collectively referred to as CH). The variable regions of both light (VL) and heavy (VH) chains determine binding recognition and specificity to the antigen. The constant region domains of the light (CL) and heavy

-7-

(CH) chains confer important biological properties, such as antibody chain association, secretion, trans-placental mobility, complement binding, and binding to Fc receptors (FcR). The Fv fragment is the N-terminal part of the Fab fragment of an immunoglobulin and consists of the variable portions of one light chain and one heavy chain. The immunoglobulin can be of any type (e.g. IgG, IgE, IgM, IgD, and IgA), class (e.g. IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass, or allotype (e.g. human G1m1, G1m2, Gm3, non-G1m1 [that, is any allotype other than G1m1], G1m17, G2m23, G3m21, G3m28, G3m1.1, G3m5, G3m13, G3m14, G3m10, G3m15, G3m16, G3m6, G3m24, G3m26, G3m27, A2m1, A2m2, Km1, Km2 and Km3) of immunoglobulin molecule. Preferred allotypes for administration include a non-G1m1 allotype (nG1m1), such as G1m17.1, G1m3, G1m3.1, G1m3.2 or G1m3.1.2. More preferably, the allotype is selected from the group consisting of the G1m17.1 or G1m3 allotype. The antibody may be engineered in the Fc-domain to enhance or nihilate binding to Fc-gamma receptors, as summarized by Saunders et al. *Front. Immunol.* **2019**, *10*, doi: 10.3389/fimmu.2019.01296 and Ward et al., *Mol. Immunol.* **2015**, *67*, 131–141. For example, the combination of Leu234Ala and Leu235Ala (commonly called LALA mutations) eliminate FcγRIIa binding. Elimination of binding to Fc-gamma receptors can also be achieved by mutation of the N297 amino acid to any other amino acid except asparagine, by mutation of the T299 amino acid to any other amino acid except threonine or serine, or by enzymatic deglycosylation or trimming of the fully glycosylated antibody with for example PNGase F or an endoglycosidase. The immunoglobulins can be derived from any species, including human, murine, or rabbit origin. Each chain contains distinct sequence domains.

[0032] A percentage of “sequence identity” may be determined by comparing the two sequences, optimally aligned over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. A sequence “at least 85% identical to a reference sequence” is a sequence having, on its entire length, 85%, or more, for instance 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity with the entire length of the reference sequence.

[0033] The term “CDR” refers to complementarity-determining region: the specificity of the antibody resides in the structural complementarity between the antibody combining site and the antigenic determinant. Antibody combining sites are made up of residues that are primarily from the hypervariable or complementarity determining regions (CDRs). Occasionally, residues from non-hypervariable or framework regions (FR) influence the overall domain structure and hence the combining site. Complementarity Determining Regions or CDRs therefore refer to amino acid sequences which together define the binding affinity and specificity of the natural Fv region of a native immunoglobulin binding site. The light and heavy chains of an immunoglobulin each have three CDRs, designated CDR1-L, CDR2-L, CDR3-L and CDR1-H, CDR2-H, CDR3-H, respectively. A conventional antibody antigen-binding site, therefore, includes six CDRs, comprising the CDR set from each of a heavy and a light chain V region. “CDR”

[0034] The term “monoclonal antibody” or “mAb” as used herein refers to an antibody molecule of a single amino acid sequence, which is directed against a specific antigen, and is not to be

construed as requiring production of the antibody by any particular method. A monoclonal antibody may be produced by a single clone of B cells or hybridoma, but may also be recombinant, i.e. produced by protein engineering.

[0035] The term “chimeric antibody” refers to an engineered antibody which, in its broadest sense, contains one or more regions from one antibody and one or more regions from one or more other antibodies. In an embodiment, a chimeric antibody comprises a VH domain and a VL domain of an antibody derived from a non-human animal, in association with a CH domain and a CL domain of another antibody, in an embodiment, a human antibody. As the non-human animal, any animal such as mouse, rat, hamster, rabbit or the like can be used. A chimeric antibody may also denote a multispecific antibody having specificity for at least two different antigens.

[0036] The term “humanised antibody” refers to an antibody which is wholly or partially of non-human origin and which has been modified to replace certain amino acids, for instance in the framework regions of the VH and VL domains, in order to avoid or minimize an immune response in humans. The constant domains of a humanized antibody are most of the time human CH and CL domains. “Fragments” of (conventional) antibodies comprise a portion of an intact antibody, in particular the antigen binding region or variable region of the intact antibody. Examples of antibody fragments include Fv, Fab, F(ab')₂, Fab', dsFv, (dsFv)₂, scFv, sc(Fv)₂, diabodies, bispecific and multispecific antibodies formed from antibody fragments. A fragment of a conventional antibody may also be a single domain antibody, such as a heavy chain antibody or VHH.

[0037] As used herein “PTK7” also known as colon carcinoma kinase 4 (CCK4) designates the highly conserved member of the pseudokinase family of receptor tyrosine kinases, without observable kinase activity across various species.

Description of figures

[0038] Figure 1 shows a representative set of reactive groups (F) that when present in a biomolecule lead to connecting group Z (**1a-1h**) upon reaction with reactive group Q. Functional groups F can be naturally present or may be artificially introduced (engineered) into a biomolecule at any position of choice.

[0039] Figure 2 schematically displays how an antibody conjugate can be obtained from any monoclonal antibody in a two-stage process. In the first stage, an azido-modified UPD-Gal or UDP-GalNAc may be attached to the monoclonal antibody in a one pot process involving (a) trimming of the glycan by an endoglycosidase (to the core GlcNAc) and (b) attachment of the azido-sugar under the action of a glycosyltransferase (a galactosyltransferase or a mutant thereof or a GalNAc-transferase), thereby generating a β -glycosidic 1-4 linkage between the azido-modified GalNAc and GlcNAc. In the second stage, the azido-modified antibody is reacted with an appropriately functionalized cyclooctyne, thereby generating the antibody conjugate.

[0040] Figure 3 shows several structures of derivatives of UDP sugars of galactosamine, which may be modified with e.g. a 3-mercaptopropionyl group (**2a**), an azidoacetyl group (**2b**), or an azidodifluoroacetyl group (**2c**) at the 2-position, or with an azido group at the 6-position of N-acetyl galactosamine (**2d**).

[0041] Figure 4 shows cyclooctynes (A-T) suitable for metal-free click chemistry, which are preferred options for reactive moiety Q.

[0042] Figure 5 shows a set of exemplary toxic payloads for conjugation to various PTK7-targeting monoclonal antibodies according to the invention. Point of attachment of a linker (to an amino group present in the payload) is indicated with an arrow. Preferred conjugates according to the present invention contain the payloads, including point of attachment, as shown in figure 5.

[0043] Figure 6 shows the structures of various camptothecins, which are preferred payloads in the context of the present invention.

[0044] Figure 7 depicts the structures of the BCN-linker drugs used to prepare ADCs via click chemistry to the azido-sugar remodeled antibody, with the following payloads (3 = exatecan, 4 = MMAE, 5a = calicheamicin γ_1^I , 5b = glycine-calicheamicin γ_1^I , 6 = 6-aminohexanoyl-maytansinoid).

[0045] Figure 8A shows the binding of cofetuzumab to hPTK7 and cPTK7 corrected for background. Figure 8B shows binding of all cofetuzumab-based ADCs to hPTK7, comparable to the corresponding mAb.

[0046] Figure 9 shows the in vitro potency data of A431 cells treated with increasing concentrations of the ADCs cofetuzumab-4, cofetuzumab-5b, and negative control B12-3. Both cofetuzumab ADCs show cell killing with IC₅₀ values of 3.9 nM and 0.1 nM respectively.

[0047] Figures 10A and 10C show the in vivo efficacy data on the PTK7 expressing NCI-H446 tumor model over time (Fig. 10A: up to t = 14 days; Fig. 10C: up to t = 44 days). Cofetuzumab-3 leads to tumor regression with a clear dose-response (2.59, 5.18 and 10.36 mg/kg). Administration of cofetuzumab-5b ADC also shows a dose-response (2.59 and 5.18 mg/kg) leading to tumor growth delay. Other ADCs including cofetuzumab-Aur0101 show only marginal tumor growth delay (similar to isotype control ADC B12-3. Figures 10B and 10D show the body weight of mice over time (Fig. 10B: up to t = 14 days; Fig. 10D: up to t = 28 or 44 days).

[0048] Figure 11 shows the binding of 4D5-3, 12C6a-3, 12C6-3 and 7C8-3 to hPTK7, cPTK7 and rPTK7. 12C6-3 shows relatively the highest binding affinity to both human and cyno PTK7. 12C6a-3 does not show any binding at all. 4D5-3 and 7C8-3 show similar binding to hPTK7 and cPTK7, however, 7C8 clearly has higher binding affinity to rPTK7.

[0049] Figure 12A shows the in vivo efficacy data on the PTK7 expressing NCI-H446 tumor model over time. 12C6-3 does not show tumor growth inhibition as it is similar to the vehicle. Both 7C8-3 and 4D5-3 show marginal tumor growth delay. Cofetuzumab-3 shows tumor growth inhibition, but also early regrowth of the tumor. 12C6-3 shows 99.7% tumor growth inhibition until day 40. Figure 12B shows the body weight of mice over time. Figure 12C shows the survival in a Kaplan-Meier plot of the NCI-H446 model over time.

[0050] Figure 13A shows the in vivo efficacy data on the PTK7 expressing BR1282 PDX tumor model over time. 12C6-3 has a better tumor growth inhibition rate than all 7C8-3, 4D5-3 and 12C6a-3 and even compared to cofetuzumab-Aur0101. When comparing to cofetuzumab-3, both ADCs show a similar response and equally good tumor growth inhibition. Figure 13B shows the body weight of mice over time.

[0051] Figure 14A shows the in vivo efficacy data on the PTK7 expressing NCI-H446 tumor model


-10-

over time. Olaparib as single treatment shows no significant effect. Both 12C6-3 and 12C6-9 show tumor growth inhibition/stasis. The combination of Olaparib with 12C6-3, however, shows a clear additive effect resulting in the complete response of 6 out of 8 mice at day 21, with the study still ongoing. Figure 14B shows the body weight of mice over time.

- 5 **[0052]** Figure 15 shows the in vivo tolerability data of 12C6-3 in female Sprague Dawley rats over time. In all concentrations no toxicity and body weight loss was observed. Therefore the maximum tolerated dose is higher than 150 mg/kg.

The invention

- 10 **[0053]** In a first aspect, the invention concerns antibody-conjugates of general structure (1)




wherein:

- AB is an antibody capable of targeting PTK7-expressing tumours;
- 15 - L is a linker that links Z to D;
- Z is a connecting group;
- L^6 is $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-\text{S}-(\text{L}^7)_w-$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-\text{N}(\text{H})\text{C}(\text{O})\text{CH}_2-$, $-\text{N}(\text{H})\text{C}(\text{O})\text{CF}_2-$ or $-\text{CH}_2-$;
- 20 - D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
- b is 0 or 1;
- 25 - x is 1 or 2; and
- y is 1, 2, 3 or 4.

[0054] Also contemplated within the present invention are salts, preferably pharmaceutically acceptable salts, of the antibody-conjugate according to structure (1).

30

- [0055]** In a second aspect, the invention concerns a process for preparing the antibody-conjugate according to the invention, comprising reacting the compound according to general structure (2) with an antibody (3). The compound according to general structure (2) comprises a reactive moiety Q and the antibody a reactive moiety F which is capable of reacting with Q in a conjugation reaction, wherein Q and F react to form connecting group Z. In this reaction, a conjugate according to general structure (1) is formed. The process according to this aspect this concerns the following bioconjugation reaction:



40

[0056] Here below, the antibody-conjugate according to structure (1) is first defined. The structural features of the antibody-conjugate according to structure (1) also apply to the compound according to structure (2) and the antibody according to structure (3), as those are unchanged in the conjugation reaction except for reactive moieties F and Q, which are transformed into connecting group Z upon reaction of the compound according to structure (2) with an antibody according to structure (3).

[0057] In a third aspect, the invention concerns the application antibody-conjugate according to structure (1), for targeting PTK7-expressing cells. Related thereto, the invention concerns the first medical use and second medical use of the antibody-conjugate according to structure (1).

[0058] As will be understood by the skilled person, the definition of the chemical moieties, as well as their preferred embodiments, apply to all aspects of the invention.

Antibody-conjugate of general structure (1)

[0059] In a first aspect, the invention concerns antibody-conjugates of general structure (1):



wherein:

- AB is an antibody capable of targeting PTK7-expressing tumours;
- b is 0 or 1;
- L^6 is $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-\text{S}-(\text{L}^7)_w-$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-\text{N}(\text{H})\text{C}(\text{O})\text{CH}_2-$, $-\text{N}(\text{H})\text{C}(\text{O})\text{CF}_2-$ or $-\text{CH}_2-$;
- Z is a connecting group;
- L is a linker that links Z to D;
- D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
- x is 1 or 2; and
- y is 1, 2, 3 or 4.

Antibody AB

[0060] The antibody-conjugate according to the invention contains an antibody that is capable of targeting PTK7-expressing cells, in particular tumour cells. PTK7 is a known target for cancer treatment. The term “expressing” is used as common in the art, and refers to overexpression of the target with respect to the expression in healthy tissue. Antibodies capable of targeting PTK7-expressing tumours may also be referred to as anti-PTK7 antibodies, PTK7-targeting antibodies or PTK7-binding antibodies. Anti-PTK7 antibodies selectively bind to PTK7-expressing cells. Anti-PTK7 antibodies are known in the art, and any suitable one can be used in the context of the present invention.

[0061] The Fc regions of these antibodies may have one or more mutations, such as 0 – 10 mutations or 0 – 5 mutations. Especially preferred are mutations that change binding to the FcRn receptor to modulate half-life of the antibody. For example, inclusion of mutations Met-to-Tyr, Ser-to-Thr and Thr-to-Glu in the region of amino acids 254 – 260 of the heavy chain, often called YTE, in the IgG1 Fc results in a ~11-fold higher binding of the antibody to human FcRn, thereby increasing the circulation half-life with a factor ~3.5. For example, the YTE mutations for cofetuzumab are Met254Tyr, Ser256Thr, and Thr258Glu. Therefore, in one embodiment, the antibody has an apparent human FcRn binding affinity $K_{D,app}$ of below 2.5×10^{-6} M, preferably in the range of $0.05 - 0.99 \times 10^{-6}$ M, more preferably in the range of $0.1 - 0.49 \times 10^{-6}$ M, most preferably in the range of $0.2 - 0.4 \times 10^{-6}$ M. The apparent binding affinity $K_{D,app}$ may be determined according to Mackness et al. *MABS*, **2019**, 11(7), 1276-1288. In a preferred embodiment, the antibody AB is the YTE mutant of the preferred antibodies defined here above or below.

[0062] In a preferred embodiment, the antibody is cofetuzumab. Cofetuzumab may also be defined as containing a light chain sequence according to SEQ ID No. 8 and a heavy chain sequence according to SEQ ID No. 7, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0063] In a preferred embodiment, the antibody is 12C6. 12C6 may also be defined as containing a light chain sequence according to SEQ ID No. 20 and a heavy chain sequence according to SEQ ID No. 19, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0064] In a preferred embodiment, the antibody is 12C6a. 12C6a may also be defined as containing a light chain sequence according to SEQ ID No. 22 and a heavy chain sequence according to SEQ ID No. 19, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0065] In a preferred embodiment, the antibody is 4D5. 4D5 may also be defined as containing a light chain sequence according to SEQ ID No. 16 and a heavy chain sequence according to SEQ ID No. 15, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody

according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

- 5 **[0066]** In a preferred embodiment, the antibody is 7C8. 7C8 may also be defined as containing a light chain sequence according to SEQ ID No. 25 and a heavy chain sequence according to SEQ ID No. 26, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of
- 10 calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

- [0067]** In a preferred embodiment, the antibody is Hu23. Hu23 may also be defined as containing a light chain sequence according to SEQ ID No. 33 and a heavy chain sequence according to SEQ
- 15 ID No. 31, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan
- 20 as payload D.

- [0068]** In a preferred embodiment, the antibody is Hu58. Hu58 may also be defined as containing a light chain sequence according to SEQ ID No. 37 and a heavy chain sequence according to SEQ ID No. 35, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody
- 25 according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

- [0069]** Thus, in a preferred embodiment, the antibody is selected from cofetuzumab, 12C6, 12C6a, 4D5, 7C8, Hu23 and Hu58, all as defined above, more preferably from cofetuzumab, 12C6, 4D5 and 7C8, even more preferably from cofetuzumab and 12C6. In one embodiment, the antibody is cofetuzumab. In one embodiment, the antibody is 12C6.
- 30

- [0070]** Preferred antibodies are defined by their V_L and V_H domains, which together form the variable domain that binds to the antigen. Thus, in a preferred embodiment, the antibody contains
- 35 a V_L domain selected from the group consisting of SEQ ID No. 2, 6, 9, 11, 14, 18, 21, 24, 32 and 36, and a V_H domain selected from the group consisting of SEQ ID No. 1, 5, 10, 12, 13, 17, 23, 30 and 34, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or
- 40 even 100 %. In an especially preferred embodiment, the antibody contains a V_L domain of SEQ ID

No. 6, and a V_H domain of SEQ ID No. 5, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %.

[0071] The aforementioned sequence identities refer to the complete sequence of the V_L and V_H domains. While the entire sequences of these domains allow for some variation in the sequence without jeopardizing the binding to PTK7, it is preferred that the complementarity-determining regions (CDRs) have a higher sequence identity, to ensure that the binding to PTK7 is not significantly jeopardized. The location of the CDRs is given in the tables below. Thus, it is preferred that the antibody contains a V_L domain selected from the group consisting of SEQ ID No. 2, 6, 9, 11, 14, 18, 21, 24, 32 and 36, and a V_H domain selected from the group consisting of SEQ ID No. 1, 5, 10, 12, 13, 17, 23, 30 and 34, preferably a V_L domain of SEQ ID No. 6, and a V_H domain of SEQ ID No. 5, wherein the sequence identity of the CDR is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. The skilled person understands that the V_L and V_H domains identified above can be combined with a suitable constant domain to form a complete antibody.

[0072] Preferred V_L domains

SEQ ID No.	CDR (amino acid numbers within the sequence)
2	23-34 50-56 89-97
6	24-38 54-60 93-101
9	24-39 55-61 94-102
11	24-34 50-56 89-97
14	24-34 50-56 89-97
18	24-25 51-57 90-99
21	24-34 50-56 89-97
24	24-34 50-56 89-98

[0073] Preferred V_H domains

SEQ ID No.	CDR (amino acid numbers within the sequence)
1	26-37 52-67 100-109
5	26-35 50-65 97-108
10	31-35 50-66 99-107
12	31-35 50-66 99-109
13	31-35 50-66 99-104
17	31-35 50-65 98-101
23	31-35 50-65 98-101

[0074] In an especially preferred embodiment, the antibody contains a V_L domain of SEQ ID No. 2 and a V_H domain of SEQ ID No. 1, or a V_L domain of SEQ ID No. 6 and a V_H domain of SEQ ID No. 5, or a V_L domain of SEQ ID No. 9 and a V_H domain of SEQ ID No. 10, or a V_L domain of SEQ

-15-

ID No. 11 and a V_H domain of SEQ ID No. 12, or a V_L domain of SEQ ID No. 14 and a V_H domain of SEQ ID No. 13, or a V_L domain of SEQ ID No. 18 or 21 and a V_H domain of SEQ ID No. 17, or a V_L domain of SEQ ID No. 24 and a V_H domain of SEQ ID No. 23, or a V_L domain of SEQ ID No. 32 and a V_H domain of SEQ ID No. 30, or a V_L domain of SEQ ID No. 36 and a V_H domain of SEQ ID No. 34. Herein, the sequence identities as defined above for the complete sequence as well as for the CDR apply.

[0075] Alternatively, the antibody is defined by its light and heavy chains, which together form the antibody. Thus, in a preferred embodiment, the antibody contains a light chain selected from the group consisting of SEQ ID No. 4, 8, 16, 20, 22, 26, 33 and 37, and a heavy chain selected from the group consisting of SEQ ID No. 3, 7, 15, 19, 25, 31 and 35, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %. In a further preferred embodiment, the antibody contains a light chain of SEQ ID No. 8, and a heavy chain of SEQ ID No. 7, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %.

[0076] The aforementioned sequence identities refer to the complete sequence of the light and heavy chains. While the entire sequences of these chains allow for some variation in the sequence without jeopardizing the binding to PTK7, it is preferred that the CDRs have a higher sequence identity, to ensure that the binding to PTK7 is not significantly jeopardized. The location of the CDRs is given in the tables below. Thus, it is preferred that the antibody contains a light chain selected from the group consisting of SEQ ID No. 4, 8, 16, 20, 22, 26, 33 and 37, and a heavy chain selected from the group consisting of SEQ ID No. 3, 7, 15, 19, 25, 31 and 35, wherein the sequence identity of the CDR is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.

[0077] Preferred light chains

SEQ ID No.	CDR (amino acid numbers within the sequence)
4	23-34 50-56 89-97
8	24-38 54-60 93-101
16	24-34 50-56 89-97
20	24-25 51-57 90-99
22	24-34 50-56 89-97
26	24-34 50-56 89-98

[0078] Preferred heavy chains

SEQ ID No.	CDR (amino acid numbers within the sequence)
3	26-37 52-67 100-109
7	26-35 50-65 97-108

15	31-35 50-66 99-104
19	31-35 50-65 98-101
25	31-35 50-65 99-115

[0079] In an especially preferred embodiment, the antibody contains a light chain of SEQ ID No. 4 and a heavy chain of SEQ ID No. 3, or a light chain of SEQ ID No. 8 and a heavy chain of SEQ ID No. 7, or a light chain of SEQ ID No. 16 and a heavy chain of SEQ ID No. 15, or a light chain of SEQ ID No. 20 or 22 and a heavy chain of SEQ ID No. 19, or a light chain of SEQ ID No. 26 and a heavy chain of SEQ ID No. 25, or a light chain of SEQ ID No. 33 and a heavy chain of SEQ ID No. 31, or a light chain of SEQ ID No. 37 and a heavy chain of SEQ ID No. 35. Herein, the sequence identities as defined above for the complete sequence as well as for the CDR apply.

10 *Linker L⁶*

[0080] In case reactive group F is directly connected to the antibody, or even part of the antibody structure, linker L⁶ that connects AB to F (for antibodies of structure (3)) or AB to Z (for conjugates of structure (1)) is absent and b = 0. This is for example the case for cysteine conjugation and lysine conjugation. Alternatively, reactive group F may also be introduced onto the antibody using a linker L⁶ that connects AB to F (for antibodies of structure (3)) or AB to Z (for conjugates of structure (1)), in which case L⁶ is present and b = 1. In case L⁶ is present, reactive group F is typically introduced at the glycan of the antibody. This is for example the case for conjugation via an artificially introduced reactive group F, such as for example using transglutaminase or by enzymatic glycan modification (e.g. glycosyltransferase or α -1,3-mannosyl-glycoprotein-2- β -N-acetylglucosaminyl-transferase). For example, a modified sugar residue S(F)_x may be introduced at the glycan, extending the glycan with one monosaccharide residue S, which introduces x reactive groups F on the glycan of an antibody. In a most preferred embodiment, conjugation occurs via the glycan of the antibody and b = 1. The site of conjugation is preferably at the heavy chain of the antibody.

[0081] If present, L⁶ is a linker that links AB to F or Z, and is represented by $\text{--GlcNAc(Fuc)}_w\text{--(G)}_j\text{--S--(L}^7\text{)}_{w'}$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L⁷ is $\text{--N(H)C(O)CH}_2\text{--}$, $\text{--N(H)C(O)CF}_2\text{--}$ or $\text{--CH}_2\text{--}$. Typically, L⁶ is at least partly formed by the glycan of the antibody. All recombinant antibodies, generated in mammalian host systems, contain the conserved N-glycosylation site at the asparagine residue at or close to position 297 of the heavy chain, which is modified by a glycan of the complex type. This naturally occurring glycosylation site of antibodies is preferably used, but other glycosylation sites, including artificially introduced ones, may also be used for the connection of linker L⁶. Thus, in a preferred embodiment, L⁶ is connected to an amino acid of the antibody which is located at a position in the range of 250 – 350 of the heavy chain, preferably in the range of 280 – 310 of the heavy chain, more preferably in the range of 295 – 300 of the heavy chain, most preferably at position 297 of the heavy chain.

[0082] The $\text{--GlcNAc(Fuc)}_w\text{--(G)}_j\text{--}$ of L⁶ is the glycan of the antibody, or part thereof. The $\text{--GlcNAc(Fuc)}_w\text{--(G)}_j\text{--}$ of the glycan thus typically originates from the original antibody, wherein

GlcNAc is an *N*-acetylglucosamine moiety and Fuc is a fucose moiety. Fuc is typically bound to GlcNAc via an α -1,6-glycosidic bond. Normally, antibodies may ($w = 1$) or may not be fucosylated ($w = 0$). In the context of the present invention, the presence of a fucosyl moiety is irrelevant, and similar effects are obtained with fucosylated ($w = 1$) and non-fucosylated ($w = 0$) antibody
 5 conjugates. The GlcNAc residue may also be referred to as the core-GlcNAc residue and is the monosaccharide that is directly attached to the peptide part of the antibody.

[0083] S may be directly connected to the core-GlcNAc(Fuc)_w moiety, i.e. $j = 0$, meaning that the remainder of the glycan is removed from the core-GlcNAc(Fuc)_w moiety before S is attached. Such trimming of glycans is well-known in the art and can be achieved by the action of an
 10 endoglycosidase. Alternatively, there are one or more monosaccharide residues present in between the core-GlcNAc(Fuc)_w moiety and S, i.e. j is an integer in the range of 1 – 10, preferably $j = 2 - 5$. In a preferred embodiment, (G)_j is an oligosaccharide fraction comprising j monosaccharide residues G, wherein j is an integer in the range of 2 – 5. (G)_j is connected to the GlcNAc moiety of GlcNAc(Fuc)_w, typically via a β -1,4 bond. In a preferred embodiment, j is 3, 4 or 5. Although any
 15 monosaccharide that may be present in a glycan may be employed as G, each G is preferably individually selected from the group consisting of galactose, glucose, *N*-acetylgalactosamine, *N*-acetylglucosamine, mannose and *N*-acetylneuraminic acid. More preferred options for G are galactose, *N*-acetylglucosamine, mannose. The inventors found that antibody-conjugates having j below 4 show no or hardly any binding to the Fc-gamma receptor, while antibody conjugates having
 20 j in the range of 4 – 10 do bind to the Fc-gamma receptor. Thus, by selecting a certain value for j , the desired extent of binding to the Fc-gamma receptor can be obtained. It is thus preferred that $j = 0, 3, 4, 5, 6, 7, 8, 9$ or 10, more preferably $j = 0, 3, 4$ or 5, most preferably the antibody is trimmed and $j = 0$.

[0084] S is a sugar or sugar derivative. The term “sugar derivative” is herein used to indicate a
 25 derivative of a monosaccharide sugar, i.e. a monosaccharide sugar comprising substituents and/or functional groups. Suitable examples for S include glucose (Glc), galactose (Gal), mannose (Man), fucose (Fuc), amino sugars and sugar acids, e.g. glucosamine (GlcNH₂), galactosamine (GalNH₂), *N*-acetylglucosamine (GlcNAc), *N*-acetylgalactosamine (GalNAc), sialic acid (Sia) which is also referred to as *N*-acetylneuraminic acid (NeuNAc), and *N*-acetylmuramic acid (MurNAc), glucuronic
 30 acid (GlcA) and iduronic acid (IdoA). Preferably, S is selected from Glc, Gal, GlcNAc and GalNAc. In an especially preferred embodiment, S is GalNAc.

[0085] x is an integer that denotes the number of connecting groups Z (for conjugate (1)) or reactive groups F (for antibody (3)) that are attached to sugar (derivative) S. Thus, the antibody according to the invention contains a moiety S comprising x reactive moieties F. Each of these
 35 reactive moieties F are reacted with a reactive moiety Q of the compound according to general structure (2), such that x connecting groups Z are formed and x compounds according to general structure (2) are attached to a single occurrence of S. x is 1 or 2, preferably $x = 1$.

[0086] Connecting group Z (for conjugate (1)) or reactive group F (for antibody (3)) may be attached directly to S, or there may be a linker L⁷ present in between S and Z or F. L⁷ is a linker
 40 that links S with Z. L⁷ may be present ($w' = 1$ or 2) or absent ($w' = 0$). Typically, each moiety Z may

-18-

be connected to S via a linker L^7 , thus in one embodiment $w' = 0$ of x . Preferably, L^7 is absent and each connecting moiety Z is directly attached to S. If present, L^7 may be selected from $-N(H)C(O)CH_2-$, $-N(H)C(O)CF_2-$ or $-CH_2-$. In a preferred embodiment, $x = 1$ and $w' = 0$ or 1, most preferably $x = 1$ and $w' = 0$.

- 5 **[0087]** y is an integer that denotes the number of sugar(s) (derivative(s)) S, each having x reactive groups F or connected to x connecting groups Z, that are connected to the antibody. y is 1, 2, 3 or 4, preferably $y = 2$ or 4, most preferably $y = 2$. Thus, the antibody contains y moieties S, each of which comprises x reactive moieties F. Each of these reactive moieties F are reacted with reactive moiety Q of the compound according to general structure (2), such that $x + y$ connecting groups Z
- 10 are formed and $x + y$ compounds according to general structure (2) are attached to a single antibody. Each compound according to general structure (2) may contain multiple payloads, e.g. by virtue of branching nitrogen atom N^* in L. It is preferred that each compound according to general structure (2) contains 1 or 2 occurrences of D, most preferably 2 occurrences of D. In an especially preferred embodiment, linker L^1 contains a branching nitrogen atom N^* to which a second
- 15 occurrence of D is connected.

- [0088]** The amount of payload (D) molecules attached to a single antibody is known in the art as the DAR (drug-antibody ratio). In the context of the present invention, it is preferred that DAR is an integer in the range 1 – 8, more preferably 2 or 4, most preferably DAR = 4. Alternatively worded, the DAR is preferably an integer in the range $(x + y)$ to $[(x + y) \times 2]$, most preferably DAR = $[(x + y)$
- 20 $\times 2]$. With preferred values for x of 1 and y of 2, the DAR is preferably 4. It will be appreciated that these are theoretical DAR values, and in practice the DAR may slightly deviate from this value, by virtue of incomplete conjugation. Typically, the conjugates are obtained as a stochastic mixture of antibody-drug conjugates, with DAR values varying between individual conjugates, and depending on the conjugation technique used the DAR may have a broad distribution (e.g. DAR = 0 – 10) or a
- 25 narrow distribution (e.g. DAR = 3-4). In case of such mixture, DAR often refers to the average DAR of the mixture. This is well-known in the art of bioconjugation. However, in case the conjugation occurs via the glycan (i.e. $b = 1$ and L^6 is present), the antibody-conjugates according to the invention have a close-to-theoretical DAR. For example, when the theoretical DAR is 4, DAR values above 3.6 or even above 3.8 are readily obtained, indicating that most antibodies in the reaction
- 30 mixture have reacted completely and have a DAR of 4.

Connecting group Z

- [0089]** Z is a connecting group, which covalently connects both parts of the conjugate according to the invention. The term “connecting group” herein refers to the structural element, resulting from
- 35 the reaction between Q and F, connecting one part of the conjugate with another part of the same conjugate. As will be understood by the person skilled in the art, the nature of a connecting group depends on the type of reaction with which the connection between the parts of said compound is obtained. As an example, when the carboxyl group of $R-C(O)-OH$ is reacted with the amino group of H_2N-R' to form $R-C(O)-N(H)-R'$, R is connected to R' via connecting group Z, and Z may be

represented by the group -C(O)-N(H)-. Since connecting group Z originates from the reaction between Q and F, it can take any form.

[0090] Since more than one reactive moiety F can be present or introduced in an antibody, the antibody-conjugate according to the present invention may contain per biomolecule more than one payload D, such as 1 – 8 payloads D, preferably 1, 2, 3 or 4 payloads D, more preferably 2 or 4 payloads D. The number of payloads is typically an even integer, in view of the symmetric nature of antibodies. In other words, when one side of the antibody is functionalized with F, the symmetrical counterpart will also be functionalized. Alternatively, in case naturally occurring thiol groups of the cysteine residues of a protein are used as F, the value of m can be anything and may vary between individual conjugates.

[0091] In a compound according to structure (1), connecting group Z connects D via linker L to AB, optionally via L⁶. Numerous reactions are known in the art for the attachment of a reactive group Q to a reactive group F. Consequently, a wide variety of connecting groups Z may be present in the conjugate according to the invention. In one embodiment, the reactive group Q is selected from the options described above, preferably as depicted in Figure 1, and complementary reactive groups F and the thus obtained connecting groups Z are known to a person skilled in the art. Several examples of suitable combinations of F and Q, and of connecting group Z³ that will be present in a bioconjugate when a linker-conjugate comprising Q is conjugated to a biomolecule comprising a complementary reactive group F, are shown in Figure 1.

[0092] For example, when F comprises or is a thiol group, complementary groups Q include N-maleimidyl groups and alkenyl groups, and the corresponding connecting groups Z are as shown in Figure 1. When F comprises or is a thiol group, complementary groups Q also include allenamide groups.

[0093] For example, when F comprises or is an amino group, complementary groups Q include ketone groups and activated ester groups, and the corresponding connecting groups Z are as shown in Figure 1.

[0094] For example, when F comprises or is a ketone group, complementary groups Q include (O-alkyl)hydroxylamino groups and hydrazine groups, and the corresponding connecting groups Z are as shown in Figure 1.

[0095] For example, when F comprises or is an alkynyl group, complementary groups Q include azido groups, and the corresponding connecting group Z is as shown in Figure 1.

[0096] For example, when F comprises or is an azido group, complementary groups Q include alkynyl groups, and the corresponding connecting group Z is as shown in Figure 1.

[0097] For example, when F comprises or is a cyclopropenyl group, a trans-cyclooctene group or a cyclooctyne group, complementary groups Q include tetrazinyl groups, and the corresponding connecting group Z is as shown in Figure 1. In these particular cases, Z is only an intermediate structure and will expel N₂, thereby generating a dihydropyridazine (from the reaction with alkene) or pyridazine (from the reaction with alkyne).

[0098] Additional suitable combinations of F and Q, and the nature of resulting connecting group Z³ are known to a person skilled in the art, and are e.g. described in G.T. Hermanson, "Bioconjugate


-20-

Techniques", Elsevier, 3rd Ed. 2013 (ISBN:978-0-12-382239-0), in particular in Chapter 3, pages 229 - 258, incorporated by reference. A list of complementary reactive groups suitable for bioconjugation processes is disclosed in Table 3.1, pages 230 - 232 of Chapter 3 of G.T. Hermanson, "Bioconjugate Techniques", Elsevier, 3rd Ed. 2013 (ISBN:978-0-12-382239-0), and the content of this Table is expressly incorporated by reference herein.

[0099] In a preferred embodiment, connecting group Z is obtained by a cycloaddition or a nucleophilic reaction, preferably wherein the cycloaddition is a [4+2] cycloaddition or a 1,3-dipolar cycloaddition or the nucleophilic reaction is a Michael addition or a nucleophilic substitution. Such a cycloaddition or nucleophilic reaction occurs via a reactive group F, connected to S, and reactive group Q, connected to D via L. Conjugation reactions via cycloadditions or nucleophilic reactions are known to the skilled person, and the skilled person is capable of selecting appropriate reaction partners F and Q, and will understand the nature of the resulting connecting group Z.

[0100] In a first preferred embodiment, Z is formed by a cycloaddition. Preferred cycloadditions are a (4+2)-cycloaddition (e.g. a Diels-Alder reaction) or a (3+2)-cycloaddition (e.g. a 1,3-dipolar cycloaddition). Preferably, the conjugation is the Diels-Alder reaction or the 1,3-dipolar cycloaddition. The preferred Diels-Alder reaction is the inverse-electron demand Diels-Alder cycloaddition. In another preferred embodiment, the 1,3-dipolar cycloaddition is used, more preferably the alkyne-azide cycloaddition, and most preferably wherein Q is or comprises an alkyne group and F is an azido group. Cycloadditions, such as Diels-Alder reactions and 1,3-dipolar cycloadditions are known in the art, and the skilled person knows how to perform them.

[0101] Preferably, Z contains a moiety selected from the group consisting of a triazole, a cyclohexene, a cyclohexadiene, a [2.2.2]-bicyclooctadiene, a [2.2.2]-bicyclooctene, an isoxazoline, an isoxazolidine, a pyrazoline, a piperazine, a thioether, an amide or an imide group. Triazole moieties are especially preferred to be present in Z. In one embodiment, Z comprises a (hetero)cycloalkene moiety, i.e. formed from Q comprising a (hetero)cycloalkyne moiety. In an alternative embodiment, Z comprises a (hetero)cycloalkane moiety, i.e. formed from Q comprising a (hetero)cycloalkene moiety. In a preferred embodiment, Z has the structure (Z1):



30

(Z1)

[0102] Herein, the bond depicted as --- is a single bond or a double bond. Furthermore:


- ring Z is obtained by a cycloaddition, preferably ring Z is selected from (Za) – (Zj) defined below, wherein the carbon atoms labelled with ** correspond to the two carbon atoms of the bond depicted as --- of (Z1) to which ring Z is fused;
- R¹⁵ is independently selected from the group consisting of hydrogen, halogen, -OR¹⁶, -NO₂, -CN, -S(O)₂R¹⁶, -S(O)₃⁽⁻⁾, C₁ – C₂₄ alkyl groups, C₆ – C₂₄ (hetero)aryl groups, C₇ – C₂₄

35


-21-

- alkyl(hetero)aryl groups and C₇ – C₂₄ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R¹⁵ may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R¹⁶ is independently selected from the group consisting of hydrogen, halogen, C₁ – C₂₄ alkyl groups, C₈ – C₂₄ (hetero)aryl groups, C₇ – C₂₄ alkyl(hetero)aryl groups and C₇ – C₂₄ (hetero)arylalkyl groups;
- 5 - Y² is C(R³¹)₂, O, S, S⁽⁺⁾R³¹, S(O)R³¹, S(O)=NR³¹ or NR³¹, wherein S⁽⁺⁾ is a cationic sulphur atom counterbalanced by B⁽⁻⁾, wherein B⁽⁻⁾ is an anion, and wherein each R³¹ individually is R¹⁵ or a connection with D, connected via L;
- 10 - u is 0, 1, 2, 3, 4 or 5;
- u' is 0, 1, 2, 3, 4 or 5, wherein u + u' = 0, 1, 2, 3, 4, 5, 6, 7 or 8;
- v = an integer in the range 8 – 16;
- Ring A is formed by the cycloaddition, and is preferably selected from (Za) – (Zj).
- 15 **[0103]** In case the bond depicted as --- is a double bond, it is preferred that u + u' = 4, 5, 6, 7 or 8. Preferably, the wavy bond labelled with * is connected to S and the wavy bond labelled with ** is connected to L.


- [0104]** It is especially preferred that Z comprises a (hetero)cycloalkene moiety, i.e. the bond depicted as --- is a double bond. In a preferred embodiment, Z is selected from the structures (Z2) – (Z20), depicted here below:
- 20



-22-




- [0105]** Herein, the connection to L is depicted with the wavy bond. B⁽⁻⁾ is an anion, preferably a pharmaceutically acceptable anion. Ring Z is formed by the cycloaddition reaction, and preferably is a triazole, a cyclohexene, a cyclohexadiene, a [2.2.2]-bicyclooctadiene, a [2.2.2]-bicyclooctene, an isoxazoline, an isoxazolidine, a pyrazoline or a piperazine. Most preferably, ring Z is a triazole ring.
- 5 Ring Z may have the structure selected from (Za) – (Zm) depicted below, wherein the carbon atoms labelled with ** correspond to the two carbon atoms of the (hetero)cycloalkane ring of (Z2) – (Z20) to which ring Z is fused. Preferred rings Z are selected from (Za) – (Zj), more preferably from (Za), (Zd) and (Zh), most preferably ring Z has structure (Za). Since the connecting group Z is formed by reaction with a (hetero)cycloalkyne in the context of the present embodiment, the bound
- 10 depicted above as $\equiv$ is a double bond.



-23-

[0106] In a further preferred embodiment, Z is selected from the structures (Z21) – (Z38) and (Z38a), depicted here below:



[0107] Herein, the connection to L is depicted with the wavy bond. In structure (Z38), $B^{(-)}$ is an anion, preferably a pharmaceutically acceptable anion. Ring Z is selected from structures (Za) - (Zm), preferably from structures (Za) - (Zj), as defined above.

[0108] In a preferred embodiment, Z comprises a (hetero)cyclooctene moiety or a (hetero)cycloheptene moiety, preferably according to structure (Z8), (Z26), (Z27), (Z28) or (Z37) or (Z38a), more preferably according to structure (Z8), (Z26), (Z27), (Z28) or (Z37), which are optionally substituted. Each of these preferred options for Z are further defined here below.

[0109] Thus, in a preferred embodiment, Z comprises a heterocycloheptene moiety according to structure (Z37), which is optionally substituted. Preferably, the heterocycloheptene moiety according to structure (Z37) is not substituted.


[0110] In a preferred embodiment, Z comprises a (hetero)cyclooctene moiety according to structure (Z8), more preferably according to (Z29), which is optionally substituted. Preferably, the cyclooctene moiety according to structure (Z8) or (Z29) is not substituted. In the context of the

-24-


present embodiment, Z preferably comprises a (hetero)cyclooctene moiety according to structure (Z39) as shown below, wherein V is $(\text{CH}_2)_l$ and l is an integer in the range of 0 to 10, preferably in the range of 0 to 6. More preferably, l is 0, 1, 2, 3 or 4, more preferably l is 0, 1 or 2 and most preferably l is 0 or 1. In the context of group (Z39), l is most preferably 1. Most preferably, Z is according to structure (Z42), defined further below.

[0111] In an alternative preferred embodiment, Z comprises a (hetero)cyclooctene moiety according to structure (Z26), (Z27) or (Z28), which are optionally substituted. In the context of the present embodiment, Z preferably comprises a (hetero)cyclooctene moiety according to structure (Z40) or (Z41) as shown below, wherein Y^1 is O or NR^{11} , wherein R^{11} is independently selected from the group consisting of hydrogen, a linear or branched $\text{C}_1 - \text{C}_{12}$ alkyl group or a $\text{C}_4 - \text{C}_{12}$ (hetero)aryl group. The aromatic rings in (Z40) are optionally O-sulfonylated at one or more positions, whereas the rings of (Z41) may be halogenated at one or more positions. Preferably, the (hetero)cyclooctene moiety according to structure (Z40) or (Z41) is not further substituted. Most preferably, Z is according to structure (Z43), defined further below.

[0112] In an alternative preferred embodiment, Z comprises a heterocycloheptenyl group and is according to structure (Z37).



[0113] In an especially preferred embodiment, Z comprises a cyclooctenyl group and is according to structure (Z42):



Herein:

- the bond labelled with * is connected to S and the wavy bond labelled with ** is connected to L;
- R^{15} is independently selected from the group consisting of hydrogen, halogen, $-\text{OR}^{16}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{S}(\text{O})_2\text{R}^{16}$, $-\text{S}(\text{O})_3^{(-)}$, $\text{C}_1 - \text{C}_{24}$ alkyl groups, $\text{C}_5 - \text{C}_{24}$ (hetero)aryl groups, $\text{C}_7 - \text{C}_{24}$ alkyl(hetero)aryl groups and $\text{C}_7 - \text{C}_{24}$ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R^{15} may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R^{16} is independently selected from the group consisting of hydrogen, halogen, $\text{C}_1 - \text{C}_{24}$ alkyl groups, $\text{C}_6 - \text{C}_{24}$ (hetero)aryl groups, $\text{C}_7 - \text{C}_{24}$ alkyl(hetero)aryl groups

-25-

and C₇ – C₂₄ (hetero)arylalkyl groups;


- R¹⁸ is independently selected from the group consisting of hydrogen, halogen, C₁ – C₂₄ alkyl groups, C₆ – C₂₄ (hetero)aryl groups, C₇ – C₂₄ alkyl(hetero)aryl groups and C₇ – C₂₄ (hetero)arylalkyl groups;

- 5 - R¹⁹ is selected from the group consisting of hydrogen, halogen, C₁ – C₂₄ alkyl groups, C₆ – C₂₄ (hetero)aryl groups, C₇ – C₂₄ alkyl(hetero)aryl groups and C₇ – C₂₄ (hetero)arylalkyl groups, the alkyl groups optionally being interrupted by one of more hetero-atoms selected from the group consisting of O, N and S, wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are independently optionally substituted, or R¹⁹ is a second
10 occurrence of Z (or Q) or D connected via a spacer moiety; and
- l is an integer in the range 0 to 10.

[0114] In a preferred embodiment of the group according to structure (Z42), R¹⁵ is independently selected from the group consisting of hydrogen, halogen, -OR¹⁶, C₁ – C₆ alkyl groups, C₅ – C₆ (hetero)aryl groups, wherein R¹⁶ is hydrogen or C₁ – C₆ alkyl, more preferably R¹⁵ is independently
15 selected from the group consisting of hydrogen and C₁ – C₆ alkyl, most preferably all R¹⁵ are H. In a preferred embodiment of the group according to structure (Z42), R¹⁸ is independently selected from the group consisting of hydrogen, C₁ – C₆ alkyl groups, most preferably both R¹⁸ are H. In a preferred embodiment of the group according to structure (Z42), R¹⁹ is H. In a preferred embodiment of the group according to structure (Z42), l is 0 or 1, more preferably l is 1.

20

[0115] In an especially preferred embodiment, Z comprises a (hetero)cyclooctynyl group and is according to structure (Z43):



25 Herein:



- the bond labelled with * is connected to S and the wavy bond labelled with ** is connected to L;
- R¹⁵ is independently selected from the group consisting of hydrogen, halogen, -OR¹⁶, -NO₂, -CN, -S(O)₂R¹⁶, -S(O)₃(⁻), C₁ – C₂₄ alkyl groups, C₅ – C₂₄ (hetero)aryl groups, C₇ – C₂₄ alkyl(hetero)aryl groups and C₇ – C₂₄ (hetero)arylalkyl groups and wherein the alkyl groups,
30 (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R¹⁵ may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R¹⁶ is independently selected from the group consisting of hydrogen, halogen, C₁ – C₂₄ alkyl groups, C₆ – C₂₄ (hetero)aryl groups, C₇ – C₂₄ alkyl(hetero)aryl groups and C₇ – C₂₄ (hetero)arylalkyl groups;
35

-26-





- Y is N or CR¹⁵.

[0116] In a preferred embodiment of the group according to structure (Z43), R¹⁵ is independently selected from the group consisting of hydrogen, halogen, -OR¹⁶, -S(O)₃⁽⁻⁾, C₁ – C₆ alkyl groups, C₅ – C₆ (hetero)aryl groups, wherein R¹⁶ is hydrogen or C₁ – C₆ alkyl, more preferably R¹⁵ is independently selected from the group consisting of hydrogen and -S(O)₃⁽⁻⁾. In a preferred embodiment of the group according to structure (Z43), Y is N or CH, more preferably Y = N.


[0117] In an especially preferred embodiment, Z comprises a heterocycloheptynyl group and is according to structure (Z37) or (Z38a), preferably according to structure (Z37), wherein ring Z is a triazole:




[0118] In an alternative preferred embodiment, Z comprises a (hetero)cycloalkane moiety, i.e. the bond depicted as — is a single bond. The (hetero)cycloalkane group may also be referred to as a heterocycloalkanyl group or a cycloalkanyl group, preferably a cycloalkanyl group, wherein the (hetero)cycloalkanyl group is optionally substituted. Preferably, the (hetero)cycloalkanyl group is a (hetero)cyclopropanyl group, a (hetero)cyclobutanyl group, a norbornane group, a norbornene group, a (hetero)cycloheptanyl group, a (hetero)cyclooctanyl group, a (hetero)cyclononyl group or a (hetero)cyclodecanyl group, which may all optionally be substituted. Especially preferred are (hetero)cyclopropanyl groups, (hetero)cycloheptanyl group or (hetero)cyclooctanyl groups, wherein the (hetero)cyclopropanyl group, the *trans*-(hetero)cycloheptanyl group or the (hetero)cyclooctanyl group is optionally substituted. Preferably, Z comprises a cyclopropanyl moiety according to structure (Z44), a heterocyclobutane moiety according to structure (Z45), a norbornane or norbornene group according to structure (Z46), a (hetero)cycloheptanyl moiety according to structure (Z47) or a (hetero)cyclooctanyl moiety according to structure (Z48). Herein, Y³ is selected from C(R²³)₂, NR²³ or O, wherein each R²³ is individually hydrogen, C₁ – C₆ alkyl or is connected to L, optionally via a spacer, and the bond labelled — is a single or double bond. In a further preferred embodiment, the cyclopropanyl group is according to structure (Z49). In another preferred embodiment, the (hetero)cycloheptane group is according to structure (Z50) or (Z51). In another preferred embodiment, the (hetero)cyclooctane group is according to structure (Z52), (Z53), (Z54), (Z55) or (Z56).



-27-




[0119] Herein, the R group(s) on Si in (Z50) and (Z51) are typically alkyl or aryl, preferably C₁-C₆ alkyl. Ring Z is typically selected from structures (Zn) – (Zu), wherein the carbon atoms labelled with ** correspond to the two carbon atoms of the (hetero)cycloalkane ring of (Z44) – (Z56) to which ring Z is fused, and the carbon a carbon labelled with * is directly connected to the peptide chain of the antibody. Preferred rings Z are selected from (Zo) – (Zr). Since the connecting group Z is formed by reaction with a (hetero)cycloalkene in the context of the present embodiment, the bond depicted above as --- is a single bond.



[0120] In a second preferred embodiment, Z is formed by a nucleophilic reaction, preferably by a nucleophilic substitution or a Michael addition, preferably by a Michael addition. A preferred Michael reaction is the thiol-maleimide ligation, most preferably wherein Q is maleimide and F is a thiol group. Preferably, the thiol is present in the sidechain of a cysteine residue. In a preferred embodiment, connection group Z comprises a succinimidyl ring or its ring-opened succinic acid amide derivative. Preferred options for connection group Z comprise a moiety selected from (Z57) – (Z71) depicted here below.

-28-



[0121] Herein, the wavy bond(s) labelled with an * is connected to the antibody Ab, optionally via a linker, and the wavy bond without label to the payload, optionally via a linker. In addition, R²⁹ is C₁₋₁₂ alkyl, preferably C₁₋₄ alkyl, most preferably ethyl, and X¹ is O or S, preferably X¹ = O. The nitrogen atom labelled with ** in (Z67)-(Z71) corresponds to the nitrogen atom of the side chain of a lysine residue of the antibody. The carbon atoms of the phenyl group of (Z69) and (Z70) are optionally substituted, preferably optionally fluorinated.

[0122] In a preferred embodiment, connection group Z comprise a moiety selected from (Z1) – (Z71).

10 Linker L

[0123] Linker L connects payload D with connecting group Z (in the conjugate according to structure (1)) or connects payload D with reactive group Q (in the compound according to structure (2)). Linkers are known in the art and may be cleavable or non-cleavable. Linker L preferably contains a self-immolative group or cleavable linker, comprising a peptide spacer and a *para*-aminobenzyloxycarbonyl (PABC) moiety or derivative thereof.

[0124] In a preferred embodiment, linker L as the structure $-(L^1)_n-(L^2)_o-(L^3)_p-(L^4)_q-$, wherein (L⁴)_q is connect to payload D and (L¹)_n is connected to Z or Q. Herein L¹, L², L³ and L⁴ are linkers or linking units and each of n, o, p and q are individually 0 or 1, wherein n + o + p + q is at least 1. In a preferred embodiment, at least linkers L¹ and L² are present (i.e. n = 1; o = 1; p = 0 or 1; q = 0 or 1), more preferably linkers L¹, L² and L³ are present and L⁴ is either present or not (i.e. n = 1; o = 1;

-29-

$p = 1$; $q = 0$ or 1). In one embodiment, linkers L^1 , L^2 , L^3 and L^4 are present (i.e. $n = 1$; $o = 1$; $p = 1$; $q = 1$). In one embodiment, linkers L^1 , L^2 and L^3 are present and L^4 is not (i.e. $n = 1$; $o = 1$; $p = 1$; $q = 0$).


- [0125]** A linker, especially linker L^1 , may contain one or more branch-points for attachment of multiple payloads to a single connecting group. In a preferred embodiment, the linker of the conjugate according to the invention contains a branching moiety. A "branching moiety" in the context of the present invention refers to a moiety that is embedded in a linker connecting three moieties. In other words, the branching moiety comprises at least three bonds to other moieties, typically one bond connecting to Z or Q, one bond to the payload D and one bond to a second payload D. The branching moiety, if present, is preferably embedded in linker L^1 , more preferably part of Sp^2 or as the nitrogen atom of NR^{13} . Any moiety that contains at least three bonds to other moieties is suitable as branching moiety in the context of the present invention. In a preferred embodiment, the branching moiety is selected from a carbon atom, a nitrogen atom, a phosphorus atom, a (hetero)aromatic ring, a (hetero)cycle or a polycyclic moiety. Most preferably, the branching moiety is a nitrogen atom.

Linker L^1


- [0126]** Linker L^1 is either absent ($n = 0$) or present ($n = 1$). Preferably, linker L^1 is present and $n = 1$. L^1 may for example be selected from the group consisting of linear or branched C_1 - C_{200} alkylene groups, C_2 - C_{200} alkenylene groups, C_2 - C_{200} alkynylene groups, C_3 - C_{200} cycloalkylene groups, C_5 - C_{200} cycloalkenylene groups, C_8 - C_{200} cycloalkynylene groups, C_7 - C_{200} alkylarylene groups, C_7 - C_{200} arylalkylene groups, C_8 - C_{200} arylalkenylene groups, C_9 - C_{200} arylalkynylene groups. Optionally the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups may be substituted, and optionally said groups may be interrupted by one or more heteroatoms, preferably 1 to 100 heteroatoms, said heteroatoms preferably being selected from the group consisting of O, $S(O)_y$ and NR^{21} , wherein y is 0, 1 or 2, preferably $y = 2$, and R^{21} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups.

- [0127]** In a preferred embodiment, linker L^1 contains a polar group. Such a polar group may be selected from (poly)ethylene glycoldiamines (e.g. 1,8-diamino-3,6-dioxaoctane or equivalents comprising longer ethylene glycol chains), (poly)ethylene glycol or (poly)ethylene oxide chains, (poly)propylene glycol or (poly)propylene oxide chains and 1, z' -diaminoalkanes (wherein z' is the number of carbon atoms in the alkane, preferably $z' = 1 - 10$), $-(O)_a-C(O)-NH-S(O)_2-NR^{13}-$ (as further defined below, see structure (23)), $-C(S(O)_3^{(-)})-$, $-C(C(O)_2^{(-)})-$, $-S(O)_2-$, $-P(O)_2^{(-)}$, $-O(CH_2CH_2O)_t-$, $-NR^{30}(CH_2CH_2NR^{30})_t-$, and the following two structures:

-30-



- [0128]** The polar group may also contain an amino acid, preferably selected from Arg, Glu, Asp, Ser and Thr. Herein, a and R^{13} are further defined below for structure (23). t is an integer in the range of integer in the range of 0 – 15, preferably 1 – 10, more preferably 2 – 5, most preferably t = 2 or 4. Each R^{30} is individually H, C_{1-12} alkyl, C_{1-12} aryl, C_{1-12} alkaryl or C_{1-12} aralkyl. Linker L^1 may contain more than one such polar group, such as at least two polar groups. The polar group may also be present in a branch of linker L^1 , which branches off a branching moiety as defined elsewhere. Preferable, a nitrogen or carbon atom is used as branching moiety. It is especially preferred to have a $-O(CH_2CH_2O)_t-$ polar group present in a branch.
- [0129]** In a preferred embodiment, Linker L^1 is or comprises a sulfamide group, preferably a sulfamide group according to structure (23):




- [0130]** The wavy lines represent the connection to the remainder of the compound, typically to Q and L^2 , L^3 , L^4 or D, preferably to Q and L^2 . Preferably, the $(O)_aC(O)$ moiety is connected to Q and the NR^{13} moiety to L^2 , L^3 , L^4 or D, preferably to L^2 .
- [0131]** In structure (23), a = 0 or 1, preferably a = 1, and R^{13} is selected from the group consisting of hydrogen, $C_1 - C_{24}$ alkyl groups, $C_3 - C_{24}$ cycloalkyl groups, $C_2 - C_{24}$ (hetero)aryl groups, $C_3 - C_{24}$ alkyl(hetero)aryl groups and $C_3 - C_{24}$ (hetero)arylalkyl groups, the $C_1 - C_{24}$ alkyl groups, $C_3 - C_{24}$ cycloalkyl groups, $C_2 - C_{24}$ (hetero)aryl groups, $C_3 - C_{24}$ alkyl(hetero)aryl groups and $C_3 - C_{24}$ (hetero)arylalkyl groups optionally substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR^{14} wherein R^{14} is independently selected from the group consisting of hydrogen and $C_1 - C_4$ alkyl groups, or R^{13} is D connected to N via a spacer moiety, preferably Sp^2 as defined below, in one embodiment D is connected to N via $-(B)_e-(A)-(B)_g-C(O)-$.
- [0132]** In a preferred embodiment, R^{13} is hydrogen or a $C_1 - C_{20}$ alkyl group, more preferably R^{13} is hydrogen or a $C_1 - C_{16}$ alkyl group, even more preferably R^{13} is hydrogen or a $C_1 - C_{10}$ alkyl group, wherein the alkyl group is optionally substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR^{14} , preferably O, wherein R^{14} is independently selected from the group consisting of hydrogen and $C_1 - C_4$ alkyl groups. In a preferred embodiment, R^{13} is hydrogen. In another preferred embodiment, R^{13} is a $C_1 - C_{20}$ alkyl group, more preferably a $C_1 - C_{16}$ alkyl group, even more preferably a $C_1 - C_{10}$ alkyl group, wherein the alkyl group is optionally interrupted by one or more O-atoms, and wherein the alkyl group is optionally substituted with an -OH group, preferably a terminal -OH group. In this embodiment it is further preferred that R^{13} is a (poly)ethylene glycol chain comprising a terminal -OH group. In another preferred embodiment, R^{13} is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl

-31-

and t-butyl, more preferably from the group consisting of hydrogen, methyl, ethyl, n-propyl and i-propyl, and even more preferably from the group consisting of hydrogen, methyl and ethyl. Yet even more preferably, R^{13} is hydrogen or methyl, and most preferably R^{13} is hydrogen.

[0133] In a preferred embodiment, L^1 is according to structure (24):



(24)

[0134] Herein, a and R^{13} are as defined above, Sp^1 and Sp^2 are independently spacer moieties and b and c are independently 0 or 1. Preferably, $b = 0$ or 1 and $c = 1$, more preferably $b = 0$ and $c = 1$. In one embodiment, spacers Sp^1 and Sp^2 are independently selected from the group consisting of linear or branched C_1 - C_{200} alkylene groups, C_2 - C_{200} alkenylene groups, C_2 - C_{200} alkynylene groups, C_3 - C_{200} cycloalkylene groups, C_5 - C_{200} cycloalkenylene groups, C_8 - C_{200} cycloalkynylene groups, C_7 - C_{200} alkylarylene groups, C_7 - C_{200} arylalkylene groups, C_8 - C_{200} arylalkenylene groups and C_9 - C_{200} arylalkynylene groups, the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR^{16} , wherein R^{16} is independently selected from the group consisting of hydrogen, C_1 - C_{24} alkyl groups, C_2 - C_{24} alkenyl groups, C_2 - C_{24} alkynyl groups and C_3 - C_{24} cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted. When the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups are interrupted by one or more heteroatoms as defined above, it is preferred that said groups are interrupted by one or more O-atoms, and/or by one or more S-S groups.

[0135] More preferably, spacer moieties Sp^1 and Sp^2 , if present, are independently selected from the group consisting of linear or branched C_1 - C_{100} alkylene groups, C_2 - C_{100} alkenylene groups, C_2 - C_{100} alkynylene groups, C_3 - C_{100} cycloalkylene groups, C_5 - C_{100} cycloalkenylene groups, C_8 - C_{100} cycloalkynylene groups, C_7 - C_{100} alkylarylene groups, C_7 - C_{100} arylalkylene groups, C_8 - C_{100} arylalkenylene groups and C_9 - C_{100} arylalkynylene groups, the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR^{16} , wherein R^{16} is independently selected from the group consisting of hydrogen, C_1 - C_{24} alkyl groups, C_2 - C_{24} alkenyl groups, C_2 - C_{24} alkynyl groups and C_3 - C_{24} cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted.

[0136] Even more preferably, spacer moieties Sp^1 and Sp^2 , if present, are independently selected from the group consisting of linear or branched C_1 - C_{50} alkylene groups, C_2 - C_{50} alkenylene groups,

C₂-C₅₀ alkynylene groups, C₃-C₅₀ cycloalkylene groups, C₅-C₅₀ cycloalkenylene groups, C₈-C₅₀ cycloalkynylene groups, C₇-C₅₀ alkylarylene groups, C₇-C₅₀ arylalkylene groups, C₈-C₅₀ arylalkenylene groups and C₉-C₅₀ arylalkynylene groups, the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, wherein R¹⁶ is independently selected from the group consisting of hydrogen, C₁ - C₂₄ alkyl groups, C₂ - C₂₄ alkenyl groups, C₂ - C₂₄ alkynyl groups and C₃ - C₂₄ cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted.

[0137] Yet even more preferably, spacer moieties Sp¹ and Sp², if present, are independently selected from the group consisting of linear or branched C₁-C₂₀ alkylene groups, C₂-C₂₀ alkenylene groups, C₂-C₂₀ alkynylene groups, C₃-C₂₀ cycloalkylene groups, C₅-C₂₀ cycloalkenylene groups, C₈-C₂₀ cycloalkynylene groups, C₇-C₂₀ alkylarylene groups, C₇-C₂₀ arylalkylene groups, C₈-C₂₀ arylalkenylene groups and C₉-C₂₀ arylalkynylene groups, the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, wherein R¹⁶ is independently selected from the group consisting of hydrogen, C₁ - C₂₄ alkyl groups, C₂ - C₂₄ alkenyl groups, C₂ - C₂₄ alkynyl groups and C₃ - C₂₄ cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted.

[0138] In these preferred embodiments it is further preferred that the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups are unsubstituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, preferably O, wherein R¹⁶ is independently selected from the group consisting of hydrogen and C₁ - C₄ alkyl groups, preferably hydrogen or methyl.

[0139] Most preferably, spacer moieties Sp¹ and Sp², if present, are independently selected from the group consisting of linear or branched C₁-C₂₀ alkylene groups, the alkylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, wherein R¹⁶ is independently selected from the group consisting of hydrogen, C₁ - C₂₄ alkyl groups, C₂ - C₂₄ alkenyl groups, C₂ - C₂₄ alkynyl groups and C₃ - C₂₄ cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted. In this embodiment, it is further preferred that the alkylene groups are unsubstituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, preferably O and/or S-S, wherein R¹⁶ is independently selected from the group consisting of hydrogen and C₁ - C₄ alkyl groups, preferably hydrogen or methyl.

[0140] Preferred spacer moieties Sp¹ and Sp² thus include -(CH₂)_r-, -(CH₂CH₂)_r-, -(CH₂CH₂O)_r-, -(OCH₂CH₂)_r-, -(CH₂CH₂O)_rCH₂CH₂-, -CH₂CH₂(OCH₂CH₂)_r-, -(CH₂CH₂CH₂O)_r-, -(OCH₂CH₂CH₂)_r-,

-33-

5 $-(\text{CH}_2\text{CH}_2\text{CH}_2\text{O})_r\text{CH}_2\text{CH}_2\text{CH}_2-$ and $-\text{CH}_2\text{CH}_2\text{CH}_2(\text{OCH}_2\text{CH}_2\text{CH}_2)_r-$, wherein r is an integer in the range of 1 to 50, preferably in the range of 1 to 40, more preferably in the range of 1 to 30, even more preferably in the range of 1 to 20 and yet even more preferably in the range of 1 to 15. More preferably n is 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10, more preferably 1, 2, 3, 4, 5, 6, 7 or 8, even more preferably 1, 2, 3, 4, 5 or 6, yet even more preferably 1, 2, 3 or 4.

[0141] Alternatively, preferred linkers L^1 may be represented by $-(W)_k-(A)_d-(B)_e-(A)_f-(C(O))_g-$, wherein:

- $d = 0$ or 1 , preferably $d = 1$;
- e is an integer in the range $0 - 10$, preferably $e = 0, 1, 2, 3, 4, 5$ or 6 , preferably an integer in the range $1 - 10$, most preferably $e = 1, 2, 3$ or 4 ;
- $f = 0$ or 1 , preferably $f = 0$;
- wherein $d + e + f$ is at least 1 , preferably in the range $1 - 5$; and preferably wherein $d + f$ is at least 1 , preferably $d + f = 1$.
- $g = 0$ or 1 , preferably $g = 1$;
- $k = 0$ or 1 , preferably $k = 1$;
- A is a sulfamide group according to structure (23);
- B is a $-\text{CH}_2-\text{CH}_2-\text{O}-$ or a $-\text{O}-\text{CH}_2-\text{CH}_2-$ moiety, or $(B)_e$ is a $-(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$ or a $-(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-$ moiety, wherein $e1$ is defined the same way as e ;
- W is $-\text{OC}(\text{O})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{OC}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})\text{O}-$, $-\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})-$, $-\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$ or $-(4\text{-Ph})\text{CH}_2\text{NHC}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$, preferably wherein W is $-\text{OC}(\text{O})\text{NH}-$, $-\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$ or $-\text{C}(\text{O})\text{NH}-$, and wherein m is an integer in the range $0 - 10$, preferably $m = 0, 1, 2, 3, 4, 5$ or 6 , most preferably $m = 2$ or 3 ;
- preferably wherein L^1 is connected to Q via $(W)_k$ and to L^2, L^3, L^4 or D , preferably to L^2 , via $(C(O))_g$, preferably via $C(O)$.

25 **[0142]** In the context of the present embodiment, the wavy lines in structure (23) represent the connection to the adjacent groups such as $(W)_k$, $(B)_e$ and $(C(O))_g$. It is preferred that A is according to structure (23), wherein $a = 1$ and $R^{13} = \text{H}$ or a $\text{C}_1 - \text{C}_{20}$ alkyl group, more preferably $R^{13} = \text{H}$ or methyl, most preferably $R^{13} = \text{H}$.

[0143] Preferred linkers L^1 have structure $-(W)_k-(A)_d-(B)_e-(A)_f-(C(O))_g-$, wherein:

- (a) $k = 0$; $d = 1$; $g = 1$; $f = 0$; $B = -\text{CH}_2-\text{CH}_2-\text{O}-$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.
- (b) $k = 1$; $W = -\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$; $m = 2$; $d = 0$; $(B)_e = -(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$; $f = 0$; $g = 1$; $e1 = 1, 2, 3$ or 4 , preferably $e1 = 1$.
- (c) $k = 1$; $W = -\text{OC}(\text{O})\text{NH}-$; $d = 0$; $B = -\text{CH}_2-\text{CH}_2-\text{O}-$; $g = 1$; $f = 0$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.
- (d) $k = 1$; $W = -\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$; $m = 2$; $d = 0$; $(B)_e = -(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$; $f = 0$; $g = 1$; $e1 = 1, 2, 3$ or 4 , preferably $e1 = 4$.
- (e) $k = 1$; $W = -\text{OC}(\text{O})\text{NH}-$; $d = 0$; $(B)_e = -(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$; $g = 1$; $f = 0$; $e1 = 1, 2, 3$ or 4 , preferably $e1 = 4$.
- (f) $k = 1$; $W = -(4\text{-Ph})\text{CH}_2\text{NHC}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$, $m = 3$; $d = 0$; $(B)_e = -(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$; $g = 1$; $f = 0$; $e1 = 1, 2, 3$ or 4 , preferably $e1 = 4$.

-34-

(g) $k = 0$; $d = 0$; $g = 1$; $f = 0$; $B = -CH_2-CH_2-O-$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.

(h) $k = 1$; $W = -C(O)NH-$; $d = 0$; $g = 1$; $f = 0$; $B = -CH_2-CH_2-O-$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.

[0144] In a preferred embodiment, linker L^1 comprises a branching nitrogen atom, which is located in the backbone between Q or Z and $(L^2)_o$ and which contains a further moiety D as substituent, which is preferably linked to the branching nitrogen atom via a linker. An example of a branching nitrogen atom is the nitrogen atom NR^{13} in structure (23), wherein R^{13} is connected to a second occurrence of D via a spacer moiety. Alternatively, a branching nitrogen atoms may be located within L^1 according to structure $-(W)_k-(A)_d-(B)_e-(A)-(C(O))_g-$. In one embodiment, L^1 is represented by $-(W)_k-(A)_d-(B)_e-(A)-(C(O))_g-N^*[-(A)_d-(B)_e-(A)-(C(O))_g-]_2$, wherein A, B, W, d, e, f, g and k are as defined above and individually selected for each occurrence, and N^* is the branching nitrogen atoms, to which two instances of $-(A)_d-(B)_e-(A)-(C(O))_g-$ are connected. Herein, both $(C(O))_g$ moieties are connected to $-(L^2)_o-(L^3)_p-(L^4)_q-D$, wherein L^2 , L^3 , L^4 , o, p, q and D are as defined above and are each selected individually. In a preferred embodiment, each of L^2 , L^3 , L^4 , o, p, q and D are the same for both moieties connected to $(C(O))_g$.

[0145] Preferred linkers L^1 comprising a branching nitrogen atom have structure $-(W)_k-(A)_d-(B)_e-(A)-(C(O))_g-N^*[-(A')_d-(B')_e-(A')_f-(C(O))_g-]_2$ wherein:

- (i) $k = d = g = e' = 1$; $f = d' = g' = 0$; $W = -C(O)-$; $B = B' = -CH_2-CH_2-O-$; A is according to structure (23) with $a = 0$ and $R^{13} = H$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.
- (j) $k = d = g = e' = g' = 1$; $f = d' = 0$; $W = -C(O)-$; $B = B' = -CH_2-CH_2-O-$; A is according to structure (23) with $a = 0$ and $R^{13} = H$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.

Linker L^2


[0146] Linker L^2 is either absent ($o = 0$) or present ($o = 1$). Preferably, linker L^2 is present and $o = 1$. Linker L^2 is a peptide spacer. The peptide spacer is preferably defined by $(NH-CR^{17}-CO)_n$, wherein R^{17} represents an amino acid side chain as known in the art. Herein, the amino acid may be a natural or a synthetic amino acid. Preferably, the amino acid(s) are all in their L-configuration. n is an integer in the range of 1 – 5, preferably in the range of 2 – 5. Thus, the peptide spacer preferably contains 1 – 5 amino acids. Preferably, the peptide is a dipeptide ($n = 2$), tripeptide ($n = 3$) or tetrapeptide ($n = 4$), most preferably the peptide spacer is a dipeptide. Although any peptide spacer may be used, preferably the peptide spacer is selected from Val-Cit, Val-Ala, Val-Lys, Val-Arg, AcLys-Val-Cit, AcLys-Val-Ala, Glu-Val-Ala, Asp-Val-Ala, iGlu-Val-Ala, Glu-Val-Cit, Asp-Val-Cit, iGlu-Val-Cit, Phe-Cit, Phe-Ala, Phe-Lys, Phe-Arg, Ala-Lys, Leu-Cit, Ile-Cit, Trp-Cit, Ala-Ala-Asn, Ala-Asn, Gly-Gly-Phe-Gly and Lys, more preferably Val-Cit, Val-Ala, Glu-Val-Ala, Val-Lys, Phe-Cit, Phe-Ala, Phe-Lys, Ala-Ala-Asn, more preferably Val-Cit, Val-Ala, Ala-Ala-Asn, most preferably Val-Cit or Val-Ala. Herein, AcLys is acetyllysine and iGlu is isoglutamate. In one embodiment, $L^2 = Val-Cit$. In one embodiment, $L^2 = Val-Ala$.

[0147] R^{17} represents the amino acid side chain, preferably selected from the side chains of alanine, cysteine, aspartic acid, glutamic acid, phenylalanine, glycine, histidine, isoleucine, lysine, acetyllysine, leucine, methionine, asparagine, pyrrolysine, proline, glutamine, arginine, serine,

-35-

threonine, selenocysteine, valine, tryptophan, tyrosine and citrulline. Preferred amino acid side chains are those of Val, Cit, Ala, Lys, Arg, AcLys, Phe, Leu, Ile, Trp, Glu, Asp and Asn, more preferably from the side chains of Val, Cit, Ala, Glu and Lys. Alternatively worded, R¹⁷ is preferably selected from CH₃ (Ala), CH₂CH(CH₃)₂ (Leu), CH₂CH₂CH₂NHC(O)NH₂ (Cit), CH₂CH₂CH₂CH₂NH₂ (Lys), CH₂CH₂CH₂NHC(O)CH₃ (AcLys), CH₂CH₂CH₂NHC(=NH)NH₂ (Arg), CH₂Ph (Phe), CH(CH₃)₂ (Val), CH(CH₃)CH₂CH₃ (Ile), CH₂C(O)NH₂ (Asn), CH₂CH₂C(O)OH (Glu), CH₂C(O)OH (Asp) and CH₂(1*H*-indol-3-yl) (Trp). Especially preferred embodiments of R¹⁷ are CH₃ (Ala), CH₂CH₂CH₂NHC(O)NH₂ (Cit), CH₂CH₂CH₂CH₂NH₂ (Lys), CH₂CH₂C(O)OH (Glu) and CH(CH₃)₂ (Val). Most preferably, R¹⁷ is CH₃ (Ala), CH₂CH₂CH₂NHC(O)NH₂ (Cit), CH₂CH₂CH₂CH₂NH₂ (Lys), or CH(CH₃)₂ (Val).


[0148] In an especially preferred embodiment, the peptide spacer may be represented by general structure (L3):



[0149] Herein, R¹⁷ is as defined above, preferably R¹⁷ is CH₃ (Val) or CH₂CH₂CH₂NHC(O)NH₂ (Cit). The wavy lines indicate the connection to (L¹)_n and (L³)_p, preferably L² according to structure (L3) is connected to (L¹)_n via NH and to (L³)_p via C(O).

Linker L³

[0150] Linker L³ is either absent (p = 0) or present (p = 1). Preferably, linker L³ is present and p = 1. Linker L³ is a self-cleavable spacer, also referred to as self-immolative spacer. Preferably, L³ is para-aminobenzyloxycarbonyl (PABC) derivative, more preferably a PABC derivative according to structure (L4):



[0151] Herein, the wavy lines indicate the connection to Q or Z, L¹ or L², and to L⁴ or D. Typically, the PABC derivative is connected via NH to Q, Z, L¹ or L², preferably to L², and via O to L⁴ or D.

[0152] A is a 5- or 6-membered aromatic or heteroaromatic ring, preferably a 6-membered aromatic or heteroaromatic ring. Suitable 5-membered rings are oxazole, thiazole and furan.

Suitable 6-membered rings are phenyl and pyridyl. In a preferred embodiment, A is 1,4-phenyl, 2,5-pyridyl or 3,6-pyridyl. Most preferably, A is 1,4-phenyl.

[0153] R²¹ is selected from H, R²⁶, C(O)OH and C(O)R²⁶, wherein R²⁶ is C₁ – C₂₄ (hetero)alkyl groups, C₃ – C₁₀ (hetero)cycloalkyl groups, C₂ – C₁₀ (hetero)aryl groups, C₃ – C₁₀ alkyl(hetero)aryl groups and C₃ – C₁₀ (hetero)arylalkyl groups, which are optionally substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR²⁸ wherein R²⁸ is independently

-36-

selected from the group consisting of hydrogen and C₁ – C₄ alkyl groups. Preferably, R²⁶ is C₃ – C₁₀ (hetero)cycloalkyl or polyalkylene glycol. The polyalkylene glycol is preferably a polyethylene glycol or a polypropylene glycol, more preferably –(CH₂CH₂O)_sH or –(CH₂CH₂CH₂O)_sH. The polyalkylene glycol is most preferably a polyethylene glycol, preferably –(CH₂CH₂O)_sH, wherein s is an integer in the range 1 – 10, preferably 1 – 5, most preferably s = 1, 2, 3 or 4. More preferably, R²¹ is H or C(O)R²⁶, wherein R²⁶ = 4-methyl-piperazine or morpholine. Most preferably, R²¹ is H.

Linker L⁴

[0154] Linker L⁴ is either absent (q = 0) or present (q = 1). Preferably, linker L⁴ is present and q = 1. Linker L⁴ is selected from:

- an aminoalkanoic acid spacer according to the structure –NR²²–(C_z-alkylene)–C(O)–, wherein z is an integer in the range 1 – 20 and R²² is H or C₁ – C₄ alkyl;
- an ethyleneglycol spacer according to the structure –NR²²–(CH₂–CH₂–O)_{e6}–(CH₂)_{e7}–C(O)–, wherein e₆ is an integer in the range 1 – 10, e₇ is an integer in the range 1 – 3 and R²² is H or C₁ – C₄ alkyl; and
- a diamine spacer according to the structure –NR²²–(C_z-alkylene)–NR²²–(C(O))_h–, wherein h is 0 or 1, z is an integer in the range 1 – 20 and R²² is H or C₁ – C₄ alkyl.

[0155] Linker L⁴ may be an aminoalkanoic acid spacer, i.e. –NR²²–(C_z-alkylene)–C(O)–, wherein z is an integer in the range 1 to 20, preferably 1 – 10, most preferably 1 – 6. Herein, the aminoalkanoic acid spacer is typically connected to L³ via the nitrogen atom and to D via the carbonyl moiety. Preferred linkers L⁴ are selected from 6-aminohexanoic acid (Ahx, z = 5), β-alanine (z = 2) and glycine (Gly, z = 1), even more preferably 6-aminohexanoic acid or glycine. In one embodiment, L⁴ = 6-aminohexanoic acid. In one embodiment, L⁴ = glycine. Herein, R²² is H or C₁ – C₄ alkyl, preferably R²² is H or methyl, most preferably R²² is H.

[0156] Alternatively, linker L⁴ may be an ethyleneglycol spacer according to the structure –NR²²–(CH₂–CH₂–O)_{e6}–(CH₂)_{e7}–C(O)–, wherein e₆ is an integer in the range 1 – 10, preferably e₆ is in the range 2 – 6, and e₇ is an integer in the range 1 – 3, preferably e₇ is 2. Herein, R²² is H or C₁ – C₄ alkyl, preferably R²² is H or methyl, most preferably R²² is H.

[0157] Alternatively, linker L⁴ may be a diamine spacer according to the structure –NR²²–(C_z-alkylene)–NR²²–(C(O))_h–, wherein h is 0 or 1, z is an integer in the range 1 – 20, preferably an integer in the range 2 – 6, even more preferably z = 2 or 5, most preferably z = 2. R²² is H or C₁ – C₄ alkyl. Herein, R²² is H or C₁ – C₄ alkyl, preferably R²² is H or methyl, most preferably R²² is methyl. Herein, h is preferably 1, in which case linker L⁴ is especially suited for conjugation via a phenolic hydroxyl group present on payload D.

Payload D

[0158] D represents the target molecule D that is or is to be connected to the antibody, which is also referred to in the art as the payload. D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiassterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine

-37-

dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof. Alternatively, D is defined as a pharmaceutically active substance, such as an anti-cancer agent, preferably a cytotoxin. Preferably, D is selected from the group consisting of anthracyclines, camptothecins, maytansinoids, enediynes, amanitins, auristatins and pyrrolobenzodiazepine dimers, more preferably D is selected from the group consisting of enediynes, auristatins and camptothecins. In one embodiment, D is an enediyne. In one embodiment, D is an auristatin. In one embodiment, D is a camptothecin. Most preferably, D is a camptothecin.

[0159] In a preferred embodiment, the enediyne is selected from the group consisting of calicheamicins, esperamicins, shishijimicins and namenamicins, more preferably calicheamicin. In another preferred embodiment, the auristatin is selected from the group consisting of MMAD, MMAE, MMAF or PF-06380101, more preferably MMAE or PF-06380101. In another preferred embodiment, the camptothecin is selected from the compounds depicted in Figure 6, preferably from the group consisting of topotecan, silatecan, cositecan, exatecan, exatecan-S, DXd, SN-38, lurtotecan, gimatecan, belotecan, rubitecan, AMDCPT and G-AMDCPT, more preferably exatecan or DXd, most preferably exatecan.


[0160] In an especially preferred embodiment, D is selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably from the group consisting of calicheamicin, MMAE, and exatecan, most preferably D is exatecan.

[0161] The compound according to general structure (2) may comprise more than one moiety D. When more than one cytotoxin D is present the cytotoxins D may be the same or different, typically they are the same. In a preferred embodiment, the compound according to general structure (2) contains 1 or 2 occurrences of D, most preferably 2 occurrence of D. Typically, the second occurrence of D is present within linker L, which may contain a branching moiety, typically a branching nitrogen atom, that is connected to the second occurrence of D. Preferably, both occurrences of D are connected to the branching moiety via the same linker. Likewise, the antibody-conjugate according to structure (1) may contain more than one moiety D per connecting group Z.

Preferred antibody-conjugates

[0162] Preferred antibody-conjugates according to the first aspect are selected from the group consisting of compounds (I) – (III), more preferably (II) or (III), most preferably (II). More preferred antibody-conjugates are selected from (X) – (XIII). In one especially preferred embodiment, the antibody-conjugates is selected from (Xa), (Xib), (XIlg), (XIlh) and (XIIle). In an even more preferred embodiment, the antibody-conjugates is selected from (XI) and (XIII), more preferably (Xib) or (XIIle), more preferably the antibody-conjugates is according to (XIII), most preferably according to (XIIle). The structures of these antibody-conjugates are defined here below.

[0163] Antibody-conjugate (I) has the following structure:



wherein:


-38-

- AB, L⁶, Z, D, x and y are as defined above;
- L¹ is a linker represented by $-(A)_d-(B)_e-(A)-(C(O))_g-$, as defined above;
- L² is Val-Cit or Val-Ala;
- L³ is the PABC derivative according to structure (L4);
- 5 - L⁴ is $-N-(C_z\text{-alkylene})-C(O)-$ or $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z and R²² are as defined above;
- q = 0 or 1.

[0164] In the context of antibody-conjugate (I), it is preferred that for L¹, d = 1 (A according to structure (23), it is preferred that a = 1 and R¹³ = H), e = 2, f = 0 and g = 1. In the context of antibody-conjugate (I), it is preferred that L² = Val-Cit. In the context of antibody-conjugate (I), it is preferred that for L³, R²¹ = H. In the context of antibody-conjugate (I), it is preferred that in case q = 1, then z = 1 or 5.

[0165] In a particularly preferred embodiment, the antibody-conjugate according to structure (I) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0166] Antibody-conjugate (II) has the following structure:




wherein:

- AB, L⁶, Z, D, x and y are as defined above;
- L¹ is a linker represented by $-(A)-(B)_e-(C(O))-$, as defined above;
- L² is Val-Cit or Val-Ala;
- 25 - L³ is the PABC derivative according to structure (L4), wherein R²¹ = H.

[0167] In the context of antibody-conjugate (II), it is preferred that for L¹, e = 2, A according to structure (23), it is preferred that a = 1 and R¹³ = H. In the context of antibody-conjugate (II), it is preferred that L² = Val-Cit.

[0168] In a particularly preferred embodiment, the antibody-conjugate according to structure (II) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0169] Antibody-conjugate (III) has the following structure:



wherein:

- AB, L⁶, Z, D, x and y are as defined above;
- L¹ is a linker represented by $-(A)-(B)_e-(C(O))-$, as defined above;
- 40 - L² is Val-Cit or Val-Ala;


-39-

- L^3 is the PABC derivative according to structure (L4), wherein $R^{21} = H$;
- L^4 is $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein R^{22} is as defined above and z is an integer in the range 1 to 6.

[0170] In the context of antibody-conjugate (III), it is preferred that for L^1 , $e = 2$, and with $a = 1$ and $R^{13} = H$. In the context of antibody-conjugate (III), it is preferred that $L^2 = \text{Val-Cit}$. In the context of antibody-conjugate (III), it is preferred that $z = 2$.

[0171] In a particularly preferred embodiment, the antibody-conjugate according to structure (III) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0172] Antibody-conjugate (X) has a linker-payload moiety according to the following structure:




wherein:

- the wavy line indicates the connection to Z;
- L^2 , o and D are as defined above.

[0173] L^2 may be present or absent, preferably L^2 is present and $o = 1$. For preferred antibody-conjugate (Xa), L^2 is according to structure (L3) and R^{17} is CH_3 . For preferred antibody-conjugate (Xb), L^2 is according to structure (L3) and R^{17} is $\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC(O)NH}_2$. The antibody-conjugate (X) preferably has structure (Xa).

[0174] In a particularly preferred embodiment, the antibody-conjugate according to structure (X) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0175] Antibody-conjugate (XI) has a linker-payload moiety according to the following structure:



wherein:

- the wavy line indicates the connection to Z;
- L^2 , o and D are as defined above.


[0176] L^2 may be present or absent, preferably L^2 is present and $o = 1$. For preferred antibody-

-40-

conjugate (XIa), L^2 is according to structure (L3) and R^{17} is CH_3 . For preferred antibody-conjugate (XIb), L^2 is according to structure (L3) and R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$. The antibody-conjugate (XI) preferably has structure (XIb).

- [0177] In a particularly preferred embodiment, the antibody-conjugate according to structure (XI) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with calicheamicin as payload D.

[0178] Antibody-conjugate (XII) has a linker-payload moiety according to the following structure:



(XII)

wherein:

- the wavy line indicates the connection to Z;
- L^2 , L^4 , o, q and D are as defined above.

- [0179] L^2 may be present or absent, preferably L^2 is present and o = 1. For preferred antibody-conjugate (XIIa), L^2 is according to structure (L3) and R^{17} is CH_3 . For preferred antibody-conjugate (XIIb), L^2 is according to structure (L3) and R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$. Preferably, R^{17} = $CH_2CH_2CH_2NHC(O)NH_2$.

- [0180] L^4 may be present or absent. For preferred antibody-conjugate (XIIc), q = 0 and L^4 is absent. For preferred antibody-conjugate (XIId), q = 1 and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 20, and R^{22} is H or $C_1 - C_4$ alkyl.

- [0181] For preferred antibody-conjugate (XIIe), L^2 is according to structure (L3), R^{17} is CH_3 , q = 0 and L^4 is absent. For preferred antibody-conjugate (XIIf), L^2 is according to structure (L3), R^{17} is CH_3 , q = 1 and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 10, preferably wherein z = 2, and R^{22} is H or $C_1 - C_4$ alkyl.

- [0182] For preferred antibody-conjugate (XIIf), L^2 is according to structure (L3), R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$, q = 0 and L^4 is absent. For preferred antibody-conjugate (XIIf), L^2 is according to structure (L3), R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$, q = 1 and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 20, and R^{22} is H or $C_1 - C_4$ alkyl.


-41-

[0183] In the context of antibody-conjugate (XII), it is preferred that z is an integer in the range of 1 – 10, more preferably $z = 2 - 6$, most preferably $z = 2$. In the context of antibody-conjugate (XII), it is further preferred that R^{22} is H or CH_3 .

[0184] In the context of antibody-conjugate (XII), structures (XIlg) and (XIIf) are most preferred.

- 5 **[0185]** In a particularly preferred embodiment, the antibody-conjugate according to structure (XII) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

- 10 **[0186]** Antibody-conjugate (XIII) has a linker-payload moiety according to the following structure:



(XIII)

wherein:

- the wavy line indicates the connection to Z;
 - L^2 , L^4 , o , q and D are as defined above.
- 15 **[0187]** L^2 may be present or absent, preferably L^2 is present and $o = 1$. For preferred antibody-conjugate (XIIIa), L^2 is according to structure (L3) and R^{17} is CH_3 . For preferred antibody-conjugate (XIIIb), L^2 is according to structure (L3) and R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$. Preferably, $R^{17} = CH_3$.
- 20 **[0188]** L^4 may be present or absent. For preferred antibody-conjugate (XIIIc), $q = 0$ and L^4 is absent. For preferred antibody-conjugate (XIId), $q = 1$ and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 20, and R^{22} is H or $C_1 - C_4$ alkyl.
- 25 **[0189]** For preferred antibody-conjugate (XIIIe), L^2 is according to structure (L3), R^{17} is CH_3 , $q = 0$ and L^4 is absent. For preferred antibody-conjugate (XIIIff), L^2 is according to structure (L3), R^{17} is CH_3 , $q = 1$ and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 10, preferably wherein $z = 2$, and R^{22} is H or $C_1 - C_4$ alkyl.
- 30 **[0190]** For preferred antibody-conjugate (XIIIg), L^2 is according to structure (L3), R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$, $q = 0$ and L^4 is absent. For preferred antibody-conjugate (XIIIh), L^2 is according to structure (L3), R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$, $q = 1$ and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 20, and R^{22} is H or $C_1 - C_4$ alkyl.

-42-

[0191] In the context of antibody-conjugate (XIII), it is preferred that z is an integer in the range of 1 – 10, more preferably $z = 2 - 6$, most preferably $z = 2$. In the context of antibody-conjugate (XIII), it is further preferred that R^{22} is H or CH_3 .

[0192] In the context of antibody-conjugate (XIII), structure (XIIIe) is most preferred.

5 **[0193]** In a particularly preferred embodiment, the antibody-conjugate according to structure (XIII) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

10 **[0194]** In one particularly preferred embodiment, antibody-conjugate according to the invention is according to structure (XIb) as defined above, and the antibody is cofetuzumab as defined above as defined above, and the payload is calicheamicin.

[0195] In one particularly preferred embodiment, antibody-conjugate according to the invention is according to structure (XIIIe) as defined above, and the antibody is cofetuzumab as defined above,
15 and the payload is exatecan.

[0196] It is further preferred that these preferred antibody-conjugates are conjugated through the glycan, i.e. $b = 1$, more preferably a trimmed glycan, i.e. $j = 0$. Herein, it is further preferred that $S = \text{GalNAc}$ and $w' = 0$. Herein, it is further preferred that connecting group Z is formed by an azide-alkyne cycloaddition, preferably connecting group $Z = (Z39)$, wherein ring $Z = (Za)$ and $V = CH_2$.

20 Herein, it is further preferred that $x = 1$. Herein, it is further preferred that $y = 2$, more preferably that $x = 1$ and $y = 2$.

[0197] In a most preferred embodiment, the antibody-conjugate according to the invention is according to structure (XIIIe) as defined above, and the antibody is enfortumab YTE as defined above and the payload is exatecan, wherein $b = 1$, $e = 0$, $S = \text{GalNAc}$, $w' = 0$, connecting group $Z = (Z39)$, wherein ring $Z = (Za)$ and $V = CH_2$, $x = 1$ and $y = 2$.

25 = (Z39), wherein ring $Z = (Za)$ and $V = CH_2$, $x = 1$ and $y = 2$.

Compound according to general structure (2)

[0198] The compound has general structure (2):



30 (2)

wherein:

- Q is a reactive moiety;
- L is a linker that links Z to D;
- D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enedynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;

35 **[0199]** The compound of general structure (2) may also be referred to as a "linker-drug construct", for containing linker L and payload D of the final conjugate. Compounds according to general
40 formula (2) can be prepared by the skilled person using standard organic synthesis techniques, and

as exemplified in the examples. Linker L and payload D are defined above in the context of the conjugate according to structure (1).

Reactive moiety Q

5 **[0200]** The compound according to general structure (2) comprises a reactive moiety Q. In the context of the present invention, the term "reactive moiety" may refer to a chemical moiety that comprises a reactive group, but also to a reactive group itself. For example, a cyclooctynyl group is a reactive group comprising a reactive group, namely a C-C triple bond. Similarly, an *N*-maleimidyl group is a reactive group, comprising a C-C double bond as a reactive group. However, a reactive
10 group, for example an azido reactive group, a thiol reactive group or an alkynyl reactive group, may herein also be referred to as a reactive moiety.

[0201] Q serves as chemical handle for the connection to S(F)_x. In other words, Q is reactive towards and complementary to F. Herein, a reactive group is denoted as "complementary" to a reactive group when said reactive group reacts with said reactive group selectively, optionally in the
15 presence of other functional groups. Complementary reactive and functional groups are known to a person skilled in the art, and are described in more detail below. As such, the compound according to general structure (2) is conveniently used in a conjugation reaction, wherein a chemical reaction between Q and F takes place, thereby forming an antibody-conjugate comprising a covalent connection between payload D and the antibody.

20 **[0202]** The exact nature of Q, and F, depends on the type of conjugation reaction that is employed. The skilled person will be able to select the appropriate combination of Q and F. Preferably, Q, and thus also F, is reactive in a cycloaddition or a nucleophilic reaction. Thus, Q preferably comprises a click probe, a thiol, a thiol-reactive moiety, an amine or an amine-reactive moiety, more preferably Q is a click probe, a thiol-reactive moiety or an amine-reactive moiety, most preferably Q is a click
25 probe. The click probe is reactive in a cycloaddition (click reaction) and is preferably selected from an azide, a tetrazine, a triazine, a nitron, a nitrile oxide, a nitrile imine, a diazo compound, an *ortho*-quinone, a dioxothiophene, a sydnone, an alkene moiety and an alkyne moiety. Preferably, the click probe comprises or is an alkene moiety or an alkyne moiety, more preferably wherein the alkene is a (hetero)cycloalkene and/or the alkyne is a terminal alkyne or a (hetero)cycloalkyne. Typical thiol-
30 reactive moieties are selected from maleimide moiety, a haloacetamide moiety, an allenamide moiety, a phosphonamidite moiety, a cyanoethynyl moiety, a vinylsulfone, a vinylpyridine moiety or a methylsulfonylphenyloxadiazole moiety. Most preferably, the thiol-reactive moiety comprises or is a maleimide moiety. Typical amine-reactive moieties are selected from *N*-hydroxysuccinimidyl esters and other activated esters, *p*-nitrophenyl carbonates and other activated carbonates,
35 isocyanates, isothiocyanates, haloacetamides and benzyl halides. In a preferred embodiment, Q is selected from an alkene moiety, an alkyne moiety, a thiol-reactive moiety or an amine-reactive moiety, more preferably an alkene moiety or an alkyne moiety, even more preferably an alkyne moiety. Herein, the alkene is preferably a (hetero)cycloalkene and the alkyne is preferably a terminal alkyne or a (hetero)cycloalkyne. Most preferably, Q is a cyclic (hetero)alkyne moiety. Each
40 of these moieties are further defined here below.

[0203] Thus, in an especially preferred embodiment, Q comprises a cyclic (hetero)alkyne moiety. The alkynyl group may also be referred to as a (hetero)cycloalkynyl group, i.e. a heterocycloalkynyl group or a cycloalkynyl group, wherein the (hetero)cycloalkynyl group is optionally substituted.

- 5 Preferably, the (hetero)cycloalkynyl group is a (hetero)cycloheptynyl group, a (hetero)cyclooctynyl group, a (hetero)cyclononyl group or a (hetero)cyclodecynyl group. Herein, the (hetero)cycloalkynes may optionally be substituted. Preferably, the (hetero)cycloalkynyl group is an optionally substituted (hetero)cycloheptynyl group or an optionally substituted (hetero)cyclooctynyl group. Most preferably, the (hetero)cycloalkynyl group is a (hetero)cyclooctynyl group, wherein the
- 10 (hetero)cyclooctynyl group is optionally substituted.

[0204] In an especially preferred embodiment, Q comprises a (hetero)cycloalkynyl or (hetero)cycloalkenyl group and is according to structure (Q1):



15 Herein:

- the bond depicted as == is a double bond or a triple bond;
- R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $-NO_2$, $-CN$, $-S(O)_2R^{16}$, $-S(O)_3^{(-)}$, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R^{15} may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R^{16} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups;
- Y^2 is $C(R^{31})_2$, O, S, $S^{(+)}R^{31}$, $S(O)R^{31}$, $S(O)=NR^{31}$ or NR^{31} , wherein $S^{(+)}$ is a cationic sulphur atom counterbalanced by $B^{(-)}$, wherein $B^{(-)}$ is an anion, and wherein each R^{31} individually is R^{15} or a connection with D, connected via L;
- u is 0, 1, 2, 3, 4 or 5;
- u' is 0, 1, 2, 3, 4 or 5, wherein $u + u' = 0, 1, 2, 3, 4, 5, 6, 7$ or 8;
- v = an integer in the range 0 – 16;

[0205] Typically, $v = (u + u') \times 2$ (when the connection to L, depicted by the wavy bond, is via Y^2) or $[(u + u') \times 2] - 1$ (when the connection to L, depicted by the wavy bond, is via one of the carbon atoms).

- 35 **[0206]** In a preferred embodiment of structure (Q1), reactive group Q comprises a (hetero)cycloalkynyl group and is according to structure (Q1a):

-45-




Herein,

- R^{15} and Y^2 are as defined above
- 5 - u is 0, 1, 2, 3, 4 or 5;
- u' is 0, 1, 2, 3, 4 or 5, wherein $u + u' = 4, 5, 6, 7$ or 8;
- v = an integer in the range 8 – 16.

[0207] In a preferred embodiment, $u + u' = 4, 5$ or 6, more preferably $u + u' = 5$. In a preferred embodiment, $v = 8, 9$ or 10, more preferably $v = 9$ or 10, most preferably $v = 10$.


10

[0208] In a preferred embodiment, Q is a (hetero)cycloalkynyl group selected from the group consisting of (Q2) – (Q20) and (Q20a) depicted here below.



[0209] Herein, the connection to L, depicted with the wavy bond, may be to any available carbon or nitrogen atom of Q. The nitrogen atom of (Q10), (Q13), (Q14) and (Q15) may bear the connection to L, or may contain a hydrogen atom or be optionally functionalized. B⁽⁻⁾ is an anion, which is preferably selected from ⁽⁻⁾OTf, Cl⁽⁻⁾, Br⁽⁻⁾ or I⁽⁻⁾, most preferably B⁽⁻⁾ is ⁽⁻⁾OTf. In the conjugation reaction, B⁽⁻⁾ does not need to be a pharmaceutically acceptable anion, since B⁽⁻⁾ will exchange with the anions present in the reaction mixture anyway. In case (Q19) is used for Q, the negatively charged counter-ion is preferably pharmaceutically acceptable upon isolation of the conjugate according to the invention, such that the conjugate is readily useable as medicament.

[0210] In a further preferred embodiment, Q is a (hetero)cycloalkynyl group selected from the group consisting of (Q21) – (Q38) and (Q38a) depicted here below.




[0211] In structure (Q38), B⁽⁻⁾ is an anion, which is preferably selected from ⁽⁻⁾OTf, Cl⁽⁻⁾, Br⁽⁻⁾ or I⁽⁻⁾, most preferably B⁽⁻⁾ is ⁽⁻⁾OTf.

[0212] In a preferred embodiment, Q comprises a (hetero)cyclooctyne moiety or a (hetero)cycloheptyne moiety, preferably according to structure (Q8), (Q26), (Q27), (Q28), (Q37) or (Q38a), more preferably according to structure (Q8), (Q26), (Q27), (Q28) or (Q37), which are optionally substituted. Each of these preferred options for Q are further defined here below.


[0213] Thus, in a preferred embodiment, Q comprises a heterocycloheptyne moiety according to structure (Q37), also referred to as a TMTHSI, which is optionally substituted. Preferably, the heterocycloheptyne moiety according to structure (Q37) is not substituted.

-47-

- [0214]** In an alternative preferred embodiment, Q comprises a cyclooctyne moiety according to structure (Q8), more preferably according to (Q29), also referred to as a bicyclo[6.1.0]non-4-yn-9-yl] group (BCN group), which is optionally substituted. Preferably, the cyclooctyne moiety according to structure (Q8) or (Q29) is not substituted. In the context of the present embodiment, Q preferably is a (hetero)cyclooctyne moiety according to structure (Q39) as shown below, wherein V is (CH₂)_l and l is an integer in the range of 0 to 10, preferably in the range of 0 to 6. More preferably, l is 0, 1, 2, 3 or 4, more preferably l is 0, 1 or 2 and most preferably l is 0 or 1. In the context of group (Q39), l is most preferably 1. Most preferably, Q is according to structure (Q42), defined further below.
- [0215]** In an alternative preferred embodiment, Q comprises a (hetero)cyclooctyne moiety according to structure (Q26), (Q27) or (Q28), also referred to as a DIBO, DIBAC, DBCO or ADIBO group, which are optionally substituted. In the context of the present embodiment, Q preferably is a (hetero)cyclooctyne moiety according to structure (Q40) or (Q41) as shown below, wherein Y¹ is O or NR¹¹, wherein R¹¹ is independently selected from the group consisting of hydrogen, a linear or branched C₁ - C₁₂ alkyl group or a C₄ - C₁₂ (hetero)aryl group. The aromatic rings in (Q40) are optionally O-sulfonylated at one or more positions, whereas the rings of (Q41) may be halogenated at one or more positions. Preferably, the (hetero)cyclooctyne moiety according to structure (Q40) or (Q41) is not further substituted. Most preferably, Q is according to structure (Q43), defined further below.
- [0216]** In an alternative preferred embodiment, Q comprises a heterocycloheptynyl group and is according to structure (Q37).



[0217] In an especially preferred embodiment, Q comprises a cyclooctynyl group and is according to structure (Q42):




Herein:

- R¹⁵ is independently selected from the group consisting of hydrogen, halogen, -OR¹⁶, -NO₂, -CN, -S(O)₂R¹⁶, -S(O)₃⁽⁻⁾, C₁ - C₂₄ alkyl groups, C₅ - C₂₄ (hetero)aryl groups, C₇ - C₂₄ alkyl(hetero)aryl groups and C₇ - C₂₄ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally

-48-

- substituted, wherein two substituents R^{15} may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R^{16} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups;
- 5 - R^{18} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups;
- 10 - R^{19} is selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups, the alkyl groups optionally being interrupted by one of more hetero-atoms selected from the group consisting of O, N and S, wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are independently optionally substituted, or R^{19} is a second occurrence of Q or D connected via a spacer moiety; and
- 15 - I is an integer in the range 0 to 10.

- [0218]** In a preferred embodiment of the reactive group according to structure (Q42), R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $C_1 - C_6$ alkyl groups, $C_5 - C_6$ (hetero)aryl groups, wherein R^{16} is hydrogen or $C_1 - C_6$ alkyl, more preferably R^{15} is independently selected from the group consisting of hydrogen and $C_1 - C_6$ alkyl, most preferably all
- 20 R^{15} are H. In a preferred embodiment of the reactive group according to structure (Q42), R^{18} is independently selected from the group consisting of hydrogen, $C_1 - C_6$ alkyl groups, most preferably both R^{18} are H. In a preferred embodiment of the reactive group according to structure (Q42), R^{19} is H. In a preferred embodiment of the reactive group according to structure (Q42), I is 0 or 1, more preferably I is 1.
- 25 **[0219]** In an especially preferred embodiment, Q comprises a (hetero)cyclooctynyl group and is according to structure (Q43):



Herein:



- 30 - R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $-NO_2$, $-CN$, $-S(O)_2R^{16}$, $-S(O)_3^{(-)}$, $C_1 - C_{24}$ alkyl groups, $C_5 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R^{15} may be linked together to form an optionally
- 35 substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R^{16} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups

and C₇ – C₂₄ (hetero)arylalkyl groups;


- Y is N or CR¹⁵.

[0220] In a preferred embodiment of the reactive group according to structure (Q43), R¹⁵ is independently selected from the group consisting of hydrogen, halogen, -OR¹⁶, -S(O)₃⁽⁻⁾, C₁ – C₈ alkyl groups, C₅ – C₈ (hetero)aryl groups, wherein R¹⁶ is hydrogen or C₁ – C₈ alkyl, more preferably R¹⁵ is independently selected from the group consisting of hydrogen and -S(O)₃⁽⁻⁾. In a preferred embodiment of the reactive group according to structure (Q43), Y is N or CH, more preferably Y = N.

[0221] In an especially preferred embodiment, Q comprises a heterocycloheptynyl group and is according to structure (Q37) or (Q38a), preferably according to structure (Q37)




[0222] In an alternative preferred embodiment, Q comprises a cyclic alkene moiety. The alkenyl group Q may also be referred to as a (hetero)cycloalkenyl group, i.e. a heterocycloalkenyl group or a cycloalkenyl group, preferably a cycloalkenyl group, wherein the (hetero)cycloalkenyl group is optionally substituted. Preferably, the (hetero)cycloalkenyl group is a (hetero)cyclopropenyl group, a (hetero)cyclobutenyl group, a norbornene group, a norbornadiene group, a *trans*-(hetero)cycloheptenyl group, a *trans*-(hetero)cyclooctenyl group, a *trans*-(hetero)cyclononenyl group or a *trans*-(hetero)cyclodecenyl group, which may all optionally be substituted. Especially preferred are (hetero)cyclopropenyl groups, *trans*-(hetero)cycloheptenyl group or *trans*-(hetero)cyclooctenyl groups, wherein the (hetero)cyclopropenyl group, the *trans*-(hetero)cycloheptenyl group or the *trans*-(hetero)cyclooctenyl group is optionally substituted. Preferably, Q comprises a cyclopropenyl moiety according to structure (Q44), a heterocyclobutene moiety according to structure (Q45), a norbornene or norbornadiene group according to structure (Q46), a *trans*-(hetero)cycloheptenyl moiety according to structure (Q47) or a *trans*-(hetero)cyclooctenyl moiety according to structure (Q48). Herein, Y³ is selected from C(R²³)₂, NR²³ or O, wherein each R²³ is individually hydrogen, C₁ – C₈ alkyl or is connected to L, optionally via a spacer, and the bond labelled — — is a single or double bond. In a further preferred embodiment, the cyclopropenyl group is according to structure (Q49). In another preferred embodiment, the *trans*-(hetero)cycloheptene group is according to structure (Q50) or (Q51). In another preferred embodiment, the *trans*-(hetero)cyclooctene group is according to structure (Q52), (Q53), (Q54), (Q55) or (Q56).




-50-




(Q49)




(Q50)




(Q51)



(Q52)




(Q53)



(Q54)



(Q55)




(Q56)


[0223] Herein, the R group(s) on Si in (Q50) and (Q51) are typically alkyl or aryl, preferably C₁-C₆ alkyl.

- [0224]** In an alternative preferred embodiment, Q is a thiol-reactive probe. In this embodiment, Q is a reactive group compatible with cysteine conjugation. Such probes are known in the art and may be selected from the group consisting of a maleimide moiety, a haloacetamide moiety, an allenamide moiety, a phosphonamidite moiety, a cyanoethynyl moiety, a vinylsulfone, a vinylpyridine moiety or a methylsulfonylphenyloxadiazole moiety. Most preferably, Q comprises or is a maleimide moiety. Reagents may be monoalkylation type or may be a cross-linker for reaction with two cysteine side-chains.


[0225] In a further preferred embodiment, probe Q is selected from the group consisting of (Q57) – (Q71) depicted here below.




(Q57)




(Q58)




(Q59)




(Q60)




(Q61)




(Q62)




(Q63)




(Q64)




(Q65)




(Q66)




(Q67)




(Q68)



(Q69)



(Q70)




(Q71)

-51-

wherein:


- X^6 is H, halogen, PhS, MeS, preferably a halogen, such as Cl, Br, I;
- X^7 is halogen, PhS, MeS, preferably a halogen, such as Cl, Br, I;
- R^{24} is H or C_{1-12} alkyl, preferably H or C_{1-6} alkyl;
- 5 - R^{25} is H, C_{1-12} alkyl, C_{1-12} aryl, C_{1-12} alkaryl or C_{1-12} aralkyl, preferably H or *para*-methylphenyl;
- wherein the aromatic ring of (Q61) and (Q63) may optionally be a heteroaromatic ring, such as a phenyl or pyridine ring.

- 10 **[0226]** In a preferred embodiment of thiol-reactive probe (Q57), the probe Q is selected from the group consisting of (Q72) – (Q74) depicted here below.



wherein:

- R^{27} is C_{1-12} alkyl, C_{1-12} aryl, C_{1-12} alkaryl or C_{1-12} aralkyl;
 - t is an integer in the range of 0 – 15, preferably 1 – 10.
- 15 **[0227]** In an alternative preferred embodiment, Q is an amine-reactive probe. In this embodiment, Q is a reactive group compatible with lysine conjugation. Such probes are known in the art and may be selected from the group consisting of *N*-hydroxysuccinimidyl groups, isocyanate groups, isothiocyanate groups and benzoyl halide groups. Most preferably, Q comprises or is an *N*-hydroxysuccinimidyl esters or a *p*-nitrophenyl carbonate moiety.
- 20 **[0228]** In a further preferred embodiment, probe Q is selected from the group consisting of (Q75) – (Q79) depicted here below.




Herein, X^2 is halogen, preferably F.

[0229] In a preferred embodiment, Q is selected from the group consisting of (Q1) – (Q79).

25 Antibody according to general structure (3)

[0230] The antibody has general structure (3):



wherein:

- 30 - AB is an antibody capable of targeting PTK7-expressing tumours;

-52-

- b is 0 or 1;
- L^6 is $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-\text{S}-(\text{L}^7)_w-$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-\text{N}(\text{H})\text{C}(\text{O})\text{CH}_2-$, $-\text{N}(\text{H})\text{C}(\text{O})\text{CF}_2-$ or $-\text{CH}_2-$;
- 5 - F is a reactive moiety;
- x is 1 or 2; and
- y is 1, 2, 3 or 4.

[0231] The antibody of general structure (3) may also be referred to as a “(modified) antibody”, for being an antibody containing reactive groups F, wherein the reactive groups F are naturally present or the antibody is modified to incorporate the reactive groups F. The (modified) antibody according to general formula (2) can be prepared by the skilled person using standard organic and/or enzymatic synthesis techniques, and as exemplified in the examples. Antibody AB, linker L^6 , b, x and y are defined above in the context of the conjugate according to structure (1).


15 *Reactive moiety F*

[0232] F is reactive towards Q in the conjugation reaction defined below, preferably wherein the conjugation reaction is a cycloaddition or a nucleophilic reaction. As the skilled person will understand, the options for F are the same as those for Q, provided that F and Q are reactive towards each other. Thus, F preferably comprises a click probe, a thiol, a thiol-reactive moiety, an amine or an amine-reactive moiety, more preferably F is a click probe, a thiol or an amine, most preferably F is a click probe. The click probe is reactive in a cycloaddition (click reaction) and is preferably selected from an azide, a tetrazine, a triazine, a nitron, a nitrile oxide, a nitrile imine, a diazo compound, an *ortho*-quinone, a dioxothiophene, a sydnone, an alkene moiety and an alkyne moiety. Preferably, the click probe comprises or is an azide, a tetrazine, a triazine, a nitron, a nitrile oxide, a nitrile imine, a diazo compound, an *ortho*-quinone, a dioxothiophene or a sydnone, most preferably an azide. Typical thiol-reactive moieties are selected from maleimide moiety, a haloacetamide moiety, an allenamide moiety, a phosphonamidite moiety, a cyanoethynyl moiety, an *ortho*-quinone moiety, a vinylsulfone, a vinylpyridine moiety or a methylsulfonylphenyloxadiazole moiety. Most preferably, the thiol-reactive moiety comprises or is a maleimide moiety. Typical amine-reactive moieties are selected from N-hydroxysuccinimidyl esters, isocyanates, isothiocyanates and benzyl halides. In a preferred embodiment, F is a click probe or a thiol, more preferably F is an azide or a thiol, most preferably F is an azide.

[0233] More than one reactive group F may be present in the antibody. The reactive group F in the antibody may be naturally present or may be placed in the antibody by a specific technique, for example a (bio)chemical or a genetic technique. The reactive group that is placed in the antibody is prepared by chemical synthesis, for example an azide or a terminal alkyne. Methods of preparing modified antibodies are known in the art, e.g. from WO 2014/065661, WO 2016/170186 and WO 2016/053107, which are incorporated herein by reference. From the same documents, the conjugation reaction between the modified antibody and a linker-drug construct is known to the skilled person.

-53-

[0234] Preferably, F is a click probe reactive towards a (hetero)cycloalkene and/or a (hetero)cycloalkyne, and is typically selected from the group consisting of azide, tetrazine, triazine, nitron, nitrile oxide, nitrile imine, diazo compound, *ortho*-quinone, dioxothiophene and sydnone. Preferred structures for the reactive group are structures (F1) – (F10) depicted here below.



5

[0235] Herein, the wavy bond represents the connection to the payload. For (F3), (F4), (F8) and (F9), the payload can be connected to any one of the wavy bonds. The other wavy bond may then be connected to an R group selected from hydrogen, C₁ – C₂₄ alkyl groups, C₂ – C₂₄ acyl groups, C₃ – C₂₄ cycloalkyl groups, C₂ – C₂₄ (hetero)aryl groups, C₃ – C₂₄ alkyl(hetero)aryl groups, C₃ – C₂₄ (hetero)arylalkyl groups and C₁ – C₂₄ sulfonyl groups, each of which (except hydrogen) may optionally be substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR³² wherein R³² is independently selected from the group consisting of hydrogen and C₁ – C₄ alkyl groups. The skilled person understands which R groups may be applied for each of the groups F. For example, the R group connected to the nitrogen atom of (F3) may be selected from alkyl and aryl, and the R group connected to the carbon atom of (F3) may be selected from hydrogen, alkyl, aryl, acyl and sulfonyl. Preferably, the reactive moiety F is selected from azides or tetrazines. Most preferably, the reactive moiety F is an azide.

[0236] In a second preferred embodiment, F is a thiol or precursor thereof. Thiol or precursor thereof F is used in the conjugation reaction to connect the linker-drug construct to the (modified) antibody. F is reactive towards thiol-reactive probe Q in a thiol ligation. The thiol preferably the thiol of the side chain of a cysteine amino acid, which are naturally present within the antibody AB, in which case linker L⁶ is not present (b = 0), although it may also be synthetically introduced, optionally via a linker L⁶. Thiol precursors in the context of bioconjugation are known in the art, and include disulfides, which may be naturally occurring disulfide bridges present in the antibody or synthetically introduced disulfides, which are reduced as known in the art. Preferably, F is a thiol group of a cysteine side chain.

[0237] In a third preferred embodiment, F is an amine or precursor thereof, preferably an amine. Amine or precursor thereof F is used in the conjugation reaction to connect the linker-drug construct to the (modified) antibody. F is reactive towards amine-reactive probe Q in nucleophilic substitution. The amine is typically a primary amine, preferably the amine of the side chain of a lysine amino acid, which are naturally present within the antibody AB, in which case linker L⁶ is not present (b = 0), although it may also be synthetically introduced, optionally via a linker L⁶. Preferably, F is a primary amine group of a lysine side chain.

30

Process for synthesising the antibody-conjugate according to general structure (1)

[0238] In a further aspect, the present invention relates to a process for the preparation of the antibody-conjugate according to the invention, the process comprising the step of reacting Q of the compound according to the invention with a reactive group F of an antibody. The compound according to general structure (2), and preferred embodiments thereof, are described in more detail above. The present process occurs under conditions such that Q is reacted with F to covalently link the antibody AB (3) to the payload D. In the process according to the invention, Q reacts with F, forming a covalent connection between the antibody and the compound according to the invention. Complementary reactive groups Q and reactive groups F are known to the skilled person and are described in more detail below.

[0239] Any conjugation technique known in the art can be employed to prepare the multifunctional antibody constructs according to the invention. Suitable conjugation techniques include thiol ligation, lysine ligation, cycloadditions (e.g. copper-catalysed click reaction, strain-promoted azide-alkyne cycloaddition, strain-promoted quinone-alkyne cycloaddition). Preferred conjugation techniques used in the context of the present invention include nucleophilic reactions and cycloadditions, preferably wherein the cycloaddition is a [4+2] cycloaddition or a [3+2] cycloaddition and the nucleophilic reaction is a Michael addition or a nucleophilic substitution. Suitable conjugation techniques are for example disclosed in G.T. Hermanson, "Bioconjugate Techniques", Elsevier, 3rd Ed. 2013 (ISBN:978-0-12-382239-0), WO 2014/065661, van Geel et al., *Bioconj. Chem.* **2015**, 26, 2233–2242, PCT/EP2021/050594, PCT/EP2021/050598 and NL 2026947.

[0240] Thus, in a preferred embodiment of the conjugation process according to the invention, conjugation is accomplished via a nucleophilic reaction, such as a nucleophilic substitution or a Michael reaction. A preferred nucleophilic reaction is the acylation of a primary amino group with an activated ester. A preferred Michael reaction is the maleimide-thiol reaction, which is widely employed in bioconjugation.


[0241] Thus, in a preferred embodiment of the conjugation process according to the invention, conjugation is accomplished via a cycloaddition. Preferred cycloadditions are a (4+2)-cycloaddition (e.g. a Diels-Alder reaction) or a (3+2)-cycloaddition (e.g. a 1,3-dipolar cycloaddition). Preferably, the conjugation reaction is the Diels-Alder reaction or the 1,3-dipolar cycloaddition. The preferred Diels-Alder reaction is the inverse-electron demand Diels-Alder cycloaddition. In another preferred embodiment, the 1,3-dipolar cycloaddition is used, more preferably the alkyne-azide cycloaddition, and most preferably wherein Q is or comprises an alkyne group and F is an azido group. Cycloadditions, such as Diels-Alder reactions and 1,3-dipolar cycloadditions are known in the art, and the skilled person knows how to perform them.

[0242] The process according to the present aspect preferably concerns a click reaction, more preferably a 1,3-dipolar cycloaddition, most preferably an alkyne/azide cycloaddition. Most preferably, Q is or comprises an alkyne group and F is an azido group. Click reactions, such as 1,3-dipolar cycloadditions, are known in the art, and the skilled person knows how to perform them.

[0243] Thus, the process for preparing the antibody-conjugate according to the invention according to this aspect comprises reacting the modified antibody of structure (3) with a compound according to structure (2), to obtain the antibody-conjugate of structure (1).

[0244] In a preferred embodiment, the process for preparing the antibody-conjugate according to the invention comprises:

- (i) contacting an antibody comprising y core *N*-acetylglucosamine (GlcNAc) moieties, wherein $y = 1, 2, 3$ or 4 , with a compound of the formula $S(F)_x-P$ in the presence of a catalyst, wherein $S(F)_x$ is a sugar derivative comprising x reactive groups F capable of reacting with a reactive group Q , x is 1 or 2 and P is a nucleoside mono- or diphosphate, and wherein the catalyst is capable of transferring the $S(F)_x$ moiety to the core-GlcNAc moiety, to obtain a modified antibody according to Formula (26):



wherein

- AB is an antibody capable of targeting PTK7-expressing tumours;
- Fuc is fucose;
- w is 0 or 1 ; and

- (ii) reacting the modified antibody with a compound according to structure (2):



wherein:

- Q is a reactive moiety;
 - L is a linker that links Z to D ;
 - D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
- to obtain the antibody-conjugate according to structure (1).

Step (i)

[0245] In step (i), an antibody comprising $1, 2, 3$ or 4 core *N*-acetylglucosamine moieties is contacted with a compound of the formula $S(F)_x-P$ in the presence of a catalyst, wherein $S(F)_x$ is a sugar derivative comprising x reactive groups F capable of reacting with a reactive group Q , x is 1 or 2 and P is a nucleoside mono- or diphosphate, and wherein the catalyst is capable of transferring the $S(F)_x$ moiety to the core-GlcNAc moiety. Herein, the antibody is typically an antibody that has been trimmed to a core-GlcNAc residue as described further below. Step (i) affords a modified antibody according to Formula (26).

[0246] The starting material, i.e. the antibody comprising a core-GlcNAc substituent, is known in the art and can be prepared by methods known by the skilled person. In one embodiment, the process according to the invention further comprises the deglycosylation of an antibody glycan

having a core N-acetylglucosamine, in the presence of an endoglycosidase, in order to obtain an antibody comprising a core N-acetylglucosamine substituent, wherein said core N-acetylglucosamine and said core N-acetylglucosamine substituent are optionally fucosylated. Depending on the nature of the glycan, a suitable endoglycosidase may be selected. The endoglycosidase is preferably selected from the group consisting of EndoS, EndoA, EndoE, EfEndo18A, EndoF, EndoM, EndoD, EndoH, EndoT and EndoSH and/or a combination thereof, the selection of which depends on the nature of the glycan. EndoSH is described in PCT/EP2017/052792, see Examples 1 – 3, and SEQ. ID No: 1, which is incorporated by reference herein.

[0247] Structural features S and x are defined above for the antibody-conjugate according to the invention, which equally applies to the present aspect. Compounds of the formula $S(F)_xP$, wherein a nucleoside monophosphate or a nucleoside diphosphate P is linked to a sugar derivative $S(F)_x$, are known in the art. For example Wang *et al.*, *Chem. Eur. J.* **2010**, *16*, 13343–13345, Piller *et al.*, *ACS Chem. Biol.* **2012**, *7*, 753, Piller *et al.*, *Bioorg. Med. Chem. Lett.* **2005**, *15*, 5459–5462 and WO 2009/102820, all incorporated by reference herein, disclose a number of compounds $S(F)_xP$ and their syntheses. In a preferred embodiment nucleoside mono- or diphosphate P in $S(F)_xP$ is selected from the group consisting of uridine diphosphate (UDP), guanosine diphosphate (GDP), thymidine diphosphate (TDP), cytidine diphosphate (CDP) and cytidine monophosphate (CMP), more preferably P is selected from the group consisting of uridine diphosphate (UDP), guanosine diphosphate (GDP) and cytidine diphosphate (CDP), most preferably P = UDP. Preferably, $S(F)_xP$ is selected from the group consisting of GalNAz-UDP, F₂-GalNAz-UDP (N-(azidodifluoro)acetylglactosamine), 6-AzGal-UDP, 6-AzGalNAc-UDP (6-azido-6-deoxy-N-acetylglactosamine-UDP), 4-AzGalNAz-UDP, 6-AzGalNAz-UDP, GlcNAz-UDP, 6-AzGlc-UDP, 6-AzGlcNAz-UDP and 2-(but-3-ynic acid amido)-2-deoxy-galactose-UDP. Most preferably, $S(F)_xP$ is GalNAz-UDP or 6-AzGalNAc-UDP.

[0248] Suitable catalyst that are capable of transferring the $S(F)_x$ moiety to the core-GlcNAc moiety are known in the art. A suitable catalyst is a catalyst wherefore the specific sugar derivative nucleotide $S(F)_xP$ in that specific process is a substrate. More specifically, the catalyst catalyses the formation of a $\beta(1,4)$ -glycosidic bond. Preferably, the catalyst is selected from the group of galactosyltransferases and N-acetylglactosaminyltransferases, more preferably from the group of $\beta(1,4)$ -N-acetylglactosaminyltransferases (GalNAcT) and $\beta(1,4)$ -galactosyltransferases (GalT), most preferably from the group of $\beta(1,4)$ -N-acetylglactosaminyltransferases having a mutant catalytic domain. Suitable catalysts and mutants thereof are disclosed in WO 2014/065661, WO 2016/022027 and WO 2016/170186, all incorporated herein by reference. In one embodiment, the catalyst is a wild-type galactosyltransferase or N-acetylglactosaminyltransferase, preferably an N-acetylglactosaminyltransferase. In an alternative embodiment, the catalyst is a mutant galactosyltransferase or N-acetylglactosaminyltransferases, preferably a mutant N-acetylglactosaminyltransferase. Mutant enzymes described in WO 2016/022027 and WO 2016/170186 are especially preferred. These galactosyltransferase (mutant) enzyme catalysts are able to recognize internal sugars and sugar derivatives as an acceptor. Thus, sugar derivative $S(F)_x$

-57-

is linked to the core-GlcNAc substituent in step (i), irrespective of whether said GlcNAc is fucosylated or not.

[0249] Step (i) is preferably performed in a suitable buffer solution, such as for example phosphate, buffered saline (e.g. phosphate-buffered saline, tris-buffered saline), citrate, HEPES, tris and glycine. Suitable buffers are known in the art. Preferably, the buffer solution is phosphate-buffered saline (PBS) or tris buffer. Step (i) is preferably performed at a temperature in the range of about 4 to about 50 °C, more preferably in the range of about 10 to about 45 °C, even more preferably in the range of about 20 to about 40 °C, and most preferably in the range of about 30 to about 37 °C. Step (i) is preferably performed a pH in the range of about 5 to about 9, preferably in the range of about 5.5 to about 8.5, more preferably in the range of about 6 to about 8. Most preferably, step (i) is performed at a pH in the range of about 7 to about 8.

Step (ii)

[0250] In step (ii), the modified antibody is reacted with a compound according to general structure (2), comprising a reactive group Q capable of reacting with reactive group F and a payload D, to obtain the antibody-conjugate according to structure (1), containing connecting group Z resulting from the reaction between Q and F. Such reaction occurs under condition such that reactive group Q is reacted with the reactive group F of the biomolecule to covalently link the antibody to the compound according to general structure (2). Step (ii) may also be referred to as the conjugation reaction.

[0251] In a preferred embodiment, in step (ii) an azide on an azide-modified antibody reacts with an alkynyl group, preferably a terminal alkynyl group, or a (hetero)cycloalkynyl group of the compound according to general structure (2), via a cycloaddition reaction. This cycloaddition reaction of a molecule comprising an azide with a molecule comprising a terminal alkynyl group or a (hetero)cycloalkynyl group is one of the reactions that is known in the art as "click chemistry". In the case of a linker-conjugate comprising a terminal alkynyl group, said cycloaddition reaction needs to be performed in the presence of a suitable catalyst, preferably a Cu(I) catalyst. However, in a preferred embodiment, the linker-conjugate comprises a (hetero)cycloalkynyl group, more preferably a strained (hetero)cycloalkynyl group. When the (hetero)cycloalkynyl is a strained (hetero)cycloalkynyl group, the presence of a catalyst is not required, and said reaction may even occur spontaneously by a reaction called strain-promoted azide-alkyne cycloaddition (SPAAC). This is one of the reactions known in the art as "metal-free click chemistry".

Application

[0252] The invention further concerns a method for the treatment of a subject in need thereof, comprising the administration of the antibody-conjugate according to the invention as defined above. The subject in need thereof is typically a cancer patient. The use of antibody-conjugates, such as antibody-drug conjugates, is well-known in the field of cancer treatment, and the antibody-conjugates according to the invention are especially suited in this respect. The method as described is typically suited for the treatment of cancer. In the method according to this aspect, the antibody-

conjugate is typically administered in a therapeutically effective dose. The present aspect of the invention can also be worded as an antibody-conjugate according to the invention for use in the treatment of a subject in need thereof, preferably for the treatment of cancer. In other words, this aspect concerns the use of an antibody-conjugate according to the invention for the preparation of
5 a medicament or pharmaceutical composition for use in the treatment of a subject in need thereof, preferably for use in the treatment of cancer. In the present context, treatment of cancer is envisioned to encompass treating, imaging, diagnosing, preventing the proliferation of, containing and reducing tumours.

[0238] This aspect of the present invention may also be worded as a method for targeting PTK7-expressing cells, in particular PTK7-expressing tumour cells, comprising contacting the antibody-conjugate according to the invention with cells that may possibly be PTK7-expressing. The method according to this aspect is thus suitable to determine whether the cells are PTK7-expressing. These PTK7-expressing cells may be present in a subject, in which case the method comprises administering to a subject in need thereof the antibody-conjugate according to the invention. In a
15 preferred embodiment, the cells that may possibly be PTK7-expressing are PTK7-expressing cells. The targeting of PTK7-expressing cells preferably includes one or more of treating, imaging, diagnosing, preventing the proliferation of, containing and reducing PTK7-expressing cells, in particular PTK7-expressing tumour cells. The method according to this embodiment may be medical or non-medical. Non-medical methods according to the present aspect may be directed to
20 *in vitro* or *ex vivo* targeting PTK7-expressing cells, wherein the cells that may possibly be PTK7-expressing are present in a sample, e.g. taken from a patient. Such a non-medical method is typically used for the diagnosis of cancer, in particular PTK7-positive cancer.

[0253] In the context of the present invention, the subject may suffer from a disorder selected from colon cancer, lung cancer, breast cancer, ovarian cancer and esophageal cancer. Thus, the
25 treatment of a subject in need thereof preferably refers to the treatment of colon cancer, lung cancer, breast cancer, ovarian cancer and esophageal cancer.

[0254] The inventors have surprisingly found that the antibody-conjugates according to the invention are superior to conventional PTK7-targeting antibody-conjugates in terms of safety and/or efficacy, such that the therapeutic index of the antibody-conjugate according to the invention is
30 increased with respect to conventional PTK7-targeting antibody-conjugates.


Mode of conjugation

[0255] In the context of the present invention, the "mode of conjugation" refers to the process that is used to conjugate a payload D to an antibody AB, as well as to the structural features of the
35 resulting antibody-conjugate, in particular of the linker that connects the payload to the antibody, that are a direct consequence of the process of conjugation. Thus, in one embodiment, the mode of conjugation refers to a process for conjugation a payload to an antibody. In an alternative embodiment, the mode of conjugation refers to structural features of the linker and/or to the attachment point of the linker to the antibody that are a direct consequence of the process for
40 conjugation a payload to an antibody.

-59-

[0256] In a further aspect, the invention concerns the use of a mode of conjugation for increasing the therapeutic index of an antibody-conjugate in the treatment of PTK7-expressing tumours, wherein the mode of conjugation is being used to connect antibody AB with payload D via a linker L. The mode of conjugation mode of conjugation comprises:

- 5 (i) contacting an antibody comprising y core *N*-acetylglucosamine (GlcNAc) moieties, wherein $y = 1, 2, 3$ or 4 , with a compound of the formula $S(F)_x-P$ in the presence of a catalyst, wherein $S(F)_x$ is a sugar derivative comprising x reactive groups F capable of reacting with a reactive group Q , x is 1 or 2 and P is a nucleoside mono- or diphosphate, and wherein the catalyst is capable of transferring the $S(F)_x$ moiety to the core-GlcNAc moiety, to obtain a modified antibody according to Formula (26):



wherein

- 15 - AB is an antibody capable of targeting PTK7-expressing tumours;
 - Fuc is fucose;
 - w is 0 or 1 ; and
 (ii) reacting the modified antibody with a compound according to structure (2):



20 wherein:

- Q is a reactive moiety;
 - L is a linker that links Z to D ;
 - D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolbenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
 25 to obtain the antibody-conjugate according to structure (1)

[0257] Preferably, increasing the therapeutic index of an antibody-conjugate is selected from:

- (a) increasing the therapeutic efficacy of the antibody-conjugate; and/or
 30 (b) increasing the tolerability of the antibody-conjugate.

[0258] Increase in therapeutic efficacy of the antibody-conjugates according to the invention may take the form of a reduction in tumour size and/or a prolonged period of regression, when compared to conventional PTK7-targeting ADC. Increase in tolerability of the antibody-conjugates according to the invention may take the form of a reduction in signs of toxicity, compared to administration of
 35 a PTK7-targeting ADC made with a conventional technology. The reduction in signs may also be referred to as a reduction in symptoms or side-effects of cancer treatment, and may involve one or more clinical signs such as reduced reduction in body weight, reduced reduction in mobility, reduced reduction in food intake and/or one or more toxicity parameters, such as improved blood chemistry, hematology, and/or histopathology.

40

Examples**[0259]** General procedure for transient expression and purification of monoclonal antibodies:

Various IgGs (cofetuzumab, 4D5, 7C8 or B12) were transiently expressed in CHO K1 cells by Evitria (Zurich, Switzerland) at 1L, 100 mL, 100 mL and 5 L scale respectively. Other IgGs (12C6 and 12C6a) were transiently expressed in ExpiCHO-S(TM) cells (Gibco) at 500 mL scale. The non-fucosylated 12C6 was transiently expressed in CHO cells using the ProBioGen's GlymaxX® technology by Evitria. The supernatant was purified using a HiTrap MabSelect sure column. The supernatant was loaded onto the column followed by washing with at least 10 column volumes of 25 mM Tris pH 7.5, 150 mM NaCl (TBS). Retained protein was eluted with 0.1 M AcOH pH 2.7. The eluted product was immediately neutralized with 2.5 M Tris-HCl pH 8.8 and dialyzed against 20 mM histidine, 150 mM NaCl, pH 7.5. Next the IgG was concentrated (>20 mg/mL) using a Vivaspinn Turbo 15 ultrafiltration unit (Sartorius). The sequences of the IgGs are given here below:

[0260] cofetuzumab (I) light chain (SEQ ID No. 8):

EIVLTQSPATLSLSPGERATLSCRASESVDSYGKSFMHWYQQKPGQAPRLLIYRASNLESGIPAR
FSGSGSGTDFTLTISLEPEDFAVYYCQQSNEDPWTFGGGKLEIKRTVAAPSVFIFPPSDEQLK
SGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHK
VYACEVTHQGLSSPVTKSFNRGEC

[0261] cofetuzumab (I) heavy chain (SEQ ID No. 7):

QVQLVQSGPEVKKPGASVKVSKASGYTFTDYAVHWVRQAPGKRLEWIGVISTYNDYTYNNQD
FKGRVTMTRDTSASTAYMELSRLRSEDYAVYYCARGNSYFYALDYWGQGTSTVTVSSASTKGPS
VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTP
SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMIS
RTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK
EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWES
NGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG

[0262] 4D5 (II) light chain (SEQ ID No. 16):

EIVLTQSPDFQSVTPKEKVTITCRASQSIGSSHLHWYQQKPDQSPKLLIKYASQSFSGVPSRFSGS
GSGTDFTLTINSLEAEDAAAYCHQSSSLPITFGQGTREIKRTVAAPSVFIFPPSDEQLKSGTASV
VCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEV
THQGLSSPVTKSFNRGEC

[0263] 4D5 (II) heavy chain (SEQ ID No. 15):

QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAFHWVRQAPGKGLEWAVISYDGSIKYYADS
VKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARTYYFDYWGQGTSLTVTVSSASTKGPSVFPLA
PSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSL
GTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPE
VTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKC
KVSNNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQ
PENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG

[0264] 7C8 (III) light chain (SEQ ID No. 25):

EIVLTQSPATLSLSPGERATLSCRASQSVSIYLA~~WY~~QQKPGQAPRLLIYDASNRATGIPARFSGSG
SGTDFTLTISLEPEDFAVYYCQQRSNWPPFTFGPGTKVDIKRTVAAPSVFIFPPSDEQLKSGTAS
VVCLLNNFYPPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSSSTLTLSKADYEKHKVYACE
5 VTHQGLSSPVTKSFNRGEC

[0265] 7C8 (III) heavy chain (SEQ ID No. 26):

QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYGMHWVRQAPGKGLEWVAVIMDDGSNKYYVD
SVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARD~~DY~~YSGGSFN~~SY~~YGT~~DV~~WVGQGT~~TV~~TVS
SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT~~SV~~WNSGALTSGVHTFPAVLQSSGLY
10 SLSSV~~TV~~VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPP
KPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT~~TV~~L
HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYP
SDIAVEWESNGQPENNYKTTPVLDSDGSFFLYSKLTVDKSRWQQGNV~~F~~SCSV~~M~~HEALHNHYT
QKSLSLSPG

15 **[0266]** 12C6 (IV) light chain (SEQ ID No. 20):

EIVLTQSPGTLSPGERATLSCRASQSVSSSYLA~~WY~~QQKPGQAPRLLIYGA~~SS~~RATGIPDRFSG
SGSGTDFTLTISRLEPEDFAVYYCQ~~QY~~GSSPMYTFGQGT~~KL~~EIKRTVAAPSVFIFPPSDEQLKSGT
ASVVCLLNNFYPPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSSSTLTLSKADYEKHKVYA
CEVTHQGLSSPVTKSFNRGEC

20 **[0267]** 12C6 (IV) heavy chain (SEQ ID No. 19):

VQLVQSGGGLVHPGGSRLSCAGSGFTFSTYLMYWRQAPGK~~LE~~WVSAIGSGGDTYYADSVK
GRFTISRDN~~AK~~NSLYLQMNSLRAEDMAVYYCARG~~LY~~WGQGT~~LV~~TVSS
ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT~~SV~~WNSGALTSGVHTFPAVLQSSGLYS
LSSV~~TV~~VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPK
25 PKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT~~TV~~LH
QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPS
DIAVEWESNGQPENNYKTTPVLDSDGSFFLYSKLTVDKSRWQQGNV~~F~~SCSV~~M~~HEALHNHYTQ
KSLSLSPG

[0268] 12C6a (V) light chain (SEQ ID No. 22):

30 DIQMTQSPSSLSASVGDRVTITCRASQGISSWLA~~WY~~QQKPEKAPKSLIYAASSLQSGVPSRFSGS
GSGTDFTLTISLQPEDFATYYCQ~~QY~~NSYPYTFGQGT~~KL~~EIK
RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPPREAKVQWKVDNALQSGNSQESVTEQDSKDS
TYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

[0269] 12C6a (V) heavy chain: see 12C6 heavy chain (SEQ ID No. 19)

35 **[0270]** B12 (VI) light chain (SEQ ID No. 28):

EIVLTQSPGTLSPGERATFSCRSSHSIRSRRAWYQHKPGQAPRLVIHGVSNRASGISDRFSG
SGSGTDFTLTITRVEPEDFALYYCQVYGASSYTFGQGT~~KL~~ERKRTVAAPSVFIFPPSDEQLKSGT
ASVVCLLNNFYPPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSSSTLTLSKADYEKHKVYA
CEVTHQGLSSPVTKSFNRGEC

[0271] B12 (VI) heavy chain (SEQ ID No. 27):

QVQLVQSGAEVKKPGASVKVSCQASGYRFSNFIHWVRQAPGQRFEWMGWINPYNGNKEFSA
 KFQDRVTFADTSANTAYMELRSLRSADTAVYYCARVGPYSWDDSPQDNYYMDVWGKGTTVIV
 SSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL
 5 YSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFP
 PKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV
 LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFY
 PSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHY
 TQKSLSLSPGK

10 **[0272]** General procedure for analytical RP-UPLC (DTT treated samples): Prior to RP-UPLC analysis, IgG (10 μ L, 1 mg/mL in PBS pH 7.4) was added to 12.5 mM DTT, 100 mM TrisHCl pH 8.0 (40 μ L) and incubated for 15 minutes at 37 °C. The reaction was quenched by adding 49% acetonitrile, 49% water, 2% formic acid (50 μ L). RP-UPLC analysis was performed on a Waters Acquity UPLC-SQD. The sample (5 μ L) was injected with 0.4 mL/min onto Bioresolve RP
 15 mAb 2.1*150 mm 2.7 μ m (Waters) with a column temperature of 70 °C. A linear gradient was applied in 9 minutes from 30 to 54% acetonitrile in 0.1% TFA and water. Absorbance of eluted peaks was measured at 215 nm followed by automated integration (MassLynx, Waters) to determine reaction conversion.

[0273] General procedure for mass spectral analysis of (modified) monoclonal antibodies: Prior to
 20 mass spectral analysis, IgG was treated with IdeS, which allows analysis of the Fc/2 fragment. For analysis of both light and heavy chain, a solution of 20 μ g (modified) IgG was incubated for 5 minutes at 37 °C with 100 mM DTT in a total volume of 4 μ L. If present, azide-functionalities are reduced to amines under these conditions. For analysis of the Fc/2 fragment, a solution of 20 μ g (modified) IgG was incubated for 1 hour at 37 °C with IdeS/Fabricator™ (1.25 U/ μ L) in phosphate-
 25 buffered saline (PBS) pH 6.6 in a total volume of 10 μ L. Samples were diluted to 80 μ L followed by analysis electrospray ionization time-of-flight (ESI-TOF) on a JEOL AccuTOF. Deconvoluted spectra were obtained using Magtran software.

[0274] General procedure for enzymatic remodeling of IgG to mAb-(6-N₃-GalNAc)₂: IgG (15 mg/mL) was incubated with 1% w/w EndoSH (as described in PCT/EP2017/052792, see Examples
 30 1 – 3, and SEQ. ID No: 1, which is incorporated by reference herein), 3% w/w His-TnGalNAcT (as described in PCT/EP2016/059194, see Examples 3 and 4, and SEQ. ID No: 49, which is incorporated by reference herein), 0.01% AP (Roche) and UDP 6-N₃-GalNAc (compound **2d** in Figure 3, 25 eq compared to IgG) in 6 mM MnCl₂ and TBS for 16 hours at 30 °C. Next, the functionalized IgG was purified using a HiTrap MabSelect sure 5 mL column. After loading of the
 35 reaction mixture the column was washed with TBS+0.2% triton and TBS. The IgG was eluted with 0.1 M AcOH pH 2.7 and neutralized with 2.5 M Tris-HCl pH 8.8. After three times dialysis to 20 mM histidine, 150 mM NaCl pH 7.5, the IgG was concentrated to 15-20 mg/mL using a Vivaspin Turbo 15 ultrafiltration unit (Sartorius).

Preparation of azide-functionalized antibodies: Examples 1 – 7:*Example 1: Preparation of cofetuzumab-(6-N₃-GalNAc)₂ (I-N₃)*

- [0275]** According to the general procedure for enzymatic remodeling, cofetuzumab was converted to cofetuzumab-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24329 Da, approximately 90% of total Fc/2), corresponding to the expected product.

Example 2: Preparation of B12-(6-N₃-GalNAc)₂ (VI-N₃)

- [0276]** According to the general procedure for enzymatic remodeling, B12 was converted to B12-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24330 Da, approximately 90% of total Fc/2), corresponding to the expected product.

Example 3: Preparation of 4D5-(6-N₃-GalNAc)₂ (II-N₃)

- [0277]** According to the general procedure for enzymatic remodeling, 4D5 was converted to 4D5-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24332 Da, approximately 90% of total Fc/2), corresponding to the expected product and one minor Fc/2 product (observed mass 24186 Da, approximately 10% of total Fc/2), corresponding to the non-fucosylated product.

Example 4: Preparation of 7C8-(6-N₃-GalNAc)₂ (III-N₃)

- [0278]** According to the general procedure for enzymatic remodeling, 7C8 was converted to 7C8-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24332 Da, approximately 90% of total Fc/2), corresponding to the expected product and one minor Fc/2 product (observed mass 24186 Da, approximately 10% of total Fc/2), corresponding to the non-fucosylated product.

Example 5: Preparation of 12C6-(6-N₃-GalNAc)₂ (IV-N₃)

- [0279]** According to the general procedure for enzymatic remodeling, 12C6 was converted to 12C6-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24331 Da, approximately 90% of total Fc/2), corresponding to the expected product and one minor Fc/2 product (observed mass 24186 Da, approximately 10% of total Fc/2), corresponding to the non-fucosylated product.

Example 6: Preparation of 12C6a-(6-N₃-GalNAc)₂ (V-N₃)

- [0280]** According to the general procedure for enzymatic remodeling, 12C6a was converted to 12C6a-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24332 Da, approximately 90% of total Fc/2), corresponding to the


-64-

expected product and one minor Fc/2 product (observed mass 24186 Da, approximately 10% of total Fc/2), corresponding to the non-fucosylated product.

Example 7: Preparation of aFuc-12C6-(6-N₃-GalNAc)₂ (aFuc-IV-N₃)

- 5 **[0281]** According to the general procedure for enzymatic remodeling, aFuc-12C6 was converted to aFuc-12C6-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24187 Da, approximately 95% of total Fc/2), corresponding to the expected product.

10 Examples 8-12: Synthesis of linker conjugates **3**, **4**, **5b** and **9**



*Example 8: Preparation of compound **11***


- [0282]** Compound **10** (163 mg, 240 μmol) was added to a mixture of exatecan mesylate (125 mg, 235 μmol) and DIPEA (61 mg, 82 μL, 0.47 mmol) in dry DMF (0.9 mL). After 20 h, the reaction mixture was diluted to 9 mL DCM and purified by gradient column chromatography (0 → 40% MeOH/DCM) to afford **11** (155 mg, 159 μmol, 68%). LCMS (ESI+) calculated for C₅₅H₅₄FN₆O₁₀⁺ (M+H)⁺ 977.39, found 977.72. In addition to **11**, free base of exatecan (82.4 mg, 189 μmol, 20%) was recovered. LCMS (ESI+) calculated for C₂₄H₂₃FN₃O₄⁺ (M+H)⁺ 436.46, found 436.54.

20 *Example 9. Preparation of compound **3***

[0283] The synthesis of BCN-HS-(va-PABC-Ex)₂ (**3**) is also described in PCT/EP2021/075401 (example 4), incorporated herein. To a solution of compound **11** (155 mg, 159 μmol) in DMF (1.6

-65-

mL) were added Et₃N (73 mg, 101 μ L, 0.72 mmol) and a solution of compound **12** (65 mg, 72 μ mol) in DMF (1.4 mL). The reaction mixture was stirred for 18 h, diluted with DCM (20 mL) and purified by gradient column chromatography (0 $\rightarrow$ 40% MeOH/DCM) to afford **3** as a pale-yellow solid (94 mg, 44 μ mol, 28%). LCMS (ESI⁺) calculated for C₁₀₂H₁₁₈F₂N₁₆O₂₉S₂²⁺ (M/2+H)⁺ 1066.88, found 1067.12.




Example 10: Preparation of linker-conjugate **4**

[0284] The synthesis of BCN-HS-(vc-PABC-MMAE)₂ (**4**) is also described in PCT/EP2017/052791 (example 25), incorporated herein. To a solution of **14** (27 mg, 33 μ mol) in DMF (400 μ L) were added triethylamine (22 μ L; 16 mg; 158 μ mol) and a solution of vc-PABC-MMAE.TFA **13** (96 mg; 78 μ mol) in DMF (1.0 mL). The mixture was left standing for 19 h and 2,2'-(ethylenedioxy)bis(ethylamine) (37 μ L, 38 mg, 253 μ mol) was added. After 2 h, the reaction mixture was diluted with DMF (100 μ L) and purified by RP HPLC (C18, 30% $\rightarrow$ 90% MeCN (1% AcOH) in water (1% AcOH). The desired product **4** was obtained as a colourless film (41 mg, 14.7 μ mol, 45%). LCMS (ESI⁺) calculated for C₁₃₈H₂₁₉N₂₃O₃₅S₂²⁺ (M+2H)⁺ 1395.79 found 1396.31.

Example 11: Synthesis of linker conjugate **5b**

[0285] The synthesis of BCN-HS-vc-PABC-CM (**5b**) was prepared according to the procedures described in WO2019110725, incorporated herein. To a solution of **15** (9.0 mg, 4.88 μ mol, compound **138** in WO2019110725) in DMF (445 μ L) were added a solution of **16** (12.8 mg, 24.4 μ mol, compound **3** in WO2019110725) in DMF (45.8 μ L) and a 50% v/v Et₃N solution in DMF (8.98 μ L, 32.2 μ mol). The mixture was left standing for 4 h and then diluted with DCM to a total volume of 4.5 mL. This mixture was purified by silica gel chromatography (0 $\rightarrow$ 20% MeOH in DCM). The desired product **5b** was obtained as a colorless oil (5.0 mg, 2.24 μ mole, 46%). LCMS (ESI⁺) calculated for C₉₆H₁₃₆IN₉O₃₅S₄²⁺ (M+2H)⁺ 1115.35, found 1115.40.

-66-



Example 12. Preparation of compound 9

- [0286]** A solution of BCN-HS-PEG₂-b-(Glu(OFm)-OH)₂ (**8**, 12.1 mg, 10 µmol, 1.0 eq) dissolved in anhydrous DMF (180 µL) was added to a solution of NH₂-Val-Ala-PABC-exatecan (**5b**, Fmoc-deprotected **5**, 19 mg, 25 µmol, 2.5 eq) in anhydrous DCM (180 µL), DIPEA (11 µL, 63 µmol, 6.2 eq) and HATU (8.9 mg, 23 µmol, 2.3 eq). After stirring for 2 h at room temperature, the reaction mixture was further diluted with DCM (800 µL) and purified by flash column chromatography over silica gel (0% → 20% MeOH in DCM) to give the product as a clear oil (difficult to determine yield due to presence of DMF). LCMS (ESI+) calculated for C₁₄₀H₁₅₀F₂N₁₇O₃₃S⁺ (M/2+H⁺) 1334.01, found 1334.79.
- [0287]** This compound was dissolved in DMF (300 µL) and triethylamine (21 µL, 150 µmol, 15 eq) was added. After 17 h at room temperature, the reaction mixture was diluted with DCM (700 µL) and purified by flash column chromatography over silica gel (0% → 45% MeOH in DCM) to give compound **9** in 44% yield as a yellow solid (10.2 mg, 4.4 µL). LCMS (ESI+) calculated for C₁₁₂H₁₃₀F₂N₁₇O₃₃S⁺ (M/2+H⁺) 1156.2, found 1156.74.

15

Examples 13 – 24: Conjugation of linker-payloads to (modified) monoclonal antibodies*Example 13: Production of cofetuzumab-Aur0101*

- [0288]** Cofetuzumab (14.4 mg, 4.34 mg/ml in 20 mM histidine, 6% sucrose) was charged with 5% pH adjust buffer (0.5 M Tris, 25 mM EDTA, pH 8.5) and 2.3 molar equivalents TCEP (1 mM in water) and incubated at room temperature for 2 hours. The reduced antibody was charged with DMA to give 10% v/v and 6 molar equivalents Auristatin0101 (2 mM in DMA) and incubated at room temperature for 1 hour. The reaction was quenched using 6 molar equivalents of NAC (100 mM in water). The conjugate was purified by preparative discontinuous diafiltration in Vivaspins into 20 mM histidine / 6% sucrose. 0.04% Tween-20 was added before filter sterilization. The average DAR was measured and was 4.3.

25

*Example 14: Conjugation of cofetuzumab(6-N₃-GalNAc)₂ with BCN-HS-PEG₂-HS-(va-PAB-Ex)₂ **3** to obtain conjugate cofetuzumab -3*

- [0289]** To a solution of cofetuzumab(6-N₃-GalNAc)₂ (1664 µL, 32.0 mg, 19.37 mg/ml in TBS pH 7.5) was added sodium deoxycholate (322 µL, 110 mM) and BCN-HS-PEG₂-HS-(va-PAB-Ex)₂ **3** (86 µL, 10 mM solution in DMF) and propylene glycol (881 µL). The reaction was incubated overnight at rt. Next the conjugate was purified on a HiLoad 16/600 Superdex200 PG column (Cytiva) on an AKTA Pure (Cytiva). To remove the excess of free payload, 25 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated overnight. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 µm, Corning). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26461 Da, approximately 90% of total

35

40

-68-

Fc/2), corresponding to the conjugate **cofetuzumab-3**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.86.

*Example 15: Conjugation of cofetuzumab(6-N₃-GalNAc)₂ with BCN-HS-(vc-PABC-MMAE)₂ **4** to obtain conjugate **cofetuzumab-4***

5 **[0290]** To a solution of cofetuzumab(6-N₃-GalNAc)₂ (463 µL, 9.0 mg, 19.37 mg/ml in TBS pH 7.5) was added sodium deoxycholate (60 µL, 110 mM) and BCN-HS-(vc-PABC-MMAE)₂ **4** (1.8 µL, 100 mM solution in DMF) and DMF (58 µL). The reaction was incubated overnight at rt. Next the conjugate was purified on a HiLoad 10/300 Superdex200 PG column (Cytiva) on an AKTA Pure
10 (Cytiva). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 27119 Da, approximately 90% of total Fc/2), corresponding to the conjugate **cofetuzumab-4**. RP-UPLC analysis of the
15 sample under reducing conditions showed an average DAR of 3.90.

*Example 16: Conjugation of cofetuzumab(6-N₃-GalNAc)₂ with BCN-HS-vc-PABC-calicheamicin **5b** to obtain conjugate **cofetuzumab-5b***

[0291] To a solution of cofetuzumab(6-N₃-GalNAc)₂ (730 µL, 14.0 mg, 19.37 mg/ml in TBS pH 7.5) was added sodium deoxycholate (94 µL, 110 mM) and BCN-HS-vc-PABC-calicheamicin **5b** (7 µL,
20 40 mM solution in DMF) and DMF (87 µL). The reaction was incubated overnight at rt. Next the conjugate was purified on a HiLoad 10/300 Superdex200 PG column (Cytiva) on an AKTA Pure (Cytiva). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0,
25 and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26714 Da, approximately 90% of total Fc/2), corresponding to the conjugate **cofetuzumab-5b**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 1.93.

30 *Example 17: Conjugation of B12(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** to obtain conjugate **B12-3***

[0292] To a solution of B12(6-N₃-GalNAc)₂ (6.33 mL, 150.0 mg, 23.71 mg/ml in TBS pH 7.5) was added BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** (495 µL, 10 mM solution in DMF) and propylene glycol (7.0 mL). The reaction was incubated overnight at rt. Next the conjugate was purified on a HiLoad
35 26/600 Superdex200 PG column (Cytiva) on an AKTA Pure (Cytiva). To remove the excess of free payload, 150 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated overnight. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 µm, Corning). Subsequently the solution was dialysed to 20 mM histidine, 6% sucrose buffer pH 6.0 for two hours at rt and overnight at 4°C. The solution was concentrated using a
40 Vivaspin Turbo 4 10 kDa MWCO ultrafiltration unit (Sartorius) and 0.04% Tween-20 was added

before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26462 Da, approximately 90% of total Fc/2), corresponding to the conjugate **B12-3**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.75.

5

*Example 18: Conjugation of 4D5(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** to obtain conjugate **4D5-3***

[0293] To a solution of 4D5(6-N₃-GalNAc)₂ (424 µL, 8.8 mg, 20.74 mg/ml in TBS pH 7.5) was added sodium deoxycholate (88 µL, 110 mM) and BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** (23.4 µL, 10 mM solution in DMF) and propylene glycol (240 µL). The reaction was incubated overnight at rt. Next the conjugate was purified using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with PBS pH 7.4 on an AKTA Pure (Cytiva). To remove the excess of free payload, 12 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated overnight. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 µm, Corning). Subsequently the solution was dialyzed three times to 20 mM histidine, 6% sucrose buffer pH 6.0, and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26465 Da, approximately 90% of total Fc/2), corresponding to the conjugate **4D5-3**. And one minor Fc/2 product (observed mass 26318 Da, approximately 10% of total Fc/2), corresponding to the non-fucosylated product. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.91.

*Example 19: Conjugation of 7C8(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** to obtain conjugate **7C8-3***

[0294] To a solution of 7C8(6-N₃-GalNAc)₂ (313 µL, 6.9 mg, 22.14 mg/ml in TBS pH 7.5) was added BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** (23.1 µL, 10 mM solution in DMF) and propylene glycol (254 µL). The reaction was incubated overnight at rt. Next the conjugate was purified on a Superdex200 10/300 increase column (Cytiva) on an AKTA Pure (Cytiva). To remove the excess of free payload, 11 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated overnight. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 µm, Corning). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26466 Da, approximately 90% of total Fc/2), corresponding to the conjugate **7C8-3**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.95.

*Example 20: Conjugation of 12C6(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** to obtain conjugate **12C6-3***

[0295] To a solution of 12C6(6-N₃-GalNAc)₂ (481 µL, 10.1 mg, 21.01 mg/ml in TBS pH 7.5) was

40

-70-

added BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** (33.7 μ L, 10 mM solution in DMF) and propylene glycol (370 μ L). The reaction was incubated overnight at rt. Next the conjugate was purified using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with PBS pH 7.4 on an AKTA Pure (Cytiva). To remove the excess of free payload, 12 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated overnight. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 μ m, Corning). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26466 Da, approximately 95% of total Fc/2), corresponding to the conjugate **12C6-3**. And one minor Fc/2 product (observed mass 26318 Da, approximately 5% of total Fc/2), corresponding to the non-fucosylated product. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.85.

*Example 21: Conjugation of 12C6a(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** to obtain conjugate **12C6a-3***

[0296] To a solution of 12C6a(6-N₃-GalNAc)₂ (484 μ L, 10.9 mg, 22.53 mg/ml in TBS pH 7.5) was added BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** (36.3 μ L, 10 mM solution in DMF) and propylene glycol (400 μ L). The reaction was incubated overnight at rt. Next the conjugate was purified on a Superdex200 10/300 increase column (Cytiva) on an AKTA Pure (Cytiva). To remove the excess of free payload, 12 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated overnight. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 μ m, Corning). Subsequently the solution was dialyzed three times to 20 mM histidine, 6% sucrose buffer pH 6.0, and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26466 Da, approximately 95% of total Fc/2), corresponding to the conjugate **12C6a-3**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.90.

*Example 22: Conjugation of 12C6(6-N₃-GalNAc)₂ with BCN-HS-PEG2-(eva-PAB-Ex)₂ **9** to obtain conjugate **12C6-9***

[0297] To a solution of 12C6(6-N₃-GalNAc)₂ (340 μ L, 10 mg, 29.45 mg/ml in TBS pH 7.5) was added sodium deoxycholate (100 μ L, 110 mM) and BCN-HS-PEG2-(eva-PAB-Ex)₂ **9** (20 μ L, 10 mM solution in DMF) and propylene glycol (280 μ L). The reaction was incubated overnight at rt. Next the conjugate was purified using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.2M NaOH and equilibrated with PBS pH 7.4 on an AKTA Pure (Cytiva). To remove the excess of free payload, 12 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated overnight. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 μ m, Corning). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.2M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the

-71-

sample after IdeS treatment showed one major Fc/2 product (observed mass 26466 Da, approximately 95% of total Fc/2), corresponding to the conjugate **12C6-9**. And one minor Fc/2 product (observed mass 26643 Da, approximately 5% of total Fc/2), corresponding to the non-fucosylated product. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.84.

*Example 23: Conjugation of aFuc-12C6(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** to obtain conjugate aFuc-12C6-3*

[0298] To a solution of aFuc-12C6(6-N₃-GalNAc)₂ (549 µL, 13 mg, 23.69 mg/ml in TBS pH 7.5) was added sodium deoxycholate (130 µL, 110 mM) and BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** (34.7 µL, 10 mM solution in DMF) and propylene glycol (355 µL). The reaction was incubated overnight at rt. Next the conjugate was purified using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.2M NaOH and equilibrated with PBS pH 7.4 on an AKTA Pure (Cytiva). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.2M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26318 Da, approximately 95% of total Fc/2), corresponding to the conjugate **aFuc-12C6-3**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.74.

*Example 24: Conjugation of aFuc-12C6(6-N₃-GalNAc)₂ with BCN-HS-PEG2-(eva-PAB-Ex)₂ **9** to obtain conjugate aFuc-12C6-9*

[0299] To a solution of aFuc-12C6(6-N₃-GalNAc)₂ (59 µL, 1.4 mg, 23.69 mg/ml in TBS pH 7.5) was added sodium deoxycholate (14 µL, 110 mM) and BCN-HS-PEG2-(eva-PAB-Ex)₂ **9** (2.8 µL, 10 mM solution in DMF) and propylene glycol (39 µL). The reaction was incubated overnight at rt. Next the conjugate was purified using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.2M NaOH and equilibrated with PBS pH 7.4 on an AKTA Pure (Cytiva). RP-UPLC analysis of the conjugate **aFuc-12C6-9** sample under reducing conditions showed an average DAR of 3.81.

Examples 25 – 28: In vitro assays

Example 25. hPTK7, cPTK7 and rPTK7 binding assay to mAbs and ADCs using ELISA

[0300] Nickel NTA plates (Pierce™ Nickel coated plated, ThermoScientific™) were washed three times prior to use. Human PTK7 (CCK4 protein, ECD, His Tag, Sino Biological), cynomolgus PTK7 (CCK4 protein, ECD, His Tag, Sino Biological) and Rat PTK7 (His Tag, Acro Biosystems) were dissolved at a concentration of 0.05 µg/mL in 0.1% BSA in PBS (PBA). 100 µL was added to each well and incubated while shaking for 1 hour at room temperature. After removal, the plate was washed 3x with 0.05% Tween-20 in PBS (washing buffer). ADCs were diluted in 0.1% PBA to a final concentration of 8 µg/mL and 100 µL was added to each well (in triplo). ADCs were incubated for 1h at room temperature. Prior to the addition of 100 µL 1:1000 dilution of secondary antibody

-72-

(Goat anti-human IgG, HRP conjugate, Invitrogen) the plate was washed 3x with washing buffer. The plate was incubated again for 1h at room temperature and subsequently washed 3x with washing buffer and 3x with PBS. Finally, 100 μ L TMB ELISA substrate (1 Step™ Turbo TMB ELISA substrate, ThermoScientific™) was added and incubated for 1 minute. The absorbance of the colorimetric signal was measured with Infinite® M1000 (Tecan) at 652 nm. Data was plotted corrected for the background (see Figure 8A, 8B and Figure 11).

Example 26. Human and mouse serum stability

[0301] Stability of ADCs in mice and human plasma was tested. Prior to the assay, the plasma was depleted from all IgG using CaptivA® Protein A agarose (1 mL agarose/mL serum). ADCs were added to the depleted human/mouse serum to a final concentration of 0.1 mg/mL followed by incubation at 37°C. At each time point, 0.5 mL was snap frozen and stored at -80°C until further analysis. To isolate the ADCs after incubation, 20 μ L CaptivA® Protein A agarose resin was added to the samples and incubated for 1 hour at room temperature. The resin was washed three times with PBS and subsequently 0.1 M Glycine-HCl pH 2.7 (0.4 mL) was added to elute the ADCs. After elution, the samples were immediately neutralized with 1.0 M Tris pH 8.0 (0.1 mL). The samples were spin-filtered against PBS for three times using Amicon Ultra spin-filter 0.5 mL MWCO 10 kDa (Merck Millipore) and the volume was reduced to 40 μ L, yielding a final ADC concentration of approximately 1 mg/mL. Samples were analyzed on RP-UPLC (DTT reduced) to determine the DAR, tables below.

[0302] Table 1. Human plasma:

	DAR t=0	DAR t=3	DAR t=7	DAR t=14
cofetuzumab-Aur0101	4.21*	3.61	- <i>t</i>	2.67 (63%)
cofetuzumab-3	3.91	3.88	3.80	3.80 (97%)
cofetuzumab-4	3.93	3.91	3.90	3.88 (99%)
12C6-3	3.79*	3.76	3.69	3.62 (96%)
12C6-9	3.86	3.81	3.73	3.67 (95%)
aFuc-12C6-3	3.73	3.73	3.65	3.61 (97%)
aFuc-12C6-9	3.76	3.74	3.68	3.62 (96%)

[0303] Table 2. Mice plasma:

	DAR t=0	DAR t=1	DAR t=3	DAR t=7
cofetuzumab-Aur0101	4.21*	3.64	3.26	2.46 (58%)
cofetuzumab-3	3.90	3.89	3.88	3.86 (99%)
cofetuzumab-4	3.90	3.78	3.81	3.57 (92%)
12C6-3	3.79	3.73	3.74	3.70 (98%)
12C6-9	3.85	3.85	3.85	3.82 (99%)
aFuc-12C6-3	3.73	3.74	3.68	3.65 (98%)

-73-

aFuc-12C6-9	3.77	-†	3.75	3.74 (99%)
-------------	------	----	------	------------

-† signal too low

* Direct measurement (not through isolation from plasma with protA agarose resin).

Between parentheses is % of t=0

5 *Example 27: In vitro cytotoxicity*

- [0304]** A431 (PTK7+, ATCC CRL-1555) cells were plated in 96-well plates (10000 cells in 150 μ L/well) in DMEM (ATCC) supplemented with 10% fetal bovine serum (FBS) (Invitrogen) and incubated overnight in a humidified atmosphere at 37°C and 5% CO₂. ADCs were added in triplo in a square root of 10 dilution series to obtain a final concentration ranging from 100nM – 0.01nM.
- 10 The cells were incubated for 5 days in a humidified atmosphere at 37°C and 5% CO₂. The culture medium was replaced by 0.01 mg/mL resazurin (Sigma Aldrich) in DMEM supplemented with 10% FBS (200 μ L/well). After approximately 4 hours in a humidified atmosphere at 37°C and 5% CO₂ the fluorescence was detected with a fluorescence plate reader (Infinite® M1000 Tecan) at 560 nm excitation and 590 nm emission. The relative fluorescent units (RFU) were normalized to cell
- 15 viability percentage by setting wells without cells at 0% viability and wells with untreated cells at 100% viability (see Figure 9). IC₅₀ values for ADCs on A431 cells were calculated by non-linear regression using Graphpad prism software and are shown in the table below:

[0305] Table 3. IC₅₀ values for several ADCs and mAbs for A431 cells

	IC ₅₀ (nM)
cofetuzumab-4	3.9
cofetuzumab-5b	0.10
B12-3	>100

20

Example 28. Binding affinity of 12C6-3 and 12C6 naked mAb with FcγRs using Biacore

- [0306]** Binding of test antibodies to high and low affinity human Fc gamma receptors was assessed by single cycle analysis using a Biacore T200 (serial no. 1909913) instrument running Biacore T200 Evaluation Software (Cytiva, Uppsala, Sweden) running at a flow rate of 30 μ L/min. The human Fc
- 25 receptors, FcγRI, FcγRIIA (both 167R and 167H polymorphisms), FcγRIIB, FcγRIIIA (both 176F and 176V polymorphisms) and FcγRIIIB were obtained from Sino Biological (Beijing, China). FcγR were captured on a CM5 sensor chip pre-coupled using a His capture kit (Cytiva, Uppsala, Sweden) using standard amine chemistry.
- [0307]** At the start of each cycle His-tagged Fc gamma receptors (ligand) diluted in HBS-P+ (Cytiva, Uppsala, Sweden) were loaded to a specific RU level. A five point, three-fold dilution range of test antibody (analyte) without regeneration between each concentration was used for each
- 30 receptor tested. Test antibodies were passed over the chip in increasing concentrations at 30 μ L/min followed by a single dissociation step. Following dissociation, the chip was regenerated with injection of Glycine pH 1.5. The signal from the reference channel Fc1 (blank) was subtracted from
- 35 that of the Fc loaded with receptor to correct for differences in non-specific binding to the reference

-74-

surface. Sensorgrams were analysed for 1:1 kinetics for the high affinity Fc gamma receptor hFcγRI and by steady state binding for the low affinity Fc gamma receptors.

- [0308]** Table 4. Overview table of the binding of test antibodies and ADC to different human Fcγ receptors. Relative scale of binding (KD) is indicated: +++++, 10⁻⁹ to 10⁻¹⁰ M range; +++++, 10⁻⁸ M range; +++, 10⁻⁷ M range; ++, 10⁻⁶ M, +, 10⁻⁵ M range as determined by Biacore.

	hCD16A		hCD16B	hCD32A		hCD32B	hCD64
	FcγRIIIA _{176F}	FcγRIIIA _{176V}	FcγRIIIB	FcγRIIA _{167R}	FcγRIIA _{167H}	FcγRIIB	FcγRI
Trastuzumab	+++	+++	++	++	++	+	+++++
12C6	++	+++	++	++	++	++	+++++
12C6-3	-	-	+/-	-	+/-	+/-	+++

Examples 29 – 34: CDX evaluation and in vivo study

10

Example 29: IHC staining of NCI-H446 for PTK7 (performed at Crown Bioscience Taicang, China)

- [0225]** Formalin-fixed paraffin-embedded (FFPE) blocks were sectioned at 4 μm thickness on a manual rotary microtome onto a glass slide. The slides were baked at 60°C for 30 min. A Bond RX autostainer (Leica) was used for staining with the following procedure: the slides were first treated with dewax solution (Leica) for 0.5 min at 72 °C, then at rt and moved to alcohol, three times at rt. Next the slide was washed with Bond Wash buffer (Leica) for three times. Epitope retrieval was performed in Bond ER2 solution (Leica) for 20min at 100°C after a fast rinse in the same solution. The slide was subsequently washed shortly for four times in Bond wash buffer and subsequently for 3 minutes at rt. Peroxide block was performed for 10 minutes at rt after which it was washed again for three times. Subsequently, the primary antibody rabbit anti-human PTK7 (PA5-82070, Thermo Fisher, 0.25 μg/mL) was added and incubated for 1h at rt. Followed by one quick wash and three times for 2 minutes in Bond wash buffer. The secondary antibody, goat anti-rabbit-HRP (Leica, ready to use), was added and incubated for 20 minutes at rt, followed by one quick wash and three times for 2 minutes in Bond wash buffer. The slides were rinsed with deionized water before the incubation with 3,3'-diaminobenzidine (DAB) Refine for 5 minutes at rt. After washing with deionized water for three times, the slides were treated with hematoxylin as counterstain for 10 minutes. Lastly all slides were washed with deionized water and bond wash buffer.

- [0226]** All stained sections were scanned with NanoZoomer-HT 2.0 Image system for 40x magnification. All the images of non-necrotic tumor regions were analyzed with HALO™ platform. The intensity of membrane positive staining was scored at four levels, 0 (negative), 1+ (weak staining), 2+ (medium staining), 3+ (strong staining). The percentages of cells at different intensity levels were evaluated with H-score. H-Score = (% at 0)×0 + (% at 1)×1 + (% at 2)×2 + (% at 3)×3 (H-Score range is 0 to 300). See Table 4 for the H-score of the NCI-H446 lung cancer xenograft cell line.

35

[0227] Table 4. IHC score NCI-H446

Total cells	Positive cells (%)	Weak positive cells (%)	Moderate positive cells (%)	Strong positive cells (%)	Negative cells (%)	H-Score
370387	96.97	10.55	74.04	12.37	3.03	195.75

Example 30: In vivo efficacy study (performed at Crown Bioscience, Zhongshan, China)

- 5 [0228] NCI-H446, lung cancer xenograft model cell line was maintained in vitro using RPMI-1640 medium supplemented with 10% FBS in humidified cell culture incubator at 37°C with standard 5% CO₂ specs. The cells in an exponential growth phase were harvested and counted for tumor inoculation.
- 10 [0229] BALB/c nude mice (female) of 7–9 weeks old received a subcutaneous injection in the right front flank region with 5×10^6 tumor cells in 0.1 ml of PBS mixed with Matrigel (1:1) for tumor development. When tumors reached an average size of 100-200 mm³, randomization was performed into 10 groups of 8 mice and treatment began. Randomization will be performed based on “Matched distribution” method (Study Director™ software, version 3.1.399.19). The date of randomization will be denoted as day 0.
- 15 [0230] Electronic caliper measurement were performed 2 times a week. Daily observations for clinical signs, food and water consumption, behavioral changes, animals were weighed 2 times per week. The endpoint of the experiment is a tumor volume of 3,000 mm³, body weight loss over 20% or 28 days, whichever comes first.
- 20 [0231] Randomization: The randomization was performed when the mean tumor size reached approximately 100-200 mm³. Totally 80 mice were enrolled in NCI-H446 model study and randomly allocated to 10 groups, with 8 mice per group. Randomization will be performed based on “Matched distribution” method (Study Director™ software, version 3.1.399.19). The date of randomization will be denoted as day 0.
- 25 [0232] Test article administration: The test article administration was performed via intravenous injection through tail vein, and the dosing volume was 10 mL/kg. Treatment was initiated on the same day of randomization. Dosing was conducted in a Laminar Flow Cabinet.
- 30 [0233] Observation and data collection. After tumor cells inoculation, the animals are checked daily for morbidity and mortality. At the time of routine monitoring, the animals are checked for any adverse effects of tumor growth and treatments on normal behavior such as mobility, visual estimation of food and water consumption, body weight gain/loss, eye/hair matting and any other abnormal effects. Tumor volumes are measured every 3-4 days in two dimensions using an caliper, and the volume data are expressed in mm³ using the formula: $V = (L \times W \times W)/2$, where V is tumor volume, L is tumor length (the longest tumor dimension) and W is tumor width (the longest tumor dimension perpendicular to L). Dosing as well as tumor and body weight measurements will be
- 35 conducted in a Laminar Flow Cabinet.

[0234] Table 5. Overview of test items and dose levels included in in vivo study

Test item	Dose levels (mg/kg)	Group (N=8)	Administration	Dosing Schedule
vehicle	–	1	IV	Single dose
cofetuzumab-Aur0101	2.33	2	IV	Single dose
cofetuzumab-3	2.59	3	IV	Single dose
	5.18	4	IV	Single dose
	10.36	5	IV	Single dose
cofetuzumab-4	1.03	6	IV	Single dose
	2.56	7	IV	Single dose
cofetuzumab-5b	2.59	8	IV	Single dose
	5.18	9	IV	Single dose
B12-3	10.84	10	IV	Single dose

[0235] Tumor volume over time and mice body weight in the efficacy study with the test items above are depicted in Figures 10A – 10D.

5

Example 31: In vivo efficacy study

[0309] NCI-H446, lung cancer xenograft model cell line was maintained in vitro using RPMI-1640 medium supplemented with 10% FBS in humidified cell culture incubator at 37°C with standard 5% CO₂ specs. The cells in an exponential growth phase were harvested and counted for tumor inoculation.

10

[0310] BALB/c nude mice (female) of 7–9 weeks old received a subcutaneous injection in the right front flank region with 5×10^6 tumor cells in 0.1 ml of PBS mixed with Matrigel (1:1) for tumor development. When tumors reached an average size of 100-200 mm³, randomization was performed into 6 groups of 8 mice and treatment began. Randomization will be performed based on “Matched distribution” method (Study Director™ software, version 3.1.399.19). The date of randomization will be denoted as day 0.

15

[0311] Electronic caliper measurement were performed 2 times a week. Daily observations for clinical signs, food and water consumption, behavioral changes, animals were weighed 2 times per week. The endpoint of the experiment is a tumor volume of 3,000 mm³, body weight loss over 20% or 77 days, whichever comes first.

20

[0312] Randomization: The randomization was performed when the mean tumor size reached approximately 136 mm³. Totally 48 mice were enrolled in NCI-H446 model study and randomly allocated to 6 groups, with 8 mice per group. Randomization will be performed based on “Matched distribution” method (Study Director™ software, version 3.1.399.19). The date of randomization will be denoted as day 0.

25

[0313] Test article administration: The test article administration was performed via intravenous injection through tail vein, and the dosing volume was 10 mL/kg. Treatment was initiated on the same day of randomization. Dosing was conducted in a Laminar Flow Cabinet.

[0314] Observation and data collection: After tumor cells inoculation, the animals are checked daily for morbidity and mortality. At the time of routine monitoring, the animals are checked for any adverse effects of tumor growth and treatments on normal behavior such as mobility, visual estimation of food and water consumption, body weight gain/loss, eye/hair matting and any other abnormal effects. Tumor volumes are measured every 3-4 days in two dimensions using an caliper, and the volume data are expressed in mm³ using the formula: $V = (L \times W \times W)/2$, where V is tumor volume, L is tumor length (the longest tumor dimension) and W is tumor width (the longest tumor dimension perpendicular to L). Dosing as well as tumor and body weight measurements will be conducted in a Laminar Flow Cabinet.

10

[0315] Table 6. Overview of test items and dose levels included in in vivo study

Test item	Dose levels (mg/kg)	Group (N=8)	Administration	Dosing Schedule
vehicle	—	1	IV	Single dose
cofetuzumab-3	5	2	IV	Single dose
12C6a-3	5	3	IV	Single dose
12C6-3	5	4	IV	Single dose
7C8-3	5	5	IV	Single dose
4D5-3	5	6	IV	Single dose

[0316] Tumor volume over time and mice body weight in the efficacy study and the corresponding Kaplan-Meier plot with the test items above are depicted in Figures 12A, 12B and 12C.

15

Example 32: In vivo efficacy study

[0317] BALB/c nude mice (female) of 7–9 weeks old received a tumor fragment of BR1282, a breast cancer patient derived xenograft model cell line. The tumor fragment (2-3 mm in diameter) was inoculated subcutaneously the right upper flank region. When tumors reached an average size of 100-200 mm³, randomization was performed into 9 groups of 8 mice and treatment began. Randomization will be performed based on “Matched distribution” method (Study Director™ software, version 3.1.399.19). The date of randomization will be denoted as day 0.

[0318] Electronic caliper measurement were performed 2 times a week. Daily observations for clinical signs, food and water consumption, behavioral changes, animals were weighed 2 times per week. The endpoint of the experiment is a tumor volume of 3,000 mm³, body weight loss over 20% or 39 days, whichever comes first.

[0319] Randomization: The randomization was performed when the mean tumor size reached approximately 135 mm³. Totally 72 mice were enrolled in BR1282 model study and randomly allocated to 9 groups, with 8 mice per group. Randomization will be performed based on “Matched distribution” method (Study Director™ software, version 3.1.399.19). The date of randomization will be denoted as day 0.

30

[0320] Test article administration: The test article administration was performed via intravenous injection through tail vein, and the dosing volume was 10 mL/kg. Treatment was initiated on the same day of randomization. Dosing was conducted in a Laminar Flow Cabinet.

- [0321]** Observation and data collection: After tumor cells inoculation, the animals are checked daily for morbidity and mortality. At the time of routine monitoring, the animals are checked for any adverse effects of tumor growth and treatments on normal behavior such as mobility, visual estimation of food and water consumption, body weight gain/loss, eye/hair matting and any other abnormal effects. Tumor volumes are measured every 3-4 days in two dimensions using a caliper, and the volume data are expressed in mm³ using the formula: $V = (L \times W \times W)/2$, where V is tumor volume, L is tumor length (the longest tumor dimension) and W is tumor width (the longest tumor dimension perpendicular to L). Dosing as well as tumor and body weight measurements will be conducted in a Laminar Flow Cabinet.

[0322] Table 7. Overview of test items and dose levels included in in vivo study

Test item	Dose levels (mg/kg)	Group (N=8)	Administration	Dosing Schedule
vehicle	—	1	IV	Single dose
cofetuzumab-Aur0101	4	2	IV	Single dose
cofetuzumab-3	4	3	IV	Single dose
	8	4	IV	Single dose
12C6a-3	8	5	IV	Single dose
12C6-3	4	6	IV	Single dose
	8	7	IV	Single dose
7C8-3	4	8	IV	Single dose
4D5	4	9	IV	Single dose

- [0323]** Tumor volume over time and mice body weight in the efficacy study with the test items above are depicted in Figures 13A and 13B.

Example 33: In vivo efficacy study

- [0324]** NCI-H446, lung cancer xenograft model cell line was maintained in vitro using RPMI-1640 medium supplemented with 10% FBS in humidified cell culture incubator at 37°C with standard 5% CO₂ specs. The cells in an exponential growth phase were harvested and counted for tumor inoculation.

- [0325]** BALB/c nude mice (female) of 7–9 weeks old received a subcutaneous injection in the right front flank region with 5×10^6 tumor cells in 0.1 ml of PBS mixed with Matrigel (1:1) for tumor development. When tumors reached an average size of 100-200 mm³, randomization was performed into 5 groups of 8 mice and treatment began. Randomization will be performed based on “Matched distribution” method (Study Director™ software, version 3.1.399.19). The date of randomization will be denoted as day 0.

[0326] Electronic caliper measurement were performed 2 times a week. Daily observations for clinical signs, food and water consumption, behavioral changes, animals were weighed 2 times per week. The endpoint of the experiment is a tumor volume of 3,000 mm³, body weight loss over 20% or 28 days, whichever comes first.

- 5 **[0327]** Randomization: The randomization was performed when the mean tumor size reached approximately 100-200 mm³. Totally 40 mice were enrolled in NCI-H446 model study and randomly allocated to 5 groups, with 8 mice per group. Randomization will be performed based on "Matched distribution" method (Study Director™ software, version 3.1.399.19). The date of randomization will be denoted as day 0.
- 10 **[0328]** Test article administration: The ADC test article administration was performed via intravenous injection through tail vein, and the dosing volume was 10 mL/kg. For the ADC single dose only, initiated on the same day of randomization. Olaparib administration was performed orally at a dose of 50 mg/kg QD x 21days, also initiated on the same day of randomization. Dosing was conducted in a Laminar Flow Cabinet.
- 15 **[0329]** Observation and data collection: After tumor cells inoculation, the animals are checked daily for morbidity and mortality. At the time of routine monitoring, the animals are checked for any adverse effects of tumor growth and treatments on normal behavior such as mobility, visual estimation of food and water consumption, body weight gain/loss, eye/hair matting and any other abnormal effects. Tumor volumes are measured every 3-4 days in two dimensions using an caliper,
- 20 and the volume data are expressed in mm³ using the formula: $V = (L \times W \times W)/2$, where V is tumor volume, L is tumor length (the longest tumor dimension) and W is tumor width (the longest tumor dimension perpendicular to L). Dosing as well as tumor and body weight measurements will be conducted in a Laminar Flow Cabinet.

- 25 **[0330]** Table 8. Overview of test items and dose levels included in in vivo study

Test item	Dose levels (mg/kg)	Dose level Olaparib (mg/kg)	Group (N=8)	Administration	Dosing Schedule
vehicle	–	-	1	IV	Single dose
12C6-3	2.5	-	2	IV	Single dose
12C6-3	2.5	50	3	IV+PO	Single dose + QD x 21 days
vehicle	-	50	4	PO	QD x 21 days
12C6-9	2.5	-	5	IV	Single dose

[0331] Tumor volume over time and mice body weight in the efficacy study with the test items above are depicted in Figures 14A and 14B.

Example 34: In vivo tolerability study

- 30 **[0332]** Sprague Dawley rats (female) of 6-7 weeks old acclimatized for a week before being

enrolled in the study.

[0333] Randomization: In the tolerability study totally 15 rats were enrolled and randomly allocated to 5 study groups with 3 rats per group. Randomization was performed based on "Matched distribution" method (StudyDirector™ software, version 3.1.399.19). The date of randomization was denoted as day 0.


[0334] Test Article Administration: The treatment was initiated on the same day of randomization (day 0) per study design. Chosen dose levels were 75, 100, 125 and 150 mg/kg, single dose. All animals were dosed via slow intravenous injection (10 mL/kg).

[0335] Observation and Data Collection After randomization: The animals were checked daily for morbidity and mortality. During routine monitoring, the animals were checked for any effects of treatments on behavior such as mobility, food and water consumption, body weight gain/loss, eye/hair matting and any other abnormalities. Mortality and observed clinical signs were recorded for individual animals in detail. Dosing as well as body weight measurements were conducted in a Laminar Flow Cabinet. The body weights were measured daily by using StudyDirector™ software (version 3.1.399.19).

[0336] Body weight of the rats over time with 12C6-3 as test item at several dose levels are depicted in Figure 15.

Claims

1. An antibody-conjugate according to general structure (1):



5 wherein:

- AB is an antibody capable of targeting PTK7-expressing tumours;
- L is a linker that links Z to D;
- Z is a connecting group;
- L^6 is $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-\text{S}-(\text{L}^7)_w-$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-\text{N}(\text{H})\text{C}(\text{O})\text{CH}_2-$, $-\text{N}(\text{H})\text{C}(\text{O})\text{CF}_2-$ or $-\text{CH}_2-$;
- D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
- b is 0 or 1;
- x is 1 or 2; and
- y is 1, 2, 3 or 4.

- 20 2. The antibody-conjugate according to claim 1, wherein D is selected from the group consisting of anthracyclines, camptothecins, maytansinoids, enediynes, auristatins and pyrrolobenzodiazepine dimers, preferably wherein:

- D is an enediyne, preferably selected from the group consisting of calicheamicins, esperamicins, shishijimicins and namenamicins, more preferably calicheamicin; or
 - D is an auristatin, preferably MMAD, MMAE, MMAF or PF-06380101, more preferably MMAE or PF-06380101; or
 - D is a camptothecin, preferably selected from the group consisting of topotecan, silatecan, cositecan, exatecan, exatecan-S, DXd, SN-38, lurtotecan, gimatecan, belotecan, rubitecan, AMDCPT and G-AMDCPT, more preferably exatecan or DXd, most preferably exatecan,
- 30 more preferably D is calicheamicin or exatecan, more preferably D is exatecan.

3. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody has an Fc region which contains one or more mutations.


- 35 4. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody contains a V_L domain having at least 70 % sequence identity with a sequence selected from the group consisting of SEQ ID No. 2, 6, 9, 11, 14, 18, 21, 24, 32 and 36, and a V_H domain having at least 70 % sequence identity with a sequence selected from the group consisting of SEQ ID No. 1, 5, 10, 12, 13, 17, 23, 30 and 34, preferably wherein the antibody has a complementarity-determining region (CDR) having at least 90 % sequence identity.
- 40

-82-

5. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is cofetuzumab, and/or contains a light chain sequence according to SEQ ID No. 8 and a heavy chain sequence according to SEQ ID No. 7, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.
6. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is 12C6, and/or contains a light chain sequence according to SEQ ID No. 20 and a heavy chain sequence according to SEQ ID No. 19, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.
7. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is 12C6a, and/or contains a light chain sequence according to SEQ ID No. 22 and a heavy chain sequence according to SEQ ID No. 19, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.
8. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is 4D5, and/or contains a light chain sequence according to SEQ ID No. 16 and a heavy chain sequence according to SEQ ID No. 15, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.
9. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is 7C8, and/or contains a light chain sequence according to SEQ ID No. 25 and a heavy chain sequence according to SEQ ID No. 26, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.
10. The antibody-conjugate according to any one of claims 1 – 9, wherein linker L has the structure
- $$-(L^1)_n-(L^2)_o-(L^3)_p-(L^4)_q-$$
- wherein:
- L^1 , L^2 , L^3 and L^4 are each individually linkers that together link Z to D;
 - n, o, p and q are each individually 0 or 1, provided that $n + o + p + q = 1, 2, 3$ or 4, preferably wherein $n = o = p = 1$;
- and wherein:
- (a) linker L^1 is represented by:
- $$-(W)_k-(A)_d-(B)_e-(A)_f-(B)_g-(C(O))_g-, \text{ or}$$
- $$-(W)_k-(A)_d-(B)_e-(A)_f-(C(O))_g-N^*[-(A)_d-(B)_e-(A)_f-(C(O))_g-]_2,$$
- wherein:
- d and d' are individually 0 or 1;
 - e and e' are individually an integer in the range 1 – 10;
 - f and f' are individually 0, or 1;
 - g and g' are individually an integer in the range 0 – 10;

-83-

- k = 0 or 1 with the proviso that if k = 1 then d = 0;
- A is a sulfamide group according to structure (23)




(23)

- 5 wherein a = 0 or 1, and R¹³ is selected from the group consisting of hydrogen, C₁ – C₂₄ alkyl groups, C₃ – C₂₄ cycloalkyl groups, C₂ – C₂₄ (hetero)aryl groups, C₃ – C₂₄ alkyl(hetero)aryl groups and C₃ – C₂₄ (hetero)arylalkyl groups, the C₁ – C₂₄ alkyl groups, C₃ – C₂₄ cycloalkyl groups, C₂ – C₂₄ (hetero)aryl groups, C₃ – C₂₄ alkyl(hetero)aryl groups and C₃ – C₂₄ (hetero)arylalkyl groups optionally substituted and optionally
- 10 interrupted by one or more heteroatoms selected from O, S and NR¹⁴ wherein R¹⁴ is independently selected from the group consisting of hydrogen and C₁ – C₄ alkyl groups, or R¹³ is D connected to N via a spacer moiety, preferably wherein the spacer moiety is –(B)_g–(C(O))_g–(L²)_o–(L³)_p–(L⁴)_q–;
- W is –OC(O)–, –C(O)O–, –C(O)NH–, –NHC(O)–, –OC(O)NH–, –NHC(O)O–, –C(O)(CH₂)_mC(O)–, –C(O)(CH₂)_mC(O)NH– or –(4-Ph)CH₂NHC(O)(CH₂)_mC(O)NH–,
- 15 wherein m is an integer in the range 0 – 10;
- B is a –CH₂–CH₂–O– or a –O–CH₂–CH₂– moiety, or (B)_e is a –(CH₂–CH₂–O)_{e1}–CH₂–CH₂– moiety, wherein e1 is an integer in the range 1 – 10;
 - N* is a branching nitrogen atom, to which two instances of –(A)_d–(B)_e–(A)_f–(C(O))_g– are connected and both (C(O))_g moieties are connected to –(L²)_o–(L³)_p–(L⁴)_q–D, wherein L², L³, L⁴, o, p, q and D are each selected individually;
- 20

and/or

- (b) linker L² is a peptide spacer, preferably comprising 1 – 5 amino acids, more preferably a dipeptide, tripeptide or tetrapeptide spacer, most preferably wherein L² is represented by
- 25 general structure (L3):




(L3)

wherein R¹⁷ = CH₃ or CH₂CH₂CH₂NHC(O)NH₂;

and/or

- (c) linker L³ is a self-immolative spacer, preferably a para-aminobenzyloxycarbonyl (PABC) derivative according to structure (L4).
- 30



(L4)

-84-

wherein R^{21} is H, R^{26} or $C(O)R^{26}$, wherein R^{26} is $C_1 - C_{24}$ (hetero)alkyl groups, $C_3 - C_{10}$ (hetero)cycloalkyl groups, $C_2 - C_{10}$ (hetero)aryl groups, $C_3 - C_{10}$ alkyl(hetero)aryl groups and $C_3 - C_{10}$ (hetero)arylalkyl groups, which are optionally substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR^{28} wherein R^{28} is independently selected from the group consisting of hydrogen and $C_1 - C_4$ alkyl groups, preferably wherein R^{21} is H or $C(O)R^{26}$, wherein $R^{26} = 4$ -methyl-piperazine or morpholine, most preferably wherein R^{21} is H;

and/or


(d) linker L^4 is an aminoalkanoic acid spacer according to the structure $-NR^{22}-(C_x-alkylene)-C(O)-$, wherein x is an integer in the range 1 – 20 and R^{22} is H or $C_1 - C_4$ alkyl; or

linker L^4 is an ethyleneglycol spacer according to the structure $-NR^{22}-(CH_2-CH_2-O)_{e6}-(CH_2)_{e7}-C(O)-$, wherein $e6$ is an integer in the range 1 – 10, $e7$ is an integer in the range 1 – 3 and R^{22} is H or $C_1 - C_4$ alkyl; or


linker L^4 is a diamine spacer according to the structure $-NR^{22}-(C_x-alkylene)-NR^{22}-C(O))_h-$, wherein h is 0 or 1, x is an integer in the range 1 – 10 and R^{22} is H or $C_1 - C_4$ alkyl.

11. The antibody-conjugate according to any one of claims 1 – 10, wherein $b = 1$ and conjugation is via the glycan of the antibody, preferably wherein $j = 0$.

12. The antibody-conjugate according to any one of claims 1 – 11, wherein Z–L–D has a structure selected from the group consisting of (X) – (XIII):




(X)




(XI)

-85-




(XII)




(XIII)

5 wherein the wavy line indicates the connection to Z, and L^2 , L^4 , o and q are as defined in claim 10,

preferably wherein Z-L-D has a structure selected from the group consisting of (Xa), (Xlb), (XIIg), (XIIle) and (XIIh):




(Xa)




(Xlb)

10


-86-



(XIlg)



(XIIIe)



(XIIIh)

wherein the wavy line indicates the connection to Z, and R²² is as defined in claim 10.

13. The antibody-conjugate according to claim 12, which is according to structure (XII) or (XIII), and wherein $o = 1$ and the antibody is labetuzumab as defined in claim 5, preferably wherein $q = 0$.
14. The antibody-conjugate according to claim 13, which is according to structure (XII) or (XIII), wherein $q = 0$, $o = 1$, L^2 is a peptide spacer containing 1 – 5 amino acids, the antibody is labetuzumab as defined in claim 5 and the payload is exatecan.
15. The antibody-conjugate according to claim 1, which is according to structure (XIIIe) as defined in claim 12, and wherein the antibody is cofetuzumab as defined in claim 5 and the payload is exatecan.
16. A process for preparing the antibody-conjugate according to any one of claims 1 – 15, comprising:
- contacting an antibody comprising y core *N*-acetylglucosamine (GlcNAc) moieties, wherein $y = 1, 2, 3$ or 4 , with a compound of the formula $S(F)_xP$ in the presence of a catalyst, wherein $S(F)_x$ is a sugar derivative comprising x reactive groups F capable of reacting with a reactive group Q , x is 1 or 2 and P is a nucleoside mono- or diphosphate, and wherein the catalyst is capable of transferring the $S(F)_x$ moiety to the core-GlcNAc moiety, to obtain a modified antibody according to Formula (26):

$$AB-[GlcNAc(Fuc)_w-S\{F\}_x]_y$$

(26)

wherein

 - AB is an antibody capable of targeting PTK7-expressing tumours;
 - Fuc is fucose;
 - w is 0 or 1 ; and
 - reacting the modified antibody with a compound according to structure (2):


$$Q-L-D$$

(2)

wherein:

 - Q is a reactive moiety;
 - L is a linker that links Z to D ;
 - D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
- to obtain the antibody-conjugate according to structure (1).

17. The process according to claim 16, wherein the reaction of step (ii) is a nucleophilic reaction or a cycloaddition, preferably a 1,3-dipolar cycloaddition, and preferably wherein Q comprises or is an alkyne moiety and F comprises or is an azide moiety.
- 5 18. The process according to claim 16 or 17, wherein Q is a click probe comprising a (hetero)cycloalkyne moiety or a (hetero)cycloalkene moiety, preferably wherein the click probe Q is selected from the group consisting of (Q21) – (Q38) and (Q44) – (Q56):



wherein

- B is an anion; and
- the R group(s) on Si in (Q50) and (Q51) are alkyl or aryl;

5 and wherein F is a click probe selected from the group consisting of azide, tetrazine, triazine, nitron, nitrile oxide, nitrile imine, diazo compound, *ortho*-quinone, dioxothiophene and sydnone, preferably F is an azide moiety.

- 10 19. Method for targeting PTK7-expressing cells, contacting the antibody-conjugate according to any one of claims 1 – 15 with cells that may possibly be PTK7-expressing, preferably wherein the cells are PTK7-expressing tumour cells.
- 15 20. The method according to claim 19, wherein the targeting PTK7-expressing cells includes one or more of treating, imaging, diagnosing, preventing the proliferation of, containing and reducing PTK7-expressing cells, in particular PTK7-expressing tumour cells.
21. The method according to claim 19 or 20, wherein the subject suffers from colon cancer, lung cancer, breast cancer, ovarian cancer and esophageal cancer.