

TITLE OF THE INVENTION

Substituted Ureas and Methods of Making and Using Same

CROSS-REFERENCE TO RELATED APPLICATIONS

5 The present application claims priority under 35 U.S.C. § 119(e) to U.S. Provisional Patent Application No. 62/382,530, filed September 1, 2016, which is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

10 Pain is defined as an unpleasant sensory and emotional experience. Pain, however, can be informative and useful. For example, nociceptive pain is often indicative of injury (*e.g.*, tissue damage), and such pain typically evokes escape or protective behaviors in animals, including humans. However, inflammation, cellular and neuronal damage and other processes resulting from injury or disease can lead to states of chronic pathological pain.

15 Hyperalgesia is a condition in which enhanced sensitivity to noxious stimuli is present, and thus the perception of pain is exaggerated. Allodynia is a condition in which normally non-noxious stimuli become painful. Persistent or chronic pain, manifested as hyperalgesia and/or allodynia, remains challenging to treat. Many patients do not respond to existing therapeutics, or have their pain poorly managed (*i.e.*, inadequate relief), or experience relief

20 of an inadequate duration.

 Chronic pain contributes to over \$600 billion worth of healthcare expenditures annually, more than the yearly cost of cancer, heart disease, and diabetes combined. Neuropathic pain affects between 6 and 10% of the population, and is associated with decreased quality of life and socioeconomic burdens exceeding all other chronic pain

25 disorders. A recent meta-analysis of more than 200 neuropathic pain clinical trials indicates that the number of drugs needed to achieve even 50% pain relief in this population is between 4 and 10. This lack of efficacy has a profound influence on patient quality of life and is a source of frustration for caregivers. Many existing neuropathic pain therapeutics either have unknown mechanisms or are thought to reduce pain by reducing neuronal excitability.

30 Administration of opioids to treat pain is a well-recognized and commonly employed therapy in medicine. Mu opioids from natural sources have used by humans for millennia for a wide variety of purposes, including the relief of dysentery and pain. Opium alkaloids were not isolated until the 1800s, and synthetic opioids, whether recapitulations of nature or newly designed molecules, came much later. Mu opioid agonists are considered gold standard

analgesic agents, and are widely used for the treatment of a variety of mostly moderate-to-severe pain conditions in humans.

However, tachyphylaxis, tolerance to opioids and opioid-induced hyperalgesia can often result during the course of therapy. In such patients, increasingly higher doses of opioids are needed to provide an acceptable level of pain relief and, in doing so, the patient is thereby subjected to a higher risk of adverse side effects and safety concerns, which include respiratory depression, constipation, nausea and vomiting. Prolonged opioid therapy to treat chronic pain states may subject the patient to develop dependence on opioids, suffer opioid withdrawal on discontinuation of treatment, and some patients may be more susceptible to engage in abuse of these medications. These phenomena present significant clinical challenges for pain treatment.

G-Protein coupled receptors are membrane-bound proteins that contain seven transmembrane domains, an extracellular N-terminus, intracellular loops and an intracellular C-terminus. The GPCR family constitutes the largest class of cell surface receptors in the human genome (~800 different members), and small molecules that modulate GPCRs account for nearly one third of all marketed drugs.

Historically, from a conceptual perspective, GPCRs were thought to behave as switches (off/on), whereupon agonist binding induces a single conformational change to recruit G-protein coupling, thereby producing second-messengers such as cAMP to alter cellular functions (*e.g.*, polarization (neurons), contractility (myocytes), transcription, translation, and so forth). Antagonists block agonist access, thereby providing a means to prevent GPCR activation, which is also useful therapeutically depending upon the specific GPCR.

Upon ligand binding, GPCRs can not only interact with G-proteins as described above, but may also recruit cytosolic proteins of the arrestin family. Arrestin recruitment often occurs as a result of phosphorylation by a family of G-protein receptor kinases (GRKs). One role of the β -arrestin pathway, in contrast to G-protein binding, is to turn off signaling, possibly serving as an evolutionary 'brake' to avoid deleterious effects of constitutive signaling. β -Arrestins accomplish signal termination in several ways: 1) G-protein uncoupling causes desensitization (loss of signaling); 2) degradation of the second messenger(s) turns down/off signaling; and 3) receptor internalization precludes agonist binding and signaling since the receptor is not exposed/present on the cell membrane. Furthermore, there are non-classical signaling modalities (*e.g.*, MAP kinase) that can be


activated by β -arrestins.

There is a thus a need in the art for novel compounds and/or compositions that can be used to treat pain and/or reduce hyperalgesia and allodynia. In certain embodiments, the compounds and/or compositions do not induce significant (or any at all) respiratory
 5 depression, constipation, and/or tolerance. The present invention addresses this unmet need.

DETAILED DESCRIPTION OF THE INVENTION



The invention relates in one aspect to compounds, and compositions comprising such compounds, that can be used to treat and/or prevent pain in a subject in need thereof. In
 10 certain embodiments, the subject is a mammal. In other embodiments, the mammal is a dog or cat. In yet other embodiments, the mammal is human.

The invention includes a compound of formula (I), or a salt, solvate, enantiomer,



diastereoisomer and/or tautomer thereof:

(I), wherein: A is selected



, and B² is H; or B¹ and B² are independently selected from the group consisting of -CH₃ and -CH₂CH₃, or B¹ and B² combine to form a divalent substituent selected from the group consisting of -CH₂CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂CH₂-, and -CH₂CH₂OCH₂CH₂-; R¹ and R² are independently selected from the group consisting of H,
 20 CH₃, and CH₃ substituted with at least one selected from the group consisting of fluoro, chloro, cyano, hydroxyl and nitro; R³ is selected from the group consisting of H and CH₃; X is selected from the group consisting of S, O, and N-R³; R³ is selected from the group

consisting of hydrogen, alkyl and substituted alkyl; each occurrence of R is independently selected from the group consisting of fluoro, chloro, bromo, iodo, cyano, nitro, hydroxyl, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, and substituted phenyl; each occurrence of m is independently selected from the group consisting of 0, 1, 2 and 3; n is selected from the group consisting of 1, 2 and 3; and each occurrence of p is independently selected from the group consisting of 0, 1, 2 and 3.

In certain embodiments, the compounds of the invention bind to the mu opioid receptor (MOR). In certain embodiments, the compounds of the invention activate and/or act as agonists of the mu opioid receptor. In certain embodiments, upon binding to the MOR, the compounds of the invention induce certain secondary messenger signaling manifested as intracellular responses, such as but not limited to G-protein signaling and β -arrestin recruitment. In certain embodiments, upon binding to the MOR, the compounds of the invention decrease cyclic adenosine monophosphate (cAMP) levels. In certain embodiments, upon binding to the MOR, the compounds of the invention activate or behave as agonists of the MOR, induce intracellular responses such as but not limited to G-protein signaling and produce a decrease in cAMP.

In certain embodiments, the compounds of the invention do not significantly induce secondary messenger signaling, such as, but not limited to, recruitment, binding to, and/or association with β -arrestins. More specifically, compounds of this invention have half-maximal effective concentration (EC_{50}) values for β -arrestin that are higher than approximately 1 micromolar ($EC_{50} > 1 \mu M$) and/or exhibit β -arrestin activity of less than, or equal to 20% of the maximal response produced by the synthetic opioid peptide known as DAMGO (H-Tyr-D-Ala-Gly-NMe-Phe-Gly-OH) as defined by the assays described herein. In certain embodiments, the compounds of the invention induce intracellular responses such as G-protein recruitment and binding, such as for the proteins known as G_i , decrease cAMP levels, and do not significantly induce recruitment, binding to, and/or association with β -arrestins.

In certain embodiments, the compounds of the invention induce intracellular responses such as G-protein recruitment and binding such as for the proteins known as G_i , produce an decrease in cAMP, and do not significantly induce recruitment, binding to, and/or association with β -arrestins. In yet other embodiments, the compounds of the invention are G-protein biased agonists.

In certain embodiments, the compounds of the invention provide relief from, and/or

alleviate, pain in a subject. In other embodiments, the compounds of the invention bind to the MOR in a biased manner, decreasing cAMP levels with minimal or no recruitment, association and/or interaction with β -arrestins.

In certain embodiments, the compounds of the invention diminish pain without
5 producing respiratory depression, such as that caused by morphine and other well-known and widely used opioids and narcotics. In certain embodiments, the compounds of the invention diminish pain without producing constipation, such as that caused by morphine and other well-known and widely used opioids and narcotics. In certain embodiments, the compounds of the invention diminish pain without producing nausea, such as that caused by morphine
10 and other well-known and widely used opioids and narcotics. In certain embodiments, the compounds of the invention diminish pain without producing significant (or any at all) tolerance, such as that caused by morphine and other well-known and widely used opioids and narcotics. In certain embodiments, the compounds of the invention diminish pain without producing significant (or any at all) tachyphylaxis, such as that caused by morphine
15 and other well-known and widely used opioids and narcotics. In certain embodiments, the compounds of the invention diminish pain without producing emesis, such as that caused by morphine and other well-known and widely used opioids and narcotics. In certain embodiments, the compounds of the invention diminish pain without producing adverse effects associated with withdrawal, such as that caused by morphine and other well-known
20 and widely used opioids and narcotics. In certain embodiments, the compounds of the invention diminish pain without producing dependence, such as that caused by morphine and other well-known and widely used opioids and narcotics. In certain embodiments, the compounds of the invention diminish pain without producing one or more of the following phenomena: nausea, emesis, constipation, respiratory depression, tolerance, tachyphylaxis, dependence and/or addiction.
25

The invention further provides methods of administering a compound of the invention to provide pain relief. In certain embodiments, the pain comprises chronic pain. In other embodiments, the pain comprises neuropathic pain. In yet other embodiments, the pain comprises nociceptive pain. In yet other embodiments, the pain comprises hyperalgesia. In
30 yet other embodiments, the pain comprises allodynia.

Prior to the present day understanding of the β -arrestin pathways, many GPCR ligands were developed pursuing optimization of 7-transmembrane (7-TM) receptor binding and not surprisingly, many widely used therapeutics are now known to be promiscuous,

engaging both G-protein and β -arrestin pathways. For example, morphine, the classic, prototypic mu opioid receptor agonist, produces analgesia through G-protein signaling. However, this effect is therapeutically ‘offset’ by β -arrestin driven effects that produce receptor internalization and desensitization, which result in tolerance that necessitates
 5 increasing doses to maintain analgesic effect. In addition, β -arrestin driven signaling leads to the production of side effects such as constipation and respiratory depression.

Consistently, mice lacking β -arrestin2 exhibit enhanced and prolonged analgesia with very little tolerance, as well as attenuated respiratory depression and acute constipation, as compared to wild type. This provides evidence that MOR agonists that are selective for G-
 10 protein signaling, but devoid of β -arrestin-mediated effects, can provide morphine-like analgesia without the classic adverse effect profile. The biased MOR agonist, oliceridine or N-[(3-methoxy thiophen-2-yl)methyl]-2-[(9R)-9-pyridin-2-yl-6-oxaspiro[4.5]decan-9-yl]ethanamine, which is selective for G-protein signaling ($EC_{50} = 8$ nM, $E_{max} = 83\%$ vs morphine) with only minor β -arrestin recruitment (14% E_{max}) in cell-based assays, exhibited
 15 decreased on-target adverse effects (such as nausea, and decrease in respiratory drive) in healthy volunteers and produced a higher level of analgesia in patients following bunionectomy. This work demonstrates the power of biased ligands for GPCR signaling, which maintain/enhance desired therapeutic benefits while removing adverse side effects.

In certain embodiments, the ability of a compound to activate G-protein or β -arrestin
 20 is not an on/off switch for a given pathway, but rather a continuous function. To capture this, each compound is measured for activity in cell assays that are linked to β -arrestin recruitment and G-protein signaling. By comparing the activity between these two measures (accounting for both EC_{50} and maximal response reached, E_{max}) a relative activity (RA) value is determined.

$$\Delta RA = \text{LOG} \left(\frac{EMAX_{\beta\text{-Arr}}}{EC50_{\beta\text{-Arr}}} \right) - \text{LOG} \left(\frac{EMAX_{G\text{-Protein}}}{EC50_{G\text{-Protein}}} \right)$$

25 In the case of MOR it is desirable to have high activation of the G-protein pathway while reducing activation of β -arrestin. In the case of the formula above, this means a lower RA.

Definitions

30 As used herein, each of the following terms has the meaning associated with it in this section. Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which

this invention belongs. Generally, the nomenclature used herein and the laboratory procedures in animal pharmacology, pharmaceutical science, separation science, and organic chemistry are those well-known and commonly employed in the art.

As used herein, the articles “a” and “an” refer to one or to more than one (*i.e.* to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

As used herein, the term “about” is understood by persons of ordinary skill in the art and varies to some extent on the context in which it is used. As used herein when referring to a measurable value such as an amount, a temporal duration, and the like, the term “about” is meant to encompass variations of $\pm 20\%$ or $\pm 10\%$, more preferably $\pm 5\%$, even more preferably $\pm 1\%$, and still more preferably $\pm 0.1\%$ from the specified value, as such variations are appropriate to perform the disclosed methods.

In one aspect, the terms “co-administered” and “co-administration” as relating to a subject refer to administering to the subject a compound of the invention or salt thereof along with a compound that may also treat any disease or disorder contemplated herein and/or with a compound that is useful in treating other medical conditions but which in themselves may cause or facilitate any disease or disorder contemplated herein. In certain embodiments, the co-administered compounds are administered separately, or in any kind of combination as part of a single therapeutic approach. The co-administered compound may be formulated in any kind of combinations as mixtures of solids and liquids under a variety of solid, gel, and liquid formulations, and as a solution.

As used herein, the term “CYP450” as applied to enzymes refers to cytochrome P450 family of enzymes.

As used herein, a “disease” is a state of health of a subject wherein the subject cannot maintain homeostasis, and wherein if the disease is not ameliorated then the subject’s health continues to deteriorate.

As used herein, a “disorder” in a subject is a state of health in which the subject is able to maintain homeostasis, but in which the subject’s state of health is less favorable than it would be in the absence of the disorder. Left untreated, a disorder does not necessarily cause a further decrease in the subject’s state of health.

As used herein, the term “ED₅₀” refers to the effective dose of a formulation that produces 50% of the maximal effect in subjects that are administered that formulation.

As used herein, an “effective amount,” “therapeutically effective amount” or “pharmaceutically effective amount” of a compound is that amount of compound that is

sufficient to provide a beneficial effect to the subject to which the compound is administered.

“Instructional material,” as that term is used herein, includes a publication, a recording, a diagram, or any other medium of expression that can be used to communicate the usefulness of the composition and/or compound of the invention in a kit. The instructional material of the kit may, for example, be affixed to a container that contains the compound and/or composition of the invention or be shipped together with a container that contains the compound and/or composition. Alternatively, the instructional material may be shipped separately from the container with the intention that the recipient uses the instructional material and the compound cooperatively. Delivery of the instructional material may be, for example, by physical delivery of the publication or other medium of expression communicating the usefulness of the kit, or may alternatively be achieved by electronic transmission, for example by means of a computer, such as by electronic mail, or download from a website.

As used herein, the term “pharmaceutical composition” or “composition” refers to a mixture of at least one compound useful within the invention with a pharmaceutically acceptable carrier. The pharmaceutical composition facilitates administration of the compound to a subject.

As used herein, the term “pharmaceutically acceptable” refers to a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compound useful within the invention, and is relatively non-toxic, *i.e.*, the material may be administered to a subject without causing undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

As used herein, the term “pharmaceutically acceptable carrier” means a pharmaceutically acceptable material, composition or carrier, such as a liquid or solid filler, stabilizer, dispersing agent, suspending agent, diluent, excipient, thickening agent, solvent or encapsulating material, involved in carrying or transporting a compound useful within the invention within or to the subject such that it may perform its intended function. Typically, such constructs are carried or transported from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulation, including the compound useful within the invention, and not injurious to the subject. Some examples of materials that may serve as pharmaceutically acceptable carriers include: sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth;

malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; surface active agents; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; and other non-toxic compatible substances employed in pharmaceutical formulations. As used herein, "pharmaceutically acceptable carrier" also includes any and all coatings, antibacterial and antifungal agents, and absorption delaying agents, and the like that are compatible with the activity of the compound useful within the invention, and are physiologically acceptable to the subject. Supplementary active compounds may also be incorporated into the compositions. The "pharmaceutically acceptable carrier" may further include a pharmaceutically acceptable salt of the compound useful within the invention. Other additional ingredients that may be included in the pharmaceutical compositions used in the practice of the invention are known in the art and described, for example in Remington's Pharmaceutical Sciences (Genaro, Ed., Mack Publishing Co., 1985, Easton, PA), which is incorporated herein by reference.

As used herein, the language "pharmaceutically acceptable salt" refers to a salt of the administered compound prepared from pharmaceutically acceptable non-toxic acids and bases, including inorganic acids, inorganic bases, organic acids, inorganic bases, solvates, hydrates, and clathrates thereof.

The term "prevent," "preventing" or "prevention," as used herein, means avoiding or delaying the onset of symptoms associated with a disease or condition in a subject that has not developed such symptoms at the time the administering of an agent or compound commences. Disease, condition and disorder are used interchangeably herein.

By the term "specifically bind" or "specifically binds," as used herein, is meant that a first molecule preferentially binds to a second molecule (*e.g.*, a particular receptor or enzyme), but does not necessarily bind only to that second molecule.

As used herein, a "subject" may be a human or non-human mammal or a bird. Non-human mammals include, for example, livestock and pets, such as ovine, bovine, porcine, canine, feline and murine mammals. In certain embodiments, the subject is human.

The term "treat," "treating" or "treatment," as used herein, means reducing the frequency or severity with which symptoms of a disease or condition are experienced by a subject by virtue of administering an agent or compound to the subject.

As used herein, the term “alkyl,” by itself or as part of another substituent means, unless otherwise stated, a straight or branched chain hydrocarbon having the number of carbon atoms designated (*i.e.*, C₁-C₁₀ means one to ten carbon atoms) and includes straight, branched chain, or cyclic substituent groups. Examples include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *tert*-butyl, pentyl, neopentyl, hexyl, and cyclopropylmethyl. Most preferred is (C₁-C₆)alkyl, such as, but not limited to, ethyl, methyl, isopropyl, isobutyl, *n*-pentyl, *n*-hexyl and cyclopropylmethyl.

As used herein, the term “alkylene” by itself or as part of another substituent means, unless otherwise stated, a straight or branched hydrocarbon group having the number of carbon atoms designated (*i.e.*, C₁-C₁₀ means one to ten carbon atoms) and includes straight, branched chain, or cyclic substituent groups, wherein the group has two open valencies. Examples include methylene, 1,2-ethylene, 1,1-ethylene, 1,1-propylene, 1,2-propylene and 1,3-propylene.

As used herein, the term “cycloalkyl,” by itself or as part of another substituent means, unless otherwise stated, a cyclic chain hydrocarbon having the number of carbon atoms designated (*i.e.*, C₃-C₆ means a cyclic group comprising a ring group consisting of three to six carbon atoms) and includes straight, branched chain or cyclic substituent groups. Examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Most preferred is (C₃-C₆)cycloalkyl, such as, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

As used herein, the term “alkenyl,” employed alone or in combination with other terms, means, unless otherwise stated, a stable mono-unsaturated or di-unsaturated straight chain or branched chain hydrocarbon group having the stated number of carbon atoms. Examples include vinyl, propenyl (or allyl), crotyl, isopentenyl, butadienyl, 1,3-pentadienyl, 1,4-pentadienyl, and the higher homologs and isomers. A functional group representing an alkene is exemplified by -CH₂-CH=CH₂.

As used herein, the term “alkynyl,” employed alone or in combination with other terms, means, unless otherwise stated, a stable straight chain or branched chain hydrocarbon group with a triple carbon-carbon bond, having the stated number of carbon atoms. Non-limiting examples include ethynyl and propynyl, and the higher homologs and isomers. The term “propargylic” refers to a group exemplified by -CH₂-C≡CH. The term “homopropargylic” refers to a group exemplified by -CH₂CH₂-C≡CH. The term “substituted propargylic” refers to a group exemplified by -CR₂-C≡CR, wherein each occurrence of R is independently H, alkyl, substituted alkyl, alkenyl or substituted alkenyl, with the proviso that

at least one R group is not hydrogen. The term “substituted homopropargylic” refers to a group exemplified by $-\text{CR}_2\text{CR}_2-\text{C}\equiv\text{CR}$, wherein each occurrence of R is independently H, alkyl, substituted alkyl, alkenyl or substituted alkenyl, with the proviso that at least one R group is not hydrogen.

5 As used herein, the term “substituted alkyl,” “substituted cycloalkyl,” “substituted alkenyl” or “substituted alkynyl” means alkyl, cycloalkyl, alkenyl or alkynyl, as defined above, substituted by one, two or three substituents selected from the group consisting of halogen, alkoxy, tetrahydro-2-H-pyranyl, $-\text{NH}_2$, $-\text{N}(\text{CH}_3)_2$, (1-methyl-imidazol-2-yl), pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, $-\text{C}(=\text{O})\text{OH}$, trifluoromethyl, $-\text{C}\equiv\text{N}$, $-\text{C}(=\text{O})\text{O}(\text{C}_1-\text{C}_4)\text{alkyl}$, $-\text{C}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{O})\text{NH}(\text{C}_1-\text{C}_4)\text{alkyl}$, $-\text{C}(=\text{O})\text{N}((\text{C}_1-\text{C}_4)\text{alkyl})_2$, $-\text{SO}_2\text{NH}_2$, $-\text{C}(=\text{NH})\text{NH}_2$, and $-\text{NO}_2$, preferably containing one or two substituents selected from halogen, $-\text{OH}$, alkoxy, $-\text{NH}_2$, trifluoromethyl, $-\text{N}(\text{CH}_3)_2$, and $-\text{C}(=\text{O})\text{OH}$, more preferably selected from halogen, alkoxy and $-\text{OH}$. Examples of substituted alkyls include, but are not limited to, 2,2-difluoropropyl, 2-carboxycyclopentyl and 3-chloropropyl. In certain embodiments, the substituted alkyl is not substituted with a hydroxy group.

As used herein, the term “alkoxy” employed alone or in combination with other terms means, unless otherwise stated, an alkyl group having the designated number of carbon atoms, as defined above, connected to the rest of the molecule via an oxygen atom, such as, for example, methoxy, ethoxy, 1-propoxy, 2-propoxy (isopropoxy) and the higher homologs and isomers. Preferred are $(\text{C}_1-\text{C}_3)\text{alkoxy}$, such as, but not limited to, ethoxy and methoxy.

As used herein, the term “halo” or “halogen” alone or as part of another substituent means, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom, preferably, fluorine, chlorine, or bromine, more preferably, fluorine or chlorine.

As used herein, the term “heteroalkyl” by itself or in combination with another term means, unless otherwise stated, a stable straight or branched chain alkyl group consisting of the stated number of carbon atoms and one or two heteroatoms selected from the group consisting of O, N, and S, and wherein the nitrogen and sulfur atoms may be optionally oxidized and the nitrogen heteroatom may be optionally quaternized. The heteroatom(s) may be placed at any position of the heteroalkyl group, including between the rest of the heteroalkyl group and the fragment to which it is attached, as well as attached to the most distal carbon atom in the heteroalkyl group. Examples include: $-\text{O}-\text{CH}_2-\text{CH}_2-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_3$, $-\text{CH}_2-\text{S}-\text{CH}_2-\text{CH}_3$, and $-\text{CH}_2\text{CH}_2-\text{S}(=\text{O})-\text{CH}_3$. Up to two heteroatoms may be consecutive, such as, for example, $-\text{CH}_2-\text{NH}-\text{OCH}_3$, or $-\text{CH}_2-\text{CH}_2-\text{S}-\text{S}-\text{CH}_3$.

As used herein, the term “heteroalkenyl” by itself or in combination with another term means, unless otherwise stated, a stable straight or branched chain monounsaturated or di-unsaturated hydrocarbon group consisting of the stated number of carbon atoms and one or two heteroatoms selected from the group consisting of O, N, and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. Up to two heteroatoms may be placed consecutively. Examples include -CH=CH-O-CH₃, -CH=CH-CH₂-OH, -CH₂-CH=N-OCH₃, -CH=CH-N(CH₃)-CH₃, and -CH₂-CH=CH-CH₂-SH.

As used herein, the term “aromatic” refers to a carbocycle or heterocycle with one or more polyunsaturated rings and having aromatic character, *i.e.* having (4n+2) delocalized π (pi) electrons, where n is an integer.

As used herein, the term “aryl,” employed alone or in combination with other terms, means, unless otherwise stated, a carbocyclic aromatic system containing one or more rings (typically one, two or three rings) wherein such rings may be attached together in a pendent manner, such as a biphenyl, or may be fused, such as naphthalene. Examples include phenyl, anthracyl, and naphthyl. Preferred are phenyl and naphthyl, most preferred is phenyl.

As used herein, the term “aryl-(C₁-C₃)alkyl” means a functional group wherein a one to three carbon alkylene chain is attached to an aryl group, *e.g.*, -CH₂CH₂-phenyl or -CH₂-phenyl (benzyl). Preferred is aryl-CH₂- and aryl-CH(CH₃)-. The term “substituted aryl-(C₁-C₃)alkyl” means an aryl-(C₁-C₃)alkyl functional group in which the aryl group is substituted. Preferred is substituted aryl(CH₂)-. Similarly, the term “heteroaryl-(C₁-C₃)alkyl” means a functional group wherein a one to three carbon alkylene chain is attached to a heteroaryl group, *e.g.*, -CH₂CH₂-pyridyl. Preferred is heteroaryl-(CH₂)-. The term “substituted heteroaryl-(C₁-C₃)alkyl” means a heteroaryl-(C₁-C₃)alkyl functional group in which the heteroaryl group is substituted. Preferred is substituted heteroaryl-(CH₂)-.

As used herein, the term “heterocycle” or “heterocyclyl” or “heterocyclic” by itself or as part of another substituent means, unless otherwise stated, an unsubstituted or substituted, stable, mono- or multi-cyclic heterocyclic ring system that consists of carbon atoms and at least one heteroatom selected from the group consisting of N, O, and S, and wherein the nitrogen and sulfur heteroatoms may be optionally oxidized, and the nitrogen atom may be optionally quaternized. The heterocyclic system may be attached, unless otherwise stated, at any heteroatom or carbon atom that affords a stable structure. A heterocycle may be aromatic or non-aromatic in nature. In certain embodiments, the heterocycle is a heteroaryl.

As used herein, the term “heteroaryl” or “heteroaromatic” refers to a heterocycle having aromatic character. A polycyclic heteroaryl may include one or more rings that are partially saturated. Examples include tetrahydroquinoline and 2,3-dihydrobenzofuryl.

Examples of non-aromatic heterocycles include monocyclic groups such as aziridine, 5 oxirane, thiirane, azetidine, oxetane, thietane, pyrrolidine, pyrroline, imidazoline, pyrazolidine, dioxolane, sulfolane, 2,3-dihydrofuran, 2,5-dihydrofuran, tetrahydrofuran, thiophane, piperidine, 1,2,3,6-tetrahydropyridine, 1,4-dihydropyridine, piperazine, morpholine, thiomorpholine, pyran, 2,3-dihydropyran, tetrahydropyran, 1,4-dioxane, 1,3-dioxane, homopiperazine, homopiperidine, 1,3-dioxepane, 4,7-dihydro-1,3-dioxepin and 10 hexamethyleneoxide.

Examples of heteroaryl groups include pyridyl, pyrazinyl, pyrimidinyl (such as, but not limited to, 2- and 4-pyrimidinyl), pyridazinyl, thienyl, furyl, pyrrolyl, imidazolyl, thiazolyl, oxazolyl, pyrazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,3,4-triazolyl, tetrazolyl, 1,2,3-thiadiazolyl, 1,2,3-oxadiazolyl, 1,3,4-thiadiazolyl and 1,3,4-oxadiazolyl.

Examples of polycyclic heterocycles include indolyl (such as, but not limited to, 3-, 4-, 15 , 5-, 6- and 7-indolyl), indolinyl, quinolyl, tetrahydroquinolyl, isoquinolinyl (such as, but not limited to, 1- and 5-isoquinolinyl), 1,2,3,4-tetrahydroisoquinolinyl, cinnolinyl, quinoxalinyl (such as, but not limited to, 2- and 5-quinoxalinyl), quinazolinyl, phthalazinyl, 1,8-naphthyridinyl, 1,4-benzodioxanyl, coumarin, dihydrocoumarin, 1,5-naphthyridinyl, 20 benzofuryl (such as, but not limited to, 3-, 4-, 5-, 6- and 7-benzofuryl), 2,3-dihydrobenzofuryl, 1,2-benzisoxazolyl, benzothienyl (such as, but not limited to, 3-, 4-, 5-, 6-, and 7-benzothienyl), benzoxazolyl, benzothiazolyl (such as, but not limited to, 2-benzothiazolyl and 5-benzothiazolyl), purinyl, benzimidazolyl, benztriazolyl, thioxanthinyl, carbazolyl, carbolinyl, acridinyl, pyrrolizidinyl, and quinolizidinyl.

25 The aforementioned listing of heterocyclyl and heteroaryl moieties is intended to be representative and not limiting.

As used herein, the term “substituted” means that an atom or group of atoms has replaced hydrogen as the substituent attached to another group.

For aryl, aryl-(C₁-C₃)alkyl and heterocyclyl groups, the term “substituted” as applied 30 to the rings of these groups refers to any level of substitution, namely mono-, di-, tri-, tetra-, or penta-substitution, where such substitution is permitted. The substituents are independently selected, and substitution may be at any chemically accessible position. In certain embodiments, the substituents vary in number between one and four. In other embodiments, the substituents vary in number between one and three. In yet other


embodiments, the substituents vary in number between one and two. In yet other embodiments, the substituents are independently selected from the group consisting of C₁₋₆ alkyl, -OH, C₁₋₆ alkoxy, halo, amino, acetamido and nitro. As used herein, where a substituent is an alkyl or alkoxy group, the carbon chain may be branched, straight or cyclic, with straight being preferred.

The following abbreviations are used herein: Ar, argon; β -arrestin, *beta*-arrestin-2 (protein); Boc₂O, di-tert-butyl dicarbonate; C, carbon; cAMP, cyclic adenosine monophosphate; CHCl₃, chloroform; DCM (or CH₂Cl₂), methylene chloride (dichloromethane); DIPEA, N,N-diisopropylethylamine; DMF, N,N-dimethylformamide; E_{MAX}, maximal response reached; ESI, electrospray ionization; Et₂O, (di)ethyl ether; EtOAc, ethyl acetate; GCMS, gas chromatogram - mass spectrometry; H₂, hydrogen gas; HCl, hydrochloric acid; iPrOH, isopropanol; K₂CO₃, potassium carbonate; KOH, potassium hydroxide; LCMS, liquid chromatography - mass spectrometry; MeCN (or CH₃CN), acetonitrile; MHz, megahertz; MgSO₄, magnesium sulfate; MOR, mu opioid receptor; MS, mass spectrometry; NaHCO₃, sodium bicarbonate; Na₂SO₄, sodium sulfate; nc, not calculated; nd, not done (not conducted); NMR, nuclear magnetic resonance; Pd, palladium (metal); PE, petroleum ether; ppm, parts per million; RA, relative activity; Ra / Ni, Raney Nickel; TEA, trimethylamine.



Throughout this disclosure, various aspects of the invention may be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub-ranges as well as individual numerical values within that range and, when appropriate, partial integers of the numerical values within ranges. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed sub-ranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 2.7, 3, 4, 5, 5.3, and 6. This applies regardless of the breadth of the range.


Compounds and Compositions




The invention includes a compound of formula (I), or a salt, solvate, enantiomer, diastereoisomer and/or tautomer thereof.



(I), wherein:

A is selected from the group consisting of , ,



B¹ is selected from the group consisting of , , and , and B² is H;

5 or B¹ and B² are independently selected from the group consisting of -CH₃ and -CH₂CH₃, or

B¹ and B² combine to form a divalent substituent selected from the group consisting of -CH₂CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂CH₂-, and -CH₂CH₂OCH₂CH₂-;

R¹ and R² are independently selected from the group consisting of H, CH₃, and CH₃

10 substituted with at least one selected from the group consisting of fluoro, chloro, cyano, hydroxyl, and nitro;

R³ is selected from the group consisting of H and CH₃;

X is selected from the group consisting of S, O, and N-R³;

R³ is selected from the group consisting of hydrogen, alkyl, and substituted alkyl;

15 each occurrence of R is independently selected from the group consisting of fluoro, chloro, bromo, iodo, cyano, nitro, hydroxyl, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, and substituted phenyl;

each occurrence of m is independently selected from the group consisting of 0, 1, 2, and 3;

n is selected from the group consisting of 1, 2, and 3; and

20 each occurrence of p is independently selected from the group consisting of 0, 1, 2, and 3.

In certain embodiments, the alkyl is C₁-C₆ alkyl. In other embodiments, the substituted alkyl is C₁-C₆ alkyl substituted with at least one selected from the group

consisting of fluoro, chloro, bromo, iodo, cyano, nitro, and hydroxyl. In yet other embodiments, the alkenyl is C₂-C₆ alkenyl. In yet other embodiments, the substituted alkenyl is C₂-C₆ alkenyl substituted with at least one selected from the group consisting of hydrogen, alkyl, fluoro, chloro, phenyl, and substituted phenyl. In yet other embodiments, the alkynyl is C₂-C₆ alkynyl. In yet other embodiments, the substituted alkynyl is C₂-C₆ alkynyl substituted with at least one selected from the group consisting of hydrogen, alkyl, phenyl, and substituted phenyl. In yet other embodiments, the substituted phenyl is phenyl substituted with at least one selected from the group consisting of alkyl, fluoro, chloro, bromo, iodo, cyano, nitro, hydroxyl, carboxylic acid, carboxylalkyl, and carboxamide.

In certain embodiments, A is isoindolin-2-yl. In other embodiments, A is 5-fluoroisoindolin-2-yl. In yet other embodiments, A is 5-chloroisoindolin-2-yl. In yet other embodiments, A is 5-methoxyisoindolin-2-yl. In yet other embodiments, A is 5-methylisoindolin-2-yl. In yet other embodiments, A is 5-hydroxyisoindolin-2-yl. In yet other embodiments, A is 5-cyanoisoindolin-2-yl. In other embodiments, A is 4-fluoroisoindolin-2-yl. In yet other embodiments, A is 4-chloroisoindolin-2-yl. In yet other embodiments, A is 4-methoxyisoindolin-2-yl. In yet other embodiments, A is 4-methylisoindolin-2-yl. In yet other embodiments, A is 4-hydroxyisoindolin-2-yl. In yet other embodiments, A is 4-cyanoisoindolin-2-yl. In yet other embodiments, A is 5,6-difluoroisoindolin-2-yl. In yet other embodiments, A is 5,6-dichloroisoindolin-2-yl. In yet other embodiments, A is 6-chloro-5-fluoroisoindolin-2-yl. In yet other embodiments, A is 5-chloro-6-fluoroisoindolin-2-yl.

In certain embodiments, A is 1,2,3,4-tetrahydroquinolin-1-yl. In other embodiments, A is 5-fluoro-1,2,3,4-tetrahydroquinolin-1-yl. In other embodiments, A is 6-fluoro-1,2,3,4-tetrahydroquinolin-1-yl. In other embodiments, A is 7-fluoro-1,2,3,4-tetrahydroquinolin-1-yl. In other embodiments, A is 8-fluoro-1,2,3,4-tetrahydroquinolin-1-yl. In other embodiments, A is 5-hydroxy-1,2,3,4-tetrahydroquinolin-1-yl. In other embodiments, A is 6-hydroxy-1,2,3,4-tetrahydroquinolin-1-yl. In other embodiments, A is 7-hydroxy-1,2,3,4-tetrahydroquinolin-1-yl. In other embodiments, A is 8-hydroxy-1,2,3,4-tetrahydroquinolin-1-yl.

In certain embodiments, A is 1,2,3,4-tetrahydronaphthalen-1-yl. In other embodiments, A is 1,2,3,4-tetrahydronaphthalen-2-yl. In yet other embodiments, A is 5-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl. In yet other embodiments, A is 6-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl. In yet other embodiments, A is 7-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl. In yet other embodiments, A is 8-fluoro-1,2,3,4-tetrahydronaphthalen-1-yl. In yet other embodiments, A is 5-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl. In yet other

embodiments, A is 6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl. In yet other embodiments, A is 7-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl. In yet other embodiments, A is 8-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl. In yet other embodiments, A is 5-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl. In yet other embodiments, A is 6-fluoro-1,2,3,4-tetrahydro
5 naphthalen-2-yl. In yet other embodiments, A is 7-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl. In yet other embodiments, A is 8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl. In yet other embodiments, A is 5-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl. In yet other embodiments, A is 6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl. In yet other embodiments, A is 7-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl. In yet other embodiments, A is 8-hydroxy-
10 1,2,3,4-tetrahydronaphthalen-2-yl.

In certain embodiments, A is 3-phenyl- pyrrolidin-1-yl. In other embodiments, A is 3-substituted phenyl- pyrrolidin-1-yl.

In certain embodiments, B¹ is 3-thienyl. In other embodiments, B¹ is 2-thienyl. In yet other embodiments, B¹ is 2-chloro-4-fluorophenyl. In yet other embodiments, B¹ is 4-chloro-
15 2-fluorophenyl. In yet other embodiments, B¹ is 4-methoxyphenyl. In yet other embodiments, B¹ is 4-hydroxyphenyl. In yet other embodiments, B¹ is phenyl. In yet other embodiments, B² is H.

In certain embodiments, B¹ is -CH₃ and B² is -CH₃. In other embodiments, B¹ is -CH₃ and B² is -CH₂CH₃. In yet other embodiments, B¹ is -CH₂CH₃ and B² is -CH₃. In yet other
20 embodiments, B¹ is -CH₂CH₃ and B¹ is -CH₂CH₃.

In certain embodiments, B¹ and B² combine to form -CH₂CH₂CH₂CH₂-. In other embodiments, B¹ and B² combine to form -CH₂CH₂CH₂CH₂CH₂-. In yet other embodiments, B¹ and B² combine to form -CH₂CH₂OCH₂CH₂-.

In certain embodiments, R is fluoro. In other embodiments, R is chloro. In yet other
25 embodiments, R is bromo. In yet other embodiments, R is iodo. In yet other embodiments, R is cyano. In yet other embodiments, R is nitro. In yet other embodiments, R is hydroxyl. In yet other embodiments, R is alkyl. In yet other embodiments, R is substituted alkyl. In yet other embodiments, R is alkenyl. In yet other embodiments, R is substituted alkenyl. In yet other embodiments, R is alkynyl. In yet other embodiments, R is substituted alkynyl. In
30 yet other embodiments, R is phenyl. In yet other embodiments, R is substituted phenyl.

In certain embodiments, in A at least one R is fluoro. In other embodiments, in A at least one R is chloro. In yet other embodiments, in A at least one R is bromo. In yet other embodiments, in A at least one R is iodo. In yet other embodiments, in A at least one R is cyano. In yet other embodiments, in A at least one R is nitro. In yet other embodiments, in

A at least one R is hydroxyl. In yet other embodiments, in A at least one R is alkyl. In yet other embodiments, in A at least one R is substituted alkyl. In yet other embodiments, in A at least one R is alkenyl. In yet other embodiments, in A at least one R is substituted alkenyl. In yet other embodiments, in A at least one R is alkynyl. In yet other
 5 embodiments, in A at least one R is substituted alkynyl. In yet other embodiments, in A at least one R is phenyl. In yet other embodiments, in A at least one R is substituted phenyl.

In certain embodiments, in B¹ at least one R is fluoro. In other embodiments, in B¹ at least one R is chloro. In yet other embodiments, in B¹ at least one R is bromo. In yet other embodiments, in B¹ at least one R is iodo. In yet other embodiments, in B¹ at least one R is cyano. In yet other embodiments, in B¹ at least one R is nitro. In yet other embodiments, in B¹ at least one R is hydroxyl. In yet other embodiments, in B¹ at least one R is alkyl. In yet other embodiments, in B¹ at least one R is substituted alkyl. In yet other embodiments, in B¹ at least one R is alkenyl. In yet other embodiments, in B¹ at least one R is substituted alkenyl. In yet other embodiments, in B¹ at least one R is alkynyl. In yet other
 10 embodiments, in B¹ at least one R is substituted alkynyl. In yet other embodiments, in B¹ at least one R is phenyl. In yet other embodiments, in B¹ at least one R is substituted phenyl.

In certain embodiments, R¹ is H. In other embodiments, R¹ is CH₃. In yet other embodiments, R² is H. In yet other embodiments, R² is CH₃.

In certain embodiments, m is 0. In other embodiments, m is 1. In other
 20 embodiments, m is 2. In other embodiments, m is 3. In other embodiments, n is 1. In other embodiments, n is 2. In other embodiments, n is 3. In other embodiments, p is 0. In other embodiments, p is 1. In other embodiments, p is 2. In other embodiments, p is 3.

In certain embodiments, the compound of the invention is at least one selected from the group consisting of: N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-isoindoline-2-
 25 carboxamide; N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide; (-)-ent-N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide; (+)-ent-N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide; N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-4-fluoroisoindoline-2-carboxamide; 5-Chloro-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-
 30 carboxamide; N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methoxyisoindoline-2-carboxamide; N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5,6-difluoroisoindoline-2-carboxamide; N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide; N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-7-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide; N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-

dihydroquinoline-1(2H)-carboxamide; 1-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3-
 (1,2,3,4-tetrahydronaphthalen-2-yl)urea; 1-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3-
 (1,2,3,4-tetrahydronaphthalen-1-yl)urea; N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-
 methylisoindoline-2-carboxamide; 3-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-1-methyl-
 5 1-(1,2,3,4-tetrahydronaphthalen-1-yl)urea; N-(2-(2-Chloro-4-fluorophenyl)-2-(dimethyl
 amino)ethyl)isoindoline-2-carboxamide; N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)
 isoindoline-2-carboxamide; N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoro
 isoindoline-2-carboxamide; N-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)-5-
 fluoroisoindoline-2-carboxamide; N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-
 10 carboxamide; 1-(2-(Dimethylamino)-2-phenylethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-
 yl)urea; N-(2-(Dimethylamino)-2-phenylethyl)-5-fluoroisoindoline-2-carboxamide; 3-(2-
 Dimethylamino-2-thiophen-3-yl-ethyl)-1-methyl-1-(S)-1,2,3,4-tetrahydro-naphthalen-1-yl-
 urea; (R)-3-Phenyl-pyrrolidine-1-carboxylic acid (2-dimethylamino-2-thiophen-3-yl-ethyl)-
 amide; (S)-3-Phenyl-pyrrolidine-1-carboxylic acid (2-dimethylamino-2-thiophen-3-yl-ethyl)-
 15 amide; (S)-3-Phenyl-pyrrolidine-1-carboxylic acid [2-dimethylamino-2-(4-methoxy-phenyl)-
 ethyl]-amide; (R)-3-Phenyl-pyrrolidine-1-carboxylic acid [2-dimethylamino-2-(4-methoxy-
 phenyl)-ethyl]-amide; 4-Cyano-1,3-dihydro-isoindole-2-carboxylic acid (2-dimethylamino-
 2-thiophen-3-yl-ethyl)-amide; 3-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-1-methyl-1-(R)-
 1,2,3,4-tetrahydro-naphthalen-1-yl-urea; 5-Fluoro-1,3-dihydro-isoindole-2-carboxylic acid
 20 (1-dimethylamino-cyclohexylmethyl)-amide; 1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-
 (R)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea; 7-Hydroxy-3,4-dihydro-2H-quinoline-1-
 carboxylic acid (2-dimethylamino-2-thiophen-3-yl-ethyl)-amide; 1-(2-Dimethylamino-2-
 thiophen-3-yl-ethyl)-3-((R)-7-hydroxy-1,2,3,4-tetrahydro-naphthalen-2-yl)-urea; 1-(2-
 Dimethylamino-2-thiophen-3-yl-ethyl)-3-(S)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea; 1-(2-
 25 Dimethylamino-2-thiophen-3-yl-ethyl)-3-(S)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea; 6-
 Hydroxy-3,4-dihydro-2H-quinoline-1-carboxylic acid (2-dimethylamino-2-thiophen-3-yl-
 ethyl)-amide; 1,3-Dihydro-isoindole-2-carboxylic acid (1-dimethylamino-cyclohexyl
 methyl)-amide; (R)-1-((1-(Dimethylamino)cyclohexyl)methyl)-3-(1,2,3,4-tetrahydro
 naphthalen-2-yl)urea; (S)-1-((1-(Dimethylamino)cyclohexyl)methyl)-3-(1,2,3,4-tetrahydro
 30 naphthalen-2-yl)urea; (-)-N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide;
 (+)-N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide; 1-(2-Dimethylamino-
 2-thiophen-3-yl-ethyl)-3-(5-hydroxy-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea; 1-(2-Dimethyl
 amino-2-thiophen-3-yl-ethyl)-3-((S)-7-hydroxy-1,2,3,4-tetrahydro-naphthalen-2-yl)-urea; 5-
 Fluoro-1,3-dihydro-isoindole-2-carboxylic acid (4-dimethylamino-tetrahydro-pyran-4-

ylmethyl)-amide; 5-Fluoro-1,3-dihydro-isoindole-2-carboxylic acid (2-dimethylamino-2-methyl-propyl)-amide; 5-Fluoro-1,3-dihydro-isoindole-2-carboxylic acid (1-dimethylamino-cyclopentylmethyl)-amide; 1,3-Dihydro-isoindole-2-carboxylic acid (1-dimethylamino-cyclopentylmethyl)-amide; 1-(1-Dimethylamino-cyclopentylmethyl)-3-(S)-1,2,3,4-

5 tetrahydronaphthalen-2-yl-urea; 1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(R)-1,2,3,4-tetrahydronaphthalen-2-yl-urea; 1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(S)-1,2,3,4-tetrahydronaphthalen-2-yl-urea; N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide; N-(2-(Dimethylamino)-2-phenylethyl)-3,4-dihydroquinoline-1(2H)-carboxamide; N-(2-(Dimethylamino)-2-phenylethyl)-6-fluoro-3,4-

10 dihydroquinoline-1(2H)-carboxamide; 1-((R)-3-(dimethylamino)-3-(thiophen-3-yl)propyl)-3-((R)-1,2,3,4-tetrahydronaphthalen-2-yl)urea; 1-((R)-3-(dimethylamino)-3-(thiophen-3-yl)propyl)-3-((S)-1,2,3,4-tetrahydronaphthalen-2-yl)urea; 1-((S)-3-(dimethylamino)-3-(thiophen-3-yl)propyl)-3-((R)-1,2,3,4-tetrahydronaphthalen-2-yl)urea; 1-((S)-3-(dimethylamino)-3-(thiophen-3-yl)propyl)-3-((S)-1,2,3,4-tetrahydronaphthalen-2-yl)urea;

15 (R)-N-((R)-3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-3-phenylpyrrolidine-1-carboxamide; (S)-N-((R)-3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-3-phenylpyrrolidine-1-carboxamide; (S)-N-((S)-3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-3-phenylpyrrolidine-1-carboxamide; (-)-N-(2-(Methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide; (+)-N-(2-(Methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide; (-)

20)-4-Fluoro-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide; (+)-4-Fluoro-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide; (-)-5-Fluoro-N-(2-(4-methoxyphenyl)-2-(methylamino)ethyl)isoindoline-2-carboxamide; (+)-5-Fluoro-N-(2-(4-methoxyphenyl)-2-(methylamino)ethyl)isoindoline-2-carboxamide; (R)-N-(3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisoindoline-2-carboxamide; (S)-N-(3-

25 (Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisoindoline-2-carboxamide; N-(2-(Dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-fluoroisoindoline-2-carboxamide; N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-hydroxyisoindoline-2-carboxamide; N-(2-(dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-hydroxyisoindoline-2-carboxamide; 5-Fluoro-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide; N-(2-

30 (Methylamino)-2-phenylethyl)isoindoline-2-carboxamide; (-)-N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide; (+)-N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide; or a salt, solvate, enantiomer, diastereoisomer and/or tautomer thereof.

The compounds of the invention may possess one or more stereocenters, and each stereocenter may exist independently in either the (*R*) or (*S*) configuration. In certain

embodiments, compounds described herein are present in optically active or racemic forms. It is to be understood that the compounds described herein encompass racemic, optically-active, regioisomeric and stereoisomeric forms, or combinations thereof that possess the therapeutically useful properties described herein. Preparation of optically active forms is achieved in any suitable manner, including by way of non-limiting example, by resolution of the racemic form with recrystallization techniques, synthesis from optically-active starting materials, chiral synthesis, or chromatographic separation using a chiral stationary phase. In certain embodiments, a mixture of one or more isomer is utilized as the therapeutic compound described herein. In other embodiments, compounds described herein contain one or more chiral centers. These compounds are prepared by any means, including stereoselective synthesis, enantioselective synthesis and/or separation of a mixture of enantiomers and/or diastereomers. Resolution of compounds and isomers thereof is achieved by any means including, by way of non-limiting example, chemical processes, enzymatic processes, fractional crystallization, distillation, and chromatography. All possible stereochemical configurations of a given compound containing chiral center(s) are contemplated. All possible mixtures enriched with a particular enantiomer or diastereomer(s) are contemplated. All pure individual enantiomers or diastereomers are contemplated.

The methods and formulations described herein include the use of N-oxides (if appropriate), crystalline forms (also known as polymorphs), solvates, amorphous phases, and/or pharmaceutically acceptable salts of compounds having the structure of any compound of the invention, as well as metabolites and active metabolites of these compounds having the same type of activity. Solvates include water, ether (*e.g.*, tetrahydrofuran, methyl tert-butyl ether) or alcohol (*e.g.*, ethanol) solvates, acetates and the like. In certain embodiments, the compounds described herein exist in solvated forms with pharmaceutically acceptable solvents such as water, and ethanol. In other embodiments, the compounds described herein exist in unsolvated form.

In certain embodiments, the compounds of the invention may exist as tautomers. "Tautomerization" is a form of isomerization involving the migration of a proton accompanied by changes in bond order, often the interchange of a single bond with an adjacent double bond. Where tautomerization is possible, (*e.g.* in solution), a chemical equilibrium of tautomers can be reached. One well known example of tautomerization is between a ketone and its corresponding enol. Heterocycles may form tautomers such as the interconversion of pyrrolidinone and hydroxypyrrole. All tautomers are included within the scope of the compounds presented herein.

In certain embodiments, compounds described herein are prepared as prodrugs. A “prodrug” refers to an agent that is converted into the parent drug *in vivo*. In certain embodiments, upon *in vivo* administration, a prodrug is chemically converted to the biologically, pharmaceutically or therapeutically active form of the compound. In other
5 embodiments, a prodrug is enzymatically metabolized by one or more steps or processes to the biologically, pharmaceutically or therapeutically active form of the compound.

In certain embodiments, sites on, for example, the aromatic ring portion of compounds of the invention is susceptible to various metabolic reactions. Incorporation of appropriate substituents on the aromatic ring structures may reduce, minimize or eliminate
10 this metabolic pathway. In certain embodiments, the appropriate substituent to decrease or eliminate the susceptibility of the aromatic ring to metabolic reactions is, by way of example only, a deuterium, a halogen, or an alkyl group.

Compounds described herein also include isotopically labeled compounds wherein one or more atoms is replaced by an atom having the same atomic number, but an atomic
15 mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes suitable for inclusion in the compounds described herein include and are not limited to ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{36}Cl , ^{18}F , ^{123}I , ^{125}I , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{32}P , and ^{35}S . In certain embodiments, isotopically labeled compounds are useful in drug and/or substrate tissue distribution studies. In other embodiments, substitution with heavier isotopes such as
20 deuterium affords greater metabolic stability (for example, increased *in vivo* half-life or reduced dosage requirements). In yet other embodiments, substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , is useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. Isotopically-labeled compounds are prepared by any suitable method or by processes using an appropriate isotopically-labeled
25 reagent in place of the non-labeled reagent otherwise employed.

In certain embodiments, the compounds described herein are labeled by other means, including, but not limited to, the use of chromophores or fluorescent moieties, bioluminescent labels, or chemiluminescent labels.

Compounds of the invention can in certain embodiments form acids or bases. In
30 certain embodiments, the invention contemplates acid addition salts. In other embodiments, the invention contemplates base addition salts. In yet other embodiments, the invention contemplates pharmaceutically acceptable acid addition salts. In yet other embodiments, the invention contemplates pharmaceutically acceptable base addition salts. Pharmaceutically acceptable salts refer to salts of those bases or acids that are not toxic or otherwise

biologically undesirable.

Suitable pharmaceutically acceptable acid addition salts may be prepared from an inorganic acid or from an organic acid. Examples of inorganic acids include hydrochloric, hydrobromic, hydriodic, nitric, carbonic, sulfuric (including sulfate and hydrogen sulfate), and phosphoric acids (including hydrogen phosphate and dihydrogen phosphate).

Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, examples of which include formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, malonic, saccharin, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, 4-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, trifluoromethanesulfonic, 2-hydroxyethanesulfonic, p-toluenesulfonic, sulfanilic, cyclohexylaminosulfonic, stearic, alginic, β -hydroxybutyric, salicylic, galactaric and galacturonic acid.

Suitable pharmaceutically acceptable base addition salts of compounds of the invention include, for example, metallic salts including alkali metal, alkaline earth metal and transition metal salts such as, for example, calcium, magnesium, potassium, sodium, lithium and copper, iron and zinc salts. Pharmaceutically acceptable base addition salts also include organic salts made from basic amines such as, for example, N,N'-dibenzylethylene-diamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (*N*-methylglucamine) and procaine. All of these salts may be prepared from the corresponding compound by reacting, for example, the appropriate acid or base with the compound.


The compounds described herein, and other related compounds having different substituents are synthesized using techniques and materials described herein and as described, for example, in Fieser & Fieser's Reagents for Organic Synthesis, Volumes 1-17 (John Wiley and Sons, 1991); Rodd's Chemistry of Carbon Compounds, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989); Organic Reactions, Volumes 1-40 (John Wiley and Sons, 1991), Larock's Comprehensive Organic Transformations (VCH Publishers Inc., 1989), March, Advanced Organic Chemistry 4th Ed., (Wiley 1992); Carey & Sundberg, Advanced Organic Chemistry 4th Ed., Vols. A and B (Plenum 2000,2001), and Green & Wuts, Protective Groups in Organic Synthesis 3rd Ed., (Wiley 1999) (all of which are incorporated by reference for such disclosure). General methods for the preparation of compound as described herein are modified by the use of appropriate reagents and conditions, for the introduction of the various moieties found in the formula as provided herein.

Compounds described herein are synthesized using any suitable procedures starting

from compounds that are available from commercial sources, or are prepared using procedures described herein.

In certain embodiments, reactive functional groups, such as hydroxyl, amino, imino, thio or carboxy groups, are protected in order to avoid their unwanted participation in reactions. Protecting groups are used to block some or all of the reactive moieties and prevent such groups from participating in chemical reactions until the protective group is removed. In other embodiments, each protective group is removable by a different means. Protective groups that are cleaved under totally disparate reaction conditions fulfill the requirement of differential removal. In certain embodiments, protective groups are removed by acid, base, reducing conditions (such as, for example, hydrogenolysis), and/or oxidative conditions. Groups such as trityl, dimethoxytrityl, acetal and t-butyl dimethylsilyl are acid labile and are used to protect carboxy and hydroxy reactive moieties in the presence of amino groups protected with Cbz groups, which are removable by hydrogenolysis, and Fmoc groups, which are base labile. Carboxylic acid and hydroxy reactive moieties are blocked with base labile groups such as, but not limited to, methyl, ethyl, and acetyl, in the presence of amines that are blocked with acid labile groups, such as t-butyl carbamate, or with carbamates that are both acid and base stable but hydrolytically removable. In certain embodiments, carboxylic acid and hydroxy reactive moieties are blocked with hydrolytically removable protective groups such as the benzyl group, while amine groups capable of hydrogen bonding with acids are blocked with base labile groups such as Fmoc. Carboxylic acid reactive moieties are protected by conversion to simple ester compounds as exemplified herein, which include conversion to alkyl esters, or are blocked with oxidatively-removable protective groups such as 2,4-dimethoxybenzyl, while co-existing amino groups are blocked with fluoride labile silyl carbamates. Allyl blocking groups are useful in the presence of acid- and base- protecting groups since the former are stable and are subsequently removed by metal or pi-acid catalysts. For example, an allyl-blocked carboxylic acid is deprotected with a palladium-catalyzed reaction in the presence of acid labile t-butyl carbamate or base-labile acetate amine protecting groups. Yet another form of protecting group is a resin to which a compound or intermediate is attached. As long as the residue is attached to the resin, that functional group is blocked and does not react. Once released from the resin, the functional group is available to react.


Typically blocking/protecting groups may be selected from:



Other protecting groups, plus a detailed description of techniques applicable to the creation of protecting groups and their removal are described in Greene & Wuts, Protective Groups in Organic Synthesis, 3rd Ed., John Wiley & Sons, New York, NY, 1999, and Kocienski, Protective Groups, Thieme Verlag, New York, NY, 1994, which are incorporated herein by reference for such disclosure.


In certain embodiments, compounds of the invention can be prepared according to the following general schemes.

In certain embodiments, an isoindoline is reacted with a similar activating agent, such as but not limited to, para-nitrophenyl-chloroformate to afford an electrophilic carbamate, which then undergoes coupling with a N^1, N^1 -dialkyl-1-(aryl)ethane-1,2-diamine, such as but not limited to N^1, N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine (**Scheme 1**).




Scheme 1


In certain embodiments, a N^1, N^1 -dialkyl-1-(aryl)ethane-1,2-diamine, such as but not limited to N^1, N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine is reacted with an activating agent, such as but not limited to para-nitrophenyl-chloroformate to afford an electrophilic carbamate, which undergoes coupling with an isoindoline (**Scheme 2**).

**Scheme 2**


In certain embodiments, a tetrahydroquinoline is reacted with an activating agent, such as but not limited to para-nitrophenyl-chloroformate, to afford an electrophilic carbamate, which undergoes coupling with a N^1,N^1 -dialkyl-1-(aryl)ethane-1,2-diamine, such as but not limited to N^1,N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine (**Scheme 3**).

**Scheme 3**


In certain embodiments, a N^1,N^1 -dialkyl-1-(aryl)ethane-1,2-diamine, such as but not limited to N^1,N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine, is reacted with an activating agent, such as but not limited to para-nitrophenyl-chloroformate, to afford an electrophilic carbamate, which undergoes coupling with a tetrahydroquinoline (**Scheme 4**).

**Scheme 4**


In certain embodiments, an aminotetralin is reacted with an activating agent, such as but not limited to para-nitrophenyl-chloroformate, to afford an electrophilic carbamate, which undergoes coupling with a N^1,N^1 -dialkyl-1-(aryl)ethane-1,2-diamine, such as but not limited to N^1,N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine (**Scheme 5**).

**Scheme 5**

In certain embodiments, a N^1,N^1 -dialkyl-1-(aryl)ethane-1,2-diamine, such as but not limited to N^1,N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine, is reacted with an activating agent, such as but not limited to para-nitrophenyl-chloroformate, to afford an electrophilic carbamate, which undergoes coupling with an aminotetralin (**Scheme 6**).

**Scheme 6**

In certain embodiments, an optically-pure (tert-butyl)sulfinamide (**i**) is reacted with an aryl-carboxaldehyde or aralkyl-carboxaldehyde, in the presence of an organic base such as but not limited to pyrrolidine, to produce a N-aralkylene-sulfinamide (**ii**). Subsequent reaction with the (carbon-based) anion of acetonitrile produces the diastereomeric N-(2-cyano-1-(aryl)alkyl)-2-sulfinamide (**iii**) which are separated by a method such as but not limited to chromatography or crystallization. Hydrolysis of the sulfinamide affords optically-enriched 3-amino-3-aryl)propanenitrile (**iv**). Reductive alkylation, using for example an aldehyde and a reducing agent, such as but not limited to (sodium) triacetoxyborohydride, affords the corresponding optically-enriched 3-(dialkylamino)-3-aryl-propanenitrile (**v**). Treatment of (**v**) with a strong reducing agent, such as but not limited to lithium aluminum hydride, produces a N^1,N^1 -dialkyl-1-(aryl)propane-1,3-diamine (**vi**). The diamine is activated by reaction with a chloroformate such as but not limited to para-nitrophenyl chloroformate to afford carbamate (**vii**), which is reacted with an isoindoline, tetrahydroisoquinoline, or aminotetralin, to produce compounds of this invention (**Scheme 7**).



Scheme 7

Administration/Dosage/Formulations

5 The invention also encompasses pharmaceutical compositions and methods of their use. These pharmaceutical compositions may comprise an active ingredient (which can be one or more compounds of the invention, or pharmaceutically acceptable salts thereof) optionally in combination with one or more pharmaceutically acceptable agents. The compositions set forth herein can be used alone or in combination with additional compounds

10 to produce additive, complementary, or synergistic effects.

 The regimen of administration may affect what constitutes an effective amount. The therapeutic formulations may be administered to the subject either prior to or after the onset of a disease or disorder contemplated herein. Further, several divided dosages, as well as staggered dosages may be administered daily or sequentially, or the dose may be

15 continuously infused, or may be a bolus injection. Further, the dosages of the therapeutic formulations may be proportionally increased or decreased as indicated by the exigencies of the therapeutic or prophylactic situation.

 Administration of the compositions of the present invention to a patient, preferably a mammal, more preferably a human, may be carried out using known procedures, at dosages

20 and for periods of time effective to treat a disease or disorder contemplated herein. An

effective amount of the therapeutic compound necessary to achieve a therapeutic effect may vary according to factors such as the state of the disease or disorder in the patient; the age, sex, and weight of the patient; and the ability of the therapeutic compound to treat a disease or disorder contemplated herein. Dosage regimens may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation. A non-limiting example of an effective dose range for a therapeutic compound of the invention is from about 1 and 5,000 mg/kg of body weight/per day. One of ordinary skill in the art would be able to study the relevant factors and make the determination regarding the effective amount of the therapeutic compound without undue experimentation.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

In particular, the selected dosage level depends upon a variety of factors including the activity of the particular compound employed, the time of administration, the rate of excretion of the compound, the duration of the treatment, other drugs, compounds or materials used in combination with the compound, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

A medical doctor, *e.g.*, physician or veterinarian, having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect, and gradually increase the dosage until the desired effect is achieved.

In particular embodiments, it is especially advantageous to formulate the compound in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the patients to be treated; each unit containing a predetermined quantity of therapeutic compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical vehicle. The dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the therapeutic compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding/formulating such a

therapeutic compound for the treatment of a disease or disorder contemplated herein.

In certain embodiments, the compositions of the invention are formulated using one or more pharmaceutically acceptable excipients or carriers. In certain embodiments, the pharmaceutical compositions of the invention comprise a therapeutically effective amount of
5 a compound of the invention and a pharmaceutically acceptable carrier.

The carrier may be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity may be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the
10 required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms may be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it is preferable to include isotonic agents, for example, sugars, sodium chloride, or polyalcohols such as mannitol and sorbitol, in the composition. Prolonged absorption of the
15 injectable compositions may be brought about by including in the composition an agent that delays absorption, for example, aluminum monostearate or gelatin.

In certain embodiments, the compositions of the invention are administered to the patient in dosages that range from one to five times per day or more. In other embodiments, the compositions of the invention are administered to the patient in range of dosages that
20 include, but are not limited to, once every day, every two, days, every three days to once a week, and once every two weeks. It is readily apparent to one skilled in the art that the frequency of administration of the various combination compositions of the invention varies from individual to individual depending on many factors including, but not limited to, age, disease or disorder to be treated, gender, overall health, and other factors. Thus, the invention
25 should not be construed to be limited to any particular dosage regime and the precise dosage and composition to be administered to any patient is determined by the attending physical taking all other factors about the patient into account.

Compounds of the invention for administration may be in the range of from about 1 µg to about 10,000 mg, about 20 µg to about 9,500 mg, about 40 µg to about 9,000 mg, about
30 75 µg to about 8,500 mg, about 150 µg to about 7,500 mg, about 200 µg to about 7,000 mg, about 350 µg to about 6,000 mg, about 500 µg to about 5,000 mg, about 750 µg to about 4,000 mg, about 1 mg to about 3,000 mg, about 10 mg to about 2,500 mg, about 20 mg to about 2,000 mg, about 25 mg to about 1,500 mg, about 30 mg to about 1,000 mg, about 40 mg to about 900 mg, about 50 mg to about 800 mg, about 60 mg to about 750 mg, about 70

mg to about 600 mg, about 80 mg to about 500 mg, and any and all whole or partial increments there between.

In certain embodiments, the dose of a compound of the invention is from about 1 mg and about 2,500 mg. In other embodiments, a dose of a compound of the invention used in compositions described herein is less than about 10,000 mg, or less than about 8,000 mg, or less than about 6,000 mg, or less than about 5,000 mg, or less than about 3,000 mg, or less than about 2,000 mg, or less than about 1,000 mg, or less than about 500 mg, or less than about 200 mg, or less than about 50 mg. Similarly, in other embodiments, a dose of a second compound as described herein is less than about 1,000 mg, or less than about 800 mg, or less than about 600 mg, or less than about 500 mg, or less than about 400 mg, or less than about 300 mg, or less than about 200 mg, or less than about 100 mg, or less than about 50 mg, or less than about 40 mg, or less than about 30 mg, or less than about 25 mg, or less than about 20 mg, or less than about 15 mg, or less than about 10 mg, or less than about 5 mg, or less than about 2 mg, or less than about 1 mg, or less than about 0.5 mg, and any and all whole or partial increments thereof.

In certain embodiments, the present invention is directed to a packaged pharmaceutical composition comprising a container holding a therapeutically effective amount of a compound of the invention, alone or in combination with a second pharmaceutical agent; and instructions for using the compound to treat, prevent, or reduce one or more symptoms of a disease or disorder contemplated herein.

Formulations may be employed in admixtures with conventional excipients, *i.e.*, pharmaceutically acceptable organic or inorganic carrier substances suitable for oral, parenteral, nasal, intravenous, subcutaneous, enteral, or any other suitable mode of administration, known to the art. The pharmaceutical preparations may be sterilized and if desired mixed with auxiliary agents, *e.g.*, lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure buffers, coloring, flavoring and/or aromatic substances and the like. They may also be combined where desired with other active agents, *e.g.*, other analgesic agents.

Routes of administration of any of the compositions of the invention include oral, nasal, rectal, intravaginal, parenteral, buccal, sublingual or topical. The compounds for use in the invention may be formulated for administration by any suitable route, such as for oral or parenteral, for example, transdermal, transmucosal (*e.g.*, sublingual, lingual, (trans)buccal, (trans)urethral, vaginal (*e.g.*, trans- and perivaginally), (intra)nasal and (trans)rectal), intravesical, intrapulmonary, intraduodenal, intragastrical, intrathecal, subcutaneous,

intramuscular, intradermal, intra-arterial, intravenous, intrabronchial, inhalation, and topical administration.

Suitable compositions and dosage forms include, for example, tablets, capsules, caplets, pills, gel caps, troches, dispersions, suspensions, solutions, syrups, granules, beads, transdermal patches, gels, powders, pellets, magmas, lozenges, creams, pastes, plasters, lotions, discs, suppositories, liquid sprays for nasal or oral administration, dry powder or aerosolized formulations for inhalation, compositions and formulations for intravesical administration and the like. It should be understood that the formulations and compositions that would be useful in the present invention are not limited to the particular formulations and compositions that are described herein.

Oral Administration

For oral application, particularly suitable are tablets, dragees, liquids, drops, suppositories, or capsules, caplets and gelcaps. The compositions intended for oral use may be prepared according to any method known in the art and such compositions may contain one or more agents selected from the group consisting of inert, non-toxic pharmaceutically excipients that are suitable for the manufacture of tablets. Such excipients include, for example an inert diluent such as lactose; granulating and disintegrating agents such as cornstarch; binding agents such as starch; and lubricating agents such as magnesium stearate. The tablets may be uncoated or they may be coated by known techniques for elegance or to delay the release of the active ingredients. Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert diluent.

For oral administration, the compounds of the invention may be in the form of tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.*, polyvinylpyrrolidone, hydroxypropylcellulose or hydroxypropyl methylcellulose); fillers (*e.g.*, cornstarch, lactose, microcrystalline cellulose or calcium phosphate); lubricants (*e.g.*, magnesium stearate, talc, or silica); disintegrates (*e.g.*, sodium starch glycollate); or wetting agents (*e.g.*, sodium lauryl sulphate). If desired, the tablets may be coated using suitable methods and coating materials such as OPADRY™ film coating systems available from Colorcon, West Point, Pa. (*e.g.*, OPADRY™ OY Type, OYC Type, Organic Enteric OY-P Type, Aqueous Enteric OY-A Type, OY-PM Type and OPADRY™ White, 32K18400). Liquid preparation for oral administration may be in the form of solutions, syrups or suspensions. The liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (*e.g.*, sorbitol syrup, methyl cellulose or hydrogenated edible fats); emulsifying agent (*e.g.*, lecithin or

acacia); non-aqueous vehicles (*e.g.*, almond oil, oily esters or ethyl alcohol); and preservatives (*e.g.*, methyl or propyl p-hydroxy benzoates or sorbic acid).

Granulating techniques are well known in the pharmaceutical art for modifying starting powders or other particulate materials of an active ingredient. The powders are typically mixed with a binder material into larger permanent free-flowing agglomerates or granules referred to as a “granulation.” For example, solvent-using “wet” granulation processes are generally characterized in that the powders are combined with a binder material and moistened with water or an organic solvent under conditions resulting in the formation of a wet granulated mass from which the solvent must then be evaporated.

Melt granulation generally consists in the use of materials that are solid or semi-solid at room temperature (*i.e.* having a relatively low softening or melting point range) to promote granulation of powdered or other materials, essentially in the absence of added water or other liquid solvents. The low melting solids, when heated to a temperature in the melting point range, liquefy to act as a binder or granulating medium. The liquefied solid spreads itself over the surface of powdered materials with which it is contacted, and on cooling, forms a solid granulated mass in which the initial materials are bound together. The resulting melt granulation may then be provided to a tablet press or be encapsulated for preparing the oral dosage form. Melt granulation improves the dissolution rate and bioavailability of an active (*i.e.* drug) by forming a solid dispersion or solid solution.

U.S. Patent No. 5,169,645 discloses directly compressible wax-containing granules having improved flow properties. The granules are obtained when waxes are admixed in the melt with certain flow improving additives, followed by cooling and granulation of the admixture. In certain embodiments, only the wax itself melts in the melt combination of the wax(es) and additives(s), and in other cases both the wax(es) and the additives(s) melt.

The present invention also includes a multi-layer tablet comprising a layer providing for the delayed release of one or more compounds of the invention, and a further layer providing for the immediate release of a medication for treatment of diseases or disorders. Using a wax/pH-sensitive polymer mix, a gastric insoluble composition may be obtained in which the active ingredient is entrapped, ensuring its delayed release.

Parenteral Administration

For parenteral administration, the compounds of the invention may be formulated for injection or infusion, for example, intravenous, intramuscular or subcutaneous injection or infusion, or for administration in a bolus dose and/or continuous infusion. Suspensions, solutions or emulsions in an oily or aqueous vehicle, optionally containing other formulatory

agents such as suspending, stabilizing and/or dispersing agents may be used.

Additional Administration Forms

Additional dosage forms of this invention include dosage forms as described in U.S. Patents Nos. 6,340,475; 6,488,962; 6,451,808; 5,972,389; 5,582,837; and 5,007,790.

5 Additional dosage forms of this invention also include dosage forms as described in U.S. Patent Applications Nos. 20030147952; 20030104062; 20030104053; 20030044466; 20030039688; and 20020051820. Additional dosage forms of this invention also include dosage forms as described in PCT Applications Nos. WO 03/35041; WO 03/35040; WO 03/35029; WO 03/35177; WO 03/35039; WO 02/96404; WO 02/32416; WO 01/97783; WO 10 01/56544; WO 01/32217; WO 98/55107; WO 98/11879; WO 97/47285; WO 93/18755; and WO 90/11757.

Controlled Release Formulations and Drug Delivery Systems

In certain embodiments, the formulations of the present invention may be, but are not limited to, short-term, rapid-offset, as well as controlled, for example, sustained release, 15 delayed release and pulsatile release formulations.

The term sustained release is used in its conventional sense to refer to a drug formulation that provides for gradual release of a drug over an extended period of time, and that may, although not necessarily, result in substantially constant blood levels of a drug over an extended time period. The period of time may be as long as a month or more and should 20 be a release that is longer than the same amount of agent administered in bolus form.

For sustained release, the compounds may be formulated with a suitable polymer or hydrophobic material that provides sustained release properties to the compounds. As such, the compounds for use the method of the invention may be administered in the form of microparticles, for example, by injection or in the form of wafers or discs by implantation.

25 In certain embodiments, the compounds of the invention are administered to a patient, alone or in combination with another pharmaceutical agent, using a sustained release formulation.

The term delayed release is used herein in its conventional sense to refer to a drug formulation that provides for an initial release of the drug after some delay following drug 30 administration and that may, although not necessarily, include a delay of from about 10 minutes up to about 12 hours.

The term pulsatile release is used herein in its conventional sense to refer to a drug formulation that provides release of the drug in such a way as to produce pulsed plasma profiles of the drug after drug administration.

The term immediate release is used in its conventional sense to refer to a drug formulation that provides for release of the drug immediately after drug administration.

As used herein, short-term refers to any period of time up to and including about 8 hours, about 7 hours, about 6 hours, about 5 hours, about 4 hours, about 3 hours, about 2 hours, about 1 hour, about 40 minutes, about 20 minutes, or about 10 minutes and any or all whole or partial increments thereof after drug administration after drug administration.

As used herein, rapid-offset refers to any period of time up to and including about 8 hours, about 7 hours, about 6 hours, about 5 hours, about 4 hours, about 3 hours, about 2 hours, about 1 hour, about 40 minutes, about 20 minutes, or about 10 minutes, and any and all whole or partial increments thereof after drug administration.

Dosing

The therapeutically effective amount or dose of a compound of the present invention depends on the age, sex and weight of the patient, the current medical condition of the patient and the progression of a disease or disorder contemplated herein in the patient being treated. The skilled artisan is able to determine appropriate dosages depending on these and other factors.

A suitable dose of a compound of the present invention may be in the range of from about 0.01 mg to about 5,000 mg per day, such as from about 0.1 mg to about 1,000 mg, for example, from about 1 mg to about 500 mg, such as about 5 mg to about 250 mg per day. The dose may be administered in a single dosage or in multiple dosages, for example from 1 to 4 or more times per day. When multiple dosages are used, the amount of each dosage may be the same or different. For example, a dose of 1 mg per day may be administered as two 0.5 mg doses, with about a 12-hour interval between doses.

It is understood that the amount of compound dosed per day may be administered, in non-limiting examples, every day, every other day, every 2 days, every 3 days, every 4 days, or every 5 days. For example, with every other day administration, a 5 mg per day dose may be initiated on Monday with a first subsequent 5 mg per day dose administered on Wednesday, a second subsequent 5 mg per day dose administered on Friday, and so on.

In the case wherein the patient's status does improve, upon the doctor's discretion the administration of the inhibitor of the invention is optionally given continuously; alternatively, the dose of drug being administered is temporarily reduced or temporarily suspended for a certain length of time (*i.e.*, a "drug holiday"). The length of the drug holiday optionally varies between 2 days and 1 year, including by way of example only, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, 35 days, 50 days, 70

days, 100 days, 120 days, 150 days, 180 days, 200 days, 250 days, 280 days, 300 days, 320 days, 350 days, or 365 days. The dose reduction during a drug holiday includes from 10%-100%, including, by way of example only, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.

5 Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, the dosage or the frequency of administration, or both, is reduced, as a function of the viral load, to a level at which the improved disease is retained. In certain embodiments, patients require intermittent treatment on a long-term basis upon any recurrence of symptoms and/or infection.

10 The compounds for use in the method of the invention may be formulated in unit dosage form. The term "unit dosage form" refers to physically discrete units suitable as unitary dosage for patients undergoing treatment, with each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, optionally in association with a suitable pharmaceutical carrier. The unit dosage form may be for a single
15 daily dose or one of multiple daily doses (*e.g.*, about 1 to 4 or more times per day). When multiple daily doses are used, the unit dosage form may be the same or different for each dose.

 Toxicity and therapeutic efficacy of such therapeutic regimens are optionally determined in cell cultures or experimental animals, including, but not limited to, the
20 determination of the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between the toxic and therapeutic effects is the therapeutic index, which is expressed as the ratio between LD₅₀ and ED₅₀. The data obtained from cell culture assays and animal studies are optionally used in formulating a range of dosage for use in human. The dosage of such compounds lies
25 preferably within a range of circulating concentrations that include the ED₅₀ with minimal toxicity. The dosage optionally varies within this range depending upon the dosage form employed and the route of administration utilized.

 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures, embodiments,
30 claims, and examples described herein. Such equivalents were considered to be within the scope of this invention and covered by the claims appended hereto. For example, it should be understood, that modifications in reaction conditions, including but not limited to reaction times, reaction size/volume, and experimental reagents, such as solvents, catalysts, pressures, atmospheric conditions, *e.g.*, nitrogen atmosphere, and reducing/oxidizing agents, with art-

recognized alternatives and using no more than routine experimentation, are within the scope of the present application.

It is to be understood that wherever values and ranges are provided herein, all values and ranges encompassed by these values and ranges, are meant to be encompassed within the scope of the present invention. Moreover, all values that fall within these ranges, as well as the upper or lower limits of a range of values, are also contemplated by the present application.

The following examples further illustrate aspects of the present invention. However, they are in no way a limitation of the teachings or disclosure of the present invention as set forth herein.

EXPERIMENTAL EXAMPLES

The invention is now described with reference to the following Examples. These Examples are provided for the purpose of illustration only and the invention should in no way be construed as being limited to these Examples, but rather should be construed to encompass any and all variations which become evident as a result of the teaching provided herein.

Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. The following working examples therefore, specifically point out the preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

Example 1: G-Protein Assay Measuring cAMP


G-protein signaling was measured via second messenger cAMP modulation. Detection of cAMP modulation was accomplished in the PathHunter human OPRM1 (μ , MOR) Arrestin CHO-K1 cell line using the Dynamic2 cAMP Kit from Cisbio. The E_{MAX} values of the chemiluminescent signal are all normalized to DAMGO, which is defined as 100%.

Example 2: β -Arrestin Assay

The Discoverx PathHunter® β -Arrestin assay is used to measure β -arrestin-2 activity. The technology is based on Enzyme Fragment Complementation (EFC) with β -galactosidase

(β -Gal) as the reporter. The enzyme is split into two inactive complementary portions (EA for Enzyme Acceptor, and ED for Enzyme Donor) expressed as fusion proteins in the cell. EA is fused to β -arrestin-2, and ED is fused to the C-terminus of the GPCR of interest. When a ligand binds and the GPCR is activated, β -arrestin-2 is recruited to the receptor resulting in an ED/EA complementation, restoring β -Gal activity, which is measured using chemiluminescence. For this assay, the PathHunter human OPRM1 (μ , MOR) Arrestin CHO-K1 cell line was used. The E_{MAX} values of the chemiluminescent signal were all normalized to DAMGO, which is defined as 100%.

Example 3: N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-isoindoline-2-carboxamide (1) and corresponding hydrochloride salt (1a)




(a) N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (1)

A mixture of N^1,N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine (150 mg, 0.88 mmol) and N,N -diisopropylethylamine (180 μ L, 1.05 mmol) in CH_2Cl_2 (10 mL) was purged with argon and cooled to 0 $^{\circ}C$. A solution of 4-nitrophenylchloroformate (213 mg, 1.05 mmol) in CH_2Cl_2 (5 mL) was added and resulting solution was stirred at 0 $^{\circ}C$ for 1 h. After this time, isoindoline hydrochloride (137 mg, 0.88 mmol) was added followed by N,N -diisopropylethylamine (300 μ L, 1.77 mmol), and the resulting mixture was stirred for 16 h at room temperature. A saturated $NaHCO_3$ solution (10 mL) was added and the resulting suspension was extracted with CH_2Cl_2 (3 x 10 mL). The combined organic extracts were washed with water (30 mL) and then dried over solid anhydrous $MgSO_4$. After filtration, the volatiles were removed, and the residue was purified by flash chromatography using eluent from CH_2Cl_2 to $CH_2Cl_2/MeOH$ (10:1) to give N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl) isoindoline-2-carboxamide (**1**) as an oil (170 mg, 61% yield). 300 MHz 1H -NMR ($CDCl_3$, ppm): 7.31 (dd, $J=5.0, 3.0$ Hz, 1H) 7.28-7.23 (m, 4H) 7.10 (dd, $J=3.0, 1.3$ Hz, 1H) 7.01 (dd, $J=5.0, 1.3$ Hz, 1H) 5.01-4.89 (m, 1H) 4.73-4.56 (m, 4H) 3.74-3.63 (m, 2H) 3.61-3.49 (m, 1H) 2.21 (s, 6H). ESI-MS (m/z): 316 [$M+H$] $^{+}$.

(b) N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride (1a)

A 2M HCl/diethyl ether (260 μ L, 0.51 mmol) was added to the solution of N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (**1**) (160 mg, 0.51 mmol) in diethyl ether/MeOH (45 mL/0.5 mL). The mixture was stirred for 2 h at ambient temperature. The resultant precipitate were filtered and dried over P₂O₅ *in vacuo* at 65 °C for 24 h to give N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride (**1a**) (140 mg, 78% yield). 400 MHz ¹H-NMR (CD₃OD, ppm): 7.85-7.79 (m, 1H) 7.66 (dd, J=5.0, 2.9 Hz, 1H) 7.35-7.28 (m, 5H) 4.78-4.65 (m, 5H) 4.13 (dd, J=15.1, 8.9 Hz, 1H) 3.65 (dd, J=15.1, 4.8 Hz, 1H) 2.95-2.70 (m, 6H). ESI-MS (m/z): 316 [M+H]⁺; melting point: 145-150 °C (dec.).


Example 4: N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (2) and corresponding hydrochloride salt (2a)



A mixture of N¹,N¹-dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine (300 mg, 1.76 mmol) and N,N-diisopropylethylamine (360 μ L, 2.11 mmol) in CH₂Cl₂ (10 mL) was purged with argon and cooled to 0 °C. A solution of 4-nitrophenylchloroformate (**2**) (426 mg, 2.11 mmol) in CH₂Cl₂ (5 mL) was added and resulting solution was stirred at 0 °C for 1 h. After this time, 5-fluoroisoindoline hydrochloride (306 mg, 1.76 mmol) was added, followed by N,N-diisopropylethylamine (610 μ L, 3.52 mmol), and the resulting mixture was stirred for 16 h at room temperature. A saturated NaHCO₃ solution (10 mL) was added and the resulting suspension was extracted with CH₂Cl₂ (3 x 10 mL). The combined organic extracts were washed with water (30 mL) and dried over solid anhydrous MgSO₄. After filtration, the volatiles were removed, and the residue was purified by flash chromatography using eluent from CH₂Cl₂ to CH₂Cl₂/MeOH (10:1) to give crude product as a colorless foam. This material was solidified with dry ethyl ether to give pure *rac*-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (**2**) as gray powder (300 mg, 51% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.32 (dd, J=5.0, 3.0 Hz, 1H) 7.24-7.17 (m, 1H) 7.11 (dd, J=3.0, 1.3 Hz, 1H) 7.02 (dd, J=5.0, 1.3 Hz, 1H) 6.99-6.92 (m, 2H) 5.03-4.87 (m, 1H) 4.72-4.55 (m, 4H) 3.77-3.49 (m, 3H) 2.21 (s, 6H). ESI-MS (m/z): 334 [M+H]⁺.

Example 5: (-)-ent-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (2-ent-(-)) and corresponding hydrochloride salt (2a-ent-(-))

Example 6: (+)-ent-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (2-ent-(+)) and corresponding hydrochloride salt (2a-ent-(+))



(a) (-)-N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (**2-ent-(-)**) and

(+)-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (**2-ent-(+)**)

rac-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (**2**) was separated into its enantiomers on a chiral stationary phase chromatography using (Daicel Chiralpak® AY-H 250 mm x 10 mm) column and Hex/EtOH (7:3) as mobile phase on Shimadzu LC-8A to obtain (-)-N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (**2-ent-(-)**) and (+)-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (**2-ent-(+)**). *R_t*(-) enantiomer= 58 min, *R_t*(+) enantiomer= 82 min at 1mL/min flow rate.


(b) (-)-N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide hydrochloride (**2a-ent-(-)**)

A solution of 2M HCl/diethyl ether (75 μ L, 0.15 mmol) was added to the solution of (-)-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide (**2-ent-(-)**) (50 mg, 0.15 mmol) in diethyl ether/MeOH (20 mL/0.5 mL). The mixture was stirred for 2 h at ambient temperature and the resultant precipitate was filtered and dried over P₂O₅ *in vacuo* at 65 °C for 24 h to give (-)-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide hydrochloride (**2a-ent-(-)**) (45 mg, 81% yield). 400 MHz ¹H-NMR (CD₃OD, ppm): 7.82 (dd, *J*=2.9, 1.4 Hz, 1H) 7.66 (dd, *J*=5.0, 2.9 Hz, 1H) 7.36-7.30 (m, 2H) 7.12-7.02 (m, 2H) 4.74 (dd, *J*=8.8, 4.7 Hz, 1H) 4.78-4.62 (m, 4H) 4.12 (dd, *J*=15.1, 8.8 Hz, 1H) 3.65 (dd, *J*=15.1, 4.7 Hz, 1H) 2.88 (s, 3H) 2.77 (s, 3H). ESI-MS (*m/z*): 334 [M+H]⁺; melting point: 150-155 °C (dec.).

(c) (+)-N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisoindoline-2-carboxamide hydrochloride (**2a-ent-(+)**)

A solution of 2M HCl/diethyl ether (60 μ L, 0.12 mmol) was added to the solution of (+)-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisindoline-2-carboxamide (**2a-ent-(+)**) (40 mg, 0.12 mmol) in diethyl ether/MeOH (15 mL/0.5 mL). The mixture was stirred for 2 h at ambient temperature and the resultant precipitate were filtered and dried over P₂O₅ *in vacuo* at 65 °C for 24 h to give (+)-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-fluoroisindoline-2-carboxamide hydrochloride (**2a-ent-(+)**) (35 mg, 79% yield). 400 MHz ¹H-NMR (CD₃OD, ppm): 7.81 (dd, J=2.9, 1.4 Hz, 1H) 7.66 (dd, J=5.0, 2.9 Hz, 1H) 7.35-7.29 (m, 2H) 7.12-7.02 (m, 2H) 4.74 (dd, J=8.8, 4.7 Hz, 1H) 4.78-4.62 (m, 4H) 4.12 (dd, J=15.1, 8.8 Hz, 1H) 3.65 (dd, J=15.1, 4.7 Hz, 1H) 2.88 (s, 3H) 2.78 (s, 3H). ESI-MS (m/z): 334 [M+H]⁺; melting point: 150-155 °C (dec.).

Example 7: N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-4-fluoroisindoline-2-carboxamide (3) and corresponding hydrochloride salt (3a)




(a) N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-4-fluoroisindoline-2-carboxamide (**3**)

4-Fluoroisindoline hydrochloride (183 mg, 1.05 mmol) and N¹,N¹-dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH₂Cl₂ using the procedure described for compound (**1**) to afford N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-4-fluoroisindoline-2-carboxamide (**3**) (120 mg, 41% yield). ESI-MS (m/z): 334 [M+H]⁺

(b) N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-4-fluoroisindoline-2-carboxamide hydrochloride (**3a**)

N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-4-fluoroisindoline-2-carboxamide (**3**) (125 mg, 0.37 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (20 mL/0.5 mL) using procedure described for compound (**1a**) to produce N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-4-fluoroisindoline-2-carboxamide hydrochloride (**3a**). 400 MHz ¹H-NMR (CD₃OD, ppm): 7.82 (dd, J=2.9, 1.4 Hz, 1H) 7.66 (dd, J=5.0, 2.9 Hz, 1H) 7.39-7.30 (m, 1H) 7.32 (dd, J=5.0, 1.4 Hz, 1H) 7.18-7.12 (m, 1H) 7.07-7.00 (m, 1H) 4.82-4.65 (m, 5H) 4.12 (dd, J=15.1, 9.0 Hz, 1H) 3.65 (dd, J=15.1, 4.9 Hz, 1H) 2.83 (s, 6H). ESI-MS (m/z): 334 [M+H]⁺; melting point: 150-155 °C (dec.).

Example 8: 5-chloro-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (4) and corresponding hydrochloride salt 4(a)




(a) 5-Chloro-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (4)

5 5-Chloroisoindoline hydrochloride (167 mg, 0.88 mmol) and N¹,N¹-dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH₂Cl₂ using the procedure described for compound (1) to afford 5-chloro-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (4) (110 mg, 36% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.33 (dd, J=5.0, 2.9 Hz, 1H) 7.28-7.23 (m, 2H) 7.21-7.17 (m, 1H) 7.14-7.10 (m, 1H) 7.02 (dd, J=5.0, 1.3 Hz, 1H) 5.05-4.91 (m, 1H) 4.72-4.57 (m, 4H) 3.78-3.63 (m, 2H) 3.62-3.52 (m, 1H) 2.24 (s, 6H). ESI-MS (m/z): 350, 352 [M+H]⁺

(b) 5-Chloro-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride (4a)

15 5-Chloro-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (4) (90 mg, 0.26 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (20 mL/3 mL) using procedure described for compound (1a) to produce 5-chloro-N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride (4a) (85 mg, 85% yield). 400 MHz ¹H-NMR (CD₃OD, ppm): 7.85-7.79 (m, 1H) 7.66 (dd, J=5.0, 2.9 Hz, 1H) 7.38-7.35 (m, 1H) 7.34-7.29 (m, 3H) 4.76-4.62 (m, 5H) 4.11 (dd, J=15.0, 8.7 Hz, 1H) 3.65 (dd, J=15.0, 4.9 Hz, 1H) 2.82 (s, 6H). ESI-MS (m/z): 350, 352 [M+H]⁺; melting point: 210-215 °C (dec.).

Example 9: N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methoxyisoindoline-2-carboxamide (5) and corresponding hydrochloride salt (5a)



(a) N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methoxyisoindoline-2-carboxamide (5)


5-Methoxyisoindoline hydrochloride (163 mg, 0.88 mmol) and N¹,N¹-dimethyl-1-

(thiophen-3-yl)ethane-1,2-diamine were reacted in CH_2Cl_2 using the procedure described for compound (1) to afford N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methoxyisoindoline-2-carboxamide (5) (190 mg, 62% yield). 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.32 (dd, $J=5.0$, 2.9 Hz, 1H) 7.15 (d, $J=8.3$ Hz, 1H) 7.12 (dd, $J=2.9$, 1.3 Hz, 1H) 7.02 (dd, $J=5.0$, 1.3 Hz, 1H) 6.83 (dd, $J=8.3$, 2.5 Hz, 1H) 6.79 (d, $J=2.5$ Hz, 1H) 5.03-4.94 (m, 1H) 4.71-4.53 (m, 4H) 3.80 (s, 3H) 3.75-3.70 (m, 1H) 3.67 (dd, $J=8.7$, 3.8 Hz, 1H) 3.59-3.52 (m, 1H) 2.22 (s, 6H). ESI-MS (m/z): 346 $[\text{M}+\text{H}]^+$

(b) N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methoxyisoindoline-2-carboxamide hydrochloride (5a)

N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methoxyisoindoline-2-carboxamide (5) (135 mg, 0.39 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (25 mL/5 mL) using procedure described for compound (1a) to produce N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methoxyisoindoline-2-carboxamide hydrochloride (5a) (102 mg, 68% yield). 400 MHz ^1H -NMR (CD_3OD , ppm): 7.82 (dd, $J=2.9$, 1.4 Hz, 1H) 7.66 (dd, $J=5.1$, 2.9 Hz, 1H) 7.32 (dd, $J=5.1$, 1.4 Hz, 1H) 7.23-7.18 (m, 1H) 6.90-6.85 (m, 2H) 4.75 (dd, $J=9.0$, 4.8 Hz, 1H) 4.72-4.57 (m, 4H) 4.12 (dd, $J=15.1$, 9.0 Hz, 1H) 3.79 (s, 3H) 3.65 (dd, $J=15.1$, 4.8 Hz, 1H) 2.88 (s, 3H) 2.78 (s, 3H). ESI-MS (m/z): 346 $[\text{M}+\text{H}]^+$; melting point: 210-215 $^\circ\text{C}$ (dec.).

Example 10: N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5,6-difluoroisoindoline-2-carboxamide (6) and corresponding hydrochloride salt (6a)




(a) N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5,6-difluoroisoindoline-2-carboxamide (6)

5,6-Difluoroisoindoline hydrochloride (150 mg, 0.78 mmol) and N^1, N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH_2Cl_2 using the procedure described for compound (1) to afford N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5,6-difluoroisoindoline-2-carboxamide (6) (90 mg, 33% yield). 400 MHz ^1H -NMR (CDCl_3 , ppm): 7.32 (dd, $J=4.9$, 2.9 Hz, 1H) 7.11 (dd, $J=2.9$, 1.2 Hz, 1H) 7.07 (dd, $J=8.5$, 8.5 Hz, 2H) 7.01 (dd, $J=5.0$, 1.2 Hz, 1H) 4.98-4.91 (m, 1H) 4.66-4.57 (m, 4H) 3.73-3.61 (m, 2H) 3.59-3.52 (m, 1H) 2.20 (s, 6H). ESI-MS (m/z): 352 $[\text{M}+\text{H}]^+$.

(b) *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5,6-difluoroisoindoline-2-carboxamide hydrochloride (**6a**)

5,6-Difluoro-1,3-dihydro-isoindole-2-carboxylic acid (2-dimethylamino-2-thiophen-3-yl-ethyl)-amide (**6**) (85 mg, 0.24 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using procedure described for compound (**1a**) to produce *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5,6-difluoroisoindoline-2-carboxamide hydrochloride (**6a**) (80 mg, 85% yield). 400 MHz ¹H-NMR (CD₃OD, ppm): 7.80 (dd, J=2.9, 1.2 Hz, 1H) 7.66 (dd, J = 5.0, 2.9 Hz, 1H) 7.32 (dd, J=5.0, 1.2 Hz, 1H) 7.26 (dd, J=8.8, 8.8 Hz, 2H) 4.73 (dd, J=8.8, 4.9 Hz, 1H) 4.67 (s, 4H) 4.11 (dd, J=15.1, 8.8 Hz, 1H) 3.65 (dd, J=15.1, 4.9 Hz, 1H) 2.82 (s, 6H). ESI-MS (m/z): 352 [M+H]⁺; melting point: 210-212 °C.

Example 11: *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide (7**) and corresponding hydrochloride salt (**7a**)**



(a) *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-dihydroquinoline-1(2H)-carboxamide (**7**)


6-Fluoro-1,2,3,4-tetrahydroquinoline hydrochloride (165 mg, 0.88 mmol) and *N*¹,*N*¹-dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH₂Cl₂ using the procedure described for compound (**1**) to afford *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-dihydroquinoline-1(2H)-carboxamide (**7**) (120 mg, 39% yield). 400 MHz ¹H-NMR (CDCl₃, ppm): 7.31 (dd, J=5.0, 2.9 Hz, 1H) 7.09-7.04 (m, 1H) 7.00-6.92 (m, 2H) 6.83 (dd, J=8.9, 2.9 Hz, 1H) 6.75 (ddd, J=8.5, 8.5, 3.0 Hz, 1H) 5.43 (s, 1H) 3.77-3.61 (m, 4H) 3.53-3.46 (m, 1H, 2.70 (t, J=6.7 Hz, 2H) 2.17 (s, 6H) 1.93-1.84 (m, 2H). ESI-MS (m/z): 348 [M+H]⁺.

(b) *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide hydrochloride (**7a**)

N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-dihydroquinoline-1(2H)-carboxamide (**7**) (120 mg, 0.35 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using procedure described for compound (**1a**) to produce *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide hydrochloride (**7a**) (133 mg,

100% yield). 400 MHz ^1H -NMR (DMSO- d_6 , ppm): 10.48 (s, 1H) 7.86 (dd, $J=2.9, 0.9$ Hz, 1H) 7.71 (dd, $J=5.0, 2.9$ Hz, 1H) 7.37 (dd, $J=5.0, 0.9$ Hz, 1H) 7.27 (dd, $J=9.0, 5.4$ Hz, 1H) 7.01 (t, $J=5.4$ Hz, 1H) 6.95 (dd, $J=9.3, 3.0$ Hz, 1H) 6.87 (ddd, $J=8.7, 8.7, 3.0$ Hz, 1H) 4.74-4.65 (m, 1H) 3.98-3.86 (m, 1H) 3.65-3.48 (m, 3H) 2.74 (d, $J=4.4$ Hz, 3H) 2.66 (t, $J=6.3$ Hz, 2H) 2.59 (d, $J=4.4$ Hz, 3H) 1.82-1.72 (m, 2H). ESI-MS (m/z): 348 $[\text{M}+\text{H}]^+$.

Example 12: N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-7-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide (8) and corresponding hydrochloride salt (8a)



10 (a) *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-7-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide (8)


7-Fluoro-1,2,3,4-tetrahydroquinoline hydrochloride (165 mg, 0.88 mmol) and N^1, N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH_2Cl_2 using the procedure described for compound (1) to afford *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-7-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide (8) (70 mg, 23% yield). 300 MHz ^1H -NMR (CDCl $_3$, ppm): 7.31 (dd, $J=5.0, 2.9$ Hz, 1H) 7.09 (dd, $J=2.9, 1.3$ Hz, 1H) 7.06-7.01 (m, 1H) 7.00-6.93 (m, 2H) 6.69 (ddd, $J=8.3, 8.3, 2.6$ Hz, 1H) 5.75-4.64 (m, 1H) 3.75-3.62 (m, 4H) 3.59-3.51 (m, 1H) 2.71-2.65 (m, 2H) 2.18 (s, 6H) 1.88 (quintet, $J=6.4$ Hz, 2H). ESI-MS (m/z): 348 $[\text{M}+\text{H}]^+$.

20 (b) *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-7-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide (8a)

N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-7-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide (8) (52 mg, 0.15 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using the procedure described for compound (1a) to *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-7-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide (8a) (45 mg, 78% yield). 400 MHz ^1H -NMR (DMSO- d_6 , ppm): 10.15 (s, 1H) 7.88 (s, 1H) 7.71 (dd, $J=4.8, 2.9$ Hz, 1H) 7.36 (d, $J=4.8$ Hz, 1H) 7.30 (dd, $J=12.3, 2.6$ Hz, 1H) 7.25-7.17 (m, 1H) 7.13-7.06 (m, 1H) 6.74 (ddd, $J=8.4, 8.4, 2.7$ Hz, 1H) 4.74 (s, 1H) 4.02-3.86 (m, 1H) 3.65-3.48 (m, 3H) 2.80-2.70 (m, 3H) 2.69-2.59 (m, 5H) 1.84-1.75 (m, 2H). ESI-MS (m/z): 348 $[\text{M}+\text{H}]^+$.

30

Example 13: N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-dihydroquinoline-1(2H)-carboxamide (9) and corresponding hydrochloride salt (9a)




(a) *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-dihydroquinoline-1(2H)-carboxamide
5 (9)

1,2,3,4-Tetrahydroquinoline (132 μ L, 1.06 mmol) and N^1,N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH_2Cl_2 using procedure described for compound (1) to afford *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-dihydroquinoline-1(2H)-carboxamide (9) as an orange oil (126 mg, 36% yield). 400 MHz ^1H -NMR (CDCl_3 , ppm):
10 7.31 (dd, $J=5.0, 2.9$ Hz, 1H) 7.11 (d, $J=7.0$ Hz, 1H) 7.09-7.05 (m, 1H) 7.04 (dd, $J=7.0, 1.8$ Hz, 1H) 7.01-6.95 (m, 3H) 5.57 (s, 1H) 3.82-3.61 (m, 4H) 3.54-3.45 (m, 1H), 2.76-2.68 (m, 2H), 2.17 (s, 6H), 1.94-1.86 (m, 2H). ESI-MS (m/z): 330 $[\text{M}+\text{H}]^+$.

(b) *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-dihydroquinoline-1(2H)-carboxamide hydrochloride (9a)

15 *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-dihydroquinoline-1(2H)-carboxamide (9) (120 mg, 0.36 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using procedure described for compound (1a) to produce *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3,4-dihydroquinoline-1(2H)-carboxamide hydrochloride (9a) (120 mg, 90% yield). 400 MHz ^1H -NMR (CD_3OD , ppm): 7.79 (dd, $J=2.9, 1.3$ Hz, 1H) 7.67 (dd, $J=5.0, 2.9$
20 Hz, 1H) 7.30 (dd, $J=5.0, 1.3$ Hz 1H) 7.16-7.12 (m, 1H) 7.12-7.05 (m, 2H) 7.05-7.00 (m, 1H) 4.74 (t, $J=7.1$ Hz, 1H) 4.09 (dd, $J=14.5, 7.6$ Hz, 1H) 3.72 - 3.55 (m, 3H) 2.83 (s, 6H) 2.72 (t, $J=6.5$ Hz, 2H) 1.91 (quintet, $J=6.5$ Hz, 2H). ESI-MS (m/z): 330 $[\text{M}+\text{H}]^+$.

Example 14: 1-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydro naphthalen-2-yl)urea (10) and corresponding hydrochloride salt (10a)




(a) *[1-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea (10)*

1,2,3,4-Tetrahydronaphthalen-2-amine (173 mg, 1.17 mmol) and N¹,N¹-dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH₂Cl₂ using procedure described for compound (1) to afford [1-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea (10) (135 mg, 33% yield). 400 MHz ¹H-NMR (CDCl₃, ppm): 7.30 (dd, J=5.0, 2.9 Hz, 1H) 7.16-7.03 (m, 5H) 7.00-6.95 (m, 1H) 4.90-4.81 (m, 1H) 4.77-4.62 (m, 1H) 4.15-4.01 (m, 1H) 3.67 (dd, J=8.8, 5.6 Hz, 1H) 3.62-3.52 (m, 1H) 3.48-3.39 (m, 1H) 3.12 (dd, J=16.4, 5.6 Hz, 1H) 2.96-2.80 (m, 2H) 2.62 (dd, J=16.4, 7.7 Hz, 1H) 2.14 (s, 6H) 2.10-2.00 (m, 1H) 1.83-1.70 (m, 1H). ESI-MS (m/z): 344 [M+H]⁺.

(b) *[1-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea hydrochloride (10a)*

[1-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea (10) (120 mg, 0.35 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH using the procedure described for compound (1a) to produce [1-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea hydrochloride (10a) (105 mg, 79% yield). 400 MHz ¹H-NMR (CD₃OD, ppm): 7.77 (dd, J=2.9, 1.3 Hz, 1H) 7.65 (dd, J=5.1, 2.9 Hz, 1H) 7.29 (dd, J=5.1, 1.3 Hz, 1H) 7.09-7.00 (m, 4H) 4.68-4.62 (m, 1H) 4.06-3.92 (m, 2H) 3.66-3.58 (m, 1H) 3.04 (dd, J=16.2, 5.1 Hz, 1H) 2.96-2.82 (m, 6H) 2.73 (s, 3H) 2.64 (dd, J=16.2, 8.6 Hz, 1H) 2.09-1.98 (m, 1H) 1.78-1.66 (m, 1H). ESI-MS (m/z): 344 [M+H]⁺; melting point: 151-153 °C.

Example 15: 1-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-1-yl)urea (11) and corresponding hydrochloride salt (11a)




(a) 1-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-1-yl)urea (11)

1,2,3,4-Tetrahydronaphthalen-1-amine (173 mg, 0.94 mmol) and N¹,N¹-dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH₂Cl₂ using the procedure described for compound (1) to afford 1-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-1-yl)urea (11) (155 mg, 48% yield). 400 MHz ¹H-NMR (CDCl₃, ppm): 7.36-7.31 (m, 1H) 7.30 (dd, J=5.0, 2.9 Hz, 1H) 7.18-7.12 (m, 2H) 7.10-7.04 (m, 2H) 6.97 (dd, J=5.0, 1.3 Hz, 1H) 5.00-4.91 (m, 1H) 4.91-4.77 (m, 1H) 3.71-3.64 (m, 1H) 3.63-3.54 (m, 1H) 3.49-3.40 (m, 1H) 2.85-2.67 (m, 2H) 2.13 (s, 3H) 2.12 (s, 3H) 2.07-1.94 (m, 1H) 1.88-1.67 (m, 4H). ESI-MS (m/z): 344 [M+H]⁺.

(b) 1-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-1-yl)urea hydrochloride (11a)

1-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-1-yl)urea (11) (141 mg, 0.41 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH using the procedure described for compound (1) to produce 1-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-3-(1,2,3,4-tetrahydronaphthalen-1-yl)urea hydrochloride (11a) (115 mg, 74% yield). 400 MHz ¹H NMR (CD₃OD, ppm): 7.80-7.77 (m, 1H) 7.68-7.64 (m, 1H) 7.32-7.29 (m, 1H) 7.24-7.20 (m, 1H) 7.15-7.09 (m, 2H) 7.09-7.04 (m, 1H) 5.92-4.82 (m, 2H) 4.69 (ddd, J=10.2, 7.5, 6.2 Hz, 1H) 4.05 (ddd, J=14.8, 7.5, 1.3 Hz, 1H) 3.67 (ddd, J=14.8, 8.3, 6.2 Hz, 1H) 3.01-2.65 (m, 8H) 2.07-1.95 (m, 1H) 1.94-1.70 (m, 3H). ESI-MS (m/z): 344 [M+H]⁺; melting point: 179-181 °C.

Example 16: N-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methylisoindoline-2-carboxamide hydrochloride (12) and corresponding hydrochloride salt (12a)




(a) *N*-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methylisoindoline-2-carboxamide (**12**)

5-Methylisoindoline (140 mg, 1.06 mmol) and N^1,N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH_2Cl_2 using the procedure described for compound (**1**) to afford *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methylisoindoline-2-carboxamide (**12**) (140 mg, 40% yield). 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.32 (dd, $J=5.0$, 2.9 Hz, 1H) 7.16-7.13 (m, 1H) 7.12 (dd, $J=2.9$, 1.3 Hz, 1H) 7.10-7.07 (m, 2H) 7.02 (dd, $J=5.0$, 1.3 Hz, 1H) 5.01-4.88 (m, 1H) 4.70-4.54 (m, 4H) 3.77-3.70 (m, 1H) (dd, $J=8.5$, 3.7 Hz, 1H) 3.62-3.50 (m, 1H) 2.36 (s, 3H) 2.22 (s, 6H). ESI-MS (m/z): 352 $[\text{M}+\text{H}]^+$.

(b) *N*-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methylisoindoline-2-carboxamide hydrochloride (**12a**)

N-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methylisoindoline-2-carboxamide (**12**) (120 mg, 0.36 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using the procedure described for compound (**1**) to produce *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methylisoindoline-2-carboxamide hydrochloride (**12a**) (110 mg, 82% yield). 300 MHz ^1H -NMR (CD_3OD , ppm): 7.81 (dd, $J=2.9$, 1.4 Hz, 1H) 7.66 (dd, $J=5.0$, 2.9 Hz, 1H) 7.35-7.30 (m, 1H) 7.22-7.16 (m, 1H) 7.16-7.09 (m, 2H) 4.75 (dd, $J=9.0$, 4.7 Hz, 1H) 4.71-4.58 (m, 4H) 4.12 (dd, $J=15.1$, 9.0 Hz, 1H) 3.64 (dd, $J=15.1$, 4.7 Hz, 1H) 2.82 (s, 6H) 2.35 (s, 3H). ESI-MS (m/z): 330 $[\text{M}+\text{H}]^+$; melting point: 200-202 $^\circ\text{C}$ (dec.).

Example 17: 3-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-1-methyl-1-(1,2,3,4-tetrahydronaphthalen-1-yl)urea (13**) and corresponding hydrochloride salt (**13a**)**




(a) 3-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-1-methyl-1-(1,2,3,4-tetrahydronaphthalen-1-yl)urea (**13**)

N-Methyl-1,2,3,4-tetrahydronaphthalen-1-amine (150 mg, 0.93 mmol) and N^1,N^1 -dimethyl-1-(thiophen-3-yl)ethane-1,2-diamine were reacted in CH_2Cl_2 using procedure described for compound **(1)** to afford 3-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-1-methyl-1-(1,2,3,4-tetrahydronaphthalen-1-yl)urea **(13)** (64 mg, 19% yield). 400 MHz ^1H -NMR (CDCl_3 , ppm): 7.31 (dd, $J=4.9, 2.9$ Hz, 1H) 7.16-7.04 (m, 5H), 7.03-6.98 (m, 1H) 5.58-5.38 (m, 1H) 5.10-4.92 (m, 1H) 3.74-3.62 (m, 2H) 3.61-3.49 (m, 1H) 2.84-2.71 (m, 2H) 2.60-2.50 (m, 3H) 2.23-2.13 (m, 6H) 2.09-1.92 (m, 2H) 1.87-1.49 (m, 2H). ESI-MS (m/z): 358 $[\text{M}+\text{H}]^+$.

(b) 3-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-1-methyl-1-(1,2,3,4-tetrahydronaphthalen-1-yl)urea hydrochloride (**13a**)

3-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-1-methyl-1-(1,2,3,4-tetrahydronaphthalen-1-yl)urea **(13)** (60 mg, 0.17 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using procedure described for compound **(1)** to produce 3-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-1-methyl-1-(1,2,3,4-tetrahydronaphthalen-1-yl)urea hydrochloride (**13a**) (55 mg, 83% yield). 400 MHz ^1H -NMR (CD_3OD , ppm): 7.83-7.75 (m, 1H) 7.71-7.64 (m, 1H) 7.31 (dd, $J=5.0, 1.2$ Hz, 1H) 7.17-7.05 (m, 3H) 7.03-6.91 (m, 1H) 5.46 (s, 1H) 4.75 (ddd, $J=14.3, 7.8, 5.6$ Hz, 1H) 4.16-4.01 (m, 1H) 3.69 (ddd, $J=14.7, 5.6, 3.5$ Hz, 1H) 2.98-2.72 (m, 8H) 2.56-2.53 (m, 3H) 2.07-1.93 (m, 2H) 1.88 - 1.71 (m, 2H). ESI-MS (m/z): 358 $[\text{M}+\text{H}]^+$; melting point: 188-190 $^\circ\text{C}$.

Example 18: N-(2-(2-chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)isoindoline-2-carboxamide (14) and corresponding hydrochloride salt (14a)



(a) *N*-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)isoindoline-2-carboxamide (**14**)


Isoindoline (99 mg, 0.83 mmol) and 1-(2-chloro-4-fluorophenyl)- N^1,N^1 -dimethylethane-1,2-diamine were reacted in CH_2Cl_2 using procedure described for compound **(1)** to afford *N*-(2-(2-chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)isoindoline-2-carboxamide (**14**) (185 mg, 62% yield). 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.45 (dd, $J=8.6, 6.2$ Hz, 1H) 7.31-7.23 (m, 3H) 7.14 (dd, $J=8.6, 2.6$ Hz, 1H) 7.05-6.97 (m, 1H) 4.71-4.54 (m, 5H) 4.04 (t, $J=6.3$ Hz, 1H) 3.82-3.72 (m, 1H) 3.49 (ddd, $J=13.3, 6.3, 5.2$ Hz, 1H) 2.26 (s,

6H). ESI-MS (m/z): 362, 364 [M+H]⁺.

(b) *N*-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)isoindoline-2-carboxamide hydrochloride (**14a**)

N-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)isoindoline-2-carboxamide (**14**) (170 mg, 0.47 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using procedure described for compound (**1**) to produce N-(2-(2-chloro-4-fluorophenyl)-2-(dimethylamino) ethyl)-isoindoline-2-carboxamide hydrochloride (**14b**) (160 mg, 86% yield). 300 MHz ¹H-NMR (CD₃OD, ppm): 7.74 (dd, J=8.5, 5.8 Hz, 1H) 7.48 (dd, J=8.5, 2.7 Hz, 1H) 7.37-7.25 (m, 5H) 5.17-5.05 (m, 1H) 4.73-4.54 (m, 4H) 4.10 (dd, J=14.8, 6.8 Hz, 1H) 3.73 (dd, J=14.8, 5.9 Hz, 1H) 3.11-2.73 (m, 6H). ESI-MS (m/z): 362, 364 [M+H]⁺; melting point: 147-149 °C.

Example 19: N-(2-(dimethylamino)-2-(4-methoxyphenyl)ethyl)isoindoline-2-carboxamide (15**) and corresponding hydrochloride salt (**15a**)**



(a) *N*-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)isoindoline-2-carboxamide (**15**)


Isoindoline (123 mg, 1.03 mmol) and 1-(4-methoxyphenyl)-N¹,N¹-dimethylethane-1,2-diamine were reacted in CH₂Cl₂ using procedure described for compound (**1**) to afford N-(2-(dimethylamino)-2-(4-methoxyphenyl)ethyl)isoindoline-2-carboxamide (**15**) (190 mg, 54% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.30-7.22 (m, 4H) 7.21-7.14 (m, 2H) 6.92-6.85 (m, 2H) 4.78-4.55 (m, 5H) 3.85-3.74 (m, 1H) 3.81 (s, 3H) 3.51-3.37 (m, 2H) 2.20 (s, 6H). ESI-MS (m/z): 340 [M+H]⁺.

(b) *N*-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)isoindoline-2-carboxamide hydrochloride (**15a**)

N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)isoindoline-2-carboxamide (**15**) (165 mg, 0.49 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (25 mL/2 mL) using procedure described for compound (**1**) to produce N-(2-(dimethylamino)-2-(4-methoxyphenyl)-ethyl)isoindoline-2-carboxamide hydrochloride (**15a**) (148 mg, 81% yield). 300 MHz ¹H-NMR (CD₃OD, ppm): 7.51-7.42 (m, 2H) 7.36-7.23 (m, 4H) 7.11-7.03

(m, 2H) 4.77-4.59 (m, 4H) 4.55 (dd, J=8.7, 5.0 Hz, 1H) 4.16 (dd, J=15.0, 8.7 Hz, 1H) 3.83 (s, 3H) 3.66 (dd, J=15.0, 5.0 Hz, 1H) 2.81 (s, 6H). ESI-MS (m/z): 340 [M+H]⁺; melting point: 149-151 °C (dec.).

5 **Example 20: N-(2-(dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide (16) and corresponding hydrochloride salt (16a)**




(a) *N*-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide (16)

10 5-Fluoroisindoline hydrochloride (115 mg, 0.59 mmol) and 1-(4-methoxyphenyl)-N¹,N¹-dimethylethane-1,2-diamine were reacted in CH₂Cl₂ using procedure described for compound (1) to afford N-(2-(dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide (16) (120 mg, 81% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.23-7.14 (m, 3H) 7.01-6.92 (m, 2H) 6.92-6.86 (m, 2H) 4.80-4.71 (m, 1H) 4.68-4.50 (m, 4H) 3.84-3.72 (m, 1H) 3.81 (s, 3H) 3.51-3.37 (m, 2H) 2.20 (s, 6H). ESI-MS (m/z): 358 [M+H]⁺.

(b) *N*-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide hydrochloride (16a)

20 N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide (16) (115 mg, 0.32 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (25 mL/0.5 mL) using procedure described for compound (1) to produce N-(2-(dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide hydrochloride (16a) (104 mg, 82% yield). 300 MHz ¹H-NMR (CD₃OD, ppm): 7.50-7.42 (m, 2H) 7.31 (dd, J=8.3, 5.0 Hz, 1H) 7.12-6.99 (m, 4H) 4.77-4.57 (m, 4H) 4.53 (dd, J=8.6, 5.1 Hz, 1H) 4.14 (dd, J=15.0, 8.6 Hz, 1H) 3.83 (s, 3H) 3.65 (dd, J=15.0, 5.1 Hz, 1H) 2.87 (s, 3H) 2.75 (s, 3H). ESI-MS (m/z): 358 [M+H]⁺; melting point: 145-149 °C (dec.).

Example 21: N-(2-(2-chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)-5-fluoroisindoline-2-carboxamide (17) and corresponding hydrochloride salt (17a)




(a) *N*-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)-5-fluoroisoindoline-2-carboxamide (**17**)

5 5-Fluoroisoindoline hydrochloride (71 mg, 0.41 mmol) and 1-(2-chloro-4-fluorophenyl)- N^1,N^1 -dimethylethane-1,2-diamine were reacted in CH_2Cl_2 using procedure described for compound (**1**) to afford *N*-(2-(2-chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)-5-fluoroisoindoline-2-carboxamide (**17**) (135 mg, 87% yield). 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.44 (dd, $J=8.7, 6.3$ Hz, 1H) 7.20 (dd, $J=8.0, 5.0$ Hz, 1H) 7.14 (dd, $J=8.7, 2.6$ Hz, 1H) 7.06-6.92 (m, 3H) 4.70-4.51 (m, 5H) 4.05 (t, $J=6.3$ Hz, 1H) 3.80-3.70 (m, 1H) 3.54-3.43 (m, 1H) 2.26 (s, 6H). ESI-MS (m/z): 380, 382 $[\text{M}+\text{H}]^+$.

(b) *N*-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)-5-fluoroisoindoline-2-carboxamide hydrochloride (**17a**)

N-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)-5-fluoroisoindoline-2-carboxamide (**17**) (120 mg, 0.34 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using procedure described for compound (**1**) to produce *N*-(2-(2-chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)-5-fluoroisoindoline-2-carboxamide hydrochloride (**17a**). 300 MHz ^1H -NMR (CD_3OD , ppm): 7.75 (dd, $J=8.5, 5.8$ Hz, 1H) 7.49 (dd, $J=8.5, 2.7$ Hz, 1H) 7.37-7.25 (m, 2H) 7.10-6.98 (m, 2H) 5.17-5.05 (m, 1H) 4.73-4.54 (m, 4H) 4.10 (dd, $J=14.8, 6.8$ Hz, 1H) 3.74 (dd, $J=14.8, 5.9$ Hz, 1H) 3.16-2.58 (m, 6H). ESI-MS (m/z): 380, 382 $[\text{M}+\text{H}]^+$; melting point: 173-174 $^\circ\text{C}$ (dec.).

Example 22: *N*-(2-(dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide (18**) and corresponding hydrochloride salt (**18a**)**



25 (a) *N*-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide (**18**)


Isoindoline (100 mg, 0.84 mmol) and N^1,N^1 -dimethyl-1-phenylethane-1,2-diamine

were reacted in CH₂Cl₂ using procedure described for compound (1) to afford N-(2-(dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide (18) (120 mg, 46% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.40-7.22 (m, 9H) 4.80-4.70 (m, 1H) 4.70-4.56 (m, 4H) 3.89-3.77 (m, 1H) 3.56-3.42 (m, 3H) 2.24 (s, 6H). ESI-MS (m/z): 310 [M+H]⁺.

5 (b) *N*-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride (18a)

N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide (18) (109 mg, 0.35 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using procedure described for compound (1) to produce N-(2-(dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride (18a) (100 mg, 82% yield). 300 MHz ¹H-NMR (CD₃OD, ppm): 7.62-7.47 (m, 5H) 7.38-7.22 (m, 4H) 4.77-4.61 (m, 4H) 4.57 (dd, J=8.1, 5.1 Hz, 1H) 4.16 (dd, J=15.0, 8.1 Hz, 1H) 3.71 (dd, J=15.0, 5.1 Hz, 1H) 2.86 (s, 6H). ESI-MS (m/z): 310 [M+H]⁺; melting point: 107-108 °C.

15 **Example 23: 1-(2-(dimethylamino)-2-phenylethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea (19) and corresponding hydrochloride salt (19a)**



(a) *1*-(2-(Dimethylamino)-2-phenylethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea (19)

1,2,3,4-Tetrahydronaphthalen-2-amine (80 mg, 0.54 mmol) and N¹,N¹-dimethyl-1-phenylethane-1,2-diamine were reacted in CH₂Cl₂ using procedure described for compound (1) to afford 1-(2-(dimethylamino)-2-phenylethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea (19) (155 mg, 85% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.38-7.28 (m, 3H) 7.23-7.17 (m, 2H) 7.14-7.01 (m, 4H) 4.76-4.60 (m, 2H) 4.11-3.96 (m, 1H) 3.77-3.63 (m, 1H) 3.50-3.34 (m, 2H) 3.06-3.02 (m, 1H) 2.91-2.80 (m, 2H) 2.60 (dd, J=16.3, 7.9 Hz, 1H) 2.16 (s, 6H) 2.08-1.95 (m, 1H) 1.82-1.65 (m, 1H). ESI-MS (m/z): 338 [M+H]⁺.


25 (b) *1*-(2-(Dimethylamino)-2-phenylethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea hydrochloride (19a)

1-(2-(Dimethylamino)-2-phenylethyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea (19) (150 mg, 0.44 mmol) was treated with 2M HCl /diethyl ether in diethyl ether using procedure described for compound (1) to produce 1-(2-(dimethylamino)-2-phenylethyl)-3-(1,2,3,4-

tetrahydro-naphthalen-2-yl)urea hydrochloride (**19a**) (120 mg, 85% yield). 300 MHz ^1H -NMR (CD_3OD , ppm): 7.59-7.43 (m, 5H) 7.13-6.96 (m, 4H) 4.48 (td, $J=6.6$, 2.1 Hz, 1H) 4.10-3.90 (m, 2H) 3.69 (ddd, $J=14.8$, 6.6, 3.6 Hz, 1H) 3.02 (dd, $J=16.3$, 5.2 Hz, 1H) 3.02-2.59 (m, 6H) 2.97-2.67 (m, 2H) 2.62 (dd, $J=16.3$, 8.5 Hz, 1H) 2.08-1.95 (m, 1H) 1.79-1.62 (m, 1H).

5 ESI-MS (m/z): 338 $[\text{M}+\text{H}]^+$; melting point: 160-161 $^\circ\text{C}$.

Example 24: N-(2-(dimethylamino)-2-phenylethyl)-5-fluoroisoindoline-2-carboxamide (20) and corresponding hydrochloride salt (20a)



10 (a) *N*-(2-(Dimethylamino)-2-phenylethyl)-5-fluoroisoindoline-2-carboxamide (**20**)

A mixture of N^1, N^1 -dimethyl-1-phenylethane-1,2-diamine (176 mg, 1.07 mmol) and *N,N*-diisopropylethylamine (260 μL , 1.50 mmol) in CH_2Cl_2 (10 mL) was purged with argon and cooled to -70°C . A solution of 4-nitrophenylchloroformate (**2**) (216 mg, 1.07 mmol) in CH_2Cl_2 (5 mL) was added and resulting solution was stirred at -70°C for 30 min. After this
 15 time, 5-fluoroisoindoline hydrochloride (130 mg, 0.75 mmol) was added followed by *N,N*-diisopropylethylamine (390 μL , 2.24 mmol), and the resulting mixture was stirred for 16 h at room temperature. A saturated NaHCO_3 solution (10 mL) was added and the resulting suspension was extracted with CH_2Cl_2 (3 x 10 mL). The combined organic extracts were washed with water (30 mL) and dried over solid anhydrous MgSO_4 . After filtration, the
 20 volatiles were removed, and the residue was purified by flash chromatography using eluent from CH_2Cl_2 to $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (10:1) to give *N*-(2-(dimethylamino)-2-phenylethyl)-5-fluoroisoindoline-2-carboxamide (**20**) as an oil (178 mg, 73% yield). 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.40-7.30 (m, 3H) 7.30-7.24 (m, 2H) 7.19 (dd, $J=8.1$, 4.9 Hz, 1H) 7.02-6.91 (m, 2H) 4.81-4.69 (m, 1H) 4.69-4.49 (m, 4H) 3.91-3.72 (m, 1H) 3.56-3.41 (m, 2H) 2.23 (s,
 25 6H). ESI-MS (m/z): 328 $[\text{M}+\text{H}]^+$.


(b) *N*-(2-(Dimethylamino)-2-phenylethyl)-5-fluoroisoindoline-2-carboxamide hydrochloride (**20a**)

N-(2-(Dimethylamino)-2-phenylethyl)-5-fluoroisoindoline-2-carboxamide (**20**) (174 mg, 0.53 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH using
 30 procedure described for compound (**1**) to produce *N*-(2-(dimethylamino)-2-phenylethyl)-5-

fluoroisindoline-2-carboxamide hydrochloride (**20a**) (185 mg, 96% yield). 300 MHz ^1H -NMR (CD_3OD , ppm): 7.54 (s, 5H) 7.31 (dd, $J=8.2, 5.2$ Hz, 1H) 7.15-6.96 (m, 2H) 4.71-4.61 (m, 4H) 4.58 (dd, $J=8.1, 5.1$ Hz, 1H), 4.15 (dd, $J=15.0, 8.1$ Hz, 1H) 3.71 (dd, $J=15.0, 5.1$ Hz, 1H) 2.85 (s, 6H). ESI-MS (m/z): 328 $[\text{M}+\text{H}]^+$; melting point: 148-150 °C.

5

Example 25: *N*-(2-(Dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide (21**) and corresponding hydrochloride salt (**21a**)**



(a) *N*-(2-(Dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide (**21**)

10 A 1M $\text{BBr}_3/\text{CH}_2\text{Cl}_2$ solution (2.80 mL, 2.80 mmol) was added dropwise to the solution of *N*-(2-(dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide (**16**) (100 mg, 0.28 mmol) in CH_2Cl_2 (3 mL) at 0 °C under an argon atmosphere. The reaction mixture was stirred at room temperature for 20 h. After this time, MeOH (3 mL) was added, and the mixture was stirred for 30 min. The volatiles were removed *in vacuo*, and the resultant residue was purified by flash chromatography using gradient elution from CH_2Cl_2 to $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (4:1) to give *N*-(2-(dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide (**21**) (70 mg, 73% yield). 300 MHz ^1H -NMR (CD_3OD , ppm): 7.34-7.21 (m, 3H) 7.10-6.97 (m, 2H) 6.90-6.82 (m, 2H) 4.72-4.48 (m, 4H) 4.17-4.06 (m, 1H) 4.01 (dd, $J=14.0, 7.8$ Hz, 1H) 3.55 (dd, $J=14.0, 5.9$ Hz) 2.56 (s, 6H); ESI-MS (m/z): 344 $[\text{M}+\text{H}]^+$.


20

(b) *N*-(2-(Dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide hydrochloride (**21a**)

25 A 2M HCl/diethyl ether solution (100 μL , 0.20 mmol) was added to the solution of *N*-(2-(dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-fluoroisindoline-2-carboxamide (**21**) (70 mg, 0.20 mmol) in diethyl ether/MeOH (5 mL/1.5 mL). The mixture was stirred for 2 h at ambient temperature. The resultant precipitate were filtered and dried over P_2O_5 *in vacuo* at 65 °C for 24 h to give *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)isindoline-2-carboxamide hydrochloride (**21a**) (65 mg, 84% yield). 300 MHz ^1H -NMR (CD_3OD , ppm): 7.40-7.26 (m, 3H) 7.14-6.97 (m, 2H) 6.97-6.84 (m, 2H) 4.77-4.58 (m, 4H) 4.50 (dd, $J=8.8,$

5.0 Hz, 1H) 4.14 (dd, J=15.0, 8.8 Hz, 1H) 3.63 (dd, J=15.0, 5.0 Hz, 1H) 2.84 (s, 3H) 2.75 (s, 3H); ESI-MS (m/z): 344 [M+H]⁺; melting point: 180-185 °C (dec.).


Example 26: *N*-(2-(Dimethylamino)-2-(thiophen-3-yl)ethyl)-5-hydroxyisoindoline-2-carboxamide hydrobromide (22a)



A 1M BBr₃/CH₂Cl₂ solution (1.45 mL, 1.45 mmol) was added dropwise to the solution of *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-methoxyisoindoline-2-carboxamide (**5**) (50 mg, 0.14 mmol) in CH₂Cl₂ (10 mL) at 0 °C under argon atmosphere.

The reaction mixture was stirred at room temperature for 24 h. After this time, MeOH (10 mL) was added and the mixture was stirred for 30 min. The volatiles were removed *in vacuo*, and the resultant oily residue was treated with CH₂Cl₂. The resultant precipitate were filtered, washed with CH₂Cl₂, then with Et₂O, and dried to give *N*-(2-(dimethylamino)-2-(thiophen-3-yl)ethyl)-5-hydroxyisoindoline-2-carboxamide hydrobromide (**22a**) (70 mg, 73% yield). 300 MHz ¹H-NMR (DMSO-d₆, ppm): 10.2-8.6 (br s, 1H) 9.47 (s, 1H) 7.92 (dd, J=2.9, 1.2 Hz, 1H) 7.73 (dd, J=5.0, 2.9 Hz, 1H) 7.36 (dd, J=5.0, 1.2 Hz, 1H) 7.15-7.08 (m, 1H) 6.80-6.67 (m, 3H) 4.76-4.66 (m, 1H) 4.60-4.43 (m, 4H) 3.90 (ddd, J=15.0, 8.8, 6.2 Hz, 1H) 3.55-3.42 (m, 1H) 2.71 (d, J=4.8 Hz, 3H) 2.64 (d, J=4.8 Hz, 3H); ESI-MS (m/z): 332 [M+H]⁺; melting point: 134 °C (dec.).

Example 27: *N*-(2-(dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-methoxyisoindoline-2-carboxamide (23) and *N*-(2-(dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-hydroxyisoindoline-2-carboxamide hydrobromide (24a)




(a) *N*-(2-(dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-methoxyisoindoline-2-carboxamide (**23**)

A mixture of 1-(4-methoxyphenyl)-*N,N'*-dimethylethane-1,2-diamine (78 mg, 0.40 mmol) and *N,N*-diisopropylethylamine (70 μ L, 0.40 mmol) in CH_2Cl_2 (10 mL) was purged with argon and cooled to 0 °C. A solution of 4-nitrophenylchloroformate (81 mg, 0.40 mmol) in CH_2Cl_2 (5 mL) was added, and resulting solution was stirred at -70 °C for 30 min. After this time, 5-methoxyisoindoline hydrochloride (50 mg, 0.27 mmol) was added, followed by *N,N*-diisopropylethylamine (93 μ L, 0.54 mmol), and the resulting mixture was stirred for 16 h at room temperature. A saturated NaHCO_3 solution (10 mL) was added, and the resulting suspension was extracted with EtOAc (3 x 10 mL). The combined organic extracts were washed with water (30 mL), then with a brine solution (20 mL) and dried over solid anhydrous Na_2SO_4 . After filtration, the volatiles were removed and the residue was purified by flash chromatography using eluent from CH_2Cl_2 to $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (1:1) to give *N*-(2-(dimethyl amino)-2-(4-methoxy phenyl)ethyl)-5-methoxyisoindoline-2-carboxamide (**23**) (79 mg, 79% yield). 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.23-7.10 (m, 2H) 6.94-6.86 (m, 2H) 6.86-6.76 (m, 3H) 4.64-4.52 (m, 4H) 3.86-3.77 (m, 1H) 3.82 (s, 3H) 3.80 (s, 3H) 3.53-3.46 (m, 2H) 2.20 (s, 6H); ESI-MS (m/z): 370 $[\text{M}+\text{H}]^+$.

(b) *N*-(2-(dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-hydroxyisoindoline-2-carboxamide hydrobromide (**24a**)

N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-methoxyisoindoline-2-carboxamide (**23**) (75 mg, 0.20 mmol) and 1M $\text{BBr}_3/\text{CH}_2\text{Cl}_2$ were reacted in CH_2Cl_2 using procedure described for compound (**22a**) to obtain *N*-(2-(dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-hydroxyisoindoline-2-carboxamide hydrobromide (**24a**) (65 mg, 94% yield). 300 MHz ^1H -NMR ($\text{DMSO}-d_6$, ppm): 10.7-8.7 (br s, 2H) 9.79 (br s, 1H) 9.52-9.30 (m, 1H) 7.41-7.27 (m, 2H) 7.15-7.05 (m, 1H) 6.91-6.78 (m, 2H) 6.77-6.61 (m, 3H) 4.60-4.37 (m, 5H) 3.91 (ddd, $J=14.8, 8.5, 6.3$ Hz, 1H) 3.47 (ddd, $J=14.8, 4.9, 4.9$ Hz, 1H) 2.69 (d, $J=4.8$ Hz, 3H) 2.59 (d, $J=4.8$ Hz, 3H); ESI-MS (m/z): 342 $[\text{M}+\text{H}]^+$; melting point: 185 °C (dec.).

Example 28: (*R*)-N-(3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisoindoline-2-carboxamide (30-*R*) and (*R*)-N-(3-(dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisoindoline-2-carboxamide hydrochloride (30a-*R*)



Scheme 8

(a) (*R*)-2-Methyl-*N*-(thiophen-3-ylmethylene)propane-2-sulfinamide (**25-*R***)

A mixture of thiophene-3-carboxaldehyde (1.50 g, 13.38 mmol), (*R*)-(+)-2-methyl-2-propanesulfinamide (1.62 g, 13.38 mmol), pyrrolidine (111 μ L, 1.34 mmol), and molecular sieves (3Å) (0.1 g) in CH₂Cl₂ (18 mL) was heated at 50 °C for 10 h in a closed vial. After cooling, the volatiles were removed in vacuo, and the resultant residue was purified by flash chromatography using CH₂Cl₂ as an eluent to give (*R*)-2-methyl-*N*-(thiophen-3-ylmethylene)propane-2-sulfinamide (**25-*R***) (2.68 g, 93% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 8.58 (s, 1H) 7.86 (dd, *J*=2.9, 1.1 Hz, 1H) 7.58 (dd, *J*=5.1, 1.1 Hz, 1H) 7.37 (ddd, *J*=5.1, 2.9, 0.6 Hz, 1H) 1.25 (s, 9H); ESI-MS (*m/z*): 216 [M+H]⁺.

(b) (*R*)-*N*-((*R*)-2-Cyano-1-(thiophen-3-yl)ethyl)-2-methylpropane-2-sulfinamide (**26-*R,R***) and (*R*)-*N*-((*S*)-2-cyano-1-(thiophen-3-yl)ethyl)-2-methylpropane-2-sulfinamide (**26-*R,S***)

An oven dried flask with *N,N*-diisopropylamine (1.16 mL, 8.27 mmol) in THF (10 mL) was purged with argon and cooled to -78°C (Dry Ice / acetone bath). A solution of *n*-butyllithium in hexane (3.75 mL, 2.2 M) was added, and the mixture was stirred at ambient temperature for 20 min. The reaction mixture was cooled to -78°C and a solution of acetonitrile (0.43 mL, 8.26 mmol) in THF (5 mL) was added. After stirring at -78°C for 10

min. a solution of (*R*)-2-methyl-N-(thiophen-3-ylmethylene)propane-2-sulfinamide (**25-R**) (1.78 g, 8.26 mmol) in THF (10 mL) was added dropwise. The reaction mixture was stirred at -78°C for 20 min, after which time a saturated aqueous NH₄Cl (30 mL) was added and the mixture was allowed to reach room temperature. The mixture was extracted with EtOAc (3 x 20 mL). The combined organic extracts were washed with water (50 mL), then with a brine solution (50 mL) and dried over solid anhydrous Na₂SO₄. After filtration, the volatiles were removed in vacuo and the resultant residue was purified by flash chromatography using eluent gradient from CH₂Cl₂ to CH₂Cl₂/Et₂O (1:4) to give (*R*)-N-((*R*)-2-cyano-1-(thiophen-3-yl)ethyl)-2-methylpropane-2-sulfinamide (**26-R,R**) (0.88 g, 41% yield) and (*R*)-N-((*S*)-2-cyano-1-(thiophen-3-yl)ethyl)-2-methylpropane-2-sulfinamide (**26-R,S**) (1.00 g, 47% yield). (**26-R,R**): 300 MHz ¹H-NMR (CDCl₃, ppm): 7.42-7.33 (m, 2H) 7.21-7.14 (m, 1H) 4.89-4.81 (m, 1H) 3.63 (d, J=4.0 Hz, 1H) 3.01 (dd, J=16.8, 5.8 Hz, 1H) 2.85 (dd, J=16.8, 5.2 Hz, 1H) 1.29 (s, 9H); ESI-MS (m/z): 257 [M+H]⁺.

(**26-R,S**): 300 MHz ¹H-NMR (CDCl₃, ppm): 7.41-7.30 (m, 2H) 7.10 (dd, J=4.9, 1.5 Hz, 1H) 4.87-4.78 (m, 1H) 3.76 (d, J=6.5 Hz, 1H) 3.03 (dd, J=16.8, 5.8 Hz, 1H) 2.98 (dd, J=16.8, 5.4 Hz, 1H) 1.29 (s, 9H); ESI-MS (m/z): 257 [M+H]⁺.

(c) (*R*)-3-Amino-3-(thiophen-3-yl)propanenitrile hydrogen chloride (**27-R**)

A 4N HCl /1,4-dioxane solution (2.1 mL, 8.40 mmol) was added to a solution of (*R*)-N-((*R*)-2-cyano-1-(thiophen-3-yl)ethyl)-2-methylpropane-2-sulfinamide (**26-R,R**) (540 mg, 2.11 mmol) in THF (20 mL). The mixture was stirred at ambient temperature for 1h. The solvents were removed under reduced pressure. The crude product was suspended in diethyl ether (30 mL) and EtOH (0.5 mL) was added. The solvents were decanted, and the remaining precipitate was washed with diethyl ether (2 x 30 mL) and dried to give (*R*)-3-amino-3-(thiophen-3-yl)propanenitrile hydrogen chloride (**27-R**) (390 mg, 98% yield). 400 MHz ¹H-NMR (DMSO-d₆, ppm): 9.02 (s, 3H) 7.81 (dd, J=3.0, 1.3 Hz, 1H) 7.81 (dd, J=5.1, 3.0 Hz, 1H) 7.40 (dd, J=5.1, 1.3 Hz, 1H) 4.80 (dd, J=8.6, 5.2 Hz, 1H) 3.46-3.23 (m, 2H); ESI-MS (m/z): 153 [M+H]⁺.

(d) (*R*)-3-(Dimethylamino)-3-(thiophen-3-yl)propanenitrile (**28-R**)

To a suspension of (*R*)-3-amino-3-(thiophen-3-yl)propanenitrile hydrogen chloride (**27-R**) (410 mg, 2.17 mmol) and formaldehyde solution in water (37%) (0.77 mL, 9.56 mmol) in MeCN (20 mL), solid Na(AcO)₃BH (2.30 g, 10.86 mmol) was added in portions. The reaction mixture was stirred at ambient temperature for 1 h, and then quenched by adding a saturated aqueous Na₂CO₃ solution (20 mL). The resultant mixture was extracted with CH₂Cl₂ (3 x 15 mL). The combined organic extracts were dried over Na₂SO₄ and

concentrated in *vacuo*. Product purified by flash chromatography using eluent gradient from CH₂Cl₂ to CH₂Cl₂/Et₂O (1:4) to give (*R*)-3-(dimethylamino)-3-(thiophen-3-yl)propanenitrile (**28-R**) (370 mg, 95% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.34 (dd, J=5.0, 2.9 Hz, 1H) 7.20-7.15 (m, 1H) 7.06 (dd, J=5.0, 1.4 Hz, 1H) 3.87-3.80 (m, 1H) 2.84 (dd, J=16.7, 6.3 Hz, 1H) 2.76 (dd, J=16.7, 7.3 Hz, 1H) 2.23 (s, 6H). ESI-MS (m/z): 181 [M+H]⁺.

(*e*) (*R*)-*N*¹,*N*¹-Dimethyl-1-(thiophen-3-yl)propane-1,3-diamine (**29-R**)

A solution of (*R*)-3-(dimethylamino)-3-(thiophen-3-yl)propanenitrile (**28-R**) (370 mg, 2.05 mmol) in diethyl ether (10 mL) was added dropwise (1 mL/min) to a suspension of lithium aluminum hydride (230 mg, 6.16 mmol) in diethyl ether (20 mL) at 0°C under an argon atmosphere. The mixture was heated at reflux for 90 min, then cooled in an ice cold water bath and lastly quenched by addition of water (370 µL), 2N aqueous NaOH (370 µL) and water (1.11 mL). The resultant white precipitate was filtered through a Celite plug, and the plug was washed with diethyl ether (50 mL). The filtrate was evaporated to give (*R*)-*N*¹,*N*¹-dimethyl-1-(thiophen-3-yl)propane-1,3-diamine (**29-R**) (375 mg, 99% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.30-7.26 (m, 1H) 7.07-7.01 (m, 1H) 6.99 (dd, J=4.9, 1.3 Hz, 1H) 3.56 (dd, J=8.8, 6.0 Hz, 1H) 2.66-2.58 (m, 2H) 2.16 (s, 6H) 2.09-1.93 (m, 1H) 1.92-1.78 (m, 1H); ESI-MS (m/z): 185 [M+H]⁺.

(*f*) (*R*)-*N*-(3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide (**30-R**)

4-Fluoroisindoline hydrochloride (80 mg, 0.46 mmol) and (*R*)-*N*¹,*N*¹-dimethyl-1-(thiophen-3-yl)propane-1,3-diamine (0.12 g, 0.66 mmol) were reacted in CH₂Cl₂ using procedure described for compound (**3**) to afford (*R*)-*N*-(3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide (**30-R**) (135 mg, 87% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.31 (dd, J=4.9, 2.9 Hz, 1H) 7.20 (dd, J=8.1, 5.0 Hz, 1H) 7.10 (dd, J=2.9, 1.1 Hz, 1H) 7.01 (dd, J=4.9, 1.1 Hz, 1H) 7.00-6.93 (m, 2H) 5.96 (t, J=4.7 Hz, 1H) 4.66-4.55 (m, 4H) 3.80-3.67 (m, 1H) 3.54-3.41 (m, 1H) 3.38-3.26 (m, 1H) 2.24 (s, 6H) 2.23-2.14 (m, 1H) 2.03-1.89 (m, 1H). ESI-MS (m/z): 348 [M+H]⁺. [α_D^{20}]: - 7.2 (1.35, acetone).

(*e*) (*R*)-*N*-(3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide hydrochloride (**30a-R**)

A 2M HCl/diethyl ether solution (194 µL, 0.39 mmol) was added to the solution of (*R*)-*N*-(3-(dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide (**30-R**) (135 mg, 0.39 mmol) in diethyl ether/MeOH (25 mL/1 mL). The mixture was stirred at ambient temperature for 2 h. The solvents were removed via decantation. The resultant precipitate was washed with diethyl ether (2 x 30 mL) and dried over P₂O₅ in *vacuo* at 65 °C

for 24 h to give (*R*)-*N*-(3-(dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide hydrochloride (**30a-R**) (138 mg, 93% yield). 300 MHz ¹H-NMR (CD₃OD, ppm): 7.86-7.72 (m, 1H) 7.60 (dd, *J*=5.1, 2.9 Hz, 1H) 7.38-7.24 (m, 2H) 7.11-6.97 (m, 2H) 4.72-4.47 (m, 5H) 3.40-3.11 (m, 2H) 2.77 (s, 3H) 2.73 (s, 3H) 2.51-2.23 (m, 2H); ESI-MS (m/z): 348 [M+H]⁺.

Example 29: (S)-N-(3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide (30-S) and (S)-N-(3-(dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide hydrochloride (30a-S)

10 (a) (*S*)-3-Amino-3-(thiophen-3-yl)propanenitrile hydrochloride (**27-S**)
(*R*)-*N*-((*S*)-2-Cyano-1-(thiophen-3-yl)ethyl)-2-methylpropane-2-sulfinamide (**26-R,S**) (650 mg, 2.53 mmol) and 4N HCl /1,4-dioxane (2.1 mL, 8.42 mmol) were reacted in THF using procedure described for compound (**27-R**) to afford the desired compound (**27-S**) (470 mg) which was used in the next step without purification.

15 (b) (*S*)-3-(Dimethylamino)-3-(thiophen-3-yl)propanenitrile (**28-S**)
(*S*)-3-Amino-3-(thiophen-3-yl)propanenitrile hydrochloride (**27-S**) (470 mg, 2.49 mmol) and formaldehyde solution in water (37%) (0.89 mL, 10.96 mmol) were reacted in MeCN using procedure described for compound (**28-R**) to afford the desired compound (**28-S**) (185 mg, 38% yield in 2 steps from (**26-R,S**)). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.34 (dd, *J*=5.0, 2.9 Hz, 1H) 7.20-7.15 (m, 1H) 7.06 (dd, *J*=5.0, 1.3 Hz, 1H) 3.87-3.79 (m, 1H) 2.84 (dd, *J*=16.7, 6.3 Hz, 1H) 2.76 (dd, *J*=16.7, 7.3 Hz, 1H) 2.23 (s, 6H); ESI-MS (m/z): 181 [M+H]⁺.

20 (c) (*S*)-*N*¹,*N*¹-Dimethyl-1-(thiophen-3-yl)propane-1,3-diamine (**29-S**)
(*S*)-3-(Dimethylamino)-3-(thiophen-3-yl)propanenitrile **23-S** (220 mg, 1.22 mmol) and lithium aluminium hydride were reacted in diethyl ether using procedure described for compound (**28-R**) to afford the desired compound (**28-S**) (185 mg, 82% yield). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.30-7.26 (m, 1H) 7.04-7.01 (m, 1H) 6.98 (dd, *J*=5.0, 1.3 Hz, 1H) 3.56 (dd, *J*=8.8, 6.0 Hz, 1H) 2.66-2.58 (m, 2H) 2.16 (s, 6H) 2.09-1.93 (m, 1H) 1.92-1.78 (m, 1H); ESI-MS (m/z): 185 [M+H]⁺.

30 (d) (*S*)-*N*-(3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide (**30-S**)


4-Fluoroisindoline hydrochloride (82 mg, 0.47 mmol) and (*S*)-*N*¹,*N*¹-dimethyl-1-(thiophen-3-yl)propane-1,3-diamine were reacted in CH₂Cl₂ using procedure described for compound (**3**) to afford the desired compound (**30-S**) (135 mg, 82% yield). 300 MHz ¹H-

NMR (CDCl₃, ppm): 7.30 (dd, J=4.9, 2.9 Hz, 1H) 7.20 (dd, J=8.1, 5.0 Hz, 1H) 7.10 (dd, J=2.9, 1.1 Hz, 1H) 7.00 (dd, J=4.9, 1.1 Hz, 1H) 7.00-6.93 (m, 2H) 5.96 (t, J=4.7 Hz, 1H) 4.66-4.55 (m, 4H) 3.77-3.67 (m, 1H) 3.55-3.43 (m, 1H) 3.38-3.25 (m, 1H) 2.24 (s, 6H) 2.23-2.14 (m, 1H) 2.03-1.89 (m, 1H); ESI-MS (m/z): 348 [M+H]⁺.

5 (e) (S)-N-(3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide hydrochloride (**26-S**)

(S)-N-(3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide (**30-S**) (130 mg, 0.37 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (25 mL/2 mL) using procedure described for compound (**25-R**) to produce (S)-N-(3-(dimethyl amino)-3-(thiophen-3-yl)propyl)-5-fluoroisindoline-2-carboxamide hydrochloride (**30a-S**) (131 mg, 91% yield). 300 MHz ¹H-NMR (CD₃OD, ppm): 7.79-7.72 (m, 1H) 7.61 (dd, J=5.0, 2.9 Hz, 1H) 7.35-7.25 (m, 2H) 7.11-7.00 (m, 2H) 4.67-4.51 (m, 5H) 3.40-3.17 (m, 2H) 2.75 (s, 6H) 2.50-2.21 (m, 2H); ESI-MS (m/z): 348 [M+H]⁺.

15 **Example 30: tert-Butyl (2-(5-fluoroisindoline-2-carboxamido)-1-(thiophen-3-yl)ethyl)(methyl)carbamate (34)**



Scheme 9

20 (a) 2-(Methylamino)-2-(thiophen-3-yl)acetonitrile (**31**)

A methylamine solution (40%, aqueous) (4.62 mL, 53.50 mmol) was added to solution of thiophene-3-carboxaldehyde (3.00 g, 26.75 mmol) in methanol (30 mL) and reaction mixture was stirred for 1h at room temperature. After this time, TMSCN (5.35 mL, 40.132 mmol) was added and stirring was continued for 16 h at room temperature. A

saturated aqueous NH_4Cl solution (7 mL) was added, and the resulting suspension was extracted with CH_2Cl_2 (3 x 20 mL). The combined organic extracts were washed with water (30 mL) and dried over solid anhydrous MgSO_4 . After filtration, the volatiles were removed to give 2-(methylamino)-2-(thiophen-3-yl)acetonitrile (**31**) (3.78 g, 93% yield), which was used in the next step without purification. 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.48-7.44 (m, 1H) 7.37 (dd, $J=5.0, 3.0$ Hz, 1H) 7.17 (dd, $J=5.0, 1.4$ Hz, 1H), 4.80 (s, 1H), 2.58 (s, 3H); ESI-MS (m/z): 153 $[\text{M}+\text{H}]^+$.

(b) *tert*-Butyl (cyano(thiophen-3-yl)methyl)(methyl)carbamate (**32**)

Di-*tert*-butyl decarbonate (Boc_2O) (4.75 mL, 20.67 mmol) in methanol (10 mL) was added dropwise to the solution of 2-(methylamino)-2-(thiophen-3-yl)acetonitrile (**31**) (2.42 g, 15.90 mmol) and triethylamine (4.43 mL, 31.80 mmol) in methanol (25 mL). The reaction mixture was stirred for 16 h at room temperature. After (rotary) evaporation, the residue was dissolved in CH_2Cl_2 (50 mL), washed with a 1N HCl solution (30 mL), then with water (30 mL) and dried over solid anhydrous MgSO_4 . After filtration, the volatiles were removed to give *tert*-butyl (cyano(thiophen-3-yl)methyl)(methyl)carbamate (**32**) (3.75 g, 94% yield), which was used in the next step without purification. 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.48-7.44 (m, 1H) 7.39 (dd, $J=5.0, 3.0$ Hz, 1H) 7.01 (dd, $J=5.0, 1.4$ Hz, 1H) 6.65-6.31 (m, 1H) 2.77 (s, 3H) 1.51 (s, 9H); ESI-MS (m/z): 253 $[\text{M}+\text{H}]^+$.

(c) *tert*-Butyl (2-amino-1-(thiophen-3-yl)ethyl)(methyl)carbamate (**33**)


A steel vessel was charged with a solution of *tert*-butyl (cyano(thiophen-3-yl)methyl)(methyl)-carbamate (**32**) (1.05 g, 4.16 mmol) in ethanol (20 mL) and NH_4OH (25%) (10 mL). A Raney-Nickel slurry in H_2O (700 mg) was added, and hydrogen gas was added to a pressure of 10 atm. The vessel was sealed and stirred at room temperature for 15 h. After depressurization, the mixture was filtered through Celite and the Celite pad was washed with ethanol (100 mL). The filtrate was evaporated, and the residue dissolved in CH_2Cl_2 (50 mL) and dried over solid anhydrous MgSO_4 . After filtration, the volatiles were removed, and the residue was purified by flash chromatography using eluent from CH_2Cl_2 to $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (1:1) to give *tert*-butyl (2-amino-1-(thiophen-3-yl)ethyl)(methyl)carbamate (**33**) (740 mg, 69% yield). 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.30 (dd, $J=5.0, 2.9$ Hz, 1H) 7.11-7.05 (m, 1H) 6.98 (dd, $J=5.0, 1.4$ Hz, 1H) 5.63-5.05 (m, 2H) 3.84-3.63 (m, 1H) 3.27-3.15 (m, 1H), 3.15-3.02 (m, 1H) 2.62 (s, 3H) 1.50 (s, 9H). ESI-MS (m/z): 257 $[\text{M}+\text{H}]^+$.

(d) *tert*-Butyl (2-(5-fluoroisoindoline-2-carboxamido)-1-(thiophen-3-yl)ethyl)(methyl)carbamate (**34**)

A mixture of *tert*-butyl (2-amino-1-(thiophen-3-yl)ethyl)(methyl)carbamate (**33**) (310

mg, 1.21 mmol) and *N,N*-diisopropylethylamine (300 μ L, 1.73 mmol) in CH_2Cl_2 (15 mL) was purged with argon and cooled to 0 $^\circ\text{C}$. A solution of 4-nitrophenylchloroformate (244 mg, 1.21 mmol) in CH_2Cl_2 (5 mL) was added, and resulting reaction mixture was stirred at -70 $^\circ\text{C}$ for 30 min. After this time, 5-fluoroisoindoline hydrochloride (150 mg, 0.86 mmol) was added, followed by *N,N*-diisopropylethylamine (450 μ L, 2.60 mmol). The resulting mixture was stirred for 20 h at room temperature. A saturated NaHCO_3 solution (10 mL) was added, and the resulting suspension was extracted with CH_2Cl_2 (3 x 15 mL). The combined organic extracts were washed with water (30 mL), then with a brine solution (20 mL) and dried over solid anhydrous Na_2SO_4 . After filtration, the volatiles were removed *in vacuo*, and the residue was purified by flash chromatography using eluent from CH_2Cl_2 to $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (1:1), then reverse phase chromatography using eluent from $\text{H}_2\text{O}/\text{MeCN}$ (9:1) MeCN , to give *tert*-butyl (2-(5-fluoroisoindoline-2-carboxamido)-1-(thiophen-3-yl)ethyl)(methyl)carbamate (**34**) (150 mg, 41% yield). 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.32 (dd, $J=5.0, 2.9$ Hz, 1H) 7.24-7.14 (m, 2H) 7.05-6.91 (m, 3H) 5.63-5.41 (m, 1H) 5.09-4.91 (m, 1H) 4.68 (s, 2H) 4.64 (s, 2H) 4.15-3.88 (m, 1H) 3.77-3.57 (m, 1H) 2.56 (s, 3H) 1.45 (s, 9H); ESI-MS (m/z): 420 $[\text{M}+\text{H}]^+$; melting point: 182-184 $^\circ\text{C}$.

Example 31: 5-Fluoro-*N*-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (35**) and corresponding hydrochloride salt (**35a**)**



*(a) 5-Fluoro-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (**35**)*


A 4M HCl/1,4-dioxane solution (343 μ L, 1.37 mmol) was added to *tert*-butyl (2-(5-fluoroisoindoline-2-carboxamido)-1-(thiophen-3-yl)ethyl)(methyl)carbamate (**34**) (115 mg, 0.27 mmol) in 1,4-dioxane (5 mL), and heated at 80 $^\circ\text{C}$ for 4h. A saturated NaHCO_3 solution (10 mL) was added, and the resulting suspension was extracted with CH_2Cl_2 (3 x 10 mL). The combined organic extracts were washed with water (30 mL) and dried over solid anhydrous MgSO_4 . After filtration, the volatiles were removed *in vacuo*, and the residue was purified by flash chromatography using eluent from CH_2Cl_2 to $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (10:1) to give 5-fluoro-*N*-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (**35**) (44 mg, 50% yield). 300 MHz ^1H -NMR (CDCl_3 , ppm): 7.34 (dd, $J=5.0, 2.9$ Hz, 1H) 7.24-7.16 (m,

2H) 7.08 (dd, J=5.0, 1.3 Hz, 1H) 7.02-6.92 (m, 2H) 4.89-4.78 (m, 1H) 4.74-4.54 (m, 4H) 3.89 (dd, J = 7.0, 5.5 Hz, 1H) 3.64-3.42 (m, 2H) 2.39 (s, 3H); ESI-MS (m/z): 320 [M+H]⁺.

(b) 5-Fluoro-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride (**35a**)

5 5-Fluoro-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide (**35**) (44 mg, 0.14 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (10 mL/1 mL) using procedure described for compound (**2**) to produce 5-fluoro-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride (**35a**) (45 mg, 92% yield). 300 MHz ¹H-NMR (CD₃OD, ppm): 7.67 (dd, J=2.9, 1.3 Hz, 1H) 7.63 (dd, J= 5.0, 2.9 Hz, 1H) 7.32 (dd, J=8.3, 5.0 Hz, 1H) 7.26 (dd, J=5.0, 1.3 Hz, 1H) 7.14- 6.98 (m, 2H) 4.73-4.62 (m, 4H) 4.54 (dd, J=7.7, 4.5 Hz, 1H) 3.83 (dd, J=14.8, 7.7 Hz, 1H) 3.66 (dd, J= 14.8, 4.5 Hz, 1H) 2.59 (s, 3H); ESI-MS (m/z): 320 [M+H]⁺.

Example 32: N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide (38**), (-)-N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide (**38-ent-A**), (+)-N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide (**38-ent-B**), and corresponding hydrochloride salts**



(a) *tert*-butyl (2-amino-1-phenylethyl)(methyl)carbamate (**36**)

20 *tert*-Butyl (2-amino-1-phenylethyl)(methyl)carbamate (**36**) was prepared from benzaldehyde using procedures described to make compound (**33**). 300 MHz ¹H-NMR (CDCl₃, ppm): 7.40-7.18 (m, 6H) 5.24 (s, 1H) 3.30-3.08 (m, 2H) 2.63 (s, 3H) 1.49 (s, 9H); ESI-MS (m/z): 251 [M+H]⁺.

(b) *tert*-butyl (2-(isoindoline-2-carboxamido)-1-phenylethyl)(methyl)carbamate (**37**)

25 Isoindoline (500 mg, 4.20 mmol) and *tert*-butyl (2-amino-1-phenylethyl)(methyl)carbamate (**36**) were reacted in CH₂Cl₂ using procedure described for compound (**34**) to obtain *tert*-butyl (2-(isoindoline-2-carboxamido)-1-phenylethyl)(methyl)carbamate (**37**) (1.49 g, 90% yield). 400 MHz ¹H-NMR (CDCl₃, ppm): 7.41-7.33 (m, 2H) 7.33-7.29 (m, 2H) 7.29-7.23 (m, 5H) 5.62-5.41 (m, 1H) 5.19-5.00 (m, 1H) 4.71 (s, 4H) 4.19-3.93 (m, 1H) 3.84-3.62 (m, 1H) 2.55 (s, 3H) 1.46 (s, 9H); ESI-MS (m/z): 396 [M+H]⁺.

30

(c) *N*-(2-(methylamino)-2-phenylethyl)isoindoline-2-carboxamide (**38**)

Trifluoroacetic acid (1.00 mL, 13.02 mmol) was added to *tert*-butyl (2-(isoindoline-2-carboxamido)-1-phenylethyl)(methyl)carbamate (**37**) (1.49 g, 3.77 mmol) in CH₂Cl₂ (15 mL), and stirred at room temperature for 2h. Volatiles were removed, saturated NaHCO₃ solution
5 (100 mL) was added, and the resulting suspension was extracted with CH₂Cl₂ (3x30 mL).

Combined organic extracts were washed with water (30 mL) and dried over MgSO₄.

Volatiles were removed, and the residue was purified by flash chromatography using eluent from CH₂Cl₂ to CH₂Cl₂/MeOH (10:1) to give *N*-(2-(methylamino)-2-phenylethyl)isoindoline-2-carboxamide (**42a**) (530 mg, 48% yield). 400 MHz ¹H-NMR (CDCl₃, ppm): 7.59-7.53 (m,
10 2H) 7.47-7.34 (m, 3H) 7.25-7.19 (m, 4H) 6.00-5.84 (m, 1H) 4.84-4.64 (m, 4H) 4.21 (dd, *J*=8.4, 4.2 Hz, 1H) 3.89-3.76 (m, 2H) 2.51 (s, 3H). ESI-MS (*m/z*): 296 [M+H]⁺.

(-)-*N*-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide (38-ent-A)

(+)-*N*-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide (38-ent-B)

The enantiomers were separated on chiral stationary phase chromatography using
15 (Chiralpak® IC 250x30mm, 5µm column and CH₂Cl₂/MeOH/ethanolamine (99:1:0.1) as mobile phase at 40mL/min flow rate: Rt(-)-enantiomer (**38-ent-A**) = 11 min, and Rt(+)-enantiomer (**38-ent-B**) = 19 min. (-)-Enantiomer (**38-ent-A**) [α_D^{20}]: - 12.7 (1.00, acetone); (+)-enantiomer (**38-ent-B**) [α_D^{20}]: +13.2 (1.02, acetone).

20 (d) *rac-N*-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride (**38a**)

N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide (**38**) (56 mg, 0.19 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (10 mL/1 mL) using procedure described for compound (**2a**) to produce *N*-(2-(methylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride (**38a**) (63 mg, 100% yield). 400 MHz
25 ¹H-NMR (CD₃OD, ppm): 7.56-7.45 (m, 5H) 7.34-7.27 (m, 4H) 4.69 (s, 4H) 4.37 (dd, *J*=7.8, 4.5 Hz, 1H) 3.85 (dd, *J*=14.8, 7.8 Hz, 1H) 3.67 (dd, *J*=14.8, 4.5 Hz, 1H) 2.59 (s, 3H); ESI-MS (*m/z*): 296 [M+H]⁺.

(e) **(-)-*N*-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride (38a-ent-A)**

30 (-)-*N*-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide (**38-ent-A**) (115 mg, 0.39 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (12 mL/0.1 mL) using procedure described for compound (**2a**) to produce (-)-*N*-(2-(methylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride (**38a-ent-A**) (102 mg, 79% yield). 400




MHz $^1\text{H-NMR}$ (CD_3OD , ppm): 7.56-7.45 (m, 5H) 7.34-7.27 (m, 4H) 4.69 (s, 4H) 4.36 (dd, $J=7.8, 4.5$ Hz, 1H) 3.84 (dd, $J=14.8, 7.8$ Hz, 1H) 3.66 (dd, $J=14.8, 4.5$ Hz, 1H) 2.58 (s, 3H); ESI-MS (m/z): 296 $[\text{M}+\text{H}]^+$.







(f) (+)-*N*-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride (**38a-ent-B**)







(+)-*N*-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide (**38-ent-B**) (125 mg, 0.42 mmol) was treated with 2M HCl /diethyl ether in diethyl ether/MeOH (10 mL/0.5 mL) using procedure described for compound (**2a**) to produce (+)-*N*-(2-(methylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride (**38a-ent-B**) (109 mg, 78% yield). 400 MHz $^1\text{H-NMR}$ (CD_3OD , ppm): 7.54-7.43 (m, 5H) 7.34-7.27 (m, 4H) 4.69 (s, 4H) 4.30 (dd, $J=7.8, 4.5$ Hz, 1H) 3.81 (dd, $J=14.8, 7.8$ Hz, 1H) 3.63 (dd, $J=14.8, 4.5$ Hz, 1H) 2.55 (s, 3H); ESI-MS (m/z): 296 $[\text{M}+\text{H}]^+$; $[\alpha]_D^{20}$: +27.9 (0.66, methanol).






Other examples:







Using the synthetic methods illustrated above, the following compounds of this invention were prepared:







Cpd #	Name (free base structures shown)	MP	HMR	MS / $[\alpha]$
39a	3-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-1-methyl-1-(S)-1,2,3,4-tetrahydro-naphthalen-1-yl-urea hydrochloride 	hygroscopic powder	300 MHz $^1\text{H-NMR}$ (CD_3OD , ppm) 7.79 (d, $J=2.6$ Hz, 1H) 7.69-7.64 (m, 1H) 7.31 (dd, $J=5.1, 1.0$ Hz, 1H) 7.18-7.07 (m, 3H) 7.02-6.93 (m, 1H) 5.46 (s, 1H) 4.81-4.69 (m, 1H), 4.15-4.02 (m, 1H) 3.80-3.62 (m, 1H) 3.01-2.69 (m, 8H) 2.56 (s, 1.3H) 2.54 (s, 1.7H) 2.09-1.90 (m, 2H) 1.87-1.69 (m, 2H)	358 $[\text{M}+\text{H}]^+$
40a	(R)-3-Phenyl-pyrrolidine-1-carboxylic acid (2-dimethylamino-2-thiophen-3-yl-ethyl)-amide hydrochloride 	foam	300 MHz $^1\text{H-NMR}$ (CD_3OD , ppm) 7.81-7.76 (m, 1H) 7.65 (dd, $J=5.0, 2.9$ Hz, 1H) 7.39-7.15 (m, 6H) 4.75-4.62 (m, 1H) 4.06 (ddd, $J=14.9, 8.7, 2.7$ Hz, 1H) 3.89-3.73 (m, 1H) 3.67-3.54 (m, 2H) 3.52-3.37 (m, 2H) 3.38-3.24 (m, 1H, overlapped with MeOD) 2.96-2.67 (m, 6H) 2.40-2.26 (m, 1H) 2.15-1.97 (m, 1H)	344 $[\text{M}+\text{H}]^+$
41a	(S)-3-Phenyl-pyrrolidine-1-carboxylic acid (2-dimethylamino-2-thiophen-3-yl-ethyl)-amide hydrochloride 	hygroscopic powder	300 MHz $^1\text{H-NMR}$ (CD_3OD , ppm) 7.82-7.75 (m, 1H) 7.65 (dd, $J=4.9, 2.8$ Hz, 1H) 7.36-7.18 (m, 6H) 4.76-4.65 (m, 1H) 4.15-3.97 (m, 1H) 3.89-3.73 (m, 1H) 3.69-3.52 (m, 2H) 3.52-3.38 (m, 2H) 3.38-3.22 (m, 1H, overlapped with methanol) 2.88 (s, 3H) 2.75 (s, 3H) 2.41-2.24 (m, 1H) 2.16-1.99 (m, 1H)	344 $[\text{M}+\text{H}]^+$







42a	(S)-3-Phenyl-pyrrolidine-1-carboxylic acid [2-dimethylamino-2-(4-methoxy-phenyl)-ethyl]-amide hydrochloride 	foam	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.48-7.38 (m, 2H) 7.37-7.18 (m, 5H) 7.12-7.00 (m, 2H) 4.46 (dd, J=7.6, 5.8 Hz, 1H) 4.08 (ddd, J = 14.8, 8.4, 3.0 Hz, 1H) 3.88-3.70 (m, 1H) 3.84 (s, 3H) 3.65-3.52 (m, 2H) 3.51-3.37 (m, 2H) 3.37-3.25 (m, 1H, overlapped with MeOD) 2.77 (s, 6H) 2.37-2.27(m, 1H) 2.13-1.97 (m, 1H)	368 [M+H] ⁺
43a	(R)-3-Phenyl-pyrrolidine-1-carboxylic acid [2-dimethylamino-2-(4-methoxy-phenyl)-ethyl]-amide hydrochloride 	foam	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.48-7.38 (m, 2H) 7.37-7.18 (m, 5H) 7.12-7.00 (m, 2H) 4.52-4.41 (m, 1H) 4.08 (ddd, J=14.8, 8.4, 3.1 Hz, 1H) 3.88-3.70 (m, 1H) 3.84 (s, 3H) 3.67-3.51 (m, 2H) 3.49-3.38 (m, 2H) 3.37-3.25 (m, 1H, overlapped with methanol) 2.77 (s, 6H) 2.37-2.27 (m, 1H) 2.13-1.97 (m, 1H)	368 [M+H] ⁺
44a	4-Cyano-1,3-dihydro-isoindole-2-carboxylic acid (2-dimethylamino-2-thiophen-3-yl-ethyl)-amide hydrochloride 	159-161	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.82 (dd, J=2.9, 1.3 Hz, 1H) 7.75-7.61 (m, 3H) 7.56-7.47 (m, 1H) 7.33 (dd, J=5.0, 1.3 Hz, 1H) 4.93-4.87 (m, 2H) 4.83-4.78 (m, 2H) 4.78 (dd, J=9.0, 5.1 Hz, 1H) 4.13 (dd, J=15.1, 9.0 Hz, 1H) 3.66 (dd, J=15.1, 5.0 Hz, 1H) 2.83 (s, 6H)	341 [M+H] ⁺
45a	3-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-1-methyl-1-(R)-1,2,3,4-tetrahydro-naphthalen-1-yl-urea hydrochloride 	foam	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.83-7.77 (m, 1H) 7.70-7.63 (m, 1H) 7.32 (d, J=5.0, 1H) 7.17-7.06 (m, 3H) 7.02-6.93 (m, 1H) 5.46 (s, 1H) 4.82-4.70 (m, 1H) 4.09 (ddd, J=14.5, 7.6, 6.1 Hz, 1H) 3.77-3.64 (m, 1H) 3.01-2.62 (m, 8H) 2.57-2.51 (m, 3H) 2.09-1.91 (m, 2H) 1.88-1.68 (m, 2H).	358 [M+H] ⁺
46a	5-Fluoro-1,3-dihydro-isoindole-2-carboxylic acid (1-dimethylamino-cyclohexylmethyl)-amide hydrochloride 	foam	400 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.33 (dd, J=8.4, 5.0 Hz, 1H) 7.12-7.01 (m, 2H) 4.75-4.69 (m, 4H) 3.69 (s, 2H) 2.86 (s, 6H) 1.97-1.86 (m, 2H) 1.85-1.55 (m, 7H) 1.38-1.26 (m, 1H).	320 [M+H] ⁺
47a	1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(R)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea hydrochloride diast-A 	amorphous powder	400 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.53 (dd, J=5.0, 2.9 Hz, 1H) 7.50 (s, 1H) 7.17 (dd, J=5.0, 1.3 Hz, 1H) 7.07-7.04 (m, 3H) 7.04-7.00 (m, 1H) 4.19-4.08 (m, 1H) 3.99-3.89 (m, 1H) 3.85 (dd, J=14.1, 7.1 Hz, 1H) 3.50	344 [M+H] ⁺







			(dd, J=14.1, 6.8 Hz, 1H) 3.02 (dd, J=16.2, 5.0 Hz, 1H) 2.89-2.82 (m, 2H) 2.60 (dd, J=16.2, 8.6 Hz, 1H) 2.49 (s, 6H) 2.06-1.95 (m, 1H) 1.77-1.63 (m, 1H).	
48a	1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(R)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea hydrochloride diast-B 	amorphous powder	400 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.77 (dd, J=2.9, 1.4 Hz, 1H) 7.66 (dd, J=5.0, 2.9 Hz, 1H) 7.28 (dd, J=5.0, 1.4 Hz, 1H) 7.08-7.05 (m, 3H) 7.05-7.01 (m, 1H) 4.64 (dd, J=7.5, 6.0 Hz, 1H) 4.00 (dd, J=14.9, 7.5 Hz, 1H) 4.02-3.94 (m, 1H) 3.62 (dd, J=14.9, 6.0 Hz, 1H) 3.04 (dd, J=16.2, 5.1 Hz, 1H) 2.90 (s, 3H) 2.91-2.85 (m, 2H) 2.73 (s, 3H) 2.64 (dd, J=16.2, 8.5 Hz, 1H) 2.11-1.96 (m, 1H) 1.80-1.66 (m, 1H).	344 [M+H] ⁺
49a	7-Hydroxy-3,4-dihydro-2H-quinoline-1-carboxylic acid (2-dimethylamino-2-thiophen-3-yl-ethyl)-amide hydrochloride 	foam	400 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.76 (dd, J=2.9, 1.3 Hz, 1H) 7.64 (dd, J=5.1, 2.9 Hz, 1H) 7.28 (dd, J=5.1, 1.3 Hz, 1H) 6.97-6.91 (m, 1H) 6.75 (d, J=2.4 Hz, 1H) 6.52 (dd, J=8.3, 2.4 Hz, 1H) 4.71 (dd, J=8.0, 6.0 Hz, 1H) 4.07 (dd, J=14.6, 8.0 Hz, 1H) 3.66 (dd, J=14.6, 6.0 Hz, 1H) 3.64-3.57 (m, 2H) 2.81 (s, 6H) 2.63 (t, J=6.5 Hz, 2H) 1.93-1.81 (m, 2H).	346 [M+H] ⁺
50a	1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(R)-7-hydroxy-1,2,3,4-tetrahydro-naphthalen-2-yl-urea hydrochloride 	foam	400 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.77 (dd, J=3.0, 1.3 Hz, 1H) 7.65 (dd, J=5.0, 3.0 Hz, 1H) 7.29 (dd, J=5.0, 1.3 Hz, 1H) 6.88 (d, J=8.3 Hz, 1H) 6.55 (dd, J=8.3, 2.6 Hz, 1H) 6.47 (d, J=2.6 Hz, 1H) 4.70-4.60 (m, 1H) 4.08-3.88 (m, 2H) 3.67-3.56 (m, 1H) 2.95 (dd, J=16.1, 4.9 Hz, 1H) 2.89 (s, 3H) 2.80-2.74 (m, 2H) 2.74 (s, 3H) 2.57 (dd, J=16.1, 8.4 Hz, 1H) 2.07-1.92 (m, 1H) 1.77-1.60 (m, 1H).	360 [M+H] ⁺
51a	1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(S)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea hydrochloride diast B 	amorphous powder	400 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.78-7.73 (m, 1H) 7.65 (dd, J=5.0, 3.0 Hz, 1H) 7.31-7.25 (m, 1H) 7.11-6.99 (m, 4H) 4.69-4.59 (m, 1H) 4.06-3.91 (m, 1H) 4.00 (dd, J=14.9, 7.5 Hz, 1H) 3.62 (dd, J=14.9, 6.0 Hz, 1H) 3.04 (dd, J=16.2, 5.1 Hz, 1H) 2.96-2.70 (m, 8H) 2.64 (dd, J=16.2, 8.5 Hz, 1H) 2.13-1.95 (m, 1H) 1.80-1.65 (m, 1H).	344 [M+H] ⁺
52a	1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(S)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea hydrochloride diast A 	amorphous powder	400 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.77 (dd, J=2.9, 1.4 Hz, 1H) 7.66 (dd, J=5.0, 2.9 Hz, 1H) 7.28 (dd, J=5.0, 1.4 Hz, 1H) 7.11-6.99 (m, 4H) 4.64 (dd, J=7.5, 6.0 Hz, 1H) 4.00 (dd, J=14.9, 7.5 Hz, 1H) 4.02-3.90 (m, 1H) 3.62 (dd, J=14.9, 6.0 Hz, 1H) 3.04 (dd, J=16.2, 5.1 Hz, 1H) 2.96-2.70 (m, 8H) 2.64 (dd, J=16.2, 8.5 Hz, 1H) 2.10-1.98 (m, 1H) 1.80-1.62	344 [M+H] ⁺




			(m, 1H).	
53a	6-Hydroxy-3,4-dihydro-2H-quinoline-1-carboxylic acid (2-dimethylamino-2-thiophen-3-yl-ethyl)-amide hydrochloride 	163 (dec.)	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.76 (dd, J=3.0, 1.3 Hz, 1H) 7.67 (dd, J=5.1, 3.0 Hz, 1H) 7.28 (dd, J=5.1, 1.3 Hz, 1H) 6.86 (d, J=8.6 Hz, 1H) 6.59 (d, J=2.8 Hz, 1H) 6.54 (dd, J=8.6, 2.8 Hz, 1H) 4.70 (t, J=7.1 Hz, 1H) 4.06 (dd, J=14.5, 7.5 Hz, 1H) 3.73-3.44 (m, 3H) 2.82 (s, 6H) 2.64 (t, J=6.7 Hz, 2H) 1.94-1.79 (m, 2H).	346 [M+H] ⁺
54a	1,3-Dihydro-isoindole-2-carboxylic acid (1-dimethylamino-cyclohexylmethyl)-amide hydrochloride 	225-227	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.32 (s, 4H) 4.75 (s, 4H) 3.70 (s, 2H) 2.89 (s, 6H) 2.00-1.89 (m, 2H) 1.88-1.52 (m, 7H) 1.42-1.24 (m, 1H).	302 [M+H] ⁺
55a	(R)-1-((1-(Dimethylamino)cyclohexyl)methyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea hydrochloride 	192-194	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.15-6.98 (m, 4H) 4.06-3.93 (m, 1H) 3.63 (s, 2H) 3.06 (dd, J=16.3, 5.1 Hz, 1H) 2.93-2.80 (m, 2H) 2.86 (s, 6H) 2.66 (dd, J=16.3, 8.4 Hz, 1H) 2.12-1.99 (m, 1H) 1.96-1.86 (m, 2H) 1.85-1.44 (m, 8H) 1.41-1.20 (m, 1H).	330 [M+H]
56a	(S)-1-((1-(Dimethylamino)cyclohexyl)methyl)-3-(1,2,3,4-tetrahydronaphthalen-2-yl)urea hydrochloride 	191-193	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.11-6.99 (m, 4H) 4.06-3.93 (m, 1H) 3.63 (s, 2H) 3.06 (dd, J=16.2, 5.1 Hz, 1H) 2.93-2.80 (m, 2H) 2.87 (s, 6H) 2.67 (dd, J=16.2, 8.4 Hz, 1H) 2.13-1.98 (m, 1H) 1.96-1.86 (m, 2H) 1.85-1.44 (m, 8H) 1.40-1.22 (m, 1H).	330 [M+H] ⁺
57a	(-)-N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride 	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.57-7.52 (m, 5H) 7.37-7.25 (m, 4H) 4.76-4.62 (m, 4H) 4.58 (dd, J=8.2, 5.1 Hz, 1H) 4.16 (dd, J=15.1, 8.2 Hz, 1H) 3.72 (dd, J=15.1, 5.1 Hz, 1H) 2.94 (s, 3H) 2.77 (s, 3H).	310 [M+H] ⁺ [α] = - 2.4 (0.86, acetone), free amine
58a	(+)-N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide hydrochloride	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.57-7.52 (m, 5H) 7.37-7.25 (m, 4H) 4.76-4.62 (m, 4H) 4.57 (dd, J=8.1, 5.1 Hz, 1H) 4.16 (dd, J=15.0, 8.1 Hz, 1H) 3.71 (dd, J=15.0, 5.1 Hz, 1H)	310 [M+H] ⁺ [α] = + 3.4 (0.86,

			3.05-2.62 (m, 6H).	acetone), free amine
59a	1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(5-hydroxy-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea hydrochloride 	foam	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.82-7.74 (m, 1H) 7.66 (dd, J=5.0, 2.9 Hz, 1H) 7.30 (dd, J=5.0, 1.3 Hz, 1H) 6.95 (dd, J=7.9, 7.9 Hz, 1H) 6.73 (d, J=7.9 Hz, 1H) 6.62 (d, J=7.9 Hz, 1H) 4.68, (q, J=6.6 Hz, 1H) 4.04 (dd, J=14.6, 7.5 Hz, 1H) 3.76-3.52 (m, 2H) 2.91 (s, 3H) 2.75 (s, 3H) 2.69-2.50 (m, 2H) 1.96-1.82 (m, 2H) 1.82-1.71 (m, 2H).	360 [M+H] ⁺
60a	1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-((S)-7-hydroxy-1,2,3,4-tetrahydro-naphthalen-2-yl)-urea hydrochloride 	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.76 (dd, J=2.9, 1.3 Hz, 1H) 7.65 (dd, J=5.1, 2.9 Hz, 1H) 7.28 (dd, J=5.1, 1.3 Hz, 1H) 6.88 (d, J=8.2 Hz, 1H) 6.55 (dd, J=8.2, 2.6 Hz, 1H) 6.47 (d, J=2.6 Hz, 1H) 4.64 (ddd, J=8.0, 6.0, 2.3 Hz, 1H) 4.08-3.87 (m, 2H) 3.61 (ddd, J=14.8, 6.0, 4.0 Hz) 2.95 (dd, J=16.3, 5.1 Hz, 1H) 2.90-2.65 (m, 8H) 2.56 (dd, J=16.3, 8.4 Hz, 1H) 2.07-1.91 (m, 1H) 1.79-1.61 (m, 1H).	360 [M+H] ⁺
61a	5-Fluoro-1,3-dihydro-isoindole-2-carboxylic acid (4-dimethylamino-tetrahydro-pyran-4-ylmethyl)-amide hydrochloride 	241-243	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.33 (dd, J=8.4, 5.0 Hz, 1H) 7.13-6.99 (m, 2H) 4.77-4.69 (m, 4H) 4.04-3.94 (m, 2H) 3.84 (s, 2H) 3.76-3.64 (m, 2H) 2.93 (s, 6H) 2.07-1.85 (m, 4H).	322 [M+H] ⁺
62a	5-Fluoro-1,3-dihydro-isoindole-2-carboxylic acid (2-dimethylamino-2-methyl-propyl)-amide hydrochloride 	200-202	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.33 (dd, J = 8.4, 4.9 Hz, 1H) 7.13-6.99 (m, 2H) 4.76-4.69 (m, 4H) 3.53 (s, 2H) 2.89 (s, 6H) 1.39 (s, 6H).	280 [M+H] ⁺
63a	5-Fluoro-1,3-dihydro-isoindole-2-carboxylic acid (1-dimethylamino-cyclopentylmethyl)-amide hydrochloride 	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.33 (dd, J= 8.4, 4.9 Hz, 1H) 7.12-7.01 (m, 2H) 4.76-4.65 (m, 4H) 3.59 (s, 2H) 2.95 (s, 6H) 2.09-1.74 (m, 8H).	306 [M+H]

64a	1,3-Dihydro-isoindole-2-carboxylic acid (1-dimethylamino-cyclopentylmethyl)-amide hydrochloride 	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.39-7.22 (m, 4H) 4.74 (s, 4H) 3.59 (s, 2H) 2.95 (s, 6H) 2.09-1.76 (m, 8H).	288 [M+H] ⁺
65a	1-(1-Dimethylamino-cyclopentylmethyl)-3-(S)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea hydrochloride 	204-206	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.12-6.99 (m, 4H) 4.05-3.91 (m, 1H) 3.51 (s, 2H) 3.05 (dd, J=16.2, 5.1 Hz, 1H) 2.95-2.83 (m, 2H) 2.91 (s, 6H) 2.66 (dd, J=16.2, 8.5 Hz, 1H) 2.14-1.98 (m, 1H) 1.98-1.66 (m, 9H).	316 [M+H] ⁺
66a	1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(R)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea hydrochloride 	164-166	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.77 (dd, J=2.9, 1.1 Hz, 1H) 7.65 (dd, J=5.0, 2.9 Hz, 1H) 7.29 (dd, J=5.0, 1.1 Hz, 1H) 7.11-6.98 (m, 4H) 4.70-4.60 (m, 1H) 4.08-3.94 (m, 2H) 3.62 (ddd, J=14.8, 5.8, 3.8 Hz, 1H) 3.04 (dd, J=16.2, 4.9 Hz, 1H) 2.95-2.69 (m, 8H) 2.64 (dd, J=16.2, 8.6 Hz, 1H) 2.10-1.09 (m, 1H) 1.80-1.64 (m, 1H).	344 [M+H] ⁺
67a	1-(2-Dimethylamino-2-thiophen-3-yl-ethyl)-3-(S)-1,2,3,4-tetrahydro-naphthalen-2-yl-urea hydrochloride 	165-167	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.80-7.73 (m, 1H) 7.65 (dd, J=5.0, 2.9 Hz, 1H) 7.29 (dd, J=5.0, 1.1 Hz, 1H) 7.11-6.98 (m, 4H) 4.69-4.59 (m, 1H) 4.07-3.90 (m, 2H) 3.62 (ddd, J=14.8, 5.8, 3.8 Hz, 1H) 3.04 (dd, J=16.3, 5.0 Hz, 1H) 2.91-2.84 (m, 2H) 2.80 (s, 6H) 2.64 (dd, J=16.3, 8.7 Hz, 1H) 2.10-1.97 (m, 1H) 1.80-1.64 (m, 1H).	344 [M+H] ⁺
68a	N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide hydrochloride 	145-146	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.47-7.39 (m, 2H) 7.12-6.99 (m, 3H) 6.89 (dd, J=9.0, 3.0 Hz, 1H) 6.83-6.71 (m, 2H) 4.52 (t, J=7.2, 1H) 4.15-4.04 (m, 1H) 3.86 (s, 3H) 3.76-3.47 (m, 3H) 2.91 (s, 3H) 2.72 (s, 3H) 2.80-2.60 (m, 2H) 1.88 (pentet, J=6.5 Hz, 2H).	372 [M+H] ⁺
69a	N-(2-(Dimethylamino)-2-phenylethyl)-3,4-dihydroquinoline-1(2H)-carboxamide hydrochloride 	very hygroscopic foam	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.61-7.46 (m, 5H) 7.17-7.09 (m, 1H) 7.07-6.91 (m, 3H) 4.57 (t, J=7.0, 1H) 4.12 (dd, J=14.4, 7.0 Hz, 1H) 3.74 (dd, J=14.4, 7.0 Hz, 1H) 3.69-3.44 (m, 2H) 2.86 (s, 6H) 2.74-2.59 (m, 2H) 1.88 (pentet, J=6.5 Hz, 2H).	324 [M+H] ⁺

70a	N-(2-(Dimethylamino)-2-phenylethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide hydrochloride 	very hygroscopic foam	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.61-7.46 (m, 5H) 7.00 (dd, J=9.0, 5.1 Hz, 1H) 6.88 (dd, J=9.0, 3.0 Hz, 1H) 6.78 (ddd, J=9.0, 9.0, 3.0 Hz, 1H) 4.56 (t, J=7.0, 1H) 4.10 (dd, J=14.4, 7.0 Hz, 1H) 3.73 (dd, J=14.4, 7.0 Hz, 1H) 3.69-3.48 (m, 2H) 2.85 (s, 6H) 2.74-2.62 (m, 2H) 1.88 (pentet, J=6.4 Hz, 2H).	342 [M+H] ⁺
71a	1-((R)-3-(dimethylamino)-3-(thiophen-3-yl)propyl)-3-((R)-1,2,3,4-tetrahydronaphthalen-2-yl)urea hydrochloride 	112-114	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.79-7.74 (m, 1H) 7.64 (dd, J=5.0, 2.9 Hz, 1H) 7.33-7.27 (m, 1H) 7.16-6.99 (m, 4H) 4.55 (dd, J=8.9, 6.5 Hz, 1H) 4.01-3.86 (m, 1H) 3.37-3.13 (m, 1H) 3.12-2.97 (m, 2H) 2.93-2.82 (m, 2H) 2.78 (s, 3H) 2.72 (s, 3H) 2.65 (dd, J=16.3, 8.9 Hz, 1H) 2.45-2.14 (m, 2H) 2.11-1.96 (m, 1H), 1.80-1.62 (m, 1H).	358 [M+H] ⁺
72a	1-((R)-3-(dimethylamino)-3-(thiophen-3-yl)propyl)-3-((S)-1,2,3,4-tetrahydronaphthalen-2-yl)urea hydrochloride 	147-149	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.77 (dd, J=3.0, 1.3 Hz, 1H) 7.63 (dd, J=5.1, 3.0 Hz, 1H) 7.30 (dd, J=5.1, 1.3 Hz, 1H) 7.11-6.98 (m, 4H) 4.55 (dd, J=9.2, 6.2 Hz, 1H) 4.01-3.86 (m, 1H) 3.37-3.13 (m, 1H) 3.12-2.96 (m, 2H) 2.94-2.84 (m, 2H) 2.74 (s, 6H) 2.63 (dd, J=16.2, 8.7 Hz, 1H) 2.45-2.14 (m, 2H) 2.11-1.96 (m, 1H) 1.81-1.64 (m, 1H).	358 [M+H] ⁺
73a	1-((S)-3-(dimethylamino)-3-(thiophen-3-yl)propyl)-3-((R)-1,2,3,4-tetrahydronaphthalen-2-yl)urea hydrochloride 	147-148	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.77 (dd, J=2.9, 1.4 Hz, 1H) 7.64 (dd, J=5.0, 2.9 Hz, 1H) 7.30 (dd, J=5.0, 1.4 Hz, 1H) 7.11-7.00 (m, 4H) 4.55 (dd, J=9.1, 6.4 Hz, 1H) 4.01-3.86 (m, 1H) 3.36-3.16 (m, 1H) 3.13-2.98 (m, 2H) 2.94-2.84 (m, 2H) 2.78 (s, 3H) 2.73 (s, 3H) 2.63 (dd, J=16.2, 8.7 Hz, 1H) 2.44-2.17 (m, 2H) 2.11-1.96 (m, 1H) 1.83-1.64 (m, 1H).	358 [M+H] ⁺
74a	1-((S)-3-(dimethylamino)-3-(thiophen-3-yl)propyl)-3-((S)-1,2,3,4-tetrahydronaphthalen-2-yl)urea hydrochloride 	144-146	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.82-7.74 (m, 1H) 7.67-7.59 (m, 1H) 7.31 (d, J=5.0, 1H) 7.11-7.00 (m, 4H) 4.62-4.49 (m, 1H) 4.01-3.86 (m, 1H) 3.36-3.15 (m, 1H) 3.11-2.95 (m, 2H) 2.93-2.82 (m, 2H) 2.78 (s, 3H) 2.73 (s, 3H) 2.65 (dd, J=16.2, 8.8 Hz, 1H) 2.44-2.17 (m, 2H) 2.11-1.96 (m, 1H) 1.80-1.64 (m, 1H).	358 [M+H] ⁺
75a	(R)-N-((R)-3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-3-phenylpyrrolidine-1-carboxamide hydrochloride 	hygroscopic foam	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.81-7.73 (m, 1H) 7.63 (dd, J=5.1, 2.9 Hz, 1H) 7.39-7.18 (m, 6H) 4.58 (dd, J=9.0, 6.3 Hz, 1H) 3.81-3.68 (m, 1H) 3.63-3.49 (m, 1H) 3.49-3.11 (m, 5H) 2.77 (s, 3H) 2.72 (s, 3H) 2.46-2.20 (m, 3H) 2.13-1.96 (m, 1H).	358 [M+H] ⁺

				
76a	(S)-N-((R)-3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-3-phenylpyrrolidine-1-carboxamide hydrochloride 	hygroscopic foam	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.77 (dd, J=3.0, 1.3 Hz, 1H) 7.64 (dd, J=5.1, 3.0 Hz, 1H) 7.39-7.18 (m, 6H) 4.58 (dd, J=9.0, 6.3 Hz, 1H) 3.81-3.68 (m, 1H) 3.62-3.51 (m, 1H) 3.49-3.11 (m, 5H) 2.76 (s, 3H) 2.73 (s, 3H) 2.47-2.18 (m, 3H) 2.15-1.97 (m, 1H).	358 [M+H] ⁺
77a	(S)-N-((S)-3-(Dimethylamino)-3-(thiophen-3-yl)propyl)-3-phenylpyrrolidine-1-carboxamide hydrochloride 	hygroscopic foam	300 MHz ¹ H-NMR (CD ₃ OD, ppm) 7.77 (dd, J=3.0, 1.3 Hz, 1H) 7.64 (dd, J=5.1, 3.0 Hz, 1H) 7.35-7.20 (m, 6H) 4.58 (dd, J=9.0, 6.3 Hz, 1H) 3.81-3.68 (m, 1H) 3.62-3.51 (m, 1H) 3.51-3.11 (m, 5H) 2.76 (s, 3H) 2.73 (s, 3H) 2.45-2.21 (m, 3H) 2.15-1.97 (m, 1H).	358 [M+H] ⁺
78a	(-)-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride 	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm): 7.69-7.64 (m, 1H) 7.62 (dd, J=5.0, 3.0 Hz, 1H) 7.36-7.23 (m, 4H) 7.26 (dd, J=5.0, 1.2 Hz, 1H) 4.70 (s, 4H) 4.53 (dd, J=7.8, 4.5 Hz, 1H) 3.83 (dd, J=14.8, 7.8 Hz, 1H) 3.66 (dd, J=14.8, 4.5 Hz, 1H) 2.57 (s, 3H).	302 [M+H] ⁺
79a	(+)-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride 	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm): 7.67-7.64 (m, 1H) 7.62 (dd, J=5.0, 3.0 Hz, 1H) 7.35-7.29 (m, 4H) 7.26 (dd, J=5.0, 1.2 Hz, 1H) 4.70 (s, 4H) 4.51 (dd, J=7.8, 4.5 Hz, 1H) 3.82 (dd, J=14.8, 7.8 Hz, 1H) 3.66 (dd, J=14.8, 4.5 Hz, 1H) 2.57 (s, 3H).	302 [M+H] ⁺
80a	(-)-4-fluoro-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride 	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm): 7.69-7.65 (m, 1H), 7.63 (dd, J=5.0, 3.1 Hz, 1H) 7.40-7.29 (m, 1H) 7.26 (dd, J=5.0, 1.2 Hz, 1H) 7.19-7.11 (m, 1H) 7.08-6.99 (m, 1H) 4.75 (s, 4H) 4.53 (dd, J=7.8, 4.5 Hz, 1H) 3.83 (dd, J=14.8, 7.8 Hz, 1H) 3.66 (dd, J=14.8, 4.5 Hz, 1H) 2.58 (s, 3H).	320 [M+H] ⁺

81a	(+)-4-fluoro-N-(2-(methylamino)-2-(thiophen-3-yl)ethyl)isoindoline-2-carboxamide hydrochloride  (+)	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm): 7.70-7.65 (m, 1H), 7.63 (dd, J=5.0, 3.0 Hz, 1H) 7.40-7.30 (m, 1H) 7.27 (dd, J=5.0, 1.2 Hz, 1H) 7.19-7.11 (m, 1H) 7.09-6.98 (m, 1H) 4.75 (s, 4H) 4.54 (dd, J=7.8, 4.5 Hz, 1H) 3.83 (dd, J=14.8, 7.8 Hz, 1H) 3.66 (dd, J=14.8, 4.5 Hz, 1H) 2.58 (s, 3H).	320 [M+H] ⁺
82a	(-)-5-fluoro-N-(2-(4-methoxyphenyl)-2-(methylamino)ethyl)isoindoline-2-carboxamide hydrochloride  (-)	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm): 7.34-7.36 (m, 2H) 7.32 (dd, J=8.3, 5.0 Hz, 1H) 7.12-6.99 (m, 4H) 4.77-4.57 (m, 4H) 4.32 (dd, J=8.1, 4.6 Hz, 1H) 3.84 (dd, J=14.8, 8.1 Hz, 1H) 3.82 (s, 3H) 3.61 (dd, J=14.8, 4.6, 1H) 2.55 (s, 3H).	344 [M+H] ⁺
83a	(+)-5-fluoro-N-(2-(4-methoxyphenyl)-2-(methylamino)ethyl)isoindoline-2-carboxamide hydrochloride  (+)	amorphous powder	300 MHz ¹ H-NMR (CD ₃ OD, ppm): 7.50-7.39 (m, 2H) 7.34 (dd, J=8.3, 5.0 Hz, 1H) 7.16-7.01 (m, 4H) 4.79-4.58 (m, 4H) 4.34 (dd, J=8.1, 4.6 Hz, 1H) 3.87 (dd, J=14.8, 8.1 Hz, 1H) 3.85 (s, 3H) 3.64 (dd, J=14.8, 4.6, 1H) 2.55 (s, 3H).	344 [M+H] ⁺

Example 33: Bias factor of selected compounds of the invention

$$\Delta RA = \text{LOG} \left(\frac{EMAX_{\beta\text{-Arr}}}{EC50_{\beta\text{-Arr}}} \right) - \text{LOG} \left(\frac{EMAX_{G\text{-Protein}}}{EC50_{G\text{-Protein}}} \right)$$

CMPD_ID; EX #	cAMP EC ₅₀ (uM)	cAMP E _{max} (%)	β-Arrestin EC ₅₀ (uM)	β-Arrestin E _{max} (%)	ΔRA of compound	Bias Factor (ΔRA of compound relative to ΔRA of DAMGO) x10
DAMGO	0.00	108.2	0.06	98.33	-1.48	0.00
Morphine	0.06	119.5	0.31	28.80	-1.33	1.51
(1a); Ex 3	0.020	69.2	0.186	5.67	-2.06	-5.75
(2a-ent-(-)); Ex 5	0.012, 0.015	55.5, 30.4	>100	<1	<-2.96	<-14.83

(2a-ent-(+)); Ex 6	0.018, 0.058	99.5, 95.4	1.08, 1.01	6.57, 13.7	-2.40	-9.24
(3a); Ex 7	0.014	94.2	0.310	21.90	-1.99	-5.07
(4a), Ex 8	>100	20.7	>100	<1	NA	NA
(5a); Ex 9	>100	5.00	>100	<1	NA	NA
(6a); Ex 10	0.009	21.9	>100	<1	<-2.96	<-14.83
(7a); Ex 11	0.407	92.1	>100	<1	<-2.96	<-14.83
(8a); Ex 12	0.730	86.9	>100	6.18	<-2.96	<-14.83
(9a); Ex 13	0.512	75.7	4.36	5.11	-2.10	-6.20
(10a); Ex 14	0.004	85.5	>100	3.33	<-2.96	<-14.83
(11a); Ex 15	<0.001	94.3	0.021	111.9	-1.24	2.32
(12a); Ex 16	>100	6.33	>100	2.16	NA	NA
(13a) Ex 17	0.068	95.3	1.68	11.3	-2.3	-8.39
(14a) Ex 18	0.056	102.0	3.96	61.2	-2.07	-5.91
(15a) Ex 19	0.015	108.5	1.37	44.2	-2.35	-8.71
(16a) Ex 20	0.026	91.9	0.95	12.2	-2.44	-9.60
(17a) Ex 21	0.043	110.6	5.12	27.0	-2.69	-12.08
(18a) Ex 22	0.023	100.4	0.73	6.91	-2.66	-11.84
(19a) Ex 23	0.032	86.1	>100	2.34	<-2.96	<-14.83
(20a) Ex 24	0.031	70.8	>100	3.65	<-2.96	<-14.83
(68a)	0.167	94.6	14.589	39.9	-2.32	-8.36
(69a)	0.563	109.6	8.556	8.0	-2.32	-8.38
(70a)	0.512	90.3	10.967	5.3	-2.56	-10.82
(30-R) Ex 28	0.603	109.4	5.729	9.2	-2.05	-5.73
(30-S) Ex 29	0.688	96.1	>100	9.9	<-2.96	<-14.83
(66)	0.008	80.5	>100	4.1	<-2.96	<-14.83
(67)	0.013	50.9	>100	2.6	<-2.96	<-14.83
(34) Ex 30	>100	2.6	>100	<1	NA	NA
(35) Ex 31	1.171	26.4	>100	<1	<-2.96	<-14.83
(72)	0.197	111.9	2.611	39.1	-1.58	-0.99

(71)	0.177	71.2	1.823	10.5	-1.85	-3.66
(39)	0.118	93.4	0.644	10.9	-1.67	-1.90
(40)	0.152	53.5	>100	3.3	<-2.96	<-14.83
(73)	0.422	13.6	>100	<1	<-2.96	<-14.83
(42)	7.429	21.9	>100	<1	<-2.96	<-14.83
(43)	1.716	54.2	>100	<1	<-2.96	<-14.83
(41)	>100	9.6	>100	<1	NA	NA
(74)	>100	11.2	>100	<1	NA	NA
(44)	0.162	51.4	>100	5.3	<-2.96	<-14.83
(45)	0.276	101.4	>100	8.1	<-2.96	<-14.83
(75)	2.088	113.5	16.956	12.2	-1.88	-3.97
(76)	1.761	33.0	>100	<1	<-2.96	<-14.83
(21) Ex 25	>100	4.5	>100	<1	NA	NA
(77)	1.097	33.2	>100	<1	<-2.96	<-14.83
(46)	0.522	108.7	10.075	13.6	-2.19	-7.08
(47)	0.010	93.0	>100	<1	<-2.96	<-14.83
(48)	0.260	31.8	>100	<1	<-2.96	<-14.83
(22) Ex 26	0.322	16.3	>100	<1	<-2.96	<-14.83
(24) Ex 27	>100	<10	>100	<1	<-2.96	<-14.83
(49)	0.998	97.0	>100	<1	<-2.96	<-14.83
(50)	0.036	48.3	>100	<1	<-2.96	<-14.83
(51)	0.029	64.9	>100	<1	<-2.96	<-14.83
(52)	0.227	12.2	>100	<1	<-2.96	<-14.83
(53)	0.267	119.7	3.790	20.1	-1.93	-4.47
(54)	0.396	113.4	4.424	20.8	-1.78	-3.04
(55)	2.853	36.6	>100	<1	<-2.96	<-14.83
(56)	7.737	43.3	>100	<1	<-2.96	<-14.83
(57)	0.072	89.4	>100	<1	<-2.96	<-14.83
(58)	0.132	109.1	1.887	35.2	-1.65	-1.66
(59)	0.006	108.2	0.160	61.0	-1.67	-1.95
(60)	0.676	27.5	>100	<1	<-2.96	<-14.83
(61)	1.174	103.9	>100	14.5	<-2.96	<-14.83
(62)	21.009	81.8	>100	<1	<-2.96	<-14.83
(63)	2.511	87.9	>100	<1	<-2.96	<-14.83
(64)	2.151	88.7	>100	<1	<-2.96	<-14.83
(65)	>100	<10	>100	<1	NA	NA
(38) Ex 32	0.045	95.9	1.674	6.6	-2.74	-12.56


(38-ent-A) Ex 32	0.379	81.6	>100	3.7	<-2.96	<-14.83
(38-ent-B) Ex 32	0.840	97.6	>100	5.9	<-2.96	<-14.83
(78)	0.362	104.0	>100	12.8	<-2.96	<-14.83
(79)	0.087	64.2	>100	1.7	<-2.96	<-14.83
(80)	0.041	102.9	1.396	25.2	-2.14	-6.64
(81)	0.014	91.6	>100	8.2	<-2.96	<-14.83
(82)	0.457	83.2	>100	2.7	<-2.96	<-14.83
(83)	0.057	100.1	>100	13.4	<-2.96	<-14.83

The disclosures of each and every patent, patent application, and publication cited herein are hereby incorporated herein by reference in their entirety.



While this invention has been disclosed with reference to specific embodiments, it is
5 apparent that other embodiments and variations of this invention may be devised by others skilled in the art without departing from the true spirit and scope of the invention. The appended claims are intended to be construed to include all such embodiments and equivalent variations.


CLAIMS


1. (Currently amended) A compound of formula (I), or a salt, solvate, enantiomer, diastereoisomer or tautomer thereof:



wherein:

A is selected from the group consisting of , , and




B¹ is (R)_p, and B² is H;

R¹ and R² are independently selected from the group consisting of H, CH₃, and CH₃ substituted with at least one selected from the group consisting of fluoro, chloro, cyano, hydroxyl and nitro;


R³ is selected from the group consisting of H and CH₃;


each occurrence of R is independently selected from the group consisting of fluoro, chloro, bromo, iodo, cyano, nitro, hydroxyl, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, and substituted phenyl;

each occurrence of m is independently selected from the group consisting of 0, 1, 2 and 3;


n is selected from the group consisting of 1, 2 and 3; and

each occurrence of p is independently selected from the group consisting of 0, 1, 2 and 3.

2. The compound of claim 1, wherein A is .



3. The compound of claim 1, wherein A is



4. The compound of claim 1, wherein A is

5. The compound of claim 1, wherein A is selected from the group consisting of isoindolin-2-yl, 5-fluoroisoindolin-2-yl, 5-chloroisoindolin-2-yl, 5-methoxyisoindolin-2-yl, 5-methylisoindolin-2-yl, 5-hydroxyisoindolin-2-yl, 5-cyanoisoindolin-2-yl, 4-fluoroisoindolin-2-yl, 4-chloroisoindolin-2-yl, 4-methoxyisoindolin-2-yl, 4-methylisoindolin-2-yl, 4-hydroxyisoindolin-2-yl, 4-cyanoisoindolin-2-yl, 5,6-difluoroisoindolin-2-yl, 5,6-dichloroisoindolin-2-yl, 6-chloro-5-fluoroisoindolin-2-yl, 5-chloro-6-fluoroisoindolin-2-yl, 1,2,3,4-tetrahydroquinolin-1-yl, 5-fluoro-1,2,3,4-tetrahydroquinolin-1-yl, 6-fluoro-1,2,3,4-tetrahydroquinolin-1-yl, 7-fluoro-1,2,3,4-tetrahydroquinolin-1-yl, 8-fluoro-1,2,3,4-tetrahydroquinolin-1-yl, 5-hydroxy-1,2,3,4-tetrahydroquinolin-1-yl, 6-hydroxy-1,2,3,4-tetrahydroquinolin-1-yl, 7-hydroxy-1,2,3,4-tetrahydroquinolin-1-yl, 8-hydroxy-1,2,3,4-tetrahydroquinolin-1-yl, pyrrolidin-1-yl, and 3-phenyl- pyrrolidin-1-yl.

6. The compound of claim 1, wherein B¹ is selected from the group consisting of 2-chloro-4-fluorophenyl, 4-chloro-2-fluorophenyl, 4-methoxyphenyl, 4-hydroxyphenyl, and phenyl.

7. A compound, or a salt, solvate, enantiomer, diastereoisomer or tautomer thereof, selected from the group consisting of:

N-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoroisoindoline-2-carboxamide;
N-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)-5-fluoroisoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-phenylethyl)-5-fluoroisoindoline-2-carboxamide;
(S)-3-Phenyl-pyrrolidine-1-carboxylic acid [2-dimethylamino-2-(4-methoxy-phenyl)-ethyl]-amide;
(R)-3-Phenyl-pyrrolidine-1-carboxylic acid [2-dimethylamino-2-(4-methoxy-phenyl)-ethyl]-amide;
(-)-N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide;
(+)-N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide;
N-(2-(Dimethylamino)-2-phenylethyl)-3,4-dihydroquinoline-1(2H)-carboxamide;
N-(2-(Dimethylamino)-2-phenylethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide;
(-)-5-Fluoro-N-(2-(4-methoxyphenyl)-2-(methylamino)ethyl)isoindoline-2-carboxamide;
(+)-5-Fluoro-N-(2-(4-methoxyphenyl)-2-(methylamino)ethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-fluoroisoindoline-2-carboxamide;
N-(2-(dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-hydroxyisoindoline-2-carboxamide;
N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide;
(-)-N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide; and
(+)-N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide.

8. A pharmaceutical composition comprising at least one pharmaceutically acceptable carrier and at least one compound of any one of claims 1-7.

9. A compound of any one of claims 1-7 and/or a pharmaceutical composition of claim 14

for use in the treatment and/or prevention of pain.

10. Use of a compound of any one of claims 1-7 and/or a pharmaceutical composition of claim 14 in the manufacture of a medicament for the treatment and/or prevention of pain.

11. The use of claim 10, wherein the at least one compound is selected from the group consisting of:

N-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-5-fluoroisoindoline-2-carboxamide;
N-(2-(2-Chloro-4-fluorophenyl)-2-(dimethylamino)ethyl)-5-fluoroisoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-phenylethyl)-5-fluoroisoindoline-2-carboxamide;
(S)-3-Phenyl-pyrrolidine-1-carboxylic acid [2-dimethylamino-2-(4-methoxy-phenyl)-ethyl]-amide;
(R)-3-Phenyl-pyrrolidine-1-carboxylic acid [2-dimethylamino-2-(4-methoxy-phenyl)-ethyl]-amide;
(-)-N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide;
(+)-N-(2-(Dimethylamino)-2-phenylethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-(4-methoxyphenyl)ethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide;
N-(2-(Dimethylamino)-2-phenylethyl)-3,4-dihydroquinoline-1(2H)-carboxamide;
N-(2-(Dimethylamino)-2-phenylethyl)-6-fluoro-3,4-dihydroquinoline-1(2H)-carboxamide;
(-)-5-Fluoro-N-(2-(4-methoxyphenyl)-2-(methylamino)ethyl)isoindoline-2-carboxamide;
(+)-5-Fluoro-N-(2-(4-methoxyphenyl)-2-(methylamino)ethyl)isoindoline-2-carboxamide;
N-(2-(Dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-fluoroisoindoline-2-carboxamide;
N-(2-(dimethylamino)-2-(4-hydroxyphenyl)ethyl)-5-hydroxyisoindoline-2-carboxamide;
N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide;
(-)-N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide; and
(+)-N-(2-(Methylamino)-2-phenylethyl)isoindoline-2-carboxamide.

12. The use of claim 10, wherein the at least one compound is a MOR agonist.
13. The use of claim 10, wherein the at least one compound decreases cyclic adenosine monophosphate (cAMP) levels in the subject.
14. The use of claim 10, wherein the at least one compound does not significantly induce recruitment, binding to, or association with a β -arrestin.
15. The use of claim 10, wherein the at least one compound does not significantly cause at least one side effect selected from the group consisting of tachyphylaxis, respiratory depression, constipation, nausea, emesis, withdrawal, dependence, and addiction.
16. The use of claim 10, wherein the pain comprises chronic pain, neuropathic pain, nociceptive pain, hyperalgesia, and/or allodynia.