protein underlying a loss-of-function disease. In some aspects, the amyloid-forming protein is selected from the group consisting of Hsp27, αA-crystallin, αB-crystallin, βB2-crystallin, βB1-crystallin, γD-crystallin, Hsp22, Hsp20, tau, Alpha-synuclein, IAPP, beta-amyloid, PrP, Huntingtin, Calcitonin, Atrial natriuretic factor, Apolipoprotein AI, Serum amyloid A, Medin, Prolactin, Transthyretin, Lysozyme, Beta 2 microglobulin, Gelsolin, Keratoepithelin, Cystatin, Immunoglobulin light chain AL, and S-IBM. In other aspects, the protein underlying a loss-of-function disease is selected from the group consisting of mutant β-glucosidase, cystic fibrosis transmembrane receptor, hexosaminidase A, hexosaminidase B, β-galactosidase, and alpha-glucosidase.

**[0010]** Other features and advantages of the present disclosure will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments of the disclosure, are given by way of illustration only, because various changes and modifications within the spirit and scope of the disclosure will become apparent to those skilled in the art from this detailed description. The entire document is intended to be related as a unified disclosure, and it should be understood that all combinations of features described herein are contemplated, even if the combination of features are not found together in the same sentence, or paragraph, or section of this document. In addition to the foregoing, the invention includes, as an additional aspect, all embodiments of the invention narrower in scope in any way than the variations specifically mentioned above. For example, if aspects of the invention are described as "comprising" a feature, embodiments also are contemplated "consisting of or "consisting essentially of" the feature.

**[0011]** In one embodiment, this invention discloses the use of a composition in the manufacture of medicament to treat and/or prevent vision disorders in a subject, said composition comprises a pharmaceutically acceptable ophthalmic carrier and a pharmaceutically effective amount of lanosterol. Said subject is having or at risk of developing a vision disorder that affects the normal structure of the lens in the eye. Said subject may be selected from the group consisting of amphibians, reptiles, avians, and mammals; wherein said mammal may be selected from the group consisting of rodents, cats, dogs, pigs, horses and humans. In another embodiment, said vision disorder is selected from the group consisting of cataract, congenital cataracts, cortical opacities, posterior subcapsular cataract, presbyopia nuclear sclerosis, retinal degenerative disorder, Refsum disease, Smith-Lemli-Opitz syndrome, Schnyder crystalline corneal dystrophy, drusen, age-related macular degeneration, and diabetic retinopathy, and lanosterol inhibits crystallin protein aggregation.

**[0012]** In yet another embodiment, this invention discloses the use of a composition in the manufacture of medicament to treat cataract or blindness/impaired vision in a subject, said composition comprises a pharmaceutically acceptable ophthalmic carrier and a pharmaceutically effective amount of lanosterol, wherein said lanosterol dissolves lens crystallin protein aggregate(s) in the eye of said subject; wherein the lens crystallin protein is any of α–crystallin, β–crystallin or γ–crystallin. The above mentioned composition may be formulated as an ophthalmic solution, an ophthalmic ointment, an ophthalmic wash, an intraocular infusion solution, a wash for anterior chamber, an internal medicine, an injection, or preservative for extracted cornea.

**[0013]** In yet another embodiment, this invention discloses a method for dissolving amyloid-like fibrils of crystallin proteins, comprising the step of contacting the amyloid-like fibrils with lanosterol in a sufficient amount and duration so as to dissolve the amyloid-like fibrils of crystalline proteins, wherein the method may be performed in situ, in vitro or in vivo. The method may be performed on a subject selected from the group consisting of amphibians, reptiles, avians, and mammals; wherein said mammal may be selected from the group consisting of rodents, cats, dogs, pigs, horses and humans.

**[0014]** In another embodiment, this invention discloses a kit for treating and/or preventing vision disorders that affect the normal structure of the eye in a subject, comprising a formulation of a pharmaceutically effective amount of lanosterol, a pharmaceutically acceptable carrier and instructions for administering said formulation such that said administration treats and/or prevents said vision disorder. In yet another embodiment, this invention discloses an ophthalmic pharmaceutical composition for treating and/or preventing vision disorders in a subject, said composition comprises a pharmaceutically acceptable ophthalmic carrier and a pharmaceutically effective amount of lanosterol; wherein said composition may be formulated as an ophthalmic solution, an ophthalmic ointment, an ophthalmic wash, an intraocular infusion solution, a wash for anterior chamber, an internal medicine, an injection, or preservative for extracted cornea.

**[0015]** In another embodiment, this invention discloses a method for identifying and/or treating a subject at risk of developing cataract or blindness/impaired vision associated with formation of lens crystallin protein aggregate(s) in an eye, comprising: a) assaying for amount of lanosterol synthase activity in the subject; b) determining whether the amount of lanosterol synthase activity is less than that of a control population without cataract or blindness/impaired vision, wherein an amount of lanosterol synthase activity less than that of a control population

What is claimed is:

1. Use of a composition in the manufacture of medicament to treat and/or prevent vision disorders in a subject, said composition comprises a pharmaceutically acceptable ophthalmic carrier and a pharmaceutically effective amount of lanosterol.

2. The use of claim 1, wherein said subject is having or at risk of developing a vision disorder that affects the normal structure of the lens in the eye.

3. The use of claim 1, wherein said vision disorder is selected from the group consisting of cataract, congenital cataracts, cortical opacities, posterior subcapsular cataract, presbyopia nuclear sclerosis, retinal degenerative disorder, Refsum disease, Smith-Lemli-Opitz syndrome, Schnyder crystalline corneal dystrophy, drusen, age-related macular degeneration, and diabetic retinopathy.

4. The use of claim 1, wherein said lanosterol inhibits crystallin protein aggregation.

5. Use of a composition in the manufacture of medicament to treat cataract or blindness/impaired vision in a subject, said composition comprises a pharmaceutically acceptable ophthalmic carrier and a pharmaceutically effective amount of lanosterol, wherein said lanosterol dissolves lens crystallin protein aggregate(s) in the eye of said subject.

6. The use of claim 4, wherein the lens crystallin protein is any of α-crystallin, β-crystallin or γ-crystallin.

7. The use of any one of claims 1-6, wherein said composition is formulated as an ophthalmic solution, an ophthalmic ointment, an ophthalmic wash, an intraocular infusion solution, a wash for anterior chamber, an internal medicine, an injection, or preservative for extracted cornea.

8. The use of any one of claims 1-6, wherein said subject is selected from the group consisting of amphibians, reptiles, avians, and mammals.