

(BL)

PROCESSES AND INTERMEDIATES FOR PREPARING BENZYL EPOXIDESBACKGROUND OF THE INVENTION

This application claims priority from U.S. Provisional  
5 Application Serial No. 60/285,772, filed April 23, 2001.

1. Field of the Invention

The invention provides processes for preparing benzyl  
substituted epoxides useful in the preparation of biologically  
10 active compounds, as well as intermediates useful in those  
processes.

2. Description of the Related Art

International Publication WO02/02512 discloses various  
15 hydroxyethylamines useful in treating Alzheimer's disease. A  
common intermediate in most of the products is an N-protected  
epoxide. As a result, there exists a need for processes and  
intermediates that efficiently produce N-protected epoxides.




\*G00002\*



\*159159\*

SUMMARY OF INVENTION

In a broad aspect, the invention provides compounds of the formula II



II

where


R is phenyl optionally substituted with 1, 2, 3, or 4 groups independently selected from:

- (A) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or three substituents independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, hydroxy, thio, -NR<sub>10</sub>R<sub>11</sub> where R<sub>10</sub> and R<sub>11</sub> are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; cyano, trifluoromethyl, and C<sub>1</sub>-C<sub>3</sub> alkoxy,
- (B) C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl,
- (C) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3 fluoro,
- (D) -NR<sub>12</sub>R<sub>13</sub> where at each occurrence R<sub>12</sub> and R<sub>13</sub> are the same or different and represent:
  - (a) -H,
  - (b) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one of:
    - (i) -OH,
    - (ii) -NH<sub>2</sub>,
    - (iii) phenyl,
  - (c) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with 1, 2, or 3 independently selected halogens,
  - (d) -C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl; and
- (E) C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NR<sub>10</sub>R<sub>11</sub>, -C(O)NR<sub>10</sub>R<sub>11</sub>, or -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl);

R<sub>1</sub> is selected from:

(I) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one halogen;  
 (II) -CH<sub>2</sub>-CH=CH<sub>2</sub>;  
 (III) phenyl optionally substituted with one nitro,  
 halogen, or cyano; and  
 5 (IV) benzyl optionally substituted on the phenyl with  
 nitro, halogen, or cyano; and  
 R<sub>30</sub> represents hydrogen or PROT, where PROT is a nitrogen  
 protecting group.

10 In another aspect, the invention provides amino alcohols  
 of the formula:




(IV-unprotected)

where

15 R is as defined for Formula II above; and  
 R<sub>2</sub> is:  
 chloro, bromo, or  
 -Si(R<sub>21</sub>)<sub>3</sub> where each R<sub>21</sub> is independently  
 C<sub>1</sub>-C<sub>5</sub> alkyl,  
 20 -N(R<sub>23</sub>)(R<sub>24</sub>) where R<sub>23</sub> and R<sub>24</sub> are the same or  
 different and represent  
 C<sub>1</sub>-C<sub>5</sub> alkyl,  
 or where NR<sub>23</sub>R<sub>24</sub> represents piperidinyl,  
 piperazinyl, or morpholinyl,  
 25 phenyl optionally substituted with 1, 2, or 3 of C<sub>1</sub>-  
 C<sub>2</sub> alkyl, with the proviso that at least one of  
 the R<sub>21</sub> groups is optionally substituted phenyl.


In another aspect the invention provides an unprotected  
 30 epoxide of formula V-unprotected



V-unprotected

where R is as defined for Formula II.


5 The invention further provides a compound of formula XI



XI

where X<sub>1</sub> is chloro, bromo, or imidazolyl; and R and PROT are as defined above for Formula II.

10 In another aspect, the invention provides a compound of formula XIV




XIV

where R, PROT and R<sub>1</sub> are as defined above for Formula II.


15

In still another aspect, the invention provides a process for the preparation of an ester of formula II, where



the process comprises

20 (1) esterifying a protected amino acid of the formula I




I

where PROT and R are as defined above with an alkylating agent in the presence of a base.

5

In a related aspect, the invention provides another process for preparing esters of formula II:



II


10 where R<sub>1</sub> is an optionally substituted phenyl, the process comprising:

(a) forming a mixture of a protected amino acid of formula I and an activating agent;

15 (b) contacting the mixture of (a) with a phenol optionally substituted on the phenyl ring with nitro, halogen, or cyano.

In another related aspect, the invention provides a process for the preparation of an ester of the formula II, which comprises treating a compound of formula XIV with 20 hydrogen in the presence of a hydrogenation catalyst at a pressure of from 1 atmosphere to about 100 psi, and preferably in a suitable solvent.

The invention also provides a process for preparing a 25 compound of formula III



III

where PROT, R, and R<sub>2</sub> are as defined above for formula IV-unprotected, which comprises:

- 5 (a) forming a mixture of an ester of formula II and a dihalogenated methane, R<sub>2</sub>CH<sub>2</sub>X<sup>2</sup>, where R<sub>2</sub> is as defined above and where X<sup>2</sup> is -Br or -I;
- (b) adding a strong base having a pK<sub>b</sub> of greater than about 30 to the mixture from (a);
- 10 (c) acidifying the mixture of (b).

In a related aspect, the invention provides another method for preparing a compound of formula III where PROT, R, and R<sub>2</sub> are as defined above. This method comprises:

- 15 (a) forming a mixture of an acid R<sub>2</sub>-CH<sub>2</sub>-COOH and a base, preferably a strong base having a pK<sub>b</sub> of greater than about 30;
- (b) treating the mixture of (a) with an ester of formula II; and
- (c) acidifying the mixture from (b).

20

In a still further related aspect, the invention provides a process for preparing ketones of formula III, wherein R<sub>2</sub> is Cl or Br, which comprises contacting a compound of formula XI with LiCH<sub>2</sub>Cl or LiCH<sub>2</sub>Br.


25

In yet another aspect, the invention provides a process for the preparation of a compound of formula XI. This process comprises contacting a protected (S) amino acid of formula (I) with thionyl chloride, phosphorous trichloride, oxalyl chloride, phosphorous tribromide, triphenylphosphorous

30

dibromide, oxalyl bromide, 1,2-phenylenetrichlorophosphate, 2,4,6-trichloro-1,3,5-triazine or CDI.

In still yet another aspect, the invention provides a process for the preparation of a compound of formula (XX)



wherein

R<sub>57</sub> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, or benzyl;

R<sub>1</sub> is -(CH<sub>2</sub>)<sub>1-2</sub>-S(O)<sub>0-2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), -CH<sub>2</sub>-CH<sub>2</sub>-S(O)<sub>0-2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), or

C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -F, -Cl, -Br, -I, -OH, =O, -SH, -C≡N, -CF<sub>3</sub>, -C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, mono- or dialkylamino, -N(R)C(O)R'-, -OC(=O)-amino and -OC(=O)-mono- or dialkylamino, or

C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl, each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, and mono- or dialkylamino, or

aryl, heteroaryl, heterocyclyl, -C<sub>1</sub>-C<sub>6</sub> alkyl-aryl, -C<sub>1</sub>-C<sub>6</sub> alkyl-heteroaryl, or -C<sub>1</sub>-C<sub>6</sub> alkyl-heterocyclyl, where the ring portions of each are optionally substituted with 1, 2, 3, or 4 groups independently selected from halogen, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -NR<sub>7</sub>R'<sub>7</sub>, -C(=O)-(C<sub>1</sub>-C<sub>4</sub>) alkyl, -SO<sub>2</sub>-amino, -SO<sub>2</sub>-mono or dialkylamino, -C(=O)-amino, -C(=O)-mono or

dialkylamino,  $-\text{SO}_2-(\text{C}_1-\text{C}_4)$  alkyl,  $-\text{CO}_2\text{R}$ ,  $-\text{N}(\text{R})\text{COR}'$ , or  $-\text{N}(\text{R})\text{SO}_2\text{R}'$  or

$-\text{C}_1-\text{C}_6$  alkoxy optionally substituted with 1, 2, or 3 groups which are independently selected from halogen, or

$\text{C}_3-\text{C}_7$  cycloalkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ ,  $-\text{OH}$ ,  $-\text{SH}$ ,  $-\text{C}\equiv\text{N}$ ,  $-\text{CF}_3$ ,  $\text{C}_1-\text{C}_3$  alkoxy, amino,  $-\text{C}_1-\text{C}_6$  alkyl and mono- or dialkylamino, or

$\text{C}_1-\text{C}_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ ,  $-\text{OH}$ ,  $-\text{SH}$ ,  $-\text{C}\equiv\text{N}$ ,  $-\text{CF}_3$ ,  $-\text{C}_1-\text{C}_3$  alkoxy, amino, mono- or dialkylamino and  $-\text{C}_1-\text{C}_3$  alkyl, or

$\text{C}_2-\text{C}_6$  alkenyl or  $\text{C}_2-\text{C}_6$  alkynyl, each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ ,  $-\text{OH}$ ,  $-\text{SH}$ ,  $-\text{C}\equiv\text{N}$ ,  $-\text{CF}_3$ ,  $\text{C}_1-\text{C}_3$  alkoxy, amino,  $-\text{C}_1-\text{C}_6$  alkyl and mono- or dialkylamino; and the heterocyclyl group is optionally further substituted with oxo;

$\text{R}_7$  and  $\text{R}_7'$  are independently H or  $-\text{C}_1-\text{C}_6$  alkyl;

$\text{R}_2$  and  $\text{R}_3$  are independently selected from the group consisting of H;  $\text{C}_1-\text{C}_6$  alkyl optionally substituted with one, two or three substituents independently selected from the group consisting of  $\text{C}_1-\text{C}_3$  alkyl, halogen,  $-\text{OH}$ ,  $-\text{SH}$ ,  $-\text{C}\equiv\text{N}$ ,  $-\text{CF}_3$ ,  $\text{C}_1-\text{C}_3$  alkoxy, and  $-\text{NR}_{30}\text{R}_{31}$ ;  $-(\text{CH}_2)_{0-4}\text{-aryl}$ ;  $-(\text{CH}_2)_{0-4}\text{-heteroaryl}$ ;  $-(\text{CH}_2)_{0-4}\text{-heterocycle}$ ;  $\text{C}_2-\text{C}_6$  alkenyl optionally substituted with one, two or three substituents independently selected from the group consisting of  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{OH}$ ,  $-\text{SH}$ ,  $-\text{C}\equiv\text{N}$ ,  $-\text{CF}_3$ ,  $\text{C}_1-\text{C}_3$  alkoxy, and  $-\text{NR}_{30}\text{R}_{31}$ ;  $\text{C}_2-\text{C}_6$  alkynyl optionally substituted with one, two or three

substituents independently selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, and -NR<sub>30</sub>R<sub>31</sub>; and -(CH<sub>2</sub>)<sub>0-4</sub>-C<sub>3</sub>-C<sub>7</sub> cycloalkyl, wherein the cycloalkyl group is optionally substituted with one, two or three substituents independently selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, and -NR<sub>30</sub>R<sub>31</sub>;

or

R<sub>2</sub>, R<sub>3</sub> and the carbon to which they are attached form a carbocycle of three, four, five, six, or seven carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced by a heteroatom independently selected from the group consisting of -O-, -S-, -SO<sub>2</sub>-, and -NR<sub>22</sub>-; wherein R<sub>30</sub> and R<sub>31</sub> at each occurrence are independently H, or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sub>22</sub> is selected from the group consisting of -H, -C<sub>1</sub>-C<sub>6</sub> alkyl, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, amino C<sub>1</sub>-C<sub>6</sub> alkyl; halo C<sub>1</sub>-C<sub>6</sub> alkyl; -C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl, -C<sub>1</sub>-C<sub>6</sub> alkyl chain with one double bond and one triple bond, aryl, heteroaryl, and heterocycloalkyl;

R<sub>C</sub> is selected from the group consisting of C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of R<sub>205</sub>, -OC=O NR<sub>235</sub>R<sub>240</sub>, -S(=O)<sub>0-2</sub> R<sub>235</sub>, -NR<sub>235</sub>C=O NR<sub>235</sub>R<sub>240</sub>, -C=O NR<sub>235</sub>R<sub>240</sub>, and -S(=O)<sub>2</sub> NR<sub>235</sub>R<sub>240</sub>; -(CH<sub>2</sub>)<sub>0-3</sub>-(C<sub>3</sub>-C<sub>8</sub>) cycloalkyl wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of R<sub>205</sub>, -CO<sub>2</sub>H, and -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl); -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocycloalkyl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-heteroaryl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-heterocycloalkyl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-aryl;

- (CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-aryl; - (CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-heterocycloalkyl; - (CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-heteroaryl; - (CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocycloalkyl-heteroaryl; - (CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocycloalkyl-heterocycloalkyl; - (CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocycloalkyl-aryl; - [C(R<sub>255</sub>)(R<sub>260</sub>)]<sub>1-3</sub>-CO-N-(R<sub>255</sub>)<sub>2</sub>; - CH(aryl)<sub>2</sub>; -CH(heteroaryl)<sub>2</sub>; -CH(heterocycloalkyl)<sub>2</sub>; -CH(aryl)(heteroaryl); cyclopentyl, cyclohexyl, or cycloheptyl ring fused to aryl, heteroaryl, or heterocycloalkyl wherein one carbon of the cyclopentyl, cyclohexyl, or cycloheptyl is optionally replaced with NH, NR<sub>215</sub>, O, or S(=O)<sub>0-2</sub>, and wherein the cyclopentyl, cyclohexyl, or -cycloheptyl group can be optionally substituted with 1 or 2 groups that are independently R<sub>205</sub> or =O; -CO-NR<sub>235</sub>R<sub>240</sub>; or -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl); C<sub>2</sub>-C<sub>10</sub> alkenyl optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; C<sub>2</sub>-C<sub>10</sub> alkynyl optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; - (CH<sub>2</sub>)<sub>0-1</sub>-CH((CH<sub>2</sub>)<sub>0-6</sub>-OH)-(CH<sub>2</sub>)<sub>0-1</sub>-aryl; - (CH<sub>2</sub>)<sub>0-1</sub>-CHR<sub>C-6</sub>-(CH<sub>2</sub>)<sub>0-1</sub>-heteroaryl; -CH(-aryl or -heteroaryl)-CO-O(C<sub>1</sub>-C<sub>4</sub> alkyl); -CH(-CH<sub>2</sub>-OH)-CH(OH)-phenyl-NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl)-OH, -CH<sub>2</sub>-NH-CH<sub>2</sub>-CH(-O-CH<sub>2</sub>-CH<sub>3</sub>)<sub>2</sub>, -H, and - (CH<sub>2</sub>)<sub>0-6</sub>-C(=NR<sub>235</sub>)(NR<sub>235</sub>R<sub>240</sub>); wherein each aryl is optionally substituted with 1, 2, or 3 R<sub>200</sub>; each heteroaryl is optionally substituted with 1, 2, 3, or 4 R<sub>200</sub>; each heterocycloalkyl is optionally substituted with 1, 2, 3, or 4 R<sub>210</sub>; R<sub>200</sub> at each occurrence is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; OH; -NO<sub>2</sub>; halogen; -CO<sub>2</sub>H; C≡N; - (CH<sub>2</sub>)<sub>0-4</sub>-CO-NR<sub>220</sub>R<sub>225</sub>; - (CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>1</sub>-C<sub>12</sub> alkyl); - (CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>2</sub>-C<sub>12</sub> alkenyl); - (CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>2</sub>-C<sub>12</sub> alkynyl); - (CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl); - (CH<sub>2</sub>)<sub>0-4</sub>-CO-aryl; - (CH<sub>2</sub>)<sub>0-4</sub>-CO-heteroaryl; - (CH<sub>2</sub>)<sub>0-4</sub>-CO-

heterocycloalkyl;  $-(CH_2)_{0-4}-CO-O-R_{215}$ ;  $-(CH_2)_{0-4}-SO_2-$   
 $NR_{220}R_{225}$ ;  $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$ ;  $-(CH_2)_{0-4}-SO_2-(C_1-C_{12}$   
 $\text{alkyl})$ ;  $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$ ;  $-(CH_2)_{0-4}-N(H$   
 $\text{or } R_{215})-CO-O-R_{215}$ ;  $-(CH_2)_{0-4}-N(H \text{ or } R_{215})-CO-N(R_{215})_2$ ;  
 5  $-(CH_2)_{0-4}-N-CS-N(R_{215})_2$ ;  $-(CH_2)_{0-4}-N(-H \text{ or } R_{215})-CO-R_{220}$ ;  
 $-(CH_2)_{0-4}-NR_{220}R_{225}$ ;  $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$ ;  $-(CH_2)_{0-4}-O-P(O)-(OR_{240})_2$ ;  
 $-(CH_2)_{0-4}-O-CO-N(R_{215})_2$ ;  $-(CH_2)_{0-4}-O-CS-N(R_{215})_2$ ;  
 $-(CH_2)_{0-4}-O-(R_{215})_2$ ;  $-(CH_2)_{0-4}-O-(R_{215})_2-COOH$ ;  
 $-(CH_2)_{0-4}-S-(R_{215})_2$ ;  $-(CH_2)_{0-4}-O-(C_1-C_6 \text{ alkyl optionally$   
 10  $\text{substituted with 1, 2, 3, or 5 -F})$ ;  $C_3-C_7 \text{ cycloalkyl}$ ;  
 $C_2-C_6 \text{ alkenyl optionally substituted with 1 or 2 } R_{205}$   
 $\text{groups}$ ;  $C_2-C_6 \text{ alkynyl optionally substituted with 1}$   
 $\text{or 2 } R_{205} \text{ groups}$ ;  $-(CH_2)_{0-4}-N(H \text{ or } R_{215})-SO_2-R_{220}$ ; and  
 $-(CH_2)_{0-4}-C_3-C_7 \text{ cycloalkyl}$ ;

15 wherein each aryl group at each occurrence is  
 optionally substituted with 1, 2, or 3 groups  
 that are independently  $R_{205}$ ,  $R_{210}$  or  $C_1-C_6 \text{ alkyl}$   
 substituted with 1, 2, or 3 groups that are  
 independently  $R_{205}$  or  $R_{210}$ ;

20 wherein each heterocycloalkyl group at each  
 occurrence is optionally substituted with 1, 2,  
 or 3 groups that are independently  $R_{210}$ ;

wherein each heteroaryl group at each occurrence is  
 optionally substituted with 1, 2, or 3 groups  
 25 that are independently  $R_{205}$ ,  $R_{210}$ , or  $C_1-C_6 \text{ alkyl}$   
 substituted with 1, 2, or 3 groups that are  
 independently  $R_{205}$  or  $R_{210}$ ;

$R_{205}$  at each occurrence is independently selected from the  
 group consisting of  $C_1-C_6 \text{ alkyl}$ , halogen,  $-OH$ ,  $-O-$   
 30  $\text{phenyl}$ ,  $-SH$ ,  $-C\equiv N$ ,  $-CF_3$ ,  $C_1-C_6 \text{ alkoxy}$ ,  $NH_2$ ,  $NH(C_1-C_6$   
 $\text{alkyl})$ , and  $N-(C_1-C_6 \text{ alkyl})(C_1-C_6 \text{ alkyl})$ ;

$R_{210}$  at each occurrence is independently selected from the  
 group consisting of  $C_1-C_6 \text{ alkyl optionally}$

substituted with 1, 2, or 3 R<sub>205</sub> groups; C<sub>2</sub>-C<sub>6</sub> alkenyl optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; C<sub>2</sub>-C<sub>6</sub> alkynyl optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; halogen; C<sub>1</sub>-C<sub>6</sub> alkoxy; C<sub>1</sub>-C<sub>6</sub> haloalkoxy; -NR<sub>220</sub>R<sub>225</sub>; OH; C≡N; C<sub>3</sub>-C<sub>7</sub> cycloalkyl optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; -CO-(C<sub>1</sub>-C<sub>4</sub> alkyl); -SO<sub>2</sub>-NR<sub>235</sub>R<sub>240</sub>; -CO-NR<sub>235</sub>R<sub>240</sub>; -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl); and =O; wherein

R<sub>215</sub> at each occurrence is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, -(CH<sub>2</sub>)<sub>0-2</sub>-(aryl), C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, and -(CH<sub>2</sub>)<sub>0-2</sub>-(heteroaryl), -(CH<sub>2</sub>)<sub>0-2</sub>-(heterocycloalkyl); wherein the aryl group at each occurrence is optionally substituted with 1, 2, or 3 groups that are independently R<sub>205</sub> or R<sub>210</sub>; wherein the heterocycloalkyl group at each occurrence is optionally substituted with 1, 2, or 3 R<sub>210</sub>; wherein each heteroaryl group at each occurrence is optionally substituted with 1, 2, or 3 R<sub>210</sub>;

R<sub>220</sub> and R<sub>225</sub> at each occurrence are independently selected from the group consisting of -H, -C<sub>1</sub>-C<sub>6</sub> alkyl, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, amino C<sub>1</sub>-C<sub>6</sub> alkyl; halo C<sub>1</sub>-C<sub>6</sub> alkyl; -C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl, -C<sub>1</sub>-C<sub>6</sub> alkyl chain with one double bond and one triple bond, -aryl, -heteroaryl, and -heterocycloalkyl; wherein the aryl group at each occurrence is optionally substituted with 1, 2, or 3 groups that are independently R<sub>205</sub> or R<sub>210</sub>;

wherein the heterocycloalkyl group at each occurrence is optionally substituted with 1, 2, or 3  $R_{210}$ ;

wherein each heteroaryl group at each occurrence is optionally substituted with 1, 2, or 3  $R_{210}$ ;

$R_{235}$  and  $R_{240}$  at each occurrence are independently H, or  $C_1$ - $C_6$  alkyl;

$R_{245}$  and  $R_{250}$  at each occurrence are independently selected from the group consisting of H,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  hydroxyalkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  haloalkoxy,  $-(CH_2)_{0-4}$ - $C_3$ - $C_7$  cycloalkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl, and phenyl; or

$R_{245}$  and  $R_{250}$  are taken together with the carbon to which they are attached to form a carbocycle of 3, 4, 5, 6, or 7 carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO<sub>2</sub>-, and -NR<sub>220</sub>-;

$R_{255}$  and  $R_{260}$  at each occurrence are independently selected from the group consisting of H;  $C_1$ - $C_6$  alkyl optionally substituted with 1, 2, or 3  $R_{205}$  groups;  $C_2$ - $C_6$  alkenyl optionally substituted with 1, 2, or 3  $R_{205}$  groups;  $C_2$ - $C_6$  alkynyl optionally substituted with 1, 2, or 3  $R_{205}$  groups;  $-(CH_2)_{1-2}$ -S(O)<sub>0-2</sub>-( $C_1$ - $C_6$  alkyl);  $-(CH_2)_{0-4}$ - $C_3$ - $C_7$  cycloalkyl optionally substituted with 1, 2, or 3  $R_{205}$  groups;  $-(C_1$ - $C_4$  alkyl)-aryl;  $-(C_1$ - $C_4$  alkyl)-heteroaryl;  $-(C_1$ - $C_4$  alkyl)-heterocycloalkyl; -aryl; -heteroaryl; -heterocycloalkyl;  $-(CH_2)_{1-4}$ - $R_{265}$ - $(CH_2)_{0-4}$ -aryl;  $-(CH_2)_{1-4}$ - $R_{265}$ - $(CH_2)_{0-4}$ -heteroaryl; and;  $-(CH_2)_{1-4}$ - $R_{265}$ - $(CH_2)_{0-4}$ -heterocycloalkyl; wherein  $R_{265}$  at each occurrence is independently -O-, -S- or -N( $C_1$ - $C_6$  alkyl)-;

each aryl or phenyl is optionally substituted with 1, 2, or 3 groups that are independently  $R_{205}$ ,  $R_{210}$ ,

or C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 1, 2, or 3  
groups that are independently R<sub>205</sub> or R<sub>210</sub>;  
each heteroaryl is optionally substituted with 1, 2,  
3, or 4 R<sub>200</sub>,

5 each heterocycloalkyl is optionally substituted with  
1, 2, 3, or 4 R<sub>210</sub>;

R<sub>N</sub> is -C(=O)-(CRR')<sub>0-6</sub>R<sub>100</sub>, R'<sub>100</sub>, -SO<sub>2</sub>R'<sub>100</sub>, -(CRR')<sub>1-6</sub>R'<sub>100</sub>,  
-C(=O)-(CRR')-O-R'<sub>100</sub>, -C(=O)-(CRR')-S-R'<sub>100</sub> or -C(=O)-  
(CRR')-NR<sub>100</sub>-R'<sub>100</sub>;

10 R<sub>100</sub> and R'<sub>100</sub> are independently aryl, heteroaryl, -aryl-W-aryl,  
-aryl-W-heteroaryl, -aryl-W-heterocyclyl, -heteroaryl-W-  
aryl, -heteroaryl-W-heteroaryl, -heteroaryl-W-  
heterocyclyl, -heterocyclyl-W-aryl, -heterocyclyl-W-  
heteroaryl, -heterocyclyl-W-heterocyclyl, -C(=O)-CH[(CH<sub>2</sub>)<sub>0-2</sub>-  
15 -O-R<sub>7</sub>]- (CH<sub>2</sub>)<sub>0-2</sub>-aryl, -C(=O)-CH[(CH<sub>2</sub>)<sub>0-2</sub>-O-R<sub>7</sub>]- (CH<sub>2</sub>)<sub>0-2</sub>-  
heterocyclyl, or -C(=O)-CH[(CH<sub>2</sub>)<sub>0-2</sub>-O-R<sub>7</sub>]- (CH<sub>2</sub>)<sub>0-2</sub>-  
heteroaryl, where the ring portions of each are optionally  
substituted with 1, 2, or 3 groups independently selected  
from

20 -OR, -NO<sub>2</sub>, halogen, -C≡N, -SR, -SO<sub>2</sub>R<sub>145</sub>, -C(=O)R, -OCF<sub>3</sub>,  
-CF<sub>3</sub>, -O-P(=O)(OR)(OR'), -N(R)(COR'), -N(R)(SO<sub>2</sub>R<sub>145</sub>),  
-(CH<sub>2</sub>)<sub>0-4</sub>-CO-NR<sub>105</sub>R'<sub>105</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-O-(CH<sub>2</sub>)<sub>0-4</sub>-CONRR',  
-(CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>1</sub>-C<sub>12</sub> alkyl), -(CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>2</sub>-C<sub>12</sub>  
alkenyl), -(CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>2</sub>-C<sub>12</sub> alkynyl), -(CH<sub>2</sub>)<sub>0-4</sub>-CO-  
25 (C<sub>3</sub>-C<sub>7</sub> cycloalkyl), -(CH<sub>2</sub>)<sub>0-4</sub>-R<sub>110</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-R<sub>120</sub>,  
-(CH<sub>2</sub>)<sub>0-4</sub>-R<sub>130</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-CO-R<sub>110</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-CO-R<sub>120</sub>,  
-(CH<sub>2</sub>)<sub>0-4</sub>-CO-R<sub>130</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-CO-R<sub>140</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-CO-O-R<sub>150</sub>,  
-(CH<sub>2</sub>)<sub>0-4</sub>-SO<sub>2</sub>-NR<sub>105</sub>R'<sub>105</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-SO-(C<sub>1</sub>-C<sub>8</sub> alkyl),  
-(CH<sub>2</sub>)<sub>0-4</sub>-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>12</sub> alkyl), -(CH<sub>2</sub>)<sub>0-4</sub>-SO<sub>2</sub>-(C<sub>3</sub>-C<sub>7</sub>  
30 cycloalkyl), -(CH<sub>2</sub>)<sub>0-4</sub>-N(H or R<sub>150</sub>)-CO-O-R<sub>150</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-  
N(H or R<sub>150</sub>)-CO-N(R<sub>150</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-N(H or R<sub>150</sub>)-CS-  
N(R<sub>150</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-N(-H or R<sub>150</sub>)-CO-R<sub>105</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-  
NR<sub>105</sub>R'<sub>105</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-R<sub>140</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-O-CO-(C<sub>1</sub>-C<sub>6</sub> alkyl),

- (CH<sub>2</sub>)<sub>0-4</sub>-O-P(O)-(O-R<sub>110</sub>)<sub>2</sub>, - (CH<sub>2</sub>)<sub>0-4</sub>-O-CO-N(R<sub>150</sub>)<sub>2</sub>,  
 - (CH<sub>2</sub>)<sub>0-4</sub>-O-CS-N(R<sub>150</sub>)<sub>2</sub>, - (CH<sub>2</sub>)<sub>0-4</sub>-O-(R<sub>150</sub>), - (CH<sub>2</sub>)<sub>0-4</sub>-O-  
 (R<sub>155</sub>)-COOH, - (CH<sub>2</sub>)<sub>0-4</sub>-S-(R<sub>150</sub>), C<sub>3</sub>-C<sub>7</sub> cycloalkyl,  
 - (CH<sub>2</sub>)<sub>0-4</sub>-N(-H or R<sub>150</sub>)-SO<sub>2</sub>-R<sub>7</sub>, or - (CH<sub>2</sub>)<sub>0-4</sub>-C<sub>3</sub>-C<sub>7</sub>  
 5 cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, or

R<sub>100</sub> is C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 R<sub>115</sub>  
 groups, wherein

R<sub>115</sub> at each occurrence is independently halogen, -OH,  
 -CO<sub>2</sub>R, -C<sub>1</sub>-C<sub>6</sub> thioalkoxy, -CO<sub>2</sub>-phenyl, -NR<sub>105</sub>R'<sub>7</sub>, -SO<sub>2</sub>-  
 10 (C<sub>1</sub>-C<sub>8</sub> alkyl), -C(=O)R<sub>180</sub>, R<sub>180</sub>, -CONR<sub>105</sub>R'<sub>105</sub>,  
 -SO<sub>2</sub>NR<sub>105</sub>R'<sub>105</sub>, -NH-CO-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NH-C(=O)-OH, -NH-  
 C(=O)-OR, -NH-C(=O)-O-phenyl, -O-C(=O)-(C<sub>1</sub>-C<sub>6</sub> alkyl),  
 -O-C(=O)-amino, -O-C(=O)-mono- or dialkylamino, -O-  
 C(=O)-phenyl, -O-(C<sub>1</sub>-C<sub>6</sub> alkyl)-CO<sub>2</sub>H, -NH-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub>  
 15 alkyl), C<sub>1</sub>-C<sub>6</sub> alkoxy or C<sub>1</sub>-C<sub>6</sub> haloalkoxy; or

R<sub>100</sub> is -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl) or -(C<sub>1</sub>-C<sub>6</sub> alkyl)-S-(C<sub>1</sub>-C<sub>6</sub>  
 alkyl), each of which is optionally substituted with 1, 2,  
 or 3 R<sub>115</sub> groups, or

R<sub>100</sub> is -(C<sub>3</sub>-C<sub>8</sub> cycloalkyl) optionally substituted with 1, 2, or  
 20 3 R<sub>115</sub> groups;

R and R' independently are hydrogen; C<sub>1</sub>-C<sub>6</sub> alkyl optionally  
 substituted with 1, 2, or 3 groups that are independently  
 F, Cl, Br, or I; or -(C<sub>1</sub>-C<sub>6</sub>)-R<sub>110</sub>;

W is -(CH<sub>2</sub>)<sub>0-4</sub>-, -O-, -S(O)<sub>0-2</sub>-, -N(R<sub>135</sub>)-, or -C(O)-;

25 R<sub>7</sub> and R'<sub>7</sub> are independently selected from the group consisting  
 of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub>  
 alkynyl, aryl, heteroaryl, and heterocyclyl,

R<sub>105</sub> and R'<sub>105</sub> are the same or different and represent -H, -R<sub>110</sub>,  
 -R<sub>120</sub>, -C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl),  
 30 -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub>  
 alkynyl, or -C<sub>1</sub>-C<sub>6</sub> alkyl chain with one double bond and  
 one triple bond, or  
 -C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with -OH or -NH<sub>2</sub>; or,

- C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen;
- R<sub>135</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(CH<sub>2</sub>)<sub>0-2</sub>-(aryl), -(CH<sub>2</sub>)<sub>0-2</sub>-(heteroaryl), or  
5 -(CH<sub>2</sub>)<sub>0-2</sub>-(heterocyclyl),
- R<sub>140</sub> is heterocyclyl optionally substituted with 1, 2, 3, or 4 groups independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub>  
10 alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and =O;
- R<sub>145</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl or CF<sub>3</sub>;
- R<sub>150</sub> is hydrogen, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub>  
15 cycloalkyl), C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> alkyl with one double bond and one triple bond, -R<sub>110</sub>, -R<sub>120</sub>, or C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, 3, or 4 groups independently selected from -OH, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, R<sub>110</sub>, and halogen;
- 20 R<sub>155</sub> is C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl), C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> alkyl with one double bond and one triple bond, -R<sub>110</sub>, -R<sub>120</sub>, or C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, 3, or 4 groups independently selected from -OH, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>3</sub>  
25 alkoxy, and halogen;
- R<sub>180</sub> is selected from morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl, each of which is  
30 optionally substituted with 1, 2, 3, or 4 groups independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-

C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and =O; R<sub>110</sub> is aryl optionally substituted with 1 or 2 R<sub>125</sub> groups, wherein,

R<sub>125</sub> at each occurrence is independently halogen, amino,

5 mono- or dialkylamino, -OH, -C≡N, -SO<sub>2</sub>-NH<sub>2</sub>, -SO<sub>2</sub>-NH-C<sub>1</sub>-C<sub>6</sub> alkyl, -SO<sub>2</sub>-N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO-NH<sub>2</sub>, -CO-NH-C<sub>1</sub>-C<sub>6</sub> alkyl, or -CO-N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>; or C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl, each of which is optionally substituted with 1, 2, or 3  
10 groups that are independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, and mono- and dialkylamino; or C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with one, two or three of halogen;

15 R<sub>120</sub> is heteroaryl, which is optionally substituted with 1 or 2 R<sub>125</sub> groups; and

R<sub>130</sub> is heterocyclyl optionally substituted with 1 or 2 R<sub>125</sub> groups;

comprising

20 (a) reducing a ketone of formula III to generate an alcohol of formula IV; and

(b) treating the alcohol of formula IV with a base to generate an epoxide.

25 In another aspect, the invention provides a process of preparing a compound of formula (XX), further comprising contacting the epoxide with an amine of formula R<sub>C</sub>NH(R<sub>57</sub>) to yield a protected amine of formula VII-1. further comprising deprotecting the protected amine of formula (VII-1.)

30 In another aspect, the invention provides a process of preparing a compound of formula (XX), further comprising deprotecting the protected amine of formula (VII-1) to generate a an amine or its acid addition salt of formula (VIII-1.)


In another aspect, the invention provides a process of preparing a compound of formula (XX), further comprising the deprotected amine of formula VIII-1 and forming an amide using the amine and a compound of the formula  $R_NZ$ , wherein Z is  $CO_2H$ ,  $COCl$ ,  $-SO_2Cl$ , a halogen, -O-mesylate, -O-tosylate, -O-nosylate, 5 -O-brosylate, -O-trifluoromethanesulfonate, or CO-imidazolyl. In a more preferred embodiment, Z is  $CO_2H$ . In an equally preferred embodiment, Z is  $COCl$ . In yet another equally preferred embodiment, Z is CO-imidazolyl. In yet another 10 equally preferred embodiment, Z is  $-SO_2Cl$ . In yet another equally preferred embodiment, Z is -O-mesylate, -O-tosylate, -O-nosylate, -O-brosylate, or -O-trifluoromethanesulfonate.

DETAILED DESCRIPTION OF THE INVENTION

Preferred compounds of Formula II include those where  $R_{30}$  is PROT and R is phenyl substituted with up to two groups  $R_p$  and  $R_q$ , where  $R_p$  and  $R_q$  independently represent

- 5 (A)  $C_1$ - $C_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $C_1$ - $C_3$  alkyl, halogen, hydroxy,  $-R_{11}$  where  $R_{10}$  and  $R_{11}$  are independently hydrogen or  $C_1$ - $C_6$  alkyl, trifluoromethyl, and  $C_1$ - $C_3$  alkoxy,
- 10 (B) halogen, hydroxy, cyano,  $C_1$ - $C_6$  alkoxy optionally substituted with 1, 2, or 3 fluoro, or
- (C)  $-NR_{12}R_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent hydrogen or alkyl.

15 More preferred compounds of Formula II are those of Formula II-A



II-A


where PROT and  $R_1$  are defined as above.

20 Such compounds, i.e., compounds of Formula II-A, are preferred in the processes of the invention employing the esters.

Preferred  $R_1$  groups in II-A are  $C_1$ - $C_6$  alkyl groups optionally substituted with one of bromo or chloro. More preferred are  $C_1$ - $C_4$  groups optionally substituted with bromo or

25 chloro, most preferably chloro.

Preferred compounds of Formula III include those of Formula III-A



III-A


where R, PROT and R<sub>2</sub> are as defined above with respect to Formula II.

- 5 Such compounds, i.e., compounds of Formula III-A, are preferred in the processes of the invention employing compounds of Formula III.

10 More preferred compounds of Formula III-A include those where R is phenyl substituted with up to two groups R<sub>p</sub> and R<sub>q</sub>, where R<sub>p</sub> and R<sub>q</sub> independently represent

- (A) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or three substituents independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, hydroxy, -NR<sub>10</sub>R<sub>11</sub> where R<sub>10</sub> and R<sub>11</sub> are  
 15 independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, trifluoromethyl, and C<sub>1</sub>-C<sub>3</sub> alkoxy,  
 (B) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3 fluoro, or  
 (C) -NR<sub>12</sub>R<sub>13</sub> where at each occurrence R<sub>12</sub> and R<sub>13</sub> are the  
 20 same or different and represent hydrogen or alkyl.

Particularly preferred compounds of Formula III-A are those where R<sub>p</sub> and R<sub>q</sub> independently represent C<sub>1</sub>-C<sub>2</sub> alkyl, halogen, hydroxy, or C<sub>1</sub>-C<sub>2</sub> alkoxy. Still other particularly preferred compounds of Formula III-A include those where R<sub>p</sub> and  
 25 R<sub>q</sub> independently represent halogen. A particularly preferred group of compounds represented by Formula III-A are those of Formula III-B




III-B

Preferred PROT groups are *t*-butoxycarbonyl, benzyloxycarbonyl, formyl, trityl, phthalimido, trichloroacetyl, chloroacetyl, bromoacetyl, iodoacetyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-ethoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, 2-(4-xenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(*p*-toluyl)prop-2-yloxycarbonyl, cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, (trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, isobornyloxycarbonyl, 1-piperidyloxycarbonyl, 9-fluorenylmethyl carbonate, -CH-CH=CH<sub>2</sub> and (-N=)CH-phenyl.

It is preferred that the nitrogen protecting group (PROT) be *t*-butoxycarbonyl (BOC) or benzyloxycarbonyl (CBZ), it is more preferred that PROT be *t*-butoxycarbonyl. One skilled in

the art will understand the preferred methods of introducing a t-butoxycarbonyl or benzyloxycarbonyl groups and may additionally consult T. W. Green and P. G. M. Wuts in "Protective Groups in Organic Chemistry, 3rd edition" John Wiley & Sons, Inc. New York, N.Y., 1999 for guidance.


Preferred compounds of Formula IV-unprotected include those of Formula IV-A-unprotected



(IV-A-unprotected)

where R and R<sub>2</sub> are defined as for Formula IV-unprotected. Such compounds, i.e., compounds of Formula IV-A-unprotected, are preferred for use in the processes of the invention employing compounds of formula IV-unprotected.

Preferred compounds of Formula IV-A-unprotected include those of Formula IV-B-unprotected



IV-B-unprotected


where R<sub>p</sub> and R<sub>q</sub> independently represent

- (A) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or three substituents independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, hydroxy, -NR<sub>10</sub>R<sub>11</sub> where R<sub>10</sub> and R<sub>11</sub> are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, trifluoromethyl, and C<sub>1</sub>-C<sub>3</sub> alkoxy,
- (B) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3 fluoro, or

(C)  $-NR_{12}R_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent hydrogen or alkyl.

Preferred  $R_p$  and  $R_q$  groups in Formula IV-B-unprotected are independently selected halogens. More preferably,  $R_p$  and  $R_q$  are fluorine atoms. Particularly preferred compounds of IV-B-unprotected are those where  $R_p$  and  $R_q$  are fluorine atoms in the 3- and 5-positions with respect to the point of attachment of the phenyl ring to the parent methylene.

Preferred compounds of Formula V-unprotected include those of Formula V-A-unprotected



Formula V-A-unprotected

where  $R$  is defined as above for Formula V-unprotected. Such compounds, i.e., compounds of Formula V-A-unprotected, are preferred in the processes of the invention employing compounds of Formula V-unprotected.

Preferred compounds of Formula V-A-unprotected include those where  $R$  is phenyl substituted with up to two groups  $R_p$  and  $R_q$ , where  $R_p$  and  $R_q$  independently represent

(A)  $C_1-C_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $C_1-C_3$  alkyl, halogen, hydroxy,  $-NR_{10}R_{11}$  where  $R_{10}$  and  $R_{11}$  are independently hydrogen or  $C_1-C_6$  alkyl, trifluoromethyl, and  $C_1-C_3$  alkoxy,


(B) halogen, hydroxy, cyano,  $C_1-C_6$  alkoxy optionally substituted with 1, 2, or 3 fluoro, or

(C)  $-NR_{12}R_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent hydrogen or alkyl.

Preferred  $R_p$  and  $R_q$  groups in Formula V-A-unprotected are independently selected halogens. More preferably,  $R_p$  and  $R_q$  are fluorine atoms. Particularly preferred compounds of V-A-unprotected are those where  $R_p$  and  $R_q$  are fluorine atoms in the 3- and 5-positions with respect to the point of attachment of the phenyl ring to the parent methylene.

Preferred compounds of Formula XI include those of Formula

10 XI-A



XI-A

where R and  $X_1$  are as defined for Formula XI.

Such compounds, i.e., compounds of Formula XI-A, are preferred  
15 in the processes of the invention employing compounds of Formula XI.

Preferred compounds of Formula XI-A include those where R is phenyl substituted with up to two groups  $R_p$  and  $R_q$ , where  $R_p$  and  $R_q$  independently represent  
20

- (A)  $C_1$ - $C_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $C_1$ - $C_3$  alkyl, halogen, hydroxy,  $-\text{NR}_{10}\text{R}_{11}$  where  $R_{10}$  and  $R_{11}$  are independently hydrogen or  $C_1$ - $C_6$  alkyl, trifluoromethyl, and  $C_1$ - $C_3$  alkoxy,  
25
- (B) halogen, hydroxy, cyano,  $C_1$ - $C_6$  alkoxy optionally substituted with 1, 2, or 3 fluoro, or
- (C)  $-\text{NR}_{12}\text{R}_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent hydrogen or alkyl.

30


Preferred  $R_p$  and  $R_q$  groups in Formula XI-A are independently selected halogens. More preferably,  $R_p$  and  $R_q$  are fluorine atoms. Particularly preferred compounds of XI-A are those where  $R_p$  and  $R_q$  are fluorine atoms in the 3- and 5-positions with respect to the point of attachment of the phenyl ring to the parent methylene.

Preferred compounds of Formula XIV include those where R is phenyl substituted with up to two groups  $R_p$  and  $R_q$ , where  $R_p$  and  $R_q$  independently represent

- (A)  $C_1$ - $C_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $C_1$ - $C_3$  alkyl, halogen, hydroxy,  $-NR_{10}R_{11}$  where  $R_{10}$  and  $R_{11}$  are independently hydrogen or  $C_1$ - $C_6$  alkyl, trifluoromethyl, and  $C_1$ - $C_3$  alkoxy,
- (B) halogen, hydroxy, cyano,  $C_1$ - $C_6$  alkoxy optionally substituted with 1, 2, or 3 fluoro, or
- (C)  $-NR_{12}R_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent hydrogen or alkyl.

Preferred  $R_p$  and  $R_q$  groups in Formula XIV are independently selected halogens. More preferably,  $R_p$  and  $R_q$  are fluorine atoms. Particularly preferred compounds of XIV are those where  $R_p$  and  $R_q$  are fluorine atoms in the 3- and 5-positions with respect to the point of attachment of the phenyl ring to the parent methylene.

Other preferred compounds of Formula II are those of Formula XV



## XV

i.e., compounds of Formula II where  $R_{30}$  is hydrogen. In compounds of Formula XV,  $R_1$  is as defined above with respect to Formula II.

5

Preferred alkylating agents for the esterification of I to II include

(a)  $X_4-C_1-C_4$  alkyl optionally substituted with one of iodo, bromo, or chloro, preferably chloro;

10

(a) dimethylsulfate;

(b)  $X_4-CH_2-CH=CH_2$ ,

(c)  $X_4-CH_2$ -phenyl where the phenyl ring is optionally substituted with nitro, halogen, cyano; and

where  $X_4$  is iodo, bromo, chloro, -O-tosylate, -O-mesylate

15

or -O-triflate.

Preferred compounds of Formula I are those having (S) stereochemistry and where R is phenyl substituted with two halogen atoms, preferably fluorine atoms. Preferably the phenyl is substituted in the 3- and 5- positions, more preferably with fluorine atoms in the 3- and 5-positions.

20

The following representative compounds are listed to give the reader an understanding of the compounds of formula X that may be prepared using the invention. Unless indicated otherwise, all names herein are generated using the Advanced Chemistry Development Inc. (ACD) nomenclature program, IUPAC Name Batch Version 4.5 or Version 5.09.

25

$N^1-((1S,2R)-1-benzyl-2-hydroxy-3-\{[4-(trifluoromethyl)benzyl]amino\}propyl)-N^3,N^3-dipropylisophthalamide$

30

$N^1-\{(1S,2R)-1-benzyl-3-[(2,3-dichlorobenzyl)amino]-2-hydroxypropyl\}-N^3,N^3-dipropylisophthalamide$

$N^1$ -{(1S,2R)-1-benzyl-3-[(3,5-dichlorobenzyl)amino]-2-hydroxypropyl}- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -{(1S,2R)-1-benzyl-3-[(3,5-difluorobenzyl)amino]-2-hydroxypropyl}- $N^3,N^3$ -dipropylisophthalamide

5  $N^1$ -{(1S,2R)-1-benzyl-2-hydroxy-3-{[4-(trifluoromethoxy)benzyl]amino}propyl}- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[2-(isobutylamino)-1-methyl-2-oxoethyl]amino}propyl}- $N^3,N^3$ -  
10 dipropylisophthalamide

$N^1$ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1S)-2-(isobutylamino)-1-methyl-2-oxoethyl]amino}propyl}- $N^3,N^3$ -dipropylisophthalamide

$N^3$ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[(1S)-2-(isobutylamino)-1-methyl-2-oxoethyl]amino}propyl}- $N^5,N^5$ -  
15 dipropyl-3,5-pyridinedicarboxamide

$N^1$ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[2-(isobutylamino)-1,1-dimethyl-2-oxoethyl]amino}propyl}-5-methyl- $N^3,N^3$ -dipropylisophthalamide

20  $N^1$ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[2-(isobutylamino)-2-oxoethyl]amino}propyl}-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1S)-1-[(isobutylamino)carbonyl]propyl}amino)propyl]-5-methyl- $N^3,N^3$ -  
25 dipropylisophthalamide

$N^1$ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1R)-1-[(isobutylamino)carbonyl]propyl}amino)propyl]-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -[(1S,2R)-1-(3,5-difluorobenzyl)-3-(ethylamino)-2-hydroxypropyl]-5-methyl- $N^3,N^3$ -dipropylisophthalamide  
30

$N^1$ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-(isobutylamino)propyl]-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(isobutylamino)-2-methyl-3-oxopropyl]amino}propyl)-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[4-(dimethylamino)benzyl]amino}-2-hydroxypropyl)-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -[(1S,2R)-3-{[(1S)-1-benzyl-2-(isobutylamino)-2-oxoethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1S)-1-[(isobutylamino)carbonyl]-2-methylpropyl}amino)propyl]-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -((1S,2R)-1-(3,5-difluorobenzyl)-3-{[2-(dimethylamino)ethyl]amino}-2-hydroxypropyl)-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-pyridinylmethyl)amino]propyl}-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -[(1S,2R)-3-{[(1S)-1-[(benzyloxy)methyl]-2-(isobutylamino)-2-oxoethyl]amino}-1-(3,5-difluorobenzyl)-2-hydroxypropyl]-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(1-methyl-1-phenylethyl)amino]propyl}-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -[(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-({(1R)-1-[(isobutylamino)carbonyl]-2-methylpropyl}amino)propyl]-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -((1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-{[3-(trifluoromethoxy)benzyl]amino}propyl)-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -{(1S,2R)-1-(3,5-difluorobenzyl)-3-[(3-fluorobenzyl)amino]-2-hydroxypropyl}-5-methyl- $N^3,N^3$ -dipropylisophthalamide

$N^1$ -{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-isopropoxybenzyl)amino]propyl}-5-methyl- $N^3,N^3$ -dipropylisophthalamide

5        Schemes 1-7 generally represent the processes of the invention; while these schemes employ various preferred compounds of the invention as intermediates and starting materials, it is to be understood that the processes are also applicable to compounds not having the specific stereochemistry  
10 or substituent patterns depicted in the schemes. In summary:

      Scheme 1 generally sets forth the process for the preparation of the N-protected epoxide V from known amino acid (0). Epoxides of Formula V are useful as intermediates in the production of biologically active compounds, e.g.,  
15 pharmaceuticals for the treatment of Alzheimer's Disease.

      Scheme 2 discloses a process for the transformation of an epoxide V to the desired compounds of Formula X.

      Scheme 3 discloses an alternative process for the conversion of the protected amino acid (I) to the corresponding  
20 ketone (III).

      Scheme 4 discloses an alternative process to prepare the ester (II).


      Scheme 5 discloses a process to change the protecting group for the ester (II).

25        Scheme 6 discloses a process to prepare the unprotected alcohol (IV-unprotected) and the unprotected epoxide (V-unprotected).


      Scheme 7 discloses a process for the transformation of an epoxide V-1 to the desired compounds of Formula X-1.

30        Representative examples of methods for preparing compounds of the invention are set forth below.


## SCHEME 1



## SCHEME 2




SCHEME 3




5


SCHEME 4




## SCHEME 5



## SCHEME 6



SCHEME 7



5        The epoxides of formula V have two chiral centers; thus, compounds of Formula V can exist as any of four stereoisomers, i.e., two pairs of diastereomers. While biologically active end products result from all stereoisomers, the (S,S) configuration is particularly preferred. One of these chiral centers in the epoxide (V) is derived from the starting amino acid (0). Therefore, it is preferred to start with the amino acid (0) containing the desired enantiomeric center rather than to start with a mixture and have to perform a resolution to obtain the desired (S)-enantiomer of the amino acid (0).

15

      SCHEME 1 depicts the conversion of amino acid (0) to N-protected amino epoxide (V). Protection of the free amino group of the preferably (S)-amino acid (0) with a nitrogen protecting group (PROT) yields the protected amino acid (I) having the same stereochemistry. Nitrogen protecting groups are well known to those skilled in the art, see for example,

20

"Nitrogen Protecting Groups in Organic Synthesis", John Wiley and sons, New York, N.Y., 1981, Chapter 7; "Nitrogen Protecting Groups in Organic Chemistry", Plenum Press, New York, N.Y., 1973, Chapter 2. See also, T. W. Green and P. G. M. Wuts in  
5 "Protective Groups in Organic Chemistry, 3rd edition" John Wiley & Sons, Inc. New York, N.Y., 1999. When the nitrogen protecting group is no longer needed, it may be removed. Suitable methods are known to those skilled in the art. The reader's attention is again directed to the references  
10 mentioned above.

The protected amino acid (I) (preferably of (S) stereochemistry) is then converted to the corresponding protected ester (II) (retaining the preferred (S) stereochemistry). This conversion can be accomplished in a  
15 variety of ways.

When  $R_1$  is (a)  $C_1$ - $C_4$  alkyl optionally substituted with one -Cl, (b)  $-CH_2-CH=CH_2$ , or (c) phenyl optionally substituted with one nitro, halogen, or cyano  
conversion of I to II comprises:

20 (1) esterifying a protected amino acid of the formula I with an alkylating agent in the presence of a base.

Suitable alkylating agents include

(a) those represented by the formula  $X_4-C_1-C_4$  alkyl optionally substituted with one -Cl where  $X_4$  is iodo,  
25 bromo, chloro, -O-tosylate, -O-mesylate or -O-triflate;

(b) dimethylsulfate

(c)  $X_4-CH_2-CH=CH_2$ , where  $X_4$  is as defined above

(d) a benzyl substituted on the methyl group with  $X_4$  where  
30  $X_4$  is defined as above and where the phenyl ring is optionally substituted with nitro, halogen, or cyano.

While a variety of bases are suitable for this esterification, the base is preferably hydroxide, carbonate,

bicarbonate, LDA, *n*-(C<sub>1</sub>-C<sub>8</sub> alkyl)lithium, LiHMDS, NaHMDS or KHMDS. More preferably, the base is hydroxide, carbonate or bicarbonate. An even more preferred base is carbonate.

Preferred alkylating agents are dimethylsulfate, methyl iodide, and methyl triflate. More preferably the alkylating agent is dimethylsulfate. When the base is LDA, *n*-(C<sub>1</sub>-C<sub>8</sub> alkyl)lithium, LiHMDS, NaHMDS or KHMDS, a solution of the ester is preferably cooled to from about -78°C, and more preferably about -20°C, to about 25°C prior to the addition of the base. After addition of the alkylating agent, the mixture is preferably heated to about 20°C to about 50°C. Heating is particularly useful when the alkylating agent is dimethylsulfate.

Alternatively, when R<sub>1</sub> is an optionally substituted benzyl group, the esterification can be accomplished by

(a) contacting a protected amino acid of formula (I) with an activating agent, i.e., activating the amino acid or forming an activated amino acid; and

(b) adding to the mixture of (a) a phenol optionally substituted on the phenyl ring with nitro, halogen, or cyano.

The use of activating agents, such as for example, alkyl chloroformates such as isobutyl chloroformate, CDI, and DCC, in esterification of acids with alcohols is well known to those skilled in the art. Preferred activating agents herein are CDI and DCC. Preferred esters in this process are those where R<sub>1</sub> is methyl or ethyl, more preferably methyl. A particularly preferred ester is (2S)-2-[(*tert*-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid methyl ester.

SCHEME 4 shows an alternate route to ester (II). See also EXAMPLES 9 and 10. In the process of SCHEME 4, preferred 3,5-difluorobenzaldehyde (XII), which is commercially available from, for example, Aldrich, Milwaukee, Wisconsin, USA, is reacted with the phosphorous compound (XIII), where X<sub>3</sub> is a

suitable leaving group, to produce olefin (XIV). Suitable leaving groups are known to those skilled in the art. A particularly preferred olefin (XIV) is methyl (2Z)-2-[[[(benzyloxy)carbonyl]-3-(3,5-difluorophenyl)-2-propenonate.

5       The phosphorous compounds (XIII) are known to those skilled in the art.  $X_3$  is preferably a  $C_1$ - $C_3$  alkyl group, more preferably methyl. The aldehyde (XII) and the phosphorous compound (XIII) are typically combined in a polar aprotic organic solvent, such as THF, MTBE, dioxane, ether or DME, and  
10   the resulting mixture, preferably a solution, is then cooled to about 0°. A base such as DBU or TMG is added and the contents of the mixture warmed to about 20-25°C and stirred until the reaction is complete, i.e., preferably to greater than about 90%, more preferably about 95%, and most preferably about 99%,  
15   conversion. Once the reaction is complete, the E- and Z-olefin isomers (XIV) are preferably separated since the Z isomer has the olefin stereochemistry preferred, and in some situations necessary, to yield the desired product. The separation is accomplished by methods known to those skilled in the art, such  
20   as, for example, by silica gel chromatography.

Next the olefin (XIV) is hydrogenated with a suitable hydrogenation catalyst to obtain the desired ester (II). The reaction may be conducted at pressures of from about 1 to about 100 psi. A variety of suitable catalysts will be recognized by  
25   those having ordinary skill in the art. An example of a class of suitable catalysts is represented by the formula  $[Rh(\text{diene})L]^+X^-$  where

Rh is rhodium;

diene is cyclooctadiene and norbornadiene;

30   L is a ligand selected from the group consisting of  
DIPMAP, MeDuPhos, EtDuPhos, Binaphane, f-Binaphane, Me-KetalPhos, Me-f-KetalPhos, Et-f-KetalPhos, BINAP,

DIOP, BPPFA, BPPM, CHIRAPHOS, PROPHOS, NORPHOS, CYCLOPHOS, BDPP, DEGPPOS, PNNP and

$X^-$  is  $ClO_4^-$ ,  $BF_4^-$ ,  $CF_3-SO_3^-$ ,  $Cl^-$ ,  $Br^-$ ,  $PF_6^-$  and  $SbF_6^-$ .

5 This class is preferred for use in this process aspect of the invention, particularly when L is DIPMAP or EtDuPhos.

Those skilled in the art will recognize suitable specific procedures for this reduction, i.e., hydrogenation. Generally, olefin XIV is first dissolved in the solvent, either in the reaction vessel or the solution is later transferred to the  
10 vessel. Hydrogen and the desired catalyst are then introduced into the vessel. The hydrogen is typically added under pressure, e.g., from about 25-75 psi of hydrogen. The catalyst can be added neat or as a solution of the catalyst in, for example, methanol.

15 Some hydrogenation reactions will give racemic ester (II). Since the preferred stereochemistry of the ester (II) is (S)-, it is preferable to use the Z-olefin (XIV) with an appropriate hydrogenation catalyst. Suitable solvents for the hydrogenation include polar solvents such as THF and various  
20 alcohols, preferably  $C_1-C_5$  alcohols, and most preferably methanol, ethanol, isopropanol. Another preferred solvent is THF. The solvent is preferably degassed. Further, it is preferable to purge the reaction vessel after dissolving the olefin (XIV) in the solvent and before introducing the  
25 catalyst.

The hydrogenation is preferably a chiral hydrogenation and is performed in a temperature range of from about  $0^\circ$  to about reflux; it is preferred that the reaction be performed in the temperature range from about  $0^\circ$  to about room temperature (20-  
30  $25^\circ$ ). The chiral hydrogenation is performed under a pressure of from about one atmosphere to about 100 psig. It is preferred that the chiral hydrogenation be performed under a pressure of from about 1 atmosphere to about 70 psig; it is

more preferred that the chiral hydrogenation be performed under a pressure of from about 10 psig to about 40 psig. The ester (II) is obtained in greater than 90% enantiomeric purity, preferably in greater than 95% enantiomeric purity.

5 Hydrogenation can be performed in a variety of fashions, such as, for example, in batch mode or in a continuous mode.

SCHEME 5 and EXAMPLES 11 and 12 disclose another alternate process to prepare ester II. The process of SCHEME 5 permits

10 the changing of one nitrogen protecting group for another and in addition provides the free amine XV. For example, if one has a "BOC"-protected ester (II) and desires a "CBZ"-protected ester (II), the "BOC"-protected ester (II) is typically reacted with an acid such as hydrochloric acid in a suitable solvent

15 such as methanol at temperatures of from about -20° to reflux to give the free amine (XV). Preferably the amine XV is, methyl (2S)-2-amino-3-(3,5-difluorophenyl)propionate. The free amine (XV) is then protected with a different nitrogen protecting group, such as "CBZ" to produce the corresponding

20 and desired "CBZ"-protected ester (II).

The protected ester (II), preferably of (S)-stereochemistry, is then converted to the corresponding preferably (S)-protected ketone (III) by any one of a number of processes.

25  $R_2$  is preferably -Cl or -Br, more preferably  $R_2$  is -Cl. One of the processes for the transformation of the (S)-protected ester (II) to the corresponding (S)-protected ketone (III) is exemplified in EXAMPLE 16.

Generally, the protected ester (II) of preferably (S)-

30 stereochemistry is combined with the dihalogenatedmethane reagent and to this mixture is then added a suitable base. It is preferable to add the base to the mixture of ester and dihalogenatedmethane rather than the other way around. Next,

to the resulting base/ester/dihalogenatedmethane mixture is added a second portion of base. It is preferred to add the second portion of base to the existing mixture. Finally, the base/ester/dihalogenatedmethane is treated with acid. It is preferred that  $X^2$  be  $-I$ . It is preferred that about 1 to about 1.5 equivalents of  $R_2CH_2X^2$  be used.

The strong base should have a  $pK_b$  of greater than about 30. It is preferred that the strong base be selected from the group consisting of LDA,  $(C_1-C_8 \text{ alkyl})\text{lithium}$ , LiHMDS, NaHMDS and KHMDS; it is more preferred that the strong base be LDA. It is preferred that strong base be present in an amount of from about 2 to about 2.5 equivalents.

Examples of the second base include compounds selected from the group consisting of  $(C_1-C_4)\text{alkyl lithium}$ , phenyl lithium,  $(C_1-C_4)\text{alkyl-Grignard}$  and phenyl-Grignard. It is preferred that the second base be selected from the group consisting of phenyl lithium, *n*-butyl lithium, methyl magnesium bromide, methyl magnesium chloride, phenyl magnesium bromide or phenyl magnesium chloride; it is more preferred that the second base is *n*-butyl lithium. It is preferred that the second base be present in an amount of from about 1 to about 1.5 equivalents.

Suitable acids are those, which have a  $pK_a$  of less than about 10. It is preferred the acid be selected from the group consisting of acetic, sulfuric, hydrochloric, citric, phosphoric, benzoic acids and mixtures thereof; it is more preferred that the acid be hydrochloric or acetic acid.

A variety of solvents are operable for the process; the preferred solvent for the process is THF. The reaction can be performed in the temperature range from about  $-80^\circ$  to about  $-50^\circ$ ; it is preferred to perform the reaction in the temperature range of from about  $-75^\circ$  to about  $-65^\circ$ . It is preferred that

the ketone (III) is tert-butyl (1S)-3-chloro-1-(3,5-difluorobenzyl)-2-oxopropylcarbamate.

The process of transforming the (S)-protected ester (II) to the corresponding (S)-protected ketone (III) can also be performed without the addition of a second base, see EXAMPLE 2. This process requires the presence of excess  $\text{CH}_2(\text{R}_2)\text{X}^2$  and three or more equivalents of strong base, which has a  $\text{pK}_b$  of greater than about 30 followed by adding acid.

In addition, the (S)-protected ester (II) and also be transformed to the corresponding ketone (III) in a process which comprises:

(1) contacting  $\text{R}_2\text{-CH}_2\text{-COOH}$  with a strong base which has a  $\text{pK}_b$  of greater than about 30;

(2) contacting the mixture of step (1) with an ester of formula (II); and

(3) contacting the mixture of step (2) with an acid.

In this process it is preferred that the strong base is selected from the group consisting of LDA,  $(\text{C}_1\text{-C}_8\text{ alkyl})\text{lithium}$ , LiHMDS, NaHMDS and KHMDS; it is more preferred that the base is LDA. It is preferred that from about 2 to about 2.5 equivalents of the strong base be used. The same acids as discussed above are operable here also.

SCHEME 3 and EXAMPLE 15 sets forth an alternative way of preparing the ketone (III) from the amino acid (I). This process first transforms the amino acid (I) to the intermediate (XI) and then transforms the intermediate (XI) to the desired ketone (III). The transformation of the amino acid (I) to the intermediate (XI) comprises:

(1) contacting a protected amino acid of formula (I) with a reagent selected from the group consisting of thionyl chloride,  $\text{SO}_2\text{Cl}_2$ , phosphorous trichloride, oxalyl chloride, phosphorous tribromide, triphenylphosphorous dibromide, oxalyl bromide, 1,2-phenylenetrichlorophosphate and 2,4,6-trichloro-

1,3,5-triazine. It is preferred that the reagent is thionyl chloride or oxalyl chloride. The intermediate (XI) is not isolated. It is preferred that the intermediate (XI) is *t*-butyl-(1*S*)-2-chloro-1-[3,5-difluorobenzyl]-2-oxoethylcarbamate. Intermediate (IX) is then transformed to the desired ketone (III) in a process which comprises:

(1) contacting a carbonyl compound of formula (XI) where  $X_1$  is -Cl, -Br and imidazolyl with  $LiCH_2Cl$  or  $LiCH_2Br$ . This compound is then reacted with the anion derived from the  $CH_2R_2X^2$  reagent. Various solvents are operable as is known to those skilled in the art; the preferred solvent is THF. The reaction should be performed in the cold, in a temperature range of from about  $-78^\circ$  to about  $-50^\circ$ .

The (S)-protected ketone (III) is then reduced to the corresponding (S)-alcohol (IV) or (IV-a) by means known to those skilled in the art for reduction of a ketone to the corresponding secondary alcohol, see EXAMPLE 3. In addition, European Patent Application EP 0 963 972 A2 and International Publication WO02/02512 of PCT/US01/21012 disclose alternate reagents which are operable and work well in the reduction. The reductions are carried out for a period of time between about 1 hour and about 3 days at temperatures ranging from about  $-78^\circ$  to elevated temperature up to the reflux point of the solvent employed. It is preferred to conduct the reduction between about  $-78^\circ$  and about  $0^\circ$ . If borane is used, it may be employed as a complex, for example, borane-methyl sulfide complex, borane-piperidine complex, or borane-tetrahydrofuran complex. The preferred combination of reducing agents and reaction conditions needed are known to those skilled in the art, see for example, Larock, R.C. in Comprehensive Organic Transformations, VCH Publishers, 1989.

The reduction of the (S)-protected compound (III) to the corresponding alcohol (IV) produces a second chiral center and

produces a mixture of diastereomers at the second center, (S, R/S)-alcohol (IV). This diastereomeric mixture is then separated by means known to those skilled in the art such as selective low-temperature recrystallization or chromatographic separation, most preferably by recrystallization, column chromatography or by employing commercially available chiral columns.

In another embodiment, the diastereomeric mixture produced by the non-selective reduction of the (S)-protected compound (III) is not separated but is directly converted into the epoxide. The epoxide diastereomers may then be separated into by means well known in the art. Or, the epoxide diastereomers may be reacted with the amine,  $R_cNH(R_{57})$  to form compounds analogous to structure (VII-1, where  $R_c$  is 3-methoxybenzyl and  $R_{57}$  is H.) The diastereomers may be separated at this point, or further transformations may be carried out before the diastereomers are separated. For example, the separation of the diastereomers can be carried out after deprotecting the alcohol (VII) to form the free amine (VIII), or the separation may be carried out after the amine (VIII) is converted into structure (X.)

Alternatively, the (S)-protected compound (III) may be reduced to selectively form the S or the R alcohol as illustrated in scheme I where the S alcohol is selectively formed. The selective reduction will decrease the need for the separation of the diastereomers as discussed above and will increase the amount of the desired isomer that is formed. Ideally, a single diastereomer is formed during the reduction of the ketone to the alcohol and a separation is not necessary.

The alcohol (IV) is transformed to the corresponding epoxide (V) by means known to those skilled in the art, see Scheme 6 (above) and EXAMPLE 4. The stereochemistry of the (S)-(IV) center is maintained in forming the epoxide (V). A preferred means is by reaction with base, for example, but not

limited to, hydroxide ion generated from sodium hydroxide, potassium hydroxide, lithium hydroxide and the like. Reaction conditions include the use of C<sub>1</sub>-C<sub>6</sub> alcohol solvents; ethanol is preferred. Reactions are conducted at temperatures ranging  
5 from about -45° up to the reflux temperature of the alcohol employed; preferred temperature ranges are between about -20° and about 40°.

The protected epoxides of amino acids (V) are known to those skilled in the art as intermediates in the preparation of  
10 pharmaceutical agents useful as renin and HIV inhibitors, see for example US Patents 5,482,947, 5,508,294, 5,510,349, 5,510,388, 5,521,219, 5,583,238, 5,610,190, 5,639,769, 5,760,064 and 5,965,588. In addition, the protected epoxides (V) are intermediates useful in producing pharmaceuticals  
15 agents to treat Alzheimer's disease. The epoxides (V) are transformed to useful compounds by the process of SCHEMES 2 and 3 and EXAMPLES 5 thru 8. The preferred compound is the compound of EXAMPLE 8.

The unprotected epoxide (V-unprotected) is useful in the  
20 same way. It can readily be protected to form the epoxide (V) or it can be reacted unprotected. In some instance the free amino group may interfere in the subsequent reactions but in others it will work quite well. In some instance it will be possible to put the N-terminal end on first and then open the  
25 epoxide to produce the desired compounds (X).

The compounds (X) are amines and as such form salts when reacted with acids. Pharmaceutically acceptable salts are preferred over the corresponding compounds (X) since they often produce compounds, which are more water soluble, stable and/or  
30 more crystalline. Pharmaceutically acceptable salts are any salt which retains the activity of the parent compound and does not impart any deleterious or undesirable effect on the subject to whom it is administered and in the context in which it is

administered. Pharmaceutically acceptable salts include salts of both inorganic and organic acids. The preferred pharmaceutically acceptable salts include salts of the following acids hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, phosphoric, citric, methanesulfonic,  $\text{CH}_3-(\text{CH}_2)_{n_1}-\text{COOH}$  where  $n_1$  is 0 thru 4,  $\text{HOOC}-(\text{CH}_2)_{n_1}-\text{COOH}$  where  $n_1$  is as defined above,  $\text{HOOC}-\text{CH}=\text{CH}-\text{COOH}$ ,  $\phi-\text{COOH}$ . For other acceptable salts, see *Int. J. Pharm.*, 33, 201-217 (1986).

The compounds (X) and pharmaceutically acceptable salts thereof are useful for treating humans who have Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating patients with mild cognitive impairment (MCI) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, diffuse Lewy body type of Alzheimer's disease. The compounds are preferably used in the treatment, prevention and/or alleviation of Alzheimer's disease.

#### DEFINITIONS AND CONVENTIONS

The definitions and explanations below are for the terms as used throughout this entire document including both the specification and the claims.

By "alkyl" and "C<sub>1</sub>-C<sub>6</sub> alkyl" in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, such as, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, 5 hexyl, 2-hexyl, 3-hexyl, and 3-methylpentyl. It is understood that in cases where an alkyl chain of a substituent (e.g. of an alkyl, alkoxy or alkenyl group) is shorter or longer than 6 carbons, it will be so indicated in the second "C" as, for example, "C<sub>1</sub>-C<sub>10</sub>" indicates a maximum of 10 carbons.

10 By "alkoxy" and "C<sub>1</sub>-C<sub>6</sub> alkoxy" in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, attached through at least one divalent oxygen atom, such as, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, pentoxy, isopentoxy, 15 neopentoxy, hexoxy, and 3-methylpentoxy.

By the term "halogen" in the present invention is meant fluorine, bromine, chlorine, and iodine.

"Alkenyl" and "C<sub>2</sub>-C<sub>6</sub> alkenyl" means straight and branched hydrocarbon radicals having from 2 to 6 carbon atoms and from 20 one to three double bonds and includes, for example, ethenyl, propenyl, 1-but-3-enyl, 1-pent-3-enyl, 1-hex-5-enyl and the like.

"Alkynyl" and "C<sub>2</sub>-C<sub>6</sub> alkynyl" means straight and branched hydrocarbon radicals having from 2 to 6 carbon atoms and one or 25 two triple bonds and includes ethynyl, propynyl, butynyl, pentyn-2-yl and the like.

As used herein, the term "cycloalkyl" refers to saturated carbocyclic radicals having three to twelve carbon atoms. The cycloalkyl can be monocyclic, or a polycyclic fused system. 30 Examples of such radicals include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Preferred cycloalkyl groups are cyclopentyl, cyclohexyl, and cycloheptyl. The cycloalkyl groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions with various groups.

For example, such cycloalkyl groups may be optionally substituted with, for example, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl.

By "aryl" is meant an aromatic carbocyclic group having a single ring (e.g., phenyl), multiple rings (e.g., biphenyl), or multiple condensed rings in which at least one is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl), which is optionally mono-, di-, or trisubstituted. Preferred aryl groups of the present invention are phenyl, 1-naphthyl, 2-naphthyl, indanyl, indenyl, dihydronaphthyl, tetralinyl or 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl. The aryl groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such aryl groups may be optionally substituted with, for example, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl.

By "heteroaryl" is meant one or more aromatic ring systems of 5-, 6-, or 7-membered rings which includes fused ring systems of 9-11 atoms containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur. Preferred heteroaryl groups of the present invention include pyridinyl, pyrimidinyl, quinolinyl, benzothienyl, indolyl, indolinyl, pyridazinyl, pyrazinyl, isoindolyl, isoquinolyl, quinazolinyl, quinoxalinyl, phthalazinyl, imidazolyl, isoxazolyl, pyrazolyl, oxazolyl, thiazolyl, indoliziny, indazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, furanyl, thienyl, pyrrolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, oxazolopyridinyl, imidazopyridinyl,

isothiazolyl, naphthyridinyl, cinnolinyl, carbazolyl, beta-carbolinyl, isochromanlyl, chromanlyl, tetrahydroisoquinolinyl, isoindolinyl, isobenzotetrahydrofuranlyl, isobenzotetrahydrothienyl, isobenzothienyl, benzoxazolyl, pyridopyridinyl, benzotetrahydrofuranlyl, benzotetrahydrothienyl, purinyl, benzodioxolyl, triazinyl, phenoxazinyl, phenothiazinyl, pteridinyl, benzothiazolyl, imidazopyridinyl, imidazothiazolyl, dihydrobenzisoxazinyl, benzisoxazinyl, benzoxazinyl, dihydrobenzisothiazinyl, benzopyranlyl, benzothiopyranlyl, coumarinyl, isocoumarinyl, chromonyl, chromanonyl, pyridinyl-N-oxide, tetrahydroquinolinyl, dihydroquinolinyl, dihydroquinolinonyl, dihydroisoquinolinonyl, dihydrocoumarinyl, dihydroisocoumarinyl, isoindolinonyl, benzodioxanlyl, benzoxazolinonyl, pyrrolyl N-oxide,, pyrimidinyl N-oxide, pyridazinyl N-oxide, pyrazinyl N-oxide, quinolinyl N-oxide, indolyl N-oxide, indolinyl N-oxide, isoquinolyl N-oxide, quinazolinyl N-oxide, quinoxalinyl N-oxide, phthalazinyl N-oxide, imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-oxide, thiazolyl N-oxide, indolizinyl N-oxide, indazolyl N-oxide, benzothiazolyl N-oxide, benzimidazolyl N-oxide, pyrrolyl N-oxide, oxadiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl N-oxide, tetrazolyl N-oxide, benzothiopyranlyl S-oxide, benzothiopyranlyl S,S-dioxide. The heteroaryl groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions with various groups. For example, such heteroaryl groups may be optionally substituted with, for example, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl.

By "heterocycle", "heterocycloalkyl" or "heterocyclyl" is meant one or more carbocyclic ring systems of 4-, 5-, 6-, or 7-membered rings which includes fused ring systems of 9-11 atoms

containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur. Preferred heterocycles of the present invention include morpholinyl, thiomorpholinyl, thiomorpholinyl S-oxide, thiomorpholinyl S,S-dioxide, 5 piperazinyl, homopiperazinyl, pyrrolidinyl, pyrrolinyl, tetrahydropyranyl, piperidinyl, tetrahydrofuranyl, tetrahydrothienyl, homopiperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S,S-dioxide, oxazolidinonyl, dihydropyrazolyl, dihydropyrrolyl, 10 dihydropyrazinyl, dihydropyridinyl, dihydropyrimidinyl, dihydrofuryl, dihydropyranyl, tetrahydrothienyl S-oxide, tetrahydrothienyl S,S-dioxide and homothiomorpholinyl S-oxide. The heterocycle groups herein are unsubstituted or, as specified, substituted in one or more substitutable positions 15 with various groups. For example, such heterocycle groups may be optionally substituted with, for example, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub>alkenyl, C<sub>2</sub>-C<sub>6</sub>alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>- 20 C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or =O.

All temperatures are in degrees Celsius.

TLC refers to thin-layer chromatography.

25 HPLC refers to high pressure liquid chromatography.

THF refers to tetrahydrofuran.

psig refers to pounds of pressure per square inch.

CDI refers to 1,1'-carbonyldiimidazole.

DCC refers to dicyclohexylcarbodiimide.

30 TMG refers to 1,1,3,3-tetramethylguanidine.

DMF refers to dimethylformamide.

DBU refers to 1,8-diazabicyclo[5.4.0]undec-7-ene.

DBN refers to 1,5-diazabicyclo[4.3.0]non-5-ene.

LDA refers to lithium diisopropylamide.

LiHMDS, refers to lithium bis(trimethylsilyl)amide.

NaHMDS refers to sodium bis(trimethylsilyl)amide.

KHMDS refers to potassium bis(trimethylsilyl)amide.

5 BOC refers to t-butoxycarbonyl; 1,1-dimethylethoxy carbonyl;  $(\text{CH}_3)_3\text{C}-\text{O}-\text{CO}-$ .

Hunig's base refers to DIPEA, diisopropylethylamine,  $[(\text{CH}_3)_2\text{CH}]_2-\text{N}-\text{CH}_2\text{CH}_3$ .

DMAP refers to dimethylaminopyridine,  $(\text{CH}_3)_2\text{N}$ -pyridin-1-yl.

10 Saline refers to an aqueous saturated sodium chloride solution.

Chromatography (column and flash chromatography) refers to purification/separation of compounds expressed as (support; eluent). It is understood that the appropriate fractions are  
15 pooled and concentrated to give the desired compound(s).

CMR refers to C-13 magnetic resonance spectroscopy, chemical shifts are reported in ppm ( $\delta$ ) downfield from TMS.

NMR refers to nuclear (proton) magnetic resonance spectroscopy, chemical shifts are reported in ppm ( $\delta$ ) downfield  
20 from TMS.

TMS refers to trimethylsilyl.

$-\phi$  refers to phenyl ( $\text{C}_6\text{H}_5$ ).

MS refers to mass spectrometry expressed as m/e, m/z or mass/charge unit.  $[\text{M} + \text{H}]^+$  refers to the positive ion of a  
25 parent plus a hydrogen atom. EI refers to electron impact. CI refers to chemical ionization. FAB refers to fast atom bombardment.

ESMS refers to electrospray mass spectrometry.

HRMS refers to high resolution mass spectrometry.

30 Pharmaceutically acceptable refers to those properties and/or substances which are acceptable to the patient from a pharmacological/toxicological point of view and to the manufacturing pharmaceutical chemist from a physical/chemical

point of view regarding composition, formulation, stability, patient acceptance and bioavailability.

Pharmaceutically acceptable anion salts include salts of the following acids methanesulfonic, hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, benzoic, citric, tartaric, fumaric, maleic,  $\text{CH}_3-(\text{CH}_2)_n-\text{COOH}$  where  $n$  is 0 thru 4,  $\text{HOOC}-(\text{CH}_2)_N-\text{COOH}$  where  $n$  is as defined above.

-O-mesylate refers to -O-methanesulfonic acid.

-O-tosylate refers to -O-toluenesulfonic acid.

-O-triflate refers to -O-trifluoroacetic acid.

When solvent pairs are used, the ratios of solvents used are volume/volume (v/v).

When the solubility of a solid in a solvent is used the ratio of the solid to the solvent is weight/volume (wt/v).

DIPMAP refers to (R,R)-1,2-bis[(o-methoxyphenyl)-phenylphosphine]ethane.

MeDuPhos refers to 1,2-bis ((2S,5S)-2,5-dimethylphospholano)benzene.

EtDuPhos refers to 1,2-bis ((2S,5S)-2,5-dimethylphospholano)benzene.

Binaphane refers to (S,S)-1,2-Bis{(S)-4,5-dihydro-3H-dinaphtho[2,1-c:1',2'-e]phosphepino}benzene.

f-Binaphane refers to (R,R)-1,1'-Bis{(R)-4,5-dihydro-3H-dinaphtho[2,1-c:1',2'-e]phosphepino}ferrocene; "f" refers to ferrocenyl.

Me-KetalPhos refers to 1,2-Bis-[(2S,3S,4S,5S)-3,4-O-isopropylidene-3,4-dihydroxy-2,5-dimethyl]benzene.

Me-f-KetalPhos refers to 1,1'-Bis-[(2S,3S,4S,5S)-2,5-dimethyl-3,4-O-isopropylidene-3,4-dihydroxyphospholanyl]ferrocene.

Et-f-KetalPhos refers to 1,1'-Bis-[(2S,3S,4S,5S)-2,5-diethyl-3,4-O-isopropylidene-3,4-dihydroxyphospholanyl]ferrocene

BINAP refers to *R*-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl.

DIOP refers to (*R,R*)-2,3-*O*-isopropylidene-2,3-dihydroxy-1,4-bis(diphenylphosphino)-butane.

5 BPPFA refers to *R*-1-[(*S*)-1'2-bisdiphenylphosphino]ferrocenyl]-ethyl dimethylamine.

BPPM refers to (2*S*,4*S*)-*N*-butoxycarbonyl-4-diphenylphosphino-2-diphenylphosphinomethylpyrrolidine.

CHIRAPHOS refers to (*S,S*)-2,3-10 bis(diphenylphosphino)butane.

PROPHOS refers to (*S*)-1,2-bis(diphenylphosphino)propane.

NORPHOS refers to (*R,R*)-5,6-bis(diphenylphosphino)-2-norbornene.

15 CYCLOPHOS refers to *R*-1-cyclohexyl-1,2-bis(diphenylphosphino)ethane.

BDPP refers to (2*S*,4*S*)-bis(diphenylphosphino)pentane.

DEGPPOS refers to 1-substituted (*S,S*)-3,4-bis(diphenylphosphino)-pyrrolidine.

20 PNNP refers to *N,N'*-bis(diphenylphosphino)-*N,N'*-bis[(*R*)-1-phenyl]ethylenediamine.

Thionyl chloride refers to SOCl<sub>2</sub>.

Phosphorous trichloride refers to PCl<sub>3</sub>.

Oxalyl chloride refers to (COCl)<sub>2</sub>.


Phosphorous tribromide refers to PBr<sub>3</sub>.

25 Triphenylphosphorous dibromide refers to φ<sub>3</sub>PBr<sub>2</sub>.


Oxalyl bromide refers to (COBr)<sub>2</sub>.

Ether refers to diethylether.

1,2-Phenylenetrichlorophosphate refers to



30 2,4,6-trichloro-1,3,5-triazine refers to



MTBE refers to methyl *t*-butyl ether.

DME refers to dimethoxyethane.

5        The disclosures in this application of all articles and references, including patents, are incorporated herein by reference.

10        The invention is illustrated further by the following examples, which are not to be construed as limiting the invention in scope or spirit to the specific procedures described in them.


15        The starting materials and various intermediates may be obtained from commercial sources, prepared from commercially available organic compounds, or prepared using well-known synthetic methods.

#### EXAMPLES

20        The following detailed examples describe how to prepare the various compounds and/or perform the various processes of the invention and are to be construed as merely illustrative, and not limitations of the preceding disclosure. Those skilled in the art will recognize appropriate variations from the procedures both as to reactants and as to reaction conditions and techniques.


25

EXAMPLE 1        (2S)-2-[(*tert*-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid methyl ester (II)



To a 1-L 3-neck round bottom flask equipped with a magnetic stirrer, nitrogen inlet and thermocouple is added  
5 (2S)-2-[(tert-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid (I, 40 g, 0.133 moles, 1 equivalent) followed by THF (240 mL). Lithium hydroxide monohydrate (5.6 g, 0.133 moles, 1 equivalent) is added in a single portion and is allowed to stir for 30 min at which time,  
10 the contents are cooled to 0°. Once cooled, dimethyl sulfate (12.6 mL, 0.133 moles, 1 equivalent) is added dropwise via syringe and then stirred for 30 min. The mixture is then heated to about 50° and monitored (by HPLC) until 90% conversion had been achieved. At that time, the mixture is  
15 cooled to below 20° (solids form). The mixture is then poured into sodium bicarbonate (200 mL), stirred for 15 min then extracted with methyl t-butyl ether (200 mL). The phases are separated and the aqueous layer is extracted with methyl t-butyl ether (2 x 200 mL). The combined organic phases are  
20 washed with water (400 mL) dried over sodium sulfate, filtered and concentrated under reduced pressure to give a solid. This material is then recrystallized from hexanes to give the title compound, mp = 81°; NMR (DMSO-d<sub>6</sub>) δ 7.51, 7.15-7.25, 4.43, 3.81, 3.00-3.26 and 1.49; CMR (DMSO-d<sub>6</sub>) δ 172.43, 163.74,  
25 161.20, 155.67, 142.58, 112.70, 120.23, 78.69, 54.71, 52.24, 39.25 and 28.37.

EXAMPLE 2                      *tert*-butyl (1*S*)-3-chloro-1-(3,5-difluorobenzyl)-2-oxopropylcarbamate (III)




5

To a 1-L 3-neck round bottom flask equipped with a magnetic stirrer, nitrogen inlet, thermocouple and additional funnel is added (2*S*)-2-[(*tert*-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid methyl ester (II, EXAMPLE 1, 10.0 g, 0.0317 moles, 1 equivalent) followed by THF (175 mL) then cooled to -78°. Once the mixture is cooled, iodochloromethane (9.25 mL, 0.127 moles, 4 equivalents) is added in one portion via syringe. The addition funnel is charged with LDA (79 mL, 0.158 moles, 5 equivalents, 2.0 M in heptane/THF) and is subsequently added dropwise to the mixture keeping the internal temperature below -70°. Once the addition is complete, the contents are stirred for 15 min at which time acetic acid (47.2 mL, 0.824 moles, 26 equivalents) is added dropwise via the addition funnel keeping the internal temperature below -65°. Once this addition is complete, the mixture is stirred for 15 min then warmed to 0° and poured into water (500 mL), saline (500 mL) and methyl *t*-butyl ether (500 mL) then transferred to a separatory funnel. The phases are separated and the aqueous phase is extracted with methyl *t*-butyl ether (2 x 250 mL). The combined organic phases are washed with saturated sodium bicarbonate (500 mL), sodium sulfite (500 mL) and water (500 mL). The organic phase is then dried over sodium sulfate, filtered and concentrated under reduced pressure to give a

25

solid. The solid is recrystallized from heptane/*i*-propyl alcohol (10/1) to give the title compound, mp = 139°; NMR (DMSO- $d_6$ )  $\delta$  7.47, 7.06-7.14, 4.78, 4.49, 3.20, 2.82 and 1.40; CMR (DMSO- $d_6$ )  $\delta$  200.87, 163.74, 161.20, 142.74, 112.80, 102.13, 79.04, 58.97, 47.72, 34.95 and 28.30.

EXAMPLE 3      *tert*-butyl (1*S*,2*S*)-3-chloro-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (IV)




10

To a 250 mL 3-neck round bottom flask equipped with magnetic stir bar, nitrogen inlet and thermocouple, is added *tert*-butyl (1*S*)-3-chloro-1-(3,5-difluorobenzyl)-2-oxopropylcarbamate (III, EXAMPLE 2, 4.4 g, 0.0132 moles, 1 equivalent) followed by THF (20 mL) and ethanol (30 mL) then cooled to -78°. Once the mixture is cooled, sodium borohydride (2.0 g, 0.0527 moles, 4 equivalents) is added as a solid portion wise over 30 min keeping the internal temperature below -70°. Once this addition is complete, the contents are stirred for 2 hr at -78° then warmed to 0° and stirred an additional 1 hr. The mixture is quenched by the addition of saturated potassium bisulfate (15 mL) and water (15 mL). This slurry is stirred for 30 min at 20-25° then concentrated under reduced pressure to half its volume. The mixture is then cooled to 0° and stirred for 30 min. After this time, the resultant solids are collected by filtration and washed with water (2 x 50 mL) then dried under reduced pressure at 50° to give crude product. A syn/anti ratio of 4-9:1 has been observed. The desired

product is recrystallized from hexanes/ethanol (25/1) to give the title compound, mp = 149°; NMR (DMSO- $d_6$ )  $\delta$  6.89-7.16, 5.61, 3.64-3.83, 3.19, 2.69 and 1.41; CMR (DMSO- $d_6$ )  $\delta$  163.67, 161.24, 155.44, 112.70, 101.55, 78.04, 72.99, 54.29, 48.24, 35.97 and 28.37.

EXAMPLE 4      *tert*-Butyl (1*S*)-2-(3,5-difluorophenyl)-1-[(2*S*)-oxirany]ethylcarbamate (V)



To a 250 mL 3-neck round bottom flask equipped with magnetic stir bar, nitrogen inlet and thermocouple, is added *tert*-butyl (1*S*,2*S*)-3-chloro-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (IV, EXAMPLE 3, 3.5 g, 0.010 moles, 1 equivalent) followed by absolute ethanol (60 mL) and cooled to 0°. To this mixture is added potassium hydroxide (0.73 g, 0.013 moles, 1.25 equivalents) dissolved in absolute ethanol (10 mL) over 1 hr and the resulting suspension is warmed to 15-20° and stirred for 1 hr. At this time, water (100 mL) is added and the reaction contents are cooled to -5° and stirred for 30 min. The solids are collected by filtration and washed with cold water (2 x 25 mL) then dried under reduced pressure at 45° to give the title compound, mp = 133°; NMR (DMSO- $d_6$ )  $\delta$  7.03, 3.61, 2.68-2.98 and 1.33; CMR (DMSO- $d_6$ )  $\delta$  163.72, 161.29, 155.55, 143.35, 112.65, 101.80, 78.17, 53.42, 52.71, 44.90, 36.98 and 28.36.

The *anti*-diastereomer mp = 101°.

EXAMPLE 5        tert-Butyl (1S, 2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propylcarbamate (VII)

5        tert-Butyl (1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxirany]ethylcarbamate (V, EXAMPLE 4, 245 mg, 0.82 mmol) is suspended in isopropyl alcohol (6 mL) and 3-methoxybenzylamine (160  $\mu$ L, 1.22 mmol) is added with stirring at 20-25°. This mixture is heated to gentle reflux (bath temp 85°) under  
10        nitrogen for 2 hr, whereupon the resulting mixture is concentrated under reduced pressure to give the title compound. The title compound is purified by flash chromatography (2-5% methanol/methylene chloride; gradient elution) to give purified title compound.

15

EXAMPLE 6        (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol (VIII)

20        tert-Butyl (1S, 2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propylcarbamate (VII, EXAMPLE 5, 258 mg, 0.59 mmol) is dissolved in methylene chloride (1 mL) at 20-25°, and trifluoroacetic acid (1 mL) is added with stirring under nitrogen. The mixture is stirred at 20-25° for 1 hr, whereupon the mixture is concentrated under reduced pressure to  
25        give the title compound. The title compound is used in the next reaction without further purification.


EXAMPLE 7        N<sup>1</sup>-{(1S,2R)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-methoxybenzyl)amino]propyl}-5-methyl-N<sup>3</sup>,N<sup>3</sup>-  
30        dipropylisophthalamide (X) (2R,3S)-3-amino-4-(3,5-difluorophenyl)-1-[(3-methoxybenzyl)amino]-2-butanol (VIII, EXAMPLE 6) is dissolved in anhydrous DMF (3 mL) and cooled to 0°. Triethylamine (500  $\mu$ L, 3.6 mmol) and 5-

methyl-*N*, *N*-dipropylisophthalamide (IX, 156 mg, 0.59 mmol) are added with stirring. The mixture is warmed to 20-25° briefly to allow for complete dissolution of the carboxylic acid, before recooling to 0°. 1-Hydroxybenzotriazole (157 mg, 1.2 mmol) is added with stirring, followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (229 mg, 1.2 mmol). The resulting mixture is stirred at 0° for 5 min, then warmed to 20-25° for 15 hr. The mixture is then quenched with aqueous citric acid (10%), and the mixture extracted three times with ethyl acetate. The combined organic extracts are washed with saturated sodium bicarbonate, saline, dried over sodium sulfate, filtered and concentrated under reduced pressure to give the title compound in crude form. This material is purified by flash chromatography (2-10% methanol/methylene chloride gradient elution) to give purified title compound, MS (ES)  $MH^+ = 582.3$ .

EXAMPLE 8  $N^1$ -{(1*S*,2*R*)-1-(3,5-difluorobenzyl)-2-hydroxy-3-[(3-iodobenzyl)amino]propyl}-5-methyl- $N^3,N^3$ -dipropylisophthalamide (X)


Following the general procedure of EXAMPLES 5, 6 and 7 and making non-critical variations but using 3-iodobenzylamine, the title compound is obtained.

EXAMPLE 9 Methyl (2*Z*)-2-[[benzyloxy]carbonyl]-3-(3,5-difluorophenyl)-2-propenonate (XIV)



3,5-Difluorobenzaldehyde (XII, 2.87 g, 0.02 moles, 1 equivalent) and THF (100 mL) are mixed and cooled to about 0°. N-(Benzyloxycarbonyl)phosphonyl-glycinetrimethylester (XIII, 8.7 g, 0.026 moles, 1.3 equivalents) is added to the 3,5-difluorobenzaldehyde (XII)/THF mixture. This is followed by 1,1,3,3-tetramethyl guanidine (4.0 mL, 0.032 moles, 1.56 equivalents) added dropwise. The reaction is stirred for 5 min at 0° then allowed to warm to 20-25°. After 2 hr, the reaction is complete (by TLC analysis) at which time water (100 mL) and ethyl acetate (100 mL) are added. The phases are separated and the aqueous phase is extracted with ethyl acetate (100 mL) and the combined organic phases are washed with saline (100 mL), dried over sodium sulfate, filtered and concentrated under reduced pressure to give a crude solid. The solid is purified by silica gel chromatography (ethyl acetate/hexanes; 15/85) to give the title compound, mp = 112°; NMR (CDCl<sub>3</sub>) δ 7.19, 7.06, 6.86, 6.15, 6.43, 4.97 and 3.69; CMR (CDCl<sub>3</sub>) δ 165.56, 164.54, 164.41, 162.07, 137.39, 136.02, 128.97, 128.80, 128.62, 128.57, 128.47, 126.25, 112.57, 112.38, 105.22, 104.97, 104.72, 68.17 and 53.33. Additional material is recovered that is a mixture of E and Z olefins.


EXAMPLE 10 methyl (2S)-2-[[ (benzyloxy)carbonyl]amino]-3-(3,5-difluorophenyl)propanoate (II)



Methyl (2Z)-2-[[ (benzyloxy)carbonyl]amino]-3-(3,5-difluorophenyl)-2-propenonate (XIV, XIV, EXAMPLE 9, 0.100 g, 0.228 mmol) and degassed methanol (10 ml) are mixed in a 100 mL Hastelloy bomb. The mixture is purged three times with hydrogen

(60 psig) and then stirred at 60 psig hydrogen for 60 min at 20-25°. Then (R,R)-DIPAP)Rh (5.2 mg, 3 mole%) is dissolved in methanol (1 mL, degassed) is added and the system purged with hydrogen (3 x 60 psig). The contents are then stirred at 20  
5 psig hydrogen at 25° overnight at which time the reaction is complete as determined by HPLC. The system is then purged and filtered to remove the catalyst and the solvent is removed under reduced pressure to give the title compound.


10 EXAMPLE 11 Methyl (2S)-2-amino-3-(3,5-difluorophenyl)propanoate (XV)



(2S)-2-[(tert-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid methyl ester (II, EXAMPLE 1, 0.60  
15 g (0.002 moles, 1 equivalent), methanol (20 mL) and hydrochloric acid (3N, 20 mL) are mixed. The mixture is then heated to 50° and stirred until complete as measured by HPLC. When the reaction is complete, the contents are cooled to 20-25° and the pH of the mixture is adjusted to 8 with saturated  
20 sodium bicarbonate and then concentrated under reduced pressure. This mixture is extracted with ethyl acetate (2 x 20 mL) and the combined organic phases are dried over sodium sulfate, filtered and concentrated, HPLC (Retention time = 2.89 min; Zorbax RX-C8 acetonitrile/0.05M potassium dihydrogen  
25 phosphate, 60/40; 1.0 mL/min,  $\lambda$ =210 nm.

This material is carried on without further purification into the next step.

EXAMPLE 12 Methyl (2S)-2-[[ (benzyloxy) carbonyl] amino]-3-(3,5-difluorophenyl)propanoate (II)



5

Methyl (2S)-2-amino-3-(3,5-difluorophenyl)propanoate (XV, EXAMPLE 11, 0.300 g, 1.40 mmol, 1 equivalent) and water (10 mL) are mixed. Sodium carbonate (0.15 g, 1.40 mmol, 1 equivalent) of is added followed by benzylchloroformate (0.2 mL, 0.24 g, 1.4 mmol, 1 equivalent) and the mixture stirred at 20-25° until complete as measured by HPLC. Once the reaction is complete, ethyl acetate (20 mL) is added and the phases separated. The aqueous phase is extracted with ethyl acetate (2 x 20 mL), and the combined organic phases are dried over sodium sulfate, filtered, and concentrated. The concentrate is crystallized from hexanes/ethyl acetate to give the title compound, mp = 54°; NMR (DMSO-*d*<sub>6</sub>) δ 7.84, 7.28, 7.06, 4.98, 4.35, 3.68, 3.12 and 2.88.

20 EXAMPLE 13 (2S)-2-[(*tert*-butoxycarbonyl) amino]-3-(3,5-difluorophenyl)propanoic acid methyl ester (II)

(2S)-2-[(*tert*-butoxycarbonyl) amino]-3-(3,5-difluorophenyl)propanoic acid (I, 5.0 g, 0.017 moles, 1.0 equivalent) and potassium carbonate (2.5 g, 0.018 moles, 1.1 equivalent) are mixed in THF (100 mL). To this heterogeneous mixture is then added dimethyl sulfate (1.6 mL, 2.1 g, 0.017 moles, 1.0 equivalent) and the contents were then stirred at 20-25° overnight. Once the reaction is complete as measured by

HPLC, ammonium hydroxide (10%, 20 mL) is added and allowed to stir for 1 hr at which time the contents are extracted with ethyl acetate (3 x 50 mL). The combined organic phases are washed with water (50 mL) and saline (50 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure to give the title compound.

EXAMPLE 14      (2S)-2-[(tert-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid methyl ester (II)

(2S)-2-[(tert-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid (I, 5.0 g, 0.017 moles, 1.0 equivalent) and potassium carbonate (2.5 g, 0.018 moles, 1.1 equivalent) and DMF (100 mL) are mixed. To this heterogeneous mixture is then added dimethyl sulfate (1.6 mL, 2.1 g, 0.017 moles, 1.0 equivalent) and the contents are then stirred at 20-25° overnight. Once the reaction is complete as measured by HPLC, ammonium hydroxide (10%, 20 mL) is added and allowed to stir for 1 hr. The contents are stirred for 30 min then cooled to 0° and filtered. The solids are washed with cold water (20 mL) and dried under reduced pressure to give the title compound.

EXAMPLE 15      tert-butyl (1S)-3-chloro-1-(3,5-difluorobenzyl)-2-oxopropylcarbamate (III)


(2S)-2-[(tert-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid (I) is dissolved in THF and stirred at 20-25°. Oxalyl chloride (1 equivalent) is added and the mixture stirred for about 15 min to give t-butyl-(1S)-2-chloro-1-[3,5-difluorobenzyl]-2-oxoethylcarbamate (XI). The mixture is cooled to <0° and LiCHCl<sub>2</sub> (greater than 2 equivalents) is added. The mixture is stirred until the

reaction is complete. The reaction is quenched with water and the product is extracted into ethyl acetate. The combined organic phases are washed with saline, dried over sodium sulfate and concentrated under reduced pressure to give the  
5 title compound.

EXAMPLE 16      *tert*-butyl (1*S*)-3-chloro-1-(3,5-difluorobenzyl)-  
2-oxopropylcarbamate (III)

10      ICH<sub>2</sub>Cl (3.54 g, 1.46 mL, 19.82 mmol, 1.25 equivalent)  
and THF (5 mL) are added to (2*S*)-2-[(*tert*-  
butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid  
methyl ester (II, EXAMPLE 1, 5 g, 15.86 mmol, 1 equivalent).  
The mixture is cooled to -78° and LDA (22.3 mL, 44.60 mmol,  
15 2.25 equivalents, 2.0M) is added dropwise maintaining an  
internal temperature below -60°. Once the addition is  
complete, the contents are stirred for 30 min at -78° at which  
time *n*-butyllithium (15.3 mL, 19.82 mmol, 1.25 equivalents;  
1.3M in hexanes) is added dropwise maintaining an internal  
20 temperature below about -60°. The reaction is stirred for 30  
min then quenched into 0° hydrochloric acid (1N). Ethyl  
acetate is added and the phases are separated and the aqueous  
phase is extracted with ethyl acetate. The combined organic  
phases are washed with saturated sodium bicarbonate, dried over  
25 sodium sulfate, filtered and concentrated under reduced  
pressure to give the title compound, NMR (DMSO-*d*<sub>6</sub>) δ 7.47,  
7.06-7.14, 4.78, 4.49, 3.20, 2.82 and 1.40; CMR (DMSO-*d*<sub>6</sub>) δ  
200.87, 163.74, 161.20, 142.74, 112.80, 102.13, 79.04, 58.97,  
47.72, 34.95 and 28.30.

30      EXAMPLE 17      (2*S*,3*S*)-3-amino-1-chloro-4-(3,5-  
difluorophenyl)butan-2-ol



tert-butyl (1S,2S)-3-

chloro-1-(3,5-difluorobenzyl)-2-hydroxypropylcarbamate (IV,

EXAMPLE 3, 1.0 gm, 2.98 mmol) and Dowex50WX2-400 resin (4.6 gm,

23.8 mmol) and methanol (25 mL) are mixed. The mixture is then

5 placed over a J-Kim shaker with heating at 50° for 2 hr. ESMS

analysis indicates no starting material left in the mixture.

The reaction contents are filtered through a sintered funnel

and the resin washed with methanol (25 mL) and

methanol/methylene chloride (1/1, 25 mL). The resulting

10 mixture is eluted with ammonia in methanol (2N, 2 x 25 mL).


The eluate is concentrated under reduced pressure to give the

title compound, ESMS = 236.1.

#### EXAMPLE 18

(1S)-2-(3,5-difluorophenyl)-1-[(2S)-oxiran-

15 2-yl]ethylamine



(1S,2S)-3-chloro-1-(3,5-difluorobenzyl)-2-

hydroxypropylamine (EXAMPLE 17, 33mg, 0.14 mmol) and absolute

ethanol (1.5 mL) are mixed. Potassium hydroxide (9.8 mg, 0.175

20 mmol) in absolute ethanol (0.5 mL) is added to this mixture and

the resulting mixture is stirred at 20-25 deg for 30 min. At


this time ESMS indicates formation of the product (MH<sup>+</sup> =

200.1). Water (2 mL) is added and mixture is concentrated under reduced pressure to half the volume and then diluted with ethyl acetate (15 mL). The organic phase is separated and the aqueous phase is extracted with ethyl acetate (2 x 10 mL). The  
5 organic phases are combined, washed with saline and dried over anhydrous magnesium sulfate. The solvent is removed under reduced pressure to give the title compound,  $MH^+ = 200.1$ .

The invention and the manner and process of making and using it, are now described in such full, clear, concise and  
10 exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the invention and that modifications may be made therein without departing from the spirit or scope of the invention as set  
15 forth in the claims. To particularly point out and distinctly claim the subject matter regarded as the invention, the following claims conclude this specification.

What is claimed is:

1. A compound of the formula:



where

- 5 R is phenyl optionally substituted with 1, 2, 3, or 4 groups independently selected from:

- (A) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or three substituents independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, hydroxy, thio, -NR<sub>10</sub>R<sub>11</sub> where R<sub>10</sub> and R<sub>11</sub> are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, cyano, trifluoromethyl, and C<sub>1</sub>-C<sub>3</sub> alkoxy,
- 10 (B) C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl,
- (C) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3 fluoro,
- 15 (D) -NR<sub>12</sub>R<sub>13</sub> where at each occurrence R<sub>12</sub> and R<sub>13</sub> are the same or different and represent:
- (a) -H,
- (b) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one of:
- 20 (i) -OH,
- (ii) -NH<sub>2</sub>,
- (iii) phenyl,
- (c) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with 1, 2, or 3 independently selected halogens,
- (d) -C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl; and
- 25 (E) C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NR<sub>10</sub>R<sub>11</sub>, -C(O)NR<sub>10</sub>R<sub>11</sub>, or -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl);

R<sub>1</sub> is selected from:


- 30 (I) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one halogen;
- (II) -CH<sub>2</sub>-CH=CH<sub>2</sub>;

(III) phenyl optionally substituted with one nitro,  
halogen, or cyano; and

(IV) benzyl optionally substituted on phenyl with nitro,  
halogen, or cyano; and

5  $R_{30}$  represents hydrogen or PROT, where PROT is a nitrogen  
protecting group.

2. An ester of the formula (II)



10 II

$R_1$  is selected from:

(I)  $C_1$ - $C_6$  alkyl optionally substituted with one halogen;

(II)  $-CH_2-CH=CH_2$ ;

(III) phenyl optionally substituted with one nitro,  
15 halogen, or cyano; and

(IV) benzyl optionally substituted on phenyl with nitro,  
halogen, or cyano; and

PROT is a nitrogen protecting group.

20 3. An ester according to claim 1 where PROT is t-  
butoxycarbonyl.

4. An ester according to claim 1 where PROT is  
benzyloxycarbonyl.

25

5. An ester according to claim 1 where  $R_1$  is  $C_1$ - $C_2$  alkyl.


6. An ester according to claim 5 where  $R_1$  is  $C_1$  alkyl.

7. An ester according to claim 1 which is selected from the group consisting of

(2S)-2-[(tert-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid methyl ester and

5 methyl (2S)-2-[[ (benzyloxy) carbonyl] amino]-3-(3,5-difluorophenyl)propanoate.

8. A compound of the formula



10 where R<sub>2</sub> is:

chloro, bromo, or

-Si(R<sub>21</sub>)<sub>3</sub> where each R<sub>21</sub> is independently

C<sub>1</sub>-C<sub>5</sub> alkyl,

-N(R<sub>23</sub>)(R<sub>24</sub>) where R<sub>23</sub> and R<sub>24</sub> are the same or

15 different and represent

C<sub>1</sub>-C<sub>5</sub> alkyl,

or where NR<sub>23</sub>R<sub>24</sub> represents piperidinyl,

piperazinyl, or morpholinyl,

phenyl optionally substituted with 1, 2, or 3 of C<sub>1</sub>-

20 C<sub>2</sub> alkyl, with the proviso that at least one of


the R<sub>21</sub> groups is optionally substituted phenyl.

9. A compound according to claim 8 where R<sub>2</sub> is -Cl.


25 10. A compound according to claim 8 where R<sub>2</sub> is -Br.

11. A compound according to claim 8 which is (2S,3S)-3-amino-1-chloro-4-(3,5-difluorophenyl)butan-2-ol.

12. A compound of the formula:



13. A compound of the formula:



5

where

X<sub>1</sub> is -Cl, -Br or imidazolyl; and

PROT is a nitrogen protecting group.


10 14. A compound according to claim 13 where PROT is t-butoxycarbonyl or benzyloxycarbonyl.

15. A compound according to claim 13 where X<sub>1</sub> is -Cl.

15 16. A compound (XI) according to claim 13 where the compound is t-butyl-(1S)-2-chloro-1-[3,5-difluorobenzyl]-2-oxoethylcarbamate.

17. A compound of the formula:

20



where

PROT is a nitrogen protecting group; and

R<sub>1</sub> is selected from:

- 5 (I) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one chloro;  
 (II) -CH<sub>2</sub>-CH=CH<sub>2</sub>;  
 (III) phenyl optionally substituted with one nitro,  
 halogen, or cyano; and  
 (IV) benzyl optionally substituted on phenyl with nitro,  
 10 halogen, or cyano.

18. A compound according to claim 17 where the PROT is t-butoxycarbonyl or benzyloxycarbonyl.


- 15 19. A compound according to claim 17 where R<sub>1</sub> is C<sub>1</sub>-C<sub>2</sub> alkyl.

20. A compound according to claim 19 where R<sub>1</sub> is C<sub>1</sub> alkyl.

20

21. A compound according to claim 17 which is methyl (2Z)-2-[[ (benzyloxy) carbonyl]-3-(3,5-difluorophenyl)-2-propenonate.

- 25 22. A compound of the formula:



where

R<sub>1</sub> is selected from:

- (I) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one chloro;

(II)  $-\text{CH}_2-\text{CH}=\text{CH}_2$ ;

(III) phenyl optionally substituted with one nitro, halogen, or cyano; and


(IV) benzyl optionally substituted on phenyl with nitro, halogen, or cyano.

23. A compound according to claim 22 where  $R_1$  is  $\text{C}_1$ - $\text{C}_2$  alkyl.

24. A compound according to claim 23 where  $R_1$  is  $\text{C}_1$  alkyl.

25. A compound according to claim 22 which is methyl (2S)-2-amino-3-(3,5-difluorophenyl)propanoate.

26. A process for the preparation of an ester of the formula:



where

$R_1$  is selected from the group consisting of:


(I)  $\text{C}_1$ - $\text{C}_4$  alkyl optionally substituted with one  $-\text{Cl}$ ;

(II)  $-\text{CH}_2-\text{CH}=\text{CH}_2$ , and

(III) benzyl optionally substituted on phenyl with nitro, halogen, or cyano; and

PROT is a nitrogen protecting group, which process comprises:

(a) contacting a protected amino acid of the formula (I)



(I)

where PROT is as defined above, with a base and

(b) contacting the mixture of (a) with an alkylating agent  
5 of the formula

(a)  $X_4-C_1-C_4$  alkyl optionally substituted with one of  
iodo, bromo, or chloro;

(a') dimethylsulfate;

(b)  $X_4-CH_2-CH=CH_2$ ,

10 (c)  $X_4$ -benzyl where the phenyl ring is optionally  
substituted with nitro, halogen, cyano; and  
 $X_4$  is iodo, bromo, chloro, -O-tosylate, -O-mesylate  
or -O-triflate.

15 27. A process according to claim 26 where PROT is *t*-  
butoxycarbonyl or benzyloxycarbonyl.

28. A process according to claim 26 where the base is  
hydroxide, carbonate, bicarbonate, LDA, *n*-( $C_1-C_8$  alkyl)lithium,  
20 LiHMDS, NaHMDS or KHMDS.

29. A process according to claim 28 where the base is  
hydroxide, carbonate, or bicarbonate.

25 30. A process according to claim 29 where the base is  
carbonate.

31. A process according to claim 26 where the alkylating  
agent is dimethylsulfate, methyl iodide or methyl triflate.

32. A process according to claim 31 where the alkylating agent is dimethylsulfate.


5 33. A process according to claim 26 where when the base is LDA, *n*-(C<sub>1</sub>-C<sub>8</sub> alkyl)lithium, LiHMDS or KHDMS, and the mixture of (a) is cooled to a range of from about -78° to about 25° prior to the addition of the base.

10 34. A process according to claim 33 where the mixture of (a) is cooled to a range of from about -20° to about 25° prior to the addition of the base.

15 35. A process according to claim 26 where the mixture of (b) is heated from about 20° to about 50°.

36. A process according to claim 26 where the ester (II) is (2S)-2-[(*tert*-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid methyl ester.

37. A process for the preparation of an ester of the formula:




25 where

R<sub>1</sub> is phenyl optionally substituted with one of nitro, halogen, or cyano; and

PROT is a nitrogen protecting group, which process comprises:

(1) contacting a protected amino acid of the formula (I)



(I)

where PROT is as defined above, with an activating agent

(2) contacting the mixture of (1) with a phenoxy compound of the formula

(d) HO- $\phi$  where  $\phi$  is optionally substituted with one:

(A) -NO<sub>2</sub>,

(B) -F, -Cl, -Br, -I,


(C) -C $\equiv$ N.

38. A process according to claim 37 where PROT is t-butoxycarbonyl and benzyloxycarbonyl.

39. A process according to claim 37 where the activating agent is CDI.

40. A process according to claim 37 where the activating agent is DCC.

41. A process for the preparation of a ketone of formula III:



III

where

PROT is a nitrogen protecting group, and

$R_2$  is:

chloro, bromo, or

-Si( $R_{21}$ )<sub>3</sub> where each  $R_{21}$  is independently

C<sub>1</sub>-C<sub>5</sub> alkyl,

-N( $R_{23}$ )( $R_{24}$ ) where  $R_{23}$  and  $R_{24}$  are the same or different and represent

C<sub>1</sub>-C<sub>5</sub> alkyl, or


where NR<sub>23</sub>R<sub>24</sub> represents piperidinyl,

piperazinyl, or morpholinyl,

phenyl optionally substituted with 1, 2, or 3 of C<sub>1</sub>-C<sub>2</sub> alkyl, with the proviso that at least one of the  $R_{21}$  groups is optionally substituted phenyl,

which process comprises:

(a) forming a mixture of an ester of formula II and a dihalogenated methane,  $R_2CH_2X^2$ , where  $R_2$  is as defined above and where  $X^2$  is -Br or -I;



(II)

where

$R_1$  is selected from the group consisting of:

(I) C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with one -Cl;

(II) -CH<sub>2</sub>-CH=CH<sub>2</sub>,

(III) phenyl optionally substituted with one nitro, halogen, or cyano; and

(IV) benzyl optionally substituted on phenyl with nitro, halogen, or cyano; and

PROT is as defined above,

- (b) adding a base to the mixture from (a);
- (c) acidifying the mixture of (b).

5        42. A process according to claim 41 where PROT is t-butoxycarbonyl or benzyloxycarbonyl.

43. A process according to claim 41 where  $R_1$  is  $C_1$ - $C_2$  alkyl.

10

44. A process according to claim 43 where  $R_1$  is  $C_1$  alkyl.

45. A process according to claim 41 where  $R_2$  is chloro or bromo.

15

46. A process according to claim 45 where  $R_2$  is chloro.

47. A process according to claim 41 where  $CH_2R_2X^2$  is present in an amount of from about 1 to about 1.5 equivalents based on the amount of ester II.

20

48. A process according to claim 41 where  $X^2$  is iodo.

49. A process according to claim 41 where the strong base is LDA, ( $C_1$ - $C_8$  alkyl)lithium, LiHMDS, NaHMDS or KHMDS.

25

50. A process according to claim 49 where the strong base is LDA.

51. A process according to claim 41 where the strong base is present in an amount of from about 2 to about 2.5 equivalents based on the amount of ester II.

30

52. A process according to claim 41 where a second portion of base is added, where the second portion of base is (C<sub>1</sub>-C<sub>4</sub>)alkyl lithium, phenyl lithium, (C<sub>1</sub>-C<sub>4</sub>)alkyl-Grignard or phenyl-Grignard.

5

53. A process according to claim 52 where the second base is phenyl lithium, *n*-butyl lithium, sec-butyllithium, tert-butyllithium, methyllithium, methyl magnesium bromide, methyl magnesium chloride, phenyl magnesium bromide or phenyl magnesium chloride.

10

54. A process according to claim 53 where the second base is *n*-butyl lithium.

15

55. A process according to claim 41 where amount of the second is from about 1 to about 1.5 equivalents based on the amount of the ester II.

56. A process according to claim 41 where the acidifying is carried out using an acid having a pK<sub>a</sub> of less than about 10.

20

57. A process according to claim 56 where the acid is selected from the group consisting of acetic, sulfuric, hydrochloric, citric, phosphoric, nitric, paratoluenesulfonic, and benzoic acids and mixtures thereof.


25

58. A process according to claim 57 where the acid is hydrochloric acid or acetic acid.

30

59. A process for according to claim 41 where the ketone (III) is *tert*-butyl (1*S*)-3-chloro-1-(3,5-difluorobenzyl)-2-oxopropylcarbamate.

60. A process for the preparation of a ketone of formula (III)



(III)


where PROT is a nitrogen protecting group, and

where R<sub>2</sub> is -Cl or -Br,

which process comprises:

(a) contacting an acid R<sub>2</sub>-CH<sub>2</sub>-COOH, where R<sub>2</sub> is as defined above, with a base;

(b) contacting the mixture of (a) with an ester of formula (II)



(II)

where R<sub>1</sub> is selected from the group consisting of:

(I) C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with one -Cl;

(II) -CH<sub>2</sub>-CH=CH<sub>2</sub>,

(III) phenyl optionally substituted with one:

(A) -NO<sub>2</sub>,

(B) -F, -Cl, -Br, -I,

(C) -C≡N, and

(IV) -CH<sub>2</sub>-φ where the -φ ring is optionally substituted with

(A) -NO<sub>2</sub>,

(B) -F, -Cl, -Br, -I,

(C) -C≡N and

where PROT is as defined above; and

(c) acidifying the mixture of (b).

5

61. A process according to claim 60 where PROT is t-butoxycarbonyl or benzyloxycarbonyl.

62. A process according to claim 60 where R<sub>1</sub> is C<sub>1</sub>-C<sub>2</sub>  
10 alkyl.

63. A process according to claim 62 where R<sub>1</sub> is C<sub>1</sub> alkyl.

64. A process according to claim 60 where R<sub>2</sub> is chloro or  
15 bromo.

65. A process according to claim 64 where R<sub>2</sub> is chloro.

66. A process according to claim 60 where the base is a  
20 strong base and is LDA, (C<sub>1</sub>-C<sub>8</sub> alkyl)lithium, LiHMDS, NaHMDS or  
KHMDS.

67. A process according to claim 66 where the strong base  
is LDA.

25

68. A process according to claim 60 where the strong base  
is present in an amount of from about 3 to about 3.5  
equivalents based on the amount of ester II.


69. A process according to claim 60 where the acidifying  
is carried out using an acid having a pK<sub>a</sub> of less than about  
30 10.

70. A process according to claim 69 where the acid is selected from the group consisting of acetic, sulfuric, hydrochloric, citric, phosphoric and benzoic acids and mixtures thereof.

5

71. A process according to claim 70 where the acid is hydrochloric or acetic acid.

72. A process for the preparation of a compound of formula (XI)




(XI)

where  $X_1$  is where  $X_1$  is -Cl, -Br or imidazolyl;

where PROT is a nitrogen protecting group,

15 which process comprises

(1) contacting a protected amino acid of formula (I)



(I)

where PROT is as defined above,

20 with thionyl chloride,  $\text{SO}_2\text{Cl}_2$ , phosphorous trichloride, oxalyl chloride, phosphorous tribromide, triphenylphosphorous dibromide, oxalyl bromide, 1,2-phenylenetrichlorophosphate, 2,4,6-trichloro-1,3,5-triazine or CDI.


73. A process according to claim 72 where PROT is *t*-butoxycarbonyl or benzyloxycarbonyl.

74. A process according to claim 72 where X<sub>1</sub> is chloro.

75. A process according to claim 72 where the compound of formula XI is contacted with thionyl chloride or oxalyl chloride.

76. A process according to claim 72 where the process produces *t*-butyl-(1*S*)-2-chloro-1-[3,5-difluorobenzyl]-2-oxoethylcarbamate.

77. A process for the preparation of ketone of formula (III)




(III)

where

R<sub>2</sub> is -Cl or -Br; and

PROT is a nitrogen protecting group,  
which process comprises:

(1) contacting a compound of formula (XI)



(XI)

where

$X_1$  is -Cl, -Br and imidazolyl; and

PROT is as defined above,

with  $\text{LiCH}_2\text{Cl}$  or  $\text{LiCH}_2\text{Br}$ .

5

78. A process according to claim 77 where PROT is *t*-butoxycarbonyl or benzyloxycarbonyl.

79. A process according to claim 77 where  $X_1$  is chloro.


10

80. A process according to claim 77 where  $R_2$  is chloro.

81. A process according to claim 77 where the process produces *tert*-butyl (1*S*)-3-chloro-1-(3,5-difluorobenzyl)-2-oxopropylcarbamate.

15

82. A process for the preparation of an ester of the formula (II)



(II)

20

where

$R_1$  is selected from the group consisting of:

(I)  $\text{C}_1\text{-C}_4$  alkyl optionally substituted with one -Cl;

(II)  $-\text{CH}_2\text{-CH=CH}_2$ ,

(III) phenyl optionally substituted with one:

(A)  $-\text{NO}_2$ ,

(B) -F, -Cl, -Br, -I,

(C)  $-\text{C}\equiv\text{N}$ ,

25

(IV)  $-\text{CH}_2-\phi$  where the  $-\phi$  ring is optionally substituted with


(A)  $-\text{NO}_2$ ,

(B)  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ ,

5 (C)  $-\text{C}\equiv\text{N}$ ; and

PROT is a nitrogen protecting group,  
which process comprises:

(b) treating a compound of formula XIV



XIV

10 where  $R_1$  and PROT are as defined above; and  
in a solvent with hydrogen in the presence of a hydrogenation  
catalyst at a pressure of from 1 atmosphere to about 100 psi.

15 83. A process according to claim 82 where the PROT is *t*-  
butoxycarbonyl or benzyloxycarbonyl.

84. A process according to claim 82 where  $R_1$  is  $\text{C}_1$ - $\text{C}_2$   
alkyl.

20 85. A process according to claim 84 where  $R_1$  is  $\text{C}_1$  alkyl.

86. A process according to claim 82 where the solvent of  
step (a) is degassed.

25 87. A process according to claim 82 where the reaction  
vessel is purged of oxygen after step (1) and before step (2).

88. A process according to claim 69 where the hydrogenation catalyst is a compound of the formula  $[\text{Rh}(\text{diene})\text{L}]^+\text{X}^-$

where Rh is rhodium;

5 where diene is cyclooctadiene and nonbornadiene;

where L is DIPMAP, MeDuPhos, EtDuPhos, Binaphane, f-Binaphane, Me-KetalPhos, Me-f-KetalPhos, Et-f-KetalPhos, BINAP, DIOP, BPPFA, BPPM, CHIRAPHOS, PROPHOS, NORPHOS, CYCLOPHOS, BDPP, DEGPPOS, PNNP and

10 where  $\text{X}^-$  is  $\text{ClO}_4^-$ ,  $\text{BF}_4^-$ ,  $\text{CF}_3\text{-SO}_3^-$ ,  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{PF}_6^-$  and  $\text{SbF}_6^-$ .

89. A process according to claim 88 where the hydrogenation catalyst is DIPMAP.

15 90. A process according to claim 88 where the hydrogenation catalyst is EtDuPhos.

91. A process according to claim 82 where the reaction temperature is from about  $0^\circ$  to about reflux.

20

92. A process according to claim 91 where the reaction temperature is from about  $0^\circ$  to about  $25^\circ$ .

25 93. A process according to claim 82 where the reaction pressure is from about 1 atmosphere to about 70 psig.

94. A process according to claim 93 where the reaction pressure is from about 10 psig to about 40 psig.


30 95. A process according to claim 82 where the ester (II) is obtained in greater than 90% enantiomeric purity.

96. A process according to claim 95 where the

ester (II) is obtained in greater than 95% enantiomeric purity.

97. A process according to claim 82 where the  
5 ester (II) is methyl (2S)-2-{[(benzyloxy)carbonyl]amino}-3-(3,5-difluorophenyl)propanoate.

98. A process for preparing an epoxide of formula V-R



10

comprising


(a) converting an ester of Formula II into a ketone of formula III;

15 (b) reducing the ketone to the corresponding alcohol of formula IV; and

(c) treating the alcohol with a base to yield the epoxide.

99. A process according to claim 98 further comprising  
20 esterifying an acid of formula (O) to generate the ester of formula II.

100. A process for preparing a compound of formula (XX)



25

wherein

R<sub>57</sub> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, or benzyl;

$R_1$  is  $-(CH_2)_{1-2}-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$ ,  $-CH_2-CH_2-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$ , or

$C_1-C_6$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -F, -Cl, -Br, -I, -OH, =O, -SH,  $-C\equiv N$ ,  $-CF_3$ ,  $-C_1-C_3$  alkoxy, amino, mono- or dialkylamino,  $-N(R)C(O)R'$ -,  $-OC(=O)$ -amino and  $-OC(=O)$ -mono- or dialkylamino, or

$C_2-C_6$  alkenyl or  $C_2-C_6$  alkynyl, each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen, -F, -Cl, -Br, -I, -OH, -SH,  $-C\equiv N$ ,  $-CF_3$ ,  $C_1-C_3$  alkoxy, amino, and mono- or dialkylamino, or

aryl, heteroaryl, heterocyclyl,  $-C_1-C_6$  alkyl-aryl,  $-C_1-C_6$  alkyl-heteroaryl, or  $-C_1-C_6$  alkyl-heterocyclyl, where the ring portions of each are optionally substituted with 1, 2, 3, or 4 groups independently selected from halogen, -F, -Cl, -Br, -I, -OH, -SH,  $-C\equiv N$ ,  $-NR_7R'_7$ ,  $-C(=O)-(C_1-C_4)$  alkyl,  $-SO_2$ -amino,  $-SO_2$ -mono or dialkylamino,  $-C(=O)$ -amino,  $-C(=O)$ -mono or dialkylamino,  $-SO_2-(C_1-C_4)$  alkyl,  $-CO_2R$ ,  $-N(R)COR'$ , or  $-N(R)SO_2R'$  or

$-C_1-C_6$  alkoxy optionally substituted with 1, 2, or 3 groups which are independently a selected from halogen, or

$C_3-C_7$  cycloalkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -F, -Cl, -Br, -I, -OH, -SH,  $-C\equiv N$ ,  $-CF_3$ ,  $C_1-C_3$  alkoxy, amino,  $-C_1-C_6$  alkyl and mono- or dialkylamino, or

$C_1-C_{10}$  alkyl optionally substituted with 1, 2, or 3 groups independently selected from halogen, -F, -Cl, -Br, -I, -OH, -SH,  $-C\equiv N$ ,  $-CF_3$ ,  $-C_1-C_3$

alkoxy, amino, mono- or dialkylamino and -C<sub>1</sub>-C<sub>3</sub> alkyl, or

C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl, each of which is optionally substituted with 1, 2, or 3 groups independently selected from halogen, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, -C<sub>1</sub>-C<sub>6</sub> alkyl and mono- or dialkylamino; and the heterocyclyl group is optionally further substituted with oxo;

R<sub>7</sub> and R<sub>7</sub>' are independently H or -C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sub>2</sub> and R<sub>3</sub> are independently selected from the group consisting of H; C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or three substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, and -NR<sub>30</sub>R<sub>31</sub>; -(CH<sub>2</sub>)<sub>0-4</sub>-aryl; -(CH<sub>2</sub>)<sub>0-4</sub>-heteroaryl; -(CH<sub>2</sub>)<sub>0-4</sub>-heterocycle; C<sub>2</sub>-C<sub>6</sub> alkenyl optionally substituted with one, two or three substituents independently selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, and -NR<sub>30</sub>R<sub>31</sub>; C<sub>2</sub>-C<sub>6</sub> alkynyl optionally substituted with one, two or three substituents independently selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, and -NR<sub>30</sub>R<sub>31</sub>; and -(CH<sub>2</sub>)<sub>0-4</sub>-C<sub>3</sub>-C<sub>7</sub> cycloalkyl, wherein the cycloalkyl group is optionally substituted with one, two or three substituents independently selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, and -NR<sub>30</sub>R<sub>31</sub>;

or

R<sub>2</sub>, R<sub>3</sub> and the carbon to which they are attached form a carbocycle of three, four, five, six, or seven carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced by a heteroatom independently selected from the group consisting of -O-, -S-, -SO<sub>2</sub>-, and -NR<sub>22</sub>-; wherein

R<sub>30</sub> and R<sub>31</sub> at each occurrence are independently H, or C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sub>22</sub> is selected from the group consisting of -H, -C<sub>1</sub>-C<sub>6</sub> alkyl, hydroxy C<sub>1</sub>-C<sub>6</sub> alkyl, amino C<sub>1</sub>-C<sub>6</sub> alkyl; halo C<sub>1</sub>-C<sub>6</sub> alkyl; -C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl, -C<sub>1</sub>-C<sub>6</sub> alkyl chain with one double bond and one triple bond, aryl, heteroaryl, and heterocycloalkyl;

10 R<sub>C</sub> is selected from the group consisting of C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of R<sub>205</sub>, -OC(=O)NR<sub>235</sub>R<sub>240</sub>, -S(=O)<sub>0-2</sub>R<sub>235</sub>, -NR<sub>235</sub>C(=O)NR<sub>235</sub>R<sub>240</sub>, -C(=O)NR<sub>235</sub>R<sub>240</sub>, and -S(=O)<sub>2</sub>NR<sub>235</sub>R<sub>240</sub>; -(CH<sub>2</sub>)<sub>0-3</sub>-(C<sub>3</sub>-C<sub>8</sub>) cycloalkyl  
 15 wherein the cycloalkyl is optionally substituted with 1, 2, or 3 groups independently selected from the group consisting of R<sub>205</sub>, -CO<sub>2</sub>H, and -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl);  
 -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl, -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocycloalkyl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-heteroaryl; -  
 20 (CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-heterocycloalkyl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-aryl-aryl;  
 -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-aryl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-heterocycloalkyl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heteroaryl-heteroaryl; -  
 (CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocycloalkyl-heteroaryl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-heterocycloalkyl-heterocycloalkyl; -(CR<sub>245</sub>R<sub>250</sub>)<sub>0-4</sub>-  
 25 heterocycloalkyl-aryl; -[C(R<sub>255</sub>)(R<sub>260</sub>)]<sub>1-3</sub>-CO-N-(R<sub>255</sub>)<sub>2</sub>; -CH(aryl)<sub>2</sub>; -CH(heteroaryl)<sub>2</sub>; -CH(heterocycloalkyl)<sub>2</sub>;  
 -CH(aryl)(heteroaryl); cyclopentyl, cyclohexyl, or cycloheptyl ring fused to aryl, heteroaryl, or heterocycloalkyl wherein one carbon of the cyclopentyl,  
 30 cyclohexyl, or cycloheptyl is optionally replaced with NH, NR<sub>215</sub>, O, or S(=O)<sub>0-2</sub>, and wherein the cyclopentyl, cyclohexyl, or -cycloheptyl group can be optionally substituted with 1 or 2 groups that are independently R<sub>205</sub> or =O; -CO-NR<sub>235</sub>R<sub>240</sub>; or -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl); C<sub>2</sub>-C<sub>10</sub> alkenyl

optionally substituted with 1, 2, or 3  $R_{205}$  groups;  $C_2$ - $C_{10}$  alkynyl optionally substituted with 1, 2, or 3  $R_{205}$  groups;  $-(CH_2)_{0-1}-CH((CH_2)_{0-6}-OH)-(CH_2)_{0-1}$ -aryl;  $-(CH_2)_{0-1}-CHRC_6-(CH_2)_{0-1}$ -heteroaryl;  $-CH(-aryl \text{ or } -heteroaryl)-CO-O(C_1-C_4 \text{ alkyl})$ ;  
 5  $-CH(-CH_2-OH)-CH(OH)$ -phenyl- $NO_2$ ,  $(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})-OH$ ,  $-CH_2-NH-CH_2-CH(-O-CH_2-CH_3)_2$ ,  $-H$ , and  $-(CH_2)_{0-6}-C(=NR_{235})(NR_{235}R_{240})$ ; wherein

each aryl is optionally substituted with 1, 2, or 3  $R_{200}$ ;  
 each heteroaryl is optionally substituted with 1, 2, 3, or  
 10 4  $R_{200}$ ;

each heterocycloalkyl is optionally substituted with 1, 2, 3, or 4  $R_{210}$ ;

$R_{200}$  at each occurrence is independently selected from the group consisting of  $C_1$ - $C_6$  alkyl optionally  
 15 substituted with 1, 2, or 3  $R_{205}$  groups;  $OH$ ;  $-NO_2$ ; halogen;  $-CO_2H$ ;  $C\equiv N$ ;  $-(CH_2)_{0-4}-CO-NR_{220}R_{225}$ ;  $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$ ;  $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl})$ ;  $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl})$ ;  $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$ ;  
 20  $-(CH_2)_{0-4}-CO$ -aryl;  $-(CH_2)_{0-4}-CO$ -heteroaryl;  $-(CH_2)_{0-4}-CO$ -heterocycloalkyl;  $-(CH_2)_{0-4}-CO-O-R_{215}$ ;  $-(CH_2)_{0-4}-SO_2-NR_{220}R_{225}$ ;  $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$ ;  $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$ ;  $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$ ;  $-(CH_2)_{0-4}-N(H \text{ or } R_{215})-CO-O-R_{215}$ ;  $-(CH_2)_{0-4}-N(H \text{ or } R_{215})-CO-N(R_{215})_2$ ;  
 25  $-(CH_2)_{0-4}-N-CS-N(R_{215})_2$ ;  $-(CH_2)_{0-4}-N(-H \text{ or } R_{215})-CO-R_{220}$ ;  $-(CH_2)_{0-4}-NR_{220}R_{225}$ ;  $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$ ;  $-(CH_2)_{0-4}-O-P(O)-(OR_{240})_2$ ;  $-(CH_2)_{0-4}-O-CO-N(R_{215})_2$ ;  $-(CH_2)_{0-4}-O-CS-N(R_{215})_2$ ;  $-(CH_2)_{0-4}-O-(R_{215})_2$ ;  $-(CH_2)_{0-4}-O-(R_{215})_2-COOH$ ;  
 30  $-(CH_2)_{0-4}-S-(R_{215})_2$ ;  $-(CH_2)_{0-4}-O-(C_1-C_6 \text{ alkyl})$  optionally substituted with 1, 2, 3, or 5  $-F$ ;  $C_3$ - $C_7$  cycloalkyl;  $C_2$ - $C_6$  alkenyl optionally substituted with 1 or 2  $R_{205}$  groups;  $C_2$ - $C_6$  alkynyl optionally substituted with 1 or 2  $R_{205}$  groups;  $-(CH_2)_{0-4}-N(H \text{ or } R_{215})-SO_2-R_{220}$ ; and  $-(CH_2)_{0-4}-C_3-C_7 \text{ cycloalkyl}$ ;

wherein each aryl group at each occurrence is  
optionally substituted with 1, 2, or 3 groups  
that are independently  $R_{205}$ ,  $R_{210}$  or  $C_1$ - $C_6$  alkyl  
substituted with 1, 2, or 3 groups that are  
independently  $R_{205}$  or  $R_{210}$ ;

wherein each heterocycloalkyl group at each  
occurrence is optionally substituted with 1, 2,  
or 3 groups that are independently  $R_{210}$ ;

wherein each heteroaryl group at each occurrence is  
optionally substituted with 1, 2, or 3 groups  
that are independently  $R_{205}$ ,  $R_{210}$ , or  $C_1$ - $C_6$  alkyl  
substituted with 1, 2, or 3 groups that are  
independently  $R_{205}$  or  $R_{210}$ ;

$R_{205}$  at each occurrence is independently selected from the  
group consisting of  $C_1$ - $C_6$  alkyl, halogen, -OH, -O-  
phenyl, -SH, -C $\equiv$ N, -CF $_3$ ,  $C_1$ - $C_6$  alkoxy, NH $_2$ , NH( $C_1$ - $C_6$   
alkyl), and N-( $C_1$ - $C_6$  alkyl)( $C_1$ - $C_6$  alkyl);

$R_{210}$  at each occurrence is independently selected from the  
group consisting of  $C_1$ - $C_6$  alkyl optionally  
substituted with 1, 2, or 3  $R_{205}$  groups;  $C_2$ - $C_6$  alkenyl  
optionally substituted with 1, 2, or 3  $R_{205}$  groups;  
 $C_2$ - $C_6$  alkynyl optionally substituted with 1, 2, or 3  
 $R_{205}$  groups; halogen;  $C_1$ - $C_6$  alkoxy;  $C_1$ - $C_6$  haloalkoxy;  
-NR $_{220}$ R $_{225}$ ; OH; C $\equiv$ N;  $C_3$ - $C_7$  cycloalkyl optionally  
substituted with 1, 2, or 3  $R_{205}$  groups; -CO-( $C_1$ - $C_4$   
alkyl); -SO $_2$ -NR $_{235}$ R $_{240}$ ; -CO-NR $_{235}$ R $_{240}$ ; -SO $_2$ -( $C_1$ - $C_4$  alkyl);  
and =O; wherein

$R_{215}$  at each occurrence is independently selected from  
the group consisting of  $C_1$ - $C_6$  alkyl, -(CH $_2$ ) $_{0-2}$ -  
(aryl),  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl,  $C_3$ - $C_7$   
cycloalkyl, and -(CH $_2$ ) $_{0-2}$ -(heteroaryl), -(CH $_2$ ) $_{0-2}$ -  
(heterocycloalkyl); wherein the aryl group at  
each occurrence is optionally substituted with

1, 2, or 3 groups that are independently  $R_{205}$  or  $R_{210}$ ; wherein the heterocycloalkyl group at each occurrence is optionally substituted with 1, 2, or 3  $R_{210}$ ; wherein each heteroaryl group at each occurrence is optionally substituted with 1, 2, or 3  $R_{210}$ ;

$R_{220}$  and  $R_{225}$  at each occurrence are independently selected from the group consisting of -H, - $C_1$ - $C_6$  alkyl, hydroxy  $C_1$ - $C_6$  alkyl, amino  $C_1$ - $C_6$  alkyl; halo  $C_1$ - $C_6$  alkyl; - $C_3$ - $C_7$  cycloalkyl, -( $C_1$ - $C_2$  alkyl)-( $C_3$ - $C_7$  cycloalkyl), -( $C_1$ - $C_6$  alkyl)-O-( $C_1$ - $C_3$  alkyl), - $C_2$ - $C_6$  alkenyl, - $C_2$ - $C_6$  alkynyl, - $C_1$ - $C_6$  alkyl chain with one double bond and one triple bond, -aryl, -heteroaryl, and -heterocycloalkyl; wherein the aryl group at each occurrence is optionally substituted with 1, 2, or 3 groups that are independently  $R_{205}$  or  $R_{210}$ ; wherein the heterocycloalkyl group at each occurrence is optionally substituted with 1, 2, or 3  $R_{210}$ ; wherein each heteroaryl group at each occurrence is optionally substituted with 1, 2, or 3  $R_{210}$ ;

$R_{235}$  and  $R_{240}$  at each occurrence are independently H, or  $C_1$ - $C_6$  alkyl;

$R_{245}$  and  $R_{250}$  at each occurrence are independently selected from the group consisting of H,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  hydroxyalkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  haloalkoxy, -( $CH_2$ )<sub>0-4</sub>- $C_3$ - $C_7$  cycloalkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl, and phenyl; or

$R_{245}$  and  $R_{250}$  are taken together with the carbon to which they are attached to form a carbocycle of 3, 4, 5, 6, or 7 carbon atoms, optionally where one carbon atom

is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO<sub>2</sub>-, and -NR<sub>220</sub>-;

R<sub>255</sub> and R<sub>260</sub> at each occurrence are independently selected from the group consisting of H; C<sub>1</sub>-C<sub>6</sub> alkyl

optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; C<sub>2</sub>-C<sub>6</sub> alkenyl optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; C<sub>2</sub>-C<sub>6</sub> alkynyl optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; -(CH<sub>2</sub>)<sub>1-2</sub>-S(O)<sub>0-2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl); -(CH<sub>2</sub>)<sub>0-4</sub>-C<sub>3</sub>-C<sub>7</sub> cycloalkyl optionally substituted with 1, 2, or 3 R<sub>205</sub> groups; -(C<sub>1</sub>-C<sub>4</sub> alkyl)-aryl; -(C<sub>1</sub>-C<sub>4</sub> alkyl)-heteroaryl; -(C<sub>1</sub>-C<sub>4</sub> alkyl)-heterocycloalkyl; -aryl; -heteroaryl; -heterocycloalkyl; -(CH<sub>2</sub>)<sub>1-4</sub>-R<sub>265</sub>-(CH<sub>2</sub>)<sub>0-4</sub>-aryl; -(CH<sub>2</sub>)<sub>1-4</sub>-R<sub>265</sub>-(CH<sub>2</sub>)<sub>0-4</sub>-heteroaryl; and; -(CH<sub>2</sub>)<sub>1-4</sub>-R<sub>265</sub>-(CH<sub>2</sub>)<sub>0-4</sub>-heterocycloalkyl; wherein

R<sub>265</sub> at each occurrence is independently -O-, -S- or -N(C<sub>1</sub>-C<sub>6</sub> alkyl)-;

each aryl or phenyl is optionally substituted with 1, 2, or 3 groups that are independently R<sub>205</sub>, R<sub>210</sub>, or C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 1, 2, or 3

groups that are independently R<sub>205</sub> or R<sub>210</sub>;

each heteroaryl is optionally substituted with 1, 2, 3, or 4 R<sub>200</sub>,

each heterocycloalkyl is optionally substituted with 1, 2, 3, or 4 R<sub>210</sub>;

R<sub>N</sub> is -C(=O)-(CRR')<sub>0-6</sub>R<sub>100</sub>, R'<sub>100</sub>, -SO<sub>2</sub>R'<sub>100</sub>, -(CRR')<sub>1-6</sub>R'<sub>100</sub>, -C(=O)-(CRR')-O-R'<sub>100</sub>, -C(=O)-(CRR')-S-R'<sub>100</sub> or -C(=O)-(CRR')-NR<sub>100</sub>-R'<sub>100</sub>;

R<sub>100</sub> and R'<sub>100</sub> are independently aryl, heteroaryl, -aryl-W-aryl, -aryl-W-heteroaryl, -aryl-W-heterocyclyl, -heteroaryl-W-aryl, -heteroaryl-W-heteroaryl, -heteroaryl-W-heterocyclyl, -heterocyclyl-W-aryl, -heterocyclyl-W-heteroaryl, -heterocyclyl-W-heterocyclyl, -C(=O)-CH[(CH<sub>2</sub>)<sub>0-2</sub>-O-R<sub>7</sub>]- (CH<sub>2</sub>)<sub>0-2</sub>-aryl, -C(=O)-CH[(CH<sub>2</sub>)<sub>0-2</sub>-O-R<sub>7</sub>]- (CH<sub>2</sub>)<sub>0-2</sub>-heterocyclyl, or -C(=O)-CH[(CH<sub>2</sub>)<sub>0-2</sub>-O-R<sub>7</sub>]- (CH<sub>2</sub>)<sub>0-2</sub>-

heteroaryl, where the ring portions of each are optionally substituted with 1, 2, or 3 groups independently selected from

-OR, -NO<sub>2</sub>, halogen, -C≡N, -SR, -SO<sub>2</sub>R<sub>145</sub>, -C(=O)R, -OCF<sub>3</sub>,  
 5 -CF<sub>3</sub>, -O-P(=O)(OR)(OR'), -N(R)(COR'), -N(R)(SO<sub>2</sub>R<sub>145</sub>),  
 -(CH<sub>2</sub>)<sub>0-4</sub>-CO-NR<sub>105</sub>R'<sub>105</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-O-(CH<sub>2</sub>)<sub>0-4</sub>-CONRR',  
 -(CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>1</sub>-C<sub>12</sub> alkyl), -(CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>2</sub>-C<sub>12</sub>  
 alkenyl), -(CH<sub>2</sub>)<sub>0-4</sub>-CO-(C<sub>2</sub>-C<sub>12</sub> alkynyl), -(CH<sub>2</sub>)<sub>0-4</sub>-CO-  
 (C<sub>3</sub>-C<sub>7</sub> cycloalkyl), -(CH<sub>2</sub>)<sub>0-4</sub>-R<sub>110</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-R<sub>120</sub>,  
 10 -(CH<sub>2</sub>)<sub>0-4</sub>-R<sub>130</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-CO-R<sub>110</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-CO-R<sub>120</sub>,  
 -(CH<sub>2</sub>)<sub>0-4</sub>-CO-R<sub>130</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-CO-R<sub>140</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-CO-O-R<sub>150</sub>,  
 -(CH<sub>2</sub>)<sub>0-4</sub>-SO<sub>2</sub>-NR<sub>105</sub>R'<sub>105</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-SO-(C<sub>1</sub>-C<sub>8</sub> alkyl),  
 -(CH<sub>2</sub>)<sub>0-4</sub>-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>12</sub> alkyl), -(CH<sub>2</sub>)<sub>0-4</sub>-SO<sub>2</sub>-(C<sub>3</sub>-C<sub>7</sub>  
 cycloalkyl), -(CH<sub>2</sub>)<sub>0-4</sub>-N(H or R<sub>150</sub>)-CO-O-R<sub>150</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-  
 15 N(H or R<sub>150</sub>)-CO-N(R<sub>150</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-N(H or R<sub>150</sub>)-CS-  
 N(R<sub>150</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-N(-H or R<sub>150</sub>)-CO-R<sub>105</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-  
 NR<sub>105</sub>R'<sub>105</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-R<sub>140</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-O-CO-(C<sub>1</sub>-C<sub>6</sub> alkyl),  
 -(CH<sub>2</sub>)<sub>0-4</sub>-O-P(O)-(O-R<sub>110</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-O-CO-N(R<sub>150</sub>)<sub>2</sub>,  
 -(CH<sub>2</sub>)<sub>0-4</sub>-O-CS-N(R<sub>150</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-4</sub>-O-(R<sub>150</sub>), -(CH<sub>2</sub>)<sub>0-4</sub>-O-  
 20 (R<sub>155</sub>)-COOH, -(CH<sub>2</sub>)<sub>0-4</sub>-S-(R<sub>150</sub>), C<sub>3</sub>-C<sub>7</sub> cycloalkyl,  
 -(CH<sub>2</sub>)<sub>0-4</sub>-N(-H or R<sub>150</sub>)-SO<sub>2</sub>-R<sub>7</sub>, or -(CH<sub>2</sub>)<sub>0-4</sub>-C<sub>3</sub>-C<sub>7</sub>  
 cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, or

R<sub>100</sub> is C<sub>1</sub>-C<sub>10</sub> alkyl optionally substituted with 1, 2, or 3 R<sub>115</sub> groups, wherein

25 R<sub>115</sub> at each occurrence is independently halogen, -OH,  
 -CO<sub>2</sub>R, -C<sub>1</sub>-C<sub>6</sub> thioalkoxy, -CO<sub>2</sub>-phenyl, -NR<sub>105</sub>R'<sub>7</sub>, -SO<sub>2</sub>-  
 (C<sub>1</sub>-C<sub>8</sub> alkyl), -C(=O)R<sub>180</sub>, R<sub>180</sub>, -CONR<sub>105</sub>R'<sub>105</sub>,  
 -SO<sub>2</sub>NR<sub>105</sub>R'<sub>105</sub>, -NH-CO-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NH-C(=O)-OH, -NH-  
 C(=O)-OR, -NH-C(=O)-O-phenyl, -O-C(=O)-(C<sub>1</sub>-C<sub>6</sub> alkyl),  
 30 -O-C(=O)-amino, -O-C(=O)-mono- or dialkylamino, -O-  
 C(=O)-phenyl, -O-(C<sub>1</sub>-C<sub>6</sub> alkyl)-CO<sub>2</sub>H, -NH-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub>  
 alkyl), C<sub>1</sub>-C<sub>6</sub> alkoxy or C<sub>1</sub>-C<sub>6</sub> haloalkoxy; or

R<sub>100</sub> is -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>6</sub> alkyl) or -(C<sub>1</sub>-C<sub>6</sub> alkyl)-S-(C<sub>1</sub>-C<sub>6</sub> alkyl), each of which is optionally substituted with 1, 2, or 3 R<sub>115</sub> groups, or

R<sub>100</sub> is -(C<sub>3</sub>-C<sub>8</sub> cycloalkyl) optionally substituted with 1, 2, or 3 R<sub>115</sub> groups;

R and R' independently are hydrogen; C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, or 3 groups that are independently F, Cl, Br, or I; or -(C<sub>1</sub>-C<sub>6</sub>)-R<sub>110</sub>;

W is -(CH<sub>2</sub>)<sub>0-4</sub>-, -O-, -S(O)<sub>0-2</sub>-, -N(R<sub>135</sub>)-, or -C(O)-;

R<sub>7</sub> and R<sub>7</sub>' are independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, aryl, heteroaryl, and heterocyclyl,

R<sub>105</sub> and R'<sub>105</sub> are the same or different and represent -H, -R<sub>110</sub>, -R<sub>120</sub>, -C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl, or -C<sub>1</sub>-C<sub>6</sub> alkyl chain with one double bond and one triple bond, or

-C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with -OH or -NH<sub>2</sub>; or,

-C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, or 3 groups

independently selected from halogen;

R<sub>135</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(CH<sub>2</sub>)<sub>0-2</sub>-(aryl), -(CH<sub>2</sub>)<sub>0-2</sub>-(heteroaryl), or -(CH<sub>2</sub>)<sub>0-2</sub>-(heterocyclyl),

R<sub>140</sub> is heterocyclyl optionally substituted with 1, 2, 3, or 4 groups independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and =O;

R<sub>145</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl or CF<sub>3</sub>;

R<sub>150</sub> is hydrogen, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl), C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> alkyl with one double bond and one triple bond, -R<sub>110</sub>, -R<sub>120</sub>, or

C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, 3, or 4 groups independently selected from -OH, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, R<sub>110</sub>, and halogen;

R<sub>155</sub> is C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>7</sub> cycloalkyl), C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> alkyl with one double bond and one triple bond, -R<sub>110</sub>, -R<sub>120</sub>, or

C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 1, 2, 3, or 4 groups independently selected from -OH, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, and halogen;

R<sub>180</sub> is selected from morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl, each of which is optionally substituted with 1, 2, 3, or 4 groups independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, cyano, nitro, amino, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> haloalkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, and =O; R<sub>110</sub> is aryl optionally substituted with 1 or 2 R<sub>125</sub> groups, wherein,

R<sub>125</sub> at each occurrence is independently halogen, amino, mono- or dialkylamino, -OH, -C≡N, -SO<sub>2</sub>-NH<sub>2</sub>, -SO<sub>2</sub>-NH-C<sub>1</sub>-C<sub>6</sub> alkyl, -SO<sub>2</sub>-N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO-NH<sub>2</sub>, -CO-NH-C<sub>1</sub>-C<sub>6</sub> alkyl, or -CO-N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>; or C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, -OH, -SH, -C≡N, -CF<sub>3</sub>, C<sub>1</sub>-C<sub>3</sub> alkoxy, amino, and mono- and dialkylamino; or C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with one, two or three of halogen;

$R_{120}$  is heteroaryl, which is optionally substituted with 1 or 2  $R_{125}$  groups; and

$R_{130}$  is heterocyclyl optionally substituted with 1 or 2  $R_{125}$  groups;

5 comprising

(a) reducing a ketone of formula III to generate an alcohol of formula IV; and

(b) treating the alcohol of formula IV with a base to generate an epoxide.

10


101. A process according to claim 100, further comprising contacting the epoxide with an amine of formula  $R_6NH(R_{57})$  to yield a protected amine of formula VII-1.

15

102. A process according to claim 101, further comprising forming a deprotected amine of formula VIII-1 and forming an amide using the amine of formula VIII-1 and a compound of the  $R_NCOZ$ , wherein Z is OH, Cl, or imidazolyl.

20

103. An amino alcohol of the formula:



where

R is phenyl optionally substituted with 1, 2, 3, or 4 groups independently selected from:

25

(A)  $C_1$ - $C_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $C_1$ - $C_3$  alkyl, halogen, hydroxy, thio,  $-NR_{10}R_{11}$  where  $R_{10}$  and  $R_{11}$  are independently hydrogen or  $C_1$ - $C_6$  alkyl, cyano, trifluoromethyl, and  $C_1$ - $C_3$  alkoxy,

30

(B)  $C_2$ - $C_6$  alkenyl or  $C_2$ - $C_6$  alkynyl,

(C) halogen, hydroxy, cyano,  $C_1-C_6$  alkoxy optionally substituted with 1, 2, or 3 fluoro,

(D)  $-NR_{12}R_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent:

(a)  $-H$ ,

(b)  $-C_1-C_8$  alkyl optionally substituted with one of:

(i)  $-OH$ ,

(ii)  $-NH_2$ ,

(iii) phenyl,

(c)  $-C_1-C_8$  alkyl optionally substituted with 1, 2, or 3 independently selected halogens,

(d)  $-C_3-C_8$  cycloalkyl,  $-(C_1-C_2$  alkyl) $-(C_3-C_8$  cycloalkyl),  $-(C_1-C_6$  alkyl) $-O-(C_1-C_3$  alkyl),  $-C_2-C_6$  alkenyl,  $-C_2-C_6$  alkynyl; and

(E)  $C_3-C_7$  cycloalkyl,  $-C(O)(C_1-C_4$  alkyl),  $-SO_2NR_{10}R_{11}$ ,  $-C(O)NR_{10}R_{11}$ , or  $-SO_2(C_1-C_4$  alkyl); and

$R_2$  is:

chloro, bromo, or

$-Si(R_{21})_3$  where each  $R_{21}$  is independently

$C_1-C_5$  alkyl,

$-N(R_{23})(R_{24})$  where  $R_{23}$  and  $R_{24}$  are the same or different and represent


$C_1-C_5$  alkyl,

or where  $NR_{23}R_{24}$  represents piperidinyl,

piperazinyl, or morpholinyl,

phenyl optionally substituted with 1, 2, or 3 of  $C_1-C_2$  alkyl, with the proviso that at least one of the  $R_{21}$  groups is optionally substituted phenyl.

104. A compound of the formula



where

R is phenyl optionally substituted with 1, 2, 3, or 4 groups independently selected from:

(A) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or three substituents independently selected from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, hydroxy, thio, -NR<sub>10</sub>R<sub>11</sub> where R<sub>10</sub> and R<sub>11</sub> are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, cyano, trifluoromethyl, and C<sub>1</sub>-C<sub>3</sub> alkoxy,

(B) C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl,

(C) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3 fluoro,

(D) -NR<sub>12</sub>R<sub>13</sub> where at each occurrence R<sub>12</sub> and R<sub>13</sub> are the same or different and represent:

(a) -H,

(b) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one of:

(i) -OH,

(ii) -NH<sub>2</sub>,


(iii) phenyl,

(c) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with 1, 2, or 3 independently selected halogens,

(d) -C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl; and

(E) C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NR<sub>10</sub>R<sub>11</sub>, -C(O)NR<sub>10</sub>R<sub>11</sub>, or -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl).

105. A compound of the formula



where


X<sub>1</sub> is chloro, bromo, or imidazolyl, or

X<sub>1</sub> is -CH<sub>2</sub>-R<sub>2</sub>, wherein

$R_2$  is chloro, bromo, or  
-Si( $R_{21}$ )<sub>3</sub> where each  $R_{21}$  is independently  
C<sub>1</sub>-C<sub>5</sub> alkyl,  
-N( $R_{23}$ )( $R_{24}$ ) where  $R_{23}$  and  $R_{24}$  are the same or  
5 different and represent  
C<sub>1</sub>-C<sub>5</sub> alkyl,  
or where  $NR_{23}R_{24}$  represents piperidinyl,  
piperazinyl, or morpholinyl,  
phenyl optionally substituted with 1, 2, or 3 of C<sub>1</sub>-  
10 C<sub>2</sub> alkyl, with the proviso that at least one of  
the  $R_{21}$  groups is optionally substituted phenyl;  
R is phenyl optionally substituted with 1, 2, 3, or 4 groups  
independently selected from:  
(A) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or  
15 three substituents independently selected from C<sub>1</sub>-C<sub>3</sub>  
alkyl, halogen, hydroxy, thio, -NR<sub>10</sub>R<sub>11</sub> where  $R_{10}$  and  
 $R_{11}$  are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, cyano,  
trifluoromethyl, and C<sub>1</sub>-C<sub>3</sub> alkoxy,  
(B) C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl,  
20 (C) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally  
substituted with 1, 2, or 3 fluoro,  
(D) -NR<sub>12</sub>R<sub>13</sub> where at each occurrence  $R_{12}$  and  $R_{13}$  are the  
same or different and represent:  
(a) -H,  
25 (b) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one of:  
(i) -OH,  
(ii) -NH<sub>2</sub>,  
(iii) phenyl,  
(c) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with 1, 2, or  
30 3 independently selected halogens,  
(d) -C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>8</sub>  
cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-  
C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl; and

(E) C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NR<sub>10</sub>R<sub>11</sub>,  
 -C(O)NR<sub>10</sub>R<sub>11</sub>, or -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl); and  
 PROT is a nitrogen protecting group.

5 106. A compound of the formula



wherein

R<sub>1</sub> is selected from:

(I) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one halogen;

10 (II) -CH<sub>2</sub>-CH=CH<sub>2</sub>;

(III) phenyl optionally substituted with one nitro,  
 halogen, or cyano; and

(IV) benzyl optionally substituted on phenyl with nitro,  
 halogen, or cyano

15 R is phenyl optionally substituted with 1, 2, 3, or 4 groups  
 independently selected from:

(A) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or  
 three substituents independently selected from C<sub>1</sub>-C<sub>3</sub>  
 alkyl, halogen, hydroxy, thio, -NR<sub>10</sub>R<sub>11</sub> where R<sub>10</sub> and  
 20 R<sub>11</sub> are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, cyano,  
 trifluoromethyl, and C<sub>1</sub>-C<sub>3</sub> alkoxy,

(B) C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl,

(C) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally  
 substituted with 1, 2, or 3 fluoro,

25 (D) -NR<sub>12</sub>R<sub>13</sub> where at each occurrence R<sub>12</sub> and R<sub>13</sub> are the  
 same or different and represent:

(a) -H,

(b) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one of:


(i) -OH,

30 (ii) -NH<sub>2</sub>,

(iii) phenyl,

- (c)  $-C_1-C_8$  alkyl optionally substituted with 1, 2, or 3 independently selected halogens,
- (d)  $-C_3-C_8$  cycloalkyl,  $-(C_1-C_2 \text{ alkyl})-(C_3-C_8 \text{ cycloalkyl})$ ,  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$ ,  $-C_2-C_6$  alkenyl,  $-C_2-C_6$  alkynyl; and
- (E)  $C_3-C_7$  cycloalkyl,  $-C(O)(C_1-C_4 \text{ alkyl})$ ,  $-SO_2NR_{10}R_{11}$ ,  $-C(O)NR_{10}R_{11}$ , or  $-SO_2(C_1-C_4 \text{ alkyl})$ ; and
- PROT is a nitrogen protecting group.

107. A process for preparing compounds of the formula




wherein

$R_1$  is selected from:

- (I)  $C_1-C_6$  alkyl optionally substituted with one halogen;
- (II)  $-\text{CH}_2-\text{CH}=\text{CH}_2$ ;
- (III) phenyl optionally substituted with one nitro, halogen, or cyano; and
- (IV) benzyl optionally substituted on phenyl with nitro, halogen, or cyano;

the process comprises

- (1) esterifying a protected amino acid of the formula



wherein

$R$  is phenyl optionally substituted with 1, 2, 3, or 4 groups independently selected from:

- (A)  $C_1-C_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $C_1-C_3$  alkyl, halogen, hydroxy, thio,  $-NR_{10}R_{11}$  where  $R_{10}$  and

$R_{11}$  are independently hydrogen or  $C_1-C_6$  alkyl, cyano, trifluoromethyl, and  $C_1-C_3$  alkoxy,

(B)  $C_2-C_6$  alkenyl or  $C_2-C_6$  alkynyl,

(C) halogen, hydroxy, cyano,  $C_1-C_6$  alkoxy optionally substituted with 1, 2, or 3 fluoro,

(D)  $-NR_{12}R_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent:

(a)  $-H$ ,

(b)  $-C_1-C_8$  alkyl optionally substituted with one of:

(i)  $-OH$ ,

(ii)  $-NH_2$ ,

(iii) phenyl,

(c)  $-C_1-C_8$  alkyl optionally substituted with 1, 2, or 3 independently selected halogens,


(d)  $-C_3-C_8$  cycloalkyl,  $-(C_1-C_2$  alkyl) $-(C_3-C_8$  cycloalkyl),  $-(C_1-C_6$  alkyl) $-O-(C_1-C_3$  alkyl),  $-C_2-C_6$  alkenyl,  $-C_2-C_6$  alkynyl; and

(E)  $C_3-C_7$  cycloalkyl,  $-C(O)(C_1-C_4$  alkyl),  $-SO_2NR_{10}R_{11}$ ,  $-C(O)NR_{10}R_{11}$ , or  $-SO_2(C_1-C_4$  alkyl); and

PROT is a nitrogen protecting group;

with an alkylating agent in the presence of a base.

108. A process for preparing compounds of the formula:



where  $R_1$  is an optionally substituted phenyl;

$R$  is phenyl optionally substituted with 1, 2, 3, or 4 groups independently selected from:

(A)  $C_1-C_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $C_1-C_3$  alkyl, halogen, hydroxy, thio,  $-NR_{10}R_{11}$  where  $R_{10}$  and

$R_{11}$  are independently hydrogen or  $C_1-C_6$  alkyl, cyano, trifluoromethyl, and  $C_1-C_3$  alkoxy,

(B)  $C_2-C_6$  alkenyl or  $C_2-C_6$  alkynyl,

(C) halogen, hydroxy, cyano,  $C_1-C_6$  alkoxy optionally substituted with 1, 2, or 3 fluoro,

(D)  $-NR_{12}R_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent:

(a)  $-H$ ,

(b)  $-C_1-C_8$  alkyl optionally substituted with one of:

(i)  $-OH$ ,

(ii)  $-NH_2$ ,

(iii) phenyl,


(c)  $-C_1-C_8$  alkyl optionally substituted with 1, 2, or 3 independently selected halogens,

(d)  $-C_3-C_8$  cycloalkyl,  $-(C_1-C_2 \text{ alkyl})-(C_3-C_8 \text{ cycloalkyl})$ ,  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$ ,  $-C_2-C_6$  alkenyl,  $-C_2-C_6$  alkynyl; and

(E)  $C_3-C_7$  cycloalkyl,  $-C(O)(C_1-C_4 \text{ alkyl})$ ,  $-SO_2NR_{10}R_{11}$ ,  $-C(O)NR_{10}R_{11}$ , or  $-SO_2(C_1-C_4 \text{ alkyl})$ ; and


PROT is a nitrogen protecting group;  
the process comprising:

(a) forming a mixture of an activating agent and a protected amino acid of the formula



(b) contacting the mixture of (a) with a phenol optionally substituted on the phenyl ring with nitro, halogen, or cyano.

109. A method for preparing a compound of the formula



where  $R_1$  is selected from:

(I)  $C_1$ - $C_6$  alkyl optionally substituted with one halogen;

5 (II)  $-\text{CH}_2-\text{CH}=\text{CH}_2$ ;

(III) phenyl optionally substituted with one nitro,  
halogen, or cyano; and

(IV) benzyl optionally substituted on phenyl with nitro,  
halogen, or cyano;

10 R is phenyl optionally substituted with 1, 2, 3, or 4 groups  
independently selected from:

(A)  $C_1$ - $C_6$  alkyl optionally substituted with one, two or  
three substituents independently selected from  $C_1$ - $C_3$   
alkyl, halogen, hydroxy, thio,  $-\text{NR}_{10}\text{R}_{11}$  where  $R_{10}$  and  
15  $R_{11}$  are independently hydrogen or  $C_1$ - $C_6$  alkyl, cyano,  
trifluoromethyl, and  $C_1$ - $C_3$  alkoxy,

(B)  $C_2$ - $C_6$  alkenyl or  $C_2$ - $C_6$  alkynyl,

(C) halogen, hydroxy, cyano,  $C_1$ - $C_6$  alkoxy optionally  
substituted with 1, 2, or 3 fluoro,

20 (D)  $-\text{NR}_{12}\text{R}_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the  
same or different and represent:

(a)  $-\text{H}$ ,

(b)  $-\text{C}_1$ - $\text{C}_8$  alkyl optionally substituted with one of:

(i)  $-\text{OH}$ ,

25 (ii)  $-\text{NH}_2$ ,

(iii) phenyl,

(c)  $-\text{C}_1$ - $\text{C}_8$  alkyl optionally substituted with 1, 2, or  
3 independently selected halogens,


(d)  $-\text{C}_3$ - $\text{C}_8$  cycloalkyl,  $-(\text{C}_1$ - $\text{C}_2$  alkyl)-( $\text{C}_3$ - $\text{C}_8$   
30 cycloalkyl),  $-(\text{C}_1$ - $\text{C}_6$  alkyl)-O-( $\text{C}_1$ - $\text{C}_3$  alkyl),  $-\text{C}_2$ -  
 $\text{C}_6$  alkenyl,  $-\text{C}_2$ - $\text{C}_6$  alkynyl; and

(E) C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NR<sub>10</sub>R<sub>11</sub>,  
-C(O)NR<sub>10</sub>R<sub>11</sub>, or -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl); and

PROT is a nitrogen protecting group;

the process comprising:

5 A) reducing a compound of the formula




with a reducing agent.

110. A method according to claim 109 wherein the reducing  
10 agent is NaBH<sub>4</sub>, NaCNBH<sub>3</sub>, or hydrogen and a catalyst.

111. A method according to claim 110 wherein the reducing  
agent is hydrogen and a catalyst.

15 112. A method according to claim 111 wherein the  
hydrogenation is carried out at a pressure of from 1 atmosphere  
to about 100 psi.

113. A method for preparing a compound of the formula



20 where

R is phenyl optionally substituted with 1, 2, 3, or 4 groups  
independently selected from:

25 (A) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or  
three substituents independently selected from C<sub>1</sub>-C<sub>3</sub>  
alkyl, halogen, hydroxy, thio, -NR<sub>10</sub>R<sub>11</sub> where R<sub>10</sub> and  
R<sub>11</sub> are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, cyano,  
trifluoromethyl, and C<sub>1</sub>-C<sub>3</sub> alkoxy,

- (B) C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl,  
(C) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3 fluoro,  
(D) -NR<sub>12</sub>R<sub>13</sub> where at each occurrence R<sub>12</sub> and R<sub>13</sub> are the same or different and represent:  
5 (a) -H,  
(b) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one of:  
(i) -OH,  
(ii) -NH<sub>2</sub>,  
10 (iii) phenyl,  
(c) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with 1, 2, or 3 independently selected halogens,  
(d) -C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl; and  
15 (E) C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NR<sub>10</sub>R<sub>11</sub>, -C(O)NR<sub>10</sub>R<sub>11</sub>, or -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl);

R<sub>2</sub> is:

chloro, bromo, or

20 -Si(R<sub>21</sub>)<sub>3</sub> where each R<sub>21</sub> is independently

C<sub>1</sub>-C<sub>5</sub> alkyl,

-N(R<sub>23</sub>)(R<sub>24</sub>) where R<sub>23</sub> and R<sub>24</sub> are the same or different and represent

C<sub>1</sub>-C<sub>5</sub> alkyl,

25 or where NR<sub>23</sub>R<sub>24</sub> represents piperidinyl,

piperazinyl, or morpholinyl,


phenyl optionally substituted with 1, 2, or 3 of C<sub>1</sub>-C<sub>2</sub> alkyl, with the proviso that at least one of the R<sub>21</sub> groups is optionally substituted phenyl;

30 and

PROT is a nitrogen protecting group;

which comprises:

- (a) forming a mixture of an ester of the formula



wherein

$\text{R}_1$  is selected from:

- (I)  $\text{C}_1\text{-C}_6$  alkyl optionally substituted with one halogen;
- (II)  $-\text{CH}_2-\text{CH}=\text{CH}_2$ ;
- (III) phenyl optionally substituted with one nitro, halogen, or cyano; and
- (IV) benzyl optionally substituted on phenyl with nitro, halogen, or cyano;

and  $\text{R}_2\text{CH}_2\text{X}^2$ , where


$\text{R}_2$  is as defined above, and

$\text{X}^2$  is  $-\text{Br}$  or  $-\text{I}$ ;

(b) adding a strong base having a  $\text{pK}_b$  of greater than about 30 to the mixture from (a);

(c) acidifying the mixture of (b).

114. A method for preparing a compound of the formula



where

$\text{R}$  is phenyl optionally substituted with 1, 2, 3, or 4 groups independently selected from:

- (A)  $\text{C}_1\text{-C}_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $\text{C}_1\text{-C}_3$  alkyl, halogen, hydroxy, thio,  $-\text{NR}_{10}\text{R}_{11}$  where  $\text{R}_{10}$  and  $\text{R}_{11}$  are independently hydrogen or  $\text{C}_1\text{-C}_6$  alkyl, cyano, trifluoromethyl, and  $\text{C}_1\text{-C}_3$  alkoxy,
- (B)  $\text{C}_2\text{-C}_6$  alkenyl or  $\text{C}_2\text{-C}_6$  alkynyl,

(C) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted with 1, 2, or 3 fluoro,

(D) -NR<sub>12</sub>R<sub>13</sub> where at each occurrence R<sub>12</sub> and R<sub>13</sub> are the same or different and represent:

5

(a) -H,

(b) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one of:

(i) -OH,

(ii) -NH<sub>2</sub>,

(iii) phenyl,

10

(c) -C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with 1, 2, or 3 independently selected halogens,

(d) -C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>2</sub> alkyl)-(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkyl)-O-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>2</sub>-C<sub>6</sub> alkenyl, -C<sub>2</sub>-C<sub>6</sub> alkynyl; and

15

(E) C<sub>3</sub>-C<sub>7</sub> cycloalkyl, -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NR<sub>10</sub>R<sub>11</sub>, -C(O)NR<sub>10</sub>R<sub>11</sub>, or -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl);

R<sub>2</sub> is:

chloro, bromo, or

-Si(R<sub>21</sub>)<sub>3</sub> where each R<sub>21</sub> is independently

20

C<sub>1</sub>-C<sub>5</sub> alkyl,

-N(R<sub>23</sub>)(R<sub>24</sub>) where R<sub>23</sub> and R<sub>24</sub> are the same or different and represent

C<sub>1</sub>-C<sub>5</sub> alkyl,

or where NR<sub>23</sub>R<sub>24</sub> represents piperidinyl,

25

piperazinyl, or morpholinyl,

phenyl optionally substituted with 1, 2, or 3 of C<sub>1</sub>-C<sub>2</sub> alkyl, with the proviso that at least one of the R<sub>21</sub> groups is optionally substituted phenyl;

and

30


PROT is a nitrogen protecting group;

which comprises:

(a) forming a mixture of an acid R<sub>2</sub>-CH<sub>2</sub>-COOH and a base,

(b) treating the mixture of (a) with an ester of the

formula



wherein R, and PROT are defined above, and

R<sub>1</sub> is selected from:

(I) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one  
5 halogen;

(II) -CH<sub>2</sub>-CH=CH<sub>2</sub>;


(III) phenyl optionally substituted with one nitro,  
halogen, or cyano; and

(IV) benzyl optionally substituted on phenyl with  
10 nitro, halogen, or cyano;

(c) acidifying the mixture from (b).

115. A method according to claim 114 wherein, the base has  
a pK<sub>b</sub> 30 or greater.

116. A method for preparing a compound of the formula



wherein

R is phenyl optionally substituted with 1, 2, 3, or 4 groups  
20 independently selected from:

(A) C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one, two or  
three substituents independently selected from C<sub>1</sub>-C<sub>3</sub>  
alkyl, halogen, hydroxy, thio, -NR<sub>10</sub>R<sub>11</sub> where R<sub>10</sub> and  
R<sub>11</sub> are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, cyano,  
25 trifluoromethyl, and C<sub>1</sub>-C<sub>3</sub> alkoxy,

(B) C<sub>2</sub>-C<sub>6</sub> alkenyl or C<sub>2</sub>-C<sub>6</sub> alkynyl,

(C) halogen, hydroxy, cyano, C<sub>1</sub>-C<sub>6</sub> alkoxy optionally  
substituted with 1, 2, or 3 fluoro,

(D)  $-NR_{12}R_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent:

(a)  $-H$ ,

(b)  $-C_1-C_8$  alkyl optionally substituted with one of:

(i)  $-OH$ ,

(ii)  $-NH_2$ ,

(iii) phenyl,

(c)  $-C_1-C_8$  alkyl optionally substituted with 1, 2, or 3 independently selected halogens,

(d)  $-C_3-C_8$  cycloalkyl,  $-(C_1-C_2 \text{ alkyl})-(C_3-C_8 \text{ cycloalkyl})$ ,  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$ ,  $-C_2-C_6$  alkenyl,  $-C_2-C_6$  alkynyl; and


(E)  $C_3-C_7$  cycloalkyl,  $-C(O)(C_1-C_4 \text{ alkyl})$ ,  $-SO_2NR_{10}R_{11}$ ,  $-C(O)NR_{10}R_{11}$ , or  $-SO_2(C_1-C_4 \text{ alkyl})$ ;

$R_2$  is Cl or Br;

and

PROT is a nitrogen protecting group;

which comprises contacting a compound of the formula




wherein

$X_1$  is chloro, bromo, or imidazolyl;

with  $\text{LiCH}_2\text{Cl}$  or  $\text{LiCH}_2\text{Br}$ .

117. A method for preparing a compound of the formula



wherein

$X_1$  is chloro, bromo, or imidazolyl;


$R$  is phenyl optionally substituted with 1, 2, 3, or 4 groups independently selected from:

- (A)  $C_1-C_6$  alkyl optionally substituted with one, two or three substituents independently selected from  $C_1-C_3$  alkyl, halogen, hydroxy, thio,  $-NR_{10}R_{11}$  where  $R_{10}$  and  $R_{11}$  are independently hydrogen or  $C_1-C_6$  alkyl, cyano, trifluoromethyl, and  $C_1-C_3$  alkoxy,
- (B)  $C_2-C_6$  alkenyl or  $C_2-C_6$  alkynyl,
- (C) halogen, hydroxy, cyano,  $C_1-C_6$  alkoxy optionally substituted with 1, 2, or 3 fluoro,
- (D)  $-NR_{12}R_{13}$  where at each occurrence  $R_{12}$  and  $R_{13}$  are the same or different and represent:
- (a)  $-H$ ,
- (b)  $-C_1-C_8$  alkyl optionally substituted with one of:
- (i)  $-OH$ ,
- (ii)  $-NH_2$ ,
- (iii) phenyl,
- (c)  $-C_1-C_8$  alkyl optionally substituted with 1, 2, or 3 independently selected halogens,
- (d)  $-C_3-C_8$  cycloalkyl,  $-(C_1-C_2 \text{ alkyl})-(C_3-C_8 \text{ cycloalkyl})$ ,  $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$ ,  $-C_2-C_6$  alkenyl,  $-C_2-C_6$  alkynyl; and
- (E)  $C_3-C_7$  cycloalkyl,  $-C(O)(C_1-C_4 \text{ alkyl})$ ,  $-SO_2NR_{10}R_{11}$ ,  $-C(O)NR_{10}R_{11}$ , or  $-SO_2(C_1-C_4 \text{ alkyl})$ ; and

PROT is a nitrogen protecting group;

which comprises

contacting a compound of the formula



with thionyl chloride, phosphorous trichloride, oxalyl chloride, phosphorous tribromide, triphenylphosphorous dibromide, oxalyl bromide, 1,2-phenylenetrichlorophosphate, 2,4,6-trichloro-1,3,5-triazine or carbonyldiimidazole.