

DESCRIPTION

Title of Invention: COMPOSITION FOR ENHANCING COGNITIVE
FUNCTION, COMPOSITION FOR REMEDYING ANXIETY SYMPTOMS, AND
COMPOSITION FOR SUPPRESSING CEREBRAL ATROPHY

5 [Technical Field]
[0001]

The present invention relates to a composition for
improving cognitive function, a composition for improving
anxiety-like symptoms, and a composition for suppressing
10 cerebral atrophy.

[Background Art]
[0002]

With the recent rapid increase in the elderly population,
the number of patients with dementia increased rapidly, and it
15 has been reported that the population with dementia reached
8.4% of the population of those aged 65 or over in 2015.

According to the Ministry of Health, Labour and Welfare
of Japan, dementia is defined to mean "a state in which various
mental functions that once developed normally after the birth
20 have chronically declined and disappeared to the extent that
daily life and social life cannot be managed".

Dementia includes dementia caused by various diseases
such as Alzheimer-type dementia, frontotemporal dementia (Pick
disease etc.), Lewy body dementia, and cerebrovascular dementia.
25 While aging is the greatest risk factor, its cause is often
unclear.

However, in any type of dementia, cognitive dysfunction
such as memory disorder, disorientation and the like are
observed as core symptoms, and behavioral and psychological
30 symptoms such as behavior abnormality, mental symptoms and the
like are also commonly observed, and progression of the above-
mentioned symptoms has serious effects such as increased burden
of nursing care and the like for not only patients but also
their families.

35 [0003]

Currently, acetylcholinesterase inhibitors such as donepezil hydrochloride, and NMDA (N-methyl-D-aspartate) receptor antagonists such as memantine have been approved as therapeutic drugs for dementia. However, these therapeutic
5 drugs for dementia are basically for Alzheimer-type dementia, are symptomatic treatment drugs, and can merely suppress progression of the symptoms somewhat.
[0004]

Regarding the prevention of dementia, the importance of
10 meal and exercise in preventing and improving a decline in cognitive function is known.

Regarding meals, the effectiveness of antioxidants such as vitamin C, vitamin E, β -carotene and ω -3 long-chain unsaturated fatty acids has been reported, and the effects of
15 ingestion of ω -3 long-chain unsaturated fatty acid, melatonin and tryptophan on mild cognitive impairment of the elderly people have been studied (non-patent document 1). Furthermore, there are found a report stating that lysine improves cognitive function of healthy humans (patent document 1), reports
20 suggesting the effect of L-arginine and lysine on Alzheimer-type dementia and anxiety-like symptoms (non-patent documents 2 to 5), and a report suggesting improvement of a decline in working memory and a decline in cognitive function by tyrosine (non-patent document 6).

25 Regarding exercise, it has been found that an increase in cerebral blood flow due to exercise can improve physical activity and prevent Alzheimer's disease.

However, some of the above-mentioned ingredients ingested in meals do not have sufficient preventive or improving effect
30 on a decline in cognitive function, and many require future verification of the effectiveness.

Furthermore, some middle-aged and elderly people who have an increased risk of a decline in cognitive function often have difficulty in exercising due to disease or the like, or may
35 have difficulty in continuing exercise due to a decline in

physical function.

[0005]

Therefore, it is difficult to say that a component having an effective preventive or improving effect on the decline in
5 cognitive function, which is highly safe and can be ingested continuously, has been found.

[0006]

It is known that the proportion of anxiety-like symptoms such as spiritless, apathy, anxiety, depression and the like
10 increases in elderly people, and such anxiety-like symptoms are seen as a risk factor for dementia before the onset of dementia.

Furthermore, it is also known that the above-mentioned anxiety-like symptoms are seen as behavioral and psychological symptoms of dementia.

15 [Document List]

[patent document]

[0007]

patent document 1: EP2367547(A1)

[non-patent documents]

20 [0008]

non-patent document 1: Anglo Food Industry Hi-Tech 22 (4) 23-24 (2011)

non-patent document 2: The American Journal of Medicine 108 (5) 439 (2000 Apr 1)

25 non-patent document 3: Neuropsychiatric Disease and Treatment 6 707-710 (2010)

non-patent document 4: Biomedical Research 28 (2) 85-90 (2007)

non-patent document 5: The Journal of Nutrition 132 (12) 3744-3746 (2002)

30 non-patent document 6: Amino Acids 45 (5) 1035-1045 (2013)

[Summary of Invention]

[Problems to be Solved by the Invention]

[0009]

Accordingly, the present invention aims to provide a
35 composition for improving cognitive function, which has an

effective preventive or improving effect on a decline in cognitive function, is highly safe, and enables continuous ingestion or administration.

The present invention aims to provide a composition for
5 improving anxiety-like symptoms mentioned above.

[Means of Solving the Problems]

[0010]

The present inventors have conducted intensive studies in an attempt to solve the above-mentioned problems and found that
10 a composition containing one or more kinds selected from the group consisting of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan has an improving effect of cognitive function
15 and an improving effect of anxiety-like symptoms, and that the aforementioned composition has an effect of suppressing cerebral atrophy characteristically observed in neurodegenerative diseases such as Alzheimer-type dementia and the like, which resulted in the completion of the present
20 invention.

[0011]

That is, the present invention relates to the following.

[1] A composition for improving cognitive function comprising one or more kinds selected from the group consisting of 25 mol%
25 - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

[2] The composition of [1], further comprising 10 mol% - 30 mol% of lysine relative to the total content of leucine, lysine,
30 valine, isoleucine, phenylalanine, histidine and tryptophan.

[3] The composition of [1] or [2], further comprising one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan.

[4] The composition of [3], comprising one or more kinds
35 selected from the group consisting of 2 mol% - 7 mol% of valine,

5 mol% - 15 mol% of isoleucine, 2 mol% - 10 mol% of histidine and 0.1 mol% - 2 mol% of tryptophan relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

5 [5] The composition of [3] or [4], comprising the following amino acids at the following contents relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan:

leucine 25 mol% - 45 mol%,
10 phenylalanine 20 mol% - 40 mol%,
lysine 10 mol% - 30 mol%,
valine 2 mol% - 7 mol%,
isoleucine 5 mol% - 15 mol%,
histidine 2 mol% - 10 mol%, and
15 tryptophan 0.1 mol% - 2 mol%.

[6] The composition of any of [1] to [5], wherein the composition is a pharmaceutical product.

[7] The composition of any of [1] to [5], wherein the composition is a food.

20 [8] A method for improving cognitive function of a subject animal in need of improving cognitive function, comprising allowing the animal to ingest a composition comprising one or more kinds selected from the group consisting of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine
25 relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan, in an amount effective for improving cognitive function of the subject animal, or administering said amount of the composition to the subject animal.

30 [9] The method of [8], wherein the composition further comprising 10 mol% - 30 mol% of lysine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

[10] The method of [8] or [9], wherein the composition further
35 comprising one or more kinds selected from the group consisting

of valine, isoleucine, histidine and tryptophan.

[11] The method of [10], wherein the composition comprising one or more kinds selected from the group consisting of 2 mol% - 7 mol% of valine, 5 mol% - 15 mol% of isoleucine, 2 mol% - 10 mol% of histidine and 0.1 mol% - 2 mol% of tryptophan relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

[12] The method of [10] or [11], wherein the composition comprising the following amino acids at the following contents relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan:

leucine 25 mol% - 45 mol%,
phenylalanine 20 mol% - 40 mol%,
lysine 10 mol% - 30 mol%,
valine 2 mol% - 7 mol%,
isoleucine 5 mol% - 15 mol%,
histidine 2 mol% - 10 mol%, and
tryptophan 0.1 mol% - 2 mol%.

[13] A composition for improving an anxiety-like symptom, comprising one or more kinds selected from the group consisting of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

[14] The composition of [13], further comprising 10 mol% - 30 mol% of lysine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

[15] The composition of [13] or [14], further comprising one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan.

[16] The composition of [15], comprising one or more kinds selected from the group consisting of 2 mol% - 7 mol% of valine, 5 mol% - 15 mol% of isoleucine, 2 mol% - 10 mol% of histidine and 0.1 mol% - 2 mol% of tryptophan relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

[17] The composition of [15] or [16], comprising the following amino acids at the following contents relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan:

5 leucine 25 mol% - 45 mol%,
 phenylalanine 20 mol% - 40 mol%,
 lysine 10 mol% - 30 mol%,
 valine 2 mol% - 7 mol%,
 isoleucine 5 mol% - 15 mol%,
10 histidine 2 mol% - 10 mol%, and
 tryptophan 0.1 mol% - 2 mol%.

[18] The composition of any of [13] to [17], wherein the composition is a pharmaceutical product.

[19] The composition of any of [13] to [17], wherein the
15 composition is a food.

[20] A method for improving an anxiety-like symptom of a subject animal in need of improving an anxiety-like symptom, comprising allowing the animal to ingest a composition comprising one or more kinds selected from the group consisting
20 of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan, in an amount effective for improving the anxiety-like symptom of the subject animal, or administering said amount of the
25 composition to the subject animal.

[21] The method of [20], wherein the composition further comprising 10 mol% - 30 mol% of lysine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

30 [22] The method of [20] or [21], wherein the composition further comprising one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan.

[23] The method of [22], wherein the composition comprising one or more kinds selected from the group consisting of 2 mol% - 7
35 mol% of valine, 5 mol% - 15 mol% of isoleucine, 2 mol% - 10

mol% of histidine and 0.1 mol% - 2 mol% of tryptophan relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

[24] The method of [22] or [23], wherein the composition
5 comprising the following amino acids at the following contents relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan:

leucine 25 mol% - 45 mol%,
phenylalanine 20 mol% - 40 mol%,
10 lysine 10 mol% - 30 mol%,
valine 2 mol% - 7 mol%,
isoleucine 5 mol% - 15 mol%,
histidine 2 mol% - 10 mol%, and
tryptophan 0.1 mol% - 2 mol%.

15 [25] A composition for suppressing cerebral atrophy, comprising one or more kinds selected from the group consisting of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

20 [26] The composition of [25], further comprising 10 mol% - 30 mol% of lysine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

[27] The composition of [25] or [26], further comprising one or more kinds selected from the group consisting of valine,
25 isoleucine, histidine and tryptophan.

[28] The composition of [27], comprising one or more kinds selected from the group consisting of 2 mol% - 7 mol% of valine, 5 mol% - 15 mol% of isoleucine, 2 mol% - 10 mol% of histidine and 0.1 mol% - 2 mol% of tryptophan relative to the total
30 content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

[29] The composition of [27] or [28], comprising the following amino acids at the following contents relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine,
35 histidine and tryptophan:

leucine 25 mol% - 45 mol%,
phenylalanine 20 mol% - 40 mol%,
lysine 10 mol% - 30 mol%,
valine 2 mol% - 7 mol%,
5 isoleucine 5 mol% - 15 mol%,
histidine 2 mol% - 10 mol%, and
tryptophan 0.1 mol% - 2 mol%.

[30] The composition of any of [25] to [29], wherein the composition is a pharmaceutical product.

10 [31] The composition of any of [25] to [29], wherein the composition is a food.

[32] A method for suppressing cerebral atrophy of a subject animal in need of suppressing cerebral atrophy, comprising allowing the animal to ingest a composition comprising one or
15 more kinds selected from the group consisting of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan, in an amount effective for suppressing cerebral atrophy of the
20 subject animal, or administering said amount of the composition to the subject animal.

[33] The method of [32], wherein the composition further comprising 10 mol% - 30 mol% of lysine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine,
25 histidine and tryptophan.

[34] The method of [32] or [33], wherein the composition further comprising one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan.

[35] The method of [34], wherein the composition comprising one
30 or more kinds selected from the group consisting of 2 mol% - 7 mol% of valine, 5 mol% - 15 mol% of isoleucine, 2 mol% - 10 mol% of histidine and 0.1 mol% - 2 mol% of tryptophan relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.

35 [36] The method of [34] or [35], wherein the composition

comprising the following amino acids at the following contents relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan:

leucine 25 mol% - 45 mol%,
5 phenylalanine 20 mol% - 40 mol%,
lysine 10 mol% - 30 mol%,
valine 2 mol% - 7 mol%,
isoleucine 5 mol% - 15 mol%,
histidine 2 mol% - 10 mol%, and
10 tryptophan 0.1 mol% - 2 mol%.

[Effects of the Invention]

[0012]

The present invention can provide a composition for improving cognitive function that can effectively prevent or
15 improve a decline in cognitive function.

Also, the present invention can provide a composition for improving anxiety-like symptoms. The composition for improving anxiety-like symptoms of the present invention is particularly effective for the improvement of anxiety-like symptoms that
20 appear along with a decline in cognitive function or before and after a decline in cognitive function.

Furthermore, the present invention can provide a composition for suppressing cerebral atrophy. The composition for suppressing cerebral atrophy of the present invention
25 suppresses cerebral atrophy observed in neurodegenerative diseases such as Alzheimer-type dementia and the like, and physiological cerebral atrophy that appears and progresses with aging, and can improve a decline in cognitive function caused by cerebral atrophy.

30 Particularly, the composition for improving cognitive function, the composition for improving anxiety-like symptoms, and the composition for suppressing cerebral atrophy of the present invention are highly safe and suitable for continuous ingestion or administration.

35 [Brief Description of Drawings]

[0013]

Fig. 1 shows the effects of the composition of Example 1 of the present invention on transfer latency time and achievement by 1st ES in the passive avoidance test in Experimental Example 1. In the Figure, "N.S." indicates no significant difference at critical rate (p)=5%.

Fig. 2 shows the effects of the composition of Example 1 of the present invention on transfer latency time and achievement by 1st ES in the passive avoidance test (reproducibility test) in Experimental Example 2. In the Figure, "n.s." indicates no significant difference at critical rate (p)=5%.

Fig. 3 shows the effects of the composition of Example 1 of the present invention on the number of entries in open arms, which is an index of anxiety and agitation symptoms, and the total distance moved, which is an index of hyperactivity, in the elevated plus maze test in Experimental Example 3. In the Figure, "N.S." indicates no significant difference at critical rate (p)=5%.

Fig. 4 shows the effects of the composition of Example 1 of the present invention on the number of entries in open arms, which is an index of anxiety and agitation symptoms, in the elevated plus maze test (reproducibility test) in Experimental Example 4.

Fig. 5 shows the effects of the composition of Example 1 of the present invention on the intracerebral neurotransmitter concentration in Experimental Example 5. In the Figure, "n.s." indicates no significant difference at critical rate (p)=5%.

Fig. 6 shows the effects of the composition of Example 1 of the present invention on the expression of brain-derived neurotrophic factor (BDNF) gene in Experimental Example 6. In the Figure, "n.s." indicates no significant difference at critical rate (p)=0.05.

Fig. 7 shows the effects of the composition of Example 1 of the present invention on the volume of cerebral cortex in

Experimental Example 7. In the Figure, "n.s." indicates no significant difference at critical rate ($p=0.05$), "*" indicates presence of significant difference at $p<0.05$, and "***" indicates presence of significant difference at $p<0.01$.

5 Fig. 8 shows the effects of the ingestion of the composition of Example 2 on the time to complete the task in the TMT-B test in Experimental Example 8. In the Figure, error bar shows 95% confidence interval.

[Description of Embodiments]

10 [0014]

The composition for improving cognitive function of the present invention (hereinafter to be also referred to as "the composition of the present invention" in the present specification) contains one or more kinds selected from the
15 group consisting of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan (hereinafter to be also referred to as "seven amino acids" in the present specification), that is, one or
20 both of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of seven amino acids.

[0015]

As used herein, the "cognitive function" in the present
25 specification refers to higher-order functions of the brain, such as memory, judgment, calculation, comprehension, learning, thinking, language, performance, attention, orientation, execution, delayed recall and the like, and "improvement of cognitive function" refers to prevention or improvement of a
30 decline in such cognitive function.

The decline in the cognitive function that may be improved by the composition of the present invention includes a decline in cognitive function due to Alzheimer-type dementia, and dementia caused by various diseases and lesions such as
35 frontotemporal lobar degeneration (Pick disease etc.), Lewy

body disease, cerebrovascular diseases and the like, a decline in cognitive function due to aging, a decline in cognitive function observed in healthy humans, for example, memory decline, reduction in attention, a decline in thinking skills
5 and the like.

[0016]

The "leucine" and "phenylalanine" to be used may be any of an L form, a D form and a DL form. An L form and a DL form are preferably used, and an L form is more preferably used.

10 [0017]

The "leucine" and "phenylalanine" can be used not only in a free form but also a salt form. The term "leucine" and "phenylalanine" in the present specification are concepts each encompassing even a salt. The salt form is not particularly
15 limited as long as it is a pharmacologically acceptable salt, and acid addition salt, salt with base and the like can be mentioned.

Concrete examples include salts with inorganic bases, salts with organic bases, salts with inorganic acids, salts
20 with organic acids, salts with amino acid and the like.

[0018]

Examples of the salts with inorganic bases include salts with alkali metals such as lithium, sodium, potassium and the like, salts with alkaline earth metals such as magnesium,
25 calcium and the like, ammonium salt and the like.

Examples of the salts with organic bases include salts with alkanolamine such as monoethanolamine, diethanolamine, triethanolamine and the like, salts with heterocyclic amine such as morpholine, piperidine and the like, and the like.

30 Examples of the salts with inorganic acids include salts with hydrohalic acid (hydrochloric acid, hydrobromic acid, hydroiodic acid etc.), sulfuric acid, nitric acid, phosphoric acid and the like.

Examples of the salts with organic acids include salts
35 with monocarboxylic acid such as formic acid, acetic acid,

propanoic acid and the like; salts with saturated dicarboxylic acid such as oxalic acid, malonic acid, malic acid, succinic acid and the like; salts with unsaturated dicarboxylic acid such as maleic acid, fumaric acid and the like; salts with
5 tricarboxylic acid such as citric acid and the like; salts with keto acid such as α -ketoglutaric acid and the like.

Examples of the salts with amino acid include salts with aliphatic amino acid such as glycine, alanine and the like; salts with aromatic amino acid such as tyrosine and the like;
10 salts with basic amino acid such as arginine and the like; salts with acidic amino acid such as aspartic acid, glutamic acid and the like; salts with amino acid forming lactam such as pyroglutamic acid and the like.

[0019]

15 Each of the above-mentioned salts may be a hydrate (salt hydrate), and examples of the hydrate include 1 hydrate to 6 hydrate and the like.

[0020]

In the present invention, one kind each of "leucine" and
20 "phenylalanine" in the above-mentioned free form or salt form may be used singly, or two or more kinds thereof may be used in combination.

For the object of the present invention, a free form, hydrochloride and the like of each of "leucine" and
25 "phenylalanine" are preferable.

[0021]

In the composition of the present invention, the content of leucine relative to the total content of seven amino acids is 25 mol% - 45 mol%, preferably 30 mol% - 40 mol%, more
30 preferably 35 mol% - 40 mol%.

In addition, the content of phenylalanine is 20 mol% - 40 mol%, preferably 20 mol% - 35 mol%, more preferably 25 mol% - 30 mol%.

[0022]

35 From the aspect of the cognitive function improving

effect, the composition of the present invention preferably contains 25 mol% - 45 mol% of leucine and 20 mol% - 40 mol% of phenylalanine.

[0023]

5 The composition of the present invention may contain 10 mol% - 30 mol% of lysine relative to the total content of seven amino acids in addition to one or more kinds selected from the group consisting of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of seven
10 amino acids.

[0024]

 The "lysine" may be any of an L form, a D form and a DL form, as in the case of the above-mentioned leucine etc. An L form and a DL form are preferably used, and an L form is more
15 preferably used.

 The "lysine" can be used not only in a free form but also a salt form, and it is a concept encompassing even a salt. Examples of the salt form include salts with inorganic bases, salts with organic bases, salts with inorganic acids, salts
20 with organic acids, salts with amino acid and the like as mentioned above.

 For the purpose of the present invention, a free form, hydrochloride or the like of "lysine" is preferable.

[0025]

25 In the composition of the present invention, the content of lysine relative to the total content of seven amino acids is 10 mol% - 30 mol%, preferably 15 mol% - 25 mol%, more preferably 15 mol% - 20 mol%.

[0026]

30 From the aspect of the cognitive function improving effect, the composition of the present invention more preferably contains 25 mol% - 45 mol% of leucine, 20 mol% - 40 mol% of phenylalanine, and 10 mol% - 30 mol% of lysine relative to the total content of seven amino acids.

35 [0027]

The composition of the present invention may further contain one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan, in addition to one or more kinds selected from the group consisting of leucine and phenylalanine at the above-mentioned molar composition ratio, or one or more kinds selected from the group consisting of leucine and phenylalanine, and lysine at the above-mentioned molar composition ratio.

[0028]

The "valine", "isoleucine", "histidine" and "tryptophan" may be any of an L form, a D form and a DL form, as in the case of the above-mentioned leucine etc. An L form and a DL form are preferably used, and an L form is more preferably used.

Each of "valine", "isoleucine", "histidine" and "tryptophan" can be used not only in a free form but also a salt form, and each is a concept encompassing even a salt. Examples of the salt form include salts with inorganic bases, salts with organic bases, salts with inorganic acids, salts with organic acids, salts with amino acid and the like as mentioned above.

For the purpose of the present invention, a free form, hydrochloride or the like of each of "valine", "isoleucine", "histidine" and "tryptophan" is preferable.

[0029]

The composition of the present invention preferably contains 2 mol% - 7 mol%, more preferably 4 mol% - 6 mol%, of valine relative to the total content of seven amino acids.

Isoleucine is preferably contained at 5 mol% - 15 mol%, more preferably 7 mol% - 13 mol%, relative to the total content of seven amino acids.

Histidine is preferably contained at 2 mol% - 10 mol%, more preferably 2.5 mol% - 8 mol%, relative to the total content of seven amino acids.

Tryptophan is preferably contained at 0.1 mol% - 2 mol%, more preferably 0.3 mol% - 1 mol%, relative to the total

content of seven amino acids.

[0030]

From the aspect of the cognitive function improving effect, the composition of the present invention more preferably contains 25 mol% - 45 mol% of leucine, 20 mol% - 40 mol% of phenylalanine, and 10 mol% - 30 mol% of lysine, as well as 2 mol% - 7 mol% of valine, 5 mol% - 15 mol% of isoleucine, 2 mol% - 10 mol% of histidine and 0.1 mol% - 2 mol% of tryptophan, relative to the total content of seven amino acids.

10 [0031]

In the present invention, the above-mentioned each amino acid in a free form or salt form to be used may be extracted from animals, plants or the like, which are naturally present, and purified, or obtained by a chemical synthesis method, a fermentation method, an enzyme method or a gene recombinant method and the like. Commercially available products provided by each company may also be utilized.

[0032]

The composition of the present invention may further contain, in addition to the above-mentioned amino acids, other nutrition components such as carbohydrates, lipid, protein, essential amino acid (methionine, threonine) other than the above-mentioned amino acid, non-essential amino acid, vitamin, mineral and the like.

25 [0033]

The composition of the present invention can be formulated into various forms such as liquids (e.g., solution, suspension, emulsion and the like); semi-solid (e.g., gel, cream and the like); solid (e.g., powder, granule, tablet, capsule and the like), and the like by adding 10 mol% - 30 mol% of lysine, or further, one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan, other nutrition components, pharmaceutically acceptable additives, and the like to one or more kinds selected from the group consisting of 25 mol% - 45 mol% of

leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of seven amino acids, as necessary and according to a formulating means well known in the field of preparations, for example, the methods described in the Japanese

5 Pharmacopoeia XVII General Rules for preparations [3]
Monographs for Preparations and the like.
[0034]

The above-mentioned pharmaceutically acceptable additive can be appropriately selected according to the form of the
10 composition of the present invention and, for example, excipient, binder, disintegrant, lubricant, coating agent, base, solvent, solubilizing agents, solubilizer, emulsifier, dispersing agent, suspending agent, stabilizer, thickener, soothing agent, isotonicity agent, pH adjuster, antioxidant,
15 antiseptic, preservative, corrigent, sweetener, flavor, colorant and the like can be mentioned.
[0035]

To be specific, examples of the excipient include magnesium carbonate, saccharides (glucose, lactose, cornstarch
20 etc.), sugar alcohol (sorbitol, mannitol etc.) and the like.

Examples of the binder include gelatin, pregelatinized starch, partly pregelatinized starch, cellulose and a derivative thereof (crystalline cellulose, hydroxypropylcellulose etc.) and the like.

25 Examples of the disintegrant include crospovidone, povidone, crystalline cellulose and the like.

Examples of the lubricant include talc, magnesium stearate and the like.

Examples of the coating agent include methacrylic acid-
30 methyl methacrylate copolymer, methacrylic acid-ethyl acrylate copolymer, methyl methacrylate-butyl methacrylate-dimethylaminoethyl methacrylate copolymer, ethyl acrylate-methyl methacrylate-trimethylammonioethylmethacrylate chloride copolymer and the like.

35 [0036]

Examples of the base include animal and plant fats and oils (olive oil, cacao butter, beef tallow, sesame oil, hydrogenated oil, castor oil etc.), wax (Carnauba wax, beeswax etc.), polyethylene glycol and the like.

5 Examples of the solvent include purified water, water for injection, monovalent alcohol (ethanol etc.), polyhydric alcohol (glycerol etc.) and the like.

Examples of the solubilizing agent include propylene glycol, medium-chain triglyceride and the like.

10 [0037]

Examples of the solubilizer, emulsifier, dispersing agent and suspending agent include surfactant and the like such as sorbitan fatty acid ester, glycerol fatty acid ester, polyoxyethylene sorbitan fatty acid ester (polysorbate 20, polysorbate 80 etc.), polyoxyethylene hydrogenated castor oil, sucrose fatty acid ester and the like.

[0038]

Examples of the stabilizer include adipic acid, β -cyclodextrin, ethylenediamine, sodium edetate and the like.

20 Examples of the thickener include water-soluble polymer (sodium polyacrylate, carboxyvinyl polymer etc.), polysaccharides (sodium alginate, xanthan gum, tragacanth etc.) and the like.

Examples of the soothing agent include ethyl
25 aminobenzoate, chlorobutanol, propylene glycol, benzyl alcohol and the like.

Examples of the isotonicity agent include potassium chloride, sodium chloride, sorbitol, saline and the like.

Examples of the pH adjuster include hydrochloric acid, sulfuric acid, acetic acid, citric acid, lactic acid, sodium
30 hydroxide, potassium hydroxide and the like.

[0039]

Examples of the antioxidant include dibutylhydroxytoluene (BHT), butylhydroxyanisole (BHA), dl- α -tocopherol, erythorbic
35 acid and the like.

Examples of the antiseptic and preservative include paraben (methylparaben etc.), benzyl alcohol, sodium dehydroacetate, sorbic acid and the like.

[0040]

5 Examples of the corrigent include ascorbic acid, erythritol, sodium L-glutamate and the like.

Examples of the sweetener include aspartame, licorice extract, saccharin and the like.

10 Examples of the flavor include l-menthol, d-camphor, vanillin and the like.

Examples of the colorant include tar pigment (Food Color Red No. 2, Food Color Blue No. 1, Food Color yellow No. 4 etc.), inorganic pigment (red iron oxide, yellow iron oxide, black iron oxide etc.), natural dye (turmeric extract, β -carotene, sodium copper-chlorophyllin etc.) and the like.

[0041]

In the present invention, one or more kinds of the above-mentioned additives can be used.

[0042]

20 The content of "one or more kinds selected from the group consisting of leucine and phenylalanine", "one or more kinds selected from the group consisting of leucine and phenylalanine, and lysine", "one or more kinds selected from the group consisting of leucine and phenylalanine, and one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan", or "one or more kinds selected from the group consisting of leucine and phenylalanine, lysine, and one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan" in the composition of the present invention is generally 30 wt% - 100 wt%, preferably 50 wt% - 100 wt%, relative to the total amount of the composition.

35 The above-mentioned content of "one or more kinds selected from the group consisting of leucine and phenylalanine" and the like is calculated as a content of the amino acid converted to a free form when any amino acid is

contained in the form of a salt.

[0043]

The composition of the present invention has an effect of improving anxiety-like symptoms in addition to the cognitive
5 function improving effect, and can also be provided as a composition for improving such anxiety-like symptoms.

[0044]

As used herein, the "anxiety-like symptoms" refers to symptoms such as spiritless, apathy, depressive symptoms
10 (anxiety, depression etc.), restlessness, sleep disorder due to anxiety, excitatory symptoms (agitation, attention deficit, hyperactivity, abnormal behavior etc.) and the like.

The "improvement of anxiety-like symptoms" refers to preventing the expression of the above-mentioned anxiety-like
15 symptoms or reducing the symptoms.

The composition of the present invention is effective for the improvement of the above-mentioned anxiety-like symptoms that appear along with a decline in cognitive function or before and after a decline in cognitive function, namely,
20 anxiety-like symptoms that appear before the onset of dementia and are known as a risk factor for dementia, and anxiety-like symptoms observed as behavioral and psychological symptoms of dementia.

[0045]

25 In addition, the composition of the present invention has an effect of suppressing cerebral atrophy, and can also be provided as a composition for suppressing cerebral atrophy.

[0046]

As used herein, "cerebral atrophy" refers to cerebral
30 atrophy characteristically observed in patients with neurodegenerative diseases such as Alzheimer-type dementia, frontotemporal dementia and the like, and physiological cerebral atrophy observed with aging.

The "suppression of cerebral atrophy" refers to
35 suppression of cerebral atrophy that progresses due to the

above-mentioned neurodegenerative diseases and the physiological cerebral atrophy observed with aging.

The composition of the present invention can suppress cerebral atrophy caused by the loss of nerve cells in the
5 above-mentioned neurodegenerative diseases, and can improve a decline in cognitive function caused by such cerebral atrophy. In addition, the composition can also suppress physiological cerebral atrophy that appears and progresses along with aging, and can improve a decline in cognitive function due to such
10 physiological cerebral atrophy, for example, memory decline, reduction in attention, a decline in thinking skills and the like observed in healthy humans along with aging.
[0047]

The daily ingestion amount or dose of the composition of
15 the present invention is appropriately determined according to the sex, age of the subject to be applied to (hereinafter to be also referred to as the "application subject" in the present specification), conditions such as a decline in cognitive function, and intracerebral lesion and the like observed in the
20 application subject or the state of anxiety-like symptoms and their degree, the form of the composition of the present invention, the administration method and the like. When application subject is a human adult, it is generally 0.5 g - 22 g, preferably 1 g - 20 g, more preferably 2.5 g - 16 g,
25 further preferably 3 g - 16 g, still more preferably 5 g - 16 g, as the total amount of one or more kinds selected from the group consisting of leucine and phenylalanine (i.e., either or both of leucine and phenylalanine) (the total amount converted to the amount of free form), the total amount of either or both
30 of leucine and phenylalanine, and lysine (the total amount converted to the amount of free form), the total amount of either or both of leucine and phenylalanine, and one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan (the total amount converted to the
35 amount of free form), or the total amount of either or both of

leucine and phenylalanine, lysine, and one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan (the total amount converted to the amount of free form).

5 The above-mentioned amount can be ingested or administered at once or in several portions (e.g., 2 - 4 portions) per day.

[0048]

From the aspect of effectively increasing the blood
10 concentration of the application subject to enhance the cognitive function improving effect, a single ingestion amount or dose of the composition when the application subject is a human adult is generally 0.3 g - 10 g, preferably 0.5 g - 10 g, more preferably 1 g - 8 g, further preferably 2 g - 6 g, still
15 more preferably 2.5 g - 6 g, as the total amount of one or more kinds selected from the group consisting of leucine and phenylalanine (i.e., either or both of leucine and phenylalanine) (the total amount converted to the amount of free form), the total amount of either or both of leucine and
20 phenylalanine, and lysine (the total amount converted to the amount of free form), the total amount of either or both of leucine and phenylalanine, and one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan (the total amount converted to the amount of free
25 form), or the total amount of either or both of leucine and phenylalanine, lysine, and one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan (the total amount converted to the amount of free form).

30 [0049]

In addition, the ingestion or dosing period of the composition of the present invention is also appropriately determined according to the condition and symptoms of the application subject, and the like. Considering that a decline
35 in cognitive function occurs along with aging and various

diseases, intracerebral lesions such as cerebral atrophy and the like, and the like, and chronically progresses, or that anxiety-like symptoms appear along with a decline in cognitive function, or before and after the decline in cognitive function, and persist for a long time, continuous ingestion or administration for a long period of time is preferable to improve cognitive function or anxiety-like symptoms.
[0050]

The composition of the present invention can be formulated as a unit package form. In the present specification, the "unit package form" means a form of one or more units with a particular amount (e.g., ingestion amount or dose per one time etc.) as one unit is/are filled in one container or packed in a package. For example, a unit package form with ingestion amount or dose per one time as one unit is referred to as "unit package form for ingestion amount or dose per one time". A container or package used for the unit package form can be appropriately selected according to the form and the like of the composition of the present invention. For example, paper container or bag, plastic container or bag, pouch, aluminum can, steel can, glass bottle, pet bottle, PTP (press through pack) package sheet and the like can be mentioned.
[0051]

The application subject of the composition of the present invention includes, for example, mammals (e.g., human, monkey, mouse, rat, guinea pig, hamster, rabbit, cat, dog, bovine, horse, donkey, swine, sheep, etc.), birds (e.g., duck, chicken, goose, turkey, etc.) and the like.

When the composition of the present invention is applied to an application subject animal (hereinafter to be also simply referred to as "subject animal") other than human, the ingestion amount or dose of the composition of the present invention can be appropriately set according to the kind, sex, body weight and the like of the subject animal.

[0052]

The composition of the present invention has a cognitive function improving effect, and is effective for the prophylaxis or improvement of various symptoms or disorders caused by a
5 decline in cognitive function such as a decline in learning ability and memory, learning and memory disorder, and the like.

In addition, the composition of the present invention has an improving effect on anxiety-like symptoms, and is effective for the prophylaxis of the expression of anxiety-like symptoms
10 such as anxiety, agitation and the like or reduction of the aforementioned symptoms.

Particularly, it is effective for preventing the expression of anxiety-like symptoms that appear along with a decline in cognitive function or before and after the decline
15 of cognitive function, or for reducing the aforementioned symptoms.

Furthermore, the composition of the present invention has an effect of suppressing cerebral atrophy, cerebral atrophy observed in neurodegenerative diseases such as Alzheimer-type
20 dementia and the like, cerebral atrophy that appears and progresses along with aging, and the like, and is effective for improving a decline in cognitive function caused by these.

Particularly, since the composition of the present invention contains amino acids, which are contained in foods
25 and have abundant food experience, as active ingredients, it is highly safe, suitable for continuous ingestion or administration, and thus suitable for preventing or improving a decline in cognitive that chronically progresses for a long time along with aging and the like, improving anxiety-like
30 symptoms that persist for a long time, or suppressing the appearance and progression of cerebral atrophy.

[0053]

Therefore, the composition of the present invention can be preferably ingested by or administered to those exhibiting
35 symptoms or disorders due to a decline in cognitive function,

elderly people and middle- or late middle-aged persons requiring improvement of cognitive function, and the like.

In addition, the composition of the present invention can be preferably ingested by or administered to those exhibiting
5 the above-mentioned anxiety-like symptoms, and elderly people and middle- or late middle-aged persons who may develop the above-mentioned anxiety-like symptoms, and the like.

Furthermore, the composition of the present invention can be preferably ingested by or administered to patients with
10 neurodegenerative diseases in which appearance and progression of cerebral atrophy are observed, such as Alzheimer-type dementia and the like, elderly people who are highly likely to show appearance and progression of cerebral atrophy, and the like.

15 [0054]

The composition of the present invention can be provided as a pharmaceutical product (hereinafter to be also referred to as "the pharmaceutical product of the present invention" in the present specification) directly or by further adding the above-
20 mentioned pharmaceutically acceptable additives.

The pharmaceutical product of the present invention can have a dosage form of oral preparation such as tablet, coated tablet, chewable tablet, pill, (micro)capsule, granule, fine granule, powder, elixir, lemonade, syrup, suspension, emulsion,
25 oral jelly and the like, injection such as solution, suspension, emulsion and the like, solid injection to be used by dissolving or suspending when in use, injectable preparation such as transfusion, sustainable injection and the like, tubal liquid, and the like.

30 [0055]

The pharmaceutical product of the present invention may contain an anti-dementia drug as long as the characteristics of the present invention are not impaired.

Examples of the anti-dementia drug include
35 acetylcholinesterase inhibitors such as donepezil hydrochloride,

galanthamine, rivastigmine and the like; and NMDA receptor antagonists such as memantine and the like, and these can be used according to general dosage and administration.

[0056]

5 The pharmaceutical product of the present invention can be preferably administered to patients exhibiting symptoms or disorders due to a decline in cognitive function, patients who may develop the aforementioned symptoms or disorders, elderly people and middle- or late middle-aged persons exhibiting a
10 decline in cognitive function, and the like.

 In addition, the pharmaceutical product of the present invention can be preferably administered to patients exhibiting anxiety-like symptoms that appear along with a decline in cognitive function or before and after a decline in cognitive
15 function, elderly people and middle- or late middle-aged persons exhibiting the aforementioned anxiety-like symptoms, and the like.

 Furthermore, the pharmaceutical product of the present invention can be preferably ingested by or administered to
20 patients with neurodegenerative diseases in which appearance and progression of cerebral atrophy are observed, such as Alzheimer-type dementia and the like, elderly people and middle- or late middle-aged persons who may be affected with the aforementioned neurodegenerative diseases, elderly people
25 who are highly likely to show appearance and progression of cerebral atrophy, and the like.

 The pharmaceutical product of the present invention is administered per day to the above-mentioned application subject such that the total amount of one or more kinds selected from
30 the group consisting of leucine and phenylalanine (i.e., either or both of leucine and phenylalanine) (the total amount converted to the amount of free form), the total amount of either or both of leucine and phenylalanine, and lysine (the total amount converted to the amount of free form), the total
35 amount of either or both of leucine and phenylalanine, and one

or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan (the total amount converted to the amount of free form), or the total amount of either or both of leucine and phenylalanine, lysine, and one or
5 more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan (the total amount converted to the amount of free form) would be the above-mentioned daily dose.

[0057]

10 Furthermore, the composition of the present invention can be ingested by adding to various foods. The food to which the composition of the present invention is added is not particularly limited, and may be any as long as it is a food in the form generally served for meals or dessert.

15 For example, the composition of the present invention is added to drinks such as beverage water and the like, and a suitable flavor is added when desired, whereby a drink can be provided.

More specifically, the composition of the present
20 invention can be added, for example, to beverage water such as fruit juice drinks, sport drinks and the like; dairy products such as milk, yogurt and the like; confectionery such as jelly, chocolate, candy, biscuit and the like, and the like.

[0058]

25 The composition of the present invention is preferably added to the above-mentioned various foods in the amounts to be ingested per day such that the total amount of one or more kinds selected from the group consisting of leucine and phenylalanine (i.e., either or both of leucine and
30 phenylalanine) (the total amount converted to the amount of free form), the total amount of either or both of leucine and phenylalanine, and lysine (the total amount converted to the amount of free form), the total amount of either or both of leucine and phenylalanine, and one or more kinds selected from
35 the group consisting of valine, isoleucine, histidine and

tryptophan (the total amount converted to the amount of free form), or the total amount of either or both of leucine and phenylalanine, lysine, and one or more kinds selected from the group consisting of valine, isoleucine, histidine and

5 tryptophan (the total amount converted to the amount of free form) would be the above-mentioned ingestion amount per day.
[0059]

The composition of the present invention can be provided as a food (hereinafter to be also referred to as "the food of
10 the present invention" in the present specification) directly or by adding general food additives as necessary and according to a general food production technique.

The food of the present invention can be prepared as various forms such as liquid (e.g., solution, suspension,
15 emulsified liquid and the like); semi-solid (e.g., gel, cream and the like); solid (e.g., powder, granule, sheet, capsule, tablet and the like), and the like.

Furthermore, the food of the present invention can be prepared as various food forms such as beverage water (fruit
20 juice drinks, sport drinks, coffee drinks, tea drinks etc.), dairy product (lactic fermenting beverage, fermented milk, butter, cheese, yogurt, processed milk, defatted milk etc.), meat product (ham, sausage, hamburger etc.), fish meat processed seafood paste product (fish cake, tube-shaped fish
25 sausage, deep-fried ball of fish paste etc.), egg product (rolled Japanese-style egg omelette, steamed egg custard etc.), confectionery (cookie, jelly, chewing gum, candy, snack food, frozen dessert etc.), bread, noodles, pickle, dried fish, food boiled in soy sauce, soup, seasoning and the like by adding the
30 composition of the present invention to various food starting materials and adding general food additives as necessary. It may also be a bottled food, canned food or retort pouch food.
[0060]

As the above-mentioned food additive, manufacturing agent
35 (brine, binding agent etc.), thickening stabilizer (xanthan gum,

sodium carboxymethylcellulose etc.), gelation agent (gelatin, agar, carrageenan etc.), gum base (vinyl acetate resin, jelutong, chicle etc.), emulsifier (glycerol fatty acid ester, sucrose fatty acid ester, saponin, lecithin etc.), preservative
5 (benzoic acid, sodium benzoate, sorbic acid, potassium sorbate, ϵ -polylysine etc.), antioxidant (ascorbic acid, erythorbic acid, catechin etc.), glazing agent (shellac, paraffin wax, beeswax etc.), fungicide (thiabendazole, fludioxonil etc.), leavening agent (sodium hydrogen carbonate, glucono- δ -lactone, alum etc.),
10 sweetener (aspartame, acesulfame potassium, licorice extract etc.), bittering agent (caffeine, naringin, worm wood extract etc.), acidulant (citric acid, tartaric acid, lactic acid etc.), seasoning (sodium L-glutamate, disodium 5'-inosinate etc.), colorant (annatto dye, turmeric dye, gardenia dye etc.), flavor
15 (synthetic flavor such as ethyl acetoacetate, anisaldehyde and the like, natural flavor such as orange, lavender and the like) and the like can be mentioned.

In the present invention, one or more kinds of the above-mentioned food additives can be used.

20 [0061]

The food of the present invention can be preferably ingested by elderly people and middle- or late middle-aged persons showing a decline in cognitive function, elderly people and middle- or late middle-aged persons who may show a decline
25 in cognitive function, and further, a wide range of subjects for the purpose of preventing a decline in cognitive function.

In addition, the food of the present invention can be ingested by elderly people and middle- or late middle-aged persons exhibiting anxiety-like symptoms that appear along with
30 a decline in cognitive function or before and after a decline in cognitive function, elderly people and middle- or late middle-aged persons who may express the aforementioned anxiety-like symptoms, and further, a wide range of subjects for the purpose of preventing expression of the aforementioned anxiety-
35 like symptoms.

Furthermore, the food of the present invention can be widely ingested by patients with neurodegenerative diseases in which appearance and progression of cerebral atrophy are observed, such as Alzheimer-type dementia and the like, elderly
5 people and middle- or late middle-aged persons who may be affected with the aforementioned neurodegenerative diseases, elderly people who are highly likely to show appearance and progression of cerebral atrophy, and the like, for the purpose of suppressing the appearance and progression of cerebral
10 atrophy.

[0062]

Therefore, the food of the present invention can also be provided as food with health claims such as food for specified health uses, food with nutrient function claims, food with
15 function claims and the like, food for special dietary uses such as food for sick people, food for the elderly and the like, health supplement, and the like for improving cognitive function or anxiety-like symptoms, or for suppressing cerebral atrophy.

20 [0063]

The food of the present invention is preferably ingested per day by the above-mentioned application subject such that the total amount of one or more kinds selected from the group consisting of leucine and phenylalanine (i.e., either or both
25 of leucine and phenylalanine) (the total amount converted to the amount of free form), the total amount of either or both of leucine and phenylalanine, and lysine (the total amount converted to the amount of free form), the total amount of either or both of leucine and phenylalanine, and one or more
30 kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan (the total amount converted to the amount of free form), or the total amount of either or both of leucine and phenylalanine, lysine, and one or more kinds selected from the group consisting of valine, isoleucine,
35 histidine and tryptophan (the total amount converted to the

amount of free form) would be the above-mentioned ingestion amount per day.

[0064]

Furthermore, the present invention also provides a method
5 for improving cognitive function in a subject animal in need of improvement of cognitive function (hereinafter to be also referred to as the "method of the present invention" in the present specification).

[0065]

10 The method of the present invention includes allowing a subject animal in need of improving cognitive function to ingest a composition containing one or more kinds selected from the group consisting of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content
15 of seven amino acids, in an amount effective for improving cognitive function of the subject animal, or administering said amount of the composition to the animal.

In the method of the present invention, ingestion or administration of a composition containing both leucine and
20 phenylalanine at the above-mentioned composition ratio is preferable, ingestion or administration of a composition further containing 10 mol% - 30 mol% of lysine relative to the total content of seven amino acids in addition to one or more kinds selected from the group consisting of leucine and
25 phenylalanine at the above-mentioned composition ratio is more preferable, and ingestion or administration of a composition containing leucine, phenylalanine and lysine at the above-mentioned composition ratio is further preferable.

In the method of the present invention, ingestion or
30 administration of a composition further containing one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan in addition to one or more kinds selected from the group consisting of leucine and phenylalanine at the above-mentioned composition ratio, or one or more kinds
35 selected from the group consisting of leucine and phenylalanine,

and lysine at the above-mentioned composition ratio is preferable.

From the aspect of a cognitive function improving effect, in the method of the present invention, ingestion or
5 administration of a composition containing leucine, phenylalanine and lysine at the above-mentioned composition ratio, and one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan is more preferable, and ingestion or administration of a composition
10 containing leucine, phenylalanine and lysine at the above-mentioned composition ratio, and valine, isoleucine, histidine and tryptophan is particularly preferable.

[0066]

In the method of the present invention, the content of
15 valine in the composition to be ingested by or administered to a subject animal is preferably 2 mol% - 7 mol%, more preferably 4 mol% - 6 mol%, relative to the total content of seven amino acids.

The content of isoleucine in the composition to be
20 ingested by or administered to a subject animal is preferably 5 mol% - 15 mol%, more preferably 7 mol% - 13 mol%, relative to the total content of seven amino acids.

The content of histidine in the composition to be
ingested by or administered to a subject animal is preferably 2
25 mol% - 10 mol%, more preferably 2.5 mol% - 8 mol%, relative to the total content of seven amino acids.

The content of tryptophan in the composition to be
ingested by or administered to a subject animal is preferably
0.1 mol% - 2 mol%, more preferably 0.3 mol% - 1 mol%, relative
30 to the total content of seven amino acids.

[0067]

The subject animal in the method of the present invention includes mammal (e.g., human, monkey, mouse, rat, guinea pig, hamster, rabbit, cat, dog, bovine, horse, donkey, swine, sheep
35 etc.), birds (e.g., duck, chicken, goose, turkey etc.) and the

like.

[0068]

The method of the present invention is effective for the prophylaxis or improvement of various symptoms or disorders
5 caused by a decline in cognitive function such as a decline in learning ability and memory, learning and memory disorder, and the like.

In addition, the method of the present invention is effective for improving anxiety-like symptoms, namely, for the
10 prophylaxis of the expression of anxiety-like symptoms such as anxiety, agitation and the like or reduction of the aforementioned anxiety-like symptoms. It is particularly effective for preventing the expression of anxiety-like symptoms that appear along with a decline in cognitive function
15 or before and after the decline of cognitive function, or for reducing the aforementioned anxiety-like symptoms.

Furthermore, the method of the present invention is effective for suppressing cerebral atrophy observed in neurodegenerative diseases such as Alzheimer-type dementia and
20 the like, cerebral atrophy that appears and progresses along with aging, and the like, and thus effective for improving a decline in cognitive function caused by these.

Particularly, since the method of the present invention uses amino acids, which are contained in foods and have
25 abundant food experience, as an active ingredient, it has high safety and can be applied continuously.

[0069]

In the case of human, the method of the present invention is preferably applied to patients exhibiting symptoms and
30 disorders due to a decline in cognitive function, and elderly people, and middle- or late middle-aged persons requiring prevention of a decline in cognitive function such as elderly people, and middle- or late middle-aged persons who may show a decline in cognitive function.

35 In addition, the method of the present invention is

preferably applied to elderly people and middle- or late middle-aged persons exhibiting anxiety-like symptoms, and elderly people and middle- or late middle-aged persons who may express anxiety-like symptoms, and particularly preferably applied to elderly people and middle- or late middle-aged persons exhibiting anxiety-like symptoms that appear along with a decline in cognitive function or before and after a decline in cognitive function, and elderly people and middle- or late middle-aged persons who may express the aforementioned anxiety-like symptoms.

Furthermore, the method of the present invention is preferably applied to patients with neurodegenerative disease who are known to show cerebral atrophy such as Alzheimer-type dementia and the like, elderly people and middle- or late middle-aged persons who may be affected with the aforementioned neurodegenerative diseases, elderly people who may show cerebral atrophy, and the like.

[0070]

The effective amount of the composition containing one or more kinds selected from the group consisting of leucine and phenylalanine at the above-mentioned composition ratio in the method of the present invention is appropriately determined according to the kind, age, sex, the condition or level of a decline in cognitive function, intracerebral lesion or anxiety-like symptoms and the like of the subject animal. An amount similar to the above-mentioned ingestion amount or dose of the composition of the present invention for a human or a subject animal other than human can be ingested or administered at the frequency and period mentioned above.

[0071]

As an ingestion or administration method of the composition containing one or more kinds selected from the group consisting of leucine and phenylalanine at the above-mentioned composition ratio and the like in the method of the present invention, oral ingestion or oral administration,

enteral tube administration, administration by infusion and the like can be mentioned. Oral ingestion or oral administration is preferable since convenient ingestion is possible without the need to perform under the guidance and supervision of a
5 doctor at a medical institution.

[Example]

[0072]

The present invention is explained in further detail in the following by referring to Examples.

10 [0073]

[Example 1] Composition for improving cognitive function

To afford the composition shown in Table 1, given amounts of respective components were weighed and mixed to prepare the composition for improving cognitive function of Example 1
15 (hereinafter to be referred to as "the composition of Example 1"). Similarly, a composition with the composition ratio shown in Table 1 was prepared and used as Comparative Example 1.

[0074]

[Table 1]

component	composition ratio relative to total content of amino acids (mol%)	
	Example 1	Comparative Example 1
L-leucine	36.10	20.32
L-lysine hydrochloride	18.42	20.54
L-valine	5.02	4.06
L-isoleucine	9.96	3.63
L-phenylalanine	26.09	17.72
L-histidine hydrochloride	3.89	33.13
L-tryptophan	0.52	0.60
total amount	100	100

20

[0075]

[Experimental Example 1] Study of effect on learning and memory functions

Using C57Bl/6j aged mice (58-week-old to 61-week-old,

male) (purchased from CHARLES RIVER LABORATORIES JAPAN, INC.), an effect of the compositions of Example 1 and Comparative Example 1 on learning and memory functions was studied.

C57Bl/6j aged mice were divided into 4 groups as shown in Table 2 (n=12/group). To a group allowed to freely ingest a normal protein diet (containing 20 wt% casein) was administered a vehicle alone (NPD+Veh.). The other 3 groups were allowed to freely ingest a low-protein diet (containing 5 wt% casein), and respectively used as a group administered with a vehicle alone (LPD+Veh.), a group administered with the composition of Example 1 (LPD+E1), and a group administered with the composition of Comparative Example 1 (LPD+C1). In LPD+E1 and LPD+C1, the composition of Example 1 (E1) and the composition of Comparative Example 1 (C1) at 1 g/kg body weight were each dissolved in 0.5 wt% methylcellulose aqueous solution, and continuously administered by oral gavage (cycle of 2 days off after administration for 5 consecutive days) 2 times daily (morning and evening) for 2 months. A 0.5 wt% methylcellulose aqueous solution (Vehicle) was similarly administered to NPD+Veh. and LPD+Veh.

[0076]

[Table 2]

group	diet	administered sample
NPD+Veh.	diet containing 20 wt% casein	vehicle (0.5 wt% methylcellulose aqueous solution)
LPD+Veh.	diet containing 5 wt% casein	vehicle (0.5 wt% methylcellulose aqueous solution)
LPD+C1	diet containing 5 wt% casein	sample containing the composition of Comparative Example 1
LPD+E1	diet containing 5 wt% casein	sample containing the composition of Example 1

[0077]

The mice of each group shown in Table 2 were subjected to a passive avoidance test as follows after 1-month administration and 2-month administration of a vehicle or each composition of Example 1 and Comparative Example 1.

(i) Using an apparatus composed of light/dark compartments with an electrical stimulation device in the dark compartment, mice were placed in the light compartment. When the mice entered the dark compartment, the door was closed simultaneously with the entry, and an electrical stimulation (1 mA, 1 sec) was applied. The entry into the dark compartment was defined to be the time point when the base of the tail entered the dark compartment.

(ii) (i) was tried once, and 24 hr later, the mice were placed in a light compartment, and the time until the mice entered a dark compartment (transfer latency time) was measured. In addition, the proportion of individuals that stayed in the light compartment for a maximum evaluation time of 480 seconds, that is, the proportion of individuals that memorized that the dark compartment is scary by one electrical stimulation (achievement by 1st ES) was determined.

[0078]

A vehicle or the composition of Example 1 or Comparative Example 1 was administered for one month, then a passive avoidance test was performed, and the time required by the mice in each group until they enter the dark compartment (transfer latency time) was measured. The results are shown in Fig. 1 as mean + standard error of the mean of each of 12 mice.

In addition, the proportion of individuals that memorized that the dark compartment is scary by one electrical stimulation (achievement by 1st ES) is also shown in Fig. 1. The numerical values in the Figure show the number of mice in each group that remained in the light compartment for 300 sec, after 24 hr from the first trial.

As for the time required for the entry into a dark compartment (transfer latency time), which was measured 24 hr after the first trial, one-way analysis of variance and Dunnett's multiple comparison test were performed with respect to the group raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.). As for the proportion of

individuals that memorized that the dark compartment is scary by one electrical stimulation (achievement by 1st ES), a chi-square test was performed with respect to NPD+Veh.

[0079]

5 As shown in Fig. 1, the time required for the entry into a dark compartment (transfer latency time) 24 hr after the first trial, drastically decreased in the group raised on a low-protein diet and administered with a vehicle alone (LPD+Veh.) as compared to the group raised on a normal protein
10 diet and administered with a vehicle alone (NPD+Veh.) ($p < 0.05$).

In the group orally administered with the composition of Example 1 of the present invention in addition to a low-protein diet (LPD+E1), it was observed that the transfer latency time was recovered to such an extent that no significant difference
15 was observed between LPD+E1 and NPD+Veh.

On the other hand, the group that received oral administration of the composition of Comparative Example 1 in addition to a low-protein diet (LPD+C1), the transfer latency time was not recovered.

20 The proportion of individuals that memorized that the dark compartment is scary by one electrical stimulation (achievement by 1st ES) decreased in LPD+Veh. as compared to NPD+Veh., and was recovered in LPD+E1 to the same level as NPD+Veh. LPD+C1 did not show such recovery.

25 [0080]

[Experimental Example 2] Study of effect on learning and memory functions (reproducibility test)

Using C57Bl/6j mature mice (19-week-old, male) (purchased from CHARLES RIVER LABORATORIES JAPAN, INC.), the effect of the
30 composition of Example 1 on the learning and memory functions was studied.

C57Bl/6j mature mice were divided into 4 groups (n=12/group) shown in Table 3. To a group raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.),
35 a group raised on a low-protein diet and administered with a

vehicle alone (LPD+Veh.), a group raised on a low-protein diet and administered with the composition of Example 1 once per day (1 g/kg body weight per administration) (LPD+E1×1), and a group raised on a low-protein diet and administered with the composition of Example 1 twice per day (1 g/kg body weight per administration) (LPD+E1×2) was administered a vehicle or the composition of Example 1 for 1 to 1.5 months, and a passive avoidance test was performed in the same manner as in Experimental Example 1, and the time required until entry into the dark compartment (transfer latency time) 24 hr after the first trial, and the proportion of individuals that memorized that the dark compartment is scary by one electrical stimulation (achievement by 1st ES) were determined. The transfer latency time and achievement by 1st ES in the passive avoidance test performed after administration for one month are shown in Fig. 2 as mean + standard error of the mean of each of 12 mice. As for the transfer latency time, one-way analysis of variance and Dunnett's multiple comparison test were performed with respect to the group raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.). As for the achievement by 1st ES, a chi-square test was performed with respect to NPD+Veh.

[0081]

[Table 3]

group	diet	administered sample
NPD+Veh.	diet containing 20 wt% casein	vehicle (0.5 wt% methylcellulose aqueous solution)
LPD+Veh.	diet containing 5 wt% casein	vehicle (0.5 wt% methylcellulose aqueous solution)
LPD+E1×1	diet containing 5 wt% casein	sample containing the composition of Example 1, administered once per day
LPD+E1×2	diet containing 5 wt% casein	sample containing the composition of Example 1, administered twice per day

25

[0082]

As shown in Fig. 2, the transfer latency time and the achievement by 1st ES showed a significant ($p < 0.05$ and $p < 0.1$)

decrease also in mature mice in the group raised on a low-protein diet and administered with a vehicle alone (LPD+Veh.) as compared to the group raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.). In the group
5 raised on a low-protein diet and administered with the composition of Example 1 (LPD+E1×1 and LPD+E1×2), it was found that both the transfer latency time and the achievement by 1st ES that decreased by raising on a low-protein diet were recovered to such an extent that no significant difference was
10 observed from when raised on a normal protein diet.

The sample containing the composition of Example 1 of the present invention, even when administered once per day, showed the same level of effect as that achieved when administered twice per day.

15 [0083]

From the results of the above-mentioned Experimental Examples 1 and 2, it was suggested that the composition of Example 1 of the present invention suppresses and improves a decrease in the learning and memory functions caused by the
20 ingestion of a low-protein diet.

[0084]

[Experimental Example 3] Study of effect on anxiety, agitation and hyperactivity symptoms

Using C57Bl/6j aged mice (58-week-old to 61-week-old,
25 male) (purchased from CHARLES RIVER LABORATORIES JAPAN, INC.), an effect of the compositions of Example 1 of the present invention and the composition of Comparative Example 1 on anxiety, agitation and hyperactivity symptoms was evaluated by an elevated plus maze test.

30 C57Bl/6j aged mice were grouped as shown in Table 2 (n=12/group), and each group was subjected to an elevated plus maze test as follows after 1-month administration and 2-month administration of a vehicle or each composition of Example 1 and Comparative Example 1.

35 That is, the mice in each group were placed in the

neutral zone of the elevated plus maze, and the behavior of the mice was observed for 8 min. The number of times the mouse entered the runway without a wall (open arm) and the distance moved within the track were analyzed using video tracking behavior analysis software Smart3.0 (Panlab), and the number of entries in open arms (rate of entries in open arms) and the total distance moved were calculated. The rate of entries in open arms becomes an index of anxiety and agitation symptoms, and the total distance moved becomes an index of hyperactivity symptoms. The rate of entries in open arms after 2 months of administration and the total distance moved after 1 month of administration are shown in Fig. 3 as mean + standard error of the mean. As for the number of entries in open arms, one-way analysis of variance and Dunnett's multiple comparison test were performed between the group raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.) and other groups. As for the total distance moved, one-way analysis of variance and Dunnett's multiple comparison test were performed between NPD+Veh. and the group raised on a low-protein diet and administered with a vehicle alone (LPD+Veh.), between LPD+Veh. and each of the group raised on a low-protein diet and administered with the composition of Example 1 of the present invention (LPD+E1), and the group raised on a low-protein diet and administered with the composition of Comparative Example 1 (LPD+C1).

[0085]

As shown in Fig. 3, both the number of entries in open arms, and the total distance moved showed a significant ($p < 0.05$) increase in the group raised on a low-protein diet and administered with a vehicle alone (LPD+Veh.) as compared to the group raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.).

On the other hand, in the group raised on a low-protein diet and administered with the composition of Example 1 of the present invention (LPD+E1), the number of entries in open arms

showed an increase of a level without a significant difference from that of NPD+Veh., and the total distance moved significantly decreased ($p<0.05$) as compared to LPD+Veh.

The same results as in LPD+E1 were also obtained in the group raised on a low-protein diet and administered with the composition of Comparative Example 1 (LPD+C1).

[0086]

[Experimental Example 4] Study of effect on anxiety, agitation and hyperactivity symptoms (reproducibility test)

Using C57Bl/6j mature mice (19-week-old, male) (purchased from CHARLES RIVER LABORATORIES JAPAN, INC.), the effect of the composition of Example 1 on agitation, anxiety and hyperactivity symptoms was evaluated by an elevated plus maze test.

C57Bl/6j mature mice were grouped ($n=12$ /group) as shown in Table 3. After 1-month to 1.5-month administration of the composition of Example 1 and a vehicle, the mice in each group were placed in the neutral zone of the elevated plus maze, and the behavior of the mice was observed for 8 min. Similar to the case of Experimental Example 3, the number of entries in open arms and the total distance moved were determined, and the evaluation results of the number of entries in open arms after 1-month administration are shown in Fig. 4 as mean + standard error of the mean.

As for the total distance moved, the group raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.) showed the same level of increase as the group raised on a low-protein diet and administered with a vehicle alone (LPD+Veh.), and the effect of the composition of Example 1 could not be evaluated.

As for the results shown in Fig. 4, one-way analysis of variance and Dunnett's multiple comparison test were performed between NPD+Veh. and LPD+Veh., between LPD+Veh. and each of the groups raised on a low-protein diet and administered with the composition of Example 1 once per day (1 g/kg body weight per

administration) (LPD+E1×1), and the group raised on a low-protein diet and administered with the composition of Example 1 twice per day (1 g/kg body weight per administration) (LPD+E1×2).

5 [0087]

As shown in Fig. 4, the number of entries in open arms to be the index of anxiety and agitation significantly ($p<0.05$) increased in LPD+Veh. as compared to NPD+Veh.

On the other hand, a decrease in the number of entries in
10 open arms was found in the group raised on a low-protein diet and administered with the composition of Example 1 once per day (LPD+E1×1), and the number of entries in open arms significantly ($p<0.05$) decreased in the group raised on a low-protein diet and administered with the composition of Example 1
15 twice per day (LPD+E1×2). That is, it was shown that the decrease in the number of entries in open arms due to the administration of the composition of Example 1 depends on the dose of the composition of Example 1.

[0088]

20 The results of Experimental Example 3 suggest that the composition of Example 1 of the present invention suppresses anxiety and agitation symptoms and hyperactivity, and the results of Experimental Example 4 suggest that the composition of Example 1 of the present invention suppresses anxiety and
25 agitation symptoms in a dose-dependent manner.

[0089]

[Experimental Example 5] Study of effect on intracerebral neurotransmitter concentration

In Experimental Example 1 and Experimental Example 3,
30 respective evaluations were performed by a passive avoidance test and an elevated plus maze test, after which the mice in each group were dissected, and the concentration of the neurotransmitters (dopamine, norepinephrine, glutamic acid) in the prefrontal cortex was measured using the high performance
35 liquid chromatograph (HPLC) method and the liquid

chromatography mass spectrometry (LC/MS) method. As for the measurement results of the concentrations of dopamine and glutamic acid, one-way analysis of variance and Tukey's multiple comparison test were performed between the group
5 raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.) and the group raised on a low-protein diet and administered with a vehicle alone (LPD+Veh.), between LPD+Veh. and each of the group raised on a low-protein diet and administered with the composition of Example 1 (LPD+E1), and
10 the group raised on a low-protein diet and administered with the composition of Comparative Example 1 (LPD+C1), and as for the measurement results of the norepinephrine concentration, between NPD+Veh. and LPD+Veh., and between NPD+Veh. and each group of LPD+E1 and LPD+C1. The results are shown in Fig. 5 as
15 mean + standard error of the mean.

[0090]

As shown in Fig. 5, it was found that the concentration of each of intracerebral dopamine, norepinephrine and glutamic acid decreased in the group raised on a low-protein diet and
20 administered with a vehicle alone (LPD+Veh.) as compared to the group raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.) (significant at $p<0.05$, significant at $p<0.01$, and significant tendency at $p=0.21$, respectively).

On the other hand, a significant increase in the
25 intracerebral dopamine concentration and glutamic acid concentration was found (significant at $p<0.001$ and $p<0.05$, respectively) in the group raised on a low-protein diet and administered with the composition of Example 1 (LPD+E1) as compared to the LPD+Veh. group, and it was found that the
30 intracerebral norepinephrine concentration increased to a level without significant difference from the NPD+Veh. group.

[0091]

Dopamine, norepinephrine and glutamic acid that decreased in the group raised on a low-protein diet in Experimental
35 Example 5 are intracerebral neurotransmitters known to decrease

in dementia patients.

From the results of Experimental Example 5, it was suggested that the composition of the present invention may improve a decrease in the intracerebral neurotransmitters and
5 improve cognitive function.

[0092]

[Experimental Example 6] Study of influence on brain-derived neurotrophic factor (BDNF) gene expression

Brain-derived neurotrophic factor (BDNF) is one of the
10 proteins belonging to the neurotrophic factor family and has various functions in the cranial nervous system, such as survival of nerve cells, formation of neural network, and expression of higher brain functions such as memory consolidation and the like. In addition, a decrease in BDNF
15 expression level is observed in various neurodegenerative diseases including dementia and psychiatric diseases, and a drug that induces the expression of BDNF or a composition containing the same is expected to show an effect of improving brain function impaired by neurodegenerative diseases or
20 psychiatric diseases.

Thus, an influence of the above-mentioned composition of Example 1 of the present invention on the expression of BDNF gene was studied.

In the above-mentioned Experimental Example 1 and
25 Experimental Example 3, respective evaluations were performed by a passive avoidance test and an elevated plus maze test, after which the mice in each group were dissected to obtain cerebral cortex tissue, and the expression level of the BDNF gene in the cerebral cortex tissue was studied by measuring
30 changes in the mRNA expression level by the RT-qPCR method.

After the completion of respective evaluations in Experimental Example 1 and Experimental Example 3, RNA was extracted using an RNA extraction kit ("RNeasy Mini Kit" (QUIAGEN)) from the cerebral cortex tissue of the mice
35 (n=6/group) of each group shown in Table 4, and a reverse

transcription reaction was performed using a reverse transcription reaction premix reagent ("Prime Script (trade mark) RT Master Mix" (Takara Bio Inc.)). Then, changes in the BDNF mRNA expression level of the obtained DNA product were
 5 analyzed using a real-time PCR kit ("SYBR (registered trade mark) Select Master Mix" (Applied Biosystems)).

As for the measurement results, the BDNF expression level was compared between the group raised on a normal protein diet and administered with a vehicle alone (NPD+Veh.) and the group
 10 raised on a low-protein diet and administered with a vehicle alone (LPD+Veh.), and the group raised on a low-protein diet and administered with the composition of Example 1 (LPD+E1), and the ratio to NPD+Veh. was determined and shown in Fig. 6 as mean + standard error of the mean of 6 mice in each group.
 15 Statistical analysis of comparison among 3 groups was performed by one-way analysis of variance and Dunnett's multiple comparison test.

[0093]

[Table 4]

group	diet	administered sample
NPD+Veh.	diet containing 20 wt% casein	vehicle (0.5 wt% methylcellulose aqueous solution) (twice per day)
LPD+Veh.	diet containing 5 wt% casein	vehicle (0.5 wt% methylcellulose aqueous solution) (twice per day)
LPD+E1	diet containing 5 wt% casein	solution of the composition of Example 1 in 0.5 wt% methylcellulose aqueous solution (1 g/Kg body weight per administration, twice per day)

20

[0094]

As shown in Fig. 6, the expression level of the BDNF gene was significantly ($p < 0.05$) high in the group raised on a low-protein diet and administered with the composition of Example 1
 25 (LPD+E1), as compared to the group raised on a low-protein diet and administered with a vehicle alone (LPD+Veh.).

[0095]

The above-mentioned results of Experimental Example 6

suggest that the composition of Example 1 of the present invention may induce BDNF gene expression and improve lowered cognitive function.

[0096]

5 [Experimental Example 7] Study of influence on cerebral atrophy in Alzheimer-type dementia model mouse

In many patients with neurodegenerative diseases including Alzheimer-type dementia, aggregates of fibrotic tau protein are observed in nerve cells and glial cells. rTg4510
10 transgenic mouse is a mouse transformed to specifically express such tau protein in nerve cells, and is known to exhibit cerebral atrophy with aging. Thus, it is used as a model mouse for Alzheimer-type dementia.

Accordingly, an influence of the composition of Example 1
15 on cerebral atrophy was studied using rTg4510 mouse and its littermate control mouse.

The 3-month-old male rTg4510 mice and littermate control mice were divided into 3 groups (n=3-4/group) as shown in Table 5, and raised allowing free ingestion of a normal protein diet
20 (containing 20 wt% casein). To one of groups of rTg4510 mice was administered by oral gavage the composition of Example 1 dissolved in 0.5 wt% methylcellulose aqueous solution twice per day (1 g/kg body weight per administration) (rTg4510+E1), and a vehicle (0.5 wt% methylcellulose aqueous solution) alone was
25 administered to the other group of rTg4510 mice and the littermate control mice group (rTg4510+Veh. and Ctrl+Veh.).

At the start of the test and after administration of the vehicle or the composition of Example 1 for 3.5 months, brain magnetic resonance imaging data of each mouse was obtained
30 using 7.0 Tesla MRI for small animals (Bruker), and the volume of cerebral cortex was calculated using image analysis software "Pmod 3.7" (Pmod Technologies Ltd.). The amount of change in the cerebral cortex volume after administration of the vehicle or the composition of Example 1 for 3.5 months relative to the
35 cerebral cortex volume at the start of the test (3-month-old)

was determined in each group, and shown in Fig. 7 as mean $\pm$ standard error of the mean of 3 to 4 mice. Statistical analysis was performed by one-way analysis of variance and Tukey's multiple comparison test of the comparison among 3

5 groups.

[0097]

[Table 5]

group	diet	administered sample
Ctrl+Veh.	diet containing 20 wt% casein	vehicle (0.5 wt% methylcellulose aqueous solution) (twice per day)
rTg4510+Veh.	diet containing 20 wt% casein	vehicle (0.5 wt% methylcellulose aqueous solution) (twice per day)
rTg4510+E1	diet containing 20 wt% casein	solution of the composition of Example 1 in 0.5 wt% methylcellulose aqueous solution (1 g/Kg body weight per administration, twice per day)

[0098]

10 As shown in Fig. 7, a significant ($p < 0.01$) decrease in the cerebral cortex volume was found in the rTg4510 mouse group administered with a vehicle alone (rTg4510+Veh.) as compared to the littermate control mouse group administered with a vehicle alone (Ctrl+Veh.).

15 On the other hand, the rTg4510 mouse group administered with the composition of Example 1 (rTg4510+E1) showed significant ($p < 0.05$) improvement of a decrease in the cerebral cortex volume as compared to the rTg4510 mouse group administered with a vehicle alone (rTg4510+Veh.), and did not
20 show a significant difference in the cerebral cortex volume as compared to the littermate control mouse group administered with a vehicle alone (Ctrl+Veh.).

[0099]

The above-mentioned results of Experimental Example 7
25 suggest that the composition of Example 1 of the present invention may suppress cerebral atrophy observed in Alzheimer-type dementia patients and the like.

[0100]

[Example 2] Composition for improving cognitive function

To afford the composition shown in Table 6, given amounts of respective components were weighed and mixed to prepare a composition for improving cognitive function of Example 2

5 (hereinafter sometimes to be referred to as "the composition of Example 2").

[0101]

[Table 6]

component	composition ratio relative to total content of amino acids (mol%)
L-leucine	36.05
L-lysine hydrochloride	18.43
L-valine	5.06
L-isoleucine	9.97
L-phenylalanine	26.05
L-histidine hydrochloride	3.92
L-tryptophan	0.52
total amount	100

10 [0102]

[Experimental Example 8] Study of effect of composition of Example 2 on cognitive function in human

A human clinical test was performed as follows to examine the effect of the composition of Example 2 on cognitive

15 function in humans.

(1) Preparation of test food

As shown in Table 7, granules containing the composition of Example 2 (3.0 g) (Active 1), granules containing the composition of Example 2 (1.5 g) (Active 2), and the
20 aforementioned granules containing dextrin, cornstarch, and lactose instead of the composition of Example 2 and not containing either amino acid or protein (Placebo 1) were prepared according to a conventional method.

[0103]

25 [Table 7]

test food	total amount (g)	component
Active 1	4.0	composition of Example 2 (3.0 g) + excipient (1.0 g)
Active 2	4.0	composition of Example 2 (1.5 g) + excipient (2.5 g)
Placebo 1	4.0	containing dextrin, cornstarch and lactose instead of composition of Example 2 and not containing either amino acid or protein

[0104]

(2) Test subject

The test subjects were 105 men and women at the age of 55
5 years or higher (54 males, 51 females, average age=64.3±5.0
years old) with normal cognitive function, that is, no
pathological abnormality such as dementia was observed, but the
cognitive function test by the Japanese version of the Mini
Mental State Examination (MMSE-J) was 26 points or more, and
10 the test by the Japanese version of the Montreal Cognitive
Assessment (MoCA-J) was less than 30 points, and they were
aware of forgetfulness, or pointed out to be forgetful by
others.

[0105]

15 (3) Clinical test

The subjects were randomly divided into 3 groups (test
food medium-dose group that ingests Active 1, test food low-
dose group that ingests Active 2, and control food group that
ingests Placebo 1), cognitive function test (Trail Making Test
20 B (TMT-B)) was performed before ingestion of each test food,
and the time to complete the task was measured and used as the
baseline value. Thereafter, the test foods shown in Table 7
were ingested when the subjects were hungry (2 packets a day,
one packet in the morning and one packet in the afternoon)
25 under double blind test. TMT-B was performed in the same
manner 12 weeks after the ingestion, and the time to complete
the task was measured.

[0106]

In the test subjects in each group of the above-mentioned clinical test, no significant difference was found among the 3 groups in sex, age, years of schooling, and the score of the Japanese version of the WHO-5 Well-Being Index (WHO-5-J) and each score of the Tokyo Metropolitan Institute of Gerontology Index of Competence, which relate to cognitive function, as shown in Table 8.

[0107]

[Table 8]

	medium-dose group	low-dose group	control group	<i>P</i>
	n=33	n=35	n=35	
sex; proportion of male (males)	16 (48.5)	18 (51.4)	18 (51.4)	n.s.
age (years old)	64.1 (5.9)	64.1 (5.0)	64.5 (4.2)	n.s.
years of schooling (years)	14.6 (2.4)	15.1 (1.8)	14.7 (2.2)	n.s.
score in WHO-5 Well-Being Index (points)	13.0 (4.0)	13.3 (3.9)	12.7 (3.0)	n.s.
Tokyo Metropolitan Institute of Gerontology Index of Competence				
score of instrumental ADL (points)	5.0 (0.0)	5.0 (0.0)	5.0 (0.0)	n.s.
score of intellectual ADL (points)	3.8 (0.5)	3.7 (0.6)	3.5 (0.7)	n.s.
score of social ADL (points)	3.7 (0.5)	3.3 (1.0)	3.4 (0.9)	n.s.

*; Numerical values in parentheses show standard deviation values.

[0108]

(4) Results

The amount of change in TMT-B in each group, that is, the value obtained by subtracting the baseline value from the value after 12 weeks was calculated and shown in Fig. 8.

TMT-B was developed as a test to evaluate frontal lobe dysfunction, in which test the time to draw lines to connect

numbers 1 to 13 and hiragana letters (Japanese syllabary letters) "A" to "Shi" alternating between numbers and hiragana letters in ascending order is measured. It is a test capable of particularly evaluating attention function, executive
5 function (flexibility) and the like. Accordingly, the results of this test can be interpreted as the evaluation results of whether or not one can perform work efficiently in daily life and the like. Therefore, as the amount of change in TMT-B turns into a more negative value, it means that the attention
10 function, executive function (flexibility) and the like have been improved more.

In Fig. 8, the mean value of TMT-B change (seconds) in the control food group (Placebo 1) is -2.44, and standard deviation is 25.77; the mean value of the test food low-dose
15 group (Active 2) is -0.44, and standard deviation is 23.49; and the mean value of the test food medium-dose group (Active 1) is -17.00, and standard deviation is 33.20. An unpaired t-test was performed between Placebo 1 and each group of Active 1 and Active 2.

20 As a result, no significant difference was found between Placebo 1 and Active 2, but a significant difference ($P=0.049$) was found between Placebo 1 and Active 1.

[0109]

From the above-mentioned results of Experimental Example
25 8, an intervention effect of the ingestion of the test food medium-dose was shown in, among the cognitive functions, attention function, executive function and the like in which the TMT-B test result becomes an index.

Therefore, it was suggested that the composition of the
30 present invention is effective in improving a decline in cognitive function that does not lead to pathological cognitive dysfunction such as dementia, but is observed in healthy humans, for example, reduction in attention, decline in thinking skills, decline in working memory (function of temporarily storing and
35 managing the information) and the like.

[Industrial Applicability]

[0110]

As described in detail above, a composition for improving cognitive function can be provided according to the present
5 invention.

The composition for improving cognitive function of the present invention effectively improves cognitive function and is effective for preventing or improving various symptoms or disorders caused by a decline in cognitive function.

10 In addition, the composition for improving cognitive function of the present invention also functions as a composition for improving anxiety-like symptoms and is effective for preventing or reducing the expression of anxiety-like symptoms that appear along with a decline in cognitive
15 function or before and after the decline of cognitive function.

Furthermore, the composition for improving cognitive function of the present invention also functions as a composition for suppressing cerebral atrophy and is effective for suppressing cerebral atrophy observed in neurodegenerative
20 diseases such as Alzheimer-type dementia and the like, and physiological cerebral atrophy that appears and progresses with aging, and improving a decline in cognitive function caused by such cerebral atrophy.

Particularly, the composition for improving cognitive
25 function of the present invention is highly safe and suitable for continuous ingestion or administration.

[0111]

This application is based on a patent application No. 2018-038950 filed in Japan, the contents of which are
30 incorporated by reference in full herein.

CLAIMS

1. A composition for improving cognitive function comprising one or more kinds selected from the group consisting of 25 mol%
5 - 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.
2. The composition according to claim 1, further comprising 10
10 mol% - 30 mol% of lysine relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.
3. The composition according to claim 1 or 2, further
15 comprising one or more kinds selected from the group consisting of valine, isoleucine, histidine and tryptophan.
4. The composition according to claim 3, comprising one or more kinds selected from the group consisting of 2 mol% - 7 mol% of
20 valine, 5 mol% - 15 mol% of isoleucine, 2 mol% - 10 mol% of histidine and 0.1 mol% - 2 mol% of tryptophan relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan.
- 25 5. The composition according to claim 3 or 4, comprising the following amino acids at the following contents relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan:
leucine 25 mol% - 45 mol%,
30 phenylalanine 20 mol% - 40 mol%,
lysine 10 mol% - 30 mol%,
valine 2 mol% - 7 mol%,
isoleucine 5 mol% - 15 mol%,
histidine 2 mol% - 10 mol%, and
35 tryptophan 0.1 mol% - 2 mol%.

6. The composition according to any one of claims 1 to 5,
wherein the composition is a pharmaceutical product.

5 7. The composition according to any one of claims 1 to 5,
wherein the composition is a food.

8. A composition for improving an anxiety-like symptom,
comprising one or more kinds selected from the group consisting
10 of 25 mol% - 45 mol% of leucine, and 20 mol% - 40 mol% of
phenylalanine relative to the total content of leucine, lysine,
valine, isoleucine, phenylalanine, histidine and tryptophan.

9. The composition according to claim 8, further comprising 10
15 mol% - 30 mol% of lysine relative to the total content of
leucine, lysine, valine, isoleucine, phenylalanine, histidine
and tryptophan.

10. The composition according to claim 8 or 9, further
20 comprising one or more kinds selected from the group consisting
of valine, isoleucine, histidine and tryptophan.

11. The composition according to claim 10, comprising one or
more kinds selected from the group consisting of 2 mol% - 7
25 mol% of valine, 5 mol% - 15 mol% of isoleucine, 2 mol% - 10
mol% of histidine and 0.1 mol% - 2 mol% of tryptophan relative
to the total content of leucine, lysine, valine, isoleucine,
phenylalanine, histidine and tryptophan.

30 12. The composition according to claim 10 or 11, comprising the
following amino acids at the following contents relative to the
total content of leucine, lysine, valine, isoleucine,
phenylalanine, histidine and tryptophan:

leucine 25 mol% - 45 mol%,
35 phenylalanine 20 mol% - 40 mol%,

lysine 10 mol% - 30 mol%,
valine 2 mol% - 7 mol%,
isoleucine 5 mol% - 15 mol%,
histidine 2 mol% - 10 mol%, and
5 tryptophan 0.1 mol% - 2 mol%.

13. The composition according to any one of claims 8 to 12,
wherein the composition is a pharmaceutical product.

10 14. The composition according to any one of claims 8 to 12,
wherein the composition is a food.

15 15. A composition for suppressing cerebral atrophy, comprising
one or more kinds selected from the group consisting of 25 mol%
- 45 mol% of leucine, and 20 mol% - 40 mol% of phenylalanine
relative to the total content of leucine, lysine, valine,
isoleucine, phenylalanine, histidine and tryptophan.

16. The composition according to claim 15, further comprising
20 10 mol% - 30 mol% of lysine relative to the total content of
leucine, lysine, valine, isoleucine, phenylalanine, histidine
and tryptophan.

17. The composition according to claim 15 or 16, further
25 comprising one or more kinds selected from the group consisting
of valine, isoleucine, histidine and tryptophan.

18. The composition according to claim 17, comprising one or
more kinds selected from the group consisting of 2 mol% - 7
30 mol% of valine, 5 mol% - 15 mol% of isoleucine, 2 mol% - 10
mol% of histidine and 0.1 mol% - 2 mol% of tryptophan relative
to the total content of leucine, lysine, valine, isoleucine,
phenylalanine, histidine and tryptophan.

35 19. The composition according to claim 17 or 18, comprising the

following amino acids at the following contents relative to the total content of leucine, lysine, valine, isoleucine, phenylalanine, histidine and tryptophan:

leucine 25 mol% - 45 mol%,
5 phenylalanine 20 mol% - 40 mol%,
lysine 10 mol% - 30 mol%,
valine 2 mol% - 7 mol%,
isoleucine 5 mol% - 15 mol%,
histidine 2 mol% - 10 mol%, and
10 tryptophan 0.1 mol% - 2 mol%.

20. The composition according to any one of claims 15 to 19, wherein the composition is a pharmaceutical product.

15 21. The composition according to any one of claims 15 to 19, wherein the composition is a food.