

ANTIBODY-CONJUGATED 8-SULFONYL-BENZAZEPINE COMPOUNDS AND THEIR USES

CROSS REFERENCE TO RELATED APPLICATIONS

- 5 This non-provisional application claims the benefit of priority to U.S. Provisional Application No. 63/308,275, filed 9 February 2022, which is incorporated by reference in its entirety.

SEQUENCE LISTING

- The instant application contains a Sequence Listing which has been submitted in XML
 10 format and is hereby incorporated by reference in its entirety. Said XML copy, created on 24 Jan 2023, is named 17019.023WO1-TW1 and is 13,070 bytes in size.

FIELD OF THE INVENTION


The invention relates generally to an immunoconjugate comprising an antibody conjugated to one or more 8-sulfonyl-benzazepine molecules.

15 BACKGROUND OF THE INVENTION

New compositions and methods for the delivery of antibodies and immune adjuvants are needed in order to reach inaccessible tumors and/or to expand treatment options for cancer patients and other subjects. The invention provides such compositions and methods.

SUMMARY OF THE INVENTION

- 20 The invention is generally directed to an immunoconjugate comprising an antibody selected from anti-PD-L1, anti-HER2, anti-CEA, and anti-TROP2, covalently attached by a linker to one or more 8-sulfonyl-benzazepine TLR agonist moieties having the formula:



- where one of R¹, R², R³ and R⁴ is attached to L. The various substituents are defined
 25 herein.

Another aspect of the invention is a method of preparing an immunoconjugate by conjugation of one or more 8-sulfonyl-benzazepine-linker compounds with an antibody selected from anti-PD-L1, anti-HER2, anti-CEA, and anti-TROP2.

Another aspect of the invention is a pharmaceutical composition comprising a therapeutically effective amount of an immunoconjugate comprising an antibody selected from anti-PD-L1, anti-HER2, anti-CEA, and anti-TROP2, covalently attached by a linker to one or more 8-sulfonyl-benzazepine moieties, and one or more pharmaceutically acceptable diluent, vehicle, carrier or excipient.

Another aspect of the invention is an 8-sulfonyl-benzazepine-linker compound.

Another aspect of the invention is a method for treating cancer comprising administering a therapeutically effective amount of an immunoconjugate comprising an antibody selected from anti-PD-L1, anti-HER2, anti-CEA, and anti-TROP2, covalently attached to one or more 8-sulfonyl-benzazepine moieties by a linker.

Another aspect of the invention is a use of an immunoconjugate comprising an antibody selected from anti-PD-L1, anti-HER2, anti-CEA, and anti-TROP2, covalently attached to one or more 8-sulfonyl-benzazepine moieties by a linker in the treatment of an illness, in particular cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a graph of secreted TNF α (Tumor Necrosis Factor alpha) cytokine levels after incubation of varying concentrations of immunoconjugate IC-3, comparator CIC-1, and naked antibody TROP2 with a co-culture of cancer cells with a cDC-enriched primary cell isolate.

Figure 2 shows a graph of secreted TNF α (Tumor Necrosis Factor alpha) cytokine levels after incubation of varying concentrations of immunoconjugate IC-3, comparator CIC-2, and naked antibody TROP2 with a co-culture of cancer cells with a cDC-enriched primary cell isolate.

DETAILED DESCRIPTION OF THE INVENTION

Reference will now be made in detail to certain embodiments of the invention, examples of which are illustrated in the accompanying structures and formulas. While the invention will be described in conjunction with the enumerated embodiments, it will be understood that they are not intended to limit the invention to those embodiments. On the contrary, the invention is intended to cover all alternatives, modifications, and equivalents, which may be included within the scope of the invention as defined by the claims.

One skilled in the art will recognize many methods and materials similar or equivalent to those described herein, which could be used in the practice of the present invention. The invention is in no way limited to the methods and materials described.

DEFINITIONS

5 The term “immunoconjugate” or “immune-stimulating antibody conjugate” refers to an antibody construct that is covalently bonded to an adjuvant moiety via a linker. the term “adjuvant” refers to a substance capable of eliciting an immune response in a subject exposed to the adjuvant.

 “Adjuvant moiety” refers to an adjuvant that is covalently bonded to an antibody
10 construct, e.g., through a linker, as described herein. The adjuvant moiety can elicit the immune response while bonded to the antibody construct or after cleavage (e.g., enzymatic cleavage) from the antibody construct following administration of an immunoconjugate to the subject.

 “Adjuvant” refers to a substance capable of eliciting an immune response in a subject exposed to the adjuvant.

15 The terms “Toll-like receptor” and “TLR” refer to any member of a family of highly-conserved mammalian proteins which recognizes pathogen-associated molecular patterns and acts as key signaling elements in innate immunity. TLR polypeptides share a characteristic structure that includes an extracellular domain that has leucine-rich repeats, a transmembrane domain, and an intracellular domain that is involved in TLR signaling.

20 The terms “Toll-like receptor 7” and “TLR7” refer to nucleic acids or polypeptides sharing at least about 70%, about 80%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or more sequence identity to a publicly-available TLR7 sequence, e.g., GenBank accession number AAZ99026 for human TLR7 polypeptide, or GenBank accession number AAK62676 for murine TLR7 polypeptide.

25 The terms “Toll-like receptor 8” and “TLR8” refer to nucleic acids or polypeptides sharing at least about 70%, about 80%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or more sequence identity to a publicly-available TLR7 sequence, e.g., GenBank accession number AAZ95441 for human TLR8 polypeptide, or GenBank accession number AAK62677 for murine TLR8 polypeptide.

30 A “TLR agonist” is a compound that binds, directly or indirectly, to a TLR (e.g., TLR7 and/or TLR8) to induce TLR signaling. Any detectable difference in TLR signaling can indicate that an agonist stimulates or activates a TLR. Signaling differences can be manifested, for example, as changes in the expression of target genes, in the phosphorylation of signal transduction components, in the intracellular localization of downstream elements such as

nuclear factor- κ B (NF- κ B), in the association of certain components (such as IL-1 receptor associated kinase (IRAK)) with other proteins or intracellular structures, or in the biochemical activity of components such as kinases (such as mitogen-activated protein kinase (MAPK)).

“Antibody” refers to a polypeptide comprising an antigen binding region (including the complementarity determining region (CDRs)) from an immunoglobulin gene or fragments thereof. The term “antibody” specifically encompasses monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments that exhibit the desired biological activity. An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 25 kDa) and one “heavy” chain (about 50-70 kDa) connected by disulfide bonds. Each chain is composed of structural domains, which are referred to as immunoglobulin domains. These domains are classified into different categories by size and function, e.g., variable domains or regions on the light and heavy chains (V_L and V_H , respectively) and constant domains or regions on the light and heavy chains (C_L and C_H , respectively). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids, referred to as the paratope, primarily responsible for antigen recognition, i.e., the antigen binding domain. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively. IgG antibodies are large molecules of about 150 kDa composed of four peptide chains. IgG antibodies contain two identical class γ heavy chains of about 50 kDa and two identical light chains of about 25 kDa, thus a tetrameric quaternary structure. The two heavy chains are linked to each other and to a light chain each by disulfide bonds. The resulting tetramer has two identical halves, which together form the Y-like shape. Each end of the fork contains an identical antigen binding domain. There are four IgG subclasses (IgG1, IgG2, IgG3, and IgG4) in humans, named in order of their abundance in serum (i.e., IgG1 is the most abundant). Typically, the antigen binding domain of an antibody will be most critical in specificity and affinity of binding to cancer cells.

“Bispecific” antibodies (bsAbs) are antibodies that bind two distinct epitopes to cancer (Suurs F.V. et al (2019) *Pharmacology & Therapeutics* 201: 103-119). Bispecific antibodies may engage immune cells to destroy tumor cells, deliver payloads to tumors, and/or block tumor signaling pathways. An antibody that targets a particular antigen includes a bispecific or multispecific antibody with at least one antigen binding region that targets the particular antigen. In some embodiments, the targeted monoclonal antibody is a bispecific antibody with at least one antigen binding region that targets tumor cells. Such antigens include but are not limited to:

mesothelin, prostate specific membrane antigen (PSMA), HER2, TROP2, CEA, EGFR, 5T4, Nectin4, CD19, CD20, CD22, CD30, CD70, B7H3, B7H4 (also known as 08E), protein tyrosine kinase 7 (PTK7), glypican-3, RGI, fucosyl-GMI, CTLA-4, and CD44 (WO 2017/196598).

In some embodiments, the antibody construct is an antigen-binding antibody “fragment,” which comprises at least an antigen-binding region of an antibody, alone or with other components that together constitute the antibody construct. Many different types of antibody “fragments” are known in the art, including, for instance, (i) a Fab fragment, which is a monovalent fragment consisting of the V_L , V_H , C_L , and CH_1 domains, (ii) a $F(ab')_2$ fragment, which is a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region, (iii) a Fv fragment consisting of the V_L and V_H domains of a single arm of an antibody, (iv) a Fab' fragment, which results from breaking the disulfide bridge of an $F(ab')_2$ fragment using mild reducing conditions, (v) a disulfide-stabilized Fv fragment (dsFv), and (vi) a single chain Fv (scFv), which is a monovalent molecule consisting of the two domains of the Fv fragment (i.e., V_L and V_H) joined by a synthetic linker which enables the two domains to be synthesized as a single polypeptide chain. In some embodiments, the antibody construct is an antibody or a fusion protein comprising (i) an antigen binding domain and (ii) an Fc domain.

The antibody or antibody fragment can be part of a larger construct, for example, a conjugate or fusion construct of the antibody fragment to additional regions. For instance, in some embodiments, the antibody fragment can be fused to an Fc region as described herein. In other embodiments, the antibody fragment (e.g., a Fab or scFv) can be part of a chimeric antigen receptor or chimeric T-cell receptor, for instance, by fusing to a transmembrane domain (optionally with an intervening linker or “stalk” (e.g., hinge region)) and optional intercellular signaling domain. For instance, the antibody fragment can be fused to the gamma and/or delta chains of a t-cell receptor, so as to provide a T-cell receptor like construct that binds PD-L1. In yet another embodiment, the antibody fragment is part of a bispecific T-cell engager (BiTEs) comprising a CD1 or CD3 binding domain and linker.

In some embodiments, the antibody construct comprises an Fc domain. In certain embodiments, the antibody construct is an antibody. In certain embodiments, the antibody construct is a fusion protein. The antigen binding domain can be a single-chain variable region fragment (scFv). A single-chain variable region fragment (scFv), which is a truncated Fab fragment including the variable (V) domain of an antibody heavy chain linked to a V domain of a light antibody chain via a synthetic peptide, can be generated using routine recombinant DNA technology techniques. Similarly, disulfide-stabilized variable region fragments (dsFv) can be prepared by recombinant DNA technology. The antibody construct or antigen binding domain may comprise one or more variable regions (e.g., two variable regions) of an antigen binding

domain of an anti-CEA antibody, each variable region comprising a CDR1, a CDR2, and a CDR3.

“Cysteine-mutant antibody” is an antibody in which one or more amino acid residues of an antibody are substituted with cysteine residues. A cysteine-mutant antibody may be prepared from the parent antibody by antibody engineering methods (Junutula, et al., (2008b) *Nature Biotech.*, 26(8):925-932; Doman et al. (2009) *Blood* 114(13):2721-2729; US 7521541; US 7723485; US 2012/0121615; WO 2009/052249). Cysteine residues provide for site-specific conjugation of an adjuvant such as a TLR agonist to the antibody through the reactive cysteine thiol groups at the engineered cysteine sites but do not perturb immunoglobulin folding and assembly or alter antigen binding and effector functions. Cysteine-mutant antibodies can be conjugated to the TLR agonist-linker compound with uniform stoichiometry of the immunoconjugate (e.g., up to two TLR agonist moieties per antibody in an antibody that has a single engineered, mutant cysteine site). The TLR agonist-linker compound has a reactive electrophilic group to react specifically with the free cysteine thiol groups of the cysteine-mutant antibody.

“Epitope” means any antigenic determinant or epitopic determinant of an antigen to which an antigen binding domain binds (i.e., at the paratope of the antigen binding domain). Antigenic determinants usually consist of chemically active surface groupings of molecules, such as amino acids or sugar side chains, and usually have specific three dimensional structural characteristics, as well as specific charge characteristics.

The terms “Fc receptor” or “FcR” refer to a receptor that binds to the Fc region of an antibody. There are three main classes of Fc receptors: (1) FcγR which bind to IgG, (2) FcαR which binds to IgA, and (3) FcεR which binds to IgE. The FcγR family includes several members, such as FcγI (CD64), FcγRIIA (CD32A), FcγRIIB (CD32B), FcγRIIIA (CD16A), and FcγRIIIB (CD16B). The Fcγ receptors differ in their affinity for IgG and also have different affinities for the IgG subclasses (e.g., IgG1, IgG2, IgG3, and IgG4).

Nucleic acid or amino acid sequence “identity,” as referenced herein, can be determined by comparing a nucleic acid or amino acid sequence of interest to a reference nucleic acid or amino acid sequence. The percent identity is the number of nucleotides or amino acid residues that are the same (i.e., that are identical) as between the optimally aligned sequence of interest and the reference sequence divided by the length of the longest sequence (i.e., the length of either the sequence of interest or the reference sequence, whichever is longer). Alignment of sequences and calculation of percent identity can be performed using available software programs. Examples of such programs include CLUSTAL-W, T-Coffee, and ALIGN (for alignment of nucleic acid and amino acid sequences), BLAST programs (e.g., BLAST 2.1,

BL2SEQ, BLASTp, BLASTn, and the like) and FASTA programs (e.g., FASTA3x, FASTM, and SSEARCH) (for sequence alignment and sequence similarity searches). Sequence alignment algorithms also are disclosed in, for example, Altschul et al., *J. Molecular Biol.*, 215(3): 403-410 (1990), Beigert et al., *Proc. Natl. Acad. Sci. USA*, 106(10): 3770-3775 (2009),
 5 Durbin et al., eds., *Biological Sequence Analysis: Probabilistic Models of Proteins and Nucleic Acids*, Cambridge University Press, Cambridge, UK (2009), Soding, *Bioinformatics*, 21(7): 951-960 (2005), Altschul et al., *Nucleic Acids Res.*, 25(17): 3389-3402 (1997), and Gusfield, *Algorithms on Strings, Trees and Sequences*, Cambridge University Press, Cambridge UK (1997)). Percent (%) identity of sequences can be also calculated, for example, as $100 \times$
 10 $[(\text{identical positions})/\min(\text{TGA}, \text{TGB})]$, where TGA and TGB are the sum of the number of residues and internal gap positions in peptide sequences A and B in the alignment that minimizes TGA and TGB. See, e.g., Russell et al., *J. Mol Biol.*, 244: 332-350 (1994).

The “antibody construct” or “binding agent” comprises Ig heavy and light chain variable region polypeptides that together form the antigen binding site. Each of the heavy and light
 15 chain variable regions are polypeptides comprising three complementarity determining regions (CDR1, CDR2, and CDR3) connected by framework regions. The antibody construct can be any of a variety of types of binding agents known in the art that comprise Ig heavy and light chains. For instance, the binding agent can be an antibody, an antigen-binding antibody “fragment,” or a T-cell receptor.

20 “Biosimilar” refers to an approved antibody construct that has active properties similar to, for example, a PD-L1-targeting antibody construct previously approved such as atezolizumab (TECENTRIQ™, Genentech, Inc.), durvalumab (IMFINZI™, AstraZeneca), and avelumab (BAVENCIO™, EMD Serono, Pfizer); a HER2-targeting antibody construct previously approved such as trastuzumab (HERCEPTIN™, Genentech, Inc.), and pertuzumab
 25 (PERJETA™, Genentech, Inc.); or a CEA-targeting antibody such as labetuzumab (CEA-CIDE™, MN-14, hMN14, Immunomedics) CAS Reg. No. 219649-07-7).

“Biobetter” refers to an approved antibody construct that is an improvement of a previously approved antibody construct, such as atezolizumab, durvalumab, avelumab, trastuzumab, pertuzumab, and labetuzumab. The biobetter can have one or more modifications
 30 (e.g., an altered glycan profile, or a unique epitope) over the previously approved antibody construct.

“Amino acid” refers to any monomeric unit that can be incorporated into a peptide, polypeptide, or protein. Amino acids include naturally-occurring α -amino acids and their stereoisomers, as well as unnatural (non-naturally occurring) amino acids and their
 35 stereoisomers. “Stereoisomers” of a given amino acid refer to isomers having the same

molecular formula and intramolecular bonds but different three-dimensional arrangements of bonds and atoms (e.g., an L-amino acid and the corresponding D-amino acid). The amino acids can be glycosylated (e.g., *N*-linked glycans, *O*-linked glycans, phosphoglycans, *C*-linked glycans, or glypication) or deglycosylated. Amino acids may be referred to herein by either the
5 commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission.

Naturally-occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline, γ -carboxyglutamate, and *O*-phosphoserine. Naturally-occurring α -amino acids include, without limitation, alanine (Ala),
10 cysteine (Cys), aspartic acid (Asp), glutamic acid (Glu), phenylalanine (Phe), glycine (Gly), histidine (His), isoleucine (Ile), arginine (Arg), lysine (Lys), leucine (Leu), methionine (Met), asparagine (Asn), proline (Pro), glutamine (Gln), serine (Ser), threonine (Thr), valine (Val), tryptophan (Trp), tyrosine (Tyr), and combinations thereof. Stereoisomers of naturally-occurring α -amino acids include, without limitation, D-alanine (D-Ala), D-cysteine (D-Cys),
15 D-aspartic acid (D-Asp), D-glutamic acid (D-Glu), D-phenylalanine (D-Phe), D-histidine (D-His), D-isoleucine (D-Ile), D-arginine (D-Arg), D-lysine (D-Lys), D-leucine (D-Leu), D-methionine (D-Met), D-asparagine (D-Asn), D-proline (D-Pro), D-glutamine (D-Gln), D-serine (D-Ser), D-threonine (D-Thr), D-valine (D-Val), D-tryptophan (D-Trp), D-tyrosine (D-Tyr), and combinations thereof.

20 Naturally-occurring amino acids include those formed in proteins by post-translational modification, such as citrulline (Cit).

Unnatural (non-naturally occurring) amino acids include, without limitation, amino acid analogs, amino acid mimetics, synthetic amino acids, *N*-substituted glycines, and *N*-methyl amino acids in either the L- or D-configuration that function in a manner similar to the naturally-
25 occurring amino acids. For example, "amino acid analogs" can be unnatural amino acids that have the same basic chemical structure as naturally-occurring amino acids (i.e., a carbon that is bonded to a hydrogen, a carboxyl group, an amino group) but have modified side-chain groups or modified peptide backbones, e.g., homoserine, norleucine, methionine sulfoxide, and methionine methyl sulfonium. "Amino acid mimetics" refer to chemical compounds that have a
30 structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally-occurring amino acid.

"Linker" refers to a functional group that covalently bonds two or more moieties in a compound or material. For example, the linking moiety can serve to covalently bond an adjuvant moiety to an antibody construct in an immunoconjugate.

“Linking moiety” refers to a functional group that covalently bonds two or more moieties in a compound or material. For example, the linking moiety can serve to covalently bond an adjuvant moiety to an antibody in an immunoconjugate. Useful bonds for connecting linking moieties to proteins and other materials include, but are not limited to, amides, amines, esters, carbamates, ureas, thioethers, thiocarbamates, thiocarbonates, and thioureas.

“Divalent” refers to a chemical moiety that contains two points of attachment for linking two functional groups; polyvalent linking moieties can have additional points of attachment for linking further functional groups. Divalent radicals may be denoted with the suffix “diyl”. For example, divalent linking moieties include divalent polymer moieties such as divalent poly(ethylene glycol), divalent cycloalkyl, divalent heterocycloalkyl, divalent aryl, and divalent heteroaryl group. A “divalent cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group” refers to a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl group having two points of attachment for covalently linking two moieties in a molecule or material. Cycloalkyl, heterocycloalkyl, aryl, or heteroaryl groups can be substituted or unsubstituted. Cycloalkyl, heterocycloalkyl, aryl, or heteroaryl groups can be substituted with one or more groups selected from halo, hydroxy, amino, alkylamino, amido, acyl, nitro, cyano, and alkoxy.

A wavy line (“”) represents a point of attachment of the specified chemical moiety. If the specified chemical moiety has two wavy lines (“”) present, it will be understood that the chemical moiety can be used bilaterally, i.e., as read from left to right or from right to left. In some embodiments, a specified moiety having two wavy lines (“”) present is considered to be used as read from left to right.

“Alkyl” refers to a straight (linear) or branched, saturated, aliphatic radical having the number of carbon atoms indicated. Alkyl can include any number of carbons, for example from one to twelve. Examples of alkyl groups include, but are not limited to, methyl (Me, -CH₃), ethyl (Et, -CH₂CH₃), 1-propyl (n-Pr, n-propyl, -CH₂CH₂CH₃), 2-propyl (i-Pr, i-propyl, -CH(CH₃)₂), 1-butyl (n-Bu, n-butyl, -CH₂CH₂CH₂CH₃), 2-methyl-1-propyl (i-Bu, i-butyl, -CH₂CH(CH₃)₂), 2-butyl (s-Bu, s-butyl, -CH(CH₃)CH₂CH₃), 2-methyl-2-propyl (t-Bu, t-butyl, -C(CH₃)₃), 1-pentyl (n-pentyl, -CH₂CH₂CH₂CH₂CH₃), 2-pentyl (-CH(CH₃)CH₂CH₂CH₃), 3-pentyl (-CH(CH₂CH₃)₂), 2-methyl-2-butyl (-C(CH₃)₂CH₂CH₃), 3-methyl-2-butyl (-CH(CH₃)CH(CH₃)₂), 3-methyl-1-butyl (-CH₂CH₂CH(CH₃)₂), 2-methyl-1-butyl (-CH₂CH(CH₃)CH₂CH₃), 1-hexyl (-CH₂CH₂CH₂CH₂CH₂CH₃), 2-hexyl (-CH(CH₃)CH₂CH₂CH₂CH₃), 3-hexyl (-CH(CH₂CH₃)(CH₂CH₂CH₃)), 2-methyl-2-pentyl (-C(CH₃)₂CH₂CH₂CH₃), 3-methyl-2-pentyl (-CH(CH₃)CH(CH₃)CH₂CH₃), 4-methyl-2-pentyl (-CH(CH₃)CH₂CH(CH₃)₂), 3-methyl-3-pentyl (-C(CH₃)(CH₂CH₃)₂), 2-methyl-3-pentyl (-CH(CH₂CH₃)CH(CH₃)₂), 2,3-dimethyl-2-butyl (-C(CH₃)₂CH(CH₃)₂), 3,3-dimethyl-2-butyl (-CH(CH₃)C(CH₃)₃), 1-heptyl, 1-octyl, and the like.

Alkyl groups can be substituted or unsubstituted. “Substituted alkyl” groups can be substituted with one or more groups selected from halo, hydroxy, amino, oxo ($=O$), alkylamino, amido, acyl, nitro, cyano, and alkoxy.

The term “alkyldiyl” refers to a divalent alkyl radical. Examples of alkyldiyl groups include, but are not limited to, methylene ($-CH_2-$), ethylene ($-CH_2CH_2-$), propylene ($-CH_2CH_2CH_2-$), and the like. An alkyldiyl group may also be referred to as an “alkylene” group.

“Alkenyl” refers to a straight (linear) or branched, unsaturated, aliphatic radical having the number of carbon atoms indicated and at least one carbon-carbon double bond, *sp*². Alkenyl can include from two to about 12 or more carbons atoms. Alkenyl groups are radicals having “cis” and “trans” orientations, or alternatively, “E” and “Z” orientations. Examples include, but are not limited to, ethylenyl or vinyl ($-CH=CH_2$), allyl ($-CH_2CH=CH_2$), butenyl, pentenyl, and isomers thereof. Alkenyl groups can be substituted or unsubstituted. “Substituted alkenyl” groups can be substituted with one or more groups selected from halo, hydroxy, amino, oxo ($=O$), alkylamino, amido, acyl, nitro, cyano, and alkoxy.

The terms “alkenylene” or “alkenyldiyl” refer to a linear or branched-chain divalent hydrocarbon radical. Examples include, but are not limited to, ethylenylene or vinylene ($-CH=CH-$), allyl ($-CH_2CH=CH-$), and the like.

“Alkynyl” refers to a straight (linear) or branched, unsaturated, aliphatic radical having the number of carbon atoms indicated and at least one carbon-carbon triple bond, *sp*. Alkynyl can include from two to about 12 or more carbons atoms. For example, C₂-C₆ alkynyl includes, but is not limited to ethynyl ($-C\equiv CH$), propynyl (propargyl, $-CH_2C\equiv CH$), butynyl, pentynyl, hexynyl, and isomers thereof. Alkynyl groups can be substituted or unsubstituted. “Substituted alkynyl” groups can be substituted with one or more groups selected from halo, hydroxy, amino, oxo ($=O$), alkylamino, amido, acyl, nitro, cyano, and alkoxy.

The term “alkynylene” or “alkynyldiyl” refer to a divalent alkynyl radical.

The terms “carbocycle”, “carbocyclyl”, “carbocyclic ring” and “cycloalkyl” refer to a saturated or partially unsaturated, monocyclic, fused bicyclic, or bridged polycyclic ring assembly containing from 3 to 12 ring atoms, or the number of atoms indicated. Saturated monocyclic carbocyclic rings include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cyclooctyl. Saturated bicyclic and polycyclic carbocyclic rings include, for example, norbornane, [2.2.2] bicyclooctane, decahydronaphthalene and adamantane. Carbocyclic groups can also be partially unsaturated, having one or more double or triple bonds in the ring. Representative carbocyclic groups that are partially unsaturated include, but are not limited to, cyclobutene, cyclopentene, cyclohexene, cyclohexadiene (1,3- and 1,4-isomers),

cycloheptene, cycloheptadiene, cyclooctene, cyclooctadiene (1,3-, 1,4- and 1,5-isomers), norbornene, and norbornadiene.

The term “cycloalkyldiyl” refers to a divalent cycloalkyl radical.

“Aryl” refers to a monovalent aromatic hydrocarbon radical of 6-20 carbon atoms (C₆–C₂₀) derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. Aryl groups can be monocyclic, fused to form bicyclic or tricyclic groups, or linked by a bond to form a biaryl group. Representative aryl groups include phenyl, naphthyl and biphenyl. Other aryl groups include benzyl, having a methylene linking group. Some aryl groups have from 6 to 12 ring members, such as phenyl, naphthyl or biphenyl. Other aryl groups have from 6 to 10 ring members, such as phenyl or naphthyl.

The terms “arylene” or “aryldiyl” mean a divalent aromatic hydrocarbon radical of 6-20 carbon atoms (C₆–C₂₀) derived by the removal of two hydrogen atom from a two carbon atoms of a parent aromatic ring system. Some aryldiyl groups are represented in the exemplary structures as “Ar”. Aryldiyl includes bicyclic radicals comprising an aromatic ring fused to a saturated, partially unsaturated ring, or aromatic carbocyclic ring. Typical aryldiyl groups include, but are not limited to, radicals derived from benzene (phenyldiyl), substituted benzenes, naphthalene, anthracene, biphenylene, indenylene, indanylene, 1,2-dihydronaphthalene, 1,2,3,4-tetrahydronaphthyl, and the like. Aryldiyl groups are also referred to as “arylene”, and are optionally substituted with one or more substituents described herein.

The terms “heterocycle,” “heterocyclyl” and “heterocyclic ring” are used interchangeably herein and refer to a saturated or a partially unsaturated (i.e., having one or more double and/or triple bonds within the ring) carbocyclic radical of 3 to about 20 ring atoms in which at least one ring atom is a heteroatom selected from nitrogen, oxygen, phosphorus and sulfur, the remaining ring atoms being C, where one or more ring atoms is optionally substituted independently with one or more substituents described below. A heterocycle may be a monocycle having 3 to 7 ring members (2 to 6 carbon atoms and 1 to 4 heteroatoms selected from N, O, P, and S) or a bicycle having 7 to 10 ring members (4 to 9 carbon atoms and 1 to 6 heteroatoms selected from N, O, P, and S), for example: a bicyclo [4,5], [5,5], [5,6], or [6,6] system. Heterocycles are described in Paquette, Leo A.; “Principles of Modern Heterocyclic Chemistry” (W.A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; “The Chemistry of Heterocyclic Compounds, A series of Monographs” (John Wiley & Sons, New York, 1950 to present), in particular Volumes 13, 14, 16, 19, and 28; and J. Am. Chem. Soc. (1960) 82:5566. “Heterocyclyl” also includes radicals where heterocycle radicals are fused with a saturated, partially unsaturated ring, or aromatic carbocyclic or heterocyclic ring. Examples of heterocyclic rings include, but are not limited to, morpholin-4-yl, piperidin-1-yl, piperazinyl,

piperazin-4-yl-2-one, piperazin-4-yl-3-one, pyrrolidin-1-yl, thiomorpholin-4-yl, S-dioxothiomorpholin-4-yl, azocan-1-yl, azetidin-1-yl, octahydropyrido[1,2-a]pyrazin-2-yl, [1,4]diazepan-1-yl, pyrrolidinyl, tetrahydrofuran-yl, dihydrofuran-yl, tetrahydrothienyl, tetrahydropyran-yl, dihydropyran-yl, tetrahydrothiopyran-yl, piperidino, morpholino, thiomorpholino, thioxanyl, piperazinyl, homopiperazinyl, azetidiny, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepinyl, diazepinyl, thiazepinyl, 2-pyrrolinyl, 3-pyrrolinyl, indolinyl, 2H-pyran-yl, 4H-pyran-yl, dioxanyl, 1,3-dioxolan-yl, pyrazolinyl, dithianyl, dithiolan-yl, dihydropyran-yl, dihydrothienyl, dihydrofuran-yl, pyrazolidinylimidazoliny, imidazolidinyl, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl, azabicyclo[2.2.2]hexanyl, 3H-indolyl quinoliziny and N-pyridyl ureas. Spiro heterocycl

moieties are also included within the scope of this definition. Examples of spiro heterocycl moieties include azaspiro[2.5]octanyl and azaspiro[2.4]heptanyl. Examples of a heterocyclic group wherein 2 ring atoms are substituted with oxo (=O) moieties are pyrimidinonyl and 1,1-dioxo-thiomorpholinyl. The heterocycle groups herein are optionally substituted independently with one or more substituents described herein.

The term "heterocyclyldiyl" refers to a divalent, saturated or a partially unsaturated (i.e., having one or more double and/or triple bonds within the ring) carbocyclic radical of 3 to about 20 ring atoms in which at least one ring atom is a heteroatom selected from nitrogen, oxygen, phosphorus and sulfur, the remaining ring atoms being C, where one or more ring atoms is optionally substituted independently with one or more substituents as described. Examples of 5-membered and 6-membered heterocyclyldiyls include morpholinyl, piperidinyl, piperazinyl, pyrrolidinyl, dioxanyl, thiomorpholinyl, and S-dioxothiomorpholinyl.

The term "heteroaryl" refers to a monovalent aromatic radical of 5-, 6-, or 7-membered rings, and includes fused ring systems (at least one of which is aromatic) of 5-20 atoms, containing one or more heteroatoms independently selected from nitrogen, oxygen, and sulfur. Examples of heteroaryl groups are pyridinyl (including, for example, 2-hydroxypyridinyl), imidazolyl, imidazopyridinyl, pyrimidinyl (including, for example, 4-hydroxypyrimidinyl), pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxadiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, indolyl, benzimidazolyl, benzofuran-yl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, and furopyridinyl. Heteroaryl groups are optionally substituted independently with one or more substituents described herein.

The term “heteroarylidiyl” refers to a divalent aromatic radical of 5-, 6-, or 7-membered rings, and includes fused ring systems (at least one of which is aromatic) of 5-20 atoms, containing one or more heteroatoms independently selected from nitrogen, oxygen, and sulfur. Examples of 5-membered and 6-membered heteroarylidiyls include pyridyldiyl, imidazolylidiyl, 5 pyrimidinylidiyl, pyrazolylidiyl, triazolylidiyl, pyrazinylidiyl, tetrazolylidiyl, furyldiyl, thienylidiyl, isoxazolylidiyl, thiazolylidiyl, oxadiazolylidiyl, oxazolylidiyl, isothiazolylidiyl, and pyrrolyldiyl.

The heterocycle or heteroaryl groups may be carbon (carbon-linked), or nitrogen (nitrogen-linked) bonded where such is possible. By way of example and not limitation, carbon 10 bonded heterocycles or heteroaryls are bonded at position 2, 3, 4, 5, or 6 of a pyridine, position 3, 4, 5, or 6 of a pyridazine, position 2, 4, 5, or 6 of a pyrimidine, position 2, 3, 5, or 6 of a pyrazine, position 2, 3, 4, or 5 of a furan, tetrahydrofuran, thiofuran, thiophene, pyrrole or tetrahydropyrrole, position 2, 4, or 5 of an oxazole, imidazole or thiazole, position 3, 4, or 5 of an isoxazole, pyrazole, or isothiazole, position 2 or 3 of an aziridine, position 2, 3, or 4 of an 15 azetidine, position 2, 3, 4, 5, 6, 7, or 8 of a quinoline or position 1, 3, 4, 5, 6, 7, or 8 of an isoquinoline.

By way of example and not limitation, nitrogen bonded heterocycles or heteroaryls are bonded at position 1 of an aziridine, azetidine, pyrrole, pyrrolidine, 2-pyrroline, 3-pyrroline, imidazole, imidazolidine, 2-imidazoline, 3-imidazoline, pyrazole, pyrazoline, 2-pyrazoline, 3- 20 pyrazoline, piperidine, piperazine, indole, indoline, 1H-indazole, position 2 of a isoindole, or isoindoline, position 4 of a morpholine, and position 9 of a carbazole, or β -carboline.

The terms “halo” and “halogen,” by themselves or as part of another substituent, refer to a fluorine, chlorine, bromine, or iodine atom.

The term “carbonyl,” by itself or as part of another substituent, refers to C(=O) or – 25 C(=O)–, i.e., a carbon atom double-bonded to oxygen and bound to two other groups in the moiety having the carbonyl.

As used herein, the phrase “quaternary ammonium salt” refers to a tertiary amine that has been quaternized with an alkyl substituent (e.g., a C₁-C₄ alkyl such as methyl, ethyl, propyl, or butyl).

30 The terms “treat,” “treatment,” and “treating” refer to any indicia of success in the treatment or amelioration of an injury, pathology, condition (e.g., cancer), or symptom (e.g., cognitive impairment), including any objective or subjective parameter such as abatement; remission; diminishing of symptoms or making the symptom, injury, pathology, or condition more tolerable to the patient; reduction in the rate of symptom progression; decreasing the 35 frequency or duration of the symptom or condition; or, in some situations, preventing the onset

of the symptom. The treatment or amelioration of symptoms can be based on any objective or subjective parameter, including, for example, the result of a physical examination.

The terms “cancer,” “neoplasm,” and “tumor” are used herein to refer to cells which exhibit autonomous, unregulated growth, such that the cells exhibit an aberrant growth phenotype characterized by a significant loss of control over cell proliferation. Cells of interest for detection, analysis, and/or treatment in the context of the invention include cancer cells (e.g., cancer cells from an individual with cancer), malignant cancer cells, pre-metastatic cancer cells, metastatic cancer cells, and non-metastatic cancer cells. Cancers of virtually every tissue are known. The phrase “cancer burden” refers to the quantum of cancer cells or cancer volume in a subject. Reducing cancer burden accordingly refers to reducing the number of cancer cells or the cancer cell volume in a subject. The term “cancer cell” as used herein refers to any cell that is a cancer cell (e.g., from any of the cancers for which an individual can be treated, e.g., isolated from an individual having cancer) or is derived from a cancer cell, e.g., clone of a cancer cell. For example, a cancer cell can be from an established cancer cell line, can be a primary cell isolated from an individual with cancer, can be a progeny cell from a primary cell isolated from an individual with cancer, and the like. In some embodiments, the term can also refer to a portion of a cancer cell, such as a sub-cellular portion, a cell membrane portion, or a cell lysate of a cancer cell. Many types of cancers are known to those of skill in the art, including solid tumors such as carcinomas, sarcomas, glioblastomas, melanomas, lymphomas, and myelomas, and circulating cancers such as leukemias.

As used herein, the term “cancer” includes any form of cancer, including but not limited to, solid tumor cancers (e.g., skin, lung, prostate, breast, gastric, bladder, colon, ovarian, pancreas, kidney, liver, glioblastoma, medulloblastoma, leiomyosarcoma, head & neck squamous cell carcinomas, melanomas, and neuroendocrine) and liquid cancers (e.g., hematological cancers); carcinomas; soft tissue tumors; sarcomas; teratomas; melanomas; leukemias; lymphomas; and brain cancers, including minimal residual disease, and including both primary and metastatic tumors.

“PD-L1 expression” refers to a cell that has a PD-L1 receptor on the cell’s surface. As used herein “PD-L1 overexpression” refers to a cell that has more PD-L1 receptors as compared to corresponding non-cancer cell.

“HER2” refers to the protein human epidermal growth factor receptor 2.

“HER2 expression” refers to a cell that has a HER2 receptor on the cell’s surface. For example, a cell may have from about 20,000 to about 50,000 HER2 receptors on the cell’s surface. As used herein “HER2 overexpression” refers to a cell that has more than about 50,000 HER2 receptors. For example, a cell 2, 5, 10, 100, 1,000, 10,000, 100,000, or 1,000,000 times

the number of HER2 receptors as compared to corresponding non-cancer cell (e.g., about 1 or 2 million HER2 receptors). It is estimated that HER2 is overexpressed in about 25% to about 30% of breast cancers.

5 The “pathology” of cancer includes all phenomena that compromise the well-being of the patient. This includes, without limitation, abnormal or uncontrollable cell growth, metastasis, interference with the normal functioning of neighboring cells, release of cytokines or other secretory products at abnormal levels, suppression or aggravation of inflammatory or immunological response, neoplasia, premalignancy, malignancy, and invasion of surrounding or distant tissues or organs, such as lymph nodes.

10 As used herein, the phrases “cancer recurrence” and “tumor recurrence,” and grammatical variants thereof, refer to further growth of neoplastic or cancerous cells after diagnosis of cancer. Particularly, recurrence may occur when further cancerous cell growth occurs in the cancerous tissue. “Tumor spread,” similarly, occurs when the cells of a tumor disseminate into local or distant tissues and organs, therefore, tumor spread encompasses tumor
15 metastasis. “Tumor invasion” occurs when the tumor growth spread out locally to compromise the function of involved tissues by compression, destruction, or prevention of normal organ function.

As used herein, the term “metastasis” refers to the growth of a cancerous tumor in an organ or body part, which is not directly connected to the organ of the original cancerous tumor.
20 Metastasis will be understood to include micrometastasis, which is the presence of an undetectable amount of cancerous cells in an organ or body part that is not directly connected to the organ of the original cancerous tumor. Metastasis can also be defined as several steps of a process, such as the departure of cancer cells from an original tumor site, and migration and/or invasion of cancer cells to other parts of the body.

25 The phrases “effective amount” and “therapeutically effective amount” refer to a dose or amount of a substance such as an immunoconjugate that produces therapeutic effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques (see, e.g., Lieberman, *Pharmaceutical Dosage Forms* (vols. 1-3, 1992); Lloyd, *The Art, Science and Technology of
30 Pharmaceutical Compounding* (1999); Pickar, *Dosage Calculations* (1999); Goodman & Gilman’s *The Pharmacological Basis of Therapeutics*, 11th Edition (McGraw-Hill, 2006); and Remington: *The Science and Practice of Pharmacy*, 22nd Edition, (Pharmaceutical Press, London, 2012)). In the case of cancer, the therapeutically effective amount of the immunoconjugate may reduce the number of cancer cells; reduce the tumor size; inhibit (i.e.,
35 slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit

(i.e., slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the cancer. To the extent the immunoconjugate may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic. For cancer therapy, efficacy can, for example, be measured by assessing the time to disease progression (TTP) and/or determining the response rate (RR)

5 “Recipient,” “individual,” “subject,” “host,” and “patient” are used interchangeably and refer to any mammalian subject for whom diagnosis, treatment, or therapy is desired (e.g., humans). “Mammal” for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, sheep, goats, pigs, camels, etc. In certain embodiments, the mammal is human.

 The phrase “synergistic adjuvant” or “synergistic combination” in the context of this invention includes the combination of two immune modulators such as a receptor agonist, cytokine, and adjuvant polypeptide, that in combination elicit a synergistic effect on immunity relative to either administered alone. Particularly, the immunoconjugates disclosed herein comprise synergistic combinations of the claimed adjuvant and antibody construct. These synergistic combinations upon administration elicit a greater effect on immunity, e.g., relative to when the antibody construct or adjuvant is administered in the absence of the other moiety. Further, a decreased amount of the immunoconjugate may be administered (as measured by the total number of antibody constructs or the total number of adjuvants administered as part of the immunoconjugate) compared to when either the antibody construct or adjuvant is administered alone.

 As used herein, the term “administering” refers to parenteral, intravenous, intraperitoneal, intramuscular, intratumoral, intralesional, intranasal, or subcutaneous administration, oral administration, administration as a suppository, topical contact, intrathecal administration, or the implantation of a slow-release device, e.g., a mini-osmotic pump, to the subject.

 The terms “about” and “around,” as used herein to modify a numerical value, indicate a close range surrounding the numerical value. Thus, if “X” is the value, “about X” or “around X” indicates a value of from 0.9X to 1.1X, e.g., from 0.95X to 1.05X or from 0.99X to 1.01X. A reference to “about X” or “around X” specifically indicates at least the values X, 0.95X, 0.96X, 0.97X, 0.98X, 0.99X, 1.01X, 1.02X, 1.03X, 1.04X, and 1.05X. Accordingly, “about X” and “around X” are intended to teach and provide written description support for a claim limitation of, e.g., “0.98X.”

ANTIBODIES

In an exemplary embodiment, the immunoconjugates of the invention comprise an antibody construct that comprises an antigen binding domain that specifically recognizes and binds PD-L1.

5 Programmed Death-Ligand 1 (PD-L1, cluster of differentiation 274, CD274, B7-homolog 1, or B7-H1) belongs to the B7 protein superfamily, and is a ligand of programmed cell death protein 1 (PD-1, PDCD1, cluster of differentiation 279, or CD279). PD-L1 can also interact with B7.1 (CD80) and such interaction is believed to inhibit T cell priming. The PD-L1/PD-1 axis plays a large role in suppressing the adaptive immune response. More
10 specifically, it is believed that engagement of PD-L1 with its receptor, PD-1, delivers a signal that inhibits activation and proliferation of T-cells. Agents that bind to PD-L1 and prevent the ligand from binding to the PD-1 receptor prevent this immunosuppression, and can, therefore, enhance an immune response when desired, such as for the treatment of cancers, or infections. PD-L1/PD-1 pathway also contributes to preventing autoimmunity and therefore agonistic
15 agents against PD-L1 or agents that deliver immune inhibitory payloads may help treatment of autoimmune disorders.

Several antibodies targeting PD-L1 have been developed for the treatment of cancer, including atezolizumab (TECENTRIQ™), durvalumab (IMFINZI™), and avelumab (BAVENCIO™). Nevertheless, there continues to be a need for new PD-L1 antibody
20 constructs, including agents that bind PD-L1 with high affinity and effectively prevent PD-L1/PD-1 signaling and agents that can deliver therapeutic payloads to PD-L1 expressing cells. In addition, there is a need for new PD-L1-binding agents to treat autoimmune disorders and infections.

A method is provided of delivering a TLR agonist payload to a cell expressing PD-L1
25 comprising administering to the cell, or mammal comprising the cell, an immunoconjugate comprising an anti-PD-L1 antibody covalently attached to a linker which is covalently attached to one or more TLR agonist moieties.

Also provided is a method for enhancing or reducing or inhibiting an immune response in a mammal, and a method for treating a disease, disorder, or condition in a mammal that is
30 responsive to PD-L1 inhibition, which methods comprise administering a PD-L1 immunoconjugate thereof, to the mammal.

The invention provides a PD-L1 antibody comprising an immunoglobulin heavy chain variable region polypeptide and an immunoglobulin light chain variable region polypeptide. The PD-L1 antibody specifically binds PD-L1. The binding specificity of the antibody allows for
35 targeting PD-L1 expressing cells, for instance, to deliver therapeutic payloads to such cells. In

some embodiments, the PD-L1 antibody binds to human PD-L1. However, antibodies that bind to any PD-L1 fragment, homolog or paralog also are encompassed.

In some embodiments, the PD-L1 antibody binds PD-L1 without substantially inhibiting or preventing PD-L1 from binding to its receptor, PD-1. However, in other embodiments, the PD-L1 antibody can completely or partially block (inhibit or prevent) binding of PD-L1 to its receptor, PD-1, such that the antibody can be used to inhibit PD-L1/PD-1 signaling (e.g., for therapeutic purposes). The antibody or antigen-binding antibody fragment can be monospecific for PD-L1, or can be bispecific or multi-specific. For instance, in bivalent or multivalent antibodies or antibody fragments, the binding domains can be different targeting different epitopes of the same antigen or targeting different antigens. Methods of constructing multivalent binding constructs are known in the art. Bispecific and multispecific antibodies are known in the art. Furthermore, a diabody, triabody, or tetrabody can be provided, which is a dimer, trimer, or tetramer of polypeptide chains each comprising a V_H connected to a V_L by a peptide linker that is too short to allow pairing between the V_H and V_L on the same polypeptide chain, thereby driving the pairing between the complementary domains on different V_H - V_L polypeptide chains to generate a multimeric molecule having two, three, or four functional antigen binding sites. Also, bis-scFv fragments, which are small scFv fragments with two different variable domains can be generated to produce bispecific bis-scFv fragments capable of binding two different epitopes. Fab dimers (Fab2) and Fab trimers (Fab3) can be produced using genetic engineering methods to create multispecific constructs based on Fab fragments.

The PD-L1 antibody can be, or can be obtained from, a human antibody, a non-human antibody, a humanized antibody, or a chimeric antibody, or corresponding antibody fragments. A "chimeric" antibody is an antibody or fragment thereof typically comprising human constant regions and non-human variable regions. A "humanized" antibody is a monoclonal antibody typically comprising a human antibody scaffold but with non-human origin amino acids or sequences in at least one CDR (e.g., 1, 2, 3, 4, 5, or all six CDRs).

The PD-L1 antibody can be internalizing, as described in WO 2021/150701 and incorporated by reference herein, or the PD-L1 antibody can be non-internalizing, as described in WO 2021/150702 and incorporated by reference herein.

In an exemplary embodiment, the immunoconjugates of the invention comprise an antibody construct that comprises an antigen binding domain that specifically recognizes and binds HER2.

A number of anti-HER2 monoclonal antibodies are approved and under clinical development (Costa, RLB et al (2020) *Breast Cancer* 6(10):1-11.

In certain embodiments, immunoconjugates of the invention comprise an anti-HER2 antibody such as those prepared by the methods of Example 201. In one embodiment of the invention, an anti-HER2 antibody of an immunoconjugate of the invention comprises a humanized anti-HER2 antibody, e.g., huMAb4D5-1, huMAb4D5-2, huMAb4D5-3, huMAb4D5-4, huMAb4D5-5, huMAb4D5-6, huMAb4D5-7 and huMAb4D5-8, as described in Table 3 of US 5821337, which is specifically incorporated by reference herein. Those antibodies contain human framework regions with the complementarity-determining regions of a murine antibody (4D5) that binds to HER2. The humanized antibody huMAb4D5-8 is also referred to as trastuzumab, commercially available under the tradename HERCEPTIN™ (Genentech, Inc.).

Trastuzumab (CAS 180288-69-1, huMAb4D5-8, rhuMAb HER2, HERCEPTIN®, Genentech, Inc.) is a recombinant DNA-derived, IgG1 kappa, monoclonal antibody that is a humanized version of a murine anti-HER2 antibody (4D5) that selectively binds with high affinity in a cell-based assay ($K_d = 5$ nM) to the extracellular domain of HER2 (US 5677171; US 5821337; US 6054297; US 6165464; US 6339142; US 6407213; US 6639055; US 6719971; US 6800738; US 7074404; Coussens et al (1985) *Science* 230:1132-9; Slamon et al (1989) *Science* 244:707-12; Slamon et al (2001) *New Engl. J. Med.* 344:783-792).

In an embodiment of the invention, the antibody construct or antigen binding domain comprises the CDR regions of trastuzumab. In an embodiment of the invention, the anti-HER2 antibody further comprises the framework regions of the trastuzumab. In an embodiment of the invention, the anti-HER2 antibody further comprises one or both variable regions of trastuzumab.

In another embodiment of the invention, an anti-HER2 antibody of an immunoconjugate of the invention comprises a humanized anti-HER2 antibody, e.g., humanized 2C4, as described in US 7862817. An exemplary humanized 2C4 antibody is pertuzumab (CAS Reg. No. 380610-27-5), PERJETA™ (Genentech, Inc.). Pertuzumab is a HER dimerization inhibitor (HDI) and functions to inhibit the ability of HER2 to form active heterodimers or homodimers with other HER receptors (such as EGFR/HER1, HER2, HER3 and HER4). See, for example, Harari and Yarden, *Oncogene* 19:6102-14 (2000); Yarden and Sliwkowski. *Nat Rev Mol Cell Biol* 2:127-37 (2001); Sliwkowski *Nat Struct Biol* 10:158-9 (2003); Cho et al. *Nature* 421:756-60 (2003); and Malik et al. *Pro Am Soc Cancer Res* 44:176-7 (2003). PERJETA™ is approved for the treatment of breast cancer.

In an embodiment of the invention, the antibody construct or antigen binding domain comprises the CDR regions of pertuzumab. In an embodiment of the invention, the anti-HER2 antibody further comprises the framework regions of the pertuzumab. In an embodiment of the

invention, the anti-HER2 antibody further comprises one or both variable regions of pertuzumab.

Margetuximab (MGAH22, MARGENZA™, MacroGenics, Inc.), CAS Reg. No. 1350624-75-7, is an FDA-approved anti-HER2 monoclonal antibody. The Fc region of
5 margetuximab is optimized for increased binding to the activating Fc gamma Rs but decreased binding to the inhibitory Fc gamma Rs on immune effector cells (Nordstrom, JL, et al (2011) *Breast Cancer Res.* 13(6):R123; Rugo, HS, et al (2021) *JAMA Oncol.*;7(4):573-584; Markham, A. (2021) *Drugs* 81:599-604). Margetuximab is approved by the FDA for treatment of patients with relapsed or refractory advanced breast cancer whose tumors express HER2 at the 2+ level
10 by immunohistochemistry and lack evidence of HER2 gene amplification by FISH.

HT-19 is another anti-HER2 monoclonal antibody that binds to an epitope in human HER2 distinct from the epitope of trastuzumab or pertuzumab. HT-19 was shown to inhibit HER2 signaling comparable to trastuzumab and enhance HER2 degradation in combination with trastuzumab and pertuzumab. XMT-1522 is an antibody-drug conjugate comprising the HT-19
15 antibody (Bergstrom D. A. et al., (2015) *Cancer Res.*; 75:LB-231).

In an exemplary embodiment, the immunoconjugates of the invention comprise an antibody construct that comprises an antigen binding domain that specifically recognizes and binds CEA. Carcinoembryonic antigen-related cell adhesion molecule 5 (CEACAM5) also known as CD66e (Cluster of Differentiation 66e), is a member of the carcinoembryonic antigen
20 (CEA) gene family.

Elevated expression of carcinoembryonic antigen (CEA, CD66e, CEACAM5) has been implicated in various biological aspects of neoplasia, especially tumor cell adhesion, metastasis, the blocking of cellular immune mechanisms, and having antiapoptosis functions. CEA is also used as a blood marker for many carcinomas. Labetuzumab (CEA-CIDE™, Immunomedics,
25 CAS Reg. No. 219649-07-7), also known as MN-14 and hMN14, is a humanized IgG1 monoclonal antibody and has been studied for the treatment of colorectal cancer (Blumenthal, R. et al (2005) *Cancer Immunology Immunotherapy* 54(4):315-327). Labetuzumab conjugated to a camptothecin analog (labetuzumab govitecan, IMMU-130) targets carcinoembryonic antigen-related cell adhesion mol. 5 (CEACAM5) and is being studied in patients with relapsed or
30 refractory metastatic colorectal cancer (Sharkey, R. et al, (2018), *Molecular Cancer Therapeutics* 17(1):196-203; Cardillo, T. et al (2018) *Molecular Cancer Therapeutics* 17(1):150-160). In an embodiment of the invention, the CEA-targeting antibody construct or antigen binding domain comprises the Variable light chain (VL kappa) of hMN-14/labetuzumab as disclosed in US 6676924, which is incorporated by reference herein for this purpose.

In an exemplary embodiment, the immunoconjugates of the invention comprise an antibody construct that comprises an antigen binding domain that specifically recognizes and binds TROP2. Tumor-associated calcium signal transducer 2 (TROP-2) is a transmembrane glycoprotein encoded by the TACSTD2 gene (Linnenbach AJ, et al (1993) *Mol Cell Biol.* 13(3): 1507-15; Calabrese G, et al (2001) *Cytogenet Cell Genet.* 92(1-2): 164-5). TROP2 is an intracellular calcium signal transducer that is differentially expressed in many cancers and signals cells for self-renewal, proliferation, invasion, and survival. TROP2 is considered a stem cell marker and is expressed in many normal tissues, though in contrast, it is overexpressed in many cancers (Ohmachi T, et al., (2006) *Clin. Cancer Res.*, 12(10), 3057-3063; Muhlmann G, et al., (2009) *J. Clin. Pathol.*, 62(2), 152-158; Fong D, et al., (2008) *Br. J. Cancer*, 99(8), 1290-1295; Fong D, et al., (2008) *Mod. Pathol.*, 21(2), 186-191; Ning S, et al., (2013) *Neurol. Sci.*, 34(10), 1745-1750). Overexpression of TROP2 is of prognostic significance. Several ligands have been proposed that interact with TROP2. TROP2 signals the cells via different pathways and it is transcriptionally regulated by a complex network of several transcription factors.

Human TROP2 (TACSTD2: tumor-associated calcium signal transducer 2, GA733-1, EGP-1, MIS1; hereinafter, referred to as hTROP2) is a single-pass transmembrane type 1 cell membrane protein consisting of 323 amino acid residues. While the presence of a cell membrane protein involved in immune resistance, which is common to human trophoblasts and cancer cells (Faulk W P, et al., *Proc. Natl. Acad. Sci.* 75(4):1947-1951 (1978)), has previously been suggested, an antigen molecule recognized by a monoclonal antibody against a cell membrane protein in a human choriocarcinoma cell line was identified and designated as TROP2 as one of the molecules expressed in human trophoblasts (Lipinski M, et al., *Proc. Natl. Acad. Sci.* 78(8), 5147-5150 (1981)). This molecule was also designated as tumor antigen GA733-1 recognized by a mouse monoclonal antibody GA733 (Linnenbach A J, et al., *Proc. Natl. Acad. Sci.* 86(1), 27-31 (1989)) obtained by immunization with a gastric cancer cell line or an epithelial glycoprotein (EGP-1; Basu A, et al., *Int. J. Cancer*, 62 (4), 472-479 (1995)) recognized by a mouse monoclonal antibody RS7-3G11 obtained by immunization with non-small cell lung cancer cells. In 1995, however, the TROP2 gene was cloned, and all of these molecules were confirmed to be identical molecules (Fornaro M, et al., *Int. J. Cancer*, 62(5), 610-618 (1995)). The DNA sequence and amino acid sequence of hTROP2 are available on a public database and can be referred to, for example, under Accession Nos. NM_002353 and NP_002344 (NCBI).

In response to such information suggesting the association with cancer, a plurality of anti-hTROP2 antibodies have been established so far and studied for their antitumor effects. Among these antibodies, there is disclosed, for example, an unconjugated antibody that exhibits in itself antitumor activity in nude mouse xenograft models (WO 2008/144891; WO

2011/145744; WO 2011/155579; WO 2013/077458) as well as an antibody that exhibits antitumor activity as ADC with a cytotoxic drug (WO 2003/074566; WO 2011/068845; WO 2013/068946; US 7999083). However, the strength or coverage of their activity is still insufficient, and there are unsatisfied medical needs for hTROP2 as a therapeutic target.

- 5 TROP2 expression in cancer cells has been correlated with drug resistance. Several strategies target TROP2 on cancer cells that include antibodies, antibody fusion proteins, chemical inhibitors, nanoparticles, etc. The in vitro studies and pre-clinical studies, using these various therapeutic treatments, have resulted in significant inhibition of tumor cell growth both in vitro and in vivo in mice. Clinical studies have explored the potential application of TROP2
10 as both a prognostic biomarker and as a therapeutic target to reverse resistance.

- Sacituzumab govitecan (TRODELVY®, Immunomedics, IMMU-132), an antibody-drug conjugate comprising a TROP2-directed antibody linked to a topoisomerase inhibitor drug, is indicated for the treatment of metastatic triple-negative breast cancer (mTNBC) in adult patients that have received at least two prior therapies. The TROP2 antibody in sacituzumab govitecan is
15 conjugated to SN-38, the active metabolite of irinotecan (US 2016/0297890; WO 2015/098099).

In an embodiment of the invention, the TROP2-targeting antibody construct or antigen binding domain comprises the light chain CDR (complementarity determining region) of hRS7 (humanized RS7), (US 7238785, incorporated by reference herein).

- In an embodiment of the invention, the TROP2-targeting antibody construct or antigen
20 binding domain comprises the light chain CDR (complementarity determining region) or light chain framework (LFR) sequences selected from SEQ ID NO. 1-7.

Region	Sequence	Residues (Kabat)	Length	SEQ ID NO.
LFR1	DIQMTQSPSSLSASVGDRVTITC	1-23	23	1
CDR-L1	KASQDVSTAVA	24-34	11	2
LFR2	WYQQKPGKAPKLLIY	35-49	15	3
CDR-L2	SASYRYT	50-56	7	4
LFR3	GVPSRFSGSGSGTDFTLTISSLQPEDFAVYYC	57-88	32	5
CDR-L3	QQHYITPLT	89-97	9	6
LFR4	FGQGTKLEIK	98-107	10	7

In an embodiment of the invention, the TROP2-targeting antibody construct or antigen binding domain comprises the heavy chain CDR (complementarity determining region) or heavy chain framework (HFR) sequences selected from SEQ ID NO. 8-14.

Region	Sequence	Residues (Kabat)	Length	SEQ ID NO.
HFR1	QVQLVQSGAEVKKPGASVKVSKASGYTFT	1-30	30	8
CDR-H1	TAGMQ	31-35	5	9
HFR2	WVRQAPGQGLEWMG	36-49	14	10
CDR-H2	WINTHSGVPKYAEDFKG	50-66	17	11
HFR3	RVTISADTSTSTAYLQLSSLKSEDTAVYYCAR	67-98	32	12
CDR-H3	SGFGSSYWYFDV	99-110	12	13
HFR4	WGQGTILVTVSS	111-121	11	14

5

In some embodiments, the antibodies in the immunoconjugates contain a modified Fc region, wherein the modification modulates the binding of the Fc region to one or more Fc receptors.

In some embodiments, the Fc region is modified by inclusion of a transforming growth factor beta 1 (TGF β 1) receptor, or a fragment thereof, that is capable of binding TGF β 1. For example, the receptor can be TGF β receptor II (TGF β RII). In some embodiments, the TGF β receptor is a human TGF β receptor. In some embodiments, the IgG has a C-terminal fusion to a TGF β RII extracellular domain (ECD) as described in US 9676863, incorporated herein. An "Fc linker" may be used to attach the IgG to the TGF β RII extracellular domain. The Fc linker may be a short, flexible peptide that allows for the proper three-dimensional folding of the molecule while maintaining the binding-specificity to the targets. In some embodiments, the N-terminus of the TGF β receptor is fused to the Fc of the antibody construct (with or without an Fc linker). In some embodiments, the C-terminus of the antibody construct heavy chain is fused to the TGF β receptor (with or without an Fc linker). In some embodiments, the C-terminal lysine residue of the antibody construct heavy chain is mutated to alanine.

In some embodiments, the antibodies in the immunoconjugates are glycosylated.

In some embodiments, the antibody in the immunoconjugates is a cysteine-engineered antibody which provides for site-specific conjugation of an adjuvant, label, or drug moiety to the antibody through cysteine substitutions at sites where the engineered cysteines are available for conjugation but do not perturb immunoglobulin folding and assembly or alter antigen binding

and effector functions (Junutula, et al., 2008b *Nature Biotech.*, 26(8):925-932; Dorman et al. (2009) *Blood* 114(13):2721-2729; US 7521541; US 7723485; US 2012/0121615; WO 2009/052249). Cysteine-engineered antibodies can be conjugated to the 8-sulfonyl-2-aminobenzazepine adjuvant moiety as an 8-sulfonyl-2-aminobenzazepine-linker compound with uniform stoichiometry (e.g., up to two 8-sulfonyl-2-aminobenzazepine moieties per antibody in an antibody that has a single engineered cysteine site).

8-SULFONYL-2-AMINO BENZAZEPINE ADJUVANT COMPOUNDS



The immunoconjugate of the invention comprises an 8-sulfonyl-2-aminobenzazepine adjuvant moiety. The adjuvant moiety described herein is a compound that elicits an immune response (i.e., an immunostimulatory agent). Generally, the adjuvant moiety described herein is a TLR agonist. TLRs are type-I transmembrane proteins that are responsible for the initiation of innate immune responses in vertebrates. TLRs recognize a variety of pathogen-associated molecular patterns from bacteria, viruses, and fungi and act as a first line of defense against invading pathogens. TLRs elicit overlapping yet distinct biological responses due to differences in cellular expression and in the signaling pathways that they initiate. Once engaged (e.g., by a natural stimulus or a synthetic TLR agonist), TLRs initiate a signal transduction cascade leading to activation of nuclear factor- κ B (NF- κ B) via the adapter protein myeloid differentiation primary response gene 88 (MyD88) and recruitment of the IL-1 receptor associated kinase (IRAK). Phosphorylation of IRAK then leads to recruitment of TNF-receptor associated factor 6 (TRAF6), which results in the phosphorylation of the NF- κ B inhibitor I- κ B. As a result, NF- κ B enters the cell nucleus and initiates transcription of genes whose promoters contain NF- κ B binding sites, such as cytokines. Additional modes of regulation for TLR signaling include TIR-domain containing adapter-inducing interferon- β (TRIF)-dependent induction of TNF-receptor associated factor 6 (TRAF6) and activation of MyD88 independent pathways via TRIF and TRAF3, leading to the phosphorylation of interferon response factor three (IRF3). Similarly, the MyD88 dependent pathway also activates several IRF family members, including IRF5 and IRF7 whereas the TRIF dependent pathway also activates the NF- κ B pathway.






Typically, the adjuvant moiety described herein is a TLR7 and/or TLR8 agonist. TLR7 and TLR8 are both expressed in monocytes and dendritic cells. In humans, TLR7 is also expressed in plasmacytoid dendritic cells (pDCs) and B cells. TLR8 is expressed mostly in cells of myeloid origin, i.e., monocytes, granulocytes, and myeloid dendritic cells. TLR7 and TLR8 are capable of detecting the presence of "foreign" single-stranded RNA within a cell, as a means to respond to viral invasion. Treatment of TLR8-expressing cells, with TLR8 agonists can result in production of high levels of IL-12, IFN- γ , IL-1, TNF- α , IL-6, and other inflammatory






cytokines. Similarly, stimulation of TLR7-expressing cells, such as pDCs, with TLR7 agonists can result in production of high levels of IFN- α and other inflammatory cytokines. TLR7/TLR8 engagement and resulting cytokine production can activate dendritic cells and other antigen-presenting cells, driving diverse innate and acquired immune response mechanisms leading to tumor destruction.






Exemplary 8-sulfonyl-2-aminobenzazepine compounds (8SO2Bz) of the invention were synthesized, purified, and characterized by mass spectrometry and shown to have the expected mass. Additional experimental procedures are found in the Examples. Activity against HEK293 NFKB reporter cells expressing human TLR7 or human TLR8 was measured according to Example 202. Certain 8-sulfonyl-2-aminobenzazepine compounds demonstrate the surprising and unexpected property of TLR8 agonist selectivity which may predict useful therapeutic activity to treat cancer and other disorders.

Table 1: 8-Sulfonyl-2-aminobenzazepine compounds (8SO2Bz)

8SO2Bz No.	Structure	MW	HEK293 hTLR7 EC50 (nM)	HEK293 hTLR8 EC50 (nM)
8SO2Bz-1		485.6	>9000	1056
8SO2Bz-2		509.6	395	<2.3

8SO2Bz-3		380.5	>5000	202
8SO2Bz-4		394.5		
8SO2Bz-5		552.7		
8SO2Bz-6		564.7		
8SO2Bz-7		484.5		

8SO2Bz-8		440.5		
8SO2Bz-9		460.6		
8SO2Bz-10		424.5		
8SO2Bz-11		438.5		
8SO2Bz-12		483.5		

8SO2Bz-13		482.6		
8SO2Bz-14		481.6		
8SO2Bz-15		438.5		
8SO2Bz-16		440.5		
8SO2Bz-17		440.5		

8-SULFONYL-2-AMINOBENZAZEPINE-LINKER COMPOUNDS


The immunoconjugates of the invention are prepared by conjugation of an antibody with a 8-sulfonyl-2-aminobenzazepine-linker compound, 8SO2Bz-L. The 8-sulfonyl-2-aminobenzazepine-linker compounds comprise a 8-sulfonyl-2-aminobenzazepine (8SO2Bz) moiety covalently attached to a linker unit. The linker units comprise functional groups and subunits which affect stability, permeability, solubility, and other pharmacokinetic, safety, and efficacy properties of the immunoconjugates. The linker unit comprises a polyethyleneoxy (PEG) group. The linker unit includes a reactive functional group which reacts, i.e. conjugates, with a reactive functional group of the antibody. For example, a nucleophilic group such as a lysine side chain amino of the antibody reacts with an electrophilic reactive functional group of the 8SO2Bz-L compound to form the immunoconjugate. Also, for example, a cysteine thiol of the antibody reacts with a maleimide or bromoacetamide group of the 8SO2Bz-L linker compound to form the immunoconjugate.

Reactive electrophilic functional groups (Q in Formula II) suitable for the 8SO2Bz-L linker compounds include, but are not limited to, N-hydroxysuccinimidyl (NHS) esters and N-hydroxysulfosuccinimidyl (sulfo-NHS) esters (amine reactive); carbodiimides (amine and carboxyl reactive); hydroxymethyl phosphines (amine reactive); maleimides (thiol reactive); halogenated acetamides such as N-iodoacetamides (thiol reactive); aryl azides (primary amine reactive); fluorinated aryl azides (reactive via carbon-hydrogen (C-H) insertion); pentafluorophenyl (PFP) esters (amine reactive); tetrafluorophenyl (TFP) esters (amine reactive); imidoesters (amine reactive); isocyanates (hydroxyl reactive); vinyl sulfones (thiol, amine, and hydroxyl reactive); pyridyl disulfides (thiol reactive); and benzophenone derivatives (reactive via C-H bond insertion). Further reagents include, but are not limited, to those described in Hermanson, *Bioconjugate Techniques* 2nd Edition, Academic Press, 2008.

The invention provides solutions to the limitations and challenges to the design, preparation and use of immunoconjugates. Some linkers may be labile in the blood stream, thereby releasing unacceptable amounts of the adjuvant/drug prior to internalization in a target cell (Khot, A. et al (2015) *Bioanalysis* 7(13):1633–1648). Other linkers may provide stability in the bloodstream, but intracellular release effectiveness may be negatively impacted. Linkers that provide for desired intracellular release typically have poor stability in the bloodstream. Alternatively stated, bloodstream stability and intracellular release are typically inversely related. In addition, in standard conjugation processes, the amount of adjuvant/drug moiety loaded on the antibody, i.e. drug loading, the amount of aggregate that is formed in the conjugation reaction, and the yield of final purified conjugate that can be obtained are interrelated. For example, aggregate formation is generally positively correlated to the number

of equivalents of adjuvant/drug moiety and derivatives thereof conjugated to the antibody. Under high drug loading, formed aggregates must be removed for therapeutic applications. As a result, drug loading-mediated aggregate formation decreases immunoconjugate yield and can render process scale-up difficult.

- 5 Exemplary embodiments include a 8-sulfonyl-2-aminobenzazepine-linker compound of Formula II:



wherein

- R^1 , R^2 , R^3 , and R^4 are independently selected from the group consisting of H, C₁-C₁₂ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₁₂ carbocyclyl, C₆-C₂₀ aryl, C₂-C₉ heterocyclyl, and C₁-C₂₀ heteroaryl, where alkyl, alkenyl, alkynyl, carbocyclyl, aryl, heterocyclyl, and heteroaryl are independently and optionally substituted with one or more groups selected from:

- (C₁-C₁₂ alkylidyl)-N(R⁵)-*;
- (C₁-C₁₂ alkylidyl)-N(R⁵)₂;
- 15 -(C₁-C₁₂ alkylidyl)-OR⁵;
- (C₃-C₁₂ carbocyclyl);
- (C₃-C₁₂ carbocyclyl)-*;
- (C₃-C₁₂ carbocyclyl)-(C₁-C₁₂ alkylidyl)-NR⁵-*;
- (C₃-C₁₂ carbocyclyl)-(C₁-C₁₂ alkylidyl)-N(R⁵)₂;
- 20 -(C₃-C₁₂ carbocyclyl)-NR⁵-C(=NR⁵)NR⁵-*;
- (C₆-C₂₀ aryl);
- (C₆-C₂₀ arylidyl)-*;
- (C₆-C₂₀ arylidyl)-N(R⁵)-*;
- (C₆-C₂₀ arylidyl)-(C₁-C₁₂ alkylidyl)-N(R⁵)-*;
- 25 -(C₆-C₂₀ arylidyl)-(C₁-C₁₂ alkylidyl)-(C₂-C₂₀ heterocyclidyl)-*;
- (C₆-C₂₀ arylidyl)-(C₁-C₁₂ alkylidyl)-N(R⁵)₂;
- (C₆-C₂₀ arylidyl)-(C₁-C₁₂ alkylidyl)-NR⁵-C(=NR^{5a})N(R⁵)-*;
- (C₂-C₂₀ heterocyclyl);
- (C₂-C₂₀ heterocyclyl)-*;
- 30 -(C₂-C₉ heterocyclyl)-(C₁-C₁₂ alkylidyl)-NR⁵-*;

- $-(C_2-C_9 \text{ heterocyclyl})-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)_2$;
 $-(C_2-C_9 \text{ heterocyclyl})-C(=O)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
 $-(C_2-C_9 \text{ heterocyclyl})-NR^5-C(=NR^{5a})NR^5-*$;
 $-(C_2-C_9 \text{ heterocyclyl})-NR^5-(C_6-C_{20} \text{ arylldiyl})-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
5 $-(C_2-C_9 \text{ heterocyclyl})-(C_6-C_{20} \text{ arylldiyl})-*$;
 $-(C_1-C_{20} \text{ heteroaryl})$;
 $-(C_1-C_{20} \text{ heteroarylldiyl})-*$;
 $-(C_1-C_{20} \text{ heteroaryl})-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
 $-(C_1-C_{20} \text{ heteroaryl})-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)_2$;
10 $-(C_1-C_{20} \text{ heteroaryl})-NR^5-C(=NR^{5a})N(R^5)-*$;
 $-(C_1-C_{20} \text{ heteroaryl})-N(R^5)C(=O)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
 $-C(=O)-*$;
 $-C(=O)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
 $-C(=O)-(C_2-C_{20} \text{ heterocyclldiyl})-*$;
15 $-C(=O)N(R^5)_2$;
 $-C(=O)N(R^5)-*$;
 $-C(=O)N(R^5)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)C(=O)R^5$;
 $-C(=O)N(R^5)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)C(=O)N(R^5)_2$;
 $-C(=O)NR^5-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)CO_2R^5$;
20 $-C(=O)NR^5-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)C(=NR^{5a})N(R^5)_2$;
 $-C(=O)NR^5-(C_1-C_{12} \text{ alkylldiyl})-NR^5C(=NR^{5a})R^5$;
 $-C(=O)NR^5-(C_1-C_8 \text{ alkylldiyl})-NR^5(C_2-C_5 \text{ heteroaryl})$;
 $-C(=O)NR^5-(C_1-C_{20} \text{ heteroarylldiyl})-N(R^5)-*$;
 $-C(=O)NR^5-(C_1-C_{20} \text{ heteroarylldiyl})-*$;
25 $-C(=O)NR^5-(C_1-C_{20} \text{ heteroarylldiyl})-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)_2$;
 $-C(=O)NR^5-(C_1-C_{20} \text{ heteroarylldiyl})-(C_2-C_{20} \text{ heterocyclldiyl})-C(=O)NR^5-(C_1-C_{12} \text{ alkylldiyl})-NR^5-*$;
 $-N(R^5)_2$;
 $-N(R^5)-*$;
30 $-N(R^5)C(=O)R^5$;
 $-N(R^5)C(=O)-*$;
 $-N(R^5)C(=O)N(R^5)_2$;
 $-N(R^5)C(=O)N(R^5)-*$;


- $-\text{N}(\text{R}^5)\text{CO}_2\text{R}^5$;
 $-\text{NR}^5\text{C}(=\text{NR}^{5a})\text{N}(\text{R}^5)_2$;
 $-\text{NR}^5\text{C}(=\text{NR}^{5a})\text{N}(\text{R}^5)-*$;
 $-\text{NR}^5\text{C}(=\text{NR}^{5a})\text{R}^5$;
5 $-\text{N}(\text{R}^5)\text{C}(=\text{O})-(\text{C}_1\text{-C}_{12} \text{ alkylldiyl})-\text{N}(\text{R}^5)-*$;
 $-\text{N}(\text{R}^5)-(\text{C}_2\text{-C}_5 \text{ heteroaryl})$;
 $-\text{N}(\text{R}^5)-\text{S}(=\text{O})_2-(\text{C}_1\text{-C}_{12} \text{ alkyl})$;
 $-\text{O}-(\text{C}_1\text{-C}_{12} \text{ alkyl})$;
 $-\text{O}-(\text{C}_1\text{-C}_{12} \text{ alkylldiyl})-\text{N}(\text{R}^5)_2$;
10 $-\text{O}-(\text{C}_1\text{-C}_{12} \text{ alkylldiyl})-\text{N}(\text{R}^5)-*$;
 $-\text{O}-\text{C}(=\text{O})\text{N}(\text{R}^5)_2$;
 $-\text{O}-\text{C}(=\text{O})\text{N}(\text{R}^5)-*$;
 $-\text{O}-(\text{R}^5)-*$;
 $-\text{OR}^5$;
15 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20} \text{ heterocyclldiyl})-*$;
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20} \text{ heterocyclldiyl})-(\text{C}_1\text{-C}_{12} \text{ alkylldiyl})-\text{N}(\text{R}^5)_2$;
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20} \text{ heterocyclldiyl})-(\text{C}_1\text{-C}_{12} \text{ alkylldiyl})-\text{NR}^5-*$; and
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20} \text{ heterocyclldiyl})-(\text{C}_1\text{-C}_{12} \text{ alkylldiyl})-\text{OH}$;
or R^2 and R^3 together form a 5- or 6-membered heterocycll ring;
20 $\text{X}^1, \text{X}^2, \text{X}^3$, and X^4 are independently selected from the group consisting of a bond,
 $\text{C}(=\text{O}), \text{C}(=\text{O})\text{N}(\text{R}^5), \text{O}, \text{N}(\text{R}^5), \text{S}, \text{S}(\text{O})_2$, and $\text{S}(\text{O})_2\text{N}(\text{R}^5)$;
 R^5 is independently selected from the group consisting of H, $\text{C}_6\text{-C}_{20}$ aryl, $\text{C}_3\text{-C}_{12}$
carbocycll, $\text{C}_6\text{-C}_{20}$ arylldiyl, $\text{C}_1\text{-C}_{12}$ alkyl, and $\text{C}_1\text{-C}_{12}$ alkylldiyl, or two R^5 groups together form
a 5- or 6-membered heterocycll ring;
25 R^{5a} is selected from the group consisting of $\text{C}_6\text{-C}_{20}$ aryl and $\text{C}_1\text{-C}_{20}$ heteroaryl;
where the asterisk * indicates the attachment site of L, and where one of $\text{R}^1, \text{R}^2, \text{R}^3$ and
 R^4 is attached to L;
L is the linker selected from the group consisting of:
 $\text{Q}-\text{C}(=\text{O})-\text{PEG}-$;
30 $\text{Q}-\text{C}(=\text{O})-\text{PEG}-\text{C}(=\text{O})\text{N}(\text{R}^6)-(\text{C}_1\text{-C}_{12} \text{ alkylldiyl})-\text{C}(=\text{O})-\text{Gluc}-$;
 $\text{Q}-\text{C}(=\text{O})-\text{PEG}-\text{O}-$;
 $\text{Q}-\text{C}(=\text{O})-\text{PEG}-\text{O}-\text{C}(=\text{O})-$;
 $\text{Q}-\text{C}(=\text{O})-\text{PEG}-\text{C}(=\text{O})-$;

- $Q-C(=O)-PEG-C(=O)-PEP-$;
 $Q-C(=O)-PEG-N(R^6)-$;
 $Q-C(=O)-PEG-N(R^6)-C(=O)-$;
 $Q-C(=O)-PEG-N(R^6)-PEG-C(=O)-PEP-$;
5 $Q-C(=O)-PEG-N^1(R^6)_2-PEG-C(=O)-PEP-$;
 $Q-C(=O)-PEG-C(=O)-PEP-N(R^6)-(C_1-C_{12} \text{ alkylidyl})-$;
 $Q-C(=O)-PEG-C(=O)-PEP-N(R^6)-(C_1-C_{12} \text{ alkylidyl})N(R^6)C(=O)-(C_2-C_5$
 $\text{monoheterocyclidyl})-$;
 $Q-C(=O)-PEG-SS-(C_1-C_{12} \text{ alkylidyl})-OC(=O)-$;
10 $Q-C(=O)-PEG-SS-(C_1-C_{12} \text{ alkylidyl})-C(=O)-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-C(=O)N(R^6)-(C_1-C_{12} \text{ alkylidyl})-C(=O)-Gluc-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-O-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-O-C(=O)-$;
15 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-C(=O)-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-N(R^5)-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-N(PEG-CO_2H)-PEG-N(R^5)-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-C(=O)N(PEG-CO_2H)-PEG-N(R^5)-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-N(R^5)-C(=O)-$;
20 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-N(PEG-CO_2H)-PEG-C(=O)-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-C(=O)N(PEG-CO_2H)-PEG-C(=O)-$;
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-C(=O)-PEP-$; and
 $Q-(CH_2)_m-C(=O)N(R^6)-PEG-SS-(C_1-C_{12} \text{ alkylidyl})-OC(=O)-$;


R^6 is independently H or C_1-C_6 alkyl;

- 25 PEG has the formula: $-(CH_2CH_2O)_n-(CH_2)_m-$; m is an integer from 1 to 5, and n is an integer from 2 to 50;

$Gluc$ has the formula:




PEP has the formula:



- where AA is independently selected from a natural or unnatural amino acid side chain, or one or more of AA, and an adjacent nitrogen atom form a 5-membered ring proline amino acid, and the wavy line indicates a point of attachment;

Cyc is selected from C₆-C₂₀ arylidyl and C₁-C₂₀ heteroarylidyl, optionally substituted with one or more groups selected from F, Cl, NO₂, -OH, -OCH₃, and a glucuronic acid having the structure:



- R⁷ is selected from the group consisting of -CH(R⁸)O-, -CH₂-, -CH₂N(R⁸)-, and -CH(R⁸)O-C(=O)-, where R⁸ is selected from H, C₁-C₆ alkyl, C(=O)-C₁-C₆ alkyl, and -C(=O)N(R⁹)₂, where R⁹ is independently selected from the group consisting of H, C₁-C₁₂ alkyl, and -(CH₂CH₂O)_n-(CH₂)_m-OH, where m is an integer from 1 to 5, and n is an integer from 2 to 50, or two R⁹ groups together form a 5- or 6-membered heterocyclyl ring;

- y is an integer from 2 to 12;


z is 0 or 1;

Q is selected from the group consisting of N-hydroxysuccinimidyl, N-hydroxysulfosuccinimidyl, maleimide, and phenoxy substituted with one or more groups independently selected from F, Cl, NO₂, and SO₃⁻; and

- alkyl, alkylidyl, alkenyl, alkenyldiyl, alkynyl, alkynyldiyl, aryl, arylidyl, carbocyclyl, carbocyclidyl, heterocyclyl, heterocyclidyl, heteroaryl, and heteroaryldiyl are independently and optionally substituted with one or more groups independently selected from F, Cl, Br, I, -CN, -CH₃, -CH₂CH₃, -CH=CH₂, -C≡CH, -C≡CCH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH(CH₃)₂, -CH₂OH, -CH₂OCH₃, -CH₂CH₂OH, -C(CH₃)₂OH, -CH(OH)CH(CH₃)₂, -C(CH₃)₂CH₂OH, -CH₂CH₂SO₂CH₃, -CH₂OP(O)(OH)₂, -CH₂F, -CHF₂, -CF₃, -CH₂CF₃, -CH₂CHF₂, -CH(CH₃)CN, -C(CH₃)₂CN, -CH₂CN, -CH₂NH₂, -CH₂NHSO₂CH₃, -CH₂NHCH₃, -CH₂N(CH₃)₂, -CO₂H, -COCH₃, -CO₂CH₃, -CO₂C(CH₃)₃, -COCH(OH)CH₃, -CONH₂, -

- CONHCH₃, -CON(CH₃)₂, -C(CH₃)₂CONH₂, -NH₂, -NHCH₃, -N(CH₃)₂, -NHCOCH₃, -N(CH₃)COCH₃, -NHS(O)₂CH₃, -N(CH₃)C(CH₃)₂CONH₂, -N(CH₃)CH₂CH₂S(O)₂CH₃, -NHC(=NH)H, -NHC(=NH)CH₃, -NHC(=NH)NH₂, -NHC(=O)NH₂, -NO₂, =O, -OH, -OCH₃, -OCH₂CH₃, -OCH₂CH₂OCH₃, -OCH₂CH₂OH, -OCH₂CH₂N(CH₃)₂, -O(CH₂CH₂O)_n-
- 5 (CH₂)_mCO₂H, -O(CH₂CH₂O)_nH, -OCH₂F, -OCHF₂, -OCF₃, -OP(O)(OH)₂, -S(O)₂N(CH₃)₂, -SCH₃, -S(O)₂CH₃, and -S(O)₃H.

An exemplary embodiment of the 8-sulfonyl-2-aminobenzazepine-linker compound of Formula II includes wherein Q is selected from:








- 10 An exemplary embodiment of the 8-sulfonyl-2-aminobenzazepine-linker compound of Formula II includes wherein Q is phenoxy substituted with one or more F.




An exemplary embodiment of the 8-sulfonyl-2-aminobenzazepine-linker compound of Formula II includes wherein Q is 2,3,5,6-tetrafluorophenoxy.





- An exemplary embodiment of the 8-sulfonyl-2-aminobenzazepine-linker (8SO2BzL)
- 15 compound is selected from Table 2a. Each compound was synthesized, purified, and characterized by mass spectrometry and shown to have the mass indicated. Additional experimental procedures are found in the Examples. The 8-sulfonyl-2-aminobenzazepine-linker compounds of Table 2a demonstrate the surprising and unexpected property of TLR8 agonist selectivity which may predict useful therapeutic activity to treat cancer and other disorders. The
- 20 8-sulfonyl-2-aminobenzazepine-linker intermediate, Formula II compounds of Table 2a are used in conjugation with antibodies by the methods of Example 201 to form the Immunoconjugates of Table 3a.




Table 2a 8-Sulfonyl-Benzazepine-Linker (8SO2BzL) compounds




8SO2BzL No.	Structure	MW
8SO2BzL-1	 The structure of 8SO2BzL-1 features a long, flexible polyether linker chain (1,3-bis(2-ethoxyethyl)oxy) that connects two distinct chemical moieties. On one end, there is a 4-((2-ethoxyethyl)carbamoyl)phenyl group. The other end is attached to a 7-amino-2-((2-ethoxyethyl)carbamoyl)-5,6-difluoro-1,2,3,4-tetrahydroquinoline-3-carboxylate derivative, which is further substituted with a 2,4,6-trifluorophenyl group via an ester linkage.	1211.3
8SO2BzL-2	 The structure of 8SO2BzL-2 consists of a long, flexible polyether linker chain (1,3-bis(2-ethoxyethyl)oxy) connecting two moieties. One end is a 4-((2-ethoxyethyl)carbamoyl)pyrrolidine-2,5-dione. The other end is attached to a 7-amino-2-((2-ethoxyethyl)carbamoyl)-5,6-difluoro-1,2,3,4-tetrahydroquinoline-3-carboxylate derivative, which is further substituted with a 2,4,6-trifluorophenyl group via an ester linkage.	1047.1




8SO2BzL-3		1001.2
8SO2BzL-4		987.1
8SO2BzL-5		1046.2

8SO2BzL-6	 Chemical structure of 8SO2BzL-6. It features a long, flexible polyether chain (1,3-bis(2-ethoxyethyl)oxy) terminated with a 4-ethyl-1H-imidazole-5-carboxamide group. The chain is linked via an amide bond to a 4-ethyl-1H-imidazole-5-carboxamide group. The imidazole ring is substituted with a 4-(dimethylsulfonyl)phenyl group and an amino group at the 2-position.	1074.2
8SO2BzL-7	 Chemical structure of 8SO2BzL-7. It features a long, flexible polyether chain (1,3-bis(2-ethoxyethyl)oxy) terminated with a 4-ethyl-1H-imidazole-5-carboxamide group. The chain is linked via an amide bond to a 4-ethyl-1H-imidazole-5-carboxamide group. The imidazole ring is substituted with a 4-(hydroxysulfonyl)phenyl group and an amino group at the 2-position.	1044.2
8SO2BzL-8	 Chemical structure of 8SO2BzL-8. It features a long, flexible polyether chain (1,3-bis(2-ethoxyethyl)oxy) terminated with a 4-ethyl-1H-imidazole-5-carboxamide group. The chain is linked via an amide bond to a 4-ethyl-1H-imidazole-5-carboxamide group. The imidazole ring is substituted with a 4-(hydroxysulfonyl)phenyl group and an amino group at the 2-position.	1045.2

8SO2BzL-9		985.2
8SO2BzL-10		999.2
8SO2BzL-11		1001.2
8SO2BzL-12		1101.3

8SO2BzL-13		1060.2
8SO2BzL-14		1015.2
8SO2BzL-15		1129.3

8SO2BzL-16		1023.2
8SO2BzL-17		1047.1
8SO2BzL-18		971.1



8SO2BzL-19		998.1
8SO2BzL-20		1058.2
8SO2BzL-21		1041.1

8SO2BzL-22		1027.1
8SO2BzL-23		1087.1

Comparator compounds from Table 2b have an activated ester, tetrafluorophenyl or sulfotetrafluorophenyl group which reacts with a lysine residue of an antibody to form an immunoconjugate with an amide bond between the antibody and the TLR-agonist-linker moiety

5 according to Example 201.


Table 2b TLR agonist-linker Comparator Compounds

Comp	Structure	MW
C-1		1163.2
C-2		1274.3

8-SULFONYL-BENZAZEPINE IMMUNOCONJUGATES

Immune-stimulating antibody conjugates, i.e. immunoconjugates, direct TLR7/8 agonists
 5 into tumors to activate tumor-infiltrating myeloid cells and initiate a broad innate and adaptive
 anti-tumor immune response (Ackerman, et al., (2021) *Nature Cancer* 2:18-33.

Exemplary embodiments of immunoconjugates comprise an antibody covalently
 attached to one or more 8-sulfonyl-2-aminobenzazepine moieties by a linker, and having
 Formula I:



or a pharmaceutically acceptable salt thereof,


wherein:

Ab is the antibody;

p is an integer from 1 to 8;

L is the linker;

5 D is the 8-sulfonyl-2-aminobenzazepine moiety having the formula:




R^1 , R^2 , R^3 , and R^4 are independently selected from the group consisting of H, C₁-C₁₂ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₁₂ carbocyclyl, C₆-C₂₀ aryl, C₂-C₉ heterocyclyl, and C₁-C₂₀ heteroaryl, where alkyl, alkenyl, alkynyl, carbocyclyl, aryl, heterocyclyl, and heteroaryl
10 are independently and optionally substituted with one or more groups selected from:

- (C₁-C₁₂ alkylidiyl)-N(R⁵)-*;
- (C₁-C₁₂ alkylidiyl)-N(R⁵)₂;
- (C₁-C₁₂ alkylidiyl)-OR⁵;
- (C₃-C₁₂ carbocyclyl);
- 15 -(C₃-C₁₂ carbocyclyl)-*;
- (C₃-C₁₂ carbocyclyl)-(C₁-C₁₂ alkylidiyl)-NR⁵-*;
- (C₃-C₁₂ carbocyclyl)-(C₁-C₁₂ alkylidiyl)-N(R⁵)₂;
- (C₃-C₁₂ carbocyclyl)-NR⁵-C(=NR⁵)NR⁵-*;
- (C₆-C₂₀ aryl);
- 20 -(C₆-C₂₀ arylidiyl)-*;
- (C₆-C₂₀ arylidiyl)-N(R⁵)-*;
- (C₆-C₂₀ arylidiyl)-(C₁-C₁₂ alkylidiyl)-N(R⁵)-*;
- (C₆-C₂₀ arylidiyl)-(C₁-C₁₂ alkylidiyl)-(C₂-C₂₀ heterocyclidiyl)-*;
- (C₆-C₂₀ arylidiyl)-(C₁-C₁₂ alkylidiyl)-N(R⁵)₂;
- 25 -(C₆-C₂₀ arylidiyl)-(C₁-C₁₂ alkylidiyl)-NR⁵-C(=NR^{5a})N(R⁵)-*;
- (C₂-C₂₀ heterocyclyl);
- (C₂-C₂₀ heterocyclyl)-*;
- (C₂-C₉ heterocyclyl)-(C₁-C₁₂ alkylidiyl)-NR⁵-*;
- (C₂-C₉ heterocyclyl)-(C₁-C₁₂ alkylidiyl)-N(R⁵)₂;


- $-(C_2-C_9 \text{ heterocyclyl})-C(=O)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
 $-(C_2-C_9 \text{ heterocyclyl})-NR^5-C(=NR^{5a})NR^5-*$;
 $-(C_2-C_9 \text{ heterocyclyl})-NR^5-(C_6-C_{20} \text{ arylldiyl})-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
 $-(C_2-C_9 \text{ heterocyclyl})-(C_6-C_{20} \text{ arylldiyl})-*$;
5 $-(C_1-C_{20} \text{ heteroaryl})$;
 $-(C_1-C_{20} \text{ heteroaryl})-*$;
 $-(C_1-C_{20} \text{ heteroaryl})-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
 $-(C_1-C_{20} \text{ heteroaryl})-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)_2$;
 $-(C_1-C_{20} \text{ heteroaryl})-NR^5-C(=NR^{5a})N(R^5)-*$;
10 $-(C_1-C_{20} \text{ heteroaryl})-N(R^5)C(=O)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
 $-C(=O)-*$;
 $-C(=O)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)-*$;
 $-C(=O)-(C_2-C_{20} \text{ heterocyclldiyl})-*$;
 $-C(=O)N(R^5)_2$;
15 $-C(=O)N(R^5)-*$;
 $-C(=O)N(R^5)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)C(=O)R^5$;
 $-C(=O)N(R^5)-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)C(=O)N(R^5)_2$;
 $-C(=O)NR^5-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)CO_2R^5$;
 $-C(=O)NR^5-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)C(=NR^{5a})N(R^5)_2$;
20 $-C(=O)NR^5-(C_1-C_{12} \text{ alkylldiyl})-NR^5C(=NR^{5a})R^5$;
 $-C(=O)NR^5-(C_1-C_8 \text{ alkylldiyl})-NR^5(C_2-C_5 \text{ heteroaryl})$;
 $-C(=O)NR^5-(C_1-C_{20} \text{ heteroarylldiyl})-N(R^5)-*$;
 $-C(=O)NR^5-(C_1-C_{20} \text{ heteroarylldiyl})-*$;
 $-C(=O)NR^5-(C_1-C_{20} \text{ heteroarylldiyl})-(C_1-C_{12} \text{ alkylldiyl})-N(R^5)_2$;
25 $-C(=O)NR^5-(C_1-C_{20} \text{ heteroarylldiyl})-(C_2-C_{20} \text{ heterocyclldiyl})-C(=O)NR^5-(C_1-C_{12} \text{ alkylldiyl})-NR^5-*$;
 $-N(R^5)_2$;
 $-N(R^5)-*$;
 $-N(R^5)C(=O)R^5$;
30 $-N(R^5)C(=O)-*$;
 $-N(R^5)C(=O)N(R^5)_2$;
 $-N(R^5)C(=O)N(R^5)-*$;
 $-N(R^5)CO_2R^5$;

- $-\text{NR}^5\text{C}(=\text{NR}^{5a})\text{N}(\text{R}^5)_2$;
 $-\text{NR}^5\text{C}(=\text{NR}^{5a})\text{N}(\text{R}^5)-*$;
 $-\text{NR}^5\text{C}(=\text{NR}^{5a})\text{R}^5$;
 $-\text{N}(\text{R}^5)\text{C}(=\text{O})-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{N}(\text{R}^5)-*$;
5 $-\text{N}(\text{R}^5)-(\text{C}_2\text{-C}_5\text{ heteroaryl})$;
 $-\text{N}(\text{R}^5)-\text{S}(=\text{O})_2-(\text{C}_1\text{-C}_{12}\text{ alkyl})$;
 $-\text{O}-(\text{C}_1\text{-C}_{12}\text{ alkyl})$;
 $-\text{O}-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{N}(\text{R}^5)_2$;
 $-\text{O}-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{N}(\text{R}^5)-*$;
10 $-\text{O}-\text{C}(=\text{O})\text{N}(\text{R}^5)_2$;
 $-\text{O}-\text{C}(=\text{O})\text{N}(\text{R}^5)-*$;
 $-\text{O}-(\text{R}^5)-*$;
 $-\text{OR}^5$;
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20}\text{ heterocyclidyl})-*$;
15 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20}\text{ heterocyclidyl})-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{N}(\text{R}^5)_2$;
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20}\text{ heterocyclidyl})-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{NR}^5-*$; and
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20}\text{ heterocyclidyl})-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{OH}$;
or R^2 and R^3 together form a 5- or 6-membered heterocyclidyl ring;
 X^1 , X^2 , X^3 , and X^4 are independently selected from the group consisting of a bond,
20 $\text{C}(=\text{O})$, $\text{C}(=\text{O})\text{N}(\text{R}^5)$, O , $\text{N}(\text{R}^5)$, S , $\text{S}(\text{O})_2$, and $\text{S}(\text{O})_2\text{N}(\text{R}^5)$;
 R^5 is independently selected from the group consisting of H, $\text{C}_6\text{-C}_{20}$ aryl, $\text{C}_3\text{-C}_{12}$ carbocyclidyl, $\text{C}_6\text{-C}_{20}$ arylidyl, $\text{C}_1\text{-C}_{12}$ alkyl, and $\text{C}_1\text{-C}_{12}$ alkylidyl, or two R^5 groups together form a 5- or 6-membered heterocyclidyl ring;
 R^{5a} is selected from the group consisting of $\text{C}_6\text{-C}_{20}$ aryl and $\text{C}_1\text{-C}_{20}$ heteroaryl;
25 where the asterisk * indicates the attachment site of L, and where one of R^1 , R^2 , R^3 and R^4 is attached to L;
L is the linker selected from the group consisting of:
 $-\text{C}(=\text{O})-\text{PEG}-$;
 $-\text{C}(=\text{O})-\text{PEG}-\text{C}(=\text{O})\text{N}(\text{R}^6)-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{C}(=\text{O})-\text{Gluc}-$;
30 $-\text{C}(=\text{O})-\text{PEG}-\text{O}-$;
 $-\text{C}(=\text{O})-\text{PEG}-\text{O}-\text{C}(=\text{O})-$;
 $-\text{C}(=\text{O})-\text{PEG}-\text{C}(=\text{O})-$;
 $-\text{C}(=\text{O})-\text{PEG}-\text{C}(=\text{O})-\text{PEP}-$;

- $-\text{C}(=\text{O})-\text{PEG}-\text{N}(\text{R}^6)-;$
 $-\text{C}(=\text{O})-\text{PEG}-\text{N}(\text{R}^6)-\text{C}(=\text{O})-;$
 $-\text{C}(=\text{O})-\text{PEG}-\text{N}(\text{R}^6)-\text{PEG}-\text{C}(=\text{O})-\text{PEP}-;$
 $-\text{C}(=\text{O})-\text{PEG}-\text{N}^+(\text{R}^6)_2-\text{PEG}-\text{C}(=\text{O})-\text{PEP}-;$
5 $-\text{C}(=\text{O})-\text{PEG}-\text{C}(=\text{O})-\text{PEP}-\text{N}(\text{R}^6)-(\text{C}_1-\text{C}_{12} \text{ alkylidyl})-;$
 $-\text{C}(=\text{O})-\text{PEG}-\text{C}(=\text{O})-\text{PEP}-\text{N}(\text{R}^6)-(\text{C}_1-\text{C}_{12} \text{ alkylidyl})\text{N}(\text{R}^6)\text{C}(=\text{O})-(\text{C}_2-\text{C}_5$
 $\text{monoheterocyclidyl})-;$
 $-\text{C}(=\text{O})-\text{PEG}-\text{SS}-(\text{C}_1-\text{C}_{12} \text{ alkylidyl})-\text{OC}(=\text{O})-;$
 $-\text{C}(=\text{O})-\text{PEG}-\text{SS}-(\text{C}_1-\text{C}_{12} \text{ alkylidyl})-\text{C}(=\text{O})-;$
10 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-;$
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{C}(=\text{O})\text{N}(\text{R}^6)-(\text{C}_1-\text{C}_{12}$
 $\text{alkylidyl})-\text{C}(=\text{O})-\text{Gluc}-;$
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{O}-;$
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{O}-\text{C}(=\text{O})-;$
15 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{C}(=\text{O})-;$
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{N}(\text{R}^5)-;$
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{N}(\text{PEG}-\text{CO}_2\text{H})-\text{PEG}-\text{N}(\text{R}^5)-;$
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{C}(=\text{O})\text{N}(\text{PEG}-\text{CO}_2\text{H})-\text{PEG}-\text{N}(\text{R}^5)-;$
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{N}(\text{R}^5)-\text{C}(=\text{O})-;$
20 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{N}(\text{PEG}-\text{CO}_2\text{H})-\text{PEG}-\text{C}(=\text{O})-;$
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{C}(=\text{O})\text{N}(\text{PEG}-\text{CO}_2\text{H})-\text{PEG}-\text{C}(=\text{O})-;$
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{C}(=\text{O})-\text{PEP}-;$ and
 $-\text{succinimidyl}-(\text{CH}_2)_m-\text{C}(=\text{O})\text{N}(\text{R}^6)-\text{PEG}-\text{SS}-(\text{C}_1-\text{C}_{12} \text{ alkylidyl})-\text{OC}(=\text{O})-;$
 R^6 is independently H or C_1-C_6 alkyl;
25 PEG has the formula: $-(\text{CH}_2\text{CH}_2\text{O})_n-(\text{CH}_2)_m-$; m is an integer from 1 to 5, and n is an
integer from 2 to 50;
Gluc has the formula:




PEP has the formula:



- where AA is independently selected from a natural or unnatural amino acid side chain, or one or more of AA, and an adjacent nitrogen atom form a 5-membered ring proline amino acid, and the wavy line indicates a point of attachment;

Cyc is selected from C₆-C₂₀ aryldiyl and C₁-C₂₀ heteroaryldiyl, optionally substituted with one or more groups selected from F, Cl, NO₂, -OH, -OCH₃, and a glucuronic acid having the structure:



- R⁷ is selected from the group consisting of -CH(R⁸)O-, -CH₂-, -CH₂N(R⁸)-, and -CH(R⁸)O-C(=O)-, where R⁸ is selected from H, C₁-C₆ alkyl, C(=O)-C₁-C₆ alkyl, and -C(=O)N(R⁹)₂, where R⁹ is independently selected from the group consisting of H, C₁-C₁₂ alkyl, and -(CH₂CH₂O)_n-(CH₂)_m-OH, where m is an integer from 1 to 5, and n is an integer from 2 to 50, or two R⁹ groups together form a 5- or 6-membered heterocyclyl ring;

- y is an integer from 2 to 12;

z is 0 or 1; and

alkyl, alkylidiyl, alkenyl, alkenyldiyl, alkynyl, alkynyldiyl, aryl, aryldiyl, carbocyclyl, carbocyclidiyl, heterocyclyl, heterocyclidiyl, heteroaryl, and heteroaryldiyl are independently and optionally substituted with one or more groups independently selected from F, Cl, Br, I, -

- CN, -CH₃, -CH₂CH₃, -CH=CH₂, -C≡CH, -C≡CCH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH(CH₃)₂, -CH₂OH, -CH₂OCH₃, -CH₂CH₂OH, -C(CH₃)₂OH, -CH(OH)CH(CH₃)₂, -C(CH₃)₂CH₂OH, -CH₂CH₂SO₂CH₃, -CH₂OP(O)(OH)₂, -CH₂F, -CHF₂, -CF₃, -CH₂CF₃, -CH₂CHF₂, -CH(CH₃)CN, -C(CH₃)₂CN, -CH₂CN, -CH₂NH₂, -CH₂NHSO₂CH₃, -CH₂NHCH₃, -CH₂N(CH₃)₂, -CO₂H, -COCH₃, -CO₂CH₃, -CO₂C(CH₃)₃, -COCH(OH)CH₃, -CONH₂, -CONHCH₃, -CON(CH₃)₂, -C(CH₃)₂CONH₂, -NH₂, -NHCH₃, -N(CH₃)₂, -NHCOCH₃, -N(CH₃)COCH₃, -NHS(O)₂CH₃, -N(CH₃)C(CH₃)₂CONH₂, -N(CH₃)CH₂CH₂S(O)₂CH₃, -NHC(=NH)H, -NHC(=NH)CH₃, -NHC(=NH)NH₂, -NHC(=O)NH₂, -NO₂, =O, -OH, -OCH₃, -OCH₂CH₃, -OCH₂CH₂OCH₃, -OCH₂CH₂OH, -OCH₂CH₂N(CH₃)₂, -O(CH₂CH₂O)_n-

$(\text{CH}_2)_m\text{CO}_2\text{H}$, $-\text{O}(\text{CH}_2\text{CH}_2\text{O})_n\text{H}$, $-\text{OCH}_2\text{F}$, $-\text{OCHF}_2$, $-\text{OCF}_3$, $-\text{OP}(\text{O})(\text{OH})_2$, $-\text{S}(\text{O})_2\text{N}(\text{CH}_3)_2$, $-\text{SCH}_3$, $-\text{S}(\text{O})_2\text{CH}_3$, and $-\text{S}(\text{O})_3\text{H}$.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein X^1 is a bond, and R^1 is H.

- 5 An exemplary embodiment of the immunoconjugate of Formula I includes wherein X^2 is a bond, and R^2 is C_1 - C_8 alkyl.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein X^2 and X^3 are each a bond, and R^2 and R^3 are independently selected from C_1 - C_8 alkyl, $-\text{O}-(\text{C}_1$ - C_{12} alkyl), $-(\text{C}_1$ - C_{12} alkylidyl)- OR^5 , $-(\text{C}_1$ - C_8 alkylidyl)- $\text{N}(\text{R}^5)\text{CO}_2\text{R}^5$, $-(\text{C}_1$ - C_{12} alkyl)-

- 10 $\text{OC}(\text{O})\text{N}(\text{R}^5)_2$, $-\text{O}-(\text{C}_1$ - C_{12} alkyl)- $\text{N}(\text{R}^5)\text{CO}_2\text{R}^5$, and $-\text{O}-(\text{C}_1$ - C_{12} alkyl)- $\text{OC}(\text{O})\text{N}(\text{R}^5)_2$.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein R^2 is C_1 - C_8 alkyl and R^3 is $-(\text{C}_1$ - C_8 alkylidyl)- $\text{N}(\text{R}^5)\text{CO}_2\text{R}^4$.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein R^2 is $-\text{CH}_2\text{CH}_2\text{CH}_3$ and R^3 is selected from $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHCO}_2(\text{t-Bu})$, $-\text{OCH}_2\text{CH}_2\text{NHCO}_2(\text{cyclobutyl})$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHCO}_2(\text{cyclobutyl})$.

- 15 $\text{OCH}_2\text{CH}_2\text{NHCO}_2(\text{cyclobutyl})$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NHCO}_2(\text{cyclobutyl})$.


An exemplary embodiment of the immunoconjugate of Formula I includes wherein R^2 and R^3 are each independently selected from $-\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CF}_3$, $-\text{OCH}_2\text{CH}_2\text{OH}$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein R^2

- 20 and R^3 are each $-\text{CH}_2\text{CH}_2\text{CH}_3$.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein R^2 is $-\text{CH}_2\text{CH}_2\text{CH}_3$ and R^3 is $-\text{OCH}_2\text{CH}_3$.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein X^3 - R^3 is selected from the group consisting of:




An exemplary embodiment of the immunoconjugate of Formula I includes wherein X^4 is a bond, and R^4 is H.

- 5 An exemplary embodiment of the immunoconjugate of Formula I includes wherein R^1 is attached to L.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein R^2 or R^3 is attached to L.

- 10 An exemplary embodiment of the immunoconjugate of Formula I includes wherein X^3 - R^3 -L is selected from the group consisting of:



where the wavy line indicates the point of attachment to N.

- 5 An exemplary embodiment of the immunoconjugate of Formula I includes wherein R^4 is C_1 - C_{12} alkyl.


An exemplary embodiment of the immunoconjugate of Formula I includes wherein R^4 is $-(C_1$ - C_{12} alkyldiyl)- $N(R^5)-*$; where the asterisk * indicates the attachment site of L.

- An exemplary embodiment of the immunoconjugate of Formula I includes wherein L is
10 $-C(=O)-PEG-$ or $-C(=O)-PEG-C(=O)-$.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein L is attached to a cysteine thiol of the antibody.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein for the PEG, m is 1 or 2, and n is an integer from 2 to 10; or wherein n is 10.

- 15 An exemplary embodiment of the immunoconjugate of Formula I includes wherein L comprises PEP and PEP is a dipeptide and has the formula:




- An exemplary embodiment of the immun conjugate of Formula I includes wherein AA₁ and AA₂ are independently selected from H, -CH₃, -CH(CH₃)₂, -CH₂(C₆H₅), -CH₂CH₂CH₂CH₂NH₂, -CH₂CH₂CH₂NHC(NH)NH₂, -CHCH(CH₃)CH₃, -CH₂SO₃H, and -CH₂CH₂CH₂NHC(O)NH₂; or AA₁ and AA₂ form a 5-membered ring proline amino acid.

An exemplary embodiment of the immun conjugate of Formula I includes wherein AA₁ is -CH(CH₃)₂, and AA₂ is -CH₂CH₂CH₂NHC(O)NH₂.


An exemplary embodiment of the immun conjugate of Formula I includes wherein AA₁ and AA₂ are independently selected from GlcNAc aspartic acid, -CH₂SO₃H, and -CH₂OPO₃H.

- 10 An exemplary embodiment of the immun conjugate of Formula I includes wherein PEP has the formula:




wherein AA₁ and AA₂ are independently selected from a side chain of a naturally-occurring amino acid.

- 15 An exemplary embodiment of the immun conjugate of Formula I includes wherein L comprises PEP and PEP is a tripeptide and has the formula:



An exemplary embodiment of the immun conjugate of Formula I includes wherein L comprises PEP and PEP is a tetrapeptide and has the formula:



20

An exemplary embodiment of the immun conjugate of Formula I includes wherein:

AA₁ is selected from the group consisting of Abu, Ala, and Val;


AA₂ is selected from the group consisting of Nle(O-Bzl), Oic and Pro;

AA₃ is selected from the group consisting of Ala and Met(O)₂; and


AA₄ is selected from the group consisting of Oic, Arg(NO₂), Bpa, and Nle(O-Bzl).

- 5 An exemplary embodiment of the immunoconjugate of Formula I includes wherein L comprises PEP and PEP is selected from the group consisting of Ala-Pro-Val, Asn-Pro-Val, Ala-Ala-Val, Ala-Ala-Pro-Ala, Ala-Ala-Pro-Val, and Ala-Ala-Pro-Nva.

An exemplary embodiment of the immunoconjugate of Formula I includes wherein L comprises PEP and PEP is selected from the structures:



An exemplary embodiment of the immunoconjugate of Formula I includes wherein L is selected from the structures:



where the wavy line indicates the attachment to R⁵.

The invention includes all reasonable combinations, and permutations of the features, of the Formula I embodiments.

5 In certain embodiments, the immunoconjugate compounds of the invention include those with immunostimulatory activity. The immunoconjugates of the invention selectively deliver an effective dose of a 8-sulfonyl-2-aminobenzazepine (8SO2Bz) drug to tumor tissue, whereby greater selectivity (*i.e.*, a lower efficacious dose) may be achieved while increasing the therapeutic index (“therapeutic window”) relative to unconjugated 8SO2Bz.

10 Each immunoconjugate of Tables 3a and 3b was prepared according to the methods of Example 201, purified by HPLC, and characterized by mass spectroscopy.

Table 3a 8-Sulfonyl-2-Aminobenzazepine Immunoconjugates (IC)

Immunoconjugate No.	8SO2BzL Table 2a	Antibody Antigen	DAR	cDC Activation (IL12p70 Secretion) – EC ₅₀ (nM)
IC-1	8SO2BzL-1	trastuzumab HER2	2.5	
IC-2	8SO2BzL-2	TROP2	3.9	0.8
IC-3	8SO2BzL-3	TROP2	3.9	0.4
IC-4	8SO2BzL-4	TROP2	3.8	

IC-5	8SO2BzL-5	TROP2	3.7	0.4
IC-6	8SO2BzL-6	TROP2	2.6	0.6
IC-7	8SO2BzL-7	TROP2	3.9	27.3
IC-8	8SO2BzL-8	TROP2	3.6	4.2
IC-9	8SO2BzL-9	TROP2	4.0	2.4
IC-10	8SO2BzL-10	TROP2	4.3	1.5
IC-11	8SO2BzL-8	TROP2	3.8	mouse
IC-12	8SO2BzL-8	TROP2	3.8	mouse
IC-13	8SO2BzL-8	rituximab CD20	3.2	
IC-14	8SO2BzL-8	TROP2	3.3	mouse
IC-15	8SO2BzL-7	avelumab PD-L1	4.0	
IC-16	8SO2BzL-8	avelumab PD-L1	3.8	
IC-17	8SO2BzL-5	110-G1f PD-L1	3.1, 4.0	0.3
IC-18	8SO2BzL-4	110-G1f PD-L1	3.2, 3.4, 3.5, 3.9	0.4
IC-19	8SO2BzL-3	110-G1f PD-L1	3.1	0.4
IC-20	8SO2BzL-2	110-G1f PD-L1	3.5	0.2
IC-21	8SO2BzL-7	anti-M PD-L1	3.3	
IC-22	8SO2BzL-7	rat IgG2b isotype control	3.3	
IC-23	8SO2BzL-11	trastuzumab HER2	3.3	
IC-24	8SO2BzL-12	trastuzumab HER2	3.8	
IC-25	8SO2BzL-14	trastuzumab HER2	3.4	
IC-26	8SO2BzL-15	trastuzumab	3.5	1.5

		HER2		
IC-27	8SO2BzL-16	trastuzumab HER2	3.5, 3.6	1.6
IC-28	8SO2BzL-18	110-G1f PD-L1	3.5	0.4
IC-29	8SO2BzL-19	110-G1f PD-L1	3.6	0.6
IC-30	8SO2BzL-20	110-G1f PD-L1	3.5	1.3
IC-31	8SO2BzL-17	trastuzumab HER2	3.2	
IC-32	8SO2BzL-2	trastuzumab HER2	3.3	1.7
IC-33	8SO2BzL-3	trastuzumab HER2	3.5	0.5
IC-34	8SO2BzL-4	trastuzumab HER2	3.7	0.9
IC-35	8SO2BzL-5	trastuzumab HER2	3.5	0.9
IC-36	8SO2BzL-10	trastuzumab HER2	3.7	
IC-37	8SO2BzL-23	trastuzumab HER2	2.1	1.3
IC-38	8SO2BzL-22	trastuzumab HER2	1.9	1.8
IC-39	8SO2BzL-21	trastuzumab HER2	2.8	1.6
IC-40	8SO2BzL-9	trastuzumab HER2	4.0	
IC-41	8SO2BzL-7	trastuzumab HER2	4.0	
IC-42	8SO2BzL-8	trastuzumab HER2	3.4, 4.0	
IC-43	8SO2BzL-6	trastuzumab HER2	3.8	

IC-44	8SO2BzL-7	hIgG1 isotype control	3.8	
-------	-----------	--------------------------	-----	--

Table 3b Comparator Immunoconjugates (CIC)

Comparator Immunoconjugate No.	Comparator compound Table 2b	Antibody	DAR	cDC Activation (IL12p70 Secretion) – EC ₅₀ (nM)
CIC-1	C-1	TROP2	2.4	0.8
CIC-2	C-2	TROP2	2.5	0.7

Figure 1 shows a graph of secreted TNFα (Tumor Necrosis Factor alpha) cytokine levels after incubation of varying concentrations of immunoconjugate IC-3, comparator CIC-1, and naked antibody TROP2 with a co-culture of cancer cells with a cDC-enriched primary cell isolate. Figure 2 shows a graph of secreted TNFα (Tumor Necrosis Factor alpha) cytokine levels after incubation of varying concentrations of immunoconjugate IC-3, comparator CIC-2, and naked antibody TROP2 with a co-culture of cancer cells with a cDC-enriched primary cell isolate.

Secreted cytokine levels in supernatants were determined using a LegendPlex cytokine bead array kit. TROP2-targeted immunoconjugates IC-2, CIC-1, and CIC-2 induce secretion of cytokine TNFα (alpha), relevant to mounting an immune response to cancer and demonstrate the activation of myeloid cells when exposed to TROP2-expressing tumor cells. The 8-sulfonyl-2-aminobenzazepine immunoconjugate IC-3 stimulated higher levels of TNFα than the comparator immunoconjugates CIC-1 and CIC-2. Notably, the 8-sulfonyl-2-aminobenzazepine payload represents a more efficient payload providing increased activity, while decreasing molecular weight and hydrophobicity. Naked antibody TROP2 does not induce myeloid activation, demonstrating the dependence on the TLR7/8 activating payload.

Drug loading is represented by p, the number of 8-sulfonyl-2-aminobenzazepine (8SO2Bz) moieties per antibody in an immunoconjugate of Formula I, and as measured (DAR) in the exemplary Immunoconjugates of Table 3a. Drug (8SO2Bz) loading may range from 1 to about 8 drug moieties (D) per antibody. Immunoconjugates of Formula I include mixtures or collections of antibodies conjugated with a range of drug moieties, from 1 to about 8. In some embodiments, the number of drug moieties that can be conjugated to an antibody is limited by the number of reactive or available amino acid side chain residues such as lysine and cysteine. In some embodiments, free cysteine residues are introduced into the antibody amino acid

sequence by the methods described herein. In such aspects, p may be 1, 2, 3, 4, 5, 6, 7, or 8, and ranges thereof, such as from 1 to 8 or from 2 to 5. In any such aspect, p and n are equal (i.e., p = n = 1, 2, 3, 4, 5, 6, 7, or 8, or some range there between). Exemplary immunoconjugates of Formula I include, but are not limited to, antibodies that have 1, 2, 3, or 4 engineered cysteine amino acids (Lyon, R. et al. (2012) *Methods in Enzym.* 502:123-138). In some embodiments, one or more free cysteine residues are already present in an antibody forming intra-chain and inter-chain disulfide bonds (native disulfide groups), without the use of engineering, in which case the existing free, reduced cysteine residues may be used to conjugate the antibody to a drug. In some embodiments, an antibody is exposed to reducing conditions prior to conjugation of the antibody in order to generate one or more free cysteine residues.

For some immunoconjugates, p may be limited by the number of attachment sites on the antibody. For example, where the attachment is a cysteine thiol, as in certain exemplary embodiments described herein, an antibody may have only one or a limited number of cysteine thiol groups, or may have only one or a limited number of sufficiently reactive thiol groups, to which the drug may be attached. In other embodiments, one or more lysine amino groups in the antibody may be available and reactive for conjugation with a 8SO₂Bz-linker compound of Formula II. In certain embodiments, higher drug loading, e.g. p > 5, may cause aggregation, insolubility, toxicity, or loss of cellular permeability of certain antibody-drug conjugates. In certain embodiments, the average drug loading for an immunoconjugate ranges from 1 to about 8; from about 2 to about 6; or from about 3 to about 5. In certain embodiments, an antibody is subjected to denaturing conditions to reveal reactive nucleophilic groups such as lysine or cysteine.

The loading (drug/antibody ratio) of an immunoconjugate may be controlled in different ways, and for example, by: (i) limiting the molar excess of the 8SO₂Bz-linker intermediate compound relative to antibody, (ii) limiting the conjugation reaction time or temperature, and (iii) partial or limiting reductive denaturing conditions for optimized antibody reactivity.

It is to be understood that where more than one nucleophilic group of the antibody reacts with a drug, then the resulting product is a mixture of immunoconjugate compounds with a distribution of one or more drug moieties attached to an antibody. The average number of drugs per antibody may be calculated from the mixture by a dual ELISA antibody assay, which is specific for antibody and specific for the drug. Individual immunoconjugate molecules may be identified in the mixture by mass spectroscopy and separated by HPLC, e.g. hydrophobic interaction chromatography (see, e.g., McDonagh et al. (2006) *Prot. Engr. Design & Selection* 19(7):299-307; Hamblett et al. (2004) *Clin. Cancer Res.* 10:7063-7070; Hamblett, K.J., et al. "Effect of drug loading on the pharmacology, pharmacokinetics, and toxicity of an anti-CD30

antibody-drug conjugate,” Abstract No. 624, American Association for Cancer Research, 2004 Annual Meeting, March 27-31, 2004, Proceedings of the AACR, Volume 45, March 2004; Alley, S.C., et al. “Controlling the location of drug attachment in antibody-drug conjugates,” Abstract No. 627, American Association for Cancer Research, 2004 Annual Meeting, March 27-31, 2004, Proceedings of the AACR, Volume 45, March 2004). In certain embodiments, a homogeneous immunoconjugate with a single loading value may be isolated from the conjugation mixture by electrophoresis or chromatography.

Assessment of Immunoconjugate Activity *In Vitro* may be conducted according to the methods of Example 203.

10 COMPOSITIONS OF IMMUNOCONJUGATES

The invention provides a composition, e.g., a pharmaceutically or pharmacologically acceptable composition or formulation, comprising a plurality of immunoconjugates as described herein and optionally a carrier therefor, e.g., a pharmaceutically or pharmacologically acceptable carrier. The immunoconjugates can be the same or different in the composition, i.e., the composition can comprise immunoconjugates that have the same number of adjuvants linked to the same positions on the antibody construct and/or immunoconjugates that have the same number of 8-sulfonyl-2-aminobenzazepine (8SO2Bz) adjuvants linked to different positions on the antibody construct, that have different numbers of 8SO2Bz adjuvants linked to the same positions on the antibody construct, or that have different numbers of 8SO2Bz adjuvants linked to different positions on the antibody construct.

In an exemplary embodiment, a composition comprising the immunoconjugate compounds comprises a mixture of the immunoconjugate compounds, wherein the average drug (8SO2Bz) loading per antibody (DAR) in the mixture of immunoconjugate compounds is about 2 to about 5.

A composition of immunoconjugates of the invention can have an average adjuvant to antibody construct ratio (DAR) of about 0.4 to about 10. A skilled artisan will recognize that the number of 8-sulfonyl-2-aminobenzazepine adjuvants conjugated to the antibody construct may vary from immunoconjugate to immunoconjugate in a composition comprising multiple immunoconjugates of the invention and thus the adjuvant to antibody construct (e.g., antibody) ratio can be measured as an average which may be referred to as the drug to antibody ratio (DAR). The adjuvant to antibody construct (e.g., antibody) ratio can be assessed by any suitable means, many of which are known in the art, including conventional means such as mass spectrometry, ELISA assay, and HPLC. The quantitative distribution of immunoconjugates in a composition in terms of p may also be determined. In some instances, separation, purification,

and characterization of homogeneous immunoconjugates where p is a certain value from immunoconjugates with other drug loadings may be achieved by means such as reverse phase HPLC or electrophoresis.

In some embodiments, the composition further comprises one or more pharmaceutically or pharmacologically acceptable excipients. For example, the immunoconjugates of the invention can be formulated for parenteral administration, such as IV administration or administration into a body cavity or lumen of an organ. Alternatively, the immunoconjugates can be injected intra-tumorally. Compositions for injection will commonly comprise a solution of the immunoconjugate dissolved in a pharmaceutically acceptable carrier. Among the acceptable vehicles and solvents that can be employed are water and an isotonic solution of one or more salts such as sodium chloride, e.g., Ringer's solution. In addition, sterile fixed oils can conventionally be employed as a solvent or suspending medium. For this purpose, any bland fixed oil can be employed, including synthetic monoglycerides or diglycerides. In addition, fatty acids such as oleic acid can likewise be used in the preparation of injectables. These compositions desirably are sterile and generally free of undesirable matter. These compositions can be sterilized by conventional, well known sterilization techniques. The compositions can contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents, e.g., sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate and the like.

The composition can contain any suitable concentration of the immunoconjugate. The concentration of the immunoconjugate in the composition can vary widely, and will be selected primarily based on fluid volumes, viscosities, body weight, and the like, in accordance with the particular mode of administration selected and the patient's needs. In certain embodiments, the concentration of an immunoconjugate in a solution formulation for injection will range from about 0.1% (w/w) to about 10% (w/w).

METHOD OF TREATING CANCER WITH IMMUNOCONJUGATES

The invention provides a method for treating cancer. The method includes administering a therapeutically effective amount of an immunoconjugate as described herein (e.g., as a composition as described herein) to a subject in need thereof, e.g., a subject that has cancer and is in need of treatment for the cancer. The method includes administering a therapeutically effective amount of an immunoconjugate (IC) selected from Table 3a.

It is contemplated that the immunoconjugate of the present invention may be used to treat various hyperproliferative diseases or disorders, e.g. characterized by the overexpression of

a tumor antigen. Exemplary hyperproliferative disorders include benign or malignant solid tumors and hematological disorders such as leukemia and lymphoid malignancies.

In another aspect, an immunoconjugate for use as a medicament is provided. In certain embodiments, the invention provides an immunoconjugate for use in a method of treating an individual comprising administering to the individual an effective amount of the immunoconjugate. In one such embodiment, the method further comprises administering to the individual an effective amount of at least one additional therapeutic agent, *e.g.*, as described herein.

In a further aspect, the invention provides for the use of an immunoconjugate in the manufacture or preparation of a medicament. In one embodiment, the medicament is for treatment of cancer, the method comprising administering to an individual having cancer an effective amount of the medicament. In one such embodiment, the method further comprises administering to the individual an effective amount of at least one additional therapeutic agent, *e.g.*, as described herein.

Carcinomas are malignancies that originate in the epithelial tissues. Epithelial cells cover the external surface of the body, line the internal cavities, and form the lining of glandular tissues. Examples of carcinomas include, but are not limited to, adenocarcinoma (cancer that begins in glandular (secretory) cells such as cancers of the breast, pancreas, lung, prostate, stomach, gastroesophageal junction, and colon) adrenocortical carcinoma; hepatocellular carcinoma; renal cell carcinoma; ovarian carcinoma; carcinoma in situ; ductal carcinoma; carcinoma of the breast; basal cell carcinoma; squamous cell carcinoma; transitional cell carcinoma; colon carcinoma; nasopharyngeal carcinoma; multilocular cystic renal cell carcinoma; oat cell carcinoma; large cell lung carcinoma; small cell lung carcinoma; non-small cell lung carcinoma; and the like. Carcinomas may be found in prostate, pancreas, colon, brain (usually as secondary metastases), lung, breast, and skin. In some embodiments, methods for treating non-small cell lung carcinoma include administering an immunoconjugate containing an antibody construct that is capable of binding a tumor-associated antigen.

Soft tissue tumors are a highly diverse group of rare tumors that are derived from connective tissue. Examples of soft tissue tumors include, but are not limited to, alveolar soft part sarcoma; angiomatoid fibrous histiocytoma; chondromyxoid fibroma; skeletal chondrosarcoma; extraskeletal myxoid chondrosarcoma; clear cell sarcoma; desmoplastic small round-cell tumor; dermatofibrosarcoma protuberans; endometrial stromal tumor; Ewing's sarcoma; fibromatosis (Desmoid); infantile fibrosarcoma; gastrointestinal stromal tumor; bone giant cell tumor; tenosynovial giant cell tumor; inflammatory myofibroblastic tumor; uterine leiomyoma; leiomyosarcoma; lipoblastoma; typical lipoma; spindle cell or pleomorphic lipoma;

atypical lipoma; chondroid lipoma; well-differentiated liposarcoma; myxoid/round cell liposarcoma; pleomorphic liposarcoma; myxoid malignant fibrous histiocyoma; high-grade malignant fibrous histiocyoma; myxofibrosarcoma; malignant peripheral nerve sheath tumor; mesothelioma; neuroblastoma; osteochondroma; osteosarcoma; primitive neuroectodermal tumor; alveolar rhabdomyosarcoma; embryonal rhabdomyosarcoma; benign or malignant schwannoma; synovial sarcoma; Evan's tumor; nodular fasciitis; desmoid-type fibromatosis; solitary fibrous tumor; dermatofibrosarcoma protuberans (DFSP); angiosarcoma; epithelioid hemangioendothelioma; tenosynovial giant cell tumor (TGCT); pigmented villonodular synovitis (PVNS); fibrous dysplasia; myxofibrosarcoma; fibrosarcoma; synovial sarcoma; malignant peripheral nerve sheath tumor; neurofibroma; pleomorphic adenoma of soft tissue; and neoplasias derived from fibroblasts, myofibroblasts, histiocytes, vascular cells/endothelial cells, and nerve sheath cells.

A sarcoma is a rare type of cancer that arises in cells of mesenchymal origin, e.g., in bone or in the soft tissues of the body, including cartilage, fat, muscle, blood vessels, fibrous tissue, or other connective or supportive tissue. Different types of sarcoma are based on where the cancer forms. For example, osteosarcoma forms in bone, liposarcoma forms in fat, and rhabdomyosarcoma forms in muscle. Examples of sarcomas include, but are not limited to, primitive neuroectodermal tumor (PNET) of the thoracopulmonary region (Askin's tumor); sarcoma botryoides; chondrosarcoma; malignant hemangioendothelioma; malignant schwannoma; osteosarcoma; and soft tissue sarcomas (e.g., alveolar soft part sarcoma; angiosarcoma; cystosarcoma phyllodes; dermatofibrosarcoma protuberans (DFSP); desmoid tumor; desmoplastic small round cell tumor; epithelioid sarcoma; extraskelatal chondrosarcoma; extraskelatal osteosarcoma; fibrosarcoma; gastrointestinal stromal tumor (GIST); hemangiopericytoma; hemangiosarcoma (more commonly referred to as "angiosarcoma"); Kaposi's sarcoma; leiomyosarcoma; liposarcoma; lymphangiosarcoma; malignant peripheral nerve sheath tumor (MPNST); neurofibrosarcoma; synovial sarcoma; and undifferentiated pleomorphic sarcoma).

A teratoma is a type of germ cell tumor that may contain several different types of tissue (e.g., can include tissues derived from any and/or all of the three germ layers: endoderm, mesoderm, and ectoderm), including, for example, hair, muscle, and bone. Teratomas occur most often in the ovaries in women, the testicles in men, and the tailbone in children.

Melanoma is a form of cancer that begins in melanocytes (cells that make the pigment melanin). Melanoma may begin in a mole (skin melanoma), but can also begin in other pigmented tissues, such as in the eye or in the intestines.

Merkel cell carcinoma is a rare type of skin cancer that usually appears as a flesh-colored or bluish-red nodule on the face, head or neck. Merkel cell carcinoma is also called neuroendocrine carcinoma of the skin. In some embodiments, methods for treating Merkel cell carcinoma include administering an immunoconjugate containing an antibody construct that is capable of binding, for example, CEA (e.g., labetuzumab, biosimilars thereof, or biobetters thereof). In some embodiments, the Merkel cell carcinoma has metastasized when administration occurs.

Leukemias are cancers that start in blood-forming tissue, such as the bone marrow, and cause large numbers of abnormal blood cells to be produced and enter the bloodstream. For example, leukemias can originate in bone marrow-derived cells that normally mature in the bloodstream. Leukemias are named for how quickly the disease develops and progresses (e.g., acute versus chronic) and for the type of white blood cell that is affected (e.g., myeloid versus lymphoid). Myeloid leukemias are also called myelogenous or myeloblastic leukemias. Lymphoid leukemias are also called lymphoblastic or lymphocytic leukemia. Lymphoid leukemia cells may collect in the lymph nodes, which can become swollen. Examples of leukemias include, but are not limited to, Acute myeloid leukemia (AML), Acute lymphoblastic leukemia (ALL), Chronic myeloid leukemia (CML), and Chronic lymphocytic leukemia (CLL).

Lymphomas are cancers that begin in cells of the immune system. For example, lymphomas can originate in bone marrow-derived cells that normally mature in the lymphatic system. There are two basic categories of lymphomas. One category of lymphoma is Hodgkin lymphoma (HL), which is marked by the presence of a type of cell called the Reed-Sternberg cell. There are currently 6 recognized types of HL. Examples of Hodgkin lymphomas include nodular sclerosis classical Hodgkin lymphoma (CHL), mixed cellularity CHL, lymphocyte-depletion CHL, lymphocyte-rich CHL, and nodular lymphocyte predominant HL.

The other category of lymphoma is non-Hodgkin lymphomas (NHL), which includes a large, diverse group of cancers of immune system cells. Non-Hodgkin lymphomas can be further divided into cancers that have an indolent (slow-growing) course and those that have an aggressive (fast-growing) course. There are currently 61 recognized types of NHL. Examples of non-Hodgkin lymphomas include, but are not limited to, AIDS-related Lymphomas, anaplastic large-cell lymphoma, angioimmunoblastic lymphoma, blastic NK-cell lymphoma, Burkitt's lymphoma, Burkitt-like lymphoma (small non-cleaved cell lymphoma), chronic lymphocytic leukemia/small lymphocytic lymphoma, cutaneous T-Cell lymphoma, diffuse large B-Cell lymphoma, enteropathy-type T-Cell lymphoma, follicular lymphoma, hepatosplenic gamma-delta T-Cell lymphomas, T-Cell leukemias, lymphoblastic lymphoma, mantle cell lymphoma, marginal zone lymphoma, nasal T-Cell lymphoma, pediatric lymphoma, peripheral T-Cell

lymphomas, primary central nervous system lymphoma, transformed lymphomas, treatment-related T-Cell lymphomas, and Waldenstrom's macroglobulinemia.

Brain cancers include any cancer of the brain tissues. Examples of brain cancers include, but are not limited to, gliomas (e.g., glioblastomas, astrocytomas, oligodendrogliomas, 5 ependymomas, and the like), meningiomas, pituitary adenomas, and vestibular schwannomas, primitive neuroectodermal tumors (medulloblastomas).

Immunoconjugates of the invention can be used either alone or in combination with other agents in a therapy. For instance, an immunoconjugate may be co-administered with at least one additional therapeutic agent, such as a chemotherapeutic agent. Such combination therapies 10 encompass combined administration (where two or more therapeutic agents are included in the same or separate formulations), and separate administration, in which case, administration of the immunoconjugate can occur prior to, simultaneously, and/or following, administration of the additional therapeutic agent and/or adjuvant. Immunoconjugates can also be used in combination with radiation therapy.

15 The immunoconjugates of the invention (and any additional therapeutic agent) can be administered by any suitable means, including oral, parenteral, intrapulmonary, and intranasal, and, if desired for local treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. Dosing can be by any suitable route, e.g. by injections, such as intravenous or subcutaneous injections, 20 depending in part on whether the administration is brief or chronic. Various dosing schedules including but not limited to single or multiple administrations over various time-points, bolus administration, and pulse infusion are contemplated herein.

The immunoconjugate is administered to a subject in need thereof in any therapeutically effective amount using any suitable dosing regimen, such as the dosing regimens utilized for 25 labetuzumab, biosimilars thereof, and biobetters thereof. For example, the methods can include administering the immunoconjugate to provide a dose of from about 100 ng/kg to about 50 mg/kg to the subject. The immunoconjugate dose can range from about 5 mg/kg to about 50 mg/kg, from about 10 µg/kg to about 5 mg/kg, or from about 100 µg/kg to about 1 mg/kg. The immunoconjugate dose can be about 100, 200, 300, 400, or 500 µg/kg. The immunoconjugate 30 dose can be about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 mg/kg. The immunoconjugate dose can also be outside of these ranges, depending on the particular conjugate as well as the type and severity of the cancer being treated. Frequency of administration can range from a single dose to multiple doses per week, or more frequently. In some embodiments, the immunoconjugate is administered from about once per month to about five times per week. In some embodiments, 35 the immunoconjugate is administered once per week.

In another aspect, the invention provides a method for preventing cancer. The method comprises administering a therapeutically effective amount of an immunoconjugate (e.g., as a composition as described above) to a subject. In certain embodiments, the subject is susceptible to a certain cancer to be prevented.


5 Some embodiments of the invention provide methods for treating cancer as described above, wherein the cancer is breast cancer. Breast cancer can originate from different areas in the breast, and a number of different types of breast cancer have been characterized. For example, the immunoconjugates of the invention can be used for treating ductal carcinoma *in situ*; invasive ductal carcinoma (e.g., tubular carcinoma, medullary carcinoma, mucinous carcinoma; papillary carcinoma; or cribriform carcinoma of the breast); lobular carcinoma *in situ*; invasive lobular carcinoma; inflammatory breast cancer; and other forms of breast cancer such as triple negative (test negative for estrogen receptors, progesterone receptors, and excess HER2 protein) breast cancer. In some embodiments, methods for treating breast cancer include administering an immunoconjugate containing an antibody construct that is capable of binding a tumor-associated antigen (TAA), or tumors over-expressing a TAA


15 In some embodiments, the cancer is susceptible to a pro-inflammatory response induced by TLR7 and/or TLR8.

20 In some embodiments, a therapeutically effective amount of an immunoconjugate is administered to a patient in need to treat cervical cancer, endometrial cancer, ovarian cancer, prostate cancer, pancreatic cancer, esophageal cancer, bladder cancer, urinary tract cancer, urothelial carcinoma, lung cancer, non-small cell lung cancer, Merkel cell carcinoma, colon cancer, colorectal cancer, gastric cancer, or breast cancer. The Merkel cell carcinoma cancer may be metastatic Merkel cell carcinoma. The breast cancer may be triple-negative breast cancer. The esophageal cancer may be gastroesophageal junction adenocarcinoma.

25 EXAMPLES

Example L-1 Synthesis of 4-[3-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[4-[[2-amino-4-[ethoxy(propyl) carbamoyl]-3H-1-benzazepin-8-yl)sulfonyl]benzoyl]amino]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]propanoyloxy]-2,3,5,6-tetrafluoro-benzenesulfonic acid, 8SO2BzL-1





Preparation of 8-bromo-N-ethoxy-N-propyl-2-(tritylamino)-3H-1-benzazepine-4-carboxamide, 8SO2BzL-1b

To a mixture of 8-bromo-2-(tritylamino)-3H-1-benzazepine-4-carboxylic acid, 8SO2BzL-1a (1 g, 1.91 mmol, 1 *eq*) and N-ethoxypropan-1-amine (320 mg, 2.29 mmol, 1.2 *eq*, HCl) in DCM (15 mL) and DMA (5 mL) was added EDCI (1.10 g, 5.73 mmol, 3.0 *eq*) in one portion at 25°C, and then stirred at 25°C for 0.5 h. The mixture was concentrated to remove DCM. Then the residue was diluted with aq. NaHCO₃ until the pH to between 8 and 9. The mixture was extracted with EtOAc (30 mL x 3). The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (column height: 250 mm, diameter: 100 mm, 100-200 mesh silica gel, Petroleum ether/Ethyl acetate=1/0,3/1) to afford 8SO2BzL-1b (0.7 g, 1.15 mmol, 60.21% yield) as white solid. ¹H NMR (CDCl₃, 400 MHz) δ 7.38-7.15 (m, 15H), 7.06-6.94 (m, 2H), 6.72 (s, 1H), 6.16 (s, 1H), 4.03-3.83 (m, 2H), 3.73 (t, J=7.2 Hz, 2H), 2.78 (s, 2H), 1.89-1.64 (m, 2H), 1.25 (t, J=7.2 Hz, 3H), 0.99 (t, J=7.2 Hz, 3H). LC/MS [M+H]⁺ 608.2 (calculated); LC/MS [M+H]⁺ 608.2 (observed).

Preparation of methyl 4-[[4-[ethoxy(propyl)carbamoyl]-2-(tritylamino)-3H-1-benzazepin-8-yl]sulfanyl]benzoate, 8SO2BzL-1c

To a mixture of 8SO2BzL-1b (0.35 g, 575 μmol, 1.0 *eq*) and methyl 4-sulfanylbenzoate (116 mg, 690 μmol, 1.2 *eq*) in DMF (4 mL) was added dicyclohexyl[2',4',6'-tris(propan-2-yl)[1,1'-biphenyl]phosphane, XPhos, CAS Reg. No. 564483-18-7, Huang, X., et al (2003) *J. Am. Chem. Soc.* 125(22):6653-6655; Bruno, N.C. et al, (2013) *Chemical Science*, 4(3):916-920, (82.3 mg, 173 μmol, 0.3 *eq*), Cs₂CO₃ (375 mg, 1.15 mmol, 2.0 *eq*) and [2-(2-aminophenyl)phenyl]-chloro-palladium;dicyclohexyl-[3-(2,4,6-triisopropyl phenyl)phenyl]phosphane, Pd-Xphos-G2, CAS Reg. No. 1310584-14-5 (226 mg, 288 μmol, 0.5 *eq*) in one portion at 25 °C under N₂, and then stirred at 120 °C for 12 h. The mixture was

diluted water (20 mL) and extracted with EtOAc (10 mL x 3). The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (column height: 250 mm, diameter: 100 mm, 100-200 mesh silica gel, Petroleum ether/Ethyl acetate=1/0, 3/1) to afford 8SO2BzL-1c (0.25 g, 359.26 umol, 62.47% yield). LC/MS [M+H]⁺ 696.3 (calculated); LC/MS [M+H]⁺ 696.2 (observed).

Preparation of methyl 4-[[4-[ethoxy(propyl)carbamoyl]-2-(tritylamino)-3H-1-benzazepin-8-yl]sulfonyl]benzoate, 8SO2BzL-1d

To a mixture of 8SO2BzL-1c (0.25 g, 359 umol, 1.0 eq) in DCM (2 mL), THF (2 mL) and H₂O (2 mL) was added potassium peroxydisulfate, KHSO₅, Oxone (663 mg, 1.08 mmol, 3.0 eq) in one portion at 25°C, and then stirred at 25 °C for 12 h. The mixture was diluted with water and extracted with EtOAc (20 mL x 3). The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (column height: 250 mm, diameter: 100 mm, 100-200 mesh silica gel, Petroleum ether/Ethyl acetate=1/0, 1/1) to afford 8SO2BzL-1d (0.2 g, 274.78 umol, 76.48% yield) as yellow oil.

LC/MS [M+H]⁺ 728.27 (calculated); LC/MS [M+H]⁺ 728.2 (observed).

Preparation of methyl 4-[[2-amino-4-[ethoxy(propyl)carbamoyl]-3H-1-benzazepin-8-yl]sulfonyl]benzoate, 8SO2BzL-1e

To a mixture of 8SO2BzL-1d (0.1 g, 137 umol, 1.0 eq) in DCM (4 mL) was added TFA (313 mg, 2.75 mmol, 203 uL, 20.0 eq) in one portion at 25°C, and then stirred at 50 °C for 12 h.

The mixture was concentrated in vacuum to give a residue, the residue was purified by prep-HPLC(column: Phenomenex Luna 80*30mm*3um;mobile phase: [water(0.1%TFA)-acetonitrile, ACN];B%: 15%-35%,8min) to give 8SO2BzL-1e (0.046 g, 94.74 umol, 68.96% yield) as white solid. ¹H NMR (CDCl₃, 400 MHz) δ 8.21 (d, J = 8.4 Hz, 2H), 8.10 (d, J = 8.4 Hz, 2H), 8.04 (s, 1H), 7.95-7.87 (m, 1H), 7.57 (d, J = 8.0 Hz, 1H), 7.49 (s, 1H), 6.97 (d, J = 1.2 Hz, 1H), 3.95 (s, 3H), 3.93-3.82 (m, 2H), 3.70 (t, J = 7.2 Hz, 2H), 3.23 (s, 2H), 1.82-1.66 (m, 2H), 1.21 (t, J = 7.2 Hz, 3H), 0.96 (t, J = 7.2 Hz, 3H). LC/MS [M+H]⁺ 485.16 (calculated); LC/MS [M+H]⁺ 486.1 (observed).

Preparation of 4-[[2-amino-4-[ethoxy(propyl)carbamoyl]-3H-1-benzazepin-8-yl]sulfonyl]benzoic acid, 8SO2BzL-1f

To a mixture of 8SO2BzL-1e (0.24 g, 494 umol, 1.0 eq) in MeOH (2 mL), H₂O (2 mL) and THF (2 mL) was added LiOH.H₂O (62.2 mg, 1.48 mmol, 3.0 eq) in one portion at 25°C, and then stirred at 25°C for 2 h. The mixture was quenched with HCl (1 M) to adjust pH to between 6 and 7 and the aqueous phase was extracted with EtOAc (10 mL x 3). The combined organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated to give 8SO2BzL-1f (0.23 g, crude) as yellow oil. ¹H NMR (DMSO, 400 MHz) δ 8.16-8.06 (m, 4H),

7.61-7.54 (m, 2H), 7.49 (dd, $J = 1.6, 8.0$ Hz, 1H), 7.11 (s, 1H), 3.82 (q, $J = 7.2$ Hz, 2H), 3.59 (t, $J = 7.2$ Hz, 2H), 2.89 (s, 2H), 1.70-1.55 (m, 2H), 1.01 (t, $J = 7.2$ Hz, 3H), 0.89 (t, $J = 7.2$ Hz, 3H).
LC/MS [M+H]⁺ 472.1 (calculated); LC/MS [M+H]⁺ 472.1 (observed).

Preparation of tert-butyl 3-[2-[2-[2-[2-[2-[2-[2-[2-[2-[4-[[2-amino-4-[ethoxy(propyl) 5 carbamoyl]-3H-1-benzazepin-8-yl)sulfonyl]benzoyl]amino]ethoxy]ethoxy]ethoxy]ethoxy] ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]propanoate. 8SO2BzL-1g

To a mixture of 8SO2BzL-1f (0.2 g, 424 umol, 1.0 eq) in DMF (4 mL) was added tert-butyl 3-[2-[2-[2-[2-[2-[2-(2-aminoethoxy)ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy] propanoate (248 mg, 424 umol, 1.0 eq), DIEA (164 mg, 1.27 mmol, 222 uL, 3.0 eq), and Hexafluorophosphate Azabenzotriazole Tetramethyl Uronium, HATU, CAS Reg. No. 148893-10-1 (177 mg, 467 umol, 1.1 eq) in one portion at 25°C, and then stirred at 25°C for 0.5 h. The mixture was diluted with water and extracted with EtOAc (30 mL x 3). The organic layer was washed with brine, dried over Na2SO4, filtered and concentrated to give 8SO2BzL-1g (0.5 g, crude) as yellow oil. LC/MS [M+H]⁺ 1039.5 (calculated); LC/MS [M+H]⁺ 1039.5(observed).

Preparation of 3-[2-[2-[2-[2-[2-[2-[2-[2-[2-[4-[[[2-amino-4-[ethoxy(propyl)
carbamoyl]-3H-1-benzazepin-8-yl)sulfonyl]benzoyl]amino]ethoxy]ethoxy]ethoxy]
ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]propanoic acid, 8SO₂BzL-1h

To a mixture of 8SO2BzL-1g (0.5 g, 481 μ mol, 1.0 *eq*) in CH₃CN (1 mL) and H₂O (3 mL) was added TFA (439 mg, 3.85 mmol, 285 μ L, 8.0 *eq*) in one portion at 25°C, and then stirred at 80 °C for 1 hours. The mixture was concentrated to give a residue. The residue was purified by prep-HPLC(column: Phenomenex Luna 80*30mm*3 μ m; mobile phase: [water(0.1%TFA)-ACN];B%: 10%-35%,8min) to give 8SO2BzL-1h (0.15 g, 152.57 μ mol, 31.71% yield) as yellow oil. ¹H NMR (MeOD, 400 MHz) δ 8.14-8.08 (m, 2H), 8.06-7.95 (m, 4H), 7.78 (d, J = 8.4 Hz, 1H), 7.40 (s, 1H), 3.94 (q, J = 6.8 Hz, 2H), 3.75-3.70 (m, 4H), 3.65-3.50 (m, 40H), 3.40 (s, 2H), 2.53 (t, J = 6.4 Hz, 2H), 1.82-1.69 (m, 2H), 1.16 (t, J = 7.2 Hz, 3H), 0.98 (t, J = 7.2 Hz, 3H). LC/MS [M+H]⁺ 982.45 (calculated); LC/MS [M+H]⁺ 983.4 (observed).


Preparation of 8SO2BzL-1

To a mixture of 8SO2BzL-1h (0.15 g, 152.57 μmol , 1.0 *eq*) in DCM (3 mL) and DMA (0.5 mL) was added sodium,2,3,5,6-tetrafluoro-4-hydroxy-benzenesulfonate (164 mg, 610 μmol , 4.0 *eq*) and EDCI (146 mg, 763 μmol , 5.0 *eq*) in one portion at 25°C, and then stirred at 25°C for 0.5 h. The mixture was concentrated to give a residue. The residue was purified by prep-HPLC(column: Phenomenex Luna 80*30mm*3 μm ,mobile phase: [water(0.1%TFA)-ACN];B%: 15%-40%,8min) to give 8SO2BzL-1 (58.9 mg, 48.63 μmol , 31.87% yield) as light yellow solid.

¹H NMR (MeOD, 400 MHz) δ 8.11-8.06 (m, 2H), 8.05-7.99 (m, 3H), 7.95 (dd, J = 2.0, 8.4 Hz,

1H), 7.77 (d, J = 8.4 Hz, 1H), 7.39 (s, 1H), 3.93 (q, J = 7.2 Hz, 2H), 3.86 (t, J = 6.0 Hz, 2H), 3.71 (t, J = 7.2 Hz, 2H), 3.67-3.53 (m, 34H), 3.53-3.48 (m, 6H), 3.42 (s, 2H), 2.97 (t, J = 6.0 Hz, 2H), 1.83-1.65 (m, 2H), 1.15 (t, J = 7.2 Hz, 3H), 0.97 (t, J = 7.2 Hz, 3H). LC/MS [M+H] 1211.4 (calculated); LC/MS [M+H] 1211.3 (observed).

Example L-2 Synthesis of 2-amino-4-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-(2,5-dioxopyrrol-1-yl)acetyl]amino]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxycarbonyl]amino]ethoxy-propyl-carbamoyl]-3H-1-benzazepine-8-sulfonic acid. 8SO2BzL-2



Preparation of 2-amino-4-[2-(tert-butoxycarbonylamino)ethoxy-propyl -carbamoyl]-3H-1-benzazepine-8-sulfonic acid, 8SO2BzL-2b

To a mixture of tert-butyl N-[2-[(2-amino-8-benzylsulfanyl-3H-1-benzazepine-4-carbonyl)-propyl-amino]oxyethyl]carbamate, 8SO2BzL-2a (0.15 g, 286 μ mol, 1.0 eq) in H₂O (0.15 mL) and AcOH (0.5 mL) was added N-chlorosuccinimide, NCS (153 mg, 1.14 mmol, 4.0 eq) in one portion at 25 °C, and then stirred at 25 °C for 1 h. The mixture was diluted with aq. NaHCO₃ to adjust pH between 7 and 8. Then the mixture was extracted with EtOAc (10 mL x 3). The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by prep-HPLC (column: Phenomenex Luna 80*30mm*3 μ m; mobile

phase: [water(TFA)-ACN];B%: 5%-35%,8min) to give 8SO2BzL-2b (11.5 mg, 23.83 μ mol, 8.34% yield) as yellow solid. ^1H NMR (DMSO, 400 MHz) δ 11.87 (s, 1H), 9.78 (s, 1H), 8.81 (s, 1H), 7.66 (s, 1H), 7.55 (s, 2H), 7.34-7.27 (m, 1H), 3.92-3.78 (m, 2H), 3.63 (t, J = 7.2 Hz, 2H), 3.31 (s, 2H), 3.16-3.01 (m, 2H), 1.70-1.60 (m, 2H), 1.36 (s, 9H), 0.89(t, J = 7.2 Hz, 3H). LC/MS [M+H] 483.2(calculated); LC/MS [M+H] 483.1 (observed).


Preparation of 2-amino-4-[2-aminoethoxy(propyl)carbamoyl]-3H-1-benzazepine-8-sulfonic acid, 8SO2BzL-2c


To a solution of 8SO2BzL-2b (150 mg, 310.85 μ mol, 1 eq) in EtOAc (10.0 mL) was added HCl/EtOAc (4 M, 20.0 mL, 257 eq), and then stirred at 25 °C for 1 h. The mixture was concentrated to give 8SO2BzL-2c (200 mg, crude) as white solid. LC/MS [M+H] 383.1(calculated); LC/MS [M+H] 383.2 (observed).

Preparation of 8SO2BzL-2

To a solution of 8SO2BzL-2c (70.0 mg, 167.11 μ mol, 1 eq, HCl) in DMF (1.00 mL) was added DIEA (90.0 mg, 668 μ mol, 120 μ L, 4 eq) and 2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-(2,5-dioxypyrrol-1-yl)acetyl]amino]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethyl (4-nitrophenyl) carbonate (70.0 mg, 83.5 μ mol, 0.5 eq), and then stirred at 0 °C for 1 h. The mixture was filtered and purified by prep-HPLC (column: Phenomenex Luna 80*30mm*3 μ m;mobile phase: [water(TFA)-ACN];B%: 1%-30%,8min) to give 8SO2BzL-2 (15 mg, 12.92 μ mol, 7.73% yield, TFA) as light yellow oil. ^1H NMR (MeOD, 400 MHz) δ 7.89-7.77 (m, 2H), 7.66 (d, J = 8.0 Hz, 1H), 7.42 (s, 1H), 6.89 (s, 2H), 4.17 (s, 2H), 3.97 (br t, J = 4.8 Hz, 2H), 3.86-3.79 (m, 2H), 3.75 (t, J = 7.2 Hz, 2H), 3.66-3.58 (m, 38H), 3.56-3.49 (m, 4H), 3.42-3.35 (m, 4H), 1.81-1.73 (m, 2H), 1.00 (t, J = 7.6 Hz, 3H). LC/MS [M+H] 1047.4(calculated); LC/MS [M+H] 1047.7 (observed).

Example L-3 Synthesis of 2-amino-8-(N-(1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl)-N-methylsulfamoyl)-N-ethoxy-N-propyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-3





Preparation of 2-amino-N-ethoxy-8-((4-methoxybenzyl)thio)-N-propyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-3b

- 5 To a mixture of 2-amino-8-bromo-N-ethoxy-N-propyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-3a (0.96 g, 1.99 mmol) and 4-methoxy- α -toluenethiol (0.37 g, 2.39 mmol) in dioxane (10 mL), was added 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene, Xantphos, CAS Reg. No. 161265-03-8 (0.06 g, 0.10 mmol), Pd₂(dba)₃ (0.05 g, 0.05 mmol), and then triethylamine (0.56 mL, 3.99 mmol). The mixture was heated to reflux for 1 h then cooled.
- 10 The solvent was removed by evaporation and the crude product was purified by silica gel chromatography using a gradient of 1-10% MeOH/DCM over 12 column volumes to give 8SO2BzL-3b (0.72 g, 82%).

Preparation of 2-amino-4-(ethoxy(propyl)carbamoyl)-3H-benzo[b]azepine-8-sulfonyl chloride, 8SO2BzL-3c

- 15 To a solution of 8SO2BzL-3b (0.72 g, 1.64 mmol) in acetonitrile/water (9:1), 10 mL at 10 deg C was added in portions N-chlorosuccinimide (0.66 g, 4.91 mmol). After addition was complete, stirred for an additional 20 min to give 8SO2BzL-3c crude product solution, used as is in the next step.

Preparation of tert-butyl (2-((2-amino-4-(ethoxy(propyl)carbamoyl)-3H-benzo[b]azepin-8-yl)sulfonyl)-5,8,11,14,17,20,23,26,29,32-decaoxa-2-azatetatriacontan-34-yl)carbamate, 8SO2BzL-3d

5 An aliquot of 8SO2BzL-3c (1.00 mL, 0.16 mmol) was added dropwise to a stirring mixture of *tert*-butyl (5,8,11,14,17,20,23,26,29,32-decaoxa-2-azatetatriacontan-34-yl)carbamate (0.12 g, 0.19 mmol) and triethylamine (0.09 mL, 0.64 mmol) in acetonitrile (3 mL). After 15 min the reaction was concentrated and purified by reverse phase chromatography using a gradient of 10-90% ACN/water (+0.1% TFA) over 10 min to obtain 8SO2BzL-3d (0.08 g, 51%).


10 Preparation of 2-amino-8-(N-(32-amino-3,6,9,12,15,18,21,24,27,30-decaoxadotriacontyl)-N-methylsulfamoyl)-N-ethoxy-N-propyl-3H-benzo[b]azepine-4-carboxamide hydrogen chloride, 8SO2BzL-3e

To a solution of 8SO2BzL-3d (0.08 g, 0.08 mmol) in ACN (3 mL) was added aq. HCl (6M, 3 mL) and the mixture was stirred at room temp. for 45 min. The solvent was removed
15 and the isolated syrup was azeotroped with ACN (3 mL) to provide the HCl salt of 8SO2BzL-3e (0.06 g, 85%) as a hazy, white film.

Preparation of 8SO2BzL-3

To a solution of 8SO2BzL-3e HCl (0.06 g, 0.07 mmol) in DMF (3 mL) was added triethylamine (0.04 mL, 0.28 mmol). 2,5-Dioxopyrrolidin-1-yl 2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)acetate (0.02 g, 0.08 mmol) was added in parts. After completion of addition, acetic
20 acid (9 μ L) was added and the solvent was removed by vacuum. After purification by reverse phase HPLC, 8SO2BzL-3 (0.03 g, 48%) was obtained as a clear oil after evaporation of solvent. LC/MS [M+H]⁺ 1001.48 (calculated); LC/MS [M+H]⁺ 1074.88 (observed).

Example L-4 Synthesis of 2-amino-8-(N-(1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl)sulfamoyl)-N-ethoxy-N-propyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-4
25



Preparation of tert-butyl (32-((2-amino-4-(ethoxy(propyl)carbamoyl)-3H-benzo[b]azepine)-8-sulfonyl)-3,6,9,12,15,18,21,24,27,30-decaoxadotriacontyl)carbamate, 8SO2BzL-4b

To a solution of tert-butyl (32-amino-3,6,9,12,15,18,21,24,27,30-decaoxadotriacontyl)carbamate, Boc-amino-PEG10-amine (0.10 g, 0.16 mmol) and DIPEA (0.14 mL, 0.80 mmol) in DMF (4 mL) was added a solution of 2-amino-4-(ethoxy(propyl)carbamoyl)-3H-benzo[b]azepine-8-sulfonyl chloride, 8SO2BzL-4a (0.16 M, 1.00 mL, 0.16 mmol) in DMF. After 20 min the reaction was concentrated and then purified on reverse phase HPLC using a gradient of 10-90% ACN/water over 10 min to give 8SO2BzL-4b (0.07 g, 47%) after removal of solvent.

Preparation of 2-amino-8-(N-(32-amino-3,6,9,12,15,18,21,24,27,30-decaoxadotriacontyl)sulfamoyl)-N-ethoxy-N-propyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-4c


To a solution of 8SO2BzL-4b (0.07 g, 0.07 mmol) in acetonitrile (3 mL) was added aq. HCl (6M, 3 mL) and the mixture was stirred at RT for 45 min. The solvent was removed and the isolated syrup was azeotroped with ACN (3 mL) to provide 8SO2BzL-4c HCl salt (0.06 g, 82%) as a hazy, white film.

5 Preparation of 8SO2BzL-4


To a solution of 8SO2BzL-4c HCl (0.06 g, 0.06 mmol) in DMF (3 mL) was added triethylamine (0.03 mL, 0.25 mmol). 2,5-Dioxopyrrolidin-1-yl 2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)acetate (0.02 g, 0.07 mmol) was added in parts. After completion of addition, acetic acid (9 μ L) was added and the solvent was removed by vacuum. After purification by reverse
 10 phase HPLC, 8SO2BzL-4 (0.04 g, 69%) was obtained after solvent removal. LC/MS [M+H]⁺ 987.45 (calculated); LC/MS [M+H]⁺ 987.86 (observed).

Example L-5

Synthesis of 1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl 2-((2-amino-N-propyl-8-sulfamoyl-3H-benzo[b]azepine-4-carboxamido)oxy)ethyl)carbamate, 8SO2BzL-5



15



Preparation of tert-butyl (2-((2-amino-8-((4-methoxybenzyl)thio)-N-propyl-3H-benzo[b]azepine-4-carboxamido)oxy)ethyl)carbamate, 8SO2BzL-b

To a mixture of tert-butyl (2-((2-amino-8-bromo-N-propyl-3H-benzo[b]azepine-4-carboxamido)oxy)ethyl)carbamate, 8SO2BzL-5a (0.96 g, 1.99 mmol) and 4-methoxy- α -toluenethiol (0.37 g, 2.39 mmol) in dioxane (10 mL), added Xantphos (0.06 g, 0.10 mmol), Pd₂(dba)₃ (0.05 g, 0.05 mmol), then triethylamine (0.56 mL, 3.99 mmol). The mixture was heated to reflux for 1 h then cooled. The solvent was removed by evaporation and the crude product was purified by silica gel chromatography using a gradient of 1-10% MeOH/DCM over 12 column volume to give 8SO2BzL-5b (0.78 g, 71%) as a yellow solid.

Preparation of tert-butyl (2-((2-amino-8-(chlorosulfonyl)-N-propyl-3H-benzo[b]azepine-4-carboxamido)oxy)ethyl)carbamate, 8SO2BzL-5c

To a solution of 8SO2BzL-5b (0.78 g, 1.41 mmol) in acetonitrile/water (9:1, 10 mL), at 0 deg C was added in three equal portions N-chlorosuccinimide (0.56 g, 4.22 mmol). After the addition was complete, the product 8SO2BzL-5c was used as is without further purification.

Preparation of tert-butyl (2-((2-amino-N-propyl-8-sulfamoyl-3H-benzo[b]azepine-4-carboxamido)oxy)ethyl)carbamate, 8SO2BzL-5d

To a solution of 8SO2BzL-5c (2.00 mL, 0.28 mmol) in acetonitrile/water (9:1) at 0 deg was added ammonium hydroxide solution (0.20 mL, 1.66 mmol). After 10 minutes the solvent was removed, and the crude product was purified by reverse phase HPLC using a gradient of 10-90% acetonitrile/water to give 8SO2BzL-5d (0.06 g, 41%) as a yellow film after evaporation of solvent.

Preparation of 8SO2BzL-5e


A solution of 8SO2BzL-5e (0.06 g, 0.11 mmol) in acetonitrile (2 mL) and 6 N HCl (2.00 mL, 12.00 mmol) was stirred at room temperature for 45 minutes. The solvent was removed by vacuum to give 8SO2BzL-5e as the HCl salt (0.05 g, 101%).


Preparation of 8SO2BzL-5

- 5 To a solution of 8SO2BzL-5e (0.04 g, 0.10 mmol) in DMF (4 mL) at room temperature was added triethylamine (0.06 mL, 0.40 mmol). To this mixture was added a solution of 1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl (4-nitrophenyl) carbonate, PNPC-PEG10-Mal (0.08 g, 0.10 mmol) in DMF (2 mL). After 20 min acetic acid (6 uL) was added and reaction was concentrated under vacuum and purified by reverse phase HPLC using a gradient of 10-90% ACN/water (+0.1% TFA) over 10 min to give 8SO2BzL-5 (0.04 g, 37%) after concentration of pure fractions. LC/MS [M+H]⁺ 1046.45 (calculated); LC/MS [M+H]⁺ 1046.88 (observed).

Example L-6

- Synthesis of 1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl (2-((2-amino-8-(N,N-dimethylsulfamoyl)-N-propyl-3H-benzo[b]azepine-4-carboxamido)oxy)ethyl)carbamate, 8SO2BzL-6





Preparation of tert-butyl N-[2-[(2-amino-8-benzylsulfanyl-3H-1-benzazepine-4-carbonyl)-propyl-amino]oxyethyl]carbamate, 8SO2BzL-6b

To a mixture of tert-butyl N-[2-[(2-amino-8-bromo-3H-1-benzazepine-4-carbonyl)-propyl-amino]oxyethyl]carbamate, 8SO2BzL-6a (0.5 g, 1.04 mmol, 1.0 *eq*) and benzylthiol, benzylmercaptan, phenylmethanethiol, BnSH, CAS Reg. No. 100-53-8 (155 mg, 1.25 mmol, 146.05 μ L, 1.2 *eq*) in dioxane (15 mL) was added 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene, Xantphos, CAS Reg. No. 161265-03-8 (120 mg, 208 μ mol, 0.2 *eq*) tris(dibenzylideneacetone)dipalladium, Pd₂(dba)₃, CAS Reg. No. 51364-51-3 (190 mg, 208 μ mol, 0.2 *eq*) and diisopropylethylamine, DIEA (268 mg, 2.08 mmol, 362 μ L, 2.0 *eq*) in one portion at 25 °C under N₂, and then stirred at 110°C for 2 h. The mixture was diluted with water (20 mL) and extracted with EtOAc (10 mL x 3). The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The mixture was further purification by prep-HPLC (column: Phenomenex Luna 80*30mm*3 μ m; mobile phase: [water(TFA)-ACN]; B%: 25%-55%, 8min) to give 8SO2BzL-6b (0.5 g, 952.97 μ mol, 91.75% yield) as yellow solid. ¹H NMR (MeOD, 400 MHz) δ 7.49 (d, J = 8.4 Hz, 1H), 7.43-7.38 (m, 3H), 7.36-7.22 (m, 5H), 4.30 (s, 2H), 3.91 (t, J = 5.2 Hz, 2H), 3.73 (t, J = 7.2 Hz, 2H), 3.32 (s, 2H), 3.24 (t, J = 5.2 Hz, 2H), 1.81-1.70 (m, 2H), 1.34 (s, 9H), 0.98 (t, J = 7.2 Hz, 3H). LC/MS [M+H] 525.2 (calculated); LC/MS [M+H] 525.2 (observed).

Preparation of tert-butyl N-[2-[[2-amino-8-(dimethylsulfamoyl)-3H-1-benzazepine-4-carbonyl]-propyl-amino]oxyethyl]carbamate, 8SO2BzL-6c

To a solution of 8SO2BzL-6b (50.0 mg, 95.30 μ mol, 1 *eq*) in CH₃CN (1.00 mL) and H₂O (0.10 mL) was added AcOH (60.0 mg, 953 μ mol, 50.0 μ L, 10 *eq*), N-chlorosuccinimide, NCS

(50.0 mg, 381 μ mol, 4 eq) at 25°C, and then stirred at this temperature for 10 min, then N-methylmethanamine;hydrochloride, dimethylamine HCl (80.0 mg, 953 μ mol, 10 eq) and DIEA (250 mg, 1.91 mmol, 330 μ L, 20 eq) was added. The mixture was stirred at 0 °C for another 1h. The mixture was filtered and purified by prep-HPLC (column: Phenomenex Luna

5 80*30mm*3 μ m;mobile phase: [water(TFA)-ACN];B%: 10%-40%,8min) to give 8SO2BzL-6c (12 mg, 23.55 μ mol, 24.71% yield) as white solid. ¹H NMR (MeOD, 400 MHz) δ 7.92-7.72 (m, 3H), 7.50 (s, 1H), 3.94 (t, J = 5.2 Hz, 2H), 3.75 (t, J = 7.2 Hz, 2H), 3.44 (s, 2H), 3.26 (br t, J = 5.2 Hz, 2H), 2.77 (s, 6H), 1.77 (sxt, J = 7.2 Hz, 2H), 1.37 (s, 9H), 0.99 (t, J = 7.2 Hz, 3H). LC/MS [M+H] 510.2 (calculated); LC/MS [M+H] 510.3 (observed).


10 Preparation of 2-amino-N-(2-aminoethoxy)-8-(N,N-dimethylsulfamoyl)-N-propyl-3H-benzo[b]azepine-4-carboxamide hydrogen chloride, 8SO2BzL-6d

A solution of 8SO2BzL-6c (0.04 g, 0.08 mmol) in acetonitrile (2 mL) and 6 N HCl (1.41 mL, 8.46 mmol) was stirred at room temperature for 45 minutes. The solvent was removed by vacuum to give 8SO2BzL-6d (0.04 g, 100%).

15 Preparation of 8SO2BzL-6

To a solution of 8SO2BzL-6d (0.04 g, 0.10 mmol) in DMF (4 mL) at room temperature added triethylamine (0.04 mL, 0.28 mmol). To this mixture was added a solution 1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl (4-nitrophenyl)carbonate (0.06 g, 0.07 mmol) in DMF (2 mL). After 20 min acetic acid (6
20 μ L) was added and reaction was concentrated under vacuum and purified by reverse phase HPLC using a gradient of 10-90% ACB/water (+0.1% TFA) over 10 min to give 8SO2BzL-6 (0.04 g, 51%) after concentration of pure fractions. LC/MS [M+H] 1074.48 (calculated); LC/MS [M+H] 1074.90 (observed).

Example L-9 Synthesis of 2-amino-8-(N-(1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-
25 1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl)sulfamoyl)-N,N-dipropyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-9



Preparation of 2-amino-8-((4-methoxybenzyl)thio)-N,N-dipropyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-9b

- 5 To a mixture of 2-amino-8-bromo-N,N-dipropyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-9a (0.96 g, 1.99 mmol) and 4-methoxy- α -toluenethiol (0.37 g, 2.39 mmol) in dioxane (10 mL), added Xantphos (0.06 g, 0.10 mmol), $\text{Pd}_2(\text{dba})_3$ (0.05 g, 0.05 mmol), then triethylamine (0.56 mL, 3.99 mmol). The mixture was heated to reflux for 1 h then cooled. The solvent was removed by evaporation and the crude product was purified by silica gel

chromatography using a gradient of 1-10% MeOH/DCM over 12 column volumes to give 8SO2BzL-9b (0.69 g, 79%).

Preparation of 2-amino-4-(dipropylcarbamoyl)-3H-benzo[b]azepine-8-sulfonyl chloride, 8SO2BzL-9c

- 5 To a solution of 8SO2BzL-9b (0.68 g, 1.55 mmol) in acetonitrile/water (9:1), 10 mL at 10 deg C was added N-chlorosuccinimide, NCS in thirds (0.62 g, 4.66 mmol). After addition was complete, stirred for an additional 20 min. to give 8SO2BzL-9c.

Preparation of tert-butyl (32-((2-amino-4-(dipropylcarbamoyl)-3H-benzo[b]azepine)-8-sulfonamido)-3,6,9,12,15,18,21,24,27,30-decaoxadotriacontyl)carbamate, 8SO2BzL-9d

- 10 An aliquot of 8SO2BzL-9c (1.00 mL, 0.15 mmol) obtained previously was added drop-wise to a stirring mixture of *tert*-butyl (32-amino-3,6,9,12,15,18,21,24,27,30-decaoxadotriacontyl)carbamate (0.11 g, 0.18 mmol) and triethylamine (0.08 mL, 0.60 mmol) in acetonitrile (3 mL). After 15 min the reaction was concentrated and purified by reverse phase chromatography using a gradient of 10-90% ACN/water (+0.1% TFA) over 10 min to 15 8SO2BzL-9d (0.07 g, 51%).


Preparation of 2-amino-8-(N-(32-amino-3,6,9,12,15,18,21,24,27,30-decaoxadotriacontyl)sulfamoyl)-N,N-dipropyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-9e

- To a solution of 8SO2BzL-9d (0.07 g, 0.08 mmol) in acetonitrile (3 mL) was added aq. 20 HCl (6M, 3 mL) and the mixture was stirred at RT for 45 min. The solvent was removed and the isolated syrup was azeotroped with acetonitrile (3 mL) to provide 8SO2BzL-9e as the HCl salt (0.06 g, 86%) as a hazy, white film.

Preparation of 8SO2BzL-9

- To a solution of 8SO2BzL-9e HCl (0.06 g, 0.06 mmol) in DMF (3 mL) was added 25 triethylamine (0.03 mL, 0.25 mmol). 2,5-Dioxopyrrolidin-1-yl 2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)acetate (0.02 g, 0.07 mmol) was added in parts. After completion of addition, acetic acid (9 uL) was added and the solvent was removed by vacuum. After purification by reverse phase HPLC to give 8SO2BzL-9 (0.03 g, 43%) after evaporation. LC/MS [M+H] 985.47 (calculated); LC/MS [M+H] =985.88 (observed).

- 30 Example L-10 Synthesis of 2-amino-8-(N-(1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl)-N-methylsulfamoyl)-N,N-dipropyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-10



Preparation of tert-butyl (2-((2-amino-4-(dipropylcarbamoyl)-3H-benzo[b]azepin-8-yl)sulfonyl)-5,8,11,14,17,20,23,26,29,32-decaoxa-2-azatetatriacontan-34-yl)carbamate,

5 8SO2BzL-10b

An solution of 2-amino-4-(dipropylcarbamoyl)-3H-benzo[b]azepine-8-sulfonyl chloride, 8SO2BzL-10a (1.00 mL, 0.15 mmol) in acetonitrile was added dropwise to a stirring mixture of tert-butyl (5,8,11,14,17,20,23,26,29,32-decaoxa-2-azatetatriacontan-34-yl)carbamate (0.11 g, 0.18 mmol) and triethylamine (0.08 mL, 0.60 mmol) in acetonitrile (3 mL). After 15 min the reaction was concentrated and purified by reverse phase chromatography using a gradient of 10-90% ACN/water (+0.1% TFA) over 10 min to obtain 8SO2BzL-10b (0.05 g, 37%).

Preparation of 2-amino-8-(N-(32-amino-3,6,9,12,15,18,21,24,27,30-decaoxadotriacontyl)-N-methylsulfamoyl)-N,N-dipropyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-10c

To a solution of 8SO2BzL-10b (0.05 g, 0.05 mmol) in acetonitrile (3 mL) was added aq. HCl (6M, 3 mL) and the mixture was stirred at room temperature, RT for 45 min. The solvent was removed and the isolated syrup was azeotroped with ACN (3 mL) to provide 8SO2BzL-10c as the HCl salt (0.04 g, 91%) as a hazy, white film.

Preparation of 8SO2BzL-10


To a solution of 8SO2BzL-10c HCl (0.04 g, 0.05 mmol) in DMF (3 mL) was added triethylamine (0.03 mL, 0.20 mmol). 2,5-Dioxopyrrolidin-1-yl 2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)acetate (0.02 g, 0.06 mmol) was added in parts. After completion of addition, acetic


acid (9 μ L) was added and the solvent was removed by vacuum. After purification by reverse phase HPLC, 8SO2BzL-10 (0.02 g, 42%) was obtained after evaporation. LC/MS [M+H]⁺ 999.49 (calculated); LC/MS [M+H]⁺ 999.92 (observed).

Example L-12

Synthesis of 2-amino-6-[5-[3-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-

5 (2,5-dioxopyrrol-1-yl)acetyl]amino[ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]propanoylamino]pentyl]-4-[ethoxy(propyl)carbamoyl]-3H-1-benzazepine-8-sulfonic acid,
8SO₂BzL-12





Preparation of 2-amino-8-bromo-N-ethoxy-6-iodo-N-propyl-3H-1-benzazepine-4-carboxamide, 8SO2BzL-12b

To a solution of 2-amino-8-bromo-6-iodo-3H-1-benzazepine-4-carboxylic acid, 8SO2BzL-12a (2.0 g, 4.91 mmol, 1.0 eq) in DCM (20 mL) and DMA (10 mL) was added methanesulfonic acid, $\text{CH}_3\text{SO}_3\text{H}$ (472 mg, 4.91 mmol, 350 μL , 1.0 eq), N-ethoxypropan-1-amine (823 mg, 5.90 mmol, 1.2 eq, HCl) and EDCI (3.77 g, 19.7 mmol, 4 eq). The mixture was stirred at 25°C for 2 hrs. The pH of the reaction mixture was adjusted to ~9 with sat. Na_2CO_3 . The aqueous phase was extracted with ethyl acetate (50 mL x 3). The combined organic phase was washed with brine (20 mL x 2), dried with anhydrous Na_2SO_4 , filtered and concentrated in vacuum. The crude product was triturated with EtOAc at 25 °C for 10 min to afford 8SO2BzL-12b (1.24 g, 2.52 mmol, 51.3% yield) as yellow solid. LC/MS $[\text{M}+\text{H}]$ 491.97 (calculated); LC/MS $[\text{M}+\text{H}]$ 491.9 (observed).

Preparation of tert-butyl N-[5-[2-amino-8-bromo-4-[ethoxy(propyl)carbamoyl]-3H-1-benzazepin-6-yl]pent-4-ynyl]carbamate, 8SO2BzL-12c

A mixture of 8SO2BzL-12b (800 mg, 1.63 mmol, 1.0 eq), tert-butyl N-pent-4-ynylcarbamate (328 mg, 1.79 mmol, 1.1 eq), $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (114 mg, 163 μmol , 0.1 eq), cuprous iodide, CuI (61.9 mg, 325 μmol , 0.2 eq) in DMF (16 mL) and Et_3N (6 mL) was degassed and purged with N_2 for 3 times, then stirred at 80 °C for 2 hrs under N_2 atmosphere. The mixture was poured into ice-water (w/w = 1/1) (20 mL) and stirred for 10 min. The aqueous phase was extracted with ethyl acetate (30 mL x 3). The combined organic phase was washed with brine (20 mL x 2), dried with anhydrous Na_2SO_4 , filtered and concentrated in vacuum. The residue was purified by silica gel chromatography (column height: 250 mm, diameter: 100 mm, 100-200 mesh silica gel, Petroleum ether/Ethyl acetate=1/0, 2/1) to afford 8SO2BzL-12c (690 mg, 1.26

mmol, 77.5% yield) as yellow oil. ¹H NMR (MeOD, 400MHz) δ 7.53 (s, 1H), 7.27 (s, 2H), 3.93 (q, J = 7.2 Hz, 2H), 3.74 (t, J = 7.2 Hz, 2H), 3.19 (t, J = 7.2 Hz, 2H), 2.90-2.83 (m, 2H), 2.51 (t, J = 7.2 Hz, 2H), 1.82-1.70 (m, 4H), 1.43 (s, 9H), 1.17 (t, J = 7.2 Hz, 3H), 0.98 (t, J = 7.2 Hz, 3H). LC/MS [M+H] 547.2 (calculated); LC/MS [M+H] 547.2 (observed).

- 5 Preparation of tert-butyl N-[5-[2-amino-8-benzylsulfanyl-4-[ethoxy(propyl) carbamoyl]-3H-1-benzazepin-6-yl]pent-4-ynyl]carbamate, 8SO2BzL-12d

A mixture of 8SO2BzL-12c (350 mg, 639 μmol, 1.0 eq), phenylmethanethiol, BnSH (0.25 g, 2.01 mmol, 236 μL, 3.15 eq), DIEA (165 mg, 1.28 mmol, 223 μL, 2.0 eq), Xantphos (74.0 mg, 128 μmol, 0.2 eq) and Pd₂(dba)₃ (117 mg, 128 μmol, 0.2 eq) in dioxane (10 mL) was degassed and purged with N₂ for 3 times, then stirred at 110 °C for 1 hr under N₂ atmosphere. The residue was poured into ice-water (w/w = 1/1) (10 mL) and stirred for 5 min. The aqueous phase was extracted with ethyl acetate (10 mL x 3). The combined organic phase was washed with brine (10mL x 1), dried with anhydrous Na₂SO₄, filtered and concentrated in vacuum. The residue was purified by silica gel chromatography (column height: 250 mm, diameter: 100 mm, 15 100-200 mesh silica gel, Petroleum ether/Ethyl acetate=1/0, 0/1) to afford 8SO2BzL-12d (300 mg, 508 μmol, 79.4% yield) as yellow solid. ¹H NMR (MeOD, 400MHz) δ 7.56 (s, 1H), 7.37-7.34 (m, 2H), 7.30-7.19 (m, 3H), 7.09 (d, J = 1.6 Hz, 1H), 7.03 (d, J = 1.6 Hz, 1H), 4.20 (s, 2H), 3.93 (q, J = 7.2 Hz, 2H), 3.73 (t, J = 7.2 Hz, 2H), 3.30 (s, 2H), 3.19 (t, J = 6.8 Hz, 2H), 2.49 (t, J = 7.2 Hz, 2H), 1.81-1.72 (m, 4H), 1.43 (s, 9H), 1.16 (t, J = 7.2 Hz, 3H), 0.98 (t, J = 7.2 Hz, 3H). 20 LC/MS [M+H] 591.3 (calculated); LC/MS [M+H] 591.3 (observed).

Preparation of tert-butyl N-[5-[2-amino-8-chlorosulfonyl-4-[ethoxy(propyl) carbamoyl]-3H-1-benzazepin-6-yl]pent-4-ynyl]carbamate, 8SO2BzL-12e

To a solution of 8SO2BzL-12d (300 mg, 508 μmol, 1.0 eq) in MeCN (6 mL) and H₂O (0.6 mL) was added AcOH (305 mg, 5.08 mmol, 290 μL, 10 eq) and NCS (271 mg, 2.03 mmol, 25 4.0 eq), and then stirred at 25 °C for 1 hr. The reaction mixture was poured into ice-water (w/w = 1/1) (10 mL) and stirred for 5 min. The aqueous phase was extracted with ethyl acetate (10 mL x 3), the combined organic phase was dried with anhydrous Na₂SO₄, filtered and concentrated in vacuum to give the crude product 8SO2BzL-12e (250 mg, 441 μmol, 86.8% yield) as yellow oil which was used in the next step without further purification. LC/MS [M+H] 567.2(calculated); 30 LC/MS [M+H] 567.3 (observed).

Preparation of 2-amino-6-(5-aminopent-1-ynyl)-4-[ethoxy(propyl)carbamoyl]-3H-1-benzazepine-8-sulfonic acid, 8SO2BzL-12f

A solution of 8SO2BzL-12e (250 mg, 441 μmol, 1.0 eq) in MeCN (2.5 mL) and H₂O (13 mL) was stirred at 100 °C for 1 hr. The mixture was concentrated in vacuum. The crude 35 product 8SO2BzL-12f (200 mg, 412 μmol, 93.5% yield, HCl) as yellow solid was used in the

next step without further purification. LC/MS [M+H] 449.18(calculated); LC/MS [M+H] 449.1 (observed).

Preparation of 2-amino-6-[5-(tert-butoxycarbonylamino)pent-1-ynyl]-4-
[ethoxy(propyl)carbamoyl]-3H-1-benzazepine-8-sulfonic acid, 8SO2BzL-12g

To a solution of 8SO2BzL-12f (200 mg, 446 μ mol, 1.0 eq) in THF (5 mL) and H₂O (5 mL) was added NaHCO₃ (112 mg, 1.34 mmol, 52 μ L, 3.0 eq) and Boc₂O (146 mg, 669 μ mol, 154 μ L, 1.5 eq), and then stirred at 25 °C for 1 hr. The mixture was concentrated under reduced pressure at 30°C. The residue was purified by prep-HPLC(column: Phenomenex Luna 80*30mm*3 μ m;mobile phase: [water(TFA)-ACN].B%: 10%-40%,8min) to afford 8SO2BzL-12g (100 mg, 182 μ mol, 40.9% yield) as yellow solid. ¹H NMR (MeOD, 400MHz) δ 7.84 (d, J = 1.2 Hz, 1H), 7.71 (d, J = 1.2 Hz, 1H), 7.63 (s, 1H), 3.98 (d, J = 7.2 Hz, 2H), 3.76 (t, J = 6.8 Hz, 2H), 3.38 (s, 2H), 3.18 (t, J = 6.8 Hz, 2H), 2.54 (t, J = 7.2 Hz, 2H), 1.81-1.76 (m, 4H), 1.43 (s, 9H), 1.20 (t, J = 7.2 Hz, 3H), 1.00 (t, J = 7.2 Hz, 3H). LC/MS [M+H]⁺ 549.2 (calculated); LC/MS [M+H]⁺ 549.3 (observed).

15 Preparation of 2-amino-6-[5-(tert-butoxycarbonylamino)pentyl]-4-[ethoxy(propyl)carbamoyl]-3H-1-benzazepine-8-sulfonic acid. 8SO2BzL-12h

A mixture of 8SO2BzL-12g (100 mg, 182 μ mol, 1.0 eq), Pd(OH)₂/C (64.0 mg, 91.1 μ mol, 20% purity, 0.5 eq) in MeOH (10 mL) was degassed and purged with H₂ (367 μ g, 182 μ mol, 1 eq) for 3 times, and then stirred at 25°C for 1 hr under H₂ (30 psi) atmosphere. The mixture was filtered. The residue was purified by prep-HPLC (column: Phenomenex Luna 80*30mm*3 μ m, mobile phase: [water(TFA)-ACN]:B%: 5%-55%, 8min) to afford 8SO2BzL-12b (72 mg, 130 μ mol, 71.5% yield) as white solid. ¹H NMR (MeOD, 400MHz) δ 7.70 (d, J = 1.2 Hz, 1H), 7.65 (d, J = 1.2 Hz, 1H), 7.46 (s, 1H), 3.98 (q, J = 7.2 Hz, 2H), 3.76 (t, J = 6.8 Hz, 2H), 3.36 (s, 2H), 3.02 (t, J = 6.8 Hz, 2H), 2.84 (t, J = 8.0 Hz, 2H), 1.83-1.74 (m, 2H), 1.71-1.59 (m, 2H), 1.52-1.34 (m, 13H), 1.20 (t, J = 7.2 Hz, 3H), 1.01 (t, J = 7.2 Hz, 3H). LC/MS [M+H] 553.26 (calculated); LC/MS [M+H] 553.2 (observed).

Preparation of 2-amino-6-(5-aminopentyl)-4-[ethoxy(propyl)carbamoyl]-3H-1-benzazepine-8-sulfonic acid. 8SO2BzL-12i

To a solution of 8SO2BzL-12h (60 mg, 108 μ mol, 1 eq) in EtOAc (2 mL) was added
30 HCl/EtOAc (4 M, 1 mL, 37 eq), and then stirred at 20 °C for 1 hr. The mixture was
concentrated in vacuum to afford 8SO2BzL-12i (50 mg, 102 μ mol, 94.2% yield, HCl) as white
solid. LC/MS [M+H] 453.2 (calculated); LC/MS [M+H] 453.2 (observed).


Preparation of 8SO2BzL-12

To a solution of 3-[2-[2-[2-[2-[2-[2-[2-[2-[[2-(2,5-dioxopyrrol-1-yl)acetyl]

35 amino]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]propanoic

acid (68.2 mg, 102 μ mol, 1.0 eq) in DMF (1 mL) was added DIEA (66.1 mg, 511 μ mol, 89 μ L, 5.0 eq), 8SO2BzL-12i (50 mg, 102 μ mol, 1 eq, HCl) and HATU (38.9 mg, 102 μ mol, 1.0 eq). And it was stirred at 25°C for 0.5 hr. The mixture was filtered and purified by prep-HPLC (column: Phenomenex Luna 80*30mm*3 μ m; mobile phase: [water(TFA)-ACN]; B%: 5%-35%, 8min) to afford 8SO2BzL-12 (14 mg, 12.7 μ mol, 12.43% yield) as yellow oil. ¹H NMR (MeOD, 400MHz) δ 7.72 (s, 1H), 7.68 (s, 1H), 7.48 (s, 1H), 6.91 (s, 2H), 4.19 (s, 2H), 4.01 (q, J = 7.2 Hz, 2H), 3.80-3.55 (m, 42H), 3.43-3.35 (m, 4H), 3.19 (t, J = 6.8 Hz, 2H), 2.88 (t, J = 8.0 Hz, 2H), 2.42 (t, J = 6.4 Hz, 2H), 1.80 (q, J = 7.2 Hz, 2H), 1.74-1.64 (m, 2H), 1.60-1.51 (m, 2H), 1.45-1.43 (m, 2H), 1.23 (t, J = 7.2 Hz, 3H), 1.02 (t, J = 7.2 Hz, 3H). LC/MS [M+H] 1101.5 (calculated); LC/MS [M+H] 1101.9 (observed).

Example L-13 Synthesis of 1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl 2-((2-amino-8-(N-methylsulfamoyl)-N-propyl-3H-benzo[b]azepine-4-carboxamido)oxy)ethyl)carbamate, 8SO2BzL-13



Preparation of tert-butyl (2-((2-amino-8-(N-methylsulfamoyl)-N-propyl-3H-benzo[b]azepine-4-carboxamido)oxy)ethyl)carbamate, 8SO2BzL-13b

To a solution of tert-butyl (2-((2-amino-8-(chlorosulfonyl)-N-propyl-3H-benzo[b]azepine-4-carboxamido)oxy)ethyl)carbamate, 8SO2BzL-13a (0.14 M, 2.00 mL, 0.28 mmol) in acetonitrile/water (9:1) at 0 deg was added a methylamine solution (2 M in THF, 0.70 mL, 1.40 mmol). After 10 minutes the solvent was removed and the crude product was purified by reverse phase HPLC using a gradient of 10-90% acetonitrile/water to give 8SO2BzL-13b (0.09 g, 68%) as a yellow film after evaporation of solvent.

Preparation of 2-amino-N-(2-aminoethoxy)-8-(N-methylsulfamoyl)-N-propyl-3H-benzo[b]azepine-4-carboxamide, 8SO2BzL-13c


A solution of 8SO2BzL-13b (0.09 g, 0.19 mmol) in acetonitrile (2 mL) and 6 N HCl (2.00 mL, 12.00 mmol) was stirred at room temperature for 45 minutes. The solvent was removed by vacuum to give 8SO2BzL-13c as the HCl salt (0.08 g, 99%).

Preparation of 8SO2BzL-13

To a solution of 8SO2BzL-13c HCl (0.03 g, 0.06 mmol) in DMF (4 mL) at room temperature added triethylamine (0.04 mL, 0.26 mmol). To this mixture was added a solution of 1-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-2-oxo-6,9,12,15,18,21,24,27,30,33-decaoxa-3-azapentatriacontan-35-yl (4-nitrophenyl) carbonate, PNPC-PEG10-mal (0.05 g, 0.06 mmol) in DMF (2 mL). After 20 min., acetic acid (6 uL) was added and reaction was concentrated under vacuum and purified by reverse phase HPLC using a gradient of 10-90% ACN/water (+0.1% TFA) over 10 min. to give 8SO2BzL-13 (0.04 g, 52%) after concentration of pure fractions. LC/MS [M+H] 1060.47 (calculated); LC/MS [M+H] 1060.89 (observed).

Example L-15 Synthesis of 2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-(2,5-dioxopyrrol-1-

yl)acetyl]amino]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy] N-[2-[2-amino-8-(thiazol-2-ylsulfamoyl)-3H-1-benzazepine-4-carbonyl]-propyl-amino]oxyethyl]carbamate, 8SO2BzL-15



Preparation of ethyl 8-bromo-2-(tritylamino)-3H-1-benzazepine-4-carboxylate,

5 8SO2BzL-15b

A mixture of ethyl 2-amino-8-bromo-3H-1-benzazepine-4-carboxylate, 8SO2BzL-15a (5 g, 16.1 mmol, 1 *eq*), TrtCl (6.76 g, 24.2 mmol, 1.5 *eq*), TEA (4.91 g, 48.5 mmol, 6.75 mL, 3 *eq*)

and DMAP (395 mg, 3.23 mmol, 0.2 *eq*) in DCM (50 mL) was degassed and purged with N₂ for 3 times, and then stirred at 40°C for 16 h under N₂ atmosphere. The reaction mixture was quenched by addition H₂O (50 mL) and extracted with EtOAc (100 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether:Ethyl acetate=100:1 to 0:1) to give 8SO2BzL-15b (7.8 g, 14.1 mmol, 87.4% yield) as a yellow oil. LC/MS [M+H] 551.1 (calculated); LC/MS [M+H] 551.1 (observed).

Preparation of ethyl 8-benzylsulfanyl-2-(tritylamino)-3H-1-benzazepine-4-carboxylate, 8SO2BzL-15c

A mixture of 8SO2BzL-15b (3 g, 5.44 mmol, 1 *eq*), DIEA (1.41 g, 10.8 mmol, 1.90 mL, 2 *eq*), (1E,4E)-1,5-diphenylpenta-1,4-dien-3-one;palladium, Pd₂(dba)₃, CAS Reg. No. 51364-51-3 (996 mg, 1.09 mmol, 0.2 *eq*) and (5-diphenylphosphanyl-9,9-dimethyl-xanthen-4-yl)-diphenyl-phosphane, Xphos (629 mg, 1.09 mmol, 0.2 *eq*) in dioxane (30 mL) was degassed and purged with N₂ for 3 times, then phenylmethanethiol, BnSH (1.35 g, 10.8 mmol, 1.27 mL, 2 *eq*) was added, the mixture was stirred at 110°C for 1 h under N₂ atmosphere. The reaction mixture was quenched by addition H₂O (50 mL) and extracted with EtOAc (100 mL x 2). The combined organic layers were washed with brine (100 mL) dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether:Ethyl acetate=100:1 to 0:18) to give 8SO2BzL-15c (2.3 g, 3.87 mmol, 71.1% yield) as a yellow oil.

Preparation of ethyl 8-chlorosulfonyl-2-(tritylamino)-3H-1-benzazepine-4-carboxylate, 8SO2BzL-15d

A mixture of 8SO2BzL-15c (3 g, 5.04 mmol, 1 *eq*), NCS (2.69 g, 20.2 mmol, 4 *eq*), AcOH (3.03 g, 50.4 mmol, 2.88 mL, 10 *eq*) in MeCN (30 mL) and H₂O (3 mL) was stirred at 25°C for 1 h. The reaction mixture was quenched by addition H₂O (50 mL), and extracted with EtOAc (50 mL). The combined organic layers were wash with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether:Ethyl acetate=99:1 to 0:1) to give 8SO2BzL-15d (2 g, 3.50 mmol, 69.4% yield) as a yellow solid. LC/MS [M+H] 571.1 (calculated); LC/MS [M+H] 571.2 (observed).

Preparation of ethyl 8-(thiazol-2-ylsulfamoyl)-2-(tritylamino)-3H-1-benzazepine-4-carboxylate, 8SO2BzL-15e

A mixture of 8SO2BzL-15d (1.5 g, 2.63 mmol, 1 *eq*) and 1-methylimidazole (258 mg, 3.15 mmol, 251 μ L, 1.2 *eq*) in MeCN (30 mL) was degassed and purged with N₂ for 3 times and then stirred at 25°C for 2 h under N₂ atmosphere. Then 4,5-dihydrothiazol-2-amine (1.07 g,

10.5 mmol, 4 *eq*) was added and the result mixture was stirred at 25°C for another 16 h under N₂ atmosphere. The reaction mixture was quenched by addition H₂O (50 mL) and extracted with EtOAc (50 mL). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was
5 purified by column chromatography (SiO₂, Petroleum ether:Ethyl acetate=100:1 to 0:1) to give 8SO2BzL-15e (0.5 g, 787 μ mol, 29.9% yield) as a yellow solid. ¹H NMR (MeOD, 400 MHz) δ 7.71 (s, 1H), 7.23 (m, 20H), 6.75 (d, J = 4.8 Hz, 1H), 4.36 (q, J = 7.2 Hz, 2H), 2.98 (s, 2H), 1.38 (t, J = 7.2 Hz, 3H). LC/MS [M+H]⁺ 635.17 (calculated); LC/MS [M+H]⁺ 635.1 (observed).

Preparation of 8-(N-(thiazol-2-yl)sulfamoyl)-2-(tritylamino)-3H-benzo[b]azepine-4-
10 carboxylic acid, 8SO2BzL-15f

A mixture of 8SO2BzL-15e (0.5 g, 787.69 μ mol, 1 *eq*), LiOH.H₂O (264 mg, 6.30 mmol, 8 *eq*) in H₂O (4 mL) and THF (4 mL) was stirred at 25 °C for 3 hr. The reaction solution was quenched by added 2M HCl to adjust pH = ~6, then filtered to give 8SO2BzL-15f (0.45 g, 699.66 μ mol, 88.82% yield, HCl) as a white solid. LC/MS [M+H]⁺ 607.1 (calculated); LC/MS
15 [M+H]⁺ 607.2 (observed).

Preparation of tert-butyl N-[2-[propyl-[8-(thiazol-2-yl)sulfamoyl)-2-(tritylamino)-3H-1-benzazepine-4-carbonyl]amino]oxyethyl]carbamate, 8SO2BzL-15g

A mixture of 8SO2BzL-15f (0.42 g, 692 μ mol, 1 *eq*), *tert*-butyl N-[2-(propylamino)oxyethyl]carbamate (181 mg, 830 μ mol, 1.2 *eq*), methanesulfonic acid (133 mg, 1.38 mmol, 98.5 μ L, 2 *eq*), EDCI (663 mg, 3.46 mmol, 5 *eq*) in DMA (5 mL) and DCM (5 mL)
20 was degassed and purged with N₂ for 3 times, and then stirred at 25°C for 2 hr under N₂ atmosphere. The reaction mixture was quenched by addition Na₂HCO₃ (3 mL) until pH about 7, and extracted with EtOAc (5 mL*3). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄ filtered and concentrated under reduced pressure to give a residue. The
25 residue was purified by column chromatography (SiO₂, Petroleum ether:Ethyl acetate=99:1 to 20:80) to give 8SO2BzL-15g (0.5 g, 619 μ mol, 89.5% yield) as a yellow solid. LC/MS [M+H]⁺ 807.3 (calculated); LC/MS [M+H]⁺ 807.3 (observed).

Preparation of 2-amino-N-(2-aminoethoxy)-N-propyl-8-(thiazol-2-yl)sulfamoyl)-3H-1-benzazepine-4-carboxamide, 8SO2BzL-15h

30 A mixture of 8SO2BzL-15g (0.5 g, 619 μ mol, 1 *eq*), TFA (5.65 g, 49.5 mmol, 3.67 mL, 80 *eq*) in DCM (10 mL) was stirred at 25 °C for 16 h. The reaction mixture was quenched by addition H₂O (5 mL), and extracted with MTBE (10 mL) (5 mL x 2) to remove the excess TFA. The combined water layers was concentrated under reduced pressure to give 8SO2BzL-15h (0.25 g, 432 μ mol, 69.7% yield, TFA) as a white solid. LC/MS [M+H]⁺ 465.1 (calculated);
35 LC/MS [M+H]⁺ 465.1 (observed).

Preparation of 8SO2BzL-15

A mixture of 8SO2BzL-15h (0.2 g, 288 μ mol, 1 *eq*, 2TFA), 2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-(2,5-dioxopyrrol-1-yl)acetyl]amino]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy] (4-nitrophenyl)carbonate (232 mg, 288 μ mol, 1 *eq*), DIEA (111 mg, 866 μ mol, 150 μ L, 3 *eq*) in DMF (0.5 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 0°C for 1 h under N₂ atmosphere. The reaction solution was quenched with TFA until pH = ~6. The residue was purified by prep-HPLC (column: Phenomenex Luna 80 x 30mm x 3 μ m; mobile phase: [water (TFA)-ACN]; B%: 15%-40%, 8min) to give 8SO2BzL-15 (15.0 mg, 13.2 μ mol, 4.6% yield) as a white solid. ¹H NMR (MeOD-d₄, 400 MHz) δ 7.97 (s, 1H), 7.88 (d, J = 2.0 Hz, 1H), 7.86 (d, J = 2.0 Hz, 1H), 7.39 (s, 1H), 7.18 (d, J = 4.8 Hz, 1H), 6.90 (s, 2H), 6.81 (d, J = 4.8 Hz, 1H), 4.17 (s, 2H), 3.74 (m, 2H), 3.67 (m, 2H), 3.59 (m, 42H), 3.38 (m, 6H), 1.80-1.72 (m, 2H) 1.00 (t, J = 7.2 Hz, 3H). LC/MS [M+H]⁺ 1129.4 (calculated); LC/MS [M+H]⁺ 1129.5 (observed).

Example 201 Preparation of Immunoconjugates (IC)

To prepare a lysine-conjugated Immunoconjugate, an antibody is buffer exchanged into a conjugation buffer containing 100 mM boric acid, 50 mM sodium chloride, 1 mM ethylenediaminetetraacetic acid at pH 8.3, using G-25 SEPHADEXTM desalting columns (Sigma-Aldrich, St. Louis, MO) or ZebaTM Spin Desalting Columns (Thermo Fisher Scientific). The eluates are then each adjusted to a concentration of about 1-10 mg/ml using the buffer and then sterile filtered. The antibody is pre-warmed to 20-30 °C and rapidly mixed with 2-20 (e.g., 7-10) molar equivalents of a tetrafluorophenyl (TFP) or sulfonic tetrafluorophenyl (sulfoTFP) ester, 8-sulfonyl-2-aminobenzazepine-linker (8SO2Bz-L) compound of Formula II dissolved in dimethylsulfoxide (DMSO) or dimethylacetamide (DMA) to a concentration of 5 to 20 mM. The reaction is allowed to proceed for about 16 hours at 30 °C and the immunoconjugate (IC) is separated from reactants by running over two successive G-25 desalting columns or ZebaTM Spin Desalting Columns equilibrated in phosphate buffered saline (PBS) at pH 7.2 to provide the Immunoconjugate (IC) of Tables 3a and 3b. Adjuvant-antibody ratio (DAR) is determined by liquid chromatography mass spectrometry analysis using a C4 reverse phase column on an ACQUITYTM UPLC H-class (Waters Corporation, Milford, MA) connected to a XEVOTM G2-XS TOF mass spectrometer (Waters Corporation).

To prepare a cysteine-conjugated Immunoconjugate, an antibody is buffer exchanged into a conjugation buffer containing PBS, pH 7.2 with 2 mM EDTA using ZebaTM Spin Desalting Columns (Thermo Fisher Scientific). The interchain disulfides are reduced using 2-4 molar excess of Tris (2-carboxyethyl) phosphine (TCEP) or dithiothreitol (DTT) at 37 °C for 30

- min -- 2 hours. Excess TCEP or DTT was removed using a Zeba™ Spin Desalting column pre-equilibrated with the conjugation buffer. The concentration of the buffer-exchanged antibody was adjusted to approximately 5 to 20 mg/ml using the conjugation buffer and sterile-filtered. The maleimide-8SO2Bz-L compound is either dissolved in dimethylsulfoxide (DMSO) or
- 5 dimethylacetamide (DMA) to a concentration of 5 to 20 mM. For conjugation, the antibody is mixed with 10 to 20 molar equivalents of maleimide-8SO2Bz-L. In some instances, additional DMA or DMSO up to 20% (v/v), was added to improve the solubility of the maleimide-8SO2Bz-L in the conjugation buffer. The reaction is allowed to proceed for approximately 30 min to 4 hours at 20 °C. The resulting conjugate is purified away from the unreacted maleimide-
- 10 8SO2Bz-L using two successive Zeba™ Spin Desalting Columns. The columns are pre-equilibrated with phosphate-buffered saline (PBS), pH 7.2. Adjuvant to antibody ratio (DAR) is estimated by liquid chromatography mass spectrometry analysis using a C4 reverse phase column on an ACQUITY™ UPLC H-class (Waters Corporation, Milford, MA) connected to a XEVO™ G2-XS TOF mass spectrometer (Waters Corporation).
- 15 For conjugation, the antibody may be dissolved in an aqueous buffer system known in the art that will not adversely impact the stability or antigen-binding specificity of the antibody. Phosphate buffered saline may be used. The 8SO2Bz-L compound is dissolved in a solvent system comprising at least one polar aprotic solvent as described elsewhere herein. In some such aspects, 8SO2Bz-L is dissolved to a concentration of about 5 mM, about 10 mM, about 20 mM,
- 20 about 30 mM, about 40 mM or about 50 mM, and ranges thereof such as from about 5 mM to about 50mM or from about 10 mM to about 30 mM in pH 8 Tris buffer (e.g., 50 mM Tris). In some aspects, the 8-sulfonyl-2-aminobenzazepine-linker intermediate is dissolved in DMSO (dimethylsulfoxide), DMA (dimethylacetamide), acetonitrile, or another suitable dipolar aprotic solvent.
- 25 Alternatively in the conjugation reaction, an equivalent excess of 8SO2Bz-L solution may be diluted and combined with antibody solution. The 8SO2Bz-L solution may suitably be diluted with at least one polar aprotic solvent and at least one polar protic solvent, examples of which include water, methanol, ethanol, n-propanol, and acetic acid. The molar equivalents of 8SO2Bz-L intermediate to antibody may be about 1.5:1, about 3:1, about 5:1, about 10:1, about
- 30 15:1, or about 20:1, and ranges thereof, such as from about 1.5:1 to about 20:1 from about 1.5:1 to about 15:1, from about 1.5:1 to about 10:1, from about 3:1 to about 15:1, from about 3:1 to about 10:1, from about 5:1 to about 15:1 or from about 5:1 to about 10:1. The reaction may suitably be monitored for completion by methods known in the art, such as LC-MS. The conjugation reaction is typically complete in a range from about 1 hour to about 16 hours. After
- 35 the reaction is complete, a reagent may be added to the reaction mixture to quench the reaction.

If antibody thiol groups are reacting with a thiol-reactive group such as maleimide of the 8SO2Bz-L linker intermediate, unreacted antibody thiol groups may be reacted with a capping reagent. An example of a suitable capping reagent is ethylmaleimide.

Following conjugation, the immunoconjugates may be purified and separated from unconjugated reactants and/or conjugate aggregates by purification methods known in the art such as, for example and not limited to, size exclusion chromatography, hydrophobic interaction chromatography, ion exchange chromatography, chromatofocusing, ultrafiltration, centrifugal ultrafiltration, tangential flow filtration, and combinations thereof. For instance, purification may be preceded by diluting the immunoconjugate, such in 20 mM sodium succinate, pH 5. The diluted solution is applied to a cation exchange column followed by washing with, e.g., at least 10 column volumes of 20 mM sodium succinate, pH 5. The conjugate may be suitably eluted with a buffer such as PBS.

Example 202 HEK Reporter Assay

HEK293 reporter cells expressing human TLR7 or human TLR8 were purchased from Invivogen and vendor protocols were followed for cellular propagation and experimentation. Briefly, cells were grown to 80-85% confluence at 5% CO₂ in DMEM supplemented with 10% FBS, Zeocin, and Blasticidin. Cells were then seeded in 96-well flat plates at 4x10⁴ cells/well with substrate containing HEK detection medium and immunostimulatory molecules. Activity was measured using a plate reader at 620-655 nm wavelength.

20 Example 203 Assessment of Immunoconjugate Activity *In Vitro*

This example shows that Immunoconjugates of the invention are effective at eliciting immune activation, and therefore are useful for the treatment of cancer.

a) Isolation of Human Antigen Presenting Cells: Human myeloid antigen presenting cells (APCs) were negatively selected from human peripheral blood obtained from healthy blood donors (Stanford Blood Center, Palo Alto, California) by density gradient centrifugation using a ROSETTESEP™ Human Monocyte Enrichment Cocktail (Stem Cell Technologies, Vancouver, Canada) containing monoclonal antibodies against CD14, CD16, CD40, CD86, CD123, and HLA-DR. Immature APCs were subsequently purified to >90% purity via negative selection using an EASYSEP™ Human Monocyte Enrichment Kit (Stem Cell Technologies) without CD16 depletion containing monoclonal antibodies against CD14, CD16, CD40, CD86, CD123, and HLA-DR.

b) Myeloid APC Activation Assay: 2 x 10⁵ APCs are incubated in 96-well plates (Corning, Corning, NY) containing iscove's modified dulbecco's medium, IMDM (Lonza)

supplemented with 10% FBS, 100 U/mL penicillin, 100 µg/mL (micrograms per milliliter) streptomycin, 2 mM L-glutamine, sodium pyruvate, non-essential amino acids, and where indicated, various concentrations of unconjugated (naked) antibodies and immunoconjugates (IC) of the invention (as prepared according to the Example above). Cell-free supernatants are
5 analyzed after 18 hours via ELISA to measure TNFα secretion as a readout of a proinflammatory response.

c) PBMC Activation Assay: Human peripheral blood mononuclear cells were isolated from human peripheral blood obtained from healthy blood donors (Stanford Blood Center, Palo Alto, California) by density gradient centrifugation. PBMCs were incubated in 96-
10 well plates (Corning, Corning, NY) in a co-culture with CEA-expressing tumor cells (e.g. MKN-45, HPAF-II) at a 10:1 effector to target cell ratio. Cells were stimulated with various concentrations of unconjugated (naked) antibodies and immunoconjugates of the invention (as prepared according to the Example above). Cell-free supernatants were analyzed by cytokine bead array using a LegendPlex™ kit according to manufacturer's guidelines (BioLegend®, San
15 Diego, CA).

d) Isolation of Human Conventional Dendritic Cells: Human conventional dendritic cells (cDCs) were negatively selected from human peripheral blood obtained from healthy blood donors (Stanford Blood Center, Palo Alto, California) by density gradient centrifugation. Briefly, cells are first enriched by using a ROSETTESEP™ Human CD3 Depletion Cocktail
20 (Stem Cell Technologies, Vancouver, Canada) to remove T cells from the cell preparation. cDCs are then further enriched via negative selection using an EASYSEP™ Human Myeloid DC Enrichment Kit (Stem Cell Technologies).

e) cDC Activation Assay: 8×10^4 APCs were co-cultured with tumor cells expressing the ISAC target antigen at a 10:1 effector (cDC) to target (tumor cell) ratio. Cells
25 were incubated in 96-well plates (Corning, Corning, NY) containing RPMI-1640 medium supplemented with 10% FBS, and where indicated, various concentrations of the indicated immunoconjugate of the invention (as prepared according to the example above). Following overnight incubation of about 18 hours, cell-free supernatants were collected and analyzed for cytokine secretion (including TNFα) using a BioLegend LEGENDPLEX cytokine bead array.

30 Activation of myeloid cell types can be measured using various screen assays in addition to the assay described in which different myeloid populations are utilized. These may include the following: monocytes isolated from healthy donor blood, M-CSF differentiated Macrophages, GM-CSF differentiated Macrophages, GM-CSF+IL-4 monocyte-derived


Dendritic Cells, conventional Dendritic Cells (cDCs) isolated from healthy donor blood, and myeloid cells polarized to an immunosuppressive state (also referred to as myeloid derived suppressor cells or MDSCs). Examples of MDSC polarized cells include monocytes differentiated toward immunosuppressive state such as M2a MΦ (IL4/IL13), M2c MΦ (IL10/TGFβ), GM-CSF/IL6 MDSCs and tumor-educated monocytes (TEM). TEM differentiation can be performed using tumor-conditioned media (e.g. 786.O, MDA-MB-231, HCC1954). Primary tumor-associated myeloid cells may also include primary cells present in dissociated tumor cell suspensions (Discovery Life Sciences).

Assessment of activation of the described populations of myeloid cells may be performed as a mono-culture or as a co-culture with cells expressing the antigen of interest which the immunoconjugate (IC) may bind to via the CDR region of the antibody. Following incubation for 18-48 hours, activation may be assessed by upregulation of cell surface co-stimulatory molecules using flow cytometry or by measurement of secreted proinflammatory cytokines. For cytokine measurement, cell-free supernatant is harvested and analyzed by cytokine bead array (e.g. LegendPlex from Biolegend) using flow cytometry.

All references, including publications, patent applications, and patents, cited herein are hereby incorporated by reference to the same extent as if each reference were individually and specifically indicated to be incorporated by reference and were set forth in its entirety herein.

CLAIMS:

1. An immunoconjugate comprising an antibody covalently attached to one or more 8-sulfonyl-2-aminobenzazepine moieties by a linker, and having Formula I:



or a pharmaceutically acceptable salt thereof,


wherein:

Ab is the antibody wherein the antibody binds to a target selected from PD-L1, HER2, CEA; and TROP2,

p is an integer from 1 to 8;

L is the linker;

D is the 8-sulfonyl-2-aminobenzazepine moiety having the formula:



R^1 , R^2 , R^3 , and R^4 are independently selected from the group consisting of H, C₁-C₁₂ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₁₂ carbocyclyl, C₆-C₂₀ aryl, C₂-C₉ heterocyclyl, and C₁-C₂₀ heteroaryl, each of which are independently and optionally substituted with one or more groups selected from:

- (C₁-C₁₂ alkylidyl)-N(R⁵)-*;
- (C₁-C₁₂ alkylidyl)-N(R⁵)₂;
- (C₁-C₁₂ alkylidyl)-OR⁵;
- (C₃-C₁₂ carbocyclyl);
- (C₃-C₁₂ carbocyclyl)-*;
- (C₃-C₁₂ carbocyclyl)-(C₁-C₁₂ alkylidyl)-NR⁵-*;
- (C₃-C₁₂ carbocyclyl)-(C₁-C₁₂ alkylidyl)-N(R⁵)₂;
- (C₃-C₁₂ carbocyclyl)-NR⁵-C(=NR⁵)NR⁵-*;
- (C₆-C₂₀ aryl);
- (C₆-C₂₀ arylidyl)-*;
- (C₆-C₂₀ arylidyl)-N(R⁵)-*;
- (C₆-C₂₀ arylidyl)-(C₁-C₁₂ alkylidyl)-N(R⁵)-*;

--(C₆-C₂₀ arylldiyl)--(C₁-C₁₂ alkylldiyl)--(C₂-C₂₀ heterocyclldiyl)--*;
 --(C₆-C₂₀ arylldiyl)--(C₁-C₁₂ alkylldiyl)--N(R⁵)₂;
 --(C₆-C₂₀ arylldiyl)--(C₁-C₁₂ alkylldiyl)--NR⁵--C(=NR^{5a})N(R⁵)--*;
 --(C₂-C₂₀ heterocyclldiyl);
 --(C₂-C₂₀ heterocyclldiyl)--*;
 --(C₂-C₉ heterocyclldiyl)--(C₁-C₁₂ alkylldiyl)--NR⁵--*;
 --(C₂-C₉ heterocyclldiyl)--(C₁-C₁₂ alkylldiyl)--N(R⁵)₂;
 --(C₂-C₉ heterocyclldiyl)--C(=O)--(C₁-C₁₂ alkylldiyl)--N(R⁵)--*;
 --(C₂-C₉ heterocyclldiyl)--NR⁵--C(=NR^{5a})NR⁵--*;
 --(C₂-C₉ heterocyclldiyl)--NR⁵--(C₆-C₂₀ arylldiyl)--(C₁-C₁₂ alkylldiyl)--N(R⁵)--*;
 --(C₂-C₉ heterocyclldiyl)--(C₆-C₂₀ arylldiyl)--*;
 --(C₁-C₂₀ heteroaryl);
 --(C₁-C₂₀ heteroaryl)--*;
 --(C₁-C₂₀ heteroaryl)--(C₁-C₁₂ alkylldiyl)--N(R⁵)--*;
 --(C₁-C₂₀ heteroaryl)--(C₁-C₁₂ alkylldiyl)--N(R⁵)₂;
 --(C₁-C₂₀ heteroaryl)--NR⁵--C(=NR^{5a})N(R⁵)--*;
 --(C₁-C₂₀ heteroaryl)--N(R⁵)C(=O)--(C₁-C₁₂ alkylldiyl)--N(R⁵)--*;
 --C(=O)--*;
 --C(=O)--(C₁-C₁₂ alkylldiyl)--N(R⁵)--*;
 --C(=O)--(C₂-C₂₀ heterocyclldiyl)--*;
 --C(=O)N(R⁵)₂;
 --C(=O)N(R⁵)--*;
 --C(=O)N(R⁵)--(C₁-C₁₂ alkylldiyl)--N(R⁵)C(=O)R⁵;
 --C(=O)N(R⁵)--(C₁-C₁₂ alkylldiyl)--N(R⁵)C(=O)N(R⁵)₂;
 --C(=O)NR⁵--(C₁-C₁₂ alkylldiyl)--N(R⁵)CO₂R⁵;
 --C(=O)NR⁵--(C₁-C₁₂ alkylldiyl)--N(R⁵)C(=NR^{5a})N(R⁵)₂;
 --C(=O)NR⁵--(C₁-C₁₂ alkylldiyl)--NR⁵C(=NR^{5a})R⁵;
 --C(=O)NR⁵--(C₁-C₈ alkylldiyl)--NR⁵(C₂-C₅ heteroaryl);
 --C(=O)NR⁵--(C₁-C₂₀ heteroarylldiyl)--N(R⁵)--*;
 --C(=O)NR⁵--(C₁-C₂₀ heteroarylldiyl)--*;
 --C(=O)NR⁵--(C₁-C₂₀ heteroarylldiyl)--(C₁-C₁₂ alkylldiyl)--N(R⁵)₂;
 --C(=O)NR⁵--(C₁-C₂₀ heteroarylldiyl)--(C₂-C₂₀ heterocyclldiyl)--C(=O)NR⁵--(C₁-C₁₂
 alkylldiyl)--NR⁵--*;

$-\text{N}(\text{R}^5)_2$;
 $-\text{N}(\text{R}^5)-*$;
 $-\text{N}(\text{R}^5)\text{C}(=\text{O})\text{R}^5$;
 $-\text{N}(\text{R}^5)\text{C}(=\text{O})-*$;
 $-\text{N}(\text{R}^5)\text{C}(=\text{O})\text{N}(\text{R}^5)_2$;
 $-\text{N}(\text{R}^5)\text{C}(=\text{O})\text{N}(\text{R}^5)-*$;
 $-\text{N}(\text{R}^5)\text{CO}_2\text{R}^5$;
 $-\text{NR}^5\text{C}(=\text{NR}^{5a})\text{N}(\text{R}^5)_2$;
 $-\text{NR}^5\text{C}(=\text{NR}^{5a})\text{N}(\text{R}^5)-*$;
 $-\text{NR}^5\text{C}(=\text{NR}^{5a})\text{R}^5$;
 $-\text{N}(\text{R}^5)\text{C}(=\text{O})-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{N}(\text{R}^5)-*$;
 $-\text{N}(\text{R}^5)-(\text{C}_2\text{-C}_5\text{ heteroaryl})$;
 $-\text{N}(\text{R}^5)-\text{S}(=\text{O})_2-(\text{C}_1\text{-C}_{12}\text{ alkyl})$;
 $-\text{O}-(\text{C}_1\text{-C}_{12}\text{ alkyl})$;
 $-\text{O}-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{N}(\text{R}^5)_2$;
 $-\text{O}-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{N}(\text{R}^5)-*$;
 $-\text{O}-\text{C}(=\text{O})\text{N}(\text{R}^5)_2$;
 $-\text{O}-\text{C}(=\text{O})\text{N}(\text{R}^5)-*$;
 $-\text{O}-(\text{R}^5)-*$;
 $-\text{OR}^5$;
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20}\text{ heterocyclidyl})-*$;
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20}\text{ heterocyclidyl})-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{N}(\text{R}^5)_2$;
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20}\text{ heterocyclidyl})-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{NR}^5-*$; and
 $-\text{S}(=\text{O})_2-(\text{C}_2\text{-C}_{20}\text{ heterocyclidyl})-(\text{C}_1\text{-C}_{12}\text{ alkylidyl})-\text{OH}$;
 or R^2 and R^3 together form a 5- or 6-membered heterocyclyl ring;

X^1 , X^2 , X^3 and X^4 are independently selected from the group consisting of a bond, $\text{C}(=\text{O})$, $\text{C}(=\text{O})\text{N}(\text{R}^5)$, O , $\text{N}(\text{R}^5)$, S , $\text{S}(\text{O})_2$, and $\text{S}(\text{O})_2\text{N}(\text{R}^5)$;

R^5 is independently selected from the group consisting of H, $\text{C}_6\text{-C}_{20}$ aryl, $\text{C}_3\text{-C}_{12}$ carbocyclyl, $\text{C}_6\text{-C}_{20}$ arylidyl, $\text{C}_1\text{-C}_{12}$ alkyl, and $\text{C}_1\text{-C}_{12}$ alkylidyl, or two R^5 groups together form a 5- or 6-membered heterocyclyl ring;

R^{5a} is selected from the group consisting of $\text{C}_6\text{-C}_{20}$ aryl and $\text{C}_1\text{-C}_{20}$ heteroaryl;

where the asterisk * indicates the attachment site of L, and where one of R^1 , R^2 , R^3 and R^4 is attached to L;


L is the linker selected from the group consisting of:

- C(=O)—PEG—;
- C(=O)—PEG—C(=O)N(R⁶)—(C₁-C₁₂ alkylidyl)—C(=O)—Gluc—;
- C(=O)—PEG—O—;
- C(=O)—PEG—O—C(=O)—;
- C(=O)—PEG—C(=O)—;
- C(=O)—PEG—C(=O)—PEP—;
- C(=O)—PEG—N(R⁶)—;
- C(=O)—PEG—N(R⁶)—C(=O)—;
- C(=O)—PEG—N(R⁶)—PEG—C(=O)—PEP—;
- C(=O)—PEG—N⁺(R⁶)₂—PEG—C(=O)—PEP—;
- C(=O)—PEG—C(=O)—PEP—N(R⁶)—(C₁-C₁₂ alkylidyl)—;
- C(=O)—PEG—C(=O)—PEP—N(R⁶)—(C₁-C₁₂ alkylidyl)N(R⁶)C(=O)—(C₂-C₅ monoheterocyclydiyl)—;
- C(=O)—PEG—SS—(C₁-C₁₂ alkylidyl)—OC(=O)—;
- C(=O)—PEG—SS—(C₁-C₁₂ alkylidyl)—C(=O)—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—C(=O)N(R⁶)—(C₁-C₁₂ alkylidyl)—C(=O)—Gluc—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—O—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—O—C(=O)—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—C(=O)—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—N(R⁵)—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—N(PEG—CO₂H)—PEG—N(R⁵)—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—C(=O)N(PEG—CO₂H)—PEG—N(R⁵)—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—N(R⁵)—C(=O)—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—N(PEG—CO₂H)—PEG—C(=O)—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—C(=O)N(PEG—CO₂H)—PEG—C(=O)—;
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—C(=O)—PEP—; and
- succinimidyl—(CH₂)_m—C(=O)N(R⁶)—PEG—SS—(C₁-C₁₂ alkylidyl)—OC(=O)—;


R⁶ is independently H or C₁-C₆ alkyl;

PEG has the formula: —(CH₂CH₂O)_n—(CH₂)_m—; m is an integer from 1 to 5, and n is an integer from 2 to 50;

Gluc has the formula:




PEP has the formula:



where AA is independently selected from a natural or unnatural amino acid side chain, or one or more of AA, and an adjacent nitrogen atom form a 5-membered ring proline amino acid, and the wavy line indicates a point of attachment;

Cyc is selected from C₆-C₂₀ arylidiyl and C₁-C₂₀ heteroaryldiyl, optionally substituted with one or more groups selected from F, Cl, NO₂, -OH, -OCH₃, and a glucuronic acid having the structure:



R⁷ is selected from the group consisting of -CH(R⁸)O-, -CH₂-, -CH₂N(R⁸)-, and -CH(R⁸)O-C(=O)-, where R⁸ is selected from H, C₁-C₆ alkyl, C(=O)-C₁-C₆ alkyl, and -C(=O)N(R⁹)₂, where R⁹ is independently selected from the group consisting of H, C₁-C₁₂ alkyl, and -(CH₂CH₂O)_m-(CH₂)_m-OH, where m is an integer from 1 to 5, and n is an integer from 2 to 50, or two R⁹ groups together form a 5- or 6-membered heterocycl ring;

y is an integer from 2 to 12;

z is 0 or 1; and

alkyl, alkylidiyl, alkenyl, alkenyldiyl, alkynyl, alkynyldiyl, aryl, aryldiyl, carbocycl, carbocyclidiyl, heterocycl, heterocyclidiyl, heteroaryl, and heteroaryldiyl are independently and optionally substituted with one or more groups independently selected from F, Cl, Br, I, -CN, -CH₃, -CH₂CH₃, -CH=CH₂, -C≡CH, -C≡CCH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH(CH₃)₂, -CH₂OH, -CH₂OCH₃, -CH₂CH₂OH, -C(CH₃)₂OH, -CH(OH)CH(CH₃)₂, -

$C(CH_3)_2CH_2OH$, $-CH_2CH_2SO_2CH_3$, $-CH_2OP(O)(OH)_2$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CF_3$, $-CH_2CHF_2$, $-CH(CH_3)CN$, $-C(CH_3)_2CN$, $-CH_2CN$, $-CH_2NH_2$, $-CH_2NHSO_2CH_3$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CO_2H$, $-COCH_3$, $-CO_2CH_3$, $-CO_2C(CH_3)_3$, $-COCH(OH)CH_3$, $-CONH_2$, $-CONHCH_3$, $-CON(CH_3)_2$, $-C(CH_3)_2CONH_2$, $-NH_2$, $-NHCH_3$, $-N(CH_3)_2$, $-NHCOCH_3$, $-N(CH_3)COCH_3$, $-NHS(O)_2CH_3$, $-N(CH_3)C(CH_3)_2CONH_2$, $-N(CH_3)CH_2CH_2S(O)_2CH_3$, $-NHC(=NH)H$, $-NHC(=NH)CH_3$, $-NHC(=NH)NH_2$, $-NHC(=O)NH_2$, $-NO_2$, $=O$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-OCH_2CH_2OCH_3$, $-OCH_2CH_2OH$, $-OCH_2CH_2N(CH_3)_2$, $-O(CH_2CH_2O)_n-$, $(CH_2)_mCO_2H$, $-O(CH_2CH_2O)_nH$, $-OCH_2F$, $-OCHF_2$, $-OCF_3$, $-OP(O)(OH)_2$, $-S(O)_2N(CH_3)_2$, $-SCH_3$, $-S(O)_2CH_3$, and $-S(O)_3H$.

2. The immunoconjugate of claim 1 wherein the antibody is an antibody construct that has an antigen binding domain that binds PD-L1.

3. The immunoconjugate of claim 2 wherein the antibody is selected from the group consisting of atezolizumab, durvalumab, and avelumab, or a biosimilar or a biobetter thereof.

4. The immunoconjugate of claim 1 wherein the antibody is an antibody construct that has an antigen binding domain that binds HER2.

5. The immunoconjugate of claim 4 wherein the antibody is selected from the group consisting of trastuzumab, pertuzumab, margetuximab, and HT-19, or a biosimilar or a biobetter thereof.

6. The immunoconjugate of claim 1 wherein the antibody is an antibody construct that has an antigen binding domain that binds CEA.

7. The immunoconjugate of claim 6 wherein the antibody is labetuzumab, or a biosimilar or a biobetter thereof.

8. The immunoconjugate of claim 1 wherein the antibody is an antibody construct that has an antigen binding domain that binds TROP2.

9. The immunoconjugate of claim 8 wherein the antibody is sacituzumab, or a biosimilar or a biobetter thereof.

10. The immunoconjugate of any one of claims 1 to 9 wherein X^1 is a bond, and R^1 is H.

11. The immunoconjugate of any one of claims 1 to 9 wherein X^2 is a bond, and R^2 is C_1 - C_8 alkyl.

12. The immunoconjugate of any one of claims 1 to 9 wherein X^2 and X^3 are each a bond, and R^2 and R^3 are independently selected from C_1 - C_8 alkyl, $-O-(C_1-C_{12}$ alkyl), $-(C_1-C_{12}$

alkyldiyl)-OR⁵, -(C₁-C₈ alkyldiyl)-N(R⁵)CO₂R⁵, -(C₁-C₁₂ alkyl)-OC(O)N(R⁵)₂, -O-(C₁-C₁₂ alkyl)-N(R⁵)CO₂R⁵, and -O-(C₁-C₁₂ alkyl)-OC(O)N(R⁵)₂.

13. The immunoconjugate of claim 12 wherein R² is C₁-C₈ alkyl and R³ is -(C₁-C₈ alkyldiyl)-N(R⁵)CO₂R⁴.


14. The immunoconjugate of claim 12 wherein R² is -CH₂CH₂CH₃ and R³ is selected from -CH₂CH₂CH₂NHCO₂(t-Bu), -OCH₂CH₂NHCO₂(cyclobutyl), and -CH₂CH₂CH₂NHCO₂(cyclobutyl).


15. The immunoconjugate of claim 12 wherein R² and R³ are each independently selected from -CH₂CH₂CH₃, -OCH₂CH₃, -OCH₂CF₃, -CH₂CH₂CF₃, -OCH₂CH₂OH, and -CH₂CH₂CH₂OH.

16. The immunoconjugate of claim 12 wherein R² and R³ are each -CH₂CH₂CH₃.


17. The immunoconjugate of claim 12 wherein R² is -CH₂CH₂CH₃ and R³ is -OCH₂CH₃.


18. The immunoconjugate of any one of claims 1 to 9 wherein X³-R³ is selected from the group consisting of:






19. The immunoconjugate of any one of claims 1 to 9 wherein X^4 is a bond, and R^4 is H.
20. The immunoconjugate of any one of claims 1 to 9 where R^1 is attached to L.
21. The immunoconjugate of any one of claims 1 to 9 where R^2 or R^3 is attached to L.
22. The immunoconjugate of claim 21 wherein X^3-R^3-L is selected from the group consisting of:






where the wavy line indicates the point of attachment to N.

23. The immunoconjugate of any one of claims 1 to 9 wherein R^4 is C_1 - C_{12} alkyl.
24. The immunoconjugate of any one of claims 1 to 9 wherein R^4 is $-(C_1-C_{12}$ alkyl)diyl)- $N(R^5)-*$; where the asterisk * indicates the attachment site of L.
25. The immunoconjugate of any one of claims 1 to 9 wherein L is $-C(=O)-PEG-$ or $-C(=O)-PEG-C(=O)-$.
26. The immunoconjugate of any one of claims 1 to 9 wherein L is attached to a cysteine thiol of the antibody.
27. The immunoconjugate of any one of claims 1 to 9 wherein for the PEG, m is 1 or 2, and n is an integer from 2 to 10.
28. The immunoconjugate of claim 27 wherein n is 10.
29. The immunoconjugate of any one of claims 1 to 9 wherein L comprises PEP and PEP is a dipeptide and has the formula:




30. The immunoconjugate of claim 29 wherein AA_1 and AA_2 are independently selected from H, $-CH_3$, $-CH(CH_3)_2$, $-CH_2(C_6H_5)$, $-CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, $-CHCH(CH_3)CH_3$, $-CH_2SO_3H$, and $-CH_2CH_2CH_2NHC(O)NH_2$; or AA_1 and AA_2 form a 5-membered ring proline amino acid.
31. The immunoconjugate of claim 29 wherein AA_1 is $-CH(CH_3)_2$, and AA_2 is $-CH_2CH_2CH_2NHC(O)NH_2$.
32. The immunoconjugate of claim 29 wherein AA_1 and AA_2 are independently selected from GlcNAc aspartic acid, $-CH_2SO_3H$, and $-CH_2OPO_3H$.
33. The immunoconjugate of claim 29 wherein PEP has the formula:




wherein AA₁ and AA₂ are independently selected from a side chain of a naturally-occurring amino acid.

34. The immunoconjugate of any one of claims 1 to 9 wherein L comprises PEP and PEP is a tripeptide and has the formula:




35. The immunoconjugate of any one of claims 1 to 9 wherein L comprises PEP and PEP is a tetrapeptide and has the formula:




36. The immunoconjugate of claim 35 wherein
 AA₁ is selected from the group consisting of Abu, Ala, and Val;
 AA₂ is selected from the group consisting of Nle(O-Bzl), Oic and Pro;
 AA₃ is selected from the group consisting of Ala and Met(O)₂; and
 AA₄ is selected from the group consisting of Oic, Arg(NO₂), Bpa, and Nle(O-Bzl).

37. The immunoconjugate of any one of claims 1 to 9 wherein L comprises PEP and PEP is selected from the group consisting of Ala-Pro-Val, Asn-Pro-Val, Ala-Ala-Val, Ala-Ala-Pro-Ala, Ala-Ala-Pro-Val, and Ala-Ala-Pro-Nva.

38. The immunoconjugate of any one of claims 1 to 9 wherein L comprises PEP and PEP is selected from the structures:



39. The immunoconjugate of any one of claims 1 to 9 wherein L is selected from the structures:



where the wavy line indicates the attachment to R⁵.

40. An 8-sulfonyl-2-aminobenzazepine-linker compound selected from Table 2a.
41. An immunoconjugate prepared by conjugation of an antibody with an 8-sulfonyl-2-aminobenzazepine-linker compound selected from Table 2a.
42. A pharmaceutical composition comprising a therapeutically effective amount of an immunoconjugate according to any one of claims 1 to 39, and one or more pharmaceutically acceptable diluent, vehicle, carrier or excipient.
43. A method for treating cancer comprising administering a therapeutically effective amount of an immunoconjugate according to any one of claims 1 to 39 to a patient in need thereof, wherein the cancer is selected from cervical cancer, endometrial cancer, ovarian cancer, prostate cancer, pancreatic cancer, esophageal cancer, bladder cancer, urinary tract cancer, urothelial carcinoma, lung cancer, non-small cell lung cancer, Merkel cell carcinoma, colon cancer, colorectal cancer, gastric cancer, and breast cancer.
44. The method of claim 43, wherein the cancer is susceptible to a pro-inflammatory response induced by TLR7 and/or TLR8 agonism.
45. The method of claim 43, wherein the cancer is selected from triple-negative breast cancer, metastatic Merkel cell carcinoma, and gastroesophageal junction adenocarcinoma.

46. The method of claim 43, wherein the immunoconjugate is administered to the patient intravenously, intratumorally, or subcutaneously.

47. The method of claim 43, wherein the immunoconjugate is administered to the patient at a dose of about 0.01 to 20 mg per kg of body weight.

48. Use of an immunoconjugate according to any one of claims 1 to 39 for treating cancer, wherein the cancer is selected from cervical cancer, endometrial cancer, ovarian cancer, prostate cancer, pancreatic cancer, esophageal cancer, bladder cancer, urinary tract cancer, urothelial carcinoma, lung cancer, non-small cell lung cancer, Merkel cell carcinoma, colon cancer, colorectal cancer, gastric cancer, and breast cancer.

49. A method of preparing an immunoconjugate of Formula I of claim 1 wherein the 8-sulfonyl-2-amino-thienoazepine-linker compound of claim 40 is conjugated with the antibody.