
WHAT IS CLAIMED IS:

1. A method for the inhibition of HIV reverse transcriptase, the treatment or prophylaxis of HIV infection and/or the treatment, prophylaxis and/or delay in the onset or progression of AIDS or ARC in a human subject in need thereof, which comprises orally administering to the human subject an effective amount of a compound of structural Formula I



I,

or a pharmaceutically acceptable salt or co-crystal thereof, wherein X is –F or –Cl, wherein the dosing regimen is less frequent than once-daily.

2. The method of claim 1 for the inhibition of HIV reverse transcriptase, the treatment of HIV infection, and the treatment, prophylaxis and/or delay in the onset or progression of AIDS or ARC in an HIV-infected human subject which comprises orally administering to the subject an effective amount of the compound of Formula I or a pharmaceutically acceptable salt or co-crystal thereof.

3. The method of claim 2 wherein the dosing regimen is a continuous dosing regimen selected from twice-weekly dosing, once-weekly dosing, bi-weekly dosing, twice-monthly dosing or once-monthly dosing.

4. The method of claim 1 for the prophylaxis of HIV infection in a human subject not infected with HIV which comprises orally administering to the subject an effective amount of a compound of Formula I or a pharmaceutically acceptable salt or co-crystal thereof, wherein the dosing regimen has a dosing interval range from about once every 3 days to about once per month.

5. The method of claim 4 wherein the dosing regimen is selected from twice-weekly dosing, once-weekly dosing, bi-weekly dosing, twice-monthly dosing or once-monthly dosing.

6. The method of claim 3 or 5 wherein the dosing regimen is once-weekly.

7. The method of claim 6 wherein the dosage amout is from 1 mg to 50 mg at each dosing interval.

8. The method of claim 3 or 5 wherein the dosing regimen is bi-weekly or twice monthly.

9. The method of claim 8 wherein the unit dosage amount is from 2 mg to 100 mg at each dosing interval.

10. The method of claim 3 or 5 wherein the dosing regimen is once-monthly.

11. The method of claim 10 wherein the dosage amount is from 4 mg to 200 mg at each dosing interval.

12. The method of claim 1 for the prophylaxis of HIV infection which comprises administering to a subject not infected with HIV an effective amount of the compound of structural Formula I or a pharmaceutically acceptable salt or co-crystal thereof.

13. The method of claim 1 for the inhibition of HIV reverse transcriptase, treatment of HIV infection. and the treatment, prophylaxis and/or delay in the onset or progression of AIDS or ARC, which comprises parenterally administering to a subject infected with HIV an effective amount of a compound of Formula I, or a pharmaceutically acceptable salt or co-crystal thereof.