

5 **BENZODIOXOLE, BENZOFURAN,
DIHYDROBENZOFURAN, AND BENZODIOXANE
MELATONERGIC AGENTS**


10

BACKGROUND OF THE INVENTION

15 The invention pertains to novel substituted benzodioxoles,
benzofurans, dihydrobenzofurans, benzodioxanes and related
derivatives having drug and bio-affecting properties and to their
preparation, pharmaceutical formulations and use. In particular, the
invention concerns benzodioxoles, benzofurans, dihydrobenzofurans
and related derivatives bearing substituted amino methyl cyclopropyl
20 groups. These compounds possess melatonergic properties that should
make them useful in treating certain medical disorders.

25 Melatonin (N-acetyl-5-methoxytryptamine) is a hormone which is
synthesized and secreted primarily by the pineal gland. Melatonin levels
show a cyclical, circadian pattern with highest levels occurring during the
dark period of a circadian light-dark cycle. Melatonin is involved in the
transduction of photoperiodic information and appears to modulate a
variety of neural and endocrine functions in vertebrates, including the
regulation of reproduction, body weight and metabolism in photoperiodic
30 mammals, the control of circadian rhythms and the modulation of retinal
physiology.

- 2 -

Melatonin

Recent evidence demonstrates that melatonin exerts its biological effects through specific receptors. Use of the biologically active, radiolabelled agonist [¹²⁵I]-2-iodomelatonin has led to the identification of high affinity melatonin receptors in the CNS of a variety of species. The sequences of two cloned human melatonin receptors have been reported [Reppert, *et al.*, *Proc. Natl. Acad. Sci.* **92**, p. 8734-8738, (1995) and Reppert, *et al.*, *Neuron* **13**, p. 1177-1185, (1994)]. In mammalian brain, autoradiographic studies have localized the distribution of melatonin receptors to a few specific structures. Although there are significant differences in melatonin receptor distribution even between closely related species, in general the highest binding site density occurs in discrete nuclei of the hypothalamus. In humans, specific [¹²⁵I]-2-iodomelatonin binding within the hypothalamus is completely localized to the suprachiasmatic nucleus, strongly suggesting the melatonin receptors are located within the human biological clock.

Exogenous melatonin administration has been found to synchronize circadian rhythms in rats (Cassone, *et al.*, *J. Biol. Rhythms*, **1**:219-229, 1986). In humans, administration of melatonin has been used to treat jet-lag related sleep disturbances, considered to be caused by desynchronization of circadian rhythms (Arendt, *et al.*, *Br. Med. J.*, **292**:1170, 1986). Further, the use of a single dose of melatonin to induce sleep in humans has been claimed by Wurtman in International Patent Application WO 94/07487, published on April 14, 1994.


Thus, melatonin agonists should be particularly useful for the treatment of sleep disorders and other chronobiological disorders. Melatonin agonists would also be useful for the further study of melatonin receptor interactions as well as in the treatment of conditions

affected by melatonin activity, such as depression, jet-lag, work-shift syndrome, sleep disorders, glaucoma, reproduction, cancer, premenstrual syndrome, immune disorders, inflammatory articular diseases and neuroendocrine disorders.

5

Aside from simple indole derivatives of melatonin itself, various bicyclic structures have been prepared and their use as melatonin ligands disclosed. In general these bicyclic amide structures can be represented as:

10




wherein Z is an aryl or heteroaryl system attached by a two carbon bridge to the amide group. Some specific examples follow.

15

Yous, et al. in European Patent Application EP-527,687A, published on February 17, 1993, disclose as melatonin ligands arylethylamines i,

20




wherein Ar' is, inter alia, a substituted or unsubstituted benzo[b]thiophen-3-yl, benzimidazol-1-yl, benzo[b]furan-3-yl, 1,2-benzisoxazol-3-yl, 1,2-benzisothiazol-3-yl, or indazol-3-yl radical; R₁ is, inter alia, an alkyl or cycloalkyl group; and R₂ is hydrogen or lower alkyl.


25

Yous, et al. in European Patent Application EP-506,539A, published on September 30, 1992, claim ligands ii,

- 4 -




wherein A is oxygen or sulfur; X is a methylene group or a bond; and R is
 5 H or lower alkyl when p is 1 and B is defined by the radical iii,



10 wherein R₁ is hydrogen or lower alkyl and R₂ is, inter alia, hydrogen, lower alkyl or cycloalkyl. Alternatively, R is defined by the radical iii when p is 0 or 1 and B is lower alkoxy.

Several naphthalene derivatives have also been disclosed as
 15 melatonin ligands.

Andrieux, et al. in European Patent Application EP-447,285A, published on September 18, 1991, claim amidoalkylnaphthalenes iv,




20

wherein R is lower alkyl; R₁ is hydrogen or lower alkyl; and R₂ is, inter alia, hydrogen, lower alkyl, or cycloalkyl.

25

Yous, et al. in European Patent Application EP-562,956A, published on September 29, 1993, disclose amide and urea naphthalene derivatives v,


5



V

in which R is hydrogen or OR₄ wherein R₄ is, inter alia, hydrogen, alkyl,
 5 cycloalkyl, or cycloalkylalkyl; R₁ is hydrogen or COOR₅ wherein R₅ is
 hydrogen or alkyl; R₂ is hydrogen or alkyl; X is NH or a bond; and R₃ is,
inter alia, alkyl, alkenyl, or cycloalkyl.

Lesieur, et al. in European Patent Application EP-530,087A,
 10 published on March 3, 1993, disclose naphthylethylureas and
 naphthylethylthioureas vi,




vi

15

in which R is hydrogen or OR₃ wherein R₃ is, inter alia, hydrogen, lower
 alkyl, or cycloalkyl; R₁ is hydrogen or lower alkyl; X is oxygen or sulfur;
 and R₂ is, inter alia, lower alkyl or cycloalkyl.

20 Langlois, et al., in Australian Patent Application AU-A-48729/93
 disclose arylalkyl(thio)amides vii as melatonergic ligands,




vii


25

wherein R₁ is hydrogen or lower alkyl; R₂ is hydrogen, halogen, or lower
 alkyl; R₃ and R₄ are identical or different groups including, inter alia,

- hydrogen, halogen, or lower alkyl or R_3 and R_4 , together with the benzene ring which carries them, form a ring-system E_3 chosen from, inter alia, naphthalene, on the understanding that the portion of the ring-system E_3 formed by R_3 and R_4 and the two carbon atoms of the
- 5 benzene ring which carry them is unhydrogenated or partially hydrogenated; R_5 is hydrogen or lower alkyl; and R_6 is,




- 10 wherein X is sulfur or oxygen and R_7 is, inter alia, lower alkyl or alkenyl. Compound viii is included as a specific example,

viii

15

Horn and Dubocovich in European Patent Application EP-420,064A, published on April 3, 1991, disclose 2-amidotetralins ix as melatonin ligands,


ix

20

wherein R_1 is, inter alia, hydrogen, lower alkyl, or lower alkoxy; R_2 is, inter alia, hydrogen, halogen, or lower alkoxy; R_3 is, inter alia, hydrogen,

or lower alkyl; R_4 is, inter alia, lower alkyl, haloalkyl or cycloalkyl; and R_5 is hydrogen, hydroxyl, halogen, oxo, aryl, lower alkyl or alkylaryl.


- Copinga et al, in J. Med. Chem., 36, p. 2891-2898 (1993),
 5 discusses amidomethoxytetralins of structure x and their melatonergic properties.



10

In structure x, R_1 is H or OCH_3 and R_2 is alkyl, haloalkyl, phenylalkyl or phenyl.


- Lesieur et al, in EP-708,099A, published April 24, 1996, disclose
 15 compounds of structure xi, which are useful for the treatment of diseases caused by a melatonin imbalance.



20

wherein --- is a single or double bond; $R_1 = \text{Me}$ or MeNH ; and $X-Y =$
 $-\text{CH}(\text{Me})-\text{CH}_2-$, $\text{CH}_2\text{CH}(\text{OH})-$ or $(\text{CH}_2)_3-$.


- North et al., in International Application WO 95/29173, published
 25 November 2, 1995, disclose naphthalene derivatives of structure xii:



xii

wherein R_1 is a group of the formula $CR_3R_4(CH_2)_pNR_5COR_6$; R_2 is
 5 hydrogen, halogen, C_{1-6} alkyl, OR_7 or CO_2R_7 ; and may be the same or
 different substituent when q is 2; R_3 , R_4 and R_5 , which may be the same
 or different, are hydrogen or C_{1-6} alkyl; R_6 is C_{1-6} alkyl or C_{3-7} cycloalkyl;
 R_7 is hydrogen or C_{1-6} alkyl; n is zero, 1 or 2; p is an integer of 1, 2, 3 or
 4; q is 1 or 2; and the dotted lines indicate the absence or presence of an
 10 additional bond. The North *et al.* compounds are taught to treat
 chronobiological disorders.

In International Application WO 95/17405, published on June 29,
 1995, North *et al.*, disclose compounds of structure *xiii* and teach their
 15 use in the treatment of conditions related to the melatonin system.



xiii

20 wherein R_1 is hydrogen, halogen or C_{1-6} alkyl; R_2 is a group of formula
 $-CR_3R_4(CH_2)_pNR_5COR_6$; R_3 , R_4 and R_5 , which may be the same or
 different, are hydrogen or C_{1-6} alkyl; R_6 is C_{1-6} alkyl or C_{3-7} cycloalkyl; n
 is an integer of 2, 3 or 4; and p is an integer of 1, 2, 3 or 4.

25 The foregoing disclosures do not teach or suggest the novel
 melatonergic benzodioxole, benzofuran or dihydrobenzofurans of the


present invention. The novel compounds of the present invention display melatonergic agonist activity.

5

SUMMARY OF THE INVENTION

The invention provides a novel series of compounds of Formula I

10



15


wherein R, R¹, R², Q¹, Q², X, Y, Z and m are as defined below, including hydrates and solvates thereof which bind to human melatonergic receptors and therefore are useful as melatonergic agents in the treatment of sleep disorders, seasonal depression, shifts in circadian cycles, melancholia, stress, appetite regulation, benign prostatic hyperplasia and related conditions.

20

DETAILED DESCRIPTION OF THE INVENTION

The invention provides a novel series of compounds of Formula I and solvates thereof having the formula:

25



wherein

- Q¹ and Q² each are independently hydrogen or halogen;
X is CH₂, CH or oxygen;
Y is CR³, CR³R⁴ or (CH₂)_n, with n = 1-4;
5 Z is CH₂, CH or oxygen;
R is hydrogen, halogen or C₁₋₄ alkyl in both cases;
m is 1 or 2;
R¹ is C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₃ haloalkyl, C₁₋₆
alkylamino, C₂₋₆ alkenyl, C₁₋₄ alkoxy(C₁₋₄)alkyl, C₁₋₄
10 alkylthio(C₁₋₄)alkyl or C₁₋₄ trifluoromethylalkyl;
R² is hydrogen or C₁₋₄ alkyl; and
R³ and R⁴ each are independently hydrogen or C₁₋₄ alkyl.

- The present invention also provides a method for the treatment of
15 sleep disorders and related conditions, which comprises administering a
therapeutically effective amount of a compound of Formula I or a solvate
or hydrate thereof.

- Q¹ and Q² are selected from H and halogen (i.e., bromine,
20 chlorine, iodine or fluorine). It is most preferred that Q¹ and Q² be H.

X may be CH₂, CH (when a double bond is present) or oxygen.

- Y is CR³ (when a double bond is present), CR³R⁴ or -(CH₂)_n- and
25 n is 1 through 4, but is preferably 1 or 2.

Z may be CH₂, CH (when a double bond is present) or oxygen,
with oxygen being most preferred.

- 30 When X and Y are CH₂ and Z is oxygen or Z and Y are CH₂ and X
is oxygen, the compound is a dihydrobenzofuran. When X and Y are CH
and Z is oxygen or Z and Y are CH and X is oxygen, the compound is a

benzofuran. When X and Z are oxygen and Y is CH₂, the compound is a benzodioxole. When X and Z are oxygen and Y is (CH₂)₂, the compound is benzodioxane. Compounds in which X and Y are CH₂ and Z is oxygen are preferred.

5

Both R groups are the same moiety. Useful R groups include hydrogen, halogen and C₁₋₄ alkyl. R is preferably hydrogen.

m is 1 or 2, with m = 1 preferred.

10

R¹ is one of several types of groups. R¹ is selected from is C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₃ haloalkyl, C₁₋₆ alkylamino, C₂₋₆ alkenyl, C₁₋₄ alkoxy(C₁₋₄)alkyl, C₁₋₄ alkylthio(C₁₋₄)alkyl or C₁₋₄ trifluoromethylalkyl.

R¹ is preferably C₁₋₆ alkyl or C₃₋₆ cycloalkyl.

15

R² is hydrogen or C₁₋₄ alkyl. R² is preferably hydrogen.

R³ and R⁴ are hydrogen or C₁₋₄ alkyl. It is preferred that R³ and R⁴ both be hydrogen. It is also preferred that R³ is hydrogen and R⁴ is methyl. When R³ is hydrogen and R⁴ is methyl, both enantiomers and racemate are preferred.

20

"Alkyl" means a monovalent straight or branched chain group of the formula C_xH_{2x+1}, with x being the number of carbon atoms.

25

"Y — X" and "Y — Z" refer to a single bond or double bond attachment when defined by the substituents X, Y, and Z.

"Cycloalkyl" groups are monovalent cyclic moieties containing at least 3 carbon atoms and conforming to the formula C_xH_(2x-1), with x being the number of carbon atoms present. The cyclopropyl group is a preferred cycloalkyl moiety.

30

"Haloalkyl" includes straight and branched chain hydrocarbon radicals bearing from 1 to 3 halogen moieties. "Halogen" means F, Cl, Br or I. Preferred halogens in haloalkyl moieties of R₁ include F and Cl.

- 5 "Alkylamino" refers to -NH-alkyl substituents containing 1 to 6 carbon atoms, preferably -NHCH₃ or -NHCH₂CH₃ groups.

Preferred compounds have IC₅₀ values of 250 nM or less in melatonergic binding tests described herein.

10

One group of preferred compounds include the benzofurans of Formula I wherein the group, -X-Y-Z-, consists of -CH = CH-O- and -CH = CCH₃-O-.

15

Some preferred compounds of this group include:

- (-)-(trans)-N-[[2-(2-benzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane
carboxamide;
(-)-(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
(trans)-N-[[2-(2-methyl-benzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
20 (trans)-N-[[2-(2-methyl-benzofuran-4-yl)cycloprop-1-yl]methyl]-
propanamide;
(trans)-N-[[2-(2-methyl-benzofuran-4-yl)cycloprop-1-yl]methyl]-
butanamide;
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
25 (trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]-2-
methylpropanamide;
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane
30 carboxamide; and
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]-N-methylurea.

A second group of preferred compounds include the dihydrobenzo-furans of Formula I wherein the group, -X-Y-Z-, consists of -CH₂-CH₂-O-, -CH₂-C(CH₃)₂-O- and -CH₂-CH(CH₃)-O-.

- 5 Some preferred compounds of this second group include:
- (+)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
- 10 (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-N-methylurea;
- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]methoxyacetamide;
- 15 (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropanecarboxamide
- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]trifluoroacetamide;
- 20 (trans)-N-[[2-(2,3-dihydro-2,2-dimethyl-benzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (trans)-N-[[2-(2,3-dihydro-2,2-dimethyl-benzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
- 25 (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- 30 (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-2-methylpropanamide;

- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]chloroacetamide;
- (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- 5 (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]-2-methyl]propanamide;
- (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
- (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]methoxyacetamide;
- 10 (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide;
- (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- 15 (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- 20 (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- (trans)-N-[[2-(2,3-dihydro-5,7-diiodobenzofuran-4-yl)cyclopropyl]methyl]propanamide; and
- 25 (trans)-N-[[2-(2,3-dihydro-5-iodobenzofuran-4-yl)cyclopropyl]methyl]propanamide.

A third group of preferred compounds include the benzodioxoles of Formula I wherein the group, -X-Y-Z-, consists of -O-CH₂-O-.

30

Some preferred compounds in the third group include:

- (trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]acetamide;
- (-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]acetamide;

- (trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]propanamide;
 (-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]
 propanamide;
 (trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]butanamide;
 5 (-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]
 butanamide;
 (trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]cyclopropane
 carboxamide;
 (-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]
 10 cyclopropane carboxamide;
 (trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]-2-
 methylpropanamide;
 (-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]-2-
 methylpropanamide;
 15 (trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]-N'-ethylurea;
 and
 (-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]-N'-
 ethylurea.

- 20 Another group of preferred compounds include the
 benzodioxanes of Formula I wherein the group, -X-Y-Z-, consists of
 -O-(CH₂)₂-O-.

Some preferred benzodioxanes include:

- 25 (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-
 yl]methyl]propanamide;
 (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-
 yl]methyl]acetamide;
 (+)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-
 30 yl]methyl]-N-methylurea
 (+)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-
 yl]methyl]-2-methoxyacetamide;
 (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-
 yl]methyl]cyclopropanecarboxamide;

- (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]butanamide;
 (+)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]propanamide;
 5 (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]trifluoroacetamide;
 (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]-3,3,3-trifluoropropanamide;
 (+)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]propanamide;
 10 (trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]acetamide;
 (trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]propanamide;
 15 (trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]butanamide;
 (trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]-2-methylpropanamide;
 (trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide;
 20 (trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]-N-methylurea; and
 (trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]-N-ethylurea.

25

Still another group of preferred compounds include the benzofurans of Formula I wherein the group, -X-Y-Z, consists of -O-CH=CH- and -O-CCH₃=CH-.

30

Some preferred compounds of this group include:

- (trans)-N-[[2-(benzofuran-7-yl)cycloprop-1-yl]methyl]acetamide;
 (trans)-N-[[2-(benzofuran-7-yl)cycloprop-1-yl]methyl]propanamide;
 (trans)-N-[[2-(benzofuran-7-yl)cycloprop-1-yl]methyl]butanamide; and

(trans)-N-[[2-(benzofuran-7-yl)cycloprop-1-yl]methyl]-cyclopropanecarboxamide.


Still yet another group of preferred compounds include the
5 benzopyrans of Formula I wherein the group, -X-Y-Z, consists of
-CH₂-(CH₂)₂-O-.

Some preferred compounds of this group include:


- (trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]propanamide;
10 (trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide;
(-)-(trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]propanamide;
15 (-)-(trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]acetamide;
(trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]butanamide; and
(+)-(trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]propanamide.
20

Additionally, compounds of Formula I encompass all pharmaceutically acceptable solvates, particularly hydrates, thereof. The present invention also encompasses diastereomers as well as optical
25 isomers, e.g. mixtures of enantiomers including racemic mixtures, as well as individual enantiomers and diastereomers, which arise as a consequence of structural asymmetry in certain compounds of Formula I. Separation of the individual isomers or selective synthesis of the individual isomers is accomplished by application of various methods
30 which are well known to practitioners in the art.

Compounds of Formula I can be prepared using the overall processes shown in the following Reaction Schemes:

Reaction Scheme 1


- The syntheses of the 4-aryl-propenoic acid derivatives, **2** and **3**, are shown in Reaction Scheme 1. The starting aldehydes, **1**, can be prepared by methods well known to those skilled in the art.
- Condensation of malonic acid with the aldehydes, **1**, in solvents such as pyridine with catalysts such as piperidine or pyrrolidine, gives the 4-aryl-propenoic acid, **2**. Subsequent conversion of the acid to the acid chloride using reagents such as thionyl chloride, phosphoryl chloride, or the like, followed by reaction with N,O-dimethyl hydroxylamine gives the amide intermediate **3** in good yields. Alternatively, aldehyde **1** can be converted directly to amide **3** using reagents such as diethyl (N-methoxy-N-methyl-carbamoylmethyl)phosphonate with a strong base such as sodium hydride.

Reaction Scheme 2


The conversion of the amide intermediate 3 to the racemic, trans-cyclopropane carboxaldehyde intermediate, 4, is shown in Reaction Scheme 2. Intermediate 3 was allowed to react with cyclopropanating reagents such as trimethylsulfoxonium iodide and sodium hydride in solvents such as DMF, THF, or the like. Subsequent reduction using reagents such as LAH in solvents such as THF, ethyl ether, or the like, gives the racemic, trans-cyclopropane carboxaldehyde intermediates, 4.

Reaction Scheme 3

10



Racemic cyclopropane intermediate 5 (R = halogen) can be prepared from intermediate 2 as shown in Reaction Scheme 3. Intermediate 2 was converted to the corresponding allylic alcohol by treatment with reducing agents such as sodium borohydride plus iodine in solvents such as THF. Subsequent acylation using reagents such as acetic anhydride in pyridine or acetyl chloride gave the allylic acetate which was allowed to react with cyclopropanating reagents such as sodium chloro-difluoroacetate in diglyme to provide the racemic, trans-cyclopropane acetate intermediates, 5.

Reaction Scheme 4

- 5 The conversion of the acid 2 to the chiral cyclopropane carboxaldehyde intermediate, (-)-(trans)-4, is shown in Reaction Scheme 4. Intermediate 2 is condensed with (-)-2,10-camphorsultam under standard conditions, and then cyclopropanated in the presence of catalysts such as palladium acetate using diazomethane generated from reagents such as 1-methyl-3-nitro-1-nitrosoguanidine. Subsequent
- 10 reduction using reagents such as LAH in solvents such as THF, followed by oxidation of the alcohol intermediates using reagents such as DMSO/oxalyl chloride, or PCC, gives the cyclopropane carboxaldehyde intermediate, (-)-(trans)-4, in good yields. The enantiomer, (+)-(trans)-4,
- 15 can also be obtained employing a similar procedure using (+)-2,10-camphorsultam in place of (-)-2,10-camphorsultam.


When it is desired to prepare compounds of Formula I wherein $m = 2$, the alcohol intermediate may be activated in the conventional

20 manner such as with mesyl chloride and treated with sodium cyanide


followed by reduction of the nitrile group with a reducing agent such as LAH to produce the amine intermediate 6.

Reaction Scheme 5


5



Reaction Scheme 5 shows the conversion of intermediates 4 and 5 to the amine intermediate, 7, and the subsequent conversion of 6 or 7 to compounds of Formula I. The carboxaldehyde intermediate, 4, is condensed with hydroxylamine and then reduced with reagents such as LAH to give the amine intermediate, 7. The acetate intermediate 5 is hydrolyzed with potassium hydroxide to the alcohol, converted to the mesylate with methane sulfonyl chloride and triethyl amine in CH_2Cl_2 and then converted to the azide by treatment with sodium azide in solvents such as DMF. Subsequent reduction of the azide group with a reducing agent such as LAH produced the amine intermediate 7. Further reaction of 6 or 7 with acylating reagents gives compounds of Formula I. Suitable acylating agents include carboxylic acid halides, anhydrides, acyl imidazoles, alkyl isocyanates, alkyl isothiocyanates, and carboxylic acids in the presence of condensing agents, such as carbonyl imidazole, carbodiimides, and the like.

Reaction Scheme 6

- 5 Reaction Scheme 6 shows the alkylation of secondary amides of Formula I ($R^2 = H$) to give tertiary amides of Formula I ($R^2 = \text{alkyl}$). The secondary amide is reacted with a base such as sodium hydride, potassium tert-butoxide, or the like, and then reacted with an alkylating reagent such as alkyl halides, alkyl sulfonate esters, or the like to
- 10 produce tertiary amides of Formula I.

Reaction Scheme 7

- 15 Reaction Scheme 7 shows the halogenation of compounds of Formula I. The carboxamides, I ($Q^1 = Q^2 = H$), are reacted with excess amounts of halogenating agents such as iodine, N-bromosuccinimide, or the like to give the dihalo-compounds of Formula I ($Q^1 = Q^2 = \text{halogen}$).
- 20 Alternatively, a stoichiometric amount of these halogenating agents can be used to give the monohalo-compounds of Formula I ($Q^1 = H$, $Q^2 = \text{halogen}$; or $Q^1 = \text{halogen}$, $Q^2 = H$). In both cases, additives such as lead IV tetraacetate can be used to facilitate the reaction.

Biological Activity of the Compounds

The compounds of the invention are melatonergic agents. They have been found to bind human melatonergic receptors expressed in a stable cell line with good affinity. Further, the compounds are agonists as determined by their ability, like melatonin, to block the forskolin-stimulated accumulation of cAMP in certain cells. Due to these properties, the compounds and compositions of the invention should be useful as sedatives, chronobiotic agents, anxiolytics, antipsychotics, analgesics, and the like. Specifically, these agents should find use in the treatment of stress, sleep disorders, seasonal depression, appetite regulation, shifts in circadian cycles, melancholia, benign prostatic hyperplasia and related conditions.

15 Melatonergic Receptor Binding Activity

1. Reagents:

- (a) TME = 50 mM Tris buffer containing 12.5 mM MgCl_2 , and 2 mM EDTA, pH 7.4 at 37°C with concentrated HCl.
- (b) Wash buffer: 20 mM Tris base containing 2 mM MgCl_2 , pH 7.4 at room temperature.
- (c) 10^{-4} M melatonin (10^{-5} M final concentration).
- (d) 2-[^{125}I]-iodomelatonin, 0.1 M final concentration

2. Membrane Homogenates:

The melatonin ML_{1a} receptor cDNA was subcloned into pcDNA3 and introduced into NIH-3T3 cells using Lipofectamine. Transformed NIH-3T3 cells resistant to geneticin (G-418) were isolated, and single colonies expressing high levels of 2-[^{125}I]-iodomelatonin binding were isolated. Cells are maintained in DMEM supplemented with 10% calf serum and G-418 (0.5 g/liter). Cells are grown to confluency in T-175

flasks, scraped using Hank's balanced salt solution, and frozen at -80°C. For preparing membrane homogenates, pellets are thawed on ice, and resuspended in TME buffer in the presence of 10 µg/ml aprotinin and leupeptin, and 100 µM phenylmethylsulfonylfluoride. The cells were
5 then homogenized using a dounce homogenizer, and centrifuged. The resulting pellet was resuspended with dounce homogenizer in TME (supplemented with the above protease inhibitors) and frozen. On the day of assay, the small aliquot was thawed on ice and resuspended in ice cold TME (1:50-1:100 v/v) and held on ice until assayed.


10

3. Incubation: 37°C for 1 hour. Reaction is terminated by filtration. Filters are washed 3 times.

4. References: Reppert, et al., Neuron, 13, p. 1177-1185 (1994).

15 The binding data for some compounds of Formula I are shown in Table 1.

Table 1: Melatonin Binding for some Compounds of Formula I



Ex. No.	R	R ¹	X-Y-Z	Melatonin Binding Affinity (IC ₅₀) ^a
2	H	Et	CH ₂ -CH ₂ -O	+++
3	H	Me	O-CH ₂ -O	+++
11	H	iPr	O-CH ₂ -O	++
18	H	cPr	O-(CH ₂) ₂ -O	++
23	H	Et	CH=CH-O	+++
29	H	Et	CH ₂ -CH ₂ -O	+++
30	H	nPr	CH ₂ -CH ₂ -O	+++
42	H	Et	CH ₂ -(CH ₂) ₂ -O	++
51	H	NHEt	O-(CH ₂) ₂ -O	+
52	H	vinyl	O-(CH ₂) ₂ -O	+++
54	H	CF ₃	O-(CH ₂) ₂ -O	+++
60	H	Me	O-(CH ₂) ₃ -O	++
61	H	Et	CH ₂ -(CH ₂) ₃ -O	+
67	H	cPr	O-CH ₂ -CH ₂	++
76	H	nPr	CH=CMe-O	+++
80	H	cPr	O-CH=CH	++
82	H	Et	CH ₂ -(CHMe)-O	+++
88	H	Et	CH ₂ -(CHMe)-O	+++
95	F	Et	CH=(CMe)-O	++
96	H	Et	CH ₂ -(CH ₂) ₂ -O	+++

a = IC₅₀ values for ML_{1a} human melatonin receptor binding

+ = 250 nM > IC₅₀ > 100 nM

++ = 100 nM > IC₅₀ > 10 nM

+++ = 10 nM > IC₅₀

The compounds of the present invention have affinity for receptors of the endogenous pineal hormone, melatonin, as determined in a receptor binding assays, as described above in Table 1 for the ML_{1a} (human) receptors. Melatonin is involved in the regulation of a variety of biological rhythms and exerts its biological effects via interaction with specific receptors. There is evidence that administration of melatonin agonists are of clinical utility in the treatment of various conditions regulated by melatonin activity. Such conditions include depression, jet-lag, work-shift syndrome, sleep disorders, glaucoma, some disorders associated with reproduction, cancer, benign prostatic hyperplasia, immune disorders and neuroendocrine disorders.

For therapeutic use, the pharmacologically active compounds of Formula I will normally be administered as a pharmaceutical composition comprising as the (or an) essential active ingredient at least one such compound in association with a solid or liquid pharmaceutically acceptable carrier and, optionally, with pharmaceutically acceptable adjuvants and excipients employing standard and conventional techniques.

The pharmaceutical compositions include suitable dosage forms for oral, parenteral (including subcutaneous, intramuscular, intradermal and intravenous) transdermal, bronchial or nasal administration. Thus, if a solid carrier is used, the preparation may be tableted, placed in a hard gelatin capsule in powder or pellet form, or in the form of a troche or lozenge. The solid carrier may contain conventional excipients such as binding agents, fillers, tableting lubricants, disintegrants, wetting agents and the like. The tablet may, if desired, be film coated by conventional techniques. If a liquid carrier is employed, the preparation may be in the form of a syrup, emulsion, soft gelatin capsule, sterile vehicle for injection, an aqueous or non-aqueous liquid suspension, or may be a dry product for reconstitution with water or other suitable vehicle before use. Liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, wetting agents, non-aqueous vehicle (including edible oils), preservatives, as well as flavoring and/or coloring agents. For parenteral administration, a vehicle normally will

comprise sterile water, at least in large part, although saline solutions, glucose solutions and like may be utilized. Injectable suspensions also may be used, in which case conventional suspending agents may be employed. Conventional preservatives, buffering agents and the like
5 also may be added to the parenteral dosage forms. Particularly useful is the administration of a compound of Formula I in oral dosage formulations. The pharmaceutical compositions are prepared by conventional techniques appropriate to the desired preparation containing appropriate amounts of the active ingredient, that is, the
10 compound of Formula I according to the invention. See, for example, Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA, 17th edition, 1985.

In making pharmaceutical compositions containing compounds of
15 the present invention, the active ingredient(s) will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a vehicle, excipient or medium for the active ingredient.
20 Thus, the composition can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing for example up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile
25 packaged powders.

Some examples of suitable carriers and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate,
30 microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, methyl cellulose, methyl- and propylhydroxybenzoates, talc, magnesium stearate and mineral oil. The formulations can additionally include lubricating agents, wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The

compositions of the invention may be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient.

5 The dosage of the compounds of Formula I to achieve a therapeutic effect will depend not only on such factors as the age, weight and sex of the patient and mode of administration, but also on the degree of melatonergic activity desired and the potency of the particular compound being utilized for the particular disorder or condition
10 concerned. It is also contemplated that the treatment and dosage of the particular compound may be administered in unit dosage form and that the unit dosage form would be adjusted accordingly by one skilled in the art to reflect the relative level of activity. The decision as to the particular dosage to be employed (and the number of times to be administered per
15 day) is within the discretion of the physician, and may be varied by titration of the dosage to the particular circumstances of this invention to produce the desired therapeutic effect.

 The compositions are preferably formulated in a unit dosage form,
20 each dosage containing from about 0.1 to 100 mg, more usually 1 to 10 mg, of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in
25 association with the required pharmaceutical carrier.

 These active compounds are effective over a wide dosage range. For example, dosages per day will normally fall within the range of about 0.1 to 500 mg. In the treatment of adult humans, the range of about 0.1
30 to 10 mg/day, in single or divided doses, is preferred. Generally, the compounds of the invention may be used in treating sleep and related disorders in a manner similar to that used for melatonin.

However, it will be understood that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances including the condition to be treated, the choice of compound to be administered, the chosen route of administration, the age, weight, and response of the individual patient, and the severity of the patient's symptoms.

The compounds which constitute this invention, their methods of preparation and their biologic actions will appear more fully from consideration of the following examples, which are given for the purpose of illustration only and are not to be construed as limiting the invention in sphere or scope.

DESCRIPTION OF SPECIFIC EMBODIMENTS

In the following examples, used to illustrate the foregoing synthetic processes, all temperatures are expressed in degrees Celsius and melting points are uncorrected. Proton magnetic resonance (^1H NMR) spectra were determined in the solvents indicated and chemical shifts are reported in δ units downfield from the internal standard tetramethylsilane (TMS) and interproton coupling constants are reported in Hertz (Hz). Splitting patterns are designated as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad peak; dd, doublet of doublet; bd, broad doublet; dt, doublet of triplet; bs, broad singlet; dq, doublet of quartet. The infrared (IR) spectral descriptions include only absorption wave numbers (cm^{-1}) having functional group identification value. The IR determinations were employed using the compound neat as a film or by employing potassium bromide (KBr) as diluent. Optical rotations $[\alpha]_D^{25}$ were determined in the solvents and concentration indicated. The elemental analyses are reported as percent by weight.

Preparation of Intermediates of Formula 1

5

Preparation 1**Benzofuran-4-carboxaldehyde****Step 1: N-Methoxy-N-methyl-benzofuran-4-carboxamide**

10

A mixture of benzofuran-4-carboxylic acid [Eissenstat, *et al.*, J. Medicinal Chemistry, **38** (16) 3094-3105 (1995)] (2.8 g, 17.4 mmol) and thionyl chloride (25 mL) was heated to reflux for 2 h and then concentrated in vacuo. The solid residue was dissolved in ethyl acetate (50 mL) and a solution of N,O-dimethylhydroxylamine hydrochloride (2.8 g) in saturated NaHCO₃ (60 mL) was added with stirring. After stirring for 1.5 h, the ethyl acetate layer was separated. The aqueous layer was extracted with ethyl acetate. The ethyl acetate extracts were combined, washed with saturated NaHCO₃ and concentrated in vacuo to give an oil (3.2 g, 95.4%).

20

Step 2: Benzofuran-4-carboxaldehyde

A solution of N-methoxy-N-methyl-benzofuran-4-carboxamide (3.2 g, 16.6 mmol) in THF (100 mL) was cooled to -45°C and then LAH (0.7 g, 18.7 mmol) was added. The mixture was stirred for 15 min, allowed to warm to -5°C, and then re-cooled to -45°C. Saturated KHSO₄ (25 mL) was added with vigorous stirring, and the mixture was allowed to warm to room temperature. The precipitate was filtered and washed with acetone. The filtrate was concentrated in vacuo to give an oil (2.3 g, 94%).

25
30

Preparation 2

2,3-Dihydrobenzofuran-4-carboxaldehyde

5 Step 1: 2,3-Dihydrobenzofuran-4-carboxylic acid

Benzofuran-4-carboxylic acid (10.0 g, 61.7 mmol) was hydrogenated (60 psi) in acetic acid (100 mL) over 10% Pd/C (2 g) for 12 hr. The mixture was filtered and the filtrate was diluted with water (500 mL) to give 2,3-dihydrobenzofuran-4-carboxylic acid as a white powder (8.4 g, 83%). A
10 sample was recrystallized from isopropanol to give fine white needles (mp: 185.5-187.5°C).

15 Step 2: (2,3-Dihydrobenzofuran-4-yl)methanol

A solution of 2,3-dihydrobenzofuran-4-carboxylic acid (10 g, 61 mmol) in THF (100 mL) was stirred as LAH (4.64 g, 122 mmol) was slowly added. The mixture was heated to reflux for 30 min. The mixture was cooled and quenched cautiously with ethyl acetate and then with 1N HCl
20 (150 mL). The mixture was then made acidic with 12 N HCl until all the inorganic precipitate dissolved. The organic layer was separated, and the inorganic layer was extracted twice with ethyl acetate. The organic layers were combined, washed twice with brine, and then concentrated in vacuo. This oil was Kugelrohr distilled to a clear oil that crystallized
25 upon cooling (8.53 g, 87.6%).

Step 3: 2,3-Dihydrobenzofuran-4-carboxaldehyde

DMSO (8.10 mL, 114 mmol) was added at -78°C to a stirred solution of
30 oxalyl chloride in CH₂Cl₂ (40 mL of a 2M solution). A solution of (2,3-dihydrobenzofuran-4-yl)methanol (8.53 g, 56.9 mmol) in CH₂Cl₂ (35 mL) was added dropwise, and the solution stirred at -78°C for 30 min. Triethyl amine (33 mL, 228 mmol) was added cautiously to quench the reaction. The resulting suspension was stirred at room temperature for

30 min and diluted with CH₂Cl₂ (100 mL). The organic layer was washed three times with water, and twice with brine, and then concentrated in vacuo to an oil (8.42 g, 100%) that was used without purification.

5

Preparation 3

2,3,4,5-Tetrahydrobenzoxepin-6-carboxaldehyde

10 Step 1: Ethyl 2-allyl-3-benzyloxybenzoate

A mixture of ethyl 2-allyl-3-hydroxybenzoate (20.6 g, 100 mmol), benzyl bromide (18 g, 105 mmol), and potassium carbonate (17 g, 123 mmol), was heated in DMF to 100°C for 18 hr. The mixture was cooled and
15 diluted with water (500 ml), and extracted with ethyl acetate three times. The ethyl acetate extracts were dried over brine and concentrated in vacuo to a tan oil (29.6 g, 100%).

20 Step 2: Ethyl 2-(3-hydroxypropyl)-3-benzyloxybenzoate

A solution of ethyl 2-allyl-3-benzyloxybenzoate (29.6 g, 100 mmol) in THF (300 ml) was cooled to -10°C under nitrogen. A solution of borane THF complex (110 ml of 1M, 110 mmol) was added dropwise, and the reaction mixture was allowed to warm to room temperature and stirred
25 for 1 hr. A solution of hydrogen peroxide (12 ml) in saturated NaHCO₃ (200 ml) was then slowly added, and the mixture was stirred for 30 min. The mixture was extracted with ethyl acetate twice. The ethyl acetate extracts were washed with water, dried over brine, and then concentrated in vacuo to give a clear oil (27.6 g, 88%).

30

Step 3: Ethyl 2-(3-methanesulfonyloxypropyl)-3-benzyloxybenzoate

A solution of ethyl 2-(3-hydroxypropyl)-3-benzyloxybenzoate (10.19 g, 32.5 mmol), and triethyl amine (4.05 g, 40 mmol) in methylene chloride

(100 ml) was cooled in an ice bath as methane sulfonyl chloride (2.79 ml, 36 mmol) was slowly added. The ice bath was removed and the reaction allowed to warm to room temperature over a 1 hr period. the mixture was diluted with water, and the methylene chloride layer was separated. The methylene chloride layer was washed with water twice, 1 N HCl twice, and then filtered through a pad of silica gel to give a clear solution. This solution was concentrated in vacuo to give a clear oil (12.4 g, 98%).

10 **Step 4: Ethyl 2-(3-cyanopropyl)-3-benzyloxybenzoate**

A mixture of ethyl 2-(3-methanesulfonyloxypropyl)-3-benzyloxybenzoate (19.42 g, 31.7 mmol) and potassium cyanide (2.28 g, 35 mmol) in DMF (50 ml) was heated at 100°C for 8 hr. The mixture was cooled, diluted with water (250 ml), and extracted three times with ethyl acetate. The ethyl acetate extracts were dried with brine, and concentrated in vacuo. The crude product was chromatographed on silica gel using methylene chloride as the eluent to give the product as an oil (5.25 g, 53%).

20 **Step 5: 4-[2-Benzyloxy-6-carboxyphenyl]butanoic acid**

A mixture of ethyl 2-(3-cyanopropyl)-3-benzyloxybenzoate (5.25 g, 16.9 mmol) in 5 N sodium hydroxide (75 ml) was heated to reflux for 18 hr. The solution was cooled and made acidic with 12 N HCl. The white precipitate was collected and air dried (5 g, 94%).

Step 6: 4-[2-Hydroxy-6-carboxy-phenyl]butanoic acid

A solution of 4-[2-benzyloxy-6-carboxy-phenyl]butanoic acid (5 g, 15.9 mmol) in ethanol was hydrogenated at 60 psi for 4 hr over 10% Pd/C (1 g). The mixture was filtered, and the filtrate was concentrated in vacuo to give a white solid (3.95 g, 99%)

Step 7: Ethyl 4-[2-ethoxycarbonyl-6-hydroxyphenyl]butanoic acid

A solution of 4-[2-hydroxy-6-carboxyphenyl]butanoic acid (7.84 g, 35 mmol) in ethanol and sulfuric acid (10 drops) was heated to reflux for 18
5 hr. The solution was neutralized with sodium bicarbonate and cooled. The mixture was filtered and the filtrate was concentrated in vacuo. The resulting oil was dissolved in ethyl acetate, washed with water, dried with brine, and the concentrated in vacuo to give the product as an oil (9.06 g, 92.4%).

10

Step 8: 4-[2-Ethoxycarbonyl-6-hydroxyphenyl]butanoic acid

Ethyl 4-[2-ethoxycarbonyl-6-hydroxyphenyl]butanoic acid (9.06 g, 32.4 mmol) was dissolved in a room temperature solution of potassium
15 hydroxide (3.62g, 64.6 mmol) in water (68 ml). The solution was stirred at room temperature for 2 hr, and then made acidic with 12 N HCl. The mixture was extracted twice with ethyl acetate. The ethyl acetate extracts were combined and extracted with 10% aqueous potassium carbonate three times. These aqueous extracts were made acidic with 12 N HCl
20 and extracted twice with ethyl acetate. The ethyl acetate extracts were concentrated in vacuo to a white solid (6.4 g, 79%).

Step 9: 4-[2-Ethoxycarbonyl-6-hydroxyphenyl]butan-1-ol

25 Borane THF complex (56 ml of 1M in THF, 56 mmol) was slowly added to a solution of 4-[2-ethoxycarbonyl-6-hydroxyphenyl]butanoic acid in THF at 0°C. The solution was stirred for 2 hr and allowed to warm to room temperature. The reaction was quenched slowly with aqueous acetic acid (5 ml of 50%) and stirred until the bubbling subsided. The mixture
30 was concentrated in vacuo to a slurry, which was dissolved in cold 10% potassium carbonate. The solution was extracted with ethyl acetate twice. The organic extracts were dried with brine and concentrated in vacuo to a tan oil that solidified upon standing (5.14 g, 84.9%).

Step 10: Ethyl 2,3,4,5-tetrahydrobenzoxepin-6-carboxylate

A solution of 4-[2-ethoxycarbonyl-6-hydroxyphenyl]butan-1-ol (5.14 g, 21.6 mmol) in THF (20 ml) was added dropwise to an ice bath cooled
5 solution of triphenyl phosphine (7.37 g, 28.1 mmol) and diethyl azodicarboxylate (4.89 g, 28.1 mmol) in THF (60 ml). The reaction was stirred for 18 hr and then concentrated in vacuo. The residue was purified by chromatography on silica gel using methylene chloride as the eluent to give the product as an oil (2.85 g, 60%).

10

Step 11: 2,3,4,5-Tetrahydrobenzoxepin-6-methanol

A solution of ethyl 2,3,4,5-tetrahydrobenzoxepin-4-carboxylate (2.85 g, 13 mmol) in THF (30 ml) was stirred as LAH (1 g, 26 mmol) was slowly
15 added. The mixture was stirred for 18 hr, and then quenched by the sequential addition of water (1 ml), 15% sodium hydroxide (1 ml), and water (3 ml). The mixture was filtered and the filter cake washed with ethanol. The filtrate was concentrated in vacuo to an oil (2.3 g, 100%).

Step 12: 2,3,4,5-Tetrahydrobenzoxepin-6-carboxaldehyde

DMSO (1.85 ml, 26 mmol) was added slowly to a -78°C solution of oxalyl chloride in methylene chloride (10 ml of 2 M, 20 mmol). To this solution was added slowly a solution of 2,3,4,5-tetrahydrobenzoxepin-4-
25 methanol (2.3 g, 13 mmol) in methylene chloride, and the mixture was stirred 30 min at -78°C. Triethyl amine (7.53 ml, 52 mmol) was added slowly to the reaction mixture, which was then allowed to warm to room temperature. The mixture was diluted with methylene chloride (50 ml) and washed with water and then brine. The organic layer was
30 concentrated in vacuo to an oil (2.28 g, 100%).

Preparation 4

Benzofuran-7-carboxaldehyde

5 **Step 1: Methyl 2-allyloxybenzoate**

A mixture of allyl bromide (152.4 g, 1.27 mol), methyl salicylate (162.44 g, 1.06 mol), and potassium carbonate (219.75 g, 1.59 mol) in THF (600 ml) and DMF (600 ml) was heated to reflux for 6 hr. The mixture was
10 poured into water (3 L) and extracted with ethyl acetate three times. The ethyl acetate extracts were combined and washed with water and dried with brine. The ethyl acetate extracts were concentrated in vacuo to a yellow oil that was vacuum distilled to a clear oil (163.45 g, 80%).

15 **Step 2: Methyl 3-allyl-salicylate**

Methyl 2-allyloxybenzoate (163.5 g, 848 mmol) was heated to 220°C for 1 hr, and then vacuum distilled to give the product (163.5 g, 100%).

20 **Step 3: Methyl benzofuran-7-carboxylate**

Ozone was bubbled through a solution of methyl 3-allyl-salicylate (30 g, 156 mmol) at -78°C for 2 hr until no starting material was present on TLC. The reaction was quenched with dimethyl sulfide and stirred at
25 room temperature for 18 hr. The mixture was concentrated in vacuo, and the residue was dissolved in ether. The ether solution was washed with brine three times, and then concentrated in vacuo to a green oil. This oil was dissolved in toluene and heated to reflux with sulfuric acid (0.5 ml) for 4 hr. Sodium carbonate (5 g) was added, and the mixture was cooled
30 and then filtered. The filtrate was concentrated in vacuo to a dark oil that was purified by chromatography on silica gel using methylene chloride as the eluent to give the product as a green oil (12 g, 44%).

Step 4: Benzofuran-7-methanol

A solution of methyl benzofuran-7-carboxylate (5.34 g, 30 mmol) was added to a suspension of LAH (2.31 g, 61 mmol) in THF and then heated
5 to reflux for 30 min. The reaction was quenched with ethyl acetate and water. The mixture was made acidic with 12 N HCl until all the precipitate dissolved. The ethyl acetate layer was separated, washed with water, dried with brine, and concentrated in vacuo to give a yellow oil (4.03 g, 91%).

10

Step 5: Benzofuran-7-carboxaldehyde

DMSO (2.87 ml, 40 mmol) was added slowly to a -78°C solution of oxalyl chloride in methylene chloride (20 ml of 2 M, 40 mmol). To this solution
15 was added slowly a solution of benzofuran-7-methanol (4.03 g, 27 mmol) in methylene chloride, and the mixture was stirred 30 min at -78°C. Triethyl amine (30 ml) was added slowly to the reaction mixture, which was then allowed to warm to room temperature. The mixture was diluted with methylene chloride and washed with water and then with
20 brine. The organic layer was concentrated in vacuo to an oil (3.16 g, 80%).

Preparation 5**25 2,3-Dihydro-benzofuran-7-carboxaldehyde****Step 1: Methyl 2,3-dihydro-benzofuran-7-carboxylate**

Methyl benzofuran-7-carboxylate (12 g, 68 mmol) was hydrogenated at
30 60 psi over 10% Pd/C (2 g) in acetic acid (60 ml) for 18 hr. The mixture was filtered, and the filtrate was concentrated in vacuo to give a pale green oil (12 g, 100%)

Step 2: 2,3-Dihydro-benzofuran-7-methanol

Methyl 2,3-dihydro-benzofuran-7-carboxylate (12 g, 68 mmol) was reduced with LAH (5.14 g, 136 mmol) in THF similar to the above procedures to give the product as a dark oil (8.13 g, 80%).

Step 3: 2,3-Dihydro-benzofuran-7-carboxaldehyde

A solution of 2,3-dihydro-benzofuran-7-methanol (8.13 g, 54.5 mmol) was oxidized using DMSO, oxalyl chloride, and triethyl amine similar to the above procedures to give the product as an oil (7.7 g, 95%).

Preparation 6**15 2,2-Dimethyl-2,3-dihydro-benzofuran-4-carboxaldehyde****Step 1: Ethyl-3-(2-methylpropenyloxy)benzoate**

A mixture of 2-methyl-3-chloropropene (64.2 g, 710 mmol), ethyl 3-hydroxybenzoate (48.21 g, 590 mmol), and potassium carbonate (122.3 g, 890 mmol) in THF (600 ml) and DMF (600 ml) was heated to reflux for 6 hr. The mixture was poured into water (3 L) and extracted with ethyl acetate three times. The ethyl acetate extracts were combined and washed with water and dried with brine. The ethyl acetate extracts were concentrated in vacuo to a yellow oil that was vacuum distilled to a clear oil (112.9 g, 87%).

Step 2: Ethyl 2,2-dimethyl-2,3-dihydro-benzofuran-4-carboxylate

30 A mixture of ethyl -3-(2-methylpropenyloxy)benzoate (23.6 g, 107 mmol) was stirred with SCX resin (2 g) at 220°C for 1 hr. The crude product was purified by chromatography on silica gel using methylene chloride as the eluent to give a clear oil (8.7 g, 37%).

Step 3: 2,2-Dimethyl-2,3-dihydro-benzofuran-4-carboxylic acid

Ethyl 2,2-dimethyl-2,3-dihydro-benzofuran-4-carboxylic acid (10 g, 45 mmol) was heated to reflux with sodium hydroxide (16.3 ml of 10 N, 163 mmol) in ethanol (50 ml) for 2 hr. The mixture was concentrated in vacuo, diluted with water and made acidic with 12 N HCl. The precipitate was filtered and air dried (8.6 g, 100%).

Step 4: 2,2-Dimethyl-2,3-dihydro-benzofuran-4-methanol

10

2,2-Dimethyl-2,3-dihydro-benzofuran-4-carboxylic acid (8.6 g, 45 mmol) was reduced with LAH (3.41 g, 89 mmol) similar to the above procedures to give the product as a clear oil (7.35 g, 93%).

Step 5: 2,2-Dimethyl-2,3-dihydro-benzofuran-4-carboxaldehyde

15

2,2-Dimethyl-2,3-dihydro-benzofuran-4-methanol (7.35 g, 41 mmol) was oxidized using DMSO, oxalyl chloride, and triethyl amine similar to the above procedures to give the product as a clear oil (8.87 g, 100%).

20

Preparation 7**2-Methyl-benzofuran-4-carboxaldehyde****Step 1: Ethyl 2-methyl-benzofuran-4-carboxylate**

25

Ozone was bubbled through a solution of ethyl -3-(2-methylpropenyloxy) benzoate (10 g, 45 mmol) at -78°C for 2 hr until no starting material was present on TLC. The reaction was quenched and the product isolated similar to the above procedures to give the product as a white solid (8.05 g, 87%).

30

Step 2: 2-Methyl-benzofuran-4-methanol

Ethyl 2-methyl-benzofuran-4-carboxylic acid (5.11 g, 26.6 mmol) was reduced with LAH (2 g, 53 mmol) similar to the above procedures to give the product as a clear oil (3.34 g, 78%).

Step 3: 2-Methyl-benzofuran-4-carboxaldehyde

2-Methyl-benzofuran-4-methanol (3.34 g, 21 mmol) was oxidized using DMSO, oxalyl chloride, and triethyl amine similar to the above procedures to give the product as a clear oil (3.14 g, 93%).

Preparation 8**2.3-Dihydro-1,4-benzodioxin-5-carboxaldehyde**

2,3-Dihydroxybenzaldehyde (58 g, 420 mmol) was added to a refluxing mixture of dibromoethane (107.4 g, 570 mmol), sodium hydroxide (35.7 g, 890 mmol) and tetrabutyl ammonium bromide (3 g) in water (50 mL). After heating at reflux for 4 h, the mixture was cooled and the organic layer separated, washed with base, dried over sodium sulfate and concentrated in vacuo. The residue was Kugelrohr distilled at 135° to give the product (48 g, 70%) which solidified on standing (mp 61-62°C).
Anal. Calc'd: C, 65.85; H, 4.91.
Found: C, 65.73; H, 4.86.

Preparation 9**2H-3,4-Dihydro-1,5-benzodioxapin-6-carboxaldehyde**

2,3-Dihydroxybenzaldehyde and 1,3-dibromopropane (107.4 g, 570 mmol), sodium hydroxide (35.7 g, 890 mmol) were reacted as described above to give an oil (43%).

Preparation 10

(2H-3,4-Dihydrobenzopyran-5-yl)carboxaldehyde

5 Step 1: (2H-3,4-Dihydrobenzopyran-5-yl)carboxylic acid

Borane THF complex (55 ml of 1M, 55 mmol) was added to a solution of ethyl 2-allyl-3-hydroxybenzoate (10.3 g, 50 mmol) in THF at -10°C. The reaction was allowed to warm to room temperature and stirred for 1 hr. A
10 saturated sodium bicarbonate (100 ml) was slowly added then hydrogen peroxide (6 ml) was added dropwise. The mixture was stirred for 30 min, diluted with ethyl acetate (150 ml). The ethyl acetate layer was separated, washed with water, dried with brine, and concentrated in vacuo to a clear oil. This material was dissolved in THF (100 ml) and
15 added slowly to a solution of triphenylphosphine (20.85 g, 79.5 mmol) and diethyl azodicarboxylate (13.84 g, 79.5 mmol) in THF (250 ml). The solution was stirred for 18 hr and then concentrated in vacuo. The residue was distilled in vacuo to give an oil. This oil was purified by chromatography on silica gel using ethyl acetate/methylene chloride
20 (1:1) as the eluent to give a clear oil (11.3 g). This oil was heated to reflux for 2 hr with sodium hydroxide (25 ml of 10 N, 250 mmol) and water (50 ml). The reaction was cooled and filtered. The filtrate was made acidic with 12 N hydrochloric acid to give a white precipitate (7.96 g, 89%).

25

Step 2: (2H-3,4-Dihydrobenzopyran-5-yl)methanol

(2H,3,4-Dihydrobenzopyran-5-yl)carboxylic acid was reduced with LAH by the above procedures to give the product (97%).

30

Step 3: (2H-3,4-Dihydrobenzopyran-5-yl)carboxaldehyde

(2H-3,4-Dihydrobenzopyran-5-yl)methanol was oxidized using DMSO, oxalyl chloride, and triethyl amine similar to the above procedures to
5 give the product as a clear oil (100%)

Preparation 11**2-Methyl-2,3-dihydrobenzofuran-4-carboxaldehyde**

10

Step 1. Ethyl 2-methyl-2,3-dihydrobenzofuran-4-carboxylate

Ethyl 2-methylbenzofuran-4-carboxylate (10 g), obtained as described in preparation 7, was hydrogenated (60 psi) in acetic acid (100 mL) over
15 10% Pd/C (2 g) for 24 h. The mixture was filtered through Celite and the filter pad washed well with ethyl acetate. The filtrate was concentrated in vacuo to obtain ethyl 2-methyl-2,3-dihydrobenzofuran-4-carboxylate.

Step 2. 2-Methyl-2,3-dihydrobenzofuran-4-methanol

20

A solution of ethyl 2-methyl-2,3-dihydrobenzofuran-4-carboxylate (10 g) in THF (100 mL) was stirred as LAH (4.64 g, 122 mmol) was slowly added. The mixture was heated to reflux for 30 min. The mixture was cooled and quenched cautiously with ethyl acetate and then with 1N HCl
25 (150 mL). The mixture was then made acidic with 12N HCl until all the inorganic precipitate dissolved. The organic layer was separated, and the inorganic layer was extracted twice with ethyl acetate. The organic layers were combined, washed twice with brine, and then concentrated in vacuo to provide the desired alcohol.

30

Step 3. 2-Methyl-2,3-dihydrobenzofuran-4-carboxaldehyde

To a solution of oxalyl chloride(51.6 mL of 2 M solution, 103.36 mmol) in dichloromethane(200 mL) at - 78°C under N₂ was added dropwise

DMSO(9.2 mL, 129.2 mmol) during 10 min. After stirring for 20 min, a solution of alcohol(10.6 g, 64.6 mmol) in dichloromethane(50 mL) was added dropwise during 10 min. The resulting mixture was stirred for 1 h. Triethylamine(36 mL, 258.4 mmol) was added and allowed to warm to room temperature, and stirred for 0.5 h. The reaction was quenched with water(30 mL), washed with brine, dried over MgSO₄, concentrated in vacuo to give a residue. The residue was purified by flash chromatography over silical gel, elution with 6% ethyl acetate in hexane, to give 8.5 g (81%) of the aldehyde as an oil.

10

Preparation of intermediates of Formula 2

15

Preparation 12

(trans)-3-(2,3-Dihydrobenzodioxin-5-yl)propenoic acid

A mixture of 2,3-dihydrobenzodioxin-5-carboxaldehyde (Morishima, et al., Eur. Pat. Appl. 309,766, 5 Apr 89) (9.25 g, 56.4 mmol), malonic acid (11.73 g, 112.8 mmol), pyrrolidine (1 mL), and pyridine (25 mL) was heated to reflux for 2 hr, cooled, and then poured into ice water (300 mL). The white precipitate was filtered, washed with 1N HCl, and air dried (9.83 g, 84.6%).

25

Preparation 13

The following compounds of Formula 2, (a) - (l) were prepared by the general procedure described for the compound of Preparation 12.

30

(a) (trans)-3-(2,3-Dihydrobenzofuran-4-yl)propenoic acid

2,3-Dihydrobenzofuran-4-carboxaldehyde was subjected to the above procedure to give a pale yellow powder that was recrystallized from
5 Isopropanol to give white flakes (95.3%, mp 205-207°C).

Anal. Calc'd for $C_{11}H_{10}O_3$: C, 69.46; H, 5.30.

Found: C, 69.36; H, 5.17.

10 (b) (trans)-3-(Benzofuran-4-yl)propenoic acid

Benzofuran-4-carboxaldehyde was subjected to the above procedure to give a white powder (83%).

15 (c) (trans)-3-(1,3-benzodioxol-4-yl)propenoic acid

1,3-Benzodioxole-4-carboxaldehyde was subjected to the above procedure to give a white powder (99%).

20 (d) (trans)-3-(2,3,4,5-tetrahydrobenzoxepin-6-yl)propenoic acid

2,3,4,5-Tetrahydrobenzoxepin-4-carboxaldehyde was subjected to the above procedure to give this product as a white solid (89%).

25 (e) (trans)-3-(Benzofuran-7-yl)propenoic acid

Benzofuran-7-carboxaldehyde was subjected to the above procedure to give this product as a white solid (100%).

30 (f) (trans)-3-(2,3-Dihydro-benzofuran-7-yl)propenoic acid

2,3-Dihydro-benzofuran-7-carboxaldehyde was subjected to the above procedure to give this product as a white solid (100%).

(g) (trans)-3-(2,2-Dimethyl-2,3-dihydro-benzofuran-4-yl)propenoic acid

2,2-Dimethyl-2,3-dihydro-benzofuran-4-carboxaldehyde was subjected to the above procedure to give this product as a white solid (81%).

5

(h) (trans)-3-(2-Methyl-benzofuran-4-yl)propenoic acid

2-Methyl-benzofuran-4-carboxaldehyde was subjected to the above procedure to give this product as a white solid (97%).

10

(i) (trans)-3-(2,3-Dihydro-1,4-benzodioxin-5-yl)propenoic acid

2,3-Dihydro-1,4-benzodioxan-5-carboxaldehyde was reacted with malonic acid as described above to give a white solid (90%).

15

(j) (trans)-3-(2H-3,4-Dihydro-1,5-benzodioxapin-6-yl)propenoic acid

2H-3,4-Dihydro-1,5-benzodioxapin-6-carboxaldehyde was reacted with malonic acid as described above to give a white solid (74%).

20

(k) (trans)-3-(2H-2,3-Dihydrobenzopyran-4-yl)propenoic acid

(2H-2,3-Dihydrobenzopyran-4-yl)carboxaldehyde was reacted with malonic acid by the above procedures to give the white solid (98%).

25

(l) (trans)-3-(2-Methyl-2,3-dihydrobenzofuran-4-yl)propenoic acid

2-Methyl-2,3-dihydrobenzofuran-4-carboxaldehyde was reacted with malonic acid by the above procedures to give the desired acid (92%).

30

Preparation of intermediates of Formula 3**Preparation 14****5 (trans)-N-Methoxy-N-methyl-3-(2,3-Dihydrobenzodioxin-5-yl)propenamide**

A mixture of (trans)-3-(2,3-dihydrobenzodioxin-5-yl)propenoic acid (9.83 g, 47.7 mmol), thionyl chloride (20 mL), and CH₂Cl₂ (75 mL) was
10 heated to reflux for 1 hr, and then concentrated in vacuo to give a yellow-green solid. This material was dissolved in ethyl acetate (75 mL), and a solution of N,O-dimethylhydroxylamine hydrochloride (9.5 g) in saturated Na₂CO₃ (100 mL) was added with stirring. The mixture was stirred for
15 90 min and then diluted with water and ethyl acetate. The ethyl acetate layer was separated and washed twice with water and twice with saturated Na₂CO₃. The ethyl acetate layer was concentrated in vacuo to give a brown oil. This crude product was chromatographed on silica gel using 1:1 ethyl acetate/methylene chloride as the eluent to give a clear oil that crystallized upon standing (11.1 g, 93.1%).

20

Preparation 15

The following compounds of Formula 3, (a) - (k) were prepared by the general procedure described for the compound of Preparation 14.

25

(a) (trans)-N-Methoxy-N-methyl-3-(benzofuran-4-yl)propenamide

(trans)-3-(Benzofuran-4-yl)-propenoic acid was subjected to the above procedure to give an oil (97.7%).

30

(b) (trans)-N-Methoxy-N-methyl-3-(2,3,4,5-tetrahydrobenzoxepin-6-yl)propenamide

(trans)-3-(2,3,4,5-Tetrahydrobenzoxepin-6-yl)-propenoic acid was subjected to the above procedures to give this product (88.6%).

(c) (trans)-N-Methoxy-N-methyl-3-(benzofuran-7-yl)propenamide

5

(trans)-3-(Benzofuran-7-yl)propenoic acid was subjected to the above procedures to give this product (90%).

(d) (trans)-N-Methoxy-N-methyl-3-(2,3-dihydro-benzofuran-7-yl)propenamide

10

(trans)-3-(2,3-Dihydro-benzofuran-7-yl)propenoic acid was subjected to the above procedures to give this product (68%).

15 (e) (trans)-N-Methoxy-N-methyl-3-(2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)propenamide

(trans)-3-(2,2-Dimethyl-2,3-dihydro-benzofuran-4-yl)propenoic acid was subjected to the above procedures to give this product (92%).

20

(f) (trans)-N-Methoxy-N-methyl-3-(2-methyl-benzofuran-4-yl)propenamide

25 (trans)-3-(2-Methyl-benzofuran-4-yl)propenoic acid was subjected to the above procedures to give this product (100%).

(g) (trans)-N-Methoxy-N-methyl-3-(2,3-dihydro-1,4-benzodioxin-5-yl)propenamide

30 (trans)-3-(2,3-Dihydro-1,4-benzodioxan-5-yl)propenoic acid was subjected to the above procedures to give this product (85%).

(h) (trans)-N-Methoxy-N-methyl-3-(2H-3,4-dihydro-1,5-benzodioxapin-6-yl)propenamide

(trans)-2H-3-(3,4-Dihydro-1,5-benzodioxapin-6-yl)propenoic acid was subjected to the above procedures to give this product (86%).

5 (i) (trans)-N-Methoxy-N-methyl-3-(2H-2,3-dihydrobenzopyran-4-yl)propenamide

(trans)-3-(2H-2,3-Dihydrobenzopyran-4-yl)propenoic acid was subjected to the above procedures to give this product (quantitative yield).

10

(j) (trans)-N-Methoxy-N-methyl-3-(1,3-benzodioxol-4-yl)-propenamide

1,3-Benzodioxole-4-carboxaldehyde was subjected to the above procedure to give a red oil (100%).

15

(k) (trans)-N-Methoxy-N-methyl-3-(2,3-dihydrobenzofuran-4-yl)propenamide

Diethyl (N-methoxy-N-methyl-carbamoylmethyl) phosphonate (4.0 g, 16.7 mmol) was added dropwise to a suspension of sodium hydride (671 mg, 60% dispersion in mineral oil, 16.7 mmol) in THF (75 mL) at 0°C. A solution of 2,3-dihydrobenzofuran-4-carboxaldehyde (3.0 g, 15.2 mmol) in THF (25 mL) was added dropwise. The resulting suspension was allowed to warm to room temperature. After 18 h, water (60 mL) was added and the solution was extracted three times with ethyl acetate. The organic extracts were combined, washed with water and brine, dried over K₂CO₃, and concentrated to give a pale red oil, 3.5 g (100%).

20

25

Preparation of intermediates of Formula 4**Preparation 16****5 (±)-(trans)-2-(2,3-Dihydrobenzofuran-4-yl)cyclopropane-
carboxaldehyde****Step 1: (±)-(trans)-N-Methoxy-N-methyl-2-(2,3-dihydrobenzofuran-4-
yl)cyclopropanecarboxamide**

10

Trimethylsulfoxonium iodide (9.9 g, 45 mmol) was added in small portions to a suspension of sodium hydride (1.8 g, 45 mmol) in DMF (120 mL). After the foaming had subsided (10 min), a solution of (trans)-N-methoxy-N-methyl-3-(2,3-dihydrobenzofuran-4-yl)propenamide (3.5 g, 15 mmol) in DMF (60 mL) was added dropwise, with the temperature maintained between 35-40°C. The mixture was stirred for 3 h at room temperature. Saturated NH₄Cl (50 mL) was added dropwise and the mixture was extracted three times with ethyl acetate. The organic extracts were combined, washed with H₂O and brine, dried over K₂CO₃, and concentrated in vacuo to give a white wax (3.7 g, 100%).

20

**Step 2: (±)-(trans)- 2-(2,3-Dihydrobenzofuran-4-yl)cyclopropane-
carboxaldehyde**

25 A solution of (±)-(trans)-N-methoxy-N-methyl-2-(2,3-dihydrobenzofuran-4-yl)cyclopropanecarboxamide (3.7 g, 15 mmol) in THF (10 mL) was added dropwise to a rapidly stirred suspension of LAH (683 mg, 18 mmol) in THF (50 mL) at -45°C, maintaining the temperature below -40°C throughout. The cooling bath was removed, the reaction was
30 allowed to warm to 5°C, and then the reaction was immediately recooled to -45°C. Potassium hydrogen sulfate (3.4 g, 25.5 mmol) in H₂O (50 mL) was cautiously added dropwise, the temperature maintained below -30°C throughout. The cooling bath was removed and the suspension

was stirred at room temperature for 30 min. The mixture was filtered through Celite and the filter cake was washed with ether. The combined filtrates were then washed with cold 1N HCl, 1N NaOH, and brine. The filtrates were dried over MgSO_4 , and concentrated in vacuo to give a
5 clear oil (2.6 g, 99%).

Preparation 17

The following compounds of Formula 4 (a) - (k) were prepared by the
10 general procedure described for the compound of Preparation 16.

(a) (±)-(trans)-2-(Benzofuran-4-yl)cyclopropanecarboxaldehyde

(trans)-N-Methoxy-N-methyl-3-(benzofuran-4-yl)propenamide was
15 subjected to the above procedure to give an oil (93.3%).

(b) (±)-(trans)-2-(1,3-Benzodioxol-4-yl)cyclopropanecarboxaldehyde

(trans)-N-Methoxy-N-methyl-3-(1,3-benzodioxol-4-yl)propenamide was
20 subjected to the above procedure to give a clear oil (100%).

(c) (±)-(trans)-2-(2,3-Dihydrobenzodioxan-5-yl)cyclopropane-
carboxaldehyde

25 (trans)-N-Methoxy-N-methyl-3-(2,3-dihydrobenzodioxan-5-yl)propenamide was subjected to the above procedure to give an orange oil (90%).

(d) (±)-(trans)-2-(2,3,4,5-Tetrahydrobenzoxepin-6-yl)cyclopropane-
30 carboxaldehyde

(trans)-N-Methoxy-N-methyl-3-(2,3,4,5-tetrahydrobenzoxepin-6-yl)propenamide was subjected to the above procedures to give this product (36%).

(e) (±)-(trans)-2-(Benzofuran-7-yl)cyclopropanecarboxaldehyde

(trans)-N-Methoxy-N-methyl-3-(benzofuran-7-yl)propenamide was subjected to the above procedures to give this product (67%).

5

(f) (±)-(trans)-2-(2,3-Dihydro-benzofuran-7-yl)cyclopropane-carboxaldehyde

(trans)-N-Methoxy-N-methyl-3-(2,3-dihydro-benzofuran-7-yl)propenamide was subjected to the above procedures to give this product (55%).

10

(g) (±)-(trans)-2-(2,2-Dimethyl-2,3-dihydro-benzofuran-4-yl)cyclopropanecarboxaldehyde

15

(trans)-N-Methoxy-N-methyl-3-(2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)propenamide was subjected to the above procedures to give this product (64%).

20

(h) (±)-(trans)-2-(2-Methyl-benzofuran-4-yl)cyclopropane-carboxaldehyde

(trans)-N-Methoxy-N-methyl-3-(2-methyl-benzofuran-4-yl)propenamide was subjected to the above procedures to give this product (100%).

25

(i) (trans)-2-(2,3-dihydro-1,4-benzodioxan-5-yl)cyclopropane-carboxaldehyde

(trans)-N-Methoxy-N-methyl-3-(2,3-dihydro-1,4-benzodioxan-5-yl)propenamide was subjected to the above procedures to give this product (79%).

30

(j) (trans)-2-(2H-3,4-dihydro-1,5-benzodioxapin-6-yl)cyclopropane-carboxaldehyde

(trans)-N-Methoxy-N-methyl-3-(2H-3,4-dihydro-1,5-benzodioxapin-6-yl)propenamide was subjected to the above procedures to give an oil (55%).

5

(k) (±)-(trans)-2-(2H-2,3-Dihydrobenzopyran-4-yl)cyclopropane-carboxaldehyde

(trans)-N-Methoxy-N-methyl-3-(2H-2,3-dihydrobenzopyran-4-yl)propenamide was subjected to the above procedure to give this product (86%).

10

Preparation 18

15 (-)-(trans)-2-(2,3-Dihydrobenzofuran-4-yl)cyclopropane-carboxaldehyde

20 Step 1: (-)-(trans)-N-[3-(2,3-Dihydrobenzofuran-4-yl)-propenoyl]-2,10-camphorsultam

To a solution of (-)-2,10-camphorsultam (8.15 g, 37.9 mmol) in 50 mL toluene at 0°C was added sodium hydride (1.67 g, 41.7 mmol). After stirring for 0.33 h at 0°C and 0.5 h at 20°C and recooling to 0°C, a solution of 3-(2,3-dihydrobenzofuran-4-yl)-2-propenoyl chloride (37.9 mmol), prepared in situ from the corresponding acid and thionyl chloride (75 mL), in toluene (50 mL), was added dropwise. After stirring for 18 h at 20°C, the mixture was diluted with ethyl acetate and washed with water, 1N HCl, and 1N NaOH. The organic solution was dried and concentrated in vacuo to give 15.8 g of crude product. Recrystallization from ethanol-methanol (600 mL, 1:1) gave the product (13.5 g, 92%, mp 199.5-200°C).

25

30

**Step 2: (-)-N-[[~~(trans)~~-2-(2,3-Dihydrobenzofuran-4-yl)-cyclopropyl]-
carbonyl]-2,10-camphorsultam**

1-Methyl-3-nitro-1-nitrosoguanidine (23.88g 163 mmol) was added in
5 portions to a mixture of 10 N sodium hydroxide (60 mL) and ether (200
mL) at 0°C. The mixture was shaken vigorously for 0.25 h and the ether
layer carefully decanted into a solution of (-)-N-[3-(2,3-
dihydrobenzofuran-4-yl)-2-propenoyl]-2,10-camphorsultam (9.67 g, 25
mmol) and palladium acetate (35 mg) in methylene chloride (200 mL).
10 After stirring for 18 h, acetic acid (5 mL) was added to the reaction and
the mixture stirred for 0.5 h. The mixture was washed with 1N HCl, 1N
NaOH and brine. The solution was dried, concentrated in vacuo and the
residue crystallized twice from ethanol to give the product (6.67 g,
66.5%, mp 157-159°C).

15

**Step 3: (-)-(trans)-2-(2,3-Dihydrobenzofuran-4-yl)cyclopropane-
methanol**

A solution of (-)-N-[(trans)-2-(2,3-dihydrobenzofuran-4-yl)cyclo-
20 propanecarbonyl]-2,10-camphorsultam (4.3 g, 10.7 mmol) in THF (50
mL) was added dropwise to a mixture of LAH (0.81 g, 21.4 mmol) in THF
(50 mL) at -45°C. The mixture was stirred for 2 hr while it warmed to
10°C. The mixture was recooled to -40°C and hydrolyzed by the
addition of saturated KHSO₄ (20 mL). The mixture was stirred at room
25 temperature for 30 minutes and filtered. The precipitate was washed
twice with acetone. The combined filtrate and acetone washes were
concentrated in vacuo. The gummy residue was dissolved in ether,
washed with 1N NaOH and 1N HCl, and then dried in vacuo to give the
product (2.0 g, 98.4%).

30

**Step 4: (-)-(trans)-2-(2,3-Dihydrobenzofuran-4-yl)cyclopropane-
carboxaldehyde**

DMSO (1.6 g, 21 mmol) was added to oxalyl chloride in CH₂Cl₂ (7.4 mL of 2 M solution, 14.8 mmole) at -78°C. The (-)-(trans)-2-(2,3-dihydrobenzofuran-4-yl)-cyclopropylmethanol (2.0 g, 10.5 mmol) in CH₂Cl₂ (15 mL) was added. The mixture was stirred for 20 min and then triethylamine (4.24 g, 42 mmol) was added. The mixture was warmed to room temperature and stirred for 30 min. The mixture was diluted with CH₂Cl₂ and washed with water, 1N HCl, and then 1N NaOH. The organic layer was dried and concentrated in vacuo to give the aldehyde product (1.98 g, 100%).

10

Preparation 19

(-)-(trans)-2-(2,3-Dihydro-1,4-benzodioxan-5-yl)cyclopropanecarboxaldehyde

15

Step 1: (-)-(trans)-N-[3-(2,3-Dihydro-1,4-benzodioxan-5-yl)propenoyl]-2,10-camphorsultam

(trans)-3-(2,3-Dihydro-1,4-benzodioxan-5-yl)propenoic acid was subjected to the above procedure to give (88%, mp 187-188°C).

20

Step 2: (-)-N-(trans)-[2-(2,3-Dihydro-1,4-benzodioxan-5-yl)cycloprop-1-yl]carbonyl]-2,10-camphorsultam

25

(-)-(trans)-N-[3-(2,3-Dihydro-1,4-benzodioxan-5-yl)propenoyl]-2,10-camphorsultam was subjected to the above procedure to give (84%, mp 214-215°C, $[\alpha]_D^{25} = -138.9$).

30

Step 3: (-)-(trans)-2-(2,3-Dihydro-1,4-benzodioxan-5-yl)-1-cyclopropanemethanol

(-)-N-(trans)-[[3-(2,3-Dihydro-1,4-benzodioxan-5-yl)cyclopropyl]carbonyl]-2,10-camphorsultam was reduced with LAH as described above to give the product as an oil (100%)

5 **Step 4: (-)-(trans)-2-(2,3-Dihydro-1,4-benzodioxan-5-yl)-1-cyclopropanecarboxaldehyde**

(-)-(trans)-2-(2,3-Dihydro-1,4-benzodioxan-5-yl)-1-cyclopropanemethanol was oxidized under as described above to give the aldehyde
10 as an oil (100%) which was immediately used in the next reaction.

Preparation 20

15 **(+)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)cyclopropane-carboxaldehyde and (-)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)cyclopropane-carboxaldehyde**

Step 1: (trans)-N-[3-(2-Methyl-2,3-dihydrobenzofuran-4-yl)propenoyl]-2,10-camphorsultam

20

(trans)-N-[3-(2-Methyl-2,3-dihydrobenzofuran-4-yl)propenoic acid was subjected to the above procedure to give the desired sultam (95% yield for two steps).

25 **Step 2: N-(trans)-[[2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]carbonyl]-2,10-camphorsultam**

(trans)-N-[3-(2-Methyl-2,3-dihydrobenzofuran-4-yl)propenoyl]-2,10-camphorsultam was subjected to the above procedure to give the
30 desired cyclopropane derivative (61%).

Step 3: (trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)-1-cyclopropanemethanol

N-(trans)-[[2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]carbonyl]-2,10-camphorsultam was reduced with LAH as described above to give the desired alcohol as a mixture of diastereomers (96%).

- 5 The mixture of diastereomeric alcohols was separated by chiral HPLC (Chiracel OD, iPrOH/hexanes) to give both (+)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)-1-cyclopropanemethanol and (-)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)-1-cyclopropanemethanol.

- 10 **Step 4: (+)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)-1-cyclopropanecarboxaldehyde and (-)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)-1-cyclopropanecarboxaldehyde**

- 15 The separated (+)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)-1-cyclopropanemethanol and the separated (-)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)-1-cyclopropanemethanol were separately oxidized as described above to give the corresponding desired aldehydes (98% and 97% yield, respectively).

20

Preparation of intermediates of Formula 5:

Preparation 21

25

2,2-Difluoro-3-(2-methyl-4-benzofuranyl)cyclopropane-methanol, O-acetate

30

Step 1. (trans)-3-(2-Methyl-4-benzofuranyl)-2-propen-1-ol

A solution of (trans)-3-(2-methyl-4-benzofuranyl)-2-propenoic acid (2.53 g, 10.9 mmol) in 30 mL anhydrous THF was added slowly to a mixture of sodium borohydride (592 mg, 15.6 mmol) in 15 mL of anhydrous THF at RT. Once the evolution of hydrogen had ceased, the reaction mixture

was cooled to 0°C and a solution of I₂ (1.72 g, 6.76 mmol) in 15 mL THF was added. The reaction mixture was stirred at 0°C for 1h, allowed to warm to RT and stirred at RT for an additional 3h. The reaction was then quenched with 3N HCl and the aqueous layer extracted with ether. The combined organic layers were washed with 3N NaOH and brine, then dried over MgSO₄ and concentrated in vacuo. Purification on silica gel (ethyl acetate/hexanes) afforded a mixture of the allylic alcohol and the saturated alcohol (10:1) in a combined yield of 46%. The mixture was carried through the next step.

10

Step 2. (trans)-3-(2-Methyl-4-benzofuranyl)-2-propen-1-ol, O-acetate

Acetic anhydride (1.20 mL, 12.6 mmol) was added to a solution of the mixture containing (trans)-3-(2-methyl-4-benzofuranyl)-2-propen-1-ol (as obtained in the previous step) in 10 mL pyridine at -5°C. The reaction mixture was stirred at RT for 20 h then diluted with ether. The crude mixture was washed with 1 N HCl, saturated NaHCO₃, water, and brine then dried over MgSO₄, filtered and dried in vacuo. Purification on silica gel (ethyl acetate/hexanes) afforded a mixture of the allylic acetate and the saturated acetate in a combined yield of 73%.

20

Step 3. 2,2-Difluoro-3-(2-methyl-4-benzofuranyl)-cyclopropanemethanol, O-acetate

To a refluxing solution of (trans)-3-(2-methyl-4-benzofuranyl)-2-propen-1-ol, O-acetate (0.1 g, 4.35 mmol) in 10 ml of diglyme was slowly added a solution of ClF₂CCOONa (5.0 g, 32.6 mmo) in 10 ml of diglyme. The reaction solution was stirring at 165°C for 1.5 h, cooled to room temperature and filtered through Celite. The filtrate was diluted with ether and washed with large amounts of water. The aqueous layer was extracted with ether. The combined organic layers were washed with water and brine solution and dried over MgSO₄. Solvent was removed in vacuo. Purification on silica gel (ethyl acetate/hexanes) afforded 2,2-

30

5 difluoro-3-(2-methyl-4-benzofuranyl)cyclopropanemethanol, O-acetate (560 mg, 56%) as a colorless liquid. ¹H NMR (300 MHz, CDCl₃) δ 2.12 (s, 3H), 2.25-2.36 (m, 1H), 2.46 (s, 3H), 2.76-2.88 (m, 1H), 4.27-4.45 (m, 2H), 6.47 (s, 1H), 7.00 (d, 1H, J = 7.5 Hz), 7.15 (t, 1H, J = 7.5 Hz), 7.33 (d, 1H, J = 7.5 Hz). ¹³C NMR (75 MHz, CDCl₃) δ 14.4, 18.4, 27.8 (t, J = 10.4 Hz), 29.8 (t, J = 10.4 Hz), 61.2, 101.2, 110.4, 113.5 (t, J = 286.7 Hz), 121.7, 123.3, 124.3, 129.7, 154.8, 156.2, 171.1.

10 Preparation of intermediates of Formula Z:

Preparation 22

15 (±)-(trans)-2-(2,3-dihydrobenzofuran-4-yl)cyclopropane-methanamine

A solution of (±)-(trans)- 2-(2,3-dihydrobenzofuran-4-yl)-cyclopropane-carboxaldehyde (2.6 g, 15 mmol), hydroxylamine hydrochloride (3.13 g, 45 mmol), ethanol (60 mL), water (40 mL), and 10 N NaOH (4.5 mL, 45 mmol) was heated to reflux for 4 h. The solution was cooled to room temperature, diluted with water and ethyl acetate. The ethyl acetate layer was separated and washed sequentially with H₂O and brine. The ethyl acetate extract was dried over K₂CO₃ and concentrated in vacuo. The residue was dissolved in THF (50 mL) and added dropwise to a - 45°C suspension of LAH (1.06 g, 28 mmol) in THF (100 mL), maintaining the temperature below -40°C. The reaction was warmed to room temperature and stirred for 2 h. The reaction was recooled to -45°C, and 1N HCl (50 mL) was cautiously added dropwise. The cooling bath was removed and the suspension was stirred at room temperature for 30 min. The resulting paste was then diluted with ethyl ether and extracted with 1N HCl. The acid extracts were combined, washed with ethyl ether, made basic with 50% NaOH, and extracted with dichloromethane. The

dichloromethane extracts were combined, washed with brine, dried over K_2CO_3 , and concentrated in vacuo to give a clear oil (900 mg, 40%).

Preparation 23

5

The following compounds of Formula Z, (a) - (m) were prepared by the general procedure described for the compound in Preparation 22.

(a) (±)-(trans)-2-(Benzofuran-4-yl)-cyclopropanemethanamine

10

(±)-(trans)-2-(Benzofuran-4-yl)cyclopropanecarboxaldehyde was subjected to the above procedure to give an oil (73.5%).

(b) (±)-(trans)-2-(1,3-Benzodioxol-4-yl)cyclopropanemethanamine

15

(±)-(trans)-2-(1,3-Benzodioxol-4-yl)cyclopropanecarboxaldehyde was reacted by the above procedure to give a red oil (61%).

(c) (±)-(trans)-2-(2,3-Dihydrobenzodioxan-5-yl)cyclopropane-methanamine

20

(±)-(trans)-2-(2,3-Dihydrobenzodioxan-5-yl)cyclopropane-carboxaldehyde was reacted by the above procedure to give an orange oil (91.1%).

25

(d) (±)-(trans)-2-(2,3,4,5-tetrahydrobenzoxepin-6-yl)cyclopropane-methanamine

30

(±)-(trans)-2-(2,3,4,5-tetrahydrobenzoxepin-6-yl)cyclopropane-carboxaldehyde was reacted by the above procedures to give this product (50%).

(e) (±)-(trans)-2-(Benzofuran-7-yl)cyclopropanemethanamine

(±)-(trans)-2-(Benzofuran-7-yl)cyclopropanecarboxaldehyde was reacted by the above procedures to give this product (66%).

5

(f) (±)-(trans)-2-(2,3-Dihydro-benzofuran-7-yl)cyclopropane-methanamine

(±)-(trans)-2-(2,3-Dihydro-benzofuran-7-yl)cyclopropanecarboxaldehyde was reacted by the above procedures to give this product (87%).

10

(g) (±)-(trans)-2-(2,2-Dimethyl-2,3-dihydro-benzofuran-4-yl)cyclopropanemethanamine

(±)-(trans)-2-(2,2-Dimethyl-2,3-dihydro-benzofuran-4-yl)cyclopropanecarboxaldehyde was reacted by the above procedures to give this product (60%).

15

(h) (±)-(trans)-2-(2-Methyl-benzofuran-4-yl)cyclopropanemethanamine

20

(±)-(trans)-2-(2-Methyl-benzofuran-4-yl)cyclopropanecarboxaldehyde was reacted by the above procedures to give this product (63%).

(i) (trans)-2-(2,3-Dihydro-1,4-benzodioxan-5-yl)cyclopropane-methanamine

25

(trans)-2-(2,3-dihydro-1,4-benzodioxan-5-yl)cyclopropanecarboxaldehyde was reacted by the above procedures to give the amine (67%) which was converted to the fumarate (mp 183-184°C).

30

(j) (trans)-2-(2H-3,4-Dihydro-1,5-benzodioxapin-6-yl)cyclopropane-methanamine

trans-(trans)-2-(2H-3,4-Dihydro-1,5-benzodioxapin-6-yl)cyclopropane-carboxaldehyde was reacted by the above procedures to give the amine which was converted to the fumarate (65%, mp 152-153°C).

Anal. Calc'd for 0.3 H₂O: C, 59.92; H, 6.39; N 4.11.

5 Found: C, 50.78; H, 6.33; N, 4.01.

(k) (±)-(trans)-2-(2H-2,3-Dihydrobenzopyran-4-yl)cyclopropane-methanamine

10 (±)-(trans)-2-(2H-2,3-Dihydrobenzopyran-4-yl)cyclopropane-carboxaldehyde was reacted by the above procedure to give this product (42%).

15 (l) (-)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)cyclopropane-methanamine

(-)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)cyclopropane-carboxaldehyde was subjected to the above procedure to give the desired amine (71%).

20

(m) (+)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)cyclopropane-methanamine

25 (+)-(trans)-2-(2-Methyl-2,3-dihydrobenzofuran-4-yl)cyclopropane-carboxaldehyde was subjected to the above procedure to give the desired amine (59%).

Preparation 24

30

(-)-(trans)-2-(2,3-Dihydrobenzofuran-4-yl)cyclopropane-methanamine

A mixture of (-)-(trans)-2-(2,3-dihydrobenzofuran-4-yl)cyclopropane-carboxaldehyde (1.98 g, 10.5 mmol), hydroxylamine hydrochloride (2.29 g, 33 mmol), and 30% NaOH (3.5 mL, 35 mmol), in 5:1 ethanol/water (50 mL) was heated on a steam bath for 2 h. The solution
5 was concentrated in vacuo, and the residue mixed with water. The mixture was extracted with CH₂Cl₂. The organic extracts were dried and concentrated in vacuo to give a solid which NMR analysis showed to be a mixture of the cis and trans oximes. This material was dissolved in THF (20 mL) and added to solution of alane in THF [prepared from LAH
10 (1.14 g, 30 mmol) and H₂SO₄ (1.47 g, 15 mmol) at 0°C]. The reaction was stirred for 18 h, and quenched successively with water (1.15 mL), 15% NaOH (1.15 mL), and then water (3.45 mL). The mixture was filtered and the filtrate was concentrated in vacuo. The residue was mixed with ether and washed with water and then 1N HCl. The acid
15 washes were made basic and extracted with CH₂Cl₂. The extracts were dried and concentrated in vacuo to give the amine product (1.4 g, 70.5%). The amine was converted to the fumarate salt in ethanol (mp: 197-198°C).

Anal. Calc'd for C₁₂H₁₅NO • C₄H₄O₄: C, 62.94; H, 6.27; N, 4.59.

20

Found: C, 62.87; H, 6.31; N, 4.52.

Preparation 25

25 (-)-(trans)-2-(1,3-Benzodioxol-4-yl)cyclopropane-methanamine

(-)-(trans)-2-(1,3-Benzodioxol-4-yl)cyclopropanecarboxaldehyde was subjected to the above general procedure described in Preparation 24 to give a beige solid (52.6%).

30

Preparation 26

(+)-trans-2-[(2,3-Dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methanamine

5

Step 1: (-)-(trans)-2-(2,3-Dihydro-1,4-benzodioxin-5-yl)-1-cyclopropanecarboxaldehyde oxime

10 (-)-(trans)-2-(2,3-Dihydro-1,4-benzodioxin-5-yl)-1-cyclopropanecarboxaldehyde was reacted with hydroxyl amine as described above to give the oxime as a mixture of isomers.

Step 2: (+)-trans-2-[(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methanamine

15

The above oxime was reduced with alane as described above to give the amine which was converted to the fumarate (80%, mp 173-174°C for fumarate, $[\alpha]_D^{25} = 6.15$).

Anal. Calc'd for 0.5 H₂O: C, 58.36; H, 6.10; N 4.24.

20 Found: C, 58.36; H, 6.09; N, 4.24.

Preparation 27

2,2-Difluoro-3-(2-methyl-4-benzofuranyl)cyclopropanemethanamine

25

Step 1. 2,2-Difluoro-3-(2-methyl-4-benzofuranyl)cyclopropanemethanol

30 To a stirred solution of 2,2-difluoro-3-(2-methyl-4-benzofuranyl)-cyclopropanemethanol, O-acetate (560 mg, 2.0 mmol) in MeOH/THF (10 ml, 3/1) was added powdered KOH (560 mg, 10.0 mmol). The resulting solution was stirred at RT for 2 h. Solvent was removed in vacuo. The

residue was diluted with ether and water. The aqueous layer was extracted with ether and the combined organic layers were washed with saturated NaHCO_3 , water and brine solution. Solvent was removed in vacuo. Purification on silica gel (ethyl acetate/hexanes) afforded 2,2-difluoro-3-(2-methyl-4-benzofuranyl)cyclopropanemethanol (420 mg, 88%) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 1.67 (bs, 1H), 2.19-2.31 (m, 1H), 2.46 (s, 3H), 2.73-2.80 (m, 1H), 3.95-3.98 (m, 2H), 6.51 (s, 1H), 7.02 (d, 1H, $J = 7.5$ Hz), 7.15 (t, 1H, $J = 7.5$ Hz), 7.32 (d, 1H, $J = 7.5$ Hz). ^{13}C NMR (75 MHz, CDCl_3) δ 14.4, 29.3 (t, $J = 10.4$ Hz), 31.3 (t, $J = 10.4$ Hz), 60.1, 101.3, 110.2, 114.1 (t, $J = 286.7$ Hz), 121.7, 123.3, 124.7, 129.8, 154.7, 156.1.

Step 2. 4-[2-(Azidomethyl)-3,3-difluorocyclopranyl]-2-methylbenzofuran

Triethylamine (916 ml, 6.55 mmol) followed by $\text{CH}_3\text{SO}_2\text{Cl}$ (355 ml, 4.59 mmol) were added to a solution of 2,2-difluoro-3-(2-methyl-4-benzofuranyl)cyclopropanemethanol (390 mg, 1.64 mmol) in 15 ml of CH_2Cl_2 . The reaction was stirred at RT for 0.5 h and diluted with CH_2Cl_2 . The organic layer was washed with water and NaHCO_3 and dried over anhydrous K_2CO_3 . Removal of solvent in vacuo afforded crude mesylate, which was used immediately in the next step. A solution of the mesylate and NaN_3 (213 mg, 3.27 mmol) in 12 ml of CH_2Cl_2 was heated to 70°C for 2.5 h. The resulting solution was cooled to RT and diluted with EtOAc and water. The aqueous layer was extracted with EtOAc. The combined organic layers were washed with water and dried over MgSO_4 . Removal of solvent in vacuo afforded 4-[2-(azidomethyl)-3,3-difluorocyclopranyl]-2-methylbenzofuran (422 mg, 98%). ^1H NMR (300 MHz, CDCl_3) δ 2.16-2.28 (m, 1H), 2.47 (s, 3H), 2.74-2.81 (m, 1H), 3.52-3.70 (m, 2H), 6.49 (s, 1H), 7.02 (d, 1H, $J = 7.5$ Hz), 7.16 (t, 1H, $J = 7.5$ Hz), 7.34 (d, 1H, $J = 7.5$ Hz).

Step 3. 2,2-Difluoro-3-(2-methyl-4-benzofuranyl)cyclopropane-methanamine

A solution of 4-[2-(azidomethyl)-3,3-difluorocyclopranyl]-2-methylbenzofuran (203 mg, 0.78 mmol) in 4 ml of THF was added dropwise to a solution of LAH (1.0 M solution in THF, 1.56 ml, 1.56 mmol) at -30°C. The resulting solution was allowed to warm up to RT and stirred at RT for 3 h. A solution of KHSO₄ (130 mg, 0.96 mmol) in 1 ml of water was added to the reaction at -30°C. After stirring at RT for 20 min, the reaction solution was filtered and the filtrate was diluted with CH₂Cl₂. The solution was adjusted to pH = 10 by addition of NH₄OH. The aqueous layer was extracted with CH₂Cl₂. The combined organic layers were washed with brine and dried over K₂CO₃. Solvent was removed in vacuo to afford 2,2-difluoro-3-(2-methyl-4-benzofuranyl)cyclopropane-methanamine (182 mg, 99%) as colorless liquid.

EXAMPLES

The following examples illustrate the preparation of the compounds of the invention by following the general procedures described above.

Example 1

(±)-(trans)-N-[[2-(2,3-Dihydrofuran-4-yl)cycloprop-1-yl]methyl]acetamide

Acetic anhydride (0.14 mL, 1.8 mmol) was added dropwise to a stirred solution of (±)-(trans)-2-(2,3-dihydrobenzofuran-4-yl)-cyclopropanemethanamine (300 mg, 1.6 mmol) and triethylamine (0.67 mL, 4.8 mmol) in dry dichloromethane (15 mL) at 0°C. The resulting suspension was warmed to room temperature and stirred for

18 h. The mixture was concentrated in vacuo and the residue was purified by flash chromatography (silica gel, CH₂Cl₂ then 2% EtOAc/CH₂Cl₂) to afford 200 mg (54%) of a clear oil.

IR (NaCl Film): 3287, 2923, 1651, 1553, 1459 cm⁻¹.

5 Anal. Calc'd for C₁₄H₁₇NO₂•0.3 H₂O: C, 71.04; H, 7.50; N, 5.92.

Found: C, 70.79; H, 7.41; N, 5.58.

Example 2

10 (-)-(trans)-N-[[2-(2,3-Dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide

This compound was prepared similar to the above procedure using propionyl chloride and (-)-(trans)-2-(2,3-dihydrobenzofuran-4-yl)-
15 cyclopropanemethanamine to give an oil that solidified upon standing to an off-white solid (61%, mp: 71-72°C).

IR (NaCl Film): 3298, 1645, 1548, 1459, 1235 cm⁻¹.


[α]_D²⁵: -17.3°

Anal. Calc'd for C₁₅H₁₉NO₂: C, 73.44; H, 7.87; N, 5.71.

20 Found: C, 73.28; H, 7.68; N, 5.58.

The following compounds of Formula I were prepared by these general methods:

Table 2: Physical Data for Compounds of Formula I.



Ex.	X-Y-Z	R ¹	[α] _D ²⁵	Elemental Analyses			Empirical Formula	mp °C	desc.	% yield
				Calculated over Found						
3	O-CH ₂ -O	Me	±	65.91 66.11	6.55 6.28	5.91 5.92	C ₁₃ H ₁₅ NO ₃ •0.2 H ₂ O	—	oil	61
4	O-CH ₂ -O	Me	-31.5°	66.42 66.45	6.52 6.70	5.96 5.86	C ₁₃ H ₁₅ NO ₃ •0.1 H ₂ O	56-58	white solid	53
5	O-CH ₂ -O	Et	±	68.00 67.78	6.93 6.69	5.66 5.56	C ₁₄ H ₁₇ NO ₃	84-86	beige solid	41
6	O-CH ₂ -O	Et	-41.4°	68.00 68.19	6.93 6.76	5.66 5.58	C ₁₄ H ₁₇ NO ₃	101-103	white solid	56
7	O-CH ₂ -O	n-Pr	±	68.94 68.83	7.33 7.23	5.36 5.24	C ₁₅ H ₁₉ NO ₃	75-77	waxy solid	49
8	O-CH ₂ -O	n-Pr	-32.6°	68.94 68.66	7.33 7.32	5.36 5.52	C ₁₅ H ₁₉ NO ₃	82-84	white solid	58
9	O-CH ₂ -O	c-Pr	±	69.48 69.22	6.61 6.78	5.40 5.31	C ₁₅ H ₁₇ NO ₃	127-129	beige solid	38
10	O-CH ₂ -O	c-Pr	-32.6°	69.00 68.69	6.64 6.56	5.36 5.42	C ₁₅ H ₁₇ NO ₃ •0.1 H ₂ O	148-150	white solid	77

Ex.	X-Y-Z	R ¹	[α] _D ²⁵	Elemental Analyses Calculated over Found			Empirical Formula	mp °C	desc.	% yield
				%C	%H	%N				
11	O-CH ₂ -O	i-Pr	±	68.94 68.75	7.33 7.33	5.36 5.51	C ₁₅ H ₁₉ NO ₃	114-115	solid	50
12	O-CH ₂ -O	i-Pr	—	68.94 68.79	7.33 7.35	5.36 5.28	C ₁₅ H ₁₉ NO ₃	127-129	beige solid	54
13	O-CH ₂ -O	NHEt	±	64.10 63.90	6.92 6.93	10.68 10.63	C ₁₄ H ₁₈ N ₂ O ₃	121-122	white solid	65
14	O-CH ₂ -O	NHEt	-26.9°	64.11 63.82	6.92 6.88	10.68 10.59	C ₁₄ H ₁₈ N ₂ O ₃	137-139	white solid	85
15	O-(CH ₂) ₂ -O	Me	±	65.60 65.75	6.66 6.45	5.35 5.27	C ₁₄ H ₁₇ NO ₃ •0.15 CH ₂ Cl ₂	—	amber oil	59
16	O-(CH ₂) ₂ -O	Et	±	67.36 67.47	7.14 7.21	5.17 5.18	C ₁₅ H ₁₉ NO ₃ •0.1 CH ₂ Cl ₂	67-69	beige solid	44
17	O-(CH ₂) ₂ -O	n-Pr	±	68.26 68.21	7.50 7.31	4.91 4.81	C ₁₆ H ₂₁ NO ₃ •0.1 CH ₂ Cl ₂	—	amber oil	49
18	O-(CH ₂) ₂ -O	c-Pr	±	70.31 69.89	7.01 6.74	5.12 5.10	C ₁₆ H ₁₉ NO ₃	96-98	white solid	68
19	O-(CH ₂) ₂ -O	i-Pr	±	69.01 68.88	7.59 7.71	5.00 4.95	C ₁₆ H ₂₁ NO ₃ •0.05 CH ₂ Cl ₂	100-102	beige solid	49
20	O-(CH ₂) ₂ -O	NHMe	±	63.23 63.42	6.98 7.31	10.54 10.59	C ₁₄ H ₁₈ N ₂ O ₃ •0.2 H ₂ O	105-107	white solid	77
21	O-(CH ₂) ₂ -O	NHEt	±	65.20 64.92	7.29 7.41	10.14 10.01	C ₁₅ H ₂₀ N ₂ O ₃	92-94	beige solid	68
22	CH=CH-O	Me	±	71.65 71.68	6.70 6.69	5.97 5.67	C ₁₄ H ₁₅ NO ₁ •0.3 H ₂ O	88-89	white solid	31

Elemental Analyses											
Calculated over Found											
Ex.	X-Y-Z	R ¹	[α] _D ²⁵	%C	%H	%N	Empirical Formula	mp °C	desc.	% yield	
23	CH=CHO	Et	±	74.04 73.80	7.04 6.96	5.76 5.70	C ₁₅ H ₁₇ NO ₂	84-85	white solid	57	
24	CH=CHO	n-Pr	±	74.68 74.42	7.44 7.40	5.44 5.32	C ₁₆ H ₁₉ NO ₂	67-68	white solid	52	
25	CH=CHO	c-Pr	±	73.71 73.45	6.81 6.81	5.37 5.12	C ₁₆ H ₁₇ NO ₂ •0.3 H ₂ O	96-98	beige solid	63	
26	CH=CHO	i-Pr	±	73.39 73.22	7.51 7.62	5.35 5.24	C ₁₆ H ₁₉ NO ₂ •0.25 H ₂ O	—	amber oil	74	
27	CH=CHO	NHMe	±	67.15 66.97	6.43 6.58	11.03 10.74	C ₁₄ H ₁₆ N ₂ O ₂ •0.1 CH ₂ Cl ₂	97-98	white solid	58	
28	CH ₂ -CH ₂ -O	Me	±	71.04 70.79	7.50 7.41	5.92 5.58	C ₁₄ H ₁₇ NO ₂ •0.3 H ₂ O	—	oil	54	
29	CH ₂ -CH ₂ -O	Et	±	72.11 72.09	7.87 7.96	5.61 5.44	C ₁₅ H ₁₉ NO ₂ •0.25 H ₂ O	—	oil	64	
30	CH ₂ -CH ₂ -O	n-Pr	±	73.59 73.45	8.18 8.40	5.36 5.19	C ₁₆ H ₂₁ NO ₂ •0.1 H ₂ O	—	oil	75	
31	CH=CHO	cPr	-35.17	74.37 74.52	6.63 6.54	5.38 5.40	C ₁₆ H ₁₇ NO ₂ •0.05 C ₄ H ₈ O	112-114	beige solid	54.7	
32	CH=CHO	Et	-35.51	73.08 73.31	6.96 6.52	5.66 5.67	C ₁₅ H ₁₇ NO ₂ •0.05 CH ₂ Cl ₂	46-48	tacky beige solid	62.8	
33	CH ₂ CH ₂ O	Et	18.41	73.44 73.05	7.81 7.84	5.71 5.62	C ₁₅ H ₁₉ NO ₂	73.5-74	white solid	37.5	

Ex.	X-Y-Z	R ¹	[α] _D ²⁵	Elemental Analyses			Empirical Formula	mp °C	desc.	% yield
				Calculated over Found						
				%C	%H	%N				
34	CH ₂ CH ₂ O	Me	—	72.70 72.57	7.41 7.58	6.06 6.00	C ₁₄ H ₁₇ NO ₂	80-82	off-white solid	65.8
35	CH ₂ CH ₂ O	nPr	—	71.61 72.01	8.26 8.39	5.22 4.77	C ₁₆ H ₂₁ NO ₂ •0.5 H ₂ O	—	amber oil	95.0
36	CH ₂ CH ₂ O	NHMe	—	67.77 67.54	7.39 7.55	11.29 11.41	C ₁₄ H ₁₈ N ₂ O ₂ •0.10 H ₂ O	113-116	off-white solid	53.8
37	CH ₂ CH ₂ O	CH ₂ OMe	—	67.78 67.53	7.39 7.55	5.27 5.11	C ₁₅ H ₁₉ NO ₃ •0.25 H ₂ O	—	amber oil	94.2
38	CH ₂ CH ₂ O	cPr	-8.38	74.16 73.93	7.47 7.39	5.41 5.38	C ₁₆ H ₁₉ NO ₂ •0.10 H ₂ O	111-115	off-white solid	62.5
39	CH ₂ CH ₂ O	CF ₃	—	58.95 58.83	4.95 4.94	4.91 4.76	C ₁₄ H ₁₄ F ₃ NO ₂	90-91	white solid	43.9
40	CH ₂ -(CH ₂) ₂ -O	Et	±	74.10 73.84	8.16 8.09	5.40 5.22	C ₁₆ H ₂₁ NO ₂	56-57	white solid	29
41	CH ₂ -(CH ₂) ₂ -O	cPr	±	75.25 74.98	7.80 7.75	5.16 5.01	C ₁₇ H ₂₁ NO ₂	132-134	white solid	40
42	CH ₂ -(CH ₂) ₂ -O	Et	-24.47	73.59 73.59	8.18 8.18	5.36 5.36	C ₁₆ H ₂₁ NO ₂ •0.10 H ₂ O	83-84	white solid	32
43	CH ₂ -(CH ₂) ₂ -O	Me	-39.15	73.06 72.83	7.82 7.88	5.68 5.57	C ₁₅ H ₁₉ NO ₂ •0.07 H ₂ O	131-132	white solid	13
44	CH ₂ -(CH ₂) ₂ -O	cPr	-35.28	74.75 74.42	7.82 7.68	5.13 5.04	C ₁₇ H ₂₁ NO ₂ •0.10 H ₂ O	157-158	white solid	14
45	O-(CH ₂) ₂ -O	Et	-9.80	68.94 68.82	7.33 7.46	5.36 5.30	C ₁₅ H ₁₉ NO ₃	90-91	white solid	78.9

Ex.	X-Y-Z	R ¹	[α] _D ²⁵	Elemental Analyses			Empirical Formula	mp °C	desc.	% yield
				Calculated over Found						
				%C	%H	%N				
46	O-(CH ₂) ₂ -O	Me	—	68.00 67.86	6.93 6.96	5.66 5.66	C ₁₄ H ₁₇ NO ₃	79-81	beige solid	82.9
47	O-(CH ₂) ₂ -O	NHMe	3.71	63.24 63.37	6.97 7.29	10.53 10.41	C ₁₄ H ₁₈ N ₂ O ₃ •0.2 H ₂ O	101-103	off-white solid	83.8
48	O-(CH ₂) ₂ -O	CH ₂ OMe	25.19	64.97 64.93	6.91 6.88	5.05 5.01	C ₁₅ H ₁₉ NO ₄	96-97	off-white solid	64.4
49	O-(CH ₂) ₂ -O	cPr	-10.51	70.31 70.29	7.01 7.02	5.12 5.08	C ₁₆ H ₁₉ NO ₃	122-123	white solid	75.4
50	O-(CH ₂) ₂ -O	nPr	-20.24	69.79 69.99	7.69 7.74	5.09 5.02	C ₁₆ H ₂₁ NO ₃	58-59	off-white solid	32.3
51	O-(CH ₂) ₂ -O	NHEt	0.0	65.20 64.79	7.30 7.49	10.14 9.66	C ₁₅ H ₂₀ N ₂ O ₃	132-133	off-white solid	70.0
52	O-(CH ₂) ₂ -O	vinyl	42.27	68.60 68.32	6.94 6.81	5.13 4.74	C ₁₅ H ₁₇ NO ₃ •0.30 C ₂ H ₆ O	—	amber oil	84.7
53	O-(CH ₂) ₂ -O	iPr	-13.23	68.89 68.91	7.73 7.72	5.02 5.00	C ₁₆ H ₂₀ NO ₃ •0.2 H ₂ O	123-124	white solid	54.7
54	O-(CH ₂) ₂ -O	CF ₃	—	55.82 55.79	4.68 4.68	4.65 4.65	C ₁₄ H ₁₄ F ₃ NO ₃	119-121	beige solid	44.2
55	O-(CH ₂) ₂ -O	CH ₂ CF ₃	-12.22	57.14 57.01	5.12 5.07	4.44 4.43	C ₁₅ H ₁₆ F ₃ NO ₃	136.5-137.5	off-white solid	65.2
56	O-(CH ₂) ₂ -O	iso-propenyl	—	67.63 67.56	7.17 7.11	4.93 4.52	C ₁₆ H ₁₉ NO ₃ •0.6 H ₂ O	144-150	off-white solid	42.7
57	O-(CH ₂) ₂ -O	Et	2.63	68.94 68.58	7.33 7.38	5.36 5.02	C ₁₅ H ₁₉ NO ₃	77-78	white solid	13
58	O-(CH ₂) ₂ -O	nPr	3.07	69.79 69.58	7.69 7.89	5.09 5.12	C ₁₆ H ₂₁ NO ₃	40-41	white solid	14

Elemental Analyses											
Calculated over Found											
Ex.	X-Y-Z	R ¹	[α] _D ²⁵	%C	%H	%N	Empirical Formula	mp °C	desc.	% yield	
59	O-(CH ₂) ₃ -O	Et	±	69.79 69.53	7.69 7.69	5.09 5.00	C ₁₆ H ₂₁ NO ₃	75-77	white solid	93.1	
60	O-(CH ₂) ₃ -O	Me	±	68.94 68.67	7.33 7.12	5.36 5.27	C ₁₅ H ₁₉ NO ₃	73-75	solid	80.9	
61	CH ₂ -(CH ₂) ₃ -O	Et	±	74.69 74.74	8.48 8.75	5.12 5.02	C ₁₇ H ₂₃ NO ₂	102.5-105	white solid	65.0	
62	CH ₂ -(CH ₂) ₃ -O	nPr	±	75.22 75.36	8.77 9.06	4.87 4.77	C ₁₈ H ₂₅ NO ₂	95-97	white solid	84.5	
63	CH ₂ -(CH ₂) ₃ -O	Me	±	74.10 73.82	8.16 8.08	5.40 5.27	C ₁₆ H ₂₁ NO ₂	70-72.5	white solid	61.0	
64	O-CH ₂ -CH ₂	Me	±	71.65 71.62	7.32 7.54	5.95 5.93	C ₁₄ H ₁₇ NO ₂ ·0.05 CH ₂ Cl ₂	53-55	white solid	53	
65	O-CH ₂ -CH ₂	Et	±	73.44 73.20	7.81 7.49	5.71 5.63	C ₁₅ H ₁₉ NO ₂	90-92	white solid	72	
66	O-CH ₂ -CH ₂	nPr	±	74.10 73.91	8.16 8.14	5.40 5.40	C ₁₆ H ₂₁ NO ₂	88-90	white solid	50	
67	O-CH ₂ -CH ₂	cPr	±	74.68 74.42	7.44 7.41	5.44 5.31	C ₁₆ H ₁₉ NO ₂	125-127	white solid	88	
68	O-CH ₂ -CH ₂	NHMe	±	67.53 67.42	7.30 7.51	11.22 11.05	C ₁₄ H ₁₈ N ₂ O ₂ ·0.04 CH ₂ Cl ₂	108-110	yellow solid	68	
69	CH ₂ -CMe ₂ O	Me	±	73.52 73.45	8.11 8.09	5.35 5.34	C ₁₆ H ₂₁ NO ₂ ·0.2 CH ₂ Cl ₂	—	clear oil	68	
70	CH ₂ -CMe ₂ -O	Et	±	74.69 74.41	8.48 8.34	5.12 4.99	C ₁₇ H ₂₃ NO ₂	63-65	white solid	72	
71	CH ₂ -CMe ₂ O	nPr	±	75.22 74.99	8.77 8.54	4.87 4.72	C ₁₈ H ₂₅ NO ₂	—	clear oil	49	

Ex.	X-Y-Z	R ¹	[α] _D ²⁵	Elemental Analyses			Empirical Formula	mp °C	desc.	% yield
				Calculated over Found						
				%C	%H	%N				
72	CH ₂ -CMe ₂ O	iPr	±	75.22 75.35	8.77 8.66	4.87 4.76	C ₁₈ H ₂₅ NO ₂	110-112	white solid	65
73	CH ₂ -CMe ₂ O	cPr	±	75.76 75.53	8.12 7.81	4.91 4.80	C ₁₈ H ₂₃ NO ₂	99-101	white solid	44
74	CH=CMe-O	Me	±	74.05 73.79	7.04 7.08	5.76 5.50	C ₁₅ H ₁₇ NO ₂	—	clear oil	63
75	CH=CMe-O	Et	±	73.89 73.61	7.38 7.38	5.37 5.09	C ₁₆ H ₁₉ NO ₂ •0.4 CH ₂ Cl ₂	—	yellow oil	72
76	CH=CMe-O	nPr	±	75.25 74.89	7.80 7.58	5.16 5.01	C ₁₇ H ₂₁ NO ₂	—	clear oil	81
77	O-CH=CH	Me	±	72.48 72.32	6.53 6.64	6.02 5.94	C ₁₇ H ₁₅ NO ₂ •0.04 CH ₂ Cl ₂	—	clear oil	58
78	O-CH=CH	Et	±	72.82 72.82	6.95 7.08	5.64 5.42	C ₁₅ H ₁₇ NO ₂ •0.6 CH ₂ Cl ₂	53-55	white powder	49
79	O-CH=CH	nPr	±	74.68 74.31	7.44 7.53	5.44 5.22	C ₁₆ H ₁₉ NO ₂	—	clear oil	51
80	O-CH=CH	cPr	±	75.27 74.99	6.71 6.76	5.49 5.21	C ₁₆ H ₁₇ NO ₂	110-112	white solid	58
81	CH ₂ (CHMe)O	Me	9.5	72.12 72.29	7.87 8.24	5.61 5.37	C ₁₅ H ₁₉ NO ₂ /0.25 H ₂ O	—	oil	70
82	CH ₂ (CHMe)O	Et	15	74.10 73.75	8.16 8.31	5.40 5.24	C ₁₆ H ₂₁ NO ₂	85-86	white solid	70
83	CH ₂ (CHMe)O	nPr	14.3	74.69 74.62	8.48 8.48	5.12 5.04	C ₁₇ H ₂₃ NO ₂	62.5- 63.5	white solid	65
84	CH ₂ (CHMe)O	cPr	13.39	75.25 74.89	7.80 7.81	5.16 4.89	C ₁₇ H ₂₁ NO ₂	126-127	white solid	73

Elemental Analyses											
Calculated over Found											
Ex.	X-Y-Z	R ¹	[α] _D ²⁵	%C	%H	%N	Empirical Formula	mp °C	desc.	% yield	
85	CH ₂ (CHMe)O	NHEt	30.5	70.04 69.80	8.08 8.11	10.21 9.95	C ₁₆ H ₂₂ N ₂ O ₂	118-119	white solid	65	
86	CH ₂ (CHMe)O	iPr	12.7	74.69 74.63	8.48 8.38	5.12 5.01	C ₁₇ H ₂₃ NO ₂	104.5- 105.5	white solid	60	
87	CH ₂ (CHMe)O	CH ₂ Cl	22.7	64.40 64.15	6.49 6.50	5.01 4.82	C ₁₅ H ₁₈ ClNO ₂	73.5- 74.5	white solid	70	
88	CH ₂ (CHMe)O	Et	-48.6	72.10 71.94	8.24 8.19	5.25 5.43	C ₁₆ H ₂₁ NO ₂ /0.4 H ₂ O	98-99	white solid	71	
89	CH ₂ (CHMe)O	iPr	-49.2	73.24 73.22	8.53 8.37	5.02 5.08	C ₁₇ H ₂₃ NO ₂ /0.3 H ₂ O	106-107	white solid	72	
90	CH ₂ (CHMe)O	Me	-55.7	73.44 73.42	7.81 7.90	5.71 5.65	C ₁₅ H ₁₉ NO ₂	117-118	white solid	67	
91	CH ₂ (CHMe)O	CH ₂ OMe	-31	69.79 69.86	7.69 7.82	5.09 5.05	C ₁₆ H ₂₁ NO ₃	75-76	white solid	61	
92	CH ₂ (CHMe)O	cPr	-48.6	75.25 75.13	7.80 7.87	5.16 5.06	C ₁₇ H ₂₁ NO ₂	120-121	white solid	75	
93	CH ₂ (CHMe)O	NHEt	-25.53	70.04 69.83	8.08 8.10	10.21 10.09	C ₁₆ H ₂₂ N ₂ O ₂	115-116	white solid	61	
94	CH ₂ (CHMe)O	nPr	-45.6	74.69 74.46	8.48 8.34	5.12 4.83	C ₁₇ H ₂₃ NO ₂	74-75	white solid	69	

Example 95**N-[[2-(2-Methyl-4-benzofuranyl)-3,3-difluorocyclopropanyl]-methyl]propanamide**

5

Obtained 95% of a white solid. m.p. 55-56°C.

Anal. Calc'd for C₁₆H₁₇F₂NO₂: C, 65.52; H, 5.84; N, 4.78.

Found: C, 66.15; H, 5.95; N, 4.73.

10

Example 96**(+)-(trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]propanamide**

15

Obtained 14% of a white solid. m.p. 81-82°C. $[\alpha]_D^{25} = 16.29$. Purity = 95% by HPLC. ¹H NMR (300 MHz, CDCl₃) δ 6.98 (t, J=7.8 Hz, 1H), 6.65 (d, J=7.9 Hz, 1H), 6.50 (d, J=7.5 Hz, 1H), 5.57 (s, 1H), 4.13 (t, J=4.5 Hz, 2H), 3.49-3.30 (m, 1H), 3.23-3.16 (m, 1H), 2.84-2.78 (m, 2H), 2.21 (q, J=7.6 Hz, 2H), 2.07-1.99 (m, 2H), 1.75-1.69 (m, 1H), 1.26-1.09 (m, 4H), 0.92-0.81 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 173.9, 155.2, 141.1, 126.1, 117.6, 115.1, 66.0, 43.9, 30.0, 22.6, 20.9, 19.6, 12.7, 10.1.

20

Example 97

25

(+)-(trans)-N-[[2(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]butanamide

Obtained 10% of a yellow oil. Purity = 87% by NMR. ¹H NMR (300 MHz, CDCl₃) δ 6.95 (d, J=7.8 Hz, 1H), 6.60 (d, J=7.4 Hz, 1H), 6.50 (d, J=7.9 Hz, 1H), 5.58 (s, 1H), 4.07 (t, J=4.7 Hz, 2H), 3.44-3.35 (m, 1H), 3.25-3.11 (m, 1H), 2.78-2.73 (m, 2H), 2.13-2.08 (t, J=7.3 Hz, 2H), 2.02-1.94 (m, 2H),

30

1.70-1.55 (m, 3H), 1.23-1.09 (m, 1H), 0.91-0.75 (m, 5H); ^{13}C NMR (75 MHz, CDCl_3) δ 155.1, 141.0, 126.8, 122.1, 117.5, 115.1, 65.9, 43.8, 38.9, 22.5, 20.8, 19.5, 19.3, 13.9, 12.7.

5

Example 98

(+)-(trans)-N-[[2-(2,3-Dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]acetamide

- 10 Obtained 18% of an oil. $[\alpha]_{\text{D}}^{25} = 5.89$. Purity = 99% by HPLC. ^1H NMR (300 MHz, CDCl_3) δ 6.77-6.71 (m, 2H), 6.50-6.44 (m, 1H), 6.04 (s, 1H), 4.36-4.33 (m, 2H), 4.29-4.26 (m, 2H), 3.70-3.62 (m, 1H), 2.87-2.79 (m, 1H), 1.99 (s, 3H), 1.86-1.80 (m, 1H), 1.15-1.00 (m, 2H), 0.87-0.81 (m, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 143.3, 130.4, 120.9, 118.5, 115.4, 15 64.6, 64.1, 44.3, 23.4, 20.8, 16.3, 11.8.

Example 99

- N-[[2-(2-Methyl-4-benzofuranyl)-3,3-difluorocyclopropanyl]-methyl]acetamide**
- 20

Obtained 90% of a colorless liquid. Purity = 98% by HPLC. ^1H NMR (300 MHz, CDCl_3) δ 2.01 (s, 3H), 2.17-2.29 (m, 1H), 2.45 (s, 3H), 2.70 (m, 1H), 3.29 (m, 1H), 3.92 (m, 1H), 5.89 (bs, 1H), 6.44 (s, 1H), 6.97 (d, J=7.5 Hz, 1H), 7.13 (t, J=7.5 Hz, 1H), 7.31 (d, J=7.5 Hz, 1H). ^{13}C NMR (75 25 MHz, CDCl_3) δ 14.2, 23.3, 28.4 (t, J=10.0 Hz), 29.7 (t, J=10.0 Hz), 37.3, 101.7, 110.2, 114.0 (t, J=288 Hz), 121.3, 123.1, 124.4, 129.5, 154.6, 156.0, 170.4.

Example 100**(trans)-N-Methyl-N-[[2-(2,3-dihydrobenzofuran-4-yl)cyclopropyl]methyl]propanamide**

5

To a solution of (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cyclopropyl]methyl]propanamide (120 mg, 0.6 mmol) in THF (1.5 mL) was added sodium hydride (30 mg of 60%, 0.75 mmol). The mixture was allowed to stir for 0.5 h and iodomethane (140 mg, 1 mmol) was added.

10 After stirring for 18 h, the solution was concentrated in vacuo. The residue was dissolved in acetonitrile and washed with hexane. The acetonitrile solution was concentrated in vacuo and the residue purified by chromatography on silica eluting with ethyl acetate-hexane (2:8) to give the product as an amber oil (90 mg, 69%).

15

Anal. Calc'd for $C_{16}H_{21}NO_2 \cdot 0.2H_2O$: C, 73.08; H, 8.20; N 5.33.

Found: C, 72.91; H, 8.24; N, 5.23.

Example 101

20

(trans)-N-[[2-(2,3-Dihydro-5,7-diiodobenzofuran-4-yl)cyclopropyl]methyl]propanamide

To a solution of (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cyclopropyl]methyl]propanamide (0.245 mg, 1 mmol) and lead IV tetraacetate (530 mg, 1.2 mmol) in 1.5 mL acetic acid was added iodine (300 mg, 1.2 mmol). The solution was allowed to stir for 15 min during which time the reaction solidified. The mixture was diluted with methylene chloride and the resulting solution was washed with water and saturated sodium bicarbonate solution and dried. The solution was concentrated in vacuo and the residue crystallized from isopropyl ether to give the product (500 mg, 99%, mp 198-199°C).

Anal. Calc'd for $C_{15}H_{17}I_2NO_2$: C, 36.24; H, 3.45; N 2.82.

Found: C, 36.29; H, 3.59; N, 2.96.

Example 102

5

(trans)-N-[[2-(2,3-Dihydro-5-iodobenzofuran-4-yl)cyclopropyl]methyl]propanamide

To a solution of 0.12 g (0.6 mmol) (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cyclopropyl]methyl]propanamide and 0.27 g (0.6 mmol) of lead IV tetraacetate in 1 mL acetic acid was added 0.075 g (0.25 mmol) iodine. The solution was allowed to stir for 1.5 h and was diluted with methylene chloride. The resulting solution was washed with water and saturated sodium bicarbonate solution and dried. The solution was concentrated in vacuo and the residue purified by chromatography on silica eluting with ethyl acetate-hexane, 3:7 containing 0.2% methanol to give the product (25 mg, 11%, mp 148-149°C).

Anal. Calc'd for $C_{15}H_{18}INO_2 \cdot 0.1H_2O$: C, 48.30; H, 4.92; N 3.75.

Found: C, 47.94; H, 4.67; N, 3.70.

Reasonable variations, such as those which would occur to a skilled artisan, can be made herein without departing from the scope of the invention.


25



We claim:

1. A compound of Formula I or a pharmaceutically acceptable solvate thereof having the Formula:

5



wherein

- 10 Q¹ and Q² each are independently hydrogen or halogen;
 X is CH₂, CH or oxygen;
 Y is CR³, CR³R⁴ or (CH₂)_n, with n = 1-4;
 Z is CH₂, CH or oxygen;
 R is hydrogen, halogen or C₁₋₄ alkyl in both cases;
 m is 1 or 2;
15 R¹ is C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₃ haloalkyl, C₁₋₆
 alkylamino, C₂₋₆ alkenyl, C₁₋₄ alkoxy(C₁₋₄)alkyl, C₁₋₄
 alkylthio(C₁₋₄)alkyl or C₁₋₄ trifluoromethylalkyl;
 R² is hydrogen or C₁₋₄ alkyl; and
 R³ and R⁴ each are independently hydrogen or C₁₋₄ alkyl.

20

2. A compound of claim 1 wherein Q¹ and Q² are independently hydrogen or iodo and m is 1.
3. A compound of claim 2 wherein R¹ is C₁₋₆ alkyl, C₃₋₄ cycloalkyl, C₁₋₃ haloalkyl, C₂₋₃ alkenyl, C₁₋₂ alkoxy(C₁₋₂)alkyl or C₁₋₂ trifluoromethylalkyl and R² is hydrogen.
- 25 4. A compound of claim 3 wherein X is CH₂ and Z is oxygen.



5. A compound of claim 4 selected from the group consisting of:
- (+)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
- 5 (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]methoxyacetamide;
- 10 (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropanecarboxamide;
- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]trifluoroacetamide;
- (trans)-N-[[2-(2,3-dihydro-2,2-dimethyl-benzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- 15 (trans)-N-[[2-(2,3-dihydro-2,2-dimethyl-benzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
- 20 (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide;
- 25 (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]-2-methylpropanamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]chloroacetamide;
- 30 (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]-2-methylpropanamide;



- (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
(-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]methoxyacetamide;
5 (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
(-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
10 (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
15 (trans)-N-[[2-(2,3-dihydro-5,7-diiodobenzofuran-4-yl)cyclopropyl]methyl]propanamide;
(trans)-N-[[2-(2,3-dihydro-5-iodobenzofuran-4-yl)cyclopropyl]methyl]propanamide;
20 (trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]propanamide;
(trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide;
(trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]butanamide; and
25 (+)-(trans)-N-[[2-(3,4-dihydro-2H-1-benzopyran-5-yl)cycloprop-1-yl]methyl]propanamide.

6. A compound of claim 5, selected from the group consisting of:
30 (+)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
(-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;



- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropanecarboxamide
- 5 (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- 10 (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide;
- (+)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]chloroacetamide;
- 15 (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
- (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide;
- 20 (-)-(trans)-N-[[2-(2-methyl-2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
- 25 (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
- (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
- 30 (trans)-N-[[2-(2,3-dihydro-5,7-diiodobenzofuran-4-yl)cyclopropyl]methyl]propanamide; and
- (trans)-N-[[2-(2,3-dihydro-5-iodobenzofuran-4-yl)cyclopropyl]methyl]propanamide.



7. The compound of claim 6, which is (trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide.
- 5 8. The compound of claim 6, which is (-)-(trans)-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-1-yl]methyl]propanamide.
9. A compound of claim 3 wherein X is oxygen, Y is (CH₂)_n in which n is 1 or 2 and Z is oxygen.
- 10
10. A compound of claim 9, selected from the group consisting of:
(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]acetamide;
(-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]acetamide;
(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]propanamide;
15 (-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]
propanamide;
(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]butanamide;
(-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]
butanamide;
20 (trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]cyclopropane
carboxamide;
(-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]
cyclopropane carboxamide;
(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]-2-
25 methylpropanamide;
(-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]-2-
methylpropanamide;
(-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-
yl]methyl]propanamide;
30 (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-
yl]methyl]acetamide;
(-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-
yl]methyl]cyclopropanecarboxamide;



- (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]butanamide;
(+)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]propenamide;
5 (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]trifluoroacetamide;
(-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]-3,3,3-trifluoropropanamide;
(+)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]propanamide;
10 (trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]acetamide;
(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]propanamide;
15 (trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]butanamide;
(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]-2-methylpropanamide; and
(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]cyclopropane carboxamide.
20

11. A compound of claim 10, selected from the group consisting of:
(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]acetamide;
(-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]acetamide;
25 (trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]propanamide;
and (-)-(trans)-N-[[2-(1,3-benzodioxol-4-yl)cycloprop-1-yl]methyl]propanamide.

12. A compound of claim 10, selected from the group consisting of:
30 (-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]propanamide;
(-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]acetamide;



(-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]cyclopropanecarboxamide; and
(-)-(trans)-N-[[2-(2,3-dihydro-1,4-benzodioxin-5-yl)cycloprop-1-yl]methyl]butanamide.

5

13. A compound of claim 3 wherein X is CH, Y is CR³ and Z is oxygen.

14. A compound of claim 13 selected from the group consisting of:

- 10 (-)-(trans)-N-[[2-(2-benzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane
carboxamide;
(-)-(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
(trans)-N-[[2-(2-methyl-benzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
(trans)-N-[[2-(2-methyl-benzofuran-4-yl)cycloprop-1-yl]methyl]-
propanamide;
15 (trans)-N-[[2-(2-methyl-benzofuran-4-yl)cycloprop-1-yl]methyl]-
butanamide;
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]propanamide;
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]butanamide;
20 (trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]-2-
methylpropanamide; and
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]cyclopropane
carboxamide.

- 25 15. A compound of claim 14, selected from the group consisting of:
(trans)-N-[[2-(2-methyl-benzofuran-4-yl)cycloprop-1-yl]methyl]acetamide;
(trans)-N-[[2-(2-methyl-benzofuran-4-yl)cycloprop-1-yl]methyl]-
propanamide;
(trans)-N-[[2-(2-methyl-benzofuran-4-yl)cycloprop-1-yl]methyl]-
30 butanamide; and
(trans)-N-[[2-(benzofuran-4-yl)cycloprop-1-yl]methyl]propanamide.



16. A method of treating sleep disorders in a patient in need of such treatment comprising administering to said patient a therapeutic amount of a compound of claim 1.
- 5 17. A composition useful for treating sleep disorders comprising a therapeutic amount of a compound of claim 1 and a suitable amount of a pharmaceutically acceptable carrier.
- 10 18. A method of treating circadian rhythm-related disorders in a patient in need of such treatment comprising administering to said patient a therapeutic amount of a compound of claim 1.
- 15 19. A composition useful for treating circadian rhythm-related disorders comprising a therapeutic amount of a compound of claim 1 and a suitable amount of a pharmaceutically acceptable carrier.