


In one aspect, there is provided a compound having the general formula (I), optionally in the form of a pharmaceutically acceptable salt, solvate, , tautomer, racemate, enantiomer, or diastereomer or mixture thereof,



(I)

wherein

X¹⁰ is NR¹⁵, N(R¹⁵)C(O), C(O)NR¹⁵, O, C(O), C(O)O, OC(O); N(R¹⁵)SO₂, SO₂N(R¹⁵), S, SO, or SO₂;

R¹⁰ is -H, a -C₁₋₆ alkyl group or a -C(O)-C₁₋₆ alkyl group;

R¹¹ is -H, a -C₁₋₆ alkyl group, or a -C₁₋₆ alkyl group which is substituted by one or more halogen atoms;

R¹² is -H, a -C₁₋₆ alkyl group, or a -C₁₋₆ alkyl group which is substituted by one or more halogen atoms;

or wherein **R¹¹** and **R¹²** can be joined together to form a 3- to 7-membered carbo- or heterocyclic ring;

R¹³ is -R¹⁶, or -X¹⁰-R¹⁶;

R¹⁴ is -H, -(optionally substituted C₁₋₆ alkyl), -(optionally substituted C₃₋₇ cycloalkyl), -(optionally substituted aryl), -C₁₋₄ alkyl-(optionally substituted C₃₋₇ cycloalkyl), or -C₁₋₄ alkyl-(optionally substituted aryl);

It is understood that throughout the present specification the term "a compound having the general formula (I)" encompasses pharmaceutically acceptable salts, solvates, polymorphs, prodrugs, codrugs, cocrystals, tautomers, racemates, enantiomers, or diastereomers or mixtures thereof unless mentioned otherwise.

5

A further embodiment of the present invention relates to a pharmaceutical composition comprising a compound having the general formula (I) and optionally one or more pharmaceutically acceptable excipient(s) and/or carrier(s).

- 10 Thus, in another aspect, there is provided a pharmaceutical composition comprising:
a compound having the general formula (I) as defined herein, optionally in the form of a pharmaceutically acceptable salt, solvate, tautomer, racemate, enantiomer, or diastereomer or mixture thereof,
and optionally one or more pharmaceutically acceptable excipient(s) and/or carrier(s).

15

The compounds having the general formula (I) are useful for treating, ameliorating or preventing viral diseases.

- 20 Thus, in another aspect, there is provided a compound having the general formula (I) as defined herein, optionally in the form of a pharmaceutically acceptable salt, solvate, polymorph, cocrystal, tautomer, racemate, enantiomer, or diastereomer or mixture thereof, wherein the compound is for use in the treatment, amelioration or prevention of a viral disease caused by Orthomyxoviridae.

- 25 In yet another aspect, there is also provided use of a compound having the general formula (I) as defined herein, optionally in the form of a pharmaceutically acceptable salt, solvate, tautomer, racemate, enantiomer, or diastereomer or mixture thereof in the manufacture of a medicament for treating, ameliorating or preventing a viral disease caused by Orthomyxoviridae.

30

In yet another aspect, there is also provided use of a compound having the general formula (I) as defined herein, optionally in the form of a pharmaceutically acceptable salt, solvate, tautomer, racemate, enantiomer, or diastereomer or mixture thereof in the manufacture of a medicament for treating, ameliorating or preventing influenza.

It has been surprisingly found that the compounds according to the present invention which have the bulky group on the left hand ring have improved properties compared to the compounds disclosed in EP-A-2 444 400. In particular, the interaction with protein could be optimized resulting in better binding properties. Furthermore, shifting of the bulky group should avoid problems due to a chiral center and due to the planarization through shift from sp³ to sp². In addition, the crucial vector for additional hydrophobic interactions may be stabilized and improved.


Detailed description of the invention

Before the present invention is described in detail below, it is to be understood that this invention is not limited to the particular methodology, protocols and reagents described herein as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art.

Preferably, the terms used herein are defined as described in "A multilingual glossary of biotechnological terms: (IUPAC Recommendations)", Leuenberger, H.G.W, Nagel, B. and Kölbl, H. eds. (1995), Helvetica Chimica Acta, CH-4010 Basel, Switzerland.

CLAIMS

1. A compound having the general formula (I), optionally in the form of a pharmaceutically acceptable salt, solvate, tautomer, racemate, enantiomer, or diastereomer or mixture thereof,



wherein

X¹⁰ is NR¹⁵, N(R¹⁵)C(O), C(O)NR¹⁵, O, C(O), C(O)O, OC(O); N(R¹⁵)SO₂, SO₂N(R¹⁵), S, SO, or SO₂;

R¹⁰ is -H, a -C₁₋₆ alkyl group or a -C(O)-C₁₋₆ alkyl group;

R¹¹ is -H, a -C₁₋₆ alkyl group, or a -C₁₋₆ alkyl group which is substituted by one or more halogen atoms;

R¹² is -H, a -C₁₋₆ alkyl group, or a -C₁₋₆ alkyl group which is substituted by one or more halogen atoms;


or wherein **R¹¹** and **R¹²** can be joined together to form a 3- to 7-membered carbo- or heterocyclic ring;

R¹³ is -R¹⁶, or -X¹⁰-R¹⁶;

R¹⁴ is -H, -(optionally substituted C₁₋₆ alkyl), -(optionally substituted C₃₋₇ cycloalkyl), -(optionally substituted aryl), -C₁₋₄ alkyl-(optionally substituted C₃₋₇ cycloalkyl), or -C₁₋₄ alkyl-(optionally substituted aryl);

R^{15} is $-H$, $-(\text{optionally substituted } C_{1-6} \text{ alkyl})$, $-(\text{optionally substituted } C_{3-7} \text{ cycloalkyl})$, $-(\text{optionally substituted aryl})$, $-C_{1-4} \text{ alkyl}-(\text{optionally substituted } C_{3-7} \text{ cycloalkyl})$, or $-C_{1-4} \text{ alkyl}-(\text{optionally substituted aryl})$;

R^{16} is selected from



wherein

X is absent, CH_2 , NH , $C(O)NH$, S or O ;

Y is CH_2 ;

Z is O or S ; and

R is independently selected from $-H$, $-C_{1-6} \text{ alkyl}$, $-CF_3$, $-Hal$, $-CN$, $-OH$, and $-O-C_{1-6} \text{ alkyl}$;

R^{17} is $-H$, $-C_{1-6} \text{ alkyl}$, or $-(CH_2CH_2O)_rH$;

R^{18} is $-H$, or $-C_{1-6} \text{ alkyl}$;

R^{**} is independently selected from $-C_{1-6} \text{ alkyl}$, $-C(O)-C_{1-6} \text{ alkyl}$, $-Hal$, $-CF_3$, $-CN$, $-COOR^{17}$, $-OR^{17}$, $-(CH_2)_qNR^{17}R^{18}$, $-C(O)-NR^{17}R^{18}$, and $-NR^{17}-C(O)-C_{1-6} \text{ alkyl}$;

R^* is independently selected from $-H$, $-C_{1-6} \text{ alkyl}$, and $-C_{3-7} \text{ cycloalkyl}$;

q is 0 to 4; and

r is 1 to 3;

wherein the alkyl group, aryl group and/or cycloalkyl group in R^{14} and R^{15} can be optionally substituted with one or more substituents R^{**} .

2. The compound according to claim 1, wherein R^{13} is $-R^{16}$.
3. The compound according to claim 1, wherein R^{13} is $-X^{10}-R^{16}$ and X^{10} is $N(R^{15})SO_2$.
4. The compound according to any of claims 1 to 3, wherein R^{11} and R^{12} are $-H$.
5. The compound according to any of claims 1 to 4, wherein R^{10} is $-H$, or $-(\text{optionally substituted } C_{1-6} \text{ alkyl})$.
6. The compound according to any of claims 1 to 5, wherein R^{14} is $-H$, $-(\text{optionally substituted } C_{1-6} \text{ alkyl})$, or $-(\text{optionally substituted aryl})$.
7. A pharmaceutical composition comprising:
a compound having the general formula (I) as defined in any of claims 1 to 6, optionally in the form of a pharmaceutically acceptable salt, solvate, tautomer, racemate, enantiomer, or diastereomer or mixture thereof,
and optionally one or more pharmaceutically acceptable excipient(s) and/or carrier(s).
8. The pharmaceutical composition according to claim 7, which additionally comprises at least one further medicament which is selected from the group consisting of a polymerase inhibitor which is different from the compound having the general formula (I); neuramidase inhibitor; M2 channel inhibitor; alpha glucosidase inhibitor; ligand of another influenza target; antibiotics, anti-inflammatory agents, lipoxygenase inhibitors, EP ligands, bradykinin ligands, and cannabinoid ligands.
9. A compound having the general formula (I) as defined in any of claims 1 to 6, optionally in the form of a pharmaceutically acceptable salt, solvate, tautomer, racemate, enantiomer, or diastereomer or mixture thereof,
wherein the compound is for use in the treatment, amelioration or prevention of a viral disease caused by Orthomyxoviridae.

10. Use of a compound having the general formula (I) as defined in any of claims 1 to 6, optionally in the form of a pharmaceutically acceptable salt, solvate, tautomer, racemate, enantiomer, or diastereomer or mixture thereof in the manufacture of a medicament for treating, ameliorating or preventing a viral disease caused by Orthomyxoviridae.
11. Use of a compound having the general formula (I) as defined in any of claims 1 to 6, optionally in the form of a pharmaceutically acceptable salt, solvate, tautomer, racemate, enantiomer, or diastereomer or mixture thereof in the manufacture of a medicament for treating, ameliorating or preventing influenza.
12. The compound for use according to claim 9, wherein at least one further medicament which is selected from the group consisting of a polymerase inhibitor which is different from the compound having the general formula (I); neuramidase inhibitor; M2 channel inhibitor; alpha glucosidase inhibitor; ligand of another influenza target; antibiotics, anti-inflammatory agents, lipoxygenase inhibitors, EP ligands, bradykinin ligands, and cannabinoid ligands is to be administered concurrently with, sequentially with or separately from the compound having the general formula (I).
13. The use according to claim 10, wherein at least one further medicament which is selected from the group consisting of a polymerase inhibitor which is different from the compound having the general formula (I); neuramidase inhibitor; M2 channel inhibitor; alpha glucosidase inhibitor; ligand of another influenza target; antibiotics, anti-inflammatory agents, lipoxygenase inhibitors, EP ligands, bradykinin ligands, and cannabinoid ligands is to be administered concurrently with, sequentially with or separately from the compound having the general formula (I).