

OLIGONUCLEOTIDE ANALOGUES INCORPORATING
5-AZA-CYTOSINE THEREIN

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] The present invention relates to design, synthesis and application of oligonucleotide analogues which are useful as therapeutics, diagnostics as well as research reagents. Oligonucleotide analogues are provided that incorporate an analogue of cytosine, 5-aza-cytosine, in the oligonucleotide sequence, e.g., in the form of 5-aza-2'-deoxycytidine or 5-aza-cytidine. Such analogues can be used for modulation of DNA methylation, especially for effective inhibition of methylation of cytosine at the C-5 position by more specifically targeting the CpG islets of the human genome. Methods for synthesizing these oligonucleotide analogues and for modulating C-5 cytosine methylation are provided. In particular, phosphoramidite building blocks and oligonucleotides containing decitabine (5-aza-2'-deoxycytidine; D), DpG-rich (Decitabine-phosphodiester linkage-Guanosine) islets and derivatives, are provided. Also provided are methods for preparing, formulating and administering these compounds or compositions as therapeutics to a host in need thereof.

Description of Related Art

[0002] Decitabine is currently being developed as a new pharmaceutical for the treatment of chronic myelogenous leukemia (CML), myelodysplastic syndrome (MDS), non-small cell lung (NSCL) cancer, sickle-cell anemia, and acute myelogenous leukemia (AML). Two isomeric forms of decitabine can be distinguished. The β -anomer is the active form, which is shown in Figure 1. Decitabine possesses multiple pharmacological characteristics. At the molecular level, it is incorporated into DNA during the S phase of cell cycle. At the cellular level, decitabine can induce cell differentiation and exert hematological toxicity. Despite having a short half-life *in vivo*, decitabine has an excellent tissue distribution.

[0003] One of the functions of decitabine is its ability to specifically and potently inhibit DNA methylation. DNA methylation is an epigenetic effect common to many systems. This modification involves the covalent modification of cytosine at the C-5 position (1a''). Methylation patterns are stably maintained at CpG dinucleotides by a family of DNA methyltransferases that recognize hemimethylated DNA after DNA replication. Inside the cell, decitabine is first converted into its active form, the phosphorylated 5-aza-deoxycytidine, by deoxycytidine kinase, which is primarily synthesized during the S phase of the cell cycle. The affinity of decitabine for the catalytic site of deoxycytidine kinase is similar to the natural substrate, deoxycytidine (Momparler et al. 1985 Mol. Pharmacol. 30:287-299). After conversion to its triphosphate form by deoxycytidine kinase, decitabine is incorporated into replicating DNA at a rate similar to that of the natural substrate, dCTP (Bouchard and Momparler 1983 Mol. Pharmacol. 24:109-114).

[0004] CpG-rich sequences of housekeeping genes are generally protected from methylation in normal cells. In cancerous cells, aberrant hypermethylation in promoter region CpG-islands of tumor suppressor genes is one of the most common events associated with progression of the tumorigenic phenotype. Each class of differentiated cells has its own distinct methylation pattern. Incorporation of decitabine into the DNA strand has a hypomethylation effect. After chromosomal duplication, in order to conserve this pattern of methylation, the 5-methylcytosine on the parental strand serves to direct methylation on the complementary daughter DNA strand. Substituting the carbon at the C-5 position of the cytosine for nitrogen interferes with this normal process of DNA methylation. The

PCT/US06/37313
replacement of cytosine with decitabine at a specific site of methylation produces an irreversible inactivation of DNA methyltransferases. Decitabine behaves faithfully as a cytosine residue until DNA methyltransferase enzymes attempt to transfer a methyl group to the hemimethylated DNA strands of the daughter cells. At this step the DNA methyltransferase enzyme is covalently trapped by decitabine in the DNA and cannot further silence (methylate) additional cytosine residues (Juttermann et al. 1994 Proc. Natl. Acad. Sci. USA 91:11797-11801). This unique mechanism of action of decitabine allows genes silenced (that were once methylated) from previous rounds of cell division to be re-expressed. The active trap is present in the hemimethylated DNA up to 48 hours after decitabine treatment. After further DNA synthesis and cell cycle division, progeny strands from the hemimethylated DNA result in DNA strands that are completely unmethylated at these sites (Jones P. 2001 Nature 409: 141, 143-4). By specifically inhibiting DNA methyltransferases, the enzyme required for methylation, aberrant methylation of the tumor suppressor genes could be reversed.

[0005] Despite its proven antileukemic effects in CML, MDS, and AML, the potential application of decitabine has been hampered by delayed and prolonged myelosuppression. Lower doses of decitabine, given over a longer period of time, have minimized myelosuppression to manageable levels without compromising its ability to suppress cancer via its hypomethylation effect. At higher doses, the associated toxicity was prohibitive. However, treatment of hematologic and solid tumors at maximally tolerated doses of decitabine has been ineffective. The cause of myelosuppression is not clear. It is plausible that since decitabine is randomly and extensively incorporated into the DNA of S phase cells, including bone marrow cells that are involved in normal hematopoiesis, the severe DNA damage due to the instability of decitabine leads to necrosis. Since incorporation of decitabine is not restricted to only the CpG-rich sequences, the DNA can break, due to the instability of decitabine, and require repair at numerous sites outside of the CpG islands.

[0006] Decitabine and azacitidine are unstable in aqueous media and undergo hydrolytic degradation. In acidic medium, decitabine is hydrolyzed at room temperature to 5-azacytosine and 2-deoxyribose. In neutral medium at room temperature, the opening of the triazine ring takes place at the 6-position to form the transient intermediate formyl derivative, which further degrades to the amidino-urea derivative and formic acid (Piskala, A.; Synackova, M.; Tomankova, H.; Fiedler, P.; Zizkowsky, V. *Nucleic Acids Res.* 1978, 4, s109-s-113.). This hydrolysis at the 6-position occurs in acidic and basic aqueous media at even faster rates.

[0007] In view of the chemical instability and toxicities associated with decitabine, there exists a need to develop not only more stable derivatives of decitabine but superior hypomethylating agents, where incorporation is localized to the CpG islands as much as possible or hypomethylation is achieved without significantly affecting the integrity of the DNA.

SUMMARY OF THE INVENTION

[0008] The present disclosure provides oligonucleotide analogues that incorporate 5-aza-cytosine in the oligonucleotide sequence, e.g., in the form of 5-aza-2'-deoxycytidine (decitabine) or 5-aza-cytidine.

[0009] In one aspect, there is provided an isolated or synthetic oligonucleotide analogue, its salt or ester, comprising one 5-aza-cytosine as a base residue in the sequence of the dinucleotide analogue, wherein the dinucleotide is linked with a phospholinker consisting of one phosphorous and wherein the dinucleotide analogue has a general formula of: a. Z-L-G, or G-L-Z, wherein Z is 5-aza-cytosine; G is guanine; and L is a chemical linker covalently linking Z and G; or b. 5'-DpG-3' or 5'GpD-3', wherein D is decitabine; p is a phospholinker; and G is deoxyguanosine.


[0009a] In a second aspect, there is provided a pharmaceutical composition comprising an dinucleotide analogue as defined above or its pharmaceutically acceptable salt or ester, and a pharmaceutically-acceptable carrier.

[0009b] In a third aspect, there is provided a kit, comprising: a first vessel containing an dinucleotide analogue as defined above, its salt or ester thereof.

[0009c] In a fourth aspect, there is provided a method for inhibiting DNA methylation in a cell in vitro, comprising: contacting the cell with the dinucleotide analogue, its salt or ester thereof as defined above, such that DNA methylation activity of the cell is inhibited.

[0009d] In a fifth aspect, there is provided the use of a pharmaceutical composition comprising a therapeutically-effective amount of the dinucleotide analogue, its salt or ester thereof as defined above, and a pharmaceutically-acceptable carrier, in the manufacture of a medicament for treating a disease associated with aberrant DNA methylation.


[0010] In an embodiment, the oligonucleotide analogue has a general formula:



PCT/US2006/037313
 wherein Z is 5-aza-cytosine; G is guanine; and L is a chemical linker covalently linking Z and G. The oligonucleotide analogue optionally has more than 30%, 35%, or 40% guanine residues in the sequence of the oligonucleotide analogue.

[0011] In particular embodiments, the oligonucleotide analogue is selected from the group consisting of 5'-DpG-3', 5'-GpD-3', 5'-DpGpD-3', 5'-GpGpD-3', 5'-GpDpG-3', 5'-GpDpD-3', 5'-DpDpG-3', 5'-DpGpG-3', 5'-GpDpD-3', 5'-DpGpA-3', 5'-DpGpDpG-3', 5'-DpGpGpD-3', 5'-GpDpGpD-3', 5'-GpDpDpG-3', 5'-DpGpDpGpA-3', wherein D is decitabine; p is a phospholinker; A is 2'-deoxyadenosine, and G is 2'-deoxyguanosine.

[0012] In another aspect of the invention, an isolated or synthetic oligonucleotide analogue is provided which comprises, 2 or more copies of a dinucleotide analogue having the general formula:




wherein Z is 5-aza-cytosine; G is guanine; and L is a chemical linker covalently linking Z and G.

[0013] Optionally, the oligonucleotide analogue comprises less than 10, 8, 6, or 4 copies of the dinucleotide analogue $-Z-L-G-$, or $-G-L-Z-$.

[0014] In particular embodiments, the oligonucleotide analogue comprises a segment selected from the group consisting of 5'-DpG-3', 5'-GpD-3', 5'-DpGpD-3', 5'-GpGpD-3', 5'-GpDpG-3', 5'-GpDpD-3', 5'-DpDpG-3', 5'-DpGpG-3', 5'-GpDpD-3', 5'-DpGpA-3', 5'-DpGpDpG-3', 5'-DpGpGpD-3', 5'-GpDpGpD-3', 5'-GpDpDpG-3', 5'-DpGpDpGpA-3', wherein D is decitabine; p is a phospholinker; A is 2'-deoxyadenosine, and G is 2'-deoxyguanosine.

[0015] Disclosed is an isolated or synthetic oligonucleotide analogue having at least 6 bases in length is provided, which comprises one or more 5-aza-cytosine residues in the sequence of the oligonucleotide analogue and has at least 75% sequence homology with a segment of a gene, preferably the 5'-untranslated region of a gene, such as the promoter of the gene.

[0016] In an embodiment, the oligonucleotide analogue has a general formula:



wherein Z is 5-aza-cytosine; G is guanine; and L is a chemical linker covalently linking Z and G. The oligonucleotide analogue optionally has more than 30%, 35%, or 40% guanine residues in the sequence of the oligonucleotide analogue.

[0017] In particular embodiments, the oligonucleotide analogue comprises a segment selected from the group consisting of 5'-DpG-3', 5'-GpD-3', 5'-DpGpD-3', 5'-GpGpD-3', 5'-GpDpG-3', 5'-GpDpD-3', 5'-DpDpG-3', 5'-DpGpG-3', 5'-GpDpD-3', 5'-DpGpA-3', 5'-DpGpDpG-3', 5'-DpGpGpD-3', 5'-GpDpGpD-3', 5'-GpDpDpG-3', 5'-DpGpDpGpA-3', wherein D is decitabine; p is a phospholinker; A is 2'-deoxyadenosine, and G is 2'-deoxyguanosine.

[0018] Also disclosed is an oligonucleotide analogue is provided that binds an allosteric site on DNA methyltransferases thereby inhibiting DNA methyltransferases.

[0019] In one embodiment, the oligonucleotide analogue has a sequence of

[0020] 5'-CTGGATCCTTGCCCCGCCCCCTTGAATTC-3' (SEQ ID NO:25),

[0021] 5'-GGGAATTCAAATGACGTCAAAAAGGATCCAG-3' (SEQ ID NO:26);

[0022] 5'-CCTACCCACCCCTGGATCCTTGCCCCGCCCCCTTGAATTC-3' (SEQ ID NO:27);

[0023] CCTCCAC-3' (SEQ ID NO:27);

[0024] 5'-ATCCTTGCCCCGCCCCCTTGAAT-3' (SEQ ID NO:28); or



[0025] 5'-TTGCCCCGCCCCCTT (SEQ ID NO:29), wherein at least one of the cytosine residues in SEQ ID NOs: 25-29 is substituted with 5-aza-cytosine.

[0026] For example, the oligonucleotide analogue may be

[0027] 5'-CTGGATCCTTGCCCDGCCCCCTGAATTCCC-3' (SEQ ID NO:30), wherein one of the 14 cytosine residues in SEQ ID NO:25 at nucleotide position 15 is substituted with 5-aza-cytosine.


[0028] Also disclosed is an oligonucleotide analogue is provided that is at least 6 nucleotide long, has at least one 5-aza-cytosine as a base residue and adopts a hairpin conformation at ambient temperature, such as 20-25°C, in aqueous solution, such as water, saline, or a buffer comprising 20mM HEPES (pH 7), 12% glycerol, 1mM EDTA, 4mM dithiothreitol, 0.1% Nonidet P-40, and 3 mM Mg Cl₂.

[0029] In one embodiment, the oligonucleotide analogue has the following general secondary structure:



wherein N is any nucleotide; N' is a nucleotide complementary to N; Z is 5-aza-cytosine as a base residue; G is guanine as a base residue; l, n, or m is an integer; nucleotide N_n, N_m, N'_n, and N'_m are positioned in the stem region of the hairpin; and N_l is positioned in the loop region of the hairpin. Preferably, l, n, or m is an integer greater than 2, 3, 4, or 5. Optionally, l is 2, 3, 4, 5, or 6. Also optionally, if N_n, N_m, or N_l has one or more cytosine residues, the cytosine residue is substituted with 5-aza-cytosine.

[0030] In a particular embodiment, the oligonucleotide analogue (SEQ ID NO:31) has the following hairpin conformation:



wherein D is decitabine, A is adenosine or 2'-deoxyadenosine, T is thymidine or 2'-deoxythymidine, and C at nucleotide position 21 is optionally substituted with 5-methyl-2'-deoxycytidine.

[0031] In any of above embodiments, the oligonucleotide analogue can be single-stranded or double-stranded.

When the oligonucleotide analogue is double-stranded, the first strand is the oligonucleotide analogue, and the second strand may be an oligonucleotide with sequence complementary to that of the first strand without the cytosine residue being replaced with 5-aza-cytosine. For example, the first strand may be 5'-TTDGDGAA-3' (SEQ ID NO: 32) wherein D is decitabine; whereas the second strand may be 5'-TTCGCGAA-3' (SEQ ID NO: 33).

[0032] Optionally, when the second strand of oligonucleotide comprises one or more cytosine residues, and at least one of the cytosine residues is substituted with 5-methyl-cytosine.

[0033] Also optionally, when the first strand has a segment of 5'-Z-L-G-3', and the second strand comprises a segment of 3'-G-L-C'-5' that matches with the segment of 5'-Z-L-G-3' in the first strand, wherein Z is 5-aza-cytosine; G is guanine; L is a chemical linker covalently linking Z and G, or G and C'; and C' is 5-methyl-cytosine.

[0034] Also optionally, when the first strand has a segment of 5'-G-L-Z-3', and the second strand comprises a segment of 3'-C'-L-G-5' that matches with the segment of 5'-G-L-Z-3' in the first strand, wherein Z is 5-aza-cytosine; G is guanine; L is a chemical linker covalently linking Z and G, or G and C'; and C' is 5-methyl-cytosine.

[0035] The present disclosure also provides methods for synthesizing the oligonucleotide analogues and for modulating nucleic acid methylation. Also provided are phosphoramidite building blocks for synthesizing the

PCT/US2006/37313
oligonucleotide analogues, formulating and administering these compounds or compositions to treat conditions, such as cancer and hematological disorders.

INCORPORATION BY REFERENCE

- 5 [0036] All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

- 10 [0037] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:
- [0038] Figure 1 shows structure of decitabine, D (1a), cytosine, C (1a'), and 5-methyl cytosine, mC (1a'').
- 15 [0039] Figure 2A depicts examples of decitabine phosphoramidite building blocks.
- [0040] Figure 2B depicts decitabine phosphoramidite building block 1d, where R₁ = phenoxyacetyl.
- [0041] Figure 3A depicts 3'- and 5'-O-capped and controlled-pore glass 3'-linked decitabine derivatives.
- [0042] Figure 3B depicts protected decitabine 3'-linked onto controlled-pore glass support.
- [0043] Figure 4 depicts examples of 3'-O-capped decitabine derivatives.
- 20 [0044] Figure 5 illustrates a standard cycle for oligonucleotide synthesis.
- [0045] Figure 6A shows synthesis schemes of GpD dinucleotides and tetranucleotides, where X⁺ is a counter ion.
- [0046] Figure 6B depicts a model synthesis cycle of GpD dinucleotide (2a) and DpGpGpD tetranucleotide (3b).
- [0047] Figure 7 depicts DpGpGpD tetranucleotide.
- [0048] Figure 8 shows synthesis schemes of GpD dinucleotides and tetranucleotides.
- 25 [0049] Figure 9 depicts GpDpG trinucleotide and GpDpDpG tetranucleotide.
- [0050] Figure 10 depicts protected decitabine 3'-linked onto poly (ethylene glycol).
- [0051] Figure 11 shows synthesis schemes of 3'- and 5'-O-capped GpD dinucleotide.
- [0052] Figure 12 shows synthesis schemes of 3'- and 5'-O-capped DpG dinucleotide.
- [0053] Figure 13 depicts GpD and DpG dinucleotides with nuclease resistant phosphothioate linkage.
- 30 [0054] Figure 14 depicts GpDpGpD and DpGpGpD tetranucleotides with nuclease resistant phosphothioate linkage.
- [0055] Figure 15 depicts DpGpDpG and GpDpDpG tetranucleotides with nuclease resistant phosphothioate linkage.
- [0056] Figure 16 depicts GpD and DpG dinucleotides with cytidine deaminase resistant 4-amino groups.
- 35 [0057] Figure 17 depicts GpD and DpG dinucleotides with cytidine deaminase resistant 4-amino groups and nuclease resistant phosphothioate linkage.
- [0058] Figure 18 depicts GpDpGpD and DpGpGpD tetranucleotides with cytidine deaminase resistant 4-amino groups.
- [0059] Figure 19 depicts GpDpGpD and DpGpGpD tetranucleotides with cytidine deaminase resistant 4-amino groups and nuclease resistant phosphothioate linkage.
- 40

PCT/US2006/37313

[0060] Figure 20 depicts Cap-O-GpD-O-Cap and Cap-O-DpG-O-Cap dinucleotides with cytidine deaminase resistant 4-amino groups.

[0061] Figure 21 depicts Cap-O-GpD-O-Cap and Cap-O-DpG-O-Cap dinucleotides with cytidine deaminase resistant 4-amino groups and nuclease resistant phosphothioate linkage.

5 [0062] Figure 22 depicts DpGpDpG and GpDpDpG tetranucleotides with cytidine deaminase resistant 4-amino groups.

[0063] Figure 23 depicts DpGpDpG and GpDpDpG tetranucleotides with cytidine deaminase resistant 4-amino groups and nuclease resistant phosphothioate linkage

[0064] Figure 24A depicts -DpG- islets with natural phosphodiester backbone or modified backbones.

10 [0065] Figure 24B depicts a -DpG- islet peptide backbone.

[0066] Figure 25 schematically illustrates a cell-based GFP assay for DNA methylation Panel a) shows control cells; and panel b) cells treated with oligonucleotides of the present invention and expressing GFP.

[0067] Figure 26 lists examples of inventive oligonucleotide analogues specifically targeting the promoter region of P15, BRAC1 or P16, where D can be decitabine or decitabine analogues and px = p for natural phosphate linkage, px = ps for phosphorothioate linkage, px = bp for boranophosphate, px = mp for methylphosphonate linkage.

[0068] Figure 27 lists the sequence of P15 promoter region and examples of segments thereof, based on which DpG and GpD rich oligonucleotide analogues can be made.

[0069] Figure 28 lists the sequence of P16 promoter region and examples of segments thereof, based on which DpG and GpD rich oligonucleotide analogues can be made.

[0070] Figure 29 lists the sequence of BRCA1 promoter region and examples of segments thereof, based on which DpG and GpD rich oligonucleotide analogues can be made.

[0071] Figure 30 is a mass spectrum of GpD (2a) triethylammonium salt.

[0072] Figure 31 is a mass spectrum of DpG (2b) triethylammonium salt.

25 [0073] Figure 32 is a mass spectrum of DpGpGpD (3b) triethylammonium salt.

[0074] Figure 33 is a mass spectrum of GpDpG (3c') triethylammonium salt.

[0075] Figure 34 is a mass spectrum of DpGpDpG (3c) triethylammonium salt.

[0076] Figure 35 is a mass spectrum of phosphorothioate linked DpG (2f) triethylammonium salt.

[0077] Figure 36 is a mass spectrum of DpG (2b) sodium salt.

30 [0078] Figure 37 is a mass spectrum of HEG-DpG (2d) triethylammonium salt.

[0079] Figure 38 is a mass spectrum of decitabine phosphoramidite building block (1d; R₁ = phenoxyacetyl).

DETAILED DESCRIPTION OF THE INVENTION

35 [0080] The present invention provides oligonucleotide analogues which incorporate an analogue of cytosine, 5-aza-cytosine, in the oligonucleotide sequence, e.g., in the form of 5-aza-2'-deoxycytidine (also known as decitabine) or 5-aza-cytidine (5-azaC). It is believed that incorporation of one or more residues of 5-aza-cytosine into an oligonucleotide would have a DNA hypomethylation effect as replacement of cytosine with decitabine at a specific site of methylation produces an irreversible inactivation of DNA methyltransferase. Preferably decitabine is

40 incorporated into the oligonucleotide 5'-adjacent to a guanine residue to form a DpG islet in order to more specifically target the CpG islets of the human genome.

PCT/US2006/37313
[0081]

The invention is aimed to overcome potential toxicities associated with conventional hypomethylating agents such as decitabine and 5-aza-cytidine. Compared to the free nucleoside forms, which are randomly and extensively incorporated into the whole genome, the inventive compounds could act as primers and are incorporated mainly into the CpG-rich islands of the DNA during replication. Preferably, the inventive compounds act as primers and are incorporated specifically into the CpG-rich islands of the promoters of therapeutically or diagnostically important genes, such as the tumor suppressor genes. The inventive compounds could form temporarily hemimethylated stands with the parental strand and function as the active trap of DNA methyltransferases without being incorporated. The inventive compounds may also directly occupy and trap DNA methyltransferases without being incorporated into the genome. Since DNA modification is localized to the CpG-rich islands in the promoter regions of tumor suppressor genes, when the inventive compounds are incorporated, the active trap is optimally placed and overall stability of the greater genome remains uncompromised.

[0082] By modulating DNA methylation, the inventive compounds can be used as therapeutics, diagnostics as well as research reagents, especially in the areas of cancer and hematological disorders. Aberrant transcriptional silencing of a number of genes, such as tumor suppressor genes, is directly related to pathogenesis of cancer and other diseases. Due to methylation of cancer-related genes, expression of these genes is suppressed or completely silenced. Meanwhile, expression of these genes is required for induction of growth arrest, differentiation, and/or apoptotic cell death of transformed cells. Inaction of these genes in the transformed cells leads to uncontrolled proliferation of these cells, which eventually results in cancer. Thus, by using the inventive compounds to actively trap DNA methyltransferases directly and without being incorporated into the genome or incorporated into the CpG-rich islands of these genes, transcription of the genes can be reactivated through inhibition of methylation of the promoters, thereby resulting suppression of cancer cell proliferation.

[0083] The compounds of the present invention can also be useful for research and diagnostics, because some embodiments of the inventive compounds can hybridize to a 5'-untranslated region or promoter sequence of a gene, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the oligonucleotide analogues of the invention with the promoter sequence can be detected by means known in the art. Such means may include conjugation or non-covalently binding of an enzyme to the oligonucleotide analogue, radiolabelling of the oligonucleotide analogue or any other suitable detection means. Kits using such detection means for modulating activity of the promoter of the gene in a sample may also be prepared.

[0084] The present invention also provides methods for synthesizing these oligonucleotide analogues and for modulating C-5 cytosine methylation. In particular, phosphoramidite building blocks and oligonucleotides containing decitabine (5-aza-2'-deoxycytidine; D), DpG-rich (Decitabine-phosphodiester linkage-Guanosine) islets and derivative, are provided. Also provided are methods for preparing, formulating and administering these compounds or compositions as therapeutics to a host in need thereof. The inventive compounds, methods of synthesis, formulation of pharmaceutical compositions, preparation of vessels and kits, and use of the compounds or compositions for treating diseases or conditions are described in detail below.

1. Oligonucleotide Analogues of the Present Invention

[0085] In general the oligonucleotide analogue of the present invention has one or more residues of 5-aza-cytosine (hereinafter abbreviated as "Z") incorporated into an oligonucleotide sequence.

PCT/US2006/037313
[0086]

In one aspect of the invention, an isolated or synthetic oligonucleotide analogue having 12 or less bases in length is provided, which comprises one or more 5-aza-cytosine residues in the sequence of the oligonucleotide analogue.

[0087] In an embodiment, the oligonucleotide analogue has a general formula:



wherein Z is 5-aza-cytosine; G is guanine; and L is a chemical linker covalently linking Z and G. The oligonucleotide analogue optionally has more than 30%, 35%, or 40% guanine residues in the sequence of the oligonucleotide analogue.

10 [0088] In particular embodiments, the oligonucleotide analogue is selected from the group consisting of 5'-DpG-3', 5'-GpD-3', 5'-DpGpD-3', 5'-GpGpD-3', 5'-GpDpG-3', 5'-GpDpD-3', 5'-DpDpG-3', 5'-DpGpG-3', 5'-GpDpD-3', 5'-DpGpA-3', 5'-DpGpDpG-3', 5'-DpGpGpD-3', 5'-GpDpGpD-3', 5'-GpDpDpG-3', 5'-DpGpDpGpA-3', wherein D is decitabine; p is a phospholinker; A is 2'-deoxyadenosine, and G is 2'-deoxyguanosine.

[0089] In another aspect of the invention, an isolated or synthetic oligonucleotide analogue is provided which comprises, 2 or more copies of a dinucleotide analogue having the general formula:



wherein Z is 5-aza-cytosine; G is guanine; and L is a chemical linker covalently linking Z and G.

[0090] Optionally, the oligonucleotide analogue comprises less than 10, 8, 6, or 4 copies of the dinucleotide analogue $-Z-L-G-, \text{ or } -G-L-Z-.$

20 [0091] In particular embodiments, the oligonucleotide analogue comprises a segment selected from the group consisting of 5'-DpG-3', 5'-GpD-3', 5'-DpGpD-3', 5'-GpGpD-3', 5'-GpDpG-3', 5'-GpDpD-3', 5'-DpDpG-3', 5'-DpGpG-3', 5'-GpDpD-3', 5'-DpGpA-3', 5'-DpGpDpG-3', 5'-DpGpGpD-3', 5'-GpDpGpD-3', 5'-GpDpDpG-3', 5'-DpGpDpGpA-3', wherein D is decitabine; p is a phospholinker; A is 2'-deoxyadenosine, and G is 2'-deoxyguanosine.

25 [0092] In yet another aspect of the invention, an isolated or synthetic oligonucleotide analogue having at least 6 bases in length is provided, which comprises one or more 5-aza-cytosine residues in the sequence of the oligonucleotide analogue and has at least 75% sequence homology with a segment of a gene, preferably the 5'-untranslated region of a gene, such as the promoter of the gene.

[0093] In an embodiment, the oligonucleotide analogue has a general formula:



wherein Z is 5-aza-cytosine; G is guanine; and L is a chemical linker covalently linking Z and G. The oligonucleotide analogue optionally has more than 30%, 35%, or 40% guanine residues in the sequence of the oligonucleotide analogue.

35 [0094] In particular embodiments, the oligonucleotide analogue comprises a segment selected from the group consisting of 5'-DpG-3', 5'-GpD-3', 5'-DpGpD-3', 5'-GpGpD-3', 5'-GpDpG-3', 5'-GpDpD-3', 5'-DpDpG-3', 5'-DpGpG-3', 5'-GpDpD-3', 5'-DpGpA-3', 5'-DpGpDpG-3', 5'-DpGpGpD-3', 5'-GpDpGpD-3', 5'-GpDpDpG-3', 5'-DpGpDpGpA-3', wherein D is decitabine; p is a phospholinker; A is 2'-deoxyadenosine, and G is 2'-deoxyguanosine.

40 [0095] The gene is preferably a mammalian gene, and more preferably a human gene, and most preferably a human tumor suppressor gene. Examples of the human gene include, but are not limited to, VHL (the Von Hippel-Landau gene involved in Renal Cell Carcinoma); P16/INK4A (involved in lymphoma); E-cadherin (involved in metastasis of breast, thyroid, gastric cancer); hMLH1 (involved in DNA repair in colon, gastric, and endometrial cancer); BRCA1 (involved in DNA repair in breast and ovarian cancer); LKB1 (involved in colon and breast

PCT/US2006/037313
cancer); P15/INK4B (involved in leukemia such as AML and ALL); ER (estrogen receptor, involved in breast,

colon cancer and leukemia); O6-MGMT (involved in DNA repair in brain, colon, lung cancer and lymphoma); GST-pi (involved in breast, prostate, and renal cancer); TIMP-3 (tissue metalloprotease, involved in colon, renal, and brain cancer metastasis); DAPK1 (DAP kinase, involved in apoptosis of B-cell lymphoma cells); P73 (involved in apoptosis of lymphomas cells); AR (androgen receptor, involved in prostate cancer); RAR-beta (retinoic acid receptor-beta, involved in prostate cancer); Endothelin-B receptor (involved in prostate cancer); Rb (involved in cell cycle regulation of retinoblastoma); P14ARF (involved in cell cycle regulation); RASSF1 (involved in signal transduction); APC (involved in signal transduction); Caspase-8 (involved in apoptosis); TERT (involved in senescence); TERC (involved in senescence); TMS-1 (involved in apoptosis); SOCS-1 (involved in growth factor response of hepatocarcinoma); PITX2 (hepatocarcinoma breast cancer); MINT1; MINT2; GPR37; SDC4; MYOD1; MDR1; THBS1; PTC1; and pMDR1, as described in Santini et al. (2001) Ann. of Intern. Med. 134:573-586, which is herein incorporated by reference in its entirety. Nucleotide sequences of these genes can be retrieved from the website of the National Center for Biotechnology Information (NCBI).

[0096] As examples, the promoter sequences of the tumor suppressor genes, p15, p16, and BRCA1, are shown in Figure 27, 28, and 29, respectively. Examples of oligonucleotide analogues with at least 75% sequence homology with a segment of p15, p16, and BRCA1 are shown in Figure 27, 28, and 29, respectively.

[0097] It is appreciated by skilled artisans in the field of nucleic acids that the higher degree of sequence homology of a tester polynucleotide with its target polynucleotide, the higher stringency of the condition which the tester polynucleotide can remain hybridized to the target polynucleotide. Thus, the oligonucleotide analogues of the present invention that are designed to target a specific gene are those oligonucleotide analogues that hybridize to the target gene under very low to very high stringency conditions.

[0098] For oligonucleotide analogues of at least about 100 nucleotides in length, very low to very high stringency conditions are defined as prehybridization and hybridization at 42° C. in 5× SSPE, 0.3% SDS, 200 µg/ml sheared and denatured salmon sperm DNA, and either 25% formamide for very low and low stringencies, 35% formamide for medium and medium-high stringencies, or 50% formamide for high and very high stringencies, following standard Southern blotting procedures. The carrier material is finally washed three times each for 15 minutes using 2× SSC, 0.2% SDS preferably at least at 45° C (very low stringency), more preferably at least at 50° C (low stringency), more preferably at least at 55° C (medium stringency), more preferably at least at 60° C (medium-high stringency), even more preferably at least at 65° C. (high stringency), and most preferably at least at 70° C (very high stringency).

[0099] For shorter oligonucleotide analogues which are about 50 nucleotides to about 100 nucleotides in length, stringency conditions are defined as prehybridization, hybridization, and washing post-hybridization at 5° C to 10° C below the calculated T_m using the calculation according to Bolton and McCarthy (1962, *Proceedings of the National Academy of Sciences USA* 48:1390) in 0.9 M NaCl, 0.09 M Tris-HCl pH 7.6, 6 mM EDTA, 0.5% NP-40, 1× Denhardt's solution, 1 mM sodium pyrophosphate, 1 mM sodium monobasic phosphate, 0.1 mM ATP, and 0.2 mg of yeast RNA per ml following standard Southern blotting procedures. The carrier material is washed once in 6× SSC plus 0.1% SDS for 15 minutes and twice each for 15 minutes using 6× SSC at 5° C to 10° C below the calculated T_m .

[00100] Another non-limiting examples of high stringency conditions include a hybridization solution containing, e.g., about 5X SSC, 0.5% SDS, 100 µg/ml denatured salmon sperm DNA and 50% formamide, at 42° C. Blots can be washed at high stringency conditions that allow, e.g., for less than 5% bp mismatch (e.g., wash twice in 0.1% SSC and 0.1% SDS for 30 min at 65° C), i.e., selecting sequences having 95% or greater sequence identity.

PCT/US2006/037313

[00101] Yet another non-limiting example of high stringency conditions includes a final wash at 65° C in aqueous buffer containing 30 mM NaCl and 0.5% SDS. Another example of high stringent conditions is hybridization in 7% SDS, 0.5 M NaPO₄, pH 7, 1 mM EDTA at 50° C, e.g., overnight, followed by one or more washes with a 1% SDS solution at 42° C. Whereas high stringency washes can allow for less than 5% mismatch, reduced or low stringency conditions can permit up to 20% nucleotide mismatch. Hybridization at low stringency can be accomplished as above, but using lower formamide conditions, lower temperatures and/or lower salt concentrations, as well as longer periods of incubation time.

[00102] In a preferred embodiment, the oligonucleotide analogue is capable of hybridizing with the target gene under low stringency conditions. In a more preferred embodiment, the oligonucleotide analogue is capable of hybridizing with the target gene under medium stringency conditions. In a most preferred embodiment, the oligonucleotide analogue is capable of hybridizing with the target gene under high stringency conditions.

[00103] In yet another aspect of the invention, an oligonucleotide analogue is provided that binds an allosteric site on DNA methyltransferase thereby inhibiting DNA methyltransferase. The inhibition of DNA methyltransferase prevents the methylation of DNA thereby treating the disorder associated with aberrant DNA methylation, such as cancer and hematological disorders.

[00104] In one embodiment, the oligonucleotide analogue has a sequence of

[00105] 5'-CTGGATCCTTGCCCCGCCCCCTTGAATTCCC-3' (SEQ ID NO:25);

[00106] 5'-GGGAATTCAAATGACGTCAAAAGGATCCAG-3' (SEQ ID NO:26);

[00107] 5'-CCTACCCACCCTGGATCCTTGCCCCGCCCCCTTGAATTCCCAA

[00108] CCTCCAC-3' (SEQ ID NO:27);

[00109] 5'-ATCCTTGCCCCGCCCCCTTGAAT-3' (SEQ ID NO:28); or


[00110] 5'-TTGCCCCGCCCCCTT (SEQ ID NO:29), wherein at least one of the cytosine residues in SEQ ID NOs: 25-28 is substituted with 5-aza-cytosine. For example, the oligonucleotide analogue may be


[00111] 5'-CTGGATCCTTGCCCDGCCCTTGAATTCCC-3' (SEQ ID NO:30)

wherein one of the 14 cytosine residues in SEQ ID NO:25 at nucleotide position 15 is substituted with 5-aza-cytosine. Other examples of oligonucleotides that bind to DNA methyltransferase can be modified according to the present invention by substituting at least one of the cytosine residues can be found in WO 99/12027, which is herein incorporated by reference in its entirety. The assays for testing the activity of the oligonucleotide analogues of the present invention in binding and inhibiting activity of DNA methyltransferase can also be found in WO 99/12027.

[00112] In yet another aspect of the invention, an oligonucleotide analogue is provided that is at least 6 nucleotide long, has at least one 5-aza-cytosine as a base residue and adopts a hairpin conformation at ambient temperature, such as 20-25°C, in aqueous solution, such as water, saline, or a buffer comprising 20 mM HEPES (pH 7), 12% glycerol, 1 mM EDTA, 4mM dithiothreitol, 0.1% Nonidet P-40, and 3 mM MgCl₂. It is believed that by adopting a hairpin conformation, the oligonucleotide analogue better mimics the double-stranded DNA substrate for DNA methyltransferase than a single-stranded oligonucleotide, thus inhibiting the activity of DNA methyltransferase more effectively.


[00113] In one embodiment, the oligonucleotide analogue has the following general secondary structure:






wherein N is any nucleotide; N' is a nucleotide complementary to N; Z is 5-aza-cytosine as a base residue; G is guanine as a base residue; l, n, or m is an integer; nucleotide N_n, N_m, N'_n, and N'_m are positioned in the stem region of the hairpin; and N_l is positioned in the loop region of the hairpin. Preferably, l, n, or m is an integer greater than 2, 3, 4, or 5. Optionally, l is 2, 3, 4, 5, or 6. Also optionally, if N_n, N_m, or N_l has one or more cytosine residues, the cytosine residue is substituted with 5-aza-cytosine.

[00114] In a particular embodiment, the oligonucleotide analogue has the following general secondary structure:



wherein C' is 5-methyl-cytidine.

[00115] In another particular embodiment, the oligonucleotide analogue (SEQ ID NO:31) has the following hairpin conformation:



wherein D is decitabine A is adenosine or 2'-deoxyadenosine, T is thymidine or 2'-deoxythymidine, and C at nucleotide position 21 is optionally substituted with 5-methyl-2'-deoxycytidine.

[00116] In any of above embodiments, the oligonucleotide analogue can be single-stranded or double-stranded. When the oligonucleotide analogue is double-stranded, the first strand is the oligonucleotide analogue, and the second strand may be an oligonucleotide with sequence complementary to that of the first strand without the cytosine residue being replaced with 5-aza-cytosine. For example, the first strand may be 5'-TTDGDGAA-3' (SEQ ID NO: 32) wherein D is decitabine; whereas the second strand may be 5'-TTCGCGAA-3' (SEQ ID NO: 33). Optionally, at least one of the cytosine residues in either the first or second strand may be substituted with 5-methyl-cytosine.

[00117] In any of above embodiments, the linker between Z and G residues or between any two of the base residues in the oligonucleotide analogue is preferably a sugar phosphodiester linkage. Preferably the linker is a phosphodiester linkage via 2'-deoxyribose or ribose, as in the natural sugar phosphodiester backbone in DNA and RNA, respectively. Optionally, to enhance the resistance to nuclease degradation in vivo, the natural phosphodiester linker —O-P(=O)(O⁻)-O-CH₂— can be modified to be a phosphorothioate linker —O-P(=O)(S⁻)-O-CH₂—, boranophosphate or methylphosphonate linker; the 2'-hydroxyl group of ribose can be modified to be a 2'-methoxy group, 2'-methoxyethyl group, or 2'-fluoro group. Examples of such oligonucleotide analogues with unnatural backbones are shown Figure 24A where decitabine is linked to guanosine through ribose phosphate backbone. Also optionally, the natural sugar phosphodiester backbone can be replaced with a protein nucleotide (PNA) backbone where the backbone is made from repeating N-(2-aminoethyl)-glycine units linked by peptide bonds. An example of such an oligonucleotide analogue with PNA backbone is shown in Figure 24B where 5-aza-cytosine is linked to guanine via a PNA backbone. Other types of linkers for oligonucleotides designed to be more

PCT/US06/37313

resistant to nuclease degradation than the natural are described US Patent Nos. 6,900, 540 and 6,900,301, which are herein incorporated by reference.

[00118] The oligonucleotide analogues of the present invention may be ones isolated from biological sources, such as tissues, cells and body fluid, and preferably purified to a substantial degree of purity, more preferably of at least 80% purity, and most preferably of at least 95% of purity. The oligonucleotide analogues may also be synthetic ones that are non-naturally occurring oligonucleotide comprising a 5-aza-cytidine, e.g., chemically or enzymatically synthesized in vitro.

[00119] The oligonucleotide analogues of the present invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

[00120] The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotide analogues of the invention may be prepared by forming one or more ester bond with any of the hydroxyl groups in the sugar ring using an organic compound containing a carboxyl group, or as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published Dec. 9, 1993 or in WO 94/26764 and U.S. Pat. No. 5,770,713 to Imbach et al.

[00121] The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

[00122] Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," J. of Pharma Sci., 1977, 66, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention.

[00123] As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic

PCT/US2006/37313

acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

[00124] For oligonucleotide analogues of the present invention, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

[00125] The invention also embraces isolated compounds. An isolated compound refers to a compound which represents at least 10%, preferably 20%, more preferably 50% and most preferably 80% of the compound present in the mixture, and exhibits a detectable (i.e. statistically significant) inhibitory activity of DNA methylation when tested in biological assays such as the combined bisulfite restriction analysis or COBRA (Xiong, Z.; Laird, P. W. Nucleic Acids Res. 1997, 25, 2532-2534) and radiolabeled methyl incorporation assay (Francis, K. T.; Thompson, R. W.; Krumdieck, C. L. Am. J. Clin. Nutr. 1977, 30, 2028-2032).

2. Pharmaceutical Formulations of the Present Invention

[00126] According to the present invention, the oligonucleotide analogues or compounds of the present invention can be formulated into pharmaceutically acceptable compositions for treating various diseases and conditions.

[00127] The pharmaceutically-acceptable compositions of the present invention comprise one or more compounds of the invention in association with one or more nontoxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants and/or excipients, collectively referred to herein as "carrier" materials, and if desired other active ingredients.

[00128] The compounds of the present invention are administered by any route, preferably in the form of a pharmaceutical composition adapted to such a route, as illustrated below and are dependent on the condition being treated. The compounds and compositions can be, for example, administered orally, parenterally, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery (for example by a catheter or stent), subcutaneously, intraadiposally, intraarticularly, or intrathecally.

[00129] The pharmaceutical formulation may optionally further include an excipient added in an amount sufficient to enhance the stability of the composition, maintain the product in solution, or prevent side effects (e.g., potential ulceration, vascular irritation or extravasation) associated with the administration of the inventive formulation. Examples of excipients include, but are not limited to, mannitol, sorbitol, lactose, dextrose, cyclodextrin such as, α -, β -, and γ -cyclodextrin, and modified, amorphous cyclodextrin such as hydroxypropyl-, hydroxyethyl-, glucosyl-,

PCT/US06/37313
maltosyl-, maltotriosyl-, carboxyamidomethyl-, carboxymethyl-, sulfobutylether-, and diethylamino-substituted α , β , and γ -cyclodextrin. Cyclodextrins such as Encapsin® from Janssen Pharmaceuticals or equivalent may be used for this purpose.

[00130] For oral administration, the pharmaceutical compositions can be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a therapeutically-effective amount of the active ingredient. Examples of such dosage units are tablets and capsules. For therapeutic purposes, the tablets and capsules which can contain, in addition to the active ingredient, conventional carriers such as binding agents, for example, acacia gum, gelatin, polyvinylpyrrolidone, sorbitol, or tragacanth; fillers, for example, calcium phosphate, glycine, lactose, maize-starch, sorbitol, or sucrose; lubricants, for example, magnesium stearate, polyethylene glycol, silica, or talc; disintegrants, for example, potato starch, flavoring or coloring agents, or acceptable wetting agents. Oral liquid preparations generally are in the form of aqueous or oily solutions, suspensions, emulsions, syrups or elixirs may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous agents, preservatives, coloring agents and flavoring agents. Examples of additives for liquid preparations include acacia, almond oil, ethyl alcohol, fractionated coconut oil, gelatin, glucose syrup, glycerin, hydrogenated edible fats, lecithin, methyl cellulose, methyl or propyl parahydroxybenzoate, propylene glycol, sorbitol, or sorbic acid.

[00131] For topical use the compounds of the present invention can also be prepared in suitable forms to be applied to the skin, or mucus membranes of the nose and throat, and can take the form of creams, ointments, liquid sprays or inhalants, lozenges, or throat paints. Such topical formulations further can include chemical compounds such as dimethylsulfoxide (DMSO) to facilitate surface penetration of the active ingredient.

[00132] For application to the eyes or ears, the compounds of the present invention can be presented in liquid or semi-liquid form formulated in hydrophobic or hydrophilic bases as ointments, creams, lotions, paints or powders.

[00133] For rectal administration the compounds of the present invention can be administered in the form of suppositories admixed with conventional carriers such as cocoa butter, wax or other glyceride.

[00134] Alternatively, the compounds of the present invention can be in powder form for reconstitution in the appropriate pharmaceutically acceptable carrier at the time of delivery.

[00135] The pharmaceutical compositions can be administered via injection. Formulations for parenteral administration can be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions or suspensions can be prepared from sterile powders or granules having one or more of the carriers mentioned for use in the formulations for oral administration. The compounds can be dissolved in polyethylene glycol, propylene glycol, ethanol, corn oil, benzyl alcohol, sodium chloride, and/or various buffers.

[00136] The oligonucleotide analogues of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μm in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa., 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and

~~PCT/US06/37313~~

dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase.

Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

[00137] Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

[00138] Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199).

Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

[00139] Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

[00140] A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

[00141] Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers

PCT/US06/37313
(for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form

colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

[00142] Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

[00143] The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

[00144] In one embodiment of the present invention, the oligonucleotide analogues are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa., 1985, p. 271).

[00145] The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

~~PCT/US06/37313~~
[00146]

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

[00147] Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., Pharmaceutical Research, 1994, 11, 1385-1390; Ritschel, Meth. Find. Exp. Clin. Pharmacol., 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., Pharmaceutical Research, 1994, 11, 1385; Ho et al., J. Pharm. Sci., 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

[00148] Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories—surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p. 92). Each of these classes has been discussed above.

[00149] Besides microemulsions there are many organized surfactant structures that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

[00150] Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes

PCT/US06/37313

possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages in vivo.

[00151] In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

[00152] Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

[00153] Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

[00154] Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

[00155] Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

[00156] Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).

[00157] One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

[00158] Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome.TM. I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome.TM. II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the

PCT/US2006/037313

dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. S.T.P. Pharma. Sci., 1994, 4, 6, 466).

[00159] Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G_{sub}.M1, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., FEBS Letters, 1987, 223, 42; Wu et al., Cancer Research, 1993, 53, 3765).

[00160] Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (Ann. N.Y. Acad. Sci., 1987, 507, 64) reported the ability of monosialoganglioside G_{sub}.M1, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (Proc. Natl. Acad. Sci. U.S.A., 1988, 85, 6949). U.S. Pat. No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G_{sub}.M1 or a galactocerebroside sulfate ester. U.S. Pat. No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

[00161] Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (Bull. Chem. Soc. Jpn., 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C_{sub}.12 15G, that contains a PEG moiety. Illum et al. (FEBS Lett., 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Pat. Nos. 4,426,330 and 4,534,899). Klivanov et al. (FEBS Lett., 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (Biochimica et Biophysica Acta, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Pat. Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Pat. No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Pat. No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.). Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Pat. Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

[00162] A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Pat. No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Pat. No. 5,665,710 to Rahman et al. describes certain methods of encapsulating

oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

[00163] Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

[00164] Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

[00165] If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

[00166] If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

[00167] If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

[00168] If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

[00169] The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

[00170] In a particular embodiment, the compounds of the present invention can be formulated into a pharmaceutically acceptable composition comprising the compound solvated in non-aqueous solvent that includes glycerin, propylene glycol, polyethylene glycol, or combinations thereof. It is believed that the compounds will be stable in such pharmaceutical formulations so that the pharmaceutical formulations may be stored for a prolonged period of time prior to use.

PCT/US2006/037313
[00171]

In current clinical treatment with decitabine, to minimize drug decomposition decitabine is supplied as lyophilized powder and reconstituted in a cold aqueous solution containing water in at least 40% vol of the solvent, such as WFI, and diluted in cold infusion fluids prior to administration. Such a formulation and treatment regimen suffers from a few drawbacks. First, refrigeration of decitabine in cold solution becomes essential, which is burdensome in handling and economically less desirable than a formulation that can sustain storage at higher temperatures. Second, due to rapid decomposition of decitabine in aqueous solution, the reconstituted decitabine solution may only be infused to a patient for a maximum of 3 hr if the solution has been stored in the refrigerator for less than 7 hr. In addition, infusion of cold fluid can cause great discomfort and pain to the patient, which induces the patient's resistance to such a regimen.

[00172] By modifying the triazine ring and/or the ribose ring of decitabine and by formulating the compound with non-aqueous solvent, the pharmaceutical formulations can circumvent the above-listed problems associated with the current clinical treatment with decitabine. These formulations of the inventive compounds are believed to be more chemically stable than decitabine formulated in aqueous solutions containing water in at least 40% vol. of the solvent.

[00173] In a preferred embodiment, the inventive formulation contains less than 40% water in the solvent, optionally less than 20% water in the solvent, optionally less than 10% water in the solvent, or optionally less than 1% water in the solvent. In one variation, the pharmaceutical formulation is stored in a substantially anhydrous form. Optionally, a drying agent may be added to the pharmaceutical formulation to absorb water.

[00174] Owing to the enhanced stability, the inventive formulation may be stored and transported at ambient temperature, thereby significantly reducing the cost of handling the drug. Further, the inventive formulation may be conveniently stored for a long time before being administered to the patient. In addition, the inventive formulation may be diluted with regular infusion fluid (without chilling) and administered to a patient at room temperature, thereby avoiding causing patients' discomfort associated with infusion of cold fluid.

[00175] In another embodiment, the inventive compound is dissolved in glycerin at different concentrations. For example, the formulation may optionally comprise between 0.1 and 200; between 1 and 100; between 1 and 50; between 2 and 50; between 2 and 100; between 5 and 100; between 10 and 100 or between 20 and 100 mg inventive compound per ml of glycerin. Specific examples of the inventive compound per glycerin concentrations include but are not limited to 2, 5, 10, 20, 22, 25, 30, 40 and 50 mg/ml.

[00176] Different grades of glycerin (synonyms: 1,2,3-propanetriol; glycerol; glycol alcohol; glycerol anhydrous) may be used to prepare the formulations. Preferably, glycerin with chemical purity higher than 90% is used to prepare the formulations.

[00177] In another embodiment, the inventive compound is dissolved in propylene glycol at different concentrations. For example, the formulation may optionally comprise between 0.1 and 200; between 0.1 and 100; between 0.1 and 50; between 2 and 50; between 2 and 100; between 5 and 100; between 10 and 100 or between 20 and 100 mg inventive compound per ml of propylene glycol. Specific examples of decitabine per propylene glycol concentrations include but are not limited to 2, 5, 10, 20, 22, 25, 30, 40 and 50 mg/ml.

[00178] In yet another embodiment, the inventive compound is dissolved in a solvent combining glycerin and propylene glycol at different concentrations. The concentration of propylene glycol in the solvent is between 0.1-99.9%, optionally between 1-90%, between 10-80%, or between 50-70%.

[00179] In yet another embodiment, the inventive compound is dissolved at different concentrations in a solvent combining glycerin and polyethylene glycol (PEG) such as PEG300, PEG400 and PEG1000. The concentration of

PCT/US2006/037313

polyethylene glycol in the solvent is between 0.1-99.9%, optionally between 1-90%, between 10-80%, or between 50-70%.

[00180] In yet another embodiment, the inventive compound is dissolved at different concentrations in a solvent combining propylene glycol, polyethylene glycol and glycerin. The concentration of propylene glycol in the solvent is between 0.1-99.9%, optionally between 1-90%, between 10-60%, or between 20-40%; and the concentration of polyethylene glycol in the solvent is between 0.1-99.9%, optionally between 1-90%, between 10-80%, or between 50-70%.

[00181] It is believed and experimentally proven that addition of propylene glycol can further improve chemical stability, reduce viscosity of the formulations and facilitate dissolution of the inventive compound in the solvent.

[00182] The pharmaceutical formulation may further comprise an acidifying agent added to the formulation in a proportion such that the formulation has a resulting pH between about 4 and 8. The acidifying agent may be an organic acid. Examples of organic acid include, but are not limited to, ascorbic acid, citric acid, tartaric acid, lactic acid, oxalic acid, formic acid, benzene sulphonic acid, benzoic acid, maleic acid, glutamic acid, succinic acid, aspartic acid, diatrizoic acid, and acetic acid. The acidifying agent may also be an inorganic acid, such as hydrochloric acid, sulphuric acid, phosphoric acid, and nitric acid.

[00183] It is believed that adding an acidifying agent to the formulation to maintain a relatively neutral pH (e.g., within pH 4-8) facilitates ready dissolution of the inventive compound in the solvent and enhances long-term stability of the formulation. In alkaline solution, there is a rapid reversible decomposition of decitabine to N-(formylamidino)-N'- β -D-2-deoxyribofuranosylurea, which decomposes irreversibly to form 1- β -D-2'-deoxyribofuranosyl-3-guanylylurea. The first stage of the hydrolytic degradation involves the formation of N-amidinium-N'-(2-deoxy- β -D-erythropentofuranosyl)urea formate (AUF). The second phase of the degradation at an elevated temperature involves formation of guanidine. In acidic solution, N-(formylamidino)-N'- β -D-2-deoxyribofuranosylurea and some unidentified compounds are formed. In strongly acidic solution (at pH <2.2) 5-azacytosine is produced. Thus, maintaining a relative neutral pH may be advantageous for the formulation comprising the analogs and derivatives of decitabine.

[00184] In a variation, the acidifying agent is ascorbic acid at a concentration of 0.01-0.2 mg/ml of the solvent, optionally 0.04-0.1 mg/ml or 0.03-0.07 mg/ml of the solvent.

[00185] The pH of the pharmaceutical formulation may be adjusted to be between pH 4 and pH 8, preferably between pH 5 and pH 7, and more preferably between pH 5.5 and pH 6.8.

[00186] The pharmaceutical formulation is preferably at least 80%, 90%, 95% or more stable upon storage at 25°C for 7, 14, 21, 28 or more days. The pharmaceutical formulation is also preferably at least 80%, 90%, 95% or more stable upon storage at 40°C for 7, 14, 21, 28 or more days.

[00187] In one embodiment, the pharmaceutical formulation of the present invention is prepared by taking glycerin and dissolving the inventive compound in the glycerin. This may be done, for example, by adding the inventive compound to the glycerin or by adding the glycerin to decitabine. By their admixture, the pharmaceutical formulation is formed.

[00188] Optionally, the method further comprises additional steps to increase the rate at which the inventive compound is solvated by the glycerin. Examples of additional steps that may be performed include, but are not limited to, agitation, heating, extension of solvation period, and application of micronized inventive compound and the combinations thereof.

[00189] In one variation, agitation is applied. Examples of agitation include but are not limited to, mechanical agitation, sonication, conventional mixing, conventional stirring and the combinations thereof. For example,

~~PCT/US06/37313~~

mechanical agitation of the formulations may be performed according to manufacturer's protocols by Silverson homogenizer manufactured by Silverson Machines Inc., (East Longmeadow, MA).

[00190] In another variation, heat may be applied. Optionally, the formulations may be heated in a water bath. Preferably, the temperature of the heated formulations may be less than 70°C, more preferably, between 25°C and 40°C. As an example, the formulation may be heated to 37°C.

[00191] In yet another variation, the inventive compound is solvated in glycerin over an extended period of time.

[00192] In yet another variation, a micronized form of the inventive compound may also be employed to enhance solvation kinetics. Optionally, micronization may be performed by a milling process. As an example, micronization may be performed by milling process performed Mastersizer using an Air Jet Mill, manufactured by IncFluid Energy Aljet Inc. (Boise, ID/Telford, PA).

[00193] Optionally, the method further comprises adjusting the pH of the pharmaceutical formulations by commonly used methods. In one variation, pH is adjusted by addition of acid, such as ascorbic acid, or base, such as sodium hydroxide. In another variation, pH is adjusted and stabilized by addition of buffered solutions, such as solution of (Ethylenedinitrilo) tetraacetic acid disodium salt (EDTA). As decitabine is known to be pH-sensitive, adjusting the pH of the pharmaceutical formulations to approximately pH 7 may increase the stability of therapeutic component.

[00194] Optionally, the method further comprises separation of non-dissolved inventive compound from the pharmaceutical formulations. Separation may be performed by any suitable technique. For example, a suitable separation method may include one or more of filtration, sedimentation, and centrifugation of the pharmaceutical formulations. Clogging that may be caused by non-dissolved particles of the inventive compound, may become an obstacle for administration of the pharmaceutical formulations and a potential hazard for the patient. The separation of non-dissolved inventive compound from the pharmaceutical formulations may facilitate administration and enhance safety of the therapeutic product.

[00195] Optionally, the method further comprises sterilization of the pharmaceutical formulations. Sterilization may be performed by any suitable technique. For example, a suitable sterilization method may include one or more of sterile filtration, chemical, irradiation, heat, and addition of a chemical disinfectant to the pharmaceutical formulation.

[00196] As noted, decitabine is unstable in water and hence it may be desirable to reduce the water content of the glycerin used for formulating the inventive compound. Accordingly, prior to the dissolution and/or sterilization step, the glycerin may be dried. Such drying of glycerin or the solution of the inventive compound in glycerin may be achieved by the addition of a pharmaceutically acceptable drying agent to the glycerin. The glycerin or the inventive formulations may be dried, for example by filtering it through a layer comprising a drying agent.

[00197] Optionally, the method may further comprise adding one or more members of the group selected from drying agents, buffering agents, antioxidants, stabilizers, antimicrobials, and pharmaceutically inactive agents. In one variation, antioxidants such as ascorbic acid, ascorbate salts and mixtures thereof may be added. In another variation stabilizers like glycols may be added.

3. Vessels or Kits Containing Inventive Compounds or Formulations

[00198] The pharmaceutical formulations, described in this invention, may be contained in a sterilized vessel such as syringes, vials or ampoules of various sizes and capacities. The sterilized vessel may optionally contain between 1-50 ml, 1-25 ml or 1-20 ml or 1-10 ml of the formulations. Sterilized vessels maintain sterility of the

PCT/US06/37313
pharmaceutical formulations, facilitate transportation and storage, and allow administration of the pharmaceutical formulations without prior sterilization step.

[00199] The present invention also provides a kit for administering the inventive compound to a host in need thereof. In one embodiment, the kit comprises the inventive compound in a solid, preferably powder form, and a non-aqueous diluent that comprises glycerin, propylene glycol, polyethylene glycol, or combinations thereof. Mixing of the solid decitabine and the diluent preferably results in the formation of a pharmaceutical formulation according to the present invention. For example, the kit may comprise a first vessel comprising the inventive compound in a solid form; and a vessel container comprising a diluent that comprises glycerin; wherein adding the diluent to the solid inventive compound results in the formation of a pharmaceutical formulation for administering the inventive compound. Mixing the solid the inventive compound and diluent may optionally form a pharmaceutical formulation that comprises between 0.1 and 200 mg of the inventive compound per ml of the diluent, optionally between 0.1 and 100, between 2 mg and 50 mg, 5 mg and 30 mg, between 10 mg and 25 mg per ml of the solvent.

[00200] According to the embodiment, the diluent is a combination of propylene glycol and glycerin, wherein the concentration of propylene glycol in the solvent is between 0.1-99.9%, optionally between 1-90%, between 10-60%, or between 20-40%.

[00201] Also according to the embodiment, the diluent is a combination of polyethylene glycol and glycerin, wherein the concentration of polyethylene glycol in the solvent is between 0.1-99.9%, optionally between 1-90%, between 10-60%, or between 20-40%.

[00202] Also according to the embodiment, the diluent is a combination of propylene glycol, polyethylene glycol and glycerin, wherein the concentration of propylene glycol in the solvent is between 0.1-99.9%, optionally between 1-90%, between 10-60%, or between 20-40%; and the concentration of polyethylene glycol in the solvent is between 0.1-99.9%, optionally between 1-90%, between 10-60%, or between 20-40%.

[00203] The diluent also optionally comprises 40%, 20%, 10%, 5%, 2% or less water. In one variation, the diluent is anhydrous and may optionally further comprise a drying agent. The diluent may also optionally comprise one or more drying agents, glycols, antioxidants and/or antimicrobials.

[00204] The kit may optionally further include instructions. The instructions may describe how the solid the inventive compound and the diluent should be mixed to form a pharmaceutical formulation. The instructions may also describe how to administer the resulting pharmaceutical formulation to a patient. It is noted that the instructions may optionally describe the administration methods according to the present invention.

[00205] The diluent and the inventive compound may be contained in separate vessels. The vessels may come in different sizes. For example, the vessel may comprise between 1 and 50, 1 and 25, 1 and 20, or 1 and 10 ml of the diluent.

[00206] The pharmaceutical formulations provided in vessels or kits may be in a form that is suitable for direct administration or may be in a concentrated form that requires dilution relative to what is administered to the patient. For example, pharmaceutical formulations described in this invention, may be in a form that is suitable for direct administration via infusion.

[00207] The methods and kits described herein provide flexibility wherein stability and therapeutic effect of the pharmaceutical formulations comprising the inventive compound may be further enhanced or complemented.

4. ~~PCT/US06/37313~~
 Methods for Administering Inventive Compounds/Compositions

[00208] The compounds/formulations of the present invention can be administered by any route, preferably in the form of a pharmaceutical composition adapted to such a route, as illustrated below and are dependent on the condition being treated. The compounds or formulations can be, for example, administered orally, parenterally, topically, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery (for example by catheter or stent), subcutaneously, intraadiposally, intraarticularly, or intrathecally. The compounds and/or compositions according to the invention may also be administered or co-administered in slow release dosage forms.

[00209] The compounds and/or compositions of this invention may be administered or co-administered in any conventional dosage form. Co-administration in the context of this invention is defined to mean the administration of more than one therapeutic agent in the course of a coordinated treatment to achieve an improved clinical outcome. Such co-administration may also be coextensive, that is, occurring during overlapping periods of time.

[00210] The inventive compound or the composition containing the inventive compound may be administered into a host such as a patient at a dose of 0.1-1000 mg/m², optionally 1-200 mg/m², optionally 1-50 mg/m², optionally 1-40 mg/m², optionally 1-30 mg/m², optionally 1-20 mg/m², or optionally 5-30 mg/m².

[00211] For example, the compound/composition of the present invention may be supplied as sterile powder for injection, together with buffering salt such as potassium dihydrogen and pH modifier such as sodium hydroxide. This formulation is preferably stored at 2-8°C, which should keep the drug stable for at least 2 years. This powder formulation may be reconstituted with 10 ml of sterile water for injection. This solution may be further diluted with infusion fluid known in the art, such as 0.9% sodium chloride injection, 5% dextrose injection and lactated ringer's injection. It is preferred that the reconstituted and diluted solutions be used within 4-6 hours for delivery of maximum potency.

[00212] In a preferred embodiment, the inventive compound/composition is administered to a patient by injection, such as subcutaneous injection, bolus i.v. injection, continuous i.v. infusion and i.v. infusion over 1 hour.

Optionally the inventive compound/composition is administered to a patient via an 1-24 hour i.v. infusion per day for 3-5 days per treatment cycle at a dose of 0.1-1000 mg/m² per day, optionally at a dose of 1-100 mg/m² per day, optionally at a dose of 2-50 mg/m² per day, optionally at a dose of 10-30 mg/m² per day, or optionally at a dose of 5-20 mg/m² per day.

[00213] For decitabine or azacitidine, the dosage below 50 mg/m² is considered to be much lower than that used in conventional chemotherapy for cancer. By using such a low dose of the analog/derivative of decitabine or azacitidine, transcriptional activity of genes silenced in the cancer cells by aberrant methylation can be activated to trigger downstream signal transduction, leading to cell growth arrest, differentiation and apoptosis, which eventually results in death of these cancer cells. This low dosage, however, should have less systemic cytotoxic effect on normal cells, and thus have fewer side effects on the patient being treated.

[00214] The pharmaceutical formulations may be co-administered in any conventional form with one or more members selected from the group comprising infusion fluids, therapeutic compounds, nutritious fluids, anti-microbial fluids, buffering and stabilizing agents.

[00215] As described above, the inventive compounds can be formulated in a liquid form by solvating the inventive compound in a non-aqueous solvent such as glycerin. The pharmaceutical liquid formulations provide the further advantage of being directly administrable, (e.g., without further dilution) and thus can be stored in a stable form until administration. Further, because glycerin can be readily mixed with water, the formulations can be easily and

readily further diluted just prior to administration. For example, the pharmaceutical formulations can be diluted with water 180, 60, 40, 30, 20, 10, 5, 2, 1 minute or less before administration to a patient.

[00216] Patients may receive the pharmaceutical formulations intravenously. The preferred route of administration is by intravenous infusion. Optionally, the pharmaceutical formulations of the current invention may be infused directly, without prior reconstitution.

[00217] In one embodiment, the pharmaceutical formulation is infused through a connector, such as a Y site connector, that has three arms, each connected to a tube. As an example, Baxter® Y-connectors of various sizes can be used. A vessel containing the pharmaceutical formulation is attached to a tube further attached to one arm of the connector. Infusion fluids, such as 0.9% sodium chloride, or 5% dextrose, or 5% glucose, or Lactated Ringer's, are infused through a tube attached to the other arm of the Y-site connector. The infusion fluids and the pharmaceutical formulations are mixed inside the Y site connector. The resulting mixture is infused into the patient through a tube connected to the third arm of the Y site connector. The advantage of this administration approach over the prior art is that the inventive compound is mixed with infusion fluids before it enters the patient's body, thus reducing the time when decomposition of the inventive compound may occur due to contact with water. For example, the inventive compound is mixed less than 10, 5, 2 or 1 minutes before entering the patient's body.

[00218] Patients may be infused with the pharmaceutical formulations for 1, 2, 3, 4, 5 or more hours, as a result of the enhanced stability of the formulations. Prolonged periods of infusion enable flexible schedules of administration of therapeutic formulations.

[00219] Alternatively or in addition, speed and volume of the infusion can be regulated according to the patient's needs. The regulation of the infusion of the pharmaceutical formulations can be performed according to existing protocols.

[00220] The pharmaceutical formulations may be co-infused in any conventional form with one or more member selected from the group comprising infusion fluids, therapeutic compounds, nutritious fluids, anti-microbial fluids, buffering and stabilizing agents. Optionally, therapeutic components including, but are not limited to, anti-neoplastic agents, alkylating agents, agents that are members of the retinoids superfamily, antibiotic agents, hormonal agents, plant-derived agents, biologic agents, interleukins, interferons, cytokines, immuno-modulating agents, and monoclonal antibodies, may be co-infused with the inventive formulations.

[00221] Co-infusion in the context of this invention is defined to mean the infusion of more than one therapeutic agents in a course of coordinated treatment to achieve an improved clinical outcome. Such co-infusion may be simultaneous, overlapping, or sequential. In one particular example, co-infusion of the pharmaceutical formulations and infusion fluids may be performed through Y-type connector.

[00222] The pharmacokinetics and metabolism of intravenously administered the pharmaceutical formulations resemble the pharmacokinetics and metabolism of intravenously administered the inventive compound.

[00223] In humans, decitabine displayed a distribution phase with a half-life of 7 minutes and a terminal half-life on the order of 10-35 minutes as measured by bioassay. The volume of distribution is about 4.6 L/kg. The short plasma half-life is due to rapid inactivation of decitabine by deamination by liver cytidine deaminase. Clearance in humans is high, on the order of 126 mL/min/kg. The mean area under the plasma curve in a total of 5 patients was 408 $\mu\text{g}/\text{h}/\text{L}$ with a peak plasma concentration of 2.01 μM . In patients decitabine concentrations were about 0.4 $\mu\text{g}/\text{mL}$ (2 μM) when administered at 100 mg/m^2 as a 3-hour infusion. During a longer infusion time (up to 40 hours) plasma concentration was about 0.1 to 0.4 $\mu\text{g}/\text{mL}$. With infusion times of 40-60 hours, at an infusion rate of 1 $\text{mg}/\text{kg}/\text{h}$, plasma concentrations of 0.43-0.76 $\mu\text{g}/\text{mL}$ were achieved. The steady-state plasma concentration at an infusion rate of 1 $\text{mg}/\text{kg}/\text{h}$ is estimated to be 0.2-0.5 $\mu\text{g}/\text{mL}$. The half-life after discontinuing the infusion is 12-20

PCT/US2006/037313

min. The steady-state plasma concentration of decitabine was estimated to be 0.31-0.39 $\mu\text{g/mL}$ during a 6-hour infusion of 100 mg/m^2 . The range of concentrations during a 600 mg/m^2 infusion was 0.41-16 $\mu\text{g/mL}$. Penetration of decitabine into the cerebrospinal fluid in man reaches 14-21% of the plasma concentration at the end of a 36-hour intravenous infusion. Urinary excretion of unchanged decitabine is low, ranging from less than 0.01% to 0.9% of the total dose, and there is no relationship between excretion and dose or plasma drug levels. High clearance values and a total urinary excretion of less than 1% of the administered dose suggest that decitabine is eliminated rapidly and largely by metabolic processes.

[00224] Owing to their enhanced stability in comparison with decitabine, the inventive compounds/compositions can enjoy longer shelf life when stored and circumvent problems associated with clinical use of decitabine. For example, the inventive compounds may be supplied as lyophilized powder, optionally with an excipient (e.g., cyclodextrin), acid (e.g., ascorbic acid), alkaline (sodium hydroxide), or buffer salt (monobasic potassium dihydrogen phosphate). The lyophilized powder can be reconstituted with sterile water for injection, e.g., i.v., i.p., i.m., or subcutaneously. Optionally, the powder can be reconstituted with aqueous or non-aqueous solvent comprising a water miscible solvent such as glycerin, propylene glycol, ethanol and PEG. The resulting solution may be administered directly to the patient, or diluted further with infusion fluid, such as 0.9% Sodium Chloride; 5% Dextrose; 5% Glucose; and Lactated Ringer's infusion fluid.

[00225] The inventive compounds/compositions may be stored under ambient conditions or in a controlled environment, such as under refrigeration (2-8°C; 36-46°F). Due to their superior stability in comparison with decitabine, the inventive compounds/compositions can be stored at room temperature, reconstituted with injection fluid, and administered to the patient without prior cooling of the drug solution.

[00226] In addition, due to their enhanced chemical stability, the inventive compound/composition should have a longer plasma half-life compared to that of decitabine. Thus, the inventive compound/composition may be administered to the patient at a lower dose and/or less frequently than that for decitabine.

5. Combination Therapy with Inventive Pharmaceutical Compositions

[00227] The compounds or pharmaceutical formulations of the present invention may be used in conjunction with inhibitors of histone deacetylase (HDAC) to further modulate transcription of genes, e.g., to reestablish transcription of genes silenced by hypermethylation and acetylation of histones, in a synergistic manner.

[00228] HDAC plays important roles in transcription silencing of genes. The amount of acetylation on the histones is controlled by the opposing activities of two types of enzymes, histone acetyl transferase (HATs) and histone deacetylases (HDACs). Substrates for these enzymes include ϵ -amino groups of lysine residues located in the amino-terminal tails of the histones H3, H4, H2A, and H2B. These amino acid residues are acetylated by HATs and deacetylated by HDACs. With the removal of the acetyl groups from the histone lysine by HDACs, a positive charge is restored to the lysine residue, thereby condensing the structure of nucleosome and silencing the genes contained within. Thus, to activate these genes silenced by deacetylase of histones, the activity of HDACs should be inhibited. With the inhibition of HDAC, histones are acetylated and the DNA that is tightly wrapped around a deacetylated histone core relaxes. The opening of DNA conformation leads to expression of specific genes.

[00229] In addition to deacetylation of histones, HDACs may also regulated gene expression by deacetylating transcription factors, such as p53 (a tumor suppressor gene), GATA-1, TFIIIE, and TFIIF. Gu and Roeder (1997) Cell 90:595-606 (p53); and Boyes et al. (1998) Nature 396:594-598 (GATA-1). HDACs also participate in cell cycle regulation, for example, by transcription repression which is mediated by RB tumor suppressor proteins recruiting HDACs. Brehm et al. (1998) Nature 391:597-601. Thus, inhibition of HDACs should activate expression

PCT/US06/37313

of tumor suppressor genes such as p53 and RB and as a result promote cell growth arrest, differentiation and apoptosis induced by these genes.

[00230] As described above, aberrant transcriptional silencing of a number of genes, such as tumor suppressor genes, is directly related to pathogenesis of cancer and other diseases. Methylation of cytosine residues in DNA and removal of acetyl groups from histones are the two primary mechanisms for gene silencing. Due to methylation and/or histone deacetylase of cancer-related genes, expression of these genes is suppressed or completely silenced. Meanwhile, expression of these genes is required for induction of growth arrest, differentiation, and/or apoptotic cell death of transformed cells. Inaction of these genes in the transformed cells leads to uncontrolled proliferation of these cells, which eventually results in cancer.

[00231] By combining the inventive compounds/compositions with HDAC inhibitors, genes required for induction of growth arrest, differentiation and cell death of transformed cells can be reactivated effectively. The inventive compounds/compositions inhibit methylation of DNA for the genes, especially in the regulatory region, thus resulting in activation of transcription of the gene. Meanwhile, HDAC inhibitors inhibit deacetylase of the histones in the nucleosomal core of the gene, thus resulting in net increase of the acetylation of histones, which, in turn, activates transcription of the gene. By exploiting these two complementary mechanisms, the combination therapy may reestablish gene transcription more effectively and, ideally, in a synergistic manner. A combination therapy having synergistic effects should require a less amount of each inhibitor than it being used alone, thus reducing potential side effects associated systemic administration of high dosages of the inhibitors and improving therapeutic index.

[00232] Many anticancer agents exert their anti-cancer effects by triggering signal transduction cascades involving proteins encoded by these tumor suppressor genes. With insufficient expression of these genes in cancer cells, the anti-cancer effects of these anti-neoplastic agents may be severely reduced or completely eradicated. Through reactivation or re-expression of these genes that are epigenetically silenced by DNA methylation and histone deacetylase, the intrinsic defense mechanisms of the body are mobilized to combat the disease by restoration of the tumor-suppressing functions to cancer cells in response to signals sent by the anti-cancer agent administered. Such stimulation of the intrinsic tumor suppressing functions of the body should lead to the requirement of lower dosage of the anticancer agent, thus resulting in a higher therapeutic index (i.e., greater efficacy and lower toxicity) of the agent.

[00233] Inhibitors of HDACs include, but are not limited to, the following structural classes: 1) hydroxamic acids, 2) cyclic peptides, 3) benzamides, and 4) short-chain fatty acids.

[00234] Examples of hydroxamic acids and hydroxamic acid derivatives, but are not limited to, trichostatin A (TSA), suberoylanilide hydroxamic acid (SAHA), oxamflatin, suberic bishydroxamic acid (SBHA), m-carboxy-cinnamic acid bishydroxamic acid (CBHA), and pyroxamide. TSA was isolated as an antifungi antibiotic (Tsuji et al (1976) J. Antibiot (Tokyo) 29:1-6) and found to be a potent inhibitor of mammalian HDAC (Yoshida et al. (1990) J. Biol. Chem. 265:17174-17179). The finding that TSA-resistant cell lines have an altered HDAC evidences that this enzyme is an important target for TSA. Other hydroxamic acid-based HDAC inhibitors, SAHA, SBHA, and CBHA are synthetic compounds that are able to inhibit HDAC at micromolar concentration or lower in vitro or in vivo. Glick et al. (1999) Cancer Res. 59:4392-4399. These hydroxamic acid-based HDAC inhibitors all possess an essential structural feature: a polar hydroxamic terminal linked through a hydrophobic methylene spacer (e.g. 6 carbon at length) to another polar site which is attached to a terminal hydrophobic moiety (e.g., benzene ring). Compounds developed having such essential features also fall within the scope of the hydroxamic acids that may be used as HDAC inhibitors.

PCT/US2006/037313
[00235]

Cyclic peptides used as HDAC inhibitors are mainly cyclic tetrapeptides. Examples of cyclic peptides include, but are not limited to, trapoxin A, apicidin and FR901228. Trapoxin A is a cyclic tetrapeptide that contains a 2-amino-8-oxo-9,10-epoxy-decanoyl (AOE) moiety. Kijima et al. (1993) J. Biol. Chem. 268:22429-22435. Apicidin is a fungal metabolite that exhibits potent, broad-spectrum antiprotozoal activity and inhibits HDAC activity at nanomolar concentrations. Darkin-Rattray et al. (1996) Proc. Natl. Acad. Sci. USA. 93:13143-13147. FR901228 is a depsipeptide that is isolated from *Chromobacterium violaceum*, and has been shown to inhibit HDAC activity at micromolar concentrations.

[00236] Examples of benzamides include but are not limited to MS-27-275. Saito et al. (1990) Proc. Natl. Acad. Sci. USA. 96:4592-4597. Examples of short-chain fatty acids include but are not limited to butyrates (e.g., butyric acid, arginine butyrate and phenylbutyrate (PB)). Newmark et al. (1994) Cancer Lett. 78:1-5; and Carducci et al. (1997) Anticancer Res. 17:3972-3973. In addition, depudecin which has been shown to inhibit HDAC at micromolar concentrations (Kwon et al. (1998) Proc. Natl. Acad. Sci. USA. 95:3356-3361) also falls within the scope of histone deacetylase inhibitor of the present invention.

[00237] The compounds or pharmaceutical formulations of the present invention may also be used in conjunction with other therapeutic components including but not limiting to anti-neoplastic agents, alkylating agents, agents that are members of the retinoids superfamily, antibiotic agents, hormonal agents, plant-derived agents, biologic agents, interleukins, interferons, cytokines, immuno-modulating agents, and monoclonal antibodies.

[00238] In one embodiment, an alkylating agent is used in combination with and/or added to the inventive compound/formulation. Examples of alkylating agents include, but are not limited to bischloroethylamines (nitrogen mustards, e.g. chlorambucil, cyclophosphamide, ifosfamide, mechlorethamine, melphalan, uracil mustard), aziridines (e.g. thiotepa), alkyl alkane sulfonates (e.g. busulfan), nitrosoureas (e.g. carmustine, lomustine, streptozocin), nonclassic alkylating agents (altretamine, dacarbazine, and procarbazine), platinum compounds (carboplatin and cisplatin).

[00239] In another embodiment, cisplatin, carboplatin or cyclophosphamide is used in combination with and/or added to the inventive compound/formulation.

[00240] In another embodiment, a member of the retinoids superfamily is used in combination with and/or added to the inventive compound/formulation. Retinoids are a family of structurally and functionally related molecules that are derived or related to vitamin A (all-trans-retinol). Examples of retinoid include, but are not limited to, all-trans-retinol, all-trans-retinoic acid (tretinoin), 13-cis retinoic acid (isotretinoin) and 9-cis-retinoic acid.

[00241] In yet another embodiment, a hormonal agent is used in combination with and/or added to the inventive compound/formulation. Examples of such a hormonal agent are synthetic estrogens (e.g. diethylstilbestrol), antiestrogens (e.g. tamoxifen, toremifene, flouxymesterol and raloxifene), antiandrogens (bicalutamide, nilutamide, flutamide), aromatase inhibitors (e.g., aminoglutethimide, anastrozole and tetrazole), ketoconazole, goserelin acetate, leuprolide, megestrol acetate and mifepristone.

[00242] In yet another embodiment, a plant-derived agent is used in combination with and/or added to the inventive compound/formulation. Examples of plant-derived agents include, but are not limited to, vinca alkaloids (e.g., vincristine, vinblastine, vindesine, vinzolidine and vinorelbine), camptothecin (20(S)-camptothecin, 9-nitro-20(S)-camptothecin, and 9-amino-20(S)-camptothecin), podophyllotoxins (e.g., etoposide (VP-16) and teniposide (VM-26)), and taxanes (e.g., paclitaxel and docetaxel).

[00243] In yet another embodiment, a biologic agent is used in combination with and/or added to the inventive compound/formulation, such as immuno-modulating proteins such as cytokines, monoclonal antibodies against tumor antigens, tumor suppressor genes, and cancer vaccines.

PCT/US06/37313

[00244] Examples of interleukins that may be used in combination with and/or added to the inventive

compound/formulation include, but are not limited to, interleukin 2 (IL-2), and interleukin 4 (IL-4), interleukin 12 (IL-12). Examples of interferons that may be used in conjunction with decitabine - glycerin formulations include, but are not limited to, interferon α , interferon β (fibroblast interferon) and interferon γ (fibroblast interferon).

- 5 Examples of such cytokines include, but are not limited to erythropoietin (epoietin), granulocyte-CSF (filgrastim), and granulocyte, macrophage-CSF (sargramostim). Immuno-modulating agents other than cytokines include, but are not limited to bacillus Calmette-Guerin, levamisole, and octreotide.

[00245] Example of monoclonal antibodies against tumor antigens that can be used in conjunction with the inventive formulations include, but are not limited to, HERCEPTIN® (Trastuzumab), RITUXAN® (Rituximab),
10 MYLOTARG® (anti-CD33), and CAMPATH® (anti-CD52).

6. Indications for Compounds or Pharmaceutical Compositions of the Present Invention

[00246] The pharmaceutical formulations according to the present invention may be used to treat a wide variety of diseases that are sensitive to the treatment with decitabine.

- 15 [00247] Preferable indications that may be treated using the pharmaceutical formulations of the present invention include those involving undesirable or uncontrolled cell proliferation. Such indications include benign tumors, various types of cancers such as primary tumors and tumor metastasis, restenosis (e.g. coronary, carotid, and cerebral lesions), hematological disorders, abnormal stimulation of endothelial cells (atherosclerosis), insults to body tissue due to surgery, abnormal wound healing, abnormal angiogenesis, diseases that produce fibrosis of tissue,
20 repetitive motion disorders, disorders of tissues that are not highly vascularized, and proliferative responses associated with organ transplants.

[00248] Generally, cells in a benign tumor retain their differentiated features and do not divide in a completely uncontrolled manner. A benign tumor is usually localized and nonmetastatic. Specific types benign tumors that can be treated using the present invention include hemangiomas, hepatocellular adenoma, cavernous haemangioma,
25 focal nodular hyperplasia, acoustic neuromas, neurofibroma, bile duct adenoma, bile duct cystadenoma, fibroma, lipomas, leiomyomas, mesotheliomas, teratomas, myxomas, nodular regenerative hyperplasia, trachomas and pyogenic granulomas.

[00249] In a malignant tumor cells become undifferentiated, do not respond to the body's growth control signals, and multiply in an uncontrolled manner. The malignant tumor is invasive and capable of spreading to distant sites
30 (metastasizing). Malignant tumors are generally divided into two categories: primary and secondary. Primary tumors arise directly from the tissue in which they are found. A secondary tumor, or metastasis, is a tumor which is originated elsewhere in the body but has now spread to a distant organ. The common routes for metastasis are direct growth into adjacent structures, spread through the vascular or lymphatic systems, and tracking along tissue planes and body spaces (peritoneal fluid, cerebrospinal fluid, etc.)

35 [00250] Specific types of cancers or malignant tumors, either primary or secondary, that can be treated using this invention include breast cancer, skin cancer, bone cancer, prostate cancer, liver cancer, lung cancer, brain cancer, cancer of the larynx, gall bladder, pancreas, rectum, parathyroid, thyroid, adrenal, neural tissue, head and neck, colon, stomach, bronchi, kidneys, basal cell carcinoma, squamous cell carcinoma of both ulcerating and papillary type, metastatic skin carcinoma, osteo sarcoma, Ewing's sarcoma, veticulum cell sarcoma, myeloma, giant cell
40 tumor, small-cell lung tumor, gallstones, islet cell tumor, primary brain tumor, acute and chronic lymphocytic and granulocytic tumors, hairy-cell tumor, adenoma, hyperplasia, medullary carcinoma, pheochromocytoma, mucosal neuromas, intestinal ganglioneuromas, hyperplastic corneal nerve tumor, marfanoid habitus tumor, Wilm's tumor,

PCT/US2006/037313
seminoma, ovarian tumor, leiomyoma tumor, cervical dysplasia and in situ carcinoma, neuroblastoma, retinoblastoma, soft tissue sarcoma, malignant carcinoid, topical skin lesion, mycosis fungoide, rhabdomyosarcoma, Kaposi's sarcoma, osteogenic and other sarcoma, malignant hypercalcemia, renal cell tumor, polycythemia vera, adenocarcinoma, glioblastoma multiforma, leukemias, lymphomas, malignant melanomas, epidermoid carcinomas, and other carcinomas and sarcomas.

[00251] Hematologic disorders include abnormal growth of blood cells which can lead to dysplastic changes in blood cells and hematologic malignancies such as various leukemias. Examples of hematologic disorders include but are not limited to acute myeloid leukemia, acute promyelocytic leukemia, acute lymphoblastic leukemia, chronic myelogenous leukemia, the myelodysplastic syndromes, and sickle cell anemia.

[00252] Acute myeloid leukemia (AML) is the most common type of acute leukemia that occurs in adults. Several inherited genetic disorders and immunodeficiency states are associated with an increased risk of AML. These include disorders with defects in DNA stability, leading to random chromosomal breakage, such as Bloom's syndrome, Fanconi's anemia, Li-Fraumeni kindreds, ataxia-telangiectasia, and X-linked agammaglobulinemia.

[00253] Acute promyelocytic leukemia (APML) represents a distinct subgroup of AML. This subtype is characterized by promyelocytic blasts containing the 15;17 chromosomal translocation. This translocation leads to the generation of the fusion transcript comprised of the retinoic acid receptor and a sequence PML.

[00254] Acute lymphoblastic leukemia (ALL) is a heterogeneous disease with distinct clinical features displayed by various subtypes. Reoccurring cytogenetic abnormalities have been demonstrated in ALL. The most common cytogenetic abnormality is the 9;22 translocation. The resultant Philadelphia chromosome represents poor prognosis of the patient.

[00255] Chronic myelogenous leukemia (CML) is a clonal myeloproliferative disorder of a pluripotent stem cell. CML is characterized by a specific chromosomal abnormality involving the translocation of chromosomes 9 and 22, creating the Philadelphia chromosome. Ionizing radiation is associated with the development of CML.

[00256] The myelodysplastic syndromes (MDS) are heterogeneous clonal hematopoietic stem cell disorders grouped together because of the presence of dysplastic changes in one or more of the hematopoietic lineages including dysplastic changes in the myeloid, erythroid, and megakaryocytic series. These changes result in cytopenias in one or more of the three lineages. Patients afflicted with MDS typically develop complications related to anemia, neutropenia (infections), or thrombocytopenia (bleeding). Generally, from about 10% to about 70% of patients with MDS develop acute leukemia.

[00257] Treatment of abnormal cell proliferation due to insults to body tissue during surgery may be possible for a variety of surgical procedures, including joint surgery, bowel surgery, and cheloid scarring. Diseases that produce fibrotic tissue include emphysema. Repetitive motion disorders that may be treated using the present invention include carpal tunnel syndrome. An example of cell proliferative disorders that may be treated using the invention is a bone tumor.

[00258] The proliferative responses associated with organ transplantation that may be treated using this invention include those proliferative responses contributing to potential organ rejections or associated complications. Specifically, these proliferative responses may occur during transplantation of the heart, lung, liver, kidney, and other body organs or organ systems.

[00259] Abnormal angiogenesis that may be may be treated using this invention include those abnormal angiogenesis accompanying rheumatoid arthritis, ischemic-reperfusion related brain edema and injury, cortical ischemia, ovarian hyperplasia and hypervascularity, (polycystic ovary syndrome), endometriosis, psoriasis, diabetic

WO 2007/041071
PCT/US2006/37313
retinopathy, and other ocular angiogenic diseases such as retinopathy of prematurity (retrolental fibroplastic), muscular degeneration, corneal graft rejection, neurovascular glaucoma and Oster Webber syndrome.

[00260] Diseases associated with abnormal angiogenesis require or induce vascular growth. For example, corneal angiogenesis involves three phases: a pre-vascular latent period, active neovascularization, and vascular maturation and regression. The identity and mechanism of various angiogenic factors, including elements of the inflammatory response, such as leukocytes, platelets, cytokines, and eicosanoids, or unidentified plasma constituents have yet to be revealed.

[00261] In another embodiment, the pharmaceutical formulations of the present invention may be used for treating diseases associated with undesired or abnormal angiogenesis. The method comprises administering to a patient suffering from undesired or abnormal angiogenesis the pharmaceutical formulations of the present invention alone, or in combination with anti-neoplastic agent whose activity as an anti-neoplastic agent in vivo is adversely affected by high levels of DNA methylation. The particular dosage of these agents required to inhibit angiogenesis and/or angiogenic diseases may depend on the severity of the condition, the route of administration, and related factors that can be decided by the attending physician. Generally, accepted and effective daily doses are the amount sufficient to effectively inhibit angiogenesis and/or angiogenic diseases.

[00262] According to this embodiment, the pharmaceutical formulations of the present invention may be used to treat a variety of diseases associated with undesirable angiogenesis such as retinal/choroidal neovascularization and corneal neovascularization. Examples of retinal/choroidal neovascularization include, but are not limited to, Best's diseases, myopia, optic pits, Stargardt's diseases, Paget's disease, vein occlusion, artery occlusion, sickle cell anemia, sarcoid, syphilis, pseudoxanthoma elasticum carotid obstructive diseases, chronic uveitis/vitritis, mycobacterial infections, Lyme's disease, systemic lupus erythematosus, retinopathy of prematurity, Eales disease, diabetic retinopathy, macular degeneration, Behcet's diseases, infections causing a retinitis or chorioiditis, presumed ocular histoplasmosis, pars planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, trauma and post-laser complications, diseases associated with rubeosis (neovascularization of the angle) and diseases caused by the abnormal proliferation of fibrovascular or fibrous tissue including all forms of proliferative vitreoretinopathy. Examples of corneal neovascularization include, but are not limited to, epidemic keratoconjunctivitis, Vitamin A deficiency, contact lens overwear, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, Sjögren's, acne rosacea, phlyctenulosis, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, Mooren ulcer, Terrien's marginal degeneration, marginal keratolysis, polyarteritis, Wegener sarcoidosis, Scleritis, periphthoid radial keratotomy, neovascular glaucoma and retrolental fibroplasia, syphilis, Mycobacteria infections, lipid degeneration, chemical burns, bacterial ulcers, fungal ulcers, Herpes simplex infections, Herpes zoster infections, protozoan infections and Kaposi sarcoma.

[00263] In yet another embodiment, the pharmaceutical formulations of the present invention may be used for treating chronic inflammatory diseases associated with abnormal angiogenesis. The method comprises administering to a patient suffering from a chronic inflammatory disease associated with abnormal angiogenesis the pharmaceutical formulations of the present invention alone, or in combination with an anti-neoplastic agent whose activity as an anti-neoplastic agent in vivo is adversely affected by high levels of DNA methylation. The chronic inflammation depends on continuous formation of capillary sprouts to maintain an influx of inflammatory cells. The influx and presence of the inflammatory cells produce granulomas and thus, maintains the chronic inflammatory state. Inhibition of angiogenesis using the pharmaceutical formulations of the present invention may prevent the formation of the granulomas, thereby alleviating the disease. Examples of chronic inflammatory disease include,

~~PCT/US06/37313~~
but are not limited to, inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, psoriasis, sarcoidosis, and rheumatoid arthritis.

[00264] Inflammatory bowel diseases such as Crohn's disease and ulcerative colitis are characterized by chronic inflammation and angiogenesis at various sites in the gastrointestinal tract. For example, Crohn's disease occurs as a chronic transmural inflammatory disease that most commonly affects the distal ileum and colon but may also occur in any part of the gastrointestinal tract from the mouth to the anus and perianal area. Patients with Crohn's disease generally have chronic diarrhea associated with abdominal pain, fever, anorexia, weight loss and abdominal swelling. Ulcerative colitis is also a chronic, nonspecific, inflammatory and ulcerative disease arising in the colonic mucosa and is characterized by the presence of bloody diarrhea. These inflammatory bowel diseases are generally caused by chronic granulomatous inflammation throughout the gastrointestinal tract, involving new capillary sprouts surrounded by a cylinder of inflammatory cells. Inhibition of angiogenesis by the pharmaceutical formulations of the present invention should inhibit the formation of the sprouts and prevent the formation of granulomas. The inflammatory bowel diseases also exhibit extra intestinal manifestations, such as skin lesions. Such lesions are characterized by inflammation and angiogenesis and can occur at many sites other the gastrointestinal tract. Inhibition of angiogenesis by the pharmaceutical formulations of the present invention should reduce the influx of inflammatory cells and prevent the lesion formation.

[00265] Sarcoidosis, another chronic inflammatory disease, is characterized as a multi-system granulomatous disorder. The granulomas of this disease can form anywhere in the body and, thus, the symptoms depend on the site of the granulomas and whether the disease is active. The granulomas are created by the angiogenic capillary sprouts providing a constant supply of inflammatory cells. By using the pharmaceutical formulations of the present invention to inhibit angiogenesis, such granulomas formation can be inhibited. Psoriasis, also a chronic and recurrent inflammatory disease, is characterized by papules and plaques of various sizes. Treatment using the pharmaceutical formulations of the present invention should prevent the formation of new blood vessels necessary to maintain the characteristic lesions and provide the patient relief from the symptoms.

[00266] Rheumatoid arthritis (RA) is also a chronic inflammatory disease characterized by non-specific inflammation of the peripheral joints. It is believed that the blood vessels in the synovial lining of the joints undergo angiogenesis. In addition to forming new vascular networks, the endothelial cells release factors and reactive oxygen species that lead to pannus growth and cartilage destruction. The factors involved in angiogenesis may actively contribute to, and help maintain, the chronically inflamed state of rheumatoid arthritis. Treatment using the pharmaceutical formulations of the present invention alone or in conjunction with other anti-RA agents may prevent the formation of new blood vessels necessary to maintain the chronic inflammation and provide the RA patient relief from the symptoms.

[00267] In yet another embodiment, the pharmaceutical formulations of the present invention may be used for treating diseases associated with abnormal hemoglobin synthesis. The method comprises administering the pharmaceutical formulations of the present invention to a patient suffering from disease associated with abnormal hemoglobin synthesis. Decitabine-containing formulations stimulate fetal hemoglobin synthesis because the mechanism of incorporation into DNA is associated with DNA hypomethylation. Examples of diseases associated with abnormal hemoglobin synthesis include, but are not limited to, sickle cell anemia and β -thalassemia.

[00268] In yet another embodiment, the pharmaceutical formulations of the present invention may be used to control intracellular gene expression. The method comprises administering the pharmaceutical formulations of the present invention to a patient suffering from disease associated with abnormal levels of gene expression. DNA methylation is associated with the control of gene expression. Specifically, methylation in or near promoters inhibit

PCT/US06/37313

transcription while demethylation restores expression. Examples of the possible applications of the described mechanisms include, but are not limited to, therapeutically modulated growth inhibition, induction of apoptosis, and cell differentiation.

[00269] Gene activation facilitated by the pharmaceutical formulations of the present invention may induce differentiation of cells for therapeutic purposes. Cellular differentiation is induced through the mechanism of hypomethylation. Examples of morphological and functional differentiation include, but are not limited to differentiation towards formation of muscle cells, myotubes, cells of erythroid and lymphoid lineages.

[00270] Although exemplary embodiments of the present invention have been described and depicted, it will be apparent to the artisan of ordinary skill that a number of changes, modifications, or alterations to the invention as described herein may be made, none of which depart from the spirit of the present invention. All such changes, modifications, and alterations should therefore be seen as within the scope of the present invention.

EXAMPLES

1. Synthesis of phosphoramidite building blocks and 3'-O-capped derivatives

[00271] The present invention also provides effective chemical methods for synthesis of the following novel phosphoramidite building blocks (Figure 2A).

[00272] The 4-amine functional group of 1a can be protected via transformation into various protective groups (R₁), such as carbamates with methyl, ethyl, 9-fluorenylmethyl, 9-(2-sulfo)fluorenylmethyl, 9-(2,7-dibromo)fluorenylmethyl, 17-tetrabenz[a,c,g,i]fluorenylmethyl, 2-chloro-3-indenylmethyl, benz[f]inden-3-ylmethyl, 2,7-di-*tert*-[9-(10,10-dioxo-10,10,10-tetrahydrothioxanthyl)methyl, 1,1-dioxobenzo[b]thiophene-2-ylmethyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-phenylethyl, 1-(1-adamantyl)-1-methylethyl, 2-chloroethyl, 1,1-dimethyl-2-haloethyl, 1,1-dimethyl-2,2-dibromomethyl, 1,1-dimethyl-2,2,2-trichloroethyl, 1-methyl-1-(4-biphenyl)ethyl, 1-(3,5-di-*tert*-butylphenyl)-1-methylethyl, 2-(2'- and 4'-pyridyl)ethyl, 2,2-bis(4'-nitrophenyl)ethyl, *N*-(2-pivaloylamino)-1,1-dimethylethyl, 2-[(2-nitrophenyl)dithio]-1-phenylethyl, 2-(*N,N*-dicyclohexylcarboxamido)ethyl, *t*-butyl, 1-adamantyl, 2-adamantyl, vinyl, allyl, 1-isopropylallyl, cinnamyl, 4-nitrocinnamyl, 3-(3'-pyridyl)prop-2-enyl, 8-quinolyl, *N*-hydroxypiperidinyl, alkyl dithio, benzyl, *p*-methoxybenzyl, *p*-nitrobenzyl, *p*-bromobenzyl, *p*-chlorobenzyl, 2,4-dichlorobenzyl, 4-methylsulfinylbenzyl, 9-anthrylmethyl, diphenylmethyl, 2-methylthioethyl, 2-methylsulfonylethyl, 2-(*p*-toluenesulfonyl)ethyl, [2-(1,3-dithianyl)]methyl, 4-methylthiophenyl, 2,4-dimethylthiophenyl, 2-phosphonioethyl, 1-methyl-1-(triphenylphosphonio)ethyl, 1,1-dimethyl-2-cyanoethyl, 2-dansylethyl, 4-phenylacetoxybenzyl, 4-azidobenzyl, 4-azidomethoxybenzyl, *m*-chloro-*p*-acyloxybenzyl, *p*-(dihydroxyboryl)benzyl, 5-benzisoxazolylmethyl, 2-(trifluoroethyl)-6-chromonylmethyl, *m*-nitrophenyl, 3,5-dimethoxybenzyl, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl, α -methylnitropiperonyl, *o*-nitrophenyl, 3,4-dimethoxy-6-nitrobenzyl, phenyl(*o*-nitrophenyl)ethyl, 2-(2-nitrophenyl)ethyl, 6-nitroveratryl, 4-methoxyphenacyl, 3',5'-dimethoxybenzoin, *t*-amyl, *S*-benzylthio, butynyl, *p*-cyanobenzyl, cyclohexyl, cyclopentyl, cyclopropylmethyl, *p*-decyloxybenzyl, diisopropylmethyl, 2,2-dimethoxycarbonylvinyl, *o*-(*N,N*-dimethylcarboxamido)benzyl, 1,1-dimethyl-3-(*N,N*-dimethylcarboxamido)propyl, 1,1-dimethylpropynyl, 2-furanylmethyl, 2-iodoethyl, isobornyl, isobutyl, isonicotinyl, *p*-(*p*'-methoxyphenylazo)benzyl, 1-methylcyclobutyl, 1-methylcyclohexyl, 1-methyl-1-cyclopropylmethyl, 1-methyl-1-(*p*-phenylazophenyl)ethyl, 1-methyl-1-phenylethyl, 1-methyl-1-(4'-pyridyl)ethyl, phenyl, *p*-(phenylazo)benzyl, 2,4,6-tri-*t*-butylphenyl, 4-(trimethylammonium)benzyl, 2,4,6-trimethylbenzyl; ureas with phenothiazinyl-(10)-carbonyl, *N'*-*p*-toluenesulfonylamino carbonyl, *N'*-phenylaminothiocarbonyl; amides such as formamide, acetamide, phenoxyacetamide, trichloroacetamide,

PCT/US06/37313

trifluoroacetamide, phenylacetamide, 3-phenylpropanamide, pent-4-enamide, *o*-nitrophenylacetamide, *o*-

nitrophenoxycetamide, 3-(*o*-nitrophenyl)propanamide, 2-methyl-2-(*o*-nitrophenoxy)propanamide, 3-methyl-3-

nitrobutanamide, *o*-nitrocinnamide, 3-(4-*t*-butyl-2,6-dinitrophenyl)-2,2-dimethylpropanamide, *o*-

(benzoyloxymethyl)benzamide, 2-[(*t*-butyldiphenylsiloxy)methyl]methylbenzamide, 3-(3',6'-dioxo-2',4',5'-

trimethylcyclohexa-1',4'-diene)-3,3-dimethylpropionamide, *o*-hydroxy-*trans*-cinnamide, acetoacetamide, *p*-

toluenesulfonamide, and benzenesulfonamide. The 4-*O*-methoxy (1b) and 4-*S*-methylthio (1c) analogs of decitabine

can be obtained by modifying the published procedure for decitabine synthesis by separating the α and β anomers of 1-(2-deoxy-3,5-di-*O*-*p*-chlorobenzoyl or benzoyl-*D*-ribofuranosyl)-4-methoxy or methylthio-1,3,5-triazin-2(*H*)-one and removing the 3,5-protective groups without treatment with methanolic ammonia. Pliml and Sorm (1964)

Collect. Czech. Chem. Commun. 29: 2576-2577; Piskala and Sorm (1978) Nucleic Acid Chemistry (by Townsend and Tipson, Wiley, 1978), pp.443-449.

[00273] Protection of the 5'-OH is achieved by dissolving the 4-amino protected decitabine and 4-methoxy and 4-methylthio analogs (1b, 1c) in anhydrous pyridine (5mL/mmol) before adding dimethoxytrityl chloride (1.1 equivalents).

[00274] For example, decitabine (1.2 g) was twice co-distilled with anhydrous pyridine and dissolved in 20 ml dry DMF. Hexamethyldisilazane (2.8 mL) was added. The solution was stirred and left overnight. The solvent was evaporated *in vacuo*, and the remaining residue was dissolved in toluene and evaporated twice. The 3',5'-di trimethylsilyl 5-aza-2'-deoxycytidine (R_f = 0.67, 4:1 dichloromethane/methanol) was co-distilled twice with dry pyridine (~10 mL) and dissolved in dry pyridine (20 mL). Phenoxyacetic anhydride (1.5 g) was added, and the resulting solution was stirred for 1 hour. A further 0.18g phenoxyacetic anhydride (0.18 g) was added and stirred for another hour. The reaction mixture was evaporated *in vacuo* to dryness and co-distilled (3x) with toluene. The residue was dissolved in dichloromethane (~50 mL) and extracted with 1M aqueous NaHCO₃ solution (~50 mL), which was re-extracted with dichloromethane (~20 mL). The combined organic phases were dried over sodium sulfate and reduced *in vacuo* to yield crude 3',5'-di trimethylsilyl-N-phenoxyacetyl 5-aza-2'-deoxycytidine (3g; R_f =

0.82, 9:1 dichloromethane/methanol). The crude material was dissolved in anhydrous DMF (20 mL) and transferred to a 50 mL plastic falcon tube, and TAS-F (2.4 g) was added (gas evolved). The reaction proceeded for 4 hours at 22 °C (the vial was not fully closed to reduce pressure built up). The DMF was evaporated *in vacuo* and the remaining residue was subjected to column chromatography (30 g silica gel, 2.5 cm column, 99:1 to 9:1 dichloromethane/methanol). A white solid N-phenoxyacetyl 5-aza-2'-deoxycytidine (0.81 g; R_f = 0.26, 9:1 dichloromethane/methanol) was obtained. This compound (0.6 g) was twice co-distilled with anhydrous pyridine and dissolved in anhydrous pyridine (20 mL) before dimethoxytrityl chloride (0.9 g) was added and stirred for 2 hours at 22 °C. Solvents were removed *in vacuo* and co-distilled (3x) with toluene. The residue was dissolved in dichloromethane (50 mL) and extracted with 1M aqueous NaHCO₃ solution (~50 mL), which was re-extracted with dichloromethane (~20 mL). The combined organic phases were dried over sodium sulfate and reduced *in vacuo*.

The residue was subjected to silica gel chromatography (dichloromethane-100% to 95:5 dichloromethane/methanol), which yielded 5'-dimethoxytrityl-N-phenoxyacetyl 5-aza-2'-deoxycytidine (0.35 g, 0.53 mmole, 32%, R_f = 0.49, 9:1 dichloromethane/methanol). This intermediate (0.3 g) was dissolved in dry acetonitrile (2 mL) before 0.3 M benzylthiotetrazole (0.9 mL) solution in dry acetonitrile and cyanoethyltetraisopropyl phosphordiamidite (0.17 mL) were added. The mixture was stirred at 22 °C for 1.5 hours. TLC (2:1 ethyl acetate/hexanes + 2% TEA) showed a diastereoisomeric mixture with R_f = 0.27 and 0.36. Solvent was removed *in vacuo* and the remaining residue subjected to column chromatography (20 g silica gel, 2.5 cm column, 9:1 hexanes/ethyl acetate +2% TEA (300 mL), 1:1 hexanes/ethyl acetate +1% TEA (200 mL), 1:2 hexanes/ethyl acetate +0% TEA (250 mL). The decitabine

WO 2007/041071
PCT/US2006/037313

phosphoramidite building 1d, where R_1 = phenoxyacetyl (0.297 g, 0.34 mmol, 76%) eluted with the 1:2 hexanes/ethyl acetate. ESI-MS of 1d (calculated exact mass for $C_{46}H_{53}N_6O_9P$ is 864.36) exhibited m/z 864.1 and 966.4 $[M+NEt_3+H]^+$; ^{31}P NMR ($CDCl_3$, 500MHz) exhibited 149.17 and 149.0 ppm; 1H NMR ($CDCl_3$, 500MHz) exhibited chemical shifts (ppm) 8.63 & 8.59 (1H, doublet, H-6), 7.4-6.6 (18H, multiplet, aromatic DMTr/Pac), 6.05 (1H, triplet, H-1'), 4.79 (2H, singlet, CH_2 of Pac), 4.59 (1H, singlet, H-4'), 4.25 to 4.20 (1H, doublet, H-3'), 3.8-3.7 (1H, multiplet, P-O- CH_2), 3.70 (3H, singlet, CH_3O of DMTr), 3.68 (3H, singlet, CH_3O of DMTr), 3.6-3.48 (3H, multiplet, two CH 's of isopropyl and one P-O- CH_2), 3.36-3.27 (2H, multiplet, H-5'), 2.80 (1H, singlet, H-2'), 2.53 (1H, singlet, H-2'), 2.40 (2H, multiplet, CH_2CN), 1.1 (12H, CH_3 of isopropyl).

[00275] In addition, minor modification of published procedures allow access to 3'- and 5'-O-capped derivatives (Figure 3A, 1g, 1h, 1i, 1j, 1k, 1l) (Bagnall, Bell and Pearson (1978) J. of Fluorine Chem. 11: 93-107), where the cap can be alkyl groups, esters and fatty acid esters, glycol derivatives such as ethylene and propylene glycols; and protected decitabine 3'-linked onto controlled-pore glass support (Figure 3B, 1m, 1n, 1o). Alul, Singman, Zhan and Letsinger (1991) 19: 1527-1532.

[00276] Other decitabine derivatives have the 3'-OH protected with esters (which include but are not limited to acetyl, benzoyl, and halobenzoyl; and fatty acids) and ethers (which include but are not limited to *p*-nitrophenylethyl, methoxymethyl, methylthiomethyl, (phenyldimethylsilyl)methoxymethyl, benzyloxymethyl, *p*-methoxybenzyloxymethyl, *p*-nitrobenzyloxymethyl, *o*-nitrobenzyloxymethyl, (4-methoxyphenoxy)methyl, *t*-butoxymethyl, 4-pentenylloxymethyl, siloxymethyl, 2-methoxyethoxymethyl, 2,2,2-trichloroethoxymethyl, bis(2-chloroethoxy)methyl, 2-(trimethylsilyl)ethoxymethyl, menthoxymethyl, tetrahydropyranyl, tetrahydrofuranyl, and 1-(2-fluorophenyl)-4-methoxypiperidin-4-yl), glycol derivatives such as ethylene and propylene glycos, as shown in Figure 4.

2. Synthesis of DpG and GpD dinucleotides and tetranucleotides on solid support

[00277] The DpG and GpD dinucleotides and tetranucleotides can be synthesized by standard procedures (Figure 5) with slight modification for increased coupling times (>2 minutes). Beaucage and Caruthers (1981) Tet. Lett. 22: 1859-1862; McBride and Caruthers (1983) Tet. Lett. 24: 245-248. Synthesis of GpD dinucleotide 2a and DpGpGpD tetranucleotide 3a can be initiated with the coupling of 1m, 1n or 1o with similarly based protected 5'-O-DMTr 2'-deoxyguanosine-3'-O-cyanoethyl-*N,N*-diisopropylphosphoramidite and 5'-O-DMTr 2'-deoxy-5-azacytidine-3'-O-cyanoethyl-*N,N*-diisopropylphosphoramidites (1d, 1e or 1f), as shown in Figure 6A. Subsequent release from the solid support (such as controlled pore glass, CPG) and removal of carbamate protective groups with bases such as DBU/pyridine (or acetonitrile), and methanolic ammonia for removal of the 4-O-methoxy and 4-O-methylthio protective groups, yield the desired oligonucleotides with the last DMTr group on or off. DpGpGpD (3b) can be similarly obtained (Figure 7).

[00278] For example (Figure 6A and 6B), an Amersham AKTA Oligopilot 10 system is loaded with a protected decitabine-linked CPG solid support 1m (where R_1 = phenoxyacetyl), which is coupled with 2-2.5 equivalents of *tert*-butyl phenoxyacetyl 2'-deoxyguanosine phosphoramidite in presence of 60% of 0.3 M benzylthiotetrazole activator (in acetonitrile) for 2.5 minutes. The CPG solid support containing protected GpD dinucleotide is treated with 20 mL of 50 mM K_2CO_3 in methanol for 1 hour and 20 minutes. The coupled product is oxidized with 2 M *tert*-butylhydroperoxide in dry acetonitrile (prepared by dissolving *tert*-butylhydroperoxide in 80% *tert*-butylperoxide) for 5 minutes. The dimethoxy trityl protective group is removed with 3% dichloroacetic acid in toluene. The CPG solid support is washed with dry methanol; the filtrate is neutralized by addition of 2 mL of 1 M acetic acid in methanol. The solution is concentrated by rotary evaporation; the residue is taken up in 200 mM

WO 2007/041071
PCT/US2006/037313

triethylammonium acetate (pH 6.9), washed with acetonitrile (500 μ L of 50% aqueous acetonitrile), and filtered through a syringe filter. The GpD dinucleotide is subsequently purified by the ÄKTA Explorer 100 HPLC with a Gemini C18 preparative column (Phenomenex), 250x21.2 mm, 10 μ m with guard column (Phenomenex),

50x21.2mm, 10 μ m, with 50 mM triethylammonium acetate (pH 7) in MilliQ water (Mobile Phase A) and 80%

5 acetonitrile in MilliQ water (Mobile Phase B), with 2% to 20/25% Mobile Phase B in column volumes. The ESI-MS (-ve) of GpD dinucleotide 2a, where X^+ = triethylammonium (calculated exact mass for the neutral compound $C_{18}H_{24}N_9O_{10}P$ is 557.14), exhibited m/z 556.1 $[M-H]^-$ and 1113.1 for $[2M-H]^-$ (see mass spectrum in Figure 30).

[00279] When the cycle is repeated three times with 2-2.5 equivalents of *tert*-butyl phenoxyacetyl 2'-

deoxyguanosine or phenoxyacetyl 5-aza-2'-deoxycytidine phosphoramidite in presence of 60% of 0.3 M

10 benzylthiotetrazole activator (in acetonitrile) for 2.5 minutes and 10 minutes, respectively, (Figure 6b and 7), the DpGpGpD tetranucleotide 3b is obtained, where X^+ = triethylammonium (calculated exact mass for the neutral compound $C_{36}H_{47}N_{18}O_{22}P_3$ is 1176.23), exhibited m/z 587.2 for $[M-2H]^{2-}$ and 1175.2 $[M-H]^-$ (Figure 32).

[00280] In addition, DpG dinucleotide 2b and GpDpGpD 3c can be synthesized by coupling 1s (where R_1 = carbamate protective group) with phosphoramidite building blocks 1d, 1e or 1f (Figure 8). GpDpDpG and GpDpG (3c') can likewise be obtained (Figure 9).

[00281] For example, when a protected 2'-deoxyguanosine-linked CPG solid support 1s (where R_1 = *tert*-butyl phenoxyacetyl), which is coupled with 2-2.5 equivalents of phenoxyacetyl decitabine phosphoramidite (Figure 2A, 1d, where R_1 = phenoxyacetyl; see mass spectrum in Figure 38) in the presence of 60% of 0.3 M benzylthiotetrazole

20 activator (in acetonitrile) for 10 minutes. The CPG-solid support containing protected DpG dinucleotide is treated with 20 mL of 50 mM K_2CO_3 in methanol for 1 hour and 20 minutes. The coupled product is oxidized, protective

group removed, washed, filtered, and purified as described for GpD dinucleotide. The ESI-MS (-ve) of DpG dinucleotide 2b, where X^+ = triethylammonium (calculated exact mass for the neutral compound $C_{18}H_{24}N_9O_{10}P$ is

557.14), exhibited m/z 556.1 $[M-H]^-$ and 1113.1 for $[2M-H]^-$ (see mass spectrum in Figure 31). The DpG

25 dinucleotide 2b, where X^+ = sodium, is obtained by re-dissolving the triethylammonium salt in 4 mL water, 0.2 mL 2M $NaClO_4$ solution. When 36 mL acetone is added, the dinucleotide precipitates. The solution is kept at -20°C for several hours and centrifuged at 4000 rpm for 20 minutes. The supernatant is discarded and the solid is washed with 30 mL acetone followed by an additional centrifugation at 4000 rpm for 20 minutes. The precipitate is dissolved in water and freeze dried, which exhibited m/z 556.0 $[M-H]^-$ (see mass spectrum in Figure 36).

[00282] When the cycle is repeated twice with 2-2.5 equivalents of *tert*-butyl phenoxyacetyl 2'-deoxyguanosine or 30 phenoxyacetyl 5-aza-2'-deoxycytidine phosphoramidite in presence of 60% of 0.3 M benzylthiotetrazole activator (in acetonitrile) for 2.5 minutes and 10 minutes, respectively, the GpDpG trinucleotide 3c' is obtained, where X^+ = triethylammonium (calculated exact mass for the neutral compound $C_{28}H_{36}N_{14}O_{16}P_2$ is 886.2), which exhibited m/z 885.16 $[M-H]^-$ (see mass spectrum Figure 33).

[00283] When cycle is repeated three times, the DpGpDpG tetranucleotide 3c is obtained, where X^+ =

35 triethylammonium (calculated exact mass for the neutral compound $C_{36}H_{47}N_{18}O_{22}P_3$ is 1176.23), which exhibited m/z 587.4 for $[M-2H]^{2-}$ and 1175.4 $[M-H]^-$ (see mass spectrum in Figure 34).

[00284] When the phosphite triester, newly formed during the coupling step, is converted to the corresponding phosphorothioate triester with 5% phenylacetyl disulfide (PADS) in dichloroethane/sym collidine 4/1 (v/v), 4.3 mL solution (3.6 column volumes), flow rate 50cm/h (contact time 3 column volumes), the phosphorothioate derivative 40 of 2b can be obtained. The sulfurization is completed within 3 minutes, at which time excess reagent is removed from the reaction vessel by washing with acetonitrile. Subsequent deprotection and purification, as described for 2a, gives phosphorothioate DpG (Sp & Rp, Figure 13, 2e), where X^+ = triethylammonium (calculated exact mass for the

WO 2007/041071
 PCT/US2006/037313
 neutral compound $C_{18}H_{24}N_9O_9P_2S$ is 573.12), which exhibited m/z 571.9 for $[M-H]^-$ (see mass spectrum in Figure 35).

[00285] When cycle is repeated once with DMT hexaethylenglycol phosphoramidite (60% activator, 7 min coupling time), followed by standard oxidation and purification as described for 2a, the HEG-DpG dinucleotide 2d is obtained (Figure 12), where X^+ = triethylammonium and Cap = hexaethyleneglycol phosphate (calculated exact mass for the neutral compound $C_{30}H_{49}N_9O_{19}P_2$ is 901.71), which exhibited m/z 900.4 $[M-H]^-$ (see mass spectrum in Figure 37).

3. Inhibition of DNA Methylation by DpG and GpD di-, tri- and tetranucleotides

[00286] The demethylating activity of DpG and GpD di-, tri-, and tetranucleotides were tested in a cell-based GFP (green fluorescent protein) assay. This assay, which is schematically illustrated in Figure 25, has a GFP gene regulated by the CMV promoter and is sensitive to the methylation of CpG sites within the promoter. A decrease in methylation resulting from exposure to a methylation inhibitor leads to GFP expression and is readily scored. Specifically, the CMV-EE210 cell line containing the epigenetically silenced GFP transgene was used to assay for reactivation of GFP expression by flow cytometry. CMV-EE210 was made by transfecting NIH 3T3 cells with the pTR-UF/UF1/UF2 plasmid (Zolotuhin et al., 1996), which is comprised of pBS(+) (Stratagene, Inc.) containing a cytomegalovirus (CMV) promoter driving a humanized GFP gene adapted for expression in mammalian cells. After transfection, high-level GFP expressing cells were initially selected by FACS analysis and sorting using a MoFlo cytometer (Cytomation, Inc.). Decitabine, potent inhibitor of mammalian DNMT1, was used as a positive control.

To screen for reactivation of CMV-EE210, decitabine (at 1 μ M) or a test compound (at a concentration of 30-50 μ M) was added to complete medium (phenol red free DMEM (Gibco, Life Technologies) supplemented with 10% fetal bovine serum (Hyclone)). Cells were then seeded to 30% confluence (~5000 cell/well) in 96 well plate containing the test compounds and grown for three days in at 37°C in 5% CO_2 . The plates were examined under a fluorescent microscope using a 450-490 excitation filter (I3 filter cube, Leica, Deerfield IL). Wells were scored g1 positive if (10%) of viable cells express GFP, g2 positive if 30% of viable cells express GFP and g3 if >75% of the viable cells express GFP. GFP 50 is the concentration of an inhibitor that (like an IC_{50}) is the dose at which the GFP expression level goes from g3 to g1/2. Table 1 lists the results of the test for decitabine, DpG, GpD, GpDpG, DpGpGpD and DpGpDpG as DNA methylation inhibitors. As shown in Table 1, all of the 5 oligonucleotide analogues tested were able to inhibit DNA methylation effectively at low concentrations, resulting in reactivation of the transcription of the GFP gene.

Table 1: Preliminary screening of demethylating activity

Compound	GFP Expression Level	IC 50 (nM)
Decitabine	g3	500
DpG	g3	400
GpD	g3	700
GpDpG	g3	1800
DpGpGpD	g3	1100
DpGpDpG	g3	1400

4. ~~PCT/US06/37313~~ Synthesis of DpG and GpD dinucleotides and tetranucleotides in solution

[00287] For the synthesis of these oligonucleotides in large scale, the use of soluble polymeric supports is desirable. Bayer and Mutter, (1972) *Nature* 237: 512-513; Bonora (1995) *Appl. Biochem. Biotechnol.* 54: 3-17. The polymer support poly (ethylene glycol) or PEG allows synthetic process to be carried out in a homogeneous phase and assures an easy intermediate purification step through simple precipitation-and-filtration procedures. Harris *Poly(ethylene glycol) Chemistry. Biotechnical and Biomedical Applications*, J. M. Harris (Ed.), Plenum Press, New York (USA), 1992, pp. 1-14 [book citation]. For example, 3'-linked derivatives such as 1t, 1u, or 1v (Figure 10) can be easily adapted to the standard phosphoramidite-based chemistry employed in the preceding solid-phase procedures to give the DpG and GpD di- and tetranucleotides 2a, 2b, 3a, 3b, 3c, 3d.

[00288] Alternatively, the novel DpG dinucleotide 2a can be prepared in solution by coupling derivatives 1p, 1q, or 1r with similarly based protected 5'-O-DMTr 2'-deoxyguanosine-3'-O-cyanoethyl-*N,N*-diisopropylphosphoramidite and GpD from the coupling of similarly 3'-protected 2'-deoxyguanosine with 5'-O-DMTr 2'-deoxy-5-aza-cytidine-3'-O-cyanoethyl-*N,N*-diisopropylphosphoramidites (1d, 1e or 1f) in acetonitrile and/or dichloromethane, followed by oxidation with iodine/water, deprotection of the base protective groups, and removal of the DMTr group (as in the standard cycle for oligonucleotide synthesis).

[00289] In addition, the novel DpG (2c) dinucleotide with the terminal 3'-OH and 5'-OH capped with methyl group can be prepared by coupling 3'-O-methyl derivative 1g, 1h or 1i with a 5'-O-methyl derivative of 2'-deoxyguanosine-3'-O-cyanoethyl-*N,N*-diisopropylphosphoramidite 1w (Figure 11), followed by oxidation with iodine/water, deprotection of the base protective groups, and removal of the DMTr group (as in the standard cycle for oligonucleotide synthesis). Dinucleotide GpD (2d) can likewise be prepared by coupling 3'-O-methyl 2'-deoxyguanosine derivative 1x with 5'-O-methyl derivative of 2'-5-azacytidine-3'-O-cyanoethyl-*N,N*-diisopropylphosphoramidite 1j, 1k or 1l (Figure 12).

5. Synthesis of DpG and GpD oligonucleotides resistant to cytidine deaminases and nucleases

[00290] In general, oligonucleotides in biological fluids are subject to nuclease degradation. Stein and Cheng (1993) *Science* 261: 1004-1012; Cohen (1994) *Adv. Pharmacol.* 25: 319-339. To increase stability and resistance to nuclease degradation phosphorothioate dinucleotide and tetranucleotide derivatives such as 2e, 2f, 3e, 3f, 3g, and 3h (Figure 13, 14 and 15) are also made, where the internucleotide non-bridging oxygen is replaced with sulfur. Standard phosphoramidite protocols are used, except for the substitution of bis(*O,O*-diisopropoxy phosphinothioyl) disulfide (S-tetra) for iodine during the oxidation step. Zon and Stec (1991) In Eckstein, F. (ed.), *Phosphorothioate Analogues' in Oligonucleotides and Their Analogs: A Practical Approach*. IRL Press, pp. 87-108; Zon, G. (1990) In Hancock, W.S. (ed.), *High Performance Liquid Chromatography in Biotechnology*. Wiley, New York, Ch.14, pp. 310-397 [book citations]; Stec, Uznanski, Wilk, Hirschbein, Fearon, and Bergot (1993) *Tet. Lett.* 34: 5317-5320; Iyer, Phillips, Egan, Regan, and Beaucage (1990) *J. Org. Chem.* 55: 4693-4699.

[00291] Another potential hindrance to the application of these oligonucleotides as pharmaceuticals is the ubiquitous presence of cytidine deaminase (CDA) since deamination of decitabine results in total loss of activity. Momparler, Cote and Eliopoulos (1997) *Leukemia* 11 (Suppl.1): 1-6; Chabot, Bouchard and Momparler (1983) *Biochem. Pharmacol.* 32: 1327-1328; Laliberte, Marquez and Momparler (1992) *Cancer Chemother. Pharmacol.* 30: 7-11. To address this problem, the oligonucleotides containing decitabine derivatives with the 4-NH₂ is replaced by 4-NR₃R₄ (where R₃ and R₄ can be alkyl, alkyl amine, and alkyl alcohol) are also prepared to give derivatives such as 2g, 2h, 2i, 2j, 2k, 2l, 2m, 2n, 3i, 3j, 3k, 3l, 3m, 3n, 3o, and 3p (Figure 16, 17, 18, 19, 20 and 21). Standard phosphoramidite protocols are used, except for the substitution of alkyl amines, alkyl diamines, and

PCT/US2006/037313

hydroxyl amines for ammonia in methanol during the removal of 4-methoxy and 4-methylthio. Since secondary and tertiary amines, diamines, and hydroxyl amines make worse leaving groups than ammonia, these derivatives are more difficult to deaminate.

5
6. Synthesis of DpG and GpD-rich oligonucleotides based on the CpG islands of the promoter regions of cancer related genes such as P15 (CDKN2B), BRCA1, and P16 (CDKN2A)

[00292] Oligonucleotide analogues rich in DpG and GpD islets that range in length from 5 to 100 bases can be prepared, where D can be decitabine or decitabine analogues. Unlike the above described DpG and GpD
10 dinucleotides and tetranucleotides, these relatively longer DpG and GpD-rich oligonucleotide analogues not only function restrictively within the CpG islands of the promoter regions but specific to a segment within the promoter region sequence for cancer related genes such as P15 (CDKN2B), P16 (CDKN2A) and BRCA1. For examples, 8-mer, 10-mer, and 12-mer DpG and GpD-rich oligonucleotide analogues (Figure 26) based on the P15, P16, and
15 BRCA1 promoter region sequences (Figure 27, 28 and 29, respectively) can be prepared by using phosphoramidite building block 1d, 1e or 1f in a standard solid phase oligonucleotide synthesis. More examples of oligonucleotides that can be modified to incorporate 5-aza-cytosine therein are listed in Figures 27, 28 and 29. These oligonucleotide analogues can function like primers and get incorporated into replicating DNA only at that specific segment of the promoter region sequence of P15, P16 or BRCA1, thus effectively and selectively inhibiting methylation of the promoter region.

20 [00293] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and
25 that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMS

1. An isolated or synthetic dinucleotide analogue, its salt or ester, comprising one 5-aza-cytosine as a base residue in the sequence of the dinucleotide analogue, wherein the dinucleotide is linked with a phospholinker consisting of one phosphorus and wherein the dinucleotide analogue has a general formula of:

5'-DpG-3' or 5'GpD-3', wherein D is decitabine; p is a phospholinker; and G is deoxyguanosine.

2. The dinucleotide analogue of claim 1, wherein the phospholinker p is a sugar phosphorodiester linker.

3. The dinucleotide analogue of claim 1, wherein the phospholinker p is a 2'-deoxyribose phosphorodiester linker.

4. The dinucleotide analogue of claim 1, wherein the phospholinker p is a 2'-deoxyribose phosphorothioate linker.

5. The dinucleotide analogue of claim 1, wherein the phospholinker p is a 2'-deoxyribose boranophosphate linker.

6. The dinucleotide analogue of claim 1, wherein the phospholinker p is a 2'-deoxyribose methylphosphonate linker.

7. The dinucleotide analogue of claim 1, wherein the salt is formed with an acid selected from the group consisting of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, carboxylic, sulfonic, sulfo or phospho acids, acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid, isonicotinic acid, amino acid, glutamic acid, aspartic acid, phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane- 1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid, and ascorbic acid.

8. The dinucleotide analogue of claim 1, wherein the salt is hydrochloride, mesylate, EDTA, sulfite, L-Aspartate, maleate, phosphate, L-Glutamate, (+)-L-Tartrate, citrate, L-Lactate, succinate, acetate, hexanoate, butyrate, or propionate salt.

9. The dinucleotide analogue of claim 1, wherein the salt is a sodium, calcium, lithium, potassium, ammonium, or trialkylammonium salt.

10. The dinucleotide analogue of claim 9, wherein the trialkylammonium salt is trimethylammonium, triethylammonium or tripropylammonium salt.

11. A pharmaceutical composition comprising a dinucleotide analogue of claim 1 or its pharmaceutically acceptable salt or ester, and a pharmaceutically-acceptable carrier.
12. The pharmaceutical composition according to claim 11, the dinucleotide analogue of claim 1 or its pharmaceutically acceptable salt or ester is in solid form.
13. The pharmaceutical composition according to claim 11, wherein the pharmaceutically-acceptable carrier is an aqueous solution.
14. The pharmaceutical composition according to claim 13, wherein the aqueous solution comprises a water miscible non-aqueous solvent selected from the group consisting of ethanol, glycerin, propylene glycol, polyethylene glycol, and combinations thereof.

15. The pharmaceutical composition according to claim 11, wherein the pharmaceutically-acceptable carrier is a solution that has less than 40% of water.
16. The pharmaceutical composition according to claim 11, wherein the pharmaceutically-acceptable carrier is a solution that has less than 5% of water.
17. The pharmaceutical composition according to claim 16, wherein the pharmaceutically-acceptable carrier is a solution comprising ethanol, glycerin, propylene glycol, polyethylene glycol, or a combination thereof.
18. The pharmaceutical composition according to claim 11, wherein the pharmaceutically-acceptable carrier is liposome.
19. The pharmaceutical composition according to claim 18 wherein the liposome is cationic liposome.
20. The pharmaceutical composition according to claim 11, wherein the pharmaceutical composition is in a form of emulsion.
21. The pharmaceutical composition according to claim 11, wherein the emulsion comprises cationic emulsifier.
22. A kit, comprising: a first vessel containing a dinucleotide analogue of claim 1 its salt or ester thereof.
23. The kit of claim 22, further comprising: a second vessel containing a diluent comprising water, glycerin, propylene glycol, polyethylene glycol or combinations thereof.
24. The kit of claim 23, further comprising: a written instruction describing how to administer the dinucleotide analogue, its salt or ester thereof to a host in need thereof.
25. A method for inhibiting DNA methylation in a cell in vitro, comprising: contacting the cell with the dinucleotide analogue, its salt or ester thereof of claim 1, such that DNA methylation activity of the cell is inhibited.
26. The method of claim 25, wherein the cell resides in a cell culture in vitro.
27. The method of claim 25, wherein the cell resides in a tissue section in vitro.

28. The method of claim 25, wherein the step of contacting includes contacting the cell with a biologically effective amount of the dinucleotide analogue, its salt or ester thereof of claim 1 such that at least 50% of the DNA methylation activity of the cell is inhibited.
29. Use of a pharmaceutical composition comprising a therapeutically-effective amount of the dinucleotide analogue, its salt or ester thereof of claim 1, and a pharmaceutically-acceptable carrier, in the manufacture of a medicament for treating a disease associated with aberrant DNA methylation.
30. The use according to claim 29, wherein the medicament is to be administered orally, parenterally, topically, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery, subcutaneously, intraadiposally, intraarticularly, or intrathecally.
31. The use according to claim 29, wherein the medicament is to be administered intravenously.
32. The use according to claim 29, wherein the medicament is to be administered subcutaneously.
33. The use according to claim 29, wherein a second therapeutic agent is to be administered in combination with the medicament.
34. The use according to claim 33, wherein the second therapeutic agent is selected from the group consisting of histone deacetylase inhibitors, antibiotic agents, alkylating agents, retinoids, hormonal agents, plant-derived agents, biologic agents, interleukins, interferons, cytokines, immuno-modulating agents, and monoclonal antibodies.
35. The use according to claim 34, wherein the histone deacetylase inhibitor is selected from the group consisting of trichostatin A, suberoylanilide hydroxamic acid, oxamflatin, suberic bishydroxamic acid, m-carboxy-cinnamic acid bishydroxamic acid, pyroxamide, trapoxin A, apicidin, depsipeptide, N-(2-amimophenyl)-4-[N-(pyridin-3-ylmethoxycarbonyl)aminomethyl] benzamide, butyric acid, phenylbutyrate and arginine butyrate.
36. The use according to claim 29, wherein the disease associated with aberrant DNA methylation is selected from the group consisting of hematological disorders, benign tumor and cancer.
37. The use according to claim 36, wherein the hematological disorder is selected from the group consisting of acute myeloid leukemia, acute promyelocyte leukemia, acute lymphoblastic leukemia, chronic myelogenous leukemia, myelodysplastic syndromes, and sickle cell anemia.

38. The use according to claim 36, wherein the cancer is selected from group consisting of breast cancer, skin cancer, bone cancer, prostate cancer, liver cancer, lung cancer, non-small cell lung cancer, brain cancer, cancer of the larynx, gall bladder, pancreas, rectum, parathyroid, thyroid, adrenal, neural tissue, head and neck, colon, stomach, bronchi, and kidney cancer, basal cell carcinoma, squamous cell carcinoma of both ulcerating and papillary type, metastatic skin carcinoma, osteo sarcoma, Ewing's sarcoma, veticulum cell sarcoma, myeloma, giant cell tumor, small-cell lung tumor, gallstones, islet cell tumor, primary brain tumor, acute and chronic lymphocytic and granulocytic tumors, hairy-cell tumor, adenoma, hyperplasia, medullary carcinoma, pheochromocytoma, mucosal neuronms, intestinal ganglioneuromas, hyperplastic corneal nerve tumor, marfanoid habitus tumor, Wilm's tumor, seminoma, ovarian tumor, leiomyomater tumor, cervical dysplasia and in situ carcinoma, neuroblastoma, retinoblastoma, soft tissue sarcoma, malignant carcinoid, topical skin lesion, mycosis fungoide, rhabdomyosarcoma, Kaposi's sarcoma, osteogenic sarcoma, malignant hypercalcemia, renal cell tumor, polycythemia vera, adenocarcinoma, glioblastoma multiforma, leukemias, lymphomas, malignant melanomas, and epidermoid carcinomas.

39. The dinucleotide analogue of claim 1, wherein the dinucleotide analogue is 5'-DpG-3', wherein D is decitabine; p is a phospholinker; and G is deoxyguanosine.

12636-350.601 Sequence Listing.txt

SEQUENCE LISTING

<110> SUPERGEN, INC.

Phiasivongsa, Pasit

Redkar, Sanjeev

<120> OLIGONUCLEOTIDE ANALOGUES INCORPORATING 5-AZA-CYTOSINE THEREIN

<130> 12636-350.601

<160> 33

<170> PatentIn version 3.1

<210> 1

<211> 8

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide analogue targeting a segment of P15 promoter regi
on

<220>

<221> misc_feature

<222> (5)..(5)

<223> n = 5-aza-cytosine as base residue

<220>

<221> misc_feature

<222> (3)..(3)

<223> n = 5-aza-cytosine as base residue

<400> 1

ttngngaa

8

<210> 2

<211> 8

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide analogue targeting a segment of P15 promoter regi
on

<220>

<221> misc_feature

<222> (6)..(6)

<223> n = 5-aza-cytosine as base residue

<400> 2

tgctngt

8

<210> 3

12636-350.601 Sequence Listing.txt

<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide analogue targeting a segment of BRCA1 promoter region

<220>
<221> misc_feature
<222> (4)..(4)
<223> n = 5-aza-cytosine as base residue

<220>
<221> misc_feature
<222> (9)..(9)
<223> n = 5-aza-cytosine as base residue

<400> 3
aggnacagna 10

<210> 4
<211> 8
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide analogue targeting a segment of BRCA1 promoter region

<220>
<221> misc_feature
<222> (4)..(4)
<223> n = 5-aza-cytosine as base residue

<220>
<221> misc_feature
<222> (7)..(7)
<223> n = 5-aza-cytosine as base residue

<400> 4
gtgnagna 8

<210> 5
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide analogue targeting a segment of P16 promoter region

<220>
<221> misc_feature

12636-350.601 Sequence Listing.txt

<222> (3)..(3)

<223> n = 5-aza-cytosine as base residue

<220>

<221> misc_feature

<222> (7)..(7)

<223> n = 5-aza-cytosine as base residue

<220>

<221> misc_feature

<222> (10)..(10)

<223> n = 5-aza-cytosine as base residue

<400> 5

aangggnggn gg

12

<210> 6

<211> 10

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide analogue targeting a segment of P16 promoter regi

on

<220>

<221> misc_feature

<222> (3)..(3)

<223> n = 5-aza-cytosine as base residue

<220>

<221> misc_feature

<222> (6)..(6)

<223> n = 5-aza-cytosine as base residue

<220>

<221> misc_feature

<222> (8)..(8)

<223> n = 5-aza-cytosine as base residue

<400> 6

cangggnggg

10

<210> 7

<211> 2144

<212> DNA

<213> homo sapien

<400> 7

tttttcattc agtcaacttg ctctcggaag ctcacacatc tgcctcgtgc aagattctca 60

gtcattttac ttatgcatig gtctttcc taccaccatt cttatgtcc ccctcaaaga 120

aaaacattat ctccatttc ctatcaact ccaaacagct ttcatttttc tgacatattt 180

12636-350.601 Sequence Listing.txt

actacctaag aaaatggctc aagaattggg tagactatct tgtcctaact ttctgataa 240
gtttcagaga aactcaaagg tcaaaacaag agcataagag taaaggtaga gaaattaaga 300
aactgaagac taggaaatgg ggggtgggat gggaaagaaa aagaattgt tataatgcta 360
cccggtccc ttccctgtcc aggtggattt cagctctgtt gaggtctgt cagtagatat 420
tcagccctaa ccagcacttc catggtggtg gcactccac tgcctttaa aagaagagc 480
ttttttaat tctacagga tttgggggat gaggagtcag agctaaggta tctaaaaaa 540
aacatgtgaa gactctcatt ttgcaatca caagcaattg cctctctgtt aagactttgt 600
cttctcagc actccgaacc aaaatgattc tgtaaacaaa aattgttcac tttaggaga 660
ggtcaccta tgcagtctc caccaaagt ttaggcaac aaatccataa ctgcggttc 720
tcttctatc caatgtagca tccgtgaaa tgttttaaat atttaagta ataatgttg 780
attcaaactc acctaggaag attaggaagg ggaaaaaag cacttggcat ttaactctc 840
agaagagaat ttaatgacag gttcagcctg ttaatgaca agcccagcac cacacccctc 900
tctatgatg ttctattt actgcataaa ttctttat tactcatgat aaataaaaat 960
aagatacctg acaagtggg ttaaatagg taagagtga acaaagatt tactgtaca 1020
atatgatgaa actgggatct cagattcta aagtataatt ttttgtct tatgtgtgcc 1080
aggtggcac tctaatctc gaactagttt tttctctt taagggtgt atccataatg 1140
caaaaatgga aagaattaa aagcacgcg aaaacatgat tctcgggatt ttctctatt 1200
tttatggtg actaattcaa acagaaagac acatccaaga gaaaattgct aagtttgata 1260
caagtatga aactgtgaa gcccaagtac tgcctgggga tgaattaac ttgatgaca 1320
ggtcagagc tgcgcttc agacatcta agaaagacg agttatttg aatgacttc 1380
tctcgtcac aaggagacca ccaacgtctc cacagtga acaactggct ggctgaagga 1440
acagaaatcc tctgtccgc ctactgggga ttaggagctg agggcagtgg tgaacattcc 1500
caaaatatta gcctggctt tactggacat ccagcgagca gtgcagccag cattctggc 1560
ggctccctg ccagctctt ggcgatgag tctagcatc ttgggcagg ctccccgc 1620
ctcgtgacgc gtcggcccg gcctggcctc ccggcgatca cagcggacag gggcgggagc 1680
ctaaggggtt ggggagacgc cggcccttg gccagctga aaacggaatt cttgccggc 1740
tggctccca ctctccaga gcgaggcgg gcagtgaaga ctccgcagc cgtccgcacc 1800
ctcggccag agcggcttg agctcggctg cgtccgcgt aggcgtttt tccagaagc 1860

5/11

12636-350.601 Sequence Listing.txt

aatccaggcg cgcccgtgg ttcttgagcg ccaggaaaag cccggagcta acgaccggcc 1920
gctcgccac tgcacggggc cccaagccgc agaaggacga cgggagggtg atgaagctga 1980
gcccagggtc ctaggaagg agagagtgcg ccggagcagc gtgggaaaga agggaagagt 2040
gtcgttaagt ttacggccaa cggtggaatg tccgggccgc tgcgcgtctg ggggctgcgg 2100
aatgcgcgag gagaacaagg gcatgccagc tgggggcggc agcg 2144

<210> 8
<211> 8
<212> DNA
<213> homo sapien

<400> 8
ttcgcgaa 8

<210> 9
<211> 8
<212> DNA
<213> homo-sapien

<400> 9
tgcctcgt 8

<210> 10
<211> 8
<212> DNA
<213> homo sapien<400> 10
tgccggct 8

<210> 11
<211> 8
<212> DNA
<213> homo sapien<400> 11
cggeccgg 8

<210> 12
<211> 8
<212> DNA
<213> homo sapien<400> 12
gctcggct 8

<210> 13
<211> 987
<212> DNA
<213> homo sapien

12636-350.601 Sequence Listing.txt

<400> 13
cggagagggg gagaacagac aacgggaggc ggggagcagc atggagccgg cggcggggag 60
cagcatggag ccttcggctg actggctggc cacggccggc gcccggggtc gggtagagga 120
ggtcggggcg ctgctggagg cggggggcgt gcccaacgca cgaatagtt acggtcggag 180
gccgatccag gtcgatgata tgggcagcgc ccgagtggcg gagctgctgc tgctccacgg 240
cgcggagccc aactgcgcg accccggcac tctaccgga cccgtgcacg acgctgccc 300
ggagggcttc ctggacagc tgggtgtgct gcaccggggc ggggcgcggc tggacgtgcg 360
cgatgcctgg ggccgtctgc ccgtggacct ggctgaggag ctgggccatc gcgatgtgc 420
acgttacctg cgcgcggctg cggggggcac cagaggcagt aacatgccc gcatagatgc 480
cgcggaaggt cctcagaca tcccgatg aaagaaccag agaggctctg agaaacctg 540
ggaaacttag atcatcagtc accgaaggtc ctacagggcc acaactgccc cggccacaac 600
ccaccccgct ttctagttt tcatttagaa aatagagctt taaaaatgt cctgccttt 660
aacgtagata taagccttc ccactaccg taaatgtcca ttatatcat ttttatata 720
ttctataaa aatgtaaaaa agaaaaacac cgctctgcc ttctactgt gttggagttt 780
tctggagtga gcactcacg cctaagcgca cattcatgtg ggcatitctt gcgagcctcg 840
cagcctccgg aagctgtoga ctcatgaca agcattttgt gaactagga agctcagggg 900
ggttactggc ttctcttag tcacactgct agcaaatggc agaaccaaag ctcaaataaa 960
aataaataa tttcattca ttcactc 987

<210> 14
<211> 12
<212> DNA
<213> homo sapien

<400> 14
aacgggaggc gg 12

<210> 15
<211> 10
<212> DNA
<213> homo sapien

<400> 15
cacggcgcgg 10

<210> 16
<211> 8

12636-350.601 Sequence Listing.txt

<212> DNA

<213> homo sapien<400> 16

cgggcggc

8

<210> 17

<211> 8

<212> DNA

<213> homo sapien<400> 17

agcagcat

8

<210> 18

<211> 8

<212> DNA

<213> homo sapien

<400> 18

gcgccgac

8

<210> 19

<211> 649

<212> DNA

<213> homo sapien

<400> 19

ccacctaat g tactgaatt gcaaattgat agttgttcta gcagtgaaga gataaagaaa 60

aaaaagtaca accaaatgcc agtcaggcac agcagaaacc tacaactcat ggaaggtaaa 120

gaacctgcaa ctggagccaa gaagagtaac aagccaaatg aacagacaag taaaagacat 180

gacagcgata cttcccaga gctgaagta acaaatgcac ctggtcttt tactaagtgt 240

tcaaatacca gtgaacttaa agaattgtc aatcctagcc ttccaagaga agaaaaagaa 300

gagaaactag aaacagttag tgtctaataa tgctgaagac cccaaagatc tcatgttaag 360

tggagaaagg gttttgcaa ctgaaagatc ttagagagt agcagtattt catttgtacc 420

tggtagtgat tatggcactc aggaaagtat ctggttactg gaagitagca ctctagggaa 480

ggcaaaaaca gaaccaaata aatgtgtgag tcagtgtgca gcatttgaaa accccaaggg 540

actaatcat ggtgttcca aagataatag aatgacaca gaaggcttta agtatccatt 600

gggacatgaa gtaaccaca gtcgggaaac aagcatagaa atggaagaa 649

<210> 20

<211> 10

<212> DNA

<213> homo sapien

<400> 20

12636-350.601 Sequence Listing.txt
10

aggcacagca

<210> 21
<211> 8
<212> DNA
<213> homo sapien

<400> 21
gtgcagca 8

<210> 22
<211> 8
<212> DNA
<213> homo sapien

<400> 22
cagcgata 8

<210> 23
<211> 8
<212> DNA
<213> homo sapien

<400> 23
tagcagtg 8

<210> 24
<211> 8
<212> DNA
<213> homo sapien

<400> 24
aggcttta 8

<210> 25
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Sequence template for oligonucleotide analogue as DNA methyltrans
ferase inhibitor

<400> 25
ctggatcctt gccccgcccc ttgaattccc 30

<210> 26
<211> 30
<212> DNA
<213> Artificial Sequence

9/11

12636-350.601 Sequence Listing.txt

<220>

<223> Sequence template for oligonucleotide analogue as DNA methyltransferase inhibitor

<400> 26

gggaattcaa atgacgtcaa aaggatccag 30

<210> 27

<211> 50

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequence template for oligonucleotide analogue as DNA methyltransferase inhibitor

<400> 27

cctaccacc cttgatcctt gccccgcccc ttgaattccc aacctccac 50

<210> 28

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequence template for oligonucleotide analogue as DNA methyltransferase inhibitor

<400> 28

atccttgccc cgccccttga at 22

<210> 29

<211> 14

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequence template for oligonucleotide analogue as DNA methyltransferase inhibitor

<400> 29

ttgccccgcc cctt 14

<210> 30

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide analogue as DNA methyltransferase

12636-350.601 Sequence Listing.txt

<220>

<221> misc_feature

<222> (15)..(15)

<223> n = 5-aza-cytosine as base residue

<400> 30

ctggatcctt gccngcccc ttgaattccc

30

<210> 31

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide analogue with hairpin conformation

<220>

<221> misc_feature

<222> (6)..(6)

<223> n = 5-aza-cytosine as base residue

<220>

<221> misc_feature

<222> (11)..(11)

<223> n = 5-aza-cytosine as base residue

<400> 31

ctgaanggat ngttcgatc cggtcag

27

<210> 32

<211> 8

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide analogue

<220>

<221> misc_feature

<222> (3)..(3)

<223> n = 5-aza-cytosine as base residue

<220>

<221> misc_feature<222> (5)..(5)

<223> n = 5-aza-cytosine as base residue

<400> 32

ttngngaa

8

<210> 33

<211> 8

<212> DNA

<213> Artificial Sequence

12636-350.601 Sequence Listing.txt

<220>

<223> Complementary strand of SEQ ID:30

<400> 33

ttcgcgaa

8

1/42




FIGURE 1

2/42




FIGURE 2A

3/42




FIGURE 2B

4/42




FIGURE 3A

5/42







FIGURE 3B

6/42

**FIGURE 4**

7/42




FIGURE 5

8/42




FIGURE 6A

9/42




FIGURE 6B

10/42




FIGURE 7

11/42




FIGURE 8

13/42





FIGURE 10

14/42




FIGURE 11

15/42




FIGURE 12

16/42




FIGURE 13

17/42





FIGURE 14

18/42




FIGURE 15

19/42





FIGURE 16

20/42





FIGURE 17

21/42




FIGURE 18

22/42




FIGURE 19

23/42




FIGURE 20

24/42




FIGURE 21

25/42




FIGURE 22

26/42






FIGURE 23


27/42




**-DpG- Islet
with Natural Phosphodiester Backbone**



**-DpG- Islet
with Phosphorothioate Backbone**




**-DpG- Islet with 2'-Methoxy Ribose
Phosphodiester Backbone**



**-DpG- Islet with 2'-Fluoro Ribose
Phosphodiester Backbone**

FIGURE 24A

28/42



-DpnaG- Islet with Peptide Backbone

FIGURE 24B

29/42




FIGURE 25

30/42

SEQ ID NO:1	TpTpDpxGpDpxGpApA
SEQ ID NO:2	TpGpxCpCpTpDpxGpT
SEQ ID NO:3	ApGpGpxDpApCpApGpxDpA
SEQ ID NO:4	GpTpGpxDpApGpxDpA
SEQ ID NO:5	ApApDpxGpGpGpxDpGpGpxDpGpG
SEQ ID NO:6	CpApDpxGpGpxDpGpxDpGpG

FIGURE 26

31/42

P15 PROMOTER REGION SEQUENCE (SEQ ID NO:7)

tttttcattcagtcgaacttgccttcgcgaagctcacacatctgcctcgtgcaagattctcagtcatttttac
 ttagtcatttggttcttttcctatcaccattctttatgtcccccctcaaagaaaaacattatcttccatttccctatcaact
 ccaaacagcttttcattttttctgacataatttactacctaagaaaaatgggtcaagaattgggttagactatcttgccttaact
 tttctgataagtttcagagaaactcaaagggtcaaaacaagagcataagagtaaaggtagagaaattaagaaactgaagac
 taggaaatgggggttgggatgggaaagaaaaagaaattggttataatgctaccoggttcccttccctgtccaggtggattt
 cagctctgttgaggctctgtcagtagatattcagccctaaccagcacttccatgggtgggtggcacttccactgcctttaa
 aagaaagagctttttttaattctacagggatttgggggatgaggagtcagagctaagggtatcctaaaaaaacatgtgaa
 gactctcattttgcaatacacaaagcaattgccctcctgttaagactttgtcttccctcagcactccgaacaaaaatgattc
 tgtaacaaaaaattgttcacttttaggagaggtccacttatgcagtttccctacccaaagttttttaggcaacaaatccataa
 ctgcggttctcttccatccaatgtagcatccgctgaaatgttttaaatatttttaagtaataaatgttgattcaaactc
 acctaggaagatttaggaaggggaaaaaaagcacttggcattttaaatcttcagaagagaatttaattgacaggttcagcctg
 tttaatgacaagcccagcaccacacccctctcttatgatgtttcattattactgcataaatttcccttattactcatgat
 aaataaaaaataagatacctgacaaagtgggttttaaataggtaagagtgcaacaaagatttactgtacaaatatgatgaa
 actgggatctcagattccttaaagtataatttttttgtcttatgtgtgcccaggttggcactctcaatctcgaactagttt
 tttctcttttaaggggtgtatccataatgcaaaaatggaaagaattaaaaagcacacgcaaaacatgattctcgggatt
 tttctctatttttatgggttgactaattcaaacagaaagacacatccaagagaaaattgctaagtttgatacaagttatga
 aacttgtgaagcccaagtactgcctggggatgaatttaacttgtatgacaggtgcagagctgtcgttttcagacatctta
 agaaagacggagttattttgatgacttctcctcggtcacaggggagccaccaacgtctccacagtgaacccaactgggt
 ggctgaaggaacagaaatcctctgctccgcctactggggatttaggagctgagggcagtggtgaacattcccaaaatatta
 gccttggttttactggacatccagcagcagtgagccagcattcctggcggtccctggcccagctctctggcgcatgcg
 tcttagcatctttgggcaggttccccgcctcgtgacgcgtcgcccggttggcctcccggtgatcacagcggacag
 gggcgaggcctaagggggtggggagacgcccggcccttggccagctgaaaacggaattctttgcccgttggtcccca
 ctctgccagagcagggcggggcagtgaggactccgcgacgcgtccgcacccctggggccagagcggctttgagctcggctg
 cgtccgcgttaggcgttttttccagaagcaatccaggcgccgcctggttcttgagcgccaggaaaagcccgaggacta
 acgaccggccgctcggccactgcacggggccccaagccgcagaaaggacgagggagggtaatgaagctgagccaggtct
 cctaggaaggagagagtgcccgaggcagcgtgggaaagaagggaagagtgctcgtaagtttacggccaacggtggatta
 tccggggcgctgcgcgtctgggggtgcggaatgcgcgaggagaacaagggtcatgccagtgggggcgccgagcg

SEQ ID NO:8 TTCGCGAA
 SEQ ID NO:9 TGCCTCGT
 SEQ ID NO:10 TGCCGGCT
 SEQ ID NO:11 CGGCCCGG
 SEQ ID NO:12 GCTCGGCT

FIGURE 27

32/42

P16 PROMOTER REGION SEQUENCE (SEQ ID NO:13)

cggagaggggggagaacagacaacgggcgggcggggagcagcatggagccggcgggcggggagcagcatggag
ccttcggctgactggctggccacggccgcccgggggtcgggtagaggaggtgcgggcgctgctggaggcggggcgct
gcccacgcacccaatagttacggtcggaggccgatccaggtcatgatgatgggcagcggcccgagtggcgggagctgctgc
tgctccacggcgcgaggcccaactgcgcgaccccgccactctcaccgacccgtgcacgacgctgccgggagggcttc
ctggacacgctgggtggtgctgcacggggcgggcggggtggacgtgcgcgatgcctggggccgtctgccgtggacct
ggctgaggagctgggcatcgcatgtgcacgggtacctgcgcggcgctgcggggggcaccagaggcagtaacctatgcc
gcatagatgccgggaaggtccctcagacatcccgattgaaagaaccagagaggctctgagaaacctcgggaaacttag
atcatcagtcacccaaggtcctacagggccacaactgcccccgccacaacccaccccgcttctgtagttttcatttagaa
aatagagcttttaaaaatgtccttgccttttaacgtagatataagccttccccactaccgtaaatgtccatttatatcat
tttttatataattottataaaaatgtaaaaaagaaaaacaccgcttctgccttttcaactgtgttgagttt
tctggagtgagcactcacgccctaagcgacattcatgtgggcattttcttgagagcctcgagcctccggaagctgtcga
cttcatgacaagcattttgtgaactaggggaagctcaggggggttactggcttctcttgagtcacaactgctagcaaattggc
agaaccaaagctcaaataaaaataaaataattttcattcattcactc

SEQ ID NO:14 AACGGGCGGCGG
SEQ ID NO:15 CACGGCGCGG
SEQ ID NO:16 CGGGCGGC
SEQ ID NO:17 AGCAGCAT
SEQ ID NO:18 GCGCCGAC

FIGURE 28

33/42

BRCA1 PROMOTER REGION SEQUENCE (SEQ ID NO:19)

ccacctaatgtactgaattgcaaattgatagttgttctagcagtgagagataaaagaaaaaaagtaca
accaaagtccagtcagggcacagcagaaaacctacaactcatggaaggtaaagaacctgcaactggagccaagaagagtaac
aagccaaatgaacagacaagtaaaagacatgacagcgatactttcccagagctgaagttaacaaatgcacctgggttcttt
tactaagtgttcaaataccagtgaaacttaagaatttgtcaatcctagccttccaagagaagaaaaagaagagaaactag
aaacagttagtgtcctaataatgctgaagaccccaaagatctcatgttaagtggagaaagggttttgcaactgaaagatc
tgtagagagtagcagttatttcattgggtacctgggtactgattatggcactcaggaaagtatctcgttactggaagttagca
ctctaggggaaggcaaaaacagaaccaaataaatgtgtgagtcagtggtgcagcatttgaaaacccaagggaactaattcat
ggttgttccaaagataatagaaatgacacagaaggctttaagtatccattgggacatgaagttaaccacagtcgggaaac
aagcatagaaatggaagaa

SEQ ID NO:20 AGGCACAGCA
SEQ ID NO:21 GTGCAGCA
SEQ ID NO:22 CAGCGATA
SEQ ID NO:23 TAGCAGTG
SEQ ID NO:24 AGGCTTTA

FIGURE 29

34/42




FIGURE 30

35/42




FIGURE 31

36/42




FIGURE 32

37/42




FIGURE 33

38/42



FIGURE 34

39/42




FIGURE 35

40/42




FIGURE 36

41/42




FIGURE 37

42/42




FIGURE 38