

INFLUENZA VIRUS VACCINES AND USES THEREOF

INTRODUCTION

The invention relates to the field of medicine. Provided herein are influenza
5 hemagglutinin stem domain polypeptides, methods for providing hemagglutinin stem
domain polypeptides, compositions comprising the same, vaccines comprising the same and
methods of their use, in particular in the detection, prevention and/or treatment of influenza.

BACKGROUND

10 Influenza viruses are major human pathogens, causing a respiratory disease
(commonly referred to as “influenza” or “the flu”) that ranges in severity from sub-
clinical infection to primary viral pneumonia which can result in death. The clinical
effects of infection vary with the virulence of the influenza strain and the exposure,
history, age, and immune status of the host. Every year it is estimated that approximately
15 1 billion people worldwide undergo infection with influenza virus, leading to severe
illness in 3-5 million cases and an estimated 300,000 to 500,000 of influenza related
deaths. The bulk of these infections can be attributed to influenza A viruses carrying H1
or H3 hemagglutinin subtypes, with a smaller contribution from Influenza B viruses, and
therefore representatives of all three are included in the seasonal vaccine. The current
20 immunization practice relies on early identification of circulating influenza viruses to
allow for timely production of an effective seasonal influenza vaccine. Apart from the
inherent difficulties in predicting the strains that will be dominant during the next season,
antiviral resistance and immune escape also play a role in failure of current vaccines to
prevent morbidity and mortality. In addition to this the possibility of a pandemic caused
25 by a highly virulent viral strain originating from animal reservoirs and reassorted to
increase human to human spread, poses a significant and realistic threat to global health.

Influenza A viruses are widely distributed in nature and can infect a variety of birds
and mammals. Influenza viruses are enveloped RNA viruses that belong to the family of
Orthomyxoviridae. Their genomes consist of eight single-stranded RNA segments that code
30 for 11 different proteins, one nucleoprotein (NP), three polymerase proteins (PA, PB1, and
PB2), two matrix proteins (M1 and M2), three non-structural proteins (NS1, NS2, and PB1-

F2), and two external glycoproteins: hemagglutinin (HA) and neuraminidase (NA). The viruses are classified on the basis of differences in antigenic structure of the HA and NA proteins, with their different combinations representing unique virus subtypes that are further classified into specific influenza virus strains. Although all known subtypes can be found in birds, currently circulating human influenza A subtypes are H1N1 and H3N2. Phylogenetic analysis has demonstrated a subdivision of hemagglutinins into two main groups: *inter alia* the H1, H2, H5 and H9 subtypes in phylogenetic group 1 and *inter alia* the H3, H4 and H7 subtypes in phylogenetic group 2.

The influenza type B virus strains are strictly human. The antigenic variation in HA within the influenza type B virus strains is smaller than those observed within the type A strains. Two genetically and antigenically distinct lineages of influenza B virus are circulating in humans, as represented by the B/Yamagata/16/88 (also referred to as B/Yamagata) and B/Victoria/2/87 (B/Victoria) lineages (Ferguson et al., 2003). Although the spectrum of disease caused by influenza B viruses is generally milder than that caused by influenza A viruses, severe illness requiring hospitalization is still frequently observed with influenza B infection.

It is known that antibodies that neutralize the influenza virus are primarily directed against hemagglutinin (HA). Hemagglutinin or HA is a trimeric glycoprotein that is anchored to the viral coat and has a dual function: it is responsible for binding to the cell surface receptor sialic acid and, after uptake, it mediates the fusion of the viral and endosomal membrane leading to release of the viral RNA in the cytosol of the cell. HA comprises a large head domain and a smaller stem domain. Attachment to the viral membrane is mediated by a C-terminal anchoring sequence connected to the stem domain. The protein is post-translationally cleaved in a designated loop to yield two polypeptides, HA1 and HA2 (the full sequence is referred to as HA0). The membrane distal head region is mainly derived from HA1 and the membrane proximal stem region primarily from HA2 (FIG. 1).

The reason that the seasonal influenza vaccine must be updated every year is the large variability of the virus. In the hemagglutinin molecule this variation is particularly manifested in the head domain where antigenic drift and shift have resulted in a large number of different variants. Since this is also the area that is immunodominant, most

neutralizing antibodies are directed against this domain and act by interfering with receptor binding. The combination of immunodominance and large variation of the head domain also explains why infection with a particular strain does not lead to immunity to other strains: the antibodies elicited by the first infection only recognize a limited number of strains closely related to the virus of the primary infection.

Recently, influenza hemagglutinin stem domain polypeptides, lacking all or substantially all of the influenza hemagglutinin globular head domain, have been described and used to generate an immune response to one or more conserved epitopes of the stem domain polypeptide. It is believed that epitopes of the stem domain polypeptide are less immunogenic than the highly immunogenic regions of a globular head domain, thus the absence of a globular head domain in the stem domain polypeptide might allow an immune response against one or more epitopes of the stem domain polypeptide to develop (Steel et al., 2010). Steel et al. thus have created a new molecule by deleting amino acid residue 53 to 276 of HA1 of the A/Puerto Rico/8/1934 (H1N1) and A/Hong Kong/1968 (H3N2) strains from the HA primary sequence, and replacing this by a short flexible linking sequence GGGG. Vaccination of mice with the H3 HK68 construct did not elicit antisera that were cross-reactive with group 1 HAs. In addition, as shown in PCT/EP2012/073706, the stem domain polypeptides were highly unstable and did not adopt the correct conformation as proven by the lack of binding of antibodies that were shown to bind to conserved epitopes in the stem region.

In addition, Bommakanti et al. (2010) described an HA2 based polypeptide comprising amino acid residues 1-172 of HA2, a 7-amino acid linker (GSAGSAG), amino acid residues 7-46 of HA1, a 6-amino acid linker GSAGSA, followed by residues 290-321 of HA1, with the mutations V297T, I300E, Y302T and C305T in HA1. The design was based on the sequence of H3 HA (A/Hong Kong/1968). The polypeptide did only provide cross-protection against another influenza virus strain within the H3 subtype (A/Phil/2/82 but not against an H1 subtype (A/PR/8/34). In a more recent paper by Bommakanti et al (2012) a stem domain sequence based on HA from H1N1 A/Puerto Rico/8/1934 (H1HA0HA6) is described. In this polypeptide the equivalent of residues 55 to 302 have been deleted and mutations I311T, V314T, I316N, C319S, F406D, F409T, and L416D have been made. Both the H3 and HA based polypeptides were expressed in

E. coli and therefore lack the glycans that are a part of the naturally occurring HA proteins. When expressed in E.coli the polypeptide is recovered mainly as high molecular weight aggregates and a minor monomeric fraction. The polypeptide binds CR6261 with two apparent dissociation constants of 9 and 0.2 μ M. The authors show that mice can survive a challenge with 1 LD90 of the homologous H1N1 A/Puerto Rico/8/1934 virus after immunization (twice, 4 week interval) with 20 μ g of protein adjuvanted with 100 μ g of CpG7909. The authors also describe circularly permuted polypeptides comparable to those described above for A/Hong Kong/1/1968 derived polypeptides. These polypeptides are derived from HA's from H1N1 A/Puerto Rico/8/1934, H1N1 A/North Carolina/20/99 or H1N1 A/California/07/2009 and can provide partial protection in a mild challenge (1LD90) model in mice of H1N1 A/Puerto Rico/8/1934 (i.e. within the same subtype). Sera from guinea pigs immunized with these polypeptides did not exhibit detectable levels of neutralization when tested in a neutralization assay.

More recently Lu et al (2013) also described soluble stem domain polypeptides derived from the HA of H1N1 A/California/05/2009. In the final design the equivalent of residues 54-303 (numbering according to SEQ ID NO: 1) have been deleted (the leader sequence, residues 1-17 is also not present) and two mutations have been introduced in the B-loop of the protein, i.e. F407D, and L413D. Furthermore the polypeptide contained a C-terminal trimerization domain (foldon). In addition, two intermonomer disulfide bridges were introduced, one in the area of the trimeric foldon domain, and one at position 430 and 431. The polypeptide is produced in an E.coli based cell free system, (and thus lacks the glycans that are part of the naturally occurring HA proteins) and is recovered in a denatured form, which needs to be refolded prior to use. No immunological or protection from influenza challenge data were shown, so immunogenicity and efficacy of this polypeptide is not known.

In a recent paper Mallajosyula et al (2014) also report a stem domain polypeptide. In this design, based on the HA from H1N1 A/Puerto Rico/8/1934 not only a large part of the HA1 sequence is deleted (residue 42 to 289, numbering according to SEQ ID NO: 1), but also large part of the N- and C-terminal sequences of HA2 (residues 344 to 383 and 457 to 565, respectively). It is noteworthy that in H3 HA proteins the deleted part contains broadly neutralizing epitopes, e.g. those of CR8020 and CR8043. The

polypeptide again contains a foldon trimerization domain at the C-terminus and is also produced in *E. coli*, so lacks the naturally occurring glycans on viral HA. The polypeptide binds the broadly neutralizing antibodies and is CR6261, F10 and FI6v3. The polypeptide was also tested in an influenza challenge model (1LD90 of H1N1 A/Puerto Rico/8/1934) and could protect mice from death. Equivalent polypeptides derived from HA of H1N1 A/New Caledonia/20/1999 and H1N1 A/California/04/2009 could also partially protect. An equivalent polypeptide derived from H5N1 A/ Viet Nam/1203/2004 only gave limited protection in this challenge model. Moreover, only one influenza strain was used to challenge the animals with a relatively low dose administered (1-2 LD90), so protection against multiple influenza strains, a prerequisite for a universal vaccine has not been established.

There thus still exists a need for a safe and effective universal vaccine that stimulates the production of a robust, broadly neutralizing antibody response and that offers protection against a broad set of current and future influenza virus strains (both seasonal and pandemic), in particular providing protection against one or more influenza A virus subtypes within phylogenetic group 1 and/or group 2, for effective prevention and therapy of influenza.

SUMMARY

Provided herein are influenza hemagglutinin stem domain polypeptides, methods for providing stem domain polypeptides, compositions comprising the same, vaccines comprising the same and methods of their use.

In a first aspect, the present invention provides novel immunogenic multimeric polypeptides comprising an influenza hemagglutinin stem domain and lacking the globular head, referred to as influenza hemagglutinin (HA) stem domain polypeptides. The multimeric polypeptides are capable of inducing an immune response when administered to a subject, in particular a human subject. The polypeptides of the invention present conserved epitopes of the membrane proximal stem domain HA molecule to the immune system in the absence of dominant epitopes that are present in the membrane distal head domain. To this end, part of the primary sequence of the HA0 protein making up the head domain is removed and the remaining amino acid sequence is

reconnected, either directly or, in some embodiments, by introducing a short flexible linking sequence ('linker') to restore the continuity of the amino acid chain. The resulting sequence is further modified by introducing specific mutations that stabilize the native 3-dimensional structure of the remaining part of the HA0 molecule. The polypeptides do not comprise the full-length HA1 and/or HA2 of an influenza virus.

The present invention provides novel multimeric influenza hemagglutinin stem domain polypeptides, wherein said multimeric polypeptides comprise at least a first and a second influenza hemagglutinin stem domain monomer, said first and second monomer each comprising: (a) an influenza hemagglutinin HA1 domain that comprises an HA1 N-terminal stem segment, covalently linked by a linking sequence of 0-50 amino acid residues to an HA1 C-terminal stem segment, wherein said HA1 C-terminal segment is linked to (b) an influenza hemagglutinin HA2 domain, wherein said HA1 N-terminal segment comprises the amino acids 1-x of HA1, preferably the amino acids p-x of HA1, and wherein the HA1 C-terminal stem segment comprises the amino acids y-C-terminal amino acid of HA1, and (c) wherein the polypeptide comprises no protease cleavage site at the junction between HA1 and HA2; and (d) wherein the first monomer is linked to said second monomer by a disulfide bridge between the amino acid on position 411 of the first monomer and the amino acid on position 419 of the second monomer.

According to the invention, the disulfide bridge thus forms a covalent cross-link between individual monomers in a multimer.

In certain embodiments, the multimeric polypeptide is trimeric, i.e. comprises three monomers. According to the invention, each monomer is linked to another monomer by the disulfide bridge between the amino acid on position 411 of one monomer to the amino acid on position 419 of another monomer. It is noted that the numbering used is in relation to SEQ ID NO: 1. A person skilled in the art will be able to determine the equivalent positions in other HA sequences.

In certain embodiments, the HA1 and HA2 domains are derived from an influenza A virus subtype derived from phylogenetic group 1.

In certain embodiments, the HA1 and HA2 domains are derived from an influenza A virus subtype comprising HA of the H1 subtype.

In certain embodiments, x= the amino acid on position 52 of SEQ ID NO: 1 (or an equivalent position in another hemagglutinin), p= the amino acid on position 18 of SEQ ID NO: 1 (or an equivalent position in another hemagglutinin) and y= the amino acid on position 321 of SEQ ID NO: 1 (or an equivalent position in another hemagglutinin). In certain embodiments, the HA1 N- terminal stem segment thus comprises the amino acids 1-52 of HA1, and the HA1 C- terminal stem segment comprises the amino acids 321-end (i.e. the C-terminal amino acid of HA1) of HA1. Thus, in certain embodiments, the deletion in the HA1 segment comprises the amino acid sequence from the amino acid at position 53 up to and including the amino acid at position 320. In certain embodiments, the polypeptides do not comprise the signal sequence. In certain embodiments, the HA1 N-terminal segment thus comprises the amino acid 18-52 of HA1, wherein p is the first amino acid of the mature HA molecule (e.g. p=18 in case of SEQ ID NO: 1).

In certain embodiments, the HA1 and HA2 domains are derived from an influenza A virus subtype derived from phylogenetic group 2.

In certain embodiments, the HA1 and HA2 domains are derived from an influenza A virus subtype comprising HA of the H3 subtype.

The multimeric polypeptides of the invention thus comprise at least two monomers, each monomer comprising a HA1 domain, said HA1 domain comprising a HA1 N-terminal segment, linked to, either directly or through a linking sequence to a HA1 C-terminal segment, and a HA2 domain. In certain embodiments, the N-terminal amino acid of the HA2 domain is directly linked to the C-terminal amino acid of the HA1 C-terminal segment.

In certain embodiments, the polypeptides comprise the complete HA2 domain, i.e. the HA2 domain including the transmembrane domain and the intracellular sequence. In certain embodiments, the HA2 domains of the monomers have been truncated. Thus, in certain embodiments, the polypeptides of the invention do not contain the intracellular sequences of HA and the transmembrane domain. In certain embodiments, the amino acid sequence from position (or the equivalent of) 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, or 530 of the HA2 domain to the C-terminus of the HA2 domain has been removed.

According to the invention, the C-terminal amino acid of the HA1 C-terminal stem segment is linked to the N-terminal amino acid of the HA2 domain, thus forming a junction between the HA1 and HA2 domain. The polypeptides of the invention do not comprise a protease cleavage site at the junction between the HA1 and HA2 domain. In certain
5 embodiments, the C-terminal amino acid residue of the HA1 C-terminal stem segment (amino acid 343 in SEQ ID NO: 1) is any amino acid other than arginine (R) or lysine (K), preferably glutamine (Q).

The influenza hemagglutinin stem domain monomers in the immunogenic polypeptides of the invention are substantially smaller than HA0, preferably lacking all or
10 substantially all of the globular head of HA. Preferably, the immunogenic monomers are no more than 360, preferably no more than 350, 340, 330, 320, 310, 305, 300, 295, 290, 285, 280, 275, or 270 amino acids in length. In certain embodiments, the immunogenic polypeptides are from about 250 to about 350, preferably from about 260 to about 340, preferably from about 270 to about 330.

15 The polypeptides of the invention do not comprise the full length HA1.

In certain embodiments, the polypeptides are glycosylated.

According to the invention, the polypeptides further comprise one or more additional mutations in the HA1 and/or HA2 domain, as compared to the amino acid sequence of the wild-type HA, in particular the HA on which the HA 1 and HA2 domains
20 are based.

In certain embodiments, the polypeptides of the invention comprise the conserved stem domain epitopes of the group 1 cross-neutralizing antibody CR6261 (as disclosed in WO2008/028946) and/or of the antibody CR9114 (as described in WO2013/007770), an antibody capable of binding to and neutralizing both group 1 and group 2 influenza A
25 viruses, as well as influenza B viruses. It is thus another aspect of the invention to provide HA stem domain polypeptides, wherein said polypeptides stably present the epitopes of the antibody CR6261 and/or CR9114, as indicated by binding of said antibody or antibodies to said polypeptides.

In certain embodiments, the polypeptides do not bind to CR8020 and CR8057
30 (described in WO 2010/130636), which are monoclonal antibodies that bind to H3 influenza viruses only. The influenza hemagglutinin stem domain polypeptides provided

herein are suitable for use in immunogenic compositions (e.g. vaccines) capable of generating immune responses against a plurality of influenza virus A and/or B strains.

In certain embodiments, the polypeptides of the invention comprise the conserved stem domain epitopes of the group 2 cross-neutralizing antibody CR8020 (described in WO
5 2010/130636).

In certain embodiments, the influenza hemagglutinin stem domain polypeptides are capable of generating immune responses against influenza A virus strains of phylogenetic group 1 and/or group 2, in particular against influenza virus strains of both phylogenetic group 1 and group 2. In an embodiment, the polypeptides are capable of generating an
10 immune response against homologous influenza virus strains. In an embodiment, the polypeptides are capable of generating an immune response against heterologous influenza virus strains of the same and/or different subtypes. In a further embodiment, the polypeptides are capable of generating an immune response to influenza virus strains of both phylogenetic group 1 and group 2 and influenza B virus strains.

15 The polypeptides according to the invention may be used e.g. in stand alone therapy and/or prophylaxis and/or diagnosis of a disease or condition caused by an influenza virus, in particular a phylogenetic group 1 or 2 influenza A virus and/or an influenza B virus, or in combination with other prophylactic and/or therapeutic treatments, such as (existing or future) vaccines, antiviral agents and/or monoclonal
20 antibodies.

In a further aspect, the present invention provides nucleic acid molecules encoding the influenza HA stem domain polypeptides. In yet another aspect, the invention provides vectors comprising the nucleic acids encoding the immunogenic polypeptides.

In a further aspect, the invention provides methods for inducing an immune
25 response in a subject, the method comprising administering to the subject a polypeptide and/or nucleic acid molecule according to the invention.

In another aspect, the invention provides immunogenic compositions comprising a polypeptide and/or a nucleic acid molecule according to the invention. The immunogenic compositions provided herein can be in any form that allows for the compositions to be
30 administered to a subject, e.g. mice, ferrets or humans. In a specific embodiment, the immunogenic compositions are suitable for human administration. The polypeptides,

nucleic acid molecules and compositions may be used in methods of preventing and/or treating an influenza virus disease and/or for diagnostic purposes. The compositions may further comprise a pharmaceutically acceptable carrier or excipient. In certain embodiments, the compositions described herein comprise, or are administered in combination with, an adjuvant.

In another aspect, the invention provides polypeptides, nucleic acids and/or immunogenic compositions for use as a vaccine. The invention in particular relates to immunogenic polypeptides, nucleic acids, and/or immunogenic compositions for use as a vaccine in the prevention and/or treatment of a disease or condition caused by an influenza virus A subtype of phylogenetic group 1 and/or 2 and/or influenza B virus.

The various embodiments and uses of the polypeptides according to the invention will become clear from the following detailed description of the invention.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows a model of the HA monomer in the pre-fusion state as present in the native trimer. HA 1 is shown in light grey, HA2 is shown in dark grey. Helix A (an important part of the epitope of CR6261) and helix CD (part of the trimer interface) are indicated, as is the loop connecting these secondary structure elements. The C-terminus of HA1 and the N-terminus of HA2 are also indicated. The fusion peptide is located at the N-terminus of HA2.

FIG. 2. CR9114 sandwich ELISA results for:

- (A) purified soluble HA from H1N1 A/Brisbane/59/2007 in trimeric and monomeric form
- (B) medium obtained from cultures expressing s127H1 (SEQ ID NO: 66) , s127H1-t2 (SEQ ID NO: 91); data for FL HA trimer and monomer are also show for reasons of comparison
- (C) double cysteine mutants variants of 55G7-t2 (SEQ ID NO: 166 to 176)
- (D) double cysteine variants of 127H1-t2 (SEQ ID NO: 177 to 187)

FIG. 3. Western blot of media of cultures expressing double cysteine variants of 55G7 (A, B) and 127H1-t2 (C,D) under reducing (A, C) and under non-reducing conditions (B, D). The numbers above each lane refer to the cluster of cysteine mutations as listed in table 10.

5

FIG. 4. Structural model of a polypeptide of the invention, indicating the positions of residues 411 and 419 that are mutated to cysteine in 127H1-t2-cl18. The model is created by deletion of residues 53 to 320 from the structure of the full length HA of H1N1 A/California/04/2009 (PDB 3LZG).

10

FIG. 5. (A). Elution profile from the preparative size exclusion column (Superdex 200) during purification of s127H1-t2-cl18long (SEQ ID NO: 181) with an additional C-terminal his-tag. Fractions are indicate below the figure.

(B) SDS-PAGE analysis of the fractions collected from the size exclusion column under non-reducing (left) and reducing conditions (right) . Numbers above the lanes correspond to figure 5A , M denotes a molecular weight marker.

15

(C) Native PAGE of fractions 1-4. Numbers above the lanes correspond to figure 5A , M denotes a molecular weight marker.

(D) Western blot of fractions 1-4, using a polyclonal anti-his-tag antibody for detection.

20

Numbers above the lanes correspond to figure 5A , M denotes a molecular weight marker.

FIG. 6. Elisa results for binding of broadly neutralizing antibodies CR6261 and CR9114 to the polypeptides in fraction 1 to 4 as indicated in Figure 5. Binding to a ployclonal anti-H1 HA serum and the group 2specific monoclonal antibody CR8020 are also included. For reasons of comparison results for a soluble HA from H1N1 A/Brisbane/59/2007 in monomeric and trimeric form are also shown.

25

FIG. 7. CR9114 sandwich ELISA results for fractions 1 to 4 as indicated in figure 5. For reasons of comparison results for a soluble HA from H1N1 A/Brisbane/59/2007 in monomeric and trimeric form are also shown.

30

FIG. 8. SEC-MALS results for the polypeptide of fraction 3 in the absence and presence of Fab fragments of CR8020 (top panel), CR6261 or CR9114 (both bottom panel).

Complex formation with Fab fragments of CR9114 and CR6261 leads to an increase in molecular mass and shift of the peak.

FIG 9. Binding of broadly neutralizing antibodies CR6261 and CR9114 to s127H1-t2-cl18long as determined by bilayer interferometry. A and C show individual binding curves for immobilized monoclonal antibodies exposed to varying concentrations of s127H1-t2-cl18long; B and D show the steady state analysis used to estimate K_d .

FIG. 10. Survival (A), relative body weight loss (B) and clinical score (C) for the negative (PBS, 3 immunizations at 3 weeks intervals) and positive control (15 mg/kg CR6261, 1 day before challenge) groups. Mice were challenged four week after the last immunization with a lethal dose (25xLD50) of H1N1 A/Puerto Rico/8/34 and monitored for 21 days. Error bars indicate 95% confidence interval (B) or interquartile range (C).

FIG 11. Survival for groups immunized 1 time (A), 2 times (B) or 3 times (C) with 30 μ g s127H1-t2-cl18long in the presence of 10 μ g Matrix-M. Mice were challenged four week after the last immunization with a lethal dose (25xLD50) of H1N1 A/Puerto Rico/8/34 and monitored for 21 days. For reasons of comparison the negative control group (PBS) is also shown.

FIG 12. Relative body weight change for groups immunized 1 time (A), 2 times (B) or 3 times with 30 μ g s127H1-t2-cl18long in the presence of 10 μ g Matrix-M. Mice were challenged four week after the last immunization with a lethal dose (25xLD50) of H1N1 A/Puerto Rico/8/34 and monitored for 21 days. For reasons of comparison the negative control group (PBS) is also shown. Error bars indicate 95% confidence interval.

FIG 13. Clinical scores for groups immunized 1 time (A), 2 times (B) or 3 times with 30 μ g s127H1-t2-cl18long in the presence 10 μ g Matrix-M. Mice were challenged four week

after the last immunization with a lethal dose (25xLD₅₀) of H1N1 A/Puerto Rico/8/34 and monitored for 21 days. For reasons of comparison the negative control group (PBS) is also shown. Error bars indicate interquartile range.

5 FIG 14. ELISA results for pre-challenge serum (4 weeks after the final immunization) of the negative control and experimental groups using s127H1-t2-cl18long (A) or a soluble form of Full length HA (B) as the antigen. Bars represent median.

FIG 15. The antibodies induced 4 weeks after the final immunization (pre-challenge
10 timepoint) after immunization with Matrix-M adjuvated polypeptide of the invention s127H1-t2-cl18long are capable of competing with CR9114 for binding to full length HA from H1N1 A/Brisbane/59/07 in a competition ELISA (top). For reasons of comparison competition levels by unlabeled CR9114 (i.e. self-competition) and the non-binding monoclonal antibodies CR8020, both serially diluted from 5 µg/ml starting concentration,
15 are indicated in a separate graph bottom. Bars represent median.

FIG 16. (A) Survival for the negative (PBS, 3 immunizations at 3 weeks intervals) and positive control (15 mg/kg CR6261, 1 day before challenge) groups. Mice were challenged four week after the last immunization with a lethal dose (12.5xLD₅₀) of
20 H5N1 A/Hong Kong/156/97. (B) Survival, (C) relative body weight change and (D) median clinical scores for the group immunized 3 times with 30 µg s127H1-t2-cl18long in the presence of 10 µg Matrix-M. Error bars indicate 95% confidence interval (C) or interquartile range (D) .Mice were challenged four week after the last immunization with a lethal dose (12.5xLD₅₀) of H5N1 A/Hong Kong/156/97 and monitored for 21 days. For
25 reasons of comparison the negative control group (PBS) is also shown in B, C, D.

FIG 17. Elisa results for sera from mice immunized 3 times with polypeptide of the invention s127H1-t2-cl18long as described in example 7 using full length HA's from a number of group I (H1, H5 and H9) and group II (H3 and H7) influenza strains as the
30 antigen.

FIG 18. (A) Survival for the negative (PBS, 3 immunizations at 3 weeks intervals) and positive control (15 mg/kg CR6261, 1 day before challenge) groups. Mice were challenged four week after the last immunization with a lethal dose (12.5xLD50) of H1N1 A/Brisbane/59/2007. (B) Survival, (C) relative body weight change and (D) median clinical scores for the group immunized 3 times with 30 µg s127H1-t2-cl18long in the presence of 10 µg Matrix-M. Error bars indicate 95% confidence interval (C) or interquartile range (D). Mice were challenged four week after the last immunization with a lethal dose (12.5xLD50) of H1N1 A/Brisbane/59/2007 and monitored for 21 days. For reasons of comparison the negative control group (PBS) is also shown in B, C, D.

FIG 19. Pseudoparticle neutralizations assay using sera from mice immunized with polypeptide of the invention s127H1-t2-cl18long or PBS. Neutralization is observed at high serum concentrations for serum from animals immunized with a polypeptide of the invention.

FIG 20. Antibody Dependent Cellular Cytotoxicity (ADCC) surrogate assay. Sera from mice immunized with polypeptide of the invention s127H1-t2-cl18long exhibit a 30-40 fold induction of FcγRIV signaling activity at the highest serum concentrations using target cells transfected with FL HA from H5N1 A/Hong Kong/156/97 (A) or H1N1 A/Brisbane/59/07 (B) as the source of antigen.

FIG. 21. A: CR9114 sandwich ELISA to detect multimeric forms of polypeptides of the invention (indicated by their SEQ ID NO). Culture medium was diluted in 3 fold steps and analyzed. B: CR9114 binding to polypeptides of the invention (indicated by their SEQ ID NO) by ELISA. Culture medium was diluted in 3 fold steps and analyzed. C: CR6261 binding to polypeptides of the invention (indicated by their SEQ ID NO) by ELISA. Culture medium was diluted in 3 fold steps and analyzed. D: CR8020 binding to polypeptides of the invention (indicated by their SEQ ID NO) by ELISA. Culture medium was diluted in 3 fold steps and analyzed. For the purified proteins (SEQ ID NO: 186 and the Full length HA in trimeric and monomeric form) a starting concentration of 5

µg /ml was used. All polypeptides contain a C-terminal factor X cleavable his tag sequence.

FIG. 22. Western Blot of supernatant of cultures expressing polypeptides of the invention (indicated by their SEQ ID NO) Under reducing conditions the intermonomer disulfide bridges are reduced and the polypeptides of the invention are monomeric on the gel (left panel). Under oxidizing conditions intermonomer disulfide bridges remain intact and the polypeptides of the invention appear as trimers on the gel. For SEQ ID NO: 186 purified trimer was used in this experiment.

FIG. 23. Serum IgG ELISA titers (antigen full length HA) obtained after immunization of NHP as described in example 16. HA's in panel A-F are derived from Group 1 influenza strains (identified in the panels), HA's in panel G-J are derived from Group 2 Influenza strains. Data were analyzed using a slope based weighted average approach. Open symbols denote measurements at LOD. Bars denote medians.

FIG. 24. Serum IgG CR9114 competition binding obtained after immunization of NHP as described in Example 16. FL HA from 3 different strains was used as identified in panel A-C. Data shown are group medians, error bars denote interquartile range.

FIG. 25. Surrogate ADCC activity determined using ADCC Bioassay effector cells in serum obtained after immunization of NHP as described in example 16. Cells transfected with DNA expressing FL HA from 3 different strains was used (identified in panel A-C). Data are expressed as fold induction, which is signal of each measurement compared to background signal in the absence of serum. Lines denote individual animals, symbols denote mean of duplicates per serum concentration.

FIG. 26. Microneutralization assay using serum obtained after immunization of NHP as described in Example 16. Read-out was ELISA based quantification of NP in fixed cells 16 hours after incubation with 100 TCID₅₀ H5N1 A/HK/156/97 per well in the presence

of serially diluted serum. Lines denote individual animals, symbols denote mean of duplicates per serum concentration.

FIG. 27.Area under the curve temperature increase of non-human primates included in the experiment described in Example 16. Per animal, a reference 24-hour body temperature cycle was reconstructed using an 21-day window prior to start of the immunizations. The net increase in body temperature during the 21 day post -challenge follow-up period was calculated by subtracting the reference body temperature with added upper limit of the 95% CI from the body temperature measured during the post-challenge follow-up period . The AUC of the net temperature increase was subsequently calculated at intervals of day 0-3, day 0-8 and day 0-21. Statistical analysis between treatments was performed using pairwise t-test with Tukey-Kramer adjustment for multiple comparisons. Bars denote median. No data animal J10014 (SEQ ID NO: 186 + Matrix-M group) due to data logger failure. Animal Ji0403061 (Inflexal group) died at the end of day 8 and was excluded from the day 0-21 interval calculation.

FIG. 28.Survival (A) and % body weight change (B) of mice after serum transfer and challenge with H5N1 A/Hong Kong/156/97 as described in Example 17.

FIG. 29.Full length HA (H1N1 A/Brisbane/59/2007) ELISA titers of donor mice (D) at day 70, and recipient mice (R) prior to serum transfer (day -4) or challenge (day 0). Data were analyzed using a slope based weighted average approach. Open symbols denote measurements at LOD. Bars denote medians.

FIG. 30.Survival (A) and % body weight change (B) of mice after immunization and challenge with H1N1 A/Brisbane/59/2007 as described in Example 18.

FIG. 31. A: Full length HA (H1N1 A/Brisbane/59/2007) ELISA titers of mice immunized as described in Example 18. Data were analyzed using a slope based weighted average approach. Open symbols denote measurements at LOD. Bars denote medians. B: Serum IgG CR9114 competition binding obtained after immunization mice as described in

Example 18. FL HA from H1N1 A/Brisbane/59/2007 was used as the antigen. Data shown are group medians, error bars denote interquartile range.

5 FIG. 32. Survival (A) and % body weight change (B) of mice after immunization and challenge with H1N1 A/Netherlands/602/09 as described in Example 19.

FIG. 33. A: Full length HA (H1N1 A/Brisbane/59/2007) ELISA titers of mice immunized as described in Example 19. Data were analyzed using a slope based weighted average approach. Open symbols denote measurements at LOD. Bars denote medians. B: Serum
10 IgG CR9114 competition binding obtained after immunization mice as described in example 19. FL HA from H1N1 A/Brisbane/59/2007 was used as the antigen. Data shown are group medians, error bars denote interquartile range.

15 FIG. 34. Survival (A) and % body weight change (B) of mice after immunization and challenge with H5N1 A/Hong Kong/156/97 as described in Example 20.

FIG. 35. A: Full length HA (H1N1 A/Brisbane/59/2007) ELISA titers of mice immunized as described in Example 20. Data were analyzed using a slope based weighted average approach. Open symbols denote measurements at LOD. Bars denote medians. B: Serum
20 IgG CR9114 competition binding obtained after immunization mice as described in Example 20. FL HA from H1N1 A/Brisbane/59/2007 was used as the antigen. Data shown are group medians, error bars denote interquartile range.

25 FIG. 36. Survival (A) and % body weight change (B) of mice after immunization and challenge with H1N1 A/Puerto Rico/8/34 as described in Example 21.

FIG. 37. A: Full length HA (H1N1 A/Brisbane/59/2007) ELISA titers of mice immunized as described in Example 21. Data were analyzed using a slope based weighted average approach. Open symbols denote measurements at LOD. Bars denote medians. B: Serum
30 IgG CR9114 competition binding obtained after immunization mice as described in

Example 21. FL HA from H1N1 A/Brisbane/59/2007 was used as the antigen. Data shown are group medians, error bars denote interquartile range.

FIG. 38: Western Blot (polyclonal anti H1) of Hek293F cell culture supernatant after transient transfection with polypeptides of the invention.

DEFINITIONS

Definitions of terms as used in the present invention are given below.

An amino acid according to the invention can be any of the twenty naturally occurring (or 'standard' amino acids) or variants thereof, such as e.g. D-proline (the D-enantiomer of proline), or any variants that are not naturally found in proteins, such as e.g. norleucine. The standard amino acids can be divided into several groups based on their properties. Important factors are charge, hydrophilicity or hydrophobicity, size and functional groups. These properties are important for protein structure and protein-protein interactions. Some amino acids have special properties such as cysteine, that can form covalent disulfide bonds (or disulfide bridges) to other cysteine residues, proline that forms a cycle to the polypeptide backbone, and glycine that is more flexible than other amino acids. Table 1 shows the abbreviations and properties of the standard amino acids.

The term "amino acid sequence identity" refers to the degree of identity or similarity between a pair of aligned amino acid sequences, usually expressed as a percentage. Percent identity is the percentage of amino acid residues in a candidate sequence that are identical (i.e., the amino acid residues at a given position in the alignment are the same residue) or similar (i.e., the amino acid substitution at a given position in the alignment is a conservative substitution, as discussed below), to the corresponding amino acid residue in the peptide after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence homology. Sequence homology, including percentages of sequence identity and similarity, are determined using sequence alignment techniques well-known in the art, such as by visual inspection and mathematical calculation, or more preferably, the comparison is done by comparing sequence information using a computer program. An exemplary, preferred computer program is the Genetics Computer Group

(GCG; Madison, Wis.) Wisconsin package version 10.0 program, 'GAP' (Devereux et al. (1984)).

"Conservative substitution" refers to replacement of an amino acid of one class is with another amino acid of the same class. In particular embodiments, a conservative substitution does not alter the structure or function, or both, of a polypeptide. Classes of amino acids for the purposes of conservative substitution include hydrophobic (e.g. Met, Ala, Val, Leu), neutral hydrophilic (e.g. Cys, Ser, Thr), acidic (e.g. Asp, Glu), basic (e.g. Asn, Gln, His, Lys, Arg), conformation disrupters (e.g. Gly, Pro) and aromatic (e.g. Trp, Tyr, Phe).

As used herein, the terms "disease" and "disorder" are used interchangeably to refer to a condition in a subject. In some embodiments, the condition is a viral infection, in particular an influenza virus infection. In specific embodiments, a term "disease" refers to the pathological state resulting from the presence of the virus in a cell or a subject, or by the invasion of a cell or subject by the virus. In certain embodiments, the condition is a disease in a subject, the severity of which is decreased by inducing an immune response in the subject through the administration of an immunogenic composition.

As used herein, the term "effective amount" in the context of administering a therapy to a subject refers to the amount of a therapy which has a prophylactic and/or therapeutic effect(s). In certain embodiments, an "effective amount" in the context of administration of a therapy to a subject refers to the amount of a therapy which is sufficient to achieve a reduction or amelioration of the severity of an influenza virus infection, disease or symptom associated therewith, such as, but not limited to a reduction in the duration of an influenza virus infection, disease or symptom associated therewith, the prevention of the progression of an influenza virus infection, disease or symptom associated therewith, the prevention of the development or onset or recurrence of an influenza virus infection, disease or symptom associated therewith, the prevention or reduction of the spread of an influenza virus from one subject to another subject, the reduction of hospitalization of a subject and/or hospitalization length, an increase of the survival of a subject with an influenza virus infection or disease associated therewith, elimination of an influenza virus infection or disease associated therewith, inhibition or reduction of influenza virus replication, reduction of influenza virus titer; and/or

enhancement and/or improvement of the prophylactic or therapeutic effect(s) of another therapy. In certain embodiments, the effective amount does not result in complete protection from an influenza virus disease, but results in a lower titer or reduced number of influenza viruses compared to an untreated subject. Benefits of a reduction in the titer, number or total burden of influenza virus include, but are not limited to, less severe symptoms of the infection, fewer symptoms of the infection and a reduction in the length of the disease associated with the infection.

The term "host", as used herein, is intended to refer to an organism or a cell into which a vector such as a cloning vector or an expression vector has been introduced. The organism or cell can be prokaryotic or eukaryotic. Preferably, the host comprises isolated host cells, e.g. host cells in culture. The term "host cells" merely signifies that the cells are modified for the (over)-expression of the polypeptides of the invention. It should be understood that the term host is intended to refer not only to the particular subject organism or cell but to the progeny of such an organism or cell as well. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent organism or cell, but are still included within the scope of the term "host" as used herein.

The term "included" or "including" as used herein is deemed to be followed by the words "without limitation".

As used herein, the term "infection" means the invasion by, multiplication and/or presence of a virus in a cell or a subject. In one embodiment, an infection is an "active" infection, i.e., one in which the virus is replicating in a cell or a subject. Such an infection is characterized by the spread of the virus to other cells, tissues, and/or organs, from the cells, tissues, and/or organs initially infected by the virus. An infection may also be a latent infection, i.e., one in which the virus is not replicating. In certain embodiments, an infection refers to the pathological state resulting from the presence of the virus in a cell or a subject, or by the invasion of a cell or subject by the virus.

Influenza viruses are classified into influenza virus types: genus A, B and C. The term "influenza virus subtype" as used herein refers to influenza A virus variants that are characterized by combinations of the hemagglutinin (H) and neuramidase (N) viral surface proteins. According to the present invention influenza virus subtypes may be referred to by

their H number, such as for example “influenza virus comprising HA of the H3 subtype”, “influenza virus of the H3 subtype” or “H3 influenza”, or by a combination of a H number and an N number, such as for example “influenza virus subtype H3N2” or “H3N2”. The term “subtype” specifically includes all individual “strains”, within each subtype, which
5 usually result from mutations and show different pathogenic profiles, including natural isolates as well as man-made mutants or reassortants and the like. Such strains may also be referred to as various “isolates” of a viral subtype. Accordingly, as used herein, the terms “strains” and “isolates” may be used interchangeably. The current nomenclature for human influenza virus strains or isolates includes the type (genus) of virus, i.e. A, B or C, the
10 geographical location of the first isolation, strain number and year of isolation, usually with the antigenic description of HA and NA given in brackets, e.g. A/Moscow/10/00 (H3N2). Non-human strains also include the host of origin in the nomenclature. The influenza A virus subtypes can further be classified by reference to their phylogenetic group. Phylogenetic analysis has demonstrated a subdivision of hemagglutinins into two main
15 groups: *inter alia* the H1, H2, H5 and H9 subtypes in phylogenetic group 1 (“group 1” influenza viruses) and *inter alia* the H3, H4, H7 and H10 subtypes in phylogenetic group 2 (“group 2” influenza viruses).

As used herein, the term “influenza virus disease” refers to the pathological state resulting from the presence of an influenza virus, e.g. an influenza A or B virus in a cell
20 or subject or the invasion of a cell or subject by an influenza virus. In specific embodiments, the term refers to a respiratory illness caused by an influenza virus.

As used herein, the term “nucleic acid” is intended to include DNA molecules (e.g., cDNA or genomic DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid can be single-
25 stranded or double-stranded. The nucleic acid molecules may be modified chemically or biochemically or may contain non-natural or derivatized nucleotide bases, as will be readily appreciated by those of skill in the art. Such modifications include, for example, labels, methylation, substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as uncharged linkages (e.g., methyl
30 phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.), charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), pendent moieties (e.g., polypeptides),

intercalators (e.g., acridine, psoralen, etc.), chelators, alkylators, and modified linkages (e.g., alpha anomeric nucleic acids, etc.). A reference to a nucleic acid sequence encompasses its complement unless otherwise specified. Thus, a reference to a nucleic acid molecule having a particular sequence should be understood to encompass its
5 complementary strand, with its complementary sequence. The complementary strand is also useful, e.g., for anti-sense therapy, hybridization probes and PCR primers.

As used herein, in certain embodiments the numbering of the amino acids in HA is based on the numbering of amino acids in HA0 of a wild type influenza virus, e.g. the numbering of the amino acids of the H1N1 influenza strain A/Brisbane/59/2007 (SEQ ID
10 NO: 1). As used in the present invention, the wording "the amino acid at position "x" in HA" thus means the amino acid corresponding to the amino acid at position x in HA0 of the particular wild type influenza virus, e.g. A/Brisbane/59/2007 (SEQ ID NO: 1; wherein the amino acids of the HA2 domain have been indicated in italics). It will be understood by the skilled person that equivalent amino acids in other influenza virus
15 strains and/or subtypes can be determined by multiple sequence alignment. Note that, in the numbering system used throughout this application 1 refers to the N-terminal amino acid of an immature HA0 protein (SEQ ID NO: 1). The mature sequence starts e.g. on position 18 of SEQ ID NO: 1. It will be understood by the skilled person that the leader sequence (or signal sequence) that directs transport of a protein during production (e.g.
20 corresponding to amino acids 1-17 of SEQ ID NO: 1), generally is not present in the final polypeptide, that is e.g. used in a vaccine. In certain embodiments, the polypeptides according to the invention thus comprise an amino acid sequence without the leader sequence, i.e. the amino acid sequence is based on the amino acid sequence of HA0 without the signal sequence.

25 "Polypeptide" refers to a polymer of amino acids linked by amide bonds as is known to those of skill in the art. As used herein, the term can refer to a single polypeptide chain linked by covalent amide bonds. The term can also refer to multiple polypeptide chains associated by non-covalent interactions such as ionic contacts, hydrogen bonds, Van der Waals contacts and hydrophobic contacts. Those of skill in the
30 art will recognize that the term includes polypeptides that have been modified, for example by post-translational processing such as signal peptide cleavage, disulfide bond

formation, glycosylation (e.g., N-linked and O-linked glycosylation), protease cleavage and lipid modification (e.g. S-palmitoylation).

"Stem domain polypeptide" refers to a polypeptide that comprises one or more polypeptide chains that make up a stem domain of a naturally-occurring (or wild-type) hemagglutinin (HA). Typically, a stem domain polypeptide is a single polypeptide chain (i.e. corresponding to the stem domain of a hemagglutinin HA0 polypeptide) or two polypeptide chains (i.e. corresponding to the stem domain of a hemagglutinin HA1 polypeptide in association with a hemagglutinin HA2 polypeptide). According to the invention, a stem domain polypeptide comprises one or more mutations as compared to the wild-type HA molecule, in particular one or more amino acid residues of the wild-type HA may have been substituted by other amino acids, not naturally occurring on the corresponding position in a particular wild-type HA. Stem domain polypeptides according to the invention can furthermore comprise one or more linking sequences, as described below.

The term "vector" denotes a nucleic acid molecule into which a second nucleic acid molecule can be inserted for introduction into a host where it will be replicated, and in some cases expressed. In other words, a vector is capable of transporting a nucleic acid molecule to which it has been linked. Cloning as well as expression vectors are contemplated by the term "vector", as used herein. Vectors include, but are not limited to, plasmids, cosmids, bacterial artificial chromosomes (BAC) and yeast artificial chromosomes (YAC) and vectors derived from bacteriophages or plant or animal (including human) viruses. Vectors comprise an origin of replication recognized by the proposed host and in case of expression vectors, promoter and other regulatory regions recognized by the host. Certain vectors are capable of autonomous replication in a host into which they are introduced (e.g., vectors having a bacterial origin of replication can replicate in bacteria). Other vectors can be integrated into the genome of a host upon introduction into the host, and thereby are replicated along with the host genome. As used herein, the term "wild-type" in the context of a virus refers to influenza viruses that are prevalent, circulating naturally and producing typical outbreaks of disease.

DETAILED DESCRIPTION

Influenza viruses have a significant impact on global public health, causing millions of cases of severe illness each year, thousands of deaths, and considerable economic losses. Current trivalent influenza vaccines elicit a potent neutralizing antibody response to the vaccine strains and closely related isolates, but rarely extend to more diverged strains within a subtype or to other subtypes. In addition, selection of the appropriate vaccine strains presents many challenges and frequently results in sub-optimal protection. Furthermore, predicting the subtype of the next pandemic virus, including when and where it will arise, is currently impossible.

Hemagglutinin (HA) is the major envelope glycoprotein from influenza A viruses which is the major target of neutralizing antibodies. Hemagglutinin has two main functions during the entry process. First, hemagglutinin mediates attachment of the virus to the surface of target cells through interactions with sialic acid receptors. Second, after endocytosis of the virus, hemagglutinin subsequently triggers the fusion of the viral and endosomal membranes to release its genome into the cytoplasm of the target cell. HA comprises a large ectodomain of ~500 amino acids that is cleaved by host- derived enzymes to generate 2 polypeptides that remain linked by a disulfide bond. The majority of the N-terminal fragment (HA1, 320-330 amino acids) forms a membrane-distal globular domain that contains the receptor-binding site and most determinants recognized by virus-neutralizing antibodies. The smaller C-terminal portion (HA2, ~180 amino acids) forms a stem-like structure that anchors the globular domain to the cellular or viral membrane. The degree of sequence homology between subtypes is smaller in the HA1 polypeptides (34% - 59% homology between subtypes) than in the HA2 polypeptide (51%- 80% homology). The most conserved region is the sequence around the cleavage site, particularly the HA2 N- terminal 23 amino acids, which is conserved among all influenza A virus subtypes (Lorieau et al., 2010). Part of this region is exposed as a surface loop in the HA precursor molecule (HA0), but becomes inaccessible when HA0 is cleaved into HA1 and HA2.

Most neutralizing antibodies bind to the loops that surround the receptor binding site and interfere with receptor binding and attachment. Since these loops are highly variable, most antibodies targeting these regions are strain-specific, explaining why current vaccines elicit such limited, strain-specific immunity. Recently, however, fully human monoclonal

antibodies against influenza virus hemagglutinin with broad cross-neutralizing potency were generated. Functional and structural analysis have revealed that these antibodies interfere with the membrane fusion process and are directed against highly conserved epitopes in the stem domain of the influenza HA protein (Throsby et al., 2008; Ekiert et al.
5 2009, WO 2008/028946, WO2010/130636, WO 2013/007770).

Stem domain polypeptides stably presenting the epitopes of these antibodies are described in the co-pending patent application PCT/EP2012/073706. At least some of the stem domain polypeptides described herein stably present the epitope of CR6261 and/or CR9114 and are immunogenic in mice. Further immunogenic stem domain polypeptides
10 stably presenting the epitope of CR6261 and/or CR9114 have been described in co-pending patent application PCT/EP2014/060997.

According to the present invention new multimeric HA stem domain polypeptides have been designed presenting these epitopes. These polypeptides can be used to create a universal epitope-based vaccine inducing protection against a broad range of influenza
15 strains. Like in the previously described stem domain polypeptides, the highly variable and immunodominant part, i.e. the head domain, is first removed from the full length HA molecule to create a stem domain polypeptide, also called mini-HA, in order to redirect the immune response towards the stem domain where the epitopes for the broadly neutralizing antibodies are located. The broadly neutralizing antibodies mentioned above
20 were used to confirm the presence of the neutralizing epitopes.

In certain embodiments, the new HA stem polypeptides of the present invention form stable trimers in solution while exposing the epitopes of the neutralizing antibodies CR6261 and/or CR9114 and/or CR8020. The additional interactions between individual monomers further stabilize the protein and the epitopes, leading to better presentation of
25 these epitopes when the polypeptides are used in vaccine.

The stem domain polypeptides of this invention are capable of presenting the conserved epitopes of the membrane proximal stem domain HA molecule to the immune system in the absence of dominant epitopes that are present in the membrane distal head domain. To this end, part of the primary sequence of the HA0 protein making up the head
30 domain is removed and reconnected, either directly or, in some embodiments, by introducing a short flexible linking sequence ('linker') to restore the continuity of the

polypeptide chain. The resulting polypeptide sequence is further modified by introducing specific mutations that stabilize the native 3-dimensional structure of the remaining part of the HA0 molecule.

The present invention provides novel multimeric influenza hemagglutinin stem domain polypeptides, wherein said multimeric polypeptides comprise at least a first and a second influenza hemagglutinin stem domain monomer, said first and second monomer each comprising: (a) an influenza hemagglutinin HA1 domain that comprises an HA1 N-terminal stem segment, covalently linked by a linking sequence of 0-50 amino acid residues to an HA1 C-terminal stem segment, wherein said HA1 C-terminal segment is linked to (b) an influenza hemagglutinin HA2 domain, wherein said HA1 N-terminal segment comprises the amino acids 1-x of HA1, preferably the amino acids p-x of HA1, and wherein the HA1 C-terminal stem segment comprises the amino acids y-C-terminal amino acid of HA1, and (c) wherein the polypeptide comprises no protease cleavage site at the junction between the HA1 and HA2 domains; and (d) wherein the first monomer is linked to said second monomer by a disulfide bridge between the amino acid on position 411 of the first monomer and the amino acid on position 419 of the second monomer.

According to the invention, the disulfide bridge thus forms a covalent cross-link between individual monomers in a multimer.

In certain embodiments, the multimeric polypeptide is trimeric, i.e. comprises three influenza stem domain monomers. According to the invention, each monomer is linked to another monomer by the disulfide bridge between the amino acid on position 411 of one monomer to the amino acid on position 419 of another monomer.

According to the present invention, it has surprisingly been found that the introduction of a novel disulfide bridge between the amino acid positions 411 and 419 (numbering according to SEQ ID NO: 1 or equivalent residues in hemagglutinin of other influenza viruses) of at least two different monomers results in a dimeric or, preferably, a trimeric polypeptide. In contrast to other published trimeric HA-stem structures this intermonomer disulfide linked trimer expresses well and folds spontaneously. It therefore does not require refolding procedures to reach its three-dimensional structure as has been described (Lu et al 2013).

In certain embodiments, the HA1 and HA2 domains are derived from an influenza A virus subtype derived from phylogenetic group 1.

In certain embodiments, the HA1 and HA2 domains are derived from an influenza A virus subtype comprising HA of the H1 subtype. The polypeptides of the invention do
5 not comprise the full length HA1.

In certain embodiments, the HA1 and HA2 domains are derived from an influenza A virus subtype derived from phylogenetic group 2.

In certain embodiments, the HA1 and HA2 domains are derived from an influenza A virus subtype comprising HA of the H3 subtype.

10 In certain embodiments, the influenza hemagglutinin stem domain monomers are substantially smaller than HA0, preferably lacking all or substantially all of the globular head of HA. Preferably, the immunogenic polypeptides are no more than 360, preferably no more than 350, 340, 330, 320, 310, 305, 300, 295, 290, 285, 280, 275, or 270 amino acids in length. In certain embodiments, the immunogenic polypeptides are from about 250 to
15 about 350, preferably from about 260 to about 340, preferably from about 270 to about 330 amino acids in length.

According to the invention, the "HA1 N-terminal segment" refers to a polypeptide segment that corresponds to the amino-terminal portion of the HA1 domain of an influenza hemagglutinin (HA) molecule. In certain embodiments, the HA1 N-terminal polypeptide
20 segment comprises the amino acids from position 1 to position x of the HA1 domain, wherein amino acid on position x is an amino acid residue within HA1. The term "HA1 C-terminal segment" refers to a polypeptide segment that corresponds to the carboxy-terminal portion of an influenza hemagglutinin HA1 domain. In certain embodiments, the HA1 C-terminal polypeptide segment comprises the amino acids from position y to and including
25 the C-terminal amino acid of the HA1 domain, wherein the amino acid on position y is an amino acid residue within HA1. According to the invention y is greater than x, thus a segment of the HA1 domain between the HA1 N-terminal segment and the HA1 C-terminal segment, i.e. between the amino acid on position x and the amino acid on position y of HA1, has been deleted, and in some embodiments, replaced by a linking sequence.

30 In certain embodiments the HA1 N-terminal stem segment comprises the amino acids 1-x of HA1, and the HA1 C-terminal stem segment comprises the amino acids y-end

of HA1. Thus, in certain embodiments, the deletion in the HA1 segment comprises the amino acid sequence from the amino acid at position x+1 up to and including the amino acid at position y-1.

In certain embodiments, the polypeptides do not comprise the signal sequence. Thus
5 in certain embodiments, the HA1 N-terminal segment comprises the amino acid p-x of HA1, wherein p is the first amino acid of the mature HA molecule (e.g. p=18 in case of SEQ ID NO: 1). The skilled person will be able to prepare the polypeptides described herein without the signal peptides (e.g. amino acids 1-17 of SEQ ID NO: 1).

It is again noted that the numbering of amino acid positions used herein refers to
10 SEQ ID NO: 1. The skilled person will be able to determine the equivalent positions in other hemagglutinin sequences.

In certain embodiments, the polypeptides comprise the complete HA2 domain, thus including the transmembrane and intracellular sequences. In other embodiments, the polypeptides of the invention do not comprise the intracellular sequences of HA and the
15 transmembrane domain. Thus, in certain embodiments the polypeptides comprise a truncated HA2 domain. In certain embodiments, the intracellular and transmembrane sequence, e.g. the amino acid sequence from position (or the equivalent of) 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, or 530 of the HA2 domain to the C-terminus of the HA2 domain has been removed, to produce a soluble
20 polypeptide following expression in a cell.

According to the invention, the hemagglutinin stem domain polypeptides are resistant to protease cleavage at the junction between the HA1 and HA2 domain, i.e. do not comprise a protease cleavage site at this junction between HA1 and HA2. It is known to those of skill in the art that the Arg (R) - Gly (G) sequence spanning HA1 and HA2 is a
25 recognition site for trypsin and trypsin-like proteases and is typically cleaved for hemagglutinin activation. Since the HA stem domain polypeptides described herein should not be activated, the influenza hemagglutinin stem domain polypeptides of the invention are resistant to protease cleavage. According to the invention, thus the protease cleavage site is removed or the protease site spanning HA1 and HA2 is mutated to a sequence that is
30 resistant to protease cleavage. According to the invention, removal of the cleavage site between HA1 and HA2 can be achieved by mutation of R (in a small number of cases K) to

Q at the P1 position (see e.g. Sun et al, 2010 for an explanation of the nomenclature of the cleavage site (position 343 in SEQ ID NO: 1). In certain embodiments, the C-terminal amino acid residue of the HA1 C-terminal stem segment is any amino acid other than arginine (R) or lysine (K). In certain embodiments, the HA1 C-terminal amino acid is
5 glutamine (Q), serine (S), threonine (T), asparagine (N), aspartic acid (D) or glutamic acid (E). In certain embodiments, the C-terminal amino acid residue of the HA1 C-terminal stem segment is glutamine (Q).

In certain embodiments, the polypeptides are glycosylated.

In certain embodiments, the influenza hemagglutinin stem domain polypeptides are
10 based on HA of influenza viruses of the H1 subtype. With “based on” it is meant that the N-terminal segments, and/or C-terminal segments of the HA1 domain and/or the HA2 domains have at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% amino acid sequence identity with the corresponding N-terminal and/or C-terminal segments of HA1 and/or the HA2 domains of any naturally occurring influenza hemagglutinin of a H1
15 subtype known to those of skill in the art or later discovered.

In certain embodiments, the polypeptides are based on H1 HA, i.e. HA comprising an amino acid sequence from an influenza virus of the H1 subtype, in particular from the influenza virus A/Brisbane/59/2007 (H1N1) (SEQ ID NO:1), as described below. It will be understood by the skilled person that also other influenza A viruses comprising HA of the
20 H1 subtype may be used according to the invention. In certain embodiments, the polypeptides comprise hemagglutinin stem domains based on HA of an influenza A H1 virus selected from Table 2.

In certain embodiments, the polypeptides comprise a HA1 N-terminal polypeptide segment comprising the amino acids from position 1 to position x of an H1 HA1 domain,
25 wherein x is any amino acid between the amino acid on position 46 and the amino acid on position 60, such as the amino acid on position 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, or 59, preferably wherein x is 52, 53, 55 or 59. Preferably, the polypeptides comprise a HA1 N-terminal segment without the signal sequence, i.e. a HA1 N-terminal segment comprising the amino acids from position 18 (e.g. for H1 HA, such as SEQ ID NO: 1), or
30 an equivalent position in other influenza virus strains (see e.g. Table 2), to position x of the HA1 domain. In certain embodiments, the HA1 N-terminal segment thus comprises the

amino acids from position p (wherein p=18 for H1 HA in SEQ ID NO: 1 or an equivalent position on other H1 HAs), to position x of the HA1 domain.

In certain embodiments, the HA1 C-terminal polypeptide segment comprises the amino acids from position y to and including the C-terminal amino acid of an H1 HA1 domain, wherein y is any amino acid between the amino acid on positions 290 and the
5 amino acid on position 325 of H1 HA1, preferably wherein y is 291, 303, 318, or 321.

In certain embodiments, x= the amino acid on position 52 of SEQ ID NO: 1 (or an equivalent position in another hemagglutinin), p= the amino acid on position 18 of SEQ ID NO: 1 (or an equivalent position in another hemagglutinin) and y= the amino acid on
10 position 321 of SEQ ID NO: 1 (or an equivalent position in another hemagglutinin). In certain embodiments, the HA1 N- terminal stem segment thus comprises the amino acids 1-52 of HA1, and the HA1 C- terminal stem segment comprises the amino acids 321-end (i.e. the C-terminal amino acid of HA1) of HA1. Thus, in certain embodiments, the deletion in the HA1 segment comprises the amino acid sequence from the amino acid at position 53 up
15 to and including the amino acid at position 320. In certain embodiments, the polypeptides do not comprise the signal sequence. In certain embodiments, the HA1 N-terminal segment thus comprises the amino acid 18-52 of HA1, wherein p is the first amino acid of the mature HA molecule (e.g. p=18 in case of SEQ ID NO: 1).

In certain embodiments, the HA1 N- terminal stem segment thus comprises the
20 amino acid residues 1-52 of HA1, preferably the amino acid residues 18-52 of HA1, and the HA1 C- terminal stem segment comprises the amino acid residues 321-343 of HA1. In certain embodiments, the HA1 N- terminal stem segment consists of the amino acid residues 1-52 of HA1, preferably the amino acid residues 18-52 of HA1, and the HA1 C-terminal stem segment consists of the amino acid residues 321-343 of HA1.

25 According to the invention, the stem polypeptides comprise one or more mutations, i.e. amino acid substitutions, in the HA1 domain and/or the HA2 domain of the individual monomers, as compared to the amino acid sequence of corresponding wild-type influenza virus HA1 and/or HA2 domains, i.e. the influenza virus on which the stem polypeptides are based.

30 In certain embodiments, the HA2 domain comprises one or more mutations in the HA2 amino acid sequence connecting the C-terminal residue of helix A to the N-terminal

residue of helix CD (FIG. 1). The H1 HA2 amino acid sequence connecting the C-terminal residue of helix A and the N-terminal residue of helix CD comprises the amino acid sequence comprising residues 402-418 of influenza HA (numbering according to SEQ ID NO: 1), comprising the amino acid sequence

5 MNTQFTAVGKEFN(H/K)LE(K/R) (SEQ ID NO: 7).

In certain embodiments, the amino acid sequence connecting the C-terminal residue of helix A to the N-terminal residue of helix CD, i.e. the region comprising the amino acid residues 402-418 of influenza HA of serotype H1 (numbering according to SEQ ID NO: 1) comprises the amino acid sequence

10 X₁NTQX₂TAX₃GKEX₄N(H/K)X₅E(K/R) (SEQ ID NO: 8).

In certain embodiments, one or more of the amino acids on position 402, 406, 409, 413 and 416 (numbering refers to SEQ ID NO: 1), i.e. one or more of the amino acids X₁, X₂, X₃, X₄ and X₅ have been mutated, i.e. comprise an amino acid that is not occurring at those positions in a wild-type influenza virus on which the stem polypeptide

15 is based.

In certain embodiments, the mutated amino acid on position 402, i.e. X₁, is an amino acid selected from the group consisting of M, E, K, V, R and T.

In certain embodiments, the mutated amino acid on position 406, i.e. X₂, is an amino acid selected from the group consisting of F, I, N, T, H, L and Y, preferably I, L or

20 Y.

In certain embodiments, the mutated amino acid on position 409, i.e. X₃, is an amino acid selected from the group consisting of V, A, G, I, R, F and S, preferably A, I or F.

In certain embodiments, the mutated amino acid on position 413, i.e. X₄, is an amino acid selected from the group consisting of F, I, N, S, T, Y, E, K, M, and V, preferably I, Y, M or V.

25

In certain embodiments, the mutated amino acid on position 416, i.e. X₅, is an amino acid selected from the group consisting of L, H, I, N, R, preferably I.

Combinations of one or more of these mutations are also possible.

30 In certain embodiments, X₁ is M, X₂ is Y, X₃ is I, X₄ is Y and X₅ is S.

According to the invention, the stem polypeptides comprise one or more additional mutations, i.e. amino acid substitutions, in the HA1 domain and/or the HA2 domain, as compared to the amino acid sequence of corresponding wild-type influenza virus HA1 and/or HA2 domains, i.e. the influenza virus on which the stem polypeptides are based.

In certain embodiments, one or more amino acid residues close to the HA0 cleavage site (residue 343 in SEQ ID NO: 1) have been mutated. In certain embodiments, one or more of the amino acid residues on position 337, 340, 352, or 353 of SEQ ID NO: 1, or equivalent positions in other influenza viruses, have been mutated, i.e. are substituted by an amino acid that is not occurring at the corresponding position in the amino acid sequence of the HA of the wild-type influenza virus on which the stem polypeptide is based. Table 6 shows the the naturally occurring amino acid variation.

In certain embodiments, the polypeptides of the invention comprise at least one mutation on position 352 of SEQ ID NO: 1, or on an equivalent position of other influenza viruses.

In certain embodiments, the polypeptides of the invention comprise at least one mutation on position 353 of SEQ ID NO: 1, or on an equivalent position of other influenza viruses.

In certain embodiments, the polypeptides of the invention comprise at least one mutation on position 337 of SEQ ID NO: 1, or on an equivalent position of other influenza viruses.

In certain embodiments, the polypeptides of the invention comprise at least one mutation on position 340 of SEQ ID NO: 1, or on an equivalent position of other influenza viruses.

In certain embodiments, the mutated amino acid residue on position 337 (HA1 domain) is selected from the group consisting of I, E, K, V, A, and T.

In certain embodiments, the mutated amino acid residue on position 340 (HA1 domain) is selected from the group consisting of I, K, R, T, F, N, S and Y.

In certain embodiments, the mutated amino acid residue on position 352 (HA2 domain) is selected from the group consisting of D, V, Y, A, I, N, S, and T.

In certain embodiments, the mutated amino acid residue on position 353 (HA2 domain) is selected from the group consisting of K, R, T, E, G, and V.

In certain embodiments the mutated amino acid introduces a consensus N-glycosylation e.g. N-X-T/S (where X is any naturally occurring amino acid except P) in the sequence as is for example the case for I340N in SEQ ID NO: 6.

In certain embodiments, the mutated amino acid is an amino acid that does not naturally occur in sequences of the same subtype.

In certain embodiments, the mutated amino acid residue on position 337 (HA1 domain) is K.

In certain embodiments, the mutated amino acid residue on position 340 (HA1 domain) is K.

In certain embodiments, the mutated amino acid residue on position 352 (HA2 domain) is F.

In certain embodiments, the mutated amino acid residue on position 353 (HA2 domain) is T.

It is again noted that throughout this application the numbering of the amino acids is based on the numbering of amino acids in H1 HA0, in particular the numbering of the amino acids of the H1N1 influenza strain A/Brisbane/59/2007 (SEQ ID NO: 1). The skilled person will be able to determine the equivalent amino acids in HA of other influenza viruses and thus will be able to determine equivalent mutations, see e.g. Table 2 for the sequence alignment of different H1 influenza viruses.

According to the invention, the polypeptides further may comprise an additional disulfide bridge between the amino acid on position 324 and the amino acid on position 436. Thus, according to the invention at least one additional disulfide bridge has been introduced in the stem domain polypeptides, preferably between amino acids of (or the equivalent of) position 324 and 436 in H1 A/Brisbane/59/2007 (SEQ ID NO: 1). In certain embodiments, the polypeptides thus further comprise the mutation R324C in the HA1 domain and T436C in the HA2 domain. Equivalent positions can be easily determined by those skilled in the art by aligning the sequences using a suitable algorithm such as Clustal, Muscle etc. Engineered disulfide bridges are created by mutating at least one (if the other is already a cysteine), but usually two residues that are

spatially close into cysteine, that will spontaneously or by active oxidation form a covalent bond between the sulfur atoms of these residues.

In certain embodiments, the influenza hemagglutinin stem domain polypeptides are based on HA of influenza viruses of the H3 subtype. With “based on” it is meant that the N-terminal segments, and/or C-terminal segments of the HA1 domain and/or the HA2 domains have at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% amino acid sequence identity with the corresponding N-terminal and/or C-terminal segments of HA1 and/or the HA2 domains of any naturally occurring influenza hemagglutinin of a H3 subtype known to those of skill in the art or later discovered.

In certain embodiments, the polypeptides are based on H3 HA, i.e. HA comprising an amino acid sequence from an influenza virus of the H3N2 virus A/Hong Kong/1/1968 (SEQ ID NO: 237), as described below. It will be understood by the skilled person that also other influenza A viruses comprising HA of the H3 subtype may be used according to the invention.

In certain embodiments, the amino acid sequence CMKQIEDKIEEIESK (SEQ ID NO: 193) has been introduced at positions 419-433 of SEQ ID NO: 1 (or equivalent positions in different HAs) or the amino acid sequence RMCQIEDKIEEIESKQK (SEQ ID NO: 194) has been introduced at position 417-433 of SEQ ID NO: 1 (or equivalent positions in different HAs).

In certain embodiments, the polypeptides further comprise one or more additional mutations in the HA1 and/or HA2 domain, as compared to the amino acid sequence of the HA of which the HA 1 and HA2 domains are derived. Thus, the stability of the stem polypeptides is further increased.

In certain embodiments, the polypeptides selectively bind to the antibodies CR6261 and/or CR9114. In an embodiment, the polypeptide does not bind to the antibodies CR8020 and/or CR8057. In an embodiment, CR6261 comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 9 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 10; CR9114 comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 11 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 12. In an embodiment, CR8057 comprises a heavy chain variable region comprising

the amino acid sequence of SEQ ID NO: 13 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 14. CR8020 comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 17 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 18.

5 In certain embodiments, the polypeptides do selectively bind to the antibody CR8020.

 According to the present invention, in certain embodiments multimeric polypeptides thus are provided that mimic the specific epitopes of CR6261, CR9114, and/or CR8020, and that can be used as immunogenic polypeptides, e.g. to elicit cross-
10 neutralizing antibodies when administered in vivo, either alone, or in combination with other prophylactic and/or therapeutic treatments. With “cross-neutralizing antibodies”, antibodies are meant that are capable of neutralizing at least two, preferably at least three, four, or five different subtypes of influenza A viruses of phylogenetic group 1, and/or at least two, preferably at least three, four, or five different subtypes of influenza A viruses
15 of phylogenetic group 2, and/or at least two, different subtypes of influenza B viruses, in particular at least all virus strains that are neutralized by CR6261 and CR9114.

 As described above, the polypeptides comprise at least two monomers, wherein said monomers each comprise an influenza hemagglutinin HA1 domain that comprises an HA1 N- terminal stem segment that is covalently linked by a linking sequence of 0- 50
20 amino acid residues to the HA1 C- terminal stem segment. The linking sequence does not occur in naturally occurring, or wild-type, HA. In certain embodiments, the linker is a peptide that comprises one amino acid residue, two or less amino acid residues, three or less amino acid residues, four or less amino acid residues, five or less amino acid residues, ten or less amino acid residues, 15 or less amino acid residues, or 20 or less
25 amino acid residues or 30 or less amino acid residues or 40 or less amino acid residues or 50 or less amino acid residues. In a specific embodiment, the linking sequence is a sequence selected from the group consisting of G, GS, GGG, GSG, GSA, GSGS, GSAG, GGGG, GSAGS, GSGSG, GSAGSA, GSAGSAG, and GSGSGSG.

 In certain embodiments, the HA1 N-terminal segment is directly linked to the HA1
30 C-terminal segment, i.e. the polypeptides do not comprise a linking sequence.

Influenza HA in its native form exists as a trimer on the cell or virus membrane. In certain embodiments the intracellular and transmembrane sequence is removed so that a secreted (soluble) polypeptide is produced following expression in cells. Methods to express and purify secreted ectodomains of HA have been described (see e.g. Dopheide et al 2009; Ekiert et al 2009, 2011; Stevens et al 2004, 2006; Wilson et al 1981). A person skilled in the art will understand that these methods can also be applied directly to stem domain polypeptides of the invention in order to achieve expression of secreted (soluble) polypeptide. Therefore these polypeptides are also encompassed in the invention.

In certain embodiments, the polypeptides of the invention contain the intracellular sequences of HA and the transmembrane domain. In other embodiments, the intracellular and transmembrane sequences, e.g. the amino acid sequence from position (or the equivalent of) 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, or 530 of the HA2 domain to the C-terminus of the HA2 domain (numbering according to SEQ ID NO: 1) have been removed to produce a soluble polypeptide following expression in cells.

In certain embodiments, the C-terminal part of the HA2 domain from position 519 to the C-terminal amino acid has been deleted. In further embodiments, the C-terminal part of the HA2 domain from position 530 to the C-terminal amino acid has been deleted.

Optionally, a his-tag sequence (HHHHHH (SEQ ID NO: 15) or HHHHHHH (SEQ ID NO: 16)) may be linked to the (optionally truncated) HA2 domain, for purification purposes, optionally connected through a linker. Optionally the linker may contain a proteolytic cleavage site to enzymatically remove the his-tag after purification.

In certain embodiments, the polypeptides are further stabilized by introducing a sequence known to form trimeric structures, i.e.

GYIPEAPRDGQAYVRKDGEWVLLSTFL (SEQ ID NO: 3) at the C-terminus of HA2, optionally connected through a linker. Thus, in certain embodiments, the C-terminal part of the HA2 domain has been replaced by the amino acid sequence

GYIPEAPRDGQAYVRKDGEWVLLSTFL (SEQ ID NO: 3), optionally connected through a linker. The linker may optionally contain a cleavage site for processing afterwards according to protocols well known to those skilled in the art. To facilitate purification of the soluble form a tag sequence may be added, e.g. a his tag (HHHHHHH

(SEQ ID NO: 15) or HHHHHH (SEQ ID NO: 16)) or FLAG tag (DYKDDDDK) (SEQ ID NO: 22) or a combination of these, optionally connected via short linkers. The linker may optionally contain (part of) a proteolytic cleavage site, e.g. IEGR (SEQ ID NO: 24) (Factor X) or LVPRGS (SEQ ID NO: 23) (thrombin) for processing afterwards
 5 according to protocols well known to those skilled in the art. The processed proteins are also encompassed in the invention.

In certain embodiments, the C-terminal part of the HA2 domain from position 519-565 has been deleted (numbering according to SEQ ID NO: 1) and replaced by SGRDYKDDDDKL VPRGSPGSGYIPEAPRDGQAYVRKDGEWVLLSTFLGHHHHH
 10 H (SEQ ID NO: 4).

In certain embodiments, the C-terminal part of the HA2 domain from position 530-565 has been deleted (numbering according to SEQ ID NO: 1) and replaced by SGRDYKDDDDKL VPRGSPGSGYIPEAPRDGQAYVRKDGEWVLLSTFLGHHHHH
 H (SEQ ID NO: 4).

15 The native HA exists as a trimer on the cell surface. Most of the interactions between the individual monomers that keep the trimer together are located in the head domain while in the stem domain trimerization is mediated by the formation of a trimeric coiled coil motif. After removal of the head the tertiary structure is destabilized and therefore modifications are needed in order to increase protein stability. By strengthening
 20 the helical propensity of the helix CD a more stable protein can be created.

In the polypeptides described in the co-pending application PCT/EP2014/060997, the sequence MKQIEDKIEEIESKQ (SEQ ID NO: 5), derived from yeast transcriptional activator protein GCN4 and known to trimerize was introduced in the CD helix at (the equivalent of) position 419-433. This sequence has a high propensity to form helical
 25 secondary structures and can enhance in this way overall stability of the polypeptides of the invention.

It was further shown that the stability and multimerization state of the polypeptide is dependent on the exact location and sequence of the GCN4 derived sequence in the primary sequence of the polypeptides of the invention.

In preferred embodiments, the amino acid sequence CMKQIEDKIEEIESK (SEQ ID NO: 193) has been introduced at positions 419-433 or wherein sequence RMCQIEDKIEEIESKQK (SEQ ID NO: 194) has been introduced at position 417-433.

In the research that led to the present invention, polypeptides s74H9 (SEQ ID NO: 65), s127H1 (SEQ ID NO: 66), s71H2 (SEQ ID NO: 71), s86B4 (SEQ ID NO: 67), s115A1 (SEQ ID NO: 70), s2201C9 (SEQ ID NO: 77), s55G7 (SEQ ID NO: 68), s113E7 (SEQ ID NO: 78), s6E12 (SEQ ID NO: 69), s181H9 (SEQ ID NO: 76) were modified, using techniques of molecular biology well known to those skilled in the art, to create sequences s74H9-t2 (SEQ ID NO: 93), s127H1-t2 (SEQ ID NO: 91), s71H2-t2 (SEQ ID NO: 97), s86B4-t2 (SEQ ID NO: 92), s115A1-t2 (SEQ ID NO: 96), s220C9-t2 (SEQ ID NO: 99), s55G7-t2 (SEQ ID NO: 95), s113E7-t2 (SEQ ID NO: 100), s6E12-t2 (SEQ ID NO: 94), s181H9-t2 (SEQ ID NO: 98) containing sequence RMKQIEDKIEEIESK (SEQ ID NO: 20) at position 419-433.

In a similar manner, polypeptides s74H9-t3 (SEQ ID NO: 123), s127H1-t3 (SEQ ID NO: 121), s71H2-t3 (SEQ ID NO: 127), s86B4-t3 (SEQ ID NO: 122), s115A1-t3 (SEQ ID NO: 126), s2201C9-t3 (SEQ ID NO: 129), s55G7-t3 (SEQ ID NO: 125), s113E7-t3 (SEQ ID NO: 130), s6E12-t3 (SEQ ID NO: 124), s181H9-t3 (SEQ ID NO: 128) containing sequence RMKQIEDKIEEIESKQK (SEQ ID NO: 21) at position 417-433 were created.

According to the invention, a disulfide bridge between the amino acid on position 411 of a first monomer and the amino acid on position 419 of a second monomer has been introduced by mutating the amino acids on positions 411 and 419 to a cysteine. Thus, in certain embodiments, the amino acid sequence CMKQIEDKIEEIESK (SEQ ID NO: 193) has been introduced at positions 419-433 or the amino acid sequence RMCQIEDKIEEIESKQK (SEQ ID NO: 194) has been introduced at position 417-433.

As described above, applicants have previously identified broadly neutralizing antibodies isolated from primary human B-cells from vaccinated individuals some of which were specific for group 1 (e.g. CR6261, as described in WO 2008/028946) and some of which were specific for group 2 influenza viruses (e.g. CR8020 as described in WO 2010/130636). Detailed analysis of the epitopes of these monoclonal antibodies has revealed the reason for the lack of cross-reactivity of these specific antibodies. In both

cases the presence of glycans in group 1 or group 2 HA molecules on different positions at least partly explained the fact that the antibodies are group-specific. With the identification of CR9114-like antibodies that cross-react with many group 1 and 2 HA molecules, as described below, it has become clear that it is possible for the human
5 immune system to elicit very broad neutralizing antibodies against influenza viruses. However, given the need for a yearly vaccination scheme these antibodies are apparently not, or only to a very low extent elicited following infection or vaccination with (seasonal) influenza viruses of subtypes H1 and/or H3.

According to the present invention multimeric polypeptides are provided that
10 mimic the specific epitopes of CR6261 and/or CR9114, and/or CR8020, and that can be used as immunogenic polypeptides, e.g. to elicit cross-neutralizing antibodies when administered in vivo, either alone, or in combination with other prophylactic and/or therapeutic treatments. With “cross-neutralizing antibodies”, antibodies are meant that are capable of neutralizing at least two, preferably at least three, four, or five different
15 subtypes of influenza A viruses of phylogenetic group 1, and/or at least two, preferably at least three, four, or five different subtypes of influenza A viruses of phylogenetic group 2, and/or at least two, different subtypes of influenza B viruses, in particular at least all virus strains that are neutralized by CR6261 and/or CR9114, and/or CR8020.

In certain embodiments, the polypeptides selectively bind to the antibodies
20 CR6261 and/or CR9114. In certain embodiments, the polypeptide does not bind to the antibody CR8057. CR6261 comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 9 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 10; CR9114 comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 11 and a light chain variable
25 region comprising the amino acid sequence of SEQ ID NO: 12; CR8020 comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 17 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 18. CR8057 comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 13 and a light chain variable region comprising the amino acid sequence of
30 SEQ ID NO: 14.

In certain embodiments, the polypeptides of the present invention are trimeric.

In certain embodiments, the polypeptide monomers comprise the amino acid sequence:

DTICIGYHANNSTDTVDTVLEKNVTVTHTSVNLLENG GGGGKYVCSAKLRMVTGL
 RNX₁PSX₂QSQGLFGAIAGX₃X₄EGGWTGMVDGWYGYHHQNEQSGSYAADQKS
 5 TQNAINGITNKVNSVIEKX₅NTQX₆TAX₇GCEX₈NKX₉ERCMKQIEDKIEEIESKIWC
 YNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFEFYHKCNDE
 CMESVKNGTYDYPKYSEESKLNREKIDGVSGRDYKDDDDKL VPRGSPGSGYIPE
 APRDGQAYVRKDGEWVLLSTFLGHHHHHH (SEQ ID NO: 145),

wherein X₁ is an amino acid selected from the group consisting of E, I, K, V, A, and T;

10 X₂ is an amino acid selected from the group consisting of I, K, R, T, F, N, S and Y;

X₃ is an amino acid selected from the group consisting of D, F, V, Y, A, I, N, S, and T;

X₄ is an amino acid selected from the group consisting of I, K, R, T, E, G and V;

X₅ is an amino acid selected from the group consisting of M, E, K, V, R, T;

X₆ is an amino acid selected from the group consisting of F, I, N, S, T, Y, H, and L;

15 X₇ is an amino acid selected from the group consisting of A, G, I, R, T, V, F, and S;

X₈ is an amino acid selected from the group consisting of F, I, N, S, T, Y, G, E, K, M, and V; and

X₉ is an amino acid selected from the group consisting of H, I, L, N, R, and S.

In certain embodiments, the polypeptide monomers comprise the amino acid sequence:

DTICIGYHANNSTDTVDTVLEKNVTVTHTSVNLLENG GGGGKYVCSAKLRMVTGL
 RNX₁PSX₂QSQGLFGAIAGX₃X₄EGGWTGMVDGWYGYHHQNEQSGSYAADQKS
 TQNAINGITNKVNSVIEKX₅NTQX₆TAX₇GCEX₈NKX₉ERCMKQIEDKIEEIESKIWC
 YNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFEFYHKCNDE
 25 CMESVKNGTYDYPKYSEESKLNREKIDG (SEQ ID NO: 146),

wherein X₁ is an amino acid selected from the group consisting of E, I, K, V, A, and T;

X₂ is an amino acid selected from the group consisting of I, K, R, T, F, N, S and Y;

X₃ is an amino acid selected from the group consisting of D, F, V, Y, A, I, N, S, and T;

X₄ is an amino acid selected from the group consisting of I, K, R, T, E, G and V;

30 X₅ is an amino acid selected from the group consisting of M, E, K, V, R, T;

X₆ is an amino acid selected from the group consisting of F, I, N, S, T, Y, H, and L;

X₇ is an amino acid selected from the group consisting of A, G, I, R, T, V, F, and S;

X₈ is an amino acid selected from the group consisting of F, I, N, S, T, Y, G, E, K, M, and V; and

X₉ is an amino acid selected from the group consisting of H, I, L, N, R, and S.

- 5 In certain embodiments, the polypeptide monomers comprise the amino acid sequence:

DTICIGYHANNSTDTVDTVLEKNVTVTHTSVNLLENG GGGKYVCSAKLRMVTGL
 RNX₁PSX₂QSQGLFGAIAGX₃X₄EGGWTGMVDGWYGYHHQNEQSGYAADQKS
 TQNAINGITNKNVNSVIEKX₅NTQX₆TAX₇GCEX₈NKX₉ERCMKQIEDKIEEIESKIWC
 10 YNAELLVLL ENERTLDFHDSNVKNLYEKVKSQLKNNAKEIGNGC FEFYHKCNDE
 CMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG (SEQ ID NO: 147),

wherein X₁ is an amino acid selected from the group consisting of E, I, K, V, A, and T;

X₂ is an amino acid selected from the group consisting of I, K, R, T, F, N, S and Y;

X₃ is an amino acid selected from the group consisting of D, F, V, Y, A, I, N, S, and T;

- 15 X₄ is an amino acid selected from the group consisting of I, K, R, T, E, G and V;

X₅ is an amino acid selected from the group consisting of M, E, K, V, R, T;

X₆ is an amino acid selected from the group consisting of F, I, N, S, T, Y, H, and L;

X₇ is an amino acid selected from the group consisting of A, G, I, R, T, V, F, and S;

X₈ is an amino acid selected from the group consisting of F, I, N, S, T, Y, G, E, K, M and

- 20 V; and

X₉ is an amino acid selected from the group consisting of H, I, L, N, R, and S.

In certain embodiments, the polypeptide monomers comprise the amino acid sequence:

- DTICIGYHANNSTDTVDTVLEKNVTVTHTSVNLLENG GGGKYVCSAKLRMVTGL
 25 RNX₁PSX₂QSQGLFGAIAGX₃X₄EGGWTGMVDGWYGYHHQNEQSGYAADQKS
 TQNAINGITNKNVNSVIEKX₅NTQX₆TAX₇GCEX₈NKX₉ERCMKQIEDKIEEIESKIWC
 YNAELLVLL ENERTLDFHDSNVKNLYEKVKSQLKNNAKEIGNGC FEFYHKCNDE
 CMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVS
 LGAISFWMC SNGSLQCRICI (SEQ ID NO: 148),

- 30 wherein X₁ is an amino acid selected from the group consisting of E, I, K, V, A, and T;

X₂ is an amino acid selected from the group consisting of I, K, R, T, F, N, S and Y;

X₃ is an amino acid selected from the group consisting of D, F, V, Y, A, I, N, S, and T;

X₄ is an amino acid selected from the group consisting of I, K, R, T, E, G and V;

X₅ is an amino acid selected from the group consisting of M, E, K, V, R, T;

X₆ is an amino acid selected from the group consisting of F, I, N, S, T, Y, H, and L;

5 X₇ is an amino acid selected from the group consisting of A, G, I, R, T, V, F, and S;

X₈ is an amino acid selected from the group consisting of F, I, N, S, T, Y, G, E, K, M and V; and

X₉ is an amino acid selected from the group consisting of H, I, L, N, R, and S.

In certain embodiments, X₁ is K, X₂ is K, X₃ is F, X₄ is T, X₅ is M., X₆ is Y, X₇ is I;

10 X₈ is Y and X₉ is S.

The influenza hemagglutinin stem domain polypeptides can be prepared according to any technique deemed suitable to one of skill, including techniques described below.

Thus, the immunogenic polypeptides of the invention may be synthesized as DNA
15 sequences by standard methods known in the art and cloned and subsequently expressed, *in vitro or in vivo*, using suitable restriction enzymes and methods known in the art. The present invention thus also relates to nucleic acid molecules encoding the above described polypeptides. The invention further relates to vectors comprising the nucleic acids encoding the polypeptides of the invention. In certain embodiments, a nucleic acid molecule
20 according to the invention is part of a vector, e.g. a plasmid. Such vectors can easily be manipulated by methods well known to the person skilled in the art, and can for instance be designed for being capable of replication in prokaryotic and/or eukaryotic cells. In addition, many vectors can directly or in the form of an isolated desired fragment there from be used for transformation of eukaryotic cells and will integrate in whole or in part into the genome
25 of such cells, resulting in stable host cells comprising the desired nucleic acid in their genome. The vector used can be any vector that is suitable for cloning DNA and that can be used for transcription of a nucleic acid of interest. When host cells are used it is preferred that the vector is an integrating vector. Alternatively, the vector may be an episomally replicating vector.

30 The person skilled in the art is capable of choosing suitable expression vectors, and inserting the nucleic acid sequences of the invention in a functional manner. To obtain

expression of nucleic acid sequences encoding polypeptides, it is well known to those skilled in the art that sequences capable of driving expression can be functionally linked to the nucleic acid sequences encoding the polypeptide, resulting in recombinant nucleic acid molecules encoding a protein or polypeptide in expressible format. In general, the promoter sequence is placed upstream of the sequences that should be expressed. Many expression vectors are available in the art, e.g. the pcDNA and pEF vector series of Invitrogen, pMSCV and pTK-Hyg from BD Sciences, pCMV-Script from Stratagene, etc, which can be used to obtain suitable promoters and/or transcription terminator sequences, polyA sequences, and the like. Where the sequence encoding the polypeptide of interest is properly inserted with reference to sequences governing the transcription and translation of the encoded polypeptide, the resulting expression cassette is useful to produce the polypeptide of interest, referred to as expression. Sequences driving expression may include promoters, enhancers and the like, and combinations thereof. These should be capable of functioning in the host cell, thereby driving expression of the nucleic acid sequences that are functionally linked to them. The person skilled in the art is aware that various promoters can be used to obtain expression of a gene in host cells. Promoters can be constitutive or regulated, and can be obtained from various sources, including viruses, prokaryotic, or eukaryotic sources, or artificially designed. Expression of nucleic acids of interest may be from the natural promoter or derivative thereof or from an entirely heterologous promoter (Kaufman, 2000). Some well-known and much used promoters for expression in eukaryotic cells comprise promoters derived from viruses, such as adenovirus, e.g. the E1A promoter, promoters derived from cytomegalovirus (CMV), such as the CMV immediate early (IE) promoter (referred to herein as the CMV promoter) (obtainable for instance from pcDNA, Invitrogen), promoters derived from Simian Virus 40 (SV40) (Das et al, 1985), and the like. Suitable promoters can also be derived from eukaryotic cells, such as methallothionein (MT) promoters, elongation factor 1 α (EF-1 α) promoter (Gill et al., 2001), ubiquitin C or UB6 promoter (Gill et al., 2001), actin promoter, an immunoglobulin promoter, heat shock promoters, and the like. Testing for promoter function and strength of a promoter is a matter of routine for a person skilled in the art, and in general may for instance encompass cloning a test gene such as lacZ, luciferase, GFP, etc. behind the promoter sequence, and test for expression of the test gene. Of course, promoters may be altered by deletion,

addition, mutation of sequences therein, and tested for functionality, to find new, attenuated, or improved promoter sequences. According to the present invention, strong promoters that give high transcription levels in the eukaryotic cells of choice are preferred.

The constructs may be transfected into eukaryotic cells (e.g. plant, fungal, yeast or
5 animal cells) or suitable prokaryotic expression systems like *E. coli* using methods that are well known to persons skilled in the art. In some cases a suitable 'tag' sequence (such as for example, but not limited to, a his-, myc-, strep-, or flag-tag) or complete protein (such as for example, but not limited to, maltose binding protein or glutathione S transferase) may be added to the sequences of the invention to allow for purification and/or identification of the
10 polypeptides from the cells or supernatant. Optionally a sequence containing a specific proteolytic site can be included to afterwards remove the tag by proteolytic digestion.

Purified polypeptides can be analyzed by spectroscopic methods known in the art (e.g. circular dichroism spectroscopy, Fourier Transform Infrared spectroscopy and NMR spectroscopy or X-ray crystallography) to investigate the presence of desired structures like
15 helices and beta sheets. ELISA, Octet and FACS and the like can be used to investigate binding of the polypeptides of the invention to the broadly neutralizing antibodies described before (CR6261, CR9114, CR8057). Thus, polypeptides according to the invention having the correct conformation can be selected.

The invention further relates to compositions comprising a therapeutically effective
20 amount of at least one of the polypeptides and/or nucleic acids of the invention. The compositions preferably are immunogenic compositions. The compositions preferably further comprise a pharmaceutically acceptable carrier. In the present context, the term "pharmaceutically acceptable" means that the carrier, at the dosages and concentrations employed, will not cause unwanted or harmful effects in the subjects to which they are
25 administered. Such pharmaceutically acceptable carriers and excipients are well known in the art (see Remington's Pharmaceutical Sciences, 18th edition, A. R. Gennaro, Ed., Mack Publishing Company [1990]; Pharmaceutical Formulation Development of Peptides and Proteins, S. Frokjaer and L. Hovgaard, Eds., Taylor & Francis [2000]; and Handbook of Pharmaceutical Excipients, 3rd edition, A. Kibbe, Ed., Pharmaceutical Press [2000]). The
30 term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the composition is administered. Saline solutions and aqueous dextrose and glycerol solutions can e.g. be

employed as liquid carriers, particularly for injectable solutions. The exact formulation should suit the mode of administration. The polypeptides and/or nucleic acid molecules preferably are formulated and administered as a sterile solution. Sterile solutions are prepared by sterile filtration or by other methods known per se in the art. The solutions can
5 then be lyophilized or filled into pharmaceutical dosage containers. The pH of the solution generally is in the range of pH 3.0 to 9.5, e.g. pH 5.0 to 7.5.

The invention also relates to influenza HA stem domain polypeptides, nucleic acid molecules and/or vectors as described above for use in inducing an immune response against influenza HA protein. The invention also relates to methods for inducing an
10 immune response in a subject, the method comprising administering to a subject, a polypeptide, nucleic acid molecule and/or immunogenic composition as described above. A subject according to the invention preferably is a mammal that is capable of being infected with an infectious disease-causing agent, in particular an influenza virus, or otherwise can benefit from the induction of an immune response, such subject for
15 instance being a rodent, e.g. a mouse, a ferret, or a domestic or farm animal, or a non-human-primate, or a human. Preferably, the subject is a human subject. The invention thus provides methods for inducing an immune response to an influenza virus hemagglutinin (HA), in particular of a group 1 and/or group 2 influenza A virus, such as an influenza virus comprising HA of the H1, H2, H3, H4, H5, H7 and/or H10 subtype,
20 and/or of an influenza B virus, in a subject utilizing the polypeptides, nucleic acids and/or immunogenic compositions described herein. In some embodiments, the invention provides methods for inducing an immune response to an influenza virus comprising HA of the H1 subtype, in a subject utilizing the polypeptides, nucleic acids and/or immunogenic compositions described herein.

25 In some embodiments, the immune response induced is effective to prevent and/or treat an influenza virus infection caused by a group 1 and/or group 2 influenza A virus subtypes and/or influenza B viruses. In some embodiments, the immune response induced by the polypeptides, nucleic acids and/or immunogenic compositions described herein is effective to prevent and/or treat an influenza A and/or B virus infection caused by two,
30 three, four, five or six subtypes of influenza A and/or B viruses. In some embodiments,

the immune response induced is effective to prevent and/or treat an influenza virus infection caused by an influenza virus comprising HA of the H1 subtype.

Since it is well known that small proteins and/or nucleic acid molecules do not always efficiently induce a potent immune response it may be necessary to increase the immunogenicity of the polypeptides and/or nucleic acid molecules by adding an adjuvant. In certain embodiments, the immunogenic compositions described herein comprise, or are administered in combination with, an adjuvant. The adjuvant for administration in combination with a composition described herein may be administered before, concomitantly with, or after administration of said composition. Examples of suitable adjuvants include aluminium salts such as aluminium hydroxide and/or aluminium phosphate; oil-emulsion compositions (or oil-in-water compositions), including squalene-water emulsions, such as MF59 (see e.g. WO 90/14837); saponin formulations, such as for example QS21 and Immunostimulating Complexes (ISCOMS) (see e.g. US 5,057,540; WO 90/03184, WO 96/11711, WO 2004/004762, WO 2005/002620); bacterial or microbial derivatives, examples of which are monophosphoryl lipid A (MPL), 3-O-deacylated MPL (3dMPL), CpG-motif containing oligonucleotides, ADP-ribosylating bacterial toxins or mutants thereof, such as E. coli heat labile enterotoxin LT, cholera toxin CT, pertussis toxin PT, or tetanus toxoid TT, Matrix M (Isconova). In addition, known immunopotentiating technologies may be used, such as fusing the polypeptides of the invention to proteins known in the art to enhance immune response (e.g. tetanus toxoid, CRM197, rCTB, bacterial flagellins or others) or including the polypeptides in virosomes, or combinations thereof. Other non-limiting examples that can be used are e.g. disclosed by Coffman et al. (2010).

In an embodiment, the influenza hemagglutinin stem domain polypeptides of the invention are incorporated into viral-like particle (VLP) vectors. VLPs generally comprise a viral polypeptide(s) typically derived from a structural protein(s) of a virus. Preferably, the VLPs are not capable of replicating. In certain embodiments, the VLPs may lack the complete genome of a virus or comprise a portion of the genome of a virus. In some embodiments, the VLPs are not capable of infecting a cell. In some embodiments, the VLPs express on their surface one or more of viral (e.g., virus surface glycoprotein) or non-viral (e.g., antibody or protein) targeting moieties known to one skilled in the art.

In a specific embodiment, the polypeptide of the invention is incorporated into a virosome. A virosome containing a polypeptide according to the invention may be produced using techniques known to those skilled in the art. For example, a virosome may be produced by disrupting a purified virus, extracting the genome, and reassembling
5 particles with the viral proteins (e.g., an influenza hemagglutinin stem domain polypeptide) and lipids to form lipid particles containing viral proteins.

The invention also relates to the above-described polypeptides, nucleic acids, vectors and/or immunogenic compositions for inducing an immune response in a subject against influenza HA, in particular for use as a vaccine. The influenza hemagglutinin
10 stem domain polypeptides, nucleic acids encoding such polypeptides, or vectors comprising such nucleic acids or polypeptides described herein thus may be used to elicit neutralizing antibodies against influenza viruses, for example, against the stem region of influenza virus hemagglutinin. The invention in particular relates to polypeptides, nucleic acids, and/or immunogenic compositions as described above for use as a vaccine in the
15 prevention and/or treatment of a disease or condition caused by an influenza A virus of phylogenetic group 1 and/or phylogenetic group 2 and/or an influenza B virus. In an embodiment, the vaccine may be used in the prevention and/or treatment of diseases caused by two, three, four, five, six or more different subtypes of phylogenetic group 1 and/or 2 and/or influenza B viruses. In an embodiment, the vaccine may be used in the
20 prevention and/or treatment of influenza infection caused by an influenza virus comprising HA of the H1 subtype.

The polypeptides of the invention may be used after synthesis *in vitro* or in a suitable cellular expression system, including bacterial and eukaryotic cells, or alternatively, may be expressed *in vivo* in a subject in need thereof, by expressing a
25 nucleic acid coding for the immunogenic polypeptide. Such nucleic acid vaccines may take any form, including naked DNA, plasmids, or viral vectors including adenoviral vectors.

Administration of the polypeptides, nucleic acid molecules, vectors and/or immunogenic compositions according to the invention can be performed using standard
30 routes of administration. Non-limiting examples include parenteral administration, such as intravenous, intradermal, transdermal, intramuscular, subcutaneous, etc, or mucosal

administration, e.g. intranasal, oral, and the like. The skilled person will be capable to determine the various possibilities to administer the polypeptides, nucleic acid molecules, and/or immunogenic compositions according to the invention, in order to induce an immune response. In certain embodiments, the polypeptide, nucleic acid molecule, and/or immunogenic composition (or vaccine) is administered more than one time, i.e. in a so-called homologous prime-boost regimen. In certain embodiments where the polypeptide, nucleic acid molecule, and/or immunogenic composition is administered more than once, the administration of the second dose can be performed after a time interval of, for example, one week or more after the administration of the first dose, two weeks or more after the administration of the first dose, three weeks or more after the administration of the first dose, one month or more after the administration of the first dose, six weeks or more after the administration of the first dose, two months or more after the administration of the first dose, 3 months or more after the administration of the first dose, 4 months or more after the administration of the first dose, etc, up to several years after the administration of the first dose of the polypeptide, nucleic acid molecule, and/or immunogenic composition. It is also possible to administer the vaccine more than twice, e.g. three times, four times, etc, so that the first priming administration is followed by more than one boosting administration. In other embodiments, the polypeptide, nucleic acid molecule, vectors and/or composition according to the invention is administered only once.

The polypeptides, nucleic acid molecules, vectors and/or compositions may also be administered, either as prime, or as boost, in a heterologous prime-boost regimen.

The invention further provides methods for preventing and/or treating an influenza virus disease in a subject utilizing the polypeptides, nucleic acids and/or compositions described herein. In a specific embodiment, a method for preventing and/or treating an influenza virus disease in a subject comprises administering to a subject in need thereof an effective amount of a polypeptide, nucleic acid molecule, vector and/or composition, as described above. A therapeutically effective amount refers to an amount of the polypeptide, nucleic acid, and/or composition as defined herein, that is effective for preventing, ameliorating and/or treating a disease or condition resulting from infection by a group 1 or 2 influenza A virus, and/or an influenza B virus, preferably a disease resulting from infection by an influenza A virus comprising HA of the H1 subtype and/or an influenza A

virus comprising HA of the H3 subtype. Prevention encompasses inhibiting or reducing the spread of influenza virus or inhibiting or reducing the onset, development or progression of one or more of the symptoms associated with infection by an influenza virus. Amelioration as used in herein may refer to the reduction of visible or perceptible disease symptoms, viremia, or any other measurable manifestation of influenza infection.

Those in need of treatment include those already inflicted with a condition resulting from infection with a group 1 or a group 2 influenza A virus, or an influenza B virus, as well as those in which infection with influenza virus is to be prevented. The polypeptides, nucleic acid molecules, vectors and/or compositions of the invention thus may be administered to a naive subject, i.e., a subject that does not have a disease caused by influenza virus infection or has not been and is not currently infected with an influenza virus infection, or to subjects that already are and/or have been infected with an influenza virus.

In an embodiment, prevention and/or treatment may be targeted at patient groups that are susceptible to influenza virus infection. Such patient groups include, but are not limited to e.g., the elderly (e.g. ≥ 50 years old, ≥ 60 years old, and preferably ≥ 65 years old), the young (e.g. ≤ 5 years old, ≤ 1 year old), hospitalized patients and patients who have been treated with an antiviral compound but have shown an inadequate antiviral response.

In another embodiment, the polypeptides, nucleic acids molecules and/or vectors may be administered to a subject in combination with one or more other active agents, such as existing, or future influenza vaccines, monoclonal antibodies and/or antiviral agents, and/or antibacterial, and/or immunomodulatory agents. The one or more other active agents may be beneficial in the treatment and/or prevention of an influenza virus disease or may ameliorate a symptom or condition associated with an influenza virus disease. In some embodiments, the one or more other active agents are pain relievers, anti-fever medications, or therapies that alleviate or assist with breathing.

Dosage regimens of the polypeptides and/or nucleic acid molecules of the invention can be adjusted to provide the optimum desired response (e.g., a therapeutic response). A suitable dosage range may for instance be 0.1-100 mg/kg body weight, preferably 1-50 mg/kg body weight, preferably 0.5-15 mg/kg body weight. The precise dosage of the

polypeptides and/or nucleic acid molecules to be employed will e.g. depend on the route of administration, and the seriousness of the infection or disease caused by it, and should be decided according to the judgment of the practitioner and each subject's circumstances. For example, effective doses vary depending target site, physiological state of the patient
5 (including age, body weight, health), and whether treatment is prophylactic or therapeutic. Usually, the patient is a human but non-human mammals including transgenic mammals can also be treated. Treatment dosages are optimally titrated to optimize safety and efficacy.

The polypeptides of the invention may also be used to verify binding of monoclonal
10 antibodies identified as potential therapeutic candidates. In addition, the polypeptides of the invention may be used as diagnostic tool, for example to test the immune status of an individual by establishing whether there are antibodies in the serum of such individual capable of binding to the polypeptide of the invention. The invention thus also relates to an in vitro diagnostic method for detecting the presence of an influenza infection in a patient
15 said method comprising the steps of a) contacting a biological sample obtained from said patient with a polypeptide according to the invention; and b) detecting the presence of antibody-antigen complexes.

The polypeptides of the invention may also be used to identify new binding molecules or improve existing binding molecules, such as monoclonal antibodies and
20 antiviral agents.

The invention is further illustrated in the following examples and figures. The examples are not intended to limit the scope of the invention in any way.

EXAMPLES

25

Example 1: Stem based polypeptides disclosed in PCT/EP2014/060997

PCT/EP2012/073706 discloses influenza hemagglutinin stem domain polypeptides, compositions and vaccines and methods of their use in the field of prevention and/or treatment of influenza. PCT/EP2014/060997 discloses further
30 sequences of stem domain polypeptides derived from the full length HA of H1N1 A/Brisbane/59/2007 (SEQ ID NO: 1), which were obtained by site-directed mutation of

H1-mini2-cluster1+5+6-GCN4 (SEQ ID NO: 2) and which stably presented the broadly neutralizing epitope of CR6261 (Throsby et al, 2009; Ekiert et al 2010) and/or CR9114.

H1-mini2-cluster1+5+6-GCN4 (SEQ ID NO: 2) was derived from the full length HA of H1N1 A/Brisbane/59/2007 (SEQ ID NO: 1) by taking the following steps:

- 5 1. Removal of the cleavage site in HA0. Cleavage of wild type HA at this site results in HA1 and HA2. The removal can be achieved by mutation of R to Q at the P1 position (see e.g. Sun et al, 2010 for an explanation of the nomenclature of the cleavage site (position 343 in SEQ ID NO: 1).
2. Removal of the head domain by deleting amino acids 53 to 320 from SEQ ID NO: 1.
- 10 The remaining N- and C-terminal parts of the sequence were joined by a four residue flexible linker, GGGG.
3. Increasing the solubility of the loop (between the A-helix and the CD helix) formed by (the equivalent of) residues 402 to 418 in H1 A/Brisbane/59/2007 (SEQ ID NO: 1) in order to both increase the stability of the pre-fusion conformation and to
- 15 destabilize the post-fusion conformation of the modified HA. In H1-mini2-cluster1+5+6-GCN4 (SEQ ID NO: 2) mutations F406S, V409T, F413G and L416S (numbering refers to SEQ ID NO: 1) were introduced
4. Introducing a disulfide bridge between amino acids at position 324 and 436 in H1 A/Brisbane/59/2007; this is achieved by introducing mutations R324C and Y436C.
- 20 (numbering refers to SEQ ID NO: 1)
5. Introducing the GCN4 derived sequence MKQIEDKIEEIESKQ (SEQ ID NO: 5), that is known to trimerize, at position 419-433 (numbering refers to SEQ ID NO: 1).

In certain embodiments, the sequence of the transmembrane and intracellular domain was deleted from position (or the equivalent thereof, as determined from sequence

25 alignment) 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 526, 527, 528, 529, or 530 of HA2 to the C-terminus of HA2 (numbering according to SEQ ID NO: 1) so that a secreted (soluble) polypeptide was produced following expression in cells.

The soluble polypeptide was further stabilized by introducing a sequence known to form trimeric structures, i.e. the foldon sequence GYIPEAPRDGQAYVRKDGEWVLLSTFL

30 (SEQ ID NO: 3), optionally connected through a short linker, as described above. The linker may optionally contain a cleavage site for processing afterwards according to

protocols well known to those skilled in the art. To facilitate purification and detection of the soluble form a tag sequence may be optionally added, e.g. a histidine tag (HHHHHHH (SEQ ID NO: 20) or HHHHHH (SEQ ID NO: 21) or a FLAG tag (DYKDDDDK; SEQ ID NO: 22) or combination of these, optionally connected via short
5 linkers. The linker may optionally contain (part of) a proteolytic cleavage site, e.g. LVPRGS (SEQ ID NO: 23) (thrombin) or IEGR (SEQ ID NO: 24) (Factor X) for processing afterwards according to protocols well known to those skilled in the art. The processed proteins are also encompassed in the invention.

An example of such a C-terminal sequence combining FLAG-tag, thrombin
10 cleavage site, foldon, and His sequences is SEQ ID NO: 4 FLAG- thrombin-foldon-His. This sequence was combined with a soluble form of H1-mini2-cluster1+5+6-GCN4 (SEQ ID NO: 2) sequence to create the parental sequence (SEQ ID NO: 6) that was used to create novel polypeptides of the invention by mutagenesis. This sequence does not contain the leader sequence corresponding to amino acids 1-17 of SEQ ID NO: 1 and 2.

15 Stem domain polypeptides thus were created by deleting the part of the hemagglutinin sequence that encodes the head domain of the molecule and reconnecting the N- and C-terminal parts of the sequence on either side of the deletion through a linker as described in PCT/2012/073706 and above. The removal of the head domain leaves part of the molecule that was previously shielded from the aqueous solvent exposed,
20 potentially destabilizing the structure of the polypeptides of the invention. For this reason residues in the B-loop (in particular amino acid residue 406 (F and S in SEQ ID NO: 1 and 2, respectively), 409 (V and T) 413 (F and G) and 416 (L and S) were mutated in various combinations using parental sequence SEQ ID NO: 6 as the starting point. SEQ ID NO: 6 was created from H1-mini2-cluster1+5+6-GCN4 (SEQ ID NO: 2) by removing
25 the leader sequence, and replacing residues 520-565 with a Flag-thrombin-foldon--his sequence (SEQ ID NO: 4).

Similarly, in the area around the fusion peptide a number of hydrophobic residues are exposed to the solvent, caused by the fact that, unlike the native full length HA, the polypeptides cannot be cleaved and undergo the associated conformational change that
30 buries the hydrophobic fusion peptide in the interior of the protein. To address this issue

some or all of the residues I337, I340, F352 and I353 in SEQ ID NO: 2 were also mutated.

This way, the soluble forms of HA stem polypeptides 74H9 (SEQ ID NO: 57), 127H1 (SEQ ID NO: 55), 71H2 (SEQ ID NO: 61), 86B4 (SEQ ID NO: 56), 115A1 (SEQ ID NO: 60), 2201C9 (SEQ ID NO: 63), 55G7 (SEQ ID NO: 59), 113E7 (SEQ ID NO: 64), 6E12 (SEQ ID NO: 58), 181H9 (SEQ ID NO: 62) were created as part of a library.

DNA sequences encoding the polypeptides described above were transformed into *Pichia pastoris* or transfected into HEK293F cells using protocols well known to persons skilled in the art. Constructs used for expression in mammalian cells contained the HA leader sequence (residue 1-17 in SEQ ID NO: 1 and 2), whereas in constructs used for expression in *P. pastoris* the HA leader sequence was replaced with the yeast alpha factor leader sequence (SEQ ID NO: 7). In this way expressed protein are directed towards the cell culture medium thus allowing binding and expression to be determined without further purification of the polypeptides of the invention. All sequences contained the FLAG-foldon-HIS C-terminal sequence (SEQ ID NO: 4).

Monoclonal antibody binding (CR6261, CR9114, CR8020) to polypeptides of the invention was determined by ELISA. To this end ELISA plates were treated overnight with a 2 µg/ml monoclonal antibody solution (20 µl/well) at 4 °C. After removal of the antibody solution the remaining surface was blocked with 4% solution of non-fat dry milk powder in PBS for a minimum of 1 h at room temperature. After washing of the plates, 20 µl of cell culture medium (neat or diluted) was added to each well and incubated for at least 1 h at room temperature. ELISA plates were then washed and 20 µl of anti-FLAG-HRP antibody solution (Sigma A8952, 2000 times diluted in 4% non-fat dry milk in PBS-Tween) was added. After incubation (1h at room temperature) plates were washed once more, and 20 µl luminescent substrate (Thermoscientific C#34078) was added to develop the signal. Alternatively, a colorimetric detection method can be used to develop the signal.

Expression of polypeptides of the invention was determined from a homogeneous time-resolved fluorescence assay (for a general description see e.g. Degorce et al., Curr. Chem. Genomics 2009 3: 22-32). To this end a mixture of Terbium (Tb) labeled anti-FLAG monoclonal antibody (donor) and Alexa488 labeled anti-His monoclonal antibody

(acceptor) (HTRF solution) was prepared by adding 210.5 µl Anti-FLAG-TB (stock solution 26 µg/ml) and 1.68 ml of anti-HIS-488 (stock solution 50 µg/ml) to 80 ml of a 1 to 1 mixture of culture medium and 50 mM HEPES + 0.1% BSA. 19 µl of HTRF solution was added to each well of a ELISA plate and 1 µl of culture medium was added. Upon
5 excitation and after a delay to allow interfering short-lived background signals arising from other compounds (proteins, media components etc) to decay, the ratio of fluorescence emission at 520 and 665 nm was determined. This is a measure of total protein content in the sample and is used to normalize the mAb binding signals between different experiments.

10 The polypeptides listed in Table 3 and 4 were expressed in P.Pastoris following protocols well known to those skilled in the art. Culture medium was collected and binding to CR6261 binding of and expression of the stem domain polypeptides was determined as described above. Since the response in the binding assay scales with the concentration of expresses protein, ELISA binding signal was normalized for protein
15 expression by comparing the ratio of binding signal over the signal in the HTRF assay for each expressed sequence. All expressed protein exhibit higher ratio's of CR626 binding to HTRF signal compared to the parental sequence of SEQ ID NO: 6.

In addition, the ratio of CR6261 binding to HTRF signals was calculated and compared to the ratio calculated for the parental sequence SEQ ID NO: 6. The results are
20 listed in column 5 of Tables 3 and 4; all expressed proteins exhibit higher ratios, indicating that the stem polypeptides described above show increased binding of CR6261.

Example 2: Design and characterization of further polypeptides

25 The polypeptides described above contain sequence RMKQIEDKIEEIESKQ, derived from yeast transcriptional activator protein GCN4, in the CD helix. This sequence has a high propensity to form helical secondary structures and can enhance in this way overall stability of the polypeptide of the invention. It has surprisingly been found that stability and aggregation state of the hemagglutinin stem polypeptides is dependent on the exact location
30 and sequence of the GCN4 derived sequence in the primary sequence of the polypeptides.

In this example, we describe a novel set of polypeptides wherein sequence RMKQIEDKIEEIESK (SEQ ID NO: 20) has been introduced at position 419-433

(numbering according to SEQ ID NO: 1; for example SEQ ID NO. 81 to 110) or sequence RMKQIEDKIEEIESKQK (SEQ ID NO: 21) has been introduced at position 417-433 (for example SEQ ID NO 111 to 140).

To this end, the polypeptides described in Example 1, i.e 74H9 (SEQ ID NO: 57),
 5 127H1 (SEQ ID NO: 55), 71H2 (SEQ ID NO: 61), 86B4 (SEQ ID NO: 56), 115A1 (SEQ ID NO: 60), 2201C9 (SEQ ID NO: 63), 55G7 (SEQ ID NO: 59), 113E7 (SEQ ID NO: 64), 6E12 (SEQ ID NO: 58), 181H9 (SEQ ID NO: 62) were modified, using techniques of molecular biology well known to those skilled in the art, to create sequences 74H9-t2 (SEQ ID NO: 83), 127H1-t2 (SEQ ID NO: 81), 71H2-t2 (SEQ ID NO: 87), 86B4-t2 (SEQ ID NO: 82), 115A1-t2 (SEQ ID NO: 86), 220C9-t2 (SEQ ID NO: 89), 55G7-t2 (SEQ ID NO: 85),
 10 113E7-t2 (SEQ ID NO: 90), 6E12-t2 (SEQ ID NO: 84), 181H9-t2 (SEQ ID NO: 88) containing sequence RMKQIEDKIEEIESK (SEQ ID NO: 20) at position 419-433.

In a similar manner sequences 74H9-t3 (SEQ ID NO: 113), 127H1-t3 (SEQ ID NO: 111), 71H2-t3 (SEQ ID NO: 117), 86B4-t3 (SEQ ID NO: 112), 115A1-t3 (SEQ ID NO: 116), 2201C9-t3 (SEQ ID NO: 119), 55G7-t3 (SEQ ID NO: 115), 113E7-t3 (SEQ ID NO: 120), 6E12-t3 (SEQ ID NO: 114), 181H9-t3 (SEQ ID NO: 118) containing sequence RMKQIEDKIEEIESKQK (SEQ ID NO: 21) at position 417-433 were created.

Polypeptides can also be created on the basis of the sequence of HA molecules from different viral strains. SEQ ID NO: 195 - 201 for example describe polypeptides based on the
 20 HA sequence of the H1N1 A/California/07/09 strain.

As described before, soluble polypeptides can be created by removing the C-terminal part of the HA based sequences for example from residue 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, or 530 of the HA2 domain to the C-terminus of the HA2 domain (numbering according to SEQ ID NO: 1).

25 The polypeptides can further be stabilized by introducing a sequence known to form trimeric structures, i.e GYIPEAPRDGQAYVRKDGWVLLSTFL (SEQ ID NO: 3), optionally connected through a linker. The linker may optionally contain a cleavage site for processing afterwards according to protocols well known to those skilled in the art. To facilitate purification of the soluble form a tag sequence may be added, e.g. a his tag
 30 (HHHHHHH (SEQ ID NO: 15) or HHHHHH (SEQ ID NO: 16)) or FLAG tag (DYKDDDDK) (SEQ ID NO: 22) or a combination of these, optionally connected via short linkers. The linker may optionally contain (part of) a proteolytic cleavage site, e.g. IEGR

(SEQ ID NO: 24) (Factor X) or LVPRGS (SEQ ID NO: 23) (thrombin) for processing afterwards according to protocols well known to those skilled in the art.

Soluble forms of the polypeptides of SEQ ID NO 55-64 and 81-90 were created by replacement of the equivalent of residue 519-565 (numbering refers to SEQ ID NO: 1) with
5 sequence RSLVPRGSPGHHHHHH, containing both a modified thrombin cleavage site and a 6 histidine tag (SEQ ID NO: 21) and were expressed in HEK293F cells following protocols well known to those skilled in the art.

For reasons of comparison, soluble forms of H1-mini2-cluster1+5+6-GCN4t2 (SEQ ID NO:52) and H1-mini2-cluster1+5+6-GCN4t3 (SEQ ID NO: 53)), described in
10 PCT/EP2012/073706 were also included in the experiments. Culture medium was collected and binding to CR6261, CR9114 was detected by a sandwich ELISA, using coated mAb CR6261 or CR9114 to capture the polypeptide directly from the culture medium and a Horse Radish Peroxidase (HRP) conjugated antibody directed against the C-terminal his-tag for detection purposes. Alternatively, biotinylated CR9114 in combination with HRP-conjugated
15 streptavidin was used for detection of CR9114 captured polypeptides in a sandwich ELISA. This format allows the detection of the presence of multimeric forms of polypeptides. All polypeptides tested were capable of binding to CR9114 and CR6261 as determined by ELISA. Increased levels of multimerization as detected by the CR9114 capture – biotinylated CR9114 detection sandwich ELISA were observed for s55G7-t2 (SEQ ID NO: 95), s86B4-t2
20 (SEQ ID NO: 92), s115A1-t2 (SEQ ID NO: 96), s127H1-t2 (SEQ ID NO: 91), s113E7-t2 (SEQ ID NO: 100), s220C9-t2 (SEQ ID NO: 99), s71H2-t3 (SEQ ID NO: 127), s127H1-t3 (SEQ ID NO: 121), s74H9-t3 (SEQ ID NO: 123).

In order to obtain a highly pure preparations of the polypeptides for further characterization, HEK293F cells were transfected with expression vector pcDNA2004
25 containing the genes encoding soluble forms of 127H1-t2 (SEQ ID NO: 81), 86B4-t2 (SEQ ID NO: 82) and 55G7-t2 (SEQ ID NO: 85). It will be understood by the skilled person that the leader sequence (or signal sequence) that directs transport of a protein during production (corresponding to amino acids 1-17 of SEQ ID NO: 1) will not be present in the secreted final polypeptide.

30 To produce the polypeptides 1.0×10^6 vc/mL were seeded by spinning down HEK293F cells (Invitrogen) at 300 g for 5 min and resuspending in 300 mL pre-warmed Freestyle™ medium per SF1000 flask. This culture was incubated for 1 hour at 37 °C, 10% CO2 at 110 rpm in a multitron incubator. After 1 hour the plasmid DNA was pipetted in 9.9

mL Optimem medium to a concentration of 1.0 µg/mL in the 300 mL culture volume. In parallel 440 µL 293fectin® was pipetted in 9.9 mL Optimem medium and incubated for 5 minutes at room temperature. After 5 minutes the plasmid DNA/Optimem mix was added to the 293fectin®/Optimem mix and incubated at room temperature for 20 minutes. After the incubation the plasmid DNA/293fectin® mix was added drop wise to the cell suspension. The transfected cultured was incubated at 37 °C, 10% CO₂ and 110 rpm in a multitrion incubator. At day 7 cells were separated from the culture medium by centrifugation (30 minutes at 3000 g), while the supernatant containing the soluble polypeptides was filtrated over a 0.2 µm bottle top filter for further processing.

For purification purposes 1500 ml (s127H1_t2), 1800 ml (s86B4_t2), and 2400 ml (s55G7_t2) of culture supernatant was applied to a 24 ml Ni Sepharose HP column, pre-equilibrated in wash buffer (20 mM TRIS, 500 mM NaCl, pH 7.8). Following a washing step with 10 mM Imidazole in wash buffer the bound polypeptides were eluted with a step-wise gradient of 300 mM imidazole in wash buffer. The elution peaks were collected, concentrated, and applied to a size exclusion column for further purification (Superdex 200For 55G7-t2 and 127H1-t2 fractions were collected and pooled. analyzed by SDS-PAGE), ELISA and analytical size exclusion chromatography combined with multi-angle light scattering to estimate molecular mass (SEC-MALS). ELISA results confirmed binding of the polypeptides to CR6261 and CR9114, but not CR8020. SEC-MALS results are summarized in table 9.

Table 8 indicates that polypeptide s127H1-t2 has a high yield (~30 mg protein/l culture supernatant) compared to 55G7-t2 and 86B4-t2. The majority of the protein exhibits a molecular weight of 62 kDa, which is in between what is expected for a monomer or a dimer. To confirm the aggregation state of the protein the SEC-MALS experiment was repeated in the presence of Fab-fragments derived from CR6261, CR9114 and CR8020. Results are summarized in Table 8.

The results show that the soluble form of polypeptide s127H1-t2 forms a complex (as evidenced by the shift of the peak in SEC chromatogram) in the presence of the Fab fragments from CR6261 and CR9114, but not with CR8020. This is in line with the specificity of the binding reactions of the Fab fragments, since CR6261 and CR9114 bind to HA's derived from group 1, whereas CR8020 does not. The size of the complex is listed in table s, and indicates that polypeptide s127H1-t2 binds one to two Fab fragments, indicating that at least part of the population of purified polypeptide s127H1-t2 is in dimeric form.

To further analyze the binding reaction between polypeptide 127H1-t2 and mAb's CR6261 and CR9114, as well as to confirm the presence of the conformational epitopes of CR6261 and CR9114 the complexation of these antibodies with the purified protein was studied by biolayer interferometry (Octet Red³⁸⁴, Forte Bio). To this end, biotinylated
5 CR6261, CR9114 and CR8020 were immobilized on streptavidin coated sensors, which subsequently were exposed first to a solution of the purified polypeptide to measure the rate of association and then to a wash solution to measure the rate of dissociation.

The immobilized CR6261 and CR9114 both recognize the as evidenced by a clear responses after exposure to the soluble form of 127H1-t2. In conclusion polypeptide
10 s127H1-t2 is produced in high quantities and is capable of binding broadly neutralizing monoclonal antibodies CR6261 and CR9114 with high affinity, confirming the presence of the corresponding neutralizing epitopes in this stem domain polypeptide. The polypeptide has a propensity to form dimeric structures.

15 Example 3: Disulfide stabilized trimers of the present invention

One way to improve the presentation of neutralizing epitopes on immunogen in a vaccine is to engineer additional interactions between monomer immunogens in order to create multimeric species with increased stability compared to the monomer. A disadvantage of this method is that by bringing together the monomeric immunogens important epitopes
20 can potentially be covered by the next protomer. Therefore care should be taken to avoid this. Polypeptides of the invention described here are derived from the hemagglutinin molecule of influenza. This molecule is a trimer in its native state on the viral membrane. Here we describe modified polypeptides of the invention that form stable trimers in solution while exposing the epitopes of neutralizing mAb CR6261 and CR9114.

25 To generate a trimeric polypeptide of the invention stabilizing interactions promoting trimerization of monomeric species of HA-stem based polypeptides were designed, focusing in particular on creating covalent disulfide bridges between individual monomers in the trimer. To this end the three dimensional structures of FL HA from H1N1 A/South Carolina/1./18 (PDB 1RD8) in its uncleaved state and H1N1 A/California/04/2009 (PDB:
30 3LZG) were analyzed to identify areas where the proximity of another monomer and the conformational features of the protein could potentially allow the formation of an intermonomer disulfide bridge. Eleven pairs of residues for which the inter-residue distance

was less than 3.5 Å were identified (table 9). Care was taken to ensure that for each pair the residues were located on a different protomer (monomer) in the trimeric structure. The equivalent of these residues (as determined from sequence alignment) in polypeptide 127H1-t2 (SEQ ID NO: 160 to 170 and 55G7-t2 (SEQ ID NO: 149-159) were mutated into cysteine, with the intent to form trimeric polypeptides covalently linked through formation of disulfide bridges. Taking into account the 3-fold symmetry of the trimeric HA molecule successful designs lead to formation of three interprotomer disulfide bridges, covalently connecting each of the monomers in the trimer to the two other monomers.

To test for the presence of the designed disulfide bridges and the neutralizing epitopes of CR9114 and CR6261, soluble forms of the designed polypeptides were expressed in HEK293F cells. Soluble forms were created by deleting the equivalent of residue 530-565 (numbering refers to SEQ ID NO: 1) from SEQ ID NO: 149 to 170 to create polypeptides SEQ ID NO: 171 to 192. An additional sequence EGRHHHHHHH (SEQ ID NO: 19) was added at the C-terminus, in effect introducing a 7 histidine purification tag preceded by a Factor X proteolytic cleavage site to aid with purification and/or detection.

To produce the polypeptides 1.0×10^6 vc/mL were seeded by spinning down HEK293F cells (Invitrogen) at 300 g for 5 min and resuspending in 30 mL pre-warmed Freestyle™ medium per SF250 flask. This culture was incubated for 1 hour at 37 °C, 10% CO2 at 110 rpm in a multitron incubator. After 1 hour the plasmid DNA was pipetted in 1 mL Optimem medium to a concentration of 1.0 µg/mL in the 30 mL culture volume. In parallel 44 µL 293fectin® was pipetted in 1 mL Optimem medium and incubated for 5 minutes at room temperature. After 5 minutes the plasmid DNA/Optimem mix was added to the 293fectin®/Optimem mix and incubated at room temperature for 20 minutes. After the incubation the plasmid DNA/293fectin® mix was added drop wise to the cell suspension. The transfected cultured was incubated at 37 °C, 10% CO2 and 110 rpm in a multitron incubator. At day 7 cells were separated from the culture medium by centrifugation (30 minutes at 3000 g), while the supernatant containing the soluble polypeptides of the invention was filtrated over a 0.2 µm bottle top filter for further processing.

Culture medium was collected and was analyzed in a sandwich ELISA for the presence of multimeric forms of polypeptides that present two or more epitopes of the broadly neutralizing antibody CR9114. First, CR9114 coated plates were used to capture the expressed polypeptides directly from the culture medium. Second, biotinylated CR9114 in combination with HRP-conjugated streptavidin was used for detection of CR9114 captured

polypeptides of the invention. As a control soluble purified full length HA from H1N1 A/Brisbane/59/2007 in trimeric and monomeric form was included in the assay (Fig 2A). The results are shown in figure 2.

The trimeric full length HA shows a clear signal at low dilution, but at a concentration of approximately 0.0001 µg/ml and lower this signal is no longer detectable (Fig. 2A). For the monomeric full length HA a signal is also observed at low dilution but intensity is much lower and the signal is no longer detectable at approximately 0.02 µg/ml and lower. Most likely the signal is caused by some residual trimer that could not be separated from monomer during purification or has formed from the monomer over time.

Soluble forms of H1 mini2-cluster1+5+6-GCN4 (SEQ ID NO: 52) (Fig 2A) and, 127H1 (SEQ ID NO:55) Fig 2B only show low intensity signals, indicating low concentrations or the absence of multimeric polypeptides exhibiting the epitope of CR9114 in solution. A soluble forms of 127H1-t2 (SEQ ID NO: 81) Fig 2B exhibits a clear response in this assay, indicating the presence of some multimeric species but the intensity of the observed signals are low compared to the full length HA trimer.

The results for the soluble polypeptides based on 55G7-t2 with additional introduced cysteines are shown in figure 2C. For most peptides no or only very low signals are observed, indicating that no multimeric species presenting the epitope of CR9114 is present in the culture medium. A notable exception is polypeptide s55G7-t2-cl18long (SEQ ID NO: 175; additional cysteines introduced at positions 411 and 419; numbering refers to SEQ ID NO: 1). The only other polypeptide that shows a detectable response is s55G7-t2-cl14long (SEQ ID NO: 171; additional cysteines introduced at position 423 and 424) but signals are lower and disappear at lower dilution.

The results for the polypeptides based on 127H1-t2 are shown in figure 2D. In this case a clear response is observed for the polypeptides s127H1-t2-cl14long (SEQ ID NO: 182; additional cysteines at position 423 and 424), s127H1-t2-cl15long (SEQ ID NO: 183; additional cysteines at 430 and 431), s127H1-t2-cl17long (SEQ ID NO: 185; additional cysteines at 405 and 429), and s127H1-t2-cl24long (SEQ ID NO: 191; additional cysteines at 344 and 467), and to a lesser extent for s127H1-t2-cl19long (SEQ ID NO: 187; additional cysteines at 38 and 390) and s127H1-t2-cl23long (SEQ ID NO: 190; additional cysteines at 342 and 460). A low but detectable response is observed for s127H1-t2-cl16long (SEQ ID NO: 184; additional cysteines at 404 and 433). However, as in the case of 55G7-t2 the best

result is obtained for variant s127H1-t2-cl18long with additional cysteines introduced at positions 411 and 419 (SEQ ID NO: 186)

To further characterize the polypeptides of the invention with additional cysteines the culture supernatant was analysed by Western Blot using protocols well established in the art.

5 For detection purposes a polyclonal antibody directed against the HA-protein of H1N1 (A/California/04/2009) was used. For a trimer under non-reducing conditions, i.e. when disulfide bridges are intact a protein band at ~90 kDa or above (depending on extent of glycosylation) is expected, whereas under reducing conditions a band close to 35 kD (corresponding to the glycosylated monomeric polypeptide of the invention) is expected. The results are shown in Figure 3A and B. For additional cysteine containing variants of s55G7-t2 under reducing conditions strong signals are observed for s55G7-t2-cl18long and s55G7-t2-cl22long, and to a lesser extent s55G7-cl14long. Under non-reducing conditions a smear of proteins of different sizes above 100 kDa was observed for s55G7-t2-cl18long and s55G7-t2-cl22long, indicating that covalently cross-linked stem domain polypeptides are present in these samples. A smear is also observed for s55G7-cl14long, but intensity is lower than observed for s55G7-cl18 and s55G7-cl22.

The results for the Western Blots of the additional cysteine containing polypeptides derived from 127H1-t2 under reducing conditions (Fig 3C) indicate strong signals for s127H1-t2-cl14long, s127H1-t2-cl15long, s127H1-t2-cl16long, s127H1-t2-cl17long, and s127H1-t2-cl18long. For s127H1-t2-cl17long and s127H1-t2-cl18long a defined protein band close to 100 kDa is observed under non-reducing conditions (Fig 3D), indicative of the presence of covalently cross-linked stem domain polypeptides. Polypeptides s127H1-t2-cl14long and s127H1-t2-cl15long also show some intensity around 100 kDa on the Western blot but signal is not as strong as for s127H1-t2-cl17long, and in particular s127H1-t2-cl18long. The disulfide bridge (cl18) in this construct connects the B-loop of one monomer to the top of the CD-helix of another monomer as indicated in figure 4 and gave the strongest results in both in the background of 55G7-t2 and 127H1-t2.

Lu et al (PNAS 2013) describe a HA stem domain polypeptide that contains multiple intermonomer disulphide bridges. Their construct is produced in an *E.coli* based cell free system and is in contrast to the proteins described here an unfolded protein and needs to be refolded. The stem based polypeptide by Lu et al contains a foldon trimerization domain at the C-terminus and monomers are covalently connected through multiple disulphide bonds. Disulfide bonds described are located either in the foldon trimerization domain or in the HA

derived part of the HA-stem polypeptide. Four potential disulphide bonds in the HA derived part of the polypeptide are described, including cysteines at position 423 and 424 (cluster 14) and 430 and 431 (cluster 15). In the described stem domain polypeptide the best results were obtained with cysteines at positions 430 and 431 (cluster 15), although trimerization could also be observed for cysteines at position 423 and 424 (cluster 14). In both cases an additional disulfide bridge was present in the C-terminal foldon domain. Surprisingly, results here show that in the absence of a disulfide linked C-terminal trimerization domain an engineered disulfide covalently connecting two different monomers through cysteines at position 411 and 419 leads to higher amounts of trimeric stem domain polypeptide. In conclusion we have shown that introduction of strategically placed disulphide pairs can lead to multimerization of HA stem based polypeptides. In particular simultaneous introduction of cysteines at position 411 and 419 (cluster 18) leads to formation of multimeric species in solution as evidenced from the Western Blot and sandwich Elisa results.

Example 4: Purification and characterization of trimeric polypeptide of the invention

To further characterize the polypeptide of the invention 127H1-t2-cl18, the protein was purified. To facilitate purification the transmembrane and cytosolic domain at the C-terminal end of the protein can be removed as described above to create a soluble version of the protein.. It will be understood by the skilled person that the leader sequence (or signal sequence) that directs transport of a protein during production (corresponding to amino acids 1-17 of SEQ ID NO: 1) will not be present in the secreted final polypeptide. A non-limiting example of a soluble polypeptides of the invention is s127H-t2-cl18long (SEQ ID NO: 186) .

In order to obtain a highly pure preparation of a polypeptide of the invention, HEK293F cells were transfected with expression vector pcDNA2004 containing the gene encoding of polypeptide of the invention s127H1-t2-cl18long (SEQ ID NO: 186) containing an additional C-terminal his tag sequence (EGRHHHHHHH) and cultured for 7 days following protocols well established in the art. For purification purposes 300 ml of culture supernatant was applied to a 5 ml His Trap column, pre-equilibrated in wash buffer (20 mM TRIS, 500 mM NaCl, pH 7.8). Following a washing step with 10 mM Imidaze in wash buffer the bound polypeptide of the invention was eluted with a step-wise gradient of 300 mM imidazole in wash buffer. The elution peaks were collected, buffer exchanged, concentrated, and applied to a size exclusion column for further purification (Superdex 200). The elution

profile is shown in figure 5A, fractions 1-4 were collected as indicated in the figure and analyzed by SDS-PAGE (Fig 5B), NATIVE PAGE (Fig 5C) and Western blot (Fig 5D).

Under non-reducing conditions SDS PAGE shows a clear band for fraction 2 and 3 between 100 and 150 kD, as expected for a covalent linked trimeric polypeptide of the invention, whereas fraction 4 shows a diffuse band centered around 37 kD, close to the size expected for a monomeric polypeptide of the invention. The variation in size is a result from the variation in the extent of glycosylation of the polypeptide and has been observed for other stem domain polypeptides derived from HA. Upon reduction of disulfide bridges the major band in fraction 3 shifts to ca 37 kD, very similar to the band observed for fraction 4 indicating that reduction leads to monomerization. For fraction 2 the shift cannot clearly be discerned. Fraction 1 shows proteins of a range of sizes without a clear major band.

Native PAGE (non-reducing conditions) shows a clear difference in size between the protein in fraction 3 and 4, with major bands between 146 - 240 kD, and below 66 kD, respectively. For fraction 2 a weak signal between 146 and 240 kD is observed, whereas no protein can be detected for fraction 1, probably due to formation of large aggregates.

The Western blot data (non-reducing conditions) using a polyclonal anti-His for detection confirm that the major band in fraction 3 is histidine tagged material since a clear band is observed between 100 and 150 kD. For fraction 2 a weak signal is observed around the expected size of a trimer, but higher order oligomers are also detected, whereas for fraction 4 a weak and diffuse signal is observed around 37 kD. These data confirm that the major bands in fraction 2, 3 and 4 are derived from H1 HA. No signal was observed for the protein in fraction 1.

The presence of the neutralizing epitopes of CR6261, CR9114 and CR8020 was determined by ELISA, using coating of anti-His-tag monoclonal antibody to capture the his-tagged polypeptide of the invention. After binding of the mAb under study a secondary antibody conjugated to horse radish peroxidase was used for detection. As controls monomeric and trimeric full length H1 HA (antigen) as well as polyclonal serum directed against H1 HA (for detection) were included. The results are shown in figure 6. Binding to mAb CR6261 and CR9114 is clearly observed for fraction 2, 3 and 4, as well as for the monomeric and trimeric FL HA. For fraction 1 only a weak signal is detected for CR9114 binding, and hardly any signal for binding to CR6261. In none of the fractions binding to CR8020 (mAb specific for HA from group 2) is observed. The monomeric FL HA is recognized by the polyclonal anti-H1 HA, but the response to the trimeric FL HA is much

less, possibly due to occlusion of some of the epitopes in the trimer vs the monomer . A similar pattern is observed between the results of fraction 3 (trimeric on SDS-PAGE) and fraction 4 (monomeric on SDS-PAGE), in agreement with the presence in fraction 3 of a well-folded trimeric form of the polypeptide of the invention in solution.

5 Fractions 1-4 were also tested in a CR9114 sandwich ELISA to detect multimeric polypeptides of the invention as described above (see figure 7). For reasons of comparison monomeric and trimeric FL HA were again included in the experiment. Fraction 3 exhibits a response very similar to the trimeric FL HA, in agreement with a trimeric form of the polypeptide of the invention, whereas the response for fraction 2 and 4 is intermediate
10 between the monomeric and trimeric FL HA.

Complex formation between Fab fragments of CR6261, CR9114 and CR8020 and s127H1-t2-cl18long was studied by analytical size exclusion chromatography combined with multi-angle light scattering to estimate molecular mass (SEC-MALS) for the protein in fraction 3 (Figure 8). The results show that the polypeptide present in fraction 3 has a
15 molecular weight of ca. 110 kD, in line with formation of a trimer (calculated monomeric molecular weight based on amino acid sequence, excluding glycosylation is 29.2 kD). Polypeptide of the invention s127H1-t2-cl18long forms a complex (as evidenced by the shift of the peak in SEC chromatogram) in the presence of the Fab fragments from CR6261 and CR9114, but not with CR8020. This is in agreement with the specificity of the binding
20 reactions of the Fab fragments, since CR6261 and CR9114 bind to HA's derived from group 1, whereas CR8020 does not. The size of the complex formed are ca 215 and 248 kD with Fab fragments of CR6261 and CR9114, respectively and indicate that polypeptide 127H1-t2-cl18 can bind 3 Fab fragments (the expected molecular weight for a trimer in complex with 3 Fab fragments is ca 250 kD; molecular masses as derived from the SEC-MALS experiments
25 are summarized in table 10)

Example 5: Characterization of polypeptides of the invention

To further analyze the binding reaction between polypeptide of the invention 127H1-t2-cl18 and mAb's CR6261 and CR9114 as well as to reconfirm the presence of the
30 conformational epitopes of CR6261 and CR9114 the complexation of these antibodies with purified protein was studied by biolayer interferometry (Octet Red³⁸⁴, Forte Bio). To this end, biotinylated CR6261, CR9114 and CR8020 were immobilized on streptavidin coated sensors,

which subsequently were exposed first to a solution of the purified polypeptide of the invention s127H1-t2-cl18 to measure the rate of association and then to a wash solution to measure the rate of dissociation. The results are shown in figure 9.

The immobilized CR6261 and CR9114 both recognize the polypeptide of the invention as evidenced by the clear responses after exposure to the soluble form of s127H1-t2-cl18 (Figure 9A and B). To estimate the dissociation constant for the binding interaction a titration was performed using a 2-fold dilution series. Sensors containing immobilized CR6261 or CR9114 were exposed to soluble s127H1-t2-cl18 long solutions at concentrations of 10, 5, 2.5, 1.3 and 0.63, 0.31 and 0.16 nM, respectively, and the final response after 6600 seconds recorded. The responses were plotted as a function of the stem domain polypeptide concentration, and a fit to a steady state binding model was performed, yielding a dissociation constant K_d of 0.7 nM for the CR6261/stem domain polypeptide complex and 0.5 nM for the CR9114 complex (Figure 9C and D).

In conclusion polypeptide of the invention s127H1-t2-cl18 forms a covalent trimer that is capable of binding broadly neutralizing monoclonal antibodies CR6261 and CR9114 with high avidity, confirming the presence of the corresponding neutralizing epitopes in this stem domain polypeptide. The stoichiometry of the binding reaction in solution is 1:3, indicating that the neutralizing epitopes are present in each individual monomer of the trimer.

Example 6: Evaluation of protective efficacy of a polypeptide of the invention in a lethal influenza challenge model

In order to further evaluate the protective efficacy of polypeptides of the invention s127H1-t2-cl18 long (SEQ ID NO: 186) in a lethal influenza challenge model, groups of 8-14 female BALB/c mice (age 6-8 weeks) were immunized 1, 2 and 3 times at 3 week intervals with 30 µg of purified s127H1-t2-cl18 long adjuvated with 10 µg Matrix-M. As a positive control for the challenge model, broadly neutralizing antibody monoclonal antibody CR6261 (15 mg/kg) was administered i.v. 1 day prior to challenge, while immunization with PBS served as a negative control. Four weeks after the last immunization mice were challenged with 25xLD50 heterologous challenge virus (H1N1 A/Puerto Rico/8/34) and monitored daily (survival, weight, clinical scores) for 3 weeks. Pre-challenge serum (obtained 4 weeks after final immunization) was tested in ELISA assays for binding to polypeptide of the invention s127H1-t2-cl18 long that was used for

immunization (to verify correct immunization), binding to soluble H1N1 A/Brisbane/59/07 full length HA (to verify recognition of full length HA) and competition with the broadly neutralizing antibody monoclonal antibody CR9114 for binding to full length HA (to determine whether induced antibodies bind at close proximity to the broadly neutralizing CR9114 epitope). The results are shown in Figures 10-15.

The results show that the experiment is valid since all mice in the PBS control group succumb to infection at day 7 post challenge, whereas the positive control group (15mg/kg CR6261, 1 day before challenge) is fully protected (Figure 10A). Nine out of 10 mice immunized once with s127H1-t2-cl18long (SEQ ID NO: 186) survived the lethal challenge, and all mice survived after two or three immunizations (Figure 11). In addition, bodyweight loss was negligible for animals that were immunized two or three times (Figure 12), and no (2 or 3 times) or minimal (1 time) clinical signs were observed (Figure 13). Compared to the PBS control group, a statistical significant increased survival proportion, increased survival time, a decrease of body weight loss and reduced clinical scores were observed for all groups immunized with polypeptide of the invention s127H1-t2-cl18long

The ELISA data from pre-challenge timepoints 4 week after the final immunization using s127H1-t2-cl18long (Figure 14A) or the soluble full length HA (Figure 14B) as the antigen indicate that the polypeptide of the invention s127H1-t2-cl18long is immunogenic and induces antibodies that are capable of recognizing full length HA even after one immunization, although levels are substantially higher after two and three immunizations.

To further understand the immunological response to the immunization a competition binding ELISA was performed. To this end plate bound full length HA was incubated with serial diluted serum samples, after which CR9114-biotin at a predetermined titrated concentration was added. After further incubation, the amount of CR9114-biotin bound was quantified using streptavidin -conjugated horse radish peroxidase following protocols well known in the art. Data were analysed using robust linear regression of OD versus log dilution, expressed as 'slope OD' ($\Delta OD/10$ fold dilution). The data show that levels of antibodies that are capable of competing for

binding with the broadly neutralizing antibody CR9114 are induced by immunization with adjuvated polypeptides of the invention. After two immunizations the levels are clearly above background, and they continue to rise after the third immunization (Figure 15, top). As a comparison levels induced by unlabeled CR9114 (i.e. self-competition) and the non-binding monoclonal antibodies CR8020, both serially diluted from 5 µg/ml starting concentration are indicated in a separate graph (Figure 15, bottom).

In conclusion we have shown that immunization with polypeptide of the invention s127H1-t2118long (SEQ ID NO: 186) can protect mice against lethal infection with influenza, even after a single immunization round. The polypeptide is immunogenic and induces antibodies that can bind to full length HA. At least part of the induced antibodies bind at, or close to, the epitope of the broadly neutralizing epitope of monoclonal antibody CR9114.

Example 7: Evaluation of protective efficacy of a polypeptide of the invention in a lethal heterosubtypic H5N1 influenza challenge model

In order to further evaluate the protective efficacy of polypeptides of the invention s127H1-t2-cl18long (SEQ ID NO: 186) in a lethal H5N1 influenza challenge model, groups of 8-12 female BALB/c mice (age 6-8 weeks) were immunized 3 times at 3 week intervals with 30 µg of purified s127H1-t2-cl18long adjuvated with 10 µg Matrix-M. As a positive control for the challenge model, broadly neutralizing antibody monoclonal antibody CR6261 (15 mg/kg) was administered i.v. 1 day prior to challenge, while immunization with PBS served as a negative control. Four weeks after the last immunization mice were challenged with 12.5xLD50 heterosubtypic challenge virus (H5N1 A/Hong Kong/156/97) and monitored daily (survival, weight, clinical scores) for 3 weeks. The results are shown in Figure 16.

The results show that the experiment is valid since all mice in the PBS control group succumb to infection between day 8-10 post challenge, whereas the positive control group (15mg/kg CR6261, 1 day before challenge) is fully protected (Figure 16A). Ten out of 10 mice immunized with s127H1-t2-cl18long (SEQ ID NO: 186) survive the lethal challenge (Figure 16B). In addition bodyweight loss is negligible for these animals (Fig 16C) and no clinical symptoms were observed during the follow-up period (Figure

16D). Compared to the PBS control group, a statistical significant increased survival proportion, increased survival time, a decrease of body weight loss and reduced clinical scores are observed for the group immunized with polypeptide of the invention s127H1-t2-cl18long.

5 In conclusion we have shown that immunization with polypeptide of the invention s127H1-t2cl18long (SEQ ID NO: 186) can protect mice against lethal infection with a heterosubtypic H5N1 influenza strain.

10 Example 8: Evaluation of the breadth of binding of sera elicited through immunization with a polypeptide of the invention

 The results described in Example 6 indicate that polypeptide of the invention s127H1-t2-cl18long (SEQ ID NO: 186) is immunogenic and can elicit antibodies that are capable of recognizing FL HA from the strain used as the basis for design of polypeptides of the invention. Sera from mice immunized 3 times as described in example 7 were also
15 tested for binding against full length HA's from a number of other group 1 (H1, H5 and H9) and group II (H3 and H7) influenza strains by ELISA following protocols well known in the art (Figure 17). The results demonstrate that antibodies induced with polypeptide of the invention s127H1-t2-cl18long (SEQ ID NO: 186) efficiently recognize epitopes present in the native sequences of FL HA and that the epitopes to which the
20 antibodies bind are conserved among different group 1 (in particular H1, H5 and H9) and even some group 2 influenza strains (e.g. H7).

Example 9: Evaluation of protective efficacy of a polypeptide of the invention in a lethal H1N1 A/Brisbane/59/2007 influenza challenge model

25 In order to further evaluate the protective efficacy of polypeptides of the invention s127H1-t2-cl18long (SEQ ID NO: 186) in a lethal H1N1 influenza challenge model, groups of 8-18 female BALB/c mice (age 6-8 weeks) were immunized 3 times at 3 week intervals with 30 µg of purified s127H1-t2-cl18long adjuvated with 10 µg Matrix-M. As a positive control for the challenge model, broadly neutralizing antibody monoclonal
30 antibody CR6261 (15 mg/kg) was administered i.v. 1 day prior to challenge, while immunization with PBS served as a negative control. Four weeks after the last

immunization mice were challenged with 12.5xLD50 challenge virus (H1N1 A/Brisbane/59/2007) and monitored daily (survival, weight, clinical scores) for 3 weeks.

The results show that the experiment is valid since all mice in the PBS control group succumb to infection between day 7-10 post challenge, whereas the positive control group (15mg/kg CR6261, 1 day before challenge) is fully protected (Figure 18A). Ten out of 10 mice immunized with s127H1-t2-cl18long (SEQ ID NO: 186) survive the lethal challenge (Fig 18B). In addition bodyweight loss is ca 10% on average 4 days post infection (Fig 18C), but animals recover fully within 10 days. Clinical symptoms also peak at 4 days post infection but are absent from day 9 post infection onwards (Fig 18D).

Compared to the PBS control group, a statistical significant increased survival proportion, increased survival time, a decrease of body weight loss and reduced clinical scores are observed for the group immunized with polypeptide of the invention s127H1-t2-cl18long

In conclusion we have shown that immunization with polypeptide of the invention s127H1-t2cl18long (SEQ ID NO: 186) can protect mice against lethal infection with

H1N1 A/Brisbane/59/2007.

Example 10: Evaluation of the presence of influenza neutralizing antibodies in sera of mice immunized with the polypeptide of the invention

To further investigate antibody-mediated effector mechanisms that play a role in protection against influenza, pre-challenge sera were tested in pseudoparticles neutralization assay (Alberini et al 2009) using the pseudoparticles derived from H5N1 A/Vietnam/1194/04 as described below.

Pseudoparticle neutralization assay

Pseudoparticles expressing FL HA were generated as previously described (Temperton et al., 2007). Neutralizing antibodies were determined using a single transduction round of HEK293 cells with H5 A/Vietnam/1194/04 pseudoparticles encoding luciferase reporter gene, as described previously (Alberini et al 2009), with a few modifications. Briefly, heat-inactivated (30 minutes at 56°C) pre-challenge serum samples were 3-fold serially diluted in growth medium (MEM Eagle with EBSS (Lonza, Basel, Switzerland) supplemented with 2 mM L-Glutamine (Lonza), 1% Non-Essential

Amino Acid Solution (Lonza), 100 U/ml Pen/Strep (Lonza) and 10% FBS (Euroclone, Pero, Italy)) in triplicate in 96-well flat bottom culture plates and a titrated number of H5 A/Vietnam/1194/04 pseudoparticles (yielding 10^6 Relative Luminescence Units (RLU) post-infection) was added. After 1h incubation at 37°C, 5% CO₂ 10^4 HEK293 cells were
 5 added per well. After 48h incubation at 37°C, 5% CO₂ luciferase substrate (Britelie Plus, Perkin Elmer, Waltham, MA) was added and luminescence measured using a luminometer (Mithras LB 940, Berthold Technologies, Germany) according to manufacturers' instructions.

Pre challenge sera obtained from animals immunized with polypeptide of the
 10 invention s127H1-t2118long (SEQ ID NO: 186) as described in examples 6, 7 and 8 showed detectable neutralization at high serum concentrations using the pseudoparticle neutralization assay (Figure 19). This demonstrates the ability of the polypeptide of the invention to elicit broadly neutralizing antibodies when used as an immunogen. Besides direct virus neutralization, Fc-mediated effector mechanisms, such as Antibody
 15 Dependent Cellular Cytotoxicity (ADCC) and Antibody Dependent Cellular Phagocytosis (ADCP), contribute substantially to protection against influenza, with stem-directed bnAbs being particularly effective in these mechanisms (DiLillo et al. 2014) In order to test whether the antibodies elicited after immunization with polypeptide of the invention s127H1-t2118long (SEQ ID NO: 186) were capable of inducing ADCC, we
 20 tested pre-challenge sera using an ADCC surrogate assay (*Parekh et al., 2012*; *A. Schnueriger et al., 2012*; *Cheng et al 2014*) adapted for mouse as described below.

Antibody Dependent Cellular Cytotoxicity (ADCC) surrogate assay

Human lung carcinoma-derived A549 epithelial cells (ATCC CCL-185) were
 25 maintained in Dulbecco's modified eagle medium (DMEM) medium supplemented with 10% heat inactivated fetal calf serum at 37°C, 10% CO₂. Two days before the experiment, A549 cells were transfected with plasmid DNA encoding H5 A/Hong Kong/156/97 HA or H1 A/Brisbane/59/2007 HA using Lipofectamine 2000 (Invitrogen) in Opti-MEM (Invitrogen). One day before the assay, transfected cells were harvested
 30 and seeded in white 96-well plates (Costar) for ADCC, and black clear bottom 96-well plate (BD Falcon) for imaging. After 24 hours, samples were diluted in assay buffer (4%

ultra-low IgG FBS (Gibco) in RPMI 1640 (Gibco)) and heat inactivated for 30 minutes at 56°C, followed by serial dilution in assay buffer. For the ADCC bioassay, A549 cells were replenished with fresh assay buffer and antibody dilutions and ADCC Bioassay Jurkat effector cells expressing mouse Fc gamma receptor IV (FcγRIV; Promega) were added to the cells and incubated for 6 hours at 37°C at a target-effector ratio of 1:4.5. Cells were equilibrated to room temperature for 15 min before Bio-Glo Luciferase System substrate (Promega) was added. Luminescence was read out after 10 minutes on a Synergy Neo (Biotek). Data are expressed as fold induction of signal in the absence of serum.

Using this assay pre-challenge sera obtained from animals immunized with polypeptide of the invention s127H1-t2118long (SEQ ID NO: 186) as described in examples 6, 7 and 8 were tested for FcγRIV signaling activity using target cells transfected with FL HA from H5N1 A/Hong Kong/156/97 or H1N1 A/Brisbane/59/07 as the source of antigen (Figure 20). In both cases a 30 to 40 fold induction at highest serum concentration tested is observed demonstrating the ability of the polypeptide of the invention to elicit antibodies that activate FcγRIV signaling, indicative for ADCC/ADCP effector function in mice.

These results shown in examples 6-9 show the capability of polypeptide of the invention s127H1-t2118long (SEQ ID NO: 186) is able to elicit stem-targeting, neutralizing and ADCC-mediating antibodies and protect mice against a lethal challenge with homologous, heterologous and heterosubtypic group I influenza strains.

Example 11: Design of a polypeptide of the invention of based on the HA from H1N1 A/California/07/09.

Examples 1 to 10 describe polypeptides of the invention based on the HA from H1N1 A/Brisbane/59/2007. Similar polypeptides can also be designed based on other HA from other influenza strains such as H1N1 A/California/07/09. So following the procedures outlined above the polypeptides of the invention described in SEQ ID NO: 202 and 203 were created.

Example 12: *Design of further polypeptides of the invention based on the HA from H1 strains.*

Examples 1 to 11 describe polypeptides of the invention based on the HA from H1N1 A/Brisbane/59/2007 (SEQ ID NO: 1) and H1N1 A/California/07/09 (SEQ ID NO: 252). Similar polypeptides can also be designed based on HA from other H1 influenza strains, and these are also included in the invention. Non-limiting examples are the HA's of H1N1 A/Texas/UR06-0526/07 (SEQ ID NO: 205), H1N1 A/New York/629/95 (SEQ ID NO: 206) and H1N1 A/AA_Marton/43 (SEQ ID NO: 204). So following the procedures outlined above stem domain polypeptides of the invention containing engineered disulfide bridges between cysteines at position 411 and 419 (cluster 18) described in SEQ ID NO: 207 to 216 were created. SEQ ID NO: 202, 203, 213 and 214 contain an additional lysine mutation introduced at position 415 (numbering refers to SEQ ID NO: 1) to create a B-loop sequence according to SEQ ID NO: 8; these sequences are also included in the invention.

Example 13: *Expression and characterization of polypeptides of the invention*

To test for the presence of the designed disulfide bridges and the neutralizing epitopes of CR9114 and CR6261, soluble forms of the designed polypeptides were expressed in HEK293F cells. Soluble forms were created by deleting the equivalent of amino acid residues 530-565 (numbering refers to SEQ ID NO: 1) from SEQ ID NO: 202, 207, 209, 213, to create polypeptides of the invention SEQ ID NO: 203, 208, 210 and 214, respectively. It is noted that these sequences describe the processed form of the polypeptides, ie, after removal of the leader sequence. An additional sequence EGRHHHHHHH (SEQ ID NO: 19) or in the case of SEQ ID NO: 208 EGRHHHHHHH was added at the C-terminus, in effect introducing a 7 or 6 histidine purification tag preceded by a Factor X proteolytic cleavage site to aid with purification and/or detection.

To produce the polypeptides 1.0×10^6 vc/mL were seeded by spinning down HEK293F cells (Invitrogen) at 300 g for 5 min and resuspending in 30 mL pre-warmed Freestyle™ medium per SF250 flask. This culture was incubated for 1 hour at 37 °C, 10% CO₂ at 110 rpm in a multitron incubator. After 1 hour the plasmid DNA was pipetted in 1 mL Optimem medium to a concentration of 1.0 µg/mL in the 30 mL culture

volume. In parallel 44 μ L 293fectin® was pipetted in 1 mL Optimem medium and incubated for 5 minutes at room temperature. After 5 minutes the plasmid DNA/Optimem mix was added to the 293fectin®/Optimem mix and incubated at room temperature for 20 minutes. After the incubation the plasmid DNA/293fectin® mix was added drop wise to the cell suspension. The transfected cultured was incubated at 37 °C, 10% CO₂ and 110 rpm in a multitron incubator. At day 7 cells were separated from the culture medium by centrifugation (30 minutes at 3000 g), while the supernatant containing the soluble polypeptides of the invention was filtrated over a 0.2 μ m bottle top filter for further processing.

Culture medium was collected and was analyzed in a sandwich ELISA for the presence of multimeric forms of polypeptides that present two or more epitopes of the broadly neutralizing antibody CR9114. First, CR9114 coated plates were used to capture the expressed polypeptides directly from the culture medium. Second, biotinylated CR9114 in combination with HRP-conjugated streptavidin was used for detection of CR9114 captured polypeptides of the invention. For reasons of comparison soluble purified full length HA from H1N1 A/Brisbane/59/2007 in trimeric and monomeric form as well as purified trimeric s127H1-t2-cl18long (SEQ ID NO: 186) was included in the assay (starting concentration 5 μ g/ml). The results are shown in figure 21A.

The trimeric full length HA shows a clear signal at low dilution, but at a dilution of 1 in 6561 or higher (i.e. lower concentration) this signal is no longer detectable (Fig. 21A). For the monomeric full length HA a signal is also observed at low dilution but intensity is much lower and the signal is no longer detectable at a dilution of 1 in 729. Most likely the signal is caused by some residual trimer that could not be separated from monomer during purification or has formed from the monomer over time. Purified trimeric s127H1-t2-cl18long (SEQ ID NO: 186) containing an additonal C-terminal his tag sequence (EGRHHHHHHH), (starting concentration 5 μ g/ml) also results in a clear signal in this assay which becomes undetectable at a concentration of 1 in 19683. The culture media containing polypeptides of the invention SEQ ID NO: 203, 208, 210 and 214 also show clear signals in this assay, indicating the presence of multimeric polypeptides of the invention. The strongest responses are observed for the polypeptide derived from A/California/07/09 (SEQ ID NO: 203). The weaker response in the assay

observed for the polypeptide of the invention derived from H1N1 A/AA_Marton/1943 (SEQ ID NO: 214) is a result from the lower expression of this particular polypeptide of the invention as evidenced from the results from a western blot of the culture media described below.

5 To further characterize the polypeptides of the invention containing additional cysteines, the culture supernatants were analysed by Western Blot using protocols well established in the art. For detection purposes a polyclonal antibody directed against the HA-protein of H1N1 (A/California/04/2009) was used. For a disulfide linked trimer under non-reducing conditions, i.e. when disulfide bridges are intact, a protein band at
10 ~90 kDa or above (depending on extent of glycosylation) is expected, whereas under reducing conditions a band close to 35 kD (corresponding to the glycosylated monomeric polypeptide of the invention) is expected. Fig 22 shows the results for the polypeptides of the invention derived from the HA from strains H1N1 A/Texas/ (SEQ ID NO: 208), H1N1 A/New York/629/95 (SEQ ID NO: 210), H1N1 A/California/07/09 (SEQ ID NO:
15 203) and H1N1 A/AA_Marton/1943 (SEQ ID NO: 214). The Western Blot under reducing conditions shows that all four polypeptides of the invention express, albeit to different levels, with the highest expression observed for H1N1 A/California/07/09 derived polypeptide of the invention (SEQ ID NO: 203) and the lowest for H1N1 A/AA_Marton/1943 derived polypeptide (SEQ ID NO: 214). Under these conditions the
20 polypeptides run as monomers in the gel similar to the the purified trimeric s127H1-t2-cl18long (SEQ ID NO: 186) containing an additional C-terminal his tag sequence (EGRHHHHHHH). Under non-reducing conditions a band between 100 and 150 kD, indicative of the presence of trimeric polypeptide of the invention is observed for all polypeptides of the invention (including s127H1-t2-cl18long (SEQ ID NO: 186)). For the
25 polypeptide derived from H1N1 A/AA_Marton/1943 (SEQ ID NO: 214) the monomeric band is no longer visible, and a band at the expected height for a trimer appears, albeit at low intensity due to the lower expression of this polypeptide. In addition, dimeric forms running at about 75 kD can also be observed. The polypeptides of the invention are somewhat heterogeneous in size due to variation in the extent of glycosylation of
30 individual polypeptides of the invention.

To further confirm the presence of the neutralizing epitopes of CR6261 and CR9114 culture supernatants were analyzed by ELISA. First, plates coated with an antibody directed to the his-tag on the soluble polypeptides of the invention were used to capture the expressed polypeptides directly from the culture medium. Second, CR9114 or CR6261 were added and HRP-conjugated goat anti-human antibody was used for detection of CR9114 or CR6261 binding to polypeptides of the invention. As a positive control soluble purified full length HA from H1N1 A/Brisbane/59/2007 in trimeric and monomeric form were included in the assay (Fig 21B, C). As a negative control an ELISA using mAb CR8020, specific for HA from group 2, was also performed. The results are shown in figure 21D. The polypeptides of the invention show clear response in the ELISA's with CR9114 and CR6261, with the highest response observed for the polypeptide derived from A/California/07/09 (SEQ ID NO: 203). Purified soluble full length HA (both monomeric and trimeric) also show strong responses. As expected, no response was observed in the ELISA with CR8020, confirming the specificity of the binding of CR6261 and CR9114 to the polypeptides of the invention.

In conclusion the results above show that polypeptides of the invention derived from H1 HA sequences as described above can form trimeric species and contain the neutralizing epitopes of CR6261 and CR9114.

Example 14: Design of further polypeptides of the invention based on the HA from Influenza group 1 strains.

Examples 1 to 13 describe polypeptides of the invention based on the HA from H1N1. Similar polypeptides can also be designed based on HA from other group 1 influenza strains, for example those containing H2, H5 and H9 HA. These polypeptides are also included in the invention. Non-limiting examples of such strains are for example H2N2 A/Adachi/2/1957, H2N2 A/Singapore/1/1957, H5N1 A/Vietnam/1203/2004 and H9N2 A/Hong Kong/69955/2008. So following the procedures outlined above the polypeptides of the invention containing engineered disulfide bridges between cysteines at position 411 and 419 (cluster 18) described in SEQ ID NO: 218 to 221, SEQ ID NO: 223 to 226, SEQ ID NO: 228 to 231, and SEQ ID NO: 233 to 236 were created.

It should be noted that the full length HA from H5N1 A/Vietnam/1203/2004 (SEQ ID NO: 227) contains a polybasic cleavage site, i.e. sequence RRRKTR at position 341-346 in SEQ ID NO: 227 directly preceding the fusion peptide sequence. In polypeptides of the invention 228-231 the polybasic cleavage site has been removed and replaced by a single glutamine (Q) residue to remove the complete cleavage site. These sequences are also encompassed in the invention.

SEQ ID NO: 219, 221, 224 and 226 contain further mutations at positions 407 and 414-416 (numbering according to SEQ ID NO: 1; please note that in the H2 sequences residues at positions 9, 10 and 139 are deleted compared to SEQ ID NO: 1) to create a B-loop sequence according to SEQ ID NO: 8 with the exception of residue 418 that is a cysteine.

SEQ ID NO: 230 and 231 contain additional mutations at positions 407 (E407T) and 415 (N415S) (numbering according to SEQ ID NO: 1). These mutations create a B-loop according to SEQ ID NO: 8 with the exception of residue 418 that is a cysteine.

SEQ ID NO: 236 and 236 have been further modified to contain the sequence MNTQYTAIGCEYNKSE (i.e a sequence according to SEQ ID NO: 8 with the exception of residue 418 that is a cysteine).

Example 15: Design of disulfide stabilized trimers of the present invention based on the HA from Influenza group 2 strains.

Examples 1 to 14 describe polypeptides of the invention based on the HA from group 1 strains. Disulfide cross-linked polypeptide can also be designed on the basis of HA sequences from group 2, such as for example H3 and H7. These polypeptides are also included in the invention. Non-limiting examples of such strains are for example H3N2 A/Hong Kong/1/1968 and H7.

One way to improve the presentation of neutralizing epitopes on an immunogen in a vaccine is to engineer additional interactions between monomer immunogens in order to create multimeric species with increased stability compared to the monomer. A disadvantage of this method is that by bringing together the monomeric immunogens important epitopes can potentially be covered by the next protomer. Therefore care should be taken to avoid this.

Here we describe modified polypeptides of the invention that form stable trimers in solution while exposing the epitopes of neutralizing mAb CR8020 and CR8043.

To generate a trimeric polypeptide of the invention stabilizing interactions promoting trimerization of monomeric species of HA-stem based polypeptides were
5 designed, focusing in particular on creating covalent disulfide bridges between individual monomers in the trimer. To this end the three dimensional structure of FL HA from H3N2 A/Hong Kong//1/1968 (PDB: 3SDY) was analyzed to identify areas where the proximity of another monomer and the conformational features of the protein could potentially allow the formation of an intermonomer disulfide bridge. Six pairs of residues
10 were identified (table 11). Care was taken to ensure that for each pair the residues were located on a different protomer (monomer) in the trimeric structure. The equivalent of these residues (as determined from sequence alignment) in stem domain polypeptides based on group 2 are then mutated into cysteine to create polypeptides of the invention so that polypeptides covalently linked through formation of disulfide bridges can be formed.
15 Taking into account the 3-fold symmetry of the trimeric HA molecule successful designs lead to formation of three interprotomer disulfide bridges, covalently connecting each of the monomers in the trimer to the two other monomers.

WO2013/079473 describes stem domain polypeptides based on HA from group 2 strains capable of binding the broadly neutralizing antibodies CR8020, CR8043. The
20 disulfide pairs from table 11 were introduced in polypeptide H3 HK mini2a-linker+c19 +10+11+12+GCN4T (SEQ ID NO: 238 here; SEQ ID NO: 130 in WO2013/079473) and H3 HK mini2a-linker+c19 +10+11+12+GCN4T-CG7-1 (SEQ ID NO: 239 here; SEQ ID NO: 174 in WO2013/079473), derived from the HA of H3N2 A/Hong Kong/1/1968 (SEQ ID NO: 237) to arrive at polypeptides of the invention 240 to 251.

25 The sequences of SEQ ID NO: 240 to 251 contain the HA leader sequence. The person skilled in the art will understand that in the mature protein the leader sequence has been cleaved off and is no longer present. The processed sequences are also included in the invention.

Soluble forms of the polypeptides of the invention can be created by deletion of
30 the C-terminal transmembrane region and cytoplasmic domain. The deletion can for example include the residues from 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535,

536 or 537 to the C-terminus. These polypeptides are also included in the invention. In some cases a C-terminal trimerization sequence, optionally connected through a short linker, optionally containing a proteolytic cleavage site can be added. An example of a trimerization domain is the foldon sequence of SEQ ID NO: 3. The processed sequences are also included in the invention.

Example 16: *Validation of a NHP pH1N1 challenge model for use in influenza vaccine protective efficacy evaluation and immunogenicity and protective efficacy of H1 mini-HA #4900 in NHP*

10 To be able to study immunogenicity and protective efficacy of UFV vaccine candidates in an alternative model, a non-human primate (NHP) challenge model in Cynomolgus macaques (*Macaca fascicularis*) using a pandemic H1N1 strain (A/Mexico/InDRE4487/2009) has previously been established at BPRC (Rijswijk, The Netherlands). This model is based on a model published in literature by
15 Safronetz et al (2011) *J Virol.* 85:1214 However, it needed to be established whether the model could be used for evaluation of protective efficacy of influenza vaccines.

The primary objectives of this study were

- a) to evaluate if a previously established pH1N1 challenge model in Cynomolgus macaques can be used to measure vaccine-mediated protective efficacy, using a seasonal vaccine (Inflexal ® V, season 2013/2014; Inflexal13/14) containing 15 µg FL HA of a
20 H1N1 strain homologous to the challenge strain used.
- b) to evaluate immunogenicity of s127H1-t2-cl18long (SEQ ID NO:186) in non-human primates.

A secondary objective was to evaluate the protective efficacy of s127H1-t2-cl18long (SEQ ID NO:186) in this pH1N1 NHP challenge model.
25

A cohort of male cynomolgus macaques were pre-screened on the presence of serum antibodies against Alpha herpes virus, STL V, SIV, SRV and flu A NP, HAI titers against the challenge virus (allowing a maximum. titer of 1/10), as well as tested for tuberculosis by Mantoux and blood test. Some animals screened had been part of a
30 previous CR8020 PK study. After screening, suitable animals were randomly allocated to 3 groups of 6 animals each, using a randomized block design taken, age, weight, HAI

titer and inclusion in PK study into account. Dataloggers measuring body temperature with an 15 minute interval were implanted abdominally, followed by an 1 month recovery period after which the immunization regimen started. One group received 2 intramuscular (i.m.) immunizations with the human dose (0.5 mL) of Inflexal V 13/14, containing 15 µg FL HA of H1N1 A/California/07/09, which is the official guideline immunization regimen for naive children advocated by healthcare agencies (CDC, RIVM). The second group received 3 i.m. immunizations with 150 µg s127H1-t2-cl18long (SEQ ID NO:186; containing an additional His-tag) protein adjuvanted with 50 µg Matrix-M in a volume of 0.5 mL. The third group was a negative control group which was administered 3 times 0.5 mL PBS i.m. All immunizations were performed with a 4-week immunization interval. Four weeks after the final immunizations animals were challenged intrabronchially with 4×10^6 TCID₅₀ H1N1 A/Mexico/InDRE4487/2009, which was the dose established during setup of the model. During the 21 day follow-up period, clinical signs were recorded daily. Animals were anesthetized on day 1, 2, 4, 6, 8, 10, 14 and 21 during which bodyweight was measured, tracheal swabs taken to determine viral load by qPCR. At the end of the study, dataloggers were removed and data analysed.

To verify immunogenicity of the administered vaccines, serum was isolated on day of immunization, as well as 5 days before the challenge. For both vaccination treatments the pre-challenge serum response was analysed for breadth of binding to a panel of influenza A group 1 and 2 FL HA capability of vaccines to induce antibodies bind at close proximity to the CR9114 epitope, using a CR9114 competition ELISA with response expressed as % competition, surrogate ADCC activity of vaccine induced antibodies (see below) and neutralisation activity of vaccine-induced antibodies using both a microneutralisation assay against the heterosubtypic H5N1 A/HK/156/97 as well a HAI assay that detects virus neutralisation mediated by FL HA head epitopes, using H1N1 A/California/07/09. The latter strain is homologous to the vaccine and challenge strains.

A surrogate ADCC activity was determined using ADCC Bioassay effector cells (a stable Jurkat cell expressing human Fc gamma receptor IIIA (FcγRIIIA), human CD3γ, and an NFAT- response element regulating a luciferase reporter gene (Promega) and

A549 target cells transiently transfected with DNA encoding FL HA of H1N1 and H5N1 strains.

Results

- 5 - No significant differences between treatment groups in tracheal viral load AUC were seen (not shown).
- Significant differences were seen in increase of body temperature (fever) post-challenge between Matrix-M adjuvanted polypeptide of the invention SEQ ID NO: 186 and PBS group during the day 0-3 ($p=0.010$) and day 0-8 ($p=0.047$) intervals (Fig. 27).
- 10 - Significant differences were seen in increase of body temperature post-challenge between Inflexal13/14 group and PBS group during the day 0-3 ($p=0.033$) interval (Fig. 27).
- Bodyweight and clinical signs were not informative due to the mild symptoms, likely predominantly determined by repeated anesthesia, and therefore not shown.
- 15 - Animal Ji0403061 (Inflexal13/14 group) died day 8 post challenge. Autopsy by veterinary pathologist determined viral pneumonia as cause of death, consistent with the high tracheal viral load up till time of death.
- An immunization regimen of 3x 150 μ g polypeptide of the invention SEQ ID NO: 186 + 50 μ g Matrix-M was immunogenic:
 - 20 • Induction of group 1 HA binding antibodies (Fig. 23)
 - Induction of CR9114 competing antibodies against 3 different group 1 HA proteins (Fig. 24)
 - Induction of antibodies that neutralizes a heterosubtypic H5N1 strain using the micro neutralization assay (Fig. 26).
 - 25 • Induction of antibodies capable of ADCC effector functions using 3 different group 1 HA proteins as target HA (Fig. 25)

Conclusion: Demonstration of vaccine mediated intervention of disease in the H1N1 influenza challenge model in Cynomolgus macaques has only been partially successful, as only fever was significantly reduced in the first 3 days after challenge. Three immunizations with 3x 150 µg H1 mini-HA #4900 + 50 µg Matrix-M is immunogenic in non-human primates. The induced antibodies are capable of binding HA and ADCC effector functions for all Group 1 full length HA's tested, and also neutralize a heterosubtypic H5 strain.

Example 17: Protection from lethal challenge with H5N1 A/Hong Kong/156/97 by passive transfer of serum from mice immunized with polypeptides of the invention

To determine the contribution of antibodies induced by polypeptides of the invention to protection observed, serum transfer studies were performed.

The aim of this study was to assess whether passive transfer (multiple dosing) of serum from mice immunized three times with s127H1-t2-cl18long (SEQ ID NO:186) in the presence of Matrix-M confers protection to a lethal challenge with H5N1 Influenza A/Hong Kong/156/97.

Groups of female BALB/c donor mice (age 6-8 weeks) were immunized 3 times at a 3 week interval with 30 µg s127H1-t2-cl18long (SEQ ID NO:186) containing a C-terminal His-tag adjuvanted with 10 µg Matrix-M or immunized with PBS. Four weeks after the last immunization (d70) serum was isolated, pooled per group and transferred in recipient mice (female BALB/c, age 6-8 weeks, n=10 per group). Each mouse received 400 µl serum i.p. on three consecutive days before challenge (d-3, -2 and -1). As a positive control for the challenge model CR6261 (15 mg/kg) was administered 1 day prior to challenge (n=8), while injection with PBS served as a negative control (n=8). On day 0, mice were challenged with 12.5xLD50 H5N1 A/Hong Kong/156/97 and monitored (survival, weight, clinical scores) for 3 weeks.

To verify immunogenicity of H1 mini-HA variants in donor mice and assess HA-specific antibody levels after transfer of serum into recipient mice, pooled serum samples of terminal bleeds (d70) of donor mice, pooled serum samples of naïve recipient mice before serum transfer (d-4) as well as individual serum samples of recipient mice after 3

serum transfers just prior to challenge (d0), were tested in ELISA for binding to FL HA from H1N1 A/Brisbane/59/07.

Results

- 5 - Survival percentages for the experimental groups are reported in Table 12
- Experiment is valid; all mice in the PBS control group succumb to infection at or before day 13 post challenge (median 9.5 days), whereas the positive control group (15mg/kg CR6261, 1 day before challenge) is fully protected ($p < 0.001$).
- 10 - Three serum transfers of serum from Matrix-M adjuvanted s127H1-t2-cl18long (SEQ ID NO:186) immunized mice into naïve recipient mice leads to significant increase in survival proportion ($p < 0.001$)(Fig 28A), increase in survival time ($p < 0.001$) (Fig 28A), decrease in bodyweight loss ($p < 0.001$) (Fig 28B) and reduction in clinical score ($p < 0.001$)(not shown), compared to the PBS serum transfer control group.
- 15 - FL HA A/Brisbane/59/07 specific antibody titers after three serum transfers are similar to levels obtained after active immunization. (Fig 29)

Conclusion: Serum components (most likely antibodies) induced by 3 times immunization with Matrix-M adjuvanted H1 mini-HA #4900 are sufficient to protect mice against lethal challenge with heterosubtypic H5N1 A/Hong Kong/156/97.

20

Example 18: Evaluation of protective efficacy in a lethal H1N1 A/Brisbane/59/07 mouse model

In order to provide further evidence for the use of polypeptides of the invention as vaccines the protective efficacy of three additional polypeptides against lethal challenge with Influenza was evaluated. To this end polypeptides of the invention SEQ ID NO: 203 and 254 (both containing C-terminal his-tags) were transiently expressed in HEK293F cells and purified as described above. In addition, polypeptide of the invention SEQ ID NO: 186 (also containing an additional C-terminal his-tag) was expressed in SF9 insect cells following procedures well known to those skilled in the art (see e.g. M. Cox,

Development of an influenza virus vaccine using the baculovirus-insect cell expression system, PhD thesis, Wageningen University Dec 2009) and purified as described above.

The aim of this study was to determine protective efficacy of polypeptides of the invention with Matrix-M in a H1N1 A/Brisbane/59/07 challenge model compared to a
5 PBS control group.

Groups of 10 female BALB/c mice (age 6-8 weeks) were immunized 3 times at a 3 week interval with 30 µg polypeptide of the invention adjuvanted with 10 µg Matrix-M. As a positive control for the challenge model CR6261 (15 mg/kg) was administered 1 day prior to challenge (n=8), while injection with PBS served as a negative control (n=16).

10 Four weeks after the last immunization mice were challenged with 12.5xLD50 challenge virus and monitored (survival, weight, clinical scores) for 3 weeks.

To verify immunogenicity of polypeptides of the invention, pre-challenge sera (day -1) were tested in ELISA assays for binding to FL HA from H1N1 A/Brisbane/59/07. To determine whether polypeptide of the invention induced antibodies
15 bind at close proximity to the CR9114 epitope, a CR9114 competition ELISA was performed. Competition data were visualized as ‘% competition’, defined as $(A - P)/A \times 100$, where A is the maximum OD signal of CR9114 binding to FL HA when no serum is present and P is the OD signal of CR9114 binding to FL HA in presence of serum at a given dilution or expressed using the slope OD metric to be able to quantify
20 responses.

Results

- Survival percentages for the experimental groups are reported in Table 12.
- Experiment is valid; all mice in the PBS control group (n=16) succumb to infection at or before day 10 post challenge (median 8 days), whereas the positive
25 control group (n=8, 15mg/kg CR6261, 1 day before challenge) is fully protected ($p < 0.001$).
- Three immunizations with polypeptides of the invention SEQ ID NO: 186, 203 and 254 adjuvanted with Matrix-M lead to significant increase in survival proportion ($p < 0.001$)(Fig 30A), increase in survival time ($p < 0.001$) (Fig. 30A),

decrease in bodyweight loss ($p < 0.001$) (Fig 30B) and reduction in clinical score ($p < 0.001$) (not shown), compared to the PBS control group.

- Pre-challenge IgG antibody titers to H1N1 A/Brisbane/59/07 FL HA induced by polypeptides of the invention are significantly higher compared to PBS for all polypeptides of the invention tested ($p \leq 0.001$) (Fig. 31A).

- IgG antibody titers to H1N1 A/Brisbane/59/07 FL HA plateau after two immunizations for all polypeptides of the invention tested (not shown).

- Matrix-M adjuvanted polypeptides of the invention SEQ ID NO: 186, 203 and 254 induce significantly higher CR9114 competition titers compared to PBS ($p < 0.001$) (Fig. 31B).

- Matrix-M adjuvanted polypeptide of the invention SEQ ID NO: 203 (background H1N1 A/California/07/2009) induces significantly lower CR9114 competing antibodies compared to other polypeptides of the invention tested ($p \leq 0.013$) when H1N1 A/Brisbane/59/07 FL HA is used as a target antigen (Fig. 31B).

Conclusion: Matrix-M adjuvanted additional polypeptide of the invention SEQ ID NO: 186, 203 and 254 confer protection against lethal challenge with H1N1 A/Brisbane/59/07.

Example 19: Evaluation of protective efficacy in a lethal H1N1 A/NL/602/09 mouse model

In order to provide further evidence for the use of polypeptides of the invention as vaccines the protective efficacy of three additional polypeptides against lethal challenge with Influenza was evaluated. To this end polypeptides of the invention SEQ ID NO: 203 and 254 (both containing C-terminal his-tags) were transiently expressed in HEK293F cells and purified as described above. In addition polypeptide of the invention SEQ ID NO: 186 (also containing an additional C-terminal his-tag) was expressed in SF9 insect cells following procedures well known to those skilled in the art (supra) and purified as described above.

The aim of this study was to determine protective efficacy of additional trimeric H1 mini-HA variants adjuvanted with Matrix-M in a H1N1 A/NL/602/09 challenge model compared to a PBS control group.

Groups of 10 female BALB/c mice (age 6-8 weeks) were immunized 3 times at a 3 week interval with 30 µg polypeptide of the invention adjuvanted with 10 µg Matrix-M. As a positive control for the challenge model CR6261 (15 mg/kg) was administered 1 day prior to challenge (n=8), while injection with PBS served as a negative control (n=16). Four weeks after the last immunization mice were challenged with 12.5xLD50 challenge virus and monitored (survival, weight, clinical scores) for 3 weeks.

To verify immunogenicity of polypeptides of the invention, pre-challenge sera (day -1) are tested in ELISA assays for binding to FL HA from H1N1 A/Brisbane/59/07. To determine whether mini-HA induced antibodies bind at close proximity to the CR9114 epitope, a CR9114 competition ELISA was performed. Competition data were visualized as '% competition', defined as $(A-P)/A \times 100$, where A is the maximum OD signal of CR9114 binding to FL HA when no serum is present and P is the OD signal of CR9114 binding to FL HA in presence of serum at a given dilution or expressed using the slope OD metric to be able to quantify responses; for reference CR9114 and CR8020 (starting concentration 5 mg/ml) solutions were included.

Results

- Survival percentages for the experimental groups are reported in Table 12
- Experiment is valid; all mice in the PBS control group (n=16) succumb to infection at or before day 8 post challenge (median 6 days), whereas the positive control group (n=8, 15mg/kg CR6261, 1 day before challenge) is fully protected ($p < 0.001$).
- Three immunizations with polypeptides of the invention SEQ ID NO: 186, 203 and 254 adjuvanted with Matrix-M lead to significant increase in survival proportion ($p \leq 0.004$) (Fig 32A), increase in survival time ($p \leq 0.001$) (Fig. 32A) and reduction in clinical score ($p < 0.001$) (not shown), compared to the PBS control group. For polypeptides of the invention SEQ ID NO: 203 a significant reduction in body weight AUC is also observed ($p < 0.001$) (Fig. 32B).

- Pre-challenge IgG antibody titers to H1N1 A/Brisbane/59/07 FL HA induced by polypeptides of the invention SEQ ID NO: 186, 203 and 254 are significantly higher compared to PBS ($p \leq 0.001$) (Fig. 33A).
- Matrix-M adjuvanted polypeptides of the invention SEQ ID NO: 186, 203 and 254 induce significantly higher CR9114 competition titers compared to PBS ($p < 0.001$) (Fig. 33B).
- Matrix-M adjuvanted trimeric polypeptide of the invention SEQ ID NO: 203 (background H1N1 A/California/07/2009) induces significantly lower CR9114 competing antibodies compared to variants derived from H1N1 A/Brisbane/59/2007 ($p \leq 0.002$) when H1N1 A/Brisbane/59/07 FL HA is used as a target antigen (Fig 33B).

Conclusion: Matrix-M adjuvanted polypeptides of the invention SEQ ID NO: 186, 203 and 254 confer protection against lethal challenge with H1N1 A/NL/602/09.

Example 20: H1 mHA trimer candidate evaluation II in a H5N1 A/Hong Kong/156/97 mouse model

In order to provide further evidence for the use of polypeptides of the invention as vaccines the protective efficacy of three additional polypeptides against lethal challenge with Influenza was evaluated. To this end polypeptides of the invention 203 and 254 (both containing C-terminal his-tags) were transiently expressed in HEK293F cells and purified as described above. In addition polypeptide of the invention SEQ ID NO: 186 (also containing an additional C-terminal his-tag) was expressed in SF9 insect cells following procedures well known to those skilled in the art and purified as described above.

The aim of this study was to determine protective efficacy of polypeptides of the invention SEQ ID NO: 186, 203 and 254 adjuvanted with Matrix-M in a H5N1 A/Hong Kong/156/97 challenge model compared to a PBS control group.

Groups of 10 female BALB/c mice (age 6-8 weeks) were immunized 3 times at a 3 week interval with 30 μ g polypeptide of the invention adjuvanted with 10 μ g Matrix-M.

As a positive control for the challenge model CR6261 (15 mg/kg) was administered 1 day

prior to challenge (n=8), while injection with PBS served as a negative control (n=16). Four weeks after the last immunization mice were challenged with 12.5xLD50 challenge virus and monitored (survival, weight, clinical scores) for 3 weeks.

To verify immunogenicity of polypeptides of the invention SEQ ID NO: 186, 203 and 254, pre-challenge sera (day -1) are tested in ELISA assays for binding to FL HA from H1N1 A/Brisbane/59/07. To determine whether mini-HA induced antibodies bind at close proximity to the CR9114 epitope, a CR9114 competition ELISA was performed. Competition data were visualized as '% competition', defined as $(A-P)/A \times 100$, where A is the maximum OD signal of CR9114 binding to FL HA when no serum is present and P is the OD signal of CR9114 binding to FL HA in presence of serum at a given dilution or expressed using the slope OD metric to be able to quantify responses; for reference CR9114 and CR8020 (starting concentration 5 µg/ml) solutions were included.

Results:

- Survival percentages for the experimental groups are reported in Table 12.
- Experiment is valid; 15 out of 16 mice in the PBS control group succumb to infection at or before day 9 post challenge (median 9 days), whereas the positive control group (n=8, 15mg/kg CR6261, 1 day before challenge) is fully protected ($p < 0.001$).
- Three immunizations with polypeptides of the invention SEQ ID NO: 186, 203 and 254 adjuvanted with Matrix-M lead to significant increase in survival proportion ($p < 0.001$) (Fig 34A), increase in survival time ($p < 0.001$) (Fig. 34A), decrease in bodyweight loss ($p < 0.001$) (Fig 34B) and reduction in clinical score ($p < 0.001$) (not shown), compared to the PBS control group.
- Pre-challenge IgG antibody titers to H1N1 A/Brisbane/59/07 FL HA induced by polypeptides of the invention SEQ ID NO: 186, 203 and 254 are significantly higher compared to PBS ($p \leq 0.001$) (Fig. 35A).
- Matrix-M adjuvanted polypeptides of the invention SEQ ID NO: 186, 203 and 254 induce significantly higher CR9114 competition titers compared to PBS ($p < 0.001$) (Fig. 35B).

- Matrix-M adjuvanted polypeptides of the invention SEQ ID NO: 203 (background H1N1 A/California/07/2009) induces significantly lower CR9114 competing antibodies compared to polypeptides of the invention SEQ ID NO: 186 and 254 (background H1N1 A/Brisbane/59/2007) when H1N1 A/Brisbane/59/07 FL HA is used as a target antigen ($p < 0.001$) (Fig. 35B).

Conclusion: Matrix-M adjuvanted polypeptides of the invention SEQ ID NO: 186, 203 and 254 confer protection against lethal challenge with heterosubtypic H5N1 A/Hong Kong/156/97.

10 Example 21: H1 mHA trimer candidate evaluation II in a H1N1 A/Puerto Rico/8/34 mouse model

In order to provide further evidence for the use of polypeptides of the invention as vaccines the protective efficacy of three additional polypeptides against lethal challenge with Influenza was evaluated. To this end polypeptides of the invention 203 and 254 (both containing C-terminal his-tags) were transiently expressed in HEK293F cells and purified as described above. In addition polypeptide of the invention SEQ ID NO: 186 (also containing an additional C-terminal his-tag) was expressed in SF9 insect cells following procedures well known to those skilled in the art and purified as described above.

20 The aim of this study was to determine protective efficacy of polypeptides of the invention SEQ ID NO: 186, 203 and 254 adjuvanted with Matrix-M in a H1N1 A/Puerto Rico/8/1934 challenge model compared to a PBS control group.

Groups of 10 female BALB/c mice (age 6-8 weeks) were immunized 3 times at a 3 week interval with 30 μ g of polypeptides of the invention SEQ ID NO: 186, 203 and 254 adjuvanted with 10 μ g Matrix-M. As a positive control for the challenge model CR6261 (15 mg/kg) was administered 1 day prior to challenge ($n=8$), while 3 immunizations with PBS served as a negative control ($n=16$). Four weeks after the last immunization mice were challenged with 25xLD50 challenge virus and monitored (survival, weight, clinical scores) for 3 weeks.

30 To verify immunogenicity of polypeptides of the invention, pre-challenge sera (day -1) are tested in ELISA assay for binding to FL HA from H1N1 A/Brisbane/59/07.

To determine whether mini-HA induced antibodies bind at close proximity to the CR9114 epitope, a CR9114 competition ELISA was performed. Competition data were visualized as ‘% competition’, defined as $(A-P)/Ax100$, where A is the maximum OD signal of CR9114 binding to FL HA when no serum is present and P is the OD signal of CR9114 binding to FL HA in presence of serum at a given dilution or expressed using the slope OD metric to be able to quantify responses; for reference CR9114 and CR8020 (starting concentration 5 µg/ml) solutions were included.

Results

- Survival percentages for the experimental groups are reported in Table 12.
- Experiment is valid; all mice in the PBS control group (n=16) succumb to infection at or before day 9 post challenge (median 8 days), whereas the positive control group (n=8, 15mg/kg CR6261, 1 day before challenge) is fully protected ($p<0.001$).
- Three immunizations with polypeptides of the invention SEQ ID NO: 186, 203 and 254 adjuvanted with Matrix-M lead to significant increase in survival proportion ($p<0.001$) (Fig 36A), increase in survival time ($p<0.001$) (Fig. 36A), decrease in bodyweight loss ($p<0.001$) (Fig. 37B) and reduction in clinical score ($p<0.001$) (not shown), compared to the PBS control group.
- Pre-challenge IgG antibody titers to H1N1 A/Brisbane/59/07 FL HA induced by H1 mini-HA variants are significantly higher compared to PBS for polypeptides of the invention SEQ ID NO: 186, 203 and 254 ($p<0.001$) (Fig. 37A).
- Matrix-M adjuvanted polypeptides of the invention SEQ ID NO: 186, 203 and 254 induce significantly higher CR9114 competition titers compared to PBS ($p<0.001$) (Fig. 37B).
- Matrix-M adjuvanted polypeptide of the invention SEQ ID NO: 203 (background H1N1 A/California/07/2009) induces significantly lower CR9114 competing antibodies compared to variants based on H1N1 A/Brisbane/59/2007 when H1N1 A/Brisbane/59/07 FL HA is used as a target antigen ($p\leq 0.001$) (Fig. 37B).

Conclusion: Matrix-M adjuvanted polypeptides of the invention SEQ ID NO: 186, 203 and 254 confer protection against lethal challenge with H1N1 A/Puerto Rico/8/34.

Example 22: Polypeptides of the invention in different H1 sequence backgrounds

5 Next to s127H1-t2-cl18long (SEQ ID NO: 181) two variants, Tex_s127H1-t2-cl18long and NY_s127H1-t2-cl18long, were produced (SEQ ID NO: 208 and 210) that share the same design features, however, are based on the HA originating from strains H1N1 A/Texas/UR06-0526/07 (SEQ ID NO: 205) and A/New York/629/95 (SEQ ID NO: 206). In this experiment a C-terminal Factor X cleavage site and 6 histidine tag are
10 present in both proteins. For expression and purification a protocol similar to described in Example 2 (was used with the exception the procedure was started out with 0.6 l culture supernatant.

 Purified polypeptides were further analyzed in a sandwich ELISA (as described in Example 3) for the presence of multimeric forms of polypeptides that present two or
15 more epitopes of the broadly neutralizing antibody CR9114. In short, coated mAb CR9114 was used to capture the purified proteins which were subsequently incubated with biotinylated CR914 or CR6261 and binding was assessed by HRP-conjugated streptavidin. Production yields and Multimer sandwich ELISA results are shown in table 13.

20 Results shown in table 13 indicate that all variants result in multimeric protein with desired binding characteristics. Polypeptide Tex_s127H1-t2-cl18 (SEQ ID NO: 208) has a similar yield compared to s127H1-t2-cl18long whereas polypeptide NY_s127H1-t2-cl18 (SEQ ID NO: 210) has a ~4 fold lower yield compared to s127H1-t2-cl18long (SEQ ID NO: 181). All polypeptides tested were capable of binding to CR9114 and
25 CR6261 similarly to s127H1-t2-cl18long as determined by ELISA which indicates the presence of multimerization. Taken together, the results shown here demonstrate the successful generation of HA stem based polypeptides using H1 sequence of different phylogenetic origin.

Example 23: Polypeptides of the invention with additional c-terminal trimerization domain

Polypeptides of the invention can be expressed with a variety of c-termini. Besides different length (resulting from alternative truncations of the transmembrane domain of HA) also tags for detection and purification as well as functional domains can be added without affecting the antigen structure. The constructs below demonstrate the addition of a foldon trimerization domain (flanked by a Flag- and a His-tag) to short (deleting residue 520 to the C-terminus; numbering according to SEQ ID NO: 1) or long (deleting residue 520 to the C-terminus) versions of polypeptides of the invention derived from FL HA from H1 A/Brisbane/59/2007 (SEQ ID NO: 1) or H1 A/California/07/2009 (SEQ ID NO: 252).

The constructs were transiently expressed in HEK293F cells (as described in Example 2) and the polypeptides of the invention present in the filtered culture supernatant investigated. The levels of expression and trimerization were first assessed by SDS-PAGE and Western Blot (as described in Example 2, see Figure 38). Culture medium was further analyzed in a sandwich ELISA (as described in Example 3) for the presence of multimeric forms of polypeptides that present two or more epitopes of the broadly neutralizing antibody CR9114. In short, coated mAb CR9114 was used to capture the purified proteins which were subsequently incubated with biotinylated CR914 and binding was assessed by HRP-conjugated streptavidin (see Table 14). Lastly, homogeneous binding studies were performed to confirm the expression level of the polypeptides of the invention as well as to determine their binding strength to well characterized monoclonal antibodies (IgG) with known epitopes on the stem of HA. Hereto an AlphaLISA setup (Perkin Elmer) was established which relied on the c-terminal Flag- and His-tags as well as the human monoclonal IgGs CR9114 and CR6261. All reagents were diluted in buffer containing PBS, 0.05% Tween-20, and 0.5mg/ml BSA.

To determine the expression level using the AlphaLISA setup, the filtered cell culture supernatants containing the polypeptides of the invention (were diluted 40 times in the presence of an anti-His donor bead and an anti-Flag acceptor bead, both from Perkin Elmer and at 10 µg/mL in a final volume of 25 µL. The homogeneous mixture

was incubated for 1h at RT. If the excited donor bead (680 nm) and acceptor bead both bind the respective c-terminal tags and come in close proximity (~100 nm), an energy transfer (singlet oxygen) can be measured as a luminescence signal of the acceptor bead (615 nm). The signal intensity in this homogeneous assay format is directly proportional to the amount of protein in solution. Averages of AlphaLISA signal intensities for the expressed polypeptides of the invention are shown in Table 14.

The interaction between the polypeptides of the invention and the IgGs was detected after 1h incubation with either CR9114 or CR6261 (1 or 2 nM final concentration respectively, in 25 μ L) at RT with two beads, an anti-His donor bead recognizing HA (10 μ g/mL) and an anti-Fc acceptor bead (10 μ g/mL) recognizing the IgG used. After an additional hour of incubation the AlphaLISA signal of the acceptor bead was measured. The signal intensity in this homogeneous assay format is directly proportional to the binding strength (affinity/avidity) between both binding partners and is thereby a measure for the integrity and quality of the mini-HA epitope. Averages of AlphaLISA signal intensities for the binding of CR9114 and CR6261 are shown in Table 14.

The parent constructs SEQ ID NO: 164 (H1 A/Brisbane/59/2007 background) and SEQ ID NO: 211 (H1 A/California/07/2009 background) already contain a GCN4 trimerization domain embedded in the C-helix. The insertion of an additional foldon trimerization domain at the c-terminus of soluble versions of these polypeptides (also included in the invention) is possible and results in similar or better expression levels. The additional trimerization domain may further improve the polypeptide of the invention with respect to the level of trimerization as evident by the Western Blot and multimer ELISA results. The binding of broadly neutralizing IgGs CR9114 and CR6261 is equal or better compared to polypeptides of the invention with only one trimerization domain.

Table 1. Standard amino acids, abbreviations and properties

Amino Acid	3-Letter	1-Letter	Side chain polarity	Side chain charge (pH 7.4)
alanine	Ala	A	nonpolar	Neutral
arginine	Arg	R	polar	Positive
asparagine	Asn	N	polar	Neutral
aspartic acid	Asp	D	polar	Negative
cysteine	Cys	C	nonpolar	Neutral
glutamic acid	Glu	E	polar	Negative
glutamine	Gln	Q	polar	Neutral
glycine	Gly	G	nonpolar	Neutral
histidine	His	H	polar	positive (10%) neutral(90%)
isoleucine	Ile	I	nonpolar	Neutral
leucine	Leu	L	nonpolar	Neutral
lysine	Lys	K	polar	Positive
methionine	Met	M	nonpolar	Neutral
phenylalanine	Phe	F	nonpolar	Neutral
proline	Pro	P	nonpolar	Neutral
serine	Ser	S	polar	Neutral
threonine	Thr	T	polar	Neutral
tryptophan	Trp	W	nonpolar	Neutral
tyrosine	Tyr	Y	polar	Neutral
valine	Val	V	nonpolar	Neutral

Table 2. Sequence alignment of H1 sequences according to particular embodiments of the invention

5	1. A/Solomon Islands/6/2003 (H1N1)	(SEQ ID NO: 25)
	2. A/Brisbane/59/2007 (H1N1)	(SEQ ID NO: 1)
	3. A/New Caledonia/20/1999(H1N1)	(SEQ ID NO: 26)
	4. A/California/07/2009 (H1N1)	(SEQ ID NO: 27)
	5. A/swine/Hubei/S1/2009(H1N1)	(SEQ ID NO: 28)
10	6. A/swine/Haseluenne/IDT2617/2003(H1N1)	(SEQ ID NO: 29)
	7. A/NewYork/8/2006(H1N1)	(SEQ ID NO: 30)
	8. A/SolomonIslands/3/2006(H1N1)	(SEQ ID NO: 31)
	9. A/NewYork/146/2000(H1N1)	(SEQ ID NO: 32)
	10. A/NewYork/653/1996(H1N1)	(SEQ ID NO: 33)
15	11. A/Beijing/262/1995(H1N1)	(SEQ ID NO: 34)
	12. A/Texas/36/1991(H1N1)	(SEQ ID NO: 35)
	13. A/Singapore/6/1986(H1N1)	(SEQ ID NO: 36)
	14. A/Chile/1/1983(H1N1)	(SEQ ID NO: 37)
	15. A/Baylor/11515/1982(H1N1)	(SEQ ID NO: 38)
20	16. A/Brazil/11/1978(H1N1)	(SEQ ID NO: 39)
	17. A/USSR/90/1977(H1N1)	(SEQ ID NO: 40)
	18. A/NewJersey/8/1976(H1N1)	(SEQ ID NO: 41)
	19. A/Denver/1957(H1N1)	(SEQ ID NO: 42)
	20. A/Albany/4835/1948(H1N1)	(SEQ ID NO: 43)
25	21. A/FortMonmouth/1/1947(H1N1)	(SEQ ID NO: 44)
	22. A/Cameron/1946(H1N1)	(SEQ ID NO: 45)
	23. A/Weiss/1943(H1N1)	(SEQ ID NO: 46)
	24. A/Iowa/1943(H1N1)	(SEQ ID NO: 47)
	25. A/Bellamy/1942(H1N1)	(SEQ ID NO: 48)
30	26. A/PuertoRico/8/1934(H1N1)	(SEQ ID NO: 49)
	27. A/WSN/1933(H1N1)	(SEQ ID NO: 50)
	28. A/SouthCarolina/1/1918(H1N1)	(SEQ ID NO: 51)
35	1. MKVKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCL 60	
	2. MKVKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL ENSHNGKLCL 60	
	3. MKAKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCL 60	
	4. MKAILVLLY TFATANADTL CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDKHNGLCK 60	
	5. MEAKLFVLC AFTALKADTF CVGYHANYST HTVDTILEKN VTVTHSVNLL ENSHNGKLCS 60	
40	6. MEAKLFVLC AFTALKADTI CVGYHANNST DTVDTILEKN VTVTHSVNLL ENNHNGKLCS 60	
	7. MKVKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCL 60	
	8. MKVKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCL 60	
	9. MKAKLLVLLC AFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	10. MKAKLLVLLC AFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
45	11. MKAKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCL 60	
	12. MKAKLLVLLC AFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	13. MKAKLLVLLC AFTATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	14. MKAKLLVLLC ALSATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDNHNGKLCK 60	
	15. MKAKLLVLLC ALSATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
50	16. MKAKLLVLLC ALSATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	17. MKAKLLVLLC ALSATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	18. MKAKLLVLLC AFTATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	19. MKAKLLVLLC ALSATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	20. MKAKLLVLLC ALSATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
55	21. MKAKLLVLLC ALTATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	22. MKAKLLVLLC ALSATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	23. MKARLLVLLC ALAATDADTI CIGYHANNST DTVDTILEKN VTVTHSVNLL EDSHNGKLCR 60	
	24. MKARLLVLLC ALAATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL EDSHNGKLCR 60	
	25. MKARLLVLLC AIAATDADTI CIGYHANNST DTVDTILEKN VTVTHSVNLL EDSHNGKLCR 60	

5	26.	MKANLLVLLC	ALAAADADTI	CIGYHANNST	DTVDTVLEKN	VTVTHSVNLL	EDSHNGKLCR	60
	27.	MKAKLLVLLY	AFVATDADTI	CIGYHANNST	DTVDTIFEKN	VAVTHSVNLL	EDRHNKGLCK	60
	28.	MEARLLVLLC	AFAATNADTI	CIGYHANNST	DTVDTVLEKN	VTVTHSVNLL	EDSHNGKLCR	60
		*:. *::**	:: :: ***:	*****	*****::**	*:*****	*: *****	
10	1.	LKGIAPQLG	NCSVAGWILG	NPECELLISR	ESWSYIVEKP	NPENGTCYPG	HFADYEELRE	120
	2.	LKGIAPQLG	NCSVAGWILG	NPECELLISK	ESWSYIVEKP	NPENGTCYPG	HFADYEELRE	120
	3.	LKGIAPQLG	NCSVAGWILG	NPECELLISK	ESWSYIVETP	NPENGTCYPG	YFADYEELRE	120
	4.	LRGVAPLHLG	KCNIAGWILG	NPECESLSTA	SSWSYIVETP	SSDNGTCYPG	DFIDYEELRE	120
	5.	LNGKIPLQLG	NCNVAGWILG	NPKCDLLTA	NSSSYIIETS	KSKNGACYPG	EFADYEELKE	120
	6.	LNGKAPQLG	NCNVAGWILG	NPECDLLTV	DSWSYIIETS	NSKNGACYPG	EFADYEELRE	120
	7.	LKGIAPQLG	NCSVAGWILG	NPECELLISK	ESWSYIVETP	NPENGTCYPG	YFADYEELRE	120
	8.	LKGIAPQLG	NCSVAGWILG	NPECELLISR	ESWSYIVEKP	NPENGTCYPG	HFADYEELRE	120
15	9.	LKGTAPQLG	NCSIAGWILG	NPECESLFSK	ESWSYIAETP	NPKNGTCYPG	YFADYEELRE	120
	10.	LKGTAPQLG	NCSVAGWILG	NPECESLFSK	ESWSYIAETP	NPENGTCYPG	YFADYEELRE	120
	11.	LKGIAPQLG	NCSVAGWILG	NPECESLISK	ESWSYIVETP	NPENGTCYPG	YFADYEELRE	120
	12.	LKGIAPQLG	NCSVAGWILG	NPKCESLFSK	ESWSYIAETP	NPENGTCYPG	YFADYEELRE	120
20	13.	LKGIAPQLG	NCSIAGWILG	NPECESLFSK	KWSYIAETP	NSENGTCYPG	YFADYEELRE	120
	14.	LKGIAPQLG	KCSIAGWILG	NPECESLFSK	KWSYIAETP	NSENGTCYPG	YFADYEELRE	120
	15.	LKGIAPQLG	KCSIAGWILG	NPECESLFSK	KWSYIAETP	NSENGTCYPG	YFADYEELRE	120
	16.	LKGIAPQLG	KCSIAGWILG	NPECESLFSK	KWSYIAETP	NSENGTCYPG	YFADYEELRE	120
25	17.	LKGIAPQLG	KCNIAGWILG	NPECESLFSK	KWSYIAETP	NSENGTCYPG	YFADYEELRE	120
	18.	LKGIAPQLG	NCSIAGWILG	NPECESLFSK	KWSYIAETP	NSENGTCYPG	YFADYEELRE	120
	19.	LKKGAPQLG	NCNIAGWVLG	NPECESLLSN	RSWSYIAETP	NSENGTCYPG	DFADYEELRE	120
	20.	LKGIAPQLG	KCNIAGWILG	NPECESLFSK	KWSYIAETP	NSENGTCYPG	YFADYEELRE	120
	21.	LKGIAPQLG	KCNIAGWILG	NPECESLLSK	RSWSYIAETP	NSENGACYPG	DFADYEELRE	120
	22.	LKGIAPQLG	KCNIAGWILG	NPECESLLSE	RSWSYIVEIP	NSENGTCYPG	DFADYEELRE	120
	23.	LKGIAPQLG	KCNIAGWILG	NPECESLLSE	RSWSYIVETP	NSENGTCYPG	DFADYEELRE	120
	24.	LKGIAPQLG	KCNIAGWILG	NPECESLLSE	RSWSYIVETP	NSENGTCYPG	DFADYEELRE	120
30	25.	LKGIAPQLG	KCNIAGWILG	NPECESLLSE	RSWSYIVETP	NSENGTCYPG	DFADYEELRE	120
	26.	LKGIAPQLG	KCNIAGWLLG	NPECDPLLPV	RSWSYIVETP	NSENGICYPG	DFADYEELRE	120
	27.	LKGIAPQLG	KCNITGWLLG	NPECDLLPA	RSWSYIVETP	NSENGACYPG	DFADYEELRE	120
	28.	LKGIAPQLG	KCNIAGWLLG	NPECDLLTA	SSWSYIVETS	NSENGTCYPG	DFADYEELRE	120
35		*:* ***:**	:*::~**:**	**:*: *	*****.*	...** **	* *****	
40	1.	QLSSVSSFER	FEIFPKESSW	PNHTTT-GVS	ASCSHNGESS	FYKNLLWLTG	KNGLYPNLSK	179
	2.	QLSSVSSFER	FEIFPKESSW	PNHTVT-GVS	ASCSHNGESS	FYRNLLWLTG	KNGLYPNLSK	179
	3.	QLSSVSSFER	FEIFPKESSW	PNHTVT-GVS	ASCSHNGKSS	FYRNLLWLTG	KNGLYPNLSK	179
	4.	QLSSVSSFER	FEIFPKTSSW	PNHDSNKGVT	AACPHAGAKS	FYKNLIWLVK	KNGSYPKLSK	180
	5.	QLSTVSSFER	FEIFPKAISW	PDHDATRGTT	VACSHSGVNS	FYRNLLSTVK	KNGSYPKLSK	180
	6.	QLSTVSSFER	FEIFPKATSW	PNHDTTRGTT	ISCSHSGANS	FYRNLLWIVK	KNGSYPKLSK	180
	7.	QLSSVSSFER	FEIFPKESSW	PNHTVT-GVS	ASCSHNGKSS	FYRNLLWLTG	KNGLYPNLSK	179
	8.	QLSSVSSFER	FEIFPKESSW	PNHTTT-GVS	ASCSHNGESS	FYKNLLWLTG	KNGLYPNLSK	179
45	9.	QLSSVSSFER	FEIFPKDSSW	PNHTVTKGVT	ASCSHNGKSS	FYKNLLWLTE	KNGLYPNLSK	180
	10.	QLSSVSSFER	FEIFPKESSW	PNHTVTKGVT	ASCSHNGKSS	FYKNLLWLTE	KNGLYPNLSK	180
	11.	QLSSVSSFER	FEIFPKESSW	PNHTVT-GVT	ASCSHNGKSS	FYRNLLWLTE	KNGLYPNLSN	179
	12.	QLSSVSSFER	FEIFPKESSW	PNHTVTKGVT	TSCSHNGKSS	FYRNLLWLTK	KNGLYPNVSK	180
50	13.	QLSSVSSFER	FEIFPKESSW	PNHTVTKGVT	ASCSHKGRSS	FYRNLLWLTK	KNGSYPNLSK	180
	14.	QLSSVSSFER	FEIFPKESSW	PKHNVTKGVT	AACSHKGKSS	FYRNLLWLTE	KNGSYPNLSK	180
	15.	QLSSVSSFER	FEIFPKESSW	PKHSVTRGVT	ASCSHKGKSS	FYRNLLWLTE	KNGSYPNLSK	180
	16.	QLSSVSSFER	FEIFPKERSW	PKHNITRGVT	ASCSHKGKSS	FYRNLLWLTE	KNGSYPNLSK	180
55	17.	QLSSVSSFER	FEIFPKERSW	PKHNVTRGVT	ASCSHKGKSS	FYRNLLWLTE	KNGSYPNLSK	180
	18.	QLSSVSSFER	FEIFPKESSW	PNHTVTKGVT	ASCSHKGRSS	FYRNLLWLTK	KNGSYPNLSK	180
	19.	QLSSVSSFER	FEIFPKERSW	PNHTTR-GVT	AACPHARKSS	FYKNLVWLTE	ANGSYPNLSR	179
	20.	QLSSVSSFER	FEIFPKERSW	PKHNITRGVT	AACSHKGKSS	FYRNLLWLTE	KNGSYPNLNK	180
60	21.	QLSSVSSFER	FEIFPKERSW	PKHNITRGVT	AACSHAGKSS	FYKNLLWLTE	TDGSYPKLSK	180
	22.	QLSSVSSFER	FEIFPKGRSW	PEHNIDIGVT	AACSHAGKSS	FYKNLLWLTE	KDGSYPNLNK	180
	23.	QLSSVSSFER	FEIFPKESSW	PKHNTARGVT	AACSHAGKSS	FYRNLLWLTE	KDGSYPNLKN	180
	24.	QLSSVSSFER	FEIFSKESSW	PKHTTG-GVT	AACSHAGKSS	FYRNLLWLTE	KDGSYPNLNN	180
	25.	QLSSVTSFER	FEIFPKETSW	PKHNTTKGVT	AACSHAGKCS	FYRNLLWLTE	KDGSYPNLNN	179
	26.	QLSSVSSFER	FEIFPKESSW	PNHNTN-GVT	AACSHAGKSS	FYRNLLWLTE	KEGSYPKLKN	179
	27.	QLSSVSSFER	FEIFPKESSW	PNHNTN-GVT	AACSHAGKSS	FYRNLLWLTE	KEGSYPKLKN	179
	28.	QLSSVSSFER	FEIFPKESSW	PNHNTN-GVT	AACSHAGKSS	FYRNLLWLTE	KEGSYPKLKN	179

27. QLSSVSSLER FEIFPKESSW PNHTFN-GVT VSCSHRGKSS FYRNLLWLTK KGDSYPKLTN 179
 28. QLSSVSSF EK FEIFPKTSSW PNHETTKGVT AACSYAGASS FYRNLLWLTK KGSSYPKL SK 180
 *****:*.**: *****.* ** *:.* **: .:*.**: * **:*.**:*. . **::..

5

1. SYANNKEKEV LVLWGVHHP NIGDQRALYH KENAYVSVVS SHYSRKFTPE IAKRPKVRDQ 239
 2. SYANNKEKEV LVLWGVHHP NIGNQKALYH TENAYVSVVS SHYSRKFTPE IAKRPKVRDQ 239
 3. SYVNNKEKEV LVLWGVHHP NIGNQRALYH TENAYVSVVS SHYSRRFTPE IAKRPKVRDQ 239
 4. SYINDKGKEV LVLWGIHHP TSADQQSLYQ NADAYVSVGS SRYSKKFKPE IAKRPKVRXX 240
 5. SYTNNKGKEV LVIWGVHHP TDSVQQTLYQ NKHTYVSVGS SKYYKRFTPE IVARPKVRGQ 240
 6. SYTNNKGKEV LVIWGVHHP TDSVQQTLYQ NNHTYVSVGS SKYYQRFTPE IVTRPKVRGQ 240
 7. SYANNKEKEV LVLWGVHHP NIGDQRALYH TENAYVSVVS SHYSRRFTPE IAKRPKVRDQ 239
 8. SYANNKEKEV LVLWGVHHP NIGDQRALYH KENAYVSVVS SHYSRKFTPE IAKRPKVRDQ 239
 9. SYVNNKGKEV LVLWGVHHP NMGDQRAIYH KENAYVSVLS SHYSRRFTPE IAKRPKVRDQ 240
 10. SYVNNKEKEV LVLWGVHHP NIGDQRAIYH TENAYVSVVS SHYSRRFTPE IAKRPKVRDQ 240
 11. SYVNNKEKEV LVLWGVHHP NIRDQRAIYH TENAYVSVVS SHYSRRFTPE IAKRPKVRGQ 239
 12. SYVNNKEKEV LVLWGVHHP NIGDQRAIYH TENAYVSVVS SHYSRRFTPE IAKRPKVRDQ 240
 13. SYVNNKEKEV LVLWGVHHP NIGDQRAIYH TENAYVSVVS SHYNRRFTPE IAKRPKVRDQ 240
 14. SYVNNKEKEV LVLWGVHHP NIEDQKTIYR KENAYVSVVS SHYNRRFTPE IAKRPKVRN 240
 15. SYVNDKEKEV LVLWGVHHP NIEDQKTIYR KENAYVSVVS SHYNRRFTPE IAKRPKVRDQ 240
 16. SYVNNKEKEV LVLWGVHHP NIEDQKTIYR KENAYVSVVS SNYNRRFTPE IAKRPKVRGQ 240
 17. SYVNNKEKEV LVLWGVHHP NIEDQKTIYR KENAYVSVVS SNYNRRFTPE IAKRPKVRGQ 240
 18. SYVNNKEKEV LVLWGVHHP NIGDQRAIYH TENAYVSVVS SHYNRRFTPE IAKRPKVRDQ 240
 19. SYVNNKEKEV LVLWGVHHP NIEEQRALYR KDNAYVSVVS SNYNRRFTPE IAKRPKVRDQ 239
 20. SYVNNKEKEV LVLWGVHHP NIEDQKTIYR KENAYVSVVS SNYNRRFTPE IAKRPKVRGQ 240
 21. SYVNNKEKEV LVLWGVHHP NIEDQKTIYR KENAYVSVVS SNYNRRFTPE IAKRPKVRDQ 240
 22. SYVNNKEKEV LILWGVHHP NIENQKTIYR KENAYVSVVS SNYNRRFTPE IAKRPKVRGQ 240
 23. SYVNNKGKEV LVLWGVHHP SIKEQQTLYQ KENAYVSVVS SNYNRRFTPE IAKRPKVRDQ 240
 24. SYVNNKGKEV LVLWGVHHP NIKDQQTLYQ KENAYVSVVS SNYNRRFTPE IAKRPKVRGQ 239
 25. SYVNNKGKEV LVLWGVHHP NIKDQQTLYQ KENAYVSVVS SNYNRRFTPE IAKRPKVRGQ 240
 26. SYVNNKGKEV LVLWGIHHP NSKEQONLY NENAYVSVVT SNYNRRFTPE IAKRPKVRDQ 239
 27. SYVNNKGKEV LVLWGVHHP SSDEQQSLYS NGNAYVSVAS SNYNRRFTPE IAKRPKVRDQ 239
 28. SYVNNKGKEV LVLWGVHHP TGTDDQSLYQ NADAYVSVGS SKYNRRFTPE IAKRPKVRDQ 240
 ** *:.*: *** *:***:***. . **:.* .:*** *:.*:*.** *:***:

35

1. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIINSNAPMD ECDACQTPQ 299
 2. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIINSNAPMD KCDACQTPQ 299
 3. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNAPMD ECDACQTPQ 299
 4. EGRMYYWTL VEPGDKITFE ATGNLVVPRY AFAMERNAGS GIIISDTPVH DCNTTCQTPK 300
 5. AGRMYYWTL FDQGDITIFE ATGNLIAPWH AFALKKGSSS GIMLSDAQVH NCTTKCQTPH 300
 6. AGRMYYWTL LDQGDITIFE ATGNLIAPWH AFALNKGPS GIMISDAHVH NCTTKCQTPH 300
 7. EGRINYYWTL LEPGDTIIFE ANGNIAPRF AFALSRGFGS GIITSNAPMD ECDACQTPQ 299
 8. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIINSNAPMD ECDACQTPQ 299
 9. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIIISNASMG ECDACQTPQ 300
 10. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNASMG ECDACQTPQ 300
 11. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNAPMN ECDACQTPQ 299
 12. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNASMD ECDACQTPQ 300
 13. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNASMD ECDACQTPQ 300
 14. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNASMD ECDACQTPQ 300
 15. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNVSM ECDACQTPQ 300
 16. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNASMD ECDTKCQTPQ 300
 17. AGRINYYWTL LEPGDTIIFE ANGNIAPWH AFALNRGFGS GIITSNASMD ECDTKCQTPQ 300
 18. EGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNASMD ECDACQTPQ 300
 19. SGRMYYWTL LEPGDTIIFE ATGNLIAPRY AFALSRGFGS GIITSNAPLD ECDTKCQTPQ 299
 20. AGRINYYWTL LEPGDTIIFE ANGNIAPWH AFALSRGFGS GIITSNASMD ECDTKCQTPQ 300
 21. AGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALSRDFGS GIITSNASMD ECDTKCQTPQ 300
 22. AGRINYYWTL LEPGDTIIFE ANGNIAPRY AFALNRGIGS GIITSNASMD ECDTKCQTPQ 300
 23. AGRMYYWTL LEPGDTIIFE ANGNIAPRY AFALSRGFGS GIITSNASMH ECDTKCQTPQ 300
 24. AGRINYYWTL LKPGDTIMFE ANGNIAPRY AFALSRGFGS GIITSNASMH ECDTKCQTPQ 299

60

5	25.	AGRMNYYWTL	LEPGDTIIFE	ANGNLIAPWY	AFALSRGFGS	GIITSNASMH	ECNTKCQTPQ	300
	26.	AGRMNYYWTL	LKPGDTIIFE	ANGNLIAPMY	AFALRRGFGS	GIITSNASMH	ECNTKCQTPL	299
	27.	HGRMNYWTL	LEPGDTIIFE	ATGNLIAPWY	AFALSRGFES	GIITSNASMH	ECNTKCQTPQ	299
	28.	AGRMNYYWTL	LEPGDTITFE	ATGNLIAPWY	AFALNRGSGS	GIITS DAPVH	DCNTKCQTPH	300
		::*** ::***.* ** *.***:.* . ***: *. * *** *:..: .*:..****						
10	1.	GAINSSLPFQ	NVHPVTIGEC	PKYVRS AKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	359
	2.	GAINSSLPFQ	NVHPVTIGEC	PKYVRS AKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	359
	3.	GAINSSLPFQ	NVHPVTIGEC	PKYVRS AKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	359
	4.	GAINTSLPFQ	NIHPITIGKC	PKYVKSTKLR	LATGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
15	5.	GALKNNLPLQ	NVHLFTIGEC	PKYVKSTQLR	MATGLRNIPS	IQSRGLFGAI	AGFIEGGRTG	360
	6.	GALKSNLPLQ	NVHPSTIGEC	PKYVKSTQLR	MATGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	7.	GAINSSLPFQ	NVHPVTIGEC	PKYVRS AKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	359
	8.	GAINSSLPFQ	NVHPVTIGEC	PKYVRS AKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	359
20	9.	GAINSSLPFQ	NVHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	10.	GAINSSLPFQ	NVHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	11.	GAINSSLPFQ	NVHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	359
	12.	GAINSSLPFQ	NVHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
25	13.	GAINSSLPFQ	NVHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	14.	GAINSSLPFQ	NIHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	15.	GAINSSLPFQ	NVHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	16.	GAINSSLPFQ	NVHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
30	17.	GAINSSLPFQ	NIHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	18.	GAINSSLPFQ	NVHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	19.	GAINSSLPFQ	NIHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	VQSRGLFGAI	AGFIEGGWTG	359
	20.	GAINSSLPFQ	NIHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
35	21.	GAINSSLPFQ	NIHPVTIGEC	PKYVKSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	22.	GAINSSLPFQ	NIHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWDG	360
	23.	GAINSSLPFQ	NIHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	24.	GAINSSLPFQ	NIHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	359
40	25.	GAINSSLPFQ	NIHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
	26.	GAINSSLPYQ	NIHPVTIGEC	PKYVRS AKLR	MVTGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	359
	27.	GSINSNLPFQ	NIHPVTIGEC	PKYVRSTKLR	MVTGLRNIPS	IQYRGLFGAI	AGFIEGGWTG	359
	28.	GAINSSLPFQ	NIHPVTIGEC	PKYVRSTKLR	MATGLRNIPS	IQSRGLFGAI	AGFIEGGWTG	360
		*:***:.**: **:..***: **:***:.**: **:..***:.**: **:..***:.**: **:..***:.**: *						
45	1.	MVDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	419
	2.	MVDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	419
	3.	MVDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	419
	4.	MVDGWYGYHH	QNEQSGGYAA	DLKSTQNAID	EITNKVNSVI	EKMNTQFTAV	GKEFNHLEKR	420
50	5.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQIAID	GINKNKANSVI	GKMNIQLTSV	GKEFNSLEKR	420
	6.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQIAID	GINKNKANSII	EKMNTQFTSV	GKEFNLEKR	420
	7.	MVDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	419
	8.	MVDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	419
55	9.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSII	EKMNTQFTAV	GKEFNKLEKR	420
	10.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAID	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	420
	11.	MMDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	419
	12.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	420
60	13.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	420
	14.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSII	EKMNTQFTAV	GKEFNKLEKR	420
	15.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	420
	16.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	420
65	17.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	420
	18.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLERR	420
	19.	MMDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	419
	20.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	420
70	21.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	WITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	420
	22.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	420
	23.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	420
	24.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	419
75	25.	MIDGWYGYHH	QNEQSGGYAA	DQKSTQNAIN	GITNKVNSVI	EKMNTQFTAV	GKEFNKLEKR	420

26. MIDGWYGYHH QNEQGSYGAA DQKSTQNAIN GITNKNVSVI EKMNIQFTAV GKEFNKLEKR 419
 27. MIDGWYGYHH QNEQGSYGAA DQKSTQNAIN GITNKNVSVI EKMNTQFTAV GKEFNNLEKR 419
 28. MIDGWYGYHH QNEQGSYGAA DQKSTQNAID GITNKNVSVI EKMNTQFTAV GKEFNNLERR 420
 *:***** ***** * *****: *****:* ***** *****:***:*

5

1. MENLNKKVDD GFIDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 479
 2. MENLNKKVDD GFIDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 479
 3. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 479
 4. IENLNKKVDD GFLDIWTYNA ELLVLENER TLDYHDSNVK NLYEKVRSQK KNNAKEIGNG 480
 5. KENLNKTVDD RFLDVWTFNA ELLVLENER TLEFHDNLNK SLYEKVKSQK RNNDEKEIGNG 480
 6. IENLNKKVDD GFLDVWTFNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK RNNAKEIGNG 480
 7. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 479
 8. MENLNKKVDD GFIDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 479
 9. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 10. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 11. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 479
 12. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 13. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 14. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 15. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 16. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 17. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 18. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 19. MENLNKKVDD GFMDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK RNNAKELGNG 479
 20. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 21. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK RNNAKEIGNG 480
 22. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK RNNAKEIGNG 480
 23. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK RNNAKEIGNG 480
 24. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK RNNAKEIGNG 479
 25. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK RNNAKEIGNG 480
 26. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 479
 27. MENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 479
 28. IENLNKKVDD GFLDIWTYNA ELLVLENER TLDHFHDSNVK NLYEKVKSQK KNNAKEIGNG 480
 :****:**** **:***** ***:**** * ***:*** **: *****:.* ** :*****:***

35

1. CFEFYHKCND ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 539
 2. CFEFYHKCND ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 539
 3. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 539
 4. CFEFYHKCDN TCMESVKNGT YDYPKYSEEA KLNREEIDGV KLESTRIYQI LAIYSTVASS 540
 5. CFEFYHKRDN ECLECVKNGT YNYPKYSEES KFNREEIVGV KLESMGIHQI LAIYSTVASS 540
 6. CFEFYHKCDN ECMESVKNGT YNYPKYSEES KLNREKIDGV KLESMGVHQI LAIYSTVASS 540
 7. CFEFYHKCND ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 539
 8. CFEFYHKCND ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 539
 9. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 10. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 11. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 539
 12. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 13. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 14. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 15. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 16. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 17. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 18. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 19. CFEFYHKCDN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYRI LAIYSTVASS 539
 20. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 21. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 22. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 23. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540
 24. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 539
 25. CFEFYHKCNN ECMESVKNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 540

60

26. CFEFYHKCDN ECMESVRNGT YDYPKYSEES KLNREKVDGV KLESMGIYQI LAIYSTVASS 539
27. CFEFYHKCDN ECMESVRNGT YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS 539
28. CFEFYHKCDD ACMESVRNGT YDYPKYSEES KLNREEIDGV KLESMGVYQI LAIYSTVASS 540
*****: : *****:*** *****: : ***** .:*** ***** : : *****.***

5

1. LVLLVSLGAI SFWMCSNGSL QCRICI 565
2. LVLLVSLGAI SFWMCSNGSL QCRICI 565
3. LVLLVSLGAI SFWMCSNGSL QCRICI 565
4. LVLLVSLGAI SFWMCSNGSL QCRICI 566
5. LVLLVSLGAI SFWMCSNGSL QCRICI 566
6. LVLLVSLGAI SFWMCSNGSL QCRICI 566
7. LVLLVSLGAI SFWMCSNGSL QCRICI 565
8. LVLLVSLGAI SFWMCSNGSL QCRICI 565
9. LVLLVSLGAI SFWMCSNGSL QCRICI 566
10. LVLLVSLGAI SFWMCSNGSL QCRICI 566
11. LVLLVSLGAI SFWMCSNGSL QCRICI 565
12. LVLLVSLGAI SFWMCSNGSL QCRICI 566
13. LVLLVSLGAI SFWMCSNGSL QCRICI 566
14. LVLLVSLGAI SFWMCSNGSL QCRICI 566
15. LVLLVSLGAI SFWMCSNGSL QCRICI 566
16. LVLLVSLGAI SFWMCSNGSL QCRICI 566
17. LVLLVSLGAI SFWMCSNGSL QCRICI 566
18. LVLLVSLGAI SFWMCSNGSL QCRICI 566
19. LVLLVSLGAI SFWMCSNGSL QCRICI 565
20. LVLLVSLGAI SFWMCSNGSL QCRICI 566
21. LVLLVSLGAI SFWMCSNGSL QCRICI 566
22. LVLLVSLGAI SFWMCSNGSL QCRICI 566
23. LVLLVSLGAI SFWMCSNGSL QCRICI 566
24. LVLLVSLGAI SFWMCSNGSL QCRICI 565
25. LVLLVSLGAI SFWMCSNGSL QCRICI 566
26. LVLLVSLGAI SFWMCSNGSL QCRICI 565
27. LVLLVSLGAI SFWMCSNGSL QCRICI 565
28. LVLLVSLGAI SFWMCSNGSL QCRICI 566
:** *****

35

Table 3. Polypeptides expressed in *P. pastoris*.

Expression and CR6261 binding were determined as described and the ratio of binding and expression signals calculated.

SET1					Fusion peptide area				B-loop				
clone	CR6261 binding signal	HTRF signal	ratio	fold increase of ratio over parental H1 mini-HA	337	340	352	353	402	406	409	413	416
					E, I, K, V	I, K, R, T	D, F, V, Y	I, K, R, T	E, K, M, V	F, I, N, S, T, Y	A, G, I, R, T, V	F, I, N, S, T, Y	H, I, L, N, R, S
239E11	1076944	1492	721,81	121,52	K	I	Y	T	M	F	I	N	R
127H1	800024	6572	121,73	20,49	K	K	F	T	M	Y	I	Y	S
171E5	879704	11508	76,44	12,87	K	T	F	T	M	I	A	F	S
239D2	570424	9279	61,47	10,35	K	K	F	T	M	I	V	F	N
247B2	414984	7583	54,73	9,21	K	I	Y	T	V	Y	I	F	S
253D4	395824	7546	52,45	8,83	K	T	F	T	M	Y	A	Y	H
252F5	421824	8621	48,93	8,24	V	K	Y	T	M	Y	V	Y	N
220C9	1086064	22606	48,04	8,09	K	T	F	T	M	F	T	Y	L
125D3	139824	2937	47,61	8,02	K	K	F	T	M	Y	G	T	H
137C11	416504	9167	45,44	7,65	V	K	F	T	M	Y	I	N	H
131B5	844344	20419	41,35	6,96	K	T	F	T	M	I	V	Y	H
233F11	583024	14389	40,52	6,82	K	K	Y	T	M	T	I	G	S
234C5	377864	9465	39,92	6,72	I	I	Y	T	M	F	T	N	L
115A1	1176904	30389	38,73	6,52	K	K	Y	T	M	I	V	Y	I
185G7	505864	13560	37,31	6,28	K	K	Y	T	M	I	V	I	S
275D4	327344	9030	36,25	6,10	K	K	Y	T	M	T	T	S	S
244B8	273744	7757	35,29	5,94	I	T	Y	T	M	Y	A	I	S
252B8	284984	8252	34,54	5,81	K	I	Y	T	M	S	I	N	L
213C11	667024	20624	32,34	5,44	V	K	Y	T	M	I	V	F	H
174G3	491184	15320	32,06	5,40	K	T	Y	K	V	S	G	Y	L
125D10	133904	4241	31,57	5,31	K	I	Y	T	M	Y	V	N	R
127A7	233064	7498	31,08	5,23	E	T	Y	T	M	I	I	I	L
304G11	110504	3588	30,8	5,19	K	K	Y	K	M	F	T	F	S
162A11	364024	11939	30,49	5,13	V	K	Y	T	M	F	A	F	I
271F10	315304	10348	30,47	5,13	I	K	Y	T	M	I	A	I	L
218G11	958504	33710	28,43	4,79	I	T	Y	I	M	I	I	I	N
251C8	269544	9634	27,98	4,71	K	T	Y	K	M	Y	I	N	L
258A6	165624	6004	27,59	4,64	I	T	Y	T	M	Y	T	F	H
134A4	456304	17366	26,28	4,42	K	I	Y	I	M	I	A	Y	N
214C11	317904	12120	26,23	4,42	E	I	Y	T	M	Y	V	S	S
182G8	399864	15262	26,2	4,41	K	K	Y	T	M	T	V	I	I
113E7	966064	38018	25,41	4,28	K	K	F	T	M	Y	T	I	H
230G9	854584	34093	25,07	4,22	K	K	Y	T	M	Y	T	F	R
222G4	419064	16996	24,66	4,15	K	T	F	I	V	I	I	Y	L
182D7	418944	17096	24,51	4,13	I	T	Y	T	M	I	I	F	N
272H2	263264	10844	24,28	4,09	K	T	Y	T	M	S	A	N	H
191C8	309064	12753	24,23	4,08	I	T	Y	T	V	I	A	F	I
123C10	237824	9843	24,16	4,07	K	I	Y	K	M	F	A	T	L
284B9	1663504	70812	23,49	3,95	K	T	Y	R	M	I	R	T	L
134A3	531784	23414	22,71	3,82	K	K	F	I	M	I	I	N	S
188F4	287384	12888	22,3	3,75	K	K	Y	T	M	S	V	T	H
189B7	336344	15207	22,12	3,72	E	T	F	T	M	Y	V	F	N
148D5	329144	14994	21,95	3,70	E	T	Y	I	M	F	G	S	H
194C8	242304	11113	21,8	3,67	I	T	F	T	M	F	V	F	I
188A8	279144	13001	21,47	3,61	K	T	Y	K	M	F	V	S	I
162B3	279584	13159	21,25	3,58	V	T	Y	T	M	Y	T	N	N
204C5	832784	39330	21,17	3,56	V	K	F	T	V	I	I	Y	L
216E5	334904	15873	21,1	3,55	V	T	F	T	M	F	R	Y	R
129C2	199464	9486	21,03	3,54	V	R	Y	I	M	I	I	Y	S
286E8	158704	7662	20,71	3,49	E	I	F	T	M	F	I	Y	S
264G4	180504	8751	20,63	3,47	K	R	Y	T	V	I	V	F	S
214C4	302264	14709	20,55	3,46	I	I	F	T	V	F	A	S	S
125A8	212224	10327	20,55	3,46	K	I	F	T	V	I	V	Y	I

Table 4 Polypeptides expressed in *P. pastoris*.

Expression and CR6261 binding were determined as described and the ratio of binding and expression signals calculated.

5

Set 2					Fusion peptide area				B-loop				
clone	CR6261 binding signal	HTRF signal	ratio	fold increase of ratio over parental SEQ ID NO: 6	337	340	352	353	402	406	409	413	416
					A, E, I, K, T, V	F, I, N, S, T, Y	A, D, F, I, N, S, T, V, Y	E, G, I, K, R, V	M, R, T	F, H, L, Y	F, I, S, T	E, K, M, V	I, L, R, S
86B4	1077144	13862	77,7	13,08	K	N	Y	K	M	F	I	M	I
7A7	987824	13452	73,43	12,36	T	N	Y	V	M	Y	F	E	R
55G7	616184	8767	70,28	11,83	K	N	Y	V	M	Y	I	M	L
71H2	1109984	16750	66,27	11,16	K	N	F	K	M	L	I	V	S
86B3	900904	14448	62,35	10,50	K	N	Y	K	M	L	I	V	R
71A4	1064144	17597	60,47	10,18	T	N	Y	V	M	Y	F	E	R
51G3	460304	7773	59,22	9,97	T	I	F	V	M	L	F	E	S
84B8	582144	10091	57,69	9,71	K	N	Y	I	M	F	F	M	S
79C2	364184	7116	51,18	8,62	T	N	Y	R	M	F	T	V	S
69G8	481344	9479	50,78	8,55	I	N	F	R	M	L	I	V	L
79D5	702584	13981	50,25	8,46	A	N	F	K	M	L	F	V	L
54H4	291744	5857	49,81	8,39	K	I	Y	K	M	L	I	E	L
11H6	427384	9146	46,73	7,87	K	N	Y	E	M	F	T	E	S
90A9	413664	9025	45,84	7,72	K	S	Y	V	M	Y	T	V	S
75G5	1011384	26695	37,89	6,38	E	S	Y	V	M	L	F	E	R
8A10	360104	9630	37,39	6,29	K	N	Y	V	M	L	I	V	R
72D4	329944	8881	37,15	6,25	V	N	F	R	M	F	S	M	S
74H9	1283144	35494	36,15	6,09	K	N	F	K	M	Y	F	M	S
88C5	471424	13355	35,3	5,94	K	N	Y	R	M	L	I	V	R
61A9	383064	10864	35,26	5,94	T	N	F	R	M	F	F	E	L
86H9	457344	13340	34,28	5,77	K	N	F	G	M	F	T	V	S
71D3	1573024	46711	33,68	5,67	I	S	Y	V	M	F	I	V	L
9C6	270984	8235	32,91	5,54	K	T	Y	V	M	Y	T	K	I
81F11	317824	9964	31,9	5,37	K	I	F	V	M	F	F	V	S
84E10	255064	7996	31,9	5,37	I	N	F	R	M	F	S	V	S
71C4	1350144	44339	30,45	5,13	K	N	F	G	M	F	I	V	S
84D3	84424	2920	28,91	4,87	E	N	F	K	M	L	I	E	S
96H8	205904	7224	28,5	4,80	K	Y	Y	K	M	F	I	M	S
85A7	235704	8416	28,01	4,72	K	N	Y	E	M	L	F	V	R
50G10	264144	9470	27,89	4,70	T	N	F	E	M	F	F	V	S
6A1	299824	10912	27,48	4,63	A	N	F	R	M	F	F	M	S
91C4	1157424	44837	25,81	4,35	K	N	F	G	M	L	I	M	R
2C4	258264	10139	25,47	4,29	I	N	F	V	M	F	I	V	L
63C3	188184	7625	24,68	4,15	E	T	Y	K	M	L	F	V	L
850	196024	8115	24,16	4,07	K	N	V	G	M	F	F	V	I
67C10	306104	12907	23,72	3,99	E	T	F	V	M	F	F	M	L
10F9	165984	7113	23,34	3,93	I	I	Y	V	M	Y	F	E	R
4C1	385504	16548	23,3	3,92	K	N	S	V	M	F	I	E	I
86G3	183944	7995	23,01	3,87	T	S	Y	V	M	F	T	V	L
51G10	215264	9727	22,13	3,73	A	N	Y	R	M	F	I	K	S
58A5	90744	4142	21,91	3,69	V	T	F	R	M	L	I	M	S
56F8	235344	10823	21,74	3,66	I	N	F	E	M	F	T	E	L
67C11	209184	9856	21,22	3,57	K	Y	Y	I	M	F	F	E	I
91C8	333584	16012	20,83	3,51	K	N	F	G	M	L	I	K	S
48B11	302864	14946	20,26	3,41	I	N	A	G	M	L	I	E	S
78F11	84104	4155	20,24	3,41	I	I	F	R	M	Y	F	E	I

Table 5. Polypeptides expressed in HEK293F.

Expression and CR6261 binding were determined as described and the ratio of binding and expression signals calculated. The mutations included in each clone are indicated in Table 4 and 5.

Clone	CR6261 binding signal	HTRF signal	ratio	fold increase of ratio over parental SEQ ID NO: 6
127H1	24150000	327363	73,77	4,25
86B4	19970680	334887	59,63	3,44
171E5	6625080	235511	28,13	1,62
7A7	6191080	242461	25,53	1,47
71H2	21080360	336346	62,67	3,61
220C9	8493560	162872	52,15	3,00
131B5	5725640	139561	41,03	2,36
115A1	9557640	175377	54,50	3,14
74H9	26144240	344988	75,78	4,37
71C4	6413600	214495	29,90	1,72
91C4	8442400	245138	34,44	1,98
113E7	13005960	260748	49,88	2,87
6E12	15326000	309443	49,53	2,85
181H9	11892520	324690	36,63	2,11
SEQ ID NO: 6 AV	5661550	326077	17,36	1,00

Table 6. Naturally occurring sequence variation at the indicated positions in % of total number of sequences for each subtype

Position	amino acid	H1	H3	H5	H7
337	V	67	99	19	100
	I	32	1	2	
	T	0,8		3	
	S			73	
	Y			0,1	
	N			0,5	
	A			2	
	G			0,1	
340	I	99		21	98
	V	0,43			
	T	0,03	0,5		
	K		97		
	R		2	47	
	G			29	
	E			0,3	
	S				2
352	F	100	100	100	100
353	I	99,9	100	100	100
	L	0,1			
402	M	100		100	
	T		99,8		100
	S		0,02		

Table 7. Purification and strength of mAb binding of polypeptides

	SEQ ID NO:	Volume supernatant (ml)	Yield (mg/l of culture)	Purity from HP- SEC (%)	K_d^{app} CR6261 (nM)	K_d^{app} CR9114 (nM)
s127H1	35	1376	9.0	100.0	130	10
s86B4	36	1380	9.0	96.0	150	13
s55G7	37	1460	18.1	100.0	150	9
s74H9	34	1335	11.3	99.7	130	10
s6E12	38	1479	13.1	90.8	390	34

Table 8. Molecular weights as determined by SEC-MALS for polypeptides of the invention and their complexes with Fab fragments of CR6261 and CR9114. Theoretical (theor) values are estimated on the basis of the sequence of the polypeptide of the invention (assuming a monomer) and an additional contribution of approximately 10 kDa from attached glycans. The molecular weights of the Fab fragments of CR6261, CR9114 and CR8020 were also determined by SEC-MALS, and were 48, 49 and 47 kDa, respectively.

	SEQ ID NO:	MW (kDa)		MW complex with CR6261 (kDa)		MW complex with CR9114 (kDa)	
		Theor	Observed	Theor	Observed	Theor	Observed
s127H1	35	40	39	87	74	86	83
s86B4	36	40	40	88	75	87	83
s55G7	37	40	40	90	66	87	80
s74H9	34	40	41	89	72	88	83
s6E12	38	40	40	88	67	87	80

Table 9. Disulfide bridges designed and tested

Cluster	Cysteines introduced at position
14	423 and 424
15	430 and 431
16	404 and 433
17	405 and 429
18	411 and 419
19	38 and 390
21	39 and 393
22	36 and 394
23	342 and 460
24	344 and 467
25	344 and 464

Table 10. Molecular weights as determined from SEC MALS experiments. Theoretically

5 expected values for trimeric FL or s127H1-t2-cl18 and the trimeric FL or s127H1-t2-cl18 complex with Fab fragments (3 per trimer) are given between brackets

Construct Name	Mw (kDa)		***K _d ^{app} (nM)		
	Trimer	Trimer protein complex with		Protein complex with	
		CRF9114	CRF6261	CR9114	CR6261
s127H1-t2-cl18long	108 (120)	241 (246)	216 (255)	0.5	0.5

*** K_d^{app} calculated from steady state Octet measurements (see figure 9)

Table 11. Design of interprotomer disulfide bridges for stem domain polypeptides derived from HA of group 2 Influenza strains

Cluster	Cysteines introduced at position	Remarks
14	425 and 426	Numbering refers to the sequence of full length HA from H3N2 A/Hong Kong/1/1968 (SEQ ID NO: 237)
15	432 and 433	
17	407 and 431	
18	413 and 421	
30	410 and 428	
31	411 and 428	

5

Table 12. Summary of survival proportion data (in %) reported in examples 17-21

subtype	Challenge strain	SEQ ID NO: 186	SEQ ID NO: 254	SEQ ID NO: 203	SEQ ID NO: 186*	Example no:
H5N1	A/HK/156/97	100				17 (serum transfer)
H1N1	A/Bris/59/07		100	100	100	18
H1N1	A/NL/602/09		60	100	70	19
H1N1	A/PR/8/34		100	100	100	21
H5N1	A/HK/156/97		100	100	100	20

Survival % are reported at day 21 post challenge (end of follow-up) and are all significantly different from the negative control group receiving PBS.

*expressed in Sf9 insect cells;

10

Table 13. The protein yield and the EC50 values of purified material in the sandwich ELISA

Construct	Yield (mg/l culture supernatant)	Antibody	EC50 (μ g/ml)
s127H1-t2-cl18long SEQ ID NO: 181	~11.1	CR9114	0.026
		CR6261	0.139
Tex_s127H1-t2-cl18 SEQ ID NO: 208	~13.7	CR9114	0.045
		CR6261	0.369
NY_s127H1-t2-cl18 SEQ ID NO: 210	~2.7	CR9114	0.070
		CR6261	0.455

Table 14: Characterization of polypeptides of the invention with additional c-terminal trimerization domain.

Construct SEQ ID NO:	Expression AlphaLISA (counts)	Expression (intensity of trimer band in Western blot)	Multimer ELISA (Log ₁₀ EC ₅₀ of supernatant dilution)	CR9114 binding AlphaLISA (counts)	CR6261 binding AlphaLISA (counts)
186		Very good	4.43	2.71E+05	1.55E+05
255	1.38E+06	Very good	4.81	8.79E+05	5.72E+05
256	1.33E+06	Very good	4.72	7.14E+05	4.75E+05
257	1.67E+06	Very good	4.86	8.61E+05	5.87E+05
258	1.54E+06	Very good	4.82	8.42E+05	6.25E+05

REFERENCES

- Alberini et al. (2009), *Vaccine* 27: 5998–6003.
- Bommakanti et al. (2010), *PNAS* 107(31): 13701-13706.
- 5 Bommakanti et al. (2012), *J Virol* 86: 13434.
- Cheng et al. (2014), *J. Immunol. Methods* 1–13. (doi:10.1016/j.jim.2014.07.010)
- Coffman et al. (2010), *Immunity* 33: 492.
- Devereux et al. (1984), *Nucl. Acids Res.* 12: 387.
- DiLillo et al. (2014), *Nat Med* 20, 143.
- 10 Dopheide TA, Ward CW. (1981) *J Gen Virol.* 367–370
- Ekiert et al. (2009), *Science* 324:246.
- Ekiert et al. (2011), *Science* 333: 844.
- Ferguson et al. (2003), *Nature* 422: 428-443.
- Lorieau et al. 2010, *Proc. Natl. Acad. Sci. USA*, 107: 11341.
- 15 Lu et al. (2013), www.pnas.org/cgi/doi/10.1073/pnas.1308701110.
- Mallajosyula et al. (2014)., www.pnas.org/cgi/doi/10.1073/pnas.1402766111.
- Parekh et al. (2012), *mAbs* 4: 310.
- Safronetz et al (2011) *J Virol.* 85:1214
- Schnueriger et al. (2011), *Molecular immunology* 48: 1512.
- 20 Steel et al. (2010), *mBio* 1(1): 1-9.
- Steven et al. (2004) *Science* 303: 1866.
- Steven et al. (2006) *Science* 312: 404.
- Temperton et al. (2007) *Viruses* 1: 105–12.
- Throsby et al. (2008), *Plos One* 12(3): 1-15.
- 25 Wilson et al (1981) *Nature* 289: 366.

SEQUENCES

SEQ ID NO 1: H1 Full length (A/Brisbane/59/2007)

MKVKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL 50
 5 ENSHNGKLCL LKGIAPLQLG NCSVAGWILG NPECELLISK ESWSYIVEKP 100
 NPENGTCTPG HFADYEELRE QLSSVSSFER FEIFPKESSW PNHTVTGVSA 150
 SCSHNGESSF YRNLLWLTGK NGLYPNLSKS YANNKEKEVL VLWGVHHPN 200
 IGDQKALYHT ENAYVSVVSS HYSRKFTPEI AKRPKVRDQE GRINYWTLL 250
 EPGDTIIFEA NGNLIAPRYA FALSRGFGSG IINSNAPMDK CDAKCQTPQG 300
 10 AINSSLPFQN VHPVTIGECF KYVRSKLRM VTGLRN**IPSI** QSRGLFGAIA 350
GFIEGGWTGM VDGWYGYHHQ NEQSGGYAAD QKSTQNAING ITNKVNSVIE 400
KMNTQFTAVG **KEFNKL**ERM ENLNKKVDDG FIDIWYNAE LLVLENERT 450
 LDFHDSNVKN LYEKVKSQK NNAKEIGNGC FEFYHKCNDE CMESVKNQTY 500
 DYPKYSEESK LNREKIDGVK LESMGVYQIL AIYSTVASSL VLLVSLGAIS 550
 15 FWMCSNGSLQ CRICI 565

SEQ ID NO: 2: H1-mini2-cluster1+5+6-GCN4

MKVKLLVLLC TFTATYA DTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL 50
 ENGGGGKYVC SAKLRMVTGL RN**IPSI**QSOG LFGAIA**GFI** GGWTGMVDGW 100
 20 YGYHHQNEQG SGYAADQKST QNAINGITNK VNSVIE**KMNT** **QSTATGKEGN** 150
KSERMKQIED KIEEIESKQI WCYNALLVL LENERTLDFH DSNVKNLYEK 200
 VKSQLKNNAK EIGNGCFEFY HKCNDECMES VKNGTYDYPK YSEESKLNRE 250
 KIDGVKLESM GVYQILAIYS TVASSLVLLV SLGAISFMMC SNGSLQCRIC 300
 I 301

25

SEQ ID NO: 3: foldon

GYIPEAPRDGQAYVRKDGEWVLLSTFL

SEQ ID NO: 4: FLAG-thrombin-foldon-HIS

30 SGRDYKDDDDKLVPRGSPGSGYIPEAPRDGQAYVRKDGEWVLLSTFLGHHHHHH

SEQ ID NO: 5:

MKQIEDKIEEIESKQ

35 SEQ ID NO: 6: H1-mini2-cluster1+5+6-GCN4 without leader
 sequence and with FLAG-thrombin-foldon-HIS

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLENGGGGGKYVCSAKLRMVTGLRNIPSIQ
 SQGLFGAIA**GFI**EGGWTGMVDGWYGYHHQNEQSGGYAADQKSTQNAINGITNKVNSVIEK
 MNTQSTATGKEGNKSERMKQIEDKIEEIESKQIWCYNALLVLENERTLDFHDSNVKNL
 40 YEKVKSQLKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVSG
 RDYKDDDDKLVPRGSPGSGYIPEAPRDGQAYVRKDGEWVLLSTFLGHHHHHH

SEQ ID NO 7: H1 consensus sequence residue 402-418
(numbering according to SEQ ID NO:1)
402 MNTQFTAVG KEFN(H/K)LE(K/R) 418

5 >SC09-114 VH PROTEIN (SEQ ID NO: 11)
QVQLVQSGAEVKKPGSSVKVSCKSSGGTSNNYAISWVRQAPGQGLDWMGGISPIFGSTAY
AQKFQGRVTISADIFSN TAYMELNSLTSEDTAVYFCARHGNYYYSGMDVWGQGTTVTVS
S

10 >SC09-114 VL PROTEIN (SEQ ID NO: 12)
SYVLTQPPAVSGTPGQRVTISCSGSDSNIGRRSVN WYQQFPGTAPKLLIYSNDQRPSVVP
DRFSGSKSGTSASLAISGLQSEDEAEYYCAAWDDSLKGAVFGGGTQLTVL

15 >CR6261 VH PROTEIN (SEQ ID NO: 9)
E V Q L V E S G A E V K K P G S S V K V S C K A S G G P F R
S Y A I S W V R Q A P G Q G P E W M G G I I P I F G T T K Y
A P K F Q G R V T I T A D D F A G T V Y M E L S S L R S E D
T A M Y Y C A K H M G Y Q V R E T M D V W G K G T T V T V S
20 S

>CR6261 VL PROTEIN (SEQ ID NO: 10)
Q S V L T Q P P S V S A A P G Q K V T I S C S G S S S N I G
N D Y V S W Y Q Q L P G T A P K L L I Y D N N K R P S G I P
25 D R F S G S K S G T S A T L G I T G L Q T G D E A N Y Y C A
T W D R R P T A Y V V F G G G T K L T V L G

>SC08-057 VH PROTEIN (SEQ ID NO: 13)
30 EVQLVESGGGLVQPGGSLRLSCAASGFTD S VIFMSWVRQAPGKGLECVSIIYIDDSTYYA
DSVKGRFTISRHN S MGT V FLEMNSLRPDDTAVYYCATESGDFGDQTGPYHYAMDV

>SC08-057 VL PROTEIN (SEQ ID NO: 14)
35 QSALTQPASVSGSPGQSITISCTGSSGDIGGYNAVSWYQHHPGKAPKLMIYEVTSRPSGV
SDRFSASRSGDTASLTVSGLQAEDEAHYYCCSFADSNILI

>SC08-020 VH PROTEIN (SEQ ID NO: 17)
40 QVQLQQSGAEVKT P GASVKVSCKASGYTFTRFGVSWIRQAPQG LEWIGWISAYNGDTYYAQKFQ
ARVTMTTDTSTTTAYMEMRSLRSDDTAVYYCAREPPLFYSSWSLDN

>SC08-020 VL PROTEIN (SEQ ID NO: 18)
EIVXTQSPGTL S LSPGERATLSCRASQSVSMNYLAWFQQKPGQAPRLLIYGASRRATGIPDRISG
45 SSGTDFTLTISRLEPADFAVYYCQYGTSPRT

SEQ ID NO: 51: H1-mini2-cluster1+5+6-GCN4t2

MKVKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL 50
 ENGGGGKYVC SAKLRMVTGL RNIPSIQSQG LFGAIAGFIE GGWTGMVDGW 100
 YGYHHQNEQG SGYAADQKST QNAINGITNK VNSVIEKMNT QSTATGKEGN 150
 5 KSERRMKQIE DKIEEIESKI WCYNALLVL LENERTLDFH DSNVKNLYEK 200
 VKSQLKNNNAK EIGNGCFEFY HKCNDECMES VKNGTYDYPK YSEESKLNRE 250
 KIDGVKLESM GYQILAIYS TVASSLVLLV SLGAISFWMC SNGSLQCRIC 300
 I 301

10

SEQ ID NO: 52: H1-mini2-cluster1+5+6-GCN4t3

MKVKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL 50
 ENGGGGKYVC SAKLRMVTGL RNIPSIQSQG LFGAIAGFIE GGWTGMVDGW 100
 YGYHHQNEQG SGYAADQKST QNAINGITNK VNSVIEKMNT QSTATGKEGN 150
 15 KSRMKQIEDK IEEIESKQKI WCYNALLVL LENERTLDFH DSNVKNLYEK 200
 VKSQLKNNNAK EIGNGCFEFY HKCNDECMES VKNGTYDYPK YSEESKLNRE 250
 KIDGVKLESM GYQILAIYS TVASSLVLLV SLGAISFWMC SNGSLQCRIC 300
 I 301

20

SEQ ID NO: 55: 127H1

MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
 MVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYGAAADQKSTQNAINGITNK
 VNSVIEKMNTQYTAIGKEYNKSERMKQIEDKIEEIESKQIWCYNALLVLEENERTLDFHDSNVK
 25 NLYEKVKSQLKNNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
 GYQILAIYSTVASSLVLLVSLGAISFWMC SNGSLQCRICI

SEQ ID NO: 56 : 86B4

MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
 30 MVTGLRNKPSNQSQGLFGAIAGYKEGGWTGMVDGWYGYHHQNEQGSYGAAADQKSTQNAINGITNK
 VNSVIEKMNTQFTAIGKEMNKIERMKQIEDKIEEIESKQIWCYNALLVLEENERTLDFHDSNVK
 NLYEKVKSQLKNNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
 GYQILAIYSTVASSLVLLVSLGAISFWMC SNGSLQCRICI

35

SEQ ID NO: 57 : 74H9

MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
 MVTGLRNKPSNQSQGLFGAIAGFKEGGWTGMVDGWYGYHHQNEQGSYGAAADQKSTQNAINGITNK
 VNSVIEKMNTQYTAFGKEMNKSERMKQIEDKIEEIESKQIWCYNALLVLEENERTLDFHDSNVK
 40 NLYEKVKSQLKNNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
 GYQILAIYSTVASSLVLLVSLGAISFWMC SNGSLQCRICI

SEQ ID NO: 58 : 6E12

MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
 45 MVTGLRNEPSNQSQGLFGAIAGFIEGGWTGMVDGWYGYHHQNEQGSYGAAADQKSTQNAINGITNK
 VNSVIEKMNTQLTAFGKEVNKLERMKQIEDKIEEIESKQIWCYNALLVLEENERTLDFHDSNVK
 NLYEKVKSQLKNNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
 GYQILAIYSTVASSLVLLVSLGAISFWMC SNGSLQCRICI

SEQ ID NO: 59 : 55G7

5 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQYTAIGKEMNKLERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 60 : 115A1

10 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSKQSQGLFGAIAAGYTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQITAVGKEYNKIERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDG
VKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

15 SEQ ID NO: 61 : 71H2

20 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQLTAIGKEVNKSERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

25 SEQ ID NO: 62 : 181H9

30 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNVPSKQSQGLFGAIAAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQFTAVGKEFNKNERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

35 SEQ ID NO: 63 : 220C9

40 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSTQSQGLFGAIAAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQFTATGKEYNKLERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

45 SEQ ID NO: 64 : 113E7

50 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSKQSQGLFGAIAAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQYTATGKEINKHERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 65 : s74H9

50 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAFGK
EMNKSERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 66: s127H1

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
EYNKSERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
5 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 67: s86B4

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTAIGK
10 EMNKIERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 68: s55G7

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
15 EMNKLERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 69: s6E12

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNEPSNQSQGLF
GAIAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQLTAFGK
20 EVNKLERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 70: s115A1

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGYTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQITAVGK
25 EYNKIERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 71: s71H2

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQLTAIGK
30 EVNKSERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 76 : s181H9

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNVPSKQSQGLF
GAIAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTAVGK
40 EFNKNERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 77 : s220C9

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSTQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQFTATGK
EYNKLERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
5 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 78 : s113E7

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQYTATGK
10 EINKHERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 72: s74H9-long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
15 GAIAGFKEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQYTAFGK
EMNKSERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 73: s127H1-long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
20 GAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
EYNKSERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 74: s86B4-long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
25 GAIAGYKEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQFTAIGK
EMNKIERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG
30

SEQ ID NO: 75: s55G7-long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
35 GAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
EMNKLERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 144: s6E12-long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNEPSNQSQGLF
40 GAIAGFIEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQLTAFGK
EVNKLERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG
45

SEQ ID NO: 79: s115A1ong

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGYTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQITAVGK
EYNKIERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
5 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 80: s71H21ong

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
10 GAIAGFKEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQLTAIGK
EVNKSERMKQIEDKIEEIESKQIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 81: 127H1-t2

15 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNK
VNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
20 GYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 82 : 86B4-t2

25 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAGYKEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNK
VNSVIEKMNTQFTAIGKEMNKIERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 83 : 74H9-t2

30 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAGFKEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNK
VNSVIEKMNTQYTAFGKEMNKSEERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
35 GYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 84 : 6E12-t2

40 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNEPSNQSQGLFGAIAGFIEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNK
VNSVIEKMNTQLTAFGKEVNKLERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 85 : 55G7-t2

45 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNK
VNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
50 GYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 86 : 115A1-t2

5 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSKQSQGLFGAIAAGYTEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
VNSVIEKMNTQITAVGKEYNKIERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 87 : 71H2-t2

10 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAAGFKEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
VNSVIEKMNTQLTAIGKEVNKSERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 88 : 181H9-t2

15 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNVPSKQSQGLFGAIAAGFIEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
VNSVIEKMNTQFTAVGKEFNKNERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
20 GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 89 : 220C9-t2

25 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSTQSQGLFGAIAAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
VNSVIEKMNTQFTATGKEYNKLERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 90 : 113E7-t2

30 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSKQSQGLFGAIAAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
VNSVIEKMNTQYTATGKEINKHERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
35 GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 91: s127H1-t2

40 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
EYNKSERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
NGNCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHH

SEQ ID NO: 92 : s86B4-t2

45 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYKEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTAGK
EMNKIERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
NGNCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHH
50

SEQ ID NO: 93 : s74H9-t2

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAFGK
EMNKSERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
5 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 94 : s6E12-t2

10 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNEPSNQSQGLF
GAIAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQLTAFGK
EVNKLERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 95 : s55G7-t2

15 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
EMNKLERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 96 : s115A1-t2

20 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGYTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQITAVGK
EYNKIERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
25 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 97 : s71H2-t2

30 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQLTAIGK
EVNKSERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 98 : s181H9-t2

35 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNVPSKQSQGLF
GAIAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTAVGK
EFNKNERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 99 : s220C9-t2

40 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSTQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTATGK
EYNKLERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
45 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 100 : s113E7-t2

50 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTATGK
EINKHERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 101: s127H1-t2long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
5 EYNKSERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 102 : s86B4-t2long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTAIGK
10 EMNKIERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 103 : s74H9-t2long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAFGK
15 EMNKIERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
20 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 104 : s6E12-t2long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNEPSNQSQGLF
GAIAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQLTAFGK
25 EVNKLERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 105 : s55G7-t2long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
30 EMNKLERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 106 : s115A1-t2long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGYTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQITAVGK
35 EYNKIERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 107 : s71H2-t2long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQLTAIGK
40 EVNKSERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
45 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 108 : s181H9-t2long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNVPSKQSQGLF
GAIAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTAIGK
50 EFNKNERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 109 :s220C9-t2long

5 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSTQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTATGK
EYNKLERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

10 SEQ ID NO: 110 : s113E7-t2long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTATGK
EINKHERRMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
15 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 111: 127H1-t3

20 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQYTAIGKEYNKS RMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

25 SEQ ID NO: 112 : 86B4-t3

MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAGYKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQFTAIGKEMNKIRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
30 GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 113 : 74H9-t3

35 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQYTAFGKEMNKS RMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

40 SEQ ID NO: 114 : 6E12-t3

MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNEPSNQSQGLFGAIAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQLTAFGKEVNKL RMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVK
45 NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 115 : 55G7-t3

5 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQYTAIGKEMNKLRMKQIEDKIEEIESKQKIWCYNAELLVLENERLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 116 : 115A1-t3

10 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSKQSQGLFGAIAAGYTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQITAVGKEYNKIRMKQIEDKIEEIESKQKIWCYNAELLVLENERLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 117 : 71H2-t3

15 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSNQSQGLFGAIAAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQLTAIGKEVNKSRMKQIEDKIEEIESKQKIWCYNAELLVLENERLDFHDSNVK
20 NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 118 : 181H9-t3

25 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNVPSKQSQGLFGAIAAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQFTAVGKEFNKNRMKQIEDKIEEIESKQKIWCYNAELLVLENERLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 119 : 220C9-t3

30 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSTQSQGLFGAIAAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQFTATGKEYNKLRMKQIEDKIEEIESKQKIWCYNAELLVLENERLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
35 GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 120 : 113E7-t3

40 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSKQSQGLFGAIAAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQYTATGKEINKHRMKQIEDKIEEIESKQKIWCYNAELLVLENERLDFHDSNVK
NLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 121: s127H1-t3

45 DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
EYNKSRMKQIEDKIEEIESKQKIWCYNAELLVLENERLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

50

SEQ ID NO: 122 : s86B4-t3

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYKEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQFTAIGK
EMNKIRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
5 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 123 : s74H9-t3

10 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQYTAFGK
EMNKS RMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

15 SEQ ID NO: 124 : s6E12-t3

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNEPSNQSQGLF
GAIAGFIEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQLTAFGK
EVNKL RMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
20 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 125 : s55G7-t3

25 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
EMNKL RMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 126 : s115A1-t3

30 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGYTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQITAVGK
EYNKIRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 127 : s71H2-t3

35 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQLTAIGK
EVNKS RMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 128 : s181H9-t3

40 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNVPSKQSQGLF
GAIAGFIEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQFTAIGK
EFNKNRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

45 SEQ ID NO: 129 : s220C9-t3

50 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSTQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQFTATGK
EYNKL RMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHHH

SEQ ID NO: 130 : s113E7-t3

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTATGK
EINKHRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
5 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGRSLVPRGSPGHHHHH

SEQ ID NO: 131: s127H1-t3long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
10 EYNKSRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 132 : s86B4-t3long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTAIGK
EMNKIRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
15 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 133 : s74H9-t3long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAFGK
20 EMNKSRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 134 : s6E12-t3long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNEPSNQSQGLF
GAIAGFIEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQLTAFGK
30 EVNKLRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 135 : s55G7-t3long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGK
35 EMNKLRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 136 : s115A1-t3long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
GAIAGYTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQITAVGK
40 EYNKIRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 137 : s71H2-t3long

DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSNQSQGLF
GAIAGFKEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQLTAIGK
50 EVNKSARMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 138 : s181H9-t3long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNVPSKQSQGLF
 5 GAIAGFIEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQFTAVGK
 EFNKNRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 139 : s220C9-t3long

10 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSTQSQGLF
 GAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQFTATGK
 EYNKLRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

15 SEQ ID NO: 140 : s113E7-t3long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
 GAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQYTATGK
 EINKHRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
 20 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 141 : s181H9long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNVPSKQSQGLF
 25 GAIAGFIEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQFTAVGK
 EFNKNRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 142 : s220C9long

30 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSTQSQGLF
 GAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQFTATGK
 EYNKLRMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 143 : s113E7long

35 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
 GAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQYTATGK
 EINKHERMKQIEDKIEEIESKQKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKLNNAKEI
 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG

40

SEQ ID NO: 149 : 55G7-t2-cl14

MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC
 SAKLRMVTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
 QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQCCDKIEEIESKIWCYNAELLVL
 45 LENERTLDFHDSNVKNLYEKVKSQKLNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPK
 YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 150 : 55G7-t2-cl15

5 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEECCSKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 151 : 55G7-t2-cl16

10 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 152 : 55G7-t2-cl17

15 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTCYTAIGKEMNKLERRMKQIEDKIECIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
20 YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 153 : 55G7-t2-cl18

25 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGCEMNKLERCMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 154 : 55G7-t2-cl19

30 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLCKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINCITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
35 YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 155 : 55G7-t2-cl21

40 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLECNVTVTTHSVNLLLENGGGGKYVC
SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITCKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 156 : 55G7-t2-cl22

45 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTCLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNCVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

50

SEQ ID NO: 157 : 55G7-t2-c123

5 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSNQCGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKCLYEKVKSQLKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 158 : 55G7-t2-c124

10 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSNQSQCLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKCQLKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 159 : 55G7-t2-c125

15 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSNQSQCLFGAIAGYVEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEVCVKSQLKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
20 YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 160 : 127H1-t2-c114

25 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQCCDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQLKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 161 : 127H1-t2-c115

30 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEECCSKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQLKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
35 YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 162 : 127H1-t2-c116

40 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNCQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQLKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 163 : 127H1-t2-c117

45 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTCYTAIGKEYNKSERRMKQIEDKIECIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQLKNNAKEIGNGCFFYHKNDECMESEVKNGTYDYPK
50 YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMCNNGSLQCRICI

SEQ ID NO: 164 : 127H1-t2-cl18

5 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECME SVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMC SNGLQCRICI

SEQ ID NO: 165 : 127H1-t2-cl19

10 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLCKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINCITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECME SVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMC SNGLQCRICI

SEQ ID NO: 166 : 127H1-t2-cl21

15 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLECNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITCKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECME SVKNGTYDYPK
20 YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMC SNGLQCRICI

SEQ ID NO: 167 : 127H1-t2-cl22

25 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTCLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNCVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECME SVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMC SNGLQCRICI

SEQ ID NO: 168 : 127H1-t2-cl23

30 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQCQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKCLYEKVKSQLKNNAKEIGNGCFFYHKNDECME SVKNGTYDYPK
35 YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMC SNGLQCRICI

SEQ ID NO: 169 : 127H1-t2-cl24

40 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQCLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKCQLKNNAKEIGNGCFFYHKNDECME SVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMC SNGLQCRICI

SEQ ID NO: 170 : 127H1-t2-cl25

45 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQCLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYECVKSQKNNAKEIGNGCFFYHKNDECME SVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAISFWMC SNGLQCRICI

50

SEQ ID NO: 171 : s55G7-t2-cl14long
DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQCCDKIEEIESKIWCYNAELLVL
5 LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKCNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 172 : s55G7-t2-cl15long
DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC
10 SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEECCSKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKCNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

15 SEQ ID NO: 173 : s55G7-t2-cl16long
DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST
QNAINGITNKVNSVIEKMNCQYTAIGKEMNKLERRMKQIEDKIEEIESCIWCYNAELLVL
20 LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKCNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 174 : s55G7-t2-cl17long
DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST
25 QNAINGITNKVNSVIEKMNTCYTAIGKEMNKLERRMKQIEDKIECIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKCNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

30 SEQ ID NO: 175 : s55G7-t2-cl18long
DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGCEMNKLERCMKQIEDKIEEIESKIWCYNAELLVL
35 LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKCNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 176 : s55G7-t2-cl19long
DTICIGYHANNSTDTVDTVLCKNVTVTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST
40 QNAINCITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKCNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 177 : s55G7-t2-cl21long
DTICIGYHANNSTDTVDTVLECNVTVTHSVNLLLENGGGGKYVC
45 SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST
QNAINGITCKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL
LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKCNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 178 : s55G7-t2-cl22long

DTICIGYHANNSTDTVDTCLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST

5 QNAINGITNCVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL

LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 179 : s55G7-t2-cl23long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

10 SAKLRMTGLRNKPSNQSQGLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST

QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL

LENERTLDFHDSNVKCLYEKVKSQLKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

15 SEQ ID NO: 180 : s55G7-t2-cl24long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSNQSQCLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST

QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL

20 LENERTLDFHDSNVKNLYEKVKCQLKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 181 : s55G7-t2-cl25long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSNQSQCLFGAIAGYVEGGWTGMVDGWYGYHHQNEQSGYAADQKST

25 QNAINGITNKVNSVIEKMNTQYTAIGKEMNKLERRMKQIEDKIEEIESKIWCYNAELLVL

LENERTLDFHDSNVKNLYECVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

30 SEQ ID NO: 182 : s127H1-t2-cl14long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKST

QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQCCDKIEEIESKIWCYNAELLVL

35 LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 183 : s127H1-t2-cl15long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKST

40 QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEECCSKIWCYNAELLVL

LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 184 : s127H1-t2-cl16long

45 DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKST

QNAINGITNKVNSVIEKMNCQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL

LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

50

SEQ ID NO: 185 : s127H1-t2-cl17long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKST

QNAINGITNKVNSVIEKMNTCYTAIGKEYNKSERRMKQIEDKIECIESKIWCYNAELLVL

5 LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 186 : s127H1-t2-cl18long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKST

QNAINGITNKVNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVL

LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 187 : s127H1-t2-cl19long

DTICIGYHANNSTDTVDTVLCNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKST

QNAINCITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL

20 LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 188 : s127H1-t2-cl21long

DTICIGYHANNSTDTVDTVLECNVTVTHSVNLLLENGGGGKYVC

25 SAKLRMTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKST

QNAINGITCKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL

LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 189 : s127H1-t2-cl22long

DTICIGYHANNSTDTVDTCLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKST

QNAINGITNCVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL

LENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGCFFYHKNDECMESVKNGTYDYPK

35 YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 190 : s127H1-t2-cl23long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSKQCQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKST

40 QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL

LENERTLDFHDSNVKCLYEKVKSQLKNNAKEIGNGCFFYHKNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 191 : s127H1-t2-cl24 long

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVC

SAKLRMVTGLRNKPSKQSQCLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKST

QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL

LENERTLDFHDSNVKNLYEKVKCQLKNNAKEIGNGCFFYHKNDECMESVKNGTYDYPK

YSEESKLNREKIDGVKLESMGVYQIEG

50

SEQ ID NO: 192 : s127H1-t2-cl24long
DTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLENGGGGKYVC
SAKLRMVTGLRNKPSKQSQCLFGAIAAGFTEGGWTGMVDGWYGYHHQNEQGSgyAADQKST
QNAINGITNKVNSVIEKMNTQYTAIGKEYNKSERRMKQIEDKIEEIESKIWCYNAELLVL
5 LENERTLDFHDSNVKNLYECVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPK
YSEESKLNREKIDGVKLESMGVYQIEG

SEQ ID NO: 193: CMKQIEDKIEEIESK

10 SEQ ID NO: 194: RMCQIEDKIEEIESKQK

SEQ ID NO: 195: smH1 Cali3964-55G7
MKAILVLLYTFATANADTLCIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLR
LATGLRNKPSNQSQGLFGAIAAGYVEGGWTGMVDGWYGYHHQNEQGSgyAADLKSTQNAIDEITNK
15 VNSVIEKMNTQYTAIGKEMNHLERMKQIEDKIEEIESKQIWCYNAELLVLENERTLDYHDSNVK
NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGRSLVPR
GSPGHHHHHH

SEQ ID NO: 196: smH1 Cali3964-86B4
20 MKAILVLLYTFATANADTLCIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLR
LATGLRNKPSNQSQGLFGAIAAGYKEGGWTGMVDGWYGYHHQNEQGSgyAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGKEMNHIERMKQIEDKIEEIESKQIWCYNAELLVLENERTLDYHDSNVK
NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGRSLVPR
GSPGHHHHHH

25 SEQ ID NO: 197: smH1 Cali3964-127H1
MKAILVLLYTFATANADTLCIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLR
LATGLRNKPSKQSQGLFGAIAAGFTEGGWTGMVDGWYGYHHQNEQGSgyAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGKEYNHSERRMKQIEDKIEEIESKQIWCYNAELLVLENERTLDYHDSNVK
30 NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGRSLVPR
GSPGHHHHHH

SEQ ID NO: 198: _smH1 Cali3964-55G7-t2
35 MKAILVLLYTFATANADTLCIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLR
LATGLRNKPSNQSQGLFGAIAAGYVEGGWTGMVDGWYGYHHQNEQGSgyAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGKEMNHLERRMKQIEDKIEEIESKIWCYNAELLVLENERTLDYHDSNVK
NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGRSLVPR
GSPGHHHHHH

40 SEQ ID NO: 199: _smH1 Cali3964-86B4-t2
MKAILVLLYTFATANADTLCIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLR
LATGLRNKPSNQSQGLFGAIAAGYKEGGWTGMVDGWYGYHHQNEQGSgyAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGKEMNHIERRMKQIEDKIEEIESKIWCYNAELLVLENERTLDYHDSNVK
45 NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGRSLVPR
GSPGHHHHHH

SEQ ID NO: 200: smH1 Cali3964-127H1-t2
50 MKAILVLLYTFATANADTLCIGYHANNSTDTVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLR
LATGLRNKPSKQSQGLFGAIAAGFTEGGWTGMVDGWYGYHHQNEQGSgyAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGKEYNHSERRMKQIEDKIEEIESKIWCYNAELLVLENERTLDYHDSNVK

NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGRSLVPR
GSPGHHHHH

SEQ ID NO: 201: mH1 Cali3964-127H1-t2

5 MKAILVVLLYTFATANADTLCIGYHANNSTDTVDTVLEKNVTVTHSVNLLLEDGGGGKYVCSTKLRLATGLRNKPSKQSQGLFGAIAAGFTTEGGWTGMVDGWYGYHHQNEQSGYAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGKEYNHSEERRMKQIEDKIEEIESKIWCYNAELLVLENERLTDYHDSNVK
NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGVKLEST
10 RIYQILAIYSTVASSLVVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 202: mH1 Cali3964-127H1-t2-cl18

15 MKAILVVLLYTFATANADTLCIGYHANNSTDTVDTVLEKNVTVTHSVNLLLEDGGGGKYVCSTKLRLATGLRNKPSKQSQGLFGAIAAGFTTEGGWTGMVDGWYGYHHQNEQSGYAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLENERLTDYHDSNVK
NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGVKLEST
RIYQILAIYSTVASSLVVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 203: smH1 Cali3964-127H1-t2-cl18long

20 DTLCIGYHANNSTDTVDTVLEKNVTVTHSVNLLLEDGGGGKYVCSTKLRLATGLRNKPSKQSQGLFGAIAAGFTTEGGWTGMVDGWYGYHHQNEQSGYAADLKSTQNAIDEITNKVNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLENERLTDYHDSNVKNLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGVKLESTRIYQIEG

SEQ ID NO: 204: FL HA H1N1 A/AA_Marton/43

30 MKARLLVLLC ALAATDADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL 050
EDSHNGKLCR LKGIAPLQLG KCNIAGWILG NPECESLLSE RSWSYIVETP 100
NSENCTCYPG DFIDYEELRE QLSSVSSFER FEIFSKESSW PKHNTTRGVT 150
AACSHAGKSS FYRNLLWLTE KDGSYPNLNN SYVNKKGKEV LVLWGVHHP 200
NIKDQQTLYQ KENAYVSVVS SNYNRRFTPE IAERP KVRGQ AGRMNYWTL 250
LKPGDTIMFE ANGNIAPWY AFALSRGFGS GIITSNASMH ECDTKCQTPQ 300
GAINSSLPFQ NIHPVTIGEC PKYVRSTKLRL MVTGLRNIPS IQSRGLFGAI 350
AGFIEGGWTG MIDGWYGYHH QNEQSGYAA DQKSTQNAIN GITNKVNSVI 400
35 EKMNTQFTAV GKEFNNLEKR MENLNKKVDD GFLDIWTYNA ELLVLENER 450
TLDFHDSNVK NLYEKVKNQL RNNAKEIGNG CFEFYHKCNN ECMESVKNGT 500
YDYPKYSEES KLNREKIDG V KLESMGVYQI LAIYSTVASS LVLLVSLGAI 550
SFWMCSNGSL QCRICI 565

SEQ ID NO: 205: FL HA H1N1 A/Texas/UR06-0526/07

40 MKVKLLVLLC TFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL 050
EDSHNGKLCR LKGTAPLQLG NCSVAGWILG NPECELLISK ESWSYIVETP 100
NPENCTCYPG YFADYEELRE QLSSVSSFER FEIFPKESSW PNHTVTGVSA 150
SCSHNGKSSF YRNLLWLTKG NGLYPNLSKS YANNKEKEVL VLWGVHHPN 200
45 IGDQRALYHT ENAYVSVVSS HYSRRFTPEI AKRPKVRDQE GRINYYWTLL 250

EPGDTIIFEA NGNLIAPRFA FALSRRGFGSG IITSNAPMGE CDAKCQTPQG 300
 AINSSLFPQN VHPVTIGECF KYVRSACLRLM VTGLRNIPSI QSRGLFGAIA 350
 GFIEGGWTGM VDGWYGYHHQ NEQGSGYAAD QKSTQNAING ITNKVNSVIE 400
 KMNTQFTAVG KEFNKLERRM ENLNKKVDDG FLDIWTYNAE LLVLLNERT 450
 5 LDFHDSNVKN LYEKVKVKNL NNAKEIGNGC FEFYHKCNDE CMESVKNGTY 500
 DYPKYSEESK LNREKIDGVK LESMGVYQIL AIYSTVASSL VLLISLGAIS 550
 FWMCSNGSLQ CRICI 565

SEQ ID NO: 206: H1N1 A/New York/629/95

10 MKVKLLVLLC AFTATYADTI CIGYHANNST DTVDTVLEKN VTVTHSVNLL 050
 EDSHNGKLCR LKGTAQLQLG NCSVAGWILG NPECESLFSK ESWSYIAETP 100
 NPENGTCYPG YFADYEELRE QLSSVSSFER FEIFPKESSW PNHTVTKGVT 150
 ASCSHNGKSS FYKNLLWLTE KNGLYPNLSK SYVNNKEKEV LVLWGVHHP 200
 NIGDQRAIYH TENAYVSVVS SHYSRRFTPE IAKRPKVRDQ EGRINYWT 250
 15 LEPGDTIIFE ANGNLIAPWY AFALSRRGFGS GIITSNASMS ECDACQTPQ 300
 GAINSSLFPQ NVHPVTIGEC PKYVRSTKLR MVTGLRNIPS IQSRGLFGAI 350
 AGFIEGGWTG MIDGWYGYHH QNEQGSGYAA DQKSTQNAID GITNKVNSVI 400
 EKMNTQFTAV GKEFNKLERR MENLNKKVDD GFLDIWTYNA ELLVLLNER 450
 TLDFHDSNVK NLYEKVKVKNL KNNAKEIGNC CFEFYHKCNN ECMESVKNGT 500
 20 YDYPKYSEES KLNREKIDGV KLESMGVYQI LAIYSTVASS LVLLVSLGAI 550
 SFWMCSNGSL QCRICI 566

SEQ ID NO: 207: H1 mini Texas 127H1_t2+c118

25 MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLEDGGGGKYVCSAKLR
 MVTGLRNKPSKQSQGLFGAIAAGFTGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
 VNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLLNERTLDFHDSNVK
 NLYEKVKVKNLKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
 GYQILAIYSTVASSLVLLISLGAISFWMCSNGSLQCRICI

30 SEQ ID NO: 208: #5123_sH1 mini Texas 127H1_t2+c118

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLEDGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
 GAIAGFTGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGC
 EYNKSERCMKQIEDKIEEIESKIWCYNAELLVLLNERTLDFHDSNVKNLYEKVKVKNLKNNAKEI
 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQI

35

SEQ ID NO: 209: H1 mini NY 127H1_t2+c118

MKAKLLVLLCAFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLEDGGGGKYVCSTKLR
 MVTGLRNKPSKQSQGLFGAIAAGFTGGWTGMIDGWYGYHHQNEQGSGYAADQKSTQNAIDGITNK
 VNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLLNERTLDFHDSNVK

NLYEKVKTKQLKNNAKEIGNGCFEFYHKCNNECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQI LAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 210: #5124_sH1 mini NY 127H1_t2+c118

5 DTICIGYHANNSTDVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMIDGWYGYHHQNEQGSYAADQKSTQNAIDGITNKVNSVIEKMNTQYTAIGC
EYNKSERCMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKTKQLKNNAKEI
GNGCFEFYHKCNNECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQI

SEQ ID NO: 211: H1 mini Cal 127H1_t2+c118

10 MKAILVLLYTFATANADTLCIGYHANNSTDVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKL
LATGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGCEYNHSEKCMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDYHDSNVK
NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGVKLEST
RIYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

15

SEQ ID NO: 212: sH1 mini Cal 127H1_t2+c118

DTLCIGYHANNSTDVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLRLATGLRNKPSKQSQGLF
GAIAGFTEGGWTGMVDGWYGYHHQNEQGSYAADLKSTQNAIDEITNKVNSVIEKMNTQYTAIGC
EYNHSEKCMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDYHDSNVKNLYEKVRSQKNNAKEI
20 GNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGVKLESTRIYQI

SEQ ID NO: 213: H1 mini Mart 127H1_t2+c118+loop

MKARLLVLLCALAATDADTICIGYHANNSTDVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKL
MVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMIDGWYGYHHQNEQGSYAADQKSTQNAINGITNK
25 VNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
NLYEKVKNQLRNNAKEIGNGCFEFYHKCNNECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 214: #5126 sH1 mini Mart 127H1_t2+c118+loop

30 DTICIGYHANNSTDVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLRMVTGLRNKPSKQSQGLF
GAIAGFTEGGWTGMIDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGC
EYNKSERCMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVKNLYEKVKNQLRNNAKEI
GNGCFEFYHKCNNECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQI

35 SEQ ID NO: 215: H1 mini Mart 127H1_t2+c118

MKARLLVLLCALAATDADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLEDGGGGKYVCSTKLR
 MVTGLRNKPSKQSQGLFGAIAAGFTEGGWTGMIDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
 VNSVIEKMNTQYTAIGCEYNNSERCMKQIEDKIEEIESKIWCYNAELLVLENERTLDFHDSNVK
 NLYEKVKNQLRNNAKEIGNGCFFEFYHKCNNECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
 5 GVIYQILAIYSTVASSLVLLVSLGAISFWMCSNGSLQCRICI

SEQ ID NO: 216: sH1 mini Mart 127H1_t2+c118

10 MKARLLVLLCALAATDADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLEDGGGGKYVCSTKLR
 MVTGLRNKPSKQSQGLFGAIAAGFTEGGWTGMIDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
 VNSVIEKMNTQYTAIGCEYNNSERCMKQIEDKIEEIESKIWCYNAELLVLENERTLDFHDSNVK
 NLYEKVKNQLRNNAKEIGNGCFFEFYHKCNNECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
 GVIYQ

SEQ ID NO: 217: FL HA H2N2 A/Adachi/2/57

15 MAIIYLILLF TAVRGDQICI GYHANNSTEK VDTILERNVT VTHAKDILEK 050
 THNGKLCKLN GIPPELGDC SIAGWLLGNP ECDRLSVPE WSYIMEKENP 100
 RNGLCYPGSF NDYEELKHLL SSVKHFEKVK ILPKDRWTQH TTTGGSQACA 150
 VSGNPSFFRN MVWLTKKGS YPAKGSYNN TSGEQMLIIW GVHHPIDETE 200
 QRTLYQNVGT YVSVGTSTLN KRSTPEIATR PKVNGLGSRM EFSWTLLDMW 250
 20 DTINFESTGN LIAPEYGFKI SKRGSSGIMK TEGTLENCET KCQTPLGAIN 300
 TTLPFHNVHP LTIGECPKYV KSEKLVLATG LRNVQIESR GLFGAIAAGFI 350
 EGGWQGMVDG WYGYHHSNDQ GSGYAADKES TQKAFDGITN KVNSVIEKMN 400
 TQFEAVGKEF GNLERLENL NKKMEDGFLD VWTYNAELLV LMENERTLDF 450
 HDSNVKNLYD KVRMQLRDNV KELGNGCFFEF YHKCDDECMN SVKNGTYDYP 500
 25 KYEEESKLN NR NEIKGVKLSS MGVIYQI LAIY ATVAGSLSLA IMMAGISFWM 550
 CSNGSLQCRI CI 562

SEQ ID NO: 218: H2 mini Adachi 127H1_t2+c118

30 MAIIYLILLFTAVRGDQICIGYHANNSTEKVDITILERNVTVTHAKDILENGGGGGKYVCSEKLVLA
 TGLRNKPQKESQGLFGAIAAGFTEGGWQGMVDGWYGYHHSNDQSGYAADKESTQKAFDGITNKVN
 SVIEKMNTQYEATGCEYGNLERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNL
 YDKVRMQLRDNVKELGNGCFFEFYHKCDDECMNSVKNGTYDYPKYEEESKLN RNEIKGVKLSSMGV
 YQI LAIYATVAGSLSLAIMMAGISFWMCSNGSLQCRICI

35 SEQ ID NO: 219: H2 mini Adachi 127H1_t2+c118+loop

MAIIYLILLFTAVRGDQICIGYHANNSTEKVDITILERNVTVTHAKDILENGGGGGKYVCSEKLVLA
 TGLRNKPQKESQGLFGAIAAGFTEGGWQGMVDGWYGYHHSNDQSGYAADKESTQKAFDGITNKVN
 SVIEKMNTQY**TAYGCEYNK**SERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNL

YDKVRMQLRDNVKELGNGCFEFYHKCDDECMNSVKNGTYDYPKYEEESKLNREIKGVKLSSMGV
YQILAIYATVAGSLSLAIMMAGISFWMCSNGSLQCRICI

SEQ ID NO: 220: #5119 sH2 mini Adachi 127H1_t2+cl18

5 DQICIGYHANNSTEKVDITLERNVTVTTHAKDILENGGGGKYVCSEKLVLATGLRNKPQKESQGLF
GAIAGFTEGGWQGMVDGWYGYHHSNDQSGSYAADKESTQKAFDGITNKVNSVIEKMNTQYEATGC
EYGNLERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNLYDKVRMQLRDNVKEL
GNGCFEFYHKCDDECMNSVKNGTYDYPKYEEESKLNREIKGVKLSSMGVYQI

10 SEQ ID NO: 221: #5120 sH2 mini Adachi 127H1_t2+cl18+loop

DQICIGYHANNSTEKVDITLERNVTVTTHAKDILENGGGGKYVCSEKLVLATGLRNKPQKESQGLF
GAIAGFTEGGWQGMVDGWYGYHHSNDQSGSYAADKESTQKAFDGITNKVNSVIEKMNTQY**TAYGC**
EY**NKS**SERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNLYDKVRMQLRDNVKEL
GNGCFEFYHKCDDECMNSVKNGTYDYPKYEEESKLNREIKGVKLSSMGVYQI

15

SEQ ID NO: 222: FL HA H2N2 A/Singapore/1/1957

	MAIIYLILLF	TAVRGDQICI	GYHANNSTEK	VDITLERNVT	VTHAKDILEK	050
	THNGKLCKLN	GIPPLELGDC	SIAGWLLGNP	ECDRLLSVPE	WSYIMEKENP	100
	RDGLCYPGSF	NDYEELKHL	SSVKHFEKVK	ILPKDRWTQH	TTTGGSRACA	150
20	VSGNPSFFRN	MVWLTEKGSN	YPVAKGSYNN	TSGEQMLIIW	GVHHPNDEKE	200
	QRTLYQNVGT	YVSVGTSTLN	KRSTPDIA	TRPKVNGLGSRM	EFSTWTLDDMW	250
	DTINFESTGN	LIAPYGFYKI	SKRGSSGIMK	TEGTLENCET	KCQTPLGAIN	300
	TTLPFHNVHP	LTIGECPKYV	KSEKLVLATG	LRNVPQIESR	GLFGAIAGFI	350
	EGGWQGMIDG	WYGYHHSNDQ	GSYAADKES	TQKAFDGITN	KVNSVIEKMN	400
25	TQFEAVGKEF	SNLERLENL	NKKMEDGFLD	VWTYNAELLV	LMENERTLDF	450
	HDSNVKNLYD	KVRMQLRDNV	KELGNGCFEF	YHKCDDECMN	SVKNGTYDYP	500
	KYEEESKLN	REIKGVKLSS	MGVYQILAIY	ATVAGSLSLA	IMMAGISFWM	550
	CSNGSLQCRI	CI				562

30 SEQ ID NO: 223: H2 mini Sing 127H1_t2+cl18

MAIIYLILLFTAVRGDQICIGYHANNSTEKVDITLERNVTVTTHAKDILENGGGGKYVCSEKLVLA
TGLRNKPQKESQGLFGAIAGFTEGGWQGMVDGWYGYHHSNDQSGSYAADKESTQKAFDGITNKVN
SVIEKMNTQYEAIGCEYSNLERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNL
YDKVRMQLRDNVKELGNGCFEFYHKCDDECMNSVKNGTYDYPKYEEESKLNREIKGVKLSSMGV
35 YQI LAIYATVAGSLSLAIMMAGISFWMCSNGSLQCRICI

SEQ ID NO: 224: H2 mini Sing 127H1_t2+cl18+loop

MAIIYLILLFTAVRGDQICIGYHANNSTEKVDITILERNVTVTTHAKDILENGGGGKYVCSEKLVLA
TGLRNKPQKESQGLFGAIAGFTEGGWQGMVDGWYGYHHSNDQSGYAADKESTQKAFDGITNKVN
SVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNL
YDKVRMQLRDNVKELGNGCFEFYHKCDDECMNSVKNGTYDYPKYEEESKLNREIKGVKLSSMGV
5 YQILAIYATVAGSLSLAIMMAGISFWMCSNGSLQCRICI

SEQ ID NO: 225: #5121 sH2 mini Sing 127H1_t2+c118

DQICIGYHANNSTEKVDITILERNVTVTTHAKDILENGGGGKYVCSEKLVLATGLRNKPQKESQGLF
GAIAGFTEGGWQGMVDGWYGYHHSNDQSGYAADKESTQKAFDGITNKVNSVIEKMNTQYEAIGC
10 EYNKSERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNLYDKVRMQLRDNVKEL
GNGCFEFYHKCDDECMNSVKNGTYDYPKYEEESKLNREIKGVKLSSMGVYQI

SEQ ID NO: 226: #5122 sH2 mini Sing 127H1_t2+c118+loop

DQICIGYHANNSTEKVDITILERNVTVTTHAKDILENGGGGKYVCSEKLVLATGLRNKPQKESQGLF
15 GAIAGFTEGGWQGMVDGWYGYHHSNDQSGYAADKESTQKAFDGITNKVNSVIEKMNTQYTAIGC
EYNKSERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNLYDKVRMQLRDNVKEL
GNGCFEFYHKCDDECMNSVKNGTYDYPKYEEESKLNREIKGVKLSSMGVYQI

SEQ ID NO: 227: FL HA H5N1 A/Vietnam/1203/2004

MEKIVLLFAIVSLVKSDQICIGYHANNSTEQVDTIMEKNVTVTTHAQDILEKKHNGKLCDLDGVKP
20 LILRDCSVAGWLLGNPMCDEFINVPESYIVEKANPVNDLCYPGDFNDYEELKHLLSRINHFEDI
QIIPKSSWSSHEASLGVSACPYQGKSSFFRNVVWLIKKNSTYPTIKRSYNNNTNQEDLLVLWGIH
HPNDAAEQTKLYQNPTTYISVGTSTLNQRLVPRIATRISKVNGQSGRMEFFWTILKPNDAINFESN
GNFIAPEYAYKIVKKG DSTIMKSELEYGNCNTKCQTPMGAINSSMPFHNIHPLTIGECPKYVKS
RLVLATGLRNSPQRERRRKT RGLFGAIAGFIEGGWQGMVDGWYGYHHSNEQSGYAADKESTQKA
25 IDGVTNKVNSIIDKMNTQFEAVGREFNLERRIENLNKKMEDGFLDVWTYNAELLVLMENERTLD
FHDSNVKNLYDKVRLQLRDNKELGNGCFEFYHKCDNECMESVRNGTYDYPQYSEEARLKREEIS
GVKLESIGIYQILSIYSTVASSLALAIMVAGLSLWMCSNGSLQCRICI

SEQ ID NO: 228: H5 mini VN/1203/2004

MEKIVLLFAIVSLVKSDQICIGYHANNSTEQVDTIMEKNVTVTTHAQDILEKGGGGKYVCSNRLVL
30 ATGLRNKPQKESQGLFGAIAGFTEGGWQGMVDGWYGYHHSNEQSGYAADKESTQKAIDGVTNKV
NSIIDKMNTQYEAIGCEYNNSECMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKN
LYDKVRLQLRDNKELGNGCFEFYHKCDNECMESVRNGTYDYPQYSEEARLKREEISGVKLESIG
IYQILSIYSTVASSLALAIMVAGLSLWMCSNGSLQCRICI

35

SEQ ID NO: 229: sH5 mini VN/1203/2004

DQICIGYHANNSTEQVDTIMEKNVTVTTHAQDILEKGGGGKYVCSNRLVLATGLRNKPQKESQGLF
 GAIAGFTEGGWQGMVDGWYGYHHSNEQSGYAADKESTQKAIDGVTNKVNSIIDKMNTQYEAIGC
 EYNNSERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNLYDKVRLQLRDNAKEL
 5 GNGCFEFYHKCDNECMESVRNGTYDYPQYSEEARLKREEISGVKLESIGIYQI

SEQ ID NO: 230: H5 mini VN/1203/2004+loop

MEKIVLLFAIVSLVKSDQICIGYHANNSTEQVDTIMEKNVTVTTHAQDILEKGGGGKYVCSNRLVL
 ATGLRNKPQKESQGLFGAIAGFTEGGWQGMVDGWYGYHHSNEQSGYAADKESTQKAIDGVTNKV
 10 NSIIDKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKN
 LYDKVRLQLRDNAKELGNGCFEFYHKCDNECMESVRNGTYDYPQYSEEARLKREEISGVKLESIG
 IYQILSIYSTVASSLALAIMVAGLSLWMCNSGSLQCRICI

SEQ ID NO: 231: sH5 mini VN/1203/2004+loop

DQICIGYHANNSTEQVDTIMEKNVTVTTHAQDILEKGGGGKYVCSNRLVLATGLRNKPQKESQGLF
 GAIAGFTEGGWQGMVDGWYGYHHSNEQSGYAADKESTQKAIDGVTNKVNSIIDKMNTQYTAIGC
 EYNKSERCMKQIEDKIEEIESKIWCYNAELLVLMENERTLDFHDSNVKNLYDKVRLQLRDNAKEL
 15 GNGCFEFYHKCDNECMESVRNGTYDYPQYSEEARLKREEISGVKLESIGIYQI

20 SEQ ID NO: 232: FL HA H9N2 A/Hong_Kong/69955/2008

METVSLITMLLVATVINADKICIGYQSTNSTETVDTLTENNVPVTHAKELLHTEHNGMLCATNLG
 YPLILDCTIEGLIYGNPSCDLLLGGREWSYIVERPSAVNGLCYPGNVENLEELRSLFSSASSYQ
 RIQIFPDTIWNVSYSGTSKACSDSFYRSMRWLTQKNNAYPIQDAQYTNNREKNILFMWGINHPPT
 DTAQTNLTYTRDTTTSVATEEINRTFKPLIGRPLVNGLQGRIDYYWSVLKPGQTLRIRSNGNLI
 25 APWYGHILSGESHGRILKTDLKRGSCTVQCQTEKGGLNTTLPFQNVSKYAFGNCSKYVGVKSLKL
 AVGLRNVPSRSSRGLFGAIAGFIEGGWSGLVAGWYGFQHSNDQGVGMAADRSTQKAIDKITSKV
 NNIVDKMNKQYEIIDHEFSEVETRLNMINNKIDDQIQDIWAYNAELLVLLNQKTLDEHDANVNN
 LYNKVKRALGSNAVEDGKGCFELYHKCDDQCMETIRNGTYNRRRYQEESKLERQKIEGVKLESEG
 TYKILTIYSTVASSLVIAMGFAAFLFWAMNSGSCRCNICI

30 SEQ ID NO: 233: H9 mini HK/69955/2009

METVSLITMLLVATVINADKICIGYQSTNSTETVDTLTENNVPVTHAKELLHGGGGKYVCVKSLLK
 LAVGLRNKPSKSSQGLFGAIAGFTEGGWSGLVAGWYGFQHSNDQGVGMAADRSTQKAIDKITSK
 VNNIVDKMNKQYEIIDCEYSESERCMKQIEDKIEEIESKIWCYNAELLVLLNQKTLDEHDANVN
 NLYNKVKRALGSNAVEDGKGCFELYHKCDDQCMETIRNGTYNRRRYQEESKLERQKIEGVKLESE
 35 GTYKILTIYSTVASSLVIAMGFAAFLFWAMNSGSCRCNICI

SEQ ID NO: 234: sH9 mini HK/69955/2009

DKICIGYQSTNSTETVDTLTENNVPVTHAKELLHGGGGKYVCVKSLLAVGLRNKPSKSSQGLFG
AIAGFTEGGWSGLVAGWYGFQHSNDQGVGMAARDSTQKAIDKITSKVNNIVDKMKNQYEIIDCE
YSESERCMKQIEDKIEEIESKIWCYNAELLVLENQKTLDEHDANVNNLYNKVKRALGSSNAVEDG
5 KGCFELYHKCDDQCMETIRNGTYNRRRYQEESKLERQKIEGVKLESEGTYKI

SEQ ID NO: 235: H9 mini HK/69955/2009+loop

METVSLITMLLVATVINADKICIGYQSTNSTETVDTLTENNVPVTHAKELLHGGGGKYVCVKSLL
LAVGLRNKPSKSSQGLFGAIAGFTEGGWSGLVAGWYGFQHSNDQGVGMAARDSTQKAIDKITSK
10 VNNIVDKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLENQKTLDEHDANVN
NLYNKVKRALGSSNAVEDGKGCFELYHKCDDQCMETIRNGTYNRRRYQEESKLERQKIEGVKLESE
GTYKILTIYSTVASSLVIAMGFAAFLFWAMSNNGSCRCNICI

SEQ ID NO: 236: sH9 mini HK/69955/2009+loop

15 DKICIGYQSTNSTETVDTLTENNVPVTHAKELLHGGGGKYVCVKSLLAVGLRNKPSKSSQGLFG
AIAGFTEGGWSGLVAGWYGFQHSNDQGVGMAARDSTQKAIDKITSKVNNIVDKMNTQYTAIGCE
YNKSERCMKQIEDKIEEIESKIWCYNAELLVLENQKTLDEHDANVNNLYNKVKRALGSSNAVEDG
KGCFELYHKCDDQCMETIRNGTYNRRRYQEESKLERQKIEGVKLESEGTYKI

20

SEQ ID NO: 237: Full length HA H3N2 A/Hong Kong/1/1968

MKTIIALSIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSSSTG
KICNNPHRILDGIDCTLIDALLGDPHCDVFQNETWDLFVERSKAFSNCYPYDVPDYASLRSLVAS
SGTLEFITEGFTWTGVTQNGGSNACKRGP GSGFFSRLNWLTKSGSTYPVLNVTMPNNDNFDKLYI
25 WGVHHPSTNQEQTSLYVQASGRVTVSTRRSQQTII PNIGSRPWVRGLSSRISYWTIVKPGDVLV
INSNGNLIAPRGYFKMRTGKSSIMRSDAPIDTCISECITPNGSIPNDKPFQNVNKITYGACPKYV
KQNTLKLATGMRNVPEKQTRGLFGAIAGFIENGWEGMIDGWYGF RHQNSEGTGQAADLKSTQAAI
DQINGKLN RVIEKTNEKFHQIEKEFSEVEGRIQDLEKYVEDTKIDLWSYNAELLVALENQHTIDL
TDSEMKNLFEKTRRQLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQIKG
30 VELKSGYKDWILWISFAISCFLLCVLLGFIMWACQRGNIRCNICI

SEQ ID NO: 238: #2999 H3 HK68 mini2-cl9+10+11+12-GCN4t

MKTIIALSIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGF RHQNSEGTGQAADLKSTQ
35 AAIDQINGKLN RVREKTNEKSHQTEKESSEGEGRMKQIEDKIEEIESKLCYNAELLVALENQHT

IDLTDSEMNKLF EKTRRQLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
 IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQ RGNIRCNICI

SEQ ID NO: 239: #3801 H3 HK mini2a-linker+c19 +10+11+12+GCN4T-
 CG7-1

5 MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
 KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
 AAIDQINGKLN RVREKTNEKSHQTEKES SNATGRMKQIEDKIEEIESKLWCYNAELLVALENQHT
 IDLTDSEMNKLF EKTRRQLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
 IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQ RGNIRCNICI

10 SEQ ID NO: 240: #2999-c114

MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
 KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
 AAIDQINGKLN RVREKTNEKSHQTEKESSEGEGRMKQCCDKIEEIESKLWCYNAELLVALENQHT
 IDLTDSEMNKLF EKTRRQLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
 15 IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQ RGNIRCNICI

SEQ ID NO: 241: #2999 c115

MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
 KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
 AAIDQINGKLN RVREKTNEKSHQTEKESSEGEGRMKQIEDKIEECCSKLWCYNAELLVALENQHT
 20 IDLTDSEMNKLF EKTRRQLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
 IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQ RGNIRCNICI

SEQ ID NO: 242: #2999 c117

MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
 KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
 25 AAIDQINGKLN RVREKTNECSHQTEKESSEGEGRMKQIEDKIECIESKLWCYNAELLVALENQHT
 IDLTDSEMNKLF EKTRRQLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
 IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQ RGNIRCNICI

SEQ ID NO: 243: #2999 c118

MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
 KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
 30 AAIDQINGKLN RVREKTNEKSHQTECESSEGE GCMKQIEDKIEEIESKLWCYNAELLVALENQHT
 IDLTDSEMNKLF EKTRRQLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
 IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQ RGNIRCNICI

SEQ ID NO: 244: #2999 c130

35 MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
 KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ

AAIDQINGKLN RVREKTNEKSHCTEKESSEGEGRMKQIEDCIEEIESKLWCYNAELLVALENQHT
IDLTDSEMNKLF EKTRRLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQRGNIRCNICI

SEQ ID NO: 245: #2999 c131

5 MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
AAIDQINGKLN RVREKTNEKSHQCEKESSEGEGRMKQIEDCIEEIESKLWCYNAELLVALENQHT
IDLTDSEMNKLF EKTRRLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQRGNIRCNICI

10

SEQ ID NO: 246: #3801-c114

MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
AAIDQINGKLN RVREKTNEKSHQTEKESSEGEGRMKQCCDKIEEIESKLWCYNAELLVALENQHT
15 IDLTDSEMNKLF EKTRRLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQRGNIRCNICI

SEQ ID NO: 247: #3801 c115

MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
20 AAIDQINGKLN RVREKTNEKSHQTEKESSEGEGRMKQIEDKIEECCSKLWCYNAELLVALENQHT
IDLTDSEMNKLF EKTRRLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQRGNIRCNICI

SEQ ID NO: 248: #3801 c117

MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
25 AAIDQINGKLN RVREKTNEKSHQTEKESSEGEGRMKQIEDKIEECIESKLWCYNAELLVALENQHT
IDLTDSEMNKLF EKTRRLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQRGNIRCNICI

SEQ ID NO: 249: #3801 c118

MKTIIALS YIFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
30 AAIDQINGKLN RVREKTNEKSHQTECESSGEGRMKQIEDKIEEIESKLWCYNAELLVALENQHT
IDLTDSEMNKLF EKTRRLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQRGNIRCNICI

35 SEQ ID NO: 250: #3801 c130

MKTIIALSIFYFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
 KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
 AAIDQINGKLNVRREKTNEKSHCTEKESS**NAT**GRMKQIEDCIEEIESKLWCYNAELLVALENQHT
 IDLTDSEMNKLFKTRRQLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
 5 IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQRGNIRCNICI

SEQ ID NO: 251: #3801 c131

MKTIIALSIFYFCLALGQDLPGNDNSTATLCLGHHAVPNGTLVKTITDDQIEVTNATELVQSGGGG
 KYVCQNTLKLATGMRNVPEKQTQGLFGAIAGFIENGWEGMIDGWYGFRHQNSEGTGQAADLKSTQ
 AAIDQINGKLNVRREKTNEKSHQCEKESS**NAT**GRMKQIEDCIEEIESKLWCYNAELLVALENQHT
 10 IDLTDSEMNKLFKTRRQLRENAEDMGNGCFKIYHKCDNACIESIRNGTYDHDVYRDEALNNRFQ
 IKGVELKSGYKDWILWISFAISCFLLCVVLLGFIMWACQRGNIRCNICI

SEQ ID NO: 252: _FL HA H1N1 A/California/07/2009

MKAILVLLYTFATANADTLICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLEDKHNGKLCKLRGVA
 15 PLHLGKCNIAGWILGNPECESLSTASSWSYIVETPSSDNGTCYPGDFIDYEELREQLSSVSSFER
 FEIFPKTSSWPNHDSNKGVTAACPHAGAKSFYKNLIWLVKKNSYPKLSKSYINDKGKEVLVLWG
 IHPSTADQQSLYQNADAYVFGSSRSYKFKPEIAIRPKVRDQEGRMNYYWTLVEPGDKITFE
 ATGNLVVPRYAFAMERNAGSGIIISDTPVHDCNTTCQTPKGAINSTLPPFQNIHPITIGKCPKYVK
 STKLRLATGLRNIPSIQSRGLFGAIAGFIEGGWTGMVDGWYGYHHQNEQGSGYAADLKSTQNAID
 20 EITNKVNSVIEKMNTQFTAVGKEFNHLEKRIENLNKKVDDGFLDIWTYNAELLVLENERLTLDYH
 DSNVKNLYEKVRSQLKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGV
 KLESTRIYQILAIYSTVASSLVLVSLGAISFWMCNSGSLQCRICI

SEQ ID NO: 253 H1 mini-HA 127H1-t2 S73L

MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
 25 MVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
 VNSVIEKMNTQYTAIGCEYNKLERCMKQIEDKIEEIESKIWCYNAELLVLENERLTDFHDSNVK
 NLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
 GYVQILAIYSTVASSLVLLVSLGAISFWMCNSGSLQCRICI

SEQ ID NO: 254 sH1 mini-HA 127H1-t2longS73L (#5114)

DTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLRMVTGLRNKPSKQSQGLF
 GAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNKVNSVIEKMNTQYTAIGC
 EYNKLERCMKQIEDKIEEIESKIWCYNAELLVLENERLTDFHDSNVKNLYEKVKSQKNNAKEI
 35 GNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQI

SEQ ID NO: 255: UFV150124

MKVKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLLENGGGGKYVCSAKLR
 MVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQGSGYAADQKSTQNAINGITNK
 40 VNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLENERLTDFHDSNVK
 NLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVSGRDY
 KDDDDKPGSGYIPEAPRDGQAYVRKDGEWVLLSTFLGHHHHHH

SEQ ID NO: 256: UFV150125

MKVKLLVLLCTFTATYADTICIGYHANNSTDVDTVLEKNVTVTTHSVNLLLENGGGGKYVCSAKLR
MVTGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADQKSTQNAINGITNK
VNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDFHDSNVK
5 NLYEKVKSQKNNAKEIGNGCFEFYHKCNDECMESVKNGTYDYPKYSEESKLNREKIDGVKLESM
GVYQIEGRAAADYKDDDDKPGSGYIPEAPRDGQAYVRKDGEWVLLSTFLGHHHHHH

SEQ ID NO: 257: UFV150134

MKAILVLLLYTFATANADTLCIGYHANNSTDVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLR
10 LATGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDYHDSNVK
NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGVSGRDY
KDDDDKPGSGYIPEAPRDGQAYVRKDGEWVLLSTFLGHHHHHH

SEQ ID NO: 258: UFV150135

MKAILVLLLYTFATANADTLCIGYHANNSTDVDTVLEKNVTVTTHSVNLLLEDGGGGKYVCSTKLR
15 LATGLRNKPSKQSQGLFGAIAGFTEGGWTGMVDGWYGYHHQNEQSGYAADLKSTQNAIDEITNK
VNSVIEKMNTQYTAIGCEYNKSERCMKQIEDKIEEIESKIWCYNAELLVLLLENERTLDYHDSNVK
NLYEKVRSQKNNAKEIGNGCFEFYHKCDNTCMESVKNGTYDYPKYSEEAKLNREEIDGVKLEST
20 RIYQIEGRAAADYKDDDDKPGSGYIPEAPRDGQAYVRKDGEWVLLSTFLGHHHHHH

CLAIMS

1. A multimeric influenza hemagglutinin stem domain polypeptide, wherein the multimeric influenza hemagglutinin stem domain polypeptide comprises at least a first and a second influenza hemagglutinin stem domain monomer, said first and second monomer each comprising: (a) an influenza hemagglutinin HA1 domain that comprises an HA1 amino (N)-terminal stem segment, wherein said HA1 N-terminal segment comprises the amino acids 1-52 of HA1, preferably the amino acids 18-52 of HA1, covalently linked by a linking sequence of less than 10 residues to an HA1 carboxy (C)-terminal stem segment, comprising the amino acids from position 321 to the C-terminal amino acid of HA1, wherein said HA1 C-terminal segment is linked to (b) an influenza hemagglutinin HA2 domain, and
 - (c) wherein the polypeptide comprises no protease cleavage site at the junction between the HA1 and HA2 domains; and
 - (d) wherein the first monomer is linked to said second monomer by a disulfide bridge between the amino acid on position 411 of the first monomer and the amino acid on position 419 of the second monomer, and wherein in each influenza hemagglutinin stem domain monomer the region comprising the amino acid residues 402-418 comprises the amino acid sequence $X_1NTQX_2TAX_3GKEX_4N(H/K)X_5E(K/R)$ (SEQ ID NO: 8), wherein X_1 is M, X_2 is Y, X_3 is I, X_4 is Y and X_5 is S; and wherein in each influenza hemagglutinin stem domain monomer the amino acid sequence CMKQIEDKIEEIESK (SEQ ID NO: 193) has been introduced at positions 419-433; and wherein the amino acid residue on position 337 is K; the amino acid residue on position 340 is K, the amino acid residue on position 352 is F and the amino acid residue on position 353 is T; wherein the HA1 and HA2 domains are derived from an influenza A virus subtype derived from phylogenetic group 1, preferably from an influenza A virus subtype comprising HA of the H1 subtype, and wherein the numbering of the amino acid positions is in relation to SEQ ID NO: 1.
2. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 1, wherein the multimeric polypeptide is trimeric.

3. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 1, wherein the HA2 domain of each influenza hemagglutinin stem domain monomer has been truncated.
4. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 3, wherein the C-terminal part of the HA2 domain from position 519 to the C-terminal amino acid has been deleted.
5. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 3, wherein the C-terminal part of the HA2 domain from position 530 to the C-terminal amino acid has been deleted.
6. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 3, wherein the C-terminal part of the HA2 domain has been replaced by the amino acid sequence GYIPEAPRDGQAYVRKDGEWVLLSTFL (SEQ ID NO: 3), optionally connected through a linker.
7. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 1, wherein the C-terminal amino acid residue of the HA1 C-terminal stem segment of each influenza hemagglutinin stem domain monomer is any amino acid other than arginine (R) or lysine (K), preferably glutamine (Q).
8. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 1, wherein the polypeptide comprises one or more further mutations in the HA1 domain and/or the HA2 domains.
9. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 1, wherein the polypeptide selectively binds to the antibodies CR6261 and/or CR9114.
10. A nucleic acid molecule encoding the polypeptide of claim 1.

11. A vector comprising the nucleic acid molecule of claim 10.
12. A composition comprising the polypeptide according to claim 1 and a pharmaceutically acceptable carrier.
13. A composition comprising the nucleic acid molecule according to claim 10 and a pharmaceutically acceptable carrier.
14. The polypeptide according to claim 1 for use in inducing an immune response against influenza virus in a subject in need thereof, wherein the polypeptide is to be administered to the subject.
15. Use of the polypeptide according to claim 1 in the manufacture of a medicament for inducing an immune response against influenza virus in a subject in need thereof, wherein the medicament is to be administered to the subject.
16. The nucleic acid molecule according to claim 10 for use in inducing an immune response against influenza virus in a subject in need thereof, wherein the nucleic acid molecule is to be administered to the subject.
17. Use of the nucleic acid molecule according to claim 10 in the manufacture of a medicament for inducing an immune response against influenza virus in a subject in need thereof, wherein the medicament is to be administered to the subject.
18. The vector according to claim 11 for use in inducing an immune response against influenza virus in a subject in need thereof, wherein the medicament is to be administered to the subject.
19. Use of the vector according to claim 11 in the manufacture of a medicament for inducing an immune response against influenza virus in a subject in need thereof, wherein the medicament is to be administered to the subject.

20. A composition comprising the vector according to claim 11 and a pharmaceutically acceptable carrier.

21. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 1, wherein each polypeptide monomer comprises the amino acid sequence:

DTICIGYHANNSTDVDTVLEKNVTVTHTSVNLLENG GGGK YVCSAKLRMVTGL
 RNX₁PSX₂QSQGLFGAIAGX₃X₄EGGWTGMVDGWYGYHHQNEQSGSYAADQKS
 TQNAINGITNKVNSVIEKX₅NTQX₆TAX₇GCEX₈NKX₉ERCMKQIEDKIEEIESKIWC
 YNAELLVLLNERTLDFHDSNVKNLYEKVKSQLKNNAKEIGNGC FEFYHKCNDE
 CMESVKNGTYDYPKYSEESKLNREKIDG (SEQ ID NO: 146), wherein X₁ is K, X₂ is K, X₃ is F, X₄ is T, X₅ is M, X₆ is Y, X₇ is I, X₈ is Y and X₉ is S.

22. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 1, wherein each polypeptide monomer comprises the amino acid sequence:

DTICIGYHANNSTDVDTVLEKNVTVTHTSVNLLENG GGGK YVCSAKLRMVTGL
 RNX₁PSX₂QSQGLFGAIAGX₃X₄EGGWTGMVDGWYGYHHQNEQSGSYAADQKS
 TQNAINGITNKVNSVIEKX₅NTQX₆TAX₇GCEX₈NKX₉ERCMKQIEDKIEEIESKIWC
 YNAELLVLLNERTLDFHDSNVKNLYEKVKSQLKNNAKEIGNGC FEFYHKCNDE
 CMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQIEG (SEQ ID NO: 147),
 wherein X₁ is K, X₂ is K, X₃ is F, X₄ is T, X₅ is M, X₆ is Y, X₇ is I, X₈ is Y and X₉ is S.

23. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 1, wherein each polypeptide monomer comprises the amino acid sequence:

DTICIGYHANNSTDVDTVLEKNVTVTHTSVNLLENG GGGK YVCSAKLRMVTGL
 RNX₁PSX₂QSQGLFGAIAGX₃X₄EGGWTGMVDGWYGYHHQNEQSGSYAADQKS
 TQNAINGITNKVNSVIEKX₅NTQX₆TAX₇GCEX₈NKX₉ERCMKQIEDKIEEIESKIWC
 YNAELLVLLNERTLDFHDSNVKNLYEKVKSQLKNNAKEIGNGC FEFYHKCNDE
 CMESVKNGTYDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVS
 LGAISFWMC SNGSLQCRICI (SEQ ID NO: 148), wherein X₁ is K, X₂ is K, X₃ is F,
 X₄ is T, X₅ is M, X₆ is Y, X₇ is I, X₈ is Y and X₉ is S.

24. The multimeric influenza hemagglutinin stem domain polypeptide according to claim 1, wherein each polypeptide monomer comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 164, 186, 202, 203, 207-216, 218-226, and 254-258.