

Description

The present invention relates to a novel derivative of bornane-3-spiro-1'-cyclopentane and a perfumery composition containing it.

In recent years, synthetic perfumes have been given much weight as the base material for perfumeries getting rid of dependence on the natural resources common in the prior art in order to satisfy various requirements such as stationary supply, invariability in the quality, inexpensiveness and the like.


Notwithstanding the above, natural perfumes are still given much weight concerning the woody note scent so that supply of synthetic substances with woody note scent is wished in consideration of the increase of demand in the future.

However, difficulties are posed in many of the cases of industrial production of the compounds with woody note scent since many of such compounds have a polycyclic structure with complexity as typically represented by sesquiterpenes. In particular, synthesis of tricyclic compounds was not easy among such polycyclic compounds.

EP-A-208 994 describes bornane-3-spiro-1'-cyclopentanes and perfume compositions comprising the same. The compounds are synthesised from camphor and have woody-note.

SUMMARY OF THE INVENTION

The present invention is a compound with woody note scent useful as a basic component for formulating various types of perfumeries which is a derivative of bornane-3-spiro-1'-cyclopentane represented by the general formula




in which R_1 represents a hydrogen atom or a hydroxyl group and R_2 represents a methoxyl group, an ethoxyl group, a formyloxyl group or an acetoxyl group in the cases in which R_1 is a hydrogen atom or represents a methyl group or an ethyl group in the cases in which R_1 is a hydroxyl group.

The present invention also includes a perfumery composition containing the above derivative of bornane-3-spiro-1'-cyclopentane.

DETAILED DESCRIPTION OF THE INVENTION

In view of the above-described present status, the inventors have successfully completed the present invention by finding a novel compound with distinctive scent prepared by using camphor in a stationary condition in the price and resources as the starting compound as the consequence of extensive studies based on a conception that the tricyclic compounds would be synthesized with a considerable easiness when bicyclo-monoterpenes with a great availability are used in the synthesis.


That is, the present invention provides a derivative of bornane-3-spiro-1'-cyclopentane represented by the formula



in which R_1 represents a hydrogen atom or a hydroxyl group, and R_2 represents a methoxyl group, an

ethoxyl group, a formyloxyl group or an acetoxyl group in the cases in which R_1 is a hydrogen atom or represents a methyl group or an ethyl group in the cases in which R_1 is a hydroxyl group, and the present invention also provides a perfumery composition containing the above compound.

The compound (I) of the invention is manufactured according to the following process using bornane-3-spiro-1'-cyclopentan-2-one (II) (hereinafter referred to as a ketone derivative) or bornane-3-spiro-1'-cyclopentan-2-ol (III) (hereinafter referred to as an alcoholic derivative) prepared from camphor by the method described for example in Japan Kokai Tokkyo Koho No. 62-16442.



(X represents a halogen atom in the formula)

The compounds of the formula (I) in which R_1 represents a hydrogen atom are prepared by any of the following methods including:

- a method of obtaining ether derivatives (Ia) or (Ib) by the ether synthesis of Williamson in which an alkoxide obtained from an alcoholic derivative (III) by a treatment with sodium hydride is reacted with a halide such as methyl iodide, ethyl iodide and the like;
- a method of obtaining a formate ester (Ic) by the reaction of an alcoholic derivative (III) with an excessive amount of formic acid at room temperature; and
- a method of obtaining an acetic ester (Id) by the reaction of an alcoholic derivative (III) with an acetylating agent in the presence of a base.

Also, the compounds of the formula (I) in which R_1 represents a hydroxyl group are prepared by the

method of obtaining (Ie) or (If) by the reaction of a ketone derivative (II) with a lithium reagent carried out in accordance with the method described in the book "Shin Jikken Kagaku Koza (New Lecture Courses of Experimental Chemistry), Vol. 14", pages 515 to 520.

The compound of formula (I) is obtained as a mixture of two distinctive species ----- different in the steric configuration of the substituting groups in the position 3 when prepared by way of one of the procedures of the above.

The compounds (I) of the invention emit distinctive scent different from that of the ketone derivatives (II) or the alcoholic derivatives (III) by the introduction of the group R₂ with simultaneous decrease of volatility and improvement in lingering of fragrance. That is, the compounds of (Ia) and (Ib) hardly emit amber-like odor characteristic in the alcoholic derivatives (III) giving grassy, metallic, moldy woody or faintly resinous note scent by the compounds (Ia) or slightly woody, herbal or ether-like note scent by the compounds (Ib). The compounds (Ic) and (Id) do not emit fragrance of cedarwood note characteristic in the alcoholic derivatives (III) altogether giving amber-grease note or woody note scent by the compounds (Ic) or sweetishly woody, camphoraceous or faintly flower-like note scent by the compounds (Id). Furthermore, the compounds (Ie) and (If) do not emit camphoraceous note characteristic in the ketone derivatives (II) giving earthy, metallic woody note or patchouli alcohol-like scent by the compounds (Ie) or slightly woody, amber-like or faintly earthy note scent by the compounds (If).

As mentioned in the above, each of the compounds of the invention emits a characteristic odor based on the woody note scent so that they can be widely used in the products which require perfuming such as high class perfumery compositions, liquid perfumes, soap, shampoo, hair rinse, detergents, cosmetics, atomizable spray agents, aromatics and the like as used alone or as a basic material for formulation of various perfumeries.

The invention is further explained referring to the examples in the following.

Example 1

Synthesis of spiro[2-methoxybornane-3,1'-cyclopentane] (Ia):


A solution consisting of 1.2 g (0.03 moles) of sodium hydride and 10 ml of xylene is added dropwise with a solution consisting of 5 g (0.024 moles) of spiro[bornane-3,1'-cyclopentan-2-ol] and 5 ml of xylene at 100°C while being stirred until the cease of hydrogen gas evolution. Thereafter, a mixture solution consisting of 5 g (0.035 moles) of methyl iodide and 5 ml of xylene is added to the solution being stirred dropwise within 15 minutes followed by continual stirring for 3 hours while the solution is kept at 100°C. After being cooled to room temperature, the solution is subjected to phase separation with added water and furthermore to washing 3 times with water. A purified product is obtained by fractional distillation under a reduced pressure following to removal of the solvent by distillation.

amount of the product: 5 g (yield 93.7%)

IR (liquid thin layer, cm⁻¹): 1120 (ν_{C-O})

¹H-NMR (CDCl₃ solvent, TMS internal standard, δ ppm):

3.30 (singlet, -OCH₃)

2.53 (singlet, )

2.38 to 1.88 (broad multiplet, )

1.85 to 1.20 (complicated multiplet, -CH₂-, 12H)

1.01 (singlet, CH₃)

0.85 (singlet, CH₃)

0.79 (singlet, CH₃)

MS [m/e (relative intensity)] :

222(M⁺, 22), 177(67), 113(34), 112(100), 108(35), 95(53), 85(83), 81(28), 55(28), 41(33)

elementary analysis:

Found	C; 81.19%,	H; 11.83%
Calculated	C; 81.02%,	H; 11.78%

Example 2

Synthesis of spiro[2-ethoxybornane-3,1'-cyclopentane] (Ib) :


A solution consisting of 1.2 g (0.03 moles) of sodium hydride and 10 ml of xylene is added dropwise with a solution consisting of 5 g (0.024 moles) of spiro[bornane-3,1'-cyclopentan-2-ol] and 5 ml of xylene at 100 °C while being stirred until the cease of hydrogen gas evolution. Thereafter, the above reaction mixture is added with a solution consisting of 5 g (0.032 moles) of ethyl iodide and 5 ml of xylene while being stirred at 100 °C within 15 minutes followed by being kept at the same temperature for 3 hours with continued stirring. After being cooled to room temperature, the solution is subjected to phase separation with added water and furthermore to washing 3 times with water. A purified product is obtained by fractional distillation under a reduced pressure following to removal of the solvent by distillation.

amount of the product: 5.1 g (yield 89.9%)

IR (liquid thin layer, cm^{-1}) : 1120 ($\nu_{\text{C-O}}$)

$^1\text{H-NMR}$ (CDCl_3 solvent, TMS internal standard, δ ppm) :

3.40 (quartet, $J = 7.0$ Hz, $-\text{OCH}_2-$)

2.63 (singlet, )

2.45 to 1.88 (broad multiplet, )

1.85 to 1.24 (complicated multiplet, $-\text{CH}_2-$, 12H)

1.12 (triplet, $J = 7.2$ Hz, $-\text{OCH}_2-\text{CH}_3$)

1.03 (singlet, CH_3)

0.83 (singlet, CH_3)

0.78 (singlet, CH_3)

MS [m/e (relative intensity)] :

236(M^+ , 23), 177(77), 153(35), 127(41), 126(72), 108(46), 99(100), 95(53), 71(40), 41(44)

elementary analysis :

Found	C; 81.35%,	H; 12.10%
Calculated	C; 81.29%,	H; 11.94%

Example 3

Synthesis of spiro [2-formyloxybornane-3,1'-cyclopentane] (Ic) :

A solution consisting of 20 g (0.087 moles) of spiro[bornane-3,1'-cyclopentan-2-ol] and 200 ml of formic acid is stirred for 3 hours while being chilled using ice water. The solution is subjected to phase separation with added ethyl ether and water followed by successive washing treatments of the organic phase using a saturated aqueous solution of sodium hydrogen carbonate and water in this order. After desiccation of the organic phase using anhydrous magnesium sulfate, the solid matter is separated by filtration and the filtrate is subjected to distillation under a reduced pressure for removal of the solvent. The residue is purified by column chromatography (the developing solvent: n-hexane/ethyl acetate = 30/1) and a purified product is

obtained after removal of the solvent by distillation.

amount of the product: 9 g (yield 43.8%)

IR (liquid thin layer, cm^{-1}):


1730 ($\nu_{\text{C=O}}$), 1160, 1180 ($\nu_{\text{C-O}}$)

$^1\text{H-NMR}$ (CDCl_3 solvent, TMS internal standard, δ ppm):

8.12 (singlet, $\text{O} - \text{C} - \text{H}$)



4.55 (singlet, OCH)



2.18 to 1.22 (complicated multiplet,

$(-\text{CH}_2-, 12\text{H})(-\text{CH}-, 1\text{H})$)

1.08 (singlet, CH_3)

0.82 (singlet, CH_3)

0.78 (singlet, CH_3)

MS [m/e (relative intensity)]:

236(M^+ , 1), 110(31), 147(35), 41(59), 95(100)

elementary analysis:

Found	C; 75.95%,	H; 10.42%
Calculated	C; 76.23%,	H; 10.23%

Example 4

Synthesis of spiro[2-acetoxymorphane-3,1'-cyclopentane] (Id):

A solution consisting of 5 g (0.024 moles) of spiro[morphane-3,1'-cyclopentan-2-ol], 5 g of acetic anhydride and 5 ml of pyridine is stirred for 3 hours at room temperature. The solution is subjected to phase separation with added ethyl ether and water followed by successive washing treatments of the organic phase using 1N hydrochloric acid, a saturated aqueous solution of sodium hydrogen carbonate and water in this order. After desiccation of the organic phase using anhydrous magnesium sulfate, the solid matter is separated by filtration and a purified product is obtained from the filtrate subjected to removal of the solvent by distillation under a reduced pressure.


amount of the product: 6 g (yield 95.4%)

IR (liquid thin layer, cm^{-1}):

1745 ($\nu_{\text{C=O}}$), 1242 ($\nu_{\text{C-O}}$)

$^1\text{H-NMR}$ (CDCl_3 solvent, TMS internal standard, δ ppm):

4.46 (singlet, OCH_3)



2.03 (singlet, $-\text{OC}-\text{CH}_3$)



1.90 to 1.20 (complicated multiplet, 13H)

1.08 (singlet, CH₃)

0.80 (singlet, CH₃)

0.75 (singlet, CH₃)

5 MS [m/e (relative intensity)]:

250(M⁺, 1), 190(50), 175(37), 147(47), 140(38), 108(62), 98(100), 95(62), 43(80), 41(43)

elementary analysis:

Found	C; 77.01%,	H; 10.23%
Calculated	C; 76.75%,	H; 10.47%

15 Example 5

Synthesis of spiro[2-methylbornane-3,1'-cyclopentan-2-ol] (Ie):

20 A solution consisting of 10 g (0.048 moles) of spiro[bornane-3,1'-cyclopentan-2-one] and 10 ml of desiccated ethyl ether is refrigerated to -78°C under an inert gas atmosphere. The solution is added with 39 ml (0.062 moles) of methyl lithium (1.6 molar solution in ethyl ether) dropwise with stirring within 1 hour while being kept at the same temperature followed by temperature elevation back to room temperature. After continual stirring for 3 hours at room temperature, the solution is subjected to phase separation with added ethyl ether and water followed by successive washing treatments of the organic phase using a saturated aqueous solution of sodium hydrogen carbonate and then water with subsequent desiccation with anhydrous magnesium sulfate. By removal of the solvent by distillation, 10.4 g of a viscous liquid is obtained. A pure product is obtained from the liquid purified by column chromatography [the developing solvent: n-hexane (containing 3% of ethyl acetate)] with subsequent removal of the solvent by distillation under a reduced pressure.


30 amount of the product: 8 g (yield 74.2%)

IR (liquid thin layer, cm⁻¹): 3450 to 3690 (ν_{O-H})

¹H-NMR (CDCl₃ solvent, TMS internal standard, δ ppm):

2.3 (triplet, J = 7.0 Hz, >CH-)

1.79 to 1.27 (complicated multiplet, 12H)

35 1.26 (singlet, )

40 1.17 (singlet, CH₃)

0.86 (singlet, CH₃)

0.82 (singlet, CH₃)

45 MS [m/e (relative intensity)]:

222(M⁺, 1), 112(100), 95(49), 43(100), 41(92)

elementary analysis:

Found	C; 81.15%,	H; 11.53%
Calculated	C; 81.02%,	H; 11.78%

55 Example 6

Synthesis of spiro [2-ethylbornane-3,1'-cyclopentan-2-ol] (If):

A solution consisting of 1.3 g (0.056 moles) of lithium (30%, a mixture with liquid paraffin) and 10 ml of

desiccated ethyl ether is added with a solution consisting of 4.4 g (0.028 moles) of ethyl iodide, 5.0 g (0.024 moles) of spiro[2-bornane-3,1'-cyclopentan-2-one] and 20 ml of desiccated ethyl ether dropwise within 30 minutes. After being stirred for 30 minutes at room temperature, the solution is added with ice-chilled water, extracted with ethyl ether, washed successively with 10% sulfuric acid and water, desiccated with anhydrous magnesium sulfate and subjected to removal of the solvent by distillation thus to give 5.2 g of a viscous liquid. A pure product is obtained from the liquid purified by column chromatography (the developing solvent: n-hexane/ethyl acetate = 25/1) with subsequent removal of the solvent by distillation under a reduced pressure.

amount of the product: 4 g (yield 69.8%)


IR (liquid thin layer, cm^{-1}):

3450 to 3650 (broad absorption peak, $\nu_{\text{O-H}}$)

$^1\text{H-NMR}$ (CDCl_3 solvent, TMS internal standard, δ ppm):

0.53 to 0.96 (complicated multiplet, CH_3 , 9H)

1.07 (singlet, CH_3)

1.13 to 2.23 (complicated multiplet, $-\text{CH}_2-$, , 15H)

MS [m/e (relative intensity)]:

236(M^+ , 2), 126(100), 123(32), 109(36), 99(51), 97(60), 95(59), 69(43), 57(52), 41(36)

elementary analysis:

Found	C; 80.15%,	H; 12.05%
Calculated	C; 81.29%,	H; 11.94%

Example 7

	<u>forest note perfumery compositions</u>	parts by weight
5	lemon oil	50
	bergamot base	80
10	pine needle base	113
	aldehyde C-12MNA	2
	isobornyl acetate	50
15	p-t-butyl cyclohexyl acetate	50
	acetyl cedrene	30
20	far balsam absolute	5
	treemoss absolute 50% DEP ¹⁾	20
	citronellol	50
25	geranyol	100
	hexyl cinnamic aldehyde	120
30	methyl dihydro jasmonate	50
	Lyrar ²⁾	30
	Galaxolide 50 DEP ³⁾	150
35		
	total	900

40 1) 50% DEP: 50% solution in diethyl phthalate

2) Lyrar: a trade name for IFF Company

45 4-(4-hydroxy-4-methylpentyl)-3-cyclohexene-
1-carboxyaldehyde

50 3) Glaxolide 50 DEP:

a trade name for IFF Company 1,3,4,6,7,8-
hexahydro-4,6,6,7,8,8-hexamethyl-cyclopenta-
55 γ -2-benzopyran

By compounding 100 parts by weight of the compound (Ic) of the present invention with 900 parts by weight of the above perfume, a forest note perfumery composition rich in woody and amber notes and with a feeling of abundance was obtained.

Exampel 8

<u>perfumes for cologne for men</u>	parts by weight
bergamot oil Italy BGF ¹⁾	50
lemon oil Sunkist BGF	30
pine oil China	20
aromoase oil	10
basil oil	5
lavender oil	15
2,4-dimethyl-3-cyclohexene-1-carboxyaldehyde	5
jasmine base	40
patchouli oil	10
acetyl cedrene	200
oakmoss absolute 50% DEP ²⁾	30
Galaxolide 50 DEP ³⁾	80
amber base	20
thyme oil red	5
eugenol	20
dipropyleneglycol	260
total	800

1) BGF: bergaptene-free grade

2) 50% DEP: 50% solution in diethyl phthalate

3) Galaxolide 50 DEP:


a trade name for IFF company

1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethyl-
cyclopenta-γ-2-benzopyran

By compounding 200 parts by weight of the compound (Ie) of the present invention with 800 parts by weight of the above perfume, a perfumery for cologne for men with increased woody note rich in aromatic feeling was obtained.

Claims


1. A derivative of bornane-3-spiro-1'-cyclopentane represented by the general formula



(I)

in which R₁ represents a hydrogen atom or a hydroxyl group and R₂ represents a methoxyl group, an ethoxyl group, a formyloxyl group or an acetoxyl group in the cases in which said R₁ is a hydrogen atom or represents a methyl group or an ethyl group in the cases in which said R₁ is a hydroxyl group.

2. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R₁ is a hydrogen atom and said R₂ is a methoxyl group being spiro[2-methoxybornane-3,1'-cyclopentane].
3. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R₁ is a hydrogen atom and said R₂ is an ethoxyl group being spiro[2-ethoxybornane-3,1'-cyclopentane].
4. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R₁ is a hydrogen atom and said R₂ is a formyloxyl group being spiro[2-formyloxybornane-3,1'-cyclopentane].
5. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R₁ is a hydrogen atom and said R₂ is an acetoxyl group being spiro[2-acetoxybornane-3,1'-cyclopentane].
6. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R₁ is a hydroxyl group and said R₂ is a methyl group being spiro[2-methylbornane-3,1'-cyclopentan-2-ol].
7. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R₁ is a hydroxyl group and said R₂ is an ethyl group being spiro[2-ethylbornane-3,1'-cyclopentan-2-ol].
8. A perfumery composition comprising a derivative of bornane-3-spiro-1'-cyclopentane represented by the general formula



(I)

in which R₁ represents a hydrogen atom or a hydroxyl group and R₂ represents a methoxyl group, an




1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethyl-
cyclopenta-γ-2-benzopyran

By compounding 200 parts by weight of the compound (Ie) of the present invention with 800 parts by weight of the above perfume, a perfumery for cologne for men with increased woody note rich in aromatic feeling was obtained.

Claims


1. A derivative of bornane-3-spiro-1'-cyclopentane represented by the general formula



(I)

in which R_1 represents a hydrogen atom or a hydroxyl group and R_2 represents a methoxyl group, an ethoxyl group, a formyloxyl group or an acetoxy group in the cases in which said R_1 is a hydrogen atom or represents a methyl group or an ethyl group in the cases in which said R_1 is a hydroxyl group.

2. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R_1 is a hydrogen atom and said R_2 is a methoxyl group being spiro[2-methoxybornane-3,1'-cyclopentane].
3. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R_1 is a hydrogen atom and said R_2 is an ethoxyl group being spiro[2-ethoxybornane-3,1'-cyclopentane].
4. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R_1 is a hydrogen atom and said R_2 is a formyloxyl group being spiro[2-formyloxybornane-3,1'-cyclopentane].
5. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R_1 is a hydrogen atom and said R_2 is an acetoxy group being spiro[2-acetoxybornane-3,1'-cyclopentane].
6. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R_1 is a hydroxyl group and said R_2 is a methyl group being spiro[2-methylbornane-3,1'-cyclopentan-2-ol].
7. The derivative of bornane-3-spiro-1'-cyclopentane according to Claim 1, wherein said R_1 is a hydroxyl group and said R_2 is an ethyl group being spiro[2-ethylbornane-3,1'-cyclopentan-2-ol].
8. A perfumery composition comprising a derivative of bornane-3-spiro-1'-cyclopentane represented by the general formula



(I)


in which R_1 represents a hydrogen atom or a hydroxyl group and R_2 represents a methoxyl group, an



ethoxyl group, a formyloxyl group or an acetoxyl group in the cases in which said R_1 is a hydrogen atom or represents a methyl group or an ethyl group in the cases in which said R_1 is a hydroxyl group.

Revendications


1. Dérivé de bornane-3-spiro-1'-cyclopentane, représenté par la formule générale :



(I)

dans laquelle :

- R_1 représente un atome d'hydrogène ou un groupe hydroxyle ; et
 - R_2 représente un groupe méthoxyle, un groupe éthoxyle, un groupe formyloxyle ou un groupe acétoxyle dans les cas où ledit R_1 est un atome d'hydrogène ou bien représente un groupe méthyle ou un groupe éthyle dans les cas où ledit R_1 représente un groupe hydroxyle.
2. Dérivé de bornane-3-spiro-1'-cyclopentane selon la revendication 1, dans lequel ledit R_1 est un atome d'hydrogène et ledit R_2 est un groupe méthoxyle, et qui est le spiro [méthoxy-2 bornane-3,1'-cyclopentane].
 3. Dérivé de bornane-3-spiro-1'-cyclopentane selon la revendication 1, dans lequel ledit R_1 est un atome d'hydrogène et ledit R_2 est un groupe éthoxyle, et qui est le spiro [éthoxy-2 bornane-3,1'-cyclopentane].
 4. Dérivé de bornane-3-spiro-1'-cyclopentane selon la revendication 1, dans lequel ledit R_1 est un atome d'hydrogène et ledit R_2 est un groupe formyloxyle, et qui est le spiro [formyloxy-2 bornane-3,1'-cyclopentane].
 5. Dérivé de bornane-3-spiro-1'-cyclopentane selon la revendication 1, dans lequel ledit R_1 est un atome d'hydrogène et ledit R_2 est un groupe acétoxyle, et qui est le spiro [acétoxy-2 bornane-3,1'-cyclopentane].
 6. Dérivé de bornane-3-spiro-1'-cyclopentane selon la revendication 1, dans lequel ledit R_1 est un groupe hydroxyle et ledit R_2 est un groupe méthyle, et qui est le spiro [méthyl-2 bornane-3,1'-cyclopentanol-2].
 7. Dérivé de bornane-3-spiro-1'-cyclopentane selon la revendication 1, dans lequel ledit R_1 est un groupe hydroxyle et ledit R_2 est un groupe éthyle, et qui est le spiro [éthyl-2 bornane-3,1'-cyclopentanol-2].
 8. Composition pour produits de parfumerie comprenant un dérivé de bornane-3-spiro-1'-cyclopentane, représenté par la formule générale :



(I)


dans laquelle :



- R₁ représente un atome d'hydrogène ou un groupe hydroxyle ; et
- R₂ représente un groupe méthoxyle, un groupe éthoxyle, un groupe formyloxyle ou un groupe acétoxyle, dans les cas où ledit R₁ est un atome d'hydrogène ou bien représente un groupe méthyle ou un groupe éthyle dans les cas où ledit R₁ est un groupe hydroxyle.

Patentansprüche

1. Derivat Von Borman-3-spiro-1'-cyclopentan, dargestellt durch die allgemeine Formel




(I)

(I)

in der R₁ für ein Wasserstoffatom oder eine Hydroxylgruppe steht und R₂ für eine Methoxylgruppe, eine Ethoxylgruppe, eine Formyloxylgruppe oder eine Acetoxylgruppe steht in den Fällen, in denen R₁ ein Wasserstoffatom ist, oder für eine Methylgruppe oder eine Ethylgruppe steht in den Fällen, in denen R₁ eine Hydroxylgruppe ist.

2. Derivat von Borman-3-spiro-1'-cyclopentan gemäß Anspruch 1, wobei R₁ für ein Wasserstoffatom steht und R₂ für eine Methoxylgruppe steht, nämlich Spiro[2-methoxyborman-3,1'-cyclopentan].
3. Derivat von Borman-3-spiro-1'-cyclopentan gemäß Anspruch 1, wobei R₁ für ein Wasserstoffatom steht und R₂ eine Ethoxylgruppe ist, nämlich Spiro[2-ethoxyborman-3,1'-cyclopentan].
4. Derivat von Borman-3-spiro-1'-cyclopentan gemäß Anspruch 1, wobei R₁ für ein Wasserstoffatom steht und R₂ eine Formyloxylgruppe ist, nämlich Spiro[2-formyloxyborman-3,1'-cyclopentan].
5. Derivat von Borman-3-spiro-1'-cyclopentan gemäß Anspruch 1, wobei R₁ für ein Wasserstoffatom steht und R₂ eine Acetoxylgruppe ist, nämlich Spiro[2-acetoxyborman-3,1'-cyclopentan].
6. Derivat von Borman-3-spiro-1'-cyclopentan gemäß Anspruch 1, wobei R₁ für eine Hydroxylgruppe steht und R₂ eine Methylgruppe ist, nämlich Spiro[2-methylborman-3,1'-cyclopentan-2-ol].
7. Derivat von Borman-3-spiro-1'-cyclopentan gemäß Anspruch 1, wobei R₁ für eine Hydroxylgruppe steht und R₂ eine Ethylgruppe ist, nämlich Spiro[2-ethylborman-3,1'-cyclopentan-2-ol].
8. Parfümzusammensetzung, umfassend ein Derivat von Borman-3-spiro-1'-cyclopentan, dargestellt durch die allgemeine Formel



(I)

(I)

in der R₁ für ein Wasserstoffatom oder eine Hydroxylgruppe steht und R₂ für eine Methoxylgruppe, eine Ethoxylgruppe, eine Formyloxylgruppe oder eine Acetoxylgruppe steht in den Fällen, in denen R₁ ein Wasserstoffatom ist, oder für eine Methylgruppe oder eine Ethylgruppe steht in den Fällen, in denen R₁ eine Hydroxylgruppe ist.